



US011795498B2

(12) **United States Patent**
Frisen et al.

(10) **Patent No.:** **US 11,795,498 B2**
(45) **Date of Patent:** **Oct. 24, 2023**

(54) **METHODS OF DETECTING ANALYTES**

G16B 50/20 (2019.02); *G16B 50/30* (2019.02); *C12Y 600/00* (2013.01)

(71) Applicant: **10x Genomics Sweden AB**, Stockholm (SE)

(58) **Field of Classification Search**
CPC C12Q 1/682; C12Q 1/6837; C12Q 1/6806; C12Q 1/6816; C12Q 1/6827; C12Q 1/6841; C12Q 1/6844; C12Q 1/6874; C12Q 1/6876; G01N 1/30; G01N 1/42; C12N 15/1065

(72) Inventors: **Jonas Frisen**, Stockholm (SE); **Patrik Stahl**, Stockholm (SE); **Joakim Lundeborg**, Stockholm (SE)

See application file for complete search history.

(73) Assignee: **10x Genomics Sweden AB**, Stockholm (SE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(56) **References Cited**

U.S. PATENT DOCUMENTS

(21) Appl. No.: **18/170,285**

4,514,388 A	4/1985	Psaledakis
4,574,729 A	3/1986	Wells
4,683,195 A	7/1987	Mullis
4,683,202 A	7/1987	Mullis
4,800,159 A	1/1989	Mullis
4,883,867 A	11/1989	Lee
4,965,188 A	10/1990	Mullis
4,968,601 A	11/1990	Jacobson et al.
4,988,617 A	1/1991	Landegren et al.
5,002,882 A	3/1991	Lunnen
5,061,049 A	10/1991	Hornbeck
5,130,238 A	7/1992	Malek
5,183,053 A	2/1993	Yeh et al.

(22) Filed: **Feb. 16, 2023**

(65) **Prior Publication Data**

US 2023/0242973 A1 Aug. 3, 2023

Related U.S. Application Data

(63) Continuation of application No. 18/047,092, filed on Oct. 17, 2022, which is a continuation of application No. 17/704,830, filed on Mar. 25, 2022, now Pat. No. 11,479,809, which is a continuation of application No. 17/474,922, filed on Sep. 14, 2021, now Pat. No. 11,352,659, which is a continuation of application No. 16/013,654, filed on Jun. 20, 2018, now abandoned, which is a continuation of application No. 14/111,482, filed as application No. PCT/EP2012/056823 on Apr. 13, 2012, now Pat. No. 10,030,261.

(Continued)

FOREIGN PATENT DOCUMENTS

AU	2003200718	10/2006
CN	1273609	11/2000

(Continued)

OTHER PUBLICATIONS

(30) **Foreign Application Priority Data**

U.S. Appl. No. 16/353,937, filed Mar. 14, 2019, Frisen et al.
(Continued)

Apr. 13, 2011 (GB) 1106254

Primary Examiner — David C Thomas

(51) **Int. Cl.**

<i>C12Q 1/6837</i>	(2018.01)
<i>C12Q 1/6841</i>	(2018.01)
<i>C12Q 1/6844</i>	(2018.01)
<i>G16B 50/30</i>	(2019.01)
<i>G16B 50/20</i>	(2019.01)
<i>G16B 30/00</i>	(2019.01)
<i>C12Q 1/6806</i>	(2018.01)
<i>C12Q 1/6816</i>	(2018.01)
<i>C12Q 1/6827</i>	(2018.01)
<i>C12Q 1/6876</i>	(2018.01)
<i>C12N 15/10</i>	(2006.01)
<i>C12Q 1/682</i>	(2018.01)
<i>G01N 1/30</i>	(2006.01)
<i>G01N 1/42</i>	(2006.01)

(74) *Attorney, Agent, or Firm* — Fish & Richardson P.C.

(52) **U.S. Cl.**

CPC *C12Q 1/6837* (2013.01); *C12N 15/1065* (2013.01); *C12Q 1/682* (2013.01); *C12Q 1/6806* (2013.01); *C12Q 1/6816* (2013.01); *C12Q 1/6827* (2013.01); *C12Q 1/6841* (2013.01); *C12Q 1/6844* (2013.01); *C12Q 1/6876* (2013.01); *G01N 1/30* (2013.01); *G01N 1/42* (2013.01); *G16B 30/00* (2019.02);

(57) **ABSTRACT**

Localized detection of RNA in a tissue sample that includes cells is accomplished on an array. The array include a number of features on a substrate. Each feature includes a different capture probe immobilized such that the capture probe has a free 3' end. Each feature occupies a distinct position on the array and has an area of less than about 1 mm². Each capture probe is a nucleic acid molecule, which includes a positional domain including a nucleotide sequence unique to a particular feature, and a capture domain including a nucleotide sequence complementary to the RNA to be detected. The capture domain can be at a position 3' of the positional domain.

42 Claims, 28 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,308,751	A	5/1994	Ohkawa	6,391,937	B1	5/2002	Beuhler et al.
5,321,130	A	6/1994	Yue	6,401,267	B1	6/2002	Drmanac
5,410,030	A	4/1995	Yue	6,404,907	B1	6/2002	Gilchrist
5,436,134	A	7/1995	Haugland	6,416,950	B1	7/2002	Lohse
5,455,166	A	10/1995	Walker	6,432,360	B1	8/2002	Church et al.
5,472,881	A	12/1995	Beebe et al.	6,485,982	B1	11/2002	Charlton
5,494,810	A	2/1996	Barany et al.	6,503,713	B1	1/2003	Rana
5,503,980	A	4/1996	Cantor	6,506,561	B1	1/2003	Cheval et al.
5,512,439	A	4/1996	Hornes	6,518,018	B1	2/2003	Szostak
5,512,462	A	4/1996	Cheng	6,534,266	B1	3/2003	Singer
5,559,032	A	9/1996	Porneroy	6,544,732	B1	4/2003	Chee
5,582,977	A	12/1996	Yue	6,544,790	B1	4/2003	Sabatini
5,589,173	A	12/1996	O'Brien	6,565,727	B1	5/2003	Shenderov
5,599,675	A	2/1997	Brenner	6,573,043	B1	6/2003	Cohen et al.
5,610,287	A	3/1997	Nikiforov et al.	6,579,695	B1	6/2003	Lambalot
5,641,658	A	6/1997	Adams	6,620,584	B1	9/2003	Chee
5,648,245	A	7/1997	Fire et al.	6,632,641	B1	10/2003	Brennan
5,658,751	A	8/1997	Yue	6,673,620	B1	1/2004	Loeffler
5,695,940	A	12/1997	Drmanac et al.	6,677,160	B1	1/2004	Stockman et al.
5,716,825	A	2/1998	Hancock et al.	6,699,710	B1	3/2004	Kononen
5,750,341	A	5/1998	Macevicz	6,737,236	B1	5/2004	Picken et al.
5,763,175	A	6/1998	Brenner	6,770,441	B2	8/2004	Dickinson
5,807,522	A	9/1998	Brown et al.	6,773,566	B2	8/2004	Shenderov
5,830,711	A	11/1998	Barany et al.	6,773,886	B2	8/2004	Kaufman
5,837,832	A	11/1998	Chee et al.	6,787,308	B2	9/2004	Balasubramanian
5,837,860	A	11/1998	Anderson et al.	6,797,470	B2	9/2004	Barany et al.
5,854,033	A	12/1998	Lizardi	6,800,453	B2	10/2004	Labaer
5,863,753	A	1/1999	Haugland	6,812,005	B2	11/2004	Fan et al.
5,866,377	A	2/1999	Kim et al.	6,828,100	B1	12/2004	Ronaghi
5,871,921	A	2/1999	Landegren et al.	6,833,246	B2	12/2004	Balasubramanian
5,912,148	A	6/1999	Eggerding	6,852,487	B1	2/2005	Barany et al.
5,919,626	A	7/1999	Shi et al.	6,859,570	B2	2/2005	Walt
5,925,545	A	7/1999	Reznikoff et al.	6,864,052	B1	3/2005	Drmanac
5,928,906	A	7/1999	Koester et al.	6,867,028	B2	3/2005	Janulaitis
5,958,775	A	9/1999	Wickstrom et al.	6,872,816	B1	3/2005	Hall et al.
5,965,443	A	10/1999	Reznikoff et al.	6,875,572	B2	4/2005	Prudent et al.
6,013,440	A	1/2000	Lipshutz	6,878,515	B1	4/2005	Landegren
6,027,889	A	2/2000	Barany et al.	6,890,741	B2	5/2005	Fan et al.
6,054,274	A	4/2000	Sampson et al.	6,897,023	B2	5/2005	Fu
6,060,240	A	5/2000	Kamb et al.	6,911,132	B2	6/2005	Pamula
6,083,761	A	7/2000	Kedar et al.	6,911,345	B2	6/2005	Quake
6,130,073	A	10/2000	Eggerding	6,913,881	B1	7/2005	Aizenstein et al.
6,136,592	A	10/2000	Leighton	6,913,921	B2	7/2005	Fischer
6,143,496	A	11/2000	Brown	6,942,968	B1	9/2005	Dickinson et al.
6,153,389	A	11/2000	Haarer	6,969,488	B2	11/2005	Bridgham
6,157,432	A	12/2000	Helbing	6,969,589	B2	11/2005	Patil
6,159,736	A	12/2000	Reznikoff et al.	6,977,033	B2	12/2005	Becker
6,165,714	A	12/2000	Lane et al.	7,001,792	B2	2/2006	Sauer et al.
6,172,218	B1	1/2001	Brenner	7,011,944	B2	3/2006	Prudent et al.
6,210,891	B1	4/2001	Nyren	7,052,244	B2	5/2006	Fouillet
6,210,894	B1	4/2001	Brennan	7,057,026	B2	6/2006	Barnes
6,214,587	B1	4/2001	Dattagupta	7,083,980	B2	8/2006	Reznikoff et al.
6,221,591	B1	4/2001	Aerts	7,098,041	B2	8/2006	Kaylor et al.
6,221,654	B1	4/2001	Quake	7,115,400	B1	10/2006	Adessi
6,251,639	B1	6/2001	Kurn	7,118,883	B2	10/2006	Inoue
6,258,558	B1	7/2001	Szostak	7,128,893	B2	10/2006	Leamon et al.
6,258,568	B1	7/2001	Nyren	7,163,612	B2	1/2007	Sterling
6,261,804	B1	7/2001	Szostak	7,166,431	B2	1/2007	Chee et al.
6,265,552	B1	7/2001	Schatz	7,192,735	B2	3/2007	Lambalot
6,266,459	B1	7/2001	Walt	7,211,414	B2	5/2007	Hardin
6,268,148	B1	7/2001	Barany et al.	7,223,371	B2	5/2007	Hayenga et al.
6,274,320	B1	8/2001	Rothberg	7,229,769	B2	6/2007	Kozlov
6,281,804	B1	8/2001	Haller	7,244,559	B2	7/2007	Rothberg
6,291,180	B1	9/2001	Chu	7,255,994	B2	8/2007	Lao
6,291,187	B1	9/2001	Kingsmore et al.	7,258,976	B2	8/2007	Mitsubishi
6,300,063	B1	10/2001	Lipshutz et al.	7,259,258	B2	8/2007	Kozlov et al.
6,306,597	B1	10/2001	Macevicz	7,264,929	B2	9/2007	Rothberg
6,309,824	B1	10/2001	Drmanac	7,270,950	B2	9/2007	Szostak
6,323,009	B1	11/2001	Lasken et al.	7,282,328	B2	10/2007	Kong et al.
6,337,472	B1	1/2002	Garner et al.	7,297,518	B2	11/2007	Quake
6,344,316	B1	2/2002	Lockhart	7,315,019	B2	1/2008	Turner
6,344,329	B1	2/2002	Lizardi et al.	7,328,979	B2	2/2008	Decre
6,348,990	B1	2/2002	Igasaki et al.	7,329,492	B2	2/2008	Hardin
6,355,431	B1	3/2002	Chee	7,358,047	B2	4/2008	Hafner et al.
6,368,801	B1	4/2002	Faruqi	7,361,488	B2	4/2008	Fan et al.
				7,375,234	B2	5/2008	Sharpless et al.
				7,378,242	B2	5/2008	Hurt
				7,393,665	B2	7/2008	Brenner
				7,405,281	B2	7/2008	Xu

(56)

References Cited

U.S. PATENT DOCUMENTS

7,407,757 B2	8/2008	Brenner	8,519,115 B2	8/2013	Webster et al.
7,427,678 B2	9/2008	Pieken et al.	8,551,710 B2	10/2013	Bernitz et al.
7,456,012 B2	11/2008	Ryttzen et al.	8,568,979 B2	10/2013	Stuelpnagel et al.
7,462,449 B2	12/2008	Quake	RE44,596 E	11/2013	Stroun et al.
7,473,767 B2	1/2009	Dimitrov	8,586,310 B2	11/2013	Mitra
7,499,806 B2	3/2009	Kermani et al.	8,597,891 B2	12/2013	Barany et al.
7,501,245 B2	3/2009	Quake	8,603,743 B2	12/2013	Liu et al.
7,534,991 B2	5/2009	Miller et al.	8,604,182 B2	12/2013	Luo et al.
7,537,897 B2	5/2009	Brenner	8,614,073 B2	12/2013	Van Eijk
7,544,473 B2	6/2009	Brennan	8,624,016 B2	1/2014	Barany et al.
7,547,380 B2	6/2009	Velev	8,637,242 B2	1/2014	Shen
7,555,155 B2	6/2009	Levenson et al.	8,658,361 B2	2/2014	Wu et al.
7,561,336 B2	7/2009	Osaka et al.	8,685,889 B2	4/2014	Van Eijk
7,563,576 B2	7/2009	Chee	8,741,564 B2	6/2014	Seligmann
7,579,153 B2	8/2009	Brenner	8,741,606 B2	6/2014	Casbon
7,582,420 B2	9/2009	Oliphant et al.	8,748,103 B2	6/2014	Faham et al.
7,595,883 B1	9/2009	El Gamal	8,771,950 B2	7/2014	Church et al.
7,601,492 B2	10/2009	Fu et al.	8,778,849 B2	7/2014	Bowen
7,601,498 B2	10/2009	Mao	8,785,353 B2	7/2014	Van Eijk
7,608,434 B2	10/2009	Reznikoff et al.	8,790,873 B2	7/2014	Namsaraev et al.
7,611,869 B2	11/2009	Fan	8,809,238 B2	8/2014	Livak et al.
7,635,566 B2	12/2009	Brenner	8,815,512 B2	8/2014	Van Eijk
7,641,779 B2	1/2010	Becker	8,835,358 B2	9/2014	Fodor
7,655,898 B2	2/2010	Miller	8,865,410 B2	10/2014	Shendure
7,666,612 B2	2/2010	Johnsson	8,895,249 B2	11/2014	Shen
7,674,589 B2	3/2010	Cohen et al.	8,906,626 B2	12/2014	Oliphant et al.
7,674,752 B2	3/2010	He	8,911,945 B2	12/2014	Van Eijk
7,700,286 B2	4/2010	Stroun et al.	8,936,912 B2	1/2015	Mitra
7,709,198 B2	5/2010	Luo et al.	8,951,726 B2	2/2015	Luo et al.
7,754,429 B2	7/2010	Rigatti	8,951,728 B2	2/2015	Rasmussen
7,776,547 B2	8/2010	Roth	8,951,781 B2	2/2015	Reed
7,776,567 B2	8/2010	Mao	8,986,926 B2	3/2015	Ferree et al.
7,785,869 B2	8/2010	Belgrader et al.	9,005,891 B2	4/2015	Sinicropi et al.
7,803,943 B2	9/2010	Mao	9,005,935 B2	4/2015	Belyaev
7,844,940 B2	11/2010	Shin et al.	9,023,768 B2	5/2015	Van Eijk
7,848,553 B2	12/2010	Hertel et al.	9,062,348 B1	6/2015	Van Eijk
7,858,321 B2	12/2010	Glezer	9,080,210 B2	7/2015	Van Eijk
7,888,009 B2	2/2011	Barany et al.	9,085,798 B2	7/2015	Chee
7,892,747 B2	2/2011	Barany et al.	9,121,069 B2	9/2015	Lo
7,910,304 B2	3/2011	Drmanac	9,163,283 B2	10/2015	Chee et al.
7,914,981 B2	3/2011	Barany et al.	9,194,001 B2	11/2015	Brenner
7,941,279 B2	5/2011	Hwang et al.	9,201,063 B2	12/2015	Sood et al.
7,955,794 B2	6/2011	Shen et al.	9,217,176 B2	12/2015	Faham et al.
7,960,119 B2	6/2011	Chee	9,273,349 B2	3/2016	Nguyen et al.
7,960,120 B2	6/2011	Rigatti	9,290,808 B2	3/2016	Fodor
7,985,565 B2	7/2011	Mayer et al.	9,290,809 B2	3/2016	Fodor
8,003,354 B2	8/2011	Shen et al.	9,309,556 B2	4/2016	Myllykangas et al.
8,030,477 B2	10/2011	Cerrina et al.	9,328,383 B2	5/2016	Van Eijk
8,076,063 B2	12/2011	Fan	9,334,536 B2	5/2016	Van Eijk
8,092,784 B2	1/2012	Mao	9,340,830 B2	5/2016	Lipson
8,124,751 B2	2/2012	Pierce et al.	9,371,563 B2	6/2016	Geiss et al.
8,148,068 B2	4/2012	Brenner	9,371,598 B2	6/2016	Chee
8,148,518 B2	4/2012	Buchanan	9,376,716 B2	6/2016	Van Eijk
8,198,028 B2	6/2012	Rigatti et al.	9,376,717 B2	6/2016	Gao et al.
8,199,999 B2	6/2012	Hoyt et al.	9,376,719 B2	6/2016	Van Eijk
8,206,917 B2	6/2012	Chee	9,404,156 B2	8/2016	Hicks
8,207,093 B2	6/2012	Szostak	9,416,409 B2	8/2016	Hayden
8,268,554 B2	9/2012	Schallmeiner	9,447,459 B2	9/2016	Van Eijk
8,278,034 B2	10/2012	Muraca	9,453,256 B2	9/2016	Van Eijk
8,288,103 B2	10/2012	Oliphant	9,493,820 B2	11/2016	Van Eijk
8,288,122 B2	10/2012	O'Leary et al.	9,494,588 B2	11/2016	Springer et al.
8,330,087 B2	12/2012	Domenicali	9,506,061 B2	11/2016	Brown
8,337,851 B2	12/2012	Aukerman	9,512,422 B2	12/2016	Barnard et al.
8,343,500 B2	1/2013	Wraith	9,512,487 B2	12/2016	Faham et al.
8,383,338 B2	2/2013	Kitzman	9,518,980 B2	12/2016	Looger et al.
8,415,102 B2	4/2013	Geiss et al.	9,541,504 B2	1/2017	Hoyt
8,431,691 B2	4/2013	McKernan et al.	9,557,330 B2	1/2017	Siciliano et al.
8,460,865 B2	6/2013	Chee	9,574,230 B2	2/2017	Van Eijk
8,462,981 B2	6/2013	Determan et al.	9,593,365 B2	3/2017	Frisen et al.
8,481,257 B2	7/2013	Van Eijk	9,598,728 B2	3/2017	Barany et al.
8,481,258 B2	7/2013	Church et al.	9,624,538 B2	4/2017	Church et al.
8,481,292 B2	7/2013	Casbon	9,644,204 B2	5/2017	Hindson et al.
8,481,698 B2	7/2013	Lieberman et al.	9,657,335 B2	5/2017	Van Eijk
8,486,625 B2	7/2013	Gunderson	9,670,542 B2	6/2017	Van Eijk
8,507,204 B2	8/2013	Pierce et al.	9,694,361 B2	7/2017	Bharadwaj
			9,702,004 B2	7/2017	Van Eijk
			9,714,446 B2	7/2017	Webster et al.
			9,714,937 B2	7/2017	Dunaway
			9,727,810 B2	8/2017	Fodor et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

9,745,627 B2	8/2017	Van Eijk	11,332,790 B2	5/2022	Chell et al.
9,777,324 B2	10/2017	Van Eijk	11,352,659 B2	6/2022	Frisen et al.
9,816,134 B2	11/2017	Namsaraev	11,359,228 B2	6/2022	Chee et al.
9,834,814 B2	12/2017	Peter et al.	11,365,442 B2	6/2022	Chee
9,850,536 B2	12/2017	Oliphant et al.	11,371,086 B2	6/2022	Chee
9,868,979 B2	1/2018	Chee et al.	11,384,386 B2	7/2022	Chee
9,879,313 B2	1/2018	Chee et al.	11,390,912 B2	7/2022	Frisen et al.
9,889,422 B2	2/2018	Smith et al.	11,401,545 B2	8/2022	Chee
9,896,721 B2	2/2018	Van Eijk	11,407,992 B2	8/2022	Dadhwal
9,898,576 B2	2/2018	Van Eijk	11,408,029 B2	8/2022	Katirae et al.
9,898,577 B2	2/2018	Van Eijk	11,434,524 B2	9/2022	Ramachandran Iyer et al.
9,902,950 B2	2/2018	Church et al.	11,479,809 B2	10/2022	Frisen et al.
9,902,991 B2	2/2018	Sinicropi et al.	11,479,810 B1	10/2022	Chee
9,958,454 B2	5/2018	Kozlov et al.	11,492,612 B1	11/2022	Dadhwal
10,002,316 B2	6/2018	Fodor et al.	11,505,828 B2	11/2022	Chell et al.
10,023,907 B2	7/2018	Van Eijk	11,512,308 B2	11/2022	Gallant et al.
10,030,261 B2	7/2018	Frisen et al.	11,519,022 B2	12/2022	Chee
10,032,064 B2	7/2018	Hoyt	11,519,033 B2	12/2022	Schnall-Levin et al.
10,053,723 B2	8/2018	Hindson et al.	11,535,887 B2	12/2022	Gallant et al.
10,072,104 B2	9/2018	Winnik et al.	11,542,543 B2	1/2023	Chee
10,095,832 B2	10/2018	Van Eijk	11,549,138 B2	1/2023	Chee
10,138,509 B2	11/2018	Church et al.	11,560,587 B2	1/2023	Chee
10,144,966 B2	12/2018	Cantor	11,560,592 B2	1/2023	Chew et al.
10,179,932 B2	1/2019	Church et al.	11,560,593 B2	1/2023	Chell et al.
10,196,691 B2	2/2019	Harkin et al.	11,592,447 B2	2/2023	Uytingco et al.
10,221,461 B2	3/2019	Robins et al.	11,608,498 B2	3/2023	Gallant et al.
10,227,639 B2	3/2019	Levner et al.	11,608,520 B2	3/2023	Galonska et al.
10,246,752 B2	4/2019	Faham et al.	11,613,773 B2	3/2023	Frisen et al.
10,266,876 B2	4/2019	Cai et al.	11,618,897 B2	4/2023	Kim et al.
10,267,808 B2	4/2019	Cai	11,618,918 B2	4/2023	Chee et al.
10,273,541 B2	4/2019	Hindson et al.	11,624,063 B2	4/2023	Dadhwal
10,308,982 B2	6/2019	Chee	11,624,086 B2	4/2023	Uytingco et al.
10,357,771 B2	7/2019	Bharadwaj	11,634,756 B2	4/2023	Chee
10,465,235 B2	11/2019	Gullberg et al.	11,649,485 B2	5/2023	Yin et al.
10,472,669 B2	11/2019	Chee	11,661,626 B2	5/2023	Katirae et al.
10,480,022 B2	11/2019	Chee	11,680,260 B2	6/2023	Kim et al.
10,494,662 B2	12/2019	Church et al.	11,692,218 B2	7/2023	Engblom et al.
10,494,667 B2	12/2019	Chee	11,702,693 B2	7/2023	Bharadwaj
10,501,791 B2	12/2019	Church et al.	11,702,698 B2	7/2023	Stoeckius
10,545,075 B2	1/2020	Deisseroth et al.	2001/0039029 A1	11/2001	Nemori et al.
10,580,128 B2	3/2020	Miller	2001/0055764 A1	12/2001	Empedocles et al.
10,612,079 B2	4/2020	Chee	2002/0006477 A1	1/2002	Shishido et al.
10,619,196 B1	4/2020	Chee	2002/0040275 A1	4/2002	Cravatt
10,640,826 B2	5/2020	Church et al.	2002/0045169 A1	4/2002	Shoemaker et al.
10,662,467 B2	5/2020	Chee	2002/0045272 A1	4/2002	McDevitt et al.
10,662,468 B2	5/2020	Chee	2002/0048766 A1	4/2002	Doyle et al.
10,669,569 B2	6/2020	Gullberg et al.	2002/0051986 A1	5/2002	Baez et al.
10,697,013 B1	6/2020	Brenner et al.	2002/0055100 A1	5/2002	Kawashima
10,725,027 B2	7/2020	Bell	2002/0058250 A1	5/2002	Firth
10,767,223 B1	9/2020	Brenner et al.	2002/0064779 A1	5/2002	Landegren
10,774,372 B2	9/2020	Chee et al.	2002/0086441 A1	7/2002	Baranov et al.
10,774,374 B2	9/2020	Frisen et al.	2002/0132246 A1	9/2002	Kallioniemi et al.
10,787,701 B2	9/2020	Chee	2002/0137031 A1	9/2002	Wolber
10,829,803 B2	11/2020	Terbruggen et al.	2002/0150909 A1	10/2002	Stuelpnagel
10,858,702 B2	12/2020	Lucero et al.	2002/0164611 A1	11/2002	Bamdad
10,914,730 B2	2/2021	Chee et al.	2002/0168645 A1	11/2002	Taylor
10,927,403 B2	2/2021	Chee et al.	2003/0017451 A1	1/2003	Wang et al.
10,961,566 B2	3/2021	Chee	2003/0022207 A1	1/2003	Balasubramanian
10,962,532 B2	3/2021	Chee	2003/0040035 A1	2/2003	Slamon
10,964,001 B2	3/2021	Miller	2003/0064398 A1	4/2003	Barnes
10,982,268 B2	4/2021	Chee	2003/0073086 A1	4/2003	Guire et al.
10,983,113 B2	4/2021	Chee	2003/0087232 A1	5/2003	Christians
10,996,219 B2	5/2021	Chee et al.	2003/0096323 A1	5/2003	James
11,001,878 B1	5/2021	Chee et al.	2003/0113713 A1	6/2003	Glezer
11,001,879 B1	5/2021	Chee et al.	2003/0124595 A1	7/2003	Lizardi
11,008,607 B2	5/2021	Chee	2003/0134279 A1	7/2003	Isola et al.
11,046,996 B1	6/2021	Chee et al.	2003/0138879 A1	7/2003	Lambalot
11,067,567 B2	7/2021	Chee	2003/0148335 A1	8/2003	Shen et al.
11,156,603 B2	10/2021	Chee	2003/0153850 A1	8/2003	Davis et al.
11,162,132 B2	11/2021	Frisen et al.	2003/0162216 A1	8/2003	Gold
11,208,684 B2	12/2021	Chee	2003/0165948 A1	9/2003	Alsmadi et al.
11,286,515 B2	3/2022	Chee et al.	2003/0170637 A1	9/2003	Pirrung et al.
11,293,917 B2	4/2022	Chee	2003/0175844 A1	9/2003	Nadler et al.
11,299,774 B2	4/2022	Frisen et al.	2003/0175947 A1	9/2003	Liu et al.
11,313,856 B2	4/2022	Chee	2003/0190744 A1	10/2003	McGarry et al.
			2003/0205632 A1	11/2003	Kim et al.
			2003/0211489 A1	11/2003	Shen et al.
			2003/0215936 A1	11/2003	Kallioniemi et al.
			2003/0224419 A1	12/2003	Corcoran

(56)		References Cited					
		U.S. PATENT DOCUMENTS					
2003/0232348	A1	12/2003	Jones et al.	2006/0216775	A1	9/2006	Burkart et al.
2003/0232382	A1	12/2003	Brennan	2006/0228758	A1	10/2006	Muchhal et al.
2003/0235535	A1	12/2003	Zhou	2006/0240439	A1	10/2006	Smith et al.
2003/0235852	A1	12/2003	Roberts	2006/0263789	A1	11/2006	Kincaid
2003/0235854	A1	12/2003	Chan et al.	2006/0275782	A1	12/2006	Gunderson et al.
2004/0002090	A1	1/2004	Mayer et al.	2006/0275799	A1	12/2006	Banerjee et al.
2004/0019005	A1	1/2004	Van Ness	2006/0281109	A1	12/2006	Barr Ost et al.
2004/0023320	A1	2/2004	Steiner et al.	2007/0003954	A1	1/2007	Kodadek et al.
2004/0033499	A1	2/2004	Ilisley et al.	2007/0014810	A1	1/2007	Baker et al.
2004/0050699	A1	3/2004	Goncalves	2007/0020625	A1	1/2007	Duchaud et al.
2004/0067492	A1	4/2004	Peng et al.	2007/0020640	A1	1/2007	McCloskey et al.
2004/0067493	A1	4/2004	Matsuzaki	2007/0020669	A1	1/2007	Ericsson
2004/0082059	A1	4/2004	Webb	2007/0023292	A1	2/2007	Kim et al.
2004/0096853	A1	5/2004	Mayer	2007/0026430	A1	2/2007	Andersen et al.
2004/0106110	A1	6/2004	Balasubramanian	2007/0036511	A1	2/2007	Lundquist et al.
2004/0112442	A1	6/2004	Maerkl	2007/0048812	A1	3/2007	Moravec et al.
2004/0121456	A1	6/2004	Fischer	2007/0054288	A1	3/2007	Su et al.
2004/0175822	A1	9/2004	Timperman et al.	2007/0087360	A1	4/2007	Boyd
2004/0219588	A1	11/2004	Furuta	2007/0099208	A1	5/2007	Drmanac et al.
2004/0224326	A1	11/2004	Kim et al.	2007/0116612	A1	5/2007	Williamson
2004/0235103	A1	11/2004	Reznikoff et al.	2007/0128071	A1	6/2007	Shea et al.
2004/0248325	A1	12/2004	Bukusoglu et al.	2007/0128624	A1	6/2007	Gormley et al.
2004/0259105	A1	12/2004	Fan et al.	2007/0128656	A1	6/2007	Agrawal
2005/0003431	A1	1/2005	Wucherpfennig	2007/0134723	A1	6/2007	Kozlov et al.
2005/0014203	A1	1/2005	Darfler et al.	2007/0141718	A1	6/2007	Bui et al.
2005/0019776	A1	1/2005	Callow et al.	2007/0161020	A1	7/2007	Luo et al.
2005/0019842	A1	1/2005	Prober et al.	2007/0161029	A1	7/2007	Li et al.
2005/0026188	A1	2/2005	Van Kessel	2007/0166705	A1	7/2007	Milton et al.
2005/0037362	A1	2/2005	Remacle et al.	2007/0166725	A1	7/2007	McBride et al.
2005/0037393	A1	2/2005	Gunderson et al.	2007/0172873	A1	7/2007	Brenner et al.
2005/0042695	A1	2/2005	Meares et al.	2007/0178503	A1	8/2007	Jiang
2005/0048580	A1	3/2005	Labaer	2007/0207482	A1	9/2007	Church et al.
2005/0064460	A1	3/2005	Holliger et al.	2007/0215466	A1	9/2007	Okada
2005/0079520	A1	4/2005	Wu	2007/0231823	A1	10/2007	McKernan
2005/0095627	A1	5/2005	Kolman et al.	2007/0243634	A1	10/2007	Pamula et al.
2005/0100900	A1	5/2005	Kawashima et al.	2007/0251824	A1	11/2007	Patton
2005/0116161	A1	6/2005	Hafeman et al.	2007/0254305	A1	11/2007	Paik et al.
2005/0130173	A1	6/2005	Leamon et al.	2007/0264656	A1	11/2007	Kawamura
2005/0130188	A1	6/2005	Walt	2007/0269805	A1	11/2007	Hogers
2005/0136414	A1	6/2005	Gunderson et al.	2007/0280517	A1	12/2007	De La Torre-Bueno et al.
2005/0164292	A1	7/2005	Farooqui	2008/0003586	A1	1/2008	Hyde et al.
2005/0179746	A1	8/2005	Roux et al.	2008/0009420	A1	1/2008	Schroth et al.
2005/0191656	A1	9/2005	Drmanac et al.	2008/0032301	A1	2/2008	Rank et al.
2005/0191698	A1	9/2005	Chee et al.	2008/0038734	A1	2/2008	Sorge et al.
2005/0196786	A1	9/2005	Levy	2008/0047835	A1	2/2008	MacConnell
2005/0202433	A1	9/2005	Van Beuningen	2008/0071071	A1	3/2008	LaBaer et al.
2005/0226780	A1	10/2005	Sandell et al.	2008/0108082	A1	5/2008	Rank et al.
2005/0227271	A1	10/2005	Kwon	2008/0108804	A1	5/2008	Hayashizaki et al.
2005/0239119	A1	10/2005	Tsukada et al.	2008/0124252	A1	5/2008	Marchand et al.
2005/0239192	A1	10/2005	Nasarabadi et al.	2008/0124810	A1	5/2008	Terbrueggen et al.
2005/0244850	A1	11/2005	Huang	2008/0128627	A1	6/2008	Lundquist et al.
2005/0255548	A1	11/2005	Lipovsek et al.	2008/0132429	A1	6/2008	Perov et al.
2005/0257284	A1	11/2005	Nakajima et al.	2008/0145616	A1	6/2008	Gharib et al.
2005/0260653	A1	11/2005	LaBaer	2008/0153086	A1	6/2008	Wong
2005/0266417	A1	12/2005	Barany et al.	2008/0160580	A1	7/2008	Adessi et al.
2006/0003394	A1	1/2006	Song	2008/0199929	A1	8/2008	Yeung et al.
2006/0039823	A1	2/2006	Yamakawa et al.	2008/0220434	A1	9/2008	Thomas
2006/0046313	A1	3/2006	Roth	2008/0220981	A1	9/2008	McGregor
2006/0063160	A1	3/2006	West et al.	2008/0261204	A1	10/2008	Lexow
2006/0079453	A1	4/2006	Sidney et al.	2008/0280773	A1	11/2008	Fedurco et al.
2006/0084078	A1	4/2006	Zhao	2008/0286795	A1	11/2008	Kawashima et al.
2006/0105352	A1	5/2006	Qiao et al.	2008/0293046	A1	11/2008	Allawi et al.
2006/0110739	A1	5/2006	Heyduk	2008/0293591	A1	11/2008	Taussig et al.
2006/0134669	A1	6/2006	Casasanta	2008/0312103	A1	12/2008	Nemoto et al.
2006/0154286	A1	7/2006	Kong et al.	2009/0005252	A1	1/2009	Drmanac et al.
2006/0164490	A1	7/2006	Kim et al.	2009/0006002	A1	1/2009	Honisch et al.
2006/0183150	A1	8/2006	Cohen et al.	2009/0018024	A1	1/2009	Church et al.
2006/0188875	A1	8/2006	Cox et al.	2009/0023148	A1	1/2009	Moyle et al.
2006/0188901	A1	8/2006	Barnes et al.	2009/0026082	A1	1/2009	Rothberg et al.
2006/0188906	A1	8/2006	Kim et al.	2009/0036323	A1	2/2009	van Eijk et al.
2006/0194331	A1	8/2006	Pamula et al.	2009/0062148	A1	3/2009	Goldberg
2006/0199183	A1	9/2006	Valat et al.	2009/0082212	A1	3/2009	Williams
2006/0199207	A1	9/2006	Matysiak	2009/0099041	A1	4/2009	Church et al.
2006/0211001	A1	9/2006	Yu et al.	2009/0105959	A1	4/2009	Braverman et al.
2006/0216721	A1	9/2006	Kozlov et al.	2009/0117573	A1	5/2009	Fu et al.
				2009/0127589	A1	5/2009	Rothberg et al.
				2009/0155781	A1	6/2009	Drmanac et al.
				2009/0169089	A1	7/2009	Hunt et al.
				2009/0170713	A1	7/2009	van Eijk et al.

(56)		References Cited					
		U.S. PATENT DOCUMENTS					
2009/0192044	A1	7/2009	Fouillet	2012/0065081	A1	3/2012	Chee
2009/0197326	A1	8/2009	El Gamal et al.	2012/0077693	A1	3/2012	Cazalis et al.
2009/0202998	A1	8/2009	Schlumpberger et al.	2012/0129248	A1	5/2012	Chee et al.
2009/0215633	A1	8/2009	van Eijk et al.	2012/0135871	A1	5/2012	van Eijk et al.
2009/0233802	A1	9/2009	Bignell et al.	2012/0142014	A1	6/2012	Cai
2009/0239232	A1	9/2009	Kurn	2012/0157322	A1	6/2012	Myllykangas
2009/0253163	A1	10/2009	Xie et al.	2012/0160683	A1	6/2012	Ye et al.
2009/0253581	A1	10/2009	van Eijk et al.	2012/0195810	A1	8/2012	Cohen et al.
2009/0253582	A1	10/2009	Pena et al.	2012/0196297	A1	8/2012	Yost et al.
2009/0264299	A1	10/2009	Drmanac et al.	2012/0202698	A1	8/2012	van Eijk et al.
2009/0280487	A1	11/2009	Hung et al.	2012/0202704	A1	8/2012	Fan et al.
2009/0283407	A1	11/2009	Van Eijk	2012/0220479	A1	8/2012	Ericsson et al.
2009/0289184	A1	11/2009	Deininger	2012/0245053	A1	9/2012	Shirai et al.
2009/0291854	A1	11/2009	Weisinger-Mayr et al.	2012/0252702	A1	10/2012	Muratani et al.
2009/0305237	A1	12/2009	Cantor et al.	2012/0258871	A1	10/2012	Kozlov et al.
2009/0312193	A1	12/2009	Kim et al.	2012/0270305	A1	10/2012	Reed et al.
2009/0321262	A1	12/2009	Adachi et al.	2012/0270748	A1	10/2012	Chee et al.
2010/0009871	A1	1/2010	Reed et al.	2012/0279954	A1	11/2012	Ceremony et al.
2010/0014537	A1	1/2010	Jacquet et al.	2012/0289414	A1	11/2012	Mitra et al.
2010/0035249	A1	2/2010	Hayashizaki et al.	2012/0301925	A1	11/2012	Belyaev
2010/0055733	A1	3/2010	Lutolf et al.	2012/0322099	A1	12/2012	Lapen et al.
2010/0069263	A1	3/2010	Shendure et al.	2013/0005594	A1	1/2013	Terbrueggen et al.
2010/0096266	A1	4/2010	Kim et al.	2013/0005600	A1	1/2013	Olek
2010/0099103	A1	4/2010	Hsieh et al.	2013/0035239	A1	2/2013	Kong et al.
2010/0105052	A1	4/2010	Drmanac et al.	2013/0052331	A1	2/2013	Kram et al.
2010/0105112	A1	4/2010	Heltze et al.	2013/0065768	A1	3/2013	Zheng et al.
2010/0111768	A1	5/2010	Banerjee et al.	2013/0065788	A1	3/2013	Glezer et al.
2010/0113302	A1	5/2010	Williams	2013/0079232	A1	3/2013	Kain et al.
2010/0120043	A1	5/2010	Sood et al.	2013/0096033	A1	4/2013	Routenberg
2010/0120097	A1	5/2010	Matz et al.	2013/0109595	A1	5/2013	Routenberg
2010/0120098	A1	5/2010	Grunenwald et al.	2013/0122516	A1	5/2013	Hong et al.
2010/0126862	A1	5/2010	Sabin et al.	2013/0171621	A1	7/2013	Luo et al.
2010/0129874	A1	5/2010	Mitra et al.	2013/0202718	A1	8/2013	Pepin et al.
2010/0137143	A1	6/2010	Rothberg et al.	2013/0211249	A1	8/2013	Barnett et al.
2010/0145037	A1	6/2010	Makarov et al.	2013/0244884	A1	9/2013	Jacobson et al.
2010/0151464	A1	6/2010	Stuelpnagel et al.	2013/0260372	A1	10/2013	Buermann et al.
2010/0151511	A1	6/2010	Gereenizer et al.	2013/0261019	A1	10/2013	Lin et al.
2010/0159446	A1	6/2010	Haff et al.	2013/0296174	A1	11/2013	Peumans
2010/0173384	A1	7/2010	Johnsson et al.	2013/0302801	A1	11/2013	Asbury et al.
2010/0184614	A1	7/2010	Ye et al.	2013/0338042	A1	12/2013	Shen et al.
2010/0184618	A1	7/2010	Namsaraev et al.	2014/0065609	A1	3/2014	Hicks et al.
2010/0210475	A1	8/2010	Lee et al.	2014/0066318	A1	3/2014	Frisen et al.
2010/0227329	A1	9/2010	Cuppens	2014/0079923	A1	3/2014	George et al.
2010/0267590	A1	10/2010	Grudzien et al.	2014/0080715	A1	3/2014	Lo et al.
2010/0273219	A1	10/2010	May et al.	2014/0121118	A1	5/2014	Warner
2010/0273679	A1	10/2010	Cuppoletti et al.	2014/0155274	A1	6/2014	Xie et al.
2010/0282617	A1	11/2010	Rothberg et al.	2014/0155295	A1	6/2014	Hindson et al.
2010/0303722	A1	12/2010	Jin et al.	2014/0213533	A1	7/2014	Suthanthiran et al.
2011/0024511	A1	2/2011	Rietzler et al.	2014/0227705	A1	8/2014	Vogelstein et al.
2011/0027772	A1	2/2011	Ahn et al.	2014/0243224	A1	8/2014	Barnard et al.
2011/0028685	A1	2/2011	Purkayastha et al.	2014/0274731	A1	9/2014	Raymond et al.
2011/0033854	A1	2/2011	Drmanac et al.	2014/0296081	A1	10/2014	Diehn et al.
2011/0045462	A1	2/2011	Fu et al.	2014/0342921	A1	11/2014	Weiner
2011/0048951	A1	3/2011	Wu	2014/0378345	A1	12/2014	Hindson et al.
2011/0059436	A1	3/2011	Hardin et al.	2014/0378350	A1	12/2014	Hindson et al.
2011/0059865	A1	3/2011	Smith et al.	2015/0005447	A1	1/2015	Berti et al.
2011/0086774	A1	4/2011	Dunaway	2015/0051085	A1	2/2015	Vogelstein et al.
2011/0111409	A1	5/2011	Sinicropi et al.	2015/0072867	A1	3/2015	Soldatov
2011/0151451	A1	6/2011	Lemaire et al.	2015/0087027	A1	3/2015	Makarov et al.
2011/0152111	A1	6/2011	Fan et al.	2015/0219618	A1	8/2015	Krishnan et al.
2011/0172115	A1	7/2011	Thompson et al.	2015/0246336	A1	9/2015	Somoza et al.
2011/0177518	A1	7/2011	Kartalov et al.	2015/0344942	A1	12/2015	Frisen et al.
2011/0201515	A1	8/2011	Webster et al.	2015/0376609	A1	12/2015	Hindson et al.
2011/0207134	A1	8/2011	Faham et al.	2016/0003812	A1	1/2016	Porreca et al.
2011/0223613	A1	9/2011	Gut	2016/0019337	A1	1/2016	Roberts et al.
2011/0244448	A1	10/2011	Shirai et al.	2016/0024555	A1	1/2016	Church et al.
2011/0245101	A1	10/2011	Chee et al.	2016/0024576	A1	1/2016	Chee
2011/0245111	A1	10/2011	Chee	2016/0032282	A1	2/2016	Vigneault et al.
2011/0275077	A1	11/2011	James	2016/0033496	A1	2/2016	Chou et al.
2011/0287435	A1	11/2011	Grunenwald et al.	2016/0138091	A1	5/2016	Chee et al.
2012/0010091	A1*	1/2012	Linnarson C12N 15/1065 506/7	2016/0145677	A1	5/2016	Chee et al.
2012/0021930	A1	1/2012	Schoen et al.	2016/0298180	A1	10/2016	Chee
2012/0046175	A1	2/2012	Rodesch et al.	2016/0304952	A1	10/2016	Boyden et al.
2012/0046178	A1	2/2012	Van Den Boom et al.	2016/0333403	A1	11/2016	Chee
				2016/0376642	A1	12/2016	Landegren et al.
				2017/0029872	A1	2/2017	Bhattacharyya et al.
				2017/0058339	A1	3/2017	Chee
				2017/0058340	A1	3/2017	Chee
				2017/0058345	A1	3/2017	Chee

(56)		References Cited					
		U.S. PATENT DOCUMENTS					
2017/0088881	A1	3/2017	Chee	2021/0223227	A1	7/2021	Stoeckius
2017/0166962	A1	6/2017	van Eijk et al.	2021/0230681	A1	7/2021	Patterson et al.
2017/0242020	A1	8/2017	Yamauchi et al.	2021/0230692	A1	7/2021	Daugharthy et al.
2017/0343545	A1	11/2017	Hadrup et al.	2021/0237022	A1	8/2021	Bava
2018/0094316	A1	4/2018	Scott et al.	2021/0238581	A1	8/2021	Mikkelsen et al.
2018/0105808	A1	4/2018	Mikkelsen et al.	2021/0238664	A1	8/2021	Bava
2018/0112209	A1	4/2018	Eshoo et al.	2021/0238675	A1	8/2021	Bava
2018/0179591	A1	6/2018	van Eijk	2021/0238680	A1	8/2021	Bava
2018/0180601	A1	6/2018	Pedersen et al.	2021/0247316	A1	8/2021	Bava
2018/0201980	A1	7/2018	Chee et al.	2021/0255175	A1	8/2021	Chee et al.
2018/0216162	A1	8/2018	Belhocine et al.	2021/0262019	A1	8/2021	Alvarado Martinez et al.
2018/0217094	A1	8/2018	Herr et al.	2021/0269864	A1	9/2021	Chee
2018/0247017	A1	8/2018	van Eijk et al.	2021/0270822	A1	9/2021	Chee
2018/0291439	A1	10/2018	van Eijk et al.	2021/0285036	A1	9/2021	Yin et al.
2019/0017106	A1	1/2019	Frisen et al.	2021/0285046	A1	9/2021	Chell et al.
2019/0024153	A1	1/2019	Frisen et al.	2021/0292748	A1	9/2021	Frisen et al.
2019/0024154	A1	1/2019	Frisen et al.	2021/0292822	A1	9/2021	Frisen et al.
2019/0064173	A1	2/2019	Bharadwaj et al.	2021/0317510	A1	10/2021	Chee et al.
2019/0177777	A1	6/2019	Chee	2021/0317524	A1	10/2021	Lucero et al.
2019/0177778	A1	6/2019	Chee	2021/0324457	A1	10/2021	Ramachandran Iyer et al.
2019/0177789	A1	6/2019	Hindson et al.	2021/0332424	A1	10/2021	Schnall-Levin
2019/0177800	A1	6/2019	Boutet et al.	2021/0332425	A1	10/2021	Pfeiffer et al.
2019/0203275	A1	7/2019	Friesen et al.	2021/0348221	A1	11/2021	Chell et al.
2019/0264268	A1	8/2019	Frisen et al.	2022/0002791	A1	1/2022	Frisen et al.
2019/0271030	A1	9/2019	Chee	2022/0003755	A1	1/2022	Chee
2019/0271031	A1	9/2019	Chee	2022/0010367	A1	1/2022	Ramachandran Iyer et al.
2019/0300943	A1	10/2019	Chee et al.	2022/0017951	A1	1/2022	Ramachandran Iyer et al.
2019/0300944	A1	10/2019	Chee et al.	2022/0025446	A1	1/2022	Shah
2019/0300945	A1	10/2019	Chee et al.	2022/0025447	A1	1/2022	Tentori et al.
2019/0309353	A1	10/2019	Chee	2022/0033888	A1	2/2022	Schnall-Levin et al.
2019/0309354	A1	10/2019	Chee	2022/0049293	A1	2/2022	Frenz et al.
2019/0309355	A1	10/2019	Chee	2022/0049294	A1	2/2022	Uytingco et al.
2019/0323071	A1	10/2019	Chee	2022/0064630	A1	3/2022	Bent et al.
2019/0367982	A1	12/2019	Belhocine et al.	2022/0081728	A1	3/2022	Williams
2019/0367997	A1	12/2019	Bent et al.	2022/0090058	A1	3/2022	Frisen et al.
2020/0048690	A1	2/2020	Chee	2022/0090175	A1	3/2022	Uytingco et al.
2020/0063195	A1	2/2020	Chee	2022/0090181	A1	3/2022	Gallant et al.
2020/0063196	A1	2/2020	Chee	2022/0098576	A1	3/2022	Dadhwal
2020/0109443	A1	4/2020	Chee	2022/0098661	A1	3/2022	Chew et al.
2020/0140934	A1	5/2020	Chee	2022/0106632	A1	4/2022	Galonska et al.
2020/0140935	A1	5/2020	Chee	2022/0106633	A1	4/2022	Engblom et al.
2020/0208205	A1	7/2020	Chee	2022/0112486	A1	4/2022	Ramachandran Iyer et al.
2020/0208206	A1	7/2020	Chee	2022/0112545	A1	4/2022	Chee
2020/0224256	A1	7/2020	Chee	2022/0119869	A1	4/2022	Ramachandran Iyer et al.
2020/0277663	A1	9/2020	Ramachandran Iyer et al.	2022/0127659	A1	4/2022	Frisen et al.
2020/0277664	A1	9/2020	Frenz	2022/0127666	A1	4/2022	Katirae et al.
2020/0283852	A1	9/2020	Oliphant et al.	2022/0127672	A1	4/2022	Stoeckius
2020/0299757	A1	9/2020	Chee et al.	2022/0145361	A1	5/2022	Frenz et al.
2020/0325531	A1	10/2020	Chee	2022/0154255	A1	5/2022	Chee et al.
2020/0332368	A1	10/2020	Ferree et al.	2022/0170083	A1	6/2022	Khaled et al.
2020/0354774	A1	11/2020	Church et al.	2022/0195422	A1	6/2022	Gallant et al.
2020/0370106	A1	11/2020	Chee	2022/0195505	A1	6/2022	Frisen et al.
2020/0399687	A1	12/2020	Frisen et al.	2022/0196644	A1	6/2022	Chee
2020/0407781	A1	12/2020	Schnall-Levin et al.	2022/0213526	A1	7/2022	Frisen et al.
2021/0002713	A1	1/2021	Chee et al.	2022/0241780	A1	8/2022	Tentori et al.
2021/0010068	A1	1/2021	Chee et al.	2022/0267844	A1	8/2022	Ramachandran Iyer et al.
2021/0010070	A1	1/2021	Schnall-Levin et al.	2022/0282329	A1	9/2022	Chell et al.
2021/0017583	A1	1/2021	Chee et al.	2022/0290217	A1	9/2022	Frenz et al.
2021/0017586	A1	1/2021	Chee	2022/0290219	A1	9/2022	Chee
2021/0062249	A1	3/2021	Chee	2022/0298560	A1	9/2022	Frisen et al.
2021/0123095	A1	4/2021	Chee	2022/0325325	A1	10/2022	Chee et al.
2021/0130883	A1	5/2021	Chee et al.	2022/0326251	A1	10/2022	Uytingco et al.
2021/0130884	A1	5/2021	Chee et al.	2022/0333171	A1	10/2022	Chee
2021/0140982	A1	5/2021	Uytingco	2022/0333191	A1	10/2022	Mikkelsen et al.
2021/0172007	A1	6/2021	Chee et al.	2022/0333192	A1	10/2022	Uytingco
2021/0189475	A1	6/2021	Tentori et al.	2022/0333195	A1	10/2022	Schnall-Levin et al.
2021/0190770	A1	6/2021	Delaney et al.	2022/0334031	A1	10/2022	Delaney et al.
2021/0198741	A1	7/2021	Williams	2022/0348905	A1	11/2022	Dadhwal
2021/0199660	A1	7/2021	Williams et al.	2022/0348992	A1	11/2022	Stoeckius et al.
2021/0207202	A1	7/2021	Chee	2022/0356464	A1	11/2022	Kim et al.
2021/0214785	A1	7/2021	Stoeckius	2022/0364163	A1	11/2022	Stahl et al.
2021/0222235	A1	7/2021	Chee	2022/0389491	A1	12/2022	Chee
2021/0222241	A1	7/2021	Bharadwaj	2022/0389503	A1	12/2022	Mikkelsen et al.
2021/0222242	A1	7/2021	Ramachandran Iyer	2022/0389504	A1	12/2022	Chew et al.
2021/0222253	A1	7/2021	Uytingco	2022/0403455	A1	12/2022	Ramachandran Iyer et al.
				2022/0404245	A1	12/2022	Chell et al.
				2023/0002812	A1	1/2023	Stoeckius et al.
				2023/0014008	A1	1/2023	Shastri
				2023/0033960	A1	2/2023	Gallant et al.

(56) References Cited			JP	2007-074967	3/2007
U.S. PATENT DOCUMENTS			JP	2009-036694	2/2009
			JP	2011-182702	9/2011
			JP	2013-544498	12/2013
			JP	2014-217381	11/2014
2023/0034039	A1	2/2023	KR	10-2004-0019276	3/2004
2023/0034216	A1	2/2023	KR	10-2009-0000812	1/2009
2023/0040363	A1	2/2023	KR	10-2009-0081260	7/2009
2023/0042088	A1	2/2023	RU	2145635	2/2000
2023/0042817	A1	2/2023	RU	2270254	2/2006
2023/0047782	A1	2/2023	RU	2410439	1/2011
2023/0056549	A1	2/2023	WO	WO 1989/010977	11/1989
2023/0064372	A1	3/2023	WO	WO 1991/006678	5/1991
2023/0069046	A1	3/2023	WO	WO 1993/004199	3/1993
2023/0077364	A1	3/2023	WO	WO 1995/023875	9/1995
2023/0080543	A1	3/2023	WO	WO 1995/025116	9/1995
2023/0081381	A1	3/2023	WO	WO 1995/035505	12/1995
2023/0100497	A1	3/2023	WO	WO 1997/031256	8/1997
2023/0107023	A1	4/2023	WO	WO 1998/044151	10/1998
2023/0111225	A1	4/2023	WO	WO 1999/032654	7/1999
2023/0113230	A1	4/2023	WO	WO 1999/044062	9/1999
2023/0126825	A1	4/2023	WO	WO 1999/044063	9/1999
2023/0129552	A1	4/2023	WO	WO 1999/063385	12/1999
2023/0135010	A1	5/2023	WO	WO 1999/067641	12/1999
2023/0143569	A1	5/2023	WO	WO 2000/017390	3/2000
2023/0145575	A1	5/2023	WO	WO 2000/018957	4/2000
2023/0147726	A1	5/2023	WO	WO 2000/024940	5/2000
2023/0151412	A1	5/2023	WO	WO 2000/063437	10/2000
2023/0159994	A1	5/2023	WO	WO 2001/006012	1/2001
2023/0159995	A1	5/2023	WO	WO 2001/007915	2/2001
2023/0160008	A1	5/2023	WO	WO 2001/009363	2/2001
2023/0175045	A1	6/2023	WO	WO 2001/012862	2/2001
2023/0183785	A1	6/2023	WO	WO 2001/042796	6/2001
2023/0194469	A1	6/2023	WO	WO 2001/046402	6/2001
2023/0194470	A1	6/2023	WO	WO 2001/059161	8/2001
2023/0203478	A1	6/2023	WO	WO 2001/090415	11/2001
2023/0183684	A1	7/2023	WO	WO 2001/096608	12/2001
2023/0212650	A1	7/2023	WO	WO 2002/024952	3/2002
2023/0212655	A1	7/2023	WO	WO 2002/040874	5/2002
2023/0220368	A1	7/2023	WO	WO 2002/059355	8/2002
2023/0220454	A1	7/2023	WO	WO 2002/059364	8/2002
2023/0220455	A1	7/2023	WO	WO 2002/077283	10/2002
2023/0227811	A1	7/2023	WO	WO 2002/088396	11/2002
2023/0228762	A1	7/2023	WO	WO 2003/002979	1/2003
2023/0242976	A1	8/2023	WO	WO 2003/003810	1/2003
FOREIGN PATENT DOCUMENTS			WO	WO 2003/008538	1/2003
			WO	WO 2003/010176	2/2003
			WO	WO 2003/102233	12/2003
			WO	WO 2003/106973	12/2003
CN	1425133	6/2003	WO	WO 2004/015080	2/2004
CN	1537953	10/2004	WO	WO 2004/018497	3/2004
CN	1680604	10/2005	WO	WO 2004/028955	4/2004
CN	1749752	3/2006	WO	WO 2004/055159	7/2004
CN	1898398	1/2007	WO	WO 2004/067759	8/2004
CN	1981188	6/2007	WO	WO 2004/081225	9/2004
CN	101142325	3/2008	WO	WO 2004/108268	12/2004
CN	101221182	7/2008	WO	WO 2005/007814	1/2005
CN	101522915	9/2009	WO	WO 2005/010145	2/2005
CN	102851369	1/2013	WO	WO 2005/026387	3/2005
DE	102008025656	12/2009	WO	WO 2005/042759	5/2005
EP	0901631	3/1999	WO	WO 2005/065814	7/2005
EP	0961110	12/1999	WO	WO 2005/067648	7/2005
EP	1712623	10/2006	WO	WO 2005/113804	12/2005
EP	1782737	5/2007	WO	WO 2006/020515	2/2006
EP	1878502	1/2008	WO	WO 2006/056861	6/2006
EP	1910562	4/2008	WO	WO 2006/064199	6/2006
EP	1923471	5/2008	WO	WO 2006/065597	6/2006
EP	1929039	6/2008	WO	WO 2006/074351	7/2006
EP	1966393	9/2008	WO	WO 2006/084130	8/2006
EP	2002017	12/2008	WO	WO 2006/117541	11/2006
EP	2292788	3/2011	WO	WO 2006/124771	11/2006
EP	2302070	3/2011	WO	WO 2006/137733	12/2006
EP	2350648	8/2011	WO	WO 2007/000669	1/2007
EP	2363504	9/2011	WO	WO 2007/010251	1/2007
EP	2580351	4/2013	WO	WO 2007/030373	3/2007
EP	2789696	10/2014	WO	WO 2007/037678	4/2007
EP	2881465	6/2015	WO	WO 2007/041689	4/2007
EP	2963127	1/2016	WO	WO 2007/060599	5/2007
EP	3045544	7/2016	WO	WO 2007/073165	6/2007
EP	3239304	11/2017			
JP	2007-014297	1/2007			

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2007/073171	6/2007	WO	WO 2020/123305	6/2020
WO	WO 2007/073271	6/2007	WO	WO 2020/123309	6/2020
WO	WO 2007/076128	7/2007	WO	WO 2020/123311	6/2020
WO	WO 2007/076726	7/2007	WO	WO 2020/123316	6/2020
WO	WO 2007/114693	10/2007	WO	WO 2020/123317	6/2020
WO	WO 2007/120241	10/2007	WO	WO 2020/123318	6/2020
WO	WO 2007/123744	11/2007	WO	WO 2020/123319	6/2020
WO	WO 2007/139766	12/2007	WO	WO 2020/167862	8/2020
WO	WO 2007/145612	12/2007	WO	WO 2020/176882	9/2020
WO	WO 2008/022332	2/2008	WO	WO 2020/190509	9/2020
WO	WO 2008/069906	6/2008	WO	WO 2020/198071	10/2020
WO	WO 2008/093098	8/2008	WO	WO 2020/219901	10/2020
WO	WO 2008/157801	12/2008	WO	WO 2021/091611	5/2021
WO	WO 2009/032167	3/2009	WO	WO 2021/092433	5/2021
WO	WO 2009/036525	3/2009	WO	WO 2021/097255	5/2021
WO	WO 2009/152928	12/2009	WO	WO 2021/133842	7/2021
WO	WO 2009/156725	12/2009	WO	WO 2021/133849	7/2021
WO	WO 2010/019826	2/2010	WO	WO 2021/142233	7/2021
WO	WO 2010/027870	3/2010	WO	WO 2021/168261	8/2021
WO	WO 2010/053587	5/2010	WO	WO 2021/168278	8/2021
WO	WO 2010/060439	6/2010	WO	WO 2021/207610	10/2021
WO	WO 2010/126614	11/2010	WO	WO 2021/216708	10/2021
WO	WO 2010/127186	11/2010	WO	WO 2021/225900	11/2021
WO	WO 2011/008502	1/2011	WO	WO 2021/236625	11/2021
WO	WO 2011/014879	2/2011	WO	WO 2021/236929	11/2021
WO	WO 2011/062933	5/2011	WO	WO 2021/237056	11/2021
WO	WO 2011/071943	6/2011	WO	WO 2021/237087	11/2021
WO	WO 2011/094669	8/2011	WO	WO 2021/242834	12/2021
WO	WO 2011/102903	8/2011	WO	WO 2021/247543	12/2021
WO	WO 2011/127006	10/2011	WO	WO 2021/247568	12/2021
WO	WO 2011/127099	10/2011	WO	WO 2021/252499	12/2021
WO	WO 2011/143583	11/2011	WO	WO 2021/252576	12/2021
WO	WO 2011/155833	12/2011	WO	WO 2021/252591	12/2021
WO	WO 2012/022975	2/2012	WO	WO 2021/263111	12/2021
WO	WO 2012/048341	4/2012	WO	WO 2022/025965	2/2022
WO	WO 2012/049316	4/2012	WO	WO 2022/060798	3/2022
WO	WO 2012/058096	5/2012	WO	WO 2022/060953	3/2022
WO	WO 2012/061832	5/2012	WO	WO 2022/087273	4/2022
WO	WO 2012/071428	5/2012	WO	WO 2022/098810	5/2022
WO	WO 2012/083225	6/2012	WO	WO 2022/099037	5/2022
WO	WO 2012/129242	9/2012	WO	WO 2022/103712	5/2022
WO	WO 2012/139110	10/2012	WO	WO 2022/109181	5/2022
WO	WO 2012/140224	10/2012	WO	WO 2022/140028	6/2022
WO	WO 2012/142213	10/2012	WO	WO 2022/147005	7/2022
WO	WO 2012/148477	11/2012	WO	WO 2022/147296	7/2022
WO	WO 2012/159089	11/2012	WO	WO 2022/164615	8/2022
WO	WO 2012/168003	12/2012	WO	WO 2022/178267	8/2022
WO	WO 2013/033271	3/2013	WO	WO 2022/198068	9/2022
WO	WO 2013/090390	6/2013	WO	WO 2022/221425	10/2022
WO	WO 2013/123442	8/2013	WO	WO 2022/226057	10/2022
WO	WO 2013/131962	9/2013	WO	WO 2022/236054	11/2022
WO	WO 2013/138510	9/2013	WO	WO 2022/256503	12/2022
WO	WO 2013/142389	9/2013	WO	WO 2022/271820	12/2022
WO	WO 2013/150082	10/2013	WO	WO 2023/287765	1/2023
WO	WO 2013/150083	10/2013	WO	WO 2023/018799	2/2023
WO	WO 2013/155119	10/2013	WO	WO 2023/034489	3/2023
WO	WO 2013/158936	10/2013	WO	WO 2023/076345	5/2023
WO	WO 2014/044724	3/2014	WO	WO 2023/086880	5/2023
WO	WO 2014/060483	4/2014	WO	WO 2023/102118	6/2023
WO	WO 2014/085725	6/2014			
WO	WO 2014/128129	8/2014			
WO	WO 2014/130576	8/2014			
WO	WO 2014/142841	9/2014			
WO	WO 2014/144713	9/2014			
WO	WO 2014/152397	9/2014			
WO	WO 2014/163886	10/2014			
WO	WO 2014/210223	12/2014			
WO	WO 2014/210225	12/2014			
WO	WO 2014/210353	12/2014			
WO	WO 2016/007839	1/2016			
WO	WO 2016/162309	10/2016			
WO	WO 2017/013170	1/2017			
WO	WO 2018/045181	3/2018			
WO	WO 2018/045186	3/2018			
WO	WO 2020/076979	4/2020			

OTHER PUBLICATIONS

U.S. Appl. No. 17/707,189, filed Mar. 29, 2022, Chell et al.
U.S. Appl. No. 60/416,118, filed Oct. 3, 2002, Fan et al.
U.S. Appl. No. 61/839,313, filed Jun. 25, 2013, Chee et al.
U.S. Appl. No. 61/839,320, filed Jun. 25, 2013, Chee et al.
U.S. Appl. No. 61/902,105, filed Nov. 8, 2013, Kozlov et al.
U.S. Appl. No. 62/839,575, filed Apr. 26, 2019, Bent et al.
[No Author Listed], "10x Genomics Begins Shipments of Visium Spatial Gene Expression Solution," 10x Genomics, Nov. 26, 2019, 2 pages.
[No Author Listed], "FY 2020 United States Patent and Trademark Office Performance and Accountability Report," USPTO, 2021, 28 pages.
[No Author Listed], "HuSNP Mapping Assay User's Manual," Affymetrix Part No. 90094 (Affymetrix, Santa Clara, Calif.), GeneChip, 2000, 104 pages.

(56)

References Cited

OTHER PUBLICATIONS

- [No Author Listed], "Microarray technologies have excellent possibilities in genomics-related researches," *Science Tools From Amersham Pharmacia Biotech*, 1998, 3(4): 8 pages (with English Translation). 10xGenomics.com, [online], "Visium Spatial Gene Expression Reagent Kits—User Guide," Jun. 2020, retrieved on May 25, 2021, retrieved from URL <https://assets.ctfassets.net/an68im79xiti/3GGHfH3RWpd1bFVhlpexR/8baa08d9007157592b65b2cdc7130990/CG000239_VisiumSpatialGeneExpression_UserGuide_RevD.pdf>, 70 pages.
- Adessi et al., "Solid phase DNA amplification: characterisation of primer attachment and amplification mechanisms," *Nucl. Acids Res.*, Oct. 2000, 28(20):E87, 8 pages.
- Affymetrix, "GeneChip Human Genome U133 Set," retrieved from the Internet: on the World Wide Web at affymetrix.com/support/technical/datasheets/hgu133_datasheet.pdf, retrieved on Feb. 26, 2003.
- Affymetrix, "Human Genome U95Av2," Internet Citation, retrieved from the internet: on the World Wide Web affymetrix.com, retrieved on Oct. 2, 2002.
- Agasti et al., "Photocleavable DNA Barcode-Antibody Conjugates Allow Sensitive and Multiplexed Protein Analysis in Single Cells," *Journal of the American Chemical Society*, Oct. 23, 2012, 134(45):18499-18502.
- Agbavwe et al., "Efficiency, error and yield in light-directed maskless synthesis of DNA microarrays," *Journal of Nanobiotechnology*, Dec. 2011, 9:57, 17 pages.
- Ahern et al., "Biochemical, Reagents Kits Offer Scientists Good Return on Investment," *The Scientist*, 1995, 9(15):20, 7 pages.
- Ahern, "Biochemical, Reagents Kits Offer Scientists Good Return on Investment," *The Scientist*, Jul. 1995, 9(15):20, 7 pages.
- Ahlfen et al., "Determinants of RNA quality from FFPE samples," *PLoS One*, Dec. 2007, 2(12):e1261, 7 pages.
- Akeroyd, "Click chemistry for the preparation of advanced macromolecular architectures," Stellenbosch University, PhD Dissertation, Mar. 2010, 138 pages.
- Albretsen et al., "Applications of magnetic beads with covalently attached oligonucleotides in hybridization: Isolation and detection of specific measles virus mRNA from a crude cell lysate," *Anal. Biochem.* 189: 40-50, 1990.
- Albretsen et al., "Optimal conditions for hybridization with oligonucleotides: a study with myc-oncogene DNA probes," *Anal Biochem.*, Apr. 1988, 170(1):193-202.
- Allawi and SantaLucia, "Thermodynamics and NMR of Internal GAT Mismatches in DNA," *Biochemistry*, 1996, 36:10581-10594.
- Altaras et al., "Production and formulation of adenovirus vectors," *Adv Biochem Eng Biotechnol.*, Nov. 2005, 99:193-260.
- Altschul et al., "Basic local alignment search tool," *J. Mol. Biol.*, Oct. 5, 1990, 215(3):403-410.
- Anderson et al., "Microarrayed Compound Screening to Identify Activators and Inhibitors of AMP-Activated Protein Kinase," *J. of Biomolecular Screening*, 2004, 9:112.
- Andersson et al., "Analysis of protein expression in cell microarrays: a tool for antibody-based proteomics," *J Histochem Cytochem.*, 4(12): 1413-1423, 2006.
- Andresen et al., "Deciphering the Antibodyome—Peptide Arrays for Serum Antibody Biomarker Diagnostics," *Current Proteomics*, 6(1), 1-12, 2009.
- Andresen et al., "Helicase-dependent amplification: use in OnChip amplification and potential for point-of-care diagnostics," *Expert Rev Mol Diagn.*, Oct. 2009, 9(7):645-650.
- Angenendt et al., "Cell-free Protein expression and functional assay in a nanowell chip format," *Analytical Chemistry*, 2004, 76(7):1844-49.
- Angenendt et al., "Generation of High Density Protein Microarrays by Cell-free in Situ Expression of Unpurified PCR Products," *Molecular and Cellular Proteomics*, (2006) Ch. 5.9, pp. 1658-1666.
- Appella, "Non-natural nucleic acids for synthetic biology," *Current Opinion in Chemical Biology*, Dec. 2009, 13(5-6): 687-696.
- Armani et al., "2D-PCR: a method of mapping DNA in tissue sections," *Lab on a Chip*, 2009, 9(24):3526-34.
- Asp et al., "Spatial detection of fetal marker genes expressed at low level in adult human heart tissue," *Scientific Reports*, 7: 12941, 10 pages, 2017.
- Atkinson and Wells, "An Updated Protocol for High Throughput Plant Tissue Sectioning," *Front Plant Sci*, 8: 1721, 2017.
- Atkinson, Overview of Translation: Lecture Manuscript, U of Texas (2000) DD. 6.1-6.8.
- Azioune et al., "Simple and rapid process for single cell micro-patterning," *Lab Chip*, Jun. 2009, 9(11):1640-1642.
- Bains et al., "A Novel Method for Nucleic Acid Sequence Determination," *Journal of Theoretical Biology*, 1988, 135(3), 303-7.
- Baird et al., "Rapid SNP Discovery and Genetic Mapping Using Sequenced RAD markers," *PLOS One*, 2008, 3(10):e3376.
- Baner et al., "Signal amplification of padlock probes by rolling circle replication," *Nucleic Acids Res.*, 1998, 26(22):5073-5078.
- Barbie et al., "Systematic RNA interference reveals that oncogenic KRAS-driven cancers require TBK1," *Nature*, Nov. 2009, 462(7269):108-12.
- Barnes, "PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates.," *Proc. Natl. Acad. Sci USA*, 91: 2216-2220, 1994.
- Baugh et al., "Quantitative analysis of mRNA amplification by in vitro transcription," *Nucleic Acids Res.*, 2001, 29:5:e29.
- Beattie et al., "Advances in genosensor research," *Clin Chem.*, May 1995, 41(5):700-6.
- Beier et al., "Versatile derivatisation of solid support media for covalent bonding on DNA-microchips," *Nucleic Acids Res.*, May 1999, 27(9):1970-7.
- Bell, "A Simple Way to Treat PCR Products Prior to Sequencing Using ExoSAP-IT," *Biotechniques*, 2008, vol. 44, No. 6.
- Bentley et al., "Accurate whole human genome sequencing using reversible terminator chemistry," *Nature*, 2008, 456:53-59.
- Berger et al., "Universal bases for hybridization, replication and chain termination," *Nucleic Acid Res.*, Aug. 2000, 28(15):2911-2914.
- Bessmertnykh et al., "Efficient Palladium-Catalyzed Synthesis of Aminopyridyl Phosphonates from Bromopyridines and Diethyl Phosphite," *Synthesis*, 2008, 10:1575-1579.
- Bielas et al., "Human cancers express a mutator phenotype," *Proc. Natl. Acad. Sci. USA*, 2006, 103(48): 18238-18242.
- Bielas et al., "Quantification of random genomic mutations," *Nat. Methods*, 2005, 2(4):285-290.
- biol.www.edu [online], "Principles of Di-Base Sequencing and the Advantages of Color Space Analysis in the SOLiD System," 2008, retrieved on Mar. 11, 2022, retrieved from URL <<http://biol.wvu.edu/young/470/stuff/abi-solid.pdf>>, 4 pages.
- Birney, et al., "Identification and analysis of functional elements in 1% of the human genome by the Encode pilot project," *Nature*, 2007, 447:799-816.
- Blair et al., "Microarray temperature optimization using hybridization kinetics," *Methods Mol Biol.*, 2009, 529:171-96.
- Blanchard et al., "High-density oligonucleotide arrays," *Biosensors & Bioelectronics*, 1996, 11(6-7):687-690.
- Blanco et al., "A practical approach to FRET-based PNA fluorescence in situ hybridization," *Methods*, Dec. 2010, 52(4):343-51.
- Blandini et al., "Animal models of Parkinson's disease," *FEBS J.*, Apr. 2012, 279(7):1156-66.
- Blokzijl et al., "Profiling protein expression and interactions: proximity ligation as a tool for personalized medicine," *J Intern. Med.*, 2010, 268:232-245.
- Blow, "Tissue Issues," *Nature*, 448(23), 959-962, 2007.
- Boeke et al., "Transcription and reverse transcription of retrotransposons," *Annu Rev Microbiol*, 1989, 43:403-34.
- Bonfield et al., "The application of numerical estimates of base calling accuracy to DNA sequencing projects," *Nucleic Acids Research*, 1995, 23(8):1406-1410.
- Bootman et al., "Loading fluorescent Ca²⁺ indicators into living cells," *Cold Spring Harb Protoc.*, Feb. 2013, 2013(2):122-5.
- Bos et al., "In Vitro Evaluation of DNA-DNA Hybridization as a Two-Step Approach in Radioimmunotherapy of Cancer," *Cancer Res.*, Jul. 1, 1994, 54(13):3479-3486.

(56)

References Cited

OTHER PUBLICATIONS

- Boulé et al., "Terminal deoxynucleotidyl transferase indiscriminately incorporates ribonucleotides and deoxyribonucleotides," *J Biol Chem.*, Aug. 2001, 276(33):31388-93.
- Boulgakov et al., "From Space to Sequence and Back Again: Iterative DNA Proximity Ligation and its Applications to DNA-Based Imaging", bioRxiv, 24 pages, 2018.
- Boutros et al., "The art and design of genetic screens: RNA interference," *Nat Rev Genet.*, Jul. 2008, 9(7):554-66.
- Bowtell, "The genesis and evolution of high-grade serous ovarian cancer," *Nat. Rev. Cancer*, 2010, (11):803-808 Abstract.
- Brandon et al., "Mitochondrial mutations in cancer," *Oncogene*, 2006, 25(34):4647-4662.
- Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: physical separation of differentially expressed cDNAs," *Proc. Natl. Acad. Sci. USA*, 2000, 97, 1665-1670.
- Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays", *Nat. Biotech.* 18: 630-634, 2000.
- Brockman et al., "Quality scores and SNP detection in sequencing-by-synthesis systems," *Methods*, 2008, 18:763-770.
- Brow, "35—The Cleavase I enzyme for mutation and polymorphism scanning," *PCR Applications Protocols for Functional Genomics*, 1999, pp. 537-550.
- Brown et al., "Retroviral integration: structure of the initial covalent product and its precursor, and a role for the viral IN protein," *Proc Natl Acad Sci USA*, Apr. 1989, 86(8):2525-9.
- Bullard et al., "Direct comparison of nick-joining activity of the nucleic acid ligases from bacteriophage T4," *Biochem. J.* 2006, 398:135-144.
- Burgess, "A space for transcriptomics", *Nature Reviews Genetics* | Published online Jul. 15, 2016; doi:10.1038/nrg.2016.94.
- Burgess, "Finding structure in gene expression", *Nature Reviews Genetics* | Published online Apr. 13, 2018; doi:10.1038/nrg.2018.19.
- Burns et al., "Well-less, gel-permeation formats for ultra-HTS," *DDT*, 2001, 6(12):S40-S47.
- Burton et al., "Coverslip Mounted-Immersion Cycled in Situ RT-PCR for the Localization of mRNA in Tissue Sections," *Biotechniques*, 1998, 24; pp. 92-100.
- Calvert, "Materials science. Printing cells," *Science*, Oct. 2007, 318(5848):208-209.
- Cardona et al., "TrakEM2 0.9a User Manual," Sep. 8, 2011, retrieved on Jul. 29, 2022, retrieved from URL <https://www.ini.uzh.ch/~acardona/trakem2_manual.html>, 38 pages.
- Carlson et al., "Function and Structure of a Prokaryotic Formylglycine-generating Enzyme," *J. of Biological Chemistry*, 2008, 283(29):20117-125.
- Carter et al., "Stabilization of an optical microscope to 0.1 nm in three dimensions," *Applied Optics*, 2007, 46:421-427.
- Cerritelli et al., "Ribonuclease H: the enzymes in eukaryotes," *FEBS Journal*, Mar. 2009, 276(6):1494-505.
- Cerutti et al., "Generation of sequence-specific, high affinity anti-DNA antibodies," *Journal of Biological Chemistry*, 2001, 276(16):12769-12773.
- Cha et al., "Specificity, Efficiency and Fidelity of PCR," *Genome Res.*, 1993, 3:518-29.
- Chandra et al., "Cell-free synthesis-based protein microarrays and their applications," *Proteomics*, 2009, 5(6):717-30.
- Chatterjee et al., "Protein Microarray On-Demand: A Novel Protein Microarray System," *PLoS One*, 2008, 3(9):e3265.
- Chatterjee, et al., "Mitochondrial DNA mutations in human cancer. *Oncogene*," 2006, 25(34):4663-4674.
- Chen et al., "A Homogeneous, Ligase-mediated DNA diagnostic test," *Genome research*, 1998, 8(5):549-556.
- Chen et al., "DNA hybridization detection in a microfluidic Channel using two fluorescently labelled nucleic acid probes," *Biosensors and Bioelectronics*, 2008, 23:1878-1882.
- Chen et al., "Expansion Microscopy," 2015, *Science*. 347(6221):543-548.
- Chen et al., "Geometric control of cell life and death," *Science*, May 1997, 276(5317):1425-1428.
- Chen et al., "Gray-scale photolithography using microfluidic photomasks," *PNAS*, Feb. 2003, 100(4):1499-1504.
- Chen et al., "Parallel single nucleotide polymorphism genotyping by surface invasive cleavage with universal detection," *Anal Chem.*, Apr. 2005, 77(8):2400-5.
- Chen et al., "Spatially resolved, highly multiplexed RNA profiling in single cells", *Science* 348(6233), 36 pages, 2015.
- Cheng et al., "Sensitive Detection of Small Molecules by Competitive Immunomagnetic-Proximity Ligation Assay," *Anal Chem*, 2012, 84:2129-2132.
- Cheng, "The Contrast Formation in Optical Microscopy," *Handbook of Biological Confocal Microscopy*, 2006, Chapter 8, pp. 162-206.
- Chester et al., "Dimethyl sulfoxide-mediated primer Tm reduction: a method for analyzing the role of renaturation temperature in the polymerase chain reaction," *Anal Biochem*, Mar. 1993, 209(2):284-90.
- Chial, "DNA Sequencing Technologies Key to the Human Genome Project," *Nature Education*, 2008, 1(1):219, 7 pages.
- Chiang et al., "NFkappaB translocation in human microvessel endothelial cells using a four-compartment subcellular protein redistribution assay," *J Biochem Biophys Methods*, Nov. 2000, 46(1-2):53-68.
- Chrisey et al., "Covalent attachment of synthetic DNA to self-assembled monolayer films," *Nucleic Acids Res.*, Aug. 1996, 24(15):3031-9.
- Chung et al., "Imaging single-cell signaling dynamics with a deterministic high-density single-cell trap array," *Anal Chem*, Sep. 2011, 83(18):7044-7052.
- Chung et al., "Structural and molecular interrogation of intact biological systems," *Nature*, May 16, 2013, 497:332-337.
- Ciaccio et al., "Systems analysis of EGF receptor signaling dynamics with microwestern arrays," *Nat Methods*, Feb. 2010, 7(2):148-55.
- Cockroft et al., "A single-molecule nanopore device detects DNA polymerase activity with single-nucleotide resolution," *J Am Chem Soc.*, Jan. 2008, 130(3):818-20.
- Colegio et al., "In vitro transposition system for efficient generation of random mutants of *Campylobacter jejuni*," *J Bacteriol.*, Apr. 2001, 183(7):2384-8.
- Condina et al., "A sensitive magnetic bead method for the detection and identification of tyrosine phosphorylation in proteins by MALDI-TOF/TOF MS," *Proteomics*, 2009, 9:3047-3057.
- Constantine et al., "Use of genechip high-density oligonucleotide arrays for gene expression monitoring," *Life Science News, Amersham Life Science*; 11-14, 1998.
- Cook et al., "The effects of secondary structure and O2 on the formation of direct strand breaks upon UV irradiation of 5-bromodeoxyuridine-containing oligonucleotides," *Chem Biol.*, Jul. 1999, 6(7):451-9.
- Copeland et al., "Mitochondrial DNA Alterations in Cancer," *Cancer Invest.*, 2002, 557-569.
- Corees et al., "An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues", *Nat. Methods*. vol. 14(10):959-962, 2017.
- Cornett et al., "MALDI imaging mass spectrometry: molecular snapshots of biochemical systems," *Nature Methods*, 2007, 4(10):828-833.
- Cox et al., "Tissue subcellular fractionation and protein extraction for use in mass-spectrometry-based proteomics," *Nat Protoc.*, 2006, 1(4):1872-8.
- Craig, "Transposon Tn7," *Curr Top Microbiol Immunol.*, 1996, 204:27-48.
- Craig, "V(D)J recombination and transposition: closer than expected," *Science*, Mar. 1996, 271(5255):1512, 1 page.
- Crisalli et al., "Importance of ortho Proton Donors in Catalysis of Hydrazone Formation," *Org. Lett.*, 2013, 15(7):1646-1649.
- Crosetto et al., "Spatially resolved transcriptomics and beyond," *Nature Review Genetics*, 2015, 57-66.

(56)

References Cited

OTHER PUBLICATIONS

- Cujec et al. "Selection of v-abl tyrosine kinase substrate sequences from randomized peptide and cellular proteomic libraries using mRNA display," *Chemistry and Biology*, 2002, 9:253-264.
- Curtis et al., "Adhesion of cells to polystyrene surfaces," *J Cell Biol.*, Nov. 1983, 97(5):1500-1506.
- Czarnik, "Encoding methods for combinatorial chemistry," *Curr Opin Chem Biol.*, Jun. 1997, 1(1):60-6.
- Dahl et al., "Circle-to-circle amplification for precise and sensitive DNA analysis," *Proc. Natl. Acad. Sci.*, 2004, 101:4548-4553.
- Daley et al., "Predicting the molecular complexity of sequencing libraries," *Nature Methods*, Apr. 2013, 10:325-327.
- Dalma-Weiszhausz et al., "The affymetrix GeneChip platform: an overview," *Methods Enzymol.*, 2006, 410:3-28.
- Dandapani et al., "Selecting, Acquiring, and Using Small Molecule Libraries for High-Throughput Screening," *Curr Protoc Chem Biol.*, Sep. 2012, 4:177-191.
- Darmanis, et al., "ProteinSeq: High-Performance Proteomic Analyses by Proximity, Ligation and Next Generation Sequencing," *PLoS One*, 2011, 6(9):e25583.
- Darnell, Jr., "Reflections on the history of pre-mRNA processing and highlights of current knowledge: A unified picture," *RNA*, Feb. 2013, 19:443-460, 19 pages.
- Daubendiek et al., "Rolling-Circle RNA Synthesis: Circular Oligonucleotides as Efficient Substrates for T7 RNA Polymerase," *J. Am. Chem. Soc.*, 1995, 117:77818-7819.
- Davies et al., "How best to identify chromosomal interactions: a comparison of approaches," *Nat. Methods* 14, 125-134, 2017.
- Dawson et al., "Genetic animal models of Parkinson's disease," *Neuron*, Jun. 2010, 66(5):646-661.
- De Clercq, "A 40-year journey in search of selective antiviral chemotherapy," *Annu Rev Pharmacol Toxicol.*, 2011, 51:1-24.
- Deamer et al., "Characterization of nucleic acids by nanopore analysis," *Acc Chem Res.*, Oct. 2002, 35(10):817-25.
- Deamer et al., "Nanopores and nucleic acids: prospects for ultrarapid sequencing," *Trends Biotechnol.*, Apr. 2000, 18(4):147-51.
- Dean et al., "Comprehensive human genome amplification using multiple displacement amplification," *Proc Natl. Acad. Sci. USA* 99:5261-66, 2002.
- Dean et al., "Rapid Amplification of Plasmid and Phage DNA Using Phi29 DNA Polymerase and Multiply-Primed Rolling Circle Amplification," *Genome Research*, Jun. 2001, 11:1095-1099.
- Deibel et al., "Biochemical properties of purified human terminal deoxynucleotidyltransferase," *J Biol Chem.*, May 1980, 255(9):4206-12.
- Deo et al., "Detection of mammalian microRNA expression by in situ hybridization with RNA oligonucleotides," *Dev Dyn.*, Sep. 2006, 235(9):2538-48.
- Devine et al., "Efficient integration of artificial transposons into plasmid targets in vitro: a useful tool for DNA mapping, sequencing and genetic analysis," *Nucleic Acids Res.*, Sep. 1994, 22(18):3765-72.
- Dhindsa et al., "Virtual Electrowetting Channels: Electronic Liquid Transport with Continuous Channel Functionality," *Lab Chip*, 2010, 10:832-836.
- Diez-Roux et al., "A high-resolution anatomical atlas of the transcriptome in the mouse embryo," *PLoS Biol.*, Jan. 2011, 9(1):e1000582, 14 pages.
- Ding et al., "On-chip manipulation of single microparticles, cells, and organisms using surface acoustic waves," *PNAS*, Jul. 2012, 109(28):11105-11109.
- Doddridge et al., "UV-induced strand break damage in single stranded bromodeoxyuridine-containing DNA oligonucleotides," *Chem Commun.*, 1998, p. 1997-1998.
- Dressman et al., "Transforming single DNA molecules into fluorescent magnetic particles for detection and enumeration of genetic variations," *Proc. Natl. Acad. Sci. USA*, 2003, 100:8817-8822.
- Drmanac et al., "Accurate sequencing by hybridization for DNA diagnostics and individual genomics," *Nature Biotechnology*, 16:54-58, 1998.
- Druley et al., "Quantification of rare allelic variants from pooled genomic DNA," *Nat. Methods* 6: 263-65, 2009.
- Duhr et al., "Why molecules move along a temperature gradient," *Proc Natl Acad Sci USA*, Dec. 2006, 103(52):19678-19682.
- Duncan et al., "Affinity chromatography of a sequence-specific DNA binding protein using teflon-linked oligonucleotides," *Anal. Biochem.* 169: 104-108, 1988.
- Dundas et al., "Reference genes for measuring mRNA expression," *Theory Biosci.*, May 17, 2012, 131:215-223.
- Eagen, "Principles of Chromosome Architecture Revealed by Hi-C", *Trends in Biochemical Sciences*, vol. 43, No. 6, 10 pages, 2018.
- Eberwine et al., "Analysis of gene expression in single live neurons," *Proc. Natl. Acad. Sci.*, USA 89, 3010-3014, 1992.
- Eberwine, "Amplification of mRNA Populations Using aRNA Generated from Immobilized Oligo(dT)-T7 Primed cDNA," *BioTechniques* 20 (4), 584, 1996.
- Ebihara et al., "Molecular detection of dermatophytes and nondermatophytes in onychomycosis by nested polymerase chain reaction based on 28S ribosomal RNA gene sequences," *Br J Dermatol.*, Nov. 2009, 161(5):1038-44.
- Eguluz et al., "Multitissue array review: A chronological description of tissue array techniques, applications and procedures," *Pathology Research and Practice*, 2006, 202:561-568.
- Ekins et al., "Microarrays: their origins and applications," *Trends in Biotechnology*, Jun. 1999, 17(6):217-218.
- Eldridge et al. "An in vitro selection strategy for conferring protease resistance to ligand binding peptides," *Protein Eng Des Sel.*, 22(11): 691-698, 2009.
- Ellington et al., "Antibody-based protein multiplex platforms: technical and operational challenges," *Clin Chem* 56(2): 186-193, 2010.
- Emmert-Buck et al., "Laser capture microdissection," *Science*, Nov. 1996, 274(5289):998-1001.
- Eng et al., "Profiling the transcriptome by RNA SPOTs," *Nat Methods.*, 14(12): 1153-1155, 2017.
- Ergin et al., "Proteomic Analysis of PAXgene-Fixed Tissues," *J Proteome Res.*, 2010, 9(10):5188-96.
- Escholarship.org [online], "Methods and devices for fabricating and assembling DNA and protein arrays for high-throughput analyses [electronic resource]," 2010, retrieved on Jun. 8, 2022, retrieved from URL <<https://escholarship.org/uc/item/6tf7p46s>>, 155 pages.
- Espina et al., "Laser-capture microdissection," *Nat Protoc*, 2006, 1(2):586-603.
- Evers et al., "The effect of formaldehyde fixation on RNA: optimization of formaldehyde adduct removal," *J Mol Diagn.*, May 2011, 13(3):282-8.
- Examination Search Report issued in Australian Appln. No. 2012241730, dated Jun. 27, 2016, 3 pages.
- Extended European Search Report in European Appln. No. 18208823. 7, dated May 6, 2019, 12 pages.
- Extended European Search Report in European Appln. No. 21163827. 5, dated Sep. 24, 2021, 12 pages.
- Extended European Search Report issued in European Appln. No. 19204655.5, dated Jun. 8, 2020, 13 pages.
- Fahy et al., "Design and synthesis of polyacrylamide-based oligonucleotide supports for use in nucleic acid diagnostics," *Nucleic Acids Res.*, Apr. 1993, 21(8):1819-26.
- Falconnet et al., "Surface engineering approaches to micropattern surfaces for cell-based assays," *Biomaterials*, Jun. 2006, 27(16):3044-3063.
- Fan et al., "Highly parallel SNP genotyping," *Cold Spring Symp. Quant. Biol.*, 68: 69-78, 2003.
- Fang et al., "Fluoride-cleavable biotinylation phosphoramidite for 5'-end-labeling and affinity purification of synthetic oligonucleotides," *Nucleic Acids Res.*, Jan. 2003, 31(2):708-715.
- Faruqi et al., "High-throughput genotyping of single nucleotide polymorphisms with rolling circle amplification," *BMC Genomics*, Aug. 2001, 2:4, 10 pages.
- Ferreira et al., "Photocrosslinkable Polymers for Biomedical Applications," *Biomedical Engineering—Frontiers and Challenges*, Prof. Reza, 2011, 22 pages.
- Fire and Xu, "Rolling replication of short DNA circles," *Proc. Natl. Acad. Sci.*, 92: 4641-4645, 1995.

(56)

References Cited

OTHER PUBLICATIONS

- Fischer et al., "Hematoxylin and eosin staining of tissue and cell sections," CSH Protoc., May 2008, 3(5):1-3.
- Flanigon et al., "Multiplex protein detection with DNA readout via mass spectrometry," N. Biotechnol., 30(2): 153-158, 2013.
- Fodor et al., "Light-Directed, Spatially Addressable Parallel Chemical Synthesis," Science, 251(4995), 767-773, 1995.
- Folch et al., "Microfabricated elastomeric stencils for micropatterning cell cultures," J Biomed Mater Res, Nov. 2000, 52(2):346-353.
- Forcucci et al., "All-plastic miniature fluorescence microscope for point-of-care readout of bead-based bioassays," J. Biomed Opt. 20 (10): 105010, 15 pages, 2015.
- Fredriksson et al., "Multiplexed protein detection by proximity ligation for cancer detection," Nature Methods, 4(4): 327-29, 2007.
- Fredriksson et al., "Multiplexed proximity ligation assays to profile putative plasma biomarkers relevant to pancreatic and ovarian cancer," Clin. Chem., 5(3): 582-89, 2008.
- Fredriksson et al., "Protein detection using proximity-dependent DNA ligation assays," Nature Biotech., 20: 473-77, 2002.
- Frese et al., "Formylglycine Aldehyde Tag-Protein Engineering through a Novel Posttranslational Modification," ChemBioChem., 10: 425-27, 2009.
- Fu et al., "Counting individual DNA molecules by the stochastic attachment of diverse labels," PNAS, 108: 9026-9031, 2011.
- Fu et al., "Repeat subtraction-mediated sequence capture from a complex genome," Plant J., Jun. 2010, 62(5):898-909.
- Fullwood et al., "Next-generation DNA sequencing of paired-end tags (PET) for transcriptome and genome analyses," Genome Res., 19: 521-532, 2009.
- Galon et al., "The immune score as a new possible approach for the classification of cancer," J Transl Med., Jan. 2012, 10:1, 4 pages.
- Gamper et al., "Gene expression profile of bladder tissue of patients with ulcerative interstitial cystitis," BMC Genomics, Apr. 28, 2009, 10(199):1-17.
- Gans et al., "Inkjet Printing of Polymers: State of the Art and Future Developments," Advanced Materials, Feb. 2004, 16(3):203-213.
- Gao et al., "High density peptide microarrays. In situ synthesis and applications," Molecular Diversity, 8, 177-187, 2004.
- Gao et al., "Q&A: Expansion microscopy," BMC Biology, 15: 50, 9 pages, 2017.
- Geiss et al., "Direct multiplexed measurement of gene expression with color-coded probe pairs," nature biotechnology, 2008, 26(3):317-325.
- Genbank Accession No. AC009495.1, "*Homo sapiens* clone NH0490102, *** Sequencing in Progress *** , 12 unordered pieces," Aug. 24, 1999, 53 pages.
- Genbank Accession No. AC009495.5, "*Homo sapiens* BAC clone RP11-49012 from 2, complete sequence," Apr. 21, 2005, 32 pages.
- Genbank Accession No. AC037198.2, "*Homo sapiens* chromosome 15 clone CTD-2033D15 map 15q14, *** Sequencing in Progress *** , 62 unordered pieces," Apr. 25, 2000, 39 pages.
- Genbank Accession No. AC087379.2, "*Homo sapiens* chromosome 11 clone RP11-396020 map 11, *** Sequencing in Progress *** , 5 ordered pieces," Jul. 6, 2002, 47 pages.
- Genbank Accession No. AC087741.1, "*Homo sapiens* chromosome 17 clone RP11-334C17 map 17, Low-Pass Sequence Sampling," Jan. 22, 2001, 18 pages.
- Genbank Accession No. AC100826.1, "*Homo sapiens* chromosome 15 clone RP11-279F6 map 15, Low-Pass Sequence Sampling," Nov. 22, 2001, 21 pages.
- Genbank Accession No. AL445433.14, "Human DNA sequence from clone RP11-234N17 on chromosome 1, complete sequence," Jan. 24, 2013, 32 pages.
- Genbank Accession No. AL445524.1, "*Homo sapiens* chromosome 1 clone RP11-295G20, Working Draft Sequence, 19 unordered pieces," Oct. 14, 2000, 47 pages.
- Genome.ucsc.edu, [online], "Genome Browser Gateway," 2000, retrieved on Jun. 11, 2021, retrieved from URL<<https://genome.ucsc.edu/cgi-bin/hgGateway>>, 3 pages.
- Genomeweb.com, [online], Han, "10x Genomics Describes Capabilities of Visium Spatial Genomics Product Ahead of Release," Oct. 18, 2019, retrieved on Mar. 10, 2021, retrieved from URL<<https://www.genomeweb.com/sequencing/10x-genomics-describes-capabilities-visium-spatial-genomics-product-ahead-release#YO7-Dr6SljV>>, 2 pages.
- Gerard et al., "Excess dNTPs minimize RNA hydrolysis during reverse transcription," Biotechniques, Nov. 2002, 33(5):984, 986, 988, 990.
- Gerdtsen et al., "Evaluation of Solid Supports for Slide- and Well-Based Recombinant Antibody Microarrays", Microarrays (2016) 5:16, 2016.
- Giacomello et al., "Spatially resolved transcriptome profiling in model plant species", Nature Plants 3, 17061, 11 pages, 2017.
- Giam et al., "Scanning probe-enabled nanocombinatorics define the relationship between fibronectin feature size and stem cell fate," PNAS, Mar. 2012, 109(12):4377-4382.
- Gibson et al., "Enzymatic assembly of DNA molecules up to several hundred kilobases," Nat Methods., May 2009, 6(5):343-5.
- Gibson-Corley et al., "Principles for valid histopathologic scoring in research," Vet Pathol., Nov. 2013, 50(6):1007-15.
- Gilar et al., "Study of phosphorothioate-modified oligonucleotide resistance to 3'-exonuclease using capillary electrophoresis," J Chromatogr B Biomed Sci Appl., Aug. 28, 1998, 714(1):13-20.
- Gill et al., "Nucleic acid isothermal amplification technologies: a review," Nucleosides Nucleotides Nucleic Acids, Mar. 2008, 27(3):224-43.
- Gilles et al., "Single nucleotide polymorphic discrimination by an electronic dot blot assay on semiconductor microchips," Nat Biotechnol., Apr. 1999, 17(4):365-70.
- Glass et al., "SIMPLE: a sequential immunoperoxidase labeling and erasing method," J. Histochem. Cytochem., Oct. 2009, 57(10):899-905.
- Gloor, "Gene targeting in *Drosophila*," Methods Mol Biol., 2004, 260:97-114.
- Gnanaprasagam, "Unlocking the molecular archive: the emerging use of formalin-fixed paraffin-embedded tissue for biomarker research in urological cancer," BJU International, 105, 274-278, 2009.
- Goebel et al., "Development of a sensitive and specific in situ hybridization technique for the cellular localization of antisense oligodeoxynucleotide drugs in tissue sections," Toxicologic Pathology, Jun. 2007, 35(4):541-548.
- Goldkom and Prockop, "A simple and efficient enzymatic method for covalent attachment of DNA to cellulose. Application for hybridization-restriction analysis and for in vitro synthesis of DNA probes," Nucleic Acids Res. 14:9171-9191, 1986.
- Goldmeyer et al., "Development of a novel one-tube isothermal reverse transcription thermophilic helicase-dependent amplification platform for rapid RNA detection," Journal of Molecular Diagnostics, American Society for Investigative Pathology and the Association for Molecular Pathology, Nov. 1, 2007, 9(5):639-644.
- Goransson et al., "A single molecule array for digital targeted molecular analyses," Nucleic Acids Res., Nov. 25, 2009, 37(1):e7, 9 pages.
- Goryshin et al., "Tn5 in vitro transposition," J Biol Chem., Mar. 1998, 273(13):7367-74.
- Gotz et al., "Animal models of Alzheimer's disease and frontotemporal dementia," Nat Rev Neurosci., Jul. 2008, 9(7):532-44.
- Grant et al., "Pathways and mechanisms of endocytic recycling," Nat. Rev. Mol. Cell Biol., Sep. 2009, 10(9):597-608.
- Grokhovskiy, "Specificity of DNA cleavage by ultrasound," Molecular Biology, 2006, 40(2):276-283.
- Grünweller et al., "Locked Nucleic Acid Oligonucleotides," BioDrugs, Jul. 2007, 21(4): 235-243.
- Gu et al., "Protein tag-mediated conjugation of oligonucleotides to recombinant affinity binders for proximity ligation," N Biotechnol., 30(2): 144-152, 2013.
- Gudjonsson et al., "Myoepithelial cells: their origin and function in breast morphogenesis and neoplasia," J Mammary Gland Biol Neoplasia, Jul. 2005, 10(3):261-72.
- Gunderson et al., "Decoding Randomly Ordered DNA Arrays," Genome Research 14: 870-877, 2004.

(56)

References Cited

OTHER PUBLICATIONS

- Guo et al., "Direct fluorescence analysis of genetic polymorphisms by hybridization with oligonucleotide arrays on glass supports," *Nucleic Acids Res.*, Dec. 1994, 22(24):5456-65.
- Hafner et al., "Identification of microRNAs and other small regulatory RNAs using cDNA library sequencing," *Methods*, Jan. 2008, 44(1):3-12.
- Hahnke et al., "Strip-tease on glass: validation of an improved stripping procedure for in situ microarrays," *J Biotechnol.*, Jan. 2007, 128(1):1-13.
- Hajduk et al., "Drug discovery: A question of library design," *Nature*, Feb. 2011, 470(7332):42-43.
- Halova et al., "Mast cell chemotaxis—chemoattractants and signaling pathways," *Front Immunol.*, May 2012, 3:119, 20 pages.
- Hamaguchi et al., "Direct reverse transcription-PCR on oligo(dT)-immobilized polypropylene microplates after capturing total mRNA from crude cell lysates," *Clin Chem.*, Nov. 1998, 44(11):2256-63.
- Hammond et al., "Profiling cellular protein complexes by proximity ligation with dual tag microarray readout," *PLoS ONE*, 2012, 7(7):e40405.
- Han et al., "3C and 3C-based techniques: the powerful tools for spatial genome organization deciphering," *Molecular Cytogenetics* (2018) 11:21, 10 pages, 2018.
- Hanauer et al., "Separation of nanoparticles by gel electrophoresis according to size and shape," *Nano Lett.*, Sep. 2007, 7(9):2881-5.
- Hardenbol et al., "Highly multiplexed molecular inversion probe genotyping: over 10,000 targeted SNPs genotyped in a single tube assay," *Genome Res.*, Feb. 2005, 15(2):269-75.
- Hardenbol et al., "Multiplexed genotyping with sequence-tagged molecular inversion probes," *Nature Biotechnol.*, Jun. 2003, 21(6):673-678.
- Harris et al., "Chloroplast ribosomes and protein synthesis," *Microbiol. Mol. Biol. Rev.*, Dec. 1, 1994, 58(4): 700-754.
- Harris et al., "The design and application of target-focused compound libraries," *Comb Chem High Throughput Screen*, Jul. 2011, 14(6):521-531.
- Harrow et al., "GenCode: The reference human genome annotation for the Encode Project," *Genome Res.*, Sep. 2012, 22(9):1760-1774.
- Hattersley et al., "Development of a microfluidic device for the maintenance and interrogation of viable tissue biopsies," *Lab Chip.*, Nov. 2008, 8(11):1842-6.
- Hayes et al., "Electrophoresis of proteins and nucleic acids: I-Theory," *BMJ*, Sep. 1989, 299(6703):843-6.
- He et al., "In situ synthesis of protein arrays," *Current Opinion in Biotechnology*, 19:4-9, 2008.
- He et al., "Printing protein arrays from DNA arrays," *Nature Methods*, 5:175-77, 2008.
- He, "Cell-free protein synthesis: applications in proteomics and biotechnology," *New Biotechnology* 25: 126-132, 2008.
- Healy, "Nanopore-based single-molecule DNA analysis," *Nanomedicine (Lond)*, Aug. 2007, 2(4):459-81.
- Hedskog et al., "Dynamics of HIV-1 Quasispecies during Antiviral Treatment Dissected using Ultra-Deep Pyrosequencing," *PLoS One*, 5(7):e11345, 2010.
- Hein et al., "Click Chemistry, A Powerful Tool for Pharmaceutical Sciences", *Pharm Res.*, 25(10): 2216-2230, 2008.
- Hejatko et al., "In Situ Hybridization Techniques for mRNA Detection in Whole Mount *Arabidopsis* Samples," *Nature Protocols*, 2006, 1(4):1939-1946.
- Hendrickson et al., "High sensitivity multianalyte immunoassay using covalent DNA-labeled antibodies and polymerase chain reaction," *Nucleic Acid Research*, Feb. 11, 1995, 23(3):522-529.
- Hessner et al., "Genotyping of factor V G1691A (Leiden) without the use of PCR by invasive cleavage of oligonucleotide probes," *Clin Chem.*, Aug. 2000, 46(8 Pt 1):1051-6.
- Hiatt et al., "Parallel, tag-directed assembly of locally-derived short sequence reads," *Nature Methods*, 7(2): 119-25, 2010.
- Hlubek et al., "Heterogeneous expression of Wnt/beta-catenin target genes within colorectal cancer," *Int J Cancer.*, Nov. 2007, 121(9):1941-8.
- Ho et al., "Bacteriophage T4 RNA ligase 2 (gp24.1) exemplifies a family of RNA ligases found in all phylogenetic domains," *PNAS*, Oct. 2002, 99(20):12709-14.
- Ho et al., "Characterization of an ATP-Dependent DNA Ligase Encoded by Chlorella Virus PBCV-1," *Journal of Virology*, Mar. 1997, 71(3):1931-1937.
- Hober et al., "Human protein atlas and the use of microarray technologies," *Curr Opin Biotechnol.*, Feb. 2008, 19(1):30-35.
- Holmström et al., "A highly sensitive and fast nonradioactive method for detection of polymerase chain reaction products," *Anal Biochem*, Mar. 1993, 209(2):278-83.
- Hoyer et al., "Electrostatic spraying: a novel technique for preparation of polymer coatings on electrodes," *Anal Chem*, Nov. 1996, 68(21):3840-4.
- Hsuih et al., "Novel, Ligation-Dependent PCR Assay for Detection of Hepatitis C Virus in Serum," *Journal of Clinical Microbiology*, Mar. 1996, 34(3):501-507.
- Hu et al., "High reproducibility using sodium hydroxide-stripped long oligonucleotide DNA microarrays," *Biotechniques*, Jan. 2005, 38(1):121-4.
- Hughes et al., "Microfluidic Western blotting," *PNAS*, Dec. 2012, 109(52):21450-21455.
- Hycultbiotech.com, [online], "Immunohistochemistry, Paraffin" Apr. 2010, retrieved on Apr. 16, 2020, retrieved from URL <https://www.hycultbiotech.com/media/wysiwyg/Protocol_Immunohistochemistry_Paraffin_2.pdf>, 3 pages.
- Hytönen et al., "Design and construction of highly stable, protease-resistant chimeric avidins," *J Biol Chem.*, Mar. 2005, 280(11):10228-33.
- Ichikawa et al., "In vitro transposition of transposon Tn3," *J Biol. Chem.*, Nov. 1990, 265(31):18829-32, Abstract.
- Illumina.com [online], "Array-Based Gene Expression Analysis," 2011, retrieved on Dec. 13, 2021, retrieved from URL <https://www.illumina.com/documents/products/datasheets/datasheet_gene_exp_analysis.pdf>, 5 pages.
- Im et al., "An Introduction to Performing Immunofluorescence Staining," *Biobanking: Methods and Protocols*, Method in Molecular Biology, Yong (ed.), 2019, 1897, Chapter 26, 299-311.
- Imbeaud et al., "Towards standardization of RNA quality assessment using user-independent classifiers of microcapillary electrophoresis traces," *Nucleic Acids Res.*, Mar. 2005, 33(6):e56, 12 pages.
- Inoue and Wittbrodt, "One for All—A Highly Efficient and Versatile Method for Fluorescent Immunostaining in Fish Embryos," *PLoS One* 6, e19713, 2011.
- Invitrogen, Immune Response Biomarker Profiling Service Report, Invitrogen, 2009, 1-33.
- Jabara et al., Accurate sampling and deep sequencing of the HIV-1 protease gene using a Primer ID. *PNAS* 108(50); 20166-20171, 2011.
- Jain, "Transport of molecules, particles, and cells in solid tumors," *Annu. Rev. Biomed. Eng.*, 1999, 1:241-263.
- Jamur and Oliver, "Permeabilization of cell membranes," *Method Mal. Biol.*, 588: 63-66, 2010.
- Jawhar et al., "Tissue Microarray: A rapidly evolving diagnostic and research tool," *Annals of Saudi Medicine*, Mar. 2009, 29(2):123-7.
- Jemt et al., "An automated approach to prepare tissue-derived spatially barcoded RNA-sequencing libraries", *Scientific Reports*, 6: 37137, 10 pages, 2016.
- Jennane et al., "Photolithography of self-assembled monolayers: optimization of protecting groups by an electroanalytical method," *Can. J Chem.*, Dec. 1996, 74(12):2509-2517.
- Jensen et al., "Zinc fixation preserves flow cytometry scatter and fluorescence parameters and allows simultaneous analysis of DNA content and synthesis, and intracellular and surface epitopes," *Cytometry A.*, Aug. 2010, 77(8):798-804.
- Jones et al., Comparative lesion sequencing provides insights into tumor evolution. *Proc. Natl. Acad. Sci. USA* 105(11): 4283-4288, 2008.
- Joos et al., "Covalent attachment of hybridizable oligonucleotides to glass supports," *Anal Biochem.*, Apr. 1997, 247(1):96-101.
- Jucá et al., "Effect of dimethyl sulfoxide on reverse transcriptase activity," *Braz. J. Med. Biol. Res.*, Mar. 1995, 28(3):285-90.

(56)

References Cited

OTHER PUBLICATIONS

- Kainkaryam et al., "Pooling in high-throughput drug screening" *Curr Opin Drug Discov Devel.*, May 2009, 12(3):339-50.
- Kanehisa, "Use of statistical criteria for screening potential homologies in nucleic acid sequences" *Nucleic Acids Res.* 12: 203-213, 1984.
- Kap et al., "Histological Assessment of PAXgene Tissue Fixation and Stabilization Reagents," *PLoS One* 6, e27704, 10 pages, 2011.
- Kapteyn et al., "Incorporation of Non-Natural Nucleotides Into Template-Switching Oligonucleotides Reduces Background and Improves cDNA Synthesis From Very Small RNA Samples," *BMC Genomics*, 2010, 11(413): 1-9.
- Karlin et al., "Applications and statistics for multiple high-scoring segments in molecular sequences," *Proc. Natl. Acad. Sci.*, Jun. 15, 1993, 90:5873-7.
- Kelleher et al., "Characterization of RNA Strand Displacement Synthesis by Moloney Murine Leukemia Virus Reverse Transcriptase," *J Biol Chem*, Apr. 1998, 273(16):9976-86.
- Kibbe, "OligoCalc: an online oligonucleotide properties calculator," *Nucleic Acids Res.*, Jul. 2007, 35:W43-6.
- Kirby et al., "Cryptic plasmids of *Mycobacterium avium*: Tn552 to the rescue," *Mol Microbiol.*, Jan. 2002, 43(1):173-86.
- Kleckner et al., "Tn10 and IS10 transposition and chromosome rearrangements: mechanism and regulation in vivo and in vitro," *Curr Top Microbiol Immunol.*, 1996, 204:49-82.
- Koch et al., "Photochemical immobilization of anthraquinone conjugated oligonucleotides and PCR amplicons on solid surfaces," *Bioconjugate Chem.*, Jul. 2000, 11(4):474-483.
- Kokkat et al., "Archived formalin-fixed paraffin-embedded (FFPE) blocks: A valuable underexploited resource for extraction of DNA, RNA, and protein," *Apr.* 2013, 11(2):101-6.
- Kolb et al., "Click Chemistry: Diverse Chemical Function from a Few Good Reactions," *Angew. Chem. Int. Ed.*, 40(11): 2004-2021, 2001.
- Kolbert et al., "Ribosomal DNA sequencing as a tool for identification of bacterial pathogens," *Curr Opin Microbiol*, Jun. 1999, 2(3):299-305.
- Kolovos et al., "Investigation of the spatial structure and interactions of the genome at sub-kilobasepair resolution using T2C," *Nat. Protoc.* 13, 459-477, 2018.
- König et al., "iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution," *Nat Struct Mol Biol.*, Jul. 2010, 17(7):909-915.
- Korbel et al., "Paired-End Mapping Reveals Extensive Structural Variation in the Human Genome," *Science*, 318(5849): 420-426, 2007.
- Korlach et al., "Selective aluminum passivation for targeted immobilization of single DNA polymerase molecules in zero-mode waveguide nanostructures," *Proc. Natl. Acad. Sci. USA* 105, 1176-1181, 2008.
- Kozlov et al., "A High-Complexity Multiplexed Solution-Phase Assay for Profiling Protease Activity on Microarrays," *Comb Chem High Throughput Screen*, 11: 24-35, 2008.
- Kozlov et al., "A Highly Scalable Peptide-Based Assay System for Proteomics," *PLoS ONE*, 7(6): e37441, 2012.
- Kozlov et al., "A Method for Rapid Protease Substrate Evaluation and Optimization," *Comb Chem High Throughput Screen*, 9: 481-87, 2006.
- Kristensen et al., "High-Throughput Methods for Detection of Genetic Variation," *BioTechniques*, Feb. 2001, 30(2):318-332.
- Kuijpers et al. "Specific recognition of antibody-oligonucleotide conjugates by radiolabeled antisense nucleotides: a novel approach for two-step radioimmunotherapy of cancer," *Bioconjugate Chem.*, Jan. 1, 1993, 4(1):94-102.
- Kumar et al., "Template-directed oligonucleotide strand ligation, covalent intramolecular DNA circularization and catenation using click chemistry," *J Am Chem Soc.*, May 2007, 129(21):6859-64.
- Kurz et al., "cDNA-Protein Fusions: Covalent Protein-Gene Conjugates for the In Vitro Selection of Peptides and Proteins," *ChemBioChem.*, 2: 666-72, 2001.
- Kwok, "High-throughput genotyping assay approaches," *Pharmacogenomics*, Feb. 2000, 1(1):95-100.
- Kwon, "Single-molecule fluorescence in situ hybridization: Quantitative imaging of single RNA molecules," Department of Biomedical Engineering, Oregon Health & Science University, Feb. 2013, 46:65-72.
- LaFerla et al., "Animal models of Alzheimer disease," *Cold Spring Harb Perspect Med.*, Nov. 2012, 2(11):a006320, 14 pages.
- Lage et al., "Whole Genome Analysis of Genetic Alterations in Small DNA Samples Using Hyperbranched Strand Displacement Amplification and Array—CGH," *Genome Research* 13: 294-307, 2003.
- Lampe et al., "A purified mariner transposase is sufficient to mediate transposition in vitro," *EMBO J.*, Oct. 1996, 15(19):5470-9.
- Lamture et al., "Direct detection of nucleic acid hybridization on the surface of a charge coupled device," *Nucleic Acid Res.*, Jun. 1994, 22(11):2121-5.
- Landegren et al., "Reading bits of genetic information: methods for single-nucleotide polymorphism analysis," *Genome Res.*, Aug. 1998, 8(8):769-76.
- Langdale et al., "A rapid method of gene detection using DNA bound to Sephacryl", *Gene* 36: 201-210, 1985.
- Larman et al., "Autoantigen discovery with a synthetic human peptidome," *Nature Biotechnology*, doi:10.1038/nbt.1856, vol. 29, No. 6, pp. 535-541, 2011.
- Larsen et al., "Characterization of a recombinantly expressed proteinase K-like enzyme from a psychrotrophic *Serratia* sp.," *FEBS J.*, Jan. 2006, 273(1):47-60.
- Larsson et al., "In situ detection and genotyping of individual mRNA molecules," *Nat Methods*, May 2010, 7(5):395-7.
- Larsson et al., "In situ genotyping individual DNA molecules by target-primed rolling-circle amplification of padlock probes," *Nat Methods*, Dec. 2004, 1(3):227-32.
- Lassmann et al., A Novel Approach for Reliable Microarray Analysis of Microdissected Tumor Cells From Formalin-Fixed and Paraffin-Embedded Colorectal Cancer Resection Specimens, *J Mol Med*, 87, 211-224, 2009.
- Laurell et al., "Chip integrated strategies for acoustic separation and manipulation of cells and particles," *Chem. Soc. Rev.*, Mar. 2007, 36(3):492-506.
- Le Reste et al., "Characterization of dark quencher chromophores as nonfluorescent acceptors for single-molecule FRET," *Biophysical Journal*, Jun. 2012, 102(11):2658-2668.
- Lee et al., "A novel COL3A1 gene mutation in patient with aortic dissected aneurysm and cervical artery dissections," *Heart Vessels*, Mar. 2008, 23(2):144-8.
- Lee et al., "Cytokines in cancer immunotherapy," *Cancers (Basel)*, Oct. 2011, 3(4):3856-3893.
- Lee et al., "Fluorescent in situ sequencing (FISSEQ) of RNA for gene expression profiling in intact cells and tissues", *Nature Protec*, 10(3): 442-458, 2015.
- Lee et al., "Highly multiplexed subcellular RNA sequencing in situ", *Science*, 343(6177): 1360-1363, 2014.
- Lee et al., "Improving the efficiency of genomic loci capture using oligonucleotide arrays for high throughput resequencing," *BMC Genomics*, Dec. 2009, 10:646, 12 pages.
- Lee et al., "Protein nanoarrays generated by dip-pen nanolithography," *Science*, Mar. 2002, 295(5560):1702-1705.
- Lein et al., "The promise of spatial transcriptomics for neuroscience in the era of molecular cell typing", *Science* 358, 64-69, 2017.
- Lenard, "Viral Membranes," *Encyclopedia of Virology*, Jul. 2008, pp. 308-314.
- Leriche et al., "Cleavable linkers in chemical biology," *Bioorganic & Medicinal Chemistry*, 20: 571-582, 2012.
- Levene et al., "Zero-Mode Waveguides for Single-Molecule Analysis at High Concentrations," *Science* 299, 682-686, 2003.
- Li et al., "A photocleavable fluorescent nucleotide for DNA sequencing and analysis," *Proc. Natl. Acad. Sci.*, 100: 414-419, 2003.
- Li et al., "DNA molecules and configurations in a solid-state nanopore microscope," *Nat Mater.*, Sep. 2003, 2(9):611-5.
- Li et al., "RASL-seq for Massively Parallel and Quantitative Analysis of Gene Expression," *Curr Protoc Mol Biol.*, Apr. 2012, 4(13):1-10.

(56)

References Cited

OTHER PUBLICATIONS

- Lin et al., "Microfluidic cell trap array for controlled positioning of single cells on adhesive micropatterns," *Lab Chip*, Feb. 2013, 13(4):714-721.
- Linnarsson, "Recent advances in DNA sequencing methods—general principles of sample preparation," *Experimental Cell Research*, 316: 1339-1343, 2010.
- Liu et al. "Barcoded oligonucleotides ligated on RNA amplified for multiplexed and parallel in situ analyses," *Nucleic Acids Res.*, Mar. 8, 2021, 49(10):e58, 15 pages.
- Liu et al., "Method for Quantitative Proteomics Research by Using Metal Element Chelated Tags Coupled with Mass Spectrometry," *Analytical Chemistry*, 2006, 78:6614-6621.
- Liu et al., "Surface and interface control on photochemically initiated immobilization," *J Am Chem Soc.*, Nov. 2006, 128(43):14067-72.
- Liu et al., An integrated and sensitive detection platform for biosensing application based on Fe@Au magnetic nanoparticles as bead array carries Biosensors and Bioelectronics, 2010, 26(4):1442-1448.
- Lizardi et al., "Mutation detection and single-molecule counting using isothermal rolling-circle amplification," *Nat. Genet.* 19: 225-232, 1998.
- Lopez-Otin et al., "Protease degradomics: a new challenge for proteomics," *Nat Rev Mol Cell Biol.*, Jul. 2002, 3(7):509-19.
- Lovatt et al., "Transcriptome in vivo analysis (TIVA) of spatially defined single cells in live tissue.," *Nature Methods* 11, 190-196, 2014.
- Lu et al., "A microfluidic electroporation device for cell lysis," *Lab Chip*, Jan. 2005, 5(1):23-29.
- Lubeck et al., "Single cell systems biology by super-resolution imaging and combinatorial labeling," *Nature Methods*, Jan. 2013, 9(7):743-748, 18 pages.
- Lund et al., "Assessment of methods for covalent binding of nucleic acids to magnetic beads, Dynabeads, and the characteristics of the bound nucleic acids in hybridization reactions.," *Nucleic Acids Res.*, 16: 10861-80, 1988.
- Lundberg et al., "High-fidelity amplification using a thermostable DNA polymerase isolated from *Pyrococcus furiosus*," *Gene*, 108(1): 1-6, 1991.
- Lundberg et al., "Homogeneous antibody-based proximity extension assays provide sensitive and specific detection of low-abundant proteins in human blood," *Nucleic Acids Res.*, 39(15): e102, 2011.
- Lundberg et al., "Multiplexed homogeneous proximity ligation assays for high-throughput protein biomarker research in serological material," *Mol Cell Proteomics*, 10(4): M110.004978, 2011.
- Lundin et al., "Increased throughput by parallelization of library preparation for massive sequencing," *PLoS One*, Apr. 2010, 5(4):e10029, 7 pages.
- Lundquist et al., "Parallel confocal detection of single molecules in real time," *Opt. Lett.* 33, 1026-1028, 2008.
- Lyamichev et al., "Invader assay for SNP genotyping," *Methods Mol Biol.*, 2003, 212:229-40.
- Lyamichev et al., "Polymorphism identification and quantitative detection of genomic DNA by invasive cleavage of oligonucleotide probes," *Nat Biotechnol.*, Mar. 1999, 17(3):292-6.
- Lyck, et al., "Immunohistochemical Markers for Quantitative Studies of Neurons and Glia in Human Neocortex," *J Histochem Cytochem* 56, 201-21, 2008.
- Lykidis et al., "Novel zinc-based fixative for high quality DNA, RNA and protein analysis," *Nucleic Acids Res.*, Jun. 2007, 35(12):e85, 10 pages.
- MacBeath et al., "Printing proteins as microarrays for high-throughput function determination," *Science*, Sep. 2000, 289(5485):1760-1763.
- MacIntyre, "Unmasking antigens for immunohistochemistry.," *Br J Biomed Sci.* 58, 190-6, 2001.
- Macosko et al., "Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets," *Cell* 161, 1202-1214, May 2015.
- Magaki et al., "An introduction to Performance of Immunohistochemistry," *Biobanking: Methods and Protocols*, Method in Molecular Biology, Yong (ed.), 2019, 1897, Chapter 25, 289-298.
- Malkov et al., "Multiplexed measurements of gene signatures in different analytes using the Nanostring nCounter™ Assay System." *BMC research notes.*, 2009, 2:80.
- Maniatis et al., "Spatiotemporal Dynamics of Molecular Pathology in Amyotrophic Lateral Sclerosis", 54 pages, 2018.
- Marras, "Selection of fluorophore and quencher pairs for fluorescent nucleic acid hybridization probes," *Methods Mol Biol.*, 2006, 335:3-16.
- Martin, "Cutadapt removes adapter sequences from high-throughput sequencing reads," *EMBnet Journal*, 2011, 17(1):10-12.
- Marx, "Method of the Year 2020: spatially resolved transcriptomics," *Jan. 2021*, 18(1):1 pages.
- Marx, "Method of the Year: spatially resolved transcriptomics," *Nat Methods*, Jan. 2021, 18(1):9-14.
- Massey et al., "Fluorescence resonance energy transfer (FRET) for DNA biosensors: FRET pairs and Förster distances for various dye-DNA conjugates," *Anal Chim Acta.*, May 2006, 568(1-2):181-9.
- Masuda et al., "Analysis of chemical modification of RNA from formalin-fixed samples and optimization of molecular biology applications for such samples," *Nucleic Acids Research*, Nov. 1999, 27(22):4436-4443.
- Materna et al., "High accuracy, high-resolution prevalence measurement for the majority of locally expressed regulatory genes in early sea urchin development," *Gene Expr Patterns.*, 2010, 10(4-5):177-184.
- Mattheyses et al., "Imaging with total internal reflection fluorescence microscopy for the cell biologist," *J Cell Sci.*, Nov. 2010, 123(Pt 21):3621-3628.
- Mauleon et al., "Precise Spatial and Temporal Control of Oxygen within In Vitro Brain Slices via Microfluidic Gas Channels," *PLoS One*, Aug. 2012, 7(8):e43309, 9 pages.
- McCloskey et al., "Encoding PCR Products with Batch-stamps and Barcodes," *Biochem. Genet.* 45: 761-767, 2007.
- Mcgee, "Structure and Analysis of Affymetrix Arrays," *UTSW Microarray Analysis Course*, Oct. 28, 2005, 68 pages.
- McKernan et al., "Sequence and structural variation in a human genome uncovered by short-read, massively parallel ligation sequencing using two-base encoding." *Genome Res.*, 19: 1527-41, 2009.
- Megason et al., "Imaging in Systems Biology," *Cell* 130, Sep. 7, 2007, pp. 784-795.
- Metzker "Sequencing technologies—the next generation," *Nature Reviews Genetics*, 11: 31-46, 2010.
- Meyer et al., "Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity," *BMC Evol. Biol.*, Mar. 2010, 10:70, 12 pages.
- Micke et al., "Biobanking of fresh frozen tissue: RNA is stable in nonfixed surgical specimens," *Lab Invest.*, Feb. 2006, 86(2):202-11.
- Miele et al., "Mapping cis- and trans-chromatin interaction networks using chromosome conformation capture (3C)," *Methods Mol Biol.*, 2009, 464:105-21.
- Miller et al., "Basic Concepts of Microarrays and Potential Applications in Clinical Microbiology," *Clinical Microbiology Reviews*, vol. 22, No. 4, pp. 611-633, 2009.
- Miner et al., "Molecular barcodes detect redundancy and contamination in hairpin-bisulfite PCR," *Nucleic Acids Res.*, Sep. 2004, 32(17):e135, 4 pages.
- Mir et al., "Sequencing by cyclic ligation and cleavage (CycliC) directly on a microarray captured template," *Nucleic Acids Research*, 37(1):e5, 8 pages, 2009.
- Mishra, "Three-dimensional genome architecture and emerging technologies: looping in disease", *Genome Medicine*, 9: 87, 14 pages, 2017.
- Mitra et al., "Digital genotyping and haplotyping with polymerase colonies," *Proc. Natl. Acad. Sci. USA*, May 2003, 100(10):5926-5931.
- Mitra et al., "Fluorescent in situ sequencing on polymerase colonies," *Anal Biochem*, Sep. 2003, 320(1):55-65.

(56)

References Cited

OTHER PUBLICATIONS

- Mitra et al., "In situ localized amplification and contact replication of many individual DNA molecules," *Nucleic Acids Res.*, Dec. 1999, 27(24):e34, 6 pages.
- Mitsuhashi et al., "Gene manipulation on plastic plates," *Nature* 357: 519-520, 1992.
- Mizusawa et al., "A bacteriophage lambda vector for cloning with BamHI and Sau3 A," *Gene*, 20: 317-322, 1982.
- Mlecinek et al., "Histopathologic-based prognostic factors of colorectal cancers are associated with the state of the local immune reaction," *J Clin Oncol.*, Feb. 2011, 29(6):610-8.
- Moncada et al., "Building a tumor atlas: integrating single-cell RNA-Seq data with spatial transcriptomics in pancreatic ductal adenocarcinoma", Institute for Computational Medicine, bioRxiv, 28 pages, 2018.
- Moor et al., "Spatial transcriptomics: paving the way for tissue-level systems biology", *Science Direct, Current Opinion in Biotechnology*, 46: 126-133, 2017.
- Morlan et al., "Selective depletion of rRNA enables whole transcriptome profiling of archival fixed tissue," *PLoS One*, Aug. 2012, 7(8):e42882, 8 pages.
- Mortazavi et al., "Mapping and quantifying mammalian transcriptomes by RNA-Seq," *Nature Methods*, 5(7): 621-8, 2008.
- Moshrefzadeh et al., "Nonuniform photobleaching of dyed polymers for optical waveguides," *Applied Physics Letters*, 1993, 62:16-18.
- Motea et al., "Terminal deoxynucleotidyl transferase: the story of a misguided DNA polymerase," *Biochim Biophys Acta.*, May 2010, 1804(5):1151-66.
- Mueller et al., "RNA Integrity Number (RIN)-Standardization of RNA Quality Control," *Agilent Technologies*, 2004, 8 pages.
- Nadji et al., "Immunohistochemistry of tissue prepared by a molecular-friendly fixation and processing system," *Appl Immunohistochem Mol Morphol.*, Sep. 2005, 13(3):277-82.
- Nagahara et al., "Neuroprotective effects of brain-derived neurotrophic factor in rodent and primate models of Alzheimer's disease," *Nat Med.*, Mar. 2009, 15(3):331-337.
- Nagai et al., "Site-specific DNA cleavage by antisense oligonucleotides covalently linked to phenazine di-N-oxide," *J Biol. Chem.*, Dec. 1991, 266(35):23994-4002.
- Nakamura et al., "Biocompatible inkjet printing technique for designed seeding of individual living cells," *Tissue Eng*, Nov. 2005, 11(11-12):1658-1666.
- Nakao et al., "Myosin heavy chain gene expression in human heart failure," *J Clin Invest.*, Nov. 1997, 100(9):2362-70.
- Nallur et al., "Signal amplification by rolling circle amplification on DNA microarrays," *Nucleic Acids Res.*, Dec. 1, 2001, 29(23):e118, 9 pages.
- Nam et al., "Nanoparticle-Based Bio-Bar Codes for the Ultrasensitive Detection of Proteins," *Science*, Sep. 26, 2003, 301(5641):1884-1886.
- Nandakumar et al., "How an RNA Ligase Discriminates RNA versus DNA Damage," *Molecular Cell*, 2004, 16(2):211-221.
- Nandakumar et al., "RNA Substrate Specificity and Structure-guided Mutational Analysis of Bacteriophage T4 RNA Ligase 2," *Journal of Biological Chemistry*, Jul. 2004, 279(30):31337-31347.
- Nawy, "Spatial transcriptomics", *Nature Methods*, vol. 15, No. 1, 2018.
- Ng et al., "Gene identification signature (GIS) analysis for transcriptome characterization and genome annotation," *Nature Methods*, 2(2): 105-111, 2005.
- Ng et al., "Massively parallel sequencing and rare disease," *Human Molec. Genetics*, 19(2): R119-R124, 2010.
- Ng et al., "Multiplex sequencing of paired-end ditags (MS-PET): a strategy for the ultra-high-throughput analysis of transcriptomes and genomes," *Nucleic Acids Research*, Jul. 2006, 34(12): e84, 10 pages.
- Nichols et al., "RNA Ligases," *Curr Protoc Mol Biol.*, Oct. 2008, 84(1):3.15.1-3.15.4.
- Nicholson, "Diffusion and related transport mechanisms in brain tissue," *Rep. Prog. Phys.*, Jun. 2001, 64(7):815-884.
- Niedringhaus et al., "Landscape of next-generation sequencing technologies," *Anal Chem.*, Jun. 2011, 83(12):4327-41.
- Niemeyer, "The developments of semisynthetic DNA-protein conjugates," *Trends Biotechnol.*, Sep. 2002, 20(9): 395-401.
- Nikiforov et al., "The use of 96-well polystyrene plates for DNA hybridization-based assays: an evaluation of different approaches to oligonucleotide immobilization," *Anal Biochem*, May 1995, 227(1):201-9.
- Niklas et al., "Selective permeabilization for the high-throughput measurement of compartmented enzyme activities in mammalian cells," *Anal Biochem*, Sep. 2011, 416(2):218-27.
- Nilsson et al., "Padlock Probes: Circularizing Oligonucleotides for Localized DNA Detection," *Science*, Sep. 30, 1994, 265(5181):2085-2088.
- Nilsson et al., "RNA-templated DNA ligation for transcript analysis," *Nucleic Acids Res.*, Jan. 2001, 29(2):578-81.
- Nuovo, "In situ PCR: protocols and applications.," *Genome Res*, Feb. 1995, 4 (4):151-167.
- Ohtsubo et al., "Bacterial insertion sequences," *Curr Top Microbiol Immunol.*, 1996, 204:1-26.
- Oleinikov et al., "Self-assembling protein arrays using electronic semiconductor microchips and in vitro translation," *J Proteome Res*, May-Jun. 2003, 2(3): 313-319.
- Olink Proteomics AB, *Proseek® Multiplex 96x96 User Manual*, 2016, 12 pages.
- Olivier, "The Invader assay for SNP genotyping," *Mutat. Res.*, Jun. 2005, 573(1-2):103-110.
- Oren et al., "Selective lysis of bacteria but not mammalian cells by diastereomers of melittin: structure-function study," *Biochemistry*, Feb. 1997, 36(7):1826-35.
- Osada et al., "Epitope mapping using ribosome display in a reconstituted cell-free protein synthesis system," *J Biochem*, May 2009, 145(5): 693-700.
- O-Shannessy et al., "Detection and quantitation of hexa-histidine-tagged recombinant proteins on western blots and by a surface plasmon resonance biosensor technique," *Anal Biochem*, 229(1): 119-124, 1995.
- Ostuni et al., "Patterning Mammalian Cells Using Elastomeric Membranes," *Langmuir*, Aug. 2000, 16(20):7811-7819.
- Ozsolak et al., "Digital transcriptome profiling from attomole-level RNA samples," *Genome Res.*, Apr. 2010, 20(4):519-25.
- Palamanda et al., "Evaluation of CYP1A1 and CYP2B1/2 m-RNA Induction in Rat Liver Slices Using the NanoString® Technology: A Novel Tool for Drug Discovery Lead Optimization," *Drug metabolism letters*, Nov. 3, 2009, 3(3):171-175.
- Pandey et al., "Inhibition of terminal deoxynucleotidyl transferase by adenine dinucleotides. Unique inhibitory action of Ap5A," *FEBS Lett.*, Mar. 1987, 213(1):204-8.
- Park et al., "Cancer gene therapy using adeno-associated virus vectors," *Front Biosci.*, Jan. 2008, 13:2653-59.
- Park et al., "The Estimation of Breast Cancer Disease-Probability by Difference of Individual Susceptibility," *Cancer Res. Treat.*, Feb. 2003, 35(1):35-51, Abstract.
- Patil et al., "DNA-based therapeutics and DNA delivery systems: a comprehensive review," *AAPS J*, Apr. 2005, 7(1):E61-77.
- Patton et al., "Rainbow's end: the quest for multiplexed fluorescence quantitative analysis in proteomics." *Current Opinion in Chemical Biology*, Feb. 1, 2002, 6(1):63-69.
- PCT International Preliminary Report on Patentability in International Appln. No. PCT/EP2012/056823, dated Oct. 15, 2013, 7 pages.
- PCT International Preliminary Report on Patentability in International Appln. No. PCT/EP2013/071645, dated Apr. 21, 2015, 6 pages.
- PCT International Search Report and Written Opinion in International Appln. No. PCT/EP2012/056823, dated Jun. 29, 2012, 12 pages.
- PCT International Search Report and Written Opinion issued in International Appln. No. PCT/EP2013/071645, dated Nov. 28, 2013, 10 pages.

(56)

References Cited

OTHER PUBLICATIONS

- Pearson et al., "Improved tools for biological sequence comparison," *Proc. Natl. Acad. Sci.*, May 1988, 85:2444-2448.
- Pellestor et al., "The peptide nucleic acids (PNAs), powerful tools for molecular genetics and cytogenetics," *Eur J Hum Genet.*, Sep. 2004, 12(9):694-700.
- Pemov et al., "DNA analysis with multiplex microarray-enhanced PCR," *Nucl. Acids Res.*, Jan. 2005, 33(2):e11, 9 pages.
- Penland et al., "RNA expression analysis of formalin-fixed paraffin-embedded tumors," *Laboratory Investigation*, Apr. 2007, 87(4):383-391.
- Perler et al., "Intervening sequences in an Archaea DNA polymerase gene," *PNAS USA*, Jun. 1992, 89(12): 5577-5581.
- Perocchi et al., "Antisense artifacts in transcriptome microarray experiments are resolved by actinomycin D," *Nucleic Acids Res.*, 2007, 35(19):e128, 7 pages.
- Pettersen et al., "Generations of sequencing technologies," *Genomics*, 2009, 105-111.
- Pipenburg et al., "DNA detection using recombination proteins," *PLoS Biol.*, Jul. 2006, 4(7):e204, 7 pages.
- Pirici et al., "Antibody elution method for multiple immunohistochemistry on primary antibodies raised in the same species and of the same subtypem," *J. Histochem. Cytochem.*, Jun. 2009, 57(6):567-75.
- Piston et al., "Fluorescent protein FRET: the good, the bad and the ugly," *Trends Biochem Sci.*, Sep. 2007, 32(9):407-14.
- Plasterk, "The Tc1/mariner transposon family," *Curr Top Microbiol Immunol.*, 1996, 204:125-43.
- Pluen et al., "Diffusion of macromolecules in agarose gels: comparison of linear and globular configurations," *Biophys J.*, Jul. 1999, 77(1):542-552.
- Polsky-Cynkin et al., "Use of DNA Immobilized on Plastic and Agarose Supports to Detect DNA by Sandwich Hybridization," *Clin. Chem.* 31: 1438-1443, 1985.
- Porreca et al., "Polony DNA sequencing," *Curr Protoc Mol Biol.*, Nov. 2006, Chapter 7, Unit 7.8, pp. 7.8.1-7.8.22.
- Promega, "GoScript™ Reverse Transcription System—Technical Manual," promega.com, revised Dec. 2012, 24 pages.
- U.S. Appl. No. 61/267,363, filed Dec. 7, 2009 (Year: 2009).
- Punwaney et al., "Human papillomavirus may be common within nasopharyngeal carcinoma of Caucasian Americans: investigation of Epstein-Barr virus and human papillomavirus in eastern and western nasopharyngeal carcinoma using ligation-dependent polymerase chain reaction," *Head & Neck*, Jan. 1999, 21(1):21-29.
- Raab et al., "Human tRNA genes function as chromatin insulators," *EMBO J.*, Jan. 2012, 31(2):330-50.
- Raj et al., "Imaging individual mRNA molecules using multiple singly labeled probes," *Nature Methods*, Oct. 2008, 5(10):877-879, 9 pages.
- Rajeswari et al., "Multiple pathogen biomarker detection using an encoded bead array in droplet PCR," *J. Microbiol Methods*, Aug. 2017, 139:22-28.
- Ramachandran et al., "Next-generation high-density self-assembling functional protein arrays," *Nature Methods*, Jun. 2008, 5(6):535-538.
- Ramanujan et al., "Diffusion and convection in collagen gels: implications for transport in the tumor interstitium," *Biophys. J.*, Sep. 2002, 83(3):1650-1660.
- Ranki et al., "Sandwich hybridization as a convenient method for the detection of nucleic acids in crude samples," *Gene* 21: 77-85, cellulose, 1983.
- Raouane et al., "Lipid conjugated oligonucleotides: a useful strategy for delivery," *Bioconjug Chem.*, Jun. 2012, 23(6):1091-104.
- Razonable, "Antiviral drugs for viruses other than human immunodeficiency virus," *Mayo Clinic Proceedings*, Oct. 2011, 86(10):1009-26.
- Reijenga et al., "Buffer Capacity, Ionic Strength and Heat Dissipation in Capillary Electrophoresis," *Journal of Chromatography A*, Sep. 13, 1996, 744(1-2):147-153.
- Reinartz et al., "Massively parallel signature sequencing (MPSS) as a tool for in-depth quantitative gene expression profiling in all organisms," *Brief Funct Genomic Proteomic*, Feb. 2002, 1(1):95-104.
- Rettig et al., "Large-scale single-cell trapping and imaging using microwell arrays," *Anal Chem*, Sep. 2005, 77(17):5628-5634.
- Reznikoff, "Tn5 as a model for understanding DNA transposition," *Mol Microbiol.*, Mar. 2003, 47(5):1199-206.
- Ristova et al., "Study of hydrogenated amorphous silicon thin films as a potential sensor for He—Ne laser light detection," *Applied Surface Science*, Sep. 2003, 218(1-4):44-53.
- Roberts et al., "RNA-peptide fusions for the in vitro selection of peptides and proteins," *PNAS USA*, Nov. 1997, 94: 12297-122302.
- Robinson et al., "Small-sample estimation of negative binomial dispersion, with applications to SAGE data," *Biostatistics*, Apr. 2008, 9(2):321-332.
- Rogers et al., "Immobilization of oligonucleotides onto a glass support via disulfide bonds: A method for preparation of DNA microarrays," *Anal Biochem.*, Jan. 1999, 266(1):23-30.
- Rogers et al., "Use of a novel cross-linking method to modify adenovirus tropism," *Gene Ther.*, Dec. 1997, 4(12):1387-92.
- Ronaghi et al., "A sequencing method based on real-time pyrophosphate," *Science*, Jul. 1998, 281(5375): 363-365.
- Ronaghi et al., "Real-time DNA sequencing using detection of pyrophosphate release," *Analytical Biochemistry*, Nov. 1996, 242(1): 84-89.
- Ronaghi, "Pyrosequencing sheds light on DNA sequencing," *Genome Res*, Jan. 2001, 11(1):3-11.
- Rosenthal et al., "Cell patterning chip for controlling the stem cell microenvironment," *Biomaterials*, Jul. 2007, 28(21):3208-3216.
- Rouillard et al., "OligoArray 2.0: design of oligonucleotide probes for DNA microarrays using a thermodynamic approach," *Nuc. Acid Research*, Jun. 2003, 31(12): 3057-3062.
- Rountenberg et al., "Microfluidic probe: a new tool for integrating microfluidic environments and electronic wafer-probing," *Lab Chip*, Oct. 2009, 10(1):123-127.
- Rubin et al., "Whole-genome resequencing reveals loci under selection during chicken domestication," *Nature*, Mar. 2010, 464: 587-591.
- Rubina et al., "Hydrogel-based protein microchips: manufacturing, properties, and applications," *Biotechniques*, May 2003, 34(5):1008-14.
- Running et al., "A procedure for productive coupling of synthetic oligonucleotides to polystyrene microtiter wells for hybridization capture," *Biotechniques*, Mar. 1990, 8(3):276-279.
- Rush et al., "New Aldehyde Tag Sequences Identified by Screening Formylglycine Generating Enzymes in Vitro and in Vivo," *J. of American Chemical Society*, Aug. 2008, 130(37): 12240-12241.
- Rusk, "Spatial transcriptomics," *Nature Methods*, Sep. 2016, 13(9):710-711.
- Russell et al., "Molecular mechanisms of late endosome morphology, identity and sorting," *Curr. Opin. Cell Bio.*, Aug. 2006, 18(4):422-428.
- Salem et al., "Multidimensional transcriptomics provides detailed information about immune cell distribution and identity in HER2+ breast tumors," *bioRxiv*, 41 pages, 2018.
- San Paulo et al., "High-resolution imaging of antibodies by tapping-mode atomic force microscopy: attractive and repulsive tip-sample interaction regimes," *Biophys J.*, Mar. 2000, 78(3):1599-1605.
- Sano et al., "Immuno-PCR: Very Sensitive Antigen Detection by Means of Specific Antibody-DNA Conjugates," *Science*, Oct. 2, 1992, 258(5079):120-122.
- Saxonov, "Mastering Biology to Advance Human Health," *10x Genomics*, Sep. 2020, 41 pages.
- Schaus et al., "A DNA nanoscope via auto-cycling proximity recording," *Nat. Commun.* 8, 696, 10 pages, 2017.
- Schellings et al., "Absence of SPARC results in increased cardiac rupture and dysfunction after acute myocardial infarction," *J Exp Med.*, Jan. 2009, 206(1):113-23.
- Schena et al., "Quantitative monitoring of gene expression patterns with a complementary DNA microarray," *Science*, Oct. 1995, 270(5235):467-470.

(56)

References Cited

OTHER PUBLICATIONS

- Schena et al., "Entering the Postgenome Era," *Science*, 1995, 270:368-9, 371.
- Schlapak et al., "Glass surfaces grafted with high-density poly (ethylene glycol) as substrates for DNA oligonucleotide microarrays," *Langmuir*, Jan. 2006, 22: 277-285.
- Schmitt et al., "Detection of ultra-rare mutations by next-generation sequencing," *PNAS* (2012) 109:14508-14523.
- Scholz et al., "The Molecular Chaperone Hsp90 Is Required for Signal Transduction by Wild-Type Hck and Maintenance of Its Constitutively Active Counterpart1," *Cell Growth Differ.*, Aug. 2001, 12(8):409-417.
- Schouten et al., "Relative quantification of 40 nucleic acid sequences by multiplex ligation-dependent probe amplification," *Nucleic Acids Res.*, Jun. 2002, 30(12):e57, 13 pages.
- Schroeder et al., "The RIN: an RNA integrity No. for assigning integrity values to RNA measurements," *BMC Molecular Biology*, Jan. 2006, 7:3, 14 pages.
- Schwartz et al., "Capturing native long-range contiguity by in situ library construction and optical sequencing," *PNAS*, Nov. 13, 2012, 109(46):18749-18754.
- Schweitzer et al., "Immunoassays with rolling circle DNA amplification: A versatile platform for ultrasensitive antigen detection," *Proc. Natl Acad. Sci. USA*, May 22, 2000, 97:10113-119.
- Schweitzer et al., "Multiplexed protein profiling on microarrays by rolling-circle amplification," *Nature Biotechnology*, Apr. 2002, 20(4):359-365.
- Schwers et al., "A high-sensitivity, medium-density, and target amplification-free planar waveguide microarray system for gene expression analysis of formalin-fixed and paraffin-embedded tissue," *Clin. Chem.*, Nov. 2009, 55(11):1995-2003.
- ScienceDirect.com [online], "Plant Fibers," Definition, 2011, retrieved on Apr. 13, 2022, retrieved from URL<<https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/plant-fibers>>, 9 pages.
- Sekar et al., "Fluorescence resonance energy transfer (FRET) microscopy imaging of live cell protein localizations," *J Cell Biol.*, Mar. 2003, 160(5):629-33.
- Sergeeva et al., "Display technologies: Application for the discovery of drug and gene delivery agents," *Advanced Drug Delivery Reviews* (2006) 58(15):1622-1654.
- Seuryck-Servoss et al., "Evaluation of Surface Chemistries for Antibody Microarrays," *Anal Biochem.*, 371(1): 105-115, 2007.
- Shalon et al., "A DNA microarray system for analyzing complex DNA samples using two-color fluorescent probe hybridization," *Genome Res.*, Jul. 1996, 6(7):639-45.
- Shelbourne et al., "Fast copper-free click DNA ligation by the ring-strain promoted alkyne-azide cycloaddition reaction," *Chem. Commun.*, 47: 6257-6259, 2011.
- Shendure et al., "Accurate Multiplex Polony Sequencing of an Evolved Bacterial Genome," *Science*, 2005, 309:1728-1732.
- Shi et al., "The MicroArray Quality Control (MAQC) project shows inter- and intraplatform reproducibility of gene expression measurements," *Nature Biotechnology*, 2006, 24(9):1151-61.
- Shi, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies," *Clin. Chem.*, Feb. 2001, 47(2):164-172.
- Shibata et al., "Detection of human papilloma virus in paraffin-embedded tissue using the polymerase chain reaction," *J Exp Med.*, Jan. 1988, 167(1):225-30.
- Shirai et al., "Novel Tools for Analyzing Gene Expressions in Single Cells," The 5th International Workshop on Approaches to Single-Cell Analysis, The University of Tokyo, Mar. 3-4, 2011, 1 page.
- Shoemaker et al., "Quantitative phenotypic analysis of yeast deletion mutants using a highly parallel molecular bar-coding strategy," *Nature genetics* (1996) 14:450-456.
- Shults et al., "A multiplexed protein kinase assay," *Chem Bio Chem* (2007) 8:933-942.
- Sievertzon et al., "Transcriptome analysis in primary neural stem cells using a tag cDNA amplification method," *BMC Neuroscience*, Dec. 2005, 6: 28.
- Simonis et al., "Nuclear organization of active and inactive chromatin domains uncovered by chromosome conformation capture-on-chip (4C)," *Nat Genet.*, Nov. 2006, 38(11):1348-54.
- Slomovic et al., "Addition of poly(A) and poly(A)-rich tails during RNA degradation in the cytoplasm of human cells," *Proc Natl Acad Sci USA*, Apr. 2010, 107(16):7407-12.
- Slonim and Yanai, "Getting started in gene expression microarray analysis," *Plos Computational Biology*, 2009, 5(10):e1000543.
- Smolock et al., "Ribosomal Protein L17, Rpl17, is an Inhibitor of Vascular Smooth Muscle Growth and Carotid Intima Formation," *Circulation*, Nov. 2012, 126(20):2418-2427.
- Soderberg et al. "Characterizing proteins and their interactions in cells and tissues using the in situ proximity ligation assay," *Methods*, Jul. 2008, 45(3):227-232.
- Soderberg et al., "Direct observation of individual endogenous protein complexes in situ by proximity ligation," *Nature Methods*, 2006, 3:995-1000.
- Son et al., "A platform for ultrasensitive and selective multiplexed marker protein assay toward early-stage cancer diagnosis," *Nanomedicine*, Feb. 7, 2007, 2(1):79-82.
- Soni and Meller, "Progress toward ultrafast DNA sequencing using solid-state nanopores," *Clin Chem.*, 2007, 53: 1996-2001.
- Spieß et al., "A highly efficient method for long-chain cDNA synthesis using trehalose and betaine," *Anal. Biochem.*, Feb. 2002, 301(2):168-74.
- Spitale et al., "Structural imprints in vivo decode RNA regulatory mechanisms," *Nature*, 519(7544): 486-90, 2015.
- Spurgeon et al., "High Throughput Gene Expression Measurement with Real Time PCR in a Microfluidic Dynamic Array," *Plos ONE*, 2008, 3(2):e1662.
- Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Science*, Jul. 2016, 353(6294):78-82.
- Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Science*, Jun. 2016, Supplementary Materials, 353(6294):78-82, 41 pages.
- Stevens Jr. et al., "Enhancement of phosphoprotein analysis using a fluorescent affinity tag and mass spectrometry," *Rapid Commun Mass Spectrom*, 2005, 19(15):2157-62.
- Stimpson et al., "Real-time detection of DNA hybridization and melting on oligonucleotide arrays by using optical wave guides," *Proc Natl Acad Sci USA*, Jul. 1995, 92(14):6379-83.
- Stoddart et al., "Single-nucleotide discrimination in immobilized DNA oligonucleotides with a biological nanopore," *PNAS U S A.*, May 2009, 106(19):7702-7707.
- Strell et al., "Placing RNA in context and space—methods for spatially resolved transcriptomics," *The FEBS Journal*, 14 pages, 2017.
- Stroh et al., "Quantum dots spectrally distinguish multiple species within the tumor milieu in vivo," *Nat Med.*, Jun. 2005, 11(6):678-82.
- Subramanian et al., "Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles," *PNAS*, Oct. 2005, 102(43):15545-15550.
- Suh et al., "A simple soft lithographic route to fabrication of poly(ethylene glycol) microstructures for protein and cell patterning," *Biomaterials*, Feb. 2004, 25(3):557-563.
- Sumitomo et al., "Ca²⁺ ion transport through channels formed by -hemolysin analyzed using a microwell array on a Si substrate," *Biosensors and Bioelectronics*, 2012, 31(1):445-450.
- Summersgill et al., "Fluorescence In Situ Hybridization Analysis of Formalin Fixed Paraffin Embedded Tissues, Including Tissue Microarrays," Chapter 4, Bridger, J. Ed., *Methods in Molecular Biology* 659, 2010, 51-70, 2010.
- Sun et al., "Direct immobilization of DNA probes on non-modified plastics by UV irradiation and integration in microfluidic devices for rapid bioassay," *Anal. Bio. Chem.*, 402: 741-748, 2012.
- Surzhik et al., "Template-dependent biosynthesis of poly(G) x poly(C) and its antiviral activity in vitro and in vivo," *Antiviral Res.*, May 1988, 38(2):131-40.
- Sutherland et al., "Utility of formaldehyde cross-linking and mass spectrometry in the study of protein-protein interactions," *J. Mass Spectrom.*, Jun. 2008, 43(6):699-715.

(56)

References Cited

OTHER PUBLICATIONS

- Swartz et al., "Interstitial flow and its effects in soft tissues," *Annu Rev Biomed Eng.*, 2007, 9:229-56.
- Syková et al., "Diffusion in brain extracellular space," *Physiol Rev.*, Oct. 2008, 88(4):1277-340.
- Tai et al., "Replication-competent retrovirus vectors for cancer gene therapy," *Front Biosci.*, Jan. 2008, 13:3083-95.
- Takahashi et al., "In Vitro Selection of Protein and Peptide Libraries Using mRNA Display," *Nucleic Acid and Peptide Aptamers: Methods and Protocols (2009) 535:293-314 (Ch.17)*.
- Tan et al., "Polyethylene peel-off arrays to probe the role of cell-cell interactions in tumour angiogenesis," *Integr Biol (Camb)*, Oct. 2009, 1(10):587-594.
- Tang et al., "RNA-Seq analysis to capture the transcriptome landscape of a single cell," *Nat Protoc.*, 5: 516-35, 2010.
- Taniguchi et al., "Quantitative analysis of gene expression in a single cell by qPCR," *Nature Methods*, 6, pp. 503-506, 2009.
- Tawfik et al., "Man-made cell-like compartments for molecular evolution," *Nat Biotechnol.*, Jul. 1998, 16(7):652-6.
- Taylor et al., "Microfluidic local perfusion chambers for the visualization and manipulation of synapses," *Neuron.*, Apr. 2010, 66(1):57-68, 25 pages.
- Taylor et al., "Mitochondrial DNA mutations in human disease," *Nature Reviews Genetics*. May 2005, 6(5):389-402.
- Tedpella.com, [online], "Glass Microscope Slides," 2021, retrieved on Jun. 9, 2021, retrieved from URL <https://www.tedpella.com/histo_html/slides.htm>, 7 pages.
- Tegtmeier et al., "Alternative Interactions of the SV40 A Protein with DNA," *Virology*, 1981, 115:75-87.
- Tentori et al., "Detection of Isoforms Differing by a Single Charge Unit in Individual Cells," *Chem. Int. Ed.*, 55, 12431, 2016.
- Thacker et al., "Alkaline Hydrolysis—Carcass Disposal: A Comprehensive Review," *National Agriculture Biosecurity Center*, Aug. 2004, Chapter 6, pp. 1-12.
- Thiery et al., "Multiplex target protein imaging in tissue sections by mass spectrometry—TAMSIM," *Rapid Commun. Mass Spectrom.*, 2007, 21:823-829.
- Thorne et al., "In vivo diffusion analysis with quantum dots and dextrans predicts the width of brain extracellular space," *Proc Natl Acad Sci USA*, Apr. 2006, 103(14):5567-5572.
- Thornton, "High rate thick film growth." *Annual review of materials science*, Aug. 1977, 7(1):239-60.
- Tian et al., "Antigen peptide-based immunosensors for rapid detection of antibodies and antigens," *Anal Chem* May 26, 2009, 81(13):5218-5225.
- Tijssen et al., "Overview of principles of hybridization and the strategy of nucleic acid assays" in *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, 1993, 24(Chapter 2), 65 pages.
- Timofeev et al., "Regioselective immobilization of short oligonucleotides to acrylic copolymer gels," *Nucleic Acids Res.*, Aug. 1996, 24(16):3142-8.
- Tolbert et al., "New Methods for Proteomic Research: Preparation of Proteins with N-Terminal Cysteines for Labeling and Conjugation," *Angewandte Chemie International Edition*, Jun. 17, 2002, 41(12):2171-4.
- Toseland, "Fluorescent labeling and modification of proteins," *J Chem Biol.*, Apr. 2013, 6(3):85-95.
- Totet et al., "Immunocompetent infants as a human reservoir for *Pneumocystis jirovecii*: rapid screening by non-invasive sampling and real-time PCR at the mitochondrial large subunit rRNA gene," *J Eukaryot Microbiol.*, 2003, pp. 668-669.
- Toubanaki et al., "Dry-reagent disposable biosensor for visual genotyping of single nucleotide polymorphisms by oligonucleotide ligation reaction: application to pharmacogenetic analysis," *Hum Mutat.*, Aug. 2008, 29(8):1071-8.
- Toy et al., "A Simple Plastic Perfusion Chamber for Continuous Maintenance and Cinematography of Tissue Cultures," *Experimental Cell Research*, 1958, 14:97-103.
- Tseng et al., "Magnetic nanoparticle-mediated massively parallel mechanical modulation of single-cell behavior," *Nat Methods*, Nov. 2012, 9(11):1113-1119.
- Twyman et al., "Techniques Patents for SNP Genotyping," *Pharmacogenomics*, Jan. 2003, 4(1):67-79.
- Tzanetakis et al., "The use of reverse transcriptase for efficient first- and second-strand cDNA synthesis from single- and double-stranded RNA templates," *J Virol Methods*, Mar. 2005, 24(1-2):73-7.
- Ueno et al., "cDNA Display: Rapid Stabilization of mRNA Display," *Antibody-Drug Conjugates, Methods in Molecular Biology*, Jan. 2012, pp. 113-135.
- Ulery et al., "Biomedical Applications of Biodegradable Polymers," *J Polym Sci B Polym Phys.*, Jun. 2011, 49(12):832-864.
- U.S. Appl. No. 60/416,118 Fan et al., "Multiplex Nucleic Acid Analysis Using Archived or Fixed Samples," filed Oct. 3, 2002, 22 pages.
- Valencia et al., "mRNA-Display-Based Selections for Proteins with Desired Functions: A Protease-Substrate Case Study," *Biotechnology progress*, May 2008, 24(3):561-9.
- Valley et al., "Optoelectronic tweezers as a tool for parallel single-cell manipulation and stimulation," *IEEE Trans Biomed Circuits Syst.*, Dec. 2009, 3(6):424-31.
- Van Gelder et al., "Amplified RNA synthesized from limited quantities of heterogeneous cDNA," *Proc. Natl. Acad. Sci. USA* 87, 1663-1667, 1990.
- Van Ness et al., "A versatile solid support system for oligodeoxynucleotide probe-based hybridization assays," *Nucleic Acids Res.* 19: 3345-3350, 1991.
- Vandenbroucke et al., "Quantification of splice variants using real-time PCR," *Nucleic Acids Research*, 2001, 29(13):e68, 7 pages.
- Vandernoot et al., "cDNA normalization by hydroxyapatite chromatography to enrich transcriptome diversity in RNA-seq applications," *Biotechniques*, Dec. 2012, 53(6):373-80.
- Vasiliskov et al., "Fabrication of microarray of gel-immobilized compounds on a chip by copolymerization," *Biotechniques*, Sep. 1999, 27(3):592-606.
- Velculescu et al., "Serial analysis of gene expression," *Science*, Oct. 20, 1995, 270(5235):484-7.
- Vermesh et al., "High-density, multiplexed patterning of cells at single-cell resolution for tissue engineering and other applications," *Angew Chem Int Ed Engl*, Aug. 2011, 50(32):7378-7380.
- Vickovic et al., "High-definition spatial transcriptomics for in situ tissue profiling," *Nat Methods*, Oct. 2019, 16(10):987-990.
- Vickovic et al., "Massive and parallel expression profiling using microarrayed single-cell sequencing," *Nature Communications*, 2016, 7(13182):1-9.
- Vincent et al., "Helicase-dependent isothermal DNA amplification," *EMBO Rep.*, Aug. 2004, 5(8):795-800.
- Viollet et al., "T4 RNA ligase 2 truncated active site mutants: improved tools for RNA analysis," *BMC Biotechnol.*, Jul. 2011, 11:72, 14 pages.
- Vogelstein et al., "Digital PCR," *Proceedings of the National Academy of Sciences*, Aug. 3, 1999, 96:9236-9241.
- Wade et al., "Genome sequence, comparative analysis, and population genetics of the domestic horse," *Science.*, 326: 865-7, 2009.
- Waichman et al., "Functional immobilization and patterning of proteins by an enzymatic transfer reaction," *Analytical chemistry*, Jan. 21, 2010, 82(4):1478-85.
- Walker et al., "Strand displacement amplification—an isothermal, in vitro DNA amplification technique," *Nucleic acids research*. 1992 Apr. 11, 1992, 20(7):1691-1696.
- Walker et al., Ed., "Chapter 1: Basic Techniques in Molecular Biology," *Medical Biomethods Handbook*, Humana Press, Totowa, New Jersey, 2005, 19 pages.
- Wang et al., "Mutations in NEXN, a Z-disc gene, are associated with hypertrophic cardiomyopathy," *Am J Hum Genet.*, Nov. 2010, 87(5):687-93.
- Wang et al., "Paramagnetic microspheres with core-shell structures," *Journal of Materials Science*, Apr. 2012, 47(16):5946-54.
- Wang et al., "Single cell analysis: the new frontier in 'omics'," *Trends Biotechnol.*, 28: 281-90, 2010.

(56)

References Cited

OTHER PUBLICATIONS

- Wang et al., "Three-dimensional intact-tissue sequencing of single-cell transcriptional states", *Science*, 361(6400), 22 pages, 2018.
- Wang et al., "High-fidelity mRNA amplification for gene profiling." *Nature biotechnology*, Apr. 2000, 18(4):457-459.
- Wang, "RNA amplification for successful gene profiling analysis," *J Transl Med.*, Jul. 2005, 3:28, 11 pages.
- Watanabe et al., "Cellular networks involved in the influenza virus life cycle," *Cell Host & Microbe*, Jun. 2010, 7(6):427-39.
- Waxman et al., "De-regulation of common housekeeping genes in hepatocellular carcinoma," *BMC Genomics*, 2007, 1-9.
- Weichhart et al., "Functional selection of vaccine candidate peptides from *Staphylococcus aureus* whole-genome expression libraries in vitro," *Infection and Immunity*, 2003, 71 (8):4333-4641.
- Weinreich et al., "Evidence that the cis Preference of the Tn5 Transposase is Caused by Nonproductive Multimerization," *Genes and Development*, Oct. 1994, 8(19):2363-2374.
- Weinstein et al., "DNA microscopy: Optics-free spatio-genetic imaging by a stand-alone chemical reaction", *bioRxiv*, 41 pages, 2018.
- Wheeler et al., "Microfluidic device for single-cell analysis," *Analytical Chemistry*, Jul. 2003, 75(14):3581-3586.
- Wiedmann et al., "Ligase chain reaction (LCR)—overview and applications," *PCR Methods Appl.*, Feb. 1994, 3(4):S51-64.
- Wikipedia.org [online], "Random hexamer," Jan. 2012, Retrieved on Jan. 21, 2022, retrieved from URL <https://en.wikipedia.org/w/index.php?title=Random_hexamer&oldid=473042236>, 1 page.
- Williams, "RAC reviews serious adverse event associated with AAV therapy trial," *Mol Ther.*, Dec. 2007, 15(12):2053-54.
- Willi-Monnerat et al., "Comprehensive spatiotemporal transcriptomic analyses of the ganglionic eminences demonstrate the uniqueness of its caudal subdivision," *Molecular and Cellular Neuroscience* 37(4):845-856, 2008.
- Wilson et al., "New transposon delivery plasmids for insertional mutagenesis in *Bacillus anthracis*," *J Microbiol Methods*, Dec. 2007, 71(3):332-5.
- Wolf et al., "Rapid hybridization kinetics of DNA attached to submicron latex particles", *Nucleic Acids Res.* 15: 2911-2926, 1987.
- Wong et al., "Direct Site-Selective Covalent Protein Immobilization Catalyzed by a Phosphopantetheinyl Transferase," *J. Am. Chem Soc.*, 2008, 130:12456-64.
- Woo et al., "A Comparison of cDNA, Oligonucleotide, and Affymetrix GeneChip Gene Expression Microarray Platforms," *Journal of Biomolecular Techniques*, 2004, 15(4), 276-284.
- Wood et al., "Single cell trapping and DNA damage analysis using microwell arrays," *PNAS*, Jun. 2010, 107(22):10008-10013.
- Worthington et al., "Cloning of random oligonucleotides to create single-insert plasmid libraries," *Analyt. Biochem*, 2001, 294:169-175.
- Wright et al., "Reusable, reversibly sealable parylene membranes for cell and protein patterning," *J Biomed Mater Res A.*, May 2008, 85(2):530-538.
- Wu et al., "Detection DNA Point Mutation with Rolling-Circle Amplification Chip," *Bioinformatics and Biomedical Engineering (ICBBE)*, 2010 4th International Conference on IEEE, Piscataway, NJ, USA, Jun. 18, 2010, 1-4 pages.
- Xiao et al., "Direct determination of haplotypes from single DNA molecules," *Nature Methods*, 2009, 6(3):199-201.
- Xie et al., "CryoFISH: Fluorescence In Situ Hybridization on Ultrathin Cryosections," *Fluorescence in situ Hybridization (FISH)*, Jul. 2010, pp. 221-230.
- Yan et al., "Decorin gene delivery inhibits cardiac fibrosis in spontaneously hypertensive rats by modulation of transforming growth factor-beta/Smad and p38 mitogen-activated protein kinase signaling pathways," *Hum Gene Ther.*, Oct. 2009, 20(10):1190-200.
- Yasukawa et al., "Effects of organic solvents on the reverse transcription reaction catalyzed by reverse transcriptases from avian myeloblastosis virus and Moloney murine leukemia virus," *Biosci Biotechnol Biochem.*, 2010, 74(9):1925-30.
- Yeakley et al., "Profiling alternative splicing on fiber-optic arrays," *Nature Biotechnology*, Apr. 2002, 20(4):353-358.
- Yershov et al., "DNA analysis and diagnostics on oligonucleotide microchips," *Proc. Natl. Acad. Sci. USA*, May 1996, 93(10):4913-4918.
- Yet et al., "Cardiac-specific expression of heme oxygenase-1 protects against ischemia and reperfusion injury in transgenic mice," *Circ Res.*, Jul. 2001, 89(2):168-73.
- Yin et al., "Genetically encoded short peptide tag for versatile protein labeling by Sfp phosphopantetheinyl transferase," *PNAS*, 2005, 102(44):15815-20.
- Yonezawa et al., "DNA display for in vitro selection of diverse peptide libraries," *Nucleic Acids Research*, 2003, 31 (19):e118.
- Yusof et al., "Inkjet-like printing of single-cells," *Lab Chip*, Jul. 2011, 11(14):2447-2454.
- Zhang et al., "A novel mechanism of transposon-mediated gene activation," *PLoS Genet.*, Oct. 2009, 5(10):e1000689, 10 pages.
- Zhang et al., "Assembling DNA through Affinity Binding to Achieve Ultrasensitive Protein Detection," *Angew Chem Int Ed* (2013) 52:2-10.
- Zhang et al., "Binding-induced DNA assembly and its application to yoctomole detection of proteins," *Anal Chem* (2012) 84(2):877-884.
- Zhang et al., "Single-base mutational analysis of cancer and genetic diseases using membrane bound modified oligonucleotides," *Nucleic Acids Res.*, Jul. 1991, 19(14):3929-33.
- Zhang et al., "Stripping custom microRNA microarrays and the lessons learned about probe-slide interactions," *Anal Biochem.*, Mar. 2009, 386(2):222-7.
- Zheng et al., "Origins of human mitochondrial point mutations as DNA polymerase mediated errors. *Mutat. Res.* 599(1-2): 11-20, 2006.
- Zheng, "Spectroscopy-based quantitative fluorescence resonance energy transfer analysis," *Methods Mol Biol.*, 2006, 337:65-77.
- Zhou et al., "Genetically encoded short peptide tags for orthogonal protein labeling by Sfp and AcpS phosphopantetheinyl transferases," *ACS Chemical Biol.*, 2007 2 5 : 337-346.
- Zhu et al., "Reverse Transcriptase Template Switching: A Smart Approach for Full-Length cDNA Library Construction," *BioTechniques*, 2001, 30(4): 892-897.
- Zieba et al., "Bright-field microscopy visualization of proteins and protein complexes by in situ proximity ligation with peroxidase detection," *Clin Chem*, Jan. 2010, 56(1):99-110.
- Zilberman et al., "Genome-wide analysis of DNA methylation patterns," *Development* (2007) 134: 3959-3965.
- Zlobec et al., "Next-generation tissue microarray (ngTMA) increases the quality of biomarker studies: an example using CD3, CD8, and CD45RO in the tumor microenvironment of six different solid tumor types," *Journal of Translational Medicine*, 2013 11(104):1-7.
- Zuker, "Mfold web server for nucleic acid folding and hybridization prediction," *Nucleic Acids Res.*, Jul. 2003, 31(13):3406-15.
- Akatsuka et al., "Rapid screening of T-cell receptor (TCR) variable gene usage by multiplex PCR: Application for assessment of clonal composition," *Tissue Antigens*, Jan. 5, 1999, 53:122-134.
- Akatsuka et al., "T cell receptor clonal diversity following allogeneic marrow grafting," *Human Immunology*, Jun.-Jul. 1996, 48:125-134.
- Almog et al., "The crystal structures of the psychrophilic subtilisin S41 and the mesophilic subtilisin Sph reveal the same calcium-loaded state," *Proteins*, Feb. 1, 2009, 74(2):489-496.
- Bibikova et al., "Quantitative gene expression profiling in formalin-fixed paraffin-embedded tissues using universal bead arrays," *The American Journal of Pathology*, Nov. 1, 2004, 165(5):1799-1807.
- Bowen et al., "Concurrent V(D)J recombination and DNA end instability increase interchromosomal trans-rearrangements in ATM-deficient thymocytes," *Nucleic Acids Research*, Apr. 1, 2013, 41(8):4535-4548.
- Cai et al., "Glutathione-mediated shedding of PEG layers based on disulfide-linked cationomers for DNA delivery," *J. Mater. Chem.*, Sep. 20, 2011, 21(38):14639-14645.
- Choi et al., "Multiplexed detection of mRNA using porosity-tuned hydrogel microparticles," *Analytical chemistry*, Sep. 28, 2012, 84(21):9370-9378.

(56)

References Cited

OTHER PUBLICATIONS

- Fan et al., "A versatile assay for high-throughput gene expression profiling on universal array matrices," *Genome Research*, May 1, 2004, 14(5):878-885.
- Ha et al., "Self-assembly hollow nanosphere for enzyme encapsulation," *Soft Matter*, Feb. 11, 2010, 6, 1405-1408, 10 pages.
- Holscher et al., "Application of Laser-Assisted Microdissection for Tissue and Cell-Specific Analysis of RNA," *Progress in Botany*, Jan. 2008, 69(3):141-167.
- Ju et al., "Supramolecular dendrimer capsules by cooperative binding," *Chem. Commun.*, Jan. 7, 2011, 47(1):268-270, 8 pages.
- Kim, "Development of Microdevices for Applications to Bioanalysis," Dissertation for the degree of Doctor of Philosophy, University of Texas at Austin, Aug. 2007, 176 pages.
- Kuiper et al., "Enzymes containing porous polymersomes as nano reaction vessels for cascade reactions," *Org. Biomol. Chem.*, Oct. 15, 2008, 6(23):4315-4318.
- Kung et al., "Long Noncoding RNAs: Past, Present, and Future," *Genetics*, Mar. 1, 2013, 193(3):651-669.
- Kwon et al., "Polyelectrolyte Gels—Fundamentals and Applications," Nov. 10, 2006, *Polymer Journal*, 38, pp. 1211-1219.
- Miller et al., "Rapid and Efficient Enzyme Encapsulation in a Dendrimer Silica Nanocomposite," *Macromolecular Bioscience*, Oct. 25, 2006, 6(10):839-845.
- Rahimi et al., "Synthesis and Characterization of Thermo-Sensitive Nanoparticles for Drug Delivery Applications," *J. Biomed. Nanotechnol.* Dec. 2008, 4(4):482-490, 19 pages.
- Robins et al., "Comprehensive assessment of T-cell receptor β -chain diversity in $\alpha\beta$ T cells," *Blood*, Nov. 5, 2009, 114(19):4099-4107.
- Villa et al., "Partial V(D)J Recombination Activity Leads to Omenn Syndrome," *Cell*, May 29, 1998, 93:885-896.
- Williams et al., "Disc electrophoresis in polyacrylamide gels: extension to new conditions of pH and buffer," *Annals of the New York Academy of Sciences*, Dec. 1964, 121(2):373-381.
- Mabruk et al., "In situ hybridization: detecting viral nucleic acid in formalin-fixed, paraffin-embedded tissue samples," *Expert Rev. Mol. Diagn.*, 2004, 4(5):653-661.
- Belton et al., "Hi-C: A comprehensive technique to capture the conformation of genomes," *Methods*, Nov. 2012, 58(3):268-276, 16 pages.
- Deininger et al., "Allograft inflammatory factor-1 defines a distinct subset of infiltrating macrophages/microglial cells in rat and human gliomas," *Acta Neuropathol.*, Dec. 2000, 100(6):673-680. Extended European Search Report in European Appln. No. 22208440. 2, dated Jun. 6, 2023, 10 pages.
- Fan et al., "Illumina Universal Bead Arrays," *Methods in Enzymology*, 2006, 410:57-73.
- Jeffers, "A Basic Subroutine for Geary's Contiguity Ratio," *J. Royal Stat. Society, Series D*, Dec. 1973, 22(4):299-302.
- Li et al., "Beyond Moran's I: Testing for Spatial Dependence Based on the Spatial Autoregressive Model," *Geographical Analysis*, Sep. 18, 2007, 39(4):357-375.
- Liu et al., "Preparation and Characterization of Temperature-Sensitive Poly(N-isopropylacrylamide)-b-poly(D,L-lactide) Microspheres for Protein Delivery," *Biomacromolecules*, 2003, 4(6):1784-1793.
- Manz et al., "Phylogenetic Composition, Spatial Structure, and Dynamics of Lotic Bacterial Biofilms Investigated by Fluorescent in situ Hybridization and Confocal Laser Scanning Microscopy," *Microb Ecol.* May 1999, 37(4):225-237.
- Zhou et al., "Analysis of the expression profile of Dickkopf-1 gene in human glioma and the associate with tumor malignancy," *Journal of Experimental & Clinical Cancer Research*, Oct. 28, 2010, 29(138):1-7.
- Altman et al., "Phenotypic Analysis of Antigen-Specific T Lymphocytes," *Science*, Oct. 4, 1996, 274(5284):94-96.
- Andersen et al., "Parallel Detection of Antigen-Specific T Cell Responses by Combinatorial Encoding of MHC Multimers," *Nature Protocols*, Apr. 12, 2012, 7(5):891-902.
- Bakker et al., "Conditional MHC class I ligands and peptide exchange technology for the human MHC gene products HLA-A1, -A3, -A11, and -B7," *Proc Natl Acad Sci USA*, Mar. 11, 2008, 105(10):3825-3830.
- Hadrup et al., "Parallel detection of antigen-specific T-cell responses by multidimensional encoding of MHC multimers," *Nat. Methods.*, Jul. 2009, 6(7), 520-526.
- Soen et al., "Detection and Characterization of Cellular Immune Responses Using Peptide-MHC Microarrays," *PLOS Biology*, Dec. 22, 2003, 1(3):429-438.

* cited by examiner




FIG. 1




FIG. 2




FIG. 4




FIG. 5




FIG. 6

Lane 7 blast results	
E-value	Hits
	1,00E-06
Barcode	Hits
Tag1	2165
Tag2	1223
Tag3	1338
Tag4	3605
Tag5	763
Tag6	1083
Tag7	1175
Tag8	1350
Tag9	616
Tag10	710
Tag11	1602
Tag12	1359
Tag13	1404
Tag14	1124
Tag15	1077
Tag16	1238
Tag17	1122
Tag18	1547
Tag19	1401
Tag20	1354




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11





FIG. 12




Overall Results for sample 3 : R6 3/1 2011

Number of peaks found: 1

Peak table for sample 3 : R6 3/1 2011

Peak	Size [bp]	Conc. [ng/μl]	Molarity [nmol/l]	Observations
1	◀50	8.30	251.5	Lower Marker
2	386	22.87	89.8	
3	▶10,380	4.20	0.6	Upper Marker

FIG. 13



Overall Results for sample 4 : R8 3/1 2011

Number of peaks found: 1

Peak table for sample 4 : R8 3/1 2011

Peak	Size [bp]	Conc. [ng/μl]	Molarity [nmol/l]	Observations
1	◀50	8.30	251.5	Lower Marker
2	676	21.53	48.2	
3	▶10,380	4.20	0.6	Upper Marker

FIG. 14





FIG. 15






FIG. 16



Overall Results for Sample 3 : 1B-200bp fragments-Genomic intern (168bp)

Number of peaks found: 2


FIG. 17



Overall Results for Sample 7 : 1F-700bp fragments-Genomic intern (168bp)

Number of peaks found: 2


FIG. 18



Overall Results for Sample 4 : 2b-200bp fragments-F Genomic+P (Random)

Number of peaks found: 2

FIG. 19



Overall Results for Sample 8 : 2F-700bp fragments-F Genomic+P (Random)

Number of peaks found: 2

FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27






FIG. 28

Chip A9



Chip A1






FIG. 29

Amplification with A431_2+2_FW_A and RE_A primers



Positive Control: 330bp

FIG. 30




FIG. 31

METHODS OF DETECTING ANALYTES**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 18/047,092, filed on Oct. 17, 2022, which is a continuation of U.S. patent application Ser. No. 17/704,830, now U.S. Pat. No. 11,479,809, filed on Mar. 25, 2022, which is a continuation of U.S. patent application Ser. No. 17/474,922, now U.S. Pat. No. 11,352,659, filed on Sep. 14, 2021, which is a continuation of U.S. patent application Ser. No. 16/013,654, filed on Jun. 20, 2018, which is a continuation of U.S. patent application Ser. No. 14/111,482, now U.S. Pat. No. 10,030,261, filed on Oct. 11, 2013, which is a U.S. National Phase Application under 35 U.S.C. § 371 of International Patent Application No. PCT/EP2012/056823, filed Apr. 13, 2012, which claims benefit to U.K. Patent Application No. 1106254.4, filed on Apr. 13, 2011, the entire contents of the foregoing applications are incorporated herein by reference.

SEQUENCE LISTING IN ELECTRONIC FORMAT

This application contains a Sequence Listing that has been submitted electronically as an XML file named 47706-0011011_SL_ST26.xml. The XML file, created on Feb. 16, 2023, is 127,615 bytes in size. The material in the XML file is hereby incorporated by reference in its entirety.

The present invention relates generally to the localized or spatial detection of nucleic acid in a tissue sample. The nucleic acid may be RNA or DNA. Thus, the present invention provides methods for detecting and/or analysing RNA, e.g. RNA transcripts or genomic DNA, so as to obtain spatial information about the localisation, distribution or expression of genes, or indeed about the localisation or distribution of any genomic variation (not necessarily in a gene) in a tissue sample, for example in an individual cell. The present invention thus enables spatial genomics and spatial transcriptomics.

More particularly, the present invention relates to a method for determining and/or analysing a transcriptome or genome and especially the global transcriptome or genome, of a tissue sample. In particular the method relates to a quantitative and/or qualitative method for analysing the distribution, location or expression of genomic sequences in a tissue sample wherein the spatial expression or distribution or location pattern within the tissue sample is retained. Thus, the new method provides a process for performing “spatial transcriptomics” or “spatial genomics”, which enables the user to determine simultaneously the expression pattern, or the location/distribution pattern of the genes expressed or genes or genomic loci present in a tissue sample.

The invention is particularly based on array technology coupled with high throughput DNA sequencing technologies, which allows the nucleic acid molecule (e.g. RNA or DNA molecules) in the tissue sample, particularly mRNA or DNA, to be captured and labelled with a positional tag. This step is followed by synthesis of DNA molecules which are sequenced and analysed to determine which genes are expressed in any and all parts of the tissue sample. Advantageously, the individual, separate and specific transcriptome of each cell in the tissue sample may be obtained at the same time. Hence, the methods of the invention may be said to provide highly parallel comprehensive transcriptome signatures from individual cells within a tissue sample without

losing spatial information within said investigated tissue sample. The invention also provides an array for performing the method of the invention and methods for making the arrays of the invention.

5 The human body comprises over 100 trillion cells and is organized into more than 250 different organs and tissues. The development and organization of complex organs, such as the brain, are far from understood and there is a need to dissect the expression of genes expressed in such tissues using quantitative methods to investigate and determine the genes that control the development and function of such tissues. The organs are in themselves a mixture of differentiated cells that enable all bodily functions, such as nutrient transport, defence etc. to be coordinated and maintained. Consequently, cell function is dependent on the position of the cell within a particular tissue structure and the interactions it shares with other cells within that tissue, both directly and indirectly. Hence, there is a need to disentangle how these interactions influence each cell within a tissue at the transcriptional level.

Recent findings by deep RNA sequencing have demonstrated that a majority of the transcripts can be detected in a human cell line and that a large fraction (75%) of the human protein-coding genes are expressed in most tissues. Similarly, a detailed study of 1% of the human genome showed that chromosomes are ubiquitously transcribed and that the majority of all bases are included in primary transcripts. The transcription machinery can therefore be described as promiscuous at a global level.

It is well-known that transcripts are merely a proxy for protein abundance, because the rates of RNA translation, degradation etc will influence the amount of protein produced from any one transcript. In this respect, a recent antibody-based analysis of human organs and tissues suggests that tissue specificity is achieved by precise regulation of protein levels in space and time, and that different tissues in the body acquire their unique characteristics by controlling not which proteins are expressed but how much of each is produced.

However, in subsequent global studies transcriptome and proteome correlations have been compared demonstrating that the majority of all genes were shown to be expressed. Interestingly, there was shown to be a high correlation between changes in RNA and protein levels for individual gene products which is indicative of the biological usefulness of studying the transcriptome in individual cells in the context of the functional role of proteins.

Indeed, analysis of the histology and expression pattern in tissues is a cornerstone in biomedical research and diagnostics. Histology, utilizing different staining techniques, first established the basic structural organization of healthy organs and the changes that take place in common pathologies more than a century ago. Developments in this field resulted in the possibility of studying protein distribution by immunohistochemistry and gene expression by in situ hybridization.

However, the parallel development of increasingly advanced histological and gene expression techniques has resulted in the separation of imaging and transcriptome analysis and, until the methods of the present invention, there has not been any feasible method available for global transcriptome analysis with spatial resolution.

As an alternative, or in addition, to in situ techniques, methods have developed for the in vitro analysis of proteins and nucleic acids, i.e. by extracting molecules from whole

tissue samples, single cell types, or even single cells, and quantifying specific molecules in said extracts, e.g. by ELISA, qPCR etc.

Recent developments in the analysis of gene expression have resulted in the possibility of assessing the complete transcriptome of tissues using microarrays or RNA sequencing, and such developments have been instrumental in our understanding of biological processes and for diagnostics. However, transcriptome analysis typically is performed on mRNA extracted from whole tissues (or even whole organisms), and methods for collecting smaller tissue areas or individual cells for transcriptome analysis are typically labour intensive, costly and have low precision.

Hence, the majority of gene expression studies based on microarrays or next generation sequencing of RNA use a representative sample containing many cells. Thus the results represent the average expression levels of the investigated genes. The separation of cells that are phenotypically different has been used in some cases together with the global gene expression platforms (Tang F et al, *Nat Protoc.* 2010; 5: 516-35; Wang D & Bodovitz S, *Trends Biotechnol.* 2010; 28:281-90) and resulted in very precise information about cell-to-cell variations. However, high throughput methods to study transcriptional activity with high resolution in intact tissues have not, until now, been available.

Thus, existing techniques for the analysis of gene expression patterns provide spatial transcriptional information only for one or a handful of genes at a time or offer transcriptional information for all of the genes in a sample at the cost of losing positional information. Hence, it is evident that methods to determine simultaneously, separately and specifically the transcriptome of each cell in a sample are required, i.e. to enable global gene expression analysis in tissue samples that yields transcriptomic information with spatial resolution, and the present invention addresses this need.

The novel approach of the methods and products of the present invention utilizes now well established array and sequencing technology to yield transcriptional information for all of the genes in a sample, whilst retaining the positional information for each transcript. It will be evident to the person of skill in the art that this represents a milestone in the life sciences. The new technology opens a new field of so-called "spatial transcriptomics", which is likely to have profound consequences for our understanding of tissue development and tissue and cellular function in all multicellular organisms. It will be apparent that such techniques will be particularly useful in our understanding of the cause and progress of disease states and in developing effective treatments for such diseases, e.g. cancer. The methods of the invention will also find uses in the diagnosis of numerous medical conditions.

Whilst initially conceived with the aim of transcriptome analysis in mind, as described in detail below, the principles and methods of the present invention may be applied also to the analysis of DNA and hence for genomic analyses also ("spatial genomics"). Accordingly, at its broadest the invention pertains to the detection and/or analysis of nucleic acid in general.

Array technology, particularly microarrays, arose from research at Stanford University where small amounts of DNA oligonucleotides were successfully attached to a glass surface in an ordered arrangement, a so-called "array", and used it to monitor the transcription of 45 genes (Schemm M et al, *Science.* 1995; 270: 368-9, 371).

Since then, researchers around the world have published more than 30,000 papers using microarray technology. Mul-

multiple types of microarray have been developed for various applications, e.g. to detect single nucleotide polymorphisms (SNPs) or to genotype or re-sequence mutant genomes, and an important use of microarray technology has been for the investigation of gene expression. Indeed, the gene expression microarray was created as a means to analyze the level of expressed genetic material in a particular sample, with the real gain being the possibility to compare expression levels of many genes simultaneously. Several commercial microarray platforms are available for these types of experiments but it has also been possible to create custom made gene expression arrays.

Whilst the use of microarrays in gene expression studies is now commonplace, it is evident that new and more comprehensive so-called "next-generation DNA sequencing" (NGS) technologies are starting to replace DNA microarrays for many applications, e.g. in-depth transcriptome analysis.

The development of NGS technologies for ultra-fast genome sequencing represents a milestone in the life sciences (Pettersson E et al, *Genomics.* 2009; 93: 105-11). These new technologies have dramatically decreased the cost of DNA sequencing and enabled the determination of the genome of higher organisms at an unprecedented rate, including those of specific individuals (Wade C M et al *Science.* 2009; 326: 865-7; Rubin J et al, *Nature* 2010; 464: 587-91). The new advances in high-throughput genomics have reshaped the biological research landscape and in addition to complete characterization of genomes it is possible also to study the full transcriptome in a digital and quantitative fashion. The bioinformatics tools to visualize and integrate these comprehensive sets of data have also been significantly improved during recent years.

However, it has surprisingly been found that a unique combination of histological, microarray and NGS techniques can yield comprehensive transcriptional or genomic information from multiple cells in a tissue sample which information is characterised by a two-dimensional spatial resolution. Thus, at one extreme the methods of the present invention can be used to analyse the expression of a single gene in a single cell in a sample, whilst retaining the cell within its context in the tissue sample. At the other extreme, and in a preferred aspect of the invention, the methods can be used to determine the expression of every gene in each and every cell, or substantially all cells, in a sample simultaneously, i.e. the global spatial expression pattern of a tissue sample. It will be apparent that the methods of the invention also enable intermediate analyses to be performed.

In its simplest form, the invention may be illustrated by the following summary. The invention requires reverse transcription (RT) primers, which comprise also unique positional tags (domains), to be arrayed on an object substrate, e.g. a glass slide, to generate an "array". The unique positional tags correspond to the location of the RT primers on the array (the features of the array). Thin tissue sections are placed onto the array and a reverse transcription reaction is performed in the tissue section on the object slide. The RT primers, to which the RNA in the tissue sample binds (or hybridizes), are extended using the bound RNA as a template to obtain cDNA, which is therefore bound to the surface of the array. As consequence of the unique positional tags in the RT primers, each cDNA strand carries information about the position of the template RNA in the tissue section. The tissue section may be visualised or imaged, e.g. stained and photographed, before or after the cDNA synthesis step to enable the positional tag in the cDNA molecule to be correlated with a position within the tissue sample. The

cDNA is sequenced, which results in a transcriptome with exact positional information. A schematic of the process is shown in FIG. 1. The sequence data can then be matched to a position in the tissue sample, which enables the visualization, e.g. using a computer, of the sequence data together with the tissue section, for instance to display the expression pattern of any gene of interest across the tissue (FIG. 2). Similarly, it would be possible to mark different areas of the tissue section on the computer screen and obtain information on differentially expressed genes between any selected areas of interest. It will be evident that the methods of the invention result in data that is in stark contrast to the data obtained using current methods to study mRNA populations. For example, methods based on in situ hybridization provide only relative information of single mRNA transcripts. Thus, the methods of the present invention have clear advantages over current in situ technologies. The global gene expression information obtainable from the methods of the invention also allows co-expression information and quantitative estimates of transcript abundance. It will be evident that this is a generally applicable strategy available for the analysis of any tissue in any species, e.g. animal, plant, fungus.

As noted above, and described in more detail below, it will be evident that this basic methodology could readily be extended to the analysis of genomic DNA, e.g. to identify cells within a tissue sample that comprise one or more specific mutations. For instance, the genomic DNA may be fragmented and allowed to hybridise to primers (equivalent to the RT primers described above), which are capable of capturing the fragmented DNA (e.g. an adapter with a sequence that is complementary to the primer may be ligated to the fragmented DNA or the fragmented DNA may be extended e.g. using an enzyme to incorporate additional nucleotides at the end of the sequence, e.g. a poly-A tail, to generate a sequence that is complementary to the primer) and priming the synthesis of complementary strands to the capture molecules. The remaining steps of the analysis may be as described above. Hence, the specific embodiments of the invention described below in the context of transcriptome analysis may also be employed in methods of analysing genomic DNA, where appropriate.

It will be seen from the above explanation that there is an immense value in coupling positional information to transcriptome or genome information. For instance, it enables global gene expression mapping at high resolution, which will find utility in numerous applications, including e.g. cancer research and diagnostics.

Furthermore, it is evident that the methods described herein differ significantly from the previously described methods for analysis of the global transcriptome of a tissue sample and these differences result in numerous advantages. The present invention is predicated on the surprising discovery that the use of tissue sections does not interfere with synthesis of DNA (e.g. cDNA) primed by primers (e.g. reverse transcription primers) that are coupled to the surface of an array.

Thus, in its first and broadest aspect, the present invention provides a method for localized detection of nucleic acid in a tissue sample comprising:

- (a) providing an array comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a primer for a primer extension or ligation reaction, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

- (i) a positional domain that corresponds to the position of the capture probe on the array, and
- (ii) a capture domain;
- (b) contacting said array with a tissue sample such that the position of a capture probe on the array may be correlated with a position in the tissue sample and allowing nucleic acid of the tissue sample to hybridise to the capture domain in said capture probes;
- (c) generating DNA molecules from the captured nucleic acid molecules using said capture probes as extension or ligation primers, wherein said extended or ligated DNA molecules are tagged by virtue of the positional domain;
- (d) optionally generating a complementary strand of said tagged DNA and/or optionally amplifying said tagged DNA;
- (e) releasing at least part of the tagged DNA molecules and/or their complements or amplicons from the surface of the array, wherein said part includes the positional domain or a complement thereof;
- (f) directly or indirectly analysing the sequence of the released DNA molecules.

The methods of the invention represent a significant advance over other methods for spatial transcriptomics known in the art. For example the methods described herein result in a global and spatial profile of all transcripts in the tissue sample. Moreover, the expression of every gene can be quantified for each position or feature on the array, which enables a multiplicity of analyses to be performed based on data from a single assay. Thus, the methods of the present invention make it possible to detect and/or quantify the spatial expression of all genes in single tissue sample. Moreover, as the abundance of the transcripts is not visualised directly, e.g. by fluorescence, akin to a standard microarray, it is possible to measure the expression of genes in a single sample simultaneously even wherein said transcripts are present at vastly different concentrations in the same sample.

Accordingly, in a second and more particular aspect, the present invention can be seen to provide a method for determining and/or analysing a transcriptome of a tissue sample comprising:

- (a) providing an array comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':
 - (i) a positional domain that corresponds to the position of the capture probe on the array, and
 - (ii) a capture domain;
- (b) contacting said array with a tissue sample such that the position of a capture probe on the array may be correlated with a position in the tissue sample and allowing RNA of the tissue sample to hybridise to the capture domain in said capture probes;
- (c) generating cDNA molecules from the captured RNA molecules using said capture probes as RT primers, and optionally amplifying said cDNA molecules;
- (d) releasing at least part of the cDNA molecules and/or optionally their amplicons from the surface of the array, wherein said released molecule may be a first strand and/or second strand cDNA molecule or an amplicon thereof and wherein said part includes the positional domain or a complement thereof;

(e) directly or indirectly analysing the sequence of the released molecules.

As described in more detail below, any method of nucleic acid analysis may be used in the analysis step. Typically this may involve sequencing, but it is not necessary to perform an actual sequence determination. For example sequence-specific methods of analysis may be used. For example a sequence-specific amplification reaction may be performed, for example using primers which are specific for the positional domain and/or for a specific target sequence, e.g. a particular target DNA to be detected (i.e. corresponding to a particular cDNA/RNA or gene etc.). An exemplary analysis method is a sequence-specific PCR reaction.

The sequence analysis information obtained in step (e) may be used to obtain spatial information as to the RNA in the sample. In other words the sequence analysis information may provide information as to the location of the RNA in the sample. This spatial information may be derived from the nature of the sequence analysis information determined, for example it may reveal the presence of a particular RNA which may itself be spatially informative in the context of the tissue sample used, and/or the spatial information (e.g. spatial localisation) may be derived from the position of the tissue sample on the array, coupled with the sequencing information. Thus, the method may involve simply correlating the sequence analysis information to a position in the tissue sample e.g. by virtue of the positional tag and its correlation to a position in the tissue sample. However, as described above, spatial information may conveniently be obtained by correlating the sequence analysis data to an image of the tissue sample and this represents one preferred embodiment of the invention. Accordingly, in a preferred embodiment the method also includes a step of:

(f) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (c).

In its broadest sense, the method of the invention may be used for localized detection of a nucleic acid in a tissue sample. Thus, in one embodiment, the method of the invention may be used for determining and/or analysing all of the transcriptome or genome of a tissue sample e.g. the global transcriptome of a tissue sample. However, the method is not limited to this and encompasses determining and/or analysing all or part of the transcriptome or genome. Thus, the method may involve determining and/or analysing a part or subset of the transcriptome or genome, e.g. a transcriptome corresponding to a subset of genes, e.g. a set of particular genes, for example related to a particular disease or condition, tissue type etc.

Viewed from another aspect, the method steps set out above can be seen as providing a method of obtaining a spatially defined transcriptome or genome, and in particular the spatially defined global transcriptome or genome, of a tissue sample.

Alternatively viewed, the method of the invention may be seen as a method for localized or spatial detection of nucleic acid, whether DNA or RNA in a tissue sample, or for localized or spatial determination and/or analysis of nucleic acid (DNA or RNA) in a tissue sample. In particular, the method may be used for the localized or spatial detection or determination and/or analysis of gene expression or genomic variation in a tissue sample. The localized/spatial detection/determination/analysis means that the RNA or DNA may be localized to its native position or location within a cell or tissue in the tissue sample. Thus for example, the RNA or DNA may be localized to a cell or group of cells, or type of cells in the sample, or to particular regions of areas within

a tissue sample. The native location or position of the RNA or DNA (or in other words, the location or position of the RNA or DNA in the tissue sample), e.g. an expressed gene or genomic locus, may be determined.

The invention can also be seen to provide an array for use in the methods of the invention comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

- (i) a positional domain that corresponds to the position of the capture probe on the array, and
- (ii) a capture domain to capture RNA of a tissue sample that is contacted with said array.

In a related aspect, the present invention also provides use of an array, comprising a substrate on which multiple species of capture probe are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

- (i) a positional domain that corresponds to the position of the capture probe on the array; and
- (ii) a capture domain; to capture RNA of a tissue sample that is contacted with said array.

Preferably, said use is for determining and/or analysing a transcriptome and in particular the global transcriptome, of a tissue sample and further comprises steps of:

- (a) generating cDNA molecules from the captured RNA molecules using said capture probes as RT primers and optionally amplifying said cDNA molecules;
- (b) releasing at least part of the cDNA molecules and/or optionally their amplicons from the surface of the array, wherein said released molecule may be a first strand and/or second strand cDNA molecule or an amplicon thereof and wherein said part includes the positional domain or a complement thereof;
- (c) directly or indirectly analysing the sequence of the released molecules; and optionally
- (d) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (a).

It will be seen therefore that the array of the present invention may be used to capture RNA, e.g. mRNA of a tissue sample that is contacted with said array. The array may also be used for determining and/or analysing a partial or global transcriptome of a tissue sample or for obtaining a spatially defined partial or global transcriptome of a tissue sample. The methods of the invention may thus be considered as methods of quantifying the spatial expression of one or more genes in a tissue sample. Expressed another way, the methods of the present invention may be used to detect the spatial expression of one or more genes in a tissue sample. In yet another way, the methods of the present invention may be used to determine simultaneously the expression of one or more genes at one or more positions within a tissue sample. Still further, the methods may be seen as methods for partial or global transcriptome analysis of a tissue sample with two-dimensional spatial resolution.

The RNA may be any RNA molecule which may occur in a cell. Thus it may be mRNA, tRNA, rRNA, viral RNA, small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), microRNA (miRNA), small interfering RNA

(siRNA), piwi-interacting RNA (piRNA), ribozymal RNA, antisense RNA or non-coding RNA. Preferably however it is mRNA.

Step (c) in the method above (corresponding to step (a) in the preferred statement of use set out above) of generating cDNA from the captured RNA will be seen as relating to the synthesis of the cDNA. This will involve a step of reverse transcription of the captured RNA, extending the capture probe, which functions as the RT primer, using the captured RNA as template. Such a step generates so-called first strand cDNA. As will be described in more detail below, second strand cDNA synthesis may optionally take place on the array, or it may take place in a separate step, after release of first strand cDNA from the array. As also described in more detail below, in certain embodiments second strand synthesis may occur in the first step of amplification of a released first strand cDNA molecule.

Arrays for use in the context of nucleic acid analysis in general, and DNA analysis in particular, are discussed and described below. Specific details and embodiments described herein in relation to arrays and capture probes for use in the context of RNA, apply equally (where appropriate) to all such arrays, including those for use with DNA.

As used herein the term “multiple” means two or more, or at least two, e.g. 3, 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 400, 500, 1000, 2000, 5000, 10,000, or more etc. Thus for example, the number of capture probes may be any integer in any range between any two of the aforementioned numbers. It will be appreciated however that it is envisaged that conventional-type arrays with many hundreds, thousands, tens of thousands, hundreds of thousands or even millions of capture probes may be used.

Thus, the methods outlined herein utilise high density nucleic acid arrays comprising “capture probes” for capturing and labelling transcripts from all of the single cells within a tissue sample e.g. a thin tissue sample slice, or “section”. The tissue samples or sections for analysis are produced in a highly parallelized fashion, such that the spatial information in the section is retained. The captured RNA (preferably mRNA) molecules for each cell, or “transcriptomes”, are transcribed into cDNA and the resultant cDNA molecules are analyzed, for example by high throughput sequencing. The resultant data may be correlated to images of the original tissue samples e.g. sections through so-called barcode sequences (or ID tags, defined herein as positional domains) incorporated into the arrayed nucleic acid probes.

High density nucleic acid arrays or microarrays are a core component of the spatial transcriptome labelling method described herein. A microarray is a multiplex technology used in molecular biology. A typical microarray consists of an arrayed series of microscopic spots of oligonucleotides (hundreds of thousands of spots, generally tens of thousands, can be incorporated on a single array). The distinct position of each nucleic acid (oligonucleotide) spot (each species of oligonucleotide/nucleic acid molecule) is known as a “feature” (and hence in the methods set out above each species of capture probe may be viewed as a specific feature of the array; each feature occupies a distinct position on the array), and typically each separate feature contains in the region of picomoles (10^{-12} moles) of a specific DNA sequence (a “species”), which are known as “probes” (or “reporters”). Typically, these can be a short section of a gene or other nucleic acid element to which a cDNA or cRNA sample (or “target”) can hybridize under high-stringency hybridization

conditions. However, as described below, the probes of the present invention differ from the probes of standard microarrays.

In gene expression microarrays, probe-target hybridization is usually detected and quantified by detection of visual signal, e.g. a fluorophore, silver ion, or chemiluminescence-label, which has been incorporated into all of the targets. The intensity of the visual signal correlates to the relative abundance of each target nucleic acid in the sample. Since an array can contain tens of thousands of probes, a microarray experiment can accomplish many genetic tests in parallel.

In standard microarrays, the probes are attached to a solid surface or substrate by a covalent bond to a chemical matrix, e.g. epoxy-silane, amino-silane, lysine, polyacrylamide etc. The substrate typically is a glass, plastic or silicon chip or slide, although other microarray platforms are known, e.g. microscopic beads.

The probes may be attached to the array of the invention by any suitable means. In a preferred embodiment the probes are immobilized to the substrate of the array by chemical immobilization. This may be an interaction between the substrate (support material) and the probe based on a chemical reaction. Such a chemical reaction typically does not rely on the input of energy via heat or light, but can be enhanced by either applying heat, e.g. a certain optimal temperature for a chemical reaction, or light of certain wavelength. For example, a chemical immobilization may take place between functional groups on the substrate and corresponding functional elements on the probes. Such corresponding functional elements in the probes may either be an inherent chemical group of the probe, e.g. a hydroxyl group or be additionally introduced. An example of such a functional group is an amine group. Typically, the probe to be immobilized comprises a functional amine group or is chemically modified in order to comprise a functional amine group. Means and methods for such a chemical modification are well known.

The localization of said functional group within the probe to be immobilized may be used in order to control and shape the binding behaviour and/or orientation of the probe, e.g. the functional group may be placed at the 5' or 3' end of the probe or within sequence of the probe. A typical substrate for a probe to be immobilized comprises moieties which are capable of binding to such probes, e.g. to amine-functionalized nucleic acids. Examples of such substrates are carboxy, aldehyde or epoxy substrates. Such materials are known to the person skilled in the art. Functional groups, which impart a connecting reaction between probes which are chemically reactive by the introduction of an amine group, and array substrates are known to the person skilled in the art.

Alternative substrates on which probes may be immobilized may have to be chemically activated, e.g. by the activation of functional groups, available on the array substrate. The term “activated substrate” relates to a material in which interacting or reactive chemical functional groups were established or enabled by chemical modification procedures as known to the person skilled in the art. For example, a substrate comprising carboxyl groups has to be activated before use. Furthermore, there are substrates available that contain functional groups that can react with specific moieties already present in the nucleic acid probes.

Alternatively, the probes may be synthesized directly on the substrate. Suitable methods for such an approach are known to the person skilled in the art. Examples are manufacture techniques developed by Agilent Inc., Affymetrix Inc., Roche Nimblegen Inc. or Flexgen BV. Typically, lasers

and a set of mirrors that specifically activate the spots where nucleotide additions are to take place are used. Such an approach may provide, for example, spot sizes (i.e. features) of around 30 μm or larger.

The substrate therefore may be any suitable substrate known to the person skilled in the art. The substrate may have any suitable form or format, e.g. it may be flat, curved, e.g. convexly or concavely curved towards the area where the interaction between the tissue sample and the substrate takes place. Particularly preferred is the where the substrate is a flat, i.e. planar, chip or slide.

Typically, the substrate is a solid support and thereby allows for an accurate and traceable positioning of the probes on the substrate. An example of a substrate is a solid material or a substrate comprising functional chemical groups, e.g. amine groups or amine-functionalized groups. A substrate envisaged by the present invention is a non-porous substrate. Preferred non-porous substrates are glass, silicon, poly-L-lysine coated material, nitrocellulose, polystyrene, cyclic olefin copolymers (COCs), cyclic olefin polymers (COPs), polypropylene, polyethylene and polycarbonate.

Any suitable material known to the person skilled in the art may be used. Typically, glass or polystyrene is used. Polystyrene is a hydrophobic material suitable for binding negatively charged macromolecules because it normally contains few hydrophilic groups. For nucleic acids immobilized on glass slides, it is furthermore known that by increasing the hydrophobicity of the glass surface the nucleic acid immobilization may be increased. Such an enhancement may permit a relatively more densely packed formation. In addition to a coating or surface treatment with poly-L-lysine, the substrate, in particular glass, may be treated by silanation, e.g. with epoxy-silane or amino-silane or by silylation or by a treatment with polyacrylamide.

A number of standard arrays are commercially available and both the number and size of the features may be varied. In the present invention, the arrangement of the features may be altered to correspond to the size and/or density of the cells present in different tissues or organisms. For instance, animal cells typically have a cross-section in the region of 1-100 μm , whereas the cross-section of plant cells typically may range from 1-10000 μm . Hence, Nimblegen® arrays, which are available with up to 2.1 million features, or 4.2 million features, and feature sizes of 13 micrometers, may be preferred for tissue samples from an animal or fungus, whereas other formats, e.g. with 8x130 k features, may be sufficient for plant tissue samples. Commercial arrays are also available or known for use in the context of sequence analysis and in particular in the context of NGS technologies. Such arrays may also be used as the array surface in the context of the present invention e.g. an Illumina bead array. In addition to commercially available arrays, which can themselves be customized, it is possible to make custom or non-standard "in-house" arrays and methods for generating arrays are well-established. The methods of the invention may utilise both standard and non-standard arrays that comprise probes as defined below.

The probes on a microarray may be immobilized, i.e. attached or bound, to the array preferably via the 5' or 3' end, depending on the chemical matrix of the array. Typically, for commercially available arrays, the probes are attached via a 3' linkage, thereby leaving a free 5' end. However, arrays comprising probes attached to the substrate via a 5' linkage, thereby leaving a free 3' end, are available and may be synthesized using standard techniques that are well known in the art and are described elsewhere herein.

The covalent linkage used to couple a nucleic acid probe to an array substrate may be viewed as both a direct and indirect linkage, in that the although the probe is attached by a "direct" covalent bond, there may be a chemical moiety or linker separating the "first" nucleotide of the nucleic acid probe from the, e.g. glass or silicon, substrate i.e. an indirect linkage. For the purposes of the present invention probes that are immobilized to the substrate by a covalent bond and/or chemical linker are generally seen to be immobilized or attached directly to the substrate.

As will be described in more detail below, the capture probes of the invention may be immobilized on, or interact with, the array directly or indirectly. Thus the capture probes need not bind directly to the array, but may interact indirectly, for example by binding to a molecule which itself binds directly or indirectly to the array (e.g. the capture probe may interact with (e.g. bind or hybridize to) a binding partner for the capture probe, i.e. a surface probe, which is itself bound to the array directly or indirectly). Generally speaking, however, the capture probe will be, directly or indirectly (by one or more intermediaries), bound to, or immobilized on, the array.

The use, method and array of the invention may comprise probes that are immobilized via their 5' or 3' end. However, when the capture probe is immobilized directly to the array substrate, it may be immobilized only such that the 3' end of the capture probe is free to be extended, e.g. it is immobilized by its 5' end. The capture probe may be immobilized indirectly, such that it has a free, i.e. extendible, 3' end.

By extended or extendible 3' end, it is meant that further nucleotides may be added to the most 3' nucleotide of the nucleic acid molecule, e.g. capture probe, to extend the length of the nucleic acid molecule, i.e. the standard polymerization reaction utilized to extend nucleic acid molecules, e.g. templated polymerization catalyzed by a polymerase.

Thus, in one embodiment, the array comprises probes that are immobilized directly via their 3' end, so-called surface probes, which are defined below. Each species of surface probe comprises a region of complementarity to each species of capture probe, such that the capture probe may hybridize to the surface probe, resulting in the capture probe comprising a free extendible 3' end. In a preferred aspect of the invention, when the array comprises surface probes, the capture probes are synthesized in situ on the array.

The array probes may be made up of ribonucleotides and/or deoxyribonucleotides as well as synthetic nucleotide residues that are capable of participating in Watson-Crick type or analogous base pair interactions. Thus, the nucleic acid domain may be DNA or RNA or any modification thereof e.g. PNA or other derivatives containing non-nucleotide backbones. However, in the context of transcriptome analysis the capture domain of the capture probe must be capable of priming a reverse transcription reaction to generate cDNA that is complementary to the captured RNA molecules. As described below in more detail, in the context of genome analysis, the capture domain of the capture probe must be capable of binding to the DNA fragments, which may comprise binding to a binding domain that has been added to the fragmented DNA. In some embodiments, the capture domain of the capture probe may prime a DNA extension (polymerase) reaction to generate DNA that is complementary to the captured DNA molecules. In other embodiments, the capture domain may template a ligation reaction between the captured DNA molecules and a surface probe that is directly or indirectly immobilised on the substrate. In yet other embodiments, the capture domain may be ligated to one strand of the captured DNA molecules.

In a preferred embodiment of the invention at least the capture domain of the capture probe comprises or consists of deoxyribonucleotides (dNTPs). In a particularly preferred embodiment the whole of the capture probe comprises or consists of deoxyribonucleotides.

In a preferred embodiment of the invention the capture probes are immobilized on the substrate of the array directly, i.e. by their 5' end, resulting in a free extendible 3' end.

The capture probes of the invention comprise at least two domains, a capture domain and a positional domain (or a feature identification tag or domain; the positional domain may alternatively be defined as an identification (ID) domain or tag, or as a positional tag). The capture probe may further comprise a universal domain as defined further below. Where the capture probe is indirectly attached to the array surface via hybridization to a surface probe, the capture probe requires a sequence (e.g. a portion or domain) which is complementary to the surface probe. Such a complementary sequence may be complementary to a positional/identification domain and/or a universal domain on the surface probe. In other words the positional domain and/or universal domain may constitute the region or portion of the probe which is complementary to the surface probe. However, the capture probe may also comprise an additional domain (or region, portion or sequence) which is complementary to the surface probe. For ease of synthesis, as described in more detail below, such a surface probe-complementary region may be provided as part, or as an extension of the capture domain (such a part or extension not itself being used for, or capable of, binding to the target nucleic acid, e.g. RNA).

The capture domain is typically located at the 3' end of the capture probe and comprises a free 3' end that can be extended, e.g. by template dependent polymerization. The capture domain comprises a nucleotide sequence that is capable of hybridizing to nucleic acid, e.g. RNA (preferably mRNA) present in the cells of the tissue sample contact with the array.

Advantageously, the capture domain may be selected or designed to bind (or put more generally may be capable of binding) selectively or specifically to the particular nucleic acid, e.g. RNA it is desired to detect or analyse. For example the capture domain may be selected or designed for the selective capture of mRNA. As is well known in the art, this may be on the basis of hybridisation to the poly-A tail of mRNA. Thus, in a preferred embodiment the capture domain comprises a poly-T DNA oligonucleotide, i.e. a series of consecutive deoxythymidine residues linked by phosphodiester bonds, which is capable of hybridizing to the poly-A tail of mRNA. Alternatively, the capture domain may comprise nucleotides which are functionally or structurally analogous to poly-T i.e., are capable of binding selectively to poly-A, for example a poly-U oligonucleotide or an oligonucleotide comprised of deoxythymidine analogues, wherein said oligonucleotide retains the functional property of binding to poly-A. In a particularly preferred embodiment the capture domain, or more particularly the poly-T element of the capture domain, comprises at least 10 nucleotides, preferably at least 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 nucleotides. In a further embodiment, the capture domain, or more particularly the poly-T element of the capture domain comprises at least 25, 30 or 35 nucleotides.

Random sequences may also be used in the capture of nucleic acid, as is known in the art, e.g. random hexamers or similar sequences, and hence such random sequences may be used to form all or a part of the capture domain. For example, random sequences may be used in conjunction with poly-T (or poly-T analogue etc.) sequences. Thus

where a capture domain comprises a poly-T (or a "poly-T-like") oligonucleotide, it may also comprise a random oligonucleotide sequence. This may for example be located 5' or 3' of the poly-T sequence, e.g. at the 3' end of the capture probe, but the positioning of such a random sequence is not critical. Such a construct may facilitate the capturing of the initial part of the poly-A of mRNA. Alternatively, the capture domain may be an entirely random sequence. Degenerate capture domains may also be used, according to principles known in the art.

The capture domain may be capable of binding selectively to a desired sub-type or subset of nucleic acid, e.g. RNA, for example a particular type of RNA such mRNA or rRNA etc. as listed above, or to a particular subset of a given type of RNA, for example, a particular mRNA species e.g. corresponding to a particular gene or group of genes. Such a capture probe may be selected or designed based on sequence of the RNA it is desired to capture. Thus it may be a sequence-specific capture probe, specific for a particular RNA target or group of targets (target group etc). Thus, it may be based on a particular gene sequence or particular motif sequence or common/conserved sequence etc., according to principles well known in the art.

In embodiments where the capture probe is immobilized on the substrate of the array indirectly, e.g. via hybridization to a surface probe, the capture domain may further comprise an upstream sequence (5' to the sequence that hybridizes to the nucleic acid, e.g. RNA of the tissue sample) that is capable of hybridizing to 5' end of the surface probe. Alone, the capture domain of the capture probe may be seen as a capture domain oligonucleotide, which may be used in the synthesis of the capture probe in embodiments where the capture probe is immobilized on the array indirectly.

The positional domain (feature identification domain or tag) of the capture probe is located directly or indirectly upstream, i.e. closer to the 5' end of the capture probe nucleic acid molecule, of the capture domain. Preferably the positional domain is directly adjacent to the capture domain, i.e. there is no intermediate sequence between the capture domain and the positional domain. In some embodiments the positional domain forms the 5' end of the capture probe, which may be immobilized directly or indirectly on the substrate of the array.

As discussed above, each feature (distinct position) of the array comprises a spot of a species of nucleic acid probe, wherein the positional domain at each feature is unique. Thus, a "species" of capture probe is defined with reference to its positional domain; a single species of capture probe will have the same positional domain. However, it is not required that each member of a species of capture probe has the same sequence in its entirety. In particular, since the capture domain may be or may comprise a random or degenerate sequence, the capture domains of individual probes within a species may vary. Accordingly, in some embodiments where the capture domains of the capture probes are the same, each feature comprises a single probe sequence. However in other embodiments where the capture probe varies, members of a species of probe will not have the exact same sequence, although the sequence of the positional domain of each member in the species will be the same. What is required is that each feature or position of the array carries a capture probe of a single species (specifically each feature or position carries a capture probe which has an identical positional tag, i.e. there is a single positional domain at each feature or position). Each species has a different positional domain which identifies the species. However, each member of a species, may in some cases, as

described in more detail herein, have a different capture domain, as the capture domain may be random or degenerate or may have a random or degenerate component. This means that within a given feature, or position, the capture domain of the probes may differ.

Thus in some, but not necessarily in all embodiments, the nucleotide sequence of any one probe molecule immobilized at a particular feature is the same as the other probe molecules immobilized at the same feature, but the nucleotide sequence of the probes at each feature is different, distinct or distinguishable from the probes immobilized at every other feature. Preferably each feature comprises a different species of probe. However, in some embodiments it may be advantageous for a group of features to comprise the same species of probe, i.e. effectively to produce a feature covering an area of the array that is greater than a single feature, e.g. to lower the resolution of the array. In other embodiments of the array, the nucleotide sequence of the positional domain of any one probe molecule immobilized at a particular feature may be the same as the other probe molecules immobilized at the same feature but the capture domain may vary. The capture domain may nonetheless be designed to capture the same type of molecule, e.g. mRNA in general.

The positional domain (or tag) of the capture probe comprises the sequence which is unique to each feature and acts as a positional or spatial marker (the identification tag). In this way each region or domain of the tissue sample, e.g. each cell in the tissue, will be identifiable by spatial resolution across the array linking the nucleic acid, e.g. RNA (e.g. the transcripts) from a certain cell to a unique positional domain sequence in the capture probe. By virtue of the positional domain a capture probe in the array may be correlated to a position in the tissue sample, for example it may be correlated to a cell in the sample. Thus, the positional domain of the capture domain may be seen as a nucleic acid tag (identification tag).

Any suitable sequence may be used as the positional domain in the capture probes of the invention. By a suitable sequence, it is meant that the positional domain should not interfere with (i.e. inhibit or distort) the interaction between the RNA of the tissue sample and the capture domain of the capture probe. For example, the positional domain should be designed such that nucleic acid molecules in the tissue sample do not hybridize specifically to the positional domain. Preferably, the nucleic acid sequence of the positional domain of the capture probes has less than 80% sequence identity to the nucleic acid sequences in the tissue sample. Preferably, the positional domain of the capture probe has less than 70%, 60%, 50% or less than 40% sequence identity across a substantial part of the nucleic acids molecules in the tissue sample. Sequence identity may be determined by any appropriate method known in the art, e.g. the using BLAST alignment algorithm.

In a preferred embodiment the positional domain of each species of capture probe contains a unique barcode sequence. The barcode sequences may be generated using random sequence generation. The randomly generated sequences may be followed by stringent filtering by mapping to the genomes of all common reference species and with pre-set Tm intervals, GC content and a defined distance of difference to the other barcode sequences to ensure that the barcode sequences will not interfere with the capture of the nucleic acid, e.g. RNA from the tissue sample and will be distinguishable from each other without difficulty.

As mentioned above, and in a preferred embodiment, the capture probe comprises also a universal domain (or linker

domain or tag). The universal domain of the capture probe is located directly or indirectly upstream, i.e. closer to the 5' end of the capture probe nucleic acid molecule, of the positional domain. Preferably the universal domain is directly adjacent to the positional domain, i.e. there is no intermediate sequence between the positional domain and the universal domain. In embodiments where the capture probe comprises a universal domain, the domain will form the 5' end of the capture probe, which may be immobilized directly or indirectly on the substrate of the array.

The universal domain may be utilized in a number of ways in the methods and uses of the invention. For example, the methods of the invention comprise a step of releasing (e.g. removing) at least part of the synthesised (i.e. extended or ligated) nucleic acid, e.g. cDNA molecules from the surface of the array. As described elsewhere herein, this may be achieved in a number of ways, of which one comprises cleaving the nucleic acid, e.g. cDNA molecule from the surface of the array. Thus, the universal domain may itself comprise a cleavage domain, i.e. a sequence that can be cleaved specifically, either chemically or preferably enzymatically.

Thus, the cleavage domain may comprise a sequence that is recognised by one or more enzymes capable of cleaving a nucleic acid molecule, i.e. capable of breaking the phosphodiester linkage between two or more nucleotides. For instance, the cleavage domain may comprise a restriction endonuclease (restriction enzyme) recognition sequence. Restriction enzymes cut double-stranded or single stranded DNA at specific recognition nucleotide sequences known as restriction sites and suitable enzymes are well known in the art. For example, it is particularly advantageous to use rare-cutting restriction enzymes, i.e. enzymes with a long recognition site (at least 8 base pairs in length), to reduce the possibility of cleaving elsewhere in the nucleic acid, e.g. cDNA molecule. In this respect, it will be seen that removing or releasing at least part of the nucleic acid, e.g. cDNA molecule requires releasing a part comprising the positional domain of the nucleic acid, e.g. cDNA and all of the sequence downstream of the domain, i.e. all of the sequence that is 3' to the positional domain. Hence, cleavage of the nucleic acid, e.g. cDNA molecule should take place 5' to the positional domain.

By way of example, the cleavage domain may comprise a poly-U sequence which may be cleaved by a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase Endonuclease VIII, commercially known as the USER™ enzyme.

A further example of a cleavage domain can be utilised in embodiments where the capture probe is immobilized to the array substrate indirectly, i.e. via a surface probe. The cleavage domain may comprise one or more mismatch nucleotides, i.e. when the complementary parts of the surface probe and the capture probe are not 100% complementary. Such a mismatch is recognised, e.g. by the MutY and T7 endonuclease I enzymes, which results in cleavage of the nucleic acid molecule at the position of the mismatch.

In some embodiments of the invention, the positional domain of the capture probe comprises a cleavage domain, wherein the said cleavage domain is located at the 5' end of the positional domain.

The universal domain may comprise also an amplification domain. This may be in addition to, or instead of, a cleavage domain. In some embodiments of the invention, as described elsewhere herein, it may be advantageous to amplify the nucleic acid, e.g. cDNA molecules, for example after they have been released (e.g. removed or cleaved) from the array

substrate. It will be appreciated however, that the initial cycle of amplification, or indeed any or all further cycles of amplification may also take place in situ on the array. The amplification domain comprises a distinct sequence to which an amplification primer may hybridize. The amplification domain of the universal domain of the capture probe is preferably identical for each species of capture probe. Hence a single amplification reaction will be sufficient to amplify all of the nucleic acid, e.g. cDNA molecules (which may or may not be released from the array substrate prior to amplification).

Any suitable sequence may be used as the amplification domain in the capture probes of the invention. By a suitable sequence, it is meant that the amplification domain should not interfere with (i.e. inhibit or distort) the interaction between the nucleic acid, e.g. RNA of the tissue sample and the capture domain of the capture probe. Furthermore, the amplification domain should comprise a sequence that is not the same or substantially the same as any sequence in the nucleic acid, e.g. RNA of the tissue sample, such that the primer used in the amplification reaction can hybridize only to the amplification domain under the amplification conditions of the reaction.

For example, the amplification domain should be designed such that nucleic acid molecules in the tissue sample do not hybridize specifically to the amplification domain or the complementary sequence of the amplification domain. Preferably, the nucleic acid sequence of the amplification domain of the capture probes and the complement thereof has less than 80% sequence identity to the nucleic acid sequences in the tissue sample. Preferably, the positional domain of the capture probe has less than 70%, 60%, 50% or less than 40% sequence identity across a substantial part of the nucleic acid molecules in the tissue sample. Sequence identity may be determined by any appropriate method known in the art, e.g. the using BLAST alignment algorithm.

Thus, alone, the universal domain of the capture probe may be seen as a universal domain oligonucleotide, which may be used in the synthesis of the capture probe in embodiments where the capture probe is immobilized on the array indirectly.

In one representative embodiment of the invention only the positional domain of each species of capture probe is unique. Hence, the capture domains and universal domains (if present) are in one embodiment the same for every species of capture probe for any particular array to ensure that the capture of the nucleic acid, e.g. RNA from the tissue sample is uniform across the array. However, as discussed above, in some embodiments the capture domains may differ by virtue of including random or degenerate sequences.

In embodiments where the capture probe is immobilized on the substrate of the array indirectly, e.g. via hybridisation to a surface probe, the capture probe may be synthesised on the array as described below.

The surface probes are immobilized on the substrate of the array directly by or at, e.g. their 3' end. Each species of surface probe is unique to each feature (distinct position) of the array and is partly complementary to the capture probe, defined above.

Hence the surface probe comprises at its 5' end a domain (complementary capture domain) that is complementary to a part of the capture domain that does not bind to the nucleic acid, e.g. RNA of the tissue sample. In other words, it comprises a domain that can hybridize to at least part of a capture domain oligonucleotide. The surface probe further comprises a domain (complementary positional domain or

complementary feature identification domain) that is complementary to the positional domain of the capture probe. The complementary positional domain is located directly or indirectly downstream (i.e. at the 3' end) of the complementary capture domain, i.e. there may be an intermediary or linker sequence separating the complementary positional domain and the complementary capture domain. In embodiments where the capture probe is synthesized on the array surface, the surface probes of the array always comprise a domain (complementary universal domain) at the 3' end of the surface probe, i.e. directly or indirectly downstream of the positional domain, which is complementary to the universal domain of the capture probe. In other words, it comprises a domain that can hybridize to at least part of the universal domain oligonucleotide.

In some embodiments of the invention the sequence of the surface probe shows 100% complementarity or sequence identity to the positional and universal domains and to the part of the capture domain that does not bind to the nucleic acid, e.g. RNA of the tissue sample. In other embodiments the sequence of the surface probe may show less than 100% sequence identity to the domains of the capture probe, e.g. less than 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91% or 90%. In a particularly preferred embodiment of the invention, the complementary universal domain shares less than 100% sequence identity to the universal domain of the capture probe.

In one embodiment of the invention, the capture probe is synthesized or generated on the substrate of the array. In a representative embodiment (see FIG. 3), the array comprises surface probes as defined above. Oligonucleotides that correspond to the capture domain and universal domain of the capture probe are contacted with the array and allowed to hybridize to the complementary domains of the surface probes. Excess oligonucleotides may be removed by washing the array under standard hybridization conditions. The resultant array comprises partially single stranded probes, wherein both the 5' and 3' ends of the surface probe are double stranded and the complementary positional domain is single stranded. The array may be treated with a polymerase enzyme to extend the 3' end of the universal domain oligonucleotide, in a template dependent manner, so as to synthesize the positional domain of the capture probe. The 3' end of the synthesized positional domain is then ligated, e.g. using a ligase enzyme, to the 5' end of the capture domain oligonucleotide to generate the capture probe. It will be understood in this regard that the 5' end of the capture domain oligonucleotide is phosphorylated to enable ligation to take place. As each species of surface probe comprises a unique complementary positional domain, each species of capture probe will comprise a unique positional domain.

The term "hybridisation" or "hybridises" as used herein refers to the formation of a duplex between nucleotide sequences which are sufficiently complementary to form duplexes via Watson-Crick base pairing. Two nucleotide sequences are "complementary" to one another when those molecules share base pair organization homology. "Complementary" nucleotide sequences will combine with specificity to form a stable duplex under appropriate hybridization conditions. For instance, two sequences are complementary when a section of a first sequence can bind to a section of a second sequence in an anti-parallel sense wherein the 3'-end of each sequence binds to the 5'-end of the other sequence and each A, T(U), G and C of one sequence is then aligned with a T(U), A, C and G, respectively, of the other sequence. RNA sequences can also include complementary G=U or U=G base pairs. Thus, two sequences need not have perfect

homology to be “complementary” under the invention. Usually two sequences are sufficiently complementary when at least about 90% (preferably at least about 95%) of the nucleotides share base pair organization over a defined length of the molecule. The domains of the capture and surface probes thus contain a region of complementarity. Furthermore the capture domain of the capture probe contains a region of complementarity for the nucleic acid, e.g. RNA (preferably mRNA) of the tissue sample.

The capture probe may also be synthesised on the array substrate using polymerase extension (similarly to as described above) and a terminal transferase enzyme to add a “tail” which may constitute the capture domain. This is described further in Example 7 below. The use of terminal transferases to add nucleotide sequences to the end of an oligonucleotide is known in the art, e.g. to introduce a homopolymeric tail e.g. a poly-T tail. Accordingly, in such a synthesis an oligonucleotide that corresponds to the universal domain of the capture probe may be contacted with the array and allowed to hybridize to the complementary domain of the surface probes. Excess oligonucleotides may be removed by washing the array under standard hybridization conditions. The resultant array comprises partially single stranded probes, wherein the 5' ends of the surface probes are double stranded and the complementary positional domain is single stranded. The array may be treated with a polymerase enzyme to extend the 3' end of the universal domain oligonucleotide, in a template dependent manner, so as to synthesize the positional domain of the capture probe. The capture domain, e.g. comprising a poly-T sequence may then be introduced using a terminal transferase to add a poly-T tail to generate the capture probe.

The typical array of, and for use in the methods of, the invention may contain multiple spots, or “features”. A feature may be defined as an area or distinct position on the array substrate at which a single species of capture probe is immobilized. Hence each feature will comprise a multiplicity of probe molecules, of the same species. It will be understood in this context that whilst it is encompassed that each capture probe of the same species may have the same sequence, this need not necessarily be the case. Each species of capture probe will have the same positional domain (i.e. each member of a species and hence each probe in a feature will be identically “tagged”), but the sequence of each member of the feature (species) may differ, because the sequence of a capture domain may differ. As described above, random or degenerate capture domains may be used. Thus the capture probes within a feature may comprise different random or degenerate sequences. The number and density of the features on the array will determine the resolution of the array, i.e. the level of detail at which the transcriptome or genome of the tissue sample can be analysed. Hence, a higher density of features will typically increase the resolution of the array.

As discussed above, the size and number of the features on the array of the invention will depend on the nature of the tissue sample and required resolution. Thus, if it is desirable to determine a transcriptome or genome only for regions of cells within a tissue sample (or the sample contains large cells) then the number and/or density of features on the array may be reduced (i.e. lower than the possible maximum number of features) and/or the size of the features may be increased (i.e. the area of each feature may be greater than the smallest possible feature), e.g. an array comprising few large features. Alternatively, if it is desirable to determine a transcriptome or genome of individual cells within a sample, it may be necessary to use the maximum number of features

possible, which would necessitate using the smallest possible feature size, e.g. an array comprising many small features.

Whilst single cell resolution may be a preferred and advantageous feature of the present invention, it is not essential to achieve this, and resolution at the cell group level is also of interest, for example to detect or distinguish a particular cell type or tissue region, e.g. normal vs tumour cells.

In representative embodiments of the invention, an array may contain at least 2, 5, 10, 50, 100, 500, 750, 1000, 1500, 3000, 5000, 10000, 20000, 40000, 50000, 75000, 100000, 150000, 200000, 300000, 400000, 500000, 750000, 800000, 1000000, 1200000, 1500000, 1750000, 2000000, 2100000, 3000000, 3500000, 4000000 or 4200000 features. Whilst 4200000 represents the maximum number of features presently available on a commercial array, it is envisaged that arrays with features in excess of this may be prepared and such arrays are of interest in the present invention. As noted above, feature size may be decreased and this may allow greater numbers of features to be accommodated within the same or a similar area. By way of example, these features may be comprised in an area of less than about 20 cm², 10 cm², 5 cm², 1 cm², 1 mm², or 100 μm².

Thus, in some embodiments of the invention the area of each feature may be from about 1 μm², 2 μm², 3 μm², 4 μm², 5 μm², 10 μm², 12 μm², 15 μm², 20 μm², 50 μm², 75 μm², 100 μm², 150 μm², 200 μm², 250 μm², 300 μm², 400 μm², or 500 μm².

It will be evident that a tissue sample from any organism could be used in the methods of the invention, e.g. plant, animal or fungal. The array of the invention allows the capture of any nucleic acid, e.g. mRNA molecules, which are present in cells that are capable of transcription and/or translation. The arrays and methods of the invention are particularly suitable for isolating and analysing the transcriptome or genome of cells within a sample, wherein spatial resolution of the transcriptomes or genomes is desirable, e.g. where the cells are interconnected or in contact directly with adjacent cells. However, it will be apparent to a person of skill in the art that the methods of the invention may also be useful for the analysis of the transcriptome or genome of different cells or cell types within a sample even if said cells do not interact directly, e.g. a blood sample. In other words, the cells do not need to present in the context of a tissue and can be applied to the array as single cells (e.g. cells isolated from a non-fixed tissue). Such single cells, whilst not necessarily fixed to a certain position in a tissue, are nonetheless applied to a certain position on the array and can be individually identified. Thus, in the context of analysing cells that do not interact directly, or are not present in a tissue context, the spatial properties of the described methods may be applied to obtaining or retrieving unique or independent transcriptome or genome information from individual cells.

The sample may thus be a harvested or biopsied tissue sample, or possibly a cultured sample. Representative samples include clinical samples e.g. whole blood or blood-derived products, blood cells, tissues, biopsies, or cultured tissues or cells etc. including cell suspensions. Artificial tissues may for example be prepared from cell suspension (including for example blood cells). Cells may be captured in a matrix (for example a gel matrix e.g. agar, agarose, etc) and may then be sectioned in a conventional way. Such procedures are known in the art in the context of immunohistochemistry (see e.g. Andersson et al 2006, *J. Histochem. Cytochem.* 54(12): 1413-23. Epub 2006 Sep. 6).

The mode of tissue preparation and how the resulting sample is handled may effect the transcriptomic or genomic analysis of the methods of the invention. Moreover, various tissue samples will have different physical characteristics and it is well within the skill of a person in the art to perform the necessary manipulations to yield a tissue sample for use with the methods of the invention. However, it is evident from the disclosures herein that any method of sample preparation may be used to obtain a tissue sample that is suitable for use in the methods of the invention. For instance any layer of cells with a thickness of approximately 1 cell or less may be used in the methods of the invention. In one embodiment, the thickness of the tissue sample may be less than 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2 or 0.1 of the cross-section of a cell. However, since as noted above, the present invention is not limited to single cell resolution and hence it is not a requirement that the tissue sample has a thickness of one cell diameter or less; thicker tissue samples may if desired be used. For example cryostat sections may be used, which may be e.g. 10-20 μm thick.

The tissue sample may be prepared in any convenient or desired way and the invention is not restricted to any particular type of tissue preparation. Fresh, frozen, fixed or unfixed tissues may be used. Any desired convenient procedure may be used for fixing or embedding the tissue sample, as described and known in the art. Thus any known fixatives or embedding materials may be used.

As a first representative example of a tissue sample for use in the invention, the tissue may prepared by deep freezing at temperature suitable to maintain or preserve the integrity (i.e. the physical characteristics) of the tissue structure, e.g. less than -20°C . and preferably less than -25 , -30 , -40 , -50 , -60 , -70 or -80°C . The frozen tissue sample may be sectioned, i.e. thinly sliced, onto the array surface by any suitable means. For example, the tissue sample may be prepared using a chilled microtome, a cryostat, set at a temperature suitable to maintain both the structural integrity of the tissue sample and the chemical properties of the nucleic acids in the sample, e.g. to less than -15°C . and preferably less than -20 or -25°C . Thus, the sample should be treated so as to minimize the degeneration or degradation of the nucleic acid, e.g. RNA in the tissue. Such conditions are well-established in the art and the extent of any degradation may be monitored through nucleic acid extraction, e.g. total RNA extraction and subsequent quality analysis at various stages of the preparation of the tissue sample.

In a second representative example, the tissue may be prepared using standard methods of formalin-fixation and paraffin-embedding (FFPE), which are well-established in the art. Following fixation of the tissue sample and embedding in a paraffin or resin block, the tissue samples may sectioned, i.e. thinly sliced, onto the array. As noted above, other fixatives and/or embedding materials can be used.

It will be apparent that the tissue sample section will need to be treated to remove the embedding material e.g. to deparaffinize, i.e. to remove the paraffin or resin, from the sample prior to carrying out the methods of the invention. This may be achieved by any suitable method and the removal of paraffin or resin or other material from tissue samples is well established in the art, e.g. by incubating the sample (on the surface of the array) in an appropriate solvent e.g. xylene, e.g. twice for 10 minutes, followed by an ethanol rinse, e.g. 99.5% ethanol for 2 minutes, 96% ethanol for 2 minutes, and 70% ethanol for 2 minutes.

It will be evident to the skilled person that the RNA in tissue sections prepared using methods of FFPE or other methods of fixing and embedding is more likely to be

partially degraded than in the case of frozen tissue. However, without wishing to be bound by any particular theory, it is believed that this may be advantageous in the methods of the invention. For instance, if the RNA in the sample is partially degraded the average length of the RNA polynucleotides will be less and more randomized than a non-degraded sample. It is postulated therefore that partially degraded RNA would result in less bias in the various processing steps, described elsewhere herein, e.g. ligation of adaptors (amplification domains), amplification of the cDNA molecules and sequencing thereof.

Hence, in one embodiment of the invention the tissue sample, i.e. the section of the tissue sample contacted with the array, is prepared using FFPE or other methods of fixing and embedding. In other words the sample may be fixed, e.g. fixed and embedded. In an alternative embodiment of the invention the tissue sample is prepared by deep-freezing. In another embodiment a touch imprint of a tissue may be used, according to procedures known in the art. In other embodiments an unfixed sample may be used.

The thickness of the tissue sample section for use in the methods of the invention may be dependent on the method used to prepare the sample and the physical characteristics of the tissue. Thus, any suitable section thickness may be used in the methods of the invention. In representative embodiments of the invention the thickness of the tissue sample section will be at least 0.1 μm , further preferably at least 0.2, 0.3, 0.4, 0.5, 0.7, 1.0, 1.5, 2, 3, 4, 5, 6, 7, 8, 9 or 10 μm . In other embodiments the thickness of the tissue sample section is at least 10, 12, 13, 14, 15, 20, 30, 40 or 50 μm . However, the thickness is not critical and these are representative values only. Thicker samples may be used if desired or convenient e.g. 70 or 100 μm or more. Typically, the thickness of the tissue sample section is between 1-100 μm , 1-50 μm , 1-30 μm , 1-25 μm , 1-20 μm , 1-15 μm , 1-10 μm , 2-8 μm , 3-7 μm or 4-6 μm , but as mentioned above thicker samples may be used.

On contact of the tissue sample section with the array, e.g. following removal of the embedding material e.g. deparaffinization, the nucleic acid, e.g. RNA molecules in the tissue sample will bind to the immobilized capture probes on the array. In some embodiments it may be advantageous to facilitate the hybridization of the nucleic acid, e.g. RNA molecules to the capture probes. Typically, facilitating the hybridization comprises modifying the conditions under which hybridization occurs. The primary conditions that can be modified are the time and temperature of the incubation of the tissue section on the array prior to the reverse transcription step, which is described elsewhere herein.

For instance, on contacting the tissue sample section with the array, the array may be incubated for at least 1 hour to allow the nucleic acid, e.g. RNA to hybridize to the capture probes. Preferably the array may be incubated for at least 2, 3, 5, 10, 12, 15, 20, 22 or 24 hours or until the tissue sample section has dried. The array incubation time is not critical and any convenient or desired time may be used. Typical array incubations may be up to 72 hours. Thus, the incubation may occur at any suitable temperature, for instance at room temperature, although in a preferred embodiment the tissue sample section is incubated on the array at a temperature of at least 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36 or 37°C . Incubation temperatures of up to 55°C . are commonplace in the art. In a particularly preferred embodiment the tissue sample section is allowed to dry on the array at 37°C . for 24 hours. Once the tissue sample section has dried the array may be stored at room temperature before performing the reverse transcription step. It will be under-

stood that the if the tissue sample section is allowed to dry on the surface of the array, it will need to be rehydrated before further manipulation of the captured nucleic acid can be achieved, e.g. the step of reverse transcribing the captured RNA.

Hence, the method of the invention may comprise a further step of rehydrating the tissue sample after contacting the sample with the array.

In some embodiments it may be advantageous to block (e.g. mask or modify) the capture probes prior to contacting the tissue sample with the array, particularly when the nucleic acid in the tissue sample is subject to a process of modification prior to its capture on the array. Specifically, it may be advantageous to block or modify the free 3' end of the capture probe. In a particular embodiment, the nucleic acid in the tissue sample, e.g. fragmented genomic DNA, may be modified such that it can be captured by the capture probe. For instance, and as described in more detail below, an adaptor sequence (comprising a binding domain capable of binding to the capture domain of the capture probe) may be added to the end of the nucleic acid, e.g. fragmented genomic DNA. This may be achieved by, e.g. ligation of an adaptor or extension of the nucleic acid, e.g. using an enzyme to incorporate additional nucleotides at the end of the sequence, e.g. a poly-A tail. It is necessary to block or modify the capture probes, particularly the free 3' end of the capture probe, prior to contacting the tissue sample with the array to avoid modification of the capture probes, e.g. to avoid the addition of a poly-A tail to the free 3' end of the capture probes. Preferably the incorporation of a blocking domain may be incorporated into the capture probe when it is synthesised. However, the blocking domain may be incorporated to the capture probe after its synthesis.

In some embodiments the capture probes may be blocked by any suitable and reversible means that would prevent modification of the capture domains during the process of modifying the nucleic acid of the tissue sample, which occurs after the tissue sample has been contacted with the array. In other words, the capture probes may be reversibly masked or modified such that the capture domain of the capture probe does not comprise a free 3' end, i.e. such that the 3' end is removed or modified, or made inaccessible so that the capture probe is not susceptible to the process which is used to modify the nucleic acid of the tissue sample, e.g. ligation or extension, or the additional nucleotides may be removed to reveal and/or restore the 3' end of the capture domain of the capture probe.

For example, blocking probes may be hybridised to the capture probes to mask the free 3' end of the capture domain, e.g. hairpin probes or partially double stranded probes, suitable examples of which are known in the art. The free 3' end of the capture domain may be blocked by chemical modification, e.g. addition of an azidomethyl group as a chemically reversible capping moiety such that the capture probes do not comprise a free 3' end. Suitable alternative capping moieties are well known in the art, e.g. the terminal nucleotide of the capture domain could be a reversible terminator nucleotide, which could be included in the capture probe during or after probe synthesis.

Alternatively or additionally, the capture domain of the capture probe could be modified so as to allow the removal of any modifications of the capture probe, e.g. additional nucleotides, that occur when the nucleic acid molecules of the tissue sample are modified. For instance, the capture probes may comprise an additional sequence downstream of the capture domain, i.e. 3' to capture domain, namely a blocking domain. This could be in the form of, e.g. a

restriction endonuclease recognition sequence or a sequence of nucleotides cleavable by specific enzyme activities, e.g. uracil. Following the modification of the nucleic acid of the tissue sample, the capture probes could be subjected to an enzymatic cleavage, which would allow the removal of the blocking domain and any of the additional nucleotides that are added to the 3' end of the capture probe during the modification process. The removal of the blocking domain would reveal and/or restore the free 3' end of the capture domain of the capture probe. The blocking domain could be synthesised as part of the capture probe or could be added to the capture probe in situ (i.e. as a modification of an existing array), e.g. by ligation of the blocking domain.

The capture probes may be blocked using any combination of the blocking mechanisms described above.

Once the nucleic acid of the tissue sample, e.g. fragmented genomic DNA, has been modified to enable it to hybridise to the capture domain of the capture probe, the capture probe must be unblocked, e.g. by dissociation of the blocking oligonucleotide, removal of the capping moiety and/or blocking domain.

In order to correlate the sequence analysis or transcriptome or genome information obtained from each feature of the array with the region (i.e. an area or cell) of the tissue sample the tissue sample is oriented in relation to the features on the array. In other words, the tissue sample is placed on the array such that the position of a capture probe on the array may be correlated with a position in the tissue sample. Thus it may be identified where in the tissue sample the position of each species of capture probe (or each feature of the array) corresponds. In other words, it may be identified to which location in the tissue sample the position of each species of capture probe corresponds. This may be done by virtue of positional markers present on the array, as described below. Conveniently, but not necessarily, the tissue sample may be imaged following its contact with the array. This may be performed before or after the nucleic acid of the tissue sample is processed, e.g. before or after the cDNA generation step of the method, in particular the step of generating the first strand cDNA by reverse transcription. In a preferred embodiment the tissue sample is imaged prior to the release of the captured and synthesised (i.e. extended or ligated) DNA, e.g. cDNA, from the array. In a particularly preferred embodiment the tissue is imaged after the nucleic acid of the tissue sample has been processed, e.g. after the reverse transcription step, and any residual tissue is removed (e.g. washed) from the array prior to the release of molecules, e.g. of the cDNA from the array. In some embodiments, the step of processing the captured nucleic acid, e.g. the reverse transcription step, may act to remove residual tissue from the array surface, e.g. when using tissue preparing by deep-freezing. In such a case, imaging of the tissue sample may take place prior to the processing step, e.g. the cDNA synthesis step. Generally speaking, imaging may take place at any time after contacting the tissue sample with the area, but before any step which degrades or removes the tissue sample. As noted above, this may depend on the tissue sample.

Advantageously, the array may comprise markers to facilitate the orientation of the tissue sample or the image thereof in relation to the features of the array. Any suitable means for marking the array may be used such that they are detectable when the tissue sample is imaged. For instance, a molecule, e.g. a fluorescent molecule, that generates a signal, preferably a visible signal, may be immobilized directly or indirectly on the surface of the array. Preferably, the array comprises at least two markers in distinct positions

on the surface of the array, further preferably at least 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 20, 30, 40, 50, 60, 70, 80, 90 or 100 markers. Conveniently several hundred or even several thousand markers may be used. The markers may be provided in a pattern, for example make up an outer edge of the array, e.g. an entire outer row of the features of an array. Other informative patterns may be used, e.g. lines sectioning the array. This may facilitate aligning an image of the tissue sample to an array, or indeed generally in correlating the features of the array to the tissue sample. Thus, the marker may be an immobilized molecule to which a signal giving molecule may interact to generate a signal. In a representative example, the array may comprise a marker feature, e.g. a nucleic acid probe immobilized on the substrate of array, to which a labelled nucleic acid may hybridize. For instance, the labelled nucleic acid molecule, or marker nucleic acid, may be linked or coupled to a chemical moiety capable of fluorescing when subjected to light of a specific wavelength (or range of wavelengths), i.e. excited. Such a marker nucleic acid molecule may be contacted with the array before, contemporaneously with or after the tissue sample is stained in order to visualize or image the tissue sample. However, the marker must be detectable when the tissue sample is imaged. Thus, in a preferred embodiment the marker may be detected using the same imaging conditions used to visualize the tissue sample.

In a particularly preferred embodiment of the invention, the array comprises marker features to which a labelled, preferably fluorescently labelled, marker nucleic acid molecule, e.g. oligonucleotide, is hybridized.

The step of imaging the tissue may use any convenient histological means known in the art, e.g. light, bright field, dark field, phase contrast, fluorescence, reflection, interference, confocal microscopy or a combination thereof. Typically the tissue sample is stained prior to visualization to provide contrast between the different regions, e.g. cells, of the tissue sample. The type of stain used will be dependent on the type of tissue and the region of the cells to be stained. Such staining protocols are known in the art. In some embodiments more than one stain may be used to visualize (image) different aspects of the tissue sample, e.g. different regions of the tissue sample, specific cell structures (e.g. organelles) or different cell types. In other embodiments, the tissue sample may be visualized or imaged without staining the sample, e.g. if the tissue sample contains already pigments that provide sufficient contrast or if particular forms of microscopy are used.

In a preferred embodiment, the tissue sample is visualized or imaged using fluorescence microscopy.

The tissue sample, i.e. any residual tissue that remains in contact with the array substrate following the reverse transcription step and optionally imaging, if imaging is desired and was not carried out before reverse transcription, preferably is removed prior to the step of releasing the cDNA molecules from the array. Thus, the methods of the invention may comprise a step of washing the array. Removal of the residual tissue sample may be performed using any suitable means and will be dependent on the tissue sample. In the simplest embodiment, the array may be washed with water. The water may contain various additives, e.g. surfactants (e.g. detergents), enzymes etc to facilitate to removal of the tissue. In some embodiments, the array is washed with a solution comprising a proteinase enzyme (and suitable buffer) e.g. proteinase K. In other embodiments, the solution may comprise also or alternatively cellulase, hemicellulase or chitinase enzymes, e.g. if the tissue sample is from a plant or fungal source. In further embodiments, the temperature of

the solution used to wash the array may be, e.g. at least 30° C., preferably at least 35, 40, 45, 50 or 55° C. It will be evident that the wash solution should minimize the disruption of the immobilized nucleic acid molecules. For instance, in some embodiments the nucleic acid molecules may be immobilized on the substrate of the array indirectly, e.g. via hybridization of the capture probe and the RNA and/or the capture probe and the surface probe, thus the wash step should not interfere with the interaction between the molecules immobilized on the array, i.e. should not cause the nucleic acid molecules to be denatured.

Following the step of contacting the array with a tissue sample, under conditions sufficient to allow hybridization to occur between the nucleic acid, e.g. RNA (preferably mRNA), of the tissue sample to the capture probes, the step of securing (acquiring) the hybridized nucleic acid takes place. Securing or acquiring the captured nucleic acid involves a covalent attachment of a complementary strand of the hybridized nucleic acid to the capture probe (i.e. via a nucleotide bond, a phosphodiester bond between juxtaposed 3'-hydroxyl and 5'-phosphate termini of two immediately adjacent nucleotides), thereby tagging or marking the captured nucleic acid with the positional domain specific to the feature on which the nucleic acid is captured.

In some embodiments, securing the hybridized nucleic acid, e.g. a single stranded nucleic acid, may involve extending the capture probe to produce a copy of the captured nucleic acid, e.g. generating cDNA from the captured (hybridized) RNA. It will be understood that this refers to the synthesis of a complementary strand of the hybridized nucleic acid, e.g. generating cDNA based on the captured RNA template (the RNA hybridized to the capture domain of the capture probe). Thus, in an initial step of extending the capture probe, e.g. the cDNA generation, the captured (hybridized) nucleic acid, e.g. RNA acts as a template for the extension, e.g. reverse transcription, step. In other embodiments, as described below, securing the hybridized nucleic acid, e.g. partially double stranded DNA, may involve covalently coupling the hybridized nucleic acid, e.g. fragmented DNA, to the capture probe, e.g. ligating to the capture probe the complementary strand of the nucleic acid hybridized to the capture probe, in a ligation reaction.

Reverse transcription concerns the step of synthesizing cDNA (complementary or copy DNA) from RNA, preferably mRNA (messenger RNA), by reverse transcriptase. Thus cDNA can be considered to be a copy of the RNA present in a cell at the time at which the tissue sample was taken, i.e. it represents all or some of the genes that were expressed in said cell at the time of isolation.

The capture probe, specifically the capture domain of the capture probe, acts as a primer for producing the complementary strand of the nucleic acid hybridized to the capture probe, e.g. a primer for reverse transcription. Hence, the nucleic acid, e.g. cDNA, molecules generated by the extension reaction, e.g. reverse transcription reaction, incorporate the sequence of the capture probe, i.e. the extension reaction, e.g. reverse transcription reaction, may be seen as a way of labelling indirectly the nucleic acid, e.g. transcripts, of the tissue sample that are in contact with each feature of the array. As mentioned above, each species of capture probe comprises a positional domain (feature identification tag) that represents a unique sequence for each feature of the array. Thus, all of the nucleic acid, e.g. cDNA, molecules synthesized at a specific feature will comprise the same nucleic acid "tag".

The nucleic acid, e.g. cDNA, molecules synthesized at each feature of the array may represent the genome of, or

genes expressed from, the region or area of the tissue sample in contact with that feature, e.g. a tissue or cell type or group or sub-group thereof, and may further represent genes expressed under specific conditions, e.g. at a particular time, in a specific environment, at a stage of development or in response to stimulus etc. Hence, the cDNA at any single feature may represent the genes expressed in a single cell, or if the feature is in contact with the sample at a cell junction, the cDNA may represent the genes expressed in more than one cell. Similarly, if a single cell is in contact with multiple features, then each feature may represent a proportion of the genes expressed in said cell. Similarly, in embodiments in which the captured nucleic acid is DNA, any single feature may be representative of the genome of a single cell or more than one cell. Alternatively, the genome of a single cell may be represented by multiple features.

The step of extending the capture probe, e.g. reverse transcription, may be performed using any suitable enzymes and protocol of which many exist in the art, as described in detail below. However, it will be evident that it is not necessary to provide a primer for the synthesis of the first nucleic acid, e.g. cDNA, strand because the capture domain of the capture probe acts as the primer, e.g. reverse transcription primer.

Preferably, in the context of the present invention the secured nucleic acid (i.e. the nucleic acid covalently attached to the capture probe), e.g. cDNA is treated to comprise double stranded DNA. However, in some embodiments, the captured DNA may already comprise double stranded DNA, e.g. where partially double stranded fragmented DNA is ligated to the capture probe. Treatment of the captured nucleic acid to produce double stranded DNA may be achieved in a single reaction to generate only a second DNA, e.g. cDNA, strand, i.e. to produce double stranded DNA molecules without increasing the number of double stranded DNA molecules, or in an amplification reaction to generate multiple copies of the second strand, which may be in the form of single stranded DNA (e.g. linear amplification) or double stranded DNA, e.g. cDNA (e.g. exponential amplification).

The step of second strand DNA, e.g. cDNA, synthesis may take place in situ on the array, either as a discrete step of second strand synthesis, for example using random primers as described in more detail below, or in the initial step of an amplification reaction. Alternatively, the first strand DNA, e.g. cDNA (the strand comprising, i.e. incorporating, the capture probe) may be released from the array and second strand synthesis, whether as a discrete step or in an amplification reaction may occur subsequently, e.g. in a reaction carried out in solution.

Where second strand synthesis takes place on the array (i.e. in situ) the method may include an optional step of removing the captured nucleic acid, e.g. RNA before the second strand synthesis, for example using an RNA digesting enzyme (RNase) e.g. RNase H. Procedures for this are well known and described in the art. However, this is generally not necessary, and in most cases the RNA degrades naturally. Removal of the tissue sample from the array will generally remove the RNA from the array. RNase H can be used if desired to increase the robustness of RNA removal.

For instance, in tissue samples that comprise large amounts of RNA, the step of generating the double stranded cDNA may yield a sufficient amount of cDNA that it may be sequenced directly (following release from the array). In this case, second strand cDNA synthesis may be achieved by any means known in the art and as described below. The second strand synthesis reaction may be performed on the array

directly, i.e. whilst the cDNA is immobilized on the array, or preferably after the cDNA has been released from the array substrate, as described below.

In other embodiments it will be necessary to enhance, i.e. amplify, the amount of secured nucleic acid, e.g. synthesized cDNA to yield quantities that are sufficient for DNA sequencing. In this embodiment, the first strand of the secured nucleic acid, e.g. cDNA molecules, which comprise also the capture probe of the features of the array, acts as a template for the amplification reaction, e.g. a polymerase chain reaction. The first reaction product of the amplification will be a second strand of DNA, e.g. cDNA, which itself will act as a template for further cycles of the amplification reaction.

In either of the above described embodiments, the second strand of DNA, e.g. cDNA, will comprise a complement of the capture probe. If the capture probe comprises a universal domain, and particularly an amplification domain within the universal domain, then this may be used for the subsequent amplification of the DNA, e.g. cDNA, e.g. the amplification reaction may comprise a primer with the same sequence as the amplification domain, i.e. a primer that is complementary (i.e. hybridizes) to the complement of the amplification domain. In view of the fact that the amplification domain is upstream of the positional domain of the capture probe (in the secured nucleic acid, e.g. the first cDNA strand), the complement of the positional domain will be incorporated in the second strand of the DNA, e.g. cDNA molecules.

In embodiments where the second strand of DNA, e.g. cDNA is generated in a single reaction, the second strand synthesis may be achieved by any suitable means. For instance, the first strand cDNA, preferably, but not necessarily, released from the array substrate, may be incubated with random primers, e.g. hexamer primers, and a DNA polymerase, preferably a strand displacement polymerase, e.g. klenow (exo), under conditions sufficient for templated DNA synthesis to occur. This process will yield double stranded cDNA molecules of varying lengths and is unlikely to yield full-length cDNA molecules, i.e. cDNA molecules that correspond to entire mRNA from which they were synthesized. The random primers will hybridise to the first strand cDNA molecules at a random position, i.e. within the sequence rather than at the end of the sequence.

If it is desirable to generate full-length DNA, e.g. cDNA, molecules, i.e. molecules that correspond to the whole of the captured nucleic acid, e.g. RNA molecule (if the nucleic acid, e.g. RNA, was partially degraded in the tissue sample then the captured nucleic acid, e.g. RNA, molecules will not be "full-length" transcripts or the same length as the initial fragments of genomic DNA), then the 3' end of the secured nucleic acid, e.g. first strand cDNA, molecules may be modified. For example, a linker or adaptor may be ligated to the 3' end of the cDNA molecules. This may be achieved using single stranded ligation enzymes such as T4 RNA ligase or Circligase™ (Epicentre Biotechnologies).

Alternatively, a helper probe (a partially double stranded DNA molecule capable of hybridising to the 3' end of the first strand cDNA molecule), may be ligated to the 3' end of the secured nucleic acid, e.g. first strand cDNA, molecule using a double stranded ligation enzyme such as T4 DNA ligase. Other enzymes appropriate for the ligation step are known in the art and include, e.g. Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9° N) DNA ligase (9° N™ DNA ligase, New England Biolabs), and Ampligase™ (Epicentre Biotechnologies). The helper probe comprises also a specific sequence from which the second strand DNA, e.g. cDNA, synthesis may be primed using a primer that is

complementary to the part of the helper probe that is ligated to the secured nucleic acid, e.g. first cDNA strand. A further alternative comprises the use of a terminal transferase active enzyme to incorporate a polynucleotide tail, e.g. a poly-A tail, at the 3' end of the secured nucleic acid, e.g. first strand of cDNA, molecules. The second strand synthesis may be primed using a poly-T primer, which may also comprise a specific amplification domain for further amplification. Other methods for generating "full-length" double stranded DNA, e.g. cDNA, molecules (or maximal length second strand synthesis) are well-established in the art.

In some embodiments, second strand synthesis may use a method of template switching, e.g. using the SMART™ technology from Clontech®. SMART (Switching Mechanism at 5' End of RNA Template) technology is well established in the art and is based that the discovery that reverse transcriptase enzymes, e.g. Superscript® II (Invitrogen), are capable of adding a few nucleotides at the 3' end of an extended cDNA molecule, i.e. to produce a DNA/RNA hybrid with a single stranded DNA overhang at the 3' end. The DNA overhang may provide a target sequence to which an oligonucleotide probe can hybridise to provide an additional template for further extension of the cDNA molecule. Advantageously, the oligonucleotide probe that hybridises to the cDNA overhang contains an amplification domain sequence, the complement of which is incorporated into the synthesised first strand cDNA product. Primers containing the amplification domain sequence, which will hybridise to the complementary amplification domain sequence incorporated into the cDNA first strand, can be added to the reaction mix to prime second strand synthesis using a suitable polymerase enzyme and the cDNA first strand as a template. This method avoids the need to ligate adaptors to the 3' end of the cDNA first strand. Whilst template switching was originally developed for full-length mRNAs, which have a 5' cap structure, it has since been demonstrated to work equally well with truncated mRNAs without the cap structure. Thus, template switching may be used in the methods of the invention to generate full length and/or partial or truncated cDNA molecules. Thus, in a preferred embodiment of the invention, the second strand synthesis may utilise, or be achieved by, template switching. In a particularly preferred embodiment, the template switching reaction, i.e. the further extension of the cDNA first strand to incorporate the complementary amplification domain, is performed in situ (whilst the capture probe is still attached, directly or indirectly, to the array). Preferably, the second strand synthesis reaction is also performed in situ.

In embodiments where it may be necessary or advantageous to enhance, enrich or amplify the DNA, e.g. cDNA molecules, amplification domains may be incorporated in the DNA, e.g. cDNA molecules. As discussed above, a first amplification domain may be incorporated into the secured nucleic acid molecules, e.g. the first strand of the cDNA molecules, when the capture probe comprises a universal domain comprising an amplification domain. In these embodiments, the second strand synthesis may incorporate a second amplification domain. For example, the primers used to generate the second strand cDNA, e.g. random hexamer primers, poly-T primer, the primer that is complementary to the helper probe, may comprise at their 5' end an amplification domain, i.e. a nucleotide sequence to which an amplification primer may hybridize. Thus, the resultant double stranded DNA may comprise an amplification domain at or towards each 5' end of the double stranded DNA, e.g. cDNA molecules. These amplification domains may be used as targets for primers used in an amplification

reaction, e.g. PCR. Alternatively, the linker or adaptor which is ligated to the 3' end of the secured nucleic acid molecules, e.g. first strand cDNA molecules, may comprise a second universal domain comprising a second amplification domain. Similarly, a second amplification domain may be incorporated into the first strand cDNA molecules by template switching.

In embodiments where the capture probe does not comprise a universal domain, particularly comprising an amplification domain, the second strand of the cDNA molecules may be synthesised in accordance with the above description. The resultant double stranded DNA molecules may be modified to incorporate an amplification domain at the 5' end of the first DNA, e.g. cDNA strand (a first amplification domain) and, if not incorporated in the second strand DNA, e.g. cDNA synthesis step, at the 5' end of the second DNA, e.g. cDNA strand (a second amplification domain). Such amplification domains may be incorporated, e.g. by ligating double stranded adaptors to the ends of the DNA, e.g. cDNA molecules. Enzymes appropriate for the ligation step are known in the art and include, e.g. Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9° N) DNA ligase (9° N™ DNA ligase, New England Biolabs), Ampligase™ (Epicentre Biotechnologies) and T4 DNA ligase. In a preferred embodiment the first and second amplification domains comprise different sequences.

From the above, it is therefore apparent that universal domains, which may comprise an amplification domain, may be added to the secured (i.e. extended or ligated) DNA molecules, for example to the cDNA molecules, or their complements (e.g. second strand) by various methods and techniques and combinations of such techniques known in the art e.g. by use of primers which include such a domain, ligation of adaptors, use of terminal transferase enzymes and/or by template switching methods. As is clear from the discussion herein, such domains may be added before or after release of the DNA molecules from the array.

It will be apparent from the above description that all of the DNA, e.g. cDNA molecules from a single array that have been synthesized by the methods of the invention may all comprise the same first and second amplification domains. Consequently, a single amplification reaction, e.g. PCR, may be sufficient to amplify all of the DNA, e.g. cDNA molecules. Thus in a preferred embodiment, the method of the invention may comprise a step of amplifying the DNA, e.g. cDNA molecules. In one embodiment the amplification step is performed after the release of the DNA, e.g. cDNA molecules from the substrate of the array. In other embodiments amplification may be performed on the array (i.e. in situ on the array). It is known in the art that amplification reactions may be carried out on arrays and on-chip thermocyclers exist for carrying out such reactions. Thus, in one embodiment arrays which are known in the art as sequencing platforms or for use in any form of sequence analysis (e.g. in or by next generation sequencing technologies) may be used as the basis of the arrays of the present invention (e.g. Illumina bead arrays etc.)

For the synthesis of the second strand of DNA, e.g. cDNA it is preferable to use a strand displacement polymerase (e.g. 029 DNA polymerase, Bst (exo⁻) DNA polymerase, klenow (exo⁻) DNA polymerase) if the cDNA released from the substrate of the array comprises a partially double stranded nucleic acid molecule. For instance, the released nucleic acids will be at least partially double stranded (e.g. DNA:DNA, DNA:RNA or DNA:DNA/RNA hybrid) in embodiments where the capture probe is immobilized indirectly on the substrate of the array via a surface probe and the step of

releasing the DNA, e.g. cDNA molecules comprises a cleavage step. The strand displacement polymerase is necessary to ensure that the second cDNA strand synthesis incorporates the complement of the positional domain (feature identification domain) into the second DNA, e.g. cDNA strand.

It will be evident that the step of releasing at least part of the DNA, e.g. cDNA molecules or their amplicons from the surface or substrate of the array may be achieved using a number of methods. The primary aim of the release step is to yield molecules into which the positional domain of the capture probe (or its complement) is incorporated (or included), such that the DNA, e.g. cDNA molecules or their amplicons are "tagged" according to their feature (or position) on the array. The release step thus removes DNA, e.g. cDNA molecules or amplicons thereof from the array, which DNA, e.g. cDNA molecules or amplicons include the positional domain or its complement (by virtue of it having been incorporated into the secured nucleic acid, e.g. the first strand cDNA by, e.g. extension of the capture probe, and optionally copied in the second strand DNA if second strand synthesis takes place on the array, or copied into amplicons if amplification takes place on the array). Hence, in order to yield sequence analysis data that can be correlated with the various regions in the tissue sample it is essential that the released molecules comprise the positional domain of the capture probe (or its complement).

Since the released molecule may be a first and/or second strand DNA, e.g. cDNA molecule or amplicon, and since the capture probe may be immobilised indirectly on the array, it will be understood that whilst the release step may comprise a step of cleaving a DNA, e.g. cDNA molecule from the array, the release step does not require a step of nucleic acid cleavage; a DNA, e.g. cDNA molecule or an amplicon may simply be released by denaturing a double-stranded molecule, for example releasing the second cDNA strand from the first cDNA strand, or releasing an amplicon from its template or releasing the first strand cDNA molecule (i.e. the extended capture probe) from a surface probe. Accordingly, a DNA, e.g. cDNA molecule may be released from the array by nucleic acid cleavage and/or by denaturation (e.g. by heating to denature a double-stranded molecule). Where amplification is carried out in situ on the array, this will of course encompass releasing amplicons by denaturation in the cycling reaction.

In some embodiments, the DNA, e.g. cDNA molecules are released by enzymatic cleavage of a cleavage domain, which may be located in the universal domain or positional domain of the capture probe. As mentioned above, the cleavage domain must be located upstream (at the 5' end) of the positional domain, such that the released DNA, e.g. cDNA molecules comprise the positional (identification) domain. Suitable enzymes for nucleic acid cleavage include restriction endonucleases, e.g. RsaI. Other enzymes, e.g. a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase-lyase Endonuclease VIII (USER™ enzyme) or a combination of the MutY and T7 endonuclease I enzymes, are preferred embodiments of the methods of the invention.

In an alternative embodiment, the DNA, e.g. cDNA molecules may be released from the surface or substrate of the array by physical means. For instance, in embodiments where the capture probe is indirectly immobilized on the substrate of the array, e.g. via hybridization to the surface probe, it may be sufficient to disrupt the interaction between the nucleic acid molecules. Methods for disrupting the interaction between nucleic acid molecules, e.g. denaturing double stranded nucleic acid molecules, are well known in

the art. A straightforward method for releasing the DNA, e.g. cDNA molecules (i.e. of stripping the array of the synthesized DNA, e.g. cDNA molecules) is to use a solution that interferes with the hydrogen bonds of the double stranded molecules. In a preferred embodiment of the invention, the DNA, e.g. cDNA molecules may be released by applying heated water, e.g. water or buffer of at least 85° C., preferably at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99° C. As an alternative or addition to the use of a temperature sufficient to disrupt the hydrogen bonding, the solution may comprise salts, surfactants etc. that may further destabilize the interaction between the nucleic acid molecules, resulting in the release of the DNA, e.g. cDNA molecules.

It will be understood that the application of a high temperature solution, e.g. 90-99° C. water may be sufficient to disrupt a covalent bond used to immobilize the capture probe or surface probe to the array substrate. Hence, in a preferred embodiment, the DNA, e.g. cDNA molecules may be released by applying hot water to the array to disrupt covalently immobilized capture or surface probes.

It is implicit that the released DNA, e.g. cDNA molecules (the solution comprising the released DNA, e.g. cDNA molecules) are collected for further manipulation, e.g. second strand synthesis and/or amplification. Nevertheless, the method of the invention may be seen to comprise a step of collecting or recovering the released DNA, e.g. cDNA molecules. As noted above, in the context of in situ amplification the released molecules may include amplicons of the secured nucleic acid, e.g. cDNA.

In embodiments of methods of the invention, it may be desirable to remove any unextended or unligated capture probes. This may be, for example, after the step of releasing DNA molecules from the array. Any desired or convenient method may be used for such removal including, for example, use of an enzyme to degrade the unextended or unligated probes, e.g. exonuclease.

The DNA, e.g. cDNA molecules, or amplicons, that have been released from the array, which may have been modified as discussed above, are analysed to investigate (e.g. determine their sequence, although as noted above actual sequence determination is not required—any method of analysing the sequence may be used). Thus, any method of nucleic acid analysis may be used. The step of sequence analysis may identify the positional domain and hence allow the analysed molecule to be localized to a position in the tissue sample. Similarly, the nature or identity of the analysed molecule may be determined. In this way the nucleic acid, e.g. RNA at given position in the array, and hence in the tissue sample may be determined. Hence the analysis step may include or use any method which identifies the analysed molecule (and hence the "target" molecule) and its positional domain. Generally such a method will be a sequence-specific method. For example, the method may use sequence-specific primers or probes, particularly primers or probes specific for the positional domain and/or for a specific nucleic acid molecule to be detected or analysed e.g. a DNA molecule corresponding to a nucleic acid, e.g. RNA or cDNA molecule to be detected. Typically in such a method sequence-specific amplification primers e.g. PCR primers may be used.

In some embodiments it may be desirable to analyse a subset or family of target related molecules, e.g. all of the sequences that encode a particular group of proteins which share sequence similarity and/or conserved domains, e.g. a family of receptors. Hence, the amplification and/or analysis methods described herein may use degenerate or gene family specific primers or probes that hybridise to a subset

of the captured nucleic acids or nucleic acids derived therefrom, e.g. amplicons. In a particularly preferred embodiment, the amplification and/or analysis methods may utilise a universal primer (i.e. a primer common to all of the captured sequences) in combination with a degenerate or gene family specific primer specific for a subset of target molecules.

Thus in one embodiment, amplification-based, especially PCR-based methods of sequence analysis are used.

However, the steps of modifying and/or amplifying the released DNA, e.g. cDNA molecules may introduce additional components into the sample, e.g. enzymes, primers, nucleotides etc. Hence, the methods of the invention may further comprise a step of purifying the sample comprising the released DNA, e.g. cDNA molecules or amplicons prior to the sequence analysis, e.g. to remove oligonucleotide primers, nucleotides, salts etc that may interfere with the sequencing reactions. Any suitable method of purifying the DNA, e.g. cDNA molecules may be used.

As noted above, sequence analysis of the released DNA molecules may be direct or indirect. Thus the sequence analysis substrate (which may be viewed as the molecule which is subjected to the sequence analysis step or process) may directly be the molecule which is released from the array or it may be a molecule which is derived therefrom. Thus, for example in the context of sequence analysis step which involves a sequencing reaction, the sequencing template may be the molecule which is released from the array or it may be a molecule derived therefrom. For example, a first and/or second strand DNA, e.g. cDNA molecule released from the array may be directly subjected to sequence analysis (e.g. sequencing), i.e. may directly take part in the sequence analysis reaction or process (e.g. the sequencing reaction or sequencing process, or be the molecule which is sequenced or otherwise identified). In the context of in situ amplification the released molecule may be an amplicon. Alternatively, the released molecule may be subjected to a step of second strand synthesis or amplification before sequence analysis (e.g. sequencing or identification by other means). The sequence analysis substrate (e.g. template) may thus be an amplicon or a second strand of a molecule which is directly released from the array.

Both strands of a double stranded molecule may be subjected to sequence analysis (e.g. sequenced) but the invention is not limited to this and single stranded molecules (e.g. cDNA) may be analysed (e.g. sequenced). For example various sequencing technologies may be used for single molecule sequencing, e.g. the Helicos or Pacbio technologies, or nanopore sequencing technologies which are being developed. Thus, in one embodiment the first strand of DNA, e.g. cDNA may be subjected to sequencing. The first strand DNA, e.g. cDNA may need to be modified at the 3' end to enable single molecule sequencing. This may be done by procedures analogous to those for handling the second DNA, e.g. cDNA strand. Such procedures are known in the art.

In a preferred aspect of the invention the sequence analysis will identify or reveal a portion of captured nucleic acid, e.g. RNA sequence and the sequence of the positional domain. The sequence of the positional domain (or tag) will identify the feature to which the nucleic acid, e.g. mRNA molecule was captured. The sequence of the captured nucleic acid, e.g. RNA molecule may be compared with a sequence database of the organism from which the sample originated to determine the gene to which it corresponds. By determining which region (e.g. cell) of the tissue sample was in contact with the feature, it is possible to determine which

region of the tissue sample was expressing said gene (or contained the gene, e.g. in the case of spatial genomics). This analysis may be achieved for all of the DNA, e.g. cDNA molecules generated by the methods of the invention, yielding a spatial transcriptome or genome of the tissue sample.

By way of a representative example, sequencing data may be analysed to sort the sequences into specific species of capture probe, i.e. according to the sequence of the positional domain. This may be achieved by, e.g. using the FastX toolkit FASTQ Barcode splitter tool to sort the sequences into individual files for the respective capture probe positional domain (tag) sequences. The sequences of each species, i.e. from each feature, may be analyzed to determine the identity of the transcripts. For instance, the sequences may be identified using e.g. Blastn software, to compare the sequences to one or more genome databases, preferably the database for the organism from which the tissue sample was obtained. The identity of the database sequence with the greatest similarity to the sequence generated by the methods of the invention will be assigned to said sequence. In general, only hits with a certainty of at least 1×10^{-6} , preferably 1×10^{-7} , 1×10^{-8} , or 1×10^{-9} will be considered to have been successfully identified.

It will be apparent that any nucleic acid sequencing method may be utilised in the methods of the invention. However, the so-called "next generation sequencing" techniques will find particular utility in the present invention. High-throughput sequencing is particularly useful in the methods of the invention because it enables a large number of nucleic acids to be partially sequenced in a very short period of time. In view of the recent explosion in the number of fully or partially sequenced genomes, it is not essential to sequence the full length of the generated DNA, e.g. cDNA molecules to determine the gene to which each molecule corresponds. For example, the first 100 nucleotides from each end of the DNA, e.g. cDNA molecules should be sufficient to identify both the feature to which the nucleic acid, e.g. mRNA was captured (i.e. its location on the array) and the gene expressed. The sequence reaction from the "capture probe end" of the DNA, e.g. cDNA molecules yields the sequence of the positional domain and at least about 20 bases, preferably 30 or 40 bases of transcript specific sequence data. The sequence reaction from the "non-capture probe end" may yield at least about 70 bases, preferably 80, 90, or 100 bases of transcript specific sequence data.

As a representative example, the sequencing reaction may be based on reversible dye-terminators, such as used in the Illumina™ technology. For example, DNA molecules are first attached to primers on, e.g. a glass or silicon slide and amplified so that local clonal colonies are formed (bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed away. Unlike pyrosequencing, the DNA can only be extended one nucleotide at a time. A camera takes images of the fluorescently labelled nucleotides then the dye along with the terminal 3' blocker is chemically removed from the DNA, allowing a next cycle. This may be repeated until the required sequence data is obtained. Using this technology, thousands of nucleic acids may be sequenced simultaneously on a single slide.

Other high-throughput sequencing techniques may be equally suitable for the methods of the invention, e.g. pyrosequencing. In this method the DNA is amplified inside water droplets in an oil solution (emulsion PCR), with each droplet containing a single DNA template attached to a single primer-coated bead that then forms a clonal colony.

The sequencing machine contains many picolitre-volume wells each containing a single bead and sequencing enzymes. Pyrosequencing uses luciferase to generate light for detection of the individual nucleotides added to the nascent DNA and the combined data are used to generate sequence read-outs.

An example of a technology in development is based on the detection of hydrogen ions that are released during the polymerisation of DNA. A microwell containing a template DNA strand to be sequenced is flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide it is incorporated into the growing complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogen ions and a proportionally higher electronic signal.

Thus, it is clear that future sequencing formats are slowly being made available, and with shorter run times as one of the main features of those platforms it will be evident that other sequencing technologies will be useful in the methods of the invention.

An essential feature of the present invention, as described above, is a step of securing a complementary strand of the captured nucleic acid molecules to the capture probe, e.g. reverse transcribing the captured RNA molecules. The reverse transcription reaction is well known in the art and in representative reverse transcription reactions, the reaction mixture includes a reverse transcriptase, dNTPs and a suitable buffer. The reaction mixture may comprise other components, e.g. RNase inhibitor(s). The primers and template are the capture domain of the capture probe and the captured RNA molecules are described above. In the subject methods, each dNTP will typically be present in an amount ranging from about 10 to 5000 μ M, usually from about 20 to 1000 μ M. It will be evident that an equivalent reaction may be performed to generate a complementary strand of a captured DNA molecule, using an enzyme with DNA polymerase activity. Reactions of this type are well known in the art and are described in more detail below.

The desired reverse transcriptase activity may be provided by one or more distinct enzymes, wherein suitable examples are: M-MLV, MuLV, AMV, HIV, ArrayScript™, MultiScribe™, ThermoScript™, and SuperScript® I, II, and III enzymes.

The reverse transcriptase reaction may be carried out at any suitable temperature, which will be dependent on the properties of the enzyme. Typically, reverse transcriptase reactions are performed between 37-55° C., although temperatures outside of this range may also be appropriate. The reaction time may be as little as 1, 2, 3, 4 or 5 minutes or as much as 48 hours. Typically the reaction will be carried out for between 5-120 minutes, preferably 5-60, 5-45 or 5-30 minutes or 1-10 or 1-5 minutes according to choice. The reaction time is not critical and any desired reaction time may be used.

As indicated above, certain embodiments of the methods include an amplification step, where the copy number of generated DNA, e.g. cDNA molecules is increased, e.g., in order to enrich the sample to obtain a better representation of the nucleic acids, e.g. transcripts captured from the tissue sample. The amplification may be linear or exponential, as desired, where representative amplification protocols of interest include, but are not limited to: polymerase chain reaction (PCR); isothermal amplification, etc.

The polymerase chain reaction (PCR) is well known in the art, being described in U.S. Pat. Nos. 4,683,202; 4,683,195; 4,800,159; 4,965,188 and 5,512,462, the disclosures of which are herein incorporated by reference. In representative PCR amplification reactions, the reaction mixture that includes the above released DNA, e.g. cDNA molecules from the array, which are combined with one or more primers that are employed in the primer extension reaction, e.g., the PCR primers that hybridize to the first and/or second amplification domains (such as forward and reverse primers employed in geometric (or exponential) amplification or a single primer employed in a linear amplification). The oligonucleotide primers with which the released DNA, e.g. cDNA molecules (hereinafter referred to as template DNA for convenience) is contacted will be of sufficient length to provide for hybridization to complementary template DNA under annealing conditions (described in greater detail below). The length of the primers will depend on the length of the amplification domains, but will generally be at least 10 bp in length, usually at least 15 bp in length and more usually at least 16 bp in length and may be as long as 30 bp in length or longer, where the length of the primers will generally range from 18 to 50 bp in length, usually from about 20 to 35 bp in length. The template DNA may be contacted with a single primer or a set of two primers (forward and reverse primers), depending on whether primer extension, linear or exponential amplification of the template DNA is desired.

In addition to the above components, the reaction mixture produced in the subject methods typically includes a polymerase and deoxyribonucleoside triphosphates (dNTPs). The desired polymerase activity may be provided by one or more distinct polymerase enzymes. In many embodiments, the reaction mixture includes at least a Family A polymerase, where representative Family A polymerases of interest include, but are not limited to: *Thermus aquaticus* polymerases, including the naturally occurring polymerase (Taq) and derivatives and homologues thereof, such as KlenTaq (as described in Barnes et al, Proc. Natl. Acad. Sci USA (1994) 91:2216-2220); *Thermus thermophilus* polymerases, including the naturally occurring polymerase (Tth) and derivatives and homologues thereof, and the like. In certain embodiments where the amplification reaction that is carried out is a high fidelity reaction, the reaction mixture may further include a polymerase enzyme having 3'-5' exonuclease activity, e.g., as may be provided by a Family B polymerase, where Family B polymerases of interest include, but are not limited to: *Thermococcus litoralis* DNA polymerase (Vent) as described in Perler et al., Proc. Natl. Acad. Sci. USA (1992) 89:5577-5581; *Pyrococcus* species GB-D (Deep Vent); *Pyrococcus furiosus* DNA polymerase (Pfu) as described in Lundberg et al., Gene (1991) 108:1-6, *Pyrococcus woesei* (Pwo) and the like. Where the reaction mixture includes both a Family A and Family B polymerase, the Family A polymerase may be present in the reaction mixture in an amount greater than the Family B polymerase, where the difference in activity will usually be at least 10-fold, and more usually at least about 100-fold. Usually the reaction mixture will include four different types of dNTPs corresponding to the four naturally occurring bases present, i.e. dATP, dTTP, dCTP and dGTP. In the subject methods, each dNTP will typically be present in an amount ranging from about 10 to 5000 μ M, usually from about 20 to 1000 μ M.

The reaction mixtures prepared in the reverse transcriptase and/or amplification steps of the subject methods may further include an aqueous buffer medium that includes

a source of monovalent ions, a source of divalent cations and a buffering agent. Any convenient source of monovalent ions, such as KCl, K-acetate, NH₄-acetate, K-glutamate, NH₄Cl, ammonium sulphate, and the like may be employed. The divalent cation may be magnesium, manganese, zinc and the like, where the cation will typically be magnesium. Any convenient source of magnesium cation may be employed, including MgCl₂, Mg-acetate, and the like. The amount of Mg²⁺ present in the buffer may range from 0.5 to 10 mM, but will preferably range from about 3 to 6 mM, and will ideally be at about 5 mM. Representative buffering agents or salts that may be present in the buffer include Tris, Tricine, HEPES, MOPS and the like, where the amount of buffering agent will typically range from about 5 to 150 mM, usually from about 10 to 100 mM, and more usually from about 20 to 50 mM, where in certain preferred embodiments the buffering agent will be present in an amount sufficient to provide a pH ranging from about 6.0 to 9.5, where most preferred is pH 7.3 at 72° C. Other agents which may be present in the buffer medium include chelating agents, such as EDTA, EGTA and the like.

In preparing the reverse transcriptase, DNA extension or amplification reaction mixture of the steps of the subject methods, the various constituent components may be combined in any convenient order. For example, in the amplification reaction the buffer may be combined with primer, polymerase and then template DNA, or all of the various constituent components may be combined at the same time to produce the reaction mixture.

As discussed above, a preferred embodiment of the invention the DNA, e.g. cDNA molecules may be modified by the addition of amplification domains to the ends of the nucleic acid molecules, which may involve a ligation reaction. A ligation reaction is also required for the in situ synthesis of the capture probe on the array, when the capture probe is immobilized indirectly on the array surface.

As is known in the art, ligases catalyze the formation of a phosphodiester bond between juxtaposed 3'-hydroxyl and 5'-phosphate termini of two immediately adjacent nucleic acids. Any convenient ligase may be employed, where representative ligases of interest include, but are not limited to: Temperature sensitive and thermostable ligases. Temperature sensitive ligases include, but are not limited to, bacteriophage T4 DNA ligase, bacteriophage T7 ligase, and *E. coli* ligase. Thermostable ligases include, but are not limited to, Taq ligase, Tth ligase, and Pfu ligase. Thermostable ligase may be obtained from thermophilic or hyperthermophilic organisms, including but not limited to, prokaryotic, eukaryotic, or archaeal organisms. Certain RNA ligases may also be employed in the methods of the invention.

In this ligation step, a suitable ligase and any reagents that are necessary and/or desirable are combined with the reaction mixture and maintained under conditions sufficient for ligation of the relevant oligonucleotides to occur. Ligation reaction conditions are well known to those of skill in the art. During ligation, the reaction mixture in certain embodiments may be maintained at a temperature ranging from about 4° C. to about 50° C., such as from about 20° C. to about 37° C. for a period of time ranging from about 5 seconds to about 16 hours, such as from about 1 minute to about 1 hour. In yet other embodiments, the reaction mixture may be maintained at a temperature ranging from about 35° C. to about 45° C., such as from about 37° C. to about 42° C., e.g., at or about 38° C., 39° C., 40° C. or 41° C., for a period of time ranging from about 5 seconds to about 16 hours, such as from about 1 minute to about 1 hour, including from about 2 minutes to

about 8 hours. In a representative embodiment, the ligation reaction mixture includes 50 mM Tris pH7.5, 10 mM MgCl₂, 10 mM DTT, 1 mM ATP, 25 mg/ml BSA, 0.25 units/ml RNase inhibitor, and T4 DNA ligase at 0.125 units/ml. In yet another representative embodiment, 2.125 mM magnesium ion, 0.2 units/ml RNase inhibitor; and 0.125 units/ml DNA ligase are employed. The amount of adaptor in the reaction will be dependent on the concentration of the DNA, e.g. cDNA in the sample and will generally be present at between 10-100 times the molar amount of DNA, e.g. cDNA.

By way of a representative example the method of the invention may comprise the following steps:

(a) contacting an array with a tissue sample, wherein the array comprises a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

(i) a positional domain that corresponds to the position of the capture probe on the array, and

(ii) a capture domain;

such that RNA of the tissue sample hybridises to said capture probes;

(b) imaging the tissue sample on the array;

(c) reverse transcribing the captured mRNA molecules to generate cDNA molecules;

(d) washing the array to remove residual tissue;

(e) releasing at least part of the cDNA molecules from the surface of the array;

(f) performing second strand cDNA synthesis on the released cDNA molecules;

and

(g) analysing the sequence of (e.g. sequencing) the cDNA molecules.

By way of an alternative representative example the method of the invention may comprise the following steps:

(a) contacting an array with a tissue sample, wherein the array comprises a substrate on which at least two species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

(i) a positional domain that corresponds to the position of the capture probe on the array, and

(ii) a capture domain;

such that RNA of the tissue sample hybridises to said capture probes;

(b) optionally rehydrating the tissue sample;

(c) reverse transcribing the captured mRNA molecules to generate first strand cDNA molecules and optionally synthesising second strand cDNA molecules;

(d) imaging the tissue sample on the array;

(e) washing the array to remove residual tissue;

(f) releasing at least part of the cDNA molecules from the surface of the array;

(g) amplifying the released cDNA molecules;

and

(h) analysing the sequence of (e.g. sequencing) the amplified cDNA molecules.

By way of yet a further representative example the method of the invention may comprise the following steps:

(a) contacting an array with a tissue sample, wherein the array comprises a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a reverse transcriptase (RT) primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

(i) a positional domain that corresponds to the position of the capture probe on the array, and

(ii) a capture domain;

such that RNA of the tissue sample hybridises to said capture probes;

(b) optionally imaging the tissue sample on the array;

(c) reverse transcribing the captured mRNA molecules to generate cDNA molecules;

(d) optionally imaging the tissue sample on the array if not already performed as step (b):

(e) washing the array to remove residual tissue;

(f) releasing at least part of the cDNA molecules from the surface of the array;

(g) performing second strand cDNA synthesis on the released cDNA molecules;

(h) amplifying the double stranded cDNA molecules;

(i) optionally purifying the cDNA molecules to remove components that may interfere with the sequencing reaction;

and

(j) analysing the sequence of (e.g. sequencing) the amplified cDNA molecules.

The present invention includes any suitable combination of the steps in the above described methods. It will be understood that the invention also encompasses variations of these methods, for example where amplification is performed in situ on the array. Also encompassed are methods which omit the imaging step.

The invention may also be seen to include a method for making or producing an array (i) for use in capturing mRNA from a tissue sample that is contacted with said array; or (ii) for use in determining and/or analysing a (e.g. the partial or global) transcriptome of a tissue sample, said method comprising immobilizing, directly or indirectly, multiple species of capture probe to an array substrate, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

(i) a positional domain that corresponds to the position of the capture probe on the array; and

(ii) a capture domain.

The method of producing an array of the invention may be further defined such that each species of capture probe is immobilized as a feature on the array.

The method of immobilizing the capture probes on the array may be achieved using any suitable means as described herein. Where the capture probes are immobilized on the array indirectly the capture probe may be synthesized on the array. Said method may comprise any one or more of the following steps:

(a) immobilizing directly or indirectly multiple surface probes to an array substrate, wherein the surface probes comprise:

(i) a domain capable of hybridizing to part of the capture domain oligonucleotide (a part not involved in capturing the nucleic acid, e.g. RNA);

(ii) a complementary positional domain; and

(iii) a complementary universal domain;

(b) hybridizing to the surface probes immobilized on the array capture domain oligonucleotides and universal domain oligonucleotides;

(c) extending the universal domain oligonucleotides, by templated polymerisation, to generate the positional domain of the capture probe; and

(d) ligating the positional domain to the capture domain oligonucleotide to produce the capture oligonucleotide.

Ligation in step (d) may occur simultaneously with extension in step (c). Thus it need not be carried out in a separate step, although this is course encompassed if desired.

The features of the array produced by the above method of producing the array of the invention, may be further defined in accordance with the above description.

Although the invention is described above with reference to detection or analysis of RNA, and transcriptome analysis or detection, it will be appreciated that the principles described can be applied analogously to the detection or analysis of DNA in cells and to genomic studies. Thus, more broadly viewed, the invention can be seen as being generally applicable to the detection of nucleic acids in general and in a further more particular aspect, as providing methods for the analysis or detection of DNA. Spatial information may be valuable also in a genomics context i.e. detection and/or analysis of a DNA molecule with spatial resolution. This may be achieved by genomic tagging according to the present invention. Such localized or spatial detection methods may be useful for example in the context of studying genomic variations in different cells or regions of a tissue, for example comparing normal and diseased cells or tissues (e.g. normal vs tumour cells or tissues) or in studying genomic changes in disease progression etc. For example, tumour tissues may comprise a heterogeneous population of cells which may differ in the genomic variants they contain (e.g. mutations and/or other genetic aberrations, for example chromosomal rearrangements, chromosomal amplifications/deletions/insertions etc.). The detection of genomic variations, or different genomic loci, in different cells in a localized way may be useful in such a context, e.g. to study the spatial distribution of genomic variations. A principal utility of such a method would be in tumour analysis. In the context of the present invention, an array may be prepared which is designed, for example, to capture the genome of an entire cell on one feature. Different cells in the tissue sample may thus be compared. Of course the invention is not limited to such a design and other variations may be possible, wherein the DNA is detected in a localized way and the position of the DNA captured on the array is correlated to a position or location in the tissue sample.

Accordingly, in a more general aspect, the present invention can be seen to provide a method for localized detection of nucleic acid in a tissue sample comprising:

(a) providing an array comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a primer for a primer extension or ligation reaction, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

(i) a positional domain that corresponds to the position of the capture probe on the array, and

(ii) a capture domain;

(b) contacting said array with a tissue sample such that the position of a capture probe on the array may be correlated with a position in the tissue sample and

41

- allowing nucleic acid of the tissue sample to hybridise to the capture domain in said capture probes;
- (c) generating DNA molecules from the captured nucleic acid molecules using said capture probes as extension or ligation primers, wherein said extended or ligated DNA molecules are tagged by virtue of the positional domain;
 - (d) optionally generating a complementary strand of said tagged DNA and/or optionally amplifying said tagged DNA;
 - (e) releasing at least part of the tagged DNA molecules and/or their complements or amplicons from the surface of the array, wherein said part includes the positional domain or a complement thereof;
 - (f) directly or indirectly analysing the sequence of (e.g. sequencing) the released DNA molecules.

As described in more detail above, any method of nucleic acid analysis may be used in the analysis step. Typically this may involve sequencing, but it is not necessary to perform an actual sequence determination. For example sequence-specific methods of analysis may be used. For example a sequence-specific amplification reaction may be performed, for example using primers which are specific for the positional domain and/or for a specific target sequence, e.g. a particular target DNA to be detected (i.e. corresponding to a particular cDNA/RNA or gene or gene variant or genomic locus or genomic variant etc.). An exemplary analysis method is a sequence-specific PCR reaction.

The sequence analysis (e.g. sequencing) information obtained in step (f) may be used to obtain spatial information as to the nucleic acid in the sample. In other words the sequence analysis information may provide information as to the location of the nucleic acid in the sample. This spatial information may be derived from the nature of the sequence analysis information obtained e.g. from a sequence determined or identified, for example it may reveal the presence of a particular nucleic acid molecule which may itself be spatially informative in the context of the tissue sample used, and/or the spatial information (e.g. spatial localisation) may be derived from the position of the tissue sample on the array, coupled with the sequence analysis information. However, as described above, spatial information may conveniently be obtained by correlating the sequence analysis data to an image of the tissue sample and this represents one preferred embodiment of the invention.

Accordingly, in a preferred embodiment the method also includes a step of:

- (g) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (c).

The primer extension reaction referred to in step (a) may be defined as a polymerase-catalysed extension reaction and acts to acquire a complementary strand of the captured nucleic acid molecule that is covalently attached to the capture probe, i.e. by synthesising the complementary strand utilising the capture probe as a primer and the captured nucleic acid as a template. In other words it may be any primer extension reaction carried out by any polymerase enzyme. The nucleic acid may be RNA or it may be DNA. Accordingly the polymerase may be any polymerase. It may be a reverse transcriptase or it may be a DNA polymerase. The ligation reaction may be carried out by any ligase and acts to secure the complementary strand of the captured nucleic acid molecule to the capture probe, i.e. wherein the captured nucleic acid molecule (hybridised to the capture probe) is partially double stranded and the complementary strand is ligated to the capture probe.

42

One preferred embodiment of such a method is the method described above for the determination and/or analysis of a transcriptome, or for the detection of RNA. In alternative preferred embodiment the detected nucleic acid molecule is DNA. In such an embodiment the invention provides a method for localized detection of DNA in a tissue sample comprising:

- (a) providing an array comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a primer for a primer extension or ligation reaction, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':
 - (i) a positional domain that corresponds to the position of the capture probe on the array, and
 - (ii) a capture domain;
- (b) contacting said array with a tissue sample such that the position of a capture probe on the array may be correlated with a position in the tissue sample and allowing DNA of the tissue sample to hybridise to the capture domain in said capture probes;
- (c) fragmenting DNA in said tissue sample, wherein said fragmentation is carried out before, during or after contacting the array with the tissue sample in step (b);
- (d) extending said capture probes in a primer extension reaction using the captured DNA fragments as templates to generate extended DNA molecules, or ligating the captured DNA fragments to the capture probes in a ligation reaction to generate ligated DNA molecules, wherein said extended or ligated DNA molecules are tagged by virtue of the positional domain;
- (e) optionally generating a complementary strand of said tagged DNA and/or optionally amplifying said tagged DNA;
- (f) releasing at least part of the tagged DNA molecules and/or their complements and/or amplicons from the surface of the array, wherein said part includes the positional domain or a complement thereof;
- (g) directly or indirectly analysing the sequence of the released DNA molecules.

The method may further include a step of:

- (h) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (d).

In the context of spatial genomics, where the target nucleic acid is DNA the inclusion of imaging and image correlation steps may in some circumstances be preferred.

In embodiments in which DNA is captured, the DNA may be any DNA molecule which may occur in a cell. Thus it may be genomic, i.e. nuclear, DNA, mitochondrial DNA or plastid DNA, e.g. chloroplast DNA. In a preferred embodiment, the DNA is genomic DNA.

It will be understood that where fragmentation is carried out after the contacting in step (b), i.e. after the tissue sample is placed on the array, fragmentation occurs before the DNA is hybridised to the capture domain. In other words the DNA fragments are hybridised (or more particularly, allowed to hybridise) to the capture domain in said capture probes.

Advantageously, but not necessarily, in a particular embodiment of this aspect of the invention, the DNA fragments of the tissue sample may be provided with a binding domain to enable or facilitate their capture by the capture probes on the array. Accordingly, the binding domain is capable of hybridising to the capture domain of the capture probe. Such a binding domain may thus be

regarded as a complement of the capture domain (i.e. it may be viewed as a complementary capture domain), although absolute complementarity between the capture and binding domains is not required, merely that the binding domain is sufficiently complementary to allow a productive hybridisation to take place, i.e. that the DNA fragments in the tissue sample are able to hybridise to the capture domain of the capture probes. Provision of such a binding domain may ensure that DNA in the sample does not bind to the capture probes until after the fragmentation step. The binding domain may be provided to the DNA fragments by procedures well known in the art, for example by ligation of adaptor or linker sequences which may contain the binding domain. For example a linker sequence with a protruding end may be used. The binding domain may be present in the single-stranded portion of such a linker, such that following ligation of the linker to the DNA fragments, the single-stranded portion containing the binding domain is available for hybridisation to the capture domain of the capture probes. Alternatively and in a preferred embodiment, the binding domain may be introduced by using a terminal transferase enzyme to introduce a polynucleotide tail e.g. a homopolymeric tail such as a poly-A domain. This may be carried out using a procedure analogous to that described above for introducing a universal domain in the context of the RNA methods. Thus, in advantageous embodiments a common binding domain may be introduced. In other words, a binding domain which is common to all the DNA fragments and which may be used to achieve the capture of the fragments on the array.

Where a tailing reaction is carried out to introduce a (common) binding domain, the capture probes on the array may be protected from the tailing reaction, i.e. the capture probes may be blocked or masked as described above. This may be achieved for example by hybridising a blocking oligonucleotide to the capture probe e.g. to the protruding end (e.g. single stranded portion) of the capture probe. Where the capture domain comprises a poly-T sequence for example, such a blocking oligonucleotide may be a poly-A oligonucleotide. The blocking oligonucleotide may have a blocked 3' end (i.e. an end incapable of being extended, or tailed). The capture probes may also be protected, i.e. blocked, by chemical and/or enzymatic modifications, as described in detail above.

Where the binding domain is provided by ligation of a linker as described above, it will be understood that rather than extending the capture probe to generate a complementary copy of the captured DNA fragment which comprises the positional tag of the capture probe primer, the DNA fragment may be ligated to the 3' end of the capture probe. As noted above ligation requires that the 5' end to be ligated is phosphorylated. Accordingly, in one embodiment, the 5' end of the added linker, namely the end which is to be ligated to the capture probe (i.e. the non-protruding end of the linker added to the DNA fragments) will be phosphorylated. In such a ligation embodiment, it will accordingly be seen that a linker may be ligated to double stranded DNA fragments, said linker having a single stranded protruding 3' end which contains the binding domain. Upon contact with the array, the protruding end hybridises to the capture domain of the capture probes. This hybridisation brings the 3' end of the capture probe into juxtaposition for ligation to the 5' (non-protruding) end of the added linker. The capture probe, and hence the positional domain, is thus incorporated into the captured DNA fragment by this ligation. Such an embodiment is shown schematically in FIG. 21.

Thus, the method of this aspect of the invention may in a more particular embodiment comprise:

- (a) providing an array comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a primer for a primer extension or ligation reaction, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':
 - (i) a positional domain that corresponds to the position of the capture probe on the array, and
 - (ii) a capture domain;
 - (b) contacting said array with a tissue sample such that the position of a capture probe on the array may be correlated with a position in the tissue sample;
 - (c) fragmenting DNA in said tissue sample, wherein said fragmentation is carried out before, during or after contacting the array with the tissue sample in step (b);
 - (d) providing said DNA fragments with a binding domain which is capable of hybridising to said capture domain;
 - (e) allowing said DNA fragments to hybridise to the capture domain in said capture probes;
 - (f) extending said capture probes in a primer extension reaction using the captured DNA fragments as templates to generate extended DNA molecules, or ligating the captured DNA fragments to the capture probes in a ligation reaction to generate ligated DNA molecules, wherein said extended or ligated DNA molecules are tagged by virtue of the positional domain;
 - (g) optionally generating a complementary strand of said tagged DNA and/or optionally amplifying the tagged DNA;
 - (h) releasing at least part of the tagged DNA molecules and/or their complements and/or amplicons from the surface of the array, wherein said part includes the positional domain or a complement thereof;
 - (i) directly or indirectly analysing the sequence of the released DNA molecules.
- The method may optionally include a further step of
- (j) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (f).

In the methods of nucleic acid or DNA detection set out above, the optional step of generating a complementary copy of the tagged nucleic acid/DNA or of amplifying the tagged DNA, may involve the use of a strand displacing polymerase enzyme, according to the principles explained above in the context of the RNA/transcriptome analysis/detection methods. Suitable strand displacing polymerases are discussed above. This is to ensure that the positional domain is copied into the complementary copy or amplicon. This will particularly be the case where the capture probe is immobilized on the array by hybridisation to a surface probe.

However, the use of a strand displacing polymerase in this step is not essential. For example a non-strand displacing polymerase may be used together with ligation of an oligonucleotide which hybridises to the positional domain. Such a procedure is analogous to that described above for the synthesis of capture probes on the array.

In one embodiment, the method of the invention may be used for determining and/or analysing all of the genome of a tissue sample e.g. the global genome of a tissue sample. However, the method is not limited to this and encompasses determining and/or analysing all or part of the genome. Thus, the method may involve determining and/or analysing

a part or subset of the genome, e.g. a partial genome corresponding to a subset or group of genes or of chromosomes, e.g. a set of particular genes or chromosomes or a particular region or part of the genome, for example related to a particular disease or condition, tissue type etc. Thus, the method may be used to detect or analyse genomic sequences or genomic loci from tumour tissue as compared to normal tissue, or even within different types of cell in a tissue sample. The presence or absence, or the distribution or location of different genomic variants or loci in different cells, groups of cells, tissues or parts or types of tissue may be examined.

Viewed from another aspect, the method steps set out above can be seen as providing a method of obtaining spatial information regarding the nucleic acids, e.g. genomic sequences, variants or loci of a tissue sample. Put another way, the methods of the invention may be used for the labelling (or tagging) of genomes, particularly individual or spatially distributed genomes.

Alternatively viewed, the method of the invention may be seen as a method for spatial detection of DNA in a tissue sample, or a method for detecting DNA with spatial resolution, or for localized or spatial determination and/or analysis of DNA in a tissue sample. In particular, the method may be used for the localized or spatial detection or determination and/or analysis of genes or genomic sequences or genomic variants or loci (e.g. distribution of genomic variants or loci) in a tissue sample. The localized/spatial detection/determination/analysis means that the DNA may be localized to its native position or location within a cell or tissue in the tissue sample. Thus for example, the DNA may be localized to a cell or group of cells, or type of cells in the sample, or to particular regions of areas within a tissue sample. The native location or position of the DNA (or in other words, the location or position of the DNA in the tissue sample), e.g. a genomic variant or locus, may be determined.

It will be seen therefore that the array of the present invention may be used to capture nucleic acid, e.g. DNA of a tissue sample that is contacted with said array. The array may also be used for determining and/or analysing a partial or global genome of a tissue sample or for obtaining a spatially defined partial or global genome of a tissue sample. The methods of the invention may thus be considered as methods of quantifying the spatial distribution of one or more genomic sequences (or variants or loci) in a tissue sample. Expressed another way, the methods of the present invention may be used to detect the spatial distribution of one or more genomic sequences or genomic variants or genomic loci in a tissue sample. In yet another way, the methods of the present invention may be used to determine simultaneously the location or distribution of one or more genomic sequences or genomic variants or genomic loci at one or more positions within a tissue sample. Still further, the methods may be seen as methods for partial or global analysis of the nucleic acid e.g. DNA of a tissue sample with spatial resolution e.g. two-dimensional spatial resolution.

The invention can also be seen to provide an array for use in the methods of the invention comprising a substrate on which multiple species of capture probes are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as an extension or ligation primer, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

- (i) a positional domain that corresponds to the position of the capture probe on the array, and

- (ii) a capture domain to capture nucleic acid of a tissue sample that is contacted with said array.

In one aspect the nucleic acid molecule to be captured is DNA. The capture domain may be specific to a particular DNA to be detected, or to a particular class or group of DNAs, e.g. by virtue of specific hybridisation to a specific sequence of motif in the target DNA e.g. a conserved sequence, by analogy to the methods described in the context of RNA detection above. Alternatively the DNA to be captured may be provided with a binding domain, e.g. a common binding domain as described above, which binding domain may be recognised by the capture domain of the capture probes. Thus, as noted above, the binding domain may for example be a homopolymeric sequence e.g. poly-A. Again such a binding domain may be provided according to or analogously to the principles and methods described above in relation to the methods for RNA/transcriptome analysis or detection. In such a case, the capture domain may be complementary to the binding domain introduced into the DNA molecules of the tissue sample.

As also described in the RNA context above, the capture domain may be a random or degenerate sequence. Thus, DNA may be captured non-specifically by binding to a random or degenerate capture domain or to a capture domain which comprises at least partially a random or degenerate sequence.

In a related aspect, the present invention also provides use of an array, comprising a substrate on which multiple species of capture probe are directly or indirectly immobilized such that each species occupies a distinct position on the array and is oriented to have a free 3' end to enable said probe to function as a primer for a primer extension or ligation reaction, wherein each species of said capture probe comprises a nucleic acid molecule with 5' to 3':

- (i) a positional domain that corresponds to the position of the capture probe on the array; and

- (ii) a capture domain;

to capture nucleic acid, e.g. DNA or RNA, of a tissue sample that is contacted with said array.

Preferably, said use is for localized detection of nucleic acid in a tissue sample and further comprises steps of:

- (a) generating DNA molecules from the captured nucleic acid molecules using said capture probes as extension or ligation primers, wherein said extended or ligated molecules are tagged by virtue of the positional domain;
- (b) optionally generating a complementary strand of said tagged nucleic acid and/or amplifying said tagged nucleic acid;
- (c) releasing at least part of the tagged DNA molecules and/or their complements or amplicons from the surface of the array, wherein said part includes the positional domain or a complement thereof;
- (d) directly or indirectly analysing the sequence of the released DNA molecules; and optionally
- (e) correlating said sequence analysis information with an image of said tissue sample, wherein the tissue sample is imaged before or after step (a).

The step of fragmenting DNA in a tissue sample may be carried out using any desired procedure known in the art. Thus physical methods of fragmentation may be used e.g. sonication or ultrasound treatment. Chemical methods are also known. Enzymatic methods of fragmentation may also be used, e.g. with endonucleases, for example restriction enzymes. Again methods and enzymes for this are well known in the art. Fragmentation may be done before during or after preparing the tissue sample for placing on an array,

e.g. preparing a tissue section. Conveniently, fragmentation may be achieved in the step of fixing tissue. Thus for example, formalin fixation will result in fragmentation of DNA. Other fixatives may produce similar results.

In terms of the detail of preparing and using the arrays in these aspects of the invention, it will be understood that the description and detail given above in the context of RNA methods applies analogously to the more general nucleic acid detection and DNA detection methods set out herein. Thus, all aspects and details discussed above apply analogously. For example, the discussion of reverse transcriptase primers and reactions etc may be applied analogously to any aspect of the extension primers, polymerase reactions etc. referred to above. Likewise, references and to first and second strand cDNA synthesis may be applied analogously to the tagged DNA molecule and its complement. Methods of sequence analysis as discussed above may be used.

By way of example, the capture domain may be as described for the capture probes above. A poly-T or poly-T-containing capture domain may be used for example where the DNA fragments are provided with a binding domain comprising a poly-A sequence.

The capture probes/tagged DNA molecules (i.e. the tagged extended or ligated molecules) may be provided with universal domains as described above, e.g. for amplification and/or cleavage.

The invention will be further described with reference to the following non-limiting Examples with reference to the following drawings in which:

FIG. 1 shows the overall concept using arrayed "bar-coded" oligo-dT probes to capture mRNA from tissue sections for transcriptome analysis.

FIG. 2 shows the a schematic for the visualization of transcript abundance for corresponding tissue sections.

FIG. 3 shows 3' to 5' surface probe composition and synthesis of 5' to 3' oriented capture probes that are indirectly immobilized at the array surface.

FIG. 4 shows a bar chart demonstrating the efficiency of enzymatic cleavage (USER or RsaI) from in-house manufactured arrays and by 99° C. water from Agilent manufactured arrays, as measured by hybridization of fluorescently labelled probes to the array surface after probe release.

FIG. 5 shows a fluorescent image captured after 99° C. water mediated release of DNA surface probes from commercial arrays manufactured by Agilent. A fluorescent detection probe was hybridized after hot water treatment. Top array is an untreated control.

FIG. 6 shows a fixated mouse brain tissue section on top of the transcriptome capture array post cDNA synthesis and treated with cytoplasmic (top) and nucleic stains (middle), respectively, and merged image showing both stains (bottom).

FIG. 7 shows a table that lists the reads sorted for their origin across the low density in-house manufactured DNA-capture array as seen in the schematic representation.

FIG. 8 shows a FFPE mouse brain tissue with nucleic and Map2 specific stains using a barcoded microarray.

FIG. 9 shows FFPE mouse brain olfactory bulb with nucleic stain (white) and visible morphology.

FIG. 10 shows FFPE mouse brain olfactory bulb (approx 2x2 mm) with nucleic stain (white), overlaid with theoretical spotting pattern for low resolution array.

FIG. 11 shows FFPE mouse brain olfactory bulb (approx 2x2 mm) with nucleic stain (white), overlaid with theoretical spotting pattern for medium-high resolution array.

FIG. 12 shows FFPE mouse brain olfactory bulb zoomed in on glomerular area (top right of FIG. 9).

FIG. 13 shows the resulting product from a USER release using a random hexamer primer (R6) coupled to the B_handle (B_R6) during amplification; product as depicted on a bioanalyzer.

FIG. 14 shows the resulting product from a USER release using a random octamer primer (R8) coupled to the B_handle (B_R8) during amplification; product as depicted on a bioanalyzer.

FIG. 15 shows the results of an experiment performed on FFPE brain tissue covering the whole array. ID5 (left) and ID20 (right) amplified with ID specific and gene specific primers (B2M exon 4) after synthesis and release of cDNA from surface; ID5 and ID20 amplified.

FIG. 16 shows a schematic illustration of the principle of the method described in Example 4, i.e. use of microarrays with immobilized DNA oligos (capture probes) carrying spatial labeling tag sequences (positional domains). Each feature of oligos of the microarray carries a 1) a unique labeling tag (positional domain) and 2) a capture sequence (capture domain).

FIG. 17 shows the results of the spatial genomics protocol described in Example 5 carried out with genomic DNA prefragmented to mean size of 200 bp. Internal products amplified on array labeled and synthesized DNA. The detected peak is of expected size.

FIG. 18 shows the results of the spatial genomics protocol described in Example 5 carried out with genomic DNA prefragmented to mean size of 700 bp. Internal products amplified on array labeled and synthesized DNA. The detected peak is of expected size.

FIG. 19 shows the results of the spatial genomics protocol described in Example 5 carried out with genomic DNA prefragmented to mean size of 200 bp. Products amplified with one internal primer and one universal sequence contained in the surface oligo. Amplification carried out on array labeled and synthesized DNA. The expected product is a smear given that the random fragmentation and terminal transferase labeling of genomic DNA will generate a very diverse sample pool.

FIG. 20 shows the results of the spatial genomics protocol described in Example 5 carried out with genomic DNA prefragmented to mean size of 700 bp. Products amplified with one internal primer and one universal sequence contained in the surface oligo. Amplification carried out on array labeled and synthesized DNA. The expected product is a smear given that the random fragmentation and terminal transferase labeling of genomic DNA will generate a very diverse sample pool.

FIG. 21 shows a schematic illustration of the ligation of a linker to a DNA fragment to introduce a binding domain for hybridisation to a poly-T capture domain, and subsequent ligation to the capture probe.

FIG. 22 shows the composition of 5' to 3' oriented capture probes used on high-density capture arrays.

FIG. 23 shows the frame of the high-density arrays, which is used to orientate the tissue sample, visualized by hybridization of fluorescent marker probes.

FIG. 24 shows capture probes cleaved and non-cleaved from high-density array, wherein the frame probes are not cleaved since they do not contain uracil bases. Capture probes were labelled with fluorophores coupled to poly-A oligonucleotides.

FIG. 25 shows a bioanalyzer image of a prepared sequencing library with transcripts captured from mouse olfactory bulb.

FIG. 26 shows a Matlab visualization of captured transcripts from total RNA extracted from mouse olfactory bulb.

FIG. 27 shows Olfr (olfactory receptor) transcripts as visualized across the capture array using Matlab visualization after capture from mouse olfactory bulb tissue.

FIG. 28 shows a pattern of printing for in-house 41-ID-tag microarrays.

FIG. 29 shows a spatial genomics library generated from an A431 specific translocation after capture of poly-A tailed genomic fragments on capture array.

FIG. 30 shows the detection of A431 specific translocation after capture of spiked 10% and 50% poly-A tailed A431 genomic fragments into poly-A tailed U2OS genomic fragments on capture array.

FIG. 31 shows a Matlab visualization of captured ID-tagged transcripts from mouse olfactory bulb tissue on 41-ID-tag in-house arrays overlaid with the tissue image. For clarity, the specific features on which particular genes were identified have been circled.

EXAMPLE 1

Preparation of the Array

The following experiments demonstrate how oligonucleotide probes may be attached to an array substrate by either

the 5' or 3' end to yield an array with capture probes capable of hybridizing to mRNA.

Preparation of In-House Printed Microarray with 5' to 3' Oriented Probes

20 RNA-capture oligonucleotides with individual tag sequences (Tag 1-20, Table 1) were spotted on glass slides to function as capture probes. The probes were synthesized with a 5'-terminus amino linker with a C6 spacer. All probes were synthesized by Sigma-Aldrich (St. Louis, MO, USA). The RNA-capture probes were suspended at a concentration of 20 μM in 150 mM sodium phosphate, pH 8.5 and were spotted using a Nanoplotter NP2.1/E (Gesim, Grosserkmannsdorf, Germany) onto CodeLink™ Activated microarray slides (7.5 cm×2.5 cm; Surmodics, Eden Prairie, MN, USA). After printing, surface blocking was performed according to the manufacturer's instructions. The probes were printed in 16 identical arrays on the slide, and each array contained a pre-defined printing pattern. The 16 sub-arrays were separated during hybridization by a 16-pad mask (ChipClip™ Schleicher & Schuell BioScience, Keene, NH, USA).

TABLE 1

Name	Sequence	5' mod	3' mod	Length
Sequences for free 3' capture probes				
TAP-ID1	UUAAGTACAAATCTCGACTGCCACTCTGAACCTTCTCCTTCTCCTTCA CCTTTTTTTTTTTTTTTTTTTVN (SEQ ID NO: 1)	Amino-C6		72
Enzymatic recog	UUAAGTACAA (SEQ ID NO: 2)			10
Universal amp handle P	ATCTCGACTGCCACTCTGAA (SEQ ID NO: 3)			20
ID1	CCTTCTCCTTCTCCTTCA (SEQ ID NO: 4)			20
Capture sequence	TTTTTTTTTTTTTTTTTTVN (SEQ ID NO: 5)			22
ID1	CCTTCTCCTTCTCCTTCA (SEQ ID NO: 6)			20
ID2	CCTTGCTGCTTCTCCTCCTC (SEQ ID NO: 7)			20
ID3	ACCTCCTCCGCCTCCTCCTC (SEQ ID NO: 8)			20
ID4	GAGACATACCACCAAGAGAC (SEQ ID NO: 9)			20
ID5	GTCCTCTATTCCGTCACCAT (SEQ ID NO: 10)			20
ID6	GACTGAGCTCGAACATATGG (SEQ ID NO: 11)			20
ID7	TGGAGGATTGACACAGAACG (SEQ ID NO: 12)			20
ID8	CCAGCCTCTCCATTACATCG (SEQ ID NO: 13)			20
ID9	AAGATCTACCAGCCAGCCAG (SEQ ID NO: 14)			20
ID10	CGAACTTCCACTGTCTCCTC (SEQ ID NO: 15)			20
ID11	TTGCGCCTTCTCCAATACAC (SEQ ID NO: 16)			20
ID12	CTCTTCTTAGCATGCCACCT (SEQ ID NO: 17)			20
ID13	ACCACTTCTGCATTACCTCC (SEQ ID NO: 18)			20
ID14	ACAGCCTCCTTCTTCCCTT (SEQ ID NO: 19)			20
ID15	AATCCTCTCCTTGCCAGTTC (SEQ ID NO: 20)			20
ID16	GATGCCTCCACCTGTAGAAC (SEQ ID NO: 21)			20
ID17	GAAGGAATGGAGATATCGC (SEQ ID NO: 22)			20

TABLE 1-continued

Name	Sequence	5' mod	3' mod	Length
ID18	GATCCAAGGACCATCGACTG (SEQ ID NO: 23)			20
ID19	CCACTGGAACCTGACAACCG (SEQ ID NO: 24)			20
ID20	CTGCTTCTTCTGGAECTA (SEQ ID NO: 25)			20
Sequences for free 5' surface probes and on-chip free 3' capture probe synthesis				
Free 5' surface probe-A	GCGTTCAGAGTGGCAGTCGAGATCACGCGGCAATCATATCGGACAGAT CGGAAGAGCGTAGTGTAG (SEQ ID NO: 26)		Amino C7	66
Free 5' surface probe-U	GCGTTCAGAGTGGCAGTCGAGATCACGCGGCAATCATATCGGACGGCT GCTGGTAAATAGAGATCA (SEQ ID NO: 27)		Amino C7	66
Nick	GCG			3
LP'	TTCAGAGTGGCAGTCGAGATCAC (SEQ ID NO: 28)			23
ID'	GCGGCAATCATATCGGAC (SEQ ID NO: 29)			18
A' 22 bp MutY mismatch	AGATCGGAAGAGCGTAGTGTAG (SEQ ID NO: 30)			22
U' 22 bp MutY mismatch	GGCTGCTGGTAAATAGAGATCA (SEQ ID NO: 31)			
Hybridized sequences for capture probe synthesis				
Illumina amp handle A	ACACTCTTCCCTACACGACGCTCTCCGATCT (SEQ ID NO: 32)			33
Universa ampl handle U	AAGTGTGGAAAGTTGATCGCTATTTACCAGCAGCC (SEQ ID NO: 33)			35
Capture_LP_Poly-dTVN	GTGATCTCGACTGCCACTCTGAATTTTTTTTTTTTTTTTTTTTVN (SEQ ID NO: 34)		Phosphorylated	45
Capture_LP_Poly-d24T	GTGATCTCGACTGCCACTCTGAATTTTTTTTTTTTTTTTTTTTTT (SEQ ID NO: 35)		Phosphorylated	47
Additional secondary universal amplification handles				
Illumina amp handle B	AGACGTGTGCTCTCCGATCT (SEQ ID NO: 36)			21
Universal amp handle X	ACGTCTGTGAATAGCCGCAT (SEQ ID NO: 37)			20
B_R6 handle (or X)	AGACGTGTGCTCTCCGATCTNNNNNNNNN (SEQ ID NO: 38)			27 (26)
B_R8 handle (or X)	AGACGTGTGCTCTCCGATCTNNNNNNNNN (SEQ ID NO: 39)			29 (28)
B_polyTVN (or X)	AGACGTGTGCTCTCCGATCTTTTTTTTTTTTTTTTTTTTTTVN (SEQ ID NO: 40)			43 (42)
B_poly24T (or X)	AGACGTGTGCTCTCCGATCTTTTTTTTTTTTTTTTTTTTTTTT (SEQ ID NO: 41)			45 (44)
Amplification handle to incorporate A handle into P handle products				
A_P handle	ACACTCTTCCCTACACGACGCTCTCCGATCTATCTCGACTGCCAC TCTGAA (SEQ ID NO: 42)			53

Preparation of In-House Printed Microarray with 3' to 5' Oriented Probes and Synthesis of 5' to 3' Oriented Capture Probes

Printing of surface probe oligonucleotides was performed as in the case with 5' to 3' oriented probes above, with an amino-C7 linker at the 3' end, as shown in Table 1.

To hybridize primers for capture probe synthesis, hybridization solution containing 4xSSC and 0.1% SDS, 2 μ M

extension primer (the universal domain oligonucleotide) and 2 μ M thread joining primer (the capture domain oligonucleotide) was incubated for 4 min at 50° C. Meanwhile the in-house array was attached to a ChipClip (Whatman). The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 μ L of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C.

2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake; 2) 0.2×SSC for 1 min at 300 rpm shake; and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry and placed back in the ChipClip.

For extension and ligation reaction (to generate the positional domain of the capture probe) 50 μL of enzyme mix containing 10× Ampligase buffer, 2.5 U AmpliTaq DNA Polymerase Stoffel Fragment (Applied Biosystems), 10 U Ampligase (Epicentre Biotechnologies), dNTPs 2 mM each (Fermentas) and water, was pipetted to each well. The array was subsequently incubated at 55° C. for 30 min. After incubation the array was washed according to the previously described array washing method but the first step has the duration of 10 min instead of 6 min.

The method is depicted in FIG. 3.

Tissue Preparation

The following experiments demonstrate how tissue sample sections may be prepared for use in the methods of the invention.

Preparation of Fresh Frozen Tissue and Sectioning onto Capture Probe Arrays

Fresh non-fixed mouse brain tissue was trimmed if necessary and frozen down in -40° C. cold isopentane and subsequently mounted for sectioning with a cryostat at 10 μm. A slice of tissue was applied onto each capture probe array to be used.

Preparation of Formalin-Fixed Paraffin-Embedded (FFPE) Tissue

Mouse brain tissue was fixed in 4% formalin at 4° C. for 24 h. After that it was incubated as follows: 3× incubation in 70% ethanol for 1 hour; 1× incubation in 80% ethanol for 1 hour; 1× incubation in 96% ethanol for 1 hour; 3× incubation in 100% ethanol for 1 hour; and 2× incubation in xylene at room temperature for 1 h.

The dehydrated samples were then incubated in liquid low melting paraffin 52-54° C. for up to 3 hours, during which the paraffin was changed once to wash out residual xylene. Finished tissue blocks were then stored at RT. Sections were then cut at 4 μm in paraffin with a microtome onto each capture probe array to be used.

The sections were dried at 37° C. on the array slides for 24 hours and stored at RT.

Deparaffinization of FFPE Tissue

Formalin fixed paraffinized mouse brain 10 μm sections attached to CodeLink slides were deparaffinised in xylene twice for: 10 min, 99.5% ethanol for 2 min; 96% ethanol for 2 min; 70% ethanol for 2 min; and were then air dried.

cDNA Synthesis

The following experiments demonstrate that mRNA captured on the array from the tissue sample sections may be used as template for cDNA synthesis.

cDNA Synthesis on Chip

A 16 well mask and Chip Clip slide holder from Whatman was attached to a CodeLink slide. The SuperScript™ III One-step RT-PCR System with Platinum® Taq DNA Polymerase from Invitrogen was used when performing the cDNA synthesis. For each reaction 25 μL 2×reaction mix (SuperScript™ III One-step RT-PCR System with Platinum® Taq DNA Polymerase, Invitrogen), 22.5 μL H₂O and 0.5 μL 100×BSA were mixed and heated to 50° C. SuperScript III/Platinum Taq enzyme mix was added to the reaction mix, 2 μL per reaction, and 50 μL of the reaction mix was added to each well on the chip. The chip was incubated at 50° C. for 30 min (Thermomixer Comfort, Eppendorf).

The reaction mix was removed from the wells and the slide was washed with: 2×SSC, 0.1% SDS at 50° C. for 10

min; 0.2×SSC at room temperature for 1 min; and 0.1×SSC at room temperature for 1 min. The chip was then spin dried.

In the case of FFPE tissue sections, the sections could now be stained and visualized before removal of the tissue, see below section on visualization.

Visualization

Hybridization of Fluorescent Marker Probes Prior to Staining

Prior to tissue application fluorescent marker probes were hybridized to features comprising marker oligonucleotides printed on the capture probe array. The fluorescent marker probes aid in the orientation of the resulting image after tissue visualization, making it possible to combine the image with the resulting expression profiles for individual capture probe "tag" (positional domain) sequences obtained after sequencing. To hybridize fluorescent probes a hybridization solution containing 4×SSC and 0.1% SDS, 2 μM detection probe (P) was incubated for 4 min at 50° C. Meanwhile the in-house array was attached to a ChipClip (Whatman). The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 μL of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry.

General Histological Staining of FFPE Tissue Sections Prior to or Post cDNA Synthesis

FFPE tissue sections immobilized on capture probe arrays were washed and rehydrated after deparaffinization prior to cDNA synthesis as described previously, or washed after cDNA synthesis as described previously. They are then treated as follows: incubate for 3 minutes in Hematoxylin; rinse with deionized water; incubate 5 minutes in tap water; rapidly dip 8 to 12 times in acid ethanol; rinse 2×1 minute in tap water; rinse 2 minutes in deionized water; incubate 30 seconds in Eosin; wash 3×5 minutes in 95% ethanol; wash 3×5 minutes in 100% ethanol; wash 3×10 minutes in xylene (can be done overnight); place coverslip on slides using DPX; dry slides in the hood overnight.

General Immunohistochemistry Staining of a Target Protein in FFPE Tissue Sections Prior to or Post cDNA Synthesis

FFPE tissue sections immobilized on capture probe arrays were washed and rehydrated after deparaffinization prior to cDNA synthesis as described previously, or washed after cDNA synthesis as described previously. They were then treated as follows without being allowed to dry during the whole staining process: sections were incubated with primary antibody (dilute primary antibody in blocking solution comprising 1× Tris Buffered Saline (50 mM Tris, 150 mM NaCl, pH 7.6), 4% donkey serum and 0.1% triton-x) in a wet chamber overnight at RT; rinse three times with 1×TBS; incubate section with matching secondary antibody conjugated to a fluorochrome (FITC, Cy3 or Cy5) in a wet chamber at RT for 1 hour. Rinse 3× with 1×TBS, remove as much as possible of TBS and mount section with ProLong Gold +DAPI (Invitrogen) and analyze with fluorescence microscope and matching filter sets.

Removal of Residual Tissue

Frozen Tissue

For fresh frozen mouse brain tissue the washing step directly following cDNA synthesis was enough to remove the tissue completely.

FFPE Tissue

The slides with attached formalin fixed paraffinized mouse brain tissue sections were attached to ChipClip slide

holders and 16 well masks (Whatman). For each 150 μ l Proteinase K Digest Buffer from the RNeasy FFPE kit (Qiagen), 10 μ l Proteinase K Solution (Qiagen) was added. 50 μ l of the final mixture was added to each well and the slide was incubated at 56° C. for 30 min.

Capture Probe (cDNA) Release

Capture Probe Release with Uracil Cleaving USER Enzyme Mixture in PCR Buffer (Covalently Attached Probes)

A 16 well mask and CodeLink slide was attached to the ChipClip holder (Whatman). 50 μ l of a mixture containing 1 \times FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ (Roche), 200 μ M dNTPs (New England Biolabs) and 0.1 U/1 μ l USER Enzyme (New England Biolabs) was heated to 37° C. and was added to each well and incubated at 37° C. for 30 min with mixing (3 seconds at 300 rpm, 6 seconds at rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released cDNA and probes was then recovered from the wells with a pipette.

Capture Probe Release with Uracil Cleaving USER Enzyme Mixture in TdT (Terminal Transferase) Buffer (Covalently Attached Probes)

50 μ l of a mixture containing: 1 \times TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com); 0.1 μ g/ μ l BSA (New England Biolabs); and 0.1 U/ μ l USER Enzyme (New England Biolabs) was heated to 37° C. and was added to each well and incubated at 37° C. for 30 min with mixing (3 seconds at 300 rpm, 6 seconds at rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released cDNA and probes was then recovered from the wells with a pipette.

Capture Probe Release with Boiling Hot Water (Covalently Attached Probes)

A 16 well mask and CodeLink slide was attached to the ChipClip holder (Whatman). 50 μ l of 99° C. water was pipetted into each well. The 99° C. water was allowed to react for 30 minutes. The reaction mixture containing the released cDNA and probes was then recovered from the wells with a pipette.

Capture Probe Release with Heated PCR Buffer (Hybridized In Situ Synthesized Capture Probes, i.e. Capture Probes Hybridized to Surface Probes)

50 μ l of a mixture containing: 1 \times TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com); 0.1 μ g/ μ l BSA (New England Biolabs); and 0.1 U/ μ l USER Enzyme (New England Biolabs) was preheated to 95° C. The mixture was then added to each well and incubated for 5 minutes at 95° C. with mixing (3 seconds at 300 rpm, 6 seconds at rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released probes was then recovered from the wells.

Capture Probe Release with Heated TdT (Terminal Transferase) Buffer (Hybridized In Situ Synthesized Capture Probes, i.e. Capture Probes Hybridized to Surface Probes)

50 μ l of a mixture containing: 1 \times TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com); 0.1 μ g/ μ l BSA (New England Biolabs); and 0.1 U/ μ l USER Enzyme (New England Biolabs) was preheated to 95° C. The mixture was then added to each well and incubated for 5 minutes at 95° C. with mixing (3 seconds at 300 rpm, 6 seconds at rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released probes was then recovered from the wells.

The efficacy of treating the array with the USER enzyme and water heated to 99° C. can be seen in FIG. 3. Enzymatic cleavage using the USER enzyme and the RsaI enzyme was performed using the “in-house” arrays described above (FIG. 4). Hot water mediated release of DNA surface probes was performed using commercial arrays manufactured by Agilent (see FIG. 5).

Probe Collection and Linker Introduction

The experiments demonstrate that first strand cDNA released from the array surface may be modified to produce double stranded DNA and subsequently amplified.

Whole Transcriptome Amplification by the Picoplex Whole Genome Amplification Kit (Capture Probe Sequences Including Positional Domain (Tag) Sequences not Retained at the Edge of the Resulting dsDNA)

Capture probes were released with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached capture probes) or with heated PCR buffer (hybridized in situ synthesized capture probes, i.e. capture probes hybridized to surface probes).

The released cDNA was amplified using the Picoplex (Rubicon Genomics) random primer whole genome amplification method, which was carried out according to manufacturers instructions.

Whole Transcriptome Amplification by dA Tailing with Terminal Transferase (TdT) (Capture Probe Sequences Including Positional Domain (Tag) Sequences Retained at the End of the Resulting dsDNA)

Capture probes were released with uracil cleaving USER enzyme mixture in TdT (terminal transferase) buffer (covalently attached capture probes) or with heated TdT (terminal transferase) buffer (hybridized in situ synthesized capture probes, i.e. capture probes hybridized to surface probes).

38 μ l of cleavage mixture was placed in a clean 0.2 ml PCR tube. The mixture contained: 1 \times TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com), 0.1 μ g/ μ l BSA (New England Biolabs); 0.1 U/ μ l USER Enzyme (New England Biolabs) (not for heated release); released cDNA (extended from surface probes); and released surface probes. To the PCR tube, 0.5 μ l RNase H (5 U/ μ l, final concentration of 0.06 U/ μ l), 1 μ l TdT (20 U/ μ l, final concentration of 0.5 U/ μ l), and 0.5 μ l dATPs (100 mM, final concentration of 1.25 mM), were added. For dA tailing, the tube was incubated in a thermocycler (Applied Biosystems) at 37° C. for 15 min followed by an inactivation of TdT at 70° C. for 10 min. After dA tailing, a PCR master mix was prepared. The mix contained: 1 \times Faststart HiFi PCR Buffer (pH 8.3) with 1.8 mM MgCl₂ (Roche); 0.2 mM of each dNTP (Fermentas); 0.2 μ M of each primer, A (complementary to the amplification domain of the capture probe) and B₋(dT)₂₄ (Eurofins MWG Operon) (complementary to the poly-A tail to be added to the 3' end of the first cDNA strand); and 0.1 U/ μ l Faststart HiFi DNA polymerase (Roche). 23 μ l of PCR Master mix was placed into nine clean 0.2 ml PCR tubes. 2 μ l of dA tailing mixture were added to eight of the tubes, while 2 μ l water (RNase/DNase free) was added to the last tube (negative control). PCR amplification was carried out with the following program: Hot start at 95° C. for 2 minutes, second strand synthesis at 50° C. for 2 minutes and 72° C. for 3 minutes, amplification with 30 PCR cycles at 95° C. for 30 seconds, 65° C. for 1 minutes, 72° C. for 3 minutes, and a final extension at 72° C. for 10 minutes.

Post-Reaction Cleanup and Analysis

Four amplification products were pooled together and were processed through a Qiaquick PCR purification column

(Qiagen) and eluted into 30 μ l EB (10 mM Tris-Cl, pH 8.5). The product was analyzed on a Bioanalyzer (Agilent). A DNA 1000 kit was used according to manufacturers instructions.

Sequencing

Illumina Sequencing

dsDNA library for Illumina sequencing using sample indexing was carried out according to manufacturers instructions. Sequencing was carried out on an HiSeq2000 platform (Illumina).

Bioinformatics

Obtaining Digital Transcriptomic Information from Sequencing Data from Whole Transcriptome Libraries Amplified Using the dA Tailing Terminal Transferase Approach

The sequencing data was sorted through the FastX toolkit FASTQ Barcode splitter tool into individual files for the respective capture probe positional domain (tag) sequences. Individually tagged sequencing data was then analyzed through mapping to the mouse genome with the Tophat mapping tool. The resulting SAM file was processed for transcript counts through the HTseq-count software.

Obtaining Digital Transcriptomic Information from Sequencing Data from Whole Transcriptome Libraries Amplified Using the Picoplex Whole Genome Amplification Kit Approach

The sequencing data was converted from FASTQ format to FASTA format using the FastX toolkit FASTQ-to-FASTA converter. The sequencing reads was aligned to the capture probe positional domain (tag) sequences using Blastn and the reads with hits better than 1×10^{-6} to one of tag sequences were sorted out to individual files for each tag sequence respectively. The file of tag sequence reads was then aligned using Blastn to the mouse transcriptome, and hits were collected.

Combining Visualization Data and Expression Profiles

The expression profiles for individual capture probe positional domain (tag) sequences are combined with the spatial information obtained from the tissue sections through staining. Thereby the transcriptomic data from the cellular compartments of the tissue section can be analyzed in a directly comparative fashion, with the availability to distinguish distinct expression features for different cellular subtypes in a given structural context

EXAMPLE 2

FIGS. 8 to 12 show successful visualisation of stained FFPE mouse brain tissue (olfactory bulb) sections on top of a bar-coded transcriptome capture array, according to the general procedure described in Example 1. As compared with the experiment with fresh frozen tissue in Example 1, FIG. 8 shows better morphology with the FFPE tissue. FIGS. 9 and 10 show how tissue may be positioned on different types of probe density arrays.

EXAMPLE 3

Whole Transcriptome Amplification by Random Primer Second Strand Synthesis Followed by Universal Handle Amplification (Capture Probe Sequences Including Tag Sequences Retained at the End of the Resulting dsDNA)

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached probes)

OR

Following capture probe release with heated PCR buffer (hybridized in situ synthesized capture probes)

1 μ l RNase H (5 U/ μ l) was added to each of two tubes, final concentration of 0.12 U/ μ l, containing 40 μ l $1 \times$ Faststart HiFi PCR Buffer (pH 8.3) with 1.8 mM $MgCl_2$ (Roche, www.roche-applied-science.com), 0.2 mM of each dNTP (Fermentas, www.fermentas.com), 0.1 μ g/ μ l BSA (New England Biolabs, www.neb.com), 0.1 U/ μ l USER Enzyme (New England Biolabs), released cDNA (extended from surface probes) and released surface probes. The tubes were incubated at 37° C. for 30 min followed by 70° C. for 20 min in a thermo cycler (Applied Biosystems, www.appliedbiosystems.com). 1 μ l Klenow Fragment (3' to 5' exo minus) (Illumina, www.illumina.com) and 1 μ l handle coupled random primer (10 μ M) (Eurofins MWG Operon, www.eurofinsdna.com) was added to the two tubes (B_R8 (octamer) to one of the tubes and B_R6 (hexamer) to the other tube), final concentration of 0.23 μ M. The two tubes were incubated at 15° C. for 15 min, 25° C. for 15 min, 37° C. for 15 min and finally 75° C. for 20 min in a thermo cycler (Applied Biosystems). After the incubation, 1 μ l of each primer, A_P and B (10 μ M) (Eurofins MWG Operon), was added to both tubes, final concentration of 0.24M each. 1 μ l Faststart HiFi DNA polymerase (5 U/ μ l) (Roche) was also added to both tubes, final concentration of 0.11 U/ μ l. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes. After the amplification, 40 μ l from each of the two tubes were purified with Qiaquick PCR purification columns (Qiagen, www.qiagen.com) and eluted into 30 μ l EB (10 mM Tris-Cl, pH 8.5). The Purified products were analyzed with a Bioanalyzer (Agilent, www.home.agilent.com), DNA 7500 kit were used. The results are shown in FIGS. 13 and 14.

This Example demonstrates the use of random hexamer and random octamer second strand synthesis, followed by amplification to generate the population from the released cDNA molecules.

EXAMPLE 4

Amplification of ID-Specific and Gene Specific Products after cDNA Synthesis and Probe Collection

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached probes).

The cleaved cDNA was amplified in final reaction volumes of 10 μ l. 7 μ l cleaved template, 1 μ l ID-specific forward primer (2 μ M), 1 μ l gene-specific reverse primer (2 μ M) and 1 μ l FastStart High Fidelity Enzyme Blend in $1.4 \times$ FastStart High Fidelity Reaction Buffer with 1.8 mM $MgCl_2$ to give a final reaction of 10 μ l with $1 \times$ FastStart High Fidelity Reaction Buffer with 1.8 mM $MgCl_2$ and 1 U FastStart High Fidelity Enzyme Blend. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes.

Primer sequences, resulting in a product of approximately 250 bp,

Beta-2 microglobulin (B2M) primer (SEQ ID NO: 43) 5'-TGGGGGTGAGAATTGCTAAG-3'

ID-1 primer (SEQ ID NO: 44) 5'-CCTTCTCCTTCTCCTTACC-3'

ID-5 primer (SEQ ID NO: 45) 5'-GTCCTCTATTCCGTACCAT-3'

ID-20 primer (SEQ ID NO: 46) 5'-CTGCTTCTTCTGGAACTCA-3'

The results are shown in FIG. 15. This shows successful amplification of ID-specific and gene-specific products using two different ID primers (i.e. specific for ID tags positioned at different locations on the microarray and the same gene specific primer from a brain tissue covering all the probes. Accordingly this experiment establishes that products may be identified by an ID tag-specific or target nucleic acid specific amplification reaction. It is further established that different ID tags may be distinguished. A second experiment, with tissue covering only half of the ID probes (i.e. capture probes) on the array resulted in a positive result (PCR product) for spots that were covered with tissue.

EXAMPLE 5

Spatial Genomics Background.

The method has as its purpose to capture DNA molecules from a tissue sample with retained spatial resolution, making it possible to determine from what part of the tissue a particular DNA fragment stems.

Method.

The principle of the method is to use microarrays with immobilized DNA oligos (capture probes) carrying spatial labeling tag sequences (positional domains). Each feature of oligos of the microarray carries a 1) a unique labeling tag (positional domain) and 2) a capture sequence (capture domain). Keeping track of where which labeling tag is geographically placed on the array surface makes it possible to extract positional information in two dimensions from each labeling tag. Fragmented genomic DNA is added to the microarray, for instance through the addition of a thin section of FFPE treated tissue. The genomic DNA in this tissue section is pre-fragmented due to the fixation treatment.

Once the tissue slice has been placed on the array, a universal tailing reaction is carried out through the use of a terminal transferase enzyme. The tailing reaction adds polydA tails to the protruding 3' ends of the genomic DNA fragments in the tissue. The oligos on the surface are blocked from tailing by terminal transferase through a hybridized and 3' blocked polydA probe.

Following the terminal transferase tailing, the genomic DNA fragments are able to hybridize to the spatially tagged oligos in their vicinity through the polydA tail meeting the polydT capture sequence on the surface oligos. After hybridization is completed a strand displacing polymerase such as Klenow exo- can use the oligo on the surface as a primer for creation of a new DNA strand complementary to the hybrid-

ized genomic DNA fragment. The new DNA strand will now also contain the positional information of the surface oligo's labeling tag.

As a last step the newly generated labeled DNA strands are cleaved from the surface through either enzymatic means, denaturation or physical means. The strands are then collected and can be subjected to downstream amplification of the entire set of strands through introduction of universal handles, amplification of specific amplicons, and/or sequencing.

FIG. 16 is a schematic illustration of this process.

Materials and Methods

Preparation of In-House Printed Microarray with 5' to 3' Oriented Probes

20 DNA-capture oligos with individual tag sequences (Table 1) were spotted on glass slides to function as capture probes. The probes were synthesized with a 5'-terminus amino linker with a C6 spacer. All probes were synthesized by Sigma-Aldrich (St. Louis, MO, USA). The DNA-capture probes were suspended at a concentration of 20 μ M in 150 mM sodium phosphate, pH 8.5 and were spotted using a Nanoplotter NP2.1/E (Gesim, Grosserkmannsdorf, Germany) onto CodeLink™ Activated microarray slides (7.5 cm \times 2.5 cm; Surmodics, Eden Prairie, MN, USA). After printing, surface blocking was performed according to the manufacturer's instructions. The probes were printed in 16 identical arrays on the slide, and each array contained a pre-defined printing pattern. The 16 sub-arrays were separated during hybridization by a 16-pad mask (ChipClip™ Schleicher & Schuell BioScience, Keene, NH, USA).

Preparation of In-House Printed Microarray with 3' to 5' Oriented Probes and Synthesis of 5' to 3' Oriented Capture Probes

Printing of oligos was performed as in the case with 5' to 3' oriented probes above.

To hybridize primers for capture probe synthesis hybridization solution containing 4 \times SSC and 0.1% SDS, 2 μ M extension primer (A_primer) and 2 μ M thread joining primer (p_poly_dT) was incubated for 4 min at 50° C. Meanwhile the in-house array was attached to a ChipClip (Whatman). The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 μ L of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2 \times SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2 \times SSC for 1 min at 300 rpm shake and 3) 0.1 \times SSC for 1 min at 300 rpm shake. The array was then spun dry and placed back in the ChipClip.

For extension and ligation 50 μ L of enzyme mix containing 10 \times Ampligase buffer, 2.5 U AmpliTaq DNA Polymerase Stoffel Fragment (Applied Biosystems), 10 U Ampligase (Epicentre Biotechnologies), dNTPs 2 mM each (Fermentas) and water, is pipetted to each well. The array is subsequently incubated at 55° C. for 30 min. After incubation the array is washed according to previously described array washing method but the first step has the duration of 10 min instead of 6 min.

Hybridization of polydA Probe for Protection of Surface Oligo Capture Sequences from dA Tailing

To hybridize a 3'-biotin blocked polydA probe for protection of the surface oligo capture sequences a hybridization solution containing 4 \times SSC and 0.1% SDS, 2 μ M 3'-bio-polydA was incubated for 4 min at 50° C. Meanwhile the in-house array was attached to a ChipClip (Whatman).

61

The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 µL of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry and placed back in the ChipClip.

Preparation of Formalin-Fixed Paraffin-Embedded (FFPE) Tissue

Mouse brain tissue was fixed in 4% formalin at 4° C. for 24 h. After that it was incubated as follows: 3× incubation in 70% ethanol for 1 hour, 1× incubation in 80% ethanol for 1 hour, 1× incubation in 96% ethanol for 1 hour, 3× incubation in 100% ethanol for 1 hour, 2× incubation in xylene at room temperature for 1 h.

The dehydrated samples were then incubated in liquid low melting paraffin 52-54° C. for up to 3 hours, during which the paraffin in changed once to wash out residual xylene. Finished tissue blocks were then stored at RT. Sections were then cut at 4 µm in paraffin with a microtome onto each capture probe array to be used.

The sections are dried at 37° C. on the array slides for 24 hours and store at RT.

Deparaffinization of FFPE Tissue

Formalin fixed paraffinized mouse brain 10 µm sections attached to CodeLink slides were deparaffinised in xylene twice for 10 min, 99.5% ethanol for 2 min, 96% ethanol for 2 min, 70% ethanol for 2 min and were then air dried.

Universal Tailing of Genomic DNA

For dA tailing a 50 µl reaction mixture containing 1×TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com), 0.1 µg/µl BSA (New England Biolabs), 1 µl TdT (20 U/µl) and 0.5 µl dATPs (100 mM) was prepared. The mixture was added to the array surface and the array was incubated in a thermo cycler (Applied Biosystems) at 37° C. for 15 min followed by an inactivation of TdT at 70° C. for 10 min. After this the temperature was lowered to 50° C. again to allow for hybridization of dA tailed genomic fragments to the surface oligo capture sequences.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry.

Extension of Labeled DNA

A 50 µl reaction mixture containing 50 µl of a mixture containing 1×Klenow buffer, 200 µM dNTPs (New England Biolabs) and 1 µl Klenow Fragment (3' to 5' exo minus) and was heated to 37° C. and was added to each well and incubated at 37° C. for 30 min with mixing (3 s. 300 rpm, 6 s. rest) (Thermomixer comfort; Eppendorf).

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry.

Removal of Residual Tissue

The slides with attached formalin fixed paraffinized mouse brain tissue sections were attached to ChipClip slide holders and 16 well masks (Whatman). For each 150 µl Proteinase K Digest Buffer from the RNeasy FFPE kit (Qiagen) 10 µl Proteinase K Solution (Qiagen) was added.

62

50 µl of the final mixture was added to each well and the slide was incubated at 56° C. for 30 min.

Capture Probe Release with Uracil Cleaving USER Enzyme Mixture in PCR Buffer (Covalently Attached Probes)

A 16 well mask and CodeLink slide was attached to the ChipClip holder (Whatman). 50 µl of a mixture containing 1×FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ (Roche), 200 µM dNTPs (New England Biolabs) and 0.1 U/1 µl USER Enzyme (New England Biolabs) was heated to 37° C. and was added to each well and incubated at 37° C. for 30 min with mixing (3 s. 300 rpm, 6 s. rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released cDNA and probes was then recovered from the wells with a pipette.

Amplification of ID-Specific and Gene Specific Products after Synthesis of Labelled DNA and Probe Collection

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached probes).

The cleaved DNA was amplified in final reaction volumes of 10 µl. 7 µl cleaved template, 1 µl ID-specific forward primer (2 µM), 1 µl gene-specific reverse primer (2 µM) and 1 µl FastStart High Fidelity Enzyme Blend in 1.4×FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ to give a final reaction of 10 µl with 1×FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ and 1 U FastStart High Fidelity Enzyme Blend. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes.

Whole Genome Amplification by Random Primer Second Strand Synthesis Followed by Universal Handle Amplification (Capture Probe Sequences Including Tag Sequences Retained at the End of the Resulting dsDNA)

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached probes).

A reaction mixture containing 40 µl 1× Faststart HiFi PCR Buffer (pH 8.3) with 1.8 mM MgCl₂ (Roche, www.roche-applied-science.com), 0.2 mM of each dNTP (Fermentas, www.fermentas.com), 0.1 µg/µl BSA (New England Biolabs, www.neb.com), 0.1 U/µl USER Enzyme (New England Biolabs), released DNA (extended from surface probes) and released surface probes. The tubes were incubated at 37° C. for 30 min followed by 70° C. for 20 min in a thermo cycler (Applied Biosystems, www.appliedbiosystems.com). 1 µl Klenow Fragment (3' to 5' exo minus) (Illumina, www.illumina.com) and 1 µl handle coupled random primer (10 µM) (Eurofins MWG Operon, www.eurofinsdna.com) was added to the tube. The tube was incubated at 15° C. for 15 min, 25° C. for 15 min, 37° C. for 15 min and finally 75° C. for 20 min in a thermo cycler (Applied Biosystems). After the incubation, 1 µl of each primer, A_P and B (10 µM) (Eurofins MWG Operon), was added to the tube. 1 µl Faststart HiFi DNA polymerase (5 U/µl) (Roche) was also added to the tube. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes. After the amplification, 40 µl from the tube was purified with Qiaquick PCR purification columns (Qiagen, www.qiagen.com) and eluted into 30 µl EB (10 mM Tris-Cl, pH 8.5). The Purified product was analyzed with a Bioanalyzer (Agilent, www.home.agilent.com), DNA 7500 kit were used.

Visualization

Hybridization of Fluorescent Marker Probes Prior to Staining

Prior to tissue application fluorescent marker probes are hybridized to designated marker sequences printed on the capture probe array. The fluorescent marker probes aid in the orientation of the resulting image after tissue visualization, making it possible to combine the image with the resulting expression profiles for individual capture probe tag sequences obtained after sequencing. To hybridize fluorescent probes a hybridization solution containing 4×SSC and 0.1% SDS, 2 μM detection probe (P) was incubated for 4 min at 50° C. Meanwhile the in-house array was attached to a ChipClip (Whatman). The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 μL of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry.

General Histological Staining of FFPE Tissue Sections Prior to or Post Synthesis of Labeled DNA

FFPE tissue sections immobilized on capture probe arrays are washed and rehydrated after deparaffinization prior to synthesis of labeled DNA as described previously. They are then treated as follows: incubate for 3 minutes in Hematoxylin, rinse with deionized water, incubate 5 minutes in tap water, rapidly dip 8 to 12 times in acid ethanol, rinse 2×1 minute in tap water, rinse 2 minutes in deionized water, incubate 30 seconds in Eosin, wash 3×5 minutes in 95% ethanol, wash 3×5 minutes in 100% ethanol, wash 3×10 minutes in xylene (can be done overnight), place coverslip on slides using DPX, dry slides in the hood overnight.

General Immunohistochemistry Staining of a Target Protein in FFPE Tissue Sections Prior to or Post Synthesis of Labeled DNA

FFPE tissue sections immobilized on capture probe arrays are washed and rehydrated after deparaffinization prior to synthesis of labeled DNA as described previously, or washed after synthesis of labeled DNA as described previously. They are then treated as follows without being let to dry during the whole staining process: Dilute primary antibody in blocking solution (1×TBS (Tris Buffered Saline (50 mM Tris, 150 mM NaCl, pH 7.6), 4% donkey serum, 0.1% triton-x), incubate sections with primary antibody in a wet chamber overnight at RT, rinse 3× with 1×TBS, incubate section with matching secondary antibody conjugated to a fluorochrome (FITC, Cy3 or Cy5) in a wet chamber at RT for 1 h, Rinse 3× with 1×TBS, remove as much as possible of TBS and mount section with ProLong Gold +DAPI (Invitrogen) and analyze with fluorescence microscope and matching filter sets.

EXAMPLE 6

This experiment was conducted following the principles of Example 5, but using fragmented genomic DNA on the array rather than tissue. The genomic DNA was pre-fragmented to a mean size of 200 bp and 700 bp respectively. This experiment shows that the principle works. Fragmented genomic DNA is very similar to FFPE tissue.

Amplification of Internal Gene Specific Products after Synthesis of Labeled DNA and Probe Collection

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached

probes) containing 1×FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ (Roche), 200 μM dNTPs (New England Biolabs) and 0.1 U/1 μl USER Enzyme (New England Biolabs).

The cleaved DNA was amplified in a final reaction volume of 50 μl. To 47 μl cleaved template was added 1 μl ID-specific forward primer (10 μM), 1 μl gene-specific reverse primer (10 μM) and 1 μl FastStart High Fidelity Enzyme Blend. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes.

Amplification of Label-Specific and Gene Specific Products after Synthesis of Labelled DNA and Probe Collection

Following capture probe release with uracil cleaving USER enzyme mixture in PCR buffer (covalently attached probes) containing 1×FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ (Roche), 200 μM dNTPs (New England Biolabs) and 0.1 U/1 μl USER Enzyme (New England Biolabs).

The cleaved DNA was amplified in a final reaction volume of 50 μl. To 47 μl cleaved template was added 1 μl label-specific forward primer (10 μM), 1 μl gene-specific reverse primer (10 μM) and 1 μl FastStart High Fidelity Enzyme Blend. PCR amplification were carried out in a thermo cycler (Applied Biosystems) with the following program: Hot start at 94° C. for 2 min, followed by 50 cycles at 94° C. for 15 seconds, 55° C. for 30 seconds, 68° C. for 1 minute, and a final extension at 68° C. for 5 minutes.

```
Forward-Genomic DNA Human Primer
                                     (SEQ ID NO: 47)
5'-GACTGCTCTTTTCACCCATC-3'

Reverse-Genomic DNA Human Primer
                                     (SEQ ID NO: 48)
5'-GGAGCTGCTGGTGACGGG-3'

P-label specific primer
                                     (SEQ ID NO: 49)
5'-ATCTCGACTGCCACTCTGAA-3'
```

The results are shown in FIGS. 17 to 20. The Figures show internal products amplified on the array—the detected peaks in FIGS. 17 and 18 are of the expected size. This thus demonstrates that genomic DNA may be captured and amplified. In FIGS. 19 and 20, the expected product is a smear given that the random fragmentation and terminal transferase labeling of genomic DNA will generate a very diverse sample pool.

EXAMPLE 7

Alternative Synthesis of 5' to 3' Oriented Capture Probes Using Polymerase Extension and Terminal Transferase Tailoring

To hybridize primers for capture probe synthesis hybridization solution containing 4×SSC and 0.1% SDS and 2 μM extension primer (A_primer) was incubated for 4 min at 50° C. Meanwhile the in-house array (see Example 1) was attached to a ChipClip (Whatman). The array was subsequently incubated at 50° C. for 30 min at 300 rpm shake with 50 μL of hybridization solution per well.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake,

2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry and placed back in the ChipClip.

1 μl Klenow Fragment (3' to 5' exo minus) (Illumina, www.illumina.com) together with 10×Klenow buffer, dNTPs 2 mM each (Fermentas) and water, was mixed into a 50 μl reaction and was pipetted into each well.

The array was incubated at 15° C. for 15 min, 25° C. for 15 min, 37° C. for 15 min and finally 75° C. for 20 min in an Eppendorf Thermomixer.

After incubation, the array was removed from the ChipClip and washed with the 3 following steps: 1) 50° C. 2×SSC solution with 0.1% SDS for 6 min at 300 rpm shake, 2) 0.2×SSC for 1 min at 300 rpm shake and 3) 0.1×SSC for 1 min at 300 rpm shake. The array was then spun dry and placed back in the ChipClip.

For dT tailing a 50 μl reaction mixture containing 1×TdT buffer (20 mM Tris-acetate (pH 7.9), 50 mM Potassium Acetate and 10 mM Magnesium Acetate) (New England Biolabs, www.neb.com), 0.1 μg/μl BSA (New England Biolabs), 0.5 μl RNase H (5 U/μl), 1 μl TdT (20 U/μl) and 0.5 μl dTTPs (100 mM) was prepared. The mixture was added to the array surface and the array was incubated in a thermo cycler (Applied Biosystems) at 37° C. for 15 min followed by an inactivation of TdT at 70° C. for 10 min.

EXAMPLE 8

Spatial Transcriptomics Using 5' to 3' High Probe Density Arrays and Formalin-Fixed Frozen (FF-Frozen) Tissue with USER System Cleavage and Amplification Via Terminal Transferase

Array Preparation

Pre-fabricated high-density microarrays chips were ordered from Roche-Nimblegen (Madison, WI, USA). Each capture probe array contained 135,000 features of which 132,640 features carried a capture probe comprising a unique ID-tag sequence (positional domain) and a capture region (capture domain). Each feature was 13×13 μm in size. The capture probes were composed 5' to 3' of a universal domain containing five dUTP bases (a cleavage domain) and a general amplification domain, an ID tag (positional domain) and a capture region (capture domain) (FIG. 22 and Table 2). Each array was also fitted with a frame of marker probes (FIG. 23) carrying a generic 30 bp sequence (Table 2) to enable hybridization of fluorescent probes to help with orientation during array visualization.

Tissue Preparation—Preparation of Formalin-Fixed Frozen Tissue

The animal (mouse) was perfused with 50 ml PBS and 100 ml 4% formalin solution. After excision of the olfactory bulb, the tissue was put into a 4% formalin bath for post-fixation for 24 hrs. The tissue was then sucrose treated in 30% sucrose dissolved in PBS for 24 hrs to stabilize morphology and to remove excess formalin. The tissue was frozen at a controlled rate down to -40° C. and kept at -20° C. between experiments. Similar preparation of tissue post-fixed for 3 hrs or without post-fixation was carried out for a parallel specimen. Perfusion with 2% formalin without post-fixation was also used successfully. Similarly the sucrose treatment step could be omitted. The tissue was mounted into a cryostat for sectioning at 10 μm. A slice of tissue was applied onto each capture probe array to be used. Optionally for better tissue adherence, the array chip was placed at 50° C. for 15 minutes.

Optional Control—Total RNA Preparation from Sectioned Tissue

Total RNA was extracted from a single tissue section (10 μm) using the RNeasy FFPE kit (Qiagen) according to manufacturers instructions. The total RNA obtained from the tissue section was used in control experiments for a comparison with experiments in which the RNA was captured on the array directly from the tissue section. Accordingly, in the case where totalRNA was applied to the array the staining, visualization and degradation of tissue steps were omitted.

On-Chip Reactions

The hybridization of marker probe to the frame probes, reverse transcription, nuclear staining, tissue digestion and probe cleavage reactions were all performed in a 16 well silicone gasket (ArrayIt, Sunnyvale, CA, USA) with a reaction volume of 50 μl per well. To prevent evaporation, the cassettes were covered with plate sealers (In Vitro AB, Stockholm, Sweden).

Optional—Tissue Permeabilization Prior to cDNA Synthesis

For permeabilization using Proteinase K, proteinase K (Qiagen, Hilden, Germany) was diluted to 1 μg/ml in PBS. The solution was added to the wells and the slide incubated at room temperature for 5 minutes, followed by a gradual increase to 80° C. over 10 minutes. The slide was washed briefly in PBS before the reverse transcription reaction.

Alternatively for permeabilization using microwaves, after tissue attachment, the slide was placed at the bottom of a glass jar containing 50 ml 0.2×SSC (Sigma-Aldrich) and was heated in a microwave oven for 1 minute at 800 W. Directly after microwave treatment the slide was placed onto a paper tissue and was dried for 30 minutes in a chamber protected from unnecessary air exposure. After drying, the slide was briefly dipped in water (RNase/DNase free) and finally spin-dried by a centrifuge before cDNA synthesis was initiated.

cDNA Synthesis

For the reverse transcription reaction the SuperScript III One-Step RT-PCR System with Platinum Taq (Life Technologies/Invitrogen, Carlsbad, CA, USA) was used. Reverse transcription reactions contained 1×reaction mix, 1×BSA (New England Biolabs, Ipswich, MA, USA) and 2 μl SuperScript III RT/Platinum Taq mix in a final volume of 50 μl. This solution was heated to 50° C. before application to the tissue sections and the reaction was performed at 50° C. for 30 minutes. The reverse transcription solution was subsequently removed from the wells and the slide was allowed to air dry for 2 hours.

Tissue Visualization

After cDNA synthesis, nuclear staining and hybridization of the marker probe to the frame probes (probes attached to the array substrate to enable orientation of the tissue sample on the array) was done simultaneously. A solution with DAPI at a concentration of 300 nM and marker probe at a concentration of 170 nM in PBS was prepared. This solution was added to the wells and the slide was incubated at room temperature for 5 minutes, followed by brief washing in PBS and spin drying.

Alternatively the marker probe was hybridized to the frame probes prior to placing the tissue on the array. The marker probe was then diluted to 170 nM in hybridization buffer (4×SSC, 0.1% SDS). This solution was heated to 50° C. before application to the chip and the hybridization was performed at 50° C. for 30 minutes at 300 rpm. After hybridization, the slide was washed in 2×SSC, 0.1% SDS at 50° C. and 300 rpm for 10 minutes, 0.2×SSC at 300 rpm for 1 minute and 0.1×SSC at 300 rpm for 1 minute. In that case the staining solution after cDNA synthesis only contained

67

the nuclear DAPI stain diluted to 300 nM in PBS. The solution was applied to the wells and the slide was incubated at room temperature for 5 minutes, followed by brief washing in PBS and spin drying.

The sections were microscopically examined with a Zeiss Axio Imager Z2 and processed with MetaSystems software.

Tissue Removal

The tissue sections were digested using Proteinase K diluted to 1.25 µg/µl in PKD buffer from the RNeasy FFPE Kit (both from Qiagen) at 56° C. for 30 minutes with an interval mix at 300 rpm for 3 seconds, then 6 seconds rest. The slide was subsequently washed in 2×SSC, 0.1% SDS at 50° C. and 300 rpm for 10 minutes, 0.2×SSC at 300 rpm for 1 minute and 0.1×SSC at 300 rpm for 1 minute.

Probe Release

The 16-well Hybridization Cassette with silicone gasket (ArrayIt) was preheated to 37° C. and attached to the Nimblegen slide. A volume of 50 µl of cleavage mixture preheated to 37° C., consisting of Lysis buffer at an unknown concentration (Takara), 0.1 U/µl USER Enzyme (NEB) and 0.1 µg/µl BSA was added to each of wells containing surface immobilized cDNA. After removal of bubbles the slide was sealed and incubated at 37° C. for 30 minutes in a Thermo-mixer comfort with cycled shaking at 300 rpm for 3 seconds with 6 seconds rest in between. After the incubation 45 µl cleavage mixture was collected from each of the used wells and placed into 0.2 ml PCR tubes (FIG. 24).

Library Preparation

Exonuclease Treatment

After cooling the solutions on ice for 2 minutes, Exonuclease I (NEB) was added, to remove unextended cDNA probes, to a final volume of 46.41 and a final concentration of 0.52 U/µl. The tubes were incubated in a thermo cycler (Applied Biosystems) at 37° C. for 30 minutes followed by inactivation of the exonuclease at 80° C. for 25 minutes.

dA-Tailing by Terminal Transferase

After the exonuclease step, 45 µl polyA-tailing mixture, according to manufacturers instructions consisting of TdT Buffer (Takara), 3 mM dATP (Takara) and manufacturers TdT Enzyme mix (TdT and RNase H) (Takara), was added to each of the samples. The mixtures were incubated in a thermocycler at 37° C. for 15 minutes followed by inactivation of TdT at 70° C. for 10 minutes.

Second-Strand Synthesis and PCR-Amplification

After dA-tailing, 23 µl PCR master mix was placed into four new 0.2 ml PCR tubes per sample, to each tube 2 µl sample was added as a template. The final PCRs consisted of 1×Ex Taq buffer (Takara), 200 µM of each dNTP (Takara), 600 nM A_primer (MWG), 600 nM B_dT20VN_primer (MWG) and 0.025 U/µl Ex Taq polymerase (Takara)(Table 2). A second cDNA strand was created by running one cycle in a thermocycler at 95° C. for 3 minutes, 50° C. for 2 minutes and 72° C. for 3 minutes. Then the samples were amplified by running 20 cycles (for library preparation) or 30 cycles (to confirm the presence of cDNA) at 95° C. for 30 seconds, 67° C. for 1 minute and 72° C. for 3 minutes, followed by a final extension at 72° C. for 10 minutes.

Library Cleanup

After amplification, the four PCRs (100 µl) were mixed with 500 µl binding buffer (Qiagen) and placed in a Qiaquick PCR purification column (Qiagen) and spun for 1 minute at 17,900×g in order to bind the amplified cDNA to the membrane. The membrane was then washed with wash buffer (Qiagen) containing ethanol and finally eluted into 50 µl of 10 mM Tris-Cl, pH 8.5.

The purified and concentrated sample was further purified and concentrated by CA-purification (purification by super-

68

paramagnetic beads conjugated to carboxylic acid) with an MBS robot (Magnetic Biosolutions). A final PEG concentration of 10% was used in order to remove fragments below 150-200 bp. The amplified cDNA was allowed to bind to the CA-beads (Invitrogen) for 10 min and were then eluted into 15 µl of 10 mM Tris-Cl, pH 8.5.

Library Quality Analysis

Samples amplified for 30 cycles were analyzed with an Agilent Bioanalyzer (Agilent) in order to confirm the presence of an amplified cDNA library, the DNA High Sensitivity kit or DNA 1000 kit were used depending on the amount of material.

Sequencing Library Preparation

Library Indexing

Samples amplified for 20 cycles were used further to prepare sequencing libraries. An index PCR master mix was prepared for each sample and 23 µl was placed into six 0.2 ml tubes. 2 µl of the amplified and purified cDNA was added to each of the six PCRs as template making the PCRs containing 1×Phusion master mix (Fermentas), 500 nM InPE1.0 (Illumina), 500 nM Index 1-12 (Illumina), and 0.4 nM InPE2.0 (Illumina). The samples were amplified in a thermocycler for 18 cycles at 98° C. for 30 seconds, 65° C. for 30 seconds and 72° C. for 1 minute, followed by a final extension at 72° C. for 5 minutes.

Sequencing Library Cleanup

After amplification, the six PCRs (150 µl) were mixed with 750 µl binding buffer and placed in a Qiaquick PCR purification column and spun for 1 minute at 17,900×g in order to bind the amplified cDNA to the membrane (because of the large sample volume (900 µl), the sample was split in two (each 450 µl) and was bound in two separate steps). The membrane was then washed with wash buffer containing ethanol and finally eluted into 50 µl of 10 mM Tris-Cl, pH 8.5.

The purified and concentrated sample was further purified and concentrated by CA-purification with an MBS robot. A final PEG concentration of 7.8% was used in order to remove fragments below 300-350 bp. The amplified cDNA was allowed to bind to the CA-beads for 10 min and were then eluted into 15 µl of 10 mM Tris-Cl, pH 8.5. Samples were analyzed with an Agilent Bioanalyzer in order to confirm the presence and size of the finished libraries, the DNA High Sensitivity kit or DNA 1000 kit were used according to manufacturers instructions depending on the amount of material (FIG. 25).

Sequencing

The libraries were sequenced on the Illumina Hiseq2000 or Miseq depending on desired data throughput according to manufacturers instructions. Optionally for read 2, a custom sequencing primer B_r2 was used to avoid sequencing through the homopolymeric stretch of 20 T.

Data Analysis

Read 1 was trimmed 42 bases at 5' end. Read 2 was trimmed 25 bases at 5' end (optionally no bases were trimmed from read 2 if the custom primer was used). The reads were then mapped with bowtie to the repeat masked *Mus musculus* 9 genome assembly and the output was formatted in the SAM file format. Mapped reads were extracted and annotated with UCSC refGene gene annotations. Indexes were retrieved with 'indexFinder' (an inhouse software for index retrieval). A mongo DB database was then created containing information about all caught transcripts and their respective index position on the chip.

A matlab implementation was connected to the database and allowed for spatial visualization and analysis of the data (FIG. 26).

Optionally the data visualization was overlaid with the microscopic image using the fluorescently labelled frame probes for exact alignment and enabling spatial transcriptomic data extraction.

EXAMPLE 9

Spatial Transcriptomics Using 3' to 5' High Probe Density Arrays and FFPE Tissue with MutY System Cleavage and Amplification Via TdT

Array Preparation

Pre-fabricated high-density microarrays chips were ordered from Roche-Nimblegen (Madison, WI, USA). Each used capture probe array contained 72 k features out of which 66,022 contained a unique ID-tag complementary sequence. Each feature was 16×16 μm in size. The capture probes were composed 3' to 5' in the same way as the probes used for the in-house printed 3' to 5' arrays with the exception to 3 additional bases being added to the upper (P') general handle of the probe to make it a long version of P', LP' (Table 2). Each array was also fitted with a frame of probes carrying a generic 30 bp sequence to enable hybridization of fluorescent probes to help with orientation during array visualization.

Synthesis of 5' to 3' Oriented Capture Probes

The synthesis of 5' to 3' oriented capture probes on the high-density arrays was carried out as in the case with in-house printed arrays, with the exception that the extension and ligation steps were carried out at 55° C. for 15 mins followed by 72° C. for 15 mins. The A-handle probe (Table 2) included an A/G mismatch to allow for subsequent release of probes through the MutY enzymatic system described below. The P-probe was replaced by a longer LP version to match the longer probes on the surface.

Preparation of Formalin-Fixed Paraffin-Embedded Tissue and Deparaffinization

This was carried out as described above in the in-house protocol.

cDNA Synthesis and Staining

cDNA synthesis and staining was carried out as in the protocol for 5' to 3' oriented high-density Nimblegen arrays with the exception that biotin labeled dCTPs and dATPs were added to the cDNA synthesis together with the four regular dNTPs (each was present at 25×times more than the biotin labeled ones).

Tissue Removal

Tissue removal was carried out in the same way as in the protocol for 5' to 3' oriented high-density Nimblegen arrays described in Example 8.

Probe Cleavage by MutY

A 16-well Incubation chamber with silicone gasket (ArrayIT) was preheated to 37° C. and attached to the Codelink slide. A volume of 50 μl of cleavage mixture preheated to 37° C., consisting of 1×Endonuclease VIII Buffer (NEB), 10 U/μl MutY (Trevigen), 10 U/μl Endonuclease VIII (NEB), 0.1 μg/μl BSA was added to each of wells containing surface immobilized cDNA. After removal of bubbles the slide was sealed and incubated at 37° C. for 30 minutes in a Thermomixer comfort with cycled shaking at 300 rpm for 3 seconds with 6 seconds rest in between. After the incubation, the plate sealer was removed and 40 μl cleavage mixture was collected from each of the used wells and placed into a PCR plate.

Library Preparation

Biotin-Streptavidin Mediated Library Cleanup

To remove unextended cDNA probes and to change buffer, the samples were purified by binding the biotin

labeled cDNA to streptavidin coated CI-beads (Invitrogen) and washing the beads with 0.1M NaOH (made fresh). The purification was carried out with an MBS robot (Magnetic Biosolutions), the biotin labelled cDNA was allowed to bind to the CI-beads for 10 min and was then eluted into 20 μl of water by heating the bead-water solution to 80° C. to break the biotin-streptavidin binding.

dA-Tailing by Terminal Transferase

After the purification step, 18 μl of each sample was placed into new 0.2 ml PCR tubes and mixed with 22 μl of a polyA-tailing master mix leading to a 40 μl reaction mixture according to manufacturers instructions consisting of lysis buffer (Takara, Cellamp Whole Transcriptome Amplification kit), TdT Buffer (Takara), 1.5 mM dATP (Takara) and TdT Enzyme mix (TdT and RNase H) (Takara). The mixtures were incubated in a thermocycler at 37° C. for 15 minutes followed by inactivation of TdT at 70° C. for 10 minutes.

Second-Strand Synthesis and PCR-Amplification

After dA-tailing, 23 μl PCR master mix was placed into four new 0.2 ml PCR tubes per sample, to each tube 2 μl sample was added as a template. The final PCRs consisted of 1×Ex Taq buffer (Takara), 200 μM of each dNTP (Takara), 600 nM A_primer (MWG), 600 nM B_dT20VN_primer (MWG) and 0.025 U/μl Ex Taq polymerase (Takara). A second cDNA strand was created by running one cycle in a thermo cycler at 95° C. for 3 minutes, 50° C. for 2 minutes and 72° C. for 3 minutes. Then the samples were amplified by running 20 cycles (for library preparation) or 30 cycles (to confirm the presence of cDNA) at 95° C. for 30 seconds, 67° C. for 1 minute and 72° C. for 3 minutes, followed by a final extension at 72° C. for 10 minutes.

Library Cleanup

After amplification, the four PCRs (100 μl) were mixed with 500 μl binding buffer (Qiagen) and placed in a Qiaquick PCR purification column (Qiagen) and spun for 1 minute at 17,900×g in order to bind the amplified cDNA to the membrane. The membrane was then washed with wash buffer (Qiagen) containing ethanol and finally eluted into 50 μl of 10 mM Tris-HCl, pH 8.5.

The purified and concentrated sample was further purified and concentrated by CA-purification (purification by super-paramagnetic beads conjugated to carboxylic acid) with an MBS robot (Magnetic Biosolutions). A final PEG concentration of 10% was used in order to remove fragments below 150-200 bp. The amplified cDNA was allowed to bind to the CA-beads (Invitrogen) for 10 min and were then eluted into 15 μl of 10 mM Tris-HCl, pH 8.5.

Second PCR-Amplification

The final PCRs consisted of 1×Ex Taq buffer (Takara), 200 μM of each dNTP (Takara), 600 nM A_primer (MWG), 600 nM B_primer (MWG) and 0.025 U/μl Ex Taq polymerase (Takara). The samples were heated to 95° C. for 3 minutes, and then amplified by running 10 cycles at 95° C. for 30 seconds, 65° C. for 1 minute and 72° C. for 3 minutes, followed by a final extension at 72° C. for 10 minutes.

Second Library Cleanup After amplification, the four PCRs (100 μl) were mixed with 500 μl binding buffer (Qiagen) and placed in a Qiaquick PCR purification column (Qiagen) and spun for 1 minute at 17,900×g in order to bind the amplified cDNA to the membrane. The membrane was then washed with wash buffer (Qiagen) containing ethanol and finally eluted into 50 μl of 10 mM Tris-HCl, pH 8.5.

The purified and concentrated sample was further purified and concentrated by CA-purification (purification by super-paramagnetic beads conjugated to carboxylic acid) with an MBS robot (Magnetic Biosolutions). A final PEG concen-

tration of 10% was used in order to remove fragments below 150-200 bp. The amplified cDNA was allowed to bind to the CA-beads (Invitrogen) for 10 min and were then eluted into 15 μ l of 10 mM Tris-HCl, pH 8.5.

Sequencing Library Preparation Library Indexing

Samples amplified for 20 cycles were used further to prepare sequencing libraries. An index PCR master mix was prepared for each sample and 23 μ l was placed into six 0.2 ml tubes. 2 μ l of the amplified and purified cDNA was added to each of the six PCR as template making the PCR containing 1 \times Phusion master mix (Fermentas), 500 nM InPE1.0 (Illumina), 500 nM Index 1-12 (Illumina), and 0.4 nM InPE2.0 (Illumina). The samples were amplified in a thermo cycler for 18 cycles at 98 $^{\circ}$ C. for 30 seconds, 65 $^{\circ}$ C. for 30 seconds and 72 $^{\circ}$ C. for 1 minute, followed by a final extension at 72 $^{\circ}$ C. for 5 minutes.

Sequencing Library Cleanup

After amplification, the samples was purified and concentrated by CA-purification with an MBS robot. A final PEG concentration of 7.8% was used in order to remove fragments below 300-350 bp. The amplified cDNA was allowed to bind to the CA-beads for 10 min and were then eluted into 15 μ l of 10 mM Tris-HCl, pH 8.5.

10 μ l of the amplified and purified samples were placed on a Caliper XT chip and fragments between 480 bp and 720 bp were cut out with the Caliper XT (Caliper). Samples were analyzed with an Agilent Bioanalyzer in order to confirm the presence and size of the finished libraries, the DNA High Sensitivity kit was used.

Sequencing and Data Analysis

Sequencing and Bioinformatic was carried out in the same way as in the protocol for 5' to 3' oriented high-density Nimblegen arrays described in Example 8. However, in the data analysis, read 1 was not used in the mapping of transcripts. Specific Olfr transcripts could be sorted out using the Matlab visualization tool (FIG. 27).

EXAMPLE 10

Spatial Transcriptomics Using in House Printed 41-Tag Microarray with 5' to 3' Oriented Probes and Formalin-Fixed Frozen (FF-Frozen) Tissue with Permeabilization Through ProteinaseK or Microwaving with USER System Cleavage and Amplification Via TdT

Array Preparation

In-house arrays were printed as previously described but with a pattern of 41 unique ID-tag probes with the same composition as the probes in the 5' to 3' oriented high-density array in Example 8 (FIG. 28).

All other steps were carried out in the same way as in the protocol described in Example 8.

EXAMPLE 11

Alternative Method for Performing the cDNA Synthesis Step

cDNA synthesis on chip as described above can also be combined with template switching to create a second strand by adding a template switching primer to the cDNA synthesis reaction (Table 2). The second amplification domain is introduced by coupling it to terminal bases added by the reverse transcriptase at the 3' end of the first cDNA strand, and primes the synthesis of the second strand. The library can be readily amplified directly after release of the double-stranded complex from the array surface.

EXAMPLE 12

Spatial Genomics Using in House Printed 41-Tag Microarray with 5' to 3' Oriented Probes and Fragmented Poly-A Tailed gDNA with USER System Cleavage and Amplification Via TdT-Tailing or Translocation Specific Primers

Array Preparation

In-house arrays were printed using Codelink slides (Surremodics) as previously described but with a pattern of 41 unique ID-tag probes with the same composition as the probes in the 5' to 3' oriented high-density in Example 8.

Total DNA Preparation from Cells

DNA Fragmentation

Genomic DNA (gDNA) was extracted by DNeasy kit (Qiagen) according to the manufacturer's instructions from A431 and U2OS cell lines. The DNA was fragmented to 500 bp on a Covaris sonicator (Covaris) according to manufacturer's instructions.

The sample was purified and concentrated by CA-purification (purification by super-paramagnetic beads conjugated to carboxylic acid) with an MBS robot (Magnetic Biosolutions). A final PEG concentration of 10% was used in order to remove fragments below 150-200 bp. The fragmented DNA was allowed to bind to the CA-beads (Invitrogen) for 10 min and were then eluted into 15 μ l of 10 mM Tris-HCl, pH 8.5.

Optional Control—Spiking of Different Cell Lines

Through spiking of A431 DNA into U2OS DNA different levels of capture sensitivity can be measured, such as from spiking of 1%, 10% or 50% of A431 DNA.

dA-Tailing by Terminal Transferase

A 45 μ l polyA-tailing mixture, according to manufacturer's instructions consisting of TdT Buffer (Takara), 3 mM dATP (Takara) and TdT Enzyme mix (TdT and RNase H) (Takara), was added to 0.5 μ g of fragmented DNA. The mixtures were incubated in a thermocycler at 37 $^{\circ}$ C. for 30 minutes followed by inactivation of TdT at 80 $^{\circ}$ C. for 20 minutes. The dA-tailed fragments were then cleaned through a Qiaquick (Qiagen) column according to manufacturer's instructions and the concentration was measured using the Qubit system (Invitrogen) according to manufacturer's instructions.

On-Chip Experiments

The hybridization, second strand synthesis and cleavage reactions were performed on chip in a 16 well silicone gasket (ArrayIt, Sunnyvale, CA, USA). To prevent evaporation, the cassettes were covered with plate sealers (In Vitro AB, Stockholm, Sweden).

Hybridization

117 ng of DNA was deposited onto a well on a prewarmed array (50 $^{\circ}$ C.) in a total volume of 45 μ l consisting of 1 \times NEB buffer (New England Biolabs) and 1 \times BSA. The mixture was incubated for 30 mins at 50 $^{\circ}$ C. in a Thermomixer Comfort (Eppendorf) fitted with an MTP block at 300 rpm shake.

Second Strand Synthesis

Without removing the hybridization mixture, 15 μ l of a Klenow extension reaction mixture consisting of 1 \times NEB buffer 1.5 μ l Klenow polymerase, and 3.75 μ l dNTPs (2 mM each) was added to the well. The reaction mixture was incubated in a Thermomixer Comfort (Eppendorf) 37 $^{\circ}$ C. for 30 mins without shaking.

The slide was subsequently washed in 2 \times SSC, 0.1% SDS at 50 $^{\circ}$ C. and 300 rpm for 10 minutes, 0.2 \times SSC at 300 rpm for 1 minute and 0.1 \times SSC at 300 rpm for 1 minute.

Probe Release

A volume of 50 μ l of a mixture containing 1 \times FastStart High Fidelity Reaction Buffer with 1.8 mM MgCl₂ (Roche), 200 μ M dNTPs (New England Biolabs), 1 \times BSA and 0.1 U/1 μ l USER Enzyme (New England Biolabs) was heated to 37° C. and was added to each well and incubated at 37° C. for 30 min with mixing (3 seconds at 300 rpm, 6 seconds at rest) (Thermomixer comfort; Eppendorf). The reaction mixture containing the released DNA which was then recovered from the wells with a pipette.

Library Preparation

Amplification Reaction

Amplification was carried out in 10 μ l reactions consisting of 7.5 μ l released sample, 1 μ l of each primer and 0.5 μ l enzyme (Roche, FastStart HiFi PCR system). The reaction was cycled as 94° C. for 2 mins, one cycle of 94° C. 15 sec, 55° C. for 2 mins, 72° C. for 2 mins, 30 cycles of 94° C. for 15 secs, 65° C. for 30 secs, 72° C. for 90 secs, and a final elongation at 72° C. for 5 mins.

In the preparation of a library for sequencing the two primers consisted of the surface probe A-handle and either of a specific translocation primer (for A431) or a specific SNP primer coupled to the B-handle (Table 2).

Library Cleanup

The purified and concentrated sample was further purified and concentrated by CA-purification (purification by superparamagnetic beads conjugated to carboxylic acid) with an MBS robot (Magnetic Biosolutions). A final PEG concentration of 10% was used in order to remove fragments below 150-200 bp. The amplified DNA was allowed to bind to the CA-beads (Invitrogen) for 10 min and was then eluted into 15 μ l of 10 mM Tris-HCl, pH 8.5.

Library Quality Analysis

Samples were analyzed with an Agilent Bioanalyzer (Agilent) in order to confirm the presence of an amplified DNA library, the DNA High Sensitivity kit or DNA 1000 kit were used depending on the amount of material.

Library Indexing

Samples amplified for 20 cycles were used further to prepare sequencing libraries. An index PCR master mix was prepared for each sample and 23 μ l was placed into six 0.2 ml tubes. 2 μ l of the amplified and purified cDNA was added to each of the six PCRs as template making the PCRs containing 1 \times Phusion master mix (Fermentas), 500 nM InPE1.0 (Illumina), 500 nM Index 1-12 (Illumina), and 0.4 nM InPE2.0 (Illumina). The samples were amplified in a thermo cycler for 18 cycles at 98° C. for 30 seconds, 65° C. for 30 seconds and 72° C. for 1 minute, followed by a final extension at 72° C. for 5 minutes.

Sequencing Library Cleanup

The purified and concentrated sample was further purified and concentrated by CA-purification with an MBS robot. A final PEG concentration of 7.8% was used in order to remove fragments below 300-350 bp. The amplified DNA was allowed to bind to the CA-beads for 10 min and were then eluted into 15 μ l of 10 mM Tris-Cl, pH 8.5. Samples were analyzed with an Agilent Bioanalyzer in order to confirm the presence and size of the finished libraries, the DNA High Sensitivity kit or DNA 1000 kit were used according to manufacturers instructions depending on the amount of material (FIG. 29).

Sequencing

Sequencing was carried out in the same way as in the protocol for 5' to 3' oriented high-density Nimblegen arrays described in Example 8.

Data Analysis

Data analysis was carried out to determine the sensitivity of capture of the arrayed ID-capture probes. Read 2 was sorted based on its content of either of the translocation or SNP primers. These reads were then sorted per their ID contained in Read 1.

Optional Control—Direct Amplification of Cell-Line Specific Translocations

This was used to measure the capture sensitivity of spiked cell lines directly by PCR. The forward and reverse primers (Table 2) for the A431 translocations were used to try and detect the presence of the translocation in the second strand copied and released material (FIG. 30).

TABLE 2

	Oligos used for spatial transcriptomics and spatial genomics
Example 8	
Nimblegen 5' to 3' arrays with free 3' end Array probes	
	5' to 3'
Probe1 (SEQ ID NO: 50)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTGTCGATATGATGTCGGCTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe2 (SEQ ID NO: 51)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTAATGAGCGGGTTCACTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe3 (SEQ ID NO: 52)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTTGAGGCACCTCTGTTGGGATTTTTTTTTTTTTTTTTTTTTTTTTV
Probe4 (SEQ ID NO: 53)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTATGATAGTTCGCAATTCGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe5 (SEQ ID NO: 54)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTACTTGAGGGTAGAATTTTTTTTTTTTTTTTTTTTTTTTTV
Probe6 (SEQ ID NO: 55)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTATGGCCAATACTGTTACTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe4 (SEQ ID NO: 56)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTCGCTACCTGATTCGACCTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe8 (SEQ ID NO: 57)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTGCCCCACTTTCGCCGAGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe9 (SEQ ID NO: 58)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTAGCAACTTTGAGCAAGATTTTTTTTTTTTTTTTTTTTTTTTTV
Probe10 (SEQ ID NO: 59)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTGCCAATTCGGAATCCGGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe11 (SEQ ID NO: 60)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTTGCCGATCTTGCCCCAAGTAATAAATTTTTTTTTTTTTTTTTV
Probe12 (SEQ ID NO: 61)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTTCGCACTTCGCACTTCCCTAATTCGAGTTTTTTTTTTTTTTTTV
Probe13 (SEQ ID NO: 62)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTTTGCTAAATCTAACCGCCTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe14 (SEQ ID NO: 63)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTGGAAATTAATAAATCTGATGGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe15 (SEQ ID NO: 64)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTCAATACATAGTGTAAAGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe16 (SEQ ID NO: 65)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTATTGACTTTGCGCTCGCACTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe17 (SEQ ID NO: 66)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTAATGACTTTGCGCTCGCACTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe18 (SEQ ID NO: 67)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTAGTAATCTCCCAAGTTCTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe19 (SEQ ID NO: 68)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTGCGCCCTGTAAATCCGCAATTTTTTTTTTTTTTTTTTTTTTTTTV
Probe20 (SEQ ID NO: 69)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTAATGCAAGTAGATGGCTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe21 (SEQ ID NO: 70)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTCCAAGCCAGGTTTATACGTTTTTTTTTTTTTTTTTTTTTTTTV
Probe22 (SEQ ID NO: 71)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTACCTGATTTGCTGTAATAACTTTTTTTTTTTTTTTTTTTTTTTTTV
Probe23 (SEQ ID NO: 72)	UUUUUACACTCTTTCCCTACACGACGCTCTTCCGATCTCAGCGCATCTAATCCCTAATTTTTTTTTTTTTTTTTTTTTTTTTV

TABLE 2-continued

	Oligos used for spatial transcriptomics and spatial genomics
Probe24 (SEQ ID NO: 73)	UUUUUACACTCTTTCCCTACACGACGCTCTCCGATCTCCACGCGTAGGACTAGTTTTTTTTTTTTTTTTTTTTTTT
Probe25 (SEQ ID NO: 74)	UUUUUACACTCTTTCCCTACACGACGCTCTCCGATCTCCGACTCGACTAAGTAAGTAGCGCTTTTTTTTTTTTTTTTTTTT
Frame probe Layout1 (SEQ ID NO: 75)	AAAF TTCGCTGTGCTATCGCGCTTCTGTACC
Fluorescent marker probe PS_1 (SEQ ID NO: 76)	GGTACAGAGCGGATAGCAG-Cy3
Second strand synthesis and first PCR. Amplification handles A_primer (SEQ ID NO: 77)	ACACTCTTTCCCTACACGACGCTCTCCGATCT
B_dt20VN_primer (SEQ ID NO: 78)	AGACGTCGCTCTCCGACTTTTTTTTTTTTTTTTTTTT
Custom sequencing primer B_r2 (SEQ ID NO: 79)	TCA GAC GTG TGC TCT TCC GAT CTT TTT TTT TTT TTT TTT TTT T
Example 9 Nimblegen 3' to 5' arrays with free 5' end Array probes 5' to 3'	
Probe1 (SEQ ID NO: 80)	GCCTTCAGAGTGGCAGTCGAGATCAGCGGCAATCATACGGACAGATCGGAGAGCGGTAGTGTAG
Probe2 (SEQ ID NO: 81)	GCCTTCAGAGTGGCAGTCGAGATCACAAGATGAACCCGGCTCATAGATCGGAGAGCGGTAGTGTAG
Probe3 (SEQ ID NO: 82)	GCCTTCAGAGTGGCAGTCGAGATCCTCCAAAGAGTGCCTCAAGATCGGAAGAGCGGTAGTGTAG
Probe4 (SEQ ID NO: 83)	GCCTTCAGAGTGGCAGTCGAGATCACCGAATGGCGACTAATCATAGATCGGAGAGCGGTAGTGTAG
Probe5 (SEQ ID NO: 84)	GCCTTCAGAGTGGCAGTCGAGATCACAACAATCTACCCCTCAGTAGATCGGAGAGCGGTAGTGTAG
Probe6 (SEQ ID NO: 85)	GCCTTCAGAGTGGCAGTCGAGATCAGATAACAGTATTTGGCCATAGATCGGAAGAGCGGTAGTGTAG
Probe7 (SEQ ID NO: 86)	GCCTTCAGAGTGGCAGTCGAGATCACGGTCGAATCAGGGTAGCGAGATCGGAGAGCGGTAGTGTAG
Probe8 (SEQ ID NO: 87)	GCCTTCAGAGTGGCAGTCGAGATCACACTACGGCAAGTGGCGAGATCGGAGAGCGGTAGTGTAG
Probe9 (SEQ ID NO: 88)	GCCTTCAGAGTGGCAGTCGAGATCACAATCTTGCTCAAAGTTGCTAGATCGGAAGAGCGGTAGTGTAG
Probe10 (SEQ ID NO: 89)	GCCTTCAGAGTGGCAGTCGAGATCACCCGGAATCCGAATGGCAGATCGGAGAGCGGTAGTGTAG
Probe11 (SEQ ID NO: 90)	GCCTTCAGAGTGGCAGTCGAGATCATATGTAATTTACCTTTGGCGAGATCGGAGAGCGGTAGTGTAG
Probe12 (SEQ ID NO: 91)	GCCTTCAGAGTGGCAGTCGAGATCACCCTCGAATAGGAATGCGAAGATCGGAAGAGCGGTAGTGTAG
Probe13 (SEQ ID NO: 92)	GCCTTCAGAGTGGCAGTCGAGATCACGGCGTTAGATTTAGCAAGATCGGAGAGCGGTAGTGTAG
Probe14 (SEQ ID NO: 93)	GCCTTCAGAGTGGCAGTCGAGATCACCCATCAGAATTTAATCCAGATCGGAGAGCGGTAGTGTAG
Probe15 (SEQ ID NO: 94)	GCCTTCAGAGTGGCAGTCGAGATCACCTTAGCACCTATGTAATGAGATCGGAAGAGCGGTAGTGTAG

TABLE 2 - continued

Oligos used for spatial transcriptomics and spatial genomics	
Probe16 (SEQ ID NO: 95)	GCCTTCAGAGTGGCAGTCGAGATCAGTCGGAGCGCAGTCAATAGATCGGAAAGAGCGTAGTGTAG
Probe17 (SEQ ID NO: 96)	GCCTTCAGAGTGGCAGTCGAGATCAGCAACTTGGAGATACTATAGATCGGAAAGAGCGTAGTGTAG
Probe18 (SEQ ID NO: 97)	GCCTTCAGAGTGGCAGTCGAGATCAGTCGGGATTCAGGGCCACAGATCGGAAAGAGCGTAGTGTAG
Probe19 (SEQ ID NO: 98)	GCCTTCAGAGTGGCAGTCGAGATCAGTACCTACCTAAAGAGTGGCCAGATCGGAAAGAGCGTAGTGTAG
Probe20 (SEQ ID NO: 99)	GCCTTCAGAGTGGCAGTCGAGATCACAAGCCAAATCACTTGCATAGATCGGAAAGAGCGTAGTGTAG
Probe21 (SEQ ID NO: 100)	GCCTTCAGAGTGGCAGTCGAGATCACCGTATAAACCGTGGCTTGGAGATCGGAAAGAGCGTAGTGTAG
Probe22 (SEQ ID NO: 101)	GCCTTCAGAGTGGCAGTCGAGATCAGCTTATACAGCAATCAGGTAGATCGGAAAGAGCGTAGTGTAG
Probe23 (SEQ ID NO: 102)	GCCTTCAGAGTGGCAGTCGAGATCACTAGAGGATAGATGGCTGAGATCGGAAAGAGCGTAGTGTAG
Probe24 (SEQ ID NO: 103)	GCCTTCAGAGTGGCAGTCGAGATCACACTAGTCCCTACGGGTGGAAGATCGGAAAGAGCGTAGTGTAG
Probe25 (SEQ ID NO: 104)	GCCTTCAGAGTGGCAGTCGAGATCACGGCTACATACTTAGTCGAGATCGGAAAGAGCGTAGTGTAG
Frame probe	
Layout1 (SEQ ID NO: 105)	AAATTCGCTGCTATCGCGCTTCGTACC
Capture probe	
LP_poly-dTVN (SEQ ID NO: 106)	GTGATCTCGACTGCCACTCTGAATTTTTTTTTTTTTTTTTTTTTV
Amplification handle probe	
A-handle (SEQ ID NO: 107)	ACACTCTTCCCTACAGGAGGCTTCCCGATCT
Second strand synthesis and first PCR amplification handles	
A_primer (SEQ ID NO: 108)	ACACTCTTCCCTACAGGAGGCTTCCCGATCT
B_dt20VN_primer (SEQ ID NO: 109)	AGACGTGCTCTCCGATCTTTTTTTTTTTTTTTTTTTTTV
Second PCR	
A_primer (SEQ ID NO: 110)	ACACTCTTCCCTACAGGAGGCTTCCCGATCT
B_primer (SEQ ID NO: 111)	GTGACTGGAGTTCAGAGGTGGCTCTCCGATCT
Example 11	
Template switching	
Templateswitch_longB (SEQ ID NO: 112)	GTGACTGGAGTTCAGAGGTGGCTCTCCGATCTATrGrGrG
Example 12	
Spatial genomics	
A_primer (SEQ ID NO: 113)	ACACTCTTCCCTACAGGAGGCTTCCCGATCT

-continued

gagacatacc accaagagac	20
SEQ ID NO: 10	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 10	
gtcctctatt ccgtcaccat	20
SEQ ID NO: 11	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 11	
gactgagctc gaacatatgg	20
SEQ ID NO: 12	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 12	
tggaggatg acacagaacg	20
SEQ ID NO: 13	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 13	
ccagcctctc cattacatcg	20
SEQ ID NO: 14	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 14	
aagatctacc agccagccag	20
SEQ ID NO: 15	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 15	
cgaacttcca ctgtctcctc	20
SEQ ID NO: 16	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 16	
ttgcgccttc tccaatacac	20
SEQ ID NO: 17	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 17	
ctcttcttag catgccacct	20
SEQ ID NO: 18	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 18	
accacttctg cattacctcc	20
SEQ ID NO: 19	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	organism = synthetic construct

-continued

SEQUENCE: 19			
acagcctect cttcttctt			20
SEQ ID NO: 20	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 20			
aatcctctcc ttgccagttc			20
SEQ ID NO: 21	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 21			
gatgcctcca cctgtagaac			20
SEQ ID NO: 22	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 22			
gaaggaatgg aggatatcgc			20
SEQ ID NO: 23	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 23			
gatccaagga ccatcgactg			20
SEQ ID NO: 24	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 24			
ccactggaac ctgacaaccg			20
SEQ ID NO: 25	moltype = DNA	length = 20	
FEATURE	Location/Qualifiers		
source	1..20		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 25			
ctgettcttc ctggaactca			20
SEQ ID NO: 26	moltype = DNA	length = 66	
FEATURE	Location/Qualifiers		
source	1..66		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 26			
gcggtcagag tggcagtcga gatcacgcgg caatcatatc ggacagatcg gaagagcgta			60
gtgtag			66
SEQ ID NO: 27	moltype = DNA	length = 66	
FEATURE	Location/Qualifiers		
source	1..66		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 27			
gcggtcagag tggcagtcga gatcacgcgg caatcatatc ggacggctgc tggtaaatag			60
agatca			66
SEQ ID NO: 28	moltype = DNA	length = 23	
FEATURE	Location/Qualifiers		
source	1..23		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 28			
ttcagagtgg cagtcgagat cac			23
SEQ ID NO: 29	moltype = DNA	length = 18	
FEATURE	Location/Qualifiers		

-continued

source	1..18 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 29 gcggaatca tateggac		18
SEQ ID NO: 30 FEATURE source	moltype = DNA length = 22 Location/Qualifiers 1..22 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 30 agatcgaag agcgtagtgt ag		22
SEQ ID NO: 31 FEATURE source	moltype = DNA length = 22 Location/Qualifiers 1..22 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 31 ggctgctggt aaatagat ca		22
SEQ ID NO: 32 FEATURE source	moltype = DNA length = 33 Location/Qualifiers 1..33 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 32 acaactcttc cctacacgac gctctccga tct		33
SEQ ID NO: 33 FEATURE source	moltype = DNA length = 35 Location/Qualifiers 1..35 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 33 aagtgtgaa agttgatcgc tatttaccag cagcc		35
SEQ ID NO: 34 FEATURE modified_base source	moltype = DNA length = 45 Location/Qualifiers 1 mod_base = OTHER note = Phosphorylated 1..45 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 34 gtgatctcga ctgccactct gaattttttt ttttttttt tttvn		45
SEQ ID NO: 35 FEATURE modified_base source	moltype = DNA length = 47 Location/Qualifiers 1 mod_base = OTHER note = Phosphorylated 1..47 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 35 gtgatctcga ctgccactct gaattttttt ttttttttt ttttttt		47
SEQ ID NO: 36 FEATURE source	moltype = DNA length = 21 Location/Qualifiers 1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 36 agacgtgtgc tcttccgatc t		21
SEQ ID NO: 37 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 37 acgtctgtga atagccgcat		20
SEQ ID NO: 38 FEATURE source	moltype = DNA length = 29 Location/Qualifiers 1..29	

-continued

	mol_type = other DNA organism = synthetic construct	
SEQUENCE: 38		
agacgtgtgc tcttccgac tnnnnnnnn		29
SEQ ID NO: 39	moltype = DNA length = 31	
FEATURE	Location/Qualifiers	
source	1..31	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 39		
agacgtgtgc tcttccgac tnnnnnnnn n		31
SEQ ID NO: 40	moltype = DNA length = 43	
FEATURE	Location/Qualifiers	
source	1..43	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 40		
agacgtgtgc tcttccgac ttttttttt ttttttttt tvn		43
SEQ ID NO: 41	moltype = DNA length = 45	
FEATURE	Location/Qualifiers	
source	1..45	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 41		
agacgtgtgc tcttccgac ttttttttt ttttttttt ttttt		45
SEQ ID NO: 42	moltype = DNA length = 53	
FEATURE	Location/Qualifiers	
source	1..53	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 42		
acactctttc cctacacgac gctcttccga tctatctcga ctgccactct gaa		53
SEQ ID NO: 43	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 43		
tgggggtgag aattgctaag		20
SEQ ID NO: 44	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 44		
ccttctcctt ctccttcacc		20
SEQ ID NO: 45	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 45		
gtcctctatt ccgtcaccat		20
SEQ ID NO: 46	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 46		
ctgcttttc ctggaactca		20
SEQ ID NO: 47	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 47		
gactgcttt ttcaccatc		20
SEQ ID NO: 48	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	

-continued

source	1..18		
		mol_type = other DNA	
		organism = synthetic construct	
SEQUENCE: 48			
ggagctgctg gtgcaggg			18
SEQ ID NO: 49		moltype = DNA length = 20	
FEATURE		Location/Qualifiers	
source	1..20		
		mol_type = other DNA	
		organism = synthetic construct	
SEQUENCE: 49			
atctcgactg ccactctgaa			20
SEQ ID NO: 50		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	
modified_base	1..5		
		mod_base = OTHER	
		note = t is u	
SEQUENCE: 50			
tttttacact ctttccctac	acgacgctct tccgatctgt	ccgatgat	tgccgctttt
ttttttttt tttttvn			60
			78
SEQ ID NO: 51		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	
modified_base	1..5		
		mod_base = OTHER	
		note = t is u	
SEQUENCE: 51			
tttttacact ctttccctac	acgacgctct tccgatctat	gagccgggtt	catctttttt
ttttttttt tttttvn			60
			78
SEQ ID NO: 52		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	
modified_base	1..5		
		mod_base = OTHER	
		note = t is u	
SEQUENCE: 52			
tttttacact ctttccctac	acgacgctct tccgatcttg	aggcactctg	ttgggatttt
ttttttttt tttttvn			60
			78
SEQ ID NO: 53		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	
modified_base	1..5		
		mod_base = OTHER	
		note = t is u	
SEQUENCE: 53			
tttttacact ctttccctac	acgacgctct tccgatctat	gattagtcgc	cattcgctttt
ttttttttt tttttvn			60
			78
SEQ ID NO: 54		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	
modified_base	1..5		
		mod_base = OTHER	
		note = t is u	
SEQUENCE: 54			
tttttacact ctttccctac	acgacgctct tccgatctac	ttgagggtag	atgttttttt
ttttttttt tttttvn			60
			78
SEQ ID NO: 55		moltype = DNA length = 78	
FEATURE		Location/Qualifiers	
source	1..78		
		mol_type = other DNA	
		organism = synthetic construct	

-continued

```

modified_base      1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 55
tttttacct ctttccctac acgacgctct tccgatctat ggccaactact gttatctttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 56      moltype = DNA length = 78
FEATURE           Location/Qualifiers
misc_feature      1..78
                   note = Probe 7
misc_feature      78
                   note = n is a, c, g, t or u
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 56
tttttacct ctttccctac acgacgctct tccgatctcg ctaccctgat tcgacctttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 57      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 57
tttttacct ctttccctac acgacgctct tccgatctgc ccactttcgc cgtagttttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 58      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 58
tttttacct ctttccctac acgacgctct tccgatctag caactttgag caagattttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 59      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 59
tttttacct ctttccctac acgacgctct tccgatctgc caattcggaa ttccggtttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 60      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

SEQUENCE: 60
tttttacct ctttccctac acgacgctct tccgatcttc gcccaaggta atacattttt 60
tttttttttt ttttttvn                                           78

SEQ ID NO: 61      moltype = DNA length = 78
FEATURE           Location/Qualifiers
source           1..78
                   mol_type = other DNA
                   organism = synthetic construct
modified_base     1..5
                   mod_base = OTHER
                   note = t is u

```

-continued

SEQUENCE: 61
 tttttacact ctttccctac acgacgctct tccgatcttc gcatttcta ttegagtttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 62 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 62
 tttttacact ctttccctac acgacgctct tccgatcttt gctaaatcta accgcctttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 63 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 63
 tttttacact ctttccctac acgacgctct tccgatctgg aattaaattc tgatggtttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 64 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 64
 tttttacact ctttccctac acgacgctct tccgatctca ttacataggt gctaagtttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 65 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 65
 tttttacact ctttccctac acgacgctct tccgatctat tgacttgccg tcgcactttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 66 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 66
 tttttacact ctttccctac acgacgctct tccgatctat agtatctccc aagttctttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 67 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 67
 tttttacact ctttccctac acgacgctct tccgatctgt gcgctgttaa tccgcatttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 68 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78

-continued

modified_base mol_type = other DNA
 organism = synthetic construct
 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 68
 tttttacact ctttccctac acgacgctct tccgatctgc gccactcttt aggtagtttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 69 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 69
 tttttacact ctttccctac acgacgctct tccgatctta tgcaagtgat tggctttttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 70 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 70
 tttttacact ctttccctac acgacgctct tccgatctcc aagccacggt tatacgtttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 71 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 71
 tttttacact ctttccctac acgacgctct tccgatctac ctgattgctg tataactttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 72 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 72
 tttttacact ctttccctac acgacgctct tccgatctca gcgcactctat cctctatttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 73 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 73
 tttttacact ctttccctac acgacgctct tccgatcttc cacgcgtagg actagttttt 60
 tttttttttt ttttttvn 78

SEQ ID NO: 74 moltype = DNA length = 78
 FEATURE Location/Qualifiers
 source 1..78
 mol_type = other DNA
 organism = synthetic construct
 modified_base 1..5
 mod_base = OTHER
 note = t is u

SEQUENCE: 74
 tttttacact ctttccctac acgacgctct tccgatctcg actaagtatg tagcgctttt 60

-continued

tttttttttt ttttttvn	78
SEQ ID NO: 75	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 75	
aaatttcgtc tgctatcgcg cttctgtacc	30
SEQ ID NO: 76	moltype = DNA length = 21
FEATURE	Location/Qualifiers
source	1..21
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 76	
ggtacagaag cgcgatagca g	21
SEQ ID NO: 77	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 77	
acactctttc cctacacgac gctcttcoga tct	33
SEQ ID NO: 78	moltype = DNA length = 43
FEATURE	Location/Qualifiers
source	1..43
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 78	
agacgtgtgc tcttccgac ttttttttt ttttttttt tvn	43
SEQ ID NO: 79	moltype = DNA length = 43
FEATURE	Location/Qualifiers
source	1..43
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 79	
tcagacgtgt gctcttcoga tcttttttt ttttttttt ttt	43
SEQ ID NO: 80	moltype = DNA length = 66
FEATURE	Location/Qualifiers
source	1..66
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 80	
gcgttcagag tggcagtcga gatcacgcgg caatcatatc ggacagatcg gaagagcgta	60
gtgtag	66
SEQ ID NO: 81	moltype = DNA length = 66
FEATURE	Location/Qualifiers
source	1..66
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 81	
gcgttcagag tggcagtcga gatcacaaga tgaacccggc tcatagatcg gaagagcgta	60
gtgtag	66
SEQ ID NO: 82	moltype = DNA length = 66
FEATURE	Location/Qualifiers
source	1..66
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 82	
gcgttcagag tggcagtcga gatcactccc aacagagtgc ctcaagatcg gaagagcgta	60
gtgtag	66
SEQ ID NO: 83	moltype = DNA length = 66
FEATURE	Location/Qualifiers
source	1..66
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 83	
gcgttcagag tggcagtcga gatcaccgaa tggcgactaa tcatagatcg gaagagcgta	60
gtgtag	66
SEQ ID NO: 84	moltype = DNA length = 66

-continued

FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 84		
gcgttcagag tggcagtcga gatcacaac atctaccctc aagtagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 85	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 85		
gcgttcagag tggcagtcga gatcacgata acagtattgg ccatagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 86	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 86		
gcgttcagag tggcagtcga gatcacggtc gaatcagggt agcgagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 87	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 87		
gcgttcagag tggcagtcga gatcacacta cggcgaaagt gggcagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 88	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 88		
gcgttcagag tggcagtcga gatcacatct tgctcaaagt tgctagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 89	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 89		
gcgttcagag tggcagtcga gatcacccgg aattccgaat tggcagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 90	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 90		
gcgttcagag tggcagtcga gatcacatgt attaccttgg gcgaagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 91	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 91		
gcgttcagag tggcagtcga gatcacctcg aataggaat gcgaagatcg gaagagcgta		60
gtgtag		66
SEQ ID NO: 92	moltype = DNA length = 66	
FEATURE	Location/Qualifiers	
source	1..66	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 92		
gcgttcagag tggcagtcga gatcacggcg gttagattta gcaaagatcg gaagagcgta		60
gtgtag		66

-continued

cggttcagag tggcagtcga gatcacgtta tacagcaatc aggtagatcg gaagagcgta 60
 gtgtag 66

SEQ ID NO: 102 moltype = DNA length = 66
 FEATURE Location/Qualifiers
 source 1..66
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 102
 cggttcagag tggcagtcga gatcactaga ggatagatgc gctgagatcg gaagagcgta 60
 gtgtag 66

SEQ ID NO: 103 moltype = DNA length = 66
 FEATURE Location/Qualifiers
 source 1..66
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 103
 cggttcagag tggcagtcga gatcacacta gtcctacgcg tggaagatcg gaagagcgta 60
 gtgtag 66

SEQ ID NO: 104 moltype = DNA length = 66
 FEATURE Location/Qualifiers
 source 1..66
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 104
 cggttcagag tggcagtcga gatcacgcbc tacatactta gtcgagatcg gaagagcgta 60
 gtgtag 66

SEQ ID NO: 105 moltype = DNA length = 30
 FEATURE Location/Qualifiers
 source 1..30
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 105
 aaatttcgtc tgctatcgcg cttctgtacc 30

SEQ ID NO: 106 moltype = DNA length = 45
 FEATURE Location/Qualifiers
 source 1..45
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 106
 gtgatctcga ctgccactct gaattttttt tttttttttt tttvn 45

SEQ ID NO: 107 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 107
 acactctttc cctacacgac gctcttccga tct 33

SEQ ID NO: 108 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 108
 acactctttc cctacacgac gctcttccga tct 33

SEQ ID NO: 109 moltype = DNA length = 43
 FEATURE Location/Qualifiers
 source 1..43
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 109
 agacgtgtgc tcttccgacg tttttttttt tttttttttt tvn 43

SEQ ID NO: 110 moltype = DNA length = 33
 FEATURE Location/Qualifiers
 source 1..33
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 110
 acactctttc cctacacgac gctcttccga tct 33

SEQ ID NO: 111 moltype = DNA length = 34

-continued

FEATURE	Location/Qualifiers	
source	1..34	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 111		
gtgactggag ttcagacgtg tgctcttccg atct		34
SEQ ID NO: 112	moltype = DNA length = 39	
FEATURE	Location/Qualifiers	
modified_base	37..39	
	mod_base = OTHER	
	note = Ribonucleotides	
source	1..39	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 112		
gtgactggag ttcagacgtg tgctcttccg atctatggg		39
SEQ ID NO: 113	moltype = DNA length = 33	
FEATURE	Location/Qualifiers	
source	1..33	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 113		
acactctttc cctacacgac gctcttccga tct		33
SEQ ID NO: 114	moltype = DNA length = 39	
FEATURE	Location/Qualifiers	
source	1..39	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 114		
agacgtgtgc tcttccgac ttggtgcct gaggcaatg		39
SEQ ID NO: 115	moltype = DNA length = 43	
FEATURE	Location/Qualifiers	
source	1..43	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 115		
agacgtgtgc tcttccgac tctcgtaac aagcagagag aac		43
SEQ ID NO: 116	moltype = DNA length = 40	
FEATURE	Location/Qualifiers	
source	1..40	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 116		
agacgtgtgc tcttccgac ttgagaacaa gggggaagag		40
SEQ ID NO: 117	moltype = DNA length = 41	
FEATURE	Location/Qualifiers	
source	1..41	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 117		
agacgtgtgc tcttccgac tccgtgaaac aagcaggtaa c		41
SEQ ID NO: 118	moltype = DNA length = 41	
FEATURE	Location/Qualifiers	
source	1..41	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 118		
agacgtgtgc tcttccgac tcattcccac actcatcaca c		41
SEQ ID NO: 119	moltype = DNA length = 40	
FEATURE	Location/Qualifiers	
source	1..40	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 119		
agacgtgtgc tcttccgac ttcacactgg agaaagaccc		40
SEQ ID NO: 120	moltype = DNA length = 43	
FEATURE	Location/Qualifiers	
source	1..43	
	mol_type = other DNA	
	organism = synthetic construct	

-continued

```

SEQUENCE: 120
agacgtgtgc tcttccgac tggggttcag agtgattttt cag          43

SEQ ID NO: 121      moltype = DNA length = 40
FEATURE            Location/Qualifiers
source             1..40
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 121
agacgtgtgc tcttccgac ttcggttttc tttcagtgcc          40

```

What is claimed is:

1. A method for detecting an mRNA from a tissue section, wherein the method comprises:
 - (a) providing an array comprising a plurality of features on a substrate, wherein each feature occupies a distinct position on the array, wherein each feature comprises a plurality of capture probes, and wherein a capture probe of the plurality of capture probes on each feature comprises a nucleic acid molecule having the following domains:
 - (i) a positional domain comprising a nucleotide sequence unique to the feature; and
 - (ii) a capture domain comprising a nucleotide sequence complementary to the mRNA;
 - (b) hybridizing the mRNA to the capture domain of the capture probe on one or more features of the array;
 - (c) extending the capture probe using the hybridized mRNA to generate a nucleic acid molecule that comprises a cDNA and the positional domain;
 - (d) introducing an amplification domain to the nucleic acid molecule using a template switching primer;
 - (e) generating a second strand that comprises a complement of the cDNA or a portion thereof;
 - (f) determining a nucleotide sequence of i) at least a portion of the second strand or a complement thereof, and ii) the positional domain, or a complement thereof; and
 - (g) using the determined nucleotide sequences of at least a portion of the second strand or the complement thereof, and the positional domain, or the complement thereof, to detect the mRNA in the tissue section.
2. The method of claim 1, wherein the tissue section comprises a fresh-frozen tissue section.
3. The method of claim 1, wherein the tissue section comprises a formalin-fixed paraffin embedded tissue section.
4. The method of claim 1, wherein the plurality of features comprises beads.
5. The method of claim 1, wherein the array comprises at least 50,000 features.
6. The method of claim 4, wherein the array comprises at least 50,000 beads, wherein each bead comprises an area of about 75 μm^2 .
7. The method of claim 1, wherein the plurality of capture probes is directly or indirectly immobilized to the features of the array.
8. The method of claim 1, wherein the method further comprises removing the hybridized mRNA from the cDNA.
9. The method of claim 1, wherein the nucleic acid molecule is orientated such that the positional domain is 5' to the capture domain.
10. The method of claim 1, wherein the capture domain comprises a poly-T sequence.
11. The method of claim 10, wherein the poly-T sequence comprises at least 25 nucleotides or at least 30 nucleotides.
12. The method of claim 1, wherein determining the nucleotide sequence of i) at least a portion of the second strand or the complement thereof, and ii) the positional domain, or the complement thereof, comprises sequencing.
13. The method of claim 1, wherein extending the capture probe comprises a reverse transcription reaction.
14. The method of claim 1, wherein the capture probe further comprises a cleavage domain.
15. The method of claim 1, further comprising after step (c), removing unextended capture probes, optionally with an exonuclease.
16. A method for detecting mRNA in a tissue section, wherein the method comprises:
 - (a) providing an array comprising a plurality of features on a substrate, wherein each feature occupies a distinct position on the array, wherein each feature comprises a plurality of capture probes, and wherein a capture probe of the plurality of capture probes on each feature comprises a nucleic acid molecule having the following domains:
 - (i) a positional domain comprising a nucleotide sequence unique to the feature; and
 - (ii) a capture domain comprising a poly-T sequence;
 - (b) applying the tissue section to the array;
 - (c) hybridizing one or more mRNAs from the tissue section to the capture domain of one or more capture probes on one or more features of the array;
 - (d) extending at least one capture probe of the one or more capture probes using a hybridized mRNA of the one or more mRNAs to generate a nucleic acid molecule that comprises a cDNA and the positional domain;
 - (e) introducing a polynucleotide tail to the 3' end of the nucleic acid molecule that comprises the cDNA and the positional domain;
 - (f) removing the tissue section from the array;
 - (g) generating a second strand that comprises a complement of the cDNA or a portion thereof;
 - (h) determining a nucleotide sequence of i) at least a portion of the second strand, or the complement thereof, and ii) the positional domain, or a complement thereof; and
 - (i) using the determined nucleotide sequences of i) at least a portion of the second strand, or the complement thereof, and ii) the positional domain, or the complement thereof, to detect mRNA in the tissue section.
17. The method of claim 16, wherein the tissue section comprises a fresh-frozen tissue section.
18. The method of claim 16, wherein the tissue section comprises a formalin-fixed paraffin embedded tissue section.
19. The method of claim 16, wherein the plurality of features comprises beads.
20. The method of claim 16, wherein the array comprises at least 50,000 features.

21. The method of claim 19, wherein the array comprises at least 50,000 beads, wherein each bead comprises an area of about 75 μm^2 .

22. The method of claim 16, wherein the plurality of capture probes is directly or indirectly immobilized to the features of the array. 5

23. The method of claim 16, wherein the poly-T sequence comprises at least 25 nucleotides or at least 30 nucleotides.

24. The method of claim 16, wherein the nucleic acid molecule is orientated such that the positional domain is 5' to the capture domain. 10

25. The method of claim 16, wherein the capture probe further comprises a cleavage domain.

26. The method of claim 16, wherein determining the nucleotide sequence of at least a portion of the second strand or the complement thereof, and the positional domain, or the complement thereof, comprises sequencing. 15

27. The method of claim 16, wherein removing the tissue section comprises treatment with a proteinase, optionally wherein the proteinase is Proteinase K. 20

28. The method of claim 16, wherein extending the at least one capture probe comprises a reverse transcription reaction, and wherein introducing the polynucleotide tail is performed using a reverse transcriptase.

29. The method of claim 16, wherein the method further comprises hybridizing a template switch primer to the polynucleotide tail, and extending the nucleic acid molecule in step (e) using the template switch primer as a template, wherein the template switch primer comprises an amplification domain, and wherein a complement of the amplification domain is incorporated into the nucleic acid molecule. 25

30. A method for spatially tagging an mRNA from a tissue section, wherein the method comprises:

- (a) providing an array comprising a plurality of features on a substrate, wherein each feature occupies a distinct position on the array, wherein each feature comprises a plurality of capture probes, and wherein a capture probe of the plurality of capture probes on each feature comprises a nucleic acid molecule having the following domains: 35
 - (i) a positional domain comprising a nucleotide sequence unique to the feature; and
 - (ii) a capture domain comprising a nucleotide sequence complementary to the mRNA; 40

(b) hybridizing the mRNA to the capture domain of one or more capture probes on one or more features of the array;

(c) extending at least one capture probe of the one or more capture probes using the hybridized mRNA to generate a nucleic acid molecule that comprises a cDNA and the positional domain;

(d) introducing an amplification domain to the nucleic acid molecule using a template switching primer; and

(e) generating a second strand that comprises a complement of the cDNA or a portion thereof.

31. The method of claim 30, wherein the tissue section comprises a fresh-frozen tissue section.

32. The method of claim 30, wherein the tissue section comprises a formalin-fixed paraffin embedded tissue section.

33. The method of claim 30, wherein the plurality of features comprises beads, and the array comprises at least 50,000 beads, wherein each bead comprises an area of about 75 μm^2 .

34. The method of claim 30, wherein the plurality of capture probes is indirectly immobilized to the features of the array.

35. The method of claim 30, wherein the plurality of capture probes is directly immobilized to the features of the array.

36. The method of claim 30, wherein the capture domain comprises a poly-T sequence.

37. The method of claim 36, wherein the poly-T sequence comprises at least 25 nucleotides or at least 30 nucleotides.

38. The method of claim 30, wherein the method further comprises removing the hybridized mRNA from the capture domain of the capture probe.

39. The method of claim 30, wherein the capture probe further comprises a cleavage domain.

40. The method of claim 30, further comprising after step (c), removing unextended capture probes, optionally with an exonuclease.

41. The method of claim 30, wherein the method further comprises hybridizing a template switch primer to the 3' end of the nucleic acid molecule.

42. The method of claim 30, wherein the method further comprises extending the nucleic acid molecule using the template switch primer as a template.

* * * * *