

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 May 2007 (03.05.2007)

PCT

(10) International Publication Number
WO 2007/050543 A2

(51) International Patent Classification:

A61K 9/00 (2006.01) A61K 31/17 (2006.01)
A61K 9/12 (2006.01) A61K 31/19 (2006.01)

(21) International Application Number:

PCT/US2006/041366

(22) International Filing Date: 24 October 2006 (24.10.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/729,788 24 October 2005 (24.10.2005) US

(71) Applicant (for all designated States except US): **COL-
LEGIUM PHARMACEUTICAL, INC.** [US/US]; 400
Highland Corporate Drive, Cumberland, RI 02864 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **TRUMBORE,
Mark, W.** [US/US]; 9 Heywood Road, Westford, MA
01886 (US). **GURGE, Ronald, M.** [US/US]; 156 Lincoln
Street, Franklin, MA 02038 (US). **HIRSH, Jane, C.**
[US/US]; 15 Peirce Road, Wellesley, MA 02581 (US).

(74) Agents: **PABST, Patrea, L.** et al.; Pabst Patent Group
LLP, 1201 Peachtree Street, Suite 1200, Atlanta, GA 30361
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS,
LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY,
MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS,
RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: TOPICAL PHARMACEUTICAL FOAM COMPOSITION

(57) Abstract: A stable topical alcohol-free aerosol foam containing one or more keratolytic agents is provided. The foam-forming formulation is an emulsion which contains an HFA propellant and one or more keratolytic agents. The emulsion has an oil phase and an aqueous, i.e. water-containing, phase. The active agent(s) may be present in either phase of the emulsion or dispersed in the emulsion. The oil phase may consist at least in part of the HFA propellant. Either or both of the oil phase and the aqueous phase may contain one or more surfactants, emulsifiers, emulsion stabilizers, buffers, and/or other excipients. The foam is stable on the skin, for example, for at least 5 minutes at body temperature, preferably at least 20 minutes at body temperature, and disappears into the skin upon rubbing or after prolonged standing. In one embodiment, the formulation contains an HFA propellant which does not contain additional co- solvents or co-propellants. The formulations demonstrate reduced intensity of the odor and/or color associated with the keratolytic agent(s) as compared to conventional formulations containing keratolytic agents.

WO 2007/050543 A2

TOPICAL PHARMACEUTICAL FOAM COMPOSITION

FIELD OF THE INVENTION

This invention is generally in the field of pharmaceutical compositions, specifically pharmaceutical foam compositions containing
5 keratolytic agents intended for topical administration.

CROSS-REFERENCE TO RELATED APPLICATIONS

This invention claims the benefit of the priority of U.S.S.N.
60/729,788, filed October 24, 2005.

BACKGROUND OF THE INVENTION

10 Pharmaceutical foams are pressurized dosage forms containing one or more active ingredients that, upon valve actuation, emit a fine dispersion of liquid and/or solid materials in a gaseous medium. Foam formulations are generally easier to apply, are less dense, and spread more easily than other topical dosage forms. Foams may be formulated in various ways to provide
15 emollient or drying functions to the skin, depending on the formulation constituents. Accordingly, this delivery technology is a useful addition to the spectrum of formulations available for topical use. However, as yet, only a few pharmaceutical foams are commercially available. Most commercially available foam dosage forms contain corticosteroids, although some products
20 have also been used to deliver antiseptics, antifungal agents, anti-inflammatory agents, local anesthetic agents, skin emollients, and protectants (American Journal of Drug Delivery, 2003, vol. 1(1), pp. 71-75).

There is growing interest in converting non-foam topical treatments to aerosol foam or mousse formulations, which better penetrate the skin,
25 provide faster treatment and do not leave any greasy residue on skin or clothing compared with conventional ointments. Until now, the most common gas propellant used in aerosol products is chlorofluorocarbon (CFC), an ozone-depleting agent. The Montreal Protocol International Treaty, signed by 180 nations, banned the use of chlorofluorocarbons (CFCs)
30 as aerosol propellants and mandated the phasing out of CFC agents. No new or revised aerosol formulations may contain CFC propellants and alternative propellants must be used that are more environmentally friendly. Therefore,

propellants must be used that are more environmentally friendly. Therefore, manufacturers must reformulate or modify existing products to use non-CFC propellants, while maintaining important aspects of the previous formulation, such as accuracy of delivery, stability, etc. The primary CFC substitute are
5 hydrofluoroalkanes (HFA), also known as hydrofluorocarbons (HFC).

Although hydrocarbon propellants, such as propane and butane, can be used in the manufacturing of pharmaceutical foams, these propellants are not suited for human use since they are flammable. Just as is the case with CFC propellants, hydrofluoroalkanes (HFAs) that possess high chemical
10 stability can be used as a primary substitute for hydrocarbons. Examples of HFAs are 1,1,1,2,3,3,3-heptafluoropropane (HFA-134a) and 1,1,1,2-tetrafluoroethane (HFA-227). Hydrofluoroalkanes (HFAs) are also referred to as hydrofluorocarbons (HFCs) and these terms are used interchangeably.

Since replacing a component of any formulation means introducing
15 new properties, and HFAs differ in their solvating power from CFCs and hydrocarbons, providing reproducible performance of reformulated aerosols for pharmaceutical uses represents a challenging task. Co-solvents (such as ethanol) are often incorporated into the formulation in order to arrive at a stable product (Pharmaceutical Aerosols, June 2003, p. 21). Such
20 formulations, however, have a number of undesirable aspects. Alcohol co-solvents can dry and irritate the skin. U.S. Patent No. 6,126,920 suggests that the use of alcohol co-solvents can lead to burning, itching, and irritation observed in the use of topical foam for delivering betamethasone. Further, volatile alcohols are highly irritating to mucous membranes.

25 Formulations that contain volatile alcohols and/or alkanes are potential safety hazards due to the high flammability of the product. Moreover, the flammability characteristics of the product require expensive precautions during manufacturing, and may require controlled environments for storage and for disposal of containers after use. For example, WO
30 85/01876 describes the fire hazards associated with alcohol- and alkane-containing aerosol foam formulations.

Pharmaceutical foam formulations containing keratolytic agents have not been described in the literature. Keratolytic agents are agents that soften, separate, and cause desquamation (i.e. shedding or peeling) of the cornified epithelium, or horny layer, of the skin. These agents are used to expose mycelia of infecting fungi or to treat corns, warts, and certain other skin diseases. Commonly used keratolytic agents include urea, urea in combination with ammonium lactate, salicylic acid, papain, papain in combination with urea, and sulfur. Sulfur is also used in combination with sodium sulfacetamide to treat acne, rosacea, seborrheic dermatitis, eczema, xerosis, scabies, pediculosis and psoriasis.

Keratolytic agents can be administered in the form of a liquid, cream, lotion or cleanser. Topical formulations containing keratolytic agents typically have an intense color and/or strong odor. For example, sulfur containing products typically have an intense yellow color and/or a strong odor characteristic of sulfur. Urea-containing products frequently exhibit a strong ammonia odor, while papain-containing products exhibit a distinctive papain odor. Current products on the market typically contain substantial amounts of odor masking agents, such as fragrances, in order to mask or shield the odor associated with the pharmaceutical agent. However, the use of high concentrations of fragrances can be problematic. The use of fragrances in topical formulations can result in skin sensitizing reactions in which the patient develops sensitivity to the odor masking agent. Furthermore, the presence of fragrances, which are often complex mixtures of different compounds, may result in undesirable side reactions between the fragrance and the active agent(s).

There exists a need for topical keratolytic foam formulations which, once applied to the skin, have little or no odor or color and are non-staining and which contain little or no odor masking agents such as fragrances.

It is therefore an object of the invention to provide alcohol-free keratolytic topical foam aerosol formulations that use hydrofluoroalkanes (HFAs) as the propellant.

It is a further object of the invention to provide keratolytic topical foam formulations which exhibit reduced intensity of the odor and/or color associated with the keratolytic agent.

BRIEF SUMMARY OF THE INVENTION

5 A stable topical alcohol-free aerosol foam containing one or more keratolytic agents is described herein. The foam-forming formulation is an emulsion which contains an HFA propellant and one or more keratolytic agents. The formulation optionally contains one or more additional
10 pharmaceutically active agents including, but not limited to, antibiotic agents, antimicrobial agents, anti-acne agents, antibacterial agents, antifungal agents, antiviral agents, steroidal anti-inflammatory agents, non-steroidal anti-inflammatory agents, anesthetic agents, antipruriginous agents, antiprotozoal agents, anti-oxidants, chemotherapeutic agents, antidepressants, antihistamines, vitamins, sunscreens, skin-soothing agents,
15 hormones, and anti-dandruff agents.

 The emulsion contains an oil phase and an aqueous, i.e. water-containing, phase. The active agent(s) may be dissolved in either phase or dispersed in the emulsion. The oil phase may contain, at least in part, the HFA propellant. The foam is stable on the skin, preferably for at least five
20 minutes, more preferably at least twenty minutes, at body temperature, and disappears into the skin upon rubbing or after prolonged standing. The formulations demonstrate reduced intensity of the odor and/or color associated with the keratolytic agent(s) as compared to conventional formulations containing keratolytic agents.

25 The composition can further contain one or more pharmaceutically acceptable excipients such as surfactants, emollients, emulsifiers, pH stabilizing agents, chelating agents, humectants, stabilizers, preservatives, and combinations thereof, which may be present in the oil phase and/or the aqueous phase. The formulations contain little or no fragrances and/or odor
30 masking agents, thus minimize the skin sensitizing reactions that can occur due to the presence of such agents.

The inert non-flammable HFA propellant does not require the use of additional co-solvents or co-propellants. Besides their high volatility and vapor pressure, the HFA propellants have been found to provide an additional benefit in terms of reduction of apparent odor of active ingredients, such as sulfur and urea. They also mask color by the formation of stable foams; and leave depots of materials on the skin, which in practice, are less staining to linen and clothing than prior art preparations with the same active ingredients.

In one embodiment the active agent is a keratolytic agent or agents such as urea or urea in combination with ammonium lactate, salicylic acid, papain, and/or sulfur. The keratolytic agent is present in an amount from about 1% to about 60% by weight of the final composition.

In another embodiment, the formulation contains a keratolytic agent in combination with an antibiotic. The concentration of the antibiotic is from about 0.01% to about 20%, preferably from about 1% to about 15%, more preferably from about 6% to about 12% by weight of the final composition. In a preferred embodiment, the formulation contains a combination of sulfur and sulfacetamide.

In yet another embodiment, the composition comprises the enzyme papain in combination with urea. The concentration of papain is from about 0.5% to about 40%, preferably from about 1% to about 20%, more preferably from about 1% to about 10% by weight of the final composition. The concentration of urea is from about 1% to about 60%, preferably from about 2.5% to about 40%, more preferably from about 5% to about 15% by weight of the final composition.

In still another embodiment, the composition comprises the enzyme papain in combination with urea and chlorophyllin copper complex sodium. The concentration of papain is from about 0.5% to about 40%, preferably from about 1% to about 20%, more preferably from about 1% to about 10% by weight of the final composition. The concentration of urea is from about 1% to about 60%, preferably from about 2.5% to about 40%, more preferably from about 5% to about 15% by weight of the final composition. The

concentration of chlorophyllin copper complex sodium is from about 0.05% to about 5%, preferably from about 0.1% to about 3%, more preferably from about 0.3% to about 1% by weight of the final composition.

The composition can be administered as a continuous or metered
5 dose that can be applied to the skin or mucous membranes.

DETAILED DESCRIPTION OF THE INVENTION

I. Concentrate

A. Propellants

In one embodiment, the propellant is a HFA or a mixture of one or
10 more hydrofluorocarbons. Suitable hydrofluorocarbons include 1,1,1,2-tetrafluoroethane (HFA 134a); 1,1,1,2,3,3,3-heptafluoropropane (HFA 227); and mixtures and admixtures of these and other HFAs that are currently approved or may become approved for medical use are suitable. The concentration of the HFA propellant is from about 5% to about 30% by
15 weight of the concentrate, which corresponds to about 4% to about 23% by weight of the final composition.

Hydrocarbon propellants such as butane/isobutane/propane have inherent, unpleasant odors which may impart negative olfactory sensory attributes to pharmaceutical foam products. When these foam products are
20 dispensed, the propellant expands and evaporates, creating a bubble structure within the foam. It is the evaporative release of the propellant which is detected by the olfactory senses of the user. Therefore, it is advantageous to have a propellant which is odor free and essentially provides no additional base odors to the foamed composition. HFAs, particularly 1,1,1,2-
25 Tetrafluoroethane (Freon-134a, HFC-134a), have no unpleasant odors associated with them and are preferred in the compositions described herein.

Furthermore, the compositions preferably contain no volatile alcohols or hydrocarbon propellant gases, which can produce flammable or explosive vapors during use. However, small amounts of such propellants may be used
30 as adjunct propellants if required for particular formulations.

B. Keratolytic Agents

Suitable keratolytic agents include, but are not limited to, urea, salicylic acid, papain, sulfur, glycolic acid, pyruvic acid, resorcinol, N-acetylcysteine, retinoids such as retinoic acid and its derivatives (e.g., cis and trans, esters), alpha hydroxy acids, beta hydroxy acids, coal tar, and combinations thereof. The concentration of the keratolytic agent is from about 1% to about 60% by weight of the final composition.

In one embodiment, the keratolytic agent is urea. Urea is present in an amount from about 5 to about 50%, preferably from about 10% to about 50%, more preferably from about 20% to about 40% by weight of the final composition. In another embodiment, ammonium lactate is added to a urea-containing formulation. Ammonium lactate is present in an amount from about 1 to about 30%, preferably from about 5% to about 20%, more preferably from about 10% to about 15% by weight of the final composition.

In another embodiment, the keratolytic agent is salicylic acid. Salicylic acid is present in an amount from about 1 to about 30%, preferably from about 4% to about 10%.

In another embodiment, the formulation comprises the enzyme papain and, optionally, urea. Papain is a protein-cleaving enzyme derived from papaya and certain other plants. The concentration of papain is from about 0.5% to about 40%, preferably from about 1% to about 20%, more preferably from about 1% to about 10% by weight of the final composition. The concentration of urea is from about 1% to about 60%, preferably from about 2.5% to about 40%, more preferably from about 5% to about 15% by weight of the final composition.

In yet another embodiment, the composition comprises the enzyme papain in combination with urea and chlorophyllin copper complex sodium. The concentration of papain is from about 0.5% to about 40%, preferably from about 1% to about 20%, more preferably from about 1% to about 10% by weight of the final composition. The concentration of urea is from about 1% to about 60%, preferably from about 2.5% to about 40%, more preferably from about 5% to about 15% by weight of the final composition. The

concentration of chlorophyllin copper complex sodium is from about 0.05% to about 5%, preferably from about 0.1% to about 3%, more preferably from about 0.3% to about 1% by weight of the final composition.

C. Other Active Agents

5 The compositions optionally contain one or more additional pharmaceutically active agents. Suitable classes of active agents include, but are not limited to, antibiotic agents, antimicrobial agents, anti-acne agents, antibacterial agents, antifungal agents, antiviral agents, steroidal anti-inflammatory agents, non-steroidal anti-inflammatory agents, anesthetic
10 agents, antipruriginous agents, antiprotozoal agents, anti-oxidants, antihistamines, vitamins, and hormones.

i. Antibiotics

Representative antibiotics include, without limitation, benzoyl peroxide, octopirox, erythromycin, zinc, tetracyclin, triclosan, azelaic acid
15 and its derivatives, phenoxy ethanol and phenoxy propanol, ethylacetate, clindamycin and meclocycline; sebastats such as flavinoids; alpha and beta hydroxy acids; and bile salts such as scymnol sulfate and its derivatives, deoxycholate and cholate. The antibiotic can be an antifungal agent. Suitable antifungal agents include, but are not limited to, clotrimazole,
20 econazole, ketoconazole, itraconazole, miconazole, oxiconazole, sulconazole, butenafine, naftifine, terbinafine, undecylinic acid, tolnaftate, and nystatin.

In one embodiment, the formulation contains one or more keratolytic agents in combination with an antibiotic agent. The concentration of the
25 antibiotic is from about 0.01% to about 20%, preferably from about 1% to about 15%, more preferably from about 6% to about 12% by weight of the final composition.

ii. Non-steroidal anti-inflammatory agents

Representative examples of non-steroidal anti-inflammatory agents
30 include, without limitation, oxicams, such as piroxicam, isoxicam, tenoxicam, sudoxicam; salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, and fendosal; acetic acid derivatives,

such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepirac, clindanac, oxepinac, felbinac, and ketorolac; fenamates, such as mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acids; propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, piroprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic; pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone. Mixtures of these non-steroidal anti-inflammatory agents may also be employed, as well as the dermatologically acceptable salts and esters of these agents. For example, etofenamate, a flufenamic acid derivative, is particularly useful for topical application.

iii. Steroidal anti-inflammatory agents

Representative examples of steroidal anti-inflammatory drugs include, without limitation, corticosteroids such as hydrocortisone, hydroxyl-triamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionates, clobetasol valerate, desonide, desoxymethasone, desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, diflucortolone valerate, fluadrenolone, flucolorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylesters, fluocortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescinalone, dichlorisone, diflurprednate, flucoloronide, flunisolid, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate,

meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, and mixtures thereof.

iv. Anesthetics

Representative anesthetics include, but are not limited to, lidocaine,
5 bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine and phenol.

v. Antimicrobial agents

Suitable antimicrobial agents include, but are not limited to, antibacterial, antifungal, antiprotozoal and antiviral agents, such as beta-
10 lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, triclosan, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, streptomycin, tobramycin,
15 and miconazole. Also included are tetracycline hydrochloride, famesol, erythromycin estolate, erythromycin stearate (salt), amikacin sulfate, doxycycline hydrochloride, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride,
20 metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin sulfate, miconazole hydrochloride,
25 amanfadine hydrochloride, amanfadine sulfate, triclosan, octopirox, nystatin, tolnaftate, clotrimazole, anidulafungin, micafungin, voriconazole, lanoconazole, ciclopirox and mixtures thereof.

vi. Other agents

Suitable other agents include, but are not limited to, deodorant agents,
30 antiperspirants, sun screening agents, sunless tanning agents, vitamins, hair conditioning agents, anti-irritants, and combinations thereof.

Examples of skin soothing agents include, but are not limited to, aloe, avocado oil, green tea extract, hops extract, chamomile extract, colloidal oatmeal, calamine, cucumber extract, and combinations thereof.

5 Examples of vitamins include, but are not limited to, vitamins A, D, E, K, and combinations thereof.

 Examples of sunscreens include, but are not limited to, p-Aminobenzoic acid, Avobenzene, Cinoxate, Dioxybenzone, Homosalate, Menthyl anthranilate, Octocrylene, Octyl methoxycinnamate, Octyl salicylate, Oxybenzone, Padimate O, Phenylbenzimidazole sulfonic acid, 10 Sulisobenzene, Titanium dioxide, Trolamine salicylate, Zinc oxide, 4-methylbenzylidene camphor, Methylene Bis-Benzotriazolyl Tetramethylbutylphenol, Bis-Ethylhexyloxyphenol Methoxyphenyl Triazine, Terephthalylidene Dicamphor Sulfonic Acid, Drometrizole Trisiloxane, Disodium Phenyl Dibenzenimidazole Tetrasulfonate, Diethylamino 15 Hydroxybenzoyl Hexyl Benzoate, Octyl Triazone, Diethylhexyl Butamido Triazone, Polysilicone-15, and combinations thereof.

D. Excipients

 The concentrate can be in the form of an emulsion. An emulsion is a preparation of one liquid distributed in small globules throughout the body of 20 a second liquid. The dispersed liquid is the discontinuous phase, and the dispersion medium is the continuous phase. When oil is the dispersed liquid and an aqueous solution is the continuous phase, it is known as an oil-in-water emulsion, whereas when water or aqueous solution is the dispersed phase and oil or oleaginous substance is the continuous phase, it is known as 25 a water-in-oil emulsion. The oil phase may consist at least in part of the propellant. Either or both of the oil phase and the aqueous phase may contain one or more excipients such as surfactants, emulsifiers, emulsion stabilizers, anti-oxidants, emollients, humectants, chelating agents, suspending agents, thickening agents, occlusive agents, preservatives, 30 stabilizing agents, pH modifying agents, solubilizing agents, penetration enhancers, and other excipients.

Suitable emulsifiers include, but are not limited to, straight chain or branched fatty acids, polyoxyethylene sorbitan fatty acid esters, sorbitan fatty acid esters, propylene glycol stearate, glyceryl stearate, polyethylene glycol, fatty alcohols, polymeric ethylene oxide-propylene oxide block copolymers, and combinations thereof.

Suitable surfactants include, but are not limited to, anionic surfactants, non-ionic surfactants, cationic surfactants, and amphoteric surfactants. Examples of anionic surfactants include, but are not limited to, ammonium lauryl sulfate, sodium lauryl sulfate, ammonium laureth sulfate, sodium laureth sulfate, alkyl glyceryl ether sulfonate, triethylamine lauryl sulfate, triethylamine laureth sulfate, triethanolamine lauryl sulfate, triethanolamine laureth sulfate, monoethanolamine lauryl sulfate, monoethanolamine laureth sulfate, diethanolamine lauryl sulfate, diethanolamine laureth sulfate, lauric monoglyceride sodium sulfate, potassium lauryl sulfate, potassium laureth sulfate, sodium lauryl sarcosinate, sodium lauroyl sarcosinate, lauryl sarcosine, cocoyl sarcosine, ammonium cocoyl sulfate, ammonium lauroyl sulfate, sodium cocoyl sulfate, sodium lauroyl sulfate, potassium cocoyl sulfate, potassium lauryl sulfate, triethanolamine lauryl sulfate, triethanolamine lauryl sulfate, monoethanolamine cocoyl sulfate, monoethanolamine lauryl sulfate, sodium tridecyl benzene sulfonate, sodium dodecyl benzene sulfonate, sodium and ammonium salts of coconut alkyl triethylene glycol ether sulfate; tallow alkyl triethylene glycol ether sulfate, tallow alkyl hexaoxyethylene sulfate, disodium N-octadecylsulfosuccinate, disodium lauryl sulfosuccinate, diammonium lauryl sulfosuccinate, tetrasodium N-(1,2-dicarboxyethyl)-N-octadecylsulfosuccinate, diamyl ester of sodium sulfosuccinic acid, dihexyl ester of sodium sulfosuccinic acid, dioctyl esters of sodium sulfosuccinic acid, docusate sodium, and combinations thereof.

Examples of nonionic surfactants include, but are not limited to, polyoxyethylene fatty acid esters, sorbitan esters, cetyl octanoate, cocamide DEA, cocamide MEA, cocamido propyl dimethyl amine oxide, coconut fatty acid diethanol amide, coconut fatty acid monoethanol amide, diglyceryl

diisostearate, diglyceryl monoisostearate, diglyceryl monolaurate, diglyceryl
 monooleate, ethylene glycol distearate, ethylene glycol monostearate,
 ethoxylated castor oil, glyceryl monoisostearate, glyceryl monolaurate,
 glyceryl monomyristate, glyceryl monooleate, glyceryl monostearate,
 5 glyceryl tricaprilate/caprinate, glyceryl triisostearate, glyceryl trioleate, glycol
 distearate, glycol monostearate, isooctyl stearate, lauramide DEA, lauric acid
 diethanol amide, lauric acid monoethanol amide, lauric/myristic acid
 diethanol amide, lauryl dimethyl amine oxide, lauryl/myristyl amide DEA,
 lauryl/myristyl dimethyl amine oxide, methyl gluceth, methyl glucose
 10 sesquisteate, oleamide DEA, PEG-distearate, polyoxyethylene butyl ether,
 polyoxyethylene cetyl ether, polyoxyethylene lauryl amine, polyoxyethylene
 lauryl ester, polyoxyethylene lauryl ether, polyoxyethylene nonylphenyl
 ether, polyoxyethylene octyl ether, polyoxyethylene octylphenyl ether,
 polyoxyethylene oleyl amine, polyoxyethylene oleyl cetyl ether,
 15 polyoxyethylene oleyl ester, polyoxyethylene oleyl ether, polyoxyethylene
 stearyl amine, polyoxyethylene stearyl ester, polyoxyethylene stearyl ether,
 polyoxyethylene tallow amine, polyoxyethylene tridecyl ether, propylene
 glycol monostearate, sorbitan monolaurate, sorbitan monooleate, sorbitan
 monopalmitate, sorbitan monostearate, sorbitan sesquioleate, sorbitan
 20 trioleate, stearamide DEA, stearic acid diethanol amide, stearic acid
 monoethanol amide, laureth-4, and combinations thereof.

Examples of amphoteric surfactants include, but are not limited to,
 sodium N-dodecyl- β -alanine, sodium N-lauryl- β -iminodipropionate,
 myristoamphoacetate, lauryl betaine, lauryl sulfobetaine, sodium 3-dodecyl-
 25 aminopropionate, sodium 3-dodecylaminopropane sulfonate, sodium
 lauroamphoacetate, cocodimethyl carboxymethyl betaine, cocoamidopropyl
 betaine, cocobetaine, lauryl amidopropyl betaine, oleyl betaine, lauryl
 dimethyl carboxymethyl betaine, lauryl dimethyl alphacarboxyethyl betaine,
 cetyl dimethyl carboxymethyl betaine, lauryl bis-(2-hydroxyethyl)
 30 carboxymethyl betaine, stearyl bis-(2-hydroxypropyl) carboxymethyl
 betaine, oleyl dimethyl gamma-carboxypropyl betaine, lauryl bis-(2-
 hydroxypropyl)alpha-carboxyethyl betaine, oleamidopropyl betaine, coco

dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine, and combinations thereof.

Examples of cationic surfactants include, but are not limited to,
5 behenyl trimethyl ammonium chloride, bis(acyloxyethyl) hydroxyethyl methyl ammonium methosulfate, cetrimonium bromide, cetrimonium chloride, cetyl trimethyl ammonium chloride, cocamido propylamine oxide, distearyl dimethyl ammonium chloride, ditallowdimonium chloride, guar hydroxypropyltrimonium chloride, lauralkonium chloride, lauryl
10 dimethylamine oxide, lauryl dimethylbenzyl ammonium chloride, lauryl polyoxyethylene dimethylamine oxide, lauryl trimethyl ammonium chloride, laurtrimonium chloride, methyl-1-oleyl amide ethyl-2-oleyl imidazolium methyl sulfate, picolin benzyl ammonium chloride, polyquaternium, stearalkonium chloride, sterayl dimethylbenzyl ammonium chloride, stearyl
15 trimethyl ammonium chloride, trimethylglycine, and combinations thereof.

Suitable suspending agents include, but are not limited to, alginic acid, bentonite, carbomer, carboxymethylcellulose and salts thereof, colloidal oatmeal, hydroxyethylcellulose, hydroxypropylcellulose, microcrystalline cellulose, colloidal silicon dioxide, dextrin, gelatin, guar gum, xanthan gum,
20 kaolin, magnesium aluminum silicate, maltitol, triglycerides, methylcellulose, polyoxyethylene fatty acid esters, polyvinylpyrrolidone, propylene glycol alginate, sodium alginate, sorbitan fatty acid esters, tragacanth, and combinations thereof.

Suitable antioxidants include, but are not limited to, butylated
25 hydroxytoluene, alpha tocopherol, ascorbic acid, fumaric acid, malic acid, butylated hydroxyanisole, propyl gallate, sodium ascorbate, sodium metabisulfite, ascorbyl palmitate, ascorbyl acetate, ascorbyl phosphate, Vitamin A, folic acid, flavons or flavonoids, histidine, glycine, tyrosine, tryptophan, carotenoids, carotenes, alpha-Carotene, beta-Carotene, uric acid,
30 pharmaceutically acceptable salts thereof, derivatives thereof, and combinations thereof.

Suitable chelating agents include, but are not limited to, EDTA, disodium edetate, trans-1,2-diaminocyclohexane-N,N,N',N'-tetraacetic acid monohydrate, N,N-bis(2-hydroxyethyl)glycine, 1,3-diamino-2-hydroxypropane-N,N,N',N'-tetraacetic acid, 1,3-diaminopropane-N,N,N',N'-tetraacetic acid, ethylenediamine-N,N'-diacetic acid, ethylenediamine-N,N'-dipropionic acid, ethylenediamine-N,N'-bis(methylenephosphonic acid), N-(2-hydroxyethyl)ethylenediamine-N,N',N'-triacetic acid, ethylenediamine-N,N,N',N'-tetrakis(methylenephosphonic acid), O,O'-bis(2-aminoethyl)ethyleneglycol-N,N,N',N'-tetraacetic acid, N,N-bis(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid, 1,6-hexamethylenediamine-N,N,N',N'-tetraacetic acid, N-(2-hydroxyethyl)iminodiacetic acid, iminodiacetic acid, 1,2-diaminopropane-N,N,N',N'-tetraacetic acid, nitrilotriacetic acid, nitrilotripropionic acid, nitrilotris(methylenephosphoric acid), 7,19,30-trioxa-1,4,10,13,16,22,27,33-octaazabicyclo[11,11,11] pentatriacontane hexahydrobromide, triethylenetetramine-N,N,N',N'',N''',N'''-hexaacetic acid, and combinations thereof.

Suitable emollients include, but are not limited to, myristyl lactate, isopropyl palmitate, light liquid paraffin, cetearyl alcohol, lanolin, lanolin derivatives, mineral oil, petrolatum, cetyl esters wax, cholesterol, glycerol, glycerol monostearate, isopropyl myristate, lecithin, and combinations thereof thereof.

Suitable humectants include, but are not limited to, glycerin, butylene glycol, propylene glycol, sorbitol, triacetin, and combinations thereof.

The compositions described herein may further contain sufficient amounts of at least one pH modifier to ensure that the composition has a final pH of about 3 to about 11. Suitable pH modifying agents include, but are not limited to, sodium hydroxide, citric acid, hydrochloric acid, acetic acid, phosphoric acid, succinic acid, sodium hydroxide, potassium hydroxide, ammonium hydroxide, magnesium oxide, calcium carbonate, magnesium carbonate, magnesium aluminum silicates, malic acid, potassium citrate, sodium citrate, sodium phosphate, lactic acid, gluconic acid, tartaric

acid, 1,2,3,4-butane tetracarboxylic acid, fumaric acid, diethanolamine, monoethanolamine, sodium carbonate, sodium bicarbonate, triethanolamine, and combinations thereof.

Preservatives can be used to prevent the growth of fungi and other microorganisms. Suitable preservatives include, but are not limited to, benzoic acid, butylparaben, ethyl paraben, methyl paraben, propylparaben, sodium benzoate, sodium propionate, benzalkonium chloride, benzethonium chloride, benzyl alcohol, cetylpyridinium chloride, chlorobutanol, phenol, phenylethyl alcohol, thimerosal, and combinations thereof.

II. Method of Making the Formulation

A. Method of Preparing An Emulsion Concentrate

The oil phase is prepared by mixing together the surfactant(s) and emulsifier(s), and heating if necessary. The aqueous phase is prepared separately by dissolving the propylene glycol and preservatives in water with heating. The oil phase is added to the aqueous phase with continuous high shear mixing to produce a milky emulsion. The emulsion is cooled and if necessary, the pH is adjusted by the addition of a pH modifying agent.

If desired, the active agent(s) can be separately suspended or dissolved in water and/or propylene glycol and treated to eliminate any large aggregates. In a small scale operation, the mixture can be milled. The final active agent particle size is small enough to allow aerosolization, for example, less than about 20 microns in diameter, preferably less than about 10 microns, more preferably, less than about 5 microns. The liquid containing the active agent is added to the emulsion with mixing.

Alternatively, the active agent can be added directly to the water phase prior to emulsification. The formulation is brought to the final weight by the addition of water.

The concentration of the surfactant(s) in the concentrate is from about 0.5 to about 15% by weight of the final composition. The concentration of the emulsifier(s) is from about 0.5% to about 5% by weight of the final composition. The concentration of the pH modifying agent(s) is from about 0.1% to about 5% by weight of the final composition. The concentration of

the preservative(s) is from about 0.01% to about 1% by weight of the final composition.

B. Method of Preparing the Formulation

The emulsion concentrate is placed in pressure cans, preferably coated aluminum cans to prevent corrosion, such as epoxy-coated cans. The lid and dispensing apparatus are crimped in place. The can is charged with propellant to the stated level, for example, by adding 10 grams of propellant per 90 grams of emulsion. At the time of application, the mixture of the emulsion with the propellant may be insured by shaking, optionally with the aid of a mixing bead. The dispenser may be metered or unmetered (continuous). The can may be arranged for either "upside down" spraying with the valve at the bottom, or the can have a dip tube so that the foam can be sprayed while the can is upright with the valve at the top. In one embodiment, the concentration of the HFC propellant(s) is from about 5% to about 40% by weight of the final composition, more preferably about 5% to about 20% by weight of the final composition. In another embodiment, the emulsion concentrate is mixed with an HFC propellant so that the final formulation in an aerosol can comprises about 90% to about 95% of concentrate and about 5% to about 10% of propellant.

C. Foam Viscosity

In one embodiment, the foam has a sufficiently high viscosity to prevent the formulation from "running off" the site of application. The formulation also preferably has a sufficiently low viscosity that the formulation efficiently releases the actives to the treatment site. These contrasting attributes can be realized in a shear-thinning foam. "Shear-thinning" describes the rheological condition where the viscosity of a material subjected to constant shear stress decreases. The amount of decrease in viscosity is a function of the degree of shear stress applied. Upon removal of the shear stress, the viscosity again increases to the original value over time. Two values are of particular importance in developing shear-thinning formulations for hyperkeratotic conditions. These are the zero shear viscosity and the yield stress. The zero shear viscosity dictates the resistance

of the formulation to flow after application and ability of the formulation to release actives. The yield stress defines the stress level at which the material transitions from a "solid-like" poorly flowing high viscosity foam to a "liquid-like" well flowing low viscosity foam. The lower the yield stress, the easier it is to uniformly spread and rub-in the formulation over the treatment site.

Foam rheology is a function of the composition of the foam base and the expansion factor of the foam. Expansion factor is the volume a given mass of foam occupies and is the reciprocal of the foam density. Foam expansion factor is determined by the foam base composition and the composition and concentration of propellant. For a given foam base and propellant composition, changes in the expansion factor of the foam can be achieved by varying the concentration of propellant.

For topical foam products, the zero shear viscosities are preferably greater than about 30,000 cP, more preferably greater than about 45,000cP, and most preferably greater than about 60,000 cP; and also less than about 700,000 cP, more preferably less than about 500,000 cP and still more preferably less than about 300,000 cP. The preferred yield stress values are preferably greater than about 250 dynes/cm², more preferably greater than about 750 dynes/cm², and most preferably greater than about 1000 dynes/cm², and also preferably lower than about 60,000 dynes/cm², more preferably lower than about 30,000 dynes/cm², and still more preferably lower than about 10,000 dynes/cm². The preferred foam expansion factor is from 1.5 to 15 cm³/g, more preferably from 1.8 to 10 cm³/g, most preferably from 2.0 and 7.0 cm³/g.

III. Mode of Administration

a. Method of Administration to a Patient

The formulation is administered to the skin or wound of a human or animal. A selected amount of product is dispensed from the spray can, preferably onto the site to be treated. The foam can be administered into the palm of the hand. Alternatively, the foam can be applied to a wound dressing. The amount to be delivered can be determined by the prescribing

physician or as directed in the instructions for non-prescription products. Alternatively, a fixed dose using the metering dispenser can be administered. The foam is rubbed into the skin at the site to be treated. Because the foam is stable at body temperature, this step does not need to be hurried.

5 Moreover, the exact site of application can be more easily controlled. If contact with the hand is to be avoided, a glove may be worn; or, the foam can be first be applied to a wound dressing or may be left in place, wherein it will eventually collapse and deliver the active ingredient to the surface of the skin.

10

Examples

Example 1. Reduced Odor Topical Formulation Containing Sulfur and Sodium Sulfacetamide

A topical formulation containing sulfur and sodium sulfacetamide was prepared wherein the formulation exhibited diminished color. The composition of the formulation and the physical and mechanical properties
15 of the formulation are shown in Table 1.

The formulation was prepared by mixing the water and propylene glycol together and adding the methylparaben, propylparaben and sodium sulfacetamide to form a uniform solution. The solution was heated to 70°C
20 and the sulfur was dispersed in the solution with moderate stirring. Separately, cetyl alcohol, emulsifying wax, and BRIJ 76 were melted together and heated to 70°C. The water and oil phases were combined and mixed for 10 minutes at high shear to form the emulsion. The emulsion was allowed to cool to 45°C with moderate stirring at which time trolamine was
25 added to the formulation and the formulation was adjusted to 100% with water.

The final formulation consisted of 91% by weight emulsion concentrate and 9% by weight HFC 134a propellant.

Example 2. Reduced Odor Topical Formulation Containing Urea

30 A topical formulation containing urea was prepared, wherein the formulation exhibited diminished odor. The composition of the formulation

and the physical and mechanical properties of the formulation are shown in Table 1.

The formulation was prepared by mixing the water and propylene glycol together and adding the methylparaben, propylparaben and urea to form a uniform solution. The solution was heated to 70°C with moderate stirring. Separately the cetyl alcohol, emulsifying wax and BRIJ 76 were melted together and heated to 70°C. The water and oil phases were combined and mixed for 10 minutes at high shear to form the emulsion. The emulsion was allowed to cool to 45°C with moderate stirring at which time the trolamine was added and the formulation was adjusted to 100% with water.

The final formulation consisted of 91% by weight emulsion concentrate and 9% by weight HFC 134a propellant.

Table 1. Formulation Composition and Physical and Mechanical Properties of Examples 1 and 2

Ingredient	Example 1 (w/w%)	Example 2 (w/w%)
Propylene Glycol USP	10	10.1
Cetyl Alcohol, NF	0.7	0.75
Trolamine, NF	0.1	0.1
Emulsifying Wax, NF	1.5	1.5
BRIJ 76	0.5	0.5
Water	72.06	76.905
Methylparaben, USP/NF	0.11	0.11
Propylparaben, USP/NF	0.03	0.035
Na Sulfacetamide, USP	10	0
Sulfur, USP	5	0
Urea, USP	0	10
Total	100	100

Mechanical Property	Example 1	Example 2
Foam Yield Stress	2074 dynes/cm ²	2143 dynes/cm ²
Foam Flow Index	0.525	0.582
Foam Zero Shear Viscosity	105600 cP	118200 cP
Conc Yield Stress	7081 dynes/cm ²	3355 dynes/cm ²
Conc Consistency Index	17286 cP	13511 cP
Conc Flow Index	0.2818	0.397
Conc Zero Shear Viscosity	169600 cP	106400 cP
Conc Infinite Shear Viscosity	1884 cP	3017 cP

Example 3. Organoleptic Analysis of Formulation Odor and Color

The odor and color of the formulations outlined in Table 1 were measured. Samples of the sulfur and urea aerosol foams were dispensed into weighing boats in a manner similar to that used to dispense the product for use. The samples of sulfur and urea emulsion concentrates were observed for color and odor in bulk packaging under conditions similar to that in which currently marketed products are used. Panelists were asked to rate on a scale of 0 to 5 each sample for the attributes of color and odor. In the scale, 0 corresponded to no detectable odor or color and 5 corresponded to strong odor or color. The results are shown in Table 2.

Table 2. Organoleptic Analysis of Formulation Odor and Color	Sulfacetamide/Sulfur				Urea	
	Foam		Concentrate		Foam	Concentrat e
	Odor Score	Color Score	Odor Score	Color Score	Odor Score	Odor Score
Average	0.7	1.5	2.0	3.9	0.1	4.3
Std Dev	0.8	0.8	1.2	1.1	0.3	1.3
p Value	0.000 5	0.000 1	n/a	n/a	≤ 0.000 1	n/a
n	10	10	10	10	10	10

Example 4. Reduced Odor Topical Formulation containing papain and urea

5. A topical formulation containing papain and urea was prepared, wherein the formulation exhibited diminished color. The composition of the formulation is shown in Table 3.

The formulation was prepared by mixing the water and propylene glycol together and adding methylparaben, propylparaben, lactose, urea and sodium phosphate monobasic to form a uniform solution. The solution was then heated to 70°C with moderate stirring. Separately the cetyl alcohol, emulsifying wax and BRIJ 76 were melted together and heated to 70°C. The water and oil phases were combined and mixed for 10 minutes at high shear to form the emulsion. The emulsion was then allowed to cool to 45°C with moderate stirring at which time the papain was added and the formulation was adjusted to 100% with water.

The final formulation consisted of 91% by weight emulsion concentrate and 9% by weight HFC 134a propellant.

Table 3. Papain-Urea Formulation Composition

Ingredient	Example 4 (w/w%)
Propylene Glycol USP	5
Cetyl Alcohol, NF	0.7
Sodium Phosphate monobasic	0.1
Glycerol, USP	5
Lactose, USP	1
Sodium Bisulfite	1
Emulsifying Wax, NF	1.5
BRIJ 76	0.5
Water	72.26
Methylparaben, USP/NF	0.11
Propylparaben, USP/NF	0.03
Papain, USP	2.8
Urea, USP	10

Example 5: Salicylic Acid Foam Formulation

A concentrate was prepared containing the following ingredients:

5

Ingredient	% W/W
Crodafos CS 20A	1
Cetostearyl Alcohol	0.5
Crodafos CES	1
Tocopheryl Acetate USP	0.5
White Petrolatum	0.5
DI Water	66.92
Glycerin USP	5
Disodium EDTA	0.05
Aloe (Aloe Vera Gel)	0.1
5N Sodium Hydroxide solution	0.4
Salicylic Acid USP	6
Ammonium Lactate	5
5N Sodium Hydroxide solution	10.807
Propylene Glycol USP	3
Methyl Paraben	0.3
Propyl Paraben	0.03
Total	100

The concentrate and foam were prepared as described in Examples 1 and 2.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of skill in the art to which the disclosed invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are as described. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

We claim:

1. A topical aerosol foam formulation comprising:
 - (a) one or more keratolytic agents dissolved or dispersed in an oil-in-water emulsion; and
 - (b) a propellant consisting essentially of one or more hydrofluoroalkanes contacting the emulsion to produce an immediate foaming action on expulsion from a pressurized container.
2. The formulation of claim 1, wherein the keratolytic agent is selected from the group consisting of urea, salicylic acid, papain, sulfur, glycolic acid, pyruvic acid, alpha hydroxy acids, beta hydroxy acids, resorcinol, N-acetylcysteine, retinoids, retinoic acid, coal tar, derivatives thereof, and combinations thereof.
3. The formulation of claim 1, wherein the concentration of the keratolytic agent is from about 1% to 60%.
4. The formulation of claim 1, further comprising one or more active agents selected from the group comprising of antibiotic agents, antimicrobial agents, anti-acne agents, antibacterial agents, antifungal agents, antiviral agents, steroidal anti-inflammatory agents, non-steroidal anti-inflammatory agents, anesthetics, antipruriginous agents, antiprotozoal agents, anti-oxidants, antihistamines, hormones, vitamins, skin-soothing agents, sunscreens, and combinations thereof.
5. The formulation of claim 1, wherein the foam is stable for at least 5 minutes at body temperature.
6. The formulation of claim 1, wherein the foam is stable for at least 20 minutes at body temperature.
7. The formulation of claim 2, wherein the keratolytic agent is urea.
8. The formulation of claim 7, wherein the amount of urea is from about 5% to about 50% (w/w).
9. The formulation of claim 8, wherein the amount of urea is from about 10% to about 50% (w/w).
10. The formulation of claim 8, wherein the amount of urea is from about 20% to about 40% (w/w).

11. The formulation of claim 7, further comprising ammonium lactate.
12. The formulation of claim 11, wherein the amount of ammonium lactate is from about 1% to about 30% (w/w).
13. The formulation of claim 12, wherein the amount of ammonium lactate is from about 5% to about 20% (w/w).
14. The formulation of claim 12, wherein amount of ammonium lactate is from about 10% to about 15% (w/w).
15. The formulation of claim 2, wherein the keratolytic agent is papain.
16. The formulation of 15, wherein the papain is present in an amount from about 0.5% to about 40%.
17. The formulation of 16, wherein the papain is present in an amount from about 1% to about 20%.
18. The formulation of 16, wherein the papain is present in an amount from about 1% to about 10%.
19. The formulation of claim 15, further comprising urea.
20. The formulation of claim 19, wherein the amount of urea is from about 1% to about 60% (w/w).
21. The formulation of claim 20, wherein the amount of urea is from about 2.5% to about 20% (w/w).
22. The formulation of claim 21, wherein the amount of urea is from about 5% to about 15% (w/w).
23. The formulation of claim 19, further comprising chlorophyllin copper complex sodium.
24. The formulation of claim 23, wherein the amount of chlorophyllin copper complex sodium is from about 0.5% to about 40% (w/w).
25. The formulation of claim 24, wherein the amount of chlorophyllin copper complex sodium is from about 1% to about 20% (w/w).
26. The formulation of claim 24, wherein the amount of chlorophyllin copper complex sodium is from about 1% to about 10% (w/w).
27. The formulation of claim 2, wherein the one or more keratolytic agents are sulfur and sodium sulfacetamide.

28. The formulation of claim 27, wherein the amounts of sulfur and sodium sulfacetamide are each from about 0.01% to about 20% (w/w).
29. The formulation of claim 28, wherein the amounts of sulfur and sodium sulfacetamide are each from about 1% to about 15% (w/w).
30. The formulation of claim 28, wherein the amounts of sulfur and sodium sulfacetamide are each from about 6% to about 12% (w/w).
31. The formulation of claim 27, further comprising one or more agents selected from the group consisting of vitamins, skin-soothing agents, and sunscreens.
32. The formulation of claim 2, wherein the keratolytic agent is salicylic acid.
33. The formulation of claim 32, wherein the amount of salicylic acid is from about 1% to about 30% (w/w).
34. The formulation of claim 33, wherein the amount of salicylic acid is from about 4% to about 10% (w/w).
35. The formulation of claim 32, further comprising one or more agents selected from the group consisting of vitamins, skin-soothing agents, and sunscreens.
36. The formulation of claim 4, wherein the antibiotic is selected from the group consisting of sodium sulfacetamide, clindamycin, erythromycin, and metronidazole.
37. The formulation of claim 36, wherein the antibiotic is present in an amount from about 0.01% to about 20% by weight of the composition.
38. The formulation of claim 37, wherein the antibiotic is present in an amount from about 1% to about 15% by weight of the composition.
39. The formulation of claim 37, wherein the antibiotic is present in an amount from about 6% to about 12% by weight of the composition.
40. The formulation of claim 1, further comprising one or more excipients selected from the group consisting of surfactants, emollients, emulsifiers, stabilizing agents, chelating agents, antioxidants, buffers, stabilizers, preservatives and combinations thereof.

41. The formulation of claim 1, wherein the hydrofluorocarbon propellant is present in an amount from about 5% to about 30% by weight of the composition.

42. The formulation of claim 1, wherein the hydrofluorocarbon is selected from the group consisting of 1,1,1,2-tetrafluoroethane (134a); 1,1,1,2,3,3,3-heptafluoropropane (227); and combinations thereof.

43. The formulation of claim 1, wherein the yield stress of the dispensed foam is between 250 and 60,000 dynes/cm².

44. The formulation of claim 1, where the zero shear viscosity of the dispensed foam is between 15,000 and 700,000 cP.

45. The formulation of claim 1, wherein the expansion factor of the dispensed foam is between 1.5 and 15 cm³/g.

46. The formulation of claim 1, wherein the foam density is from about 0.1 g/mL to about 0.6 g/mL.

47. The formulation of claim 1, further comprising a fragrance or odor masking agent.

48. A method of making a topical foam aerosol formulation, the method comprising:

(a) making an oil in water emulsion with a predominantly, more than 50%, aqueous phase,

(b) dissolving or dispersing one or more keratolytic agents in the aqueous phase or oil phase prior to emulsification, and

(c) adding a propellant consisting essentially of a hydrofluoroalkane or a mixture of hydrofluoroalkanes, without additional co-solvents or co-propellants, to the emulsion, wherein the emulsion produces an immediate foaming action on expulsion from a pressurized container.

49. A HFA containing topical foam formulation free of volatile lower alcohols produced by the method of claim 48.