



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2477973 A1 2003/09/18

(21) **2 477 973**

**(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2003/03/07
(87) Date publication PCT/PCT Publication Date: 2003/09/18
(85) Entrée phase nationale/National Entry: 2004/09/01
(86) N° demande PCT/PCT Application No.: US 2003/007101
(87) N° publication PCT/PCT Publication No.: 2003/075850
(30) Priorités/Priorities: 2002/03/07 (60/362,080) US;
2002/06/13 (10/167,491) US

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/365, A61K 31/35, A61K 31/335

(71) **Demandeur/Applicant:**
BLANCHETTE ROCKEFELLER NEUROSCIENCES
INSTITUTE, US

(72) **Inventeurs/Inventors:**
ETCHEBERRIGARAY, RENE, US;
ALKON, DANIEL L., US

(74) **Agent:** OGILVY RENAULT

(54) Titre : **METHODES DE TRAITEMENT DE LA MALADIE D'ALZHEIMER ET AMELIORATION NEURO-COGNITIVE**
(54) Title: **METHODS FOR ALZHEIMER'S DISEASE TREATMENT AND COGNITIVE ENHANCEMENT**

(57) Abrégé/Abstract:

The present invention relates to compositions and methods to modulate α -secretase and/or to improve cognitive ability. The invention further relates to the improved/enhanced cognitive ability in diseased individuals, particularly Alzheimer's Disease patients, and treatment thereof through increased sAPP production. Macroyclic lactones (i.e. bryostatin class and neristatin class) are compounds preferred for use with the present composition. The present invention also provides methods for increasing the generation of non-amyloidogenic soluble APP comprising the activation of protein kinase C (PKC) by administering an effective amount of PKC activator(s).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 September 2003 (18.09.2003)

PCT

(10) International Publication Number
WO 2003/075850 A3

(51) International Patent Classification⁷: **A61K 31/35**, 31/335

(21) International Application Number: PCT/US2003/007101

(22) International Filing Date: 7 March 2003 (07.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/362,080 7 March 2002 (07.03.2002) US
10/167,491 13 June 2002 (13.06.2002) US

(71) Applicant (for all designated States except US):
BLANCHETTE ROCKEFELLER NEURO-SCIENCES INSTITUTE [US/US]; 9601 Medical Center Drive, 3rd Floor, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ETCHEBER-RIGARAY, Rene** [US/US]; 7604 Westlake Terrace, Bethesda, MD 20817 (US). **ALKON, Daniel, L.** [US/US]; 6701 Bonaventure Court, Bethesda, MD 20817 (US).

(74) Agents: **STOLE, Einar** et al.; Milbank, Tweed, Hadley & McCloy LLP, 1825 Eye Street, N.W., Suite 1100, Washington, DC 20006 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report: 8 January 2004

(48) Date of publication of this corrected version: 19 February 2004

(15) Information about Correction:
see PCT Gazette No. 08/2004 of 19 February 2004, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2003/075850 A3

(54) Title: METHODS FOR ALZHEIMER'S DISEASE TREATMENT AND COGNITIVE ENHANCEMENT

(57) Abstract: The present invention relates to compositions and methods to modulate α -secretase and/or to improve cognitive ability. The invention further relates the improved/enhanced cognitive ability in diseased individuals, particularly Alzheimer's Disease patients, and treatment thereof through increased sAPP production. Macrocyclic lactones (i.e. bryostatin class and neristatin class) are compounds preferred for use with the present composition. The present invention also provides methods for increasing the generation of non-amyloidogenic soluble APP comprising the activation of protein kinase C (PKC) by administering an effective amount of PKC activator(s).

Methods for Alzheimer's Disease Treatment and Cognitive Enhancement

BACKGROUND OF THE INVENTION

(i) Field of the Invention

5 The present invention relates to the modulation of α -secretase and cognitive enhancement. The invention further relates to compounds for treatment of conditions associated with amyloid processing such as Alzheimer's Disease and compositions for the treatment of such conditions.

(ii) Background of the Invention

10 Various disorders and diseases exist which affect cognition. Cognition can be generally described as including at least three different components: attention, learning, and memory. Each of these components and their respective levels affect the overall level of a subject's cognitive ability. For instance, while Alzheimer's Disease patients suffer from a loss of overall cognition and thus deterioration of each of these 15 characteristics, it is the loss of memory that is most often associated with the disease. In other diseases patients suffer from cognitive impairment that is more predominately associated with different characteristics of cognition. For instance Attention Deficit Hyperactivity Disorder (ADHD), focuses on the individual's ability to maintain an attentive state. Other conditions include general dementias associated with other 20 neurological diseases, aging, and treatment of conditions that can cause deleterious effects on mental capacity, such as cancer treatments, stroke/ischemia, and mental retardation.

25 Cognition disorders create a variety of problems for today's society. Therefore, scientists have made efforts to develop cognitive enhancers or cognition activators. The cognition enhancers or activators that have been developed are generally classified to include nootropics, vasodilators, metabolic enhancers, psychostimulants, cholinergic agents, biogenic amine drugs, and neuropeptides. Vasodilators and metabolic enhancers (e.g. dihydroergotoxine) are mainly effective in the cognition disorders induced by cerebral vessel ligation-ischemia; however, they are ineffective in clinical

use and with other types of cognition disorders. Of the developed cognition enhancers, typically only metabolic drugs are employed for clinical use, as others are still in the investigation stage. Of the nootropics for instance, piracetam activates the peripheral endocrine system, which is not appropriate for Alzheimer's disease due to the high 5 concentration of steroids produced in patients while tacrine, a cholinergic agent, has a variety of side effects including vomiting, diarrhea, and hepatotoxicity.

Ways to improve the cognitive abilities of diseased individuals have been the subject of various studies. Recently the cognitive state related to Alzheimer's Disease and different ways to improve patient's memory have been the subject of various 10 approaches and strategies. Unfortunately, these approaches and strategies only improve symptomatic and transient cognition in diseased individuals but have not addressed the progression of the disease. In the case of Alzheimer's Disease, efforts to improve cognition, typically through the cholinergic pathways or though other brain transmitter pathways, have been investigated. The primary approach relies on the inhibition of 15 acetyl cholinesterase enzymes through drug therapy. Acetyl cholinesterase is a major brain enzyme and manipulating its levels can result in various changes to other neurological functions and cause side effects.

While these and other methods may improve cognition, at least transiently, they do not modify the disease progression, or address the cause of the disease. For instance, 20 Alzheimer's Disease is typically associated with the formation of plaques through the accumulation of amyloid precursor protein. Attempts to illicit an immunological response through treatment against amyloid and plaque formation have been done in animal models, but have not been successfully extended to humans.

Furthermore, cholinesterase inhibitors only produce some symptomatic improvement for a short time and in only a fraction of the Alzheimer's Disease patients with mid to moderate symptoms and are thus only a useful treatment for a small portion 25 of the overall patient population. Even more critical is that present efforts at improving cognition do not result in treatment of the disease condition, but are merely ameliorative

of the symptoms. Current treatments do not modify the disease progression. These treatments have also included the use of a "vaccine" to treat the symptoms of Alzheimer's Disease patients which, while theoretically plausible and effective in mice tests, have been shown to cause severe adverse reactions in humans.

5 As a result, use of the cholinergic pathway for the treatment of cognitive impairment, particularly in Alzheimer's Disease, has proven to be inadequate. Additionally, the current treatments for cognitive improvement are limited to specific neurodegenerative diseases and have not proven effective in the treatment of other cognitive conditions.

10 Alzheimer's disease is associated with extensive loss of specific neuronal subpopulations in the brain with memory loss being the most universal symptom. (Katzman, R. (1986) New England Journal of Medicine 314:964). Alzheimer's disease is well characterized with regard to neuropathological changes. However, abnormalities have been reported in peripheral tissue supporting the possibility that Alzheimer's 15 disease is a systemic disorder with pathology of the central nervous system being the most prominent. (Connolly, G., Fibroblast models of neurological disorders: fluorescence measurement studies, Review, TiPS Vol. 19, 171-77 (1998)). For a discussion of Alzheimer's disease links to a genetic origin and chromosomes 1, 14, and 21 see St. George-Hyslop, P. H., et al., Science 235:885 (1987); Tanzi, Rudolph et al., 20 The Gene Defects Responsible for Familial Alzheimer's Disease, Review, Neurobiology of Disease 3, 159-168 (1996); Hardy, J., Molecular genetics of Alzheimer's disease, Acta Neurol Scand: Supplement 165: 13-17 (1996).

25 While cellular changes leading to neuronal loss and the underlying etiology of the disease remain under investigation the importance of APP metabolism is well established. The two proteins most consistently identified in the brains of patients with Alzheimer's disease to play a role in the physiology or pathophysiology of brain are β -amyloid and tau. (See Selkoe, D., Alzheimer's Disease: Genes, Proteins, and Therapy, Physiological Reviews, Vol. 81, No. 2, 2001). A discussion of the defects in β -amyloid

protein metabolism and abnormal calcium homeostasis and/or calcium activated kinases. (Etcheberrigaray et al., Calcium responses are altered in fibroblasts from Alzheimer's patients and pre-symptomatic PS1 carriers: a potential tool for early diagnosis, *Alzheimer's Reports*, Vol. 3, Nos. 5 & 6, pp. 305-312 (2000); Webb et al., 5 Protein kinase C isozymes: a review of their structure, regulation and role in regulating airways smooth muscle tone and mitogenesis, *British Journal of Pharmacology*, 130, pp 1433-52 (2000)).

Further with regard to normal and abnormal memory both K^+ and Ca^{2+} channels have been demonstrated to play key roles in memory storage and recall. For instance, 10 potassium channels have been found to change during memory storage. (Etcheberrigaray, R., et al. (1992) *Proceeding of the National Academy of Science* 89:7184; Sanchez-Andres, J. V. and Alkon, D. L. (1991) *Journal of Neurobiology* 65:796; Collin, C., et al. (1988) *Biophysics Journal* 55:955; Alkon, D. L., et al. (1985) 15 *Behavioral and Neural Biology* 44:278; Alkon, D. L. (1984) *Science* 226:1037). This observation, coupled with the almost universal symptom of memory loss in Alzheimer's patients, led to the investigation of potassium channel function as a possible site of Alzheimer's disease pathology and the effect of PKC modulation on cognition.

PKC was identified as one of the largest gene families of non-receptor serine-threonine protein kinases. Since the discovery of PKC in the early eighties by 20 Nishizuka and coworkers (Kikkawa et al., *J. Biol. Chem.*, 257, 13341 (1982), and its identification as a major receptor for phorbol esters (Ashendel et al., *Cancer Res.*, 43, 4333 (1983)), a multitude of physiological signaling mechanisms have been ascribed to this enzyme. The intense interest in PKC stems from its unique ability to be activated *in vitro* by calcium and diacylglycerol (and its phorbol ester mimetics), an effector 25 whose formation is coupled to phospholipid turnover by the action of growth and differentiation factors.

The PKC gene family consists presently of 11 genes which are divided into four subgroups: 1) classical PKC α , β_1 , β_2 (β_1 and β_2 are alternatively spliced forms of the

same gene) and γ , 2) novel PKC δ , ϵ , η and θ , 3) atypical PKC ζ , λ , η and ι and 4) PKC μ . PKC μ resembles the novel PKC isoforms but differs by having a putative transmembrane domain (reviewed by Blohe et al., *Cancer Metast. Rev.*, 13, 411 (1994); Ilug et al., *Biochem j.*, 291, 329 (1993); Kikkawa et al., *Ann. Rev. Biochem.* 58, 31 (1989)). The α , β_1 , β_2 , and γ isoforms are Ca^{2+} , phospholipid and diacylglycerol-dependent and represent the classical isoforms of PKC, whereas the other isoforms are activated by phospholipid and diacylglycerol but are not dependent on Ca^{2+} . All isoforms encompass 5 variable (V1-V5) regions, and the α , β , γ isoforms contain four (C1-C4) structural domains which are highly conserved. All isoforms except PKC α , β and γ lack the C2 domain, and the λ , η and ι isoforms also lack nine of two cysteine-rich zinc finger domains in C1 to which diacylglycerol binds. The C1 domain also contains the pseudosubstrate sequence which is highly conserved among all isoforms, and which serves an autoregulatory function by blocking the substrate-binding site to produce an inactive conformation of the enzyme (House et al., *Science*, 238, 1726 (1987)).

Because of these structural features, diverse PKC isoforms are thought to have highly specialized roles in signal transduction in response to physiological stimuli (Nishizuka, *Cancer*, 10, 1892 (1989)), as well as in neoplastic transformation and differentiation (Glazer, *Protein Kinase C*, J. F. Kuo, ed., Oxford U. Press (1994) at pages 171-198). For a discussion of known PKC modulators see PCT/US97/08141, U.S. Patent Nos. 5,652,232; 6,043,270; 6,080,784; 5,891,906; 5,962,498; 5,955,501; 5,891,870 and 5,962,504.

In view of the central role that PKC plays in signal transduction, PKC has proven to be an exciting target for the modulation of APP processing. It is well established that PKC plays a role in APP processing. Phorbol esters for instance have been shown to significantly increase the relative amount of non-amyloidogenic soluble APP (sAPP) secreted through PKC activation. Activation of PKC by phorbol ester does not appear to result in a direct phosphorylation of the APP molecule, however. Irrespective of the precise site of action, phorbol-induced PKC activation results in an

enhanced or favored α -secretase, non-amyloidogenic pathway. Therefore PKC activation is an attractive approach for influencing the production of non-deleterious sAPP and even producing beneficial sAPP and at the same time reduce the relative amount of $A\beta$ peptides. Phorbol esters, however, are not suitable compounds for 5 eventual drug development because of their tumor promotion activity. (Ibarreta, et al., Benzolactam (BL) enhances sAPP secretion in fibroblasts and in PC12 cells, NeuroReport, Vol. 10, No. 5&6, pp 1035-40 (1999)).

There is increasing evidence that the individual PKC isozymes play different, sometimes opposing, roles in biological processes, providing two directions for 10 pharmacological exploitation. One is the design of specific (preferably, isozyme specific) inhibitors of PKC. This approach is complicated by the fact that the catalytic domain is not the domain primarily responsible for the isotype specificity of PKC. The other approach is to develop isozyme-selective, regulatory site-directed PKC activators. These may provide a way to override the effect of other signal transduction pathways 15 with opposite biological effects. Alternatively, by inducing down-regulation of PKC after acute activation, PKC activators may cause long term antagonism. Bryostatin is currently in clinical trials as an anti-cancer agent. The bryostatins are known to bind to the regulatory domain of PKC and to activate the enzyme. Bryostatin is an example of isozyme-selective activators of PKC. Compounds in addition to bryostatins have been 20 found to modulate PKC. (see for example WO 97/43268)

There still exists a need for the development of methods for the treatment for improved overall cognition, either through a specific characteristic of cognitive ability or general cognition. There also still exists a need for the development of methods for the improvement of cognitive enhancement whether or not it is related to specific 25 disease state or cognitive disorder. The methods and compositions of the present invention fulfill these needs and will greatly improve the clinical treatment for Alzheimer's disease and other neurodegenerative diseases, as well as, provide for improved cognitive enhancement. The methods and compositions also provide

treatment and/or enhancement of the cognitive state through the modulation of α -secretase.

SUMMARY OF THE INVENTION

The invention relates to compounds, compositions, and methods for the treatment of conditions associated with enhancement/improvement of cognitive ability. In a preferred embodiment, the present invention further relates to compounds, compositions and methods for the treatment of conditions associated with amyloid processing, such as Alzheimer's Disease, which provides for improved/enhanced cognitive ability in the subject treated. In particular the compounds and compositions of the present invention are selected from macrocyclic lactones (i.e. bryostatin class and neristatin class).

In another aspect the invention relates to macrocyclic lactone compounds, compositions and methods that modulate α -secretase activity. Of particular interest are the bryostatin and neristatin class compounds, and of further interest is bryostatin-1.

Another aspect of the invention relates to the bryostatin and neristatin class compounds, as a PKC activator, to alter conditions associated with amyloid processing in order to enhance the α -secretase pathway to generate soluble α -amyloid precursor protein (α APP) so as to prevent β -amyloid aggregation and improve/enhance cognitive ability. Such activation, for example, can be employed in the treatment of Alzheimer's Disease. Of particular interest is bryostatin-1.

In another aspect, the invention relates to a method for treating plaque formation, such as that associated with Alzheimer's Disease, and improving/enhancing the cognitive state of the subject comprising administering to the subject an effective amount of a macrocyclic lactone to activate PKC. In a preferred embodiment, the PKC activator is of the bryostatin or neristatin class of compounds. In a more preferred embodiment the compound is bryostatin-1.

Another aspect of the invention relates to a composition for treating plaque formation and improving/enhancing cognitive ability comprising: (i) a macrocyclic

lactone in an amount effective to elevate soluble β -amyloid, generate soluble α APP and prevent β -amyloid aggregation; and (ii) a pharmaceutically effective carrier. In a preferred embodiment the composition is used to improve/enhance cognitive ability associated with Alzheimer's Disease. The macrocyclic lactone is preferably selected
5 from the bryostatin or neristatin class compounds, particularly bryostatin-1.

In one embodiment of the invention the activation of PKC isoenzymes results in improved cognitive abilities. In one embodiment the improved cognitive ability is memory. In another embodiment the improved cognitive ability is learning. In another embodiment the improved cognitive ability is attention. In another embodiment PKC's isoenzymes are activated by a macrocyclic lactone (i.e. bryostatin class and neristatin class). In particular, bryostatin-1 through 18 and neristatin is used to activate the PKC isoenzyme. In a preferred embodiment bryostatin-1 is used.

In another aspect, the invention comprises a composition of a PKC isoenzyme activator administered in an amount effective to improve cognitive abilities. In a preferred embodiment the PKC isoenzyme activator is selected from macrocyclic lactones (i.e. bryostatin class and neristatin class). In a preferred embodiment the amount of PKC activator administered is in an amount effective to increase the production of sAPP. In a more preferred embodiment the amount of composition administered does not cause myalgia.

20 In a preferred embodiment the PKC isoenzymes are activated in subjects, which are suffering or have suffered from neurological diseases, strokes or hypoxia. In a more preferred embodiment the PKC isoenzyme is activated in Alzheimer's Disease subjects or models.

25 In another embodiment of the invention the PKC activation results in the modulation of amyloid precursor protein metabolism. Further the modulation by the PKC activation results in an increase in the alpha secretase pathway. The alpha secretase pathway results in non-toxic, non-amyloidogenic fragments related to cognitive impairment. As a result the cognitive condition of the subject improves. In

another embodiment of the invention the PKC activation reduces the amyloidogenic and toxic fragments Abeta 40 and Ab42.

Another embodiment of the invention is a method of improving cognitive ability through the activation of PKC isoenzymes. In another embodiment of the invention the PKC activation occurs in “normal” subjects. In another embodiment of the invention the PKC activation occurs in subjects suffering from a disease, deteriorating cognitive faculties, or malfunctioning cognition. In a preferred embodiment the method is a method for treating Alzheimer’s Disease.

In another embodiment of the invention the modulation of PKC is through the use of a non-tumor promoting agent resulting in improved cognitive abilities. In a preferred embodiment the PKC activator is selected from bryostatin-1 through bryostatin-18 and neristatin. In a more preferred embodiment bryostatin-1 is used. In another embodiment bryostatin-1 is used in combination with a non-bryostatin class compound to improve cognitive ability and reduce side effects.

In another embodiment of the invention, the modulation of PKC through macrocyclic lactones (i.e. bryostatin class and neristatin class) is used *in vitro* for the testing of conditions associated with Alzheimer’s Disease. The *in vitro* use may include for example, the testing of fibroblast cells, blood cells, or the monitoring of ion channel conductance in cellular models.

In a preferred embodiment of the invention the compounds and compositions are administered through oral and/or injectable forms including intravenously and intraventricularly.

The present invention therefore provides a method of treating impaired memory or a learning disorder in a subject, the method comprising administering thereto a therapeutically effective amount of one of the present compounds. The present compounds can thus be used in the therapeutic treatment of clinical conditions in which memory defects or impaired learning occur. In this way memory and learning can be improved. The condition of the subject can thereby be improved.

The compositions and methods have utility in treating clinical conditions and disorders in which impaired memory or a learning disorder occurs, either as a central feature or as an associated symptom. Examples of such conditions which the present compounds can be used to treat include Alzheimer's disease, multi-infarct dementia and 5 the Lewy-body variant of Alzheimer's disease with or without association with Parkinson's disease; Creutzfeld-Jakob disease and Korsakow's disorder.

The compositions and methods can also be used to treat impaired memory or learning which is age-associated, is consequent upon electro-convulsive therapy or which is the result of brain damage caused, for example, by stroke, an anesthetic 10 accident, head trauma, hypoglycemia, carbon monoxide poisoning, lithium intoxication or a vitamin deficiency.

The compounds have the added advantage of being non-tumor promoting and already being involved in phase II clinical trials.

The invention relates to a pharmaceutical composition for enhancing cognition, 15 preventing and/or treating cognition disorders. More particularly, it relates to the pharmaceutical composition comprising macrocyclic lactones (i.e. bryostatin class and neristatin class) and their derivatives as the active ingredient for enhancing cognition, preventing and/or treating cognition disorders.

It is therefore a primary object of the invention to provide pharmaceutical 20 compositions for enhancing cognition, preventing and/or treating cognition disorders. The pharmaceutical composition comprises macrocyclic lactones, particularly the bryostatin and neristatin class, or a pharmaceutically acceptable salt or derivative thereof, and a pharmaceutically acceptable carrier or excipient.

The pharmaceutical composition according to the invention is useful in the 25 enhancement of cognition, prophylaxis and/or treatment of cognition disorders, wherein cognition disorders include, but are not limited to, disorders of learning acquisition, memory consolidation, and retrieval, as described herein.

The invention concerns a method for the treatment of amyloidosis associated with neurological diseases, including Alzheimer's disease by administering to a patient an effective amount of at least one agent that modulates or affects the phosphorylation of proteins in mammalian cells.

5 The invention also provides a method for treating Alzheimer's disease comprising administering to a patient an effective amount of a macrocyclic lactone (i.e. bryostatin class and neristatin class).

10 In another embodiment the bryostatin or neristatin class compounds may be used in the above methods in combination with different phorbol esters to prevent or reduce tumorigenic response in the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 illustrates the effect of different PKC inhibitors on sAPP α secretion with Bryostatin-1 showing greater efficacy at lower concentrations than controls and Benzolactam.

15 Fig. 2 illustrates the effect of different concentrations of Bryostatin-1 on the PKC α isozyme.

Fig. 3 illustrates the effect of different concentrations of Bryostatin-1 on sAPP α secretion.

20 Fig. 4 illustrates the amount of time required for treated rats verse controls to learn a water maze.

Fig. 5(a) illustrates the amount of time control rats spent swimming in the different quadrants.

Fig. 5(b) illustrates the amount of time treated rats spent swimming in the different quadrants.

25 Fig. 5(c) illustrates the different between the amount of time the treated rats spent in target quadrant compared to control rats.

Fig. 6 illustrates sAPP α secretion in human fibroblast cells following administration of bryostatin 0.1 nM for both controls and AD cells.

Fig. 7 illustrates an immunoblot for sAPP following administration of bryostatin in AD cells.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Memory loss and impaired learning ability are features of a range of clinical 5 conditions. For instance, loss of memory is the most common symptom of dementia states including Alzheimer's disease. Memory defects also occur with other kinds of dementia such as multi-infarct dementia (MID), a senile dementia caused by cerebrovascular deficiency, and the Lewy-body variant of Alzheimer's disease with or without association with Parkinson's disease, or Creutzfeld-Jakob disease. Loss of 10 memory is a common feature of brain-damaged patients. Brain damage may occur, for example, after a classical stroke or as a result of an anesthetic accident, head trauma, hypoglycemia, carbon monoxide poisoning, lithium intoxication, vitamin (B1, thiamine and B12) deficiency, or excessive alcohol use or Korsakow's disorder. Memory impairment may furthermore be age-associated; the ability to recall information such as 15 names, places and words seems to decrease with increasing age. Transient memory loss may also occur in patients, suffering from a major depressive disorder, after electro-convulsive therapy (ECT). Alzheimer's disease is in fact the most important clinical entity responsible for progressive dementia in ageing populations, whereas hypoxia/stroke is responsible for significant memory defects not related to neurological 20 disorders.

Individuals with Alzheimer's disease are characterized by progressive memory impairments, loss of language and visuospatial skills and behavior deficits (McKhann et al., 1986, *Neurology*, 34:939-944). The cognitive impairment of individuals with Alzheimer's disease is the result of degeneration of neuronal cells located in the cerebral cortex, hippocampus, basal forebrain and other brain regions. Histologic analyses of 25 Alzheimer's disease brains obtained at autopsy demonstrated the presence of neurofibrillary tangles (NFT) in perikarya and axons of degenerating neurons, extracellular neuritic (senile) plaques, and amyloid plaques inside and around some

blood vessels of affected brain regions. Neurofibrillary tangles are abnormal filamentous structures containing fibers (about 10 nm in diameter) that are paired in a helical fashion, therefore also called paired helical filaments. Neuritic plaques are located at degenerating nerve terminals (both axonal and dendritic), and contain a core compound of amyloid protein fibers. In summary, Alzheimer's disease is characterized by certain neuropathological features including intracellular neurofibrillary tangles, primarily composed of cytoskeletal proteins, and extracellular parenchymal and cerebrovascular amyloid. Further, there are now methods in the art for distinguishing between Alzheimer's patients, normal aged people, and people suffering from other neurodegenerative diseases, such as Parkinson's, Huntington's chorea, Wernicke-Korsakoff or schizophrenia further described for instance in U.S. patent 5,580,748 and U.S. patent 6,080582.

Alzheimer's disease (AD) is a brain disorder characterized by altered protein catabolism. Altered protein phosphorylation has been implicated in the formation of the intracellular neurofibrillary tangles found in Alzheimer's disease. A role for protein phosphorylation in the catabolism of the amyloid precursor protein (APP), from which is derived the major component of amyloid plaques found in AD, has also been investigated. A central feature of the pathology of Alzheimer's disease is the deposition of amyloid protein within plaques.

The processing of the amyloid precursor protein (APP) determines the production of fragments that later aggregate forming the amyloid deposits characteristic of Alzheimer's disease (AD), known as senile or AD plaques. Thus, APP processing is an early and key pathophysiological event in AD.

Three alternative APP processing pathways have been identified. The previously termed "normal" processing involves the participation of an enzyme that cleaves APP within the A β sequence at residue Lys16 (or between Lys16 and Leu17; APP770 nomenclature), resulting in non-amyloidogenic fragments: a large N-terminus ectodomain and a small 9 kDa membrane bound fragment. This enzyme, yet to be fully

identified, is known as α -secretase. Two additional secretases participate in APP processing. One alternative pathway involves the cleavage of APP outside the $A\beta$ domain, between Met671 and Asp672 (by β -secretase) and the participation of the endosomal-lysosomal system. An additional cleavage site occurs at the carboxyl-terminal 5 end of the $A\beta$ portion, within the plasma membrane after amino acid 39 of the $A\beta$ peptide. The secretase (γ) action produces an extracellular amino acid terminal that contains the entire $A\beta$ sequence and a cell-associated fragment of ~6kDa. Thus, processing by β and γ secretases generate potential amyloidogenic fragments since they contain the complete $A\beta$ sequence. Several lines of evidence have shown that all 10 alternative pathways occur in a given system and that soluble $A\beta$ may be a “normal product.” However, there is also evidence that the amount of circulating $A\beta$ in CSF and plasma is elevated in patients carrying the “Swedish” mutation. Moreover, cultured cells transfected with this mutation or the APP₇₁₇ mutation, secrete larger amounts of $A\beta$. More recently, carriers of other APP mutations and PS1 and PS2 mutations have 15 been shown to secrete elevated amounts of a particular form, long (42-43 amino acids) $A\beta$.

Therefore, although all alternative pathways may take place normally, an imbalance favoring amyloidogenic processing occurs in familial and perhaps sporadic AD. These enhanced amyloidogenic pathways ultimately lead to fibril and plaque 20 formation in the brains of AD patients. Thus, intervention to favor the non-amyloidogenic, α -secretase pathway effectively shifts the balance of APP processing towards a presumably non-pathogenic process that increases the relative amount of sAPP compared with the potentially toxic $A\beta$ peptides.

The PKC isoenzymes provides a critical, specific and rate limiting molecular 25 target through which a unique correlation of biochemical, biophysical, and behavioral efficacy can be demonstrated and applied to subjects to improve cognitive ability.

The present inventors have studied bryostatins as activators of protein kinase (PKC). Alterations in PKC, as well alterations in calcium regulation and potassium

(K⁺) channels are included among alterations in fibroblasts in Alzheimer's disease (AD) patients. PKC activation has been shown to restore normal K⁺ channel function, as measured by TEA-induced [Ca²⁺] elevations. Further patch-clamp data substantiates the effect of PKC activators on restoration of 113pS K⁺ channel activity. Thus PKC activator-based restoration of K⁺ channels has been established as an approach to the investigation of AD pathophysiology, and provides a useful model for AD therapeutics. (See pending applications 09/652,656 herein incorporated in its entirety.)

The use of peripheral tissues from Alzheimer's disease (AD) patients and animal neuronal cells permitted the identification of a number of cellular/molecular alterations reflecting comparable processes in the AD brain and thus, of pathophysiological relevance (Baker et al., 1988; Scott, 1993; Huang, 1994; Scheuner et al., 1996; Etcheberrigaray & Alkon, 1997; Gasparini et al., 1997). Alteration of potassium channel function has been identified in fibroblasts (Etcheberrigaray et al., 1993) and in blood cells (Bondy et al., 1996) obtained from AD patients. In addition, it was shown that β -amyloid, widely accepted as a major player in AD pathophysiology (Gandy & Greengard, 1994; Selkoe, 1994; Yankner, 1996), was capable of inducing an AD-like K⁺ channel alteration in control fibroblasts (Etcheberrigaray et al., 1994). Similar or comparable effects of β -amyloid on K⁺ channels have been reported in neurons from laboratory animals (Good et al., 1996; also for a review see Fraser et al., 1997). An earlier observation of hippocampal alterations of apamin-sensitive K⁺ channels in AD brains (as measured by apamin binding) provides additional support for the suggestion that K⁺ channels may be pathophysiological relevant in AD (Ikeda et al., 1991). Furthermore, protein kinase C (PKC) exhibits parallel changes in peripheral and brain tissues of AD patients. The levels and/or activity of this enzyme(s) were introduced in brains and fibroblasts from AD patients (Cole et al., 1988; Van Huynh et al., 1989; Govoni et al., 1993; Wang et al., 1994). Studies using immunoblotting analyses have revealed that of the various PKC isozymes, primarily the α isoform was significantly reduced in fibroblasts (Govoni et al., 1996), while both α and β isoforms are reduced in

brains of AD patients (Shimohama et al., 1993; Masliah et al., 1990). These brain PKC alterations might be an early event in the disease process (Masliah et al., 1991). It is also interesting to note that PKC activation appears to favor nonamyloidogenic processing of the amyloid precursor protein, APP (Buxbaum et al., 1990; Gillespie et al., 1992; Selkoe, 1994; Gandy & Greengard, 1994; Bergamashi et al., 1995; Desdouits et al., 1996; Efthimiopoulos et al., 1996). Thus, both PKC and K⁺ channel alterations coexist in AD, with peripheral and brain expression in AD.

The link between PKC and K⁺ channel alterations has been investigated because PKC is known to regulate ion channels, including K⁺ channels and that a defective PKC leads to defective K⁺ channels. This is important not only for the modulation of APP, but also for the role PKC and K⁺ channels play in memory establishment and recall. (e.g., see Alkon et al., 1988; Covarrubias et al., 1994; Hu et al., 1996) AD fibroblasts have been used to demonstrate both K⁺ channels and PKC defects (Etcheberrigaray et al., 1993; Govoni et al., 1993, 1996). Studies also show, fibroblasts with known dysfunctional K⁺ channels treated with PKC activators restore channel activity as monitored by the presence/absence of TEA-induced calcium elevations. Further, assays based on tetraethylammonium chloride (TEA)-induced [Ca²⁺] elevation have been used to show functional 113pS K⁺ channels that are susceptible to TEA blockade (Etcheberrigaray et al., 1993, 1994; Hirashima et al., 1996). Thus, TEA-induced [Ca²⁺] elevations and K⁺ channel activity observed in fibroblasts from control individuals are virtually absent in fibroblasts from AD patients (Etcheberrigaray et al., 1993; Hirashima et al., 1996). These studies demonstrate that the use of PKC activators can restore the responsiveness of AD fibroblast cell lines to the TEA challenge. Further, immunoblot evidence from these studies demonstrate that this restoration is related to a preferential participation of the α isoform.

The present inventors have also observed that activation of protein kinase C favors the α -secretase processing of the Alzheimer's disease (AD) amyloid precursor protein (APP), resulting in the generation of non-amyloidogenic soluble APP (sAPP).

Consequently, the relative secretion of amyloidogenic A₁₋₄₀ and A₁₋₄₂₍₃₎ is reduced. This is particularly relevant since fibroblasts and other cells expressing APP and presenilin AD mutations secrete increased amounts of total A β and/or increased ratios of A₁₋₄₂₍₃₎/A₁₋₄₀. Interestingly, PKC defects have been found in AD brain (α and β isoforms) and in fibroblasts (α -isoform) from AD patients.

Studies have shown that other PKC activators (i.e. benzolactam) with improved selectivity for the α , β and γ isoforms enhance sAPP secretion over basal levels. The sAPP secretion in benzolactam-treated AD cells was also slightly higher compared to control benzolactam-treated fibroblasts, which only showed significant increases of sAPP secretion after treatment with 10 μ M BL. It was further reported that staurosporine (a PKC inhibitor) eliminated the effects of benzolactam in both control and AD fibroblasts while related compounds also cause a ~3-fold sAPP secretion in PC12 cells. The present inventors have found that the use of bryostatin as a PKC activators to favor non-amyloidogenic APP processing is of particular therapeutic value since it is non-tumor promoting and already in stage II clinical trials.

Memories are thought to be a result of lasting synaptic modification in the brain structures related to information processing. Synapses are considered a critical site at final targets through which memory-related events realize their functional expression, whether the events involve changed gene expression and protein translation, altered kinase activities, or modified signaling cascades. A few proteins have been implicated in associative memory including Ca²⁺/calmodulin II kinases, protein kinase C, calexcitin, a 22-kDa learning-associated Ca²⁺ binding protein, and type II ryanodine receptors. The modulation of PKC through the administration of macrocyclic lactones provides a mechanism to effect synaptic modification.

The area of memory and learning impairment is rich in animal models that are able to demonstrate different features of memory and learning processes. (See, for example, Hollister, L. E., 1990, *Pharmacopsychiat.*, 23, (Suppl II) 33-36). The available animal models of memory loss and impaired learning involve measuring the

ability of animals to remember a discrete event. These tests include the Morris Water Maze and the passive avoidance procedure. In the Morris Water Maze, animals are allowed to swim in a tank divided into four quadrants, only one of which has a safety platform beneath the water. The platform is removed and the animals are tested for how long they search the correct quadrant versus the incorrect quadrants. In the passive avoidance procedure the animal remembers the distinctive environment in which a mild electric shock is delivered and avoids it on a second occasion. A variant of the passive avoidance procedure makes use of a rodent's preference for dark enclosed environments over light open ones. Further discussion can be found in Crawley, J. N., 1981, Pharmacol. Biochem. Behav., 15, 695-699; Costall, B. et al, 1987, Neuropharmacol., 26, 195-200; Costall, B. et al, 1989, Pharmacol. Biochem. Behav., 32, 777-785; Barnes, J. M. et al, 1989, Br. J. Pharmacol., 98 (Suppl) 693P; Barnes, J. M. et al, 1990, Pharmacol. Biochem. Behav., 35, 955-962.

The use of the word, "normal" is meant to include individuals who have not been diagnosed with or currently display diminished or otherwise impaired cognitive function. The different cognitive abilities may be tested and evaluated through known means well established in the art, including but not limited to tests from basic motor-spatial skills to more complex memory recall testing. Non-limiting examples of tests used for cognitive ability for non-primates include the Morris Water Maze, Radial Maze, T Maze, Eye Blink Conditioning, Delayed Recall, and Cued Recall while for primate subjects test may include Eye Blink, Delayed Recall, Cued Recall, Face Recognition, MiniMental, and ADAS-Cog. Many of these tests are typically used in the mental state assessment for patients suffering from AD. Similarly, the evaluation for animal models for similar purposes with well describe in the literature.

Of particular interest are macrocyclic lactones (i.e. bryostatin class and neristatin class) that act to stimulate PKC. Of the bryostatin class compounds, bryostatin-1 has been shown to activate PKC and proven to be devoid of tumor promotion activity. Bryostatin-1, as a PKC activator, is also particularly useful since the dose response

curve of bryostatin-1 is biphasic. Additionally, bryostatin-1 demonstrates differential regulation of PKC isozymes, including PKC α , PKC δ , and PKC ϵ . Bryostatin-1 has undergone toxicity and safety studies in animals and humans and is actively being investigated as an anti-cancer agent. Bryostatin-1's use in the studies has determined
5 that the main adverse reaction in humans is myalgia, limiting the maximum dose to 40 mg/m². The present invention has utilized concentrations of 0.1 nM of bryostatin-1 to cause a dramatic increase of sAPP secretion. Bryostatin-1 has been compared to a vehicle alone and to another PKC activator, benzolactam (BL), used at a concentration 10,000 times higher. Also bryostatin used at 0.01 nM still proved effective to increase
10 sAPP secretion. (See Figure 1). PKC translocation shows that a measure of activation is maximal at 30 min, followed by a partial decline, which remains higher than basal translocation levels up to six hours. (see Figures 2,3, & 7). The use of the PKC inhibitor staurosporin completely prevents the effect of bryostatin on sAPP secretion. The data further demonstrates that PKC activation mediates the effect of the bryostatin on sAPP
15 secretion. (see Figures 1-3)

Macrocyclic lactones, and particularly bryostatin-1 is described in U.S. Patent 4,560,774. Macrocyclic lactones and their derivatives are described elsewhere in the art for instance in U.S. Patent 6,187,568, U.S. Patent 6,043,270, U.S. Patent 5,393,897, U.S. Patent 5,072,004, U.S. Patent 5,196,447, U.S. Patent 4,833,257, and U.S. Patent
20 4,611,066. The above patents describe various compounds and various uses for macrocyclic lactones including their use as an anti-inflammatory or anti-tumor agent. Other discussions regarding bryostatin class compounds can be found in: Differential Regulation of Protein Kinase C Isozymes by Bryostatin 1 and Phorbol 12-Myristate 13-Acetate in NIH 3T3 Fibroblasts, Szallasi et al., Journal of Biological Chemistry, Vol.
25 269, No. 3, pp. 2118-24 (1994); Preclinical Pharmacology of the Natural Product Anticancer Agent Bryostatin 1, an Activator of Protein Kinase C, Zhang et al., Cancer Research 56, 802-808 (1996); Bryostatin 1, an activator of protein kinase C, inhibits tumor promotion by phorbol esters in SENCAR mouse skin, Hennings et al.,

Carcinogenesis vol. 8, no. 9, pp 1343-46 (1987); Phase II Trial of Bryostatin 1 in Patients with Relapse Low-Grade Non-Hodgkin's Lymphoma and Chronic Lymphocytic Leukemia, Varterasian et al., Clinical Cancer Research, Vol. 6, pp. 825-28 (2000); and Review Article: Chemistry and Clinical Biology of the Bryostatins, Mutter et al., Bioorganic & Medicinal Chemistry 8, pp. 1841-1860 (2000).

5 Macrocyclic lactones, including the bryostatin class, represent known compounds, originally derived from *Bugula neritina L.* While multiple uses for macrocyclic lactones, particularly the bryostatin class are known, the relationship between macrocyclic lactones and cognition enhancement was previously unknown.

10 The examples of the compounds that may be used in the present invention include macrocyclic lactones (i.e. bryostatin class and neristatin class compounds). While specific embodiments of these compounds are described in the examples and detailed description, it should be understood that the compounds disclosed in the references and derivatives thereof could also be used for the present compositions and
15 methods.

As will also be appreciated by one of ordinary skill in the art, macrocyclic lactone compounds and their derivatives, particularly the bryostatin class, are amenable to combinatorial synthetic techniques and thus libraries of the compounds can be generated to optimize pharmacological parameters, including, but not limited to efficacy
20 and safety of the compositions. Additionally, these libraries can be assayed to determine those members that preferably modulate α -secretase and/or PKC.

Combinatorial libraries high throughput screening of natural products and
25 fermentation broths has resulted in the discovery of several new drugs. At present, generation and screening of chemical diversity is being utilized extensively as a major technique for the discovery of lead compounds, and this is certainly a major fundamental advance in the area of drug discovery. Additionally, even after a "lead" compound has been identified, combinatorial techniques provide for a valuable tool for the optimization of desired biological activity. As will be appreciated, the subject

reactions readily lend themselves to the creation of combinatorial libraries of compounds for the screening of pharmaceutical, or other biological or medically-related activity or material-related qualities. A combinatorial library for the purposes of the present invention is a mixture of chemically related compounds, which may be screened 5 together for a desired property; said libraries may be in solution or covalently linked to a solid support. The preparation of many related compounds in a single reaction greatly reduces and simplifies the number of screening processes that need to be carried out. Screening for the appropriate biological property may be done by conventional methods. Thus, the present invention also provides methods for determining the ability 10 of one or more inventive compounds to bind to effectively modulate α -secretase and/or PKC.

A variety of techniques are available in the art for generating combinatorial libraries described below, but it will be understood that the present invention is not intended to be limited by the foregoing examples and descriptions. See, for example, 15 Blondelle et al. (1995) Trends Anal. Chem. 14:83; the Affymax U.S. Patents 5,359,115 and 5,362,899; the Ellman U.S. Patent 5,288,514; the Still et al. PCT publication WO 94/08051; Chen et al. (1994) JACS1 1 6:266 1; Kerr et al. (1993) JACS I 1 5:252; PCT publications W092/10092, W093/09668 and W091/07087; and the Lemer et al. PCT publication W093/20242). Accordingly, a variety of libraries on the order of about 16 to 20 1,000,000 or more diversomers can be synthesized and screened for a particular activity or property.

The present compounds can be administered by a variety of routes and in a variety of dosage forms including those for oral, rectal, parenteral (such as subcutaneous, intramuscular and intravenous), epidural, intrathecal, intra-articular, 25 topical and buccal administration. The dose range for adult human beings will depend on a number of factors including the age, weight and condition of the patient and the administration route.

For oral administration, fine powders or granules containing diluting, dispersing and/or surface-active agents may be presented in a draught, in water or a syrup, in capsules or sachets in the dry state, in a non-aqueous suspension wherein suspending agents may be included, or in a suspension in water or a syrup. Where desirable or 5 necessary, flavouring, preserving, suspending, thickening or emulsifying agents can be included.

Other compounds which may be included by admixture are, for example, medically inert ingredients, e.g. solid and liquid diluent, such as lactose, dextrose, saccharose, cellulose, starch or calcium phosphate for tablets or capsules, olive oil or 10 ethyl oleate for soft capsules and water or vegetable oil for suspensions or emulsions; lubricating agents such as silica, talc, stearic acid, magnesium or calcium stearate and/or polyethylene glycols; gelling agents such as colloidal clays; thickening agents such as gum tragacanth or sodium alginate, binding agents such as starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinylpyrrolidone; 15 disintegrating agents such as starch, alginic acid, alginates or sodium starch glycolate; effervesing mixtures; dyestuff; sweeteners; wetting agents such as lecithin, polysorbates or laurylsulphates; and other therapeutically acceptable accessory ingredients, such as humectants, preservatives, buffers and antioxidants, which are known additives for such formulations.

20 Liquid dispersions for oral administration may be syrups, emulsions or suspensions. The syrups may contain as carrier, for example, saccharose or saccharose with glycerol and/or mannitol and/or sorbitol. In particular a syrup for diabetic patients can contain as carriers only products, for example sorbitol, which do not metabolize to glucose or which metabolize only a very small amount to glucose. The suspensions and 25 the emulsions may contain a carrier, for example a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose or polyvinyl alcohol.

Suspensions or solutions for intramuscular injection may contain, together with the active compound, a pharmaceutically acceptable carrier such as sterile water, olive

oil, ethyl oleate, glycols such as propylene glycol and, if desired, a suitable amount of lidocaine hydrochloride. Solutions for intravenous injection or infusion may contain a carrier, for example, sterile water that is generally Water for Injection. Preferably, however, they may take the form of a sterile, aqueous, isotonic saline solution.

5 Alternatively, the present compounds may be encapsulated within liposomes. The present compounds may also utilize other known active agent delivery systems.

The present compounds may also be administered in pure form unassociated with other additives, in which case a capsule, sachet or tablet is the preferred dosage form.

10 Tablets and other forms of presentation provided in discrete units conveniently contain a daily dose, or an appropriate fraction thereof, of one of the present compounds. For example, units may contain from 5 mg to 500 mg, but more usually from 10 mg to 250 mg, of one of the present compounds.

15 It will be appreciated that the pharmacological activity of the compositions of the invention can be demonstrated using standard pharmacological models that are known in the art. Furthermore, it will be appreciated that the inventive compositions can be incorporated or encapsulated in a suitable polymer matrix or membrane for site-specific delivery, or can be functionalized with specific targeting agents capable of effecting site specific delivery. These techniques, as well as other drug delivery 20 techniques are well known in the art.

All books, articles, or patents references herein are incorporated by reference to the extent not inconsistent with the present disclosure. The present invention will now be described by way of examples, which are meant to illustrate, but not limit, the scope of the invention.

Examples

Example I:

Materials And Methods

Cell Culture

5 Cultured skin fibroblasts were obtained from the Coriell Cell Repositories and grown using the general guidelines established for their culture with slight modifications (Cristofalo & Carpentier, 1988; Hirashima et al., 1996). The culture medium in which cells were grown was Dulbecco's modified Eagle's medium (GIBCO) supplemented with 10% fetal calf serum (Biofluids, Inc.). Fibroblasts from control cell
10 lines (AC), cases AG07141 and AG06241, and a familial AD (FAD) case (AG06848) were utilized.

PKC Activators

15 The different tissue distributions, the apparently distinctive roles of different isozymes, and the differential involvement in pathology make it important to use pharmacological tools that are capable of preferentially targeting specific isozymes (Kozikowski et al., 1997; Hofmann, 1997). Recent research in the medicinal chemistry field has resulted in the development of several PKC activators, for instance different benzolactams and pyrrolidinones. However, the currently studied bryostatin PKC activator not only has the benefit of providing isospecific activity, but also does not
20 suffer from the set back of the previously used PKC activator, such as being tumor promoting. The bryostatin competes for the regulatory domain of PKC and engages in very specific hydrogen bond interactions within this site. Additional information on the organic chemistry and molecular modeling of this compound can be found throughout the literature.

25 **Treatment**

 Cells grown to confluence in 6 cm Petri dishes for 5-7 days. On the day of the experiment, medium was replaced with DMEM without serum and left undisturbed for 2h. Upon completion of the 2 hour serum deprivation, treatment was achieved by direct

application to the medium of Bryo, BL and DMSO at the appropriate concentrations. DMSO was less than 1% in all cases. In most cases, medium was collected and processed after 3 hours of treatment for sAPP secretion. Other time points were also used to establish a time course of secretion.

5 Immunoblot Assay

Immunoblot experiments were conducted using well-established procedures (Dunbar, 1994). Cells were grown to confluence (~90%) in 6 cm Petri dishes. Levels of isozyme in response to treatment with 0.1 nM bryostatin-1 for 5, 30, 60, and 120 minutes was quantified using procedures slightly modified from that established by 10 Racchi et al., (1994). Fibroblasts were washed twice with ice-cold PBS, scraped in PBS, and collected by low-speed centrifugation. The pellets were re-suspended in the following homogenization buffer: 20 mM Tris-HCl, pH 7.5, 2 mM EDTA, 2mM EGTA, 5 mM DTT, 0.32 M sucrose, and protease inhibitor cocktail (Sigma). Homogenates were obtained by sonication, and centrifuged at ~12,000g for 20 minutes, 15 and the supernatants were used as the cytosolic fraction. The pellets were homogenized in the same buffer containing 1.0% Triton X-100, incubated in ice for 45 minutes, and centrifuged at ~12,000g for 20 minutes. The supernatant from this batch was used as the membranous fraction. After protein determination, 20 µg of protein were diluted in 2 X electrophoresis sample buffer (Novex), boiled for 5 minutes, run on 10% 20 acrylamide gel, and transferred electrophoretically to a PVDF membrane. The membrane was saturated with 5% milk blocker by incubating it at room temperature for an hour. The primary antibody for PKC isoform (Transduction Laboratories) was diluted (1:1000) in blocking solution and incubated with the membrane overnight at 4°C. After incubation with the secondary antibody, alkaline phosphatase antimouse IgG 25 (Vector Laboratories), the membrane was developed using a chemoluminescent substrate (Vector Laboratories) per the manufacturer's instructions. The band intensities were quantified by densitometry using a BioRad GS-800 calibrated scanning densitometer and Multianalyst software (BioRad).

sAPP - Determinations.

The concentration of secreted APP was measured using conventional immunoblotting techniques, with minor modifications the protocol. Precipitated protein extracts from each dish/treatment were loaded to freshly prepared 10% acrylamide 5 Tris-HCl minigels and separated by SDPAGE. The volume of sample loaded was corrected for total cell protein per dish. Proteins were then electrophoretically transferred to PVDF membranes. Membranes were saturated with 5% non-fat dry milk to block non-specific binding. Blocked membranes were incubated overnight at 4 °C with the commercially available antibody 6E10 (1:500), which recognizes sAPP-alpha 10 in the conditioned medium (SENETEK). After washing, the membranes were incubated at room temperature with horseradish peroxidase conjugated anti-mouse IgG secondary antibody (Jackson's Laboratories). The signal was then detected using enhanced chemiluminescence followed by exposure of Hyperfilm ECL (Amersham). The band intensities were quantified by densitometry using a BioRad GS-800 calibrated 15 scanning densitometer and Multianalyst software (BioRad).

As shown in Fig. 7, Bryostatin-1 elicits a powerful response, demonstrating the activation of PKC, It should be noted the activation of PKC is easily detectable 30 minutes after delivery, following a dose of only 0.1 nM of bryostatin-1.

It is also interesting to consider the data in relation to APP metabolism and the 20 effects of its sub-products. Studies have demonstrated that PKC activation increases the amount of ratio of non-amyloidogenic (soluble APP, presumably product of the secretase) vs. amyloidogenic (A β 1-40 and/or A β 1-42) secreted fragments (Buxbaum et al., 1990; Gillespie et al., 1992; Selkoe, 1994). Without wishing to be held to this theory, one could speculate that AD cells with low PKC would have an impaired 25 secretion of sAPP and/or have increased proportion of amyloidogenic fragments. Indeed, there is evidence that some AD cell lines exhibit both defective PKC and impaired sAPP secretion (Bergamaschi et al., 1995; Govoni et al., 1996). In addition, β -amyloid has been shown to induce an AD-like K $^{+}$ channel defect in fibroblasts

(Etcheberrigaray et al., 1994) and to block K⁺ currents in cultured neurons (Good et al., 1996). Therefore, we suggest a mechanistic link such that an isozyme-specific PKC defect may lead to abnormal APP processing that, among other possible deleterious effects, alters K⁺ channel function. Recent preliminary data also suggest that, perhaps 5 in a vicious cyclical manner, β -amyloid in turn causes reductions of PKC (Favit et al., 1997).

In summary, the data suggest that the strategy to up-regulate PKC function targeting specific isozymes increases sAPP production. These studies and such a fibroblast model could be expanded and used as tools to monitor the effect of 10 compounds (bryostatin, for example) that alter potential underlying pathological processes. Further, one of ordinary skill in the art would know how to further test these samples through Ca²⁺ imaging and electrophysiology. Such compounds could then be used as bases for rational design of pharmacological agents for this disorder.

Example II

15 The effect of PKC activators on cognition was demonstrated by the Morris Water Maze paradigm. In the present example, rats were injected intraventricularly with bryostatin-1 and trained for 4 days (following standard protocols). Retention was assessed on the 5th day. Learning was measured as the reduction of escape latency from trial to trial, which was significantly lower in the treated animals. Acquisition of 20 memory was measured as time spent in the relevant quadrant (5th day). Memory or retention was significantly enhanced in treated animals, compared to sham injection animals (see Figures 4 through 5(a)-5(c)). The rats treated with bryostatin-1 showed improved cognition over control rats within 2 days of treatment (see Figure 4). Bryostatin-1 is capable of being used at concentrations to improve cognition that are 25 300 to 300,000 times lower than the concentration used to treat tumors. The above example further shows that cognitive ability can be improved in non-diseased subjects as compared to other non-diseased subjects through the administration of bryostatin-1.

Because of the previously conducted safety, toxicology and phase II clinical studies for cancer, one can conclude that the use of PKC activators, particularly bryostatin-1, would be viewed as safe and that phase II studies for AD treatment/cognitive enhancement could be expedited. Furthermore, bryostatin-1's 5 lipophilic nature provides increased blood brain barrier transport. The present invention would allow for intravenous, oral, intraventricular, and other known methods of administration.

Test of sAPP secretion experiments, PKC activation experiments, and animal behavior experiments have shown that increases in sAPP secretion follow increased 10 PKC activation and result in improved cognition in animal behavior studies.

WHAT IS CLAIMED IS:

1. A method for enhancing cognitive ability in a human or animal, comprising administering to said human or animal a PKC activator in an amount effective for enhancing cognitive ability in a pharmaceutically acceptable carrier.
- 5 2. The method of claim 1 wherein the PKC activator selectively activates PKC α , PKC δ , and PKC ϵ .
3. The method of claim 1, wherein the PKC activator is a macrocyclic lactone, a benzolactam, a pyrrolidinone or a combination thereof.
4. The method of claim 3, wherein the macrocyclic lactone is a bryostatin class or 10 neristatin class compound.
5. The method of claim 3 wherein the PKC activator is bryostatin-1 through bryostatin 18 or neristatin-1.
6. The method of claim 1, wherein the cognitive ability enhanced is learning, memory, or attention.
- 15 7. The method of claim 5, wherein the animal is a primate.
8. The method of claim 5, wherein the animal is a non-primate.
9. The method of claim 1, wherein the amount of PKC activator administered is in an amount effective to treat cognitive impairment of a neurological disease or disorder.
10. The method of claim 9, wherein the neurological disease is Alzheimer's Disease, 20 multi-infarct dementia, the Lewy-body variant of Alzheimer's Disease with or without association with Parkinson's disease; Creutzfeld-Jakob disease, Korsakow's disorder, or attention deficit hyperactivity disorder.
11. The method of claim 9, wherein the disorder is -associated with age, electro-convulsive therapy or brain damage.
- 25 12. The method of claim 11, wherein the brain damage was caused by stroke, an anesthetic accident, head trauma, hypoglycemia, carbon monoxide poisoning, lithium intoxication or a vitamin deficiency.

13. The method of claim 1, wherein the PKC activator is administered in an amount effected to cause an increase in sAPP.
14. A method for altering cellular modulation of ion channels comprising administering a PKC activator in an amount effective for altering cellular modulation of ion channels and a pharmaceutically acceptable carrier.
5
15. The method of claim 14 wherein, said modulation is *in vivo* or *in vitro* modulation.
16. The method of claim 15, wherein said ion channel is a K⁺ or Ca⁺⁺ channel.
17. A method for treating neurotumors comprising administering macrocyclic lactone in an amount effective to treat said neurotumors and a pharmaceutically acceptable carrier.
10
18. The method of claim 17 wherein, the macrocyclic lactone is a bryostatin class or neristatin class compound.
19. A method for modulating sAPP comprising administering a macrocyclic lactone in an amount effective to modulate sAPP and a pharmaceutically acceptable carrier.
15
20. The method of claim 19 wherein, the macrocyclic lactone activates PKC.
21. The method of claim 19 wherein, the macrocyclic lactone is a bryostatin class or a neristatin class compound.
22. The method of claim 19 wherein, the macrocyclic lactone is bryostatin-1 through bryostatin 18, or neristatin-1.
20
23. A method for modulating α -secretase comprising administering a macrocyclic lactone in an amount effective to modulate α -secretase and a pharmaceutically suitable carrier.
24. The method of claim 23 wherein, the macrocyclic lactone is a bryostatin class or a neristatin class compound.
25
25. The method of claim 23 wherein, the macrocyclic lactone is bryostatin-1 through bryostatin 18, or neristatin-1.
26. The method of claim 23 wherein, the macrocyclic lactone is administered *in vivo* or *in vitro*.

27. The method of claim 23 wherein, the modulation of α -secretase reduces amyloid plaque formation and enhances cognitive ability in a patient with Alzheimer's Disease.
28. A method for treating Alzheimer's Disease comprising administering to a patient a macrocyclic lactone in an amount effective to treat Alzheimer's Disease and a pharmaceutically acceptable carrier.
5
29. A method for treating Alzheimer's Disease comprising administering bryostatin-1 to a patient in an amount effective to treat Alzheimer's Disease and a pharmaceutically acceptable carrier.
- 10 30. A method for enhancing cognitive ability in a human or animal, comprising administering to said human or animal a bryostatin or a neristatin class compound in an amount effective for enhancing cognitive ability in a pharmaceutically acceptable carrier.
- 15 31. A method for enhancing cognitive ability in a human or animal, comprising administering to said human or animal bryostatin-1 in an amount effective for enhancing cognitive ability in a pharmaceutically acceptable carrier.
32. A method comprising the modulation of α -secretase through the administration of a pharmaceutically effective amount of a bryostatin or a neristatin class compound and a pharmaceutically acceptable carrier.
- 20 33. A method for modulating α -secretase comprising administering bryostatin-1 in an amount effective to modulate α -secretase and a pharmaceutically acceptable carrier.
34. A method for providing a neuroprotective effect for cell comprising administering a bryostatin or neristatin class compound in an amount effective to provided a neuroprotective effect for cells which suffer from a hypoxic event and a pharmaceutically acceptable carrier.
25
35. A method for providing a neuroprotective effect for cell comprising administering bryostatin-1 in an amount effective to provided a neuroprotective effect for cells which suffer from a hypoxic event and a pharmaceutically acceptable carrier.

36. A method for the reduction of amyloid plaque formation comprising administering bryostatin-1 in an amount effective to reduce amyloid plaque formation and a pharmaceutically acceptable carrier.

5




FIG. 1




FIG. 2





FIG. 3



BRYOSTATIN-1 (i.c.v.; 1 μ l/SITE OF 2 μ M SOLUTION; ~0.5 HR PRIOR TO THE 1ST AND 5TH TRAINING TRIALS);
10 RATS/GROUP.

FIG. 4




FIG. 5A




FIG. 5B




FIG. 5C




FIG 7

