



(86) Date de dépôt PCT/PCT Filing Date: 2001/10/02  
(87) Date publication PCT/PCT Publication Date: 2002/04/11  
(85) Entrée phase nationale/National Entry: 2003/04/01  
(86) N° demande PCT/PCT Application No.: US 2001/042461  
(87) N° publication PCT/PCT Publication No.: 2002/028185  
(30) Priorités/Priorities: 2000/10/06 (60/238,406) US;  
2001/10/01 (09/968,174) US

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> A01N 51/00, A01N 63/00

(71) Demandeur/Applicant:  
MONSANTO TECHNOLOGY LLC, US

(72) Inventeurs/Inventors:  
ASRAR, JAWED, US;  
KOHN, FRANK C., US;  
SANDERS, ERNEST F., US

(74) Agent: MCFADDEN, FINCHAM

(54) Titre : PROCEDE PERMETTANT DE REDUIRE LES PRELEVEMENTS ALIMENTAIRES DE DEPREDATEURS SUR  
LE MAIS PAR TRAITEMENT DES SEMENCES DE MAIS TRANSGENIQUES AU MOYEN DE PESTICIDE  
THIAMETHOXAM

(54) Title: METHOD FOR REDUCING PEST DAMAGE TO CORN BY TREATING TRANSGENIC CORN SEEDS WITH  
THIAMETHOXAM PESTICIDE

(57) **Abrégé/Abstract:**

A method to protect corn against feeding damage by one or more pests includes the treatment of corn seed having a transgenic event that is targeted against at least one of the pests with thiamethoxam in an amount that is effective against the same or another of the one or more pests. Seeds having such protection are also disclosed.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number  
**WO 02/28185 A3**

(51) International Patent Classification<sup>7</sup>: **A01N 51/00 //**  
(A01N 51/00, 63:00)

(21) International Application Number: PCT/US01/42461

(22) International Filing Date: 2 October 2001 (02.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/238,406 6 October 2000 (06.10.2000) US  
09/968,174 1 October 2001 (01.10.2001) US

(71) Applicant (for all designated States except US): **MON-SANTO TECHNOLOGY, LLC** [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ASRAR, Jawed** [US/US]; 14949 Royal Brook, Chesterfield, MO 63017 (US). **KOHN, Frank, C.** [US/US]; 12532 Larkwood Drive, St. Louis, MO 63146 (US). **SANDERS, Ernest, F.** [US/US]; 16 Regatta Bay Ct., Lake St. Louis, MO 63367 (US).

(74) Agent: **NELSON MULLINS RILEY & SCARBOROUGH**; Keenan Building, Third Floor, 1330 Lady Street, Columbia, SC 29201 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
20 June 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR REDUCING PEST DAMAGE TO CORN BY TREATING TRANSGENIC CORN SEEDS WITH THI-AMETHOXAM PESTICIDE

(57) Abstract: A method to protect corn against feeding damage by one or more pests includes the treatment of corn seed having a transgenic event that is targeted against at least one of the pests with thiamethoxam in an amount that is effective against the same or another of the one or more pests. Seeds having such protection are also disclosed.



**WO 02/28185 A3**

**METHOD FOR REDUCING PEST DAMAGE TO CORN  
BY TREATING TRANSGENIC CORN SEEDS WITH  
THIAMETHOXAM PESTICIDE**

CROSS-REFERENCES TO RELATED APPLICATIONS

5           This application is a non-provisional of U.S. Patent Application  
Serial No. 60/238,406, filed on October 6, 2000, and claims priority  
thereto.

BACKGROUND OF THE INVENTION

(1)   Field of the Invention:

10           The present invention relates generally to the control of pests  
that cause damage to corn plants by their feeding activities, and more  
particularly to the control of such corn plant pests by the combination of  
a corn seed having a transgenic event and the treatment of such seed  
with a thiamethoxam pesticide prior to planting the seed.

15   (2)   Description of the Related Art:

Insects and related arthropods annually destroy an estimated  
15% of agricultural crops in the United States and even more than that  
in developing countries. In addition, competing weeds and parasitic  
and saprophytic plants account for even more potential yield losses.

20           Some of this damage occurs in the soil when plant pathogens,  
insects and other such soil borne pests attack the seed after planting.  
In the production of corn, much of the rest of the damage is caused by  
rootworms -- insect pests that feed upon or otherwise damage the plant  
roots; and by cutworms, European corn borers, and other pests that  
25   feed upon or damage the above ground parts of the plant. General  
descriptions of the type and mechanisms of attack of pests on  
agricultural crops are provided by, for example, Metcalf, in *Destructive  
and Useful Insects*, (1962); and Agrios, in *Plant Pathology*, 3rd Ed.,  
Academic Press (1988).

30           Corn is the most important grain crop in the Midwestern United  
States. Among the most serious insect pests of corn in this region are  
the larval forms of three species of *Diabrotica* beetles. These include



the Western corn rootworm, *Diabrotica vergifera vergifera* LeConte, the Northern corn rootworm, *Diabrotica berberi* Smith and *Diabrotica berberi* Lawrence, and the Southern corn rootworm, *Diabrotica undecimpunctata howardi* Barber. In fact, more insecticide is used for the control of corn rootworm than for any other pest of corn, and the total acreage treated is greater than for any other pest in the United States.

Corn rootworms (CRW) overwinter in the egg stage in fields where corn was grown the previous season. The eggs hatch from late May through June. If a corn crop is not followed by another corn crop in the subsequent year, the larvae will die. Accordingly, the impact of corn rootworm is felt most directly in areas where corn is systematically followed by corn, as is typical in many areas of the Midwestern United States.

After hatching, the larvae pass through three larval stages or instars, during which they feed on the corn root system. About three weeks is required for completion of the larval stage. Damage to the corn root system caused by the feeding of larvae is the major cause of harvest losses in corn due to corn rootworm. Corn plants that fall over and lodge in the field after weakening or destruction of a major part of the root system are the cause of a major portion of this loss, since this lodged corn cannot be harvested by conventional mechanized machinery and is left in the field.

Following completion of larval development, the larvae transform into immobile pupae, and thence into the adult beetles that emerge from the soil throughout the summer, with the period of emergence depending upon the growing location. After emergence, the adult beetles feed for about two weeks before the females start laying eggs. Initially, the adults feed predominantly in the same field from which they emerged, but later will migrate to other fields. Peak adult activity normally occurs in the U.S. Corn Belt during late July or early August in fields planted to continuous corn, but activity may peak later in first

year or late maturing cornfields. Rootworm beetles begin depositing eggs in cornfields approximately two weeks after they emerge. (For more information, see, e.g., *Corn Rootworms*, Field Crops Pest Management Circular #16, Ohio Pest Management & Survey Program, The Ohio State University, Extension Division, Columbus, OH; available online at [www.ag.ohio-state.edu/~ohioline/icm-fact/fc-16.html](http://www.ag.ohio-state.edu/~ohioline/icm-fact/fc-16.html), September 13, 2000; and McGahen *et al.*, *Corn Insect Control: Corn Rootworm*, PENpages number 08801502, Factsheet available from Pennsylvania State University, State College, PA, 1989).

In present conventional agricultural practice, in cases where corn follows corn, it is normal for an insecticide to be applied to protect the corn root system from severe feeding by rootworm larvae. Conventional practice is to treat for the adult beetles or to treat for the larvae. Examples of conventional treatment formulations for adult beetles include the application of carbaryl insecticides (e.g., SEVIN® 80S at 1.0 - 2.0 lbs active/acre); fenvalerate or esfenvalerate (e.g., PYDRIN® 2.4EC at 0.1 to 0.2 lbs active/acre, or ASANA® 0.66EC at 0.03 to 0.05 lbs active/acre); malathion (57% E at 0.9 lbs active/acre); permethrin (e.g., AMBUSH® 2.0EC at 0.1 to 0.2 lbs active/acre, or POUNCE® 3.2EC at 0.1 to 0.2 lbs active ingredient/acre); or PENNCAP-M® at 0.25 - 0.5 lbs active/acre.

To treat for CRW larvae, conventional practice is to apply a soil insecticide either at or after planting, but preferably as close to egg hatching as possible. Conventional treatments include carbofuran insecticides (e.g., FURADAN® 15G at 8 oz/ 1000 ft of row); chloropyrifos (e.g., LORSBAN® 15G at 8 oz/1000 ft of row); fonophos (e.g., DYFONATE® 20G at 4.5 to 6.0 oz/1000 ft of row); phorate (e.g., THIMET® 20G at 6 oz/1000 ft of row); terbufos (e.g., COUNTER® 15G at 8 oz/1000 ft of row); or tefluthrin (e.g., FORCE® 3G at 4 to 5 oz/1000 ft of row).

Many of the chemical pesticides listed above are known to be harmful to humans and to animals in general. The environmental harm

that these pesticides cause is often exacerbated due to the practice of applying the pesticides by foliar spraying or direct application to the surface of the soil. Wind-drift, leaching, and runoff can cause the migration of a large fraction of the pesticide out of the desired zone of activity and into surface waters and direct contact with birds, animals and humans.

Because of concern about the impact of chemical pesticides on public health and the health of the environment, significant efforts have been made to find ways to reduce the amount of chemical pesticides that are used. Recently, much of this effort has focused on the development of transgenic crops that are engineered to express insect toxicants derived from microorganisms. For example, U.S. Patent No. 5,877,012 to Estruch *et al.* discloses the cloning and expression of proteins from such organisms as *Bacillus*, *Pseudomonas*, *Clavibacter* and *Rhizobium* into plants to obtain transgenic plants with resistance to such pests as black cutworms, armyworms, several borers and other insect pests. Publication WO/EP97/07089 by Privalle *et al.* teaches the transformation of monocotyledons, such as corn, with a recombinant DNA sequence encoding peroxidase for the protection of the plant from feeding by corn borers, earworms and cutworms. Jansens *et al.*, in *Crop Sci.*, 37(5):1616 - 1624 (1997), reported the production of transgenic corn containing a gene encoding a crystalline protein from *Bacillus thuringiensis* (Bt) that controlled both generations of the European corn borer. U. S. Patent Nos. 5,625,136 and 5,859,336 to Koziel *et al.* reported that the transformation of corn with a gene from *B. thuringiensis* that encoded for delta-endotoxins provided the transgenic corn with improved resistance to European corn borer. A comprehensive report of field trials of transgenic corn that expresses an insecticidal protein from *B. thuringiensis* has been provided by Armstrong *et al.*, in *Crop Science*, 35(2):550 - 557 (1995).

It was known that wild-type Bt  $\delta$ -endotoxins had low activity against coleopteran insects, and Kreig *et al.*, in 1983, reported the first



isolation of a coleopteran-toxic *B. thuringiensis* strain. (See U.S. Patent No. 4,766,203). U.S. Patent Nos. 4,797,279 and 4,910,016, also disclosed wild-type and hybrid *B. thuringiensis* strains that produced proteins having some coleopteran activity. More recently,  
5 however, more precise genetic engineering methods have shown promise in developing modified *B. thuringiensis* proteins that have significantly higher levels of corn rootworm activity than those produced by wild-type parents. (See, *e.g.*, WO 99/31248 to Ecogen, Inc. and Monsanto Company).

10 However, it is not known at present whether any transgenic event alone will be sufficiently effective to protect corn from damage by corn rootworm in heavily infested fields that are dedicated to serial corn. In fact, the total control of corn rootworm damage by any one transgenic event may not be desirable in the long term, because of the  
15 potential for the development of resistant strains of the target pest.

Another alternative to the conventional forms of pesticide application is the treatment of plant seeds with pesticides. The use of fungicides to protect seeds from attack after planting, and the use of low levels of insecticides for the protection of, for example, corn seed  
20 from wireworm, has been used for some time. Seed treatment with pesticides has the advantages providing for the protection of the seeds, while minimizing the amount of pesticide that was required and limiting the amount of contact with the pesticide and the number of different field applications that were necessary.

25 Other examples of the control of pests by applying insecticides directly to plant seed are provided in, for example, U.S. Patent No. 5,696,144, which discloses that the European corn borer caused less feeding damage to corn plants grown from seed treated with a 1-arylpyrazole compound at a rate of 500 g per quintal of seed than  
30 control plants grown from untreated seed. In addition, U.S. Patent No. 5,876,739 to Turnblad *et al.* (and its parent, U. S. Patent No. 5,849,320) disclose a method for controlling soil-borne insects which

involves treating seeds with a coating containing one or more polymeric binders and an insecticide. This reference provides a list of insecticides that it identifies as candidates for use in this coating and also names a number of potential target insects. However, while the 5,876,739 patent states that treating corn seed with a coating containing a particular insecticide protects corn roots from damage by the corn rootworm, it does not indicate or otherwise suggest that such treatment could be used with seed having a transgenic event.

The treatment of seed having a transgenic event with nitroimino- or nitroguanidino-compound pesticides has been mentioned (See, e.g., WO 99/35913), however, no guidance has been found as to the potential utility or efficacy of such treatments, or the details of how such treatments might be effected -- such as the amounts of active ingredient that would be necessary per unit amount of seed -- and no examples that would give reason to believe that the proposed treatments would actually provide suitable protection.

Therefore, although recent developments in genetic engineering of plants have improved the ability to protect plants from pests without using chemical pesticides, and while such techniques as the treatment of seeds with pesticides have reducing the harmful effects of pesticides on the environment, numerous problems remain that limit the successful application of these methods under actual field conditions. Accordingly, it would be useful to provide an improved method for the protection of plants, especially corn plants, from feeding damage by pests. It would be particularly useful if such method would reduce the required application rate of conventional chemical pesticides, and also if it would limit the number of separate field operations that were required for crop planting and cultivation.



## BRIEF SUMMARY OF THE INVENTION

Briefly, therefore, the present invention is directed to a novel method for protecting a transgenic corn plant against feeding damage by one or more pests, the method comprising providing a seed for the transgenic corn plant which seed comprises a transgenic event having activity against at least one of the one or more pests; and treating the seed with an effective amount of thiamethoxam pesticide.

The present invention is also directed to a novel seed of a transgenic corn plant that provides increased resistance to the resulting corn plant against feeding damage by one or more pests, comprising a transgenic event having activity against at least one of the one or more pests, which seed has been treated with an effective amount of thiamethoxam pesticide.

The present invention is also directed to a novel transgenic corn seed that has been treated by the method of the present invention.

Among the several advantages found to be achieved by the present invention, therefore, may be noted the provision of an improved method for the protection of plants, especially corn plants, from feeding damage by pests; the provision of such a method that reduces the required application rate of conventional chemical pesticides; and also the provision of such a method that limits the number of separate field operations that were required for crop planting and cultivation.

## DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, it has been discovered that corn plants can be protected against feeding damage by one or more pests by a method that includes providing a corn seed having a transgenic event that has activity against at least one of the pests and then treating the transgenic corn seed with an effective amount of thiamethoxam pesticide. For example, in preferred embodiments it has been found that the combination of a transgenic event having activity against corn rootworm and treatment of the seed with thiamethoxam

provides unexpectedly synergistic advantages to seeds having such treatment, including unexpectedly superior efficacy for protection against damage to the resulting corn plant by corn rootworm. In particular, it is believed that corn seeds having a transgenic event that causes the production of CryBb.11231 endotoxin in combination with the treatment of such seeds with thiamethoxam would be unexpectedly superior to either the transgenic event alone, or to treatment with thiamethoxam alone, in protecting resulting corn plants against more severe levels of damage by corn rootworm -- levels of damage that are known to reduce corn yield.

Corn plants and seeds that have been engineered to include exogenous genes derived from *Bacillus thuringiensis* that encode for the expression of Cry3  $\delta$ -endotoxins having activity against Coleopteran pests are known, as are methods for the treatment of seeds (even some transgenic seeds) with pesticides. However, it had not been realized until the present invention that certain effective amounts of thiamethoxam could be used to treat corn seeds having such Cry3 events, with the result that the combination would be unexpectedly superior in increasing the efficacy of both the pesticide and the transgenic event, and would provide the additional advantages of increasing the ability to match pesticidal activity against pest pressure, decreasing cost of treatment and/or application, increasing safety of seed handling, and decreasing environmental impact of either or both the event and the pesticide.

In particular, it is believed that in preferred embodiments the treatment of transgenic corn seeds that are capable of expressing certain modified Cry3Bb proteins with from about 100 gm to about 400 gm of thiamethoxam per 100 kg of seed can provide unexpectedly superior protection against corn rootworm. In addition, it is believed that such combinations are also effective to protect the emergent corn plants against damage by black cutworm. The seeds of the present invention are also believed to have the property of decreasing the cost

of pesticide use, because less of the pesticide can be used to obtain a required amount of protection than if the innovative method is not used. Moreover, because less pesticide is used and because it is applied prior to planting and without a separate field application, it is believed  
5 that the subject method is therefore safer to the operator and to the environment, and is potentially less expensive than conventional methods.

When it is said that some effects are "synergistic", it is meant to include the synergistic effects of the combination on the pesticidal  
10 activity (or efficacy) of the combination of the transgenic event and the pesticide. However, it is not intended that such synergistic effects be limited to the pesticidal activity, but that they should also include such unexpected advantages as increased scope of activity, advantageous activity profile as related to type and amount of damage reduction,  
15 decreased cost of pesticide and application, decreased pesticide distribution in the environment, decreased pesticide exposure of personnel who produce, handle and plant corn seeds, and other advantages known to those skilled in the art.

The present invention also provides an advantage of increasing  
20 the ability to match pesticidal activity against pest pressure. This refers to the ability to design the combination of the transgenic event and the pesticide treatment so that the seed or the resulting plant is provided with effective pesticidal activity during the period when feeding pressure from the target pest on the seed or plant reaches its  
25 maximum. By way of example, when a pesticide such as thiamethoxam is applied to a corn seed having a corn rootworm transgenic event, the pesticide can be applied in a coating designed to provide controlled release of the pesticide. The release rate can be selected so that the pesticide provides protection against such other  
30 pests as are controlled by thiamethoxam at the post emergence stage of corn, while the transgenic event provides corn rootworm protection



at a later stage of plant development -- when such protection is needed.

As used herein, the terms "pesticidal effect" and "pesticidal activity", or "activity" refer to a toxic effect against a pest. The terms  
5 "activity against (one or more) pests", also have the same meaning. When it is said that a seed or plant is "protected against feeding damage by one or more pests", it is meant that such seed or plant possesses a feature having direct or indirect action on one or more pests that results in reduced feeding damage by such pest or pests on  
10 the seeds, roots, shoots and foliage of plants having such feature as compared to the feeding damage caused under the same conditions to plants not having such feature. Such direct or indirect actions include inducing death of the pest, repelling the pest from the plant seeds, roots, shoots and/or foliage, inhibiting feeding of the pest on, or the  
15 laying of its eggs on, the plant seeds, roots, shoots and/or foliage, and inhibiting or preventing reproduction of the pest.

The term "insecticidal activity" has the same meaning as pesticidal activity, except it is limited to those instances where the pest is an insect. Except where specifically noted, when the term "pesticide"  
20 is used herein, that term refers to a chemical pesticide that is supplied externally to the seed, and it is not meant to include active agents that are produced by the particular seed or the plant that grows from the particular seed. However, the terms "pesticidal activity" and "insecticidal activity" can be used with reference to the activity of either,  
25 or both, an externally supplied pesticide and/or an agent that is produced by the seed or the plant.

One feature of the present invention is a seed of a transgenic corn plant. As used herein, the terms "transgenic corn plant" mean a corn plant or progeny thereof derived from a transformed corn plant cell  
30 or protoplast, wherein the plant DNA contains an introduced exogenous DNA molecule not originally present in a native, non-transgenic plant of the same strain.

The transgenic corn seed is one that contains an exogenous gene that encodes a pesticidal protein. Pesticidal proteins of this type are described by Schnepf *et al.*, in *Microbiology & Molecular Biology Reviews*, 62:775 - 806 (1998), and by ffrench-Constant and Bowen, in  
5 *CMSL Cell. Mol. Life Sci.*, 57:828 - 833 (2000). In one application of the invention, the pesticidal protein is an insecticidal protein.

It is preferred that the seed contains an exogenous gene derived from a strain of *Bacillus thuringiensis*, and in particular, it is preferred that the exogenous gene is one that encodes an insecticidal  $\delta$ -  
10 endotoxin derived from *B. thuringiensis*. Such  $\delta$ -endotoxins are described in WO 99/31248 and U.S. Patent No. 6,063,597, and include the Cry3 toxins. Nucleic acid segments that encode modified *B. thuringiensis* coleopteran-toxic crystal proteins that are useful in the present invention are described in U.S. Patent No. 6,060,594, and  
15 insect resistant transgenic plants that include nucleic acid sequences that encode such insecticidal proteins are discussed in U.S. Patent No. 6,023,013. It is preferred that the  $\delta$ -endotoxins of the present invention include the Cry3B proteins, and even more preferred that the  $\delta$ -endotoxins include the coleopteran-active Cry3Bb proteins. The  
20 nomenclature of the *B. thuringiensis* insecticidal crystal proteins was set forth by Höfte and Whitely, *Microbiol. Rev.*, 53:242-255, 1989. This nomenclature has been revised, and the revised nomenclature can be found at <http://epunix.biols.susx.ac.uk/Home/Neil-Crickmore/Bt/index.html>. The revised nomenclature will be used herein  
25 to describe transgenic event features and the  $\delta$ -endotoxin proteins encoded by the transgenic event.

When the terms "transgenic event" are used herein, such terms are meant to refer to the genetically engineered DNA that is described above, but also to include the protein(s) that are encoded by the  
30 modified gene. A transgenic event in a corn seed, or corn plant, therefore, includes the ability to express a protein. When it is said that a "transgenic event has activity against a pest", it is to be understood

that it is the protein that is encoded by the gene that actually has such activity when the protein is expressed and brought into contact with the pest.

WO 99/31248 and U.S. Patent Nos. 6,063,597 describe  
5 methods for genetically engineering *B. thuringiensis*  $\delta$ -endotoxin genes so that modified  $\delta$ -endotoxins can be expressed. The modified  $\delta$ -endotoxins differ from the wild-type proteins by having specific amino acid substitutions, additions or deletions as compared with the proteins produced by the wild-type organism. Such modified  $\delta$ -endotoxins are  
10 identified herein by the use of an asterisk (\*), or by reference to a specific protein by its identifying number. Thus, a genetically modified Cry3  $\delta$ -endotoxin would be expressed as Cry3\*, examples of which include, without limitation: Cry3Bb.11230, Cry3Bb.11231, Cry3Bb.11232, Cry3Bb.11233, Cry3Bb.11234, Cry3Bb.11235,  
15 Cry3Bb.11236, Cry3Bb.11237, Cry3Bb.11238, Cry3Bb.11239, Cry3Bb.11241, Cry3Bb.11242, and Cry3Bb.11098.

Some of the modified  $\delta$ -endotoxins that were described in WO 99/31248 and in U.S. Patent No. 6,063,597 were found to have enhanced activity against coleopteran insects, and in particular against  
20 *Diabrotica* spp., including corn rootworm. As used herein, the terms "enhanced activity" refer to the increased insecticidal activity of a modified toxin as compared with the activity of the same toxin without the amino acid modifications when both are tested under the same conditions. In particular, it was found that Cry3\*  $\delta$ -endotoxins had  
25 enhanced activity against corn rootworm, and are therefore preferred for use in the present invention. More preferred are Cry3B\*  $\delta$ -endotoxins, and even more preferred are Cry3Bb\*  $\delta$ -endotoxins. Even more preferred transgenic events are those that comprise the ability to express the modified  $\delta$ -endotoxins that are listed in the following table.  
30 Also shown in the table are strains of transgenic *B. thuringiensis* that include genes for expression of the respective novel endotoxins, and the date and accession number of their deposit with the Agricultural



Research Service Collection (NRRL) at 1815 N. University Street,  
Peoria, IL 91904.

STRAIN	DEPOSIT DATE	PROTEIN	ACCESSION NUMBER (NRRL NUMBER)
EG11230	5/27/97	Cry3Bb.11230	B-21768
EG11231	5/27/97	Cry3Bb.11231	B-21769
EG11232	5/27/97	Cry3Bb.11232	B-21770
EG11233	5/27/97	Cry3Bb.11233	B-21771
EG11234	5/27/97	Cry3Bb.11234	B-21772
EG11235	5/27/97	Cry3Bb.11235	B-21773
EG11236	5/27/97	Cry3Bb.11236	B-21774
EG11237	5/27/97	Cry3Bb.11237	B-21775
EG11238	5/27/97	Cry3Bb.11238	B-21776
EG11239	5/27/97	Cry3Bb.11239	B-21777
EG11241	5/27/97	Cry3Bb.11241	B-21778
EG11242	5/27/97	Cry3Bb.11242	B-21779
EG11098	11/28/97	Cry3Bb.11098	B-21903

It has also been found that a preferred use of the present  
invention is for reducing pest feeding damage when used in  
combination with seeds having transgenic events that have certain  
levels of effectiveness against such pest. To illustrate which levels of  
effectiveness are preferred, the following example will use the Iowa  
Root Rating Method (Hills and Peters, *J. Econ. Entomol.*, 64:764-765,  
1971), which measures corn rootworm feeding damage to corn roots  
on a 1 - 6 scale. In the rating, 1 = no damage or only a few minor  
feeding scars; 2 = feeding scars evident but no roots eaten off to within  
1½ inch of the plant; 3 = several roots eaten off to within 1½ inch of the  
plant, but never the equivalent of an entire node of roots is destroyed; 4  
= one root node completely destroyed; 5 = two root nodes completely  
destroyed; and 6 = three or more root nodes destroyed. A destroyed  
root is defined as a root that has been pruned to within 1½ inch of the

base. Pruned roots do not have to originate from a single node, but all pruned roots must equal the equivalent of a full node to count as a destroyed node.

5 As used herein, a transgenic event is within the preferred range of effectiveness level against a target pest if that event reduces feeding damage by that pest by a certain amount as compared with the same crop without the transgenic event, but does not prevent substantially all damage by the target pest. For example, if 10% of transgenic corn suffered corn rootworm damage of 4 or higher on the Iowa 1 - 6 Scale, 10 while 80% of non-transgenic corn suffered damage of 4 or higher, then it could be said that the damage to the transgenic corn was  $(10/80) \times 100 = 12.5\%$  of that of the non-transgenic corn. For the purposes of the present invention, it will be understood that a transgenic event in corn is within the preferred range of effectiveness level if corn having 15 such event suffers from about 5% to about 50% of the damage suffered by non-transgenic corn due to the same pest under the same conditions. It is more preferred that corn having such transgenic event suffers from about 10% to about 40% of the damage suffered by non-transgenic corn by the same pest under the same conditions, even 20 more preferred is damage of from about 15% to about 30%, and yet more preferred is damage of from about 20% to about 30% of the damage suffered by non-transgenic corn by the same pest under the same conditions. As used herein, when the term "about" is used to describe the degree of damage to corn, it is to be understood that the 25 degree of damage can be above or below the limits described by as much as 1% or 2% and still be considered to be within the ranges described. By way of example, a level of 4.5% damage would be regarded as being "about 5%".

30 Without wishing to be bound to this or any other theory, it is believed that the pesticidal seed treatment can provide significant advantages when combined with a transgenic event that provides protection that is within the preferred effectiveness range against a

target pest. In addition, it is believed that there are situations that are well known to those having skill in the art, where it is advantageous to have such transgenic events within the preferred range of effectiveness.

5           The present invention also includes seeds and plants having more than one transgenic event. Such combinations are referred to as "stacked" transgenic events. These stacked transgenic events can be events that are directed at the same target pest, or they can be directed at different target pests. In one preferred method, a seed  
10       having the ability to express a Cry 3 protein also has the ability to express at least one other insecticidal protein that is different from a Cry 3 protein.

          In another preferred method, the seed having the ability to express a Cry 3 protein also has a transgenic event that provides  
15       herbicide tolerance. It is more preferred that the transgenic event that provides herbicide tolerance is an event that provides resistance to glyphosate, N- (phosphonomethyl) glycine, including the isopropylamine salt form of such herbicide, even more preferred is the transgenic event that is effective to provide the herbicide resistance of  
20       ROUNDUP READY® plants and seeds available from Monsanto Co., St. Louis, MO.

          In the present method, a corn seed having a transgenic event is treated with thiamethoxam (3-[(2-chloro-5-thiazolyl)methyl]tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine; CAS RN 153719-23-4;  
25       and described in EP 580553, as well as in *The Pesticide Manual, Id.*, at 896).

          When an insecticide is described herein, it is to be understood that the description is intended to include salt forms of the insecticide as well as any isomeric and/or tautomeric form of the insecticide that  
30       exhibits the same insecticidal activity as the form of the insecticide that is described.



The insecticides that are useful in the present method can be of any grade or purity that pass in the trade as such insecticide. Other materials that accompany the insecticides in commercial preparations as impurities can be tolerated in the subject methods and compositions, as long as such other materials do not destabilize the composition or significantly reduce or destroy the activity of any of the insecticide components or the transgenic event against the target pest(s). One of ordinary skill in the art of the production of insecticides can readily identify those impurities that can be tolerated and those that cannot.

It has been found that the present method is useful to protect seeds and plants against a wide array of agricultural pests, including insects, mites, fungi, yeasts, molds, bacteria, nematodes, weeds, and parasitic and saprophytic plants.

When an insect is the target pest for the present invention, such pests include but are not limited to:  
from the order Lepidoptera, for example,

*Acleris* spp., *Adoxophyes* spp., *Aegeria* spp., *Agrotis* spp., *Alabama argillaceae*, *Amylois* spp., *Anticarsia gemmatilis*, *Archips* spp., *Argyrotaenia* spp., *Autographa* spp., *Busseola fusca*, *Cadra cautella*, *Carposina nipponensis*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocrocis* spp., *Cnephasia* spp., *Cochylis* spp., *Coleophora* spp., *Crocidolomia binotalis*, *Cryptophlebia leucotreta*, *Cydia* spp., *Diatraea* spp., *Diparopsis castanea*, *Earias* spp., *Ephestia* spp., *Eucosma* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Grapholita* spp., *Hedya nubiferana*, *Heliothis* spp., *Hellula undalis*, *Hyphantria cunea*, *Keiferia lycopersicella*, *Leucoptera scitella*, *Lithocollethis* spp., *Lobesia botrana*, *Lymantria* spp., *Lyonetia* spp., *Malacosoma* spp., *Mamestra brassicae*, *Manduca sexta*, *Operophtera* spp., *Ostrinia Nubilalis*, *Pammene* spp., *Pandemis* spp., *Panolis flammea*, *Pectinophora gossypiella*, *Phthorimaea operculella*, *Pieris rapae*, *Pieris* spp., *Plutella xylostella*, *Prays* spp., *Scirpophaga* spp.,

Sesamia spp., Sparganothis spp., Spodoptera spp., Synanthedon spp.,  
Thaumetopoea spp., Tortrix spp., *Trichoplusia ni* and Yponomeuta  
spp.;

from the order Coleoptera, for example,

- 5           Agriotes spp., Anthonomus spp., *Atomaria linearis*,  
*Chaetocnema tibialis*, Cosmopolites spp., Curculio spp., Dermestes  
spp., Diabrotica spp., Epilachna spp., Eremnus spp., *Leptinotarsa*  
*decemlineata*, Lissorhoptrus spp., Melolontha spp., Orycaephilus spp.,  
Otorhynchus spp., Phlyctinus spp., Popillia spp., Psylliodes spp.,  
10       Rhizopertha spp., Scarabeidae, Sitophilus spp., Sitotroga spp.,  
Tenebrio spp., Tribolium spp. and Trogoderma spp.;

from the order Orthoptera, for example,

Blatta spp., Blattella spp., Gryllotalpa spp., *Leucophaea*  
*maderae*, Locusta spp., Periplaneta spp., and Schistocerca spp.;

- 15       from the order Isoptera, for example,

Reticulitermes spp.;

from the order Psocoptera, for example,

Liposcelis spp.;

from the order Anoplura, for example,

- 20       Haematopinus spp., Linognathus spp., Pediculus spp.,  
Pemphigus spp. and Phylloxera spp.;

from the order Mallophaga, for example,

Damalinae spp. and Trichodectes spp.;

from the order Thysanoptera, for example,

- 25       Franklinella spp., Hercinothrips spp., Taeniothrips spp., *Thrips*  
*palmi*, *Thrips tabaci* and *Scirtothrips aurantii*;

from the order Heteroptera, for example,

Cimex spp., *Distantiella theobroma*, Dysdercus spp., Euchistus  
spp., Eurygaster spp., Leptocorisa spp., Nezara spp., Piesma spp.,  
30       Rhodnius spp., *Sahlbergella singularis*, Scotinophara spp. and  
Triatoma spp.;

from the order Homoptera, for example,

*Aleurothrixus floccosus*, *Aleyrodes brassicae*, *Aonidiella* spp.,  
*Aphididae*, *Aphis* spp., *Aspidiotus* spp., *Bemisia tabaci*, *Ceroplaster*  
spp., *Chrysomphalus aonidium*, *Chrysomphalus dictyospermi*, *Coccus*  
*hesperidum*, *Empoasca* spp., *Eriosoma larigerum*, *Erythroneura* spp.,  
5 *Gascardia* spp., *Laodelphax* spp., *Lacanium corni*, *Lepidosaphes* spp.,  
*Macrosiphus* spp., *Myzus* spp., *Nehotettix* spp., *Nilaparvata* spp.,  
*Paratoria* spp., *Pemphigus* spp., *Planococcus* spp., *Pseudaulacaspis*  
spp., *Pseudococcus* spp., *Psylla* spp., *Pulvinaria aethiopica*,  
*Quadraspidotus* spp., *Rhopalosiphum* spp., *Saissetia* spp.,  
10 *Scaphoideus* spp., *Schizaphis* spp., *Sitobion* spp., *Trialeurodes*  
*vaporariorum*, *Trioza erytreae* and *Unaspis citri*;  
from the order Hymenoptera, for example,

*Acromyrmex*, *Atta* spp., *Cephus* spp., *Diprion* spp., *Diprionidae*,  
*Gilpinia polytoma*, *Hoplocampa* spp., *Lasius* spp., *Monomorium*  
15 *pharaonis*, *Neodiprion* spp., *Solenopsis* spp. and *Vespa* spp.;  
from the order Diptera, for example,

*Aedes* spp., *Antherigona soccata*, *Bibio hortulanus*, *Calliphora*  
*erythrocephala*, *Ceratitis* spp., *Chrysomyia* spp., *Culex* spp., *Cuterebra*  
spp., *Dacus* spp., *Drosophila melanogaster*, *Fannia* spp., *Gastrophilus*  
20 spp., *Glossina* spp., *Hypoderma* spp., *Hyppobosca* spp., *Liriomyza*  
spp., *Lucilia* spp., *Melanagromyza* spp., *Musca* spp., *Oestrus* spp.,  
*Orseolia* spp., *Oscinella frit*, *Pegomyia hyoscyami*, *Phorbia* spp.,  
*Rhagoletis pomonella*, *Sciara* spp., *Stomoxys* spp., *Tabanus* spp.,  
*Tannia* spp. and *Tipula* spp.,  
25 from the order Siphonaptera, for example,

*Ceratophyllus* spp. und *Xenopsylla cheopis* and  
from the order Thysanura, for example,

*Lepisma saccharina*.

It has been found that the present invention is particularly  
30 effective when the insect pest is a *Diabrotica* spp., and especially when  
the pest is *Diabrotica virgifera*, *Diabrotica barberi*, or *Diabrotica*  
*undecimpunctata*.



Another application wherein the present invention is believed to be particularly effective is when the pesticide has activity against a weed or a parasitic or saprophytic plant and the transgenic event has activity against a member selected from the group consisting of

5     *Diabrotica virgifera*, *Diabrotica barberi* and *Diabrotica undecimpunctata*. This is believed to be more preferred useful when the weed or a parasitic or saprophytic plant is the African plant known as "Striga", and even more preferred when the pesticide is ROUNDUP® (available from Monsanto Company).

10             In the method of the present invention, thiamethoxam is applied to a transgenic corn seed. Although it is believed that the present method can be applied to a transgenic corn seed in any physiological state, it is preferred that the seed be in a sufficiently durable state that it incurs no damage during the treatment process. Typically, the seed

15     would be a seed that had been harvested from the field; removed from the plant; and separated from any cob, stalk, outer husk, and surrounding pulp or other non-seed plant material. The seed would preferably also be biologically stable to the extent that the treatment would cause no biological damage to the seed. In one embodiment,

20     for example, the treatment can be applied to seed corn that has been harvested, cleaned and dried to a moisture content below about 15% by weight. In an alternative embodiment, the seed can be one that has been dried and then primed with water and/or another material and then re-dried before or during the treatment with the pesticide. Within

25     the limitations just described, it is believed that the treatment can be applied to the seed at any time between harvest of the seed and sowing of the seed. As used herein, the term "unsown seed" is meant to include seed at any period between the harvest of the seed and the sowing of the seed in the ground for the purpose of germination and

30     growth of the plant.

When it is said that unsown seed is "treated" with the pesticide, such treatment is not meant to include those practices in which the

pesticide is applied to the soil, rather than to the seed. For example, such treatments as the application of the pesticide in bands, "T"-bands, or in-furrow, at the same time as the seed is sowed are not considered to be included in the present invention.

5           The pesticide, or combination of pesticides, can be applied "neat", that is, without any diluting or additional components present. However, the pesticide is typically applied to the seeds in the form of a pesticide formulation. This formulation may contain one or more other desirable components including but not limited to liquid diluents,  
10           binders to serve as a matrix for the pesticide, fillers for protecting the seeds during stress conditions, and plasticizers to improve flexibility, adhesion and/or spreadability of the coating. In addition, for oily pesticide formulations containing little or no filler, it may be desirable to add to the formulation drying agents such as calcium carbonate, kaolin  
15           or bentonite clay, perlite, diatomaceous earth or any other adsorbent material. Use of such components in seed treatments is known in the art. See, e.g., U.S. Patent No. 5,876,739. The skilled artisan can readily select desirable components to use in the pesticide formulation depending on the seed type to be treated and the particular pesticide  
20           that is selected. In addition, readily available commercial formulations of known pesticides may be used, as demonstrated in the examples below.

          The seeds may also be treated with one or more of the following ingredients: other pesticides, including compounds which act only  
25           below the ground; fungicides, such as captan, thiram, metalaxyl, (methoxam = resolved isomer of metalaxyl), fludioxonil, oxadixyl, and isomers of each of those materials, and the like; herbicides, including compounds selected from carbamates, thiocarbamates, acetamides, triazines, dinitroanilines, glycerol ethers, pyridazinones, uracils,  
30           phenoxys, ureas, and benzoic acids; herbicidal safeners such as benzoxazine, benzhydryl derivatives, N,N-diallyl dichloroacetamide, various dihaloacyl, oxazolidinyl and thiazolidinyl compounds, ethanone,

naphthalic anhydride compounds, and oxime derivatives; fertilizers; and biocontrol agents such as naturally-occurring or recombinant bacteria and fungi from the genera *Rhizobium*, *Bacillus*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Glomus*, *Gliocladium* and mycorrhizal fungi.

5 These ingredients may be added as a separate layer on the seed or alternatively may be added as part of the pesticide composition.

Preferably, the amount of the novel composition or other ingredients used in the seed treatment should not inhibit generation of the seed, or cause phytotoxic damage to the seed.

10 The pesticide formulation that is used to treat the transgenic corn seed in the present invention can be in the form of a suspension; emulsion; slurry of particles in an aqueous medium (e.g., water); wettable powder; wettable granules (dry flowable); and dry granules. If formulated as a suspension or slurry, the concentration of the active  
15 ingredient in the formulation is preferably about 0.5% to about 99% by weight (w/w), preferably 5-40%.

As mentioned above, other conventional inactive or inert ingredients can be incorporated into the formulation. Such inert ingredients include but are not limited to: conventional sticking agents,  
20 dispersing agents such as methylcellulose (Methocel A15LV or Methocel A15C, for example, serve as combined dispersant/sticking agents for use in seed treatments), polyvinyl alcohol (e.g., Elvanol 51-05), lecithin (e.g., Yelkinol P), polymeric dispersants (e.g., polyvinylpyrrolidone/vinyl acetate PVP/VA S-630), thickeners (e.g., clay  
25 thickeners such as Van Gel B to improve viscosity and reduce settling of particle suspensions), emulsion stabilizers, surfactants, antifreeze compounds (e.g., urea), dyes, colorants, and the like. Further inert ingredients useful in the present invention can be found in McCutcheon's, vol. 1, "*Emulsifiers and Detergents*," MC Publishing  
30 Company, Glen Rock, New Jersey, U.S.A., 1996. Additional inert ingredients useful in the present invention can be found in



McCutcheon's, vol. 2, "*Functional Materials*," MC Publishing Company, Glen Rock, New Jersey, U.S.A., 1996.

5 The pesticides and pesticide formulations of the present invention can be applied to seeds by any standard seed treatment methodology, including but not limited to mixing in a container (e.g., a bottle or bag), mechanical application, tumbling, spraying, and immersion. Any conventional active or inert material can be used for contacting seeds with pesticides according to the present invention, such as conventional film-coating materials including but not limited to  
10 water-based film coating materials such as Sepiret (Seppic, Inc., Fairfield, NJ) and Opacoat (Berwind Pharm. Services, Westpoint, PA).

The subject pesticides can be applied to a seed as a component of a seed coating. Seed coating methods and compositions that are known in the art are useful when they are modified by the addition of  
15 one of the embodiments of the combination of pesticides of the present invention. Such coating methods and apparatus for their application are disclosed in, for example, U.S. Patent Nos. 5,918,413, 5,891,246, 5,554,445, 5,389,399, 5,107,787, 5,080,925, 4,759,945 and 4,465,017. Seed coating compositions are disclosed, for example, in U.S. Patent  
20 Nos. 5,939,356, 5,882,713, 5,876,739, 5,849,320, 5,834,447, 5,791,084, 5,661,103, 5,622,003, 5,580,544, 5,328,942, 5,300,127, 4,735,015, 4,634,587, 4,383,391, 4,372,080, 4,339,456, 4,272,417 and 4,245,432, among others.

Useful seed coatings contain one or more binders and at least  
25 one of the subject combinations of pesticides.

Binders that are useful in the present invention preferably comprise an adhesive polymer that may be natural or synthetic and is without phytotoxic effect on the seed to be coated. The binder may be selected from polyvinyl acetates; polyvinyl acetate copolymers;  
30 ethylene vinyl acetate (EVA) copolymers; polyvinyl alcohols; polyvinyl alcohol copolymers; celluloses, including ethylcelluloses, methylcelluloses, hydroxymethylcelluloses, hydroxypropylcelluloses

and carboxymethylcellulose; polyvinylpyrrolidones; polysaccharides, including starch, modified starch, dextrans, maltodextrins, alginate and chitosans; fats; oils; proteins, including gelatin and zeins; gum arabics; shellacs; vinylidene chloride and vinylidene chloride copolymers; calcium lignosulfonates; acrylic copolymers; polyvinylacrylates; polyethylene oxide; acrylamide polymers and copolymers; polyhydroxyethyl acrylate, methylacrylamide monomers; and polychloroprene.

It is preferred that the binder be selected so that it can serve as a matrix for the subject pesticides. While the binders disclosed above may all be useful as a matrix, the specific binder will depend upon the properties of the combination of pesticides. The term "matrix", as used herein, means a continuous solid phase of one or more binder compounds throughout which is distributed as a discontinuous phase one or more of the subject pesticides. Optionally, a filler and/or other components can also be present in the matrix. The term matrix is to be understood to include what may be viewed as a matrix system, a reservoir system or a microencapsulated system. In general, a matrix system consists of pesticides of the present invention and filler uniformly dispersed within a polymer, while a reservoir system consists of a separate phase comprising the subject pesticides, that is physically dispersed within a surrounding, rate-limiting, polymeric phase. Microencapsulation includes the coating of small particles or droplets of liquid, but also to dispersions in a solid matrix.

The amount of binder in the coating can vary, but will be in the range of about 0.01 to about 25% of the weight of the seed, more preferably from about 0.05 to about 15%, and even more preferably from about 0.1% to about 10%.

As mentioned above, the matrix can optionally include a filler. The filler can be an absorbent or an inert filler, such as are known in the art, and may include woodflours, clays, activated carbon, sugars, diatomaceous earth, cereal flours, fine-grain inorganic solids, calcium

carbonate, and the like. Clays and inorganic solids, which may be used, include calcium bentonite, kaolin, china clay, talc, perlite, mica, vermiculite, silicas, quartz powder, montmorillonite and mixtures thereof. Sugars, which may be useful, include dextrin and maltodextrin. Cereal flours include wheat flour, oat flour and barley flour.

The filler is selected so that it will provide a proper microclimate for the seed, for example the filler is used to increase the loading rate of the active ingredients and to adjust the control-release of the active ingredients. The filler can aid in the production or process of coating the seed. The amount of filler can vary, but generally the weight of the filler components will be in the range of about 0.05 to about 75% of the seed weight, more preferably about 0.1 to about 50%, and even more preferably about 0.5% to 15%.

The pesticides that are useful in the coating are those pesticides that are described herein. The amount of pesticide that is used for the treatment of the seed will vary depending upon the type of seed and the type of active ingredients, but the treatment will comprise contacting the seeds with an amount of the pesticide that is pesticidally effective. When insects are the target pest, that amount will be an amount of the insecticide that is insecticidally effective. As used herein, an insecticidally effective amount means that amount of insecticide that will kill insect pests in the larvae or pupal state of growth, or will consistently reduce or retard the amount of damage produced by insect pests.

In general, the amount of thiamethoxam that is applied to the seed in the treatment will range from about 10 gm to about 2000 gm of the active ingredient of the pesticide per 100 kg of the weight of the seed. Preferably, the amount of pesticide will be within the range of about 50 gm to about 1000 gm active per 100 kg of seed, more preferably within the range of about 100 gm to about 600 gm active per 100 kg of seed, and even more preferably within the range of about



200 gm to about 500 gm of active per 100 kg of seed weight.

Alternatively, it has been found to be preferred that the amount of the pesticide be over about 60 gm of the active ingredient of the pesticide per 100 kg of the seed, and more preferably over about 80 gm per 100 kg of seed.

In preferred embodiments of the present invention the transgenic event comprises the ability to express a Cry3Bb.11231 protein or a Cry3Bb.11098 protein, and the pesticide is thiamethoxam.

The pesticides that are used in the treatment must not inhibit germination of the seed and should be efficacious in protecting the seed and/or the plant during that time in the target insect's life cycle in which it causes injury to the seed or plant. In general, the coating will be efficacious for approximately 0 to 120 days after sowing.

In the subject invention, the pesticides be applied to the seed in the form of a coating. The use of a coating is particularly effective in accommodating high pesticidal loads, as can be required to treat typically refractory pests, such as corn rootworm, while at the same time preventing unacceptable phytotoxicity due to the increased pesticidal load.

Optionally, a plasticizer can be used in the coating formulation. Plasticizers are typically used to make the film that is formed by the coating layer more flexible, to improve adhesion and spreadability, and to improve the speed of processing. Improved film flexibility is important to minimize chipping, breakage or flaking during storage, handling or sowing processes. Many plasticizers may be used, however, useful plasticizers include polyethylene glycol, glycerol, butylbenzylphthalate, glycol benzoates and related compounds. The range of plasticizer in the coating layer will be in the range of from about 0.1 to about 20% by weight.

When the pesticide used in the coating is an oily type formulation and little or no filler is present, it may be useful to hasten the drying process by drying the formulation. This optional step may

be accomplished by means well known in the art and can include the addition of calcium carbonate, kaolin or bentonite clay, perlite, diatomaceous earth, or any absorbent material that is added preferably concurrently with the pesticidal coating layer to absorb the oil or excess moisture. The amount of calcium carbonate or related compounds necessary to effectively provide a dry coating will be in the range of about 0.5 to about 10% of the weight of the seed.

The coatings formed with the pesticide are preferably of the type that are capable of effecting a slow rate of release of the pesticide by diffusion or movement through the matrix to the surrounding medium.

In addition to the coating layer, the seed may be treated with one or more of the following ingredients: other pesticides including fungicides and herbicides; herbicidal safeners; fertilizers and/or biocontrol agents. These ingredients may be added as a separate layer or alternatively may be added in the pesticidal coating layer.

The pesticide formulation may be applied to the seeds using conventional coating techniques and machines, such as fluidized bed techniques, the roller mill method, rotostatic seed treaters, and drum coaters. Other methods, such as spouted beds may also be useful. The seeds may be presized before coating. After coating, the seeds are typically dried and then transferred to a sizing machine for sizing. Such procedures are known in the art.

The pesticide-treated seeds may also be enveloped with a film overcoating to protect the pesticide coating. Such overcoatings are known in the art and may be applied using conventional fluidized bed and drum film coating techniques.

In another embodiment of the present invention, a pesticide can be introduced onto or into a seed by use of solid matrix priming. For example, a quantity of the pesticide can be mixed with a solid matrix material and then the seed can be placed into contact with the solid matrix material for a period to allow the pesticide to be introduced to the seed. The seed can then optionally be separated from the solid

matrix material and stored or used, or the mixture of solid matrix material plus seed can be stored or planted directly. Solid matrix materials which are useful in the present invention include polyacrylamide, starch, clay, silica, alumina, soil, sand, polyurea, polyacrylate, or any other material capable of absorbing or adsorbing the pesticide for a time and releasing that pesticide into or onto the seed. It is useful to make sure that the pesticide and the solid matrix material are compatible with each other. For example, the solid matrix material should be chosen so that it can release the pesticide at a reasonable rate, for example over a period of minutes, hours, or days.

The present invention further embodies imbibition as another method of treating seed with the pesticide. For example, plant seed can be combined for a period of time with a solution comprising from about 1% by weight to about 75% by weight of the pesticide in a solvent such as water. Preferably the concentration of the solution is from about 5% by weight to about 50% by weight, more preferably from about 10% by weight to about 25% by weight. During the period that the seed is combined with the solution, the seed takes up (imbibes) a portion of the pesticide. Optionally, the mixture of plant seed and solution can be agitated, for example by shaking, rolling, tumbling, or other means. After imbibition, the seed can be separated from the solution and optionally dried, for example by patting or air drying.

In yet another embodiment, a powdered pesticide can be mixed directly with seed. Optionally, a sticking agent can be used to adhere the powder to the seed surface. For example, a quantity of seed can be mixed with a sticking agent and optionally agitated to encourage uniform coating of the seed with the sticking agent. The seed coated with the sticking agent can then be mixed with the powdered pesticide. The mixture can be agitated, for example by tumbling, to encourage contact of the sticking agent with the powdered pesticide, thereby causing the powdered pesticide to stick to the seed.



The present invention also provides a transgenic corn seed that has been treated with a pesticide by the method described above.

The treated seeds of the present invention can be used for the propagation of corn plants in the same manner as conventional treated corn seed. The treated seeds can be stored, handled, sowed and tilled in the same manner as any other pesticide treated seed. Appropriate safety measures should be taken to limit contact of the treated seed with humans, food or feed materials, water and birds and wild or domestic animals.

Preferred embodiments of the invention are described in the following examples. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

The following examples describe preferred embodiments of the invention. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples. In the examples all percentages are given on a weight basis unless otherwise indicated.

#### EXAMPLE 1

Production of corn seed having a transgenic event effective against corn rootworm and treatment of such seed with thiamethoxam (Cruiser®).

Corn seeds can be prepared to express *Bacillus thuringiensis* endotoxin Cry3Bb.11231 or Cry3Bb.11098 by the methods described

for these respective events in WO 99/31248, or in U.S. Patent No. 6,023,013.

Corn seeds of the same hybrid species, with and without the respective transgenic events, can be treated with thiamethoxam (available as CRUISER® from Novartis) as follows:

A seed treatment formulation of the desired pesticide is prepared by mixing a measured amount in water as a carrier and applying the formulation for one minute at room temperature to a measured weight of corn seed in a rotostatic seed treater. The respective weights of the pesticide preparation and the corn seed are calculated to provide the desired rate of treatment of pesticide on the seed. The pesticide is mixed into sufficient water to permit efficient distribution of the formulation to all of the seeds in the batch while minimizing loss of treatment formulation due to lack of uptake of the formulation by the seeds. Treated seeds are allowed to sit uncapped for at least four hours before planting.

When the seeds are treated with thiamethoxam, a sufficient amount of CRUISER® is thoroughly mixed into water to form a seed treatment formulation, and the formulation is applied to a weight of corn seed to provide treatment levels of 300 grams thiamethoxam per 100 kg of seed (.75 mg thiamethoxam/kernel), or 400 grams thiamethoxam per 100 kg of seed (1.0 mg thiamethoxam/kernel).

#### EXAMPLE 2

Field trials for the determination of efficacy of transgenic event Cry3Bb.11231 in corn seed in combination with corn root worm pesticide seed treatments against western and northern corn rootworm.

Field trials can be run in accordance with pertinent protocols and in conformance with USDA notification requirements. The purpose of the trials is to determine the efficacy of transgenic event Cry3Bb.11231 in corn seed in combination with corn root worm seed treatments against western and northern corn root worm.

For each growing site that is selected, the plot design would included the following:

5	Row spacing:	30 inches
	Plot size:	4 rows x 20
	Plant density:	2.0 seed/foot
	Hybrid used:	LH198 x LH185 or RX670
	Replicates:	4
10	Design:	Randomized complete block
	Locations:	4
	Larvae source:	natural infestations supplemented by artificial infestation of corn rootworm eggs at 400 eggs/ft (growth stage V2)

15

The following seed treatment combinations can be used for each growing area:

	<u>No.</u>	<u>Corn Seed Type</u>	<u>Pesticide and amount (grams AI/100</u>
			<u>kg seed or mg ai/kernel)</u>
20	1	Isohybrid	None, other than low levels for wire worm protection
	2	Cry3Bb.11231	None, other than low levels for wire worm protection
25	3	Cry3Bb.11231	CRUISER® @ 300 gm AI/100 kg or .75 mg AI/kernel
	4	Cry3Bb.11231	CRUISER® @ 400 gm AI/ 100 kg or 1.0 mg AI/kernel
30	5	Isohybrid	Force® 3G @ 0.014 gm AI/ m, or 0.15 oz AI/ 1000 ft row, applied as a



- 5" band on the soil surface at the time of planting.
- 6 Isohybrid Lorsban® 15G (chlorpyrifos; available from DowElanco) @ 0.11 gm AI/m, or 1.2 oz AI/1000 ft row, applied as a 5" band on the soil surface at the time of planting.
- 5

10 All seed treatments with pesticides are carried out as described in Example 1. In seed treatment number 1 and 2, Gaucho® (imidacloprid) can be used for wire worm protection, but at levels sufficiently low that it would be expected to have no effect on corn rootworms (*i.e.*, at a treatment level of about 60 gm of active/ 100 kg seed or 0.16 mg active/kernel), otherwise, seed receiving treatment number 2 should have only transgenic event Cry3Bb.11231 and no pesticide treatment that would be expected to be effective against corn rootworm.

15

For seeds having treatments numbered 3 and 4, the pesticides are applied by the methods described in Example 1. For seeds having treatment numbers 5 and 6, commercially available Force® 3G and Lorsban® 15G are applied to the soil in a 5" band at the time of sowing to provide a comparison with standard practices. The levels of application are as shown and are within the ranges recommended for standard commercial practice.

20

25 Corn seeds to be tested should be planted and grown at several different locations according to the protocol described above.

The determination of damage by corn rootworm can be made according to the following protocol. At stage V4 - V6, an evaluation of early stand is made by counting the number of plants per acre. At stage VT-R1, an evaluation of corn rootworm damage can be carried out by methods that are well known in the industry, and damage by corn rootworm is reported according to the Iowa 1 - 6 rating system. In

30

that system, the root systems of 10 corn plants per plot are recovered and scored using the 1 - 6 rating scale, where: 1 = no injury or only a few minor feeding scars, 2 = feeding injury evident, but no roots eaten back to 1 1/2 inches of the plant, 3 = at least one root eaten off to within 1 1/2 inches of the plant, but never an entire node of roots destroyed, 4 = one node of roots eaten back to within 1 1/2 inches of the plant, 5 = two nodes (circles) of roots eaten back to within 1 1/2 inches of the plant, 6 = three nodes (circles) of roots eaten back to within 1 1/2 inches of the plant.

It is believed that the data that is obtained from these techniques would show that transgenic seeds that were treated with thiamethoxam were more resistant to corn rootworm damage than the transgenic seeds without such pesticide treatment. Moreover, it is believed that such tests would show that combination treatments (of transgenic event plus pesticide treatment) were more efficacious than conventional surface banding with either FORCE® or LORSBAN®.

The discussion of the references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinence of the cited references.

In view of the above, it will be seen that the several advantages of the invention are achieved and other advantageous results attained.

As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.

**WHAT IS CLAIMED IS:**

1. A method for protecting a transgenic corn plant against feeding damage by one or more pests, the method comprising providing a seed for the transgenic corn plant which seed  
5 comprises a transgenic event having activity against at least one of the one or more pests; and  
treating the seed with an effective amount of thiamethoxam pesticide.
2. The method according to claim 1 wherein the transgenic  
10 event comprises the ability to express a Cry 3 protein.
3. The method according to claim 2 wherein the transgenic event comprises the ability to express a Cry3\* protein.
4. The method according to claim 3 wherein the Cry3\* protein comprises a Cry3B\* protein.
- 15 5. The method according to claim 4 wherein the Cry 3B\* protein is selected from the group consisting of Cry3Bb.11230, Cry3Bb.11231, Cry3Bb.11232, Cry3Bb.11233, Cry3Bb.11234, Cry3Bb.11235, Cry3Bb.11236, Cry3Bb.11237, Cry3Bb.11238, Cry3Bb.11239, Cry3Bb.11241, Cry3Bb.11242, and Cry3Bb.11098.
- 20 6. The method according to claim 5 wherein the Cry 3B\* protein comprises Cry3Bb.11231.
7. The method according to claim 5 wherein the Cry3B\* protein comprises Cry3Bb.11098.
8. The method according to claim 2 wherein the transgenic  
25 event is capable of reducing the damage caused by corn rootworm so that the damage to transgenic corn is within the range of from about 5% to about 50% of the damage to non-transgenic corn under the same conditions, when such damage is expressed as the percent of corn plants having a score of 4 - 6 as measured by the Iowa Corn  
30 Rootworm 1 - 6 Scale.
9. The method according to claim 8 wherein the transgenic event is capable of reducing the damage caused by corn rootworm so



that the damage to transgenic corn is within the range of from about 10% to about 40% of the damage to non-transgenic corn under the same conditions, when such damage is expressed as the percent of corn plants having a score of 4 - 6 as measured by the Iowa Corn Rootworm 1 - 6 Scale.

10. The method according to claim 9 wherein the transgenic event is capable of reducing the damage caused by corn rootworm so that the damage to transgenic corn is within the range of about 15% to about 30% of the damage to non-transgenic corn under the same conditions, when such damage is expressed as the percent of corn plants having a score of 4 - 6 as measured by the Iowa Corn Rootworm 1 - 6 Scale.

11. The method according to claim 10 wherein the transgenic event is capable of reducing the damage caused by corn rootworm so that the damage to transgenic corn is within the range of about 20% to about 30% of the damage to non-transgenic corn under the same conditions, when such damage is expressed as the percent of corn plants having a score of 4 - 6 as measured by the Iowa Corn Rootworm 1 - 6 Scale.

12. The method according to claim 2 wherein the seed having the ability to express a Cry 3 protein also has the ability to express at least one other insecticidal protein that is different from a Cry 3 protein.

13. The method according to claim 2 wherein the seed having the ability to express a Cry 3 protein also has a transgenic event that provides herbicide tolerance.

14. The method according to claim 13 wherein the transgenic event that provides herbicide tolerance against glyphosate.

15. The method according to claim 2 wherein the transgenic event has activity against an insect.

16. The method according to claim 15 wherein the insect is selected from the group consisting of members of the orders of Lepidoptera, Coleoptera and Hemiptera.

5 17. The method according to claim 16 wherein the insect is a Coleopteran insect.

18. The method according to claim 17 wherein the insect is a *Diabrotica* spp.

10 19. The method according to claim 18 wherein the insect comprises at least one member selected from the group consisting of *Diabrotica virgifera*, *Diabrotica barberi* and *Diabrotica undecimpunctata*.

20. The method of claim 2 wherein the effective amount of thiamethoxam is from at least about 10 grams to about 2000 grams of the pesticide active ingredient per 100 kilograms of the seed.

15 21. The method of claim 20 wherein the effective amount of thiamethoxam is from at least about 70 grams to about 1000 grams of the pesticide active ingredient per 100 kilograms of the seed.

20 22. The method of claim 21 wherein the effective amount of thiamethoxam is from at least about 100 grams to about 600 grams of the pesticide active ingredient per 100 kilograms of the seed.

23. The method of claim 22 wherein the effective amount of the pesticide is from at least about 200 grams to about 500 grams of the pesticide active ingredient per 100 kilograms of the seed.

25 24. A seed of a transgenic corn plant that provides increased resistance to the resulting corn plant against feeding damage by one or more pests, comprising a transgenic event having activity against at least one of the one or more pests, which seed has been treated with an effective amount of thiamethoxam.

30 25. The seed according to claim 24 wherein the transgenic event comprises the capability to express a Cry3 protein derived from *Bacillus thuringiensis*.

26. The seed according to claim 25 wherein the transgenic event comprises the capability to express a Cry3Bb protein derived from *Bacillus thuringiensis*.

5 27. The seed according to claim 26 wherein the transgenic event is the capability to express a Cry3Bb.11230, Cry3Bb.11231, Cry3Bb.11232, Cry3Bb.11233, Cry3Bb.11234, Cry3Bb.11235, Cry3Bb.11236, Cry3Bb.11237, Cry3Bb.11238, Cry3Bb.11239, Cry3Bb.11241, Cry3Bb.11242, or a Cry3Bb.11098 protein derived from *Bacillus thuringiensis*

10 28. A transgenic corn seed that has been treated by the method of claim 20.