

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2006237680 B2**

(54) Title
Analysing breath samples for volatile organic compound

(51) International Patent Classification(s)
C12Q 1/04 (2006.01) **C12Q 1/22** (2006.01)

(21) Application No: **2006237680** (22) Date of Filing: **2006.04.13**

(87) WIPO No: **WO06/112733**

(30) Priority Data

(31) Number	(32) Date	(33) Country
539518	2005.04.19	NZ

(43) Publication Date: **2006.10.26**

(44) Accepted Journal Date: **2011.11.17**

(71) Applicant(s)
Otago Innovation Limited

(72) Inventor(s)
Chambers, Stephen T;Syhre, Mona;Scotter, Jennifer M

(74) Agent / Attorney
Garth Hendry, PO Box 200, Phillip, ACT, 2606

(56) Related Art
Gao, P. et al, AIHA Journal, 2002, Volume 63, pages 135-140

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 October 2006 (26.10.2006)

PCT

(10) International Publication Number
WO 2006/112733 A1

(51) International Patent Classification:
C12Q 1/04 (2006.01) *C12Q 1/22* (2006.01)

(74) Agent: BALDWIN; 342 Lambton Quay, Wellington
6001 (NZ).

(21) International Application Number:
PCT/NZ2006/000071

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 13 April 2006 (13.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
539518 19 April 2005 (19.04.2005) NZ

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): CANE-
SIS NETWORK LTD [NZ/NZ]; Corner Of Springs Road
& Gerald Street, Lincoln, Canterbury 8152 (NZ).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCOTTER, Jen-
nifer M [NZ/NZ]; South Eyre Road, Kaiapoi, Christchurch
8250 (NZ). CHAMBERS, Stephen T [NZ/NZ]; 27
Kiteroa Place, Cashmere, Christchurch, 8002 (NZ).
SYHRE, Mona [DE/NZ]; 142 Fisher Ave Beckenham,
Christchurch 8002 (NZ).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 2006/112733 A1

(54) Title: ANALYSING BREATH SAMPLES FOR VOLATILE ORGANIC COMPOUND

(57) Abstract: The invention describes the use of a biomarker pentylfuran to detect bacterial and/or fungal pathogens. Pentylfuran is released from certain pathogens and detected in the headspace gas of an *in vitro* culture or in the breath sample of a patient. Pentylfuran is particularly useful in the detection of *Aspergillus* species, especially *Aspergillus fumigatus* which is a pathogen of humans.

ANALYSING BREATH SAMPLES FOR VOLATILE ORGANIC COMPOUND

Field of the Invention

- 5 The invention relates to the determination of a unique biomarker pentylfuran and to its use to test for fungal and bacterial pathogens, including *Aspergillus fumigatus* in an individual or from a culture using analysis of breath/air samples.

Background to the invention

- 10 All micro-organisms produce by-products as a result of their normal metabolism. The ability of different organisms to metabolise different substrates in order to satisfy their energy and nutritional requirements is fundamental to laboratory microbiology, and forms the basis of many rapid identification tests. The metabolites produced by a single species can vary widely, depending upon the growth substrate, conditions (temperature, oxygen
15 availability), and the age of the culture itself.

- Amongst the many large primary and secondary metabolites produced by microbes, some organic substances are formed which readily volatilise at low temperatures. Microbial volatile organic compounds (MVOCs) have been studied extensively in agriculture and food production, as some MVOCs have important health and economic implications in
20 these fields. For example, some MVOCs have been associated with spoilage in stored crops and foodstuffs, where they may be responsible for tainted, "off" flavours, discolouring of products, or toxicity. Profiles of MVOCs are increasingly being found to be unique to the species or strain level. This should not seem surprising, as it is the

2

different biochemical abilities of different microbes that gives them most of their identifiable and distinguishing traits.

Invasive aspergillosis is one of the most problematic infections due to difficulties of
5 diagnosis and treatment. Volatile organic compounds (VOCs) have the potential to improve the specificity and sensitivity of diagnosis of this and other infections. It would be useful to identify a unique biomarker of *Aspergillus* species, particularly *Aspergillus fumigatus*, in the headspace gas of *in vitro* cultures and to detect the marker from breath samples of infected or colonised patients.

10

Object of the Invention

It is an object of the invention to provide a biomarker to detect bacterial and/or fungal pathogens such as *Aspergillus fumigatus* in a bio sample or at least to provide the public with a useful choice.

15

Summary of the Invention

The invention provides the biomarker, pentylfuran for the use in bio analysis of micro organisms such as fungal and bacterial pathogens. In particular, the invention provides the use of the biomarker pentylfuran to detect fungal species, more particularly, *Aspergillus*
20 species in a biosample.

More particularly the invention provides the use of the biomarker pentylfuran to detect *Aspergillus fumigatus* in a biosample.

The biosample is preferably a gaseous biological source.

The biosample may, most preferably, be the headspace gas of an *in vitro* culture or a breath
5 sample of a patient, or another biosample such as a sputum sample.

Other micro organisms such as *A-flavus*, *Haemophilus influenza* and *Pseudomonas aeruginosa* may also be detected. The bio sample is preferably a breath sample. Preferably the biomarker is 2-pentylfuran. However it could be 3-pentylfuran.

10

The invention provides a method of detecting *Aspergiullus* in a patient comprising:

- (a) obtaining a biosample from the breath of the patient;
- (b) analysing the biosample for the presence of pentylfuran; and
- (c) determining whether *Aspergillus* is present in the breath sample.

15

The invention also provides a method of detecting *Aspergillus* species in an *in vitro* culture comprising:

- a. obtaining a biosample from the headspace gas of the *in vitro* culture;
- b. analysing the biosample for the presence of pentylfuran; and
- 20 c. determining whether *Aspergillus* is present in the *in vitro* culture.

4

The invention also provides the use of a biomarker pentylfuran in the detection of *Aspergillus fumigatus* from a breath sample of an animal or from an air sample from a culture

- 5 The *Aspergillus* species is preferably *Aspergillus fumigatus*.

Description of the Drawings

Figure 1 shows a chest CT scan of a patient with multiple foci of air space opacity surrounded by hyperdense material.

10

Detailed Description of the Invention

The invention will now be described, by way of example only

Gas Chromatography-Mass Spectroscopy (GC-MS) combined with Solid Phase Micro

- 15 Extraction (SPME) was used to identify pentylfuran as a specific biomarker of *A.fumigatus* from cultures. Four litre breath samples were collected from patients with Cystic Fibrosis, with or without colonisation of *A.fumigatus* and other pathogens, and healthy volunteers. Breath samples were semi-quantitatively analysed by SPME / GC-MS for presence or absence of pentylfuran.

- 20 A total of 21 individuals were tested. Pentylfuran was detected from breath samples of 4/4 patients with CF and *A.fumigatus* colonisation, 3/7 patients with CF and no microbiological evidence of *A.fumigatus* and 0/10 healthy control individuals.

Materials and Methods

Strains and culture conditions

Clinical isolates of *Aspergillus flavus*, *Aspergillus fumigatus*, *Candida albicans*, *Mucor racemosus*, *Fusarium solani*, and *Cryptococcus neoformans* were used in these
5 experiments.

Organisms were grown on blood agar within 100 ml sterile glass vials stoppered with airtight aluminium caps incorporating a teflon-coated rubber septum.

Strain preparation.

Strains were grown for 72 hours on blood agar plates, then, in the case of yeast species, a
10 sterile loop of culture was removed from the plate and transferred to 5 mL of sterile water. For filamentous fungal strains, spores were harvested from the plate with sterile water containing 0.05% Tween. Five hundred microlitres of this suspension was introduced to the sealed culture vial by injecting through the septum onto the medium.

Cultures were maintained at 37°C for 5 days. Vials were flushed with 100 ml purified dry
15 air once every approximately 12 hours.

Detection of pentylfuran by GC-MS, Calibration and standardisation

Calibration curves of headspace gas analysis of serial aqueous dilutions of pentylfuran were plotted. The resulting calibration curve proved to be linear in the range 1-50 pg.

Patient demographics

Four participants with cystic fibrosis and *Aspergillus* colonisation, 6 patients with CF and no *Aspergillus* colonisation, and 10 control individuals meeting the inclusion criteria for the study were identified. Relevant demographic and clinical data are shown in table 2.

5 Solid –Phase Micro Extraction (SPME) Gas Chromatography - Mass Spectrometry (GC-MS).

A comprehensive literature review was performed, and a database of all reported MVOCs from *Aspergillus* species was created, which included the compound isolated, species and strain, culture medium and conditions, and analysis method. Headspace gases of strains
10 cultured as described above were subjected to SPME / GC-MS analysis.

Sample preparation

The conditioned SPME fibre was exposed into culture vials for 10 minutes and then desorbed directly in the injection port for 5 minutes.

GC/MS parameters

15 The temperatures of the injector, ion trap, manifold and transfer line were 250, 200, 60 and 250°C, respectively. The oven program commenced at 50°C for 2 minutes and was raised to 250°C at a rate of 10°C/min, at which temperature it was maintained for a further 2 minutes. Helium flow was set at a constant rate of 1.2 ml/min. The split vent was opened to a ratio of 1:50 after 1 minute. Fragmentation was performed in the EI-mode as full scan
20 which gave additional certainty. Further MS/MS fragmentation could be used to further increase sensitivity.

Calibration and semi-quantification

Fifty microlitres of diluted pentylfuran solutions in methanol were deposited into 20 mL headspace vials. The final calibration was made up as total amounts in the headspace vials containing 1,5,10 and 50 pg, respectively. The fibre was exposed into the headspace vials
5 for 5 min using the Combi-PAL autosampler.

Patient selection

Patients enrolled on the study included individuals with cystic fibrosis (CF) colonised chronically with aspergillus, patients with CF not colonised with aspergillus, and healthy control individuals. Ethical approval for the study was obtained from the local ethics
10 committee, and participants gave their informed consent to take part in the study. Inclusion criteria for patients to act as “positives” (colonised with *Aspergillus*) were a history of positive culture for *A. fumigatus* from lower respiratory tract specimens (sputum, BAL, tracheal aspirate, or cough swab in cases where sputum had not been made available to the microbiology laboratory). Patients needed to have a minimum of three positive results
15 within the past 12 months, and ideally one from within 1 month of testing. Patients were excluded from the study if they were currently undergoing treatment with itraconazole for ABPA.

Patients in the cystic fibrosis “uncolonised” group were selected if they had no current or past history of positive culture results for *Aspergillus* as described above, and no clinical
20 evidence of aspergillus colonisation. Healthy control individuals were recruited from laboratory staff. These participants were asked to complete a questionnaire which served to give information concerning any recent antibiotic use or evidence of respiratory or urinary tract infection. Any participants not meeting these criteria were excluded from the study.

Breath sampling

Breath samples were collected into a 4L tedlar bag, which incorporated a valve, disposable mouthpiece and septum that could be pierced for sampling. Samples were collected by asking participants to exhale through the mouth into the bag until full. The valve in the bag
5 was then closed, and samples transported immediately to the laboratory for testing.

Analysis of breath samples by GC-MS

Breath samples were analysed by GC-MS for presence and quantity of pentylfuran as described for cultures above. The conditioned SPME fibre was exposed into the collection bags for 48h and then desorbed directly in the injection port for 5 minutes.

Results

Detection of pentylfuran from laboratory cultures.

The results for testing of laboratory strains for the presence of pentylfuran are given in Table 1.

5 Table 1. Results of screening fungal and bacterial isolates by GC-MS for presence of pentylfuran.

Organism	Source	pentylfuran			
			<i>A.fumigatus</i>	Environmental	xx
<i>A.flavus</i>	Clinical (ear swab)	nd	<i>A.fumigatus</i>	Environmental	xxx
<i>A.flavus</i>	Clinical (ear swab)	nd	<i>A.fumigatus</i>	Environmental	xxx
<i>A.niger</i>	Environmental	xx	<i>A.fumigatus</i>	Environmental	xxx
<i>A.niger</i>	Environmental	xx	<i>A.fumigatus</i>	Environmental	xxx
<i>F.oxysporum</i>	Environmental	x	<i>A.fumigatus</i>	Environmental	xxx
<i>M.racemosus</i>	Clinical (sinus biopsy)	x	<i>A.fumigatus</i>	Environmental	xx
<i>C.albicans</i>	Clinical (urine)	x	<i>A.fumigatus</i>	Environmental	xx
<i>C.albicans</i>	Clinical (urine)	x	<i>H.influenzae</i>	Clinical (sputum)	x
<i>A.fumigatus</i>	Clinical (lung biopsy)	xx	<i>B.cepacia</i>	Clinical (sputum)	x
<i>A.fumigatus</i>	Clinical (sputum)	xx	<i>Ps.aeruginosa</i>	Clinical (sputum)	x
<i>A.fumigatus</i>	Clinical (sputum)	xx	<i>Ps.aeruginosa</i>	Clinical (sputum)	x
<i>A.fumigatus</i>	Clinical (BAL)	xx	<i>S.aureus</i>	Clinical (sputum)	nd
<i>A.fumigatus</i>	Clinical (sputum)	xx	<i>S.aureus</i>	Clinical (sputum)	nd
<i>A.fumigatus</i>	Clinical (sputum)	xx			
<i>A.fumigatus</i>	Clinical (sputum)	xx			
<i>A.fumigatus</i>	Clinical (sputum)	xxx			
<i>A.fumigatus</i>	Clinical (sputum)	xx			
<i>A.fumigatus</i>	Clinical (ear swab)	xx			
<i>A.fumigatus</i>	Environmental	xx			
<i>A.fumigatus</i>	Environmental	xx			

Source: Spt; sputum, BAL; bronchoalveolar lavage. pentylfuran: nd; not detected; x; low/trace levels, xx; moderate levels; xxx; high levels.

Analysis of breath samples for presence of pentylfuran

Four litre breath samples were assayed for the presence and quantity of pentylfuran.

Results are shown in table 2, along with relevant clinical and microbiological data.

Table 2 Demographic and microbiological data of patients enrolled in the study, and results of testing breath by SPME / GC-MS for presence and quantity of pentylfuran. "Aspergillus colonisation" refers to colonisation with *Aspergillus fumigatus* only. Organisms shown in bold are those shown to produce trace levels of pentylfuran.. Other organisms isolated by culture of respiratory samples is shown. Organisms shown in bold are known producers of pentylfuran.

	ID	Age	Sex	Underlying disease	<i>S.aureus</i>	<i>S.pneumoniae</i>	<i>M.catarrhalis</i>	<i>H.influenzae</i>	<i>Ps.aeruginosa</i>	<i>A.fumigatus</i>	2-pentyl furan (pg)
CF +aspergillus	12	20	F	CF				+	+	+	7
	16	20	F	CF	+			+		+	3
	9	5	F	CF				+		+	7
	30		M	CF					+	+	5
CF, no aspergillus	10	7	M	CF	+	+		+			10
	11	7	M	CF	+	+		+			nd
	17	19	M	CF	+				+		2
	19	9	M	CF				+			nd
	20	7	M	CF			+	+			nd
	21	7	M	CF		+	+	+			9

	22	8	M	CF	+	+	nd
	18	30	F	none	.		nd
	13	25	F	none			nd
	14	33	M	none			nd
	15	31	M	none			nd
	24	35	M	none			nd
	25	36	M	none			nd
	26	36	M	none			nd
Normal controls	27	44	M	none			nd
	28	37	M	none			nd
	29	57	M	none			nd

Pentylfuran was detected on the breath of all (n=4) patients colonised with *A.fumigatus*. Pentylfuran was also detected on breath samples of 3 of 7 patients without evidence of *Aspergillus* colonisation, but colonisation with other pathogens, including some shown to produce pentylfuran in modest quantities. Of the healthy individuals, one of ten subjects showed a trace level of pentylfuran on the breath.

The important message from this data set is that healthy, normal individuals appear not to produce pentylfuran (or only at baseline levels), while those with pulmonary colonisation with organisms that produce pentylfuran, do. This is the first time that the detection and quantification of a specific microbial metabolite from breath of infected/colonised individuals has been reported.

Pentylfuran is a metabolite of *Aspergillus fumigatus*, and possibly other fungal and bacterial pathogens. These *in vitro* experiments showed that low levels of pentylfuran is produced by *A.flavus*, as well as *Pseudomonas aeruginosa* and *Haemophilus influenzae*, both of which are common colonisers of the lungs of patients with CF .

Blood tests for *Aspergillus* infection are not optimal in that they are prone to sampling error, and may return false-negative results even in cases of proven disseminated infection. This may be further compounded by the administration of systemic antifungal therapy. Use of Broncho-alveolar lavage (BAL) for diagnostic purposes has had increasing support over recent years, notwithstanding that it is not an ideal procedure to be performing in a neutropenic, and often thrombocytopenic patient . However, BAL has the advantage over blood testing in that it allows direct sampling of the site of primary infection. Breath testing for metabolites of pathogens resident in the lungs may be viewed in a similar way to BAL, except that it is without the risks associated with BAL. The principle of breath testing, like BAL, is a more direct way of sampling the site of primary infection. DNA and antigen detection based assays will detect both viable and non-viable fungal cell elements. Detection of *A.fumigatus* metabolites from breath samples would suggest that the organism is metabolically active.

The results of this study show that:

1. Pentylfuran may be used as a biomarker of *Aspergillus fumigatus*, from cultures.
2. Pentylfuran can be detected from the breath of patients colonised or infected with *Aspergillus fumigatus* by GC-MS

3. Detection of pentylfuran from breath may form the basis of a useful diagnostic test for aspergillus infection.

Case Report

A group of 79 woman undergoing treatment with dexamethasone and cyclophosphamide for multiple myeloma suffered an episode of febrile neutropenia (neutrophils $0.5-0.9 \times 10^9/L$) that failed to resolve despite broad spectrum antibiotic therapy. A CT of the sinuses showed extensive disease suggestive of *Aspergillus* and the chest CT showed multiple foci of air space opacity with a surrounding halo of hyperdense material, suggestive of aspergillus infection (arrowed, figure 1). *A. fumigatus* was cultured from the sputum on 2 occasions. Nested PCR for *A. fumigatus* DNA in peripheral blood, and was negative on 6 occasions. The patient gave informed consent to participate in the study, and 3-litre breath samples were collected as described above for patients with CF. 2-pentylfuran was detected (10.1 pg) in the exhaled breath 2 days after the CT scan was performed. After treatment with voriconazole for 4 weeks, the lesions in the lungs had reduced in size, the breath test had become negative and aspergillus was no longer culturable from the sputum.

Industrial Applicability

The invention will be of use in the medical area, assisting in the detection of microbes and pathogens in patients. In particular, the ability to detect *Aspergillus* species, especially *Aspergillus fumigatus* will be of use in detecting and hence treatment of the infection caused by the microbe in patients.

2006237680 01 Nov 2011

Claims

1. Use of the biomarker pentylfuran in the bio analysis of microorganisms in a gaseous bio sample from breath of an animal.
2. Use according to claim 1 wherein the microorganisms are fungal pathogens.
3. Use according to claim 1 or claim 2 wherein the microorganisms are from the group comprising *Aspergillus* species.
4. Use according to claim 3 wherein the microorganism is an *Aspergillus* species.
5. Use according to claim 4 wherein the microorganism is *Aspergillus fumigatus*.
6. Use according to any one of claims 1 to 5 wherein pentylfuran is 2-pentylfuran.
7. Use according to any one of claims 1 to 6 wherein the animal is a human.
8. A method of detecting *Aspergillus* in a patient comprising:
 - (a) obtaining a biosample from the breath of the patient;
 - (b) analyzing the biosample for the presence of pentylfuran; and
 - (c) determining whether *Aspergillus* is present in the breath biosample.
9. The method according to claim 7 wherein the *Aspergillus* species is *Aspergillus fumigatus*.
10. Use of a biomarker pentylfuran in the detection of *Aspergillus fumigatus* from a breath sample of an animal.
11. Use according to claim 9 in which the animal is a human.
12. Use according to any one of claims 1 to 7, 10 or 11 substantially as hereinbefore described with or without reference to Figure 1.
13. The method according to claim 8 or claim 9 substantially as hereinbefore described with or without reference to Figure 1.

1/1

Figure 1

