



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>A01N 63/04, 25/00, C12R 1/645</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 95/25430</b></p> <p>(43) International Publication Date: 28 September 1995 (28.09.95)</p>
<p>(21) International Application Number: PCT/US95/03572</p> <p>(22) International Filing Date: 21 March 1995 (21.03.95)</p> <p>(30) Priority Data: 08/210,965 21 March 1994 (21.03.94) US</p> <p>(71) Applicant: UNIVERSITY OF FLORIDA [US/US]; 186 Grinter Hall, Gainesville, FL 32611 (US).</p> <p>(72) Inventors: STIMAC, Jerry, L.; Building 970, Hull Road, Gainesville, FL 32611 (US). PEREIA, Roberto; 4503 N.W. 44th Place, Gainesville, FL 32606 (US).</p> <p>(74) Agents: SALIWANCHIK, David, R. et al.; Saliwanchik &amp; Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville, FL 32606-6669 N.W. (US).</p>	<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p><b>Published</b> <i>With international search report.</i></p>	
<p>(54) Title: METHODS AND FORMULATIONS FOR CONTROL OF INSECT PESTS</p>		
<p>(57) Abstract</p>		
<p>The subject invention concerns novel biopesticides and their use to control cockroaches, carpenter ants, and pharaoh ants. Specifically, highly virulent isolates of <i>Beauveria bassiana</i> in an agricultural composition, can be used to effectively control these pests. Exemplified are <i>Beauveria bassiana</i> No. 447, ATCC 20872, and <i>Beauveria bassiana</i> SP111, ATCC 74038. Also described are unique formulations which are highly effective for delivering biocontrol agents to target pests. By using these novel compositions, target pests can be controlled without the environmental and public safety hazards presented by chemical control agents.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AT</b>	Austria	<b>GB</b>	United Kingdom	<b>MR</b>	Mauritania
<b>AU</b>	Australia	<b>GE</b>	Georgia	<b>MW</b>	Malawi
<b>BB</b>	Barbados	<b>GN</b>	Guinea	<b>NE</b>	Niger
<b>BE</b>	Belgium	<b>GR</b>	Greece	<b>NL</b>	Netherlands
<b>BF</b>	Burkina Faso	<b>HU</b>	Hungary	<b>NO</b>	Norway
<b>BG</b>	Bulgaria	<b>IE</b>	Ireland	<b>NZ</b>	New Zealand
<b>BJ</b>	Benin	<b>IT</b>	Italy	<b>PL</b>	Poland
<b>BR</b>	Brazil	<b>JP</b>	Japan	<b>PT</b>	Portugal
<b>BY</b>	Belarus	<b>KE</b>	Kenya	<b>RO</b>	Romania
<b>CA</b>	Canada	<b>KG</b>	Kyrgystan	<b>RU</b>	Russian Federation
<b>CF</b>	Central African Republic	<b>KP</b>	Democratic People's Republic of Korea	<b>SD</b>	Sudan
<b>CG</b>	Congo	<b>KR</b>	Republic of Korea	<b>SE</b>	Sweden
<b>CH</b>	Switzerland	<b>KZ</b>	Kazakhstan	<b>SI</b>	Slovenia
<b>CI</b>	Côte d'Ivoire	<b>LI</b>	Liechtenstein	<b>SK</b>	Slovakia
<b>CM</b>	Cameroon	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>CN</b>	China	<b>LK</b>	Sri Lanka	<b>TD</b>	Chad
<b>CS</b>	Czechoslovakia	<b>LV</b>	Latvia	<b>TG</b>	Togo
<b>CZ</b>	Czech Republic	<b>MD</b>	Republic of Moldova	<b>TJ</b>	Tajikistan
<b>DE</b>	Germany	<b>MG</b>	Madagascar	<b>TT</b>	Trinidad and Tobago
<b>DK</b>	Denmark	<b>ML</b>	Mali	<b>UA</b>	Ukraine
<b>ES</b>	Spain	<b>MN</b>	Mongolia	<b>US</b>	United States of America
<b>FI</b>	Finland			<b>UZ</b>	Uzbekistan
<b>FR</b>	France			<b>VN</b>	Viet Nam
<b>GA</b>	Gabon				

DESCRIPTION

Methods and formulations for control of insect pests

5

Cross-Reference to Related Applications

This application is a continuation-in-part of co-pending application Serial No. 07/999,185, filed December 30, 1992, which is a continuation of application Serial No. 07/687,362, filed April 18, 1991, now abandoned. This application is also a continuation-in-part of co-pending application Serial No. 07/999,186, filed December 30, 1992, which is a continuation of application Serial No. 07/687,361, filed April 18, 1991, now abandoned.

10

Background of the Invention

15

The development of biological control agents as alternatives to chemical insecticides for the control of important pest species is a subject of increasing interest. Concerns for the environment and exposure of man to harmful substances in air, food and water have stimulated legislation and restrictions regarding the use of chemical pesticides, particularly for pests found in the urban environment. Control of insect pests in urban areas is highly desirable but exposure to chemical pesticides in the household and from lawns and gardens is of great concern to the public. If given a choice, most people would use a non-toxic biological control rather than a toxic chemical to control insects in the urban environment. The problem is that very few biological alternatives to chemical insecticides are available for purchase and use by the public.

20

25

For most insect pests that need to be controlled in the urban environment (ants, roaches, termites, fleas, wasps, etc.) there is no biological agent available for purchase as a product.

30

Cockroaches are serious economic pests in urban areas. Because cockroaches are so closely associated with humans and commonly feed on decaying food, crumbs, or scraps, and frequent unsanitary areas such as sewage systems and septic tanks, their presence leads to suspicion of a threat to human health. Pathogenic organisms have

been isolated from cockroaches collected in domestic or peridomestic environments; however, the role of cockroaches as vectors of pathogens is controversial. Unlike many blood-feeding arthropods whose feeding behavior results in the direct transmission of pathogens to humans, cockroaches have the potential to transmit pathogens indirectly via contamination of foods or utensils used to prepare food. It has been demonstrated that cockroaches acquire pathogenic bacteria simply by walking over cultures and showed that these pathogens are subsequently transferred to foodstuffs via the normal foraging behavior of the infested cockroaches. Aside from bacterially caused food poisoning and diseases such as typhoid and dysentery, many other human illnesses and diseases associated with microorganisms isolated from cockroaches have been reported. These include paralytic polio, giardiasis, otomycosis, pneumomycosis, and various worms such as hookworm and tapeworm.

Besides the possible role of cockroaches as vectors of pathogenic microorganisms, the mere presence of these insects is known to contribute to human morbidity in other ways. Perhaps the most insidious aspect is the psychological impact of these pests in terms of the anxiety and stress related to infestation, which in some instances can take on pathologic dimensions. Further, defensive secretions among cockroach species may cause burning sensations, vertigo, or nausea in individuals who come into contact with the insects.

Current cockroach control methods in buildings include preventative and corrective approaches. Preventative measures emphasize sanitation to eliminate harborages and food sources, sealing off access routes, and the creation of inhospitable environments by the application of boric acid or sorptive dusts in wall voids during construction (Ebeling, W. [1971] *Ann. Rev. Entomol.* 16:123-158; Ebeling, W. [1978] *Urban Entomology*, Berkeley: Univ. Calif. Div. Agric. Sci. 695 pp.). However the implementation of these measures is difficult and thus limits their effectiveness (Thoms, E.M., W.H. Robinson [1987] *J. Econ. Entomol.* 80:131-135). Corrective measures used to suppress established infestations emphasize the use of insecticide applications. A commonly used technique is to spray insecticides with long residual activity in areas frequented by cockroaches at fixed time intervals (Schal, C., R.L. Hamilton [1990] *Ann. Rev. Entomol.* 35:521-551). Despite short term suppression of cockroach populations, toxic residues and the development of insecticide resistance

(Cochran, D.G. [1989] *J. Econ. Entomol.* 82:336-341) make total reliance on this technique undesirable. Alternative corrective measures such as the placement of toxic bait traps may provide sufficient control under proper conditions (Thoms & Robinson [1987], *supra*).

5           The use of natural enemies for the biological control of cockroaches has been examined to varying degrees. Although traps using biocontrol agents have been proposed, these traps are only as good as the biocontrol agent used. U.S. Patent Nos. 5,057,315 and 5,057,316. Field releases of parasitoids of the American and brown  
10           banded cockroaches resulted in rates of parasitism as high as 95% and has generated some optimism for their potential utilization (Coler, R.R., Van Driesche, R.G., Elkinton, J.S. [1984] *Environ. Entomol.* 13:603-606; Hagenbuch, B.E., R.S. Patterson, P.G. Koehler [1989] *J. Econ. Entomol.* 82:90-94). Pathogenic yeasts isolated from laboratory cockroach colonies also have been suggested as possible biological control agents, but more research is required to evaluate their potential (Archbold, E.F., M.K.  
15           Rust, D.A. Reiersen [1987] *J. Med. Entomol.* 24:269-272; Archbold, E.F., M.K. Rust, D.A. Reiersen, K.D. Atkinson [1986] *Environ. Entomol.* 15:221-226; Verrett, J.M., K.B. Green, L.M. Gamble, F.C. Crochen [1987] *J. Econ. Entomol.* 80:1205-1212). Numerous other fungi, bacteria, protozoans, and nematodes have been reported to be associated with cockroaches, but their potential as biological control agents is not  
20           significant, or has not been fully evaluated (Roth and Willis [1960] *Smithsonian Misc. Coll.* Vol. 141; Tsai, Y.H., K.M. Cahill [1970] *J. Parasitol.* 56:375-377; Zervos, S. [1983] *N.Z. J. Zool.* 10:329-334; Rahmet-Alla, M., A.F. Rowley [1989] *J. Invert. Path.* 53:190-196). Thus, there is a significant and long-felt need for a more effective and safe means for controlling cockroaches.

25           Carpenter ants, *Camponotus* spp., are distributed throughout North America. Some of the more common and/or studied species include *C. modoc* in the Pacific northwest, *C. clarithorax* in southern California, and the *C. floridanus* in Florida. *C. pennsylvanicus*, *C. noveboracensis*, and *C. abdominalis*, are found in the east (Ebeling, W. [1978] *Urban Entomology*, Univ. Calif.: Berkeley p. 209-213). Public concern  
30           over carpenter ants has been increasing due to the greater probability of structural infestations as suburban developments extend into the forest habitats of the ants.

Pestiferous species of carpenter ants may be considered nuisance pests because of their foraging activity inside homes. More significant damage occurs when carpenter ants extend their nests into sound wood. Nesting sites may be located in live and dead trees, sometimes resulting in damage to shade trees. Nests may also be established in walls and support beams of structures, or in voids within doors, walls, and furniture. Preference for moist or decaying wood has been reported, but nesting sites are not restricted to such areas. Carpenter ant populations develop relatively slowly with colonies of 300-2,000 workers being produced over a 2-year or longer period for various species. The presence of reproductives follows this slow development since their production has been reported only from well established colonies (Hansen, L.D., R.D. Akre [1985] "Biology of carpenter ants in Washington state (Hymenoptera: Formicidae: *Camponotus*)," *Melandria* 43: 62 pp.; Pricer, J.L. [1908] *Biol. Bull.* 14:177-218). Despite the slow colony growth, large colonies with satellite colonies have been found. Worker movement occurs between the main colony and the satellites, which serve as areas for further brood development and colony expansion (Hansen and Akre [1985], *supra*).

Current methods for controlling structural infestations of carpenter ants include sanitation of potential and current nest sites, minimizing access to structures (eg. preventing the contact of tree branches with a structure), and the application of insecticides to repel (perimeter spray barriers) and/or eliminate carpenter ants. The use of boric acid dust in dry, wall voids is reported to be effective for up to 20 years (Hansen and Akre, *supra*).

Recommendations for the chemical control of established structural infestations in the home are often accompanied with warnings of possible hazards to the applicator as well as children and pets. Alternative control methods such as effective biological control agents have not been found (Akre, R.D., L.D. Hansen, A.L. Antonelli [1989] *Ext. Bull. Washington State Univ. Coop. Ext. Serv.* 1989 rev. no. EB 0818, 6 pp.). A need clearly exists for a safe, effective biological control agent for carpenter ants.

Pharaoh ants, *Monomorium pharaonis*, have been described as ". . . the most persistent and difficult of all our house-infesting ants to control or eradicate" (Smith, M.R. [1965] *USDA-ARS Tech. Bull.* No. 1326, 105 pp.). It is a tropical species which has extended its range to more temperate regions by establishing colonies in

heated buildings. Pharaoh ants frequently infests buildings where food is prepared, and have been found to carry pathogenic organisms (Beatson, S.H. [1972] *Lancet* 1:425-427).

5 The difficulty in controlling pharaoh ants may be attributed to their inaccessible nesting sites, rapid population growth, and dispersion of colonies. Their small size allows establishment of colonies in any suitable location, including unusual places such as between books and in stored clothing. With multiple queen colonies, and the warm (30°C), humid (63-80% RH) conditions that favor pharaoh ants, large colonies can develop rapidly. Portions of these large colonies may disperse to form new colonies at any time, probably in response to overcrowding and unfavorable microenvironmental conditions. Unlike other ant species, pharaoh ants do not exhibit intercolony aggression. This permits the adoption of ants from other colonies and may further enhance the establishment of new colonies and reinfestations. Pharaoh ants also forage for food more than 35 m from the nest without distinct trail following, and thus make nests difficult to find and eradicate.

15 Control methods for pharaoh ants emphasize the use of insect growth regulators (IGR) or toxicants incorporated into baits. Properly implemented bait programs are effective, however it may take over a month to achieve control. Insecticide applications, while fast acting, usually do not eliminate colonies, and may be unacceptable in certain areas where toxic residues are a concern. In addition, insecticide applications are generally not compatible with bait programs. A need exists for safe and effective biological control agents for pharaoh ants.

20 A United States patent has been granted for a fungus showing high activity against fire ants, U.S. Patent No. 4,925,663. This isolate, designated *Beauveria bassiana* isolate No. 447, was deposited in a public repository. No biological activity other than the activity against fire ants had been previously reported for this isolate, nor could activity against other pests be inferred from the mere knowledge that the isolate was active against fire ants. The subject invention concerns the new uses of *B. bassiana* No. 447.

30

### Brief Summary of the Invention

The subject invention concerns the use of highly virulent *Beauveria bassiana* isolates to control certain pests, including cockroaches, carpenter ants, fire ants, and pharaoh ants. Specifically exemplified herein are formulations containing *B. bassiana* isolates No. 447 and SP111. These isolates, advantageously, show unexpectedly high virulence against certain pests, including cockroaches, carpenter ants, fire ants, and pharaoh ants, and do not produce the environmental hazards associated with chemical control agents. The fungal biopesticides described herein can be applied to each of these pests in any of their normal habitats. The fungus may be applied, for example, directly to the pests, in trays, or applied to their surroundings, or anywhere that these pests are a problem. The subject invention also includes mutants of the exemplified isolates which substantially retain the high virulence of the parent strain.

A further aspect of the subject invention concerns unique formulations which can be used to effectively deliver biocontrol agents to target pests. In a preferred embodiment a biocontrol agent is delivered in a formulation which is readily foraged by the target pest and adheres to the body of the pest. Specifically exemplified herein is a formulation which has been discovered to be non-repulsive to fire ants and other pests. This discovery is quite unexpected because pests are known to be repelled by many formulations of microbial agents. The formulation of the subject invention is particularly advantageous because it has been found to be highly effective in delivering the biocontrol agent to the target pest. The formulation of the subject invention comprises a unique blend of a food source and the fungal biocontrol agent. In a preferred embodiment, a drying agent is also used. These ingredients are presented as a dry powder.

25

### Brief Description of the Drawings

**Figure 1** shows cumulative percent mortality of carpenter ants exposed to *B. bassiana* isolate No. 447.

**Figure 2** shows cumulative percent mortality of carpenter ants exposed to *B. bassiana* isolate SP111.

**Figure 3** shows cumulative percent mortality of pharaoh ants exposed to *B. bassiana* isolate No. 447.



Figure 4 shows cumulative percent mortality of German cockroaches exposed to *B. bassiana* isolate No. 447.

Figure 5 shows cumulative percent mortality of American cockroaches exposed to *B. bassiana* isolate No. 447.

5 Figure 6 shows cumulative percent mortality of German cockroaches exposed to *B. bassiana* isolate SP111.

Figure 7 shows mortality of fire ants from the fungal formulation in comparison to chemicals for commercial traps.

10 Figure 8 shows a comparison of chemical baits from traps to fungus formulation for the control of pharaoh ants.

Figure 9 shows a comparison of chemical baits from traps to fungus formulation for the control of crazy ants.

Figure 10 shows a comparison of chemical baits from traps to fungus formulation for the control of carpenter ants.

15 Figure 11 shows a comparison of field pesticides to fungus formulation for the control of fire ants.

#### Detailed Disclosure of the Invention

20 The subject invention concerns the use of fungal biocontrol agents to control certain pests. Specifically exemplified herein is the use of *Beauveria bassiana* isolates No. 447 and SP111. *B. bassiana* SP111 is a novel isolate. A further aspect of the subject invention includes formulations which are highly effective in delivering the biocontrol agent to the target pest.

25 Biologically pure cultures of *Beauveria bassiana* No. 447 and *Beauveria bassiana* SP111, have been deposited in the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, MD 20852. The deposit information and accession numbers are as follows:

<u>Culture</u>	<u>Accession Number</u>	<u>Deposit Date</u>
<i>Beauveria bassiana</i> No. 447	ATCC 20872	December 29, 1987
30 <i>Beauveria bassiana</i> SP111	ATCC 74308	March 5, 1991

The subject cultures have been deposited under conditions that assure that access to the cultures will be available during the pendency of this patent application to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 CFR 1.14 and 35 U.S.C. 122. The deposits are available as required  
5 by foreign patent laws in countries wherein counterparts of the subject application, or its progeny, are filed. However, it should be understood that the availability of the deposits does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

Further, the subject culture deposits will be stored and made available to the  
10 public in accord with the provisions of the Budapest Treaty for the Deposit of Microorganisms, i.e., they will be stored with all the care necessary to keep them viable and uncontaminated for a period of at least five years after the most recent request for the furnishing of a sample of a deposit, and in any case, for a period of at least thirty (30) years after the date of deposit or for the enforceable life of any patent  
15 which may issue disclosing the cultures. The depositor acknowledges the duty to replace the deposit(s) should the depository be unable to furnish a sample when requested, due to the condition of a deposit. All restrictions on the availability to the public of the subject culture deposits will be irrevocably removed upon the granting of a patent disclosing them.

20 The entomopathogenic fungus *Beauveria bassiana* is an imperfect fungus (Fungi Imperfecti) in the subdivision Deuteromycotonia. The genus *Beauveria* Vuill is within the Class Deuteromycetes and is distinguished from other genera by having conidia that are borne singly, not catenulate and having the fertile portion of the conidiophore zig-zag in shape and drawn out at the tip. The species *Beauveria*  
25 *bassiana* has spherical, not ellipsoid, conidia measuring 2-3  $\mu\text{m}$  by 2-2.5  $\mu\text{m}$  and with conidiophores forming dense bunches.

For a biological control agent to be effective at a practical level to control cockroaches, carpenter ants, and pharaoh ants, it is essential that the agent not only exhibit pathogenicity against these pests, but it must also be virulent. The more  
30 virulent it is, the better it is as a biocontrol agent. Though some fungal isolates have been shown to have some pathogenicity to these pests, these isolates did not have the essential virulence to function as a biocontrol agent. There is no known way to

convert a pathogenic non-virulent fungal isolate into a pathogenic virulent isolate. Thus, the discovery of the novel isolate of the invention accomplishes a goal which has long been sought after.

Mode of action and virulence. Like most entomogenous fungi, *Beauveria bassiana* initiates infection by a germinating spore (conidium) attaching to and subsequently penetrating the cuticle of the insect host. Advantageously, and unexpectedly, the claimed *Beauveria bassiana* isolates attach very securely to the cuticle of cockroaches and ants and are typically not removed by the insect's grooming activities. This may account somewhat for the high virulence of the fungus. As the fungus penetrates the insect's cuticle, the invasive hyphae begin to enter the host's tissues and ramify through the hemocoel. Hyphal bodies or segments of the hyphae distribute throughout the hemocoel, filling the dying insect with mycelia. Emergence hyphae grow out through the insect's integument and produce spores on the external surface of the host. These spores, or conidia, are dispersed and capable of infecting new host insects. *B. bassiana* spores can be dispersed within the nest by the activities of the pests.

Formulations. The formulations of the subject invention were found to be particularly effective for the control of fire ants and other pests. In a preferred embodiment, the formulation comprises a dry powder having the fungal biocontrol agent and a food component. Preferably, the formulation further comprises a drying agent. Optionally, the formulation may also comprise an attractant. The preferred formulation is non-repellant and includes a food source so that the target pest will forage and recruit other nestmates for foraging activity. Furthermore, the formulation of the subject invention has been found to advantageously adhere to the body of the target pest, thereby facilitating colonization of the pest by the fungal biocontrol agent. The ability to adhere to the pest makes the formulation of the subject invention quite distinct from other formulations which are currently used to administer chemical pesticides.

In one embodiment, the formulation of the subject invention consists of about 25-40% peanut material, about 45-60% cornstarch, about 2-20% fungal biocontrol agent, and about 0-15% drying agent. In a specific embodiment, the formulation can comprise about 35% peanut material, about 50% cornstarch, about 5% drying agent,

and about 10% fungus. The drying agent can be any one of many materials known to those skilled in the art which are small particles but have a high surface area to volume ratio so as to effectively remove water or oils from the formulation to create a dry powder. Preferably, the drying agent can be diatomaceous earth or a synthetic calcium silicate such as Micro-Cel®.

The peanut component of the formulation is preferably prepared by grinding roasted peanuts so as to obtain a powder. To achieve a dry powder, it is best to grind the peanuts together with the cornstarch and/or a drying agent. Preferably, the components of the formulation are small particles and will pass through a 60 mesh sieve. Typically, the cornstarch and drying agent will pass through a much smaller sieve, such as at 300 mesh. Preferably, the formulation is a powder which is free flowing and does not stick together in clumps. Food sources other than peanut material or cornstarch can also be used according to the subject invention. The choice of a food source will depend upon the particular pest which is the target for control. Also, various attractants known to those skilled in the art can be used. These attractants can be, for example, pheromones or various extracts.

In a preferred embodiment, the fungal pathogen is *B. bassiana* No. 447 or *B. bassiana* SP111. However, other microbes can be used as can other biocontrol agents. For example, *Bacillus thuringiensis* can also be used with the formulation of the subject invention.

To evaluate the control achieved using the materials and methods of the subject invention, tests were conducted to compare the control of pests achieved with certain commercial pesticides. As described below, these tests demonstrated that the fungal formulations of the present invention are highly effective in controlling pests.

Following are examples which illustrate procedures, including the best mode, for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

#### Example 1 – Preparation of the Fungus

The subject fungus can be produced in trays with a rice-based medium. An isolate of fungal inoculum is used to initiate the growth of the fungus in the trays.

The initial inoculum is prepared in petri dishes. The pure spores are then transferred into jars containing sterile white rice without skins.

The medium for the trays is prepared as follows:

1. The rice is pre-cooked for 10 minutes.
- 5 2. 750 grams of cooked rice is placed in polyethylene bags and sterilized in an autoclave at 120°C for 30 minutes.
3. Within a laminar flow hood, one teaspoon of spores and rice from the inoculum jars is added to each bag of prepared sterile medium.
4. Each bag is closed tightly by folding and stapling the open end.
- 10 5. The bags are transferred to a sterile room with positive pressure, temperature at 25.0-27.0°C, relative humidity above 70%, and 16 hours photophase. This room is known as the "environment room."

After 3 days in the environment room, bags containing mycelia are selected  
15 and their contents are transferred to plastic trays. The size of the trays is such that each tray will accommodate the contents of 2-3 bags. The trays and their contents are left in the environment room for 12-15 days. At the end of the 12-15 day period, the trays are transferred to a room with a cool (10-20°C) current of clean air. The trays are left in this room until the cool air has dried the rice and fungus mixture.

20 The uncontaminated trays of rice covered with fungus can be harvested and prepared for application or storage. If the fungus will be applied to cockroaches or ants within 1-2 weeks after production, conidia can be collected by shaking and sieving. The resulting powder contains spores and some mycelia, and can be applied directly to target insects or used to prepare a formulation as a liquid, powder, or bait.

25 If the fungus is to be stored, the mixture can be mixed with cornstarch or talc and placed into sterile plastic containers sealed tightly and stored in a refrigerator at 4°C or in a room with a temperature range of 10-25°C and no direct sunlight. The high virulence of *B. bassiana* can be compromised by bacterial or fungal contamination. Therefore, throughout the preparation of the fungus, great care must  
30 be taken to maintain the sterility of all instruments and equipment.

As the following examples demonstrate, the fungus-containing product can be applied to target pests and their nests as a liquid, powder, or put out as a baited trap for the pests to forage, become infected, and carry inoculum back to the nest.

5     Example 2 — Spray Application

Spraying can be used for treating individual ants or cockroaches or small groups of these pests. A fungal suspension containing  $1.0 \times 10^7$  to  $1.0 \times 10^9$  spores per milliliter of water can be sprayed on the target pests using an airbrush or other means as an applicator.

10

Example 3 — Powder Application

A fungal spore and mycelia mixture can be mixed with cornstarch or talc and applied to the pests' surroundings as a dry powder.

15     The powder is prepared as in Example 1 above. The sieved *B. bassiana* powder which contains the rice, spores, and mycelia is mixed with cornstarch or talc. Application of this powder to the nests or directly to the pests can facilitate rapid and widespread fungal growth within the nest or on the pest.

20     The application can be accomplished using an pressurized air applicator with an attachment that distributes the mixtures into cracks and crevices of a pest-inhabited building. During and following application, pests covered with white powder will be observed. These infected pests will die within 1-5 days, and the spores they produce will be infective to other pests. There should be a marked decrease in activity within 1-3 days and death should occur within 1-2 weeks following application. Active spores will remain in the surroundings at the nest site, thereby providing inoculum to  
25     infect other roaches or ants.

Example 4 — Baited Trap Application

30     The fungal powder can be used in a trap in which entryways are laced with fungal inoculum. Preferably, fungal spores are utilized. A bait attractant contained within the trap will be foraged by cockroaches or ants and the foragers will become infected. These infected individuals will return to the nest contaminated and thereby introduce the fungal disease into the nest. A vegetable oil or other liquid substance

can be added to a bait in the trap to make it more attractive to the pests. Various attractants, including pheromone compounds, are well known to those skilled in this art. The baited traps should be placed in cabinets, along baseboards, windowsills, etc. A quantity of 0.5-2.0 grams of fungal mixture containing spores and mycelia should  
5 be contained in each trap. The number of traps used in an area will depend on the level of infestation.

Example 5 – Treatment of Carpenter Ants With *B. bassiana* No. 447

Carpenter ants (*Camponotus floridanus*), were exposed *Beauveria bassiana* No.  
10 447. Each treatment entailed exposing two groups of 50 ants each to conidia of the isolates. Ants were coated with a conidia/cornstarch mixture, by gently shaking the ants and spores together in a covered container. The control treatment consisted of cornstarch only. Ants were subsequently held in open plastic boxes that contained a nest cell (100 mm covered petri dish with the bottom dish filled with plaster that was  
15 periodically moistened with water) and honey water for food.

Mortality was recorded daily for 18 days beginning with the second day after exposure. The test was terminated after 28 days. Dead ants were individually held under high humidity and examined for sporulation to determine infection rates.

Carpenter ants exposed to isolates of *B. bassiana* sustained over 95% mortality  
20 (Figure 1). At least 49% of the dead ants developed sporulating bodies of the fungi to which they were exposed, indicating that these isolates can grow and reproduce on carpenter ants.

Example 6 – Treatment of Carpenter Ants With *B. bassiana* SP111

Carpenter ants (*Camponotus floridanus*), were exposed *Beauveria bassiana*  
25 SP111. Each treatment entailed exposing two groups of 50 ants each to conidia of the isolates. Ants were coated with a conidia/cornstarch mixture, by gently shaking the ants and spores together in a covered container. The control treatment consisted of cornstarch only. Ants were subsequently held in open plastic boxes that contained a nest cell (100 mm covered petri dish with the bottom dish filled with plaster that was  
30 periodically moistened with water) and honey water for food.

Mortality was recorded daily for 18 days beginning with the second day after exposure. The test was terminated after 28 days. Dead ants were individually held under high humidity and examined for sporulation to determine infection rates.

5 Carpenter ants exposed to *B. bassiana* SP111 sustained over 75% mortality (Figure 2). At least 49% of the dead ants developed sporulating bodies of the fungi they were exposed to, indicating that these isolates can grow and produce spores on carpenter ants.

Example 7 – Treatment of Pharaoh Ants With *B. bassiana* No. 447

10 Pharaoh ants were exposed to a mixture comprising *B. bassiana* No. 447 conidia as the active ingredient. Three colonies of approximately 100-200 ants were individually dusted with the conidia in a petri dish and allowed to crawl out into a nest cell (15 x 40 mm plastic petri dish with a plaster filled base and entrance holes in the lid). Controls consisting of three colonies were not dusted. The ant colonies  
15 were held separately in larger petri dishes along with the nest cells and honey water. Mortality was recorded daily for 25 days. Dead ants were individually surface sterilized and held under high humidity to the rate of infection.

Pharaoh ant exposure to *B. bassiana* 447 resulted in 90% mortality after 8 days (Figure 3). Furthermore, all of the dead ants were confirmed to have fungal spores,  
20 indicating that the fungus can successfully develop on pharaoh ants.

Example 8 – Treatment of Cockroaches With *B. bassiana* No. 447

For the German cockroach, *Blattella germanica*, *B. bassiana* No. 447 was tested for ability of its conidia to infect and kill the host. Groups of 50 male cockroaches  
25 were anesthetized with CO<sub>2</sub> and then dusted with conidia, within a covered container (8 oz). Controls consisted of a group of 20 cockroaches. Dusted cockroaches were transferred individually into separate petri dishes (10 x 35mm) containing moistened filter paper. Mortality was recorded from the second day after conidia application and daily thereafter. Dead cockroaches were individually held in a humidity chamber for  
30 10 days to identify sporulating fungi.



For the American cockroach, *Periplaneta americana*, *B. bassiana* No. 447 was applied by brushing the conidia onto anesthetized cockroaches. Cockroaches were then held in petri dishes as described above, at 26°C.

5 In the test with German cockroaches the fungal isolate *B. bassiana* No. 447 (Figure 4) caused 100% mortality after contact with spores. Sporulation of the fungus was evident on 82% of the dead cockroaches. For the exposures of the American cockroaches, the *B. bassiana* isolate caused 90% or more mortality after 8 days (Figure 5). Fungal sporulation occurred on all of the dead American cockroaches.

10 Example 9 — Treatment of Cockroaches With *B. bassiana* SP111

For the German cockroach, *Blattella germanica*, *B. bassiana* SP111 was tested for ability of its conidia to infect and kill the host. Groups of 50 male cockroaches were dusted with conidia, within a covered container (8 oz). Controls consisted of a group of 20 cockroaches. After exposure to conidia, cockroaches were anesthetized  
15 with CO<sub>2</sub> and transferred individually into separate petri dishes (10 x 35mm) containing moistened filter paper. Mortality was recorded beginning on the second day after conidia application and daily thereafter. Dead cockroaches were individually held in a humidity chamber for 10 days to identify sporulating fungi.

In the test with German cockroaches the fungal isolate *B. bassiana* SP111  
20 (Figure 6) caused 100% mortality after contact with spores. Sporulation of the fungus was evident on about 82% of the dead cockroaches.

Example 10 — Evaluation of *B. bassiana* Bait Formulations for Control of Fire Ants

Bait formulations with ground peanut material were offered to colonies of fire  
25 ant workers in plastic boxes (≈20 x 12 x 10 cm) containing a small dish of water and a plastic petri dish (60 mm diameter) to serve as a nest cell. Colonies were established 2-4 days before the start of the experiment to allow ants to adapt to their new environment. The formulations (0.5 g) were offered on weigh papers (1 square inch) or small dishes and left in the boxes for 3-4 days. Two controls were used:  
30 a clean control which received no formulation but only water, and a bait control which received the bait formulation without any fungus. Abbott-corrected mortality greater

than 70% at 14 days after treatment was observed for the fungal isolate containing about 10% of the *B. bassiana* No. 447.

5 Example 11 – Chemical Baits from Traps Compared to Fungal Formulations for the Control of Fire Ants

The chemical baits which were compared are from: MAX ant trap, RAID ant trap, and COMBAT ant trap. The chemical baits were removed from the traps and offered to ants on paper. The control received the same formulation as the fungus treatment but without conidia. The fungal formulation contained peanut material and cornstarch and 10% conidia of *B. bassiana* No. 447. MAX and the fungal formulations had similar mortality, although MAX caused mortality to increase much more rapidly than the fungus as expected, since the fungus requires 3-4 days to infect and kill the insect. As shown in Figure 7, COMBAT and RAID were less efficient than MAX and *Beauveria bassiana*.

15

Example 12 – Chemical Baits from Traps Compared to Fungal Formulation for the Control of Pharaoh Ants

The chemical baits which were compared are from: MAX ant trap, RAID ant trap, and COMBAT ant trap. The chemical baits were removed from the traps and offered to ants on paper. The control received the same formulation as the fungus treatment but without conidia. The fungal formulation contained peanut material and cornstarch and 10% conidia of *B. bassiana* No. 447. The results of this experiment is shown in Figure 8.

20

25 Example 13 – Chemical Baits from Traps Compared to Fungal Formulations for the Control of Crazy Ants

The chemical baits which were compared are from: MAX ant trap, RAID ant trap, and COMBAT ant trap. The chemical baits were removed from traps and offered to ants on paper. The control received the same formulation as the fungus treatment but without conidia. The fungal formulation contained peanut material and cornstarch and 10% conidia of *B. bassiana* No. 447. In all experiments, the bait with *Beauveria*

30

*bassiana* caused mortality similar or greater than that caused by chemical baits. See Figure 9 for the results of one such experiment.

5 Example 14 — Chemical Baits from Traps Compared to Fungal Formulation for Control of Carpenter Ants

The chemical baits which were compared are from: MAX ant trap, RAID ant trap, and COMBAT ant trap. The chemical baits were removed from traps and offered to ants on paper. The control received the same formulation as the fungus treatment but without conidia. The fungal formulation contained peanut material and cornstarch and 10% conidia of *B. bassiana* No. 447. As can be seen from Figure 10, the fungal formulation has performance similar to, or slightly better than, MAX and RAID baits, and only slightly less than COMBAT. Both COMBAT and the fungus had their effects delayed in relation to RAID and MAX, but the delay in fungal effect is longer than that of COMBAT.

15

Example 15 — Field Pesticides Compared to Fungal Formulations for the Control of Fire Ants

AMDRO® chemical baits were evaluated. The bait in one experiment also had 10% of an ACEPHATE fire ant powder insecticide. In each treatment, one-half gram of formulation was provided per arena on weighing paper. The formulation was removed after 4 days. The control received the same formulation as the fungus treatment but without conidia. The fungal formulation contained peanut material, cornstarch, and 10% conidia of *B. bassiana* No. 447.

25 ACEPHATE (which is not normally a bait formulation) kills the ants almost immediately. Mortality with AMDRO increases less rapidly, but by day 4, ~80% of the population was dead. Mortality with fungus increases at a slower rate but final mortality after 2-4 weeks is similar to that obtained with the chemical pesticides (Figure 11).

30

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and the scope of the appended claims.

Claims

1           1. A process for controlling cockroaches, said process comprising applying a  
2 *Beauveria bassiana* onto said cockroaches, or their surroundings, wherein said  
3 *Beauveria bassiana* is selected from the group consisting of *Beauveria bassiana* No.  
4 447, having the characteristics of ATCC 20872, and *Beauveria bassiana* SP111, having  
5 the characteristics of ATCC 74038.

1           2. The process, according to claim 1, wherein said *Beauveria bassiana* is  
2 isolate No. 447, having the characteristics of ATCC 20872.

1           3. The process, according to claim 1, wherein said *Beauveria bassiana* is  
2 isolate SP111, having the characteristics of ATCC 74038.

1           4. A process for controlling carpenter ants, said process comprising applying  
2 a *Beauveria bassiana* onto said carpenter ants, or their surroundings, wherein said  
3 *Beauveria bassiana* is selected from the group consisting of *Beauveria bassiana* No.  
4 447, having the characteristics of ATCC 20872, and *Beauveria bassiana* SP111, having  
5 the characteristics of ATCC 74038.

1           5. The process, according to claim 4, wherein said *Beauveria bassiana* is  
2 isolate No. 447, having the characteristics of ATCC 20872.

1           6. The process, according to claim 4, wherein said *Beauveria bassiana* is  
2 isolate SP111, having the characteristics of ATCC 74038.

1           7. A process for controlling pharaoh ants, said process comprising applying  
2 a *Beauveria bassiana* onto said pharaoh ants, or their surroundings, wherein said  
3 *Beauveria bassiana* is *Beauveria bassiana* No. 447, having the characteristics of ATCC  
4 20872.

1           8. A trap for applying a pesticidal fungus to a target pest, wherein said trap  
2 comprises a pesticidal amount of a *Beauveria bassiana*, wherein said *Beauveria*  
3 *bassiana* is selected from the group consisting of *Beauveria bassiana* No. 447, having  
4 the characteristics of ATCC 20872, and *Beauveria bassiana* SP111, having the  
5 characteristics of ATCC 74038.

1           9. The process, according to claim 8, wherein said *Beauveria bassiana* is  
2 isolate No. 447, having the characteristics of ATCC 20872.

1           10. The process, according to claim 8, wherein said *Beauveria bassiana* is  
2 isolate SP111, having the characteristics of ATCC 74038.

1           11. A *Beauveria bassiana* which, when in its essentially biologically pure  
2 form, has the virulence against cockroaches and carpenter ants characteristic of  
3 *Beauveria bassiana* SP111 culture deposit ATCC 74038, and mutants thereof.

1           12. A composition for the control of a cockroach or ant pest wherein said  
2 composition comprises a food source and a biocontrol agent wherein said composition  
3 is a dry powder capable of passing through a 65 mesh sieve.

1           13. The composition, according to claim 12, which comprises a peanut  
2 material, cornstarch, and a fungal biocontrol agent.

1           14. The composition, according to claim 13, wherein said biocontrol agent is  
2 selected from the group consisting of *Beauveria bassiana* No. 447, having the  
3 characteristics of ATCC 20872, and *Beauveria bassiana* SP111, having the  
4 characteristics of ATCC 74038.

1           15. The composition, according to claim 14, wherein said *Beauveria bassiana*  
2 is isolate No. 447, having the characteristics of ATCC 20872.

1           16. The composition, according to claim 14, wherein said *Beauveria bassiana*  
2 is isolate SP111, having the characteristics of ATCC 74038.

1           17. The composition, according to claim 14, which further comprises a drying  
2 agent.

1           18. The composition, according to claim 14, which comprises about 30-40%  
2 peanut material, about 45-55% cornstarch, about 10% drying agent, and about 10%  
3 *Beauveria bassiana* isolate No. 447.




Fig. 1






Fig. 2




Fig. 3




Fig. 4




Fig. 5

6/11




Fig. 6

Fig. 7





Fig. 8



9/11

Fig. 9






Fig. 10





Fig. 11



**INTERNATIONAL SEARCH REPORT**

International Application No  
PCT/US 95/03572

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A01N63/04 A01N25/00 C12R1/645

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A01N C12R

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 925 663 (J.L.STIMAC) 15 May 1990 cited in the application see examples 3, and, 4	12
Y	see claims 1, 2, 8, 12, 13	8-10
A	---	1-7
Y	WO,A,90 10389 (ECOSCIENCE LABORATORIES INC.) 20 September 1990 cited in the application	8-10
A	see claims	1-7
	--- -/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

\* Special categories of cited documents :

- |  |  |
|--|--|
| <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> | <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> |
|--|--|

Date of the actual completion of the international search

5 July 1995

Date of mailing of the international search report

12.07.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Fletcher, A

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/03572

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 70, no. 11, 17 March 1969, Columbus, Ohio, US; abstract no. 46398j, L.I.BRIKMAN ET AL 'Virulence of entomopathogenic spore-forming microorganisms in relation to the red house ant, Monomorium pharaonis' page 227 ;column 1 ; see abstract & Tr. Tsent. Nauch.-Issled. Dezinfek. Inst., 1967, No. 18(P. 2), 70-77 ----	1-7
X	US,A,4 992 275 (R.W.LUSH) 12 February 1991 see column 3, line 33 - column 4, line 10 col. 5 "Crop = Citrus Orchard Floor" ----	12
A	US,A,5 186 935 (J.W.TUCKER) 16 February 1993 see column 3, line 8 - line 50; example 1 ----	12
A	EP,A,0 190 844 (SCHERING AGROCHEMICALS LTD.) 13 August 1986 see page 1, line 2 - line 24 -----	13

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/03572

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4925663	15-05-90	NONE	
WO-A-9010389	20-09-90	US-A- 5057315	15-10-91
		AU-B- 646388	24-02-94
		AU-A- 4962490	09-10-90
		CA-A- 2046884	16-09-90
		CN-A- 1047608	12-12-90
		EP-A- 0462978	02-01-92
		JP-T- 4504413	06-08-92
		US-A- 5189831	02-03-93
		US-A- 5057316	15-10-91
		US-A- 5310552	10-05-94
		US-A- 5427784	27-06-95
US-A-4992275	12-02-91	NONE	
US-A-5186935	16-02-93	CA-A- 1286219	16-07-91
EP-A-0190844	13-08-86	AU-B- 571121	31-03-88
		AU-A- 5293186	14-08-86
		DE-A- 3682076	28-11-91
		JP-A- 61183201	15-08-86