



(22) Date de dépôt/Filing Date: 2006/09/18
(41) Mise à la disp. pub./Open to Public Insp.: 2007/03/29
(45) Date de délivrance/Issue Date: 2018/06/05
(62) Demande originale/Original Application: 2 622 911
(30) Priorité/Priority: 2005/09/16 (US60/718,170)

(51) Cl.Int./Int.Cl. *A61K 38/07*(2006.01),
A61K 9/00(2006.01), *A61P 15/00*(2006.01),
A61N 1/30(2006.01)

(72) Inventeurs/Inventors:
SZETO, HAZEL H., US;
LIU, SHAOYI, US;
CHO, SUNGHEE, US

(73) Propriétaire/Owner:
CORNELL RESEARCH FOUNDATION, INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : METHODES PERMETTANT DE REDUIRE L'EXPRESSION DE CD36

(54) Title: METHODS FOR REDUCING CD36 EXPRESSION

(57) Abrégé/Abstract:

The invention provides a method of reducing CD36 expression in a cell. The method comprises contacting the cell with an effective amount of an aromatic-cationic peptide having at least one net positive charge; a minimum of four amino acids; a maximum of about twenty amino acids; a relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) wherein $3p_m$ is the largest number that is less than or equal to $r + 1$; and a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) wherein $2a$ is the largest number that is less than or equal to $p_t + 1$, except that when a is 1, p_t may also be 1.

79434-30D5

ABSTRACT

The invention provides a method of reducing CD36 expression in a cell. The method comprises contacting the cell with an effective amount of an aromatic-cationic peptide having at least one net positive charge; a minimum of four amino acids; a maximum of about 5 twenty amino acids; a relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) wherein $3p_m$ is the largest number that is less than or equal to $r + 1$; and a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) wherein $2a$ is the largest number that is less than or equal to $p_t + 1$, except that when a is 1, p_t may also be 1.

79434-30D5

METHODS FOR REDUCING CD36 EXPRESSION

[0001] The application asserts priority to U.S. Provisional Application Serial No. 60/718,170 filed on September 16, 2005. This application is a division of Canadian Application Serial No. 2,622,911 filed September 18, 2006.

5 [0002] The invention described in this application was funded by the National Institute of Drug Abuse, Grant No. P01 DA08924, the National Institute of Neurological Diseases and Stroke, Grant No. R21 NS48295, and the National Heart, Lung and Blood Institute, Grant No. RO1 HL082511. The United States Government has certain rights in this invention.

10 **BACKGROUND OF THE INVENTION**

[0003] CD36 is a transmembrane protein of the class B scavenger receptor family. The protein is widely expressed on numerous cells, such as microvascular endothelium, macrophages, platelets, adipocytes, epithelial cells (e.g., intestinal epithelial and renal tubular cells, etc.), pancreatic islet cells and cardiac muscle. The receptor may interact 15 with multiple extracellular ligands, such as thrombospondin-1, long-chain fatty acids, and oxidized low-density lipoprotein.

20 [0004] Abnormal expression of CD36 has been implicated in a wide variety of diseases and conditions. For example, mice lacking CD36 have less atherosclerotic lesions when fed a Western diet compared to wild-type mice. Further, CD36 knock out mice were reported to be protected against acute cerebral ischemia.

[0005] Therefore, methods for reducing expression of CD36 expression are beneficial for treating a disease or condition characterized by abnormal expression of CD36.

79434-30D5

SUMMARY OF THE INVENTION

[0006] In one embodiment, the invention provides a method for reducing CD36 expression in a cell. The method comprises contacting the cell with an effective amount of an aromatic-cationic peptide.

79434-30D5

[0007] In another embodiment, the invention provides a method for reducing CD36 expression in a mammal in need thereof. The method comprises administering to the mammal an effective amount of an aromatic-cationic peptide.

[0008] In yet another embodiment, the invention provides a method for treating a disease or condition characterized by increased CD36 expression in a mammal in need thereof. The method comprises administering to the mammal an effective amount of an aromatic-cationic peptide.

[0009] In a further embodiment, the invention provides a method for treating ureteral obstruction in a mammal in need thereof. The method comprises administering to the mammal an effective amount of an aromatic-cationic peptide.

[0010] In yet a further embodiment, the invention provides a method for treating diabetic nephropathy in a mammal in need thereof. The method comprises administering to the mammal an effective amount of an aromatic-cationic peptide.

[0011] In another embodiment, the invention provides a method for reducing CD36 expression in a removed organ or tissue. The method comprises administering to the mammal an effective amount of an aromatic-cationic peptide.

[0012] The aromatic-cationic peptides useful in the methods of the present invention have at least one net positive charge; a minimum of four amino acids; a maximum of about twenty amino acids; a relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) wherein $3p_m$ is the largest number that is less than or equal to $r + 1$; and a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) wherein $2a$ is the largest number that is less than or equal to $p_t + 1$, except that when a is 1, p_t may also be 1.

79434-30D5

[0012a]

This invention as claimed relates to:

- a use of the aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ in the manufacture of a medicament for treating ureteral obstruction in a mammal in need thereof;
- use of the aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ for 5 treating ureteral obstruction in a mammal in need thereof; and
- the aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ for use in treating ureteral obstruction in a mammal in need thereof.

BRIEF DESCRIPTION OF THE FIGURES

[0013]

Figure 1. SS-31 reduced oxLDL-induced CD36 mRNA expression, 10 CD36 protein expression, and foam cell formation in mouse peritoneal macrophages.

[0014] Figure 2. SS-31 treatment reduced infarct volume and hemispheric swelling in wild-type mice subjected to acute cerebral ischemia.

[0015] Figure 3. SS-31 treatment reduced the decrease in reduced glutathione (GSH) in post-ischemic brain in wild-type mice.

[0016] Figure 4. SS-31 had no effect in reducing infarct volume or hemispheric swelling in CD36 knock-out mice subjected to acute cerebral ischemia.

[0017] Figure 5. SS-31 did not reduce GSH depletion in post-ischemic brain from CD36 knock-out mice.

[0018] Figure 6. SS-31 reduced CD36 mRNA expression in post-ischemic brain in wild-type mice.

[0019] Figure 7. SS-31 decreases CD36 expression on renal tubular cells after unilateral ureteral obstruction (UUO). Contralateral unobstructed kidney (fig. 7A); obstructed kidney in animals treated with saline (fig. 7B); and obstructed kidneys obtained from rats treated with SS-31 (fig. 7C).

[0020] Figure 8. SS-31 reduces lipid peroxidation in kidney after UUO. Tubular cells in the obstructed kidney (fig. 8B), contralateral unobstructed control (fig. 8A); obstructed kidneys from rats treated with SS-31 (fig. 8C).

[0021] Figure 9. SS-31 reduced tubular cell apoptosis in obstructed kidney after UUO. Obstructed kidney from saline-treated animals (fig. 9B); contralateral unobstructed control (fig. 9A); obstructed kidney from SS-31 treated animals (fig. 9C).

[0022] Figure 10. SS-31 reduced macrophage infiltration in obstructed kidney induced by UUO. Obstructed kidney (fig. 10B); contralateral unobstructed control (fig. 10A); rats treated with SS-31 (fig. 10C).

[0023] Figure 11. SS-31 reduced interstitial fibrosis in obstructed kidney after UUO. Obstructed kidney (fig. 11B); contralateral unobstructed control (fig. 11A); rats treated with SS-31 (fig. 11C).

[0024] Figure 12. Cold storage of isolated hearts with SS-31 or SS-20 prevented upregulation of CD36 expression. The “background” control (figs. 12A and 12B) represents two sections from a normal non-ischemic heart that were not treated with the primary anti-CD36 antibody. “Normal heart” (figs. 12C and 12D) represents two sections obtained from a non-ischemic heart. The sections from a representative heart stored in St. Thomas solution (figs. 12E and 12F) for 18 hours at 4°C showed increased CD36 staining compared to “Normal heart.” CD36 staining was significantly reduced in hearts stored with either 1 nM SS-31 (figs. 12G and 12H) or 100 nM SS-20 (figs. 12I and 12J) in St. Thomas solution.

[0025] Figure 13. SS-31 and SS-20 reduced lipid peroxidation in isolated guinea pig hearts subjected to warm reperfusion after prolonged cold ischemia. HNE staining in hearts subjected to 18 hours of cold storage in St. Thomas solution (fig. 13B) compared to non-ischemic hearts (fig. 13A). HNE staining was reduced in hearts stored in SS-31 (fig. 9C) or SS-20 (fig. 13D).

[0026] Figure 14. SS-31 and SS-20 abolished endothelial apoptosis in isolated guinea pig hearts subjected to warm reperfusion after prolonged cold ischemia. Hearts subjected to 18 hours of cold storage in St. Thomas solution (figs. 14C and 14D); non-ischemic normal hearts (figs. 14A and 14B). Apoptotic cells were not observed in hearts stored in SS-31 (figs. 14E and 14F) or SS-20 (figs. 14G and 14H).

[0027] Figure 15. SS-31 and SS-20 preserves coronary flow in isolated guinea pig hearts subjected to warm reperfusion after prolonged cold ischemia. Guinea pig hearts perfused with a cardioplegic solution (St. Thomas solution) alone or St. Thomas solution containing either 1 nM SS-31 (fig. 15A) or 100 nM SS-20 (fig. 15B) for 3 min. and then subjected to 18 hours of cold ischemia (4°C).

[0028] Figure 16. SS-31 prevented damage to proximal tubules in diabetic mice. Diabetes was induced by streptozotocin (STZ) injection for 5 d. Kidney sections obtained after 3 weeks showed loss of brush border in STZ-treated animals (fig. 16A, panel B) that was not seen in mice not treated with STZ (panel A). The loss of brush border was not seen in STZ-treated animal that received daily SS-31 (3 mg/kg) (panel C).

[0029] Figure 17. SS-31 prevented renal tubular epithelial cell apoptosis in diabetic mice. Diabetes was induced by streptozotocin (STZ) injection for 5d. Kidney sections obtained after 3 weeks showed dramatic increase in apoptotic cells in proximal tubules in STZ-treated animals (fig. 17A, panel b) that was not seen in mice not treated with STZ (fig. 17A, panel a). The STZ-induced apoptosis was not seen in mice that received daily SS-31 (3 mg/kg) (fig. 17A, panel c). The percent of apoptotic cells caused by STZ was significantly reduced by SS-31 treatment (fig. 17B).

DETAILED DESCRIPTION OF THE INVENTION

Peptides

[0030] The invention is directed to the reduction of CD36 expression by certain aromatic-cationic peptides. The aromatic-cationic peptides are water-soluble and highly polar. Despite these properties, the peptides can readily penetrate cell membranes.

[0031] The aromatic-cationic peptides useful in the present invention include a minimum of three amino acids, and preferably include a minimum of four amino acids, covalently joined by peptide bonds.

[0032] The maximum number of amino acids present in the aromatic-cationic peptides of the present invention is about twenty amino acids covalently joined by peptide bonds. Preferably, the maximum number of amino acids is about twelve, more preferably about nine, and most preferably about six. Optimally, the number of amino acids present in the peptides is four.

[0033] The amino acids of the aromatic-cationic peptides useful in the present invention can be any amino acid. As used herein, the term "amino acid" is used to refer to any organic molecule that contains at least one amino group and at least one carboxyl group. Preferably, at least one amino group is at the α position relative to a carboxyl group.

[0034] The amino acids may be naturally occurring. Naturally occurring amino acids include, for example, the twenty most common levorotatory (L) amino acids normally found in mammalian proteins, i.e., alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine

(His), isoleucine (Ileu), leucine (Leu), lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), tryptophan, (Trp), tyrosine (Tyr), and valine (Val).

[0035] Other naturally occurring amino acids include, for example, amino acids that are synthesized in metabolic processes not associated with protein synthesis. For example, the amino acids ornithine and citrulline are synthesized in mammalian metabolism during the production of urea. Another example of a naturally occurring amino acid include hydroxyproline (Hyp).

[0036] The peptides useful in the present invention optionally contain one or more non-naturally occurring amino acids. Optimally, the peptide has no amino acids that are naturally occurring. The non-naturally occurring amino acids may be levorotary (L-), dextrorotatory (D-), or mixtures thereof.

[0037] Non-naturally occurring amino acids are those amino acids that typically are not synthesized in normal metabolic processes in living organisms, and do not naturally occur in proteins. In addition, the non-naturally occurring amino acids useful in the present invention preferably are also not recognized by common proteases.

[0038] The non-naturally occurring amino acid can be present at any position in the peptide. For example, the non-naturally occurring amino acid can be at the N-terminus, the C-terminus, or at any position between the N-terminus and the C-terminus.

[0039] The non-natural amino acids may, for example, comprise alkyl, aryl, or alkylaryl groups not found in natural amino acids. Some examples of non-natural alkyl amino acids include α -aminobutyric acid, β -aminobutyric acid, γ -aminobutyric acid, δ -aminovaleric acid, and ε -aminocaproic acid. Some examples of non-natural aryl amino acids include ortho-, meta, and para-aminobenzoic acid. Some examples of non-natural alkylaryl amino acids include ortho-, meta-, and para-aminophenylacetic acid, and γ -phenyl- β -aminobutyric acid.

[0040] Non-naturally occurring amino acids include derivatives of naturally occurring amino acids. The derivatives of naturally occurring amino acids may, for example, include the addition of one or more chemical groups to the naturally occurring amino acid.

[0041] For example, one or more chemical groups can be added to one or more of the 2', 3', 4', 5', or 6' position of the aromatic ring of a phenylalanine or tyrosine residue, or the 4', 5', 6', or 7' position of the benzo ring of a tryptophan residue. The group can be any chemical group that can be added to an aromatic ring. Some examples of such groups include branched or unbranched C₁-C₄ alkyl, such as methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, or t-butyl, C₁-C₄ alkyloxy (i.e., alkoxy), amino, C₁-C₄ alkylamino and C₁-C₄ dialkylamino (e.g., methylamino, dimethylamino), nitro, hydroxyl, halo (i.e., fluoro, chloro, bromo, or iodo). Some specific examples of non-naturally occurring derivatives of naturally occurring amino acids include norvaline (Nva) and norleucine (Nle).

[0042] Another example of a modification of an amino acid in a peptide useful in the methods of the present invention is the derivatization of a carboxyl group of an aspartic acid or a glutamic acid residue of the peptide. One example of derivatization is amidation with ammonia or with a primary or secondary amine, e.g. methylamine, ethylamine, dimethylamine or diethylamine. Another example of derivatization includes esterification with, for example, methyl or ethyl alcohol.

[0043] Another such modification includes derivatization of an amino group of a lysine, arginine, or histidine residue. For example, such amino groups can be acylated. Some suitable acyl groups include, for example, a benzoyl group or an alkanoyl group comprising any of the C₁-C₄ alkyl groups mentioned above, such as an acetyl or propionyl group.

[0044] The non-naturally occurring amino acids are preferably resistant, and more preferably insensitive, to common proteases. Examples of non-naturally occurring amino acids that are resistant or insensitive to proteases include the dextrorotatory (D-) form of any of the above-mentioned naturally occurring L-amino acids, as well as L- and/or D- non-naturally occurring amino acids. The D-amino acids do not normally occur in proteins, although they are found in certain peptide antibiotics that are synthesized by means other than the normal ribosomal protein synthetic machinery of the cell. As used herein, the D-amino acids are considered to be non-naturally occurring amino acids.

[0045] In order to minimize protease sensitivity, the peptides useful in the methods of the invention should have less than five, preferably less than four, more preferably less than three, and most preferably, less than two contiguous L-amino acids

recognized by common proteases, irrespective of whether the amino acids are naturally or non-naturally occurring. Optimally, the peptide has only D-amino acids, and no L-amino acids.

[0046] If the peptide contains protease sensitive sequences of amino acids, at least one of the amino acids is preferably a non-naturally-occurring D-amino acid, thereby conferring protease resistance. An example of a protease sensitive sequence includes two or more contiguous basic amino acids that are readily cleaved by common proteases, such as endopeptidases and trypsin. Examples of basic amino acids include arginine, lysine and histidine.

[0047] It is important that the aromatic-cationic peptides have a minimum number of net positive charges at physiological pH in comparison to the total number of amino acid residues in the peptide. The minimum number of net positive charges at physiological pH will be referred to below as (p_m). The total number of amino acid residues in the peptide will be referred to below as (r).

[0048] The minimum number of net positive charges discussed below are all at physiological pH. The term “physiological pH” as used herein refers to the normal pH in the cells of the tissues and organs of the mammalian body. For instance, the physiological pH of a human is normally approximately 7.4, but normal physiological pH in mammals may be any pH from about 7.0 to about 7.8.

[0049] “Net charge” as used herein refers to the balance of the number of positive charges and the number of negative charges carried by the amino acids present in the peptide. In this specification, it is understood that net charges are measured at physiological pH. The naturally occurring amino acids that are positively charged at physiological pH include L-lysine, L-arginine, and L-histidine. The naturally occurring amino acids that are negatively charged at physiological pH include L-aspartic acid and L-glutamic acid.

[0050] Typically, a peptide has a positively charged N-terminal amino group and a negatively charged C-terminal carboxyl group. The charges cancel each other out at physiological pH.

[0051] As an example of calculating net charge, the peptide Tyr-Arg-Phe-Lys-Glu-His-Trp-Arg has one negatively charged amino acid (i.e., Glu) and four positively charged amino acids (i.e., two Arg residues, one Lys, and one His). Therefore, the above peptide has a net positive charge of three.

[0052] In one embodiment of the present invention, the aromatic-cationic peptides have a relationship between the minimum number of net positive charges at physiological pH (p_m) and the total number of amino acid residues (r) wherein $3p_m$ is the largest number that is less than or equal to $r + 1$. In this embodiment, the relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) is as follows:

(r)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(p_m)	1	1	2	2	2	3	3	3	4	4	4	5	5	5	6	6	6	7

[0053] In another embodiment, the aromatic-cationic peptides have a relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) wherein $2p_m$ is the largest number that is less than or equal to $r + 1$. In this embodiment, the relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) is as follows:

(r)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(p_m)	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10	10

[0054] In one embodiment, the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) are equal. In another embodiment, the peptides have three or four amino acid residues and a minimum of one net positive charge, preferably,

a minimum of two net positive charges and more preferably a minimum of three net positive charges.

[0055] It is also important that the aromatic-cationic peptides have a minimum number of aromatic groups in comparison to the total number of net positive charges (p_t). The minimum number of aromatic groups will be referred to below as (a).

[0056] Naturally occurring amino acids that have an aromatic group include the amino acids histidine, tryptophan, tyrosine, and phenylalanine. For example, the hexapeptide Lys-Gln-Tyr-Arg-Phe-Trp has a net positive charge of two (contributed by the lysine and arginine residues) and three aromatic groups (contributed by tyrosine, phenylalanine and tryptophan residues).

[0057] In one embodiment of the present invention, the aromatic-cationic peptides useful in the methods of the present invention have a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges at physiological pH (p_t) wherein $3a$ is the largest number that is less than or equal to $p_t + 1$, except that when p_t is 1, a may also be 1. In this embodiment, the relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) is as follows:

(p_t)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(a)	1	1	1	1	2	2	2	3	3	3	4	4	4	5	5	5	6	6	6	7

[0058] In another embodiment, the aromatic-cationic peptides have a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) wherein $2a$ is the largest number that is less than or equal to $p_t + 1$. In this embodiment, the relationship between the minimum number of aromatic amino acid residues (a) and the total number of net positive charges (p_t) is as follows:

(p_t)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(a)	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10	10

[0059] In another embodiment, the number of aromatic groups (a) and the total number of net positive charges (p_+) are equal.

[0060] Carboxyl groups, especially the terminal carboxyl group of a C-terminal amino acid, are preferably amidated with, for example, ammonia to form the C-terminal amide. Alternatively, the terminal carboxyl group of the C-terminal amino acid may be amidated with any primary or secondary amine. The primary or secondary amine may, for example, be an alkyl, especially a branched or unbranched C₁-C₄ alkyl, or an aryl amine. Accordingly, the amino acid at the C-terminus of the peptide may be converted to an amido, N-methylamido, N-ethylamido, N,N-dimethylamido, N,N-diethylamido, N-methyl-N-ethylamido, N-phenylamido or N-phenyl-N-ethylamido group.

[0061] The free carboxylate groups of the asparagine, glutamine, aspartic acid, and glutamic acid residues not occurring at the C-terminus of the aromatic-cationic peptides of the present invention may also be amidated wherever they occur within the peptide. The amidation at these internal positions may be with ammonia or any of the primary or secondary amines described above.

[0062] In one embodiment, the aromatic-cationic peptide useful in the methods of the present invention is a tripeptide having two net positive charges and at least one aromatic amino acid. In a particular embodiment, the aromatic-cationic peptide useful in the methods of the present invention is a tripeptide having two net positive charges and two aromatic amino acids.

[0063] Aromatic-cationic peptides useful in the methods of the present invention include, but are not limited to, the following peptide examples:

Lys-D-Arg-Tyr-NH₂,

Phe-D-Arg-His,

D-Tyr-Trp-Lys-NH₂,

Trp-D-Lys-Tyr-Arg-NH₂,

Tyr-His-D-Gly-Met,

Phe-Arg-D-His-Asp,

Tyr-D-Arg-Phe-Lys-Glu-NH₂,
 Met-Tyr-D-Lys-Phe-Arg,
 D-His-Glu-Lys-Tyr-D-Phe-Arg,
 Lys-D-Gln-Tyr-Arg-D-Phe-Trp-NH₂,
 Phe-D-Arg-Lys-Trp-Tyr-D-Arg-His,
 Gly-D-Phe-Lys-Tyr-His-D-Arg-Tyr-NH₂,
 Val-D-Lys-His-Tyr-D-Phe-Ser-Tyr-Arg-NH₂,
 Trp-Lys-Phe-D-Asp-Arg-Tyr-D-His-Lys,
 Lys-Trp-D-Tyr-Arg-Asn-Phe-Tyr-D-His-NH₂,
 Thr-Gly-Tyr-Arg-D-His-Phe-Trp-D-His-Lys,
 Asp-D-Trp-Lys-Tyr-D-His-Phe-Arg-D-Gly-Lys-NH₂,
 D-His-Lys-Tyr-D-Phe-Glu-D-Asp-D-His-D-Lys-Arg-Trp-NH₂,
 Ala-D-Phe-D-Arg-Tyr-Lys-D-Trp-His-D-Tyr-Gly-Phe,
 Tyr-D-His-Phe-D-Arg-Asp-Lys-D-Arg-His-Trp-D-His-Phe,
 Phe-Phe-D-Tyr-Arg-Glu-Asp-D-Lys-Arg-D-Arg-His-Phe-NH₂,
 Phe-Try-Lys-D-Arg-Trp-His-D-Lys-D-Lys-Glu-Arg-D-Tyr-Thr,
 Tyr-Asp-D-Lys-Tyr-Phe-D-Lys-D-Arg-Phe-Pro-D-Tyr-His-Lys,
 Glu-Arg-D-Lys-Tyr-D-Val-Phe-D-His-Trp-Arg-D-Gly-Tyr-Arg-D-Met-NH₂,
 Arg-D-Leu-D-Tyr-Phe-Lys-Glu-D-Lys-Arg-D-Trp-Lys-D-Phe-Tyr-D-Arg-Gly,
 D-Glu-Asp-Lys-D-Arg-D-His-Phe-Phe-D-Val-Tyr-Arg-Tyr-D-Tyr-Arg-His-Phe-NH₂,
 Asp-Arg-D-Phe-Cys-Phe-D-Arg-D-Lys-Tyr-Arg-D-Tyr-Trp-D-His-Tyr-D-Phe-Lys-Phe,
 His-Tyr-D-Arg-Trp-Lys-Phe-D-Asp-Ala-Arg-Cys-D-Tyr-His-Phe-D-Lys-Tyr-His-Ser-NH₂,
 Gly-Ala-Lys-Phe-D-Lys-Glu-Arg-Tyr-His-D-Arg-D-Arg-Asp-Tyr-Trp-D-His-Trp-His-D-Lys-Asp, and
 Thr-Tyr-Arg-D-Lys-Trp-Tyr-Glu-Asp-D-Lys-D-Arg-His-Phe-D-Tyr-Gly-Val-Ile-D-His-Arg-Tyr-Lys-NH₂.

[0064] In one embodiment, the peptides useful in the methods of the present invention have mu-opioid receptor agonist activity (i.e., they activate the mu-opioid receptor). Activation of the mu-opioid receptor typically elicits an analgesic effect.

[0065] In certain instances, an aromatic-cationic peptide having mu-opioid receptor agonist activity is preferred. For example, during short-term treatment, such as in an acute disease or condition, it may be beneficial to use an aromatic-cationic peptide that

activates the mu-opioid receptor. Such acute diseases and conditions are often associated with moderate or severe pain. In these instances, the analgesic effect of the aromatic-cationic peptide may be beneficial in the treatment regimen of the human patient or other mammal. An aromatic-cationic peptide which does not activate the mu-opioid receptor, however, may also be used with or without an analgesic, according to clinical requirements.

[0066] Alternatively, in other instances, an aromatic-cationic peptide that does not have mu-opioid receptor agonist activity is preferred. For example, during long-term treatment, such as in a chronic disease state or condition, the use of an aromatic-cationic peptide that activates the mu-opioid receptor may be contraindicated. In these instances the potentially adverse or addictive effects of the aromatic-cationic peptide may preclude the use of an aromatic-cationic peptide that activates the mu-opioid receptor in the treatment regimen of a human patient or other mammal. Potential adverse effects may include sedation, constipation and respiratory depression. In such instances an aromatic-cationic peptide that does not activate the mu-opioid receptor may be an appropriate treatment.

[0067] Peptides useful in the methods of the present invention which have mu-opioid receptor agonist activity are typically those peptides which have a tyrosine residue or a tyrosine derivative at the N-terminus (i.e., the first amino acid position). Preferred derivatives of tyrosine include 2'-methyltyrosine (Mmt); 2',6'-dimethyltyrosine (2'6'Dmt); 3',5'-dimethyltyrosine (3'5'Dmt); N,2',6'-trimethyltyrosine (Tmt); and 2'-hydroxy-6'-methyltryosine (Hmt).

[0068] In a particular preferred embodiment, a peptide that has mu-opioid receptor agonist activity has the formula Tyr-D-Arg-Phe-Lys-NH₂ (for convenience represented by the acronym: DALDA, which is referred to herein as SS-01). DALDA has a net positive charge of three, contributed by the amino acids tyrosine, arginine, and lysine and has two aromatic groups contributed by the amino acids phenylalanine and tyrosine. The tyrosine of DALDA can be a modified derivative of tyrosine such as in 2',6'-dimethyltyrosine to produce the compound having the formula 2',6'-Dmt-D-Arg-Phe-Lys-NH₂ (i.e., Dmt¹-DALDA, which is referred to herein as SS-02).

[0069] Peptides that do not have mu-opioid receptor agonist activity generally do not have a tyrosine residue or a derivative of tyrosine at the N-terminus (i.e., amino acid

position 1). The amino acid at the N-terminus can be any naturally occurring or non-naturally occurring amino acid other than tyrosine.

[0070] In one embodiment, the amino acid at the N-terminus is phenylalanine or its derivative. Preferred derivatives of phenylalanine include 2'-methylphenylalanine (Mmp), 2',6'-dimethylphenylalanine (Dmp), N,2',6'-trimethylphenylalanine (Tmp), and 2'-hydroxy-6'-methylphenylalanine (Hmp).

[0071] Another aromatic-cationic peptide that does not have mu-opioid receptor agonist activity has the formula Phe-D-Arg-Phe-Lys-NH₂ (i.e., Phe¹-DALDA, which is referred to herein as SS-20). Alternatively, the N-terminal phenylalanine can be a derivative of phenylalanine such as 2',6'-dimethylphenylalanine (2'6'Dmp). DALDA containing 2',6'-dimethylphenylalanine at amino acid position 1 has the formula 2',6'-Dmp-D-Arg-Phe-Lys-NH₂ (i.e., 2'6'Dmp¹-DALDA).

[0072] In a preferred embodiment, the amino acid sequence of Dmt¹-DALDA (SS-02) is rearranged such that Dmt is not at the N-terminus. An example of such an aromatic-cationic peptide that does not have mu-opioid receptor agonist activity has the formula D-Arg-2'6'Dmt-Lys-Phe-NH₂ (referred to in this specification as SS-31).

[0073] DALDA, Phe¹-DALDA, SS-31, and their derivatives can further include functional analogs. A peptide is considered a functional analog of DALDA, Phe¹-DALDA, or SS-31 if the analog has the same function as DALDA, Phe¹-DALDA, or SS-31. The analog may, for example, be a substitution variant of DALDA, Phe¹-DALDA, or SS-31, wherein one or more amino acids are substituted by another amino acid.

[0074] Suitable substitution variants of DALDA, Phe¹-DALDA, or SS-31 include conservative amino acid substitutions. Amino acids may be grouped according to their physicochemical characteristics as follows:

- (a) Non-polar amino acids: Ala(A) Ser(S) Thr(T) Pro(P) Gly(G);
- (b) Acidic amino acids: Asn(N) Asp(D) Glu(E) Gln(Q);
- (c) Basic amino acids: His(H) Arg(R) Lys(K);
- (d) Hydrophobic amino acids: Met(M) Leu(L) Ile(I) Val(V); and
- (e) Aromatic amino acids: Phe(F) Tyr(Y) Trp(W) His (H).

[0075] Substitutions of an amino acid in a peptide by another amino acid in the same group is referred to as a conservative substitution and may preserve the physicochemical characteristics of the original peptide. In contrast, substitutions of an amino acid in a peptide by another amino acid in a different group is generally more likely to alter the characteristics of the original peptide.

[0076] Examples of analogs useful in the practice of the present invention that activate mu-opioid receptors include, but are not limited, to the aromatic-cationic peptides shown in Table 1.

TABLE 1

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	Amino Acid Position 5 (if present)	C-Terminal Modification
Tyr	D-Arg	Phe	Lys		NH ₂
Tyr	D-Arg	Phe	Orn		NH ₂
Tyr	D-Arg	Phe	Dab		NH ₂
Tyr	D-Arg	Phe	Dap		NH ₂
2'6'Dmt	D-Arg	Phe	Lys		NH ₂
2'6'Dmt	D-Arg	Phe	Lys	Cys	NH ₂
2'6'Dmt	D-Arg	Phe	Lys-NH(CH ₂) ₂ -NH-dns		NH ₂
2'6'Dmt	D-Arg	Phe	Lys-NH(CH ₂) ₂ -NH-atn		NH ₂
2'6'Dmt	D-Arg	Phe	dnsLys		NH ₂
2'6'Dmt	D-Cit	Phe	Lys		NH ₂
2'6'Dmt	D-Cit	Phe	Ahp		NH ₂
2'6'Dmt	D-Arg	Phe	Orn		NH ₂
2'6'Dmt	D-Arg	Phe	Dab		NH ₂
2'6'Dmt	D-Arg	Phe	Dap		NH ₂
2'6'Dmt	D-Arg	Phe	Ahp(2-aminoheptanoic acid)		NH ₂
Bio-2'6'Dmt	D-Arg	Phe	Lys		NH ₂
3'5'Dmt	D-Arg	Phe	Lys		NH ₂
3'5'Dmt	D-Arg	Phe	Orn		NH ₂
3'5'Dmt	D-Arg	Phe	Dab		NH ₂
3'5'Dmt	D-Arg	Phe	Dap		NH ₂
Tyr	D-Arg	Tyr	Lys		NH ₂
Tyr	D-Arg	Tyr	Orn		NH ₂
Tyr	D-Arg	Tyr	Dab		NH ₂
Tyr	D-Arg	Tyr	Dap		NH ₂
2'6'Dmt	D-Arg	Tyr	Lys		NH ₂
2'6'Dmt	D-Arg	Tyr	Orn		NH ₂
2'6'Dmt	D-Arg	Tyr	Dab		NH ₂
2'6'Dmt	D-Arg	Tyr	Dap		NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Lys		NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Orn		NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Dab		NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Dap		NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Arg		NH ₂

3'5'Dmt	D-Arg	3'5'Dmt	Lys	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Orn	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Dab	NH ₂
Tyr	D-Lys	Phe	Dap	NH ₂
Tyr	D-Lys	Phe	Arg	NH ₂
Tyr	D-Lys	Phe	Lys	NH ₂
Tyr	D-Lys	Phe	Orn	NH ₂
2'6'Dmt	D-Lys	Phe	Dab	NH ₂
2'6'Dmt	D-Lys	Phe	Dap	NH ₂
2'6'Dmt	D-Lys	Phe	Arg	NH ₂
2'6'Dmt	D-Lys	Phe	Lys	NH ₂
3'5'Dmt	D-Lys	Phe	Orn	NH ₂
3'5'Dmt	D-Lys	Phe	Dab	NH ₂
3'5'Dmt	D-Lys	Phe	Dap	NH ₂
3'5'Dmt	D-Lys	Phe	Arg	NH ₂
Tyr	D-Lys	Tyr	Lys	NH ₂
Tyr	D-Lys	Tyr	Orn	NH ₂
Tyr	D-Lys	Tyr	Dab	NH ₂
Tyr	D-Lys	Tyr	Dap	NH ₂
2'6'Dmt	D-Lys	Tyr	Lys	NH ₂
2'6'Dmt	D-Lys	Tyr	Orn	NH ₂
2'6'Dmt	D-Lys	Tyr	Dab	NH ₂
2'6'Dmt	D-Lys	Tyr	Dap	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Lys	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Orn	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Dab	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Dap	NH ₂
2'6'Dmt	D-Arg	Phe	dnsDap	NH ₂
2'6'Dmt	D-Arg	Phe	atnDap	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Lys	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Orn	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Dab	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Dap	NH ₂
Tyr	D-Lys	Phe	Arg	NH ₂
Tyr	D-Orn	Phe	Arg	NH ₂
Tyr	D-Dab	Phe	Arg	NH ₂
Tyr	D-Dap	Phe	Arg	NH ₂
2'6'Dmt	D-Arg	Phe	Arg	NH ₂
2'6'Dmt	D-Lys	Phe	Arg	NH ₂
2'6'Dmt	D-Orn	Phe	Arg	NH ₂
2'6'Dmt	D-Dab	Phe	Arg	NH ₂
3'5'Dmt	D-Dap	Phe	Arg	NH ₂
3'5'Dmt	D-Arg	Phe	Arg	NH ₂
3'5'Dmt	D-Lys	Phe	Arg	NH ₂
3'5'Dmt	D-Orn	Phe	Arg	NH ₂
Tyr	D-Lys	Tyr	Arg	NH ₂
Tyr	D-Orn	Tyr	Arg	NH ₂
Tyr	D-Dab	Tyr	Arg	NH ₂
Tyr	D-Dap	Tyr	Arg	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Arg	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Arg	NH ₂
2'6'Dmt	D-Orn	2'6'Dmt	Arg	NH ₂

2'6'Dmt	D-Dab	2'6'Dmt	Arg	NH ₂
3'5'Dmt	D-Dap	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Orn	3'5'Dmt	Arg	NH ₂
Mmt	D-Arg	Phe	Lys	NH ₂
Mmt	D-Arg	Phe	Orn	NH ₂
Mmt	D-Arg	Phe	Dab	NH ₂
Mmt	D-Arg	Phe	Dap	NH ₂
Tmt	D-Arg	Phe	Lys	NH ₂
Tmt	D-Arg	Phe	Orn	NH ₂
Tmt	D-Arg	Phe	Dab	NH ₂
Tmt	D-Arg	Phe	Dap	NH ₂
Hmt	D-Arg	Phe	Lys	NH ₂
Hmt	D-Arg	Phe	Orn	NH ₂
Hmt	D-Arg	Phe	Dab	NH ₂
Hmt	D-Arg	Phe	Dap	NH ₂
Mmt	D-Lys	Phe	Lys	NH ₂
Mmt	D-Lys	Phe	Orn	NH ₂
Mmt	D-Lys	Phe	Dab	NH ₂
Mmt	D-Lys	Phe	Dap	NH ₂
Mmt	D-Lys	Phe	Arg	NH ₂
Tmt	D-Lys	Phe	Lys	NH ₂
Tmt	D-Lys	Phe	Orn	NH ₂
Tmt	D-Lys	Phe	Dab	NH ₂
Tmt	D-Lys	Phe	Dap	NH ₂
Tmt	D-Lys	Phe	Arg	NH ₂
Hmt	D-Lys	Phe	Lys	NH ₂
Hmt	D-Lys	Phe	Orn	NH ₂
Hmt	D-Lys	Phe	Dab	NH ₂
Hmt	D-Lys	Phe	Dap	NH ₂
Hmt	D-Lys	Phe	Arg	NH ₂
Mmt	D-Lys	Phe	Arg	NH ₂
Mmt	D-Orn	Phe	Arg	NH ₂
Mmt	D-Dab	Phe	Arg	NH ₂
Mmt	D-Dap	Phe	Arg	NH ₂
Mmt	D-Arg	Phe	Arg	NH ₂
Tmt	D-Lys	Phe	Arg	NH ₂
Tmt	D-Orn	Phe	Arg	NH ₂
Tmt	D-Dab	Phe	Arg	NH ₂
Tmt	D-Dap	Phe	Arg	NH ₂
Tmt	D-Arg	Phe	Arg	NH ₂
Hmt	D-Lys	Phe	Arg	NH ₂
Hmt	D-Orn	Phe	Arg	NH ₂
Hmt	D-Dab	Phe	Arg	NH ₂
Hmt	D-Dap	Phe	Arg	NH ₂
Hmt	D-Arg	Phe	Arg	NH ₂

Dab = diaminobutyric

Dap = diaminopropionic acid

Dmt = dimethyltyrosine

Mmt = 2'-methyltyrosine

Tmt = N, 2',6'-trimethyltyrosine

Hmt = 2'-hydroxy,6'-methyltyrosine

dnsDap = β -dansyl-L- α , β -diaminopropionic acid

atnDap = β -anthraniloyl-L- α , β -diaminopropionic acid

Bio = biotin

[0077] Examples of analogs useful in the practice of the present invention that do not activate mu-opioid receptors include, but are not limited to, the aromatic-cationic peptides shown in Table 2.

TABLE 2

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
D-Arg	Dmt	Lys	Phe	NH ₂
D-Arg	Dmt	Phe	Lys	NH ₂
D-Arg	Phe	Lys	Dmt	NH ₂
D-Arg	Phe	Dmt	Lys	NH ₂
D-Arg	Lys	Dmt	Phe	NH ₂
D-Arg	Lys	Phe	Dmt	NH ₂
Phe	Lys	Dmt	D-Arg	NH ₂
Phe	Lys	D-Arg	Dmt	NH ₂
Phe	D-Arg	Dmt	Lys	NH ₂
Phe	D-Arg	Lys	Dmt	NH ₂
Phe	Dmt	D-Arg	Lys	NH ₂
Phe	Dmt	Lys	D-Arg	NH ₂
Lys	Phe	D-Arg	Dmt	NH ₂
Lys	Phe	Dmt	D-Arg	NH ₂
Lys	Dmt	D-Arg	Phe	NH ₂
Lys	Dmt	Phe	D-Arg	NH ₂
Lys	D-Arg	Phe	Dmt	NH ₂
Lys	D-Arg	Dmt	Phe	NH ₂
D-Arg	Dmt	D-Arg	Phe	NH ₂
D-Arg	Dmt	D-Arg	Dmt	NH ₂
D-Arg	Dmt	D-Arg	Tyr	NH ₂
D-Arg	Dmt	D-Arg	Trp	NH ₂
Trp	D-Arg	Phe	Lys	NH ₂
Trp	D-Arg	Tyr	Lys	NH ₂
Trp	D-Arg	Trp	Lys	NH ₂
Trp	D-Arg	Dmt	Lys	NH ₂
D-Arg	Trp	Lys	Phe	NH ₂
D-Arg	Trp	Phe	Lys	NH ₂

D-Arg	Trp	Lys	Dmt	NH ₂
D-Arg	Trp	Dmt	Lys	NH ₂
D-Arg	Lys	Trp	Phe	NH ₂
D-Arg	Lys	Trp	Dmt	NH ₂
Cha	D-Arg	Phe	Lys	NH ₂
Ala	D-Arg	Phe	Lys	NH ₂

Cha = cyclohexyl

[0078] The amino acids of the peptides shown in table 1 and 2 may be in either the L- or the D- configuration.

Methods

[0079] The aromatic-cationic peptides described above are useful in reducing CD36 expression in a cell. For the purposes of this specification, CD36 expression in a cell is considered to be reduced if the expression of CD36 is decreased by about 10%, preferably by about 25%, more preferably by about 50%, even more preferably by about 75%. Optimally, CD36 is reduced to about normal levels in a cell.

[0080] CD36 is expressed on a wide variety of cells. Examples of such cells include macrophages, platelets, adipocytes, endothelial cells such as microvascular endothelial cells and umbilical vein endothelial cells; epithelial cells such as intestinal epithelial cells, gall bladder epithelial cells, bladder epithelial cells, bronchial epithelial cells and alveolar epithelial cells; renal tubular cells; pancreatic islet cells; hepatocytes; skeletal muscle cells; cardiac muscle cells; neuronal cells; glia cells; pancreas cells; sperm cells; etc.

[0081] For the purposes of this specification, cells expressing about 10%, typically about 25%, about typically about 50%, and even more typically about 75% more CD36 than normal cells are considered to express increased levels of CD36.

[0082] In one embodiment, the invention provides a method for reducing CD36 expression in a cell. Any cell that expresses CD36 can be used in the method of the invention, and include those mentioned above. The method for reducing CD36 expression in a cell comprises contacting the cell with an effective amount of an aromatic-cationic peptide described above.

[0083] In another embodiment, the invention provides a method for reducing CD36 expression in a mammal in need thereof. The method for reducing CD36 expression in the mammal comprises administering to the mammal an effective amount of an aromatic-cationic peptide described herein.

[0084] Mammals in need of reducing CD36 expression include, for example, mammals that have increased CD36 expression. The increased expression of CD36 is associated with various diseases and conditions. Examples of diseases and conditions characterized by increased CD36 expression include, but are not limited to atherosclerosis, inflammation, abnormal angiogenesis, abnormal lipid metabolism, abnormal removal of apoptotic cells, ischemia such as cerebral ischemia and myocardial ischemia, ischemia-reperfusion, ureteral obstruction, stroke, Alzheimer's Disease, diabetes, diabetic nephropathy and obesity. A discussion on the involvement of CD36 in atherosclerosis may be found in "Targeted disruption of the class B scavenger receptor CD36 protects against atherosclerotic lesion development in mice". Febraio M, Podrez EA, Smith JD, Hajjar DP, Hazen SL et al. J Clinical Investigation 105:1049-1056, 2000 and "CD36: a class B scavenger receptor involved in angiogenesis, atherosclerosis, inflammation, and lipid metabolism". Febraio M., Hajjar DP and Silverstein RL. Journal of Clinical Investigation 108:785-791, 2001.

[0085] Mammals in need of reducing CD36 expression also include mammals suffering from complications of diabetes. Some complications of diabetes include, in addition to nephropathy, neuropathy, retinopathy, coronary artery disease, and peripheral vascular disease associated with diabetes.

[0086] In another embodiment, the invention relates to a method for reducing CD36 expression in removed organs and tissues. The method comprises contacting the removed organ or tissue with an effective amount of an aromatic-cationic peptide described above. An organ or tissue may, for example, be removed from a donor for autologous or heterologous transplantation. Some examples of organs and tissues include heart, lungs, pancreas, kidney, liver, skin, etc.

Synthesis of the Peptides

[0087] The peptides useful in the methods of the present invention may be synthesized by any of the methods well known in the art. Suitable methods for chemically synthesizing the protein include, for example those described by Stuart and Young in "Solid Phase Peptide Synthesis," Second Edition, Pierce Chemical Company (1984), and in "Solid Phase Peptide Synthesis," Methods Enzymol. 289, Academic Press, Inc, New York (1997).

Modes of Administration

[0088] Any method known to those in the art for contacting a cell, organ or tissue with a peptide may be employed. Suitable methods include *in vitro*, *ex vivo*, or *in vivo* methods.

[0089] *In vitro* methods typically include cultured samples. For example, a cell can be placed in a reservoir (e.g., tissue culture plate), and incubated with an aromatic-cationic peptide under appropriate conditions suitable for reducing CD36 expression. Suitable incubation conditions can be readily determined by those skilled in the art.

[0090] *Ex vivo* methods typically include cells, organs or tissues removed from a mammal, such as a human. The cells, organs or tissues can, for example, be incubated with the peptide under appropriate conditions. The contacted cells, organs or tissues are normally returned to the donor, placed in a recipient, or stored for future use. Thus, the peptide is generally in a pharmaceutically acceptable carrier.

[0091] *In vivo* methods are typically limited to the administration of an aromatic-cationic peptide, such as those described above, to a mammal, preferably a human. The peptides useful in the methods of the present invention are administered to a mammal in an amount effective in reducing expression of CD36 or treating the mammal. The effective amount is determined during pre-clinical trials and clinical trials by methods familiar to physicians and clinicians.

[0092] An effective amount of a peptide useful in the methods of the present invention, preferably in a pharmaceutical composition, may be administered to a mammal in need thereof by any of a number of well-known methods for administering pharmaceutical compounds. The peptide may be administered systemically or locally.

794,4-30

[0093] In one embodiment, the peptide is administered intravenously. For example, the aromatic-cationic peptides useful in the methods of the present invention may be administered via rapid intravenous bolus injection. Preferably, however, the peptide is administered as a constant rate intravenous infusion.

[0094] The peptide may also be administered orally, topically, intranasally, intramuscularly, subcutaneously, or transdermally. In a preferred embodiment, transdermal administration of the aromatic-cationic peptides by methods of the present invention is by iontophoresis, in which the charged peptide is delivered across the skin by an electric current.

[0095] Other routes of administration include intracerebroventricularly or intrathecally. Intracerebroventricularly refers to administration into the ventricular system of the brain. Intrathecally refers to administration into the space under the arachnoid membrane of the spinal cord. Thus intracerebroventricular or intrathecal administration may be preferred for those diseases and conditions which affect the organs or tissues of the central nervous system.

[0096] The peptides useful in the methods of the invention may also be administered to mammals by sustained release, as is known in the art. Sustained release administration is a method of drug delivery to achieve a certain level of the drug over a particular period of time. The level typically is measured by serum or plasma concentration.

[0097] A description of methods for delivering a compound by controlled release can be found in international PCT Application No. WO 02/083106.

[0098] Any formulation known in the art of pharmacy is suitable for administration of the aromatic-cationic peptides useful in the methods of the present invention. For oral administration, liquid or solid formulations may be used. Some examples of formulations include tablets, gelatin capsules, pills, troches, elixirs, suspensions, syrups, wafers, chewing gum and the like. The peptides can be mixed with a suitable pharmaceutical carrier (vehicle) or excipient as understood by practitioners in the art. Examples of carriers and excipients include starch, milk, sugar, certain types of clay, gelatin, lactic acid, stearic acid or salts

79434-30

thereof, including magnesium or calcium stearate, talc, vegetable fats or oils, gums and glycols.

[0099] For systemic, intracerebroventricular, intrathecal, topical, intranasal, subcutaneous, or transdermal administration, formulations of the aromatic-cationic peptides useful in the methods of the present inventions may utilize conventional diluents, carriers, or excipients etc., such as those known in the art to deliver the peptides. For example, the formulations may comprise one or more of the following: a stabilizer, a surfactant, preferably a nonionic surfactant, and optionally a salt and/or a buffering agent. The peptide may be delivered in the form of an aqueous solution, or in a lyophilized form.

[00100] The stabilizer may, for example, be an amino acid, such as for instance, glycine; or an oligosaccharide, such as for example, sucrose, tetralose, lactose or a dextran. Alternatively, the stabilizer may be a sugar alcohol, such as for instance, mannitol; or a combination thereof. Preferably the stabilizer or combination of stabilizers constitutes from about 0.1% to about 10% weight for weight of the peptide.

[0100] The surfactant is preferably a nonionic surfactant, such as a polysorbate. Some examples of suitable surfactants include Tween²⁰, Tween⁸⁰; a polyethylene glycol or a polyoxyethylene polyoxypropylene glycol, such as Pluronic F-68 at from about 0.001% (w/v) to about 10% (w/v).

[0101] The salt or buffering agent may be any salt or buffering agent, such as for example, sodium chloride, or sodium/potassium phosphate, respectively. Preferably, the buffering agent maintains the pH of the pharmaceutical composition in the range of about 5.5 to about 7.5. The salt and/or buffering agent is also useful to maintain the osmolality at a level suitable for administration to a human or an animal. Preferably the salt or buffering agent is present at a roughly isotonic concentration of about 150mM to about 300mM.

[0102] The formulations of the peptides useful in the methods of the present invention may additionally contain one or more conventional additive. Some examples of such additives include a solubilizer such as, for example, glycerol; an antioxidant such as for example, benzalkonium chloride (a mixture of quaternary ammonium compounds, known as "quats"), benzyl alcohol, chloretone or chlorobutanol; anaesthetic agent such as for example a

*trademark

morphine derivative; or an isotonic agent etc., such as described above. As a further precaution against oxidation or other spoilage, the pharmaceutical compositions may be stored under nitrogen gas in vials sealed with impermeable stoppers.

[0103] The mammal treated in accordance with the invention can be any mammal, including, for example, farm animals, such as sheep, pigs, cows, and horses; pet animals, such as dogs and cats; laboratory animals, such as rats, mice and rabbits. In a preferred embodiment, the mammal is a human.

EXAMPLES

Example 1: SS-31 reduced oxidized low-density lipoprotein (oxLDL)-induced CD36 expression and foam cell formation in mouse peritoneal macrophages.

[0104] Atherosclerosis is thought to develop as a result of lipid uptake by vascular-wall macrophages leading to the development of foam cells and the elaboration of cytokines and chemokines resulting in smooth muscle-cell proliferation. CD36 is a scavenger receptor that mediates uptake of oxLDL into macrophages and subsequent foam-cell development. CD36 knock out mice showed reduced uptake of oxLDL and reduced atherosclerosis.

[0105] CD36 expression is regulated at the transcriptional level by various cellular stimuli, including glucose and oxLDL. Macrophages were harvested from mice peritoneal cavity and culture overnight in the absence or presence of oxLDL (50 µg/ml) for 48 h. Incubation with oxLDL significantly increased CD36 mRNA (figure 1A). Inclusion of SS-31 (10 nM or 1 µM) to the culture medium abolished the up-regulation of CD36 (figure 1A). SS-31 by itself had no effect on CD36 expression.

[0106] Expression of CD36 protein, as determined by western blot, was also significantly increased after 48 h incubation with 25 µg/ml of oxLDL (oxL) when compared to vehicle control (V) (figure 1B). Other controls included CD36 expression from mouse heart (H) and macrophages obtained from CD36 knockout mice (KO). The amount of CD36 protein was normalized to β-actin. Incubation with SS-31 (1 µM) (S) significantly reduced CD36 protein expression compared to macrophages exposed to vehicle control (V) ($P < 0.01$,

ANOVA with *posthoc* Neuman Keuls test). Concurrent incubation with SS-31 (1 μ M) also significantly inhibited the upregulation of CD36 protein expression in macrophages exposed to 25 μ g/ml oxLDL for 48h (oxL/S) (P<0.01, ANOVA with *posthoc* Neuman Keuls test).

[0107] Incubation of macrophages with oxLDL for 48 h also increased foam cell formation (figure 1C). Foam cell is indicated by oil red O which stains lipid droplets red. Inclusion of SS-31 (1 μ M) prevented oxLDL-induced foam cell formation (figure 1C).

[0108] Incubation of macrophages with oxLDL increased apoptotic cells from 6.7% to 32.8%. Concurrent treatment with SS-31 (1 nM) significantly reduced the percentage of apoptotic cells induced by oxLDL to 20.8%.

Example 2: SS-31 Protected Mice from Acute Cerebral Ischemia

[0109] Cerebral ischemia initiates a cascade of cellular and molecular events that lead to brain damage. One such event is postischemic inflammation. Using a mouse model of cerebral ischemia-reperfusion (20 min. occlusion of the middle cerebral artery), it was found that CD36 was upregulated in microglia and macrophages in the post-ischemic brain, and there was increased reactive oxygen species production. CD36 knock out mice had a profound reduction in reactive oxygen species after ischemia and improved neurological function compared to wild type mice.

[0110] Cerebral ischemia was induced by occlusion of the right middle cerebral artery for 30 min. Wild-type (WT) mice were given either saline vehicle (Veh) (ip, n=9) or SS-31 (2 mg/kg or 5 mg/kg, ip, n=6) at 0, 6, 24 and 48 h after ischemia. Mice were killed 3 days after ischemia. Brains were removed, frozen, and sectioned. Brain sections were stained by the Nissl stain. Infarct volume and hemispheric swelling was determined using an image analyzer. Data were analyzed by one-way ANOVA with posthoc analysis.

[0111] Treatment of wild type mice with SS-31 (2 mg/kg or 5 mg/kg, ip, n=6) at 0, 6, 24 and 48 hours after 30 min. occlusion of the middle cerebral artery resulted in a significant reduction in infarct volume (figure 2A) and hemispheric swelling (figure 2B) compared to saline controls. (*P<0.05 compared to Veh).

[0112] Thirty min. cerebral ischemia in WT mice resulted in significant depletion in reduced glutathione (GSH) in the ipsilateral cortex and striatum compared to the contralateral side in vehicle-treated animals (figure 3). The depletion of GSH in the ipsilateral cortex was significantly reduced in mice treated with SS-31 (2 mg/kg ip at 0, 6, 24 and 48 h) (figure 3). The depletion of GSH in the striatum was also reduced by SS-31 treatment but did not reach statistical significance.

Example 3. SS-31 mediated protection against acute cerebral ischemia mimics protection observed in CD36 knockout mice.

[0113] CD36 knockout (CD36 KO) mice were subjected to acute cerebral ischemia as described under Example 2. CD36 KO mice were given either saline vehicle (Veh) (ip, n=5) or SS-31 (2 mg/kg, i.p. n=5) at 0, 6, 24 and 48 h after 30 min ischemia.

Infarct volume (figure 4A) and hemispheric swelling (figure 4B) in CD36 KO mice were similar whether they received saline or SS-31.

[0114] Treatment of CD36 KO mice with SS31 (2 mg/kg, i.p., n=5) also failed to further prevent GSH depletion in the ipsilateral cortex caused by 30 min ischemia (figure 5).

[0115] These data suggest that the protective action of SS-31 against acute cerebral ischemia may be mediated by inhibiting the upregulation of CD36.

Example 4. SS-31 reduced CD36 mRNA expression in post-ischemic brain.

[0116] Transient occlusion of the middle cerebral artery has been shown to significantly increase the expression of CD36 mRNA in microglia and macrophages in the post-ischemic brain. Wild-type mice were given saline vehicle (Veh, i.p., n=6) or SS-31 (5 mg/kg, i.p., n=6) at 0 and 6h after 30 min ischemia, and CD36 mRNA levels were determined using real time PCR. CD36 expression was upregulated almost 6-fold in the ipsilateral brain compared to the contralateral brain in mice that received saline (figure 6). CD36 mRNA was significantly reduced in the ipsilateral brain in mice that received SS-31 treatment (figure 6).

Example 5: SS-31 Suppressed Upregulation of CD36 in Renal Tubular Cells Following Unilateral Ureteral Obstruction

[0117] Unilateral ureteral obstruction (UUO) is a common clinical disorder associated with tubular cell apoptosis, macrophage infiltration, and interstitial fibrosis. Interstitial fibrosis leads to a hypoxic environment and contributes to progressive decline in renal function despite surgical correction. CD36 has been shown to be expressed on renal tubular cells.

[0118] CD36 was found to have been upregulated in tubular cells after UUO. UUO was performed in Sprague-Dawley rats. The rats were treated with saline (ip, n=6) or SS-31 (1 mg/kg ip, n=6) one day prior to induction of UUO, and once a day for 14 days after UUO. Rats were killed, kidneys removed, embedded in paraffin and sectioned. The slides were treated with the anti-CD36 polyclonal IgG (Santa Cruz #sc-9154; 1:100 with blocking serum) at room temperature for 1.5 hours. The slides were then incubated with the second antibody conjugated with biotin (anti-rabbit IgG-B1; ABC kit, PK-6101) at room temperature

79434-30

for 30 min. The slides were then treated with avidin, developed with DAB and counterstained with 10% hematoxylin. The contralateral unobstructed kidney served as the control for each animal.

[0119] UUO resulted in tubular dilation and significant increase in expression of CD36 on the tubular cells (fig. 7). Tubular dilation was also observed in rats treated with SS-31, but there was a significant reduction in CD36 expression (fig. 7). CD36 expression (brown stain) is primarily found on tubular cells in the contralateral unobstructed kidney (fig. 7A). CD36 expression was increased in the obstructed kidney in animals treated with saline (fig. 7B), but was much reduced in obstructed kidneys obtained from rats treated with SS-31 (fig. 7C).

[0120] To determine whether SS-31 reduces lipid peroxidation in kidney after UUO, rats were treated with either saline (n=6) or SS-31 (1 mg/kg ip, n=6) one day prior to induction of UUO, and once a day for 14 days UUO. Rats were then killed, kidneys removed, embedded in paraffin and sectioned. Slides were incubated with anti-HNE rabbit IgG and a biotin-linked anti-rabbit IgG was used as secondary antibody. The slides were developed with DAB. Lipid peroxidation, which was increased by UUO, was reduced by SS-31 treatment (fig. 8). HNE stain (brown) was significantly increased in tubular cells in the obstructed kidney (fig. 8B) compared to the contralateral control (fig. 8A). Obstructed kidneys from rats treated with SS-31 showed significantly less HNE stain (Fig. 8C) compared to saline-treated rats (fig. 8B).

[0121] To determine whether SS-31 reduced tubular cell apoptosis in obstructed kidney after UUO, rats were treated with either saline (n=6) or SS-31 (1 mg/kg ip, n=6) one day prior to induction of UUO, and once a day for 14 days after UUO. Rats were then killed, kidneys removed, embedded in paraffin and sectioned. To quantitate nuclei with fragmented DNA, the TUNEL assay were performed with in situ TUNEL kit (Intergen, Purchase, NY). Slides were developed with DAB and counterstained with 10% hematoxylin. The upregulation of CD36 in saline-treated controls associated with tubular cell apoptosis was significantly inhibited by SS-31 treatment (fig. 9). Compared to the contralateral unobstructed control (fig. 9A), a significant increase in apoptotic cells was observed in the obstructed kidney from saline-treated animals (fig. 9B). The number of apoptotic cells was

significantly reduced in obstructed kidney from SS-31 treated animals (fig. 9C) (P<0.001; n=6).

[0122] Macrophage infiltration (fig. 10) and interstitial fibrosis (fig. 11) were also prevented by SS-31 treatment. Rats were treated with either saline (n=6) or SS-31 (1 mg/kg ip, n=6) one day prior to induction of UUO, and once a day for 14 days after UUO. Rats were then killed, kidneys removed, embedded in paraffin and sectioned. Slides were treated with monoclonal antibody for ED1 macrophage (1:75; Serotec). Horseradish-peroxidase-linked rabbit anti-mouse secondary antibody (Dako) was used for macrophage detection. Sections were then counterstained with 10% hematoxylin. The number of macrophages in the obstructed kidney in saline-treated rats (fig. 10B) was significantly increased compared to the contralateral unobstructed control (fig. 10A). Macrophage infiltration was significantly reduced in rats treated with SS-31 (fig. 10C) (P<0.05; t-test).

[0123] Rats were treated with either saline (n=6) or SS-31 (1 mg/kg ip, n=6) one day prior to induction of UUO, and once a day for 14 days after UUO. Rats were then killed, kidneys removed, embedded in paraffin and sectioned. Slides were stained with hematoxylin and eosin and Masson's trichome for interstitial fibrosis (blue stain). Obstructed kidneys from saline-treated rats showed increase fibrosis (fig. 11B) compared to the contralateral unobstructed control (fig. 11A); while obstructed kidneys from SS-31 treated rats showed significantly less fibrosis (P<0.05; t-test).

[0124] These results show that SS-31 suppresses the upregulation of CD36 on renal tubular cells induced by UUO.

Example 6: SS-31 and SS-20 Reduced CD36 Expression in Isolated Hearts Upon Reperfusion After Prolonged Cold Ischemic Storage

[0125] Organ transplantation requires hypothermic storage of the isolated organ for transport to the recipient. Currently, cardiac transplantation is limited by the short time of cold ischemic storage that can be tolerated before coronary blood flow is severely compromised (<4 hours). The expression of CD36 in coronary endothelium and cardiac muscles is up-regulated in isolated hearts subjected to prolonged cold ischemic storage and warm reperfusion.

[0126] Isolated guinea pig hearts were perfused with St. Thomas solution alone, or St. Thomas solution containing 1 nM SS-31, or 100 nM SS-20, for 3 min. and then stored in the same solution at 4°C for 18 hours. After ischemic storage, hearts were reperfused with 34°C Kreb-Henseleit solution for 90 min. Hearts freshly isolated from guinea pigs were used as controls.

[0127] The hearts were fixed in paraffin and sliced for immunostaining with an anti-CD36 rabbit polyclonal antibody. The results are shown in fig. 12. Two sections are shown for each treatment group. Antibody staining showed that CD36 is expressed in endothelium and cardiac muscles in normal hearts. The “background” control (figs. 12A and 12B) represents two sections from a normal non-ischemic heart that was not treated with the primary antibody. “Normal heart” (figs. 12C and 12D) represents two sections obtained from a non-ischemic heart. The sections from a representative heart stored in St. Thomas solution (figs. 12E and 12F) for 18 hours at 4°C showed increased CD36 staining compared to “Normal heart.” CD36 staining was significantly reduced in hearts stored with either 1 nM SS-31 (figs. 12G and 12H) or 100 nM SS-20 (figs. 12I and 12J) in St. Thomas solution for 18 h.

[0128] CD36 staining is increased in hearts that have undergone 18 hours of cold ischemic storage and warm reperfusion. However, hearts that were stored with either 1 nM SS-31 or 100 nM SS-20 did not show the upregulation of CD36 expression.

[0129] Lipid peroxidation in the hearts was also decreased by the aromatic-cationic peptides. Guinea pig hearts were perfused with a cardioplegic solution (St. Thomas solution) alone or St. Thomas solution containing either 1 nM SS-31 or 100 nM SS-20 for 3 min. and then subjected to 18 hours of cold ischemia (4°C). The hearts were then reperfused with Krebs Henseleit buffer at 34°C for 90 min. Immunohistochemical analysis of 4-hydroxynonenol (HNE)- modified proteins in paraffin sections from tissue slices were performed by incubation with an anti-HNE antibody (Santa Cruz) and a fluorescent secondary antibody. HNE staining was significantly increased in hearts subjected to 18 hours of cold storage in St. Thomas solution (fig. 13B) compared to non-ischemic hearts (fig. 13A). HNE staining was reduced in hearts stored in SS-31 (fig. 13C) or SS-20 (fig. 13D).

[0130] Further, the peptides dramatically reduced endothelial apoptosis (fig. 14). Guinea pig hearts were perfused with a cardioplegic solution (St. Thomas solution) alone or St. Thomas solution containing either 1 nM SS-31 or 100 nM SS-20 for 3 min. and then subjected to 18 hours of cold ischemia (4°C). The hearts were then reperfused with Krebs Henseleit buffer at 34°C for 90 min. After deparaffinization, sections were incubated with deoxynucleotidyl transferase (Tdt) with digoxigenin-dNTP for 1 hour. The reaction was stopped with terminating buffer. A fluorescent anti-digoxigenin antibody was then applied. Hearts subjected to 18 hours of cold storage in St. Thomas solution (figs. 14C and 14D) showed prominent endothelial apoptosis whereas no endothelial apoptosis was observed in non-ischemic normal hearts (figs. 14A and 140B). Apoptotic cells were not observed in hearts stored in SS-31 (figs. 14E and 14F) or SS-20 (figs. 14G and 14H).

[0131] A significant improvement of coronary blood flow after prolonged cold ischemic storage and warm reperfusion occurred (fig. 15). Guinea pig hearts were perfused with a cardioplegic solution (St. Thomas solution) alone or St. Thomas solution containing either 1 nM SS-31 (fig. 15A) or 100 nM SS-20 (fig. 15B) for 3 min. and then subjected to 18 hours of cold ischemia (4°C). The hearts were then reperfused with Krebs Henseleit buffer at 34°C for 90 min. Coronary flow was significantly reduced after prolonged ischemia compared to pre-ischemic control (expressed as 100%). Preservation in either SS-31 or SS-20 significantly restored coronary flow to approximately 80% of pre-ischemic flow.

Example 7: SS-31 prevented renal damage in diabetic mice

[0132] CD36 expression is upregulated in a variety of tissues of diabetic patients, including monocytes, heart, kidneys, and plasma. High glucose is known to upregulate the expression of CD36 by improving the translational efficiency of CD36 mRNA. Diabetic nephropathy is a common complication of type 1 and type 2 diabetes, and is associated with tubular epithelial degeneration and interstitial fibrosis. CD36 has been identified as a mediator of tubular epithelial apoptosis in diabetic nephropathy. High glucose stimulates CD36 expression and apoptosis in proximal tubular epithelial cells.

[0133] Streptozotocin (STZ) was used to induce diabetes in mice. Three groups of CD-1 mice were studied: Group I – no STZ treatment; Group II – STZ (50 mg/kg, ip) was given once a day for 5d; Group III – STZ (50 mg/kg, ip) was given once a day for 5 d, and

SS-31 (3 mg/kg, ip) was given once a day for 16 d. STZ treatment resulted in progressive increase in blood glucose. By week 3, blood glucose values were: Group I (10.6 ± 0.27 mmol/L); Group II (24.5 ± 1.15 mmol/L); Group III (21.3 ± 1.48 mmol/L). Animals were sacrificed after 3 weeks and kidney tissues preserved for histopathology. Kidney sections were examined by Periodic Schiff (PAS) staining for renal tubular brush border.

[0134] STZ treatment caused dramatic loss of brush border in proximal tubules in the renal cortex (figure 16). In mice not treated with STZ, the renal brush border in the cortex was stained red with PAS (figure 16A, see white arrows). In mice treated with STZ, the brush border was obliterated, and the tubular epithelial cells showed small condensed nuclei (figure 16B). Daily treatment with SS-31 (3 mg/kg, ip) prevented the loss of brush border in the STZ-treated mice (figure 16C), and the nuclei appeared normal (figure 16C, top and bottom panels). In general, the architecture of the proximal renal tubules was preserved in diabetic mice treated with SS-31.

[0135] STZ treatment induced significant apoptosis in tubular epithelial cells (figure 17). Kidney sections were examined for apoptosis using the TUNEL assay. After deparaffinization, sections were incubated with deoxynucleotidyl transferase (Tdt) with digoxigenin-dNTP for 1 hour. The reaction was stopped with terminating buffer. A fluorescent anti-digoxigenin antibody was then applied. Kidney sections from mice treated with STZ showed large number of apoptotic nuclei in the proximal tubules (PT) (figure 17A, panel b), compared to no apoptotic cells in mice not treated with STZ (figure 17A, panel a). Treatment with daily SS-31 dramatically reduced apoptotic cells in the proximal tubule (figure 17A, panel c). Figure 17B shows the significant decrease in tubular cell apoptosis provided by SS-31.

[0136] CD36 expression in proximal tubular epithelial cells is known to be increased by high glucose and is upregulated in diabetic models. SS-31, by reducing CD36 expression, was able to inhibit tubular cell apoptosis and loss of brush border in mice treated with STZ without affecting blood glucose.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 79434-30D5 Seq 31-OCT-16 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> Cornell Research Foundation, Inc.
<120> Methods for Reducing CD36 Expression
<130> 79434-30D5
<140> Division of CA 2,622,911
<141> 2006-09-18
<150> US60/718170
<151> 2005-09-16
<160> 25
<170> PatentIn version 3.5

<210> 1
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (5)..(5)
<223> Amidation

<400> 1
Tyr Arg Phe Lys Glu
1 5

<210> 2
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<400> 2
Met Tyr Lys Phe Arg
1 5

<210> 3
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<400> 3
His Glu Lys Tyr Phe Arg
1 5

<210> 4
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE

```

<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (6)..(6)

<400> 4
Lys Gln Tyr Arg Phe Trp
1 5

<210> 5
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

<400> 5
Phe Arg Lys Trp Tyr Arg His
1 5

<210> 6
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> Amidation

<400> 6
Gly Phe Lys Tyr His Arg Tyr
1 5

<210> 7
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (8)..(8)
<223> Amidation

<400> 7
Val Lys His Tyr Phe Ser Tyr Arg
1 5

<210> 8
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<400> 8
 Trp Lys Phe Asp Arg Tyr His Lys
 1 5

<210> 9
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (3)..(3)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (8)..(8)
 <223> D-Amino Acid

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Amidation

<400> 9
 Lys Trp Tyr Arg Asn Phe Tyr His
 1 5

<210> 10
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (5)..(5)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (8)..(8)
 <223> D-Amino Acid

<400> 10
 Thr Gly Tyr Arg His Phe Trp His Lys
 1 5

<210> 11
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (9)..(9)
<223> Amidation

<400> 11
Asp Trp Lys Tyr His Phe Arg Gly Lys
1 5

<210> 12
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

```

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (10)..(10)
<223> Amidation

<400> 12
His Lys Tyr Phe Glu Asp His Lys Arg Trp
1 5 10

<210> 13
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> D-Amino Acid

<400> 13
Ala Phe Arg Tyr Lys Trp His Tyr Gly Phe
1 5 10

<210> 14
<211> 11
<212> PRT
<213> Artificial Sequence

```

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> D-Amino Acid

<400> 14
Tyr His Phe Arg Asp Lys Arg His Trp His Phe
1 5 10

<210> 15
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (11)..(11)
<223> Amidation

<400> 15
Phe Phe Tyr Arg Glu Asp Lys Arg Arg His Phe
1 5 10

<210> 16
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> D-Amino Acid

<400> 16
Phe Tyr Lys Arg Trp His Lys Lys Glu Arg Tyr Thr
1 5 10

<210> 17
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

```

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> D-Amino Acid

<400> 17
Tyr Asp Lys Tyr Phe Lys Arg Phe Pro Tyr His Lys
1 5 10

<210> 18
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (13)..(13)
<223> Amidation

<400> 18
Glu Arg Lys Tyr Val Phe His Trp Arg Gly Tyr Arg Met
1 5 10

```

<210> 19
 <211> 14
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (2)..(2)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (3)..(3)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (13)..(13)
 <223> D-Amino Acid

<400> 19
 Arg Leu Tyr Phe Lys Glu Lys Arg Trp Lys Phe Tyr Arg Gly
 1 5 10

<210> 20
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> D-Amino Acid

<220>
<221> MOD_RES
<222> (15)..(15)
<223> Amidation

<400> 20
Glu Asp Lys Arg His Phe Phe Val Tyr Arg Tyr Tyr Arg His Phe
1 5 10 15

<210> 21
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Peptide

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> D-Amino Acid

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (12)..(12)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> D-Amino Acid

<400> 21
 Asp Arg Phe Cys Phe Arg Lys Tyr Arg Tyr Trp His Tyr Phe Lys Phe
 1 5 10 15

<210> 22
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (3)..(3)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> D-Amino Acid

<220>
 <221> MOD_RES
 <222> (17)..(17)
 <223> Amidation

<400> 22
 His Tyr Arg Trp Lys Phe Asp Ala Arg Cys Tyr His Phe Lys Tyr His
 1 5 10 15
 Ser

<210> 23
 <211> 19

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (5)..(5)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (10)..(10)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (15)..(15)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (18)..(18)
 <223> D-Amino Acid

<400> 23
 Gly Ala Lys Phe Lys Glu Arg Tyr His Arg Arg Asp Tyr Trp His Trp
 1 5 10 15
 His Lys Asp

<210> 24
 <211> 20
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE

<222> (10)..(10)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (13)..(13)
 <223> D-Amino Acid

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> D-Amino Acid

<220>
 <221> MOD_RES
 <222> (20)..(20)
 <223> Amidation

<400> 24
 Thr Tyr Arg Lys Trp Tyr Glu Asp Lys Arg His Phe Tyr Gly Val Ile
 1 5 10 15
 His Arg Tyr Lys
 20

<210> 25
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Peptide

<220>
 <221> MOD_RES
 <222> (1)..(1)
 <223> 2',6' dimethyl

<220>
 <221> MISC_FEATURE
 <222> (2)..(2)
 <223> D-Amino Acid

<220>
 <221> MOD_RES
 <222> (5)..(5)
 <223> Amidation

<400> 25
 Tyr Arg Phe Lys Cys
 1 5

79434-30D5

CLAIMS:

1. A use of the aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ in the manufacture of a medicament for treating ureteral obstruction in a mammal in need thereof.
2. Use of the aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ for treating 5 ureteral obstruction in a mammal in need thereof.
3. The use according to any one of claims 1 to 2, wherein the mammal is a human.
4. The use according to any one of claims 1 to 3, wherein the peptide is formulated for oral, topical, systemic, intravenous, subcutaneous, intramuscular, 10 intracerebroventricular, intrathecal, or transdermal administration.
5. The use according to claim 4, wherein the transdermal administration is by iontophoresis.
6. The aromatic-cationic peptide Phe-D-Arg-Phe-Lys-NH₂ for use in treating ureteral obstruction in a mammal in need thereof.

Figure 1





Figure 2







Figure 3






Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 14




Figure 15




Figure 16




Figure 17

