

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199888639 B2**
(10) Patent No. **736737**

(54) Title
Hydrogenolysis of glycerol

(51)⁶ International Patent Classification(s)
C07C 029/60 C07C 045/52
C07C 031/20 C07C 047/22

(21) Application No: **199888639** (22) Application Date: **1998 .07 .20**

(87) WIPO No: **WO99/05085**

(30) Priority Data

(31) Number	(32) Date	(33) Country
97305508	1997 .07 .23	EP

(43) Publication Date : **1999 .02 .16**

(43) Publication Journal Date : **1999 .04 .15**

(44) Accepted Journal Date : **2001 .08 .02**

(71) Applicant(s)
Shell Internationale Research Maatschappij B.V.

(72) Inventor(s)
Eit Drent; Willem Wabe Jager

(74) Agent/Attorney
DAVIES COLLISON CAVE, 1 Little Collins Street, MELBOURNE VIC 3000

OPI DATE 16/02/99 APPLN. ID 88639/98
AOJP DATE 15/04/99 PCT NUMBER PCT/EP98/04982



AU9888639

T)

(51) International Patent Classification 6 : C07C 29/60, 31/20, 45/52, 47/22		A1	(11) International Publication Number: WO 99/05085
			(43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/EP98/04982		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD; TG).	
(22) International Filing Date: 20 July 1998 (20.07.98)			
(30) Priority Data: 97305508.0 23 July 1997 (23.07.97) EP			
(71) Applicant: SHELL INTERNATIONALE RESEARCH MAATSCHAPPIJ B.V. [NL/NL]; Carel van Bylandtlaan 30, NL-2596 HR The Hague (NL).		Published <i>With international search report.</i>	
(72) Inventors: DRENT, Eit; Badhuisweg 3, NL-1031 CM Amsterdam (NL). JAGER, Willem, Wabe; Badhuisweg 3, NL-1031 CM Amsterdam (NL).			
(54) Title: HYDROGENOLYSIS OF GLYCEROL			
(57) Abstract A process for the catalytic hydrogenolysis of glycerol in the presence of a homogeneous catalyst that is based on a platinum group metal or a compound of a platinum group metal, an anion source, and a metal-complexing compound of the formula $Q^1Q^2MQ^3$ (I) or $Q^1Q^2MQMQ^3Q^4$ (II) wherein M represents phosphorus, arsenic or antimony, Q represents a group that is covalently bonded to both M's and having at least two atoms in the bridge and Q^1 to Q^4 are independently similar or dissimilar optionally substituted hydrocarbyl groups or Q^4 and Q^2 and/or Q^3 and Q^4 represent similar or dissimilar optionally substituted hydrocarbylene groups.			

HYDROGENOLYSIS OF GLYCEROL

The invention relates to a process for the catalytic hydrogenolysis of glycerol. In particular, the invention relates to the preparation of propylene glycols (1,2- and 1,3-propanediols) and/or acrolein by the hydrogenolysis of glycerol.

Propylene glycols and acrolein are valuable chemicals. For instance, 1,3-propanediol (PDO) is an attractive monomer in the preparation of polyesters and polyurethanes. It may also be used to prepare cyclic ethers that find use as solvent. Likewise, acrolein and its dimer provide a valuable starting point for the synthesis of chemicals used in textile finishing, paper treating, and the manufacture of rubber chemicals, pharmaceuticals, plasticizers and synthetic resins. Propylene glycols and acrolein may be prepared by a variety of processes. For example, PDO may be prepared by the hydroformylation of ethylene oxide, or by the hydrogenation of 3-hydroxypropionaldehyde. However, each of these processes requires chemicals as starting point that have to be prepared separately, often at considerable costs. Besides, the starting chemicals may find other higher-value uses.

The chemicals industries have realised that our petrochemical resources are not unlimited. Therefore, they focused and still focus on natural resources as starting point for their processes. For instance, US patent No. 4,642,394 describes the process for the conversion of glycerol to lower oxygenated hydrocarbons, such as 1,2- and 1,3-propanediol, by reacting glycerol with carbon monoxide and hydrogen (in a "hydrogenolysis" reaction) in the presence of a homogeneous catalyst

-2-

containing tungsten and Group VIII metal components. The examples of this patent document, however, reveal the need for elevated temperatures and pressure conditions (200 centigrade, 4600 psig). The process is therefore not quite
5 as attractive as it could be.

The art also includes examples of hydrogenolysis processes using heterogeneous catalysts. For instance, DE-A-4,302,464 describes the conversion of glycerol into 1,2-propanediol and other products (but not 1,3-propane-diol)
10 using copper chrome tablets at various elevated temperatures and pressures. US patent No. 5,326,912 employs a catalyst containing ruthenium, palladium and copper. However, glycerol is produced rather than converted.

The inventors set out to provide a process for the
15 conversion of glycerol to lower oxygenated hydrocarbons which avoids the need for these elevated temperature and pressure conditions.

Accordingly, the invention provides a process for the catalytic hydrogenolysis of glycerol in the presence of
20 hydrogen and in the presence of a homogeneous catalyst, wherein the homogeneous catalyst is based on a platinum group metal or a compound of a platinum group metal, an anion, and a metal-complexing compound of the formula $Q^1Q^2MQ^3$ (I) or $Q^1Q^2MQMQ^3Q^4$ (II) in which M represents phosphorous,
25 arsenic or antimony, Q represents a bridging group that is covalently bonded to each M and has at least two carbon atoms in the bridge and Q^1 to Q^4 are independently the same or different optionally substituted hydrocarbyl groups or Q^1 and Q^2 and/or Q^3 and Q^4 are independently the same or
30 different optionally substituted hydrocarbylene groups.



Note that the standard textbook "Advanced Organic Chemistry", by Jerry March (3rd ed., pages 392-393) in respect of hydrogenolysis of alcohols mentions that "the hydroxyl groups of most alcohols can seldom be cleaved".
5 Examples that do undergo the reaction readily are benzyl-

2000



type alcohols. In addition, 1,3-glycols are mentioned as being especially susceptible to hydrogenolysis, whereas tertiary alcohols can be reduced by catalytic hydrogenolysis when the catalyst is platinum bis(triphenylphosphine)dichloride. It is therefore surprising that glycerol, which is not a benzyl-type alcohol, may be converted into propylene glycols and/or acrolein. Moreover, it is surprising that the conversion of glycerol may be controlled to stop at the stage where the propylene glycols and/or the acrolein are produced.

In the present specification the metals of the platinum group are defined as the metals with the atomic numbers 28, 46 and 78, i.e. nickel, palladium and platinum. Of these, palladium and platinum are preferred.

Examples of suitable metal sources are metal nitrates, sulphates, sulphonates, metal salts carboxylic acids with up to 12 carbon atoms, or inorganic metal complexes, e.g. with carbon monoxide or acetylacetonate. Palladium(II) acetate and platinum(II) acetylacetonate are examples of preferred metal sources.

The anion on which the catalyst is based may be obtained from sources such as acids and salts. The anion may also stem from platinum group metal salts, in which case both catalyst components are provided by the same source.

Preferred anion sources in the catalyst systems of the present invention are strong acids, i.e., acids having a pKa value of less than 3, preferably less than 2, measured in aqueous solution at 18 °C. The anions derived from these acids are non-coordinating or weakly coordinating with the metals of the platinum group.

Typical examples of suitable anions are anions of phosphoric acid, sulphuric acid, hydrogen halides, sulphonic acids and halogenated carboxylic acids such as trifluoroacetic acid. Sulphonic acids are in particular

preferred, for example methanesulphonic acid, trifluoro-
methanesulphonic acid, tert-butanesulphonic acid,
p-toluenesulphonic acid and 2,4,6-trimethylbenzene-
sulphonic acid. Halide anions have been found parti-
cularly useful (in combination with other anions) when
water is applied as reaction solvent.

Also, complex anions are suitable, such as the anions
generated by a combination of a Lewis acid such as BF_3 ,
 AlCl_3 , SnF_2 , $\text{Sn}(\text{CF}_3\text{SO}_3)_2$, SnCl_2 or GeCl_2 , with a protic
acid, such as a sulphonic acid, e.g. $\text{CF}_3\text{SO}_3\text{H}$ or $\text{CH}_3\text{SO}_3\text{H}$
or a hydrogen halide such as HF or HCl , or a combination
of a Lewis acid with an alcohol. Examples of such complex
anions are BF_4^- , SnCl_3^- , $[\text{SnCl}_2.\text{CF}_3\text{SO}_3]^-$ and PF_6^- .

Finally, also mixtures of anions may be used; in
particular the mixture of halide anions in combination
with another anion source mentioned above.

In the metal-complexing compounds of formula (I) or
(II), (each) M is a phosphorus atom, in which case the
compounds are monophosphines or bisphosphines.

Preferably, a metal-complexing compound of the
general formula (II) is used. The bridging group in such
a compound, represented by Q, is composed of at least two
carbon atoms. Preferably the bridging group contains two
or three carbon atoms in the bridge.

The hydrocarbyl groups Q^1 to Q^4 may independently
represent various non-cyclic or cyclic groups of up to
20 carbon atoms, preferably up to 15 carbon atoms, more
preferably up to 10 carbon atoms, optionally substituted
with substituents such as alkoxy groups with 1 to
4 carbon atoms, halogen atoms or (C_1 to C_4 alkyl)amino
groups. Examples are alkyl groups such as ethyl, iso-
propyl, sec-butyl and tert-butyl groups, cycloalkyl
groups such as cyclopentyl and cyclohexyl groups, and
aryl groups such as phenyl, tolyl and naphthyl groups.



- 5 -

In view of the higher activity of the resulting catalyst system, Q^1 together with Q^2 , and in case of a compound of general formula (II) Q^3 together with Q^4 , preferably represent an optionally substituted hydrocarbylene group. The hydrocarbylene group in general comprises at least 5 ring atoms and preferably contains from 6 to 9 ring atoms. More preferably the cyclic group contains 8 ring atoms. Substituents, if any, are alkyl groups having from 1 to 4 carbon atoms. As a rule, all ring atoms are carbon atoms, but bivalent cyclic groups containing one or two heteroatoms in the ring, such as oxygen- or nitrogen atoms, are not precluded. Examples of suitable bivalent cyclic groups are 1,4-cyclohexylene, 1,4-cycloheptylene, 1,3-cycloheptylene, 1,2-cyclooctylene, 1,3-cyclooctylene, 1,4-cyclooctylene, 1,5-cyclooctylene, 2-methyl-1,5-cyclooctylene, 2,6-dimethyl-1,4-cyclooctylene and 2,6-dimethyl-1,5-cyclooctylene groups.

The preferred metal-complexing compounds of formula (II) are 1,2-bis(1,4-cyclooctylenephosphino)ethane, 1,2-bis(1,5-cyclooctylenephosphino)ethane and mixtures thereof. For the preparation of these compounds, reference is made to known techniques, for example the method disclosed in GB-A-1,127,965.

The quantity in which the catalyst system is used, is not critical and may vary within wide limits. Usual amounts range from 10^{-8} to 10^{-1} , preferably from 10^{-7} to 10^{-2} mole atom of platinum group metal per mole of ethylenically unsaturated compound. The amounts of the participants in the catalyst system are conveniently selected such that per mole atom of platinum group metal from 0.5 to 10, preferably from 1 to 6 moles of metal-complexing compound ligand are used, and from 0.5 to 15, preferably from 1 to 8 moles of anion source or a complex anion source are used.

The hydrogenolysis process is carried out under an atmosphere that comprises or is composed of hydrogen gas. For instance, an atmosphere of carbon monoxide and hydrogen is quite suitable. These gases may be present in equimolar or non-equimolar ratios, e.g. in a ratio within the range of 5:1 to 1:5.

The hydrogenolysis can be carried out at moderate reaction conditions. Hence temperatures in the range of 50 to 250 °C are recommended, preferred temperatures being in the range of 70 to 200 °C. Reaction pressures in the range of 5 to 100 bar are preferred. Lower or higher pressures may be selected, but are not considered particularly advantageous. Moreover, higher pressures require special equipment provisions.

In the process of the invention, the starting material and the formed product may act as reaction diluent. Hence, the use of a separate solvent is not necessary. However, the hydrogenolysis reaction is conveniently carried out in the additional presence of a solvent. As such, saturated hydrocarbons, e.g. paraffins and isoalkanes are recommended and furthermore alcohols, preferably having from 3 to 10 carbon atoms per molecule, such as propanol, butanol, ethylhexanol-1, nonanol-1, or in general terms the alcohols formed as hydrogenolysis product; ethers such as 2,5,8-trioxanonane (diglyme), diethylether and anisole, and ketones, such as methylbutylketone. A particularly suitable solvent or cosolvent is water. Solvents comprising or substantially consisting of sulphones are also quite suitable. Particular preferred sulphones are, for example, dialkylsulphones such as dimethylsulphone and diethylsulphone and cyclic sulphones, such as sulfolane (tetrahydrothiophene-2,2-dioxide), sulfolene, 2-methylsulfolane and 2-methyl-4-ethylsulfolane. Mixtures of solvents may also



be used, for example a mixture of a sulphone with a protic solvent, such as an alcohol or water.

The amount of solvent to be used in the process of the invention may vary considerably. The experimental results provided hereinafter are indicative for the amount of solvent preferably to be used.

The invention will be illustrated by the non-limiting examples, as described hereinafter. The abbreviations, used in the Tables have the following meanings:

BCPE = 1,2-bis(1,5-cyclooctylenephosphino)ethane

BBPE = 1,2-bis(di-sec-butylphosphino)ethane

MSA = methanesulphonic acid

TFSA = trifluoromethanesulphonic acid

Examples 1 to 5

The experiments were carried out in a 250 ml magnetically stirred autoclave. The autoclave was charged with 30 ml glycerol, sulfolane and water in the amounts disclosed in the Table, 0.25 mmol of palladium(II) acetate, 0.6 mmol of complexing compound and anions again in the amount disclosed in the table. After being flushed, the autoclave was pressurized. Subsequently, the reactor was sealed and the contents were heated to the pre-set temperature and maintained at that temperature for 10 hours. After cooling, a sample was taken from the contents of the reactor and analysed by Gas Liquid Chromatography. Further details and the results of the analysis can be found in the Table.

The calculated conversion rate is expressed as moles of product per mole atom of platinum group metal and per hour, (mol/mol.h).

Comparative Example 1

The experiment was performed substantially according to the procedure as described above, however using 1.0 g of a heterogeneous Pd on carbon catalyst (10% Pd on C, ex. Janssen Chimica), 40 ml of glycerol and an atmosphere



of pure hydrogen gas (which should afford a better yield). The results are also set out in the Table.

Comparative Example 2

5 The experiment was performed substantially according to the procedure as described above, however using 1.0 g of a heterogeneous Ru on carbon catalyst (5% Ru on C, ex. Janssen Chimica), 40 ml of glycerol and an atmosphere of pure hydrogen gas. The results are also set out in the Table.

10 Comparative Example 3 and 4

The experiment was performed substantially according to the procedure as described in example 1 of US patent No. 4,642,394 at the conditions mentioned above.

15 The autoclave was charged with 50 ml 1-methyl-2-pyrrolidinone, 20 ml glycerol, 0.58 mmol $\text{Rh}(\text{CO})_2$ acetylacetonate, and 4 mmol H_2WO_4 . The reactor is heated to about 150 °C and maintained at 60 bar of 1:2 ($\text{CO}:\text{H}_2$) synthesis gas. After 15 hours only traces of acrolein could be detected.

20 When the Group 10 metal was replaced by $\text{Pd}(\text{CO})_2$ acetylacetonate, no products could be detected at all.

Conclusions

25 Although not optimised, the examples according to the invention outperform the comparative examples. Other remarks than can be made concern the metal-complexing compound. Thus, the preferred BCPE affords a catalyst system with a higher rate than the one based on BBPE. The acidity to the anion source affects the rate and selectivity to acrolein. The rate is also improved by the presence of halide anions. Whereas the amount of water
30 allows some control in respect of selectivity.

Table I

Example No	Ligand	Anion source (mmol)	Solvent (ml)	Temp. (°C)	CO/H ₂ (bar)	Rate (mol/mol.h)	Selectivity (%) A/B/C/D
1	BCPE	MSA (5)	sulpholane (10) + water (10)	140	20/40	12.8	-/47.4/21.8/30.8
2	BCPE	MSA (2) + HCl (0.2)	sulpholane (10) + water (5)	170	30/30	31.2	1.6/61.6/15.8/21.0
3	BCPE	MSA (2) + HI (0.2)	sulpholane (10) + water (5)	170	20/40	23.5	4.2/88.8/3.5/3.5
4	BCPE	TFSA (2) + HCl (0.2)	sulpholane (10) + water (5)	175	20/40	82.5	79.3/15.9/0.8/4.0
5	BBPE	MSA (2)	water (10)	170	20/40	19.4	0.8/56.8/21.2/21.2
C1	-	TFSA (2)	water (5)	150	-/50	<5	traces A & B
C2	-	MSA (5)	water (50)	170	-/50	<5	traces B

A = acrolein; B = 1-propanol; C = 1,2-propanediol; D = 1,3-propanediol

- 9A -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a
5 stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any
10 form of suggestion that that prior art forms part of the common general knowledge in Australia.

8
10
12
14
16
18
20
22
24
26
28
30
32
34
36
38
40
42
44
46
48
50
52
54
56
58
60
62
64
66
68
70
72
74
76
78
80
82
84
86
88
90
92
94
96
98
100



- 10 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A process for the catalytic hydrogenolysis of glycerol in the presence of hydrogen and in the presence of a
5 homogeneous catalyst, wherein the homogeneous catalyst is based on a platinum group metal or a compound of a platinum group metal, an anion source, and a metal-complexing compound of the formula $Q^1Q^2MQ^3$ (I) or $Q^1Q^2MQMQ^3Q^4$ (II) in which M represents phosphorous, arsenic or antimony, Q
10 represents a group, that is composed of carbon atoms, that is covalently bonded to each M and has at least two atoms in the bridge and Q^1 to Q^4 are independently the same or different optionally substituted hydrocarbyl groups or Q^1 and Q^2 and/or Q^3 and Q^4 are independently the same or different
15 optionally substituted hydrocarbylene groups.
2. A process as claimed in claim 1, wherein the platinum group metal is platinum or palladium.
3. A process as claimed in claim 1 or 2, wherein the anion source is an acid having a pKa value of less than 3,
20 measured in aqueous solution at 18°C.
4. A process as claimed in any one of claims 1 to 3, wherein the catalyst comprises halide anions as a source of anions.
5. A process as claimed in any one of claims 1 to 4,
25 wherein each M in the metal-complexing compound is phosphorous.
6. A process as claimed in any one of claims 1 to 5, wherein a metal-complexing compound of formula (II) is used.



5 8. A process as claimed in any one of claims 1 to 7, that
is carried out at a temperature in the range of 50 to 250°C.

10. A process as claimed in any one of claims 1 to 9, that
10 is carried out in the presence of sulpholane, water or a
mixture thereof.

11. A process according to claim 1 and substantially as
hereinbefore described with reference to any one of Examples
1 to 4.

15 DATED this 12th day of April, 2001

Shell International Research Maatschappij B.V.

by DAVIES COLLISON CAVE
Patent Attorneys for the Applicant(s)

