

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number
WO 03/013517 A1

(51) International Patent Classification⁷: **A61K 31/422**,
31/4439, 31/427, A61P 25/28, 35/00, C07D 413/04,
417/04

(21) International Application Number: PCT/EP02/08634

(22) International Filing Date: 29 July 2002 (29.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/921,751 6 August 2001 (06.08.2001) US

(71) Applicant (for all designated States except US): **PHARMACIA ITALIA S.P.A.** [IT/IT]; VIA ROBERT KOCH, 1.2, I-20152 MILANO (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CAVICCHIOLI, Marcello** [IT/FR]; 25, Rue du Professeur Patel, F-69009 LYON (FR). **PEVARELLO, Paolo** [IT/IT]; Piazza San Pietro in Ciel d'Oro, 7/A, I-27100 PAVIA (IT). **SALOM, Barbara** [IT/IT]; Via Villa, 20, I-20057 VEDANO AL LAMBRO (IT). **VULPETTI, Anna** [IT/IT]; Via Volturmo Portici, 3, I-20047 Brugherio (Milano) (IT).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: AMINOISOXAZOLE DERIVATIVES ACTIVE AS KINASE INHIBITORS



(I)

(57) Abstract: Compounds (I) which are aminoisoxazole derivatives or pharmaceutically acceptable salts thereof, together with pharmaceutical compositions comprising them are disclosed; these compounds or compositions are useful in the treatment of diseases caused by and/or associated with an altered protein kinase activity such as cancer, cell proliferative disorders, Alzheimer's disease, viral infections, auto-immune diseases and neurodegenerative disorders.

AMINOISOXAZOLE DERIVATIVES ACTIVE AS KINASE INHIBITORS

5

The present invention relates to aminoisoxazole derivatives active as kinase inhibitors and, more particularly, to 3-heteroaryl-aminoisoxazole derivatives, to a process for
10 their preparation, to pharmaceutical compositions comprising them and to their use as therapeutic agents, particularly in the treatment of diseases linked to dysregulated protein kinases.

15 The malfunctioning of protein kinases (PKs) is the hallmark of numerous diseases. A large share of the oncogenes and proto-oncogenes involved in human cancers code for PKs. The enhanced activities of PKs are also implicated in many non-malignant diseases, such as benign prostate
20 hyperplasia, familial adenomatosis, polyposis, neurofibromatosis, psoriasis, vascular smooth cell proliferation associated with atherosclerosis, pulmonary fibrosis, arthritis glomerulonephritis and post-surgical stenosis and restenosis.

25 PKs are also implicated in inflammatory conditions and in the multiplication of viruses and parasites. PKs may also play a major role in the pathogenesis and development of neurodegenerative disorders.

For a general reference to PKs malfunctioning or
30 dysregulation see, for instance, Current Opinion in Chemical Biology 1999, 3, 459 - 465.

It is an object of the invention to provide compounds which are useful in therapy as agents against a host of diseases
35 caused by and/or associated to a dysregulated protein kinase activity.

- 2 -

It is another object to provide compounds which are endowed with multiple protein kinase inhibiting activity.

The present inventors have now discovered that some 3-heteroaryl-aminoisoxazole derivatives, hereinafter shortly
5 referred to as aminoisoxazole derivatives or aminoisoxazoles, are endowed with multiple protein kinase inhibiting activity and are thus useful in therapy in the treatment of diseases associated with disregulated protein kinases.

10 More specifically, the aminoisoxazoles of this invention are useful in the treatment of a variety of cancers including, but not limited to: carcinoma such as bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall-bladder, ovary, pancreas,
15 stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell-lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy
20 cell lymphoma and Burkett's lymphoma; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; tumors of the central
25 and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas; other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoxanthoma, thyroid follicular cancer and Kaposi's sarcoma.

30 Due to the key role of PKs in the regulation of cellular proliferation, these aminoisoxazoles are also useful in the treatment of a variety of cell proliferative disorders such as, for instance, benign prostate hyperplasia, familial adenomatosis, polyposis, neuro-fibromatosis, psoriasis,
35 vascular smooth cell proliferation associated with atherosclerosis, pulmonary fibrosis, arthritis

glomerulonephritis and post-surgical stenosis and restenosis.

The compounds of the invention can be useful in the treatment of Alzheimer's disease, as suggested by the fact
5 that cdk5 is involved in the phosphorylation of tau protein (*J. Biochem.*, 117, 741-749, 1995).

The compounds of this invention, as modulators of apoptosis, may also be useful in the treatment of cancer, viral infections, prevention of AIDS development in HIV-
10 infected individuals, autoimmune diseases and neurodegenerative disorders.

The compounds of this invention may be useful in inhibiting tumor angiogenesis and metastasis.

The compounds of the invention are useful as cyclin
15 dependent kinase (cdk) inhibitors and also as inhibitors of other protein kinases such as, for instance, protein kinase C in different isoforms, Met, PAK-4, PAK-5, ZC-1, STLK-2, DDR-2, Aurora 1, Aurora 2, Bub-1, PLK, Chk1, Chk2, HER2, raf1, MEK1, MAPK, EGF-R, PDGF-R, FGF-R, IGF-R, VEGF-R,
20 PI3K, weel kinase, Src, Abl, Akt, ILK, MK-2, IKK-2, Cdc7, Nek, and thus be effective in the treatment of diseases associated with other protein kinases.

Some 5-amino-isoxazole derivatives structurally related to
25 the compounds of formula (I) are known in the art, either for pharmaceutical or non pharmaceutical use.

As an example, the international patent application WO 98/47880 discloses guanidino isoxazoles useful in the treatment of autoimmune and inflammatory diseases.


30 WO 98/28282 discloses a very broad class of widely substituted isoxazole derivatives, useful in therapy as factor Xa inhibitors.

WO 97/34881 discloses isoxazole and isoxazoline derivatives, further substituted in position 5 by penta-
35 atomic heterocyclic rings, as antitumor agents.

Nitrofuryl-isoxazole derivatives are known in the art as antibacterial and antiprotozoal agents (see, for a reference, US 3,631,169 and GB 1250219).

Other 5-aminoisoxazole derivatives further substituted in position 3 by given heteroaryl groups are known in the art. See, in particular, Chemical Abstracts 79(1973):61409 which discloses 5-amino-3-(2-aminofuryl-5-yl)-4-methoxycarbonyl-isoxazole in a study to assess the stability of 3-(nitrofuryl)isoxazoles; or Indian J. Chem., Sect. B (1984), 23B(10), 926-9 which discloses 5-amino-3-(indol-3-yl)-4-ethoxycarbonyl-isoxazoles optionally substituted in the indole moiety, as end products of synthetic schemes.

Accordingly, the present invention provides a method for treating diseases caused by and/or associated with an altered protein kinase activity, by administering to a mammal in need thereof an effective amount of an aminoisoxazole derivative represented by formula (I):



wherein

R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

X is a divalent group selected from -N(R₃)- or -O-;

Y is a divalent group selected from -CH(R₃)-, -CO-, -CONH- or -SO₂-, or Y may also be a single bond when R₂ is a hydrogen atom or a C₃-C₆ cycloalkyl group;

R₁ is a hydrogen atom or a group, optionally further substituted, selected from straight or branched C₁-C₆ alkyl,

- 5 -

C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur; the said cycloalkyl, aryl or heterocyclyl groups being
5 optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

R₂ and R₃ have, each independently, the meanings above
10 reported for R₁ or represent an optionally substituted straight or branched C₂-C₆ alkenyl or alkynyl group; and the pharmaceutically acceptable salts thereof.


In a preferred embodiment of the method described above,
15 the disease caused by and/or associated with an altered protein kinase activity is selected from the group consisting of cancer, cell proliferative disorders, Alzheimer's disease, viral infections, auto-immune diseases and neurodegenerative disorders.

20 Specific types of cancer that may be treated include carcinoma, squamous cell carcinoma, hematopoietic tumors of myeloid or lymphoid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system, melanoma, seminoma, teratocarcinoma, osteosarcoma,
25 xeroderma pigmentosum, keratoxanthoma, thyroid follicular cancer and Kaposi's sarcoma.

In another preferred embodiment of the method described above, the cell proliferative disorder is selected from the group consisting of benign prostate hyperplasia, familial
30 adenomatosis polyposis, neuro-fibromatosis, psoriasis, vascular smooth cell proliferation associated with atherosclerosis, pulmonary fibrosis, arthritis glomerulonephritis and post-surgical stenosis and restenosis.

35 In addition, the method object of the present invention, also provides tumor angiogenesis and metastasis inhibition.

The present invention further provides an aminoisoxazole derivative represented by formula (I):



5 wherein

R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

X is a divalent group selected from -N(R₃)- or -O-;

Y is a divalent group selected from -CH(R₃)-, -CO-, -CONH- or -SO₂-, or Y may also be a single bond when R₂ is a hydrogen atom or a C₃-C₆ cycloalkyl group;

R₁ is a hydrogen atom or a group, optionally further substituted, selected from straight or branched C₁-C₆ alkyl, C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur; the said cycloalkyl, aryl or heterocyclyl groups being optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

R₂ and R₃ have, each independently, the meanings above reported for R₁ or represent an optionally substituted straight or branched C₂-C₆ alkenyl or alkynyl group; and the pharmaceutically acceptable salts thereof;

30 provided that:

a) R is other than nitrofuryl; and

- 7 -

b) 5-amino-3-(2-aminofuryl-5-yl)-4-methoxycarbonyl-isoxazole and 5-amino-4-ethoxycarbonyl-3-(indol-3-yl)-isoxazole, optionally further substituted at the indole moiety, being excluded.

5

The compounds of formula (I), object of the present invention, may have asymmetric carbon atoms and may therefore exist either as racemic admixtures or as individual optical isomers.

10 Accordingly, all the possible isomers and their admixtures and of both the metabolites and the pharmaceutically acceptable bio-precursors (otherwise referred to as pro-drugs) of the compounds of formula (I), as well as any therapeutic method of treatment comprising them, are also
15 within the scope of the present invention.

In the present description, unless otherwise specified, the meaning of aromatic or non-aromatic ring system is the one conventionally adopted in organic chemistry and clearly
20 encompasses carbocyclic and heterocyclic ring systems.

With the term aryl we typically intend any aromatic either carbocyclic as well as heterocyclic hydrocarbon with 1 or 2 ring moieties, either fused or linked to each other by single bonds, wherein at least one of the carbocyclic or
25 heterocyclic rings is aromatic.

Non limiting examples of aryl groups are, for instance, phenyl, indenyl, biphenyl, α - or β -naphthyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, imidazolyl, imidazopyridyl, 1,2-methylenedioxyphenyl, thiazolyl, isothiazolyl, pyrrolyl, pyrrolyl-phenyl, furyl, phenyl-furyl, benztetrahydrofuranlyl, oxazolyl, isoxazolyl, pyrazolyl, chromenyl, thienyl, benzothienyl, isoindolinyl, benzoimidazolyl, tetrazolylphenyl, pyrrolidinyl-tetrazolyl, quinolinyl, isoquinolinyl quinoxalinyl, benzofurazanyl,
30 1,2,3-triazolyl, 1-phenyl-1,2,3-triazolyl, and the like.

- 8 -

With the term 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur we specifically intend an aromatic heterocycle such as, for instance, furan, oxazole, isoxazole, pyrrole, imidazole, 5 pyrazole, thiophene, thiazole, isothiazole, thiadiazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, and the like.

With the term 5 to 7 membered aromatic or non-aromatic carbocycle we intend, unless otherwise specified, aromatic 10 or non aromatic cyclic hydrocarbons with from 5 to 7 carbon atoms which thus include saturated, partly unsaturated or fully unsaturated carbocyclic rings.

Examples of the above carbocycles according to the invention are, for instance, cyclopentane, cyclopentene, 15 cyclopentadiene, cyclohexane, cyclohexene, 1,3- or 1,4-cyclohexadiene, benzene, cycloheptane, cycloheptene, and the like.

With the term 5 to 7 membered aromatic or non-aromatic heterocycle we intend, unless otherwise specified, any 20 saturated, partly unsaturated or fully unsaturated heterocycle, hence comprising aromatic systems also referred to as heteroaryl groups, with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur.

Therefore, apart from any 5 or 6 membered heteroaryl group 25 previously reported, examples of the above heterocycles are, for instance, 2H-pyrrole, pyran, pyrrolidine, pyrroline, imidazolidine, imidazoline, pyrazolidine, pyrazoline, piperidine, piperazine, morpholine, tetrahydrofuran, oxazolidine, oxazoline, azepine, diazepine, and the like.

30 From all of the above, it is clear to the skilled man that the term aryl refers to aromatic ring systems comprising heterocyclic and carbocyclic hydrocarbons; the term heteroaryl specifically refers to aromatic heterocyclic rings; the term aromatic or non-aromatic carbocycle 35 encompasses carbocyclic aryl groups as well as saturated or partly unsaturated, hence non-aromatic, carbocycles; the

- 9 -

term aromatic or non-aromatic heterocycle refers to heteroaryl groups and saturated or partly unsaturated, hence non-aromatic, heterocyclic rings.

5. In addition, any heteroaryl R group and any one of R_1 , R_2 or R_3 , when representing, each independently, a cycloalkyl, aryl or heterocyclyl group, may be optionally condensed through any one of their available bonds with an aromatic or non-aromatic, either carbocyclic as well as heterocyclic ring, substantially as set forth above.

Examples of R groups which are further condensed are, for instance, isobenzofuran, indole, isoindole, 1H-indazole, purine, quinoline, isoquinoline, phthalazine, quinoxaline, quinazoline, cinnoline, pteridine, benzofuran, and the like, which are all bonded through the heteroaryl moiety to the isoxazole in formula (I).

Preferably, when further condensed, the heteroaryl group is condensed with benzene or pyridine rings.

20 With the term straight or branched C_1 - C_6 alkyl we intend a group such as, for instance, methyl, ethyl, n.propyl, isopropyl, n.butyl, isobutyl, sec-butyl, tert-butyl, n.pentyl, n.hexyl and the like.

With the term straight or branched C_2 - C_6 alkenyl or alkynyl we intend an unsaturated hydrocarbon chain having a double or triple bond such as, for instance, vinyl, ethynyl, 1-propenyl, allyl, 1- or 2-propynyl, 1-, 2- or 3-butenyl, pentenyl, pentynyl, hexenyl, hexynyl and the like.

With the term C_3 - C_6 cycloalkyl group we intend cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

With the term heterocyclyl group we intend any aromatic or non-aromatic heterocyclic group, as formerly indicated.

According to the meanings provided to R, R_1 , R_2 and R_3 , any of the above groups may be further optionally substituted in any of the free positions by one or more groups, for instance 1 to 6 groups such as, for instance: halogen,

- 10 -

nitro, oxo groups (=O), carboxy, cyano, straight or branched C₁-C₆ alkyl or perfluorinated alkyl, C₃-C₆ cycloalkyl, aryl, heterocyclyl, amino, alkylamino, dialkylamino, arylamino, diarylamino, ureido, alkylureido, 5 arylureido, formylamino, alkylcarbonylamino, alkenylcarbonylamino, arylcarbonylamino, alkoxycarbonylamino, sulphonamido, alkylsulphonamido, arylsulphonamido, hydroxy, straight or branched C₁-C₆ alkoxy, aryloxy, alkylcarbonyloxy, arylcarbonyloxy, 10 alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, cycloalkyloxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl; alkylthio, arylthio, alkylsulphonyl, arylsulphonyl, arylsulphonyloxy, aminosulphonyl, alkylaminosulphonyl or 15 dialkylaminosulphonyl.

In their turn, whenever appropriate, each of the above groups may also be optionally further substituted with one or more of the aforementioned groups.

When referring, in particular, to the meanings of R, R₂ or 20 R₃, these same groups may be optionally substituted, in any of the free positions, also by straight or branched C₂-C₆ alkenyl or alkynyl groups.

Unless otherwise specified, with the term perfluorinated 25 alkyl we intend an alkyl group wherein two or more hydrogen atoms are replaced by fluorine atoms such as, for instance trifluoromethyl, 2,2,2-trifluoroethyl and the like.

With the term halogen atom we intend a fluorine, chlorine, bromine or iodine atom.

30 From all of the above, it is clear to the skilled man that any group, which name has been identified as a composite name, has to be intended as construed from the single moieties. As an example, the term alkenylcarbonylamino has to be intended as referring to a carbonylamino group which 35 is further substituted by an alkenyl group; the term heterocyclyl C₁-C₆ alkyl has to be intended as an alkyl

- 11 -

group which is substituted by a heterocyclyl group, and the like.

Pharmaceutically acceptable salts of the compounds of formula (I) are the acid addition salts with inorganic or organic, e.g. nitric, hydrochloric, hydrobromic, sulfuric, perchloric, phosphoric, acetic, trifluoroacetic, propionic, glycolic, lactic, oxalic, malonic, malic, maleic, tartaric, citric, benzoic, cinnamic, mandelic, methanesulphonic, isethionic and salicylic acid, as well as the salts with inorganic or organic bases, e.g. alkali or alkaline-earth metals, especially sodium, potassium, calcium or magnesium hydroxides, carbonates or bicarbonates, acyclic or cyclic amines, preferably methylamine, ethylamine, diethylamine, triethylamine or piperidine.

A first class of preferred compounds of the invention is represented by the compounds of formula (I) wherein Y is a single bond and R_2 is a hydrogen atom, and R, R_1 and X are as above defined.

Still more preferred, within this class, are the compounds wherein X is a $-N(R_3)-$ group and R_3 is as above defined.

Another class of preferred compounds of the invention is represented by the compounds of formula (I) wherein Y is a $-CO-$ group and R, R_1 , R_2 and X are as above defined.

Still more preferred, within this class, are the compounds wherein X is $-N(R_3)-$ and R_1 and R_3 are both hydrogen atoms.

Another class of preferred compounds of the invention is represented by the compounds of formula (I) wherein Y is a $-CONH-$ group and R, R_1 , R_2 and X are as above defined.

Still more preferred, within this class, are the compounds wherein X is $-N(R_3)-$ and R_1 and R_3 are both hydrogen atoms.

- 12 -

Another class of preferred compounds of the invention is represented by the compounds of formula (I) wherein Y is a -SO₂- group and R, R₁, R₂ and X are as above defined.

Still more preferred, within this class, are the compounds
5 wherein X is -N(R₃)- and R₁ and R₃ are both hydrogen atoms.

Preferably, in any one of the above classes, when the heteroaryl group R is further condensed, it is condensed with a benzene or pyridine ring.

10 Also preferred are the compounds of formula (I) wherein any heteroaryl or heterocyclyl group has 1 or 2 heteroatoms selected among nitrogen, oxygen or sulfur.

Specific examples of compounds of formula (I), optionally
15 in the form of pharmaceutically acceptable salts, are conveniently listed in the experimental section.



As formerly indicated, the process for preparing the compounds of formula (I) represents a further object of the
20 invention.

The amino isoxazole derivatives of formula (I), object of the invention, are thus obtainable through a synthetic process comprising well known reactions carried out according to conventional techniques, as well as through a
25 new and extremely versatile solid-phase combinatorial process, being both comprised within the scope of the invention.


The compounds of formula (I) and the pharmaceutically
30 acceptable salts thereof may be thus prepared according to a process comprising:

a) reacting a compound of formula (II) with a compound of formula (III)

- 13 -




wherein R, R₁ and X are as above defined and Z represents a halogen atom or a suitable leaving group, so as to obtain a compound of formula (I)



5

and, optionally, reacting the compound of formula (I) according to any one of the alternative steps b) below

b.1) with a compound of formula (IV), (V) or (VI)



10 wherein R₂ is as above defined, W is hydroxy or a suitable leaving group and W' is a suitable leaving group, so as to obtain the compounds of formula (I) wherein Y is -CO-, -SO₂- or -CONH-, respectively, and R, R₁, R₂ and X are as above defined;

15 b.2) with a suitable aldehyde or ketone derivative of formula (VII)



wherein R₂ and R₃ are as above defined, under reductive conditions, so as to obtain a compound of formula (I) wherein Y is a group -CH(R₃)- and R, R₁, R₂, R₃ and X are as above defined;

20 b.3) with a suitable acylating agent in the presence of ammonia, so as to obtain a compound of formula (I) wherein Y is -CONH-, R₂ is hydrogen and R, R₁ and X are as above defined; and, optionally

25 c) converting the thus obtained compound of formula (I) into another compound of formula (I) and/or into a pharmaceutically acceptable salt thereof.

The above process is an analogy process which can be carried out according to well known methods.

It is clear to the person skilled in the art that if a compound of formula (I), prepared according to the above process, is obtained as an admixture of isomers, their separation into the single isomers of formula (I), carried out according to conventional techniques, is still within the scope of the present invention.

Likewise, the conversion into the free compound (I) of a corresponding salt thereof, according to well-known procedures in the art, is still within the scope of the invention.

According to step a) of the process, the compound of formula (II) is reacted with a compound of formula (III).

This reaction is carried out in the presence of a base such as triethylamine, N,N-diisopropylethylamine, 1,8-diazabicyclo[5.4.0]undec-7-ene, pyridine, lithium diisopropylamide or lithium bis(trimethylsilyl)amide, and in a suitable solvent such as, for instance, dichloromethane, chloroform, tetrahydrofuran, acetonitrile, diethyl ether, 1,4-dioxane or N,N-dimethylformamide, at a temperature ranging from about -40°C to room temperature and for a suitable time, for instance from about 30 minutes to about 96 hours.

The 5-aminoisoxazoles of formula (I) thus obtained can be easily converted into a variety of derivatives of formula (I) and/or into salts thereof.

30

According to step b.1) of the process, the compound of formula (I) is reacted with any one of the compounds of formula (IV), (V) or (VI), so as to obtain the corresponding derivative of formula (I). In this respect, it is clear to the skilled man that a carboxamido derivative of formula (I) wherein Y is (>C=O) is obtained

35

- 15 -

through reaction with a compound of formula (IV); a sulphonamido derivative of formula (I) wherein Y is (-SO₂-) is obtained through reaction with a compound of formula (V); and an ureido derivative of formula (I) wherein Y is (-NHCO-) is obtained through reaction with a compound of formula (VI).

As formerly indicated, within the compound of formula (IV) W is hydroxy or a suitable leaving group such as, for instance, a halogen atom. Preferably, W is a hydroxy group or a chlorine or bromine atom.

The reaction between a compound of formula (I) and a carboxylic acid derivative of formula (IV) wherein W is hydroxy can be carried out in the presence of a condensing agent such as, for instance, benzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphatecarbodiimide, 1,3-dicyclohexylcarbodiimide, bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, 1,3-diisopropylcarbodiimide, o-benzotriazol-1-yl-n,n,n',n'-tetra-methyluronium tetrafluoroborate, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, N-cyclohexylcarbodiimide-N'-propyloxymethyl polystyrene or N-cyclohexylcarbodiimide-N'-methyl polystyrene, in a suitable solvent such as, for instance, dichloromethane, chloroform, tetrahydrofuran, diethyl ether, 1,4-dioxane, acetonitrile, toluene or N,N-dimethylformamide, at a temperature ranging from about -10°C to reflux and for a suitable time ranging from about 30 minutes to about 96 hours.

The said reaction is optionally carried out in the presence of a suitable catalyst, for instance 4-dimethylaminopyridine, or in the presence of a further condensing agent such as N-hydroxybenzotriazole. The reaction between a compound of formula (I) and a compound of formula (IV) can be also carried out through a mixed anhydride method, that is by using an alkyl chloroformate such as ethyl, isobutyl, or isopropyl chloroformate, in the presence of a tertiary base such as triethylamine, N,N-

- 16 -

diisopropylethylamine or pyridine, in a suitable solvent such as toluene, dichloromethane, chloroform, tetrahydrofuran, acetonitrile, diethyl ether, 1,4-dioxane or N,N-dimethylformamide, and at a temperature ranging from
5 about -30°C to room temperature.

The reaction between a compound of formula (I) and a compound of formula (IV) wherein W is a suitable leaving group; for instance chlorine or bromine, can be carried out in the presence of a tertiary base such as triethylamine,
10 N,N-diisopropylethylamine or pyridine, in a suitable solvent such as toluene, dichloromethane, chloroform, diethyl ether, tetrahydrofuran, acetonitrile or N,N-dimethylformamide, and at a temperature ranging from about -10°C to reflux.

15 The said reaction is optionally carried out in the presence of a suitable catalyst, for instance 4-dimethylaminopyridine, or in the presence of a further condensing agent such as N-hydroxybenzotriazole.

As formerly indicated, within the compound of formula (V)
20 W' is a suitable leaving group such as, for instance, a halogen atom; chlorine or bromine being preferred.

The reaction between a compound of formula (I) and a sulphonyl derivative of formula (V) can be carried out in the presence of a tertiary base such as triethylamine, N,N-
25 diisopropylethylamine or pyridine, in a suitable solvent such as toluene, dichloromethane, chloroform, diethyl ether, tetrahydrofuran, acetonitrile or N,N-dimethylformamide, at a temperature ranging from about -10°C to reflux.

30 The said reaction is optionally carried out in the presence of a suitable catalyst, for instance 4-dimethylaminopyridine, or in the presence of a further condensing agent such as N-hydroxybenzotriazole.

The reaction between a compound of formula (I) and an
35 isocyanate derivative of formula (VI) can be carried out in the presence of a tertiary base such as triethylamine, N,N-

- 17 -

diisopropylethylamine or pyridine, in a suitable solvent such as toluene, dichloromethane, chloroform, diethyl ether, tetrahydrofuran, acetonitrile, or N,N-dimethylformamide, and at a temperature ranging from about
5 -10°C to reflux.

According to step b.2) of the process, the compound of formula (I) is reacted, under reductive conditions, with an aldehyde or ketone derivative of formula (VII) so as to
10 obtain the corresponding compound of formula (I) wherein Y is as above defined.

From the above, it is clear to the skilled man that by reacting an aldehyde derivative of formula (VII), for instance having R₃ as a hydrogen atom, the corresponding
15 derivative of formula (I) wherein Y is a -CH₂- group may be obtained. Likewise, by reacting a ketone derivative of formula (VII) wherein both R₂ and R₃ are other than hydrogen, the corresponding derivative of formula (I) wherein Y is a -CH(R₃)- group and R₃ is other than hydrogen,
20 may be obtained.

By analogy, if the ketone derivative of formula (VII) is represented by a compound (R₂)=O wherein R₂ is a heterocyclyl or cyclolalkyl moiety, e.g. cyclohexanone, the above reaction according to step b.2) gives rise to a
25 corresponding derivative of formula (I) wherein Y is a single bond and R₂ is just the said cyclolalkyl or heterocyclyl group.

All of these reactions are widely known in the art as reductive alkylation of amines and occur in the presence of
30 a reducing agent such as, for instance, sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride, in a suitable solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, chloroform, dichloromethane, tetrahydrofuran or acetonitrile, optionally in the presence
35 of acetic acid, methanol or ethanol as co-solvents, at a

- 18 -

temperature ranging from about -10°C to reflux and for a time varying from about 30 minutes to about 96 hours.

According to step b.3) of the process, ureido derivatives
5 of formula (I) wherein Y is $-\text{NHCO}-$ and R_2 is a hydrogen atom may be prepared by reacting the compound of formula (I) with a suitable acylating agent such as, for instance, triphosgene or trichloromethyl chloroformate, in the presence of aqueous or gaseous ammonia, according to
10 conventional techniques for preparing ureido $-\text{NHCONH}_2$ derivatives starting from the corresponding amino $-\text{NH}_2$ derivatives.

Finally, according to step c) of the process and whenever
15 desired, a given compound of formula (I) may be optionally converted into another compound of formula (I), by working according to well known methods.

Just as an example, a given 4-carboxy-isoxazole derivative of formula (I), wherein X is $-\text{O}-$ and R_1 is hydrogen, may be
20 easily converted into the corresponding ester of formula (I) wherein X is $-\text{O}-$ and R_1 is other than hydrogen, for instance an alkyl group, or into the corresponding carboxamido derivative of formula (I) wherein X is $-\text{NH}(\text{R}_3)-$ and, as an example, R_3 is a hydrogen atom and R_1 is an alkyl
25 group.

In this specific case, the above reactions may be performed according to well known methods of esterification or amidation, by reacting the given carboxy derivative of formula (I) with a suitable alcohol R_1-OH (VIII) or amino
30 derivative $\text{R}_1-\text{NH}(\text{R}_3)$ (IX), both bearing the desired R_1 and R_3 groups.

Likewise, the optional salification of a compound of formula (I) or the conversion of its salt into the free compound, may be all carried out by conventional methods.

- 19 -

The compounds of formula (II) and (III) according to the process object of the present invention are known compounds which are easily prepared according to known methods.

As an example, the compounds of formula (II) wherein Z is a chlorine atom may be prepared from the corresponding oximes, wherein Z is hydrogen, by working as described in J. Org. Chem., (1980), 3916; or J. Org. Chem., (1992), 6649.

The oximes, in their turn, are commercially available or readily obtainable from the corresponding aldehyde derivatives, according to conventional methods [see, for a reference, Org. Synth. Coll., 2, 70, 313 (1955)].

Also the compounds of formula (III) are commercially available or they may be prepared according to conventional techniques for carboxylic ester or carboxamide syntheses, by reacting cyanoacetic acid or a suitable derivative thereof with a compound of formula (VIII) or (IX), as above defined.

Likewise, all of the compounds of formula from (IV) to (IX) are known or may be prepared according to conventional methods.

As it will be really appreciated by the man skilled in the art, when preparing the compounds of formula (I) object of the invention, optional functional groups within both the starting materials or the intermediates thereof, which could give rise to unwanted side reactions, need to be properly protected according to conventional techniques. Likewise, the conversion of these latter into the free deprotected compounds may be carried out according to known procedures.

According to a preferred aspect of the process of the invention, the compounds of formula (I) may be also prepared by working on solid phase synthesis (SPS).


- 20 -

Therefore, it is a further object of the invention a process for preparing the compounds of formula (I), and the pharmaceutically acceptable salts thereof, which process comprises:

- 5 a') reacting cyanoacetic acid with a suitable polystyrenic resin of formula (X) or (XI)




wherein R_1 is as above defined and W is hydroxy or a
 10 suitable leaving group, so as to obtain the compounds of formula (XII) or (XIII), respectively




wherein R_1 is as above reported;

- b') reacting any one of the compounds of formula (XII) or
 15 (XIII) with a compound of formula (II)




wherein R and Z are as above defined, so as to obtain a compound of formula (XIV) or (XV)





- 20 c') cleaving the resin from the compounds of formula (XIV) or (XV) under acidic or basic conditions, so as to obtain a compound of formula (I) wherein X is a group -O- and R_1 is hydrogen, or X is a group -N(R_3)- wherein R_3 is hydrogen and R_1 is as above defined, respectively; or, alternatively,

- 21 -

reacting the above compounds of formula (XIV) or (XV) according to any one of the alternative steps d') below d.1') with a compound of formula (IV), (V) or (VI)



- 5 wherein R_2 is as above defined, W is hydroxy or a suitable leaving group and W' is a suitable leaving group, so as to obtain the compounds of formula (XVI) or (XVII)



- wherein Y is -CO-, -SO₂- or -CONH-, respectively, and R, R_1 and R_2 are as above defined;

d.2') with a suitable aldehyde or ketone derivative of formula (VII)



- wherein R_2 and R_3 are as above defined, under reductive conditions, so as to obtain the compounds of the above formula (XVI) or (XVII) wherein Y is a group -CH(R_3)- and R, R_1 , R_2 and R_3 are as above defined;

- d.3') with a suitable acylating agent in the presence of ammonia, so as to obtain a compound of the above formula (XVI) or (XVII) wherein Y is -CONH-, R_2 is hydrogen and R and R_1 are as above defined;

- e') cleaving the resin from the compounds of formula (XVI) or (XVII) under acidic or basic conditions, so as to obtain a compound of formula (I) wherein X is a group -O- and R_1 is hydrogen, or X is a group -N(R_3)- wherein R_3 is hydrogen and R_1 is as above defined, respectively; and, optionally,

f') converting the thus obtained compound of formula (I) into another compound of formula (I) and/or into a pharmaceutically acceptable salt thereof.

According to step a') of the process, cyanoacetic acid is reacted with a suitable polystyrenic resin of formula (X) or (XI) according to conventional operative conditions, in the presence of a base and of optional condensing agents or
5 catalysts.

Likewise, according to step b') of the process, the thus obtained compounds of formula (XII) or (XIII) are then
reacted with a compound of formula (II), substantially as set forth above per step a) of the process under
10 homogeneous conditions.

Analogous considerations apply to the alternative steps from d.1') to d.3') which are all carried out by working as described in corresponding steps from b.1) to b.3) of the process under homogeneous conditions.

15

According to any one of steps c') or e') of the process, the compounds of formula (I) can be obtained by cleaving the compounds of formula (XIV), (XV), (XVI) or (XVII) from the resin, under acidic or basic conditions.

20 The acidic cleavage can be performed in the presence of suitable acids such as, for instance, hydrochloric, trifluoroacetic, methanesulphonic or p-toluensulphonic acid, as well as by using conventional acid ion exchange resins. The reaction is carried out under conventional
25 methods, for instance by using a solution of the acid, e.g. a 10% to 100% (v/v) solution of trifluoroacetic acid in dichloromethane, at a temperature ranging from about 0°C to reflux, and for a suitable time, for instance from about 5 minutes to about 2 hours.

30 The basic cleavage can be performed under basic hydrolysis conditions, for instance in the presence of a conventional aqueous base such as sodium, potassium or lithium hydroxide. The reaction is carried out in a suitable solvent such as, for instance, N,N-dimethylformamide,
35 ethanol, methanol isopropanol or tetrahydrofuran, at a temperature comprised from about 20°C to reflux and for a

- 23 -

suitable time, for instance from about 30 minutes to about 96 hours.

Finally, according to step f'), any obtained compound of
5 formula (I) may be optionally converted into another
compound of formula (I) by working according to
conventional methods, for instance as reported in step c)
of the process under homogeneous conditions.

From all of the above it is clear to the skilled man that
10 the above optional conversion of a compound of formula (I)
into another compound of formula (I) may be also carried
out after step c'), that is after having carried out the
resin cleavage from the compounds of formula (XIV) or (XV).

15 The polystyrenic resins of formula (X) or (XI) are
commercially available and comprise, for instance, Wang
resin, trityl resin, Cl-trityl resin, Rink amide resin
Tentagel OH resin and derivatives thereof.

The resins of formula (XI), wherein R_1 is other than
20 hydrogen may be also prepared according to conventional
methods, for instance by reacting a commercially available
formyl polystyrene resin with a suitable amine derivative
 R_1-NH_2 of formula (IX) under reductive conditions, as set
forth above.

25 Cyanoacetic acid is also a known, commercially available,
compound.

From what above reported, it is clear to the man skilled in
the art that the compounds of formula (I) of the invention
30 can be advantageously prepared by performing the above
described reactions in a combinatorial fashion, for example
according to the aforementioned solid-phase-synthesis (SPS)
techniques, so as to get a combinatorial library of
compounds.


35 As an example, the compounds of formula (XII) or (XIII)
supported onto resin particles and prepared as above

- 24 -

described, may be reacted with a variety of compounds of formula (II) so as to obtain a plurality of compounds of formula (XIV) or (XV), to be further reacted with a variety of compounds, for instance of formula (IV), (V), (VI) or (VII), so as to get thousands of different compounds of formula (XVI) or (XVII), according to combinatorial chemistry methods.

These latter derivatives, in their turn, are then conveniently converted into the derivatives of formula (I) of the invention, by cleaving the resin support as formerly reported.

It is therefore a further object of the invention a library of two or more compounds of formula (I)



15

wherein

R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

X is a divalent group selected from -N(R₃)- or -O-;

Y is a divalent group selected from -CH(R₃)-, -CO-, -CONH- or -SO₂-, or Y may also be a single bond when R₂ is a hydrogen atom or a C₃-C₆ cycloalkyl group;

R₁ is a hydrogen atom or a group, optionally further substituted, selected from straight or branched C₁-C₆ alkyl, C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur; the

30

- 25 -

said cycloalkyl, aryl or heterocyclyl groups being optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

R_2 and R_3 have, each independently, the meanings above reported for R_1 or represent an optionally substituted straight or branched C_2 - C_6 alkenyl or alkynyl group; and the pharmaceutically acceptable salts thereof.

10

All of the compounds of formula (I) which are prepared according to combinatorial chemistry techniques, for instance as reported in the examples, whenever appropriate in the form of pharmaceutically acceptable salts, are herewith conveniently indicated and defined as "products by process", that is as compounds of formula (I) which are obtainable through a given process.

15

As such, it is a further object of the present invention any specific compound of formula (I) which is obtainable, for instance through a combinatorial chemistry technique, by reacting each of the derivatives of formula (II), as set forth in *table I*, with any one of the derivatives of formula (XIII) which are obtainable as above indicated from the amines of formula (IX), as set forth in *table II*, and by subsequently operating as per the process of the invention by reacting any one of the obtained derivatives of formula (XV) with any one of the derivatives of formula (IV), as set forth in *table III*.

25

30

It is a further object of the present invention any specific compound of formula (I) which is obtainable, for instance through a combinatorial chemistry technique, by reacting each of the derivatives of formula (II), as set forth in *table I*, with any one of the derivatives of formula (XIII) which are obtainable as above indicated from


35

- 26 -

the amines of formula (IX), as set forth in table II, and by subsequently operating as per the process of the invention by reacting any one of the obtained derivatives of formula (XV) with any one of the derivatives of formula (V), as set forth in table IV.

It is a further object of the present invention any specific compound of formula (I) which is obtainable, for instance through a combinatorial chemistry technique, by reacting each of the derivatives of formula (II), as set forth in table I, with any one of the derivatives of formula (XIII) which are obtainable as above indicated from the amines of formula (IX), as set forth in table II, and by subsequently operating as per the process of the invention by reacting any one of the obtained derivatives of formula (XV) with any one of the derivatives of formula (VI), as set forth in table V.

Table I: Compounds of formula (II)



20

1	3-Thiophenylhydroxyaminomethyl chloride
2	Pyridine-3-hydroxyaminomethyl chloride
3	Pyridine-2-hydroxyaminomethyl chloride
4	2-Thiophenehydroxyaminomethyl chloride
5	Pyrrole-2-hydroxyaminomethyl chloride
6	5-Methylfurfurhydroxyaminomethyl chloride
7	Indole-3-hydroxyaminomethyl chloride
8	5-bromo-2-thiophenehydroxyaminomethyl chloride
9	1-methylpyrrole-2-hydroxyaminomethyl chloride
10	4-pyridinehydroxyaminomethyl chloride
11	3-methylthiophene-2-hydroxyaminomethyl chloride
12	5-methyl-2-thiophenehydroxyaminomethyl chloride
13	6-methyl-2-pyridinehydroxyaminomethyl chloride
14	5-methoxyindole-3-hydroxyaminomethyl chloride
15	1-methylindole-3-hydroxyaminomethyl chloride
16	4-bromothiophene-2-hydroxyaminomethyl chloride
17	N-ethyl-carbazole-3-hydroxyaminomethyl chloride
18	Furfurhydroxyaminomethyl chloride

19	3-furhydroxyaminomethyl chloride
20	5-fluoroindole-3-hydroxyaminomethyl chloride
21	5-ethyl-2-thiophenehydroxyaminomethyl chloride
22	3-phenyl-1H-pyrazole-4-hydroxyaminomethyl chloride
23	Imidazole-4-hydroxyaminomethyl chloride
24	Benzo[b]furan-2-hydroxyaminomethyl chloride
25	Indole-7-hydroxyaminomethyl chloride
26	4-benzyloxyindole-3-hydroxyaminomethyl chloride
27	3-methylbenzo[b]thiophene-2-hydroxyaminomethyl chloride
28	3,5-dimethyl-1-phenylpyrazole-4-hydroxyaminomethyl chloride
29	Thiazole-2-hydroxyaminomethyl chloride
30	7-methylindole-3-hydroxyaminomethyl chloride
31	2,5-dimethyl-1-[3-(trifluoromethyl)phenyl]pyrrole-3-hydroxyaminomethyl chloride
32	5-acetoxymethyl-2-furhydroxyaminomethyl chloride
33	5-[2-(trifluoromethyl)phenyl]furfurhydroxyaminomethyl chloride
34	5-(4-chlorophenyl)furfurhydroxyaminomethyl chloride
35	5-(2-chlorophenyl)furfurhydroxyaminomethyl chloride
36	5-(4-nitrophenyl)-2-furhydroxyaminomethyl chloride
37	5-(2-nitrophenyl)furfurhydroxyaminomethyl chloride
38	5-bromo-2-furhydroxyaminomethyl chloride
39	5-(4-bromophenyl)furfurhydroxyaminomethyl chloride
40	5-(3-chlorophenyl)furfurhydroxyaminomethyl chloride
41	5-[2-chloro-5-(trifluoromethyl)phenyl]-furfurhydroxyaminomethyl chloride
42	5-(3-trifluoromethylphenyl)furan-2-hydroxyaminomethyl chloride
43	5-ethyl-2-furhydroxyaminomethyl chloride
44	5-chloro-2-thiophenehydroxyaminomethyl chloride
45	2-chloro-3-quinolinehydroxyaminomethyl chloride
46	2,4-dimethoxy-5-pyrimidinehydroxyaminomethyl chloride
47	N-hydroxy-1-methyl-1H-indole-2-carboximidoyl chloride
48	Methyl 3-[chloro(hydroxyimino)methyl]-2-methyl-1H-indole-6-carboxylate
49	N-hydroxy-6-methoxypyridine-3-carboximidoyl chloride
50	N-hydroxy-1-benzofuran-2-carboximidoyl chloride

51	5-chloro-N-hydroxyfuran-2-carboximidoyl chloride
52	N-hydroxy-1,3-thiazole-5-carboximidoyl chloride

Table II: Compounds of formula (IX)

R₁-NH₂ (IX)

1	2-aminofluorene
2	Cyclopropylamine
3	Cyclopentylamine
4	2-(2-aminoethyl)-1-methylpyrrolidine
5	2-(1-pyrrolidinyl)ethylamine
6	Furfurylamine
7	(+/-)-1-(1-naphthyl)ethylamine
8	2-aminothiazole
9	1-(2-aminophenyl)pyrrole
10	Tetrahydrofurfurylamine
11	2-amino-5-chlorobenzoxazole
12	4-aminobenzo-2,1,3-thiadiazole
13	2-aminopyrimidine
14	4-(4-aminophenyl)morpholine
15	3-amino-2-chloropyridine
16	2-aminopyridine
17	2-amino-5-chloropyridine
18	4-(aminomethyl)pyridine
19	Ethyl 4-amino-1-piperidinecarboxylate
20	3,4-ethylenedioxyaniline
21	Aniline
22	2-(methylthio)aniline
23	3,4-difluoroaniline
24	3-chloro-4-methylaniline
25	m-anisidine
26	3-aminobenzotrifluoride
27	3,4-dimethylaniline
28	4-fluoro-2-methylaniline
29	N1-(4-aminophenyl)-N1-methylacetamide
30	N,N-diethyl-p-phenylenediamine
31	4-butoxyaniline
32	2-aminopentane
33	2-methoxybenzylamine
34	4-fluorobenzylamine
35	4-chlorobenzylamine
36	N,N-dimethylethylenediamine
37	2-(4-methoxyphenyl)ethylamine
38	5-amino-2-methylisoindoline-1,3-dione
39	3-methoxypropylamine
40	Sulfapyridine
41	3-phenoxyaniline
42	2-amino-4-methoxy-6-methylpyrimidine
43	4-(difluoromethoxy)aniline
44	N-(4'-aminophenyl)-4-methyl-benzene-sulfonamide

45	1,2,3,4-Tetrahydro-1-Naphthylamine
46	1,2-Dimethylpropylamine
47	1,2-Diphenylethylamine
48	1,3-Dimethylbutylamine
49	1-Aminoindan
50	1-Methyl-3-Phenylpropylamine
51	1-Naphthalenemethylamine
52	2-(1-Cyclohexenyl)ethylamine
53	2-(3-Chlorophenyl)ethylamine
54	2-(Trifluoromethyl)Benzylamine
55	2,4-Dimethoxybenzylamine
56	2-Amino-6-Methylheptane
57	2-Aminoheptane
58	2-Aminooctane
59	2-Bromobenzylamine
60	2-Chlorobenzylamine
61	2-Ethylhexylamine
62	2-Fluorobenzylamine
63	2-Methoxyisopropylamine
64	2-Methylbenzylamine
65	2-Methylcyclohexylamine
66	3-(Di-N-Butylamino) Propylamine
67	3-(Methylthio) Propylamine
68	3-(Trifluoromethyl) Benzylamine
69	3,3-Diphenylpropylamine
70	3,4-Difluorobenzylamine
71	3,4-Dimethoxybenzylamine
72	3,4-Methylenedioxybenzylamine
73	3,5-Bis(Trifluoromethyl) Benzylamine
74	3-Aminopentane
75	3-Chlorobenzylamine
76	3-Fluorobenzylamine
77	3-Methylbenzylamine
78	3-Phenylpropylamine
79	4-(Trifluoromethoxy) Benzylamine
80	4-(Trifluoromethyl) Benzylamine
81	4-Amino-1-Benzylpiperidine
82	4-Methoxybenzylamine
83	4-Methylbenzylamine
84	4-Phenylbutylamine
85	4-Tert-Butylcyclohexylamine
86	5-Methylfurfurylamine
87	Allylamine
88	Aminomethylcyclohexane
89	Benzhydrylamine
90	Benzylamine
91	Cyclobutylamine
92	Cycloheptylamine
93	Cyclohexylamine
94	Cyclopropanemethylamine
95	DL-Alpha-Methylbenzylamine

96	Hexylamine
97	Isoamylamine
98	Isobutylamine
99	Isopropylamine
100	n-Amylamine
101	n-Butylamine
102	n-Heptylamine
103	n-Octylamine
104	n-Propylamine
105	Phenethylamine
106	Propargylamine
107	Sec-Butylamine
108	Tert-Butylamine
109	Tetrahydrofurfurylamine
110	Thiophene-2-Methylamine

Table III: Compounds of formula (IV)

R₂-COW (IV)

1.	9-fluorene-carboxylic acid;
2.	1-phenyl-1-cyclopropanecarboxylic acid;
3.	1-methylcyclopropane-1-carboxylic acid;
4.	Cyclobutanecarboxylic acid;
5.	cyclopentanecarboxylic acid;
6.	(-)-menthoxyacetic acid;
7.	1,2,3,4-tetrahydro-2-naphthoic acid;
8.	2-fluorobenzoic acid;
9.	2,5-dimethoxybenzoic acid;
10.	2-biphenylcarboxylic acid;
11.	2-(4-chlorobenzoyl)benzoic acid;
12.	2,6-dimethylbenzoic acid;
13.	3-cyanobenzoic acid;
14.	3-bromobenzoic acid;
15.	3,4-dimethoxybenzoic acid;
16.	3,4,5-trimethoxybenzoic acid;
17.	3,4-diethoxybenzoic acid;
18.	4-cyanobenzoic acid;
19.	4-iodobenzoic acid;
20.	4-diethylaminobenzoic acid;
21.	4-biphenylcarboxylic acid;
22.	3-methyl-2-oxovaleric acid;
23.	pyruvic acid;
24.	2-methylvaleric acid;
25.	tert-butylacetic acid;
26.	3-(2-methoxyphenyl)propionic acid;
27.	5-nitro-2-furoic acid;
28.	1-naphthoic acid;
29.	2-naphthoic acid;
30.	2-ketobutyric acid;

31.	pivalic acid;
32.	2,2-dimethylbutyric acid;
33.	diphenylacetic acid;
34.	N,N-dimethylglycine;
35.	2,3-dichlorophenoxyacetic acid;
36.	phenylacetic acid;
37.	2,4-dichlorophenylacetic acid;
38.	3-fluorophenylacetic acid;
39.	4-ethoxyphenylacetic acid;
40.	p-tolylacetic acid;
41.	4-pentynoic acid;
42.	mono-methyl glutarate;
43.	monomethyl adipate;
44.	6-acetamidohexanoic acid;
45.	1-pyroglutamic acid;
46.	3-furoic acid;
47.	thiophene-3-carboxylic acid;
48.	thiophene-3-acetic acid;
49.	nicotinic acid;
50.	nalidixic acid;
51.	2-nitro-4-trifluoromethylbenzoic acid;
52.	4-methyl-3-nitrobenzoic acid;
53.	3-nitrobenzoic acid;
54.	3-nitrophenylacetic acid;
55.	4-carboxybenzenesulfonamide;
56.	succinamic acid;
57.	N-(4-nitrobenzoyl)-beta-alanine;
58.	3-(phenylsulfonyl)propionic acid;
59.	2,2,3,3-tetramethylcyclopropanecarboxylic acid;
60.	2-(4-nitrophenyl)propionic acid;
61.	2,2-dimethyl-4-pentenoic acid;
62.	3-(diethylamino)propionic acid hydrochloride;
63.	4-dimethylaminobutyric acid hydrochloride;
64.	4-isopropylphenoxyacetic acid;
65.	5-benzoylpentanoic acid;
66.	4-acetamido-3-nitrobenzoic acid;
67.	d-campholic acid;
68.	2,5-dibromobenzoic acid;
69.	3-acetoxybenzoic acid;
70.	2,4,6-trimethoxyphenylacetic acid;
71.	2-benzyloxyphenylacetic acid;
72.	(3,5-dimethoxyphenyl)acetic acid;
73.	2-nitrophenoxyacetic acid;
74.	chromone-3-carboxylic acid;
75.	N-acetyl-4-fluoro-dl-phenylalanine;
76.	N-m-tolylphthalamic acid;
77.	4-acetamidobutyric acid;
78.	3-(2-thenoyl)-propionic acid;
79.	3,5-diacetamidobenzoic acid;
80.	5-acetamido-2-nitrobenzoic acid;
81.	acetic acid;

82.	5-methylhexanoic acid;
83.	N-benzoyl-L-alanine;
84.	4-bromo-3-methylbenzoic acid;
85.	4,5-dibromothiophene-2-carboxylic acid;
86.	2-acetamido-5-bromobenzoic acid;
87.	4-bromo-2-methylbenzoic acid;
88.	2-fluoro-6-iodobenzoic acid;
89.	2-furanyloxyacetic acid;
90.	N,N-dimethylsuccinamic acid;
91.	2-(2-methoxyethoxy)acetic acid;
92.	4-chloro- α -methylphenylacetic acid;
93.	1-(p-tolyl)-1-cyclopentanecarboxylic acid;
94.	picolinic acid hydrochloride;
95.	3,5-dibromobenzoic acid;
96.	5-chlorothiophene-3-acetic acid;
97.	2-nitrothiophene-4-carboxylic acid;
98.	3-chloro-2-methylbenzoic acid;
99.	2-bromo-4-fluorobenzoic acid;
100.	3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid;
101.	fenbufen;
102.	indoprofen;
103.	chrysanthemum monocarboxylic acid;
104.	6-acetoxy-2-naphthoic acid;
105.	3-methylthiopropionic acid;
106.	(r) - (+) -N-(1-phenylethyl)phthalamic acid;
107.	α -ketovaleric acid;
108.	5-methyl-1-phenylpyrazole-4-carboxylic acid;
109.	3-methyl-1-cyclohexanecarboxylic acid;
110.	3-methoxycyclohexanecarboxylic acid;
111.	dicyclohexylacetic acid;
112.	5,6-dichloronicotinic acid;
113.	4-(dimethylamino)phenylacetic acid;
114.	(r) - (+) -N-(1-phenylethyl)succinamic acid;
115.	(s) - (-) -N-(1-phenylethyl)succinamic acid;
116.	(+)-menthyloxyacetic acid;
117.	suprofen;
118.	N,N-dimethyl-L-phenylalanine;
119.	4-iodophenylacetic acid;
120.	4-(3,4-dimethoxyphenyl)butyric acid;
121.	2-fluoro-5-nitrobenzoic acid;
122.	N,N-diethyl-3,6-difluorophthalamic acid;
123.	2-bromo-5-nitrobenzoic acid;
124.	4-bromo-2-fluorobenzoic acid;
125.	5-(2-thienyl)pentanoic acid;
126.	isoxazole-5-carboxylic acid;
127.	5-nitrothiophene-2-carboxylic acid;
128.	2-(4-pyridyl)thiazole-4-carboxylic acid;
129.	2-methyl-4,4,4-trifluorobutyric acid;
130.	1-(aminocarbonyl)-1-cyclopropanecarboxylic acid;

131	1-cyanocyclopropanecarboxylic acid;
132	(s) - (-) -2-acetoxypionic acid;
133	3-(methylsulfonyl)benzoic acid;
134	2-chloro-4-methylsulfonylbenzoic acid;
135	2,6-dichloropyridine-4-carboxylic acid;
136	3-pyridinepropionic acid;
137	5-(4-chloro-2-nitrophenyl)-2-furoic acid;
138	7-chloro-1-ethyl-6-fluoro-4-oxohydroquinoline-3-carboxylic acid;
139	cis-2-(2-thiophenecarbonyl)-1-cyclohexanecarboxylic acid;
140	5-bromo-3-pyridylacetic acid;
141	5-methylisoxazole-4-carboxylic acid;
142	2,2-dimethylhexanoic acid;
143	3-carboxypropanesulfonamide;
144	6-cyanonicotinic acid;
145	(r) - (-) -2-methoxypropionic acid;
146	(s) - (+) -2-methoxypropionic acid;
147	4-(tert-butoxymethyl)benzoic acid;
148	cis-2-(benzyloxycarbonylamino)-cyclohexanecarboxylic acid;
149	cis-2-(benzyloxycarbonylamino)-4-cyclohexene-1-carboxylic acid.

Table IV: Compounds of formula (V)

R₂-SO₂W' (V)

1.	1-naphthalenesulfonyl chloride
2.	2-naphthalenesulfonyl chloride
3.	2-thiophenesulfonyl chloride
4.	8-quinolinesulfonyl chloride
5.	benzenesulfonyl chloride
6.	2,4,5-trichlorobenzenesulfonyl chloride
7.	2,5-dichlorobenzenesulfonyl chloride
8.	3,5-dichloro-2-hydroxybenzenesulfonyl chloride
9.	2-mesitylenesulfonyl chloride
10.	4-bromobenzenesulfonyl chloride
11.	4-fluorobenzenesulfonyl chloride
12.	4-chlorobenzenesulfonyl chloride
13.	pipsyl chloride
14.	4-methoxybenzenesulfonyl chloride
15.	4-tert-butylbenzenesulfonyl chloride
16.	p-toluenesulfonyl chloride
17.	isopropylsulfonyl chloride
18.	methanesulfonyl chloride
19.	alpha-toluenesulfonyl chloride
20.	ethanesulfonyl chloride
21.	1-propanesulfonyl chloride
22.	1-butan sulfonyl chloride
23.	Pentamethylbenzenesulfonyl chloride
24.	2,3,5,6-tetramethylbenzenesulfonyl chloride

25.	3-(trifluoromethyl)benzenesulphonyl chloride
26.	3,5-bis(trifluoromethyl)benzenesulfonyl chloride
27.	2,3,4-trichlorobenzenesulfonyl chloride
28.	2,5-dimethoxybenzenesulfonyl chloride
29.	4-methoxy-2,3,6-trimethylbenzenesulfonyl chloride
30.	3,4-dichlorobenzenesulfonyl chloride
31.	4,5-dibromothiophene-2-sulfonyl chloride
32.	3-chloro-4-fluorobenzenesulphonyl chloride
33.	4-ethylbenzenesulfonyl chloride
34.	4-N-propylbenzenesulfonyl chloride
35.	4-N-amylobenzenesulfonyl chloride
36.	4-isopropylbenzenesulphonyl chloride
37.	4-bromo-2,5-difluorobenzenesulphonyl chloride
38.	2-fluorobenzenesulphonyl chloride
39.	3-fluorobenzenesulphonyl chloride
40.	4-(trifluoromethoxy)benzenesulphonyl chloride
41.	4-(trifluoromethyl)benzenesulfonyl chloride
42.	2,4-difluorobenzenesulphonyl chloride
43.	2,4-dichloro-5-methylbenzenesulfonyl chloride
44.	4-chloro-2,5-dimethylbenzenesulphonyl chloride
45.	2-chlorobenzenesulfonyl chloride
46.	4-bromo-2,5-dichlorothiophene-3-sulfonyl chloride
47.	2,5-dichlorothiophene-3-sulphonyl chloride
48.	5-chlorothiophene-2-sulfonyl chloride
49.	2-(trifluoromethyl)benzenesulfonyl chloride
50.	3-chlorobenzenesulfonyl chloride
51.	3,5-dichlorobenzenesulfonyl chloride
52.	m-toluenesulfonyl chloride
53.	2-chloro-6-methylbenzenesulfonyl chloride
54.	5-bromo-2-methoxybenzenesulfonyl chloride
55.	3,4-dimethoxybenzenesulfonyl chloride
56.	2,3-dichlorobenzenesulfonyl chloride
57.	2-bromobenzenesulfonyl chloride
58.	2,3-dichlorothiophene-5-sulphonyl chloride
59.	4-phenylthiophene-2,4-disulfonyl
60.	5-phenylthiophene-2,5-disulfonyl chloride
61.	3-chloro-2-methylbenzenesulfonyl chloride
62.	2-chloro-5-(trifluoromethyl)benzenesulfonyl chloride
63.	2,6-dichlorobenzenesulfonyl chloride
64.	3-bromobenzenesulfonyl chloride
65.	2-(trifluoromethoxy)benzenesulfonyl chloride
66.	4-cyanobenzenesulfonyl chloride
67.	2-cyanobenzenesulfonyl chloride
68.	4-(N-butoxy)benzenesulfonyl chloride
69.	4-acetamido-3-chlorobenzenesulfonyl chloride
70.	3,5-dimethylisoxazole-4-sulfonyl chloride
71.	2,4-dichlorobenzenesulfonyl chloride
72.	2-chloro-4-fluorobenzenesulphonyl chloride

73.	5-fluoro-2-methylbenzenesulphonyl chloride
74.	5-chloro-2-methoxybenzenesulfonyl chloride
75.	2,4,6-trichlorobenzenesulfonyl chloride
76.	4-biphenylsulfonyl chloride
77.	5-bromothiophene-2-sulfonyl chloride
78.	2,6-difluorobenzenesulfonyl chloride
79.	4-n-butylbenzenesulfonyl chloride
80.	4-methylsulfonylbenzenesulfonyl chloride
81.	2-methylsulfonylbenzenesulfonyl chloride
82.	4-acetylbenzenesulfonyl chloride
83.	3-methoxybenzenesulphonyl chloride
84.	2-methoxy-4-methylbenzenesulphonyl chloride

Table V: Compounds of formula (VI)

R₂-NCO (VI)

1.	phenyl isocyanate
2.	2-bromophenyl isocyanate
3.	2-fluorophenyl isocyanate
4.	2,4-difluorophenyl isocyanate
5.	2,6-difluorophenyl isocyanate
6.	2-chlorophenyl isocyanate
7.	2,3-dichlorophenyl isocyanate
8.	2,4-dichlorophenyl isocyanate
9.	2,5-dichlorophenyl isocyanate
10.	2,6-dichlorophenyl isocyanate
11.	2-methoxyphenyl isocyanate
12.	2,4-dimethoxyphenyl isocyanate
13.	2,5-dimethoxyphenyl isocyanate
14.	2-ethoxyphenyl isocyanate
15.	2-(trifluoromethyl)phenyl isocyanate
16.	o-tolyl isocyanate
17.	2,6-dimethylphenyl isocyanate
18.	2-ethylphenyl isocyanate
19.	3-bromophenyl isocyanate
20.	3-fluorophenyl isocyanate
21.	3-chlorophenyl isocyanate
22.	3,4-dichlorophenyl isocyanate
23.	3-methoxyphenyl isocyanate
24.	3-(trifluoromethyl)phenyl isocyanate
25.	m-tolyl isocyanate
26.	4-bromophenyl isocyanate
27.	4-fluorophenyl isocyanate
28.	4-chlorophenyl isocyanate
29.	4-methoxyphenyl isocyanate
30.	4-(trifluoromethyl)phenyl isocyanate
31.	p-tolyl isocyanate
32.	benzoyl isocyanate
33.	tert-butyl isocyanate
34.	(S)-(-)-1-phenylethyl isocyanate

35.	Isopropyl isocyanate
36.	Ethyl isocyanate
37.	Allyl isocyanate
38.	N-propyl isocyanate
39.	n-butyl isocyanate
40.	Cyclohexyl isocyanate
41.	1-naphthyl isocyanate
42.	(R)-(-)-1-(1-naphthyl)ethyl isocyanate
43.	Benzyl isocyanate
44.	3,5-bis(trifluoromethyl)phenyl isocyanate
45.	2,5-difluorophenyl isocyanate
46.	2,4,5-trichlorophenyl isocyanate
47.	2,4,6-trichlorophenyl isocyanate
48.	2-isopropylphenyl isocyanate
49.	2,3-dimethylphenyl isocyanate
50.	4-methoxy-2-methylphenyl isocyanate
51.	2,4-dimethylphenyl isocyanate
52.	2,5-dimethylphenyl isocyanate
53.	2-ethyl-6-methylphenyl isocyanate
54.	3-cyanophenyl isocyanate
55.	5-chloro-2,4-dimethoxyphenyl isocyanate
56.	3-chloro-4-methylphenyl isocyanate
57.	3,5-dichlorophenyl isocyanate
58.	5-chloro-2-methoxyphenyl isocyanate
59.	3,4,5-trimethoxyphenyl isocyanate
60.	3,5-dimethoxyphenyl isocyanate
61.	3-(methylthio)phenyl isocyanate
62.	3-acetylphenyl isocyanate
63.	3,4-dimethylphenyl isocyanate
64.	3,5-dimethylphenyl isocyanate
65.	2-methoxy-5-methylphenyl isocyanate
66.	3-ethylphenyl isocyanate
67.	4-bromo-2-(trifluoromethyl)phenyl isocyanate
68.	4-chloro-2-(trifluoromethyl)phenyl isocyanate
69.	4-chloro-3-(trifluoromethyl)phenyl isocyanate
70.	4-iodophenyl isocyanate
71.	4-phenoxyphenyl isocyanate
72.	4-ethoxyphenyl isocyanate
73.	4-acetylphenyl isocyanate
74.	4-isopropylphenyl isocyanate
75.	4-ethylphenyl isocyanate
76.	4-n-butylphenyl isocyanate
77.	2,4,6-trimethylphenyl isocyanate
78.	2-isopropyl-6-methylphenyl isocyanate
79.	2,6-diethylphenyl isocyanate
80.	5-chloro-2-methylphenyl isocyanate
81.	4-chloro-2-methylphenyl isocyanate
82.	4-(trifluoromethoxy)phenyl isocyanate
83.	2-chloro-5-(trifluoromethyl)phenyl isocyanate
84.	2-chloro-6-methylphenyl isocyanate
85.	2,4,5-trimethylphenyl isocyanate

86.	3-chloro-2-methoxyphenyl isocyanate
87.	3-chloro-2-methylphenyl isocyanate
88.	3-chloro-4-fluorophenyl isocyanate
89.	4-bromo-2-methylphenyl isocyanate
90.	4-bromo-2,6-dimethylphenyl isocyanate
91.	2,6-dibromo-4-fluorophenyl isocyanate
92.	4-butoxyphenyl isocyanate
93.	3-fluoro-4-methylphenyl isocyanate
94.	5-fluoro-2-methylphenyl isocyanate
95.	2-biphenyllyl isocyanate
96.	4-biphenyllyl isocyanate
97.	2-bromo-4,6-difluorophenyl isocyanate
98.	(r)-(+) -1-phenylethyl isocyanate
99.	1-(1-naphthyl)ethyl isocyanate
100.	(s)-(+) -1-(1-naphthyl) ethyl isocyanate
101.	3,4-difluorophenyl isocyanate
102.	3-isopropenyl- α,α -dimethylbenzyl isocyanate
103.	2-(trifluoromethoxy)phenyl isocyanate
104.	4-benzyloxyphenyl isocyanate
105.	4-bromo-2-chlorophenyl isocyanate
106.	4-bromo-2-fluorophenyl isocyanate
107.	2-fluoro-5-methylphenyl isocyanate
108.	2,3,4-trifluorophenyl isocyanate
109.	2-(difluoromethoxy)phenyl isocyanate
110.	4-(difluoromethoxy)phenyl isocyanate
111.	2-methylbenzyl isocyanate
112.	2-chlorobenzyl isocyanate
113.	4-fluorobenzyl isocyanate
114.	4-methoxybenzyl isocyanate
115.	2,6-difluorobenzoyl isocyanate
116.	4-fluorobenzoyl isocyanate
117.	2-fluoro-3-(trifluoromethyl)phenyl isocyanate
118.	2-fluoro-5-(trifluoromethyl)phenyl isocyanate
119.	2-fluoro-6-(trifluoromethyl)phenyl isocyanate
120.	4-fluoro-2-(trifluoromethyl)phenyl isocyanate
121.	2-(tert-butyl)phenyl isocyanate
122.	3-pyridyl isocyanate

From all of the above, it is clear to the skilled man that once a library of aminoisoxazoles derivatives is thus prepared, for instance consisting of a few thousands of compounds of formula (I), the said library can be very advantageously used for screening towards given target kinases, as formerly reported.

See, for a general reference to libraries of compounds and uses thereof as tools for screening biological activities,

J. Med. Chem. 1999, 42, 2373-2382; and Bioorg. Med. Chem. Lett. 10 (2000), 223-226.

Pharmacology

5 The compounds of formula (I) are active as protein kinase inhibitors and are therefore useful, for instance, to restrict the unregulated proliferation of tumor cells.

In therapy, they may be used in the treatment of various tumors such as, for instance, carcinomas, e.g. mammary
10 carcinoma, lung carcinoma, bladder carcinoma, colon carcinoma, ovary and endometrial tumors, sarcomas, e.g. soft tissue and bone sarcomas, and the hematological malignancies such as, e.g., leukemias.

In addition, the compounds of formula (I) are also useful
15 in the treatment of other cell proliferative disorders such as psoriasis, vascular smooth cell proliferation associated with atherosclerosis and post-surgical stenosis and restenosis and in the treatment of Alzheimer's disease.

The inhibiting activity of putative protein kinase
20 inhibitors and the potency of selected compounds was determined through a method of assay based on the use of the MultiScreen-PH 96 well plate (Millipore), in which a phosphocellulose filter paper was placed at each well bottom allowing binding of positive charged substrate after
25 a washing/filtration step.

When a radioactivity labeled phosphate moiety was transferred by the ser/threo kinase to the filter-bound histone, light emitted was measured in a scintillation counter.

30 Or through a method of assay based on the use of the SPA technology (Amersham Pharmacia Biotech).

The assay consists of the transfer of radioactivity labelled phosphate moiety by the kinase to a biotinylated substrate. The resulting ³³P-labelled biotinylated product
35 is allowed to bind to streptavidin-coated SPA beads (biotin

capacity 130pmol/mg), and light emitted was measured in a scintillation counter.

Inhibition assay of cdk2/Cyclin A activity

5 **Kinase reaction:** 4 μ M in house biotinylated histone H1 (Sigma # H-5505) substrate, 10 μ M ATP (0.1 microCi $P^{33}\gamma$ -ATP), 4.2 ng Cyclin A/CDK2 complex, inhibitor in a final volume of 30 μ l buffer (TRIS HCl 10 mM pH 7.5, $MgCl_2$ 10 mM, DTT 7.5 mM + 0.2 mg/ml BSA) were added to each well of a 96
10 U bottom. After 30 min at r.t. incubation, reaction was stopped by 100 μ l PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, containing 1 mg SPA beads. Then a volume of 110 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 μ l 5M
15 CsCl were added to allow stratification of beads to the top of the plate and let stand 4 hours before radioactivity counting in the Top-Count instrument

IC50 determination: inhibitors were tested at different concentrations ranging from 0.0015 to 10 μ M. Experimental
20 data were analyzed by the computer program GraphPad Prism using the four parameter logistic equation:

$$y = \text{bottom} + (\text{top} - \text{bottom}) / (1 + 10^{((\log IC_{50} - x) * \text{slope}))})$$

where x is the logarithm of the inhibitor concentration, y is the response; y starts at bottom and goes to top with a
25 sigmoid shape.

Ki calculation:

Experimental method: Reaction was carried out in buffer (10 mM Tris, pH 7.5, 10 mM $MgCl_2$, 0.2 mg/ml BSA, 7.5 mM DTT) containing 3.7 nM enzyme, histone and ATP (constant ratio
30 of cold/labeled ATP 1/3000). Reaction was stopped with EDTA and the substrate captured on phosphomembrane (Multiscreen 96 well plates from Millipore). After extensive washing, the multiscreen plates are read on a top counter. Control (time zero) for each ATP and histone concentrations was
35 measured.

- 40 -

Experimental design: Reaction velocities are measured at different four ATP, substrate (histone) and inhibitor concentrations. An 80-point concentration matrix was designed around the respective ATP and substrate Km values, and the inhibitor IC50 values (0.3, 1, 3, 9 fold the Km or IC50 values). A preliminary time course experiment in the absence of inhibitor and at the different ATP and substrate concentrations allow the selection of a single endpoint time (10 min) in the linear range of the reaction for the Ki determination experiment.

Kinetic parameter estimates: Kinetic parameters were estimated by simultaneous nonlinear least-square regression using [Eq.1] (competitive inhibitor respect to ATP, random mechanism) using the complete data set (80 points):

15

$$v = \frac{V_m \cdot A \cdot B}{\alpha \cdot K_a \cdot K_b + \alpha \cdot K_a \cdot B + a \cdot K_b \cdot A + A \cdot B + \alpha \cdot \frac{K_a}{K_i} \cdot I \cdot (K_b + \frac{B}{\beta})} \quad [\text{Eq.1}]$$

where A=[ATP], B=[Substrate], I=[inhibitor], Vm= maximum velocity, Ka, Kb, Ki the dissociation constants of ATP, substrate and inhibitor respectively. α and β the cooperativity factor between substrate and ATP binding and substrate and inhibitor binding respectively.

In addition the selected compounds have been characterized on a panel of ser/threo kinases strictly related to cell cycle (cdk2/cyclin E, cdk1/cyclin B1, cdk5/p25, cdk4/cyclin D1), and also for specificity on MAPK, PKA, EGFR, IGF1-R, and Aurora-2.

Inhibition assay of cdk2/Cyclin E activity

Kinase reaction: 10 μ M in house biotinylated histone H1 (Sigma # H-5505) substrate, 30 μ M ATP (0.3 microCi P³³ γ -ATP), 4 ng GST-Cyclin E/CDK2 complex, inhibitor in a final volume of 30 ml buffer (TRIS HCl 10 mM pH 7.5, MgCl₂ 10 mM,

- 41 -

DTT 7.5 mM + 0.2 mg/ml BSA) were added to each well of a 96 U bottom. After 60 min at r.t. incubation, reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, containing 1 mg SPA beads. Then a volume of 110 5 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the plate and let stand 4 hours before radioactivity counting in the Top-Count instrument

10 **IC50 determination:** see above

Inhibition assay of cdk1/Cyclin B1 activity

Kinase reaction: 4 μ M in house biotinylated histone H1 (Sigma # H-5505) substrate, 20 mM ATP (0.2 microCi $P^{33}\gamma$ - 15 ATP), 3 ng Cyclin B/CDK1 complex, inhibitor in a final volume of 30 ml buffer (TRIS HCl 10 mM pH 7.5, $MgCl_2$ 10 mM, DTT 7.5 mM + 0.2 mg/ml BSA) were added to each well of a 96 U bottom. After 20 min at r.t. incubation, reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 20 500 mM ATP, containing 1 mg SPA beads. Then a volume of 110 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the Optiplate and let stand 4 hours before radioactivity 25 counting in the Top-Count instrument.

IC50 determination: see above

Inhibition assay of cdk5/p25 activity

The inhibition assay of cdk5/p25 activity was performed 30 according to the following protocol.

Kinase reaction: 10 μ M biotinylated histone H1 (Sigma # H-5505) substrate, 30 μ M ATP (0.3 microCi $P^{33}\gamma$ -ATP), 15 ng CDK5/p25 complex, inhibitor in a final volume of 30 ml buffer (TRIS HCl 10 mM pH 7.5, $MgCl_2$ 10 mM, DTT 7.5 mM + 0.2 35 mg/ml BSA) were added to each well of a 96 U bottom. After

- 42 -

30 min at r.t. incubation, reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, containing 1 mg SPA beads. Then a volume of 110 ml is transferred to Optiplat.

- 5 After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the plate and let stand 4 hours before radioactivity counting in the Top-Count instrument.

IC50 determination: see above

10

Inhibition assay of cdk4/Cyclin D1 activity

- Kinase reaction:** 0,4 uM mouse GST-Rb (769-921) (# sc-4112 from Santa Cruz) substrate, 10 µM ATP (0.5 µCi $P^{33}\gamma$ -ATP), 100 ng of baculovirus expressed GST-cdk4/GST-Cyclin D1, suitable concentrations of inhibitor in a final volume of 50 ml buffer (TRIS HCl 10 mM pH 7.5, MgCl₂ 10 mM, 7.5 mM DTT+ 0.2mg/ml BSA) were added to each well of a 96 U bottom well plate. After 40 min at 37 °C incubation, reaction was stopped by 20 ml EDTA 120 mM.

- 20 **Capture:** 60 ml were transferred from each well to MultiScreen plate, to allow substrate binding to phosphocellulose filter. Plates were then washed 3 times with 150 ml/well PBS Ca⁺⁺/Mg⁺⁺ free and filtered by MultiScreen filtration system.

- 25 **Detection:** filters were allowed to dry at 37°C, then 100 ml/well scintillant were added and ^{33}P labeled Rb fragment was detected by radioactivity counting in the Top-Count instrument.

IC50 determination: see above

30

Inhibition assay of MAPK activity

- Kinase reaction:** 10 µM in house biotinylated MBP (Sigma # M-1891) substrate, 15 µM ATP (0.15 microCi $P^{33}\gamma$ -ATP), 30 ng GST-MAPK (Upstate Biotechnology # 14-173), inhibitor in a final volume of 30 ml buffer (TRIS HCl 10 mM pH 7.5, MgCl₂

35

- 43 -

10 mM, DTT 7.5 mM + 0.2 mg/ml BSA) were added to each well of a 96 U bottom. After 30 min at r.t. incubation, reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, containing 1 mg SPA beads. Then a volume of 5 110 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the Optiplate and let stand 4 hours before radioactivity counting in the Top-Count instrument.

10 IC50 determination: see above

Inhibition assay of PKA activity

Kinase reaction: 10 μ M in house biotinylated histone H1 (Sigma # H-5505) substrate, 10 μ M ATP (0.2 microM $P^{33}\gamma$ -ATP), 15 0.45 U PKA (Sigma # 2645), inhibitor in a final volume of 30 ml buffer (TRIS HCl 10 mM pH 7.5, $MgCl_2$ 10 mM, DTT 7.5 mM + 0.2 mg/ml BSA) were added to each well of a 96 U bottom. After 90 min at r.t. incubation, reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, 20 containing 1 mg SPA beads. Then a volume of 110 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the Optiplate and let stand 4 hours before radioactivity 25 counting in the Top-Count instrument.

IC50 determination: see above

Inhibition assay of EGFR activity

Kinase reaction: 10 μ M in house biotinylated MBP (Sigma # 30 M-1891) substrate, 2 mM ATP (0.04 microCi $P^{33}\gamma$ -ATP), 36 ng insect cell expressed GST-EGFR, inhibitor in a final volume of 30 ml buffer (Hepes 50 mM pH 7.5, $MgCl_2$ 3 mM, $MnCl_2$ 3 mM, DTT 1 mM, $NaVO_3$ 3 μ M + 0.2 mg/ml BSA) were added to each well of a 96 U bottom. After 20 min at r.t. incubation,

- 44 -

reaction was stopped by 100 ml PBS + 32 mM EDTA + 0.1% Triton X-100 + 500 mM ATP, containing 1 mg SPA beads. Then a volume of 110 ml is transferred to Optiplate.

After 20 min. incubation for substrate capture, 100 ml 5M CsCl were added to allow stratification of beads to the top of the Optiplate and let stand 4 hours before radioactivity counting in the Top-Count instrument.

IC50 determination: see above

10 **Inhibition assay of IGF1-R activity**

The inhibition assay of IGF1-R activity was performed according to the following protocol.

Kinase reaction: 10 uM biotinylated MBP (Sigma cat. # M-1891) substrate, 0-20 mM inhibitor, 6 mM ATP, 1 microCi ³³P-ATP, and 22.5 ng GST-IGF1-R (pre-incubated for 30 min at room temperature with cold 60 mM cold ATP) in a final volume of 30 ml buffer (50 mM HEPES pH 7.9, 3 mM MnCl₂, 1 mM DTT, 3 mM NaVO₃) were added to each well of a 96 U bottom well plate. After incubation for 35 min at room temperature, the reaction was stopped by addition of 100 ml PBS buffer containing 32 mM EDTA, 500 mM cold ATP, 0.1% Triton X100 and 10mg/ml streptavidin coated SPA beads. After 20 min incubation, 110 mL of suspension were withdrawn and transferred into 96-well OPTIPLATES containing 100 ml of 5M CsCl. After 4 hours, the plates were read for 2 min in a Packard TOP-Count radioactivity reader.

Inhibition assay of Aurora-2 activity

30 **Kinase reaction:** 8 μM biotinylated peptide (4 repeats of LRRWSLG), 10 μM ATP (0.5 uCi P³³γ-ATP), 15 ng Aurora2, inhibitor in a final volume of 30 ml buffer (HEPES 50 mM pH 7.0, MgCl₂ 10 mM, 1 mM DTT, 0.2 mg/ml BSA, 3mM orthovanadate) were added to each well of a 96 U bottom well plate. After 30 minutes at room temperature

- 45 -

incubation, reaction was stopped and biotinylated peptide captured by adding 100 ml of bead suspension.

Stratification: 100 ml of CsCl₂ 5 M were added to each well and let stand 4 hour before radioactivity was counted in the Top-Count instrument.

IC₅₀ determination: see above

Inhibition assay of Cdc7/dbf4 activity

The inhibition assay of Cdc7/dbf4 activity was performed according to the following protocol.

The Biotin-MCM2 substrate is trans-phosphorylated by the Cdc7/Dbf4 complex in the presence of ATP traced with γ^{33} -ATP. The phosphorylated Biotin-MCM2 substrate is then captured by Streptavidin-coated SPA beads and the extent of phosphorylation evaluated by β counting.

The inhibition assay of Cdc7/dbf4 activity was performed in 96 wells plate according to the following protocol.

To each well of the plate were added :

- 10 ul substrate (biotinylated MCM2, 6 uM final concentration)
- 10 ul enzyme (Cdc7/Dbf4, 12.5 nM final concentration)
- 10 ul test compound (12 increasing concentrations in the nM to mM range to generate a dose-response curve)
- 10 ul of a mixture of cold ATP (10uM final concentration) and radioactive ATP (1/2500 molar ratio with cold ATP) was then used to start the reaction which was allowed to take place at 37°C.

Substrate, enzyme and ATP were diluted in 50 mM HEPES pH 7.9 containing 15 mM MgCl₂, 2 mM DTT, 3 mM NaVO₃, 2mM glycerophosphate and 0.2mg/ml BSA. The solvent for test compounds also contained 10% DMSO.

After incubation for 20 minutes, the reaction was stopped by adding to each well 100 ul of PBS pH 7.4 containing 50 mM EDTA, 1 mM cold ATP, 0.1% Triton X100 and 10 mg/ml streptavidin coated SPA beads.

- 46 -

After 15 minutes of incubation at room temperature to allow the biotinylated MCM2-streptavidin SPA beads interaction to occur, beads were trapped in a 96 wells filter plate (Unifilter^R GF/BTM) using a Packard Cell Harvester (Filtermate), washed with distilled water and then counted using a Top Count (Packard).

Counts were blank-subtracted and then the experimental data (each point in triplicate) were analyzed for IC50 determination using a non-linear regression analysis (Sigma Plot).

The compounds of formula (I) of the present invention, suitable for administration to a mammal, e.g. to humans, can be administered by the usual routes and the dosage level depends upon the age, weight, conditions of the patient and the administration route.

For example, a suitable dosage adopted for oral administration of a compound of formula (I) may range from about 10 to about 500 mg pro dose, from 1 to 5 times daily.

The compounds of the invention can be administered in a variety of dosage forms, e.g. orally, in the form of tablets, capsules, sugar or film coated tablets, liquid solutions or suspensions; rectally in the form of suppositories; parenterally, e.g. intramuscularly, or by intravenous and/or intrathecal and/or intraspinal injection or infusion.

In addition, the compounds of the invention can be administered either as single agents or, alternatively, in combination with known anticancer treatments such as radiation therapy or chemotherapy regimen in combination with cytostatic or cytotoxic agents, antibiotic-type agents, alkylating agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents, cyclooxygenase inhibitors (e.g. COX-2 inhibitors), metallomatrixprotease inhibitors, telomerase inhibitors,

tyrosine kinase inhibitors, anti-growth factor receptor agents, anti-HER agents, anti-EGFR agents, anti-angiogenesis agents, farnesyl transferase inhibitors, ras-raf signal transduction pathway inhibitors, cell cycle
5 inhibitors, other cdks inhibitors, tubulin binding agents, topoisomerase I inhibitors, topoisomerase II inhibitors, and the like.

As an example, the compounds of the invention can be administered in combination with one or more
10 chemotherapeutic agents such as, for instance, exemestane, formestane, anastrozole, letrozole, fadrozole, taxane and derivatives such as paclitaxel and docetaxel, encapsulated taxanes, CPT-11, camptothecin derivatives, anthracycline glycosides, e.g., doxorubicin, idarubicin, epirubicin,
15 etoposide, navelbine, vinblastine, carboplatin, cisplatin, estramustine, celecoxib, parecoxib, rofecoxib, valecoxib, tamoxifen, raloxifen, JTE 5222, Sugem SU-5416, Sugem SU-6668, Herceptin, estramustine, and the like, optionally within liposomal formulations thereof.

20 If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described above and the other pharmaceutically active agent within the approved dosage range.

Compounds of formula (I) may be used sequentially with
25 known anticancer agents when a combination formulation is inappropriate.

The present invention also includes pharmaceutical compositions comprising a compound of formula (I) or a
30 pharmaceutically acceptable salt thereof in association with a pharmaceutically acceptable excipient (which can be a carrier or a diluent).

The pharmaceutical compositions containing the compounds of the invention are usually prepared following conventional
35 methods and are administered in a pharmaceutically suitable form.

For example, the solid oral forms may contain, together with the active compound, diluents, e.g. lactose, dextrose, saccharose, sucrose, cellulose, corn starch or potato starch; lubricants, e.g. silica, talc, stearic , magnesium
5 or calcium stearate, and/or polyethylene glycols; binding agents, e.g. starches, arabic gum, gelatine, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disaggregating agents, e.g. a starch, alginic, alginates or sodium starch glycolate; effervescing
10 mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be manufactured in known manner, for
15 example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The liquid dispersions for oral administration may be e.g. syrups, emulsions and suspensions.

The syrups may contain as carrier, for example, saccharose
20 or saccharose with glycerin and/or mannitol and/or sorbitol.

The suspensions and the emulsions may contain as carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl
25 alcohol.

The suspension or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol,
30 and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusions may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions or they may contain as a
35 carrier propylene glycol.

The suppositories may contain together with the active compound a pharmaceutically acceptable carrier, e.g. cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty ester surfactant or lecithin.

5

The following examples are herewith intended to better illustrate the present invention, without posing any limitation to it.

10 General Methods

Flash chromatography was performed on silica gel (Merck grade 9385, 60Å). HPLC/MS was performed on a Waters X Terra RP 18 (4.6 x 50 mm, 3.5 mm) column using a Waters 2790 HPLC system equipped with a 996 Waters PDA detector and a
15 Micromass mod. ZQ single quadrupole mass spectrometer, equipped with an electrospray (ESI) ion source. Mobile phase A was ammonium acetate 5 mM buffer (pH 5.5 with acetic acid / acetonitrile 95:5), and Mobile phase B was H₂O / acetonitrile (5:95). Gradient from 10 to 90% B in 8
20 minutes, hold 90% B 2 min. UV detection at 220 nm and 254 nm. Flow rate 1 ml/min. Injection volume 10 µl. Full scan, mass range from 100 to 800 amu. Capillary voltage was 2.5 KV; Source temp. was 120°C; Cone was 10 V. Retention Times (HPLC r.t.) are given in minutes at 220 nm or 254 nm. Mass
25 is given as m/z ratio.

HPLC/MS was also (Method B) performed on a Hypersil C18 BDS (50 x 2.0mm, 5mm) column using a HPLC system equipped with Hewlett Packard 1312A binary pump with Gilson 215 autosampler fitted with a 1ml syringe with a Polymer Labs
30 PL1000 Evaporative Light Scattering Detector (ELSD) and a Micromass ZMD mass spectrometer operating in Electrospray positive ionisation mode. Mobile phase A was water with 0.1% of trifluoroacetic acid, and mobile phase B was acetonitrile with 0.1% of trifluoroacetic acid. Gradient
35 from 0% to 95% B in 1.8 minutes, hold 95% B 0.3 min. Flow rate 1 ml/min. Injection volume 3 µl. ELS Detector:

- 50 -

Nebuliser Temperature 80°C; Evaporation temperature 90°C; Gas Flow 1.5 l/hr. Full scan, mass range from 150 to 800 amu (@ 0.5 secs/scan, 0.1 second interscan delay). Capillary voltage was 25 V; Source temp. was 140°C.

5 Retention Times (HPLC Method B r.t.) are given in minutes (rev. ELSD); mass is given as mass found in MassLynx® Report (Waters).

When necessary compounds have been purified by Preparative HPLC on a Waters Symmetry C18 (19 x 50 mm, 5µm) column
10 using a Waters preparative HPLC 600 equipped with a 996 Waters PDA detector and a Micromass mod. ZMD single quadrupole mass spectrometer, electrospray ionisation, positive mode. Mobile phase A was water 0.01% TFA, and Mobile phase B was acetonitrile. Gradient from 10 to 90%B
15 in 8 min, hold 90%B 2 min. Flow rate 20 ml/m.

¹H-NMR spectroscopy was performed on a Mercury VX 400 operating at 400.45 MHz equipped with a 5mm double resonance probe (1H {15N-31P} ID_PFG Varian).

20

Example 1

N-(4-tert-butylphenyl)cyanoacetamide

To 100 mg of cyanoacetic acid (1.17 mmol, 1.5 eq) in 8 ml of dichloromethane (DCM), 0.130 ml of 4-tert-butyl aniline
25 (0.78 mmol) and 804 mg of N-cyclohexylcarbodiimide, N'-methylpolystyrene (1.94 mmol/g theoretical loading, 2 eq) were added. The suspension was shaken over-night then the resin was filtered off and washed with dichloromethane and dimethylformamide. The final product was obtained in 80%
30 yield (135 mg) with 100% HPLC purity (according to HPLC analysis at 220nm). [M+H]⁺ = 217; [M-H]⁻ = 215; HPLC r.t. 5,35.

¹H-NMR (DMSO-d₆), diagnostic signals (ppm): 10,2 (bs, 1H), 7,42 (d, 2H), 7,2 (d, 2H), 3,8 (s, 2H), 1,2 (s, 9H).

35

By working in an analogous way and by using the appropriate amine derivative, the following compounds were prepared:

- N-(4-fluorophenyl)cyanoacetamide [M-H]⁻ = 177; HPLC r.t. 3,2.
- 5 N-(2,4-difluorophenyl)cyanoacetamide [M-H]⁻ = 195; HPLC r.t. 3,1.
- N-(4-chlorophenyl)cyanoacetamide [M-H]⁻ = 193; HPLC r.t. 4,2.
- N-(Benzyl)cyanoacetamide [M+H]⁺ = 175 [M-H]⁻ = 173; HPLC
- 10 r.t. 2,6.
- 2-cyano-N-(5,6,7,8-tetrahydronaphthalen-1-yl)acetamide [M+H]⁺ = 215; [M-H]⁻ = 213; HPLC r.t. 4,5.
- 2-cyano-N-(2,3-dihydro-1H-inden-5-yl)acetamide [M-H]⁻ = 199; HPLC r.t. 4,5.
- 15 N-(4-(trifluoromethoxy)phenyl)cyanoacetamide [M-H]⁻ = 243; HPLC r.t. 4,9.
- N-(1-naphthyl)cyanoacetamide [M+H]⁺ = 211 [M-H]⁻ = 209; HPLC r.t. 4,0.
- N-(2-benzylphenyl)cyanoacetamide [M-H]⁻ = 249; HPLC r.t.
- 20 4,9.
- 2-cyano-N-(4-methylphenyl)acetamide [M+H]⁺ = 175; [M-H]⁻ = 173; HPLC r.t. 3,63.
- 2-cyano-N-[4-(dimethylamino)phenyl]acetamide [M+H]⁺ = 204; HPLC r.t. 3,06.
- 25 N-(3-benzoyloxyphenyl)cyanoacetamide [M+H]⁺ = 267; [M-H]⁻ = 265; HPLC r.t. 5,4.
- N-benzhydryl-2-cyanoacetamide [M-H]⁻ = 249; HPLC r.t. 4,9.
- N-(2,4-dimethoxybenzyl)cyanoacetamide [M-H]⁻ = 233; HPLC r.t. 3,28.
- 30 N-(3,4,5-trimethoxyphenyl)cyanoacetamide [M+H]⁺ = 251; [M-H]⁻ = 249; HPLC r.t. 2,79.
- 2-cyano-N-(2-methoxydibenzo[b,d]furan-3-yl)acetamide [M+H]⁺ = 281 [M-H]⁻ = 279 HPLC r.t. 5,7.
- 2-cyano-N-quinolin-8-ylacetamide [M+H]⁺ = 212 [M-H]⁻ = 210
- 35 HPLC r.t. 3,97.

- 2-cyano-N-[2-oxo-4-(trifluoromethyl)-2H-chromen-7-yl]acetamide [M-H]⁻ =295 HPLC r.t. 4,7.
- N-(3,5-dimethoxyphenyl)cyanoacetamide [M+H]⁺ = 221 [M-H]⁻ =219 HPLC r.t. 3,58.
- 5 Ethyl 3-[(cyanoacetyl)amino]benzoate [M-H]⁻ =231 HPLC r.t. 4,1.
- 2-cyano-N-[2-methoxy-5-(trifluoromethyl)phenyl]acetamide [M-H]⁻ =257 HPLC r.t. 4,2.
- 2-cyano-N-[(5-methyl-2-furyl)methyl]acetamide.
- 10 2-cyano-N-cyclobutylacetamide.
- 2-cyano-N-cyclohexylacetamide.
- 2-cyano-N-(2-methylcyclohexyl)acetamide.
- 2-cyano-N-(cyclohexylmethyl)acetamide.
- 2-cyano-N-(1,2,3,4-tetrahydronaphthalen-1-yl)acetamide.
- 15 2-cyano-N-(1-naphthylmethyl)acetamide.
- 2-cyano-N-cycloheptylacetamide.
- 2-cyano-N-(thien-2-ylmethyl)acetamide.
- N-(1,3-benzodioxol-5-ylmethyl)-2-cyanoacetamide.
- N-(1-benzylpiperidin-4-yl)-2-cyanoacetamide.
- 20 N-(tert-butyl)-2-cyanoacetamide.
- 2-cyano-N-(1,2-diphenylethyl)acetamide.
- 2-cyano-N-(1-phenylethyl)acetamide.
- 2-cyano-N-(1,2-dimethylpropyl)acetamide.
- 2-cyano-N-isopropylacetamide.
- 25 2-cyano-N-(2-methoxy-1-methylethyl)acetamide.
- 2-cyano-N-(1,3-dimethylbutyl)acetamide.
- 2-cyano-N-(1-methyl-3-phenylpropyl)acetamide.
- 2-cyano-N-(1,5-dimethylhexyl)acetamide.
- N-(sec-butyl)-2-cyanoacetamide.
- 30 2-cyano-N-(1-ethylpropyl)acetamide.
- 2-cyano-N-(1-methylhexyl)acetamide.
- 2-cyano-N-(1-methylheptyl)acetamide.
- N-benzyl-2-cyanoacetamide.
- 2-cyano-N-(2-fluorobenzyl)acetamide.
- 35 N-(2-chlorobenzyl)-2-cyanoacetamide.
- 2-cyano-N-(2-methylbenzyl)acetamide.

- 2-cyano-N-(3-fluorobenzyl)acetamide.
2-cyano-N-(3,4-dimethoxybenzyl)acetamide.
2-cyano-N-[3-(trifluoromethyl)benzyl]acetamide.
2-cyano-N-(3-methylbenzyl)acetamide.
5 2-cyano-N-(4-methoxybenzyl)acetamide.
2-cyano-N-(4-methylbenzyl)acetamide.
2-cyano-N-isobutylacetamide.
2-cyano-N-(2-ethylhexyl)acetamide.
2-cyano-N-(2-phenylethyl)acetamide.
10 2-cyano-N-prop-2-ynylacetamide.
N-allyl-2-cyanoacetamide.
2-cyano-N-(3,3-diphenylpropyl)acetamide.
2-cyano-N-isopentylacetamide.
2-cyano-N-propylacetamide.
15 2-cyano-N-[3-(dibutylamino)propyl]acetamide.
2-cyano-N-(3-phenylpropyl)acetamide.
2-cyano-N-(4-phenylbutyl)acetamide.
2-cyano-N-pentylacetamide.
2-cyano-N-hexylacetamide.
20 2-cyano-N-heptylacetamide.
2-cyano-N-octylacetamide.
N-[3,5-bis(trifluoromethyl)benzyl]-2-cyanoacetamide.
2-cyano-N-(3,4-difluorobenzyl)acetamide.
2-cyano-N-[4-(trifluoromethyl)benzyl]acetamide.
25 2-cyano-N-[2-(trifluoromethyl)benzyl]acetamide.
N-butyl-2-cyanoacetamide.
2-cyano-N-(2-cyclohex-1-en-1-ylethyl)acetamide.
N-(4-tert-butylcyclohexyl)-2-cyanoacetamide.
N-(2-bromobenzyl)-2-cyanoacetamide.
30 2-cyano-N-(cyclopropylmethyl)acetamide.
N-(3-chlorobenzyl)-2-cyanoacetamide.
2-cyano-N-[3-(methylthio)propyl]acetamide.
N-[2-(3-chlorophenyl)ethyl]-2-cyanoacetamide.
2-cyano-N-(2,4-dimethoxybenzyl)acetamide.
35 2-cyano-N-[4-(trifluoromethoxy)benzyl]acetamide.

- 54 -

Example 2**Cyanoacetic acid Rink amide**

Rink amide (100 mg, Novabiochem, 0.59 mmol/g, 0.059 mmol) was suspended in 2 ml of piperidine 20% in dimethylformamide (DMF), after 20 minutes of shaking, the resin was washed with DMF, DCM and DMF and the procedure was repeated.

To the Fmoc-protected resin 2 ml of DCM/DMF 1:1, cyanoacetic acid (20 mg, 0.236 mmol, 4 eq.), HOBt (64 mg, 0.47 mmol, 8 eq.), DIPEA (80.8 μ l, 0.47 mmol, 8 eq.) were added. The suspension was shaken 10 min and DIC (74 μ l, 0.47 mmol, 8 eq.) was added. After 16 hrs the resin was washed with DCM (x2), DMF (x2), DCM (x2), DMF (x2) and DCM (x5); a second cycle was repeated. The unreacted amino groups were capped with acetic anhydride (3 eq). The resin was washed with DCM (x2), DMF (x2), DCM (x2), DMF (x2) and DCM (x5).

Example 3**5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide**

Cyanacetamide (1.5 g, 18 mmol) and 3-thiophenylhydroxyaminomethyl chloride (3.46 g, 21 mmol) were suspended in dry THF (80 ml) and cooled to -20°C. Then lithium bis(trimethylsilyl)amide LiHMDS (41.02 ml of 1 M solution in THF, 41.02 mmol) was added, the resulting suspension was stirred at -20°C for 15 min, allowed to warm to room temperature (23°C) and stirred for 2h. The reaction was quenched with water (50 ml) and stirred for 10', the final mixture was extracted several times with ethyl acetate AcOEt and the organic phases were collected, dried over anhydrous Na₂SO₄ and solvents were removed under reduced pressure. The product was purified by crystallization from isopropanol and water or by flash chromatography over silica gel. [M+H]⁺ = 210. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm): 7.9 (d, 1H), 7.7 (dd, 1H),

7.6 (bs, 2H), 7.3 (d, 1H).

By working in an analogous way and by using the appropriate chloroxime derivative and the appropriate cyanacetic acid derivative such as ester, primary amide or secondary amide, the following compounds were prepared:

- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide
[M+H]⁺ = 210. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
7.85 (d, 1H), 7.8 (bs, 3H), 7.2 (dd, 1H).
- 10 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide.
[M+H]⁺ = 205. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
9,8 (bs, 1H), 8,7 (m, 1H), 8,05 (m, 2H), 7,95 (bs, 2H), 7,6
(m, 1H), 7,2 (bs, 1H).
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide.
15 [M+H]⁺ = 205. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
8,7 (m, 2H), 7,95 (m, 1H), 7,6 (bs, 2H), 7,5 (m, 1H).
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide.
[M+H]⁺ = 205. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
8,7 (d, 2H), 7,7 (bs, 2H), 7,5 (s, 2H).
- 20 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide.
[M+H]⁺ = 211. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
9,5 (bs, 1H), 8,1 (s, 1H), 8,0 (bs, 2H), 7,9 (d, 1H).
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid amide. [M+H]⁺ = 298. ¹H-NMR
25 (DMSO-d₆), diagnostic signals (ppm): 7,65 (bs, 2H), 7,4-7,5
(m, 5H), 2,2 (s, 3H), 2,1 (s, 3H).
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide.
[M+H]⁺ = 255. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):
10,4 (bs, 1H), 8,6 (d, 1H), 8,15 (d, 1H), 8,1 (dd, 1H) 8,15
30 (bs, 2H), 8,0 (dd, 1H), 7,9 (m, 1H), 7,75 (m, 1H).
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
butylamide. [M+H]⁺ = 279. ¹H-NMR (DMSO-d₆), diagnostic
signals (ppm): 7,8 (d, 1H), 7,6 (d, 1H), 7,4 (bs, 2H), 7,2
(dd, 1H) 3,2 (m, 2H), 1,4 (m, 2H), 1,2 (m, 2H), 0,9 (m,
35 3H).

Methyl 5-amino-3-(2-pyridinyl)-4-isoxazolecarboxylate.

[M+H]⁺ = 220. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm):

8,6 (d, 1H), 7,9 (bs, 2H), 7,85 (dd, 1H), 7,55 (d, 1H) 7,45 (dd, 1H), 3,5 (s, 3H).

- 5 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide. HPLC r.t. 4.1, [M+H]⁺ =415.
5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide. HPLC r.t. 4.58, [M+H]⁺ =315.1.
5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
10 benzylamide. HPLC r.t. 3.55, [M+H]⁺ =295.
5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 4.96, [M+H]⁺ =335.
5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
15 fluoro-phenyl)-amide. HPLC r.t. 3.86, [M+H]⁺ =299.
5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid benzhydryl-amide. HPLC r.t. 5.02, [M+H]⁺ =371.1.
5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide. HPLC r.t. 5.94, [M+H]⁺ =322.
20 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide. HPLC r.t. 6.26, [M+H]⁺ =438.
5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid benzylamide. HPLC r.t. 5.08, [M+H]⁺ =300.
5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
25 (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 6.7, [M+H]⁺ =340.
5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide. HPLC r.t. 5.57, [M+H]⁺ =304.
5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
30 benzhydryl-amide. HPLC r.t. 6.37, [M+H]⁺ =375.9.
5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide. HPLC r.t. 5.8, [M+H]⁺ =322.
5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide. HPLC r.t. 6.15, [M+H]⁺ =320.
35 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid benzylamide. HPLC r.t. 4.96, [M+H]⁺ =300.

- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 6.55,
[M+H]⁺ = 340.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-
5 fluoro-phenyl)-amide. HPLC r.t. 5.48, [M+H]⁺ = 304.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
benzhydryl-amide. HPLC r.t. 6.24, [M+H]⁺ = 375.9.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2,4-
difluoro-phenyl)-amide. HPLC r.t. 6.82, [M+H]⁺ = 323.
- 10 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
chloro-phenyl)-amide. HPLC r.t. 7.25, [M+H]⁺ = 321.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
benzylamide. HPLC r.t. 5.63, [M+H]⁺ = 301.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
15 (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 7.21,
[M+H]⁺ = 341.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
fluoro-phenyl)-amide. HPLC r.t. 6.5, [M+H]⁺ = 305.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
20 benzhydryl-amide. HPLC r.t. 6.76, [M+H]⁺ = 376.8.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2,4-
difluoro-phenyl)-amide. HPLC r.t. 6.79, [M+H]⁺ = 317.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-
chloro-phenyl)-amide. HPLC r.t. 7.07, [M+H]⁺ = 315.
- 25 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
benzylamide. HPLC r.t. 5.44, [M+H]⁺ = 295.1.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 6.89,
[M+H]⁺ = 335.
- 30 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-
fluoro-phenyl)-amide. HPLC r.t. 6.32, [M+H]⁺ = 299.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
benzhydryl-amide. HPLC r.t. 6.64, [M+H]⁺ = 370.9.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (2,4-
35 difluoro-phenyl)-amide. HPLC r.t. 4.08, [M+H]⁺ = 317.1.

- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide. HPLC r.t. 4.59, $[M+H]^+ = 315.1$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid benzylamide. HPLC r.t. 3.54, $[M+H]^+ = 295.1$.
- 5 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 5, $[M+H]^+ = 335$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide. HPLC r.t. 3.84, $[M+H]^+ = 299.1$.
- 10 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid benzhydryl-amide. HPLC r.t. 5.05, $[M+H]^+ = 371.2$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide. HPLC r.t. 6.04, $[M+H]^+ = 410$.
- 15 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide. HPLC r.t. 6.39, $[M+H]^+ = 408$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (5,6,7,8-tetrahydro-naphthalen-
- 20 1-yl)-amide. HPLC r.t. 6.73, $[M+H]^+ = 428.2$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide. HPLC r.t. 5.78, $[M+H]^+ = 392.1$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
- 25 isoxazole-4-carboxylic acid benzhydryl-amide. HPLC r.t. 6.35, $[M+H]^+ = 464.2$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide. HPLC r.t. 7.36, $[M+H]^+ = 367$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-
- 30 chloro-phenyl)-amide. HPLC r.t. 8.18, $[M+H]^+ = 365$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid benzylamide. HPLC r.t. 6.77, $[M+H]^+ = 345.1$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide. HPLC r.t. 7.99,
- 35 $[M+H]^+ = 385.1$.

- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide. HPLC r.t. 7.45, $[M+H]^+ = 349$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid benzhydryl-amide. HPLC r.t. 7.79, $[M+H]^+ = 421$.
- 5 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 4.41, $[M+H]^+ = 331.1$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 4.12, $[M+H]^+ = 295$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 5.52, $[M+H]^+ = 387.1$.
- 10 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 3.85, $[M+H]^+ = 355$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 4.77, $[M+H]^+ = 321$.
- 15 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 4, $[M+H]^+ = 341$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 6.25, $[M+H]^+ = 336$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid p-
- 20 tolylamide. HPLC r.t. 5.87, $[M+H]^+ = 300$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 6.92, $[M+H]^+ = 392$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 5.29, $[M+H]^+ = 359.9$.
- 25 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 6.48, $[M+H]^+ = 326$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 5.64, $[M+H]^+ = 346$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
- 30 naphthalen-1-ylamide. HPLC r.t. 6.14, $[M+H]^+ = 336$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 5.76, $[M+H]^+ = 300$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. , $[M+H]^+ = 392$.
- 35 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 5.18, $[M+H]^+ = 359.9$.

- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 6.37, $[M+H]^+ = 326$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 5.52, $[M+H]^+ = 346$.
- 5 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 6.95, $[M+H]^+ = 337$.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 6.73, $[M+H]^+ = 301$.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 7.7, $[M+H]^+ = 393$.
- 10 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 5.64, $[M+H]^+ = 360.8$.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 7.38, $[M+H]^+ = 327$.
- 15 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 6.37, $[M+H]^+ = 347$.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 6.75.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 6.53, $[M+H]^+ = 295$.
- 20 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 7.55, $[M+H]^+ = 387$.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 5.39, $[M+H]^+ = 417.1$.
- 25 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 7.17, $[M+H]^+ = 321$.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 6.19, $[M+H]^+ = 341$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 4.44, $[M+H]^+ = 331$.
- 30 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 4.13, $[M+H]^+ = 295$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 5.53, $[M+H]^+ = 387.1$.
- 35 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC r.t. 3.87, $[M+H]^+ = 355$.

- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 4.78, $[M+H]^+ = 321.1$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 3.99, $[M+H]^+ = 341$.
- 5 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 6.37, $[M+H]^+ = 424.1$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 6.01,
- 10 $[M+H]^+ = 388.1$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 6.99, $[M+H]^+ = 480.1$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide. HPLC
- 15 r.t. 5.31, $[M+H]^+ = 448$.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 6.54, $[M+H]^+ = 414.1$.
- 20 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 5.8, $[M+H]^+ = 434.1$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid naphthalen-1-ylamide. HPLC r.t. 7.66, $[M+H]^+ = 381$.
- 25 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid p-tolylamide. HPLC r.t. 7.69, $[M+H]^+ = 345$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide. HPLC r.t. 8.51, $[M+H]^+ = 437.1$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid 2,4-
- 30 dimethoxy-benzylamide. HPLC r.t. 6.6, $[M+H]^+ = 405$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid indan-5-ylamide. HPLC r.t. 8.35, $[M+H]^+ = 371$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide. HPLC r.t. 7.34, $[M+H]^+ = 391$.

- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 6.44, $[M+H]^+ = 401$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3,4,5-trimethoxy-phenyl)-amide. HPLC r.t. 3.27, $[M+H]^+ = 371.2$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-methoxy-5-trifluoromethyl-phenyl)-amide. HPLC r.t. 5.66, $[M+H]^+ = 379.1$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
10 quinolin-8-ylamide. HPLC r.t. 5.08, $[M+H]^+ = 332$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide. HPLC r.t. 5.5, $[M+H]^+ = 337.1$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 7.8, $[M+H]^+ = 406$.
- 15 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3,4,5-trimethoxy-phenyl)-amide. HPLC r.t. 4.87, $[M+H]^+ = 376$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide. HPLC r.t. 6.62, $[M+H]^+ = 336.9$.
- 20 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide. HPLC r.t. 7.04, $[M+H]^+ = 342$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 7.6, $[M+H]^+ = 406$.
- 25 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3,4,5-trimethoxy-phenyl)-amide. HPLC r.t. 4.78, $[M+H]^+ = 376$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2-methoxy-5-trifluoromethyl-phenyl)-amide. HPLC r.t. 6.84, $[M+H]^+ = 384$.
- 30 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide. HPLC r.t. 6.39, $[M+H]^+ = 337$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide. HPLC r.t. 6.95, $[M+H]^+ = 342$.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 8.21, $[M+H]^+ = 407$.
- 35

- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide. HPLC r.t. 6.71, [M+H]⁺ =338.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-tert-
butyl-phenyl)-amide. HPLC r.t. 7.88, [M+H]⁺ =343.
- 5 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 7.98, [M+H]⁺
=401.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3,4,5-
trimethoxy-phenyl)-amide. HPLC r.t. 5.45, [M+H]⁺ =371.
- 10 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-5-trifluoromethyl-phenyl)-amide. HPLC r.t. 7.19,
[M+H]⁺ =379.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide. HPLC r.t. 6.55, [M+H]⁺ =332.
- 15 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-tert-
butyl-phenyl)-amide. HPLC r.t. 7.68, [M+H]⁺ =337.1.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 6.37, [M+H]⁺
=401.1.
- 20 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide. HPLC r.t. 4.98, [M+H]⁺ =332.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-tert-
butyl-phenyl)-amide. HPLC r.t. 5.52, [M+H]⁺ =337.1.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
25 isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-yl)-
amide. HPLC r.t. 7.76, [M+H]⁺ =494.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2-methoxy-5-trifluoromethyl-
phenyl)-amide. HPLC r.t. 7.03, [M+H]⁺ =472.
- 30 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide.
HPLC r.t. 7.07, [M+H]⁺ =430.2.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide. HPLC r.t. 8.77, [M+H]⁺
35 =451.

- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (3,4,5-trimethoxy-phenyl)-amide. HPLC r.t. 6.61, $[M+H]^+ = 421$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid quinolin-8-ylamide. HPLC r.t. 7.14, $[M+H]^+ = 382$.
- 5 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide. HPLC r.t. 8.75, $[M+H]^+ = 387.1$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 5.13, $[M+H]^+ = 371$.
- 3-[(5-Amino-3-pyridin-4-yl-isoxazole-4-carbonyl)-amino]-
- 10 benzoic acid ethyl ester. HPLC r.t. 4.44, $[M+H]^+ = 353$.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 5.13, $[M+H]^+ = 365$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2-
- 15 benzyl-phenyl)-amide. HPLC r.t. 6.6, $[M+H]^+ = 376$.
- 3-[(5-Amino-3-thiophen-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester. HPLC r.t. 6.02, $[M+H]^+ = 358$.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 6.59, $[M+H]^+ = 370$.
- 20 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide. HPLC r.t. 5.32, $[M+H]^+ = 329$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 6.47, $[M+H]^+ = 376$.
- 25 3-[(5-Amino-3-thiophen-3-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester. HPLC r.t. 5.92, $[M+H]^+ = 357.9$.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 6.49, $[M+H]^+ = 370$.
- 30 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide. HPLC r.t. 6.23, $[M+H]^+ = 330.1$.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 7.04, $[M+H]^+ = 377$.
- 3-[(5-Amino-3-thiazol-2-yl-isoxazole-4-carbonyl)-amino]-
- 35 benzoic acid ethyl ester. HPLC r.t. 6.97, $[M+H]^+ = 358.9$.

- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 7.5, $[M+H]^+ = 371$.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide. HPLC r.t. 5.98.
- 5 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 6.74, $[M+H]^+ = 371$.
- 3-[(5-Amino-3-pyridin-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester. HPLC r.t. 6.84, $[M+H]^+ = 353$.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 7.34, $[M+H]^+ = 365$.
- 10
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide. HPLC r.t. 3.66, $[M+H]^+ = 324$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 5.16, $[M+H]^+ = 371.2$.
- 15
- 3-[(5-Amino-3-pyridin-3-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester. HPLC r.t. 4.45, $[M+H]^+ = 353.1$.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 5.15, $[M+H]^+ = 365$.
- 20
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 6.72, $[M+H]^+ = 464.2$.
- 3-{[5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carbonyl]-amino}-benzoic acid ethyl ester. HPLC r.t. 6.18, $[M+H]^+ = 446.1$.
- 25
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 6.69, $[M+H]^+ = 458.1$.
- 30
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide. HPLC r.t. 7.27, $[M+H]^+ = 374.1$.
- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide. HPLC r.t. 7.6, $[M+H]^+ = 421$.
- 3-[(5-Amino-3-quinolin-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester. HPLC r.t. 7.99, $[M+H]^+ = 403$.
- 35

- 5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide. HPLC r.t. 8.33.
- 5-amino-N-benzyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 427.11.
- 5-amino-N-(2-furylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 417.09.
- 5-amino-N-(2-phenylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 441.13.
- 5-amino-N-cyclohexyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 419.15.
- 5-amino-N-(4-methoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 457.12.
- 5-amino-N-hexyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.76, mass found = 421.16.
- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 421.12.
- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 433.07.
- 5-amino-N-cyclopentyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 405.13.
- 5-amino-N-(1,5-dimethylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.87, mass found = 449.19.
- 5-amino-N-isopentyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 407.15.

5-amino-N-(2-ethylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.87, mass found = 449.19.

5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 469.16.

5-amino-N-(2-methylcyclohexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 433.16.

10 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 563.09.

5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 495.1.

5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 455.15.

5-amino-N-isobutyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 393.13.

5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 495.1.

25 5-amino-N-(2-methoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 457.12.

5-amino-N-(2-fluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 445.1.

5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 495.1.

5-amino-N-(3,4-difluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 463.1.

- 5-amino-N-(3-chlorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.68, mass found = 461.08.
- 5-amino-N-(2-chlorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
5 HPLC (Method B) r.t. 1.67, mass found = 461.08.
- 5-amino-N-(3-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.67, mass found = 441.13.
- 10 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 477.13.
- 5-amino-N-(cyclopropylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
15 HPLC (Method B) r.t. 1.57, mass found = 391.11.
- 5-amino-N-cycloheptyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.75, mass found = 433.16.
- 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
20 HPLC (Method B) r.t. 1.61, mass found = 471.14.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.77, mass found = 445.16.
- 25 5-amino-N-(1,2-dimethylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.65, mass found = 407.15.
- 5-amino-N-(3,3-diphenylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
30 HPLC (Method B) r.t. 1.8, mass found = 531.18.
- 5-amino-N-(1,2-diphenylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.77, mass found = 517.16.
- 5-amino-N-(2,4-dimethoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
35 HPLC (Method B) r.t. 1.59, mass found = 487.14.

- 5-amino-N-[3-(dibutylamino)propyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 506.25.
- 5-amino-N-butyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 393.13.
- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found = 511.1.
- 10 5-amino-N-(1-methylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.82, mass found = 435.18.
- 5-amino-N-(2-methoxy-1-methylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 409.12.
- 15 5-amino-N-(1-methylbutyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 407.15.
- 5-amino-N-(3-fluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 445.1.
- 20 5-amino-N-(4-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 441.13.
- 25 5-amino-N-(1,3-dimethylbutyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 421.16.
- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.75, mass found = 469.16.
- 30 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 425.1.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 471.1.
- 35

- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 407.15.
- 5-amino-N-(cyclohexylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.77, mass found = 433.16.
- 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 475.09.
- 10 5-amino-N-(1-ethylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 407.15.
- 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 441.13.
- 15 5-amino-N-(2-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 441.13.
- 5-amino-N-cyclobutyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 391.11.
- 20 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.92, mass found = 449.19.
- 25 5-amino-N-(2-bromobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 505.02.
- 5-amino-N-(1-methylheptyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.89, mass found = 449.19.
- 30 5-amino-N-heptyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.84, mass found = 435.18.
- 5-amino-N-(tert-butyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 393.13.
- 35

- 5-amino-N-(3,4-dimethoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.51, mass found = 487.14.
- 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
5 HPLC (Method B) r.t. 1.72, mass found = 453.13.
- 5-amino-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide.
HPLC (Method B) r.t. 1.74, mass found = 467.15.
- 10 5-amino-N-benzyl-3-(5-bromothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 376.98.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 366.96.
15
- 5-amino-3-(5-bromothien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 391.
- 5-amino-3-(5-bromothien-2-yl)-N-(sec-butyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 343.
- 20 5-amino-3-(5-bromothien-2-yl)-N-cyclohexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 369.01.
- 5-amino-3-(5-bromothien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 406.99.
25
- 5-amino-3-(5-bromothien-2-yl)-N-isopropylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 328.98.
- 5-amino-3-(5-bromothien-2-yl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 371.03.
30
- N-allyl-5-amino-3-(5-bromothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 326.97.

- 72 -

- 5-amino-3-(5-bromothien-2-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 370.99.
- 5-amino-3-(5-bromothien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 382.94.
- 5-amino-3-(5-bromothien-2-yl)-N-cyclopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 355.
- 5-amino-3-(5-bromothien-2-yl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 399.06.
- 5-amino-3-(5-bromothien-2-yl)-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 357.01.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.79, mass found = 399.06.
- 5-amino-3-(5-bromothien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 419.03.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 383.03.
- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-bromothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 512.96.
- 5-amino-3-(5-bromothien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 444.97.
- 5-amino-3-(5-bromothien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 405.01.
- 5-amino-3-(5-bromothien-2-yl)-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 343.

- 5-amino-3-(5-bromothien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 444.97.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 406.99.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 394.97.
- 10 5-amino-3-(5-bromothien-2-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 444.97.
- 5-amino-3-(5-bromothien-2-yl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 412.96.
- 15 5-amino-3-(5-bromothien-2-yl)-N-(3-chlorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 410.94.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-chlorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 410.94.
- 20 5-amino-3-(5-bromothien-2-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 391.
- 5-amino-3-(5-bromothien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 427.
- 25 5-amino-3-(5-bromothien-2-yl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 340.98.
- 30 5-amino-3-(5-bromothien-2-yl)-N-cycloheptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 383.03.
- 5-amino-3-(5-bromothien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 421.01.
- 35

- 5-amino-3-(5-bromothien-2-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 395.03.
- 5-amino-3-(5-bromothien-2-yl)-N-prop-2-ynylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 324.95.
- 5-amino-3-(5-bromothien-2-yl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 357.01.
- 10 5-amino-3-(5-bromothien-2-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 481.05.
- 5-amino-3-(5-bromothien-2-yl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 467.03.
- 15 5-amino-3-(5-bromothien-2-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 456.12.
- 5-amino-3-(5-bromothien-2-yl)-N-butylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 343.
- 20 5-amino-3-(5-bromothien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 460.97.
- 5-amino-3-(5-bromothien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 380.98.
- 25 5-amino-3-(5-bromothien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 385.05.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 358.99.
- 30 5-amino-3-(5-bromothien-2-yl)-N-(4-tert-butylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.88, mass found = 425.08.

- 5-amino-3-(5-bromothien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 357.01.
- 5-amino-3-(5-bromothien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 394.97.
- 5-amino-3-(5-bromothien-2-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 391.
- 5-amino-3-(5-bromothien-2-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 371.03.
- 5-amino-3-(5-bromothien-2-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 419.03.
- 5-amino-3-(5-bromothien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 374.97.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-bromothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 420.97.
- 5-amino-3-(5-bromothien-2-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 357.01.
- 5-amino-3-(5-bromothien-2-yl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 383.03.
- ethyl 4-([5-amino-3-(5-bromothien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.35, mass found = 442.03.
- 5-amino-3-(5-bromothien-2-yl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 424.96.
- 5-amino-3-(5-bromothien-2-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 357.01.

- 5-amino-3-(5-bromothien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 391.
- 5-amino-3-(5-bromothien-2-yl)-N-propylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found =
- 5 328.98.
- 5-amino-3-(5-bromothien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 391.
- 5-amino-3-(5-bromothien-2-yl)-N-cyclobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found =
- 10 340.98.
- 5-amino-3-(5-bromothien-2-yl)-N-(3-methoxypropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found =
- 358.99.
- 5-amino-3-(5-bromothien-2-yl)-N-octylisoxazole-4-
- 15 carboxamide. HPLC (Method B) r.t. 1.82, mass found =
- 399.06.
- 5-amino-N-(2-bromobenzyl)-3-(5-bromothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found =
- 454.89.
- 20 5-amino-3-(5-bromothien-2-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.8, mass found =
- 399.06.
- 5-amino-3-(5-bromothien-2-yl)-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found =
- 25 385.05.
- 5-amino-3-(5-bromothien-2-yl)-N-(tert-butyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 343.
- 5-amino-3-(5-bromothien-2-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass
- 30 found = 403.
- 5-amino-3-(5-bromothien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 417.01.
- 5-amino-N-benzyl-3-(5-chlorothien-2-yl)isoxazole-4-
- 35 carboxamide. HPLC (Method B) r.t. 1.46, mass found =
- 333.03.

- 5-amino-3-(5-chlorothien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 323.01.
- 5-amino-3-(5-chlorothien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 347.05.
- 5-amino-N-(sec-butyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 299.05.
- 10 5-amino-3-(5-chlorothien-2-yl)-N-cyclohexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 325.07.
- 5-amino-3-(5-chlorothien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 363.04.
- 15 5-amino-3-(5-chlorothien-2-yl)-N-isopropylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 285.03.
- 5-amino-3-(5-chlorothien-2-yl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 327.08.
- 20 N-allyl-5-amino-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 283.02.
- 25 5-amino-3-(5-chlorothien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 338.99.
- 5-amino-3-(5-chlorothien-2-yl)-N-cyclopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 311.05.
- 30 5-amino-3-(5-chlorothien-2-yl)-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 313.07.
- 5-amino-3-(5-chlorothien-2-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.81, mass found = 355.11.
- 35

- 78 -

- 5-amino-3-(5-chlorothien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 375.08.
- 5-amino-3-(5-chlorothien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 339.08.
- 5-amino-3-(5-chlorothien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 401.02.
- 10 5-amino-3-(5-chlorothien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 351.02.
- 5-amino-3-(5-chlorothien-2-yl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 369.02.
- 15 5-amino-N-(3-chlorobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 366.99.
- 5-amino-N-(2-chlorobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 366.99.
- 20 5-amino-3-(5-chlorothien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 383.05.
- 25 5-amino-3-(5-chlorothien-2-yl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 297.03.
- 5-amino-3-(5-chlorothien-2-yl)-N-cycloheptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 339.08.
- 30 5-amino-3-(5-chlorothien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 377.06.
- 5-amino-3-(5-chlorothien-2-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 351.08.
- 35

- 79 -

- 5-amino-3-(5-chlorothien-2-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 437.1.
- 5-amino-3-(5-chlorothien-2-yl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 423.08.
- 5-amino-3-(5-chlorothien-2-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 412.17.
- 10 5-amino-N-butyl-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 299.05.
- 5-amino-3-(5-chlorothien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 417.02.
- 15 5-amino-3-(5-chlorothien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 337.03.
- 5-amino-3-(5-chlorothien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found = 341.1.
- 20 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.16, mass found = 416.11.
- 25 5-amino-3-(5-chlorothien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 315.04.
- 5-amino-3-(5-chlorothien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 313.07.
- 30 5-amino-3-(5-chlorothien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 351.02.
- 5-amino-3-(5-chlorothien-2-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 327.08.
- 35

- 5-amino-3-(5-chlorothien-2-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 375.08.
- 5-amino-3-(5-chlorothien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 331.02.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 377.02.
- 10 5-amino-3-(5-chlorothien-2-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 313.07.
- 5-amino-3-(5-chlorothien-2-yl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 339.08.
- 15 ethyl 4-([5-amino-3-(5-chlorothien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.35, mass found = 398.08.
- 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 381.01.
- 20 5-amino-3-(5-chlorothien-2-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 313.07.
- 25 5-amino-3-(5-chlorothien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 347.05.
- 5-amino-3-(5-chlorothien-2-yl)-N-propylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 285.03.
- 30 5-amino-3-(5-chlorothien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 347.05.
- 5-amino-3-(5-chlorothien-2-yl)-N-cyclobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 297.03.
- 35

5-amino-3-(5-chlorothien-2-yl)-N-(3-methoxypropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.26, mass found = 315.04.

5-amino-N-(2-bromobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 410.94.

5-amino-3-(5-chlorothien-2-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.82, mass found = 355.11.

10 5-amino-3-(5-chlorothien-2-yl)-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 341.1.
5-amino-N-(tert-butyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 299.05.

15 5-amino-3-(5-chlorothien-2-yl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 393.05.

5-amino-3-(5-chlorothien-2-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass
20 found = 359.05.

5-amino-3-(5-chlorothien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 373.07.

5-amino-N-benzyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 313.09.

5-amino-N-(2-furylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 303.07.

30 5-amino-3-(5-methylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 327.1.

5-amino-N-(sec-butyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 279.1.

5-amino-N-cyclohexyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 305.12.

5-amino-N-(4-methoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 343.1.

5-amino-N-isopropyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 265.09.

10 5-amino-N-hexyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 307.14.

N-allyl-5-amino-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 263.07.

5-amino-3-(5-methylthien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 319.04.

20 5-amino-N-cyclopentyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 291.1.

5-amino-N-(1,5-dimethylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found = 335.17.

25 5-amino-N-isopentyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 293.12.

5-amino-N-(2-ethylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found = 335.17.

30 5-amino-3-(5-methylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 355.14.

5-amino-N-(2-methylcyclohexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 319.14.

35

- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 449.06.
- 5-amino-3-(5-methylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 381.08.
- 5-amino-3-(5-methylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 341.12.
- 10 5-amino-N-isobutyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 279.1.
- 5-amino-3-(5-methylthien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 381.08.
- 15 5-amino-N-(2-methoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 343.1.
- 5-amino-N-(2-fluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 331.08.
- 20 5-amino-N-(3,4-difluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 349.07.
- 5-amino-N-(3-chlorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 347.05.
- 25 5-amino-N-cycloheptyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 319.14.
- 30 5-amino-N-(1,2-dimethylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 293.12.
- 5-amino-N-(3,3-diphenylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 417.15.
- 35

- 5-amino-N-(1,2-diphenylethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 403.14.
- 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 373.11.
- 5-amino-N-[3-(dibutylamino)propyl]-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 392.22.
- 10 5-amino-N-butyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 279.1.
- 5-amino-3-(5-methylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 397.07.
- 15 5-amino-N-[(5-methyl-2-furyl)methyl]-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 317.08.
- 5-amino-N-(1-methylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found =
- 20 321.15.
- 5-amino-N-(2-methoxy-1-methylethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 295.1.
- 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.83, mass found = 361.18.
- 25 5-amino-N-(1-methylbutyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 293.12.
- 30 5-amino-N-(3-fluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 331.08.
- 5-amino-N-(4-methylbenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found =
- 35 327.1.

- 5-amino-N-(1-methyl-3-phenylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 355.14.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 357.08.
- 5-amino-3-(5-methylthien-2-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 293.12.
- 10 5-amino-N-(cyclohexylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 319.14.
- 5-amino-3-(5-methylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 327.1.
- 15 5-amino-3-(5-methylthien-2-yl)-N-propylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 265.09.
- 5-amino-N-(2-methylbenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 327.1.
- 20 5-amino-N-cyclobutyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 277.09.
- 25 5-amino-N-(3-methoxypropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.17, mass found = 295.1.
- 5-amino-3-(5-methylthien-2-yl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.76, mass found = 335.17.
- 30 5-amino-N-(1-methylheptyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 335.17.
- 5-amino-N-heptyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 321.15.
- 35

- 5-amino-N-(tert-butyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 279.1.
- 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass
5 found = 373.11.
- 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 339.1.
- 5-amino-3-(5-methylthien-2-yl)-N-(1,2,3,4-
10 tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 353.12.
- 5-amino-N-benzyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 313.09.
- 15 5-amino-N-(2-furylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 303.07.
- 5-amino-3-(3-methylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found =
20 327.1.
- 5-amino-N-(sec-butyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 279.1.
- 5-amino-N-cyclohexyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found =
25 305.12.
- 5-amino-N-(4-methoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 343.1.
- 5-amino-N-isopropyl-3-(3-methylthien-2-yl)isoxazole-4-
30 carboxamide. HPLC (Method B) r.t. 1.25, mass found = 265.09.
- 5-amino-N-hexyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 307.14.

N-allyl-5-amino-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 263.07.

5 5-amino-3-(3-methylthien-2-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.14, mass found = 307.1.

5-amino-N-cyclopentyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 291.1.

10 5-amino-N-(1,5-dimethylhexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 335.17.

5-amino-N-isopentyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 293.12.

15 5-amino-N-(2-ethylhexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 335.17.

5-amino-3-(3-methylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 355.14.

20 5-amino-N-(2-methylcyclohexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 319.14.

5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 449.06.

5-amino-3-(3-methylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 381.08.

30 5-amino-3-(3-methylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 341.12.

5-amino-N-isobutyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 279.1.

35 5-amino-3-(3-methylthien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 381.08.

- 5-amino-N-(2-methoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 343.1.
- 5-amino-N-(2-fluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-
5 4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 331.08.
- 5-amino-3-(3-methylthien-2-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 381.08.
- 10 5-amino-N-(3,4-difluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 349.07.
- 5-amino-N-(3-chlorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found =
15 347.05.
- 5-amino-N-(2-chlorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 347.05.
- 5-amino-N-(3-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-
20 4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 327.1.
- 5-amino-3-(3-methylthien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 363.1.
- 25 5-amino-N-(cyclopropylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 277.09.
- 5-amino-N-cycloheptyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found =
30 319.14.
- 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 357.11.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass
35 found = 331.14.

- 5-amino-N-(1,2-dimethylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 293.12.
- 5-amino-N-(3,3-diphenylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 417.15.
- 5-amino-N-(2,4-dimethoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 373.11.
- 10 5-amino-N-[3-(dibutylamino)propyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.17, mass found = 392.22.
- 5-amino-N-butyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 279.1.
- 15 5-amino-3-(3-methylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 397.07.
- 5-amino-N-[(5-methyl-2-furyl)methyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 317.08.
- 20 5-amino-N-(1-benzylpiperidin-4-yl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.05, mass found = 396.16.
- 5-amino-N-(2-methoxy-1-methylethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 295.1.
- 25 5-amino-N-(4-tert-butylcyclohexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.8, mass found = 361.18.
- 30 5-amino-N-(1-methylbutyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 293.12.
- 5-amino-N-(3-fluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 331.08.
- 35

- 5-amino-N-(4-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 327.1.
- 5-amino-N-(1,3-dimethylbutyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 307.14.
- 5-amino-N-(1-methyl-3-phenylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 355.14.
- 10 5-amino-3-(3-methylthien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 311.08.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 357.08.
- 15 5-amino-3-(3-methylthien-2-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 293.12.
- 5-amino-N-(cyclohexylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 319.14.
- 20 ethyl 4-([5-amino-3-(3-methylthien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.24, mass found = 378.14.
- 25 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 361.07.
- 5-amino-N-(1-ethylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 293.12.
- 30 5-amino-3-(3-methylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 327.1.
- 5-amino-N-(2-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 327.1.
- 35

- 5-amino-N-cyclobutyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 277.09.
- 5-amino-N-benzhydryl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 389.12.
- 5-amino-3-(3-methylthien-2-yl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 335.17.
- 10 5-amino-N-(2-bromobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 391.
- 5-amino-N-(1-methylheptyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 335.17.
- 15 5-amino-N-heptyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 321.15.
- 5-amino-N-(tert-butyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 279.1.
- 20 5-amino-N-(3,4-dimethoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 373.11.
- 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 339.1.
- 25 5-amino-3-(3-methylthien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 353.12.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 383.07.
- 30 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 407.1.

- 5-amino-N-(sec-butyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 359.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclohexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 385.12.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 423.1.
- 10 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-isopropylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 345.09.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.79, mass found =
- 15 387.14.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 387.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 371.1.
- 20 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.9, mass found = 415.17.
- 25 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 373.12.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-ethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.91, mass found = 415.17.
- 30 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.77, mass found = 435.14.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 399.14.
- 35

- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.79, mass found = 529.06.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 461.08.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 421.12.
- 10 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 359.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 461.08.
- 15 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 423.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 411.08.
- 20 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 461.08.
- 25 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 429.07.
- 5-amino-N-(3-chlorobenzyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.7, mass found = 427.05.
- 30 5-amino-N-(2-chlorobenzyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 427.05.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 407.1.
- 35

- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 443.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(cyclopropylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 357.09.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cycloheptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 399.14.
- 10 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 437.11.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.79, mass found = 411.14.
- 15 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 373.12.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.8, mass found = 497.15.
- 20 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2-diphenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 483.14.
- 25 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 453.11.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 472.22.
- 30 5-amino-N-butyl-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 359.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 477.07.
- 35

- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 397.08.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.84, mass found = 401.15.
- 5-amino-N-(1-benzylpiperidin-4-yl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 476.16.
- 10 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.76, mass found = 387.14.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.77, mass found = 435.14.
- 15 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 391.08.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 373.12.
- 20 ethyl 4-[(5-amino-3-[5-(3-chlorophenyl)-2-furyl]isoxazol-4-yl)carbonyl]amino]piperidine-1-carboxylate. HPLC (Method B) r.t. 1.48, mass found = 458.14.
- 25 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 441.06.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-ethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 373.12.
- 30 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 407.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 407.1.
- 35

- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 357.09.
- 5-amino-N-benzhydryl-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.76, mass found = 469.12.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.96, mass found = 415.17.
- 10 5-amino-N-(2-bromobenzyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 471.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.93, mass found = 415.17.
- 15 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.87, mass found = 401.15.
- 5-amino-N-(tert-butyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 359.1.
- 20 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 453.11.
- 25 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 419.1.
- 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 433.12.
- 30 5-amino-N-benzyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 297.11.
- 5-amino-N-(2-furylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.25, mass found = 287.09.
- 35

- 5-amino-3-(5-methyl-2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 311.13.
- 5-amino-N-(sec-butyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 263.13.
- 5-amino-N-cyclohexyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 289.14.
- 10 5-amino-N-(4-methoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 327.12.
- 5-amino-N-hexyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 291.16.
- 15 5-amino-3-(5-methyl-2-furyl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.14, mass found = 291.12.
- 5-amino-3-(5-methyl-2-furyl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 303.07.
- 20 5-amino-N-cyclopentyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 275.13.
- 25 5-amino-N-(1,5-dimethylhexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 319.19.
- 5-amino-N-isopentyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 277.14.
- 30 5-amino-N-(2-ethylhexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 319.19.
- 5-amino-3-(5-methyl-2-furyl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 339.16.
- 35

- 5-amino-N-(2-methylcyclohexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 303.16.
- 5-amino-3-(5-methyl-2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 365.1.
- 5-amino-3-(5-methyl-2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 325.14.
- 10 5-amino-N-isobutyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 263.13.
- 5-amino-3-(5-methyl-2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 365.1.
- 15 5-amino-N-(2-methoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 327.12.
- 5-amino-N-(2-fluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 315.1.
- 20 5-amino-3-(5-methyl-2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 365.1.
- 5-amino-N-(3,4-difluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 333.09.
- 25 5-amino-N-(3-chlorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 331.07.
- 5-amino-N-(2-chlorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 331.07.
- 30 5-amino-N-(3-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 311.13.

- 5-amino-3-(5-methyl-2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 347.13.
- 5-amino-N-(cyclopropylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 261.11.
- 5-amino-N-cycloheptyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 303.16.
- 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 341.14.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 315.16.
- 5-amino-N-(1,2-dimethylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 277.14.
- 5-amino-N-(3,3-diphenylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 401.17.
- 5-amino-N-(1,2-diphenylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 387.16.
- 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 357.13.
- 5-amino-N-[3-(dibutylamino)propyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.16, mass found = 376.25.
- 5-amino-N-butyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 263.13.
- 5-amino-3-(5-methyl-2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 381.09.

- 100 -

- 5-amino-3-(5-methyl-2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 301.11.
- 5-amino-3-(5-methyl-2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 305.17.
- 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.03, mass found = 380.18.
- 10 5-amino-N-(2-methoxy-1-methylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.17, mass found = 279.12.
- 5-amino-N-(1-methylbutyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 277.14.
- 15 5-amino-N-(3-fluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 315.1.
- 5-amino-N-(4-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 311.13.
- 20 5-amino-N-(1,3-dimethylbutyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 291.16.
- 5-amino-3-(5-methyl-2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 339.16.
- 25 5-amino-3-(5-methyl-2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 295.1.
- 30 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 341.1.
- 5-amino-3-(5-methyl-2-furyl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 277.14.
- 35

- 5-amino-N-(cyclohexylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 303.16.
- ethyl 4-([5-amino-3-(5-methyl-2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.21, mass found = 362.16.
- 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 345.09.
- 10 5-amino-N-(1-ethylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 277.14.
- 5-amino-3-(5-methyl-2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 311.13.
- 15 5-amino-N-(2-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 311.13.
- 5-amino-N-cyclobutyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.26, mass found = 261.11.
- 20 5-amino-N-benzhydryl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 373.14.
- 25 5-amino-3-(5-methyl-2-furyl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 319.19.
- 5-amino-N-(2-bromobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 375.02.
- 30 5-amino-3-(5-methyl-2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 319.19.
- 5-amino-N-heptyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 305.17.
- 35

- 5-amino-N-(tert-butyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 263.13.
- 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 357.13.
- 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 323.13.
- 10 5-amino-3-(5-methyl-2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 337.14.
- 5-amino-N-benzyl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.26, mass found = 283.1.
- 15 5-amino-3-(2-furyl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.15, mass found = 273.08.
- 5-amino-3-(2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 297.11.
- 20 5-amino-N-(sec-butyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 249.11.
- 5-amino-3-(2-furyl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 313.11.
- 25 5-amino-3-(2-furyl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 277.14.
- 5-amino-3-(2-furyl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.02, mass found = 277.11.
- 30 5-amino-3-(2-furyl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 289.05.
- 5-amino-N-cyclopentyl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 261.11.

- 103 -

- 5-amino-N-(1,5-dimethylhexyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 305.17.
- 5-amino-3-(2-furyl)-N-isopentylisoxazole-4-carboxamide.
5 HPLC (Method B) r.t. 1.31, mass found = 263.13.
- 5-amino-N-(2-ethylhexyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 305.17.
- 5-amino-3-(2-furyl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found =
10 325.14.
- 5-amino-3-(2-furyl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 289.14.
- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54,
15 mass found = 419.07.
- 5-amino-3-(2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 351.08.
- 5-amino-3-(2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found =
20 311.13.
- 5-amino-3-(2-furyl)-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.21, mass found = 249.11.
- 5-amino-3-(2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found =
25 351.08.
- 5-amino-3-(2-furyl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 313.11.
- 30 5-amino-N-(2-fluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 301.09.
- 5-amino-3-(2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found =
35 351.08.

- 5-amino-N-(3,4-difluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 319.08.
- 5-amino-N-(3-chlorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 317.06.
- 5-amino-N-(2-chlorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 317.06.
- 10 5-amino-3-(2-furyl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 297.11.
- 5-amino-3-(2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 333.11.
- 15 5-amino-N-(cyclopropylmethyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.13, mass found = 247.1.
- 5-amino-N-cycloheptyl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 289.14.
- 20 5-amino-3-(2-furyl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 327.12.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 301.14.
- 25 5-amino-N-(1,2-dimethylpropyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 263.13.
- 5-amino-N-(3,3-diphenylpropyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 387.16.
- 30 5-amino-N-(1,2-diphenylethyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 373.14.

- 105 -

- 5-amino-N-(2,4-dimethoxybenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 343.12.
- 5-amino-N-[3-(dibutylamino)propyl]-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.09, mass found = 362.23.
- 5-amino-N-butyl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 249.11.
- 5-amino-3-(2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 367.08.
- 5-amino-3-(2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 287.09.
- 5-amino-3-(2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 291.16.
- 5-amino-N-(1-benzylpiperidin-4-yl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 366.17.
- 5-amino-3-(2-furyl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.05, mass found = 265.11.
- 5-amino-N-(4-tert-butylcyclohexyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 331.19.
- 5-amino-3-(2-furyl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 263.13.
- 5-amino-N-(3-fluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 301.09.
- 5-amino-3-(2-furyl)-N-(4-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 297.11.

- 5-amino-N-(1,3-dimethylbutyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 277.14.
- 5-amino-3-(2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 325.14.
- 5-amino-3-(2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.16, mass found = 281.08.
- 10 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 327.09.
- 5-amino-3-(2-furyl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 263.13.
- 15 5-amino-N-(cyclohexylmethyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 289.14.
- ethyl 4-([5-amino-3-(2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.14, mass found = 348.14.
- 20 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 331.07.
- 5-amino-N-(1-ethylpropyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 263.13.
- 25 5-amino-3-(2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 297.11.
- 30 5-amino-3-(2-furyl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 297.11.
- 5-amino-N-cyclobutyl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.15, mass found = 247.1.
- 35 5-amino-N-benzhydryl-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 359.13.

- 5-amino-3-(2-furyl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 305.17.
- 5-amino-N-(2-bromobenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found =
- 5 361.01.
- 5-amino-3-(2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 305.17.
- 5-amino-3-(2-furyl)-N-heptylisoxazole-4-carboxamide. HPLC
- 10 (Method B) r.t. 1.54, mass found = 291.16.
- 5-amino-N-(tert-butyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.25, mass found = 249.11.
- 5-amino-N-(3,4-dimethoxybenzyl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.14, mass found =
- 15 343.12.
- 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 309.11.
- 5-amino-3-(2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-
- 20 yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 323.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-benzylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 333.11.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 323.09.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found =
- 30 347.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-(sec-butyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 299.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-cyclohexylisoxazole-4-
- 35 carboxamide. HPLC (Method B) r.t. 1.57, mass found = 325.14.

- 5-amino-3-(1-benzofuran-3-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 363.12.
- 5-amino-3-(1-benzofuran-3-yl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 327.16.
- 5-amino-3-(1-benzofuran-3-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 327.12.
- 10 5-amino-3-(1-benzofuran-3-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 339.07.
- 5-amino-3-(1-benzofuran-3-yl)-N-cyclopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 311.13.
- 15 5-amino-3-(1-benzofuran-3-yl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.78, mass found = 355.19.
- 5-amino-3-(1-benzofuran-3-yl)-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 313.14.
- 20 5-amino-3-(1-benzofuran-3-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.8, mass found = 355.19.
- 5-amino-3-(1-benzofuran-3-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 375.16.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 339.16.
- 30 5-amino-3-(1-benzofuran-3-yl)-N-[3,5-bis(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 469.09.
- 5-amino-3-(1-benzofuran-3-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 401.1.
- 35

- 109 -

- 5-amino-3-(1-benzofuran-3-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 361.14.
- 5-amino-3-(1-benzofuran-3-yl)-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 299.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 401.1.
- 10 5-amino-3-(1-benzofuran-3-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 363.12.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 351.1.
- 15 5-amino-3-(1-benzofuran-3-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 401.1.
- 5-amino-3-(1-benzofuran-3-yl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 369.09.
- 20 5-amino-3-(1-benzofuran-3-yl)-N-(3-chlorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 367.07.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-(2-chlorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 367.07.
- 5-amino-3-(1-benzofuran-3-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 347.13.
- 30 5-amino-3-(1-benzofuran-3-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 383.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 297.11.
- 35

- 110 -

- 5-amino-3-(1-benzofuran-3-yl)-N-cycloheptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 339.16.
- 5-amino-3-(1-benzofuran-3-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 377.14.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 351.16.
- 10 5-amino-3-(1-benzofuran-3-yl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 313.14.
- 5-amino-3-(1-benzofuran-3-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 437.17.
- 15 5-amino-3-(1-benzofuran-3-yl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 423.16.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 393.13.
- 20 5-amino-3-(1-benzofuran-3-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 412.25.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-butylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 299.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 417.09.
- 30 5-amino-3-(1-benzofuran-3-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 337.11.
- 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 341.17.
- 35

- 111 -

- 5-amino-3-(1-benzofuran-3-yl)-N-(1-benzylpiperidin-4-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.16, mass found = 416.18.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 315.12.
- 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 313.14.
- 10 5-amino-3-(1-benzofuran-3-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 351.1.
- 5-amino-3-(1-benzofuran-3-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 327.16.
- 15 5-amino-3-(1-benzofuran-3-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 375.16.
- 5-amino-3-(1-benzofuran-3-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 331.1.
- 20 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(1-benzofuran-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 377.1.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 313.14.
- 5-amino-3-(1-benzofuran-3-yl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 339.16.
- 30 ethyl 4-([5-amino-3-(1-benzofuran-3-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.34, mass found = 398.16.
- 5-amino-3-(1-benzofuran-3-yl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 381.09.
- 35

- 5-amino-3-(1-benzofuran-3-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 313.14.
- 5-amino-3-(1-benzofuran-3-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 347.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 347.13.
- 10 5-amino-3-(1-benzofuran-3-yl)-N-cyclobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 297.11.
- 5-amino-N-benzhydryl-3-(1-benzofuran-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 409.14.
- 15 5-amino-3-(1-benzofuran-3-yl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.83, mass found = 355.19.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2-bromobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 411.02.
- 20 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.8, mass found = 355.19.
- 25 5-amino-3-(1-benzofuran-3-yl)-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 341.17.
- 5-amino-3-(1-benzofuran-3-yl)-N-(tert-butyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 299.13.
- 30 5-amino-3-(1-benzofuran-3-yl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 393.13.
- 5-amino-3-(1-benzofuran-3-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 359.13.
- 35

- 5-amino-3-(1-benzofuran-3-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 373.14.
- 5-amino-N-benzyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 327.1.
- 5-amino-3-(5-ethylthien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 317.08.
- 5-amino-3-(5-ethylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found =
- 10 341.12.
- 5-amino-N-(sec-butyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 293.12.
- 5-amino-N-cyclohexyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found =
- 15 319.14.
- 5-amino-3-(5-ethylthien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 357.11.
- 20 5-amino-3-(5-ethylthien-2-yl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 321.15.
- 5-amino-3-(5-ethylthien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t.
- 25 1.46, mass found = 333.06.
- 5-amino-N-cyclopentyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 305.12.
- 5-amino-N-(1,5-dimethylhexyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.83, mass
- 30 found = 349.18.
- 5-amino-3-(5-ethylthien-2-yl)-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 307.14.
- 5-amino-N-(2-ethylhexyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.83, mass found =
- 35 349.18.

- 5-amino-3-(5-ethylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 369.15.
- 5-amino-3-(5-ethylthien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 333.15.
- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 463.08.
- 10 5-amino-3-(5-ethylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 395.09.
- 5-amino-3-(5-ethylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 355.14.
- 15 5-amino-3-(5-ethylthien-2-yl)-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 293.12.
- 5-amino-3-(5-ethylthien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 395.09.
- 20 5-amino-3-(5-ethylthien-2-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 357.11.
- 25 5-amino-3-(5-ethylthien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 345.09.
- 5-amino-N-(3,4-difluorobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 363.09.
- 30 5-amino-N-(3-chlorobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 361.07.
- 5-amino-N-(2-chlorobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 361.07.
- 35

- 5-amino-3-(5-ethylthien-2-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 341.12.
- 5-amino-3-(5-ethylthien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 377.12.
- 5-amino-N-(cyclopropylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 291.1.
- 10 5-amino-N-cycloheptyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 333.15.
- 5-amino-3-(5-ethylthien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 371.13.
- 15 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.69, mass found = 345.15.
- 5-amino-N-(1,2-dimethylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 307.14.
- 20 5-amino-N-(3,3-diphenylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.72, mass found = 431.17.
- 25 5-amino-N-(1,2-diphenylethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.71, mass found = 417.15.
- 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 387.13.
- 30 5-amino-N-[3-(dibutylamino)propyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 406.24.
- 5-amino-N-butyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 293.12.
- 35

- 116 -

- 5-amino-3-(5-ethylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 411.09.
- 5-amino-3-(5-ethylthien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 331.1.
- 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.75, mass found = 335.17.
- 10 5-amino-3-(5-ethylthien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 309.11.
- 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.92, mass found = 375.2.
- 15 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 307.14.
- 5-amino-3-(5-ethylthien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 345.09.
- 20 5-amino-3-(5-ethylthien-2-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 341.12.
- 25 5-amino-N-(1,3-dimethylbutyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 321.15.
- 5-amino-3-(5-ethylthien-2-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.66, mass found = 369.15.
- 30 5-amino-3-(5-ethylthien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 325.09.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 371.09.
- 35

- 117 -

- 5-amino-3-(5-ethylthien-2-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 307.14.
- 5-amino-N-(cyclohexylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 333.15.
- ethyl 4-([5-amino-3-(5-ethylthien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.36, mass found = 392.15.
- 10 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 375.08.
- 5-amino-N-(1-ethylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 307.14.
- 15 5-amino-3-(5-ethylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 341.12.
- 5-amino-3-(5-ethylthien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 341.12.
- 20 5-amino-N-cyclobutyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 291.1.
- 5-amino-N-benzhydryl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 403.14.
- 25 5-amino-3-(5-ethylthien-2-yl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.86, mass found = 349.18.
- 30 5-amino-N-(2-bromobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.6, mass found = 405.01.
- 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.85, mass found = 349.18.

- 5-amino-3-(5-ethylthien-2-yl)-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.77, mass found = 335.17.
- 5-amino-N-(tert-butyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 293.12.
- 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.36, mass found = 387.13.
- 10 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 353.12.
- 5-amino-3-(5-ethylthien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC
- 15 (Method B) r.t. 1.64, mass found = 367.14.
- 5-amino-N-benzyl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 317.06.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-furylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 307.04.
- 20 5-amino-3-(5-chloro-2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 331.07.
- 5-amino-N-(sec-butyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 283.07.
- 25 5-amino-3-(5-chloro-2-furyl)-N-cyclohexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 309.09.
- 30 5-amino-3-(5-chloro-2-furyl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 347.07.
- 5-amino-3-(5-chloro-2-furyl)-N-hexylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 311.1.
- 35 5-amino-3-(5-chloro-2-furyl)-N-cyclopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 295.07.

- 5-amino-3-(5-chloro-2-furyl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.73, mass found = 339.14.
- 5-amino-3-(5-chloro-2-furyl)-N-isopentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 297.09.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-ethylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.74, mass found = 339.14.
- 10 5-amino-3-(5-chloro-2-furyl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.57, mass found = 359.1.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 323.1.
- 15 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 453.03.
- 5-amino-3-(5-chloro-2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC
- 20 (Method B) r.t. 1.51, mass found = 385.04.
- 5-amino-3-(5-chloro-2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 345.09.
- 5-amino-3-(5-chloro-2-furyl)-N-isobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found =
- 25 283.07.
- 5-amino-3-(5-chloro-2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 385.04.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-methoxybenzyl)isoxazole-
- 30 4-carboxamide. HPLC (Method B) r.t. 1.42, mass found = 347.07.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 335.05.

- 5-amino-3-(5-chloro-2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 385.04.
- 5-amino-3-(5-chloro-2-furyl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 353.04.
- 5-amino-N-(3-chlorobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 351.02.
- 10 5-amino-N-(2-chlorobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 351.02.
- 5-amino-3-(5-chloro-2-furyl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 331.07.
- 15 5-amino-3-(5-chloro-2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.53, mass found = 367.07.
- 5-amino-3-(5-chloro-2-furyl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.31, mass found = 281.06.
- 20 5-amino-3-(5-chloro-2-furyl)-N-cycloheptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 323.1.
- 5-amino-3-(5-chloro-2-furyl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 361.08.
- 25 5-amino-3-(5-chloro-2-furyl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 335.1.
- 30 5-amino-3-(5-chloro-2-furyl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 297.09.
- 5-amino-3-(5-chloro-2-furyl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.63, mass found = 421.12.
- 35

- 5-amino-3-(5-chloro-2-furyl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.64, mass found = 407.1.
- 5-amino-3-(5-chloro-2-furyl)-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 377.08.
- 5-amino-3-(5-chloro-2-furyl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 396.19.
- 10 5-amino-N-butyl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 283.07.
- 5-amino-3-(5-chloro-2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 401.04.
- 15 5-amino-3-(5-chloro-2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 321.05.
- 5-amino-3-(5-chloro-2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.67, mass found = 325.12.
- 20 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.08, mass found = 400.13.
- 25 5-amino-3-(5-chloro-2-furyl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 299.07.
- 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.83, mass found = 365.15.
- 30 5-amino-3-(5-chloro-2-furyl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 297.09.
- 5-amino-3-(5-chloro-2-furyl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 335.05.
- 35

- 5-amino-3-(5-chloro-2-furyl)-N-(4-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 331.07.
- 5-amino-3-(5-chloro-2-furyl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.55, mass found = 311.1.
- 5-amino-3-(5-chloro-2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 359.1.
- 10 5-amino-3-(5-chloro-2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 315.04.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 361.05.
- 15 5-amino-3-(5-chloro-2-furyl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.48, mass found = 297.09.
- 5-amino-3-(5-chloro-2-furyl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 323.1.
- 20 ethyl 4-([5-amino-3-(5-chloro-2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.26, mass found = 382.1.
- 25 5-amino-3-(5-chloro-2-furyl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 365.03.
- 5-amino-3-(5-chloro-2-furyl)-N-(1-ethylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.45, mass found = 297.09.
- 30 5-amino-3-(5-chloro-2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 331.07.
- 5-amino-3-(5-chloro-2-furyl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.46, mass found = 331.07.
- 35

- 5-amino-3-(5-chloro-2-furyl)-N-cyclobutylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 281.06.
- 5-amino-N-benzhydryl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.61, mass found = 393.09.
- 5-amino-3-(5-chloro-2-furyl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.77, mass found = 339.14.
- 10 5-amino-N-(2-bromobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 394.97.
- 5-amino-3-(5-chloro-2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.76, mass found = 339.14.
- 15 5-amino-3-(5-chloro-2-furyl)-N-heptylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.68, mass found = 325.12.
- 5-amino-N-(tert-butyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 283.07.
- 20 5-amino-3-(5-chloro-2-furyl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.26, mass found = 377.08.
- 25 5-amino-3-(5-chloro-2-furyl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.5, mass found = 343.07.
- 5-amino-3-(5-chloro-2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.56, mass found = 357.09.
- 30 5-amino-N-benzyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 324.12.
- 5-amino-N-(2-furylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.12, mass found = 314.1.
- 35

- 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 338.14.
- 5-amino-N-(sec-butyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.17, mass found = 290.14.
- 5-amino-N-cyclohexyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 316.15.
- 10 5-amino-N-(4-methoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 354.13.
- 5-amino-N-hexyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.41, mass found = 318.17.
- 15 5-amino-3-(6-methoxypyridin-3-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 330.08.
- 5-amino-N-cyclopentyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.2, mass found = 302.14.
- 20 5-amino-N-(1,5-dimethylhexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 346.2.
- 5-amino-N-isopentyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.29, mass found = 304.15.
- 25 5-amino-N-(2-ethylhexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.58, mass found = 346.2.
- 30 5-amino-3-(6-methoxypyridin-3-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 366.17.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 330.17.
- 35

- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.54, mass found = 460.1.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 392.11.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 352.15.
- 10 5-amino-N-isobutyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.18, mass found = 290.14.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.39, mass found = 392.11.
- 15 5-amino-N-(2-methoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 354.13.
- 5-amino-N-(2-fluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.25, mass found = 342.11.
- 20 5-amino-3-(6-methoxypyridin-3-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 392.11.
- 25 5-amino-N-(3,4-difluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 360.1.
- 5-amino-N-(3-chlorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.34, mass found = 358.08.
- 30 5-amino-N-(2-chlorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 358.08.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 338.14.
- 35

- 126 -

- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 374.14.
- 5-amino-N-(cyclopropylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.11, mass found = 288.12.
- 5-amino-N-cycloheptyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 330.17.
- 10 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 368.15.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.44, mass found = 342.17.
- 15 5-amino-N-(1,2-dimethylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 304.15.
- 5-amino-N-(3,3-diphenylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 428.18.
- 20 5-amino-N-(1,2-diphenylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.52, mass found = 414.17.
- 25 5-amino-N-(2,4-dimethoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 384.14.
- 5-amino-N-[3-(dibutylamino)propyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.11, mass found = 403.26.
- 30 5-amino-N-butyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 290.14.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 408.1.
- 35

- 127 -

- 5-amino-3-(6-methoxypyridin-3-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.22, mass found = 328.12.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.49, mass found = 332.18.
- 5-amino-N-(1-benzylpiperidin-4-yl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 407.2.
- 10 5-amino-N-(2-methoxy-1-methylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.03, mass found = 306.13.
- 5-amino-N-(4-tert-butylcyclohexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.65, mass found = 372.22.
- 15 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.28, mass found = 304.15.
- 5-amino-N-(3-fluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 342.11.
- 20 5-amino-3-(6-methoxypyridin-3-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.32, mass found = 338.14.
- 25 5-amino-N-(1,3-dimethylbutyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.37, mass found = 318.17.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.43, mass found = 366.17.
- 30 5-amino-3-(6-methoxypyridin-3-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.13, mass found = 322.11.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.21, mass found = 368.11.
- 35

- 128 -

- 5-amino-3-(6-methoxypyridin-3-yl)-N-pentylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 304.15.
- 5-amino-N-(cyclohexylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.4, mass found = 330.17.
- ethyl 4-([5-amino-3-(6-methoxypyridin-3-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate. HPLC (Method B) r.t. 1.11, mass found = 389.17.
- 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.38, mass found = 372.1.
- 5-amino-N-(1-ethylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.27, mass found = 304.15.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 338.14.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.3, mass found = 338.14.
- 5-amino-N-cyclobutyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.12, mass found = 288.12.
- 5-amino-N-benzhydryl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.47, mass found = 400.15.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-octylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.62, mass found = 346.2.
- 5-amino-N-(2-bromobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.35, mass found = 402.03.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.59, mass found = 346.2.

- 5-amino-N-heptyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.51, mass found = 332.18.
- 5-amino-N-(tert-butyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 290.14.
- 5-amino-N-(3,4-dimethoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.13, mass found = 384.14.
- 10 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.33, mass found = 350.14.
- 5-amino-3-(6-methoxypyridin-3-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC
- 15 (Method B) r.t. 1.39, mass found = 364.15.
- 5-amino-N-benzyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.87, mass found = 294.11.
- 5-amino-N-(2-furylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.74, mass found =
- 20 284.09.
- 5-amino-N-(2-phenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.93, mass found = 308.13.
- 5-amino-N-cyclohexyl-3-pyridin-3-ylisoxazole-4-carboxamide.
- 25 HPLC (Method B) r.t. 0.89, mass found = 286.14.
- 5-amino-N-(4-methoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.89, mass found = 324.12.
- 5-amino-N-hexyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC
- 30 (Method B) r.t. 1.03, mass found = 288.16.
- 5-amino-3-pyridin-3-yl-N-(thien-2-ylmethyl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 0.83, mass found = 300.07.
- 5-amino-N-cyclopentyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.79, mass found =
- 35 272.13.

- 130 -

- 5-amino-N-(1,5-dimethylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 316.19.
- 5-amino-N-isopentyl-3-pyridin-3-ylisoxazole-4-carboxamide.
5 HPLC (Method B) r.t. 0.9, mass found = 274.14.
- 5-amino-N-(2-ethylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.21, mass found = 316.19.
- 5-amino-N-(4-phenylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.1, mass found = 336.16.
10 5-amino-N-(2-methylcyclohexyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 300.16.
- 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 430.09.
15
- 5-amino-3-pyridin-3-yl-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.08, mass found = 362.1.
- 5-amino-N-(3-phenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.03, mass found = 322.14.
20
- 5-amino-N-isobutyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.79, mass found = 260.13.
- 5-amino-3-pyridin-3-yl-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.05, mass found = 362.1.
25
- 5-amino-N-(2-methoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.92, mass found = 324.12.
30
- 5-amino-N-(2-fluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.89, mass found = 312.1.
- 5-amino-3-pyridin-3-yl-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.1, mass found = 362.1.
35

- 131 -

- 5-amino-N-(3,4-difluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 330.09.
- 5-amino-N-(3-chlorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1, mass found = 328.07.
- 5-amino-N-(2-chlorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 328.07.
- 5-amino-N-(3-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 308.13.
- 5-amino-N-(1-naphthylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.04, mass found = 344.13.
- 5-amino-N-(cyclopropylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.72, mass found = 258.11.
- 5-amino-N-cycloheptyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.98, mass found = 300.16.
- 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.94, mass found = 338.14.
- 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.07, mass found = 312.16.
- 5-amino-N-(1,2-dimethylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.86, mass found = 274.14.
- 5-amino-N-(3,3-diphenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.23, mass found = 398.17.
- 5-amino-N-(1,2-diphenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.19, mass found = 384.16.

- 132 -

- 5-amino-N-(2,4-dimethoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.95, mass found = 354.13.
- 5-amino-N-[3-(dibutylamino)propyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.86, mass found = 373.25.
- 5-amino-N-butyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.79, mass found = 260.13.
- 5-amino-3-pyridin-3-yl-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.13, mass found = 378.09.
- 5-amino-N-[(5-methyl-2-furyl)methyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.84, mass found = 298.11.
- 5-amino-N-(1-methylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.1, mass found = 302.17.
- 5-amino-N-(1-methylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.87, mass found = 274.14.
- 5-amino-N-(3-fluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.92, mass found = 312.1.
- 5-amino-N-(4-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 308.13.
- 5-amino-N-(1,3-dimethylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.97, mass found = 288.16.
- 5-amino-N-(1-methyl-3-phenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.09, mass found = 336.16.
- 5-amino-N-[3-(methylthio)propyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.79, mass found = 292.1.
- 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.9, mass found = 338.1.

- 133 -

- 5-amino-N-pentyl-3-pyridin-3-ylisoxazole-4-carboxamide.
HPLC (Method B) r.t. 0.92, mass found = 274.14.
- 5-amino-N-(cyclohexylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.02, mass found =
- 5 300.16.
- ethyl 4-{[(5-amino-3-pyridin-3-ylisoxazol-4-yl)carbonyl]amino}piperidine-1-carboxylate. HPLC (Method B) r.t. 0.84, mass found = 359.16.
- 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.05, mass
- 10 found = 342.09.
- 5-amino-N-(1-ethylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.86, mass found = 274.14.
- 15 5-amino-N-(1-phenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.94, mass found = 308.13.
- 5-amino-N-(2-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.95, mass found =
- 20 308.13.
- 5-amino-N-cyclobutyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.72, mass found = 258.11.
- 5-amino-N-benzhydryl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.15, mass found = 370.14.
- 25 5-amino-N-octyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.24, mass found = 316.19.
- 5-amino-N-(2-bromobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.99, mass found = 372.02.
- 30 5-amino-N-(1-methylheptyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.21, mass found = 316.19.
- 5-amino-N-heptyl-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 1.14, mass found = 302.17.
- 35 5-amino-N-(tert-butyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.8, mass found = 260.13.

- 134 -

5-amino-N-(3,4-dimethoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.84, mass found = 354.13.

5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-pyridin-3-ylisoxazole-4-carboxamide. HPLC (Method B) r.t. 0.98, mass found = 320.13.

5-amino-3-pyridin-3-yl-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide. HPLC (Method B) r.t. 1.04, mass found = 334.14.

10

Example 4

4-(3-Methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin

4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin
15 [copoly(styrene-1%dvb) 100-200 mesh] (1,5 g, 1 eq, loading 0,94 mmol/g) is swollen in DCM and then filtered. A mixture of THF/DCM (4 : 1, 15 ml), aniline (6 eq.) and AcOH (6 eq.) were added. After 15 min NaBH(OAc)₃ (3 eq) was added and the reaction was shaken over night at room temperature.
20 After filtration, the resin was washed with methanol (x 3), DMF/DCM (1 : 1) (x 3) and DCM (x 5).

Example 5

4-(4-{[(2-Cyano-acetyl)-phenyl-amino]-methyl}-3-methoxy-phenoxy)-butyryl AM resin

Diisopropyl carbodimide (1.1 ml, 7.08 mmol) was added to a suspension of cyanoacetic acid (1.34 g, 15.79 mmol) in DCM (18.9 ml) at 0 C°, the resulting solution was stirred for 30 min, then was added to 1.4 g 4-(3-Methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin (1.32 mmol).
30 The suspension was shaken for 16 hours, then filtered and washed with dichloromethane (2x), dimethylformamide (2x) and dichloromethane (3x). A second cycle with the same procedure was performed for 16 h, then the solution was
35 drained and the resin washed with dichloromethane (2x),

- 135 -

dimethylformamide (2x), methanol (2x), dichloromethane (2x), dimethylformamide (2x), dichloromethane (5x) and dried in vacuum overnight.

5

Example 6

5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid 4-(3-methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin amide

4-(4-{[(2-Cyano-acetyl)-phenyl-amino]-methyl}-3-methoxy-phenoxy)-butyryl AM resin (0.75 g, 0.443 mmol) and
10 pyridine-2-hydroxyaminomethyl chloride (208 mg, 1.328 mmol) were suspended in dry THF (8 ml) and cooled to -20°C. Then lithium bis(trimethylsilyl)amide, LiHMDS, (2.2 ml of 1 M solution in THF, 2.2 mmol) was added, the resulting suspension was stirred at -20°C for 15 min, allowed to warm
15 to room temperature (23°C) and shaken for 2h and drained. The resin was washed with dimethylformamide, dichloromethane, dimethylformamide (2x) and dichloromethane (3x) and dried under vacuum.

By working in an analogous way starting from cyanoacetic
20 acid Rink amide the following compounds were prepared:

5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid Rink amide.

5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid Rink amide.

25 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid Rink amide.

5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid Rink amide).

30 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid Rink amide).

5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid Rink amide.

5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid Rink amide.

5-Amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid Rink amide.

Example 7

5 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
phenylamide

5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid 4-(3-methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin amide (0.75 mmol) was treated with a solution of TFA 20% in
10 anhydrous dichloromethane (10 ml). The resulting suspension was gently stirred or shaken at 22°C for 1h, then washed with anhydrous dichloromethane, methanol, dichloromethane and dried under vacuum to give a crude solid, that after purification by flash chromatography gave 120 mg of the
15 desired compound.

Analogously, by using the appropriate chloroxime derivative and the appropriate cyanacetamide derivative supported on resin, the following compounds were prepared after
20 purification by flash chromatography or preparative HPLC, when necessary:

- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
25 isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide.
5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide.
30 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide.
5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.

- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 5 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 10 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid phenylamide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
- 15 phenylamide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 20 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid phenylamide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
- 25 isoxazole-4-carboxylic acid (3-phenoxy-phenyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 4-{[5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carbonyl]-amino}-piperidine-1-carboxylic acid
- 30 ethyl ester.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (tetrahydro-furan-2-ylmethyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
- 35 phenylamide.

- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid 2-methoxy-benzylamide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid 4-chloro-benzylamide.
- 5 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid 2-methoxy-benzylamide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 4-fluoro-benzylamide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3,4-
- 10 dimethyl-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (2-dimethylamino-ethyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
- 15 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
- 20 diethylamino-phenyl)-amide.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide.
- 25 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (1-ethyl-propyl)-amide.
- 30 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3,4-dimethyl-phenyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-fluoro-2-methyl-phenyl)-amide.
- 35 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid [4-(acetyl-methyl-amino)-phenyl]-amide.

- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 2-methoxy-benzylamide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide.
- 5 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid 4-chloro-benzylamide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
- 10 isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-phenoxy-phenyl)-amide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
- 15 (pyridin-4-ylmethyl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide.
- 5-Amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid phenylamide.
- 20 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-methylsulfanyl-phenyl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3-chloro-4-methyl-phenyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-
- 25 chloro-4-methyl-phenyl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
- 30 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide.
- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
- 35 isoxazole-4-carboxylic acid cyclopropylamide.

- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide.
- 4-[(5-Amino-3-thiophen-3-yl-isoxazole-4-carbonyl)-amino]-piperidine-1-carboxylic acid ethyl ester.
- 5 4-[(5-Amino-3-pyridin-2-yl-isoxazole-4-carbonyl)-amino]-piperidine-1-carboxylic acid ethyl ester.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide.
- 10 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (furan-2-ylmethyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (furan-2-ylmethyl)-amide.
- 15 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (furan-2-ylmethyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide.
- 20 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide.
- 25 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide.
- 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid 4-fluoro-benzylamide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide.
- 30 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-difluoromethoxy-phenyl)-amide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid [4-(toluene-4-sulfonylamino)-phenyl]-amide.
- 35 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2-methylsulfanyl-phenyl)-amide.

- 141 -

- 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid phenylamide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (1-ethyl-propyl)-amide.
- 5 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid [4-(toluene-4-sulfonylamino)-phenyl]-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide.
- 10 5-Amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (3-chloro-4-methyl-phenyl)-amide.
- 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide.
- 15 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-dimethylamino-ethyl)-amide.
- 5-Amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide.
- 20 5-Amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide.
- 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide.
- 5-Amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide.
- 25 5-Amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide.

Example 8

30 5-Benzoylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide

To a suspension of 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide (25 mg, 0.123 mmol) in dichloromethane (0.25 ml), triethyl amine (0.051 ml, 0.368 mmol), 4-dimethyl aminopyridine (3 mg, 0.025 mmol) and

35

- 142 -

benzoyl chloride (0.021 ml, 0.184 mmol) were added. The final mixture was stirred at room temperature (23°C) for 16 hours, afterward volatiles were removed under reduced pressure to give a crude solid, that was purified by flash chromatography (yield 62%). HPLC r.t. 3.24, [M+H]⁺ = 309. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm): 11,5 (bs, 1H), 8,7 (d, 2H), 8 (m, 2H) 7,7(m, 2H), 7,6 (m, 1H), 7,52(bt, 2H).

By working in an analogous way and by using the appropriate isoxazole derivative and the appropriate acyl chloride derivative, the following compounds were prepared after purification by flash chromatography or preparative HPLC when necessary:

5-Benzoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.52, [M+H]⁺ = 314. ¹H-NMR (DMSO-d₆), diagnostic signals (ppm): 9,05 (bs, 1H), 8,2 (bs, 2H), 8 (d, 2H) 7,6 (m, 2H), 7,5 (m, 2H), 7,22(m, 2H).

5-(4-Methyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.85, [M+H]⁺ = 323.

5-(4-Chloro-butyrylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.64, [M+H]⁺ = 309.

5-(4-tert-Butyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 7.08, [M+H]⁺ = 365.

5-Pentanoylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.95, [M+H]⁺ = 289.

5-(4-Methyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.81, [M+H]⁺ = 323.

5-(4-Chloro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.84, [M+H]⁺ = 343.

5-Pentanoylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.96, [M+H]⁺ = 289.

5-(3-Phenyl-propionylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.74, [M+H]⁺ = 337.

5-(4-tert-Butyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.34, [M+H]⁺ = 370.

- 143 -

- 5-Pentanoylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.18, [M+H]⁺=294.
- 5-(3-Phenyl-propionylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.89, [M+H]⁺=342.
- 5 5-(4-Methyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.07, [M+H]⁺=328.
- 5-(4-tert-Butyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.33, [M+H]⁺=370.
- 5-Pentanoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.12, [M+H]⁺=294.
- 10 5-(3-Phenyl-propionylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.83, [M+H]⁺=342.
- 5-(4-Chloro-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.83, [M+H]⁺=315.
- 15 5-(4-tert-Butyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 7.25, [M+H]⁺=371.
- 5-(4-Chloro-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.3, [M+H]⁺=349.
- 5-(3-Phenyl-propionylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.69, [M+H]⁺=343.
- 20 5-(4-Methyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.76, [M+H]⁺=323.
- 5-(4-tert-Butyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.04, [M+H]⁺=365.
- 25 5-(4-Chloro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.81, [M+H]⁺=343.
- 5-Pentanoylamino-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.92, [M+H]⁺=289.
- 5-(4-tert-Butyl-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.79, [M+H]⁺=458.
- 30 5-(4-Chloro-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.9, [M+H]⁺=436.
- 35 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-pentanoylamino-isoxazole-4-carboxylic acid amide. HPLC r.t. 5, [M+H]⁺=382.

- 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl) -5- (3-phenyl-propionylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.5, [M+H]⁺=430.
- 5- (4-Chloro-butyrylamino) -3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.14, [M+H]⁺=359.
- 5-Pentanoylamino-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.47, [M+H]⁺=339.
- 5- (3-Phenyl-propionylamino) -3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.84, [M+H]⁺=387.
- 10 5- (3-Fluoro-benzoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.65, [M+H]⁺=327.
- 5- (3-Methoxy-benzoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.59, [M+H]⁺=339.
- 15 3-Pyridin-2-yl-5- (4-trifluoromethyl-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.63, [M+H]⁺=377.
- 5- (3-Fluoro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.33, [M+H]⁺=327.
- 5- (2-Methoxy-acetylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 1.76, [M+H]⁺=277.
- 20 5- (3-Methoxy-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.49, [M+H]⁺=339.
- 3-Pyridin-4-yl-5- (4-trifluoromethyl-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 3.4, [M+H]⁺=377.
- 25 5- (3-Bromo-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.9, [M+H]⁺=387.
- 5- (3-Fluoro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.64, [M+H]⁺=332.
- 30 5- (2-Methoxy-acetylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.85, [M+H]⁺=282.
- 5-Benzoylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.52, [M+H]⁺=314.
- 5- (3-Methoxy-benzoylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.77, [M+H]⁺=344.
- 35

- 3-Thiophen-2-yl-5-(4-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 4.64,
[M+H]⁺=382.
- 5-(2-Methoxy-acetylamino)-3-thiophen-3-yl-isoxazole-4-
5 carboxylic acid amide. HPLC r.t. 2.79, [M+H]⁺=282.
- 5-(3-Methoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.78, [M+H]⁺=344.
- 3-Thiophen-3-yl-5-(4-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 3.7,
10 [M+H]⁺=382.
- 5-(2-Methoxy-acetylamino)-3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.56, [M+H]⁺=283.
- 5-Benzoylamino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
amide. HPLC r.t. 5.52, [M+H]⁺=315.
- 15 5-(3-Methoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.76, [M+H]⁺=345.
- 3-Thiazol-2-yl-5-(4-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.77,
[M+H]⁺=383.
- 20 5-(3-Bromo-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 6.5, [M+H]⁺=393.
- 5-(2-Methoxy-acetylamino)-3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 1.71, [M+H]⁺=277.
- 5-(3-Methoxy-benzoylamino)-3-pyridin-3-yl-isoxazole-4-
25 carboxylic acid amide. HPLC r.t. 3.47, [M+H]⁺=339.
- 3-Pyridin-3-yl-5-(4-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 3.36,
[M+H]⁺=377.
- 5-(3-Bromo-benzoylamino)-3-pyridin-3-yl-isoxazole-4-
30 carboxylic acid amide. HPLC r.t. 3.85, [M+H]⁺=387.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(3-fluoro-
benzoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t.
5.5, [M+H]⁺=420.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-methoxy-
35 acetylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t.
3.88, [M+H]⁺=370.

- 5- (3-Bromo-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.92, [M+H]⁺=480.
- 3-Quinolin-2-yl-5- (4-trifluoromethyl-benzoylamino) -
5 isoxazole-4-carboxylic acid amide. HPLC r.t. 6.93, [M+H]⁺=427.
- 5- (Cyclopropanecarbonyl-amino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.89, [M+H]⁺=273.
- 5- (2-Phenoxy-acetylamino) -3-pyridin-4-yl-isoxazole-4-
10 carboxylic acid amide. HPLC r.t. 3.6, [M+H]⁺=339.
- 5- (2-Chloro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.19, [M+H]⁺=343.
- 5- (Cyclopropanecarbonyl-amino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2, [M+H]⁺=273.
- 15 5- (3-Methyl-but-2-enoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.81, [M+H]⁺=287.
- 5- (2,4-Difluoro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.51, [M+H]⁺=345.
- 5- [2- (3-Methoxy-phenyl) -acetylamino] -3-thiophen-2-yl-
20 isoxazole-4-carboxylic acid amide. HPLC r.t. 4.53, [M+H]⁺=358.
- 5- (2-Phenoxy-acetylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.87, [M+H]⁺=344.
- 5- (2-Chloro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-
25 carboxylic acid amide. HPLC r.t. 4.51, [M+H]⁺=348.
- 5- (Cyclopropanecarbonyl-amino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.09, [M+H]⁺=278.
- 5- (3-Methyl-but-2-enoylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.02, [M+H]⁺=292.
- 30 5- (2,4-Difluoro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.85, [M+H]⁺=350.
- 5- [2- (3-Methoxy-phenyl) -acetylamino] -3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.49, [M+H]⁺=358.
- 35 5- (2-Chloro-benzoylamino) -3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.53, [M+H]⁺=348.

- 5- (Cyclopropanecarbonyl-amino) -3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.03, [M+H]⁺=278.
- 5- (3-Methyl-but-2-enoylamino) -3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.94, [M+H]⁺=292.
- 5 5- (2,4-Difluoro-benzoylamino) -3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.87, [M+H]⁺=350.
- 5- (2-Phenoxy-acetylamino) -3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.52, [M+H]⁺=345.
- 5- (Cyclopropanecarbonyl-amino) -3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.1, [M+H]⁺=277.
- 10 5- (3-Methyl-but-2-enoylamino) -3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.95, [M+H]⁺=293.
- 5- (2,4-Difluoro-benzoylamino) -3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.62, [M+H]⁺=351.
- 15 5- (2-Phenoxy-acetylamino) -3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.56, [M+H]⁺=339.
- 5- (2-Chloro-benzoylamino) -3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.16, [M+H]⁺=343.
- 5- (Cyclopropanecarbonyl-amino) -3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 1.95, [M+H]⁺=273.
- 20 5- (2,4-Difluoro-benzoylamino) -3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.46, [M+H]⁺=345.
- 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl) -5- (2-phenoxy-acetylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.45, [M+H]⁺=432.
- 25 5- (2-Chloro-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.32, [M+H]⁺=436.
- 5- (Cyclopropanecarbonyl-amino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 4.12, [M+H]⁺=366.
- 30 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl) -5- (3-methyl-but-2-enoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 4.81, [M+H]⁺=380.

- 5- (2,4-Difluoro-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.46, [M+H]⁺=438.
- 5- (Cyclopropanecarbonyl-amino) -3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.6, [M+H]⁺=323.
- 5- (4-Fluoro-benzoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.56, [M+H]⁺=327.
- 3-Pyridin-2-yl-5- (4-trifluoromethoxy-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 6.52, [M+H]⁺=393.
- 5- (2-Methyl-benzoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.61, [M+H]⁺=323.
- 5- (4-Ethyl-benzoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.38, [M+H]⁺=337.
- 5- [(Naphthalene-1-carbonyl)-amino] -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.27, [M+H]⁺=359.
- 5- (4-Fluoro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.38, [M+H]⁺=327.
- 3-Pyridin-4-yl-5- (4-trifluoromethoxy-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 4.3, [M+H]⁺=393.
- 5- (2-Methyl-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.5, [M+H]⁺=323.
- 3-Pyridin-4-yl-5- (3-trifluoromethyl-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 3.97, [M+H]⁺=377.
- 5- (4-Ethyl-benzoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.32, [M+H]⁺=337.
- 5- [(Naphthalene-1-carbonyl)-amino] -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.2, [M+H]⁺=359.
- 5- (4-Fluoro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.66, [M+H]⁺=332.

- 3-Thiophen-2-yl-5-(4-trifluoromethoxy-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.53,
[M+H]⁺=398.
- 5 5-(2-Methyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.79, [M+H]⁺=328.
- 3-Thiophen-2-yl-5-(3-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.21,
[M+H]⁺=382.
- 10 5-(4-Ethyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.64, [M+H]⁺=342.
- 5-[(Naphthalene-1-carbonyl)-amino]-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.44,
[M+H]⁺=364.
- 15 5-(4-Fluoro-benzoylamino)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.66, [M+H]⁺=332.
- 3-Thiophen-3-yl-5-(4-trifluoromethoxy-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.59,
[M+H]⁺=398.
- 20 5-(2-Methyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.77, [M+H]⁺=328.
- 3-Thiophen-3-yl-5-(3-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.29,
[M+H]⁺=382.
- 25 5-(4-Ethyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.58, [M+H]⁺=342.
- 5-[(Naphthalene-1-carbonyl)-amino]-3-thiophen-3-yl-
isoxazole-4-carboxylic acid amide. HPLC r.t. 5.43,
[M+H]⁺=364.
- 30 5-(4-Fluoro-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.72, [M+H]⁺=333.
- 3-Thiazol-2-yl-5-(4-trifluoromethoxy-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 6.66,
[M+H]⁺=398.
- 35 5-(2-Methyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.8, [M+H]⁺=329.

- 150 -

- 3-Thiazol-2-yl-5-(3-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 6.47,
[M+H]⁺=383.
- 5-(4-Ethyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
5 carboxylic acid amide. HPLC r.t. 6.6, [M+H]⁺=343.
- 5-(4-Fluoro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.34, [M+H]⁺=327.
- 3-Pyridin-3-yl-5-(4-trifluoromethoxy-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 4.27,
10 [M+H]⁺=393.
- 5-(2-Methyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.45, [M+H]⁺=323.
- 3-Pyridin-3-yl-5-(3-trifluoromethyl-benzoylamino)-
isoxazole-4-carboxylic acid amide. HPLC r.t. 3.95,
15 [M+H]⁺=377.
- 5-(4-Ethyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.33, [M+H]⁺=337.
- 5-[(Naphthalene-1-carbonyl)-amino]-3-pyridin-3-yl-
isoxazole-4-carboxylic acid amide. HPLC r.t. 4.2,
20 [M+H]⁺=359.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(4-fluoro-
benzoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t.
5.4, [M+H]⁺=420.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-methyl-
25 benzoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t.
5.51, [M+H]⁺=416.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(4-ethyl-
benzoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t.
6.21, [M+H]⁺=430.
- 30 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-[(naphthalene-
1-carbonyl)-amino]-isoxazole-4-carboxylic acid amide. HPLC
r.t. 6.04, [M+H]⁺=452.
- 5-(2-Fluoro-benzoylamino)-3-pyridin-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.22, [M+H]⁺=327.
- 35 5-(3-Methyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.83, [M+H]⁺=323.

- 5- [2- (4-Fluoro-phenyl) -acetyl-amino] -3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide. HPLC r.t. 3.48,
[M+H]⁺=341.
- 5- (3,5-Difluoro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-
5 carboxylic acid amide. HPLC r.t. 3.32, [M+H]⁺=345.
- 5- (3-Chloro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.74, [M+H]⁺=343.
- 5- (2-Fluoro-benzoylamino) -3-pyridin-4-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.36, [M+H]⁺=327.
- 10 5- (3-Methyl-benzoylamino) -3-pyridin-4-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.78, [M+H]⁺=323.
- 5- (3,5-Difluoro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.49, [M+H]⁺=350.
- 5- (3-Chloro-benzoylamino) -3-thiophen-2-yl-isoxazole-4-
15 carboxylic acid amide. HPLC r.t. 5.05, [M+H]⁺=348.
- 5- (3,5-Difluoro-benzoylamino) -3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 4.58, [M+H]⁺=350.
- 5- (3-Methyl-benzoylamino) -3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.05, [M+H]⁺=328.
- 20 5- (3-Chloro-benzoylamino) -3-thiazol-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 6.33, [M+H]⁺=349.
- 5- (3-Chloro-benzoylamino) -3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.71, [M+H]⁺=343.
- 5- (2-Fluoro-benzoylamino) -3-pyridin-3-yl-isoxazole-4-
25 carboxylic acid amide. HPLC r.t. 3.3, [M+H]⁺=327.
- 5- (3-Methyl-benzoylamino) -3-pyridin-3-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 3.73, [M+H]⁺=323.
- 5- (3-Chloro-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-
pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t.
30 5.78, [M+H]⁺=436.
- 5-Butyrylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
amide. HPLC r.t. 4.22, [M+H]⁺=275.
- 5- [(E) - (3-Phenyl-acryloyl) amino] -3-pyridin-2-yl-isoxazole-
4-carboxylic acid amide. HPLC r.t. 5.74, [M+H]⁺=335.
- 35 5- (3,3-Dimethyl-butyrylamino) -3-pyridin-2-yl-isoxazole-4-
carboxylic acid amide. HPLC r.t. 5.2, [M+H]⁺=303.

- 5- (2-Chloro-2-phenyl-acetyl-amino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.37, [M+H]⁺=357.
- 5-Butyrylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.23, [M+H]⁺=275.
- 5 5- [(Biphenyl-4-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.94, [M+H]⁺=385.
- 5- (4-Chloromethyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.91, [M+H]⁺=355.
- 3-Pyridin-4-yl-5- (3,4,5-trimethoxy-benzoylamino)-isoxazole-10 4-carboxylic acid amide. HPLC r.t. 3.28, [M+H]⁺=399.
- 5- [(E)- (3-Phenyl-acryloyl) amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.91, [M+H]⁺=335.
- 5- (3,3-Dimethyl-butyrylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.25, [M+H]⁺=303.
- 15 5- (2-Chloro-2-phenyl-acetyl-amino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.6, [M+H]⁺=362.
- 5- (4-Nitro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.23, [M+H]⁺=359.
- 20 5-Butyrylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.38, [M+H]⁺=280.
- 5- [(Biphenyl-4-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.19, [M+H]⁺=390.
- 5- (4-Chloromethyl-benzoylamino)-3-thiophen-2-yl-isoxazole-25 4-carboxylic acid amide. HPLC r.t. 5.17, [M+H]⁺=362.
- 3-Thiophen-2-yl-5- (3,4,5-trimethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.47, [M+H]⁺=404.
- 5- [(E)- (3-Phenyl-acryloyl) amino]-3-thiophen-2-yl-isoxazole-30 4-carboxylic acid amide. HPLC r.t. 5.06, [M+H]⁺=340.
- 5- (3,3-Dimethyl-butyrylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.5, [M+H]⁺=308.
- 5- (2-Chloro-2-phenyl-acetyl-amino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.77, 35 [M+H]⁺=362.

- 5-Butyrylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.35, [M+H]⁺=380.
- 5-(3,3-Dimethyl-butyrylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.44, [M+H]⁺=308.
- 5 5-(4-Chloromethyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.13, [M+H]⁺=363.
- 5-(3,3-Dimethyl-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.41, [M+H]⁺=309.
- 5-(3,3-Dimethyl-butyrylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.23, [M+H]⁺=303.
- 10 5-Butyrylamino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.42, [M+H]⁺=368.
- 3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-[(E)-(3-phenyl-acryloyl)amino]-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.67, [M+H]⁺=428.
- 15 5-(3,3-Dimethyl-butyrylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.22, [M+H]⁺=396.
- 20 5-Butyrylamino-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.89, [M+H]⁺=325.
- 5-(3,3-Dimethyl-butyrylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.72, [M+H]⁺=353.
- 5-[(Furan-2-carbonyl)-amino]-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.45, [M+H]⁺=299.
- 25 5-[(Naphthalene-2-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.41, [M+H]⁺=359.
- 5-(4-Methoxy-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.51, [M+H]⁺=339.
- 30 5-[(Furan-2-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.42, [M+H]⁺=299.
- 5-(3-Cyclopentyl-propionylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.32, [M+H]⁺=329.

- 5- [(Naphthalene-2-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.71, [M+H]⁺=364.
- 5- (4-Methoxy-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.75, [M+H]⁺=344.
- 5- (2-Benzyloxy-acetylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.06, [M+H]⁺=358.
- 5- (2,4-Dimethoxy-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.16, [M+H]⁺=374.
- 10 5- [(Furan-2-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.67, [M+H]⁺=304.
- 5- (3-Cyclopentyl-propionylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.5, [M+H]⁺=334.
- 5- (4-Bromo-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.34, [M+H]⁺=392.
- 15 5- [(Naphthalene-2-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.71, [M+H]⁺=364.
- 5- (4-Methoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.72, [M+H]⁺=344.
- 20 5- (2,4-Dimethoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.02, [M+H]⁺=374.
- 5- [(Furan-2-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.69, [M+H]⁺=304.
- 25 5- (3-Cyclopentyl-propionylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.45, [M+H]⁺=334.
- 5- (4-Methoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.63, [M+H]⁺=345.
- 5- (2,4-Dimethoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.66, [M+H]⁺=375.
- 30 5- [(Furan-2-carbonyl)-amino]-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.61, [M+H]⁺=305.
- 5- (3-Cyclopentyl-propionylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.36, [M+H]⁺=335.
- 35 5- [(Furan-2-carbonyl)-amino]-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.39, [M+H]⁺=299.

- 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl) -5- (4-methoxy-benzoylamino) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.4, [M+H]⁺=432.
- 5- (2,4-Dimethoxy-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.5, [M+H]⁺=462.
- 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl) -5- [(furan-2-carbonyl) -amino] -isoxazole-4-carboxylic acid amide. HPLC r.t. 4.58, [M+H]⁺=392.
- 10 5- (3-Cyclopentyl-propionylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 6.05, [M+H]⁺=422.
- 5- (3,4-Dimethoxy-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.02, [M+H]⁺=462.
- 5- (4-Bromo-benzoylamino) -3- (3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl) -isoxazole-4-carboxylic acid amide. HPLC r.t. 5.99, [M+H]⁺=480.
- 5- (3-Cyclopentyl-propionylamino) -3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 7.56, [M+H]⁺=379.
- 20 5- (2-Ethyl-butyrylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.23, [M+H]⁺=303.
- 5- (2-Propyl-pentanoylamino) -3-pyridin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.35, [M+H]⁺=331.
- 25 5- (3-Methyl-butyrylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 2.79, [M+H]⁺=289.
- 5- [(Benzo[1,3]dioxole-5-carbonyl) -amino] -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.38, [M+H]⁺=353.
- 30 5- (2-Ethyl-butyrylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.17, [M+H]⁺=303.
- 5- (2-Propyl-pentanoylamino) -3-pyridin-4-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.42, [M+H]⁺=331.
- 5- (3-Methyl-butyrylamino) -3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.98, [M+H]⁺=294.
- 35

- 5-[(Benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.63, [M+H]⁺=358.
- 5- (2-Ethyl-butyrylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.42, [M+H]⁺=308.
- 5- (2-Propyl-pentanoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.65, [M+H]⁺=336.
- 5- [2-(3,4-Dimethoxy-phenyl)-acetylamino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.08, [M+H]⁺=388.
- 5- (3-Methyl-butyrylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.93, [M+H]⁺=294.
- 5-[(Benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.61, [M+H]⁺=358.
- 5- (2-Ethyl-butyrylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.38, [M+H]⁺=308.
- 5- (2-Propyl-pentanoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.6, [M+H]⁺=336.
- 5- (3-Methyl-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.01, [M+H]⁺=295.
- 5-[(Benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.56, [M+H]⁺=359.
- 5- (2-Ethyl-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.42, [M+H]⁺=309.
- 5- (2-Propyl-pentanoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.53, [M+H]⁺=337.
- 5- (2-Ethyl-butyrylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 3.14, [M+H]⁺=303.
- 5- (2-Propyl-pentanoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.38, [M+H]⁺=331.
- 3- (3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5- (3-methyl-butyrylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t. 4.85, [M+H]⁺=382.

5-[(Benzo[1,3]dioxole-5-carbonyl)-amino]-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.3, [M+H]⁺=446.

3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-ethyl-butylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t. 5.21, [M+H]⁺=396.

3-(3,5-Dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-propyl-pentanoylamino)-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.16, [M+H]⁺=424.

5-(3-Methyl-butylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.36, [M+H]⁺=339.

5-(2-Ethyl-butylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide. HPLC r.t. 6.74, [M+H]⁺=353.

15

Example 9

5-Acetylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
4-(3-methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin
amide

To a suspension of 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 4-(3-methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin amide (160 mg, 0.094 mmol) in dichloromethane (1.6 ml), triethyl amine (0.040 ml, 0.285 mmol), 4-dimethyl aminopyridine (5.8 mg, 0.048 mmol) and acetyl chloride (0.142 mmol) were added. The final mixture was shaken at room temperature (23°C) for 16 hours, afterward the resin was filtered, washed with N,N-dimethylformamide, dichloromethane, methanol, and dichloromethane (3x) dried under vacuum.

30

Example 10

5-Acetylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(4-methoxy-phenyl)-amide.

5-Acetylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 4-(3-methoxy-4-phenylaminomethyl-phenoxy)-butyryl AM resin amide (0.75 mmol) was treated with a solution of TFA 20% in

- 158 -

anhydrous dichloromethane (10 ml). The resulting suspension was gently stirred at 22°C for 1h, then washed with anhydrous dichloromethane, methanol, dichloromethane and dried under vacuum to give a crude solid, that after
5 purification by flash chromatography gave 130 mg of the desired compound. $[M+H]^+ = 358$. ^1H -NMR (DMSO- d_6), diagnostic signals (ppm): 11,4 (bs, 1H), 10,1 (bs, 1H), 7,7 (d, 1H), 7,6 (d, 1H) 7,5 (d, 2H), 7,2 (dd, 1H), 6,9 (d, 2H), 3,7 (s, 3H), 2,1 (s, 3H).

10

By working in an analogous way and by using the appropriate isoxazole resin bound derivatives and the appropriate acyl chloride the following compound has been prepared, after purification by flash chromatography or preparative HPLC
15 when necessary:

5-Phenylacetyl-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid butylamide. $[M+H]^+ = 384$. ^1H -NMR (DMSO- d_6), diagnostic signals (ppm): 11,4 (bs, 1H), 8,1 (bs, 1H), 7,7 (d, 1H) 7,6 (d, 1H), 7,3 (m, 5H), 7,1 (d, 1H), 3,7 (s, 2H), 3,1 (m, 2H)
20 1,4 (m, 2H), 1,2 (m, 2H), 0,9 (m, 3H).

Example 11

5-(3-Phenyl-ureido)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide

25 To a suspension of 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide (25 mg, 0.12 mmol) in dichloromethane (0.25 ml), phenyl isocyanate (0.130 ml, 1.2 mmol) was added. The final mixture was stirred at room temperature (23°C) for 16 hours, afterward volatiles were removed under
30 reduced pressure to give a crude solid, that was purified by flash chromatography (45% yield). $[M+H]^+ = 329$, $[M-H]^- = 327$. HPLC r.t. 4.6.

- 159 -

Example 12

5-(4-Acetylamino-benzenesulfonylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid Rink amide

To a suspension of 5-Amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid Rink amide (50 mg, 0.030 mmol) in dichloromethane (0.5 ml), diisopropylethyl amine (15.2 µl, 0.089 mmol), 4-dimethyl aminopyridine (1.8 mg, 0.015 mmol) and N-acetylbenzenesulfonyl chloride (10.3 mg, 0.044 mmol) were added. The final mixture was shaken at room temperature (23°C) for 16 hours, afterward the resin was filtered, washed with N,N-dimethylformamide, dichloromethane, methanol, dichloromethane (3x) and dried under vacuum.

Example 13


5-(4-Acetylamino-benzenesulfonylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide

5-(4-Acetylamino-benzenesulfonylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid Rink amide (0.75 mmol) was treated with a solution of TFA 20% in anhydrous dichloromethane (10 ml). The resulting suspension was gently stirred or shaken at 22°C for 1h, then washed with anhydrous dichloromethane, methanol, dichloromethane and dried under vacuum to give a crude solid, that after purification by flash chromatography gave 154 mg of the desired compound. [M+H]⁺ = 402. HPLC r.t. 3.0.

- 160 -

CLAIMS

1. A method for treating diseases caused by and/or associated with an altered protein kinase activity which comprises administering to a mammal in need thereof an effective amount of an aminoisoxazole derivative represented by formula (I):



wherein

10 R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms
15 selected among nitrogen, oxygen or sulfur;

X is a divalent group selected from -N(R₃)- or -O-;

Y is a divalent group selected from -CH(R₃)-, -CO-, -CONH- or -SO₂-, or Y may also be a single bond when R₂ is a hydrogen atom or a C₃-C₆ cycloalkyl group;

20 R₁ is a hydrogen atom or a group, optionally further substituted, selected from straight or branched C₁-C₆ alkyl, C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur; the
25 said cycloalkyl, aryl or heterocyclyl groups being optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

- 161 -

R₂ and R₃ have, each independently, the meanings above reported for R₁ or represent an optionally substituted straight or branched C₂-C₆ alkenyl or alkynyl group; and the pharmaceutically acceptable salts thereof.

5

2. The method of claim 1 wherein the disease caused by and/or associated with an altered protein kinase activity is a cell proliferative disorder selected from the group consisting of cancer, Alzheimer's disease, viral
10 infections, auto-immune diseases and neurodegenerative disorders.

3. The method of claim 2 wherein the cancer is selected from carcinoma, squamous cell carcinoma, hematopoietic
15 tumors of lymphoid or myeloid lineage, tumors of mesenchymal origin, tumors of the central and peripheral nervous system, melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoxanthoma, thyroid follicular cancer and Kaposi's sarcoma.

20


4. The method of claim 1 wherein the cell proliferative disorder is selected from benign prostate hyperplasia, familial adenomatosis, polyposis, neuro-fibromatosis, psoriasis, vascular smooth cell proliferation associated
25 with atherosclerosis, pulmonary fibrosis, arthritis glomerulonephritis and post-surgical stenosis and restenosis.

5. The method of claim 1 which provides tumor
30 angiogenesis and metastasis inhibition.

6. The method of claim 1 further comprising subjecting the mammal in need thereof to a radiation therapy or chemotherapy regimen in combination with at least one
35 cytostatic or cytotoxic agent.

- 162 -

7. The method of claim 1 wherein the mammal in need thereof is a human.
8. The method of claim 1 wherein, within the compounds of formula (I), Y is a single bond, R₂ is a hydrogen atom, and R, R₁ and X are as above defined.
9. The method of claim 1 wherein, within the compounds of formula (I), Y is a -CO- group and R, R₁, R₂ and X are as above defined.
10. The method of claim 1 wherein, within the compounds of formula (I), Y is a -CONH- group and R, R₁, R₂ and X are as above defined.
11. The method of claim 1 wherein, within the compounds of formula (I), Y is a -SO₂- group and R, R₁, R₂ and X are as above defined.
12. A method for inhibiting protein kinase activity which comprises contacting the said kinase with an effective amount of a compound as defined in claim 1.
13. An aminoisoxazole derivative represented by formula (I)



wherein

R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

- 163 -

X is a divalent group selected from -N(R₃)- or -O-;
Y is a divalent group selected from -CH(R₃)-, -CO-,
-CONH- or -SO₂-, or Y may also be a single bond when R₂ is a
hydrogen atom or a C₃-C₆ cycloalkyl group;

5 R₁ is a hydrogen atom or a group, optionally further
substituted, selected from straight or branched C₁-C₆ alkyl,
C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered
heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3
heteroatoms selected among nitrogen, oxygen or sulfur; the
10 said cycloalkyl, aryl or heterocyclyl groups being
optionally further condensed with a 5 to 7 membered
aromatic or non-aromatic carbocycle or heterocycle with
from 1 to 3 heteroatoms selected among nitrogen, oxygen or
sulfur;

15 R₂ and R₃ have, each independently, the meanings above
reported for R₁ or represent an optionally substituted
straight or branched C₂-C₆ alkenyl or alkynyl group;
and the pharmaceutically acceptable salts thereof;
provided that:

20 a) R is other than nitrofuryl; and
b) 5-amino-3-(2-aminofuryl-5-yl)-4-methoxycarbonyl-
isoxazole and 5-amino-4-ethoxycarbonyl-3-(indol-3-yl)-
isoxazole, optionally further substituted at the indole
moiety, being excluded.

25

14. A compound of formula (I) according to claim 13
wherein Y is a single bond, R₂ is a hydrogen atom, and R, R₁
and X are as defined in claim 13.

30 **15.** The compound of claim 14 wherein X is a -N(R₃)- group
and R₃ is as defined in claim 13.

16. A compound of formula (I) according to claim 13
wherein Y is a -CO- group and R, R₁, R₂ and X are as defined
35 in claim 13.

- 164 -

17. The compound of claim 16 wherein X is -N(R₃)- and R₁ and R₃ are both hydrogen atoms.

18. A compound of formula (I) according to claim 13 wherein Y is a -CONH- group and R, R₁, R₂ and X are as above
5 defined.

19. The compound of claim 18 wherein X is -N(R₃)- and R₁ and R₃ are both hydrogen atoms.

10 20. A compound of formula (I) according to claim 13 wherein Y is a -SO₂- group and R, R₁, R₂ and X are as above defined.

21. The compound of claim 20 wherein X is -N(R₃)- and R₁
15 and R₃ are both hydrogen atoms.

22. A compound of formula (I) according to claim 13 wherein any heteroaryl or heterocyclyl group has 1 or 2 heteroatoms selected among nitrogen, oxygen or sulfur.

20

23. A compound of formula (I) as defined in claim 13, optionally in the form of a pharmaceutically acceptable salt, selected from:

- 1) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
25 amide;
- 2) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
amide;
- 3) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 4) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 30 5) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 6) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 7) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid amide;
- 8) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
35 amide;

- 9) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
butylamide;
- 10) 5-acetyl-amino-3-thiophen-2-yl-isoxazole-4-carboxylic
acid (4-methoxy-phenyl)-amide;
- 5 11) methyl 5-amino-3-(2-pyridinyl)-4-isoxazolecarboxylate;
- 12) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2,4-
difluoro-phenyl)-amide;
- 13) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
chloro-phenyl)-amide;
- 10 14) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
benzylamide;
- 15) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 16) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
15 fluoro-phenyl)-amide;
- 17) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 18) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(2,4-difluoro-phenyl)-amide;
- 20 19) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(4-chloro-phenyl)-amide;
- 20) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
benzylamide;
- 21) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
25 (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 22) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(4-fluoro-phenyl)-amide;
- 23) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 30 24) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(2,4-difluoro-phenyl)-amide;
- 25) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(4-chloro-phenyl)-amide;
- 26) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
35 benzylamide;

- 27) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 28) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(4-fluoro-phenyl)-amide;
- 5 29) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 30) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
(2,4-difluoro-phenyl)-amide;
- 31) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
10 chloro-phenyl)-amide;
- 32) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
benzylamide;
- 33) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 15 34) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
fluoro-phenyl)-amide;
- 35) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 36) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
20 (2,4-difluoro-phenyl)-amide;
- 37) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-
chloro-phenyl)-amide;
- 38) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
benzylamide;
- 25 39) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 40) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-
fluoro-phenyl)-amide;
- 41) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
30 benzhydryl-amide;
- 42) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
(2,4-difluoro-phenyl)-amide;
- 43) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-
chloro-phenyl)-amide;
- 35 44) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
benzylamide;

- 45) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
(5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 46) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide;
- 5 47) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 48) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2,4-difluoro-phenyl)-amide;
- 49) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
10 isoxazole-4-carboxylic acid (4-chloro-phenyl)-amide;
- 50) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (5,6,7,8-tetrahydro-
naphthalen-1-yl)-amide;
- 51) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
15 isoxazole-4-carboxylic acid (4-fluoro-phenyl)-amide;
- 52) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid benzhydryl-amide;
- 53) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(2,4-difluoro-phenyl)-amide;
- 20 54) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(4-chloro-phenyl)-amide;
- 55) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
benzylamide;
- 56) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
25 (5,6,7,8-tetrahydro-naphthalen-1-yl)-amide;
- 57) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(4-fluoro-phenyl)-amide;
- 58) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
benzhydryl-amide;
- 30 59) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
naphthalen-1-ylamide;
- 60) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid p-
tolylamide;
- 61) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-
35 benzyloxy-phenyl)-amide;

- 62) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 63) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 5 64) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
(3,5-dimethoxy-phenyl)-amide;
- 65) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
naphthalen-1-ylamide;
- 66) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid p-
10 tolylamide;
- 67) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(3-benzyloxy-phenyl)-amide;
- 68) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 15 69) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 70) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(3,5-dimethoxy-phenyl)-amide;
- 71) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
20 naphthalen-1-ylamide;
- 72) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid p-
tolylamide;
- 73) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(3-benzyloxy-phenyl)-amide;
- 25 74) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 75) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 76) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
30 (3,5-dimethoxy-phenyl)-amide;
- 77) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
naphthalen-1-ylamide;
- 78) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid p-
tolylamide;
- 35 79) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-
benzyloxy-phenyl)-amide;

- 80) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 81) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 5 82) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
(3,5-dimethoxy-phenyl)-amide;
- 83) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
naphthalen-1-ylamide;
- 84) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid p-
10 tolylamide;
- 85) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-
benzyloxy-phenyl)-amide;
- 86) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 15 87) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 88) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
(3,5-dimethoxy-phenyl)-amide;
- 89) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
20 naphthalen-1-ylamide;
- 90) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid p-
tolylamide;
- 91) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (3-
benzyloxy-phenyl)-amide;
- 25 92) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 93) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 94) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
30 (3,5-dimethoxy-phenyl)-amide;
- 95) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid naphthalen-1-ylamide;
- 96) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid p-tolylamide;
- 35 97) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (3-benzyloxy-phenyl)-amide;

- 98) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid 2,4-dimethoxy-benzylamide;
- 99) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid indan-5-ylamide;
- 5 100) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (3,5-dimethoxy-phenyl)-amide;
- 101) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
naphthalen-1-ylamide;
- 102) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid p-
10 tolylamide;
- 103) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(3-benzyloxy-phenyl)-amide;
- 104) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
2,4-dimethoxy-benzylamide;
- 15 105) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
indan-5-ylamide;
- 106) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(3,5-dimethoxy-phenyl)-amide;
- 107) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-
20 methoxy-dibenzofuran-3-yl)-amide;
- 108) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
(3,4,5-trimethoxy-phenyl)-amide;
- 109) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-
methoxy-5-trifluoromethyl-phenyl)-amide;
- 25 110) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 111) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
tert-butyl-phenyl)-amide;
- 112) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
30 (2-methoxy-dibenzofuran-3-yl)-amide;
- 113) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(3,4,5-trimethoxy-phenyl)-amide;
- 114) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 35 115) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(4-tert-butyl-phenyl)-amide;

- 116) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(2-methoxy-dibenzofuran-3-yl)-amide;
- 117) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(3,4,5-trimethoxy-phenyl)-amide;
- 5 118) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(2-methoxy-5-trifluoromethyl-phenyl)-amide;
- 119) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 120) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
10 (4-tert-butyl-phenyl)-amide;
- 121) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide;
- 122) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 15 123) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
tert-butyl-phenyl)-amide;
- 124) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide;
- 125) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
20 (3,4,5-trimethoxy-phenyl)-amide;
- 126) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-
methoxy-5-trifluoromethyl-phenyl)-amide;
- 127) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 25 128) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-
tert-butyl-phenyl)-amide;
- 129) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (2-
methoxy-dibenzofuran-3-yl)-amide;
- 130) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
30 quinolin-8-ylamide;
- 131) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-
tert-butyl-phenyl)-amide;
- 132) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2-methoxy-dibenzofuran-3-
35 yl)-amide;

- 133) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2-methoxy-5-trifluoromethyl-
phenyl)-amide;
- 134) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
5 isoxazole-4-carboxylic acid (4-tert-butyl-phenyl)-amide;
- 135) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(2-methoxy-dibenzofuran-3-yl)-amide;
- 136) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(3,4,5-trimethoxy-phenyl)-amide;
- 10 137) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
quinolin-8-ylamide;
- 138) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(4-tert-butyl-phenyl)-amide;
- 139) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (2-
15 benzyl-phenyl)-amide;
- 140) 3-[(5-amino-3-pyridin-4-yl-isoxazole-4-carbonyl)-
amino]-benzoic acid ethyl ester;
- 141) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
trifluoromethoxy-phenyl)-amide;
- 20 142) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(2-benzyl-phenyl)-amide;
- 143) 3-[(5-amino-3-thiophen-2-yl-isoxazole-4-carbonyl)-
amino]-benzoic acid ethyl ester;
- 144) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
25 (4-trifluoromethoxy-phenyl)-amide;
- 145) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(4-dimethylamino-phenyl)-amide;
- 146) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(2-benzyl-phenyl)-amide;
- 30 147) 3-[(5-amino-3-thiophen-3-yl-isoxazole-4-carbonyl)-
amino]-benzoic acid ethyl ester;
- 148) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(4-trifluoromethoxy-phenyl)-amide;
- 149) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
35 dimethylamino-phenyl)-amide;

- 150) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide;
- 151) 3-[(5-amino-3-thiazol-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester;
- 5 152) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide;
- 153) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide;
- 154) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide;
- 10 155) 3-[(5-amino-3-pyridin-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester;
- 156) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide;
- 15 157) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide;
- 158) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide;
- 159) 3-[(5-amino-3-pyridin-3-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester;
- 20 160) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide;
- 161) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide;
- 25 162) 3-{[5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carbonyl]-amino}-benzoic acid ethyl ester;
- 163) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-trifluoromethoxy-phenyl)-amide;
- 30 164) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (4-dimethylamino-phenyl)-amide;
- 165) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid (2-benzyl-phenyl)-amide;
- 166) 3-[(5-amino-3-quinolin-2-yl-isoxazole-4-carbonyl)-amino]-benzoic acid ethyl ester;
- 35

- 167) 5-amino-3-quinolin-2-yl-isoxazole-4-carboxylic acid
(4-trifluoromethoxy-phenyl)-amide;
- 168) 5-amino-3-pyridin-2-yl-isoxazole-4-phenyl carboxamide;
- 169) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
5 isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide;
- 170) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide;
- 171) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide;
- 10 172) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
(2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide;
- 173) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2,3-dihydro-
benzo[1,4]dioxin-6-yl)-amide;
- 15 174) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid
(9H-fluoren-2-yl)-amide;
- 175) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
(9H-fluoren-2-yl)-amide;
- 176) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
20 (4-morpholin-4-yl-phenyl)-amide;
- 177) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-
morpholin-4-yl-phenyl)-amide;
- 178) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-
25 amide;
- 179) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
(pyridin-4-ylmethyl)-amide;
- 180) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
phenylamide;
- 30 181) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
phenylamide;
- 182) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide;
- 183) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
35 morpholin-4-yl-phenyl)-amide;

- 184) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid
phenylamide;
- 185) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-
methoxy-phenyl)-amide;
- 5 186) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (3-phenoxy-phenyl)-amide;
- 187) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(4-morpholin-4-yl-phenyl)-amide;
- 188) 4-{[5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
10 isoxazole-4-carbonyl]-amino}-piperidine-1-carboxylic acid
ethyl ester;
- 189) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (tetrahydro-furan-2-
ylmethyl)-amide;
- 15 190) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
phenylamide;
- 191) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid 2-
methoxy-benzylamide;
- 192) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid 4-
20 chloro-benzylamide;
- 193) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid 2-
methoxy-benzylamide;
- 194) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 4-
fluoro-benzylamide;
- 25 195) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid
(3,4-dimethyl-phenyl)-amide;
- 196) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
isoxazole-4-carboxylic acid (2-dimethylamino-ethyl)-
amide;
- 30 197) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
(9H-fluoren-2-yl)-amide;
- 198) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid
(9H-fluoren-2-yl)-amide;
- 199) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
35 (3-methoxy-propyl)-amide;

- 200) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;
- 201) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;
- 5 202) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;
- 203) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide;
- 10 204) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide;
- 205) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (1-ethyl-propyl)-amide;
- 206) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
- 15 (3,4-dimethyl-phenyl)-amide;
- 207) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-fluoro-2-methyl-phenyl)-amide;
- 208) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid [4-(acetyl-methyl-amino)-phenyl]-amide;
- 20 209) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid 2-methoxy-benzylamide;
- 210) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide;
- 211) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide;
- 25 212) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid 4-chloro-benzylamide;
- 213) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-
- 30 amide;
- 214) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-phenoxy-phenyl)-amide;
- 215) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide;
- 35 216) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide;

- 217) 5-amino-3-pyridin-3-yl-isoxazole-4-carboxylic acid phenylamide;
- 218) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (2-methylsulfanyl-phenyl)-amide;
- 5 219) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (3-chloro-4-methyl-phenyl)-amide;
- 220) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-chloro-4-methyl-phenyl)-amide;
- 221) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
10 (3-methoxy-phenyl)-amide;
- 222) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide;
- 223) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide;
- 15 224) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-methoxy-propyl)-amide;
- 225) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid cyclopropylamide;
- 226) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid
20 (3-trifluoromethyl-phenyl)-amide;
- 227) 4-[(5-amino-3-thiophen-3-yl-isoxazole-4-carbonyl)-amino]-piperidine-1-carboxylic acid ethyl ester;
- 228) 4-[(5-amino-3-pyridin-2-yl-isoxazole-4-carbonyl)-amino]-piperidine-1-carboxylic acid ethyl ester;
- 25 229) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2,3-dihydro-benzo[1,4]dioxin-6-yl)-amide;
- 230) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide;
- 231) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid
30 (furan-2-ylmethyl)-amide;
- 232) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (furan-2-ylmethyl)-amide;
- 233) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid (furan-2-ylmethyl)-amide;
- 35 234) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;

- 235) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;
- 236) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide;
- 5 237) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (9H-fluoren-2-yl)-amide;
- 238) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide;
- 239) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid 4-
10 fluoro-benzylamide;
- 240) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-butoxy-phenyl)-amide;
- 241) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-difluoromethoxy-phenyl)-amide;
- 15 242) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid [4-(toluene-4-sulfonylamino)-phenyl]-amide;
- 243) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (2-methylsulfanyl-phenyl)-amide;
- 244) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
20 isoxazole-4-carboxylic acid phenylamide;
- 245) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (1-ethyl-propyl)-amide;
- 246) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide;
- 25 247) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid [4-(toluene-4-sulfonylamino)-phenyl]-amide;
- 248) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide;
- 249) 5-amino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-
30 isoxazole-4-carboxylic acid (3-chloro-4-methyl-phenyl)-amide;
- 250) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (4-diethylamino-phenyl)-amide;
- 251) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (2-
35 dimethylamino-ethyl)-amide;

- 252) 5-amino-3-thiazol-2-yl-isoxazole-4-carboxylic acid (3-trifluoromethyl-phenyl)-amide;
- 253) 5-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid (pyridin-4-ylmethyl)-amide;
- 5 254) 5-amino-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-morpholin-4-yl-phenyl)-amide;
- 255) 5-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (3-methoxy-phenyl)-amide;
- 256) 5-amino-3-pyridin-4-yl-isoxazole-4-carboxylic acid (4-
10 butoxy-phenyl)-amide;
- 257) 5-benzoylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid phenylamide;
- 258) 5-phenylacetylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid butylamide;
- 15 259) 5-(4-methyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 260) 5-(4-chloro-butyrylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 261) 5-(4-tert-butyl-benzoylamino)-3-pyridin-2-yl-
20 isoxazole-4-carboxylic acid amide;
- 262) 5-pentanoylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 263) 5-(4-methyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 25 264) 5-(4-chloro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 265) 5-pentanoylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 266) 5-(3-phenyl-propionylamino)-3-pyridin-4-yl-isoxazole-
30 4-carboxylic acid amide;
- 267) 5-(4-tert-butyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 268) 5-pentanoylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 35 269) 5-(3-phenyl-propionylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;

- 270) 5-(4-methyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 271) 5-(4-tert-butyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 5 272) 5-pentanoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 273) 5-(3-phenyl-propionylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 274) 5-(4-chloro-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 10 275) 5-(4-tert-butyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 276) 5-(4-chloro-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 15 277) 5-(3-phenyl-propionylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 278) 5-(4-methyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 279) 5-(4-tert-butyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 20 280) 5-(4-chloro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 281) 5-pentanoylamino-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 25 282) 5-(4-tert-butyl-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 283) 5-(4-chloro-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 30 284) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-pentanoylamino-isoxazole-4-carboxylic acid amide;
- 285) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(3-phenyl-propionylamino)-isoxazole-4-carboxylic acid amide;
- 286) 5-(4-chloro-butyrylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 35

- 287) 5-pentanoylamino-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 288) 5-(3-phenyl-propionylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 5 289) 5-(3-fluoro-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 290) 5-(3-methoxy-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 291) 3-pyridin-2-yl-5-(4-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 10 292) 5-(3-fluoro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 293) 5-(2-methoxy-acetyl-amino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 15 294) 5-benzoylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 295) 5-(3-methoxy-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 296) 3-pyridin-4-yl-5-(4-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 20 297) 5-(3-bromo-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 298) 5-(3-fluoro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 25 299) 5-(2-methoxy-acetyl-amino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 300) 5-benzoylamino-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 301) 5-(3-methoxy-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 30 302) 3-thiophen-2-yl-5-(4-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 303) 5-(2-methoxy-acetyl-amino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 35 304) 5-benzoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;

- 305) 5-(3-methoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 306) 3-thiophen-3-yl-5-(4-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 5 307) 5-(2-methoxy-acetylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 308) 5-benzoylamino-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 309) 5-(3-methoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-
10 carboxylic acid amide;
- 310) 3-thiazol-2-yl-5-(4-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 311) 5-(3-bromo-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 15 312) 5-(2-methoxy-acetylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 313) 5-(3-methoxy-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 314) 3-pyridin-3-yl-5-(4-trifluoromethyl-benzoylamino)-
20 isoxazole-4-carboxylic acid amide;
- 315) 5-(3-bromo-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 316) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(3-fluoro-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 25 317) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-methoxy-acetylamino)-isoxazole-4-carboxylic acid amide;
- 318) 5-(3-bromo-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 319) 3-quinolin-2-yl-5-(4-trifluoromethyl-benzoylamino)-
30 isoxazole-4-carboxylic acid amide;
- 320) 5-(cyclopropanecarbonyl-amino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 321) 5-(2-phenoxy-acetylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 35 322) 5-(2-chloro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;

- 323) 5-(cyclopropanecarbonyl-amino)-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 324) 5-(3-methyl-but-2-enoylamino)-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 5 325) 5-(2,4-difluoro-benzoylamino)-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 326) 5-[2-(3-methoxy-phenyl)-acetylamino]-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 327) 5-(2-phenoxy-acetylamino)-3-thiophen-2-yl-isoxazole-4-
10 carboxylic acid amide;
- 328) 5-(2-chloro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide;
- 329) 5-(cyclopropanecarbonyl-amino)-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 15 330) 5-(3-methyl-but-2-enoylamino)-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 331) 5-(2,4-difluoro-benzoylamino)-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 332) 5-[2-(3-methoxy-phenyl)-acetylamino]-3-thiophen-3-yl-
20 isoxazole-4-carboxylic acid amide;
- 333) 5-(2-chloro-benzoylamino)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid amide;
- 334) 5-(cyclopropanecarbonyl-amino)-3-thiophen-3-yl-
isoxazole-4-carboxylic acid amide;
- 25 335) 5-(3-methyl-but-2-enoylamino)-3-thiophen-3-yl-
isoxazole-4-carboxylic acid amide;
- 336) 5-(2,4-difluoro-benzoylamino)-3-thiophen-3-yl-
isoxazole-4-carboxylic acid amide;
- 337) 5-(2-phenoxy-acetylamino)-3-thiazol-2-yl-isoxazole-4-
30 carboxylic acid amide;
- 338) 5-(cyclopropanecarbonyl-amino)-3-thiazol-2-yl-
isoxazole-4-carboxylic acid amide;
- 339) 5-(3-methyl-but-2-enoylamino)-3-thiazol-2-yl-
isoxazole-4-carboxylic acid amide;
- 35 340) 5-(2,4-difluoro-benzoylamino)-3-thiazol-2-yl-
isoxazole-4-carboxylic acid amide;

- 341) 5-(2-phenoxy-acetylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 342) 5-(2-chloro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 5 343) 5-(cyclopropanecarbonyl-amino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 344) 5-(2,4-difluoro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 345) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-
10 phenoxy-acetylamino)-isoxazole-4-carboxylic acid amide;
- 346) 5-(2-chloro-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 347) 5-(cyclopropanecarbonyl-amino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid
15 amide;
- 348) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(3-methyl-but-2-enoylamino)-isoxazole-4-carboxylic acid amide;
- 349) 5-(2,4-difluoro-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid
20 amide;
- 350) 5-(cyclopropanecarbonyl-amino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 351) 5-(4-fluoro-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 25 352) 3-pyridin-2-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 353) 5-(2-methyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 354) 5-(4-ethyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
30
- 355) 5-[(naphthalene-1-carbonyl)-amino]-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 356) 5-(4-fluoro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 35 357) 3-pyridin-4-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;

- 358) 5-(2-methyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 359) 3-pyridin-4-yl-5-(3-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 5 360) 5-(4-ethyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 361) 5-[(naphthalene-1-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 362) 5-(4-fluoro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 10 363) 3-thiophen-2-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 364) 5-(2-methyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 15 365) 3-thiophen-2-yl-5-(3-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 366) 5-(4-ethyl-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 367) 5-[(naphthalene-1-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 20 368) 5-(4-fluoro-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 369) 3-thiophen-3-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 25 370) 5-(2-methyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 371) 3-thiophen-3-yl-5-(3-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 372) 5-(4-ethyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 30 373) 5-[(naphthalene-1-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 374) 5-(4-fluoro-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 35 375) 3-thiazol-2-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;

- 376) 5-(2-methyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 377) 3-thiazol-2-yl-5-(3-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 5 378) 5-(4-ethyl-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 379) 5-(4-fluoro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 380) 3-pyridin-3-yl-5-(4-trifluoromethoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 10 381) 3-pyridin-3-yl-5-(3-trifluoromethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 382) 5-(2-methyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 15 383) 5-(4-ethyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 384) 5-[(naphthalene-1-carbonyl)-amino]-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 385) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(4-fluoro-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 20 386) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-methyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 387) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(4-ethyl-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 25 388) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-[(naphthalene-1-carbonyl)-amino]-isoxazole-4-carboxylic acid amide;
- 389) 5-(2-fluoro-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 30 390) 5-(3-methyl-benzoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 391) 5-[2-(4-fluoro-phenyl)-acetylamino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 392) 5-(3,5-difluoro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 35

- 393) 5-(3-chloro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 394) 5-(2-fluoro-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 5 395) 5-(3-methyl-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 396) 5-(3,5-difluoro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 397) 5-(3-chloro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 10 398) 5-(3,5-difluoro-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 399) 5-(3-methyl-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 15 400) 5-(3-chloro-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 401) 5-(3-chloro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 402) 5-(2-fluoro-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 20 403) 5-(3-methyl-benzoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 404) 5-(3-chloro-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 25 405) 5-butyrylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 406) 5-[(E)-(3-phenyl-acryloyl)amino]-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 407) 5-(3,3-dimethyl-butyrylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 30 408) 5-(2-chloro-2-phenyl-acetylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 409) 5-butyrylamino-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 35 410) 5-[(biphenyl-4-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;

- 411) 5-(4-chloromethyl-benzoylamino)-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 412) 3-pyridin-4-yl-5-(3,4,5-trimethoxy-benzoylamino)-
isoxazole-4-carboxylic acid amide;
- 5 413) 5-[(E)-(3-phenyl-acryloyl)amino]-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 414) 5-(3,3-dimethyl-butyrylamino)-3-pyridin-4-yl-
isoxazole-4-carboxylic acid amide;
- 415) 5-(2-chloro-2-phenyl-acetylamino)-3-thiophen-2-yl-
10 isoxazole-4-carboxylic acid amide;
- 416) 5-(4-nitro-benzoylamino)-3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide;
- 417) 5-butyrylamino-3-thiophen-2-yl-isoxazole-4-carboxylic
acid amide;
- 15 418) 5-[(biphenyl-4-carbonyl)-amino]-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 419) 5-(4-chloromethyl-benzoylamino)-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 420) 3-thiophen-2-yl-5-(3,4,5-trimethoxy-benzoylamino)-
20 isoxazole-4-carboxylic acid amide;
- 421) 5-[(E)-(3-phenyl-acryloyl)amino]-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 422) 5-(3,3-dimethyl-butyrylamino)-3-thiophen-2-yl-
isoxazole-4-carboxylic acid amide;
- 25 423) 5-(2-chloro-2-phenyl-acetylamino)-3-thiophen-3-yl-
isoxazole-4-carboxylic acid amide;
- 424) 5-butyrylamino-3-thiophen-3-yl-isoxazole-4-carboxylic
acid amide;
- 425) 5-(3,3-dimethyl-butyrylamino)-3-thiophen-3-yl-
30 isoxazole-4-carboxylic acid amide;
- 426) 5-(4-chloromethyl-benzoylamino)-3-thiazol-2-yl-
isoxazole-4-carboxylic acid amide;
- 427) 5-(3,3-dimethyl-butyrylamino)-3-thiazol-2-yl-
isoxazole-4-carboxylic acid amide;
- 35 428) 5-(3,3-dimethyl-butyrylamino)-3-pyridin-3-yl-
isoxazole-4-carboxylic acid amide;

- 429) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-[(E)-(3-phenyl-acryloyl)amino]-isoxazole-4-carboxylic acid amide;
- 430) 5-butyrylamino-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 5 431) 5-(3,3-dimethyl-butyrylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 432) 5-butyrylamino-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 10 433) 5-(3,3-dimethyl-butyrylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 434) 5-[(furan-2-carbonyl)-amino]-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 435) 5-[(naphthalene-2-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 15 436) 5-(4-methoxy-benzoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 437) 5-[(furan-2-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 20 438) 5-(3-cyclopentyl-propionylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 439) 5-[(naphthalene-2-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 440) 5-(4-methoxy-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 25 441) 5-(2-benzyloxy-acetyl-amino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 442) 5-(2,4-dimethoxy-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 30 443) 5-[(furan-2-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 444) 5-(3-cyclopentyl-propionylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 445) 5-(4-bromo-benzoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 35

- 446) 5-[(naphthalene-2-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 447) 5-(4-methoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 5 448) 5-(2,4-dimethoxy-benzoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 449) 5-[(furan-2-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 450) 5-(3-cyclopentyl-propionylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 10 451) 5-(4-methoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 452) 5-(2,4-dimethoxy-benzoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 15 453) 5-[(furan-2-carbonyl)-amino]-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 454) 5-(3-cyclopentyl-propionylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 455) 5-[(furan-2-carbonyl)-amino]-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 20 456) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(4-methoxy-benzoylamino)-isoxazole-4-carboxylic acid amide;
- 457) 5-(2,4-dimethoxy-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 25 458) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-[(furan-2-carbonyl)-amino]-isoxazole-4-carboxylic acid amide;
- 459) 5-(3-cyclopentyl-propionylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 30 460) 5-(3,4-dimethoxy-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 461) 5-(4-bromo-benzoylamino)-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 35

- 462) 5-(3-cyclopentyl-propionylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 463) 5-(2-ethyl-butyrylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 5 464) 5-(2-propyl-pentanoylamino)-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 465) 5-(3-methyl-butyrylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 466) 5-[(benzo[1,3]dioxole-5-carbonyl)-amino]-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 10 467) 5-(2-ethyl-butyrylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 468) 5-(2-propyl-pentanoylamino)-3-pyridin-4-yl-isoxazole-4-carboxylic acid amide;
- 15 469) 5-(3-methyl-butyrylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 470) 5-[(benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 471) 5-(2-ethyl-butyrylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 20 472) 5-(2-propyl-pentanoylamino)-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 473) 5-[2-(3,4-dimethoxy-phenyl)-acetylamino]-3-thiophen-2-yl-isoxazole-4-carboxylic acid amide;
- 25 474) 5-(3-methyl-butyrylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 475) 5-[(benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 476) 5-(2-ethyl-butyrylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 30 477) 5-(2-propyl-pentanoylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 478) 5-(3-methyl-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 35 479) 5-[(benzo[1,3]dioxole-5-carbonyl)-amino]-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;

- 480) 5-(2-ethyl-butyrylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 481) 5-(2-propyl-pentanoylamino)-3-thiazol-2-yl-isoxazole-4-carboxylic acid amide;
- 5 482) 5-(2-ethyl-butyrylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 483) 5-(2-propyl-pentanoylamino)-3-pyridin-3-yl-isoxazole-4-carboxylic acid amide;
- 484) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(3-methyl-10 butyrylamino)-isoxazole-4-carboxylic acid amide;
- 485) 5-[(benzo[1,3]dioxole-5-carbonyl)-amino]-3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-isoxazole-4-carboxylic acid amide;
- 486) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-ethyl-15 butyrylamino)-isoxazole-4-carboxylic acid amide;
- 487) 3-(3,5-dimethyl-1-phenyl-1H-pyrazol-4-yl)-5-(2-propyl-pentanoylamino)-isoxazole-4-carboxylic acid amide;
- 488) 5-(3-methyl-butyrylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 20 489) 5-(2-ethyl-butyrylamino)-3-quinolin-2-yl-isoxazole-4-carboxylic acid amide;
- 490) 5-Phenylacetyl-amino-3-thiophen-2-yl-isoxazole-4-carboxylic acid butylamide;
- 491) 5-Phenylacetyl-amino-3-thiophen-3-yl-isoxazole-4-25 carboxylic acid butylamide;
- 492) 5-Butyrylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 493) 5-[(Furan-2-carbonyl)-amino]-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 30 494) 5-Benzoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 495) 5-Phenylacetyl-amino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 496) 5-Butyrylamino-3-thiophen-3-yl-isoxazole-4-carboxylic 35 acid (4-methoxy-phenyl)-amide;

- 497) 5-[(Furan-2-carbonyl)-amino]-3-thiophen-3-yl-
isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 498) 5-Benzoylamino-3-thiophen-3-yl-isoxazole-4-carboxylic
acid (4-methoxy-phenyl)-amide;
- 5 499) 5-Phenylacetyl-amino-3-pyridin-2-yl-isoxazole-4-
carboxylic acid butylamide;
- 500) 5-Butyrylamino-3-pyridin-2-yl-isoxazole-4-carboxylic
acid butylamide;
- 501) 5-[(Furan-2-carbonyl)-amino]-3-pyridin-2-yl-isoxazole-
10 4-carboxylic acid butylamide;
- 502) 5-Benzoylamino-3-pyridin-2-yl-isoxazole-4-carboxylic
acid butylamide;
- 503) 5-Phenylacetyl-amino-3-pyridin-2-yl-isoxazole-4-
carboxylic acid (4-methoxy-phenyl)-amide;
- 15 504) 5-Butyrylamino-3-pyridin-2-yl-isoxazole-4-carboxylic
acid (4-methoxy-phenyl)-amide;
- 505) 5-[(Furan-2-carbonyl)-amino]-3-pyridin-2-yl-isoxazole-
4-carboxylic acid (4-methoxy-phenyl)-amide;
- 506) 5-Benzoylamino-3-pyridin-2-yl-isoxazole-4-carboxylic
20 acid (4-methoxy-phenyl)-amide;
- 507) 5-(4-Acetyl-amino-benzenesulfonylamino)-3-pyridin-2-yl-
isoxazole-4-carboxylic acid phenylamide;
- 508) 5-(4-Acetyl-amino-benzenesulfonylamino)-3-pyridin-2-yl-
isoxazole-4-carboxylic acid amide;
- 25 509) 5-(Butane-1-sulfonylamino)-3-pyridin-2-yl-isoxazole-4-
carboxylic acid amide;
- 510) 3-Pyridin-2-yl-5-(toluene-4-sulfonylamino)-isoxazole-
4-carboxylic acid amide;
- 511) 5-(2,5-Dichloro-thiophene-3-sulfonylamino)-3-pyridin-
30 2-yl-isoxazole-4-carboxylic acid amide;
- 512) 5-(Butane-1-sulfonylamino)-3-pyridin-2-yl-isoxazole-4-
carboxylic acid acetylamide;
- 513) 3-Pyridin-2-yl-5-(toluene-4-sulfonylamino)-isoxazole-
4-carboxylic acid benzylamide;
- 35 514) 5-(2,5-Dichloro-thiophene-3-sulfonylamino)-3-pyridin-
2-yl-isoxazole-4-carboxylic acid phenylamide;

- 515) 5-Methanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 516) 5-Benzenesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 5 517) 5-Phenylmethanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 518) 5-(Thiophene-2-sulfonylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid amide;
- 519) 5-Benzenesulfonylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 10 520) 5-Phenylmethanesulfonylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 521) 3-Pyridin-2-yl-5-(thiophene-2-sulfonylamino)-isoxazole-4-carboxylic acid amide;
- 15 522) 5-Methanesulfonylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid amide;
- 523) 5-Methanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 524) 5-Benzenesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 20 525) 5-Phenylmethanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 526) 5-(Thiophene-2-sulfonylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid butylamide;
- 25 527) 5-Methanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 528) 5-Benzenesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 529) 5-Phenylmethanesulfonylamino-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 30 530) 5-(Thiophene-2-sulfonylamino)-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 531) 5-Benzenesulfonylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid butylamide;
- 35 532) 5-Phenylmethanesulfonylamino-3-pyridin-2-yl-isoxazole-4-carboxylic acid butylamide;

- 533) 3-Pyridin-2-yl-5-(thiophene-2-sulfonylamino)-
isoxazole-4-carboxylic acid butylamide;
- 534) 5-Methanesulfonylamino-3-pyridin-2-yl-isoxazole-4-
carboxylic acid butylamide;
- 5 535) 5-Methanesulfonylamino-3-pyridin-2-yl-isoxazole-4-
carboxylic acid (4-methoxy-phenyl)-amide;
- 536) 5-Benzenesulfonylamino-3-pyridin-2-yl-isoxazole-4-
carboxylic acid (4-methoxy-phenyl)-amide;
- 537) 5-Phenylmethanesulfonylamino-3-pyridin-2-yl-isoxazole-
10 4-carboxylic acid (4-methoxy-phenyl)-amide;
- 538) 3-Pyridin-2-yl-5-(thiophene-2-sulfonylamino)-
isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 539) Phenyl 5-amino-3-(2-pyridinyl)-4-isoxazolecarboxylate
- 540) Cyclohexyl 5-amino-3-(2-pyridinyl)-4-
15 isoxazolecarboxylate
- 541) 5-Benzothiophenyl 5-amino-3-(2-pyridinyl)-4-
isoxazolecarboxylate;
- 542) Phenyl 5-acetylamino-3-(2-pyridinyl)-4-
isoxazolecarboxylate;
- 20 543) Cyclohexyl 5-benzoylamino-3-(2-pyridinyl)-4-
isoxazolecarboxylate;
- 544) 5-Benzothiophenyl 5-(2,4-dimethoxybenzoylamino)-3-(2-
pyridinyl)-4-isoxazolecarboxylate;
- 545) 5-(3-Phenyl-ureido)-3-thiophen-2-yl-isoxazole-4-
25 carboxylic acid amide;
- 546) 5-(3-Isopropyl-ureido)-3-thiophen-2-yl-isoxazole-4-
carboxylic acid amide;
- 547) 5-(3-Phenyl-ureido)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid butylamide;
- 30 548) 5-(3-Benzyl-ureido)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid butylamide;
- 549) 5-(3-Isopropyl-ureido)-3-thiophen-3-yl-isoxazole-4-
carboxylic acid butylamide;
- 550) 5-(3-Pyridin-3-yl-ureido)-3-thiophen-3-yl-isoxazole-4-
35 carboxylic acid butylamide;

- 551) 5-(3-Phenyl-ureido)-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 552) 5-(3-Benzyl-ureido)-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 5 553) 5-(3-Isopropyl-ureido)-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 554) 5-(3-Pyridin-3-yl-ureido)-3-thiophen-3-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 555) 5-(3-Phenyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid butylamide;
- 10 556) 5-(3-Benzyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid butylamide;
- 557) 5-(3-Isopropyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid butylamide;
- 15 558) 3-Pyridin-2-yl-5-(3-pyridin-3-yl-ureido)-isoxazole-4-carboxylic acid butylamide;
- 559) 5-(3-Phenyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 560) 5-(3-Benzyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 20 561) 5-(3-Isopropyl-ureido)-3-pyridin-2-yl-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 562) 3-Pyridin-2-yl-5-(3-pyridin-3-yl-ureido)-isoxazole-4-carboxylic acid (4-methoxy-phenyl)-amide;
- 25 563) 5-amino-N-benzyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 564) 5-amino-N-(2-furylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 565) 5-amino-N-(2-phenylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 30 566) 5-amino-N-cyclohexyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 567) 5-amino-N-(4-methoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 35 568) 5-amino-N-hexyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;

- 569) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 570) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 5 571) 5-amino-N-cyclopentyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 572) 5-amino-N-(1,5-dimethylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 573) 5-amino-N-isopentyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 10 574) 5-amino-N-(2-ethylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 575) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 15 576) 5-amino-N-(2-methylcyclohexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 577) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 578) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 20 579) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 580) 5-amino-N-isobutyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 25 581) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 582) 5-amino-N-(2-methoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 583) 5-amino-N-(2-fluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 30 584) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 585) 5-amino-N-(3,4-difluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 35 586) 5-amino-N-(3-chlorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;

- 587) 5-amino-N-(2-chlorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 588) 5-amino-N-(3-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 5 589) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 590) 5-amino-N-(cyclopropylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 591) 5-amino-N-cycloheptyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 10 592) 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 593) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 15 594) 5-amino-N-(1,2-dimethylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 595) 5-amino-N-(3,3-diphenylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 596) 5-amino-N-(1,2-diphenylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 20 597) 5-amino-N-(2,4-dimethoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 598) 5-amino-N-[3-(dibutylamino)propyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 25 599) 5-amino-N-butyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 600) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 601) 5-amino-N-(1-methylhexyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 30 602) 5-amino-N-(2-methoxy-1-methylethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 603) 5-amino-N-(1-methylbutyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 35 604) 5-amino-N-(3-fluorobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;

- 605) 5-amino-N-(4-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 606) 5-amino-N-(1,3-dimethylbutyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 5 607) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 608) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 609) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 10 610) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-pentylisoxazole-4-carboxamide;
- 611) 5-amino-N-(cyclohexylmethyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 15 612) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 613) 5-amino-N-(1-ethylpropyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 614) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 20 615) 5-amino-N-(2-methylbenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 616) 5-amino-N-cyclobutyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 25 617) 5-amino-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}-N-octylisoxazole-4-carboxamide;
- 618) 5-amino-N-(2-bromobenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 619) 5-amino-N-(1-methylheptyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 30 620) 5-amino-N-heptyl-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 621) 5-amino-N-(tert-butyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 35 622) 5-amino-N-(3,4-dimethoxybenzyl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;

- 623) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 624) 5-amino-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-3-{5-[3-(trifluoromethyl)phenyl]-2-furyl}isoxazole-4-carboxamide;
- 5 625) 5-amino-N-benzyl-3-(5-bromothien-2-yl)isoxazole-4-carboxamide;
- 626) 5-amino-3-(5-bromothien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide;
- 627) 5-amino-3-(5-bromothien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 10 628) 5-amino-3-(5-bromothien-2-yl)-N-(sec-butyl)isoxazole-4-carboxamide;
- 629) 5-amino-3-(5-bromothien-2-yl)-N-cyclohexylisoxazole-4-carboxamide;
- 15 630) 5-amino-3-(5-bromothien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 631) 5-amino-3-(5-bromothien-2-yl)-N-isopropylisoxazole-4-carboxamide;
- 632) 5-amino-3-(5-bromothien-2-yl)-N-hexylisoxazole-4-carboxamide;
- 20 633) N-allyl-5-amino-3-(5-bromothien-2-yl)isoxazole-4-carboxamide;
- 634) 5-amino-3-(5-bromothien-2-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 25 635) 5-amino-3-(5-bromothien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 636) 5-amino-3-(5-bromothien-2-yl)-N-cyclopentylisoxazole-4-carboxamide;
- 637) 5-amino-3-(5-bromothien-2-yl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide;
- 30 638) 5-amino-3-(5-bromothien-2-yl)-N-isopentylisoxazole-4-carboxamide;
- 639) 5-amino-3-(5-bromothien-2-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide;
- 35 640) 5-amino-3-(5-bromothien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;

- 641) 5-amino-3-(5-bromothien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 642) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-bromothien-2-yl)isoxazole-4-carboxamide;
- 5 643) 5-amino-3-(5-bromothien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 644) 5-amino-3-(5-bromothien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 645) 5-amino-3-(5-bromothien-2-yl)-N-isobutylisoxazole-4-
10 carboxamide;
- 646) 5-amino-3-(5-bromothien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 647) 5-amino-3-(5-bromothien-2-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 15 648) 5-amino-3-(5-bromothien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide;
- 649) 5-amino-3-(5-bromothien-2-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 650) 5-amino-3-(5-bromothien-2-yl)-N-(3,4-
20 difluorobenzyl)isoxazole-4-carboxamide;
- 651) 5-amino-3-(5-bromothien-2-yl)-N-(3-chlorobenzyl)isoxazole-4-carboxamide;
- 652) 5-amino-3-(5-bromothien-2-yl)-N-(2-chlorobenzyl)isoxazole-4-carboxamide;
- 25 653) 5-amino-3-(5-bromothien-2-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 654) 5-amino-3-(5-bromothien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 655) 5-amino-3-(5-bromothien-2-yl)-N-
30 (cyclopropylmethyl)isoxazole-4-carboxamide;
- 656) 5-amino-3-(5-bromothien-2-yl)-N-cycloheptylisoxazole-4-carboxamide;
- 657) 5-amino-3-(5-bromothien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 35 658) 5-amino-3-(5-bromothien-2-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide;

- 659) 5-amino-3-(5-bromothien-2-yl)-N-prop-2-ynylisoxazole-4-carboxamide;
- 660) 5-amino-3-(5-bromothien-2-yl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide;
- 5 661) 5-amino-3-(5-bromothien-2-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide;
- 662) 5-amino-3-(5-bromothien-2-yl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide;
- 663) 5-amino-3-(5-bromothien-2-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide;
- 10 664) 5-amino-3-(5-bromothien-2-yl)-N-butylisoxazole-4-carboxamide;
- 665) 5-amino-3-(5-bromothien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 15 666) 5-amino-3-(5-bromothien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 667) 5-amino-3-(5-bromothien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 668) 5-amino-3-(5-bromothien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;
- 20 669) 5-amino-3-(5-bromothien-2-yl)-N-(4-tert-butylcyclohexyl)isoxazole-4-carboxamide;
- 670) 5-amino-3-(5-bromothien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 25 671) 5-amino-3-(5-bromothien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide;
- 672) 5-amino-3-(5-bromothien-2-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide;
- 673) 5-amino-3-(5-bromothien-2-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide;
- 30 674) 5-amino-3-(5-bromothien-2-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 675) 5-amino-3-(5-bromothien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 35 676) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-bromothien-2-yl)isoxazole-4-carboxamide;

- 677) 5-amino-3-(5-bromothien-2-yl)-N-pentylisoxazole-4-carboxamide;
- 678) 5-amino-3-(5-bromothien-2-yl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide;
- 5 679) ethyl 4-([5-amino-3-(5-bromothien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 680) 5-amino-3-(5-bromothien-2-yl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide;
- 681) 5-amino-3-(5-bromothien-2-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide;
- 10 682) 5-amino-3-(5-bromothien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 683) 5-amino-3-(5-bromothien-2-yl)-N-propylisoxazole-4-carboxamide;
- 15 684) 5-amino-3-(5-bromothien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 685) 5-amino-3-(5-bromothien-2-yl)-N-cyclobutylisoxazole-4-carboxamide;
- 686) 5-amino-3-(5-bromothien-2-yl)-N-(3-methoxypropyl)isoxazole-4-carboxamide;
- 20 687) 5-amino-3-(5-bromothien-2-yl)-N-octylisoxazole-4-carboxamide;
- 688) 5-amino-N-(2-bromobenzyl)-3-(5-bromothien-2-yl)isoxazole-4-carboxamide;
- 25 689) 5-amino-3-(5-bromothien-2-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 690) 5-amino-3-(5-bromothien-2-yl)-N-heptylisoxazole-4-carboxamide;
- 691) 5-amino-3-(5-bromothien-2-yl)-N-(tert-butyl)isoxazole-4-carboxamide;
- 30 692) 5-amino-3-(5-bromothien-2-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide;
- 693) 5-amino-3-(5-bromothien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 35 694) 5-amino-N-benzyl-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;

- 695) 5-amino-3-(5-chlorothien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide;
- 696) 5-amino-3-(5-chlorothien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 5 697) 5-amino-N-(sec-butyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 698) 5-amino-3-(5-chlorothien-2-yl)-N-cyclohexylisoxazole-4-carboxamide;
- 699) 5-amino-3-(5-chlorothien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 10 700) 5-amino-3-(5-chlorothien-2-yl)-N-isopropylisoxazole-4-carboxamide;
- 701) 5-amino-3-(5-chlorothien-2-yl)-N-hexylisoxazole-4-carboxamide;
- 15 702) N-allyl-5-amino-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 703) 5-amino-3-(5-chlorothien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 704) 5-amino-3-(5-chlorothien-2-yl)-N-cyclopentylisoxazole-4-carboxamide;
- 20 705) 5-amino-3-(5-chlorothien-2-yl)-N-isopentylisoxazole-4-carboxamide;
- 706) 5-amino-3-(5-chlorothien-2-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide;
- 25 707) 5-amino-3-(5-chlorothien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 708) 5-amino-3-(5-chlorothien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 709) 5-amino-3-(5-chlorothien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 30 710) 5-amino-3-(5-chlorothien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide;
- 711) 5-amino-3-(5-chlorothien-2-yl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide;
- 35 712) 5-amino-N-(3-chlorobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;

- 713) 5-amino-N-(2-chlorobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 714) 5-amino-3-(5-chlorothien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 5 715) 5-amino-3-(5-chlorothien-2-yl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide;
- 716) 5-amino-3-(5-chlorothien-2-yl)-N-cycloheptylisoxazole-4-carboxamide;
- 717) 5-amino-3-(5-chlorothien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 10 718) 5-amino-3-(5-chlorothien-2-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide;
- 719) 5-amino-3-(5-chlorothien-2-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide;
- 15 720) 5-amino-3-(5-chlorothien-2-yl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide;
- 721) 5-amino-3-(5-chlorothien-2-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide;
- 722) 5-amino-N-butyl-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 20 723) 5-amino-3-(5-chlorothien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 724) 5-amino-3-(5-chlorothien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 25 725) 5-amino-3-(5-chlorothien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 726) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 727) 5-amino-3-(5-chlorothien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;
- 30 728) 5-amino-3-(5-chlorothien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 729) 5-amino-3-(5-chlorothien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide;
- 35 730) 5-amino-3-(5-chlorothien-2-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide;

- 731) 5-amino-3-(5-chlorothien-2-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 732) 5-amino-3-(5-chlorothien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 5 733) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 734) 5-amino-3-(5-chlorothien-2-yl)-N-pentylisoxazole-4-carboxamide;
- 735) 5-amino-3-(5-chlorothien-2-yl)-N-
- 10 (cyclohexylmethyl)isoxazole-4-carboxamide;
- 736) ethyl 4-([5-amino-3-(5-chlorothien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 737) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 15 738) 5-amino-3-(5-chlorothien-2-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide;
- 739) 5-amino-3-(5-chlorothien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 740) 5-amino-3-(5-chlorothien-2-yl)-N-propylisoxazole-4-
- 20 carboxamide;
- 741) 5-amino-3-(5-chlorothien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 742) 5-amino-3-(5-chlorothien-2-yl)-N-cyclobutylisoxazole-4-carboxamide;
- 25 743) 5-amino-3-(5-chlorothien-2-yl)-N-(3-methoxypropyl)isoxazole-4-carboxamide;
- 744) 5-amino-N-(2-bromobenzyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 745) 5-amino-3-(5-chlorothien-2-yl)-N-(1-
- 30 methylheptyl)isoxazole-4-carboxamide;
- 746) 5-amino-3-(5-chlorothien-2-yl)-N-heptylisoxazole-4-carboxamide;
- 747) 5-amino-N-(tert-butyl)-3-(5-chlorothien-2-yl)isoxazole-4-carboxamide;
- 35 748) 5-amino-3-(5-chlorothien-2-yl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide;

- 749) 5-amino-3-(5-chlorothien-2-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide;
- 750) 5-amino-3-(5-chlorothien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 5 751) 5-amino-N-benzyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 752) 5-amino-N-(2-furylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 753) 5-amino-3-(5-methylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 10 754) 5-amino-N-(sec-butyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 755) 5-amino-N-cyclohexyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 15 756) 5-amino-N-(4-methoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 757) 5-amino-N-isopropyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 758) 5-amino-N-hexyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 20 759) N-allyl-5-amino-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 760) 5-amino-3-(5-methylthien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 25 761) 5-amino-N-cyclopentyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 762) 5-amino-N-(1,5-dimethylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 763) 5-amino-N-isopentyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 30 764) 5-amino-N-(2-ethylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 765) 5-amino-3-(5-methylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 35 766) 5-amino-N-(2-methylcyclohexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;

- 767) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 768) 5-amino-3-(5-methylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 5 769) 5-amino-3-(5-methylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 770) 5-amino-N-isobutyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 771) 5-amino-3-(5-methylthien-2-yl)-N-[2-
- 10 (trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 772) 5-amino-N-(2-methoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 773) 5-amino-N-(2-fluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 15 774) 5-amino-N-(3,4-difluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 775) 5-amino-N-(3-chlorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 776) 5-amino-N-cycloheptyl-3-(5-methylthien-2-yl)isoxazole-
- 20 4-carboxamide;
- 777) 5-amino-N-(1,2-dimethylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 778) 5-amino-N-(3,3-diphenylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 25 779) 5-amino-N-(1,2-diphenylethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 780) 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 781) 5-amino-N-[3-(dibutylamino)propyl]-3-(5-methylthien-2-
- 30 yl)isoxazole-4-carboxamide;
- 782) 5-amino-N-butyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 783) 5-amino-3-(5-methylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 35 784) 5-amino-N-[(5-methyl-2-furyl)methyl]-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;

- 785) 5-amino-N-(1-methylhexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 786) 5-amino-N-(2-methoxy-1-methylethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 5 787) 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 788) 5-amino-N-(1-methylbutyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 789) 5-amino-N-(3-fluorobenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 10 790) 5-amino-N-(4-methylbenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 791) 5-amino-N-(1-methyl-3-phenylpropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 15 792) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 793) 5-amino-3-(5-methylthien-2-yl)-N-pentylisoxazole-4-carboxamide;
- 794) 5-amino-N-(cyclohexylmethyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 20 795) 5-amino-3-(5-methylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 796) 5-amino-3-(5-methylthien-2-yl)-N-propylisoxazole-4-carboxamide;
- 25 797) 5-amino-N-(2-methylbenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 798) 5-amino-N-cyclobutyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 799) 5-amino-N-(3-methoxypropyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 30 800) 5-amino-3-(5-methylthien-2-yl)-N-octylisoxazole-4-carboxamide;
- 801) 5-amino-N-(1-methylheptyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 35 802) 5-amino-N-heptyl-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;

- 803) 5-amino-N-(tert-butyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 804) 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 5 805) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-methylthien-2-yl)isoxazole-4-carboxamide;
- 806) 5-amino-3-(5-methylthien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 807) 5-amino-N-benzyl-3-(3-methylthien-2-yl)isoxazole-4-
10 carboxamide;
- 808) 5-amino-N-(2-furylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 809) 5-amino-3-(3-methylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 15 810) 5-amino-N-(sec-butyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 811) 5-amino-N-cyclohexyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 812) 5-amino-N-(4-methoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 20 813) 5-amino-N-isopropyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 814) 5-amino-N-hexyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 25 815) N-allyl-5-amino-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 816) 5-amino-3-(3-methylthien-2-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 817) 5-amino-N-cyclopentyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 30 818) 5-amino-N-(1,5-dimethylhexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 819) 5-amino-N-isopentyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 35 820) 5-amino-N-(2-ethylhexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;

- 821) 5-amino-3-(3-methylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 822) 5-amino-N-(2-methylcyclohexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 5 823) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 824) 5-amino-3-(3-methylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 825) 5-amino-3-(3-methylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 10 826) 5-amino-N-isobutyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 827) 5-amino-3-(3-methylthien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 15 828) 5-amino-N-(2-methoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 829) 5-amino-N-(2-fluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 830) 5-amino-3-(3-methylthien-2-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 20 831) 5-amino-N-(3,4-difluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 832) 5-amino-N-(3-chlorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 25 833) 5-amino-N-(2-chlorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 834) 5-amino-N-(3-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 835) 5-amino-3-(3-methylthien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 30 836) 5-amino-N-(cyclopropylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 837) 5-amino-N-cycloheptyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 35 838) 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;

- 839) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 840) 5-amino-N-(1,2-dimethylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 5 841) 5-amino-N-(3,3-diphenylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 842) 5-amino-N-(2,4-dimethoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 843) 5-amino-N-[3-(dibutylamino)propyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 10 844) 5-amino-N-butyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 845) 5-amino-3-(3-methylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 15 846) 5-amino-N-[(5-methyl-2-furyl)methyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 847) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 848) 5-amino-N-(2-methoxy-1-methylethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 20 849) 5-amino-N-(4-tert-butylcyclohexyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 850) 5-amino-N-(1-methylbutyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 25 851) 5-amino-N-(3-fluorobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 852) 5-amino-N-(4-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 853) 5-amino-N-(1,3-dimethylbutyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 30 854) 5-amino-N-(1-methyl-3-phenylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 855) 5-amino-3-(3-methylthien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 35 856) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;

- 213 -

- 857) 5-amino-3-(3-methylthien-2-yl)-N-pentylisoxazole-4-carboxamide;
- 858) 5-amino-N-(cyclohexylmethyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 5 859) ethyl 4-([5-amino-3-(3-methylthien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 860) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 861) 5-amino-N-(1-ethylpropyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 10 862) 5-amino-3-(3-methylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 863) 5-amino-N-(2-methylbenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 15 864) 5-amino-N-cyclobutyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 865) 5-amino-N-benzhydryl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 866) 5-amino-3-(3-methylthien-2-yl)-N-octylisoxazole-4-carboxamide;
- 20 867) 5-amino-N-(2-bromobenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 868) 5-amino-N-(1-methylheptyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 25 869) 5-amino-N-heptyl-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 870) 5-amino-N-(tert-butyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 871) 5-amino-N-(3,4-dimethoxybenzyl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 30 872) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methylthien-2-yl)isoxazole-4-carboxamide;
- 873) 5-amino-3-(3-methylthien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 35 874) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-furylmethyl)isoxazole-4-carboxamide;

- 875) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 876) 5-amino-N-(sec-butyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 5 877) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclohexylisoxazole-4-carboxamide;
- 878) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 879) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-
- 10 isopropylisoxazole-4-carboxamide;
- 880) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-hexylisoxazole-4-carboxamide;
- 881) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 15 882) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclopentylisoxazole-4-carboxamide;
- 883) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide;
- 884) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-
- 20 isopentylisoxazole-4-carboxamide;
- 885) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-ethylhexyl)isoxazole-4-carboxamide;
- 886) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 25 887) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 888) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 889) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-
- 30 (trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 890) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 891) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-isobutylisoxazole-4-carboxamide;
- 35 892) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;

- 893) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 894) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-fluorobenzyl)isoxazole-4-carboxamide;
- 5 895) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 896) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide;
- 897) 5-amino-N-(3-chlorobenzyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 10 898) 5-amino-N-(2-chlorobenzyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 899) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 15 900) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 901) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(cyclopropylmethyl)isoxazole-4-carboxamide;
- 902) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cycloheptylisoxazole-4-carboxamide;
- 20 903) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 904) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide;
- 25 905) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide;
- 906) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide;
- 907) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2-diphenylethyl)isoxazole-4-carboxamide;
- 30 908) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 909) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide;
- 35 910) 5-amino-N-butyl-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;

- 911) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 912) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 5 913) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 914) 5-amino-N-(1-benzylpiperidin-4-yl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 915) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,3-
10 dimethylbutyl)isoxazole-4-carboxamide;
- 916) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 917) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 15 918) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-pentylisoxazole-4-carboxamide;
- 919) ethyl 4-[(5-amino-3-[5-(3-chlorophenyl)-2-furyl]isoxazol-4-yl)carbonyl]amino]piperidine-1-carboxylate;
- 20 920) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 921) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-ethylpropyl)isoxazole-4-carboxamide;
- 922) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-
25 phenylethyl)isoxazole-4-carboxamide;
- 923) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 924) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-cyclobutylisoxazole-4-carboxamide;
- 30 925) 5-amino-N-benzhydryl-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 926) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-octylisoxazole-4-carboxamide;
- 927) 5-amino-N-(2-bromobenzyl)-3-[5-(3-chlorophenyl)-2-
35 furyl]isoxazole-4-carboxamide;

- 217 -

- 928) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 929) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-heptylisoxazole-4-carboxamide;
- 5 930) 5-amino-N-(tert-butyl)-3-[5-(3-chlorophenyl)-2-furyl]isoxazole-4-carboxamide;
- 931) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 932) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(2,3-dihydro-10 1H-inden-1-yl)isoxazole-4-carboxamide;
- 933) 5-amino-3-[5-(3-chlorophenyl)-2-furyl]-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 934) 5-amino-N-benzyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 15 935) 5-amino-N-(2-furylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 936) 5-amino-3-(5-methyl-2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 937) 5-amino-N-(sec-butyl)-3-(5-methyl-2-furyl)isoxazole-4-20 carboxamide;
- 938) 5-amino-N-cyclohexyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 939) 5-amino-N-(4-methoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 25 940) 5-amino-N-hexyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 941) 5-amino-3-(5-methyl-2-furyl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 942) 5-amino-3-(5-methyl-2-furyl)-N-(thien-2-30 ylmethyl)isoxazole-4-carboxamide;
- 943) 5-amino-N-cyclopentyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 944) 5-amino-N-(1,5-dimethylhexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 35 945) 5-amino-N-isopentyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;

- 946) 5-amino-N-(2-ethylhexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 947) 5-amino-3-(5-methyl-2-furyl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 5 948) 5-amino-N-(2-methylcyclohexyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 949) 5-amino-3-(5-methyl-2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 950) 5-amino-3-(5-methyl-2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 10 951) 5-amino-N-isobutyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 952) 5-amino-3-(5-methyl-2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 15 953) 5-amino-N-(2-methoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 954) 5-amino-N-(2-fluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 955) 5-amino-3-(5-methyl-2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 20 956) 5-amino-N-(3,4-difluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 957) 5-amino-N-(3-chlorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 25 958) 5-amino-N-(2-chlorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 959) 5-amino-N-(3-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 960) 5-amino-3-(5-methyl-2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 30 961) 5-amino-N-(cyclopropylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 962) 5-amino-N-cycloheptyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 35 963) 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;

- 964) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 965) 5-amino-N-(1,2-dimethylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 5 966) 5-amino-N-(3,3-diphenylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 967) 5-amino-N-(1,2-diphenylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 968) 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 10 969) 5-amino-N-[3-(dibutylamino)propyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 970) 5-amino-N-butyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 15 971) 5-amino-3-(5-methyl-2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 972) 5-amino-3-(5-methyl-2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 973) 5-amino-3-(5-methyl-2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 20 974) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 975) 5-amino-N-(2-methoxy-1-methylethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 25 976) 5-amino-N-(1-methylbutyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 977) 5-amino-N-(3-fluorobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 978) 5-amino-N-(4-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 30 979) 5-amino-N-(1,3-dimethylbutyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 980) 5-amino-3-(5-methyl-2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 35 981) 5-amino-3-(5-methyl-2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;

- 982) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 983) 5-amino-3-(5-methyl-2-furyl)-N-pentylisoxazole-4-carboxamide;
- 5 984) 5-amino-N-(cyclohexylmethyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 985) ethyl 4-([5-amino-3-(5-methyl-2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 986) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 10 987) 5-amino-N-(1-ethylpropyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 988) 5-amino-3-(5-methyl-2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 15 989) 5-amino-N-(2-methylbenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 990) 5-amino-N-cyclobutyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 991) 5-amino-N-benzhydryl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 20 992) 5-amino-3-(5-methyl-2-furyl)-N-octylisoxazole-4-carboxamide;
- 993) 5-amino-N-(2-bromobenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 25 994) 5-amino-3-(5-methyl-2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 995) 5-amino-N-heptyl-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 996) 5-amino-N-(tert-butyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 30 997) 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 998) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-methyl-2-furyl)isoxazole-4-carboxamide;
- 35 999) 5-amino-3-(5-methyl-2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;

- 1000) 5-amino-N-benzyl-3-(2-furyl)isoxazole-4-carboxamide;
1001) 5-amino-3-(2-furyl)-N-(2-furylmethyl)isoxazole-4-carboxamide;
1002) 5-amino-3-(2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
5 1003) 5-amino-N-(sec-butyl)-3-(2-furyl)isoxazole-4-carboxamide;
1004) 5-amino-3-(2-furyl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
10 1005) 5-amino-3-(2-furyl)-N-hexylisoxazole-4-carboxamide;
1006) 5-amino-3-(2-furyl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
1007) 5-amino-3-(2-furyl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
15 1008) 5-amino-N-cyclopentyl-3-(2-furyl)isoxazole-4-carboxamide;
1009) 5-amino-N-(1,5-dimethylhexyl)-3-(2-furyl)isoxazole-4-carboxamide;
1010) 5-amino-3-(2-furyl)-N-isopentylisoxazole-4-carboxamide;
20 1011) 5-amino-N-(2-ethylhexyl)-3-(2-furyl)isoxazole-4-carboxamide;
1012) 5-amino-3-(2-furyl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
25 1013) 5-amino-3-(2-furyl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
1014) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(2-furyl)isoxazole-4-carboxamide;
1015) 5-amino-3-(2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
30 1016) 5-amino-3-(2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
1017) 5-amino-3-(2-furyl)-N-isobutylisoxazole-4-carboxamide;
1018) 5-amino-3-(2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
35

- 1019) 5-amino-3-(2-furyl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 1020) 5-amino-N-(2-fluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 5 1021) 5-amino-3-(2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1022) 5-amino-N-(3,4-difluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1023) 5-amino-N-(3-chlorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 10 1024) 5-amino-N-(2-chlorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1025) 5-amino-3-(2-furyl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 15 1026) 5-amino-3-(2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 1027) 5-amino-N-(cyclopropylmethyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1028) 5-amino-N-cycloheptyl-3-(2-furyl)isoxazole-4-carboxamide;
- 20 1029) 5-amino-3-(2-furyl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 1030) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 25 1031) 5-amino-N-(1,2-dimethylpropyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1032) 5-amino-N-(3,3-diphenylpropyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1033) 5-amino-N-(1,2-diphenylethyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 30 1034) 5-amino-N-(2,4-dimethoxybenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1035) 5-amino-N-[3-(dibutylamino)propyl]-3-(2-furyl)isoxazole-4-carboxamide;
- 35 1036) 5-amino-N-butyl-3-(2-furyl)isoxazole-4-carboxamide;

- 1037) 5-amino-3-(2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 1038) 5-amino-3-(2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 5 1039) 5-amino-3-(2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 1040) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1041) 5-amino-3-(2-furyl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;
- 10 1042) 5-amino-N-(4-tert-butylcyclohexyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1043) 5-amino-3-(2-furyl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 15 1044) 5-amino-N-(3-fluorobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1045) 5-amino-3-(2-furyl)-N-(4-methylbenzyl)isoxazole-4-carboxamide;
- 1046) 5-amino-N-(1,3-dimethylbutyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 20 1047) 5-amino-3-(2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 1048) 5-amino-3-(2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 25 1049) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1050) 5-amino-3-(2-furyl)-N-pentylisoxazole-4-carboxamide;
- 1051) 5-amino-N-(cyclohexylmethyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 30 1052) ethyl 4-([5-amino-3-(2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 1053) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(2-furyl)isoxazole-4-carboxamide;
- 1054) 5-amino-N-(1-ethylpropyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 35

- 1055) 5-amino-3-(2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 1056) 5-amino-3-(2-furyl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 5 1057) 5-amino-N-cyclobutyl-3-(2-furyl)isoxazole-4-carboxamide;
- 1058) 5-amino-N-benzhydryl-3-(2-furyl)isoxazole-4-carboxamide;
- 1059) 5-amino-3-(2-furyl)-N-octylisoxazole-4-carboxamide;
- 10 1060) 5-amino-N-(2-bromobenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1061) 5-amino-3-(2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 1062) 5-amino-3-(2-furyl)-N-heptylisoxazole-4-carboxamide;
- 15 1063) 5-amino-N-(tert-butyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1064) 5-amino-N-(3,4-dimethoxybenzyl)-3-(2-furyl)isoxazole-4-carboxamide;
- 1065) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-furyl)isoxazole-4-carboxamide;
- 20 1066) 5-amino-3-(2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 1067) 5-amino-3-(1-benzofuran-3-yl)-N-benzylisoxazole-4-carboxamide;
- 25 1068) 5-amino-3-(1-benzofuran-3-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide;
- 1069) 5-amino-3-(1-benzofuran-3-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 1070) 5-amino-3-(1-benzofuran-3-yl)-N-(sec-butyl)isoxazole-4-carboxamide;
- 30 1071) 5-amino-3-(1-benzofuran-3-yl)-N-cyclohexylisoxazole-4-carboxamide;
- 1072) 5-amino-3-(1-benzofuran-3-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 35 1073) 5-amino-3-(1-benzofuran-3-yl)-N-hexylisoxazole-4-carboxamide;

- 1074) 5-amino-3-(1-benzofuran-3-yl)-N-(tetrahydrofuran-2-ylmethyl)isoxazole-4-carboxamide;
- 1075) 5-amino-3-(1-benzofuran-3-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 5 1076) 5-amino-3-(1-benzofuran-3-yl)-N-cyclopentylisoxazole-4-carboxamide;
- 1077) 5-amino-3-(1-benzofuran-3-yl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide;
- 1078) 5-amino-3-(1-benzofuran-3-yl)-N-isopentylisoxazole-4-
10 carboxamide;
- 1079) 5-amino-3-(1-benzofuran-3-yl)-N-(2-ethylhexyl)isoxazole-4-carboxamide;
- 1080) 5-amino-3-(1-benzofuran-3-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 15 1081) 5-amino-3-(1-benzofuran-3-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 1082) 5-amino-3-(1-benzofuran-3-yl)-N-[3,5-bis(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1083) 5-amino-3-(1-benzofuran-3-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 20 1084) 5-amino-3-(1-benzofuran-3-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 1085) 5-amino-3-(1-benzofuran-3-yl)-N-isobutylisoxazole-4-carboxamide;
- 25 1086) 5-amino-3-(1-benzofuran-3-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1087) 5-amino-3-(1-benzofuran-3-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 1088) 5-amino-3-(1-benzofuran-3-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide;
- 30 1089) 5-amino-3-(1-benzofuran-3-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1090) 5-amino-3-(1-benzofuran-3-yl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide;
- 35 1091) 5-amino-3-(1-benzofuran-3-yl)-N-(3-chlorobenzyl)isoxazole-4-carboxamide;

- 1092) 5-amino-3-(1-benzofuran-3-yl)-N-(2-chlorobenzyl)isoxazole-4-carboxamide;
- 1093) 5-amino-3-(1-benzofuran-3-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 5 1094) 5-amino-3-(1-benzofuran-3-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 1095) 5-amino-3-(1-benzofuran-3-yl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide;
- 1096) 5-amino-3-(1-benzofuran-3-yl)-N-cycloheptylisoxazole-
10 4-carboxamide;
- 1097) 5-amino-3-(1-benzofuran-3-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 1098) 5-amino-3-(1-benzofuran-3-yl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide;
- 15 1099) 5-amino-3-(1-benzofuran-3-yl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide;
- 1100) 5-amino-3-(1-benzofuran-3-yl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide;
- 1101) 5-amino-3-(1-benzofuran-3-yl)-N-(1,2-
20 diphenylethyl)isoxazole-4-carboxamide;
- 1102) 5-amino-3-(1-benzofuran-3-yl)-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 1103) 5-amino-3-(1-benzofuran-3-yl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide;
- 25 1104) 5-amino-3-(1-benzofuran-3-yl)-N-butylisoxazole-4-carboxamide;
- 1105) 5-amino-3-(1-benzofuran-3-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 1106) 5-amino-3-(1-benzofuran-3-yl)-N-[(5-methyl-2-
30 furyl)methyl]isoxazole-4-carboxamide;
- 1107) 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 1108) 5-amino-3-(1-benzofuran-3-yl)-N-(1-benzylpiperidin-4-yl)isoxazole-4-carboxamide;
- 35 1109) 5-amino-3-(1-benzofuran-3-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;

- 1110) 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 1111) 5-amino-3-(1-benzofuran-3-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide;
- 5 1112) 5-amino-3-(1-benzofuran-3-yl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide;
- 1113) 5-amino-3-(1-benzofuran-3-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 1114) 5-amino-3-(1-benzofuran-3-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 10 1115) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(1-benzofuran-3-yl)isoxazole-4-carboxamide;
- 1116) 5-amino-3-(1-benzofuran-3-yl)-N-pentylisoxazole-4-carboxamide;
- 15 1117) 5-amino-3-(1-benzofuran-3-yl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide;
- 1118) ethyl 4-([5-amino-3-(1-benzofuran-3-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 1119) 5-amino-3-(1-benzofuran-3-yl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide;
- 20 1120) 5-amino-3-(1-benzofuran-3-yl)-N-(1-ethylpropyl)isoxazole-4-carboxamide;
- 1121) 5-amino-3-(1-benzofuran-3-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 25 1122) 5-amino-3-(1-benzofuran-3-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 1123) 5-amino-3-(1-benzofuran-3-yl)-N-cyclobutylisoxazole-4-carboxamide;
- 1124) 5-amino-N-benzhydryl-3-(1-benzofuran-3-yl)isoxazole-4-carboxamide;
- 30 1125) 5-amino-3-(1-benzofuran-3-yl)-N-octylisoxazole-4-carboxamide;
- 1126) 5-amino-3-(1-benzofuran-3-yl)-N-(2-bromobenzyl)isoxazole-4-carboxamide;
- 35 1127) 5-amino-3-(1-benzofuran-3-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide;

- 1128) 5-amino-3-(1-benzofuran-3-yl)-N-heptylisoxazole-4-carboxamide;
- 1129) 5-amino-3-(1-benzofuran-3-yl)-N-(tert-butyl)isoxazole-4-carboxamide;
- 5 1130) 5-amino-3-(1-benzofuran-3-yl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 1131) 5-amino-3-(1-benzofuran-3-yl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide;
- 1132) 5-amino-3-(1-benzofuran-3-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 10 1133) 5-amino-N-benzyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1134) 5-amino-3-(5-ethylthien-2-yl)-N-(2-furylmethyl)isoxazole-4-carboxamide;
- 15 1135) 5-amino-3-(5-ethylthien-2-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 1136) 5-amino-N-(sec-butyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1137) 5-amino-N-cyclohexyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 20 1138) 5-amino-3-(5-ethylthien-2-yl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 1139) 5-amino-3-(5-ethylthien-2-yl)-N-hexylisoxazole-4-carboxamide;
- 25 1140) 5-amino-3-(5-ethylthien-2-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 1141) 5-amino-N-cyclopentyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1142) 5-amino-N-(1,5-dimethylhexyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 30 1143) 5-amino-3-(5-ethylthien-2-yl)-N-isopentylisoxazole-4-carboxamide;
- 1144) 5-amino-N-(2-ethylhexyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 35 1145) 5-amino-3-(5-ethylthien-2-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;

- 1146) 5-amino-3-(5-ethylthien-2-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 1147) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 5 1148) 5-amino-3-(5-ethylthien-2-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1149) 5-amino-3-(5-ethylthien-2-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 1150) 5-amino-3-(5-ethylthien-2-yl)-N-isobutylisoxazole-4-
10 carboxamide;
- 1151) 5-amino-3-(5-ethylthien-2-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1152) 5-amino-3-(5-ethylthien-2-yl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 15 1153) 5-amino-3-(5-ethylthien-2-yl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide;
- 1154) 5-amino-N-(3,4-difluorobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1155) 5-amino-N-(3-chlorobenzyl)-3-(5-ethylthien-2-
20 yl)isoxazole-4-carboxamide;
- 1156) 5-amino-N-(2-chlorobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1157) 5-amino-3-(5-ethylthien-2-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 25 1158) 5-amino-3-(5-ethylthien-2-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 1159) 5-amino-N-(cyclopropylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1160) 5-amino-N-cycloheptyl-3-(5-ethylthien-2-yl)isoxazole-
30 4-carboxamide;
- 1161) 5-amino-3-(5-ethylthien-2-yl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 1162) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 35 1163) 5-amino-N-(1,2-dimethylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;

- 1164) 5-amino-N-(3,3-diphenylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1165) 5-amino-N-(1,2-diphenylethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 5 1166) 5-amino-N-(2,4-dimethoxybenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1167) 5-amino-N-[3-(dibutylamino)propyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1168) 5-amino-N-butyl-3-(5-ethylthien-2-yl)isoxazole-4-
10 carboxamide;
- 1169) 5-amino-3-(5-ethylthien-2-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 1170) 5-amino-3-(5-ethylthien-2-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 15 1171) 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 1172) 5-amino-3-(5-ethylthien-2-yl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;
- 1173) 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-ethylthien-2-
20 yl)isoxazole-4-carboxamide;
- 1174) 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 1175) 5-amino-3-(5-ethylthien-2-yl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide;
- 25 1176) 5-amino-3-(5-ethylthien-2-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide;
- 1177) 5-amino-N-(1,3-dimethylbutyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1178) 5-amino-3-(5-ethylthien-2-yl)-N-(1-methyl-3-
30 phenylpropyl)isoxazole-4-carboxamide;
- 1179) 5-amino-3-(5-ethylthien-2-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 1180) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 35 1181) 5-amino-3-(5-ethylthien-2-yl)-N-pentylisoxazole-4-carboxamide;

- 1182) 5-amino-N-(cyclohexylmethyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1183) ethyl 4-([5-amino-3-(5-ethylthien-2-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 5 1184) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1185) 5-amino-N-(1-ethylpropyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1186) 5-amino-3-(5-ethylthien-2-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 10 1187) 5-amino-3-(5-ethylthien-2-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 1188) 5-amino-N-cyclobutyl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 15 1189) 5-amino-N-benzhydryl-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1190) 5-amino-3-(5-ethylthien-2-yl)-N-octylisoxazole-4-carboxamide;
- 1191) 5-amino-N-(2-bromobenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 20 1192) 5-amino-3-(5-ethylthien-2-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 1193) 5-amino-3-(5-ethylthien-2-yl)-N-heptylisoxazole-4-carboxamide;
- 25 1194) 5-amino-N-(tert-butyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1195) 5-amino-N-(3,4-dimethoxybenzyl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 1196) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-ethylthien-2-yl)isoxazole-4-carboxamide;
- 30 1197) 5-amino-3-(5-ethylthien-2-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 1198) 5-amino-N-benzyl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 35 1199) 5-amino-3-(5-chloro-2-furyl)-N-(2-furylmethyl)isoxazole-4-carboxamide;

- 1200) 5-amino-3-(5-chloro-2-furyl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 1201) 5-amino-N-(sec-butyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 5 1202) 5-amino-3-(5-chloro-2-furyl)-N-cyclohexylisoxazole-4-carboxamide;
- 1203) 5-amino-3-(5-chloro-2-furyl)-N-(4-methoxybenzyl)isoxazole-4-carboxamide;
- 1204) 5-amino-3-(5-chloro-2-furyl)-N-hexylisoxazole-4-
- 10 carboxamide;
- 1205) 5-amino-3-(5-chloro-2-furyl)-N-cyclopentylisoxazole-4-carboxamide;
- 1206) 5-amino-3-(5-chloro-2-furyl)-N-(1,5-dimethylhexyl)isoxazole-4-carboxamide;
- 15 1207) 5-amino-3-(5-chloro-2-furyl)-N-isopentylisoxazole-4-carboxamide;
- 1208) 5-amino-3-(5-chloro-2-furyl)-N-(2-ethylhexyl)isoxazole-4-carboxamide;
- 1209) 5-amino-3-(5-chloro-2-furyl)-N-(4-
- 20 phenylbutyl)isoxazole-4-carboxamide;
- 1210) 5-amino-3-(5-chloro-2-furyl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 1211) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 25 1212) 5-amino-3-(5-chloro-2-furyl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1213) 5-amino-3-(5-chloro-2-furyl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 1214) 5-amino-3-(5-chloro-2-furyl)-N-isobutylisoxazole-4-
- 30 carboxamide;
- 1215) 5-amino-3-(5-chloro-2-furyl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1216) 5-amino-3-(5-chloro-2-furyl)-N-(2-methoxybenzyl)isoxazole-4-carboxamide;
- 35 1217) 5-amino-3-(5-chloro-2-furyl)-N-(2-fluorobenzyl)isoxazole-4-carboxamide;

- 1218) 5-amino-3-(5-chloro-2-furyl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1219) 5-amino-3-(5-chloro-2-furyl)-N-(3,4-difluorobenzyl)isoxazole-4-carboxamide;
- 5 1220) 5-amino-N-(3-chlorobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1221) 5-amino-N-(2-chlorobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1222) 5-amino-3-(5-chloro-2-furyl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;
- 10 1223) 5-amino-3-(5-chloro-2-furyl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 1224) 5-amino-3-(5-chloro-2-furyl)-N-(cyclopropylmethyl)isoxazole-4-carboxamide;
- 15 1225) 5-amino-3-(5-chloro-2-furyl)-N-cycloheptylisoxazole-4-carboxamide;
- 1226) 5-amino-3-(5-chloro-2-furyl)-N-[2-(4-methoxyphenyl)ethyl]isoxazole-4-carboxamide;
- 1227) 5-amino-3-(5-chloro-2-furyl)-N-(2-cyclohex-1-en-1-ylethyl)isoxazole-4-carboxamide;
- 20 1228) 5-amino-3-(5-chloro-2-furyl)-N-(1,2-dimethylpropyl)isoxazole-4-carboxamide;
- 1229) 5-amino-3-(5-chloro-2-furyl)-N-(3,3-diphenylpropyl)isoxazole-4-carboxamide;
- 25 1230) 5-amino-3-(5-chloro-2-furyl)-N-(1,2-diphenylethyl)isoxazole-4-carboxamide;
- 1231) 5-amino-3-(5-chloro-2-furyl)-N-(2,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 1232) 5-amino-3-(5-chloro-2-furyl)-N-[3-(dibutylamino)propyl]isoxazole-4-carboxamide;
- 30 1233) 5-amino-N-butyl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1234) 5-amino-3-(5-chloro-2-furyl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 35 1235) 5-amino-3-(5-chloro-2-furyl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;

- 1236) 5-amino-3-(5-chloro-2-furyl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 1237) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 5 1238) 5-amino-3-(5-chloro-2-furyl)-N-(2-methoxy-1-methylethyl)isoxazole-4-carboxamide;
- 1239) 5-amino-N-(4-tert-butylcyclohexyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1240) 5-amino-3-(5-chloro-2-furyl)-N-(1-methylbutyl)isoxazole-4-carboxamide;
- 10 1241) 5-amino-3-(5-chloro-2-furyl)-N-(3-fluorobenzyl)isoxazole-4-carboxamide;
- 1242) 5-amino-3-(5-chloro-2-furyl)-N-(4-methylbenzyl)isoxazole-4-carboxamide;
- 15 1243) 5-amino-3-(5-chloro-2-furyl)-N-(1,3-dimethylbutyl)isoxazole-4-carboxamide;
- 1244) 5-amino-3-(5-chloro-2-furyl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 1245) 5-amino-3-(5-chloro-2-furyl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 20 1246) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1247) 5-amino-3-(5-chloro-2-furyl)-N-pentylisoxazole-4-carboxamide;
- 25 1248) 5-amino-3-(5-chloro-2-furyl)-N-(cyclohexylmethyl)isoxazole-4-carboxamide;
- 1249) ethyl 4-([5-amino-3-(5-chloro-2-furyl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 1250) 5-amino-3-(5-chloro-2-furyl)-N-[2-(3-chlorophenyl)ethyl]isoxazole-4-carboxamide;
- 30 1251) 5-amino-3-(5-chloro-2-furyl)-N-(1-ethylpropyl)isoxazole-4-carboxamide;
- 1252) 5-amino-3-(5-chloro-2-furyl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 35 1253) 5-amino-3-(5-chloro-2-furyl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;

- 1254) 5-amino-3-(5-chloro-2-furyl)-N-cyclobutylisoxazole-4-carboxamide;
- 1255) 5-amino-N-benzhydryl-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 5 1256) 5-amino-3-(5-chloro-2-furyl)-N-octylisoxazole-4-carboxamide;
- 1257) 5-amino-N-(2-bromobenzyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 1258) 5-amino-3-(5-chloro-2-furyl)-N-(1-methylheptyl)isoxazole-4-carboxamide;
- 10 1259) 5-amino-3-(5-chloro-2-furyl)-N-heptylisoxazole-4-carboxamide;
- 1260) 5-amino-N-(tert-butyl)-3-(5-chloro-2-furyl)isoxazole-4-carboxamide;
- 15 1261) 5-amino-3-(5-chloro-2-furyl)-N-(3,4-dimethoxybenzyl)isoxazole-4-carboxamide;
- 1262) 5-amino-3-(5-chloro-2-furyl)-N-(2,3-dihydro-1H-inden-1-yl)isoxazole-4-carboxamide;
- 1263) 5-amino-3-(5-chloro-2-furyl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 20 1264) 5-amino-N-benzyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1265) 5-amino-N-(2-furylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 25 1266) 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-phenylethyl)isoxazole-4-carboxamide;
- 1267) 5-amino-N-(sec-butyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1268) 5-amino-N-cyclohexyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 30 1269) 5-amino-N-(4-methoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1270) 5-amino-N-hexyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 35 1271) 5-amino-3-(6-methoxypyridin-3-yl)-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;

- 236 -

- 1272) 5-amino-N-cyclopentyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1273) 5-amino-N-(1,5-dimethylhexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 5 1274) 5-amino-N-isopentyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1275) 5-amino-N-(2-ethylhexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1276) 5-amino-3-(6-methoxypyridin-3-yl)-N-(4-phenylbutyl)isoxazole-4-carboxamide;
- 10 1277) 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-methylcyclohexyl)isoxazole-4-carboxamide;
- 1278) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 15 1279) 5-amino-3-(6-methoxypyridin-3-yl)-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1280) 5-amino-3-(6-methoxypyridin-3-yl)-N-(3-phenylpropyl)isoxazole-4-carboxamide;
- 1281) 5-amino-N-isobutyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 20 1282) 5-amino-3-(6-methoxypyridin-3-yl)-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1283) 5-amino-N-(2-methoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 25 1284) 5-amino-N-(2-fluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1285) 5-amino-3-(6-methoxypyridin-3-yl)-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1286) 5-amino-N-(3,4-difluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 30 1287) 5-amino-N-(3-chlorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1288) 5-amino-N-(2-chlorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 35 1289) 5-amino-3-(6-methoxypyridin-3-yl)-N-(3-methylbenzyl)isoxazole-4-carboxamide;

- 1290) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-naphthylmethyl)isoxazole-4-carboxamide;
- 1291) 5-amino-N-(cyclopropylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 5 1292) 5-amino-N-cycloheptyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1293) 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1294) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 10 1295) 5-amino-N-(1,2-dimethylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1296) 5-amino-N-(3,3-diphenylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 15 1297) 5-amino-N-(1,2-diphenylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1298) 5-amino-N-(2,4-dimethoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1299) 5-amino-N-[3-(dibutylamino)propyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 20 1300) 5-amino-N-butyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1301) 5-amino-3-(6-methoxypyridin-3-yl)-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 25 1302) 5-amino-3-(6-methoxypyridin-3-yl)-N-[(5-methyl-2-furyl)methyl]isoxazole-4-carboxamide;
- 1303) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylhexyl)isoxazole-4-carboxamide;
- 1304) 5-amino-N-(1-benzylpiperidin-4-yl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 30 1305) 5-amino-N-(2-methoxy-1-methylethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1306) 5-amino-N-(4-tert-butylcyclohexyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 35 1307) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylbutyl)isoxazole-4-carboxamide;

- 1308) 5-amino-N-(3-fluorobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1309) 5-amino-3-(6-methoxypyridin-3-yl)-N-(4-methylbenzyl)isoxazole-4-carboxamide;
- 5 1310) 5-amino-N-(1,3-dimethylbutyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1311) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methyl-3-phenylpropyl)isoxazole-4-carboxamide;
- 1312) 5-amino-3-(6-methoxypyridin-3-yl)-N-[3-(methylthio)propyl]isoxazole-4-carboxamide;
- 10 1313) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1314) 5-amino-3-(6-methoxypyridin-3-yl)-N-pentylisoxazole-4-carboxamide;
- 15 1315) 5-amino-N-(cyclohexylmethyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1316) ethyl 4-([5-amino-3-(6-methoxypyridin-3-yl)isoxazol-4-yl]carbonyl)amino)piperidine-1-carboxylate;
- 1317) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 20 1318) 5-amino-N-(1-ethylpropyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1319) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-phenylethyl)isoxazole-4-carboxamide;
- 25 1320) 5-amino-3-(6-methoxypyridin-3-yl)-N-(2-methylbenzyl)isoxazole-4-carboxamide;
- 1321) 5-amino-N-cyclobutyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1322) 5-amino-N-benzhydryl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 30 1323) 5-amino-3-(6-methoxypyridin-3-yl)-N-octylisoxazole-4-carboxamide;
- 1324) 5-amino-N-(2-bromobenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 35 1325) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1-methylheptyl)isoxazole-4-carboxamide;

- 1326) 5-amino-N-heptyl-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1327) 5-amino-N-(tert-butyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 5 1328) 5-amino-N-(3,4-dimethoxybenzyl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1329) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-(6-methoxypyridin-3-yl)isoxazole-4-carboxamide;
- 1330) 5-amino-3-(6-methoxypyridin-3-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide;
- 10 1331) 5-amino-N-benzyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1332) 5-amino-N-(2-furylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 15 1333) 5-amino-N-(2-phenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1334) 5-amino-N-cyclohexyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1335) 5-amino-N-(4-methoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 20 1336) 5-amino-N-hexyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1337) 5-amino-3-pyridin-3-yl-N-(thien-2-ylmethyl)isoxazole-4-carboxamide;
- 1338) 5-amino-N-cyclopentyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 25 1339) 5-amino-N-(1,5-dimethylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1340) 5-amino-N-isopentyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 30 1341) 5-amino-N-(2-ethylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1342) 5-amino-N-(4-phenylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1343) 5-amino-N-(2-methylcyclohexyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 35

- 1344) 5-amino-N-[3,5-bis(trifluoromethyl)benzyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1345) 5-amino-3-pyridin-3-yl-N-[3-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 5 1346) 5-amino-N-(3-phenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1347) 5-amino-N-isobutyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1348) 5-amino-3-pyridin-3-yl-N-[2-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 10 1349) 5-amino-N-(2-methoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1350) 5-amino-N-(2-fluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 15 1351) 5-amino-3-pyridin-3-yl-N-[4-(trifluoromethyl)benzyl]isoxazole-4-carboxamide;
- 1352) 5-amino-N-(3,4-difluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1353) 5-amino-N-(3-chlorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 20 1354) 5-amino-N-(2-chlorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1355) 5-amino-N-(3-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 25 1356) 5-amino-N-(1-naphthylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1357) 5-amino-N-(cyclopropylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1358) 5-amino-N-cycloheptyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 30 1359) 5-amino-N-[2-(4-methoxyphenyl)ethyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1360) 5-amino-N-(2-cyclohex-1-en-1-ylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 35 1361) 5-amino-N-(1,2-dimethylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide;



- 1362) 5-amino-N-(3,3-diphenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1363) 5-amino-N-(1,2-diphenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 5 1364) 5-amino-N-(2,4-dimethoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1365) 5-amino-N-[3-(dibutylamino)propyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1366) 5-amino-N-butyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 10 1367) 5-amino-3-pyridin-3-yl-N-[4-(trifluoromethoxy)benzyl]isoxazole-4-carboxamide;
- 1368) 5-amino-N-[(5-methyl-2-furyl)methyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1369) 5-amino-N-(1-methylhexyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 15 1370) 5-amino-N-(1-methylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1371) 5-amino-N-(3-fluorobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 20 1372) 5-amino-N-(4-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1373) 5-amino-N-(1,3-dimethylbutyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1374) 5-amino-N-(1-methyl-3-phenylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 25 1375) 5-amino-N-[3-(methylthio)propyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1376) 5-amino-N-(1,3-benzodioxol-5-ylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 30 1377) 5-amino-N-pentyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1378) 5-amino-N-(cyclohexylmethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1379) ethyl 4-{[(5-amino-3-pyridin-3-ylisoxazol-4-yl)carbonyl]amino}piperidine-1-carboxylate;
- 35

- 242 -


- 1380) 5-amino-N-[2-(3-chlorophenyl)ethyl]-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1381) 5-amino-N-(1-ethylpropyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 5 1382) 5-amino-N-(1-phenylethyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1383) 5-amino-N-(2-methylbenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1384) 5-amino-N-cyclobutyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 10 1385) 5-amino-N-benzhydryl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1386) 5-amino-N-octyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1387) 5-amino-N-(2-bromobenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 15 1388) 5-amino-N-(1-methylheptyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1389) 5-amino-N-heptyl-3-pyridin-3-ylisoxazole-4-carboxamide;
- 20 1390) 5-amino-N-(tert-butyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1391) 5-amino-N-(3,4-dimethoxybenzyl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 1392) 5-amino-N-(2,3-dihydro-1H-inden-1-yl)-3-pyridin-3-ylisoxazole-4-carboxamide;
- 25 1393) 5-amino-3-pyridin-3-yl-N-(1,2,3,4-tetrahydronaphthalen-1-yl)isoxazole-4-carboxamide.

24. A process for preparing the compounds of formula (I)
- 30 and the pharmaceutically acceptable salts, as defined in claim 13, which process comprises:
- a) reacting a compound of formula (II) with a compound of formula (III)

- 243 -




wherein R, R₁ and X are as above defined and Z represents a halogen atom or a suitable leaving group, so as to obtain a compound of formula (I)



5

and, optionally, reacting the compound of formula (I) according to any one of the alternative steps b) below

b.1) with a compound of formula (IV), (V) or (VI)



10 wherein R₂ is as above defined, W is hydroxy or a suitable leaving group and W' is a suitable leaving group, so as to obtain the compounds of formula (I) wherein Y is -CO-, -SO₂- or -CONH-, respectively, and R, R₁, R₂ and X are as above defined;

15 b.2) with a suitable aldehyde or ketone derivative of formula (VII)



wherein R₂ and R₃ are as above defined, under reductive conditions, so as to obtain a compound of formula (I)

20 wherein Y is a group -CH(R₃)- and R, R₁, R₂, R₃ and X are as above defined;

b.3) with a suitable acylating agent in the presence of ammonia, so as to obtain a compound of formula (I) wherein Y is -CONH-, R₂ is hydrogen and R, R₁ and X are as above defined; and, optionally

25

c) converting the thus obtained compound of formula (I) into another compound of formula (I) and/or into a pharmaceutically acceptable salt thereof.

25. The process of claim 24 wherein, within the compound of formula (IV) or (V) in step b.1), W is hydroxy or a chlorine or bromine atom and W' is a chlorine or bromine atom.

26. The process of claim 24 wherein, within step b.2), reductive conditions occur in the presence of sodium borohydride, sodium cyanoborohydride or sodium triacetoxymethylborohydride.


27. The process of claim 24 wherein, in step b.3), the acylating agent is selected from triphosgene or trichloromethyl chloroformate.

28. A process under solid phase synthesis (SPS) conditions for preparing the compounds of formula (I) and the pharmaceutically acceptable salts, as defined in claim 13, which process comprises:

a') reacting cyanoacetic acid with a suitable polystyrenic resin of formula (X) or (XI)




wherein R_1 is as above defined and W is hydroxy or a suitable leaving group, so as to obtain the compounds of formula (XII) or (XIII), respectively





wherein R_1 is as above reported;

b') reacting any one of the compounds of formula (XII) or (XIII) with a compound of formula (II)


- 245 -





wherein R and Z are as above defined, so as to obtain a compound of formula (XIV) or (XV)



- 5 c') cleaving the resin from the compounds of formula (XIV) or (XV) under acidic or basic conditions, so as to obtain a compound of formula (I) wherein X is a group -O- and R₁ is hydrogen, or X is a group -N(R₃)- wherein R₃ is hydrogen and R₁ is as above defined, respectively; or, alternatively,
- 10 reacting the above compounds of formula (XIV) or (XV) according to any one of the alternative steps d') below
- d.1') with a compound of formula (IV), (V) or (VI)



- wherein R₂ is as above defined, W is hydroxy or a suitable
- 15 leaving group and W' is a suitable leaving group, so as to obtain the compounds of formula (XVI) or (XVII)



wherein Y is -CO-, -SO₂- or -CONH-, respectively, and R, R₁ and R₂ are as above defined;

- 20 d.2') with a suitable aldehyde or ketone derivative of formula (VII)




wherein R_2 and R_3 are as above defined, under reductive conditions, so as to obtain the compounds of the above formula (XVI) or (XVII) wherein Y is a group $-\text{CH}(R_3)-$ and R , R_1 , R_2 and R_3 are as above defined;

5 d.3') with a suitable acylating agent in the presence of ammonia, so as to obtain a compound of the above formula (XVI) or (XVII) wherein Y is $-\text{CONH}-$, R_2 is hydrogen and R and R_1 are as above defined;

e') cleaving the resin from the compounds of formula (XVI) or (XVII) under acidic or basic conditions, so as to obtain
10 a compound of formula (I) wherein X is a group $-\text{O}-$ and R_1 is hydrogen, or X is a group $-\text{N}(R_3)-$ wherein R_3 is hydrogen and R_1 is as above defined, respectively; and, optionally,

f') converting the thus obtained compound of formula (I)
15 into another compound of formula (I) and/or into a pharmaceutically acceptable salt thereof.

29. A library of two or more compounds of formula (I)



20 wherein

R is an optionally substituted 5 or 6 membered heteroaryl group with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur, optionally further condensed with a 5 to 7 membered aromatic or non-aromatic
25 carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or sulfur;

X is a divalent group selected from $-\text{N}(R_3)-$ or $-\text{O}-$;

Y is a divalent group selected from $-\text{CH}(R_3)-$, $-\text{CO}-$, $-\text{CONH}-$ or $-\text{SO}_2-$, or Y may also be a single bond when R_2 is a
30 hydrogen atom or a C_3 - C_6 cycloalkyl group;

R₁ is a hydrogen atom or a group, optionally further substituted, selected from straight or branched C₁-C₆ alkyl, C₃-C₆ cycloalkyl, aryl or aryl C₁-C₆ alkyl, 5 or 6 membered heterocyclyl or heterocyclyl C₁-C₆ alkyl having from 1 to 3
5 heteroatoms selected among nitrogen, oxygen or sulfur; the said cycloalkyl, aryl or heterocyclyl groups being optionally further condensed with a 5 to 7 membered aromatic or non-aromatic carbocycle or heterocycle with from 1 to 3 heteroatoms selected among nitrogen, oxygen or
10 sulfur;

R₂ and R₃ have, each independently, the meanings above reported for R₁ or represent an optionally substituted straight or branched C₂-C₆ alkenyl or alkynyl group; and the pharmaceutically acceptable salts thereof.

15

30. Any specific compound of formula (I), as defined in claim 13, which is obtainable, for instance through a combinatorial chemistry technique as per the process of claim 28, by reacting each of the derivatives of formula
20 (II), as set forth in table I, with any one of the derivatives of formula (XIII) which are obtainable from the amines of formula (IX), as set forth in table II, and by subsequently reacting any one of the obtained derivatives of formula (XV) with any one of the derivatives of formula
25 (IV), as set forth in table III.

31. Any specific compound of formula (I), as defined in claim 13, which is obtainable, for instance through a combinatorial chemistry technique as per the process of
30 claim 28, by reacting each of the derivatives of formula (II), as set forth in table I, with any one of the derivatives of formula (XIII) which are obtainable from the amines of formula (IX), as set forth in table II, and by subsequently reacting any one of the obtained derivatives

of formula (XV) with any one of the derivatives of formula (V), as set forth in table IV.

5 32. Any specific compound of formula (I), as defined in claim 13, which is obtainable, for instance through a combinatorial chemistry technique as per the process of claim 28, by reacting each of the derivatives of formula (II), as set forth in table I, with any one of the derivatives of formula (XIII) which are obtainable from the
10 amines of formula (IX), as set forth in table II, and by subsequently reacting any one of the obtained derivatives of formula (XV) with any one of the derivatives of formula (VI), as set forth in table V.

15 33. A pharmaceutical composition comprising an effective amount of an aminoisoxazole of formula (I) as defined in claim 13 and, at least, one pharmaceutically acceptable excipient, carrier or diluent.

20 34. A pharmaceutical composition according to claim 33 further comprising one or more chemotherapeutic agents, as a combined preparation for simultaneous, separate or sequential use in anticancer therapy.

25 35. A product or kit comprising a compound of claim 13 or a pharmaceutical composition thereof as defined in claim 34, and one or more chemotherapeutic agents, as a combined preparation for simultaneous, separate or sequential use in anticancer therapy.

30

36. A compound of formula (I) or a pharmaceutically acceptable salt thereof, as defined in claim 13, for use as a medicament.

- 249 -

37. Use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, as defined in claim 13, in the manufacture of a medicament for treating diseases caused by and/or associated with an altered protein kinase activity.

5

38. Use according to claim 37 for treating tumors.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08634

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/422 A61K31/4439 A61K31/427 A61P25/28 A61P35/00
C07D413/04 C07D417/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 22610 A (JANETKA JAMES ;MASHAL ROBERT (US); VERTEX PHARMA (US); CAO JINGRON) 21 March 2002 (2002-03-21) Page 10, line 25 the linker group can be -C(O)NH- claim 1	1-38
A	WO 01 12621 A (BAKER CHRISTOPHER ;HARRINGTON EDMUND (US); BEMIS GUY (US); LEDEBOE) 22 February 2001 (2001-02-22) G is aryl where aryl is as defined on page 11, lines 3-13. claim 1	1-38
A	GB 1 250 219 A (J.R.GEIGY A.G.) 20 October 1971 (1971-10-20) cited in the application claim 1	13



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

3 December 2002

Date of mailing of the international search report

13/12/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08634

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 631 169 A (MINAMI SHINSAKU ET AL) 28 December 1971 (1971-12-28) cited in the application claim 1 -----	13
A	US 3 562 267 A (HOYLE WILLIAM ET AL) 9 February 1971 (1971-02-09) claim 1 -----	13
A	US 3 522 252 A (LEMIEUX RAYMOND URGEL ET AL) 28 July 1970 (1970-07-28) claim 1 -----	13
A	GB 1 162 257 A (DAINIPPON) 20 August 1969 (1969-08-20) claim 1 -----	13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08634

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0222610	A	21-03-2002	AU 9094001 A WO 0222610 A1	26-03-2002 21-03-2002
WO 0112621	A	22-02-2001	AU 6909600 A CN 1378541 T CZ 20020534 A3 EP 1218369 A1 NO 20020713 A SK 3572002 A3 WO 0112621 A1	13-03-2001 06-11-2002 17-07-2002 03-07-2002 12-04-2002 02-07-2002 22-02-2001
GB 1250219	A	20-10-1971	NONE	
US 3631169	A	28-12-1971	NONE	
US 3562267	A	09-02-1971	GB 1108397 A BE 685636 A CH 475270 A DE 1695085 A1 DK 118132 B FI 45045 B FR 1490038 A FR 5947 M IL 26353 A NL 6611584 A NO 121097 B SE 310679 B AT 265248 B AT 260918 B	03-04-1968 17-02-1967 15-07-1969 16-03-1972 13-07-1970 30-11-1971 28-07-1967 08-04-1968 19-07-1970 20-02-1967 18-01-1971 12-05-1969 10-10-1968 25-03-1968
US 3522252	A	28-07-1970	NONE	
GB 1162257	A	20-08-1969	DE 1670534 A1 FR 6916 M	12-11-1970 05-05-1969