



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C12N 15/89, 15/90, 15/63, 15/62, 15/85, A01K 67/027, C07K 14/75 // 14/47</p>	A1	<p>(11) International Publication Number: <b>WO 95/23868</b></p> <p>(43) International Publication Date: 8 September 1995 (08.09.95)</p>
<p>(21) International Application Number: PCT/US95/02648</p> <p>(22) International Filing Date: 1 March 1995 (01.03.95)</p> <p>(30) Priority Data: 08/206,176 3 March 1994 (03.03.94) US</p> <p>(71) Applicants: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US). PHARMACEUTICAL PROTEINS LTD. [GB/GB]; Roslin, Edinburgh, Midlothian EH25 9PP (GB).</p> <p>(72) Inventors: GARNER, Ian; 13 Lismore Avenue, Edinburgh EH8 7DW (GB). DALRYMPLE, Michael, A.; 21 North Fort Street, Edinburgh EH6 4HB (GB). PRUNKARD, Donna, E.; 3200 NW 65th Street #201, Seattle, WA 98117 (US). FOSTER, Donald, C.; 3002 NE 181st Street, Seattle, WA 98155 (US).</p> <p>(74) Agent: PARKER, Gary, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</p>		<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PRODUCTION OF FIBRINOGEN IN TRANSGENIC ANIMALS</p>		
<p>(57) Abstract</p> <p>Materials and methods for producing fibrinogen in transgenic non-human mammals are disclosed. DNA segments encoding A<math>\alpha</math>, B<math>\beta</math> and <math>\gamma</math> chains of fibrinogen are introduced into the germ line of a non-human mammal, and the mammal or its female progeny produces milk containing fibrinogen expressed from the introduced DNA segments. Non-human mammalian embryos and transgenic non-human mammals carrying DNA segments encoding heterologous fibrinogen polypeptide chains are also disclosed.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Description

## 5 PRODUCTION OF FIBRINOGEN IN TRANSGENIC ANIMALS

Background of the Invention

10 The final step in the blood coagulation cascade is the thrombin-catalyzed conversion of the soluble plasma protein fibrinogen to insoluble fibrin. Thrombin cleaves a small peptide (fibrinopeptide A) from one of the three component chains (the A $\alpha$ -chain) of fibrinogen. Fibrin monomers subsequently polymerize and are cross-linked by activated factor XIII to form a stable clot.

15 Fibrinogen is a key component of biological tissue glues (see, e.g., U.S. Patents Nos. 4,377,572 and 4,442,655), which mimic the formation of natural blood clots to promote hemostasis and repair damaged tissue. Tissue glues provide an adjunct or alternative to sutures, 20 staples and other mechanical means for wound closure. However, the principal ingredients of these products (fibrinogen, factor XIII and thrombin) are prepared from pooled human plasma by cryoprecipitation (e.g. U.S. Patents No. 4,377,572; 4,362,567; 4,909,251) or ethanol 25 precipitation (e.g. U.S. Patent No. 4,442,655) or from single donor plasma (e.g. U.S. Patent No. 4,627,879; Spotnitz et al., Am. Surg. 55: 166-168, 1989). The resultant fibrinogen/factor XIII preparation is mixed with bovine thrombin immediately before use to convert the 30 fibrinogen to fibrin and activate the factor XIII, thus initiating coagulation of the adhesive.

Commercially available adhesives are of pooled plasma origin. Because blood-derived products have been associated with the transmission of human immunodeficiency 35 virus (HIV), hepatitis virus and other etiologic agents, the acceptance and availability of such adhesives is

limited. At present they are not approved for use in the United States.

While the use of autologous plasma reduces the risk of disease transmission, autologous adhesives can only be used in elective surgery when the patient is able to donate the necessary blood in advance.

As noted above, fibrinogen consists of three polypeptide chains, each of which is present in two copies in the assembled molecule. These chains, designated the  $A\alpha$ ,  $B\beta$  and  $\gamma$ -chains, are coordinately expressed, assembled and secreted by the liver. While it might be expected that recombinant DNA technology could provide an alternative to the isolation of fibrinogen from plasma, this goal has proven to be elusive. The three fibrinogen chains have been individually expressed in *E. coli* (Lord, DNA 4: 33-38, 1985; Bolyard and Lord, Gene 66: 183-192, 1988; Bolyard and Lord, Blood 73: 1202-1206), but functional fibrinogen has not been produced in a prokaryotic system. Expression of biologically competent fibrinogen in yeast has not been reported. Cultured transfected mammalian cells have been used to express biologically active fibrinogen (Farrell et al., Blood 74: 55a, 1989; Hartwig and Danishefsky, J. Biol. Chem. 266: 6578-6585, 1991; Farrell et al., Biochemistry 30: 9414-9420, 1991), but expression levels have been so low that production of recombinant fibrinogen in commercial quantities is not feasible. Experimental evidence suggests that lower transcription rates in cultured cells as compared to liver may be a factor in the low expression rates achieved to date, but increasing the amount of fibrinogen chain mRNA in transfected BHK cells did not produce corresponding increases in fibrinogen protein secretion (Prunkard and Foster, XIV Congress of the International Society on Thrombosis and Haemostasis, 1993). These latter results suggest that proper assembly and processing of fibrinogen involves tissue-specific mechanisms not present in common laboratory cell lines.

There remains a need in the art for methods of producing large quantities of high quality fibrinogen for use in tissue adhesives and other applications. There is a further need for fibrinogen that is free of blood-borne pathogens. The present invention fulfills these needs and provides other, related advantages.

#### Summary of the Invention

It is an object of the present invention to provide commercially useful quantities of recombinant fibrinogen, particularly recombinant human fibrinogen. It is a further object of the invention to provide materials and methods for expressing fibrinogen in the mammary tissue of transgenic animals, particularly livestock animals such as cattle, sheep, pigs and goats.

Within one aspect, the present invention provides a method for producing fibrinogen comprising (a) providing a first DNA segment encoding a secretion signal operably linked to a fibrinogen A $\alpha$  chain, a second DNA segment encoding a secretion signal operably linked to a fibrinogen B $\beta$  chain, and a third DNA segment encoding a secretion signal operably linked to a fibrinogen  $\gamma$  chain, wherein each of the first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal; (b) introducing the DNA segments into a fertilized egg of a non-human mammalian species; (c) inserting the egg into an oviduct or uterus of a female of the species to obtain offspring carrying the DNA constructs; (d) breeding the offspring to produce female progeny that express the first, second and third DNA segments and produce milk containing biocompetent fibrinogen encoded by the segments; (e) collecting milk from the female progeny; and (f) recovering the fibrinogen from the milk. Within one embodiment, the egg containing the introduced segments is cultured for a period of time prior to insertion.

Within another aspect, the invention provides a method of producing fibrinogen comprising the steps of (a) incorporating a first DNA segment encoding a secretion signal operably linked to an A $\alpha$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a first gene fusion; (b) incorporating a second DNA segment encoding a secretion signal operably linked to a B $\beta$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a second gene fusion; (c) incorporating a third DNA segment encoding a secretion signal operably linked to a  $\gamma$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a third gene fusion; (d) introducing the first, second and third gene fusions into the germ line of a non-human mammal so that the DNA segments are expressed in a mammary gland of the mammal or its female progeny and biocompetent fibrinogen is secreted into milk of the mammal or its female progeny; (e) obtaining milk from the mammal or its female progeny; and (f) recovering the fibrinogen from the milk. Within preferred embodiments, the mammal is a sheep, pig, goat or bovine.

Within another aspect, the invention provides a method for producing fibrinogen comprising the steps of (a) providing a transgenic female non-human mammal carrying in its germline heterologous DNA segments encoding A $\alpha$ , B $\beta$  and  $\gamma$  chains of fibrinogen, wherein the DNA segments are expressed in a mammary gland of the mammal and fibrinogen encoded by the DNA segments is secreted into milk of the mammal; (b) collecting milk from the mammal; and (c) recovering the fibrinogen from the milk.

Within another aspect, the invention provides a non-human mammalian embryo containing in its nucleus heterologous DNA segments encoding A $\alpha$ , B $\beta$  and  $\gamma$  chains of fibrinogen. Within a related aspect, the invention provides a transgenic non-human female mammal that produces recoverable amounts of human fibrinogen in its milk.

Within another aspect, the invention provides a method for producing a transgenic offspring of a mammal comprising the steps of (a) providing a first DNA segment encoding a fibrinogen A $\alpha$  chain, a second DNA segment encoding a fibrinogen B $\beta$  chain, and a third DNA segment encoding a fibrinogen  $\gamma$  chain, wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in a mammary gland of a host female mammal and secretion into milk of the host female mammal; (b) introducing the DNA segments into a fertilized egg of a mammal of a non-human species; (c) inserting the egg into an oviduct or uterus of a female of the non-human species to obtain an offspring carrying the first, second and third DNA segments. In a related aspect, the invention provides non-human mammals produced according to this process.

Within an additional aspect, the invention provides a non-human mammal carrying its germline DNA segments encoding heterologous A $\alpha$ , B $\beta$  and  $\gamma$  chains of fibrinogen, wherein female progeny of the mammal express the DNA segments in a mammary gland to produce biocompetent fibrinogen.

These and other aspects of the invention will become evident to the skilled practitioner upon reference to the following detailed description and the attached drawings.

### Brief Description of the Drawings

Figure 1 illustrates the subcloning of a human fibrinogen A $\alpha$  chain DNA sequence.

Figure 2 is a partial restriction map of the vector Zem228. Symbols used are MT-1p, mouse metallothionein promoter; SV40t, SV40 terminator; and SV40p, SV40 promoter.

Figure 3 illustrates the subcloning of a human fibrinogen B $\beta$  chain DNA sequence.

Figure 4 illustrates the subcloning of a human fibrinogen  $\gamma$  chain DNA sequence.

Figure 5 is a partial restriction map of the vector Zem219b. Symbols used are MT-1p, mouse metallothionein promoter; hGHt, human growth hormone terminator; SV40p, SV40 promoter; DHFR, dihydrofolate reductase gene; and SV40t, SV40 terminator.

### Detailed Description of the Invention

Prior to setting forth the invention in detail, it will be helpful to define certain terms used herein:

As used herein, the term "biocompetent fibrinogen" is used to denote fibrinogen that polymerizes when treated with thrombin to form insoluble fibrin.

The term "egg" is used to denote an unfertilized ovum, a fertilized ovum prior to fusion of the pronuclei or an early stage embryo (fertilized ovum with fused pronuclei).

A "female mammal that produces milk containing biocompetent fibrinogen" is one that, following pregnancy and delivery, produces, during the lactation period, milk containing recoverable amounts of biocompetent fibrinogen. Those skilled in the art will recognized that such animals will produce milk, and therefore the fibrinogen, discontinuously.

The term "progeny" is used in its usual sense to include children and descendants.



The term "heterologous" is used to denote genetic material originating from a different species than that into which it has been introduced, or a protein produced from such genetic material.

5           Within the present invention, transgenic animal technology is employed to produce fibrinogen within the mammary glands of a host female mammal. Expression in the mammary gland and subsequent secretion of the protein of interest into the milk overcomes many difficulties  
10 encountered in isolating proteins from other sources. Milk is readily collected, available in large quantities, and well characterized biochemically. Furthermore, the major milk proteins are present in milk at high concentrations (from about 1 to 15 g/l).

15           From a commercial point of view, it is clearly preferable to use as the host a species that has a large milk yield. While smaller animals such as mice and rats can be used (and are preferred at the proof-of-concept stage), within the present invention it is preferred to  
20 use livestock mammals including, but not limited to, pigs, goats, sheep and cattle. Sheep are particularly preferred due to such factors as the previous history of transgenesis in this species, milk yield, cost and the ready availability of equipment for collecting sheep milk.  
25 See WO 88/00239 for a comparison of factors influencing the choice of host species. It is generally desirable to select a breed of host animal that has been bred for dairy use, such as East Friesland sheep, or to introduce dairy stock by breeding of the transgenic line at a later date.  
30 In any event, animals of known, good health status should be used.

Fibrinogen produced according to the present invention may be human fibrinogen or fibrinogen of a non-human animal. For medical uses, it is preferred to employ  
35 proteins native to the patient. The present invention thus provides fibrinogen for use in both human and veterinary medicine. Cloned DNA molecules encoding the

component chains of human fibrinogen are disclosed by Rixon et al. (Biochem. 22: 3237, 1983), Chung et al. (Biochem. 22: 3244, 1983), Chung et al. (Biochem. 22: 3250, 1983), Chung et al. (Adv. Exp. Med. Biol. 281: 39-48, 1990) and Chung et al. (Ann. NY Acad. Sci. 408: 449-456, 1983). Bovine fibrinogen clones are disclosed by Brown et al. (Nuc. Acids Res. 17: 6397, 1989) and Chung et al. (Proc. Natl. Acad. Sci. USA 78: 1466-1470, 1981). Other mammalian fibrinogen clones are disclosed by Murakawa et al. (Thromb. Haemost. 69: 351-360, 1993). Representative sequences of human A $\alpha$ , B $\beta$  and  $\gamma$  chain genes are shown in SEQ ID NOS: 1, 3 and 5, respectively. Those skilled in the art will recognize that allelic variants of these sequences will exist; that additional variants can be generated by amino acid substitution, deletion, or insertion; and that such variants are useful within the present invention. In general, it is preferred that any engineered variants comprise only a limited number of amino acid substitutions, deletions, or insertions, and that any substitutions are conservative. Thus, it is preferred to produce fibrinogen chain polypeptides that are at least 90%, preferably at least 95%, and more preferably 99% or more identical in sequence to the corresponding native chains. The term " $\gamma$  chain" is meant to include the alternatively spliced  $\gamma'$  chain of fibrinogen (Chung et al., Biochem. 23: 4232-4236, 1984). A human  $\gamma'$  chain amino acid sequence is shown in SEQ ID NO: 6. The shorter  $\gamma$  chain is produced by alternative splicing at nucleotides 9511 and 10054 of SEQ ID NO: 5, resulting in translation terminating after nucleotide 10065 of SEQ ID NO: 5. .

To obtain expression in the mammary gland, a transcription promoter from a milk protein gene is used. Milk protein genes include those genes encoding caseins, beta-lactoglobulin (BLG),  $\alpha$ -lactalbumin, and whey acidic protein. The beta-lactoglobulin promoter is preferred. In the case of the ovine beta-lactoglobulin gene, a region

of at least the proximal 406 bp of 5' flanking sequence of the ovine BLG gene (contained within nucleotides 3844 to 4257 of SEQ ID NO:7) will generally be used. Larger portions of the 5' flanking sequence, up to about 5 kbp, are preferred. A larger DNA segment encompassing the 5' flanking promoter region and the region encoding the 5' non-coding portion of the beta-lactoglobulin gene (contained within nucleotides 1 to 4257 of SEQ ID NO:7) is particularly preferred. See Whitelaw et al., Biochem J. 10 286: 31-39, 1992. Similar fragments of promoter DNA from other species are also suitable.

Other regions of the beta-lactoglobulin gene may also be incorporated in constructs, as may genomic regions of the gene to be expressed. It is generally accepted in the art that constructs lacking introns, for example, express poorly in comparison with those that contain such DNA sequences (see Brinster et al., Proc. Natl. Acad. Sci. USA 85: 836-840, 1988; Palmiter et al., Proc. Natl. Acad. Sci. USA 88: 478-482, 1991; Whitelaw et al., Transgenic Res. 1: 3-13, 1991; WO 89/01343; WO 91/02318). In this regard, it is generally preferred, where possible, to use genomic sequences containing all or some of the native introns of a gene encoding the protein or polypeptide of interest. Within certain embodiments of the invention, the further inclusion of at least some introns from the beta-lactoglobulin gene is preferred. One such region is a DNA segment which provides for intron splicing and RNA polyadenylation from the 3' non-coding region of the ovine beta-lactoglobulin gene. When substituted for the natural 3' non-coding sequences of a gene, this ovine beta-lactoglobulin segment can both enhance and stabilize expression levels of the protein or polypeptide of interest. Within other embodiments, the region surrounding the initiation ATG of one or more of the fibrinogen sequences is replaced with corresponding sequences from a milk specific protein gene. Such replacement provides a putative tissue-specific initiation

environment to enhance expression. It is convenient to replace the entire fibrinogen chain pre-pro and 5' non-coding sequences with those of, for example, the BLG gene, although smaller regions may be replaced.

5 For expression of fibrinogen, DNA segments encoding each of the three component polypeptide chains of fibrinogen are operably linked to additional DNA segments required for their expression to produce expression units. Such additional segments include the above-mentioned milk  
10 protein gene promoter, as well as sequences which provide for termination of transcription and polyadenylation of mRNA. The expression units will further include a DNA segment encoding a secretion signal operably linked to the segment encoding the fibrinogen polypeptide chain. The  
15 secretion signal may be a native fibrinogen secretion signal or may be that of another protein, such as a milk protein. The term "secretion signal" is used herein to denote that portion of a protein that directs it through the secretory pathway of a cell to the outside. Secretion  
20 signals are most commonly found at the amino-termini of proteins. See, for example, von Heinje, Nuc. Acids Res. 14: 4683-4690, 1986; and Meade et al., U.S. Patent No. 4,873,316, which are incorporated herein by reference.

Construction of expression units is conveniently  
25 carried out by inserting a fibrinogen chain sequence into a plasmid or phage vector containing the additional DNA segments, although the expression unit may be constructed by essentially any sequence of ligations. It is particularly convenient to provide a vector containing a  
30 DNA segment encoding a milk protein and to replace the coding sequence for the milk protein with that of a fibrinogen chain (including a secretion signal), thereby creating a gene fusion that includes the expression control sequences of the milk protein gene. In any event,  
35 cloning of the expression units in plasmids or other vectors facilitates the amplification of the fibrinogen sequences. Amplification is conveniently carried out in

bacterial (e.g. *E. coli*) host cells, thus the vectors will typically include an origin of replication and a selectable marker functional in bacterial host cells.

In view of the size of the fibrinogen chain genes it is most practical to prepare three separate expression units, mix them, and introduce the mixture into the host. However, those skilled in the art will recognize that other protocols may be followed. For example, expression units for the three chains can be introduced individually into different embryos to be combined later by breeding. In a third approach, the three expression units can be linked in a single suitable vector, such as a yeast artificial chromosome or phage P1 clone. Coding sequences for two or three chains can be combined in polycistronic expression units (see, e.g., Levinson et al., U.S. Patent No. 4,713,339).

The expression unit(s) is(are) then introduced into fertilized eggs (including early-stage embryos) of the chosen host species. Introduction of heterologous DNA can be accomplished by one of several routes, including microinjection (e.g. U.S. Patent No. 4,873,191), retroviral infection (Jaenisch, Science 240: 1468-1474, 1988) or site-directed integration using embryonic stem (ES) cells (reviewed by Bradley et al., Bio/Technology 10: 534-539, 1992). The eggs are then implanted into the oviducts or uteri of pseudopregnant females and allowed to develop to term. Offspring carrying the introduced DNA in their germ line can pass the DNA on to their progeny in the normal, Mendelian fashion, allowing the development of transgenic herds. General procedures for producing transgenic animals are known in the art. See, for example, Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, 1986; Simons et al., Bio/Technology 6: 179-183, 1988; Wall et al., Biol. Reprod. 32: 645-651, 1985; Buhler et al., Bio/Technology 8: 140-143, 1990; Ebert et al., Bio/Technology 9: 835-838, 1991; Krimpenfort et al.,

Bio/Technology 9: 844-847, 1991; Wall et al., J. Cell. Biochem. 49: 113-120, 1992; and WIPO publications WO 88/00239, WO 90/05188, WO 92/11757; and GB 87/00458, which are incorporated herein by reference. Techniques for  
5 introducing foreign DNA sequences into mammals and their germ cells were originally developed in the mouse. See, e.g., Gordon et al., Proc. Natl. Acad. Sci. USA 77: 7380-7384, 1980; Gordon and Ruddle, Science 214: 1244-1246, 1981; Palmiter and Brinster, Cell 41: 343-345, 1985;  
10 Brinster et al., Proc. Natl. Acad. Sci. USA 82: 4438-4442, 1985; and Hogan et al. (ibid.). These techniques were subsequently adapted for use with larger animals, including livestock species (see e.g., WIPO publications WO 88/00239, WO 90/05188, and WO 92/11757; and Simons et  
15 al., Bio/Technology 6: 179-183, 1988). To summarize, in the most efficient route used to date in the generation of transgenic mice or livestock, several hundred linear molecules of the DNA of interest are injected into one of the pro-nuclei of a fertilized egg. Injection of DNA into  
20 the cytoplasm of a zygote can also be employed.

It is preferred to obtain a balanced expression of each fibrinogen chain to allow for efficient formation of the mature protein. Ideally, the three expression units should be on the same DNA molecule for introduction  
25 into eggs. This approach, however, may generate technical problems at, for example, the injection and manipulation stages. For example, the size of fibrinogen expression units may necessitate the use of yeast artificial chromosomes (YACs) or phage P1 to amplify and manipulate  
30 the DNA prior to injection. If this approach is followed, segments of DNA to be injected, containing all three expression units, would be very large, thus requiring modification of the injection procedure using, for example, larger bore needles. In a more simple approach,  
35 a mixture of each individual expression unit is used. It is preferred to combine equimolar amounts of the three expression units, although those skilled in the art will

recognize that this ratio may be varied to compensate for the characteristics of a given expression unit. Some expression, generally a reduced level, will be obtained when lesser molar amounts of one or two chains are used, and expression efficiencies can generally be expected to decline in approximate proportion to the divergence from the preferred equimolar ratio. In any event, it is preferred to use a mixture having a ratio of  $A\alpha:B\beta:\gamma$  expression units in the range of 0.5-1:0.5-1:0.5-1. When the ratio is varied from equimolar, it is preferred to employ relatively more of the  $B\beta$  expression unit. Alternatively, one or a mixture of two of the expression units is introduced into individual eggs. However, animals derived by this approach will express only one or two fibrinogen chains. To generate an intact fibrinogen molecule by this approach requires a subsequent breeding program designed to combine all three expression units in individuals of a group of animals.

In general, female animals are superovulated by treatment with follicle stimulating hormone, then mated. Fertilized eggs are collected, and the heterologous DNA is injected into the eggs using known methods. See, for example, U.S. Patent No. 4,873,191; Gordon et al., Proc. Natl. Acad. Sci. USA 77: 7380-7384, 1980; Gordon and Ruddle, Science 214: 1244-1246, 1981; Palmiter and Brinster, Cell 41: 343-345, 1985; Brinster et al., Proc. Natl. Acad. Sci. USA 82: 4438-4442, 1985; Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, 1986; Simons et al. Bio/Technology 6: 179-183, 1988; Wall et al., Biol. Reprod. 32: 645-651, 1985; Buhler et al., Bio/Technology 8: 140-143, 1990; Ebert et al., Bio/Technology 9: 835-838, 1991; Krimpenfort et al., Bio/Technology 9: 844-847, 1991; Wall et al., J. Cell. Biochem. 49: 113-120, 1992; WIPO publications WO 88/00239, WO 90/05118, and WO 92/11757; and GB 87/00458, which are incorporated herein by reference.

For injection into fertilized eggs, the expression units are removed from their respective vectors by digestion with appropriate restriction enzymes. For convenience, it is preferred to design the vectors so that  
5 the expression units are removed by cleavage with enzymes that do not cut either within the expression units or elsewhere in the vectors. The expression units are recovered by conventional methods, such as electro-elution followed by phenol extraction and ethanol precipitation,  
10 sucrose density gradient centrifugation, or combinations of these approaches.

DNA is injected into eggs essentially as described in Hogan et al., *ibid.* In a typical injection, eggs in a dish of an embryo culture medium are located  
15 using a stereo zoom microscope (x50 or x63 magnification preferred). Suitable media include Hepes (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) or bicarbonate buffered media such as M2 or M16 (available from Sigma Chemical Co., St. Louis, USA) or synthetic  
20 oviduct medium (disclosed below). The eggs are secured and transferred to the center of a glass slide on an injection rig using, for example, a drummond pipette complete with capillary tube. Viewing at lower (e.g. x4) magnification is used at this stage. Using the holding  
25 pipette of the injection rig, the eggs are positioned centrally on the slide. Individual eggs are sequentially secured to the holding pipette for injection. For each injection process, the holding pipette/egg is positioned in the center of the viewing field. The injection needle  
30 is then positioned directly below the egg. Preferably using x40 Nomarski objectives, both manipulator heights are adjusted to focus both the egg and the needle. The pronuclei are located by rotating the egg and adjusting the holding pipette assembly as necessary. Once the  
35 pronucleus has been located, the height of the manipulator is altered to focus the pronuclear membrane. The injection needle is positioned below the egg such that the



needle tip is in a position below the center of the pronucleus. The position of the needle is then altered using the injection manipulator assembly to bring the needle and the pronucleus into the same focal plane. The  
5 needle is moved, via the joy stick on the injection manipulator assembly, to a position to the right of the egg. With a short, continuous jabbing movement, the pronuclear membrane is pierced to leave the needle tip inside the pronucleus. Pressure is applied to the  
10 injection needle via the glass syringe until the pronucleus swells to approximately twice its volume. At this point, the needle is slowly removed. Reverting to lower (e.g. x4) magnification, the injected egg is moved to a different area of the slide, and the process is  
15 repeated with another egg.

After the DNA is injected, the eggs may be cultured to allow the pronuclei to fuse, producing one-cell or later stage embryos. In general, the eggs are cultured at approximately the body temperature of the  
20 species used in a buffered medium containing balanced salts and serum. Surviving embryos are then transferred to pseudopregnant recipient females, typically by inserting them into the oviduct or uterus, and allowed to develop to term. During embryogenesis, the injected DNA  
25 integrates in a random fashion in the genomes of a small number of the developing embryos.

Potential transgenic offspring are screened via blood samples and/or tissue biopsies. DNA is prepared from these samples and examined for the presence of the  
30 injected construct by techniques such as polymerase chain reaction (PCR; see Mullis, U.S. Patent No. 4,683,202) and Southern blotting (Southern, J. Mol. Biol. 98:503, 1975; Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, 1982). Founder transgenic  
35 animals, or G0s, may be wholly transgenic, having transgenes in all of their cells, or mosaic, having transgenes in only a subset of cells (see, for example,

Wilkie et al., Develop. Biol. 118: 9-18, 1986). In the latter case, groups of germ cells may be wholly or partially transgenic. In the latter case, the number of transgenic progeny from a founder animal will be less than  
5 the expected 50% predicted from Mendelian principles. Founder G0 animals are grown to sexual maturity and mated to obtain offspring, or G1s. The G1s are also examined for the presence of the transgene to demonstrate transmission from founder G0 animals. In the case of male  
10 G0s, these may be mated with several non-transgenic females to generate many offspring. This increases the chances of observing transgene transmission. Female G0 founders may be mated naturally, artificially inseminated or superovulated to obtain many eggs which are transferred  
15 to surrogate mothers. The latter course gives the best chance of observing transmission in animals having a limited number of young. The above-described breeding procedures are used to obtain animals that can pass the DNA on to subsequent generations of offspring in the  
20 normal, Mendelian fashion, allowing the development of, for example, colonies (mice), flocks (sheep), or herds (pigs, goats and cattle) of transgenic animals.

The milk from lactating G0 and G1 females is examined for the expression of the heterologous protein  
25 using immunological techniques such as ELISA (see Harlow and Lane, Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, 1988) and Western blotting (Towbin et al., Proc. Natl. Acad. Sci. USA 76: 4350-4354, 1979). For a variety of reasons known in the art, expression levels  
30 of the heterologous protein will be expected to differ between individuals.

A satisfactory family of animals should satisfy three criteria: they should be derived from the same founder G0 animal; they should exhibit stable transmission  
35 of the transgene; and they should exhibit stable expression levels from generation to generation and from lactation to lactation of individual animals. These

principles have been demonstrated and discussed (Carver et al., Bio/Technology 11: 1263-1270, 1993). Animals from such a suitable family are referred to as a "line." Initially, male animals, G0 or G1, are used to derive a flock or herd of producer animals by natural or artificial insemination. In this way, many female animals containing the same transgene integration event can be quickly generated from which a supply of milk can be obtained.

The fibrinogen is recovered from milk using standard practices such as skimming, precipitation, filtration and protein chromatography techniques.

Fibrinogen produced according to the present invention is useful within human and veterinary medicine, such as in the formulation of surgical adhesives. Adhesives of this type are known in the art. See, for example, U.S. Patents No. 4,377,572; 4,442,655; 4,462,567; and 4,627,879, which are incorporated herein by reference. In general, fibrinogen and factor XIII are combined to form a first component that is mixed just prior to use with a second component containing thrombin. The thrombin converts the fibrinogen to fibrin, causing the mixture to gel, and activates the factor XIII. The activated factor XIII cross links the fibrin to strengthen and stabilize the adhesive matrix. Such adhesives typically contain from about 30 mg/ml to about 100 mg/ml fibrinogen and from about 50  $\mu$ g/ml to about 500  $\mu$ g/ml factor XIII. They may also contain additional ingredients, such as aprotinin, albumin, fibronectin, bulking agents, and solubilizers. Methods for producing factor XIII are known in the art. See, for example, U.S. Patent No. 5,204,447. The fibrinogen is also useful for coating surfaces of polymeric articles, e.g. synthetic vascular grafts, as disclosed in U.S. Patent No. 5,272,074 (incorporated herein by reference).

The invention is further illustrated by the following non-limiting examples.

ExamplesExample I

The multiple cloning site of the vector pUC18  
5 (Yanisch-Perron et al., Gene 33:103-119, 1985) was removed  
and replaced with a synthetic double stranded  
oligonucleotide (the strands of which are shown in SEQ ID  
NO: 8 and SEQ ID NO: 27) containing the restriction sites  
Pvu I/Mlu I/Eco RV/Xba I/Pvu I/Mlu I, and flanked by 5'  
10 overhangs compatible with the restriction sites Eco RI and  
Hind III. pUC18 was cleaved with both Eco RI and Hind  
III, the 5' terminal phosphate groups were removed with  
calf intestinal phosphatase, and the oligonucleotide was  
ligated into the vector backbone. The DNA sequence across  
15 the junction was confirmed by sequencing, and the new  
plasmid was called pUCPM.

The  $\beta$ -lactoglobulin (BLG) gene sequences from  
pSS1tgXS (disclosed in WIPO publication WO 88/00239) were  
excised as a Sal I-Xba I fragment and recloned into the  
20 vector pUCPM that had been cut with Sal I and Xba I to  
construct vector pUCXS. pUCXS is thus a pUC18 derivative  
containing the entire BLG gene from the Sal I site to the  
Xba I site of phage SS1 (Ali and Clark, J. Mol. Biol. 199:  
415-426, 1988).

25 The plasmid pSS1tgSE (disclosed in WIPO  
publication WO 88/00239) contains a 1290 bp BLG fragment  
flanked by Sph I and EcoR I restriction sites, a region  
spanning a unique Not I site and a single Pvu II site  
which lies in the 5' untranslated leader of the BLG mRNA.  
30 Into this Pvu II site was ligated a double stranded, 8 bp  
DNA linker (5'-GGATATCC-3') encoding the recognition site  
for the enzyme Eco RV. This plasmid was called  
pSS1tgSE/RV. DNA sequences bounded by Sph I and Not I  
restriction sites in pSS1tgSE/RV were excised by enzymatic  
35 digestion and used to replace the equivalent fragment in  
pUCXS. The resulting plasmid was called pUCXSRV. The  
sequence of the BLG insert in pUCXSRV is shown in SEQ ID

NO: 7, with the unique Eco RV site at nucleotide 4245 in the 5' untranslated leader region of the BLG gene. This site allows insertion of any additional DNA sequences under the control of the BLG promoter 3' to the transcription initiation site.

Using the primers BLGAMP3 (5'-TGG ATC CCC TGC CCG TGC CTC TGG-3'; SEQ ID NO: 9) and BLGAMP4 (5'-AAC GCG TCA TCC TCT GTG AGC CAG-3'; SEQ ID NO: 10) a PCR fragment of approximately 650 bp was produced from sequences immediately 3' to the stop codon of the BLG gene in pUCXSRV. The PCR fragment was engineered to have a BamH I site at its 5' end and an Mlu I site at its 3' end and was cloned as such into BamH I and Mlu I cut pGEM7zf(+) (Promega) to give pDAM200(+).

pUCXSRV was digested with Kpn I, and the largest, vector containing band was gel purified. This band contained the entire pUC plasmid sequences and some 3' non-coding sequences from the BLG gene. Into this backbone was ligated the small Kpn I fragment from pDAM200(+) which, in the correct orientation, effectively engineered a BamH I site at the extreme 5' end of the 2.6 Kbp of the BLG 3' flanking region. This plasmid was called pBLAC200. A 2.6 Kbp Cla I-Xba I fragment from pBLAC200 was ligated into Cla I-Xba I cut pSP72 vector (Promega), thus placing an EcoR V site immediately upstream of the BLG sequences. This plasmid was called pBLAC210.

The 2.6 Kbp Eco RV-Xba I fragment from pBLAC210 was ligated into Eco RV-Xba I cut pUCXSRV to form pMAD6. This, in effect, excised all coding and intron sequences from pUCXSRV, forming a BLG minigene consisting of 4.3 Kbp of 5' promoter and 2.6 Kbp of 3' downstream sequences flanking a unique EcoR V site. An oligonucleotide linker (ZC6839: ACTACGTAGT; SEQ ID NO: 11) was inserted into the Eco RV site of pMAD6. This modification destroyed the Eco RV site and created a Sna BI site to be used for cloning purposes. The vector was designated pMAD6-Sna. Messenger

RNA initiates upstream of the Sna BI site and terminates downstream of the Sna BI site. The precursor transcript will encode a single BLG-derived intron, intron 6, which is entirely within the 3' untranslated region of the gene.

5

### Example II

Clones encoding the individual fibrinogen chains were obtained from the laboratory of Dr. Earl W. Davie, University of Washington, Seattle. A genomic fibrinogen  $\text{A}\alpha$ -chain clone (Chung et al., 1990, *ibid.*) was obtained from the plasmid BS4. This plasmid contains the  $\text{A}\alpha$  clone inserted into the Sal I and Bam HI sites of the vector pUC18, but lacks the coding sequence for the first four amino acids of the  $\text{A}\alpha$  chain. A genomic  $\text{B}\beta$ -chain DNA (Chung et al., *ibid.*) was isolated from a lambda Charon 4A phage clone (designated  $\beta\lambda 4$ ) as two EcoRI fragments of ca. 5.6 Kbp each. The two fragments were cloned separately into pUC19 that had been digested with Eco RI and treated with calf intestinal phosphatase. The resulting clones were screened by digestion with the restriction enzyme Pvu II to distinguish plasmids with the 5' and 3'  $\text{B}\beta$  inserts (designated Beta5'RI/puc and Beta3'RI/puc, respectively). Genomic  $\gamma$ -chain clones were isolated as described by Rixon et al. (Biochemistry 24: 2077-2086, 1985). Clone py12A9 comprises 5' non-coding sequences and approximately 4535 bp of  $\gamma$ -chain coding sequence. Clone py12F3 comprises the remaining coding sequence and 3' non-coding nucleotides. Both are pBR322-based plasmids with the fibrinogen sequences inserted at the EcoRI site. These plasmids were used as templates for the respective PCR reactions.

The fibrinogen chain coding sequences were tailored for insertion into expression vectors using the polymerase chain reaction (PCR) as generally described by Mullis (U.S. Patent No. 4,683,202). This procedure removed native 5' and 3' untranslated sequences, added a 9 base sequence (CCT GCA GCC) upstream of the first ATG of

each coding sequence, supplied the first four codons for the A $\alpha$ -chain sequence, removed an internal Mlu I site in the A $\alpha$  sequence and added restriction sites to facilitate subsequent cloning steps.

5 Referring to Figure 1, the 5' end of the A $\alpha$  coding sequence was tailored in a PCR reaction containing 20 pmole for each of primers ZC6632 (SEQ ID NO: 12) and ZC6627 (SEQ ID NO: 13), approximately 10 ng of plasmid BS4 template DNA, 10  $\mu$ l of a mix containing 2.5 mM each dNTP,  
10 7.5  $\mu$ l 10x *Pyrococcus furiosus* (Pfu) DNA polymerase buffer #1 (200 mM Tris-HCl, pH 8.2, 100 mM KCl, 60 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM MgCl<sub>2</sub>, 1% Triton X-100, 100  $\mu$ g/ml nuclease free bovine serum albumin) (Stratagene, La Jolla, CA), and water to 75  $\mu$ l. The mixture was heated to 94°C in a DNA thermal  
15 cycler (Perkin-Elmer Corp., Norwalk, CT). To the heated mixture was added 25  $\mu$ l of a mixture containing 2.5  $\mu$ l 10x Pfu buffer #1, 22  $\mu$ l H<sub>2</sub>O and 1  $\mu$ l 2.5 units/ $\mu$ l Pfu DNA polymerase (Stratagene). The reactions were run in a DNA thermal cycler (Perkin-Elmer) for five cycles of 94°, 45  
20 seconds; 40°, 90 seconds; 72°, 120 seconds; 20 cycles of 94°, 45 seconds; 45°, 90 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. The 5' PCR-generated fragment was digested with Bam HI and Hind III, and the Bam HI-Hind III fragment was then ligated to an internal  
25 2.91 Kbp Hind III-Xba I fragment and Bam HI, Xba I-digested pUC18. PCR-generated exon sequences were sequenced.

Referring again to Figure 1, the 3' end of the A $\alpha$  coding sequence was tailored in a series of steps in  
30 which the Mlu I site 563 bases upstream from the stop codon of the A $\alpha$  sequence was mutated using an overlap extension PCR reaction (Ho et al., Gene 77: 51-59, 1989). In the first reaction 40 pmole of each of primers ZC6521 (SEQ ID NO: 14) and ZC6520 (SEQ ID NO: 15) were combined  
35 with approximately 10 ng of plasmid BS4 template DNA in a reaction mixture as described above. The reaction was run for 5 cycles of 94°, 45 seconds; 40°, 60 seconds; 72°, 120

seconds; 15 cycles of 94°, 45 seconds; 45°, 60 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. A second reaction was carried out in the same manner using 40 pmole of each of primers ZC6519 (SEQ ID NO: 16) and ZC6518 (SEQ ID NO: 17) and BS4 as template. The PCR-generated DNA fragments from the first and second reactions were isolated by gel electrophoresis and elution from the gel. Approximately 1/10 of each recovered reaction product was combined with 40 pmole of each of primers ZC6521 (SEQ ID NO: 14) and ZC6518 (SEQ ID NO: 17) in a PCR reaction in which the complementary 3' ends of each fragment (containing the single base change) annealed and served as a primer for the 3' extension of the complementary strand. PCR was carried out using the same reaction conditions as in the first and second 3' PCR steps. The reaction product was then digested with Xba I and Bam HI, and the Xba I-Bam HI fragment was cloned into Xba I, Bam HI-digested pUC18. PCR-generated exons were sequenced.

As shown in Figure 1, the 5' Bam HI-Xba I fragment (3.9 Kbp) and the 3' Xba I-Bam HI fragment (1.3 Kbp) were inserted into the Bam HI site of the vector Zem228. Zem228 is a pUC18 derivative comprising a Bam HI cloning site between a mouse MT-1 promoter and SV40 terminator, and a neomycin resistance marker flanked by SV40 promoter and terminator sequences. See European Patent Office Publication EP 319,944 and Fig. 2. The entire  $\alpha$  coding sequence was isolated from the Zem228 vector as an Sna BI fragment, which was inserted into the Sna BI site of the plasmid pMAD6-Sna.

Referring to Fig. 3, the 5' end of the B $\beta$ -chain was tailored by PCR using the oligonucleotides ZC6629 (SEQ ID NO: 18), ZC6630 (SEQ ID NO: 19) and ZC6625 (SEQ ID NO: 20). These primers were used in pairwise combinations (ZC6629 + ZC6625 or ZC6630 + ZC6625) to generate B $\beta$  coding sequences beginning at the first ATG codon (position 470 in SEQ ID NO: 3) (designated N1-Beta) or the third ATG



codon (position 512 in SEQ ID NO: 3) (designated N3-Beta). Approximately 5 ng of Beta5'RI/puc template DNA was combined with 20 pmole of each of the primers (N1-Beta:ZC6629, SEQ ID NO: 18 + ZC6625, SEQ ID NO: 20; or N3-Beta:ZC6630, SEQ ID NO: 19 + ZC6625, SEQ ID NO: 20) in a reaction mixture as described above. The mixtures were incubated for 5 cycles of 94°, 45 seconds; 40°, 120 seconds; (N1-Beta) or 90 seconds (N3-Beta); 72°, 120 seconds; 20 cycles of 94°, 45 seconds; 45°, 120 seconds; (N1-Beta) or 90 seconds (N3-Beta); 72°, 120 seconds; then incubated at 72° for 7 minutes. The two reaction products N1, 555 bp or N3, 510 bp) were each digested with Eco RI and Bgl II, and the fragments were ligated to the internal Bgl II-Xba I fragment and Eco RI + Xba I-digested pUC19. The 3' end of the B $\beta$  sequence was tailored in a reaction mixture as described above using the oligonucleotide primers ZC6626 (SEQ ID NO: 21) and ZC6624 (SEQ ID NO: 22) and approximately 5 ng of Beta3'RI/puc template. The mixtures were incubated for 5 cycles of 94°, 45 seconds; 40°, 90 seconds; 72°, 120 seconds; 15 cycles of 94°, 45 seconds; 45°, 90 seconds; 72°, 120 seconds; then incubated at 72° for 7 minutes. A 990 bp Bgl II-Eco RI fragment was isolated. This 3' fragment was ligated to the adjacent coding fragment (340 bp, SphI-Bgl II) and Sph I + Eco RI-digested pUC19. The 3' and 5' PCR-generated exons were sequenced. A third intermediate vector was constructed by combining two internal fragments (4285 bp Xba I-Eco RI and 383 kb Eco RI-Sph I) in Xba I + Sph I-digested pUC19. The entire B $\beta$  coding sequence (two forms) was then assembled by ligating one of the 5' Eco RI-Xba I fragments, the internal Xba I-Sph I fragment, the 3' Sph I-Eco RI fragment and Eco RI-digested vector pUC19. The B $\beta$  sequence was then isolated as a 7.6 Kbp Sna BI fragment and inserted into the Sna BI site of pMAD6-Sna.

Referring to Fig. 4, the 5' end of the gamma chain sequence was tailored by PCR using the oligonucleotide primers ZC6514 (SEQ ID NO: 23) and ZC6517

(SEQ ID NO: 24) and approximately 50 ng of py12A9 as template. The PCR reaction was run as described above using 40 pM of each primer. The reaction was run for 5 cycles of 94°, 45 seconds; 40°, 60 seconds, 72°, 120  
5 seconds, followed by 15 cycles of 94°, 45 seconds; 45°, 60 seconds; 72°, 120 seconds. The resulting 213 bp fragment was digested with Bam HI and Spe I, and the resulting restriction fragment was ligated with the adjacent  
10 downstream 4.4 kb Spe I-Eco RI fragment and Bam HI + Eco RI digested pUC19. The 3' end of the gamma chain sequence was tailored using oligonucleotide primers ZC6516 (SEQ ID NO: 25) and ZC6515 (SEQ ID NO: 26) using 40 pM of each primer, approximately 50 ng of py12F3 template and the same thermal cycling schedule as used for the 5' fragment.  
15 The resulting 500 bp fragment was digested with Spe I and Bam HI, and the resulting restriction fragment was ligated with the upstream 2.77 kb Eco RI-Spe I fragment and Eco RI + Bam HI-digested pUC19. All PCR-generated exons were sequenced. The entire  $\gamma'$ -chain coding sequence was then  
20 assembled by ligating a 4.5 Kbp Bam HI-Eco RI 5' fragment, a 1.1 Kbp Eco RI-Pst I internal fragment and a 2.14 Kbp Pst I-Xba I 3' fragment in Bam HI + Xba I-digested Zem219b. Zem219b is a pUC18-derived vector containing a mouse metallothionein promoter and a DHFR selectable  
25 marker operably linked to an SV40 promoter (Fig. 5). Plasmid Zem219b has been deposited with American Type Culture Collection as an *E. coli* XL1-blue transformant under Accession No. 68979. The entire  $\gamma'$ -chain coding sequence was then isolated as a 7.8 Kbp Sna BI fragment  
30 and inserted into the Sna BI site of pMAD6-Sna.

### Example III

Mice for initial breeding stocks (C57BL6J, CBACA) were obtained from Harlan Olac Ltd. (Bicester, UK).  
35 These were mated in pairs to produce F1 hybrid cross (B6CBAF1) for recipient female, superovulated females, stud males and vasectomized males. All animals were kept

on a 14 hour light/10 hour dark cycle and fed water and food (Special Diet Services RM3, Edinburgh, Scotland) *ad libitum*.

Transgenic mice were generated essentially as described in Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, 1986, which is incorporated herein by reference in its entirety. Female B6CBAF1 animals were superovulated at 4-5 weeks of age by an i.p. injection of pregnant mares' serum gonadotrophin (FOLLIGON, Vet-Drug, Falkirk, Scotland) (5 iu) followed by an i.p. injection of human chorionic gonadotrophin (CHORULON, Vet-Drug, Falkirk, Scotland) (5 iu) 45 hours later. They were then mated with a stud male overnight. Such females were next examined for copulation plugs. Those that had mated were sacrificed, and their eggs were collected for microinjection.

DNA was injected into the fertilized eggs as described in Hogan et al. (ibid.) Briefly, each of the vectors containing the  $A\alpha$ ,  $B\beta$  and  $\gamma$  expression units was digested with Mlu I, and the expression units were isolated by sucrose gradient centrifugation. All chemicals used were reagent grade (Sigma Chemical Co., St. Louis, MO, U.S.A.), and all solutions were sterile and nuclease-free. Solutions of 20% and 40% sucrose in 1 M NaCl, 20 mM Tris pH 8.0, 5 mM EDTA were prepared using UHP water and filter sterilized. A 30% sucrose solution was prepared by mixing equal volumes of the 20% and 40% solutions. A gradient was prepared by layering 0.5 ml steps of the 40%, 30% and 20% sucrose solutions into a 2 ml polyallomer tube and allowed to stand for one hour. 100  $\mu$ l of DNA solution (max. 8  $\mu$ g DNA) was loaded onto the top of the gradient, and the gradient was centrifuged for 17-20 hours at 26,000 rpm, 15°C in a Beckman TL100 ultracentrifuge using a TLS-55 rotor (Beckman Instruments, Fullerton, CA, USA). Gradients were fractionated by puncturing the tube bottom with a 20 ga. needle and collecting drops in a 96 well microtiter plate. 3  $\mu$ l

aliquots were analyzed on a 1% agarose mini-gel. Fractions containing the desired DNA fragment were pooled and ethanol precipitated overnight at -20°C in 0.3M sodium acetate. DNA pellets were resuspended in 50-100  $\mu$ l UHP  
5 water and quantitated by fluorimetry. The expression units were diluted in Dulbecco's phosphate buffered saline without calcium and magnesium (containing, per liter, 0.2 g KCl, 0.2 g  $\text{KH}_2\text{PO}_4$ , 8.0 g NaCl, 1.15 g  $\text{Na}_2\text{HPO}_4$ ), mixed (using either the N1-Beta or N3-Beta expression unit) in a  
10 1:1:1 molar ratio, concentration adjusted to about 6  $\mu$ g/ml, and injected into the eggs (~2 pl total DNA solution per egg).

Recipient females of 6-8 weeks of age are prepared by mating B6CBAF1 females in natural estrus with  
15 vasectomized males. Females possessing copulation plugs are then kept for transfer of microinjected eggs.

Following birth of potential transgenic animals, tail biopsies are taken, under anesthesia, at four weeks of age. Tissue samples are placed in 2 ml of tail buffer  
20 (0.3 M Na acetate, 50 mM HCl, 1.5 mM  $\text{MgCl}_2$ , 10 mM Tris-HCl, pH 8.5, 0.5% NP40, 0.5% Tween 20) containing 200  $\mu$ g/ml proteinase K (Boehringer Mannheim, Mannheim, Germany) and vortexed. The samples are shaken (250 rpm) at 55°-60° for 3 hours to overnight. DNA prepared from  
25 biopsy samples is examined for the presence of the injected constructs by PCR and Southern blotting. The digested tissue is vigorously vortexed, and 5  $\mu$ l aliquots are placed in 0.5 ml microcentrifuge tubes. Positive and negative tail samples are included as controls. Forty  $\mu$ l  
30 of silicone oil (BDH, Poole, UK) is added to each tube, and the tubes are briefly centrifuged. The tubes are incubated in the heating block of a thermal cycler (e.g. Omni-gene, Hybaid, Teddington, UK) to 95°C for 10 minutes. Following this, each tube has a 45  $\mu$ l aliquot of PCR mix  
35 added such that the final composition of each reaction mix is: 50 mM KCl; 2 mM  $\text{MgCl}_2$ ; 10 mM Tris-HCl (pH 8.3); 0.01% gelatin; 0.1% NP40, 10% DMSO; 500 nM each primer, 200  $\mu$ M

dNTPs; 0.02 U/ $\mu$ l Taq polymerase (Boehringer Mannheim, Mannheim, Germany). The tubes are then cycled through 30 repeated temperature changes as required by the particular primers used. The primers may be varied but in all cases must target the BLG promoter region. This is specific for the injected DNA fragments because the mouse does not have a BLG gene. Twelve  $\mu$ l of 5x loading buffer containing Orange G marker dye (0.25% Orange G [Sigma] 15% Ficoll type 400 [Pharmacia Biosystems Ltd., Milton Keynes, UK]) is then added to each tube, and the reaction mixtures are electrophoresed on a 1.6% agarose gel containing ethidium bromide (Sigma) until the marker dye has migrated 2/3 of the length of the gel. The gel is visualized with a UV light source emitting a wavelength of 254 nm. Transgenic mice having one or more of the injected DNA fragments are identified by this approach.

Positive tail samples are processed to obtain pure DNA. The DNA samples are screened by Southern blotting using a BLG promoter probe (nucleotides 2523-4253 of SEQ ID NO: 7). Specific cleavages with appropriate restriction enzymes (e.g. Eco RI) allow the distinction of the three constructs containing the A $\alpha$ , B $\beta$  and  $\gamma$  sequences.

Southern blot analysis of transgenic mice prepared essentially as described above demonstrated that more than 50% of progeny contained all three fibrinogen sequences. Examination of milk from positive animals by reducing SDS polyacrylamide gel electrophoresis demonstrated the presence of all three protein chains at concentrations up to 1 mg/ml. The amount of fully assembled fibrinogen was related to the ratios of individual subunits present in the milk. No apparent phenotype was associated with high concentrations of human fibrinogen in mouse milk.

#### 35 Example IV

Donor ewes are treated with an intravaginal progesterone-impregnated sponge (CHRONOGEST Goat Sponge,

Intervet, Cambridge, UK) on day 0. Sponges are left *in situ* for ten or twelve days.

Superovulation is induced by treatment of donor ewes with a total of one unit of ovine follicle stimulating hormone (OFSH) (OVAGEN, Horizon Animal  
5      Reproduction Technology Pty. Ltd., New Zealand) administered in eight intramuscular injections of 0.125 units per injection starting at 5:00 pm on day -4 and ending at 8:00 am on day 0. Donors are injected  
10     intramuscularly with 0.5 ml of a luteolytic agent (ESTRUMATE, Vet-Drug) on day -4 to cause regression of the corpus luteum, to allow return to estrus and ovulation. To synchronize ovulation, the donor animals are injected intramuscularly with 2 ml of a synthetic releasing hormone  
15     analog (RECEPTAL, Vet-Drug) at 5:00 pm on day 0.

Donors are starved of food and water for at least 12 hours before artificial insemination (A.I.). The animals are artificially inseminated by intrauterine laparoscopy under sedation and local anesthesia on day 1.  
20     Either xylazine (ROMPUN, Vet-Drug) at a dose rate of 0.05-0.1 ml per 10 kg bodyweight or ACP injection 10 mg/ml (Vet-Drug) at a dose rate of 0.1 ml per 10 kg bodyweight is injected intramuscularly approximately fifteen minutes before A.I. to provide sedation. A.I. is carried out  
25     using freshly collected semen from a Poll Dorset ram. Semen is diluted with equal parts of filtered phosphate buffered saline, and 0.2 ml of the diluted semen is injected per uterine horn. Immediately pre- or post-A.I., donors are given an intramuscular injection of AMOXPEN  
30     (Vet-Drug).

Fertilized eggs are recovered on day 2 following starvation of donors of food and water from 5:00 pm on day 1. Recovery is carried out under general anesthesia induced by an intravenous injection of 5% thiopentone sodium (INTRAVAL SODIUM, Vet-Drug) at a dose rate of 3 ml  
35     per 10 kg bodyweight. Anesthesia is maintained by inhalation of 1-2% Halothane/O<sub>2</sub>/N<sub>2</sub>O after intubation. To

recover the fertilized eggs, a laparotomy incision is made, and the uterus is exteriorized. The eggs are recovered by retrograde flushing of the oviducts with Ovum Culture Medium (Advanced Protein Products, Brierly Hill, West Midlands, UK) supplemented with bovine serum albumin of New Zealand origin. After flushing, the uterus is returned to the abdomen, and the incision is closed. Donors are allowed to recover post-operatively or are euthanized. Donors that are allowed to recover are given an intramuscular injection of Amoxypen L.A. at the manufacturer's recommended dose rate immediately pre- or post-operatively.

Plasmids containing the three fibrinogen chain expression units are digested with Mlu I, and the expression unit fragments are recovered and purified on sucrose density gradients. The fragment concentrations are determined by fluorimetry and diluted in Dulbecco's phosphate buffered saline without calcium and magnesium as described above. The concentration is adjusted to 6  $\mu\text{g/ml}$  and approximately 2  $\mu\text{l}$  of the mixture is microinjected into one pronucleus of each fertilized eggs with visible pronuclei.

All fertilized eggs surviving pronuclear microinjection are cultured in vitro at 38.5°C in an atmosphere of 5% CO<sub>2</sub>:5% O<sub>2</sub>:90% N<sub>2</sub> and about ~100% humidity in a bicarbonate buffered synthetic oviduct medium (see Table) supplemented with 20% v/v vasectomized ram serum. The serum may be heat inactivated at 56°C for 30 minutes and stored frozen at -20°C prior to use. The fertilized eggs are cultured for a suitable period of time to allow early embryo mortality (caused by the manipulation techniques) to occur. These dead or arrested embryos are discarded. Embryos having developed to 5 or 6 cell divisions are transferred to synchronized recipient ewes.

Table  
Synthetic Oviduct Medium

5	<u>Stock A (Lasts 3 Months)</u>	
	NaCl	6.29 g
	KCl	0.534 g
	KH <sub>2</sub> PO <sub>4</sub>	0.162 g
	MgSO <sub>4</sub> ·7H <sub>2</sub> O	0.182 g
10	Penicillin	0.06 g
	Sodium Lactate 60% syrup	0.6 mls
	Super H <sub>2</sub> O	99.4 mls
	<u>Stock B (Lasts 2 weeks)</u>	
15	NaHCO <sub>3</sub>	0.21 g
	Phenol red	0.001 g
	Super H <sub>2</sub> O	10 mls
	<u>Stock C (Lasts 2 weeks)</u>	
20	Sodium Pyruvate	0.051 g
	Super H <sub>2</sub> O	10 mls
	<u>Stock D (Lasts 3 months)</u>	
25	CaCl <sub>2</sub> ·2H <sub>2</sub> O	0.262 g
	Super H <sub>2</sub> O	10 mls
	<u>Stock E (Lasts 3 months)</u>	
30	Hepes	0.651 g
	Phenol red	0.001 g
	Super H <sub>2</sub> O	10 mls
	<u>To make up 10mls of Bicarbonate Buffered Medium</u>	
35	STOCK A	1 ml
	STOCK B	1 ml
	STOCK C	0.07 ml
	STOCK D	0.1 ml
	Super H <sub>2</sub> O	7.83 ml
40	Osmolarity should be 265-285 mOsm. Add 2.5 ml of heat inactivated sheep serum and filter sterilize.	
	<u>To make up 10 mls of HEPES Buffered Medium</u>	
45	STOCK A	1 ml
	STOCK B	0.2 ml
	STOCK C	0.07 ml
	STOCK D	0.1 ml
	STOCK E	0.8 ml
50	Super H <sub>2</sub> O	7.83 ml



Table, cont.

- 5 Osmolarity should be 265-285 mOsm.  
Add 2.5 ml of heat inactivated sheep serum  
and filter sterilize.

Recipient ewes are treated with an intravaginal progesterone-impregnated sponge (Chronogest Ewe Sponge or  
10 Chronogest Ewe-Lamb Sponge, Intervet) left *in situ* for 10 or 12 days. The ewes are injected intramuscularly with 1.5 ml (300 iu) of a follicle stimulating hormone substitute (P.M.S.G., Intervet) and with 0.5 ml of a luteolytic agent (Estrumate, Coopers Pitman-Moore) at  
15 sponge removal on day -1. The ewes are tested for estrus with a vasectomized ram between 8:00 am and 5:00 pm on days 0 and 1.

Embryos surviving *in vitro* culture are returned to recipients (starved from 5:00 pm on day 5 or 6) on day  
20 6 or 7. Embryo transfer is carried out under general anesthesia as described above. The uterus is exteriorized via a laparotomy incision with or without laparoscopy. Embryos are returned to one or both uterine horns only in ewes with at least one suitable corpora lutea. After  
25 replacement of the uterus, the abdomen is closed, and the recipients are allowed to recover. The animals are given an intramuscular injection of Amoxypen L.A. at the manufacturer's recommended dose rate immediately pre- or post-operatively.

30 Lambs are identified by ear tags and left with their dams for rearing. Ewes and lambs are either housed and fed complete diet concentrates and other supplements and or *ad lib.* hay, or are let out to grass.

Within the first week of life (or as soon  
35 thereafter as possible without prejudicing health), each lamb is tested for the presence of the heterologous DNA by two sampling procedures. A 10 ml blood sample is taken from the jugular vein into an EDTA vacutainer. If fit enough, the lambs also have a second 10 ml blood sample

taken within one week of the first. Tissue samples are taken by tail biopsy as soon as possible after the tail has become desensitized after the application of a rubber elastrator ring to its proximal third (usually within 200  
5 minutes after "tailing"). The tissue is placed immediately in a solution of tail buffer. Tail samples are kept at room temperature and analyzed on the day of collection. All lambs are given an intramuscular injection of Amoxypen L.A. at the manufacturer's  
10 recommended dose rate immediately post-biopsy, and the cut end of the tail is sprayed with an antibiotic spray.

DNA is extracted from sheep blood by first separating white blood cells. A 10 ml sample of blood is diluted in 20 ml of Hank's buffered saline (HBS; obtained  
15 from Sigma Chemical Co.). Ten ml of the diluted blood is layered over 5 ml of Histopaque (Sigma) in each of two 15 ml screw-capped tubes. The tubes are centrifuged at 3000 rpm (2000 x g max.), low brake for 15 minutes at room temperature. White cell interfaces are removed to a clean  
20 15 ml tube and diluted to 15 ml in HBS. The diluted cells are spun at 3000 rpm for 10 minutes at room temperature, and the cell pellet is recovered and resuspended in 2-5 ml of tail buffer.

To extract DNA from the white cells, 10% SDS is  
25 added to the resuspended cells to a final concentration of 1%, and the tube is inverted to mix the solution. One mg of fresh proteinase K solution is added, and the mixture is incubated overnight at 45°C. DNA is extracted using an equal volume of phenol/chloroform (x3) and  
30 chloroform/isoamyl alcohol (x1). The DNA is then precipitated by adding 0.1 volume of 3 M NaOAc and 2 volumes of ethanol, and the tube is inverted to mix. The precipitated DNA is spooled out using a clean glass rod with a sealed end. The spool is washed in 70% ethanol,  
35 and the DNA is allowed to partially dry, then is redissolved in TE (10 mM Tris-HCl, 1 mM EDTA, pH 7.4).

DNA samples from blood and tail are analyzed by Southern blotting using probes for the BLG promoter region and the fibrinogen chain coding regions.

From the foregoing, it will be appreciated that, 5 although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(i) APPLICANT: ZymoGenetics, Inc.  
1201 Eastlake Avenue East  
Seattle, Washington 98102  
United States of America

Pharmaceutical Proteins Ltd.  
Roslin  
Edinburgh  
Midlothian, Scotland EH25 9PP

(ii) TITLE OF INVENTION: Production of Fibrinogen in Transgenic  
Animals

(iii) NUMBER OF SEQUENCES: 27

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: ZymoGenetics, Inc.  
(B) STREET: 1201 Eastlake Avenue East  
(C) CITY: Seattle  
(D) STATE: WA  
(E) COUNTRY: USA  
(F) ZIP: 98102

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parker, Gary E  
(B) REGISTRATION NUMBER: 31-648  
(C) REFERENCE/DOCKET NUMBER: 93-15PC

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-442-6673  
(B) TELEFAX: 206-442-6678

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5943 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: Human Fibrinogen A-alpha chain

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: join(31..84, 1154..1279, 1739..1922, 3055..3200, 3786..5210)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GTCTAGGAGC CAGCCCCACC CTTAGAAAAG ATG TTT TCC ATG AGG ATC GTC TGC      54
                               Met Phe Ser Met Arg Ile Val Cys
                               1           5

CTA GTT CTA AGT GTG GTG GGC ACA GCA TGG GTATGGCCCT TTTCATTTTT      104
Leu Val Leu Ser Val Val Gly Thr Ala Trp
  10           15

TCTTCTTGCT TTCTCTCTGG TGTTTATTCC ACAAAGAGCC TGGAGGTCAG AGTCTACCTG      164

CTCTATGTCC TGACACACTC TTAGCTTTAT GACCCCAGGC CTGGGAGGAA ATTCCTGGG      224

TGGGCTTGAC ACCTCAAGAA TACAGGGTAA TATGACACCA AGAGGAAGAT CTTAGATGGA      284

TGAGAGTGTA CAACTACAAG GGAAACTTTA GCATCTGTCA TTCAGTCTTA CCACATTTTG      344

TTTTGTTTTG TTTTAAAAG GGCAAGAATT ATTTGCCATC CTTGTACCTA TAAAGCCTTG      404

GTGCATTATA ATGCTAGTTA ATGGAATAAA ACATTTTATG GTAAGATTTG TTTTCTTTAG      464

TTATTAATTT CTTGCTACTT GTCCATAATA AGCAGAACTT TTAGTGTTAG TACAGTTTTG      524

CTGAAAGGTT ATTGTTGTGT TTGTCAAGAC AGAAGAAAAA GCAAACGAAT TATCTTTGGA      584

AATATCTTTG CAGTATCAGA AGAGATTAGT TAGTAAGGCA ATACGCTTTT CCGCAGTAAT      644

```

GGTATTCTTT TAAATTATGA ATCCATCTCT AAAGGTTACA TAGAACTTG AAGGAGAGAG	704
GAACATTCAG TTAAGATAGT CTAGGTTTTT CTA CTGAAGC AGCAATTACA GGAGAAAGAG	764
CTCTACAGTA GTTTTCAACT TTCTGTCTGC AGTCATTAGT AAAAATGAAA AGGTAAAATT	824
TAACTGATTT TATAGATTCA AATAATTTTC CTTTTAGGAT GGATTCTTTA AAACTCCTAA	884
TATTTATCAA ATGCTTATTT AAGTGTACACA CACAGTTAAG AAATTTGTAC ACCTTGTCTC	944
CTTTAATTCT CATAACA ACT CCATAAAATG GGCCTAGGA TTTCCATTG AAGATAAGAA	1004
ACCTGAAGCT TGCCGAAGCC CTGTGTCTGC TCTCCTTAAT CTCTGTGAGA GTGCCATCTC	1064
TTCTGGGGA CTTGTAGGCA TGCCACTGTC TCCTCTTCTG GCTAACATTG CTGTTGCTCT	1124
CTTTTGTGTA TGTGAATGAA TCTTTAAAG ACT GCA GAT AGT GGT GAA GGT GAC	1177
Thr Ala Asp Ser Gly Glu Gly Asp	
20 25	
TTT CTA GCT GAA GGA GGA GGC GTG CGT GGC CCA AGG GTT GTG GAA AGA	1225
Phe Leu Ala Glu Gly Gly Gly Val Arg Gly Pro Arg Val Val Glu Arg	
30 35 40	
CAT CAA TCT GCC TGC AAA GAT TCA GAC TGG CCC TTC TGC TCT GAT GAA	1273
His Gln Ser Ala Cys Lys Asp Ser Asp Trp Pro Phe Cys Ser Asp Glu	
45 50 55	
GAC TGG GTAAGCAGTC AGCGGGGGAA GCAGGAGATT CCTTCCCTCT GATGCTAGAG	1329
Asp Trp	
60	
GGGCTCACAG GCTGACCTGA TTGGTCCCAG AAACTTTTTT AAATAGAAAA TAATTGAATA	1389
GTTACCTACA TAGCAAATAA AGAAAAGGAA CCTACTCCCA AGAGCACTGT TTATTTACCT	1449
CCCCA ACTCT GGATCATTAG TGGGTGAACA GACAGGATTT CAGTTGCATG CTCAGGCAAA	1509
ACCAGGCTCC TGAGTATTGT GGCCTCAATT TCCTGGCACC TATTTATGGC TAAGTGGACC	1569
CTCATTCCAG AGTTTCTCTG CGACCTCTAA CTAGTCCTCT TACCTACTTT TAAGCCA ACT	1629
TATCTGGAAG AGAAAGGGTA GGAAGAAATG GGGGCTGCAT GGAAACATGC AAAATTATTC	1689
TGAATCTGAG AGATAGATCC TTA CTGTAAAT TTTCTCCCTT CACTTTCAG AAC TAC	1744
Asn Tyr	

AAA TGC CCT TCT GGC TGC AGG ATG AAA GGG TTG ATT GAT GAA GTC AAT	1792
Lys Cys Pro Ser Gly Cys Arg Met Lys Gly Leu Ile Asp Glu Val Asn	
65 70 75	
CAA GAT TTT ACA AAC AGA ATA AAT AAG CTC AAA AAT TCA CTA TTT GAA	1840
Gln Asp Phe Thr Asn Arg Ile Asn Lys Leu Lys Asn Ser Leu Phe Glu	
80 85 90	
TAT CAG AAG AAC AAT AAG GAT TCT CAT TCG TTG ACC ACT AAT ATA ATG	1888
Tyr Gln Lys Asn Asn Lys Asp Ser His Ser Leu Thr Thr Asn Ile Met	
95 100 105 110	
GAA ATT TTG AGA GGC GAT TTT TCC TCA GCC AAT A GTAAGTATTA	1932
Glu Ile Leu Arg Gly Asp Phe Ser Ser Ala Asn	
115 120	
CATATTTACT TCTTTGACTT TATAACAGAA ACAACAAAA TCCTAAATAA ATATGATATC	1992
CGCTTATATC TATGACAATT TCATCCCAA GTACTTAGTG TAGAAACACA TACCTTCATA	2052
ATATCCCTGA AAATTTAAG AGGGAGCTTT TGTTCGTT ATTTTTCAA AGTAAAAGAT	2112
GTAACTGAG ATTGTTAAG GTCACAAAAT AAGTCAGAAT TTTGGATTAA AACAAGAATT	2172
TAAATGTGT CTTTTCAACA GTATATACTG AAAGTAGGAT GGGTCAGACT CTTTGAGTTG	2232
ATATTTTTGT TTCTGCTTG TAAAGGTGAA AACTGAGAGG TCAAGGAACT TGTTCAAAGA	2292
CACAGAGCTG GGAATTCAAC TCCCAGACTC CACTGAGCTG ATTAGGTAGA TTTTAAATT	2352
TAAATATAG GGTCAAGCTA CGTCATTCTC ACAGTCTACT CATTAGGGTT AGGAAACATT	2412
GCATTCACTC TGGGCATGGA CAGCGAGTCT AGGGAGTCCT CAGTTTCTCA AGTTTTGCTT	2472
TGCCTTTTTA CACCTTCACA AACACTTGAC ATTTAAATC AGTGATGCCA AACTAGCTG	2532
GCAAGTGAGT GATCCTGTTG ACCCAAACA GCTTAGGAAC CATTCAAAT CTATAGAGTT	2592
AAAAAGAAA GCTCATCAGT AAGAAAATCC AATATGTTCA AGTCCCTTGA TTAAGGATGT	2652
TATAAAATAA TTGAAATGCA ATCAAACCAA CTATTTTAACT TCCAAATTAC ACCTTTAAAA	2712
TTCAAAGAA AGTTCTTCTT CTATATTTCT TTGGGATTAC TAATTGCTAT TAGGACATCT	2772
TAACTGGCAT TCATGGAAGG CTGCAGGGCA TAACATTATC CAAAAGTCAA ATGCCCCATA	2832

GGTTTTGAAC TCACAGATTA AACTGTAACC AAAATAAAAT TAGGCATATT TACAAGCTAG 2892

TTTCTTTCTT TCTTTTTTCT CTTTCTTCTT TTCTTTCTTT CTTTCTTCTT TTCTTTCTTT 2952

CTTTCTTCTT TTCTCCTTCC TTCCTTCTT CTTTTCTTTT TTGCTGGCAA TTACAGACAA 3012

ATCACTCAGC AGCTACTTCA ATAACCATAT TTTCGATTTT AC CGT GAT AAT 3065  
 Asn Arg Asp Asn  
 125

ACC TAC AAC CGA GTG TCA GAG GAT CTG AGA AGC AGA ATT GAA GTC CTG 3113  
 Thr Tyr Asn Arg Val Ser Glu Asp Leu Arg Ser Arg Ile Glu Val Leu  
 130 135 140

AAG CGC AAA GTC ATA GAA AAA GTA CAG CAT ATC CAG CTT CTG CAG AAA 3161  
 Lys Arg Lys Val Ile Glu Lys Val Gln His Ile Gln Leu Leu Gln Lys  
 145 150 155

AAT GTT AGA GCT CAG TTG GTT GAT ATG AAA CGA CTG GAG GTAAGTATGT 3210  
 Asn Val Arg Ala Gln Leu Val Asp Met Lys Arg Leu Glu  
 160 165 170

GGCTGTGGTC CCGAGTGTCC TTGTTTTTGA GTAGAGGGAA AAGGAAGGCG ATAGTTATGC 3270

ACTGAGTGTC TACTATATGC AGAGAAAAGT GTTATATCCA TCATCTACCT AAAAGTAGGT 3330

ATTATTTTCC TCACTCCACA GTTGAAGAAA AAAAAATTCA GAGATATTAA GTAAATTTTC 3390

CAACGTACAT AGATAGTAAT TCAAAGCAAT GTTCAGTCCC TGTCTATTCC AAGCCATTAC 3450

ATCACCACAC CTCTGAGCCC TCAGCCTGAG TTCACCAAGG ATCATTTAAT TAGCGTTTCC 3510

TTTGAGAGGG AATAGCACCT TACTCTTGAT CCATTCTGAG GCTAAGATGA ATTAACAGC 3570

ATCCATTGCT TATCCTGGCT AGCCCTGCAA TACCCAACAT CTCTTCCACT GAGGGTGCTC 3630

GATAGGCAGA AAACAGAGAA TATTAAGTGG TAGGTCTCCG AGTCAAAAAA AATGAAACCA 3690

GTTTCCAGAA GGAAAATTAA CTACCAGGAA CTCAATAGAC GTAGTTTATG TATTTGTATC 3750

TACATTTTCT CTTTATTTTT CTCCCCTCTC TCTAG GTG GAC ATT GAT ATT AAG 3803  
 Val Asp Ile Asp Ile Lys  
 175



ATC CGA TCT TGT CGA GGG TCA TGC AGT AGG GCT TTA GCT CGT GAA GTA	3851
Ile Arg Ser Cys Arg Gly Ser Cys Ser Arg Ala Leu Ala Arg Glu Val	
180 185 190	
GAT CTG AAG GAC TAT GAA GAT CAG CAG AAG CAA CTT GAA CAG GTC ATT	3899
Asp Leu Lys Asp Tyr Glu Asp Gln Gln Lys Gln Leu Glu Gln Val Ile	
195 200 205	
GCC AAA GAC TTA CTT CCC TCT AGA GAT AGG CAA CAC TTA CCA CTG ATA	3947
Ala Lys Asp Leu Leu Pro Ser Arg Asp Arg Gln His Leu Pro Leu Ile	
210 215 220	
AAA ATG AAA CCA GTT CCA GAC TTG GTT CCC GGA AAT TTT AAG AGC CAG	3995
Lys Met Lys Pro Val Pro Asp Leu Val Pro Gly Asn Phe Lys Ser Gln	
225 230 235 240	
CTT CAG AAG GTA CCC CCA GAG TGG AAG GCA TTA ACA GAC ATG CCG CAG	4043
Leu Gln Lys Val Pro Pro Glu Trp Lys Ala Leu Thr Asp Met Pro Gln	
245 250 255	
ATG AGA ATG GAG TTA GAG AGA CCT GGT GGA AAT GAG ATT ACT CGA GGA	4091
Met Arg Met Glu Leu Glu Arg Pro Gly Gly Asn Glu Ile Thr Arg Gly	
260 265 270	
GGC TCC ACC TCT TAT GGA ACC GGA TCA GAG ACG GAA AGC CCC AGG AAC	4139
Gly Ser Thr Ser Tyr Gly Thr Gly Ser Glu Thr Glu Ser Pro Arg Asn	
275 280 285	
CCT AGC AGT GCT GGA AGC TGG AAC TCT GGG AGC TCT GGA CCT GGA AGT	4187
Pro Ser Ser Ala Gly Ser Trp Asn Ser Gly Ser Ser Gly Pro Gly Ser	
290 295 300	
ACT GGA AAC CGA AAC CCT GGG AGC TCT GGG ACT GGA GGG ACT GCA ACC	4235
Thr Gly Asn Arg Asn Pro Gly Ser Ser Gly Thr Gly Gly Thr Ala Thr	
305 310 315 320	
TGG AAA CCT GGG AGC TCT GGA CCT GGA AGT GCT GGA AGC TGG AAC TCT	4283
Trp Lys Pro Gly Ser Ser Gly Pro Gly Ser Ala Gly Ser Trp Asn Ser	
325 330 335	
GGG AGC TCT GGA ACT GGA AGT ACT GGA AAC CAA AAC CCT GGG AGC CCT	4331
Gly Ser Ser Gly Thr Gly Ser Thr Gly Asn Gln Asn Pro Gly Ser Pro	
340 345 350	
AGA CCT GGT AGT ACC GGA ACC TGG AAT CCT GGC AGC TCT GAA CGC GGA	4379
Arg Pro Gly Ser Thr Gly Thr Trp Asn Pro Gly Ser Ser Glu Arg Gly	
355 360 365	

AGT GCT GGG CAC TGG ACC TCT GAG AGC TCT GTA TCT GGT AGT ACT GGA Ser Ala Gly His Trp Thr Ser Glu Ser Ser Val Ser Gly Ser Thr Gly 370 375 380	4427
CAA TGG CAC TCT GAA TCT GGA AGT TTT AGG CCA GAT AGC CCA GGC TCT Gln Trp His Ser Glu Ser Gly Ser Phe Arg Pro Asp Ser Pro Gly Ser 385 390 395 400	4475
GGG AAC GCG AGG CCT AAC AAC CCA GAC TGG GGC ACA TTT GAA GAG GTG Gly Asn Ala Arg Pro Asn Asn Pro Asp Trp Gly Thr Phe Glu Glu Val 405 410 415	4523
TCA GGA AAT GTA AGT CCA GGG ACA AGG AGA GAG TAC CAC ACA GAA AAA Ser Gly Asn Val Ser Pro Gly Thr Arg Arg Glu Tyr His Thr Glu Lys 420 425 430	4571
CTG GTC ACT TCT AAA GGA GAT AAA GAG CTC AGG ACT GGT AAA GAG AAG Leu Val Thr Ser Lys Gly Asp Lys Glu Leu Arg Thr Gly Lys Glu Lys 435 440 445	4619
GTC ACC TCT GGT AGC ACA ACC ACC ACG CGT CGT TCA TGC TCT AAA ACC Val Thr Ser Gly Ser Thr Thr Thr Thr Arg Arg Ser Cys Ser Lys Thr 450 455 460	4667
GTT ACT AAG ACT GTT ATT GGT CCT GAT GGT CAC AAA GAA GTT ACC AAA Val Thr Lys Thr Val Ile Gly Pro Asp Gly His Lys Glu Val Thr Lys 465 470 475 480	4715
GAA GTG GTG ACC TCC GAA GAT GGT TCT GAC TGT CCC GAG GCA ATG GAT Glu Val Val Thr Ser Glu Asp Gly Ser Asp Cys Pro Glu Ala Met Asp 485 490 495	4763
TTA GGC ACA TTG TCT GGC ATA GGT ACT CTG GAT GGG TTC CGC CAT AGG Leu Gly Thr Leu Ser Gly Ile Gly Thr Leu Asp Gly Phe Arg His Arg 500 505 510	4811
CAC CCT GAT GAA GCT GCC TTC TTC GAC ACT GCC TCA ACT GGA AAA ACA His Pro Asp Glu Ala Ala Phe Phe Asp Thr Ala Ser Thr Gly Lys Thr 515 520 525	4859
TTC CCA GGT TTC TTC TCA CCT ATG TTA GGA GAG TTT GTC AGT GAG ACT Phe Pro Gly Phe Phe Ser Pro Met Leu Gly Glu Phe Val Ser Glu Thr 530 535 540	4907



ACCAGCTTAA TTTTTTTTTT AGACTGTGAT GATGTCCTCC AAACACATCC TTCAGGTACC 5907  
 CAAAGTGGCA TTTTCAATAT CAAGCTATCC GGATCC 5943

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 644 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Phe Ser Met Arg Ile Val Cys Leu Val Leu Ser Val Val Gly Thr  
 1 5 10 15  
 Ala Trp Thr Ala Asp Ser Gly Glu Gly Asp Phe Leu Ala Glu Gly Gly  
 20 25 30  
 Gly Val Arg Gly Pro Arg Val Val Glu Arg His Gln Ser Ala Cys Lys  
 35 40 45  
 Asp Ser Asp Trp Pro Phe Cys Ser Asp Glu Asp Trp Asn Tyr Lys Cys  
 50 55 60  
 Pro Ser Gly Cys Arg Met Lys Gly Leu Ile Asp Glu Val Asn Gln Asp  
 65 70 75 80  
 Phe Thr Asn Arg Ile Asn Lys Leu Lys Asn Ser Leu Phe Glu Tyr Gln  
 85 90 95  
 Lys Asn Asn Lys Asp Ser His Ser Leu Thr Thr Asn Ile Met Glu Ile  
 100 105 110  
 Leu Arg Gly Asp Phe Ser Ser Ala Asn Asn Arg Asp Asn Thr Tyr Asn  
 115 120 125  
 Arg Val Ser Glu Asp Leu Arg Ser Arg Ile Glu Val Leu Lys Arg Lys  
 130 135 140  
 Val Ile Glu Lys Val Gln His Ile Gln Leu Leu Gln Lys Asn Val Arg  
 145 150 155 160

43

Ala Gln Leu Val Asp Met Lys Arg Leu Glu Val Asp Ile Asp Ile Lys  
165 170 175

Ile Arg Ser Cys Arg Gly Ser Cys Ser Arg Ala Leu Ala Arg Glu Val  
180 185 190

Asp Leu Lys Asp Tyr Glu Asp Gln Gln Lys Gln Leu Glu Gln Val Ile  
195 200 205

Ala Lys Asp Leu Leu Pro Ser Arg Asp Arg Gln His Leu Pro Leu Ile  
210 215 220

Lys Met Lys Pro Val Pro Asp Leu Val Pro Gly Asn Phe Lys Ser Gln  
225 230 235 240

Leu Gln Lys Val Pro Pro Glu Trp Lys Ala Leu Thr Asp Met Pro Gln  
245 250 255

Met Arg Met Glu Leu Glu Arg Pro Gly Gly Asn Glu Ile Thr Arg Gly  
260 265 270

Gly Ser Thr Ser Tyr Gly Thr Gly Ser Glu Thr Glu Ser Pro Arg Asn  
275 280 285

Pro Ser Ser Ala Gly Ser Trp Asn Ser Gly Ser Ser Gly Pro Gly Ser  
290 295 300

Thr Gly Asn Arg Asn Pro Gly Ser Ser Gly Thr Gly Gly Thr Ala Thr  
305 310 315 320

Trp Lys Pro Gly Ser Ser Gly Pro Gly Ser Ala Gly Ser Trp Asn Ser  
325 330 335

Gly Ser Ser Gly Thr Gly Ser Thr Gly Asn Gln Asn Pro Gly Ser Pro  
340 345 350

Arg Pro Gly Ser Thr Gly Thr Trp Asn Pro Gly Ser Ser Glu Arg Gly  
355 360 365

Ser Ala Gly His Trp Thr Ser Glu Ser Ser Val Ser Gly Ser Thr Gly  
370 375 380

Gln Trp His Ser Glu Ser Gly Ser Phe Arg Pro Asp Ser Pro Gly Ser  
385 390 395 400

Gly Asn Ala Arg Pro Asn Asn Pro Asp Trp Gly Thr Phe Glu Glu Val  
405 410 415



## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8878 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: human fibrinogen B-beta chain

## (ix) FEATURE:

- (A) NAME/KEY: misc\_RNA
- (B) LOCATION: 1..469

## (ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 470..583

## (ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION: 584..3257

## (ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 3258..3449

## (ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION: 3450..3938

## (ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 3939..4122

## (ix) FEATURE:

- (A) NAME/KEY: intron
- (B) LOCATION: 4123..5042

## (ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 5043..5270

- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 5271..5830
  
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 5831..5944
  
- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 5945..6632
  
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 6633..6758
  
- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 6759..6966
  
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 6967..7252
  
- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 7253..7870
  
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 7871..8102
  
- (ix) FEATURE:
  - (A) NAME/KEY: 3'UTR
  - (B) LOCATION: 8103..8537
  
- (ix) FEATURE:
  - (A) NAME/KEY: misc\_RNA
  - (B) LOCATION: 8538..8878
  
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: join(470..583, 3258..3449, 3939..4122, 5043..5270, 5831..5944, 6633..6758, 6967..7252, 7871..8102)
  
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:



GAATTCATGC CCCTTTTGAA ATAGACTTAT GTCATTGTCA GAAAACATAA GCATTTATGG	60
TATATCATT A TGAGTCACG ATTTAGTGG TTGCCTTGTG AGTAGGTCAA ATTTACTAAG	120
CTTAGATTG TTTTCTCACA TATTCTTTCG GAGCTTGTGT AGTTTCCACA TTAATTTACC	180
AGAAACAAGA TACACACTCT CTTTGAGGAG TGCCCTAACT TCCCATCATT TTGTCCAATT	240
AAATGAATTG AAGAAATTTA ATGTTTCTAA ACTAGACCAA CAAAGAATAA TAGTTGTATG	300
ACAAGTAAAT AAGCTTTGCT GGAAGATGT TGCTTAAATG ATAAAATGGT TCAGCCAACA	360
AGTGAACCAA AAATTAATA TTAAC TAAGG AAAGGTAACC ATTTCTGAAG TCATTCCTAG	420
CAGAGGACTC AGATATATAT AGGATTGAAG ATCTCTCAGT TAAGTCTAC ATG AAA	475
	Met Lys
	1
AGG ATG GTT TCT TGG AGC TTC CAC AAA CTT AAA ACC ATG AAA CAT CTA	523
Arg Met Val Ser Trp Ser Phe His Lys Leu Lys Thr Met Lys His Leu	
5 10 15	
TTA TTG CTA CTA TTG TGT GTT TTT CTA GTT AAG TCC CAA GGT GTC AAC	571
Leu Leu Leu Leu Leu Cys Val Phe Leu Val Lys Ser Gln Gly Val Asn	
20 25 30	
GAC AAT GAG GAG GTGAATTTTT TAAAGCATT A TTATATTATT AGTAGTATTA	623
Asp Asn Glu Glu	
35	
TTAATATAAG ATGTAACATA ATCATATTAT GTGCTTATTT TAATGAAATT AGCATTGCTT	683
ATAGTTATGA AATGGAATTG TTAACCTCTG ACTTATTGTA TTTAAAGAAT GTTTCATAGT	743
ATTTCTTATA TAAAAACAAA GTAATTTCTT GTTTTCTAGT TTATCACCTT TGTTTTCTTA	803
AGATGAGGAT GGCTTAGCTA ATGTAAGATG TGTTTTTCTC ACTTGCTATT CTGAGTACTG	863
TGATTTTCAT TTA CTCTAG CAATACAGGA TTACAATTAA GAGGACAAGA TCTGAAAATC	923
TCACAAACTA TAAAATAATA AAAGAGCAGA ATTTTAAGAT AAAAGAACT GGTGGTAGGT	983
AGATTGTTCT TTGGTGAAGG AAGGTAATAT ATATTGTTAC TGAGATTACT ATTTATAAAA	1043
ATTATAACTA AGCCTAAAAG CAAAATACAT CAAGTGAAT GATAGAAAAT GAAATATTGC	1103

TTTTTTCAGA TGAAAAGTTC AAATTAGAGT TAGTGTGTAT TGTTATTATT AATAGTTATG	1163
AAACACGGTT CAGTCTAATT TATTTATTTG TAGAACAGTT TGTCCTCAAC TATTATTTTT	1223
GCTGACTTAT TGCTGTTAAT TTGCAGTTAC TAAAAATACA GAAATGCATT TAGGACAATG	1283
GATATTTAAG AAATTTAAAT TTTATCATCA AACGTATCAT GGCCAAATTT CTTACATATA	1343
GCATAGTATC ATTAAACTAG AAATAAGAAT ACACAATAAT ATTTAAATGA AGTGATTCAT	1403
TTCGGATCAT TATTGAGTTT CAAGGGAAC T GAGTGTGT ACTTATCAGA CTCTACATGT	1463
AAGAACATAT AGTTAATCTG GTTGTGTGTG TAAAAACATA TGGTTAATCT GGTAAAGTCT	1523
GGTTAATCAT ATTAGGTAAG AAAAATGTAA AGAATGTGTA AGACGAAATT TTTGTAAAGT	1583
ACTCTGCAAA GCACTTTCAC ATTTCTGCTT ATCAACTAAA CCTCACAGAG ATAGTTTAAT	1643
AGTTTAGGCT TTA AAAATGGA TTTTGATTAT TCAACAAGTG GCCTTCATAA TTTCTTTAAG	1703
TGTTTTTCTT TAAGTATATA CTTTCTTTAA ATATTTTTTA AAATTCCTT TTCTCTAGTA	1763
AAGCCAGACC ATCCATGCTA CCTCTCTAGT GGC ACTCTGA AATAAAAAGA AAATAGTTTT	1823
CTCTGTTATA ATTGTATTTG TAATAAGCAG ATGAATCACA TTTCTTAAAA TTTGTTTTAG	1883
AGAGGGTAAG CTCTGACTAG GACCATGACT TCAATGTGAA ATATGTATAT ATCCTCCGAA	1943
TCTTTACATA TTAAGAATGT ATATAGTCAA CTGGTTAAAC AGGAAAATCT GGAACAGCCT	2003
GGCTGGGTTT TAATCTTAGC ACCATCCTAC TAAATGTAA ATAATATTAT AATCTAATGA	2063
ATAAATGACA ATGCAATTCC AAATAGAGTT CATCTGATGA CTTCTAGACT CACAAAATTG	2123
CAAGAGAGCT CAGTTGTTGC TCAGTTGTTT CAAATCATGT CGTTTGTTAA TTTGTAATTA	2183
AGCTCCAAAG GATGTATAGC TACTGACAAA AAAAAAATG AGAATGTAGT TAATCCAAAT	2243
CAAACTTTC CTATTGCAAT GCGTATTTTC TGCTTCATTA TCCTTTAATA TAATATTTTA	2303
AGTTAGCAAG TAATTTTAAT TACAATGCAC AAGCCTTGAG AATTATTTTA AATATAAGAA	2363
AATCATAATG TTTGATAAAG AAATCATGTA AGAAATTTCA AGATAATGGT TTAACAAATA	2423
ATTTTGTGTA TAGAAGATAA GACTAAAAGT GAAATTCGAA GTGGAGAGGA CACTTAAACT	2483
GTAGTACTTG TTATGTGTGA TTCCAGTAAA AATAGTAATG AGCACTTATT ATTGCCAAGT	2543

ACTGTTCTGA GGGTACCATA TGCAATAAGT TATTTAATCC TTACAATAAT CTTGTAAGGC	2603
AGATTCAAAC TATCATTACA CTTATTTTAC AGATGAGAAA ACTGGGGCAC AGATAAAGCA	2663
ACTTGCCCAA GGTCTCATAG CTGTAAGTCA ACCCTACGGT CAAGACCTAC AAGTAGCCGA	2723
GCTCCAGAGT ACATTATGAG GGTCAAAGAT TGTCTTATTA CAAATAAATT CCAAGTAGAA	2783
TCAACCTTTA ATAAGTCTTT AATGTCTCTT AAATATGTTT ATATAGGAGT CTAATCACCA	2843
ATTCACAAAA ATGAAAGTAG GGAAATGATT AACAATAATC ATAGGAATCT AACAATCCAA	2903
GTGGCTTGAG AATATTCATT CTTCTTGACA GTATAGATTC TTTACAATTT CGTAAGTTCC	2963
AATGTATGTT TTAGGAATAT GAGGTCATTA CTATTCATAA TCTGATACAG CTTTATCCTA	3023
AGGCCTCTCT TTAAAACTA CACTGCATCA TAGCTTTTTT GTGCAGTTGG TCTTTCTACT	3083
GTTACTGAAC AGTAAGCAAC CTACAGATTC ACTATCACCA ACCAGCCAGT TGATGGATCT	3143
TAAGCAAATT ATCAAGCTTG TGATAACCTA AATTATAAAA TGAGGGTGTT GGAATAGTTA	3203
CATTCCAAAT CTTCTATAAC ACTCTGTATT ATATTTCTGC CTCATTCCTT GTAG GGT Gly	3260
TTC TTC AGT GCC CGT GGT CAT CGA CCC CTT GAC AAG AAG AGA GAA GAG	3308
Phe Phe Ser Ala Arg Gly His Arg Pro Leu Asp Lys Lys Arg Glu Glu	
40 45 50 55	
GCT CCC AGC CTG AGG CCT GCC CCA CCG CCC ATC AGT GGA GGT GGC TAT	3356
Ala Pro Ser Leu Arg Pro Ala Pro Pro Pro Ile Ser Gly Gly Gly Tyr	
60 65 70	
CGG GCT CGT CCA GCC AAA GCA GCT GCC ACT CAA AAG AAA GTA GAA AGA	3404
Arg Ala Arg Pro Ala Lys Ala Ala Ala Thr Gln Lys Lys Val Glu Arg	
75 80 85	
AAA GCC CCT GAT GCT GGA GGC TGT CTT CAC GCT GAC CCA GAC CTG	3449
Lys Ala Pro Asp Ala Gly Gly Cys Leu His Ala Asp Pro Asp Leu	
90 95 100	
GTGGGTGCAC TGATGTTTCT TGCAGTGGTG GCTCTCTCAT GCAGAGAAAG CCTGTAGTCA	3509
TGGCAGTCTG CTAATGTTTC ACTGACCCAC ATTACCATCA CTGTTATTTT GTTTGTAT	3569

TTTGAAATA AAATTCAAAA CATAACATA TTGGGCCTTT GGTTTAGGCT TTCTTTCTTG	3629
TTTTCTTTGG TCTGGGCCCA AAATTTCAAA TTAGGATATG TGGGTGCCAC CTTTCCATTT	3689
GTATTTTGCC ACTGCCTTTG TTTAGTTGGT AAAATTTTCA TAGCCCAATT ATATTTTTTC	3749
TGGGGTAAGT AATATTTTAA ATCTCTATGA GAGTATGATG ATGACTTTTCG AATTTCTGGT	3809
CTTACAGAAA ACCAAATAAT AAATTTTTAT GTTGGCTAAT CGTATCGCTG AATTTTCCTA	3869
TGTGCTATTT TAACAAATGT CCATGACCCA AATCCTTCAT CTAATGCCTG CTATTTTCTT	3929
TGTTTTTAG GGG GTG TTG TGT CCT ACA GGA TGT CAG TTG CAA GAG GCT	3977
Gly Val Leu Cys Pro Thr Gly Cys Gln Leu Gln Glu Ala	
105 110 115	
TTG CTA CAA CAG GAA AGG CCA ATC AGA AAT AGT GTT GAT GAG TTA AAT	4025
Leu Leu Gln Gln Glu Arg Pro Ile Arg Asn Ser Val Asp Glu Leu Asn	
120 125 130	
AAC AAT GTG GAA GCT GTT TCC CAG ACC TCC TCT TCT TCC TTT CAG TAC	4073
Asn Asn Val Glu Ala Val Ser Gln Thr Ser Ser Ser Ser Phe Gln Tyr	
135 140 145	
ATG TAT TTG CTG AAA GAC CTG TGG CAA AAG AGG CAG AAG CAA GTA AAA G	4122
Met Tyr Leu Leu Lys Asp Leu Trp Gln Lys Arg Gln Lys Gln Val Lys	
150 155 160	
GTAGATATCC TTGTGCTTTC CATTGATTT TCAGCTATAA AATTGGAACC GTTAGACTGC	4182
CACGAGAATG CATGGTTGTG AGAAGATTAA CATTCTGGG TTAGTGAATA GCATTCATAC	4242
GCTTTTGGGC ACCTTCCCCT GCAACTTGCC AGATAAGCAC TATTCAGCTC TTATTCCCAG	4302
TCTGACATCA GCAAGTGTGA TTTTCTATGA AAAATTCTAC TATGACTCCT TATTTTAAGT	4362
ATACAAGAAA CTTGTGACTC AGAAGATAAT ATTTACAGAG TGGAAAAAAA CCCCTAGCAT	4422
TTATAGTTTT AACATTTGAG GTTTTGAATG AGAGAGTTAT CCATAATATA TTCAATTGTG	4482
TTGTGGATAA TGACACCTAA CCTGTGAATC TTGAGGTCAG AATGTTGAGT GCTGTTGACT	4542
TGGTGGTCAG GAAACAGCTA GTGCGTGAGC CTGGCACAGG CATCTCAGTG AGTAGCATAC	4602
CCACAGTTGG AAATTTTCA AAGAAATCAA AGGAATCATG ACATCTTATA AATTTCAAGG	4662
TTCTGCTATA CTTATGTGAA ATGGATAAAT AAATCAAGCA TATCCACTCT GTAAGATTGA	4722





GGA TTT GGA AAT GTT GCA ACC AAC ACA GAT GGG AAG AAT TAC TGT GGC 6751  
 Gly Phe Gly Asn Val Ala Thr Asn Thr Asp Gly Lys Asn Tyr Cys Gly  
 305 310 315

CTA CCA G GTAACGAACA GGCATGCAAA ATAAAATCAT TCTATTTGAA ATGGGATTTT 6808  
 Leu Pro

TTTTAATTAA AAAACATTCA TTGTTGGAAG CCTGTTTTAG GCAGTTAAGA GGAGTTTCCT 6868  
 GACAAAAATG TGGAAGCTAA AGATAAGGGA AGAAAGGCAG TTTTAGTTT CCCAAAATTT 6928

TATTTTTGGT GAGAGATTTT ATTTTGTTTT TCTTTTAG GT GAA TAT TGG CTT 6980  
 Gly Glu Tyr Trp Leu  
 320

GGA AAT GAT AAA ATT AGC CAG CTT ACC AGG ATG GGA CCC ACA GAA CTT 7028  
 Gly Asn Asp Lys Ile Ser Gln Leu Thr Arg Met Gly Pro Thr Glu Leu  
 325 330 335 340

TTG ATA GAA ATG GAG GAC TGG AAA GGA GAC AAA GTA AAG GCT CAC TAT 7076  
 Leu Ile Glu Met Glu Asp Trp Lys Gly Asp Lys Val Lys Ala His Tyr  
 345 350 355

GGA GGA TTC ACT GTA CAG AAT GAA GCC AAC AAA TAC CAG ATC TCA GTG 7124  
 Gly Gly Phe Thr Val Gln Asn Glu Ala Asn Lys Tyr Gln Ile Ser Val  
 360 365 370

AAC AAA TAC AGA GGA ACA GCC GGT AAT GCC CTC ATG GAT GGA GCA TCT 7172  
 Asn Lys Tyr Arg Gly Thr Ala Gly Asn Ala Leu Met Asp Gly Ala Ser  
 375 380 385

CAG CTG ATG GGA GAA AAC AGG ACC ATG ACC ATT CAC AAC GGC ATG TTC 7220  
 Gln Leu Met Gly Glu Asn Arg Thr Met Thr Ile His Asn Gly Met Phe  
 390 395 400

TTC AGC ACG TAT GAC AGA GAC AAT GAC GGC TG GTATGTGTGG 7262  
 Phe Ser Thr Tyr Asp Arg Asp Asn Asp Gly Trp  
 405 410 415

CACTCTTTGC TCCTGCTTTA AAAATCACAC TAATATCATT ACTCAGAATC ATTAACAATA 7322

TTTTAAATAG CTACCACTTC CTGGGCACTT ACTGTCAGCC ACTGTCCTAA GCTCTTTATG 7382

CATCACTCGA AAGCATTTC ACTATAAGGT AGACATTCTT ATTCTCATTT TACAGATGAG 7442

ATTTAGAGAG ATTACGTGAT TTGTCCAATG TCACACAAC ACCCAGAGAT AAAACTAGAA 7502

TTTGAGCACA GTTACTTTCT GAATAATGAG CATTAGATA AATACCTATA TCTCTATATT	7562
CTAAAGTGTG TGTGAAAACCT TTCATTTTCA TTTCCAGGGT TCTCTGATAC TAAGGGTTGT	7622
AAAAGCTATT ATTCCAGTAT AAAGTAACAA ACACAGTCCC TAGATGGATT GCCACAAAGG	7682
CCCAGTTATC TCTCTTTCTT GCTATAGGGC ACAGGAGGTC TTTGGTGTAT TAGTGTGACT	7742
CTATGTATAG CACCCAAAGG AAAGACTACT GTGCACACGA GTGTAGCAGT CTTTTATGGG	7802
TAATCTGCAA AACGTAACCT GACCACCGTA GTTCTGTTTC TAATAACGCC AAACACATTT	7862
TCTTTCAG G TTA ACA TCA GAT CCC AGA AAA CAG TGT TCT AAA GAA GAC	7910
Leu Thr Ser Asp Pro Arg Lys Gln Cys Ser Lys Glu Asp	
420 425	
GGT GGT GGA TGG TGG TAT AAT AGA TGT CAT GCA GCC AAT CCA AAC GGC	7958
Gly Gly Gly Trp Trp Tyr Asn Arg Cys His Ala Ala Asn Pro Asn Gly	
430 435 440	
AGA TAC TAC TGG GGT GGA CAG TAC ACC TGG GAC ATG GCA AAG CAT GGC	8006
Arg Tyr Tyr Trp Gly Gly Gln Tyr Thr Trp Asp Met Ala Lys His Gly	
445 450 455 460	
ACA GAT GAT GGT GTA GTA TGG ATG AAT TGG AAG GGG TCA TGG TAC TCA	8054
Thr Asp Asp Gly Val Val Trp Met Asn Trp Lys Gly Ser Trp Tyr Ser	
465 470 475	
ATG AGG AAG ATG AGT ATG AAG ATC AGG CCC TTC TTC CCA CAG CAA TAGTCCCCAA	8109
Met Arg Lys Met Ser Met Lys Ile Arg Pro Phe Phe Pro Gln Gln	
480 485 490	
TACGTAGATT TTTGCTCTTC TGTATGTGAC AACATTTTTG TACATTATGT TATTGGAATT	8169
TTCTTTCATA CATTATATTC CTCTAAAACCT CTCAAGCAGA CGTGAGTGTG ACTTTTTGAA	8229
AAAAGTATAG GATAAATTAC ATTAAAATAG CACATGATTT TCTTTTGTTT TCTTCATTC	8289
TCTTGCTCAC CCAAGAAGTA ACAAAGTAT AGTTTTGACA GAGTTGGTGT TCATAATTC	8349
AGTTCTAGTT GATTGCGAGA ATTTTCAAAT AAGGAAGAGG GGTCTTTTAT CCTTGTCGTA	8409
GGAAAACCAT GACGGAAAGG AAAAAGTAT GTTTAAAAGT CCACTTTTAA AACTATATTT	8469
ATTTATGTAG GATCTGTCAA AGAAAACCTC CAAAAGATT TATTAATTAA ACCAGACTCT	8529



GTTGCAATAA GTTAATGTTT TCTTGTTTTG TAATCCACAC ATTCAATGAG TTAGGCTTTG 8589  
 CACTTGTAAG GAAGGAGAAG CGTTCACAAC CTCAAATAGC TAATAAACCG GTCTTGAATA 8649  
 TTTGAAGATT TAAAATCTGA CTCTAGGACG GGCACGGTGG CTCACGACTA TAATCCCAAC 8709  
 ACTTTGGGAG GCTGAGGCGG GCGGTCACAA GGTCAGGAGT TCAAGACCAG CCTGACCAAT 8769  
 ATGGTGAAC CCCATCTCTA CTAAAATAC AAAAATTAGC CAGGCGTGGT GGCAGGTGCC 8829  
 TGTAGGTCCC AGCTAGCCTG TGAGGTGGAG ATTGCATTGA GCCAAGATC 8878

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 491 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Lys Arg Met Val Ser Trp Ser Phe His Lys Leu Lys Thr Met Lys  
 1 5 10 15  
 His Leu Leu Leu Leu Leu Cys Val Phe Leu Val Lys Ser Gln Gly  
 20 25 30  
 Val Asn Asp Asn Glu Glu Gly Phe Phe Ser Ala Arg Gly His Arg Pro  
 35 40 45  
 Leu Asp Lys Lys Arg Glu Glu Ala Pro Ser Leu Arg Pro Ala Pro Pro  
 50 55 60  
 Pro Ile Ser Gly Gly Gly Tyr Arg Ala Arg Pro Ala Lys Ala Ala Ala  
 65 70 75 80  
 Thr Gln Lys Lys Val Glu Arg Lys Ala Pro Asp Ala Gly Gly Cys Leu  
 85 90 95  
 His Ala Asp Pro Asp Leu Gly Val Leu Cys Pro Thr Gly Cys Gln Leu  
 100 105 110

56

Gln Glu Ala Leu Leu Gln Gln Glu Arg Pro Ile Arg Asn Ser Val Asp  
 115 120 125

Glu Leu Asn Asn Asn Val Glu Ala Val Ser Gln Thr Ser Ser Ser Ser  
 130 135 140

Phe Gln Tyr Met Tyr Leu Leu Lys Asp Leu Trp Gln Lys Arg Gln Lys  
 145 150 155 160

Gln Val Lys Asp Asn Glu Asn Val Val Asn Glu Tyr Ser Ser Glu Leu  
 165 170 175

Glu Lys His Gln Leu Tyr Ile Asp Glu Thr Val Asn Ser Asn Ile Pro  
 180 185 190

Thr Asn Leu Arg Val Leu Arg Ser Ile Leu Glu Asn Leu Arg Ser Lys  
 195 200 205

Ile Gln Lys Leu Glu Ser Asp Val Ser Ala Gln Met Glu Tyr Cys Arg  
 210 215 220

Thr Pro Cys Thr Val Ser Cys Asn Ile Pro Val Val Ser Gly Lys Glu  
 225 230 235 240

Cys Glu Glu Ile Ile Arg Lys Gly Gly Glu Thr Ser Glu Met Tyr Leu  
 245 250 255

Ile Gln Pro Asp Ser Ser Val Lys Pro Tyr Arg Val Tyr Cys Asp Met  
 260 265 270

Asn Thr Glu Asn Gly Gly Trp Thr Val Ile Gln Asn Arg Gln Asp Gly  
 275 280 285

Ser Val Asp Phe Gly Arg Lys Trp Asp Pro Tyr Lys Gln Gly Phe Gly  
 290 295 300

Asn Val Ala Thr Asn Thr Asp Gly Lys Asn Tyr Cys Gly Leu Pro Gly  
 305 310 315 320

Glu Tyr Trp Leu Gly Asn Asp Lys Ile Ser Gln Leu Thr Arg Met Gly  
 325 330 335

Pro Thr Glu Leu Leu Ile Glu Met Glu Asp Trp Lys Gly Asp Lys Val  
 340 345 350

Lys Ala His Tyr Gly Gly Phe Thr Val Gln Asn Glu Ala Asn Lys Tyr  
 355 360 365

Gln Ile Ser Val Asn Lys Tyr Arg Gly Thr Ala Gly Asn Ala Leu Met  
 370 375 380  
 Asp Gly Ala Ser Gln Leu Met Gly Glu Asn Arg Thr Met Thr Ile His  
 385 390 395 400  
 Asn Gly Met Phe Phe Ser Thr Tyr Asp Arg Asp Asn Asp Gly Trp Leu  
 405 410 415  
 Thr Ser Asp Pro Arg Lys Gln Cys Ser Lys Glu Asp Gly Gly Gly Trp  
 420 425 430  
 Trp Tyr Asn Arg Cys His Ala Ala Asn Pro Asn Gly Arg Tyr Tyr Trp  
 435 440 445  
 Gly Gly Gln Tyr Thr Trp Asp Met Ala Lys His Gly Thr Asp Asp Gly  
 450 455 460  
 Val Val Trp Met Asn Trp Lys Gly Ser Trp Tyr Ser Met Arg Lys Met  
 465 470 475 480  
 Ser Met Lys Ile Arg Pro Phe Phe Pro Gln Gln  
 485 490

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10564 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: human fibrinogen gamma chain

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: join(1799..1876, 1973..2017, 2207..2390, 2510  
 ..2603, 4211..4341, 4645..4778, 5758..5942, 7426  
 ..7703, 9342..9571)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CTACACACTT CTTGAAGGCA AAGGCAATGC TGAAGTCACC TTTCATGTTT AAATCATATT	60
AAAAAGTTAG CAAGATGTAA TTATCAGTGT ACTATGTAAA TCTTTGTGAA TGATCAATAA	120
TTACATATTT TCATTATATA TATTTTAGTA GATAATATTT ATATACATTC AACATTCTAA	180
ATATAGAAAAG TTTACAGAGA AAAATAAAGC CTTTTTTTCC AATCCTGTCC TCCACCTCTG	240
CATCCCATTG TTCTTCACAG AGGCAACTGA TTCAAGTCAT TACATAGTTA TTGAGTGTTA	300
ACTACAATA TGTTAAGTAC AGCTATATAT GTTAGATGCC GTAGCCACAG AAATCAGTTT	360
ACAATCTAAT GCAGTGGATA CAGCATGTAT ACATATAATA TAAGGTTGCT ACAAATGCTA	420
TCTGAGGTAG AGCTGTTTGA AAGAATACTA AACTTAAAT GTTTAATTCA ACTGACTTGA	480
TTGACAACTG ATTAGCTGAG TGGAAAAGAT GGATGAGAAA GATTGTGAGA CTTAATTGGC	540
TGGTGGTATG GTGATATGAT TGACAATAAC TGCTAAGTCA GAGAGGGATA TATTAAGGAG	600
GAGAAGAAAA GCAACAAATC TGGTTTTGAT GTGTTCACTT TGTATAATT ATTGATTATT	660
TACTGAATAT GAATATTTAT CTTTGTTTTT GAGTCAATAA ATATACCTTT GTAAAGACAG	720
AATTAAGTA TTAGTATTTT TTTCAAATG GAGGCATTTT TCCCACTAAC ATATTTTCATC	780
AAAACCTATA ATAAGCTTGG TTCCAGAGGA AGAAATGAGG GATAACCAAA AATAGAGACA	840
TTAATAATAG TGTAACGCCC AGTGATAAAT CTCAATAGGC AGTGATGACA GACATGTTTT	900
CCCAAACACA AGGATGCTGT AAGGGCCAAA CAGAAATGAT GGCCCTCCC CAGCACCTCA	960
TTTTGCCCT TCCTCAGCT ATGCCTCTAC TCTCCTTTAG ATACAAGGA GGTGGATTTT	1020
TCTCTTCTCT GAGATAGCTT GATGGAACCA CAGGAACAAT GAAGTGGGCT CCTGGCTCTT	1080
TTCTCTGTGG CAGATGGGGT GCCATGCCCA CCTTCAGACA AAGGGAAGAT TGAGCTCAAA	1140
AGCTCCCTGA GAAGTGAGAG CCTATGAACA TGGTTGACAC AGAGGGACAG GAATGTATTT	1200
CCAGGGTCAT TCATTCCTGG GAATAGTGAA CTGGGACATG GGGGAAGTCA GTCTCCTCT	1260
GCCACAGCCA CAGATTAATA ATAATAATGT TAACTGATCC CTAGGCTAAA ATAATAGTGT	1320
TAACTGATCC CTAAGCTAAG AAAGTTCTTT TGTAATTCA GGTGATGGCA GCAGGACCCA	1380

TCTTAAGGAT AGACTAGGTT TGCTTAGTTC GAGGTCATAT CTGTTTGCTC TCAGCCATGT 1440  
 ACTGGAAGAA GTTGCATCAC ACAGCCTCCA GGACTGCCCT CCTCCTCACA GCAATGGATA 1500  
 ATGCTTCACT AGCCTTTGCA GATAATTTTG GATCAGAGAA AAAACCTTGA GCTGGGCCAA 1560  
 AAAGGAGGAG CTTCAACCTG TGTGCAAAAT CTGGGAACCT GACAGTATAG GTTGGGGGCC 1620  
 AGGATGAGGA AAAAGGAACG GGAAAGACCT GCCCACCTT CTGGTAAGGA GGCCCCGTGA 1680  
 TCAGCTCCAG CCATTTGCAG TCCTGGCTAT CCCAGGAGCT TACATAAAGG GACAATTGGA 1740  
 GCCTGAGAGG TGACAGTGCT GACACTACAA GGCTCGGAGC TCCGGGCACT CAGACATC 1798  
 ATG AGT TGG TCC TTG CAC CCC CGG AAT TTA ATT CTC TAC TTC TAT GCT 1846  
 Met Ser Trp Ser Leu His Pro Arg Asn Leu Ile Leu Tyr Phe Tyr Ala  
 1 5 10 15  
 CTT TTA TTT CTC TCT TCA ACA TGT GTA GCA GTAAGTGTGC TCTTCACAAA 1896  
 Leu Leu Phe Leu Ser Ser Thr Cys Val Ala  
 20 25  
 ACGTTGTTTA AAATGGAAAG CTGGAAAATA AACAGATAA TAACTAGTG AAATTTTCGT 1956  
 ATTTTTTCTC TTTTAG TAT GTT GCT ACC AGA GAC AAC TGC TGC ATC TTA 2005  
 Tyr Val Ala Thr Arg Asp Asn Cys Cys Ile Leu  
 30 35  
 GAT GAA AGA TTC GTAAGTAGTT TTTATGTTTC TCCCTTTGTG TGTGAACTGG 2057  
 Asp Glu Arg Phe  
 40  
 AGAGGGGCAG AGGAATAGAA ATAATCCCT CATAAATATC ATCTGGCACT TGTAACTTTT 2117  
 TAAAACATA GTCTAGGTTT TACCTATTTT TCTTAATAGA TTTAAGAGT AGCATCTGTC 2177  
 TACATTTTTA ATCACTGTTA TATTTTCAG GGT AGT TAT TGT CCA ACT ACC TGT 2230  
 Gly Ser Tyr Cys Pro Thr Thr Cys  
 45  
 GGC ATT GCA GAT TTC CTG TCT ACT TAT CAA ACC AAA GTA GAC AAG GAT 2278  
 Gly Ile Ala Asp Phe Leu Ser Thr Tyr Gln Thr Lys Val Asp Lys Asp  
 50 55 60 65

60

CTA CAG TCT TTG GAA GAC ATC TTA CAT CAA GTT GAA AAC AAA ACA TCA Leu Gln Ser Leu Glu Asp Ile Leu His Gln Val Glu Asn Lys Thr Ser	2326
70 75 80	
GAA GTC AAA CAG CTG ATA AAA GCA ATC CAA CTC ACT TAT AAT CCT GAT Glu Val Lys Gln Leu Ile Lys Ala Ile Gln Leu Thr Tyr Asn Pro Asp	2374
85 90 95	
GAA TCA TCA AAA CCA A GTGAGAAAAT AAAGACTACT GACCAAAAAA Glu Ser Ser Lys Pro	2420
100	
TAATAATAAT AATCTGTGAA GTTCTTTTGC TGGTGTTTTA GTTGTCTAT TTGCTTAAGG	2480
ATTTTTATGT CTCTGATCCT ATATTACAG AT ATG ATA GAC GCT GCT ACT TTG Asn Met Ile Asp Ala Ala Thr Leu	2532
105 110	
AAG TCC AGG ATA ATG TTA GAA GAA ATT ATG AAA TAT GAA GCA TCG ATT Lys Ser Arg Ile Met Leu Glu Glu Ile Met Lys Tyr Glu Ala Ser Ile	2580
115 120 125	
TTA ACA CAT GAC TCA AGT ATT CG GTAAGGATTT TTGTTTTAAT TTGCTCTGCA Leu Thr His Asp Ser Ser Ile Arg	2633
130	
AGACTGATTT AGTTTTTATT TAATATTCTA TACTTGAGTG AAAGTAATTT TTAATGTGTT	2693
TTCCCCATTT ATAATATCCC AGTGACATTA TGCCTGATTA TGGTGGAGCAT AGTAGAGATA	2753
GAAGTTTTTA GTGCAATATA AATTATACTG GGTTATAATT GCTTATTAAT AATCACATTG	2813
AAGAAAGATG TTCTAGATGT CTCAAATGC TAGTTTGACC ATATTTATCA AAAATTTTTT	2873
CCCCATCCCC CATTTATCTT ACAACATAAA ATCAATCTCA TAGGAATTTG GGTGTTGAAA	2933
ATAAAATCCT CTTTATAAAA ATGCTGACAA ATTGGTGGTT AAAAAAATTA GCAAGCAGAG	2993
GCATAGTAAG GATTTTGGCT CCTAAAGTAA ATTATATTGA ATGTGGAGCA GGAAGAAACA	3053
TGTCTTGAGA GACTAAGTGT GGCAAATATT GCAAAGCTCA TATTGATCAT TGCAGAAATGA	3113
ACCTGCATAG TCTCTTCCCT TCATTTGGAA GTGAATGTCT CTGTTAAAGC TTCTCAGGGA	3173
CTCATAAACT TTCTGAACAT AAGGTCTCAG ATACAGTTTT AATATTTTTT CCAATTTTTT	3233
TTTTCTGAAT TTTTCTCAA GCAGCTTGAG AAATTGAGAT AAATAGTAGC TAGGGAGAAG	3293

TGCCCAGGA AAGATTTCTC CTCTTTTTC TATCAGAGGG CCCTTGTTAT TATTGTTATT	3353
ATTATTACTT GCATTATTAT TGTCCATCAT TGAAGTTGAA GGAGGTTATT GTACAGAAAT	3413
TGCCTAAGAC AAGGTAGAGG GAAAACGTGG ACAAATAGTT TGTCTACCCT TTTTACTTC	3473
AAAGAAAGAA CGGTTTATGC ATTGTAGACA GTTTTCTATC ATTTTTGGAT ATTTGCAAGC	3533
CACCCTGTAA GTAACATAA AAGGAGGGT TTTACTTCCC CCAGTCCATT CCCAAAGCTA	3593
TGTAACCAGA AGCATTAAAG AAGAAAGGGG AAGTATCTGT TGTTTTATTT TACATACAAT	3653
AACGTTCCAG ATCATGTCCC TGTGTAAGTT ATATTTTAGA TTGAAGCTTA TATGTATAGC	3713
CTCAGTAGAT CCACAAGTGA AAGGTATACT CCTTCAGCAC ATGTGAATTA CTGAACTGAG	3773
CTTTTCCTGC TTCTAAAGCA TCAGGGGGTG TTCCTATTAA CCAGTCTCGC CACTCTTGCA	3833
GGTTGCTATC TGCTGTCCCT TATGCATAAA GTAAAAGCA AAATGTCAAT GACATTTGCT	3893
TATTGACAAG GACTTTGTTA TTTGTGTTGG GAGTTGAGAC AATATGCCCC ATTCTAAGTA	3953
AAAAGATTCA GGTCCACATT GTATTCCTGT TTTAATTGAT TTTTGATTT GTTTTCTTT	4013
TTCAAAAAGT TTATAATTTT AATTCATGTT AATTTAGTAA TATAATTTTA CATTTTCCTC	4073
AAGAATGGAA TAATTTATCA GAAAGCACTT CTTAAGAAAA TACTTAGCAG TTTCCAAAGA	4133
AAATATAAAA TTAATCTTCT GAAAGGAATA CTTATTTTTG TCTTCTTATT TTTGTTATCT	4193
TATGTTTCTG TTTGTAG A TAT TTG CAG GAA ATA TAT AAT TCA AAT AAT CAA	4244
	Tyr Leu Gln Glu Ile Tyr Asn Ser Asn Asn Gln
	135 140 145
AAG ATT GTT AAC CTG AAA GAG AAG GTA GCC CAG CTT GAA GCA CAG TGC	4292
Lys Ile Val Asn Leu Lys Glu Lys Val Ala Gln Leu Glu Ala Gln Cys	
	150 155 160
CAG GAA CCT TGC AAA GAC ACG GTG CAA ATC CAT GAT ATC ACT GGG AAA G	4341
Gln Glu Pro Cys Lys Asp Thr Val Gln Ile His Asp Ile Thr Gly Lys	
	165 170 175
GTAACCTGATG AAGGTTATAT TGGGATTAGG TTCATCAAAG TAAGTAATGT AAAGGAGAAA	4401
GTATGTA CTG GAAAGTATAG GAATAGTTTA GAAAGTGGCT ACCCATTAAG TCTAAGAATT	4461

TCAGTTGTCT AGACCTTTCT TGAATAGCTA AAAAAACAG TTAAAAGGA ATGCTGATGT	4521
GAAAAGTAAG AAAATTATTC TTGGAAAATG AATAGTTTAC TACATGTAA AAGCTATTTT	4581
TCAAGGCTGG CACAGTCTTA CCTGCATTTT AAACCACAGT AAAAGTCGAT TCTCCTTCTC	4641
TAG AT TGT CAA GAC ATT GCC AAT AAG GGA GCT AAA CAG AGC GGG CTT	4688
Asp Cys Gln Asp Ile Ala Asn Lys Gly Ala Lys Gln Ser Gly Leu	
180 185 190	
TAC TTT ATT AAA CCT CTG AAA GCT AAC CAG CAA TTC TTA GTC TAC TGT	4736
Tyr Phe Ile Lys Pro Leu Lys Ala Asn Gln Gln Phe Leu Val Tyr Cys	
195 200 205	
GAA ATC GAT GGG TCT GGA AAT GGA TGG ACT GTG TTT CAG AAG	4778
Glu Ile Asp Gly Ser Gly Asn Gly Trp Thr Val Phe Gln Lys	
210 215 220	
GTAATTTTTT CCCCACCATG TGTATTTAAT AAATTCCTAC ATTGTTTCTG CCATATGGCA	4838
GATACTTTTC TAAGCACCTT GTGAACCGTA GCTCATTAA TCCTTGCAAT AGCCCTAAGA	4898
GGAAGGTACT TCTGTTACTC CTATTTACAG AAAAGGAAAC TGAGGCACAC AAGGTTAAAT	4958
AACTTGCCA AGACCACATA ACTAATAAGC AACAGAGTCA GCATTTGAAC CTAGGCAGTA	5018
TAGTTTCAGA GTTTGTGACT TGA CTCTATA TTGACTGGC ACTGACTTTG TAGATTCATG	5078
GTGGCACATA ATCATAGTAC CACAGTGACA AATAAAAAGA AGGAACTCT TTTGTCAGGT	5138
AGGTCAAGAC CTGAGGTTTC CCATCACAAG ATGAGGAAGC CCAACACCAC CCCCACCAC	5198
CCCACCACCA TCACCACCCT TTCACACACC AGAGGATACA CTTGGGCTGC TCCAAGACAA	5258
GGAACCTGTG TTGCATCTGC CACTTGCTGA TACCCACTAG GAATCTTGGC TCCTTTACTT	5318
TCTGTTTACC TCCCACCACT GTTATAACTG TTTCTACAGG GGGCGCTCAG AGGGAATGAA	5378
TGGTGAAGC ATTAGTTGCC AGACACCGAT TGAGCAATGG GTTCCATCAT AAGTGTAAGA	5438
ATCAGTAATA TCCAGCTAGA GTTCTGAAGT CGTCTAGGTG TCTTTTAAAT ATTACCACTC	5498
ATTTAGAATT TATGATGTGC CAGAAACCCT CTTAAGTATT TCTCTTATAT TCTCTCTCAT	5558
GATCCTTGCA GCAACCCTAA GAAGTAACCA TCATTTTTCC TATTTGATAC ATGAGGAAAC	5618
TGAGGTAGCT TGGCCAAGAT CACTTAGTTG GGAGTTGATA GAACCAGTGC TCTGTATTTT	5678



TGACAAAATG TTGACAGCAT TCTCTTTACA TGCATTGATA GTCTATTTTC TCCTTTTGCT	5738
CTTGCAAATG TGTAATTAG AGA CTT GAT GGC AGT GTA GAT TTC AAG AAA AAC	5790
Arg Leu Asp Gly Ser Val Asp Phe Lys Lys Asn	
225 230	
TGG ATT CAA TAT AAA GAA GGA TTT GGA CAT CTG TCT CCT ACT GGC ACA	5838
Trp Ile Gln Tyr Lys Glu Gly Phe Gly His Leu Ser Pro Thr Gly Thr	
235 240 245	
ACA GAA TTT TGG CTG GGA AAT GAG AAG ATT CAT TTG ATA AGC ACA CAG	5886
Thr Glu Phe Trp Leu Gly Asn Glu Lys Ile His Leu Ile Ser Thr Gln	
250 255 260 265	
TCT GCC ATC CCA TAT GCA TTA AGA GTG GAA CTG GAA GAC TGG AAT GGC	5934
Ser Ala Ile Pro Tyr Ala Leu Arg Val Glu Leu Glu Asp Trp Asn Gly	
270 275 280	
AGA ACC AG GTACTGTTTT GAAATGACTT CCAACTTTTT ATTGTAAAGA	5982
Arg Thr Ser	
TTGCCTGGAA TGTGCACTTT CCAACTATCA ATAGACAATG GCAAATGCAG CCTGACAAAT	6042
GCAAACAGCA CATCCAGCCA CCATTTTCTC CAGGAGTCTG TTTGGTTCTT GGGCAATCCA	6102
AAAAGGTAAA TTCTATTCAG GATGAATCTA AGTGTATTGG TACAATCTAA TTACCCTGGA	6162
ACCATTCAGA GTAATAGCTA ATTACTGAAC TTTAATCAG TCCCAGGAAT TGAGCATAAA	6222
ATTATAATTT TATCTAGTCT AAATTACTAT TTCATGAAGC AGGTATTATT ATTAATCCCA	6282
TTTTATAGAT TAACTTGCTC AAAGTCACAT TGCTGATAAG TGGTAGAGGT AGAATTCAGA	6342
CTCAAGTAGT TTAACCTTAG AGCCTGTCCT CTTAACAACT ATCCTGGTTG AAAAGCAAAT	6402
ACAGCCTCTT CAGACTTCTC AGTGCCTTGA TGGCCATTTA TTCTGTCAA TCATGAGCTA	6462
CCCTAAAAGT AAACCAGCTA GCTCTTTTGA TGATCTAGAG GCTTCTTTTT GCTTGAGATA	6522
TTTGAAGGTT TTAAGCATTG TTACCTAATT AAAATGCAGA AAAATATCCA ACCCTCTTGT	6582
TATGTTTAAAG GAATAGTGAA ATATATTGTC TTCAAACACA TGGACTTTTT TTTATTGCTT	6642
GGTTGGTTTT TAATCCAGAA AGTGCTATAG TCAGTAGACC TTCTTCTAGG AAAGGACCTT	6702

CCATTTCCCA GCCACTGGAG ATTAGAAAAT AAGCTAAATA TTTTCTGGAA ATTTCTGTTC	6762
ATTCATTAAG GCCCATCCTT TCCCCACTC TATAGAAGTG TTGTCCACTT GCACAATTTT	6822
TTCCAGGAAA GAATCTCTCT AACTCCTTCA GCTCACATGC TTTGGACCAC ACAGGGAAGA	6882
CTTTGATTGT GTAATGCCCT CAGAAGCTCT CTTTCTTGCC ACTACCACAC TGATTTGAGG	6942
AAGAAAATCC CTTTAGCACC TAACCCTTCA GGTGCTATGA GTGGCTAATG GAACTGTACC	7002
TCCTTCAAGT TTTGTGCAAT AATTAAGGGT CACTCACTGT CAGATACTTT CTGTGATCTA	7062
TGATAATGTG TGTGCAACAC ATAACATTTT AATAAAAGTA GAAAATATGA AATTAGAGTC	7122
ATCTACACAT CTGGATTTGA TCTTAGAATG AAACAAGCAA AAAAGCATCC AAGTGAGTGC	7182
AATTATTAGT TTTCAGAGAT GCTTCAAAGG CTTCTAGGCC CATCCCGGGA AGTGTTAATG	7242
AGCTGTGGAC TGGTTCACAT ATCTATTGCC TCTTGCCAGA TTTGCAAAA ACTTCACTCA	7302
ATGAGCAAAT TTCAGCCTTA AGAAACAAAG TCAAAAATTC CAAGGAAGCA TCCTACGAAA	7362
GAGGGAECTT CTGAGATCCC TGAGGAGGGT CAGCATGTGA TGGTTGTATT TCCTTCTTCT	7422
CAG T ACT GCA GAC TAT GCC ATG TTC AAG GTG GGA CCT GAA GCT GAC	7468
Thr Ala Asp Tyr Ala Met Phe Lys Val Gly Pro Glu Ala Asp	
285 290 295	
AAG TAC CGC CTA ACA TAT GCC TAC TTC GCT GGT GGG GAT GCT GGA GAT	7516
Lys Tyr Arg Leu Thr Tyr Ala Tyr Phe Ala Gly Gly Asp Ala Gly Asp	
300 305 310	
GCC TTT GAT GGC TTT GAT TTT GGC GAT GAT CCT AGT GAC AAG TTT TTC	7564
Ala Phe Asp Gly Phe Asp Phe Gly Asp Asp Pro Ser Asp Lys Phe Phe	
315 320 325 330	
ACA TCC CAT AAT GGC ATG CAG TTC AGT ACC TGG GAC AAT GAC AAT GAT	7612
Thr Ser His Asn Gly Met Gln Phe Ser Thr Trp Asp Asn Asp Asn Asp	
335 340 345	
AAG TTT GAA GGC AAC TGT GCT GAA CAG GAT GGA TCT GGT TGG TGG ATG	7660
Lys Phe Glu Gly Asn Cys Ala Glu Gln Asp Gly Ser Gly Trp Trp Met	
350 355 360	
AAC AAG TGT CAC GCT GGC CAT CTC AAT GGA GTT TAT TAC CAA G	7703
Asn Lys Cys His Ala Gly His Leu Asn Gly Val Tyr Tyr Gln	
365 370 375	

GTATGTTTT CTTTCTTAGA TTCCAAGTTA ATGTATAGTG TATACTATTT TCATAAAAAA 7763  
TAATAAATAG ATATGAAGAA ATGAAGAATA ATTTATAAAG ATAGTAGGGA TTTTATCATG 7823  
TTCTTTATTT CAACTAAGTT CTTTGAAACT GGAAGTGGAT AATACCAAGT TCATGCCTAA 7883  
AATTAGCCCT TCTAAAGAAA TCCACCTGCT GCAAAATATC CAGTAGTTTG GCATTATATG 7943  
TGAAACTATC ACCATCATAG CTGGCACTGT GGGTTGTGGG ATCTCCTTTA GACATACAAC 8003  
ATAAATGATC TGGATGGATT AACATTACTA CATGGATGCT TGTGACACA TTAACCTGGC 8063  
TTCCCATGAG CTTTGTGTCA GATACACGCA GTGAACAGGT GTTTGGAGGA ACAGAATAAA 8123  
GAGAAGGCAA GCACTGGTAA GGGCAGGGGT TTGTGAAAGC TTGAGAGAAG AGACCAGTCT 8183  
GAGGACAGTA GACTTATT TTAGGATGGG GGTTGGATGA GGAGGCTATA GTTTGCTATA 8243  
AGCTTGAAT GGTGGAAC ACTGGTTTCA CTCACCTACC CAGCAGTTAT GTGTGGGGAA 8303  
GCCTTACCGA TGCTAAAGGA TCCATGTTAC AATAATGGCA TTATTTGGAA ATCCCAGTGG 8363  
TATTCCATGA ATAAAACCAC TATGAAGATA ATCCCCTCA ACAGACTCTC CGTTGGAGAA 8423  
GGACAGCAAC ACCACCCTGG GAAAGCCAAA CAGTCAGACC AGACCTGTTT AGCATCAGTA 8483  
GGACTTCCCT ACCATATCTG CTGGGTAGAT GAGTGAAACC AGTGTCCAA ACCACTCCGG 8543  
GCTTGTAGCA AACCATAGTC TCCTCATCTA CCAAGATGAG CAACCTTACC TCCTGATGTC 8603  
CTAGCCAATC ACCAACTAGG AAACCTTGCA CAGTTTATTT AAAGTAACAG TTTGATTTTC 8663  
ACAATATTTT TAAATTGGAG AACATAACT TATCTTTGCA CTCACAAACC ACATAATGAG 8723  
AAGAACTCT AAGGGAAAAT GCTTGATCTG TGTGACCCGG GCGCCATGC CAGAGCTGTA 8783  
GTTTATGCCA GTGTTGTGCT CTGACAAGCC TTTTACAGAA TTACATGAGA TCTGCTTCCC 8843  
TAGGACAAGG AGAAGGCAAA TCAACAGAGG CTGCACTTTA AAATGGAGAC ATAAAATAAC 8903  
ATGCCAGAAC CATTTCCTAA AGCTCCTCAA TCAACCAACA AAATTGTGCT TTCAAATAAC 8963  
CTGAGTTGAC CTCATCAGGA ATTTTGTGGC TCCTTCTCTT CTAACCTGCC TGAAGAAAGA 9023  
TGGTCCACAG CAGCTGAGTC CGGGATGGAT AAGCTTAGGG ACAGAGGCCA ATTAGGGAAC 9083

TTTGGGTTTC TAGCCCTACT AGTAGTGAAT AAATTTAAAG TGTGGATGTG ACTATGAGTC	9143
ACAGCACAGA TGTTGTTTAA TAATATGTTT ATTTTATAAA TTGATATTTT AGGAATCTTT	9203
GGAGATATTT TCAGTTAGCA GATAATACTA TAAATTTTAT GTAACCTGGCA ATGCACTTCG	9263
TAATAGACAG CTCTTCATAG ACTTGCAGAG GTAAAAAGAT TCCAGAATAA TGATATGTAC	9323
ATCTACGACT TGTTTTAG GT GGC ACT TAC TCA AAA GCA TCT ACT CCT AAT	9373
Gly Gly Thr Tyr Ser Lys Ala Ser Thr Pro Asn	
380 385	
GGT TAT GAT AAT GGC ATT ATT TGG GCC ACT TGG AAA ACC CGG TGG TAT	9421
Gly Tyr Asp Asn Gly Ile Ile Trp Ala Thr Trp Lys Thr Arg Trp Tyr	
390 395 400	
TCC ATG AAG AAA ACC ACT ATG AAG ATA ATC CCA TTC AAC AGA CTC ACA	9469
Ser Met Lys Lys Thr Thr Met Lys Ile Ile Pro Phe Asn Arg Leu Thr	
405 410 415	
ATT GGA GAA GGA CAG CAA CAC CAC CTG GGG GGA GCC AAA CAG GTC AGA	9517
Ile Gly Glu Gly Gln Gln His His Leu Gly Gly Ala Lys Gln Val Arg	
420 425 430 435	
CCA GAG CAC CCT GCG GAA ACA GAA TAT GAC TCA CTT TAC CCT GAG GAT	9565
Pro Glu His Pro Ala Glu Thr Glu Tyr Asp Ser Leu Tyr Pro Glu Asp	
440 445 450	
GAT TTG TAGAAAATTA ACTGCTAACT TCTATTGACC CACAAAGTTT CAGAAATTCT	9621
Asp Leu	
CTGAAAGTTT CTTCTTTTT TCTCTTACTA TATTTATTGA TTTCAAGTCT TCTATTAAGG	9681
ACATTTAGCC TTCAATGGAA ATTAAACTC ATTTAGGACT GTATTTCCAA ATTACTGATA	9741
TCAGAGTTAT TTAAAAATTG TTTATTTGAG GAGATAACAT TTCAACTTTG TTCCTAAATA	9801
TATAATAATA AAATGATTGA CTTTATTTGC ATTTTATGA CCACTTGTCA TTTATTTTGT	9861
CTTCGTAAAT TATTTTCATT ATATCAAATA TTTTAGTATG TACTTAATAA AATAGGAGAA	9921
CATTTTAGAG TTTCAAATTC CCAGGTATTT TCCTTGTTTA TTACCCCTAA ATCATTCTTA	9981
TTAATTCTT CTTTTTAAAT GGAGAAAATT ATGTCTTTTT AATATGGTTT TTGTTTTGTT	10041
ATATATTCAC AGGCTGGAGA CGTTTAAAAG ACCGTTTCAA AAGAGATTTA CTTTTTTAAA	10101

GGACTTTATC TGAACAGAGA GATATAATAT TTTTCCTATT GGACAATGGA CTTGCAAAGC 10161  
 TTCACTTCAT TTTAAGAGCA AAAGACCCCA TGTTGAAAAC TCCATAACAG TTTTATGCTG 10221  
 ATGATAATTT ATCTACATGC ATTTCAATAA ACCTTTTGTT TCCTAAGACT AGATACATGG 10281  
 TACCTTTATT GACCATTA AAACCACCAC TTTTGGCCAA TTTACCAATT ACAATTGGGC 10341  
 AACCATCAGT AGTAATTGAG TCCTCATTTT ATGCTAAATG TTATGCCTAA CTCTTTGGGA 10401  
 GTTACAAAGG AAATAGCAAT TATGGCTTTT GCCCTCTAGG AGATACAGGA CAAATACAGG 10461  
 AAAATACAGC AACCCAACT GACAATACTC TATACAAGAA CATAATCACT AAGCAGGAGT 10521  
 CACAGCCACA CAACCAAGAT GCATAGTATC CAAAGTGCAG CTG 10564

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 453 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ser Trp Ser Leu His Pro Arg Asn Leu Ile Leu Tyr Phe Tyr Ala  
 1 5 10 15  
 Leu Leu Phe Leu Ser Ser Thr Cys Val Ala Tyr Val Ala Thr Arg Asp  
 20 25 30  
 Asn Cys Cys Ile Leu Asp Glu Arg Phe Gly Ser Tyr Cys Pro Thr Thr  
 35 40 45  
 Cys Gly Ile Ala Asp Phe Leu Ser Thr Tyr Gln Thr Lys Val Asp Lys  
 50 55 60  
 Asp Leu Gln Ser Leu Glu Asp Ile Leu His Gln Val Glu Asn Lys Thr  
 65 70 75 80  
 Ser Glu Val Lys Gln Leu Ile Lys Ala Ile Gln Leu Thr Tyr Asn Pro  
 85 90 95

68

Asp Glu Ser Ser Lys Pro Asn Met Ile Asp Ala Ala Thr Leu Lys Ser  
 100 105 110

Arg Ile Met Leu Glu Glu Ile Met Lys Tyr Glu Ala Ser Ile Leu Thr  
 115 120 125

His Asp Ser Ser Ile Arg Tyr Leu Gln Glu Ile Tyr Asn Ser Asn Asn  
 130 135 140

Gln Lys Ile Val Asn Leu Lys Glu Lys Val Ala Gln Leu Glu Ala Gln  
 145 150 155 160

Cys Gln Glu Pro Cys Lys Asp Thr Val Gln Ile His Asp Ile Thr Gly  
 165 170 175

Lys Asp Cys Gln Asp Ile Ala Asn Lys Gly Ala Lys Gln Ser Gly Leu  
 180 185 190

Tyr Phe Ile Lys Pro Leu Lys Ala Asn Gln Gln Phe Leu Val Tyr Cys  
 195 200 205

Glu Ile Asp Gly Ser Gly Asn Gly Trp Thr Val Phe Gln Lys Arg Leu  
 210 215 220

Asp Gly Ser Val Asp Phe Lys Lys Asn Trp Ile Gln Tyr Lys Glu Gly  
 225 230 235 240

Phe Gly His Leu Ser Pro Thr Gly Thr Thr Glu Phe Trp Leu Gly Asn  
 245 250 255

Glu Lys Ile His Leu Ile Ser Thr Gln Ser Ala Ile Pro Tyr Ala Leu  
 260 265 270

Arg Val Glu Leu Glu Asp Trp Asn Gly Arg Thr Ser Thr Ala Asp Tyr  
 275 280 285

Ala Met Phe Lys Val Gly Pro Glu Ala Asp Lys Tyr Arg Leu Thr Tyr  
 290 295 300

Ala Tyr Phe Ala Gly Gly Asp Ala Gly Asp Ala Phe Asp Gly Phe Asp  
 305 310 315 320

Phe Gly Asp Asp Pro Ser Asp Lys Phe Phe Thr Ser His Asn Gly Met  
 325 330 335

Gln Phe Ser Thr Trp Asp Asn Asp Asn Asp Lys Phe Glu Gly Asn Cys  
 340 345 350

Ala Glu Gln Asp Gly Ser Gly Trp Trp Met Asn Lys Cys His Ala Gly  
 355 360 365

His Leu Asn Gly Val Tyr Tyr Gln Gly Gly Thr Tyr Ser Lys Ala Ser  
 370 375 380

Thr Pro Asn Gly Tyr Asp Asn Gly Ile Ile Trp Ala Thr Trp Lys Thr  
 385 390 395 400

Arg Trp Tyr Ser Met Lys Lys Thr Thr Met Lys Ile Ile Pro Phe Asn  
 405 410 415

Arg Leu Thr Ile Gly Glu Gly Gln Gln His His Leu Gly Gly Ala Lys  
 420 425 430

Gln Val Arg Pro Glu His Pro Ala Glu Thr Glu Tyr Asp Ser Leu Tyr  
 435 440 445

Pro Glu Asp Asp Leu  
 450

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10807 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (B) CLONE: ovine beta-lactoglobulin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ACGCGTGTG ACCTGCAGGT CAACGGATCT CTGTGTCTGT TTTCATGTTA GTACCACACT	60
GTTTTGGTGG CTGTAGCTTT CAGCTACAGT CTGAAGTCAT AAAGCCTGGT ACCTCCAGCT	120
CTGTTCTCTC TCAAGATTGT GTTCTGCTGT TTGGGTCTTT AGTGTCTCCA CACAATTTTT	180
AGAATTGTTT GTTCTAGTTC TGTGAAAAAT GATGCTGGTA TTTTGATAAG GATTGCATTG	240
AATCTGTA AAA GCTACAGATA TAGTCATTGG GTAGTACAGT CACTTTAACA ATATTA ACTC	300

TTCACATCTG TGAGCATGAT ATATTTTCCC CCTCTATATC ATCTTCAATT CCTCCTATCA	360
GTTTCTTTCA TTGCAGTTTT CTGAGTACAG GTCTTACACC TCCTTGGTTA GAGTCATTCC	420
TCAGTATTTT ATTCCTTTGA TACAATTGTG AATGAGGTAA TTTTCTTAGT TTCTCTTTCT	480
GATAGCTCAT TGTTAGTGTA TATATAGAAA AGCAACAGAT TTCTATGTAT TAATTTTGTA	540
TCCTGCAACA GATTTCTATG TATTAATTTT GTATCCTGCT ACTTTACGGA ATTCACTTAT	600
TAGCTTTTTG GTGACATCTT GAGGATTTTC TGAAGAAAAT GGCATGGTAT GGTAGGACAA	660
GGTGTATGT CATCTGCAAA CAGTGGCAGT TTTCTTCTT CCCTTCCAAC CTGGATTTCT	720
TTGATTTCTT TCTGTCTGAG TACGACTAGG ATTCCAATA CTATACCGAA TAAAAGTGGC	780
AAGAGTGGAC ATCCTTGCTT TATTTTCTG ACCTTAGAGG AAATGCTTTC AGTTTTTAC	840
CATTAATTAT AATGTTTACT GTGGGCTTGT CATATGTGGC CTTCAATTATA TGGAGGTCTA	900
TTCCCTCTAT ACCCACCTTG TTGAGAGTTT TTATCATAAA AGTATGTTGA ATTTTGCAA	960
AAGTTTTTCC TGCATCTATT GAGATGATTT TTA CTCTTCA ATTCATTAAT GATTTTTATT	1020
CTTCATTTTG TTAATGATTT CCATTCTTCA ATTTGTTAAC GTGGTATATC ACATTGATTG	1080
ATTTGTGGAT ACCTTTGTAT CCCTGGGATA AACCTCACTT GATCATGAGC TTTCAATGTA	1140
TTTTTGAATT CACTTTGCTA ATATTCTGTT GGGTATTTTT GCATCTCTAT TCATCAATGA	1200
TATTGGCCTA AGAAAGGTTT TGTCTGGTTT TAGTATCAGG GTGATGCTGG CCTCATAGAG	1260
AGAGTTTAGA AGCATTTCCT CCTCTTTGAT TTTTCGGAAT AGTTTGAGTA GGATAGGTAT	1320
TAACCTTCT TAAATGTTT GGGGACTTCC CTGGTGAGCC GGTGGTTGAG AATCCGCTC	1380
AGGGATGTGG GTTTGATCCC TGGTCAGGGA ACCATTAATA AGATCCCACA TGCTGCAGGC	1440
AACAAGCCCC CAAGCTGCAA CCACTGAGCT GCAACCGCTG CAGTGCCAC AGGCCACGAC	1500
CAGAGAAAGC CCACATACAG CAGGGAAGAC CCAGCACAAC CGGAAAAAGG AGTTTGGTGG	1560
AATACAGCTG TGAAGCCGTC TGGTCCTGGA CTCCTGCTTG AGGGAATTTT TAAAAATTA	1620
TTGATTCAAT TTCATTACTG GTAACCTGGTC TGTTCAATTTT TTCTATTTCT TCCGGGTTCA	1680
GTCTTGGGAG ATTGTACATG CCTAGGAATG TGTCGGTTTC TTCTAGGTTG TCCATTTTAT	1740



TGGACATGCA TGGGAGCACA CAGCACCGAC CAGCGAGACT CATGCTGGCT TCCTGGGGCC 1800  
AGGCTGGGGC CCCAAGCAGC ATGGCATCCT AGAGTGTGTG AAAGCCCACT GACCCTGCCC 1860  
AGCCCCACAA TTTCATTCTG AGAAGTGATT CTTGCTTCT GCACTTACAG GCCCAGGATC 1920  
TGACCTGCTT CTGAGGAGCA GGGGTTTTGG CAGGACGGGG AGATGCTGAG AGCCGACGGG 1980  
GGTCCAGGTC CCCTCCCAGG CCCCCCTGTC TGGGGCAGCC CTTGGGAAAG ATTGCCCCAG 2040  
TCTCCCTCCT ACAGTGGTCA GTCCCAGCTG CCCCAGGCCA GAGCTGCTTT ATTTCCGTCT 2100  
CTCTCTCTGG ATGGTATTCT CTGGAAGCTG AAGGTTCTG AAGTTATGAA TAGCTTTGCC 2160  
CTGAAGGGCA TGGTTTGTGG TCACGGTTCA CAGGAACTG GGAGACCCTG CAGCTCAGAC 2220  
GTCCCGAGAT TGGTGGCACC CAGATTTCT AAGCTCGCTG GGGAACAGGG CGCTTGTTC 2280  
TCCCTGGCTG ACCTCCCTCC TCCCTGCATC ACCCAGTTCT GAAAGCAGAG CGGTGCTGGG 2340  
GTCACAGCCT CTCGCATCTA ACGCCGGTGT CCAAACCACC CGTGCTGGTG TTCGGGGGGC 2400  
TACCTATGGG GAAGGGCTTC TCACTGCAGT GGTGCCCCC GTCCCCTCTG AGATCAGAAG 2460  
TCCCAGTCCG GACGTCAAAC AGGCCGAGCT CCCTCCAGAG GCTCCAGGGA GGGATCCTTG 2520  
CCCCCCCCT GCTGCCTCCA GCTCCTGGTG CCGCACCTT GAGCCTGATC TTGTAGACGC 2580  
CTCAGTCTAG TCTCTGCCTC CGTGTTTACA CGCCTTCTCC CCATGTCCCC TCCGTGTCCC 2640  
CGTTTTCTCT CACAAGGACA CCGGACATTA GATTAGCCCC TGTTCCAGCC TCACCTGAAC 2700  
AGCTCACATC TGTAAGACC TAGATTCAA ACAAGATTCC AACCTGAAGT TCCCGGTGGA 2760  
TGTGAGTTCT GGGGCGACAT CCTTCAACCC CATCACAGCT TGCAGTTCAT CGAAAACAT 2820  
GGAACCTGGG GTTTATCGTA AAACCCAGGT TCTTCATGAA AACTGAGCT TCGAGGCTTG 2880  
TTGCAAGAAT TAAAGGTGCT AATACAGATC AGGGCAAGGA CTGAAGCTGG CTAAGCCTCC 2940  
TCTTTCCATC ACAGGAAAGG GGGGCTGGG GCGGCTGGA GGTCTGCTCC CGTGAGTGAG 3000  
CTCTTTCTG CTACAGTCAC CAACAGTCTC TCTGGGAAGG AAACCAGAGG CCAGAGAGCA 3060  
AGCCGGAGCT AGTTTAGGAG ACCCCTGAAC CTCCACCCAA GATGCTGACC AGCCAGCGGG 3120

CCCCCTGGAA AGACCCTACA GTTCAGGGGG GAAGAGGGGC TGACCCGCCA GGTCCCTGCT	3180
ATCAGGAGAC ATCCCCGCTA TCAGGAGATT CCCCCACCTT GCTCCCGTTC CCCTATCCCA	3240
ATACGCCAC CCCACCCCTG TGATGAGCAG TTTAGTCACT TAGAATGTCA ACTGAAGGCT	3300
TTTGCATCCC CTTTGCCAGA GGCACAAGGC ACCCACAGCC TGCTGGGTAC CGACGCCCAT	3360
GTGGATTGAG CCAGGAGGCC TGTCCCTGCAC CCTCCCTGCT CGGGCCCCCT CTGTGCTCAG	3420
CAACACACCC AGCACCAGCA TTCCCCTGC TCCTGAGGTC TGCAGGCAGC TCGCTGTAGC	3480
CTGAGCGGTG TGGAGGGAAG TGTCCCTGGGA GATTTAAAT GTGAGAGGCG GGAGGTGGGA	3540
GGTTGGGCC TGTGGGCTG CCCATCCCAC GTGCCTGCAT TAGCCCCAGT GCTGCTCAGC	3600
CGTGCCCCCG CCGCAGGGT CAGGTCACTT TCCCGTCTG GGGTTATTAT GACTCTTGTC	3660
ATTGCCATTG CCATTTTTGC TACCCTAACT GGGCAGCAGG TGCTTGCAGA GCCCTCGATA	3720
CCGACCAGGT CCTCCCTCGG AGCTCGACCT GAACCCCATG TCACCCTTGC CCCAGCCTGC	3780
AGAGGGTGGG TGA CTGCAGA GATCCCTTCA CCCAAGGCCA CGGTCACATG GTTTGGAGGA	3840
GCTGGTGCCC AAGGCAGAGG CCACCCTCCA GGACACACCT GTCCCCAGTG CTGGCTCTGA	3900
CCTGTCTTG TCTAAGAGGC TGACCCCGGA AGTGTTCTG GCACTGGCAG CCAGCCTGGA	3960
CCCAGAGTCC AGACACCCAC CTGTGCCCC GCTTCTGGGG TCTACCAGGA ACCGTCTAGG	4020
CCCAGAGGGG ACTTCTGCT TGGCCTTGA TGGAAGAAGG CCTCCTATTG TCCTCGTAGA	4080
GGAAGCCACC CCGGGGCTG AGGATGAGCC AAGTGGGATT CCGGGAACCG CGTGGCTGGG	4140
GGCCAGCCC GGGCTGGCTG GCCTGCATGC CTCCTGTATA AGGCCCAAG CCTGCTGTCT	4200
CAGCCCTCCA CTCCTGCAG AGCTCAGAAG CACGACCCCA GGGATATCCC TGCAGCCATG	4260
AAGTGCCTCC TGCTTGCCT GGGCCTGGCC CTCGCCTGTG GCGTCCAGGC CATCATCGTC	4320
ACCCAGACCA TGAAAGGCCT GGACATCCAG AAGGTTTCAG GGTGCGCCG GTGGGTGAGT	4380
TGCAGGGCGG GCAGGGGAGC TGGCCTCAG AGAGCCAAGA GAGGCTGTGA CGTTGGGTTC	4440
CCATCAGTCA GCTAGGGCCA CCTGACAAAT CCCCCTGGG GCAGCTTCAA CCAGGCGTTC	4500
ACTGTCTTGC ATTCTGGAGG CTGGAAGCCC AAGATCCAGG TGTTGGCAGG GCTGGCTTCT	4560

CCTGCGGCCG CTCTCTGGGG AGCAGACGGC CGTCTTCTCC AGTCCTCTGC GCGCCCTGAT	4620
TTCCTCTTCC TGTGAGGCCA CCAGGCCTGC TGGAAACACG CCTGCCTGCG CAGCTTCACA	4680
CGACCTTTGT CATCTCTTTA AAGGCCATGT CTCCAGAGTC ATGTGTTGAA GTTCTGGGGG	4740
TTAGTGGGAC ACAGTTCAGC CCCTAAAAGA GTCTCTCTGC CCCTCAAATT TTCCCCACCT	4800
CCAGCCATGT CTCCCCAAGA TCCAAATGTT GCTACATGTG GGGGGGCTCA TCTGGGTCCC	4860
TCTTTGGGTT CAGTGTGAGT CTGGGGAGAG CATTCCCCAG GGTGCAGAGT TGGGGGGAGT	4920
ATCTCAGGGC TGCCCAGGCC GGGGTGGGAC AGAGAGCCCA CTGTGGGGCT GGGGGCCCCT	4980
TCCCACCCCC AGAGTGCAAC TCAAGGTCCC TCTCCAGGTG GCGGGGACTT GGCACCTCTT	5040
GGCTATGGCG GCCAGCGACA TCTCCCTGCT GGATGCCCAG AGTGCCCCC TGAGAGTGTA	5100
CGTGGAGGAG CTGAAGCCCA CCCCCGAGGG CAACCTGGAG ATCCTGCTGC AGAAATGGTG	5160
GGCGTCTCTC CCCAACATGG AACCCCACT CCCAGGGCT GTGGACCCCC CGGGGGGTGG	5220
GGTGCAGGAG GGACCAGGGC CCCAGGGCTG GGGGAAGAGGG CTCAGAGTTT ACTGGTACCC	5280
GGCGCTCCAC CCAAGGCTGC CCACCCAGGG CTTTTTTTTT TTTTAACTT TTATTAATT	5340
GATGCTTCAG AACATCATCA AACAAATGAA CATAAACAT TCATTTTTGT TTACTTGAA	5400
GGGGAGATAA AATCCTCTGA AGTGGAATG CATAGCAAAG ATACATACAA TGAGGCAGGT	5460
ATTCTGAATT CCCTGTTAGT CTGAGGATTA CAAGTGTATT TGAGCAACAG AGAGACATTT	5520
TCATCATTTT TAGTCTGAAC ACCTCAGTAT CTAATGAA CAAGAAGTCC TGGAAACGAA	5580
GCAGTGTGGG GATAGGCCCG TGTGAAGGCT GCTGGGAGGC AGCAGACCTG GGTCTTCGGG	5640
CTCAAGCAGT TCCCCTACC AGCCCTGTCC ACCTCAGACG GGGGTCAGGG TGCAGGAGAG	5700
AGCTGGATGG GTGTGGGGGC AGAGATGGGG ACCTGAACCC CAGGGCTGCC TTTTGGGGGT	5760
GCCTGTGGTC AAGGCTCTCC CTGACCTTTT CTCTCTGGCT TCATCTGACT TCTCCTGGCC	5820
CATCCACCCG GTCCCCTGTG GCCTGAGGTG ACAGTGAGTG CGCCGAGGCT AGTTGGCCAG	5880
CTGGCTCCTA TGCCCATGCC ACCCCCCTCC AGCCCTCCTG GGCCAGCTTC TGCCCCTGGC	5940

CCTCAGTTCA TCCTGATGAA AATGGTCCAT GCCAATGGCT CAGAAAGCAG CTGTCTTTCA	6000
GGGAGAACGG CGAGTGTGCT CAGAAGAAGA TTATTGCAGA AAAAAACCAAG ATCCCTGCGG	6060
TGTTCAAGAT CGATGGTGAG TCCGGGTCCC TGGGGGACAC CCACCACCCC CGCCCCGGG	6120
GA CTGTGGAC AGGTT CAGGG GGCTGGCGTC GGGCCCTGGG ATGCTAAGGG ACTGGTGGTG	6180
ATGAAGACAC TGCCTTGACA CCTGCTTAC TTGCCTCCCC TGCCACCTGC CCGGGGCCTT	6240
GGGGCGGTGG CCATGGGCAG GTCCCGGCTG GCGGGCTAAC CCACCAGGGT GACACCCGAG	6300
CTCTCTTTGC TGGGGGGCGG GCGGTGCTCT GGGCCCTCAG GCTGAGCTCA GGAGGTACCT	6360
GTGCCCTCCC AGGGGTAACC GAGAGCCGT GCCCACTCCA GGGGCCAGG TGCCCCACGA	6420
CCCCAGCCG CTCCACAGCT CCTTCATCTC CTGGAGACAA ACTCTGTCCG CCCTCGCTCA	6480
TTC ACTTGTT CGTCCTAAAT CCGAGATGAT AAAGCTTCGA GGGGGGGTTG GGGTTCCATC	6540
AGGGCTGCCC TTCCGCCGGG CAGCCTGGGC CACATCTGCC CTTGGCCCC TCAGGACTCA	6600
CTCTGACTGG AGGCCCTGCA CTGACTGACG CCAGGGTGCC CAGCCCAGGG TCTCTGGCGC	6660
CATCCAGCTG CACTGGGTTT GGGTGTGGT CCTGCCCCCA AGCTGCCCGG ACACCACAGG	6720
CAGCCGGGGC TGCCCACTGG CCTCGGTCAG GGTGAGCCCC AGCTGCCCCC GCTCAGGGCT	6780
TGCCCCGACA ATGACCCCAT CCTCAGGACG CACCCCTT CCCTTGCTGG GCAGTGCCA	6840
GCCCCACCCG AGATCGGGG AAGCCCTATT TCTTGACAAC TCCAGTCCCT GGGGAGGGG	6900
GCCTCAGACT GAGTGGTGAG TGTTCCAAG TCCAGGAGGT GGTGGAGGGT CCTGGCGGAT	6960
CCAGAGTTGA CAGTGAGGGC TTCCTGGGCC CCATGCGCCT GGCAGTGGCA GCAGGAAGA	7020
GGAAGCACCA TTTCAGGGT GGGGGATGCC AGAGGCGCTC CCCACCCGT CTCGCCGGG	7080
TGGTGACCCC GGGGAGCCC CGCTGGTCGT GGAGGGTGCT GGGGGCTGAC TAGCAACCCC	7140
TCCCCCCCCG TTGGA ACTCA CTTTTCTCCC GTCTTGACCG CGTCCAGCCT TGAATGAGAA	7200
CAAAGTCCTT GTGCTGGACA CCGACTACAA AAAGTACCTG CTCTTCTGCA TGGAAAACAG	7260
TGCTGAGCCC GAGCAAAGCC TGGCCTGCCA GTGCCTGGGT GGGTGCCAAC CCTGGCTGCC	7320
CAGGGAGACC AGCTGCGTGG TCCTTGCTGC AACAGGGGT GGGGGGTGGG AGCTTGATCC	7380

CCAGGAGGAG GAGGGGTGGG GGGTCCCTGA GTCCCGCCAG GAGAGAGTGG TGCATACCG 7440  
GGAGCCAGTC TGCTGTGGGC CTGTGGGTGG CTGGGGACGG GGGCCAGACA CACAGGCCGG 7500  
GAGACGGGTG GGCTGCAGAA CTGTGACTGG TGTGACCGTC GCGATGGGGC CGGTGGTCAC 7560  
TGAATCTAAC AGCCTTTGTT ACCGGGGAGT TTCAATTATT TCCCAAATA AGAACTCAGG 7620  
TACAAAGCCA TCTTTCAACT ATCACATCCT GAAAACAAAT GGCAGGTGAC ATTTTCTGTG 7680  
CCGTAGCAGT CCCACTGGGC ATTTTCAGGG CCCCTGTGCC AGGGGGGCGC GGGCATCGGC 7740  
GAGTGGAGGC TCCTGGCTGT GTCAGCCGGC CCAGGGGGAG GAAGGGACCC GGACAGCCAG 7800  
AGGTGGGGGG CAGGCTTTCC CCCTGTGACC TGCAGACCCA CTGCACTGCC CTGGGAGGAA 7860  
GGGAGGGGAA CTAGGCCAAG GGGGAAGGGC AGGTGCTCTG GAGGGCAAGG GCAGACCTGC 7920  
AGACCACCCT GGGGAGCAGG GACTGACCCC CGTCCCTGCC CCATAGTCAG GACCCCGGAG 7980  
GTGGACAACG AGGCCCTGGA GAAATTCGAC AAAGCCCTCA AGGCCCTGCC CATGCACATC 8040  
CGGCTTGCTT TCAACCCGAC CCAGCTGGAG GGTGAGCACC CAGGCCCGC CCTTCCCCAG 8100  
GGCAGGAGCC ACCCGGCCCC GGGACGACCT CCTCCATGG TGACCCCCAG CTCCCCAGG 8160  
CTCCCAGGAG GAAGGGGTGG GGTGCAGCAC CCCGTGGGGG CCCCTCCCC ACCCCCTGCC 8220  
AGGCCTCTCT TCCCAGAGTG TCCAGTCCA TCCTGACCCC CCCATGACTC TCCCTCCCC 8280  
ACAGGGCAGT GCCACGTCTA GGTGAGCCCC TGCCGGTGCC TCTGGGGTAA GCTGCCTGCC 8340  
CTGCCCCACG TCCTGGGCAC ACACATGGGG TAGGGGTCT TGGTGGGGCC TGGGACCCCA 8400  
CATCAGGCC TGGGGTCCCC CCTGTGAGAA TGGCTGGAAG CTGGGGTCCC TCCTGGCGAC 8460  
TGCAGAGCTG GCTGGCCGCG TGCCACTCTT GTGGGTGACC TGTGTCTGG CCTCACACAC 8520  
TGACCTCCTC CAGCTCCTC CAGCAGAGCT AAGGCTAAGT GAGCCAGAAT GGTACCTAAG 8580  
GGGAGGCTAG CGGTCTTCT CCCGAGGAGG GGCTGTCTG GAACCACCAG CCATGGAGAG 8640  
GCTGGCAAGG GTCTGGCAGG TGCCCAGGA ATCACAGGGG GGCCCCATGT CCATTTCAGG 8700  
GCCCCGGAGC CTTGGACTCC TCTGGGGACA GACGACGTCA CCACCGCCCC CCCCCATCA 8760

GGGGGACTAG AAGGGACCAG GACTGCAGTC ACCCTTCCTG GGACCCAGGC CCCTCCAGGC 8820  
CCCTCCTGGG GCTCCTGCTC TGGGCAGCTT CTCCTTCACC AATAAAGGCA TAAACCTGTG 8880  
CTCTCCCTTC TGAGTCTTTG CTGGACGACG GGCAGGGGGT GGAGAAGTGG TGGGGAGGGA 8940  
GTCTGGCTCA GAGGATGACA GCGGGGCTGG GATCCAGGGC GTCTGCATCA CAGTCTTGTG 9000  
ACAACTGGGG GCCCACACAC ATCACTGCGG CTCTTTGAAA CTTTCAGGAA CCAGGGAGGG 9060  
ACTCGGCAGA GACATCTGCC AGTTCACTTG GAGTGTTCAG TCAACACCCA AACTCGACAA 9120  
AGGACAGAAA GTGGAAAATG GCTGTCTCTT AGTCTAATAA ATATTGATAT GAAACTCAAG 9180  
TTGCTCATGG ATCAATATGC CTTTATGATC CAGCCAGCCA CTA CTGTCGT ATCAACTCAT 9240  
GTACCCAAAC GCACTGATCT GTCTGGCTAA TGATGAGAGA TTCCAGTAG AGAGCTGGCA 9300  
AGAGGTCACA GTGAGAACTG TCTGCACACA CAGCAGAGTC CACCAGTCAT CCTAAGGAGA 9360  
TCAGTCCTGG TGTTCAATTG AGGACTGATG TTGAAGCTGA AACTCCAATG CTTTGGCCAC 9420  
CTGATGTGAA GAGCTGACTC ATTTGAAAAG ACCCTGATGC TGGGAAAGAT TGAGGGCAGG 9480  
AGGAGAAGGG GACGACAGAG GATGAGATGG TTGGATGGCA TCACCAACAC AATGGACATG 9540  
GGTTTGGGTG GACTCCAGGA GTTGGTGATG GACAGGGAGG CCTGGCGTGC TACGGAAGCG 9600  
GTTTATGGGG TCACAAAGAC TGAGTGA CTG AACTGAGCTG AACTGAATGG AAATGAGGTA 9660  
TACAGCAAAG TGGGGATTTT TTAGATAATA AGAATATACA CATAACATAG TGTATACTCA 9720  
TATTTTTATG CATACTGAA TGCTCAGTCA CTCAGTCGTA TCTGACTCTG TGACCTATGG 9780  
ACCGTAGCCT TCCAGGTTTC TTCTGTCCAC AGAATTCTCC AAGGCAAGAA TACTGGAGTG 9840  
GGTAGCCATT TCCTCCTCCA GGGGATCCTC CCGACCCAGG GATTGAACCG GCATCTCCTG 9900  
TATTGGCAGG TGGATTCTTT ACCACTGTGC CACCAGGGAA GCCCGTGTTA CTCTCTATGT 9960  
CCCCTTAAT TACCAAAGCT GCTCCAAGAA AAAGCCCCTG TGCCCTCTGA GCTTCCCGGC 10020  
CTGCAGAGGG TGGTGGGGGT AGACTGTGAC CTGGGAACAC CCTCCCGCTT CAGGACTCCC 10080  
GGGCCACGTG ACCCACAGTC CTGCAGACAG CCGGGTAGCT CTGCTCTTCA AGGCTCATT 10140  
TCTTTAAAAA AACTGAGGT CTATTTTGTG ACTTCGCTGC CGTAACTTCT GAACATCCAG 10200

TCGGATGGAC AGGACCTCCT CCCAGGCCT CAGGGGCTTC AGGGAGCCAG CCTTCACCTA 10260  
 TGAGTCACCA GACTCTCGGG GGTGGCCCCG CCTTCAGGGT GCTCACAGTC TTCCCATCGT 10320  
 CCTGATCAAA GAGCAAGACC AATGACTTCT TAGGAGCAAG CAGACACCCA CAGGACTG 10380  
 AGGTTACCA GAGCTGAGCT GTCCTTTTGA ACCTAAAGAC ACACAGCTCT CGAAGGTTTT 10440  
 CTCTTTAATC TGGATTTAAG GCCTACTTGC CCCTCAAGAG GGAAGACAGT CCTGCATGTC 10500  
 CCCAGGACAG CCACTCGGTG GCATCCGAGG CCACTTAGTA TTATCTGACC GCACCCTGGA 10560  
 ATTAATCGGT CCAAACCTGGA CAAAACCTT GGTGGGAAGT TTCATCCAG AGGCCTCAAC 10620  
 CATCCTGCTT TGACCACCCT GCATCTTTTT TTCTTTTATG TGTATGCATG TATATATATA 10680  
 TATATATTTT TTTTTTTTTC ATTTTTTGGC TGTGCTGGCT GTTCGTTGCA GTTCGGTGCG 10740  
 CAGGCTTCTC TCTAGTTTCT CTCTAGTCTT CTCTTATCAC AGAGCAGTCT CTAGACGATC 10800  
 GACGCGT 10807

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AATTCCGATC GACGCGTCGA CGATATACTC TAGACGATCG ACGCGTA

47

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:  
(B) CLONE: BLGAMP3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGGATCCCCT GCCGGTGCCT CTGG

24

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:  
(B) CLONE: BLGAMP4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

AACGCGTCAT CCTCTGTGAG CCAG

24

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:  
(B) CLONE: ZC6839

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ACTACGTAGT

10

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid



79

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6632

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CGACGCGGAT CCTACGTACC TGCAGCCATG TTTCCATGA GG

42

- (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6627

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AGGGCTTCGG CAAGCTTCAG G

21

- (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6521

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GCCAAAGACT TACTCCCTC TAGA

24

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6520

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GCATGAACGT CGCGTGGTGG TTGTGCTACC

30

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6519

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ACCACGCGAC GTTCATGCTC TAAAACCGTT

30

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6518

81

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCTGCGGGAT CCTACGTACT AGGGGGACAG GGAAGG

36

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC6629

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CGACGCGAAT TCTACGTACC TGCAGCCATG AAAAGGATGG TTTCT

45

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

(B) CLONE: ZC6630

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGACGCGAAT TCTACGTACC TGCAGCCATG AAACATCTAT TATTG

45

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

82

- (vii) IMMEDIATE SOURCE:  
(B) CLONE: ZC6625

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GTGAGATTTT CAGATCTTGT C

21

- (2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
(B) CLONE: ZC6626

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

AAGAATTACT GTGGCCTACC A

21

- (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
(B) CLONE: ZC6624

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCTGCGGAAT TCTACGTACT ATTGCTGTGG GAA

33

- (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 base pairs  
(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6514

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CGACGCGGAT CCTACGTACC TGCAGCCATG AGTTGGTCCT TGCAC

45

- (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6517

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GTCTCTGGTA GCAACATACT A

21

- (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: ZC6516

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GGGTTTCTAG CCCTACTAGT AG

22

- (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:  
(B) CLONE: ZC6515

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GGGTTTCTAG CCCTACTAGT AG

22

- (2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 47 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

AAGCTACGCG TCGATCGTCT AGAGTATATC GTCGACGCGT CGATCGG

Claims

1. A method for producing fibrinogen comprising:  
providing a first DNA segment encoding a secretion signal operably linked to a fibrinogen A $\alpha$  chain, a second DNA segment encoding a secretion signal operably linked to a fibrinogen B $\beta$  chain, and a third DNA segment encoding a secretion signal operably linked to a fibrinogen  $\gamma$  chain, wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in the mammary gland of a host female mammal;  
introducing said DNA segments into a fertilized egg of a non-human mammalian species;  
inserting said egg into an oviduct or uterus of a female of said species to obtain offspring carrying said DNA constructs;  
breeding said offspring to produce female progeny that express said first, second and third DNA segments and produce milk containing biocompetent fibrinogen encoded by said segments;  
collecting milk from said female progeny;  
and recovering the fibrinogen from the milk.
2. A method according to claim 1 wherein said species is selected from the group consisting of sheep, pigs, goats and cattle.
3. A method according to claim 1 wherein each of said first, second and third DNA segments comprises an intron.
4. A method according to claim 1 wherein the molar ratio of said first, second and third DNA segments is within the range of 0.5-1:0.5-1:0.5-1.
5. A method according to claim 1 wherein each of said first, second and third DNA segments is operably linked to a transcription promoter selected from the group consisting

of casein,  $\beta$ -lactoglobulin,  $\alpha$ -lactalbumin and whey acidic protein gene promoters.

6. A method according to claim 1 wherein said first, second and third DNA segments are expressed under the control of a  $\beta$ -lactoglobulin promoter.

7. A method according to claim 1 wherein said introducing step comprises injecting said first, second and third DNA segments into a pronucleus of said fertilized egg.

8. A method according to claim 1 wherein said fibrinogen is human fibrinogen.

9. A method according to claim 1 wherein said second DNA segment comprises a sequence of nucleotides as shown in SEQ ID NO: 3 from nucleotide 470 to nucleotide 8100.

10. A method according to claim 1 wherein said second DNA segment comprises a sequence of nucleotides as shown in SEQ ID NO: 3 from nucleotide 512 to nucleotide 8100.

11. A method of producing fibrinogen comprising:  
incorporating a first DNA segment encoding a secretion signal operably linked to an A $\alpha$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a first gene fusion;  
incorporating a second DNA segment encoding a secretion signal operably linked to a B $\beta$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a second gene fusion;  
incorporating a third DNA segment encoding a secretion signal operably linked to a  $\gamma$  chain of fibrinogen into a  $\beta$ -lactoglobulin gene to produce a third gene fusion;  
introducing said first, second and third gene fusions into the germ line of a non-human mammal so that said DNA segments are expressed in a mammary gland of said mammal or its female progeny and biocompetent fibrinogen is secreted into milk of said mammal or its female progeny ;



obtaining milk from said mammal or its female progeny; and

recovering said fibrinogen from said milk.

12. A method according to claim 11 wherein said mammal is a sheep, pig, goat or bovine.

13. A method according to claim 11 wherein each of said first, second and third gene fusions comprises an intron.

14. A method according to claim 11 wherein the molar ratio of said first, second and third gene fusions introduced is within the range of 0.5-1:0.5-1:0.5-1.

15. A method according to claim 11 wherein said introducing step comprises injecting said first, second and third gene fusions into a pronucleus of a fertilized egg and inserting said egg into an oviduct of a pseudopregnant female to produce female offspring carrying said gene fusions in the germ line.

16. A method for producing fibrinogen comprising:  
providing a transgenic female non-human mammal carrying in its germline heterologous DNA segments encoding  $A\alpha$ ,  $B\beta$  and  $\gamma$  chains of fibrinogen, wherein said segments are expressed in a mammary gland of said mammal and fibrinogen encoded by said segments is secreted into milk of said mammal;  
collecting milk from said mammal; and  
recovering said fibrinogen from said milk.

17. A method according to claim 16 wherein said mammal is a sheep, pig, goat or bovine.

18. A non-human mammalian embryo containing in its nucleus heterologous DNA segments encoding  $A\alpha$ ,  $B\beta$  and  $\gamma$  chains of fibrinogen.

19. A transgenic non-human female mammal that produces recoverable amounts of human fibrinogen in its milk.

20. A process for producing a transgenic offspring of a mammal comprising:

providing a first DNA segment encoding a fibrinogen A $\alpha$  chain, a second DNA segment encoding a fibrinogen B $\beta$  chain, and a third DNA segment encoding a fibrinogen  $\gamma$  chain, wherein each of said first, second and third segments is operably linked to additional DNA segments required for its expression in a mammary gland of a host female mammal and secretion into milk of said host female mammal;

introducing said DNA segments into a fertilized egg of a mammal of a non-human species;

inserting said egg into an oviduct or uterus of a female of said non-human species to obtain an offspring carrying said first, second and third DNA segments.

21. A process according to claim 20 wherein said offspring is female.

22. A process according to claim 20 wherein said offspring is male.

23. A non-human mammal produced according to the process of claim 20.

24. A non-human mammal according to claim 23 wherein said mammal is female.

25. A female mammal according to claim 24 that produces milk containing biocompetent fibrinogen encoded by said DNA segments.

26. A non-human mammal according to claim 23 wherein said mammal is male.

27. A non-human mammal carrying in its germline DNA segments encoding heterologous  $A\alpha$ ,  $B\beta$  and  $\gamma$  chains of fibrinogen, wherein female progeny of said mammal express said DNA segments in a mammary gland to produce biocompetent fibrinogen.

28. A mammal according to claim 27 wherein said mammal is female.

29. A mammal according to claim 27 wherein said mammal is male.

FIGURE 1




FIGURE 2




FIGURE 3






FIGURE 4



**FIGURE 5**





# INTERNATIONAL SEARCH REPORT

In. tional Application No

PCT/US 95/02648

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC 6 C12N15/89 C12N15/90 C12N15/63 C12N15/62 C12N15/85  
 A01K67/027 C07K14/75 //C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	FIBRINOLYSIS, vol. 8, no. suppl.1, 18 September 1994 - 22 September 1994 page 102 PRUNCKARD ET AL. 'Expression of recombinant human fibrinogen in the milk of transgenic mice' see abstract nr 285	19, 27, 28
Y	---	1-18, 20-26, 29
	-/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

27 June 1995

Date of mailing of the international search report

- 3. 07. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+ 31-70) 340-3016

Authorized officer

Gac, G

INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 95/02648

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CHEMICAL ABSTRACTS, vol. 115, no. 19, 11 November 1991 Columbus, Ohio, US; abstract no. 202768k, REDMAN C ET AL. 'Recombinant production, secretion, and clotting behavior of fibrinogen, and cell line used therein' page 526; column 202762; see abstract &amp; US,A,663 380 (UNITED STATES DEPT OF HEALTH AND HUMAN SERVICES) 1 September 1991</p>	1-29
Y	<p>J. CONTROL. RELEASE, vol. 29, 1994 pages 213-221, SANG HE LEE ET AL. 'Production of biomedical proteins in the milk of transgenic dairy cows : the state of the art' proc. fourth int. symp. dispos. delivery peptide drugs, Leiden, Netherlands, 23-25 April 1993 see page 215 see page 218</p>	1-29
Y	<p>WO,A,92 11358 (THE AGRICULTURAL AND FOOD RESEARCH COUNCIL) 9 July 1992 see the whole document</p>	1-29
A	<p>WO,A,88 00239 (PHARMACEUTICAL PROTEINS LTD) 14 January 1988 cited in the application see the whole document</p>	1-29
A	<p>WO,A,90 05188 (PHARMACEUTICAL PROTEINS LIMITED) 17 May 1990 see pages 6-10,12,35-46,51-53 and claims</p>	1-10, 15-17
A	<p>WO,A,91 08216 (GENPHARM INTERNATIONAL) 13 June 1991 see the whole document</p>	1-29
A	<p>BIOCHEMISTRY, vol. 22, 1983 pages 3244-33250, CHUNG ET AL. 'Characterization of complementary deoxyribonucleic acid and genomic deoxyribonucleic acid for the beta chain of human fibrinogen' see the whole document</p>	9,10

1

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International Application No  
PCT/US 95/02648

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9211358	09-07-92	AU-A- 9139191	22-07-92
		EP-A- 0566596	27-10-93
		JP-T- 6508261	22-09-94
-----			
WO-A-8800239	14-01-88	AU-B- 605497	17-01-91
		AU-A- 7649087	29-01-88
		EP-A- 0274489	20-07-88
		JP-T- 1500162	26-01-89
		US-A- 5366894	22-11-94
		US-A- 5322775	21-06-94
-----			
WO-A-9005188	17-05-90	AU-B- 628101	10-09-92
		AU-A- 4494389	28-05-90
		EP-A- 0396699	14-11-90
		JP-T- 3505674	12-12-91
-----			
WO-A-9108216	13-06-91	AU-B- 656720	16-02-95
		AU-A- 6960891	26-06-91
		CA-A- 2075206	02-06-91
		CN-A- 1053446	31-07-91
		EP-A- 0502976	16-09-92
		OA-A- 9669	15-05-93
-----			