



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:

06.01.2021 Bulletin 2021/01

(21) Application number: **11840762.6**

(22) Date of filing: **15.11.2011**

(51) Int Cl.:

A01N 43/16 (2006.01)	A61K 31/35 (2006.01)
A61K 9/10 (2006.01)	A61P 1/00 (2006.01)
A61P 11/00 (2006.01)	A61P 13/08 (2006.01)
A61P 15/00 (2006.01)	A61P 17/00 (2006.01)
A61P 19/08 (2006.01)	A61P 35/00 (2006.01)
A61P 39/00 (2006.01)	A61P 43/00 (2006.01)

(86) International application number:

PCT/US2011/060829

(87) International publication number:

WO 2012/068140 (24.05.2012 Gazette 2012/21)

(54) **NANOPARTICLE ISOFLAVONE COMPOSITIONS & METHODS OF MAKING AND USING THE SAME**

NANOPARTIKEL-ISOFLAVONZUSAMMENSETZUNGEN SOWIE VERFAHREN ZU IHRER
HERSTELLUNG UND VERWENDUNG

COMPOSITIONS DE NANOPARTICULES D'ISOFLAVONE & LEURS PROCÉDÉS DE FABRICATION
ET D'UTILISATION

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **15.11.2010 US 946711**

(43) Date of publication of application:

25.09.2013 Bulletin 2013/39

(73) Proprietor: **HUMANETICS CORPORATION**

Eden Prairie, MN 55344 (US)

(72) Inventors:

- **ELDER, Edmund Joseph**
Madison
Wisconsin 53719 (US)
- **SACCHETTI, Mark Joseph**
Madison
Wisconsin 53717 (US)
- **TLACHAC, Randall Joseph**
Minneapolis
Minnesota 55442 (US)

• **ZENK, John L.**

Eden Prairie
Minnesota 55347 (US)

(74) Representative: **Stewart, Hazel et al**

Cleveland
10 Fetter Lane
London EC4A 1BR (GB)

(56) References cited:

WO-A1-2007/000192 WO-A1-2007/000193
US-A1- 2007 020 197 US-A1- 2007 116 510
US-A1- 2009 035 336

- **SINGH V K ET AL: "Effects of genistein
administration on cytokine induction in
whole-body gamma irradiated mice",
INTERNATIONAL IMMUNOPHARMACOLOGY,
ELSEVIER, AMSTERDAM, NL, vol. 9, no. 12, 1
November 2009 (2009-11-01), pages 1401-1410,
XP026667281, ISSN: 1567-5769, DOI:
10.1016/J.INTIMP.2009.08.012 [retrieved on
2009-08-28]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionTechnical Field

5 **[0001]** The present invention relates to compositions including genistein and methods for producing and utilizing such compositions.

Background

10 **[0002]** Genistein is a pharmaceutically active isoflavone. In the body, genistein interacts with various enzymes that have wide-ranging actions in many tissues. Therefore, the potential therapeutic impacts of genistein are diverse. However, genistein has proven difficult to formulate and deliver to subjects in a manner that achieves and maintains therapeutically effective blood plasma levels.

15 Description of the Figures**[0003]**

20 Figure 1 shows the thirty-day survival rates in groups of mice receiving a solution formulation of genistein versus the thirty-day survival rates in groups of mice receiving a genistein suspension formulation as described herein.
 Figure 2 shows the thirty-day survival rates of mice after subcutaneous administration of a genistein suspension formulation as described herein at 24, 18, 12, or 6 hr pre-irradiation.
 Figure 3 shows the thirty-day survival rates of mice after subcutaneous and intramuscular administration of a genistein suspension formulation as described herein.
 25 Figure 4 shows the free genistein concentration achieved after oral administration of a genistein suspension formulation as described herein versus that achieved by a solution formulation of genistein.
 Figure 5 shows the total genistein concentration achieved after oral administration of a genistein suspension formulation as described herein versus that achieved by a solution formulation of genistein.
 Figure 6 shows the free genistein concentration achieved after oral administration of a genistein suspension formulation as described herein versus that achieved by a non-nanoparticulate suspension formulation of genistein.
 30 Figure 7 shows the total genistein concentration achieved after oral administration of a genistein suspension formulation as described herein versus that achieved by a non-nanoparticulate suspension formulation of genistein.

Detailed Description of Preferred Embodiments

35 **[0004]** The invention is described by reference to the claims. Compositions of genistein compounds are described herein. In particular formulations, the compositions described herein are pharmaceutical formulations suitable for oral or parenteral administration. Generally, given the desired therapeutic applications for genistein, it is desirable to deliver genistein to a subject in a manner that (i) achieves a therapeutic blood plasma concentration in a relatively short period
 40 of time and (ii) maintains a therapeutic blood plasma concentration over an extended period of time. Using available formulations of genistein, however, it has been found that relatively high doses of genistein are often required to achieve and maintain therapeutic blood plasma concentrations of genistein. This is particularly true when genistein is administered orally. However, even when genistein is formulated for parenteral administration (e.g., via intravenous injection or infusion, intravascular injection, subcutaneous injection, or intramuscular injection), it is often the case that relatively large volumes
 45 of drug formulation must be delivered in order to achieve and maintain therapeutic blood plasma concentrations.
[0005] Genistein is practically insoluble in water, requiring greater than 50,000 parts water at 25° C to dissolve one part genistein. Furthermore, when delivered orally, genistein has shown poor bioavailability, which may be due, at least in part, to the compound's low water solubility. Therefore, in light of genistein's generally low bioavailability and water solubility, achieving and maintaining therapeutic blood plasma concentrations of genistein can require high doses of
 50 genistein delivered at relatively high dose frequencies. Genistein compositions with high concentrations of genistein that provide significantly increased bioavailability are described herein. Moreover, in particular formulations, the genistein compositions described herein maintain therapeutic blood plasma levels of genistein over an extended period of time.
[0006] The genistein formulations described herein are suitable for oral and parenteral administration. Additionally, the formulations described herein potentially provide several advantages. For example, because they can be used to
 55 achieve therapeutic plasma concentrations of genistein using less amounts of drug substance and, in some cases, relatively fewer doses, the formulations described herein may reduce the costs of genistein treatments as well as any potential side effects that may be associated with relatively higher doses of the compound. Moreover, because the formulations described herein enable delivery of therapeutic amounts of genistein using relatively smaller administered

amounts of formulated drug, they may ease patient compliance and expand the contexts in which administration of genistein may be utilized.

[0007] The formulations described herein also exhibit desirable stability characteristics, are scalable for commercial production and, in specific compositions, may increase the circulating half-life of genistein after administration.

[0008] Methods of treating subjects at risk for or suffering from various diseases and disorders suitable for treatment using genistein are also described herein.

I. Definitions

[0009] It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a buffer" includes a plurality of such buffers, reference to "the buffer" is a reference to one or more buffers and equivalents thereof known to those skilled in the art, and so forth.

[0010] Ranges can be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another embodiment. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also herein disclosed as "about" that particular value in addition to the value itself. For example, if the value "10" is disclosed, then "about 10" is also disclosed. It is also understood that each unit between two particular units are also disclosed. For example, if 10 and 15 are disclosed, then 11, 12, 13, and 14 are also disclosed.

[0011] As used herein, "nanoparticulate" refers to material exhibiting a volume diameter, as measured using laser light diffraction, wherein the D (0.50) of the material is 0.5 μm or less and no particles measure greater than 2 μm . Particle size analysis using laser light diffraction is a technique based on light being scattered through various angles which are directly related to the size of the particles. By measuring the angles of light scattered by the particles being analyzed and the intensity of this scattered light, a particle size distribution can be calculated. Techniques for use in analyzing particle size in the context of the present disclosure can be referred to as static light scattering, Rayleigh light scattering, low angle light scattering (LALS), multiple angle light scattering (MALS) Fraunhofer diffraction, or Mie Scattering. Measurement of particle size distributions using Mie Scattering allows for the determination of particle size distributions through the direction measurement of mass.

[0012] Two theoretical applications to the analysis of particle size by laser light diffraction are based on assumptions about the properties of the particles. Fraunhofer theory considers the following: particles are spherical, non-porous and opaque; particle diameters are greater than the wavelength of the laser light used in the analysis; and particles are distant enough from each other not to interfere in the diffraction of light, exhibit random motion, and diffract light with the same efficiency regardless of size and shape. Mie theory considers the differences in refractive index between the particles and the suspending medium, which allows the measurement technique to account for particles in the size range below the wavelength of the laser light used in the analysis. The relative amounts of different size particles are determined by measuring the intensity of light scattered at different angles. As the particles get close to or smaller than the wavelength of light, more of the light intensity is scattered to higher angles and back-scattered. Mie Scattering Theory accounts for this different behavior. In order to make particle size measurements, the light intensity pattern is measured over the full angular range. When the particle size is larger than the wavelength of the incident light, the Mie equation reduces to the Fraunhofer equation. An array of detectors, including high-angle and back-scatter detectors, and multiple light sources of different wavelengths are typically employed to allow measurement of the full size range in one analysis. Equipment suited for use in analyzing particle size by laser light diffraction is commercially available and manufactured, for example, by Horiba Instruments, Irvine, CA.

[0013] The term "volume diameter" as used herein refers to the size of a particle as measured using a laser diffraction particle size analyzer, operating in the Mie Scattering Theory diffraction mode, equipped with a suspension dispersion sample chamber (e.g., as available from Horiba Instruments, Irvine, CA, USA). For purposes of the present description, volume diameter is given as a particle size distribution defined by one or more of D (0.10), D (0.50) and D (0.90). When referred to herein, the term D (0.10) indicates the volume frequency distribution of particles for which 10% of the sample is below the referenced size, the term D (0.50) indicates the volume frequency distribution of particles for which 50% of the sample is below the referenced size, and the term D (0.90) indicates the volume frequency distribution of particles for which 90% of the sample is below the referenced size.

[0014] The term "parenteral" as used herein refers to delivery of an active agent or formulation to a subject via any route or means other than oral administration. For example, for purposes of the present disclosure, parenteral formulations include formulations and systems for topical, transdermal, and buccal delivery. The term "parenteral" as contemplated herein further encompasses delivery via suppository and compositions suited to formulation as a suppository. For pur-

poses of the present disclosure, the term "parenteral" additionally encompasses delivery via infusion or injection, such as, for example, intravenous injection, intravenous infusion, intravascular injection, subcutaneous injection, and intramuscular injection.

[0015] As used herein, "pharmaceutical composition" refers to a composition that includes genistein in combination with one or more pharmaceutically acceptable excipients or adjuvants and is suitable for oral or parenteral administration to a subject.


[0016] The term "radioprotective agent" refers to agents that protect cells or living organisms from the deleterious cellular effects that result from exposure to ionizing radiation. These deleterious cellular effects include damage to cellular DNA, such as DNA strand break, disruption in cellular function, cell death and/or carcinogenesis. More particularly, the hematopoietic system is a rapidly dividing system and is therefore centrally affected by exposure to high-dose whole body ionizing radiation. Bone marrow aplasia and the resultant leukopenia, erythropenia and thrombocytopenia predispose the animal or human to infection, hemorrhage and ultimately death. For purposes of the present disclosure, a radioprotective agent may be one that is administered prophylactically prior to potential radiation exposure, with such administration resulting in the prevention, reduction in severity, or slowing of the symptoms or effects of exposure to ionizing radiation, should such an exposure occur. Additionally, a radioprotective agent may be used as a treatment for radiation exposure, administered after exposure to ionizing radiation has occurred, with such administration resulting in mitigation (i.e., prevention, reduction in severity, slowing, halting, or reversal of symptoms or effects that are otherwise associated with exposure to a given dose of ionizing radiation).

[0017] A "subject" for purposes of this disclosure is an animal to which a formulation as described herein can be administered in order to achieve a therapeutic effect. Optionally, the subject is a human being.

[0018] "Therapeutically effective" refers to an amount of genistein or an amount of a formulation of genistein as described herein which achieves a therapeutic effect by inhibiting a disease or disorder in a patient or by prophylactically inhibiting or preventing the onset of a disease or disorder. A therapeutically effective amount may be an amount which relieves to some extent one or more symptoms of a disease or disorder in a patient; returns to normal either partially or completely one or more physiological or biochemical parameters associated with or causative of the disease or disorder; and/or reduces the likelihood of the onset of the disease or disorder.

II. Genistein Formulations

[0019] Genistein is one of several known isoflavones that are normally found in plants. The main sources of natural genistein are soybeans and other legumes. Genistein is commercially available and may be obtained in synthetic, purified form. Synthetic genistein is available, for example, as BONISTEIN from DSM Nutritional Products (DSM Nutritional Products, Inc. Parsippany, NJ). Genistein's chemical name is 5,7-dihydroxy-3-(4-hydroxyphenyl)-chromen-4-one (IUPAC). Genistein's chemical structure is shown as:



[0020] The genistein formulations described herein are suspension formulations that include nanoparticulate genistein suspended in a suspension medium formed of one or more carriers, excipients, and/or diluents. In particular formulations, the formulations are provided as pharmaceutical compositions, and the carriers, excipients and/or diluents used in forming such compositions are selected from pharmaceutically acceptable materials. Pharmaceutically acceptable carriers, excipients and diluents suited for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Maack Publishing Co. (A.R. Gennaro (Ed.) 1985). In one such composition, the formulations as disclosed herein may include a suspension comprising nanoparticulate genistein suspended within a suspension medium including a water soluble polymer and a nonionic surfactant. The genistein used in the formulations described herein may be naturally derived or synthetically produced genistein. Pharmaceutical compositions of genistein as described herein can be formulated to be simultaneously suitable for both oral and parenteral administration. Though the formulations of genistein described herein are characterized as suspensions, in some formulations, depending on the carriers, excipients and diluents included in the medium within which the nanoparticulate genistein is suspended, a measureable amount of genistein may also be dissolved within the suspension medium.

[0021] Nanoparticulate genistein suitable for use in the formulations disclosed herein may be prepared according to

known methods for producing nano-sized particles. In one composition, natural or synthetic genistein material may be nanomilled according to milling techniques known in the art. Nanomilling may optionally include wet bead milling utilizing an agitator bead mill in a horizontal grinding container for continuous dispersion and fine wet grinding. In another composition, using a bead mill such as a DYNO-mill (CB Mills, Gurnee, IL), the necessary energy for dispersion and grinding is transmitted to the grinding beads through agitator discs mounted on an agitator shaft.

[0022] In one composition, a nanoparticulate genistein composition as described herein is provided by introducing genistein suspended in a pharmaceutically acceptable suspension medium into a bead mill. In such a composition, the genistein suspension is fed into the bead mill and milled in a manner that results in a pharmaceutical genistein composition characterized by nanoparticulate genistein suspended within the pharmaceutically acceptable suspension medium. In one such composition, a genistein suspension formulation may be fed continuously through the bead mill until a suspension composition containing nanoparticulate genistein material of a defined particle size distribution is reached. For example, the genistein formulation may be nanomilled by recirculating the volume of the suspension through the bead mill, followed by one or more single passes through a bead mill to reach a pharmaceutical composition exhibiting the desired genistein particle size distribution. The particle size of the genistein material suspended within a pharmaceutical composition as described herein can be controlled by adjusting the parameters of the bead mill and the grinding conditions. For example, the particle size produced by nanomilling genistein or a genistein suspension formulation in a bead mill may be controlled by bead size, bead load/suspension weight ratio, suspension composition, agitation rate, and milling time.

[0023] Though nanomilling is generally referenced herein as a means for producing nanoparticulate material suitable for use in the formulations described herein, the nanoparticulate material can be produced other suitable techniques as well. For example, nanoparticulate genistein material as used herein can be produced using one or more known wet milling techniques, super-critical or compressed fluid techniques, hot or high-pressure homogenization, emulsification techniques, evaporative precipitation, antisolvent precipitation, microprecipitation, cryogenic techniques, complexation techniques, ultrasonication techniques, or solid dispersion techniques. Spray drying and lyophilization may be used post-processing to isolate nanoparticles resulting from an aqueous or solvent dispersion technique.

[0024] The genistein included in the formulations described herein is a nanoparticulate material as defined herein. Optionally, the compositions disclosed herein may comprise nanoparticulate genistein material exhibiting a D (0.50) of 0.2 μm or less. In one such composition, the nanoparticulate genistein material exhibits a D (0.50) of 0.2 μm or less and a D (0.90) of 0.5 μm or less. In yet another embodiment, the nanoparticulate genistein material exhibits a D (0.50) of 0.2 μm or less and a D (0.90) of 0.4 μm or less.

[0025] The nanoparticulate genistein material included in the formulations described herein is suspended within a suspension medium that includes one or more carriers, excipients and/or diluents. As described herein, in particular formulations, such carriers, excipients and diluents are selected from pharmaceutically acceptable materials to facilitate preparation of pharmaceutical compositions that can be administered to a subject at risk for or suffering from a disease or disorder, such as, for example, a disease or disorder as described herein.

[0026] One or more nonionic surfactants may be included in the suspension medium to facilitate wetting and aid in preventing agglomeration of the nanoparticulate genistein drug substance. Nonionic surfactants suitable for use in the formulations described herein may be selected from, for example, polysorbates, poloxamers, polyoxyethylene castor oil derivatives (e.g., Cremophor EL, Cremophor RH60), bile salts, lecithin, 12-Hydroxystearic acid-polyethylene glycol copolymer (e.g., Solutol HS 15), and the like. Optionally, the formulations described herein include a nonionic surfactant selected from polysorbate 80 (Tween 80), polysorbate 20 (Tween 20), Poloxamer 188, and combinations thereof. In one formulation, the total nonionic surfactant content ranges from about 0.01% to about 10% by weight (w/w). In another formulation, the total nonionic surfactant content ranges from about 0.1% to about 10% (w/w). In certain such formulations, the total amount of nonionic surfactant is selected from about 0.2% to about 5% (w/w), about 0.2% to about 1% (w/w), about 0.2% to about 1% (w/w), about 0.2% to about 0.6% (w/w), and about 0.2% to about 0.8% (w/w).

[0027] The suspension formulations described herein may include one or more water soluble polymers, which may serve, for example, to enhance the viscosity of the suspension or to stabilize nanoparticulate genistein against particle agglomeration or potential deleterious effects from other formulation components. Water soluble polymers are pharmaceutically acceptable polymers that can be dissolved or dispersed in water. Suitable water soluble polymers for use in the formulations described herein may be selected from, for example, vegetable gums, such as alginates, pectin, guar gum, and xanthan gum, modified starches, polyvinyl pyrrolidone (PVP), hypromellose (HPMC), methylcellulose, and other cellulose derivatives, such as sodium carboxymethylcellulose, hydroxypropylcellulose, and the like. Optionally, the formulations described herein may include a poloxamer, such as Poloxamer 188, as a water soluble polymer. Poloxamer 188 is both a polymer and surfactant. Optionally, the formulations described herein may include Povidone K17 as a water soluble polymer. Where one or more water soluble polymers are included in the suspension formulations described herein, in specific compositions, the total water soluble polymer content ranges from about 0.5% to about 15% (w/w). For example, in certain such compositions, the total water soluble polymer content ranges from about 1% to about 10% (w/w). In other such compositions, the total water soluble polymer content may be selected from about

5% to about 15% (w/w), about 10% to about 15% (w/w), and 12% to about 15% (w/w), about 1% to about 8% (w/w), and about 1% to about 5% (w/w).

[0028] In particular compositions, the suspension medium included in the suspension formulation includes a combination of one or more nonionic surfactants with one or more water soluble polymers. Where that is the case, the nonionic surfactant constituent and water soluble polymer constituent can be selected from the materials already described herein, including combinations of such materials. Moreover, where the suspension medium includes a combination of nonionic surfactant and water soluble polymer, the total nonionic surfactant and total water soluble polymer included in the suspension formulation can be selected from those amounts already detailed. For example, where the suspension medium includes a combination of nonionic surfactant and water soluble polymer, the total nonionic surfactant content may be selected from about 0.01% to about 10% (w/w), about 0.1% to about 10% (w/w), about 0.2% to about 5% (w/w), about 0.2% to about 1% (w/w), about 0.2% to about 1% (w/w), about 0.2% to about 0.6% (w/w), and about 0.2% to about 0.8% (w/w), and the total water soluble polymer content may be selected from about 0.5% to about 15% (w/w), about 1% to about 10% (w/w), about 5% to about 15% (w/w), about 10% to about 15% (w/w), about 12% to about 15% (w/w), about 1% to about 8% (w/w), and about 1% to about 5% (w/w). In one such composition, the nonionic surfactant constituent may be present in an amount ranging from about 0.1% to about 1% (w/w) and the water soluble polymer constituent may be present in an amount ranging from about 1% to about 15% (w/w). In another such composition, the nonionic surfactant constituent may be present in an amount ranging from about 0.2% to about 1% (w/w) and the water soluble polymer constituent may be present in an amount ranging from about 5% to about 15% (w/w). In specific suspension formulations where the suspension medium includes both a nonionic surfactant and a water soluble polymer, the nonionic surfactant may be selected from a polysorbate, such as polysorbate 80 (Tween 80) and polysorbate 20 (Tween 20), the water soluble polymer may be selected from a poloxamer, such as Poloxamer 188, and a PVP, such as Povidone K17, with the nonionic surfactant and water soluble polymer being included in the formulation at any of the relative amounts detailed herein.

[0029] The suspension medium included in the suspension formulations according to the present description may also include a carrier. For example, carriers suitable for use in the formulations described herein include pharmaceutically acceptable aqueous carriers such as, for example, sterile water, physiologically buffered saline Hank's solution, Ringer's solution, and any other suitable aqueous carrier. The suspension formulations described herein can utilize buffers such as, for example, one or more of a citrate buffer, phosphate buffer, TRIS buffer, and a borate buffer to achieve a desired pH and osmolality. For example, the typical pH range for formulating injectable pharmaceutical products is from about 2 to about 12. In some compositions, the pH of the formulation may fall in a range that more closely approximates physiologic pH. For example, in certain compositions, the suspension formulations described herein are formulated to exhibit a pH selected from a range of from about 4 to about 8 and a range of from about 5 to about 7.

[0030] The suspension formulations described herein can also include one or more diluents. Suitable diluents may be selected from, for example, pharmaceutically acceptable buffers, solvents and surfactants.

[0031] Suspension formulations prepared as described herein are suited to providing high concentration genistein formulations (i.e., formulations containing genistein in amounts of about 250 mg/mL or greater). Though genistein exhibits low to virtually no solubility in several pharmaceutically acceptable solvents, the nanoparticulate suspension formulations described herein can incorporate genistein up to and over 300 mg/ml. In specific compositions, genistein formulations as described herein may incorporate genistein in amounts ranging from about 250 mg/mL to about 500 mg/mL. In certain such compositions, the amount of genistein included in a suspension formulation as described herein may be selected from about 200 mg/ml to about 400 mg/ml, from about 250 mg/ml to about 350 mg/ml, and from about 275 mg/ml and about 325 mg/ml.

[0032] The relative amount of genistein included in the suspension formulations described herein may be varied, as desired, to achieve a formulation having a desired total content of genistein. For example, the suspension formulations as described herein may include up to about 85% (w/w) genistein. In certain such formulations, the relative amount of genistein is selected from up to about 75% (w/w), up to about 65% (w/w), and up to about 50% (w/w). Optionally, the suspension formulations described herein may include an amount of genistein selected from a range of about 40% to about 75% (w/w), a range of about 40% to about 65% (w/w), a range of about 40% to about 50% (w/w), a range of about 50% to about 75% (w/w), and a range of about 50% to about 65% (w/w).

[0033] The inventors have also found that suspension compositions prepared according to the present description can increase bioavailability of genistein relative to solution formulations. In particular, as is illustrated in the experimental examples that follow, suspension formulations prepared as described herein exhibited significantly improved relative bioavailability when compared to solution formulations prepared using, for example, pharmaceutically acceptable PEG solvent. Such a result runs counter to what would be generally expected. For example, in certain compositions, relative to a solution formulation of genistein or formulations of genistein incorporating larger sized genistein material, a suspension formulation as described herein provides an increase in peak total genistein serum concentrations of up to 300%. In particular such compositions, the increase in peak total genistein serum concentration ranges from about 50% to about 300%. In other such compositions, the increase in peak total genistein serum concentration is selected from about

50% or greater, about 75% or greater, about 100% or greater, and about 200% or greater.

[0034] The combination of high drug loading and significantly increased relative bioavailability provided by formulations described herein present several advantages. The significant jump in drug loading by the genistein suspension formulations described herein facilitates administration of therapeutically effective amounts of genistein to subjects in need thereof using much less formulated drug substance, which, in turn, can increase patient compliance and facilitate manufacture of a genistein drug product that is much better suited to administration of genistein in therapeutic contexts requiring delivery of relatively high doses of genistein. Moreover, the increase in bioavailability afforded by the genistein suspension formulations described herein provides the added benefit of reducing the amount of genistein that must be delivered to a subject in order to achieve and maintain therapeutic genistein blood plasma levels. Therefore, the formulations described herein offer a significant reduction in the relative amount of administered genistein required to achieve and maintain a therapeutic benefit, which can reduce the costs of genistein treatments, work to mitigate or avoid potential side effects that may be associated with relatively higher doses of the compound, and further decreases the amount of formulated drug substance required to achieve and maintain therapeutic efficacy.

[0035] Even further, the suspension formulations taught herein can be formulated such that that a single given formulation is suited to both oral and parenteral delivery. Where a suspension formulation as described herein is prepared for parenteral delivery it can be manufactured according to standard methods to provide a sterile composition deliverable via, for example, intravenous injection or infusion, intravascular injection, subcutaneous injection, or intramuscular injection. The suspension formulations described herein can be prepared to exhibit viscosities suited for the desired route of parenteral administration and can be manufactured and packaged in any manner suited to the desired application, including, for example, as a formulation deliverable via intravenous injection or infusion, intravascular injection, subcutaneous injection, or intramuscular injection. Optionally, the formulations described herein may be included in pre-filled syringes prepared for administration of a given dose or range of doses of genistein.

[0036] Where prepared for oral administration, the formulations may be prepared in any suitable manner and using any suitable devices for oral administration of desired doses of genistein. For example, when the formulations described herein are prepared for oral delivery, they may be prepared as a liquid suspension that can be metered to deliver a desired dose or incorporated into capsules, such as gelatin or soft capsules, suited for delivery of liquid formulations. Alternatively, formulations as described herein prepared for oral administration can be loaded into prefilled sachets or premeasured dosing cups. Genistein formulations prepared for oral administration may optionally include one or more pharmaceutically acceptable sweetening agents, preservatives, dyestuffs, flavorings, or any combination thereof.

III. Methods

[0037] The genistein suspension formulations described herein can be used to treat subjects suffering from or at risk for a disease or disorder treatable with genistein. Clinical trials, animal studies, cell-culture experiments, and epidemiological studies have provided evidence that genistein exerts various physiological effects. Examples of diseases and disorders amenable to treatment by genistein are described herein. However, the potential therapeutic applications of genistein are not limited to those described herein, and genistein formulations according to the present description can be used to treat a subject at risk for or suffering from any disease or disorder for which administration of genistein will be therapeutically effective.

[0038] As one example, genistein has displayed antitumor, antimetastatic and antiangiogenic (suppression of blood-vessel growth) properties in tissue culture and *in vivo*. Several epidemiological studies suggest that soybean consumption may contribute to lower incidence of breast, colon, prostate, thyroid, and head and neck cancers - an effect that is attributed to genistein and other isoflavones (Takimoto et al., Cancer Epidemiol Biomarkers Prev. 2003 Nov; 12(11 Pt 1): 1213-21; Wei et al., J Nutr. 2003 Nov; 133(11 Suppl 1): 3811 S-3819S; Sakar, F. H. and Y. Li, Cancer Invest. 2003; 21(5): 744-57; Magee P. J. and I. R. Roland, Br J Nutr. 2004 Apr; 91(4): 513-31; Park, O. J. and Y. J. Surh, Toxicol Lett. 2004 Apr 15; 150(1): 43-56; Messina, M. J., Nutr Re. 2003 Apr; 61(4): 117-31). Genistein has also been reported to inhibit non-Hodgkin's lymphoma, melanoma, lung cancers, and ovarian cancer (Wei et al. 2003; Mohammad et al., Mol Cancer Ther. 2003 Dec; 2(12): 1361-8; Nicosia et al., Hematol Oncol Clin North Am. 2003 Aug; 17(4): 927-43; Sun et al., Nutr Cancer. 2001; 39(1): 85-95). Tissue culture experiments suggest that genistein's cancer-fighting effects occur at dosages that are hard to attain from food alone, unless one eats very large amounts of soy products. Reliable genistein dosing therefore requires the use of concentrated supplements (Magee and Roland 2004).

[0039] The genistein formulations may, therefore, be used in methods of inhibiting the onset, development or progression of certain cancers, such as cancers selected from breast, colon, prostate, thyroid, and head and neck cancers. In one such method, a subject at risk for developing a breast, colon, prostate, thyroid, head or neck cancer is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject. The genistein formulations described herein may also be used in methods of treating cancer. In a particular method, a patient at risk for or suffering from a cancer responsive to genistein treatment, such as for example, a cancer selected from non-Hodgkin's lymphoma, melanoma, lung cancers, and ovarian cancer is identified and a

therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject.

[0040] The ability of genistein and related soy isoflavones to reduce post-menopausal bone-loss has also been shown in many studies. These substances prevent bone loss and promote bone formation, especially in the spine. Among the dosage regimens found to be effective are: 1 mg/day genistein + 0.5 mg/day daidzein + 42 mg/day other isoflavones (biochanin A and formononetin, in this case); 54 mg/day genistein; 57 mg/day isoflavones; 65 mg/day isoflavones; 90 mg/day isoflavones (Morabito et al. J Bone Miner Res. 2002 Oct; 17(10): 1904-12; Cotter A. and K. D. Cashman, Nutr Rev. 2003 Oct; 61(10): 346-51; Atkinson et al., Am J Clin Nutr. 2004 Feb; 79(2): 326-33; Setchell K. D. and E. Lydeking-Olsen, Am J Clin Nutr. 2003 Sep; 78(3 Suppl); 593S-609S; Clifton-Bligh et al., Menopause. 2001 Jul-Aug; 8(4): 259-65; Fitzpatrick, L. A., 2003 Mar 14; 44 Supl 1: S21-9). Therefore, methods for reducing post-menopausal bone-loss are also provided herein. In one example, such a method includes identifying a subject at risk for or suffering from post-menopausal bone loss and administering to the subject a therapeutically effective amount of a genistein formulation selected from any of those described herein. Alternatively, methods for promoting bone formation are also provided. In one such example, a method for promoting bone formation, such as in the spine, includes identifying a subject at risk for or suffering from loss of bone mass and administering to the subject a therapeutically effective amount of a genistein formulation selected from any of those described herein.

[0041] Genistein has also been suggested for use in treating cystic fibrosis. The main clinical symptoms of cystic fibrosis are chronic obstructive lung disease, which is responsible for most of the morbidity and mortality associated with cystic fibrosis, and pancreatic insufficiency. Cystic fibrosis (CF) is caused by a mutation in the cystic fibrosis transmembrane conductance regulator (CFTR), a plasma membrane protein. CFTR functions as a chloride channel, and about 1000 mutations of the gene coding for CFTR are currently known. The most common of these known mutations results in a deletion of a phenylalanine at position 508 of the CFTR protein. This mutation is referred to as Delta508 and is present in the majority of patients suffering from cystic fibrosis. The Delta508 mutation results in an aberrant CFTR that is not transported to the plasma membrane, but is instead degraded in the ubiquitin-proteasome pathway. One approach for developing a treatment for cystic fibrosis is to inhibit the breakdown of DeltaF508-CFTR by interfering with the chaperone proteins involved in the folding of CFTR. Genistein has been shown in *in-vitro* systems to inhibit the breakdown of DeltaF508-CFTR through interference the relevant chaperone proteins. In addition, it has been shown that it is possible to stimulate CFTR or its mutated forms, when present in the plasma membrane, using genistein (Roomans, G. M., Am J Respir Med. 2003; 2(5): 413-31).

[0042] The genistein formulations described herein may be used in treating cystic fibrosis. In an example of such a method, a subject at risk for or suffering from cystic fibrosis is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject. In a particular example, a subject at risk for or suffering from cystic fibrosis associated with DeltaF508-CFTR is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject. In each example of a method for treating cystic fibrosis described herein, the therapeutically effective amount of genistein formulation administered to the subject is sufficient to accomplish one or more of the following: inhibit the breakdown of DeltaF508-CFTR; inhibit or prevent the onset of cystic fibrosis or one or more symptoms associated with cystic fibrosis; mitigate or reduce the severity of one or more symptoms associated with cystic fibrosis; delay the progression of cystic fibrosis or the worsening of one or more symptoms associated with cystic fibrosis.

[0043] Genistein appears to increase the rate at which fats are metabolized by the body, and to decrease the rate at which they are deposited in the tissues (Goodman-Gruen, D. and D. Kritz-Silverstein, Menopause. 2003 Sep-Oct; 10(5): 427-32). Moreover, in clinical studies of humans and animals, the consumption of genistein and daidzein resulted in loss of body fat, lower fasting insulin concentrations, lower LDL and higher HDL cholesterol, and improved insulin responses to blood sugar. Cholesterol benefits were seen at dosages of 42 mg/day of genistein plus 27 mg/day of daidzein (Bhathena, S. J. and M. T. Velasquez, Am J Clin Nutr. 2002 Dec; 76(6): 1191-201; Urban et al., J Urol. 2001 Jan; 165(1): 294-300). In addition to lowering LDL and raising HDL (mentioned above), genistein prevents the oxidation of LDL, a process thought to contribute to arterial plaques (Young, S. G. and S. Parthasarathy, West J Med. 1994 Feb; 160(2): 153-54). The genistein formulations described herein can be used in methods for lowering LDL and/or raising HDL in subjects in need thereof. In one such method, a subject at risk for or suffering from a high circulating level of LDL is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject, wherein the therapeutically effective amount of genistein formulation is sufficient to lower the LDL levels or prevent or delay an increase in circulating LDL levels in the subject. In another method, a subject that could benefit from an increase in circulating levels HDL is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject, wherein the therapeutically effective amount of genistein formulation is sufficient to increase circulating HDL levels or prevent or delay decrease in circulating HDL levels in the subject.

[0044] Genistein is also a radioprotective agent. For example, genistein has been reported to increase hematopoiesis and survival in irradiated mice (Zhou, 2005; Land Auer, 2001, 2003 & 2005). The mechanism of action for this radiopro-

5 tective effect may potentially involve several of genistein's known effects including inhibition of protein tyrosine kinases (PTKs) and PTK-triggered apoptosis, inhibition of topoisomerase II, inhibition of phosphatidylinositol turnover and the second messenger system, both agonist and antagonist estrogenic effects, reduction of stress gene expression through inactivation of Y/CCA-AT binding factor, increased antioxidant activity, apoptosis, cell cycle arrest and differentiation, improved immune defenses and/or increased AKT kinase levels. The beneficial effects of genistein may also be due, in part, to its antioxidant properties, reducing free radicals and stabilizing the cell membrane structure. Further, genistein may also have a role in protecting stem cells and/or stimulating proliferation.

10 **[0045]** Genistein administered prior to, during, and/or after exposure to radiation, may be used to eliminate or reduce the severity of deleterious cellular effects caused by exposure to ionizing radiation resulting from, for example, from a nuclear explosion, a spill of radioactive material, close proximity to radioactive material, cancer radiation therapy, diagnostic tests that utilize radiation, and the like. Genistein can be used for the treatment and prevention of Acute Radiation Syndrome (ARS) (sometimes known as radiation toxicity or radiation sickness). ARS is an acute illness caused by irradiation of a substantial portion of the body by a high dose of penetrating radiation (i.e., greater than 0.7 Gray (Gy) or 70 rads, with mild symptoms possible at doses as low as 0.3 Gy or 30 rads) over a very short period of time (usually a matter of minutes). It is thought that the major cause of ARS is depletion of immature parenchymal stem cells in specific tissues.

15 **[0046]** Methods for treating radiation exposure are, therefore provided herein. In each method, a subject at risk of or that has suffered from exposure to radiation is identified and a therapeutically effective amount of a genistein formulation selected from any of those described herein is administered to the subject. In specific examples, the method of treating radiation exposure is a method for preventing ARS, wherein a subject at risk of ARS is identified and a therapeutically effective amount of a genistein formulation as described herein is administered to the subject before the subject is exposed to radiation. In other examples, the method of treating radiation exposure is a method for treating ARS, wherein a subject suffering from ARS is identified and a therapeutically effective amount of a genistein formulation as described herein is administered to the subject after the subject has suffered exposure to radiation. In yet other methods, a subject at risk of radiation exposure is identified, a therapeutically effective amount of a genistein formulation as described herein is administered to the subject prior to exposure to radiation, and, in the event the subject suffers from radiation exposure, administration of therapeutically effective amounts of genistein is continued after the radiation exposure occurs.

20 **[0047]** In additional methods, subjects at risk for or having suffered from a radiation exposure resulting from an event selected from cancer radiation therapy or a diagnostic test utilizing radiation are identified, and the subjects are administered a therapeutically effective amount of the genistein formulation. In one such method, the genistein formulation is administered to the subject prior to radiation exposure in order to prevent or reduce the severity of the deleterious effects of such exposure. In another such method, the genistein formulation is administered to the subject after radiation exposure in order to mitigate, reverse or reduce the severity of the deleterious effects of such exposure. Optionally, the methods of treating radiation exposure resulting from an event selected from cancer radiation therapy or a diagnostic test utilizing radiation in a subject may include administering a genistein formulation as described herein both before and after radiation exposure.

25 **[0048]** In each of the examples of the methods described herein, the therapeutically effective amount of genistein formulation may be administered orally or parenterally. In specific methods, where the genistein formulation is administered parenterally, it may be administered, for example, via intravenous injection or infusion, subcutaneous injection, intravascular injection, or intramuscular injection. Where the formulation is administered orally, the formulation may be prepared in any manner suitable for oral administration, such as is described herein. The dose and dosing regimen most appropriate for a given therapeutic method described herein may depend upon, for example, the subject being treated, the nature of the disease or disorder, as well as the severity of any symptoms suffered. Using formulations prepared as described herein, one of skill in the art will be able to identify the appropriate dose and dosing regimen useful for achieving therapeutic efficacy in each of the methods described herein. The genistein formulations described herein may be administered, for example, as a single dose, a regular daily dose, a two-times daily dose, a three-times daily dose, or according to another desired dosing schedule.

30 **[0049]** The total daily dose of genistein delivered using a formulation or method described herein may depend on the desired condition to be treated or the desired therapeutic effect. Optionally, a therapeutically effective amount of a genistein formulation according to the present description may be an amount sufficient to deliver a dose of genistein ranging from about 50 mg/day to about 10,000 mg/day. Optionally, the amount of genistein formulation administered to the subject is sufficient to deliver a dose of genistein selected from about 50 mg/day to about 9,000 mg/day, about 50 mg/day to about 8,000 mg/day, about 50 mg/day to about 2,000 mg/day, about 100 mg/day to about 9,000 mg/day, about 100 mg/day to about 5,000 mg/day, about 100 mg/day to about 4,000 mg/day, and about 100 mg/day to about 2,000 mg/day.

Examples*Example 1-Solubility of Genistein*

[0050] Calculated pKa's for genistein range from 7-9, with the predicted solubility increasing above pH 7 in accordance with the lowest pKa. The calculated properties were used to design the appropriate pH range for the pH-solubility profile of genistein in several acceptable cosolvents, which was established to be pH 6 - 9. Solubility of genistein was increased at higher pH, however degradation was observed at pH 9. Table 1 shows the solubility results of genistein in selected pharmaceutically acceptable cosolvents.

Table 1

Vehicle	Solubility at 25 °C (mg/mL)
Propylene Glycol	6.2
Polyethylene Glycol 300 (PEG300)	110.5
Polyethylene Glycol 400 (PEG400)	115.1
Ethanol	25.0
Dimethyl acetamide (DMA)	141.3
N-methyl pyrrolidone (NMP)	238
Citrate buffer pH 6	Not detected
Phosphate buffer pH 7	Not detected
TRIS buffer pH 8	Not detected
Borate buffer pH 9, 3 days	Not detected
Borate buffer pH 9, 7 days	0.043
Borate buffer pH 9, 14 days	0.005

[0051] The solubility of genistein in water is not detectable at pH 6-7, which implies the solubility is less than 0.02 mg/mL or ~ 0.00002 g/g H₂O (lowest concentration). Based on the pH-solubility and solubility in cosolvents, it was determined that PEG400 would be the cosolvent to achieve highest solubility. Since parenteral formulations preferably have a maximum of 50% organic component, the addition of Ethanol (EtOH), N-methylpyrrolidone (NMP) and a surfactant were considered, since these would be expected to enhance absorption from the injection site. Ethanol has the added benefit of reducing viscosity. Polysorbate 80 (Tween 80) was considered as a surfactant due to its use in approved parenteral dosage forms at levels as high as 12% (FDA Inactive Ingredients Guide), although a more common range is 0.1-1%. Solubility was further evaluated in two aqueous/organic mixtures with concentrations acceptable for a parenteral dosage form. Additionally, cyclodextrin formulations were evaluated. Solubility testing results for genistein are given in Table 2.

Table 2

Vehicle	Solubility of GENISTEIN at 25 °C (mg/mL)
10% Polysorbate 80/40% PEG400/50% 25 mM Phosphate Buffered Saline (pH 7)	11
10% Polysorbate 80/10% EtOH/40% PEG400/40% 25 mM Phosphate Buffered Saline (pH 7)	12
10% Polysorbate 80/10% NMP/40%PEG400/40% 25 mM Phosphate Buffered Saline (pH 7)	19
30% Hydroxypropyl-β-cyclodextrin	6
30% Sulfobutylether-β-cyclodextrin	8

Example 2-Nanoparticulate Genistein Composition

[0052] None of the previously evaluated genistein formulations achieved the desired level of drug loading. In an effort to achieve higher drug loading, a sterile injectable suspension was prepared according to the present description. The formulation included nanoparticulate genistein that had been nanomilled with a vehicle solution of 5% Povidone K17 (w/w), 0.2% Polysorbate 80 (w/w), in 50 mM phosphate buffered saline (61 mM sodium chloride). The quantitative composition of the formulation is listed in Table 3.

Table 3

Component	Amount	Amount per 1 L
Genistein	300 mg/mL	300 g
Polysorbate 80	2 mg/mL	2 g
Povidone K17	40 mg/mL	50 g
50 mM Sodium Phosphate/61 mM Sodium Chloride	0.948 mg/mL	948 g

[0053] The function of each component and excipient listed in Table 1 is as follows: 1) Polysorbate 80 is included as a surfactant to enable wetting and aid in preventing agglomeration of suspended genistein drug substance, 2) Povidone K17 is included as a viscosity enhancer to aid in stabilizing the genistein drug substance suspension, and 3) Sodium Phosphate Buffer, Sodium Chloride is included as the diluent and to achieve physiological osmolality and maintain pH for parenteral administration of the composition.

[0054] The composition of 50 mM Sodium Phosphate Buffer/61 mM Sodium Chloride solution is as shown in Table 4.

Table 4

Component	Amount per 1 L
NaH ₂ PO ₄ •H ₂ O	6.9 g
NaCl	3.56 g
NaOH	(pH adjust)
HCl	(pH adjust)

Example 3-Second Nanoparticulate Genistein Composition

[0055] A second nanoparticulate genistein formulation as described herein was prepared. The nanomilled genistein was achieved using wet bead milling, utilizing an agitator bead mill in a horizontal grinding container for continuous dispersion and fine wet grinding in a closed system. A DYNO®-Mill Type Multi Lab agitator bead mill was used to prepare the nanoparticulate genistein, wherein the necessary energy for dispersion and grinding was transmitted to the grinding beads via the agitator discs mounted on the agitator shaft. Material was continuously fed into the mill via a product pump. The gap setting of the dynamic gap separator, the diameter of the beads, and length of the milling period were used to determine the particle size distribution. The product was fed continuously through the mill until a defined particle size distribution was reached. Although the DYNO®-Mill Type Multi Lab agitator bead mill was utilized in this work, other high energy, wet bead milling process equipment may be utilized.

[0056] Two formulations were tested that incorporated either Polysorbate 80 or Poloxamer 188 as a wetting agent to maintain a stable particle size distribution. Povidone (Polyvinyl pyrrolidone (PVP)) K17 was used at the 5% level in the formulations as a viscosity enhancer as well as a stabilizer against particle agglomeration. The quantitative composition of the formulation is given in Table 5.

Table 5

Component	Reference to Quality Standard	Amount (mg/mL)
Genistein	SP-001	300
Polysorbate 80	USP	2
Povidone K17	USP	50

(continued)

Component	Reference to Quality Standard	Amount (mg/mL)
Sterile Water for Injection	USP	QS to 1mL

[0057] The formulation may also include the replacement of the sterile water with injection with a phosphate-buffered saline for pH control and osmolality (e.g., as provided in the formulation described in Example 2).

[0058] The formulations exhibited an excellent, reproducible and stable particle size distribution profile, with d(0.5) of less than 0.2 μm . Optical microscopy confirmed a uniform particle size in the suspension. Powder X-Ray diffraction (XRD) was performed to examine physical, crystalline changes to the genistein material as a result of the milling process or as a result of a formulation incompatibility. Analyses performed indicate that there was no change in crystal form post-milling for genistein drug substance and milled suspensions containing 0.2% (w/w) polysorbate 80 with 5% (w/w) povidone K17, and 0.2% (w/w) poloxamer 188 with 5% (w/w) povidone K17.

[0059] The nanomilled genistein suspension comprised of nanoparticulate genistein (300 mg/mL) containing 0.2% (w/w) Polysorbate 80 with 5% (w/w) Povidone K17 was placed on stability at 5°C, 30 °C at 65% RH, and 40 °C at 75% RH. The suspensions were stored in 5 mL serum vials and 20 mm PTFE-faced butyl rubber stoppers. No impurities were observed after 7 months and there was no significant change in the particle size distribution.

Example 4-In-vivo Comparison of Genistein Suspension Formulation with Genistein Solution Formulation

[0060] This experiment evaluated a nanoparticulate formulation of genistein according to the present description and compared it to administration of genistein in a PEG 400 solution formulation. The genistein suspension formulation included nanomilled genistein in 50 nM phosphate buffered saline with 0.2% (w/w) Polysorbate 80 and 5% (w/w) PVP K17. The suspension formulation exhibited a pH of 6.96, and the nanoparticulate genistein incorporated into the suspension formulation exhibited a D (0.50) of 0.126 μm and a D (0.90) of 0.253 μm . The formulations were administered via subcutaneous injection("SC") 24 hr prior to irradiation. A separate vehicle and genistein group was included for each formulation. The study was conducted at two radiation doses, either, 8.75 Gy or 9.0 Gy.

[0061] Male CD2F1 mice were exposed to bilateral whole-body irradiation at a dose of 8.75 Gy or 9.0 Gy at 0.6 Gy/min. Thirty-day survival was the endpoint for this study. The different experimental groups evaluated in this study are detailed in Table 6.

Table 6

Experimental Groups:

Group	Route	Time of SC	Gy	N	30 Day Survival (%)
At 8.75 Gy:					
1.) Vehicle, PEG-400	SC	-24 Hr pre-rad	8.75	16	38%
2.) Genistein (PEG-400)	SC	-24 Hr pre-rad	8.75	16	81%*
At 9.0 Gy:					
3.) Vehicle (Nano))	SC	-24 Hr pre-rad	8.75	16	25%
4.) Genistein (Nano)	SC	-24 Hr pre-rad	8.75	16	100%*
At 9.0 Gy:					
1.) Vehicle, PEG-400	SC	-24 Hr pre-rad	9.0	16	38%
2.) Genistein (PEG-400)	SC	-24 Hr pre-rad	9.0	16	81%*
3.) Vehicle (Nano)	SC	-24 Hr pre-rad	9.0	16	19%
4.) Genistein (Nano)	SC	-24 Hr pre-rad	9.0	16	88%*

[0062] As shown in Table 6 and FIG. 1, the thirty-day survival rates in groups receiving solution formulation of genistein (Genistein (PEG-400)) and suspension formulation of genistein (Genistein (NANO)) at 8.75 Gy were 81% and 100%, respectively. Survival rates of the control groups (Vehicle (PEG-400) and Vehicle (Nano)) were 38% and 25%, respectively. At 9.0 Gy, 30-day survival rates of the Genistein (PEG-400) group and the Genistein (NANO) group were 81% and 88% respectively. The survival rates of the control groups (Vehicle (PEG-400) and Vehicle (Nano)) were 38% and 19%, respectively. Every group that received genistein 24 hr pre-irradiation were significantly ($p < 0.05$) different from their respective control group.

Example 5-In-vivo Comparison Genistein Suspension Formulation Administered Parenterally and Genistein Suspension Formulation and Genistein Solution Formulation Given Orally

[0063] This experiment evaluated the effect of a genistein nanoparticulate suspension formulation (Genistein-IS) prepared as described in Example 4 given via intramuscular injection ("IM") compared to the effect of a PEG 400 solution formulation and the Genistein IS suspension formulation given orally. The different formulations were administered twice daily for 6 days prior to irradiation. A positive control was also included which was Genistein-IS administered IM 24 hours prior to irradiation. A separate vehicle and genistein group was included for each group. The study was conducted at one radiation dose, 9.25 Gy. Male CD2F1 mice were exposed to bilateral whole-body irradiation at a dose of 9.25 Gy at 0.6 Gy/min. Thirty-day survival was the endpoint for this study.

Table 7

Experimental Groups:

Group	Route	Time of SC	Gy	N	30 Day Survival (%)
1) Vehicle, IS	IM	-24 Hr pre-rad	9.25	20	10%
2) Genistein IS	IM	-24 Hr pre-rad	9.25	20	85%*
3) Vehicle, IS	PO	BID for 6 days pre-rad	9.25	20	15%
4) Genistein IS	PO	BID for 6 days pre-rad	9.25	20	85%*
5) Vehicle, PEG 400	PO	BID for 6 days pre-rad	9.25	20	0%
6) Genistein/PEG 400	PO	BID for 6 days pre-rad	9.25	20	80%*
p < 0.05 two-tailed Fisher Exact Test (vehicle vs. genistein)					
BID = twice daily dosing IS=Injectable Suspension					

[0064] As shown in Table 7, the thirty-day survival rates of orally administered Genistein/PEG 400 and Genistein-IS at 9.25 Gy were 80% and 85%, respectively. Survival rates of the control groups (vehicle only administration) were 0% and 15%, respectively. The positive control group, Genistein IS administered IM 24 hours prior to irradiation, had a survival percentage of 85% vs. 10% for the vehicle.

[0065] Every group that received genistein either IM or orally were significantly ($p < 0.05$) different from their respective negative control group. There was not, however, a significant difference in survival between genistein/PEG-400, and the Genistein IS formulation.

Example 6- Radioprotection Time Course Study with Vehicle Injection Suspension and Genistein Nanoparticulate Injection Suspension Subcutaneously Administered 24 Hr, 18 Hr, 12 Hr, or 6 Hr Before 9.0 Gy ⁶⁰Co radiation

[0066] Previous experiments showed statistically significant radioprotective results when a nanoparticulate genistein injectable suspension prepared according to the present description (Genistein-IS) was administered 24 hr before irradiation in a saline based vehicle. This time course study was performed to determine whether there was a time-dependent effect on radioprotective efficacy with SC administered Genistein-IS. The time dependent effects of Genistein-IS were compared to a placebo formulation (Vehicle-IS). The Genistein-IS formulation was prepared as described in Example 4.

[0067] Male CD2F1 mice were used in this experiment. All groups received a single 200 mg/kg SC administration at 24 hr, 18 hr, 12 hr, or 6 hr before irradiation. SC injections were administered in the nape of the neck using a 25 G needle in an injection volume of 0.1 ml via a 1 ml tuberculin syringe. All mice were exposed to bilateral whole-body irradiation at a dose of 9.0 Gy at 0.6 Gy/min. Thirty-day survival was the endpoint for this study.

Table 8

Experimental Groups:

Group	Route	Time of Dose	Gy	N	30-Day Survival (%)
1.) Vehicle-IS	SC	24 Hr pre-rad	9.0	16	44%
2.) Genistein-IS	SC	24 Hr pre-rad	9.0	16	88%*
3.) Vehicle-IS	SC	18 Hr pre-rad	9.0	16	13%
4.) Genistein-IS	SC	18 Hr pre-rad	9.0	16	69%*
5.) Vehicle-IS	SC	12 Hr pre-rad	9.0	16	44%

(continued)

Experimental Groups:

Group	Route	Time of Dose	Gy	N	30-Day Survival (%)
6.) Genistein-IS	SC	12 Hr pre-rad	9.0	16	81%
7.) Vehicle-IS	SC	6 Hr pre-rad	9.0	16	38%
8.) Genistein-IS	SC	6 Hr pre-rad	9.0	16	63%

* p < 0.05 two-tailed Fisher Exact Test (vehicle vs. genistein)

[0068] The results shown in Table 8 and FIG. 2 demonstrate that a single SC administration of Genistein-IS administered at 24, 18, 12, or 6 hr pre-irradiation resulted in 30-day survival rates of 88%, 69%, 81%, and 63%, respectively. The survival rates for the Vehicle-IS groups at the corresponding time points were 44%, 13%, 44%, and 38%, respectively. Genistein-IS resulted in significant radioprotection when injected either 24 or 18 hr before irradiation ($p < 0.05$).

Example 7- Effect of Subcutaneous vs. Intramuscular injection of Nanoparticulate Genistein Formulation on Radioprotective Efficacy When Administered 24 hr Before 9.25 Gy⁶⁰Co Radiation

[0069] The purpose of this experiment was to compare the radioprotective efficacy of a nanoparticulate genistein suspension formulation prepared as described herein (Genistein-IS) delivered to provide a genistein dose of 200 mg/kg when administered SC or IM. The Genistein-IS formulation was prepared as described in Example 4. Male CD2F1 mice were used in this experiment. Groups were given a single SC or IM injection of an injectable placebo suspension (Vehicle-IS) or Genistein-IS (200 mg/kg) 24 hr before irradiation. Also included in this experiment were groups that received a solution formulation of genistein in PEG 400 (Genistein) delivered to provide a genistein dose of 200 mg/kg or placebo PEG 400 formulation (PEG 400) administered SC 24 hr before irradiation.

[0070] All vehicle and Genistein-IS groups received a single 200 mg/kg SC or IM injection at 24 hr before irradiation. SC injections were administered in the nape of the neck using a 25 G needle in an injection volume of 0.1 ml via a 1 ml tuberculin syringe. Mice were administered Vehicle-IS or Genistein-IS by IM injection into the quadriceps muscle using a 25 G needle attached to a Hamilton syringe. The injection volume was 50 μ l.

[0071] Mice were exposed to bilateral whole-body irradiation at a dose of 9.25 Gy at 0.6 Gy/min. Thirty-day survival was the endpoint for this study.

Table 9

Experimental Groups:

Group	Route	Time of Dose	Gy	N	30-Day Survival (%)
1.) PEG 400	SC	24 Hr pre-rad	9.25	20	15%
2.) Genistein	SC	24 Hr pre-rad	9.25	20	75%*
3.) Vehicle IS	SC	24 Hr pre-rad	9.25	20	30%
4.) Genistein IS	SC	24 Hr pre-rad	9.25	20	85%*
5.) Vehicle IS	IM	24 Hr pre-rad	9.25	20	15%
6.) Genistein IS	IM	24 Hr pre-rad	9.25	20	75%*

* p < 0.05 two-tailed Fisher Exact Test (vehicle vs. genistein)

[0072] Survival rates for vehicle-PEG 400 and genistein PEG 400 administered SC were 15% and 75%, respectively. This resulted in statistically significant radioprotection by genistein over PEG 400 vehicle ($p < 0.05$) (shown in Table 9 and FIG. 3).

[0073] When Vehicle-IS or Genistein-IS was administered SC, 30-day survival rates were 30% and 85%, respectively. For IM administration, survival rates for Vehicle-IS and Genistein-IS, were 15% and 75%, respectively. For both SC and IM routes, genistein provided significant protection over vehicle ($p < 0.05$) (shown in Table 9 and FIG. 3).

[0074] These results demonstrate that nanoparticulate genistein formulations prepared according to the present description provide similar levels of radioprotection when administered by either the SC or IM route.

Example 8- Pharmacokinetics in Mice Following Intravenous or Intramuscular Injection of ¹⁴C-Genistein

[0075] As shown in the results of Table 10, male CD1 mice were administered a single IM dose (Group 2, nominal 200 mg/kg) or a single IV bolus dose (Group 3, nominal 50 mg/kg) of ¹⁴C-Genistein. The genistein formulation used was

a suspension formulation that includes genistein suspended in sterile water with 0.2% (w/w) Polysorbate 80 and 5% (w/w) PVP K17. The genistein material used in the suspension formulation exhibited a D (0.50) of 0.136 μm and a D(0.90) of 0.310 μm . Following dosing, the content and concentration of radioactivity in blood, plasma, excreta and carcass, and the non-compartmental pharmacokinetics of total radioactivity in whole blood and plasma were determined. Following dose administration, the extent and severity of any clinical signs was assessed. The dose level of 200 mg/kg (IM) and 50 mg/kg (IV) were well tolerated and therefore chosen for the main study.

[0076] The concentration of radioactivity in both dose formulations was measured pre and post dose by liquid scintillation spectroscopy and was similar on both occasions. The radiochemical stability of the test article in both dose formulations was assessed prior to and following the administration of the dose. The mean radiochemical stability of the test article in the dose formulation samples from the intramuscular dose formulation (Group 2) were 98.5% and 98.2%, respectively. The mean radiochemical stability values for pre and post dose samples from the intravenous dose formulation (Group 3) were 98.3% and 98.5%, respectively. Therefore, the ^{14}C -Genistein in both formulations was considered to have been radiochemically stable throughout the dosing period. No treatment-related clinical signs were observed in any of the main study male mice following a single IM dose of ^{14}C -Genistein (200 mg/kg) or a single IV dose of ^{14}C -Genistein (50 mg/kg).

[0077] Whole-blood samples were collected and plasma was obtained by centrifugation. The concentration of radioactivity in whole-blood and plasma was measured by liquid scintillation spectroscopy. Pharmacokinetic parameters were calculated from the composite concentration vs. time profiles and are presented in Table 10.

Table 10

Pharmacokinetic		Group 2 (IM)		Group 3 (IV)	
Parameter	Units	Blood	Plasma	Blood	Plasma
t_{max}	h	0.50	0.50	0	0
C_{max}	$\mu\text{g eq/mL}$	33.3	63.1	60.1	108
t_{last}	h	168	168	168	168
$\text{AUC}_{0\text{-}t_{\text{last}}}$	$\mu\text{g eq}\cdot\text{h/mL}$	222	375	99.7	116
R^2		0.977	0.943	0.850	0.999
k_{e1}	h^{-1}	0.0242	0.0145	0.0118	0.0307
$t_{1/2}$	h	28.7	47.8	58.5	22.6
$\text{AUC}_{0\text{-}\infty}$	$\mu\text{g eq}\cdot\text{h/mL}$	223	387	112	116
Extrapolation	%	0.663	3.18	11.4	0.191
V_{d}	mL/kg	NA	NA	37554	14042
CL	m L/h/kg	NA	NA	445	431
Bioavailability	%	49.7	83.5	NA	NA

[0078] For Group 2, the highest mean concentration of radioactivity in plasma and whole blood was observed at 30 minutes post dose (first time point analyzed), suggesting a rapid absorption from the IM dose. Blood to plasma ratios of less than 1 suggested that dose-related material was not particularly associated with the blood cells at any time post dose. Exposure of plasma to dose-related material was greater than that of whole blood, as measured by $\text{AUC}_{0\text{-}\infty}$, and the rate of clearance was slower (as measured by $t_{1/2}$). The systemic exposure ($\text{AUC}_{0\text{-}\infty}$) following the IM administration was relatively good with an estimation of relative bioavailability of total radioactivity of 49.7% and 83.5% for blood and plasma, respectively.

[0079] For Group 3, the highest mean concentration of radiolabelled material in plasma and whole blood was observed at 30 minutes post dose (the first time point analyzed). For the early time points (0 to 24 hours), concentrations in plasma were always higher than those in blood, as reflected by blood-to-plasma ratios of less than 1. This indicated that dose-related material was not particularly associated with the blood cells at these time points. After 24 hours, the concentrations of radioactivity in blood were always higher than those in plasma suggesting that the dose-related material was associated with the blood cells. Exposure of plasma to dose-related material was similar to whole blood, as measured by $\text{AUC}_{0\text{-}\infty}$, but the rate of clearance was faster as measured by $t_{1/2}$.

[0080] The major route of excretion following an IM dose or an IV bolus dose was via urine, with a smaller percentage recovered in feces. The recoveries in excreta following the intramuscular and intravenous doses were very similar, at approximately 52.5% to 54.0% for urine and at approximately 31.3% to 35.5% for feces. For both dose routes, excretion was relatively rapid with the majority of the dose administered excreted within 24 hours. The proportion of the administered radiolabelled material recovered in feces suggested that biliary excretion of dose-related material had occurred following both dose routes. Excretion recovery was approximate 92% and 93% for Groups 2 and 3, respectively, indicating that

excretion was essentially complete by 168 hours post dose. A small percentage of the administered radiolabelled material, for both routes, was found in the remaining carcass. Thus the overall mean mass balance of radioactivity was good, at approximately 93% to 94% of the administered dose for both Groups 2 and 3 animals.

[0081] In conclusion, male mice were administered an IM dose (200 mg/kg) or an IV bolus dose (50 mg/kg) of ^{14}C -Genistein. Concentrations of radioactivity in whole blood, plasma, excreta and carcass were determined. The highest radioactivity concentrations were observed at 30 minutes post intramuscular or intravenous bolus dose, indicating rapid absorption from the intramuscular dose. The bioavailability of dose-related material following the intramuscular dose was good, at greater than 49%. Radioactivity was excreted rapidly and urine was the major route of excretion for both dose routes. The high level of radioactivity recovered in feces following the intramuscular or intravenous bolus dose suggested that biliary excretion had occurred. The total recovery of dose-related material following both dose routes was essentially complete by 168 hours post dose.

Example 9 - Pharmacokinetics in Beagle Dogs Following Intravenous or Intramuscular Injection of ^{14}C -Genistein

[0082] Male Beagle dogs were administered a single IV bolus dose (Group 1, nominal 20 mg/kg) or a single IM dose (Group 2, nominal 20 mg/kg) of ^{14}C -Genistein (results shown in Table 11). The Genistein suspension formulation used was prepared as described in Example 8. Following dosing, the content and concentration of radioactivity in blood, plasma and excreta, and the non-compartmental pharmacokinetics of total radioactivity in whole blood and plasma were determined. The concentration of radioactivity in both dose formulations was measured pre and post dose by liquid scintillation spectroscopy and was similar on both occasions. The radiochemical stability of the test article in both dose formulations was assessed prior to and following the administration of the dose. The mean stability values for pre and post dose samples from the intravenous dose formulation (Group 1) were 100% and 99.6%, respectively. The mean stability values for pre and post dose samples from the intramuscular dose formulation (Group 2) were 99.2% and 98.9%, respectively. Therefore, the ^{14}C -Genistein in both formulations was considered to have been radiochemically stable throughout the dosing period.

[0083] Whole-blood samples were collected and plasma was obtained by centrifugation. The concentration of radioactivity in whole-blood and plasma was measured by liquid scintillation spectroscopy. Pharmacokinetic parameters were calculated from the concentration vs. time profiles and are presented in Table 11.

Table 11

Pharmacokinetic		Group 1 (IV)		Group 2 (IM)	
Parameter	Units	Blood	Plasma	Blood	Plasma
t_{\max}	h	0	0	2	2
C_{\max}	$\mu\text{g eq/mL}$	40.4	94.4	5.39	10.9
t_{last}	h	120	120	168	168
$\text{AUC}_{0-\text{tlast}}$	$\mu\text{g eq}\cdot\text{h/mL}$	117	201	108	193
k_{e1}	h^{-1}	0.0130	0.0138	a	0.0092
$t_{1/2}$	h	55.7	227	a	75.2
$\text{AUC}_{0-\text{inf}}$	$\mu\text{g eq}\cdot\text{h/mL}$	122	227	a	228
Extrapolation	%	16.8	11.3	a	15.1
V_z	mL/kg	13168	6511	NA	NA
CL	mL/h/kg	163	88.7	NA	NA
Bioavailability	%	NA	NA	92.9	96.1

^aextrapolation to $\text{AUC}_{0-\text{inf}}$ greater than 20%, therefore not reported.

NA not applicable

[0084] For Group 1 (IV bolus dose), the highest mean concentration of radiolabelled material in plasma and whole blood was observed at 15 minutes post dose (the first time point analyzed). Concentrations in plasma were always higher than those in blood, as reflected by blood to plasma ratios of less than 1. This indicated that dose-related material was not particularly associated with the blood cells at these time points. Exposure of plasma to dose-related material was greater than that of whole blood, as measured by $\text{AUC}_{0-\text{inf}}$, but the rate of clearance was similar as measured by $t_{1/2}$.

[0085] For Group 2 (IM dose), the highest mean concentration of radioactivity in plasma and whole blood was observed at 2 hours post dose, suggesting a relatively rapid absorption from the intramuscular dose. Blood to plasma ratios of less than 1 suggested that dose-related material was not particularly associated with the blood cells at any time post

dose. Exposure of plasma to dose-related material was greater than that of whole blood, as measured by $AUC_{0-t_{last}}$. The rate of clearance (as measured by $t_{1/2}$) was generally slower than the one observed following the IV bolus dose. However, the systemic exposure ($AUC_{0-t_{last}}$) following the IM administration was good with an estimation of relative bioavailability of total radioactivity of 92.9% and 96.1% for blood and plasma, respectively.

[0086] The major route of excretion following an intravenous bolus dose or an IM dose was via feces, with a smaller percentage recovered in urine. The recoveries in excreta following the IV bolus and IM doses were very similar, at approximately 48.7 to 51.9% for feces and at approximately 32.0 to 33.6% for urine. For both dose routes, excretion was relatively rapid with the majority of the dose administered excreted within 48 hours. The proportion of the administered radiolabelled material recovered in feces suggested that biliary excretion of dose-related material had occurred following both dose routes. Excretion recovery by 168 hours post dose was approximately 87.9% and 85.8% for Groups 1 and 2, respectively. Thus, the overall mean excretion mass balance of radioactivity for both groups was good, at approximately 86-88% of the administered dose.

[0087] In conclusion, male dogs were administered an IV bolus dose (20 mg/kg) or an IM dose (20 mg/kg) of ^{14}C -Genistein. Concentrations of radioactivity in whole blood, plasma and excreta were determined. Clinical signs were observed in both groups and were considered to be dose-related. The highest radioactivity concentrations in blood and plasma were observed at 15 minutes (intravenous dose) or 2 hours (intramuscular dose) post dose, indicating relatively rapid absorption from the IM dose. The bioavailability of dose-related material following the IM dose was good, at greater than 92%. Test article-related material was excreted rapidly and feces was the major route of excretion for both dose routes. The high level of radioactivity recovered in feces following the IV bolus or intramuscular dose suggested that biliary excretion had occurred. The excretion mass balance for both dose routes was considered good at greater than 85%.

Example 10 - Oral Pharmacokinetic Comparison Study

[0088] An oral bioavailability comparison of genistein solution formulation in PEG 400 vs. a genistein nanosuspension prepare according to the present description was carried out. The genistein suspension formulation was prepared as described in Example 4. Considering the limited oral bioavailability of genistein in earlier preclinical and clinical work, this experiment was designed to compare a previously used genistein solution formulation prepared with PEG 400 as the vehicle with a genistein nanosuspension formulation prepared as described herein.

[0089] Ten groups of seven mice were prepared at each time point (70 mice) for each of the two formulations (total mice = 140). A single dose of 400 mg/kg genistein was given by oral gavage and then blood was collected at 10 subsequent time points. Time points for blood collection were the following: 0, 0.5, 1, 2, 3, 4, 6, 8, 10, and 12 hours post administration.

[0090] The oral bioavailability with the nanoparticulate suspension was statistically significantly greater than that achieved in the PEG 400 solution formulation. Both free and total genistein levels were determined in each of the groups for every time point. The mean concentration of seven mice was determined for each time point and reported. At the time 0 time point, the free and total genistein concentration was below the limit of quantitation and not reported. For the mice receiving the nanoparticulate suspension, the free genistein concentrations were significantly greater at 1, 2, 4, 8, 10 and 12 hours when compared to the concentration achieved in the mice receiving the PEG 400 solution formulation. The total genistein concentrations were also significantly greater with the nanoparticulate suspension at 0.5, 1, 2, 4, 8, 10 and 12 hours when compared to the PEG 400 solution formulation. As noted in Table 12 and Table 13, and as shown in FIG. 4 and FIG. 5, the absorption and excretion curve for the nanoparticulate suspension is also much more predictable and less erratic than that achieved by the PEG 400 formulation for both the free and total genistein concentrations. Free genistein determinations after a single oral dose of 400 mg/kg for two different genistein formulations in mice.

Table 12

Mean Free Genistein Concentrations at Each Sampling Time point						
Time point	Free (ng/ml) PEG 400 (ng/ml)	Free Nanosuspension (ng/ml)	Difference	Percent	p value	
0.5 hr	1137.9	1598.6	460.7	40.5	0.062	
1 hr	514.0	999.7	485.7	94.5	0.006	
2 hr	255.1	746.4	491.3	192.6	0.000	
3 hr	403.1	634.6	231.4	57.4	0.079	
4 hr	207.6	484.3	276.7	133.2	0.002	
6 hr	361.9	313.6	-48.4	-13.4	0.791	
8 hr	436.3	172.3	-264.0	-60.5	0.023	
10 hr	239.0	59.0	-180.0	-75.3	0.000	
12 hr	113.7	28.3	-85.4	-75.1	0.034	

Table 13
Mean Total Genistein Concentrations at Each Sampling Time point

Time point	Total PEG 400 (ng/ml)	Total Nanosuspension (ng/ml)	Difference	Percent	p value
0.5 hr	15640.0	32114.3	16474.3	105.3	0.000
1 hr	7711.4	20974.3	13262.9	172.0	0.002
2 hr	4151.4	12050.0	7898.6	190.3	0.033
3 hr	6251.4	9112.9	2861.5	45.8	0.136
4 hr	2970.1	7692.9	4722.8	159.0	0.003
6 hr	6215.1	4004.3	-2210.8	-35.6	0.257
8 hr	5558.6	1982.9	-3575.7	-64.3	0.006
10 hr	3480.0	755.4	-2724.6	-78.3	0.001
12 hr	2205.0	380.3	-1824.7	-82.8	0.002

Example 11- Oral Bioavailability Comparison of Non-nanoparticulate Genistein Suspension Formulation and a Nano-particulate Genistein Formulation

[0091] In a previous experiment we were able to demonstrate improved oral bioavailability with orally administered Genistein-IS suspension formulation in comparison with a formulation of genistein dissolved in PEG-400. This experiment compared the oral bioavailability of an aqueous genistein suspension formulation that included non-nanoparticulate genistein material with that provided by the Genistein-IS formulation prepared as described in Example 4. The non-nanoparticulate genistein suspension formulation was the same as the Genistein-IS formulation, except that the genistein material exhibited a volume average particle size of 8 μm , instead of the volume average particle size of 0.13 μm exhibited by the Genistein-IS suspension formulation.

[0092] Ten groups of seven mice were prepared at each timepoint (70 mice) for each of the two formulations (total mice = 140). A single dose of 400 mg/kg was given by oral gavage and then blood was collected at 10 subsequent timepoints. Time points for blood collection were the following: 0, 0.5, 1, 2, 4, 6, 8, 10, 12 and 24 hours post administration.

[0093] Both free and total genistein levels were determined in each of the groups for every timepoint. The mean concentration of seven mice was determined for each timepoint and reported. At the time 0 timepoint, the free and total genistein concentration was below the limit of quantitation and not reported. For the Genistein-IS suspension formulation mice, the free genistein concentrations were significantly greater at 0.5, 1 & 2 hours when compared to those achieved by the non-nanoparticulate genistein formulation (See, Table 14). The total genistein concentrations were also significantly greater with the with the Genistein-IS suspension formulation at 0.5, 1 and 2 hours when compared to those achieved by the non-nanoparticulate genistein formulation (See, Table 15). As noted in FIG. 6 and FIG. 7, the absorption and excretion curve for the Genistein-IS suspension formulation is also much more predictable and less erratic than that achieved by the non-nanoparticulate genistein formulation for both the free and total genistein concentrations.

Table 14

Mean Free Genistein Concentrations at Each Sampling Timepoint					
Time point	Free Conc. 8 Micron l	Free Conc. Nanosuspension (ng/ml)	Difference	Percent	p value
0.5 hr	630.7	1494.0	863.3	136.9	0.000
1 hr	512.1	853.4	341.3	66.6	0.003
2 hr	314.4	652.6	338.1	107.5	0.003
4 hr	214.3	367.4	153.1	71.5	0.055
6 hr	136.3	168.0	31.7	23.2	0.460
8 hr	140.1	45.8	-94.3	-67.3	0.006
10 hr	50.3	21.6	-28.6	-57.0	0.284
12 hr	47.3	23.9	-23.3	-49.4	0.195
24 hr	0.0	0.0	0.0	-	-

Table 15

Mean Total Genistein Concentrations at Each Sampling Timepoint					
Timepoint	Total Conc. 8 Micron (ng/ml)	Total Conc. Nanosuspension (ng/ml)	Difference (ng/ml)	Percent	p value
0.5 hr	9561.4	32385.7	22824.3	238.7	0.000
1 hr	9762.9	25842.9	16080.0	164.7	0.000
2 hr	5247.1	15357.1	10110.0	192.7	0.007
4 hr	3230.0	5938.6	2708.6	83.9	0.055
6 hr	2230.4	2423.1	192.7	8.6	0.802
8 hr	1902.9	1007.1	-895.7	-47.1	0.070
10 hr	1002.2	426.4	-575.8	-57.5	0.253
12 hr	876.4	338.1	-538.3	-61.4	0.121
24 hr	53.5	57.7	4.3	8.0	0.715

[0094] It will be obvious to those having skill in the art that many changes may be made to the details of the above-described embodiments without departing from the underlying principles of the invention. The scope of the present invention should, therefore, be determined only by the following claims.

Claims

1. A genistein formulation for use as a medicament by oral and/or parenteral administration, the formulation comprising:
 - nanoparticulate genistein; and
 - one or more pharmaceutically acceptable excipients forming a suspension medium, wherein the one or more pharmaceutically acceptable excipients includes a polyvinylpyrrolidone (PVP) and a nonionic surfactant, wherein the nanoparticulate genistein exhibits a D (0.50) of 0.5 μm or less and no particles measure greater than 2 μm , and the one or more pharmaceutically acceptable excipients are selected such that the nanoparticulate genistein is suspended within the suspension medium.
2. A formulation for use according to claim 1, wherein the nanoparticulate genistein exhibits a D (0.50) of 0.2 μm or less.
3. A formulation for use according to claim 2, wherein the nanoparticulate genistein exhibits a D (0.50) of 0.2 μm or less and a D (0.90) of 0.5 μm or less.
4. A formulation for use according to any preceding claim, wherein the one or more pharmaceutically acceptable excipients further includes at least one of a diluent and a buffer.
5. A formulation for use according to any preceding claim, wherein the nonionic surfactant is present in an amount ranging from about 0.01% to about 10% (w/w).
6. A formulation for use according to any preceding claim, wherein the PVP is povidone K17.
7. A formulation for use according to any preceding claim, wherein the PVP is present in an amount ranging from about 0.5% to about 15% (w/w).
8. A formulation for use according to any preceding claim, wherein the amount of nonionic surfactant is selected from about 0.01% to about 10% (w/w), about 0.1% to about 10% (w/w), about 0.2% to about 5% (w/w), about 0.2% to about 1% (w/w), , about 0.2% to about 0.6% (w/w), and about 0.2% to about 0.8% (w/w), and the amount of water soluble polymer is selected from about 0.5% to about 15% (w/w), about 1% to about 10% (w/w), about 5% to about 15% (w/w), about 10% to about 15% (w/w), about 12% to about 15% (w/w), about 1% to about 8% (w/w), and about 1% to about 5% (w/w).

9. A formulation for use according to any preceding claim, wherein the nonionic surfactant is at least one of polysorbate 80 and polysorbate 20.
10. A formulation for use according to any preceding claim, wherein the genistein is present in the formulation at a concentration of 250 mg/ml or greater.
11. A formulation for use according to any preceding claim, wherein the genistein is present in the formulation at a concentration selected from a concentration of between about 250 mg/ml and about 500 mg/ml, a concentration of between about 200 mg/ml and about 400 mg/ml, a concentration of between about 250 mg/ml and about 350 mg/ml, and a concentration of between about 275 mg/ml and about 325 mg/ml.
12. A formulation for use according to any preceding claim, wherein the amount of nanoparticulate genistein is at a concentration of at least 300 mg/ml.
13. A formulation for use according to any of claims 4 to 11, wherein the formulation exhibits a pH ranging from about 2 to about 12, preferably about 4 to about 8, more preferably about 5 to about 7.
14. A formulation according to any of claims 1 to 13 for use in the treatment of acute radiation syndrome.
15. A formulation according to any of claims 1 to 13 for use according to claim 14, wherein the treatment is prophylactic and the formulation is to be administered to a subject at risk of radiation exposure.
16. A formulation according to any of claims 1 to 13 for use in the treatment of a subject that has undergone exposure to radiation as part of a therapeutic or diagnostic procedure.
17. A formulation according to any of claims 1 to 13 for use according to claim 16, wherein treatment with the formulation commences prior to the subject undergoing the therapeutic or diagnostic procedure.

Patentansprüche

1. Genistein-Formulierung für die Verwendung als Medikament durch orale und/oder parenterale Verabreichung, wobei die Formulierung Folgendes umfasst:
 - nanopartikuläres Genistein; und
 - einen oder mehrere pharmazeutisch verträgliche Hilfsstoffe, die ein Suspensionsmedium bilden, wobei der eine oder die mehreren pharmazeutisch verträglichen Hilfsstoffe ein Polyvinylpyrrolidon (PVP) und ein nichtionisches Tensid einschließen,
 - wobei das nanopartikuläre Genistein einen D (0,50) von 0,5 µm oder weniger zeigt und keine Partikel mehr als 2 µm bemessen, und der eine oder die mehreren pharmazeutisch verträglichen Hilfsstoffe derart ausgewählt sind, dass das nanopartikuläre Genistein innerhalb des Suspensionsmediums suspendiert ist.
2. Formulierung für die Verwendung nach Anspruch 1, wobei das nanopartikuläre Genistein einen D (0,50) von 0,2 µm oder weniger zeigt.
3. Formulierung für die Verwendung nach Anspruch 2, wobei das nanopartikuläre Genistein einen D (0,50) von 0,2 µm oder weniger und einen D (0,90) von 0,5 µm oder weniger zeigt.
4. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei der eine oder die mehreren pharmazeutisch verträglichen Hilfsstoffe weiter mindestens eines von einem Verdünnungsmittel und einem Puffer einschließen.
5. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das nichtionische Tensid in einer Menge im Bereich von etwa 0,01 % bis etwa 10 % (m/m) vorliegt.
6. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das PVP Povidon K17 ist.
7. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das PVP in einer Menge im Bereich

von etwa 0,5 % bis etwa 15 % (m/m) vorliegt.

8. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei die Menge des nichtionischen Tensids aus Folgenden ausgewählt ist: von etwa 0,01 % bis etwa 10 % (m/m), etwa 0,1 % bis etwa 10 % (m/m), etwa 0,2 % bis etwa 5 % (m/m), etwa 0,2 % bis etwa 1 % (m/m), etwa 0,2 % bis etwa 0,6 % (m/m) und etwa 0,2 % bis etwa 0,8 % (m/m), und die Menge an wasserlöslichem Polymer aus Folgenden ausgewählt ist: von etwa 0,5 % bis etwa 15 % (m/m), etwa 1 % bis etwa 10 % (m/m), etwa 5 % bis etwa 15 % (m/m), etwa 10 % bis etwa 15 % (m/m), etwa 12 % bis etwa 15 % (m/m), etwa 1 % bis etwa 8 % (m/m) und etwa 1 % bis etwa 5 % (m/m).
9. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das nichtionische Tensid mindestens eines von Polysorbat 80 und Polysorbat 20 ist.
10. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das Genistein in der Formulierung in einer Konzentration von 250 mg/ml oder mehr vorliegt.
11. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei das Genistein in der Formulierung in einer Konzentration vorliegt, die aus Folgenden ausgewählt ist: von einer Konzentration zwischen etwa 250 mg/ml und etwa 500 mg/ml, einer Konzentration zwischen etwa 200 mg/ml und etwa 400 mg/ml, einer Konzentration zwischen etwa 250 mg/ml und etwa 350 mg/ml und einer Konzentration zwischen etwa 275 mg/ml und etwa 325 mg/ml.
12. Formulierung für die Verwendung nach einem vorstehenden Anspruch, wobei die Menge an nanopartikulärem Genistein in einer Konzentration von mindestens 300 mg/ml vorliegt.
13. Formulierung für die Verwendung nach einem der Ansprüche 4 bis 11, wobei die Formulierung einen pH im Bereich von etwa 2 bis etwa 12, bevorzugt etwa 4 bis etwa 8, bevorzugter etwa 5 bis etwa 7 zeigt.
14. Formulierung nach einem der Ansprüche 1 bis 13 für die Verwendung bei der Behandlung von akutem Strahlensyndrom.
15. Formulierung nach einem der Ansprüche 1 bis 13 für die Verwendung nach Anspruch 14, wobei die Behandlung prophylaktisch ist und die Formulierung an ein Subjekt zu verabreichen ist, bei dem ein Risiko von Strahlenexposition besteht.
16. Formulierung nach einem der Ansprüche 1 bis 13 für die Verwendung bei der Behandlung eines Subjekts, das Strahlung als Teil eines therapeutischen oder diagnostischen Verfahrens ausgesetzt wurde.
17. Formulierung nach einem der Ansprüche 1 bis 13 für die Verwendung nach Anspruch 16, wobei eine Behandlung mit der Formulierung beginnt, bevor sich das Subjekt dem therapeutischen oder diagnostischen Verfahren unterzieht.


Revendications

1. Formulation de génistéine pour une utilisation comme médicament par administration orale et/ou parentérale, la formulation comprenant :
 de la génistéine nanoparticulaire ; et
 un ou plusieurs excipients pharmaceutiquement acceptables formant un milieu de suspension, où les un ou plusieurs excipients pharmaceutiquement acceptables comportent une polyvinylpyrrolidone (PVP) et un agent tensioactif non ionique,
 dans laquelle la génistéine nanoparticulaire présente une D (0,50) de 0,5 µm ou moins, et aucune particule ne mesure plus de 2 µm, et les un ou plusieurs excipients pharmaceutiquement acceptables sont choisis de façon à ce que la génistéine nanoparticulaire soit en suspension au sein du milieu de suspension.
2. Formulation pour une utilisation selon la revendication 1, dans laquelle la génistéine nanoparticulaire présente une D (0,50) de 0,2 µm ou moins.
3. Formulation pour une utilisation selon la revendication 2, dans laquelle la génistéine nanoparticulaire présente une

EP 2 640 188 B1

D (0,50) de 0,2 μm ou moins et une D (0,90) de 0,5 μm ou moins.

4. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle les un ou plusieurs excipients pharmaceutiquement acceptables comportent en outre au moins l'un parmi un diluent et un tampon.
5. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle l'agent tensioactif non ionique est présent selon une quantité allant d'environ 0,01% à environ 10% (p/p).
6. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la PVP est la povidone K17.
7. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la PVP est présente selon une quantité allant d'environ 0,5% à environ 15% (p/p).
8. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la quantité d'agent tensioactif non ionique est choisie d'environ 0,01% à environ 10% (p/p), d'environ 0,1% à environ 10% (p/p), d'environ 0,2% à environ 5% (p/p), d'environ 0,2% à environ 1% (p/p), d'environ 0,2% à environ 0,6% (p/p), et d'environ 0,2% à environ 0,8% (p/p), et la quantité de polymère hydrosoluble est choisie d'environ 0,5% à environ 15% (p/p), d'environ 1% à environ 10% (p/p), d'environ 5% à environ 15% (p/p), d'environ 10% à environ 15% (p/p), d'environ 12% à environ 15% (p/p), d'environ 1% à environ 8% (p/p), et d'environ 1% à environ 5% (p/p).
9. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle l'agent tensioactif non ionique est au moins l'un parmi le polysorbate 80 et le polysorbate 20.
10. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la génistéine est présente dans la formulation selon une concentration de 250 mg/ml ou plus.
11. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la génistéine est présente dans la formulation selon une concentration choisie parmi une concentration comprise entre environ 250 mg/ml et environ 500 mg/ml, une concentration comprise entre environ 200 mg/ml et environ 400 mg/ml, une concentration comprise entre environ 250 mg/ml et environ 350 mg/ml, et une concentration comprise entre environ 275 mg/ml et environ 325 mg/ml.
12. Formulation pour une utilisation selon l'une quelconque des revendications précédentes, dans laquelle la quantité de génistéine nanoparticulaire se trouve selon une concentration d'au moins 300 mg/ml.
13. Formulation pour une utilisation selon l'une quelconque des revendications 4 à 11, la formulation présentant un pH dans la plage allant d'environ 2 à environ 12, préférablement d'environ 4 à environ 8, plus préférablement d'environ 5 à environ 7.
14. Formulation selon l'une quelconque des revendications 1 à 13, pour une utilisation dans le traitement du syndrome d'irradiation aiguë.
15. Formulation selon l'une quelconque des revendications 1 à 13, pour une utilisation selon la revendication 14, le traitement étant prophylactique et la formulation étant prévue pour une administration à un sujet risquant d'être exposé à des rayonnements.
16. Formulation selon l'une quelconque des revendications 1 à 13, pour une utilisation dans le traitement d'un sujet ayant subi une exposition à des rayonnements dans le cadre d'une procédure thérapeutique ou de diagnostic.
17. Formulation selon l'une quelconque des revendications 1 à 13, pour une utilisation selon la revendication 16, le traitement par la formulation débutant avant que le sujet ne subisse la procédure thérapeutique ou de diagnostic.



*p < 0.05 two-tailed Fisher Exact Test (vehicle vs. genistein)

FIG. 1





FIG. 2



*p < 0.05 two-tailed Fisher Exact Test (vehicle vs. genistein)

FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- Remington's Pharmaceutical Sciences. Maack Publishing Co, 1985 [0020]
- **TAKIMOTO et al.** *Cancer Epidemiol Biomarkers Prev.*, November 2003, vol. 12 (11), 1213-21 [0038]
- **WEI et al.** *J Nutr.*, November 2003, vol. 133 (11), 3811 S-3819S [0038]
- **SAKAR, F. H. ; Y. LI.** *Cancer Invest.*, 2003, vol. 21 (5), 744-57 [0038]
- **MAGEE P. J. ; I. R. ROLAND.** *Br J Nutr.*, April 2004, vol. 91 (4), 513-31 [0038]
- **PARK, O. J. ; Y. J. SURH.** *Toxicol Lett.*, 15 April 2004, vol. 150 (1), 43-56 [0038]
- **MESSINA, M. J.** *Nutr Re.*, April 2003, vol. 61 (4), 117-31 [0038]
- **MOHAMMAD et al.** *Mol Cancer Ther.*, December 2003, vol. 2 (12), 1361-8 [0038]
- **NICOSIA et al.** *Hematol Oncol Clin North Am.*, August 2003, vol. 17 (4), 927-43 [0038]
- **SUN et al.** *Nutr Cancer*, 2001, vol. 39 (1), 85-95 [0038]
- **MORABITO et al.** *J Bone Miner Res.*, October 2002, vol. 17 (10), 1904-12 [0040]
- **COTTER A. ; K. D. CASHMAN.** *Nutr Rev.*, October 2003, vol. 61 (10), 346-51 [0040]
- **ATKINSON et al.** *Am J Clin Nutr.*, February 2004, vol. 79 (2), 326-33 [0040]
- **SETCHELL K. D. ; E. LYDEKING-OLSEN.** *Am J Clin Nutr.*, September 2003, vol. 78, 593S-609S [0040]
- **CLIFTON-BLIGH et al.** *Menopause*, July 2001, vol. 8 (4), 259-65 [0040]
- **ROOMANS, G. M.** *Am J Respir Med.*, 2003, vol. 2 (5), 413-31 [0041]
- **GOODMAN-GRUEN, D. ; D. KRITZ-SILVERSTEIN.** *Menopause*, September 2003, vol. 10 (5), 427-32 [0043]
- **BHATHENA, S. J. ; M. T. VELASQUEZ.** *Am J Clin Nutr.*, December 2002, vol. 76 (6), 1191-201 [0043]
- **URBAN et al.** *J Urol.*, January 2001, vol. 165 (1), 294-300 [0043]
- **YOUNG, S. G. ; S. PARTHASARATHY.** *West J Med.*, February 1994, vol. 160 (2), 153-54 [0043]