

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 June 2007 (14.06.2007)

PCT

(10) International Publication Number
WO 2007/066248 A2

(51) International Patent Classification: **Not classified**

(21) International Application Number:
PCT/IB2006/054275

(22) International Filing Date:
15 November 2006 (15.11.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/742,292 5 December 2005 (05.12.2005) US

(71) Applicant (for all designated States except US): **KONINKLIJKE PHILIPS ELECTRONICS, N.V.** [NL/NL]; Groenewoudseweg 1, NL-5621 BA Eindhoven (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHOU, Bin** [CN/US]; 1399 Meadow Ridge, San Jose, California 95131 (US). **FANG, Yaqiong** [CN/US]; 167 Evening Star Court, Milpitas, California 95035 (US). **SIMMS, Daniel** [US/US]; 250 North Frances Street, Sunnyvale, California

94086 (US). **HAQUE, Imtiyaz** [US/US]; 11531 Bianchini Lane, Cupertino, California 95014 (US).


(74) Common Representative: **KONINKLIJKE PHILIPS ELECTRONICS, N.V.**; C/o Mcknight, Douglas, B. 595 Miner Road, Cleveland, Ohio 44143 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

[Continued on next page]

(54) Title: GESTION DE MISE A JOUR DE PLAN DE SOINS



(57) Abstract: La présente invention concerne un système de gestion de plan de soins (10) qui comprend un serveur (12), lequel comprend une base de données de contenu (14) contenant un contenu relatif à la gestion de soins, laquelle comprend de multiples révisions d'au moins quelques-uns de ses contenus. Chaque révision de contenu est étiquetée avec une information de révision de contenu temporelle. Un stockage de plan de soins (18) stocke une pluralité de plans de soins de patients (21, 22, 23) associés à des patients correspondants. Chaque plan de soins de patients comprend une sélection de contenus relatifs à une gestion de soins à présenter au patient et une politique de révision indiquant quelle révision temporelle de ce contenu doit être présentée. Des interfaces utilisateurs (31, 32, 33) sont connectées de manière opérationnelle avec une interface de communication (28) du serveur (12) de façon à recevoir et présenter le contenu relatif à gestion de soins sélectionné.



FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

Declaration under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

CARE PLAN UPDATE MANAGEMENT

DESCRIPTION

The following relates to the health management arts. It finds particular application in conjunction with out-patient management of chronic illnesses such as congestive heart failure, emphysema, chronic obstructive pulmonary disease (COPD), and so forth, and will be described with particular reference thereto. It finds application more generally in conjunction with methods and apparatuses for providing extended-term health management for: chronic diseases; rehabilitation from a catastrophic event such as a stroke or an automobile accident; managing weight; controlling insomnia; redressing health-impacting lifestyle issues such as smoking or poor diet or inadequate physical exercise; avoiding potential medical conditions such as osteoporosis or tooth decay; and so forth.

Medical professionals recognize that providing extended-term health care management assistance to chronically ill patients is an important aspect of treating the chronic illness and assuring the patient a high quality of life. Extended term health management is typically performed on an out-patient basis, and is typically wholly or in large part self-administered, with occasional help from weekly therapy classes or so forth. It is well known, however, that patients often fail to adequately follow the prescribed health care plan outside of a hospital or other medical setting. This failure can result from lack of understanding of how to perform health care activities, apathy or lack of motivation due to depression, fear of failure, or so forth.

Such problems can in principle be overcome by increased one-on-one interaction between the patient and medical personnel. For example, a daily visit to the patient by a traveling nurse could help ensure that the patient is taking medications in a timely fashion and following prescribed dietary and exercise regimens. However, it is often not feasible to provide such intensive one-on-one sessions due to high cost, lack of available medical personnel, or so forth.

In some cases, the patient can choose to access a hospital website or other on-line (e.g., Internet-based) medical database to pull information relevant to the patient's care plan. However, the patient may not have Internet access, or may be unable to navigate a complex on-line medical database. Moreover, providing access to on-line databases does

nothing to help patients who are unmotivated. Other approaches that have been used include providing the patient with instructional or motivational videos. However, these approaches do not provide interactive assistance of a type likely to encourage the patient to follow care plan regimens. Moreover, passive videos are difficult to personalize so as to
5 directly address specific issues related to the patient.

Royal Philips Electronics, Cardiovascular Associates of the Delaware Valley, and Comcast Corporation have announced a cooperative effort called Motiva™ to provide a test group of chronic heart failure patients with a remote patient management broadband-enabled platform for connecting the test patients with their healthcare
10 community. The Motiva™ system provides a cable television-based interactive health care management platform, in which content such as educational video, medication scheduling, personalized encouragement and reinforcement, and so forth, is pushed to the patient based on a personalized health care plan. Feedback from the patient, for example through the use of interactive surveys, enables the Motiva™ system to adjust or personalize content to the
15 needs of each patient. The Motiva™ system can deliver personalized health care management assistance to patients on a daily or more frequent basis.

One problem that arises in maintaining such a personalized interactive care management system is timely updating of content while continuously providing personalized care management assistance to large numbers of patients. As medical
20 knowledge and state-of-the-art progresses, the content deliverable by such a system can become outdated. Accordingly, it is useful to update content, or portions of content, to reflect new knowledge, new medical technology, new information, or so forth. However, medical professionals often have differing opinions as to the advisability of following new (and possibly unproven) medical procedures or recommendations, and some doctors may
25 not want certain updated content to be delivered to their patients. Also, on an individual patient basis certain updated content may be inappropriate.

The following contemplates improvements that overcome the aforementioned limitations and others.

30 According to one aspect, a server is disclosed for a care management system. A content database contains care management-related content including multiple revisions of

at least some said content. Each content revision is tagged with temporal content revision information. A care plans storage stores a plurality of patient care plans associated with corresponding patients. Each patient care plan includes: (i) a selection of care management-related content to be presented to the corresponding patient and (ii) a revision policy indicating which temporal revision of said content should be presented to the corresponding patient. At least one communication interface is included for communicating selected care management-related content to patients. The communicated content for each patient is selected based on the corresponding patient care plan, tagged temporal content revision information, and the revision policy of the corresponding patient care plan.

According to another aspect, a care plan management system is disclosed, including a server as set forth in the preceding paragraph, and a user interface operatively connected with the at least one communication interface of the server to receive and present the selected care management-related content.

According to another aspect, a method is disclosed for managing care. Care management-related content is stored, including multiple revisions of at least some said content. Each content revision is tagged with temporal content revision information. A plurality of patient care plans associated with corresponding patients are stored. Each patient care plan includes: (i) a selection of care management-related content to be presented to the corresponding patient and (ii) a revision policy indicating which temporal revision of said content should be presented to the corresponding patient. Selected care management-related content are communicated to patients. The communicated content for each patient is selected based on the corresponding patient care plan, tagged temporal content revision information, and the revision policy of the corresponding patient care plan.

According to another aspect, a computer readable medium or media such as an optical disk, magnetic disk, magnetic tape, random access memory (RAM), read-only memory (ROM), FLASH memory, or so forth, stores a program for managing care. The program when executed by a computer, digital processor, network server, or so forth, or by a combination thereof, performs process operations comprising communicating selected care management related content to a patient. The communicated content for each patient

is selected based on a patient care plan, temporal content revision information tagged to the selected care management related content, and a revision policy of the patient care plan.

One advantage resides in balancing currency of a personalized interactive care management assistance system with the flexibility to continue delivering older content to
5 certain patients as appropriate.

Another advantage resides in enabling updating of a personalized interactive care management assistance system substantially simultaneously with continuing delivery of personalized content to patients.

Another advantage resides in providing medical personnel the option to provide, or
10 not provide, revised medical content to their patients.

Numerous additional advantages and benefits will become apparent to those of ordinary skill in the art upon reading the following detailed description of the preferred embodiments.

15 The invention may take form in various components and arrangements of components, and in various process operations and arrangements of process operations. The drawings are only for the purpose of illustrating preferred embodiments and are not to be construed as limiting the invention.

FIGURE 1 diagrammatically shows principal components of a personalized
20 interactive care management assistance system.

FIGURE 2 diagrammatically shows the content modules database of the personalized interactive care management assistance system of FIGURE 1.

FIGURE 3 diagrammatically shows an example post-cardiac event care plan suitable for assignment to overweight patients who are smokers and who have suffered a
25 traumatic cardiac event such as a heart attack.

FIGURE 4 diagrammatically shows the post-cardiac event care plan assigned to Patient #1.

FIGURE 5 diagrammatically shows the post-cardiac event care plan assigned to Patient #2.

FIGURE 6 diagrammatically shows a user interface display querying Patient #2 to select a content revision based on displayed semantic information contained in the tagged temporal content revision information.

FIGURE 7 diagrammatically shows the post-cardiac event care plan assigned to Patient #3.

FIGURE 8 diagrammatically shows a five-day period of care plan activity undertaken by Patient #3 in accordance with the post-cardiac event care plan assigned to Patient #3 shown in FIGURE 7.

With reference to FIGURE 1, a personalized interactive care management system **10** includes a server **12** for distributing care management-related content. In the example system **10**, the content is arranged in a content database **14** stored on a common data storage **16**. The server **12** can be embodied in various ways, such as by a centralized computer or computer server, a desktop computer, a distributed array of computers, or so forth. The care management-related content typically includes videos, text, surveys, questionnaires, or so forth directed toward assisting in achieving various health management tasks. For example, care management-related content may be provided that are directed toward: reducing weight; stopping smoking; learning to self-administer a medical intervention such as a medication, a biometric monitor, or so forth; learning to follow a dietary restriction such as a low-salt diet; learning to follow a dietary requirement such as a high-fiber diet; performing a physical exercise; or so forth.

To enable personalized distributing of content, each patient in the system **10** has an associated care plan that in the illustrated embodiment is stored in a care plans partition **18** that is also stored on the common system storage **16**. The example common data storage **16** is logically partitioned to define the content database **16** and the care plan storage. In the illustrated embodiment, the common data storage **16** stores a care plan **21** for Patient #1, a care plan **22** for Patient #2, and a care plan **23** for Patient #3. While only three care plans **21**, **22**, **23** associated with a corresponding three Patients #1, #2, #3 are illustrated, it is contemplated that the server **12** may store care plans for a thousand or more different patients. As used herein, the term "patient" encompasses patients recovering from surgery, stroke, heart failure, or another condition, patients suffering a chronic illness that is being

treated on an out-patient basis, or so forth. As used herein, the term "patient" also encompasses other users of the health management system **10** who may be generally healthy but who are following a health management program assisted by the system **10** to maintain fitness, control weight, avoid osteoporosis, or otherwise maintain a healthy condition or make health-related lifestyle modifications. Moreover, the use of the partitioned common data storage is one example of many suitable storage arrangements. As another example, in other contemplated embodiments separate storage media may be provided for the content database and the patient care plans.

A rules-based processor **26** determines content to be distributed to each patient based on suitable rules contained in or identified by the patient care plan of that patient. Rules suitably determine which content is presented to which patient or patients, the ordering of such content presentation, and so forth. A communication interface **28** of the server **12** communicates selected care management-related content to selected patients at respective user interfaces. For example: the communication interface **28** communicates content intended for Patient #1 to a user interface **31** that is accessible by Patient #1; the communication interface **28** communicates content intended for Patient #2 to a user interface **32** that is accessible by Patient #2; the communication interface **28** communicates content intended for Patient #3 to a user interface **33** that is accessible by Patient #3; and so forth. The user interfaces **31**, **32**, **33** can employ substantially any hardware capable of providing content presentation and capable of providing feedback to the server **12** via the communication interface **28**. For example, the user interfaces **31**, **32**, **33** can be embodied by hardware such as: a desktop computer; a laptop computer; a personal data assistant (PDA); a cellular telephone (i.e., cellphone); a television set having Internet connectivity integrally included and operated by a television remote control or other input device; a digital or analog television set having Internet connectivity provided by an add-on set-top unit and operated by a television remote control, set-top unit remote control, or other input device; or so forth. The communication interface **28** is operatively connected with each of the user interfaces **31**, **32**, **33** by a pathway or pathways **36** such as the Internet, a cable television network, a satellite television network, a cellular telephone network, or so forth. Moreover, the communication interface **28** optionally includes more than one communication interface. For example, it is contemplated for different user interfaces to connect with the communication interface **28** by different pathways each

employing different interface hardware and software. For example, the user interface **31** might be a computer operatively connected with the communication interface **28** by the Internet, while the user interface **32** might be a cellphone connected with the communication interface **28** by a cellular telephone network. To construct such an embodiment, the communication interface **28** suitably includes an Internet port component, and a cellular telephone network port. The pathway or pathways **36** are advantageously secure links because private medical information may be conveyed across the pathway or pathways **36**. However, unsecured pathways can also be used.

To provide an interactive system, the user interfaces **31, 32, 33** are optionally interactive user interfaces whereby the respective Patients #1, #2, #3 can provide input that is communicated to the server **12** via the communication interface **28**. For example, the user interface can be an interactive television set, such as a digital television set or a television set including a set-top box configured to provide interactivity. In other cases, the user interfaces **31, 32, 33** may be a computer, a laptop computer, a personal data assistant (PDA), and a cellular telephone (cellphone), each of which devices provides interactive capability. Feedback provided by the user interfaces **31, 32, 33** may include for example answers to questions posed by the content, or answers to surveys, quizzes, tests, questionnaires, or the like that assess how well the patient or user understood previously presented content. Optionally, the user interface may include one or more biometric feedback monitors each of which measures at least one biometric parameter of a patient that is communicated to the server via the communication interface **28**. Suitable biometric monitors may include, for example: a saturated blood oxygen level (SpO₂) monitor; a heart rate monitor; a blood pressure monitor; a weight scale; an electrocardiograph (ECG); or so forth.

Maintenance of the server **12** is suitably performed by an administrator via an administrator interface **40**. In some embodiments, the administrator interface **40** is suitably a network administrator account having a high level of access to the server **12**. The administrator may, for example, add new care management-related content, delete obsolete or outdated care management-related content, organize content, modify or update content flow rules, or so forth. In some embodiments, medical personnel such as doctors or nurses can directly generate and/or update the patient care plans **21, 22, 23** by directly accessing the server **12** via medical personnel interfaces **41, 42**. Medical personnel are optionally

assigned a lower level of access through a regular user account or other network account providing lower level access limited, for example, to patients of a doctor who is accessing the system **10**. For example, the first medical personnel interface **41** accesses the patient care plans **21, 22** of Patients #1 and #2 who are patients of the doctor employing the first
5 medical personnel interface **41**, while the second medical personnel interface **42** accesses the patient care plan **23** of Patient #3 who is a patient of the doctor employing the second medical personnel interface **42**. In some embodiments, medical personnel interfaces **41, 42** are omitted, and one or more system administrators perform all creation and updating of the patient care plans **21, 22, 23** via the administrator interface **40**, and in accordance with
10 instructions from the patient's physician or other medical personnel.

With reference to FIGURE 2, the care management-related content stored in the content database **16** is in some embodiments organized into self-contained content modules. FIGURE 2 diagrammatically depicts the following example content modules: weight loss module; quit smoking module; exercise module; and high fiber diet module.
15 Each module may be occasionally updated. To enable suitable selection of whether or not to present each user with the updated content, a new module revision is created for each update. Changes in module content justifying creation of a new revision may include, for example, adding new material, adding new components, revising existing material or components to reflect current medical knowledge or updated equipment, or so forth. To
20 enable suitable selection of which content revision to present to each user, each such content revision is tagged with temporal content revision information. For example, the weight loss module has an earliest revision tagged as revision 1.1, a later revision tagged as revision 1.2, and a latest revision tagged as revision 1.3. Additionally, a revision-in-progress is tagged as revision 1.4, and is not yet ready for use. The revision 1.3
25 is optionally marked as the latest revision (the latest revision of each content module is diagrammatically indicated in FIGURE 2 by a large boldfaced asterisk). Alternatively, where the tagged temporal content revision information includes ordering information such as the illustrated revision number tags 1.1, 1.2, 1.3, or as another example an alphabetical revision indicator tagging, the latest content revision can be identified as that content
30 revision whose tagged temporal content revision ordering information indicates the content revision is the latest revision. However, if this latter approach is used, then the

revision-in-progress should be numbered "0.0" or otherwise tagged to ensure that it is not inadvertently identified as the latest revision.

The other content modules illustrated as examples in FIGURE 2 also have multiple content revisions. The quit smoking module includes three content revisions tagged as revisions 6.1, 6.2, and 6.3, with revision 6.3 marked as the latest revision. The exercise module includes two content revisions tagged as revisions 3.6 and 3.7, with revision 3.7 marked as the latest revision. The high fiber diet module includes one content revision tagged as revision 8.3 and marked as the latest revision, and also includes a revision-in-progress tagged as revision 8.4 which is not yet ready for use.

With reference to FIGURE 3, each patient care plan is suitably constructed from a care plan template that is configured for a certain class of patients. FIGURE 3 illustrates an example "Post-Cardiac Event Care Plan" **100** that is suitable for assignment to patients within a class of patients who are overweight smokers and who have suffered a traumatic cardiac event such as a heart attack. The "Post-Cardiac Event Care Plan" **100** includes a list of pointers or identifiers of content modules that such a patient should access. These modules include the example weight loss, quit smoking, exercise, high fiber diet modules or others which are appropriate to overweight smokers who have suffered a traumatic cardiac event. For each listed content module, the "Post-Cardiac Event Care Plan" **100** also includes a default revision policy indicating which content revision should be presented. As illustrated, the default revision policy is to show the latest revision of each content module.

The "Post-Cardiac Event Care Plan" **100** further includes a set of intervention rules, optionally also tagged with revision information, such as the illustrated example revision 1.2 tag for the intervention rules of the "Post-Cardiac Event Care Plan" **100**. The intervention rules suitably are performed by the rules-based processor **26** of the server **12** shown in FIGURE 1 to order the presentation of the listed content modules, to automate actions on behalf of human care team members based on certain patient activities or supplied information, or so forth. For example, if the user interface includes a scale or other way to convey the patient's weight to the server **12**, then based on this weight the intervention rules may decide whether or not the patient needs to be presented with the weight loss module. Still further, the "Post-Cardiac Event Care Plan" **100** optionally includes a patient data repository template. This template may define, for example, data

partitions in the patient record for storing quiz results, biometric data supplied from the user interface, or so forth.

With continuing reference to FIGURE 3 and with further reference to FIGURE 4, when the "Post-Cardiac Event Care Plan" **100** is assigned to a specific patient, such as Patient #1, various parameters of the "Post-Cardiac Event Care Plan" **100** are optionally modified. For example, in the patient care plan **21** for Patient #1, the revision policy for the weight loss and quit smoking content modules has been modified to specify a specific content revision of those content modules. Thus, when following the patient care plan **21**, Patient #1 will be presented with weight loss module revision 1.3, which also happens to be the latest weight loss module.

However, Patient #1 will be presented with quit smoking module revision 6.2 even though revision 6.3 is the latest revision, because the patient care plan **21** for Patient #1 has modified the revision policy for the quit smoking module to call for the specific quit smoking module revision 6.2. For example, the latest quit smoking module revision 6.3 may instruct the patient to use a nicotine patch if the latest medical recommendations call for using such a nicotine patch. If this is now the most popular method for quitting smoking, then it can be expected that any future revisions of the quit smoking module will also probably call for using a nicotine patch. If, however, Patient #1's doctor does not believe that nicotine patches are a good method for quitting smoking, then Patient #1's doctor can modify the revision policy of the patient care plan **21** as shown in FIGURE 3 to call for Patient #1 to see the older quit smoking module revision 6.2 which does not instruct the patient to use a nicotine patch.

There are many other possible reasons why a particular doctor, or a particular patient, may want to specify a particular content revision. For example, Patient #1's doctor may have specified the specific weight loss control module revision 1.3, rather than keeping the default revision policy of using the latest version, because this doctor is concerned that future revisions may instruct patients in some sort of "fad" diet. By specifying the particular revision 1.3, Patient #1's doctor is assured that Patient #1 will receive the weight loss module revision 1.3 which is a known program.

Other doctors, or other patients, may have different revision policies for different content modules.

With reference to FIGURE 5, for example, the patient care plan **22** for Patient #2 employs the default latest version revision policy for the weight loss and high fiber diet modules. The patient care plan **22** for Patient #2 employs an "omit" revision policy for the quit smoking module. This optional revision policy omits the content module entirely. This
5 may be appropriate if, for example, Patient #2 does not intend to quit smoking regardless of the medical consequences of this decision.

The patient care plan **22** for Patient #2 also employs a "user query" revision policy for the exercise module. This optional revision policy asks the patient, at the time of presentation of the content module, which revision should be presented. To enable the
10 patient to make an informed choice, the tagged temporal content revision information includes semantic about the revision that is presented to the user as part of the query. For the example exercise module version 3.7, the tagged temporal content revision information includes the following semantic information: "This version employs the Jazzercise
ExerciseBike. You should select this version only if your ExerciseBike is labeled
15 'Jazzercise.'" For the example exercise module version 3.6, the tagged temporal content revision information includes the following semantic information: "This version employs the older version of ExerciseBike. You should select this version if your ExerciseBike does not include the label 'Jazzercise.'"

With continuing reference to FIGURE 5 and with further reference to FIGURE 6,
20 in accordance with the "user query" revision policy, Patient #2 is presented with a query constructed from the tagged semantic temporal content revision information, and Patient #2 is asked to select between revision 3.6 and revision 3.7. By giving Patient #2 the semantic information contained in the tagged temporal content revision information, Patient #2 can make an informed choice between revision 3.6 and revision 3.7 based on
25 which type of ExerciseBike Patient #2 has. The selected exercise module version is then presented to Patient #2. This "query user" revision policy can be useful in various situations in which the revised content reflects a change in medications, medical equipment, or so forth supplied to the patient. In some embodiments, the patient's doctor or other medical person may have the option of limiting the choices given to the patient in the
30 "user query" revision policy.

With continuing reference to FIGURE 5, the patient care plan **22** further includes an intervention rules version number 1.1, which it will be noted is an earlier rules revision

than the version number 1.2 included in the baseline "Post-Cardiac Event Care Plan" **100** shown in FIGURE 3. As the patient care plan itself, such as the intervention rules, may be revised occasionally, the care plan or components thereof such as the intervention rules include selectable revisions. In the illustrated embodiment, the care plan defaults to using the latest revision policy in which the latest revision of the intervention rules is used; accordingly, a rules revision policy is suitably only included in a specific patient care plan if, as in the example patient care plan **22**, an earlier revision is to be used.

With reference to FIGURE 7, the example patient care plan **23** for Patient #3 uses all of the default revision policies of the baseline "Post-Cardiac Event Care Plan" **100** shown in FIGURE 3, except that the revision policy for the high fiber diet module is modified to "latest version (retro)." This optional revision policy ensures that the patient receives the latest content revision by re-presenting the content module, or at least re-presenting those portions of the content module that are revised, if the content is revised after the patient has been presented with that content module.

With continuing reference to FIGURE 7 and with further reference to FIGURE 8, operation of the "latest version (retro)" revision policy is described in the context of patient care plan **23** followed by Patient #3. FIGURE 8 diagrammatically shows which content module is accessed by Patient #3 on each day of a particular five-day period running from Monday through Friday. On Monday, the weight loss module is accessed. Since the latest revision policy is in effect for this module, the latest weight loss module revision 1.3 is presented. On Tuesday, the exercise module is accessed. Again, the latest revision policy is in effect for this module, and so the latest exercise module revision 3.7 is presented. On Wednesday, the high fiber diet module is accessed. Here, the "latest version (retro)" policy is in effect. The latest revision as of Wednesday is version 8.3, since version 8.4 is still in-progress and is not yet ready for viewing. Accordingly, Patient #3 is presented with the current latest version high fiber diet module revision 8.3 on Wednesday. On Thursday, the quit smoking module is accessed. Since the latest revision policy is in effect for this module, the latest quit smoking module revision 7.0 is presented.

On Thursday night, the server **12** is updated as part of routine maintenance and updating of the server **12**. In this updating, the weight loss module is revised to version 1.4, which was previously in-progress. Accordingly, starting on Friday morning, the latest version is weight loss module revision 1.4, rather than the revision 1.3 that Patient #3

accessed on Monday. However, the latest revision policy is not retroactive. Accordingly, Patient #3 is never presented with the updated material of the weight loss module revision 1.4. In some embodiments, the in-progress modules can be designated to permit automatically changing to the updated version. This option may be appropriate where only not yet viewed sessions are updated, where additional sessions are added, and the like.

Additionally, however, the Thursday night updating includes revising the high fiber diet module from version 8.3 to version 8.4, which was previously in-progress. Accordingly, starting on Friday morning, the latest version is high fiber diet module revision 8.4, rather than the revision 8.3 that Patient #3 accessed on Wednesday. Unlike the situation for the weight loss module, the "latest version (retro)" revision policy is in effect for the high fiber diet module in the case of Patient #3. Accordingly, on Friday (or some other time subsequent to the Thursday updating) Patient #3 is presented with at least the updated components of the high fiber diet module revision 8.4. In this instant example, Patient #3's doctor may have selected the "latest version (retro)" revision policy for the high fiber diet module because that doctor is of the opinion that recent developments in medical knowledge of high fiber diets are significant advances, and so Patient #3's doctor wants to ensure that Patient #3 is presented with the latest high fiber diet information.

In the illustrated embodiment, the default revision policy is to use the latest revision. However, other default revision policies can be used. For example, other revision policies can be made default by modifying the defaults specified in the baseline "Post-Cardiac Event Care Plan" 100 shown in FIGURE 3. In some embodiments, the patient's doctor may select the default revision policy. For example, a more conservative doctor may select to use an older revision until he or she has seen and approved each new revision. In such embodiments, the doctor may receive notification by email or another automated messaging pathway each time a module assigned to one of his or her patients is revised. The doctor may also choose different revision policies for different patients, taking into account the type of patient, types of modules assigned to the patient, and other relevant information. Still further, other persons besides the doctor optionally may have the ability to set the revision policy for a given patient. For example, the patient, members of the patient's family, other doctors besides the primary doctor, or other interested parties may be given authorization to set or adjust the patient's revision policy.

Moreover, the example revision policies such as specific revision, latest revision, "user query", and "latest revision (retro)" are illustrative examples – the skilled artisan can readily construct other revision policies that are suited for specific situations. For example, another contemplated revision policy is to use information about the patient to define which revision should be presented – for example, a new revision containing subject matter related to women's health may be presented only to female patients. Another contemplated revision policy is to use the latest revision policy unless a retro-condition is met. For example, the patient may be presented with the latest revision unless the patient is an older piece of medical equipment, in which case an earlier revision that is appropriate for the older equipment is used.

The invention has been described with reference to the preferred embodiments. Obviously, modifications and alterations will occur to others upon reading and understanding the preceding detailed description. It is intended that the invention be construed as including all such modifications and alterations insofar as they come within the scope of the appended claims or the equivalents thereof.

CLAIMS

Having described the preferred embodiments, the invention is now claimed to be:

1. A server **(12)** for a patient care management system, the server comprising:
 - a content database **(14)** containing care management-related content including multiple revisions of at least some said content, each content revision being tagged with temporal content revision information;
 - a care plans storage **(18)** storing a plurality of patient care plans **(21, 22, 23)** associated with corresponding patients, each patient care plan including: (i) a selection of care management-related content to be presented to the corresponding patient and (ii) a revision policy indicating which temporal revision of said content should be presented to the corresponding patient; and
 - at least one communication interface **(28)** for communicating selected care management-related content to patients, the communicated content for each patient being selected based on the corresponding patient care plan, tagged temporal content revision information, and the revision policy of the corresponding patient care plan.
2. The server **(12)** as set forth in claim 1, wherein the revision policy is selected from a group of revision policy options including at least: (i) a specific revision, and (ii) the latest revision.
3. The server **(12)** as set forth in claim 2, wherein the tagged temporal content revision information includes an indication of which revision is the latest revision, the latest revision policy selecting that content revision indicated as the latest revision.
4. The server **(12)** as set forth in claim 2, wherein the tagged temporal content revision information includes ordering information, the latest revision policy selecting that content revision whose tagged temporal content revision ordering information indicates the content revision is the latest revision.
5. The server **(12)** as set forth in claim 2, wherein the group of revision policy options further includes: (iii) omission of the content.

6. The server (12) as set forth in claim 2, wherein the group of revision policy options further includes: (iii) latest revision retroactive, the at least one communication interface re-presenting at least an updated portion of the latest revision under the latest revision retroactive revision policy responsive to an updated latest content revision.

7. The server (12) as set forth in claim 2, wherein the group of revision policy options further includes: (iii) a user-selection policy, the at least one communication interface querying the patient via the communication interface as to which content revision to present and then presenting the user-selected content revision.

8. The server (12) as set forth in claim 7, wherein the tagged temporal content revision information includes semantic information about the revision that is presented to the user as part of the query.

9. The server (12) as set forth in claim 1, wherein the care management-related content of the content database (14) is organized into self-contained content modules, and the revision policy of each patient care plan (21, 22, 23) includes an independent revision policy for each selected content module that indicates which temporal revision of that content module should be presented to the corresponding patient.

10. The server (12) as set forth in claim 1, wherein each patient care plan (21, 22, 23) further includes:

(iii) intervention rules providing at least a temporal ordering for presentation of the care management-related content to the corresponding patient.

11. The server (12) as set forth in claim 10, wherein the intervention rules are also tagged with temporal content revision information, and each patient care plan (21, 22, 23) further includes (iii) a rules revision policy indicating which revision of intervention rules should be applied for the corresponding patient.

12. A care plan management system (10) comprising:
a server (12) as set forth in claim 1; and

a user interface **(31, 32, 33)** operatively connected with the at least one communication interface **(28)** of the server **(12)** to receive and present the selected care management-related content.

13. The care plan management system **(10)** as set forth in claim **12**, wherein the user interface **(31, 32, 33)** is an interactive user interface whereby an associated patient can provide input that is communicated to the server via the at least one communication interface **(28)**.

14. The care plan management system **(10)** as set forth in claim **12**, wherein the user interface **(31, 32, 33)** is operatively connected with the at least one communication interface **(28)** of the server **(12)** via at least one of the Internet, a cable television network, a satellite television network, and a cellular telephone network.

15. The care plan management system **(10)** as set forth in claim **12**, wherein the user interface **(31, 32, 33)** includes at least one of: a television set, a television set including a set-top box, a computer, a laptop computer, a personal data assistant (PDA), and a cellular telephone (cellphone).

16. The care plan management system **(10)** as set forth in claim **12**, wherein the user interface **(31, 32, 33)** includes at least one biometric feedback monitor that measures at least one biometric parameter of an associated patient that is communicated to the server via the at least one communication interface.

17. A method for managing care, the method comprising:
storing care management-related content including multiple revisions of at least some said content, each content revision being tagged with temporal content revision information;

storing a plurality of patient care plans **(21, 22, 23)** associated with corresponding patients, each patient care plan including: (i) a selection of care management-related content to be presented to the corresponding patient and (ii) a revision policy indicating which temporal revision of said content should be presented to the corresponding patient;
and

communicating selected care management-related content to patients, the communicated content for each patient being selected based on the corresponding patient care plan, tagged temporal content revision information, and the revision policy of the corresponding patient care plan.

18. The method as set forth in claim **17**, wherein the revision policy of each patient care plan is selected from a group of revision policy options including at least: (i) a specific revision policy, (ii) a latest revision policy, (iii) a latest revision retroactive policy, (iv) a user-query revision policy, and (v) a condition-based revision policy.

19. The method as set forth in claim **17**, wherein the revision policy of each patient care plan includes a plurality of generally different revision policies each applicable to different designated portion of the care management-related content.

20. The method as set forth in claim **19**, wherein the designated portions of the care management-related content are arranged as self-contained content modules.

21. A computer readable medium or media storing a program for managing care, the program when executed performing process operations comprising:

communicating selected care management-related content to a patient, the communicated content for each patient being selected based on a patient care plan, temporal content revision information tagged to the selected care management-related content, and a revision policy of the patient care plan.

22. The computer readable medium or media as set forth in claim **21**, wherein the program when executed further performs process operations comprising:


tagging an updated portion of the selected care management-related content with updated temporal content revision information, the communicating presenting one of the portion or the updated portion of the selected care management-related content to the patient based on the temporal content revision information and the updated temporal content revision information.


23. The computer readable medium or media as set forth in claim **21**, wherein the selected care management-related content is arranged as a plurality of self-contained content modules each tagged with temporal content revision information, and the communicating further comprises:

communicating content modules to a patient, the communicated content for each content module being selected based on temporal content revision information tagged to the content module and the revision policy of the patient care plan.



24. The computer readable medium or media as set forth in claim **23**, wherein the revision policy of the patient care plan is configured to apply a generally different revision policy for each communicated content module.

1 / 6



**Fig. 1**

**Fig. 2**

3 / 6

**Fig. 3****Fig. 4**

4 / 6

**Fig. 5****Fig. 6**

Patient #3 Care Plan	
Content module	Revision policy
Weight loss module	Latest version
Quit Smoking module	Latest version
Exercise module	Latest version
Fiber module	Latest version (retro)

Patient #3
data repository

Fig. 7

Patient #1 Care Plan Activity

Monday	Weight loss module v. 1.3
Tuesday	Exercise module v. 3.7
Wednesday	Fiber module v. 8.3
Thursday	Quit smoking module v. 7.0
Friday	<i>System Upgrade To:</i> <i>Weight loss module v 1.4</i> <i>Fiber module v. 8.4</i>
	Fiber module v. 8.4 (updated components)

Fig. 8