

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2002 (10.05.2002)

PCT

(10) International Publication Number
WO 02/36804 A2

(51) International Patent Classification⁷: **C12Q**

(21) International Application Number: PCT/US01/31365

(22) International Filing Date: 8 October 2001 (08.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/706,594 3 November 2000 (03.11.2000) US

(71) Applicant: **NEW YORK BLOOD CENTER, INC.**
[US/US]; 310 East 67th Street, New York, NY 10021 (US).

(72) Inventors: **BELYAVSKY, Alexander**; Apartment 4E, 179 Bennett Avenue, New York, NY 10040 (US). **SHMELKOV, Sergey**; Apartment 903, 100 Manhattan Avenue, Union City, NJ 07087 (US). **VISSER, Jan**; Apartment 30E, 1520 York Avenue, New York, NY 10028 (US).

(74) Agent: **ARNOLD, Craig, J.**; Amster, Rothstein & Ebenstein, 90 Park Avenue, New York, NY 10016 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR DETECTING HEMATOPOIETIC STEM CELLS

(57) **Abstract:** The present invention provides an isolated nucleic acid sequence encoding KIAA0918, and an isolated nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid that is complementary to a nucleic acid sequence encoding KIAA0918. The present invention also discloses a purified KIAA0918 protein, and a purified protein encoded by a nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid sequence that is complementary to a nucleic acid sequence encoding KIAA0918. Also provided is a method of making KIAA0918 protein. The present invention further discloses an antibody specific for KIAA0918, and a method for producing an antibody specific for KIAA0918 protein. Additionally, the present invention is directed to a vector comprising a nucleic acid sequence encoding KIAA0918, and a host cell transformed with a vector comprising a nucleic acid sequence encoding KIAA0918. Also provided is a method for detecting the presence of hematopoietic stem cells in a heterogeneous cell suspension that may contain hematopoietic stem cells, as well as a method for isolating hematopoietic stem cells from a heterogeneous cell suspension that may contain hematopoietic stem cells. Finally, the present invention discloses a method for assessing gene expression in a tissue sample.



WO 02/36804 A2

METHOD FOR DETECTING HEMATOPOIETIC STEM CELLS

Background of the Invention

[0001] In vertebrates, most tissues are composed of differentiated cells that no longer divide. Nevertheless, there are tissues which retain an 'embryonic' cell population within themselves. The cellular composition of such embryonic populations is always changing, even in adult animals. This phenomenon is most evident in the mammalian hematopoietic system. This system, which is organized hierarchically, consists of a heterogeneous mixture of many different kinds of blood cells at all stages of differentiation – some morphologically recognizable and some not [23]. Mature, functional blood cells are divided into several lines, including erythroid, lymphoid, and myeloid, each possessing its own morphology, characteristics, and function. Each blood line derives from restricted progenitor cells, which become committed to a particular line of differentiation. However, despite this diversity, the various developing blood cells and progenitor cells derive from one discrete source: the embryonic cell population of multipotential, self-renewing hematopoietic stem cells [21, 22].

[0002] A stem cell is a cell capable of extensive proliferation: it generates more stem cells (through self-renewal) in addition to its differentiated progeny [20, 21]. In mammals and birds, a multipotential hematopoietic stem cell can give rise to red blood cells (erythrocytes), white blood cells (granulocytes), macrophages, platelets, and immunocompetent cells (lymphocytes) [22]. Thus, a single hematopoietic stem cell can generate a clone containing millions of differentiated cells, as well as a few stem cells. The continuous formation of new blood cells is accomplished in bone marrow by hematopoietic stem cells. Stem cells mature into progenitor cells, which then become lineage-committed, although not yet terminally-differentiated. Once committed, progenitor cells are

-2-

no longer capable of maturing into all of the cell lineages which comprise the hematopoietic system [22].

[0003] Hematopoietic stem cells currently find use in a myriad of clinical settings. Indeed, with the recent remarkable progress in cell processing technology, there has been a rapid increase in the number of patients and types of diseases that are now treated with hematopoietic stem-cell transplantation, in both autologous and allogeneic cases. For example, autologous peripheral stem-cell support has largely replaced bone marrow transplantation as a means of regenerating the hematopoietic system of myeloma patients undergoing myeloablative chemotherapy. Stem cell transplantation is also used to treat patients with non-Hodgkin's lymphomas [25]. Moreover, peripheral blood autografting has been widely used in trials for the treatment of chemosensitive tumors [26].

[0004] Hematopoietic stem cells also may be used *in vitro* to enable the detection and assessment of growth factors associated with stem cell self-renewal and hematopoietic development. Furthermore, *in vitro* expansion of hematopoietic stem cells from various sources, including bone marrow and blood of the umbilical cord, is gaining importance, as it provides a clinically-potential graft in autologous or allogeneic transplant cases, and facilitates transduction of genes for the treatment of genetic diseases through gene therapy [24].

[0005] To be useful, however, hematopoietic stem cells first must be identified and isolated – a task made more difficult by the fact that hematopoietic stem cells comprise only a small proportion of the total cell population in bone marrow. Hematopoietic stem cells may be identified, for example, by detecting expression of specific cell-surface protein or carbohydrate antigen markers.

[0006] The CD34 antigen, which is a glycosylated transmembrane protein, has frequently been used as a marker for the identification of hematopoietic stem cells [26]. Detection of CD34 is also considered by some to

-3-

be the first step in quality assessment of hematopoietic stem cell grafts [27]. The CD34 antigen was previously indicated to be present solely on stem cells, and not on lineage-committed progenitor cells. However, recent evidence suggests that expression of CD34 on the cell membrane does not always correlate with stem cell activity [28]. Indeed, in the mouse, there is a highly quiescent population of stem cells that lacks CD34 expression, but which has full reconstituting capacity. It also has been discovered that there is a similar population of dormant CD34-negative human hematopoietic stem cells. This information clearly casts some uncertainty on the benefits of using CD34 as a marker for isolating hematopoietic stem cells [28].

[0007] While it appears that lineage-committed hematopoietic cells may display epitopic characteristics associated with hematopoietic stem cells, it is not known how many of the antigen markers associated with differentiated cells are also present on stem cells. Accordingly, it is clear that significant problems exist in connection with the identification of hematopoietic stem cells, and new methods of detection of hematopoietic stem cells are needed.

Summary of the Invention

[0008] The present invention is predicated on the discovery that the KIAA0918 gene is specifically expressed in the primitive hematopoietic Kg-1a cell line, which is close to the hematopoietic stem cell line, thereby providing a genetic marker for identifying hematopoietic stem cells. On the basis of this finding, the present invention provides an isolated nucleic acid sequence encoding KIAA0918, and an isolated nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid that is complementary to a nucleic acid sequence encoding KIAA0918.

[0009] The present invention also discloses a purified KIAA0918 protein, and a purified protein encoded by a nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid sequence that is

-4-

complementary to a nucleic acid sequence encoding KIAA0918. Also provided is a method of making KIAA0918 protein.

[0010] The present invention is further directed to an antibody specific for KIAA0918, and a method for producing an antibody specific for KIAA0918 protein.

[0011] Additionally, the present invention discloses a vector comprising a nucleic acid sequence encoding KIAA0918, and a host cell transformed with a vector comprising a nucleic acid sequence encoding KIAA0918.

[0012] Also provided in the present invention is a method for detecting the presence of hematopoietic stem cells in a heterogeneous cell suspension that may contain hematopoietic stem cells, as well as a method for isolating hematopoietic stem cells from a heterogeneous cell suspension that may contain hematopoietic stem cells.

[0013] Finally, the present invention discloses a method for assessing gene expression in a tissue sample.

[0014] Additional objects of the present invention will be apparent in view of the description which follows.

Brief Description of the Figures

[0015] Figure 1 depicts a scheme of the two-dimensional gene expression fingerprinting procedure and the computer-assisted prediction of cDNA fragment identity. The first dimension strip in the middle of the figure corresponds to the cDNA sample obtained from Kg-1 cells using NdeII as primary enzyme. ◆=biotin groups; SavB=Streptavidin beads

[0016] Figure 2 illustrates a region of the second-dimension polyacrylamide gel electrophoresis (PAGE) performed in the *quasi*-2-D format. cDNA and samples from Kg-1 (lane 1) and Kg-1a (lane 2) cell lines were loaded side-by-side. The picture presented in the figure was generated by the StyI restriction enzyme treatment. Prior to StyI, the cDNA on beads was treated sequentially with EcoRV, XbaI, ApaI, NsiI, StuI, and PstI. The arrow

indicates the SequaMark™ (Research Genetics) marker lane. Positions of selected SequaMark bands are indicated on the left side; it should be noted, though, that this marker provides 1 base resolution for lengths ranging from 40 to 400 bases.

[0017] Figure 3 sets forth a Northern blot analysis of total RNA isolated from Kg-1 and Kg-1a cell lines. Equal amounts of total RNA (20 μ g) were loaded into lanes 1 (Kg-1) and 2 (Kg-1a). Blots were hybridized with ³²P-labeled cloned probes, corresponding to fragments NB66 (EST AA428186), NB63 (EST H11385), NB114 (EST H93036), NB112 (EST AA142937), NB67 (EST AA617864), NB91 (EST AA573912), NB101 (obtained using 3' rapid amplification of cDNA ends (RACE) procedure), NB113 (EST AI144181), NB96 and NB16 (both obtained using RT-PCR with two specific primers), NB68 (EST AI050734), and NB64 (EST AA814890). Positions of 0.24- to 9.5-Kb RNA ladder bands (Gibco-BRL) are indicated on the left side. The 9.5-kb band is not shown.

[0018] Figure 4 depicts the nucleotide sequence of human KIAA0918.

[0019] Figure 5 depicts the amino acid sequence of human KIAA0918.

Detailed Description of the Invention

[0020] The present invention provides an isolated nucleic acid sequence encoding KIAA0918. KIAA0918 is a protein which has previously been identified, but for which there was previously no known use. As used herein, KIAA0918 includes, where appropriate, both a KIAA0918 protein and a "KIAA0918 analogue". Unless otherwise indicated, "protein" shall mean a protein, protein domain, polypeptide, or peptide. A "KIAA0918 analogue" may be any protein having functional similarity to the KIAA0918 protein that is 80% or greater (preferably, 90% or greater) in amino-acid-sequence homology with the KIAA0918 protein.

[0021] The nucleic acid sequence of the present invention may be genomic DNA, cDNA, RNA, or antisense RNA, and may be derived from any

species. The nucleic sequence of the present invention is preferably derived from a mammalian species, and, more preferably, from a human. The nucleic acid sequence of the present invention preferably comprises the nucleotide sequence of Figure 4 (including conservative substitutions thereof).

"Conservative substitutions" are those amino acid substitutions which are functionally equivalent to the substituted amino acid residue, either because they have similar polarity or steric arrangement, or because they belong to the same class as the substituted residue (*e.g.*, hydrophobic, acidic, or basic). The nucleic acid of the present invention also may encode a KIAA0918 protein comprising the amino acid sequence set forth in Figure 5.

[0022] The present invention further discloses an isolated nucleic acid sequence that hybridizes under high stringency conditions (*e.g.*, hybridization to filter-bound DNA in 0.5-M NaHPO₄ at 65°C and washing in 0.1X SSC/0.1% SDS at 68°C) or moderate stringency conditions (*e.g.*, washing in 0.2X SSC/0.1% SDS at 42°C) [29] to a second nucleic acid that is complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof. In addition, the present invention provides a nucleic acid sequence encoding KIAA0918 protein having one or more mutations that result in the expression of either a non-functional or mutant protein, or in a lack of expression altogether. The mutation may be generated by at least one point, insertion, rearrangement, or deletion mutation, or a combination thereof.

[0023] The present invention also provides an isolated and purified KIAA0918 protein. The KIAA0918 protein may be isolated from tissue obtained from a subject, or recombinantly produced as described below. The KIAA0918 protein of the present invention may comprise the amino acid sequence set forth in Figure 5. Alternatively, the KIAA0918 protein of the present invention may be encoded by the nucleotide sequence set forth in Figure 4.

[0024] The present invention is further directed to a purified protein encoded by a nucleic acid sequence that hybridizes under high stringency or moderate stringency conditions to a second nucleic acid sequence that is

-7-

complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof.

[0025] The present invention also provides agents that bind to KIAA0918 protein. The agent may include, without limitation, an antibody, a compound, a drug, a Fab fragment, a $F(ab')_2$ fragment, a molecule, a nucleic acid, a protein (including a growth factor), a polypeptide, a peptide, a nucleic acid (including DNA, RNA, mRNA, antisense RNA), and any combinations thereof. Moreover, an agent that binds to KIAA0918 protein may be either natural or synthetic. A Fab fragment is a univalent antigen-binding fragment of an antibody, which is produced by papain digestion. An $F(ab')_2$ fragment is a divalent antigen-binding fragment of an antibody, which is produced by pepsin digestion. Agents that bind to KIAA0918 protein may be identified or screened by contacting the protein with the agent of interest, and assessing the ability of the agent to bind to the protein.

[0026] The agent of the present invention is preferably an antibody specific for, or immunoreactive with, KIAA0918. The antibody of the present invention may be monoclonal or polyclonal. Furthermore, the antibody of the present invention may be produced by techniques well known to those skilled in the art. The antibody of the present invention may be incorporated into kits which include an appropriate labeling system, buffers, and other necessary reagents for use in a variety of detection and diagnostic applications. Labeling of the antibody of the present invention may be accomplished by standard techniques using one of the variety of different chemiluminescent and radioactive labels known in the art.

[0027] The present invention further provides a method for producing an antibody specific for KIAA0918 protein, comprising the steps of: (a) immunizing a mammal with KIAA0918 protein; and (b) purifying antibody from a tissue of the mammal or from a hybridoma made using tissue of the mammal. For example, a polyclonal antibody may be produced by immunizing a rabbit, mouse, or rat with purified KIAA0918. Thereafter, a monoclonal antibody may

-8-

be produced by removing the spleen from the immunized rabbit, mouse, or rat, and fusing the spleen cells with myeloma cells to form a hybridoma which, when grown in culture, will produce a monoclonal antibody. Also provided is an antibody produced by this method.

[0028] The present invention further discloses agents that bind to a nucleic acid encoding KIAA0918 protein. Suitable agents include, but are not limited to, an antibody, a compound, a drug, a Fab fragment, a $F(ab')_2$ fragment, a molecule, a nucleic acid, a protein, a polypeptide, a peptide, a nucleic acid (including DNA, RNA, mRNA, antisense RNA), and any combinations thereof. The agents that bind to a nucleic acid encoding KIAA0918 may inhibit or promote expression of the nucleic acid. Such agents may be discovered by a method for screening for an agent that binds to a nucleic acid encoding KIAA0918, comprising contacting the nucleic acid with an agent of interest, and assessing the ability of the agent to bind to the nucleic acid. An agent that inhibits or promotes the expression of a nucleic acid encoding KIAA0918 may be screened by contacting a host cell transformed with a vector comprising the nucleic acid, and assessing the agent's effect on expression of the nucleic acid.

[0029] The present invention also provides nucleic acid probes and mixtures thereof which hybridize to nucleic acid encoding KIAA0918 protein. Such probes may be prepared by a variety of techniques known to those skilled in the art, including, without limitation, PCR and restriction-enzyme digestion of nucleic acid encoding KIAA0918; and automated synthesis of oligonucleotides whose sequences correspond to selected portions of the nucleotide sequence of nucleic acid encoding KIAA0918, using commercially-available oligonucleotide synthesizers such as the Applied Biosystems Model 392 DNA/RNA synthesizer. The nucleic acid probes of the present invention also may be prepared so that they contain at least one point, insertion, rearrangement, or deletion mutation, or a combination thereof, to correspond to mutations of the KIAA0918 gene.

[0030] The nucleic acid probes of the present invention may be DNA or RNA, and may vary in length from about 8 nucleotides to the entire length of the nucleic acid encoding KIAA0918. Preferably, the probes are 8 to 30 nucleotides in length. Labeling of the nucleic acid probes may be accomplished using one of a number of methods known in the art, including, without limitation, PCR, nick translation, end labeling, fill-in end labeling, polynucleotide kinase exchange reaction, random priming, or SP6 polymerase (for riboprobe preparation), and one of a variety of labels, including, without limitation, radioactive labels such as ^{35}S , ^{32}P , or ^3H and nonradioactive labels such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine (ROX). Combinations of two or more nucleic probes, corresponding to different or overlapping regions of nucleic acid encoding KIAA0918, also may be included in kits for use in a variety of detection and diagnostic applications.

[0031] The present invention is further directed to a vector comprising a nucleic acid sequence encoding KIAA0918 protein. The vector of the present invention may comprise the nucleotide sequence of Figure 4 or a contiguous fragment thereof. Alternatively, the vector of the present invention may comprise a nucleic acid sequence which hybridizes under high stringency or moderate stringency conditions to a nucleic acid sequence that is complementary to the nucleotide sequence set forth in Figure 4, or to a contiguous fragment thereof.

[0032] The vector of the present invention may be constructed by inserting nucleic acid encoding KIAA0918 into a suitable vector nucleic acid operably linked to an expression control sequence, as described below. The term "inserted", as used herein, means the ligation of a foreign DNA fragment with vector DNA, by techniques such as the annealing of compatible cohesive ends generated by restriction endonuclease digestion, or by the use of blunt-end ligation techniques. Other methods of ligating DNA molecules will be apparent to one skilled in the art.

-10-

[0033] The vector of the present invention may be derived from a number of different sources, including plasmids, viral-derived nucleic acids, lytic bacteriophage derived from phage lambda, cosmids, or filamentous single-stranded bacteriophages such as M13. Depending upon the type of host cell into which the vector is introduced, vectors may be bacterial or eukaryotic. Bacterial vectors are derived from many sources, including the genomes of plasmids and phages. Eukaryotic vectors are constructed from a number of different sources, *e.g.*, yeast plasmids and viruses. Some vectors, referred to as shuttle vectors, are capable of replicating in both bacteria and eukaryotes. The nucleic acid from which the vector is derived is usually greatly reduced in size, such that only those genes essential for its autonomous replication remain. This reduction in size enables the vectors to accommodate large segments of foreign DNA. Examples of suitable vectors into which nucleic acid encoding KIAA0918 can be inserted include, but are not limited to, pCGS, pBR322, pUC18, pUC19, pHSV-106, pJS97, pJS98, M13mp18, M13mp19, pSPORT 1, pGem, pSPORT 2, pSV●SPORT 1, pBluescript II, λ ZapII, λ gt10, λ gt11, λ gt22A, and λ ZIPLOX. Other suitable vectors will be obvious to one skilled in the art.

[0034] The vector of the present invention may be introduced into a host cell. Accordingly, the present invention provides a host cell transformed with the vector of the present invention. The term "host cell", as used herein, means the bacterial or eukaryotic cell into which the vector is introduced. The term "transform" denotes the introduction of a vector into a bacterial or eukaryotic host cell. Additionally, as used herein, the term "introduction" is a general term indicating that one of a variety of means has been used to allow the vector to enter the intracellular environment of the host cell in such a way that the nucleic acid exists in stable form, and may be expressed, therein. As such, it encompasses transformation of bacterial cells, as well as transfection, transduction, and related methods in eukaryotic cells. The vector of the present invention may exist in integrated or unintegrated form within the host cell. When in unintegrated form, the vector is capable of autonomous replication.

[0035] Any one of a number of suitable bacterial or eukaryotic host cells may be transformed with the vector of the present invention. Examples of suitable host cells are known to one skilled in the art, and include, without limitation, bacterial cells such as *Escherichia coli* strains c600, c600hfl, HB101, LE392, Y1090, JM103, JM109, JM101, JM107, Y1088, Y1089, Y1090, Y1090(ZZ), DM1, PH10B, DH11S, DH125, RR1, TB1 and SURE, *Bacillus subtilis*, *Agrobacterium tumefaciens*, *Bacillus megaterium*; and eukaryotic cells such as *Pichia pastoris*, *Chlamydomonas reinhardtii*, *Cryptococcus neoformans*, *Neurospora crassa*, *Podospora anserina*, *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Uncinula necator*, cultured insect cells, cultured chicken fibroblasts, cultured hamster cells, cultured human cells such as HT1080, MCF7, and 143B, and cultured mouse cells such as EL4 and NIH3T3 cells.

[0036] Some bacterial and eukaryotic vectors have been engineered so that they are capable of expressing inserted nucleic acids to high levels within the host cell. An "expression cassette" or "expression control sequence", comprising nucleic acid encoding a KIAA0918 protein operably linked or under the control of transcriptional and translational regulatory elements (e.g., a promoter, ribosome binding site, operator, or enhancer), can be made and used for expression of KIAA0918 protein *in vitro* or *in vivo*. As used herein, "expression" refers to the ability of the vector to transcribe the inserted nucleic acid into mRNA, so that synthesis of the protein encoded by the inserted nucleic acid can occur. The choice of regulatory elements employed may vary, depending on such factors as the host cell to be transformed and the desired level of expression.

[0037] For example, in vectors used for the expression of a gene in a bacterial host cell such as *Escherichia coli*, the lac operator-promoter or the tac promoter is often used. Eukaryotic vectors use promoter-enhancer sequences of viral genes, especially those of tumor viruses. Several promoters for use in mammalian cells are known in the art and include, for example, the phosphoglycerate (PGK) promoter, the simian virus 40 (SV40) early promoter,

-12-

the Rous sarcoma virus (RSV) promoter, the adenovirus major late promoter (MLP), and the human cytomegalovirus (CMV) immediate early 1 promoter. However, any promoter that facilitates suitable expression levels can be used in the present invention. Inducible promoters (e.g., those obtained from the heat shock gene, metallothionine gene, beta-interferon gene, or steroid hormone responsive genes, including, without limitation, the lac operator-promoter in *E. coli* and metallothionine or mouse mammary tumor virus promoters in eukaryotic cells) may be useful for regulating transcription based on external stimuli.

[0038] Vectors suitable for expression in a host cell of nucleic acid encoding KIAA0918 are well-known to one skilled in the art, and include pET-3d (Novagen), pProEx-1 (Life Technologies), pFastBac 1 (Life Technologies), pSFV (Life Technologies), pcDNA II (Invitrogen), pSL301 (Invitrogen), pSE280 (Invitrogen), pSE380 (Invitrogen), pSE420 (Invitrogen), pTrcHis A,B,C (Invitrogen), pRSET A,B,C (Invitrogen), pYES2 (Invitrogen), pAC360 (Invitrogen), pVL1392 and pVL1392 (Invitrogen), pCDM8 (Invitrogen), pcDNA I (Invitrogen), pcDNA I(amp) (Invitrogen), pZeoSV (Invitrogen), pcDNA 3 (Invitrogen), pRc/CMV (Invitrogen), pRc/RSV (Invitrogen), pREP4 (Invitrogen), pREP7 (Invitrogen), pREP8 (Invitrogen), pREP9 (Invitrogen), pREP10 (Invitrogen), pCEP4 (Invitrogen), pEBVHis (Invitrogen), and λ Pop6. Other vectors will be apparent to one skilled in the art.

[0039] The vector of the present invention may be introduced into a host cell using conventional procedures known in the art, including, without limitation, electroporation, DEAE Dextran transfection, calcium phosphate transfection, monocationic liposome fusion, polycationic liposome fusion, protoplast fusion, creation of an *in vivo* electrical field, DNA-coated microprojectile bombardment, injection with recombinant replication-defective viruses, homologous recombination, *in vivo* gene therapy, *ex vivo* gene therapy, viral vectors, and naked DNA transfer, or any combination thereof. For the purposes of gene transfer into a host cell, tissue, or subject, a recombinant

-13-

vector containing nucleic acid encoding KIAA0918 may be combined with a sterile aqueous solution which is preferably isotonic with the blood of the recipient. Such formulations may be prepared by suspending the recombinant vector in water containing physiologically-compatible substances, such as sodium chloride, glycine, and the like, and having buffered pH compatible with physiological conditions, to produce an aqueous solution, then rendering the solution sterile. In a preferred embodiment of the invention, the recombinant vector is combined with a 20-25% sucrose in saline solution in preparation for introduction into a mammal.

[0040] The present invention further provides a method of making recombinant KIAA0918 protein, comprising the steps of: (a) introducing into a suitable bacterial or eukaryotic host cell a nucleic acid sequence encoding KIAA0918 (or a nucleic acid that hybridizes under high stringency conditions or moderate stringency conditions to a second nucleic acid that is complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof); (b) maintaining the host cell under conditions such that the nucleic acid sequence is expressed to produce KIAA0918 protein; and (c) recovering the recombinant KIAA0918 protein from the culture medium, from the host cells, or from cell lysate. As used herein, the term "recombinant" refers to KIAA0918 produced by purification from a host cell transformed with a vector capable of directing its expression to a high level. In the method of the present invention, a nucleic acid sequence encoding KIAA0918 may be introduced into a suitable host cell by any of the above-described methods.

[0041] A variety of methods of growing host cells transformed with a vector are known to those skilled in the art. The type of host cell (*i.e.*, bacterial or eukaryotic) is the primary determinant of both the method to be utilized and the optimization of specific parameters relating to such factors as temperature, trace nutrients, humidity, and growth time. Depending on the vector used, the host cells may have to be induced by the addition of a specific compound at a certain point in the growth cycle, in order to initiate expression of the nucleic

-14-

acid contained in the vector. Examples of compounds used to induce expression of the nucleic acid contained in the vector are known to one skilled in the art, and include, without limitation, IPTG, zinc, and dexamethasone. Using standard methods of protein isolation and purification, such as ammonium sulfate precipitation and subsequent dialysis to remove salt, followed by fractionation according to size, charge of the protein at specific pH values, affinity methods, etc., recombinant KIAA0918 may be extracted from suitable host cells transformed with a vector capable of expressing nucleic acid encoding KIAA0918.

[0042] The present invention also discloses a method for detecting the presence of hematopoietic stem cells in a heterogeneous cell suspension that may contain hematopoietic stem cells. The heterogeneous cell suspension may be, for example, adult bone marrow, embryonic yolk sac, fetal liver, adult peripheral blood, spleen, or umbilical cord blood. Moreover, the heterogeneous cell suspension may be a tissue sample obtained from any mammal (e.g., humans, domestic animals, and commercial animals), but is preferably obtained from a human. A sample of cells from a mammal may be removed using standard procedures, including biopsy and aspiration.

[0043] The method of the present invention comprises detecting cells expressing KIAA0918 in a heterogeneous cell suspension, wherein the detection of expression of KIAA0918 indicates the presence of hematopoietic stem cells. Expression of KIAA0918 may be detected using an agent reactive with KIAA0918. As used herein, "reactive" means the agent has affinity for, binds to, or is directed against KIAA0918. The agent may be an antibody, a compound, a drug, a Fab fragment, a F(ab')₂ fragment, a molecule, a nucleic acid, a protein (including a growth factor), a polypeptide, a peptide, a nucleic acid (including DNA, RNA, mRNA, antisense RNA), and any combinations thereof. Furthermore, the agent may be labeled with a detectable marker.

[0044] For example, the agent of the present invention may be a monoclonal or polyclonal antibody. Preferably, the antibody is a high-affinity

-15-

antibody, labeled with a detectable marker. The antibody of the present invention may be produced by techniques well known to those skilled in the art, including those described above, and may be labeled with detectable markers, in accordance with the above-described methods. Where the agent is an antibody immunoreactive with KIAA0918, expression of KIAA0918 may be detected from standard immunological detection techniques. For example, the method may be performed by contacting the heterogeneous cell suspension with an antibody immunoreactive with KIAA0918, under conditions permitting the antibody to bind to KIAA0918, if present in the suspension, to form an antibody-KIAA0918 complex. Appropriate binding conditions (*e.g.*, temperature, pH, and salt concentration) are readily determinable by the skilled artisan. Thereafter, the presence of the complex may be detected by a variety of immunological detection techniques, including an ELISA assay, a Western blot, flow cytometry, hemagglutination, or any other immunostaining method employing an antigen-antibody interaction.

[0045] Alternatively, the agent of the present invention may be a nucleic acid probe which hybridizes to nucleic acid encoding KIAA0918. The presence of cells expressing KIAA0918 in a heterogeneous cell suspension may be detected using hybridization analysis of nucleic acid extracted from the heterogeneous cell suspension. According to this method of the present invention, the hybridization analysis may be conducted using one or more nucleic acid probes which hybridize to nucleic acid encoding KIAA0918. The probes may be prepared by a variety of techniques known to those skilled in the art, including, without limitation, restriction enzyme digestion of nucleic acid encoding KIAA0918; and automated synthesis of oligonucleotides whose sequences correspond to selected portions of the nucleotide sequence of nucleic acid encoding KIAA0918, using commercially-available oligonucleotide synthesizers, such as the Applied Biosystems Model 392 DNA/RNA synthesizer.

[0046] The nucleic acid probes used in the present invention may be DNA or RNA, and may vary in length from about 8 nucleotides to the entire length of

-16-

the nucleic acid encoding KIAA0918. The nucleic acid used in the probes may be derived from mammalian KIAA0918. The nucleotide sequence for human KIAA0918 is depicted in Figure 4. Using this sequence as a probe, the skilled artisan could readily clone corresponding KIAA0918 cDNA from other species. In addition, the nucleic acid probes of the present invention may be labeled with one or more detectable markers. Labeling of the nucleic acid probes may be accomplished using one of a number of methods known in the art (e.g., nick translation, end labeling, fill-in end labeling, polynucleotide kinase exchange reaction, random priming, or SP6 polymerase for riboprobe preparation), along with one of a variety of labels (e.g., radioactive labels, such as ^{35}S , ^{32}P , or ^3H , or nonradioactive labels, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine (ROX)). Combinations of two or more nucleic acid probes (or primers), corresponding to different or overlapping regions of the nucleic acid encoding KIAA0918, also may be used to detect expression of KIAA0918, using, for example, PCR or RT-PCR.

[0047] The present invention also contemplates the provision of a kit for performing a method of detecting the presence of hematopoietic stem cells in a heterogeneous cell suspension that may contain hematopoietic stem cells, as described above. The kit would include a container, an agent which binds to KIAA0918 to form a complex, and a reagent or reagents capable of detecting the resulting complex. As noted above, the agent may be any of an antibody, a compound, a drug, a Fab fragment, a F(ab')_2 fragment, a molecule, a nucleic acid, a protein (including a growth factor), a polypeptide, a peptide, a nucleic acid (including DNA, RNA, mRNA, antisense RNA), and any combinations thereof. The reagent or reagents capable of detecting the complex are preferably secondary antibodies that bind selectively to one or the other of the antibody and KIAA0918, preferably KIAA0918, and which are further linked, either through a covalent linkage or by a noncovalent linkage, to a reporter molecule, including, without limitation, an enzyme, a fluorescent molecule, a light-emitting molecule, or a radioactive molecule. If antibody binding to

hematopoietic stems cells results in cell agglutination, the reagent capable of detecting the resulting complex will not be necessary.

[0048] The present invention is also directed to a method for isolating hematopoietic stem cells from a heterogeneous cell suspension that may contain hematopoietic stem cells. The heterogeneous cell suspension may be, for example, adult bone marrow, embryonic yolk sac, fetal liver, adult peripheral blood, spleen, and umbilical cord blood. Moreover, the heterogeneous cell suspension may be a tissue sample obtained from any mammal (*e.g.*, humans, domestic animals, and commercial animals), but is preferably obtained from a human. A sample of cells from a mammal may be removed using standard procedures, including biopsy and aspiration.

[0049] The method of the present invention comprises contacting the heterogeneous cell suspension with a monoclonal antibody specific for KIAA0918, and separating from the heterogeneous cell suspension cells to which the monoclonal antibody is bound. A monoclonal antibody specific for KIAA0918 may be produced in accordance with the procedures described above. The monoclonal antibody specific for KIAA0918 may be contacted with the heterogeneous cell suspension by adding the antibody to the cell suspension under conditions and for a time suitable for antibody binding. Appropriate binding conditions (*e.g.*, temperature, pH, and salt concentration) are readily determinable by the skilled artisan. Cells to which the monoclonal antibody is bound then may be separated from the heterogeneous cell suspension using one of the various methods of separation known in the art.

[0050] In the method of the present invention, the appropriate method of separation will depend on the degree of enrichment of hematopoietic stem cells which is desired, as well as the ease, efficacy, efficiency, and speed of the separation technique that is employed. Examples of methods of separation include, without limitation, affinity chromatography, flow cytometry, fluorescence-activated cell sorting, and magnetic separation using antibody-coated magnetic beads. It is also within the confines of the present invention

-18-

that hematopoietic stem cells may be separated from a heterogeneous cell suspension by passing the suspension over a solid support, such as an insoluble organic polymer in the form of a bead, gel, or plate, to which is attached a monoclonal antibody specific for KIAA0918. Examples of suitable solid supports include, without limitation, agarose, cellulose, dextran, polyacrylamide, polystyrene, or other insoluble organic polymers. The monoclonal antibody may further be attached to the solid support through a spacer molecule, if desired.

[0051] The method of the present invention further comprises enriching the concentration of hematopoietic stem cells in the cell suspension before contacting the cell suspension with the KIAA0918. The concentration of hematopoietic stem cells may be enriched, for example, through blood fractionation, density gradient separation, flow cytometry, or magnetic bead separation. Magnetic bead separation may be useful for removing numerous lineage-committed cells, such as basophils, eosinophils, mast cells, megakaryocytes, T-cells, and B-cells.

[0052] The present invention also discloses a method for assessing gene expression in a sample of tissue or cells, comprising the steps of: (a) obtaining cDNA from the sample; (b) digesting cDNA obtained from the sample with a primary restriction enzyme to produce a primary set of 3'-terminal restriction fragments; (c) separating in the first dimension, by denaturing polyacrylamide gel electrophoresis (PAGE), the 3'-terminal restriction fragments; (d) sequentially treating the 3'-terminal restriction fragments with a set of secondary restriction enzymes, to produce secondary restriction fragments; (e) separating in the second dimension, by PAGE, the secondary restriction fragments, to generate two-dimensional coordinates; and (f) assessing gene expression in the sample by assessing the two-dimensional coordinates. This method, referred to herein as two-dimensional Gene Expression Fingerprinting (2-D GEF), is based on the two-dimensional gel display of 3'-terminal cDNA restriction fragments produced by one primary (first dimension) digestion and

several sequential secondary restriction digestions. This sequence of digestion leads to subdivision of the initial cDNA fragment population into many thousands of non-overlapping subsets. As a result, many thousands of individual sequences per cDNA sample can be visualized using this approach, which is also characterized by a high reproducibility and predictable spatial location of cDNA fragments on two-dimension gels. The 2-D GEF procedure is advantageous because it permits the assessment of gene expression in two ways: (i) it allows global analysis of gene expression; and (ii) it facilitates the discovery of new genes.

[0053] In the method of the present invention, cDNA from a tissue or cell sample may be obtained by any method known to one of skill in the art, including isolation of RNA followed by first- and second-strand synthesis of cDNA. Isolation of RNA from cells is well known in the art, and may be accomplished by a number of techniques. For example, whole cell RNA may be extracted using guanidine thiocyanate; cytoplasmic RNA may be prepared by using phenol extraction methods or an RNAgents kit (Promega); and polyadenylated RNA may be selected using oligo-dT cellulose. Thereafter, double-stranded cDNA may be synthesized, extracted, and precipitated, according to methods well known in the art. First-strand cDNA synthesis may be performed using a one-base anchored 5'-biotinylated oligo(dT)-containing primer, followed by second-strand synthesis using the RNaseH-DNA polymerase I protocol [15].

[0054] In the method of the present invention, cDNA obtained from a tissue sample is digested with a primary restriction enzyme, such as a four-base-recognition primary restriction enzyme, to produce a primary set of 3'-terminal cDNA restriction fragments. This set of fragments may be selected by binding to Streptavidin beads, then ligated to a double-stranded adapter. Following PCR amplification with biotinylated oligo(dT) and adapter primers, this primary set of 3'-terminal restriction fragments is then separated in the first dimension, according to size, by denaturing PAGE.

[0055] For transfer to the second dimension, the gel lane containing biotinylated cDNA fragments may be cut into slices. The slices then may be treated in separate wells of a microliter plate. For example, they may be eluted, bound to Streptavidin beads, and rendered double-stranded by a Sequenase synthesis initiated from a highly-labeled adapter primer. The cDNA fragments then are treated sequentially with a set of restriction enzymes to liberate subpopulations of secondary restriction fragments. The secondary restriction fragments are then separated in the second dimension, by PAGE; thereafter, they may be analyzed on separate gels. Restriction enzymes suitable for the method of the present invention may be selected in such a way as to produce relatively similar numbers of liberated fragments per each round of restriction digestion. Examples of restriction enzymes suitable for primary and secondary digestion include, without limitation, ApaI, AclI, EcoRI, EcoRV, HinfI, MspI, NdeI, NsiI, PstI, SacI, StyI, and Xba.

[0056] In one embodiment of the present invention, the secondary restriction fragments may be separated in the second dimension by standard two-dimensional gene expression fingerprinting format. In a standard 2-D GEF format, intended for the global analysis of gene expression in one cDNA sample, all fragments liberated by a given enzyme are loaded onto one gel to generate a 2-D-like pattern of bands. Alternatively, the secondary restriction fragments may be separated in the second dimension by *quasi*-two-dimensional gene expression fingerprinting (*quasi*-2-D GEF). This format is preferable for comparative analysis of gene expression in two or more cell samples. In *quasi*-2-D GEF, cDNA fragments originating from the same size fraction of the first-dimension gel, but from different cell samples, are loaded side-by-side on each gel.

[0057] Following separation of the secondary restriction fragments in the second dimension, to generate two-dimensional coordinates, gene expression in the tissue sample is assessed by assessing the two-dimensional coordinates so generated. In one embodiment of the present invention, gene expression may

be assessed using computer-assisted prediction of two-dimensional coordinates of cDNA fragments. Due to the excellent resolving power of the 2-D GEF procedure, the identity of bands on the second-dimension (2-D) gels may be established on the basis of their two-dimensional coordinates. In this regard, it is possible to create software simulating the 2-D GEF procedure.

[0058] For example, sequences obtained from a gene database, such as the NCBI UniGene database, may be subjected to computer-simulated primary and sequential secondary digestions, to generate a subdatabase of predicted two-dimensional coordinates for all known human genes. It is contemplated herein that such a subdatabase would be useful for establishing the correspondence between known genes and bands on the 2-D gels. This procedure may be performed in numerous ways, including the following: (i) from database to the gel: identification of fragments on 2-D gels which correspond to selected known genes; or (ii) from gel to the database: identification of genes in the database which might correspond to selected fragments on the 2-D gels. Use of the 2-D GEF procedure in combination with computer software permits a computer-based sampling of existing databases for possible matches with a given fragment detected in a tissue sample, and permits identification of cDNA fragments solely on the basis of their two-dimensional coordinates.

[0059] In another embodiment of the present invention, gene expression may be assessed using an expressed sequence tag (EST) derived from an EST clone collection. For example, the 2-D GEF gel patterns of the present invention easily can be compared against the sequence information in an EST database for a given organism, permitting identification in digested cDNA obtained from a tissue sample of a particular EST, including one used as a genetic marker. Comparison between the 2-D GEF gel patterns and the sequence information in an EST also may reveal fragments likely to represent novel mRNAs that have not yet been characterized.

[0060] The 2-D GEF method of the present invention permits global analysis of gene expression, and facilitates the discovery of new genes. Furthermore, 2-D GEF allows for comparative assessments of gene expression by comparing the two-dimensional coordinates generated from one particular DNA sample, cell sample, or tissue sample with those generated by a second DNA sample, cell sample, or tissue sample. Moreover, as disclosed herein, the 2-D GEF procedure may be used as an *in vitro* diagnostic method for detecting the presence of a target cell in a tissue sample, by permitting identification of a genetic marker for the desired target cell. For example, detection of expression of KIAA0918 in a tissue sample may indicate the presence in the tissue sample of hematopoietic stem cells, in accordance with the claimed invention.

[0061] The present invention is described in the following Experimental Details section, which is set forth to aid in the understanding of the invention, and should not be construed to limit in any way the scope of the invention as defined in the claims which follow thereafter.

Experimental Details

1. Introduction

[0062] Analysis of differential gene expression is important for understanding a number of key biological processes, including proliferation, differentiation, and development. Various approaches for comparative analysis of gene expression, and for identification of differentially-expressed genes, have been proposed. Most widely-used procedures are based on subtractive hybridization [1, 2] or gel display of arbitrarily-primed RT-PCR fragments [3, 4]. Previously, the inventors developed the Gene Expression Fingerprinting (GEF) technique [5-7], which was based on the creation of non-overlapping sets of 3'-terminal cDNA restriction fragments that are displayed by a high-resolution polyacrylamide gel electrophoresis (PAGE). Different variants of cDNA restriction-fragment gel displays were later proposed by other authors [8-10]. Another approach to comparative analysis of gene expression, Serial Analysis of

Gene Expression (SAGE), is based on analysis of short, concatenated nucleotide sequence tags; it allows detailed analysis of thousands of transcripts [11]. Integration of principles of subtractive hybridization and 3'-end cDNA restriction-fragment analysis for genome-wide gene analysis also has been suggested [12]. With the development of micro-miniaturization techniques, and the availability of a large number of expressed sequence tag (EST) clones, microchip-based hybridization analysis of global gene expression patterns became possible [13].

[0063] In the present experiment, the inventors have developed a new procedure: two-dimensional Gene Expression Fingerprinting (2-D GEF). 2-D GEF builds on the previously-developed variant of the GEF procedure [6, 7] by introducing a first-dimension PAGE for additional subdivision of the primary set of cDNA into a large number of size fractions. This system offers a dramatically-increased resolution, as well as a number of other advantages which were validated in this experiment by analysis of global and differential gene expression in the closely-related Kg-1 and Kg-1a cell lines that were chosen as a model system.

2. Materials and Methods

[0064] All standard DNA and RNA manipulations were performed as previously described [14].

a. RNA isolation

[0065] Kg-1 and Kg-1a cell lines were obtained from the ATCC. Cell expansion was performed according to the manufacturer's recommendations. Total RNA from about 10^8 cells of each line was prepared using RNAgents kit (Promega), according to the manufacturer's protocol. The yield was 820 μg for Kg-1 cells and 656 μg for Kg-1a cells.

b. First- and second-strand synthesis

-24-

[0066] For the first-strand cDNA synthesis, 4.5 μ g of total RNA, 15 pmol of BioAdl#T15 primer (5' biotin-GGAATGCCTACCT15-A/G/C), 4 μ l of 5xSuperScript buffer (Gibco-BRL), 2 μ l of 100 mM DTT, and 1 μ l of dNTPs (10 mM each) were combined in a total volume of 19 μ l. The mixture was incubated at 75°C for 2 min, and at 46°C for 3 min. The reaction was performed with 1 μ l (200 units) of SuperScript II (Gibco-BRL) at 46°C for 1 h. For the second-strand cDNA synthesis, 3 μ l of dNTPs (10 mM each), 10 units of *E. coli* DNA ligase (Gibco-BRL), 40 units of *E. coli* DNA polymerase I (Gibco-BRL), 2 units of *E. coli* RNaseH (Gibco-BRL), and a second-strand synthesis buffer (Boehringer, Mannheim, Germany) were added to make a total volume of 100 μ l. This reaction mixture was incubated at 16°C for 2 h. Double-stranded cDNA was extracted with phenol, and precipitated with 1 volume of isopropanol.

c. Preparation of a primary set of 3' cDNA fragments

[0067] Double-stranded cDNA was digested with 7.5 units of NdeII (Boehringer, Mannheim, Germany) for 3 h at 37°C in a total volume of 30 μ l. M280 Streptavidin magnetic beads (Dyna) were washed with 1xWBT buffer (10 mM Tris-HCl, pH8.0; 170 mM NaCl; 1 mM EDTA; 0.1% Tween 20), treated with 100 μ g of yeast tRNA for 15 min at room temperature, and washed again. The beads were mixed with NdeII-digested cDNA in 1xWBT, incubated for 2 h at room temperature with agitation, then washed to remove unbound fragments. cDNA on the beads was ligated with 50 pmol of double-stranded adapter obtained by annealing of AD#MS/Cmod primer (5'-CGTGGGCTCCAAGCTTCAAATAAACC) and AD#Sau/Wmod primer (5'-GATCGGTTTATTTGAAGCTTGGAGCCCACG), using 6 units of *E. coli* DNA ligase (Boehringer, Mannheim, Germany) in a 30- μ l reaction at 16°C for 3 h. The beads were washed twice with 1xWBT and once with 1xWB (10 mM Tris-HCl, pH 8.0; 170 mM NaCl; 0.1 mM EDTA), and were finally suspended in 32 μ l of TE (10 mM Tris-Cl; 0.2 mM EDTA), pH 8.0.

[0068] PCR amplification of 3'-terminal cDNA fragments was performed in a total volume of 50 μ l using Advantage cDNA Polymerase Mix (Clontech), 17

-25-

pmol of BioAd1#T15 primer, and 17 pmol of AD#MS/Cmod primer, according to the manufacturer's recommendations. 3 μ l of cDNA template were amplified for 15 cycles. After initial preheating at 95°C for 35 sec, amplification was performed as follows: denaturation for 25 sec at 94°C, annealing for 45 sec at 60°C, and elongation for 2.5 min at 70°C.

d. First-dimension PAGE

[0069] Ad#MS/C primer (50 pmol) was labeled using 100 μ Ci of γ^{32} P-ATP (6000 Ci/mmol, Amersham) and 10 units of T4 polynucleotide kinase (Boehringer, Mannheim, Germany) in 20 μ l for 30 min at 37°C. Non-incorporated radioactivity was removed by passage through the Micro Bio-Spin 6 column (Bio-Rad).

[0070] For the first-dimension PAGE, 5 μ l of cDNA from the previous step were re-amplified, in a total volume of 200 μ l, by 8 cycles of PCR under conditions described above (except that labeled Ad#MS/C primer, diluted five-fold with cold primer, was used). Amplified cDNA was extracted with phenol and precipitated with isopropanol, then dissolved in 5 μ l of TE. 10 μ l of formamide stop solution were added, and DNA was denatured by incubation at 100° for 1.5 min and resolved by electrophoresis in 6% 1xTBE-7 M Urea denaturing polyacrylamide gel (acrylamide:bisacrylamide=19:1). A 50-bp ladder (Gibco-BRL) labeled with γ^{32} P-ATP was used as a size marker.

Electrophoresis was performed at 1000 V for 2.5 h.

[0071] After the run, one glass plate was removed; the gel was covered with Saran Wrap, and then exposed for up to 2 h at +5°C. Gel lanes encompassing cDNA sizes between 100 bp and 1000 bp were each cut into 96 equal slices, and placed in the consecutive wells of the 96-well PCR plate. The cDNA was eluted overnight at room temperature in TE (pH 8.0). Eluates (60 μ l) then were transferred to 8-tube PCR strips.

-26-

- e. Preparation of the samples for secondary restriction endonuclease digestion

[0072] 3 mg of Streptavidin beads (Dyna) were washed with 1xWBT, treated with 100 μ g of yeast tRNA for 15 min at room temperature in 1xWBT, washed once, and suspended in 1 ml of 6xWBT. Subsequent steps were taken with series of 64 samples (32 for each KG-1 and KG-1a cDNA sample). Washes and incubations were performed in 8-tube PCR strips using 8-channel micropipets. For washing, 100 μ l of buffer were placed in each well of the strip. Strips were placed on the 96-well magnet (Dyna), and agitation of beads was performed by hand or by transferring the strips in the neighboring columns of the magnet.

[0073] Eluted cDNA fragments were bound to 12 μ l of a Streptavidin bead suspension for 2 h at room temperature, followed by washing in 1xWBT. Streptavidin beads were incubated for 5 min in 45 μ l of denaturing buffer (100mM NaOH; 1mM EDTA), and washed in 1xWBT (three times), 1xWB, 15 μ l of 1xSequenase buffer, 0.2 mM dNTPs, and 5 mM DTT. Labeled Ad#MS/C primer (1.3 pmol) was annealed with cDNA template at 45°C for 25 min in a total volume of 8 μ l of 1xSequenase buffer. 2 μ l (3.25 units) of Sequenase version 20 (Amersham), diluted 8-fold in the Sequenase dilution buffer, were added, and the second-strand synthesis was performed for 6 min at room temperature and 15 min at 37°C. Beads were washed with 1xWBT and 1xWB.

- f. Restriction endonuclease cycle

[0074] Streptavidin beads with immobilized cDNA fragments were treated sequentially with 8 units each of the restriction enzymes listed below, in a total volume of 8 μ l. Incubation with restriction endonuclease was performed in a buffer provided by the manufacturer (New England Biolabs) at 37°C for 60 min, with agitation every 10 min. Supernatants containing liberated fragments were transferred to 96-well PCR plates, and beads were washed with 1xWBT (twice) and with 1xWB prior to the next enzyme treatment.

-27-

[0075] Restriction enzymes EcoRV, XbaI, ApaLI, NsiI, StuI, PstI, StyI, and MspI were used for the middle third of the first dimension (150-270 b), and SacII, EcoRV, ApaLI, NsiI, EcoRI, PstI, StyI, MspI, AvaII, and HinfI were used for the upper third (270-1000 b). The restriction enzymes and their order were selected on the basis of a computer simulation of the GEF procedure, the aim of which was to obtain a more even distribution of the number of fragments liberated per digestion cycle.

g. Second-dimension PAGE

[0076] Fragments released after each restriction enzyme treatment (7.5 μ l) were mixed with 1.5 μ l of formamide stop solution, denatured and concentrated by incubation at 100°C for 10 min with lids open, and resolved by electrophoresis in 6% denaturing 1xTBE-7M urea polyacrylamide gel. SequaMark™ (Research Genetics) was ³²P labeled according to the manufacturer's recommendations, and used as a size marker for the second-dimension (2-D) gels. To facilitate handling, the gel was covalently fixed to one of the glass plates by the Bind-Silane (Pharmacia) treatment. Gels were electrophoresed for 2.5 to 3 h at 60 W until the bromophenol blue dye had reached the end of the gel. The gel was soaked in 10% acetic acid for 45 min with agitation, air-dried, and exposed for 1 to 5 days at room temperature using Kodak BioMax film.

h. Recovery of the cDNA fragments from the gel

[0077] The cDNA fragments were eluted from the second-dimension gel in the elution buffer (150 mM NaCl; 10 mM Tris-HCl, pH 8.0; 0.2 mM EDTA; tRNA 100 μ g/ml) at 65°C for 2 h, precipitated with 3 vol of ethanol, and oligo(dG)-tailed using 20 units of terminal transferase (Gibco-BRL), 1.5 mM dGTP, and a buffer supplied by the manufacturer, for a total volume of 20 μ l. cDNA fragments were amplified by PCR using Advantage cDNA Polymerase Mix (Clontech), 20 pmol of C13 primer (5'-AAGGAATTCGGGGGGGGGGGG-3'), and 20 pmol of Ad#MS/Cmod primer for 40 cycles: 60 sec at 95°C, 60 sec at 60°C, and 60 sec at 70°C. The PCR fragments were purified through agarose gel, and

-28-

sequenced directly. For cloning, the PCR fragments were treated with EcoR1 and NdeII, and inserted into pBlueScript KS plasmid (Stratagene).

i. Northern blot analysis

[0078] 20 μ g of total RNA from KG-1 and KG-1a were resolved on a 1.2% formaldehyde agarose gel, transferred onto Hybond-N nylon membrane (Amersham), and probed with 32 P-labeled inserts carrying cloned cDNA fragments or corresponding ESTs. EST clones used for hybridization were obtained from Genome Systems, and their identities were verified by sequencing.

3. Results and Discussion

a. The 2-D GEF protocol

[0079] The scheme of the 2-D GEF procedure is presented in Figure 1. First-strand cDNA synthesis is performed using a one-base anchored 5'-biotinylated oligo(dT)-containing primer, followed by second-strand synthesis using the RNaseH-DNA polymerase I protocol [15]. After digestion with a four-base recognition primary restriction enzyme, 3'-terminal cDNA fragments are selected by binding to Streptavidin beads, and ligated to a double-stranded adapter. Following PCR amplification with biotinylated oligo(dT) and adapter primers, this primary set of 3'-terminal restriction fragments is separated according to size by denaturing PAGE.

[0080] For transfer to the second dimension, the gel lane containing biotinylated cDNA fragments is cut into 96 slices, which are then treated in separate wells of a microliter plate: they are eluted, bound to the Streptavidin beads, and rendered double-stranded by a Sequenase synthesis initiated from the highly-labeled adapter primer. cDNA fragments immobilized on the Streptavidin beads are treated sequentially with a set of restriction enzymes to liberate subpopulations of secondary restriction fragments. These fragments then are analyzed on separate gels by electrophoresis. Restriction enzymes are

selected in such a way as to produce relatively similar numbers of liberated fragments per each round of restriction digestion.

[0081] Two different formats of second-dimension gels have been established. In a standard 2-D GEF format, intended for the global analysis of gene expression in one cDNA sample, all fragments liberated by a given enzyme are loaded onto one gel to generate a 2-D-like pattern of bands (not shown). Another format, *quasi*-2-D GEF, is preferable for comparative analysis of gene expression in two or more cell samples (Fig. 2). In *quasi*-2-D GEF, cDNA fragments originating from the same size fraction of the first-dimension gel, but from different cell samples, are loaded side-by-side on each gel.

b. Testing the approach

[0082] The inventors have applied the 2-D GEF procedure to analyze gene expression in two closely-related human myeloid cell lines: Kg-1 and Kg-1a. The latter line is a more-undifferentiated subline of the former [16, 17]. Primary restriction digestion was performed with NdeII endonuclease, followed by sequential digestion with eight restriction enzymes for the middle portion of the first-dimension gel, and ten enzymes for the upper portion of the first-dimension gel (Table 1). The *quasi*-2-D variant was used for the second-dimension gels. The average number of bands per second-dimension (2-S) gel was found to be close to 200 for the middle portion (150-270 bp) and 570 for the upper portion (270-1000 bp) of the first-dimension gel. The analysis encompassed approximately 50% of the mRNA population (see below), and the number of fragments displayed on all 2-D gels together was 7,300 for each cell line. Therefore, estimates suggest that each of these cell lines contains about 15,000 independent mRNA species.

[0083] Inspection of half of the resulting eighteen 2-D gels (3,650 bands per cell line) revealed a total of 74 fragments, demonstrating substantial differential distribution (more than 3-fold difference in intensity) between the two cell lines. Therefore, the candidate differentially-expressed mRNAs constituted about 1% (37 vs. 3,650) of the total number of mRNAs expressed in

-30-

these cells. On the basis of the relative signal intensity, the majority of these candidate sequences represented moderately- to weakly-expressed RNAs.

Table 1. An example of the enzyme set for the secondary restriction digestion

Endonuclease	Individual, %	Overlap, %	Total, %
SacII (CCGC/GG)	3.12	0.00	3.12
EcoRV (GAT/ATC)	4.26	0.22	7.38
ApaLI (G/TGCAC)	6.37	1.09	13.75
NsiI (ATGCA/T)	7.40	2.25	21.15
EcoRI (G/AATTC)	5.96	3.21	27.11
PstI (CTGCA/G)	11.82	8.34	38.93
StyI (C/CWWGG)	14.42	19.24	53.35
MspI (C/CGG)	8.18	23.68	61.53
Avall (G/GWCC)	5.32	30.87	66.85
HinfI (G/ANTC)	9.65	43.83	76.50

Individual: the percentage of cDNA fragments predicted to be liberated by the current endonuclease; Overlap: the percentage of cDNA sequences sharing cleavage sites for the current and previous restriction enzymes; Total: the cumulative percentage of cDNA fragments cleaved after the current and previous restriction cycles

c. Verification of candidate sequences

[0084] Twenty-five candidate differentially-expressed sequences were recovered from the gel, amplified, and then directly sequenced. BLAST analysis [18] revealed 100% identity with ESTs for four cDNA fragments. Six cDNA fragments did not show any matches with the sequence database. One fragment revealed homology with a sequence belonging to an Alu repeat subfamily. The other fourteen fragments corresponded to known genes described in Table 2. Among these genes, those encoding beta and gamma chains of MHCII molecules were previously known to be expressed in Kg-1, but not in Kg-1a, cells [17]. Three fragments originated from highly-expressed ribosomal protein mRNAs. Sequence analysis indicated that, for at least two of them, this was apparently caused by procedural artifacts.

Table 2. Characterization of candidate sequences differentially expressed in Kg-1 and Kg-1a cell lines

Band #	Kg-1	Kg-1a	Homology	Reproducibility	Verification	Northern	RT-PCR
NB16	-	+	KIAA0918	Yes	Differential	Yes	N.D.
NB63	-	+	EST	N.D.	Differential	Yes	N.D.
NB64	-	+	EST	N.D.	Differential	Yes	N.D.
NB66	-	+	EST	N.D.	Differential	Yes	N.D.
NB67	+	-	MHCII gamma chain	N.D.	Differential	Yes	N.D.
NB68	+	-	ALDH1	N.D.	Differential	Yes	N.D.
NB69	-	+	No hits found	N.D.	Differential	No signal	Yes
NB70	-	+	NB101=Uncx4.1	N.D.	Differential	N.D.	N.D.
NB71	-	+	No hits found	N.D.	Differential	No signal	Yes
NB72	-	+	No hits found	N.D.	Non differential	Yes	N.D.
NB91	+	-	MHCII beta chain	Yes	Differential	Yes	N.D.
NB95	-	+	Alu repeats	Yes	N.D.	N.D.	N.D.
NB96	-	+	hKCa4	Yes	Differential	Yes	N.D.
NB97	-	+	ZEB	N.D.	Non differential	Yes	N.D.
NB101	-	+	mouse Uncx4.1*	Yes	Differential	Yes	Yes
NB103	+	-	No hits found	Yes	Differential	No signal	Yes
NB104	-	+	No hits found	Yes	N.D.	N.D.	N.D.
NB106	+	-	ribosomal protein L9	No	Non differential	Yes	N.D.
NB108	-	+	ribosomal protein L10	No	Non differential	Yes	N.D.
NB109	+	-	No hits found	N.D.	Differential	No signal	Yes
NB110	+	-	EST	N.D.	N.D.	N.D.	N.D.
NB111	+	-	ribosomal protein L27	No	Non differential	Yes	N.D.
NB112	+	-	claudin 10	Yes	Differential	Yes	N.D.
NB113	-	+	Jagged-1	Yes	Differential	Yes	N.D.
NB114	-	+	TALLA-1	Yes	Differential	Yes	N.D.

+ or - indicates relative expression level in two cell lines inferred from the 2-D GEF experiment. In the last two columns, Yes indicates that a reliably detectable signal on a Northern blot or in an RT-PCR experiment was obtained for a given sequence. N.D.=no data available; *=only part of the NB101 sequence shows high homology to mouse Uncx4.1

[0085] To verify the expression pattern of candidate sequences, Northern blot hybridization was performed with total RNA from KG-1 and KG-1a cell lines. Twelve of twenty-one candidate sequences confirmed the expected differential expression pattern (Fig. 3). Four sequences did not show a signal on Northern blots, but their expression patterns were confirmed by a more sensitive reverse-transcription PCR (RT-PCR) technique (data not shown). Five clones, including all three corresponding to ribosomal proteins, were not differentially

-32-

expressed. Therefore, the percentage of the false-positive fragments was approximately 25%. In summary, these results demonstrate the high efficiency and reliability of the 2-D GEF procedure for identification of differentially-expressed genes. These advantages obviate the need for duplicate samples commonly required for such techniques as differential mRNA display.

[0086] In comparing gene expression patterns in primitive human hematopoietic Kg-1 and Kg-1a cell lines, as described above, the inventors identified a number of genes expressed specifically in one cell line or the other. In particular, one of the genes was found to be identical to the previously-sequenced human KIAA0918 gene, a use for which was unknown prior to the present experiment. Prediction of the amino acid sequence for KIAA0918 indicates that its product belongs to a class of transmembrane, presumably cell surface, proteins. Because the gene is expressed in the more primitive Kg-1a cell line, which is close to a human hematopoietic stem cell line, this gene may be used as a genetic marker for identification of human hematopoietic stem cells.

d. Reproducibility of the 2-D GEF procedure

[0087] To rigorously test the reproducibility of the 2-D GEF procedure, the 2-D GEF experimental protocol, including the cDNA synthesis, was repeated, beginning with the same RNA preparation. The 2-D gels derived from the middle portion of the first dimension (150-270 pb) were compared for two independent experiments. Corresponding bands differing in intensity more than three-fold were ranked as non-reproducible. The reproducibility from gel to gel varied from 92 to 100% of total bands, with an average of 95.4%. It appears that a substantial part of the apparent non-reproducibility was caused by the difficulty in achieving a perfectly-identical subdivision of the first-dimension strip into size fractions for experiments separated in time. The reproducibility of fragments distributed differentially between the two cell lines was also high, with about 90% of the differential fragments reproducible between two independent experiments.

-33-

[**0088**] When cDNA synthesis and 2-D analysis were performed in parallel on duplicate samples, the reproducibility, as expected, increased markedly, reaching 99.5-100%. The results of the verification of candidate differentially-expressed sequences described above also support these data, since the reported low percentage of false positives (c.a. 25%) may only be obtained if the degree of non-reproducibility in the experiment was several-fold lower than the percentage of differentially-expressed sequences (1%).

e. Computer simulation of 2-D GEF

[**0089**] The excellent resolving power of the 2-D GEF procedure prompted the inventors to ascertain whether the identity of bands on the 2-D gels could be established on the basis of their two-dimensional coordinates. To this end, the inventors created software simulating the 2-D GEF procedure. Sequences obtained from the NCBI UniGene database were subjected to computer-simulated primary and sequential secondary digestions, to generate a subdatabase of predicted two-dimensional coordinates for all known human genes. This subdatabase can be used to establish the correspondence between known genes and bands on the 2-D gels. This procedure might be performed in two ways: (i) from the database to the gel: identification of fragments on 2-D gels which correspond to selected known genes; or (ii) from the gel to the database: identification of genes in the database which might correspond to selected fragments on the second-dimension gels.

[**0090**] Computer simulation of the GEF procedure was performed using sequence information from 7,459 known human genes. Approximately 97% of these cDNA sequences can be digested with NdeII. Approximately 5% of these NdeII-digested cDNA fragments are too short (less than 20-30 bp), and are poorly suited for analysis, since only a small percentage contains sites for secondary restriction enzymes. The primary cDNA fragments longer than 1,000 bp (c.a. 8% of total digested cDNA) are relatively inadequate for analysis, due to the reduced resolution of the first-dimension gel in this range of lengths. Therefore, prior to the secondary restriction digestion stage, at least 8%, and up

-34-

to 15%, of the cDNAs are excluded from the analysis. Similar estimates were obtained when Nla III or Mae III was used as the primary enzyme (up to 11.5% and 15.5%, respectively).

[0091] The computer analysis indicates that, during the secondary digestion stage, 10 to 15 digestion cycles should normally liberate 80-95% of cDNA fragments. Thus, the theoretical estimates show that approximately 70-85% of the total mRNA population can be analyzed by the 2-D GEF procedure, using one primary enzyme. However, for the nearly-complete (95% or more) analysis of cellular mRNAs, two different primary enzymes should be used. In the present experiment, the inventors have analyzed and resolved approximately 50% of the total mRNA population.

f. Prediction ability of the software

[0092] To explore the prediction power of the 2-D GEF software, sixteen fragments were recovered from the gels, amplified, and sequenced without prior cloning (Table 3). Eleven of the fragments corresponded to the predicted sequences. For one fragment, an incorrect prediction was made, due to the absence of the corresponding sequence (mitochondrial 16S ribosomal RNA) in the database. For four fragments, no correct matches in the database were found. Sequencing of these fragments revealed their identities as known genes. In most cases, the reason why sequences were not identified correctly related to flaws in the Unigene database, *e.g.*, sequence mistakes, an absence of complete 3' sequence, or an absence of the sequence in the database (Table 3). In only one case out of five (ribosomal protein L27), the failure to correctly identify the sequence was due to an artifact in the GEF procedure itself.

[0093] In an earlier 2-D GEF experiment, performed using a suboptimal cDNA synthesis protocol, twenty fragments with matches in the human cDNA database were analyzed. Fourteen fragments were found to correspond to the correct prediction. Two other bands had more than one match in the database, and corresponded to one of them. Three more bands arose due to a cDNA synthesis artifact which was later eliminated. Only one band did not correspond

-35-

to the predicted sequence in the database. The reason for this mistake was the lack of information on the 3'-untranslated region of the corresponding sequence in the cDNA database.

Table 3. Prediction of the identity of the cDNA fragments on the basis of their 2-D GEF coordinates

Band	Identity	Prediction	Source of the problem
NB1	16S mitochondrial RNA	wrong	DB: sequence absent
NB2	ribosomal protein S29	correct	
NB3	ribosomal protein L23	correct	
NB4	KIAA0178	correct	
NB5	KIAA0190	correct	
NB6	transl. controlled tumor protein	no match	DB: sequence error
NB91	MHC class 11 DR-beta-111	correct	
NB67	MHC class 11 gamma-chain	correct	
NB114	TALLA-1	correct	
NB97	zinc finger protein ZEB	no match	DB: 3' sequence absent
NB112	putative OSP-like protein	correct	
NB96	hKCa4	no match	DB: sequence absent
NB68	Aldehyde dehydrogenase 1	correct	
NB113	Jagged 1	correct	3rd GATC site cleaved
NB111	ribosomal protein L27	no match	
NB106	ribosomal protein L9	correct	

The second experiment is shown. Summary of results for first experiment: 70%=correct; 10%=more than one match; 15%=GEF artifacts (cause eliminated later); and 5%=database errors. Summary of results for second experiment: 69%=correct; 6%=GEF artifacts; and 25%=database errors. DB=database

g. Problems and further optimization

[0094] The major improvement in the present experiment arose from the use of the RNaseH/DNA polymerase cDNA synthesis protocol [15], which significantly increased reproducibility over that of previous GEF variants [6, 7]. However, the differing densities of fragments liberated from the lower and upper portions of the first dimension is one of the drawbacks of the 2-D GEF procedure. Separation into gel slices, with a regular increase in the linear size of the slice from top to bottom, will substantially alleviate this problem. Several artifacts were also detected and eliminated during the establishment of the procedure.

[0095] One potentially important, but so far rarely encountered, artifact is the annealing of the (T)-primer on oligo(dA)-containing stretches upstream of

-36-

the poly(A) tail during the first-strand synthesis. Such mispriming would lead to the appearance of bands not predicted by computer simulation of the 2-D GEF method, and might be a major source of non-reproducibility in the procedure. Increasing the temperature in the first-strand synthesis to 46°C reduces the significance of this artifact. Using yet more stringent conditions for annealing and synthesis in the presence of trehalose stabilizer [19] may essentially suppress this mispriming on all but the longest uninterrupted oligo(dA) stretches. These latter cases, though, would be easily recognized by software means, and might be incorporated in the computer analysis.

[0096] Another artifact detected in the present experiment is the incomplete digestion of double-stranded cDNA by the restriction enzymes, particularly in the vicinity of magnetic beads. Although the magnitude of this defect is low, it does represent a problem for highly-expressed genes, such as those encoding ribosomal proteins. In the current protocol, the significance of this artifact was substantially reduced; increasing the length of the oligo(dT)-containing primer is likely to further improve the situation.

h. Potential use of the 2-D GEF procedure

[0097] The 2-D GEF method represents a dramatic improvement over the one-dimensional GEF previously developed by the inventors [6]. One of the most significant advantages of 2-D GEF is the high resolution created by the additional subdivision of the cDNA population into approximately 100 size subsets at the first-dimension step. Therefore, it can be estimated, taking into account partial overlap, that each lane of the second dimension may contain, on average, 60- to 80-fold fewer individual cDNA sequences than in the original GEF procedure. The further 10- to 15-fold reduction of the cDNA complexity in the second dimension, due to sequential digestion, decreases the complexity of the population of cDNA fragments 600- to 1200-fold per average lane of the second-dimension gel.

[0098] Based on the complexity of the mRNA pool in an average cell type (~10,000-20,000 independent sequences per cell), the inventors have estimated

-37-

that, on average, 10 to 15 bands per lane might be expected for the middle portion, and approximately 20 to 40 bands per lane for the upper portion, of the first-dimension gel. Therefore, the 2-D GEF procedure should be able to resolve into individual cDNA bands the majority of RNAs expressed in a given cell type. Moreover, given that polyacrylamide gel can resolve 300-500 DNA fragments per lane, the 2-D GEF procedure provides the potential for the unique localization and correct identification of the larger part of the estimated 100,000 human mRNAs.

[0099] The high resolution provided by the 2-D GEF procedure also permits analysis of the cDNA fragments by direct sequencing without prior cloning. In comparison with other techniques for gel display of cDNA restriction fragments [8-10], the inventors' procedure provides the highest resolution, and can work reliably with small amounts of starting material. The present experiments have demonstrated that the method may be applied to starting material consisting of merely two thousand sorted cells (not shown). High reproducibility, even between completely independent experiments, provides the potential basis for sharing and comparing 2-D GEF data between different labs, and for establishing databases of 2-D coordinates for cDNA fragments.

[0100] One of the major features of 2-D GEF is its ability to facilitate prediction, using computer simulation, of the location on the gel of cDNA fragments. The inventors' data indicate that approximately 70% of cDNA fragments on the 2-D gel can be correctly identified, and that most failures are associated with database errors. In only 6% or less of the cases, the problems could be attributed to the method itself. Therefore, it may be assumed that further accumulation of sequence data and further improvement of the quality of cDNA databases will increase prediction power of the 2-D GEF software to approximately 90-95%. Finally, it should be noted that 2-D GEF provides a very sensitive technique for the detection of gene expression. Some of the sequences differentially expressed in Kg-1/Kg-1a cells were detected by PCR only, not by Northern blot analysis, thereby indicating that they belonged to the low-

-38-

abundance mRNA class. According to the inventors' estimates, messages expressed at a level of several copies per cell should be readily detected by the 2-D GEF procedure.

[0101] The technical developments of the last years, which culminated in the advent of cDNA microarrays [13], have provided an efficient procedure for gene expression analysis on a mass scale. However, these methods depend on cDNA/EST clone collections and existing sequence information and, by themselves, are poorly suited to the identification of novel expressed sequences. For the 2-D GEF approach, no such pre-existing information is necessary. Furthermore, with accumulation of the sequence information in the EST database for a given organism, the 2-D GEF gel patterns easily can be compared against the database to reveal the fragments likely to represent novel mRNAs that have not yet been characterized. Accordingly, the inventors suggest that the most efficient uses of the proposed technique may be gene expression analysis during the initial stages of a genome project for a given organism, as well as searches for novel expressed sequences.

[0102] The major bottleneck in the 2-D GEF technique so far is the necessity to perform routine and laborious steps of restriction-enzyme treatments on numerous samples. However, the inventors believe that it is exactly this feature which might provide the basis for automation of the procedure, thereby permitting the performance of an exhaustive global gene expression analysis for a given cell type within a matter of days or weeks.

References

1. Hedrick *et al.*, *Nature*, 308:149-153, 1984.
2. Sagerström *et al.*, *Annu. Rev. Biochem.*, 66:751-83, 1997.
3. Welsh *et al.*, *Nucleic Acids Res.*, 20:4965-70, 1992.
4. Liang and Pardee, *Science*, 257:967-71, 1992.
5. Ivanova *et al.*, *Molekuliarnaia Biologia.*, 28:1367-75, 1994.

6. Ivanova and Belyavsky, *Nucleic Acids Res.*, 23:2954-58, 1995.
7. Ivanova and Belyavsky, In *Gene Cloning and Analysis: Current Innovations*, Schaefer, B.C., ed. (Wymondham, U.K.: Horizon Scientific Press, 1997) 43-60.
8. Prashar and Weissman, *Proc. Natl. Acad. Sci. USA*, 93:659-63, 1996.
9. Kato, K., *Nucleic Acids Res.*, 23:3685-90, 1995.
10. Suzuki *et al.*, *Nucleic Acids Res.*, 24:289-94, 1996.
11. Velculescu *et al.*, *Science*, 270:484-87, 1995.
12. Wang and Rowley, *Proc. Natl. Acad. Sci. USA*, 95:11909-14, 1998.
13. Schena *et al.*, *Science*, 270:467-70, 1995.
14. Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Plainview, NY: Cold Spring Harbor Laboratory Press, 1989).
15. Gubler and Hoffman, *Gene*, 25:263-69, 1983.
16. Koeffler and Golde, *Science*, 200:1153-54, 1978.
17. Koeffler *et al.*, *Blood*, 56:263-72, 1980.
18. Altschul *et al.*, *Nucleic Acids Res.*, 25:3389-402, 1997.
19. Mizuno *et al.*, *Nucleic Acids Res.*, 27:1345-49, 1999.
20. Siminovitch *et al.*, *J. Cell. Comp. Physiol.*, 62:327-36, 1963.
21. Jursskova and Tkadlecek, *Nature*, 206:951-52, 1965.
22. Till and McCulloch, *Rad. Res.*, 14:213-22, 1961.
23. Dexter and Spooncer, *Annu. Rev. Cell Biol.*, 3:423-41, 1987.
24. Rao and Dravid, *Indian J. Exp. Biol.*, 37:1051-52, 1999.
25. Beers and Berkow, eds., *The Merck Manual of Diagnosis and Therapy*, 17th ed. (Whitehouse Station, NJ: Merck Research Laboratories, 1999) 961, 968.

-40-

26. Benedetti, F., *Tumori*, 82:S3-13, 1996.
27. Baech and Johnsen, *Stem Cells*, 18:76-86, 2000.
28. Dao and Nolte, *Leukemia*, 14:773-76, 2000.
29. Ausubel *et al.*, *Current Protocols in Molecular Biology* (New York: John Wiley and Sons, New York, 1997).

[0103] All publications mentioned hereinabove are hereby incorporated in their entireties. While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art, from a reading of the disclosure, that various changes in form and detail can be made without departing from the true scope of the invention in the appended claims.

-41-

What is claimed is:

1. An isolated nucleic acid sequence encoding KIAA0918.
2. The nucleic acid sequence of Claim 1, which is DNA, cDNA, or RNA.
3. The nucleic acid sequence of Claim 1, comprising the nucleotide sequence set forth in Figure 4.
4. The nucleic acid sequence of Claim 1, which encodes a protein comprising the amino acid sequence set forth in Figure 5.
5. An isolated nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid sequence that is complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof.
6. A purified KIAA0918 protein.
7. The protein of Claim 6, comprising the amino acid sequence set forth in Figure 5.
8. The protein of Claim 6, encoded by the nucleotide sequence set forth in Figure 4.
9. A purified protein encoded by a nucleic acid sequence that hybridizes under high stringency conditions to a second nucleic acid sequence that is complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof.

-42-

10. A method of making KIAA0918 protein, comprising the steps of:
 - (a) introducing into a host cell a nucleic acid sequence encoding KIAA0918;
 - (b) maintaining the host cell under conditions such that the nucleic acid sequence is expressed to produce KIAA0918 protein; and
 - (c) recovering the KIAA0918 protein.
11. An antibody specific for KIAA0918.
12. A method for producing an antibody specific for KIAA0918 protein, comprising the steps of:
 - (a) immunizing a mammal with KIAA0918 protein; and
 - (b) purifying antibody from a tissue of the mammal or from a hybridoma made using tissue of the mammal.
13. An antibody produced by the method of Claim 12.
14. A vector comprising a nucleic acid sequence encoding KIAA0918.
15. The vector of Claim 14, wherein the nucleic acid sequence comprises the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof.
16. The vector of Claim 14, wherein the nucleic acid sequence hybridizes under high stringency conditions to a nucleic acid sequence that is complementary to the nucleotide sequence set forth in Figure 4 or a contiguous fragment thereof.
17. A host cell transformed with the vector of Claim 14.

-43-

18. A method for detecting the presence of hematopoietic stem cells in a heterogeneous cell suspension that may contain hematopoietic stem cells, comprising detecting cells expressing KIAA0918 in the heterogeneous cell suspension.

19. The method of Claim 18, wherein the expression of KIAA0918 is detected using an agent reactive with KIAA0918.

20. The method of Claim 19, wherein the agent is labeled with a detectable marker.

21. The method of Claim 19, wherein the agent is an antibody.

22. The method of Claim 21, wherein the antibody is labeled with a detectable marker.

23. The method of Claim 18, wherein the expression of KIAA0918 is detected using at least one nucleic acid probe which hybridizes to nucleic acid encoding KIAA0918.

24. The method of Claim 23, wherein the nucleic acid probe is labeled with a detectable marker.

25. A method for isolating hematopoietic stem cells from a heterogeneous cell suspension that may contain hematopoietic stem cells, comprising the steps of:

- (a) contacting the heterogeneous cell suspension with a monoclonal antibody specific for KIAA0918; and
- (b) separating from the heterogeneous cell suspension cells to which the monoclonal antibody is bound.

26. The method of Claim 25, further comprising enriching the concentration of hematopoietic stem cells in the cell suspension before contacting the heterogeneous cell suspension with a monoclonal antibody specific for KIAA0918.

27. A method for assessing gene expression in a tissue sample, comprising the steps of:

- (a) obtaining cDNA from the tissue sample;
- (b) digesting said cDNA with a primary restriction enzyme to produce a primary set of 3'-terminal restriction fragments;
- (c) separating in the first dimension, by denaturing PAGE, said 3'-terminal restriction fragments;
- (d) sequentially treating said 3'-terminal restriction fragments with a set of secondary restriction enzymes to produce secondary restriction fragments;
- (e) separating in the second dimension, by PAGE, said secondary restriction fragments, to generate two-dimensional coordinates; and
- (f) assessing gene expression in the tissue sample by assessing said two-dimensional coordinates.

28. The method of Claim 27, wherein the secondary restriction fragments are separated in the second dimension by standard two-dimensional gene expression fingerprinting format.

29. The method of Claim 27, wherein the secondary restriction fragments are separated in the second dimension by *quasi*-two-dimensional gene expression fingerprinting.

30. The method of Claim 27, wherein gene expression is assessed using computer-assisted prediction of two-dimensional coordinates of cDNA fragments.

-45-

31. The method of Claim 27, wherein gene expression is assessed using an expressed sequence tag.






FIG. 1

**Fig. 2**

**Fig. 3**

1 tagacgcgga gcccaaggag gtaaaatgca cacttgctgc ccccagtaa ctttgaaca
 61 ggaccctac agaaaaatgc atagctggat gctgcagact ctacgcttg ctgtaacatc
 121 tctcgtcctt tegtgtgcag aaaccatcga ttattacggg gaaatctgtg acaatgcacg
 181 tccttgtgag gaaaaggacg gcattttaac tgtgagctgt gaaaaccggg ggatcatcag
 241 tctctctgaa attagccctc cccgtttccc aatctaccac ctctgttgtt ccggaacctt
 301 ttgaacctg ctctatccca atgagtttgt caattacact ggggcttcaa ttttgcactt
 361 aggtagcaat gttatccagg acattgagac cggggcttcc catgggctac ggggtttgag
 421 gagattgcat ctaaacata ataaactgga acttctgcga gatgataact tcttggctt
 481 ggagaacctg gagtacctac aggtcgatta caactacatc agcgtcattg aaccaatgc
 541 ttttgggaaa ctgcatttgt tgcagggtct tatctcaat gacaattttt tgtccagttt
 601 acccaacaat ctittccgtt ttgtgccctt aacgcacttg gacctccggg ggaaccggct
 661 gaaactctg cctacagtg ggctcttgca gcacatggat aaagtgttg agctacagct
 721 ggaggaaaaac cttggaatt gttctgtga gctgatctct cttaaaggatt ggttgacag
 781 catctctat tcagccctgg tgggggagt agttgtgag acccccttcc gcttacagg
 841 aagggaactg gacgaggtat ccaagcagga actttgccca aggagactta tttctgacta
 901 cgagatgagg ccgagagcgc cttgagcac cagggggtat ttacacacca ccccgcgctc
 961 agtgaattct gtggccactt ctctcttgc tgtttacaaa cccctttga agccccctaa
 1021 ggggactcgc caaccaaca agcccagggt gcgcccacc tctcgagcgc ccttaagga
 1081 cttggctac agcaactatg gcccagcat cgcctatcag accaaatccc cgggtccttt
 1141 ggagtgtccc accgctgct cttgcaacct gcagatctct gatctggcc tcaacgtaaa
 1201 ctgcaggag cgaagatcg agagcatcgc tgaactcgag ccaagccct acaatcccaa
 1261 gaaatgtat ctgacagaga actacatcgc tgcgtgcgc aggacagact tcttgaggc
 1321 caccgggctg gacctctgc accctgggaa taaccgcac tcgatgacc aggaccgcgc
 1381 ttcggggat ctacacac tgaggcgct ctactgaat ggcaacagga tcgagaggct
 1441 gagcccgag ttattctatg gctgcagag cctgcagtat ctctctcc agtacaactt
 1501 catcgcgag attcagctg gaactttga cccgttcca aaactccagc tgcattctt
 1561 gaataaacac ctctgcagg ceatgcctc aggcgtcttc tctgcttga cctctctcag
 1621 gctaaactg aggagtaacc acttcaact cttgccagt agtggagttt tggaccagct
 1681 gaagtcaact atccaaatc accctcatga caatccttg gattgtacct gtgacattgt
 1741 gggcatgaag ctgtgggtgg agcagctcaa agtggcgct ctatggagc aggtgatctg
 1801 taaggcgccc aaaaaatcg ctgagaccga catgcgctcc attaatcgg agctgctgtg
 1861 cctgactat tcagatgtag tagtttccac gccacaccc tctctatcc aggtccctgc
 1921 gaggaccagc gccgtgactc ctgcggtccg gttgaatagc accggggccc ccgagagctt
 1981 gggcgaggc ggaggggctg cgtcgtgccc cttgtctgtg ttaattctca gctcctgct
 2041 ggttttcatc atgtccgtct tctgtgccc cgggctcttc gtgctgtca tgaagcgag
 2101 gaagaagaac cagagcgacc acaccagcac caacaactcc gacgtgagct ctttaacat
 2161 gcagtacagc gtgtacggcg cggcgggcgg cagggcggc caccacacg cgcacgtgca
 2221 tcaccggcg cccgctgc ccaaggtgaa gacggcgcg ggccagctgt atgaatacat
 2281 cccccacca ctggccaca tgtcaaaaa cccatctac cgtctccag agggcaactc
 2341 cgtagaggat tacaagacc tgcacagact caaggtacc tacagcagca accaccactt
 2401 gcagcagcag cagcagccgc cgcgccacc gcagcagcca cagcagcgc ccccgccgca
 2461 gctgcagctg cagctgggg agggagagag cggggaaagc caccattgc ggagcccg
 2521 ctacagcgtc agcaccatc agcccgga ggacctgtg tgcgggtgc aggagccga
 2581 ccgttttac aggggcatt tagaaccaga caaactctgc tccaccacc ccccgga
 2641 tagctcccg gaataccca aattccctg cagcccgct gcttacatt tctcccaa
 2701 ctatgacct agacgcccc atcagtatt gcaccgggg gcaggggaca gcaggctacg
 2761 ggaaccggtg ctctacagcc cccgagtg tcttttga gaaccaacc ggaacgaata
 2821 tctggagtta aaagcaaac taaacttga gccggactac ctgaagtgc tggaaaaaca
 2881 gaccagctt agccagttt aaaagcaag aaactctt ggagctttg catttaaac
 2941 aaacaagcaa gcagacacac acagtgaaca catttgatta attgttgtt ttaacgttt
 3001 aggtgaagt gccttggcac gggattctc agcttcggtg gaagatacga aaagggtgtg
 3061 caatttctt taaaattac acgtgggaaa catttgtta aactgggcac atcatttct
 3121 ctcttgcgt gtggggcagg tgtgagaag ggcttaagg aggccattt gctgcggcg
 3181 tgacctgtga aaggtcacag tcattttgt agtggttga agtgctaaga atggtgatg
 3241 atggcagagc atagattcta ctctctctt ttgtctct cccctccc cgccttgc
 3301 ccactctct tctccctt ttaagccatg ggtgggtcta actggcttt gtggagaaat
 3361 tagcacacc caacttaat aggaatttg ttcttttt cgcctctct cttctctc
 3421 tccctccc tctcttca ttctttt ttgtttta aggatgtgt tgtatgcat
 3481 ctggacattt gaattaaaa aaagtattg tgatctgta aagatcacc atagatgtg
 3541 acaatcatt aaaaattac agctatatga tccataattg attagcaaa ataacttatt
 3601 gatgaatat acaaatatt tatttagca cctatttta tatgcacatt tagcattct
 3661 ctttcttca ctatttagc tatgatttg cagaggtgt acactgtat aggatctgca
 3721 ttctaaac tgacgtgta tcaggaggc attttcaatc attcaaatg tggagaatt
 3781 aatggctaaa tctttaaag ccaatgaac ccaccaatt gaactgcat ttctttta
 3841 gaaaacagag ctgattgtat ccaatgtat tttaaaaaat agggcaattg attggccat
 3901 tccgagagaa ttgttgcaa gtttgggtt ttattagaa atatttgaa gtattttat
 3961 taatgaacca aatgacatg ttcatttgac tactattga gccgatttc gattttta
 4021 ccaaacccag ttgatttgc acagatccac gtgtactggc acctcagaag accaaatcat
 4081 ggactgtaca agtctctata caatgtctt atcccttgg gcagcaagca atgatgata
 4141 tgacaacag gatatctga agatgggct actgtgtga cagtctata tgtatccag
 4201 cacatgtaatt ttttaataa gtttctgaat aaacttga taactatgc

Fig. 4

MHTCCPPVTLEQDLHRKMHSWMLQTLAFAVTSVLVS
CAETIDYYGEICDNACPCEEKDGILTVSCENRGIISLSEISPPRFPIYHLLLSGNLLN
RLYPNEFVNYTGASILHLGSNVIQDIETGAFHGLRGLRRLHLNHNKLELLRDDTFLGL
ENLEYLQVDYNYISVIEPNAFGKLHLLQVLILNDNLLSSLPNNLFRFVPLTHLDLRGN
RLKLLPYVGLLQHMDKVVELQLEENPWNCSCELISLKDWLDSISYSALVGDVVCETPF
RLHGRDLDEVSKQELCPRRLISDYEMRPQTPLSTTGYLHTTPASVNSVATSSSAVYKP
PLKPPKGTRQPNKPRVRPTSRQPSKDLGYSNYGPSIAYQTKSPVPLECPTACSCNLQI
SDLGLNVNCQERKIESIAELQPKPYNPKKMYLTENYIAVVRRTDFLEATGLDLLHLGN
NRISMIQDRAFGDLTNLRRLYLNGNRIERLSPELFYGLQSLQYLFLQYNLIREIQSGT
FDPVPNLQLLFLNNNLLQAMPSGVFSGLTLLRLNLRSNHFTSLPVSGVLDQLKSLIQI
DLHDNPWDCTCDIVGMKLWVEQLKVGVLVDEVICKAPKKFAETDMRSIKSELLCPDYS
DVVVSTPTPSSIQVPARTSAVTPAVRLNSTGAPASLGAGGGASSVPLSVLILSLLVF
IMSVFVAAGLFLVLMKRRKKNQSDHTSTNNSDVSSFNMQYSVYGGGGGTGGHPHAHVH
HRGPALPKVKTPAGHVYIYIPHLGHMCKNPIYRSREGNSVEDYKDLHELKVTYSSNH
HLQQQQQPPPPPPQPPQQPPPPQLQLQPGEERRESHHLRSPAYSVSTIEPREDLLSPV
QDADR FYRGILEPDKHCSTTPAGNSLPEYKFPKPCSPAAYTFSPNYDLRRPHQYLHPGA
GDSRLREPVLYSPPSAVFVEPNRNEYLELKAKLNVEPDYLEVLEKQTTFSQF

Fig. 5