



(86) Date de dépôt PCT/PCT Filing Date: 2017/09/14

(87) Date publication PCT/PCT Publication Date: 2018/03/22

(45) Date de délivrance/Issue Date: 2023/09/26

(85) Entrée phase nationale/National Entry: 2019/03/07

(86) N° demande PCT/PCT Application No.: US 2017/051610

(87) N° publication PCT/PCT Publication No.: 2018/053161

(30) Priorité/Priority: 2016/09/14 (US62/394,479)


(51) Cl.Int./Int.Cl. *C07D 401/12* (2006.01),
A61K 31/4184 (2006.01), *A61K 31/4439* (2006.01),
C07D 235/04 (2006.01)

(72) Inventeurs/Inventors:
TSENG, YUFENG JANE, US;
LIU, YU-LI, TW;
SUN, CHUNG-MING, TW;
...

(73) Propriétaires/Owners:
TSENG, YUFENG JANE, US;
NATIONAL TAIWAN UNIVERSITY, CN;
NATIONAL HEALTH RESEARCH INSTITUTES, TW; ...

(54) Titre : NOUVEAUX DERIVES DE BENZIMIDAZOLE SUBSTITUES UTILISES EN TANT QU'INHIBITEURS DE LA D-AMINO-ACIDE OXYDASE (DAAO)

(54) Title: NOVEL SUBSTITUTED BENZIMIDAZOLE DERIVATIVES AS D-AMINO ACID OXIDASE (DAAO) INHIBITORS



(57) Abrégé/Abstract:

Herein is described a compound of formula (I),

(see formula I)

wherein A, R_a, R_b, R_c, m, n, and X are as described herein; or a pharmaceutically acceptable salt thereof.


(72) Inventeurs(suite)/Inventors(continued): LAI, WEN-SUNG, TW; LIU, CHIH-MIN, TW; HWU, HAI-GWO, TW

(73) Propriétaires(suite)/Owners(continued): NATIONAL YANG MING CHIAO TUNG UNIVERSITY, TW

(74) Agent: BORDEN LADNER GERVAIS LLP

Abstract

Herein is described a compound of formula (I),



wherein A , R_a , R_b , R_c , m , n , and X are as described herein; or a pharmaceutically acceptable salt thereof.

NOVEL SUBSTITUTED BENZIMIDAZOLE DERIVATIVES AS D-AMINO ACID OXIDASE (DAAO) INHIBITORS

[0001] *This paragraph intentionally removed.*

Field of the Invention

[0002] The invention relates to D-amino acid oxidase (DAAO) inhibitors. Particularly, the present invention provides novel substituted benzimidazole derivatives used as DAAO inhibitors and for treatment and/or prevention of neurological disorders.

Background of the Invention

[0003] The aberrant regulatory mechanism of glutamate transmission on N-methyl-D-aspartic acid (NMDA) receptor has been reported as one of the neuropathology in schizophrenia. The receptor is a heterotetramer composed of two structure subunits of NMDA receptor 1 (NR1) and NR2. Modulation of the glycine binding site of NMDA receptor may improve the cognitive function and negative symptoms in schizophrenia. D-amino acid oxidase (DAAO) was found to be involved in the activation process of the NMDA receptor. The substrates of DAAO, especially the D-serine, may bind to the glycine site of the NMDA receptor as a co-agonist. This in turn may regulate the NMDA receptor in opening its calcium channel. D-serine has been found to inhibit the α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor-mediated current in rat hippocampus neurons. Thus, DAAO was hypothesized to be implicated in the pathogenesis of schizophrenia. As the NMDA receptor also involved in affective disorder, it is likely that inhibiting the DAAO may elevate the function of NMDA and improve both the symptoms of schizophrenia and depression affective disorder.

[0004] Known inhibitors of DAAO include benzoic acid, pyrrole-2-carboxylic acids, and indole-2-carboxylic acids. Indole derivatives and particularly certain indole-2-carboxylates have been described in the literature for treatment of neurodegenerative disease and neurotoxic injury. EP 396124 discloses indole-2-carboxylates and derivatives for treatment or management of neurotoxic injury resulting from a CNS disorder or traumatic event or in treatment or management of a neurodegenerative disease. U.S. Pat. Nos. 5,373,018; 5,374,649; 5,686,461; 5,962,496 and 6,100,289 disclose treatment of neurotoxic injury and neurodegenerative disease using indole

derivatives. WO 03/039540 disclose DAAO inhibitors, including indole-2-carboxylic acids, and methods of enhancing learning, memory and cognition as well as methods for treating neurodegenerative disorders. Patent Application No. WO/2005/089753 discloses benzisoxazole analogs and methods of treating mental disorders, such as Schizophrenia. WO/2015/168346
5 discloses a list of known known compounds as DAAO inhibitors

[0005] There is a need to develop candidate drugs having DAAO inhibitory effect to treat various neurological and physical disorder.

Summary of the Invention

[0006] The present invention pertains to a list of substituted benzimidazole derivatives used as
10 DAAO inhibitors and for treatment and/or prevention of neurological disorders.

[0007] The present invention provides a compound having the following formula (I), wherein each substituent is described herein.

[0008] The present invention also provides a pharmaceutical composition comprising a compound of the present invention.

15 [0009] The present invention also provides a method of inhibiting a DAAO comprising contacting a cell with a compound of the present invention.

[0010] The present invention also provides a method of treating or preventing the disease associated with DAAO inhibition in a subject comprising administering an effective amount of a compound of the present invention to the subject.

20 [0011] In some embodiments, the disease is symptom domains of schizophrenia and schizoaffective disorder, depression, Tourette Syndrome, Post-traumatic stress disorder (PTSD), Obsessive-compulsive disorder (OCD), analgesics, loss of memory and/or cognition associated with neurodegenerative diseases or loss of neuronal function characteristic of neurodegenerative diseases. Certain embodiments include mild cognitive impairment (MCI), Alzheimer's disease,
25 Parkinson's disease and schizophrenia.

Brief Description of the Drawing

[0012] Figure 1 shows that compared to the MK-801 group, different dosages of RS-D7, Drug 12083 and Prodrug 28095 can rescue the MK-801-induced hyperlocomotion.

[0013] Figure 2 shows that different dosages of RS-D7, Drug 12083 and Prodrug 28095 rescue
30 the anhedonia after acute MK-801 injection.

[0014] Figure 3 shows a significant reduction of PPI after acute MK-801 injections.

Detailed Description of the Invention

[0015] As used herein and in the appended claims, the singular forms "a," "and," and "the" include plural referents unless the context clearly dictates otherwise. When ranges are used herein for physical properties, such as molecular weight, or chemical properties, such as chemical formulae, all combinations and subcombinations of ranges and specific embodiments therein are intended to be included. The term "or" refers to "and/or" unless explicitly indicated to refer to alternatives only or unless the alternatives are mutually exclusive. The term "about" when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error). The term "comprising" (and related terms such as "comprise" or "comprises" or "having" or "including") is not intended to exclude that in other certain embodiments, for example, an embodiment of any composition of matter, composition, method, or process, or the like, described herein, may "consist of" or "consist essentially of" the described features.

[0016] Definitions

[0017] "Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to fifteen carbon atoms (e.g., C₁-C₁₅ alkyl). In certain embodiments, an alkyl comprises one to thirteen carbon atoms (e.g., C₁-C₁₀ alkyl). In certain embodiments, an alkyl comprises one to eight carbon atoms (e.g., C₁-C₈ alkyl). In other embodiments, an alkyl comprises one to five carbon atoms (e.g., C₁-C₆ alkyl). In other embodiments, an alkyl comprises one to four carbon atoms (e.g., C₁-C₄ alkyl). In other embodiments, an alkyl comprises one to three carbon atoms (e.g., C₁-C₃ alkyl). In other embodiments, an alkyl comprises one to two carbon atoms (e.g., C₁-C₂ alkyl). In other embodiments, an alkyl comprises one carbon atom (e.g., C₁ alkyl). In other embodiments, an alkyl comprises five to fifteen carbon atoms (e.g., C₅-C₁₅ alkyl). In other embodiments, an alkyl comprises five to eight carbon atoms (e.g., C₅-C₈ alkyl). In other embodiments, an alkyl comprises two to five carbon atoms (e.g., C₂-C₅ alkyl). In other embodiments, an alkyl comprises three to five carbon atoms (e.g., C₃-C₅ alkyl). In other embodiments, the alkyl group is selected from methyl, ethyl, 1-propyl (n-propyl), 1-methylethyl (iso-propyl), 1-butyl (n-butyl), 1-methylpropyl (sec-butyl), 2-methylpropyl (iso-butyl), 1,1-dimethylethyl (tert-butyl), 1-pentyl (n-pentyl). The alkyl is attached to the rest of the molecule by a single bond. Unless specifically stated otherwise in the specification, an alkyl group is optionally substituted by one or more of substituents.

[0018] "Alkoxy" refers to a radical bonded through an oxygen atom of the formula -O-alkyl, where alkyl is an alkyl chain as defined above.

[0019] "Alkenyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon double bond, and having from two to twelve carbon atoms. In certain embodiments, an alkenyl comprises two to eight carbon atoms. In other embodiments, an alkenyl comprises two to four carbon atoms. The alkenyl is attached to the rest of the molecule by a single bond, for example, ethenyl (i.e., vinyl), prop-1-enyl (i.e., allyl), but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. Unless specifically stated otherwise in the specification, an alkenyl group is optionally substituted by one or more of substituents.

[0020] "Alkynyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one carbon-carbon triple bond, having from two to twelve carbon atoms. In certain embodiments, an alkynyl comprises two to eight carbon atoms. In other embodiments, an alkynyl has two to four carbon atoms. The alkynyl is attached to the rest of the molecule by a single bond, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Unless specifically stated otherwise in the specification, an alkynyl group is optionally substituted by one or more of substituents.

[0021] "Aryl" refers to a radical derived from an aromatic monocyclic or multicyclic hydrocarbon ring system by removing a hydrogen atom from a ring carbon atom. The aromatic monocyclic or multicyclic hydrocarbon ring system contains only hydrogen and carbon from five to eighteen carbon atoms, where at least one of the rings in the ring system is fully unsaturated, i.e., it contains a cyclic, delocalized $(4n+2)$ pi-electron system in accordance with the Huckel theory. The ring system from which aryl groups are derived include, but are not limited to, groups such as benzene, fluorene, indane, indene, tetralin and naphthalene. Unless stated otherwise specifically in the specification, the term "aryl" or the prefix "ar-" (such as in "aralkyl") is meant to include aryl radicals optionally substituted by one or more substituents independently selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b--OR^a$, $-R^b-OC(O)-R^a$, $-R^b-OC(O)-OR^a$, $-R^b-OC(O)-N(R^a)_2$, $-R^b--$

$N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R-C(O)N(R^a)_2$, $-R^b--N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2), $-R^b--S(O).sub_tR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, each $R^{sup.b}$ is independently a direct bond or a straight or branched alkylene or alkenylene chain, and $R^{sup.c}$ is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0022] "Heteroaryl" refers to a radical derived from a 3- to 18-membered aromatic ring radical that comprises two to seventeen carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. As used herein, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, wherein at least one of the rings in the ring system is fully unsaturated, i.e., it contains a cyclic, delocalized $(4n+2)$ pi-electron system in accordance with the Huckel theory. Heteroaryl includes fused or bridged ring systems. The heteroatom(s) in the heteroaryl radical is optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heteroaryl is attached to the rest of the molecule through any atom of the ring(s). Examples of heteroaryls include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzindolyl, 1,3-benzodioxolyl, benzofuranyl, benzooxazolyl, benzo[d]thiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, benzo[b][1,4]oxazinyl, 1,4-benzodioxan-1-yl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyran-2-yl, benzopyran-4-yl, benzofuran-2-yl, benzofuran-3-yl, benzothiophen-2-yl, benzothiophen-3-yl, benzothieno[3,2-d]pyrimidinyl, benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl, cyclopenta[d]pyrimidinyl, 6,7-dihydro-5H-cyclopenta[4,5]thieno[2,3-d]pyrimidinyl, 5,6-dihydrobenzo[h]quinazolinyl, 5,6-dihydrobenzo[h]cinnolinyl, 6,7-dihydro-5H-benzo[6,7]cyclohepta[1,2-c]pyridazinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, furo[3,2-c]pyridinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyrimidinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyridazinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyridinyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indolizinyl, isoxazolyl, 5,8-methano-5,6,7,8-tetrahydroquinazolinyl, naphthyridinyl, 1,6-naphthyridinonyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 5,6,6a,7,8,9,10,10a-octahydrobenzo[h]quinazolinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl,

phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyrazolo[3,4-d]pyrimidinyl, pyridinyl, pyrido[3,2-d]pyrimidinyl, pyrido[3,4-d]pyrimidinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, quinazolinyl, quinoxalinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, 5,6,7,8-tetrahydroquinazolinyl, 5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidinyl, 6,7,8,9-tetrahydro-5H-cyclohepta[4,5]thieno[2,3-d]pyrimidinyl, 5,6,7,8-tetrahydropyrido[4,5-c]pyridazinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, thieno[2,3-d]pyrimidinyl, thieno[3,2-d]pyrimidinyl, thieno[2,3-c]pyridinyl, and thiophenyl (i.e. thienyl). Unless stated otherwise specifically in the specification, the term "heteroaryl" is meant to include heteroaryl radicals as defined above which are optionally substituted by one or more substituents selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, haloalkenyl, haloalkynyl, oxo, thioxo, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, -R^b--OR^a, -R^b--OC(O)-R^a, -R^b--OC(O)-OR^a, -R^b--OC(O)-N(R^a)₂, -R^b--N(R^a)₂, -R^b--C(O)R^a, -R^b--C(O)OR^a, -R^b--C(O)N(R^a)₂, -R^b--O-R-C(O)N(R^a)₂, -R^b--N(R^a)C(O)OR^a, -R^b--N(R^a)C(O)R^a, -R^b--N(R^a)S(O)_tR^a (where t is 1 or 2), -R^b--S(O)_tOR^a (where t is 1 or 2), -R^b--S(O)_tsub_tR^a (where t is 1 or 2) and -R^b--S(O)_tN(R^a)₂ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, each R^{sup.b} is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^{sup.c} is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0023] The term "pharmaceutically acceptable salt" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts of basic compounds encompassed within the term "pharmaceutically acceptable salt" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts of basic compounds of the present invention include, but are not limited to, the following: acetate, ascorbate, adipate, alginate, aspirate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, butyrate, camphorate, camphorsulfonate, camsylate, carbonate, chloride, clavulanate, citrate, cyclopentane propionate, diethylacetic,

digluconate, dihydrochloride, dodecylsulfanate, edetate, edisylate, estolate, esylate, ethanesulfonate, formic, fumarate, gluceptate, glucoheptanoate, gluconate, glutamate, glycerophosphate, glycollylarsanilate, hemisulfate, heptanoate, hexanoate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, 2-hydroxyethanesulfonate, hydroxynaphthoate, iodide, isonicotinic, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, methanesulfonate, mucate, 2-naphthalenesulfonate, napsylate, nicotinate, nitrate, N-methylglucamine ammonium salt, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, pectinate, persulfate, phosphate/diphosphate, pimelic, phenylpropionic, polygalacturonate, propionate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, thiocyanate, tosylate, triethiodide, trifluoroacetate, undeconate, valerate and the like. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof include, but are not limited to, salts derived from inorganic bases including aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, mangamous, potassium, sodium, zinc, and the like. Also included are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, cyclic amines, dicyclohexyl amines and basic ion-exchange resins, such as arginine, betaine, caffeine, choline, N,N-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like. Also, included are the basic nitrogen-containing groups that may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl; and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

[0024] The term "subject" includes living organisms such as humans, monkeys, cows, sheep, horses, pigs, cattle, goats, dogs, cats, mice, rats, cultured cells, and transgenic species thereof. In a preferred embodiment, the subject is a human.

[0025] The term "administering" includes routes of administration which allow the active ingredient of the invention to perform their intended function.

[0026] The term "treat" or "treatment" refers to a method of reducing the effects of a disease or condition. Treatment can also refer to a method of reducing the underlying cause of the disease or condition itself rather than just the symptoms. The treatment can be any reduction from native levels and can be, but is not limited to, the complete ablation of the disease, condition, or the symptoms of the disease or condition.


[0027] The term "prevent," "prevention" or "preventing" means inhibition or averting of symptoms associated with the target disease.

[0028] The phrase "therapeutically effective amount" refers to that amount of a compound, material, or composition comprising a compound of the present invention which is effective for producing a desired therapeutic effect, at a reasonable benefit/risk ratio applicable to any medical treatment.

[0029] The term "neurological disorder" refers to any undesirable condition of the central or peripheral nervous system of a mammal. The term "neurological disorder" includes neurodegenerative diseases (e.g., Alzheimer's disease, Parkinson's disease and amyotrophic lateral sclerosis), neuropsychiatric diseases (e.g. schizophrenia and anxieties, such as general anxiety disorder). Exemplary neurological disorders include MLS (cerebellar ataxia), Huntington's disease, Down syndrome, multi-infarct dementia, status epilepticus, contusive injuries (e.g. spinal cord injury and head injury), viral infection induced neurodegeneration, (e.g. AIDS, encephalopathies), epilepsy, benign forgetfulness, closed head injury, sleep disorders, depression (e.g., bipolar disorder), dementias, movement disorders, psychoses, alcoholism, post-traumatic stress disorder and the like. "Neurological disorder" also includes any undesirable condition associated with the disorder. For instance, a method of treating a neurodegenerative disorder includes methods of treating loss of memory and/or loss of cognition associated with a neurodegenerative disorder. Such method would also include treating or preventing loss of neuronal function characteristic of neurodegenerative disorder.


Compounds of the present invention

[0030] In one aspect, the present invention provides a compound of formula (I):



wherein n is 0 or 1,

X is $-S-$, $-S(=O)-$ or $-NR_n-$; wherein



A is $-CH-$, $-CR_c$ or N ;

R_a is $-C(=O)OR_{a1}$, $-OR_{a2}$, $-O-C(=O)R_{a3}$ or $-O-C(=O)-T-OR_{a4}$; wherein

R_{a1} is H or linear or branched C_{1-15} alkyl;

R_{a2} is H, linear or branched C_{1-15} alkyl, phosphonate, diarylphosphonate or an *O*-protecting group;

R_{a3} and R_{a4} are independently a protecting group, linear or branched C_{1-15} alkyl, linear or branched C_{2-15} alkenyl, $-T-C_{3-10}$ cycloalkyl, $-T-NHR_{a3p}$, $-T-C_{3-10}$ cycloalkenyl, $-T-C_{6-10}$ aryl, $-T-C_{5-10}$ heteroaryl, $-T-NH-C(=O)-O-C_{1-10}$ alkyl, $-T$ -adamantyl or $-C_{1-3}$ alkylene- C_{6-10} aryl where the alkylene is substituted with $-T-NHR_{a3p}$;

R_{a3p} is H or an *N*-protecting group;

R_b is H, linear or branched C_{1-15} alkyl, linear or branched C_{2-15} alkenyl, C_{1-3} alkoxy- C_{1-15} alkyl-, $-T'-C_{3-10}$ cycloalkyl, $-T'-C_{3-10}$ cycloalkenyl, $-T'-C_{6-10}$ aryl or $-T'-C_{5-10}$ heteroaryl;

R_c each is independently linear or branched C_{1-15} alkyl, linear or branched C_{1-15} alkoxyl, unprotected or protected hydroxyl group, or $-C_{1-10}$ alkylene- $Y-C_{6-10}$ heteroaryl wherein $-Y-$ is $-CH_2-$, $-NH-$, $-O-$ or $-S-$;

symbol $*$ represents the bonding position;

m is an integer from 0 to 4;

$-T-$ is absent, C_{1-3} alkylene or C_{2-3} alkenylene;

$-T'-$ is C_{1-3} alkylene or C_{2-3} alkenylene; and

wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;

wherein the alkyl, alkenyl, alkoxy, cycloalkyl, aryl, heteroaryl, alkylene and alkenylene are each independently unsubstituted or substituted with at least one substituent;

wherein the substituent is each independently a halogen, a protecting group, protected or unprotected amino group, nitro, nitroso, linear or branched C₁₋₁₅ alkyl, or linear or branched C₁₋₁₅ alkoxy or C₃₋₁₀ cycloalkyl; and

when R_b is H, the tautomers are included,


with the proviso that

when X is -S- or -S(=O)-, R_a is -OR_{a2} and R_{a2} is H or linear or branched C₁₋₁₅ alkyl, then A is -CH or -CR_c;

when X is -S- or -S(=O)- and R_a is -C(=O)OR_{a1}, R_b is linear or branched C₆₋₁₅ alkyl, linear or branched C₆₋₁₅ alkenyl, C₁₋₃ alkoxy-C₁₋₁₅ alkyl-, -T'-C₃₋₁₀ cycloalkyl, -T'-C₃₋₁₀ cycloalkenyl, -T'-C₆₋₁₀ aryl or -T'-C₅₋₁₀ heteroaryl;


or a pharmaceutically acceptable salt thereof.

[0031] In one embodiment, the present invention provides a compound of formula (I-a):



wherein n is 0 or 1,

X is -S-, -S(=O)- or -NR_n-; wherein



A is -CH-, -CR_c or N;

R_a is -C(=O)OR_{a1}, -OR_{a2}, -O-C(=O)R_{a3} or -O-C(=O)-T-OR_{a4}; wherein

R_{a1} is H or linear or branched C₁₋₁₅ alkyl;

R_{a2} is H, linear or branched C₁₋₁₅alkyl, diarylphosphonate or an *O*-protecting group;
R_{a3} and R_{a4} are independently a protecting group, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, -T-C₃₋₁₀cycloalkyl, -T-NHR_{a3p}, -T-C₃₋₁₀cycloalkenyl, -T-C₆₋₁₀aryl, -T-C₅₋₁₀heteroaryl, -T-NH-C(=O)-O-C₁₋₁₀alkyl or -T-adamantyl;

5 R_{a3p} is H or an *N*-protecting group;

R_b is H, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, C₁₋₃alkoxy-C₁₋₁₅alkyl-, -T'-C₃₋₁₀cycloalkyl, -T'-C₃₋₁₀cycloalkenyl, -T'-C₆₋₁₀aryl or -T'-C₅₋₁₀heteroaryl;

R_c each is independently linear or branched C₁₋₁₅alkyl, linear or branched C₁₋₁₅alkoxyl, unprotected or protected hydroxyl group, or -C₁₋₁₀alkylene-Y-C₆₋₁₀heteroaryl wherein -Y- is
10 -CH₂-, -NH-, -O- or -S-;

symbol * represents the bonding position;

m is an integer from 0 to 4;

-T- is absent, C₁₋₃alkylene or C₂₋₃alkenylene;

-T'- is C₁₋₃alkylene or C₂₋₃alkenylene; and

15 wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;

wherein the alkyl, alkenyl, alkoxy, cycloalkyl, aryl, heteroaryl, alkylene and alkenylene are each independently unsubstituted or substituted with at least one substituent;

wherein the substituent is each independently a halogen, a protecting group, protected or
20 unprotected amino group, nitro, nitroso, linear or branched C₁₋₁₅alkyl, linear or branched C₁₋₁₅alkoxy or C₃₋₁₀cycloalkyl and

when R_b is H, the tautomers are included,


with the proviso that

when X is -S- or -S(=O)-, R_a is -OR_{a2} and R_{a2} is H or linear or branched C₁₋₁₅alkyl, then A
25 is -CH or -CR_c;

when X is -S- or -S(=O)- and R_a is -C(=O)OR_{a1}, R_b is linear or branched C₆₋₁₅alkyl, linear or branched C₆₋₁₅alkenyl, C₁₋₃alkoxy-C₁₋₁₅alkyl-, -T'-C₃₋₁₀cycloalkyl, -T'-C₃₋₁₀cycloalkenyl, -T'-C₆₋₁₀aryl or -T'-C₅₋₁₀heteroaryl;


or a pharmaceutically acceptable salt thereof.

30 **[0032]** In one embodiment, the present invention provides a compound of formula (I-b),



wherein n is 0 or 1,

X is $-S-$, $-S(=O)-$ or $-NR_n-$;



R_n is H or

A is $-CH$, $-CR_c$ or N;

R_a is $-C(=O)OR_{a1}$, $-OR_{a2}$ or $-O-C(=O)R_{a3}$; wherein

R_{a1} is H or linear or branched C_{1-15} alkyl;

R_{a2} is H, linear or branched C_{1-15} alkyl, phosphonate, diarylphosphonate or an *O*-protecting group;

R_{a3} is $-T-NHR_{a3p}$, $-T-NH-C(=O)-O-C_{1-10}$ alkyl or $-C_{1-3}$ alkylene- C_{6-10} aryl where the alkylene is substituted with $-T-NHR_{a3p}$;

R_{a3p} is H or an *N*-protecting group;

R_b is H, linear or branched C_{1-15} alkyl, C_{1-3} alkoxy- C_{1-10} alkyl-, $-T'-C_{3-10}$ cycloalkyl, $-T'-C_{3-10}$ cycloalkenyl, $-T'-C_{6-10}$ aryl or $-T'-C_{5-10}$ heteroaryl;

R_c each is independently linear or branched C_{1-15} alkyl, linear or branched C_{1-15} alkoxyl, unprotected or protected hydroxyl group or $-C_{1-10}$ alkylene- $Y-C_{6-10}$ heteroaryl wherein $-Y-$ is $-CH_2-$, $-NH-$, $-O-$ or $-S-$;

symbol * represents the bonding position;

m is an integer from 0 to 4;

$-T-$ is absent, C_{1-3} alkylene or C_{2-3} alkenylene;

$-T'-$ is C_{1-3} alkylene; and

wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;

wherein the alkyl, alkenyl, alkoxy, cycloalkyl, aryl and heteroaryl are each independently unsubstituted or substituted with at least one substituent;

wherein the substituent is each independently a halogen, protected or unprotected amino group, nitro, nitroso, linear or branched C₁₋₁₅ alkyl, linear or branched C₁₋₁₅ alkoxy or C₃₋₁₀cycloalkyl; and

when R_b is H, the tautomers are included,

with the proviso that


when X is -S- or -S(=O)-, R_a is -OR_{a2} and R_{a2} is H or linear or branched C₁₋₁₅alkyl, then A is -CH or -CR_c;

when X is -S- or -S(=O)- and R_a is -C(=O)OR_{a1}, R_b is linear or branched C₆₋₁₅alkyl, linear or branched C₆₋₁₅alkenyl, C₁₋₃alkoxy-C₁₋₁₅alkyl-, -T'-C₃₋₁₀cycloalkyl, -T'-C₃₋₁₀cycloalkenyl, -T'-C₆₋₁₀ aryl or -T'-C₅₋₁₀heteroaryl;

or a pharmaceutically acceptable salt thereof.

[0033] In one embodiment, the present invention provides the compound of formula (I), wherein n is 0 or 1;

X is -S-, -S(=O)- or -NR_n-; wherein



R_n is H or

A is -CH₃, -CR_c or N;

R_a is -OR_{a2}, -O-C(=O)R_{a3} or -O-C(=O)-T-OR_{a4}; wherein

R_{a2} is H, linear or branched C₁₋₁₅alkyl, phosphonate, diarylphosphonate or an *O*-protecting group;

R_{a3} and R_{a4} are independently a protecting group, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, -T-C₃₋₁₀cycloalkyl, -T-NHR_{a3p}, -T-C₃₋₁₀cycloalkenyl, -T-C₆₋₁₀aryl, -T-C₅₋₁₀heteroaryl, -T-NH-C(=O)-O-C₁₋₁₀alkyl, -T-adamantyl or -C₁₋₃alkylene-C₆₋₁₀aryl where the alkylene is substituted with -T-NHR_{a3p};

R_{a3p} is H or an *N*-protecting group;

R_b is H, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, C₁₋₃alkoxy-C₁₋₁₅alkyl-, -T'-C₃₋₁₀cycloalkyl, -T'-C₃₋₁₀cycloalkenyl, -T'-C₆₋₁₀ aryl or -T'-C₅₋₁₀heteroaryl;

R_c each is independently linear or branched C₁₋₁₅alkyl, linear or branched C₁₋₁₅alkoxy, unprotected or protected hydroxyl group, or -C₁₋₁₀alkylene-Y-C₆₋₁₀heteroaryl wherein -Y- is -CH₂-, -NH-, -O- or -S-;

symbol * represents the bonding position;

5 m is an integer from 0 to 4;

-T- is absent, C₁₋₃alkylene or C₂₋₃alkenylene;

-T'- is C₁₋₃alkylene or C₂₋₃alkenylene; and

wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;

10 wherein the alkyl, alkenyl, alkoxy, cycloalkyl, aryl, heteroaryl, alkylene and alkenylene are each independently unsubstituted or substituted with at least one substituent;

wherein the substituent is each independently a halogen, a protecting group, protected or unprotected amino group, nitro, nitroso, linear or branched C₁₋₁₅ alkyl, or linear or branched C₁₋₁₅ alkoxy or C₃₋₁₀cycloalkyl; and

15 when R_b is H, the tautomers are included,

with the proviso that

when X is -S- or -S(=O)-, R_a is -OR_{a2} and R_{a2} is H or linear or branched C₁₋₁₅alkyl, then A is -CH or -CR_c;

or a pharmaceutically acceptable salt thereof.

20 **[0034]** In a further embodiment, the present invention provides the compound of formula (I), wherein

n is 0;

X is -S(=O)-;

A is N;

25 R_a is -OR_{a2}, -O-C(=O)R_{a3} or -O-C(=O)-T-OR_{a4}, wherein R_{a2} is H, linear or branched C₁₋₁₅alkyl, phosphonate, diarylphosphonate or an *O*-protecting group;

R_{a3} and R_{a4} are independently a protecting group, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, -T-C₃₋₁₀cycloalkyl, -T-NHR_{a3p}, -T-C₃₋₁₀cycloalkenyl, -T-C₆₋₁₀aryl, -T-C₅₋₁₀heteroaryl, -T-NH-C(=O)-O-C₁₋₁₀alkyl, -T-adamantyl or -C₁₋₃alkylene-C₆₋₁₀aryl where the

30 alkylene is substituted with -T-NHR_{a3p}; R_{a3p} is H or an *N*-protecting group;

R_b is H;

m is 3; and

R_c each is independently linear or branched C₁₋₁₅alkyl, linear or branched C₁₋₁₅alkoxy;
or a pharmaceutically acceptable salt thereof.

[0035] In one embodiment, n is 0.

5 [0036] In one embodiment, m is an integer from 0 to 3.

[0037] In some embodiments, R_a is -C(=O)OH, -C(=O)OC₁₋₄alkyl, H, -OR_{a2} wherein R_{a2} is H, linear or branched C₁₋₁₀alkyl or an *O*-protecting group; -O-C(=O)R_{a3} wherein R_{a3} is independently tert-butyl protecting group; linear or branched C₁₋₁₀alkyl unsubstituted or substituted by halogen, tert-butyl protecting group or protected amino group; linear or branched
10 C₂₋₁₀alkenyl; C₁₋₄alkoxy; C₃₋₁₀cycloalkyl; -C₁₋₃alkylene-C₃₋₁₀cycloalkyl; -C₃₋₁₀cycloalkenyl; -C₆₋₁₀aryl unsubstituted or substituted by C₁₋₁₀alkyl, nitro, C₁₋₁₅alkoxy or halogen; -C₅₋₁₀heteroaryl unsubstituted or substituted by C₁₋₁₀alkoxy; C₂₋₃alkenylene-C₆₋₁₀aryl wherein C₆₋₁₀aryl is unsubstituted or substituted by halogen; -C₁₋₃alkylene-NH-C(=O)-O-C₁₋₁₀alkyl; or adamantly; or -O-C(=O)-O-C₁₋₁₀alkyl.

15 [0038] In some embodiments, R_a is -O-C₁₋₁₀alkyl; -O-protecting group or -O-C(=O)R_{a3} wherein R_{a3} is a tert-butyl protecting group; adamantly; linear or branched C₁₋₁₀alkyl unsubstituted or substituted by halogen or a tert-butyl protecting group; C₁₋₄alkoxy; -C₆₋₁₀aryl unsubstituted or substituted by C₁₋₁₀alkyl, nitro, C₁₋₁₅alkoxy or halogen; C₃₋₁₀cycloalkyl; -C₃₋₁₀cycloalkenyl; linear or branched C₂₋₁₀alkenyl; -C₅₋₁₀heteroaryl; -C₁₋₃alkylene-C₃₋₁₀cycloalkyl; C₂₋₃alkenylene-C₆₋₁₀aryl
20 wherein C₆₋₁₀aryl is unsubstituted or substituted by halogen; -O-C(=O)-O-C₁₋₁₀alkyl. In some embodiments, R_a is -O-C₁₋₄alkyl, -O-tert-butyloxycarbonyl protecting group or -O-C(=O)R_{a3} wherein R_{a3} is a tert-butyl protecting group; adamantly; linear or branched C₁₋₈alkyl unsubstituted or substituted by halogen or a tert-butyl protecting group; C₁₋₄alkoxy; -phenyl unsubstituted or substituted by C₁₋₆alkyl, nitro, C₁₋₄alkoxy or halogen; C₃₋₆cycloalkyl; -C₃₋₆cycloalkenyl; linear or
25 branched C₂₋₆alkenyl; -C₅₋₆heteroaryl; -C₁₋₃alkylene-C₃₋₆cycloalkyl; C₂₋₃alkenylene-phenyl wherein phenyl is unsubstituted or substituted by halogen; -O-C(=O)-O-C₁₋₄alkyl. In some further embodiments, C₃₋₆cycloalkyl is cyclopropyl or cyclohexyl. In some further embodiments, -C₃₋₁₀cycloalkenyl is cyclohexenyl.


[0039] In some further embodiments, heteroaryl is pyrrolidinyl, pyrrolinyl, pyrazolidinyl,
30 imidazolidinyl, pyrazolinyl, imidazolynyl, pyrazolyl, imidazolyl, tetrahydrofuranyl, furanyl, dioxolanyl, tetrahydrothiophenyl, thiophenyl, oxazolyl, isoxazolyl, isothiazolyl, thiazolyl,

oxathiolanyl, piperidiny, pyridiny, piperaziny, pyridaziny, pyrimidiny, pyraziny, tetrahydropyrany, pyrany, dioxany, thianyl, thiopyrany, morpholiny, oxaziny or thiaziny. In further embodiments, heteroaryl is furany, isoxazoly or thiopheny.

[0040] In some embodiments, R_a is -OH, -COOH, -O-phosphate, -O- C_{1-6} alkyl or -O-C(=O)- C_{1-6} alkyl, -O-C(=O)- C_{1-4} alkylene-NH(Fmoc or Bocprotecting group), or -O-C(=O)-NH-C(=O)-O- C_{1-10} alkyl.

[0041] In some embodiments, R_c each is independently linear or branched C_{1-6} alkyl or linear or branched C_{1-6} alkoxy. In some embodiments, R_c each is independently halogen, linear or branched C_{1-6} alkyl, linear or branched C_{1-6} alkoxy, or - C_{1-10} alkenylene-Y- C_{6-10} heteroaryl; wherein




Y is S and C_{6-10} heteroaryl is unsubstituted or substituted by C_{1-15} alkyl (preferably C_{1-4} alkyl), C_{1-15} alkenyl (preferably C_{2-4} alkyl), C_{1-15} alkoxy (preferably C_{1-4} alkoxy), -OH, -NH₂, -NO₂ or halogen.










In a further embodiment, - C_{1-10} alkenylene-Y- C_{6-10} heteroaryl is




[0042] In some embodiments, the compound of the present invention is selected from the group consisting of:




NCTU-SUN-ID	M.W.	Chemical Structure
13001	488.56	




NCTU-SUN-ID	M.W.	Chemical Structure
		2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-6-yl 2-((tert-butoxycarbonyl)amino)acetate
26090	410.52	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-pentyl-3,4-dihydroquinazoline-7-carboxylic acid
21115	398.46	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylic acid
25030	429.17	




NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3,3-dimethylbutanoate)
28096	479.55	 (2-(((5-methoxy-4,6-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-ethoxybenzoate)
12093	472.56	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazol-5-yl (tert-butoxycarbonyl)glycinate
25016	415.16	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pentanoate)





NCTU-SUN-ID	M.W.	Chemical Structure
12130	508.43	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 6-bromohexanoate)
27077	401.48	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl isobutyrate
27079	449.52	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-enecarboxylate
26089	420.47	



NCTU-SUN-ID	M.W.	Chemical Structure
		3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylic acid
21129	453.12	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-fluorobenzoate)
25032	403.12	 (ethyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate)
26098	491.61	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-butylbenzoate)




NCTU-SUN-ID	M.W.	Chemical Structure
21127	449.14	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-methylbenzoate)
25017	480.11	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-nitrobenzoate)
12128	455.18	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-cyclopentylpropanoate)




NCTU-SUN-ID	M.W.	Chemical Structure
26071	460.53	 Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(4-methoxybenzyl)-1H-benzo[d]imidazole-5-carboxylate
11021	453.12	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-fluorobenzoate)
21118	444.22	




NCTU-SUN-ID	M.W.	Chemical Structure
		Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl-1H-benzo[d]imidazole-5-carboxylate
26070	406.44	 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylic acid
25029	415.16	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methylbutanoate)
12129	495.98	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (E)-3-(2-chlorophenyl)acrylate)





NCTU-SUN-ID	M.W.	Chemical Structure
11023	426.10	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl isoxazole-5-carboxylate)
26096	401.48	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl butyrate)
25027	413.14	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclobutanecarboxylate)
28092	399.13	




NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclopropanecarboxylate)
12088	469.64	 methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methylthio)-1-octyl-1H-benzo[d]imidazole-5-carboxylate
12127	415.5	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methylbutanoate)




NCTU-SUN-ID	M.W.	Chemical Structure
21117	434.54	 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylic acid
12084	481.61	 Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazole-5-carboxylate
21126	480.11	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-nitrobenzoate)




NCTU-SUN-ID	M.W.	Chemical Structure
26097	441.55	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohexanecarboxylate)
21110	412.49	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(3-methoxypropyl)-3,4-dihydroquinazoline-7-carboxylic acid
21116	448.57	 Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate




NCTU-SUN-ID	M.W.	Chemical Structure
21120	517.27	 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-propyl- 1H-benzo[d]imidazole-5-carboxylic acid
21121	579.28	 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl- 1H-benzo[d]imidazole-5-carboxylic acid
22138	328.43	 5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazole



NCTU-SUN-ID	M.W.	Chemical Structure
25015	415.16	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pivalate)
28094	449.52	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-phenylacetate)
28091	480.11	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-nitrobenzoate)
21130	413.14	





NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (Z)-2-methylbut-2-enoate)
21103	412.49	 Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylate
21122	373.11	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acetate)
11022	493.62	




NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) (3r,5r,7r)-adamantane-1-carboxylate)
11030	491.61	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) 4-(tert-butyl)benzoate)
21102	368.44	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-propyl-1H-benzo[d]imidazole-5-carboxylic acid
21132	431.15	 tert-butyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate




NCTU-SUN-ID	M.W.	Chemical Structure
12123	385.44	 (2-((((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acrylate)
13084	563.56	 2-((((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl diphenyl phosphate
12094	670.78	 2-((((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]-imidazole-5-yl (S)-2-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)-2-phenylacetate




NCTU-SUN-ID	M.W.	Chemical Structure
21105	583.73	 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(2-(cyclohex-1-en-1-yl)ethyl)-1H-benzo[d]imidazole-5-carboxylic acid
12124	399.46	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (Z)-but-2-enoate)
12122	425.46	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl furan-2-carboxylate)




NCTU-SUN-ID	M.W.	Chemical Structure
26072	583.69	 Methyl 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(furan-2-ylmethyl)-3,4-dihydroquinazoline-7-carboxylate
22140	506.14	 2,2'-(((2-methoxy-4-methyl-1,3-phenylene)bis(methylene))bis(sulfanediyl))bis(5-methoxy-1H-benzo[d]imidazole)




NCTU-SUN-ID	M.W.	Chemical Structure
21133	506.11	 2-(((3-(bromomethyl)-2-((tert-butyldimethylsilyl)oxy)-6-methylbenzyl)thio)-5-methoxy-1H-benzo[d]imidazole
21125	469.09	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-chlorobenzoate
27078	439.53	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-ene-1-carboxylate)
11020	465.13	




NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methoxybenzoate)
26076	396.49	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-pentyl-1H-benzo[d]imidazole-5-carboxylic acid
25031	403.12	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methoxyacetate)
21128	443.19	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl heptanoate





NCTU-SUN-ID	M.W.	Chemical Structure
27076	429.53	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl hexanoate
12083	451.58	 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylic acid
21119	430.20	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl-1H-benzo[d]imidazole-5-carboxylic acid




NCTU-SUN-ID	M.W.	Chemical Structure
21104	547.65	 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylic acid
21106	462.59	 Methyl 3-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate
26077	446.51	




NCTU-SUN-ID	M.W.	Chemical Structure
		2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(4-methoxybenzyl)-1H-benzo[d]imidazole-5-carboxylic acid
26066	410.52	 Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-pentyl-1H-benzo[d]imidazole-5-carboxylate
12125	413.49	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-methylbut-2-enoate)
26065	420.46	

NCTU-SUN-ID	M.W.	Chemical Structure
		Methyl 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate
26079	434.50	 Methyl 3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate
25028	441.08	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl thiophene-2-carboxylate)
26092	474.56	

NCTU-SUN-ID	M.W.	Chemical Structure
		Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(4-methoxybenzyl)-3,4-dihydroquinazoline-7-carboxylate
21124	435.13	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl benzoate)
28093	429.53	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-ethylbutanoate)
26091	424.55	 Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-pentyl-3,4-dihydroquinazoline-7-carboxylate

NCTU-SUN-ID	M.W.	Chemical Structure
22139	344.11	 5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)sulfinyl)-1H-benzo[d]imidazole
22141	314.1	 2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazol-5-ol
28087	449.14	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methylbenzoate)
21123	387.13	

NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl propionate)
12092	594.69	 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazol-5-yl (((9H-fluoren-9-yl)methoxy)carbonyl)glycinate
21098	382.46	 Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-propyl-1H-benzo[d]imidazole-5-carboxylate
21131	421.09	

NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-chloropropanoate)
12082	465.61	 Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate
28095	471.61	 (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3,5,5-trimethylhexanoate
11031	487.93	

NCTU-SUN-ID	M.W.	Chemical Structure
		(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-chloro-4-fluorobenzoate)

or a pharmaceutically acceptable salt thereof.

[0043] The present invention encompasses all stereoisomeric forms of the compounds of Formula I, Formula I-a and Formula I-b. Centers of asymmetry that are present in the compounds of Formula I, Formula I-a and Formula I-b can all independently of one another have (R) configuration or (S) configuration. When bonds to the chiral carbon are depicted as straight lines in the structural Formulas of the invention, it is understood that both the (R) and (S) configurations of the chiral carbon, and hence both enantiomers and mixtures thereof, are embraced within the Formula. When a particular configuration is depicted, that enantiomer (either (R) or (S), at that center) is intended. Similarly, when a compound name is recited without a chiral designation for a chiral carbon, it is understood that both the (R) and (S) configurations of the chiral carbon, and hence individual enantiomers and mixtures thereof, are embraced by the name.


[0044] The invention includes all possible enantiomers and diastereomers and mixtures of two or more stereoisomers, for example mixtures of enantiomers and/or diastereomers, in all ratios. Thus, enantiomers are a subject of the invention in enantiomerically pure form, both as levorotatory and as dextrorotatory antipodes, in the form of racemates and in the form of mixtures of the two enantiomers in all ratios. In the case of a cis/trans isomerism the invention includes both the cis form and the trans form as well as mixtures of these forms in all ratios. The preparation of individual stereoisomers can be carried out, if desired, by separation of a mixture by customary methods, for example by chromatography or crystallization, by the use of stereochemically uniform starting materials for the synthesis or by stereoselective synthesis. Optionally a derivatization can be carried out before a separation of stereoisomers. The separation of a mixture of stereoisomers can be carried out at an intermediate step during the synthesis of a compound of Formula I, Formula I-a and Formula I-b or it can be done on a final racemic product. Absolute stereochemistry may be determined by X-ray crystallography of crystalline products or crystalline

intermediates which are derivatized, if necessary, with a reagent containing a stereogenic center of known configuration. Where compounds of this invention are capable of tautomerization, all individual tautomers as well as mixtures thereof are included in the scope of this invention. The present invention includes all such isomers, as well as salts, solvates (including hydrates) and solvated salts of such racemates, enantiomers, diastereomers and tautomers and mixtures thereof.

[0045] General preparation procedures of the compounds of the present invention


[0046] The compounds of Formula (I) of the present invention are prepared according to general chemical synthetic procedures. The preparation of the embodiments of the compounds of the present invention is illustrated below.

[0047] Synthetic Scheme and Procedure for the Preparation of the compounds of the invention from RS-D7




R is $-R_{a3}$ or $-T-OR_{a4}$.


[0048] Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-26065 series




$\text{R}_b = \text{H}$, linear or branched C1-15alkyl, linear or branched C2-15alkenyl,
 C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl, -T'-C3-10cycloalkenyl,
 -T'-C6-10 aryl or -T'-C5-10heteroaryl




[0049] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-26070 series*




$\text{R}_b = \text{H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl-, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl}$




[0050] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-26079 series*




$R_b =$ H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl




[0051] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-26089 series*



R_b = H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl-, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl




[0052] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-12082 series*




R_b = H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl

[0053] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-12083 series*





R_b = H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl

[0054] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-12084*





R_b = H, linear or branched C1-15alkyl, linear or branched C2-15alkenyl, C1-3alkoxy-C1-15alkyl-, -T'-C3-10cycloalkyl, -T'-C3-10cycloalkenyl, -T'-C6-10 aryl or -T'-C5-10heteroaryl


[0055] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-12092 series*



[0056] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-22138 series*



[0057] *Synthetic Scheme and Procedure for the Preparation of NCTU-SUN-22139*



Utilities

[0058] The compounds of the invention are useful for treating or preventing any disease and/or condition, wherein modulation of D-serine levels, and/or its oxidative products, is effective in ameliorating symptoms. Inhibition of the enzyme can lead to increases in D-serine levels and a reduction in the formation of toxic D-serine oxidation products. Thus, the invention provides methods for the treatment or prevention of neurological disorders and methods of enhancing learning, memory and/or cognition. The invention also provides methods for the treatment or prevention of the disease mediated by DAAO inhibition; preferably, symptom domains of schizophrenia and schizoaffective disorder, depression, Tourette Syndrome, Post-traumatic stress disorder (PTSD), Obsessive-compulsive disorder (OCD), analgesics, loss of memory and/or cognition associated with neurodegenerative diseases or loss of neuronal function characteristic of neurodegenerative diseases. In some embodiments, the symptom domains of schizophrenia and schizoaffective disorder include negative, cognitive, depressive, positive and general psychopathology symptom domains. In another embodiment, the disease associated with DAAO inhibition is mild cognitive impairment (MCI), Alzheimer's disease, Parkinson's disease or schizophrenia. In some embodiments, the disease associated with DAAO inhibition is pain, ataxia

or convulsion. In some embodiments, the compounds of the invention can be used for treating or preventing loss of memory and/or cognition associated with neurodegenerative diseases (e.g., Alzheimer's disease and schizophrenia) and for preventing loss of neuronal function characteristic of neurodegenerative diseases. Further, methods are provided for the treatment or prevention of pain, ataxia and convulsion.

[0059] In some embodiment, the effective amount of the compound described herein ranges from about 0.5 mg/kg body weight to about 20 g/kg, about 1 mg/kg body weight to about 20 g/kg, about 2 mg/kg body weight to about 20 g/kg, about 4 mg/kg body weight to about 20 g/kg, about 6 mg/kg body weight to about 20 g/kg, about 8 mg/kg body weight to about 20 g/kg, about 10 mg/kg body weight to about 20 g/kg, about 12 mg/kg body weight to about 20 g/kg, about 14 mg/kg body weight to about 20 g/kg, about 16 mg/kg body weight to about 20 g/kg, about 0.5 mg/kg body weight to about 15 g/kg, about 0.5 mg/kg body weight to about 12 g/kg, about 0.5 mg/kg body weight to about 10 g/kg, about 0.5 mg/kg body weight to about 8 g/kg, about 0.5 mg/kg body weight to about 6 g/kg, about 2 mg/kg body weight to about 15 g/kg, about 2 mg/kg body weight to about 12 g/kg, about 2 mg/kg body weight to about 10 g/kg, about 2 mg/kg body weight to about 7 g/kg, about 2 mg/kg body weight to about 5 g/kg, about 5 mg/kg body weight to about 15 g/kg or about 5 mg/kg body weight to about 10 g/kg body weight.

Pharmaceutical Composition

[0060] Another aspect of the present invention provides pharmaceutical compositions which comprises a compound of Formula I (or a pharmaceutically acceptable salt or solvate thereof) and a pharmaceutically acceptable carrier. The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and pharmaceutically acceptable excipients.

[0061] The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (or a pharmaceutically acceptable salt or solvate thereof) as an active

ingredient, a pharmaceutically acceptable carrier and optionally other therapeutic ingredients or adjuvants. The compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

[0062] The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, troches, dragees, granules and powders, or in liquid dosage forms, such as elixirs, syrups, emulsions, dispersions, and suspensions. The active ingredient can also be administered parenterally, in sterile liquid dosage forms, such as dispersions, suspensions or solutions. Other dosages forms that can also be used to administer the active ingredient as an ointment, cream, drops, transdermal patch or powder for topical administration, as an ophthalmic solution or suspension formation, i.e., eye drops, for ocular administration, as an aerosol spray or powder composition for inhalation or intranasal administration, or as a cream, ointment, spray or suppository for rectal or vaginal administration.

[0063] For topical applications, the active ingredient or a pharmaceutical composition thereof can be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the active ingredient or a pharmaceutical composition thereof include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax, sugars such as lactose and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active ingredient or a pharmaceutical composition thereof suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0064] Depending on the particular condition, disorder or disease to be treated, additional therapeutic agents can be administered together with the active ingredient or a pharmaceutical composition thereof. Those additional agents can be administered sequentially in any order, as part of a multiple dosage regimen, from the active ingredient or a pharmaceutical composition thereof (consecutive or intermittent administration). Alternatively, those agents can be part of a single

dosage form, mixed together with the active ingredient or a pharmaceutical composition thereof (simultaneous or concurrent administration).

[0065] For oral administration, a pharmaceutical composition useful in the invention can take the form of solutions, suspensions, tablets, pills, capsules, powders, granules, semisolids, sustained release formulations, elixirs, aerosols, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various disintegrants such as starch, preferably potato or tapioca starch, and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient or a pharmaceutical composition thereof of this invention can be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

[0066] The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous, intramedullary and intraarticular injection and infusion. A pharmaceutical composition for parenteral injection can comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal injection purposes. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0067] The pharmaceutical compositions useful in the present invention can also contain adjuvants such as, but not limited to, preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents, such as for example, paraben, chlorobutanol, phenol
5 sorbic acid, and the like. It can also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

[0068] Administration by slow infusion is particularly useful when intrathecal or epidural routes are employed. A number of implantable or body-mountable pumps useful in delivering
10 compound at a regulated rate are known in the art.

[0069] Suspensions, in addition to the active compounds, can contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

[0070] For purposes of transdermal (e.g., topical) administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, are prepared.

[0071] The pharmaceutical compositions useful in the invention can also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known
20 in the art of pharmaceutical formulation and can be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0072] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the active ingredient or a pharmaceutical composition thereof with
25 suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the drugs.

[0073] Other pharmaceutically acceptable carriers include, but are not limited to, a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any
30 type, including but not limited to ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid,

potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.


[0074] Solid pharmaceutical excipients include, but are not limited to, starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients can be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

[0075] Methods of preparing various pharmaceutical compositions with a certain amount of active ingredient are known, or will be apparent in light of this disclosure, to those skilled in this art. Other suitable pharmaceutical excipients and their formulations are described in Remington's Pharmaceutical Sciences, edited by E. W. Martin, Mack Publishing Company, 19th ed. 1995.

[0076] Without further elaboration, it is believed that one skilled in the art can utilize the present invention to its fullest extent on the basis of the preceding description. The following examples are, therefore, to be construed as merely illustrative and not a limitation of the scope of the present invention in any way.

Examples

Example 1-1 NCTU-SUN-21122: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acetate)




[0077] To a solution of RS-D7 (0.1 g, 0.30 mmol) in DCM (10 mL) was added NaOH (0.90 mmol) and the reaction mixture was stirred for 5-10 minutes in the nitrogen. Then acetyl chloride

(0.60 mmol) was added at 0 °C (in the ice bath). After stirring for 5-10 minutes, the reaction was allowed to warm to room temperature and stirred further for 1 hour. The reaction was extracted with ethyl acetate and pure water. The organic layer was dried over MgSO₄, filtered and concentrated to give the reaction mixture. The reaction mixture was purified by silica-gel column chromatography to obtain the pure product.

[0078] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.44 (s, 1H), 7.09 (d, *J* = 8.7 Hz, 1H), 4.71 (s, 2H), 3.75 (s, 3H), 2.29 (s, 3H), 2.24 (s, 6H).

[0079] LRMS (ESI⁺) *m/z* : 374.1 (M+H)⁺.

Example 1-2 NCTU-SUN-21124: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl benzoate)




[0080] Except that acetyl chloride is replaced by benzyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0081] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.24 (s, 1H), 8.22 (d, *J* = 1.4 Hz, 1H), 8.18 (s, 1H), 7.78 – 7.71 (m, 2H), 7.65 – 7.59 (m, 3H), 7.27 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.74 (s, 2H), 3.76 (s, 3H), 2.26 (s, 3H), 2.25 (s, 3H).

[0082] LRMS (ESI⁺) *m/z* : 436.2 (M+H)⁺.


Example 1-3 NCTU-SUN-26096: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl butyrate)



[0083] Except that acetyl chloride is replaced by butyryl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0084] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.7 Hz, 1H), 7.43 (d, *J* = 2.0 Hz, 1H), 7.08 (dd, *J* = 8.7, 2.1 Hz, 1H), 4.72 (s, 1H), 3.74 (s, 3H), 2.60 (t, *J* = 7.3 Hz, 2H), 2.24 (s, 6H), 1.77 (h, *J* = 7.3 Hz, 3H), 1.04 (t, *J* = 7.4 Hz, 3H).


[0085] LRMS (ESI⁺) *m/z* : 402.1 (M+H)⁺.

Example 1-4 NCTU-SUN-26097:**(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl cyclohexanecarboxylate)**

[0086] Except that acetyl chloride is replaced by hexahydrobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0087] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.7 Hz, 1H), 7.42 (s, 1H), 7.07 (d, *J* = 8.7 Hz, 1H), 4.72 (s, 2H), 3.75 (s, 3H), 2.64 (t, *J* = 11.1 Hz, 1H), 2.24 (s, 7H), 1.88 – 1.76 (m, 2H), 1.69 (d, *J* = 12.1 Hz, 1H), 1.65 – 1.52 (m, 3H), 1.42 (q, *J* = 11.8 Hz, 2H).


[0088] LRMS (ESI⁺) *m/z* : 442.2 (M+H)⁺.

Example 1-5 NCTU-SUN-26098:**(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 4-butylbenzoate)**

[0089] Except that acetyl chloride is replaced by 4-Butylbenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0090] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.18 (s, 1H), 8.13 (d, *J* = 8.2 Hz, 2H), 7.75 (d, *J* = 8.8 Hz, 1H), 7.61 (s, 1H), 7.45 (d, *J* = 8.2 Hz, 2H), 7.25 (d, *J* = 8.7 Hz, 1H), 4.74 (s, 2H), 3.76 (s, 3H), 2.25 (d, *J* = 6.2 Hz, 6H), 1.66 (q, *J* = 7.7 Hz, 3H), 1.47 – 1.28 (m, 3H), 0.95 (t, *J* = 7.3 Hz, 3H).

[0091] LRMS (ESI⁺) *m/z* : 492.1 (M+H)⁺.

Example 1-6 NCTU-SUN-21127:**(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 3-methylbenzoate)**


[0092] Except that acetyl chloride is replaced by *m*-Toluoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0093] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.18 (s, 1H), 8.06 – 7.99 (m, 2H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.61 (d, *J* = 2.2 Hz, 1H), 7.56 (d, *J* = 7.5 Hz, 1H), 7.49 (t, *J* = 7.6 Hz, 1H), 7.25 (dd, *J* = 8.7, 2.2 Hz, 1H), 4.76 (d, *J* = 3.2 Hz, 2H), 3.74 (s, 3H), 2.47 (s, 3H), 2.25 (s, 3H), 2.23 (s, 3H).

[0094] LRMS (ESI⁺) *m/z* : 450.1 (M+H)⁺.

Example 1-7 NCTU-SUN-27076:

2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl hexanoate




[0095] Except that acetyl chloride is replaced by hexanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0096] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.7 Hz, 1H), 7.44 (s, 1H), 7.09 (d, *J* = 8.8 Hz, 1H), 4.72 (s, 2H), 3.74 (s, 3H), 2.62 (t, *J* = 7.4 Hz, 2H), 2.24 (s, 6H), 1.75 (p, *J* = 7.3 Hz, 2H), 1.41 (h, *J* = 7.9, 7.5 Hz, 6H), 0.94 (t, *J* = 6.7 Hz, 3H).

[0097] LRMS (ESI⁺) *m/z* : 430.2 (M+H)⁺.

Example 1-8 NCTU-SUN-27077:

2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl isobutyrate




[0098] Except that acetyl chloride is replaced by isobutyryl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[0099] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.69 (d, *J* = 8.7 Hz, 1H), 7.44 (d, *J* = 2.1 Hz, 1H), 7.08 (dd, *J* = 8.7, 2.1 Hz, 1H), 4.71 (s, 2H), 3.75 (s, 3H), 2.89 – 2.84 (m, 1H), 2.24 (d, *J* = 2.4 Hz, 6H), 1.31 (d, *J* = 7.0 Hz, 6H).

[00100] LRMS (ESI⁺) *m/z* : 402.2 (M+H)⁺.

Example 1-9 NCTU-SUN-27078:

(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-ene-1-carboxylate)




[00101] Except that acetyl chloride is replaced by cyclohex-3-enecarbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

5

[00102] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.66 (d, $J = 8.2$ Hz, 1H), 7.42 (s, 1H), 7.07 (dd, $J = 8.7, 1.8$ Hz, 1H), 5.74 (s, 2H), 4.78 (d, $J = 13.6$ Hz, 1H), 4.73 (d, $J = 13.7$ Hz, 1H), 3.70 (s, 3H), 2.96 – 2.80 (m, 2H), 2.54 – 2.30 (m, 3H), 2.22 (d, $J = 2.6$ Hz, 6H), 1.95 – 1.72 (m, 2H).

10 **[00103]** LRMS (ESI $^+$) m/z : 440.1 (M+H) $^+$.

Example 1-10 NCTU-SUN-27079: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-enecarboxylate




15 **[00104]** Except that acetyl chloride is replaced by 2-methylbenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00105] ^1H NMR (400 MHz, Acetone- d_6) δ 8.20 – 8.17 (m, 2H), 7.76 (d, $J = 8.7$ Hz, 1H), 7.63 (d, $J = 2.2$ Hz, 1H), 7.57 (td, $J = 7.5, 1.5$ Hz, 1H), 7.42 (dt, $J = 7.4, 3.4$ Hz, 2H), 7.27 (dd, $J = 8.7, 2.2$ Hz, 1H), 4.74 (s, 2H), 3.77 (s, 3H), 2.67 (s, 3H), 2.26 (s, 4H), 2.25 (s, 4H).

20 **[00106]** LRMS (ESI $^+$) m/z : 450.1 (M+H) $^+$.

Example 1-11 NCTU-SUN-28087: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methylbenzoate)




[00107] Except that acetyl chloride is replaced by 4-methylbenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00108] ^1H NMR (400 MHz, Acetone- d_6) δ 8.18 (s, 1H), 8.10 (d, J = 8.1 Hz, 2H), 7.73 (d, J = 8.7 Hz, 1H), 7.60 (s, 1H), 7.42 (d, J = 7.9 Hz, 2H), 7.24 (d, J = 7.1 Hz, 1H), 4.82 – 4.69 (m, 2H), 3.73 (s, 3H), 2.47 (s, 3H), 2.24 (s, 3H), 2.23 (s, 3H).

[00109] LRMS (ESI $^+$) m/z : 450.1 (M+H) $^+$.

Example 1-12 NCTU-SUN-28091: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-nitrobenzoate)




[00110] Except that acetyl chloride is replaced by 2-nitrobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00111] ^1H NMR (400 MHz, Acetone- d_6) δ 8.16 (d, J = 5.9 Hz, 2H), 8.10 (d, J = 7.3 Hz, 1H), 7.96 (td, J = 7.5, 1.8 Hz, 1H), 7.91 (td, J = 7.8, 1.8 Hz, 1H), 7.74 (dd, J = 8.7, 1.5 Hz, 1H), 7.61 (s, 1H), 7.24 (d, J = 8.4 Hz, 1H), 4.75 (dd, J = 13.7, 5.7 Hz, 2H), 3.69 (s, 3H), 2.20 (s, 6H).

[00112] LRMS (ESI $^+$) m/z : 481.2 (M+H) $^+$.

Example 1-13 NCTU-SUN-28092: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclopropanecarboxylate)




[00113] Except that acetyl chloride is replaced by cyclopropanecarbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00114] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.67 (d, J = 8.7 Hz, 1H), 7.45 (d, J = 2.2 Hz, 1H), 7.09 (dd, J = 8.8, 2.2 Hz, 1H), 4.73 (dd, J = 13.7, 2.5 Hz, 2H), 3.73 (s, 3H), 2.23 (s, 6H), 1.92 (dt, J = 12.5, 6.3 Hz, 1H), 1.08 (s, 2H), 1.06 (s, 2H).

[00115] LRMS (ESI $^+$) m/z : 400.2 (M+H) $^+$.

Example 1-14 **NCTU-SUN-28093:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-ethylbutanoate)**




[00116] Except that acetyl chloride is replaced by 2-ethylbutanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00117] ^1H NMR (400 MHz, Acetone- d_6) δ 8.15 (s, 1H), 7.64 (d, J = 8.7 Hz, 1H), 7.39 (d, J = 2.2 Hz, 1H), 7.05 (dd, J = 8.8, 2.2 Hz, 1H), 4.81 (dd, J = 28.8, 13.7 Hz, 2H), 3.65 (s, 3H), 2.51 (tt, J = 8.6, 5.5 Hz, 1H), 2.19 (s, 3H), 2.18 (s, 3H), 1.74 (m, 4H), 1.04 (t, 7.5 Hz, 6H).

[00118] LRMS (ESI $^+$) m/z : 430.2 (M+H) $^+$.

Example 1-15 **NCTU-SUN-28094:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-phenylacetate)**




[00119] Except that acetyl chloride is replaced by 2-phenylacetyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00120] ^1H NMR (400 MHz, Acetone- d_6) δ 8.16 (s, 1H), 7.66 (d, J = 8.8 Hz, 1H), 7.41 (m, 5H), 7.30 (m, 1H), 7.06 (dd, J = 8.8, 2.2 Hz, 1H), 4.74 (dd, J = 13.7, 19.2 Hz, 2H), 3.98 (s, 2H), 3.70 (s, 3H), 2.21 (s, 6H).

[00121] LRMS (ESI⁺) m/z : 450.2 (M+H)⁺.

Example 1-16 NCTU-SUN-28095: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 3,5,5-trimethylhexanoate




5 [00122] Except that acetyl chloride is replaced by 3,5,5-trimethylhexanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00123] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.15 (s, 1H), 7.63 (d, *J* = 8.8 Hz, 1H), 7.40 (s, 1H), 7.09 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.80 (dd, *J* = 29.0, 13.7 Hz, 2H), 3.65 (s, 3H), 2.61 (dd, *J* = 15.0, 6.2 Hz, 1H), 2.44 (dd, *J* = 15.0, 7.9 Hz, 1H), 2.19 (m, 1H), 2.19 (s, 3H), 2.18 (s, 3H), 1.43 (dd, *J* = 14.1, 4.0 Hz, 1H), 1.23 (dd, *J* = 14.1, 6.5 Hz, 1H), 1.12 (d, *J* = 6.7 Hz, 3H), 0.97 (s, 9H).

[00124] LRMS (ESI⁺) m/z : 472.1 (M+H)⁺.

Example 1-17 NCTU-SUN-28096: (2-(((5-methoxy-4,6-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 2-ethoxybenzoate)




15 [00125] Except that acetyl chloride is replaced by 2-ethoxybenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00126] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.18 (s, 1H), 7.92 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.71 (d, *J* = 8.8, 1H), 7.58 (m, 2H), 7.22 (dd, *J* = 8.8, 2.3 Hz, 1H), 7.18 (d, *J* = 8.5, 1H), 4.80 (dd, *J* = 23.0, 13.7 Hz, 2H), 4.19 (q, *J* = 7.0 Hz, 2H), 3.69 (s, 3H), 2.21 (s, 3H), 2.20 (s, 3H), 1.44 (t, *J* = 7.1 Hz, 3H)

[00127] LRMS (ESI⁺) m/z : 480.1 (M+H)⁺.

25 **Example 1-18** NCTU-SUN-21123: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl propionate)




[00128] Except that acetyl chloride is replaced by propionyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00129] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.68 (d, J = 8.8 Hz, 1H), 7.44 (s, 1H), 7.09 (d, J = 8.7 Hz, 1H), 4.73 (s, 2H), 3.74 (s, 3H), 2.64 (d, J = 7.6 Hz, 2H), 2.24 (s, 6H), 1.22 (t, J = 7.6 Hz, 3H).

[00130] LRMS (ESI $^+$) m/z : 388.2 (M+H) $^+$.

Example 1-19 NCTU-SUN-21125: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-chlorobenzoate)




[00131] Except that acetyl chloride is replaced by 4-chlorobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00132] ^1H NMR (400 MHz, Acetone- d_6) δ 8.33 – 8.12 (m, 1H), 7.76 (d, J = 8.7 Hz, 0H), 7.72 – 7.61 (m, 1H), 7.27 (dd, J = 8.7, 2.2 Hz, 0H), 4.74 (s, 1H), 3.76 (s, 1H), 2.25 (d, J = 6.00 Hz, 2H).

[00133] LRMS (ESI $^+$) m/z : 470.2 (M+H) $^+$.

Example 1-20 NCTU-SUN-21126: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-nitrobenzoate)




[00134] Except that acetyl chloride is replaced by 3-nitrobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00135] ^1H NMR (400 MHz, Acetone- d_6) δ 8.95 (t, J = 2.0 Hz, 1H), 8.64 – 8.56 (m, 2H), 8.18 (s, 1H), 7.95 (t, J = 8.0 Hz, 1H), 7.77 (d, J = 8.8 Hz, 1H), 7.69 (d, J = 2.2 Hz, 1H), 7.32 (dd, J = 8.8, 2.2 Hz, 1H), 4.76 (s, 2H), 3.74 (s, 3H), 2.24 (d, J = 7.0 Hz, 6H).

[00136] LRMS (ESI $^+$) m/z : 481.2 (M+H) $^+$.

5 **Example 1-21 NCTU-SUN-21128:** 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[d]imidazol-5-yl heptanoate




[00137] Except that acetyl chloride is replaced by heptanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

10 [00138] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.67 (s, 1H), 7.43 (d, J = 2.2 Hz, 1H), 7.08 (dd, J = 8.8, 2.1 Hz, 1H), 4.72 (s, 2H), 3.74 (s, 3H), 2.62 (s, 2H), 2.24 (s, 6H), 1.81 – 1.68 (m, 3H), 1.53 – 1.25 (m, 8H).

[00139] LRMS (ESI $^+$) m/z : 444.3 (M+H) $^+$.

15 **Example 1-22 NCTU-SUN-21129:** (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[d]imidazol-5-yl 4-fluorobenzoate)




[00140] Except that acetyl chloride is replaced by 4-fluorobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

20 [00141] ^1H NMR (400 MHz, Acetone- d_6) δ 8.30 (dd, J = 8.6, 5.6 Hz, 1H), 7.75 (d, J = 8.7 Hz, 1H), 7.38 (t, J = 8.8 Hz, 1H), 7.26 (d, J = 8.7 Hz, 1H), 2.24 (d, J = 6.5 Hz, 3H).

[00142] LRMS (ESI $^+$) m/z : 454.1 (M+H) $^+$.

Example 1-23 NCTU-SUN-21130: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[d]imidazol-5-yl (Z)-2-methylbut-2-enoate)




[00143] Except that acetyl chloride is replaced by (Z)-2-methylbut-2-en-1-yl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

5 **[00144]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.18 (s, 1H), 7.68 (s, 1H), 7.46 (d, $J = 2.2$ Hz, 1H), 7.19 – 7.01 (m, 2H), 4.72 (s, 2H), 3.75 (s, 3H), 2.24 (d, $J = 2.6$ Hz, 6H), 1.95 (s, 3H), 1.91 (d, $J = 7.2$ Hz, 3H).

[00145] LRMS (ESI $^+$) m/z : 414.2 (M+H) $^+$.

Example 1-24 NCTU-SUN-21131: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-chloropropanoate)




[00146] Except that acetyl chloride is replaced by 2-chloropropanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

15 **[00147]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.16 (s, 1H), 7.71 (d, $J = 8.7$ Hz, 1H), 7.54 – 7.40 (m, 1H), 7.13 (dd, $J = 8.7, 2.0$ Hz, 1H), 4.91 (d, $J = 6.8$ Hz, 1H), 4.75 (d, $J = 4.1$ Hz, 2H), 3.71 (s, 4H), 2.22 (s, 6H), 1.83 (d, $J = 6.8$ Hz, 4H).

[00148] LRMS (ESI $^+$) m/z : 422.1 (M+H) $^+$.

Example 1-25 NCTU-SUN-21132: tert-butyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate




[00149] Except that acetyl chloride is replaced by tert-butyl carbonochloridate, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00150] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.68 (d, J = 8.8 Hz, 1H), 7.49 (s, 1H), 7.15 (d, J = 8.9, 1H), 4.72 (s, 2H), 3.74 (s, 3H), 2.24 (s, 6H), 1.54 (s, 9H).

[00151] LRMS (ESI $^+$) m/z : 432.2 (M+H) $^+$.

Example 1-26 **NCTU-SUN-12124:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl (Z)-but-2-enoate)**




[00152] Except that acetyl chloride is replaced by (Z)-but-2-enoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00153] ^1H NMR (400 MHz, Acetone- d_6) δ 8.18 (s, 1H), 7.69 (d, J = 8.4 Hz, 1H), 7.47 (dd, J = 2.2, 0.4 Hz, 1H), 7.19 (dq, J = 15.5, 6.9 Hz, 1H), 7.11 (dd, J = 8.8, 2.2 Hz, 1H), 6.13 (dq, J = 15.5, 1.7 Hz, 1H), 4.76 – 4.67 (q, J = 13.6, 2H), 3.74 (s, 3H), 2.25 (s, 3H), 2.24 (s, 3H), 1.99 (dd, J = 6.9, 1.7 Hz, 3H).

[00154] LRMS (ESI $^+$) m/z : 400.2 (M+H) $^+$.

Example 1-27 **NCTU-SUN-12125:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 3-methylbut-2-enoate)**




[00155] Except that acetyl chloride is replaced by 3-methylbut-2-enoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00156] ^1H NMR (400 MHz, Acetone- d_6) δ 8.18 (s, 1H), 7.69 (d, J = 8.7 Hz, 1H), 7.45 (d, J = 2.0 Hz, 1H), 7.09 (dd, J = 8.7, 2.1 Hz, 1H), 5.98 (dt, J = 2.6, 1.3 Hz, 1H), 4.71 (dd, J = 13.6 Hz, 2H), 3.74 (s, 3H), 2.24 (s, 3H), 2.24 (s, 3H), 2.22 (d, J = 1.2 Hz, 3H), 2.02 (d, J = 1.3 Hz, 3H).

[00157] LRMS (ESI⁺) m/z : 414.2 (M+H)⁺.

Example 1-28 NCTU-SUN-12122: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl furan-2-carboxylate)




5 [00158] Except that acetyl chloride is replaced by furan-2-carbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00159] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.18 (s, 1H), 7.96 (dd, J = 1.8, 0.8 Hz, 1H), 7.74 (d, J = 8.8 Hz, 1H), 7.60 (d, J = 2.1 Hz, 1H), 7.51 (dd, J = 3.5, 0.8 Hz, 1H), 7.24 (dd, J = 8.8, 2.1 Hz, 1H), 6.77 (dd, J = 3.5, 1.8 Hz, 1H), 4.74 (q, J = 13.6 Hz, 2H), 3.75 (s, 3H), 2.25 (s, 3H), 2.24 (s, 3H).

[00160] LRMS (ESI⁺) m/z : 426.1 (M+H)⁺.

Example 1-29 NCTU-SUN-12123: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acrylate)




15 [00161] Except that acetyl chloride is replaced by acryloyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00162] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.71 (d, J = 8.8 Hz, 1H), 7.51 (d, J = 2.2 Hz, 1H), 7.14 (dd, J = 8.8, 2.2 Hz, 1H), 6.59 (dd, J = 17.3, 1.5 Hz, 1H), 6.42 (dd, J = 17.3, 10.4 Hz, 1H), 6.11 (dd, J = 10.4, 1.5 Hz, 1H), 4.73 (q, J = 13.6, 2H), 3.74 (s, 3H), 2.24 (s, 3H), 2.24 (s, 3H).

[00163] LRMS (ESI⁺) m/z : 386.1 (M+H)⁺.

Example 1-30 NCTU-SUN-12127: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methylbutanoate)




[00164] Except that acetyl chloride is replaced by 2-methylbutanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

5 **[00165]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.68 (d, J = 8.0 Hz, 1H), 7.42 (s, 1H), 7.07 (dd, J = 8.7, 1.9 Hz, 1H), 4.80 – 4.68 (q, J = 13.6 Hz, 2H), 3.72 (s, 3H), 2.75 – 2.63 (m, 1H), 2.23 (s, 6H), 1.89 – 1.77 (m, 1H), 1.71 – 1.60 (m, 1H), 1.29 (d, J = 7.0 Hz, 4H), 1.04 (t, J = 7.4 Hz, 3H).

[00166] LRMS (ESI $^+$) m/z : 416.1 (M+H) $^+$.

10 **Example 1-31** NCTU-SUN-12128: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-cyclopentylpropanoate)




[00167] Except that acetyl chloride is replaced by cyclopentanecarbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

15 **[00168]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.68 (d, J = 8.7 Hz, 1H), 7.43 (d, J = 2.1 Hz, 1H), 7.08 (dd, J = 8.8, 2.2 Hz, 1H), 4.78 – 4.66 (q, J = 13.6, 2H), 3.74 (s, 3H), 2.66 – 2.61 (m, 2H), 2.24 (s, 6H), 1.94 – 1.81 (m, 3H), 1.77 (dd, J = 14.9, 7.4 Hz, 2H), 1.69 – 1.51 (m, 5H), 1.23 – 1.13 (m, 2H).

20 **[00169]** LRMS (ESI $^+$) m/z : 456.1 (M+H) $^+$.

Example 1-32 NCTU-SUN-12129: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (E)-3-(2-chlorophenyl)acrylate)




[00170] Except that acetyl chloride is replaced by 2-chlorobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00171] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.26 (d, *J* = 16.0 Hz, 1H), 8.18 (s, 1H), 8.02 (dd, *J* = 7.6, 2.0 Hz, 1H), 7.72 (d, *J* = 7.8 Hz, 1H), 7.58 – 7.54 (m, 2H), 7.48 (m, 2H), 7.20 (dd, *J* = 8.7, 2.0 Hz, 1H), 6.88 (d, *J* = 16.0 Hz, 1H), 4.81 – 4.70 (q, *J* = 13.6, 2H), 3.73 (s, 3H), 2.24 (s, 3H), 2.23 (s, 3H).

[00172] LRMS (ESI⁺) *m/z* : 496.0 (M+H)⁺.

Example 1-33 **NCTU-SUN-12130:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 6-bromohexanoate)**




[00173] Except that acetyl chloride is replaced by 6-bromohexanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00174] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.5 Hz, 1H), 7.44 (d, *J* = 2.3 Hz, 1H), 7.09 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.77 – 4.68 (q, *J* = 13.6, 2H), 3.74 (s, 3H), 3.55 (t, *J* = 6.7 Hz, 2H), 2.66 (t, *J* = 7.4 Hz, 2H), 2.35 (s, 6H) 1.99 – 1.91 (m, 2H), 1.84-1.76 (m, 2H), 1.66 – 1.56 (m, 2H).

[00175] LRMS (ESI⁺) *m/z* : 508.1 (M+H)⁺.

Example 1-34 **NCTU-SUN-11021:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 2-fluorobenzoate)**




[00176] Except that acetyl chloride is replaced by 2-fluorobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00177] ^1H NMR (400 MHz, Acetone- d_6) δ 8.24 – 8.13 (m, 2H), 7.78 (dddd, J = 8.4, 7.4, 4.9, 1.8 Hz, 2H), 7.64 (d, J = 2.3 Hz, 1H), 7.48 – 7.32 (m, 2H), 7.28 (dd, J = 8.8, 2.2 Hz, 1H), 4.74 (s, 3H), 3.76 (s, 3H), 2.26 (s, 3H), 2.25 (s, 3H).

[00178] LRMS (ESI $^+$) m/z : 454.1 (M+H) $^+$.

5 **Example 1-35** NCTU-SUN-11020: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methoxybenzoate)




10 [00179] Except that acetyl chloride is replaced by 4-methoxybenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00180] ^1H NMR (400 MHz, Acetone- d_6) δ 8.76-7.84 (m, 3H), 7.74 (dd, J = 8.8, 0.6 Hz, 1H), 7.59 (dd, J = 2.2, 0.5 Hz, 1H), 7.23 (dd, J = 8.7, 2.2 Hz, 1H), 7.13 (d, J = 8.9 Hz, 2H), 4.74 (s, 2H), 3.94 (s, 3H), 3.75 (s, 3H), 2.25 (s, 3H), 2.24 (s, 3H).

[00181] LRMS (ESI $^+$) m/z : 466.2 (M+H) $^+$.

15 **Example 1-36** NCTU-SUN-11022: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (3r,5r,7r)-adamantane-1-carboxylate)




20 [00182] Except that acetyl chloride is replaced by adamantane-1-carbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00183] ^1H NMR (400 MHz, Acetone- d_6) δ 8.16 (s, 1H), 7.64 (d, J = 8.6 Hz, 1H), 7.37 (d, J = 2.2 Hz, 1H), 7.02 (dd, J = 8.8, 2.2 Hz, 1H), 4.83 – 4.71 (m, 2H), 3.69 (s, 3H), 2.22 (s, 3H), 2.21 (s, 3H), 2.20-1.80 (m, 15H)

[00184] LRMS (ESI $^+$) m/z : 494.2 (M+H) $^+$.

Example 1-37 **NCTU-SUN-11023:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl isoxazole-5-carboxylate)**




[00185] Except that acetyl chloride is replaced by isoxazole-5-carbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00186] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.76 (d, *J* = 1.8 Hz, 1H), 8.17 (s, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.67 (d, *J* = 2.2 Hz, 1H), 7.39 (d, *J* = 1.8 Hz, 1H), 7.30 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.86 – 4.72 (m, 2H), 3.70 (s, 3H), 2.22 (s, 3H), 2.21 (s, 3H).

[00187] LRMS (ESI⁺) *m/z* : 427.0 (M+H)⁺.

Example 1-38 **NCTU-SUN-11030:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 4-(tert-butyl)benzoate)**




[00188] Except that acetyl chloride is replaced by 4-(tert-butyl)benzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00189] ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.14 (d, *J* = 8.5 Hz, 2H), 8.11 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 1H), 7.62 (d, *J* = 8.5 Hz, 2H), 7.51 (d, *J* = 2.2 Hz, 1H), 7.21 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.78 (d, *J* = 9.4 Hz, 2H), 3.69 (s, 3H), 2.24 (s, 3H), 2.18 (s, 3H), 1.39 (s, 9H).

[00190] LRMS (ESI⁺) *m/z* : 492.1 (M+H)⁺.

Example 1-39 **NCTU-SUN-11031:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 3-chloro-4-fluorobenzoate)**




[00191] Except that acetyl chloride is replaced by 4-chloro-3-fluorobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

5 **[00192]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.34 (dd, $J = 7.2, 2.2$ Hz, 1H), 8.24 (ddd, $J = 8.7, 4.7, 2.2$ Hz, 1H), 8.18 (t, $J = 0.8$ Hz, 1H), 7.76 (dd, $J = 8.8, 0.6$ Hz, 1H), 7.65 (dd, $J = 2.3, 0.6$ Hz, 1H), 7.58 (t, $J = 8.8$ Hz, 1H), 7.28 (dd, $J = 8.8, 2.2$ Hz, 1H), 4.74 (s, 2H), 3.76 (s, 2H), 2.26 (s, 3H), 2.24 (s, 3H).

[00193] LRMS (ESI $^+$) m/z : 488.0 (M+H) $^+$.

10 **Example 1-40** NCTU-SUN-25015: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pivalate)




[00194] Except that acetyl chloride is replaced by pivaloyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

15 **[00195]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.69 (d, $J = 8.7$ Hz, 1H), 7.42 (d, $J = 2.0$ Hz, 1H), 7.07 (dd, $J = 8.7, 2.1$ Hz, 1H), 5.62 (s, 1H), 4.71 (s, 2H), 3.75 (s, 3H), 2.24 (d, $J = 2.6$ Hz, 6H), 1.37 (s, 9H).

[00196] LRMS (ESI $^+$) m/z : 416.1 (M+H) $^+$.

20 **Example 1-41** NCTU-SUN-25016: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pentanoate)




[00197] Except that acetyl chloride is replaced by pentanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00198] ^1H NMR (400 MHz, Acetone- d_6) δ 8.16 (s, 1H), 7.65 (d, J = 8.7 Hz, 1H), 7.41 (d, J = 2.1 Hz, 1H), 7.07 (dd, J = 8.9, 2.1 Hz, 1H), 5.62 (s, 1H), 4.96 – 4.55 (m, 2H), 3.69 (s, 3H), 2.62 (t, J = 7.5 Hz, 2H), 2.21 (d, J = 3.3 Hz, 6H), 1.83 – 1.64 (m, 2H), 1.46 (q, J = 7.4 Hz, 2H), 0.97 (t, J = 7.4 Hz, 3H).

[00199] LRMS (ESI $^+$) m/z : 416.1 (M+H) $^+$.

Example 1-42 **NCTU-SUN-25017:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-nitrobenzoate)**




[00200] Except that acetyl chloride is replaced by 4-nitrobenzoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00201] ^1H NMR (400 MHz, Acetone- d_6) δ 8.48 (d, J = 2.3 Hz, 4H), 8.18 (s, 1H), 7.78 (d, J = 8.8 Hz, 1H), 7.70 (d, J = 2.2 Hz, 1H), 7.33 (dd, J = 8.8, 2.3 Hz, 1H), 4.74 (s, 2H), 3.76 (s, 3H), 2.25 (d, J = 7.2 Hz, 6H).

[00202] LRMS (ESI $^+$) m/z : 481.2 (M+H) $^+$.

Example 1-43 **NCTU-SUN-25027:** **(2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclobutanecarboxylate)**




[00203] Except that acetyl chloride is replaced by cyclobutanecarbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00204] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.69 (d, J = 8.8 Hz, 1H), 7.44 (d, J = 2.2 Hz, 1H), 7.09 (dd, J = 8.8, 2.2 Hz, 1H), 4.71 (s, 2H), 3.75 (s, 3H), 3.48 (t, J = 8.6 Hz, 1H), 2.39 (dt, J = 29.5, 9.1 Hz, 4H), 2.24 (d, J = 2.9 Hz, 5H), 2.05 (m, J = 2.4 Hz, 2H).

[00205] LRMS (ESI⁺) m/z : 414.2 (M+H)⁺.

Example 1-44 NCTU-SUN-25028: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl thiophene-2-carboxylate)




5 [00206] Except that acetyl chloride is replaced by thiophene-2-carbonyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00207] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.18 (s, 1H), 8.04 (dd, *J* = 3.8, 1.4 Hz, 1H), 7.99 (dd, *J* = 5.0, 1.4 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.62 (d, *J* = 2.1 Hz, 1H), 7.32 (dd, *J* = 5.1, 3.7 Hz, 1H), 7.25 (dd, *J* = 8.8, 2.2 Hz, 1H), 4.75 (d, *J* = 2.1 Hz, 2H), 3.74 (d, *J* = 1.7 Hz, 3H), 2.24 (d, *J* = 5.4 Hz, 6H).

[00208] LRMS (ESI⁺) m/z : 442.2 (M+H)⁺.

Example 1-45 NCTU-SUN-25029: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 2-methylbutanoate)




15 [00209] Except that acetyl chloride is replaced by 2-methylbutanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

20 [00210] ¹H NMR (400 MHz, Acetone-*d*₆) δ 8.17 (s, 1H), 7.68 (d, *J* = 8.7 Hz, 1H), 7.49 – 7.32 (m, 1H), 7.07 (ddd, *J* = 8.7, 2.3, 1.0 Hz, 1H), 4.76 – 4.52 (m, 2H), 3.74 (d, *J* = 1.0 Hz, 3H), 2.69 (q, *J* = 7.1 Hz, 1H), 2.24 (s, 6H), 1.92 – 1.75 (m, 1H), 1.65 (dddd, *J* = 13.7, 7.4, 6.3, 1.1 Hz, 1H), 1.29 (dd, *J* = 7.0, 1.0 Hz, 3H), 1.04 (td, *J* = 7.4, 1.0 Hz, 3H).

[00211] LRMS (ESI⁺) m/z : 416.1 (M+H)⁺.

Example 1-46 NCTU-SUN-25030: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1*H*-benzo[*d*]imidazol-5-yl 3,3-dimethylbutanoate)




[00212] Except that acetyl chloride is replaced by 3,3-dimethylbutanoyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

5 **[00213]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.69 (dd, $J = 8.7, 0.6$ Hz, 1H), 7.43 (dd, $J = 2.2, 0.6$ Hz, 1H), 7.08 (dd, $J = 8.7, 2.2$ Hz, 1H), 4.72 (d, $J = 1.6$ Hz, 2H), 3.74 (s, 3H), 2.50 (s, 2H), 2.24 (d, $J = 1.3$ Hz, 6H), 1.15 (s, 9H).

[00214] LRMS (ESI $^+$) m/z : 430.1 (M+H) $^+$.

Example 1-47 NCTU-SUN-25031: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) 2-methoxyacetate)




[00215] Except that acetyl chloride is replaced by 2-methoxyacetyl chloride, the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

15 **[00216]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.70 (d, $J = 8.7$ Hz, 1H), 7.50 (s, 1H), 7.14 (d, $J = 8.8$ Hz, 1H), 4.72 (s, 2H), 4.36 (s, 2H), 3.75 (s, 3H), 3.49 (d, $J = 1.2$ Hz, 2H), 2.24 (d, $J = 2.5$ Hz, 6H).

[00217] LRMS (ESI $^+$) m/z : 404.0 (M+H) $^+$.

Example 1-48 NCTU-SUN-25032 (ethyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate)




[00218] Except that acetyl chloride is replaced by ethyl carbonochloridate the other reactants and preparation steps are similar to those described in Example 1 to afford the title compound.

[00219] ^1H NMR (400 MHz, Acetone- d_6) δ 8.17 (s, 1H), 7.69 (d, J = 8.8 Hz, 1H), 7.53 (d, J = 2.2 Hz, 1H), 7.18 (dd, J = 8.8, 2.2 Hz, 1H), 4.74 (d, J = 2.6 Hz, 2H), 4.30 (q, J = 7.1 Hz, 2H), 3.73 (s, 3H), 2.23 (s, 7H), 1.35 (t, J = 7.1 Hz, 4H).

[00220] LRMS (ESI $^+$) m/z : 404.0 (M+H) $^+$.

Example 2-1 26065: Methyl 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate



[00221] To a solution of 4-fluoro-3-nitrobenzoic acid **1** (5.0 g, 27.0 mmol) in dry MeOH (30 mL), H_2SO_4 (5 mL, 0.3 M) was added and the reaction mixture was heated to reflux for 12 h. The solvent was removed under reduced pressure, crude reaction mixture was dissolved in EtOAc (150 mL), washed with saturated NaHCO_3 (20 mL $\times$ 2), water (10 mL $\times$ 2) and brine (10 mL). The EtOAc layer was dried over anhydrous MgSO_4 and evaporated to get methyl 4-fluoro-3-nitrobenzoate **2** (95%) as a white solid.

[00222] Compound **2** (2.0 g, 10.2 mmol) and 2-Aminomethylfuran (3 equiv.) in dry CH_2Cl_2 (50 mL) were stirred at room temperature for 2 h. Upon completion of reaction the solvent was removed and the crude product was purified by flash column chromatography to afford nitro benzoates **3** (90%).

[00223] To a solution of compound **3** (2.0 g, 4.8 mmol) in dry MeOH (100 mL), zinc dust (15 equiv., 71.4 mmol) and ammonium formate (7.5 equiv., 35.7 mmol) were added and the resulting reaction mixture was stirred at room temperature for 30 min. Upon completion of reaction, Zn dust was filtered through a bed of celite, filtrate was evaporated and the product was dissolved in CH_2Cl_2 (100 mL). The precipitated ammonium formate was filtered off and the solvent was evaporated to furnish compound **4** (92 %).


[00224] Use DCM to dissolve compound **4** (1.0 g, 4.0 mmol) then add 1.2 equiv. CNBr to react at room temperature. After 8 hours the mixture can be extracted with DCM and water. The solvent was removed and the crude product was purified by flash column chromatography to afford **5** (60%)

5 [00225] To a solution of methyl 2-amino-1-(furan-2-ylmethyl)-1H-benzo[d]imidazole-5-carboxylate **5** (0.05 g, 0.18 mmol) in acetonitrile (10 mL) was added K_2CO_3 (0.0497 g, 0.36 mmol) and KI (0.0089 g, 0.054 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** (0.041 g, 0.22 mmol) and the reaction mixture was allowed to reflux for six hour. After 6 h, the solvent was evaporated and the reaction mixture was diluted with saturated aq. $NaHCO_3$ (10 mL) and extracted with EtOAc (3* 10 mL).

[00226] The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 8 % methanol/ EtOAc to obtain the pure product **NCTU-SUN-26065** as a white solid 0.053 g (71 %).

15 [00227] ^1H NMR (400 MHz, Chloroform-*d*) δ 8.34 (s, 1H), 7.83 (dd, J = 8.3, 1.4 Hz, 1H), 7.73 (d, J = 1.2 Hz, 1H), 7.34 (dd, J = 1.8, 0.7 Hz, 1H), 7.13 (d, J = 8.3 Hz, 1H), 6.37 (d, J = 3.3 Hz, 1H), 6.32 (dd, J = 3.2, 1.9 Hz, 1H), 5.42 (s, 2H), 5.10 (s, 2H), 3.89 (s, 3H), 3.87 (s, 3H), 2.33 (s, 3H), 2.30 (s, 3H); ^{13}C NMR (101 MHz, Chloroform-*d*) δ 166.75 , 153.88 , 142.73 , 133.38 , 128.36 , 124.86 , 124.00 , 110.60 , 109.61 , 109.04 , 108.26 , 77.22 , 61.46 , 52.24 , 38.50 , 31.90 , 29.67 , 29.33 , 22.66 , 14.64 , 14.09 , 11.39; LRMS (ESI+) : m/z 422.3 (M+H) $^+$.

20 **Example 2-2 21098: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)- 1-propyl-1H-benzo[d]imidazole-5-carboxylate**




[00228] Except that amine is replaced by propan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

25 [00229] ^1H NMR (400 MHz, Methanol-*d*₄) δ 7.96 (s, 1H), 7.91 (d, J = 8.4 Hz, 1H), 7.68 (s, 1H), 7.42 (d, J = 8.4 Hz, 1H), 5.41 (s, 2H), 4.09 (t, J = 7.4 Hz, 2H), 3.81 (d, J = 13.8 Hz, 6H), 2.34 (s, 3H), 2.20 (s, 3H), 1.85 (d, J = 7.9 Hz, 2H), 1.02 (t, J = 7.4 Hz, 3H); ^{13}C NMR (101 MHz, Methanol-

d_4) δ 164.54 , 148.32 , 134.44 , 126.25 , 124.69 , 124.43 , 109.95 , 108.43 , 59.29 , 51.27 , 45.10 , 43.91 , 29.31 , 20.86 , 11.90 , 9.78 , 9.16 ; LRMS (ESI+) : m/z 383.3 (M+H)⁺.

Example 2-3 21103: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1- (3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylate



5


[00230] Except that amine is replaced by 3-methoxypropan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

[00231] ¹H NMR (400 MHz, Methanol- d_4) δ 8.10 (s, 1H), 7.90 (t, J = 1.8 Hz, 1H), 7.67 (dt, J = 8.4, 1.9 Hz, 1H), 7.11 (dd, J = 8.4, 2.1 Hz, 1H), 4.58 (d, J = 2.1 Hz, 2H), 4.09 – 4.01 (m, 2H), 3.82 (d, J = 2.1 Hz, 3H), 3.73 (d, J = 2.1 Hz, 3H), 3.19 (d, J = 2.1 Hz, 3H), 2.19 (dd, J = 10.1, 2.1 Hz, 6H), 1.96 (p, J = 6.2 Hz, 2H); ¹³C NMR (101 MHz, Methanol- d_4) δ 167.93 , 164.20 , 155.50 , 153.88 , 147.84 , 141.23 , 138.15 , 125.49 , 124.19 , 122.68 , 121.59 , 116.14 , 106.85 , 68.04 , 59.16 , 57.48 , 50.95 , 45.26 , 38.49 , 28.17 , 11.97 , 9.08 ; LRMS (ESI+) : m/z 413.3 (M+H)⁺.

10

Example 2-4 26070: 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylic acid

15



20

[00232] To a solution of 4-fluoro-3-nitrobenzoic acid **1** (5.0 g, 27.0 mmol) in dry MeOH (30 mL), H₂SO₄ (5 mL, 0.3 M) was added and the reaction mixture was heated to reflux for 12 h. The solvent was removed under reduced pressure, crude reaction mixture was dissolved in EtOAc (150 mL), washed with saturated NaHCO₃ (20 mL $\times$ 2), water (10 mL $\times$ 2) and brine (10 mL). The EtOAc layer was dried over anhydrous MgSO₄ and evaporated to get methyl 4-fluoro-3-nitrobenzoate **2** (95%) as a white solid.

[00233] Compound **2** (2.0 g, 10.2 mmol) and furan-2-ylmethanamine (3 equiv.) in dry CH₂Cl₂ (50 mL) were stirred at room temperature for 2 h. Upon completion of reaction the solvent was removed and the crude product was purified by flash column chromatography to afford nitro benzoates **3** (90%).

[00234] To a solution of compound **3** (2.0 g, 4.8 mmol) in dry MeOH (100 mL), zinc dust (15 equiv., 71.4 mmol) and ammonium formate (7.5 equiv., 35.7 mmol) were added and the resulting reaction mixture was stirred at room temperature for 30 min. Upon completion of reaction, Zn dust was filtered through a bed of celite, filtrate was evaporated and the product was dissolved in CH₂Cl₂ (100 mL). The precipitated ammonium formate was filtered off and the solvent was evaporated to furnish compound **4** (92 %).


[00235] Use DCM to dissolve compound **4** (1.0 g, 4.0 mmol) then add 1.2 equiv. CNBr to react at room temperature. After 8 hours the mixture can be extracted with DCM and water. The solvent was removed and the crude product was purified by flash column chromatography to afford **5** (60%)

[00236] To a solution of methyl 2-amino-1-(furan-2-ylmethyl)-1H-benzo[d]imidazole-5-carboxylate **5** (0.05 g, 0.18 mmol) in acetonitrile (10 mL) was added K₂CO₃ (0.0497 g, 0.36 mmol) and KI (0.0089 g, 0.054 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** (0.041 g, 0.22 mmol) and the reaction mixture was allowed to reflux for six hour. After 6 h, the solvent was evaporated and the reaction mixture was diluted with saturated aq. NaHCO₃ (10 mL) and extracted with EtOAc (3* 10 mL). The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 8 % methanol/ EtOAc to obtain Methyl 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate **7**.

[00237] And add NaOH (0.0251 g, 0.63 mmol) to a solution of Methyl 1-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate **7** (0.053g, 0.126 mmol) in the EtOH/H₂O (1/1, 3mL) in the reflux condition. After 1 h, the solvent was evaporated and the reaction mixture was diluted with saturated aq. HCl (10 mL) and extracted with EtOAc (3* 10 mL). The combined organic phase was washed with saturated brine (10 mL). The crude product was purified by silica-gel column chromatography using 20 % methanol/ EtOAc to obtain the pure product as a white solid 0.030 g (65%).

[00238] LRMS (ESI⁺) : m/z 407.2 (M+H)⁺.


Example 2-5 26066: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-pentyl-1H-benzo[d]imidazole-5-carboxylate



[00239] Except that amine is replaced by pentan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

[00240] ^1H NMR (400 MHz, Chloroform-*d*) δ 8.08 (s, 1H), 8.01 (dd, J = 8.4, 1.2 Hz, 1H), 7.72 (s, 1H), 7.29 (d, J = 8.5 Hz, 1H), 5.92 (s, 2H), 4.49 (t, J = 7.2 Hz, 2H), 3.89 (s, 6H), 2.47 (s, 3H), 2.26 (s, 3H), 1.96 – 1.83 (m, 2H), 1.54 – 1.41 (m, 2H), 1.44 – 1.29 (m, 2H), 0.88 (t, J = 7.2 Hz, 3H); LRMS (ESI⁺) : m/z 411.2 ($M+H$)⁺.


Example 2-6 21102: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-propyl-1H-benzo[d]imidazole-5-carboxylic acid



[00241] Except that amine is replaced by propan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00242] ^1H NMR (400 MHz, Methanol-*d*₄) δ 8.04 (d, J = 8.3 Hz, 1H), 7.95 (s, 1H), 7.89 (s, 1H), 7.64 (d, J = 8.1 Hz, 1H), 5.57 (s, 2H), 4.25 (t, J = 7.3 Hz, 2H), 3.83 (s, 3H), 2.41 (s, 3H), 2.21 (s, 3H), 1.92 (d, J = 7.3 Hz, 2H), 1.07 (t, J = 7.3 Hz, 3H); ^{13}C NMR (101 MHz, Methanol-*d*₄) δ 150.03, 148.40, 130.63, 126.40, 125.74, 111.55, 109.87, 59.38, 45.34, 44.56, 20.95, 11.93, 9.68, 9.17; LRMS (ESI⁺) : m/z 369.2 ($M+H$)⁺.


Example 2-7 26071: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(4-methoxybenzyl)-1H-benzo[d]imidazole-5-carboxylate



[00243] Except that amine is replaced by (4-methoxyphenyl)methanamine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

[00244] ^1H NMR (400 MHz, Methanol- d_4) δ 8.02 (s, 1H), 7.83 (dd, $J=8.4$, 1.5 Hz, 1H), 7.68 (d, $J=1.6$ Hz, 1H), 7.29 (d, $J=8.7$ Hz, 2H), 7.23 (d, $J=8.4$ Hz, 1H), 6.92 (d, $J=8.7$ Hz, 2H), 5.43 (s, 2H), 5.29 (s, 2H), 3.83 (s, 3H), 3.81 (s, 3H), 3.77 (s, 3H), 2.36 (s, 3H), 2.24 (s, 3H); LRMS (ESI $^+$) : m/z 461.2 (M+H) $^+$.


Example 2-8 21105: 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(2-(cyclohex-1-en-1-yl)ethyl)-1H-benzo[d]imidazole-5-carboxylic acid



[00245] Except that amine is replaced by 2-(cyclohex-1-en-1-yl)ethanamine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00246] ^1H NMR (400 MHz, Methanol- d_4) δ 8.17 (s, 2H), 8.07 (s, 1H), 7.82 (d, $J=8.2$ Hz, 1H), 7.26 (d, $J=8.3$ Hz, 1H), 5.13 (s, 1H), 4.77 (s, 4H), 4.11 (t, $J=7.8$ Hz, 2H), 3.80 – 3.69 (m, 6H), 2.29 (t, $J=8.4$ Hz, 2H), 2.18 (d, $J=21.8$ Hz, 12H), 1.76 (s, 4H), 1.49 – 1.36 (m, 4H); ^{13}C NMR (101 MHz, Methanol- d_4) δ 156.95 , 145.93 , 139.17 , 131.88 , 130.28 , 125.53 , 118.06 , 117.75 , 117.05 , 115.18 , 114.79 , 110.01 , 100.70 , 51.25 , 46.58 , 35.22 , 28.52 , 19.80 , 16.65 , 14.33 , 13.64 , 3.99 , 1.48 ; LRMS (ESI $^+$) : m/z 584.3 (M+H) $^+$.


Example 2-9 21104: 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylic acid



[00247] Except that amine is replaced by 3-methoxypropan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00248] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.06 (s, 2H), 7.88 (d, $J = 1.6$ Hz, 1H), 7.68 (dd, $J = 8.4, 1.6$ Hz, 1H), 7.32 (d, $J = 8.4$ Hz, 1H), 4.65 (s, 4H), 4.17 (t, $J = 7.6$ Hz, 2H), 3.62 (s, 6H), 3.20 (t, $J = 5.9$ Hz, 2H), 3.09 (s, 3H), 2.10 (s, 6H), 2.05 (s, 6H), 1.94 – 1.88 (m, 2H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ 168.52, 163.68, 158.90, 155.26, 148.49, 141.33, 125.19, 125.04, 124.05, 122.18, 118.11, 109.15, 69.30, 60.07, 58.30, 54.87, 41.74, 29.10, 13.24, 10.68; LRMS (ESI $^+$): m/z 548.3 ($\text{M}+\text{H}$) $^+$.


Example 2-10 26076: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-pentyl-1H-benzo[d]imidazole-5-carboxylic acid



[00249] Except that amine is replaced by pentan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00250] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.88 (s, 1H), 7.81 (dd, $J = 8.2, 1.3$ Hz, 1H), 7.70 (d, $J = 1.6$ Hz, 1H), 7.39 (d, $J = 8.4$ Hz, 2H), 5.59 (s, 2H), 4.23 (t, $J = 6.9$ Hz, 2H), 3.72 (s, 3H), 2.30 (s, 3H), 2.11 (s, 3H), 1.83 (s, 2H), 1.30 (dq, $J = 6.7, 3.3$ Hz, 4H), 0.90 – 0.73 (m, 3H); LRMS (ESI $^+$): m/z 397.2 ($\text{M}+\text{H}$) $^+$.


Example 2-11 26077: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(4-methoxybenzyl)-1H-benzo[d]imidazole-5-carboxylic acid



[00251] Except that amine is replaced by (4-methoxyphenyl)methanamine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00252] LRMS (ESI⁺) : m/z 447.2 (M+H)⁺.


Example 2-12 21115: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-(3-methoxypropyl)-1H-benzo[d]imidazole-5-carboxylic acid



[00253] Except that amine is replaced by 3-methoxypropan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00254] ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.20 (s, 1H), 7.98 – 7.92 (m, 2H), 7.50 (d, *J* = 8.7 Hz, 1H), 4.86 (s, 2H), 4.34 (t, *J* = 6.7 Hz, 2H), 3.87 (s, 3H), 3.44 (t, *J* = 5.6 Hz, 2H), 3.27 (s, 3H), 2.35 (s, 3H), 2.28 (s, 3H), 2.17 – 2.10 (m, 2H); ¹³C NMR (101 MHz, Methanol-*d*₄) δ 169.05 , 152.95 , 147.73 , 136.45 , 132.73 , 127.94 , 127.04 , 126.46 , 125.88 , 114.80 , 110.19 , 69.41 , 60.81 , 58.69 , 46.10 , 41.07 , 28.74 , 13.36 , 10.48 ; LRMS (ESI⁺) : m/z 399.2 (M+H)⁺.


Example 2-13 21116: Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylate



[00255] Except that amine is replaced by 2-(cyclohex-1-en-1-yl)ethanamine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

5 **[00256]** ^1H NMR (400 MHz, Acetone- d_6) δ 8.20 (s, 1H), 7.94 (d, J = 1.4 Hz, 1H), 7.70 (dt, J = 8.2, 1.4 Hz, 1H), 7.22 (dd, J = 8.3, 1.2 Hz, 1H), 6.71 (s, 1H), 5.33 (dt, J = 4.8, 2.3 Hz, 1H), 4.68 (d, J = 3.7 Hz, 2H), 4.17 (td, J = 7.2, 1.2 Hz, 2H), 3.84 (d, J = 1.2 Hz, 3H), 3.79 (d, J = 1.2 Hz, 3H), 2.41 (t, J = 7.2 Hz, 2H), 2.28 (s, 3H), 2.23 (s, 3H), 2.04 (h, J = 1.8 Hz, 2H), 2.02 – 1.97 (m, 2H), 1.82 – 1.76 (m, 2H), 1.51 (t, J = 5.9 Hz, 2H), 1.42 – 1.37 (m, 2H); ^{13}C NMR (101 MHz, Acetone- d_6) δ 167.29 , 163.89 , 155.45 , 154.37 , 148.06 , 142.70 , 138.71 , 133.89 , 125.06 ,
10 124.16 , 123.65 , 122.50 , 120.86 , 116.95 , 107.03 , 59.48 , 50.95 , 45.16 , 41.27 , 36.41 , 29.18 , 8.99 , 28.79 , 28.11 , 24.91 , 22.61 , 21.81 , 12.36 , 9.35 .

Example 2-14 **21117:** **1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1H-benzo[d]imidazole-5-carboxylic acid**




15 **[00257]** Except that amine is replaced by 2-(cyclohex-1-en-1-yl)ethanamine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00258] ^1H NMR (400 MHz, DMSO- d_6) δ 8.15 (s, 1H), 7.70 (s, 1H), 7.56 (d, J = 8.2 Hz, 1H),
20 7.15 (d, J = 8.3 Hz, 1H), 7.08 (t, J = 5.3 Hz, 1H), 5.23 (s, 1H), 4.61 (d, J = 4.2 Hz, 2H), 4.12 (t, J

= 7.1 Hz, 2H), 3.69 (s, 3H), 2.28 – 2.13 (m, 8H), 1.90 (s, 2H), 1.72 (s, 2H), 1.38 (dq, $J = 31.9, 5.4$ Hz, 4H); ^{13}C NMR (101 MHz, DMSO- d_6) δ 168.24, 155.18, 155.12, 147.74, 141.57, 137.88, 133.55, 124.39, 123.56, 123.09, 123.01, 120.32, 115.84, 106.77, 59.43, 45.42, 40.30, 35.64, 27.52, 24.33, 21.97, 21.29, 12.55, 9.85.


5 **Example 2-15 21118: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl-1H-benzo[d]imidazole-5-carboxylate**



[00259] Except that amine is replaced by 2-phenylethanamine, the other reactants and preparation steps are similar to those described in Example 2-1 to afford the title compound.

10 [00260] ^1H NMR (400 MHz, Methanol- d_4) δ 8.16 (s, 1H), 7.89 (s, 1H), 7.63 (d, $J = 8.2$ Hz, 1H), 7.10 (dt, $J = 16.2, 7.1$ Hz, 5H), 6.99 (d, $J = 8.2$ Hz, 1H), 4.57 (s, 3H), 4.30 (t, $J = 7.0$ Hz, 2H), 3.82 (d, $J = 25.8$ Hz, 6H), 3.05 (t, $J = 7.0$ Hz, 2H), 2.29 – 2.20 (m, 6H); ^{13}C NMR (101 MHz, Methanol- d_4) δ 169.48, 165.74, 156.92, 155.57, 149.24, 142.52, 139.50, 139.30, 130.05, 129.59, 127.73, 127.01, 125.84, 123.98, 122.90, 117.45, 108.52, 60.62, 52.38, 49.85, 46.69, 44.97, 40.00, 39.79, 39.58, 39.37, 39.16, 35.62, 13.36, 10.56; LRMS (ESI+) : m/z 445.4 ($\text{M}+\text{H}$) $^+$.


15 **Example 2-16 21119: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl-1H-benzo[d]imidazole-5-carboxylic acid**



20 [00261] Except that amine is replaced by 2-phenylethanamine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00262] ^1H NMR (400 MHz, Methanol- d_4) δ 8.19 (s, 1H), 7.89 (d, J = 1.5 Hz, 1H), 7.79 (dd, J = 8.5, 1.7 Hz, 1H), 7.19 – 7.07 (m, 6H), 4.73 (s, 2H), 4.48 (t, J = 6.8 Hz, 2H), 3.87 (s, 3H), 3.16 (t, J = 6.8 Hz, 2H), 2.29 (d, J = 14.6 Hz, 6H); ^{13}C NMR (101 MHz, Methanol- d_4) δ 169.15 , 167.25 , 153.40 , 153.06 , 148.02 , 138.56 , 136.34 , 130.16 , 129.70 , 128.21 , 128.04 , 127.16 , 126.66 , 126.02 , 114.72 , 110.61 , 61.03 , 46.15 , 45.80 , 34.84 , 13.58 , 10.68 .


Example 2-17 21120: 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-propyl- 1H-benzo[d]imidazole-5-carboxylic acid



[00263] Except that amine is replaced by propan-1-amine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00264] ^1H NMR (400 MHz, DMSO- d_6) δ 8.09 (s, 2H), 7.88 (d, J = 1.6 Hz, 1H), 7.68 (dd, J = 8.3, 1.6 Hz, 1H), 7.39 (d, J = 8.4 Hz, 1H), 4.63 (s, 4H), 4.11 (t, J = 7.9 Hz, 2H), 3.62 (s, 6H), 2.08 (d, J = 17.4 Hz, 12H), 1.66 (d, J = 7.7 Hz, 2H), 0.73 (t, J = 7.3 Hz, 3H).


Example 2-18 21121: 2-(bis((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-1-phenethyl- 1H-benzo[d]imidazole-5-carboxylic acid



[00265] Except that amine is replaced by 2-phenylethanamine, the other reactants and preparation steps are similar to those described in Example 2-4 to afford the title compound.

[00266] ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.07 (s, 2H), 7.89 (d, J = 1.5 Hz, 1H), 7.67 (dd, J = 8.4, 1.7 Hz, 1H), 7.40 (d, J = 8.4 Hz, 1H), 7.19 – 7.13 (m, 3H), 7.10 – 7.05 (m, 2H), 4.64 (s, 4H), 4.39 (t, J = 8.0 Hz, 2H), 3.59 (s, 6H), 2.97 (t, J = 8.1 Hz, 2H), 2.07 (d, J = 11.5 Hz, 12H); ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ 168.15, 163.34, 154.81, 148.03, 140.97, 138.79, 138.04, 128.79, 128.39, 126.52, 124.88, 124.76, 123.79, 122.04, 117.97, 109.35, 59.74, 55.15, 45.31, 34.51, 12.91, 10.42.

Example 3-1 NCTU-SUN-26079 : Methyl 3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate



[00267] To a solution of 4-(bromomethyl)-3-nitrobenzoic acid **1** (5.0 g, 27.0 mmol) in dry MeOH/ CH_2Cl_2 (3 mL: 30 mL), was added DCC (1.2 equiv) and DMAP (0.005 equiv) and the reaction mixture was stirred at room temperature for 16 hours. The byproduct DCU was filtered off and crude was purified by flash column chromatography to get methyl 4-(bromomethyl)-3-nitrobenzoate **2** (76%) as a off-white solid.

[00268] Compound **2** (4.0 g, 14.5 mmol) and 2-aminomethylfuran (3 equiv) in dry CH_2Cl_2 (50 mL) were stirred at room temperature for 48 hours. Upon completion of reaction the solvent was removed and the crude product was purified by flash column chromatography to afford nitro benzoate **3** (82%).

[00269] To a solution of compound **3** (3.65 g, 11.9 mmol) in dry MeOH (100 mL), $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (3.5 equiv) was added and the resulting reaction mixture was refluxed for 10 minutes. Upon completion of reaction, the byproduct was filtered through a bed of celite and filtrate was evaporated. The crude product was portioned between 1 N NaOH and ethyl acetate. The aqueous layer was extracted with ethyl acetate (3 x 20 mL) and the combined layers were dried over MgSO_4 and concentrated under reduced pressure to furnish compound **4** (87%).

[00270] Use DCM to dissolve compound **4** (1.0 g, 3.8 mmol) then add 1.2 equiv. CNBr to react at room temperature. After 8 hours the mixture can be extracted with DCM and water. The solvent


was removed and the crude product was purified by flash column chromatography to afford **5** (60%)

[00271] To a solution of methyl 2-amino-3-(furan-2-ylmethyl)-3,4-dihydroquinazoline-7-carboxylate **5** (0.3 g, 1.05mmol) in acetonitrile (20 mL) was added K_2CO_3 (0.29g, 2.1 mmol) and KI (0.005 g, 0.03 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** (0.722 g, 3.89 mmol) and the reaction mixture was allowed to reflux for six hour. After 24 hours, the solvent was evaporated and the reaction mixture was diluted with saturated aq. $NaHCO_3$ (30 mL) and extracted with EtOAc (3* 30 mL).

[00272] The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 8 % methanol/ EtOAc to obtain the pure product **NCTU-SUN-26079** as a white solid 0.43 g (70 %).

[00273] ^1H NMR (400 MHz, Methanol- d_4) δ 8.16 (s, 1H), 7.81 (dd, J = 7.9, 1.4 Hz, 1H), 7.56 (dd, J = 1.8, 0.8 Hz, 1H), 7.37 (d, J = 1.4 Hz, 1H), 7.32 (d, J = 7.9 Hz, 1H), 6.60 (d, J = 3.3 Hz, 1H), 6.47 (dd, J = 3.3, 1.9 Hz, 1H), 5.28 (s, 2H), 4.88 (s, 2H), 4.61 (s, 2H), 3.88 (s, 3H), 3.83 (s, 3H), 2.45 (s, 3H), 2.28 (s, 3H) ; LRMS (ESI+) : m/z 435.3 (M+H) $^+$.


Example 3-2 21106: Methyl 3-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate



[00274] Except that amine is replaced by 2-(cyclohex-1-en-1-yl)ethanamine, the other reactants and preparation steps are similar to those described in Example 3-1 to afford the title compound.

[00275] ^1H NMR (400 MHz, Methanol- d_4) δ 8.17 (s, 1H), 7.83 (dd, J = 7.9, 1.5 Hz, 1H), 7.40 – 7.33 (m, 2H), 5.24 (s, 2H), 5.16 (s, 1H), 4.61 (s, 2H), 3.88 (s, 3H), 3.84 (s, 3H), 3.76 (s, 2H), 2.46 (s, 3H), 2.28 (s, 3H), 2.12 (d, J = 8.1 Hz, 2H), 1.99 (s, 2H), 1.63 – 1.55 (m, 4H), 1.43 – 1.36 (m, 2H); LRMS (ESI+) : m/z 314.2 (M+H) $^+$.


Example 3-3 26072: Methyl 2-(bis(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(furan-2-ylmethyl)-3,4-dihydroquinazoline-7-carboxylate



[00276] Same as described in Example 3-1 to afford the title compound.

[00277] ^1H NMR (400 MHz, Methanol- d_4) δ 8.21 – 8.19 (m, 1H), 8.07 (d, J = 0.8 Hz, 1H), 7.83 (dd, J = 7.9, 1.5 Hz, 1H), 7.57 (d, J = 1.5 Hz, 1H), 7.43 (dd, J = 1.9, 0.8 Hz, 1H), 7.29 (d, J = 7.9 Hz, 1H), 6.57 (dd, J = 3.4, 0.8 Hz, 1H), 6.41 (dd, J = 3.3, 1.9 Hz, 1H), 5.48 (d, J = 1.9 Hz, 2H), 4.93 (s, 2H), 4.81 (s, 2H), 4.62 (s, 2H), 3.86 (s, 3H), 3.85 (s, 3H), 3.77 (s, 3H), 2.44 (s, 3H), 2.26 (s, 3H), 2.22 (s, 3H), 2.20 (s, 3H); LRMS (ESI+) : m/z 584.31 ($M+H$) $^+$.


Example 3-4 26091: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-pentyl-3,4-dihydroquinazoline-7-carboxylate



[00278] Except that amine is replaced by pentan-1-amine, the other reactants and preparation steps are similar to those described in Example 3-1 to afford the title compound.

[00279] ^1H NMR (400 MHz, Methanol- d_4) δ 8.14 (s, 1H), 7.79 (d, J = 7.9 Hz, 1H), 7.39 (d, J = 7.9 Hz, 1H), 7.32 (s, 1H), 5.24 (s, 2H), 4.70 (s, 2H), 3.86 (s, 3H), 3.81 (s, 3H), 3.64 (t, J = 7.8 Hz, 2H), 2.44 (s, 3H), 2.25 (s, 3H), 1.76 (p, J = 7.9 Hz, 2H), 1.37 (dp, J = 11.3, 7.1, 6.2 Hz, 4H), 0.96 – 0.83 (m, 3H); ^{13}C NMR (101 MHz, Methanol- d_4) δ 165.84, 165.03, 155.04, 152.00, 148.85, 137.27, 130.52, 128.00, 126.46, 125.87, 125.64, 124.60, 116.16, 59.47, 51.56, 50.90, 50.56, 47.94, 28.23, 26.36, 22.07, 12.93, 12.02, 9.51; LRMS (ESI+) : m/z 425.3 ($M+H$) $^+$.


Example 3-5 26092: Methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(4-methoxybenzyl)-3,4-dihydroquinazoline-7-carboxylate



[00280] Except that amine is replaced by (4-methoxyphenyl)methanamine, the other reactants and preparation steps are similar to those described in Example 3-1 to afford the title compound.

[00281] ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.09 (s, 1H), 7.71 (dd, *J* = 7.8, 1.4 Hz, 1H), 7.57 (d, *J* = 1.4 Hz, 1H), 7.39 (d, *J* = 8.5 Hz, 2H), 7.31 (d, *J* = 7.8 Hz, 1H), 6.92 – 6.83 (m, 2H), 5.65 (s, 2H), 5.18 (s, 2H), 4.68 (s, 2H), 3.82 (s, 3H), 3.79 (s, 3H), 3.74 (s, 3H), 2.43 (s, 3H), 2.18 (s, 3H); LRMS (ESI⁺) : *m/z* 475.3 (M+H)⁺.


Example 3-6 21110: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-(3-methoxypropyl)-3,4-dihydroquinazoline-7-carboxylic acid



[00282] Except that amine is replaced by 3-methoxypropan-1-amine, the other reactants and preparation steps are similar to those described in Example 3-7 to afford the title compound.

[00283] ¹H NMR (400 MHz, Methanol-*d*₄) δ 7.84 (dd, *J* = 7.9, 1.4 Hz, 1H), 7.64 (s, 1H), 7.46 (d, *J* = 1.4 Hz, 1H), 7.40 (d, *J* = 7.9 Hz, 1H), 5.26 (s, 2H), 4.69 (s, 2H), 3.85 (s, 3H), 3.75 (t, *J* = 6.9 Hz, 2H), 3.51 (t, *J* = 5.7 Hz, 2H), 2.27 (s, 3H), 2.04 (d, *J* = 4.9 Hz, 6H), 1.29 (d, *J* = 3.5 Hz, 2H); LRMS (ESI⁺) : *m/z* 413.3 (M+H)⁺.

Example 3-7 26089: 3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylic acid



[00284] To a solution of 4-(bromomethyl)-3-nitrobenzoic acid **1** (5.0 g, 27.0 mmol) in dry MeOH/CH₂Cl₂ (3 mL: 30 mL), was added DCC (1.2 equiv) and DMAP (0.005 equiv) and the reaction mixture was stirred at room temperature for 16 h. The byproduct DCU was filtered off and crude was purified by flash column chromatography to get methyl 4-(bromomethyl)-3-nitrobenzoate **2** (76%) as an off-white solid.

[00285] Compound **2** (4.0 g, 14.5 mmol) and 2-Aminomethylfuran (3 equiv) in dry CH₂Cl₂ (50 mL) were stirred at room temperature for 48 hours. Upon completion of reaction the solvent was removed and the crude product was purified by flash column chromatography to afford nitro benzoate **3** (82%).

[00286] To a solution of compound **3** (3.65 g, 11.9 mmol) in dry MeOH (100 mL), SnCl₂·2H₂O (3.5 equiv) was added and the resulting reaction mixture was refluxed for 10 min. Upon completion of reaction, the byproduct was filtered through a bed of celite and filtrate was evaporated. The crude product was portioned between 1 N NaOH and ethyl acetate. The aqueous layer was extracted with ethyl acetate (3 x 20 mL) and the combined layers were dried over MgSO₄ and concentrated under reduced pressure to furnish compound **4** (87%).

[00287] Use DCM to dissolve compound **4** (1.0 g, 3.8 mmol) then add 1.2 equiv. CNBr to react at room temperature. After 8 hours the mixture can be extracted with DCM and water. The solvent was removed and the crude product was purified by flash column chromatography to afford **5** (60%).

[00288] To a solution of methyl 2-amino-3-(furan-2-ylmethyl)-3,4-dihydroquinazoline-7-carboxylate **5** (0.3 g, 1.05 mmol) in acetonitrile (20 mL) was added K₂CO₃ (0.29 g, 2.1 mmol) and KI (0.005 g, 0.03 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** (0.722 g, 3.89 mmol) and the reaction mixture was allowed to reflux for six hours. After 24 hours, the


solvent was evaporated and the reaction mixture was diluted with saturated aq. NaHCO₃ (30 mL) and extracted with EtOAc (3* 30 mL).

[00289] The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 8 % methanol/ EtOAc to obtain Methyl 3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate 7 0.43 g (70 %).

[00290] And add NaOH (0.198 g, 4.95 mmol) to a solution of methyl 3-(furan-2-ylmethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3,4-dihydroquinazoline-7-carboxylate 7 (0.43g, 0.99 mmol) in the EtOH/H₂O (1/1, 10mL) in the reflux condition. After 1 hour, the solvent was evaporated and the reaction mixture was diluted with saturated aq. HCl (30 mL) and extracted with EtOAc (3* 30 mL). The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 20 % methanol/ EtOAc to obtain the pure product as a white solid 0.27 g (65%).

[00291] ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.07 (s, 1H), 7.70 (d, *J* = 7.9 Hz, 1H), 7.53 (d, *J* = 1.8 Hz, 1H), 7.35 (s, 1H), 7.14 (d, *J* = 7.8 Hz, 1H), 6.57 (d, *J* = 3.2 Hz, 1H), 6.43 (dd, *J* = 3.2, 1.8 Hz, 1H), 5.22 (s, 2H), 4.84 (s, 2H), 4.53 (s, 2H), 3.82 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ¹³C NMR (101 MHz, Methanol-*d*₄) δ 171.74 , 164.87 , 155.39 , 151.73 , 148.52 , 147.49 , 143.54 , 138.56 , 136.13 , 126.27 , 125.93 , 124.64 , 124.63 , 116.56 , 110.30 , 109.86 , 59.40 , 50.46 , 47.46 , 46.94 , 46.47 , 11.97 , 9.33; LRMS (ESI⁺) : *m/z* 421.2 (M+H)⁺.


Example 3-8 26090: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)amino)-3-pentyl-3,4-dihydroquinazoline-7-carboxylic acid



[00292] Except that amine is replaced by pentan-1-amine, the other reactants and preparation steps are similar to those described in Example 3-7 to afford the title compound.

[00293] ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.03 (dd, *J* = 8.4, 1.4 Hz, 1H), 7.96 (s, 1H), 7.84 (d, *J* = 1.3 Hz, 1H), 7.55 (d, *J* = 8.4 Hz, 1H), 3.83 (s, 3H), 2.41 (s, 3H), 2.22 (s, 3H), 1.92 – 1.83 (m, 2H), 1.43 (tt, *J* = 5.7, 2.8 Hz, 4H), 1.29 (d, *J* = 4.0 Hz, 2H), 0.97 – 0.92 (m, 3H); ; LRMS (ESI⁺) : *m/z* 411.3 (M+H)⁺.

Example 4-1 NCTU-SUN-12082: Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate



[00294] To a solution of 4-fluoro-3-nitrobenzoic acid **1**, H_2SO_4 (5 mL, 0.3 M) is added and the reaction mixture is heated to reflux. The solvent is removed under reduced pressure, crude reaction mixture is dissolved in EtOAc. The EtOAc layer was dried over anhydrous MgSO_4 and evaporated to get methyl 4-fluoro-3-nitrobenzoate **2** as a white solid.

[00295] Compound **2** and 2-(cyclohex-1-en-1-yl)ethanamine were stirred at room temperature for 2 hours. Upon completion of reaction the solvent was removed and the crude product was purified to afford nitro benzoates **3**.

[00296] To a solution of compound **3**, zinc dust and ammonium formate are added and the resulting reaction mixture is stirred at room temperature. Upon completion of reaction, Zn dust is filtered and the filtrate is evaporated and the product is dissolved in CH_2Cl_2 . The precipitated ammonium formate was filtered off and the solvent was evaporated to furnish compound **4**.


[00297] To the stirred solution of compound **4** is added carbon disulfide and KOH at 50 °C in the ethanol for 8 hours. The mixture can be neutralized by acetic acid and extracted with EtOAc and water. The solvent was removed and the crude product was purified to afford **5**.

[00298] To a solution of methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-thioxo-2,3-dihydro-1H-benzo[d]imidazole-5-carboxylate **5** is added K_2CO_3 and KI followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** and the reaction mixture was allowed to reflux. The solvent is evaporated and the reaction mixture is diluted and extracted with EtOAc.

[00299] The combined organic phase was washed with saturated brine. The crude product was purified to obtain the pure product NCTU-SUN-12082 as a white solid 0.053 g (71 %).

[00300] ^1H NMR (300 MHz, Acetone- d_6) δ 8.24 (d, J = 1.2 Hz, 1H), 8.20 (s, 1H), 7.91 (dd, J = 8.5, 1.4 Hz, 1H), 7.53 (d, J = 8.5 Hz, 1H), 5.22 (s, 1H), 4.83 (s, 2H), 4.28 (t, J = 7.0 Hz, 2H), 3.91 (s, 3H), 3.80 (s, 3H), 2.47 – 2.35 (m, 5H), 2.25 (s, 3H), 1.99 (m, 2H), 1.80 (m, 2H), 1.62 – 1.38 (m, 4H).

5 **Example 4-2 12083: 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylic acid**



[00301] To a solution of 4-fluoro-3-nitrobenzoic acid **1**, H_2SO_4 (5 mL, 0.3 M) is added and the reaction mixture is heated to reflux. The solvent is removed under reduced pressure, crude reaction mixture is dissolved in EtOAc. The EtOAc layer was dried over anhydrous MgSO_4 and evaporated to get methyl 4-fluoro-3-nitrobenzoate **2** as a white solid.

[00302] Compound **2** and 2-(cyclohex-1-en-1-yl)ethanamine were stirred at room temperature for 2 h. Upon completion of reaction the solvent was removed and the crude product was purified to afford nitro benzoates **3**.

[00303] To a solution of compound **3**, zinc dust and ammonium formate are added and the resulting reaction mixture is stirred at room temperature. Upon completion of reaction, Zn dust is filtered and the filtrate is evaporated and the product is dissolved in CH_2Cl_2 . The precipitated ammonium formate was filtered off and the solvent was evaporated to furnish compound **4**.

25 [00304] To the stirred solution of compound **4** is added carbon disulfide and KOH at 50 °C in the ethanol for 8 hours. The mixture can be neutralized by acetic acid and extracted with EtOAc and water. The solvent was removed and the crude product was purified to afford **5**.


[00305] To a solution of methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-thioxo-2,3-dihydro-1H-benzo[d]imidazole-5-carboxylate **5** is added K_2CO_3 and KI followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** and the reaction mixture was allowed to reflux. The solvent is evaporated and the reaction mixture is diluted and extracted with EtOAc.

[00306] The combined organic phase was washed with saturated brine. The crude product was purified to obtain Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate **7** 0.053 g (71 %).

[00307] And add NaOH (0.0251 g, 0.63 mmol) to a solution of Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate **7** (0.053g, 0.126 mmol) in the EtOH/H₂O (1/1, 3mL) in the reflux condition. After 1 hour, the solvent was evaporated and the reaction mixture was diluted with saturated aq. HCl (10 mL) and extracted with EtOAc (3* 10 mL). The combined organic phase was washed with saturated brine (30 mL). The crude product was purified by silica-gel column chromatography using 20 % methanol/ EtOAc to obtain the pure product **NCTU-SUN-12083** as a white solid 0.030 g (65%).

[00308] ¹H NMR (300 MHz, CD₃OD) δ 8.30 (d, J = 1.4 Hz, 1H), 8.14 (s, 1H), 7.98 (dd, J = 8.5, 1.5 Hz, 1H), 7.49 (d, J = 8.5 Hz, 1H), 5.51 (s, 2H), 5.08 (s, 1H), 4.71 (s, 2H), 4.24 (t, J = 6.8 Hz, 2H), 3.79 (s, 3H), 2.44 – 2.31 (m, 5H), 2.27 (s, 3H), 2.04 – 1.89 (m, 2H), 1.88 – 1.70 (m, 2H), 1.62 – 1.39 (m, 4H).

Example 4-3 12084: Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazole-5-carboxylate



[00309] To a solution of 4-fluoro-3-nitrobenzoic acid **1**, H₂SO₄ (5 mL, 0.3 M) is added and the reaction mixture is heated to reflux. The solvent is removed under reduced pressure, crude reaction mixture is dissolved in EtOAc. The EtOAc layer was dried over anhydrous MgSO₄ and evaporated to get methyl 4-fluoro-3-nitrobenzoate **2** as a white solid.

[00310] Compound **2** and 2-(cyclohex-1-en-1-yl)ethanamine were stirred at room temperature for 2 hours. Upon completion of reaction the solvent was removed and the crude product was purified to afford nitro benzoates **3**.

[00311] To a solution of compound **3**, zinc dust and ammonium formate are added and the resulting reaction mixture is stirred at room temperature. Upon completion of reaction, Zn dust is filtered and the filtrate is evaporated and the product is dissolved in CH₂Cl₂. The precipitated ammonium formate was filtered off and the solvent was evaporated to furnish compound **4**.

5 [00312] To the stirred solution of compound **4** is added carbon disulfide and KOH at 50 °C in the ethanol for 8 hours. The mixture can be neutralized by acetic acid and extracted with EtOAc and water. The solvent was removed and the crude product was purified to afford **5**.


[00313] To a solution of methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-thioxo-2,3-dihydro-1H-benzo[d]imidazole-5-carboxylate **5** is added K₂CO₃ and KI followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **6** and the reaction mixture was allowed to reflux. The solvent is
10 evaporated and the reaction mixture is diluted and extracted with EtOAc.

[00314] The combined organic phase was washed with saturated brine. The crude product was purified to obtain Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate **7** 0.053 g (71 %).

15 [00315] And add mCPBA (0.0058 g, 0.034 mmol) to a solution of Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate **7** (0.053g, 0.0126 mmol) in the DCM/ MeOH (9/1, 4.5 mL) in the ice bath. Then, add NaHCO₃ (0.0007 g, 0.0088 mmol) and remove the ice bath. Let the crude stir at room temperature in 1 hour. The reaction mixture was washed with DCM (5 mL). The solvent was
20 evaporated and the to obtain the pure product **NCTU-SUN-12084** as a white solid 0.030 g (65%).

[00316] ¹H NMR (300 MHz, CDCl₃) δ 8.55 (s, 1H), 8.13 (s, 1H), 8.10 (dd, *J* = 8.7, 1.5 Hz, 1H), 7.43 (d, *J* = 8.8 Hz, 1H), 5.03 (q, *J* = 12.9 Hz, 3H), 4.59 – 4.35 (t, *J* = 8.3 Hz, 2H), 4.58 – 4.36 (m, 2H), 3.97 (s, 3H), 3.71 (s, 3H), 2.49 (t, *J* = 8.3 Hz, 2H), 2.30 (s, 3H), 2.22 (s, 3H), 2.03 – 1.78 (m, 4H), 1.51 (m, 4H).


25 **Example 5-1 12092: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazol-5-yl (((9H-fluorene-9-yl)methoxy)carbonyl)glycinate**



[00317] To a solution of 2-thioxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl 2-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetate **1** (0.08 g, 0.18 mmol) in ethanol (9 mL) was added NaOH (0.079 g, 0.198 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **2** (0.367 g, 0.198 mmol) and the reaction mixture was allowed to reflux for one hour. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using 2 % MeOH/DCM to obtain the pure product **NCTU-SUN-12092** as a white solid. 0.31 g, 54.5 %.

[00318] ^1H NMR (300 MHz, Acetone- d_6) δ 8.26 (s, 1H), 7.87 (d, J = 7.4 Hz, 2H), 7.74 (d, J = 7.4 Hz, 2H), 7.51 (d, J = 8.6 Hz, 1H), 7.41 (t, J = 7.3 Hz, 2H), 7.32 (t, J = 7.4 Hz, 3H), 7.14 (t, J = 6.6 Hz, 1H), 6.95 (dd, J = 8.7, 2.1 Hz, 1H), 4.67 (s, 2H), 4.40 (d, J = 7.3 Hz, 2H), 4.36 – 4.21 (m, 3H), 3.80 (s, 3H), 2.36 (s, 3H), 2.25 (s, 3H).


Example 5-2 12093: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazol-5-yl (tert-butoxycarbonyl)glycinate



[00319] Except that imidazole is replaced by 2-thioxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl 2-((tert-butoxycarbonyl)amino)acetate, the other reactants and preparation steps are similar to those described in Example 5-1 to afford the title compound.

[00320] ^1H NMR (300 MHz, Acetone) δ 8.27 (s, 1H), 7.50 (s, 1H), 7.29 (d, J = 2.1 Hz, 1H), 6.93 (dd, J = 8.6, 2.1 Hz, 1H), 6.50 (s, 1H), 4.67 (s, 2H), 4.12 (d, J = 6.2 Hz, 2H), 3.81 (s, 3H), 2.38 (s, 3H), 2.26 (s, 3H), 1.45 (s, 9H).


Example 5-3 12094: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H benzo[d]-imidazole-5-yl (S)-2-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)-2-phenylacetate



[00321] Except that imidazole is replaced by (S)-2-thioxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl 2-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)-2-phenylacetate, the other reactants and preparation steps are similar to those described in Example 5-1 to afford the title compound.

[00322] ¹H NMR (300 MHz, Acetone-d₆) δ 8.24 (s, 1H), 7.85 (d, *J* = 7.5 Hz, 2H), 7.75 (d, *J* = 7.4 Hz, 2H), 7.65 (d, *J* = 7.2 Hz, 2H), 7.52 – 7.39 (m, 6H), 7.32 (m, 2H), 7.22 (d, *J* = 1.6 Hz, 1H), 6.83 (dd, *J* = 9.1, 1.5 Hz, 1H), 5.67 (s, 1H), 4.66 (s, 2H), 4.48 – 4.25 (m, 3H), 2.35 (s, 3H), 2.25 (s, 3H).

Example 6-1 NCTU-SUN-22138: 5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazole




[00323] To a solution of 2-methoxy-1,3,4-trimethylbenzene **1** (0.3 g, 2.00 mmol) in chloroform (30 mL) was added NBS (0.177 g, 1.00 mmol) and in the light-induced reactions, two Philips "IR 250 W lamps were placed at such a distance from the reaction flask that reflux was maintained. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using hexane to obtain brominated product **2** 0.092 g, 20 %.

[00324] To a solution of brominated product **2** (0.1 g, 0.43 mmol) in ethanol (2 mL) was added NaOH (0.017 g, 0.43 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **3** (0.071 g, 0.39 mmol) and the reaction mixture was allowed to reflux for one hour. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using 2 % MeOH/DCM to obtain NCTU-SUN-22138 0.077 g, 60 %.

[00325] LRMS (ESI⁺) : *m/z* 329.2 (M+H)⁺

Example 6-2 22141: 2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazol-5-ol




[00326] To a solution of 2-methoxy-1,3,4-trimethylbenzene **1** (0.3 g, 2.00 mmol) in chloroform (30 mL) was added NBS (0.177 g, 1.00 mmol) and in the light-induced reactions, two Philips "IR 250 W lamps were placed at such a distance from the reaction flask that reflux was maintained. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using hexane to obtain brominated product **2** 0.092 g, 20 %.

[00327] To a solution of brominated product **2** (0.1 g, 0.43 mmol) in ethanol (2 mL) was added NaOH (0.017 g, 0.43 mmol) followed by 2-(chloromethyl)-4-hydroxy-3,5-dimethylpyridine **3** (0.071 g, 0.39 mmol) and the reaction mixture was allowed to reflux for one hour. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using 2 % MeOH/DCM to obtain NCTU-SUN-22138 0.077 g, 60 %.

[00328] LRMS (ESI⁺) : m/z 315.1 (M+H)⁺

Example 6-3 21133: 2-((3-(bromomethyl)-2-((tert-butyldimethylsilyl)oxy)-6-methylbenzyl)thio)-5-methoxy-1H-benzo[d]imidazole




[00329] To a solution of tert-butyldimethyl(2,3,6-trimethylphenoxy)silane **1** (1.2 g, 4.7 mmol) in chloroform (50 mL) was added NBS (1.7 g, 9.5 mmol) and in the light-induced reactions, two Philips "IR 250 W lamps were placed at such a distance from the reaction flask that reflux was maintained. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using hexane to obtain dibrominated product **2** 0.31 g, 20 %.

[00330] To a solution of dibrominated product **2** (0.3 g, 0.90 mmol) in ethanol (9 mL) was added NaOH (0.036 g, 0.90 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **3** (0.148 g, 0.82 mmol) and the reaction mixture was allowed to reflux for one hour. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using 2 % MeOH/DCM to obtain NCTU-SUN-21133 0.23 g, 60 %.

[00331] ^1H NMR (400 MHz, Chloroform-*d*) δ 7.43 (d, J = 8.8 Hz, 1H), 7.16 (s, 1H), 7.04 (d, J = 2.3 Hz, 1H), 6.82 (d, J = 2.3 Hz, 1H), 4.69 (s, 2H), 3.78 (s, 4H), 2.22 (s, 3H), 2.13 (s, 4H), 0.98 (s, 9H), 0.09 (s, 6H).

[00332] LRMS (ESI+) : m/z 507.1 ($M+H$)⁺

5 **Example 6-4 22139:5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)sulfinyl)-1H-benzo[d]imidazole**



[00333] To a solution of 2-methoxy-1,3,4-trimethylbenzene **1** (0.3 g, 2.00 mmol) in chloroform (30 mL) was added NBS (0.177 g, 1.00 mmol) and in the light-induced reactions, two Philips "IR 250 W lamps were placed at such a distance from the reaction flask that reflux was maintained. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using hexane to obtain brominated product **2** 0.092 g, 20 %.

To a solution of brominated product **2** (0.1 g, 0.43 mmol) in ethanol (2 mL) was added NaOH (0.017 g, 0.43 mmol) followed by 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine **3** (0.071 g, 0.39 mmol) and the reaction mixture was allowed to reflux for one hour. Once the reaction was completed, the solvent was evaporated and the crude product was purified by silica-gel column chromatography using 2 % MeOH/DCM to obtain 5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazole **3** 0.077 g, 60 %.

[00334] And add mCPBA (0.069g, 0.40 mmol) to a solution of 5-methoxy-2-((2-methoxy-3,6-dimethylbenzyl)thio)-1H-benzo[d]imidazole **3** (0.077g, 0.23 mmol) in the DCM/ MeOH (9/1, 10 mL) in the ice bath. Then, add NaHCO₃ (0.013g, 0.16 mmol) and remove the ice bath. Let the crude stir at room temperature in 1 h. The reaction mixture was washed with DCM (10 mL). The solvent was evaporated and the to obtain the pure product as a white solid 0.047 g (65%).

[00335] LRMS (ESI+) : m/z 345.1 ($M+H$)⁺

Example 11 DAAO enzymatic assay

[00336] The DAAO enzymatic activity assay was modified according to the report of Oguri *et al* (Oguri, S., Screening of *d*-amino acid oxidase inhibitor by a new multi-assay method. Food

chemistry **2007**, *100* (2), 616). The DAAO activity was measured by using substrate D-alanine reaction produced hydrogen peroxide (H₂O₂) to further react with 3-(4-hydroxyphenyl) propionic acid (HPPA). The HPPA were oxidized by H₂O₂ and peroxidase to become the fluorogenic dimer which was measured to represent the activity of DAAO.

[00337] For porcine kidney DAO IC₅₀ assay, the DAO substrate was prepared in 50 mM *D*-alanine (dissolved in 0.2 M Tris-HCl buffer, pH 8.3). A 100 µl of *D*-alanine solution was mixed with 4 µl (in 100% dimethyl sulfoxide, DMSO) of different concentrations of candidate compounds shown in tables below ranging from 48.83 µM, 97.66 µM, 195.31 µM, 390.63 µM, 781.25 µM, 1.56 mM, 3.13 mM, 6.25 mM, 12.50 mM, 25.00 mM, and 50.00 mM, with a final DMSO concentration of 0.167% in each reaction concentration. 10 µl of *D*-alanine and candidate compound mixture was incubated with 220 µl of Reaction Master Mix in black 96 well plate at 37°C for 5 minutes. The Reaction Master Mix contained 110 µl of 5 U/mL porcine kidney DAO (Sigma-Aldrich, USA) solution (dissolved with 0.2 M Tris-HCl buffer, pH 8.3), 1.1 mL of 15 U/mL peroxidase solution (dissolved with 0.2 M Tris-HCl buffer, pH 8.3), 1.1 mL of 20 mM HPPA solution (dissolved with 0.2 M Tris-HCl buffer, pH 8.3), and 22 ml of 2 M Tris-HCl buffer (pH 8.3) for 110 reaction assays.

[00338] Fluorescence intensity (Fs) was measured at 405 nm by irradiation excitation at 320 nm. The higher the DAO enzymatic activity was, the higher the fluorescence intensity. The fluorometric inhibition indicator (Fi) was obtained from the following equation: $Fi = (Fs - F_{Drug}) / (F_{DMSO})$, where the fluorescent drug blank (F_{Drug}) was measured in the drug mixture solution (using 0.2 M Tris HCl buffer, pH 8.3, without *D*-alanine). A DMSO blank (F_{DMSO}) was measured under a 100% DMSO solution.

[00339] Although FAD is generally included in the reaction mixture in the *D*-amino acid oxidase assay since it easily dissociates from the holoenzyme, the present method was performed without FAD. The inhibitory effect of DAO inhibitors was compared by using inhibitory concentrations leading to 50% inhibition of DAAO activity (IC₅₀). The IC₅₀ values were calculated by nonlinear regression model using GraphPad Prism, version 5 (GraphPad Software, Inc., La Jolla, CA). The results of DAO IC₅₀ assay of the candidate compounds of the invention are shown in the table below.

ID	M.W.	pkDAO Enzymatic Assay
----	------	-----------------------

		IC50 (uM)	IC50 95% Confidence Range (uM)
CBIO	169.6	0.5827	0.5165 to 0.6573
RS-D7	331.39	1.1850	0.9970 to 1.408
12083	451.5811	366.6000	302.2 to 444.8
21105	583.73	300.8000	223.0 to 405.9
26072	583.69	236.4000	211.2 to 264.5
21106	462.59	488.1000	309.4 to 769.9
21132	431.15	64.1500	60.86 to 67.61
13084	563.56	181.2000	154.3 to 212.8
12122	425.4577	12.5400	11.55 to 13.62
21122	373.1096	21.0900	16.63 to 26.75
11020	465.13	10.4400	9.549 to 11.40
12124	399.46	11.1600	10.11 to 12.32
12125	413.49	13.3100	12.26 to 14.44
25016	415.16	13.3600	12.39 to 14.42
28092	399.13	46.7000	41.85 to 52.10
28094	449.52	21.1100	19.18 to 23.23
25027	413.1409	21.0000	19.70 to 22.37

Example 12 Cell-based DAO Assay

[00340] Neuronal cell culture

[00341] The SK-N-SH neuroblastoma cell line was purchased from American Type Culture Collection (ATCC). It was cultured in MEM media (Invitrogen™/GIBCO™, Rockville, MD) supplemented with 10% fetal bovine serum, 1X NEAA (Invitrogen™/GIBCO™) at 37 °C with 5% CO2 in a humidified atmosphere. Cells were trypsinized and plated into a black 96-well plate (NUNC™ No. 237108) at a density of 125,000 cells/well in 50 µl before cell-based DAO assay.

[00342] Cell-based DAO Assay

[00343] The cellular DAO activity assay was using a method modified according to the report of Brandish *et al.* (Brandish, P.E., *et al.*, *A cell-based ultra-high-throughput screening assay for identifying inhibitors of D-amino acid oxidase. J Biomol Screen*, **2006**. 11(5): p. 481-7.). SK-N-

SH cells were suspended in assay buffers of HANKS buffer solution (Invitrogen™/GIBCO™ No. 14025-092) with 20 mM HEPES. D-serine of (final concentration 50 mM) was added in each well as the substrate for DAO enzyme. The Amplex™ Red Hydrogen Peroxide/Proxidase Assay Kit (Molecular Probes/Invitrogen™, cat. A22188) was used to measure H₂O₂ production which diffused across the cell membrane into the assay medium after DAO reaction. After seeding cells into black 96-well plate (Nunc™ No. 237108, Denmark), 50 µl of SK-N-SH cells (125,000 cells/well) was mixed with 50 µl drug solution (2.5 fold of interesting final concentration) and incubate at 37 °C with 5% CO₂ in a humidified atmosphere for 30 minutes. After 30 minutes later, 25 µl of a 5 fold mixture containing D-serine, horseradish peroxidase (HRP), and Amplex™ Red was added into wells containing 100 µl of cell-drugs mixture and incubated at 37 °C with 5% CO₂ in a humidified atmosphere for 3 hours. The final concentration of DMSO is below 1%. The fluorescence signal was then detected in SpectraMax® M2e microplate reader (Molecular Devices, USA) with excitation at 544 nm and emission at 590 nm. The optimized assay buffer contained a final concentration of 50 mM D-serine, 0.625 units HRP, and 50 µM Amplex™ Red in a 125 µl assay volume. The results of the cell-base DAO assay of the candidate compounds of the invention are shown in the table below.

ID	M.W.	Cell-base DAO assay			
		SK-N-SH 1.25*10 ⁵ cells/well (Drug treated for 3.5 hr)			
		Relative DAO activity normalize to DMSO at Final Conc.		Cell-based IC ₅₀ assay	
		100 µM	10 µM	IC ₅₀ (µM)	IC ₅₀ 95% Confidence Range (µM)
CBIO	169.6	0.83	1.00	392.30	281.4 to 547.0
RS-D7	331.39	0.69	1.05	141.00	81.07 to 245.1
12083	451.5811	0.48	0.62	74.15	27.04 to 203.4
21105	583.73	0.58	0.89	218.80	98.89 to 484.3
26072	583.69	0.74	0.87	393.00	215.3 to 717.2
21106	462.59	0.77	1.02	477.60	245.5 to 929.2

21132	431.15	0.61	0.94	174.80	137.9 to 221.7
13084	563.56	0.49	0.71	34.91	21.34 to 57.09
12122	425.4577	0.57	0.98	146.60	106.1 to 202.6
21122	373.1096	0.50	0.99	149.40	94.76 to 235.5
11020	465.13	0.67	0.97	220.90	121.5 to 401.9
12124	399.46	0.36	0.97	107.60	73.47 to 157.6
12125	413.49	0.50	0.80	122.20	61.25 to 243.7
25016	415.16	0.62	1.00	187.60	107.7 to 326.8
28092	399.13	0.58	0.94	136.40	86.51 to 215.1
28094	449.52	0.58	1.10	121.50	73.91 to 199.9
25027	413.1409	0.69	1.15	240.20	142.0 to 406.3

Example 13 Animal Studies of Potency in Treating Schizophrenia Symptoms

[00344] Drug Efficacy Screening

[00345] The NMDA-receptor antagonist MK-801-induced negative or cognitive deficits in C57BL/6 mice were in a well-established drug-induced schizophrenia mouse model and as a useful pharmacological animal model to identify if the RS-D7, its analog and its prodrug improve the symptoms through the NMDA receptor.

[00346] Animals

[00347] All wild-type (WT) mice used in this study were backcrossed onto a C57BL/6J background from the Laboratory Animal Center of National Taiwan University Hospital, and all behavioral examinations were conducted in WT mice. For acclimatizing to laboratory conditions, the mice were allowed free access to food and water and were housed in groups with a 12 hours light-dark cycle in a temperature and humidity controlled room of the Psychology Department, National Taiwan University. All animals within the age of 3 months were housed individually one week before experiment testing with food and water available ad libitum. In the beginning of the experiments, the mice were handled and weighted daily at least 1 week before the behavioral experiments. The entire animal procedures were performed according to protocols approved by the Animal Care and Use Committee established by the National Taiwan University.

[00348] Drug preparation for treating animals

[00349] MK-801 was dissolved in saline and administered at a volume of 0.01 ml/g body weight. RS-D7, Drug 12083 (analog of RS-D7) and was freshly dissolved in 1% CMC to the concentration of 2 mg/ml before usage. Prodrug 28095 was freshly dissolved in NMP:HP-beta-CD:H₂O (5:25:70) to the concentration of 2 mg/ml before usage. All animals were given vehicle (saline) or MK-801 (0.2 mg/kg, *i.p.*) 25 minutes before the behavioral experiments. Both vehicle (1% CMC or NMP:HP-beta-CD:H₂O) and experimental groups (RS-D7, Drug 12083 or Prodrug 28095) were treated after 5 minutes MK-801 administration with the appropriate dose via *P.O.* injections (at a volume of 0.01 ml/g body weight).

[00350] Behavioral experiment procedure

[00351] For investigating the treatment effect of RS-D7 for negative and cognitive symptoms, a series of three behavioral tests (run from the first to the third week), which included open field, sucrose preference test and prepulse inhibition were performed in sequence with a 1 week interval between tests.

[00352] Open field

[00353] To assess spontaneous locomotor activity, each subject was placed into the center of an open-field apparatus (25.40*25.40*40.64 cm³, Coulbourn Instruments, Whitehall, PA, USA) under dim lighting condition (60 lx). Motor activity parameters (including total travel distance and travel distance per 10 minutes) were monitored and recorded over a 60 minutes period by using Smart video tracking software (Panlab, Harvard apparatus, US). For comparing the treatment effects of different treatment groups, the percentage change of rescue effect on MK-801-induced hyperlocomotion was calculated using the following formula: % = (rescue effect of drug – MK-801 effect) X 100 % / MK-801 effect.

[00354] Compared with the saline controls, mice exhibit the hyperlocomotion in the open field after acute MK-801 injection. The injection of 200 and 400 mg/kg RS-D7 rescued the MK-801 induced hyperlocomotion in mice. However, the 100 mg/kg RS-D7 did not display the treatment effect. This result suggests that acute RS-D7 injection normalized MK-801-induced hyperlocomotion as a positive symptom of schizophrenia in the open field test. Drug 12083 alleviated MK-801 induced hyperlocomotion deficits at 20 and 40 mg/kg dose. 100 mg/kg Prodrug 28095 alleviated MK-801 induced hyperlocomotion. In conclusion, compared to the MK-801 group, different dosages of RS-D7, Drug 12083 and Prodrug 28095 can rescue the MK-801-

induced hyperlocomotion. The rescue effects of these drugs on MK-801 induced hyperlocomotion were indicated in Figure 1.

[00355] Sucrose preference test

[00356] To assess the anhedonia, one of the negative symptoms of schizophrenia, all mice were
5 underwent 4-day testing. In the beginning of the sucrose preference, all mice were deprived of water for 23 hours in all the experiment from one day before the first day. On the first day, each mouse was given free access to 2 identical bottles with water for 1 hour. Then, the 2 identical bottles were replaced that one filled with 1 % (wt/vol) sucrose solution and the other with water on the second day. On the third and fourth day, each mouse was received MK-801 and RS-D7
10 treatment before the experiment, and also free accessed to bottles for 1 hour. After the experiment, the 2 bottles were weighted to measure the 1 hour consumption of sucrose solution and water. The sucrose preference percentage (SPP) was calculated using the following formula: %SPP = sucrose solution consumption (g) X 100 % / [water consumption (g) + sucrose solution consumption (g)].

[00357] Compared with the saline controls, a significant reduction of sucrose intake in the
15 sucrose preference test was observed in testing mice after acute MK-801 injection. The injection of 100, 200 and 400 mg/kg RS-D7 rescued the MK-801 induced anhedonia deficit in mice. 20 mg/kg Drug 12083 and 200 mg/kg Prodrug 28095 also normalized MK-801 induced anhedonia. As a result, different dosages of RS-D7, Drug 12083 and Prodrug 28095 rescued the anhedonia after acute MK-801 injection. The rescue effects of these drugs on MK-801 induced anhedonia
20 were indicated in Figure 2.

[00358] Prepulse inhibition


[00359] To assess the sensorimotor gating function, each mouse was tested with the SR-LAB startle apparatus (San Diego Instruments, San Diego, CA, USA). The background noise was 72 dB during testing. Each session was initiated with a 5 minutes acclimatization period followed by
25 64 trials, consisting of pulse-alone (P-alone) trials, prepulse pulse (pp + P) trials, and no stimulation (nostim) trials. A P-alone trial was a 120 dB white noise burst of 40 msec. In the pp + P trials, the 120 dB pulse was preceded (by 100 msec) by a 20 msec of white noise prepulse burst of 78 dB (PP6), 82 dB (PP10), or 90 dB (PP18). The nostim trials consisted of background noise only. The session began and ended with a block of six presentations of the P-alone trial. Between
30 these two blocks, the rest of the 52 trials were performed pseudorandomly and separated by intertrial intervals of 15 sec on average (varying between 10 and 20 sec). PPI was calculated as a

percentage of the startle response using the formula: $\%PPI = 100 \times [(P\text{-alone score}) - (pp + P\text{ score})] / (pulse\text{-alone score})$, where the pulse-alone score was the average of the pulse-alone values from the in-between block of 52 trials.

[00360] Mice with acute MK-801 injection exhibited a profound reduction of acoustic PPI. However, the injection of 100, 200, and 400 mg/kg RS-D7, 20 and 40 mg/kg Drug 12083, and 100 and 200 mg/kg Prodrug 28095 significantly alleviated MK-801 induced PPI deficit in these mice. In other words, Mice displayed a significant reduction of PPI after acute MK-801 injections and they can be normalized by all dosages of RS-D7, Drug 12083 and Prodrug 28095. The recovery rates of MK-801 induced PPI deficit were indicated in Figure 3.

CLAIMS:

1. A compound of formula (I),



wherein n is 0 or 1,

X is $\text{S}(=\text{O})-$;

A is $-\text{CH}_2-$, $-\text{CR}_c$ or N;

R_a is $-\text{O}-\text{C}(=\text{O})\text{R}_{a3}$ or $-\text{O}-\text{C}(=\text{O})-\text{T}-\text{OR}_{a4}$; wherein

R_{a3} is a linear or branched C_{1-15} alkyl, linear or branched C_{2-15} alkenyl, $-\text{T}-\text{C}_{3-10}$ cycloalkyl, $-\text{T}-\text{NHR}_{a3p}$, $-\text{T}-\text{C}_{3-10}$ cycloalkenyl, $-\text{T}-\text{C}_{6-10}$ aryl, $-\text{T}-\text{C}_{5-10}$ heteroaryl, $-\text{T}-\text{NH}-\text{C}(=\text{O})-\text{O}-\text{C}_{1-10}$ alkyl, $-\text{T}-\text{adamantyl}$ or $-\text{C}_{1-3}$ alkylene- C_{6-10} aryl where the alkylene is substituted with $-\text{T}-\text{NHR}_{a3p}$;

R_{a4} is a linear or branched C_{1-15} alkyl, linear or branched C_{2-15} alkenyl, $-\text{T}-\text{C}_{3-10}$ cycloalkyl, $-\text{T}-\text{NHR}_{a3p}$, $-\text{T}-\text{C}_{3-10}$ cycloalkenyl, $-\text{T}-\text{C}_{6-10}$ aryl, $-\text{T}-\text{C}_{5-10}$ heteroaryl, $-\text{T}-\text{NH}-\text{C}(=\text{O})-\text{O}-\text{C}_{1-10}$ alkyl, $-\text{T}-\text{adamantyl}$ or $-\text{C}_{1-3}$ alkylene- C_{6-10} aryl where the alkylene is substituted with $-\text{T}-\text{NHR}_{a3p}$;

R_{a3p} is H or an *N*-protecting group selected from Fmoc or Boc;

R_b is H, linear or branched C_{1-15} alkyl, linear or branched C_{2-15} alkenyl, C_{1-3} alkoxy- C_{1-15} alkyl-, $-\text{T}'-\text{C}_{3-10}$ cycloalkyl, $-\text{T}'-\text{C}_{3-10}$ cycloalkenyl, $-\text{T}'-\text{C}_{6-10}$ aryl or $-\text{T}'-\text{C}_{5-10}$ heteroaryl;

R_c each is independently halogen, linear or branched C_{1-15} alkyl, linear or branched C_{1-15} alkoxy, hydroxyl group, or $-\text{C}_{1-10}$ alkylene- $\text{Y}-\text{C}_{6-10}$ heteroaryl wherein $-\text{Y}-$ is $-\text{CH}_2-$, $-\text{NH}-$, $-\text{O}-$ or $-\text{S}-$;

symbol * represents the bonding position;

m is an integer from 0 to 4;
 -T- is absent, C₁₋₃ alkylene or C₂₋₃ alkenylene;
 -T'- is C₁₋₃ alkylene or C₂₋₃ alkenylene; and
 wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;
 wherein the alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, alkylene and alkenylene are each independently unsubstituted or substituted with at least one substituent;
 wherein the substituent is each independently a halogen, amino, nitro, nitroso, linear or branched C₁₋₁₅ alkyl, or linear or branched C₁₋₁₅ alkoxy or C₃₋₁₀ cycloalkyl;
 wherein the alkoxy is unsubstituted; and
 when R_b is H, the tautomers are included,
 or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1, wherein

n is 0;
 A is N;
 R_b is H;
 m is 3; and
 R_c each is independently linear or branched C₁₋₁₅ alkyl, linear or branched C₁₋₁₅ alkoxy;
 or a pharmaceutically acceptable salt.
3. The compound of any one of Claims 1 to 2, wherein n is 0.
4. The compound of Claim 1, wherein m is an integer from 0 to 3.
5. The compound of Claim 1, wherein R_a is -O-C(=O)R_{a3} wherein R_{a3} is adamantyl; linear or branched C₁₋₁₀ alkyl unsubstituted or substituted by halogen; C₁₋₄ alkoxy; -C₆₋₁₀ aryl unsubstituted or substituted by C₁₋₁₀ alkyl, nitro, C₁₋₁₅ alkoxy or halogen; C₃₋₁₀ cycloalkyl; -C₃₋₁₀ cycloalkenyl; linear or branched C₂₋₁₀ alkenyl; -C₅₋₁₀ heteroaryl; -C₁₋₃alkylene-C₃₋₁₀

cycloalkyl; C₂₋₃ alkenylene-C₆₋₁₀ aryl wherein C₆₋₁₀ aryl is unsubstituted or substituted by halogen; or -O-C(=O)-O-C₁₋₁₀ alkyl.

6. The compound of Claim 1, wherein R_a is -O-C(=O)R_{a3} wherein R_{a3} is adamantyl; linear or branched C₁₋₈ alkyl unsubstituted or substituted by halogen or a tert-butyl protecting group; C₁₋₄ alkoxy; -phenyl unsubstituted or substituted by C₁₋₆ alkyl, nitro, C₁₋₄ alkoxy or halogen; C₃₋₆ cycloalkyl; -C₃₋₆ cycloalkenyl; linear or branched C₂₋₆ alkenyl; -C₅₋₆ heteroaryl; -C₁₋₃ alkylene-C₃₋₆ cycloalkyl; C₂₋₃ alkenylene-phenyl wherein phenyl is unsubstituted or substituted by halogen; or -O-C(=O)-O-C₁₋₄ alkyl.

7. The compound of Claim 1, wherein R_a is -O-C(=O)-C₁₋₆ alkyl, -O-C(=O)-C₁₋₄ alkenylene-NHFmoc or -O-C(=O)-C₁₋₄alkylene-NHBoc, or -O-C(=O)-NH-C(=O)-O-C₁₋₁₀ alkyl.

8. The compound of Claim 1, wherein R_c each is independently halogen, linear or branched C₁₋₆ alkyl, linear or branched C₁₋₆ alkoxy, or -C₁₋₁₀ alkenylene-Y-C₆₋₁₀ heteroaryl; wherein Y is S and C₆₋₁₀ heteroaryl is unsubstituted or substituted by C₁₋₁₅ alkyl, C₁₋₁₅ alkoxy, -OH, -NH₂, -NO₂ or halogen.

9. The compound of Claim 1, which is selected from the group consisting of:

21122: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acetate)

21124: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl benzoate)

26096: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl butyrate)

26097: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohexanecarboxylate)

26098: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-butylbenzoate)

21127: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-methylbenzoate)

27076: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl hexanoate

27077: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl isobutyrate

27078: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-ene-1-carboxylate)

27079: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclohex-3-enecarboxylate

28087: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methylbenzoate)

28091: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-nitrobenzoate)

28092: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclopropanecarboxylate)

28093: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-ethylbutanoate)

28094: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-phenylacetate)

28095: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3,5,5-trimethylhexanoate)

28096: (2-(((5-methoxy-4,6-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-ethoxybenzoate)

21123: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl propionate)

21125: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-chlorobenzoate)

21126: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-nitrobenzoate)

21128: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl heptanoate

21129: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-fluorobenzoate)

21130: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (Z)-2-methylbut-2-enoate)

21131: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-chloropropanoate)

21132: tert-butyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate

12124: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (Z)-but-2-enoate)

12125: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-methylbut-2-enoate)

12122: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl furan-2-carboxylate)

12123: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl acrylate)

12127: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methylbutanoate)

12128: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-cyclopentylpropanoate)

12129: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (E)-3-(2-chlorophenyl)acrylate)

12130: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 6-bromohexanoate)

11021: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-fluorobenzoate)

11020: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-methoxybenzoate)

11022: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl (3r,5r,7r)-adamantane-1-carboxylate)

11023: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl isoxazole-5-carboxylate)

11030: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-(tert-butyl)benzoate)

11031: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3-chloro-4-fluorobenzoate)

25015: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pivalate)

25016: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl pentanoate)

25017: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 4-nitrobenzoate)

25027: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl cyclobutanecarboxylate)

25028: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl thiophene-2-carboxylate)

25029: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methylbutanoate)

25030: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 3,3-dimethylbutanoate)

25031: (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl 2-methoxyacetate) and


25032: (ethyl (2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-5-yl) carbonate)

or a pharmaceutically acceptable salt thereof.

10. The compound of Claim 1, which is:

13001: 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)sulfinyl)-1H-benzo[d]imidazol-6-yl 2-((tert-butoxycarbonyl)amino)acetate
or a pharmaceutically acceptable salt thereof.

11. A pharmaceutical composition, comprising a compound of any one of Claims 1 to 10 in an admixture with a pharmaceutically acceptable carrier or diluent.
12. A method of inhibiting a DAAO, comprising contacting in vitro a cell with a compound of any one of Claims 1 to 10.
13. A compound of any one of claims 1 to 10 for use in the treatment or prevention of a disease associated with DAAO inhibition in a subject.
14. The compound of Claim 13, wherein the disease is symptom domains of schizophrenia and schizoaffective disorder, depression, Tourette Syndrome, Post-traumatic stress disorder (PTSD), Obsessive-compulsive disorder (OCD), analgesics, loss of memory and/or cognition associated with neurodegenerative diseases or loss of neuronal function characteristic of neurodegenerative diseases.
15. The compound of Claim 14, wherein the symptom domains of schizophrenia and schizoaffective disorder are negative, cognitive, depressive, positive or general psychopathology symptom domains.
16. The compound of Claim 13, wherein the disease is mild cognitive impairment (MCI), Alzheimer's disease, Parkinson's disease or schizophrenia.
17. The compound of Claim 13, wherein the disease associated with DAAO inhibition is pain, ataxia or convulsion.
18. A compound of formula (I),



wherein n is 0 or 1,

X is -S-;

A is N;

R_a is -C(=O)OR_{a1}; wherein

R_{a1} is H or linear or branched C₁₋₁₅alkyl;

R_b is H, linear or branched C₁₋₁₅alkyl, linear or branched C₂₋₁₅alkenyl, C₁₋₃alkoxy-C₁₋₁₅alkyl-, -T'-C₃₋₁₀cycloalkyl, -T'-C₃₋₁₀cycloalkenyl, -T'-C₆₋₁₀aryl or -T'-C₅₋₁₀heteroaryl;

R_c each is independently linear or branched C₁₋₁₅alkyl, linear or branched C₁₋₁₅alkoxyl, hydroxyl group, or -C₁₋₁₀alkylene-Y-C₆₋₁₀heteroaryl wherein -Y- is -CH₂-, -NH-, -O- or -S-;

m is an integer from 0 to 4;

-T- is absent, C₁₋₃alkylene or C₂₋₃alkenylene;

-T'- is C₁₋₃alkylene or C₂₋₃alkenylene; and

wherein the heteroaryl contains at least one heteroatom, each heteroatom being independently S, N or O;

wherein the alkyl, alkenyl, alkoxy, cycloalkyl, aryl, heteroaryl, alkylene and alkenylene are each independently unsubstituted or substituted with at least one substituent;

wherein the substituent is each independently a halogen, amino, nitro, nitroso, linear or branched C₁₋₁₅alkyl, or linear or branched C₁₋₁₅alkoxy or C₃₋₁₀cycloalkyl; and

when R_b is H, the tautomers are included,

or a pharmaceutically acceptable salt thereof.

19. The compound of Claim 18, wherein n is 0.

20. The compound of Claim 18 or Claim 19, wherein m is an integer from 0 to 3.
21. The compound of any one of Claims 18 to 20, wherein R_a is –C(=O)OH, or –C(=O)OC₁₋₄alkyl.
22. The compound of any one of Claims 18 to 21, wherein R_c each is independently halogen, linear or branched C₁₋₆alkyl, linear or branched C₁₋₆alkoxyl, or –C₁₋₁₀alkenylene-Y-C₆₋₁₀heteroaryl; wherein Y is S and C₆₋₁₀heteroaryl is unsubstituted or substituted by C₁₋₁₅alkyl, C₂₋₁₅alkenyl, C₁₋₁₅alkoxy, –OH, –NH₂, –NO₂ or halogen.
23. The compound of Claim 18, which is selected from the group consisting of:
 12082: Methyl 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylate
 12083: 1-(2-(cyclohex-1-en-1-yl)ethyl)-2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1H-benzo[d]imidazole-5-carboxylic acid, and
 12088: methyl 2-(((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)thio)-1-octyl-1H-benzo[d]imidazole-5-carboxylate,
 or a pharmaceutically acceptable salt thereof.
24. A pharmaceutical composition, comprising a compound of any one of Claims 18 to 23.
25. A use of the compound as defined in any one of Claims 18 to 23 for inhibiting a DAAO in a cell.
26. The compound of any one of Claims 1 to 23 for use in the treatment or prevention of the treatment or prevention of a disease associated with DAAO inhibition in a subject.
27. The compound of Claim 26, wherein the disease is symptom domains of schizophrenia and schizoaffective disorder, depression, Tourette Syndrome, Post-traumatic stress disorder (PTSD), Obsessive-compulsive disorder (OCD), analgesics, loss of memory and/or cognition

associated with neurodegenerative diseases or loss of neuronal function characteristic of neurodegenerative diseases.

28. The compound of Claim 27, wherein the symptom domains of schizophrenia and schizoaffective disorder comprise one or more of negative, cognitive, depressive, positive and general psychopathology symptom domains.

29. The compound of Claim 26, wherein the disease is mild cognitive impairment (MCI), Alzheimer's disease, Parkinson's disease or schizophrenia.

30. The compound of Claim 26, wherein the disease associated with DAAO inhibition is pain, ataxia or convulsion.




Fig. 1




Fig. 2




Fig. 3

