


(51) **International Patent Classification:**
C12N 9/92 (2006.01) C12P 7/06 (2006.01)**YANG, Jianjun** [US/US]; 7 Kent Drive, Hockessin, Delaware 19707 (US).(21) **International Application Number:**
PCT/US201 1/042132(74) **Agent: FELTHAM, S., Neil;** E. I. du Pont de Nemours and Company, Legal Patent Records Center, 4417 Lancaster Pike, Wilmington, Delaware 19805 (US).(22) **International Filing Date:**
28 June 2011 (28.06.2011)(25) **Filing Language:** English(26) **Publication Language:** English(30) **Priority Data:**
61/359,463 29 June 2010 (29.06.2010) US(71) **Applicant (for all designated States except US): E. I. DU PONT DE NEMOURS AND COMPANY** [US/US]; 1007 Market Street, Wilmington, Delaware 19898 (US).(72) **Inventors; and**(75) **Inventors/Applicants (for US only): KAHSAY, Robel, Y.** [ER/US]; 1304 W. 13th Street, Apt. 3, Wilmington, Delaware 19806 (US). **QI, Min** [US/US]; 11 Stuyvesant Drive, Hockessin, Delaware 19707 (US). **TAO, Luan** [US/US]; 404 Karen Lane, Wallingford, Pennsylvania 19086 (US). **VHTANEN, Paul, V.** [US/US]; 1138 Country Club Road, West Chester, Pennsylvania 19382 (US).(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) **Title:** IMPROVED XYLOSE UTILIZATION IN RECOMBINANT ZYMONONAS HAVING ADDITIONAL XYLOSE ISOMERASE ACTIVITY

Figure 1

(57) **Abstract:** *Zymomonas* expressing xylose isomerase from *A. missouriensis* was found to have improved xylose utilization, growth, and ethanol production when grown in media containing xylose. Xylose isomerases related to that of *A. missouriensis* were identified structurally through molecular phylogenetic and Profile Hidden Markov Model analyses, providing xylose isomerases that may be used to improve xylose utilization.



Published:

— with sequence listing part of description (Rule 5.2(a))

-with international search report (Art. 21(3))

TITLE**IMPROVED XYLOSE UTILIZATION IN RECOMBINANT ZYMOMONAS HAVING
ADDITIONAL XYLOSE ISOMERASE ACTIVITY**

This application claims the benefit of United States Provisional
5 Application 61/359463, filed June 29, 2010 and is incorporated by
reference in its entirety.

STATEMENT OF GOVERNMENT RIGHTS

10 This invention was made with United States government
support under Contract Nos. DE-FC36-07GO17056 awarded by the
Department of Energy. The government has certain rights in this
invention.

15 FIELD OF THE INVENTION

The invention relates to the fields of microbiology and genetic
engineering. More specifically, xylose isomerases with high activity in
Zymomonas were identified that provide for improved xylose utilization
and ethanol production.

20

BACKGROUND OF THE INVENTION

Production of ethanol by microorganisms provides an alternative
energy source to fossil fuels and is therefore an important area of current
research. It is desirable that microorganisms producing ethanol, as well as
25 other useful products, be capable of using xylose as a carbon source
since xylose is the major pentose in hydrolyzed lignocellulosic biomass.
Biomass can provide an abundantly available, low cost carbon substrate.
Zymomonas mobilis and other bacterial ethanologens which do not
naturally utilize xylose have been genetically engineered for xylose
30 utilization by introduction of genes encoding 1) xylose isomerase, which
catalyses the conversion of xylose to xylulose; 2) xylulokinase, which
phosphorylates xylulose to form xylulose 5-phosphate; 3) transketolase;
and 4) transaldolase. Typically the coding regions used were from *E. coli*
genes.

There has been success in engineering *Z. mobilis* strains for xylose metabolism (US 5514583, US 571 2133, US 6566107, WO 95/28476, Feldmann et al. (1992) Appl. Microbiol. Biotechnol. 38: 354-361, Zhang et al. (1995) Science 267:240-243), as well as a *Zymobacter palmae* strain (Yanase et al. (2007) Appl. Environ. Microbiol. 73:2592-2599). However, typically the engineered strains do not grow and produce ethanol as well on xylose as on glucose. Strains engineered for xylose utilization have been adapted by serial passage on xylose medium, resulting in strains with improved xylose utilization as described in U. S. Pat. 7,223,575 and U. S. Pat. 7,741,119. Disclosed in commonly owned and co-pending US Patent Application Publication US 20090246846A1 is engineering for improved xylose utilization by expression of *E. coli* xylose isomerase from a mutated, highly active *Zymomonas mobilis* glyceraldehyde-3-phosphate dehydrogenase gene promoter (*Pgap*).

There remains a need for strains of *Zymomonas*, and other bacterial ethanolagens, which have further improvement in xylose utilization.

SUMMARY OF THE INVENTION

The invention provides recombinant xylose-utilizing *Zymomonas* or *Zymobacter* cells that express a highly active xylose isomerase providing improved xylose utilization and ethanol production.

Accordingly, the invention provides a recombinant bacterial strain selected from the group consisting of *Zymomonas* and *Zymobacter* comprising a heterologous nucleic acid molecule encoding a polypeptide having xylose isomerase activity wherein the polypeptide is a Group I xylose isomerase and is included in the class of enzymes identified by EC 5.3.1.5, and wherein the strain utilizes xylose as a carbon source.

Xylose isomerase enzymes useful in the invention are those that have an E-value score of 1E-15 or less when queried using a Profile Hidden Markov Model prepared using SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142; the query being carried out using the hmmsearch algorithm wherein the Z parameter is set to 1 billion, or those that have at least 90% identity

to an amino acid sequence selected from the group consisting of SEQ ID
 NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38,
 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76,
 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110,
 5 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 131, 132, 133, 134,
 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, and 147
 based on the Clustal W method of alignment using the default parameters
 of GAP PENALTY=10, GAP LENGTH PENALTY=0.1, and Gonnet 250
 series of protein weight matrix; or those that have the following conserved
 10 amino acids, or 90% of the following conserved amino acids, when
 compared with the reference amino acid sequence of SEQ ID NO:66:

- a) leucine at position 226,
- b) methionine at position 223,
- c) isoleucine at position 191,
- 15 d) threonine, serine, or valine at position 195,
- e) methionine, threonine or glutamine at position 88,
- f) histidine at position 290,
- g) glutamic acid or aspartic acid at position 221,
- h) phenylalanine, valine, or leucine at position 242,
- 20 i) histidine at position 243,
- j) leucine, phenylalanine, or methionine at position 193,
- k) glutamine at position 256,
- l) glycine at position 213,
- m) proline, tyrosine, alanine, or serine at position 288, and
- 25 n) glutamine at position 249.

In another embodiment the invention provides a process for
 improving xylose utilization in a recombinant bacterial cell comprising:

- a) providing a recombinant bacterial strain selected from the group
 consisting of *Zymomonas* and *Zymobacter* comprising a xylose
 30 utilization pathway comprising a xylose isomerase not belonging to
 Group I; and
- b) introducing a heterologous nucleic acid molecule encoding a
 polypeptide having xylose isomerase activity wherein the
 polypeptide is a Group I xylose isomerase.

Alternatively the invention provides a method for the production of ethanol comprising:

- a) providing the recombinant bacterial strain of the invention; and
- b) contacting the strain of (a) with xlyose under conditions whereby the strain produces ethanol.

BRIEF DESCRIPTION OF THE FIGURES AND SEQUENCE DESCRIPTIONS

Figure 1 shows a diagram of the ethanol fermentation pathway in *Zymomonas* engineered for xylose utilization.

Figure 2 is a diagram of a phylogenetic tree of xylose isomerases showing Group I and Group II branches.

Figure 3 is a diagram of a phylogenetic tree of Group I xylose isomerases.

Figure 4 is a graph of growth curves for a control ZW641 strain and strains of ZW641 transformed with a gene having a codon-optimized coding region for expression of xylose isomerase from *A. missourinensis* (355-1 , 355-2), *L. brevis* (356-1 , 356-2), or *E. coli* (357-1 , 357-2).

Figure 5 is a graph of xylose isomerase activities in a control ZW641 strain and strains of ZW641 transformed with a gene having a codon-optimized coding region for expression of xylose isomerase from *A. missourinensis* (355-1), *L. brevis* (356-2), or *E. coli* (357-2) measured by the Cysteine-Carboazole method.

Figure 6 is a graph of growth curves for a control ZW641 strain and strains of ZW641 transformed with a gene having a codon-optimized coding region for expression of xylose isomerase from *A. missourinensis* (AMxylA), *E. coli* (ECxylA), *Geodermatophilus obscurus* (GOxylA), *Mycobacterium smegmatis* (MSxylA), *Salinispora arenicola* (SAxylA), or *Xylanimonas cellulosilytica* (XCxylA).

Figure 7 is a graph of xylose isomerase activities in a control ZW641 strain and strains of ZW641 transformed with a gene having a codon-optimized coding region for expression of xylose isomerase from *A. missourinensis* (AMxylA), *E. coli* (ECxylA), *Geodermatophilus obscurus* (GOxylA), *Mycobacterium smegmatis* (MSxylA), *Salinispora arenicola*

(SAxylA), or *Xylanimonas cellulosilytica* (XCxylA) measured by the Cysteine-Carboazole method.

Table 1 is a table of the Profile HMM for xylose isomerase Group I proteins. Table 1 is submitted herewith electronically and is incorporated
5 herein by reference.

Table 2 is a table of the E-value scores for XI proteins, each identified by a SEQ ID NO, that were queried using the Group I profile HMM. Table 2 is submitted herewith electronically and is incorporated
herein by reference

10 The invention can be more fully understood from the following detailed description and the accompanying sequence descriptions which form a part of this application.

The following sequences conform with 37 C.F.R. 1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences
15 and/or Amino Acid Sequence Disclosures - the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST.25 (1998) and the sequence listing requirements of the EPO and PCT (Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administrative Instructions). The symbols and format used for nucleotide
20 and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

Table 3. SEQ ID numbers of Coding Regions and Proteins for Group I xylose isomerases. Uniprot accession number (AC) given for proteins that
25 are seeds and NCBI GI number for given those that are not seeds.

Organism	GI or AC #	SEQ ID NO: Nucleic acid	SEQ ID NO: Amino acid
<i>Clavibacter michiganensis</i>	B0RIF1	1	2
<i>Arthrobacter chlorophenolicus</i>	220912923	3	4
<i>Actinosynnema mirum</i>	226865307	5	6
<i>Kribbella flavida</i>	227382478	7	8
<i>Mycobacterium smegmatis</i>	118469437	9	10

<i>Arthrobacter sp.</i>	60615686	11	12
<i>Actinomyces urogenitalis</i>	2274971 16	13	14
<i>Streptomyces ambofaciens</i>	126348424	15	16
<i>Salinispora arenicola</i>	159039501	17	18
<i>Streptomyces sp.</i>	38141596	19	20
<i>Meiothermus silvanus</i>	227989553	21	22
<i>Actinoplanes sp.</i>	P10654	23	24
<i>Mobiluncus curtisii</i>	227493823	25	26
<i>Herpetosiphon aurantiacus</i>	159898286	27	28
<i>Acidothermus cellulolyticus</i>	117929271	29	30
<i>Streptomyces coelicolor</i>	Q9L0B8	31	32
<i>Streptomyces avermitilis</i>	Q93HF3	33	34
<i>Nocardioopsis dassonvillei</i>	229207664	35	36
<i>Nakamurella multipartita</i>	229221673	37	38
<i>Xylanimonas cellulosilytica</i>	227427650	39	40
<i>Clavibacter michiganensis</i>	A5CPC1	41	42
<i>Salinispora tropica</i>	145596104	43	44
<i>Streptomyces sp.</i>	197764953	45	46
<i>Streptomyces pristinaespiralis</i>	197776540	47	48
<i>Roseiflexus sp.</i>	148656997	49	50
<i>Meiothermus ruber</i>	227992647	51	52
<i>Arthrobacter sp.</i>	P12070	53	54
<i>Thermobaculum terrenum</i>	227374836	55	56
<i>Janibacter sp.</i>	84495191	57	58
<i>Brachybacterium faecium</i>	237671435	59	60
<i>Beutenbergia cavernae</i>	229821786	61	62
<i>Geodermatophilus obscurus</i>	227404617	63	64
<i>Actinoplanes missouriensis</i>	P12851	65	66
<i>Streptomyces violaceusniger</i>	P09033	67	68
<i>Actinomyces odontolyticus</i>	154508186	69	70
<i>Mobiluncus mulieris</i>	227875705	71	72
<i>Cellulomonas flavigena</i>	229243977	73	74

<i>Saccharomonospora viridis</i>	229886404	75	76
<i>Streptomyces lividans</i>	Q9RFM4	77	78
<i>Frankia</i> sp.	158316430	79	80
<i>Streptosporangium roseum</i>	229851079	81	82
<i>Nocardioides</i> sp.	119716602	83	84
<i>Kribbella flavida</i>	227381155	85	86
<i>Roseiflexus castenholzii</i>	156742580	87	88
<i>Arthrobacter aurescens</i>	119964059	89	90
<i>Leifsonia xyli</i>	50954171	91	92
<i>Jonesia denitrificans</i>	227383768	93	94
<i>Streptomyces olivaceoviridis</i>	Q93RJ9	95	96
<i>Stackebrandtia nassauensis</i>	229862570	97	98
<i>Thermus thermophilus</i>	P26997	99	100
<i>Acidobacteria bacterium</i>	94967932	101	102
<i>Catenulispora acidiphila</i>	229246901	103	104
<i>Streptomyces corchorusii</i>	Q9S3Z4	105	106
<i>Streptomyces thermocyaneoviolaceus</i>	Q9L558	107	108
<i>marine actinobacterium</i>	88856315	109	110
<i>Micromonospora</i> sp.	237882534	111	112
<i>Thermobifida fusca</i>	72162004	113	114
<i>Herpetosiphon aurantiacus</i>	159897776	115	116
<i>Streptomyces griseus</i>	182434863	117	118
<i>Mycobacterium vanbaalenii</i>	120406242	119	120
<i>Streptomyces diastaticus</i>	P50910	121	122
<i>Deinococcus geothermalis</i>	94972159	123	124
<i>Arthrobacter</i> sp.	A0JXN9	125	126
<i>Streptomyces rubiginosus</i>	P24300	127	128
<i>Streptomyces murinus</i>	P37031	129	130
<i>Thermus caldophilus</i>	4930285	*	131
<i>Thermus caldophilus</i>	P56681	*	132
<i>Arthrobacter</i> sp.	231103	*	133
<i>Actinoplanes missouriensis</i>	443486	*	134

<i>Streptomyces olivochromogenes</i>	P15587	*	135
<i>Streptomyces olivochromogenes</i>	157879319	*	136
<i>Streptomyces rochei</i>	P22857	*	137
<i>Streptomyces olivochromogenes</i>	157881044	*	138
<i>Streptomyces diastaticus</i>	7766813	*	139
<i>Actinoplanes missouriensis</i>	349936	*	140
<i>Arthrobacter sp.</i>	2914276	*	141
<i>Streptomyces albus</i>	P24299	*	142
<i>Actinoplanes missouriensis</i>	443303	*	143
<i>Streptomyces diastaticus</i>	9256915	*	144
<i>Actinoplanes missouriensis</i>	443526	*	145
<i>Streptomyces rubiginosus</i>	21730246	*	146
<i>Actinoplanes missouriensis</i>	443568	*	147

* Sequence not readily available

Table 4. SEQ ID numbers of Proteins for Group II seed xylose isomerases. Uniprot accession number (AC) given for the seed proteins.

Organism	AC #	SEQ ID NO: Amino acid
<i>Salmonella enterica</i>	B4T952	148
<i>Klebsiella pneumoniae</i>	P29442	149
<i>Sinorhizobium meliloti</i>	Q92LW9	150
<i>Escherichia coli</i>	Q7A9X4	151
<i>Salmonella enterica</i>	Q5PLM6	152
<i>Xanthomonas campestris</i>	Q3BMF2	153
<i>Pectobacterium atrosepticum</i>	Q6DB05	154
<i>Rhodopirellula baltica</i>	Q7UVG2	155
<i>Xanthomonas axonopodis</i>	Q8PEW5	156
<i>Xanthomonas oryzae</i>	Q5GUF2	157
<i>Pediococcus pentosaceus</i>	Q03HN1	158
<i>Brucella suis</i>	Q8G204	159
<i>Escherichia coli</i>	Q0TBN7	160
<i>Bifidobacterium longum</i>	Q8G3Q1	161
<i>Brucella canis</i>	A9M9H3	162

<i>Burkholderia multivorans</i>	A9ARG7	163
<i>Brucella ovis</i>	A5VPA1	164
<i>Rhizobium etli</i>	B3Q0R5	165
<i>Burkholderia xenovorans</i>	Q13RB8	166
<i>Actinobacillus pleuropneumoniae</i>	A3N3K2	167
<i>Burkholderia cenocepacia</i>	B4ENA5	168
<i>Solibacter usitatus</i>	Q022S9	169
<i>Brucella abortus</i>	B2SA37	170
<i>Rhodobacter sphaeroides</i>	A4WVT8	171
<i>Thermoanaerobacter sp.</i>	B0K1L3	172
<i>Yersinia pseudotuberculosis</i>	Q1C0D3	173
<i>Xanthomonas oryzae</i>	Q5GYQ7	174
<i>Bifidobacterium longum</i>	B3DR33	175
<i>Thermoanaerobacter pseudethanolicus</i>	P22842	176
<i>Photobacterium profundum</i>	Q6LUY7	177
<i>Escherichia coli</i>	B1LJC7	178
<i>Agrobacterium tumefaciens</i>	Q8U7G6	179
<i>Tetragenococcus halophilus</i>	O82845	180
<i>Salmonella enterica</i>	B4TZ55	181
<i>Yersinia pseudotuberculosis</i>	Q8Z9Z1	182
<i>Yersinia pseudotuberculosis</i>	Q1CDB8	183
<i>Rhodobacter sphaeroides</i>	A3PNM4	184
<i>Brucella abortus</i>	Q2YMQ2	185
<i>Salmonella enterica</i>	Q8ZL90	186
<i>Bacteroides vulgatus</i>	A6L792	187
<i>Xanthomonas axonopodis</i>	Q8PLL9	188
<i>Salmonella enterica</i>	Q57IG0	189
<i>Escherichia coli</i>	B7M3I8	190
<i>Roseobacter denitrificans</i>	Q162B6	191
<i>Bacteroides fragilis</i>	Q64U20	192
<i>Enterobacter sakazakii</i>	A7MNI5	193
<i>Brucella abortus</i>	Q57EI4	194
<i>Geobacillus thermodenitrificans</i>	A4IP67	195
<i>Bacteroides thetaiotaomicron</i>	Q8A9M2	196
<i>Haemophilus influenzae</i>	A5UCZ3	197

<i>Yersinia pseudotuberculosis</i>	B2K7D2	198
<i>Xanthomonas campestris</i>	Q4UTU6	199
<i>Haemophilus somnus</i>	B0UT19	200
<i>Pseudoalteromonas atlantica</i>	Q15PG0	201
<i>Escherichia fergusonii</i>	B7LTH9	202
<i>Silicibacter</i> sp.	Q1GKQ4	203
<i>Salmonella enterica</i>	B5R4P8	204
<i>Bifidobacterium adolescentis</i>	A1A0H0	205
<i>Staphylococcus xylosus</i>	P27157	206
<i>Thermotoga maritima</i>	Q9X1Z5	207
<i>Salmonella enterica</i>	A9MUV0	208
<i>Pseudomonas syringae</i>	Q48J73	209
<i>Shigella boydii</i>	Q31V53	210
<i>Burkholderia ambifaria</i>	Q0B1U7	211
<i>Bacillus amyloliquefaciens</i>	A7Z522	212
<i>Haemophilus influenzae</i>	A5UIN7	213
<i>Bacillus megaterium</i>	O08325	214
<i>Arabidopsis thaliana</i>	Q9FKK7	215
<i>Escherichia coli</i>	Q3YVV0	216
<i>Bacteroides fragilis</i>	Q5LCV9	217
<i>Pseudomonas fluorescens</i>	Q3KDW0	218
<i>Escherichia coli</i>	B1X8I1	219
<i>Bacillus subtilis</i>	P04788	220
<i>Xanthomonas campestris</i>	Q4UNZ4	221
<i>Pseudomonas syringae</i>	Q4ZSF5	222
<i>Sinorhizobium medicae</i>	A6UD89	223
<i>Ochrobactrum anthropi</i>	A6X4G3	224
<i>Burkholderia thailandensis</i>	Q2SW40	225
<i>Salmonella enterica</i>	B5EX72	226
<i>Thermotoga</i> sp.	B1LB08	227
<i>Bacillus cereus</i>	Q739D2	228
<i>Salmonella enterica</i>	B4SWK9	229
<i>Salmonella enterica</i>	Q7C637	230
<i>Enterococcus faecalis</i>	Q7C3R3	231
<i>Thermotoga neapolitana</i>	P45687	232

<i>Escherichia coli</i>	B7MES1	233
<i>Photorhabdus luminescens</i>	Q7N4P7	234
<i>Enterobacter</i> sp.	A4W566	235
<i>Burkholderia cenocepacia</i>	B1KB47	236
<i>Bacillus licheniformis</i>	P77832	237
<i>Geobacillus stearothermophilus</i>	P54273	238
<i>Brucella abortus</i>	Q8YFX5	239
<i>Rhizobium leguminosarum</i>	Q1MBL8	240
<i>Yersinia enterocolitica</i>	A1JT10	241
<i>Serratia proteamaculans</i>	A8G7W8	242
<i>Yersinia pseudotuberculosis</i>	A7FP68	243
<i>Escherichia coli</i>	B7NEL7	244
<i>Yersinia pestis</i>	A9R5Q1	245
<i>Fervidobacterium gondwanense</i>	Q6T6K9	246
<i>Xanthomonas campestris</i>	Q8P3H1	247
<i>Rhizobium leguminosarum</i>	B5ZQV6	248
<i>Bradyrhizobium japonicum</i>	Q89VC7	249
<i>Mesorhizobium</i> sp.	Q11EH9	250
<i>Actinobacillus pleuropneumoniae</i>	B3H2X9	251
<i>Yersinia pseudotuberculosis</i>	Q663Y3	252
<i>Xanthomonas campestris</i>	Q8P9T9	253
<i>Burkholderia cenocepacia</i>	A0KE56	254
<i>Oceanobacillus iheyensis</i>	Q8ELU7	255
<i>Brucella suis</i>	B0CKM9	256
<i>Thermoanaerobacterium thermosaccharolyticum</i>	P29441	257
<i>Burkholderia phymatum</i>	B2JFE9	258
<i>Yersinia pseudotuberculosis</i>	B1JH40	259
<i>Bacillus</i> sp.	P54272	260
<i>Lactococcus lactis</i>	Q02Y75	261
<i>Novosphingobium aromaticivorans</i>	Q2GAB9	262
<i>Lactobacillus brevis</i>	P29443	263
<i>Mesorhizobium loti</i>	Q98CR8	264
<i>Escherichia coli</i>	A8A623	265
<i>Burkholderia cenocepacia</i>	Q1BG90	266
<i>Thermoanaerobacterium thermosulfurigenes</i>	P19148	267

<i>Thermotoga petrophila</i>	A5ILR5	268
<i>Lactobacillus pentosus</i>	P21938	269
<i>Lactococcus lactis</i>	Q9CFG7	270
<i>Ruminococcus flavefaciens</i>	Q9S306	271
<i>Burkholderia phytofirmans</i>	B2T929	272
<i>Salmonella enterica</i>	B5FLD6	273
<i>Lactobacillus brevis</i>	Q03TX3	274
<i>Burkholderia ambifaria</i>	B1Z405	275
<i>Salmonella enterica</i>	B5RGL6	276
<i>Bacillus halodurans</i>	Q9K993	277
<i>Bacillus clausii</i>	Q5WKJ3	278
<i>Marinomonas sp.</i>	A6VWH1	279
<i>Yersinia pseudotuberculosis</i>	A4TS63	280
<i>Actinobacillus pleuropneumoniae</i>	B0BT19	281
<i>Silicibacter pomeroyi</i>	Q5LV46	282
<i>Xanthomonas oryzae</i>	Q2NXR2	283
<i>Thermoanaerobacterium saccharolyticum</i>	P30435	284
<i>Escherichia coli</i>	B6I3D6	285
<i>Escherichia coli</i>	B5YVL8	286
<i>Escherichia coli</i>	B7NP65	287
<i>Escherichia coli</i>	B2U560	288
<i>Escherichia coli</i>	B1IZM7	289
<i>Rhizobium etli</i>	Q2K433	290
<i>Escherichia coli</i>	P00944	291
<i>Hordeum vulgare</i>	Q40082	292
<i>Dinoroseobacter shibae</i>	A8LP53	293
<i>Rhodobacter sphaeroides</i>	Q3IYM4	294
<i>Actinobacillus succinogenes</i>	A6VLM8	295
<i>Bacillus pumilus</i>	A8FE33	296
<i>Escherichia coli</i>	Q8FCE3	297
<i>Pseudomonas syringae</i>	Q880Z4	298
<i>Burkholderia vietnamiensis</i>	A4JSU5	299
<i>Escherichia coli</i>	A7ZTB2	300
<i>Haemophilus influenzae</i>	P44398	301
<i>Haemophilus influenzae</i>	Q4QLI2	302

<i>Listeria welshimeri</i>	A0AF79	303
<i>Thermoanaerobacter yonseiensis</i>	Q9KGU2	304
<i>Geobacillus kaustophilus</i>	Q5KYS6	305
<i>Mannheimia succiniciproducens</i>	Q65PY0	306

SEQ ID NO:307 is the nucleotide sequence of the LoxPw-aadA-LoxPw DNA fragment PCR product.

SEQ ID NO:308 is the coding region for the *Actinoplanes missourinesis* xylose isomerase that was codon optimized for *Zymomonas*.

SEQ ID NO:309 is the coding region for the *Lactobacillus brevis* xylose isomerase that was codon optimized for *Zymomonas*.

SEQ ID NO:310 is the coding region for the *E. coli* xylose isomerase that was codon optimized for *Zymomonas*.

SEQ ID NO:311 is the nucleotide sequence of the glyceraldehyde-3-phosphate dehydrogenase gene promoter from *Z. mobilis* (*ZmPgap*).

SEQ ID NO:312 is the nucleotide sequence of the terminator from the *Z. mobilis* L-ribulose 5 phosphate 4-epimerase gene.

SEQ ID NO:313 is the nucleotide sequence of the pARA354 plasmid.

SEQ ID NO:314-327, 329, 330, and 332-334 are nucleotide sequences of PCR and sequencing primers.

SEQ ID NO:328 is the nucleotide sequence of the LDH-L DNA fragment PCR product.

SEQ ID NO:331 is the nucleotide sequence of the LDH-R DNA fragment PCR product.

SEQ ID NO:335 is the nucleotide sequence of the codon optimized coding region for *Geodermatophilus obscurus* xylose isomerase.

SEQ ID NO:336 is the nucleotide sequence of the codon optimized coding region for *Mycobacterium smegmatis* xylose isomerase.

SEQ ID NO:337 is the nucleotide sequence of the codon optimized coding region for *Salinispora arenicola* xylose isomerase.

SEQ ID NO:338 is the nucleotide sequence of the codon optimized coding region for *Xylanimonas cellulosilytica* xylose isomerase.

SEQ ID NO:339 is the nucleotide sequence of the promoter $P_{gap\bar{s}}$ used in chimeric gene constructions.

SEQ ID NOS:340-345 are nucleotide sequences of PCR and sequencing primers.

5

DETAILED DESCRIPTION

Disclosed herein are xylose isomerase enzymes that are highly active in *Zymomonas*, and may be used to increase xylose utilization and ethanol production. Ethanol is an important compound for use in replacing fossil fuels.

10

The following definitions may be used for the interpretation of the claims and specification:

As used herein, the terms "comprises," "comprising," "includes," "including," "has," "having," "contains" or "containing," or any other variation thereof, are intended to cover a non-exclusive inclusion. For example, a composition, a mixture, process, method, article, or apparatus that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, "or" refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

15

20

Also, the indefinite articles "a" and "an" preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e. occurrences) of the element or component. Therefore "a" or "an" should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

25

The term "invention" or "present invention" as used herein is a non-limiting term and is not intended to refer to any single embodiment of the particular invention but encompasses all possible embodiments as described in the specification and the claims.

30

As used herein, the term "about" modifying the quantity of an ingredient or reactant of the invention employed refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use
5 solutions in the real world; through inadvertent error in these procedures; through differences in the manufacture, source, or purity of the ingredients employed to make the compositions or carry out the methods; and the like. The term "about" also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial
10 mixture. Whether or not modified by the term "about", the claims include equivalents to the quantities. In one embodiment, the term "about" means within 10% of the reported numerical value, preferably within 5% of the reported numerical value.

The term "carbon substrate" or "fermentable carbon substrate"
15 refers to a carbon source capable of being metabolized by host organisms of the present invention and particularly carbon sources selected from the group consisting of monosaccharides, oligosaccharides, and polysaccharides.

"Gene" refers to a nucleic acid fragment that expresses a specific
20 protein or functional RNA molecule, which may optionally include regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" or "wild type gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene,
25 comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature.
30 "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes.

The term "genetic construct" refers to a nucleic acid fragment that encodes for expression of one or more specific proteins or functional RNA molecules. In the gene construct the gene may be native, chimeric, or foreign in nature. Typically a genetic construct will comprise a "coding
5 sequence". A "coding sequence" refers to a DNA sequence that encodes a specific amino acid sequence.

"Promoter" or "Initiation control regions" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence.
10 Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of
15 development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters".

The term "expression", as used herein, refers to the transcription and stable accumulation of coding (mRNA) or functional RNA derived from
20 a gene. Expression may also refer to translation of mRNA into a polypeptide. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms.

The term "transformation" as used herein, refers to the transfer of a
25 nucleic acid fragment into a host organism, resulting in genetically stable inheritance. The transferred nucleic acid may be in the form of a plasmid maintained in the host cell, or some transferred nucleic acid may be integrated into the genome of the host cell. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or
30 "recombinant" or "transformed" organisms.

The terms "plasmid" and "vector" as used herein, refer to an extra chromosomal element often carrying genes which are not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously

replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear or circular, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction
5 which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is
10 affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

15 The term "selectable marker" means an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene's effect, i.e., resistance to an antibiotic, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic
20 acid of interest.

As used herein the term "codon degeneracy" refers to the nature in the genetic code permitting variation of the nucleotide sequence without affecting the amino acid sequence of an encoded polypeptide. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in
25 usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a gene for improved expression in a host cell, it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

The term "codon-optimized" as it refers to genes or coding regions
30 of nucleic acid molecules for transformation of various hosts, refers to the alteration of codons in the gene or coding regions of the nucleic acid molecules to reflect the typical codon usage of the host organism without altering the polypeptide encoded by the DNA.

The term "fermentable sugar" refers to oligosaccharides and

monosaccharides that can be used as a carbon source by a microorganism in a fermentation process.

The term "lignocellulosic" refers to a composition comprising both lignin and cellulose. Lignocellulosic material may also comprise hemicellulose.

The term "cellulosic" refers to a composition comprising cellulose and additional components, including hemicellulose.

The term "saccharification" refers to the production of fermentable sugars from polysaccharides.

The term "pretreated biomass" means biomass that has been subjected to thermal, physical and/or chemical pretreatment to increase accessibility of polysaccharides in the biomass prior to saccharification.

"Biomass" refers to any cellulosic or lignocellulosic material and includes materials comprising cellulose, and optionally further comprising hemicellulose, lignin, starch, oligosaccharides and/or monosaccharides. Biomass may also comprise additional components, such as protein and/or lipid. Biomass may be derived from a single source, or biomass can comprise a mixture derived from more than one source; for example, biomass could comprise a mixture of corn cobs and corn stover, or a mixture of grass and leaves. Biomass includes, but is not limited to, bioenergy crops, agricultural residues, municipal solid waste, industrial solid waste, sludge from paper manufacture, yard waste, wood and forestry waste. Examples of biomass include, but are not limited to, corn cobs, crop residues such as corn husks, corn stover, grasses, wheat, wheat straw, barley straw, hay, rice straw, switchgrass, waste paper, sugar cane bagasse, sorghum, components obtained from milling of grains, trees, branches, roots, leaves, wood chips, sawdust, shrubs and bushes, vegetables, fruits, flowers and animal manure.

"Biomass hydrolysate" refers to the product resulting from saccharification of biomass. The biomass may also be pretreated or pre-processed prior to saccharification.

The term "xylose isomerase" refers to an enzyme that catalyzes the interconversion of D-xylose and D-xylulose. Xylose isomerases (XI) belong to the group of enzymes classified as EC 5.3.1 .5.

The term "E-value", as known in the art of bioinformatics, is "Expect-value" which provides the probability that a match will occur by chance. It provides the statistical significance of the match to a sequence. The lower the E-value, the more significant the hit.

- 5 The term "Group I xylose isomerase" refers herein to a xylose isomerase protein that belongs to Group I as defined by at least one of the following criteria: a) it falls within a 50% threshold sequence identity grouping that includes the *A. missouriensis* XI that is prepared using molecular phylogenetic bioinformatics analysis as in Example 4; b) it
- 10 substantially fits the amino acids for Group I in the specificity determining positions (SDP) identified using GroupSim analysis of the Group I and Group II XI sets determined from molecular phylogenetic analysis that are given in Table 6 in Example 4; and/or c) it has an E-value of $1E-15$ or less when queried using a Profile Hidden Markov
- 15 Model prepared using SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142; where the query is carried out using the hmmsearch algorithm with the Z parameter is set to 1 billion, as in Example 4. It is understood that although "Group I" xylose isomerases are known and defined in the
- 20 literature that the definition provided herein is more precise than the literature definition and is the definition that informs the following discussion. Thus, "Group I" as used herein will refer to Applicants' definition whereas "Group II" will refer to the definitions as commonly understood in the art.
- 25 The term "heterologous" means not naturally found in the location of interest. For example, a heterologous gene refers to a gene that is not naturally found in the host organism, but that is introduced into the host organism by gene transfer. In addition, a heterologous nucleic acid molecule that is present in a chimeric gene is a nucleic acid molecule that
- 30 is not naturally found associated with the other segments of the chimeric gene, such as the nucleic acid molecules having the coding region and promoter segments not naturally being associated with each other.

As used herein, an "isolated nucleic acid molecule" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing

synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid molecule in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA.

A nucleic acid fragment is "hybridizable" to another nucleic acid
5 fragment, such as a cDNA, genomic DNA, or RNA molecule, when a single-stranded form of the nucleic acid fragment can anneal to the other nucleic acid fragment under the appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T.
10 Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory: Cold Spring Harbor, NY (1989), particularly Chapter 11 and Table 11.1 therein (entirely incorporated herein by reference). The conditions of temperature and ionic strength determine the "stringency" of the hybridization. Stringency conditions can be adjusted to screen for
15 moderately similar fragments (such as homologous sequences from distantly related organisms), to highly similar fragments (such as genes that duplicate functional enzymes from closely related organisms). Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5%
20 SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45 °C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50 °C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X
25 SSC, 0.5% SDS was increased to 60 °C. Another preferred set of highly stringent conditions uses two final washes in 0.1 X SSC, 0.1 % SDS at 65 °C. An additional set of stringent conditions include hybridization at 0.1X SSC, 0.1 % SDS, 65 °C and washes with 2X SSC, 0.1 % SDS followed by 0.1X SSC, 0.1 % SDS, for example.

30 Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the

nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability

5 (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., supra, 9.50-9.51). For hybridizations with shorter nucleic acids, i.e., oligonucleotides, the position of

10 mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., supra, 11.7-11.8). In one embodiment the length for a hybridizable nucleic acid is at least about 10 nucleotides. Preferably a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably

15 at least about 20 nucleotides; and most preferably the length is at least about 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

The term "complementary" is used to describe the relationship between

20 nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine.

The terms "homology" and "homologous" are used interchangeably herein. They refer to nucleic acid fragments wherein changes in one or

25 more nucleotide bases do not affect the ability of the nucleic acid fragment to mediate gene expression or produce a certain phenotype. These terms also refer to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially alter the functional properties of the resulting nucleic acid

30 fragment relative to the initial, unmodified fragment. It is therefore understood, as those skilled in the art will appreciate, that the invention encompasses more than the specific exemplary sequences.

Moreover, the skilled artisan recognizes that homologous nucleic acid sequences encompassed by this invention are also defined by their

ability to hybridize, under moderately stringent conditions (e.g., 0.5 X SSC, 0.1 % SDS, 60 °C) with the sequences exemplified herein, or to any portion of the nucleotide sequences disclosed herein and which are functionally equivalent to any of the nucleic acid sequences disclosed herein.

5 The term "percent identity", as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as
10 determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in: 1.) Computational Molecular Biology (Lesk, A. M., Ed.) Oxford University: NY (1988); 2.) Biocomputing: Informatics and Genome Projects (Smith, D. W., Ed.) Academic: NY
15 (1993); 3.) Computer Analysis of Sequence Data, Part I (Griffin, A. M., and Griffin, H. G., Eds.) Humana: NJ (1994); 4.) Sequence Analysis in Molecular Biology (von Heinje, G., Ed.) Academic (1987); and
 5.) Sequence Analysis Primer (Gribskov, M. and Devereux, J., Eds.) Stockton: NY (1991).

20 Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity calculations may be performed using the MegAlign™ program of the LASERGENE bioinformatics
25 computing suite (DNASTAR Inc., Madison, WI).

 Multiple alignment of the sequences is performed using the "Clustal method of alignment" which encompasses several varieties of the algorithm including the "Clustal V method of alignment" corresponding to the alignment method labeled Clustal V (described by Higgins and Sharp,
30 *CABIOS*. 5:151-153 (1989); Higgins, D.G. et al., *Comput. Appl. Biosci.*, 8:189-191 (1992)) and found in the MegAlign™ program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.). For multiple alignments, the default values correspond to GAP PENALTY=10

and GAP LENGTH PENALTY=10. Default parameters for pairwise alignments and calculation of percent identity of protein sequences using the Clustal method are KTUPLE=1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. For nucleic acids these parameters are

5 KTUPLE=2, GAP PENALTY=5, WINDOW=4 and DIAGONALS SAVED=4. After alignment of the sequences using the Clustal V program, it is possible to obtain a "percent identity" by viewing the "sequence distances" table in the same program.

Additionally the "Clustal W method of alignment" is available and

10 corresponds to the alignment method labeled Clustal W (described by Higgins and Sharp, *CABIOS*. 5:151-153 (1989); Higgins, D.G. et al., *Comput. Appl. Biosci.* 8:189-191 (1992)) and found in the MegAlign™ v6.1 program of the LASERGENE bioinformatics computing suite (DNASTAR Inc.). Default parameters for multiple alignment (GAP PENALTY=10,

15 GAP LENGTH PENALTY=0.2, Delay Divergen Seqs(%)=30, DNA Transition Weight=0.5, Protein Weight Matrix=Gonnet Series, DNA Weight Matrix=IUB). After alignment of the sequences using the Clustal W program, it is possible to obtain a "percent identity" by viewing the "sequence distances" table in the same program.

20 It is well understood by one skilled in the art that many levels of sequence identity are useful in identifying polypeptides, from other species, wherein such polypeptides have the same or similar function or activity. Useful examples of percent identities include, but are not limited to: 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, or any

25 integer percentage from 50% to 100% may be useful in describing the present invention, such as 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,

30 97%, 98% or 99%. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least 50 amino acids, preferably at least 100 amino acids, more preferably at

least 150 amino acids, still more preferably at least 200 amino acids, and most preferably at least 250 amino acids.

The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include, but is not limited to: 1.) the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI); 2.) BLASTP, BLASTN, BLASTX (Altschul et al., J. Mol. Biol., 215:403-410 (1990)); 3.) DNASTAR (DNASTAR, Inc. Madison, WI); 4.) Sequencher (Gene Codes Corporation, Ann Arbor, MI); and 5.) the FASTA program incorporating the Smith-Waterman algorithm (W. R. Pearson, Comput. Methods Genome Res., [Proc. Int. Symp.] (1994), Meeting Date 1992, 111-20. Editor(s): Suhai, Sandor. Plenum: New York, NY). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters that originally load with the software when first initialized.

Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, 2nd ed.; Cold Spring Harbor Laboratory: Cold Spring Harbor, New York, 1989 (hereinafter "Maniatis"); and by Silhavy, T. J., Bennis, M. L. and Enquist, L. W. *Experiments with Gene Fusions*; Cold Spring Harbor Laboratory: Cold Spring Harbor, New York, 1984; and by Ausubel, F. M. et al., In *Current Protocols in Molecular Biology*, published by Greene Publishing and Wiley-Interscience, 1987.

The present invention relates to engineered strains of xylose-utilizing *Zymomonas* or *Zymobacter* that have improved xylose utilization when fermented in xylose containing media. A challenge for improving ethanol production by fermentation of a biocatalyst in media that includes biomass hydrolysate, produced typically by pretreatment and

saccharification of biomass, is obtaining optimal utilization of xylose.

Xylose is one of the predominant pentose sugars in hydrolyzed lignocellulosic materials, the other being arabinose. Applicants have identified a group of xylose isomerases that when expressed in xylose-utilizing *Zymomonas* strains provide for increased efficiency in xylose utilization and higher ethanol yields, when fermentation is in xylose containing media.

Discovery Of Highly Effective Xylose Isomerases

Xylose isomerase (XI), which catalyzes the conversion of xylose to xylulose, is one of four enzymes that when expressed in a bacterial cell provide the ability to utilize xylose as a carbon source. XI along with xylulokinase, transketolase, and transaldolase provide a pathway from xylose that produces fructose -6-P and glyceraldehyde-3-P that feed into ethanol biosynthesis as shown in Figure 1. As the first enzyme of the xylose utilization pathway, xylose isomerase activity is particularly important in providing the ability to ultimately convert xylose to ethanol. Applicants have identified xylose isomerases that have higher activity when expressed in the ethanol producing bacteria, *Zymomonas*, and support enhanced xylose utilization and ethanol production, as compared to the typically used *E. coli* XI.

The xylose isomerase from *E. coli* has been used in engineering *Zymomonas* for xylose utilization. Xylose isomerases are classified in two groups based on their size, amino acid sequence similarity, and divalent cation preference (Park and Batt (2004) Applied and Environmental Microbiology 70:4318-4325). The *E. coli* XI belongs to Group II.

Applicants have discovered that XIs belonging to Group I (as defined herein) provide enhanced properties for xylose utilization and ethanol production in xylose-utilizing *Zymomonas*. The *Actinoplanes missouriensis* XI was found herein to be a Group I XI as described below. When expressed in *Zymomonas* using a codon-optimized coding sequence (SEQ ID NO:308), the XI from *Actinoplanes missouriensis* had a xylose isomerase specific activity that was higher than specific activities of similarly expressed *E. coli* XI (using a codon-optimized coding sequence: SEQ ID NO:310), or another Group II XI, that of *Lactobacillus*

brevis (codon-optimized coding sequence: SEQ ID NO:309). The codon-optimized coding sequences for the *A. missouriensis*, *E. coli*, and *L. brevis* XIs were each expressed in *Zymomonas* engineered with all four enzymes for xylose utilization, including the *E. coli* xylose isomerase expressed
5 from a non-optimized coding sequence. Strains expressing the *A. missouriensis* XI grew better in xylose containing medium, utilized more xylose, and produced more ethanol than strains expressing the XI from either *E. coli* or *L. brevis*.

Zymomonas strains expressing additional XIs that were identified
10 as belonging to Group I, as described below, were also found to grow better, utilize more xylose and produce more ethanol in xylose containing medium than strains expressing the Group II XI from *E. coli* or *L. brevis*. These were strains containing the XIs from *Geodermatophilus obscurus* (SEQ ID NO:64), *Mycobacterium smegmatis* (SEQ ID NO:10), *Salinispora arenicola* (SEQ ID NO:18), and *Xylanimonas cellulositytica* (SEQ ID
15 NO:40).

The growth enhancement, xylose utilization improvement, and
20 ethanol production improvement each may vary in extent in a Group I XI expressing strain as compared to a Group II XI expressing strain. Differences may be based on factors such as specific XI encoding gene expression properties, culturing conditions such as carbon source composition in the medium including amount of xylose and other carbon
25 sources, and strain characteristics such as additional genetic modifications involved in xylose utilization and/or ethanol production.

Group I xylose isomerases

Any XI belonging to Group I may be used in the present strains to improve xylose utilization. The Group I XIs have been distinguished from
30 Group II XIs by their length. Group I XIs were found typically to be about 380 to 390 amino acids in length while Group II XIs are typically about 440 to 460 amino acids in length. Among Group I XIs there is amino acid identity of at least about 50%, while Group II XIs have only 20-30% amino

acid identity with Group I XIs. Thus XIs can be readily classified as belonging to Group I or Group II using these structural criteria.

XIs identified in Park and Batt (*supra*) as belonging to Group I are those from *Streptomyces*, *Actinoplanes*, *Thermus*, and *Arthrobacter* while XIs identified as belonging to Group II are those from *Klebsiella*, *Escherichia*, *Lactobacillus*, *Lactococcus*, *Clostridium*, *Bacillus*, *Staphylococcus*, and *Thermoanaerobacter*.

Bioinformatics analysis was used to more fully characterize Group I as opposed to Group II XIs to identify those XIs that may be used in the present strains. Members of Group I xylose isomerases were identified using molecular phylogenetic analysis of XI amino acid sequences. The molecular phylogenetic analysis was performed on 444 XI sequences collected from a public database using multiple query sequence BLAST analysis (blastall) using 180 XI seed sequences with functional annotations retrieved from the SWISSPROT database: SEQ ID NOs:2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, 142 and 148-306.

The resulting phylogenetic tree shown in Figure 2 places the *A. missouriensis* XI in one phylogenetic grouping, called Group I, and the *E. coli* XI in a separate grouping, called Group II. Group I and Group II are labeled to coincide with the groupings of Park and Bratt (*supra*). The *Lactobacillus brevis* XI, tested herein in Example 3 and shown to have comparable effects as the XI from *E. coli*, is also in Group II. Twenty-one of the seed sequences (SEQ ID NOs:2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142) were found to belong to Group I XIs. The identified XI sequences belonging to Group I form a 50% threshold identity cluster, as described in Example 4. XI sequences belonging to Group II form a separate 50% threshold identity cluster. Each identified Group I XI is encoded by the genome of, and therefore is endogenous to, one of the following microorganisms: *Actinoplanes*, *Arthrobacter*, *Streptomyces*, *Thermus*, *Thermobaculum*, *Herpetosiphon*, *Acidobacteria*, *Roseiflexus*, *Meiothermus*, *Deinococcus*, *Meiothermus*, *Stackebrandtia*, *Kribbella*, *Xylanimonas*, *Nocardiopsis*, *Catenulispora*, *Streptosporangium*, *Geodermatophilus*, *Actinosynnema*,

Saccharomonospora, *Acidothermus*, *Tthermobifida*, *Nocardiodes*,
Janibacter, *Mycobacterium*, *Leifsonia*, *Clavibacter*, *Micromonospora*,
Salinispora, *Cellulomonas*, *Jonesia*, *Nakamurella*, *Actinomyces*,
Mobiluncus, *Brachybacterium*, *Beutengergai*, *Frankia*, and
 5 *Actinobacterium*.

Any XI that belongs to Group I XIs as determined by molecular phylogenetic analysis as described in Example 4 herein may be used in the present strains. The molecular phylogeny of the Group I XIs is shown in more detail in Figure 3. The Group I XIs in Figure 3 are listed in Table 3
 10 as having SEQ ID NOs that are even numbers between 2 and 130 and 131-147. Coding regions for these proteins are listed in Table 3 as having SEQ ID NOs that are odd numbers between 1 and 129. Any other identified XI that can be identified as belonging to Group I using molecular phylogenetic analysis as described in Example 4 herein may be used in
 15 the present strains. Alternatively, XIs that may be used in the present strains may be identified as xylose isomerase proteins with amino acid sequences having at least about 70%-75%, 75%-80%, 80-85%, 85%-90%, 90%-95%, or at least about 96%, 97%, 98%, or 99% sequence identity to any of the XI amino acid sequences of SEQ ID NOs that are
 20 even numbers between, and including, 2 and 130 and 131-147. In one embodiment XIs that may be used in the present strains may be identified as proteins with xylose isomerase activity and with amino acid sequences having at least about 70%-75%, 75%-80%, 80-85%, 85%-90%, 90%-95%, or at least about 96%, 97%, 98%, or 99% sequence identity to any of
 25 the XI amino acid sequences of SEQ ID NOs: 24, 66, 134, 140, 143, 145, and 147. Identities are based on the Clustal W method of alignment using the default parameters of GAP PENALTY=10, GAP LENGTH PENALTY=0.1, and Gonnet 250 series of protein weight matrix.

Group I and Group II XIs were further characterized using
 30 GroupSim analysis as described in Example 4 herein. Through this analysis specific amino acid positions were determined to be specificity determining positions (SDP) for distinguishing the structures of Group I and Group II XI proteins. The locations of these SDP amino acids are given here at corresponding positions in the representative Group I protein

P 12851 from *Actinoplanes missouriensis* (SEQ ID NO:66) and in the representative Group II protein P 19148 from *Thermoanaerobacterium thermosulfurigenes* (SEQ ID NO:267). The positions in the Group II protein are generally about 51 greater than in the Group I protein. The

5 corresponding positions in other Group I and II proteins can readily be identified by one skilled in the art by sequence alignment and context. The SDP identifiers distinguishing Group I and Group II XIs with a score of 0.9 or greater (where a perfect score of 1 would indicate that all proteins within the group have the listed amino acid in the specified position and

10 between groups the amino acid is always different) are the following amino acid (AA) positions in P12851 vs in P 19148:

- 1) AA 226 is leucine; AA277 is histidine
- 2) AA223 is methionine; AA274 is leucine
- 3) AA191 is isoleucine; AA242 is glutamine
- 15 4) AA195 is threonine, serine, or valine; AA246 is aspartic acid
- 5) AA88 is methionine, threonine, or guanine; AA139 is arginine or tryptophan
- 6) AA290 is histidine; AA337 is asparagine or methionine
- 7) AA221 is glutamic acid or aspartic acid; AA 272 is alanine, threonine, or
- 20 glycine
- 8) AA242 is phenylalanine, valine, or leucine; A293 is glycine, cysteine, or tryptophan
- 9) AA243 is histidine; AA294 is serine, asparagine, glycine, or leucine
- 10) AA193 is leucine, phenylalanine, or methionine; AA244 is aspartic acid
- 25 11) AA256 is glutamine; AA 308 is threonine, isoleucine, valine, leucine, methionine, tyrosine, or histidine
- 12) AA213 is glycine; AA264 is lysine, asparagine, serine, glutamic acid, alanine, leucine, arginine, or glutamine
- 13) AA288 is proline, tyrosine, alanine, or serine; AA335 is valine or
- 30 glycine
- 14) AA249 is glutamine; AA301 is aspartic acid, histidine, asparagine, arginine, serine, or alanine

Using these amino acid position identifiers, a Group I XI can be readily identified by one skilled in the art. An XI having the Group I amino

acid position identifiers substantially as listed above will be considered herein to be a Group I XI irrespective of length. Thus if substantially all of the positions indicated for Group II XIs are filled by the amino acids indicated for a Group I XI, the protein is considered to be a Group I XI regardless of length. For example, an XI having the Group I AA226 leucine identifier at the 277 position instead of histidine is a Group I XI if this pattern holds for the other amino acid identifiers. Substantially here means that there need not be a complete exact match at all positions, as indicated by the scores of 0.9 as opposed to a score of 1. A Group I XI has at least 90% of the amino acids at the SDPs matching the above list.

An additional bioinformatics analysis of Group I XIs was performed using the hmmsearch algorithm of the HMMER software package (Janelia Farm Research Campus, Ashburn, VA). The Z parameter of the hmmsearch algorithm was set to 1 billion. The output of the HMMER analysis using a set of protein sequences is a Profile Hidden Markov Model (Profile HMM). The theory behind Profile HMMs as described in Durbin et al. (Biological sequence analysis: probabilistic models of proteins and nucleic acids, Cambridge University Press, 1998) and Krogh et al. (1994 *J. Mol. Biol.* 235:1 501-1 531), both incorporated herein by reference, is characterization of a set of proteins based on the probability of each amino acid occurring at each position in the alignment of the proteins of the set.

The 21 seed sequences having known xylose isomerase activity and found to belong to Group I XIs in the molecular phylogenetic analysis described above were used as the set of proteins to prepare a profile HMM. These proteins have SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142. All of the XIs that were identified by molecular phylogenetic analysis as belonging to Group I, which are listed in Table 3, match the profile HMM prepared from Group I seed sequences with E-value scores of less than or equal to 2.2×10^{-181} , with the Z parameter set to 1 billion. All of the XIs identified by molecular phylogenetic analysis as belonging to Group II match the same profile HMM with E-value scores of greater than or equal to 1.5×10^{-07} . Table 2 of the appendix lists the E-value scores for each XI SEQ ID NO. Thus

the prepared profile HMM gives a structural characterization for functional Group I XIs and corroborates the molecular phylogenetic analysis.

Therefore any XI protein that matches the profile HMM prepared using the 21 Group I XI seed sequences described above with an E-value of $1E-15$ or less, where $1E-15$ is between the highest score for a Group I XI and the lowest score for a Group II XI, may be used in the present strains. Lower E-value scores indicate better matches.

Additionally, the Group I XI sequences described herein or those recited in the art may be used to identify other homologs in nature. For example each of the XI encoding nucleic acid fragments described herein may be used to isolate genes encoding homologous proteins. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to: 1) methods of nucleic acid hybridization; 2) methods of DNA and RNA amplification, as exemplified by various uses of nucleic acid amplification technologies [e.g., polymerase chain reaction (PCR), Mullis et al., U.S. Patent 4,683,202; ligase chain reaction (LCR), Tabor, S. et al., *Proc. Acad. Sci. USA* 82:1074 (1985); or strand displacement amplification (SDA), Walker, et al., *Proc. Natl. Acad. Sci. U.S.A.*, 89:392 (1992)]; and 3) methods of library construction and screening by complementation.

As is known in the art, there may be variations in DNA sequences encoding an amino acid sequence due to the degeneracy of the genetic code. Codons may be optimized for expression of an amino acid sequence in a target host cell to provide for optimal encoded protein expression

Group I XI expression

In the present strains, any of the Group I XIs described above may be expressed in a strain of *Zymomonas* or related ethanolagen, such as *Z. mobilis* or *Zymobacter*, along with genes encoding xylulokinase, transketolase, and transaldolase as described above, which then is capable of utilizing xylose as described below. *Zymobacter palmae* is an ethanol-producing bacterium that has been engineered for xylose utilization by expressing genes for xylose utilization as described below for

Zymomonas, using *Z. mobilis* glyceraldehyde-3-phosphate dehydrogenase and enolase promoters (Yanase et al. Applied and Environmental Microbiology (2007) 73:2592-2599).

Coding regions that may be used to express Group I XIs include
5 those listed in Table 1 as SEQ IDs with odd numbers from 1 through 129,
other sequences encoding XI proteins listed in Table 1 as SEQ IDs with
even numbers 2 through 130 and 131-147, as well as sequences identified
in the art as encoding Group I XIs using bioinformatics or experimental
methods described herein and those well known in the art.

10 For expression, a Group I XI coding region is constructed in a
chimeric gene with operably linked promoter and typically a termination
sequence. Alternatively the Group I XI coding region is constructed as part
of an operon that is operably linked to a promoter and a termination
sequence, and includes one or more additional coding regions. Promoters
15 that may be used are promoters that are expressed in *Zymomonas* or
Zymobacter cells such as the promoters of *Z. mobilis* glyceraldehyde-3-
phosphate dehydrogenase (GAP promoter or P_{gap}), including mutant more
highly active GAP promoters disclosed in US 20090246876, which is
incorporated herein by reference, that may be called superGAP promoters
20 or P_{gaps}, and *Z. mobilis* enolase (ENO promoter) genes. Termination
signals are also those that are expressed in the target cell.

A chimeric gene or operon for XI expression is typically constructed
in or transferred to a vector for further manipulations. Vectors are well
known in the art. Certain vectors are capable of replicating in a broad
25 range of host bacteria and can be transferred by conjugation. The
complete and annotated sequence of pRK404 and three related vectors:
pRK437, pRK442, and pRK442(H) are available. These derivatives have
proven to be valuable tools for genetic manipulation in gram-negative
bacteria (Scott et al., *Plasmid* 50(1):74-79 (2003)).

30 Particularly useful for expression in *Zymomonas* are vectors that
can replicate in both *E. coli* and *Zymomonas*, such as pZB188 which is
described in U.S. Pat. No. 5,514,583. Vectors may include plasmids for
autonomous replication in a cell, and plasmids for carrying constructs to
be integrated into bacterial genomes. Plasmids for DNA integration may

include transposons, regions of nucleic acid sequence homologous to the target bacterial genome, or other sequences supporting integration. An additional type of vector may be a transposome produced using, for example, a system that is commercially available from EPICENTRE®. It is well known how to choose an appropriate vector for the desired target host and the desired function.

Bacterial cells may be engineered by introducing a vector having a chimeric gene comprising a xylose isomerase coding region by well known methods, such as using freeze-thaw transformation, calcium-mediated transformation, electroporation, or conjugation. Any bacterial cell to be engineered for xylose utilization by expressing a xylose isomerase enzyme is a target host cell for transformation to engineer a strain as described herein. Particularly suitable host cells are *Zymomonas* and *Zymobacter* cells. The introduced chimeric gene may be maintained in the cell on a stably replicating plasmid, or integrated into the genome following introduction.

For engineering a strain with an integrated xylose isomerase chimeric gene or operon in the bacterial cell genome, methods may be used that are well known in the art such as homologous recombination, transposon insertion, or transposome insertion. In homologous recombination, DNA sequences flanking a target integration site are placed bounding a spectinomycin-resistance gene, or other selectable marker, and xylose isomerase chimeric gene leading to insertion of the selectable marker and the xylose isomerase chimeric gene into the target genomic site. In addition, the selectable marker may be bounded by site-specific recombination sites, so that after expression of the corresponding site-specific recombinase, the resistance gene is excised from the genome.

Engineering Of Full Xylose Utilization Pathway

In addition to transforming with a chimeric gene or operon comprising a Group I XI coding region, the present strains are also engineered for expression of three other enzymes needed for xylose utilization: xylulokinase, which phosphorylates xylulose to form xylulose 5-phosphate, and transaldolase and transketolase, two enzymes of the

pentose phosphate pathway which convert xylulose 5-phosphate to intermediates that couple pentose metabolism to the glycolytic Entner-Doudoroff pathway permitting the metabolism of xylose to ethanol (see Figure 1). Xylose utilizing *Zymomonas* strains are described in U.S. Pat. No. 5514583, U.S. Pat. No. 5712133, U.S. Pat. No. 6566107, WO 95/28476, Feldmann et al. ((1992) Appl Microbiol Biotechnol 38: 354-361), Zhang et al. ((1995) Science 267:240-243. These strains were transformed with coding sequences from *E. coli* genes, including a Group II xylose isomerase.

The Group I XI may be the sole XI expressed for xylose utilization-, or it may be expressed in addition to an expressed Group II XI such as that from *E. coli*. Thus a Group I XI may be introduced in a *Zymomonas* or *Zymobacter* strain that has a full xylose utilization pathway that includes a Group II xylose isomerase encoding gene and is capable of utilizing xylose. Alternatively, a Group I XI may be introduced in a *Zymomonas* or *Zymobacter* strain that expresses xylulokinase, transaldolase, and transketolase, and only lacks xylose isomerase activity for utilizing xylose. With introduction of a Group I XI the strain is capable of utilizing xylose.

The additional three enzymes may be expressed from individual chimeric genes or from operons including more than one coding region as well known to one skilled in the art. DNA sequences encoding these enzymes may be obtained from any of numerous microorganisms that are able to metabolize xylose, such as enteric bacteria, and some yeasts and fungi. Sources for the coding regions include *Xanthomonas*, *Klebsiella*, *Escherichia*, *Rhodobacter*, *Flavobacterium*, *Acetobacter*, *Gluconobacter*, *Rhizobium*, *Agrobacterium*, *Salmonella*, *Pseudomonads*, and *Zymomonas*. Particularly useful are the coding regions of *E. coli*.

Endogenous genes may provide part of a xylose fermentation pathway, or may be altered by any known genetic manipulation technique to provide a protein with enzyme activity useful for xylose metabolism. For example, an endogenous transketolase may complement other introduced enzyme activities in creating a xylose utilization pathway.

Examples of xylose-utilizing strains that are known and may be used include CP4(pZB5) (US 5,514,583), ATCC31821/pZB5 (US

6,566,1 07), 8b (US 200301 62271 ; Mohagheghi et al., (2004) Biotechnol. Lett. 25; 321 -325), and ZW658 (ATTCC # PTA-7858; US 7,741 ,1 19).

Zymomonas or *Zymobacter* strains that are additionally engineered to utilize other sugars that, like xylose, are not natural substrates, may
5 also be used in the present process. An example is a strain of *Z. mobilis* engineered for arabinose utilization as described in US 5^843^760, which is herein incorporated by reference. Strains may be modified in other additional ways to improve xylose utilization and ethanol production.

Fermentation Of Improved Xylose-Utilizing Strain

10 An engineered xylose-utilizing strain having a Group I xylose isomerase chimeric gene and genes or operons for expression of xylulokinase, transaldolase and transketolase may be used in fermentation to produce a product that is a natural product of the strain, or a product that the strain is engineered to produce. For example, *Zymomonas mobilis*
15 and *Zymobacter palmae* are natural ethanolagens. As an example, production of ethanol by a *Z. mobilis* strain of the invention is described.

For production of ethanol, recombinant xylose-utilizing *Z. mobilis* having a Group I xylose isomerase chimeric gene is brought in contact with medium that contains mixed sugars including xylose. Typically the
20 medium contains mixed sugars including arabinose, xylose, and glucose. The medium may contain biomass hydrolysate that includes these sugars that are derived from treated cellulosic or lignocellulosic biomass.

When the mixed sugars concentration is high such that growth is inhibited, the medium includes sorbitol, mannitol, or a mixture thereof as
25 disclosed in US 7,629,1 56. Galactitol or ribitol may replace or be combined with sorbitol or mannitol. The *Z. mobilis* grows in the medium where fermentation occurs and ethanol is produced. The fermentation is run without supplemented air, oxygen, or other gases (which may include conditions such as anaerobic, microaerobic, or microaerophilic
30 fermentation), for at least about 24 hours, and may be run for 30 or more hours. The timing to reach maximal ethanol production is variable, depending on the fermentation conditions. Typically, if inhibitors are present in the medium, a longer fermentation period is required. The

fermentations may be run at temperatures that are between about 30 ° C and about 37 ° C, at a pH of about 4.5 to about 7.5.

The present *Z. mobilis* may be grown in medium containing mixed sugars including xylose in laboratory scale fermenters, and in scaled up
5 fermentation where commercial quantities of ethanol are produced. Where commercial production of ethanol is desired, a variety of culture methodologies may be applied. For example, large-scale production from the present *Z. mobilis* strains may be produced by both batch and continuous culture methodologies. A classical batch culturing method is a
10 closed system where the composition of the medium is set at the beginning of the culture and not subjected to artificial alterations during the culturing process. Thus, at the beginning of the culturing process the medium is inoculated with the desired organism and growth or metabolic activity is permitted to occur adding nothing to the system. Typically,
15 however, a "batch" culture is batch with respect to the addition of carbon source and attempts are often made at controlling factors such as pH and oxygen concentration. In batch systems the metabolite and biomass compositions of the system change constantly up to the time the culture is terminated. Within batch cultures cells moderate through a static lag
20 phase to a high growth log phase and finally to a stationary phase where growth rate is diminished or halted. If untreated, cells in the stationary phase will eventually die. Cells in log phase are often responsible for the bulk of production of end product or intermediate in some systems. Stationary or post-exponential phase production can be obtained in other
25 systems.

A variation on the standard batch system is the Fed-Batch system. Fed-Batch culture processes are also suitable for growth of the present *Z. mobilis* strains and comprise a typical batch system with the exception that the substrate is added in increments as the culture progresses. Fed-Batch
30 systems are useful when catabolite repression is apt to inhibit the metabolism of the cells and where it is desirable to have limited amounts of substrate in the medium. Measurement of the actual substrate concentration in Fed-Batch systems is difficult and is therefore estimated on the basis of the changes of measurable factors such as pH and the

partial pressure of waste gases such as CO₂. Batch and Fed-Batch culturing methods are common and well known in the art and examples may be found in *Biotechnology: A Textbook of Industrial Microbiology*, Crueger, Crueger, and Brock, Second Edition (1989) Sinauer Associates, Inc., Sunderland, MA, or Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36, 227, (1992), herein incorporated by reference.

Commercial production of ethanol may also be accomplished with a continuous culture. Continuous cultures are open systems where a defined culture medium is added continuously to a bioreactor and an equal amount of conditioned medium is removed simultaneously for processing. Continuous cultures generally maintain the cells at a constant high liquid phase density where cells are primarily in log phase growth. Alternatively, continuous culture may be practiced with immobilized cells where carbon and nutrients are continuously added, and valuable products, by-products or waste products are continuously removed from the cell mass. Cell immobilization may be performed using a wide range of solid supports composed of natural and/or synthetic materials as is known to one skilled in the art.

Continuous or semi-continuous culture allows for the modulation of one factor or any number of factors that affect cell growth or end product concentration. For example, one method will maintain a limiting nutrient such as the carbon source or nitrogen level at a fixed rate and allow all other parameters to moderate. In other systems a number of factors affecting growth can be altered continuously while the cell concentration, measured by medium turbidity, is kept constant. Continuous systems strive to maintain steady state growth conditions and thus the cell loss due to medium being drawn off must be balanced against the cell growth rate in the culture. Methods of modulating nutrients and growth factors for continuous culture processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology and a variety of methods are detailed by Brock, *supra*.

Particularly suitable for ethanol production is a fermentation regime as follows. The desired *Z. mobilis* strain of the present invention is grown in shake flasks in semi-complex medium at about 30 °C to about 37 °C

with shaking at about 150 rpm in orbital shakers and then transferred to a 10 L seed fermentor containing similar medium. The seed culture is grown in the seed fermentor anaerobically until OD_{600} is between 3 and 6, when it is transferred to the production fermentor where the fermentation parameters are optimized for ethanol production. Typical inoculum volumes transferred from the seed tank to the production tank range from about 2% to about 20% v/v. Typical fermentation medium contains minimal medium components such as potassium phosphate (1.0 - 10.0 g/l), ammonium sulfate (0- 2.0 g/l), magnesium sulfate (0 - 5.0 g/l), a complex nitrogen source such as yeast extract or soy based products (0 - 10 g/l). A final concentration of about 5 mM sorbitol or mannitol is present in the medium. Mixed sugars including xylose and at least one additional sugar such as glucose (or sucrose), providing a carbon source, are continually added to the fermentation vessel on depletion of the initial batched carbon source (50-200 g/l) to maximize ethanol rate and titer. Carbon source feed rates are adjusted dynamically to ensure that the culture is not accumulating glucose in excess, which could lead to build up of toxic byproducts such as acetic acid. In order to maximize yield of ethanol produced from substrate utilized, biomass growth is restricted by the amount of phosphate that is either batched initially or that is fed during the course of the fermentation. The fermentation is controlled at pH 5.0 - 6.0 using caustic solution (such as ammonium hydroxide, potassium hydroxide, or sodium hydroxide) and either sulfuric or phosphoric acid. The temperature of the fermentor is controlled at 30 °C - 35 °C. In order to minimize foaming, antifoam agents (any class- silicone based, organic based etc) are added to the vessel as needed. An antibiotic, for which there is an antibiotic resistant marker in the strain, such as kanamycin, may be used optionally to minimize contamination.

Any set of conditions described above, and additionally variations in these conditions that are well known in the art, are suitable conditions for production of ethanol by a xylose-utilizing recombinant *Zymomonas* strain.

EXAMPLES

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can

5 ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various uses and conditions.

GENERAL METHODS

10 The meaning of abbreviations is as follows: "kb" means kilobase(s), "bp" means base pairs, "nt" means nucleotide(s), "hr" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "L" means liter(s), "ml" or "ml_" means milliliter(s), " μ l" means microliter(s), "Vg" means microgram(s), "ng" means nanogram(s), "mg" means

15 milligram(s), "mM" means millimolar, " μ M" means micromolar, "nm" means nanometer(s), " μ mol" means micromole(s), "pmol" means picomole(s), "Cm" means chloramphenicol, "Cm^r" means chloramphenicol resistant, "Cm^s" means chloramphenicol sensitive, "Sp^r" means spectinomycin resistance, "Sp^s" means spectinomycin sensitive, "XI" is xylose isomerase,

20 "XK" is xylulokinase, "TAL" is transaldolase, "TKT" is transketolase, "RM" means rich medium containing 10 g/L yeast extract plus 2 g/L KH_2PO_4 , "MM" means mating medium containing 10 g/L yeast extract, 5 g/L tryptone, 2.5 g/L $(\text{NH}_4)_2\text{SO}_4$ and 0.2 g/L KH_2PO_4 .

Standard recombinant DNA and molecular cloning techniques used

25 here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory: Cold Spring Harbor, NY (1989) (hereinafter "Maniatis"); and by Silhavy, T. J., Bennis, M. L. and Enquist, L. W., *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory:

30 Cold Spring Harbor, NY (1984); and by Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, published by Greene Publishing Assoc. and Wiley-Interscience, Hoboken, NJ (1987).

Example 1Construction Of Chimeric Xylose Isomerase Genes And Assembly Of
Double Crossover Suicide Vectors

Constructs were made for integration and expression in

5 *Zymomonas mobilis* of the coding regions for xylose isomerases from *Actinoplanes missouriensis* (AMxylA), *Lactobacillus brevis* (LBxylA), and *Escherichia coli* (ECxylA). The coding sequences were optimized for expression in *Z. mobilis* according to the codon bias of *Z. mobilis* ZM4 and synthesized by GenScript Corporation (Piscataway, NJ). Each was cloned

10 into pUC57 at the EcoRV site and provided as the plasmids pUC57-AMxylA (with codon optimized AMxylA coding region SEQ ID NO:308), pUC57-LBxylA (with codon optimized LBxylA coding region SEQ ID NO:309), and pUC57-ECxylA (with codon optimized ECxylA coding region SEQ ID NO:310). The optimized xylA coding sequences encode the

15 native xylose isomerases (XIs).

The xylA coding sequences were constructed into chimeric genes with the structure of P_{gap}-xylA-araD3'UTR by linking with a 305 bp promoter from the *Z. mobilis* glyceraldehyde-3-phosphate dehydrogenase gene (P_{gap}; SEQ ID NO:311) and a 166 bp terminator from the *E. coli* L-ribulose 5 phosphate 4-epimerase gene (araD3'UTR; SEQ ID NO:312).

20 For this purpose, P_{gap} and araD3'UTR overlapping fragments were synthesized by PCR. One PCR reaction consisted of 50 μ L AccuPrime Pfx SuperMix (Invitrogen, Carlsbad, CA), 1 μ L of 40 ng/ μ L pARA354 as template (SEQ ID NO:313), and 1 μ L of 10 μ M forward and reverse

25 primers. Plasmid pARA354 (SEQ ID NO:313) is described in commonly owned and co-pending US Patent Application 12/796025, which is herein incorporated by reference, and is a pBS SK(+) vector that includes a P_{gap}-araBAD operon, which is the P_{gap} promoter adjacent to coding regions for araB, araA, and araD (encoding the proteins L-ribulose kinase, L-

30 arabinose isomerase, and L-ribulose-5-phosphate-4-epimerase, respectively) from *E. coli*. The operon includes the 3' untranslated region (UTR) that is 3' to the araD coding region. pARA354 is described further below. Reactions were carried out on an Eppendorf Mastercycler

(Hamburg, Germany), following a hot start PCR program with 30 cycles of denaturing at 95°C for 30 sec / annealing at 58°C (56°C for araD3'UTR) for 30 sec / extension at 68°C for 2 min.

Primers ara98 and ara120 (SEQ ID NOS:314 and 315) produced a P_{gap}-AM overlapping fragment. Primers ara98 and ara121 (SEQ ID NOS: 314 and 316) produced a P_{gap}-EC overlapping fragment. Primers ara98 and ara122 (SEQ ID NOS: 314 and 317) produced a P_{gap}-LB overlapping fragment. In addition to the 305 bp P_{gap} sequence, all three P_{gap} overlapping PCR fragments included a 17 bp 5' sequence to add StuI and SpeI sites and a 22 bp 3' sequence to match the first 22 nucleotides of their counterpart xylA coding sequence, that were provided in the primers.

Primer ara96 and ara97 (SEQ ID NOS: 318 and 319) produced a 210-bp araD3'UTR overlapping fragment. It included a 24 bp 5' sequence with an XbaI site at the end, and a 20 bp 3' sequence providing EcoRI, HindIII, and FseI sites, in addition to the 166-bp araD3'UTR. Similar PCR was also conducted to synthesize xylA overlapping fragments, but annealing temperature was lowered to 55°C. In these reactions, a 1,229 bp AMxylA overlapping fragment was amplified from pUC57-AMxylA using primers ara114 and ara115 (SEQ ID NOS:320 and 321). A 1,367-bp ECxylA overlapping fragment was amplified from pUC57-ECxylA using primers ara116 and ara117 (SEQ ID NOS:322 and 323). A 1,394-bp LBxylA overlapping fragment was amplified from pUC57-LBxylA using primers ara118 and ara119 (SEQ ID NOS:324 and 325). All xylA overlapping fragments have an 18-bp 5' sequence matching the last 18 nucleotides of P_{gap}, a 24-bp 3' sequence providing an XbaI site and matching the first 18 nucleotides of araD3'UTR, as well as a xylA coding sequence between them. The P_{gap}, araD3'UTR, and xylA overlapping fragments were confirmed by running 5 µL of each PCR sample on an agarose gel, and then purified by using QIAquick PCR Purification Kit (Qiagen, Valencia, CA).

Overlapping fragments were linked together by overlapping PCR. The first overlapping PCR was assembled to include 50 µL AccuPrime Pfx SuperMix, 1 µL of 20 ng/µL P_{gap} overlapping fragment, 2 µL of 10 ng/µL

counterpart xylA overlapping fragment, and 1 μ L of 10 μ M forward and reverse primers as follows. Reaction was conducted by following a hot start PCR program with 30 cycles of denaturing at 95°C for 30 sec / annealing at 55°C for 30 sec / extension at 68°C for 2 min. As a result, a

5 P_{gap}-AMxylA fragment was synthesized from P_{gap}-AM and AMxylA fragments using primers ara98 and ara1 15 (SEQ ID NOs:314 and 321); a P_{gap}-ECxylA fragment was synthesized from P_{gap}-EC and ECxylA fragments using primers ara98 and ara1 17 (SEQ ID NOs:314 and 323); and a P_{gap}-LBxylA fragment was synthesized from P_{gap}-LB and LBxylA

10 fragments using primers ara98 and ara1 19 (SEQ ID NOs:314 and 325). These P_{gap}-xylA fragments were confirmed by running 5 μ L of each PCR sample on an agarose gel, and then purified by using QIAquick PCR Purification Kit.

A second overlapping PCR was assembled similarly to above. It

15 included 50 μ L AccuPrime Pfx SuperMix, 1 μ L of 20 ng/ μ L araD3'UTR overlapping fragment, 2 μ L of 10 ng/ μ L P_{gap}-xylA fragment, 1 μ L of 10 μ M primer ara97 (SEQ ID NO:319), and 1 μ L of 10 μ M primer ara98 (SEQ ID NO:318). Reaction was carried out for 30 cycles of denaturing at 95°C for 30 sec / annealing at 56°C for 30 sec / extension at 68°C for 2.5 min. Five

20 microliters of the resulting PCR product were inspected on an agarose gel. The reactions containing P_{gap}-AMxylA and P_{gap}-LBxylA produced a 1,714-bp chimeric AMxylA operon fragment (P_{gap}-AMxylA-araD3'UTR) and a 1,879-bp chimeric LBxylA operon fragment (P_{gap}-LBxylA-araD3'UTR), respectively. In both chimeric gene fragments, the first 17 nucleotides

25 provide Stul and SphI sites while the last 35 nucleotides contain FseI, HindIII, and EcoI sites. The PCR reaction containing P_{gap}-ECxylA failed to generate the chimeric ECxylA fragment (P_{gap}-ECxylA-araD3'UTR).

The chimeric genes containing AMxylA and LBxylA were each ligated into a double crossover (DCO) vector named pARA354 (SEQ ID

30 NO:313) that was described in commonly owned and co-pending US Patent Application 12/796025. pARA354 is pBS SK(+) derived plasmid (a Bluescript plasmid; Stratagene), used as a suicide vector since pBS vectors cannot replicate in *Zymomonas*, containing a P_{gap}-araBAD operon

as described above and DCO homologous recombination fragments to direct integration of a bounded fragment into the *IdhA* locus of the *Zymomonas* genome. The two *IdhA* DNA fragments of pARA354 for DCO, LDH-L and LDH-R, were synthesized by PCR using *Z. mobilis* DNA as
 5 template. The reaction used AccuPrime Mix and followed a standard PCR procedure. The LDH-L DNA fragment was synthesized using forward primer ara20 (SEQ ID NO:326) and reverse primer ara21 (SEQ ID NO:327). The resulting product was an 895-bp DNA fragment including sequence 5' to the *IdhA* coding region and nucleotides 1-493 of the *IdhA*
 10 coding region, with a 5' *SacI* site and a 3' *SpeI* site (SEQ ID NO:328). The LDH-R DNA fragment was synthesized using forward primer ara22 (SEQ ID NO:329) and reverse primer ara23 (SEQ ID NO:330). The resulting product was a 1169 bp fragment including nucleotides 494-996 of the *IdhA* coding region and sequence 3' to the *IdhA* coding region, with a 5' *EcoRI*
 15 site and a 3' *NotI* site (SEQ ID NO:331). Since LDH-L and LDH-R contained the first 493 base pairs and the remaining 503 base pairs of the *IdhA* coding sequence, respectively, pARA354 was designed to direct insertion of a DNA fragment into the *IdhA* coding sequence of *Z. mobilis* between nucleotides #493 and #494 by crossover recombination

20 pARA354 contains an *f1(+)* origin and an ampicillin resistance gene for plasmid propagation in *E. coli*. In addition, between the LDH-L and LDH-R homologous recombination fragments in pARA354 is the *aadA* marker (for spectinomycin resistance) bounded by wild type *LoxP* sites (*LoxPw*-*aadA*-*LoxPw* fragment; SEQ ID NO:307) and a *P_{gap}*-*araBAD*
 25 operon.

The PCR fragments containing *AMxylA* and *LBxylA* chimeric genes were each digested with *SpeI* and *EcoRI*, subjected to agarose gel electrophoresis, and purified by using QIAquick Gel Purification Kit (Qiagen). At the same time, pARA354 was also digested with *SpeI* and
 30 *EcoRI* to drop out the *P_{gap}*-*araBAD* operon. The *EcoRI*-pARA354-*SpeI* plasmid backbone (6,023 bp) was isolated by agarose gel electrophoresis, and purified by using a QIAquick Gel Purification Kit. The chimeric *AMxylA* and *LBxylA* genes were each constructed into the pARA354

backbone in 15 μ I standard ligation reactions that included 5 μ I of the digested AMxylA or LBxylA chimeric gene fragment, 2 μ I of the digested pARA354 backbone, 3 μ I 5x ligase buffer, and 1 μ I T4 DNA ligase (Invitrogen), resulting in a 7,714-bp DCO plasmid pARA355 and a 7,879-bp DCO plasmid pARA356, respectively. Both plasmids were propagated in DH5a *E. coli* cells and prepared by using QIAprep Spin Miniprep Kit (Qiagen).

To construct a chimeric ECxylA gene in the DCO vector, the P_{gap} -ECxylA overlapping fragment produced in the first overlapping PCR was digested with SpeI and XbaI, subjected to agarose gel electrophoresis, and purified by using QIAquick Gel Purification Kit. At the same time, pARA355 was digested with SpeI and XbaI. The XbaI-pARA355-SpeI plasmid backbone (6,220 bp) was isolated by agarose gel electrophoresis, and purified by using QIAquick Gel Purification Kit. The P_{gap} -ECxylA fragment was assembled into the pARA355 backbone in a 15 μ I standard ligation reaction as described above, including 5 μ I of the digested P_{gap} -ECxylA fragment and 2 μ I of the digested pARA355 backbone fragment. The resultant 7,852 bp DCO plasmid pARA357 was propagated in DH5a *E. coli* cells and prepared by using QIAprep Spin Miniprep Kit.

Example 2

Integration of chimeric AMxylA, LBxylA, and ECxylA genes into *Zymomonas mobilis* strain ZW641

Effects of expressing the *A. missouriensis*, *L. brevis* or *E. coli* XI in a xylose-utilizing *Zymomonas mobilis* were assayed using strain ZW641.

Preparation of the ZW641 strain is described in Example 1 of US 7,741,119, which is incorporated herein by reference. Strain X13L3 described therein was later renamed ZW641. ZW641 was prepared by sequentially integrating the two operons P_{gap} xylAB and P_{gap} taltkt, along with the chloramphenicol resistance selectable marker, into the genome of *Zymomonas mobilis* ZW1 (ATCC #31821). Transformants were further adapted for xylose utilization by growth in xylose-containing medium.

In the ZW641 integrated P_{gap} xylAB and P_{gap} taltkt operons the xylA, xylB, tal, and tkt coding regions are from *E. coli* genes. Though ZW641 has

all four genes necessary for xylose metabolism, xylose utilization is not optimal. Thus a background level of xylose utilization in ZW641 could potentially be improved by expressing an additional XI gene.

Competent cells of the ZW641 -1A strain (a ZW641 isolate) were prepared by growing seed cells overnight in MRM3G5 (1% yeast extract, 15 mM KH_2PO_4 , 4 mM MgSO_4 , and 50 g/L glucose) at 30°C with 150 rpm shaking, to an OD_{600} value near 5. The OD_{600} value was measured using a Shimadzu UV-1200 Spectrophotometer (Kyoto, Japan). Cells were harvested and resuspended in fresh medium to an OD_{600} value of 0.05.

The cells were grown under the same conditions to early to middle log phase (OD_{600} near 0.5). Cells were harvested and washed twice with ice-cold water and then once with ice-cold 10% glycerol. The resulting competent cells were collected and resuspended in ice-cold 10% glycerol to an OD_{600} value near 100. Since transformation of *Z. mobilis* requires non-methylated DNA, DCO plasmids pARA355, pARA356, and pARA357 were each transformed into *E. coli* SCS1 10 competent cells (Stratagene, La Jolla, CA). For each transformation, one colony of transformed cells was grown in 10 mL LB-Amp100 (LB broth containing 100 mg/L ampicillin) overnight at 37°C. DNA was prepared from the 10 mL culture, using QIAprep Spin DNA Miniprep Kit (Qiagen).

Approximately 1 μg non-methylated plasmid DNA was mixed with 50 μL ZW641 -1A competent cells in a 1 MM Electroporation Cuvette (VWR, West Chester, PA). The plasmid DNA was electroporated into the cells at 2.0 KV using a BT720 Transporater Plus (BTX-Genetronics, San Diego, CA). Transformed cells were recovered in 1 mL MMG5 medium (10 g/L glucose, 10 g/L yeast extract, 5 g/L tryptone, 2.5 g/L $(\text{NH}_4)_2\text{SO}_4$, 2 g/L K_2HPO_4 , and 1 mM MgSO_4) for 4 hours at 30°C and grown on MMG5-Spec250 plates (MMG5 with 250 mg/L spectinomycin and 15 g/L agar) for 3 days at 30°C, inside an anaerobic jar with an AnaeroPack (Mitsubishi Gas Chemical, New York, NY). About 20 spectinomycin-resistant colonies were obtained for each transformation. These colonies were streaked onto a fresh MMG5-Spec250 plate and their growth under the same conditions as described above indicated that the chimeric xylA gene / Spec-R

construct had been integrated into the genome of ZW641 . Integration was analyzed by PCR. One reaction included 25 μ L PCR SuperMix (Invitrogen), 0.5 μ L 10 μ M forward primer and reverse primer (as specified below), and a small amount of *Z. mobilis* cells from the colonies. Reaction was carried out on an Eppendorf Mastercycler, following a hard start PCR program with 35 cycles of denaturing at 94°C for 45 sec / annealing at 55°C for 45 sec / extension at 72°C for 1.5 min. Reactions were examined by running 5 μ L on an agarose gel. When ara46 and ara43 primers (SEQ ID NOs:332 and 333) were used in the first inspection, a 1,521-bp PCR product was amplified from most colonies. This product spans from the aadA coding region of the Spec-R marker in the plasmids to a *Z. mobilis* genomic sequence downstream of the LDH-R fragment, demonstrating the integration events mediated by the LDH-R fragment. When forward-reverse primer pairs were ara45-ara120 (SEQ ID NOs:334 and 315), ara45-ara122 (SEQ ID NOs:334 and 317), and ara45-ara121 (SEQ ID NOs:334 and 316) in the second inspection, a 1,289-bp PCR product was amplified from the colonies of ZW641-ara355, ZW641-ara356, and ZW641-ara357, respectively. These products span from a *Z. mobilis* genomic sequence upstream of the LDH-L fragment to AMxylA in pARA355, LBxylA in pARA356, or ECxylA in pARA357. These demonstrated the integration events mediated by the LDH-L fragment. Therefore, PCR evidence confirmed that the chimeric xylA operon / Spec-R construct had been integrated into the genome of ZW641-1A. The ZW641-1A cells transformed with pARA355, pARA356, and pARA357 were named as ZW641-ara355, ZW641-ara356, and ZW641-ara357, respectively.

Example 3

Characterization of AMxylA, LBxylA, and ECxylA expression in *Zymomonas mobilis* strain ZW641

ZW641 has a copy of the native *E. coli* xylA coding region, which is expressed at a low level. Strains ZW641-ara355, ZW641-ara356, and ZW641-ara357 each contain an additional copy of a codon-optimized xylA coding region: AMxylA, LBxylA, and ECxylA, respectively. Enhanced

xylose utilization, ethanol production, and growth in xylose were assayed for the strains.

To examine the growth of these new strains in media containing xylose and compare them with the parent strain ZW641 -1A, two strains (#1 and #2) of each of ZW641 -ara355, ZW641 -ara356, and ZW641 -ara357 from the MMG5-Spec250 plates described in the previous example were re-streaked onto a MMX5 plate (the same medium except glucose is replaced by xylose). ZW641 was also streaked onto the plate as a control. Cells on the plate were grown for 6 days at 30°C inside an anaerobic jar with an AnaeroPack. Growth was observed for all three sets of the new strains, but not for the ZW641 control. The #1 and #2 strains of ZW641 -ara355, which contain an additional copy of AMxylA, showed significantly more growth on the xylose medium than ZW641 -ara356 and ZW641 -ara357 strains.

To quantitatively measure the growth in xylose, these 7 strains were subjected to a 96-hour growth assay. In the assay, cells from each strain were grown overnight in 3 ml MRM3G5 in a 30°C 150 rpm shaker. Cells were harvested, washed with MRM3X10 (same as MRM3G5 but 50 g/L glucose was replaced by 100 g/L xylose), and resuspended in MRM3X10 to have a starting OD₆₀₀ value near 0.1. Twenty-five milliliters of the suspension was placed in a 50 ml screw capped VWR centrifuge tube and grown at 30°C with 150 rpm shaking for a 96-hour time course. During the time course, OD₆₀₀ value was measured at 0-, 4-, 24-, 48-, 72-, and 96-hours. The results plotted as growth curves in Figure 4 show that the second copy of xylA indeed enhanced the growth of the engineered strains in xylose containing medium. When comparing between those strains containing the second copy of xylA, ZW641 -ara355 grew significantly faster than ZW641 -ara357. It had a cell density almost twice as high as ZW641 -ara357 after 96 hours of growth. This result indicates that the xylose isomerase encoded by AMxylA may function much better than the xylose isomerase encoded by ECxylA. ZW641 -ara356 grew similarly to or slightly slower than ZW641 -ara357, indicating that the xylose isomerase encoded by LBxylA may not function better than the xylose isomerase encoded by ECxylA.

During the time course, 1 mL samples of the ZW641, ZW641 - ara355-1, ZW641 -ara356-2, and ZW641 -ara357-2 cultures were collected at the 72-hour point. They were centrifuged at 10,000x g to remove cells. The supernatant was filtered through a 0.22 μ m Costar Spin-X Centrifuge Tube Filter and analyzed by running through a BioRad Aminex HPX-A7H ion exclusion column with 0.01 N H₂SO₄ in a speed of 0.6 mL/min at 55°C on an Agilent 1100 HPLC system to determine ethanol and xylose concentrations. The results given in Table 5 show that, comparing to the basal level xylose utilization and ethanol production in ZW641, AMxylA in ZW641 -ara355 significantly promoted xylose consumption and had increased ethanol production by more than 3.5 fold. ECxylA in ZW641 -ara357 slightly increased xylose metabolism and ethanol production, while LBxylA in ZW641-ara356 offered the smallest increase in xylose utilization and did not cause a detectable change in ethanol production. These results agree with previous observations on the growth and suggest that the difference in growth between the strains was caused by the difference in xylose metabolism, which may result from a difference in xylose isomerase activity.

Table 5. Cell growth, xylose consumption, and ethanol production after 72 hours culturing at 30°C in MRM3X10.

Strain	Additional XI source	Growth (OD ₆₀₀)	Ethanol (g/L)	Xylose* (g/L)
ZW641	none	0.46	1.9	93.1
ZW641-ara355-1	<i>A. missourinensis</i>	1.14	6.8	81.7
ZW641-ara356-2	<i>L. brevis</i>	0.53	1.9	92.9
ZW641-ara357-2	<i>E. coli</i>	0.61	2.2	91.7
MRM3X10 ⁺	na [#]	na	0.0	95.8

*xylose remaining in the medium

⁺starting media

[#]na: not applicable

To determine whether xylose isomerase enzymes encoded by AMxylA, LBxylA, and ECxylA have different activities, ZW641, ZW641-ara355-1, ZW641-ara356-2, and ZW641-ara357-2 were grown overnight in MRM3G5 on a 30°C 150 rpm shaker. Cells were collected from 2 ml_

5 cultures by 10,000 x g centrifugation, washed with ice-cold Protein Extraction Buffer (10 mM triethanolamine hydrochloride, pH8.0, 10 mM MgSO₄, 1 mM DTT, and 5% glycerol), resuspended in 500 µl_ ice-cold Protein Extraction Buffer, and subjected to 3 minutes sonication for 3 times at setting 7 by using a Misonix Sonicator 4000 with microplate horn

10 (Qsonica, Newtown, CT). Cell debris was removed by 10,000 x g centrifugation. Supernatants were kept as protein extracts. Their protein concentrations were measured by using Coomassie Plus Protein Assay Reagent (Pierce, Rockford, IL), and xylose isomerase activities were measured by a modified Cysteine-Carboazole method. A 100 µl_

15 Cysteine-Carboazole assay reaction contained 10 mM triethanolamine hydrochloride buffer, pH7.0, 10 mM MgSO₄, 25 mM D-xylose, and 30 µg extracted protein. After incubation at 32 °C for 15 minutes, the reaction was stopped by adding 25 µl 50% trichloroacetic acid. Then, 3 ml ice-cold 75% sulphuric acid, 100 µl 2.4% cysteine hydrochloride solution, and 100

20 µl 0.12% carbazole ethanolic solution were sequentially added into the reaction. The mixture was kept at room temperature for 10 minutes. OD₅₄₀ value was measured on a Shimadzu UV-1200 Spectrophotometer. The corresponding D-xylulose concentration was determined based on a standard curve of D-xylulose concentration vs. OD₅₄₀ value. The standard

25 curve was developed by carrying out Cysteine-Carboazole assays containing various amounts of D-xylulose but no D-xylose and protein. Finally, one unit of xylose isomerase enzyme was defined as the activity required to produce one micromole of D-xylulose in the reaction. Specific activity was calculated as unit per milligram of protein. In the assay,

30 background OD₅₄₀ of D-xylose was measured in a blank reaction without protein. This background was subtracted from the original OD₅₄₀ reading prior to calculation of activity. Each assay was repeated 3 times. The results of the Cysteine-Carboazole assay are graphed in Figure 5 showing

specific activities of xylose isomerase in the protein extracts of ZW641 , ZW641 -ara355-1 , ZW641 -ara356-2, and ZW641 -ara357-2. Each activity bar represents an average of three parallel reactions with a standard deviation calculated based on them. This result demonstrates that

5 expression of the second copy of xylA in ZW641 introduced additional xylose isomerase activity into cells. AMxylA in ZW641 -ara355, LBxylA in ZW641 -ara356, and ECxylA in ZW641 -ara357 increased XI activity by approximately 20- fold, 3 fold, and 4 fold, respectively. Since these three xylA genes are constructed in ZW641 by the same approach, the

10 difference in the xylose isomerase specific activities in the cell extracts suggests that the xylose isomerase from *A. missouriensis* functions much better than xylose isomerases from *L. brevis* and *E. coli*. In fact, the *A. missouriensis* XI presented a specific activity of about 2.3, which was 6 times higher than *L. brevis* XI and 5 times higher than *E. coli* XI.

15 Therefore, by examining cell growth, xylose metabolism, and XI activity, this example has identified *A. missouriensis* XI as an excellent xylose isomerase for improving xylose utilization in xylose-utilizing *Z. mobilis* strains.

20

Example 4

Structural Analysis of Xylose Isomerase Enzymes

A collection of available protein sequences that are potentially xylose isomerases was prepared by first identifying a set of seed sequences that are known to have xylose isomerase (XI) activity. The

25 seed sequences were retrieved as xylose isomerases from the SWISSPROT database, which contains protein sequences that have high confidence functional annotations. There were 180 XI seed sequences retrieved from SWISSPROT (Swiss Institute of Bioinformatics):SEQ ID

30 NOs:2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, 142 and 148-306. These seed sequences were then used to search the NCBI (National Center for Biotechnology Information, Bethesda, MD) non-redundant (nr) comprehensive protein database as a group of multiple query sequences in the blastall wrapper of BLAST. A

total of 444 sequences were identified to form a set that can be described as the sequence space of XI activity proteins.

Clustering based on sequence identity and molecular phylogenetic analysis using the PHYLIP neighbor joining algorithm (as implemented in PHYLIP (Phylogeny Inference Package) version 3.5c (Felsenstein (1989) Cladistics 5:164-166) showed that the above generated sequence space for XI activity separated into two groups referred to as Group I and Group II as shown in Figure 2. The Group I set consisted of 82 sequences (SEQ ID NOs that are even numbers between 2 and 130 and 131-147), 21 of which were seeds (SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142). Similarly, the Group II set consisted of 351 members, 159 of which were seeds (SEQ ID NOs: 148-306). As shown in Figure 2, the *L. brevis* and *E. coli* XI proteins belong to Group II while the *A. missouriensis* XI protein belongs to Group I.

The following process was followed in forming the phylogenetic groups:

Starting with the 444 XI sequences:

Step 1 establishing 70% identity groups:

The longest sequence out of the 444 sequences was designated as the first master. Other sequences of the 444 sequences that have 70% or more sequence identity to the first master were grouped to form the first ref70 cluster (A). Out of the remaining sequences, the longest sequence was designated the second master and used to create the second ref70 cluster similarly (B). The grouping process was continued until every sequence was in a cluster. Some of the clusters were singletons.

Step2 merging at 70% threshold:

For every pair of clusters A and B, if a third of the sequences in A are related to sequences in B by 70% sequence identity or more and vice-versa, clusters A and B were merged. This process was continued until there were no pairs that could be merged using the 70% identity threshold.

Step3 merging at 50% threshold:

The same process as in step 2 was followed but using a sequence identity threshold of 50%.

Step4 merging at 30% threshold:

The same process as in step 2 was followed but using a sequence
5 identity threshold of 30%.

Group I represents a 50% threshold identity cluster. Group II represents a separate 50% threshold identity cluster.

There were 11 sequences that were not unambiguously assigned to Group I or Group II since they did not cluster with either group at the 50%
10 threshold of identity.

Figure 3 shows a phylogenetic tree for the Group I XIs with specific genera labeled. Group I includes XI proteins from *Arthrobacter*,
Streptomyces, *Thermus*, *Thermobaculum*, *Herpetosiphon*, *Acidobacteria*,
Roseiflexus, *Meiothermus*, *Deinococcus*, *Meiothermus*, *Stackebrandtia*,
15 *Kribbella*, *Xylanimonas*, *Nocardiopsis*, *Catenulispora*, *Streptosporangium*,
Geodermatopyilus, *Actinosynnema*, *Saccharomonospora*, *Acicothermus*,
Tthermobifida, *Nocardiodes*, *Janibacter*, *Mycobacterium*, *Leifsonia*,
Clavibacter, *Micromonospora*, *Salinispora*, *Cellulomonas*, *Jonesia*,
Nakamurella, *Actinomyces*, *Mobiluncus*, *Brachybacterium*, *Beutengergai*,
20 *Frankia*, and *Actinobacterium*.

Discriminating between Group I and Group II: Method 1

Discrimination between Group I members and Group II members was performed by GroupSim analysis (Capra and Singh (2008)
25 Bioinformatics 24: 1473-1480). The GroupSim method identifies amino acid residues that determine a protein's functional specificity. In a multiple sequence alignment (MSA) of a protein family whose sequences are divided into multiple groups, amino acid residues that distinguish between the functional groups of sequences can be identified. The method takes a
30 multiple sequence alignment (MSA) and known specificity groupings as input, and assigns a score to each amino acid position in the MSA. Higher scores indicate a greater likelihood that an amino acid position is a specificity determining position (SDP).

GroupSim analysis performed on the MSA of XI sequences that were divided into Group I and Group II by the phylogenetic analysis above identified highly discriminating positions. Listed in Table 6 are positions (Pos) having scores greater than or equal to 0.9, where a perfect score of 1.0 would indicate that all proteins within the group have the listed amino acid in the specified position and between groups the amino acid would always be different. The "residue" column gives the amino acid(s) in single letter code for the position in Group I proteins vs in Group II proteins (separated by a bar: |). The amino acid position number in column 2 is for the representative Group I protein P12581 which is the XI from *Actinoplanes missourinesis*. The amino acid position number in column 3 is for the representative Group II protein P19148 which is the XI *Thermoanaerobacterium thermosulfurigenes* (SEQ ID NO:267)

Table 6 Highly discriminating amino acid positions for Group I and Group II XIs from GroupSim analysis.

Pos in alignment	Pos in P12581 (member of Group I)	Pos in P19148 (member of Group II)	Residue (Group I Group II)
391	226	277	L H
388	223	274	M L
341	191	242	I Q
345	195	246	TSV D
228	88	139	MTG RW
462	290	337	H NM
386	221	272	ED ATG
407	242	293	FVL GCW
408	243	294	H SNGL
343	193	244	LFM D
422	256	308	Q TIVLMYH
377	213	264	G KNSEALRQ
460	288	335	PYAS VG
415	249	301	Q DHNRSA

Discriminating between Group I and Group II: Method 2

An alternative structure/function characterization of the groups of the xylose isomerase family of enzymes was performed using the HMMER software package (the theory behind profile HMMs is described in R.

- 5 Durbin, S. Eddy, A. Krogh, and G. Mitchison, Biological sequence analysis: probabilistic models of proteins and nucleic acids, Cambridge University Press, 1998; Krogh et al., 1994; J. Mol. Biol. 235:1 501-1 531), following the user guide which is available from HMMER (Janelia Farm Research Campus, Ashburn, VA).

- 10 Using a multiple sequence alignment of the 21 seed sequences in Group I (SEQ ID NOS:2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142), a profile Hidden Markov Model (HMM) was created for representing Group I members. As stated in the user guide, Profile HMMs are statistical models of multiple sequence
15 alignments. They capture position-specific information about how conserved each column of the alignment is, and which amino acid residues are most likely to occur at each position. Thus HMMs have a formal probabilistic basis. Profile HMMs for a large number of protein families are publicly available in the PFAM database (Janelia Farm
20 Research Campus, Ashburn, VA).

The Profile HMM was built as follows:

Step 1. Build a sequence alignment

- The 21 seed sequences (sequences with high confidence annotation) that are in Group I were aligned using Clustal W with default
25 parameters.

Step 2. Build a Profile HMM

- The hmmbuild program was run on the set of aligned sequences using default parameters. hmmbuild reads the multiple sequence alignment file, builds a new Profile HMM, and saves the Profile HMM to
30 file. Using this program an un-calibrated profile was generated from the multiple alignment for the set of seed sequences described above.

The following information based on the HMMER software user guide gives some description of the way that the hmmbuild program prepares a Profile HMM. A Profile HMM is capable of modeling gapped

alignments, e.g. including insertions and deletions, which lets the software describe a complete conserved domain (rather than just a small ungapped motif). Insertions and deletions are modeled using insertion (I) states and deletion (D) states. All columns that contain more than a certain fraction x of gap characters will be assigned as an insert column. By default, x is set to 0.5. Each match state has an I and a D state associated with it. HMMER calls a group of three states (M/D/I) at the same consensus position in the alignment a "node". These states are interconnected with arrows called *state transition probabilities*. M and I states are emitters, while D states are silent. The transitions are arranged so that at each node, either the M state is used (and a residue is aligned and scored) or the D state is used (and no residue is aligned, resulting in a deletion-gap character, '-'). Insertions occur between nodes, and I states have a self-transition, allowing one or more inserted residues to occur between consensus columns.

The scores of residues in a match state (i.e. match state emission scores), or in an insert state (i.e. insert state emission scores) are proportional to $\text{Log}_2(p_x / (\text{null}_x))$. Where p_x is the probability of an amino acid residue, at a particular position in the alignment, according to the Profile HMM and null_x is the probability according to the Null model. The Null model is a simple one state probabilistic model with pre-calculated set of emission probabilities for each of the 20 amino acids derived from the distribution of amino acids in the SWISSPROT release 24.

State transition scores are also calculated as log odds parameters and are proportional to $\text{Log}_2(t_x)$. Where t_x is the probability of transiting to an emitter or non-emitter state.

Step 3. Calibrate the Profile HMM

The Profile HMM was read using `hmmcalibrate` which scores a large number of synthesized random sequences with the Profile (the default number of synthetic sequences used is 5,000), fits an extreme value distribution (EVD) to the histogram of those scores, and re-saves the HMM file now including the EVD parameters. These EVD parameters (μ and λ) are used to calculate the E-values of bit scores when the profile is

searched against a protein sequence database. hmmcalibrate writes two parameters into the HMM file on a line labeled "EVD": these parameters are the μ (location) and λ (scale) parameters of an extreme value distribution (EVD) that best fits a histogram of scores calculated on

5 randomly generated sequences of about the same length and residue composition as SWISS-PROT. This calibration was done once for the Profile HMM.

The calibrated profile HMM for the Group I set is provided as Table 1 in the appendix as a Group I profile HMM Excel chart. The Profile HMM

10 is provided in a chart that gives the probability of each amino acid occurring at each position in the amino acid sequence. The highest probability is highlighted for each position. Table 7 shows a few lines of the Group I Profile HMM.

15 Table 7. A portion of the Group I Profile HMM.

HMM	A	C	D	E	F	G	H	I	K	L	M	N	P	Q	R	S	T	V	W	Y
	m->m	m->i	m->d	i->m	i->i	d->m	d->d	b->m	m->e											
	-462	*	-1868																	
1	-1476	-1441	-2702	-2503	-679	-2424	-1861	-88	-2024	240	4620	-2293	-2670	-1975	-1997	-1977	-1568	-381	-1705	-1336
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-	-24	-6482	-7525	-894	-1115	-701	-1378	-462	*											
2	421	-976	-556	-40	-1303	-1252	-182	-920	65	-1119	1573	1057	-1439	138	-354	1386	371	-651	-1514	-1000
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-	-21	-6672	-7714	-894	-1115	-701	-1378	*	*											
3	-926	-719	-2839	-2316	1366	-2394	-664	122	-1975	1177	430	-1816	-2377	-1532	-1832	-1492	-859	972	-254	2848
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-	-21	-6672	-7714	-894	-1115	-701	-1378	*	*											

The amino acids are represented by the one letter code.

The first line for each position reports the match emission scores: probability for each amino acid to be in that state (highest score is

20 highlighted). The second line reports the insert emission scores, and the third line reports on state transition scores: M->M, M->I, M->D; I->M, I->I; D^M, D^D; B^M; M^E. Table 7 shows that in the Group I profile HMM, methionine has a 4620 probability of being in the first position, the highest probability.

25 Step 4. Test the specificity and sensitivity of the built Profile HMM

The Group I profile HMM was evaluated using hmmsearch, with the Z parameter set to 1 billion, for the ability to discriminate Group I members from those of Group II. The hmmsearch program takes the hmm file for the Group I profile HMM and all the sequences from both groups and assigns

an E-value score to each sequence. This E-value score is a measure of fit to the Profile HMM, with a lower score being a better fit. The resulting score assignment list is provided in the appendix as Table 2. The Profile HMM clearly distinguished Group I members from Group II members since
 5 there was a large margin of E-value difference between the worst scoring Group I member (2.2×10^{-81}) and the best scoring Group II member (1.5×10^{-07}).

This analysis shows that the Profile HMM prepared for Group I XI proteins distinguishes Group I from Group II XI proteins. The Group I
 10 Profile HMM provides a structure that is linked to XI proteins that are functionally similar to XI of *A. missouriensis*.

Example 5

Construction of chimeric P_{gap} -xylA Genes for Group 1 XIs and Assembly 15 of Double Crossover suicide Vectors

The xylose isomerases from *Geodermatophilus obscurus* DSM 43160 (GOxylA; SEQ ID NO:64), *Mycobacterium smegmatis* str. MC2 155 (MSxylA; SEQ ID NO:10), *Salinispora arenicola* CNS-205 (SAxylA; SEQ ID NO:18), and *Xylanimonas cellulosilytica* DSM 15894 (XCxylA; SEQ ID
 20 NO:40) all belong to Group 1 xylose isomerases, based on amino acid sequence analysis described in Example 4. The sequences encoding these proteins were each optimized (SEQ ID NOs:335, 336, 337, and 338, respectively) for expression in *Z. mobilis* according to codon bias of *Z. mobilis* ZM4, and synthesized de novo as DNA fragments bounded by
 25 SpeI and XhoI sites, and with the coding region adjacent to a mutant *Z. mobilis* glyceraldehyde-3-phosphate dehydrogenase gene promoter (Pgaps; SEQ ID NO:339) by GenScript Corporation (Piscataway, NJ). The Pgaps is an improved P_{gap} that has a mutation which is a "G" to "T" change at position 116 of the natural promoter fragment (P_{gap}), that increases
 30 expression from the promoter, as disclosed in US 2009-0246876. The SpeI-Pgaps-xylA-XhoI fragments were cloned into pUC57 at the EcoRV site by GenScript Corporation (Piscataway, NJ). The resultant intermediate plasmids were called pUC57- P_{gap} sGOxylA, pUC57- P_{gap} sMSxylA, pUC57- P_{gap} sSAxylA, and pUC57- P_{gap} sXCxylA. The optimized xylA coding

sequences are GOxylA (SEQ ID NO:335), MSxylA (SEQ ID NO:336),
 SAXylA (SEQ ID NO:337), and XCxylA (SEQ ID NO:338).

Common molecular cloning methods were used to construct DCO
 suicide vectors. First, plasmid pARA356 (described in Example 1) was
 5 modified to add an XhoI site between the LBxylA and araD3'UTR
 sequences as follows. First, an araD3'UTR fragment was PCR amplified
 from pARA356 using forward primer ara368 (SEQ ID NO:345) and reverse
 primer ara97 (SEQ ID NO:319). The ara368 primer added SpeI and XhoI
 sites to the 5' end of araD3'UTR, and ara97 added HindIII, FseI, and
 10 EcoRI sites to the 3' end of araD3'UTR. The PCR product was digested
 with SpeI and EcoRI. The P_{gap}-LBxylA-araD3'UTR segment in pARA356
 has a 5' SpeI site and a 3' EcoRI site. This segment was removed by
 digestion with SpeI and EcoRI and it was replaced by the above SpeI-
 XhoI-araD3'UTR-HindIII-FseI-HindIII-EcoRI PCR product. The resulting
 15 intermediate plasmid pARA356D has same sequence as pARA356,
 except P_{gap}-LBxylA was replaced by a XhoI site.

The four P_{gap}^S-xylA fragments described above were isolated from
 the pUC57-based plasmids following SpeI and XhoI digestion, and cloned
 into the XhoI-modified pARA356D between the SpeI and XhoI sites to
 20 replace the P_{gap}-LBxylA fragment. The resulting four DCO suicide vectors
 were pARA356-GOxylA, pARA356-MSxylA, pARA356-SAxylA, and
 pARA356-XCxylA. These vectors are identical to pARA356 except that
 their chimeric xylA genes are expressed from P_{gap}^S-

As controls, AMxylA in pARA355 and ECxylA in pARA357 were
 25 used as representatives for Group I and II xylAs, respectively. However,
 since P_{gap}^S was employed to express the four new Group 1 xylA genes,
 the P_{gap} promoters controlling AMxylA in pARA355 and ECxylA in
 pARA357 were changed to P_{gap}^S. For this purpose, a 319-bp P_{gap}^S OLE-
 PCR fragment was synthesized from pARA356-XCxylA by PCR, using
 30 forward primer ara10 (SEQ ID NO:340) and reverse primer ara401 (SEQ
 ID NO:341); a 1,229-bp P_{gap}^S-AMxylA OLE-PCR fragment was
 synthesized from pARA355 by PCR, using forward primer ara402 (SEQ ID
 NO:342) and reverse primer ara403 (SEQ ID NO:343); and a 1,367-bp
 P_{gap}^S-ECxylA OLE-PCR fragment was synthesized from pARA357 by

PCR, using forward primer ara402 and reverse primer ara404 (SEQ ID NO:344). One PCR reaction consisted of 50 μ L AccuPrime Pfx SuperMix (Invitrogen, Carlsbad, CA), 1 μ L of 40 ng/ μ L DNA template, and 1 μ L of 10 μ M forward and reverse primers. Reactions were carried out on an

5 Eppendorf Mastercycler (Hamburg, Germany), following a hot start PCR program with 35 cycles of denaturing at 94 °C for 1 min/ annealing at 56 °C for 1 min / extension at 72 °C for 2 min. The P_{gap} s OLE-PCR fragment included the entire P_{gap} s, a 5' Spel site, and a 3' start codon. The P_{gap} s-AMxylA and P_{gap} s-ECxylA OLE-PCR fragments contained an AMxylA and

10 an ECxylA coding sequence, respectively. Each had a 36-nt 5' sequence that matches the last 36 nt of P_{gap} s and a 3' XhoI site. Furthermore, Spel- P_{gap} s-AMxylA-XhoI and Spel- P_{gap} s-ECxylA-XhoI fragments were synthesized by overlapping PCR (OLE-PCR). PCR reactions were set up as described above, but two templates were included. Spel- P_{gap} s-AMxylA-

15 XhoI was amplified from P_{gap} s and P_{gap} s-AMxylA OLE-PCR fragments by using forward primer ara10 and reverse primer ara403, while Spel- P_{gap} s-ECxylA-XhoI was amplified from P_{gap} s and P_{gap} s-ECxylA OLE-PCR fragments by using forward primer ara10 and reverse primer ara404. Both

20 Spel- P_{gap} s-AMxylA-XhoI and Spel- P_{gap} s-ECxylA-XhoI were digested with Spel and XhoI, subjected to agarose gel electrophoresis, and purified by using QIAquick Gel Purification Kit (Qiagen). The DNA fragments were cloned into modified pARA356 (described above) between Spel and XhoI sites to replace the P_{gap} -LBxylA fragment. The resulting two DCO suicide vectors were called pARA356-AMxylA and pARA356-ECxylA. All vectors

25 were propagated in DH5a *E. coli* cells and prepared by using a QIAprep Spin Miniprep Kit.

Example 6

Integration of Chimeric P_{nanS} -xylA Genes into ZW641 and Characterization of Their Expression

This Example describes integration and expression of P_{gap} s-xylA chimeric genes in strain ZW641, described in Example 2, and

demonstrates that the four tested Group I XIs indeed function better than Group II XIs in *Z. mobilis*.

Competent cells of strain ZW641-1A were prepared and transformed separately with pARA356-AMxylA, pARA356-ECxylA, pARA356-GOxylA,
 5 pARA356-MSxylA, pARA356-SAxylA, or pARA356-XCxylA.

Transformants were selected on MMG5-Spec250 plates and analyzed by PCR for integration of the introduced P_{gap} -xylA genes as described previously (see Example 2). The resultant strains were named ZW641 - ara356-AMxylA, ZW641-ara356-ECxylA, ZW641 -ara356-GOxylA, ZW641 - ara356-MSxylA, ZW641 -ara356-ASxylA, and ZW641 -ara356-XCxylA
 10 strains. Among these strains, ZW641 -ara356-AMxylA and ZW641- ara356-ECxylA were made as control strains since Example 3 demonstrated that the AMxylA Group 1 xylose isomerase was highly active in *Z. mobilis*, while ECxylA was the better enzyme of two tested
 15 Group II xylose isomerases.

To examine the growth of these six new strains in xylose and compare them with the parental strain ZW641-1A, all strains were subjected to a 96-hour shake flask fermentation in xylose. In the assay, each strain was grown overnight in 3 mL MRM3G5 at 30 °C with 150 rpm
 20 shaking. Cells were harvested, washed with MRM3X10 (same as MRM3G5 but 50 g/L glucose was replaced with 100 g/L xylose), and resuspended in MRM3X10 to OD₆₀₀ of about 0.1. Twenty-five milliliters of the suspension were placed in a 50 mL screw capped VWR centrifuge tube and grown at 30 °C with 150 rpm shaking for a 96-hour time course.
 25 During the time course, OD₆₀₀ was measured at 0, 24, 48, 72, and 96 hours. The resulting growth curve is shown in Figure 6. It shows that similar to ZW1-ara355 and ZW1-ara357 strains analyzed in Example 3, ZW641 -ara356-AMxylA grew to a cell density approximately three times higher than ZW641-ara356-ECxylA at end of the fermentation. ZW641 - ara356-AMxylA had an OD₆₀₀ of 3.43, while ZW641 -ara356-ECxylA
 30 reached 1.18. Both strains grew faster than ZW641. The other four strains all grew faster than ZW641 -ara356-ECxylA. Their cell densities at end of the fermentation were between those of ZW641 -ara356-AMxylA and ZW641 -ara356-ECxylA.

To measure the metabolic profile of each strain, a 1 ml sample of each culture was collected at the 72-hour point. The samples were centrifuged at 10,000x g to remove cells. The supernatant was filtered through a 0.22 μ m Costar Spin-X Centrifuge Tube Filter and analyzed by running through a BioRad Aminex HPX-A7H ion exclusion column with 0.01 N H₂SO₄ at a speed of 0.6 mL/min at 55 °C on an Agilent 1100 HPLC system to determine ethanol and xylose concentrations. The results given in Table 8 show that faster growth correlated with higher xylose utilization and more ethanol production. These results suggest that the difference in growth is due to the difference in XI activity. All strains had better growth, ethanol production and xylose utilization than the ZW641 control, and all strains with Group 1 XIs performed better than the strain with the Group II *E. coli* XI.

Table 8. Cell growth, xylose consumption, and ethanol production after 72 hours culturing at 30 °C in MRM3X10.

Strain	Growth (OD ₆₀₀)	Ethanol (g/L)	Xylose* (g/L)
ZW641	0.22	0.0	96.1
ZW641-ara356-AMxylA	3.21	26.3	41.5
ZW641-ara356-ECxylA	0.59	1.6	92.9
ZW641-ara356-GOxylA	1.14	3.0	90.0
ZW641-ara356-MSxylA	1.75	8.4	78.6
ZW641-ara356-SAxylA	1.51	5.0	86.3
ZW641-ara356-XCxylA	2.80	19.4	56.1
MRM3X10 ⁺	na [#]	0.0	96.1

*xylose remaining in the medium

⁺starting medium

[#]na: not applicable

20

To further confirm that these six strains each had an XI activity level corresponding to their growth and metabolic profiles, protein extracts were prepared and assayed for protein concentration and xylose isomerase activity as in Example 3. Specific activity was calculated as unit per milligram of protein with background OD₅₄₀ of D-xylose subtracted from

25

the original OD₅₄₀ reading prior to calculation of activity. Each assay was repeated 3 times. Figure 7 shows the resulting average specific XI activities in each protein extract. The absolute specific activities are lower than those in Example 3, most likely due to different reaction conditions.

- 5 However, comparison of relative specific activities clearly shows that the XI activities correspond to the growth rates of the strains. Faster growth is supported by higher XI activity, with the AMxylA having highest activity and ECsylA the lowest activity. All of the Group I XIs have higher average activities than the Group II ECxylA.

10

TABLE 1

HMIMER2.0 [2.2g]¹
 NAME seed-short²
 LENG 397³
 ALPH Amino⁴
 MAP yes⁵
 NSEQ 21⁶

DATE Mon Oct 19 14:08:50 2009⁷

XT -8455 -4 -1000 -1000 -8455 -4 -8455 -4

NULT -4 -8455⁸

NULE 595-1558 85 338 -294 453 -1158 197 249 902 -1085 -142 -21 -313 45 531 201 384 -1998 -644⁹

HMM	A	C	D	E	F	G	H	I	K	L	M	N	P	Q	R	S	T	V	W	Y
	m->m	m->i	m->d	i->m	i->i	d->m	d->d	b->m	m->e											
	-462 *		-1868																	
1(M)	-1476	-1441	-2702	-2503	-679	-2424	-1861	-88	-2024	240	4620	-2293	-2670	-2670	-1975	-1997	-1977	-1568	-381	-1705
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	394	45	96	359	117	-369	-294
-	-24	-6482	-7525	-894	-1115	-701	-1378	-462 *												-249
2(M)	421	-976	-556	-40	-1303	-1252	-182	-920	65	-1119	1573	1057	-1439	138	-354	-354	1386	371	-651	-1514
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
3(Y)	-926	-719	-2839	-2316	1366	-2394	-664	122	-1975	1177	430	-1816	-2377	-1532	-1832	-1492	-859	972	-254	-2848
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
4(Q)	-587	-1894	98	1129	-2216	-1318	-135	-1916	288	-1916	-1086	6	-1535	2856	-133	-486	476	-1535	-2124	-1485
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
5(P)	963	-839	-1538	-1576	-2513	-1059	-1614	-2104	-1628	-2420	-1708	-1177	3418	-1478	-1771	-1771	-442	-597	-1456	-2726
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
6(T)	-582	-1338	-802	-473	-2030	-1419	-515	-1548	1250	-1750	-1035	-569	-1741	-184	-76	-652	2793	-1225	-2043	-1605
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
7(P)	-453	-1271	-792	-423	-2113	-1308	-480	-1789	138	-1875	-1095	-508	2771	-134	1323	87	-556	-1368	-2082	-1622
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
8(E)	540	-2335	997	2666	-2648	-1254	-347	-2412	-241	-2400	-1594	99	-1622	34	-857	-639	-843	-1973	-2617	-1835
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
9(D)	-1212	-2686	3205	333	-2575	-1370	2184	-2849	-658	-2784	-2085	-32	-1828	-282	-1310	-968	-1255	-2414	-2702	-1831
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249
10(R)	-1360	-2277	-1623	-673	-2899	-2052	1299	-2378	2306	-2139	-1406	-717	-2040	268	2661	-1247	-1137	-2097	-2071	-1820
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	96	359	117	-369	-294
-	-21	-6672	-7714	-894	-1115	-701	-1378 *	*												-249

TABLE 1

11(F)	-2605	-2164	-3410	-3518	4186	-3034	-940	-1504	-3300	-1072	-1142	-2698	-3246	-2697	-2999	-2815	-2670	-1743	-274	806	11
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
12(T)	-29	-659	-1587	-1466	-2300	-927	-1414	-1919	-1399	-2228	-1440	-1037	-1583	-1249	-1574	1473	2936	-1259	-2551	-2190	12
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
13(F)	-2605	-2164	-3410	-3518	4186	-3034	-940	-1504	-3300	-1072	-1142	-2698	-3246	-2697	-2999	-2815	-2670	-1743	-274	806	13
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
14(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	14
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
15(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	15
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
16(W)	-3232	-2594	-3508	-3648	-882	-2917	-1886	-3060	-3237	-2665	-2615	-3229	-3282	-3161	-2996	-3457	-3341	-3103	6081	-495	16
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
17(T)	-665	-1148	-2040	-2098	-2442	-1424	-1936	-1850	-1956	-2305	-1815	-1645	-2071	-1911	-2001	-909	3609	-1444	-2672	-2426	17
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
18(V)	-1365	-986	-3741	-3350	-1294	-3426	-2948	1806	-3144	-231	-144	-3097	-3349	-2976	-3194	-2702	-1365	3087	-2628	-2205	18
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
19(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	19
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
20(W)	-1622	-1592	-2164	-1843	1022	-2585	-106	-1330	-1522	-1235	-864	1189	-2631	-1288	-1607	-1710	-1559	-1277	4882	2199	20
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
21(Q)	-321	-1138	-459	136	-1344	-1359	-136	-875	203	-1115	-365	-201	-1496	2306	-180	-368	1172	505	-1531	-1013	21
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
22(G)	1024	-757	-1652	-1777	-2766	3031	-1806	-2449	-1965	-2750	-1944	-1230	-1687	-1701	-2063	-356	-543	-1600	-2942	-2694	22
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
23(R)	1048	-1275	-1018	-415	-1643	-1529	-292	-1159	404	-641	-643	-522	-1703	44	2555	-628	-541	-930	-1713	-1274	23
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
24(D)	-2059	-2867	3660	-532	-3452	-1877	-1516	-3616	-1847	-882	-2414	-1318	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	24
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

[illegible]

TABLE 1

38(E)	-909	-1965	24	2744	-1246	-1505	-347	-1841	-213	-1874	-1207	-228	-1785	-120	-648	-805	-890	-1573	-1601	1568	38
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
39(A)	1463	-630	-1276	-809	-1410	236	-731	-999	-672	-1275	-522	-711	-1520	-537	-976	877	1188	793	1715	-1298	39
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
40(V)	-529	-777	-2238	-2105	-1525	-7	-1733	111	-1947	-1096	-672	-1661	-2112	-1793	-1998	-945	-758	3083	-2136	-1758	40
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
41(H)	-582	-1877	-351	773	-2233	-1449	2951	-1912	686	-1839	-988	-97	-1545	1646	1256	-467	-492	-1531	-1962	-1406	41
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
42(R)	-1587	-2371	-1979	-913	-3101	-2190	-224	-2514	2541	-2236	-1542	-885	-2171	194	2749	-1477	-1330	-2260	-2127	-1955	42
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
43(L)	-1974	-1575	-4060	-3545	-282	-3812	-2663	1310	-3204	2683	857	-3408	-3401	-2583	-3003	-3063	-1891	178	-1830	-1733	43
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
44(A)	2637	-604	-1805	-1733	-2482	-871	-1613	-2129	-1700	-2432	-1611	-1136	-1569	-1478	-1829	1682	-361	-1353	-2728	-2414	44
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
45(E)	-1003	-2473	512	2948	-2788	-831	-430	-2578	-247	-2533	-1768	24	-1719	707	-767	-793	-1016	-2149	-2707	-1955	45
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
46(L)	-1959	-1558	-4063	-3546	-291	-3808	-2666	1414	-3212	2654	850	-3406	-3399	-2589	-3010	-3056	-1876	210	-1837	-1742	46
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
47(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	47
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
48(A)	2866	-638	-2146	-1978	-1831	-1146	-1655	-629	-1782	-1443	-953	-1374	-1775	-1605	-1861	-461	-489	571	-2360	-2006	48
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
49(Y)	-2410	-2105	-2586	-2487	1214	-3046	2815	-1980	-2072	-1592	-1442	-1863	-3075	-1758	-2026	-2294	-2336	-2000	511	4049	49
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
50(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	50
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
51(V)	-1387	-977	-3821	-3371	-1118	-3516	-2833	2100	-3172	597	34	-3124	-3343	-2908	-3170	-2752	-1359	2763	-2430	-2081	51
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
52(T)	-	-	-316	-1039	-679	-633	-2149	-1803	-741	-2089	-1321	1353	-1653	-671	-1045	-444	2964	-1311	-2344	-1843	52		
-	-	-	-149	-500	233	43	-381	399	106	-626	-720	275	394	45	96	359	117	-369	-294	-249			
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
53(F)	-	-	-2065	-1675	-3615	-3327	3564	-3380	-1003	-121	-2974	1209	326	-2658	-3213	-2303	-2729	-2622	-1996	-549	-310	671	53
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
54(H)	-	-	-2323	-2419	-1868	-1956	-1368	-2396	5124	-3143	-1645	-2924	-2595	-1989	-2806	-1873	-1712	-2381	-2439	-2923	-1719	-935	54
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
55(D)	-	-	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	55
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
56(D)	-	-	-1205	-3003	2930	1455	-3203	-1271	-537	-3082	-663	-2979	-2251	1491	-1755	-188	-1433	-903	-1247	-2583	-3173	-2230	56
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
57(D)	-	-	-1425	-3134	3461	881	-3374	-1343	-726	-3296	-972	-3221	-2585	25	-1881	-410	-1787	-1106	-1504	-2808	-3355	-2435	57
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
58(L)	-	-	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	58
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
59(I)	-	-	-584	-415	-2576	-2004	811	-2186	-1087	2394	-1704	-44	421	-179	-2212	-1418	-1667	-1280	-535	1582	-1034	-637	59
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
60(P)	-	-	831	-858	-1530	-1581	-2527	-1073	-1627	-2121	-1640	-2436	-1732	-1189	3462	-1494	-1780	-462	-619	-1475	-2736	-2431	60
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
61(F)	-	-	-1404	-1445	-2265	-1857	3570	-2371	-514	-956	-1089	-826	-548	-1590	-2505	-1217	1154	-1633	-1384	-986	-276	725	61
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
62(G)	-	-	-1147	-2145	1555	-151	-3123	2941	-1064	-3044	-1224	-3073	-2375	-443	-1969	-787	-1831	-1065	-1335	-2467	-3051	-2494	62
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
63(S)	-	-	1540	-619	-1644	-1374	-2259	-896	-1302	-1897	-1273	-2159	-1334	-974	-1528	-1100	-1502	2281	1259	-1223	-2486	-2129	63
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
64(T)	-	-	-278	-1411	820	210	-1980	-1166	-168	-1685	139	-1746	-888	-45	610	235	-352	1310	1395	-1258	-1999	-1378	64
-	-	-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

TABLE 1

65(D)	775	-2108	2304	884	-2418	-1237	-217	-2176	-19	-2160	-1317	112	719	185	-599	-488	-639	-1745	-2375	-1635	65
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
66(Q)	1223	-1288	-293	188	-1719	-1202	-83	-1395	258	-1490	-649	435	-1388	1247	-204	588	1044	-1031	-1779	-1191	66
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
67(E)	-786	-1958	234	2838	-2471	-1312	-479	-2090	-260	-2221	-1480	-118	-1693	-120	-738	-684	705	-1709	-2489	-1828	67
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
68(R)	-2176	-2415	-2504	-1923	-3054	-2403	-1269	-3053	-167	-2856	-2350	-1844	-2662	-997	3887	-2228	-2131	-2807	-2501	-2452	68
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-86	-4238	-7714	-942	-1061	-701	-1378*	*													
69(E)	180	-1985	1618	1858	-2269	-1227	-60	-2029	241	-1986	-1104	146	-1438	1147	-305	-338	-450	-1591	-2174	-880	71
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
70(Q)	-93	-1588	-113	706	-1865	-138	790	-1583	816	-888	-690	129	-1311	1321	26	758	459	-1186	-1798	-1145	72
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
71(H)	-345	-877	-695	-213	-903	-685	2413	1646	31	-776	-65	-380	-1577	906	-283	-494	-289	-359	-1209	-725	73
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
72(I)	-367	-817	-3505	-3009	-998	-3128	-2307	2526	-2771	715	78	-2719	-3035	-2503	-2747	-2310	-1131	2143	-2063	-1704	74
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
73(K)	361	-1699	596	415	-2015	-63	90	-1743	1707	-1710	-809	113	-1347	537	931	-183	-235	-1322	-1892	-1235	75
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
74(R)	-320	-1614	442	360	-1920	-41	31	-1626	443	-1643	-772	621	-1381	460	2112	-229	-270	-563	-1865	-1229	76
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
75(F)	-324	-1039	-2854	-2418	3331	-2568	-1007	160	-2106	857	406	-2027	-2596	-1719	-2040	-1726	-1158	-29	-596	214	77
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
76(K)	-650	-1908	-584	44	-2320	-1534	916	-1955	2061	-1862	-1018	553	-1598	439	1858	2	-543	-1590	-1953	-1448	78
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
77(Q)	71	-1748	-305	251	-2095	-183	48	-1783	1408	-1737	-866	-20	-1452	2274	774	-328	-355	-1392	-1894	-1306	79
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
78(A)	3113	-723	-1876	-1947	-2635	-214	-1832	-2241	-1993	-2601	-1822	-1305	-1695	-1754	-2055	-360	-529	-1479	-2845	-2605	80
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
79(L)	-1818	-1523	-3759	-3310	-316	-3460	-2478	536	-2926	2733	784	-3124	-3244	-2443	-2783	-2733	616	-1782	117	-369	-294	-1623	81
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
80(D)	101	-1863	2365	425	-2206	-1222	-98	-1948	669	-1938	-1069	103	-382	321	-339	397	446	-1522	-2148	-1452	82		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
81(E)	1195	-2299	1673	2036	-2594	-1247	-290	-2366	-151	-2338	-1512	118	-1587	102	-761	-582	-772	-1923	-2546	-1769	83		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
82(T)	-194	-791	-1643	-1654	-2474	22	-1622	-2066	-1652	-2413	-1662	-1202	-1715	-1503	-1776	-414	3374	-1411	-2683	-2390	84		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
83(G)	439	-855	-1581	-1757	-2835	3187	-1850	-2546	-2010	-2843	-2063	-1269	-1756	-1754	-2106	-457	-649	-1703	-2977	-2743	85		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
84(M)	-740	-586	-2676	-2080	-191	-930	-1155	1108	-1758	1717	2785	-1802	-2271	-1414	-1692	-1385	-677	372	-1016	-731	86		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
85(K)	-131	-982	-588	-21	-1093	-1402	-93	359	1662	-893	-147	-231	-1484	210	265	-362	838	279	-1327	-833	87		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
86(V)	-521	-748	-233	-1105	-1131	-1713	-1055	404	-1134	-720	-220	-1100	-2000	-969	-1389	-914	300	2729	-1701	-1269	88		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
87(P)	-1095	-1645	-1418	-1291	-2556	-1705	-1272	-2442	-720	-2498	-1867	-1276	3654	-1049	408	-1217	-1284	-2022	-2467	-2195	89		
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249			
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*															
88(M)	-953	-1122	-1705	-1356	-305	-2019	883	-460	-763	-333	4260	-1283	-2194	-908	-858</								

TABLE 1

92(N)	-1161	-2498	1023	254	-2901	-1337	-713	-2897	-803	-2894	-2199	3546	-1842	-402	-1449	-964	-1262	-2411	-2948	-2145	94
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
93(L)	-1075	-1199	-2228	-2047	-923	-434	-1571	-385	-1753	2693	-140	-1841	-2396	-1672	-1788	-1474	-1191	-504	-1704	-1322	95
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
94(F)	-1621	-1684	-10	-1676	3875	-2340	-642	-1251	-1954	-1200	-952	-1552	-2602	-1572	-2091	-1778	-1682	-1250	-309	743	96
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
95(T)	-248	-1135	326	-166	-2058	-1097	-537	-1731	-339	-1883	-1065	-315	-1509	-198	-783	1772	2073	-1257	-2181	-1628	97
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
96(H)	265	-2079	1321	773	-2325	-1256	3495	-2082	19	-2080	-1247	103	-1537	198	-538	-495	-631	-1679	-2300	-1577	98
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
97(P)	-2089	-2190	-2567	-2787	-3372	-2253	-2663	-3607	-2895	-3590	-3215	-2612	4122	-2846	-2839	-2316	-2418	-3135	-2975	-3234	99
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
98(V)	479	-552	-2113	-1863	-1375	-3	-1443	-29	-1694	-1045	-494	-1385	-1851	-1485	-1768	-608	-468	-2820	-1913	-1556	100
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
99(F)	-2377	-1971	-2987	-2956	3952	-3100	1295	-1675	-2615	-1307	-1196	-2083	-3140	-2054	-2431	-2384	-2334	-1754	478	1631	101
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
100(K)	-186	-1836	-969	-568	-2577	-1685	-425	-2088	3138	-2110	-1410	-652	-1925	-45	429	-978	-981	-1757	-2208	-1848	102
-	-149	-498	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-86	-4238	-7714	-1698	-552	-701	-1378*	*													
101(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	103
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
102(G)	-920	-1508	-1339	-1295	-2751	3192	-1377	-2630	-937	-2716	-2012	-1241	-2085	-1160	523	-1056	-1170	-2071	-2626	-2390	109
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-102	-6672	-4094	-894	-1115	-701	-1378*	*													
103(G)	1678	-647	-1598	-1653	-2595	2694	-1650	-2259	-1777	-2558	-1746	-1116	-1581	-1527	-1899	-241	-417	-1446	-2792	-2519	110
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-22	-6592	-7635	-894	-1115	-618	-1521*	*													
104(F)	-2605	-2164	-3410	-3518	4186	-3034	-940	-1504	-3300	-1072	-1142	-2698	-3246	-2697	-2999	-2815	-2670	-1743	-274	806	111
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
105(T)	-665	-1148	-2040	-2098	-2442	-1424	-1936	-1850	-1956	-2305	-1815	-1645	-2071	-1911	-2001	-909	3609	-1444	-2672	-2426	112
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*												
106(S)	2001	-616	-1754	-1668	-2497	-871	-1578	-1641	-2157	-1619	-2446	-1109	-1565	-1427	-1789	2457	-361	-1370	-2732	-2410	113
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
107(N)	-603	-1690	-182	42	-2219	-1348	-286	-1949	226	-1968	-1162	2998	26	97	673	-546	-618	-1549	-2135	-1563	114
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
108(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	115
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
109(R)	-1150	-1864	-1429	-882	-2591	-1829	-597	-2283	393	-2219	-1523	-923	1080	-238	3277	-1166	-1141	-1945	-2211	-1924	116
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
110(W)	344	-1328	1619	227	-1582	-1248	-77	-1278	216	-1399	-584	-39	-1418	299	-257	1029	-265	-964	2113	-1101	117
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
111(V)	-1349	-988	-3690	-3306	-1278	-3360	-2895	1671	-3086	-226	-141	-3047	-3312	-2925	-3136	-2638	-1356	3125	-2597	-2168	118
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
112(R)	-2176	-2415	-2504	-1923	-3054	-2403	-1269	-3053	-167	-2856	-2350	-1844	-2662	-997	3887	-2228	-2131	-2807	-2501	-2452	119
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
113(R)	1106	-1210	-925	-462	-2109	-352	-466	-1735	167	-1836	-1045	-526	-1629	-112	2713	-468	-502	-1309	-2063	-1625	120
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
114(Y)	-1833	-1522	-3003	-2757	2546	-2889	-152	-1096	-2433	-987	-712	-1944	-2891	-1848	-2268	-2038	-243	-1114	464	3769	121
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
115(A)	2880	-913	-1413	-1186	-2083	-1183	-1104	-1652	-747	-1929	-1234	-982	-1727	-890	214	-497	-570	-1201	-2274	-1896	122
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
116(I)	-818	1001	-3111	-2515	678	-2491	-1402	1758	-2173	1686	640	-2108	-2446	-1766	-2010	-1604	-758	1207	-1150	-857	123
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
117(R)	615	-1244	-985	-490	-2134	-1326	-441	-1752	263	-1841	-1057	-543	-1650	-82	2871	238	-530	-1334	-2053	-1625	124
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
118(K)	-1518	-2307	-1691	-846	-2975	-2112	-281	-2466	3232	-2235	-1545	-856	-2148	126	1301	-1423	-1307	-2203	-2146	-1935	125
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

TABLE 1

119(V)	334	-503	-1661	-1221	-1131	-1220	-917	-428	-1040	-959	-283	-984	-1644	-868	-1235	814	1722	1832	-1545	-1167	121
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
120(I)	-1801	-1400	-4009	-3498	-409	-3715	-2658	2299	-3194	2239	733	-3319	-3361	-2625	-3020	-2951	-1730	515	-1907	-1779	127
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
121(R)	-590	-1886	-363	992	-2256	-1091	1110	-1926	695	-1849	-997	-103	-1551	997	2392	-475	499	-1544	-1970	-1419	128
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
122(N)	-525	-1683	10	202	-2130	-1278	-228	-1842	152	-1887	-1070	2687	-1544	1246	-266	-453	843	-1452	-2112	-1492	129
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
123(M)	-1469	-1138	-3630	-3124	-557	-3307	-2344	2828	-2803	593	2857	-2896	-3124	-2415	-2714	-2510	-1422	784	-1868	-1641	130
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
124(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	131
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
125(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	132
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
126(A)	2657	-677	-1768	-1791	-2628	1691	-1726	-2289	-1863	-2588	-1770	-1201	-1631	-1609	-1973	-280	-452	-1476	-2836	-2569	133
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
127(A)	2560	-636	-2232	-1987	-1562	-1335	-1616	-72	-1780	-1097	-650	-1463	-1894	-1599	-1867	-627	-539	1482	-2167	-1804	134
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
128(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	135
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
129(L)	-992	-910	-2475	-1914	-111	-2437	615	229	-1462	2357	1583	-1768	-2392	-1292	-1486	-1534	-923	-34	-1023	-591	136
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
130(G)	-302	-865	-1762	-1813	-2349	3145	-1733	-1847	-1826	-2301	-1669	-1332	-1811	-1666	-1921	-535	-673	-35	-2637	-2293	137
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
131(A)	3260	-995	-2154	-2261	-2604	-1267	-2060	-2035	-2233	-2502	-1934	-1631	-1961	-2058	-2240	-724	-871	-1499	-2817	-2619	138
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
132(K)	-495	-1865	-257	1564	-2231	-1381	33	-1927	1891	-1845	-968	-21	-1480	1087	678	-101	-414	-1517	-1970	-1379	139
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*											
133(T)	15	-567	-2088	-1580	-827	-1880	-1130	1432	-1354	83	-1415	-2081	-1179	-1499	-1015	2384	1019	-1446	-1065	140	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
134(Y)	-1352	-1117	-2812	-2425	2395	-2595	-217	-654	-2102	398	-289	-1774	-2598	-1607	-1976	-504	-1284	-635	320	3348	141
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	118	-369	-294	-249	
-	-86	-4238	-7714	-565	-1626	-701	-1378*	*													
1135(V)	-879	1157	-3282	-2943	-1304	-2350	-2295	1015	-2647	-535	-296	-2399	-2667	-2450	-2627	-1648	-1002	3123	-2267	-1857	143
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
136(M)	1321	-485	-2344	-1767	-356	-2101	-1016	611	-1491	715	-2475	-1554	-2136	-1223	-1516	-1184	-533	1286	-1047	-714	144
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
137(W)	-3232	-2594	-3508	-3648	-882	-2917	-1886	-3060	-3237	-2665	-2615	-3229	-3282	-3161	-2996	-3457	-3341	-3103	6081	-495	145
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
138(G)	-629	-1215	-1479	-1608	-2971	3100	-1817	-2847	-1898	-3004	-2246	-1366	1410	-1693	-2073	-816	-993	-2058	-2924	-2780	146
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
139(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	147
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
140(R)	-2176	-2415	-2504	-1923	-3054	-2403	-1269	-3053	-167	-2856	-2350	-1844	-2662	-997	3887	-2228	-2131	-2807	-2501	-2452	148
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
141(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	149
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
142(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865											

TABLE 1

146(D)	-1105	-2793	2411	1424	-3056	1647	-501	-2904	-567	-2827	-2074	116	-1726	-144	-1292	-838	-1143	-2419	-3024	-2136	154
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
147(A)	2023	-621	-1648	-1495	-2508	1640	-1463	-2219	-1491	-2440	-1580	-1023	-1533	-1271	-1696	1504	-328	-1400	-2709	-2377	155
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
148(A)	2499	-588	-1757	-1515	-2231	-893	-1384	-1850	-1406	-2137	-1324	-1039	-1540	-1216	-1597	1059	1273	-1189	-2483	-2146	156
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
149(K)	-1004	-1906	-751	-464	-2680	668	-524	-2356	3024	-2305	-1553	-618	-1920	-150	132	-968	-1013	-1956	-2316	-1940	157
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
150(D)	-1145	-2665	3157	357	-2959	-1353	-536	-2796	1099	-2743	-2006	-4	-1786	-182	-1024	-906	-1172	-2352	-2877	-2097	158
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
151(V)	556	-703	-2845	-2356	-868	-2476	-1724	1032	-2090	672	119	-2092	-2570	-1874	-2129	-1644	-865	2613	-1729	-1361	159
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
152(R)	307	-1186	-699	-235	-2036	-283	-362	-1698	145	-1783	-956	-359	-1520	12	2596	677	-375	-1252	-2024	-1524	160
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
153(D)	966	-1545	1122	387	-1839	-201	57	-1545	1069	-1575	-700	91	-1343	486	-66	-11	-210	-337	-1814	-1172	161
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
154(A)	2670	-635	-2197	-1970	-1633	-1272	-1616	-227	-1767	-1190	-730	-1428	-1852	-1587	-1854	-571	-520	1219	-2215	-1854	162
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
155(W)	-1671	-1400	-3325	-2868	233	-3039	-1268	0	-2363	2540	532	-2472	-2926	-1994	-2227	-2277	-1599	-418	2890	45	163
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
156(D)	-1386	-3150	3360	1221	-3366	-1320	-681	-3279	-911	-3191	-2536	63	-1851	-357	-1729	-1063	-1456	-2785	-3348	-2403	164
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
157(R)	-1492	-1910	-2040	-1238	-1130	-2207	-499	-1858	364	-1773	-1220	-1192	-2291	-345	3286	-1537	-1364	-1712	2966	-659	165
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
158(M)	-671	-553	-2451	-1885	-21	-2161	-868	387	-1554	249	3340	-1603	-2178	-1263	-1528	-1248	-615	1153	-731	1912	166
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
159(R)	-1586	-2370	-1977	-912	-3100	-2189	-223	-2513	2555	-2235	-1541	-885	-2170	194	2735	-1476	-1329	-2260	-2127	-1954	167
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*												
160(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	168
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
161(A)	2667	-659	-1524	-1392	-2402	874	-1398	-2082	-1414	-2320	-1489	-994	377	-1210	-1630	-211	-359	-1347	-2615	-2271	169
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
162(F)	-1311	-1025	-3449	-2891	2343	-2994	-1698	606	-2578	1623	733	-2548	-2836	-2068	-2410	-2134	-1242	1636	-1220	-793	170
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
163(D)	-1226	-2864	2797	438	-3143	-1293	-620	-3082	-772	-3012	-2309	2528	-1798	-288	-1527	-951	-1296	-2581	-3152	-2243	171
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
164(L)	-982	-834	-2810	-2255	1392	-2500	-1292	389	-1949	2125	736	-1995	-2463	-1577	-1892	-1612	726	161	-1047	-640	172
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
165(M)	438	-905	-2819	-2258	-264	-2565	-1490	453	-1943	2160	2219	-2061	-2520	-1607	-1918	-1683	-989	202	-1306	-1076	173
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
166(G)	2232	-685	-1742	-1778	-2655	2330	-1733	-2323	-1870	-2618	-1798	-1198	-1634	-1614	-1983	-284	-460	-1498	-2857	-2591	174
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
167(E)	823	-1941	898	1743	-2233	-330	-41	-1990	272	-1949	-1063	145	-1423	1246	-270	-312	-417	-1552	-2138	-1425	175
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
168(Y)	-2742	-2290	-3014	-3139	701	-3008	-615	-2258	-2836	-1879	-1835	-2399	-3220	-2421	-2640	-2733	-2781	-2314	18	4561	176
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
169(V)	631	-431	-2194	-1660	-618	-1885	-1021	1372	-1413	-302	239	-1431	-2038	-1190	-1486	234	-449	2166	-1192	-819	177
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
170(T)	-500	-1770	-32	1487	-2088	-1320	-71	-1768	1131	-1785	-944	-2	-1493	344	-18	-397	1857	-1398	-1998	-1382	178
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
171(D)	437	-2189	2507	985	-2514	-1237	-262	-2284	-99	-2259	-1424	112	-1561	134	-696	605	-707	-1841	-2470	-1711	179
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
172(Q)	-1116	-2167	-1119	-394	-2714	-1859	-124	-2252	1946	-2066	-1298	-521	-1894	2940	1231	-1001	-946	-1936	-2060	-1719	180
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

TABLE 1

173(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	181
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
174(Y)	-2742	-2290	-3014	-3139	701	-3008	-615	-2258	-2836	-1879	-1835	-2399	-3220	-2421	-2640	-2733	-2781	-2314	18	4361	182
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
175(D)	-827	-2247	2476	406	-2712	1546	-431	-2510	-378	-2495	-1697	868	-1641	-63	-1011	-654	-878	-1542	-2708	-1921	183
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
176(L)	-604	-460	-2538	-1944	-130	-2143	-956	1316	-1614	1768	679	-1640	-2146	-1291	-1551	-1226	89	434	-843	1377	184
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
177(R)	-1151	-2083	-1284	-523	-2658	-1893	-175	-2181	1390	-2032	-1286	-614	244	239	3031	-1068	-991	-1886	-2044	-1735	185
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
178(F)	-1690	-1368	-3472	-3107	3473	-3159	-1180	1363	-2783	394	347	-2559	-3051	-2224	-2601	-2369	-1641	22	-559	349	186
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
179(A)	3260	-995	-2154	-2261	-2604	-1267	-2060	-2035	-2233	-2502	-1934	-1631	-1961	-2058	-2240	-724	-871	-1499	-2817	-2619	187
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
180(I)	-1681	-1285	-3919	-3432	-552	-3632	-2662	2824	-3136	1690	578	-3234	-3343	-2865	-3012	-2872	-1625	767	-2003	-1807	188
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
181(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	189
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
182(P)	-2089	-2190	-2567	-2787	-3372	-2253	-2663	-3607	-2895	-3590	-3215	-2612	4122	-2846	-2839	-2316	-2418	-3135	-2975	-3234	190
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
183(K)	-1892	-2397	-1701	-1300	-3050	-2216	-948	-2829	3583	-2696	-2116	-1348	-2450	-619	78	-1865	-1815	-2562	-2460	-2321	191
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
184(P)	-2089	-2190	-2567	-2787	-3372	-2253	-2663	-3607	-2895	-3590	-3215	-2612	4122	-2846	-2839	-2316	-2418	-3135	-2975	-3234	192
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
185(N)	-1468	-2089	-762	-997	-2634	-1768	-1516	-3059	-1508	-3127	-2580	4001	-2312	-1391	-1789	-1503	-1700	-2572	-2658	-2191	193
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
186(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	194
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

-	-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
187(P)	-2089	-2190	-2567	-2787	-2787	-2787	-3372	-2253	-2663	-2895	-3607	-2895	-3590	-3215	-2612	4122	-2846	-2839	-2316	-2418	-3135	-2975	-3234	195
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	106	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
188(R)	-2176	-2415	-2504	-1923	-3054	-1923	-3054	-2403	-1269	-3053	-167	-2856	-2350	-1844	-2662	3887	997	96	359	117	-2807	-2501	-2452	196
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
189(G)	-1815	-2026	-2479	-2749	-3578	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	197	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
190(D)	-2059	-2867	3860	-532	-3452	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	198	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
191(I)	-1608	-1307	-3432	-3238	-1179	-3238	-1179	-3095	-2687	3474	-2941	-191	-206	-2987	-3216	-2818	-2915	-2635	-1653	832	-2363	-1952	199	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
192(Y)	-1651	-1320	-3317	-2937	1876	-2937	1876	-2994	-612	-279	-2586	2172	216	-2231	-2886	-1963	-2359	-2131	-1568	-559	4	2207	200	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
193(L)	-2071	-1671	-3975	-3513	1646	-3975	1646	-3704	-1951	276	-3199	2643	835	-3187	-3331	-2460	-2935	-2941	-1974	-311	-1161	-569	201	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
194(P)	1158	-816	-1549	-1568	-2492	-1549	-2492	-1042	-1594	-2082	-1610	-2398	-1676	-1163	3346	-1455	-1757	-417	-570	-1432	-2710	-2399	202	
-	-149	-500	233	43	-381	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-894	-1115	-701	-1378*	*														
195(T)	-665	-1148	-2040	-2098	-2442	-2040	-24																	

TABLE 1

200(L)	-1959	-1558	-4063	-3546	-291	-3808	-2666	1414	-3212	2654	850	-3406	-3399	-2589	-3010	-3056	-1876	210	-1837	-1742	208
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
201(A)	3260	-995	-2154	-2261	-2604	-1267	-2060	-2035	-2233	-2502	-1934	-1631	-1961	-2058	-2240	-724	-871	-1499	-2817	-2619	209
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
202(F)	-2151	-1753	-3530	-3327	3739	-3318	-783	-347	-2962	686	53	-2548	-3220	-2300	-2714	-2589	-2099	-722	-95	953	210
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
203(I)	-1413	-1022	-3801	-3407	-1219	-3492	-2985	3184	-3199	-111	-62	-3161	-3383	-2998	-3231	-2772	-1408	1813	-2596	-2206	211
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
204(E)	90	-1897	443	2010	-2185	-1230	1725	-1936	326	-1898	-1010	138	-1411	1242	-197	-292	-385	-1505	-2086	-1385	212
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
205(R)	-351	-1687	-179	891	-1996	-826	66	-1700	567	-1680	-800	50	-1394	1328	1582	-249	1128	-1307	-1867	-1245	213
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
206(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	214
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
207(E)	-1261	-2925	1287	3091	-3172	-1326	-611	-3046	-721	-2978	-2273	46	-1812	-271	-1450	-976	-1310	-2573	-3131	-2258	215
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
208(R)	-1403	-2166	-1551	-797	-2257	-2051	2088	-2287	844	-2112	-1431	-830	-2113	68	3221	-1333	-1230	-2033	-1908	-1480	216
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
209(P)	-721	-1287	-1519	-1649	-2984	1028	-1849	-2874	-1919	-3027	-2289	-1426	3581	-1736	-2085	-907	-1079	-2116	-2912	-2790	217
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
210(E)	-1251	-2967	1669	2974	-3197	-1308	-591	-3075	-714	-2993	-2283	76	-1796	-249	-1463	-958	-1299	-2592	-3159	-2259	218
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
211(L)	-904	-844	-2403	-1784	-342	-2365	-1121	1250	-1186	2028	628	-1672	-2331	-1162	772	-1456	-834	204	-1214	-915	219
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
212(Y)	-1403	-1120	-3032	-2663	2477	-2894	-320	-428	-2317	-519	-196	-1924	-2681	-1767	-2127	-1818	-1336	1473	248	3071	220
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
213(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	221
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
214(V)	-1401	-1014	-3772	-3281	-3281	-840	-3424	-2588	1734	-3048	1458	291	-3027	-3233	-2685	-2984	-2633	-1359	2492	-2135	-1864	222
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
215(N)	-1468	-2069	-762	-997	-2634	-1768	-1516	-3059	-1508	-3127	-2580	4001	-2312	-1391	-1789	-1503	-1700	-2572	-2658	-2191	223	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
216(P)	-1085	-1645	-1400	-1250	-2540	-1703	-1221	-2412	-640	-2466	-1828	-1243	3622	-987	539	-1201	-1262	-1999	-2445	-2166	224	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
217(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	225	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
218(V)	-565	-483	-2432	-1897	1185	-2061	-1097	692	-1621	-87	343	-1617	-2167	-1361	-1634	-1181	1369	2185	-1112	-695	226	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
219(G)	1216	-738	-1673	-1781	-2746	2966	-1793	-2425	-1950	-2725	-1915	-1223	-1673	-1686	-2050	-337	-522	-1578	-2929	-2677	227	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
220(H)	-2323	-2419	-1868	-1956	-1368	-2396	5124	-3143	-1645	-2924	-2595	-1989	-2806	-1873	-1712	-2381	-2439	-2923	-1719	-935	228	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
221(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	229	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
222(Q)	-600	-1482	-475	-254	-2078	-1374	-473	-1739	63	-1831	-1129	-420	-1705	3290	-202	-619	958	-1394	-2115	-1591	230	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
223(M)	-1736	-1678	-2975	-2821	-903	-2619	-2137	-342	-2336	24	4777	-2581	-2886	-2281	-2280	-2241	-1843	-654	-1928	-1574	231	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
224(A)	2907	-600	-1848	-1825	-2494	-873	-1680	-2122	-1800	-2451	-1642	-1177	-1583	-1569	-1896	972	-376	-1349	-2754	-2449	232	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
225(G)	-596	-1353	-455	-651	-2765	2967	-1238	-2694	-1243	-2817	-2057	1271	-1821	-1009	-1597	-689	-896	-1982	-2836	-2358	233	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
226(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	234	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														

TABLE 1

222(N)	-1468	-2069	-762	-997	-2634	-1768	-1516	-3059	-1508	-3127	-2580	4001	-2312	-1391	-1789	-1503	-1700	-2572	-2658	-2191	235
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
228(F)	-2605	-2164	-3410	-3518	4186	-3034	-940	-1504	-3300	-1072	-1142	-2698	-3246	-2687	-2999	-2815	-2670	-1743	-274	806	236
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
229(T)	-156	-631	-1441	-1038	-1327	-1213	-890	-577	-867	-1099	-449	-889	1925	-751	-1111	-410	1968	948	-1720	-1318	237
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
230(H)	299	-1594	-259	23	-1760	-1411	3693	-1605	285	-1670	-904	-212	-1624	1240	-25	-563	-577	-1296	-1828	-1224	238
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
231(G)	1216	-738	-1673	-1781	-2746	2966	-1793	-2425	-1950	-2725	-1915	-1223	-1673	-1686	-2050	-337	-522	-1578	-2929	-2677	239
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
232(I)	-1412	-1014	-3818	-3423	-1244	-3514	-3011	3143	-3223	-139	-82	-3177	-3397	-3025	-3258	-2792	-1405	1920	-2622	-2228	240
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
233(A)	2643	-702	-1841	-1506	-1222	-1365	-1211	-382	-1249	-749	1643	-1220	-1832	-1146	-1400	-615	-514	-252	-1806	-1421	241
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
234(Q)	-1762	-2287	-1132	-1144	-2520	-2040	-1289	-2773	-740	-2625	-2183	-1302	-2433	4141	-881	-1765	-1840	-2519	-2457	-2076	242
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
235(A)	3260	-995	-2154	-2261	-2604	-1267	-2060	-2035	-2233	-2502	-1934	-1631	-1961	-2058	-2240	-724	-871	-1499	-2817	-2619	243
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
236(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	244
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
237(W)	-1283	-1528	932	-1130	736	-2224	-138	-1253	-1094	-1220	-765	-1084	-2321	-868	-1347	-1344	-1228	-1158	4771	1985	245
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
238(A)	2026	-1366	-313	34	-1635	-1335	1574	-1366	204	-1485	-707	-193	-1544	1074	-155	-433	-425	-1066	-1749	-1178	246
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
239(G)	-831	-2052	172	754	-2493	2361	-347	-2192	1024	-2191	-1409	-103	-1691	30	-296	-704	-829	-1804	-2375	-1762	247
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
240(K)	-1892	-2397	-1701	-1300	-3050	-2216	-948	-2829	3583	-2696	-2116	-1348	-2450	-619	78	-1865	-1815	-2562	-2460	-2321	248
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*												
241(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	249
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
242(F)	-2151	-1753	-3530	-3327	3759	-3318	-783	-347	-2962	686	53	-2548	-3220	-2300	-2714	-2589	-2099	-722	-95	953	250
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
243(H)	-2323	-2419	-1868	-1956	-1368	-2396	5124	-3143	-1645	-2924	-2595	-1989	-2806	-1873	-1712	-2381	-2439	-2923	-1719	-935	251
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
244(I)	-1608	-1307	-3432	-3238	-1179	-3095	-2687	3474	-2941	-191	-206	-2987	-3216	-2818	-2915	-2635	-1653	832	-2363	-1952	252
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
245(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	253
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
246(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	254
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
247(N)	-1468	-2069	-762	-997	-2634	-1768	-1516	-3059	-1508	-3127	-2580	4001	-2312	-1391	-1789	-1503	-1700	-2572	-2658	-2191	255
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
248(G)	-1138	-1994	970	-393	-3114	3135	-1239	-3051	-1406	-3111	-2421	-642	-2033	-992	-1944	-1114	-1363	-2451	-3025	-2576	256
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
249(Q)	-1762	-2287	-1132	-1144	-2520	-2040	-1289	-2773	-740	-2625	-2183	-1302	-2433	4141	-881	-1765	-1840	-2519	-2457	-2076	257
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
250(R)	-494	-1670	-394	131	-1997	-1407	1541	-1693	597	-1689	-853	228	-1520	390	1988	1217	-426	-1339	-1858	-1299	258
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
251(G)	-454	-932	-1545	-1368	-1592	2791	-1240	-1146	-1201	-1367	2191	-1171	-1868	-1149	-1371	-692	-687	-902	-2003	-1589	259
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
252(I)	-252	-621	-1011	-601	-831	-1403	-480	2266	-483	-684	1	362	344	-369	-758	147	-276	-145	-1217	-790	260
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
253(K)	-1539	-2341	-1845	-864	-3043	-2150	-233	-2484	3078	-2225	-1527	-857	-2149	183	1839	-1433	-1300	-2224	-2127	-1935	261
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

TABLE 1

254(Y)	-2984	-2211	-3668	-3802	2833	-3558	-44	-1957	-3405	-1449	-1443	-2318	-3461	-2386	-2945	-2790	-2877	-2089	685	4009	262
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
255(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	263
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
256(Q)	-1762	-2287	-1132	-1144	-2520	-2040	-1289	-2773	-740	-2625	-2183	-1302	-2433	4141	-881	-1765	-1840	-2519	-2457	-2076	264
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
257(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	265
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
258(L)	-1238	1377	-3271	-2822	-453	-2615	-1907	366	-2410	2625	519	-2451	-2721	-2079	-2306	-1895	-1275	123	-1577	-1279	266
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
259(R)	-435	-969	-1119	-575	-1319	-375	-421	-736	74	-1090	-402	-632	-1698	-167	2179	-601	-453	1627	-1543	-1122	267
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
260(F)	-1141	-1255	-2377	-2279	3772	-2092	-569	-1041	-2100	-1019	-743	-1705	-2446	-1735	-2059	-76	-1293	-987	-121	942	268
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
261(G)	-920	-1508	-1339	-1295	-2751	3192	-1377	-2630	-937	-2716	-2012	-1241	-2085	-1160	523	-1056	-1170	-2071	-2626	-2390	269
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
262(H)	910	-1328	-303	212	-1640	-1258	2619	-1308	356	-1405	-574	-44	-733	368	434	599	-227	-980	-1697	-1118	270
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-102	-6672	-4094	-894	-1115	-701	-1378*	*													
263(G)	-998	-2008	156	1372	-2814	2761	-820	-2621	-763	-2678	-1963	-325	-1846	-517	-1251	-919	-1134	-2140	-2758	-2201	271
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-22	-6592	-7635	-894	-1115	-618	-1521*	*													
264(D)	-896	-2439	2885	448	-2718	-1275	-358	-2532	-225	-2479	-1678	1424	-1648	21	249	-685	-902	-2082	-2659	-1868	272
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
265(L)	-1226	-1270	-2434	-2156	-630	-2346	-1635	-84	-1796	2639	285	-2000	-48	-1683	-1827	-1692	-1284	-360	-1630	-1298	273
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
266(R)	261	-1516	-309	264	-1768	-1323	57	-1437	1161	-420	-634	-7	-1406	1111	361	-261	663	-1102	-1724	-1141	274
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
267(A)	1860	-968	-672	-395	-2180	-32	-670	-1885	-459	-2014	-1164	1678	-1481	-332	-871	1019	-328	-1306	-2284	-1769	275
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
268(A)	3048	-694	-1839	-1891	-1891	-2627	246	-1791	-2258	-1948	-2591	-1793	-1261	-1663	-1699	-2026	-319	-489	-1473	-2840	-2589	276
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
269(F)	-1633	-1375	-3311	-2975	3610	-2991	-973	88	-2640	335	217	-2384	-2965	-2115	-2486	-2194	-1612	560	-373	594	277	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
270(W)	-649	-554	-2266	-1731	1870	-2042	-407	-45	-1432	578	294	-1386	-2086	-1111	-1417	831	-590	6	3399	661	278	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
271(L)	-954	-821	-2749	-2189	-379	-2508	-1467	615	-1885	2189	848	-1995	-2489	-1589	-1889	-1624	721	884	-1355	-1090	279	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
272(V)	-1294	-1167	-3176	-3053	-1510	-2597	-2599	776	-2808	-665	-568	-2665	-2928	-2721	-2804	-2093	-1429	3343	-2527	-2099	280	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
273(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	281	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
274(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	282	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
275(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	283	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
276(E)	-1877	-2708	-271	3539	-3235	-1868	-1322	-3210	-1299	-3203	-2676	-849	-2342	-1076	-1736	-1723	-1961	-2857	-2976	-2640	284	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
277(N)	-343	-1453	-252	132	-2019	-1231	-165	-1718	264	-1761	-914	2012	-1464	238	635	1541	-137	-1302	-1987	-1397	285	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*														
278(A)	1989	-615	-1671	-1548	-2532	1980	-1508	-2242	-1558	-2469	-1610	-1047	-1538	-1327	-1748	1152	-334	-1411	-2738	-2415	286	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-1061	-6672	-969	-894	-1115	-701	-1378*	*														
279(F)	-62	-382	-890	-612	2072	848	-159	-157	-517	-365	131	-522	-1382	-366	-722	-310	-160	-26	-381	402	287	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-43	-5654	-6897	-894	-1115	-1298	-753*	*														
280(P)	53	-411	-960	-673	-945	-944	-591	-73	-495	-605	-126	-572	2133	-444	-723	-198	-123	1167	-1421	-973	288	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249		
-	-36	-5916	-6958	-894	-1115	-1541	-607*	*														

TABLE 1

281(N)	-403	-1803	1732	803	-2196	600	10	-1981	99	-1980	-1180	1760	-1229	381	-510	-235	-435	-1535	-2198	-1435	289
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-36	-5916	-6958	-894	-1115	-1541	-607*		*												
282(G)	-564	-931	-1093	-1247	-2240	3219	-1339	-2241	-1464	-2403	-1833	-1104	-1599	-1344	-1556	-759	-890	-1685	-2020	-2051	290
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-36	-5916	-6958	-894	-1115	-1541	-607*		*												
283(G)	1043	-280	-688	-700	-1858	2341	-847	-1455	-840	-1779	-1026	-438	-1116	-671	-1053	164	18	-830	-2063	-1703	291
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-36	-5916	-6958	-894	-1115	-1541	-607*		*												
284(P)	-290	-1038	-466	-90	-1581	-1092	-104	-1283	527	-1372	-663	-208	2166	220	1475	-341	-334	-968	-1564	-1136	292
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-36	-5916	-6958	-894	-1115	-349	-2218*		*												
285(G)	638	-1161	-471	-124	-2022	1878	-396	-1704	1009	-1808	-971	-280	-1477	-27	-463	540	-339	-1238	-2076	-1532	293
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
286(Y)	-2887	-2205	-3496	-3544	1333	-3432	-88	-2037	-3047	-1553	-1514	-2272	-3386	-2288	-2715	-2719	-2795	-2134	1494	4420	294
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
287(D)	192	-1858	1553	1374	-2155	-1216	15	-1908	361	-1866	-969	151	-1383	1382	-169	-23	260	-1470	-2052	-1350	295
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
288(G)	-1815	-2026	-2479	-2749	-3578	3649	-2701	-3848	-3053	-3865	-3355	-2493	-2700	-2875	-2985	-2041	-2188	-3149	-3082	-3440	296
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
289(P)	-2089	-2190	-2567	-2787	-3372	-2253	-2663	-3607	-2895	-3590	-3215	-2612	4122	-2846	-2839	-2316	-2418	-3135	-2975	-3234	297
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
290(R)	-1641	-2275	-1933	-1088	-2932	-2175	-424	-2500	1015	-2295	-1649	-1042	-2256	-38	3525	-1586	-1455	-2254	-2181	-1996	298
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
291(H)	-2323	-2419	-1868	-1956	-1368	-2396	5124	-3143	-1645	-2924	-2595	-1989	-2806	-1873	-1712	-2381	-2439	-2923	-1719	-935	299
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
292(F)	-2605	-2164	-3410	-3518	4136	-3034	-940	-1504	-3300	-1072	-1142	-2698	-3246	-2697	-2999	-2815	-2670	-1743	-274	806	300
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
293(D)	-2059	-2867	3860	-532	-3452	-1877	-1516	-3665	-1847	-3616	-3145	-882	-2414	-1318	-2469	-1872	-2198	-3236	-3167	-2847	301
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*		*												
294(Y)	579	-1160	-2665	-2299	2734	-2531	-229	-710	-2012	-758	-361	-1718	-2573	-1556	-1937	-1652	-1288	-684	285	3136	302
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

[illegible]

TABLE 1

308(S)	-447	-881	-1613	-1364	1517	-1419	-806	-1058	-1261	-1223	-705	-1119	-1875	-1080	-1446	2639	-648	-827	-983	-113	316
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
309(A)	3260	-995	-2154	-2261	-2604	-1267	-2060	-2035	-2233	-2502	-1934	-1631	-1961	-2058	-2240	-724	-871	-1499	-2817	-2619	317
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
310(K)	1246	-1752	-354	687	-2095	-1400	24	-1767	1628	-1730	-874	-63	-1490	459	1177	-382	-400	-1394	-1891	-1325	318
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
311(G)	1279	-1129	455	-229	-2280	2077	-665	-1991	-519	-2113	-1277	-355	-1522	-330	-976	392	-436	-1424	-2385	-1830	319
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
312(C)	-342	4308	-1550	-1431	-1885	-1211	-1326	-1596	-1300	-1974	-1267	2515	-1793	-1235	-1463	-561	-633	-1160	-2187	-1732	320
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
313(M)	-1538	-1273	-3473	-2967	-378	-3209	-2177	1411	-2583	795	4160	-2776	-3045	-2211	-2502	-2413	-1492	381	-1708	-1493	321
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
314(R)	-802	-1573	-1236	-713	-2404	-1580	-468	-2096	486	-2079	-1343	-728	-1867	-105	3143	691	-850	-1698	-2130	-1757	322
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
315(M)	-291	-866	-751	-323	-1102	-1363	-365	-625	-175	-903	2766	1884	-1580	-129	-510	-445	882	-433	-1424	-950	323
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
316(Y)	-2742	-2290	-3014	-3139	701	-3008	-615	-2258	-2836	-1879	-1835	-2399	-3220	-2421	-2640	-2733	-2781	-2314	18	4561	324
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
317(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	325
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
318(I)	-1718	-1320	-3950	-3454	-500	-3658	-2657	2675	-3157	1891	635	-3261	-3346	-2650	-3014	-2895	-1657	685	-1967	-1796	326
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
319(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	327
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
320(K)	-1892	-2397	-1701	-1300	-3050	-2216	-948	-2829	3583	-2696	-2116	-1348	-2450	-619	78	-1865	-1815	-2562	-2460	-2321	328
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
321(E)	-1027	-2709	1389	2805	-2938	-1270	-422	-2771	-394	-2690	-1910	126	-1688	847	-1061	-772	-1042	-2301	-2880	-2022	329
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

-	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*												
322(R)	-1262	-1929	-1576	-1005	-2676	-1901	-641	-2373	404	-2286	-1610	-1018	104	-286	3467	-1282	-2043	-2243	-1991	330	
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
323(A)	2703	-817	-1288	-964	-1935	-1111	-906	-1465	-558	-1746	-1009	-813	-1616	-662	368	-363	-64	-1031	-2134	-1737	331
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
324(Q)	921	-1531	-150	983	-1783	-1243	90	-1481	984	-362	-643	89	-373	1104	16	-170	-191	-1116	-1755	-1125	332
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
325(A)	3002	-721	-1722	-1692	-2428	-988	-1617	-2000	-1677	-2335	-1590	-1188	277	-1495	-1805	-343	-489	-1343	-2672	-2372	333
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
326(F)	-1141	-1255	-2377	-2279	3772	-2092	-569	-1041	-2100	-1019	-743	-1705	-2446	-1735	-2059	-76	-1293	-987	-121	942	334
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
327(R)	26	-1636	-1533	-952	-2483	-1704	-566	-2043	472	-2068	-1404	-909	-1989	-212	3357	-1021	-1002	-1707	-2177	-1875	335
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
328(A)	2206	-1366	-122	1071	-2009	-1198	-396	-1654	-158	-1799	-1003	-190	-81	-44	-610	-402	-477	-1263	-2110	-1531	336
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
329(D)	-1040	-2279	3369	219	-2914	-1294	-744	-2643	-860	-2773	-2097	-113	-1811	-431	-1522	-889	170	-2183	-2988	-2215	337
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
330(P)	-1078	-1646	-1388	-1222	-2530	-1702	-1185	-2393	-586	-2444	-1802	-1219	3599	-944	631	-1190	-1247	-1984	-2429	-2146	338
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
331(E)	-592	-2098	820	2419	-2389	-1258	-116	-2150	202	-2093	-1225	119	-1495	1056	383	-427	-554	-1711	-2269	-1550	339
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
332(V)	-864	1397	-3270	-2925	-1287	-2333	-2263	1008	-2628	-532	-284	-2379	-2649	-2425	-2605	-1626	-986	3100	-2235	-1829	340
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
333(Q)	-1116	-2159	-1088	-390	-2695	-1853	-132	-2237	1638	-2059	-1295	-519	-1896	3120	1253	-1004	-949	-1924	-2060	-1716	341
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
334(E)	77	-1884	74	2581	-2260	-1329	-186	-1946	248	-1959	-1140	-36	-1567	216	517	-508	-594	-1564	-2165	-1539	342
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													

TABLE 1

335(A)	2118	-778	-1188	-678	-992	-1464	-510	-448	-265	351	-128	-707	-1690	-333	310	-572	-378	-302	-1373	-946	343
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
336(L)	-273	-1147	-3152	-2607	-269	-2901	-1837	473	-2264	2501	1696	-2426	-2787	-1885	-2206	-2051	-1280	127	-1497	-1323	344
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
337(A)	1540	-1532	-474	87	-1918	-1382	-46	-1567	1215	-1601	-772	-145	-1504	365	1342	125	-380	-1223	-1818	-1283	345
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
338(A)	2335	-972	-614	294	-1776	-1148	-611	-1301	-417	-1587	-829	-465	-1555	-323	-786	-354	962	937	-1990	-1503	346
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
339(S)	1294	-832	-756	-278	-1210	-1235	-345	-856	-177	-1085	-331	-396	-201	-85	-555	1388	-227	-590	-1486	1213	347
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
340(R)	-570	-1686	-582	34	-1977	144	-15	-1637	1580	-1635	-817	-192	-1574	399	1931	-500	-474	-1317	-1803	1086	348
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	46	96	359	117	-369	-294	-249	
-	-307	-2420	-7714	-266	-2570	-701	-1378*	*													
341(V)	-336	-353	-1746	340	-254	-1760	-562	286	-906	1013	531	-1061	-1812	-710	-1015	-805	-277	1674	1260	-385	350
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
342(D)	352	-1357	1943	274	527	-1272	-34	-1207	263	-1335	-525	-5	-1411	741	-215	-270	262	-916	-1662	-1069	351
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
343(E)	-521	-1935	227	2230	-2246	-1250	-100	-1987	223	-1965	-1102	97	607	1057	-287	155	-487	-1568	-2166	-1473	352
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
344(L)	546	-413	-2480	-1879	-125	-2068	-906	1031	-1548	1827	714	-1574	-2081	-1226	-1484	-1149	-484	382	1382	-472	353
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
345(A)	1293	-1490	227	381	-1734	4	86	-1428	735	-1476	-609	88	-377	512	391	-161	-178	-1072	-1730	-134	354
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
346(Q)	-254	-1503	-183	1017	-1748	-1253	86	-1438	495	-297	-617	73	-1345	1273	735	190	977	-1084	-1731	-1110	355
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
347(P)	421	-948	-865	-456	-1802	-1152	-548	-1416	-196	-1606	-823	-506	2712	-237	384	-316	452	-1015	-1949	-1485	356
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
348(T)	-276	-686	-1314	-885	-1018	-1380	-716	-410	-659	414	-163	-836	-135	-591	-897	-537	2548	-256	-1462	-1045	357
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

	-	-21	-6672	-7714	-894	-1115	-701	-1378*	*												
349(L)		1151	-788	-2503	-1952	-325	-2317	-1281	427	-1656	1950	-1784	-2340	-1385	-1690	-1427	-817	257	-1248	-972	358
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
350(N)		1068	-1985	832	927	-2287	217	-111	-2043	157	-2016	1836	-1465	308	-398	-378	-498	-1611	-2219	-1499	359
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
351(D)		515	-2116	2033	1040	-2419	-1237	-204	-2180	1	-2157	1836	1413	200	-577	-479	-630	-1746	-2368	-1626	360
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-214	-6672	-2965	-894	-1115	-701	-1378*	*												
352(G)		-1483	-1738	-2118	-2361	-3240	3582	-2356	-3437	-2647	-3497	-2968	-2132	-2414	-2483	-2624	-1701	-1847	-2772	-2817	-3088
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-602	-6482	-1599	-894	-1115	-1010	-990*	*												
353(E)		-844	-1778	472	2917	-2144	-1088	-364	-1914	-199	-2018	-1432	9	-1504	-76	-613	-726	-895	-1616	-2068	-1590
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-36	-5916	-6958	-894	-1115	-1541	-607*	*												
354(S)		409	-277	-676	-434	-1615	472	-531	-1263	-415	-1493	-687	-281	-1062	-284	-715	1604	1521	-699	-1824	-1402
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-36	-5916	-6958	-894	-1115	-349	-2218*	*												
355(Y)		176	-326	-1853	-1280	-211	-1746	-587	276	-1009	1143	525	-1118	-1820	-790	-1087	-806	334	460	-736	2298
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
356(Q)		1158	-1744	813	939	-2039	-1215	47	-1776	411	-1756	-861	138	-1359	1168	-105	600	-85	-1356	-1956	-1275
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
357(D)		1036	-1772	1595	823	-2060	-1224	1065	-1797	408	-1774	-882	135	-1370	477	479	-226	-297	-1379	-1971	-1290
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
358(L)		-831	-865	-1889	480	-453	-2235	-1046	1057	-1158	2022	523	-1421	-2241	-1015	-1320	-1316	-771	284	-1297	-968
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
359(L)		-837	-924	-1941	-1334	-448	-2177	-874	129	765	2074	1446	-1328	-2171	-793	-816	-1251	-762	-40	-1238	-923
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
360(A)		2412	-1304	264	-12	-2084	-1143	-485	-1768	-315	-1902	-1096	676	-568	-145	-788	-383	-481	-1328	-2203	-1621
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												
361(D)		-1360	-3141	3267	1479	-3350	-1311	-657	-3259	-874	-3165	-2499	78	-1836	-330	-1688	-1038	-1426	-2762	-3331	-2380
-		-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249
-		-21	-6672	-7714	-894	-1115	-701	-1378*	*												

TABLE 1

362(R)	1110	-1178	-914	-425	-2040	-1274	-431	-1667	185	-1776	-982	-495	-1601	-75	2624	266	-459	-1250	-2016	-1566	371
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
363(S)	660	-658	-1132	-653	-1190	-1207	-583	-715	-506	248	-310	-639	-1546	-394	-817	1790	985	-458	-1520	-1091	372
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
364(A)	2005	-1109	-599	-215	-1913	-1164	-412	-1560	1070	-1701	-888	-346	-1504	-60	-397	906	-348	-1138	-1994	-1480	373
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
365(F)	-1666	-1475	-2647	-2273	3207	-2733	-126	-1109	-1832	-1025	-680	-1722	-2730	-1513	985	-1846	-1589	-1094	401	2740	374
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
366(E)	1231	-2217	1090	2270	-2529	-1247	-283	-2286	-130	-2276	-1451	104	-1579	108	-726	-564	-741	-1853	-2493	-1734	375
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
367(D)	-807	-2381	2435	1701	-2662	-153	-299	-2456	-176	-2404	-1577	131	-1595	93	-801	-602	340	-1999	-2603	-1807	376
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
368(F)	-2122	-1662	-3439	-3251	3476	-3201	-250	-937	-2878	738	-472	-2219	-3110	-2117	-2574	-2384	-2033	-1127	427	2382	377
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
369(D)	-106	-2081	2987	302	-2719	-1222	-536	-2478	-516	-2518	-1742	-21	945	-185	-1136	-670	-895	-1991	-2748	-1995	378
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
370(V)	1420	-434	-1909	-1374	-548	-1743	-830	417	-1135	567	297	-1213	-880	-938	-1254	-846	-374	-1727	-1093	-720	379
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
371(D)	-1069	-2768	2618	2120	-3007	-1268	-462	-2850	-491	-2770	-2001	126	-238	-97	-1197	-805	-1095	-2370	-2962	-2084	380
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
372(A)	2636	-1018	-521	219	-2069	-1094	-793	-1676	-652	-1914	-1144	-512	-1591	-518	-1012	194	-479	-1214	-2262	-1772	381
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-248	
-	-149	-3423	-7714	-422	-1979	-701	-1378*	*													
373(A)	1582	-605	-1051	-501	555	-1512	-336	-123	725	-487	179	-589	-1619	-212	-602	-527	-227	695	-1054	-619	383
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
374(A)	1620	-1501	684	281	-1928	64	-79	-1628	270	-1677	-820	6	-1408	332	916	-249	-300	-1233	-1925	-1300	384
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
375(A)	1658	-1589	-193	665	-1851	-1305	1059	-1534	504	-1565	-713	20	-1412	946	607	-274	-289	-1184	-1797	-1195	385
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	

TABLE 1

389(M)	-586	-422	-2607	-2005	-236	-2162	-1033	1738	-1674	727	2647	-1686	-2164	-1351	-269	-1247	-526	1278	-937	-619	403
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
390(E)	-86	-2659	2289	2344	-2918	-1261	-428	-2735	-423	-2675	-1894	125	-1682	-59	-1107	-758	-1026	-2265	-2876	-2019	404
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
391(H)	-1063	-1596	-1226	-723	125	-1986	3902	-1400	-6	-1381	-808	-779	-2060	-275	584	-1076	-975	-1242	-421	1920	405
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
392(L)	-2228	-1908	-3555	-3373	-604	-3255	-2515	-62	-2963	2976	358	-3207	-3275	-2644	-2802	-2988	-2232	-567	-1904	-1643	406
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
393(L)	-1311	-1120	-3138	-2575	-246	-2891	-1784	463	-2223	2450	1961	-2395	-2763	-1847	-2162	-2028	-31	113	-1452	-1278	407
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
394(G)	-290	-847	-1769	-1799	-2291	3105	-1705	-1729	-1796	-2216	-1592	-1322	-1804	-1637	-1895	-526	-654	179	-2597	-2248	408
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
395(A)	2540	-539	-1889	-1592	-1657	-1048	-1298	-901	-1422	-1478	-795	-1130	-1629	-1231	-1567	446	-331	1000	-2050	-1697	409
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-21	-6672	-7714	-894	-1115	-701	-1378*	*													
396(R)	-1262	-1929	-1576	-1005	-2676	-1901	-641	-2373	404	-2286	-1610	-1018	104	-286	3467	-1282	-1246	-2043	-2243	-1991	410
-	-149	-500	233	43	-381	399	106	-626	210	-466	-720	275	394	45	96	359	117	-369	-294	-249	
-	-30	-6155	-7197	-894	-1115	-701	-1378*	*													
397(G)	21	-791	-381	-92	-1740	1778	-286	-1423	81	-1560	-757	-172	-1269	42	694	855	-134	-959	-1829	-1336	411
-	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	
-	*	*	*	*	*	*	*	*	0												

¹Program name and version²Name of the input sequence alignment file³Length of the alignment: include indels⁴Type of residues⁵Map of the match states to the columns of the alignment⁶Number of sequences in the alignment⁷When the file was generated⁸The transition probability distribution for the null model (single G state).⁹The symbol emission probability distribution for the null model (G rate); consists of K integers. The null probability used to convert these back to model probabilities is 1/K.

TABLE 2

Table 2. E-values and XI Group membership for proteins by SEQ ID NO.

Organism	SEQ ID NO: Amino acid	GI or AC #	XI Group	E-value
<i>Streptomyces ambofaciens</i>	16	126348424	I	0
<i>Streptomyces sp.</i>	20	38141596	I	0
<i>Streptomyces coelicolor</i>	32	Q9L0B8	I	0
<i>Streptomyces avermitilis</i>	34	Q93HF3	I	0
<i>Streptomyces violaceusniger</i>	68	P09033	I	0
<i>Streptomyces olivaceoviridis</i>	96	Q93RJ9	I	0
<i>Streptomyces corchorusii</i>	106	Q9S3Z4	I	0
<i>Streptomyces thermocyaneoviolaceus</i>	108	Q9L558	I	0
<i>Streptomyces diastaticus</i>	122	P50910	I	0
<i>Streptomyces rubiginosus</i>	128	P24300	I	0
<i>Streptomyces murinus</i>	130	P37031	I	0
<i>Streptomyces olivochromogenes</i>	135	P15587	I	0
<i>Streptomyces olivochromogenes</i>	136	157879319	I	0
<i>Streptomyces olivochromogenes</i>	138	157881044	I	0
<i>Streptomyces diastaticus</i>	139	7766813	I	0
<i>Streptomyces albus</i>	142	P24299	I	0
<i>Streptomyces diastaticus</i>	144	9256915	I	0
<i>Streptomyces rubiginosus</i>	146	21730246	I	0
<i>Streptomyces pristinaespiralis</i>	48	197776540	I	1.00E-305
<i>Actinoplanes missouriensis</i>	66	P12851	I	2.70E-305
<i>Actinoplanes missouriensis</i>	140	349936	I	1.30E-302
<i>Streptomyces sp.</i>	46	197764953	I	2.90E-302
<i>Clavibacter michiganensis</i>	42	A5CPC1	I	4.30E-302
<i>Actinoplanes missouriensis</i>	134	443486	I	7.80E-302
<i>Actinoplanes missouriensis</i>	147	443568	I	7.80E-302
<i>Clavibacter michiganensis</i>	2	B0RIF1	I	1.40E-301
<i>Actinoplanes sp.</i>	24	P10654	I	2.10E-301
<i>Arthrobacter sp.</i>	126	A0JXN9	I	2.30E-301

TABLE 2

<i>Actinoplanes missouriensis</i>	145	443526		2.30E-301
<i>Actinoplanes missouriensis</i>	143	443303		4.00E-301
<i>Arthrobacter sp.</i>	54	P12070		3.50E-298
<i>Arthrobacter aurescens</i>	90	119964059		1.10E-297
<i>Micromonospora sp.</i>	112	237882534		2.00E-297
<i>Arthrobacter chlorophenolicus</i>	4	220912923		3.30E-296
<i>Arthrobacter sp.</i>	133	231103		3.20E-294
<i>Arthrobacter sp.</i>	141	2914276		1.10E-293
<i>Arthrobacter sp.</i>	12	60615686		8.70E-293
<i>Acidothermus cellulolyticus</i>	30	117929271		3.50E-292
<i>Streptomyces griseus</i>	118	182434863		1.50E-291
<i>Salinispora arenicola</i>	18	159039501		8.10E-289
<i>Geodermatophilus obscurus</i>	64	227404617		1.80E-286
<i>Catenulispora acidiphila</i>	104	229246901		5.60E-286
<i>Salinispora tropica</i>	44	145596104		1.80E-282
<i>marine actinobacterium</i>	110	88856315		8.10E-282
<i>Mycobacterium smegmatis</i>	10	118469437		5.30E-281
<i>Saccharomonospora viridis</i>	76	229886404		3.40E-278
<i>Streptomyces rochei</i>	137	P22857		3.40E-275
<i>Mycobacterium vanbaalenii</i>	120	120406242		7.00E-274
<i>Streptomyces lividans</i>	78	Q9RFM4		1.70E-272
<i>Thermus thermophilus</i>	100	P26997		4.80E-272
<i>Nakamurella multipartita</i>	38	229221673		1.00E-271

TABLE 2

<i>Thermus caldophilus</i>				1.90E-
	131	4930285		269
<i>Thermus caldophilus</i>				2.70E-
	132	P56681		269
<i>Streptosporangium roseum</i>				1.90E-
	82	229851079		264
<i>Beutenbergia cavernae</i>				9.10E-
	62	229821786		262
<i>Cellulomonas flavigena</i>				1.10E-
	74	229243977		261
<i>Kribbella flavida</i>				4.80E-
	8	227382478		261
<i>Thermobifida fusca</i>				5.90E-
	114	72162004		261
<i>Janibacter sp.</i>				1.40E-
	58	84495191		259
<i>Nocardiopsis dassonvillei</i>				1.80E-
	36	229207664		258
<i>Actinosynnema mirum</i>				4.70E-
	6	226865307		257
<i>Nocardioides sp.</i>				6.40E-
	84	119716602		256
<i>Actinomyces urogenitalis</i>				3.70E-
	14	227497116		250
<i>Meiothermus ruber</i>				1.40E-
	52	227992647		245
<i>Brachybacterium faecium</i>				3.10E-
	60	237671435		245
<i>Jonesia denitrificans</i>				1.60E-
	94	227383768		244
<i>Kribbella flavida</i>				3.80E-
	86	227381155		244
<i>Frankia sp.</i>				2.50E-
	80	158316430		242
<i>Actinomyces odontolyticus</i>				8.80E-
	70	154508186		239
<i>Meiothermus silvanus</i>				1.90E-
	22	227989553		229
<i>Leifsonia xyli</i>				2.50E-
	92	50954171		226
<i>Mobiluncus curtisii</i>				2.50E-
	26	227493823		225
<i>Mobiluncus mulieris</i>				9.70E-
	72	227875705		225
<i>Deinococcus geothermalis</i>				1.80E-
	124	94972159		224

TABLE 2

<i>Stackebrandtia nassauensis</i>				9.40E-
	98	229862570	I	224
<i>Roseiflexus sp.</i>				5.10E-
	50	148656997	I	223
<i>Thermobaculum terrenum</i>				5.60E-
	56	227374836	I	222
<i>Roseiflexus castenholzii</i>				3.10E-
	88	156742580	I	221
<i>Herpetosiphon aurantiacus</i>				1.10E-
	28	159898286	I	204
<i>Xylanimonas cellulositytica</i>				1.90E-
	40	227427650	I	204
<i>Herpetosiphon aurantiacus</i>				8.70E-
	116	159897776	I	203
<i>Acidobacteria bacterium</i>				2.20E-
	102	94967932	I	181
<i>Geobacillus kaustophilus</i>	305	Q5KYS6	II	1.50E-07
<i>Geobacillus thermodenitrificans</i>	195	A4IP67	II	3.80E-06
<i>Tetragenococcus halophilus</i>	180	O82845	II	0.00028
<i>Bacillus licheniformis</i>	237	P77832	II	0.00036
<i>Mesorhizobium sp.</i>	250	Q11EH9	II	0.00085
<i>Bacillus subtilis</i>	220	P04788	II	0.0038
<i>Thermotoga maritima</i>	207	Q9X1Z5	II	0.0049
<i>Thermotoga sp.</i>	227	B1LB08	II	0.0049
<i>Bacillus amyloliquefaciens</i>	212	A7Z522	II	0.012
<i>Thermotoga petrophila</i>	268	A5ILR5	II	0.03
<i>Bacillus sp.</i>	260	P54272	II	0.041
<i>Bifidobacterium longum</i>	175	B3DR33	II	0.068
<i>Geobacillus stearothermophilus</i>	238	P54273	II	0.072
<i>Bacillus megaterium</i>	214	O08325	II	0.28
<i>Bacillus cereus</i>	228	Q739D2	II	0.46
<i>Thermotoga neapolitana</i>	232	P45687	II	1.1
<i>Fervidobacterium gondwanense</i>	246	Q6T6K9	II	1.5
<i>Bacillus halodurans</i>	277	Q9K993	II	1.7
<i>Lactobacillus pentosus</i>	269	P21938	II	1.8
<i>Staphylococcus xylosus</i>	206	P27157	II	2.5
<i>Bifidobacterium adolescentis</i>	205	A1A0H0	II	3.9
<i>Bacillus clausii</i>	278	Q5WKJ3	II	4.2
<i>Bifidobacterium longum</i>	161	Q8G3Q1	II	11
<i>Lactococcus lactis</i>	270	Q9CFG7	II	15
<i>Bacillus pumilus</i>	296	A8FE33	II	21
<i>Thermoanaerobacterium saccharolyticum</i>	284	P30435	II	62

TABLE 2

<i>Thermoanaerobacterium</i> <i>thermosulfurigenes</i>	267	P19148	II	75
<i>Thermoanaerobacter</i> <i>pseudethanolicus</i>	176	P22842	II	1.10E+02
<i>Thermoanaerobacter</i> sp.	172	B0K1L3	II	1.40E+02
<i>Rhizobium leguminosarum</i>	240	Q1MBL8	II	2.40E+02
<i>Thermoanaerobacterium</i> <i>thermosaccharolyticum</i>	257	P29441	II	2.90E+02
<i>Lactobacillus brevis</i>	263	P29443	II	3.00E+02
<i>Lactobacillus brevis</i>	274	Q03TX3	II	3.20E+02
<i>Listeria welshimeri</i>	303	A0AF79	II	3.50E+02
<i>Bacteroides thetaiotaomicron</i>	196	Q8A9M2	II	3.90E+02
<i>Thermoanaerobacter yonseiensis</i>	304	Q9KGU2	II	4.10E+02
<i>Rhizobium etli</i>	290	Q2K433	II	4.20E+02
<i>Salmonella enterica</i>	148	B4T952	II	4.30E+02
<i>Klebsiella pneumoniae</i>	149	P29442	II	4.30E+02
<i>Sinorhizobium meliloti</i>	150	Q92LW9	II	4.30E+02
<i>Escherichia coli</i>	151	Q7A9X4	II	4.30E+02
<i>Salmonella enterica</i>	152	Q5PLM6	II	4.30E+02
<i>Xanthomonas campestris</i>	153	Q3BMF2	II	4.30E+02
<i>Pectobacterium atrosepticum</i>	154	Q6DB05	II	4.30E+02
<i>Rhodopirellula ballica</i>	155	Q7UVG2	II	4.30E+02
<i>Xanthomonas axonopodis</i>	156	Q8PEW5	II	4.30E+02
<i>Xanthomonas oryzae</i>	157	Q5GUF2	II	4.30E+02
<i>Pediococcus pentosaceus</i>	158	Q03HN1	II	4.30E+02
<i>Brucella suis</i>	159	Q8G204	II	4.30E+02
<i>Escherichia coli</i>	160	Q0TBN7	II	4.30E+02
<i>Brucella canis</i>	162	A9M9H3	II	4.30E+02
<i>Burkholderia multivorans</i>	163	A9ARG7	II	4.30E+02
<i>Brucella ovis</i>	164	A5VPA1	II	4.30E+02
<i>Rhizobium etli</i>	165	B3Q0R5	II	4.30E+02
<i>Burkholderia xenovorans</i>	166	Q13RB8	II	4.30E+02
<i>Actinobacillus pleuropneumoniae</i>	167	A3N3K2	II	4.30E+02
<i>Burkholderia cenocepacia</i>	168	B4ENA5	II	4.30E+02
<i>Solibacter usitatus</i>	169	Q022S9	II	4.30E+02
<i>Brucella abortus</i>	170	B2SA37	II	4.30E+02
<i>Rhodobacter sphaeroides</i>	171	A4WVT8	II	4.30E+02
<i>Yersinia pseudotuberculosis</i>	173	Q1C0D3	II	4.30E+02
<i>Xanthomonas oryzae</i>	174	Q5GYQ7	II	4.30E+02
<i>Photobacterium profundum</i>	177	Q6LUY7	II	4.30E+02
<i>Escherichia coli</i>	178	B1LJC7	II	4.30E+02
<i>Agrobacterium tumefaciens</i>	179	Q8U7G6	II	4.30E+02
<i>Salmonella enterica</i>	181	B4TZ55	II	4.30E+02

TABLE 2

<i>Yersinia pseudotuberculosis</i>	182	Q8Z9Z1	II	4.30E+02
<i>Yersinia pseudotuberculosis</i>	183	Q1CDB8	II	4.30E+02
<i>Rhodobacter sphaeroides</i>	184	A3PNM4	II	4.30E+02
<i>Brucella abortus</i>	185	Q2YMQ2	II	4.30E+02
<i>Salmonella enterica</i>	186	Q8ZL90	II	4.30E+02
<i>Bacteroides vulgatus</i>	187	A6L792	II	4.30E+02
<i>Xanthomonas axonopodis</i>	188	Q8PLL9	II	4.30E+02
<i>Salmonella enterica</i>	189	Q57IG0	II	4.30E+02
<i>Escherichia coli</i>	190	B7M3I8	II	4.30E+02
<i>Roseobacter denitrificans</i>	191	Q162B6	II	4.30E+02
<i>Bacteroides fragilis</i>	192	Q64U20	II	4.30E+02
<i>Enterobacter sakazakii</i>	193	A7MNI5	II	4.30E+02
<i>Brucella abortus</i>	194	Q57EI4	II	4.30E+02
<i>Haemophilus influenzae</i>	197	A5UCZ3	II	4.30E+02
<i>Yersinia pseudotuberculosis</i>	198	B2K7D2	II	4.30E+02
<i>Xanthomonas campestris</i>	199	Q4UTU6	II	4.30E+02
<i>Haemophilus somnus</i>	200	B0UT19	II	4.30E+02
<i>Pseudoalteromonas atlantica</i>	201	Q15PG0	II	4.30E+02
<i>Escherichia fergusonii</i>	202	B7LTH9	II	4.30E+02
<i>Silicibacter sp.</i>	203	Q1GKQ4	II	4.30E+02
<i>Salmonella enterica</i>	204	B5R4P8	II	4.30E+02
<i>Salmonella enterica</i>	208	A9MUV0	II	4.30E+02
<i>Pseudomonas syringae</i>	209	Q48J73	II	4.30E+02
<i>Shigella boydii</i>	210	Q31V53	II	4.30E+02
<i>Burkholderia ambifaria</i>	211	Q0B1U7	II	4.30E+02
<i>Haemophilus influenzae</i>	213	A5UIN7	II	4.30E+02
<i>Arabidopsis thaliana</i>	215	Q9FKK7	II	4.30E+02
<i>Escherichia coli</i>	216	Q3YVV0	II	4.30E+02
<i>Bacteroides fragilis</i>	217	Q5LCV9	II	4.30E+02
<i>Pseudomonas fluorescens</i>	218	Q3KDW0	II	4.30E+02
<i>Escherichia coli</i>	219	B1X8I1	II	4.30E+02
<i>Xanthomonas campestris</i>	221	Q4UNZ4	II	4.30E+02
<i>Pseudomonas syringae</i>	222	Q4ZSF5	II	4.30E+02
<i>Sinorhizobium medicae</i>	223	A6UD89	II	4.30E+02
<i>Ochrobactrum anthropi</i>	224	A6X4G3	II	4.30E+02
<i>Burkholderia thailandensis</i>	225	Q2SW40	II	4.30E+02
<i>Salmonella enterica</i>	226	B5EX72	II	4.30E+02
<i>Salmonella enterica</i>	229	B4SWK9	II	4.30E+02
<i>Salmonella enterica</i>	230	Q7C637	II	4.30E+02
<i>Enterococcus faecalis</i>	231	Q7C3R3	II	4.30E+02
<i>Escherichia coli</i>	233	B7MES1	II	4.30E+02
<i>Photobacterium luminescens</i>	234	Q7N4P7	II	4.30E+02
<i>Enterobacter sp.</i>	235	A4W566	II	4.30E+02

TABLE 2

<i>Burkholderia cenocepacia</i>	236	B1KB47		4.30E+02
<i>Brucella abortus</i>	239	Q8YFX5		4.30E+02
<i>Yersinia enterocolitica</i>	241	A1JT10		4.30E+02
<i>Serratia proteamaculans</i>	242	A8G7W8		4.30E+02
<i>Yersinia pseudotuberculosis</i>	243	A7FP68		4.30E+02
<i>Escherichia coli</i>	244	B7NEL7		4.30E+02
<i>Yersinia pestis</i>	245	A9R5Q1		4.30E+02
<i>Xanthomonas campestris</i>	247	Q8P3H1		4.30E+02
<i>Rhizobium leguminosarum</i>	248	B5ZQV6		4.30E+02
<i>Bradyrhizobium japonicum</i>	249	Q89VC7		4.30E+02
<i>Actinobacillus pleuropneumoniae</i>	251	B3H2X9		4.30E+02
<i>Yersinia pseudotuberculosis</i>	252	Q663Y3		4.30E+02
<i>Xanthomonas campestris</i>	253	Q8P9T9		4.30E+02
<i>Burkholderia cenocepacia</i>	254	A0KE56		4.30E+02
<i>Oceanobacillus iheyensis</i>	255	Q8ELU7		4.30E+02
<i>Brucella suis</i>	256	B0CKM9		4.30E+02
<i>Burkholderia phymatum</i>	258	B2JFE9		4.30E+02
<i>Yersinia pseudotuberculosis</i>	259	B1JH40		4.30E+02
<i>Lactococcus lactis</i>	261	Q02Y75		4.30E+02
<i>Novosphingobium aromaticivorans</i>	262	Q2GAB9		4.30E+02
<i>Mesorhizobium loti</i>	264	Q98CR8		4.30E+02
<i>Escherichia coli</i>	265	A8A623		4.30E+02
<i>Burkholderia cenocepacia</i>	266	Q1BG90		4.30E+02
<i>Ruminococcus flavefaciens</i>	271	Q9S306		4.30E+02
<i>Burkholderia phytomans</i>	272	B2T929		4.30E+02
<i>Salmonella enterica</i>	273	B5FLD6		4.30E+02
<i>Burkholderia ambifaria</i>	275	B1Z405		4.30E+02
<i>Salmonella enterica</i>	276	B5RGL6		4.30E+02
<i>Marinomonas sp.</i>	279	A6VWH1		4.30E+02
<i>Yersinia pseudotuberculosis</i>	280	A4TS63		4.30E+02
<i>Actinobacillus pleuropneumoniae</i>	281	B0BTI9		4.30E+02
<i>Silicibacter pomeroyi</i>	282	Q5LV46		4.30E+02
<i>Xanthomonas oryzae</i>	283	Q2NXR2		4.30E+02
<i>Escherichia coli</i>	285	B6I3D6		4.30E+02
<i>Escherichia coli</i>	286	B5YVL8		4.30E+02
<i>Escherichia coli</i>	287	B7NP65		4.30E+02
<i>Escherichia coli</i>	288	B2U560		4.30E+02
<i>Escherichia coli</i>	289	B1IZM7		4.30E+02
<i>Escherichia coli</i>	291	P00944		4.30E+02
<i>Hordeum vulgare</i>	292	Q40082		4.30E+02
<i>Dinoroseobacter shibae</i>	293	A8LP53		4.30E+02
<i>Rhodobacter sphaeroides</i>	294	Q3IYM4		4.30E+02
<i>Actinobacillus succinogenes</i>	295	A6VLM8		4.30E+02

TABLE 2

<i>Escherichia coli</i>	297	Q8FCE3	II	4.30E+02
<i>Pseudomonas syringae</i>	298	Q880Z4	II	4.30E+02
<i>Burkholderia vietnamiensis</i>	299	A4JSU5	II	4.30E+02
<i>Escherichia coli</i>	300	A7ZTB2	II	4.30E+02
<i>Haemophilus influenzae</i>	301	P44398	II	4.30E+02
<i>Haemophilus influenzae</i>	302	Q4QLI2	II	4.30E+02
<i>Mannheimia succiniciproducens</i>	306	Q65PY0	II	4.30E+02

CLAIMS

What is claimed is:

1. A recombinant bacterial strain selected from the group
5 consisting of *Zymomonas* and *Zymobacter* comprising a heterologous nucleic acid molecule encoding a polypeptide having xylose isomerase activity wherein the polypeptide is a Group I xylose isomerase and is included in the class of enzymes identified by EC 5.3.1 .5, and wherein the strain utilizes xylose as a carbon source.
- 10 2. A recombinant bacterial strain of claim 1 wherein the polypeptide having xylose isomerase activity gives an E-value score of 1E-15 or less when queried using a Profile Hidden Markov Model prepared using SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108,
15 122, 126, 128, 130, 132, 135, 137, and 142; the query being carried out using the hmmsearch algorithm wherein the Z parameter is set to 1 billion.
3. A recombinant bacterial strain of claim 1 wherein the polypeptide having xylose isomerase activity:
20 1) has the following conserved amino acids when compared with the reference amino acid sequence of SEQ ID NO:66:
 - a) leucine at position 226,
 - b) methionine at position 223,
 - c) isoleucine at position 191,
 - 25 d) threonine, serine, or valine at position 195,
 - e) methionine, threonine or quanine at position 88,
 - f) histidine at position 290,
 - g) glutamic acid or aspartic acid at position 221 ,
 - h) phenylalanine, valine, or leucine at position 242,
 - 30 i) histidine at position 243,
 - j) leucine, phenylalanine, or methionine at position 193,
 - k) glutamine at position 256,
 - l) glycine at position 213,
 - m) proline, tyrosine, alanine, or serine at position 288, and

n) glutamine at position 249; or

2) has at least 90% of the conserved amino acids of part (1)

4. A recombinant bacterial strain of claim 1 wherein the xylose
isomerase has an amino acid sequence having at least 90% identity to an
amino acid sequence selected from the group consisting of SEQ ID
NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38,
40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76,
78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110,
112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 131, 132, 133, 134,
135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, and 147
based on the Clustal W method of alignment using the default parameters
of GAP PENALTY=10, GAP LENGTH PENALTY=0.1, and Gonnet 250
series of protein weight matrix.

15

5. A recombinant bacterial strain of claim 1 wherein the xylose
isomerase is isolated from a microorganism selected from the group
consisting of *Actinoplanes*, *Arthrobacter*, *Streptomyces*, *Thermus*,
Thermobaculum, *Herpetosiphon*, *Acidobacteria*, *Roseiflexus*,
Meiothermus, *Deinococcus*, *Meiothermus*, *Stackebrandtia*, *Kribbella*,
Xylanimonas, *Nocardiopsis*, *Catenulispora*, *Streptosporangium*,
Geodermatophilus, *Actinosynnema*, *Saccharomonospora*, *Acidothermus*,
Tthermobifida, *Nocardiodes*, *Janibacter*, *Mycobacterium*, *Leifsonia*,
Clavibacter, *Micromonospora*, *Salinispora*, *Cellulomonas*, *Jonesia*,
Nakamurella, *Actinomyces*, *Mobiluncus*, *Brachybacterium*, *Beutengergai*,
Frankia, and *Actinobacterium*.

6. A recombinant bacterial strain of claim 5 wherein the xylose
isomerase is isolated from *Actinoplanes missouriensis*.

30

7. A recombinant bacterial strain of claim 4 wherein the Group I
polypeptide having xylose isomerase activity has at least 90% identity to
an amino acid sequence selected from the group consisting of SEQ ID
NOs:24, 66, 134, 140, 143, 145, and 147, based on the Clustal W method

of alignment using the default parameters of GAP PENALTY=10, GAP LENGTH PENALTY=0.1, and Gonnet 250 series of protein weight matrix.

8. A process for improving xylose utilization in a recombinant bacterial cell comprising:
- a) providing a recombinant bacterial strain selected from the group consisting of *Zymomonas* and *Zymobacter* comprising xylulokinase, transketolase, transaldolase and optionally a xylose isomerase not belonging to Group I; and
 - b) introducing a heterologous nucleic acid molecule encoding a polypeptide having xylose isomerase activity wherein the polypeptide is a Group I xylose isomerase;
- wherein xylose utilization is improved as compared to the same strain containing a xylose isomerase not belonging to Group I.

9. The process of claim 8 wherein the polypeptide having xylose isomerase activity gives an E-value score of $1E-15$ or less when queried using a Profile Hidden Markov Model prepared using SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142; the query being carried out using the hmmsearch algorithm wherein the Z parameter is set to 1 billion.

10. The process of claim 8 wherein the polypeptide having xylose isomerase activity
- 1) has the following conserved amino acids when compared with the reference amino acid sequence of SEQ ID NO:66:
 - a) leucine at position 226,
 - b) methionine at position 223,
 - c) isoleucine at position 191,
 - d) threonine, serine, or valine at position 195,
 - e) methionine, threonine or glutamine at position 88,
 - f) histidine at position 290,
 - g) glutamic acid or aspartic acid at position 221,
 - h) phenylalanine, valine, or leucine at position 242,

- i) histidine at position 243,
 - j) leucine, phenylalanine, or methionine at position 193,
 - k) glutamine at position 256,
 - l) glycine at position 213,
 - 5 m) proline, tyrosine, alanine, or serine at position 288, and
 - nj) glutamine at position 249; or
- 2) has at least 90% of the conserved amino acids of part (1).

11. A method for the production of ethanol comprising:
- 10 a) providing the recombinant bacterial strain of Claim 1; and
 - b) contacting the strain of (a) with xlyose under conditions whereby the strain produces ethanol.

12. A method according to claim 11 wherein the polypeptide having
- 15 xylose isomerase activity gives an E-value score of $1E-15$ or less when queried using a Profile Hidden Markov Model prepared using SEQ ID NOs: 2, 24, 32, 34, 42, 54, 66, 68, 78, 96, 100, 106, 108, 122, 126, 128, 130, 132, 135, 137, and 142; the query being carried out using the hmmsearch algorithm wherein the Z parameter is set to 1 billion.

20

13. A method according to claim 11 wherein the polypeptide having xylose isomerase activity:

1) has the following conserved amino acids when compared with the reference amino acid sequence of SEQ ID NO:66:

- 25 a) leucine at position 226,
- b) methionine at position 223,
- c) isoleucine at position 191,
- d) threonine, serine, or valine at position 195,
- e) methionine, threonine or quanine at position 88,
- 30 f) histidine at position 290,
- g) glutamic acid or aspartic acid at position 221,
- h) phenylalanine, valine, or leucine at position 242,
- i) histidine at position 243,
- j) leucine, phenylalanine, or methionine at position 193,

- k) glutamine at position 256,
 - l) glycine at position 213,
 - m) proline, tyrosine, alanine, or serine at position 288, and
 - n) glutamine at position 249; or
- 5 2) has at least 90% of the conserved amino acids of part (1).

14. A method according to claim 11 wherein the xylose isomerase has an amino acid sequence having at least 90% identity to an amino acid sequence selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 10 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, and 147 based on the Clustal W 15 method of alignment using the default parameters of GAP PENALTY=10, GAP LENGTH PENALTY=0.1, and Gonnet 250 series of protein weight matrix.

15. A method according to claim 11 wherein the xylose isomerase 20 is isolated from a microorganism selected from the group consisting of *Actinoplanes*, *Arthrobacter*, *Streptomyces*, *Thermus*, *Thermobaculum*, *Herpetosiphon*, *Acidobacteria*, *Roseiflexus*, *Meiothermus*, *Deinococcus*, *Meiothermus*, *Stackebrandtia*, *Kribbella*, *Xylanimonas*, *Nocardiopsis*, *Catenulispora*, *Streptosporangium*, *Geodermatophilus*, *Actinosynnema*, 25 *Saccharomonospora*, *Acidothrmus*, *Thermobifida*, *Nocardiodes*, *Janibacter*, *Mycobacterium*, *Leifsonia*, *Clavibacter*, *Micromonospora*, *Salinispora*, *Cellulomonas*, *Jonesia*, *Nakamurella*, *Actinomyces*, *Mobiluncus*, *Brachybacterium*, *Beutengergai*, *Frankia*, and *Actinobacterium*.

30

Figure 1




Figure 2

- Group I
- Group II
- ▨ *A. missouriensis*
- ▲ *L. brevis*
- ▨ *E. coli*




Figure 3

Group I

- ▲ *Arthrobacter*
- *A. missouriensis*
- *Thermus*
- *Streptomyces*




Figure 4




Figure 5




Figure 6





Figure 7



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/042132

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N9/92 C12P7/06
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal , WPI Data, BIOSIS, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2009/120731 A2 (DU PONT [US] ; ALLIANCE SUSTAINABLE ENERGY [US] ; VI ITANEN PAUL V [US] ;) 1 October 2009 (2009-10-01) cited in the appl icati on Whole document especi ally paragraph 29 and table 1.	1-15
A	----- VAN BASTELAERE ET AL: "Ki neti c studi es of Mg(2+) - , Co(2+) - and Mn(2+) -activated D-xyl ose i somerases. ", BIOCHEMICAL JOURNAL, vol . 278, no. 1, 1 August 1991 (1991-08-01) , pages 285-292 , XP55009184, ISSN : 0264-6021 the whole document ----- - / - -	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 October 2011

Date of mailing of the international search report

19/10/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Kool s , Patri ck

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2011/042132

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LIU C-Q ET AL: "Expressi on of cloned Xanthomonas d-xyl ose catabol ic genes in Zymomonas mobi lis", JOURNAL OF BIOTECHNOLOGY, ELSEVI ER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol . 7, no. 1, 1 January 1988 (1988-01-01) , pages 61-70, XP023704758, ISSN: 0168-1656, DOI : 10. 1016/0168-1656(88)90035-1 [retri eved on 1988-01-01] abstract -----	1-15
A	US 6 475 768 B1 (OTERO RICARDO ROMAN CORDERO [ZA] ET AL) 5 November 2002 (2002-11-05) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 2011/ 042132

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. (means)

☐
☐

on paper

in electronic form
 - b. (time)

☒
☐
☐

in the international application as filed

together with the international application in electronic form

subsequently to this Authority for the purpose of search
2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2011/042132

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009120731 A2	01-10-2009	AU 2009228323 A1 CA 2716585 A1 CN 102037120 A EP 2271743 A2 JP 2011515107 A US 2009246846 A1	01-10-2009 01-10-2009 27-04-2011 12-01-2011 19-05-2011 01-10-2009
US 6475768 B1	05-11-2002	AU 4136800 A WO 0061733 A1	14-11-2000 19-10-2000