

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 November 2006 (09.11.2006)

PCT

(10) International Publication Number
WO 2006/119458 A1

(51) International Patent Classification:

A61K 48/00 (2006.01) *C12N 15/74* (2006.01)
C12N 15/63 (2006.01) *C12N 15/86* (2006.01)

(21) International Application Number:

PCT/US2006/017242

(22) International Filing Date: 2 May 2006 (02.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/677,057 2 May 2005 (02.05.2005) US
60/685,808 31 May 2005 (31.05.2005) US

(71) Applicant (for all designated States except US): GENZYME CORPORATION [US/US]; 500 Kendall Street, Cambridge, Massachusetts 02142 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PASSINI, Marco, A. [US/US]; 14 Nightingale Drive, Shrewsbury, MA 01545 (US). DODGE, James [US/US]; 1 Gates Circle, Worcester, Massachusetts 01603 (US). STEWART, Gregory, R. [US/US]; 4735 Xene Lane South, Plymouth, Minnesota 55446 (US).

(74) Agents: TOUSIGNANT, Jennifer, D. et al.; GENZYME CORPORATION, 15 Pleasant Street Connector, Framingham, MA 01701-9322 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2006/119458 A1

(54) Title: GENE THERAPY FOR NEUROMETABOLIC DISORDERS

(57) Abstract: The disclosure pertains to methods and compositions for treating disorders affecting the central nervous system (CNS). These disorders include neurometabolic disorders such as lysosomal storage diseases that affect the central nervous system, e.g., Niemann-Pick A disease. They also include disorders such as Alzheimer's disease. The disclosed methods involve contacting an axonal ending of a neuron with a composition containing high titer AAV carrying a therapeutic transgene so that the AAV vector is axonally transported in a retrograde fashion and transgene product is expressed distally to the administration site.

NOT FURNISHED AT THE TIME OF PUBLICATION

NOT FURNISHED AT THE TIME OF PUBLICATION

<i>Lysosomal storage disease</i>	<i>Defective enzyme</i>
Wolman*	Acid Lipase

* CNS involvement

[004] The hallmark feature of LSD is the abnormal accumulation of metabolites in the lysosomes which leads to the formation of large numbers of distended lysosomes in the perikaryon. A major challenge to treating LSD (as opposed to treating a liver-specific enzymopathy) is the need to reverse lysosomal storage pathology in multiple separate tissues. Some LSDs can be effectively treated by intravenous infusion of the missing enzyme, known as enzyme replacement therapy (ERT). For example, Gaucher type 1 patients have only visceral disease and respond favorably to ERT with recombinant glucocerebrosidase (Cerezyme®, Genzyme Corp.). However, patients with metabolic disease that affects the CNS (e.g., type 2 or 3 Gaucher disease) do not respond to intravenous ERT because the replacement enzyme is prevented from entering the brain by the blood brain barrier (BBB). Furthermore, attempts to introduce a replacement enzyme into the brain by direct injection have been unsuccessful in part due to enzyme cytotoxicity at high local concentrations (unpublished observations) and limited parenchymal diffusion rates in the brain (Pardridge, Peptide Drug Delivery to the Brain, Raven Press, 1991).

[005] Alzheimer's disease (AD) is a disorder affecting the central nervous system (CNS) characterized by the accumulation of amyloid β -peptide (A β) due to decreased A β catabolism. As A β accumulates, it aggregates into extracellular plaques, causing impairment of synaptic function and loss of neurons. The pathology leads to dementia, loss of coordination, and death.

[006] Gene therapy is an emerging treatment modality for disorders affecting the CNS, including LSDs and Alzheimer's disease. In this approach, restoration of the normal metabolic pathway and reversal of pathology occurs by transducing affected cells with a vector carrying a healthy version or a modified version of the gene.

[007] CNS gene therapy has been facilitated by the development of viral vectors capable of effectively infecting post-mitotic neurons. For a review of viral vectors for gene delivery to the CNS, see Davidson et al. (2003) *Nature Rev.*, 4:353-364. Adeno-associated virus (AAV) vectors are considered optimal for CNS gene therapy because they have a favorable toxicity and immunogenicity profile, are able to transduce neuronal cells, and are able to mediate long-term expression in the CNS (Kaplitt et al. (1994) *Nat. Genet.*, 8:148-154; Bartlett et al. (1998) *Hum. Gene Ther.*, 9:1181-1186; and Passini et al. (2002) *J. Neurosci.*, 22:6437-6446).

[008] A therapeutic transgene product, e.g., an enzyme, can be secreted by transduced cells and subsequently taken up by other cells, in which it then alleviates pathology. This process is known as cross-correction (Neufeld et al. (1970) *Science*, 169:141-146). However, the correction of pathology, such as storage pathology in the context of LSD, is typically confined to the immediate vicinity of the injection site because of limited parenchymal diffusion of the injected vector and the secreted transgene product (Taylor et al. (1997) *Nat. Med.*, 3:771-774; Skorupa et al. (1999) *Exp. Neurol.*, 160:17-27). Thus, neuropathology affecting multiple brain regions requires widespread vector delivery, using multiple spatially distributed injections, especially in a large brain such as human. This significantly increases the risk of brain damage. In addition, some regions of the brain may be difficult to access surgically. Thus, other modes of vector transport within the CNS, besides diffusion, would be beneficial.

[009] When administered at axonal endings, some viruses are internalized and transported retrogradely along the axon to the nucleus. Neurons in one brain region are interconnected by axons to distal brain regions thereby providing a transport system for vector delivery. Studies with adenovirus, HSV, and pseudo-rabies virus have utilized trafficking properties of these viruses to deliver genes to distal structures within the brain (Soudas et al. (2001) *FASEB J.*, 15:2283-2285; Breakefield et al. (1991) *New Biol.*, 3:203-218; and deFalco et al. (2001) *Science*, 291:2608-2613).

[010] Several groups have reported that the transduction of the brain by AAV serotype 2 (AAV2) is limited to the intracranial injection site (Kaplitt et al. (1994) *Nat. Genet.*, 8:148-154; Passini et al. (2002) *J. Neurosci.*, 22:6437-6446; and Chamberlin et al. (1998) *Brain Res.*, 793:169-175). One recent report suggests that retrograde axonal transport of AAV2 can also occur in select circuits of the normal rat brain (Kaspar et al. (2002) *Mol. Ther.*, 5:50-56). However, it is not known what specific parameters were responsible for the observed axonal transport, and whether sufficient and effective axonal transport would occur in a diseased neuron that is in a state of cellular dysfunction. Indeed, lesions observed in LSD neurons have been reported to interfere with or even block axonal transport (reviewed in Walkley (1998) *Brain Pathol.*, 8:175-193), suggesting that disease-compromised neurons would not support trafficking of AAV along their axons.

[011] Therefore, there is a need in the art to develop new therapeutic methods for treating metabolic disorders that affect the CNS.

Summary of the Invention

[012] The invention provides methods and compositions for treating or preventing metabolic disorders, such as lysosomal storage diseases (LSD) or abnormal cholesterol storage function that are characterized by or associated with a risk of diminution of CNS function.

[013] The invention provides methods and compositions for treating or preventing disorders affecting the central nervous system (CNS), such as Alzheimer's disease that are characterized by or associated with a risk of diminution of CNS function.

[014] The invention further provides methods for minimally invasive targeted delivery of a transgene to select regions in the brain of an affected subject.

[015] Additional advantages of the invention will be set forth in part in the following description, and in part will be understood from the description, or may be learned by practice of the invention.

[016] Acid sphingomyelinase (ASM) knockout mice, a model of Niemann-Pick Type A disease, were administered an AAV2 vector carrying the human ASM gene (AAV-ASM) by a single intracranial injection into one hemisphere of the brain. The present invention is based, in part, on the discovery and demonstration that the injection of high titer AAV-ASM into the diseased brain results in AAV-ASM expression within multiple distal sites in a pattern consistent with the topographical organization of the projection neurons that innervate the injection site. The invention is further based, in part, on the discovery and demonstration of extensive correction of lysosomal storage pathology at the injection site and distal sites to which AAV-ASM was transported and where ASM was expressed.

[017] In another aspect, the invention provides a method to correct cholesterol storage pathology and initiate functional recovery in the ASMKO mouse after unilateral or alternatively, bilateral injection within the deep cerebellar nuclei.

[018] Further provided are the above-noted methods wherein the transgene is delivered in a recombinant AAV vector selected from the group consisting of AAV2/1, AAV2/2, AAV2/5, AAV2/7 and AAV2/8 serotype. For the purpose of illustration only, the recombinant vectors encoded functional human ASM protein in a mouse model.

[019] Accordingly, in one aspect, the present invention provides methods for treating neurometabolic disorders in mammals. The populations treated by the methods of the invention include, but are not limited to, patients having or at risk for developing a LSD, such as disorders listed in Table 1, particularly, if such disease affects the CNS. In an illustrative embodiment, the disease is Niemann-Pick A disease and/or the secondary cholesterol storage pathology commonly associated with NPA.

[020] In one aspect, the disclosed methods include administering to the CNS of an afflicted subject an AAV viral vector carrying a transgene encoding a therapeutic product and allowing the transgene to be expressed within the CNS distally from the administration site at a therapeutic level. In addition, the vector may comprise a polynucleotide encoding for a biologically active molecule effective to treat the CNS disorder. Such biologically active molecules may comprise peptides including but not limited to native or mutated versions of full-length proteins, native or mutated versions of protein fragments, synthetic polypeptides, antibodies, and antibody fragments such as Fab' molecules. Biologically active molecules may also comprise nucleotides including single-stranded or double-stranded DNA polynucleotides and single-stranded or double-stranded RNA polynucleotides. For a review of exemplary nucleotide technologies that may be used in the practice of the methods disclosed herein, see Kurreck, (2003) J., Eur. J. Biochem. 270, 1628-1644 [antisense technologies]; Yu et al., (2002) PNAS 99(9), 6047-6052 [RNA interference technologies]; and Elbashir et al., (2001) Genes Dev., 15(2):188-200 [siRNA technology].

[021] In an illustrative embodiment, the administration is accomplished by direct intraparenchymal injection of a high titer AAV vector solution into the diseased brain. Thereafter the transgene is expressed distally, contralaterally or ipsilaterally, to the administration site at a therapeutic level at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm from the administration site.

[022] In another aspect, the invention also provides a method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron *in vivo*. In some embodiments, the cellular pathology exhibited by the neuron is that of a lysosomal storage disease such as disorders listed in Table 1. In an illustrative embodiment, the disease is Niemann-Pick A disease. In other embodiments, the cellular pathology exhibited is that of Alzheimer's disease. The method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron comprises contacting an axonal ending of the disease-compromised neuron with a composition comprising an AAV viral particle comprising the recombinant AAV genome and allowing the viral particle to be

endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron. The concentration of the vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In certain embodiments, the neuron is a projection neuron and/or the distance of the axonal ending to the nucleus of the neuron is at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm.

[023] This invention provides methods and compositions to deliver a transgene to the spinal cord and/ or the brainstem region of a subject by administering a recombinant neurotropic viral vector containing the transgene to at least one region of the deep cerebellar nuclei (DCN) region of the subject's brain. The viral delivery is under conditions that favor expression of the transgene in the spinal cord and/ or the brainstem region. In an illustrative embodiment, the disease is Niemann-Pick A disease. In other embodiments, the cellular pathology exhibited is that of Alzheimer's disease.

[024] In another aspect, the invention provides methods and compositions to deliver a transgene to a subject's spinal cord by administering a recombinant neurotropic viral vector containing the transgene to the motor cortex region of the subject's brain. The delivery of the viral vector is under conditions that favor expression of the transgene in the spinal cord. Viral vectors administered to the motor cortex region are internalized by motor neurons via their cell body region and the transgene is expressed. The expressed transgene may then undergo anterograde transport to the axon terminal portion of the motor neuron, which is present in the spinal cord. Due to the nature of the motor cortex, viral vectors administered to this region of the brain may also be internalized by axon terminals of motor neurons. The viral vector also may undergo retrograde transport along the motor neuron's axon and be expressed in the cell body of the motor neuron. In an illustrative embodiment, the disease is Niemann-Pick A disease. In other embodiments, the cellular pathology exhibited is that of Alzheimer's disease.

[025] In another aspect, the invention provides a method of delivering a therapeutic transgene product to a target cell of the CNS, which is a neuron or a glial cell, in a mammal afflicted with a neurometabolic disorder, e.g., an LSD that affects the CNS. The method includes contacting an axonal ending of a neuron with a composition containing an AAV vector carrying at least a part of a gene encoding a therapeutic transgene product, allowing the viral particle to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron; allowing the therapeutic transgene product to be expressed and secreted by the neuron, and allowing the target cell to uptake the therapeutic transgene product, wherein the therapeutic transgene product thereby alleviates pathology in the target cell. In certain embodiments, the concentration of the vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml).

[026] In the methods of the invention, the therapeutic transgene encodes a biologically active molecule, expression of which in the CNS results in at least partial correction of neuropathology. In some embodiments, the therapeutic transgene product is a lysosomal hydrolase. In an illustrative embodiment, the lysosomal hydrolase is ASM. In other embodiments, the therapeutic transgene is a metalloendopeptidase, e.g., neprilysin.

[027] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention as claimed.

Brief Description of the Figures

[028] **Figure 1A** depicts a representation of a cross-section of the ASMKO mouse brain, at 5 or 15 weeks following a 2 μ l injection of high titer (9.3×10^{12} gp/ml) AAV-ASM into the hippocampus. The site of injection is shown by a vertical line; ASM mRNA expression, as detected by *in situ* hybridization, is represented by the smaller circles; and ASM protein expression, as detected by

immunohistochemical staining, is represented by the larger shaded circles. The expression pattern resulted in an extensive area of reversal of pathology (represented by the light shading) in the hippocampus and cortical regions in both hemispheres of the brain.

[029] **Figure 1B** depicts the axonal transport of AAV to distal regions of the mouse brain following a high titer AAV injection into the hippocampus as described for Figure 1A. Injection into the hippocampus (10) resulted in axonal transport of the viral vector via the intrahippocampal circuit to the contralateral hippocampus (20) and via the entorhinodentate circuit to the entorhinal cortex (30). The site of injection is shown by a vertical line.

[030] **Figure 1C** is a schematic diagram showing the connections of the intrahippocampal and entorhinodentate circuits of the mouse brain. Injection into the hippocampus (10) results in infection and transduction of cell bodies located in the cornu ammonis area 3 (CA3) and in the dentate granule cell layer (G). In addition, a subset of the injected AAV vector infects the axonal endings of the projection neurons innervating the injection site, undergoes retrograde axonal transport, and delivers the transgene to the CA3 field (CA3) and hilus (H) in the contralateral part of the hippocampus (20), and ipsilaterally in the entorhinal cortex (30).

[031] **Figure 2A** depicts a representation of a cross-section of the ASMKO mouse brain, at 5 or 15 weeks following an intrahippocampal injection of high titer AAV-ASM as described in Figure 1A. ASM mRNA expression, as detected by *in situ* hybridization, is represented by the smaller circles; and ASM protein expression, as detected by immunohistochemical staining, is represented by the larger shaded circles. The injection resulted in ASM mRNA and protein to be detected in the septum. This expression pattern resulted in an extensive area of reversal of pathology (represented by the light shading).

[032] **Figure 2B** depicts the axonal transport of AAV to distal regions of the mouse brain, following a high titer injection into the hippocampus as described in Figure 1A. Injection into the hippocampus (10) resulted in axonal transport of the

viral vector via the septohippocampal circuit from the injection site (represented by a vertical line) to the septum (40).

[033] **Figure 2C** is a schematic diagram showing the connections of the septohippocampal circuit. Injection into the hippocampus resulted in transduction to cell bodies located in the CA3 field (11). In addition, a subset of the AAV vector infects the axonal endings of the projection neurons innervating the injection site, undergoes retrograde axonal transport, and delivers the transgene to the medial septum (40).

[034] **Figure 3** depicts the axonal transport of AAV in the nigrostriatal circuit, following a high titer injection of AAV into the striatum (50) of the mouse brain. Axonal transport of AAV occurs from the injection site (represented by a vertical line) to the substantia nigra (60).

[035] **Figure 4** depicts the axonal transport of AAV in the medullocerebellar circuit, following a high titer injection of AAV-ASM into the cerebellum (70) of the ASMKO mouse brain. Axonal transport of AAV2 occurs from the injection site (represented by a vertical line) to the medulla oblongata (80).

[036] **Figure 5** depicts axonal transport of AAV in the intrahippocampal, nigrostriatal, and entorhinodentate circuits following high-titer injection of AAV7-ASM in the ipsilateral hippocampus (10). Transduced cells were detected, as determined by *in situ* hybridization, along the entire rostral-caudal axis of the contralateral hippocampus (90), medial septum (40), and entorhinal cortex (100) after AAV7-ASM injection of the ipsilateral hippocampus (represented by a vertical line).

[037] **Figures 6A through 6E** show human ASM immunopositive staining in sagittal cerebellar sections following injection of different AAV serotype vectors [(A)2/1, (B)2/2, (C)2/5, (D)2/7 and (E)2/8] encoding for human ASM into the deep cerebellar nuclei of ASMKO mice.

[038] **Figures 7A through 7E** demonstrate hASM transport to the spinal cord from the deep cerebellar nuclei. This effect was observed in mice treated with AAV2/2-ASM, AAV2/5 -ASM, AAV2/7-ASM & AAV2/8-ASM (A) hASM 10X magnification; (B) hASM 40X magnification; (C) confocal hASM; (D) confocal ChAT; and (E) confocal hASM & ChAT.

[039] **Figure 8** shows cerebellar tissue homogenate levels following injection of different AAV serotype vectors (2/1, 2/2, 2/5, 2/7 and 2/8) encoding for human ASM into the deep cerebellar nuclei (n=5/group). Groups not connected by the same letter are significantly (p<.0001) different.

[040] **Figures 9A through G** show calbindin immunopositive staining in sagittal cerebellar sections following injection of different AAV serotype vectors [(A)2/1, (B)2/2, (C)2/5, (D)2/7 and (E)2/8] encoding for human ASM into the deep cerebellar nuclei of ASMKO mice.

[041] **Figures 10A and 10B** show accelerating and rocking rotarod performance (at 14 weeks of age) in ASMKO (injected with AAV- β gal), WT, and AAV-ASM treated ASMKO mice (n=8/group). Groups not connected by the same letter are significantly different. Mice injected with AAV2/1-ASM and AAV2/8-ASM demonstrated a significantly (p< .0009) longer latency to fall than ASMKO mice injected with AAV2/1- β gal in the accelerating rotarod test. For the rocking rotarod test, mice injected with AA2/1-ASM demonstrated a significantly (p< .0001) longer latency to fall than mice injected with AAV2/1- β gal.

[042] **Figures 11A and 11B** show Rotarod performance in ASMKO (n=8), WT (n=8), and bilaterally AAV-ASM (n=5/group) treated mice (at 20 weeks of age). For both accelerating and rocking test, AAV-ASM treated mice performed significantly (p < .001) better than ASMKO AAV2/1- β gal treated mice. Performance of mice injected with AAV2/1-ASM were indistinguishable from wild type mice in both the accelerating and rocking tests.

[043] **Figure 12** shows brain sphingomyelin levels (at 20 weeks of age) in ASMKO mice bilaterally injected with AAV1- β gal or with AAV serotypes 1 and 2

encoding for hASM. Brains were divided into 5 rostrocaudal sections (SI = most rostral and S5 = most caudal. An asterisk indicates that the data point is significantly different from ASMKO mice ($p<0.01$).

[044] **Figure 13A** illustrates the connections between the deep cerebellar nuclei regions (medial, interposed, and lateral) and the spinal cord regions (cervical, thoracic, lumbar, and sacral). **Figure 13B** illustrates the connections between the deep cerebellar nuclei regions (medial, interposed, and lateral) and the brainstem regions (midbrain, pons, and medulla). The connections are represented by arrows, which start at the cell body region of a neuron and end at the axon terminal region of the neuron. For example, the three regions of the DCN each have neurons with cell bodies that send axons that terminate in the cervical region of the spinal cord while the cervical region of the spinal cord has cell bodies that send axons that terminate in either the medial or interposed regions of the DCN.

[045] **Figure 14** illustrates green fluorescent protein distribution in the brainstem, or upper motor neurons, following DCN delivery of AAV encoding for green fluorescent protein (GFP).

[046] **Figure 15** illustrates green fluorescent protein distribution in the spinal cord regions following DCN delivery of AAV encoding for green fluorescent protein (GFP).

[047] **Figure 16** shows GFP distribution within the mouse brain following bilateral delivery of a GFP expressing AAV1 vector to the deep cerebellar nuclei (DCN). In addition to the DCN, GFP positive staining was also observed in the olfactory bulbs, cerebral cortex, thalamus, brainstem, cerebellar cortex and spinal cord. All of these areas either receive projections from and/or send projections to the DCN.

Detailed Description of the Invention

[048] In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

[049] The term "transgene" refers to a polynucleotide that is introduced into a cell of and is capable of being translated and/or expressed under appropriate conditions and confers a desired property to a cell into which it was introduced, or otherwise leads to a desired therapeutic outcome.

[050] The terms "genome particles (gp)," or "genome equivalents," as used in reference to a viral titer, refer to the number of virions containing the recombinant AAV DNA genome, regardless of infectivity or functionality. The number of genome particles in a particular vector preparation can be measured by procedures such as described in the Examples herein, or for example, in Clark et al. (1999) *Hum. Gene Ther.*, 10:1031-1039; Veldwijk et al. (2002) *Mol. Ther.*, 6:272-278.

[051] The terms "infection unit (iu)," "infectious particle," or "replication unit," as used in reference to a viral titer, refer to the number of infectious and replication-competent recombinant AAV vector particles as measured by the infectious center assay, also known as replication center assay, as described, for example, in McLaughlin et al. (1988) *J. Virol.*, 62:1963-1973.

[052] The term "transducing unit (tu)" as used in reference to a viral titer, refers to the number of infectious recombinant AAV vector particles that result in the production of a functional transgene product as measured in functional assays such as described in Examples herein, or for example, in Xiao et al. (1997) *Exp. Neurobiol.*, 144:113-124; or in Fisher et al. (1996) *J. Virol.*, 70:520-532 (LFU assay).

[053] The terms "therapeutic," "therapeutically effective amount," and their cognates refer to that amount of a compound that results in prevention or delay of onset or amelioration of symptoms of in a subject or an attainment of a desired biological outcome, such as correction of neuropathology, e.g., cellular pathology associated with a lysosomal storage disease such as that described herein or in Walkley (1998) *Brain Pathol.*, 8:175-193. The term "therapeutic correction" refers to that degree of correction that results in prevention or delay of onset or amelioration of symptoms in a subject. The effective amount can be determined by methods well-known in the art and as described in the subsequent sections.

Methods and Compositions

[054] ASMKO mice are an accepted model of types A and B Niemann-Pick disease (Horinouchi et al. (1995) *Nat. Genetics*, 10:288-293; Jin et al. (2002) *J. Clin. Invest.*, 109:1183-1191; and Otterbach (1995) *Cell*, 81:1053-1061). Niemann-Pick disease (NPD) is classified as a lysosomal storage disease and is an inherited neurometabolic disorder characterized by a genetic deficiency in acid sphingomyelinase (ASM; sphingomyelin cholinephosphohydrolase, EC 3.1.3.12). The lack of functional ASM protein results in the accumulation of sphingomyelin substrate within the lysosomes of neurons and glia throughout the brain. This leads to the formation of large numbers of distended lysosomes in the perikaryon, which are a hallmark feature and the primary cellular phenotype of type A NPD. The presence of distended lysosomes correlates with the loss of normal cellular function and a progressive neurodegenerative course that leads to death of the affected individual in early childhood (The Metabolic and Molecular Bases of Inherited Diseases, eds. Scriver et al., McGraw-Hill, New York, 2001, pp. 3589-3610). Secondary cellular phenotypes (e.g., additional metabolic abnormalities) are also associated with this disease, notably the high level accumulation of cholesterol in the lysosomal compartment. Sphingomyelin has strong affinity for cholesterol, which results in the sequestering of large amounts of cholesterol in the lysosomes of ASMKO mice and human patients (Leventhal et al. (2001) *J. Biol. Chem.*, 276:44976-44983; Slotte (1997) *Subcell. Biochem.*, 28:277-293; and Viana et al. (1990) *J. Med. Genet.*, 27:499-504.)

[055] The present invention is based, in part, on the discovery and demonstration that an intrahippocampal injection of high titer AAV-ASM into the diseased brains of ASMKO mice results in expression of ASM mRNA and protein distally from the injection site in a pattern consistent with the topographical organization of the projection neurons that innervate the injection site. In addition to robust expression at the site of injection, ASM mRNA and protein are also detected in several distal regions outside of the ipsilateral (injected) hippocampus, specifically, in the contralateral hippocampal dentate gyrus and CA3, and the medial septum and entorhinal cortex. The invention is further based, in part, on the discovery and demonstration of the extensive correction of lysosomal storage pathology at the distal sites thereby allowing a larger volume of correction via a smaller number of injection sites.

[056] Accordingly, in one aspect, the present invention provides methods for treating neurometabolic disorders in mammals. The populations treated by the methods of the invention include, but are not limited to, patients having or at risk for developing a neurometabolic disorder, e.g., a LSD, such as diseases listed in Table 1, particularly, if such a disease affects the CNS. In an illustrative embodiment, the disease is type A Niemann-Pick disease. In certain embodiments, neurometabolic disorders may exclude Alzheimer's, Parkinson, Huntington, Tay Sachs, Lesch-Nyan, and Creutzfeldt-Jakob diseases. However, methods of the invention utilizing a metalloendopeptidase as a therapeutic transgene, are specifically useful to the treatment of Alzheimer's disease and amyloid-related disorders.

[057] In some embodiments, the method of treating a neurometabolic disorder comprises administration of a high titer AAV vector carrying a therapeutic transgene so that the transgene product is expressed at a therapeutic level in a second site within the CNS distal to the first site. In some embodiments, the viral titer of the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In further embodiments, the administration is accomplished by direct intraparenchymal injection of a high titer AAV vector solution into the diseased brain, thereafter the transgene is

expressed distally, contralaterally or ipsilaterally, to the administration site at a therapeutic level at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm from the administration site.

[058] The distance between the first and the second sites is defined as the minimal distance region between the site of administration (first site) and the boundary of the detectable transduction of the distal site (second site) as measured using procedures known in the art or as described in the Examples, e.g., *in situ* hybridization. Some neurons in the CNS of larger mammals may span large distances by virtue of their axonal projections. For example, in humans, some axons may span a distance of 1000 mm or greater. Thus, in various methods of the invention, AAV can be axonally transported along the entire length of the axon at such a distance to reach and transduce the parent cell body.

[059] A site of vector administration within the CNS is chosen based on the desired target region of neuropathology and the topology of brain circuits involved so long as an administration site and the target region have axonal connections. The target region can be defined, for example, using 3-D stereotaxic coordinates. In some embodiments, the administration site is chosen so that at least 0.1, 0.5, 1, 5, or 10 % of the total amount of vector injected is delivered distally at the target region of at least 1, 200, 500, or 1000 mm³. An administration site may be localized in a region innervated by projection neurons connecting distal regions of the brain. For example, the substantia nigra and ventral tegmental area send dense projections to the caudate and putamen (collectively known as the striatum). Neurons within the substantia nigra and ventral tegmentum can be targeted for transduction by retrograde transport of AAV following injection into the striatum. As another example, the hippocampus receives well-defined, predictable axonal projections from other regions of the brain. Other administration sites may be localized, for example, in the spinal cord, brainstem (medulla and pons), mesencephalon, cerebellum (including the deep cerebellar nuclei), diencephalon (thalamus, hypothalamus), telencephalon (corpus striatum, cerebral cortex, or, within the cortex, the occipital, temporal, parietal or frontal lobes), or combinations thereof.

[060] For identification of structures in the human brain, see, e.g., *The Human Brain: Surface, Three-Dimensional Sectional Anatomy With MRI, and Blood Supply*, 2nd ed., eds. Deuteron et al., Springer Vela, 1999; *Atlas of the Human Brain*, eds. Mai et al., Academic Press; 1997; and *Co-Planar Sterotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*, eds. Tamarack et al., Thyme Medical Pub., 1988. For identification of structures in the mouse brain, see, e.g., *The Mouse Brain in Sterotaxic Coordinates*, 2nd ed., Academic Press, 2000. If desired, the human brain structure can be correlated to similar structures in the brain of another mammal. For example, most mammals, including humans and rodents, show a similar topographical organization of the entorhinal-hippocampus projections, with neurons in the lateral part of both the lateral and medial entorhinal cortex projecting to the dorsal part or septal pole of the hippocampus, whereas the projection to the ventral hippocampus originates primarily from neurons in medial parts of the entorhinal cortex (*Principles of Neural Science*, 4th ed., eds Kandel et al., McGraw-Hill, 1991; *The Rat Nervous System*, 2nd ed., ed. Paxinos, Academic Press, 1995). Furthermore, layer II cells of the entorhinal cortex project to the dentate gyrus, and they terminate in the outer two-thirds of the molecular layer of the dentate gyrus. The axons from layer III cells project bilaterally to the cornu ammonis areas CA1 and CA3 of the hippocampus, terminating in the stratum lacunose molecular layer.

[061] The second (target) site can be located any region of the CNS, including the brain and the spinal cord, that contains a neurons that project to the first (administration) site. In some embodiments, the second site is in a region of the CNS chosen from the substantia nigra, the medulla oblongata, or the spinal cord.

[062] To deliver the vector specifically to a particular region of the central nervous system, especially to a particular region of the brain, it may be administered by stereotaxic microinjection. For example, on the day of surgery, patients will have the stereotaxic frame base fixed in place (screwed into the skull). The brain with stereotaxic frame base (MRI-compatible with fiduciary markings) will be imaged using high resolution MRI. The MRI images will then be

transferred to a computer that runs stereotaxic software. A series of coronal, sagittal and axial images will be used to determine the target site of vector injection, and trajectory. The software directly translates the trajectory into 3-dimensional coordinates appropriate for the stereotaxic frame. Burr holes are drilled above the entry site and the stereotaxic apparatus localized with the needle implanted at the given depth. The vector in a pharmaceutically acceptable carrier will then be injected. The AAV vector is then administrated by direct injection to the primary target site and retrogradely transported to distal target sites via axons. Additional routes of administration may be used, e.g., superficial cortical application under direct visualization, or other non-stereotaxic application.

[063] The total volume of material to be administered, and the total number of vector particles to be administered, will be determined by those skilled in the art based upon known aspects of gene therapy. Therapeutic effectiveness and safety can be tested in an appropriate animal model. For example, a variety of well-characterized animal models exist for LSDs, e.g., as described herein or in Watson et al. (2001) *Methods Mol. Med.*, 76:383-403; or Jeyakumar et al. (2002) *Neuropath. Appl. Neurobiol.*, 28:343-357.

[064] In experimental mice, the total volume of injected AAV solution is for example, between 1 to 5 μ l. For other mammals, including the human brain, volumes and delivery rates are appropriately scaled. For example, it has been demonstrated that volumes of 150 μ l can be safely injected in the primate brain (Janson et al. (2002) *Hum. Gene Ther.*, 13:1391-1412). Treatment may consist of a single injection per target site, or may be repeated along the injection tract, if necessary. Multiple injection sites can be used. For example, in some embodiments, in addition to the first administration site, a composition comprising AAV carrying a transgene is administered to another site which can be contralateral or ipsilateral to the first administration site.

[065] In another aspect, the invention provides a method of delivering a recombinant AAV genome via retrograde axonal transport to the nucleus of a disease-compromised neuron *in vivo*. In some embodiments, the cellular

pathology exhibited by a neuron is that of a LSD such as listed in Table 1 (see, e.g., Walkley (1998) *Brain Pathol.*, 8:175-193). In an illustrative embodiment, the disease is Niemann-Pick A disease. The method of delivering a recombinant AAV genome to the nucleus of a disease-compromised neuron comprises contacting an axonal ending of a disease-compromised neuron with a composition comprising an AAV viral particle comprising the recombinant AAV genome and allowing the viral particle to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron, wherein the concentration of AAV genomes in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). In certain embodiments, the neuron is a projection neuron and/or the distance from the axonal ending to the nucleus of the neuron is at least 2, 3, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mm. In various embodiments, the AAV genome is transported along the entire length of the axon, at distances varying depending on the axon length. In humans, these distances may be as much as 1000 mm or greater.

[066] In another aspect, the invention provides a method of delivering a transgene product to a target cell of the CNS, which is a neuron or a glial cell, in a mammal afflicted with a disorder, for example an LSD as listed in Table 1. The method comprises contacting an axonal ending of a neuron with a composition comprising an AAV vector carrying at least a part of a gene encoding a therapeutic transgene product; allowing the viral particles to be endocytosed and retrogradely transported intracellularly along the axon to the nucleus of the neuron; allowing the transgene product to be expressed and secreted by the neuron; and allowing a second cell to uptake the transgene product, wherein the transgene product thereby alleviates pathology in the second cell. In some embodiments the concentration of the AAV vector in the composition is at least: (a) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{12}$ gp/ml); (b) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^9$ tu/ml); or (c) 5, 6, 7, 8, 8.4, 9, 9.3, 10, 15, 20, 25, or 50 ($\times 10^{10}$ iu/ml). For example, lysosomal hydrolases can be secreted by transduced cells and subsequently taken up by another cell via mannose-6-phosphate receptor-mediated endocytosis, the second cell being

transduced or non-transduced (Sando et al. (1977) *Cell*, 12:619-627; Taylor et al. (1997) *Nat. Med.*, 3:771-774; Miranda et al. (2000) *Gene Ther.*, 7:1768-1776; and Jin et al. (2002) *J. Clin. Invest.*, 109:1183-1191).

[067] In the methods of the invention, AAV of any serotype can be used so long as the vector is capable of undergoing retrograde axonal transport in a disease-compromised brain. The serotype of the viral vector used in certain embodiments of the invention is selected from the group consisting from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, and AAV8 (see, e.g., Gao et al. (2002) *PNAS*, 99:11854-11859; and *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003). Other serotype besides those listed herein can be used. Furthermore, pseudotyped AAV vectors may also be utilized in the methods described herein. Pseudotyped AAV vectors are those which contain the genome of one AAV serotype in the capsid of a second AAV serotype; for example, an AAV vector that contains the AAV2 capsid and the AAV1 genome or an AAV vector that contains the AAV5 capsid and the AAV 2 genome. (Auricchio et al., (2001) *Hum. Mol. Genet.*, 10(26):3075-81.) However, AAV5 may be specifically excluded from the methods of the invention utilizing a metalloendopeptidase, e.g., neprilysin, as a therapeutic transgene.

[068] AAV vectors are derived from single-stranded (ss) DNA parvoviruses that are nonpathogenic for mammals (reviewed in Muzyscka (1992) *Curr. Top. Microb. Immunol.*, 158:97-129). Briefly, AAV-based vectors have the rep and cap viral genes that account for 96% of the viral genome removed, leaving the two flanking 145-basepair (bp) inverted terminal repeats (ITRs), which are used to initiate viral DNA replication, packaging and integration. In the absence of helper virus, wild-type AAV integrates into the human host-cell genome with preferential site-specificity at chromosome 19q 13.3 or it may remain expressed episomally. A single AAV particle can accommodate up to 5 kb of ssDNA, therefore leaving about 4.5 kb for a transgene and regulatory elements, which is typically sufficient. However, trans-splicing systems as described, for example, in United States Patent No. 6,544,785, may nearly double this limit.

[069] In an illustrative embodiment, AAV is AAV2 or AAV1. Adeno-associated virus of many serotypes, especially AAV2, have been extensively studied and characterized as gene therapy vectors. Those skilled in the art will be familiar with the preparation of functional AAV-based gene therapy vectors. Numerous references to various methods of AAV production, purification and preparation for administration to human subjects can be found in the extensive body of published literature (see, e.g., *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003). Additionally, AAV-based gene therapy targeted to cells of the CNS has been described in United States Patent Nos. 6,180,613 and 6,503,888.

[070] In certain methods of the invention, the vector comprises a transgene operably linked to a promoter. The transgene encodes a biologically active molecule, expression of which in the CNS results in at least partial correction of neuropathology. In some embodiments, the transgene encodes a lysosomal hydrolase. In an illustrative embodiment, the lysosomal hydrolase is ASM. The genomic and functional cDNA sequences of human ASM have been published (see, e.g., United States Patent No. 5,773,278 and 6,541,218). Other lysosomal hydrolases can be used for appropriate diseases, for example, as listed in Table 1.

[071] The invention further provides methods of treating Alzheimer's disease in mammals, including humans. In such methods, the transgene encodes a metalloendopeptidase. The metalloendopeptidase can be, for example, the amyloid-beta degrading enzyme neprilysin (EC 3.4.24.11; sequence accession number, e.g., P08473 (SWISS-PROT)), the insulin-degrading enzyme insulin (EC 3.4.24.56; sequence accession number, e.g., P14735 (SWISS-PROT)), or thimet oligopeptidase (EC 3.4.24.15; sequence accession number, e.g., P52888 (SWISS-PROT)).

[072] The level of transgene expression in eukaryotic cells is largely determined by the transcriptional promoter within the transgene expression cassette. Promoters that show long-term activity and are tissue- and even cell-specific are used in some embodiments. Nonlimiting examples of promoters include, but are

not limited to, the cytomegalovirus (CMV) promoter (Kaplitt et al. (1994) *Nat. Genet.*, 8:148-154), CMV/human β 3-globin promoter (Mandel et al. (1998) *J. Neurosci.*, 18:4271-4284), GFAP promoter (Xu et al. (2001) *Gene Ther.*, 8:1323-1332), the 1.8-kb neuron-specific enolase (NSE) promoter (Klein et al. (1998) *Exp. Neurol.*, 150:183-194), chicken beta actin (CBA) promoter (Miyazaki (1989) *Gene*, 79:269-277) and the β -glucuronidase (GUSB) promoter (Shipley et al. (1991) *Genetics*, 10:1009-1018). To prolong expression, other regulatory elements may additionally be operably linked to the transgene, such as, e.g., the Woodchuck Hepatitis Virus Post-Regulatory Element (WPRE) (Donello et al. (1998) *J. Virol.*, 72, 5085-5092) or the bovine growth hormone (BGH) polyadenylation site.

[073] For some CNS gene therapy applications, it will be necessary to control transcriptional activity. To this end, pharmacological regulation of gene expression with AAV vectors can be obtained by including various regulatory elements and drug-responsive promoters as described, for example, in Habermaet al. (1998) *Gene Ther.*, 5:1604-16011; and Ye et al. (1995) *Science*, 283:88-91.

[074] High titer AAV preparations can be produced using techniques known in the art, e.g., as described in United States Patent No. 5,658,776 and *Viral Vectors for Gene Therapy: Methods and Protocols*, ed. Machida, Humana Press, 2003.

[075] The following examples provide illustrative embodiments of the invention. One of ordinary skill in the art will recognize the numerous modifications and variations that may be performed without altering the spirit or scope of the present invention. Such modifications and variations are encompassed within the scope of the invention. The examples do not in any way limit the invention.

ExamplesTitration of Recombinant Vectors

[076] AAV vector titers were measured according to genome copy number (genome particles per milliliter). Genome particle concentrations were based on Taqman® PCR of the vector DNA as previously reported (Clark et al. (1999) Hum. Gene Ther., 10:1031-1039; Veldwijk et al. (2002) Mol. Ther., 6:272-278). Briefly, purified AAV-ASM was treated with capsid digestion buffer (50mM Tris-HCl pH 8.0, 1.0 mM EDTA, 0.5% SDS, 1.0 mg/ml proteinase K) at 50°C for 1 hour to release vector DNA. DNA samples were put through a polymerase chain reaction (PCR) with primers that anneal to specific sequences in the vector DNA, such as the promoter region, transgene, or the poly A sequence. The PCR results were then quantified by a Real-time Taqman® software, such as that provided by the Perkin Elmer-Applied Biosystems (Foster City, CA) Prism 7700 Sequence Detector System.

[077] Vectors carrying an assayable marker gene such as the β -galactosidase or green fluorescent protein gene (GFP) can be titered using an infectivity assay. Susceptible cells (e.g., HeLa, or COS cells) are transduced with the AAV and an assay is performed to determine gene expression such as staining of β -galactosidase vector-transduced cells with X-gal (5-bromo-4chloro-3-indolyl- β -D-galactopyranoside) or fluorescence microscopy for GFP-transduced cells. For example, the assay is performed as follows: 4×10^4 HeLa cells are plated in each well of a 24-well culture plate using normal growth media. After attachment, i.e., about 24 hours later, the cells are infected with Ad type 5 at a multiplicity of infection (MOI) of 10 and transduced with serial dilutions of the packaged vector and incubated at 37°C. One to three days later, before extensive cytopathic effects are observed, the appropriate assay is performed on the cells (e.g., X-gal staining or fluorescence microscopy). If a reporter gene

such as β -galactosidase is used, the cells are fixed in 2% paraformaldehyde, 0.5% glutaraldehyde and stained for β -galactosidase activity using X-gal. Vector dilutions that give well-separated cells are counted. Each positive cell represents 1 transduction unit (tu) of vector.

Correction of LSD pathology in the mouse brain

[078] Ten-week old ASMKO mice contain significant NPD pathology in the central nervous system. Identification of homozygous recessive mutants was verified by PCR. Sixteen 10-week old ASMKO mice were anesthetized with isoflurane and mounted on a stereotaxic frame, an incision was made to expose the underlying skull, and a single drill hole was made over one hemisphere of each mouse. Two microliters of high titer (9.3×10^{12} gp/ml) AAV2-CMV-ASM (Targeted Genetics, Seattle, WA) were injected into the hippocampus at a final stereotaxic coordinate of 2.0 mm rostral of bregma, 1.5 mm right of midline, and 2.0 mm ventral to pial surface. This hippocampal coordinates ensured that the AAV2 vector was exposed to neurons of the dentate gyrus and of the cornu ammonis area 3 (CA3), as well as to axonal endings of projection neurons of the contralateral hippocampus, medial septum and entorhinal cortex. The injections were performed at a rate of 0.2 μ l/minute, and a total of 1.86×10^{10} genomic particles were administered into each brain.

[079] Mice were tested by accelerating and rocking rotarod for motor function on the Smartrod (AccuScan) using methods known in the art and reproduced in Sleat et al. (2004) *J. Neurosci.* **24**:9117-9126. Figures 10A/B and Figures 11A/B graphically show the results of rotarod tests as a measurement of recovery of motor function.

[080] The mice were then sacrificed at either 5 (n = 8) or 15 (n = 8) weeks post injection (pi). Eight brains (n = 4 each at 5 and 15 weeks pi) were analyzed for ASM mRNA and protein distribution, and for the reduction of the supraphysiologic levels of cholesterol in the lysosomes. The remaining 8 brains (n=4 each at 5 and 15 weeks pi) were processed for histopathology to analyze the correction of

accumulated and distended lysosomes, which is the most direct and accurate method for determining reversal of storage pathology for LSDs.

[081] Robust transduction was detected in the injected (ipsilateral) hippocampus at 5 and 15 weeks pi. The granule cell layer and hilus of the dentate gyrus, and the pyramidal and oriens cell layers of CA3 were extensively transduced by the AAV2 vector. This impressive pattern of transduction extended to other regions of the ipsilateral hippocampus, such as the subiculum and cornu ammonis area 1 (CA1) and 2 (CA2). Immunofluorescence with an anti-human ASM monoclonal antibody confirmed the presence of ASM protein in many cells. The overall protein pattern was similar to the mRNA pattern, with some additional localized spread of protein.

[082] Human ASM mRNA and protein were also detected in regions outside of the ipsilateral hippocampus at both time points. The contralateral dentate gyrus and CA3, and the medial septum and entorhinal cortex were positive for in situ hybridization and immunofluorescence (Figures 1A and 2A). The pattern of transduction at these distal sites was consistent with the topographical organization of the projection neurons that innervate the injection site (Figures 1B and 2B). This demonstrated that AAV2 underwent retrograde axonal transport in the intrahippocampal, septohippocampal and entorhinodentate circuits of ASMKO brains, and that the viral vector was targeted to the nucleus following transport up the axons (Figures 1C and 2C). The pattern of transduction does not support parenchymal diffusion as the reason for AAV2 transport to these distal sites. If such diffusion had occurred, structures between the injected and distal sites would have been exposed to migrating virus. But these intermediate structures were negative for in situ hybridization. For example the striatum, which possesses a strong natural tropism for AAV2, was negative for gene transfer despite being in the direct path between the hippocampus and medial septum. Thus, gene transfer to the distal sites must have arisen by retrograde axonal transport, which indicates that an affected projection neuron can function as an effective highway transport system for AAV2 movement through a diseased brain.

[083] The ability of ASM to reverse the cholesterol abnormalities in the ASMKO brain additionally was investigated. Filipin is an autofluorescent molecule isolated from *Streptomyces filipinensis* that binds to cholesterol complexes (Leventhal et al. (2001) J. Biol. Chem., 276:44976-44983; and Sarna et al. (2001) Eur. J. Neurosci., 13:1-9). Uninjected ASMKO brains had high levels of filipin staining due to these abundant cholesterol complexes, whereas normal mouse brains produced no filipin staining.

[084] Injection of AAV2-CMV-ASM resulted in the complete loss of filipin staining throughout the entire ipsilateral and contralateral hippocampus, septum and entorhinal cortex at 5 and 15 weeks pi of ASMKO mice (Figures 1A and 2A). This was in stark contrast to uninjected age-matched ASMKO controls, where high levels of filipin staining were detected in these same structures. The loss of filipin staining in AAV2-injected brains demonstrates that a secondary cellular phenotype (e.g. metabolic defect) of ASM disease was corrected. This strongly suggests that ASM protein was targeted to the lysosome and interacted with the sphingomyelin-cholesterol complex. This interaction likely resulted in the release of cholesterol from sphingomyelin, and subsequent entry of cholesterol into its normal biological pathways (such as degradation or translocation to the plasma membrane (Leventhal et al. (2001) J. Biol. Chem., 276:44976-44983)) .

[085] The loss of filipin staining was observed in all cell layers and subfields of the intrahippocampal, septohippocampal and entorhinodentate circuits. The area of cholesterol correction was far greater and more extensive than the ASM protein pattern. This indicates that, following retrograde axonal transport of AAV2, projection neurons may have functioned as "enzyme pumps" and secreted ASM protein into the surrounding tissue. Significantly, only a small amount of ASM is needed to have a therapeutic effect on cholesterol accumulation within ASMKO-affected cells, an amount below the detection limit of the immunofluorescent protocol.

[086] Whether axonal transport of the AAV2-ASM vector results in the correction of the primary cellular phenotype of NPD also was evaluated. One-micron-thick histopathology brain sections demonstrated a remarkable reduction of accumulated and distended lysosomal pathology in AAV2-CMV-ASM-injected

brains at 5 weeks pi (Table 2). Reversal of pathology resulting in partial or complete restoration of normal cellular architecture occurred in all regions of the ipsilateral and the contralateral hemispheres of the hippocampus. The medial septum and the entorhinal cortex also showed a substantial reduction in storage lesions. Similar to the filipin data, the number of cells corrected was greater and more widespread than the ASM protein pattern. Reversal of pathology was evident within regions known to project to the hippocampus including the intrahippocampal, septohippocampal and entorhinodentate circuits. Overall, the volume of correction was 30-35 mm³ or more in the contralateral hippocampus, 5-8 mm³ or more in the ipsilateral entorhinal cortex, 1-2 mm³ or more in the contralateral entorhinal cortex, and 2-3 mm³ or more in the medial septum. This further supports that axonal transport of the viral vector was responsible for this therapeutic effect, because nearby structures (that do not contribute to these circuits) would have been corrected if viral distribution was mediated merely by diffusion (see, "ipsilateral striatum" and "contralateral striatum" in Table 2).

[087] To demonstrate that reversal of pathology was specific to ASM, an additional group of ASMKO mice was injected with a control vector carrying a reporter gene, AAV2-CMV-β-gal (n = 2 each at 5 and 15 weeks pi), and processed for histopathology. In all four brains, cells remained inundated with distended lysosomes, and contained other abnormalities such as cytoplasmic swelling and disorganized cellular layers.

Table 2

<i>Brain Region</i>	<i>Untreated</i>	<i>AAV2-ASM Treated</i>
Ipsilateral Hippocampus		
CA1 Field	++++	+
CA3 Field	++++	+
Dentate Granule Cell Layer	++++	+
Hilus	++++	+
Contralateral Hippocampus		
CA1 Field	++++	+
CA3 Field	++++	+
Dentate Granule Cell Layer	++++	+
Hilus	++++	+
Ipsilateral Entorhinal Cortex	++++	++
Contralateral Entorhinal Cortex	++++	+++
Ipsilateral Medial Septum	++++	++
Contralateral Medial Septum	++++	++
Ipsilateral Striatum	++++	++++
Contralateral Striatum	++++	++++

++++ high level of pathology in virtually all cells

+++ pathology in many cells, correction is observed in some cells

++ pathology in some cells, correction is observed in many cells

+ little or no pathology is in most cells, virtually every cell is corrected

[088] Thus, in accordance with the present invention, a single injection of high titer AAV2 vector is sufficient to transfer the ASM gene to structures that innervate the ASMKO affected hippocampus. The number of structures positive for AAV2 vector was greater than that demonstrated by a recent study in the normal rat hippocampus, which showed axonal transport only in the

entorhinodentate circuit (Kaspar et al. (2002) Mol. Ther., 5:50-56). The results described herein demonstrate that axonal transport can occur in projection neurons inflicted with storage pathology, and that this mode of transport results in the clearance of storage pathology in proximal structures and multiple regions distal to the injection site. We also demonstrate that axonal transport is not limited to only those circuits associated with the hippocampus. Retrograde axonal transport occurred in the nigrostriatal (Figure 3) and in the medullocerebellar (Figure 4) circuits. This demonstrates that axonal transport of AAV2 in diseased-compromised neurons is a general property of the viral vector.

[089] A similar study was performed with AAV1-ASM at the concentrations of 1- 4×10^{13} gp/ml and AAV7-ASM at the concentration of 8.4×10^{12} gp/ml. While AAV1 did not exhibit detectable retrograde axonal transport, AAV7 did undergo retrograde axonal transport, similar to AAV2, and produced correction of LSD pathology in distal regions (see Figure 5).

Injection of AAV into the Cerebellum

[090] ASMKO mice were anesthetized with isoflurane and mounted on a stereotaxic frame. Bregma was located as a reference point to determine the drilling location for injection into the deep cerebellar nuclei region of the cerebellum. Once located, an incision was made to expose the underlying skull, and a single drill hole was made into the skull without piercing the brain surface. A Hamilton syringe was lowered into the brain via the hole and AAV2-CMV-ASM was injected into the deep cerebellar nuclei at a rate of 0.5 microliters per second. Three microliters were injected for a total dose of 1×10^{10} genome particles. Mice were sacrificed 7 weeks post injection. The brains and spinal cords were evaluated for ASM mRNA expression, ASM protein expression, filipin staining, and calbindin staining.

[091] Filipin is an autofluorescent molecule that binds to cholesterol complexes. Untreated ASMKO mice have high levels of filipin staining due to abundant cholesterol complexes, which accumulate as a result of their disease. In contrast, normal mouse brains do not exhibit filipin staining.

[092] Calbindin is a marker of Purkinje cells, which are found in the cerebellum and are involved in coordinated movements. In the ASMKO mouse, Purkinje cells die off in these mice as they age, resulting in decreased coordinated movement behavior. This loss of Purkinje cells and the correlative loss of coordinated movement behavior are not observed in normal mice.

[093] Following AAV2 injection into the deep cerebellar nucleus, the cerebellum was positive for ASM mRNA, ASM protein, and calbindin staining. These results demonstrate that ability of AAV2 to transduce the cerebellum following injection into the deep cerebellar nuclei. Moreover, the cerebellar transduction and resulting ASM expression prevented Purkinje cell death as evidenced by the presence of calbindin staining in the treated mice. In AAV-ASM treated mice, expression of hASM protein was also observed throughout the brainstem, thalamus, and mescencephalon. hASM protein expression in these regions overlapped with regional clearance of filipin/cholesterol staining. Overall, in the cerebellum there was a positive relationship between ASM protein levels, filipin clearance, and Purkinje cell survival.

[094] ASM mRNA and ASM protein were also detected outside the cerebellum. Specifically, the spinal cord was positive for ASM mRNA expression as evidenced by *in situ* hybridization. The spinal cord was also positive for ASM protein as evidenced by ASM-specific immunofluorescence. These results indicate that the spinal cord was transduced following a distal injection of the AAV vector into the deep cerebellar nuclei. This pattern of transduction was consistent with the topographical organization of the projection neurons that innervate the deep cerebellar nuclei region. These results indicate that the AAV2 vector was taken up by distal spinal cord cells and expressed.

Treatment of Alzheimer's disease

[095] Alzheimer's disease (AD) is a neurodegenerative disorder characterized by the accumulation of amyloid β -peptide (A β) due to decreased A β catabolism. As A β accumulates, it aggregates into extracellular plaques, causing impairment

of synaptic function and loss of neurons. The pathology leads to dementia, loss of coordination, and death.

[096] Neprilysin is a 97 kD membrane-bound zinc metalloendopeptidase that is the rate-limiting enzyme in the normal degradation of A β . Introduction of neprilysin may decelerate the progression of the disease by removing A β pools before aggregation. Indeed, neprilysin was shown to degrade oligomeric forms of A β thereby removing existing plaques in an animal model of AD (Kanemitsu et al. (2003) *Neurosci. Lett.*, 350:113-116). Neprilysin knockout mice exhibit high levels of A β (Iwata et al. (2001) *J. Neurosci.*, 24:991-998.). Neprilysin inhibitors, such as thiorphan and phosphoramidon, increase A β levels in mouse brain (Iwata et al. (2000) *Nat. Med.*, 6:143-150). Additionally, decreased neprilysin mRNA levels were found in areas of high amyloid plaque burden in human brains, further demonstrating the link between neprilysin and AD (Yasojima et al. (2001) *Neurosci. Lett.*, 297:97-100).

[097] The areas of brain most affected by AD are the hippocampus, cortex, cerebellum, striatum and thalamus (see, e.g., Iwata et al. (2001) *supra*; Yasojima et al. (2001) *supra*). These are the same areas of the brain that show efficient retrograde axonal transport with AAV.

[098] Accordingly, AAV can be used to deliver therapeutic transgenes to regions of high plaque burden by direct injection and subsequent translocation of virus through brain circuits to our target sites. Viral vector-mediated gene transfer of neprilysin was effective in treating mouse models of AD (Marr et al. (2003) *J. Neurosci.*, 23:1992-1996; Marr et al. (2004) *J. Mol. Neurosci.*, 22:5-11; Iwata et al. (2004) *J. Neurosci.*, 24:991-998). A recent report showed that AAV5-neprilysin removed A β from the pre-synaptic terminals of the hippocampus in neprilysin-deficient mice, decelerating plaque formation at the synapses (Iwata et al. (2004) *supra*). In this report, neprilysin was found in the contralateral hippocampus but whether this is attributable to retrograde transport of virus or anterograde transport of expressed protein remains unknown.

Correction of Cholesterol Storage Pathology

[099] The following experiment, evaluated the relative ability of recombinant AAV2/1, AAV2/2, AAV2/5, AAV2/7 and AAV2/8 serotype vectors encoding human ASM (hASM) to express hASM protein, correct cholesterol storage pathology, undergo transport, rescue Purkinje cells, and initiate functional recovery in the ASMKO mouse after unilateral injection within the deep cerebellar nuclei. An additional group of ASMKO mice received bilateral injections into the DCN in order to assess whether increased transgene protein spread/expression would improve behavioral functional recovery.

[0100] Sixty-six male homozygous (-/-) acid sphingomyelinase knockout (ASMKO) mice and sixteen male wild type littermate controls were bred from heterozygote matings (+/-). Mice were genotyped by PCR following the procedure described in Gal et al. (1975) *N Engl J Med*:**293**:632-636. Mice from the original colony were backcrossed onto the C57/Bl6 strain. Animals were housed under 12:12 hour light:dark cycle and provided with food and water *ad libitum*. All procedures were performed under a protocol approved by the Institutional Animal Care and Use Committee.

[0101] After being anesthetized with isoflurane, mice (~7 weeks of age) were unilaterally injected into the deep cerebellar nuclei (DCN) (A-P: -5.75 from bregma, M-L: -1.8 from bregma, D-V: -2.6 from dura, incisor bar: 0.0) with one of the following AAV serotype vectors (n=8/vector): AAV1-CMV- β gal, AAV1-CMV-ASM, AAV2-CMV-ASM, AAV5-CMV-ASM, AAV7-CMV-ASM, and AAV8-CMV-ASM. Vectors were delivered with a 10 μ l Hamilton syringe mounted on a syringe pump at a rate of 0.5 μ l/minute for a total of 1.86×10^{10} genome particles per brain. The final injection volume for each vector was 4 μ l. One hour before and twenty-four hours after surgery mice were given ketoprofen (5 mg/kg; SC) for analgesia. Mice were killed 7 weeks post-injection (14 weeks of age). At the time of sacrifice mice were overdosed with euthasol (150 mg/kg; IP) and rapidly decapitated (n=5/group) or transcardially perfused (n=3/group). Brains from decapitated mice were rapidly removed, snap frozen in liquid nitrogen, dissected into 3 sections (right cerebral hemisphere, left cerebral hemisphere, & cerebellum) homogenized, and analyzed for hASM by ELISA. Brains and spinal

cords from perfused mice were processed for human ASM protein expression, cholesterol accumulation as detected by filipin staining, and Purkinje cell survival with calbindin staining on 50 μ m vibratome sections.

[0102] Using a similar protocol, ASMKO mice (~7 weeks of age) were injected bilaterally with AAV2/1- β gal (n=8), AAV2/1-ASM (n=5), and AAV2/2-ASM (n=5) and sacrificed at 20 weeks of age after undergoing rotarod testing. Brains were snap frozen in liquid nitrogen, bisected at midline and then divided into five sections (S1, S2, S3, S4, and S5) using a mouse brain matrix (ASI Instruments, Inc, MI.) Sections 1-4 were approximately 2 mm apart with S1 being the most rostral and S4 the most caudal. Section 5 contained the cerebellum only. The right hemisphere was used to quantify brain spingomyelin levels and the left hASM levels.

[0103] The full-length human ASM cDNA under the control of the human cytomegalovirus immediate-early (CMV) promoter, with an SV40 polyadenylation sequence, and a hybrid intron, was cloned into a plasmid containing ITRs from AAV serotype 2 (AAV2 ITR). Jin et al. (2002) *J Clin Invest.* **109**:1183-1191. Hybrid vectors were produced by triple transfection using a series of helper plasmids containing serotype specific capsid coding domains in addition to the AAV type 2 replication genes. This strategy allows the packaging of AAV2 ITR vectors into each serotype-specific virion Rabinowitz, et al. (2002) *J Virol.* **76**:791-801. With this approach the hASM recombinant genome was used to generate a series of rAAV-hASM vectors of various serotypes including AAV2/1, AAV2/2, AAV2/5, AAV2/7 and AAV2/8. Recombinant AAV vectors were purified by ion-exchange chromatography (Serotypes 2/1, 2/2 and 2/5). O'Riordan, et al. (2000) *J Gene Med* **2**: 444-54 or CsCl centrifugation (serotypes 2/8 and 2/7) Rabinowitz et al. (2002) *J. Virol.* **76**:791-801. The final titer of AAV-ASM virion particles (DNase-resistant particles), was determined by TaqMan PCR of the CMV sequence. Clark et al. (1999) *Hum. Gene Therapy* **10**:1031-1039.

[0104] Human ASM antibodies are human specific and do not cross react with mouse ASM. Coster (Corning, NY) 9018 plates coated (100 μ l/well) with monoclonal recombinant human ASM (rhASM) antibody (2 μ g/ml) diluted in 50

mM sodium carbonate buffer (pH 9.6) were incubated overnight @ 2-8°C. Excess coating antibody was removed and blocking diluent (KPL, Inc., MD) was added for 1h @ 37°C. Plates were washed with a microplate washer (Molecular Devices, CA) for two cycles. Standards, controls and samples diluted in standard dilution buffer (PBS, 0.05% Tween, 1% HP-BSA) were pipetted in duplicate and allowed to incubate for 1 h @ 37°C. Plates were washed as described above. One hundred microliters of biotinylated recombinant human ASM (rhASM) antibody (diluted 1:20K in standard dilution buffer) was added to each well, allowed to incubate for 1 h @ 37°C, and then removed with a microplate washer. Streptavidin – HRP (Pierce Biotechnology, Inc., IL) diluted 1:10K was then added (100 µl/well) and allowed to incubate for 30 min at room temperature. Plates were washed as described above and then incubated with SureBlue TMB (KPL, Inc., MD) for 15 minutes @ 36-38°C. The reaction was stopped with stop solution (KPL, Inc., MD) and absorbance values were then read at 450 nm with a Spectra Max 340 plate reader (Molecular Devices, CA). Data analysis was completed using Softmax Pro 4.3 software (Molecular Devices, CA).

[0105] The protein concentration for each sample was determined with a BCA protein assay kit (Pierce Biotechnology, Inc., IL) using bovine serum albumin as standard.

[0106] Mice were transcardially perfused with fixative containing 2% paraformaldehyde, 0.03% glutaraldehyde, 0.002% CaCl₂ in 0.1 M sodium acetate buffer at pH 6.5, followed by perfusion with the same fixative at pH 8.5. Mouse brains and spinal cords were dissected and post-fixed overnight at 4° C in pH 8.5 fixative without glutaraldehyde. The tissues were washed in 0.1 M potassium phosphate buffer, pH 7.4, embedded in 3.5% agar and cut into 50 µm sagittal sections with a vibratome.

[0107] Brains and spinal cords were vibratome-sectioned sagittally at 50 µm intervals. Sections were processed for immunofluorescence with primary antibodies against human ASM (1:200). Sections were incubated in 10% donkey serum, 0.3% Triton X-100 in PBS for 1 hour, followed by incubation with biotinylated mouse anti-human ASM in 2% donkey serum, 0.2% Triton X-100 in

PBS for 72 hours. After washing, the signal was amplified using a Tyramide Signal Amplification kit (PerkinElmer, Boston MA). Human ASM protein was visualized with a Nikon fluorescent microscope, and images were captured with a SPOT camera and Adobe Photoshop software.

[0108] Filipin Complex (Sigma, St. Louis, MO) was first diluted in 100% methanol for a stock concentration of 1 mg/ml. Stock solution is stable for 4 weeks at -20° C. After washing with PBS, the sections were incubated in the dark for three hours in a freshly made 10 µg/ml filipin solution in PBS. Sections were then washed three times with PBS. Cholesterol deposits were visualized under an ultraviolet filter on a fluorescence microscope.

[0109] Brains were processed for immunofluorescence using primary antibodies directed against the calcium binding protein, calbindin. Sections were washed with potassium phosphate buffer (KPB) and then rinsed with potassium phosphate buffered saline (KPBS). Sections were then blocked in 5% donkey serum, 0.25% Triton X-100 in KPBS for up to 3 hours and then incubated in 5% donkey serum, 0.2% Triton X-100 and mouse anti-calbindin (1:2500, Sigma, St. Louis, MO) in KPBS. After 72 hours at 4° C sections were rinsed with KPBS with 0.1% Triton X-100 three times. Secondary antibody, donkey-anti mouse CY3 (1:333, Jackson Immunoresearch Laboratories, West Grove, PA) was added in KPBS + 0.1% Triton X-100 for 90 minutes at room temperature. Sections were washed with KPB and then mounted onto gel-coated slides. Calbindin-positive cells were visualized under epifluorescence. In order to quantify Purkinje cells of the cerebellum, four full-faced, medial cerebellar sections were selected from each animal. Calbindin-immunopositive Purkinje cells were viewed under a fluorescent microscope and cell bodies were counted at a magnification of 20X. Each lobe was counted separately. Two separate focal planes were counted per lobe. Only cells in focus were counted to insure that no cell was counted twice.

[0110] Fifty (50) µm vibratome sections were first processed for immunofluorescence with antibodies directed against human ASM, as described above. The sections were then washed in PBS and stained for choline acetyltransferase (ChAT; rabbit polyclonal, 1:500, Chemicon International,

Temecula, CA) with the protocol outlined above for calbindin. Rather than using a CY3 secondary antibody, however, donkey-anti-rabbit FITC (1:200, Jackson Immunoresearch Laboratories, West Grove, PA) was used. The staining was first visualized under epifluorescence and later images were acquired with a confocal microscope.

[0111] Filipin staining was quantified as follows. Exposure-matched images were captured using a Nikon E600 wide field upright epifluorescence microscope equipped with a SPOT digital camera. The AAV2/1- β -gal group was imaged first, and that exposure was used to acquire all additional images. Each image analyzed represents a medial sagittal plane through the length of each half-brain. Morphometric analysis was performed with Metamorph software (Universal Imaging Corporation). The AAV2/1- β -gal images were thresholded; once established, the same threshold was used in all images. The following regions were manually selected by the user and analyzed separately: cerebellum, pons, medulla, midbrain, cerebral cortex, hippocampus, thalamus, hypothalamus and striatum. Integrated intensity was measured in each region, and all measurements (n= 3/group) from a given group of animals were used to generate averages. The reduction in cholesterol in the treated animals was then calculated as the percent decrease of the integrated intensity compared to the knockout β -gal injected mice.

[0112] Positive hASM immunostaining was observed throughout the cerebellum (Figure 6, Table 3), pons, medulla and spinal cord (Figure 7) following unilateral injection of AAV-ASM within the deep cerebellar nuclei.

Table 3

Areas with positive hASM staining as function of AAV serotype. * indicates positive hASM was below limit of detection, but correction of cholesterol pathology still occurred

Structure	AAV1	AAV2	AAV5	AAV7	AAV8
deep cerebellar nuclei	++++	++	+++	+++	++++
cerebellar lobules	++++	++	+++	+++	++++
pons	++	++	++	++	+
medulla	+	++	++	+++	+
spinal cord		+++	+++	++	+
thalamus	*	*	*	*	*
hypothalamus	*	*	*	*	*
hippocampus	*	*	*	*	*
striatum	*	*	*	*	*
cerebral cortex	*	*	*	*	*

[0113] Within the cerebellum mice treated with AAV2/1-ASM and had the most widespread (i.e., spread between lobules within the same sagittal section) level of hASM expression, whereas mice treated with AAV2/2-ASM had the most restricted level of Human ASM protein expression. Human ASM protein expression in mice treated with AAV2/5-ASM, AAV2/7-ASM, and AAV2/8-ASM was intermediate between these two groups. Medial – lateral spread between sagittal sections was maximal in mice treated with serotypes 1 & 8 and minimal in mice injected with serotype 2. Serotypes 5 & 7 initiated medial –lateral spread patterns intermediate between serotypes 1 and 2. Each layer of the cerebellum (i.e., molecular, Purkinje and granular) was transduced by each AAV serotype; however, an increased affinity for the molecular layer was apparent for all serotypes. Purkinje cell transduction was maximal in mice treated with serotypes 1 and 5. Mice injected with serotype 7 had the fewest number of transduced Purkinje cells. Mice treated with serotype 8 also had few transduced Purkinje cells, but had less ASM expression within the granular layer when compared to serotypes 1, 2, 5 & 7. Purkinje cells transduced with ASM appeared to have a healthy cytostructure. Quantitative analysis of AAV mediated hASM protein expression by ELISA in cerebellar tissue homogenates supports our

immunohistochemical findings. Mice injected with serotypes 1 and 8 demonstrated significantly ($p < .0001$) higher cerebellum hASM protein levels when compared to all other mice (Figure 8). Cerebellar hASM levels from mice injected with serotypes 2, 5, & 7 were not above WT levels (i.e. background). As expected Human ASM was not detected in wild type mice – the hASM antibody used in the ELISA are human specific.

[0114] An absence of functional ASM protein results in lysosomal accumulation of sphingomyelin, and subsequent secondary metabolic defects such as abnormal cholesterol trafficking. Sarna et al. Eur. J. Neurosci. **13**:1873-1880 and Leventhal et al. (2001) J. Biol. Chem. **276**:44976-4498. Free cholesterol buildup in the ASMKO mouse brain is visualized using filipin, an autofluorescent molecule isolated from streptomyces filipinensis. Wild-type mouse brains do not stain positively for filipin. In all AAV treated mice (with exception to AAV2/1- β gal) clearance of filipin staining (Table 4) overlapped with areas that were positive for hASM immunostaining indicating that each serotype vector is capable of generating a functional transgene product.

Table 4

[0115] Percent Reduction in Filipin (i.e., cholesterol) clearance as compared to ASMKO mice treated with AAV- β gal in selected brain regions following intracerebellar injection of different AAV serotypes (n=3/serotype; 2/1, 2/2, 2/5, 2/7, and 2/8) encoding for human ASM into the deep cerebellar nuclei of ASMKO mice.

	2/1	2/2	2/5	2/7	2/8
Cerebellum	96.54 \pm 2.14	93.85 \pm 1.287	86.75 \pm 9.58	96.47 \pm 1.93	99.12 \pm .66
Midbrain	96.72 \pm 1.73	53.08 \pm 22.89	65.88 \pm 24.53	73.39 \pm 22.39	91.10 \pm .105
Pons	91.31 \pm 5.80	50.07 \pm 21.26	70.96 \pm 25.60	93.15 \pm 31.20	96.72 \pm 1.20
Medulla	93.29 \pm 6.22	88.46 \pm 3.04	81.55 \pm 17.31	80.73 \pm 14.99	97.40 \pm 1.60
Thalamus	48.88 \pm 25.25	41.21 \pm 27.35	34.86 \pm 16.67	48.44 \pm 28.65	77.03 \pm 12.08
Hypothalamus	82.81 \pm 10.14	86.96 \pm 12.93	88.46 \pm 5.90	82.95 \pm 11.46	99.68 \pm .31
Cortex	27.60 \pm 24.75	73.62 \pm 14.9	55.65 \pm 28.89	76.97 \pm 14.27	98.30 \pm .34

[0116] As previously demonstrated by (Passini et al. (2003) in "Society for Neuroscience" New Orleans, LA), filipin clearance also occurred in areas anatomically connected with the injection site, but that did not stain positively for

hASM. MetaMorph analysis indicated that a reduction in filipin staining occurred throughout the entire rostral caudal axis. In the cerebellum and brainstem filipin was maximally reduced in mice treated with AAV2/1-ASM and AAV2/8-ASM, whereas in the diencephalon and cerebral cortex mice injected with AAV2/8-ASM had the best overall level of filipin clearance (Table 4). Nevertheless, these results indicate that the level of hASM required to correct cholesterol storage pathology in the ASMKO mouse CNS is minimal (i.e., below the hASM immunofluorescence limit of detection).

Table 5

[0117] Histological studies indicate that the ASMKO mouse cerebellum undergoes rapid deterioration. More specifically, Purkinje cells progressively die off between 8 and 20 weeks of age (Sarna et al. (2001) Eur. J. Neurosci. 13:1813-1880 and Stewart et al. (2002) in "Society for Neuroscience" Orland, FL). Calbindin is a widely accepted Purkinje cell marker. Positive calbindin staining in AAV-ASM treated mice would suggest that AAV mediated expression of hASM is therapeutic. Overall our results indicate that AAV mediated hASM expression in the cerebellum prevents Purkinje cell death in the ASMKO mouse (Table 5, Figure 9). As expected Purkinje cell survival did not occur in lobules I-III; mice were injected at 7 weeks of age and by 8 weeks the majority of these cells have already died off. Purkinje cell survival in lobules IV/V was maximal in mice treated with serotype 1. In lobule VI no significant Purkinje cell survival was observed in AAV treated mice. In lobule VII only mice treated with serotype 5 showed significant Purkinje cell survival. In lobule VIII again mice treated with serotype 5 as well as serotype 2 showed significant Purkinje cell survival. In lobules IX and X there were no significant differences between WT and KO mice (or between AAV treated mice) in Purkinje cell counts. This was expected, because at 14 weeks of age (i.e., age at sacrifice) Purkinje cells in these lobules are still viable in ASMKO mice. Across all lobules Purkinje cell survival was maximal in mice treated with serotypes 1, 2, & 5 and minimal in mice treated with serotypes 7 & 8. Purkinje cell survival (based on calbindin staining) in the anterior cerebellar lobules was greatest in mice that were injected with serotype 1.

[0118] Purkinje cell counts in cerebellar lobules I-X in WT and ASMKO mice following intracerebellar injection of different AAV serotypes (n=3/serotype; 2/1, 2/2, 2/5, 2/7, and 2/8) encoding for human ASM into the deep cerebellar nuclei of ASMKO mice. Numbers appearing bold italic are significantly different from KO mice (i.e., mice treated with AAV2/1- β gal) $p \leq .01$

	2/1	2/2	2/5	2/7	2/8	KO	WT
I/II	7.42 \pm 9.80	4.5 \pm 10.58	9.40 \pm 11.59	12.33 \pm 10.58	1 \pm 9.16	5.8 \pm 11.59	113 \pm 10.58
III	12.42 \pm 10.32	11.33 \pm 11.14	26.80 \pm 12.21	15.33 \pm 11.14	9.8 \pm 12.21	2 \pm 9.65	147.50 \pm 11.14
IV/V	60.57 \pm 17.28	36.5 \pm 18.67	27.80 \pm 20.45	29.66 \pm 18.67	6.8 \pm 20.45	8 \pm 16.16	220.66 \pm 18.67
VI	61.14 \pm 11.21	27.5 \pm 12.11	72.20 \pm 13.26	31.16 \pm 12.11	3.8 \pm 13.26	68.5 \pm 10.48	121.16 \pm 12.11
VII	17.42 \pm 4.15	37.66 \pm 4.49	40.60 \pm 4.91	5.33 \pm 4.49	.2 \pm 4.95	17.37 \pm 3.88	7.16 \pm 4.49
VIII	44.14 \pm 10.75	48.66 \pm 11.62	82.80 \pm 12.73	11.33 \pm 11.62	18.40 \pm 12.73	35.12 \pm 10.06	103.33 \pm 11.62
IX	126.28 \pm 19.17	102.66 \pm 20.71	136.40 \pm 22.68	60.16 \pm 20.71	84.40 \pm 22.68	108.0 \pm 17.93	144 \pm 20.71
X	89.85 \pm 12.54	76.83 \pm 13.55	93.80 \pm 14.84	48.16 \pm 13.55	64.80 \pm 14.84	87 \pm 11.73	86.66 \pm 13.55

[0119] On the accelerating rotarod test mice unilaterally injected with AAV2/1-ASM and AAV2/8-ASM demonstrated a significantly ($p < .0009$) longer latency to fall than ASMKO mice injected with AAV2/1- β gal (Figure 10). Mice injected with serotype AA2/1-ASM were not significantly different from wild type mice. Mice injected with AAV2/2-ASM and AAV2/5-ASM showed a trend for a longer latency to fall than ASMKO mice injected with AAV2/1- β gal; whereas, mice injected with AAV2/7-ASM did not. For the rocking rotarod test, only mice injected with AA2/1-ASM demonstrated a significantly ($p < .0001$) longer latency to fall than mice injected with AA2/1- β gal. In this case wild type mice performed significantly better than mice injected with AA2/1-ASM (Figure 10). ASMKO mice that received bilateral injection of either AAV2/1-ASM or AAV2/2-ASM performed significantly ($p < .001$) better than ASMKO AAV2/1- β gal treated mice for both accelerating and rocking tests (Figure 11). AAV2/1-ASM bilaterally injected mice performed comparably to wild type mice for both tests.

[0120] One way to determine if AAV generated hASM is functionally active within the ASMKO CNS is to assess its influence on cholesterol storage pathology - a secondary metabolic defect of NPA disease. In all AAV treated mice (with exception to AAV2/1- β gal) correction of cholesterol storage pathology overlapped with areas that were positive for hASM immunostaining indicating that each

serotype vector is capable of generating a functional transgene product. As previously demonstrated, correction of abnormal cholesterol metabolism correction also occurred in areas anatomically connected with the injection site, but also in regions that did not stain positively for hASM, suggesting that the level hASM required for correction of cholesterol storage pathology is minimal. Consistent with our hASM histochemical and biochemical results mice treated with serotypes 1 and 8 demonstrated a marked reduction in cholesterol storage pathology. Mice treated with serotypes 2, 5, & 7 also showed a reduction in cholesterol storage pathology, but not to the same extent as mice treated with serotypes 1 & 8.

[0121] While cholesterol storage pathology changes are illustrative, a more direct measurement of hASM enzyme activity is through the analysis of sphingomyelin levels, which is the primary substrate accumulated in the tissues of mammals with Niemann-Pick disease. Brain tissue from mice that received bilateral injections of AAV serotype 1 and 2 were assayed for sphingomyelin (SPM) levels (the tissue homogenization procedure for detection of SPM is not compatible with the detection of hASM.) A significant reduction ($p<0.01$) in SPM tissue content in the cerebellum was observed in mice treated with both serotypes 1 and 2 (Figure 12). SPM tissue content levels were also significantly reduced in sections 2, 3, and 4 (each section being 2 mm apart from the next) in mice injected with serotype 1 but not in mice injected with serotype 2.

[0122] ASMKO mice that received bilateral intracerebellar injections of AAV serotypes 1 and 2 encoding for hASM showed a significant reduction in sphingomyelin storage within the cerebellum. As observed with cholesterol storage accumulation, a significant reduction in sphingomyelin storage also occurred in regions outside the cerebellum in ASMKO mice bilaterally treated with AAV1.

[0123] Overall, these results indicate that AAV1 is preferred over serotypes 2, 5, 7, and 8 in its relative ability to initiate enzyme expression, correct storage pathology in the brain, prevent neurodegeneration (by, for example, preventing Purkinje cell death), and improve motor functional outcome. In addition, the DCN

can be exploited as an injection site to maximize enzyme expression throughout the CNS.

[0124] To evaluate the distribution of transgene expression following injection of AAV vectors into the DCN, G93A SOD1 (SOD1^{G93A} mutant mouse, referred to here as the SOD1 mouse) were injected recombinant AAV vectors encoding for green fluorescent protein (GFP). One group of mice was injected with AAV serotype 1 encoding for green fluorescent protein (AAV1-GFP) while another group was injected with AAV serotype 2 encoding for green fluorescent protein (AAV2-GFP).

[0125] The mice were injected bilaterally into the DCN with the AAV recombinant vectors using methods similar to those described above. The dose was approximately 2.0 e10 gc/ml injected per site. Mice were sacrificed about 110 days after birth and their brain and spinal cord were analyzed for GFP staining.

[0126] Green fluorescent protein distribution was observed in the brainstem (see Figure 14) and in the spinal cord regions (see Figure 15) following DCN delivery of AAV encoding for green fluorescent protein (GFP). GFP staining was also observed in the DCN as well as in the olfactory bulbs, cerebral cortex, thalamus, brainstem, cerebellar cortex and spinal cord. All of these areas either receive projections from and/or send projections to the DCN (see Figure 16).

[0127] The specification is most thoroughly understood in light of the teachings of the references cited within the specification. The embodiments within the specification provide an illustration of embodiments of the invention and should not be construed to limit the scope of the invention. The skilled artisan readily recognizes that many other embodiments are encompassed by the invention. All publications, patents, and biological sequences cited in this disclosure are incorporated by reference in their entirety. To the extent the material incorporated by reference contradicts or is inconsistent with the present specification, the present specification will supercede any such material. The citation of any references herein is not an admission that such references are prior art to the present invention.

[0128] Unless otherwise indicated, all numbers expressing quantities of ingredients, cell culture, treatment conditions, and so forth used in the specification, including claims, are to be understood as being modified in all instances by the term "about." Accordingly, unless otherwise indicated to the contrary, the numerical parameters are approximations and may very depending upon the desired properties sought to be obtained by the present invention. Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

1. A method comprising: administering a composition comprising an AAV vector encoding for a biologically active molecule to a site in the disease-compromised central nervous system of a mammal, under conditions whereby said injected AAV vectors transduce cells located at a distal site in the central nervous system and said encoded biologically active molecule is translated.
2. The method of claim 1, wherein the translated biologically active molecule is then expressed.
3. The method of claim 1, wherein the mammal has a lysosomal storage disease or abnormal cholesterol metabolism.
4. The method of claim 3, wherein the lysosomal storage disease is Niemann Pick A disease.
5. The method of claim 1, wherein the mammal is human.
6. The method of claim 1, wherein the distal site is contralateral to the administration site.
7. The method of claim 1, where the administration site is in a region of the CNS selected from the group consisting of the spinal cord, the brainstem, the hippocampus, the striatum, the medulla, the pons, the mesencephalon, the cerebellum, the thalamus, the hypothalamus, the cerebral cortex, the occipital lobe, the temporal lobe, the parietal lobe, and the frontal lobe.
8. The method of claim 1, wherein the administration site is in the hippocampus and the distal site is in a region of the brain selected from the group consisting of the contralateral dentate gyrus and the contralateral CA3, and the medial septum, and the entorhinal cortex.
9. The method of claim 1, wherein the administration site is in a region of the brain selected from the group consisting of the striatum and the cerebellum, and the distal site is in a region of the brain selected from the group consisting of the substantia nigra and the medulla oblongata.
10. The method of claim 9, wherein the administration site is in the deep cerebellar nuclei of the cerebellum.
11. The method of claim 1, wherein the AAV vector has an AAV serotype 1 capsid.
12. The method of claims 3 or 4, wherein the AAV vector has an AAV serotype 1 capsid.

13. The method of claim 10, wherein the AAV vector has an AAV serotype 1 capsid.
14. The method of claim 1, wherein the AAV vector is AAV1 or AAV2/1.
15. The method of claims 3 or 4, wherein the AAV vector is AAV1 or AAV2/1.
16. The method of claim 10, wherein the AAV vector is AAV1 or AAV2/1.
17. The method of claim 1, further comprising administration of a composition to a second site in the CNS of the mammal, wherein the composition comprises an AAV vector comprising a polynucleotide encoding a biologically active molecule product.
18. The method of claim 17, wherein the second administration site is contralateral to the first administration site.
19. The method of claim 2, wherein the biologically active molecule is a lysosomal hydrolase.
20. The method of claim 19, wherein the lysosomal hydrolase is any one of lysosomal hydrolases listed in Table 1.
21. The method of claim 20, wherein the lysosomal hydrolase is acid sphingomyelinase.
22. The method of claim 1, wherein the administration site is in the deep cerebellar nuclei and the distal site is the spinal cord.
23. The method of claim 1, wherein the distance between the administration site and distal site is at least 2 mm.
24. The method of claim 1, wherein the concentration of the AAV vector in the composition is at least 5 X 10¹² gp/ml.
25. A method comprising: administering a composition comprising an AAV vector encoding for a metalloendopeptidase molecule to a site in the central nervous system of a mammal with Alzheimer's disease, under conditions whereby said injected AAV vectors transduce cells located at a distal site in the central nervous system and said encoded metalloendopeptidase is expressed.
26. The method of claim 25, wherein the metalloendopeptidase is selected from the group consisting of neprilysin, insulysin, and thimet oligopeptidase.

27. The method of claim 1 or 25, wherein the AAV vector is selected from the group consisting of AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, and AAV8.
28. The method of claim 25, wherein the distance between the administration site and the distal site is at least 2 mm.
29. The method of claim 25, wherein the concentration of the AAV vector in the composition is at least 5 X 10¹² gp/ml.
30. The method of claim 29, wherein the AAV is selected from a group consisting of AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, and AAV8.
31. The method of claim 29, wherein the AAV is a recombinant AAV vector.
32. The method of claim 29, wherein the recombinant AAV vector is selected from the group consisting of AAV2/1, AAV2/2, AAV2/5, AAV2/7 and AAV2/8 serotype vectors.




FIG. 1A




FIG. 1C




FIG. 1B




FIG. 2A




FIG. 2C






FIG. 2B






FIG. 5




Figure 6




Figure 7




FIGURE 8




Figure 9





Figure 5

FIGURE 10




Figure 6

FIGURE 11




FIGURE 12

Figure 13A






Figure 13B





AAV1-GFP

Figure 14




Figure 15




Figure 16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US06/17242

A. CLASSIFICATION OF SUBJECT MATTER

IPC: A61K 48/00(2006.01);C12N 15/63(2006.01),15/74(2006.01),15/86(2006.01)

USPC: 514/44;435/320.1,455,456

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44;435/320.1,455,456

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2004/0258666 A1 (PASSINI et al.) 23 December 2004 (23.12.2004), see entire document.	1-32
Y	US 2003/0165481 A1 (HERSH) 4 September 2003 (04.09.2003), see entire document.	25-32
X	BOSCH, A. et al. Long-term and significant correction of brain lesions in adult	1-3,5-7,19-20,23
---	mucopolysaccharidosis type VII mice using recombinant AAV vectors. Mol. Ther. January	-----
Y	2000, Vol. 1, No. 1, pages 63-70, see entire document.	4, 8-18,21-22,24
Y	DATTO, J. et al. Distribution of AAV-1 GFP with respect to time, distance traveled, and expression following injection to the midbrain. Society for Neuroscience Abstract Viewer. 2002. Online. Program No. 623.4, see abstract.	1-32



Further documents are listed in the continuation of Box C.



See patent family annex.

*	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

15 August 2006 (15.08.2006)

Date of mailing of the international search report

15 SEP 2006

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner of Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (571) 273-3201

Authorized officer

Anne Marie S. Wehbe

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

PCT/US06/17242

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LIU, Y. et al. High level of transduction of mammalian brain by adeno-associated type 1 virus vector. Society for Neuroscience Abstract Viewer. 2002. Online. Program No. 902.2, see abstract.	11-16, 32

INTERNATIONAL SEARCH REPORT

PCT/US06/17242

Continuation of B. FIELDS SEARCHED Item 3:
DIALOG-medicine database; BRS/EAST-USPatful, EPO, JPO, Derwint, PG-Pubs
search terms: aav,raav,adeno associated, brain,cns, stereotactic,metalloendopeptidase,lysosomal storage disease,acid sphingomyelinase,nepriyisn,insulysin,thimet,hippocampus