

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 241/00	A2	(11) International Publication Number: WO 99/02502 (43) International Publication Date: 21 January 1999 (21.01.99)
(21) International Application Number: PCT/EP98/04973 (22) International Filing Date: 9 July 1998 (09.07.98) (30) Priority Data: 9714530.4 11 July 1997 (11.07.97) GB 9724530.2 19 November 1997 (19.11.97) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BROMIDGE, Steven, Mark [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). MOSS, Stephen, Frederik [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: WATERS, David, Martin; SmithKline Beecham plc, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NOVEL COMPOUNDS (57) Abstract The invention relates to novel compounds having pharmacological activity, processes for their preparation, to compositions containing them and to their use in the treatment of CNS disorders.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


NOVEL COMPOUNDS

This invention relates to novel compounds having pharmacological activity, processes for their preparation, to compositions containing them and to their use in the treatment of CNS disorders.

- 5 EPA 0 021 580 and EPA 0 076 072 describe sulphonamide derivatives which are disclosed as having antiarrhythmic activity. A structurally distinct class of compounds has now been discovered, which have been found to have 5HT₆ receptor antagonist activity. 5HT₆ receptor antagonists are believed to be of potential use in the treatment of certain CNS disorders such as anxiety, depression, epilepsy, obsessive
10 compulsive disorders, migraine, Alzheimers disease, sleep disorders (including disturbances of Circadian rhythm), feeding disorders such as anorexia and bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Compounds of the invention are also
15 expected to be of utility for cognitive memory enhancement. Compounds of the invention are also expected to be of use in the treatment of certain GI (gastrointestinal) disorders such as IBS (Irritable Bowel Syndrome).

The present invention therefore provides, in a first aspect, a compound of formula (I) or a salt thereof:

20



(I)

25 wherein:

P is phenyl, naphthyl, anthracenyl, a bicyclic heterocyclic ring, a tricyclic heteroaromatic ring or is a 5 to 7-membered heterocyclic ring each containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulphur;

A is a single bond, a C₁₋₆alkylene or a C₁₋₆alkenylene group;

30 B is SO₂;

R¹ is halogen, C₁₋₆alkyl optionally substituted by one or more fluorine atoms, C₃₋₆cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkanoyl, C₁₋₆alkoxy, OCF₃, hydroxy, hydroxyC₁₋₆alkyl, hydroxyC₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkoxy, nitro, cyano, NR¹⁰R¹¹ where R¹⁰ and R¹¹ are independently hydrogen, C₁₋₆alkyl or optionally

- substituted phenyl, SR^{11} where R^{11} is as defined above or R^1 is optionally substituted phenyl, naphthyl, a bicyclic heterocyclic ring or is a 5 to 7-membered heterocyclic ring each containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulphur, or R^1 together with a second R^1 substituent forms a group $-O-CH_2-O-$, OCH_2CH_2O- , $-CH_2CH_2CH_2-$ or $-CH_2CH_2CH_2CH_2-$,
 5 n is 0, 1, 2, 3, 4, 5 or 6;
 R^2 is hydrogen, C_{1-6} alkyl, aryl C_{1-6} alkyl or together with group P form a 5 to 8 membered ring optionally substituted with one or more C_{1-6} alkyl groups;
 R^3 is hydrogen, halogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkanoyl, C_{1-6} alkoxy
 10 optionally substituted with one or more fluorine atoms, hydroxy, hydroxy C_{1-6} alkyl, hydroxy C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkoxy, , nitro, trifluoromethyl, cyano or aryl or together with the group R^5 forms a group $(CH_2)_2O$ or $(CH_2)_3O$ optionally substituted with 1 or more C_{1-6} alkyl groups;
 R^4 is $-X(CH_2)_p-R^6$ where X is a single bond, CH_2 , O, NH or N-alkyl and p is 0 to 6
 15 and R^6 is an optionally substituted 4- to 7-membered heterocyclic ring containing 1 to 3 heteroatoms selected from nitrogen, sulphur or oxygen, or R^6 is NR^7R^8 where R^7 and R^8 are independently hydrogen, C_{1-6} alkyl or aryl C_{1-6} alkyl; and
 R^5 is a group R^3 or together with R^3 forms a group $(CH_2)_2O$ or $(CH_2)_3O$ optionally substituted with 1 or more C_{1-6} alkyl groups.

20

- C_{1-6} Alkyl groups, whether alone or as part of another group, may be straight chain or branched. As used herein the term aryl includes phenyl and naphthyl. When a group is defined as "optionally substituted", unless otherwise stated, suitable substituents include halogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkanoyl, C_{1-6} alkoxy
 25 optionally substituted with one or more fluorine atoms, hydroxy, hydroxy C_{1-6} alkyl, hydroxy C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkoxy, nitro, trifluoromethyl, or cyano.

- When P is a bicyclic heterocyclic ring, suitable examples include benzothiophene, indole, quinoline or isoquinoline. Bicyclic heterocyclic rings can also be partially saturated. When P is a tricyclic heteroaromatic ring suitable examples
 30 include dibenzofuran. Suitable 5 to 7-membered heterocyclic rings include thienyl, furyl, pyrrolyl, triazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, isothiazolyl, isoxazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyrrolidinyl and pyrazinyl. The heterocyclic rings can be linked to the remainder of the molecule via any suitable carbon atom or, when present, a nitrogen atom. Preferably P is phenyl or naphthyl.

- 35 Suitably A is a single bond, a methylene or ethylene group or a $-CH=CH-$ group. Preferably A is a single bond or methylene.

When R^1 is a bicyclic heterocyclic ring or a 5 to 7 membered heterocyclic ring suitable examples include those listed in the definition of P.

It will be appreciated that when R^1 combines with a second R^1 substituent the two substituents must be attached to adjacent atoms on the ring P. Thus, when P is phenyl, groups such as methylenedioxyphenyl, ethylenedioxyphenyl, indane and tetrahydronaphthalene are within the scope of this invention.

5 Suitably R^1 is hydrogen, halogen, phenyl, C_{1-6} alkoxy most preferably OMe, SR^{11} most preferably SMe or C_{1-6} alkyl optionally substituted by one or more fluorine atoms, for example methyl or trifluoromethyl. Preferably R^1 is halogen. Preferably n is 0, 1, 2, 3 or 4.

Suitably R^2 is hydrogen, methyl or together with group P form a 5 or 6-
10 membered ring. It will be appreciated that when groups P and R^2 are linked together the latter must be attached to the adjacent carbon atom on the ring P i.e. with an ortho relationship with respect to group A.

It will be appreciated that when R^3/R^5 groups are linked together the two groups must be attached to adjacent carbon atoms of the phenyl ring. Preferably R^3 is a group R^5 , in particular hydrogen.
15

Preferably R^4 is meta with respect to the substituent B. Preferably X is a bond, p is 0 and R^6 is an optionally substituted 5- to 7-membered heterocyclic ring. The heterocyclic rings can be linked to the remainder of the molecule via a carbon atom or, when present, a nitrogen atom. Optional substituents for these rings, which can be
20 present on carbon and/or nitrogen atoms, include C_{1-6} alkyl, in particular methyl or NR^9R^{10} where R^9 and R^{10} are independently hydrogen or C_{1-6} alkyl. More preferably R^4 is an optionally substituted piperazine. Most preferably R^4 is N-methyl piperazine or NH-piperazine.

Preferably R^5 is para with respect to the substituent B. Suitably R^5 is C_{1-6} alkoxy. Preferably R^5 is methoxy.
25

Particular compounds of the invention include:

4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-naphthalen-1-ylbenzenesulfonamide,
N-(4-Chloronaphthalen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene
30 sulfonamide,
N-(3-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide,
N-(3,4-Dichlorobenzyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide,
6-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-
tetrahydroisoquinoline,
35 1-[4-Methoxy-3-(4-methyl-piperazin-1-yl)-benzene-sulfonyl]-6-trifluoromethyl-2,3-
dihydro-1*H*-indole,
N-(3-Chlorophenyl)-4-methoxy-*N*-methyl-3-(4-methylpiperazin-1-yl)-
benzenesulfonamide,

- 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline,
N-(3-Iodo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
5 *N*-(5-Iodo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3,4-Methylenedioxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
6-Chloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methyl-2,3-
10 dihydro-1*H*-indole,
7,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
7,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
15 5-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
8-Chloro-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
6,7-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-
20 tetrahydroisoquinoline,
2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-phenyl-1,2,3,4-tetrahydroisoquinoline,
8-Bromo-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
25 5,6-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
5,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
6-Iodo-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methylthio-2,3-
30 dihydro-1*H*-indole,
N-(3,4-Dichlorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
6,7,8-Trimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
35 2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-6-pyridin-3-yl-2,3-dihydro-1*H*-indole,
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-5-pyridin-3-yl-2,3-dihydro-1*H*-indole,

- 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,5-tetrahydropyrrolo[2,3-f]indole,
1-(4-Methoxy-3-piperazin-1-ylbenzenesulfonyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline,
- 5 *N*-(3-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide
N-(2-Fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(2-Trifluoromethoxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-
- 10 benzenesulfonamide,
N-(4-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(9,10-Dioxo-9,10-dihydroanthracen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Hydroxymethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-
- 15 benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(2-methylsulfanylphenyl)-benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(5,6,7,8-tetrahydronaphthalene-1-yl)-benzenesulfonamide,
- 20 *N*-(2-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
4-Methoxy-*N*-(2-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3,4-Dimethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
4-Methoxy-*N*-(2-methoxy-6-methylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
- 25 *N*-(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
8-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
N-(2-Chloro-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-
- 30 benzenesulfonamide,
N-(2-Trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-quinolin-7-ylbenzenesulfonamide (E47)
N-(4-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
- 35 *N*-(3-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3-Bromo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,

- 4-Methoxy-*N*-(2-methoxydibenzofuran-3-yl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(4-Cyclohexylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
 5 *N*-(2-Chloro-4-iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Bromo-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-[4-(4-Chlorophenyl)thiazol-2-yl]-4-methoxy-3-(4-methylpiperazin-1-yl)-
 10 benzenesulfonamide,
 4-Methoxy-*N*-(3-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3-Chloro-4-bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
 15 *N*-(2-Acetylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-phenylaminophenyl)-benzenesulfonamide,
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-pentyloxyphenyl)-benzenesulfonamide,
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-vinylphenyl)benzenesulfonamide,
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(2-pyrrol-1-ylphenyl)-benzenesulfonamide,
 20 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-[4-(4-nitrophenylsulfanyl)phenyl]-benzenesulfonamide,
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(3-oxazol-5-ylphenyl)-benzenesulfonamide,
N-(4-Bromo-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
 25 4-Methoxy-*N*-(2,3-dimethylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(4-Chloro-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-trifluoromethyl-2,3-dihydro-1*H*-indole,
 30 7-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
 5,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
 5,7-Dichloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
 35 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-6-methyl-1,2,3,4-tetrahydroquinoline,

- 6-Fluoro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
5-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulphonyl]-2,3-dihydro -
1*H*-isindole hydrochloride,
- 5 N-(2-Isopropylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide hydrochloride,
N-(4-Chloronaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-N-naphthalen-1-yl-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Chloro-2-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-Indan-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 10 N-(2-Fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-N-(2-methylsulfonylphenyl)-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-3-piperazin-1-yl-N-(2-trifluoromethylphenyl)benzenesulfonamide,
4-Methoxy-N-(2-methylphenyl)-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Ethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 15 4-Methoxy-3-piperazin-1-yl-N-(3-trifluoromethylphenyl)benzenesulfonamide,
N-(3,4-Dimethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3,4-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 20 N-(3,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Chloro-3-fluoro-4-methylphenyl)-4-methoxy-3-piperazin-1-yl-
benzenesulfonamide,
N-(4-Chloro-3-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 25 N-Benzol[1,3]dioxol-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Bromo-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Chloro-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 30 N-(4-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Isopropenylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-N-(2-methyl-5-nitrophenyl)-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-*tert*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
- 35 N-(4-Isopropylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Hexylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,4-Dibromonaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-N-(4-methoxybiphenyl-3-yl)-3-piperazin-1-ylbenzenesulfonamide,

- N*-(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide,
N-Biphenyl-2-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Benzylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Propylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
5 *N*-(2-*sec*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-*tert*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Iodo-2-methylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide,
6-Chloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-2,3-dihydro-1*H*-
10 indole hydrochloride,
6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methylsulfanyl-2,3-dihydro-
1*H*-indole,
6-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline,
8-Chloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-
15 tetrahydroisoquinoline,
1-(4-Methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-6-trifluoromethyl-2,3-
dihydro-1*H*-indole,
5,8-Dimethoxy-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-
tetrahydroisoquinoline hydrochloride,
20 5,8-Dichloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-
tetrahydroisoquinoline hydrochloride,
N-(3-Iodo-4-methylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide,
5,7-Dichloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-
tetrahydroquinoline,
25 *N*-(2-Chloro-3,5-difluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Chloro-2-trifluoromethoxyphenyl)-4-methoxy-3-piperazin-1-
ylbenzenesulfonamide,
N-(2,4,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Chloro-2-methoxyphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
30 *N*-(4-Chloro-2-trifluoromethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide ,
N-(3-Bromo-2,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,3,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Bromo-2,3-dihydro-benzofuran-7-yl)-4-methoxy-3-piperazin-1-
35 ylbenzenesulfonamide,
N-(2-Bromo-3,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Bromo-5,6-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dibromo-3-fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide

N-(2,5-Dibromo-3-chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide

N-(2,3,5-Tribromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide

6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1H-indole,

5-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1H-indole,


- 5 7-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline
and pharmaceutically acceptable salts thereof.

The compounds of the formula (I) can form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric, lactic, mandelic, tartaric and
10 methanesulphonic.

Compounds of formula (I) may also form solvates such as hydrates, and the invention also extends to these forms. When referred to herein, it is understood that the term 'compound of formula (I)' also includes these forms.

Certain compounds of formula (I) are capable of existing in stereoisomeric
15 forms including diastereomers and enantiomers and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis. The invention also extends to any tautomeric forms and mixtures thereof.

20 The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises the coupling of a compound of formula (II):




25

(II)

in which R^1 , R^2 , n , P , and A are as defined in formula (I) or protected derivatives thereof with a compound of formula (III):

30



(III)

in which B, R³, R⁴ and R⁵ are as defined in formula (I) or protected derivatives thereof and L is a leaving group and optionally thereafter:

- removing any protecting groups,
- forming a pharmaceutically acceptable salt.

5 Suitable leaving groups include halogen, in particular chloro. The reaction of a compounds of formulae (II) and (III) is carried out by mixing the two reagents together, optionally in an inert solvent such as dichloromethane with or without the addition of a suitable base such as triethylamine [or pyridine].

10 Those skilled in the art will appreciate that it may be necessary to protect certain groups. Suitable protecting groups and methods for their attachment and removal are conventional in the art of organic chemistry, such as those described in Greene T.W. 'Protective groups in organic synthesis' New York, Wiley (1981).

15 Compounds of formulae (II) and (III) are commercially available or may be prepared according to known methods or analogous to known methods. For example to prepare compounds of formulae (I) where R³ is H, R⁵ is OMe and R⁴ is 1-piperazine a suitable protecting group in intermediates of formulae (III) was found to be trichloroacetyl. Thus reacting 1-(2-methoxyphenyl) piperazine with trichloroacetyl chloride in a suitable solvent such as dichloromethane in the presence of a base such as diisopropylethylamine afforded 2-(4-trichloroacetyl-piperazin-1-yl) anisole. On
20 treatment with chlorosulfonic acid at 0°C in a suitable inert solvent such as dichloromethane 3-(4-trichloroacetyl-piperazin-1-yl)-4-methoxybenzenesulfonyl chloride was obtained. Coupling of this compound with compounds of formulae (II) as described above followed by treatment with 20% aqueous potassium hydroxide afforded the required compound.

25 Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

30 Compounds of formula (I) and their pharmaceutically acceptable salts have 5HT₆ receptor antagonist activity and are believed to be of potential use in the treatment of certain CNS disorders such as anxiety, depression, epilepsy, obsessive compulsive disorders, migraine, Alzheimers disease, sleep disorders (including disturbances of Circadian rhythm), feeding disorders such as anorexia and bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus. Compounds of the invention are also
35 expected to be of use in the treatment of certain GI disorders such as IBS.

 Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of the above disorders.

The invention further provides a method of treatment or prophylaxis of the above disorders, in mammals including humans, which comprises administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

5 In another aspect, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment or prophylaxis of the above disorders.

The present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof,
10 and a pharmaceutically acceptable carrier.

A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable
15 powders, injectable or infusable solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated
20 according to methods well known in normal pharmaceutical practice.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents,
25 emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The compound, depending on the vehicle and concentration used, can be
30 either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed
35 under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a

surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 20.0 mg, for example 0.2 to 5 mg; and such unit doses may be administered more than once a day, for example two or three a day, so that the total daily dosage is in the range of about 0.5 to 100 mg; and such therapy may extend for a number of weeks or months.

When administered in accordance with the invention, no unacceptable toxicological effects are expected with the compounds of the invention.

The following Descriptions and Examples illustrate the preparation of compounds of the invention.

Description 1

2-(4-Methylpiperazin-1-yl)anisole (D1)

To an ice-cooled, stirred suspension of lithium aluminium hydride (7.9g, 0.21mol) in dry tetrahydrofuran (150ml) was added a solution of 1-(2-methoxyphenyl) piperazine (10g, 52 mmol) in dry tetrahydrofuran (150ml) over 0.5h under argon. A solution of ethyl formate (12.6ml, 0.156mol) in dry tetrahydrofuran (25ml) was added to the cold mixture over 0.25h and the resulting suspension was stirred for a further 2h at room temperature. Dilute sodium hydroxide solution (15%, 8ml) was slowly added to the cooled mixture, followed by water (24ml) and the whole left to stir for 0.25h. The mixture was filtered and the filtrate concentrated to an oil which was partitioned between dichloromethane and water. The organic phase was dried and concentrated to an oil which was purified by column chromatography on silica gel eluting with a methanol/dichloromethane gradient to afford the title compound as a colourless oil (5.7g, 53%)

δ_H (250 MHz, $CDCl_3$), 2.36 (3H, s), 2.63 (4H, br s), 3.10 (4H, br s), 3.86 (3H, s), 6.84-7.03 (4H, m).

Description 2

3-(4-Methylpiperazin-1-yl)-4-methoxybenzene sulfonyl chloride (D2)

2-(4-Methylpiperazin-1-yl)anisole (200mg, 1mmol) was added in portions over ten minutes to ice-cooled, stirred chlorosulfonic acid (1.2ml) under argon. The resulting

brown solution was stirred at 0°C for 0.25h and then at ambient temperature for a further 1.25h. The solution was slowly poured onto crushed ice (50g).

Dichloromethane (50ml) was added to the mixture followed by saturated sodium carbonate solution until pH10 was attained in the aqueous phase. The layers were

5 separated and the aqueous phase further extracted with dichloromethane. The combined organic phases were dried (Na₂SO₄) and concentrated to an oil. The oil was stirred with hexane (4ml) to give the title compound as a cream solid (210mg, 71%).

10 δ_H (250 MHz, CDCl₃) 2.43 (3H, s), 2.71 (4H, br t, J = 4.2), 3.20 (4H, br t, J = 4.2), 4.00 (3H, s), 6.97 (1H, d, J 8.7), 7.48 (1H, d, J 2.2), 7.72 (1H, dd, J 2.2, 8.7), MS: m/z (MH⁺) = 305.

Description 3

2-(4-Trichloroacetyl piperazin-1-yl) anisole (D3)

15 A solution of 1-(2-methoxyphenyl) piperazine (7.0g) in dichloromethane (30ml) was added over 0.25h to a stirred solution of trichloroacetyl chloride (4.06ml) in dichloromethane (40ml) at room temperature under argon. Diisopropylethylamine (5.95ml) was then added and the whole was stirred for 18h. The reaction mixture was washed with water (2 x 100ml), dried (Na₂SO₄) and concentrated to give the title compound (D3) as an oil (11.2g, 91%). MH⁺ 337/339.

Description 4

3-(4-Trichloroacetyl piperazin-1-yl)-4-methoxybenzenesulfonyl chloride (D4)

25 A solution of 2-(4-trichloroacetyl piperazin-1-yl) anisole (D3) (10g) in dichloromethane (115ml) was added over 0.3h to ice-cooled chlorosulfonic acid (52ml). After 0.5h at 0°C then 1h at ambient temperature, the solution was poured onto a mixture of ice-water (500g) and dichloromethane (500ml) with rapid stirring. The layers were separated and the organic phase was washed with water (2 x 800ml), dried (MgSO₄) and concentrated to give the title compound (D4) as a foam (6.0g, 46%). MH⁺ 435/437.

Description 5

6-Iodo-2,3-dihydro-1H-indole (D5)

This compound was prepared as previously described (Heterocycles, 1987, 26, 2817)

Description 6

5-Iodo-2,3-dihydro-1H-indole (D6)

35 This compound was prepared as previously described (Chem. Pharm. Bull. 1987, 35, 3146.)

Description 7**3,5-Dibromoaniline (D7)**

A suspension of 3,5-dibromo nitrobenzene (J. Amer. Chem Soc., 1950, 72, 793) (1.0g, 3.6mmol) in methanol (30ml) was added in portions to a stirred mixture of iron powder (0.52g, 9.3mmol) in a saturated solution of ammonium chloride (50ml) at 60°C. The mixture was heated at reflux for 2h, filtered and the filtrate extracted with dichloromethane (2 x 70ml). The organic extract was dried (Na₂SO₄) and concentrated *in vacuo* to give the title compound (D7) as an oil (0.787g, 87%), MH⁺ 250/252.

Description 8**2-Bromo-3,5-dichloro-4-nitroaniline (D8)**

A solution of N-bromosuccinimide (2.6g, 14.5mmol) in N,N-dimethylformamide (DMF) (70ml) was added over 20 min to a stirred solution of 2,5-dichloro-4-nitroaniline (3.0g, 14.5mmol) in DMF (30ml) at room temperature under argon. After stirring for 18h, the mixture was poured into water (1l) and extracted with dichloromethane (500ml). The organic extract was washed with water (5 x 500ml), dried (MgSO₄) and concentrated to an oil which was purified by column chromatography over silica gel eluting with ethyl acetate/hexane gradient to give the title compound (D8) as a yellow solid (1.0g, 24%), MH⁺ 285/287.

Description 9**3-Bromo-2,5-dichloro nitrobenzene (D9)**

Concentrated sulphuric acid (2.2ml) was slowly added to a suspension of 2-bromo-3,5-dichloro-4-nitroaniline (D8) (0.9g, 3.1mmol) in ethanol (20ml). The resulting solution was heated to reflux and crushed sodium nitrite (478mg, 6.9mmol) was added in two portions. After 0.5h at reflux, the mixture was cooled, diluted with dichloromethane (50ml) and saturated sodium hydrogen carbonate solution (50ml) was added. The layers were separated and the organic phase dried (Na₂SO₄) and concentrated to an oil which was purified by column chromatography over silica gel eluting with a gradient of ethyl acetate/hexane to give the title compound (D9) as an orange solid (0.67g, 80%), MH⁺ 269/271.

Description 10**3-Bromo-2,5-dichloroaniline (D10)**

3-Bromo-2,5-dichloro nitrobenzene (D9) was treated with iron powder in the manner described in Description 7 to give the title compound (D10) as a solid (77%), MH^+ 240/242.

5 **Description 11**

2,3,6-trichloro-4-nitroaniline (D11)

To a suspension of 2,5-dichloro-4-nitroaniline (4.0g, 19.3mmol) in ethanol (50ml) was added concentrated hydrochloric acid (20ml) and water (20ml). The mixture was heated to 50°C and 27.5% hydrogen peroxide (6ml) added over 15 min. The mixture
10 was maintained at this temperature for a further 2h, cooled to room temperature, and the solid filtered and washed with water (2 x 20ml) to give the title compound (D11) (4.1g, 88%), MH^+ 241/243.

Description 12

15 **2,3,5-Trichloro nitrobenzene (D12)**

2,3,6-Trichloro-4-nitroaniline (D11) was deaminated as described in Description 9 to give the title compound (D12) (64%), MH^+ 226/228.

Description 13

20 **2,3,5-Trichloroaniline (D13)**

2,3,5-Trichloro nitrobenzene (D12) was reduced with iron powder as described in Description 7 to give the title compound (D13) (68%), MH^+ 196/198.

Description 14

25 **7-Amino-5-bromo-2,3-dihydrobenzofuran (D14)**

Concentrated sulfuric acid (8.8ml) was added over 5 min to a stirred mixture of 5-bromo-2,3-dihydrobenzofuran-7-carboxylic acid (0.55g, 2.3mmol) in chloroform (27ml) at 45°C. Sodium azide (0.737g, 11.3mmol) was then added portion-wise over 0.5h and the temperature was maintained for a further 1h after which time the mixture
30 was poured onto ice (100g) and extracted with chloroform (2 x 50ml). The aqueous phase was basified to pH 12 with 40% sodium hydroxide solution and extracted with chloroform (2 x 50ml). The extract was dried (Na_2SO_4), concentrated, and the residue purified by column chromatography over silica gel eluting with a gradient of acetone/toluene to give the title compound (D14) as a solid (63mg, 13%), MH^+
35 214/216.

Description 15

7-Bromo-1,2,3,4-tetrahydroquinoline (D15)

A solution of 7-bromoquinoline (J. Amer. Chem. Soc., 1947, **69**, 705) (362mg, 1.74mmol) in glacial acetic acid (10ml) was treated portion-wise with sodium cyanoborohydride (437mg, 7.0mmol) under argon at room temperature. After 18h at this temperature, the mixture was cooled in an ice bath and to it was added water (35ml) and 50% aq. sodium hydroxide until pH 14 was attained. The mixture was extracted with dichloromethane and the organic phase washed with saturated sodium chloride solution, dried (Na₂SO₄) and concentrated *in vacuo* to a residue which was purified by column chromatography over silica gel eluting with a gradient of methanol/dichloromethane to give the title compound (D15) (168mg, 46%). MH⁺ 212/214.

Description 16

3-(4-Methyl-1-piperazin-1-yl)-4-methoxybenzenesulphonamide (D16)

The title compound (D16) was prepared in 35% yield by treating the sulfonyl chloride (D2) with excess aqueous ammonia in acetone.

Many of the intermediates used in the preparation of the compounds of this invention can be prepared by known procedures. These are shown in Table A.

Table A

Description No.	Compound Name	Reference
Description 17	6-Trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole (D17)	J. Med. Chem. 1998, 41(10), 1598-1612
Description 18	6-Chloro-5-methyl-2,3-dihydro-1 <i>H</i> -indole (D18)	WO-95/01976
Description 19	7,8-Dichloro-1,2,3,4-tetrahydroisoquinoline (D19)	J. Org. Chem. 1980, 45(10), 1950-3
Description 20	7,8-Dimethoxy-1,2,3,4-tetrahydroisoquinoline (D20)	J. Org. Chem. 1968, 33(2), 494-503
Description 21	5-Bromo-1,2,3,4-tetrahydroisoquinoline (D21)	WO-95/13274
Description 22	8-Chloro-7-methoxy-	J. Med. Chem. 1982,

	1,2,3,4-tetrahydroisoquinoline (D22)	25(10), 1235-40
Description 23	6,7-Dimethoxy-1,2,3,4-tetrahydroisoquinoline (D23)	J. Org. Chem. 1968, 33(2), 494-503
Description 24	7-phenyl-1,2,3,4-tetrahydroisoquinoline (D24)	J. Pharm. Sci. 1970, 59(1), 59-62
Description 25	8-Bromo-7-methoxy-1,2,3,4-tetrahydroisoquinoline (D25)	J. Het. Chem. 1978, 15(3), 429-32
Description 26	5,6-Dichloro-1,2,3,4-tetrahydroisoquinoline (D26)	J. Med. Chem. 1980, 25(3), 506-11
Description 27	5,8-Dimethoxy-1,2,3,4-tetrahydroisoquinoline (D27)	J. Med. Chem. 1981, 24(12), 1432-7
Description 28	6-Iodo-5-methylthio-2,3-dihydro-1 <i>H</i> -indole (D28)	WO-95/01976
Description 29	6,7,8-Trimethoxy-1,2,3,4-tetrahydroisoquinoline (D29)	Heterocycles 1989, 29(6), 2817-22
Description 30	6-pyridin-3-yl-2,3-dihydro-1 <i>H</i> -indole (D30)	FR-2530246 CA:101:23351
Description 31	5-pyridin-3-yl-2,3-dihydro-1 <i>H</i> -indole (D31)	FR-2530246 CA:101:23351
Description 32	1,2,3,5-tetrahydropyrrolo-[2,3- <i>f</i>]indole (D32)	J. Med. Chem. 1996, 39(25), 4966-77
Description 33	3-Fluoro-5-pyridin-3-ylaniline (D33)	WO-96/23783
Description 34	8-Chloro-1,2,3,4-tetrahydroisoquinoline (D34)	J. Med. Chem. 1980, 23(5), 506-11

Description 35	5-Trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole (D35)	WO-97/48700
Description 36	7-Bromo-1,2,3,4-tetrahydroisoquinoline (D36)	WO-98/06699
Description 37	5,8-Dichloro-1,2,3,4-tetrahydroisoquinoline (D37)	J. Med. Chem. 1980, 23(5), 506-11
Description 38	5,7-Dichloro-1,2,3,4-tetrahydroquinoline (D38)	J. Med. Chem. 1980, 23(5), 506-11
Description 39	6-Fluoro-1,2,3,4-tetrahydroquinoline (D39)	JP-55040616
Description 40	6-Bromo-1,2,3,4-tetrahydroquinoline (D40)	EP-702004
Description 41	8-Chloro-1,2,3,4-tetrahydroisoquinoline (D41)	J. Med. Chem. 1980, 23(5), 506-11
Description 42	5-Methyl-6-trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole (D42)	WO-97/48700
Description 43	5,7-Dichloro-1,2,3,4-tetrahydroquinoline (D43)	JP-55040616

Example 1**4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-naphthalen-1-ylbenzenesulfonamide hydrochloride (E1)**

5 1-Naphthylamine (29mg, 0.2mmol) was added to a stirred solution of 3-(4-methyl-1-piperazinyl)-4-methoxybenzene sulfonyl chloride (D2) (60mg, 0.2mmol) in acetone (1ml) at ambient temperature. After stirring for 18h, the precipitate was filtered off and washed with acetone and diethyl ether to afford the title compound (E1) as a

10 cream solid (60mg, 67%).

δ_H (250 MHz, DMSO- d_6) 2.88 (3H, s), 2.90-2.97 (2H, m), 3.15-3.30 (2H, m), 3.37-3.56 (4H, m), 3.89 (3H, s), 7.11 (1H, d, $J = 8.8$), 7.22-7.59 (6H, m), 7.86 (1H, d, $J = 8.3$), 7.96 (1H, d, $J = 9.0$), 8.10 (1H, d, $J = 7.0$), 10.18 (1H, s), 10.55 (1H, br s). MS: m/z ($MH^+ - HCl$) = 412.

5

The compounds in Tables 1 and 2 were prepared in a similar manner to Example 1 using 3-(4-methylpiperazin-1-yl)-4-methoxybenzene sulfonyl chloride (D2) and the appropriate amine. All amines are either commercially available or can be prepared by methods as referenced in Table A above. If required purification by recrystallisation or
10 alternatively by aqueous basic (K_2CO_3) workup followed by column chromatography was carried out.

Table 1

Compound	δ_H (250 MHz, DMSO- d_6)	MS (MH^+)
<i>N</i> -(4-Chloronaphthalen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide hydrochloride (E2)	2.75 (3H, d, $J=4.5$), 2.70-2.84 (2H, m), 3.02-3.15 (2H, m), 3.25-3.45 (4H, m), 3.74 (3H, s), 6.95 (1H, d, $J=8.5$), 7.07-7.13 (2H, m), 7.21 (1H, d, $J=8.5$), 7.48-7.62 (3H, m), 8.04 (1H, t, $J=7.2$), 10.11 (1H, s), 10.20 (1H, br s).	446
<i>N</i> -(3-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide hydrochloride (E3)	2.68 (3H, d $J=3.5$), 2.78-2.88 (2H, br m), 3.0-3.10 (2H, m), 3.30-3.40 (4H, m), 3.71 (3H, s), 6.96-7.32 (7H, m), 10.31 (1H, s), 10.47 (1H, br s).	440
<i>N</i> -(3,4-Dichlorobenzyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide hydrochloride (E4)	2.80 (3H, s), 3.06-3.55 (8H, m), 3.86 (3H, s), 4.00 (2H, d $J=6.4$), 7.06 (1H, d, $J=8.6$), 7.18-7.22 (2H, m), 7.35-7.40 (2H, m), 7.50 (1H, d, $J=8.3$), 8.18 (1H, s), 10.83 (1H, s).	444
6-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E5)	2.68 (3H, s), 2.89-3.31 (12H, m), 3.74 (3H, s), 4.12 (2H, t, $J=16.5$), 7.01-7.11 (6H, m), 7.38-7.48 (2H, m), 9.15 (1H, s), 10.91 (1H, s).	436
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzene-	2.64 (3H, s), 2.80-3.03 (6H, m), 3.20-3.39 (4H, m), 3.68 (3H, s), 3.74-3.81	456

sulfonyl]-6-trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E6)	(2H, t, J=8.6), 6.95 (1H, d, J=2.2), 7.02-7.07 (1H, d, J=13.0), 7.21-7.22 (2H, d, J=2.1), 7.30-7.35 (1H, dd, J=2.2, 8.6), 7.50 (1H, s), 10.65 (1H, s).	
<i>N</i> -(3-Chlorophenyl)-4-methoxy- <i>N</i> -methyl-3-(4-methylpiperazin-1-yl)-benzenesulfonamide (E7)	2.34 (3H, s), 2.57 (4H, s), 2.97 (4H, s), 3.10 (3H, s), 3.93 (3H, s), 6.87-6.90 (2H, m), 7.04-7.10 (2H, m), 7.18-7.31 (2H, m).	410
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline (E8)	1.59-1.69 (2H, m), 2.33 (3H, s), 2.41-2.52 (6H, br s), 2.90 (4H, s), 3.80 (2H, t, J=5.7), 3.89 (3H, s), 6.83 (1H, d, J=8.6), 6.94 (1H, d, J=2.2), 7.06-7.10 (1H, d, J=7.9), 7.29 (1H, s), 7.35 (1H, dd, J=2.3, 8.5), 8.16 (1H, s).	470

Table 2

Compound	MS (MH ⁺)
<i>N</i> -(3-Iodo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E9)	502
<i>N</i> -(5-Iodo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E10)	502
<i>N</i> -(3,4-Methylenedioxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E11)	406
6-Chloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methyl-2,3-dihydro-1 <i>H</i> -indole (E12)	436
7,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E13)	470/472
7,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E14)	462
5-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline (E15)	480/482
8-Chloro-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline (E16)	466
6,7-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-	462

benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E17)	
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-phenyl-1,2,3,4-tetrahydroisoquinoline hydrochloride (E18)	478
8-Bromo-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline (E19)	510/512
5,6-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E20)	470/472
5,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E21)	462
6-Iodo-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methylthio-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E22)	560
<i>N</i> -(3,4-Dichlorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E23)	430/432
<i>N</i> -(3-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E24)	488
6,7,8-Trimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline (E25)	492
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-6-pyridin-3-yl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E26)	465
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-5-pyridin-3-yl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E27)	465
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,5-tetrahydropyrrolo[2,3- <i>f</i>]indole hydrochloride (E28)	427
<i>N</i> -(2-Fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E31)	380
<i>N</i> -(2-Trifluoromethoxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E32)	446
<i>N</i> -(2-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E33)	454/456
<i>N</i> -(4-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E34)	488
<i>N</i> -(9,10-Dioxo-9,10-dihydroanthracen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E35)	492
<i>N</i> -(2-Hydroxymethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E36)	392
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(2-methylsulfanylphenyl)-benzenesulfonamide hydrochloride (E37)	408

4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(5,6,7,8-tetrahydronaphthalene-1-yl)-benzenesulfonamide hydrochloride (E38)	416
<i>N</i> -(2-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E39)	390
4-Methoxy- <i>N</i> -(2-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E40)	376
<i>N</i> -(3,4-Dimethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E41)	390
4-Methoxy- <i>N</i> -(2-methoxy-6-methylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E42)	406
<i>N</i> -(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E43)	457
8-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E44)	436/438
<i>N</i> -(2-Chloro-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E45)	414/416
<i>N</i> -(2-Trifluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E46)	430
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -quinolin-7-ylbenzenesulfonamide hydrochloride (E47)	413
<i>N</i> -(4-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E48)	440/442
<i>N</i> -(3-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E49)	454/456
<i>N</i> -(3-Bromo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E50)	454/456
4-Methoxy- <i>N</i> -(2-methoxydibenzofuran-3-yl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E51)	482
<i>N</i> -(4-Cyclohexylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E52)	444
<i>N</i> -(2-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E53)	488
<i>N</i> -(2-Chloro-4-iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E54)	522/524
<i>N</i> -(2-Bromo-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E55)	458/460
<i>N</i> -[4-(4-Chlorophenyl)thiazol-2-yl]-4-methoxy-3-(4-methylpiperazin-1-	479/481

yl)-benzenesulfonamide hydrochloride (E56)	
4-Methoxy- <i>N</i> -(3-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E57)	376
<i>N</i> -(3-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E58)	390
<i>N</i> -(3-Chloro-4-bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E59)	474/476
<i>N</i> -(2-Acetylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide hydrochloride (E60)	404
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(4-phenylaminophenyl)-benzenesulfonamide hydrochloride (E61)	453
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(4-pentyloxyphenyl)-benzenesulfonamide hydrochloride (E62)	448
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(4-vinylphenyl)benzenesulfonamide hydrochloride (E63)	388
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(2-pyrrol-1-ylphenyl)-benzenesulfonamide hydrochloride (E64)	427
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -[4-(4-nitrophenylsulfanyl)phenyl]-benzenesulfonamide hydrochloride (E65)	515
4-Methoxy-3-(4-methylpiperazin-1-yl)- <i>N</i> -(3-oxazol-5-ylphenyl)-benzenesulfonamide hydrochloride (E66)	429
<i>N</i> -(4-Bromo-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E67)	508/510
4-Methoxy- <i>N</i> -(2,3-dimethylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E68)	390
<i>N</i> -(4-Chloro-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide hydrochloride (E69)	464/466
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E70)	456
7-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E71)	480/482
5,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride (E72)	470/472
5,7-Dichloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline hydrochloride (E73)	470/472
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline (E75)	402

1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-6-methyl-1,2,3,4-tetrahydroquinoline (E76)	416
6-Fluoro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline (E77)	434

Example 78**5-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulphonyl]-2,3-dihydro -1*H*-isoindole hydrochloride (E78).**

- 5 Sodium hydride (20mg of a 60% dispersion in mineral oil, 0.5mmol) was added to a solution of 3- (4-methyl-1-piperazin-1-yl)-4-methoxybenzenesulphonamide (D16) (53mg, 0.19mmol) in DMF (2ml) in one portion at room temperature under argon. Stirring was continued for 1hr before a solution of 1,2 *bis* bromomethyl-4-chlorobenzene (110mg, 0.37mmol) in DMF (0.5ml) was added. The reaction was
- 10 heated at 60°C for 3hrs, cooled and then partitioned between water and dichloromethane. The organic phase was dried over sodium sulphate and concentrated *in vacuo*. The residue was purified by chromatography on silica gel to afford material which was converted to the title compound by treatment with 1M ethereal HCl. MS: m/z (MH⁺) = 422/424.

15

Example 29**1-(4-Methoxy-3-piperazin-1-ylbenzenesulfonyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline hydrochloride (E29)**

- A solution of 1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline (E8) (70mg, 0.15 mmol) and 1-chloro-ethyl chloroformate (0.08ml, 0.75 mmol) in 1,2-dichloroethane (2ml) was heated under reflux for 18h, and then was cooled to ambient temperature.
- N,N-Diisopropylethylamine (0.05ml, 0.26mmol) was added and the resulting solution was heated under reflux for 2h. The solvent was removed, the residue was dissolved
- 25 in methanol (4ml), and the reaction was heated under reflux for 18h. The solvent was partially removed, dichloromethane (20ml) was added, and the solution was washed with saturated aqueous sodium hydrogen carbonate (10ml), dried (MgSO₄), and then evaporated. The residue was purified by column chromatography on silica gel eluting with a methanol-dichloromethane gradient to afford the sulfonamide derivative as a
- 30 yellowish oil. The oil was dissolved in acetone (0.5ml) and 1M solution of hydrogen chloride in diethyl ether (0.1ml) was added. The solution was evaporated, and the residue was coevaporated with dry benzene (3 x 2ml) to give the title compound (E25) as a cream solid (38mg, 52%).

δ_H (250MHz, DMSO- d_6), 1.58 (2H, m), 2.51 (2H, m), 3.01 (4H, br s), 3.16 (4H, br s), 3.79 (2H, t, $J=5.80$), 3.85 (3H, s), 6.83 (1H, d, $J=2.10$), 7.13 (1H, d, $J=8.74$), 7.33 (2H, m), 7.44 (1H, d, $J=8.11$), 7.95 (1H, s), 9.1 (2H, br s). MS: m/z (MH^+) = 456.

5 Example 30

N-(3-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E30)

The title compound was prepared using a similar procedure to that of example E29 using *N*-(3-bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide (E3) and 1-chloroethyl chloroformate.

10 δ_H (250MHz, DMSO- d_6), 2.84 (8H, m), 3.81 (3H, s), 7.01-7.20 (5H, m), 7.23 (1H, m), 7.36 (1H, m). MS: m/z (MH^+) = 426.

Example 79

N-(2-Isopropylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide

15 hydrochloride (E79)

Pyridine (0.28ml) was added to a stirred solution of 2-isopropylaniline (98mg) and 3-(4-trichloroacetyl)piperazin-1-yl)-4-methoxybenzenesulfonyl chloride (D4) (300mg) in dichloromethane (4ml) at room temperature. After 18h the solution was washed with 1M hydrochloric acid (5ml) then water (5ml). The organic phase was stirred
20 vigorously with 20% aq. potassium hydroxide (0.5ml) for 18h. A 10% aqueous solution of KH_2PO_4 (8ml) was then added to the mixture and after 0.25h stirring the layers were separated. The organic layer was dried (Na_2SO_4), acidified with 1M ethereal hydrogen chloride (2ml) and concentrated to an oil which was stirred with acetone/diethyl ether to afford the title compound (E79) as a white solid (0.224g,
25 83%). MH^+ 390.

The compounds in Table 3 were prepared in a similar manner to Example 79 using 3-(4-trichloroacetyl)piperazin-1-yl)-4-methoxybenzenesulfonyl chloride (D4) and the appropriate amine. All amines are either commercially available or can be prepared by
30 methods described above.

Table 3

Compound	MS (MH^+)
<i>N</i> -(4-Chloronaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E80)	432/434
4-Methoxy- <i>N</i> -naphthalen-1-yl-3-piperazin-1-ylbenzenesulfonamide	398

hydrochloride (E81)	
<i>N</i> -(3-Chloro-2-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E82)	396/398
<i>N</i> -Indan-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E83)	388
<i>N</i> -(2-Fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E84)	366
4-Methoxy- <i>N</i> -(2-methylsulfanylphenyl)-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E85)	394
4-Methoxy-3-piperazin-1-yl- <i>N</i> -(2-trifluoromethylphenyl)benzenesulfonamide hydrochloride (E86)	416
4-Methoxy- <i>N</i> -(2-methylphenyl)-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E87)	362
<i>N</i> -(2-Ethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E88)	376
4-Methoxy-3-piperazin-1-yl- <i>N</i> -(3-trifluoromethylphenyl)benzenesulfonamide hydrochloride (E89)	416
<i>N</i> -(3,4-Dimethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E90)	376
<i>N</i> -(2-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E91)	426/428
<i>N</i> -(3,4-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E92)	416/418
<i>N</i> -(3-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E93)	474
<i>N</i> -(3,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E94)	416/418
<i>N</i> -(3-Chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E95)	382/384
<i>N</i> -(2-Chloro-3-fluoro-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E96)	414/416
<i>N</i> -(4-Chloro-3-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E97)	396/398
<i>N</i> -Benzol[1,3]dioxol-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E98)	392
<i>N</i> -(2-Bromo-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E99)	440/442

<i>N</i> -(2,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E100)	504/506
<i>N</i> -(2,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E101)	416/418
<i>N</i> -(2-Chloro-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E102)	396/398
<i>N</i> -(4-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E103)	426/428
<i>N</i> -(2-Isopropenylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E104)	388
4-Methoxy- <i>N</i> -(2-methyl-5-nitrophenyl)-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E105)	407
<i>N</i> -(4-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E106)	474
<i>N</i> -(4- <i>tert</i> -Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E107)	404
<i>N</i> -(4-Isopropylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E108)	390
<i>N</i> -(4-Hexylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E109)	432
<i>N</i> -(2,4-Dibromonaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E110)	554/556
4-Methoxy- <i>N</i> -(4-methoxybiphenyl-3-yl)-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E111)	454
<i>N</i> -(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E112)	443
<i>N</i> -Biphenyl-2-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E113)	424
<i>N</i> -(2-Benzylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E114)	438
<i>N</i> -(2-Propylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E115)	390
<i>N</i> -(2- <i>sec</i> -Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E116)	404
<i>N</i> -(2- <i>tert</i> -Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E117)	404
<i>N</i> -(2-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide	404

hydrochloride (E118)	
<i>N</i> -(5-Iodo-2-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E119)	488
<i>N</i> -(2-Chloro-3,5-difluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E129)	418/420
<i>N</i> -(4-Chloro-2-trifluoromethoxyphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E130)	466/468
<i>N</i> -(2,4,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E131)	450/452
<i>N</i> -(5-Chloro-2-methoxyphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E132)	412/414
<i>N</i> -(4-Chloro-2-trifluoromethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E133)	450/452
<i>N</i> -(3,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E134)	506/508
<i>N</i> -(3-Bromo-2,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E135)	494/496
<i>N</i> -(2,3,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E136)	450/452
<i>N</i> -(5-Bromo-2,3-dihydro-benzofuran-7-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide hydrochloride (E137)	468/470
<i>N</i> -(2-Bromo-3,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E138)	
<i>N</i> -(3-Bromo-5,6-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E139)	
<i>N</i> -(2,5-Dibromo-3-fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E140)	
<i>N</i> -(2,5-Dibromo-3-chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E141)	
<i>N</i> -(2,3,5-Tribromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide (E142)	

Example 120**6-Chloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-2,3-dihydro-1H-indole hydrochloride (E120)**

- 5 Pyridine (0.28ml) was added to a stirred solution of 6-chloro-5-methyl-2,3-dihydro-1H-indole (WO-9501976) (0.116g) and 3-(4-trichloroacetyl piperazin-1-yl)-4-

methoxybenzenesulfonyl chloride (D4) (300mg) in dichloromethane (4ml) at room temperature. After 18h the solution was washed with 1M hydrochloric acid (5ml), water (5ml), dried (MgSO₄) and concentrated to an oil. The oil was dissolved in 1,4-dioxan (13ml) and a 0.15M potassium hydroxide solution (6.5ml) was added. The solution was stirred at room temperature for 4h then concentrated to remove the organic solvent, diluted with water (10ml) and the solution extracted with dichloromethane (20ml). The organic phase was dried (Na₂SO₄), acidified with 1M ethereal hydrogen chloride (2ml), concentrated to a solid and stirred with acetone to afford the title compound (0.175g, 55%). MH⁺ 422/424.

10

The compounds in Table 4 were prepared in a similar manner to Example 120 using 3-(4-trichloroacetyl)piperazin-1-yl)-4-methoxybenzenesulfonyl chloride (D4) and the appropriate amine. All amines are either commercially available or can be prepared by methods described above.

15

Table 4

Compound	MS (MH ⁺)
6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methylsulfanyl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E121)	546
6-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline hydrochloride (E122)	466/468
8-Chloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride (E123)	422/424
1-(4-Methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-6-trifluoromethyl-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E124)	456
6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E143)	500
5-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1 <i>H</i> -indole hydrochloride (E144)	500
7-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline hydrochloride (E145)	466/468

Example 125

20 **5,8-Dimethoxy-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride (E125)**

A solution of 1-chloroethylchloroformate (0.12ml;1.11mmol) and 5,8-dimethoxy-2-[4-methoxy-3-(4-methyl-1-piperazinyl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline (E21) (111mg;0.223mmol) in 1,2-dichloroethane (3ml) was refluxed for 0.75h, cooled, diluted with diisopropylethylamine (0.19ml;1.11mmol) and refluxed for a further 2.5hrs. The solution was concentrated to a residue which was re-dissolved in methanol, refluxed for 1hr and then stirred at room temperature for 24h. The mixture was concentrated, and the residue partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was dried, concentrated to a residue and purified by column chromatography on silica gel using a methanol/dichloromethane solvent gradient. The title compound was prepared by dissolving the pure free base material from chromatography in acetone/dichloromethane and acidifying with 1M ethereal HCl (53mg, 49%). MH^+ 456/458.

Example 126

5,8-Dichloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride (E126)

The title compound was prepared from its N-methylpiperazine analogue (E72) by a method as described in E125, but the reaction was kept at ambient temperature and the diisopropylethylamine was added at the start of the reaction. MH^+ 456/458.

Example 127

N-(3-Iodo-4-methylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide hydrochloride (E127)

The title compound was prepared from its N-methylpiperazine analogue (E9) by a method as described in E125. MH^+ 488.

Example 128

5,7-Dichloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline hydrochloride (E128)

The title compound was prepared from its N-methylpiperazine analogue (E73) by a method as described in E125. MH^+ 456/458.

Pharmacological data

Method for assay of 5-HT₆ antagonistic activity:

The test compounds were dissolved in polyethylene glycol:dimethyl sulphoxide (1:1) at 1 or 10mM and diluted to 0.1mM using 5mM tris buffer (pH 7.7 @ 25°C).

Dissolution was assisted by addition of 0.02ml 5M HCl plus heating to 40°C and sonication for 10 minutes. Serial dilutions of drugs in the same buffer were carried out using either a TECAN 5052 or Biomek 2000 Workstation. Samples of the diluted test compounds (0.05ml) were mixed with 0.05ml of radio-ligand [³H]-LSD prepared in the incubation buffer, and 0.4ml of a suspension of a preparation of the washed membranes of HeLa_5HT6 cells (acquired from Dr. D. Sibley, NIH, Bethesda, see Ref 1)(see Table 1), also in the incubation buffer. The details of the incubation conditions for each assay are shown in Table 2. The incubation buffer was 50mM Trizma (Sigma, UK) pH7.7 @ 25°C, 4mM MgCl₂.

10

After incubation at 37°C, the mixtures were filtered using a Packard Filtermate in Packard TopCount format. Filters were washed with 4 x 1ml aliquots of ice-cold incubation buffer. Filters were dried and impregnated with 0.04ml of Microscint 20 (Packard). IC₅₀ values were estimated from the counts per minute using a four parameter logistic curve fit within EXCEL (2). K_i values were calculated using the method of Cheng and Prusoff (3). pIC₅₀ and pK_i are the negative log₁₀ of the molar IC₅₀ and K_i respectively.

15

Table 1 Details of the methods used to prepare membranes for binding assays

1st resuspension cells/ml	spin / resuspension 1, 2, 3	Incubation before final spin	protein conc. in stored aliquots	cells /ml in stored aliquots
7 x 10 ⁷	Yes	20min at 37°C	4mg/ml	1.0 x 10 ⁸

Table 2 Summary of receptor binding assay conditions

protein (ug/sample)	radio-ligand [³ H]-LSD (nM)	Specific Activity (Ci/mmol)	Non-Specific Definition	Kd (nM)
40	2.0	83	Methiothepin	3.1

20

References

1. MONSMA, F.J., SHEN, Y., WARD, R.P., HAMBLIN, M.W., SIBLEY, D.R.. 1993. Cloning and expression of a novel serotonin receptor with high affinity for tricyclic psychotropic drugs. *Mol. Pharmacol.*, **43**, 320-327.

25


2. BOWEN, W.P., JERMAN, J.C.. 1995. Nonlinear regression using spreadsheets. *Trends in Pharmacol. Sci.*, **16**, 413-417.
3. CHENG, Y.C., PRUSSOF, W.H.. 1973. Relationship between inhibition constant (K_i) and the concentration of inhibitor which causes 50% inhibition (IC₅₀) of an enzymatic reaction. *Biochem. Pharmacol.*, **92**, 881-894.

All compounds tested showed good selective 5-HT₆ receptor antagonist activity, having pK_i values 7.5 - 9.5 at human cloned 5-HT₆ receptors. Particularly preferred compounds demonstrated pK_i > 8.5 and selectivity > 100. Examples of such compounds include:

3, 8, 21, 29, 32, 37-39, 41, 44, 45, 53, 54, 57-59, 63, 67, 69, 72, 73, 79, 85, 88, 89, 91, 93-95, 100, 104, 107, 113, 117-119, 121-128, 131, 132, 134-143, 145.

Claims:

1. A compound of formula (I) or a salt thereof:



- 10 wherein:

P is phenyl, naphthyl, anthracenyl, a bicyclic heterocyclic ring, a tricyclic heteroaromatic ring or is a 5 to 7-membered heterocyclic ring each containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulphur;

A is a single bond, a C₁₋₆alkylene or a C₁₋₆alkenylene group;

- 15 B is SO₂;

R¹ is halogen, C₁₋₆alkyl optionally substituted by one or more fluorine atoms, C₃₋₆cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkanoyl, C₁₋₆alkoxy, OCF₃, hydroxy, hydroxyC₁₋₆alkyl, hydroxyC₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkoxy, nitro, cyano, NR¹⁰R¹¹ where R¹⁰ and R¹¹ are independently hydrogen, C₁₋₆alkyl or optionally substituted phenyl, SR¹¹ where R¹¹ is as defined above or R¹ is optionally substituted phenyl, naphthyl, a bicyclic heterocyclic ring or is a 5 to 7-membered heterocyclic ring each containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulphur, or R¹ together with a second R¹ substituent forms a group -O-CH₂-O-, OCH₂CH₂O-, -CH₂CH₂CH₂- or -CH₂CH₂CH₂CH₂-,

- 25 n is 0, 1, 2, 3, 4, 5 or 6;

R² is hydrogen, C₁₋₆ alkyl, aryl C₁₋₆ alkyl or together with group P form a 5 to 8 membered ring optionally substituted with one or more C₁₋₆alkyl groups;

R³ is hydrogen, halogen, C₁₋₆alkyl, C₃₋₆cycloalkyl, C₁₋₆alkanoyl, C₁₋₆alkoxy optionally substituted with one or more fluorine atoms, hydroxy, hydroxyC₁₋₆alkyl,

- 30 hydroxyC₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkoxy, , nitro, trifluoromethyl, cyano or aryl or together with the group R⁵ forms a group (CH₂)₂O or (CH₂)₃O optionally substituted with 1 or more C₁₋₆alkyl groups;

R⁴ is -X(CH₂)_p-R⁶ where X is a single bond, CH₂, O, NH or N-alkyl and p is 0 to 6 and R⁶ is an optionally substituted 4- to 7-membered heterocyclic ring containing 1 to

3 heteroatoms selected from nitrogen, sulphur or oxygen, or R⁶ is NR⁷R⁸ where R⁷ and R⁸ are independently hydrogen, C₁₋₆ alkyl or aryl C₁₋₆ alkyl; and R⁵ is a group R³ or together with R³ forms a group (CH₂)₂O or (CH₂)₃O optionally substituted with 1 or more C₁₋₆alkyl groups.

5

2. A compound according to claim 1 in which P is phenyl or naphthyl.
3. A compound according to claim 1 or 2 in which R¹ is hydrogen, halogen, C₁₋₆alkoxy or C₁₋₆alkyl optionally substituted by one or more halogen atoms.
4. A compound according to any one of claims 1 to 3 in which R⁴ is an optionally substituted piperazine ring.
- 10 5. A compound according to any one of claims 1 to 4 in which R⁵ is methoxy.
6. A compound according to claim 1 which is:
 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-naphthalen-1-ylbenzenesulfonamide,
 15 *N*-(4-Chloronaphthalen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide,
N-(3-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide,
N-(3,4-Dichlorobenzyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzene sulfonamide,
 6-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-
 20 tetrahydroisoquinoline,
 1-[4-Methoxy-3-(4-methyl-piperazin-1-yl)-benzene-sulfonyl]-6-trifluoromethyl-2,3-dihydro-1*H*-indole,
N-(3-Chlorophenyl)-4-methoxy-*N*-methyl-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
 25 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline,
N-(3-Iodo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(5-Iodo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-
 30 benzenesulfonamide,
N-(3,4-Methylenedioxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
 6-Chloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methyl-2,3-dihydro-1*H*-indole,
 35 7,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
 7,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,

- 5-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
8-Chloro-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
5 6,7-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-7-phenyl-1,2,3,4-tetrahydroisoquinoline,
8-Bromo-7-methoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-
10 1,2,3,4-tetrahydroisoquinoline,
5,6-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
5,8-Dimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
15 6-Iodo-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-methylthio-2,3-dihydro-1*H*-indole,
N-(3,4-Dichlorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
6,7,8-Trimethoxy-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,4-
20 tetrahydroisoquinoline,
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-6-pyridin-3-yl-2,3-dihydro-1*H*-indole,
2-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-5-pyridin-3-yl-2,3-dihydro-1*H*-indole,
25 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonyl]-1,2,3,5-tetrahydropyrrolo[2,3-*f*]indole,
1-(4-Methoxy-3-piperazin-1-yl)benzenesulfonyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline,
N-(3-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
30 *N*-(2-Fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(2-Trifluoromethoxyphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
35 *N*-(4-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(9,10-Dioxo-9,10-dihydroanthracen-1-yl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,


- N*-(2-Hydroxymethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(2-methylsulfanylphenyl)-benzenesulfonamide,
5 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(5,6,7,8-tetrahydronaphthalene-1-yl)-benzenesulfonamide,
N-(2-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
4-Methoxy-*N*-(2-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3,4-Dimethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
10 4-Methoxy-*N*-(2-methoxy-6-methylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
8-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-
15 tetrahydroisoquinoline,
N-(2-Chloro-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
20 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-quinolin-7-ylbenzenesulfonamide,
N-(4-Bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3-Bromo-4-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(3-Bromo-2-methylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-
25 benzenesulfonamide,
4-Methoxy-*N*-(2-methoxydibenzofuran-3-yl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(4-Cyclohexylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
30 *N*-(2-Chloro-4-iodophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Bromo-4-fluorophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-[4-(4-Chlorophenyl)thiazol-2-yl]-4-methoxy-3-(4-methylpiperazin-1-yl)-
35 benzenesulfonamide,
4-Methoxy-*N*-(3-methylphenyl)-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
N-(3-Ethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,

- N*-(3-Chloro-4-bromophenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(2-Acetylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-phenylaminophenyl)-benzenesulfonamide,
5 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-pentyloxyphenyl)-benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(4-vinylphenyl)benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(2-pyrrol-1-ylphenyl)-benzenesulfonamide,
4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-[4-(4-nitrophenylsulfanyl)phenyl]-benzenesulfonamide,
10 4-Methoxy-3-(4-methylpiperazin-1-yl)-*N*-(3-oxazol-5-ylphenyl)-benzenesulfonamide,
N-(4-Bromo-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
4-Methoxy-*N*-(2,3-dimethylphenyl)-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
N-(4-Chloro-3-trifluoromethylphenyl)-4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulfonamide,
15 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-5-trifluoromethyl-2,3-dihydro-1*H*-indole,
7-Bromo-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
20 5,8-Dichloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroisoquinoline,
5,7-Dichloro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
25 1-[4-Methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-6-methyl-1,2,3,4-tetrahydroquinoline,
6-Fluoro-1-[4-methoxy-3-(4-methylpiperazin-1-yl)benzenesulfonyl]-1,2,3,4-tetrahydroquinoline,
5-Chloro-2-[4-methoxy-3-(4-methylpiperazin-1-yl)-benzenesulphonyl]-2,3-dihydro -
30 1*H*-isoindole hydrochloride,
N-(2-Isopropylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide hydrochloride,
N-(4-Chloronaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-*N*-naphthalen-1-yl-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Chloro-2-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
35 *N*-Indan-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-*N*-(2-methylsulfanylphenyl)-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-3-piperazin-1-yl-*N*-(2-trifluoromethylphenyl)benzenesulfonamide,

- 4-Methoxy-*N*-(2-methylphenyl)-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Ethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-3-piperazin-1-yl-*N*-(3-trifluoromethylphenyl)benzenesulfonamide,
N-(3,4-Dimethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
5 *N*-(2-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3,4-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
10 *N*-(2-Chloro-3-fluoro-4-methylphenyl)-4-methoxy-3-piperazin-1-yl-
benzenesulfonamide,
N-(4-Chloro-3-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-Benzol[1,3]dioxol-5-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Bromo-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
15 *N*-(2,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Chloro-4-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Bromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Isopropenylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
20 4-Methoxy-*N*-(2-methyl-5-nitrophenyl)-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Iodophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-*tert*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Isopropylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Hexylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
25 *N*-(2,4-Dibromonaphthalen-1-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
4-Methoxy-*N*-(4-methoxybiphenyl-3-yl)-3-piperazin-1-ylbenzenesulfonamide,
N-(3-Fluoro-5-pyridin-3-ylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-Biphenyl-2-yl-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Benzylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
30 *N*-(2-Propylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-*sec*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-*tert*-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Butylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Iodo-2-methylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
35 6-Chloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-2,3-dihydro-1*H*-
indole hydrochloride,
6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methylsulfanyl-2,3-dihydro-
1*H*-indole,


- 6-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline,
 8-Chloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline,
 1-(4-Methoxy-3-piperazin-1-ylbenzenesulfonyl)-5-methyl-6-trifluoromethyl-2,3-dihydro-1*H*-indole,
 5,8-Dimethoxy-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride,
 5,8-Dichloro-2-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroisoquinoline hydrochloride,
 10 *N*-(3-Iodo-4-methylphenyl)-4-methoxy-3-piperazin-1-yl-benzenesulfonamide,
 5,7-Dichloro-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline,
N-(2-Chloro-3,5-difluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Chloro-2-trifluoromethoxyphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
 15 *N*-(2,4,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Chloro-2-methoxyphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(4-Chloro-2-trifluoromethylphenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(3,5-Dibromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide ,
 20 *N*-(3-Bromo-2,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,3,5-Trichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(5-Bromo-2,3-dihydro-benzofuran-7-yl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2-Bromo-3,5-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
 25 *N*-(3-Bromo-5,6-dichlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dibromo-3-fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,5-Dibromo-3-chlorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
N-(2,3,5-Tribromophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulfonamide,
 30 6-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1*H*-indole,
 5-Iodo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-2,3-dihydro-1*H*-indole,
 7-Bromo-1-(4-methoxy-3-piperazin-1-ylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline
 and pharmaceutically acceptable salts thereof.
7. A compound according to any one of claims 1 to 6 for use in therapy.
- 35 8. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 6 and a pharmaceutically acceptable carrier or excipient.

9. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises the coupling of a compound of formula (II):



(II)

in which R^1 , R^2 , n , P , and A are as defined in formula (I) or protected derivatives thereof with a compound of formula (III):



(III)

in which B , R^3 , R^4 and R^5 are as defined in formula (I) or protected derivatives thereof and L is a leaving group and optionally thereafter

- removing any protecting groups
- forming a pharmaceutically acceptable salt:

10. Use of a compound according to any one of the claims 1 to 6 for the manufacture of a medicament for the treatment of anxiety and/or depression.