



(86) Date de dépôt PCT/PCT Filing Date: 2003/02/28
(87) Date publication PCT/PCT Publication Date: 2003/09/12
(85) Entrée phase nationale/National Entry: 2004/08/11
(86) N° demande PCT/PCT Application No.: US 2003/006193
(87) N° publication PCT/PCT Publication No.: 2003/073991
(30) Priorités/Priorities: 2002/03/01 (60/361,258) US;
2002/08/27 (60/406,171) US; 2003/02/13 (60/447,393) US


(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/12, A61K 31/7105,
A61K 31/7088, A61K 31/58, A61K 31/575, A61K 31/573,
A61K 31/57, A61K 31/56, A61K 31/5575, A61P 19/10,
C12Q 1/02, C07K 14/47, G01N 33/53

(71) Demandeur/Applicant:
CELLTECH R & D, INC., US


(72) Inventeurs/Inventors:
KUNG SUTHERLAND, MAY S., US;
GEOGHEGAN, JAMES C., US;
YU, CHANGPU, US;
LATHAM, JOHN, US

(74) Agent: OYEN WIGGS GREEN & MUTALA


(54) Titre : PROCEDES DESTINES A ACCROITRE OU A REDUIRE LA DENSITE OSSEUSE
(54) Title: METHODS TO INCREASE OR DECREASE BONE DENSITY




A



C



B



D

(57) Abrégé/Abstract:

The SOST gene gives rise to sclerostin, a protein that leads to apoptosis of bone progenitor cells. The invention provides antagonists to the sclerostin protein, and methods for identifying new sclerostin antagonists. The invention also provides molecules that can depress expression of the SOST gene, as well as methods for identifying such molecules. Such molecules and antagonists are useful for increasing bone mineralization in mammals, for example, in the treatment of osteoporosis.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
12 September 2003 (12.09.2003)

PCT

(10) International Publication Number
WO 03/073991 A2(51) International Patent Classification⁷:**A61K**

(71) Applicants and

(21) International Application Number: PCT/US03/06193

(22) International Filing Date: 28 February 2003 (28.02.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/361,258	1 March 2002 (01.03.2002)	US
60/406,171	27 August 2002 (27.08.2002)	US
60/447,393	13 February 2003 (13.02.2003)	US

(72) Inventors: **KUNG SUTHERLAND, May, S.** [CA/US]; 614 213th Street SW, Bothell, WA 98021-8100 (US). **GE-OGHEGAN, James, C.** [US/US]; Mr. James C. Geoghegan, 502 13th Ave. East, Apt. 418, Seattle, WA 98102 (US). **YU, Changpu** [US/US]; 15520 30th Ave. SE, Mill Creek, WA 98012 (US). **LATHAM, John** [US/US]; 2409 10th Ave W., Seattle, WA 98119 (US).


(74) Agents: **STEFFEY, Charles, E.** et al.; P.O. Box 2938, Minneapolis, MN 55402 (US).

(71) Applicant (for all designated States except US): **CELL-TECH R & D, INC.** [US/US]; 1631 220th Street S.E., Bothell, WA 98021 (US).


(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,

[Continued on next page]


(54) Title: METHODS TO INCREASE OR DECREASE BONE DENSITY




A



C



B



D

(57) Abstract: The SOST gene gives rise to sclerostin, a protein that leads to apoptosis of bone progenitor cells. The invention provides antagonists to the sclerostin protein, and methods for identifying new sclerostin antagonists. The invention also provides molecules that can depress expression of the SOST gene, as well as methods for identifying such molecules. Such molecules and antagonists are useful for increasing bone mineralization in mammals, for example, in the treatment of osteoporosis.

WO 03/073991 A2

WO 03/073991 A2

SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHODS TO INCREASE OR DECREASE BONE DENSITY

This application claims priority from U.S. Application Ser. No.
5 60/361,258 filed March 1, 2002, from U.S. Application Ser. No. 60/406,171
filed August 27, 2002 and from U.S. Application Ser. No. _____ filed
February 13, 2003.

FIELD OF THE INVENTION

10 The invention provides compositions and methods for modulating bone
density. Methods for identifying new compounds that can modulate bone
density are also provided.

BACKGROUND OF THE INVENTION

15 Osteoporosis is a bone disorder characterized by the loss of bone mass,
which leads to fragility and porosity of the bone of man. As a result, patients
suffering from osteoporosis have an increased fracture risk of the bones.
Postmenopausal women are particularly at risk for osteoporosis as a result of
reduced levels of estrogen production. When administered at low levels,
20 estrogens have a beneficial effect on the loss of bone. However, estrogen
replacement therapy can have unwanted side effects including an increased risk
of blood clots, breast carcinomas, endometrium hyperplasia, and an increased
risk of endometrium carcinomas. The remaining current therapies provide little
in terms of generating new bone for osteoporotic patients. Hence, a need exists
25 for an alternative treatment of osteoporosis.

Sclerosteosis is a genetic disease resulting in increased bone formation
and the development of strong skeletons in humans. Sclerosteosis is caused by a
loss of function or null mutation in the SOST gene (Beighton et al. 1976;
Brunkow et al. 2001; Balesman et al. 2001). The majority of affected
30 individuals have been reported in the Afrikaner population of South Africa,
where a high incidence of the disorder occurs as a result of a founder effect.

Homozygosity mapping in Afrikaner families along with analysis of historical recombinants localized sclerosteosis to an interval of approximately 2 cM between the loci D17S1787 and D17S930 on chromosome 17q12-q21. Affected Afrikaners carry a nonsense mutation near the amino terminus of the encoded protein, whereas an unrelated affected person of Senegalese origin carries two splicing mutations within the single intron of the gene.

The SOST gene encodes a protein called sclerostin that shares some sequence similarity with a class of cystine knot-containing factors including dan, cerberus, gremlin, prdc, and caronte. The sclerostin protein gene is thought to interact with one or more of the bone morphogenetic proteins (BMPs) (Brunkow et al, 2001). Bone morphogenetic proteins are members of the transforming growth factor (TGF- β) superfamily that have been shown to play a role in influencing cell proliferation, differentiation and apoptosis of many tissue types including bone. Bone morphogenetic proteins can induce *de novo* cartilage and bone formation, and appear to be essential for skeletal development during mammalian embryogenesis (Wang 1993). Early in the process of fracture healing the concentration of bone morphogenetic protein-4 (BMP-4) increases dramatically (Nakase et al. 1994 and Bostrom et al. 1995). In vivo experiments indicate that up-regulation of BMP-4 transcription may promote bone healing in mammals (Fang et al. 1996). Bone morphogenetic proteins have been reported to induce the differentiation of cells of the mesenchymal lineage to osteogenic cells as well as to enhance the expression of osteoblastic phenotypic markers in committed cells (Gazzerro et al. 1998, Nifuji & Noda, 1999). The activities of bone morphogenetic proteins in osteoblastic cells appear to be modulated by proteins such as noggin and gremlin that function as bone morphogenetic protein antagonists by binding and inactivating bone morphogenetic proteins (Yamaguchi et al. 2000).


However, the cascade of events leading to bone mineralization and the factors that control bone density are not completely understood. A need exists for factors that can modulate the differentiation of osteoblastic cells, promote bone mineralization and improve bone density.

SUMMARY OF THE INVENTION

According to the invention, the sclerostin protein encoded by the SOST gene is a unique bone morphogenetic protein antagonist and an important
 5 regulator of bone matrix formation. The SOST gene is expressed at high levels by committed, mature osteoblasts but at low or non-existent levels in immature and undifferentiated osteoblasts. Also according to the invention, the sclerostin protein blocks the ability of osteoprogenitor cells to differentiate, to deposit mineral and to express proteins characteristic of fully mature osteoblasts, for
 10 example, collagen and alkaline phosphatase. The expression of SOST is modulated by steroids and is regulated by bone morphogenetic proteins, for example, bone morphogenetic factors 2, 4 and 6. While bone morphogenetic factors increase the activity of osteoblastic markers such as alkaline phosphatase, the SOST gene product (sclerostin) blocks the activity of such osteoblastic
 15 markers. Moreover, bone morphogenetic factors can reverse the sclerostin-induced decrease in osteoblastic activity.

The invention provides pharmaceutical compositions for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a compound of the formula I:

20




wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl. In some embodiments, the R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ groups can

separately be carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl. In other embodiments, R_1 is carbonyl or hydroxy and/or R_8 is fluorine, hydrogen or halo.

The invention further provides a pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a glucocorticoid. The glucocorticoid can be, for example,
 5 fluocinolone acetonide, triamcinolone or dexamethasone.


The invention also provides a pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a prostaglandin comprising formula II:



10

wherein R_{10} and R_{11} are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl, aryl or aryloxy. R_{10} and R_{11} can also separately be carbonyl, hydroxy, hydrogen, or lower hydroxy
 15 alkyl. The prostaglandin can be, for example, prostaglandin E2.

The invention also provides a pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a bile salt of formula III:



wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈ and R₉ are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl. R₁, R₂, R₃, R₄, R₅, R₆, R₇ and R₉ can also separately be carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl. An example of such a bile salt is ursodeoxycholic acid.

The invention also provides a pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a prostaglandin such as prostaglandin E₂, a bile salt such as ursodeoxycholic acid, or a glucocorticoid such as fluocinolone acetonide, triamcinolone or dexamethasone.

The invention further provides a method for decreasing SOST expression in a mammal that comprises administering to the mammal a therapeutically effective amount of a prostaglandin such as prostaglandin E₂, a bile salt such as ursodeoxycholic acid, or a glucocorticoid such as fluocinolone acetonide, triamcinolone or dexamethasone.

The invention also provides a method for increasing bone density in a mammal that comprises administering a therapeutically effective amount of ursodeoxycholic acid, fluocinolone acetonide, triamcinolone, prostaglandin E₂ or dexamethasone to the mammal. In one embodiment, the methods of the invention can be practiced with a steroid or a glucocorticoid hormone. Examples of such glucocorticoid hormones include cortisol, dexamethasone and structurally related compounds. An example of a therapeutically effective

amount of a glucocorticoid hormone is a blood concentration of about 0.01 micromolar to about 1 micromolar.

The invention further provides a pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective
5 amount of an anti-sense nucleic acid capable of hybridizing to SEQ ID NO:1, 2, 4, 5 or 7.

The invention also provides a pharmaceutical composition that comprises a therapeutically effective amount of a sclerostin antagonist. Such a sclerostin antagonist can be an antibody that can bind a sclerostin polypeptide having SEQ
10 ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8. Another example of such a sclerostin antagonist is a peptide of a bone morphogenetic protein that binds to sclerostin.

The invention further provides a method for identifying a factor that decreases SOST expression comprising: providing a cell that comprises a
15 nucleic acid construct comprising a SOST promoter having SEQ ID NO:4 that is operably linked to a nucleic acid encoding a detectable marker; contacting the cell with a test sample containing a factor; and detecting whether a factor in the test sample decreases expression of the detectable marker relative to a control that comprises the cell that has not been contacted with a test sample. For
20 example, such a cell line is any cell line responsive to a bone morphogenetic protein. Another example of such a cell is a C2C12 cell line having ATCC Deposit No. CRL-1772 or a C3H10T1/2 cell line having ATCC Deposit No. CCL-226 or a human mesenchymal (hMSC) cell. Another element that can be present in the nucleic acid construct is an osteoblastic-specific steroid response
25 element.

The invention also provides a method for identifying a factor that decreases SOST expression comprising contacting a cell with a test sample and detecting whether a factor in the test sample decreases endogenous SOST expression. In this method, SOST expression can be from a nucleic acid
30 comprising SEQ ID NO:1 or SEQ ID NO:2 or SEQ ID NO:5 or SEQ ID NO:7. In another embodiment of this method, SOST expression can be from a nucleic

acid encoding a protein comprising SEQ ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8.

The invention further provides a method for identifying an antagonist for sclerostin comprising contacting a cell with both sclerostin and a test sample and
5 detecting whether the test sample prevents apoptosis of the cell.

The invention also provides a method for identifying a molecule that binds to sclerostin comprising contacting a test sample with a sclerostin polypeptide comprising SEQ ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8 and determining whether a molecule in the test sample binds to the polypeptide.
10

DESCRIPTION OF THE FIGURES

The twenty-seven figures of the application illustrate the results of experiments that define the importance of SOST and its gene product, sclerostin, in the regulation of bone homeostasis. In general, Figures 1 to 9 show that
15 SOST is expressed by mature osteoblastic cells and that its expression is modulated by important bone growth factors and steroids. Figures 10 to 19 demonstrate the inhibitory effect of sclerostin on the activity of osteoblastic markers. Finally, Figures 20 to 27 show that sclerostin affects the viability and the survival of osteoblastic bone cells.

20 Figure 1A provides a Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) analysis of SOST expression compared to the expression of other factors. The RNA used as a template for RT-PCR was prepared from primary human osteoblasts, undifferentiated hMSC cells and hMSC cells differentiated to yield osteoblasts, chondrocytes and adipocytes. RT-PCR amplification products
25 were separated and visualized on an agarose gel. The cell type from which the RNA template was obtained is shown at the bottom of the gel shown in Figure 1A. Several sets of primers were used in separate RT-PCR reactions to compare the level of SOST expression in the various cell types with the level of expression of characteristic phenotypic markers for various cell types, including
30 parathyroid hormone receptor (PTHr), collagen (types I, X and IX), PPAR γ 2 (a characteristic phenotypic marker for adipocytes) and Defender Against Death

(DAD, a housekeeping gene). As can be observed in Figure 1A, undifferentiated hMSC cells and adipocytes expressed negligible levels of SOST. In contrast, the cell types capable of mineralization that express markers associated with the osteoblast and hypertrophic chondrocyte phenotype (e.g. type I collagen and type X collagen, respectively) did express SOST.

Figure 1B illustrates that SOST is expressed at higher levels in mature osteoblasts. Human MSC cells were grown in regular growth medium (Undifferentiated) or Osteoblast-Inducing medium (Differentiated) for 21 days. Matching cultures were stopped at 7 or 21 days and stained histochemically for alkaline phosphatase (ALP) or collagen (Masson Trichrome stain). Other cultures were processed for RT-PCR analysis of SOST expression. hMSC cells that had differentiated into osteoblasts were identified by enhanced histochemical staining for alkaline phosphatase (ALP) and collagen (Figure 1B, purple and blue stains respectively in the original photographs of the differentiated cells in the right panel at 21 days) and deposition of mineral (negligible at Day 7 versus 437 ± 60 μg calcium/mg protein at Day 28). Undifferentiated hMSC cells expressed negligible levels of SOST mRNA as well as negligible levels of alkaline phosphatase activity, and low collagen and mineral deposition (Figure 1B, left two panels). In contrast, the expression of alkaline phosphatase, collagen and SOST increased with time in cells grown in osteoblast-inducing media.

Figure 2 illustrates that SOST expression is up-regulated by bone morphogenetic proteins (BMPs). RNA was isolated for RT-PCR analysis of SOST expression from hMSC cells treated with various factors, including insulin-like growth factor-1 (IGF-1, 10 or 50 ng/ml), 1,25-dihydroxyvitamin D₃ (vit D, 10^{-7}M), dexamethasone (DEX, 10^{-7}M), BMP-6 (300 ng/ml), or BMP-4 (50 or 300 ng/ml), for 72 hrs. RNA was also isolated for RT-PCR analysis of SOST expression from cells grown for 7 or 21 days in regular growth media (Control) or Osteoblast-Inducing media (Induced). As illustrated, while untreated hMSC cells ("vehicle") strongly expressed type I collagen, such cells had negligible levels of PTHr and SOST (Vehicle and Control, 21 days). These

results indicate that untreated hMSC cells are in an early stage of osteoblast lineage, but are committed to osteoblastogenesis. Treatment of these cells with DEX, BMPs, IGF-1 (IGF-1 at 50 ng/ml) or long-term culture in osteoblast-inducing media (Induced, 21 days) advanced the stage of differentiation and induced PTHr expression. Several factors enhanced the expression of SOST, including BMP-4, BMP-6, and IGF-1 (50 ng/ml). Figure 3 illustrates that bone morphogenetic proteins up-regulate SOST expression and also provides a time course of such BMP-induced SOST expression in hMSC and primary human osteoblasts. Cells were treated with BMP-2 (100 ng/ml) or BMP-4 (50 ng/ml) for 24 to 120 hrs prior to harvesting for RNA and RT-PCR analyses. As shown, maximum levels of SOST expression were achieved within 72 to 120 hrs after the start of treatment.

Figure 4 illustrates the effects of bone morphogenetic proteins 2, 4 and 6 on the expression of BMP antagonists gremlin, noggin, and SOST in hMSC cells. hMSC cells were treated with BMPs 2, 4 and 6 (100 ng /ml) for 72 hrs prior to harvesting RNA and performing RT-PCR analyses. Two cDNA controls were run for the noggin analysis. While the basal levels of SOST and noggin were negligible in undifferentiated hMSC cells, the expression of SOST and noggin was up-regulated by BMPs-2, 4 and 6. In contrast, gremlin was constitutively expressed in undifferentiated hMSC cells and its expression was only modestly increased by the bone morphogenetic proteins.

Figure 5 illustrates that steroids can modulate the effect of bone morphogenetic protein-4 on SOST expression. RNA was isolated from hMSC cells treated for 72 hrs with 1,25-dihydroxyvitamin D3 (vit D, 10^{-7} M), retinoic acid (RA, 10^{-6} M), BMP-4 (250 ng/ml), dexamethasone (DEX, 10^{-7} M), or a combination of BMP-4 with these steroids. The isolated RNA was then analyzed for SOST by RT-PCR. BMP-4 alone enhanced SOST expression. Retinoic acid and vitamin D enhanced the levels of SOST when added in combination with BMP-4 -- the levels of SOST were greater when retinoic acid or vitamin D was present with BMP-4 than when in cells were treated with

BMP-4 alone. Dexamethasone abolished the stimulatory effect of BMP-4 on SOST expression.

Figure 6 provides a comparison of the effects of steroids on the expression of gremlin, noggin, and SOST in hMSC cells. hMSC cells were
5 treated for 72 hrs with BMP-4 (100 ng/ml), retinoic acid (RA, 10^{-6} M),
dexamethasone (DEX, 10^{-7} M), and 1,25-dihydroxyvitamin D3 (vit D, 10^{-7} M)
or a combination of BMP-4 with retinoic acid, dexamethasone or vitamin D.
RNA was then harvested from the cells and RT-PCR analyses were performed.
BMP-4 enhanced the expression of gremlin and noggin. The steroids had small
10 effects on BMP-induced noggin and gremlin expression. As illustrated, BMP-4
also significantly increased the levels of SOST expression. This effect on SOST
expression was enhanced by the presence of vitamin D or retinoic acid but was
depressed by the presence of dexamethasone.

Figure 7 provides a photograph of a gel illustrating the effects of
15 glucocorticoid analogs and a bile salt on the expression of SOST in hMSC cells.
hMSC cells were treated with BMP-6 (250 ng/ml) in the absence or presence of
dexamethasone (DEX, at 10^{-7} M), triamcinolone (Triam, at 10^{-7} M),
ursodeoxycholic acid (Urso-1 at 1×10^{-6} M or Urso-2 at 3×10^{-6} M) for 72 hrs.
RNA was isolated and analyzed by RT-PCR for SOST. The DAD gene was
20 used as a control for DNA loading. Dexamethasone, triamcinolone,
ursodeoxycholic acid blocked the induction of SOST by BMP-6.

Figure 8 provides a photograph of a gel illustrating the effects of
glucocorticoid analogs and a prostaglandin on the expression of SOST in hMSC
cells. hMSC cells were treated with BMP-6 (250 ng/ml) in the absence or
25 presence of dexamethasone (DEX at 10^{-7} M), tauroursodeoxycholic acid (Tauro
at 2×10^{-6} M), fluocinolone (Fluo at 10^{-7} M), lovastatin (Lova at 10^{-7} M), ICI
182,780 (ICI at 10^{-6} M), 17β -estradiol (Estrogen at 10^{-6} M) and prostaglandin E2
(PGE2 at 10^{-6} M) for 72 hrs. RNA was isolated and analyzed by RT-PCR for
SOST. The DAD gene was used as a control for DNA loading. DEX, Fluo and
30 PGE2 blocked the induction of SOST by BMP-6.

Figure 9A provides chemical structure of glucocorticoid, dexamethasone, that blocks BMP-induced SOST expression.

Figure 9B provides chemical structure of a glucocorticoid, triamcinolone, that blocks BMP-induced SOST expression.

5 Figure 9C provides chemical structure of a glucocorticoid, fluocinolone acetonide, that blocks BMP-induced SOST expression.

Figure 9D provides chemical structure of a bile salt, ursodeoxycholic acid, that blocks BMP-induced SOST expression.

10 Figure 9E provides chemical structure of an eicosanoid, prostaglandin E2, that blocks BMP-induced SOST expression.

Figure 10 provides a comparison of the effects of human sclerostin and mouse noggin proteins on alkaline phosphatase (ALP) activity in differentiating human MSC cells. ALP activity is a phenotypic marker for osteoblast differentiation. Human MSC cells were cultured for 7 days in Osteoblast-
15 Inducing media containing noggin (1 or 5 $\mu\text{g/ml}$) or partially-purified sclerostin protein (1 or 5 $\mu\text{g/ml}$). Cells were then harvested and analyzed for ALP activity. After 7 days of treatment, mouse noggin at 1 $\mu\text{g/ml}$ had no effect on ALP activity, whereas mouse noggin at 5 $\mu\text{g/ml}$ significantly increased ALP activity ($p < 0.001$ to vehicle-treated cells). Under the same conditions, partially purified
20 preparations of sclerostin consistently decreased ALP activity when administered at either 1 or 5 $\mu\text{g/ml}$.

Figure 11 illustrates that human sclerostin also decreased alkaline phosphatase activity in primary cultures of human osteoblasts. Primary human osteoblasts were grown in regular growth media or in Inducing media (to
25 promote differentiation) and then treated with Sclerostin (20 $\mu\text{g/ml}$) or Control (protein purified from Sf9 conditioned media) for 7 days prior to histochemical staining for alkaline phosphatase activity. Osteoblasts grown in regular growth media expressed low, basal levels of ALP ("No Treatment," far right panel). These levels increased significantly when cells were grown in Inducing media
30 (fourth panel, purple stain). When a similar group of cells were treated with sclerostin, there was a marked decrease in the amount of ALP expressed, as

evidenced by the reduction in purple stain (third panel, "Sclerostin + Inducing Media").

Figure 12A illustrates that human sclerostin decreases alkaline phosphatase activity in hMSC cells in a dose-dependent manner. hMSC cells were plated in Osteoblast-Inducing media. Partially-purified sclerostin protein or proteins purified from Sf9 conditioned media (control) were added the next day. Cultures were maintained for 7 days prior to analysis for alkaline phosphatase activity. As illustrated, increasing concentrations of sclerostin give rise to decreasing levels of alkaline phosphatase activity.

Figure 12B illustrates that human sclerostin decreases mineralization (calcium deposition) in hMSC cells in a dose-dependent manner. hMSC cells were plated in Osteoblast-Inducing media. Partially-purified sclerostin protein or proteins purified from Sf9 conditioned media (control) were added the next day. Cultures were maintained for 8 days, treated with sclerostin or control proteins and maintained for another 13 days before testing for mineralization. As illustrated, increasing concentrations of sclerostin give rise to diminished mineralization.

Figure 13 illustrates that partially purified preparations of human sclerostin reduce the synthesis of type I collagen by about 50 % in cells treated with 10 $\mu\text{g}/\text{ml}$ of sclerostin ($p < 0.001$ to control). hMSC cells were cultured in Osteoblast-Inducing media and treated with partially-purified sclerostin protein or proteins purified from Sf9 conditioned media (Control) for 21 days. Sclerostin or Control at the indicated concentrations was added 8 days after plating. Figure 14 illustrates that human sclerostin protein decreases expression of osteoblastic markers in cultures of differentiating hMSC cells. hMSC cells were plated in Osteoblast-Inducing media. After 3 days, partially purified sclerostin protein (10 $\mu\text{g}/\text{ml}$) or proteins purified from Sf9 conditioned media (control) were added. Cells were harvested after 40 hrs and RNA was prepared for RT-PCR analyses of osteoblastic markers. As can be seen, RNA levels of markers for adipocytes (PPAR γ 2) or undifferentiated mesenchymal cells (endoglin) were not affected by treatment with partially purified preparations of

sclerostin protein. On the other hand, RNA levels for osteoblast phenotypic markers such as PTHr, Type I collagen, BMP-2 and BMP-6 were significantly reduced in cells treated with partially purified preparations of sclerostin protein compared to cells treated with control.

5 Figure 15 illustrates that human sclerostin blocks BMP-6-stimulated activity as well as basal alkaline phosphatase activity in cultures of hMSC. Human MSC were plated in osteoblast-inducing media and treated with vehicle or BMP-6 (500 ng/ml) as well as increasing concentrations of human sclerostin-Flag protein. BMP-6 increased alkaline phosphatase activity. Sclerostin
10 decreased BMP-induced alkaline phosphatase activity as well as basal levels of alkaline phosphatase activity in a dose dependent manner ($p < 0.0001$).

 Figure 16 illustrates that BMP-6 reverses the inhibitory effect of sclerostin on alkaline phosphatase in cultures of hMSC. hMSC were plated in osteoblast-inducing media and treated with sclerostin (10 μ g/ml) and increasing
15 amounts of BMP-6. While sclerostin decreased basal levels of ALP, BMP-6 reversed this effect in a dose-dependent manner ($p < 0.001$).

 Figures 17A-17D illustrate that human sclerostin decreases alkaline phosphatase activity in mouse mesenchymal cells. Mouse mesenchymal C3H10T1/2 cells having ATCC Deposit No. CCL-226 were plated in regular
20 growth media. Human sclerostin, sclerostin immuno-depleted with an anti-Flag M2 antibody and mouse noggin were pre-incubated with 500 ng/ml BMP-6 for 1 hr prior to addition to cells. Cells were then incubated for 72 hrs with the BMP-6, sclerostin, immuno-depleted sclerostin or noggin and tested for alkaline
25 phosphatase activity. The stimulatory effect of BMP-6 (Figure 17A) on alkaline phosphatase activity in the C3H10T1/2 cells was antagonized by noggin (Figure 17B) and by sclerostin. To verify the specificity of the sclerostin response (Figure 17C), sclerostin protein that had been synthesized with a Flag-tag was immuno-depleted by pre-treatment with an anti-Flag M2 antibody/ agarose
30 complex prior to addition to the cells. As can be seen in Figure 17D, pre-treatment of the sclerostin preparation with an anti-Flag antibody completely

abolished the antagonism of BMP-6 in the C3H10T1/2 cells, indicating that a sclerostin-specific response exists.

Figure 18 provides a model of sclerostin activity. Bone morphogenetic proteins enhance the osteoblastic phenotype in osteoprogenitor cells by increasing the expression of osteoblastic markers (ALP, Collagen). Bone morphogenetic proteins also increase SOST levels. However, SOST feeds back to decrease Bone morphogenetic protein levels, resulting in decreased expression of osteoblastic markers. Steroids such as dexamethasone (and possibly estrogen, or androgen) bind to nuclear receptors (HR) and interact with the transcriptional machinery to reduce the production of SOST and therefore decrease sclerostin activity.

Figure 19 illustrates screening assays provided by the invention for identifying factors that may down-regulate SOST expression. In both assays, the premise is that compounds bind to glucocorticoid receptor (GR) and interact with the BMP machinery to down-regulate activity of the SOST promoter, resulting in less SOST being transcribed. An example of a control compound is dexamethasone (DEX). In Scenario A, bone cells that are responsive to bone morphogenetic proteins and to glucocorticoids may be used to screen chemical libraries for compounds that would block the BMP-induced increase in SOST expression as measured by RT-PCR assays for SOST expression. Compounds selected to be of interest would be those that decrease the expression of SOST. In Scenario B, cultured cells that had been transfected with a DNA construct may be used to screen chemical libraries of compounds. The DNA construct is comprised of the SOST promoter sequence, a glucocorticoid response element (GRE) and a detectable marker (such as luciferase), for example, the DNA construct would have the structure:

GRE -- SOST promoter – Luciferase

Compounds selected to be of interest would be those that decrease the expression of this DNA construct. Successful candidates from these screening assays can be tested in mesenchymal or osteoblastic cells to see if they can block endogenous sclerostin expression and/or increase osteoblastic activity in treated

cells. Figure 20 graphically illustrates that exposure to increasing amounts of human sclerostin decreases the viability of hMSC cells. hMSC cells were plated in Osteoblast-Inducing media and treated with baculo-expressed partially purified preparation of Sclerostin or proteins purified from Sf9 conditioned media (control) for 1 week. Sclerostin and Control were refreshed with each media change. After one week, cells were lysed and processed with Promega's CellTiterGlo Viability assay to detect what percentage of cells were viable. The viability of control and sclerostin-treated cells is plotted on the y-axis as the percentage of signal from vehicle-treated cells.

Figure 21 shows that sclerostin, but not other BMP antagonists, increases the apoptosis of hMSC. hMSC were plated in osteoblast-inducing media and treated with 20 µg/ml sclerostin, inactivated sclerostin or one the BMP antagonists noggin, chordin, gremlin and twisted gastrulation (Tsg). Caspase activity was used as a measure of apoptosis. Sclerostin increased caspase activity whereas an inactive preparation of sclerostin or the BMP antagonists had no significant effect on caspase activity.

Figure 22A graphically illustrates that exposure to increasing amounts of human sclerostin increases the apoptosis of hMSC cells in a dose-dependent manner. The amount of caspase activity was used as a measure of apoptosis. hMSC cells were plated in Osteoblast-Inducing media and treated with the indicated amounts of a partially purified preparation of sclerostin or proteins purified from Sf9 conditioned media (control) for 24 hrs. As shown, increasing amounts of sclerostin lead to increased caspase activity.

Figure 22B graphically illustrates that exposure to increasing amounts of human sclerostin increases the apoptosis of hMSC cells in a dose-dependent manner. The amount of histone-DNA fragments as determined by ELISA (Roche) was used as a measure of apoptosis. hMSC cells were plated in Osteoblast-Inducing media and treated with the indicated amounts of a partially purified preparation of sclerostin or proteins purified from Sf9 conditioned media (control) for 24 hrs. As shown, increasing amounts of sclerostin lead to greater amounts of histone-DNA fragments.

Figure 23 illustrates that hMSC cells exposed to human sclerostin undergo apoptosis, as detected by Tunel staining for apoptotic nuclei. In contrast, hMSC cells exposed to vehicle or conditioned media show little evidence of apoptosis. hMSC cells were plated in Osteoblast-Inducing media and
5 treated with a partially purified preparation of Sclerostin (20 μ g/ml), proteins purified from Sf9 conditioned media (Control), or dexamethasone (DEX, 1 μ M, positive control) for 24 hrs. Tunel staining for apoptotic nuclei (arrows point to positively stained nuclei) was performed using Roche's In Situ Cell Death Detection kit. Magnification, Bright field, 100X.

10 Figure 24A shows the time course of effect of human sclerostin on the apoptosis of hMSC cells. hMSC cells were plated in Osteoblast-Inducing Media and treated with sclerostin. Cells were harvested and assayed for caspase activity at the indicated times. Caspase activity remained elevated 72 hrs after treatment of hMSC cells with sclerostin.

15 Figure 24B illustrates that rat sclerostin also induced the apoptosis of hMSC cells. Human MSC cells were plated in Osteoblast-Inducing media and treated with human sclerostin or rat sclerostin for 24 hrs prior to assay for caspase activity (Roche).

Figure 25 shows that commercially available inhibitors of caspase
20 activity can block the sclerostin-induced apoptosis of hMSC cells. hMSC cells were plated in Osteoblast-Inducing media and treated with human sclerostin or proteins purified from Sf9 conditioned media (Control), in the presence or absence of caspase inhibitors for 24 hrs. Caspase activity was determined by Roche ELISA. Caspase inhibitors (Cas-1, Cas-3, Calbiochem) alone or in
25 combination with each other efficiently blocked the sclerostin-induced increase in caspase activity. Caspase inhibitors also blocked caspase activity induced by rat sclerostin in these cells.

Figure 26 provides a schematic diagram illustrating how sclerostin may induce apoptosis in maturing osteoblast cells. Mesenchymal cells develop along
30 a defined temporal pathway from inducible progenitor cells to mature osteoblasts and finally, into lining cells or osteocytes. As these cells progress, they express

phenotypic markers characteristic of each stage of differentiation (e.g. type I collagen for pre-osteoblasts, mineralization for mature osteoblasts). Sclerostin is highly expressed by maturing osteoblasts. Sclerostin regulates mineralization and hence bone formation by a negative feedback mechanism. Sclerostin
5 decreases osteoblastogenesis by increasing the apoptosis of cells at early stages of osteoblast differentiation. Signals from osteocytes / lining cells regulate this entire process. Sclerostin may be an integral connection for osteocytic regulation of bone remodeling.

Figure 27 provides one possible assay for sclerostin antagonists. In this
10 assay, a variety of cells may be tested separately. In one assay, immortalized or transformed bone cells can be cultured in the presence of sclerostin and a test compound or test sample. In another assay, primary osteoblastic or mesenchymal cells that can be induced to differentiate are exposed to a test compound or test sample. After a period of incubation, apoptosis is measured.
15 Factors that reduce apoptosis are candidate sclerostin antagonists that can be further characterized in a number of ways as illustrated herein.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides compositions and methods for modulating bone
20 density in a mammal. In particular, the invention provides factors that diminish SOST expression and antagonists of the SOST gene product, sclerostin. According to the invention, sclerostin decreases bone mineralization, collagen synthesis and alkaline phosphatase (ALP) activity in maturing bone cells in both direct and indirect ways. One way that sclerostin regulates bone density and
25 bone formation is by acting as an antagonist of bone morphogenetic proteins (BMPs). Another way that sclerostin regulates bone density and formation is through apoptosis of cells involved in bone mineralization and bone formation.

By modulating or diminishing the amount of sclerostin or by inhibiting its action, bone morphogenetic proteins can perform essential functions needed
30 for the commitment and differentiation of osteoprogenitor cells and pre-osteoblastic cells. Moreover, by administering antagonists of sclerostin-

mediated apoptosis, the survival and longevity of cells involved in bone mineralization can be increased. Such compositions and procedures provide therapeutic strategies for increasing bone mineralization and formation.

The invention further provides assays for identifying factors that
5 diminish SOST expression and/or antagonists that counteract the effects of sclerostin.

Bone cell development

A small population of pluripotent mesenchymal/stromal cells, called
10 inducible osteoprogenitor cells, can be induced to differentiate into osteoblastic cells (Pittenger, MF et al. 1999). These inducible osteoprogenitor cells develop and express certain phenotypic markers in a defined, sequential manner (Pockwinse et al. 1992; Lian et al. 1999). Osteoprogenitor cells express type I collagen whereas committed pre-osteoblasts and osteoblasts express many of the
15 phenotypic markers that are typically identified with a cell of the osteoblast lineage. These markers include type I collagen, alkaline phosphatase, parathyroid hormone receptor (PTHr) and osteopontin. In the terminal stage of osteoblast differentiation, osteocytes are surrounded by deposits of mineral as well as matrix proteins such as CD44 and osteocalcin. Therefore, the
20 development, growth and differentiation of osteoblastic precursor cells into mature osteoblasts occur in a defined, time-dependent manner (Pockwinse et al., 1992).

The expression of phenotypic markers that are characteristic of osteoblasts at different stages of differentiation is regulated by a variety of
25 factors, including the bone morphogenetic proteins (BMPs). It is thought that BMP-6 may initiate osteoblast differentiation at an early osteoprogenitor stage but that other bone morphogenetic proteins or growth factors may be necessary to continue the differentiation process (Gitelman et al., 1995; Boden et al., 1997). Certain studies indicate that BMP-2 and BMP-4 are needed for the
30 commitment and differentiation of osteoprogenitor and pre-osteoblastic cells to osteoblasts (Katagiri et al., 1990; Oreffo et al. 1999; Suzawa et al., 1999).

Antagonists of bone morphogenetic proteins, for example, noggin, gremlin, chordin and the Dan / cereberus family of proteins, have also been identified. However, the role that bone morphogenetic protein antagonists play in the regulation and modulation of bone morphogenetic protein action in osteoblastic cells is not well understood.

SOST and Sclerostin

The sequence of the human SOST gene is approximately 22 kb and can be found as Genbank Accession No. AF326736 in the NCBI nucleotide database (SEQ ID NO:1). See website at ncbi.nlm.nih.gov/Genbank/GenbankOverview. Co-owned and co-assigned U.S. Patent Nos. 6,395,511 and 6,489,445, incorporated by reference herein, provide additional sequence information and a further description of the properties of the SOST gene and its gene product (called beer therein).

The human SOST cDNA has Genbank Accession No. NM025237, and is provided for the convenience of the reader as follows (SEQ ID NO:2).

```

1  AGAGCCTGTG  CTACTGGAAG  GTGGCGTGCC  CTCCTCTGGC
41  TGGTACCATG  CAGCTCCCAC  TGGCCCTGTG  TCTCGTCTGC
81  CTGCTGGTAC  ACACAGCCTT  CCGTGTAGTG  GAGGGCCAGG
20  121  GGTGGCAGGC  GTTCAAGAAT  GATGCCACGG  AAATCATCCC
161  CGAGCTCGGA  GAGTACCCCG  AGCCTCCACC  GGAGCTGGAG
201  AACAACAAGA  CCATGAACCG  GGCGGAGAAC  GGAGGGCGGC
241  CTCCCCACCA  CCCCTTTGAG  ACCAAAGACG  TGTCCGAGTA
281  CAGCTGCCGC  GAGCTGCACT  TCACCCGCTA  CGTGACCGAT
25  321  GGGCCGTGCC  GCAGCGCCAA  GCCGGTCACC  GAGCTGGTGT
361  GCTCCGGCCA  GTGCGGCCCC  GCGCGCCTGC  TGCCCAACGC
401  CATCGGCCGC  GGCAAGTGGT  GGCGACCTAG  TGGGCCCCGAC
441  TTCCGCTGCA  TCCCCGACCG  CTACCGCGCG  CAGCGCGTGC
481  AGCTGCTGTG  TCCCGGTGGT  GAGGCGCCGC  GCGCGCGCAA
30  521  GGTGCGCCTG  GTGGCCTCGT  GCAAGTGCAA  GCGCCTCACC
561  CGCTTCCACA  ACCAGTCGGA  GCTCAAGGAC  TTCGGGACCG
601  AGGCCGCTCG  GCCGCAGAAG  GGCCGGAAGC  CGCGGCCCCG
641  CGCCCGGAGC  GCCAAAGCCA  ACCAGGCCGA  GCTGGAGAAC
681  GCCTACTAGA  GCCCGCCCCG  GCCCCTCCCC  ACCGGCGGGC
35  721  GCCCCGGCCC  TGAACCCGCG  CCCACATTT  CTGTCCTCTG
761  CGCGTGGTTT  GATTGTTTAT  ATTTCATTGT  AAATGCCTGC
801  AACCAGGGC  AGGGGGCTGA  GACCTTCCAG  GCCCTGAGGA
841  ATCCCGGGCG  CCGGCAAGGC  CCCCTCAGC  CCGCCAGCTG
881  AGGGGTCCCA  CGGGGCAGGG  GAGGGAATTG  AGAGTCACAG

```

```

    921 ACACTGAGCC ACGCAGCCCC GCCTCTGGGG CCGCCTACCT
    961 TTGCTGGTCC CACTTCAGAG GAGGCAGAAA TGGAAGCATT
1001 TTCACCGCCC TGGGGTTTTA AGGGAGCGGT GTGGGAGTGG
1041 GAAAGTCCAG GGACTGGTTA AGAAAGTTGG ATAAGATTCC
  5 1081 CCCTTGCACC TCGCTGCCCA TCAGAAAGCC TGAGGCGTGC
    1121 CCAGAGCACA AGACTGGGGG CAACTGTAGA TGTGGTTTCT
    1161 AGTCCTGGCT CTGCCACTAA CTTGCTGTGT AACCTTGAAC
    1201 TACACAATTC TCCTTCGGGA CCTCAATTTC CACTTTGTAA
    1241 AATGAGGGTG GAGGTGGGAA TAGGATCTCG AGGAGACTAT
10 1281 TGGCATATGA TTCCAAGGAC TCCAGTGCCT TTTGAATGGG
    1321 CAGAGGTGAG AGAGAGAGAG AGAAAGAGAG AGAATGAATG
    1361 CAGTTGCATT GATTCAGTGC CAAGGTCACT TCCAGAATTC
    1401 AGAGTTGTGA TGCTCTCTTC TGACAGCCAA AGATGAAAAA
    1441 CAAACAGAAA AAAAAAAGTA AAGAGTCTAT TTATGGCTGA
15 1481 CATATTTACG GCTGACAAAC TCCTGGAAGA AGCTATGCTG
    1521 CTTCCCAGCC TGGCTTCCCC GGATGTTTGG CTACCTCCAC
    1561 CCCTCCATCT CAAAGAAATA ACATCATCCA TTGGGGTAGA
    1601 AAAGGAGAGG GTCCGAGGGT GGTGGGAGGG ATAGAAATCA
    1641 CATCCGCCCC AACTTCCCAA AGAGCAGCAT CCCTCCCCCG
20 1681 ACCCATAGCC ATGTTTTTAA GTCACCTTCC GAAGAGAAGT
    1721 GAAAGGTTCA AGGACACTGG CCTTGCAGGC CCGAGGGAGC
    1761 AGCCATCACA AACTCACAGA CCAGCACATC CCTTTTGAGA
    1801 CACCGCCTTC TGCCCACCAC TCACGGACAC ATTTCTGCCT
    1841 AGAAAACAGC TTCTTACTGC TCTTACATGT GATGGCATAT
25 1881 CTTACACTAA AAGAATATTA TTGGGGGAAA AACTACAAGT
    1921 GCTGTACATA TGCTGAGAAA CTGCAGAGCA TAATAGCTGC
    1961 CACCCAAAAA TCTTTTTTGAA AATCATTTCC AGACAACCTC
    2001 TTA CTTTCTG TG TAGTTTTT AATTGTTAAA AAAAAAAGT
    2041 TTAAACAGA AGCACATGAC ATATGAAAGC CTGCAGGACT
30 2081 GGTCGTTTTT TTGGCAATTC TTCCACGTGG GACTTGTCCA
    2121 CAAGAATGAA AGTAGTGGTT TTTAAAGAGT TAAGTTACAT
    2161 ATTTATTTTC TCACTTAAGT TATTTATGCA AAAGTTTTTC
    2201 TTGTAGAGAA TGACAATGTT AATATTGCTT TATGAATTAA
    2241 CAGTCTGTTC TTCCAGAGTC CAGAGACATT GTTAATAAAG
35 2281 ACAATGAATC ATGACCGAAA GAAAAA AAAA
    2321 AAA

```

The SOST gene gives rise to the sclerostin protein. The human sclerostin protein has Genbank Accession No. NP079513 and is provided below for convenient reference as SEQ ID NO:3.

```

  1 MQLPLALCLV CLLVHTAFRV VEGQGWQAFK NDATEIIPEL
  41 GEYPEPPPEL ENNKTMNRAE NGGRPPHHPF ETKDVSEYSC
  81 RELHFTRYVT DGPCRSAPV TELVCSGQCG PARLLPNAIG
121 RGKWWRPSGP DFRCIPDRYR AQRVQLLCPG GEAPRARKVR
45 161 LVASCKCKRL TRFHNQSELK DFGTEAARPQ KGRKPRPRAR

```

201 SAKANQAELE NAY

The promoter for the human SOST gene is found within nucleotides 7960-11960 of SEQ ID NO:1. This promoter sequence is provided below as

5 SEQ ID NO:4.

```

7960 T
7961 GAATGCCACA GCTTGTGAGC AGCCAGCAGA AGGCAGGTAA
8001 AAGGCCTTCC TCACTGTCCT CGAAGGAACC AACTCTGCCA
8041 ATGTCTTGAA CTTGCACTCC TAACCTCCAG CTCTGTGAGA
10 8081 CGACTTCTGT TGTTTATGCC ACCCAATTTG TGGTACTTAG
8121 TTACGGCAGC CTCATCAAAG TACCACCTAT GGGAGCCTCT
8161 ATTTTGCAGG TGAGGGCGGG GACTGGGCTG AGTTTTCTGG
8201 AAAACAGCCC TGCAATACCC TCATCAGACC ACCAAACTCT
8241 TCACACTCCC TCAGACACAG CATTCACTTC CAGAAATAAC
15 8281 TCTAAAGTTT TGTTTTGTTT TTTTAAACTT TGTGGAATAC
8321 TACTCAGCCA AAAAAAAAAA AAAAGAAAAG AAAAAGGAAC
8361 ATGTTACTGA TATGTACAAC TTGGATAATG GAAATAATGC
8401 TGAGTGAAAA AAAAATCCCC AAAGGCTACA TACTAATTGA
8441 TTCCATTTAT ATAACCTTTT TTTTTTTTTT TTGAGACAGT
20 8481 CTCACTCTGT CACCCAGGCT GGAGCACAGT GCGCGGATCT
8521 CAGCTCACTG CAACTTCCGC CTCCTGAGTT CAAGCGATTC
8561 TCTTGCCCTA GCTTCCCAAG TAGCTGGGAT TACAGGTGCG
8601 TGCCACCATG CCCAGCTAAT TTTTGTATTT TTAGTAGAGA
8641 CAGGGTTTCG CCATGTTGGC CAGGCTGGTC TCGAACTCCT
25 8681 GACCTCAAGT GATCTGCCTG CCTTGGCTTC CCAAAGTGCT
8721 GGGATTACAT GAGTGAGCCA CCGCACCTGG TCGCATTTAT
8761 GTAACAATTT TGAAGTGAAA AAAAAAATGA CAGAAATGGA
8801 GATTAGATGA GTAGTTGCCA GGGGTTAGTT GTGGGAGGGA
8841 GGGAAAAGGA GGAAGGAGG TGGGCAACAG GAGAAAGACT
30 8881 TGTGGTCACA GAGCTGTGCT TTATCTTGAC TGTGGTGGAT
8921 CCCCAAATTT ACCCGTGACA AGATTGCATA GAACTAAGTA
8961 TACACACACG TGAATGCGTG TGCACGCACA CAGTAGAGGT
9001 TAAGCCACGG GAGATGTGGG TAAGATTGGT AAATTGTGTC
9041 AATATCAATA TCCTAGTTGT GATATTGTCC TATAGTTTCG
35 9081 CAAGGTGTTA TTGTTGTGGG AAAGTGGATA AAGGATACAC
9121 GGAGTCTGCA TTTTCTTTTT TTTATTTTAA TTTTTTGAGA
9161 CGGGGTCTCA CTCTGTCAAC CAGGCTGGAG TGCAGTGGCC
9201 CAAGTATGGC TCACTGCAGC CTCGACCTCA ACCTCAAGTG
9241 AACCTCCAC CTCAGCCTCC CAAGTAGCTA AGACCACAGG
40 9281 CGTGCGACCC CATGCCCAGC TAATTTTAA ATTTTGTGTA
9321 GAGACTAGGC CTCACCATGT TGCCAGGCT TTTATTTCTT
9361 ATAAGTATAT TTAAATTTAT AATTATATCC ACATTTTAAA
9401 ATTTTAATTT AAAAAATTAC TCTGAGGCCG GGCATTGTGG
9441 CTCATGCCTC TAATCCCCAG CACCTTGGGA GGCCGAGTTG
45 9481 GGCAGATCAC CCGAGGTCAG AAGTTCGAGA CCAGCCTGAG

```

	9521	TAACATGGAG	AAACCCCCGT	CTCTACTAAA	AATACAAAAT
	9561	TGGCCGGGCG	TGGTGGTGCA	TTCCTGTAAT	CCCAGCTACT
	9601	CGGGAGGCTG	AGGCAGGAGA	ATTGCTTGAA	CCCAGGAGGT
	9641	GGAGGTTGCA	GTGAGCCAAG	ATTGTGCCAC	TGCACTGCAG
5	9681	CCTGGGCCAC	AGAGAGAATC	TGCCAAAAAA	AGAAAAAAA
	9721	AAAAATTTCA	GCCGTACAAG	GATGTTCATA	GCAACCCTGC
	9761	TGGAAATAGG	AAAAAAAATT	GGAAATAACC	TAAACTACTC
	9801	ACAATAGGAA	TCAGCTAAAA	CCCTGGGGGT	TTAATTCCAG
	9841	GGAATACTGT	GAACAATGAC	AAGTTTGTGG	ACTGAGTAAA
10	9881	AATAAACAGC	TGTCAATGAC	TTAACATTAA	ATGAAACAGC
	9921	AGAAGATGTC	ACAGCAGGTT	CTCGCTGAGC	CATTCAGAGG
	9961	GGTGTGGATC	ATTTAGAGGT	TCAAGTCCAC	TGGATTCTTC
	10001	TTTTTCCTTT	TAATATTACT	TCACTTCCAA	ATAAGGAAAG
	10041	GAAAGGAAAG	GAAATCACGT	CCAGTCCTGA	GACTTGCCAT
15	10081	CCTGCAGTCA	CCCCTCCTTT	TGTCTCCAGC	AGGTGGCAGA
	10121	CGCGTTCCAG	GGATGAATCC	CACTGCCTCT	GTTTAATGCA
	10161	GACGGTCCAG	CCGCTCCCAA	CAGCAGGTGG	GGCTATAAGC
	10201	ATCCATCCTA	CCTGCTCAAG	GAACCCAGGC	ATCAGAACTG
	10241	CTCTCTCCCA	AGTCCATTGC	AAGAAGGCAG	TCGTCTGGTC
20	10281	ATGAGAGGGT	TAACAGTCCA	CATTCCAGAG	CAAGGGAAAA
	10321	GGAGGCTGGA	GGGTCATAGA	CAAGGGGAGG	TGGTGCGGAG
	10361	GGCCAGCTTC	TCACAACACT	ACCGGCTCTG	CTGGGAGAGA
	10401	TAGATCACCC	CCAACAATGG	CCACAGCTGT	TTTCATCTGC
	10441	CCTGAAGGAA	ACTGACTTAG	GAAGCAGGTA	TCAGAGAGGG
25	10481	CCCTTCCTGA	GGGGGCTTCT	GTCTGGCTTG	TAAAAC TGTC
	10521	AGAGCAGCTG	CATTCATGTG	TCGGATGATG	GATGATGGAA
	10561	AGGACAGTCG	GCTGCAGATG	GACACAGCGA	CTTGCAAGTT
	10601	GAGGCAGGTG	GCAAAGGACT	TGCAGAGGCT	CTGCAGGTGG
	10641	GGCATGCTGA	TTCATTGCCC	AGTTAAAATA	CCAGAGGATC
30	10681	TGGGCAGCCT	CTTCACAGGA	GCTGCTTGTC	CTCAAACAAT
	10721	CTGTCTTCAA	TGAAAGATTC	CTCTGGCCTT	CCTTTCTCTT
	10761	CTTGCACCTC	AGGTGTGAAT	CCTTCTCCCC	CACGCCTCTA
	10801	CCTGCGCCCC	CGCCCCCGGC	CCCGGCCCTG	TGTGGCTCAT
	10841	TATATGCAGG	GCCAAGGCAG	CATTTTCTCT	TAGCTTCTTT
35	10881	GTGACCAGTT	GGTCCTGGGA	TGGCTTCATG	GAACACATCC
	10921	TGTGGTGTGC	ACCAATGAAG	CTTTCCATAC	AGGACTCAAA
	10961	ACTGTTTTTG	AAAAATGTAA	CCAGCTGGAA	GACAAGAAAA
	11001	TAAAATGTCA	GCACTAAAAA	CGCTGGCTGT	GGCTTTTGCT
	11041	AAGGAAAGGA	ATTTGGTGTT	GTCTTCTCAC	ACACACAGAC
40	11081	TGGTTGGGGA	AATGACTGTC	TTCAGCACAT	CACCCTGCGA
	11121	GCCACAGTGA	GTGCCCTGGC	TCAGAAGTGC	CTGTCACAGT
	11161	GCACAGGATC	CCTGAGGAGC	ATGAGCTGGG	ATTTCTCTCTG
	11201	TGCTGTCCAT	CACAGGAGCC	TGAGTGACCA	GCGCATCCTC
	11241	GATTTGTAAC	CAGAATCCTG	CCCTCTCTCC	CAAGCGGGCA
45	11281	CCCTTGCTCT	GACCCTCTAG	TTCTCTCTCT	TGCCTTCCAG
	11321	AGAATACCAA	GAGAGGCTTT	CTTGGTTAGG	ACAATGAATG
	11361	CTGAGACTTG	TGGAGTTGGG	ACCAATGGGA	TTTCTTTAAA

```

11401 AGCATCTTTT TGCCTCTGGC TGGGTCTATG GGGGTCAAAC
11441 AGAAACACCT TGGGCCATTT GTTGGTGGGG TGACAAATGA
11481 ACTTGGCCTG AGAAATGGAA TAGGCCGGGC TCAGCCCCGC
11521 GAAGCACTCA GAACTGCACA TTTTCTTTGT TGAGCGGGTC
5 11561 CACAGTTTGT TTTGAGAATG CCCGAGGGCC CAGGGAGACA
11601 GACAATTAAA AGCCGGAGCT CATTTTGATA TCTGAAAACC
11641 ACAGCCGCCA GCACGTGGGA GGTGCCGGAG AGCAGGCTTG
11681 GGCCTTGCC T CACACGCCCC CTCTCTCTGG GTCACCTGGG
11721 AGTGCCAGCA GCAATTTGGA AGTTTGCTGA GCTAGAGGAG
10 11761 AAGTCTTTGG GGAGGGTTTG CTCTGAGCAC ACCCCTTTCC
11801 CTCCCTCCGG GGCTGAGGGA AACATGGGAC CAGCCCTGCC
11841 CCAGCCTGTC CTCATTGGCT GGCATGAAGC AGAGAGGGGC
11881 TTAAAAAGG CGACCGTGTC TCGGCTGGAG ACCAGAGCCT
11921 GTGCTACTGG AAGGTGGCGT GCCCTCCTCT GGCTGGTACC
15 11960 ATGCAGCTCC CACTGGCCCT GTGTCTCGTC TGCCTGCTG

```

The invention can also be practiced with variants of the SOST gene and with variants of the sclerostin protein. For example, the invention can be practiced with the mouse or rat SOST gene or with the mouse or rat sclerostin protein. The mouse SOST cDNA has Genbank Accession No. NM024449, and is provided below as SEQ ID NO:5 for easy reference.

```

1 ATGCAGCCCT CACTAGCCCC GTGCCTCATC TGCCTACTTG
41 TGCACGCTGC CTTCTGTGCT GTGGAGGGCC AGGGGTGGCA
25 81 AGCCTTCAGG AATGATGCCA CAGAGGTCAT CCCAGGGCTT
121 GGAGAGTACC CCGAGCCTCC TCCTGAGAAC AACCAGACCA
161 TGAACCGGGC GGAGAATGGA GGCAGACCTC CCCACCATCC
201 CTATGACGCC AAAGATGTGT CCGAGTACAG CTGCCGCGAG
241 CTGCACTACA CCCGCTTCCT GACAGACGGC CCATGCCGCA
30 281 GCGCCAAGCC GGTCACCGAG TTGGTGTGCT CCGGCCAGTG
321 CGGCCCCGCG CGGCTGCTGC CCAACGCCAT CGGGCGCGTG
361 AAGTGGTGGC GCCCGAACGG ACCGGATTTC CGCTGCATCC
401 CGGATCGCTA CCGCGCGCAG CGGGTGCAGC TGCTGTGCCC
441 CGGGGGCGCG GCGCCGCGCT CGCGCAAGGT GCGTCTGGTG
35 481 GCCTCGTGCA AGTGCAAGCG CCTCACCCGC TTCCACAACC
521 AGTCGGAGCT CAAGGACTTC GGGCCGGAGA CCGCGCGGCC
561 GCAGAAGGGT CGCAAGCCGC GGCCCGGCGC CCGGGGAGCC
601 AAAGCCAACC AGGCGGAGCT GGAGAACGCC TACTAG

```

The mouse sclerostin protein has Genbank Accession No. NP077769, and is provided below as SEQ ID NO:6 for easy reference.

1 MQPSLAPCLI CLLVHAAFCA VEGQGWQAFR NDATEVIPGL
 41 GEYPEPPPEN NQTMNRAENG GRPPHHPYDA KDVSEYSCRE
 81 LHYTRFLTDG PCRSAPVTE LVCSGQCGPA RLLPNAIGRV
 121 KWWRPNGPDF RCIPDRYRAQ RVQLLCPGGA APRSRKVRLV
 5 161 ASCKCKRLTR FHNQSELKDF GPETARPQKG RKPRPGARGA
 201 KANQAELENA Y

The rat SOST cDNA has Genbank Accession No. NM030584, and is provided below as SEQ ID NO:7 for easy reference.

10

1 GAGGACCGAG TGCCCTTCCT CCTTCTGGCA CCATGCAGCT
 41 CTCACTAGCC CCTTGCCTTG CCTGCCTGCT TGTACATGCA
 81 GCCTTCGTTG CTGTGGAGAG CCAGGGGTGG CAAGCCTTCA
 121 AGAATGATGC CACAGAAATC ATCCCGGGAC TCAGAGAGTA
 15 161 CCCAGAGCCT CCTCAGGAAC TAGAGAACAA CCAGACCATG
 201 AACCGGGCCG AGAACGGAGG CAGACCCCCC CACCATCCTT
 241 ATGACACCAA AGACGTGTCC GAGTACAGCT GCCGCGAGCT
 281 GCACTACACC CGCTTCGTGA CCGACGGCCC GTGCCGCAGT
 321 GCCAAGCCGG TCACCGAGTT GGTGTGCTCG GGCCAGTGCG
 20 361 GCGGCGCGCG GCTGCTGCCC AACGCCATCG GGCGCGTGAA
 401 GTGGTGGCGC CCGAACGGAC CCGACTTCCG CTGCATCCCG
 441 GATCGCTACC GCGCGCAGCG GGTGCAGCTG CTGTGCCCCG
 481 GCGGCGCGGC GCCGCGCTCG CGCAAGGTGC GTCTGGTGGC
 521 CTCGTGCAAG TGCAAGCGCC TCACCCGCTT CCACAACCAG
 25 561 TCGGAGCTCA AGGACTTCGG ACCTGAGACC GCGCGGCCGC
 601 AGAAGGGTCG CAAGCCGCGG CCGCGCGCCC GGGGAGCCAA
 641 AGCCAACCAG GCGGAGCTGG AGAACGCCTA CTAG

The rat sclerostin protein has Genbank Accession No. NP085073, and is provided below as SEQ ID NO:8 for easy reference.

30

1 MQLSLAPCLA CLLVHAAFVA VESQGWQAFK NDATEIIPGL
 41 REYPEPPQEL ENNQTMNRAE NGGRPPHHPY DTKDVSEYSC
 81 RELHYTRFVT DGPCRSAPV TELVCSGQCG PARLLPNAIG
 35 121 RVKWWRPNGP DFRCIPDRYR AQRVQLLCPG GAAPRSRKVR
 161 LVASCKCKRL TRFHNQSELK DFGPETARPQ KGRKPRPRAR
 201 GAKANQAELE NAY

Other SOST nucleic acids and sclerostin proteins can be used for the practice of the invention, including the vervet SOST cDNA [accession number AF326742] and the bovine SOST cDNA [accession number AF326738].

5 According to the invention, the expression of the SOST gene increases with the maturation and differentiation of osteoblasts as well as with the onset of mineralization. Hence, SOST expression can be used as a marker for osteoblast differentiation.

 However, factors that increase SOST expression lead to decreased bone
10 mineralization because the SOST gene product, sclerostin, is an antagonist of bone morphogenetic proteins and causes apoptosis of osteoblasts. Sclerostin decreased the expression of many phenotypic markers of pre-osteoblasts / osteoblasts (ALP, type I collagen, PTHr) when it was added to cultures of differentiating hMSC cells and primary cultures of osteoblastic cells. Moreover,
15 sclerostin interacted with and antagonized the activity of BMP-6, thereby inhibiting osteoblast differentiation.

 According to the invention, sclerostin plays an integral role in the regulation of bone matrix formation and mineralization. Although several proteins are thought to function as antagonists of bone morphogenetic protein
20 action (for example, noggin and gremlin), it appears that when the sclerostin function is lost that function cannot be replaced by noggin or any of the other members of the DAN family of antagonists.

 A schematic diagram of how sclerostin to affects bone cell biology is depicted in Figure 26. Sclerostin exerts its effects on pre-osteoblasts,
25 osteoprogenitor, and mature osteoblasts cells. By depriving pre-osteoblasts and osteoprogenitor cells of factors such as bone morphogenetic proteins, these cells are prevented from completing their differentiation into mature osteoblasts. Thus, fewer osteoblasts are available to provide mineralization. In addition, these proteins play important anti-apoptotic roles, which are potentially impacted
30 by increasing exposure to sclerostin. As a result, alkaline phosphatase activity and synthesis of type I collagen are reduced in pre-osteoblasts and

osteoprogenitor cells, while mineralization is affected in mature osteoblasts treated with sclerostin. Reducing the expression of SOST or the amount of sclerostin available to developing osteoblasts permits more cells to progress through differentiation to produce more mature osteoblasts, more synthesis of type I collagen and greater deposition of mineral in bones.

Several factors enhance the expression of SOST, including insulin-like growth factor-1 and bone morphogenetic proteins such as BMP-2, BMP-4, and BMP-6. When bone morphogenetic proteins are present, retinoic acid and vitamin D also enhanced the levels of SOST expression. Such factors can therefore lead to diminished collagen synthesis and less mineralization within bones through increased synthesis of sclerostin.

In contrast, SOST expression levels decrease when dexamethasone is present with BMP-4 or BMP-6. Therefore, dexamethasone tends to abolish the stimulatory effect of BMP-4 and BMP-6 on SOST expression. Other chemical entities including glucocorticoid analogs, bile salts and prostaglandins also modulate the effects of bone morphogenetic proteins on SOST expression, as described below.


Modulating SOST/Sclerostin

According to the invention, any agents that modulate the ability of sclerostin to decrease osteoblastic activity are useful in increasing bone formation, modulating bone resorption and augmenting bone mineralization. Such agents can act directly or indirectly on SOST or sclerostin. Such agents can act at the transcriptional, translational or protein level to modulate the ability of sclerostin to decrease osteoblastic activity.

The term “modulate” or “modulating” means changing, that is increasing or decreasing. Hence, while agents that can decrease SOST expression or sclerostin activity are often used in the compositions and methods of the invention, agents that also increase SOST expression or sclerostin activity are also encompassed within the scope of the invention.

Agents useful for modulating SOST expression and sclerostin activity include, but are not limited to, steroids, alkaloids, terpenoids, peptoids, peptides, nucleic acids, anti-sense nucleic acids and synthetic chemicals. In some embodiments, the SOST antagonist or agonist can bind to a glucocorticoid receptor. For example, dexamethasone tends to abolish the stimulatory effect of BMP-4 and BMP-6 on SOST expression. Other chemical entities including glucocorticoid analogs, bile salts and prostaglandins also modulate the effects of bone morphogenetic proteins on SOST expression.

In one embodiment of the invention, steroids of the formula I can be used for modulating the expression of sclerostin.




wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl. Preferred R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , and R_7 groups are carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl. Preferably, R_1 is carbonyl or hydroxy and R_2 is lower alkyl. Also preferably, R_8 is a fluorine, hydrogen or halo group.

The following general definitions are used, unless otherwise described: halo is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc. denote both straight and branched groups; but reference to an individual radical such as "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" being specifically referred to. Aryl denotes a phenyl

radical or an ortho-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic. Aryloxy means aryl-O-.


More specifically, lower alkyl means (C₁-C₆) alkyl. Such (C₁-C₆) alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, or hexyl. Lower alkoxy generally means (C₁-C₆) alkoxy; such (C₁-C₆) alkoxy can, for example, be methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, pentoxy, 3-pentoxy, or hexyloxy. Preferred lower alkyl groups are (C₁-C₃) alkyl including methyl ethyl, propyl, isopropyl and the like. More preferred lower alkyl groups are methyl. Lower acyl refers to a carbonyl group attached to a lower alkyl group (e.g., -CO-CH₃). Lower hydroxy alkyl refers to a hydroxy group attached to a lower alkyl or lower alkylene group (e.g. -CH₂-CH₂-OH).

One example of an effective SOST/sclerostin antagonist is dexamethasone. Dexamethasone is a synthetic version of the glucocorticoid hormone, cortisol, which has been used as an anti-inflammatory agent. The structure of dexamethasone is provided below.




Cortisol and other glucocorticoids and molecules with structures similar to dexamethasone are also contemplated by the invention as antagonists of sclerostin. Androgens and estrogens are also contemplated as antagonists of sclerostin. Other chemical entities including glucocorticoid analogs, bile salts and prostaglandins also modulate the effects of bone morphogenetic proteins on SOST expression.

Exemplary bile salts can have the following structure:



wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ groups are as described above, and R₉ is carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl. Preferably, R₉ is hydrogen, hydroxyl, lower acyl, lower alkyl, or lower hydroxy alkyl.

Exemplary prostaglandins or eicosanoids can have the following structure:



wherein R₁₀ and R₁₁ are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl, aryl or aryloxy.

Compounds or molecules that stimulate the degradation of SOST transcripts or sclerostin can also be used to diminish the levels of sclerostin and enhance bone mineralization.

In another embodiment, the invention provides anti-sense RNA or DNA molecules to modulate SOST expression, sclerostin translation and/or the degradation of SOST transcripts. For example, an anti-sense RNA or DNA that

can hybridize to a nucleic acid having any one of SEQ ID NO:1, 2, 4, 5 or 7 can be used as an anti-sense RNA or DNA for diminishing the expression of sclerostin.

The degradation of SOST mRNA may also be increased upon exposure to small duplexes of synthetic double-stranded RNA through the use of RNA interference (siRNA or RNAi) technology (Scherr, M. et al. 2003; Martinez, L.A. et al. 2002). A process is therefore provided for inhibiting expression of a target gene in a cell. The process comprises introduction of RNA with partial or fully double-stranded character into the cell or into the extracellular environment. Inhibition is specific to SOST RNA because a nucleotide sequence from a portion of the SOST gene is chosen to produce inhibitory RNA. This process is effective in producing inhibition of gene expression.

SiRNAs were designed using the guidelines provided by Ambion (Austin, TX). Briefly, the SOST cDNA sequence was scanned for target sequences that had AA dinucleotides. Sense and anti-sense oligonucleotides were generated to these targets (AA + 3' adjacent 19 nucleotides) that contained a G/C content of 35 to 55%. These sequences were then compared to others in the human genome database to minimize homology to other known coding sequences (Blast search).

The target and siRNA sequences designed are provided below.

Target sequence 1: AAGAATGATGCCACGGAAATC (SEQ ID NO:9)
 Position in gene sequence: 140
 GC content: 42.9%
 Sense strand siRNA: GAAUGAUGCCACGGAAAUCtt (SEQ ID NO:10)
 Antisense strand siRNA: GAUUUCCGUGGCAUCAUUCtt (SEQ ID NO:11)

Target sequence 3: AATGATGCCACGGAAATCATC (SEQ ID NO:12)
 Position in gene sequence: 143
 GC content: 42.9%
 Sense strand siRNA: UGAUGCCACGGAAAUCAUCtt (SEQ ID NO:13)
 Antisense strand siRNA: GAUGAUUCCGUGGCAUCAtt (SEQ ID NO:14)

Target sequence 5: AACACAAGACCATGAACCGG (SEQ ID NO:15)
 Position in gene sequence: 209
 GC content: 47.6%

- Sense strand siRNA: CAACAAGACCAUGAACCGGtt (SEQ ID NO:16)
 Antisense strand siRNA: CCGGUUCAUGGUCUUGUUGtt (SEQ ID NO:17)
- 5 Target sequence 27: AATTGAGAGTCACAGACACTG (SEQ ID NO:18)
 Position in gene sequence: 950
 GC content: 42.9%
 Sense strand siRNA: UUGAGAGUCACAGACACUGtt (SEQ ID NO:19)
 Antisense strand siRNA: CAGUGUCUGUGACUCUCAAtt (SEQ ID NO:20)
- 10 Target sequence 28: AAATGGAAGCATTTTCACCGC (SEQ ID NO:21)
 Position in gene sequence: 1035
 GC content: 42.9%
 Sense strand siRNA: AUGGAAGCAUUUUCACCGCtt (SEQ ID NO:22)
 Antisense strand siRNA: GCGGUGAAA AUGCUUCCAUtt (SEQ ID NO:23)
- 15 Target sequence 30: AAAGTCCAGGGACTGGTTAAG (SEQ ID NO:24)
 Position in gene sequence: 1093
 GC content: 47.6%
 Sense strand siRNA: AGUCCAGGGACUGGUUAAGtt (SEQ ID NO:25)
 20 Antisense strand siRNA: CUUAACCAGUCCCUGGACUtt (SEQ ID NO:26)
- Target sequence 31: AAGAAAGTTGGATAAGATTCC (SEQ ID NO:27)
 Position in gene sequence: 1111
 GC content: 33.3%
 25 Sense strand siRNA: GAAAGUUGGAUAAGAUUCCtt (SEQ ID NO:28)
 Antisense strand siRNA: GGAAUCUUAUCCAACUUUCtt (SEQ ID NO:29)
- Target sequence 36: AACTGTAGATGTGGTTTCTAG (SEQ ID NO:30)
 Position in gene sequence: 1201
 30 GC content: 38.1%
 Sense strand siRNA: CUGUAGAUGUGGUUUCUAGtt (SEQ ID NO:31)
 Antisense strand siRNA: CUAGAAACCACAUCUACAGtt (SEQ ID NO:32)
- Target sequence 40: AATTCTCCTTCGGGACCTCAA (SEQ ID NO:33)
 35 Position in gene sequence: 1269
 GC content: 47.6%
 Sense strand siRNA: UUCUCCUUCGGGACCUCAAAtt (SEQ ID NO:34)
 Antisense strand siRNA: UUGAGGUCCCGAAGGAGAAAtt (SEQ ID NO:35)
- 40 Target sequence 47: AAAGAGAGAGAATGAATGCAG (SEQ ID NO:36)
 Position in gene sequence: 1414
 GC content: 38.1%
 Sense strand siRNA: AGAGAGAGAAUGAAUGCAGtt (SEQ ID NO:37)
 Antisense strand siRNA: CUGCAUUCAUUCUCUCUCUtt (SEQ ID NO:38)
- 45

Target sequence 63: AAGAAGCTATGCTGCTTCCCA (SEQ ID NO:39)

Position in gene sequence: 1590

GC content: 47.6%

Sense strand siRNA: GAAGCUAUGCUGCUUCCCA_{tt} (SEQ ID NO:40)

5 Antisense strand siRNA: UGGGAAGCAGCAUAGCUU (SEQ ID NO:41)

Target sequence 70: AAATCACATCCGCCCAACTT (SEQ ID NO:42)

Position in gene sequence: 1726

GC content: 47.6%

10 Sense strand siRNA: AUCACAUCCGCCCAACUU_{tt} (SEQ ID NO:43)

Antisense strand siRNA: AAGUUGGGGCGGAUGUGAU_{tt} (SEQ ID NO:44)

Hence, the invention provides a pharmaceutical composition that includes a carrier and a siRNA comprising SEQ ID NO:11, SEQ ID NO:14, 15 SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41 or SEQ ID NO:44, wherein the siRNA can modulate SOST expression. The invention also provides a pharmaceutical composition comprising a carrier and a siRNA that is selectively hybridizable under stringent conditions to an RNA 20 derived from a DNA comprising SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:39, or SEQ ID NO:42, wherein the siRNA can modulate SOST expression.

Mixtures and combinations of such siRNA molecules are also 25 contemplated by the invention. These compositions can be used in the methods of the invention, for example, for treating or preventing apoptosis of bone-related cells or for treating or preventing loss of bone density. These compositions are also useful for modulating (e.g. decreasing) SOST expression.

The siRNA provided herein can selectively hybridize to RNA *in vivo* or 30 *in vitro*. A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under physiological conditions or under moderate stringency hybridization and wash conditions. In some embodiments the siRNA is selectively hybridizable to an RNA (e.g. a SOST RNA) under physiological 35 conditions. Hybridization under physiological conditions can be measured as a

practical matter by observing interference with the function of the RNA.

Alternatively, hybridization under physiological conditions can be detected *in vitro* by testing for siRNA hybridization using the temperature (e.g. 37 °C) and salt conditions that exist *in vivo*.

- 5 Moreover, as an initial matter, other *in vitro* hybridization conditions can be utilized to characterize siRNA interactions. Exemplary *in vitro* conditions include hybridization conducted as described in the Bio-Rad Labs ZetaProbe manual (Bio-Rad Labs, Hercules, Calif.); Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, (1989), or
- 10 Sambrook et al., Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Laboratory Press, (2001)), expressly incorporated by reference herein. For example, hybridization can be conducted in 1 mM EDTA, 0.25 M Na₂ HPO₄ and 7% SDS at 42 °C, followed by washing at 42 °C in 1 mM EDTA, 40 mM NaPO₄, 5% SDS, and 1 mM EDTA, 40 mM NaPO₄, 1% SDS.
- 15 Hybridization can also be conducted in 1 mM EDTA, 0.25 M Na₂ HPO₄ and 7% SDS at 60 °C, followed by washing in 1 mM EDTA, 40 mM NaPO₄, 5% SDS, and 1 mM EDTA, 40 mM NaPO₄, 1% SDS. Washing can also be conducted at other temperatures, including temperatures ranging from 37 °C to at 65 °C, from 42 °C to at 65 °C, from 37 °C to at 60 °C, from 50 °C to at 65 °C, from 37 °C to at
- 20 55 °C, and other such temperatures.

 The siRNA employed in the compositions and methods of the invention may be synthesized either *in vivo* or *in vitro*. In some embodiments, the siRNA molecules are synthesized *in vitro* using methods, reagents and synthesizer equipment available to one of skill in the art. Endogenous RNA polymerases

25 within a cell may mediate transcription *in vivo*, or cloned RNA polymerase can be used for transcription *in vivo* or *in vitro*. For transcription from a transgene or an expression construct *in vivo*, a regulatory region may be used to transcribe the siRNA strands.

 Depending on the particular sequence utilized and the dose of double

30 stranded siRNA material delivered, the compositions and methods may provide partial or complete loss of function for the target gene (SOST). A reduction or

loss of gene expression in at least 99% of targeted cells has been shown for other genes. See, e.g., U.S. Patent 6,506,559. Lower doses of injected material and longer times after administration of the selected siRNA may result in inhibition in a smaller fraction of cells.

5 The siRNA may comprise one or more strands of polymerized ribonucleotide; it may include modifications to either the phosphate-sugar backbone or the nucleoside. The double-stranded siRNA structure may be formed by a single self-complementary RNA strand or two complementary RNA strands. siRNA duplex formation may be initiated either inside or outside the
10 cell. The siRNA may be introduced in an amount that allows delivery of at least one copy per cell. Higher doses of double-stranded material may yield more effective inhibition.

 Inhibition is sequence-specific in that nucleotide sequences corresponding to the duplex region of the RNA are targeted for genetic
15 inhibition. siRNA containing nucleotide sequences identical to a portion of the target gene is preferred for inhibition. However, siRNA sequences with insertions, deletions, and single point mutations relative to the target sequence have also been found to be effective for inhibition. Thus, sequence identity may optimized by alignment algorithms known in the art and calculating the percent
20 difference between the nucleotide sequences. Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a portion of the target gene transcript.

 The siRNA may be directly introduced into the cell (i.e., intracellularly); or introduced extracellularly into a cavity, interstitial space, into the circulation
25 of an organism, introduced orally, or may be introduced by bathing an organism in a solution containing siRNA. Methods for oral introduction include direct mixing of siRNA with food of the organism, as well as engineered approaches in which a species that is used as food is engineered to express an siRNA, then fed to the organism to be affected. Physical methods of introducing nucleic acids
30 include injection directly into the cell or extracellular injection into the organism of an siRNA solution.

The siRNA may also be delivered *in vitro* to cultured cells using transfection agents available in the art such as lipofectamine or by employing viral delivery vectors such as those from lentiviruses. Such *in vitro* delivery can be performed for testing purposes or for therapeutic purposes. For example,
5 cells from a patient can be treated *in vitro* and then re-administered to the patient.

The advantages of using siRNA include: the ease of introducing double-stranded siRNA into cells, the low concentration of siRNA that can be used, the stability of double-stranded siRNA, and the effectiveness of the inhibition. The
10 ability to use a low concentration of a naturally-occurring nucleic acid avoids several disadvantages of anti-sense interference.

An antagonist or agonist for sclerostin may also be a co-factor, such as a protein, peptide, carbohydrate, lipid or small molecular weight molecule, which interacts with sclerostin to regulate its activity. Other molecules contemplated as
15 agents for modulating sclerostin include antibodies targeted against sclerostin as well as molecules, compounds or peptides that mimic bone morphogenetic proteins in structure and that bind to and form inactive complexes with sclerostin. Potential polypeptide antagonists include antibodies that react with sclerostin.

20 As described herein, sclerostin is involved in controlling formation of mature osteoblasts, the primary cell type implicated in bone mineralization. A decrease in the rate of bone mineralization can lead to various bone disorders collectively referred to as osteopenias, including bone disorders such as osteoporosis, osteomyelitis, hypercalcemia, osteopenia brought on by surgery or
25 steroid administration, Paget's disease, osteonecrosis, bone loss due to rheumatoid arthritis; periodontal bone loss, immobilization, prosthetic loosening and osteolytic metastasis.

According to the invention, agents that modulate the function of sclerostin, or that modulate the expression of SOST, can be used to treat any of
30 these diseases. Preferably, SOST/sclerostin antagonists or agents that decrease

the function of SOST or sclerostin are used to treat osteoporosis, frail bone problems and related diseases where bone density is inadequate.

Conversely, an increase in bone mineralization can lead to osteopetrosis, a condition marked by excessive bone density. According to the invention, sclerostin may be used to treat conditions characterized by excessive bone density. The most common condition is osteopetrosis in which a genetic defect results in elevated bone mass and is usually fatal in the first few years of life. Osteopetrosis is preferably treated by administration of sclerostin.

The therapeutic methods for treating or preventing a bone disorder such as an inadequate rate of bone mineralization involve administering an agent that can modulate SOST (e.g. a SOST antagonist) to a mammal affected with the bone disorder. Any of the agents or antagonists described herein, or isolated by the methods provided herein, can be administered. Any bone disorder involving diminished mineralization can be treated or prevented using these methods.

The therapeutic methods for treating or preventing a bone disorder such as an excessive bone density involve administering a therapeutically effective amount of sclerostin to a mammal affected with the bone disorder. Any of the available mammalian sclerostin polypeptides (for example, those described herein) can be administered for treating or preventing excessive bone density.

20

Methods of Isolating Agents that can Modulate SOST/Sclerostin

The invention further provides screening assays that are useful for generating or identifying therapeutic agents for the treatment of bone disorders. In particular, the SOST nucleic acids and sclerostin proteins identified herein may be used in a variety of assays for detecting SOST and/or sclerostin and for identifying factors that interact with SOST nucleic acids or with the sclerostin protein.

In general, an assay for identifying sclerostin involves incubating a test sample under conditions which permit binding of sclerostin to a reporter molecule, and measuring the extent of binding. A reporter molecule can be any molecule that stably binds to sclerostin and that can be detected. For example,

30

the reporter molecule can be an anti-sclerostin antibody that is labeled with radioactive isotopes (^{125}I , ^{32}P , ^{35}S , ^3H), fluorescent dyes (fluorescein, rhodamine), enzymes and the like. It is understood that the choice of a reporter molecule will depend upon the detection system used.

5 Sclerostin may be purified or present in mixtures, such as in cultured cells, tissue samples, body fluids or culture medium. Assays may be developed that are qualitative or quantitative, with the latter being useful for determining the binding parameters (affinity constants and kinetics) of the reporter molecule to sclerostin and for quantifying levels of sclerostin in mixtures. Assays may
10 also be used to detect fragments, analogs and derivatives of sclerostin and to identify new sclerostin family members.

Binding assays may be carried out in several formats, including cell-based binding assays, solution-phase assays and immunoassays. In general, test samples or compounds are incubated with sclerostin test samples for a specified
15 period of time followed by measurement of the reporter molecule by use of microscopy, fluorimetry, a scintillation counter, or any available immunoassay. Binding can also be detected by labeling sclerostin in a competitive radioimmunoassay. Alternatively, sclerostin may be modified with an unlabeled epitope tag (e.g., biotin, peptides, His₆, FLAG, myc etc.) and bound to proteins
20 such as streptavidin, anti-peptide or anti-protein antibodies that have a detectable label as described above. Additional forms of sclerostin containing epitope tags may be used in solution and immunoassays.

Methods for identifying compounds or molecules that interact with sclerostin are also encompassed by the invention. In general, an assay for
25 identifying compounds or molecules that interact with sclerostin involves incubating sclerostin with a test sample that may contain such a compound or molecule under conditions that permit binding of the compound or molecule to sclerostin, and measuring whether binding has occurred. Sclerostin may be purified or present in mixtures, such as in cultured cells, tissue samples, body
30 fluids or culture medium. Assays may be developed that are qualitative or quantitative. Quantitative assays can be used for determining the binding

parameters (affinity constants and kinetics) of the compound or molecule for sclerostin and for quantifying levels of biologically active compounds and molecules in mixtures. Assays may also be used to evaluate the binding of a compound or molecule to fragments, analogs and derivatives of sclerostin and to
5 identify new sclerostin family members.

The compound or molecule in a test sample may be substantially purified or present in a crude mixture. Binding compounds and molecules may be nucleic acids, proteins, peptides, carbohydrates, lipids or small molecular weight organic compounds. The compounds and molecules may be further characterized by
10 their ability to increase or decrease sclerostin activity in order to determine whether they act as an agonist or an antagonist.

In another embodiment, the invention involves a method for identifying an agent that can modulate sclerostin-related apoptosis activity comprising contacting a cell with both sclerostin and a test agent and detecting whether the
15 test sample prevents apoptosis of the cell. The amount of apoptosis in cells treated with sclerostin and a test agent can be compared to the amount of apoptosis in cells treated with sclerostin only. To detect the level of apoptosis, cell viability and apoptosis assays can be performed. Many cell viability assays are available to one of skill in the art and any of these can be used. For example,
20 a CellTiterGlo Luminescence Viability Assay available from Promega can be used to assess cell viability. Apoptosis can be detected using any of the many available apoptosis assays. For example, the Homogeneous Caspase ELISA (Roche) or the Cell Death assay (Histone-Associated DNA Fragmentation, Roche) can be used. Apoptosis can also be detected by TUNEL staining using
25 Roche's In Situ Cell Death Detection kit.

SOST nucleic acids are also useful for identification of factors that interact with the SOST promoter and that modulate SOST expression. Such factors may be intracellular proteins such as DNA binding proteins that interact with regulatory sequences that control SOST transcription, for example, the
30 SOST promoter. As an example, hybrid constructs may be used that include a nucleic acid encoding the SOST promoter fused to a nucleic acid encoding a

marker protein. The SOST promoter can be found within SEQ ID NO:1. The promoter used can, for example, have sequence SEQ ID NO:4.

The marker protein can be any marker protein available to one of skill in the art. For example, the marker protein can be luciferase, green fluorescence
5 protein (GFP) or CAT.

Such constructs are used for in vitro or in vivo transcription assays to identify factors that modulate SOST expression. Factors that depress or diminish SOST expression are particularly useful. Expression or transcription levels can be assessed using any method available to one of skill in the art for
10 measuring RNA levels. For example, RNA levels can be assessed by northern analysis, reverse transcriptase analysis, reverse transcriptase coupled with polymerase chain reaction (RT-PCR) analysis and other methods. Chemical libraries can be screened using such methods for small molecule compounds that block SOST transcription.

15 Sclerostin is involved in controlling formation of mature osteoblasts, the primary cell type implicated in bone mineralization. Factors that counteract the sclerostin-induced depression of mineralization can be identified by observing whether a test agent increases mineralization in cultured osteoblasts that have been exposed to sclerostin. Hence, an assay may involve contacting an
20 osteoblast with sclerostin and a test agent and observing whether the mineralization of the cell is increased relative to a similar cell that is contacted only with sclerostin.

Many variations and modifications of the methods described herein will be apparent to one of skilled in the art. All such variations and modifications are
25 encompassed by the invention.

Antibodies

The invention provides antibody preparations directed against sclerostin, for example, antibodies capable of binding a polypeptide having SEQ ID NO:3,
30 SEQ ID NO:6 or SEQ ID NO:8.

Antibody molecules belong to a family of plasma proteins called immunoglobulins, whose basic building block, the immunoglobulin fold or domain, is used in various forms in many molecules of the immune system and other biological recognition systems. A typical immunoglobulin has four
5 polypeptide chains, containing an antigen binding region known as a variable region and a non-varying region known as the constant region.

Native antibodies and immunoglobulins are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy
10 chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (VH) followed by a number of constant domains. Each light chain has a variable domain at one end (VL) and
15 a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains (Clothia et al., J. Mol. Biol. 186, 651-66,
20 1985); Novotny and Haber, Proc. Natl. Acad. Sci. USA 82, 4592-4596 (1985).

Depending on the amino acid sequences of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are at least five (5) major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG-
25 1, IgG-2, IgG-3 and IgG-4; IgA-1 and IgA-2. The heavy chains constant domains that correspond to the different classes of immunoglobulins are called alpha (α), delta (δ), epsilon (ϵ), gamma (γ) and mu (μ), respectively. The light chains of antibodies can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino sequences of their constant
30 domain. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "variable" in the context of variable domain of antibodies, refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies. The variable domains are for binding and determine the specificity of each particular antibody for its particular antigen.

5 However, the variability is not evenly distributed through the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs) also known as hypervariable regions both in the light chain and the heavy chain variable domains.

The more highly conserved portions of variable domains are called the
10 framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close
15 to the formation of the antigen-binding site of antibodies. The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

An antibody that is contemplated for use in the present invention thus can
20 be in any of a variety of forms, including a whole immunoglobulin, an antibody fragment such as Fv, Fab, and similar fragments, a single chain antibody which includes the variable domain complementarity determining regions (CDR), and the like forms, all of which fall under the broad term "antibody", as used herein. The present invention contemplates the use of any specificity of an antibody,
25 polyclonal or monoclonal, and is not limited to antibodies that recognize and immunoreact with a specific antigen. In preferred embodiments, in the context of both the therapeutic and screening methods described below, an antibody or fragment thereof is used that is immunospecific for an antigen or epitope of the invention.

30 The term "antibody fragment" refers to a portion of a full-length antibody, generally the antigen binding or variable region. Examples of antibody

fragments include Fab, Fab', F(ab')₂ and Fv fragments. Papain digestion of antibodies produces two identical antigen binding fragments, called the Fab fragment, each with a single antigen binding site, and a residual "Fc" fragment, so-called for its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen binding fragments that are capable of cross-linking antigen, and a residual other fragment (which is termed pFc'). Additional fragments can include diabodies, linear antibodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments. As used herein, "functional fragment" with respect to antibodies, refers to Fv, F(ab) and F(ab')₂ fragments.

Antibody fragments contemplated by the invention are therefore not full-length antibodies but do have similar or improved immunological properties relative to an anti-sclerostin antibody. Such antibody fragments may be as small as about 4 amino acids, 5 amino acids, 6 amino acids, 7 amino acids, 9 amino acids, about 12 amino acids, about 15 amino acids, about 17 amino acids, about 18 amino acids, about 20 amino acids, about 25 amino acids, about 30 amino acids or more. In general, an antibody fragment of the invention can have any upper size limit so long as it binds with specificity to sclerostin, e.g. a polypeptide having SEQ ID NO:3, 6 or 8.

Antibody fragments retain some ability to selectively bind with its antigen. Some types of antibody fragments are defined as follows:

(1) Fab is the fragment that contains a monovalent antigen-binding fragment of an antibody molecule. A Fab fragment can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain.

(2) Fab' is the fragment of an antibody molecule can be obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain. Two Fab' fragments are obtained per antibody molecule. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxyl terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region.

(3) (Fab')₂ is the fragment of an antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent reduction. F(ab')₂ is a dimer of two Fab' fragments held together by two disulfide bonds.

(4) Fv is the minimum antibody fragment that contains a complete
5 antigen recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in a tight, non-covalent association (V_H-V_L dimer). It is in this configuration that the three CDRs of each variable domain interact to define an antigen binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen binding specificity to the
10 antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

(5) Single chain antibody ("SCA"), defined as a genetically engineered molecule containing the variable region of the light chain, the
15 variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule. Such single chain antibodies are also referred to as "single-chain Fv" or "sFv" antibody fragments. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the sFv to form the desired structure for antigen binding.
20 For a review of sFv see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds. Springer-Verlag, N.Y., pp. 269-315 (1994).

The term "diabodies" refers to a small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain
25 (V_H) connected to a light chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161,
30 and Hollinger et al., Proc. Natl. Acad. Sci. USA 90: 6444-6448 (1993).

Methods for preparing polyclonal antibodies are available to those skilled in the art. See, for example, Green, et al., Production of Polyclonal Antisera, in: Immunochemical Protocols (Manson, ed.), pages 1-5 (Humana Press); Coligan, et al., Production of Polyclonal Antisera in Rabbits, Rats Mice and Hamsters, in: 5 Current Protocols in Immunology, section 2.4.1 (1992), which are hereby incorporated by reference.

Methods for preparing monoclonal antibodies are likewise available to one of skill in the art. See, for example, Kohler & Milstein, Nature, 256:495 (1975); Coligan, et al., sections 2.5.1-2.6.7; and Harlow, et al., in: Antibodies: 10 A Laboratory Manual, page 726 (Cold Spring Harbor Pub. (1988)), which are hereby incorporated by reference. Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography. 15 See, e.g., Coligan, et al., sections 2.7.1-2.7.12 and sections 2.9.1-2.9.3; Barnes, et al., Purification of Immunoglobulin G (IgG), in: Methods in Molecular Biology, Vol. 10, pages 79-104 (Humana Press (1992)).

Methods of *in vitro* and *in vivo* manipulation of monoclonal antibodies are also available to those skilled in the art. For example, monoclonal antibodies 20 to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler and Milstein, Nature 256, 495 (1975), or may be made by recombinant methods, e.g., as described in U.S. Pat. No. 4,816,567. The monoclonal antibodies for use with the present invention may also be isolated from phage antibody libraries using the techniques 25 described in Clackson et al. Nature 352: 624-628 (1991), as well as in Marks et al., J. Mol Biol. 222: 581-597 (1991). Another method involves humanizing a monoclonal antibody by recombinant means to generate antibodies containing human specific and recognizable sequences. See, for review, Holmes, et al., J. Immunol., 158:2192-2201 (1997) and Vaswani, et al., Annals Allergy, Asthma 30 & Immunol., 81:105-115 (1998).

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates that the antibody preparation is a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567); Morrison et al. Proc. Natl. Acad. Sci. 81, 6851-6855 (1984).

Methods of making antibody fragments are also known in the art (see for example, Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York, (1988), incorporated herein by reference). Antibody fragments of the present invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* of DNA encoding the fragment. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S

fragment denoted $F(ab')_2$. This fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce 3.5S Fab' monovalent fragments. Alternatively, an enzymatic cleavage using pepsin produces two
5 monovalent Fab' fragments and an Fc fragment directly. These methods are described, for example, in U.S. Patents No. 4,036,945 and No. 4,331,647, and references contained therein. These patents are hereby incorporated in their entireties by reference.

Other methods of cleaving antibodies, such as separation of heavy chains
10 to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody. For example, Fv fragments comprise an association of V_H and V_L chains. This association may be non-covalent or the variable chains can be linked by an
15 intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. Preferably, the Fv fragments comprise V_H and V_L chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V_H and V_L domains connected by an oligonucleotide. The
20 structural gene is inserted into an expression vector, which is subsequently introduced into a host cell such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing sFvs are described, for example, by Whitlow, et al., Methods: a Companion to Methods in Enzymology, Vol. 2, page 97 (1991);
25 Bird, et al., *Science* 242:423-426 (1988); Ladner, et al, US Patent No. 4,946,778; and Pack, et al., Bio/Technology 11:1271-77 (1993).

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") are often involved in antigen recognition and binding. CDR
30 peptides can be obtained by cloning or constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the

polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells. See, for example, Larrick, et al., Methods: a Companion to Methods in Enzymology, Vol. 2, page 106 (1991).

The invention contemplates human and humanized forms of non-human
5 (e.g. murine) antibodies. Such humanized antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which
10 residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a nonhuman species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity.

In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized
15 antibodies may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, humanized antibodies will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR
20 regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see: Jones et al., Nature 321, 522-
25 525 (1986); Reichmann et al., Nature 332, 323-329 (1988); Presta, Curr. Op. Struct. Biol. 2, 593-596 (1992); Holmes, et al., J. Immunol., 158:2192-2201 (1997) and Vaswani, et al., Annals Allergy, Asthma & Immunol., 81:105-115 (1998).

The invention also provides methods of mutating antibodies to optimize
30 their affinity, selectivity, binding strength or other desirable property. A mutant antibody refers to an amino acid sequence variant of an antibody. In general,

one or more of the amino acid residues in the mutant antibody is different from what is present in the reference antibody. Such mutant antibodies necessarily have less than 100% sequence identity or similarity with the reference amino acid sequence. In general, mutant antibodies have at least 75% amino acid sequence identity or similarity with the amino acid sequence of either the heavy or light chain variable domain of the reference antibody. Preferably, mutant antibodies have at least 80%, more preferably at least 85%, even more preferably at least 90%, and most preferably at least 95% amino acid sequence identity or similarity with the amino acid sequence of either the heavy or light chain variable domain of the reference antibody. One method of mutating antibodies involves affinity maturation using phage display.

The invention is therefore directed to a method for selecting antibodies and/or antibody fragments or antibody polypeptides with desirable properties. Such desirable properties can include increased binding affinity or selectivity for the epitopes of the invention

The antibodies and antibody fragments of the invention are isolated antibodies and antibody fragments. An isolated antibody is one that has been identified and separated and/or recovered from a component of the environment in which it was produced. Contaminant components of its production environment are materials that would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. The term "isolated antibody" also includes antibodies within recombinant cells because at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step

If desired, the antibodies of the invention can be purified by any available procedure. For example, the antibodies can be affinity purified by binding an antibody preparation to a solid support to which the antigen used to raise the antibodies is bound. After washing off contaminants, the antibody can be eluted by known procedures. Those of skill in the art will know of various techniques common in the immunology arts for purification and/or concentration of

polyclonal antibodies, as well as monoclonal antibodies (see for example, Coligan, et al., Unit 9, Current Protocols in Immunology, Wiley Interscience, 1991, incorporated by reference).

In preferred embodiments, the antibody will be purified as measurable by
5 at least three different methods: 1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight; 2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequentator; or 3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using
10 Coomassie blue or, preferably, silver stain.

Compositions

The sclerostin antagonists and other agents (e.g. sclerostin proteins and SOST nucleic acids) of the invention, including their salts, are administered so
15 as to achieve a reduction in at least one symptom associated with a bone disorder, indication or disease.

To achieve the desired effect(s), sclerostin antagonists and other agents, may be administered as single or divided dosages, for example, of at least about 0.01 mg/kg to about 500 to 750 mg/kg, of at least about 0.01 mg/kg to about 300
20 to 500 mg/kg, at least about 0.1 mg/kg to about 100 to 300 mg/kg or at least about 1 mg/kg to about 50 to 100 mg/kg of body weight, although other dosages may provide beneficial results. The amount administered will vary depending on various factors including, but not limited to, the sclerostin antagonist or other agent chosen, the disease, the weight, the physical condition, the health, the age
25 of the mammal, whether prevention or treatment is to be achieved, and if the sclerostin antagonist or agent is chemically modified. Such factors can be readily determined by the clinician employing animal models or other test systems that are available in the art.

Administration of the therapeutic agents in accordance with the present
30 invention may be in a single dose, in multiple doses, in a continuous or intermittent manner, depending, for example, upon the recipient's physiological

condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of the therapeutic agents of the invention may be essentially continuous over a preselected period of time or may be in a series of spaced
5 doses. Both local and systemic administration is contemplated.

To prepare the composition, therapeutic agents are synthesized or otherwise obtained, purified as necessary or desired and then lyophilized and stabilized. The therapeutic agent can then be adjusted to the appropriate concentration, and optionally combined with other agents. The absolute weight
10 of a given therapeutic agent included in a unit dose can vary widely. For example, about 0.01 to about 2 g, or about 0.1 to about 500 mg, of at least one therapeutic agents of the invention, or a plurality of therapeutic agents can be administered. Alternatively, the unit dosage can vary from about 0.01 g to about 50 g, from about 0.01 g to about 35 g, from about 0.1 g to about 25 g, from about
15 0.5 g to about 12 g, from about 0.5 g to about 8 g, from about 0.5 g to about 4 g, or from about 0.5 g to about 2 g.

Daily doses of the therapeutic agents of the invention can vary as well. Such daily doses can range, for example, from about 0.1 g/day to about 50 g/day, from about 0.1 g/day to about 25 g/day, from about 0.1 g/day to about 12 g/day,
20 from about 0.5 g/day to about 8 g/day, from about 0.5 g/day to about 4 g/day, and from about 0.5 g/day to about 2 g/day.

Thus, one or more suitable unit dosage forms comprising the therapeutic agents of the invention can be administered by a variety of routes including oral, parenteral (including subcutaneous, intravenous, intramuscular and
25 intraperitoneal), rectal, dermal, transdermal, intrathoracic, intrapulmonary and intranasal (respiratory) routes. The therapeutic agents may also be formulated for sustained release (for example, using microencapsulation, see WO 94/ 07529, and U.S. Patent No.4,962,091). The formulations may, where appropriate, be conveniently presented in discrete unit dosage forms and may be prepared by
30 any of the methods well known to the pharmaceutical arts. Such methods may include the step of mixing the therapeutic agent with liquid carriers, solid

matrices, semi-solid carriers, finely divided solid carriers or combinations thereof, and then, if necessary, introducing or shaping the product into the desired delivery system.

When the therapeutic agents of the invention are prepared for oral
5 administration, they are generally combined with a pharmaceutically acceptable carrier, diluent or excipient to form a pharmaceutical formulation, or unit dosage form. For oral administration, the therapeutic agents may be present as a powder, a granular formulation, a solution, a suspension, an emulsion or in a natural or synthetic polymer or resin for ingestion of the active ingredients from
10 a chewing gum. The therapeutic agents may also be presented as a bolus, electuary or paste. Orally administered therapeutic agents of the invention can also be formulated for sustained release, e.g., the therapeutic agents can be coated, micro-encapsulated, or otherwise placed within a sustained delivery device. The total active ingredients in such formulations comprise from 0.1 to
15 99.9% by weight of the formulation.

By "pharmaceutically acceptable" it is meant a carrier, diluent, excipient, and/or salt that is compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof.

Pharmaceutical formulations containing the therapeutic agents of the
20 invention can be prepared by procedures known in the art using well-known and readily available ingredients. For example, the therapeutic agent can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, solutions, suspensions, powders, aerosols and the like. Examples of excipients, diluents, and carriers that are suitable for such
25 formulations include buffers, as well as fillers and extenders such as starch, cellulose, sugars, mannitol, and silicic derivatives. Binding agents can also be included such as carboxymethyl cellulose, hydroxymethylcellulose, hydroxypropyl methylcellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl-pyrrolidone. Moisturizing agents can be included such as
30 glycerol, disintegrating agents such as calcium carbonate and sodium bicarbonate. Agents for retarding dissolution can also be included such as

paraffin. Resorption accelerators such as quaternary ammonium compounds can also be included. Surface-active agents such as cetyl alcohol and glycerol monostearate can be included. Adsorptive carriers such as kaolin and bentonite can be added. Lubricants such as talc, calcium and magnesium stearate, and solid polyethyl glycols can also be included. Preservatives may also be added. The compositions of the invention can also contain thickening agents such as cellulose and/or cellulose derivatives. They may also contain gums such as xanthan, guar or carbo gum or gum arabic, or alternatively polyethylene glycols, bentones and montmorillonites, and the like.

For example, tablets or caplets containing the therapeutic agents of the invention can include buffering agents such as calcium carbonate, magnesium oxide and magnesium carbonate. Caplets and tablets can also include inactive ingredients such as cellulose, pre-gelatinized starch, silicon dioxide, hydroxy propyl methyl cellulose, magnesium stearate, microcrystalline cellulose, starch, talc, titanium dioxide, benzoic acid, citric acid, corn starch, mineral oil, polypropylene glycol, sodium phosphate, zinc stearate, and the like. Hard or soft gelatin capsules containing at least one therapeutic agent of the invention can contain inactive ingredients such as gelatin, microcrystalline cellulose, sodium lauryl sulfate, starch, talc, and titanium dioxide, and the like, as well as liquid vehicles such as polyethylene glycols (PEGs) and vegetable oil. Moreover, enteric-coated caplets or tablets containing one or more therapeutic agent of the invention are designed to resist disintegration in the stomach and dissolve in the more neutral to alkaline environment of the duodenum.

The therapeutic agents of the invention can also be formulated as elixirs or solutions for convenient oral administration or as solutions appropriate for parenteral administration, for instance by intramuscular, subcutaneous, intraperitoneal or intravenous routes. The pharmaceutical formulations of the therapeutic agents of the invention can also take the form of an aqueous or anhydrous solution or dispersion, or alternatively the form of an emulsion or suspension or salve.

Thus, the therapeutic agents may be formulated for parenteral administration (e.g., by injection, for example, bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion containers or in multi-dose containers. As noted
5 above, preservatives can be added to help maintain the shelf life of the dosage form. The therapeutic agents and other ingredients may form suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the therapeutic agents and other ingredients may be in powder form, obtained by
10 aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

These formulations can contain pharmaceutically acceptable carriers, vehicles and adjuvants that are well known in the art. It is possible, for example, to prepare solutions using one or more organic solvent(s) that is/are acceptable
15 from the physiological standpoint, chosen, in addition to water, from solvents such as acetone, ethanol, isopropyl alcohol, glycol ethers such as the products sold under the name "Dowanol," polyglycols and polyethylene glycols, C₁-C₄ alkyl esters of short-chain acids, ethyl or isopropyl lactate, fatty acid triglycerides such as the products marketed under the name "Miglyol," isopropyl
20 myristate, animal, mineral and vegetable oils and polysiloxanes.

It is possible to add, if necessary, an adjuvant chosen from antioxidants, surfactants, other preservatives, film-forming, keratolytic or comedolytic agents, perfumes, flavorings and colorings. Antioxidants such as t-butylhydroquinone, butylated hydroxyanisole, butylated hydroxytoluene and α -tocopherol and its
25 derivatives can be added.

Also contemplated are combination products that include one or more of the therapeutic agents of the present invention and one or more other ingredients. For example, the compositions can contain vitamins, minerals (e.g. calcium), anti-inflammatory agents and the like.

30 Additionally, the therapeutic agents are well suited to formulation as sustained release dosage forms and the like. The formulations can be so

constituted that they release the therapeutic agents, for example, in a particular part of the intestinal or respiratory tract, possibly over a period of time.

Coatings, envelopes, and protective matrices may be made, for example, from polymeric substances, such as polylactide-glycolates, liposomes,

5 microemulsions, microparticles, nanoparticles, or waxes. These coatings, envelopes, and protective matrices are useful to coat indwelling devices, e.g., tissue re-modeling devices, pins, splints, joint replacement devices and the like.

For topical administration, the therapeutic agents may be formulated as is known in the art for direct application to a target area. Forms chiefly
10 conditioned for topical application take the form, for example, of creams, milks, gels, dispersion or microemulsions, lotions thickened to a greater or lesser extent, impregnated pads, ointments or sticks, aerosol formulations (e.g., sprays or foams), soaps, detergents, lotions or cakes of soap. Other conventional forms for this purpose include wound dressings, coated bandages or other polymer
15 coverings, ointments, creams, lotions, pastes, jellies, sprays, and aerosols. Thus, the therapeutic agents of the invention can be delivered via patches or bandages for dermal administration. Alternatively, the therapeutic agent can be formulated to be part of an adhesive polymer, such as polyacrylate or acrylate/vinyl acetate copolymer. For long-term applications it might be desirable to use microporous
20 and/or breathable backing laminates, so hydration or maceration of the skin can be minimized. The backing layer can be any appropriate thickness that will provide the desired protective and support functions. A suitable thickness will generally be from about 10 to about 200 microns.

Ointments and creams may, for example, be formulated with an aqueous
25 or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. The therapeutic agents can also be delivered via iontophoresis, e.g., as disclosed in U.S. Patent Nos.
30 4,140,122; 4,383,529; or 4,051,842. The percent by weight of a therapeutic agent of the invention present in a topical formulation will depend on various

factors, but generally will be from 0.01% to 95% of the total weight of the formulation, and typically 0.1-85% by weight.

Drops, such as eye drops or nose drops, may be formulated with one or more of the therapeutic agents in an aqueous or non-aqueous base also comprising one or more dispersing agents, solubilizing agents or suspending agents. Liquid sprays are conveniently delivered from pressurized packs. Drops can be delivered via a simple eye dropper-capped bottle, or via a plastic bottle adapted to deliver liquid contents dropwise, via a specially shaped closure.

The therapeutic agents may further be formulated for topical administration in the mouth or throat. For example, the active ingredients may be formulated as a lozenge further comprising a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the composition in an inert base such as gelatin and glycerin or sucrose and acacia.

The pharmaceutical formulations of the present invention may include, as optional ingredients, pharmaceutically acceptable carriers, diluents, solubilizing or emulsifying agents, and salts of the type that are available in the art. Examples of such substances include normal saline solutions such as physiologically buffered saline solutions and water. Specific non-limiting examples of the carriers and/or diluents that are useful in the pharmaceutical formulations of the present invention include water and physiologically acceptable buffered saline solutions such as phosphate buffered saline solutions pH 7.0-8.0.

The therapeutic agents of the invention can also be administered to the respiratory tract. Thus, the present invention also provides aerosol pharmaceutical formulations and dosage forms for use in the methods of the invention. In general, such dosage forms comprise an amount of at least one of the agents of the invention effective to treat or prevent the clinical symptoms of a specific bone disorder or disease. Any statistically significant attenuation of one or more symptoms of the disorder or disease that has been treated pursuant to the method of the present invention is considered to be a treatment of such disorder or disease within the scope of the invention.

Alternatively, for administration by inhalation or insufflation, the composition may take the form of a dry powder, for example, a powder mix of the therapeutic agent and a suitable powder base such as lactose or starch. The powder composition may be presented in unit dosage form in, for example, capsules or cartridges, or, e.g., gelatin or blister packs from which the powder may be administered with the aid of an inhalator, insufflator, or a metered-dose inhaler (see, for example, the pressurized metered dose inhaler (MDI) and the dry powder inhaler disclosed in Newman, S. P. in Aerosols and the Lung, Clarke, S. W. and Davia, D. eds., pp. 197-224, Butterworths, London, England, 1984).

Therapeutic agents of the present invention can also be administered in an aqueous solution when administered in an aerosol or inhaled form. Thus, other aerosol pharmaceutical formulations may comprise, for example, a physiologically acceptable buffered saline solution containing between about 0.1 mg/ml and about 100 mg/ml of one or more of the therapeutic agents of the present invention specific for the indication or disease to be treated. Dry aerosol in the form of finely divided solid particles that are not dissolved or suspended in a liquid are also useful in the practice of the present invention. Therapeutic agents of the present invention may be formulated as dusting powders and comprise finely divided particles having an average particle size of between about 1 and 5 μm , alternatively between 2 and 3 μm . Finely divided particles may be prepared by pulverization and screen filtration using techniques well known in the art. The particles may be administered by inhaling a predetermined quantity of the finely divided material, which can be in the form of a powder. It will be appreciated that the unit content of active ingredient or ingredients contained in an individual aerosol dose of each dosage form need not in itself constitute an effective amount for treating the particular indication or disease since the necessary effective amount can be reached by administration of a plurality of dosage units. Moreover, the effective amount may be achieved using less than the dose in the dosage form, either individually, or in a series of administrations.

For administration to the upper (nasal) or lower respiratory tract by inhalation, the therapeutic agents of the invention are conveniently delivered from a nebulizer or a pressurized pack or other convenient means of delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as
5 dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Nebulizers include, but are not limited to, those described in U.S. Patent Nos. 4,624,251; 3,703,173; 3,561,444; and 4,635,627. Aerosol delivery
10 systems of the type disclosed herein are available from numerous commercial sources including Fisons Corporation (Bedford, Mass.), Schering Corp. (Kenilworth, NJ) and American Pharmoseal Co., (Valencia, CA). For intra-nasal administration, the therapeutic agent may also be administered via nose drops, a liquid spray, such as via a plastic bottle atomizer or metered-dose inhaler.
15 Typical of atomizers are the Mistometer (Wintrop) and the Medihaler (Riker).

Furthermore, the active ingredients may also be used in combination with other therapeutic agents, for example, pain relievers, anti-inflammatory agents, vitamins, minerals and the like, whether for the conditions described or some other condition.

20 The present invention further pertains to a packaged pharmaceutical composition for controlling bone disorders such as a kit or other container. The kit or container holds a therapeutically effective amount of a pharmaceutical composition for controlling bone disorders and instructions for using the pharmaceutical composition for control of the bone disorder. The
25 pharmaceutical composition includes at least one therapeutic agent of the present invention, in a therapeutically effective amount such that bone disorder is controlled.

All references cited herein are incorporated by reference in their entirety.
30 The following examples illustrate certain aspects of the invention and are not intended to limit the scope thereof.

EXAMPLES

The following materials were used in these examples.

Primary human mesenchymal cells, primary human osteoblasts and
5 corresponding media were purchased from Biowhittaker (Walkersville, MD).
Mouse mesenchymal C3H10T1/2 cells were obtained from American Type
Culture Collection (Manassas, VA) (ATCC Deposit No. CCL-226). Total RNA
prepared from abdominal adipose tissue was obtained from Biochain Institute
(Hayward, CA). General tissue culture reagents, first strand cDNA synthesis
10 kits and Platinum Taq DNA polymerase were purchased from Invitrogen
(Rockville, MD). Noggin, gremlin, chordin, twisted gastrulation, and bone
morphogenetic proteins 2, 4 and 6 were purchased from R & D Systems
(Minneapolis, MN). Strataprep Total RNA mini-prep kits were purchased from
Stratagene (San Diego, CA). Sephadex G-25 pre-packed NAP-5 columns were
15 purchased from Amersham Biosciences (Piscataway, NJ). Reagents for
determining alkaline phosphatase (ALP) activity were obtained from Pierce-
Endogen (Rockford, IL). Prolagen C ELISA kits were purchased from Quidel
Corporation (Mountainview, CA). CellTiterGlo Cell Viability kits were
obtained from Promega (Madison, WI). Homogeneous Caspase ELISA, Cell
20 Death assay (Histone-Associated DNA Fragmentation), and In Situ Cell Death
Detection kits were purchased from Roche Diagnostics (Indianapolis, IN).
Caspase inhibitors (Caspase-1, Inhibitor VI and Caspase-3, Inhibitor I) were
obtained from Calbiochem (San Diego, CA). All other reagents and chemicals
were purchased from Sigma (St. Louis, MO).

25

EXAMPLE 1: Expression of SOST Correlates with Commitment and Differentiation of Precursor Cells to Osteogenic Lineage

In this example, the SOST expression was correlated with cellular
30 differentiation and was found to increase as progenitor cells developed into
osteoblasts.

SOST Expression Assay

Human mesenchymal (hMSC) cells were cultured in regular growth media (Biowhittaker's MSCGM) or Osteoblast-inducing media (Biowhittaker's MSCGM medium supplemented with 100 nM dexamethasone, 50 µg/ml ascorbic acid, and 10 mM β-glycerophosphate). Media were refreshed twice per week. Cultures were harvested 1, 7 and 21 days after plating and RNA isolated for RT-PCR analyses of SOST. SOST expression was also determined in adipocytes and chondrocytes generated from mesenchymal cells cultured for 21 to 28 days in corresponding media (Pittenger et al.). Primary cultures of human osteoblasts were also grown in Osteoblast-inducing media for 21 days and RNA isolated for RT-PCR. SOST expression was also analyzed in preparations of total RNA isolated from abdominal adipose tissue and from cartilagenous ends of long bones.

Results

Cell culture conditions necessary for human-derived mesenchymal (hMSC) cells to differentiate into osteoblast-like cells were developed. Mesenchymal cells are a population of bone marrow-derived pluripotent cells capable of differentiating into a number of different tissue types including bone, cartilage, fat, smooth muscle, tendon, marrow stroma, and neurons (Owen, 1998).

Figure 1a provides an RT-PCR analyses of RNA prepared from primary human osteoblasts, undifferentiated hMSC cells and hMSC cells differentiated to yield osteoblastic, chondrocytic and adipocytic cells. The expression of characteristic phenotypic markers (e.g. PPARγ2 for adipocytes) in these cell types was compared with that of SOST. As can be observed in Figure 1a, undifferentiated hMSC cells and adipocytes expressed negligible levels of SOST. In contrast, the cell types capable of mineralization that express markers associated with the osteoblast phenotype (type I collagen, type X collagen, parathyroid hormone receptor (PTHr)) expressed SOST. These included cultures

of primary human osteoblasts, hMSC cells differentiated to yield osteoblastic cells and hypertrophic chondrocytes.

The expression of the SOST in hMSC cells that had differentiated into osteoblast-like cells was correlated with the stage of osteoblast differentiation by observing hMSC cells grown in osteoblast-inducing medium over a 21-day period. hMSC cells differentiated into osteoblasts as indicated by enhanced histochemical staining for ALP and collagen (Figure 1b, darker stains in right panel at 21 days respectively) and deposition of mineral (negligible at Day 7 versus 437 ± 60 μg calcium/mg protein at Day 28). Undifferentiated hMSC cells expressed negligible levels of SOST mRNA as well as negligible levels of mRNA correlating with alkaline phosphatase (ALP) activity, collagen and mineral deposition (Figure 1b, left panel). In contrast, the expression of ALP, collagen and SOST increased with time in cells grown in osteoblast-inducing media.

These results indicate that SOST expression increases in cells committed to osteoblast differentiation.

EXAMPLE 2: Regulation of SOST Expression by Growth Factors and Hormones in Human Bone Cells

In this example, the effect of growth factors and hormones on SOST expression in hMSC cells or primary human osteoblasts was observed using RT-PCR. Prior to measuring SOST expression, these cells were treated for 72 hrs with BMP-2, BMP-4, BMP-6, insulin-like growth factor-1 (IGF-1), parathyroid hormone (PTH), transforming growth factor- β (TGF- β), dexamethasone (DEX), retinoic acid (RA), and 1,25-dihydroxyvitamin D₃ (vit D). As immature osteoblasts differentiate and become capable of mineralization, they express markers associated with the osteoblast phenotype (type I collagen and parathyroid hormone receptor (PTHr)). These markers were used to ascertain whether differentiation had occurred so that the influence of growth factors and hormones could be correlated with the stage differentiation as well as the level of SOST expression.

Methods

Human mesenchymal cells and primary human osteoblasts were plated in regular growth media containing 2% FCS at a density of 10,000 cells / cm². Test reagents (10⁻⁷ M dexamethasone, 10⁻⁶ M retinoic acid, 10 ng/ml TGF-β1, PTH (10⁻⁸ M), 10 or 50 ng/ml IGF-1, 10⁻⁷ M 1,25-dihydroxyvitamin D₃, and 50 to 300 ng/ml of bone morphogenetic proteins 2, 4 or 6) were added singly or in combination on the following day. Cultures were continued for 24 to 120 hrs after which the cells were harvested for RNA isolation.

10

Results

Figures 2 to 6 show the results of these experiments. As can be seen in Figure 2, while untreated hMSC cells ("vehicle") strongly expressed type I collagen, they had negligible levels of PTHr and SOST (Vehicle and Control, 21 days). These results indicate that untreated hMSC cells are in an early stage of osteoblast lineage, but are committed to osteoblastogenesis. Treatment of these cells with dexamethasone, bone morphogenetic proteins, IGF-1 (IGF-1 at 50 ng/ml) or long-term culture in osteoblast-inducing media (Induced, 21 days) advanced the stage of differentiation and induced PTHr expression.

Several factors also enhanced the expression of SOST, including BMP-4, BMP-6, and IGF-1. Interestingly, factors such as vitamin D, PTH, TGF-β and dexamethasone by themselves had no effect on SOST expression. Bone morphogenetic proteins enhanced the expression of SOST in differentiating hMSC cells and in primary cultures of human osteoblasts.

Figure 3 provides a time course of BMP-induced expression of SOST in hMSC and primary human osteoblasts. As shown, maximum levels of SOST expression were achieved within 72 to 120 hrs after start of treatment (Figure 3).

The proteins noggin and gremlin are thought to function as BMP antagonists by binding and inactivating bone morphogenetic proteins (Yamaguchi et al. 2000). Bone morphogenetic proteins are also thought to regulate the expressions of noggin and gremlin in osteoblastic cells in culture.

30

See Gazzero et al. (1998); Pereira et al (2000); and Nifuji & Noda (1999). The effect of bone morphogenetic proteins on SOST expression was compared to the effect of bone morphogenetic proteins on noggin and gremlin expression.

Figure 4 shows that while the basal levels of SOST and noggin were negligible in undifferentiated hMSC cells, the levels of these genes were up-regulated by BMPs-2, 4 and 6. In contrast, gremlin was constitutively expressed in undifferentiated hMSC cells and its expression was only modestly increased by the bone morphogenetic proteins.

Steroids such as retinoids have been shown to regulate the differentiation of mesenchymal and osteoblastic cells (Gazit et al, 1999; Weston et al., 2000). The cellular response to steroids is modulated through interplay between the steroid hormone and TGF- β signaling pathways (Yanagi et al., 1999).

To determine whether steroids and/or the TGF- β signaling pathway affect BMP-mediated regulation of SOST expression, hMSC and osteoblastic cells were incubated with vitamin D, retinoic acid, and dexamethasone alone or in combination with BMP-4 for 72 hrs prior to RT-PCR analyses of SOST expression.

As shown in Figure 5, steroids alone (vitamin D, retinoic acid, and dexamethasone) had no effect on the basal expression of SOST in hMSC or osteoblastic cells. However, BMP-4 enhanced SOST expression (Figure 5). Interestingly, retinoic acid and vitamin D enhanced the levels of SOST when added in combination with BMP-4 such that the levels detected were greater than those found in cells treated with BMP-4 alone.

In contrast, there was a decrease in SOST expression levels when dexamethasone was added with BMP-4. Therefore, dexamethasone apparently abolished the stimulatory effect of BMP on SOST expression (Figure 5).

BMP-4 also enhanced gremlin and noggin expression (Figure 6). However, steroids affected BMP-4 induced expression of gremlin and noggin somewhat differently than BMP-4 induced expression of SOST (Figure 6). Like SOST, gremlin expression was unchanged in cells treated only with retinoic acid or vitamin D. Unlike SOST, basal levels of gremlin decreased when cells were

treated with dexamethasone alone. But the stimulatory effect of BMP-4 on gremlin expression was diminished when retinoic acid and dexamethasone were added with BMP-4. Vitamin D did not alter the effect of BMP-4 on gremlin expression.

5 In contrast, BMP-4, retinoic acid, dexamethasone and vitamin D all appeared to increase the levels of noggin in hMSC cells. BMP-4 exerted the greatest effect. The stimulatory effect of BMP-4 on noggin was not greatly altered by steroids.

10 Therefore, the effects of steroids and BMP-4 on gremlin and noggin contrasted markedly with those on SOST (Figure 6). BMP-4 significantly increased the levels of SOST expression and this effect was enhanced in the presence of vitamin D or retinoic acid but not with dexamethasone. In fact, dexamethasone significantly suppressed SOST expression.

15 **EXAMPLE 3: Compounds Affecting SOST Expression**

 The previous example illustrates that SOST expression was enhanced by bone morphogenetic proteins and modulated by steroids such as dexamethasone. In this example, human MSC cells were incubated with bone morphogenetic proteins in the presence of various test agents to determine whether other
20 steroids and compounds could affect the BMP-induction of SOST. SOST expression was monitored by reverse transcriptase polymerase chain reaction (RT-PCR).

Materials and Methods

25 Human mesenchymal cells (hMSCs) were plated in regular growth media containing 2% FCS and treated with BMP-6 either alone or in combination with the test compounds. The cells were harvested 72 hrs later and RNA was prepared for RT-PCR analysis of SOST expression.

30 **Results**

The compounds tested and the results obtained for these compounds are summarized in Table 1.

Table 1

Compound	Source	Compound Type	Concentration Tested	Results
Dexamethasone	SIGMA-Aldrich	Glucocorticoid analog	1×10^{-7} M	Inhibition
Triamcinolone	SIGMA-Aldrich	Glucocorticoid analog	1×10^{-7} M	Inhibition
Fluocinolone acetonide	SIGMA-Aldrich	Glucocorticoid analog	1×10^{-7} M	Inhibition
Ursodeoxycholic acid	SIGMA-Aldrich	Bile salt	1×10^{-6} M 3×10^{-6} M	Inhibition Inhibition
Tauroursodeoxycholic acid	SIGMA-Aldrich	Bile salt	2×10^{-7} M 2×10^{-6} M	No Effect No Effect
Prostaglandin E2	Biomol	Eicosanoid	1×10^{-7} M 1×10^{-6} M	Inhibition Inhibition
Spironolactone	SIGMA-Aldrich		1×10^{-7} M 1×10^{-6} M	No Effect No Effect
1,2-napthoquinone-4-sulfate	SIGMA-Aldrich		1×10^{-6} M 5×10^{-6} M	No Effect No Effect
17 β -estradiol	SIGMA-Aldrich	Estrogen	1×10^{-7} M 1×10^{-6} M	No Effect No Effect
ICI 182,780	TOCRIS	Estrogen receptor- α antagonist	1×10^{-6} M	No Effect
Lovastatin /Mevinolin	SIGMA-Aldrich	Statin, HMG-CoA reductase inhibitor	1×10^{-7} M	No Effect
PD98059	Calbiochem	MEK inhibitor	2×10^{-6} M	No Effect
SB203580	Calbiochem	P38 inhibitor	2×10^{-6} M	No Effect
SB202474	Calbiochem	MAPK inhibitor	2×10^{-6} M	No Effect

Compound	Source	Compound Type	Concentration Tested	Results
		control		
Parathyroid Hormone (1-34)	Biomol	Calciotropic Peptide	1×10^{-7} M	No Effect

Figures 7 and 8 illustrate the results obtained for several compounds that blocked BMP-6 mediated induction of SOST expression in hMSC cells.

Triamcinolone (Triam), a glucocorticoid analog similar to dexamethasone, blocked the stimulatory effect of BMP-6 on SOST (Figure 7). Ursodeoxycholic acid (Urso), a bile acid used to treat primary biliary cirrhosis, also blocked the BMP-6 induction of SOST in a dose-dependent manner (Figure 7). In particular, ursodeoxycholic acid at 3×10^{-6} M (Urso-2) completely blocked the induction of SOST by BMP-6 (Figure 7).

Figure 8 shows the results of treatment of hMSC cells with fluocinolone acetonide (Fluo) and prostaglandin E2 (PGE2). Both compounds effectively blocked the induction of SOST by BMP-6 whereas other compounds (tauroursodeoxycholic acid, ICI 182,780, estrogen and lovastatin) had minimal effects.

The structures for the five compounds that effectively blocked the induction of SOST expression by BMP-6 are shown in Figure 9.

These results indicate that 3 different types of glucocorticoid analogs were capable of blocking the induction of SOST expression by BMPs. Other compounds that also blocked SOST expression included bile salts (e.g.

ursodeoxycholic acid), and prostaglandins.

EXAMPLE 4: Exogenously Added Sclerostin Regulates Expression of Osteoblast Function

Individuals with heterozygous or homozygous SOST mutations have a skeletal phenotype characterized by denser and heavier bones (Beighton et al. 1976). In this example, the sclerostin gene product was added to *in vitro*

cultures of osteoblasts. Osteoblastic phenotypic markers were used to monitor the stage of differentiation at which the sclerostin protein affected osteoblastic function.

5 **Determining Effects of Sclerostin Protein on Human Mesenchymal Cells**

hMSC cells were plated in 96-well tissue culture dishes at a density of 10,000 cells/cm² in Osteoblast-Inducing medium. Partially purified preparations
10 of baculovirus-expressed sclerostin protein were prepared in sterile PBS using NAP-5 columns prior to use. Human sclerostin protein (0 to 30 µg/ml) or an equal volume of Sf9 conditioned media (Control) was added to cultures of hMSC cells at various times after plating (1 day, 8 days, 15 days, or 21 days). The effects of sclerostin on osteoblastic differentiation were assessed by
15 measuring alkaline phosphatase activity (ALP, determined in cell layers using DEAA buffer (Pierce) containing 0.5% NP-40 and 10 mM p-nitrophenylphosphate), synthesis of collagen type I (Prolagen C ELISA), and calcium deposition for mineralization (colorimetric assay of acid lysates of cell layers, Sigma).

20

Determining the Effects of Sclerostin Protein on Mouse Mesenchymal C3H10T1/2 Cells

C3H10T1/2 cells (ATCC Deposit No. CCL-226) were plated in 96-well
25 dishes at a density of 25,000 cells per well in complete growth medium (DMEM with high glucose and glutamine supplemented with 10% FCS, 1 % penicillin /streptomycin, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, 55 µM β-mercaptoethanol, and 20 mM HEPES, pH 7.3). C3H10T1/2 cells were used in a short-term (72 hr) assay to determine the effects of human sclerostin
30 protein on BMP-induced ALP activity. Partially purified preparations of baculovirus-expressed sclerostin protein were prepared in sterile PBS using NAP-5 columns prior to use. Sclerostin protein (0 to 50 µg/ml) or an equal volume of Sf9 conditioned media (control) was pre-incubated with 500 ng/ml

BMP-6 for 1 hr prior to addition to cells. For comparison, similar incubations were carried out with anti-BMP-6 antibody and noggin. Cells were harvested 72 hrs later for determination of ALP activity.

To test the specificity of the sclerostin effect, human sclerostin that was synthesized with a FLAG tag, was pre-incubated for 2 hrs at 4 °C with an anti-FLAG antibody coupled to agarose beads. The mixture was then spun down at 10,000xg for 15 minutes at 4 °C and the supernatant treated as “sclerostin protein” in the above assay.

10 Results

Figure 10 provides a comparison of the effects of human sclerostin and mouse noggin proteins on alkaline phosphatase (ALP) activity in differentiating hMSC cells. Alkaline phosphatase activity is a phenotypic marker for osteoblast differentiation. After 7 days of treatment, noggin at 1 µg/ml had no effect on alkaline phosphatase activity whereas noggin at 5 µg/ml significantly increased alkaline phosphatase activity ($p < 0.001$ to vehicle-treated cells).

In contrast, under the same conditions, partially purified preparations of human sclerostin consistently decreased alkaline phosphatase activity when administered at either 1 or 5 µg/ml (Figure 10).

Sclerostin also decreased alkaline phosphatase activity in primary cultures of human osteoblasts (Figure 11). Osteoblasts that were grown in regular growth media expressed basal levels of alkaline phosphatase (Figure 11, far right panel); these levels increased significantly when cells were grown in Inducing media (middle panel, darker stain). When a similar group of cells were treated with human sclerostin, there was a marked decrease in the amount of alkaline phosphatase expressed, as evidenced by the reduction in stain (Figure 11, Sclerostin + Inducing Media).

Sclerostin decreased alkaline phosphatase activity and mineralization in hMSC cells in a dose-dependent manner (Figure 12). Alkaline phosphatase activity and deposition of mineral (as measured by calcium assay) were significantly decreased in cells treated with increasing concentrations of partially

purified preparations of human sclerostin protein but not the control (proteins purified from conditioned Sf9 media) (Figure 12). Alkaline phosphatase activity and mineral deposition were decreased by about 50 % in cells treated with about 10 to 15 $\mu\text{g}/\text{ml}$ of the protein ($p < 0.001$ to control).

5 Synthesis of collagen type I was also decreased in a dose-dependent manner in sclerostin-treated cultures. Partially purified preparations of human sclerostin reduced the synthesis of type I collagen by about 50 % in cells treated with 10 $\mu\text{g}/\text{ml}$ of the protein ($p < 0.001$ to control, Figure 13).

10 To determine the effect of sclerostin protein on the expression of osteoblastic markers in cultures of differentiating hMSC cells, RNA was isolated after a 40-hr treatment and analyzed by RT-PCR (Figure 14). As can be seen in Figure 14, RNA levels of markers for adipocytes (PPAR γ 2) or undifferentiated mesenchymal cells (endoglin) were not affected by treatment with partially purified preparations of human sclerostin protein. On the other hand, RNA levels
15 for osteoblast phenotypic markers such as PTHr, Type I collagen, BMP-2 and BMP-6 were significantly reduced in cells treated with partially purified preparations of human sclerostin protein compared to cells treated with proteins purified from conditioned Sf9 media (control).

 BMPs modulate the activity of osteoblasts. Treatment of osteoprogenitor
20 or pre-osteoblasts with BMPs can enhance the activity of markers such as ALP (Gazzerro et al. 1998, Nifuji & Noda, 1999). In cultures of hMSC, BMP-6 increased the activity of ALP about 2-fold (Figure 15). Sclerostin reduced this BMP-induced ALP activity in a dose-dependent manner (ANOVA, $p < 0.0001$). ALP levels in cells treated with BMP-6+30 $\mu\text{g}/\text{ml}$ sclerostin were significantly
25 lower than that expressed in BMP-6 treated or vehicle-treated cells ($p < 0.0001$). Indeed, sclerostin also significantly reduced the basal levels of ALP in a dose-dependent manner (ANOVA, $p < 0.0001$).

 In Figure 16, increasing concentrations of BMP-6 partially reversed the inhibitory effect of sclerostin on alkaline phosphatase activity in hMSC cultures.

30 These findings show that sclerostin interacts with BMPs in regulating osteoblast function.

In mouse mesenchymal C3H10T1/2 cells, ALP activity was increased in a dose-dependent manner by BMP-6 (Figure 17). The stimulatory effect of BMP-6 on alkaline phosphatase activity in these cells was antagonized by mouse noggin, by partially purified preparations of human sclerostin and by a neutralizing antibody to BMP-6. Figure 17 shows that mouse noggin decreased BMP-6 stimulated alkaline phosphatase activity with an IC₅₀ of 0.3 µg/ml, data that contrast with the findings of Abe et al (2000). The anti-BMP-6 antibodies decreased BMP-6 stimulated alkaline phosphatase activity by 50 % with an IC₅₀ of 0.49 µg/ml. In addition, the partially purified preparation of human sclerostin effectively decreased BMP-6 stimulated alkaline phosphatase activity in a dose-dependent manner, with an IC₅₀ of 10 µg/ml (Figure 17).

To verify the specificity of the sclerostin response, a human sclerostin fusion protein was prepared that had a FLAG tag fused to the sclerostin coding region. Before addition to mouse mesenchymal C3H10T1 cells, the human sclerostin-FLAG preparation was immuno-depleted with an anti-Flag M2 antibody/agarose complex. As can be seen in Figure 17, pre-incubation of the sclerostin-FLAG preparation with an anti-Flag antibody/agarose bead complex completely abolished sclerostin antagonism of BMP-6 in the C3H10T1/2 cells. These data indicate that the diminution of BMP-6 induced alkaline phosphatase activity observed upon addition of partially purified sclerostin protein is specifically due to sclerostin, rather than to some unidentified co-purified substance.

EXAMPLE 5: Sclerostin Causes Apoptosis

Apoptosis is mediated by the activation of a series of cysteine proteases known as caspases (Thornberry & Lazebnik, 1998). Caspases activate proteins such as Bax that promote apoptosis as well as proteins that exhibit protective properties (for example, I-TRAF and Survivin). The initiation of the apoptosis cascade can occur through a number of different mechanisms including the recruitment of, and the ligand binding to, cell surface Death receptors such as FAS (Vaughan et al. 2002; Budd, 2002). In the present example, the role of

sclerostin in the survival of osteoblasts is investigated by incubating hMSC cells with sclerostin and measuring the levels of caspases and other pro-apoptotic factors.

5 Cell Viability and Apoptosis Assays

Human mesenchymal (hMSC) cells were plated in Osteoblast-inducing media (Biowhittaker's MSCGM medium supplemented with 100 nM dexamethasone, 50 µg/ml ascorbic acid, and 10 mM β-glycerophosphate) at a density of 10,000 cells /cm² in 96-well dishes.

10 For cell viability assays, cells were treated with a partially purified preparation of baculovirus-expressed human sclerostin (0 to 20 µg/ml) or an equal volume of a protein preparation purified from Sf9 conditioned media (control) for 1 week. Sclerostin and control were refreshed upon media change. Cells were then lysed and processed using Promega's CellTiterGlo

15 Luminescence Viability Assay.

For apoptosis assays, cells were treated with sclerostin or commercially available preparations of the BMP antagonists noggin, chordin, gremlin or twisted gastrulation (Tsg) for 24 hours prior to assay for caspase activity. In some experiments, cells were treated for various times with a partially purified
20 preparation of baculovirus-expressed human sclerostin (0 to 30 µg/ml) or an equal volume of a protein preparation purified from Sf9 conditioned media (control) in the absence or presence of Calbiochem's caspase inhibitors. Cells were harvested and processed for apoptosis assays using the Homogeneous Caspase ELISA (Roche) or Cell Death (Histone-Associated DNA
25 Fragmentation, Roche).

For apoptosis determination by TUNEL staining, cells were treated for 24 hrs with a partially purified preparation of baculovirus-expressed human sclerostin (20 µg/ml) or an equal volume of a protein preparation purified from Sf9 conditioned media (control). Cells were fixed with paraformaldehyde and
30 processed using Roche's In Situ Cell Death Detection kit.

In another study, hMSC were treated with vehicle or sclerostin for 6 hrs or 48 hrs. Cells were harvested, RNA prepared and used for the analysis of a commercially available Apoptosis Gene Array (SuperArray, Bethesda, MD).

5 Results

Figure 20 shows the effect of a 7-day treatment of hMSC cells with a partially purified preparation of human sclerostin. Cells were lysed and processed with Promega's CellTiter Glo Viability assay. At this point in time, these cells represent committed osteoprogenitors / pre-osteoblasts. They express
10 high levels of Type I collagen and low levels of alkaline phosphatase. Treatment with increasing concentrations of human sclerostin significantly reduced the number of hMSC cells remaining in the culture compared to cells treated with the control (proteins purified from Sf9 conditioned media).

Treatment of hSMC with sclerostin significantly increased caspase
15 activity (Figure 21). An inactive preparation of sclerostin (as determined by lack of effects in other biological and biochemical assays) did not significantly affect caspase activity. When commercially available BMP antagonists (noggin, chordin, gremlin, Tsg) were added to hSMC under the same conditions, no increase in caspase activity was observed. These findings show that sclerostin
20 selectively increases the apoptosis of human bone cells.

Figure 22 illustrates that increasing concentrations of partially purified human sclerostin significantly elevated the activities of caspases as well as the amounts of histone-associated DNA fragments. DNA fragmentation can also be detected by TUNEL staining using commercially available reagents. Increased
25 staining for apoptotic nuclei can be seen in sclerostin-treated as compared to control (proteins purified from Sf9 conditioned media)-treated cells in Figure 23 (see arrows). Sclerostin decreases the survival of osteoblastic cells by promoting their apoptosis.

Sclerostin treatment of hMSC cells elevated caspase activity for up to 72
30 hrs (Figure 24a). Rat sclerostin also increased the apoptosis of hMSC cells but to a lesser extent than the human protein (Figure 24b).

The induction of caspase activity by sclerostin in hMSC cells could be effectively decreased by commercially available caspase inhibitors (Figure 25). Caspase-1 inhibitor and Caspase-3 inhibitor effectively blocked the induction of caspase activity by human sclerostin. The combination of the two inhibitors was
5 more potent than either one alone.

To further investigate the effects of sclerostin on the apoptosis of hMSC, RNA was harvested from vehicle and sclerostin-treated cells and used the RNA to probe a cDNA expression array containing apoptosis genes. The results for vehicle and sclerostin-treated samples were compared and data (corrected for
10 background) were expressed as a ratio of sclerostin to vehicle. The data is summarized in the table below.

Incubation Time	Gene	Function	Ratio
6 hrs	BAX	Proapoptotic	1.53±0.2
	Bcl10	Proapoptotic	4.09±2.13
	Hrk	Proapoptotic	1.59±0.39
	Myd88	Proapoptotic	1.4±0.31
	Caspase 3	Proapoptotic	2.46±0.69
	Caspase 4	Proapoptotic	1.83±1.12
	Caspase 7	Proapoptotic	1.76±0.18
	XIAP	Anti-apoptotic	2.01±0.64
	Survivin, IAP2	Anti-apoptotic	1.32±0.48
	Bruce	Anti-apoptotic	1.45±0.75
48 hrs	Bak1	Proapoptotic	2.36±0.74
	Blk	Proapoptotic	1.83±0.12
	Myd88	Proapoptotic	2.64±1.3
	Bax	Proapoptotic	1.64±0.3
	Bcl10	Proapoptotic	1.84±0.76

	I-TRAF	Anti-apoptotic	2.05±0.54
	Casper	Anti-apoptotic	2.68±0.28
	Mdm2	Anti-apoptotic	1.66±0.34

In hMSC treated for 6 hrs with sclerostin, there was a significant 2 to 4-fold increase in the expression of apoptosis-associated genes such as caspase 3, caspase 4, caspase 7 and the Bcl-2 family members, Bax and Hrk as well as genes involved in cell survival (XIAP, survivin, and Bruce which have been reported to inhibit caspases 3, 7 and 9). By 48 hrs, there was increased expression of proapoptotic members of the Bcl-2 family as well as anti-apoptotic genes (Casper, I-TRAF). Thus treatment of hSMC with sclerostin triggers the induction of proapoptotic as well as anti-apoptotic genes.

In Figure 25, two commercially available caspase inhibitors (Calbiochem's Caspase-1, Inhibitor VI which blocks caspases 1 and 4, and Caspase-3, Inhibitor I which blocks caspases 3, 6, 7, 8 and 10) blocked sclerostin-induced caspase activities in hMSC cultures. These findings, in addition to those from the apoptosis array, suggest that sclerostin enhances the apoptosis of osteoblastic cells via a FAS-mediated pathway.

Summary

The data shown in these examples demonstrate that sclerostin interacts with BMPs to modulate the activity of osteoblastic cells. This ability of sclerostin to interact with important growth factors is also likely the basis by which it modulates the survival of osteoblasts. By making the growth factors unavailable for cell function, sclerostin may increase the apoptosis of bone cells. Thus agents that block the ability of sclerostin to decrease cell survival and osteoblastic activity offer promise as agents useful in restoring lost bone.

EXAMPLE 6: Small Interfering RNAs (SiRNAs) to Block Expression of SOST

SiRNAs were designed using the guidelines provided by Ambion (Austin, TX). Briefly, the SOST cDNA sequence was scanned for target sequences that had AA dinucleotides. Sense and anti-sense oligonucleotides
 5 were generated to these targets (AA + 3' adjacent 19 nucleotides) that contained a G/C content of 35 to 55%. These sequences were then compared to others in the human genome database to minimize homology to other known coding sequences (Blast search).

The target and siRNA sequences designed are provided below.

10

Target sequence 1: AAGAATGATGCCACGGAAATC (SEQ ID NO:9)

Position in gene sequence: 140

GC content: 42.9%

Sense strand siRNA: GAAUGAUGCCACGGAAAUCtt (SEQ ID NO:10)

15 Antisense strand siRNA: GAUUUCCGUGGCAUCAUUCtt (SEQ ID NO:11)

Target sequence 3: AATGATGCCACGGAAATCATC (SEQ ID NO:12)

Position in gene sequence: 143

GC content: 42.9%

20 Sense strand siRNA: UGAUGCCACGGAAAUCAUCtt (SEQ ID NO:13)

Antisense strand siRNA: GAUGAUUCCGUGGCAUCAtt (SEQ ID NO:14)

Target sequence 5: AACACAAGACCATGAACCGG (SEQ ID NO:15)

Position in gene sequence: 209

25 GC content: 47.6%

Sense strand siRNA: CAACAAGACCAUGAACCGGtt (SEQ ID NO:16)

Antisense strand siRNA: CCGGUUCAUGGUCUUGUUGtt (SEQ ID NO:17)

Target sequence 27: AATTGAGAGTCACAGACACTG (SEQ ID NO:18)

30 Position in gene sequence: 950

GC content: 42.9%

Sense strand siRNA: UUGAGAGUCACAGACACUGtt (SEQ ID NO:19)

Antisense strand siRNA: CAGUGUCUGUGACUCUCAAtt (SEQ ID NO:20)

35 Target sequence 28: AAATGGAAGCATTTTCACCGC (SEQ ID NO:21)

Position in gene sequence: 1035

GC content: 42.9%

Sense strand siRNA: AUGGAAGCAUUUUCACCGCtt (SEQ ID NO:22)

Antisense strand siRNA: GCGGUGAAAAUGCUUCCAUtt (SEQ ID NO:23)

40

Target sequence 30: AAAGTCCAGGGACTGGTTAAG (SEQ ID NO:24)

Position in gene sequence: 1093

- GC content: 47.6%
 Sense strand siRNA: AGUCCAGGGACUGGUUAAGtt (SEQ ID NO:25)
 Antisense strand siRNA: CUUAACCAGUCCCUGGACUtt (SEQ ID NO:26)
- 5 Target sequence 31: AAGAAAGTTGGATAAGATTCC (SEQ ID NO:27)
 Position in gene sequence: 1111
 GC content: 33.3%
 Sense strand siRNA: GAAAGUUGGAUAAGAUUCCtt (SEQ ID NO:28)
 Antisense strand siRNA: GGAAUCUUAUCCAACUUUCtt (SEQ ID NO:29)
- 10 Target sequence 36: AACTGTAGATGTGGTTTCTAG (SEQ ID NO:30)
 Position in gene sequence: 1201
 GC content: 38.1%
 Sense strand siRNA: CUGUAGAUGUGGUUUCUAGtt (SEQ ID NO:31)
- 15 Antisense strand siRNA: CUAGAAACCACAUCUACAGtt (SEQ ID NO:32)
- Target sequence 40: AATTCTCCTTCGGGACCTCAA (SEQ ID NO:33)
 Position in gene sequence: 1269
 GC content: 47.6%
- 20 Sense strand siRNA: UUCUCCUUCGGGACCUCAAtt (SEQ ID NO:34)
 Antisense strand siRNA: UUGAGGUCCCGAAGGAGAAtt (SEQ ID NO:35)
- Target sequence 47: AAAGAGAGAGAATGAATGCAG (SEQ ID NO:36)
 Position in gene sequence: 1414
- 25 GC content: 38.1%
 Sense strand siRNA: AGAGAGAGAAUGAAUGCAGtt (SEQ ID NO:37)
 Antisense strand siRNA: CUGCAUUCAUUCUCUCUCUtt (SEQ ID NO:38)
- Target sequence 63: AAGAAGCTATGCTGCTTCCCA (SEQ ID NO:39)
 Position in gene sequence: 1590
- 30 GC content: 47.6%
 Sense strand siRNA: GAAGCUAUGCUGCUUCCCAtt (SEQ ID NO:40)
 Antisense strand siRNA: UGGGAAGCAGCAUAGCUU (SEQ ID NO:41)
- Target sequence 70: AAATCACATCCGCCCCAACTT (SEQ ID NO:42)
 Position in gene sequence: 1726
 GC content: 47.6%
 Sense strand siRNA: AUCACAUCCGCCCCAACUUtt (SEQ ID NO:43)
- 40 Antisense strand siRNA: AAGUUGGGGCGGAUGUGAUtt (SEQ ID NO:44)

These siRNAs will be tested in vitro and in vivo to ascertain how well they modulate SOST RNA levels.

References

- Abe, E et al. 2000 J Bone Min Res 15:663
- Beighton, P et al. 1976 Ann Int Med 84:393
- Beresford & Owen, 1998 Marrow Stromal Cell Culture, Cambridge University
5 Press
- Boden, S et al., 1997 Endocrinology 138: 2820
- Bostrom et al., J. Orthopaed. Res. 13, 357, 1995.
- Brunkow, M et al. 2001 Amer J Hum Genet 68: 577-589.
- Budd, RC 2002 J Clin Invest 109: 437.
- 10 Cooper, MS et al. J Endocrinology 1999, 163: 159.
- Fang et al., Proc. Natl. Acad. Sci. USA. 93, 5753, 1996.
- Gazzero, E et al. 1998 J Clin Invest 102:2106.
- Gazit, D et al, 1999 Mol Endocrinol 7: 189.
- Gitelman, SE et al., 1995 Cell Growth & Differ 6: 827.
- 15 Hofmann et al. FEBS Letters 1998, 441: 441.
- Lian, JB et al. 1999 in Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism, 4th edition, MJ Favus (ed), Lippincott, Philadelphia, pg. 14.
- Lu, TT et al. J Biol Chem 2001, 276: 37735.
- 20 Katagiri, T et al., 1990 Biochem Biophys Res Commun 172: 295.
- Manolagas, SC 2000 Endocrine Rev 21: 115.
- Martinez, LA et al. 2002 PNAS 99: 14849.
- Miura et al. J Biol Chem 2001, 276: 47371.
- Nakase et al., J. Bone Miner. Res. 9, 651, 1994.
- 25 Nifuji, A & Noda, M 1999 J Bone Min Res 14: 2057.
- Oreffo, ROC et al. 1999 J Cell Biochem 75: 382.
- Pelletier, JP et al. Am J Ther 1996, 3:115.
- Pereira, R et al. 2000 Endocrinology 141: 4558.
- Pittenger, MF et al. 1999 Science 284: 143.
- 30 Pockwinse, S et al., 1992 J Cell Biochem 49:310.
- Scherr, M et al. Curr Med Chem 2003 10:245.

- Song, CZ et al. Proc Natl Acad Sci USA 1999, 96:11776.
- Suzawa, M et al., 1999 Endocrinology 140: 2125.
- Takiguchi, T et al. J Periodontal Res 1999, 34:431.
- Tanaka, H & Makino, I. BBRC 1992, 188: 942.
- 5 Thornberry, NA & Lazebnik, Y 1998 Science 281: 1312.
- Vaughan, ATM et al. 2002 Apoptosis 7: 173
- Wang, Trends Biotechnol. 11, 379, 1993.
- Weston, A. et al. 2000 J Cell Biol 148: 679.
- Winkler, DG et al. 2000 J Bone Min Res 16 (Suppl.1): S322.
- 10 Yanagi, Y et al., 1999 J Biol Chem 7: 12971.
- Yamaguchi, A et al. 2000 Endocrine Rev 21: 393

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

I

SEQUENCE LISTING

<110> Celltech R&D, Inc.
 5 Sutherland, May S. Kung
 Geoghegan, James Charles
 Yu, Changpu
 Latham, John
 Celltech R&D, Inc.
 10
 <120> Methods to Increase or Decrease Bone Density

 <130> 1427.005WO1

 15<150> US 60/361,258
 <151> 2002-03-01

 <150> US 60/406,171
 <151> 2002-08-27
 20
 <150> US Ser. No. Unknown
 <151> 2003-02-13

 <160> 44
 25
 <170> FastSEQ for Windows Version 4.0

 <210> 1
 <211> 21501
 30<212> DNA
 <213> Homo sapiens

 <400> 1
 acccccatcc ccagctccca gggcctctgc cacgcgggct tctttttttt tttgagacag 60
 35agtctcgctc tgtcaccag gctggagcac aatctcagca caatctcacc tcaccgcaac 120
 ctctgcctcc tgggttcaag caattctctg gcctcagcct cctgagtagc tgggattata 180
 ggctcgcgcc accacacca gctatTTTTT tgtatcttta gtaaagacgg gatttcacta 240
 tgttggccag tctggtctcg aactcctgac cttgtgatcc agttgccttg gcctcccaaa 300
 gtgctgggat tacaggcgtg agccactgtg cccagcccct tgcaggcttc tgacagcaca 360
 40gatggTTTTT cccgctttcc tactctgttg gctcatccac ctcatatgtg tgtgctgcct 420
 ccgtgttgct tctgctaagt ggcagtctgg tcattctttg tatttgtcct ttctactgtg 480
 gatgggcctt tgggggtttc taactgtggg ccatgatgtg cagtgttgcc tgtgaacatt 540
 ctggcccatg ttcgtgaatg tgaatgcgta tctctatttg acctgggcct gggagtggga 600
 ttgctgggcc agaggggatg cacgtgttca gcttgaggag ataccgccgt gccgcacagc 660

	tttccaaagt	ggttgtgcca	attcacgcct	cacctcccc	agccacgcag	aagagttcca	720
	gttgcccat	atcttcacca	gcactttggg	ttttccatcc	ttttcatttt	agccattctg	780
	acaggcacac	agtggcgtga	cattgtgctt	ttaagtggca	cttcacggaa	gagtcttgac	840
	caacgccatg	gtgtgatggg	acgccccag	ctcaccacgt	ctccccgtt	tcttacaggc	900
5	cggtgtgctcg	tccactgccg	ggaaggttat	agccgctccc	caacgctagt	tatcgcctac	960
	ctcatgatgc	ggcagaagat	ggacgtcaag	tctgccctga	gcacgtgag	gcagaaccgt	1020
	gagatcggcc	ccaacgatgg	cttcctggcc	cagctctgcc	agctcaatga	cagactagcc	1080
	aaggaggga	agttgaaacc	ctagggcacc	cccaccgcct	ctgctcgaga	ggtccgtggg	1140
	ggaggccgtg	ggcaaagggtg	tcccgagctg	ccatgttttag	gaaacacact	gtaccctgct	1200
10	cccagcatca	caaggcactt	gtctacaagt	gtgtcccaac	acagtcctgg	gccactttcc	1260
	ccaccctggg	gagcacataa	agaagcttgc	caaggggggc	gtccttgctc	cccagttgtc	1320
	ctgtttctgt	aacttatgat	gtcttttccc	tgagatgggg	gctcagaggg	ggaaggcctg	1380
	tggcctgcat	gcttcccgat	ggcccacggc	aggaggtgtg	tggaaagtga	aggcctaaga	1440
	tgctcacaga	ggtccctcat	gacctccctt	ccccaaactcc	cgaatcctct	cttgagtgtg	1500
15	gacctcaaca	ccttgagccc	tagtaaagga	actatgcaaa	tgcaggccac	tctccccacc	1560
	acgtctgtgc	cccgcactgt	ccccacagcc	ttccacaccc	tgtgcatagg	cagccctctc	1620
	acgtcttgag	gtccgaagct	gggggtggggg	tgtccgtcag	ttattagtg	atggagattc	1680
	ccacagcaag	gctgcatttg	aatgatttcc	ttaggatgaa	tggtccttac	acaaagaggc	1740
	cttgtgggca	aacctggaga	accctcctaa	atccatagag	ttttcaaaat	gtgaatcttt	1800
20	ggaagccttg	agttcagaat	ctgctgctct	ggaatatattc	ccttcgatct	tatctcagtc	1860
	acttcgtttt	tgagaagagt	gatgccttgg	gcacgtcttt	ttttttttct	tttttagaaa	1920
	acagggagtt	gaagtccaac	ctattttaaaa	accccaccat	ttggagaatt	acaagggttt	1980
	tgtcctgaat	tgtagtgttg	gcaagcccaa	gccactcgtg	ctaactgctt	tttgtctcgg	2040
	ttgctattcc	aagaacagaa	ggaggaagtt	ggccaattac	agcgtgtgtg	catggatgtg	2100
25	tgtggggggc	gtgcctctca	gaaacgcggc	cagaagacaa	gcagggaagt	gaaagggtccc	2160
	aggcacacac	cctgcccatt	gcagggtggc	cttacagctc	tctggtgcca	gcacgggatc	2220
	cctgaagtga	ctcagccagg	cagacatgag	acatggcgga	gtgtccaaat	ggatccttta	2280
	ttggtggtag	agcaaaaaaa	cccaaacacg	ataaaccttt	caaaagactt	tctaaggatg	2340
	atattggaat	gcaccagccc	tcacatgtgt	atgcacattt	gccagaatat	aagagttttg	2400
30	tttttaaatac	agtcttggtt	ggattttacg	ttattgttat	tatggaaagt	gattgtgatg	2460
	ctatttatct	tcagggtcac	tctgggcaaa	gagaagggtcc	tcagccatgc	ccccagcacc	2520
	ttgcacatag	gtgtctgata	aaagtttaag	aaattaaaca	ctttttgagc	accaaata	2580
	tatagggcat	tgttctggtg	ggtgtgtcac	gctcccagaa	gactgaattt	atggtaggat	2640
	cactcgcaag	gccttgtgaa	ggagtcttac	ctaaaacaaa	agaaatatca	gggacttttg	2700
35	ttgactatth	acaactcagt	tttacattta	aattcaggca	gtgttaatat	gccaaggtag	2760
	ggaatgtgcc	tttttcagag	ttggccagga	gctcctggct	gggacacgga	gaggcaggtg	2820
	tggcgtaagg	cctcactccc	ggctgtgaag	gtctctgac	acacagaagc	agccctgccc	2880
	agcctggtca	tttgctgtcc	gcttttctct	gtgaccacag	cagccctgaa	caaccagtat	2940
	gtgtcttctt	ctccagatag	tgaaaaagg	gtccagataa	accacactaa	gtgaaatggc	3000
40	catcctctaa	actgggtacc	tcactgcaca	gcttctaggt	agccttccaa	cttaatactaa	3060
	cttgagcctc	acagtaaccc	tgtaaagtta	gtagagcttg	ttcttgtatt	gtgacctttt	3120
	ttaaaaaaaaa	ggaactgagg	ttcagaatga	ttaagggcct	ggccccaggg	gttgtccagc	3180

tccataaggt	ggagctgggc	aagatttttg	gtttgctgct	ccctgaagct	ggattctttc	3240
atacgatact	ctttctcaag	aagggggctc	cctgggatct	ccaggtgtac	tgcacttacc	3300
ctcaatccag	ccccggagaa	gcaagtgaag	aggggtgggtc	cctcataggc	tagaatgtgc	3360
agctctttct	ccaggtggga	tgtagcacc	caaagtagag	ctttctgctc	tgctcctgga	3420
5aaaggctagg	gagctggggc	tggggctccc	ctcccatgac	caggcagtgg	tcaccccatg	3480
ggacaggcac	agctacttac	gcgaacacag	cagggttggtg	tggctggcta	actaggacct	3540
ctcgaaagtc	tctgtggggg	catgagggag	aaaaggccat	tgggagaatt	actgccttta	3600
ctttgggact	acttttatgc	tgataacttg	ggatttcttg	atagtccttc	accctgaaa	3660
ccccgtat	acttaacaag	atttagctct	tagttcttca	agtaaaatta	aagtctcttg	3720
10tgtaagagcc	aacacatgcc	cagctgcgga	tgggagctgt	tcctggacag	ccttctactg	3780
cctgggaagt	gatggaacag	gaactcaggg	tgcccttacc	ccctccccag	acctgttccc	3840
tttctttgac	tgacagagca	ccatccaggc	aaaattagag	cgccaaatgg	ttttcttctc	3900
aatcttaaag	cagtatacct	ttccacaggc	tcgtctgtgt	ccctgccact	ctgagttatc	3960
cagaaaccac	cacctacaaa	tgaggggact	catctagaag	acctctaagg	tccccttttg	4020
15gctctgaggg	gtctctaata	atccccactt	ggaattcagc	accgcaagga	aattatgggt	4080
atgtgagcca	taatatgatg	gccagcaggt	ggcgctgcct	tccacccatg	gtgatggatg	4140
gtttggaaag	ggaatgttgg	tgcccttttg	gccacaagtt	aagatgctac	tgttttaaaag	4200
gaaaaaaaaa	aaaaaaagta	ctgatcttca	atatgaagac	atgagctttt	ctcgcaggaa	4260
attttctttt	tcacagaact	gggtgtcagga	atcactgaag	ggctaaccgt	gatagtcctt	4320
20gcaagtaagt	caagggttta	tcctgattgg	aaatagaaga	catttccggt	tgagagaaca	4380
gattcgttgg	aagcttaact	tttgttgcct	cttaacgcca	ccaaatttta	gggtaatttg	4440
attatgaaag	agtgaatttt	tctggacaga	aaaggagag	ctaccaaatt	gtttttttct	4500
ttttaaaagg	aagtttaattg	tccgttgtat	cacaaatcag	tgtaaaaaca	ccagaacttt	4560
agccaaaata	aatgtcttac	attacaaagg	tattgttttt	ttgtccttct	tatcacagtt	4620
25ggatattctt	tacgttttta	tgcttagctt	tttttgtttg	ggcttagctt	ttatatattgc	4680
gatttctaac	tttttaaaaa	taatcatcta	aaacatagcc	gttctgagct	agttataggt	4740
tccatgatgg	cactgtttat	gcagtagata	ttgatgaaca	cctttccagt	accagaaatg	4800
ttctggtagg	aatattctta	ggtagaatgg	ccataatggg	cctgaatgaa	ggaggtaagg	4860
tttgtgcttc	taagaaagca	ggggactagg	gtgttcatct	caaggtagcc	tcacgtgatg	4920
30actgggtgac	attttgagat	ttgagtgatc	ctgcaaataga	acagtcctta	aacatatccc	4980
cttccccccag	atgccttaaa	ttccactatt	gggtgctatgt	tcctttgaat	aaaaaacctt	5040
agagaaagta	gggtgcctttt	cctcccagac	agaagtctgg	gcagagaaag	cctctcgagg	5100
ttcccatctc	ctctatcaca	tgaagcaatt	ggactcgctt	cctatttgct	aagactctga	5160
caatgtaatc	tttatggttc	tacacaaaat	cttaaggcag	agcactaaaa	tataaaacag	5220
35gtaaaagt	tttaagtataa	tttttaaaaa	atttttcatt	aaaaataata	ctatatgtac	5280
atagtaatac	acttaaacaa	tgcaaaaggc	ttatacggaa	aagtgaagtca	cttgctcctt	5340
caatctatct	cccataagc	gtcatatcct	taaaggaaaa	tgtcaagcat	atataagcac	5400
acatatgtgc	atacgtgtgt	atatacagag	atatctctgc	tttggtgttt	gttttgctta	5460
gcttatatta	tggttattgt	ttttattgct	ctacccctt	catcttaaca	gctacatagc	5520
40attccattgt	gtatggatgt	gtaaataaag	acaaggatgt	ttggacatcc	ctcaatgcag	5580
ggtagtgcct	aaaagcagaa	gtttgccctc	ctaattccaga	tatttccctt	ttaccaccta	5640
caagtgggag	caaaatctgt	tttaaaatgc	agccaataag	gagaatcctc	agtttccatt	5700

aattagagat	aggggcctcc	tccagaacca	agtagagaat	gaaaccggat	gacctttgag	5760
agtccttcca	gatcaaagat	tctatgatcc	caaggagtca	aaggagcata	atatactttc	5820
caaattttggc	ctgtcattca	aagtctggtc	caaatgccac	atcttaccat	aaggatccca	5880
gacctcctccc	aggggaaaga	gattgttctc	tcctttgaag	gcccataaga	aaattgtcct	5940
5tttgaaaaaa	tgcattctgg	gaggatgtgg	aacaacagga	actctttggt	tcagtgggga	6000
atgcaaaatg	gtacagccac	tttgaggagc	agtttgagc	ttagtttctt	gcaaaactac	6060
acatactttt	attgcacaat	ccagcagtca	tgctccttgg	tatttattca	aatgatttga	6120
aaacttaacg	ttcgtataaa	aacctgcaca	cagctgggca	tggtggctca	tgctgtaat	6180
cccagcactt	tgggaggctg	aggtgggcag	atcacttgag	gttaggagtt	tgagaccata	6240
10ctggccaaca	tggggaaacc	ccttctctac	taaaaataca	aaaattagcc	aggtgtggtg	6300
gtgctcacct	gtaatcccac	ctactcggga	ggctgagcca	ggagaatcac	ttgaaccacg	6360
gaggcggagg	ttgcagtgag	ctgagatcac	ggcactgcac	cccagcctgg	gagacagagt	6420
aagactccgt	cttaagacaa	aacaaaacaa	aaaacctgta	caaatgtttt	ataccacctc	6480
tattaataat	tgccagagct	tgaaagcaac	caagatgtcc	ttcagtagat	gacagtaggt	6540
15gactagataa	accgtgatac	atcaagacga	tggagtatta	ctcagtgtca	aaaaaaaaaa	6600
taagctgttc	atgccatgaa	aagacatgga	ggaaccccaa	atgaatatta	ctaactgaaa	6660
gaagccaatc	tgaaaagact	atgtatgggt	atattccaac	tctatatgac	atcctggaaa	6720
aggcaaaaat	aaaggtacag	taaaaagatc	agtggttgcc	aggggttggg	gaaagtcggg	6780
gagggatgaa	taggcagagc	agatagtttt	tttgacaaat	gaaaataactg	cccatctgta	6840
20tgatactata	atggtagatg	tatgtcatta	tacatttgtc	caaagccata	ggatgtacaa	6900
cactagtgtc	gtgatgtaaa	ccatggactt	caggtgataa	tgatgtgtca	tgtaggtctc	6960
tcaatcatag	caaatgcacc	actctgggtg	cgggatgttg	ataacggggg	cagctgtaca	7020
tgtgtgggca	caggacataa	atgagaaatc	tctgtaccat	ccactcagtt	ttgctgtgaa	7080
cctaaaagtg	ctccagaaaa	aaaataaagt	tttttttaga	aaaaagcatc	tgaacctttt	7140
25tttcagtggc	attgatcacc	ttccatctgg	tagtggagtt	gtttggatgc	ggaaattatt	7200
tcttccatgg	tatgcgagct	cctggagagg	gagtgcgtgt	cccccttgct	ttattaaacg	7260
tttggcgagt	gaacatcaga	agaagcaact	cgacagaaac	aaaagctggg	aaataatcac	7320
atggaaaagc	agcttttctg	gaaattaagt	ttcaactgtt	aaattcaatg	tttaaaaaaa	7380
tcggctgggt	gcgggtggctc	ccgcctgcaa	tcccagcact	ttatgggatt	gtgggaacac	7440
30ctgatgtcag	gagttcaaga	ccagcctggc	cgacatggcg	aaaccccatc	tctacaaaac	7500
aaacaaaaaa	aatgcaaaa	attagctggg	tgtggtggcg	ggagtctgta	gtcctagcta	7560
ctcgggaggc	tgaggcagga	gaactgcttg	aaccaggag	gtggagggtta	tatgagccaa	7620
gatcacacca	ctgcactccc	gcctgggtga	cagagcgaga	ctctgtctca	aaaaaaaaaa	7680
aaaaaatcaa	catcccaaac	ccagtgttga	cagtcttatt	tatgggttga	actgcattcc	7740
35tcccaaatgc	atatattgaa	gtcctaagtc	acagtacctg	agaatgtgac	cttattttgga	7800
aatagggcta	ttacagatgt	aactcggtta	gatgagggtca	ttaggctggg	cctaattctaa	7860
tatgcctgac	ctccttagaa	aaaagggaaa	attcgaacac	agagagaaca	ccatgtgaag	7920
atgaaggcag	ggattggggg	tgacacttct	gcaagccagt	gaatgccaca	gcttgtcagc	7980
agccagcaga	aggcaggtaa	aaggccttcc	tactgtcct	cgaagggaacc	aactctgcca	8040
40atgtcttgaa	cttgcaactcc	taacctccag	ctctgtgaga	cgacttctgt	tgtttatgcc	8100
acccaatttg	tgggtacttag	ttacggcagc	ctcatcaaag	taccacctat	gggagcctct	8160
atthttgcagg	tgagggcggg	gactgggctg	agthtttctgg	aaaacagccc	tgcaataccc	8220

	tcacagacc	accaaactct	tcacactccc	tcagacacag	cattcacttc	cagaaataac	8280
	tctaaagttt	tgttttgttt	ttttaaactt	tgtggaatac	tactcagcca	aaaaaaaaaa	8340
	aaaagaaaag	aaaaaggaac	atggttactga	tatgtacaac	ttggataatg	gaaataaatgc	8400
	tgagtgaaaa	aaaaatcccc	aaaggctaca	tactaattga	ttccatttat	ataacctttt	8460
5	ttttttttttt	ttgagacagt	ctcactctgt	cacccaggct	ggagcacagt	ggcgcgatct	8520
	cagctcactg	caacttccgc	ctcctgagtt	caagcgattc	tcttgcctta	gcttcccaag	8580
	tagctgggat	tacaggtgcg	tgccaccatg	cccagctaata	ttttgtattt	ttagtagaga	8640
	cagggtttcg	ccatgttggc	caggctggtc	tcgaactcct	gacctcaagt	gatctgcctg	8700
	ccttggcttc	ccaaagtgct	gggattacat	gagtgaacca	ccgcacctgg	tcgcatttat	8760
10	gtaacaattt	tgaagtgaag	aaaaaaatga	cagaaatgga	gattagatga	gtagttgcca	8820
	gggggttagtt	gtgggaggga	gggaaaagga	gggaaggagg	tgggcaacag	gagaaagact	8880
	tgtggtcaca	gagctgtgct	ttatcttgac	tgtggtggat	cccaaattt	acccgtgaca	8940
	agattgcata	gaactaagta	tacacacacg	tgaatgcgtg	tgcacgcaca	cagtagaggt	9000
	taagccacgg	gagatgtggg	taagattggt	aaattgtgtc	aatatcaata	tcctagttgt	9060
15	gatattgtcc	tatagtttcg	caaggtgtta	ttgttgtggg	aaactggata	aaggatacac	9120
	ggagtctgca	ttttcttttt	tttattttta	ttttttgaga	cgggggtctca	ctctgtcaac	9180
	caggctggag	tgcagtggcc	caagtatggc	tactgcagc	ctcgacctca	acctcaagtg	9240
	aacctccac	ctcagcctcc	caagtagcta	agaccacagg	cgtgcgaccc	catgcccagc	9300
	taatttttaa	attttttgta	gagactaggc	ctcaccatgt	tgcccaggct	tttatttctt	9360
20	ataagtatat	ttaaatttat	aattatatcc	acatttttaa	atttttaattt	aaaaaattac	9420
	tctgaggccg	ggcattgtgg	ctcatgcctc	taatccccag	caccttggga	ggccgagttg	9480
	ggcagatcac	ccgaggtcag	aagttcgaga	ccagcctgag	taacatggag	aaacccccgt	9540
	ctctactaaa	aatacaaaat	tggccggggc	tgggtggtgca	ttcctgtaat	cccagctact	9600
	cgggaggctg	aggcaggaga	attgcttgaa	cccaggagggt	ggagggttgca	gtgagccaag	9660
25	attgtgccac	tgcactgcag	cctgggcccac	agagagaatc	tgccaaaaaa	agaaaaaaa	9720
	aaaaatttca	gccgtacaag	gatgttcata	gcaaccctgc	tggaaatagg	aaaaaaaatt	9780
	ggaaataacc	taaactactc	acaataggaa	tcagctaaaa	ccctgggggt	ttaattccag	9840
	ggaatactgt	gaacaatgac	aagtttgtgg	actgagtaaa	aataaacagc	tgtcaatgac	9900
	ttaacattaa	atgaaacagc	agaagatgtc	acagcagggt	ctcgctgagc	cattcagagg	9960
30	ggtgtggatc	atttagaggt	tcaagtccac	tggattcttc	tttttccttt	taatattact	10020
	tcacttccaa	ataaggaaag	gaaaggaaag	gaaatcacgt	ccagtcctga	gacttgccat	10080
	cctgcagtca	cccctccttt	tgtctccagc	aggtggcaga	cgcgttccag	ggatgaatcc	10140
	cactgcctct	gtttaatgca	gacgggtccag	ccgctcccaa	cagcagggtg	ggctataagc	10200
	atccatccta	cctgctcaag	gaacccaggc	atcagaactg	ctctctccca	agtccattgc	10260
35	aagaaggcag	tcgtctgggt	atgagagggt	taacagtcca	cattccagag	caagggaaaa	10320
	ggaggctgga	gggtcataga	caaggggagg	tgggtgcggag	ggccagcttc	tcacaacact	10380
	accggctctg	ctgggagaga	tagatcaccc	ccaacaatgg	ccacagctgt	tttcatctgc	10440
	cctgaaggaa	actgacttag	gaagcaggta	tcagagaggg	cccttcctga	gggggcttct	10500
	gtctggcttg	taaaactgtc	agagcagctg	cattcatgtg	tcggatgatg	gatgatggaa	10560
40	aggacagtcg	gctgcagatg	gacacagcga	cttgcaagtt	gaggcagggt	gcaaaggact	10620
	tgcagaggct	ctgcagggtg	ggcatgctga	ttcattgccc	agttaaaata	ccagaggatc	10680
	tgggcagcct	cttcacagga	gctgcttggt	ctcaaacaat	ctgtcttcaa	tgaaagattc	10740

	ctctggcctt	cctttctctt	cttgccacctc	aggtgtgaat	ccttctcccc	cacgcctcta	10800
	cctgcgcccc	cgcgcgcgc	cccggccctg	tgtggctcat	tatatgcagg	gccaaggcag	10860
	cattttctct	tagcttcttt	gtgaccagtt	ggctctggga	tggcttcatg	gaacacatcc	10920
	tgtgggtgtgc	accaatgaag	ctttccatac	aggactcaaa	actgtttttg	aaaaatgtaa	10980
5	ccagctggaa	gacaagaaaa	taaaatgtca	gcactaaaaa	cgctggctgt	ggcttttgct	11040
	aaggaaagga	atttgggtgtt	gtcttctcac	acacacagac	tggttgggga	aatgactgtc	11100
	ttcagcacat	caccctgcga	gccacagtga	gtgccctggc	tcagaagtgc	ctgtcacagt	11160
	gcacaggatc	cctgaggagc	atgagctggg	atttcctctg	tgtgtccat	cacaggagcc	11220
	tgagtgacca	gcgcatectc	gatttgtaac	cagaatcctg	ccctctctcc	caagcgggca	11280
10	cccttgctct	gaccctctag	ttctctctct	tgccttccag	agaataccaa	gagaggcttt	11340
	cttggttagg	acaatgaatg	ctgagacttg	tggagtggg	accaatggga	tttctttaa	11400
	agcatctttt	tgcctctggc	tgggtctatg	ggggtcaaac	agaaacacct	tgggccatth	11460
	gttgggtggg	tgacaaatga	acttggcctg	agaaatggaa	taggccgggc	tcagccccgc	11520
	gaagcactca	gaactgcaca	ttttctttgt	tgagcgggtc	cacagtthgt	tttgagaatg	11580
15	cccgaggggc	caggagagaca	gacaattaaa	agccggagct	cattttgata	tctgaaaacc	11640
	acagccgcca	gcacgtggga	ggtgccggag	agcaggcttg	ggccttgctt	cacacgcccc	11700
	ctctctctgg	gtcacctggg	agtgccagca	gcaatttgga	agtttgctga	gctagaggag	11760
	aagtcttttg	ggagggtttg	ctctgagcac	acccttttcc	ctccctccgg	ggctgaggga	11820
	aacatgggac	cagccctgcc	ccagcctgtc	ctcattggct	ggcatgaagc	agagaggggc	11880
20	tttaaaaagg	cgaccgtgtc	tcggctggag	accagagcct	gtgctactgg	aagggtggcgt	11940
	gccctcctct	ggctgggtacc	atgcagctcc	cactggccct	gtgtctcgtc	tgcctgctgg	12000
	tacacacagc	cttccgtgta	gtggagggcc	aggggtggca	ggcgttcaag	aatgatgcca	12060
	cggaaatcat	ccccgagctc	ggagagtacc	ccgagcctcc	accggagctg	gagaacaaca	12120
	agaccatgaa	ccgggcggag	aacggagggc	ggcctcccca	ccaccctttt	gagaccaaa	12180
25	gtatgggggtg	gaggagagaa	ttcttagtaa	aagatcctgg	ggaggthtta	gaaacttctc	12240
	tttgggaggc	ttggaagact	ggggtagacc	cagtgaagat	tgttggcctc	tgccagcact	12300
	ggtcgaggaa	cagtcttgcc	tggagggtgg	ggaagaatgg	ctcgctgggtg	cagccttcaa	12360
	attcagggtgc	agaggcatga	ggcaacagac	gctgggtgaga	gccaggggca	gggaggagcg	12420
	tgggggtggtg	aggggtatgg	atcaggggcat	cagaacaggc	tcaggggctc	agaaaagaaa	12480
30	aggthttcaa	gaatctcctc	ctgggaatat	aggagccacg	tccagctgct	ggtaccactg	12540
	ggaagggaac	aaggtaaggg	agcctcccat	ccacagaaca	gcacctgtgg	ggcaccggac	12600
	actctatgct	ggtgggtggct	gtccccacca	cacagacca	catcatggaa	tccccaggag	12660
	gtgaaccccc	agctcgaagg	ggaagaaaca	ggttccaggc	actcagtaac	ttggtagtga	12720
	gaagagctga	ggtgtgaacc	tggthtgatc	caactgcaag	atagccctgg	tgtgtggggg	12780
35	ggtgtggggg	acagatctcc	acaaagcagt	ggggagggaag	gccagagagg	caccctgca	12840
	gtgtgcattg	cccatggcct	gccaggggag	ctggcacttg	aaggaaatgg	agthtttcggc	12900
	acagthtttag	cccctgacat	gggtgcagct	gagtccaggc	cctggagggg	agagcagcat	12960
	cctctgtgca	ggagtaggga	catctgtcct	cagcagccac	cccagtccca	accttgccctc	13020
	attccagggg	agggagaagg	aagagggaacc	ctgggttcct	ggtcaggcct	gcacagagaa	13080
40	gccaggtga	cagtgtgcat	ctggctctat	aattggcagg	aatcctgagg	ccatgggggc	13140
	gtctgaaatg	acacttcaga	ctaagagctt	ccctgtcctc	tggccattat	ccaggtggca	13200
	gagaagtcca	ctgcccaggc	tcctggaccc	cagccctccc	cgcctcaca	cctgttggga	13260

ctatggggtg	ctaaaaaggg	caactgcatg	ggaggccagc	caggaccctc	cgtcttcaaa	13320
atggaggaca	agggcgccctc	ccccacagc	tccccttcta	ggcaagggtca	gctgggctcc	13380
agcgactgcc	tgaagggctg	taaggaaccc	aaacacaaaa	tgtccacctt	gctggactcc	13440
cacgagaggc	cacagcccct	gaggaagcca	catgctcaaa	acaaagtcat	gatctgcaga	13500
5ggaagtgcct	ggcctagggg	cgctattctc	gaaaagccgc	aaaatgcccc	cttccctggg	13560
caaatgcccc	cctgaccaca	cacacattcc	agccctgcag	aggtgaggat	gcaaaccagc	13620
ccacagacca	gaaagcagcc	ccagacgatg	gcagtggcca	catctcccct	gctgtgcttg	13680
ctcttcagag	tgggggtggg	gggtggcctt	ctctgtcccc	tctctggttt	ggtcttaaga	13740
ctatttttca	ttctttcttg	tcacattgga	actatcccca	tgaaaccttt	gggggtggac	13800
10tggtactcac	acgacgacca	gctatttaaa	aagctcccac	ccatctaagt	ccaccatagg	13860
agacatggtc	aaggtgtgtg	caggggatca	ggccaggcct	cggagcccaa	tctctgcctg	13920
cccagggagt	atcaccatga	ggcgcccatt	cagataacac	agaacaagaa	atgtgcccag	13980
cagagagcca	ggtcaatggt	tgtggcagct	gaacctgtag	gttttgggtc	agagctcagg	14040
gcccctatgg	taggaaagta	acgacagtaa	aaagcagccc	tcagctccat	ccccagccc	14100
15agcctcccat	ggatgctcga	acgcagagcc	tccactcttg	ccggagccaa	aaggtgctgg	14160
gaccccaggg	aagtggagtc	cggagatgca	gcccagcctt	ttgggcaagt	tcttttctct	14220
ggctgggcct	cagtattctc	attgataatg	aggggggttg	acacactgcc	tttgattcct	14280
ttcaagtcta	atgaattcct	gtcctgatca	cctccccttc	agtccctcgc	ctccacagca	14340
gctgccctga	tttattacct	tcaattaacc	tctactcctt	tctccatccc	ctgtccaccc	14400
20ctcccaagtg	gctggaaaag	gaatttgga	gaagccagag	ccaggcagaa	ggtgtgctga	14460
gtacttacct	tgcccaggcc	agggaccctg	cggcacaagt	gtggcttaaa	tcataagaag	14520
accccagaag	agaaatgata	ataataatac	ataacagccg	acgctttcag	ctatatgtgc	14580
caaatggtat	tttctgcatt	gcgtgtgtaa	tggattaact	cgcaatgctt	ggggcgggccc	14640
attttgacga	caggaagaag	agagagggtta	aggaacttgc	ccaagatgac	acctgcagtg	14700
25agcgatggag	ccctggtggt	tgaaccccag	cagtcatttg	gctccgaggg	gacaggggtgc	14760
gcaggagagc	tttccaccag	ctctagagca	tctgggacct	tcctgcaata	gatgttcagg	14820
ggcaaaaagcc	tctggagaca	ggcttgga	aagcagggct	gggggtggaga	gagacggggcc	14880
ggtccagggc	aggggtggcc	aggcgggcgg	ccaccctcac	gcgcgcctct	ctccacagac	14940
gtgtccgagt	acagctgccg	cgagctgcac	ttcaccgcgt	acgtgaccga	tgggcccgtgc	15000
30cgcagcgcca	agccggtcac	cgagctgggt	tgctccggcc	agtgcggccc	ggcgcgcctg	15060
ctgcccacacg	ccatcgggccg	cggcaagtgg	tggcgacctt	gtgggcccga	cttccgctgc	15120
atccccgacc	gctaccgcgc	gcagcgcgtg	cagctgctgt	gtcccgggtg	tgaggcgccg	15180
cgcgcgcgca	aggtgcgcct	gggtggcctc	tgcaagtgca	agcgcctcac	ccgcttccac	15240
aaccagtcgg	agctcaagga	cttcgggacc	gaggccgctc	ggccgcagaa	gggccggaag	15300
35ccgcggcccc	gcgcccggag	cgccaaagcc	aaccaggccg	agctggagaa	cgcctactag	15360
agcccgcccc	cgcccctccc	caccggcggg	cgccccggcc	ctgaaccgcg	gccccacatt	15420
tctgtcctct	gcgcgtgggt	tgattgttta	tatttctattg	taaatgcctg	caaccagggg	15480
cagggggctg	agaccttcca	ggccctgagg	aatcccgggc	gccggcaagg	ccccctcag	15540
cccgccagct	gaggggtccc	acggggcagg	ggagggaatt	gagagtcaca	gacactgagc	15600
40cacgcagccc	cgctctggg	gccgcctacc	tttgctggtc	ccacttcaga	ggaggcagaa	15660
atggaagcat	tttcaccgcc	ctgggggtttt	aaggagcg	tgtgggagtg	ggaaagtcca	15720
gggactgggt	aagaaagttg	gataagattc	ccccttgcac	ctcgctgccc	atcagaaagc	15780

ctgaggcgtg	cccagagcac	aagactgggg	gcaactgtag	atgtgggttc	tagtcctggc	15840
tctgccacta	acttgctgtg	taaccttgaa	ctacacaatt	ctccttcggg	acctcaattt	15900
ccactttgta	aaatgagggg	ggagggtggga	ataggatctc	gaggagacta	ttggcatatg	15960
attccaagga	ctccagtgcc	ttttgaatgg	gcagaggtga	gagagagaga	gagaaagaga	16020
5gagaatgaat	gcagttgcat	tgattcagtg	ccaagggtcac	ttccagaatt	cagagttgtg	16080
atgctctctt	ctgacagcca	aagatgaaaa	acaaacagaa	aaaaaaaaagt	aaagagtcta	16140
tttatggctg	acataatttac	ggctgacaaa	ctcctggaag	aagctatgct	gcttcccagc	16200
ctggcttccc	cggatgtttg	gctacctcca	cccctccatc	tcaaagaaat	aacatcatcc	16260
attggggtag	aaaaggagag	ggtccgaggg	tggtgggagg	gatagaaatc	acatccgccc	16320
10caacttccca	aagagcagca	tccctcccc	gacccatagc	catgttttaa	agtcaccttc	16380
cgaagagaag	tgaaagggtc	aaggacactg	gccttgacag	cccgaggagg	cagccatcac	16440
aaactcacag	accagcacat	cccttttgag	acaccgcctt	ctgcccacca	ctcacggaca	16500
catttctgcc	tagaaaacag	cttcttactg	ctcttacatg	tgatggcata	tcttacacta	16560
aaagaatatt	attgggggaa	aaactacaag	tgctgtacat	atgctgagaa	actgcagagc	16620
15ataatagctg	ccacccaaaa	atctttttga	aatcatttc	cagacaacct	cttactttct	16680
gtgtagtttt	taattgttaa	aaaaaaaaag	ttttaaacag	aagcacatga	catatgaaag	16740
cctgcaggac	tggtcgtttt	tttggaatt	cttccacgtg	ggacttgctc	acaagaatga	16800
aagtagtggt	ttttaagag	ttaagttaca	tatttatatt	ctcacttaag	ttatttatgc	16860
aaaagttttt	cttgtagaga	atgacaatgt	taatattgct	ttatgaatta	acagtctggt	16920
20cttccagagt	ccagagacat	tgttaataaa	gacaatgaat	catgaccgaa	aggatgtggt	16980
ctcattttgt	caaccacaca	tgacgtcatt	tctgtcaaag	ttgacaccct	tctcttggtc	17040
actagagctc	caaccttgga	cacacctttg	actgctctct	ggtggccctt	gtggcaatta	17100
tgtcttcctt	tgaaaagtca	tgtttatccc	ttcctttcca	aaccagacc	gcatttcttc	17160
accagggca	tggttaataac	ctcagccttg	tatcctttta	gcagcctccc	ctccatgctg	17220
25gcttccaaaa	tgctgttctc	attgtatcac	tccctgctc	aaaagccttc	catagctccc	17280
ccttgcccag	gatcaagtgc	agtttcccta	tctgacatgg	gaggccttct	ctgcttgact	17340
cccacctccc	actccaccaa	gcttctact	gactccaaat	ggtcatgcag	atccctgctt	17400
ccttagtttg	ccatccacac	ttagcacccc	caataactaa	tcctctttct	ttaggattca	17460
cattacttgt	catctcttcc	cctaaccttc	cagagatggt	ccaatctccc	atgatccctc	17520
30tctcctctga	ggttccagcc	ccttttgtct	acaccactac	tttggttcct	aattctgttt	17580
tccatttgac	agtcattcat	ggaggaccag	cctggccaag	tcctgcttag	tactggcata	17640
gacaacacaa	agccaagtac	aattcaggac	cagctcacag	gaaacttcat	cttcttcgaa	17700
gtgtggattt	gatgcctcct	gggtagaaat	gtaggatctt	caaaagtggg	ccagcctcct	17760
gcacttctct	caaagtctcg	cctccccaag	gtgtcttaat	agtgtggat	gctagctgag	17820
35ttagcatctt	cagatgaaga	gtaaccctaa	agttactctt	cagttgccct	aagggtgggat	17880
ggtcaactgg	aaagctttta	attaagtcca	gcctaccttg	ggggaacca	ccccacaaa	17940
gaaagctgag	gtccctcctg	atgacttgct	agtttaacta	ccaataaccc	acttgaatta	18000
atcatcatca	tcaagtcttt	gatagggtgtg	agtgggtatc	agtggccggg	cccttcctgg	18060
ggctccagcc	cccgaggagg	cctcagtgag	cccctgcaga	aaatccatgc	atcatgagtg	18120
40tctcagggcc	cagaatatga	gagcaggtag	gaaacagaga	catcttccat	ccctgagagg	18180
cagtgcgggc	cagtgggtgg	ggacacgggc	tctgggtcag	gtttgtgttg	tttgtttggt	18240
tgttttgaga	cagagtctcg	ctctattgcc	caggctggag	tgcagtgtca	caatctcggc	18300

	ttactgcaac	ttctgccttc	ccggattcaa	gtgattctcc	tgcctcagcc	tccagagtag	18360
	ctgggattac	aggtgcgtgc	caccacgcct	ggctaatttt	tgtattttttg	atagagacgg	18420
	ggtttcacca	tgttgccag	gctagtctcg	aactcttgac	ctcaagtgat	ctgcctgcct	18480
	cggcctccca	aagtgctggg	attacaggcg	tgagccacca	cacccagccc	caggttggtg	18540
5	tttgaatctg	aggagactga	agcaccaagg	ggttaaagt	tttgcccaca	gccatacttg	18600
	ggctcagttc	cttgccctac	ccctcacttg	agctgcttag	aacctgggtg	gcacatgggc	18660
	aataaccagg	tcacactggt	ttgtaccaag	tgttatggga	atccaagata	ggagtaattt	18720
	gctctgtgga	ggggatgagg	gatagtgggt	agggaaagct	tcacaaagt	ggtgttgctt	18780
	agagattttc	cagggtggaga	agggggcttc	taggcagaag	gcatagccca	agcaaagact	18840
10	gcaagtgcac	ggctgctcat	gggtagaaga	gaatccacca	ttcctcaaca	tgtaccgagt	18900
	ccttgccatg	tgcaaggcaa	catgggggta	ccaggaattc	caagcaatgt	ccaaacctag	18960
	ggtctgcttt	ctgggacctg	aagatacagg	atggatcagc	ccaggctgca	atcccattac	19020
	cacgaggggg	aaaaaacct	gaaggctaaa	ttgtaggctg	ggtagaggt	tatttatgga	19080
	aagttatatt	ctacctacat	ggggtctata	agcctggcgc	caatcagaaa	aggaacaaac	19140
15	aacagaccta	gctgggaggg	gcagcatttt	gttgtagggg	gcggggcaca	tgttctgggg	19200
	gtacagccag	actcagggt	tgtattaata	gtctgagagt	aagacagaca	gagggataga	19260
	aggaaatagg	tccctttctc	tctctctctc	tctctctctc	tactctctc	tctctctcac	19320
	acacacacac	agacacacac	acacgctctg	taggggtcta	cttatgctcc	aagtacaaat	19380
	caggccacat	ttacacaagg	aggtaaagga	aaagaacgtt	ggaggagcca	caggacccca	19440
20	aaattccctg	tttcccttga	atcaggcagg	acttacgcag	ctgggagggg	ggagagcctg	19500
	cagaagccac	ctgcgagtaa	gccaagttca	gagtcacaga	caccaaagc	tggtgccatg	19560
	tcccacaccc	gccacctcc	cacctgctcc	ttgacacagc	cctgtgctcc	acaaccggc	19620
	tcccagatca	ttgattatag	ctctggggcc	tgaccgctcc	ttcctgccac	atccccaccc	19680
	cattcttgga	acctgcctc	tgtcttctcc	cttgtccaag	ggcaggcaag	ggctcagcta	19740
25	ttgggcagct	ttgaccaaca	gctgaggctc	cttttgtggc	tggagatgca	ggaggcaggg	19800
	gaatattcct	cttagtcaat	gcgaccatgt	gcctgggttg	cccagggtgg	tctcgtttac	19860
	acctgtaggc	caagcgtaat	tattaacagc	tcccacttct	actctaaaaa	atgacccaat	19920
	ctgggcagta	aattatatgg	tgcccatgct	attaagagct	gcaacttgct	gggcgtggtg	19980
	gctcacacct	gtaatcccag	tactttggga	cgtcaaggcg	ggtggatcac	ctgagggtcac	20040
30	gagttagaga	ctggcctggc	cagcatggca	aaaccccatc	tttactaaaa	atacaaaaat	20100
	tagcaaggca	tggtggcatg	cacctgtaat	cccagggtact	cgggaggctg	agacaggaga	20160
	atggcttgaa	cccaggaggc	agagggtgca	gtgagccaag	attgtgccac	tgccctccag	20220
	ccctggcaac	agagcaagac	ttcatctcaa	aagaaaaagg	atactgtcaa	tactgcagg	20280
	aagaacctag	gtaatgaatg	aggagaagag	aggggctgag	tcaccatagt	ggcagcaccg	20340
35	actcctgcag	gaaaggcgag	acactgggtc	atgggtactg	aagggtgcc	tgaatgacgt	20400
	tctgcttttag	agaccgaacc	tgagccctga	aagtgcctgc	ctgttcatgg	gtgagagact	20460
	aaattcatca	ttccttggca	ggtactgaat	cctttcttac	ggctgcctc	caatgcccac	20520
	tttccctaca	attgtctggg	gtgcctaagc	ttctgcccac	caagagggcc	agagctggca	20580
	gcgagcagct	gcaggtagga	gagataggta	cccataaggg	aggtgggaaa	gagagatgga	20640
40	aggagagggg	tgagagcac	acacctcccc	tgctgacaa	cttcctgagg	gctgggtcatg	20700
	ccagcagatt	taaggcgag	gcaggggaga	tggggcgga	gaggaagtga	aaaaggagag	20760
	ggtggggatg	gagaggaaga	gagggtgatc	attcattcat	tccattgcta	ctgactggat	20820

10

gccagctgtg	agccaggcac	caccctagct	ctgggcatgt	ggttgtaatc	ttggagcctc	20880
atggagctca	cagggagtgc	tggcaaggag	atggataatg	gacggataac	aaataaacat	20940
ttagtacaat	gtccgggaat	ggaaagttct	cgaaagaaaa	ataaagctgg	tgagcatata	21000
gacagccctg	aaggcggcca	ggccaggcat	ttctgaggag	gtggcatttg	agctaagacc	21060
5aaaatgtggt	gggagaggga	gccacacaag	gatctggggg	tgtgtgcact	gggaggggga	21120
gcagcaagtg	caaagggcct	gaaggatatg	agggggtgga	gaagggacac	cactaattgc	21180
cttgggtgcc	tttcaggcgt	ctccatgcca	aggctctaac	ccctgatgtg	gaagccaaca	21240
tagtaattat	aattatagtg	aagaggggtg	tgtgtgtgtg	tgtgtgtgtg	tgtgtgtgtg	21300
tgattgaaag	agagagagaa	tgagtctctg	taccttctct	tgggtctgta	tgttgaaaac	21360
10ctaagtgacc	ctaaagaaga	atatttgcat	attattaatt	atgtagctat	acatgttttt	21420
tttgagacaa	agtcttgctt	tgtcacccag	actggagtgc	agtggcagca	atcatggctc	21480
actgcagcct	ctgcctccca	g				21501

15

<210> 2

<211> 2323

<212> DNA

<213> Homo sapiens

20

<400> 2

agagcctgtg	ctactggaag	gtggcgtgcc	ctcctctggc	tggtaccatg	cagctcccac	60
tggccctgtg	tctcgtctgc	ctgctggtac	acacagcctt	ccgtgtagtg	gagggccagg	120
ggtggcaggc	gttcaagaat	gatgccacgg	aatcatccc	cgagctcgga	gagtaccccg	180
25agcctccacc	ggagctggag	aacaacaaga	ccatgaaccg	ggcggagAAC	ggagggcggc	240
ctccccacca	cccctttgag	accaaagacg	tgtccgagta	cagctgccgc	gagctgcact	300
tcacccgcta	cgtgaccgat	gggcccgtgc	gcagcgccaa	gccggtcacc	gagctgggtg	360
gctccggcca	gtgcggcccc	gcgcgcctgc	tgcccaacgc	catcggccgc	ggcaagtgg	420
ggcgacctag	tgggcccga	ttccgctgca	tccccgaccg	ctaccgcgcg	cagcgcgtgc	480
30agctgctgtg	tcccgggtgg	gaggcgccgc	gcgcgcgcaa	ggtgcgcctg	gtggcctcgt	540
gcaagtgcaa	gcgcctcacc	cgcttccaca	accagtcgga	gctcaaggac	ttcgggaccg	600
aggccgctcg	gccgcagaag	ggccggaagc	cgcgcccccg	cgccccgagc	gccaaagcca	660
accaggccga	gctggagAAC	gcctactaga	gcccgcgccg	gcccctcccc	accggcgggc	720
gccccggccc	tgaaccgcg	ccccacattt	ctgtcctctg	cgcggtggtt	gattgtttat	780
35atttcattgt	aatgcctgc	aaccaggggc	agggggctga	gaccttccag	gccctgagga	840
atcccgggcg	ccggcaaggc	ccccctcagc	ccgccagctg	aggggtccca	cggggcaggg	900
gaggggaattg	agagtcacag	acactgagcc	acgcagcccc	gcctctgggg	ccgcctacct	960
ttgctgggtcc	cacttcagag	gaggcagaaa	tgaagcatt	ttcaccgccc	tggggtttta	1020
agggagcgg	gtgggagtgg	gaaagtccag	ggactgggtta	agaaagttag	ataagattcc	1080
40cccttgaccc	tcgctgcccc	tcagaaagcc	tgaggcgtgc	ccagagcaca	agactggggg	1140
caactgtaga	tgtggtttct	agtcctggct	ctgccactaa	cttgctgtgt	aaccttgAAC	1200
tacacaattc	tccttcggga	cctcaatttc	cactttgtaa	aatgaggggtg	gaggtgggaa	1260

11

```

taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct tttgaatggg 1320
cagaggtgag agagagagag agaaagagag agaatgaatg cagttgcatt gattcagtgc 1380
caaggtcact tccagaattc agagttgtga tgctctcttc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaaagta aagagtctat ttatggctga catattttacg gctgacaaac 1500
5tcctggaaga agctatgctg cttcccagcc tggcttcccc ggatgtttgg ctacctccac 1560
ccctccatct caaagaaata acatcatcca ttggggtaga aaaggagagg gtccgagggt 1620
ggtgggaggg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
acccatagcc atgtttttaa gtcaccttcc gaagagaagt gaaaggttca aggacactgg 1740
ccttgcaggc ccgaggggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
10caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc cacccaaaaa tctttttgaa 1980
aatcatttcc agacaacctc ttactttctg tgtagttttt aattgttaaa aaaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcgttttt ttggcaattc 2100
15ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaaagagt taagttacat 2160
atatttttcc tcaacttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag 2280
acaatgaatc atgaccgaaa gaaaaaaaaa aaaaaaaaaa aaa 2323

```

20<210> 3

<211> 213

<212> PRT

<213> Homo sapiens

25<400> 3

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Thr
  1              5              10              15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
      20              25              30
30Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
      35              40              45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
      50              55              60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
3565              70              75              80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
      85              90              95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
      100             105             110
40Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
      115             120             125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val

```

12

130	135	140
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg		
145	150	155
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln		
5	165	170
Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly		
	180	185
Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu		
	195	200
10Leu Glu Asn Ala Tyr		205
210		

<210> 4

<211> 4040

15<212> DNA

<213> Homo sapiens

<400> 4

tgaatgccac	agcttgctcag	cagccagcag	aaggcaggta	aaaggccttc	ctcactgtcc	60
20tcgaaggaac	caactctgcc	aatgtcttga	acttgcactc	ctaacctcca	gctctgtgag	120
acgacttctg	ttgttttatgc	cacccaattt	gtgggtactta	gttacggcag	cctcatcaaa	180
gtaccaccta	tgggagcctc	tatttttgcag	gtgagggcg	ggactgggct	gagttttctg	240
gaaaacagcc	ctgcaatacc	ctcatcagac	caccaaactc	ttcacactcc	ctcagacaca	300
gcattcactt	ccagaaataa	ctctaaagtt	ttgtttttgtt	tttttaaact	ttgtggaata	360
25ctactcagcc	aaaaaaaaaa	aaaaagaaaa	gaaaaaggaa	catgttactg	atatgtacaa	420
cttgataat	ggaaataatg	ctgagtga	aaaaaatccc	caaaggctac	ataactaattg	480
attccattta	tataaccttt	tttttttttt	tttgagacag	tctcactctg	tcaccagggc	540
tggagcacag	tggcgcgatc	tcagctcact	gcaacttccg	cctcctgagt	tcaagcgatt	600
ctcttgcctt	agcttcccaa	gtagctggga	ttacagggtgc	gtgccaccat	gcccagctaa	660
30tttttgtatt	tttagtagag	acagggtttc	gccatgttgg	ccaggctggg	ctcgaactcc	720
tgacctcaag	tgatctgcct	gccttggtt	cccaaagtgc	tgggattaca	tgagtgaagc	780
accgcacctg	gtcgcattta	tgtaacaatt	ttgaagtga	aaaaaaaaatg	acagaaatgg	840
agattagatg	agtagttgcc	aggggttagt	tgtgggaggg	agggaaggag	agggaaggag	900
gtgggcaaca	ggagaaagac	ttgtggtcac	agagctgtgc	tttatcttga	ctgtggtgga	960
35tccccaaatt	taccctgtgac	aagattgcat	agaactaagt	atacacacac	gtgaatgcgt	1020
gtgcacgcac	acagtagagg	ttaagccacg	ggagatgtgg	gtaagattgg	taaattgtgt	1080
caatatcaat	atcctagttg	tgatattgtc	ctatagtttc	gcaagggtgtt	attgttgtgg	1140
gaaactggat	aaaggataca	cggagtctgc	attttctttt	ttttattttt	attttttgag	1200
acggggtctc	actctgtcaa	ccaggctgga	gtgcagtggc	ccaagtatgg	ctcactgcag	1260
40cctcgacctc	aacctcaagt	gaacctcca	cctcagctc	ccaagtagct	aagaccacag	1320
gcgtgcgacc	ccatgcccag	ctaattttta	aattttttgt	agagactagg	cctcaccatg	1380
ttgcccaggc	ttttatttct	tataagtata	tttaaattta	taattatatc	cacattttta	1440

aattttaatt	taaaaaatta	ctctgaggcc	gggcattgtg	gctcatgcct	ctaatacccca	1500
gcaccttggg	aggccgagtt	gggcagatca	cccgagggtca	gaagtctgag	accagcctga	1560
gtaacatgga	gaaacccccg	tctctactaa	aaatacaaaa	ttggccgggc	gtggtggtgc	1620
attcctgtaa	tcccagctac	tcgggaggct	gaggcaggag	aattgcttga	acccaggagg	1680
5tgagggttgc	agtgagccaa	gattgtgcca	ctgcactgca	gcctgggcca	cagagagaat	1740
ctgccaaaaa	aagaaaaaaa	aaaaaatttc	agccgtacaa	ggatgttcat	agcaaccctg	1800
ctggaaatag	gaaaaaaaaat	tggaaataac	ctaaactact	cacaatagga	atcagctaaa	1860
accctggggg	tttaattcca	gggaatactg	tgaacaatga	caagtttgtg	gactgagtaa	1920
aaataaacag	ctgtcaatga	cttaacatta	aatgaaacag	cagaagatgt	cacagcaggt	1980
10tctcgtgag	ccattcagag	gggtgtggat	catttagagg	ttcaagtcca	ctggattcct	2040
ctttttcctt	ttaatatattac	ttcacttcca	aataaggaaa	ggaaaggaaa	ggaaatcacg	2100
tccagtcctg	agacttgcca	tcctgcagtc	acccctcctt	ttgtctccag	caggtggcag	2160
acgcgttcca	gggatgaatc	ccactgcctc	tgtttaatgc	agacggtcca	gccgctccca	2220
acagcaggtg	gggtataaag	catccatcct	acctgctcaa	ggaacccagg	catcagaact	2280
15gctctctccc	aagtccattg	caagaaggca	gtcgtctggt	catgagaggg	ttaacagtcc	2340
acattccaga	gcaagggaaa	aggaggctgg	agggtcatag	acaaggggag	gtggtgcgga	2400
gggccagctt	ctcacaacac	taccggctct	gctgggagag	atagatcacc	cccaacaatg	2460
gccacagctg	ttttcatctg	ccctgaagga	aactgactta	ggaagcaggt	atcagagagg	2520
gcccttcctg	agggggcttc	tgtctggctt	gtaaaactgt	cagagcagct	gcattcatgt	2580
20gtcggatgat	ggatgatgga	aaggacagtc	ggctgcagat	ggacacagcg	acttgcaagt	2640
tgaggcaggt	ggcaaaggac	ttgcagaggc	tctgcaggtg	gggcatgctg	attcattgcc	2700
cagttaaaat	accagaggat	ctgggcagcc	tcttcacagg	agctgcttgt	cctcaaacaa	2760
tctgtcttca	atgaaagatt	cctctggcct	tcctttctct	tcttgacact	caggtgtgaa	2820
tccttctccc	ccacgcctct	acctgcgccc	ccgccccccg	ccccggccct	gtgtgggtca	2880
25ttatatgcag	ggccaaggca	gcattttctc	ttagcttctt	tgtgaccagt	tggtcctggg	2940
atggcttcat	ggaacacatc	ctgtggtgtg	caccaatgaa	gctttccata	caggactcaa	3000
aactgttttt	gaaaaatgta	accagctgga	agacaagaaa	ataaaatgtc	agcactaaaa	3060
acgctggctg	tggcttttgc	taaggaaagg	aatttggtgt	tgtcttctca	cacacacaga	3120
ctggttgggg	aaatgactgt	cttcagcaca	tcaccctgcg	agccacagtg	agtgccttgg	3180
30ctcagaagtg	cctgtcacag	tgcacaggat	ccctgaggag	catgagctgg	gatttcctct	3240
gtgctgtcca	tcacaggagc	ctgagtgacc	agcgcctcct	cgatttgtaa	ccagaatcct	3300
gcctctctct	ccaagc'gggc	acccttgctc	tgaccctcta	gttctctctc	ttgccttcca	3360
gagaatacca	agagaggctt	tcttggttag	gacaatgaat	gctgagactt	gtggagttag	3420
gaccaatggg	atttctttta	aagcatcttt	ttgcctctgg	ctgggtctat	gggggtcaaa	3480
35cagaaacacc	ttgggccatt	tgttggtggg	gtgacaaatg	aacttggcct	gagaaatgga	3540
ataggccggg	ctcagccccg	cgaagcactc	agaactgcac	attttctttg	ttgagcgggt	3600
ccacagtttg	ttttgagaat	gcccaggggc	ccaggggagac	agacaattaa	aagccggagc	3660
tcattttgat	atctgaaaac	cacagccgcc	agcacgtggg	aggtgccgga	gagcaggctt	3720
gggccttgcc	tcacacgccc	cctctctctg	ggtcacctgg	gagtgccagc	agcaatttgg	3780
40aagtttgctg	agctagagga	gaagtctttg	gggagggttt	gctctgagca	cacccttttc	3840
cctccctccg	gggctgaggg	aaacatggga	ccagccctgc	cccagcctgt	cctcattggc	3900
tggcatgaag	cagagagggg	ctttaaaaaag	gcgaccgtgt	ctcggctgga	gaccagagcc	3960

14

tgtgctactg gaaggtggcg tgccctcctc tggttggtac catgcagctc ccactggccc 4020
 tgtgtctcgt ctgcctgctg 4040

<210> 5

5<211> 636

<212> DNA

<213> Mus musculus

<400> 5

10atgcagccct cactagcccc gtgcctcatc tgccctacttg tgcacgctgc cttctgtgct 60
 gtggaggggcc aggggtggca agccttcagg aatgatgccca cagaggtcat cccagggctt 120
 ggagagtacc ccgagcctcc tcctgagaac aaccagacca tgaaccgggc ggagaatgga 180
 ggcagacctc cccaccatcc ctatgacgcc aaagatgtgt ccgagtacag ctgccgcgag 240
 ctgcactaca cccgcttcct gacagacggc ccatgccgca gcgccaagcc ggtcaccgag 300
 15ttggtgtgct ccggccagtg cggccccgcg cggtgtgtgc ccaacgccat cgggcgcgctg 360
 aagtgggtggc gcccgaaagg accggatttc cgctgcatcc cggatcgcta ccgcgcgcag 420
 cgggtgcagc tgctgtgccc cggggggcgcg gcgccgcgct cgcgcaaggt gcgtctggtg 480
 gcctcgtgca agtgcaagcg cctcaccgcg ttccacaacc agtcggagct caaggacttc 540
 gggccggaga ccgcgcggcc gcagaagggt cgcaagccgc ggcccggcgc ccggggagcc 600
 20aaagccaacc aggcggagct ggagaacgcc tactag 636

<210> 6

<211> 211

<212> PRT

25<213> Mus musculus

<400> 6

Met Gln Pro Ser Leu Ala Pro Cys Leu Ile Cys Leu Leu Val His Ala
 1 5 10 15
 30Ala Phe Cys Ala Val Glu Gly Gln Gly Trp Gln Ala Phe Arg Asn Asp
 20 25 30
 Ala Thr Glu Val Ile Pro Gly Leu Gly Glu Tyr Pro Glu Pro Pro Pro
 35 40 45
 Glu Asn Asn Gln Thr Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro
 35 50 55 60
 His His Pro Tyr Asp Ala Lys Asp Val Ser Glu Tyr Ser Cys Arg Glu
 65 70 75 80
 Leu His Tyr Thr Arg Phe Leu Thr Asp Gly Pro Cys Arg Ser Ala Lys
 85 90 95
 40Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala Arg Leu
 100 105 110
 Leu Pro Asn Ala Ile Gly Arg Val Lys Trp Trp Arg Pro Asn Gly Pro

15

115	120	125
Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu		
130	135	140
Leu Cys Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val		
5145	150	155
Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln Ser Glu		
	165	170
Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly Arg Lys		
	180	185
10Pro Arg Pro Gly Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu Leu Glu		
	195	200
Asn Ala Tyr		205
210		

15<210> 7

<211> 674

<212> DNA

<213> Rattus norvegicus

20<400> 7

gaggaccgag tgcccttcct ccttctggca ccatgcagct ctcactagcc ccttgccttg	60
cctgcctgct tgtacatgca gccttcgttg ctgtggagag ccaggggtgg caagccttca	120
agaatgatgc cacagaaatc atcccgggac tcagagagta ccagagacct cctcaggaac	180
tagagaacaa ccagaccatg aaccggggccg agaacggagg cagaccccc caccatcctt	240
25atgacaccaa agacgtgtcc gagtacagct gccgcgagct gcactacacc cgcttcgtga	300
ccgacggccc gtgccgcagt gccaagccgg tcaccgagtt ggtgtgctcg ggccagtgcg	360
gccccgcgcg gctgctgccc aacgccatcg ggcgcgtagaa gtggtggcgc ccgaacggac	420
ccgacttccg ctgcatcccg gatcgctacc gcgcgcagcg ggtgcagctg ctgtgccccg.	480
gcggcgcggc gccgcgctcg cgcaaggtgc gtctggtggc ctcgtgcaag tgcaagcgcc	540
30tcacccgctt ccacaaccag tcggagctca aggacttcgg acctgagacc gcgcggccgc	600
agaagggtcg caagccgcgg ccccgcgccc ggggagccaa agccaaccag gcggagctgg	660
agaacgccta ctag	674

<210> 8

35<211> 213

<212> PRT

<213> Rattus norvegicus

<400> 8

40Met Gln Leu Ser Leu Ala Pro Cys Leu Ala Cys Leu Leu Val His Ala
1 5 10 15
Ala Phe Val Ala Val Glu Ser Gln Gly Trp Gln Ala Phe Lys Asn Asp

16

	20	25	30
	Ala Thr Glu Ile Ile Pro Gly Leu Arg Glu Tyr Pro Glu Pro Pro Gln		
	35	40	45
	Glu Leu Glu Asn Asn Gln Thr Met Asn Arg Ala Glu Asn Gly Gly Arg		
5	50	55	60
	Pro Pro His His Pro Tyr Asp Thr Lys Asp Val Ser Glu Tyr Ser Cys		
	65	70	75
	Arg Glu Leu His Tyr Thr Arg Phe Val Thr Asp Gly Pro Cys Arg Ser		
	85	90	95
10	Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala		
	100	105	110
	Arg Leu Leu Pro Asn Ala Ile Gly Arg Val Lys Trp Trp Arg Pro Asn		
	115	120	125
	Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val		
15	130	135	140
	Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg		
	145	150	155
	Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln		
	165	170	175
20	Ser Glu Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly		
	180	185	190
	Arg Lys Pro Arg Pro Arg Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu		
	195	200	205
	Leu Glu Asn Ala Tyr		
25	210		

<210> 9

<211> 21

<212> DNA

30<213> Homo sapiens

<400> 9

aagaatgatg ccacggaaat c

21

35<210> 10

<211> 21

<212> RNA

<213> Artificial Sequence

40<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

17

<221> misc_feature

<222> 20-21

<223> n = t

5<400> 10

gaaugaugcc acggaaauch n

21

<210> 11

<211> 21

10<212> RNA

<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

15

<221> misc_feature

<222> 20-21

<223> n = t

20<400> 11

gauuuccgug gcaucauuch n

21

<210> 12

<211> 21

25<212> DNA

<213> Homo sapiens

<400> 12

aatgatgcc a cggaatcat c

21

30

<210> 13

<211> 21

<212> RNA

<213> Artificial Sequence

35

<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature

40<222> 20-21

<223> n = t

WO 03/073991

PCT/US03/06193

18

<400> 13

ugaugccacg gaaaucaucn n

21

<210> 14

5<211> 21

<212> RNA

<213> Artificial Sequence

<220>

10<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature

<222> 20-21

<223> n = t

15

<400> 14

gaugauuucc guggcaucan n

21

<210> 15

20<211> 21

<212> DNA

<213> Homo sapiens

<400> 15

25aacaacaaga ccatgaaccg g

21

<210> 16

<211> 21

<212> RNA

30<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

35

<221> misc_feature

<222> 20-21

<223> n = t

40<400> 16

caacaagacc augaaccggn n

21

19

<210> 17

<211> 21

<212> RNA

<213> Artificial Sequence

5

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature

10<222> 20-21

<223> n = t

<400> 17

ccgguucaug gucuuguugn n

21

15

<210> 18

<211> 21

<212> DNA

<213> Homo sapiens

20

<400> 18

aattgagagt cacagacact g

21

<210> 19

25<211> 21

<212> RNA

<213> Artificial Sequence

<220>

30<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature

<222> 20-21

<223> n = t

35

<400> 19

uugagaguca cagacacugn n

21

<210> 20

40<211> 21

<212> RNA

<213> Artificial Sequence

20

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature

5<222> 20-21

<223> n = t

<400> 20

cagugucugu gacucucaan n

21

10

<210> 21

<211> 21

<212> DNA

<213> Homo sapiens

15

<400> 21

aaatggaagc attttcacccg c

21

<210> 22

20<211> 21

<212> RNA

<213> Artificial Sequence

<220>

25<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature

<222> 20-21

<223> n = t

30

<400> 22

auggaagcau uuucaccgcn n

21

<210> 23

35<211> 21

<212> RNA

<213> Artificial Sequence

<220>

40<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

WO 03/073991

PCT/US03/06193

21

<221> misc_feature

<222> 20-21

<223> n = t

5<400> 23

gcggugaaaa ugcuuccaun n

21

<210> 24

<211> 21

10<212> DNA

<213> Homo sapiens

<400> 24

aaagtccagg gactgggtaa g

21

15

<210> 25

<211> 21

<212> RNA

<213> Artificial Sequence

20

<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature

25<222> 20-21

<223> n = t

<400> 25

aguccaggga cugguuaagn n

21

30

<210> 26

<211> 21

<212> RNA

35<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

40<221> misc_feature

<222> 20-21

<223> n = t

22

<400> 26

cuuaaccagu ccugggacun n

21

<210> 27

5<211> 21

<212> DNA

<213> Homo sapiens

<400> 27

10aagaaagttg gataagattc c

21

<210> 28

<211> 21

<212> RNA

15<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

20<221> misc_feature

<222> 20-21

<223> n = t

<400> 28

25gaaaguugga uaagauuccn n

21

<210> 29

<211> 21

<212> RNA

30<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

35<221> misc_feature

<222> 20-21

<223> n = t

<400> 29

40ggaaucuuau ccaacuuucn n

21

23

<210> 30
 <211> 21
 <212> DNA
 <213> Homo sapiens

5

<400> 30
 aactgtagat gtggtttcta g

21

10<210> 31
 <211> 21
 <212> RNA
 <213> Artificial Sequence

15<220>

<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature
 <222> 20-21

20<223> n = t

<400> 31
 cuguagaugu gguuucuagn n

21

25<210> 32
 <211> 21
 <212> RNA
 <213> Artificial Sequence

30<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature
 <222> 20-21

35<223> n = t

<400> 32
 cuagaaacca caucuacagn n

21

40

24

<210> 33
 <211> 21
 <212> DNA
 <213> Homo sapiens

5

<400> 33
 aattctcctt cgggacctca a

21

<210> 34
 10<211> 21
 <212> RNA
 <213> Artificial Sequence

<220>

15<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature
 <222> 20-21
 <223> n = t

20

<400> 34
 uucuccuucg ggaccucaan n

21

<210> 35
 25<211> 21
 <212> RNA
 <213> Artificial Sequence

<220>

30<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature
 <222> 20-21
 <223> n = t

35

<400> 35
 uugagguccc gaaggagaan n

21

<210> 36
 40<211> 21
 <212> DNA
 <213> Homo sapiens

WO 03/073991

PCT/US03/06193

25

<400> 36

aaagagagag aatgaatgca g

21

<210> 37

5<211> 21

<212> RNA

<213> Artificial Sequence

<220>

10<223> Small Interfering RNA to Block Expression of SOST (sense strand)

<221> misc_feature

<222> 20-21

<223> n = t

15

<400> 37

agagagagaa ugaaugcagn n

21

<210> 38

20<211> 21

<212> RNA

<213> Artificial Sequence

<220>

25<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

<221> misc_feature

<222> 20-21

<223> n = t

30

<400> 38

cugcauucan ucucucucun n

21

<210> 39

35<211> 21

<212> DNA

<213> Homo sapiens

<400> 39

40aagaagctat gctgcttccc a

21

26

<210> 40
 <211> 21
 <212> RNA
 <213> Artificial Sequence
 5
 <220>
 <223> Small Interfering RNA to Block Expression of SOST (sense strand)

 <221> misc_feature
 10<222> 20-21
 <223> n = t

 <400> 40
 gaagcuaugc ugcuuccan n 21
 15
 <210> 41
 <211> 18
 <212> RNA
 <213> Artificial Sequence
 20
 <220>
 <223> Small Interfering RNA to Block Expression of SOST (antisense strand)

 <400> 41
 25ugggaagcag cauagcuu 18

 <210> 42
 <211> 21
 <212> DNA
 30<213> Homo sapiens

 <400> 42
 aaatcacatc cgccccaact t 21

 35<210> 43
 <211> 21
 <212> RNA
 <213> Artificial Sequence

 40<220>
 <223> Small Interfering RNA to Block Expression of SOST (sense strand)

27

<221> misc_feature

<222> 20-21

<223> n = t

5<400> 43

aucacauccg ccccaacuun n

21

<210> 44

<211> 21

10<212> RNA

<213> Artificial Sequence

<220>

<223> Small Interfering RNA to Block Expression of SOST (antisense strand)

15

<221> misc_feature

<222> 20-21

<223> n = t


20<400> 44

aaguuggggc ggaugugaun n

21

WHAT IS CLAIMED:


1. A pharmaceutical composition comprising a carrier and a compound of the formula I:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl; and wherein the composition can modulate SOST expression in a mammal.

2. The pharmaceutical composition of claim 1, wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 groups are each separately carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl.
3. The pharmaceutical composition of claim 1, wherein R_1 is carbonyl or hydroxy.
4. The pharmaceutical composition of claim 1, wherein R_8 is fluorine, hydrogen or halo.
5. A pharmaceutical composition for decreasing SOST expression in a mammal comprising a therapeutically effective amount of a glucocorticoid.


6. The pharmaceutical composition of claim 5, wherein the glucocorticoid can bind to the glucocorticoid receptor.
7. The pharmaceutical composition of claim 5, wherein the glucocorticoid is fluocinolone acetonide, triamcinolone or dexamethasone.
8. A pharmaceutical composition comprising a carrier and a prostaglandin comprising formula II:



wherein R_{10} and R_{11} are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl, aryl or aryloxy; and
 wherein the composition can modulate SOST expression in a mammal.

9. The pharmaceutical composition of claim 8, wherein R_1 and R_4 are each separately carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl.
10. The pharmaceutical composition of claim 8, wherein the prostaglandin is prostaglandin E2.

11. A pharmaceutical composition comprising a carrier and a bile salt of formula III:




- 5 wherein $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8$ and R_9 are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl; and wherein the composition can modulate SOST expression in a mammal.

- 10 12. The pharmaceutical composition of claim 11, wherein $R_1, R_2, R_3, R_4, R_5, R_6, R_7$ and R_9 are each separately carbonyl, hydroxy, hydrogen, or lower hydroxy alkyl.
- 15 13. The pharmaceutical composition of claim 11, wherein the bile salt is ursodeoxycholic acid.
- 20 14. A pharmaceutical composition comprising a carrier and a siRNA comprising SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41 or SEQ ID NO:44, wherein the siRNA can modulate SOST expression.
15. A pharmaceutical composition comprising a carrier and a siRNA that is selectively hybridizable under stringent conditions to an RNA


derived from a DNA comprising SEQ ID NO:9, SEQ ID NO:12,
 SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24,
 SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36,
 SEQ ID NO:39, or SEQ ID NO:42, wherein the siRNA can modulate
 SOST expression.

16. A method for modulating SOST expression in a mammal that
 comprises administering to the mammal ursodeoxycholic acid,
 fluocinolone acetonide, triamcinolone, prostaglandin E2 or
 dexamethasone.
17. A method for modulating SOST expression in a mammal that
 comprises administering to the mammal a compound of formula I:




wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are each separately
 carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower
 alkoxy, lower alkyl, or lower hydroxy alkyl.

18. A method for modulating SOST expression in a mammal that
 comprises administering to the mammal a compound of formula II:



wherein R_{10} and R_{11} are separately carbonyl, halo, fluorine, hydrogen,
hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl,
aryl or aryloxy.


19. A method for modulating SOST expression in a mammal that
comprises administering to the mammal a compound of formula III:



wherein $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8$ and R_9 are each separately
carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower
alkoxy, lower alkyl, or lower hydroxy alkyl.

20. A method for modulating SOST expression in a mammal that
comprises administering to the mammal a glucocorticoid.


21. The method of claim 20, wherein the glucocorticoid is cortisol or dexamethasone.
22. A method for modulating SOST expression in a mammal that
5 comprises administering to the mammal a siRNA comprising SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41 or SEQ ID NO:44, wherein the siRNA can modulate SOST expression.
- 10 23. A method for modulating SOST expression in a mammal that comprises administering to the mammal a siRNA that is selectively hybridizable under stringent conditions to an RNA derived from a DNA comprising SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:15,
15 SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:39, or SEQ ID NO:42, wherein the siRNA can modulate SOST expression.
24. A method for increasing bone density in a mammal that comprises
20 administering ursodeoxycholic acid, fluocinolone acetonide, triamcinolone, prostaglandin E2 or dexamethasone to the mammal.
25. A method for increasing bone density in a mammal that comprises administering to the mammal a compound of formula I:
- 25



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl.

5

26. A method for increasing bone density in a mammal that comprises administering to the mammal a compound of formula II:




10

wherein R_{10} and R_{11} are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl, aryl or aryloxy.

15

27. A method for increasing bone density in a mammal that comprises administering to the mammal a compound of formula III:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 and R_9 are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl.

5

28. A method for increasing bone density in a mammal that comprises administering to the mammal a glucocorticoid, wherein the glucocorticoid modulates SOST expression.

10

29. The method of claim 28, wherein the glucocorticoid is cortisol or dexamethasone.

15

30. A method for increasing bone density in a mammal that comprises administering to the mammal a siRNA comprising SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41 or SEQ ID NO:44, wherein the siRNA can modulate SOST expression.


20

31. A method for increasing bone density in a mammal that comprises administering to the mammal a siRNA that is selectively hybridizable under stringent conditions to an RNA derived from a DNA comprising SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:27, SEQ ID

NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:39, or SEQ ID NO:42, wherein the siRNA can modulate SOST expression.


32. A method for decreasing apoptosis of bone cells in a mammal that comprises administering ursodeoxycholic acid, flucinolone acetonide, triamcinolone, prostaglandin E2 or dexamethasone to the mammal.

33. A method for decreasing apoptosis of bone cells in a mammal that comprises administering to the mammal a compound of formula I:



wherein R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl.


34. A method for decreasing apoptosis of bone cells in a mammal that comprises administering to the mammal a compound of formula II:



wherein R_{10} and R_{11} are separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, lower hydroxy alkyl, aryl or aryloxy.

5

35. A method for decreasing apoptosis of bone cells in a mammal that comprises administering to the mammal a compound of formula III:



wherein $R_1, R_2, R_3, R_4, R_5, R_6, R_7, R_8$ and R_9 are each separately carbonyl, halo, fluorine, hydrogen, hydroxyl, lower acyl, lower alkoxy, lower alkyl, or lower hydroxy alkyl.

10

36. A method for decreasing apoptosis of bone cells in a mammal that comprises administering to the mammal a glucocorticoid, wherein the glucocorticoid modulates SOST expression.

15

37. The method of claim 36, wherein the glucocorticoid is cortisol or dexamethasone.

38. A method for decreasing apoptosis of bone cells in a mammal that comprises administering to the mammal a siRNA comprising SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID

20

NO:35, SEQ ID NO:38, SEQ ID NO:41 or SEQ ID NO:44, wherein the siRNA can modulate SOST expression.

39. A method for decreasing apoptosis of bone cells in a mammal that
5 comprises administering to the mammal a siRNA that is selectively hybridizable under stringent conditions to an RNA derived from a DNA comprising SEQ ID NO:9, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:39, or
10 SEQ ID NO:42, wherein the siRNA can modulate SOST expression.
40. A method for identifying a factor that decreases SOST expression comprising:
- (1) providing a cell that comprises a nucleic acid construct
15 comprising a SOST promoter comprising SEQ ID NO:4 that is operably linked to a nucleic acid encoding a detectable marker;
- (2) contacting the cell with a test agent; and
- (3) detecting whether the test agent decreases expression of the
20 detectable marker relative to a control that comprises a cell that has not been contacted with a test agent.
41. The method of claim 40, wherein the cell is responsive to a bone morphogenetic protein.
- 25 42. The method of claim 40, wherein the cell is a C2C12 cell from a cell line having ATCC Deposit No. CRL-1772 or a C3H10T1/2 cell from a cell line having ATCC Deposit No. CCL-226.
- 30 43. The method of claim 40, wherein the cell is a human mesenchymal (hMSC) cell.

44. The method of claim 40, wherein the nucleic acid construct further comprises a nucleic acid segment encoding an osteoblastic-specific steroid response element.
- 5
45. The method of claim 40, wherein the nucleic acid construct further comprises a nucleic acid segment encoding glucocorticoid response element.
- 10
46. A method for identifying a factor that decreases SOST expression comprising:
- 15
- (1) contacting a cell with a test agent; and
- (2) detecting whether endogenous SOST expression decreases, wherein the cell comprises a nucleic acid encoding a protein comprising SEQ ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8.
47. The method of claim 46, wherein the nucleic acid comprises SEQ ID NO:1 or SEQ ID NO:2 or SEQ ID NO:5 or SEQ ID NO:7.
- 20
48. A method for identifying an antagonist for sclerostin comprising:
- (1) contacting a cell with sclerostin and a test agent; and
- (2) detecting whether the test agent prevents apoptosis of the cell.
- 25
49. A method for identifying a molecule that binds to sclerostin comprising:
- (1) contacting a test agent with a sclerostin polypeptide comprising SEQ ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8; and
- (2) determining whether the test agent binds to the polypeptide.
- 30

50. A marker for osteoblast differentiation comprising a sclerostin polypeptide comprising SEQ ID NO:3 or SEQ ID NO:6 or SEQ ID NO:8.
- 5 51. A marker for osteoblast differentiation comprising a nucleic acid comprising SEQ ID NO:1 or SEQ ID NO:2 or SEQ ID NO:5 or SEQ ID NO:7.




FIG. 1A




FIG. 1B

3/28




FIG. 2





FIG. 3

5/28



*No bands for gremlin, noggin in absence of RT

FIG. 4

6/28




FIG. 5

7/28





FIG. 6


8/28



DEX = dexamethasone
Triam = triamcinolone
Urso = ursodeoxycholic acid (2 concentrations tested)

FIG. 7

9/28



DEX = dexamethasone
Tauro = tauroursodeoxycholic acid
Fluo = fluocinolone acetonide
Lova = lovastatin
ICI = ICI 182,780
Estrogen = 17 β -estradiol
PGE2 = prostaglandin E2

FIG. 8




FIG. 9C

FIG. 9B

FIG. 9A




FIG. 9D

FIG. 9E

11/28




FIG. 10




FIG. 11

13/28




FIG. 12A

14/28




FIG. 12B

15/28




FIG. 13

16/28




FIG. 14




FIG. 15




FIG. 16




FIG. 17A




FIG. 17B

19/28




FIG. 17C




FIG. 17D




FIG. 18




FIG. 19

22/28




FIG. 20




FIG. 21

23/28




FIG. 22A




FIG. 22B

24/28




FIG. 23

25/28




FIG. 24A





FIG. 24B






FIG. 26




FIG. 27

