



(72) TEREK, RICHARD M., US

(71) RHODE ISLAND HOSPITAL, A LIFESPAN PARTNER, US

(51) Int.Cl.⁶ C12N 15/12, C12N 15/63, G01N 33/574, C07K 14/47

(30) 1998/03/13 (09/042,225) US

(54) **GENES ASSOCIES AU CHONDROSARCOME**

(54) **CHONDROSARCOMA ASSOCIATED GENES**

(57) L'invention concerne une molécule d'acide nucléique codant un polypeptide associé au chondrosarcome, ainsi que des méthodes diagnostiques pour des patients atteints du chondrosarcome.

(57) The invention features a nucleic acid molecule encoding a chondrosarcoma associated polypeptide and methods for diagnosing patients with chondrosarcoma.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, C12N 15/63, G01N 33/574</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/46382 (43) International Publication Date: 16 September 1999 (16.09.99)</p>
<p>(21) International Application Number: PCT/US99/05348 (22) International Filing Date: 12 March 1999 (12.03.99) (30) Priority Data: 09/042,225 13 March 1998 (13.03.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/042,225 (CON) Filed on 13 March 1998 (13.03.98) (71) Applicant (for all designated States except US): RHODE ISLAND HOSPITAL, A LIFESPAN PARTNER [US/US]; 593 Eddy Street, Providence, RI 02903 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): TEREK, Richard, M. [US/US]; 129 Benefit Street, 5, Providence, RI 02903 (US). (74) Agent: BEATTIE, Ingrid, A.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: CHONDROSARCOMA ASSOCIATED GENES</p> <p>(57) Abstract</p> <p>The invention features a nucleic acid molecule encoding a chondrosarcoma associated polypeptide and methods for diagnosing patients with chondrosarcoma.</p>		

- 1 -

CHONDROSARCOMA ASSOCIATED GENESBackground of the Invention

The invention relates to bone malignancies.

5 Chondrosarcoma, which usually occurs in late adulthood and old age, is the second most common form of bone malignancy. Conventional chondrosarcoma tumors are graded from stage I through stage III, stage III being the most advanced. In addition to conventional
10 chondrosarcoma, there are other types of chondrosarcoma with distinguishing characteristics: myxoid, mesenchymal, clear cell, and dedifferentiated (spindle cell) chondrosarcoma.

Diagnosis and grading of chondrosarcoma has been
15 problematic. For example, the criteria used to distinguish benign enchondroma from low grade chondrosarcoma include parameters which are difficult to quantify such as increased cellularity and more than occasional binucleate cells. The histologic criteria are
20 not absolute, and the diagnosis is frequently made by taking into account clinical features such as pain, rate of growth, location, and radiologic features. Furthermore, the location of the tumor may affect clinical assessment. For example, lesions in the hand
25 can appear aggressive histologically and yet behave benignly. In contrast, lesions occurring in the pelvis are likely to represent a malignancy despite a relatively innocuous histologic appearance. Notwithstanding attempts to integrate clinicopathologic criteria, it has
30 not been possible to predict which tumors will metastasize or recur.

Summary of the Invention

The invention is based on the discovery of a novel gene which is differentially expressed in chondrosarcoma
35 cells. Accordingly the invention features an isolated

- 2 -

nucleic acid (e.g., genomic DNA, cDNA or synthetic DNA) encoding a chondrosarcoma associated (CSA) polypeptide such as human CSA-1. The term "chondrosarcoma associated" refers to the property of differential expression in chondrosarcoma cell compared to normal cartilage cells. For example, a CSA gene product is expressed at a detectably higher or lower level compared to the level at which it is expressed in normal cartilage cells. A CSA gene product may be expressed solely in chondrosarcoma cells (and not in normal cartilage cells).

The nucleic acid molecule contains a nucleotide sequence encoding a polypeptide having an amino acid sequence that is at least 80% identical to the amino acid sequence of CSA-1 (SEQ ID NO:2). Preferably, the nucleic acid molecule contains the nucleotide sequence of SEQ ID NO:1 or a degenerate variant thereof. For example, the nucleic acid contain the nucleotide sequence of SEQ ID NO:3. The invention also includes a nucleic acid molecule which contains a strand which hybridizes at high stringency to a DNA having the sequence of SEQ ID NO:1, or the complement thereof. A substantially pure DNA having at least 50% sequence identity (preferably at least 70%, more preferably at least 80%, and most preferably at least 90%) to SEQ ID NO:1, and encoding a polypeptide having the differential pattern of expression of a CSA-1 polypeptide is also within the invention. For expression of a CSA polypeptide, a CSA polypeptide encoding nucleic acid molecule is operably linked to regulatory sequences, e.g., a promoter.

The invention also includes a substantially pure CSA polypeptide such as human CSA-1 or a fragment thereof. CSA-1 fragments, e.g., a fragment containing the amino acid sequence of SEQ ID NO: 8), are useful as immunogens for raising anti-CSA antibodies. The CSA polypeptide preferably contains an amino acid sequence

- 3 -

that is at least 50% identical to the amino acid sequence of SEQ ID NO:2. More preferably the amino acid sequence of the polypeptide is 75%, 85%, 95%, 98%, and most preferably 100% identical to the amino acid sequence of
5 SEQ ID NO:2. A cell containing a CSA polypeptide-encoding nucleic acid molecule is also within the invention, as is a method of making a CSA polypeptide. Such a method may involve the following steps: (a) providing cell containing a CSA polypeptide-encoding
10 nucleic acid molecule, and (b) culturing it under conditions permitting expression of the nucleic acid molecule.

By "isolated nucleic acid molecule" is meant a nucleic acid molecule that is free of the genes which, in
15 the naturally-occurring genome of the organism, flank a *csa* gene. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or which exists
20 as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence. The term
25 excludes large segments of genomic DNA, e.g., such as those present in cosmid clones, which contain a gene of interest, e.g., a *csa* gene, flanked by one or more other genes which naturally flank it in a naturally-occurring genome.

30 Nucleic acid molecules include both RNA and DNA, including cDNA, genomic DNA, and synthetic (e.g., chemically synthesized) DNA. Where single-stranded, the nucleic acid molecule may be a sense strand or an antisense strand. The term therefore includes, for
35 example, a recombinant DNA which is incorporated into a

- 4 -

vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote at a site other than its natural site; or which exists as a separate molecule (e.g., a cDNA or a genomic
5 or cDNA fragment produced by polymerase chain reaction (PCR) or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

10 Hybridization is carried out using standard techniques such as those described in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, (1989). "High stringency" refers to DNA hybridization and wash conditions characterized by high
15 temperature and low salt concentration, e.g., wash conditions of 65° C at a salt concentration of approximately 0.1 X SSC. "Low" to "moderate" stringency refers to DNA hybridization and wash conditions characterized by low temperature and high salt
20 concentration, e.g. wash conditions of less than 60° C at a salt concentration of at least 1.0 X SSC. For example, high stringency conditions may include hybridization at about 42°C, and about 50% formamide; a first wash at about 65°C, about 2 X SSC, and 1% SDS; followed by a
25 second wash at about 65°C and about 0.1% x SSC. Lower stringency conditions suitable for detecting DNA sequences having about 50% sequence identity to *csa-1* gene are detected by, for example, hybridization at about 42°C in the absence of formamide; a first wash at about
30 42°C, about 6 X SSC, and about 1% SDS; and a second wash at about 50°C, about 6 X SSC, and about 1% SDS.

Where a particular polypeptide or nucleic acid molecule is said to have a specific percent identity to a reference polypeptide or nucleic acid molecule of a
35 defined length, the percent identity is relative to the

- 5 -

reference polypeptide or nucleic acid molecule. Thus, a peptide that is 50% identical to a reference polypeptide that is 100 amino acids long can be a 50 amino acid polypeptide that is completely identical to a 50 amino acid long portion of the reference polypeptide. It might also be a 100 amino acid long polypeptide which is 50% identical to the reference polypeptide over its entire length. Of course, many other polypeptides will meet the same criteria. The same rule applies for nucleic acid molecules.

For polypeptides, the length of the reference polypeptide sequence will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids, 50 amino acids, or 100 amino acids. For nucleic acids, the length of the reference nucleic acid sequence will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 100 nucleotides or 300 nucleotides.

In the case of polypeptide sequences which are less than 100% identical to a reference sequence, the non-identical positions are preferably, but not necessarily, conservative substitutions for the reference sequence. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine.

Sequence identity is measured using sequence analysis software (for example, the Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710

- 6 -

University Avenue, Madison, WI 53705), with the default parameters as specified therein.

By "promoter" is meant a minimal DNA sequence sufficient to direct transcription. Promoters may be
5 constitutive or inducible, and may be coupled to other regulatory sequences or "elements" which render promoter-dependent gene expression cell-type specific, tissue-specific or inducible by external signals or agents; such elements may be located in the 5' or 3' region of the
10 native gene, or within an intron. DNA encoding a CSA polypeptide may be operably linked to such regulatory sequences for expression of the polypeptide in prokaryotic or eukaryotic cells. By "operably linked" is meant that a coding sequence and a regulatory sequence(s)
15 are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequence(s).

A protein is substantially free of naturally
20 associated components when it is separated from those contaminants which accompany it in its natural state. (proteins and other naturally-occurring organic molecules) which naturally accompany it. Typically, the polypeptide is substantially pure when it constitutes at
25 least 60%, by weight, of the protein in the preparation. Preferably, the protein in the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, CSA-1 polypeptide. A substantially pure CSA-1 polypeptide may be obtained, for example, by
30 extraction from a natural source (e.g., a chondrosarcoma cell); by expression of a recombinant nucleic acid encoding an CSA-1 polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, e.g., column chromatography,
35 polyacrylamide gel electrophoresis, or HPLC analysis.

- 7 -

Accordingly, substantially pure polypeptides include recombinant polypeptides derived from a eukaryote but produced in *E. coli* or another prokaryote, or in a eukaryote other than that from which the polypeptide was
5 originally derived.

The invention also features CSA polypeptide binding species, such as an antibody or antibody fragment which specifically binds to a CSA polypeptide, e.g., a CSA-1-specific antibody. Antibodies specific for a CSA
10 polypeptide are useful to diagnose chondrosarcoma.

Chondrosarcoma is diagnosed by measuring expression of *csa* gene expression in patient tissue samples. Expression of CSA-1 is detectable in chondrosarcoma cells but not in normal cells (or in
15 certain other types of tumors which were tested). Thus, the use of CSA polypeptides and CSA polypeptide-encoding nucleic acid molecules in diagnosing and grading of chondrosarcoma is also within the invention. For example, a method for diagnosing the presence of a
20 chondrosarcoma cell in a tissue sample is carried out by measuring expression of a *csa* gene, e.g., a gene encoding CSA-1, in the tissue sample and a control sample such as a normal nonneoplastic cartilage cell. An increase in expression of the *csa* gene in the tissue sample compared
25 to the control sample indicates that the tissue sample contains a chondrosarcoma cell. A method of grading a chondrosarcoma tumor may be carried out by determining the level of *csa-1* gene expression in a test sample and comparing it to the level of *csa-1* gene expression in a
30 control sample. The level of expression in the test sample compared to the control sample is directly proportional to the grade or stage of tumor, i.e., the greater the level of expression of a *csa* gene the more advanced the stage of the tumor.

- 8 -

In addition to evaluating tissue biopsy samples for *csa* gene expression, *csa* gene expression may be detected *in vivo*. For example, a diagnostically effective amount of a detectably labeled CSA-1-specific binding species may be administered to a patient, followed by a determination of whether the species specifically binds to cartilage cells of the patient. Binding of the CSA-1 binding species, e.g., a CSA-1-specific antibody, antibody fragment, or non-antibody CSA-1 binding compound, to patient cells is an indication of the presence of chondrosarcoma in the patient. The level of binding correlates with the grade of the chondrosarcoma, i.e., a greater amount of binding compared to a normal control of known low grade tumor indicates that the patient's tumor is of a high grade. Similarly, a method of detecting progressive chondrosarcoma in a patient may be carried out as follows: (a) successively administering to a patient suspected of having a chondrosarcoma a diagnostically effective amount of a detectably labeled CSA-1-specific binding species (e.g., an antibody labelled with a radioisotope or a paramagnetic label), and (b) comparing the amount of the species that binds to cartilage cells of the patient in each successive administration to detect an increase of binding of the binding species over time. An increase in binding over time is an indication of progressive chondrosarcoma in said patient. Where the detecting step is quantitative, the amount of binding would correlate with and allow diagnosis of the severity of the disease. A diagnostic method carried out multiple times by repeatedly administering at spaced intervals the labelled binding species to the patient, with the administrations spaced by, e.g., a day, a week, a month, several months, or even years, is a useful method for detecting progression of disease in a patient.

- 9 -

Compounds capable of inhibiting expression of a CSA polypeptide may be therapeutically useful to treat chondrosarcoma. Accordingly, the invention includes a method of screening a candidate compound to identify a
5 compound capable of inhibiting expression of a CSA polypeptide, e.g., CSA-1, by (a) providing a chondrosarcoma cell expressing a CSA polypeptide, (b) contacting the cell with the candidate compound, and (c) determining the amount of expression of the CSA
10 polypeptide by the cell. A decrease in the amount of CSA polypeptide expression in the presence of the candidate compound compared to that in the absence of the candidate compound indicates that the compound inhibits expression of the CSA polypeptide.

15 The invention also includes a method of inhibiting the expression or activity of CSA-1. Suitable antagonists include a nucleic acid molecule that interfere with transcription or translation of *csa-1*, for example, antisense nucleic acid molecules and ribozymes.
20 Also included are antibodies or other suitable antagonistic molecules, that specifically binds CSA-1 polypeptide and "neutralize" its activity.

In addition to diagnostic methods, such as described above, the present invention encompasses
25 methods and compositions for evaluating appropriate treatment, and treatment effectiveness of malignancies associated with expression of *csa-1*. For example, the *csa-1* can be used as a probe to classify cells in terms of their level of *csa-1* expression, or as primers for
30 diagnostic PCR analysis in which mutations and allelic variation of *csa-1* can be detected.

The invention also includes non-human transgenic animals that express human CSA-1 and non-human transgenic mammal with a null mutation in its endogenous CSA-1 gene.
35 These animals can serve as new and useful models of

- 10 -

chondrosarcoma. The invention also includes a transgenic non-human mammal, e.g., a rodent such as a mouse, the germ cells and somatic cells of which contain a null mutation, e.g., a deletion, in DNA encoding a *csa* gene.

5 By "null mutation" is meant an alteration in the nucleotide sequence that renders the gene incapable of expressing a functional protein product. The mutation could be in *csa* gene regulatory regions or in the coding sequence. It can, e.g., introduce a stop codon that

10 results in production of a truncated, inactive gene product or it can be a deletion of all or a substantial portion of the coding sequence.

The invention also features an isolated nucleic acid (e.g., genomic DNA, cDNA or synthetic DNA) encoding

15 a cartilage associated (CAA) polypeptide such as human CAA-1. The term "cartilage associated" refers to the property of differential expression in cells of the cartilage lineage compared to cells of other tissue specificities. For example, a CAA gene product is

20 expressed in normal cartilage cells and chondrosarcoma cells but not cells of other tissue specificities or other tumor types.

The nucleic acid molecule contains a nucleotide sequence encoding a polypeptide having an amino acid

25 sequence that is at least 80% identical to the amino acid sequence of CAA-1 (SEQ ID NO:7). Preferably, the nucleic acid molecule contains the nucleotide sequence of SEQ ID NO:6 or a degenerate variant thereof. For example, the nucleic acid may have the nucleotide sequence of SEQ ID

30 NO:5. The invention also includes a nucleic acid molecule which contains a strand which hybridizes at high stringency to a DNA having the sequence of SEQ ID NO:6, or the complement thereof. A substantially pure DNA having at least 50% sequence identity (preferably at

35 least 70%, more preferably at least 80%, and most

- 11 -

preferably at least 90%) to SEQ ID NO:7, and encoding a polypeptide having the activity of CAA-1. By the activity of CAA-1 is meant inhibition of interferon gamma induced upregulation of HLA class II antigens. For
5 expression of a CAA polypeptide, a CAA polypeptide encoding nucleic acid molecule is operably linked to regulatory sequences, e.g., a promoter.

The invention also includes a substantially pure CAA polypeptide such as human CAA-1 or a fragment
10 thereof. The CAA-1 polypeptide preferably contains an amino acid sequence that is at least 50% identical to the amino acid sequence of SEQ ID NO:7. More preferably the amino acid sequence of the polypeptide is 75%, 85%, 95%, 98%, and most preferably 100% identical to the amino acid
15 sequence of SEQ ID NO:7. A cell containing a CAA polypeptide-encoding nucleic acid molecule is also within the invention, as is a method of making a CAA polypeptide. Such a method may involve the following steps: (a) providing cell containing a CAA polypeptide-
20 encoding nucleic acid molecule, and (b) culturing it under conditions permitting expression of the nucleic acid molecule.

The invention also features CAA polypeptide binding species, such as an antibody or antibody fragment
25 which specifically binds to a CAA polypeptide, e.g., a CAA-1-specific antibody. Antibodies specific for a CAA polypeptide are useful to for tissue typing and for therapeutic applications, e.g., to inhibit the activity of CAA-1.

30 Compounds capable of inhibiting expression of a CAA polypeptide may be therapeutically useful to treat conditions, e.g., rheumatoid arthritis, associated with undesired or pathologic joint inflammation. Accordingly, the invention includes a method of screening a candidate
35 compound to identify a compound capable of inhibiting

- 12 -

expression of a CAA polypeptide, e.g., CAA-1, by (a) providing a cell expressing a CAA polypeptide, (b) contacting the cell with the candidate compound, and (c) determining the amount of expression of the CAA polypeptide by the cell. A decrease in the amount of CAA polypeptide expression in the presence of the candidate compound compared to that in the absence of the candidate compound indicates that the compound inhibits expression of the CAA polypeptide. A method of identifying a compound which inhibits the activity of CAA-1 can be carried out as follows: (a) providing a cell expressing a CAA polypeptide, (b) contacting the cell with the candidate compound, and (c) determining the amount of HLA II expression by the cell. A decrease in the amount of HLA II expression in the presence of the candidate compound compared to that in the absence of the candidate compound indicates that the compound inhibits HLA II expression in the cell.

Methods of treating undesired inflammation such as that associated with rheumatoid arthritis and other inflammatory arthropathies are also within the invention. Such a method may be carried out by administering to a mammal in need of such therapy, e.g. a patient suffering from rheumatoid arthritis or other inflammatory arthropathies, an effective amount of a CAA-1 polypeptide. For example, the peptide is administered locally at the site of a rheumatoid lesion to reduce local inflammation and swelling.

The invention also includes a non-human transgenic mammal that expresses human CAA-1 and non-human transgenic mammal with a null mutation in its endogenous CAA-1 gene.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

- 13 -

Detailed Description

To date, no consistent genetic abnormality has been associated with chondrosarcoma. The tumors are heterogeneous and often have a number of abnormalities in gene expression. The following examples provide evidence of a novel gene, *csa-1*, that is expressed in a human chondrosarcoma cell line and in cartilaginous neoplasms but not in normal cartilage. The level of expression of a CSA-1 polypeptide correlates with the histological grade of the neoplasm, i.e., an increase in expression indicates a higher grade tumor. Detection of a CSA-1 gene product or *csa-1* transcript is a means by which to distinguish a neoplastic cell from a normal cartilage cell.

CSA polypeptides and CAA polypeptides may be used therapeutically. For example, a CAA polypeptide such as CAA-1 may function as a tumor suppressor. CAA-1 can also be administered to patients to reduce undesired inflammation such as joint inflammation in rheumatoid arthritis.

CSA-1 plays a role in potentiating chondrogenesis associated with chondrosarcoma. Transfecting the nonexpressing or normal cell lines with vectors which promote high levels of expression of a CSA polypeptide, e.g., CSA-1, followed by transformation of a low grade cell line to a high grade cell line indicates that CSA expression potentiates neoplastic growth. Inhibitors of CSA-1 expression can slow or inhibit neoplastic growth. Transformation and grading of transformed cells is evaluated by examining changes in morphology, proliferation, adhesion, and invasiveness.

Two novel genes have been cloned. CSA-1 is expressed in a tumor cell line and also in some high grade chondrosarcoma, but not normal cartilage, or low, or intermediate grade tumors. A second gene, CAA-1, is

- 14 -

expressed in normal cartilage and an intermediate grade tumor cell line, and as alternative sized messages in a high grade cell line.

Example 1: Cloning of *csa* Genes

5 Chondrosarcoma cell lines derived from human grade I, II and III chondrosarcomas were alternately cultured in monolayer cultures and in agarose suspension cultures using standard methods. Total cellular RNA was isolated using standard techniques. Three different human
10 chondrosarcoma cell lines (AQ, stage II tumor; FS, stage II tumor; and MW, stage I tumor) and normal articular cartilage obtained from amputation specimens was analyzed.

Using the differential mRNA display technique, a
15 technique that systematically amplifies mRNAs by means of RT-PCR with different sets of 5' arbitrary primers and 3' oligo-dT anchoring primers, the mRNA patterns of different cells and cell types were compared. The PCR products were resolved on a denaturing polyacrylamide
20 sequencing gel to display mRNA patterns that distinguish one cell type from another. The bands that were separated by gel electrophoresis represent the 3'-termini of the cDNAs. Therefore, a band that is present in one cell type, e.g., a chondrosarcoma cell line or
25 chondrosarcoma biopsy tissue, but not in the normal cartilage tissue or in noncartilage tissue, suggests that the gene is differentially expressed in chondrosarcomas.

cDNA was generated using reverse transcriptase and an oligo-dT primer (TTTTTTTTTTTMMN (SEQ ID NO:4), where M
30 can be C, G, or A; N can be C, G, A, or T). A PCR reaction was then carried out in triplicate with the same oligo-dT primer and a second random ten base pair primer (RNAmapp, GenHunter Corp., Brookline, MA). This combination of primers amplified approximately 100 cDNAs
35 from 100-500 base pairs long. The cDNAs were separated

- 15 -

on a sequencing gel. A band that is present in one or more of the cancer cell lines and absent in a normal cell may represent an oncogene which is being expressed in the cancer cell but not the normal cell. Conversely, a band
5 that is absent in one or more of the cancer cell lines and present in a normal cell may represent a tumor suppressor gene which is not being expressed because of a mutation or regulatory defect.

The cDNA of a differentially expressed mRNA was
10 eluted from the sequencing gel and reamplified in a PCR reaction with the same primers as was used in the differential display reaction and cloned into a pCR™ II vector (Invitrogen, San Diego, CA). Specific mRNAs that were present solely in chondrosarcoma cells were
15 identified and the corresponding cDNAs cloned. cDNAs were sequenced using the M13 forward and reverse sequencing primers, which flank the cloning site of the pCR™ vector. Some sequences were identical to known genes, e.g., cyclin D2, a cell cycle regulatory protein,
20 and PTX3, a member of the pentaxin gene family. Novel cDNAs, i.e., those without sequence similarity to known genes, were used for Northern blotting to confirm that the corresponding gene is differentially expressed in chondrosarcoma cells. Twenty such cDNA probes were used
25 to screen for differential expression of mRNA and sequenced (TABLE 1). A novel gene, *csa-1*, was found to be differentially expressed in chondrosarcoma cells compared to normal cartilage cells and other cell types such as breast, lung, and colon cells.

- 16 -

TABLE 1

	Clone	Gel Type	MW(I)	FS(II)	AQ(III)	NI	Cart
	FS1	DD	-	+	-	-	-
		N3-0	-	+	+	-	-
5		N3-1	-	+	+	+	+
	FS2	DD	-	+	-	-	-
	E1	DD	-	-	-	-	+
		N3-2	-	+	+	+	+
		N3-3	-	+	+	+	-
10	FS3	DD	+	+	+	+	-
	AQ1	DD	-	-	+	-	-
	E2	DD	-	-	-	-	+
	AQ3	DD	-	-	+	-	-
	E6	DD	-	-	-	-	+
15	AQ2	DD	-	-	+	-	-
	E7	DD	-	-	-	-	+
	FS10	DD	+	+	+	+	-
	FS11	DD	+	+	+	+	-
	AQ6	DD	-	-	+	+	+
20	FS8	DD	+	+	-	-	+
	MW1	DD	+	-	-	-	-
	MW2	DD	+	-	-	-	-
	MW3	DD	+	-	-	-	-
	FS9	DD	-	+	+	+	+
25	FS10	DD	-	+	-	-	-
	AQ5	DD	-	+	+	+	+

DD: Differential Display Gel

N: Northern Blot

- 17 -

Cloning of *csa-1*

The gene encoding CSA-1 was identified using differential display PCR as described above. FS8 is a probe corresponding to one of the differentially
5 expressed sequences identified. As shown in TABLE 1, the differential display gel from which probe FS8 was isolated indicated that this gene was not expressed in the AQ cell line. In a Northern blot assay, the probe hybridized to a message approximately 0.85 kb in size in
10 the FS cell line as well as in a high grade chondrosarcoma. No message was detected in normal cartilage, bovine growth plate, or grade 1 or 2 chondrosarcoma.

The FS8 probe (specific for CSA-1) which
15 corresponded to the 3' end of the *csa-1* gene was found to be 250 bases long. 5' Rapid Amplification of cDNA (5' RACE) was used to clone the full length gene. A gene specific primer was synthesized which is complementary to the probe was made and used as a primer to synthesize
20 cDNA using RNA from the FS cell line as a template. The RNA was digested away with Rnase H, and an anchor primer was added to the 3' end with TdT and dCTP. PCR was performed using the 3' anchor primer and a second, nested gene specific primer, thereby yielding double stranded
25 DNA which is an extension of the gene fragment from which the differential display probe was derived. The 5' RACE generated fragment was cloned and sequenced.

Expression of CSA-1 in chondrosarcoma cells was localized to the nucleus of the cells by immunostaining
30 using a rabbit polyclonal antibody specific for a CSA-1 polypeptide.

The sequence of the full length *csa-1* cDNA (TABLE 2) was found to have an open reading frame (ORF) (TABLE 3 and shown in bold in TABLE 2) encoding a fifty-two amino
35 acid gene product, CSA-1 (TABLE 4).

- 18 -

TABLE 2: CSA-1 cDNA

ACTTCCCTGGGTTACACAGCAGGGGTGGAAGCTGGATTCTTCCTGGATGGGGATCCAGA
 TGGAGGTGGAGCTGCACCCCTTGTAGAGAATGGCTGCGGGTCCCAGGCCAGGAGCTC
 CCTGCAGGGCGGGGGCTCCCACGATCGTATTGACCTCTGGAAGAAGACAGACACTTT
 5 CCCACGGGAGCTCCTCTCCAGCCAGAGCTACACTTGGCAAACCTTTGGTCCTAAATG
 ATTATTCACTGAATTGAAGAAATACGGTTTACATATCTTCCAAGTATATATGTAGGG
 TTGATTTGGGAAGCAGAACACAGCAGCCCAAATTTGCTTGTAATGTCTGCGACTACA
 GCCTGCTGGCCTGCCTTC
 ACTGTCTTGGGGGAAGCTCGGGGAGACCAGGTGGACTGGAGTAGACTGTGCAGAGAC
 10 ACTGGTCTGGTGAAGATGTCCAGGAAACCACGAGCCTCCAGCCCATTTTCCAACAAC
 CACCCATCAACACCAAAGAGGTTCCAAGACAACCCAGAAGGGAAAAGGGACCCGTC
 AAGGAAGTTCCAGGAACAAAAGGCTCTCCCTAAAAGACCACCGCTTCAAAAAAACCT
 GAGGAATGGAGTGGGCCAACACTATCCAGCCACTCTGACCAGCCGAACGAGGAACTC
 AATCAAAATGCGCCATAGCAGGACCACAAGGGCAAGGAGACCACCGCCTTCTCCAGT
 15 GCTTCCTTGGGCAGCCAGTAATCCCAGGCAAGGCCAGAGACTTCAAGTCTATCTGA
 AAAGTCTCCAGAAGTCTAACCCAGATAAATAGCCAACAGGGTGTAGAGTACGTTTT
 ACACCCAAAGGGTAATGCCCATGGTGATGGAAATAAAATGAACATGTTGTAAAATG
 AAAAAAAAAA (SEQ ID NO:3)

TABLE 3: CSA-1 coding sequence

20 ATGGCTGCGGGTCCCAGGCCAGGAGCTCCCTGCAGGGCGGGGGCTCCCACGATCGTA
 TTGACCTCTGGAAGAAGACAGACACTTTCCCACGGGAGCTCCTCTCCAGCCAGAGCT
 AACTTGGCAAACCTTTGGTCCTAAATGATTATTCACTGAATTGAAGAAA (SEQ
 ID NO:1)

TABLE 4: CSA-1 AMINO ACID SEQUENCE

25 MAAGPRPGAPCRAGAPTIVLTSGRRQTL SHGSSSPARATLGKPLVLNDYSLN (SEQ
 ID NO:2)

Cloning of caa-1

The gene encoding CAA-1 was also identified using
 differential display PCR as described above. E1 is a
 30 probe corresponding to one of the differentially
 expressed sequences identified. As shown in TABLE 1, E1

- 19 -

was expressed in normal cartilage, but not in any of the human chondrosarcoma cell lines tested by differential display PCR. A northern blot with probe E1 showed expression of a 2.2 kb message in normal articular
5 cartilage and the FS cell line. In contrast, 2 alternative sized transcripts (7.5 kb and 1.2 kb) were detected in the high grade cell line AQ. These data indicate that the *caa-1* gene may be alternatively spliced or rearranged in the AQ cell line. The pattern of
10 expression indicates that CAA-1 functions as a tumor suppressor gene.

Expression of CAA-1 expression was analyzed using Northern blotting and RT-PCR in human chondrosarcoma cell lines osteoblasts, normal cartilage, muscle, primary
15 chondrosarcoma and a panel of tumors and normal tissues. Northern blotting was performed with E1 and with a 1.5 kb 5' RACE generated portion of the gene as probes. PCR was performed using primers which amplify 306 bp of the *caa* gene.

20 Differential display of mRNA showed expression of a gene in normal cartilage and in the FS and AQ cell lines (but not in the MW cell line). The differential display probe (E1) was sequenced, found to be novel, and used for Northern blot analysis, which revealed an
25 estimated message size of 2.2 kb in normal cartilage and FS, but not MW, osteoblast, muscle or bovine growth plate.

5' RACE was performed using oligonucleotide primers complementary to the 5' end of clone E1 and
30 yielded a 1.5 kb fragment. In order to obtain the remaining 5' portion of the gene, 5' RACE was repeated with new 5' primers and an additional 0.5 kb fragment was cloned, and the overlapping gene fragments were sequenced.

- 20 -

Northern blotting was performed with the 1.5 kb fragment and a 2.0 kb message was detected in four different samples of FS RNA and in a grade II chondrosarcoma (CS) and FS RNA. The full length gene is 1955 nucleotides in length, which correlates with the message size seen with Northern blotting.

TABLE 5: CAA-1 cDNA

CACGCAAAGCAGTGTGGGTTGATTCTGAGGTGCACTGTGGGAAAGAGCTTGTCGCTG
 CGGTGTTGCTGTTGGAGACTCGATTGTTGGTGACAGCGAAAGAACGATAACAAAATG
 10 CCGGAGCGAGATAGTGAGCCGTTCTCCAACCCTTTGGCCCCCGATGGCCACGATGTG
 GATGATCCTCACTCCTTCCACCAATCAAACTCACCAATGAAGACTTCAGGAAANTN
 NTCATGACCCCCAGGGNTGCACNTACNTNTGCACCACNTTNTAANTNNNNTCACCAT
 GAGATGCCAAGGGAGTACAATGAGGATGAAGACCCAGCTGCACGAAGGAGGAAAAAG
 AAAAGTTATTATGCCAAGCTACGCCAACAAGAAATTGAGAGAGAGAGAGAGCTAGCA
 15 GAGAAGTACCGGGATCGTGCCAAGGAACGGAGAGATGGAGTGAACAAAGATTATGAA
 GAAACCGAGCTTATCAGCACCACAGCTAACTATAGGGCTGTTGGCCCCACTGCTGAG
 GCGGACAAATCAGCTRCAGNNRAGAGAAGACANWINDAHCNAGGAGTCCAAATTCTTG
 GGTGGTGACATGGAACACACCCATTTGGTGAAAGGCTTGGATTTTGNTNTGCTTCHN
 AANGTNCGAGCTGAGATTGNCMSCMNANARAAANARGAARANGNNCTGATGGNAAAN
 20 CCCMGAAAGAAACCAAGAAAGATGAGGATCCTGAAAATAAAATTGAATTTAAAACA
 CGTCTGGGCCGCAATGTTTACCGAATGCTTTTTAAGAGCAAAGCATATGAGCGGAAT
 GAGTTGTTCCCTGCCGGGCCGCATGGCCTATGTGGTAGACCTGGATGATGAGTATGCT
 GACACAGATATCCCCACCACTCTTATCCCGCAGCAAGGCTGATTGCCCCACCATGGA
 GGCCAGACCACACTGACCACAAATGACATTGTCATTAGCAAGCTGACCCAGATCCT
 25 TTCATACCTGAGGCAGGGAACCCGTAACAAGAAGCTTAAGAAGAAGGATAAAGGGAA
 GCCGGAAGAGAAGAAACCTCCTGAGGCTGACATGAATATTTTTGAAGACATTGGGGA
 TTACGTACCCTCCACAACCAAGACACCTCGGGACAAGGAGCGGGAGAGATATCGGGA
 ACGGGAGCGTGATCGGGAAAGAGACAGAGACCGTGACCGAGAGCGAGAGCGAGAACG
 AGATCGGGAACGAGAGCGAGAGCGGGACCGAGAGAGAGAAGAGGAAAAGAAGAGACA
 30 CAGCTACTTTGAGAAGCCAAAAGTAGATGATGAGCCCATGGACGTTGACAAAGGACC
 TGGGTCTACCAAGGAGTTGATCAAGTCCATCAATGAAAAGTTTGCTGGGTCTGCTGG
 CTGGGAAGGCACAGAATCGCTGAAGAAGCCAGAAGACAAAAGCAGCTGGGAGATTT
 CTTTGGCATGTCCAACAGTTATGCAGAGTGCTACCCAGCCACGATGGATGACATGGC
 TGTGGATAGTGATGAGGAGGTGGATTATAGCAAATGGACCAGGGTAACAAGAAGGG
 35 GCCCTTAGGCCGTTGGGACTTTGATACCCAGGAAGAATACAGCGAGTATATGAACAA

- 21 -

CAAAGAAGCTTTGCCCAAGGCTGCATTCCAGTATGGTATCAAAATGTCTGAAGGGCG
 GAAAACCAGGCGCTTCAAGGAAACCAATGACAAAGCAGAGCTTGATCGCCAGTGGAA
 GAAGATTAGTGCAATCATTGANGAAGAGGAAGAAGATGGAAGCTGATGGGGTTGAAG
 TCAAAGACCAAATACTAATCACTAGTTACAACCAGAGATGCTCCACAAGGATATG
 5 CTCCCCACTGTTTTCTTTCTACAATTTCCAAAGGTTGCAAGATGTTTTTTTGTGATG
 AATATAAAATTTTATTGTGTAATTACTTGGTTCATTAAAATTGGTTAACTTGCTAA
 AAAAAAA (SEQ ID NO:5)

TABLE 6: CAA-1 Coding Sequence

ATGATGAGTATGCTGACACAGATATCCCCACCACTCTTATCCCGCAGCAAGGCTGAT
 10 TGCCCCACCATGGAGGCCAGACCACACTGACCACAAATGACATTGTCATTAGCAAG
 CTGACCCAGATCCTTTCATACCTGAGGCAGGGAACCCGTAACAAGAAGCTTAAGAAG
 AAGGATAAAGGGAAGCCGGAAGAGAAGAAACCTCCTGAGGCTGACATGAATATTTTT
 GAAGACATTGGGGATTACGTACCCTCCACAACCAAGACACCTCGGGACAAGGAGCGG
 GAGAGATATCGGGAACGGGAGCGTGATCGGGAAAGAGACAGAGACCGTGACCGAGAG
 15 CGAGAGCGAGAACGAGATCGGGAACGAGAGCGAGAGCGGGACCGAGAGAGAGAAGAG
 GAAAAGAAGAGACACAGCTACTTTGAGAAGCCAAAAGTAGATGATGAGCCCATGGAC
 GTTGACAAAGGACCTGGGTCTACCAAGGAGTTGATCAAGTCCATCAATGAAAAGTTT
 GCTGGGTCTGCTGGCTGGGAAGGCACAGAATCGCTGAAGAAGCCAGAAGACAAAAAG
 CAGCTGGGAGATTTCTTTGGCATGTCCAACAGTTATGCAGAGTGCTACCCAGCCACG
 20 ATGGATGACATGGCTGTGGATAGTGATGAGGAGGTGGATTATAGCAAAATGGACCAG
 GGTAACAAGAAGGGGCCCTTAGGCCGTTGGGACTTTGATACCCAGGAAGAATACAGC
 GAGTATATGAACAACAAAGAAGCTTTGCCCAAGGCTGCATTCCAGTATGGTATCAAA
 ATGTCTGAAGGGCGGAAAACCAGGCGCTTCAAGGAAACCAATGACAAAGCAGAGCTT
 GATCGCCAGTGGAAGAAGATTAGTGCAATCATTGANGAAGAGGAAGAAGATGGAAGC
 25 TGA (SEQ ID NO:6)

- 22 -

TABLE 7: CAA-1 Amino Acid Sequence

Met Met Ser Met Leu Thr Gln Ile Ser Pro Pro Leu Leu Ser
 Arg Ser Lys Ala Asp Cys Pro Thr Met Glu Ala Gln Thr Thr
 Leu Thr Thr Asn Asp Ile Val Ile Ser Lys Leu Thr Gln Ile
 5 Leu Ser Tyr Leu Arg Gln Gly Thr Arg Asn Lys Lys Leu Lys
 Lys Lys Asp Lys Gly Lys Pro Glu Glu Lys Lys Pro Pro Glu
 Ala Asp Met Asn Ile Phe Glu Asp Ile Gly Asp Tyr Val Pro
 Ser Thr Thr Lys Thr Pro Arg Asp Lys Glu Arg Glu Arg Tyr
 Arg Glu Arg Glu Arg Asp Arg Glu Arg Asp Arg Asp Arg Asp
 10 Arg Glu Arg Glu Arg Glu Arg Asp Arg Glu Arg Glu Arg Glu
 Arg Asp Arg Glu Arg Glu Glu Glu Lys Lys Arg His Ser Tyr
 Phe Glu Lys Pro Lys Val Asp Asp Glu Pro Met Asp Val Asp
 Lys Gly Pro Gly Ser Thr Lys Glu Leu Ile Lys Ser Ile Asn
 Glu Lys Phe Ala Gly Ser Ala Gly Trp Glu Gly Thr Glu Ser
 15 Leu Lys Lys Pro Glu Asp Lys Lys Gln Leu Gly Asp Phe Phe
 Gly Met Ser Asn Ser Tyr Ala Glu Cys Tyr Pro Ala Thr Met
 Asp Asp Met Ala Val Asp Ser Asp Glu Glu Val Asp Tyr Ser
 Lys Met Asp Gln Gly Asn Lys Lys Gly Pro Leu Gly Arg Trp
 Asp Phe Asp Thr Gln Glu Glu Tyr Ser Glu Tyr Met Asn Asn
 20 Lys Glu Ala Leu Pro Lys Ala Ala Phe Gln Tyr Gly Ile Lys
 Met Ser Glu Gly Arg Lys Thr Arg Arg Phe Lys Glu Thr Asn
 Asp Lys Ala Glu Leu Asp Arg Gln Trp Lys Lys Ile Ser Ala
 Ile Ile Xaa Glu Glu Glu Glu Asp Gly Ser (SEQ ID NO:7)

The longest predicted open reading frame (ORF) is
 25 942 bp. This ORF begins with the second in frame start
 codon, and is preceded by a shorter, 126 bp ORF. The
 predicted protein for the long ORF is 314 amino acids
 with a molecular weight of 37 kDa. The estimated message
 was 2.1 kb, but only 756 bp of sequence was reported,
 30 which was the largest clone isolated from their cDNA
 library.

Expression of CAA-1 has been detected with PCR in
 4/5 normal cartilage specimens, 1/2 grade 0, 3/3 grade I,
 3/4 grade II, and 5/5 grade III chondrosarcoma.

- 23 -

Expression was not detected in colon, breast, renal cell, and gastric carcinoma and corresponding normal tissues; osteogenic and soft tissue sarcoma; and giant cell tumor.

CAA-1 functions to regulate an immune response.

5 Regulation of an immune response, e.g., inflammation, is critical for normal synovial joint physiology and tumor surveillance. HLA class II antigens are necessary for antigen presentation to T cells, and interferon gamma has been shown to upregulate HLA class II expression in many
10 different normal and tumor cells, including chondrocytes and synovial lining cells. In addition, chondrocytes have been shown to function as antigen presenting cells. CAA-1 functions as a cytokine which inhibits the interferon gamma induced upregulation of HLA class II
15 antigens. Thus, chondrocytes express a gene which modulates its own ability, as well as cells in surrounding synovium, to function as antigen presenting cells. Treating a synovial joint with a CAA-1 polypeptide decreases the expression of HLA II antigens.
20 Thus, a CAA-1 polypeptide can be administered locally to reduce pathological such as that associated with rheumatoid arthritis and other inflammatory arthropathies.

CAA-1 is also expressed in neoplastic cartilage.
25 CAA-1 inhibits interferon gamma induced upregulation of HLA Class II. Expression of CAA-1 by tumor cells may be a mechanism of escape from immunorecognition, i.e. increased CAA-1 expression diminishes the ability of the host to control tumor growth through immunologic
30 mechanisms. Treatment of chondrosarcoma by inhibiting the expression of CAA-1 or function of the CAA-1 gene product enhances the ability of the host immune system to control tumor growth through immunologic mechanisms.

- 24 -

Example 2: Production and Purification of Recombinant
CSA and CAA Polypeptides

To produce recombinant polypeptides, DNA encoding
a CSA or CAA polypeptide in an appropriate expression
5 vector is transfected into a cell. Standard methods for
transfecting cells with isolated nucleic acid are well
known to those skilled in the art of molecular biology.
For example, prokaryotic or eukaryotic cells in culture
can be transfected with the DNA of the invention
10 operatively linked to expression control sequences
appropriate for high-level expression in the cell. Such
cells are useful for producing large amounts of the CSA-1
or CAA-1, which can be purified and used, e.g., as a
therapeutic or for raising anti-CSA-1 or anti-CAA-1
15 antibodies.

For example, the recombinant gene product may be
expressed as a fusion protein and purified using a
commercially available expression and purification
system, e.g., the pFLAG expression system (IBI).
20 Recombinant polypeptides are injected into a rabbit or
rodent to produce antibodies as described below.

Example 3: Production of Antibodies Specific for CSA or
CAA Polypeptides

Antibodies specific for CSA polypeptides can be
25 obtained by techniques well known in the art. Such
antibodies can be polyclonal or monoclonal. Polyclonal
antibodies can be obtained, for example, by the methods
described in Ghose et al., Methods in Enzymology, Vol.
93, 326-327, 1983. For example, a CSA-1 polypeptide
30 (containing 23-24 amino acids, e.g., a polypeptide
containing RRQTLSHGSSSPARAC (SEQ ID NO:8) was used as an
immunogen to stimulate the production of CSA-1-reactive
polyclonal antibodies in the antisera of a rabbit.
Similar methods can be used to raise antisera in animals
35 such as goats, sheep, and rodents.

- 25 -

Monoclonal antibodies useful in the present invention can be obtained by the well known process described by Milstein and Kohler in Nature, 256:495-97, 1975, or as modified by Gerhard, Monoclonal Antibodies, 5 Plenum Press, 1980, pages 370-371. Hybridomas are screened to identify those producing antibodies that are highly specific for a CSA polypeptide. Preferably, the antibody will have an affinity of at least about 10^8 liters/mole and more preferably, an affinity of at least 10 about 10^9 liters/mole. The use of such monoclonal antibodies provides a means of obtaining greater sensitivity in the assays of the present invention compared with the use of polyclonal antibodies.

Example 4: Transgenic animals

15 CSA polypeptides can also be expressed in transgenic animals. These animals represent a model system for the study of disorders that are caused by or exacerbated by overexpression or underexpression of a CSA polypeptide, and for the development of therapeutic 20 agents that modulate the expression or activity of a CSA polypeptide.

A CSA-1 knockout animal is useful to study CSA-1 function. Immunostaining of mouse embryos with anti-CSA-1 antibody showed staining of the musculoskeletal 25 precursor. A *csa-1* transgene with a null mutation results in an animal which does not express the CSA-1 gene, a condition which may lead to developmental abnormalities since some genes expressed by tumors are also expressed during normal embryological development.

30 Alternatively, a transgenic animal overexpressing the CSA-1 is useful to study the development chondrosarcoma. Such an animal would be a useful tool for evaluating treatment of this tumor.

Transgenic animals can be farm animals (pigs, 35 goats, sheep, cows, horses, rabbits, and the like)

- 26 -

rodents (such as rats, guinea pigs, and mice), non-human primates (for example, baboons, monkeys, and chimpanzees), and domestic animals (for example, dogs and cats). Transgenic mice are especially preferred.

5 Any technique known in the art can be used to introduce a *csa-1* transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear
10 microinjection (U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., *Proc. Natl. Acad. Sci., USA* 82:6148, 1985); gene targeting into embryonic stem cells (Thompson et al., *Cell* 56:313, 1989); and electroporation of embryos (Lo, *Mol. Cell. Biol.* 3:1803, 1983).

15 When it is desired that the *csa-1* transgene be integrated into the chromosomal site of the endogenous *csa-1* gene, gene targeting is preferred. Briefly, when such a technique is to be used, vectors containing some nucleotide sequences homologous to an endogenous *csa-1*
20 gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene also can be selectively introduced into a particular cell type, thus
25 inactivating the endogenous *csa-1* gene in only that cell type (Gu et al., *Science* 265:103, 1984). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the
30 art. These techniques are useful for preparing "knock outs" having no functional *csa* or *caa* gene.

Once transgenic animals have been generated, the expression of the recombinant transgene can be assayed utilizing standard techniques. Initial screening may be
35 accomplished by Southern blot analysis or PCR techniques

- 27 -

to determine whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR. Samples of *csa* or *caa* gene-expressing tissue can also be evaluated immunocytochemically using antibodies specific for the transgene product.

10 For a review of techniques that can be used to generate and assess transgenic animals, skilled artisans can consult Gordon (*Intl. Rev. Cytol.* 115:171-229, 1989), and may obtain additional guidance from, for example: Hogan et al. "Manipulating the Mouse Embryo" (Cold Spring Harbor Press, Cold Spring Harbor, NY, 1986; Krimpenfort et al., *Bio/Technology* 9:86, 1991; Palmiter et al., *Cell* 41:343, 1985; Kraemer et al., "Genetic Manipulation of the Early Mammalian Embryo," Cold Spring Harbor Press, Cold Spring Harbor, NY, 1985; Hammer et al., *Nature* 20 315:680, 1985; Purcel et al., *Science*, 244:1281, 1986; Wagner et al., U.S. Patent No. 5,175,385; and Krimpenfort et al., U.S. Patent No. 5,175,384.

Example 5: Diagnosis of chondrosarcoma

The invention includes a method of detecting cartilaginous neoplasms in a sample of patient-derived tissue. Detection of *csa-1* expression (by measuring gene transcripts or gene products) in a patient sample compared to a control sample or CSA-1 polypeptide would predict chondrogenesis indicative of, e.g., chondrosarcoma. The diagnostic method of the invention is carried out by measuring *csa* gene expression in a tissue, e.g, a biopsy, or in a bodily fluid, e.g., blood or plasma. Detection of expression and determination of the level of gene expression is measured using methods known in the art, e.g., *in situ* hybridization, Northern

- 28 -

blot analysis, or Western blot analysis using CSA-1-specific monoclonal or polyclonal antibodies. An increase in the level of *csa-1* expression per cell in the test sample of tissue compared to the level per cell in control tissue indicates the presence of a chondrosarcoma in the test sample. For example, tissue obtained at an biopsy could be tested for CSA-1 expression, e.g., the level of CSA-1 transcript or polypeptide. An increased level of CSA-1 transcript or polypeptide (compared to normal tissue) indicates a high probability of chondrosarcoma. For example, PCR was used to detect expression *csa-1* in 15 patient-derived chondrosarcoma biopsy samples. In contrast, no *csa-1* expression was detected in 3 patient-derived normal control samples.

The methods described above can also be used to determine the grade of a tumor. Northern blotting and quantitative PCR techniques are used to determine the level of expression of *csa* gene expression. For example, elevated CSA-1 expression correlates with a higher grade of tumor.

The diagnostic procedures described above are useful to identify patients in need of therapeutic intervention to reduce or prevent chondrosarcoma.

Example 6: Methods of Therapy

Patients with chondrosarcoma can be treated by administering CSA-1 antisense nucleic acids or ribozymes. Other malignant conditions, which are characterized by a increase in CSA-1 expression may be treated in a similar manner.

Antisense therapy is used to inhibit expression of proteins, e.g., CSA-1, involved in chondrogenesis, e.g., that associated with chondrosarcoma. For example, an antisense strand of *csa-1* (either RNA or DNA) is directly introduced into the cells in a form that is capable of binding to the mRNA transcripts. Alternatively, a

- 29 -

vector-containing sequence which, which once within the target cells is transcribed into the appropriate antisense mRNA, may be administered. Antisense nucleic acids which hybridize to mRNA can decrease or inhibit production of the polypeptide product encoded by a gene by associating with the normally single-stranded mRNA transcript, thereby interfering with translation and thus, expression of the protein.

Ribozyme therapy can also be used to inhibit gene expression. Ribozymes bind to specific mRNA and then cut it at a predetermined cleavage point, thereby destroying the transcript. These RNA molecules may be used to inhibit expression of a *csa* gene involved in chondrogenesis associated with chondrosarcoma according to methods known in the art (Sullivan et al., 1994, J. Invest. Derm. 103:85S-89S; Czubayko et al., 1994, J. Biol. Chem. 269:21358-21363; Mahieu et al, 1994, Blood 84:3758-65; Kobayashi et al. 1994, Cancer Res. 54:1271-1275).

Another therapeutic approach to inhibiting the expression of proteins or polypeptides is the production of intracellularly expressed antibodies which, when expressed in a cell, bind to and prevent the transport and surface expression of target proteins. Intracellular antibodies may be expressed in a cell using known techniques (Chen et al., 1994, Hum. Gene Ther. 5:595-601).

Gene therapy may be carried out by administering to a patient a nucleic acid encoding a therapeutic polypeptide, e.g., a tumor suppressor gene such as *caa-1*, by standard vectors and/or gene delivery systems. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses,

- 30 -

adenoviruses and adeno-associated viruses, among others.

As is discussed above, undesired or pathological inflammation such as that associated with rheumatoid arthritis and other inflammatory arthropathies can be
5 treated by inhibiting CAA-1 expression. Antisense therapy and ribozyme therapy can be used to inhibit CAA-1 expression, and intracellular immunization using DNA encoding an anti-CAA-1 antibody can be used to inhibit function of the gene product.

10 A therapeutic composition may include one or more compounds, e.g., nucleic acids or immunosuppressive agents, and a pharmaceutically acceptable carrier. The therapeutic composition may also include a gene delivery system as described above. Pharmaceutically acceptable
15 carriers are biologically compatible vehicles which are suitable for administration to an animal: e.g., physiological saline. A therapeutically effective amount of a compound is an amount which is capable of producing a medically desirable result in a treated animal, e.g.,
20 inhibition of expression of a target gene, e.g., a cell surface or secreted protein, or inhibition of cell activity, e.g., proliferation, migration, antigen presentation, antibody production, or cytokine production.

25 Parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal delivery routes, may be used to deliver the compound, with intravenous administration being the preferred route. Dosages for any one patient depends upon many factors,
30 including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Dosages of the compound to be administered will vary (doses of immunosuppressive
35 agents are expected to be in the range of doses used for

- 31 -

administration of other immunosuppressive agents known in the art). A preferred dosage for intravenous administration of nucleic acids is from approximately 10^6 to 10^{22} copies of the nucleic acid molecule.

5 Alternatively, the compound may be administered via a timed-release implant placed in close proximity to diseased tissue or a surgical site after removal of neoplastic tissue.

CSA polypeptides or CAA polypeptides, e.g., a
10 CAA-1 polypeptide, may be administered to the patient intravenously in a pharmaceutically acceptable carrier such as physiological saline. Standard methods for intracellular delivery of peptides can be used, e.g. packaged in liposomes. Such methods are well known to
15 those of ordinary skill in the art. It is expected that an intravenous dosage of approximately 1 to 100 μ moles of the polypeptide of the invention would be administered per kg of body weight per day. The compositions of the invention are useful for parenteral administration, such
20 as intravenous, subcutaneous, intramuscular, and intraperitoneal. Alternatively, a CAA polypeptide, e.g., CAA-1, can be administered as an implant for slow release at the site of an inflammatory lesion.

DNA (*csa-1* encoding DNA, tumor cell-specific
25 promoters, and vectors) of the invention may be introduced into target cells of the patient by standard vectors and/or gene delivery systems. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as
30 herpes viruses, retroviruses, and adenoviruses, among others. For example, the DNA of the invention under the control of a strong constitutive promoter may be administered locally using an adenovirus delivery system.

The DNA of the invention may be administered in a
35 pharmaceutically acceptable carrier. The therapeutic

- 32 -

composition may also include a gene delivery system as described above. Pharmaceutically acceptable carriers are biologically compatible vehicles which are suitable for administration to an animal e.g., physiological saline. A therapeutically effective amount is an amount of the nucleic acid of the invention which is capable of producing a medically desirable result in a treated animal.

As is well known in the medical arts, dosage for any given patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Dosages for the compounds of the invention will vary, but a preferred dosage for intravenous administration is from approximately 10^6 to 10^{22} copies of the nucleic acid molecule. Determination of optimal dosage is well within the abilities of a pharmacologist of ordinary skill. Drugs which inhibit the CSA-1 promoter may also be administered as described above to decrease the level of expression CSA-1 in tissues.

Example 7: Identification of Compounds that Decrease *csa* or *caa* Gene Expression

A method of screening candidate compounds to identify compounds capable of inhibiting *csa* gene, e.g. *csa-1*, expression includes the following steps: providing a chondrosarcoma cell; contacting the cell with a candidate compound; and determining the amount of *csa-1* expression in the cell, e.g., by immunostaining to detect a CSA-1 polypeptide or *in situ* hybridization, PCR, or Northern blotting to detect *csa-1* transcripts. A decrease in the amount of *csa-1* expression in cells exposed to the candidate compound compared to the amount of expression in cells in the absence of compound

- 33 -

indicates that the compound inhibits expression of *csa-1* in chondrosarcoma cells.

Compounds that inhibit *csa-1* expression can also be identified by contacting the *csa-1* promoter linked to a reporter gene with a candidate compound and measuring the level of expression of the reporter gene in the presence and absence of the compound. An decreased level of expression in the presence of the compound compared to that in its absence indicates that the compound inhibits expression of *csa-1*.

The screening methods described above can also be used to identify compounds which inhibit expression of a *caa* gene such as *caa-1*.

Other embodiments are within the following claims.

WO 99/46382

PCT/US99/05348

SEQUENCE LISTING

<110> Terek, Richard M.

<120> CHONDROSARCOMA ASSOCIATED GENES

<130> 04930/021001

<140> US 09/042,225

<141> 1998-03-13

<160> 8

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 164

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)...(156)

<400> 1

atg	gct	gcg	ggt	ccc	agg	cca	gga	gct	ccc	tgc	agg	gcg	ggg	gct	ccc	48
Met	Ala	Ala	Gly	Pro	Arg	Pro	Gly	Ala	Pro	Cys	Arg	Ala	Gly	Ala	Pro	
1				5					10					15		

acg	atc	gta	ttg	acc	tct	gga	aga	aga	cag	aca	ctt	tcc	cac	ggg	agc	96
Thr	Ile	Val	Leu	Thr	Ser	Gly	Arg	Arg	Gln	Thr	Leu	Ser	His	Gly	Ser	
			20					25					30			

tcc	tct	cca	gcc	aga	gct	aca	ctt	ggc	aaa	cct	ttg	gtc	cta	aat	gat	144
Ser	Ser	Pro	Ala	Arg	Ala	Thr	Leu	Gly	Lys	Pro	Leu	Val	Leu	Asn	Asp	
		35					40					45				

tat	tca	ctg	aat	tgaagaaa												164
Tyr	Ser	Leu	Asn													
	50															

<210> 2

<211> 52

<212> PRT

<213> Homo sapiens

<400> 2

Met	Ala	Ala	Gly	Pro	Arg	Pro	Gly	Ala	Pro	Cys	Arg	Ala	Gly	Ala	Pro	
1				5					10					15		
Thr	Ile	Val	Leu	Thr	Ser	Gly	Arg	Arg	Gln	Thr	Leu	Ser	His	Gly	Ser	
			20					25					30			
Ser	Ser	Pro	Ala	Arg	Ala	Thr	Leu	Gly	Lys	Pro	Leu	Val	Leu	Asn	Asp	
		35					40					45				
Tyr	Ser	Leu	Asn													
	50															

<210> 3

<211> 884

<212> DNA

<213> Homo sapiens

WO 99/46382

PCT/US99/05348

<400> 3
 acttccctgg gttcacagca ggggtggaac tggattcttc ctggatgggg atccagatgg 60
 aggtggagct gcacccttg tagagaatgg ctgcggtcc caggccagga gctccctgca 120
 gggcgggggc tcccacgatc gtattgacct ctggaagaag acagacactt tcccacggga 180
 gctcctctcc agccagagct acaactggca aacctttggt cctaaatgat tattcactga 240
 attgaagaaa tacggtttac atatcttcca agtatatatg taggggtgat ttgggaagca 300
 gaacacagca gcccaaattt gcttgtaatg tctgcgacta cagcctgctg gcctgccttc 360
 actgtcttgg gggaaagctcg gggagaccag gtggactgga gtagactgtg cagagacact 420
 ggtctggtga agatgtccag gaaaccacga gcctccagcc cattttccaa caaccacca 480
 tcaacaccaa agaggttccc aagacaacc agaagggaaa agggaccctg caaggaagt 540
 ccaggaacaa aaggctctcc ctaaaagacc accgcttcaa aaaaacctga ggaatggagt 600
 gggccaacac tatccagcca ctctgaccag cogaacgagg aactcaatca aatgcgcca 660
 tagcaggacc acaagggcaa ggagaccacc gccttctcca gtgcttcctt gggcagccag 720
 taattcccag gcaaggccag agacttcaag tctatctgaa aagtctccag aagtctaacc 780
 ccagataaat agccaacagg gtgtagagta cgttttacac ccaaagggta atgccccatg 840
 gtgatggaaa taaatgaac atggtgtaaa atgaaaaaaaa aaaa 884

<210> 4
 <211> 14
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Artificial sequence

<221> misc feature
 <222> (1)...(14)
 <223> n = A,T,C or G

<400> 4
 tttttttttt ttnv 14

<210> 5
 <211> 1946
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc feature
 <222> (1)...(1946)
 <223> n = A,T,C or G

<400> 5
 cacgcaaagc agtgtgggtt gattctgagg tgcactgtgg gaaagagctt gtcgctgagg 60
 tgttgctgtt ggagactcga ttggtggtga cagcgaaga acgataacaa aatgccggag 120
 cgagatagt agccgttctc caaccctttg gccccgatg gccacgatgt ggatgatcct 180
 cactccttcc accaatcaaa actcaccaat gaagacttca ggaaantnnt catgaccccc 240
 agggntgcac ntacntntgc accacntntt aantnnntc accatgagat gccaagggag 300
 tacaatgagg atgaagaccc agctgcacga aggaggaaaa agaaaagtta ttatgccaag 360
 ctacgccaac aagaaattga gagagagaga gagctagcag agaagtaccg ggatcgtgcc 420
 aaggaacgga gagatggagt gaacaaagat tatgaagaaa ccgagcttat cagcaccaca 480
 gctaactata gggctggttg cccactgct gagggcgaca aatcagctrc agnnragaga 540
 agacanwnda hcnaggagtc caaattcttg ggtggtgaca tggaaacacac ccatttggtg 600
 aaaggcttgg attttgntnt gcttchnaan gtncgagctg agattgncms cmnanaraaa 660
 nargaarang nnetgatggn aaancccmg aaagaaacca agaaagatga ggatcctgaa 720
 aataaaattg aatttaaac acgtctgggc cgcaatgttt accgaatgct ttttaagagc 780
 aaagcatatg agcggaatga gttgttcttg ccgggcccga tggcctatgt ggtagacctg 840
 gatgatgagt atgctgacac agatatcccc accactctta tcccgcagca aggctgattg 900
 ccccaccatg gaggcccaga ccacactgac cacaatgac attgtcatta gcaagctgac 960
 ccagatcctt tcatacctga ggcagggaac ccgtaacaag aagcttaaga agaaggataa 1020
 agggaagccg gaagagaaga aacctcctga ggctgacatg aatatttttg aagacattgg 1080
 ggattacgta ccctccacaa ccaagacacc tcgggacaag gagcgggaga gatatcggga 1140

WO 99/46382

PCT/US99/05348

acgggagcgt	gatcgggaaa	gagacagaga	ccgtgaccga	gagcgagagc	gagaacgaga	1200
tcgggaacga	gagcgagagc	gggaccgaga	gagagaagag	gaaaagaaga	gacacagcta	1260
ctttgagaag	ccaaaagtag	atgatgagcc	catggacggt	gacaaaggac	ctgggtctac	1320
caaggagttg	atcaagtcca	tcaatgaaaa	gtttgctggg	tctgctggct	gggaaggcac	1380
agaatcgctg	aagaagccag	aagacaaaaa	gcagctggga	gatttctttg	gcatgtccaa	1440
cagttatgca	gagtgctacc	cagccacgat	ggatgacatg	gctgtggata	gtgatgagga	1500
ggtggattat	agcaaaatgg	accagggtaa	caagaagggg	cccttaggcc	gttgggactt	1560
tgatacccag	gaagaataca	gcgagtatat	gaacaacaaa	gaagctttgc	ccaaggctgc	1620
attccagtat	ggtatcaaaa	tgtctgaagg	gcggaaaacc	aggcgcttca	aggaaaccaa	1680
tgacaaagca	gagcttgatc	gccagtgga	gaagattagt	gcaatcattg	angaagagga	1740
agaagatgga	agctgatggg	gttgaagtca	aaagaccaa	ataactaatca	ctagttacaa	1800
ccagagatgc	tccacaagga	tatgctcccc	actgttttct	ttctacaatt	tccaaagggt	1860
gcaagatggt	tttttgtgat	gaatataaaa	ttttattgtg	taattacttg	gttccattaa	1920
aattggttaa	cttgctaaaa	aaaaaa				1946

<210> 6
 <211> 915
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 <222> (1)...(912)
 <221> misc_feature
 <222> (1)...(915)
 <223> n = A,T,C or G

<400> 6																
atg	atg	agt	atg	ctg	aca	cag	ata	tcc	cca	cca	ctc	tta	tcc	cgc	agc	48
Met	Met	Ser	Met	Leu	Thr	Gln	Ile	Ser	Pro	Pro	Leu	Leu	Ser	Arg	Ser	
1				5					10					15		
aag	gct	gat	tgc	ccc	acc	atg	gag	gcc	cag	acc	aca	ctg	acc	aca	aat	96
Lys	Ala	Asp	Cys	Pro	Thr	Met	Glu	Ala	Gln	Thr	Thr	Leu	Thr	Thr	Asn	
			20					25					30			
gac	att	gtc	att	agc	aag	ctg	acc	cag	atc	ctt	tca	tac	ctg	agg	cag	144
Asp	Ile	Val	Ile	Ser	Lys	Leu	Thr	Gln	Ile	Leu	Ser	Tyr	Leu	Arg	Gln	
		35					40					45				
gga	acc	cgt	aac	aag	aag	ctt	aag	aag	aag	gat	aaa	ggg	aag	ccg	gaa	192
Gly	Thr	Arg	Asn	Lys	Lys	Leu	Lys	Lys	Lys	Asp	Lys	Gly	Lys	Pro	Glu	
	50					55					60					
gag	aag	aaa	cct	cct	gag	gct	gac	atg	aat	att	ttt	gaa	gac	att	ggg	240
Glu	Lys	Lys	Pro	Pro	Glu	Ala	Asp	Met	Asn	Ile	Phe	Glu	Asp	Ile	Gly	
	65				70				75						80	
gat	tac	gta	ccc	tcc	aca	acc	aag	aca	cct	cgg	gac	aag	gag	cgg	gag	288
Asp	Tyr	Val	Pro	Ser	Thr	Thr	Lys	Thr	Pro	Arg	Asp	Lys	Glu	Arg	Glu	
				85					90					95		
aga	tat	cgg	gaa	cgg	gag	cgt	gat	cgg	gaa	aga	gac	aga	gac	cgt	gac	336
Arg	Tyr	Arg	Glu	Arg	Glu	Arg	Asp	Arg	Glu	Arg	Asp	Arg	Asp	Arg	Asp	
			100					105					110			
cga	gag	cga	gag	cga	gaa	cga	gat	cgg	gaa	cga	gag	cga	gag	cgg	gac	384
Arg	Glu	Arg	Glu	Arg	Glu	Arg	Asp	Arg	Glu	Arg	Glu	Arg	Glu	Arg	Asp	
		115					120					125				

- 34 -

What is claimed is:

1. An isolated nucleic acid molecule encoding a chondrosarcoma associated (CSA) polypeptide.
2. The nucleic acid molecule of claim 1, wherein
5 said polypeptide is human CSA-1.
3. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO:1 or a degenerate variant thereof.
4. The nucleic acid molecule of claim 1, wherein
10 said nucleic acid molecule comprises a nucleotide sequence encoding a polypeptide having a sequence that is at least 80% identical to the sequence of SEQ ID NO:2.
5. An isolated nucleic acid molecule comprising a strand which hybridizes at high stringency to a DNA
15 having the sequence of SEQ ID NO:1, or the complement thereof.
6. The nucleic acid molecule of claim 1, wherein said nucleic acid molecule is operably linked to regulatory sequences for expression of said polypeptide,
20 said regulatory sequences comprising a promoter.
7. A substantially pure CSA polypeptide.
8. The polypeptide of claim 7, wherein said polypeptide is a human CSA-1 polypeptide.
9. The polypeptide of claim 8, said polypeptide
25 comprising an amino acid sequence that is at least 50% identical to the amino acid sequence of SEQ ID NO:2.

- 35 -

10. The polypeptide of claim 9, wherein said polypeptide comprises the amino acid sequence of SEQ ID NO:2.

11. A cell comprising the nucleic acid molecule
5 of claim 1.

12. A method of making a CSA polypeptide, comprising (a) providing the cell of claim 11, and (b) culturing it under conditions permitting expression of said nucleic acid molecule.

10 13. A method for diagnosing the presence of a chondrosarcoma cell in a tissue sample, comprising measuring expression of a *csa* gene in said tissue sample and a control sample, wherein an increase in expression of said *csa* gene in said tissue sample compared to said
15 control sample indicates that said tissue sample contains a chondrosarcoma cell.

14. The method of claim 13, wherein said *csa* gene encodes a human CSA-1 polypeptide.

15. A method of grading a chondrosarcoma tumor in
20 a test sample of patient-derived tissue, comprising determining the level of *csa-1* gene expression in said test sample, and comparing it to the level of *csa-1* gene expression in a control sample, wherein the level of expression in said test sample compared to said control
25 sample is directly proportional to the grade of said tumor.

- 36 -

16. A method for detecting chondrosarcoma in a patient, comprising:

(a) administering to a patient a diagnostically effective amount of a detectably labeled
5 CSA-1-specific binding species, and

(b) determining whether said species specifically binds to cartilage cells of the patient, said binding being an indication of the presence of chondrosarcoma in the patient.

10 17. The method of claim 16, wherein the level of said binding correlates with the grade of said chondrosarcoma.

18. A method of detecting progressive chondrosarcoma in a patient, comprising

15 (a) successively administering to a patient suspected of having said chondrosarcoma a diagnostically effective amount of a detectably labeled CSA-1-specific binding species selected from the group consisting of antibodies, antibody fragments, and non-antibody CSA-1
20 binding compounds, and

(b) comparing the amount of said species that binds to cartilage cells of the patient in successive administrations to detect an increase of binding of said binding species over time, wherein said increase is an
25 indication of progressive chondrosarcoma in said patient.

- 37 -

19. A method of screening a candidate compound to identify a compound capable of inhibiting expression of a CSA polypeptide, comprising

- (a) providing a chondrosarcoma cell
5 expressing a CSA polypeptide;
- (b) contacting said cell with said candidate compound; and
- (c) determining the amount of expression of said CSA polypeptide by said cell, wherein a decrease in
10 said amount in the presence of said candidate compound compared to the amount in the absence of said candidate compound indicates that said candidate compound inhibits expression of said CSA polypeptide.

20. The method of claim 19, wherein said CSA
15 polypeptide is human CSA-1.