

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978**PUBLICATION PARTICULARS AND ABSTRACT**

(Section 32(3)(a) – Regulation 22(1)(g) and 31)

OFFICIAL APPLICATION NO.

LODGING DATE

ACCEPTANCE DATE

21	01	2007/07856	22	1 AUG 2005	43	30-5-2008
----	----	------------	----	------------	----	-----------

INTERNATIONAL CLASSIFICATION

NOT FOR PUBLICATION

51	C12N; C12P	CLASSIFIED BY: WIPO
----	------------	---------------------

FULL NAMES OF APPLICANT

71	KANEKA CORPORATION
----	--------------------

FULL NAMES OF INVENTORS

72	1. KIZAKI, NORIYUKI 2. YANO, MIHO 3. FUNAKI, MASAHIRO 4. TAKESUE, TERUAKI 5. YASOHARA, YOSHIHIKO 6. MORIKAWA, SOUICHI 7. NAKAI, TAKAHISA 8. KATAOKA, MICHIIKO 9. SHIMIZU, SAKAYU
----	--

EARLIEST PRIORITY CLAIMED

COUNTRY	NUMBER	DATE			
33	JP	31	2004-231226	32	6 AUG 2004

TITLE OF INVENTION

54	NOVEL CARBONYL REDUCTASE, GENE THEREOF AND METHOD OF USING THE SAME
----	---

57	ABSTRACT (NOT MORE THAT 150 WORDS)	NUMBER OF SHEETS	32
----	------------------------------------	------------------	----

If no classification is finished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

ABSTRACT

A polypeptide which is isolated from a microorganism belonging to the genus *Ogataea* and has an activity of asymmetrically reducing (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to form (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol; a DNA encoding this polypeptide; a transformant producing this polypeptide; and a method of producing (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol with the use of the above-described polypeptide or transformant. By using the polypeptide or the transformant as described above, an optically active alcohol such as (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol can be efficiently produced.

DESCRIPTION

NOVEL CARBONYL REDUCTASE, GENE THEREOF AND METHOD OF
USING THE SAME


Technical Field

5 【0001】

The present invention relates to a polypeptide (carbonylreductase) having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by the following formula

10 (1):


 【0002】



 【0003】

to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the following formula
15 (2):

 【0004】



 【0005】

20 , which is isolated from a microorganism having the activity, a DNA encoding the polypeptide, a vector containing the DNA, and a transformant transformed with the vector.

【0006】

The present invention also relates to a production method of an optically active alcohol, particularly (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the aforementioned formula (2),
5 using the polypeptide or the transformant.

【0007】

(2R,3S)-1-Chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol is a compound useful as a synthetic
10 starting material for a pharmaceutical agent and the like.

Background Art

【0008】

As the production method of (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, a method
15 comprising chemically reducing amino-protected (3S)-1-halo-3-amino-4-phenyl-2-butanone derivative with a reducing agent such as sodium borohydride and the like is known (patent reference 1, non-patent reference 1).
20 However, this method requires a comparatively expensive reducing agent, and the stereoselectivity thereof is not practically sufficient.

【0009】

As a method using a microorganism, moreover, a
25 method comprising allowing a microorganism belonging to the genus *Candida* and the like to act on a (3S)-1-halo-3-amino-4-phenyl-2-butanone derivative to produce a (2R,3S)-1-halo-3-amino-4-phenyl-2-butanol derivative (patent reference 2), and a method comprising bringing a
30 microorganism belonging to the genus *Rhodococcus* and the like in contact with (3S)-1-halo-2-oxo-3-protected amino-4-substituted butane to produce (2R,3S)-1-halo-2-hydroxy-3-protected amino-4-substituted butane (patent reference 3, non-patent reference 2) are known. By these
35 methods, however, the concentration of the product that

can be accumulated in the reaction mixture cannot be practically sufficient.

patent reference 1: JP-A-2-42048

patent reference 2: JP-A-9-285

5 patent reference 3: WO2002/014528

non-patent reference 1: Tetrahedron, 50, 6333 (1994)

non-patent reference 2: Tetrahedron:Asymmetry, 14, 3105 (2003)

Disclosure of the Invention

10 Problems to be Solved by the Invention

【0010】

In view of the above, the present invention aims at providing a polypeptide useful for the production of (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-
15 butanol, a DNA encoding the polypeptide, a vector containing the DNA, and a transformant transformed with the vector.

【0011】

In addition, the present invention aims at
20 providing an efficient production method of various optically active alcohols including (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, using the polypeptide or the transformant.

Means of Solving the Problems

25 【0012】

The present inventors have isolated a polypeptide having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-
30 phenyl-2-butanol from a microorganism having the activity. Moreover, they have succeeded in clarifying the base sequence of a DNA encoding the polypeptide and generating a transformant that highly produces the polypeptide utilizing the sequence. In addition, they
35 have found that various useful optically active alcohols

such as (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol can be efficiently produced by utilizing the polypeptide or the transformant, which resulted in the completion of the present invention.

5 **【0013】**

That is, the present invention is directed to a polypeptide having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol.

【0014】

Moreover, the present invention provides a DNA encoding the polypeptide. Furthermore, the present invention provides a vector containing the DNA. In addition, the present invention provides a transformant containing the vector. Further, the present invention provides a production method of optically active alcohols including (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, using the polypeptide or the transformant.

Effect of the Invention

【0015】

According to the present invention, a practical production method of useful optically active alcohols including (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol can be provided.

Brief Description of the Drawing

【0016】

Fig. 1 shows a production method and a structure of a recombinant vector pNTOM4G1.

Best Mode for Embodying the Invention

【0017】

The present invention is explained in detail in the following.

35 (3S)-1-Chloro-3-tert-butoxycarbonylamino-4-phenyl-

2-butanone described in the present specification can be prepared, for example, by the method disclosed in JP-A-62-126158 or JP-A-2-42048. In addition, unless otherwise specified, isolation of DNA, preparation of vector, gene
5 manipulation such as transformation described in the present specification can be performed by methods described in books such as Molecular Cloning 2nd Edition (Cold Spring Harbor Laboratory Press, 1989) and the like. Moreover, "%" used in the description of the
10 present specification means "%(w/v)" unless otherwise specified.

【0018】

The polypeptide of the present invention is a polypeptide having an activity to asymmetrically reduce
15 (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol. Such polypeptide can be isolated from a microorganism having the activity.

20 【0019】

While the microorganism to be the origin of the polypeptide of the present invention is not particularly limited, for example, a yeast belonging to the genus *Ogataea* can be used. Particularly preferred is *Ogataea*
25 *minuta* var. *minuta*, NBRC0975 strain. The microorganism can be obtained from the Incorporated Administrative Agency, National Institute of Technology and Evaluation, Biological Resource Center (NBRC: 2-5-8 Kazusakamatari, Kisarazu-shi 292-0818 Chiba, Japan).

30 【0020】

As a medium for cultivating a microorganism to be the origin of the polypeptide of the present invention, general liquid nutritional media including carbon sources, nitrogen sources, inorganic salts, organic
35 nutrients and the like can be used as long as the

microorganism grows.

【0021】

The polypeptide of the present invention can be isolated from the microorganism to be the origin of the polypeptide by appropriately combining generally known protein purification methods. For example, the isolation can be performed as follows.

【0022】

First, the microorganism is cultured in a suitable medium, and the cells are collected from the cultured medium by centrifugation or filtration. The obtained cells were disrupted by a physical means using an ultrasonic disintegrator, glass beads and the like, and the cellular debris was removed by centrifugation to give a cell-free extract. Then, the polypeptide of the present invention is isolated from the cell-free extract by a method such as salting-out (ammonium sulfate precipitation, sodium phosphate precipitation and the like), solvent precipitation (protein fractionation precipitation by acetone, ethanol and the like), dialysis, gel filtration chromatography, ion exchange chromatography, reversed-phase chromatography, ultrafiltration and the like, alone or in combination.

【0023】

The activity to reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone can be determined, for example, by adding a 0.2 mM substrate, (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone, a 0.25 mM coenzyme, NADPH and a crude enzyme to a 100 mM phosphate buffer (pH 6.5) containing 0.33% (v/v) dimethyl sulfoxide, reacting the mixture at 30°C for 1 min and calculating the activity from the decreasing rate of absorbance at a wavelength of 340 nm after the reaction.

35 【0024】

In addition, the determination of absolute configuration and measurement of diastereomeric excess of (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol produced by the above-mentioned reaction can be performed by high performance liquid chromatography (column: COSMOSIL 5C8-MS (ϕ 4.6 mm $\times$ 250 mm; manufactured by Nacalai Tesque), eluent: 10 mM aqueous phosphoric acid solution/acetonitrile=1/1 (v/v), flow rate: 1 ml/min, detection: 210 nm).

10 **【0025】**

The DNA of the present invention is a DNA encoding the aforementioned polypeptide of the present invention, and may be any as long as it can express the polypeptide in a host cell transformed according to the below-mentioned method, and may contain any untranslated region. Once the polypeptide can be acquired, those of ordinary skill in the art can obtain such DNA by a known method from a microorganism to be the origin of the polypeptide. For example, it can be obtained by the following method.

20 **【0026】**

First, an isolated polypeptide of the present invention is digested with a suitable endopeptidase, and the resulting peptide fragments are fractionated by reversed-phase HPLC. Then, for example, using an ABI492 protein sequencer (manufactured by Applied Biosystems), a part or the whole amino acid sequence of the peptide fragments are determined.

30 **【0027】**

Based on the information of the amino acid sequence obtained in this way, PCR (Polymerase Chain Reaction) primers for amplifying a part of a DNA encoding the polypeptide are synthesized. Then, a chromosomal DNA of a microorganism to be the origin of the polypeptide is prepared by a general DNA isolation method, for example,

the method of Visser and the like (Appl. Microbiol. Biotechnol., 53, 415 (2000)). Using the chromosomal DNA as a template and the aforementioned PCR primers, PCR is performed to amplify a part of the DNA encoding the polypeptide and determine the base sequence thereof. The base sequence can be determined, for example, using ABI373A DNA Sequencer (manufactured by Applied Biosystems) and the like.

【0028】

Once a part of the base sequence of the DNA encoding the polypeptide is clarified, the full-length sequence can be determined, for example, by i-PCR (Nucl. Acids Res., 16, 8186 (1988)).

【0029】

Examples of the DNA of the present invention obtained in this way include a DNA containing the base sequence shown in SEQ ID NO: 1 and a DNA containing the base sequence shown in SEQ ID NO: 2. In addition, a DNA that hybridizes to a DNA consisting of a base sequence complementary to the base sequence shown in SEQ ID NO: 1 or SEQ ID NO: 2 under stringent conditions, which encodes a polypeptide having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, is also encompassed in the DNA of the present invention.

【0030】

The DNA that hybridizes to a DNA consisting of a base sequence complementary to the base sequence shown in SEQ ID NO: 1 or SEQ ID NO: 2 under stringent conditions refers to a DNA that specifically forms a hybrid with a DNA having a base sequence complementary to the base sequence shown in SEQ ID NO: 1 or SEQ ID NO: 2, when colony hybridization method, plaque

hybridization method, Southern hybridization method and the like are performed.

【0031】

As used herein the stringent conditions refer to,
5 for example, the conditions under which hybridization in an aqueous solution of a composition of 75 mM trisodium citrate, 750 mM sodium chloride, 0.5% sodium dodecylsulfate, 0.1% bovine serum albumin, 0.1% polyvinylpyrrolidone and 0.1% Ficoll 400 (manufactured
10 by Amersham Biosciences) is performed at 65°C, followed by washing with an aqueous solution of a composition of 15 mM trisodium citrate, 150 mM sodium chloride, and 0.1% sodium dodecylsulfate at 60°C. Preferably, they refer to the conditions under which washing with an
15 aqueous solution of a composition of 15 mM trisodium citrate, 150 mM sodium chloride, and 0.1% sodium dodecylsulfate is performed at 65°C after hybridization under the above-mentioned conditions. More preferably, they refer to the conditions under which washing with an
20 aqueous solution of a composition of 1.5 mM trisodium citrate, 15 mM sodium chloride, and 0.1% sodium dodecylsulfate is performed at 65°C after the hybridization in the same manner as above.

【0032】

25 Examples of the polypeptide of the present invention include a polypeptide having the amino acid sequence shown in SEQ ID NO: 3 encoded by the base sequence shown in SEQ ID NO: 1 and a polypeptide having the amino acid sequence shown in SEQ ID NO: 4 encoded by
30 the base sequence shown in SEQ ID NO: 2.

【0033】

In addition, a polypeptide showing at least a certain level of homology to either of these two kinds of polypeptides and having an activity to asymmetrically
35 reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-

phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol is functionally equivalent to these two kinds of polypeptides and encompassed in the present invention.

5 **【0034】**

As used herein, the homology of the sequence is expressed, for example, by the value of Identity relative to the whole sequence when two amino acid sequences are comparatively analyzed using a homology
10 search program FASTA (W.R. Pearson & D.J. Lipman; Pro. Natl. Acad. Sci. USA, 85, 2444-2448 (1988)). Examples of the polypeptide equivalent to the polypeptide shown in SEQ ID NO: 3 or SEQ ID NO: 4 include a polypeptide showing not less than 78% (a homology between the
15 polypeptide shown in SEQ ID NO: 3 and the polypeptide shown in SEQ ID NO: 4 is 78%), preferably not less than 80%, more preferably not less than 85%, still more preferably not less than 90%, of homology to either of these two kinds of polypeptides.

20 **【0035】**

Such polypeptide can be obtained, for example, by ligating a DNA that can be hybridized to a DNA consisting of a base sequence complementary to the base sequence shown by the aforementioned SEQ ID NO: 1 or SEQ
25 ID NO: 2 under stringent conditions, into a suitable vector and introducing the vector into a suitable host cell to allow expression. Moreover, such polypeptide can be obtained, for example, by causing substitution, insertion, deletion or addition of amino acid in a
30 polypeptide having the amino acid sequence shown in SEQ ID NO: 3 or SEQ ID NO: 4, according to a known method described in Current Protocols in Molecular Biology (John Wiley and Sons, Inc., 1989) and the like.

【0036】

35 Examples of the polypeptide obtained by the latter

method include a polypeptide obtained by substituting the 42nd alanine with glycine, the 43rd glutamic acid with alanine, the 46th lysine with glutamic acid and/or the 49th asparagine with lysine in the amino acid sequence shown in SEQ ID NO: 4. A polypeptide having all these four mutations (mutant polypeptide 1) is shown in SEQ ID NO: 16, and a DNA encoding the polypeptide (mutant DNA 1) is shown in SEQ ID NO: 15.

【0037】

The vector to be used for introducing the DNA of the present invention into a host microorganism to allow expression of the DNA in the host microorganism is not particularly limited as long as the DNA can express the encoded gene in a suitable host microorganism. Examples of the vectors include plasmid vector, phage vector, cosmid vector and the like. In addition, a shuttle vector capable of gene exchange with other host cells can also be used.

【0038】

Such vector generally contains a regulator such as lacUV5 promoter, trp promoter, trc promoter, tac promoter, lpp promoter, tufB promoter, recA promoter, pL promoter and the like, and can be preferably used as an expression vector containing an expression unit operably linked to the DNA of the present invention. For example, pUCNT (WO94/03613) can be preferably used.

【0039】

The term "regulator" used in the present specification refers to a base sequence having a functional promoter and any related transcriptional element (e.g., enhancer, CCAAT box, TATA box, SPI moiety and the like).

【0040】

The term "operably linked" used in the present specification means that various regulatory elements

such as a promoter, an enhancer and the like, that control the gene expression, and a gene are connected in an operable state in a host cell. It is a well known matter to those of ordinary skill in the art that the
5 type and kind of regulator can vary depending on the host.

【0041】

Examples of the expression vector of the present invention include pNTOM3 wherein the DNA shown in SEQ ID
10 NO: 1 has been introduced into pUCNT, pNTOM4 wherein the DNA shown in SEQ ID NO: 2 has been introduced into pUCNT, pNTOM5 wherein the DNA shown in SEQ ID NO: 15 has been introduced into pUCNT, which are mentioned below, and the like.

15 【0042】

As the host cell, into which a vector containing the DNA of the present invention is introduced, bacterium, yeast, filamentous bacterium, plant cell, animal cell and the like can be used. Since introduction
20 and/or culture are/is easy, bacterium is preferable, and *Escherichia coli* is particularly preferable. The vector containing the DNA of the present invention can be introduced into a host cell by a known method. When *Escherichia coli* is used as a host cell, for example, a
25 commercially available *E. coli* HB101 competent cell (manufactured by TAKARA BIO) can be used to introduce the vector into the host cell.

【0043】

Examples of the transformant of the present invention include *E. coli* HB101 (pNTMO3) FERM BP-10368
30 wherein the aforementioned pNTMO3 has been introduced into *E. coli* HB101, *E. coli* HB101 (pNTOM4) FERM BP-10369 wherein the aforementioned pNTMO4 has been introduced into *E. coli* HB101, *E. coli* HB101 (pNTOM5) FERM BP-10370
35 wherein the aforementioned pNTMO5 has been introduced

into *E. coli* HB101, which are mentioned below, and the like. These transformants have been respectively deposited under the aforementioned accession numbers in the National Institute of Advanced Industrial Science and Technology, International Patent Organism Depositary (IPOD: Central 6, 1-1-1, Higashi, Tsukuba, Ibaraki, 305-8566 Japan) (date of original deposit in Japan on June 3, 2004, the national deposit strains were transferred to the international deposit based on Budapest Treaty on July 6, 2005).

【0044】

When a compound having a carbonyl group is asymmetrically reduced to produce an optically active alcohol using the polypeptide of the present invention, a coenzyme such as NADH, NADPH and the like becomes necessary. However, the amount of an expensive coenzyme to be used can be drastically reduced by carrying out the reaction in the co-presence of a polypeptide capable of converting the oxidized coenzyme to a reduced form (hereinafter to be referred to as a coenzyme regenerating capability) a compound to be a substrate of the polypeptide, and the polypeptide of the present invention.

【0045】

As the polypeptide having the coenzyme regenerating capability, for example, hydrogenase, formic acid dehydrogenase, alcohol dehydrogenase, aldehyde dehydrogenase, glucose-6-phosphate dehydrogenase, glucose dehydrogenase and the like can be used. Preferably, glucose dehydrogenase is used.

【0046】

The optically active alcohol can be produced using the polypeptide of the present invention as follows. First, a compound having a carbonyl group to be the substrate, a coenzyme such as NADPH and the like, and

the polypeptide are added to a suitable solvent, and the mixture is stirred under an adjusted pH to carry out the reaction. When the reaction is carried out using the polypeptide of the present invention and a polypeptide
5 having a coenzyme regenerating capability in combination, a polypeptide having a coenzyme regenerating capability (e.g., glucose dehydrogenase) and a compound (e.g., glucose) to be its substrate are further added to the above-mentioned reaction
10 composition.

【0047】

An aqueous solvent may be used for the reaction or a mixture of an aqueous solvent and an organic solvent may be used for the reaction. Examples of the organic
15 solvent include toluene, ethyl acetate, n-butyl acetate, hexane, isopropanol, diisopropyl ether, methanol, acetone, dimethyl sulfoxide and the like. The reaction is carried out at a temperature of 10°C - 70°C, and the pH of the reaction mixture is maintained at 4 - 10. The
20 reaction can be carried out by a batch process or a continuous process. For a batch process, the reaction substrate is added at a charge concentration of from 0.1% to 70% (w/v). Examples of the compound having a carbonyl group to be a substrate include (3S)-1-chloro-
25 3-tert-butoxycarbonylamino-4-phenyl-2-butanone. However, the compound is not particularly limited as long as it can be reduced under the aforementioned reaction conditions and converted to an optically active alcohol.

【0048】

30 The reaction can be carried out similarly even when a transformant containing a DNA encoding the polypeptide, or a treated product thereof is used instead of the polypeptide of the present invention. Moreover, the reaction can be carried out similarly even
35 when a transformant containing both a DNA encoding the

polypeptide of the present invention and a DNA encoding a polypeptide having a coenzyme regenerating capability, or a treated product thereof is used. The "treated product of transformant" means, for example, a cell
5 treated with a surfactant or an organic solvent, a dried cell, a disrupted cell, crude extract of a cell and the like, as well as those fixed by a known means.

【0049】

Particularly, when a transformant containing both a
10 DNA encoding the polypeptide of the present invention and a DNA encoding a polypeptide having a coenzyme regenerating capability or a treated product thereof is used, an enzyme for regenerating a coenzyme does not need to be separately prepared and added, and an
15 optically active alcohol can be produced efficiently.

【0050】

A transformant containing both a DNA encoding the polypeptide of the present invention and a DNA encoding a polypeptide having a coenzyme regenerating capability
20 can be obtained by incorporating both the DNA encoding the polypeptide of the present invention and the DNA encoding a polypeptide having a coenzyme regenerating capability into a single vector, and introducing the vector into a host cell, or respectively incorporating
25 these two kinds of DNAs into two kinds of vectors belonging to different incompatible groups, and introducing the two kinds of vectors into a single host cell.

【0051】

30 Examples of the vector incorporating both a DNA encoding the polypeptide of the present invention and a DNA encoding a polypeptide having a coenzyme regenerating capability include pNTOM3G1, pNTOM4G1 and pNTOM5G1, wherein a glucose dehydrogenase gene derived
35 from *Bacillus megaterium* has been introduced into each

of the aforementioned expression vectors pNTOM3, pNTOM4 and pNTOM5, and the like. Examples of the transformant containing both the DNA encoding the polypeptide of the present invention and the DNA encoding a polypeptide
5 having a coenzyme regenerating capability include *E. coli* HB101 (pNTOM3G1), *E. coli* HB101 (pNTOM4G1) and *E. coli* HB101 (pNTOM5G1), wherein *E. coli* HB101 has been transformed with these vectors, and the like.

【0052】

10 The activity of a polypeptide having a coenzyme regenerating capability in a transformant can be determined by a conventional method. For example, the activity of glucose dehydrogenase can be calculated from the rate of increase in the absorbance at a wavelength
15 of 340 nm, when 100 mM glucose, 2 mM coenzyme NADP or NAD, and the enzyme are added to 1M Tris-HCl buffer (pH 8.0), and the mixture is reacted at 25°C for 1 min.

【0053】

A transformant containing a DNA encoding the
20 polypeptide of the present invention can be cultured in a conventional liquid nutritional medium containing carbon sources, nitrogen sources, inorganic salts, organic nutrients and the like, as long as it grows.

【0054】

25 An optically active alcohol resulting from a reaction can be purified by a conventional method. For example, when the optically active alcohol resulting from a reaction is (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, the reaction
30 mixture is extracted with an organic solvent such as ethyl acetate, toluene and the like, and the organic solvent is evaporated under reduced pressure. Moreover, it can be purified by a treatment such as crystal precipitation, chromatography and the like.

35 【0055】

For quantitation of (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone and (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, as well as for measurement of diastereomeric excess of (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, high performance liquid chromatography (column: COSMOSIL 5C8-MS (ϕ 4.6 mm $\times$ 250 mm; manufactured by Nacalai Tesque), eluent: 10 mM aqueous phosphoric acid solution/acetonitrile=1/1, flow rate: 1 ml/min, detection: 210 nm) can be used.

【0056】

As mentioned above, according to the present invention, the polypeptide of the present invention can be efficiently produced and, utilizing the polypeptide, a superior production method of various useful optically active alcohols including (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol can be provided.

Examples

【0057】

The present invention is explained in detail in the following Examples, which are not to be construed as limitative. The detailed operation method and the like relating to the recombinant DNA technique used in the following Examples is described in the following books: Molecular Cloning 2nd Edition (Cold Spring Harbor Laboratory Press, 1989), Current Protocols in Molecular Biology (Greene Publishing Associates and Wiley-Interscience).

【0058】

30 (Example 1) purification of polypeptide

According to the following method, a polypeptide having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol was isolated and purified from the

Ogataea minuta var. *minuta* NBRC0975 strain. Unless otherwise specified, the purification was performed at 4°C.

【0059】

5 The reduction activity on (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone was determined by dissolving, in 100 mM phosphate buffer (pH 6.5) containing 0.33%(v/v) dimethyl sulfoxide, a substrate [(3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-
10 butanone] to a final concentration of 0.2 mM and a coenzyme NADPH to a final concentration of 0.25 mM, further adding a crude enzyme solution, carrying out the reaction at 30°C for 1 min and calculating from the rate of decrease in the absorbance of the reaction mixture at
15 a wavelength of 340 nm. The activity to oxidize 1 μmol of NADPH to NADP for 1 min under said reaction conditions was defined to be 1 unit.

【0060】

(culture of microorganism)

20 A liquid medium (18L) (pH 6.5) containing glucose 5%, polypeptone 0.5%, yeast extract 0.1%, dipotassium hydrogen phosphate 0.1%, potassium dihydrogen phosphate 0.2% and magnesium sulfate heptahydrate 0.02%, was prepared in a 30L jar-fermentor (manufactured by B.E.
25 Marubishi Co., Ltd.) and sterilized with vapor at 120°C for 20 min.

【0061】

To this medium was inoculated 180 ml of a cultured medium of the *Ogataea minuta* var. *minuta* NBRC0975 strain
30 precultured in the same medium in advance, and the mixture was cultured under the conditions of stirring at 250 rpm, airflow amount 5.0 NL/min, 28°C for 48 hr.

【0062】

(preparation of cell-free extract)

35 The cells were collected from the above-mentioned

cultured medium by centrifugation and washed with 2000 ml of 10 mM Tris-HCl buffer (pH 7.5) to give 907 g of the cells of the strain. The cells were suspended in 1800 ml of 10mM Tris-HCl buffer (pH 7.5) containing 0.1 mM dithiothreitol (DTT), and disrupted using a UH-600 ultrasonication dispersion machine (manufactured by SMT). The cell debris was removed from the disrupt by centrifugation to give a cell-free extract.

【0063】

10 (ammonium sulfate fractionation)

The pH of the cell-free extract obtained above was adjusted to 7.5 with aqueous ammonia, ammonium sulfate was added and dissolved to 40% saturation while maintaining the pH, and the resulting precipitate was removed by centrifugation. Ammonium sulfate was further added to the supernatant and dissolved to 60% saturation while maintaining at pH 7.5 in the same manner as above, and the resulting precipitate was collected by centrifugation. The precipitate was dissolved in 10 mM Tris-HCl buffer (pH 7.5) containing 0.1 mM DTT, and dialyzed against the same buffer overnight to give an active fraction.

【0064】

(DEAE-Sephacel column chromatography)

25 The active fraction obtained by ammonium sulfate fractionation was applied to a DEAE-Sephacel (manufactured by Amersham Biosciences) column (ϕ 29x290 mm) equilibrated in advance with 10 mM Tris-HCl buffer (pH 7.5) containing 0.1 mM DTT to allow adsorption of the active fraction. After washing the column with the same buffer, the active fraction was eluted with the linear gradient (from 0 M to 1 M) of sodium chloride. The active fraction was collected and dialyzed against the same buffer overnight.

35 【0065】

(MonoQ HR column chromatography)

The active fraction obtained by DEAE-Sephacel column chromatography was applied to a MonoQ HR 10/10 column (manufactured by Amersham Biosciences) equilibrated in advance with 10 mM Tris-HCl buffer (pH 7.5) containing 0.1 mM DTT to allow adsorption of the active fraction. After washing the column with the same buffer, the active fraction was eluted with the linear gradient (from 0 M to 1 M) of sodium chloride. The active fraction was collected and dialyzed against the same buffer overnight.

【0066】

(Phenyl-Superose column chromatography)

Sodium chloride was dissolved in the active fraction obtained by MonoQ HR column chromatography to a final concentration of 4M and the solution was applied to a Phenyl-Superose HR 10/10 column (manufactured by Amersham Biosciences) equilibrated in advance with 10 mM Tris-HCl buffer (pH 7.5) containing 4 M sodium chloride and 0.1 mM DTT to allow adsorption of the active fraction. After washing the column with the same buffer, the active fraction was eluted with the linear gradient (from 4 M to 0 M) of sodium chloride. The active fraction was collected and dialyzed against 10 mM Tris-HCl buffer (pH 7.5) containing 0.1 mM DTT overnight.

【0067】

(MonoQ HR column chromatography)

The active fraction obtained by Phenyl-Superose column chromatography was applied to a MonoQ HR 5/5 column (manufactured by Amersham Biosciences) equilibrated in advance with 10 mM Tris-HCl buffer (pH 7.5) containing 0.1 mM DTT to allow adsorption of the active fraction. After washing the column with the same buffer, the active fraction was eluted with the linear gradient (from 0 M to 1 M) of sodium chloride. The

active fraction was collected and concentrated using Centricon YM-10 (manufactured by Millipore).

【0068】

(gel filtration chromatography)

5 The active fraction obtained by the aforementioned MonoQ HR column chromatography was applied to a Superdex 200 HR 10/30 column (manufactured by Amersham Biosciences) equilibrated in advance with 10 mM Tris-HCl buffer (pH 7.5) containing 0.2 M sodium chloride and 0.1
10 mM DTT and the active fraction was eluted at a flow rate of 0.5 ml/min using the same buffer. The active fraction was collected and concentrated using Centricon YM-10 (manufactured by Millipore). This was applied to a Superdex 200 HR 10/30 column (manufactured by Amersham
15 Biosciences) equilibrated in advance with the same buffer and the active fraction was eluted at a flow rate of 0.5 ml/min using the same buffer. The active fraction was collected and used as a purified standard sample of the polypeptide.

20 【0069】

(Example 2) gene cloning

(preparation of PCR primers)

 The purified polypeptide obtained in Example 1 was denatured in the presence of 8M urea and digested with
25 lysyl endopeptidase derived from Achromobacter (manufactured by Wako Pure Chemical Industries, Ltd.). Then, the amino acid sequence of the obtained peptide fragment was determined by ABI492 protein sequencer (manufactured by Perkin Elmer). Based on the DNA
30 sequence predicted from the amino acid sequence, primer 1: 5'-acngtntayttyathgcngg-3' (SEQ ID NO: 5) and primer 2: 5'-atnggdatrtrcraaytgrtc-3' (SEQ ID NO: 6) for amplifying a part of the gene encoding the polypeptide by PCR were synthesized.

35 【0070】

(gene amplification by PCR)

Chromosomal DNA was extracted from the cells of *Ogataea minuta* var. *minuta* NBRC0975 strain cultured in the same manner as in Example 1 according to the method of Visser (Appl. Microbiol. Biotechnol., 53, 415 (2000)) and the like. Using the DNA primers 1 and 2 prepared above and the obtained chromosomal DNA as a template, PCR was performed. As a result, about 700 bp DNA fragment considered to be a part of the object gene was amplified. PCR was performed using TaKaRa Ex Taq (manufactured by TAKARA BIO) as a DNA polymerase, where the reaction conditions followed the instruction manual thereof. The DNA fragment was cloned to a plasmid pT7Blue T-Vector (manufactured by Novagen), and the base sequence thereof was analyzed using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit (manufactured by Perkin Elmer) and ABI 373A DNA Sequencer (manufactured by Perkin Elmer). As a result of the PCR, it has been clarified that two kinds of DNA fragments having different base sequences were amplified. The base sequences thereof are shown in SEQ ID NO: 7 and SEQ ID NO: 8.

【0071】

(identification of full-length sequence of object gene by i-PCR)

The chromosomal DNA of *Ogataea minuta* var. *minuta* NBRC0975 strain prepared above was completely digested with a restriction enzyme XbaI, a mixture of the obtained DNA fragments was intramolecularly cyclized with T4 ligase. Using this as a template, the full-length base sequence of the gene containing the aforementioned base sequence shown in SEQ ID NO: 7 was determined by i-PCR (Nucl. Acids Res., 16, 8186 (1988)). The results are shown in SEQ ID NO: 1. i-PCR was performed using TaKaRa LA Taq (manufactured by TAKARA

BIO) as a DNA polymerase, where the reaction conditions followed the instruction manual thereof. The full-length base sequence of the gene containing the aforementioned base sequence shown in SEQ ID NO: 8 was also determined
5 by a similar operation, and the results thereof are shown in SEQ ID NO: 2. In addition, the amino acid sequence encoded by the base sequence shown in SEQ ID NO: 1 is shown in SEQ ID NO: 3, and the amino acid sequence encoded by the base sequence shown in SEQ ID
10 NO: 2 is shown in SEQ ID NO: 4.

【0072】

(Example 3) construction of expression vector

Using primer 3: 5'-gtgcatatgaccaagactgtttatttca-3' (SEQ ID NO: 9) and primer 4: 5'-
15 gtcgaattcttattataaaatggaatgtcgaa-3' (SEQ ID NO: 10) and the chromosomal DNA of the *Ogataea minuta* var. *minuta* NBRC0975 strain obtained in Example 2 as a template, PCR was performed. As a result, a double stranded DNA wherein an NdeI recognition site is added to the
20 initiation codon and an EcoRI recognition site is added immediately after the termination codon of a gene having the base sequence shown in SEQ ID NO: 2 was obtained. PCR was performed using TaKaRa LA Taq (manufactured by TAKARA BIO) as a DNA polymerase, where the reaction
25 conditions followed the instruction manual thereof. The DNA was digested with NdeI and EcoRI, and inserted between the NdeI recognition site and the EcoRI recognition site in the downstream of the lac promoter of plasmid pUCNT (WO94/03613) to construct a recombinant
30 vector pNTOM4.

【0073】

Similarly, using primer 5: 5'-
gtgcatatggcctaagactgtctatttca-3' (SEQ ID NO: 11) and primer 6: 5'-gtcgaattcttactagaatggaatgtcaaa-3' (SEQ ID
35 NO: 12), a double stranded DNA wherein an NdeI

recognition site is added to the initiation codon and an EcoRI recognition site is added immediately after the termination codon of a gene having the base sequence shown in SEQ ID NO: 1 was obtained. The DNA was inserted
5 into plasmid pUCNT in the same manner as above to construct a recombinant vector pNTOM3.

【0074】

(Example 4) preparation of mutant gene

Using Mutan-SuperExpress Km (manufactured by TAKARA
10 BIO) and following the operation method described in the instruction manual thereof, a mutant gene wherein, in the gene having the base sequence shown in SEQ ID NO: 2, the 125th C was substituted by G, the 128th A was substituted by C, the 136th A was substituted by G, and
15 the 147th C was substituted by G, was prepared. The mutant gene encodes a polypeptide wherein, in the amino acid sequence shown in SEQ ID NO: 4, the 42nd alanine is substituted by glycine, the 43rd glutamic acid is substituted by alanine, the 46th lysine is substituted
20 by glutamic acid, and the 49th asparagine is substituted by lysine. The base sequence of the mutant gene is shown in SEQ ID NO: 15, and the amino acid sequence of the mutant polypeptide encoded by the gene is shown in SEQ ID NO: 16. The mutant gene was inserted into a vector
25 pUCNT in the same manner as in Example 3 to construct a recombinant vector pNTOM5.

【0075】

(Example 5) construction of expression vector further containing glucose dehydrogenase gene

30 Using primer 7: 5'-
gccgaattctaaggagggttaacaatgtataaa-3' (SEQ ID NO: 13) and
primer 8: 5'-gcggtcgacttatccgcgtcctgcttgg-3' (SEQ ID NO:
14) and plasmid pGDK1 (Eur. J. Biochem., 186, 389
(1989)) as a template, PCR was performed to give a
35 double stranded DNA wherein, in a glucose dehydrogenase

(hereinafter to be referred to as GDH) gene derived from *Bacillus megaterium* IAM1030 strain, an *Escherichia coli* ribosome binding sequence is added at 5 bases upstream of the initiation codon, an EcoRI cleavage point is added immediately before the ribosome binding sequence and a SalI cleavage point is added immediately after the stop codon, was obtained.

【0076】

The obtained DNA fragment was digested with EcoRI and SalI, and inserted into the EcoRI and SalI sites of pNTOM3, pNTOM4 and pNTOM5 constructed in Examples 3 and 4 to construct recombinant plasmids pNTOM3G1, pNTOM4G1 and pNTOM5G1. As an example, the production method and structure of pNTOM4G1 are shown in Fig. 1.

15 【0077】

(Example 6) preparation of transformant

Using the recombinant vectors pNTOM3, pNTOM4 and pNTOM5 constructed in Examples 3 and 4, *E. coli* HB101 competent cell (manufactured by TAKARA BIO) was transformed to give *E. coli* HB101 (pNTOM3), *E. coli* HB101 (pNTOM4) and *E. coli* HB101 (pNTOM5). These transformants have been deposited under accession numbers FERM BP-10368, FERM BP-10369 and FERM BP-10370, respectively, in the National Institute of Advanced Industrial Science and Technology, International Patent Organism Depositary (IPOD).

25 【0078】

Similarly, using the recombinant vectors pNTOM3G1, pNTOM4G1 and pNTOM5G1 constructed in Example 5, *E. coli* HB101 competent cell (manufactured by TAKARA BIO) was transformed to give *E. coli* HB101 (pNTOM3G1), *E. coli* HB101 (pNTOM4G1) and *E. coli* HB101 (pNTOM5G1).

30 【0079】

(Example 7) gene expression in transformant

Six kinds of transformants obtained in Example 6

and *E. coli* HB101 (pUCNT), which is a transformant containing a vector plasmid pUCNT, were inoculated to 50 ml of 2×YT medium (tryptone 1.6%, yeast extract 1.0%, NaCl 0.5%, pH 7.0) containing 200 µg/ml ampicillin, and
 5 cultured with shaking at 37°C for 24 hr. The cells were collected by centrifugation, and suspended in 50 ml of 100 mM phosphate buffer (pH 6.5). The cells were disrupted with a UH-50 ultrasonication homogenizer (manufactured by SMT), and cell debris was removed by
 10 centrifugation to give a cell-free extract. The (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone reduction activity and GDH activity of the cell-free extract were measured and expressed as the specific activity in Table 1. In all of the 6 kinds of
 15 transformants obtained in Example 6, expression of the (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone reduction activity was observed. In *E. coli* HB101 (pNTOM3G1), *E. coli* HB101 (pNTOM4G1) and *E. coli* HB101 (pNTOM5G1), containing a GDH gene, expression of
 20 the GDH activity was also observed. The (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone reduction activity was measured by the method described in Example 1. The GDH activity was calculated from rate of increase in the absorbance at a wavelength of 340 nm, when 0.1 M
 25 glucose, 2 mM coenzyme NADP, and a crude enzyme solution were added to 1M Tris-HCl buffer (pH 8.0), and the mixture was reacted at 25°C for 1 min. The enzyme activity to reduce 1 µmol of NADP to NADPH for 1 min under said reaction conditions was defined to be 1 unit.
 30 In addition, the protein concentration in the cell-free extract was measured using a protein assay kit (manufactured by BIO-RAD).

【0080】

Table 1

strain	CBPB* reduction activity (U/mg)	GDH activity (U/mg)
<i>E. coli</i> HB101 (pUCNT)	0.0	0
<i>E. coli</i> HB101 (pNTOM3)	6.3	0
<i>E. coli</i> HB101 (pNTOM4)	5.0	0
<i>E. coli</i> HB101 (pNTOM5)	14.0	0
<i>E. coli</i> HB101 (pNTOM3G1)	5.7	119
<i>E. coli</i> HB101 (pNTOM4G1)	3.5	157
<i>E. coli</i> HB101 (pNTOM5G1)	9.0	122

CBPB*: (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone

5

【0081】

(Example 8) reaction using transformant

Glucose dehydrogenase (manufactured by Amano Enzyme) 2000 U, glucose 2 g, NADP 6 mg, (3S)-1-chloro-3-
 10 tert-butoxycarbonylamino-4-phenyl-2-butanone 2.5 g were added to 50 ml of a cell-free extract of *E. coli* HB101 (pNTOM3) prepared in Example 7, and the mixture was stirred at 30°C for 22 hr while adjusting to pH 6.5 by dropwise addition of 5 M sodium hydroxide. After
 15 completion of the reaction, the reaction mixture was extracted with toluene, desolvated and the extract was analyzed. As a result, (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol (yield 86.3%, diastereomeric excess 96.4%d.e.) was obtained.

20 【0082】

For quantitation of (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone and (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, as

well as for measurement of diastereomeric excess of (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol, high performance liquid chromatography (column: COSMOSIL 5C8-MS (ϕ 4.6 mm $\times$ 250 mm; manufactured by Nacalai Tesque), eluent: 10 mM aqueous phosphoric acid solution/acetonitrile=1/1, flow rate: 1 ml/min, detection: 210 nm) was used.

【0083】

(Example 9) reaction using transformant

10 Glucose dehydrogenase (manufactured by Amano Enzyme) 1250 U, glucose 3 g, NADP 4 mg, (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone 0.25 g were added to 22.5 ml of a cell-free extract of *E. coli* HB101 (pNTOM4) prepared in Example 7, and the mixture was
15 stirred at 30°C while adjusting to pH 6.5 by dropwise addition of 5 M sodium hydroxide. (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone (0.25 g) was added at 2 hr, 4 hr and 6 hr from the start of the reaction and 1.25 g thereof was added at 8 hr from the
20 start of the reaction, and the mixture was reacted for 31 hr. After completion of the reaction, the reaction mixture was extracted with toluene, desolvated and the extract was analyzed in the same manner as in Example 8. As a result, (2R,3S)-1-chloro-3-tert-
25 butoxycarbonylamino-4-phenyl-2-butanol (yield 99.0%, diastereomeric excess 99.8%d.e.) was obtained.

【0084】

(Example 10) reaction using transformant

30 Glucose 3 g, NADP 3 mg, toluene 0.25 ml, and (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone 2.5 g were added to 25 ml of a cultured medium of *E. coli* HB101 (pNTOM4G1) prepared in the same manner as in Example 7, and the mixture was stirred at 30°C for 40 hr while adjusting to pH 6.5 by dropwise addition of 5 M
35 sodium hydroxide. After completion of the reaction, the

reaction mixture was extracted with toluene, desolvated and the extract was analyzed in the same manner as in Example 8. As a result, (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol (yield 99.1%,
5 diastereomeric excess 99.8%d.e.) was obtained.

【0085】

(Example 11) reaction using transformant

Glucose 6 g, NADP 1.4 mg, toluene 0.25 ml, and (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-
10 butanone 2.5 g were added to 50 ml of a cultured medium of *E. coli* HB101 (pNTOM5G1) prepared in the same manner as in Example 7, and the mixture was stirred at 30°C while adjusting to pH 6.5 by dropwise addition of 5 M sodium hydroxide. One hour after the start of the
15 reaction, 2.5 g of (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone was added, and the mixture was reacted for 20 hr. After completion of the reaction, the reaction mixture was extracted with toluene, desolvated and the extract was analyzed in the
20 same manner as in Example 8. As a result, (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol (yield 99.6%, diastereomeric excess 99.8%d.e.) was obtained.

【0086】

25 (Example 12) substrate specificity of polypeptide

In 100 mM phosphate buffer (pH 6.5) containing 0.33% (v/v) dimethyl sulfoxide were dissolved a carbonyl compound to be a substrate to a final concentration of 1 mM, and a coenzyme NADPH to a final concentration of
30 0.25 mM, respectively. Thereto was added a cell-free extract of *E. coli* HB101 (pNTOM4) or *E. coli* HB101 (pNTOM3) prepared in Example 7, and the mixture was reacted at 30°C for 1 min. The reduction activity on each carbonyl compound was calculated from the rate of
35 decrease in the absorbance of the reaction mixture at a

wavelength of 340 nm, and shown in Table 2 as a value relative to the activity on (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone as 100%. The polypeptides consisting of the amino acid sequences shown in SEQ ID NO: 3 and SEQ ID NO: 4, which were produced by the transformants, both showed reduction activity on a wide range of carbonyl compounds.

【0087】

Table 2

Substrate specificity of polypeptides produced by transformant

substrate (carbonyl compound)	relative activity (%)	
	<i>E. coli</i> HB101	
	(pNTOM3)	(pNTOM4)
CBPB*	100	100
2-acetylpyridine	29	0
3-acetylpyridine	15	0
4-acetylpyridine	98	0
acetylpyrazine	185	0
acetophenone	215	0
3'-hydroxyacetophenone	33	0
3'-nitroacetophenone	273	31
4'-chloroacetophenone	35	9
4'-fluoroacetophenone	28	1
3',4'-dimethoxyacetophenone	51	4
4'-methylacetophenone	41	1
2-hydroxyacetophenone	32	1
2-chloro-1-(3'-chlorophenyl)ethanone	1301	33
1-phenyl-2-butanone	285	2
propiophenone	183	2
benzoin	4018	5
benzylacetone	66	0

ethyl benzoylformate	296	20
ethyl 2-oxo-4-phenylbutyrate	127	106
acetone	47	0
2-butanone	22	0
2-hexanone	11	0
2-heptanone	102	0
2-octanone	808	3
chloroacetone	100	1
hydroxyacetone	51	0
diacetyl	444	5
acetylacetone	23	0
cyclopentanone	8	0
7-methoxy-2-tetoralone	599	6
1-tetoralone	146	1
ethyl 3-oxocyclopentanecarboxylate	484	12
methyl pyruvate	674	5
ethyl pyruvate	1067	23
methyl acetoacetate	8	0
ethyl acetoacetate	137	0
tert-butyl acetoacetate	1330	2
ethyl 2-methylacetoacetate	619	4
ethyl 2-chloroacetoacetate	3309	40
ethyl 2-eneacetoacetate	2013	10
ethyl 2-oxodecanoate	985	5
ethyl 4-chloroacetoacetate	4547	22
n-octyl 4-chloroacetoacetate	466	24
ethyl 4-bromoacetoacetate	476	7
ethyl 4-azideacetoacetate	659	1
ethyl 4-benzyloxyacetoacetate	1259	8
ethyl 4-acetoxyacetoacetate	465	1
benzyl acetoacetate	883	4
ethyl benzoylacetate	1233	0
ethyl 2-chloro-3-oxo-3-phenylpropionate	716	5

benzaldehyde	81	1
2-pyridinecarbaldehyde	56	1
pyridine-4-aldehyde	571	2
o-chlorobenzaldehyde	269	17
m-chlorobenzaldehyde	344	17
p-chlorobenzaldehyde	91	12
o-nitrobenzaldehyde	1621	89
m-nitrobenzaldehyde	851	29
p-nitrobenzaldehyde	1166	21
propionaldehyde	99	0
n-butylaldehyde	168	0
n-hexylaldehyde	265	3
2-phenylpropionaldehyde	479	3
3-phenylpropionaldehyde	2410	7
glutaraldehyde	80	1


CBPB*: (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone

Claims


1. A DNA of the following (a) or (b):

5 (a) a DNA comprising a base sequence shown in SEQ ID NO: 1,

(b) a DNA hybridizing to a DNA consisting of a base sequence complementary to the base sequence shown in SEQ ID NO: 1 under stringent conditions, and encoding a
10 polypeptide having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by the following formula (1):



15 to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the following formula (2):



20

2. A DNA of the following (a) or (b):

(a) a DNA comprising a base sequence shown in SEQ ID NO: 2,

(b) a DNA hybridizing to a DNA consisting of a base sequence complementary to the base sequence shown in SEQ ID NO: 2 under stringent conditions, and encoding a polypeptide having an activity to asymmetrically reduce
5 (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by the aforementioned formula (1) to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the aforementioned formula (2).

10

3. The polypeptide encoded by the DNA of claim 1 or 2 and having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by the aforementioned formula (1) to produce
15 (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the aforementioned formula (2).

4. A polypeptide of the following (a) or (b):

(a) a polypeptide comprising an amino acid sequence
20 shown in SEQ ID NO: 3,

(b) a polypeptide comprising an amino acid sequence showing not less than 78% homology with the amino acid sequence shown in SEQ ID NO: 3 and having an activity to asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by
25 the aforementioned formula (1) to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the aforementioned formula (2).

30 5. A polypeptide of the following (a) or (b):

(a) a polypeptide comprising an amino acid sequence shown in SEQ ID NO: 4,

(b) a polypeptide comprising an amino acid sequence showing not less than 78% homology with the amino acid
35 sequence shown in SEQ ID NO: 4 and having an activity to

asymmetrically reduce (3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanone represented by the aforementioned formula (1) to produce (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol
5 represented by the aforementioned formula (2).

6. A polypeptide having the amino acid sequence shown in SEQ ID NO: 4, but with at least one mutation from the following 1) to 4):

- 10 1) the 42nd alanine is substituted by glycine,
2) the 43rd glutamic acid is substituted by alanine,
3) the 46th lysine is substituted by glutamic acid,
4) the 49th asparagine is substituted by lysine.

15 7. The polypeptide of claim 6, comprising all of the mutations 1) to 4).

8. A DNA encoding the polypeptide of any one of claims 4 to 7.

20

9. A vector comprising the DNA of claim 1, 2 or 8.

10. The vector of claim 9, which is a plasmid pNTOM3 obtainable from *E. coli* HB101 (pNTOM3) FERM BP-10368.

25 11. The vector of claim 9, which is a plasmid pNTOM4 obtainable from *E. coli* HB101 (pNTOM4) FERM BP-10369.

12. The vector of claim 9, which is a plasmid pNTOM5 obtainable from *E. coli* HB101 (pNTOM5) FERM BP-10370.

13. The vector of claim 9, further comprising a DNA
30 encoding a polypeptide having a glucose dehydrogenase activity.

14. The vector of claim 13, wherein the polypeptide having a glucose dehydrogenase activity is a glucose
35 dehydrogenase derived from *Bacillus megaterium*.

15. A transformant obtained by transforming a host cell with the vector of any one of claims 9 to 14.

5 16. The transformant of claim 15, wherein the host cell is *Escherichia coli*.

17. The transformant of claim 16, which is *E. coli* HB101 (pNTOM3) FERM BP-10368.

10

18. The transformant of claim 16, which is *E. coli* HB101 (pNTOM4) FERM BP-10369.

19. The transformant of claim 16, which is *E. coli* HB101
15 (pNTOM5) FERM BP-10370.

20. A method of producing an optically active alcohol, which comprises reacting the polypeptide of any one of claims 3 to 7 or the transformant of any one of claims
20 15 to 19, or a treated product thereof, with a compound having a carbonyl group.

21. The method of claim 20, wherein the compound having a carbonyl group is (3S)-1-chloro-3-tert-
25 butoxycarbonylamino-4-phenyl-2-butanone represented by the aforementioned formula (1), and the optically active alcohol is (2R,3S)-1-chloro-3-tert-butoxycarbonylamino-4-phenyl-2-butanol represented by the aforementioned formula (2).

FIG. 1

