



US00PP13953P39

(12) **United States Plant Patent**
Crocker

(10) **Patent No.:** **US PP13,953 P3**

(45) **Date of Patent:** **Jul. 8, 2003**

(54) **PLUMBAGO AURICULATA NAMED**
'MONITE'

(52) **U.S. Cl.** **Plt./226**

(58) **Field of Search** **Plt./226**

(75) **Inventor:** **Brad Alan Crocker, Aliseo Viejo, CA**
(US)

Primary Examiner—Bruce R. Campell

(73) **Assignee:** **Monrovia Nursery Company, Azusa,**
CA (US)

Assistant Examiner—Susan B. McCormick

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

(21) **Appl. No.:** **09/826,222**

A new and distinct selection of *Plumbago* shrub, denomi-
nated as *Plumbago auriculata* named 'Monite' characterized
by its unique combination of longer bloom season of a
floriferous habit of white flowers with larger and wider
petals and a more dense compact growth habit.

(22) **Filed:** **Apr. 5, 2001**

(65) **Prior Publication Data**

US 2002/0148019 P1 Oct. 10, 2002

(51) **Int. Cl.**⁷ **A01H 5/00**

1 Drawing Sheet

1

2

BACKGROUND

This invention relates to a new and distinct selection of
Plumbago auriculata, which is native to Southern Africa,
and a member of the Leadwort family. The new variety,
Plumbago auriculata cv. 'Monite' resulted from a naturally
occurring, whole plant mutation which was discovered in
May of 1999 in a cultivated planting of *Plumbago auricu-*
lata 'Monott' (U.S. Plant Pat. No. 7,822). 'Monite' exhibited
a white flower in #1 nursery containers in Azusa, Calif. The
'Monite' selection differs from 'Monott' by possessing a
white flower versus the rich blue flower of 'Monott'.
'Monite' differs from other *Plumbago* plants by its combi-
nation of floriferous habit of white flowers and larger and
wider petals and a more dense compact growth habit.

The new plant has been asexually reproduced by cuttings
since its discovery in 1999 in Azusa, Calif. Through exten-
sive asexual propagation of this plant and observation, it has
been established that the plant is true to type and the novel,
exceptional characteristics of this new plant are fixed and
stable and reliably passed on to progeny through asexual
reproduction.

SUMMARY

Plumbago auriculata cultivar 'Monite' possesses a white
flower. Other white flowering *Plumbago* plants do exist,
however; the individual flowers do not exhibit the larger size
and fullness associated with the overlapping flower petals of
cultivar 'Monite'.

The compact growth habit of cultivar 'Monite' sets this
plant apart from other *Plumbago* plants which are generally
sprawling, mounding plants with a height of 6-8 feet and
8-10 foot spread. The growth habit of cultivar 'Monite' is
slightly smaller than 'Monott'.

The cultivar 'Monite' typically will bloom for eight to
nine months of the year in Southern California commencing
bloom before even the early cultivar 'Monott'. Comparison
of other *Plumbago* plants demonstrate cultivar 'Monite' has

the propensity to conspicuously grow less and flower more
when grown under similar conditions.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 photograph exhibits a close up of the inflorescence
of cultivar 'Monite'.

FIG. 2 photograph shows the comparison of cultivar
'Monite' on the right and cultivar 'Monott' on the left and
the similarity of flower size and contrast of flower color.

BOTANICAL DESCRIPTION

Below is a detailed description of *Plumbago auriculata*
cultivar 'Monite'. Color terminology is from The Royal
Horticultural Society Colour Charts. Measurements are
based on five year old plants grown in Southern California.

Overall habit: Mounding shrub.

Overall size: Maximum height from soil surface to top of
foliage ranges from 120 cm to 150 cm. Maximum width
of the plant ranges from 120 cm to 150 cm.

Flowering time in Southern California: Flowering com-
mences in early March and continues until November.

Root initiation: Six to eight weeks.

Flower:

Arrangement.—Terminal, spike like racemes.

Lastingness of bloom.—Averaging 2 to 3 days.

Corolla diameter.—Averaging 2.5 to 3 cm in width.

Corolla color.—White Group 155D.

Corolla.—Tubular with 5 parted lobes. Tube: typically
3 cm in length; 2 mm in width. Lobes: slightly
overlapping, obovate in shape, acute base averaging
1.5 cm long and 1 cm wide, prominent midrib,
slightly depressed. Texture: thin, glabrous.

Calyx.—Tubular, 5 parted at tip. Color: Green Group
141C. Size: averaging 1.3 cm in length, 3 mm in
width. Surface: upper half of calyx with stalked
glands, Greyed-Red Group 180C. Bracts: 3 at base,

6–7 mm long and 2 mm wide, color Green Group 141C.

Stamens.—5 in number, adhering to within throat arising above corolla tube. Anthers: length averaging 2 mm. width: less than 1 mm. color: Violet-Blue 92B. Filaments: length averaging 3 cm. width: less than 1 mm, very slender. color: White Group 155D.

Pistil.—Averaging 2 cm in length. Color: White Group 155D.

Fragrance: None detected.

Fruit: To date not observed.

Foliage:

Size.—Length: ranges from 3.5 cm to 6.5 cm. Width: ranges from 1 cm to 2 cm.

Shape.—Overall: oblong to oblong — spatulate to obtuse. Base: auricled, wrapping around stem, narrowly acute. Tip: shortly mucronate.

Margin.—Entire, slightly undulate.

Color.—Mature upper surface: Yellow Green Group 146A. Mature lower surface: Yellow Green Group 146B. Juvenile upper surface: Yellow Green Group 146B. Juvenile lower surface: Yellow Green Group 146C.

Texture.—Thin, glabrous.

Arrangement.—Alternate.

Venation.—Pattern: pinnate, craspedrome. Upper surface: Yellow-Green Group 146C. Lower surface: Yellow-Green Group 146D.

Branchlets:

Juvenile.—Lightly ribbed, 2–3 mm in diameter. color: Yellow Green Group 146B.

Mature.—3 mm and larger in diameter. color: Red-Purple Group 59C mixed with Yellow Green Group 146B.

Length.—Varies from 60 cm.

Length of internode.—Varies from 2.5 cm to 6 cm dependent upon age and sun exposure on branch.

Disease resistance: As is typical of this species, 'Monite' cultivar exhibited a strong resistance to disease under normal growing conditions.

Heat and drought hardiness: When established in the landscape 'Monte' is very tolerant of dry conditions and reflected and dry heat.

Winter hardiness: As is typical of this species, will tolerate to a 25 degree Fahrenheit temperature.

I claim:

1. A new and distinct *Plumbago auriculata* plant named 'Monite' substantially as herein shown and described.

* * * * *



FIGURE 1



FIGURE 2