



(86) **Date de dépôt PCT/PCT Filing Date:** 2014/11/21
(87) **Date publication PCT/PCT Publication Date:** 2015/05/28
(85) **Entrée phase nationale/National Entry:** 2016/05/17
(86) **N° demande PCT/PCT Application No.:** US 2014/066965
(87) **N° publication PCT/PCT Publication No.:** 2015/077655
(30) **Priorité/Priority:** 2013/11/22 (US61/907,913)

(51) **Cl.Int./Int.Cl.** **C07D 417/04** (2006.01),
A61K 31/4439 (2006.01), **A61K 31/706** (2006.01),
C07H 15/26 (2006.01)
(71) **Demandeur/Applicant:**
UNIVERSITY OF FLORIDA RESEARCH FOUNDATION,
INC., US
(72) **Inventeur/Inventor:**
BERGERON, RAYMOND J., JR., US
(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : ANALOGUES DE LA DESFERRITHIOCINE ET LEURS UTILISATIONS**
(54) **Title: DESFERRITHIOCIN ANALOGS AND USES THEREOF**





Figure 1B



(57) **Abrégé/Abstract:**

Iron overload is associated with pathological conditions such as oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or



(57) Abrégé(suite)/Abstract(continued):

neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, and reperfusion injury. The present invention provides methods and pharmaceutical compositions using desferrithiocin analogs of Formulae (A) and (J) for treating and/or preventing these pathological conditions, metal (e.g., iron, aluminum, a lanthanide, or an actinide (e.g., uranium)) overload conditions, and infectious diseases (e.g., malaria).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
28 May 2015 (28.05.2015)(10) International Publication Number
WO 2015/077655 A1(51) International Patent Classification:
A61K 31/426 (2006.01)(21) International Application Number:
PCT/US2014/066965(22) International Filing Date:
21 November 2014 (21.11.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/907,913 22 November 2013 (22.11.2013) US(71) Applicant: UNIVERSITY OF FLORIDA RESEARCH
FOUNDATION, INC. [US/US]; 223 Grinter Hall,
Gainesville, FL 32611 (US).(72) Inventor: BERGERON, Raymond, J., Jr.; 6220 NW
56th Lane, Gainesville, FL 32653-3117 (US).(74) Agent: BAKER, C., Hunter; Wolf Greenfield & Sacks,
P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: DESFERRITHIOCIN ANALOGS AND USES THEREOF





Figure 1B



(57) Abstract: Iron overload is associated with pathological conditions such as oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, and reperfusion injury. The present invention provides methods and pharmaceutical compositions using desferrithiocin analogs of Formulae (A) and (J) for treating and/or preventing these pathological conditions, metal (e.g., iron, aluminum, a lanthanide, or an actinide (e.g., uranium)) overload conditions, and infectious diseases (e.g., malaria).

WO 2015/077655 A1

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *before the expiration of the time limit for amending the*
GW, KM, ML, MR, NE, SN, TD, TG). *claims and to be republished in the event of receipt of*
Published: *amendments (Rule 48.2(h))*
— *with international search report (Art. 21(3))*

DESFERRITHIOCIN ANALOGS AND USES THEREOF

RELATED APPLICATION

[0001] The present application claims priority under 35 U.S.C. § 119(e) to U.S. provisional patent application, U.S.S.N. 61/907,913, filed November 22, 2013, which is incorporated herein by reference.

GOVERNMENT SUPPORT

[0002] This invention was made with U.S. Government support under Grant No. R37DK049108 awarded by the National Institutes of Health. The U.S. Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

[0003] Nearly all life forms require iron as a micronutrient. However, the low solubility of Fe(III) hydroxide ($K_{sp} = 1 \times 10^{-39}$) (Raymond *et al.*, “Coordination Chemistry and Microbial Iron Transport.” *Acc. Chem. Res.* **1979**, *12*, 183-190), the predominant form of the metal in the biosphere, required the development of sophisticated iron storage and transport systems in nature. Microorganisms utilize low molecular weight, ferric iron-specific chelators, siderophores (Byers *et al.*, “Microbial Iron Transport: Iron Acquisition by Pathogenic Microorganisms.” *Met. Ions Biol. Syst.* **1998**, *35*, 37-66); eukaryotes tend to employ proteins to transport and store iron (Bergeron, “Iron: A Controlling Micronutrient in Proliferative Processes.” *Trends Biochem. Sci.* **1986**, *11*, 133-136; Theil *et al.*, “Ferritin Mineralization: Ferroxidation and Beyond.” *J. Inorg. Biochem.* **1997**, *67*, 30; Ponka *et al.*, “Function and Regulation of Transferrin and Ferritin.” *Semin. Hematol.* **1998**, *35*, 35-54). Humans have evolved a highly efficient iron management system in which we absorb and excrete only about 1 mg of the metal daily; there is no mechanism for the excretion of excess metal (Brittenham, “Disorders of Iron Metabolism: Iron Deficiency and Overload.” In *Hematology: Basic Principles and Practice*; 3rd ed.; Hoffman *et al.*, Eds.; Churchill Livingstone: New York, 2000; pp. 397-428). Whether derived from transfused red blood cells (Olivieri *et al.*, “Iron-Chelating Therapy and the Treatment of Thalassemia.” *Blood* **1997**, *89*, 739-761; Vichinsky, “Current Issues with Blood Transfusions in Sickle Cell Disease.” *Semin. Hematol.* **2001**, *38*, 14-22; Kersten *et al.*, “Long-Term Treatment of Transfusional Iron Overload with the Oral Iron Chelator Deferiprone (L1): A Dutch Multicenter Trial.” *Ann.*

Hematol. **1996**, 73, 247-252) or from increased absorption of dietary iron (Conrad *et al.*, “Iron Absorption and Transport.” *Am. J. Med. Sci.* **1999**, 318, 213-229; Lieu *et al.*, “The Roles of Iron in Health and Disease.” *Mol. Aspects Med.* **2001**, 22, 1-87), without effective treatment, body iron progressively increases with deposition in the liver, heart, pancreas, and elsewhere (iron overload disease).

[0004] In patients with iron overload disease, the toxicity derives from iron’s interaction with reactive oxygen species (Graf *et al.*, “Iron-Catalyzed Hydroxyl Radical Formation. Stringent Requirement for Free Iron Coordination Site.” *J. Biol. Chem.* **1984**, 259, 3620-3624; Halliwell, “Free Radicals and Antioxidants: A Personal View.” *Nutr. Rev.* **1994**, 52, 253-265; Halliwell, “Oxidative Damage, and Chelating Agents.” In *The Development of Iron Chelators for Clinical Use*; Bergeron *et al.*, Eds.; CRC: Boca Raton, FL, 1994; pp 33-56; Koppenol, “Kinetics and Mechanism of the Fenton Reaction: Implications for Iron Toxicity.” In *Iron Chelators: New Development Strategies*; Badman *et al.*, Eds.; Saratoga: Ponte Vedra Beach, FL, 2000, pp 3-10). For example, in the presence of Fe(II), endogenous H₂O₂ is reduced to the hydroxyl radical (HO•), a very reactive species, and HO⁻, in the Fenton reaction. The hydroxyl radical reacts very quickly with a variety of cellular constituents and can initiate free radicals and radical-mediated chain processes that damage DNA and membranes as well as produce carcinogens (Halliwell, “Free Radicals and Antioxidants: A Personal View.” *Nutr. Rev.* **1994**, 52, 253-265); Babbs, “Oxygen Radicals in Ulcerative Colitis.” *Free Radical Biol. Med.* **1992**, 13, 169-181; Hazen *et al.*, “Human Neutrophils Employ the Myeloperoxidase-Hydrogen Peroxide-Chloride System to Oxidize α-Amino Acids to a Family of Reactive Aldehydes. Mechanistic Studies Identifying Labile Intermediates along the Reaction Pathway.” *J. Biol. Chem.* **1998**, 273, 4997-5005). The liberated Fe(III) is reduced back to Fe(II) via a variety of biological reductants (*e.g.*, ascorbate, glutathione), a problematic cycle.

[0005] Iron-mediated damage can be focal, as in reperfusion damage (Millán *et al.*, “Biological Signatures of Brain Damage Associated with High Serum Ferritin Levels in Patients with Acute Ischemic Stroke and Thrombolytic Treatment.” *Dis. Markers* **2008**, 25, 181-188), Parkinson’s (Zecca *et al.*, “Neuromelanin Can Protect Against Iron-Mediated Oxidative Damage in System Modeling Iron Overload of Brain Aging and Parkinson’s Disease.” *J. Neurochem.* **2008**, 106, 1866-1875), Friedreich’s ataxia (Pietrangelo, “Iron Chelation Beyond Transfusion Iron Overload.” *Am. J. Hematol.* **2007**, 82, 1142-1146), macular degeneration (Dunaief, “Iron Induced Oxidative Damage as a Potential Factor in

Age-Related Macular Degeneration: The Cogan Lecture” *Invest. Ophthalmol. Vis. Sci.* **2006**, 47, 4660-4664), and hemorrhagic stroke (Hua *et al.*, “Long-Term Effects of Experimental Intracerebral Hemorrhage: The Role of Iron.” *J. Neurosurg.* **2006**, 104, 305-312), or global, as in transfusional iron overload, *e.g.*, thalassemia (Pippard, “Iron Overload and Iron Chelation Therapy in Thalassaemia and Sickle Cell Haemoglobinopathies.” *Acta. Haematol.* **1987**, 78, 206-211), sickle cell disease (Pippard, “Iron Overload and Iron Chelation Therapy in Thalassaemia and Sickle Cell Haemoglobinopathies.” *Acta. Haematol.* **1987**, 78, 206-211; Olivieri, “Progression of Iron Overload in Sickle Cell Disease.” *Semin. Hematol.* **2001**, 38, 57-62), and myelodysplasia (Malcovati, “Impact of Transfusion Dependency and Secondary Iron Overload on the Survival of Patients with Myelodysplastic Syndromes.” *Leukemia Res.* **2007**, 31, S2-S6), with multiple organ involvement. The solution in both scenarios is the same: chelate and promote the excretion of excess unmanaged iron.

[0006] Treatment with a chelating agent capable of sequestering iron and permitting its excretion from the body is the only therapeutic approach available. Some of the iron chelating agents that are now in use or that have been clinically evaluated include desferrioxamine B mesylate (DFO^a) (*Desferal*; Novartis Pharmaceuticals Corporation: East Hanover, NJ, **2008**; www.pharma.us.novartis.com/product/pi/pdf/desferal.pdf), 1,2-dimethyl-3-hydroxy-4-pyridinone (deferiprone, L1) (Hoffbrand, “Long-Term Trial of Deferiprone in 51 Transfusion-Dependent Iron Overloaded Patients.” *Blood* **1998**, 91, 295-300; Olivieri, “Long-Term Therapy with Deferiprone.” *Acta Haematol.* **1996**, 95, 37-48; Olivieri, “Long-Term Safety and Effectiveness of Iron-Chelation Therapy with Deferiprone from Thalassemia Major.” *N. Engl. J. Med.* **1998**, 339, 417-423; Richardson, “The Controversial Role of Deferiprone in the Treatment of Thalassemia.” *J. Lab. Clin. Med.* **2001**, 137, 324-329), and 4-[3,5-bis(2-hydroxyphenyl)-1,2,4-triazol-1-yl]benzoic acid (desferasirox, ICL670A) (Nisbet-Brown *et al.*, “Effectiveness and Safety of ICL670 in Iron-Loaded Patients with Thalassemia: A Randomised, Double-Blind, Placebo-Controlled, Dose-Escalation Trial.” *Lancet*, **2003**, 361, 1597-1602; Galanello *et al.*, “Safety, Tolerability, and Pharmacokinetics of ICL670, a New Orally Active Iron-Chelating Agent in Patients with Transfusion-Dependent Iron Overload Due to β -Thalassemia.” *J. Clin. Pharmacol.* **2003**, 43, 565-572; Cappellini, “Iron-Chelating Therapy with the New Oral Agent ICL670 (Exjade).” *Best Pract. Res. Clin. Haematol.* **2005**, 18, 289-298). Each of these compounds presents with shortcomings. DFO must be given subcutaneously (sc) for protracted periods of time, *e.g.*, 12 h a day, five days a week, a serious patient compliance issue (Olivieri *et al.*, “Iron-Chelating


Therapy and the Treatment of Thalassemia.” *Blood* **1997**, 89, 739-761; Pippard, “Desferrioxamine-Induced Iron Excretion in Humans.” *Bailliere’s Clin. Haematol.* **1989**, 2, 323-343; Giardina *et al.*, “Chelation Therapy in β -Thalassemia: An Optimistic Update.” *Semin. Hematol.* **2001**, 38, 360-366). Deferiprone, while orally active, simply does not remove enough iron to maintain patients in a negative iron balance (Hoffbrand, “Long-Term Trial of Deferiprone in 51 Transfusion-Dependent Iron Overloaded Patients.” *Blood* **1998**, 91, 295-300; Olivieri, “Long-Term Therapy with Deferiprone.” *Acta Haematol.* **1996**, 95, 37-48; Olivieri, “Long-Term Safety and Effectiveness of Iron-Chelation Therapy with Deferiprone from Thalassemia Major.” *N. Engl. J. Med.* **1998**, 339, 417-423; Richardson, “The Controversial Role of Deferiprone in the Treatment of Thalassemia.” *J. Lab. Clin. Med.* **2001**, 137, 324-329). Desferasirox did not show noninferiority to DFO and is associated with numerous side effects, including some renal toxicity (Nisbet-Brown *et al.*, “Effectiveness and Safety of ICL670 in Iron-Loaded Patients with Thalassemia: A Randomised, Double-Blind, Placebo-Controlled, Dose-Escalation Trial.” *Lancet*, **2003**, 361, 1597-1602; Galanello *et al.*, “Safety, Tolerability, and Pharmacokinetics of ICL670, a New Orally Active Iron-Chelating Agent in Patients with Transfusion-Dependent Iron Overload Due to β -Thalassemia.” *J. Clin. Pharmacol.* **2003**, 43, 565-572; Cappellini, “Iron-Chelating Therapy with the New Oral Agent ICL670 (Exjade).” *Best Pract. Res. Clin. Haematol.* **2005**, 18, 289-298).

[0007] Despite the work on metal chelation agents described above, there is still a need for other chelators with more desirable properties (*e.g.*, improved physiochemical, pharmacokinetic, pharmacodynamic, and/or toxicological properties, such as absorption, distribution, metal-clearing efficiency, and toxicity) for a better treatment and/or prevention of pathological conditions in a subject.

SUMMARY OF THE INVENTION


[0008] The present invention provides novel desferrithiocin analogs based on desferrithiocin **1** (shown below), and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof, that chelate a metal. The invention also provides methods of using the inventive desferrithiocin analogs, or pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, or polymorphs thereof, and pharmaceutical compositions thereof, for the treatment and/or prevention of a pathological condition. Without wishing to be bound by any particular theory, the inventive compounds are thought to chelate iron and/or other metals (*e.g.*, aluminum, thallium, chromium,

magnesium, calcium, strontium, nickel, manganese, cobalt, copper, zinc, silver, sodium, potassium, cadmium, mercury, lead, antimony, molybdenum, tungsten, a lanthanide (*e.g.*, cerium), or an actinide (*e.g.*, uranium)). Therefore, a metal overload, metal poisoning, and other pathological conditions (*e.g.*, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, and reperfusion injury) that are associated with metal overload may be treated and/or prevented by the inventive methods. The methods of the invention may also be useful in treating and/or preventing an infectious disease (*e.g.*, malaria). Iron is usually a nutrient necessary for the growth of microorganisms. Depriving the organisms of iron by chelating and/or removing iron may contribute to the treatment and/or prevention of infectious diseases. Further provided by the invention are kits, containing one or more inventive desferrithiocin analogs, or pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, or polymorphs thereof, or pharmaceutical compositions thereof, for treating and/or preventing a pathological condition (*e.g.*, iron overload).



1


[0009] In one aspect, provided are compounds of Formula (A), and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof:



wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and k are as described herein. The compounds of Formula (A) include at least one carbohydrate moiety, wherein the C1 position of the carbohydrate moiety is attached to the pyridyl ring or carbonyl group of the compounds of Formula (A),

optionally through a linker. In certain embodiments, the linker is a polyethylene glycol (PEG) moiety.


[0010] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[0011] In certain embodiments, the compound of Formula (A) is not of Formula (I-1), (I-2), (I-3), or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof. In certain embodiments, the compound of Formula (A) is not of Formula (I-1), (I-2), or (I-3).

[0012] In certain embodiments, the compound of Formula (A) is of the formula:






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[0013] Without wishing to be bound by any particular theory, by introducing an $-OR_2$ group on the pyridinyl ring of desferrithiocin **1**, the resulting inventive compounds have lower nephrotoxicity and/or improved metal clearance properties than the parent compound **1**.

[0014] In certain embodiments, the compounds of the invention may also include one or more carbohydrate (*e.g.*, glucose, including α -D-, β -D-, α -L-, and β -L-glucose) moieties. Such compounds are expected to show superior physiochemical, pharmacokinetic, and/or pharmacodynamic properties (*e.g.*, greater solubility, permeability, and bioavailability; improved distribution, absorption, metabolism, and iron-clearing efficiency; and reduced clearance, excretion, and toxicity) compared with the parent compound **1** and/or desferrithiocin analogs that do not include a carbohydrate moiety.

[0015] In another aspect, provided are compounds of Formula (J), and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof:



wherein --- , W₁, W₂, W₃, Z, R₁₆, R₁₇, and R₂₁ are as described herein.

[0016] In another aspect, the present invention provides pharmaceutical compositions including an inventive compound, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, and optionally a pharmaceutically acceptable excipient. The pharmaceutical compositions of the invention may include a therapeutically or prophylactically effective amount of the inventive compound.

[0017] In still another aspect, the invention provides methods of using the inventive compounds, or pharmaceutical compositions thereof, for the treatment and/or prevention of a pathological condition in a subject. In certain embodiments, the pathological condition is responsive to the chelation or sequestration of a metal. In certain embodiments, the metal is iron (*e.g.*, Fe(III)). In certain embodiments, the metal is aluminum, thallium, chromium, magnesium, calcium, strontium, nickel, manganese, cobalt, copper, zinc, silver, sodium, potassium, cadmium, mercury, lead, antimony, molybdenum, tungsten, a lanthanide (*e.g.*, cerium), or an actinide (*e.g.*, uranium). In certain embodiments, the metal is a trivalent metal. In certain embodiments, the metal is a monovalent, divalent, tetravalent, pentavalent, or hexavalent metal. In certain embodiments, the subject is a human. In certain embodiments, the pathological condition is metal overload (*e.g.*, iron overload, aluminum overload, chromium overload, magnesium overload, calcium overload, strontium overload, nickel overload, manganese overload, cobalt overload, copper overload, zinc overload, silver overload, sodium overload, potassium overload, cadmium overload, mercury overload, lead overload, molybdenum overload, tungsten overload, or actinide overload (*e.g.*, uranium overload)). In certain embodiments, the pathological condition is iron overload. In certain embodiments, the pathological condition is metal poisoning (*e.g.*, iron poisoning, aluminum poisoning, thallium poisoning, chromium poisoning, magnesium poisoning, calcium

poisoning, strontium poisoning, nickel poisoning, manganese poisoning, cobalt poisoning, copper poisoning, zinc poisoning, silver poisoning, sodium poisoning, potassium poisoning, cadmium poisoning, mercury poisoning, lead poisoning, antimony poisoning, molybdenum poisoning, tungsten poisoning, lanthanide poisoning (*e.g.*, cerium poisoning), or actinide poisoning (*e.g.*, uranium poisoning). In certain embodiments, the pathological condition is oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, and reperfusion injury. In certain embodiments, the pathological condition is an infectious disease (*e.g.*, malaria). In certain embodiments, the methods of treatment and/or prevention include administering to the subject a therapeutically or prophylactically effective amount of a compound of the invention, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, or a pharmaceutical compositions thereof.

[0018] In yet another aspect, the invention provides kits for treating and/or preventing a pathological condition in a subject. The inventive kits include a first container containing a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, or a pharmaceutical compositions thereof; and instructions for administering the compound to the subject to treat and/or prevent the pathological condition. A kit may include multiple unit dosages, for example, for multiple days of treatment.

[0019] The details of one or more embodiments of the invention are set forth herein. Other features, objects, and advantages of the invention will be apparent from the Detailed Description, Figures, Examples, and Claims.

DEFINITIONS

[0020] Definitions of specific functional groups and chemical terms are described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in *Organic Chemistry*, Thomas Sorrell, University Science Books, Sausalito, 1999; Smith and March *March's Advanced Organic*

Chemistry, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

[0021] It is to be understood that compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed “isomers.” Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers.” Stereoisomers that are not mirror images of one another are termed “diastereomers,” and those that are non-superimposable mirror images of each other are termed “enantiomers”. When a compound has an asymmetric center, for example, a carbon atom of the compound is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the *R*- and *S*-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates plane polarized light and designated as dextrorotatory or levorotatory (*i.e.*, as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a “racemic mixture.” For example, the compounds described herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. Isomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred isomers can be prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); Wilen, *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972). The invention additionally encompasses compounds described herein as individual isomers substantially free of other isomers, and alternatively, as mixtures of various isomers.

[0022] Where an isomer/enantiomer is preferred, it may, in some embodiments, be provided substantially free of the corresponding enantiomer, and may also be referred to as “optically enriched” or “enantiomerically enriched.” “Optically enriched” and “enantiomerically enriched” means that a provided compound is made up of a significantly greater proportion of one enantiomer. In certain embodiments, a compound of the present invention is made up of at least about 70% by weight of a preferred enantiomer. In certain

embodiments, a compound of the present invention is made up of at least about 80% by weight of a preferred enantiomer. In certain embodiments, a compound of the present invention is made up of at least about 90% by weight of a preferred enantiomer. In other embodiments the compound is made up of at least about 95%, 98%, or 99% by weight of a preferred enantiomer. Preferred enantiomers may be isolated from racemic mixtures by any method known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts or prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); Wilen, *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972).

[0023] Unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the depicted structures that differ only in the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by ^{13}C or ^{14}C are within the scope of this invention. Such compounds are useful, for example, as analytical tools, as probes in biological assays, or as therapeutic agents in accordance with the present invention.

[0024] In a formula, $\sim$ is a single bond where the stereochemistry of the moieties immediately attached thereto is not specified, --- is absent or a single bond, and == or === is a single or double bond.

[0025] When a range of values is listed, it is intended to encompass each value and sub-range within the range. For example “C₁₋₆” is intended to encompass, C₁, C₂, C₃, C₄, C₅, C₆, C₁₋₆, C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₅, C₂₋₄, C₂₋₃, C₃₋₆, C₃₋₅, C₃₋₄, C₄₋₆, C₄₋₅, and C₅₋₆.

[0026] The terms “purified,” “substantially purified,” and “isolated” refer to a compound useful in the present invention being free of other, dissimilar compounds with which the compound is normally associated in its natural state, so that the compound comprises at least 0.5%, 1%, 5%, 10%, 20%, 50%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.9% of the mass, by weight, of a given sample or composition. In one embodiment, these terms refer to the compound comprising at least 95%, 98%, 99%, or 99.9% of the mass, by weight, of a given sample or composition.

[0027] The term “acyl” refers to a group having the general formula $-\text{C}(=\text{O})\text{R}^{\text{X1}}$, $-\text{C}(=\text{O})\text{OR}^{\text{X1}}$, $-\text{C}(=\text{O})-\text{O}-\text{C}(=\text{O})\text{R}^{\text{X1}}$, $-\text{C}(=\text{O})\text{SR}^{\text{X1}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{X1}})_2$, $-\text{C}(=\text{S})\text{R}^{\text{X1}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{X1}})_2$, and $-\text{C}(=\text{S})\text{S}(\text{R}^{\text{X1}})$, $-\text{C}(=\text{NR}^{\text{X1}})\text{R}^{\text{X1}}$, $-\text{C}(=\text{NR}^{\text{X1}})\text{OR}^{\text{X1}}$, $-\text{C}(=\text{NR}^{\text{X1}})\text{SR}^{\text{X1}}$, and $-\text{C}(=\text{NR}^{\text{X1}})\text{N}(\text{R}^{\text{X1}})_2$.

$C(=NR^{X1})N(R^{X1})_2$, wherein R^{X1} is hydrogen; halogen; substituted or unsubstituted hydroxyl; substituted or unsubstituted thiol; substituted or unsubstituted amino; substituted or unsubstituted acyl, cyclic or acyclic, substituted or unsubstituted, branched or unbranched aliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched heteroaliphatic; cyclic or acyclic, substituted or unsubstituted, branched or unbranched alkyl; cyclic or acyclic, substituted or unsubstituted, branched or unbranched alkenyl; substituted or unsubstituted alkynyl; substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, mono- or di- aliphaticamino, mono- or di- heteroaliphaticamino, mono- or di- alkylamino, mono- or di- heteroalkylamino, mono- or di-arylamino, or mono- or di-heteroarylamino; or two R^{X1} groups taken together form a 5- to 6-membered heterocyclic ring. Exemplary acyl groups include aldehydes ($-CHO$), carboxylic acids ($-CO_2H$), ketones, acyl halides, esters, amides, imines, carbonates, carbamates, and ureas. Acyl substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0028] The term “acyloxy” refers to a “substituted hydroxyl” of the formula ($-OR^i$), wherein R^i is an optionally substituted acyl group, as defined herein, and the oxygen moiety is directly attached to the parent molecule.

[0029] The term “aliphatic” includes both saturated and unsaturated, nonaromatic, straight chain (*i.e.*, unbranched), branched, acyclic, and cyclic (*i.e.*, carbocyclic) hydrocarbons, which are optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, “aliphatic” is intended herein to include, but is not limited to, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, and cycloalkynyl moieties. Thus, the term “alkyl” includes straight, branched and cyclic alkyl groups. An analogous convention applies to other generic terms such as “alkenyl”, “alkynyl”, and the like. Furthermore, the terms “alkyl”, “alkenyl”, “alkynyl”, and the like encompass both substituted and unsubstituted groups. In certain embodiments, “aliphatic” is used to indicate those

aliphatic groups (cyclic, acyclic, substituted, unsubstituted, branched or unbranched) having 1–20 carbon atoms. Aliphatic group substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0030] The term “alkyl” refers to saturated, straight- or branched-chain hydrocarbon radicals derived from a hydrocarbon moiety containing between one and twenty carbon atoms by removal of a single hydrogen atom. In some embodiments, the alkyl group employed in the invention contains 1–20 carbon atoms. In another embodiment, the alkyl group employed contains 1–15 carbon atoms. In another embodiment, the alkyl group employed contains 1–10 carbon atoms. In another embodiment, the alkyl group employed contains 1–8 carbon atoms. In another embodiment, the alkyl group employed contains 1–5 carbon atoms. Examples of alkyl radicals include, but are not limited to, methyl (*e.g.*, unsubstituted methyl (Me)), ethyl (*e.g.*, unsubstituted ethyl (Et)), propyl (*e.g.*, unsubstituted propyl (Pr)), *n*-propyl, isopropyl, butyl (*e.g.*, unsubstituted butyl (Bu)), *n*-butyl, *iso*-butyl, *sec*-butyl, *sec*-pentyl, *iso*-pentyl, *tert*-butyl, *n*-pentyl, neopentyl, *n*-hexyl, *sec*-hexyl, *n*-heptyl, *n*-octyl, *n*-decyl, *n*-undecyl, dodecyl, and the like, which may bear one or more substituents. Alkyl group substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0031] The term “alkenyl” denotes a monovalent group derived from a straight- or branched-chain hydrocarbon moiety having at least one carbon-carbon double bond by the removal of a single hydrogen atom. In certain embodiments, the alkenyl group employed in the invention contains 2–20 carbon atoms. In some embodiments, the alkenyl group employed in the invention contains 2–15 carbon atoms. In another embodiment, the alkenyl

group employed contains 2–10 carbon atoms. In still other embodiments, the alkenyl group contains 2–8 carbon atoms. In yet other embodiments, the alkenyl group contains 2–5 carbons. Alkenyl groups include, for example, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like, which may bear one or more substituents. Alkenyl group substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0032] The term “alkynyl” refers to a monovalent group derived from a straight- or branched-chain hydrocarbon having at least one carbon-carbon triple bond by the removal of a single hydrogen atom. In certain embodiments, the alkynyl group employed in the invention contains 2–20 carbon atoms. In some embodiments, the alkynyl group employed in the invention contains 2–15 carbon atoms. In another embodiment, the alkynyl group employed contains 2–10 carbon atoms. In still other embodiments, the alkynyl group contains 2–8 carbon atoms. In still other embodiments, the alkynyl group contains 2–5 carbon atoms. Representative alkynyl groups include, but are not limited to, ethynyl, 2-propynyl (propargyl), 1-propynyl, and the like, which may bear one or more substituents. Alkynyl group substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0033] Exemplary carbon atom substituents include, but are not limited to, halogen, –CN, –NO₂, –N₃, –SO₂H, –SO₃H, –OH, –OR^{aa}, –ON(R^{bb})₂, –N(R^{bb})₂, –N(R^{bb})₃⁺X[–], –N(OR^{cc})R^{bb}, –SH, –SR^{aa}, –SSR^{cc}, –C(=O)R^{aa}, –CO₂H, –CHO, –C(OR^{cc})₂, –CO₂R^{aa}, –OC(=O)R^{aa}, –OCO₂R^{aa}, –C(=O)N(R^{bb})₂, –OC(=O)N(R^{bb})₂, –NR^{bb}C(=O)R^{aa}, –NR^{bb}CO₂R^{aa}, –NR^{bb}C(=O)N(R^{bb})₂, –C(=NR^{bb})R^{aa}, –C(=NR^{bb})OR^{aa}, –OC(=NR^{bb})R^{aa}, –OC(=NR^{bb})OR^{aa}, –

$C(=NR^{bb})N(R^{bb})_2$, $-OC(=NR^{bb})N(R^{bb})_2$, $-NR^{bb}C(=NR^{bb})N(R^{bb})_2$, $-C(=O)NR^{bb}SO_2R^{aa}$, $-NR^{bb}SO_2R^{aa}$, $-SO_2N(R^{bb})_2$, $-SO_2R^{aa}$, $-SO_2OR^{aa}$, $-OSO_2R^{aa}$, $-S(=O)R^{aa}$, $-OS(=O)R^{aa}$, $-Si(R^{aa})_3$, $-OSi(R^{aa})_3$, $-C(=S)N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=S)SR^{aa}$, $-SC(=S)SR^{aa}$, $-SC(=O)SR^{aa}$, $-OC(=O)SR^{aa}$, $-SC(=O)OR^{aa}$, $-SC(=O)R^{aa}$, $-P(=O)_2R^{aa}$, $-OP(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-OP(=O)(R^{aa})_2$, $-OP(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, $-OP(=O)_2N(R^{bb})_2$, $-P(=O)(NR^{bb})_2$, $-OP(=O)(NR^{bb})_2$, $-NR^{bb}P(=O)(OR^{cc})_2$, $-NR^{bb}P(=O)(NR^{bb})_2$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-OP(R^{cc})_2$, $-OP(R^{cc})_3$, $-B(R^{aa})_2$, $-B(OR^{cc})_2$, $-BR^{aa}(OR^{cc})$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

or two geminal hydrogens on a carbon atom are replaced with the group $=O$, $=S$, $=NN(R^{bb})_2$, $=NNR^{bb}C(=O)R^{aa}$, $=NNR^{bb}C(=O)OR^{aa}$, $=NNR^{bb}S(=O)_2R^{aa}$, $=NR^{bb}$, or $=NOR^{cc}$;

each instance of R^{aa} is, independently, selected from C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, or two R^{aa} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{bb} is, independently, selected from hydrogen, $-OH$, $-OR^{aa}$, $-N(R^{cc})_2$, $-CN$, $-C(=O)R^{aa}$, $-C(=O)N(R^{cc})_2$, $-CO_2R^{aa}$, $-SO_2R^{aa}$, $-C(=NR^{cc})OR^{aa}$, $-C(=NR^{cc})N(R^{cc})_2$, $-SO_2N(R^{cc})_2$, $-SO_2R^{cc}$, $-SO_2OR^{cc}$, $-SOR^{aa}$, $-C(=S)N(R^{cc})_2$, $-C(=O)SR^{cc}$, $-C(=S)SR^{cc}$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)_2N(R^{cc})_2$, $-P(=O)(NR^{cc})_2$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, or two R^{bb} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{cc} is, independently, selected from hydrogen, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, or two R^{cc} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{dd} is, independently, selected from halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{ee}$, $-\text{ON}(\text{R}^{ff})_2$, $-\text{N}(\text{R}^{ff})_2$, $-\text{N}(\text{R}^{ff})_3^+\text{X}^-$, $-\text{N}(\text{OR}^{ee})\text{R}^{ff}$, $-\text{SH}$, $-\text{SR}^{ee}$, $-\text{SSR}^{ee}$, $-\text{C}(=\text{O})\text{R}^{ee}$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^{ee}$, $-\text{OC}(=\text{O})\text{R}^{ee}$, $-\text{OCO}_2\text{R}^{ee}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{ff})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{ff})_2$, $-\text{NR}^{ff}\text{C}(=\text{O})\text{R}^{ee}$, $-\text{NR}^{ff}\text{CO}_2\text{R}^{ee}$, $-\text{NR}^{ff}\text{C}(=\text{O})\text{N}(\text{R}^{ff})_2$, $-\text{C}(=\text{NR}^{ff})\text{OR}^{ee}$, $-\text{OC}(=\text{NR}^{ff})\text{R}^{ee}$, $-\text{OC}(=\text{NR}^{ff})\text{OR}^{ee}$, $-\text{C}(=\text{NR}^{ff})\text{N}(\text{R}^{ff})_2$, $-\text{OC}(=\text{NR}^{ff})\text{N}(\text{R}^{ff})_2$, $-\text{NR}^{ff}\text{C}(=\text{NR}^{ff})\text{N}(\text{R}^{ff})_2$, $-\text{NR}^{ff}\text{SO}_2\text{R}^{ee}$, $-\text{SO}_2\text{N}(\text{R}^{ff})_2$, $-\text{SO}_2\text{R}^{ee}$, $-\text{SO}_2\text{OR}^{ee}$, $-\text{OSO}_2\text{R}^{ee}$, $-\text{S}(=\text{O})\text{R}^{ee}$, $-\text{Si}(\text{R}^{ee})_3$, $-\text{OSi}(\text{R}^{ee})_3$, $-\text{C}(=\text{S})\text{N}(\text{R}^{ff})_2$, $-\text{C}(=\text{O})\text{SR}^{ee}$, $-\text{C}(=\text{S})\text{SR}^{ee}$, $-\text{SC}(=\text{S})\text{SR}^{ee}$, $-\text{P}(=\text{O})_2\text{R}^{ee}$, $-\text{P}(=\text{O})(\text{R}^{ee})_2$, $-\text{OP}(=\text{O})(\text{R}^{ee})_2$, $-\text{OP}(=\text{O})(\text{OR}^{ee})_2$, C_{1-6} alkyl, C_{1-6} perhaloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3–10 membered heterocyclyl, C_{6-10} aryl, 5–10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups, or two geminal R^{dd} substituents can be joined to form $=\text{O}$ or $=\text{S}$;

each instance of R^{ee} is, independently, selected from C_{1-6} alkyl, C_{1-6} perhaloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, 3–10 membered heterocyclyl, and 3–10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups;

each instance of R^{ff} is, independently, selected from hydrogen, C_{1-6} alkyl, C_{1-6} perhaloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3–10 membered heterocyclyl, C_{6-10} aryl and 5–10 membered heteroaryl, or two R^{ff} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups; and

each instance of R^{gg} is, independently, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OC}_{1-6}$ alkyl, $-\text{ON}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{N}(\text{C}_{1-6}$ alkyl) $_3^+\text{X}^-$, $-\text{NH}(\text{C}_{1-6}$ alkyl) $_2^+\text{X}^-$, $-\text{NH}_2(\text{C}_{1-6}$ alkyl) $^+\text{X}^-$, $-\text{NH}_3^+\text{X}^-$, $-\text{N}(\text{OC}_{1-6}$ alkyl)(C_{1-6} alkyl), $-\text{N}(\text{OH})(\text{C}_{1-6}$ alkyl), $-\text{NH}(\text{OH})$, $-\text{SH}$, $-\text{SC}_{1-6}$ alkyl, $-\text{SS}(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{OCO}_2(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{OC}(=\text{O})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{N}(\text{C}_{1-6}$ alkyl) $\text{C}(=\text{O})(\text{C}_{1-6}$ alkyl), $-\text{NHCO}_2(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{NHC}(=\text{O})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{NHC}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{NH})\text{O}(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{NH})(\text{C}_{1-6}$ alkyl), $-\text{OC}(=\text{NH})\text{OC}_{1-6}$ alkyl, $-\text{C}(=\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{OC}(=\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{OC}(\text{NH})\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{OC}(\text{NH})\text{NH}_2$, $-\text{NHC}(\text{NH})\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{NHC}(=\text{NH})\text{NH}_2$, $-\text{NH}\text{SO}_2(\text{C}_{1-6}$ alkyl), $-\text{SO}_2\text{N}(\text{C}_{1-6}$ alkyl) $_2$, $-\text{SO}_2\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{C}_{1-6}$ alkyl, $-\text{SO}_2\text{OC}_{1-6}$ alkyl, $-\text{OSO}_2\text{C}_{1-6}$ alkyl, $-\text{SOC}_{1-6}$ alkyl, $-\text{Si}(\text{C}_{1-6}$ alkyl) $_3$, $-\text{OSi}(\text{C}_{1-6}$ alkyl) $_3$, $-\text{C}(=\text{S})\text{N}(\text{C}_{1-6}$ alkyl) $_2$,

C(=S)NH(C₁₋₆ alkyl), C(=S)NH₂, -C(=O)S(C₁₋₆ alkyl), -C(=S)SC₁₋₆ alkyl, -SC(=S)SC₁₋₆ alkyl, -P(=O)₂(C₁₋₆ alkyl), -P(=O)(C₁₋₆ alkyl)₂, -OP(=O)(C₁₋₆ alkyl)₂, -OP(=O)(OC₁₋₆ alkyl)₂, C₁₋₆ alkyl, C₁₋₆ perhaloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ carbocyclyl, C₆₋₁₀ aryl, 3-10 membered heterocyclyl, 5-10 membered heteroaryl; or two geminal R^{gg} substituents can be joined to form =O or =S; wherein X⁻ is a counterion.

[0034] The term “amino” refers to a group of the formula (-NH₂). A “substituted amino” refers either to a mono-substituted amine (-NHR^h) or a disubstituted amine (-NR^h₂), wherein the R^h substituent is any substituent as described herein that results in the formation of a stable moiety (*e.g.*, a suitable amino protecting group; aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, amino, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted). In certain embodiments, the R^h substituents of the disubstituted amino group(-NR^h₂) form a 5- to 6-membered heterocyclic ring.

[0035] The term “alkoxy” refers to a “substituted hydroxyl” of the formula (-ORⁱ), wherein Rⁱ is an optionally substituted alkyl group as defined herein, and the oxygen moiety is directly attached to the parent molecule.

[0036] The term “alkylthioxy” refers to a “substituted thiol” of the formula (-SR^f), wherein R^f is an optionally substituted alkyl group as defined herein, and the sulfur moiety is directly attached to the parent molecule.

[0037] The term “alkylamino” refers to a “substituted amino” of the formula (-NR^h₂), wherein R^h is, independently, a hydrogen or an optionally substituted alkyl group as defined herein, and the nitrogen moiety is directly attached to the parent molecule.

[0038] The term “aryl” refer to stable aromatic mono- or polycyclic ring system having 3-20 ring atoms, of which all the ring atoms are carbon, and which may be substituted or unsubstituted. In certain embodiments of the present invention, “aryl” refers to a mono, bi, or tricyclic C₄-C₂₀ aromatic ring system having one, two, or three aromatic rings which include, but not limited to, phenyl, biphenyl, naphthyl, and the like, which may bear one or more substituents. Aryl substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino,

alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0039] The term “arylalkyl” refers to an aryl substituted alkyl group, wherein the terms “aryl” and “alkyl” are defined herein, and wherein the aryl group is attached to the alkyl group, which in turn is attached to the parent molecule. Exemplary arylalkyl groups are benzyl and phenethyl.

[0040] The term “aryloxy” refers to a “substituted hydroxyl” of the formula ($-OR^i$), wherein R^i is an optionally substituted aryl group as defined herein, and the oxygen moiety is directly attached to the parent molecule.

[0041] The term “arylamino,” refers to a “substituted amino” of the formula ($-NR^h$), wherein R^h is, independently, a hydrogen or an optionally substituted aryl group as defined herein, and the nitrogen moiety is directly attached to the parent molecule.

[0042] The term “arylthioxy” refers to a “substituted thiol” of the formula ($-SR^f$), wherein R^f is an optionally substituted aryl group as defined herein, and the sulfur moiety is directly attached to the parent molecule.

[0043] The terms “halo” and “halogen” refer to an atom selected from fluorine (fluoro, $-F$), chlorine (chloro, $-Cl$), bromine (bromo, $-Br$), and iodine (iodo, $-I$).

[0044] The term “heteroaliphatic” refers to an aliphatic moiety, as defined herein, which includes both saturated and unsaturated, nonaromatic, straight chain (*i.e.*, unbranched), branched, acyclic, cyclic (*i.e.*, heterocyclic), or polycyclic hydrocarbons, which are optionally substituted with one or more functional groups, and that contain one or more oxygen, sulfur, nitrogen, phosphorus, or silicon atoms, *e.g.*, in place of carbon atoms. In certain embodiments, heteroaliphatic moieties are substituted by independent replacement of one or more of the hydrogen atoms thereon with one or more substituents. As will be appreciated by one of ordinary skill in the art, “heteroaliphatic” is intended herein to include, but is not limited to, heteroalkyl, heteroalkenyl, heteroalkynyl, heterocycloalkyl, heterocycloalkenyl, and heterocycloalkynyl moieties. Thus, the term “heteroaliphatic” includes the terms “heteroalkyl,” “heteroalkenyl,” “heteroalkynyl,” and the like. Furthermore, the terms “heteroalkyl,” “heteroalkenyl,” “heteroalkynyl,” and the like encompass both substituted and unsubstituted groups. In certain embodiments, “heteroaliphatic” is used to indicate those heteroaliphatic groups (cyclic, acyclic, substituted, unsubstituted, branched or unbranched) having 1–20 carbon atoms. Heteroaliphatic group substituents include, but are not limited to,

any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, sulfinyl, sulfonyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0045] The term “heteroalkyl” refers to an alkyl moiety, as defined herein, which contain one or more oxygen, sulfur, nitrogen, phosphorus, or silicon atoms, *e.g.*, in place of carbon atoms.

[0046] The term “heteroalkenyl” refers to an alkenyl moiety, as defined herein, which contain one or more oxygen, sulfur, nitrogen, phosphorus, or silicon atoms, *e.g.*, in place of carbon atoms.

[0047] The term “heteroalkynyl” refers to an alkynyl moiety, as defined herein, which contain one or more oxygen, sulfur, nitrogen, phosphorus, or silicon atoms, *e.g.*, in place of carbon atoms.

[0048] The term “heteroalkylamino” refers to a “substituted amino” of the formula ($-NR^h_2$), wherein R^h is, independently, a hydrogen or an optionally substituted heteroalkyl group, as defined herein, and the nitrogen moiety is directly attached to the parent molecule.

[0049] The term “heteroalkyloxy” refers to a “substituted hydroxyl” of the formula ($-OR^i$), wherein R^i is an optionally substituted heteroalkyl group, as defined herein, and the oxygen moiety is directly attached to the parent molecule.

[0050] The term “heteroalkylthioxy” refers to a “substituted thiol” of the formula ($-SR^f$), wherein R^f is an optionally substituted heteroalkyl group, as defined herein, and the sulfur moiety is directly attached to the parent molecule.

[0051] The term “carbocyclyl” or “carbocyclic” refers to a radical of a non-aromatic cyclic hydrocarbon group having from 3 to 14 ring carbon atoms (“C₃₋₁₄ carbocyclyl”) and zero heteroatoms in the non-aromatic ring system. In some embodiments, a carbocyclyl group has 3 to 10 ring carbon atoms (“C₃₋₁₀ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 8 ring carbon atoms (“C₃₋₈ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 7 ring carbon atoms (“C₃₋₇ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 6 ring carbon atoms (“C₃₋₆ carbocyclyl”). In some embodiments, a carbocyclyl group has 4 to 6 ring carbon atoms (“C₄₋₆ carbocyclyl”). In some embodiments, a

carbocyclyl group has 5 to 6 ring carbon atoms (“C₅₋₆ carbocyclyl”). In some embodiments, a carbocyclyl group has 5 to 10 ring carbon atoms (“C₅₋₁₀ carbocyclyl”). Exemplary C₃₋₆ carbocyclyl groups include, without limitation, cyclopropyl (C₃), cyclopropenyl (C₃), cyclobutyl (C₄), cyclobutenyl (C₄), cyclopentyl (C₅), cyclopentenyl (C₅), cyclohexyl (C₆), cyclohexenyl (C₆), cyclohexadienyl (C₆), and the like. Exemplary C₃₋₈ carbocyclyl groups include, without limitation, the aforementioned C₃₋₆ carbocyclyl groups as well as cycloheptyl (C₇), cycloheptenyl (C₇), cycloheptadienyl (C₇), cycloheptatrienyl (C₇), cyclooctyl (C₈), cyclooctenyl (C₈), bicyclo[2.2.1]heptanyl (C₇), bicyclo[2.2.2]octanyl (C₈), and the like. Exemplary C₃₋₁₀ carbocyclyl groups include, without limitation, the aforementioned C₃₋₈ carbocyclyl groups as well as cyclononyl (C₉), cyclononenyl (C₉), cyclodecyl (C₁₀), cyclodecenyl (C₁₀), octahydro-1*H*-indenyl (C₉), decahydronaphthalenyl (C₁₀), spiro[4.5]decanyl (C₁₀), and the like. As the foregoing examples illustrate, in certain embodiments, the carbocyclyl group is either monocyclic (“monocyclic carbocyclyl”) or polycyclic (*e.g.*, containing a fused, bridged or spiro ring system such as a bicyclic system (“bicyclic carbocyclyl”) or tricyclic system (“tricyclic carbocyclyl”)) and can be saturated or can contain one or more carbon-carbon double or triple bonds. “Carbocyclyl” also includes ring systems wherein the carbocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups wherein the point of attachment is on the carbocyclyl ring, and in such instances, the number of carbons continue to designate the number of carbons in the carbocyclic ring system. Unless otherwise specified, each instance of a carbocyclyl group is independently unsubstituted (an “unsubstituted carbocyclyl”) or substituted (a “substituted carbocyclyl”) with one or more substituents. In certain embodiments, the carbocyclyl group is an unsubstituted C₃₋₁₄ carbocyclyl. In certain embodiments, the carbocyclyl group is a substituted C₃₋₁₄ carbocyclyl.

[0052] In some embodiments, “carbocyclyl” is a monocyclic, saturated carbocyclyl group having from 3 to 14 ring carbon atoms (“C₃₋₁₄ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 10 ring carbon atoms (“C₃₋₁₀ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 8 ring carbon atoms (“C₃₋₈ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 6 ring carbon atoms (“C₃₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 4 to 6 ring carbon atoms (“C₄₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 5 to 6 ring carbon atoms (“C₅₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 5 to 10 ring carbon atoms (“C₅₋₁₀ cycloalkyl”). Examples of C₅₋₆ cycloalkyl groups include cyclopentyl (C₅) and cyclohexyl (C₅). Examples of C₃₋₆ cycloalkyl groups include the aforementioned C₅₋₆ cycloalkyl groups as well as cyclopropyl (C₃) and

cyclobutyl (C₄). Examples of C₃₋₈ cycloalkyl groups include the aforementioned C₃₋₆ cycloalkyl groups as well as cycloheptyl (C₇) and cyclooctyl (C₈). Unless otherwise specified, each instance of a cycloalkyl group is independently unsubstituted (an “unsubstituted cycloalkyl”) or substituted (a “substituted cycloalkyl”) with one or more substituents. In certain embodiments, the cycloalkyl group is an unsubstituted C₃₋₁₄ cycloalkyl. In certain embodiments, the cycloalkyl group is a substituted C₃₋₁₄ cycloalkyl.

[0053] The term “heterocyclic,” “heterocycles,” or “heterocyclyl” refers to a cyclic heteroaliphatic group. A heterocyclic group refers to a non-aromatic, partially unsaturated or fully saturated, 3- to 12-membered ring system, which includes single rings of 3 to 8 atoms in size, and bi- and tri-cyclic ring systems which may include aromatic five- or six-membered aryl or heteroaryl groups fused to a non-aromatic ring. These heterocyclic rings include those having from one to three heteroatoms independently selected from oxygen, sulfur, and nitrogen, in which the nitrogen and sulfur heteroatoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. In certain embodiments, the term heterocyclic refers to a non-aromatic 5-, 6-, or 7-membered ring or polycyclic group wherein at least one ring atom is a heteroatom selected from O, S, and N (wherein the nitrogen and sulfur heteroatoms may be optionally oxidized), and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms. Heterocyclyl groups include, but are not limited to, a bi- or tri-cyclic group, comprising fused five, six, or seven-membered rings having between one and three heteroatoms independently selected from the oxygen, sulfur, and nitrogen, wherein (i) each 5-membered ring has 0 to 2 double bonds, each 6-membered ring has 0 to 2 double bonds, and each 7-membered ring has 0 to 3 double bonds, (ii) the nitrogen and sulfur heteroatoms may be optionally oxidized, (iii) the nitrogen heteroatom may optionally be quaternized, and (iv) any of the above heterocyclic rings may be fused to an aryl or heteroaryl ring. Exemplary heterocycles include azacyclopropanyl, azacyclobutanyl, 1,3-diazatidinyl, piperidinyl, piperazinyl, azocanyl, thiaranyl, thietanyl, tetrahydrothiophenyl, dithiolanyl, thiacyclohexanyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropuranyl, dioxanyl, oxathiolanyl, morpholinyl, thioxanyl, tetrahydronaphthyl, and the like, which may bear one or more substituents. Substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, sulfinyl, sulfonyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy,

heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0054] The term “heteroaryl” refer to stable aromatic mono- or polycyclic ring system having 3-20 ring atoms, of which one ring atom is selected from S, O, and N; zero, one, or two ring atoms are additional heteroatoms independently selected from S, O, and N; and the remaining ring atoms are carbon, the radical being joined to the rest of the molecule via any of the ring atoms. Exemplary heteroaryls include, but are not limited to pyrrolyl, pyrazolyl, imidazolyl, pyridinyl (pyridyl), pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, tetrazinyl, ppyrolizinyl, indolyl, quinolinyl, isoquinolinyl, benzoimidazolyl, indazolyl, quinolinyl, isoquinolinyl, quinolizinyl, cinnolinyl, quinazolynyl, phthalazinyl, naphthridinyl, quinoxalinyl, thiophenyl, thianaphthenyl, furanyl, benzofuranyl, benzothiazolyl, thiazolynyl, isothiazolyl, thiadiazolynyl, oxazolyl, isoxazolyl, oxadiazolyl, oxadiazolyl, and the like, which may bear one or more substituents. Heteroaryl substituents include, but are not limited to, any of the substituents described herein, that result in the formation of a stable moiety (*e.g.*, aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, sulfinyl, sulfonyl, oxo, imino, thiooxo, cyano, isocyano, amino, azido, nitro, hydroxyl, thiol, halo, aliphaticamino, heteroaliphaticamino, alkylamino, heteroalkylamino, arylamino, heteroarylamino, alkylaryl, arylalkyl, aliphaticoxy, heteroaliphaticoxy, alkyloxy, heteroalkyloxy, aryloxy, heteroaryloxy, aliphaticthioxy, heteroaliphaticthioxy, alkylthioxy, heteroalkylthioxy, arylthioxy, heteroarylthioxy, acyloxy, and the like, each of which may or may not be further substituted).

[0055] The term “heteroarylamino” refers to a “substituted amino” of the $(-NR^h_2)$, wherein R^h is, independently, hydrogen or an optionally substituted heteroaryl group, as defined herein, and the nitrogen moiety is directly attached to the parent molecule.

[0056] The term “heteroaryloxy” refers to a “substituted hydroxyl” of the formula $(-OR^i)$, wherein R^i is an optionally substituted heteroaryl group, as defined herein, and the oxygen moiety is directly attached to the parent molecule.

[0057] The term “heteroarylthioxy” refers to a “substituted thiol” of the formula $(-SR^f)$, wherein R^f is an optionally substituted heteroaryl group, as defined herein, and the sulfur moiety is directly attached to the parent molecule.

[0058] The term “hydroxy” or “hydroxyl” refers to a group of the formula $(-OH)$. A “substituted hydroxyl” refers to a group of the formula $(-OR^i)$, wherein R^i can be any substituent which results in a stable moiety (*e.g.*, a suitable hydroxyl protecting group;

aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, nitro, alkylaryl, arylalkyl, and the like, each of which may or may not be further substituted).

[0059] The term “imino” refers to a group of the formula ($=NR^f$), wherein R^f corresponds to hydrogen or any substituent as described herein, that results in the formation of a stable moiety (for example, a suitable amino protecting group; aliphatic, alkyl, alkenyl, alkynyl, heteroaliphatic, heterocyclic, aryl, heteroaryl, acyl, amino, hydroxyl, alkylaryl, arylalkyl, and the like, each of which may or may not be further substituted). In certain embodiments, imino refers to $=NH$ wherein R^f is hydrogen.

[0060] The term “nitro” refers to a group of the formula ($-NO_2$).

[0061] The term “oxo” refers to a group of the formula ($=O$).

[0062] A “protecting group” is well known in the art and include those described in detail in *Greene's Protective Groups in Organic Synthesis*, P. G. M. Wuts and T. W. Greene, 4th edition, Wiley-Interscience, 2006, the entirety of which is incorporated herein by reference. Suitable “amino-protecting groups” (also referred to as “nitrogen protecting groups”) include methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-trimethylsilylethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate (TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl carbamate (*t*-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(*N,N*-dicyclohexylcarboxamido)ethyl carbamate, *t*-butyl carbamate (BOC or Boc), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, *N*-hydroxypiperidinyl carbamate, alkylthio carbamate, benzyl carbamate (Cbz), *p*-methoxybenzyl carbamate (Moz), *p*-nitrobenzyl carbamate, *p*-bromobenzyl carbamate, *p*-chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonylethyl carbamate, 2-(*p*-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-

triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, *m*-chloro-*p*-acyloxybenzyl carbamate, *p*-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolylmethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Tcroc), *m*-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, *o*-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(*o*-nitrophenyl)methyl carbamate, phenothiazinyl-(10)-carbonyl derivative, *N'*-*p*-toluenesulfonylaminocarbonyl derivative, *N'*-phenylaminothiocarbonyl derivative, *t*-amyl carbamate, *S*-benzyl thiocarbamate, *p*-cyanobenzyl carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, *p*-decyloxybenzyl carbamate, 2,2-dimethoxycarbonylvinyl carbamate, *o*-(*N,N*-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-furanylmethyl carbamate, 2-iodoethyl carbamate, isoborynl carbamate, isobutyl carbamate, isonicotinyl carbamate, *p*-(*p'*-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(*p*-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, *p*-(phenylazo)benzyl carbamate, 2,4,6-tri-*t*-butylphenyl carbamate, 4-(trimethylammonium)benzyl carbamate, 2,4,6-trimethylbenzyl carbamate, formamide, acetamide, chloroacetamide, trichloroacetamide, trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, benzamide, *p*-phenylbenzamide, *o*-nitrophenylacetamide, *o*-nitrophenoxyacetamide, acetoacetamide, (*N'*-dithiobenzyloxycarbonylamino)acetamide, 3-(*p*-hydroxyphenyl)propanamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-nitrophenoxy)propanamide, 2-methyl-2-(*o*-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, *o*-nitrocinnamide, *N*-acetylmethionine derivative, *o*-nitrobenzamide, *o*-(benzoyloxymethyl)benzamide, 4,5-diphenyl-3-oxazolin-2-one, *N*-phthalimide, *N*-dithiasuccinimide (Dts), *N*-2,3-diphenylmaleimide, *N*-2,5-dimethylpyrrole, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, *N*-methylamine, *N*-allylamine, *N*-[2-(trimethylsilyl)ethoxy]methylamine (SEM), *N*-3-acetoxypropylamine, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyroolin-3-yl)amine, quaternary ammonium salts, *N*-benzylamine, *N*-di(4-methoxyphenyl)methylamine, *N*-5-dibenzosuberylamine, *N*-triphenylmethylamine (Tr), *N*-

[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), *N*-9-phenylfluorenylamine (PhF), *N*-2,7-dichloro-9-fluorenylmethyleneamine, *N*-ferrocenylmethylamino (Fcm), *N*-2-picolylamino *N'*-oxide, *N*-1,1-dimethylthiomethyleneamine, *N*-benzylideneamine, *N*-*p*-methoxybenzylideneamine, *N*-diphenylmethyleneamine, *N*-[(2-pyridyl)mesityl]methyleneamine, *N*-(*N'*,*N'*-dimethylaminomethylene)amine, *N*,*N'*-isopropylidenediamine, *N*-*p*-nitrobenzylideneamine, *N*-salicylideneamine, *N*-5-chlorosalicylideneamine, *N*-(5-chloro-2-hydroxyphenyl)phenylmethyleneamine, *N*-cyclohexylideneamine, *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)amine, *N*-borane derivative, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentacarbonylchromium- or tungsten)carbonyl]amine, *N*-copper chelate, *N*-zinc chelate, *N*-nitroamine, *N*-nitrosoamine, amine *N*-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, *o*-nitrobenzenesulfenamide (Nps), 2,4-dinitrobenzenesulfenamide, pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, 3-nitropyridinesulfenamide (Npys), *p*-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilylethanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacysulfonamide.

[0063] A “hydroxyl protecting group” (also referred to as an “oxygen protecting group”) is well known in the art and includes those described in detail in Greene (1999). Suitable hydroxyl protecting groups include methyl, methoxymethyl (MOM), methylthiomethyl (MTM), *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl (SMOM), benzyloxymethyl (BOM), *p*-methoxybenzyloxymethyl (PMBM), (4-methoxyphenoxy)methyl (*p*-AOM), guaiacolmethyl (GUM), *t*-butoxymethyl, 4-pentenylloxymethyl (POM), siloxymethyl, 2-methoxyethoxymethyl (MEM), 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl (SEMOR), tetrahydropyranyl (THP), 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-

methoxytetrahydropyranyl (MTHP), 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl (CTMP), 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl, 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, benzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p,p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl, α -naphthyldiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxyphenyl)diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-yl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, benzisothiazolyl S,S-dioxido, trimethylsilyl (TMS), triethylsilyl (TES), triisopropylsilyl (TIPS), dimethylisopropylsilyl (IPDMS), diethylisopropylsilyl (DEIPS), dimethylhexylsilyl, *t*-butyldimethylsilyl (TBDMS), *t*-butyldiphenylsilyl (TBDPS), tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl (DPMS), *t*-butylmethoxyphenylsilyl (TBMPS), formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio)pentanoate (levulinoyldithioacetal), pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate (mesitoate), alkyl methyl carbonate, 9-fluorenylmethyl carbonate (Fmoc), alkyl ethyl carbonate, alkyl 2,2,2-trichloroethyl carbonate (Troc), 2-(trimethylsilyl)ethyl carbonate (TMSEC), 2-(phenylsulfonyl) ethyl carbonate (Psec), 2-(triphenylphosphonio) ethyl carbonate (Peoc), alkyl isobutyl carbonate, alkyl vinyl carbonate alkyl allyl carbonate, alkyl *p*-nitrophenyl carbonate, alkyl benzyl carbonate, alkyl *p*-methoxybenzyl carbonate, alkyl 3,4-dimethoxybenzyl carbonate, alkyl *o*-nitrobenzyl carbonate, alkyl *p*-nitrobenzyl carbonate, alkyl *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl carbonate, methyl dithiocarbonate, 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-

(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxy)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenolate, *o*-(methoxycarbonyl)benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, alkyl *N*-phenylcarbamate, borate, dimethylphosphinothioyl, alkyl 2,4-dinitrophenylsulfenate, sulfate, methanesulfonate (mesylate), benzylsulfonate, and tosylate (Ts). For protecting 1,2- or 1,3-diols, protecting groups include methylene acetal, ethylidene acetal, 1-*t*-butylethylidene ketal, 1-phenylethylidene ketal, (4-methoxyphenyl)ethylidene acetal, 2,2,2-trichloroethylidene acetal, acetonide, cyclopentylidene ketal, cyclohexylidene ketal, cycloheptylidene ketal, benzylidene acetal, *p*-methoxybenzylidene acetal, 2,4-dimethoxybenzylidene ketal, 3,4-dimethoxybenzylidene acetal, 2-nitrobenzylidene acetal, methoxymethylene acetal, ethoxymethylene acetal, dimethoxymethylene ortho ester, 1-methoxyethylidene ortho ester, 1-ethoxyethylidene ortho ester, 1,2-dimethoxyethylidene ortho ester, α -methoxybenzylidene ortho ester, 1-(*N,N*-dimethylamino)ethylidene derivative, α -(*N,N'*-dimethylamino)benzylidene derivative, 2-oxacyclopentylidene ortho ester, di-*t*-butylsilylene group (DTBS), 1,3-(1,1,3,3-tetraisopropylidisiloxanylidene) derivative (TIPDS), tetra-*t*-butoxydisiloxane-1,3-diylidene derivative (TBDS), cyclic carbonates, cyclic boronates, ethyl boronate, and phenyl boronate.

[0064] In certain embodiments, the substituent present on a sulfur atom is a sulfur protecting group (also referred to as a “thiol protecting group”). Sulfur protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, and $-P(=O)(NR^{bb})_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein. Sulfur protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[0065] The term “carbohydrate” or “saccharide” refers to an aldehydic or ketonic derivative of polyhydric alcohols. Carbohydrates include compounds with relatively small molecules (*e.g.*, sugars) as well as macromolecular or polymeric substances (*e.g.*, starch, glycogen, and cellulose polysaccharides). The term “sugar” refers to monosaccharides, disaccharides, or polysaccharides. Monosaccharides are the simplest carbohydrates in that

they cannot be hydrolyzed to smaller carbohydrates. Most monosaccharides can be represented by the general formula $C_yH_{2y}O_y$ (*e.g.*, $C_6H_{12}O_6$ (a hexose such as glucose)), wherein y is an integer equal to or greater than 3. Certain polyhydric alcohols not represented by the general formula described above may also be considered monosaccharides. For example, deoxyribose is of the formula $C_5H_{10}O_4$ and is a monosaccharide. Monosaccharides usually consist of five or six carbon atoms and are referred to as pentoses and hexoses, receptively. If the monosaccharide contains an aldehyde it is referred to as an aldose; and if it contains a ketone, it is referred to as a ketose. Monosaccharides may also consist of three, four, or seven carbon atoms in an aldose or ketose form and are referred to as trioses, tetroses, and heptoses, respectively. Glyceraldehyde and dihydroxyacetone are considered to be aldotriose and ketotriose sugars, respectively. Examples of aldotetrose sugars include erythrose and threose; and ketotetrose sugars include erythrulose. Aldopentose sugars include ribose, arabinose, xylose, and lyxose; and ketopentose sugars include ribulose, arabulose, xylulose, and lyxulose. Examples of aldohexose sugars include glucose (for example, dextrose), mannose, galactose, allose, altrose, talose, gulose, and idose; and ketohexose sugars include fructose, psicose, sorbose, and tagatose. Ketoheptose sugars include sedoheptulose. Each carbon atom of a monosaccharide bearing a hydroxyl group ($-OH$), with the exception of the first and last carbons, is asymmetric, making the carbon atom a stereocenter with two possible configurations (R or S). Because of this asymmetry, a number of isomers may exist for any given monosaccharide formula. The aldohexose D-glucose, for example, has the formula $C_6H_{12}O_6$, of which all but two of its six carbons atoms are stereogenic, making D-glucose one of the 16 (*i.e.*, 2^4) possible stereoisomers. The assignment of D or L is made according to the orientation of the asymmetric carbon furthest from the carbonyl group: in a standard Fischer projection if the hydroxyl group is on the right the molecule is a D sugar, otherwise it is an L sugar. The aldehyde or ketone group of a straight-chain monosaccharide will react reversibly with a hydroxyl group on a different carbon atom to form a hemiacetal or hemiketal, forming a heterocyclic ring with an oxygen bridge between two carbon atoms. Rings with five and six atoms are called furanose and pyranose forms, respectively, and exist in equilibrium with the straight-chain form. During the conversion from the straight-chain form to the cyclic form, the carbon atom containing the carbonyl oxygen, called the anomeric carbon, becomes a stereogenic center with two possible configurations: the oxygen atom may take a position either above or below the plane of the ring. The resulting possible pair of stereoisomers is called anomers. In an α anomer, the $-OH$ substituent on the anomeric carbon rests on the opposite side (*trans*) of the ring from the $-$

CH₂OH side branch. The alternative form, in which the –CH₂OH substituent and the anomeric hydroxyl are on the same side (*cis*) of the plane of the ring, is called a β anomer. A carbohydrate including two or more joined monosaccharide units is called a disaccharide or polysaccharide (*e.g.*, a trisaccharide), respectively. The two or more monosaccharide units bound together by a covalent bond known as a glycosidic linkage formed via a dehydration reaction, resulting in the loss of a hydrogen atom from one monosaccharide and a hydroxyl group from another. Exemplary disaccharides include sucrose, lactulose, lactose, maltose, isomaltose, trehalose, cellobiose, xylobiose, laminaribiose, gentiobiose, mannobiose, melibiose, nigerose, and rutinose. Exemplary trisaccharides include, but are not limited to, isomaltotriose, nigerotriose, maltotriose, melezitose, maltotriulose, raffinose, and kestose. The term carbohydrate also includes other natural or synthetic stereoisomers of the carbohydrates described herein.

[0066] The term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and other animals without undue toxicity, irritation, allergic response, and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, Berge *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1–19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting the appropriate compound in the form of the free base with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, L-ascorbate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, formate, fumarate, gentisate, glutarate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, malonate, DL-mandelate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphonate, picrate, pivalate, propionate, pyroglutamate, succinate, sulfonate, tartrate, L-tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, *para*-toluenesulfonate (*p*-tosylate), and undecanoate. Also, basic groups in the compounds disclosed herein can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and

benzyl and phenethyl bromides. Examples of acids which can be employed to form therapeutically acceptable salts include inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, and phosphoric acid; and organic acids such as oxalic acid, maleic acid, succinic acid, and citric acid. “Basic addition salts” refer to salts derived from appropriate bases, these salts including alkali metal, alkaline earth metal, and quaternary amine salts. Hence, the present invention contemplates sodium, potassium, magnesium, and calcium salts of the compounds disclosed herein, and the like. Basic addition salts can be prepared during the final isolation and purification of the compounds, often by reacting a carboxyl group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of therapeutically acceptable salts include lithium, sodium (by using, *e.g.*, NaOH), potassium (by using, *e.g.*, KOH), calcium (by using, *e.g.*, Ca(OH)₂), magnesium (by using, *e.g.*, Mg(OH)₂ and magnesium acetate), zinc, (by using, *e.g.*, Zn(OH)₂ and zinc acetate), and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, *N,N*-dimethylaniline, *N*-methylpiperidine, *N*-methyldmorpholine, dicyclohexylamine, procaine, dibenzylamine, *N,N*-dibenzylphenethylamine, 1-phenamine, and *N,N*-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine, choline hydroxide, hydroxyethyl morpholine, hydroxyethyl pyrrolidone, imidazole, *n*-methyl-*D*-glucamine, *N,N'*-dibenzylethylenediamine, *N,N'*-diethylethanolamine, *N,N'*-dimethylethanolamine, triethanolamine, and tromethamine. Basic amino acids (*e.g.*, 1-glycine and 1-arginine) and amino acids which may be zwitterionic at neutral pH (*e.g.*, betaine (*N,N,N*-trimethylglycine)) are also contemplated.

[0067] The term “tautomer” refers to a particular isomer of a compound in which a hydrogen and double bond have changed position with respect to the other atoms of the molecule. For a pair of tautomers to exist there must be a mechanism for interconversion. Examples of tautomers include keto-enol forms, imine-enamine forms, amide-imino alcohol forms, amidine-aminidine forms, nitroso-oxime forms, thio ketone-enethiol forms, *N*-nitroso-hydroxyazo forms, nitro-*aci*-nitro forms, lactam-lactim forms, ketene-ynol forms, enamine-enamine forms, and pyridione-hydroxypyridine forms.

[0068] The term “polymorphs” refers to a crystalline form of a compound (or a salt, hydrate, or solvate thereof). All polymorphs have the same elemental composition. Different

crystalline forms usually have different X-ray diffraction patterns, infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other factors may cause one crystal form to dominate. Various polymorphs of a compound can be prepared by crystallization under different conditions.

[0069] The term “solvate” refers to forms of the compound, or a salt thereof, that are associated with a solvent, usually by a solvolysis reaction. This physical association may include hydrogen bonding. Conventional solvents include water, methanol, ethanol, acetic acid, DMSO, THF, diethyl ether, and the like. The compounds of the invention may be prepared, *e.g.*, in crystalline form, and may be solvated. Suitable solvates include pharmaceutically acceptable solvates and further include both stoichiometric solvates and non-stoichiometric solvates. In certain instances, the solvate will be capable of isolation, for example, when one or more solvent molecules are incorporated in the crystal lattice of a crystalline solid. “Solvate” encompasses both solution-phase and isolable solvates. Representative solvates include hydrates, ethanولات, and methanولات.

[0070] The term “hydrate” refers to a compound that is associated with water. Typically, the number of the water molecules contained in a hydrate of a compound is in a definite ratio to the number of the compound molecules in the hydrate. Therefore, a hydrate of a compound may be represented, for example, by the general formula $R \cdot xH_2O$, wherein R is the compound and wherein x is a number greater than 0. A given compound may form more than one type of hydrates, including, *e.g.*, monohydrates (x is 1), lower hydrates (x is a number greater than 0 and smaller than 1, *e.g.*, hemihydrates ($R \cdot 0.5H_2O$)), and polyhydrates (x is a number greater than 1, *e.g.*, dihydrates ($R \cdot 2H_2O$) and hexahydrates ($R \cdot 6H_2O$)).

[0071] The term “subject” refers to any animal. In certain embodiments, the subject is a mammal. In certain embodiments, the subject is a human (*e.g.*, a man, a woman, or a child). The human may be of either sex and may be at any stage of development. In certain embodiments, the subject has been diagnosed with the condition or disease to be treated. In other embodiments, the subject is at risk of developing the condition or disease. In certain embodiments, the subject is an experimental animal (*e.g.*, mouse, rat, rabbit, dog, pig, or primate). The experimental animal may be genetically engineered. In certain embodiments, the subject is a domesticated animal (*e.g.*, dog, cat, bird, horse, cow, goat, sheep).

[0072] The terms “administer,” “administering,” or “administration” refers to implanting, absorbing, ingesting, injecting, inhaling, or otherwise introducing an inventive compound, or a pharmaceutical composition thereof.

[0073] The terms “treatment,” “treat,” and “treating” refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a “pathological condition” (*e.g.*, a disease, disorder, or condition, or one or more signs or symptoms thereof) described herein. In some embodiments, treatment may be administered after one or more signs or symptoms have developed or have been observed. In other embodiments, treatment may be administered in the absence of signs or symptoms of the disease or condition. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (*e.g.*, in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example, to delay or prevent recurrence.

[0074] The terms “condition,” “disease,” and “disorder” are used interchangeably.

[0075] An “effective amount” of a compound of the present invention or a pharmaceutical composition thereof refers to an amount sufficient to elicit the desired biological response, *i.e.*, treating the condition. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound of the invention may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the condition being treated, the mode of administration, and the age and health of the subject. In certain embodiments, an effective amount is a therapeutically effective amount. In certain embodiments, an effective amount is a prophylactic treatment. In certain embodiments, an effective amount is the amount of a compound described herein in a single dose. In certain embodiments, an effective amount is the combined amounts of a compound described herein in multiple doses.

[0076] A “therapeutically effective amount” of a compound of the present invention or a pharmaceutical composition thereof is an amount sufficient to provide a therapeutic benefit in the treatment of a condition, *e.g.*, iron overload, or to delay or minimize one or more symptoms associated with the condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment of the condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of the condition, and/or enhances the therapeutic efficacy of another therapeutic agent. In certain embodiments, a therapeutically effective amount is an amount sufficient for chelating a metal described herein. In certain

embodiments, a therapeutically effective amount is an amount sufficient for treating a pathological condition described herein. In certain embodiments, a therapeutically effective amount is an amount sufficient for chelating a metal described herein and for treating a pathological condition described herein.

[0077] A “prophylactically effective amount” of a compound of the present invention is an amount sufficient to prevent a condition, *e.g.*, iron overload, or one or more symptoms associated with the condition or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the condition. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent. In certain embodiments, a prophylactically effective amount is an amount sufficient for chelating a metal described herein. In certain embodiments, a prophylactically effective amount is an amount sufficient for preventing a pathological condition described herein. In certain embodiments, a prophylactically effective amount is an amount sufficient for chelating a metal described herein and for preventing a pathological condition described herein.

[0078] The term “biological sample” refers to any sample including tissue samples (such as tissue sections and needle biopsies of a tissue); cell samples (*e.g.*, cytological smears (such as Pap or blood smears) or samples of cells obtained by microdissection); samples of whole organisms (such as samples of yeasts or bacteria); or cell fractions, fragments or organelles (such as obtained by lysing cells and separating the components thereof by centrifugation or otherwise). Other examples of biological samples include blood, serum, urine, semen, fecal matter, cerebrospinal fluid, interstitial fluid, mucous, tears, sweat, pus, biopsied tissue (*e.g.*, obtained by a surgical biopsy or needle biopsy), nipple aspirates, milk, vaginal fluid, saliva, swabs (such as buccal swabs), or any material containing biomolecules that is derived from a first biological sample.

[0079] “Chelation,” “chelating,” “sequestration,” or “sequestering” is the formation or presence of two or more separate coordinate bonds between a polydentate (multiple-bonded) compound and a single central atom. The polydentate compound is typically an organic compound and referred to as a “chelator,” “chelant,” “chelating agent,” “sequestrator,” “sequestering agent,” or “ligand.” The central atom is usually a metal atom or metal ion (*e.g.*, a metal atom or metal ion described herein, such as iron (*e.g.*, Fe(III)), Al(III), chromium (*e.g.*, Cr(III) or Cr(VI)), and uranium (*e.g.*, U(VI)), *etc.*). The chelator may form a

stable complex with the central atom through coordinate bonds, inactivating the central atom so that the central atom is less likely to react with other molecules or atoms.

[0080] The term “metal-clearing efficiency” or “MCE” refers to the efficacy of a given concentration of chelator in clearing a metal atom or metal ion (*e.g.*, a metal atom or metal ion described herein, such as iron (*e.g.*, Fe(III)), Al(III), chromium (*e.g.*, Cr(III) or Cr(VI)), and uranium (*e.g.*, U(VI)) from the body or one of its organs or parts.

Efficaciousness in turn concerns quantity of the metal atom or metal ion removed from a target system (*e.g.*, a whole body, an organ, or a tissue) in a unit of time. Chelators of a metal atom or metal ion are needed in one or more of three clinical situations: (1) for acute metal toxicity from ingestion or infusion of the metal atom or metal ion ; (2) to reduce total body metal secondary to transfusion or excess metal absorption; and (3) for the maintenance of metal balance after total body metal has been satisfactorily reduced and only daily dietary metal needs to be excreted. In practical terms, therefore, for chronic metal overload secondary to transfusion, the recommendation is that 0.3-0.5 mg/kg/d (*i.e.*, mg iron per kg body weight of the subject per day) need be excreted. For the maintenance treatment, 0.25-1 mg/kg/d is sufficient. Other ranges are also possible. In certain embodiments, the metal-clearing efficiency is iron-clearing efficiency or “ICE.” In certain embodiments, the metal-clearing efficiency is aluminum-clearing efficiency. In certain embodiments, the metal-clearing efficiency is chromium-clearing efficiency. In certain embodiments, the metal-clearing efficiency is uranium-clearing efficiency.

[0081] The term “focal iron overload” refers to any disease or condition that involves the accumulation of unmanaged iron in a tissue or organ. Focal iron overload typically involves less than the subject’s whole body but may involve more than one organ or tissue. Unmanaged iron in any tissue or organ is typically undesired and can be the focus of the treatments of the present invention. The treatment may involve the removal of as much iron as possible from the tissue or organ or may only involve the removal of excess iron. Examples of disease and conditions associated with focal iron overload include, but are not limited to, macular degeneration, IBD, reperfusion injury, stroke including hemorrhagic stroke, and closed head injury; however, any disease or condition of focal iron overload may be treated as described herein. In certain embodiments, the term “focal iron overload” does not include diseases or conditions associated with global iron overload (*e.g.*, global iron overload associated with chronic transfusion therapy, hereditary hemochromatosis, *etc.*). The treatment of focal iron overload may be systemic or local administration of an effective amount of an inventive compound, or a pharmaceutical composition thereof.

[0082] The term “reactive oxygen species” or “ROS” refers to molecules or ions formed by the incomplete reduction of oxygen. Reactive oxygen species include superoxide anion ($O_2^{\bullet-}$), peroxides such as hydrogen peroxide (H_2O_2), hydroxyl radical ($HO^{\bullet}$), and hypochlorous acid ($HClO$). These molecules are typically chemically reactive. Reactive oxygen species may be formed by any number of mechanisms (*e.g.*, enzymatically, by ionizing radiation, by reaction oxygen with a metal). In certain embodiments, the reactive oxygen species are formed by the reduction of oxygen by an iron ion, such as Fe^{+2} .

[0083] “Primary hemochromatosis” is a genetic disorder characterized by excessive iron accumulation that results in tissue damage. Manifestations include systemic symptoms, liver disorders, cardiomyopathy, diabetes, erectile dysfunction, and arthropathy. Normal total body iron content is about 2.5 g in women and 3.5 g in men. Because symptoms may be delayed until iron accumulation is excessive, hemochromatosis may not be recognized until total body iron content is > 10 g, or often several times greater. In women, clinical manifestations are uncommon before menopause because iron loss due to menses (and sometimes pregnancy and childbirth) tends to offset iron accumulation. One mechanism for iron overload is increased iron absorption from the gastrointestinal tract, leading to chronic deposition of iron in the tissues. Hepcidin, a liver-derived peptide, is the critical control mechanism for iron absorption. Hepcidin, along with the normal HFE gene, prevents excessive iron absorption and storage in normal people. Tissue injury in a subject with primary hemochromatosis may result from reactive free hydroxyl radicals generated when iron deposition in tissues catalyzes their formation. Other mechanisms may affect particular organs (*e.g.*, skin hyperpigmentation can result from increased melanin as well as iron accumulation).

[0084] “Secondary hemochromatosis” is a condition acquired as a consequence of another disease that causes iron overload, or blood transfusions, or both, and typically characterized by increased hepatic and total body iron content and unequivocal portal cirrhosis of the liver. Secondary hemochromatosis is usually caused by disorders of erythropoiesis (*e.g.*, thalassemia, sickle cell anemia, X-linked sideroblastic anemia, pyruvate kinase deficiency, hereditary spherocytosis, and congenital dyserythropoietic anemia (CDA)) and the treatment of these diseases with blood transfusions. After damaging the transfused erythrocytes by macrophages, iron freed from the heme is accumulated in the body (*e.g.*, in the liver, heart, or skin).

[0085] “Diabetes” or “diabetes mellitus” is a metabolic disorder in which there are high levels of glucose in the blood. Diabetes can be caused by insufficient amount of insulin

(a hormone produced by the pancreas to control blood glucose) or resistance to insulin in a subject, or both. There are three major types of diabetes: Type 1, Type 2, and gestational diabetes. Type 1 diabetes is usually diagnosed in children and young adults, and was previously known as juvenile diabetes. In type 1 diabetes, the body does not produce insulin, which may be a result of the destruction of islet cells in the pancreas. Type 2 diabetes, or non-insulin-dependent diabetes mellitus (NIDDM) or adult-onset diabetes, is the most common form of diabetes. Type 2 diabetes is characterized by high blood glucose in the context of insulin resistance and relative insulin deficiency. This is in contrast to Type 1 diabetes in which there is an absolute insulin deficiency. Obesity is thought to be one of the primary causes of Type 2 diabetes in subjects who are genetically predisposed to obesity. Gestational diabetes is characterized by high blood glucose that develops during pregnancy in a woman who does not have diabetes prior to the pregnancy. Gestational diabetes may be caused by various pregnancy hormones that may interfere with the body's response to insulin.

[0086] The term “closed head injury” refers to any injury to the head that does not penetrate the skull. Closed head injuries may result from falls, blasts, accidents including vehicular accidents, and assaults. Closed head injuries can lead to hemorrhage or brain swelling, which can result in increased intracranial pressure, which can in turn lead to permanent brain damage or even death. Various types of closed head injury include concussions, brain contusions, diffuse axonal injury, and hematomas.

[0087] “Thalassemia” is a group of inherited autosomal recessive blood disorders that originated in the Mediterranean region. In a subject with thalassemia, the genetic defect, which could be either mutation or deletion, results in reduced rate of synthesis or no synthesis of one of the globin chains that make up hemoglobin. This can cause the formation of abnormal hemoglobin molecules, thus causing anemia. There are two main types of thalassemia: alpha and beta thalassemias. Alpha thalassemia occurs when a gene or genes related to the alpha globin protein are missing or changed (*i.e.*, mutated). Beta thalassemia occurs when similar gene defects affect production of the beta globin protein. Each of alpha and beta thalassemias includes two forms: thalassemia major and thalassemia minor. Beta thalassemia major is also referred to as Cooley's anemia or Mediterranean anemia.

[0088] “Friedreich's ataxia” or “FRDA” is an inherited disease that causes progressive damage to the nervous system of a subject resulting in symptoms including muscle weakness, speech problems, and heart disease. In a subject with Friedreich's ataxia, the spinal cord and peripheral nerves degenerate and become thinner. The cerebellum, part of the brain that coordinates balance and movement, also degenerates to a lesser extent. This

damage results in awkward, unsteady movements and impaired sensory functions.

Friedreich's ataxia also causes problems in the heart and spine, and some subjects with the condition develop diabetes. However, this disorder usually does not affect cognitive functions, such as thinking and reasoning. Friedreich's ataxia is caused by a defect, which may be a result of mutation, in a gene labeled as FXN. This disorder is recessive, meaning it occurs only in someone who inherits two defective copies of the gene, one from each parent.

[0089] "Macular degeneration" is a disease that affects the retina of a subject. The retina is a thin tissue lining the back of the eye. Light-sensitive cells in the retina are responsible for converting light into electrical impulses, which are then sent via the optic nerve to the brain for interpretation. In the center of the retina is the macula. The macula contains the highest concentration of the light-sensitive cells, called cones, which are responsible for sharp, detailed, and central vision. In macular degeneration, cells in the macular region begin to die, which results in blind spots and distorted vision. Macular degeneration is the leading cause of vision loss in humans over the age of 60. There are two forms of macular degeneration: dry and wet macular degenerations. It is possible for a subject to suffer from both forms, for it to affect one or both eyes, and for the disease to progress slowly or rapidly. Dry macular degeneration is the most common type of macular degeneration, in which the photosensitive cells of the macula slowly break down. Yellow deposits called drusen (extracellular waste products from metabolism) form and accumulate under the retina between the retinal pigmented epithelium (RPE) layer and the Bruch's membrane, which supports the retina. Over time, drusen are associated with deterioration of the macula and the death of RPE and photoreceptor cells, resulting in a blurring or spotty loss of clear, straight-ahead vision. This process does not cause any pain. In the early stages of the disease, the subject may notice slightly blurry vision. However, as more and more of the cells die, central vision worsens. Dry macular degeneration may advance and cause loss of vision without turning into the wet form of the disease. However, it is also possible for the early-stage dry form to change into the wet form of macular degeneration. Wet macular degeneration occurs when abnormal blood vessels grow behind the macula as RPE and photoreceptor cells die. The Bruch's membrane begins to break down, usually near drusen deposits, and new blood vessels grow. These vessels are very fragile and can leak fluid and blood. Scarring of and severe damage to the macula may result. Straight-ahead vision can become distorted or lost entirely in a short period of time.

BRIEF DESCRIPTION OF THE DRAWINGS

[0090] *Figure 1* shows the chemical structures of certain chelating compounds.

Figure 1A illustrates that the desferrithiocin (DFT, **1**) has been structurally modified to give desazadesferrithiocin (DADFT) analogs **2-5**. *Figure 1B* shows the chemical structures of additional desazadesferrithiocin analogs (**7** and **9**) and a few examples of inventive desferrithiocin analogs (**I-1**, **I-2**, and **I-3**).

[0091] *Figure 2* shows a Job's plot of the Fe(III) complex of DFT analogs **I-1**, **I-2**, and **I-3**. Solutions containing DFT analog/Fe(III) at different ratios were prepared such that [DFT analog] + [Fe(III)] = 1.0 mM in Tris-HCl buffer at pH 7.4. The theoretical mole fraction maximum for a 2:1 DFT analog:Fe complex is 0.667 (indicated by an arrow). The observed maxima for **I-1**, **I-2**, and **I-3** are 0.669, 0.676, and 0.677, respectively. Optical density (y-axis) was determined at 498, 484, and 485 nm for **I-1**, **I-2**, and **I-3**, respectively.

[0092] *Figure 3* shows the biliary ferrokinetics of DFT analogs (**I-1**, **I-2**, and **I-3**) and DADFT analogs (**3**, **7**, and **9**) in bile duct-cannulated rats. The compounds were given po at 300 $\mu\text{mol/kg}$.

[0093] *Figure 4* is a plot showing the iron-clearing efficiency versus $\log P_{\text{app}}$ for DFT analogs **1**, **I-1**, **I-2**, and **I-3** in bile duct-cannulated rats. Compound **1** was given po at a dose of 150 $\mu\text{mol/kg}$; compounds **I-1**, **I-2**, and **I-3** were administered po at 300 $\mu\text{mol/kg}$.

[0094] *Figure 5* includes bar graphs showing the urinary excretion of kidney injury molecule-1 (Kim-1), expressed as Kim-1 (ng/kg/24 h) of rats treated with a DFT analog (**1** (*Figure 5A*), **I-1** (*Figure 5C*), **I-2** (*Figure 5D*), or **I-3** (*Figure 5F*)) or a DADFT analog (**3** (*Figure 5B*) or **7** (*Figure 5E*)). The rats were given the compounds po twice daily (BID) at a dose of 237 $\mu\text{mol/kg/dose}$ (474 $\mu\text{mol/kg/d}$) for up to 7 d. None of the rats survived the planned 7-d exposure to **1**. $N = 5$ for **1**, **I-1**, **7**, **I-2**, and **I-3**; $N = 3$ for compound **3**.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

[0095] Various desazadesferrithiocin analogs have been described for use in the treatment of iron overload, which results from transfusion therapy, high-iron diet, acute iron ingestion, or malabsorption. Such analogs may also be used to treat focal iron overload, where the local concentration of iron in a particular tissue or organ contributes to the pathological process. For instance, the unmanaged Fe^{+2} ions in a tissue or organ may result in the production of hydroxyl radicals or other reactive oxygen species that lead to tissue or cell damage. Structural modification of a desazadesferrithiocin analog by replacing the phenyl

ring of the analog with a pyridinyl ring gives rise to a desferrithiocin analog, such as desferrithiocin (**1**). Structural modification of **1** by, among other things, attaching one or more carbohydrate (*e.g.*, a sugar, such as glucose, including α -D-, β -D-, α -L-, and β -L-glucose) moieties, optionally through linkers, to **1** gives rise to novel desferrithiocin analogs of Formula (**A**) or (**J**). These inventive compounds may have one or more superior properties (such as greater solubility, permeability, and bioavailability; improved distribution, absorption, metabolism, and iron-clearing efficiency; and reduced clearance, excretion, and toxicity) compared with the parent compound **1** and/or other desferrithiocin analogs. The inventive compounds may also be efficiently delivered into cells or taken up by cells and be retained inside cells, which is desired for the treatment and/or prevention of pathological conditions in a subject using the inventive compounds. For example, the carbohydrate moieties are hydrophilic, and the inventive compounds bearing these moieties may be more soluble and/or have a greater ability to get into a cell. Moreover, the carbohydrate moieties may be recognized by membrane transport proteins that lead to the uptake of the inventive compounds into cells. As a result, the inventive compounds with carbohydrate moieties attached may be more efficiently transported into the cells of a subject. Additionally, the linker connecting the carbohydrate moiety and desferrithiocin **1** may be hydrolyzed in a cell to give rise to a desferrithiocin analog without a carbohydrate moiety. This analog may no longer be recognized by membrane transport proteins and, therefore, may be retained inside the cell. This analog may also remain in the cell because the analog is too polar to pass through the cell membrane to get out of the cell. Any linkers capable of hydrolysis under physiological conditions may be used in the present invention. For example, when –polyether– (*e.g.*, a PEG moiety) is employed as the linker, the compound of the invention is of the formula:

Desferrithiocin–polyether–carbohydrate.

In certain embodiments, one of the oxygen atoms of the polyether linker is attached to the anomeric carbon (*i.e.*, C1) of the carbohydrate moiety. The polyether–C1 bond may hydrolyze under physiological conditions, and a hydrolysis product desferrithiocin–polyether–H, which is an alcohol, may be generated.

[0096] In certain embodiments, –polyether–NHC(=O)O– is used as the linker, and the compound of the invention is of the formula:

Desferrithiocin–polyether–NHC(=O)O–carbohydrate.

The carbamate moiety --NHC(=O)O-- may hydrolyze under physiological conditions, and a positively-charged hydrolysis product desferrithiocin–polyether– NH_3^+ may be formed.

[0097] Desferrithiocin analogs of Formulae (A) and (J) are expected to be useful in the treatment and/or prevention of a wide range of pathological conditions, including, but not limited to, metal overload (*e.g.*, iron overload, aluminum overload, chromium overload, magnesium overload, calcium overload, strontium overload, nickel overload, manganese overload, cobalt overload, copper overload, zinc overload, silver overload, sodium overload, potassium overload, cadmium overload, mercury overload, lead overload, molybdenum overload, tungsten overload, or actinide overload (*e.g.*, uranium overload)), metal poisoning (*e.g.*, iron poisoning, aluminum poisoning, thallium poisoning, chromium poisoning, magnesium poisoning, calcium poisoning, strontium poisoning, nickel poisoning, manganese poisoning, cobalt poisoning, copper poisoning, zinc poisoning, silver poisoning, sodium poisoning, potassium poisoning, cadmium poisoning, mercury poisoning, lead poisoning, antimony poisoning, molybdenum poisoning, tungsten poisoning, lanthanide poisoning (*e.g.*, cerium poisoning), or actinide poisoning (*e.g.*, uranium poisoning)), oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, and reperfusion injury. Without wishing to be bound by any particular theory, the compounds of the invention are thought to chelate a metal (*e.g.*, iron, aluminum, thallium, chromium, magnesium, calcium, strontium, nickel, manganese, cobalt, copper, zinc, silver, sodium, potassium, cadmium, mercury, lead, antimony, molybdenum, tungsten, a lanthanide (*e.g.*, cerium), or an actinide (*e.g.*, uranium)). The inventive compounds may prevent iron from participating in the generation of reactive oxygen species. Moreover, the inventive compounds may act as free radical scavengers thereby limiting the damage of reactive oxygen species or other radicals.

[0098] The inventive compounds may also be useful in the treatment and/or prevention of infectious diseases (*e.g.*, malaria). Infectious diseases are caused by microbes such as bacteria, fungi, and parasites. These pathogenic microbes typically require one or more metals (*e.g.*, iron, calcium, magnesium, strontium, potassium, sodium, chromium, copper, manganese, molybdenum, zinc, and tungsten) to sustain life. For example, iron is used by cytochromes and as a cofactor for enzymes in electron-transport proteins. Without wishing to be bound by any particular theory, the compounds of the invention are thought to deprive the microbes of the iron needed for their metabolic processes by chelating iron.

[0099] The invention, therefore, provides novel carbohydrate-modified desferrithiocin analogs; as well as methods of treating and/or preventing pathological conditions, pharmaceutical compositions, uses, and kits, each involving the inventive compounds.

Compounds

[00100] Desferrithiocin (DFT) **1** (*Figure 1A*) is a natural product iron chelator isolated from *Streptomyces antibioticus* (Naegeli *et al.*, “Metabolites of Microorganisms. Part 193. Ferrithiocin.” *Helv. Chim. Acta* **1980**, 63, 1400-1406). It forms a 2:1 complex with Fe(III) with a cumulative formation constant of $4 \times 10^{29} \text{ M}^{-1}$ (Hahn *et al.*, “Coordination Chemistry of Microbial Iron Transport. 42. Structural and Spectroscopic Characterization of Diastereomeric Cr(III) and Co(III) Complexes of Desferriferrithiocin.” *J. Am. Chem. Soc.* **1990**, 112, 1854-1860; Anderegg *et al.*, “Metal Complex Formation of a New Siderophore Desferrithiocin and of Three Related Ligands.” *J. Chem. Soc., Chem. Commun.* **1990**, 1194-1196). Although the compound was shown to be an excellent deferration agent when administered orally (po) to rats (Bergeron *et al.*, “Evaluation of Desferrithiocin and Its Synthetic Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1991**, 34, 2072-2078) and primates (Bergeron *et al.*, “A Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393; Wolfe *et al.*, “A Non-Human Primate Model for the Study of Oral Iron Chelators.” *Br. J. Haematol.* **1989**, 72, 456-461), it caused severe nephrotoxicity in rats (Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, 81, 2166-2173). However, the compound’s oral activity spurred SAR studies focused on the DFT platform aimed at identifying an orally active and safe DFT analog (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Methoxylation of Desazadesferrithiocin Analogs: Enhanced Iron Clearing Efficiency.” *J. Med. Chem.* **2003**, 46, 1470-1477; Bergeron *et al.*, “Desazadesmethyldesferrithiocin Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1999**, 42, 95-108). Various desazadesferrithiocin analogs have been developed that effectively chelate and remove metals from biological systems. See International PCT Application Publications, WO 1997/036885, published October 9, 1997; WO 2000/016763, published March 30, 2000; WO 2000/012493, published March 9, 2000; WO 2004/017959, published March 4, 2004; WO 2005/034949, published April 21, 2005; WO 2005/023310, published March 17, 2005; WO 2006/107626,

published October 12, 2006; WO 2008/130395, published October 30, 2008; WO 2008/115433, published September 25, 2008; WO 2011/028255, published March 10, 2011; WO 2013/090750, published June 20, 2013; and WO 2013/090766, published June 20, 2013; each of which is incorporated herein by reference. Also see U.S. Patent Nos. US 5,840,739; US 6,864,270; US 7,144,904; US 7,879,886, US RE39,132; US 6,083,966; US 6,521,652; US 6,525,080; US 6,559,315; US 8,278,458; and US 8,324,397; each of which is incorporated herein by reference. Also see U.S. Patent Application Publications, US 2004/044220, US 2004/132789, US 2005/234113, US 2008/255081, US 2006/211746, US 2006/211773, US 2008/096974, US 2013/030028, US 2010/137346, US 2013/210870, and US 2012/184586, each of which is incorporated herein by reference.

Removal of the pyridine nitrogen of **1** provided **2** (*Figure 1A*), the parent compound of the desazadesferrithiocin (DADFT) series (Bergeron *et al.*, “Desazadesmethyldesferrithiocin Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1999**, 42, 95-108). Interestingly, although **2** was not overtly nephrotoxic, it elicited serious gastrointestinal (GI) problems (Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, 81, 2166-2173; Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Desazadesmethyldesferrithiocin Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1999**, 42, 95-108). In spite of its GI toxicity, the compound's excellent iron-clearing efficiency (ICE) and the absence of nephrotoxicity prompted further SAR studies predicated on this pharmacophore. This led to the discovery that the lipophilicity (partition between octanol and water, expressed as the log of the fraction in the octanol layer, $\log P_{app}$) (Sangster, *Octanol-Water Partition Coefficients: Fundamentals and Physical Chemistry*; John Wiley and Sons: West Sussex, England, 1997; Vol. 2) of the DADFT analogs could have a profound effect on the compound's ICE, organ distribution, and toxicity profile (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Iron Chelators and Therapeutic Uses.” In: Abraham, ed. *Burger's Medicinal Chemistry*. 6th. Wiley; New York: **2003**. pp. 479-561; Bergeron *et al.*, “Desferrithiocin Analogs and Nephrotoxicity.” *J. Med. Chem.* **2008**, 51, 5993-6004). Desferrithiocin analogs have been reported to chelate and remove iron or other metals. See International PCT Application Publications, WO 1997/036885, published October 9, 1997;

WO 2000/016763, published March 30, 2000; WO 2000/012493, published March 9, 2000; and WO 2004/017959, published March 4, 2004; each of which is incorporated herein by reference. Also see U.S. Patent Nos., US 5,840,739; US 6,864,270; US 7,144,904; US 7,879,886; US RE39,132; US 6,083,966; US 6,521,652; US 6,525,080; and US 6,559,315; each of which is incorporated herein by reference. Also see U.S. Patent Application Publications, US 2004/044220, US 2004/132789, US 2005/234113, and US 2008/255081, each of which is incorporated herein by reference.


[00101] Ultimately, it was determined that hydroxylation of DADFT and a number of different analogs at the 3'-, 4'-, or 5'-position allowed for compounds that were very efficient, orally active iron chelators with less toxicity than **1** or **2** (Bergeron *et al.*, "Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs." *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, "Iron Chelators and Therapeutic Uses." In: Abraham, ed. *Burger's Medicinal Chemistry*. 6th. Wiley; New York: **2003**. pp. 479-561). Hydroxylation had a significant effect on toxicity reduction. One of these compounds, **3**, was studied in human clinical trials by Genzyme (Galanello *et al.*, "A Dose Escalation Study of the Pharmacokinetics, Safety, and Efficacy of Deferitritin, an Oral Iron Chelator in Beta Thalassaemia Patients." *ASH Annu. Meet. Abstr.* **2007**, 110, 2669).

[00102] Compound **3** was reengineered, by introducing a 3,6,9-trioxadecyloxy group at the 4'-position, to yield **4** (*Figure 1A*) (Bergeron *et al.*, "(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity." *J. Med. Chem.* **2006**, 49, 2772-2783). This provided a remarkably efficient orally active iron chelator which, given to rats orally once or twice daily, was virtually nephrotoxicity-free (Bergeron *et al.*, "(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity." *J. Med. Chem.* **2006**, 49, 2772-2783). This turned out to also be true when a variety of polyether backbones were fixed at the 3'-, 4'-, or 5'-position of the DADFT pharmacophore (Bergeron, R. J.; Wiegand, J.; Bharti, N.; Singh, S.; Rocca, J. R. Impact of 3,6,9-Trioxadecyloxy Group on Desazadesferrithiocin Analog Iron Chelators and Organ Distribution. *J. Med. Chem.* **2007**, 50, 3302-3313; Bergeron *et al.*, "Design, Synthesis, and Testing of Non-Nephrotoxic Desazadesferrithiocin Polyether Analogs." *J. Med. Chem.* **2008**, 51, 3913-3923; Bergeron *et al.*, "The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs." *J. Med. Chem.* **2010**, 53, 2843-2853;


Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258). In fact, (*S*)-4,5-dihydro-2-[2-hydroxy-3-(3,6,9-trioxadecyloxy)]-4-methyl-4-thiazolecarboxylic acid ((*S*)-3'-(HO)-DADFT-PE, **5**; *Figure 1A*) has now been moved forward to clinical trials. Thus, it appeared as though fixing a polyether fragment to the DADFT framework was also a uniformly effective tool in further reducing the nephrotoxicity induced by **3**.

[00103] Although DFT and DADFT analogs as a class of compounds appear promising as metal chelating agents, much work remains to be done to improve these compounds' physiochemical, pharmacokinetic, pharmacodynamic, and/or toxicological properties, such as absorption, distribution, metal-clearing efficiency, and toxicity, for the purpose of providing safe and effective compounds for a better treatment and/or prevention of pathological conditions in a subject. Provided by the present invention are novel DFT analogs that include one or more carbohydrate (*e.g.*, a sugar, such as glucose, including α -D-, β -D-, α -L-, and β -L-glucose) moieties. The carbohydrate moieties may be attached directly to or through a linker moiety at different positions on the parent compound DFT **1**, for example, at the 2'-, 3'-, 4'-, and/or 5'-position on the pyridinyl ring, and/or at the carboxyl group. The compounds of the invention may be useful in the treatment and/or prevention of a variety of pathological conditions.


[00104] In one aspect of the present invention, provided are compounds of Formula (A):



and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof, wherein:



R_1 is hydrogen, alkyl, acyl, an oxygen protecting group,



R_2 is hydrogen, alkyl, acyl, an oxygen protecting group, $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-$
 R'' , or $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-(CH_2)_n-NR_{10}-C(=O)O-R''$;


each occurrence of R_3 is independently alkyl, arylalkyl, or $-OR_8$;

R_4 is hydrogen or alkyl;



R_5 is hydrogen or alkyl;

R_6 is hydrogen or alkyl;

R_7 is $-OR_9$ or $-SR_9$;



R_8 is hydrogen, alkyl, acyl, an oxygen protecting group,





R_9 is hydrogen, alkyl, , an oxygen
 protecting group when attached to an oxygen atom, or a sulfur protecting group when
 attached to a sulfur atom;

R_{10} is hydrogen, alkyl, acyl, or a nitrogen protecting group;

R' is hydrogen or an oxygen protecting group;

R'' is hydrogen, alkyl, acyl, an oxygen protecting group,




each occurrence of n is independently an integer from 1 to 8, inclusive;


k is an integer from 0 to 2, inclusive;

x is an integer from 1 to 8, inclusive; and


y is an integer from 0 to 8, inclusive.



[00105] In certain embodiments, at least one of R_1 and R_9 is



; or R_2 is $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-R''$ or $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-$



[00106] In certain embodiments, R_1 is hydrogen, alkyl, acyl, or an oxygen protecting group;

R_2 is hydrogen, alkyl, acyl, an oxygen protecting group, $-\text{[(CH}_2)_n\text{-O]}_x\text{[(CH}_2)_n\text{-O]}_y\text{-R''}$, or $-\text{[(CH}_2)_n\text{-O]}_x\text{[(CH}_2)_n\text{-O]}_y\text{-(CH}_2)_n\text{-NR}_{10}\text{-C(=O)O-R''}$;

each occurrence of R_3 is independently alkyl, arylalkyl, or $-\text{OR}_8$;

R_4 , R_5 , and R_6 are each independently hydrogen or alkyl;

R_7 is $-\text{OR}_9$ or $-\text{SR}_9$;

R_8 is hydrogen, alkyl, acyl, or an oxygen protecting group;

R_9 is hydrogen, alkyl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

R_{10} is hydrogen, alkyl, acyl, or a nitrogen protecting group;

R' is hydrogen or an oxygen protecting group;

R'' is hydrogen, alkyl, acyl, or an oxygen protecting group;


each occurrence of n is independently an integer from 1 to 8, inclusive;

k is an integer from 0 to 2, inclusive;






x is an integer from 1 to 8, inclusive; and


y is an integer from 0 to 8, inclusive.

[00107] In compounds of Formula (A), R_1 is hydrogen, alkyl, acyl, an oxygen




is hydrogen. In certain embodiments, R_1 is alkyl. In certain embodiments, R_1 is C_{1-6} alkyl. In certain embodiments, R_1 is methyl. In certain embodiments, R_1 is ethyl. In certain embodiments, R_1 is propyl. In certain embodiments, R_1 is butyl. In certain embodiments, R_1 is acyl. In certain embodiments, R_1 is acetyl. In certain embodiments, R_1 is pivaloyl. In certain embodiments, R_1 is an oxygen protecting group. In certain embodiments, R_1 is silyl. In certain embodiments, R_1 is TBDPS, TBDMS, TIPS, TES, or TMS. In certain embodiments, R_1 is MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain






In certain embodiments, R_1 is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain




embodiments, R_1 is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R_1




is

. In certain embodiments, R_1 is




. In certain




embodiments, R_1 is

. In certain embodiments, R_1 is




. In




certain embodiments, R_1 is


. In certain embodiments, R_1 is




; wherein all R' are oxygen protecting groups. In certain embodiments, R_1



is ; wherein all R' are oxygen protecting groups. In certain embodiments,



R₁ is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R₁ is





; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz.


[00108] In compounds of Formula (A), R₂ is hydrogen, alkyl, acyl, an oxygen protecting group, $-\text{[(CH}_2\text{)}_n\text{-O]}_x\text{[(CH}_2\text{)}_n\text{-O]}_y\text{-R''}$, or $-\text{[(CH}_2\text{)}_n\text{-O]}_x\text{[(CH}_2\text{)}_n\text{-O]}_y\text{-(CH}_2\text{)}_n\text{-NR}_{10}\text{-C(=O)O-R''}$. In certain embodiments, R₂ is hydrogen. In certain embodiments, R₂ is alkyl. In certain embodiments, R₂ is C₁₋₁₂ alkyl optionally substituted with one or more substituents independently selected from the group consisting of hydroxyl, alkyloxy, and $\text{-CO}_2\text{H}$. In certain embodiments, R₂ is $\text{-(CH}_2\text{)}_2\text{-OH}$, $\text{-(CH}_2\text{)}_3\text{-OH}$, $\text{-(CH}_2\text{)}_4\text{-OH}$, $\text{-(CH}_2\text{)}_5\text{-OH}$, $\text{-(CH}_2\text{)}_6\text{-OH}$, $\text{-(CH}_2\text{)}_7\text{-OH}$, or $\text{-(CH}_2\text{)}_8\text{-OH}$. In certain embodiments, R₂ is $\text{-(CH}_2\text{)}_2\text{-OCH}_3$, $\text{-(CH}_2\text{)}_3\text{-OCH}_3$, $\text{-(CH}_2\text{)}_4\text{-OCH}_3$, $\text{-(CH}_2\text{)}_5\text{-OCH}_3$, $\text{-(CH}_2\text{)}_6\text{-OCH}_3$, $\text{-(CH}_2\text{)}_7\text{-OCH}_3$, or $\text{-(CH}_2\text{)}_8\text{-OCH}_3$. In certain embodiments, R₂ is $\text{-CH}_2\text{-CO}_2\text{H}$, $\text{-(CH}_2\text{)}_2\text{-CO}_2\text{H}$, $\text{-(CH}_2\text{)}_3\text{-CO}_2\text{H}$, $\text{-(CH}_2\text{)}_4\text{-CO}_2\text{H}$, $\text{-(CH}_2\text{)}_5\text{-CO}_2\text{H}$, $\text{-(CH}_2\text{)}_6\text{-CO}_2\text{H}$, or $\text{-(CH}_2\text{)}_7\text{-CO}_2\text{H}$. In certain embodiments, R₂ is C₁₋₁₂ alkyl substituted at least with $\text{-CO}_2\text{R}_{31}$, wherein R₃₁ is substituted or unsubstituted C₁₋₆ alkyl (*e.g.*, -CH_3). In certain embodiments, R₂ is $\text{-CH}_2\text{-CO}_2\text{R}_{31}$, $\text{-(CH}_2\text{)}_2\text{-CO}_2\text{R}_{31}$, $\text{-(CH}_2\text{)}_3\text{-CO}_2\text{R}_{31}$, $\text{-(CH}_2\text{)}_4\text{-CO}_2\text{R}_{31}$, $\text{-(CH}_2\text{)}_5\text{-CO}_2\text{R}_{31}$, $\text{-(CH}_2\text{)}_6\text{-CO}_2\text{R}_{31}$, or $\text{-(CH}_2\text{)}_7\text{-CO}_2\text{R}_{31}$. In certain embodiments, R₂ is C₁₋₆ alkyl. In certain embodiments, R₂ is methyl. In certain embodiments, R₂ is ethyl. In certain embodiments, R₂ is propyl. In certain embodiments, R₂ is butyl. In certain embodiments, R₂ is acyl. In certain embodiments, R₂ is acetyl. In certain embodiments, R₂ is pivaloyl. In certain embodiments, R₂ is an oxygen protecting group. In



certain embodiments, R_2 is silyl. In certain embodiments, R_2 is TBDPS, TBDMS, TIPS, TES, or TMS. In certain embodiments, R_2 is MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R_2 is $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-R''$. In certain embodiments, R_2 is





embodiments, R_2 is  R'' . In certain embodiments, R_2 is




certain embodiments, R_2 is  . In certain

embodiments, R_2 is  or  . In


certain embodiments, R_2 is  or




alkyl (e.g., $-CH_3$). In certain embodiments, R_2 is




embodiments, R_2 is



. In certain embodiments, R_2 is




. In certain embodiments, R_2 is




. In


certain embodiments, R_2 is



. In certain embodiments, R_2 is




; wherein all R' are oxygen protecting groups. In certain



embodiments, R_2 is


; wherein all R' are oxygen protecting



groups. In certain embodiments, R_2 is

; wherein all R' are silyl,


TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In



certain embodiments, R_2 is


; wherein all R' are silyl, TBDPS,

TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain




embodiments, R_2 is


. In certain embodiments, R_2 is




. In certain embodiments, R_2 is




In certain embodiments, R_2 is




. In certain embodiments, R_2 is




. In certain embodiments, R_2 is



wherein all R' are oxygen protecting groups. In certain embodiments, R_2 is




; wherein all R' are oxygen protecting groups. In certain



embodiments, R_2 is

; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain











embodiments, R_2 is





; wherein all R' are silyl, TBDPS, TBDMS,

TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain

embodiments, R_2 is $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-(CH_2)_n-NR_{10}-C(=O)O-R''$. In certain


embodiments, R_2 is $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-(CH_2)_n-NH-C(=O)O-R''$. In certain






groups. In certain embodiments, R_2 is

all R' are oxygen protecting groups. In certain embodiments, R_2 is




TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain




embodiments, R₂ is

TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In




certain embodiments, R₂ is

. In certain




embodiments, R₂ is

. In certain




embodiments, R₂ is

. In certain




embodiments, R₂ is

. In certain embodiments,




R₂ is


. In certain embodiments, R₂ is



. In certain embodiments, R_2 is




; wherein all R' are oxygen protecting



groups. In certain embodiments, R_2 is


; wherein

all R' are oxygen protecting groups. In certain embodiments, R_2 is



; wherein all R' are silyl, TBDPS, TBDMS,

TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain



embodiments, R_2 is


; wherein all R' are silyl,

TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz.


[00109] In certain embodiments, both R_1 and R_2 are hydrogen.

[00110] In compounds of Formula (A), each occurrence of R_3 is independently alkyl, arylalkyl, or $-\text{OR}_8$. In certain embodiments, at least one occurrence of R_3 is alkyl. In certain embodiments, at least one occurrence of R_3 is C_{1-6} alkyl. In certain embodiments, at least one


occurrence of R_3 is methyl. In certain embodiments, at least one occurrence of R_3 is ethyl. In certain embodiments, at least one occurrence of R_3 is propyl. In certain embodiments, at least one occurrence of R_3 is butyl. In certain embodiments, at least one occurrence of R_3 is arylalkyl. In certain embodiments, at least one occurrence of R_3 is benzyl. In certain embodiments, at least one occurrence of R_3 is $-OR_8$. In certain embodiments, at least one occurrence of R_3 is $-OH$. In certain embodiments, at least one occurrence of R_3 is $-O$ -alkyl. In certain embodiments, at least one occurrence of R_3 is $-O$ -(C_{1-6} alkyl). In certain embodiments, at least one occurrence of R_3 is $-OMe$. In certain embodiments, at least one occurrence of R_3 is $-OEt$. In certain embodiments, at least one occurrence of R_3 is $-OPr$. In certain embodiments, at least one occurrence of R_3 is $-OBu$. In certain embodiments, at least one occurrence of R_3 is $-O$ -acyl. In certain embodiments, at least one occurrence of R_3 is $-O-C(=O)-CH_3$. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is



. In certain embodiments, at least one occurrence of R₃ is




; wherein all R' are oxygen protecting groups. In certain embodiments,




at least one occurrence of R₃ is

; wherein all R' are oxygen protecting




groups. In certain embodiments, at least one occurrence of R₃ is

wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, at least one occurrence of R₃ is




; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM,

THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, at least one occurrence




of R₃ is


. In certain embodiments, at least one occurrence of R₃ is




. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is




. In certain embodiments, at least one occurrence of R_3 is



. In certain embodiments, at least one occurrence of R_3 is




; wherein all R' are oxygen protecting groups. In certain embodiments,




at least one occurrence of R_3 is

; wherein all R' are oxygen protecting



groups. In certain embodiments, at least one occurrence of R_3 is

wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, at least one occurrence of R_3 is



; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM,

THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz.

[00111] In compounds of Formula (A), R_4 is hydrogen or alkyl. In certain embodiments, R_4 is hydrogen. In certain embodiments, R_4 is alkyl. In certain embodiments, R_4 is C_{1-6} alkyl. In certain embodiments, R_4 is methyl. In certain embodiments, R_4 is ethyl. In certain embodiments, R_4 is propyl. In certain embodiments, R_4 is butyl.

[00112] In compounds of Formula (A), R_5 is hydrogen or alkyl. In certain embodiments, R_5 is hydrogen. In certain embodiments, R_5 is alkyl. In certain embodiments, R_5 is C_{1-6} alkyl. In certain embodiments, R_5 is methyl. In certain embodiments, R_5 is ethyl. In certain embodiments, R_5 is propyl. In certain embodiments, R_5 is butyl.

[00113] In certain embodiments, R_4 and R_5 are each hydrogen. In certain embodiments, R_4 and R_5 are each alkyl. In certain embodiments, R_4 and R_5 are each C_{1-6} alkyl. In certain embodiments, R_4 and R_5 are each methyl. In certain embodiments, R_4 and R_5 are each ethyl.

[00114] In compounds of Formula (A), R_6 is hydrogen or alkyl. In certain embodiments, R_6 is hydrogen. In certain embodiments, R_6 is alkyl. In certain embodiments, R_6 is C_{1-6} alkyl. In certain embodiments, R_6 is methyl. In certain embodiments, R_6 is ethyl. In certain embodiments, R_6 is propyl. In certain embodiments, R_6 is butyl.


[00115] In certain embodiments, R_4 and R_5 are each hydrogen; and R_6 is alkyl. In certain embodiments, R_4 and R_5 are each hydrogen; and R_6 is C_{1-6} alkyl. In certain embodiments, R_4 and R_5 are each hydrogen; and R_6 is methyl. In certain embodiments, R_4 and R_5 are each hydrogen; and R_6 is ethyl. In certain embodiments, R_4 and R_5 are each

hydrogen; and R_6 is propyl. In certain embodiments, R_4 and R_5 are each hydrogen; and R_6 is butyl.


[00116] In certain embodiments, R_4 and R_5 are each alkyl; and R_6 is methyl. In certain embodiments, R_4 and R_5 are each C_{1-6} alkyl; and R_6 is methyl. In certain embodiments, R_4 and R_5 are each methyl; and R_6 is methyl. In certain embodiments, R_4 and R_5 are each ethyl; and R_6 is methyl.

[00117] In certain embodiments, R_4 and R_5 are each alkyl; and R_6 is hydrogen. In certain embodiments, R_4 and R_5 are each C_{1-6} alkyl; and R_6 is hydrogen. In certain embodiments, R_4 and R_5 are each methyl; and R_6 is hydrogen. In certain embodiments, R_4 and R_5 are each ethyl; and R_6 is hydrogen. In certain embodiments, R_4 , R_5 , and R_6 are each hydrogen.


[00118] In compounds of Formula (A), R_7 is $-OR_9$ or $-SR_9$. In certain embodiments, R_7 is $-OR_9$. In certain embodiments, R_7 is $-OH$. In certain embodiments, R_7 is $-O$ -alkyl. In certain embodiments, R_7 is $-O$ -(C_{1-6} alkyl). In certain embodiments, R_7 is $-OMe$. In certain embodiments, R_7 is $-OEt$. In certain embodiments, R_7 is $-OPr$. In certain embodiments, R_7 is



. In certain embodiments, R_7 is




. In certain




embodiments, R_7 is


. In certain embodiments, R_7 is





. In certain embodiments, R_7 is






. In certain


embodiments, R_7 is  ; wherein all R' are oxygen protecting groups. In


certain embodiments, R_7 is  ; wherein all R' are oxygen protecting

groups. In certain embodiments, R_7 is  ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain


embodiments, R_7 is  ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R_7

is  . In certain embodiments, R_7 is  . In certain


embodiments, R_7 is  . In certain embodiments, R_7 is



. In certain embodiments, R_7 is




. In certain




embodiments, R_7 is

; wherein all R' are oxygen protecting groups. In



certain embodiments, R_7 is


; wherein all R' are oxygen protecting



groups. In certain embodiments, R_7 is

; wherein all R' are silyl,


TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In



certain embodiments, R_7 is


; wherein all R' are silyl, TBDPS, TBDMS,

TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain




embodiments, R_7 is


. In certain embodiments, R_7 is



. In certain embodiments, R_7 is




. In certain




embodiments, R_7 is


. In certain embodiments, R_7 is



. In certain embodiments, R_7 is




; wherein all R'



are oxygen protecting groups. In certain embodiments, R_7 is


; wherein



all R' are oxygen protecting groups. In certain embodiments, R_7 is

;


wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl,



acetyl, pivaloyl, or Bz. In certain embodiments, R_7 is


; wherein all R'

are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl,




or Bz. In certain embodiments, R_7 is


. In certain embodiments, R_7 is



. In certain embodiments, R_7 is




. In certain




embodiments, R_7 is


. In certain embodiments, R_7 is



. In certain embodiments, R_7 is




; wherein all R'



are oxygen protecting groups. In certain embodiments, R_7 is


; wherein



all R' are oxygen protecting groups. In certain embodiments, R_7 is


;

wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl,




acetyl, pivaloyl, or Bz. In certain embodiments, R_7 is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz.


[00119] In compounds of Formula (A), R_8 is hydrogen, alkyl, acyl, an oxygen




protecting group, or . In certain embodiments, R_8 is hydrogen. In certain embodiments, R_8 is alkyl. In certain embodiments, R_8 is C_{1-6} alkyl. In certain embodiments, R_8 is methyl. In certain embodiments, R_8 is ethyl. In certain embodiments, R_8 is propyl. In certain embodiments, R_8 is butyl. In certain embodiments, R_8 is acyl. In certain embodiments, R_8 is acetyl. In certain embodiments, R_8 is pivaloyl. In certain embodiments, R_8 is an oxygen protecting group. In certain embodiments, R_8 is silyl. In certain embodiments, R_8 is TBDPS, TBDMS, TIPS, TES, or TMS. In certain embodiments, R_8 is MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain



embodiments, R_8 is




. In certain embodiments, R_8 is




In certain embodiments, R_8 is


. In certain embodiments, R_8 is



. In certain embodiments, R₈ is




. In certain




embodiments, R₈ is

; wherein all R' are oxygen protecting groups. In



certain embodiments, R₈ is


; wherein all R' are oxygen protecting groups.



In certain embodiments, R₈ is

; wherein all R' are silyl, TBDPS, TBDMS,


TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain



embodiments, R₈ is


; wherein all R' are silyl, TBDPS, TBDMS, TIPS,

TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R₈










is

. In certain embodiments, R₈ is




. In certain






[00120] In compounds of Formula (A), R_9 is hydrogen, alkyl,




, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R_9 is hydrogen. In certain embodiments, R_9 is alkyl. In certain embodiments, R_9 is C_{1-6} alkyl. In certain embodiments, R_9 is methyl. In certain embodiments, R_9 is ethyl. In certain embodiments, R_9 is propyl. In certain embodiments, R_9 is butyl. In certain embodiments, R_9




is

. In certain embodiments, R_9 is




. In certain




embodiments, R_9 is

. In certain embodiments, R_9 is




. In










certain embodiments, R_9 is

. In certain embodiments, R_9 is




; wherein all R' are oxygen protecting groups. In certain embodiments, R_9






; wherein all R' are oxygen protecting groups. In certain embodiments, R₉




is ; wherein all R' are oxygen protecting groups. In certain embodiments,



R₉ is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS,

MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R₉ is



; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM,

THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R₉ is an oxygen

protecting group when attached to an oxygen atom. In certain embodiments, when attached to

an oxygen atom, R₉ is silyl. In certain embodiments, when attached to an oxygen atom, R₉ is

TBDPS, TBDMS, TIPS, TES, or TMS. In certain embodiments, when attached to an oxygen

atom, R₉ is MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R₉ is

a sulfur protecting group when attached to a sulfur atom. In certain embodiments, when

attached to a sulfur atom, R₉ is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-

pyridine-sulfenyl, or triphenylmethyl.

[00121] In compounds of Formula (A), R₁₀ is hydrogen, alkyl, acyl, or a nitrogen

protecting group. In certain embodiments, R₁₀ is hydrogen. In certain embodiments, R₁₀ is

alkyl. In certain embodiments, R₁₀ is C₁₋₆ alkyl. In certain embodiments, R₁₀ is methyl. In


certain embodiments, R₁₀ is ethyl. In certain embodiments, R₁₀ is propyl. In certain

embodiments, R₁₀ is butyl. In certain embodiments, R₁₀ is acyl. In certain embodiments, R₁₀

is acetyl. In certain embodiments, R_{10} is pivaloyl. In certain embodiments, R_{10} is a nitrogen protecting group. In certain embodiments, R_{10} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, or Ts.






[00122] In compounds of Formula (A), at least one R' is hydrogen or an oxygen protecting group. In certain embodiments, at least one R' is hydrogen. In certain embodiments, at least one R' is an oxygen protecting group. In certain embodiments, at least one R' is silyl. In certain embodiments, at least one R' is TBDPS. In certain embodiments, at least one R' is TBDMS. In certain embodiments, at least one R' is TIPS. In certain embodiments, at least one R' is TES. In certain embodiments, at least one R' is TMS. In certain embodiments, at least one R' is MOM. In certain embodiments, at least one R' is THP. In certain embodiments, at least one R' is *t*-Bu. In certain embodiments, at least one R' is Bn. In certain embodiments, at least one R' is allyl. In certain embodiments, at least one R' is acetyl. In certain embodiments, at least one R' is pivaloyl. In certain embodiments, at least one R' is Bz. In certain embodiments, all R' are hydrogen. In certain embodiments, all R' are oxygen protecting groups. In certain embodiments, all R' are silyl. In certain embodiments, all R' are TBDPS. In certain embodiments, all R' are TBDMS. In certain embodiments, all R' are TIPS. In certain embodiments, all R' are TES. In certain embodiments, all R' are TMS. In certain embodiments, all R' are MOM. In certain embodiments, all R' are THP. In certain embodiments, all R' are *t*-Bu. In certain embodiments, all R' are Bn. In certain embodiments, all R' are allyl. In certain embodiments, all R' are acetyl. In certain embodiments, all R' are pivaloyl. In certain embodiments, all R' are Bz.


[00123] In compounds of Formula (A), R'' is hydrogen, alkyl, acyl, an oxygen




is hydrogen. In certain embodiments, R'' is alkyl. In certain embodiments, R'' is C_{1-6} alkyl. In certain embodiments, R'' is methyl. In certain embodiments, R'' is substituted methyl. In certain embodiments, R'' is $-CH_2-CO_2H$. In certain embodiments, R'' is $-CH_2-CO_2$ (substituted or unsubstituted C_{1-6} alkyl) (e.g., $-CH_2-CO_2Me$). In certain embodiments, R'' is ethyl. In certain embodiments, R'' is propyl. In certain embodiments, R'' is butyl. In certain

embodiments, R'' is acyl. In certain embodiments, R'' is acetyl. In certain embodiments, R'' is pivaloyl. In certain embodiments, R'' is an oxygen protecting group. In certain embodiments, R'' is silyl. In certain embodiments, R'' is TBDPS, TBDMS, TIPS, TES, or TMS. In certain embodiments, R'' is MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain






In certain embodiments, R'' is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain




embodiments, R'' is ; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R''




is

. In certain embodiments, R'' is




. In certain




embodiments, R'' is

. In certain embodiments, R'' is




. In




certain embodiments, R'' is


. In certain embodiments, R'' is




; wherein all R' are oxygen protecting groups. In certain embodiments, R''



; wherein all R' are oxygen protecting groups. In certain embodiments,



; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz. In certain embodiments, R'' is



; wherein all R' are silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or Bz.

[00124] In compounds of Formula (A), each occurrence of *n* is independently an integer from 1 to 8, inclusive. In certain embodiments, at least one occurrence of *n* is 1. In certain embodiments, at least one occurrence of *n* is 2. In certain embodiments, at least one occurrence of *n* is 3. In certain embodiments, at least one occurrence of *n* is 4. In certain embodiments, at least one occurrence of *n* is 5. In certain embodiments, at least one occurrence of *n* is 6. In certain embodiments, at least one occurrence of *n* is 7. In certain embodiments, at least one occurrence of *n* is 8.


[00125] In compounds of Formula (A), *k* is an integer from 0 to 2, inclusive. In certain embodiments, *k* is 0. In certain embodiments, *k* is 1. In certain embodiments, *k* is 2.

[00126] In compounds of Formula (A), *x* is an integer from 1 to 8, inclusive. In certain embodiments, *x* is 1. In certain embodiments, *x* is 2. In certain embodiments, *x* is 3. In certain embodiments, *x* is 4. In certain embodiments, *x* is 5. In certain embodiments, *x* is 6. In certain embodiments, *x* is 7. In certain embodiments, *x* is 8.

[00127] In compounds of Formula (A), *y* is an integer from 0 to 8, inclusive. In certain embodiments, *y* is 0. In certain embodiments, *y* is 1. In certain embodiments, *y* is 2. In certain embodiments, *y* is 3. In certain embodiments, *y* is 4. In certain embodiments, *y* is 5. In certain embodiments, *y* is 6. In certain embodiments, *y* is 7. In certain embodiments, *y* is 8.


[00128] In compounds of Formula (A), all combinations of n, x, and y are contemplated. In certain embodiments, n is 2; x is 0; and y is 1. In certain embodiments, n is 2; x is 0; and y is 2. In certain embodiments, n is 2; x is 0; and y is 3. In certain embodiments, n is 2; x is 0; and y is 4.

[00129] The compound of Formula (A) may have one or more chiral centers. In certain embodiments, the compound of Formula (A) is of Formula (B):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00130] In other embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00131] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00132] The group $-OR_2$ of Formula (A) may be at any position, as valency permits, of the pyridyl ring of Formula (A). In certain embodiments, $-OR_2$ of Formula (A) is at the 3'-position of the pyridyl ring of Formula (A), and the compound of Formula (A) is of Formula (C):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00133] In certain embodiments, $-OR_2$ of Formula (A) is at the 4'-position of the pyridyl ring of Formula (A), and the compound of Formula (A) is of Formula (D):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00134] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00135] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00136] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00137] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00138] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00139] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00140] In certain embodiments, the compound of Formula (A) is of Formula (E):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00141] In certain embodiments, the compound of Formula (A) is of Formula (F):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00142] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00143] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00144] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00145] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00146] In certain embodiments, the compound of Formula (A) is of Formula (G):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00147] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00148] In certain embodiments, the compound of Formula (A) is of Formula (H):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00149] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00150] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00151] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00152] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, wherein R₃₀ is H, substituted or unsubstituted C₁₋₆ alkyl (*e.g.*, -CH₃), or an oxygen protecting group; and a is 2, 3, 4, 5, 6, 7, 8, 9, or 10 (*e.g.*, 4, 5, 6, 7, or 8).

[00153] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00154] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00155] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, wherein b is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 (*e.g.*, 1, 2, 3, 4, or 5).

[00156] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.



[00157] In certain embodiments, the compound of Formula (A) is of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.



[00158] In certain embodiments, the compound of Formula (A) is of the formula:







or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00159] In certain embodiments, the compound of Formula (A) is of the formula:






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00160] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.



[00161] In certain embodiments, the compound of Formula (A) is of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.



[00162] In certain embodiments, the compound of Formula (A) is of the formula:






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00163] In certain embodiments, the compound of Formula (A) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00164] In certain embodiments, the compound of Formula (A) is of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00165] In certain embodiments, the compound of Formula (A) is not of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00166] In certain embodiments, when R_7 is $-\text{OH}$, R_2 is not H or $-(\text{CH}_2\text{CH}_2\text{O})_2\text{CH}_3$. In certain embodiments, when R_7 is $-\text{OH}$, R_2 is not H or $-\text{[(CH}_2\text{)}_n\text{-O]}_x\text{[(CH}_2\text{)}_n\text{-O]}_y\text{-R''}$. In certain embodiments, when R_7 is $-\text{OH}$, $-\text{OR}_2$ is attached to the 5'-position of the pyridinyl ring.

[00167] In another aspect, novel DFT analogs are obtained by, among other things, changing the thiazoliny ring of DFT 1 to other 5-membered rings. In certain embodiments, compounds useful in the present invention are of Formula (J):



wherein:

--- represents a single bond or double bond;

W_1 - W_3 are independently CR_{22} , NR_{23} , oxygen, or sulfur, provided that:

when W_3 is nitrogen, R_{23} is null, and

when R_{21} is methyl or hydrogen, W_1 is not sulfur;

Z is $-\text{OR}_{11}$, $-\text{NR}_{12}\text{R}_{13}$, morpholine, or optionally substituted piperazinyl;

R_{11} is $-[(CH_2)_p-O]_u-[(CH_2)_q-O]_v-R_{14}$, $-[(CH_2)_p-NH]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$, or $-[(CH_2)_p-O]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$;

R_{12} is hydrogen, alkyl, $-[(CH_2)_p-O]_u-[(CH_2)_q-O]_v-R_{14}$, $-[(CH_2)_p-NH]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$, or $-[(CH_2)_p-O]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$;

R_{13} is hydrogen or alkyl;

p and q are independently an integer from 1 to 8, inclusive;

u is an integer from 0 to 8, inclusive;

v is an integer from 1 to 8, inclusive;

R_{14} and R_{15} are independently hydrogen, alkyl, or acyl;

R_{16} is hydrogen, alkyl, or alkoxy;

R_{17} is $-OR_{18}$ or $-N(OH)R_{19}$;

R_{18} is hydrogen, alkyl, or arylalkyl;

R_{19} is alkyl or $-(CH_2)_s-N(OH)C(=O)R_{20}$;

s is an integer from 1 to 8;

R_{20} is alkyl;

R_{21} is hydrogen or alkyl;


each occurrence of R_{22} is independently null, hydrogen, or alkyl; and

each occurrence of R_{23} is independently null, hydrogen, or alkyl;


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00168] In compounds of Formula (J), Z is $-OR_{11}$, $-NR_{12}R_{13}$, morpholine, or optionally substituted piperazinyl. In certain embodiments, Z is $-OR_{11}$. In certain embodiments, Z is $-O-[(CH_2)_p-O]_u-[(CH_2)_q-O]_v-R_{14}$. In certain embodiments, Z is



certain embodiments, Z is  $\cdot$ In certain embodiments, Z is




certain embodiments, Z is  $\cdot$ In certain embodiments, Z is




NH]_u–[(CH₂)_q–NR₁₄]_v–R₁₅. In certain embodiments, Z is –O–[(CH₂)_p–O]_u–[(CH₂)_q–NR₁₄]_v–R₁₅. In certain embodiments, Z is –NR₁₂R₁₃. In certain embodiments, Z is morpholine. In certain embodiments, Z is unsubstituted piperazinyl. In certain embodiments, Z is substituted piperazinyl. In certain embodiments, Z is piperazinyl substituted at the 4-position with an optionally substituted alkyl. In certain embodiments, Z is 4-(2-hydroxyethyl)-piperazin-1-yl.


[00169] In compounds of Formula (J), R₁₁ is –[(CH₂)_p–O]_u–[(CH₂)_q–O]_v–R₁₄, –[(CH₂)_p–NH]_u–[(CH₂)_q–NR₁₄]_v–R₁₅, or –[(CH₂)_p–O]_u–[(CH₂)_q–NR₁₄]_v–R₁₅. In certain embodiments, R₁₁ is –[(CH₂)_p–O]_u–[(CH₂)_q–O]_v–R₁₄. In certain embodiments, R₁₁ is



[00170] In compounds of Formula (J), R₁₂ is hydrogen, alkyl, –[(CH₂)_p–O]_u–[(CH₂)_q–O]_v–R₁₄, –[(CH₂)_p–NH]_u–[(CH₂)_q–NR₁₄]_v–R₁₅, or –[(CH₂)_p–O]_u–[(CH₂)_q–NR₁₄]_v–R₁₅. In certain embodiments, R₁₂ is hydrogen. In certain embodiments, R₁₂ is alkyl. In certain embodiments, R₁₂ is C₁₋₆ alkyl. In certain embodiments, R₁₂ is methyl. In certain embodiments, R₁₂ is ethyl. In certain embodiments, R₁₂ is propyl. In certain embodiments, R₁₂ is butyl. In certain embodiments, R₁₂ is –[(CH₂)_p–O]_u–[(CH₂)_q–O]_v–R₁₄. In certain





certain embodiments, R_{12} is  . In certain
 embodiments, R_{12} is $-[(CH_2)_p-NH]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$. In certain embodiments, R_{12} is $-[(CH_2)_p-O]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$.

[00171] In compounds of Formula (J), R_{13} is hydrogen or alkyl. In certain
 embodiments, R_{13} is hydrogen. In certain embodiments, R_{13} is alkyl. In certain embodiments,
 R_{13} is C_{1-6} alkyl. In certain embodiments, R_{13} is methyl. In certain embodiments, R_{13} is ethyl.
 In certain embodiments, R_{13} is propyl. In certain embodiments, R_{13} is butyl.

[00172] In compounds of Formula (J), p is an integer from 1 to 8, inclusive. In certain
 embodiments, p is 1. In certain embodiments, p is 2. In certain embodiments, p is 3. In certain
 embodiments, p is 4. In certain embodiments, p is 5. In certain embodiments, p is 6. In certain
 embodiments, p is 7. In certain embodiments, p is 8.

[00173] In compounds of Formula (J), q is an integer from 1 to 8, inclusive. In certain
 embodiments, q is 1. In certain embodiments, q is 2. In certain embodiments, q is 3. In certain
 embodiments, q is 4. In certain embodiments, q is 5. In certain embodiments, q is 6. In certain
 embodiments, q is 7. In certain embodiments, q is 8.

[00174] In compounds of Formula (J), u is an integer from 0 to 8, inclusive. In certain
 embodiments, u is 0. In certain embodiments, u is 1. In certain embodiments, u is 2. In certain
 embodiments, u is 3. In certain embodiments, u is 4. In certain embodiments, u is 5. In certain
 embodiments, u is 6. In certain embodiments, u is 7. In certain embodiments, u is 8.

[00175] In compounds of Formula (J), v is an integer from 1 to 8, inclusive. In certain
 embodiments, v is 1. In certain embodiments, v is 2. In certain embodiments, v is 3. In certain
 embodiments, v is 4. In certain embodiments, v is 5. In certain embodiments, v is 6. In certain
 embodiments, v is 7. In certain embodiments, v is 8.

[00176] In compounds of Formula (J), R_{14} is hydrogen, alkyl, or acyl. In certain
 embodiments, R_{14} is hydrogen. In certain embodiments, R_{14} is alkyl. In certain embodiments,
 R_{14} is C_{1-6} alkyl. In certain embodiments, R_{14} is methyl. In certain embodiments, R_{14} is ethyl.
 In certain embodiments, R_{14} is propyl. In certain embodiments, R_{14} is butyl. In certain
 embodiments, R_{14} is acyl. In certain embodiments, R_{14} is acetyl.

[00177] In compounds of Formula (J), R_{15} is hydrogen, alkyl, or acyl. In certain
 embodiments, R_{15} is hydrogen. In certain embodiments, R_{15} is alkyl. In certain embodiments,

R₁₅ is C₁₋₆ alkyl. In certain embodiments, R₁₅ is methyl. In certain embodiments, R₁₅ is ethyl. In certain embodiments, R₁₅ is propyl. In certain embodiments, R₁₅ is butyl. In certain embodiments, R₁₅ is acyl. In certain embodiments, R₁₅ is acetyl.

[00178] In compounds of Formula (J), R₁₆ is hydrogen, alkyl, or alkoxy. In certain embodiments, R₁₆ is hydrogen. In certain embodiments, R₁₆ is alkyl. In certain embodiments, R₁₆ is C₁₋₆ alkyl. In certain embodiments, R₁₆ is methyl. In certain embodiments, R₁₆ is ethyl. In certain embodiments, R₁₆ is propyl. In certain embodiments, R₁₆ is butyl. In certain embodiments, R₁₆ is alkoxy. In certain embodiments, R₁₆ is C₁₋₆ alkoxy. In certain embodiments, R₁₆ is methoxy. In certain embodiments, R₁₆ is ethoxy. In certain embodiments, R₁₆ is propoxy. In certain embodiments, R₁₆ is butoxy.

[00179] In compounds of Formula (J), R₁₇ is –OR₁₈ or –N(OH)R₁₉. In certain embodiments, R₁₇ is –OR₁₈. In certain embodiments, R₁₇ is –OH. In certain embodiments, R₁₇ is –O–alkyl. In certain embodiments, R₁₇ is –O–(C₁₋₆ alkyl). In certain embodiments, R₁₇ is –OMe. In certain embodiments, R₁₇ is –OEt. In certain embodiments, R₁₇ is –OPr. In certain embodiments, R₁₇ is –OBu. In certain embodiments, R₁₇ is –O–arylalkyl. In certain embodiments, R₁₇ is –O–phenalkyl. In certain embodiments, R₁₇ is –O–Bn. In certain embodiments, R₁₇ is –O–phenethyl. In certain embodiments, R₁₇ is –N(OH)R₁₉. In certain embodiments, R₁₇ is –N(OH)–alkyl. In certain embodiments, R₁₇ is –N(OH)–(C₁₋₆ alkyl). In certain embodiments, R₁₇ is –N(OH)–(CH₂)_s–N(OH)C(=O)R₂₀.

[00180] In compounds of Formula (J), R₁₈ is hydrogen, alkyl, or arylalkyl. In certain embodiments, R₁₈ is hydrogen. In certain embodiments, R₁₈ is alkyl. In certain embodiments, R₁₈ is C₁₋₆ alkyl. In certain embodiments, R₁₈ is methyl. In certain embodiments, R₁₈ is ethyl. In certain embodiments, R₁₈ is propyl. In certain embodiments, R₁₈ is butyl. In certain embodiments, R₁₈ is arylalkyl. In certain embodiments, R₁₈ is aryl–(C₁₋₆ alkyl). In certain embodiments, R₁₈ is phenyl–(C₁₋₆ alkyl). In certain embodiments, R₁₈ is Bn. In certain embodiments, R₁₈ is phenethyl.

[00181] In compounds of Formula (J), R₁₉ is alkyl or –(CH₂)_s–N(OH)C(=O)R₂₀.

[00182] In certain embodiments, R₁₉ is alkyl. In certain embodiments, R₁₉ is C₁₋₆ alkyl. In certain embodiments, R₁₉ is methyl. In certain embodiments, R₁₉ is ethyl. In certain embodiments, R₁₉ is propyl. In certain embodiments, R₁₉ is butyl. In certain embodiments, R₁₉ is –(CH₂)_s–N(OH)C(=O)R₂₀.

[00183] In compounds of Formula (J), s is an integer from 1 to 8, inclusive. In certain embodiments, s is 1. In certain embodiments, s is 2. In certain embodiments, s is 3. In certain

embodiments, s is 4. In certain embodiments, s is 5. In certain embodiments, s is 6. In certain embodiments, s is 7. In certain embodiments, s is 8.


[00184] In compounds of Formula (J), R_{20} is alkyl. In certain embodiments, R_{20} is C_{1-6} alkyl. In certain embodiments, R_{20} is methyl. In certain embodiments, R_{20} is ethyl. In certain embodiments, R_{20} is propyl. In certain embodiments, R_{20} is butyl.


[00185] In compounds of Formula (J), R_{21} is hydrogen or alkyl. In certain embodiments, R_{21} is hydrogen. In certain embodiments, R_{21} is alkyl. In certain embodiments, R_{21} is C_{1-6} alkyl. In certain embodiments, R_{21} is methyl. In certain embodiments, R_{21} is ethyl. In certain embodiments, R_{21} is propyl. In certain embodiments, R_{21} is butyl.

[00186] In compounds of Formula (J), each occurrence of R_{22} is independently null, hydrogen, or alkyl. In certain embodiments, at least one R_{22} is null. In certain embodiments, at least one R_{22} is hydrogen. In certain embodiments, at least one R_{22} is alkyl. In certain embodiments, at least one R_{22} is C_{1-6} alkyl. In certain embodiments, at least one R_{22} is methyl. In certain embodiments, at least one R_{22} is ethyl. In certain embodiments, at least one R_{22} is propyl. In certain embodiments, at least one R_{22} is butyl.

[00187] In compounds of Formula (J), each occurrence of R_{23} is independently null, hydrogen, or alkyl. In certain embodiments, at least one R_{23} is null. In certain embodiments, at least one R_{23} is hydrogen. In certain embodiments, at least one R_{23} is alkyl. In certain embodiments, at least one R_{23} is C_{1-6} alkyl. In certain embodiments, at least one R_{23} is methyl. In certain embodiments, at least one R_{23} is ethyl. In certain embodiments, at least one R_{23} is propyl. In certain embodiments, at least one R_{23} is butyl.


[00188] In certain embodiments, the compound of Formula (J) is of the formula:






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00189] The compound of Formula (J) may have one or more chiral centers. In certain embodiments, the compound of Formula (J) is of Formula (K):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00190] In certain embodiments, the compound of Formula (J) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00191] In certain embodiments, the compound of Formula (J) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00192] In certain embodiments, the compound of Formula (J) is of the formula:




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00193] In certain embodiments, the compound of Formula (J) is of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00194] In certain embodiments, the compound of Formula (J) is of the formula:



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00195] The compounds of the invention may be provided in various salts forms. In certain embodiments, the inventive compounds are provided as alkali metal salts. In certain embodiments, the inventive compounds are provided as alkaline earth metal salts. In certain embodiments, when R₉ is –OH, the compound may be provided as a carboxylate salt with a positively charged counterion. In certain embodiments, the counterion is betaine, choline hydroxide, diethanolamine, diethylamine, ethanolamine, hydroxyethylmorpholine, 4-(2-hydroxyethyl morpholine), 1-(2-hydroxyethyl pyrrolidine), 1-(2-hydroxyethyl)-piperidine, 1,2-EDSA, HCl, H₂SO₄, MSA, *p*-TSA, hydroxyethyl pyrrolidine, imidazone, lysine (*e.g.*, L-lysine), arginine (*e.g.*, L-arginine), histidine (*e.g.*, L-histidine) *N*-methyl-D-glucamine

(NMG), *N, N'*-dibenzyl-ethylenediamine, *N, N'*-diethyl-ethanolamine, triethanolamine, tromethamine, calcium (*e.g.*, $\text{Ca}(\text{OH})_2$), magnesium (*e.g.*, $\text{Mg}(\text{OH})_2$, magnesium acetate), potassium (*e.g.*, KOH, potassium 2-ethylhexanoate), sodium (*e.g.*, NaOH, sodium acetate, sodium 2-ethylhexanoate), zinc (*e.g.*, $\text{Zn}(\text{OH})_2$, zinc acetate), $\text{Zn}(\text{OH})_2/\text{Mg}(\text{OH})_2$, EDA, or piperazinyl. In certain embodiments, the counterion is lysine. In certain embodiments, the counterion is *N*-methyl-D-glucamine (NMG). In certain embodiments, the counterion is tromethamine. In certain embodiments, the counterion is calcium. In certain embodiments, the counterion is magnesium. In certain embodiments, the counterion is cesium. In certain embodiments, the counterion is potassium. In certain embodiments, the counterion is sodium. In certain embodiments, the counterion is lithium. In certain embodiments, the counterion is zinc. In certain embodiments, the counterion is piperzine. In certain embodiments, the counterion is MgOH^+ . In certain embodiments, the counterion is ZnOH^+ .

[00196] In certain embodiments, a polymorph of a salt of a compound of the invention is provided. In certain embodiments, a polymorph of a magnesium salt of a compound of the invention is provided. In certain embodiments, a polymorph of a Na^+ salt of a compound of the invention is provided. In certain embodiments, a polymorph of a salt of a carboxylate compound of the invention, wherein R_9 is $-\text{OH}$, is provided. In certain embodiments, a polymorph of a magnesium salt of a carboxylate compound of the invention, wherein R_9 is $-\text{OH}$, is provided. In certain embodiments, a polymorph of a Na^+ salt of a carboxylate compound of the invention, wherein R_9 is $-\text{OH}$, is provided.


[00197] The cation and anion in a salt disclosed herein may combine in a 1:1 molar ratio. Other molar ratios (*e.g.*, 1:1.5, 1:2, 1:6, and 2:1) are also possible. Salts shown herein may be, for the sole purpose of convenience in notation, shown in a 1:1 ratio. All possible stoichiometric arrangements are encompassed by the scope of the present invention.

[00198] In certain embodiments, the compound of Formula (A) is of the Formula (I-1):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00199] In certain embodiments, a salt of a compound of the Formula (I-1) is provided. In certain embodiments, a salt of the Formula (I-1) is provided as shown in the Formula (I-1-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00200] In certain embodiments, the compound of the Formula (A) is of the Formula (I-1-ii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00201] In certain embodiments, the compound of the Formula (A) is of the Formula (I-1-iii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00202] In certain embodiments, the compound of the Formula (A) is of the Formula (I-1-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00203] In certain embodiments, the compound of the Formula (A) is of the Formula (I-2):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00204] In certain embodiments, a salt of a compound of the Formula (I-2) is provided. In certain embodiments, a salt of the Formula (I-2) is provided as shown in the Formula (I-2-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00205] In certain embodiments, the compound of the Formula (A) is of the Formula (I-2-ii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00206] In certain embodiments, the compound of the Formula (A) is of the Formula (I-2-iii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00207] In certain embodiments, the compound of the Formula (A) is of the Formula (I-2-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00208] In certain embodiments, the compound of the Formula (A) is of the Formula (I-3):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00209] In certain embodiments, a salt of a compound of the Formula (I-3) is provided. In certain embodiments, a salt of the Formula (I-3) is provided as shown in the Formula (I-3-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00210] In certain embodiments, the compound of the Formula (A) is of the Formula (I-3-ii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00211] In certain embodiments, the compound of the Formula (A) is of the Formula (I-3-iii):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.




[00212] In certain embodiments, the compound of the Formula (A) is of the Formula (I-3-iv):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00213] In certain embodiments, the compound of Formula (A) is of the formula:








or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00214] In certain embodiments, the compound of Formula (A) is of the formula:








or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00215] In certain embodiments, the salt of a compound of Formula (A) is of the formula:








or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00216] In certain embodiments, the compound of Formula (A) is of the formula:








or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00217] In certain embodiments, the compound of Formula (A) is of the formula:






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00218] In certain embodiments, the salt of a compound of Formula (A) is of the formula:







or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00219] Additional exemplary compounds of Formula (A) include, but are not limited to:









and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof.

[00220] In certain embodiments, the compound of Formula (J) is of the Formula (J-1):






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00221] In certain embodiments, a salt of a compound of the Formula (J-1) is provided. In certain embodiments, a salt of the Formula (J-1) is provided as shown in the Formula (J-1-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00222] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-1-ii) to (J-1-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00223] In certain embodiments, the compound of Formula (J) is of the Formula (J-2):





or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00224] In certain embodiments, a salt of a compound of the Formula (J-2) is provided. In certain embodiments, a salt of the Formula (J-2) is provided as shown in the Formula (J-2-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00225] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-2-ii) to (J-2-iv):






(J-2-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof


[00226] In certain embodiments, the compound of Formula (J) is of the Formula (J-3):



(J-3),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.




[00227] In certain embodiments, a salt of a compound of the Formula (J-3) is provided. In certain embodiments, a salt of the Formula (J-3) is provided as shown in the Formula (J-3-i):



(J-3-i),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00228] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-3-ii) to (J-3-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00229] In certain embodiments, the compound of Formula (J) is of the Formula (J-4):






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00230] In certain embodiments, a salt of a compound of the Formula (J-4) is provided. In certain embodiments, a salt of the Formula (J-4) is provided as shown in the Formula (J-4-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00231] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-4-ii) to (J-4-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00232] In certain embodiments, the compound of Formula (J) is of the Formula (J-5):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00233] In certain embodiments, a salt of a compound of the Formula (J-5) is provided. In certain embodiments, a salt of the Formula (J-5) is provided as shown in the Formula (J-5-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00234] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-5-ii) to (J-5-iv):






(J-5-iv)

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00235] In certain embodiments, the compound of Formula (J) is of the Formula (J-6):



(J-6),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00236] In certain embodiments, a salt of a compound of the Formula (J-6) is provided. In certain embodiments, a salt of the Formula (J-6) is provided as shown in the Formula (J-6-i):



(J-6-i),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00237] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-6-ii) to (J-6-iv):



(J-6-ii)


(J-6-iii)



(J-6-iv)

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00238] In certain embodiments, the compound of Formula (J) is of the Formula (J-7):



(J-7),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00239] In certain embodiments, a salt of a compound of the Formula (J-7) is provided. In certain embodiments, a salt of the Formula (J-7) is provided as shown in the Formula (J-7-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00240] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-7-ii) to (J-7-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00241] In certain embodiments, the compound of Formula (J) is of the Formula (J-8):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00242] In certain embodiments, a salt of a compound of the Formula (J-8) is provided. In certain embodiments, a salt of the Formula (J-8) is provided as shown in the Formula (J-8-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00243] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-8-ii) to (J-8-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00244] In certain embodiments, the compound of Formula (J) is of the Formula (J-9):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00245] In certain embodiments, a salt of a compound of the Formula (J-9) is provided. In certain embodiments, a salt of the Formula (J-9) is provided as shown in the Formula (J-9-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00246] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-9-ii) to (J-9-iv):






(J-9-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00247] In certain embodiments, the compound of Formula (J) is of the Formula (J-10):



(J-10),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00248] In certain embodiments, a salt of a compound of the Formula (J-10) is provided. In certain embodiments, a salt of the Formula (J-10) is provided as shown in the Formula (J-10-i):



(J-10-i),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00249] In certain embodiments, the compound of the Formula (A) is of any one of the Formulae (J-10-ii) to (J-10-iv):



(J-10-ii)


(J-10-iii),



(J-10-iv),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00250] In certain embodiments, the compound of Formula (J) is of the Formula (J-11):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00251] In certain embodiments, a salt of a compound of the Formula (J-11) is provided. In certain embodiments, a salt of the Formula (J-11) is provided as shown in the Formula (J-11-i):




(J-11-i),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00252] In certain embodiments, the compound of the Formula (A) is of any one of the Formulae (J-11-ii) to (J-11-iv):



(J-11-ii)




(J-11-iii),



(J-11-iv)


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof

[00253] In certain embodiments, the compound of Formula (J) is of the Formula (J-12):






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00254] In certain embodiments, a salt of a compound of the Formula (J-12) is provided. In certain embodiments, a salt of the Formula (J-12) is provided as shown in the Formula (J-12-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00255] In certain embodiments, the compound of the Formula (A) is of any one of the Formulae (J-12-ii) to (J-12-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00256] In certain embodiments, the compound of Formula (J) is of the Formula (J-13):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00257] In certain embodiments, a salt of a compound of the Formula (J-13) is provided. In certain embodiments, a salt of the Formula (J-13) is provided as shown in the Formula (J-13-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00258] In certain embodiments, the compound of the Formula (A) is of any one of the Formulae (J-13-ii) to (J-13-iv):



(J-13-ii)


(J-13-iii),



(J-13-iv),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00259] In certain embodiments, the compound of Formula (J) is of the Formula (J-14):





or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00260] In certain embodiments, a salt of a compound of the Formula (J-14) is provided. In certain embodiments, a salt of the Formula (J-14) is provided as shown in the Formula (J-14-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00261] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-14-ii) to (J-14-iv):






or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00262] In certain embodiments, the compound of Formula (J) is of the Formula (J-15):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00263] In certain embodiments, a salt of a compound of the Formula (J-15) is provided. In certain embodiments, a salt of the Formula (J-15) is provided as shown in the Formula (J-15-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00264] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-15-ii) to (J-15-iv):



(J-15-ii)


(J-15-iii),



(J-15-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00265] In certain embodiments, the compound of Formula (J) is of the Formula (J-16):



(J-16),




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00266] In certain embodiments, a salt of a compound of the Formula (J-16) is provided. In certain embodiments, a salt of the Formula (J-16) is provided as shown in the Formula (J-16-i):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00267] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-16-ii) to (J-16-iv):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00268] In certain embodiments, the compound of Formula (J) is of the Formula (J-17):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00269] In certain embodiments, a salt of a compound of the Formula (J-17) is provided. In certain embodiments, a salt of the Formula (J-17) is provided as shown in the Formula (J-17-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00270] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-17-ii) to (J-17-iv):






(J-17-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00271] In certain embodiments, the compound of Formula (J) is of the Formula (J-18):



(J-18),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00272] In certain embodiments, a salt of a compound of the Formula (J-18) is provided. In certain embodiments, a salt of the Formula (J-18) is provided as shown in the Formula (J-18-i):



(J-18-i),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00273] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-18-ii) to (J-18-iv):



(J-18-ii)


(J-18-iii),



(J-18-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00274] In certain embodiments, the compound of Formula (J) is of the Formula (J-19):



(J-19),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00275] In certain embodiments, a salt of a compound of the Formula (J-19) is provided. In certain embodiments, a salt of the Formula (J-19) is provided as shown in the Formula (J-19-i):



(J-19-i),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00276] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-19-ii) to (J-19-iv):



(J-19-ii)


(J-19-iii),



(J-19-iv),


or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00277] In certain embodiments, the compound of Formula (J) is of the Formula (J-20):




or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00278] In certain embodiments, a salt of a compound of the Formula (J-20) is provided. In certain embodiments, a salt of the Formula (J-20) is provided as shown in the Formula (J-20-i):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00279] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-20-ii) to (J-20-iv):






(J-20-iv),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.


[00280] In certain embodiments, the compound of Formula (J) is of the Formula (J-21):



(J-21),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.




[00281] In certain embodiments, a salt of a compound of the Formula (J-21) is provided. In certain embodiments, a salt of the Formula (J-21) is provided as shown in the Formula (J-21-i):



(J-21-i),

or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

[00282] In certain embodiments, the compound of the Formula (J) is of any one of the Formulae (J-21-ii) to (J-21-iv):



or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof.

Pharmaceutical Compositions, Kits, and Administration

[00283] The present invention provides pharmaceutical compositions comprising a compound of the invention, and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof, and optionally a pharmaceutically acceptable excipient. In certain embodiments, the compound of the present invention, or a pharmaceutically acceptable salt thereof, is provided in an effective amount in the pharmaceutical composition. In certain embodiments, the effective amount is a therapeutically effective amount. In certain embodiments, the effective amount is a prophylactically effective amount.

[00284] Pharmaceutical compositions described herein can be prepared by any method known in the art of pharmacology. In general, such preparatory methods include the steps of bringing the compound of the present invention (the “active ingredient”) into association with a carrier or excipient, and/or one or more other accessory ingredients, and then, if necessary

and/or desirable, shaping, and/or packaging the product into a desired single- or multi-dose unit.

[00285] Pharmaceutical compositions can be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. A “unit dose” is a discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[00286] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition of the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition may comprise between 0.1% and 100% (w/w) active ingredient.

[00287] Pharmaceutically acceptable excipients used in the manufacture of provided pharmaceutical compositions include inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and perfuming agents may also be present in the composition.

[00288] Exemplary diluents include calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, and mixtures thereof.

[00289] Exemplary granulating and/or dispersing agents include potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose, and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, and mixtures thereof.

[00290] Exemplary surface active agents and/or emulsifiers include natural emulsifiers (e.g., acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan,

pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (*e.g.*, bentonite (aluminum silicate) and Veegum (magnesium aluminum silicate)), long chain amino acid derivatives, high molecular weight alcohols (*e.g.*, stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (*e.g.*, carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (*e.g.*, carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (*e.g.*, polyoxyethylene sorbitan monolaurate (Tween 20), polyoxyethylene sorbitan (Tween 60), polyoxyethylene sorbitan monooleate (Tween 80), sorbitan monopalmitate (Span 40), sorbitan monostearate (Span 60), sorbitan tristearate (Span 65), glyceryl monooleate, sorbitan monooleate (Span 80)), polyoxyethylene esters (*e.g.*, polyoxyethylene monostearate (Myrj 45), polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and Solutol), sucrose fatty acid esters, polyethylene glycol fatty acid esters (*e.g.*, Cremophor™), polyoxyethylene ethers, (*e.g.*, polyoxyethylene lauryl ether (Brij 30)), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic F-68, Poloxamer188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, and/or mixtures thereof.

[00291] Exemplary binding agents include starch (*e.g.*, cornstarch and starch paste), gelatin, sugars (*e.g.*, sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol, *etc.*), natural and synthetic gums (*e.g.*, acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum), and larch arabogalactan), alginates, polyethylene oxide, polyethylene glycol, inorganic calcium salts, silicic acid, polymethacrylates, waxes, water, alcohol, and/or mixtures thereof.

[00292] Exemplary preservatives include antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and other preservatives.

[00293] Exemplary antioxidants include alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium

metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and sodium sulfite.

[00294] Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA) and salts and hydrates thereof (*e.g.*, sodium edetate, disodium edetate, trisodium edetate, calcium disodium edetate, dipotassium edetate, and the like), citric acid and salts and hydrates thereof (*e.g.*, citric acid monohydrate), fumaric acid and salts and hydrates thereof, malic acid and salts and hydrates thereof, phosphoric acid and salts and hydrates thereof, and tartaric acid and salts and hydrates thereof. Exemplary antimicrobial preservatives include benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and thimerosal.

[00295] Exemplary antifungal preservatives include butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and sorbic acid.

[00296] Exemplary alcohol preservatives include ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and phenylethyl alcohol.

[00297] Exemplary acidic preservatives include vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and phytic acid.

[00298] Other preservatives include tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, Glydant Plus, Phenonip, methylparaben, Germall 115, Germaben II, Neolone, Kathon, and Euxyl. In certain embodiments, the preservative is an anti-oxidant. In other embodiments, the preservative is a chelating agent.

[00299] Exemplary buffering agents include citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate,

potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, and mixtures thereof.

[00300] Exemplary lubricating agents include magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behanate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, and mixtures thereof.

[00301] Exemplary natural oils include almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary synthetic oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and mixtures thereof.

[00302] Liquid dosage forms for oral and parenteral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredients, the liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (*e.g.*, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. In certain embodiments for parenteral administration, the conjugates of the invention are mixed with solubilizing agents such as Cremophor™, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and mixtures thereof.

[00303] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00304] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00305] In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

[00306] Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing the conjugates of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

[00307] While it may be possible for the compounds disclosed herein, or pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, or polymorphs thereof, to be administered orally as they are, it is also possible to present them as a pharmaceutical formulation or dosage. Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or (a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders such as, for example,

carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, (c) humectants such as glycerol, (d) disintegrating agents such as agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, (e) solution retarding agents such as paraffin, (f) absorption accelerators such as quaternary ammonium compounds, (g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, (h) absorbents such as kaolin and bentonite clay, and (i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may comprise buffering agents.

[00308] Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00309] The active ingredient can be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active ingredient can be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may comprise buffering agents. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

[00310] Dosage forms for topical and/or transdermal administration of a compound of this invention may include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants and/or patches. Generally, the active ingredient is admixed under sterile conditions with a pharmaceutically acceptable carrier or excipient and/or any needed preservatives and/or buffers as can be required. Additionally, the present invention contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of an active ingredient to the body. Such dosage forms can be prepared, for example, by dissolving and/or dispensing the active ingredient in the proper medium. Alternatively or additionally, the rate can be controlled by either providing a rate controlling membrane and/or by dispersing the active ingredient in a polymer matrix and/or gel.

[00311] Suitable devices for use in delivering intradermal pharmaceutical compositions described herein include short needle devices such as those described in U.S. Patents 4,886,499; 5,190,521; 5,328,483; 5,527,288; 4,270,537; 5,015,235; 5,141,496; and 5,417,662. Intradermal compositions can be administered by devices which limit the effective penetration length of a needle into the skin, such as those described in international PCT Application Publication No. WO 99/34850 and functional equivalents thereof. Jet injection devices which deliver liquid vaccines to the dermis *via* a liquid jet injector and/or *via* a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Jet injection devices are described, for example, in U.S. Patents 5,480,381; 5,599,302; 5,334,144; 5,993,412; 5,649,912; 5,569,189; 5,704,911; 5,383,851; 5,893,397; 5,466,220; 5,339,163; 5,312,335; 5,503,627; 5,064,413; 5,520,639; 4,596,556; 4,790,824; 4,941,880; 4,940,460; and international PCT Application Publication Nos. WO 97/37705 and WO 97/13537. Ballistic powder/particle delivery devices which use compressed gas to accelerate vaccine in powder form through the outer layers of the skin to the dermis are suitable. Alternatively or additionally, conventional syringes can be used in the classical mantoux method of intradermal administration.

[00312] Formulations suitable for topical administration include, but are not limited to, liquid and/or semi-liquid preparations such as liniments, lotions, oil in water and/or water in oil emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions. Topically-administrable formulations may, for example, comprise from about 1% to about 10% (w/w) active ingredient, although the concentration of the active ingredient can be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical

administration may further comprise one or more of the additional ingredients described herein.

[00313] Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50 to 99.9% (w/w) of the composition, and the active ingredient may constitute 0.1 to 20% (w/w) of the composition. The propellant may further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

[00314] Pharmaceutical compositions of the invention formulated for pulmonary delivery may provide the active ingredient in the form of droplets of a solution and/or suspension. Such formulations can be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising the active ingredient, and may conveniently be administered using any nebulization and/or atomization device. Such formulations may further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. The droplets provided by this route of administration may have an average diameter in the range from about 0.1 to about 200 nanometers.

[00315] Formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutical composition of the invention. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 to 500 micrometers. Such a formulation is administered by rapid inhalation through the nasal passage from a container of the powder held close to the nares.

[00316] Formulations for nasal administration may, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of the active ingredient, and may comprise one or more of the additional ingredients described herein. A pharmaceutical composition of the invention can be prepared, packaged, and/or sold in a formulation for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may contain, for example, 0.1 to 20% (w/w) active ingredient, the balance comprising an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising the active ingredient.

Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, may have an average particle and/or droplet size in the range from about 0.1 to about 200 nanometers, and may further comprise one or more of the additional ingredients described herein.

[00317] A pharmaceutical composition of the invention can be prepared, packaged, and/or sold in a formulation for ophthalmic administration. Such formulations may, for example, be in the form of eye drops including, for example, a 0.1-1.0% (w/w) solution and/or suspension of the active ingredient in an aqueous or oily liquid carrier or excipient. Such drops may further comprise buffering agents, salts, and/or one or more other of the additional ingredients described herein. Other ophthalmically-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this invention.

[00318] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with ordinary experimentation.

[00319] Compounds provided herein are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject or organism will depend upon a variety of factors including the disease being treated and the severity of the disorder; the activity of the specific active ingredient employed; the specific composition employed; the age, body weight, general health, sex, and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific active ingredient employed; the duration of the treatment; drugs used in combination or coincidental with the specific active ingredient employed; and like factors well known in the medical arts.

[00320] The compounds and compositions provided herein can be administered by any route, including enteral (*e.g.*, oral), parenteral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal,

intravaginal, intraperitoneal, topical (as by powders, ointments, creams, and/or drops), mucosal, nasal, bucal, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; and/or as an oral spray, nasal spray, and/or aerosol. Specifically contemplated routes are oral administration, intravenous administration (*e.g.*, systemic intravenous injection), regional administration *via* blood and/or lymph supply, and/or direct administration to an affected site. The inventive compounds and compositions may also be mixed with blood *ex vivo*, and the resulting mixture may be administered (*e.g.*, intravenously) to a subject. In general the most appropriate route of administration will depend upon a variety of factors including the nature of the agent (*e.g.*, its stability in the environment of the gastrointestinal tract), and/or the condition of the subject (*e.g.*, whether the subject is able to tolerate oral administration).

[00321] The exact amount of a compound required to achieve an effective amount will vary from subject to subject, depending, for example, on species, age, and general condition of a subject, severity of the side effects or disorder, identity of the particular compound, mode of administration, and the like. An effective amount may be included in a single dose (*e.g.*, single oral dose) or multiple doses (*e.g.*, multiple oral doses). In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, any two doses of the multiple doses include different or substantially the same amounts of a compound described herein. In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is three doses a day, two doses a day, one dose a day, one dose every other day, one dose every third day, one dose every week, one dose every two weeks, one dose every three weeks, or one dose every four weeks. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is one dose per day. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is two doses per day. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is three doses per day. In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, the duration between the first dose and last dose of the multiple doses is one day, two days, four days, one week, two weeks, three weeks, one month, two months, three months, four months, six months, nine months, one year, two years, three years, four years, five years, seven years, ten years, fifteen years, twenty years, or the lifetime of the subject, tissue, or cell. In certain embodiments, the duration between the first dose and last dose of the

multiple doses is three months, six months, or one year. In certain embodiments, the duration between the first dose and last dose of the multiple doses is the lifetime of the subject, tissue, or cell. In certain embodiments, a dose (*e.g.*, a single dose, or any dose of multiple doses) described herein includes independently between 0.1 μg and 1 μg , between 0.001 mg and 0.01 mg, between 0.01 mg and 0.1 mg, between 0.1 mg and 1 mg, between 1 mg and 3 mg, between 3 mg and 10 mg, between 10 mg and 30 mg, between 30 mg and 100 mg, between 100 mg and 300 mg, between 300 mg and 1,000 mg, or between 1 g and 10 g, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 1 mg and 3 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 3 mg and 10 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 10 mg and 30 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 30 mg and 100 mg, inclusive, of a compound described herein.

[00322] It will be appreciated that dose ranges as described herein provide guidance for the administration of provided pharmaceutical compositions to an adult. The amount to be administered to, for example, a child or an adolescent can be determined by a medical practitioner or person skilled in the art and can be lower or the same as that administered to an adult.

[00323] It will be also appreciated that a compound or composition, as described herein, can be administered in combination with one or more additional therapeutically active agents. The compounds or compositions can be administered in combination with additional therapeutically active agents that improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. It will also be appreciated that the therapy employed may achieve a desired effect for the same disorder, and/or it may achieve different effects.

[00324] The compound or composition can be administered concurrently with, prior to, or subsequent to, one or more additional therapeutically active agents. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. It will further be appreciated that the additional therapeutically active agent utilized in this combination can be administered together in a single composition or administered separately in different compositions. The particular combination to employ in a regimen will take into account compatibility of the inventive compound with the additional therapeutically active agent and/or the desired therapeutic effect to be achieved. In general, it is expected that

additional therapeutically active agents in combination be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

[00325] Exemplary additional therapeutically active agents include, but are not limited to, anti-cancer agents, anti-diabetic agents, anti-inflammatory agents, immunosuppressant agents, and a pain-relieving agent. Therapeutically active agents include small organic molecules such as drug compounds (*e.g.*, compounds approved by the U.S. Food and Drug Administration as provided in the Code of Federal Regulations (CFR)), peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoproteins, mucoproteins, lipoproteins, synthetic polypeptides or proteins, small molecules linked to proteins, glycoproteins, steroids, nucleic acids, DNAs, RNAs, nucleotides, nucleosides, oligonucleotides, antisense oligonucleotides, lipids, hormones, vitamins, and cells.

[00326] Also encompassed by the invention are kits (*e.g.*, pharmaceutical packs). The kits provided may comprise an inventive pharmaceutical composition or compound and a container (*e.g.*, a vial, ampule, bottle, syringe, and/or dispenser package, or other suitable container). In some embodiments, provided kits may optionally further include a second container comprising a pharmaceutical excipient for dilution or suspension of an inventive pharmaceutical composition or compound. In some embodiments, the inventive pharmaceutical composition or compound provided in the first container and the second container are combined to form one unit dosage form.

[00327] Thus, in another aspect, provided are kits for treating and/or preventing a pathological condition of a subject. In certain embodiments, the kits include a first container comprising a compound of the present invention, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, polymorph, or composition thereof; and an instruction for administering the compound, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, polymorph, or composition thereof, to the subject to treat and/or prevent the pathological condition. In certain embodiments, the kits of the present invention include one or more additional approved therapeutic agents for use as a combination therapy. In certain embodiments, the instruction includes a notice in the form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use, or sale for human administration.

Methods of Treatment and Uses


[00328] The compounds of the invention and pharmaceutical compositions thereof are expected to be useful in the treatment and/or prevention of a pathological condition in a subject. In one aspect, provided herein are methods of treating and/or preventing a pathological condition in a subject, the methods including administering to the subject a therapeutically or prophylactically effective amount of a compound of the invention, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, and optionally a pharmaceutically acceptable excipient.

[00329] In another aspect of the present invention, provided are methods of treating and/or preventing a pathological condition, the methods including mixing blood or a component thereof (*e.g.*, red blood cells) with a therapeutically or prophylactically effective amount of a compound of the invention, or a pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, or a pharmaceutical composition of the invention; and administering to the subject the mixture of blood or a component thereof (*e.g.*, red blood cells) and the compound, or the pharmaceutically acceptable salt, tautomer, stereoisomer, solvate, hydrate, or polymorph thereof, or the pharmaceutical composition. The blood may be whole blood or a fluid comprising one or more components of whole blood (*e.g.*, red blood cells, white blood cells, plasma, clotting factors, and platelets). In certain embodiments, the mixture is administered intravenously to the subject.

[00330] In another aspect of the present invention, provided are compounds of the invention, and pharmaceutically acceptable salts, tautomers, stereoisomers, solvates, hydrates, and polymorphs thereof, for use in treatment and/or prevention of a pathological condition.

[00331] The present invention stems from the recognition that the pathogenesis of various pathological conditions, including oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and other pathological conditions, involves free iron and the generation of reactive oxygen species (ROS), including superoxide anion, hydrogen peroxide, hypochlorous acid, and hydroxyl radicals, and other longer lived, free radicals. Such radicals are now realized to be important contributors to these pathological conditions. Free iron is known to contribute to the formation of reactive oxygen species. For example, Fe^{+2} ions in biological systems react with oxygen species to produce highly reactive hydroxyl radicals via

the Fenton reaction (see scheme below). The hydroxyl radical is a highly effective oxidizing agent, reacting at a diffusion-controlled rate with most organic species, such as nucleic acids, proteins, and lipids. Furthermore, superoxide anions or a biological reductant (*e.g.*, ascorbic acid) can reduce the resulting Fe^{+3} ion back to Fe^{+2} for continued peroxide reduction, thus a problematic cycle.



[00332] Therefore, pathological conditions that lead to bleeding and/or an inflammatory response involve the possibility that reactive oxygen species will come in contact with Fe^{+2} ions to produce highly reactive and damaging hydroxyl radicals. That is, the iron released from red blood cells react with oxygen species produced by inflammatory cells such as neutrophils to produce hydroxyl radicals that cause cell and tissue injury. The solution, therefore, is chelation and removal of the unmanaged iron.

[00333] Without wishing to be bound by any particular theory, the compounds of the invention are thought to chelate or sequester a metal, and, in certain embodiments, the pathological condition is responsive to chelation or sequestration of the metal. In certain embodiments, the metal is iron (*e.g.*, Fe(II) or Fe(III)), aluminum, thallium (*e.g.*, Tl(I) or Tl(III)), chromium (*e.g.*, Cr(III) or Cr(VI)), magnesium, calcium, strontium, nickel (*e.g.*, Ni(II)), manganese (*e.g.*, Mn(II)), cobalt (*e.g.*, Co(II) or Co(III)), copper (*e.g.*, Cu(I) or Cu(II)), zinc, silver (*e.g.*, Ag(I)), sodium, potassium, cadmium (*e.g.*, Cd(II)), mercury (*e.g.*, Hg(I) or Hg(II)), lead (*e.g.*, Pb(II) or Pb(IV)), antimony (*e.g.*, Sb(III) or Sb(V)), molybdenum (*e.g.*, Mo(III) or Mo(VI)), tungsten (*e.g.*, W(VI)), a lanthanide (*e.g.*, cerium, such as Ce(III) or Ce(IV)), or an actinide (*e.g.*, uranium, such as U(VI)). In certain embodiments, the metal is a trivalent metal. In certain embodiments, the metal is iron (*e.g.*, Fe(III)). In certain embodiments, the metal is aluminum. In certain embodiments, the metal is Tl(III) , Cr(III) , Co(III) , Sb(III) , Mo(III) , or Ce(III) . In certain embodiments, the metal is a monovalent metal (*e.g.*, Tl(I) , Cu(I) , Ag(I) , Na(I) , K(I) , or Hg(I)). In certain embodiments, the metal is a divalent metal (*e.g.*, Fe(II) , Mg(II) , Ca(II) , Sr(II) , Ni(II) , Mn(II) , Co(II) , Cu(II) , Zn(II) , Cd(II) , Hg(II) , or Pb(II)). In certain embodiments, the metal is a tetravalent metal (*e.g.*, Pb(IV) or Ce(IV)). In certain embodiments, the metal is a pentavalent metal (*e.g.*, Sb(V)). In certain embodiments, the metal is a hexavalent metal (*e.g.*, Cr(VI) , Mo(VI) , W(VI) , or U(VI)).

[00334] In certain embodiments, the subject administered the inventive compound or pharmaceutical composition is a mammal. In certain embodiments, the subject is a human. In certain embodiments, the subject is a domesticated animal, such as a dog, cat, cow, pig, horse, sheep, or goat. In certain embodiments, the subject is a companion animal such as a dog or cat. In certain embodiments, the subject is a livestock animal such as a cow, pig, horse, sheep, or goat. In certain embodiments, the subject is a zoo animal. In another embodiment, the subject is an experimental animal such as a rodent or non-human primate.

[00335] The inventive compounds, pharmaceutical compositions, and methods may also be useful for the treatment and/or prevention of infectious diseases in a subject. Infectious diseases are typically caused by microbial pathogens (*e.g.*, viruses, bacteria, parasites (*e.g.*, protozoa and multicellular parasites), and fungi) into the cells (“host cells”) of a subject (“host”). Iron is an oxidant as well as a nutrient for many microorganisms. To survive and replicate, microbial pathogens must acquire iron from their host. Highly virulent microbial strains usually possess powerful mechanisms for obtaining iron from their host. Depriving the pathogenic microbes of iron may inhibit their activities and may be useful for the treatment and/or prevention of the infectious diseases caused by microbes. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a viral infection. In certain embodiments, the pathological condition is a bacterial infection. In certain embodiments, the pathological condition is a parasitic infection. In certain embodiments, the pathological condition is a protozoan infection. In certain embodiments, the pathological condition is malaria. Malaria is typically caused by parasites of the genus *Plasmodium* (phylum *Apicomplexa*), including, but not limited to, the species *P. falciparum*, *P. malariae*, *P. ovale*, *P. vivax*, and *P. knowlesi*. In certain embodiments, the pathological condition is a multicellular-parasitic infection. In certain embodiments, the pathological condition is a fungal infection.

[00336] In certain embodiments, methods are provided herein that are useful in the treatment and/or prevention of metal overload in a subject. The amount of free metal (*e.g.*, a trivalent metal, such as iron(III) or aluminum) may be elevated in the subject (*e.g.*, in the serum or in a cell), such as when there is insufficient storage capacity for the metal or an abnormality in the metal storage system that leads to metal release. In certain embodiments, the metal overload is iron overload (*e.g.*, Fe(III) overload or Fe(II) overload).

[00337] Iron overload conditions or diseases can be characterized by global iron overload or focal iron overload. Global iron overload conditions generally involve an excess

of iron in multiple tissues or excess iron located throughout an organism. Global iron overload conditions can result from excess uptake of iron by a subject, excess storage and/or retention of iron, from, for example, dietary iron or blood transfusions. One global iron overload condition is primary hemochromatosis, which is typically a genetic disorder. A second global iron overload condition is secondary hemochromatosis, which is typically the result of receiving multiple (chronic) blood transfusions. Blood transfusions are often required for subjects suffering from thalassemia or sickle cell anemia. A type of dietary iron overload is referred to as Bantu siderosis, which is associated with the ingestion of homebrewed beer with high iron content. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is global iron overload. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is focal iron overload. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is primary hemochromatosis. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is secondary hemochromatosis. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is Bantu siderosis.

[00338] In focal iron overload conditions, the excess iron is limited to one or a few cell types or tissues or a particular organ. Alternatively, symptoms associated with the excess iron are limited to a discrete organ, such as the heart, lungs, liver, pancreas, kidneys, or brain. It is believed that focal iron overload can lead to neurological or neurodegenerative disorders such as Parkinson's disease, Alzheimer's disease, Huntington's disease, neuroferritinopathy, amyotrophic lateral sclerosis, and multiple sclerosis. Pathological conditions that benefit from metal chelation or sequestration are often associated with deposition of the metal in the tissues of a subject. Deposition can occur globally or focally. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a neurological or neurodegenerative disorder. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a neurological disorder. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and

methods of the invention is a neurodegenerative disorder. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is Parkinson's disease. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is Alzheimer's disease. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is Huntington's disease. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is neuroferritinopathy. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is amyotrophic lateral sclerosis. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is multiple sclerosis.

[00339] While humans have a highly efficient iron management system in which they absorb and excrete about 1 mg of iron daily, there is no conduit for the excretion of excess metal. Transfusion-dependent anemias, like thalassemia, lead to a build up of iron in the liver, heart, pancreas, and elsewhere resulting in (i) liver disease that may progress to cirrhosis (Angelucci *et al.*, "Hepatic Iron Concentration and Total Body Iron Stores in Thalassemia Major." *N. Engl. J. Med.* **2000**, 343, 327–331; Bonkovsky *et al.*, "Iron-Induced Liver Injury." *Clin. Liver Dis.* **2000**, 4, 409–429; Peitrangelo, "Mechanism of Iron Toxicity." *Adv. Exp. Med. Biol.* **2002**, 509, 19–43), (ii) diabetes related both to iron-induced decreases in pancreatic beta -cell secretion and to increases in hepatic insulin resistance (Cario *et al.*, "Insulin Sensitivity and β -Cell Secretion in Thalassemia Major with Secondary Haemochromatosis: Assessment by Oral Glucose Tolerance Test." *Eur. J. Pediatr.* **2004**, 162, 139–146; Wojcik *et al.*, "Natural History of C282Y Homozygotes for Haemochromatosis." *Can. J. Gastroenterol.* **2002**, 16, 297–302), and (iii) heart disease. Relative excess iron has been associated with increased risk of heart disease. Cardiac failure is still the leading cause of death in thalassemia major and related forms of transfusional iron overload (Brittenham, "Disorders of Iron Metabolism: Iron Deficiency and Overload." In: Hoffman *et al.*, editors. *Hematology: Basic Principles and Practice*. 3. Churchill Livingstone; New York: 2000. pp. 397–428; Brittenham *et al.*, "Efficacy of Deferoxamine in Preventing Complications of Iron Overload in Patients with Thalassemia Major." *N. Engl. J. Med.* **1994**, 331, 567–573; Zurlo *et al.*, "Survival and Causes of Death in Thalassemia Major." *Lancet*.

1989, 2, 27–30). There is a strong correlation between serum ferritin levels, inflammatory biomarkers such as C-reactive protein and interleukin-1, and mortality in a subset of patients with peripheral arterial disease; phlebotomy and iron chelation has been used to mitigate that risk. Treatment with an iron chelator would reduce iron stores, reduce serum ferritin and potentially reduce the incidence of heart disease and stroke. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is transfusional iron overload. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is transfusion-dependent anemia. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is thalassemia. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a liver disease (*e.g.*, hepatitis B, hepatitis C, and liver cirrhosis). In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a heart disease (*e.g.*, cardiomyopathy, coronary heart disease, inflammatory heart disease, ischemic heart disease, valvular heart disease, hypertensive heart disease, and atherosclerosis). In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a pancreas disease. In certain embodiments, the pathological condition that is treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is diabetes.

[00340] Moreover, the compounds, pharmaceutical compositions, and methods of the present invention may be useful in the treatment and/or prevention of metal overload where the metal is not iron. All metals described herein are contemplated for chelation by the inventive compounds. In certain embodiments, the metal is aluminum. In certain embodiments, the metal is Tl(III), Cr(III), Co(III), Sb(III), Mo(III), or Ce(III). In certain embodiments, the metal is a monovalent metal (*e.g.*, Tl(I), Cu(I), Ag(I), Na(I), K(I), or Hg(I)). In certain embodiments, the metal is a divalent metal (*e.g.*, Fe(II), Mg(II), Ca(II), Sr(II), Ni(II), Mn(II), Co(II), Cu(II), Zn(II), Cd(II), Hg(II), or Pb(II)). In certain embodiments, the metal is a tetravalent metal (*e.g.*, Pb(IV) or Ce(IV)). In certain embodiments, the metal is a pentavalent metal (*e.g.*, Sb(V)). In certain embodiments, the metal is a hexavalent metal (*e.g.*, Cr(VI), Mo(VI), W(VI), or U(VI)).

[00341] In certain embodiments, the metal overload is aluminum overload, chromium overload, magnesium overload, calcium overload, strontium overload, nickel overload, manganese overload, cobalt overload, copper overload, zinc overload, silver overload, sodium overload, potassium overload, cadmium overload, mercury overload, lead overload, molybdenum overload, tungsten overload, or actinide overload (*e.g.*, uranium overload). In certain embodiments, the metal overload is trivalent metal overload. In certain embodiments, the metal overload is aluminum overload. In certain embodiments, the metal overload is Cr(III) overload, Mo(III) overload, or Co(III) overload). In certain embodiments, the metal overload is monovalent metal overload (*e.g.*, Cu(I) overload, Ag(I) overload, Na(I) overload, K(I) overload, or Hg(I) overload). In certain embodiments, the metal overload is divalent metal overload (*e.g.*, Mg(II) overload, Ca(II) overload, Sr(II) overload, Ni(II) overload, Mn(II) overload, Co(II) overload, Cu(II) overload, Zn(II) overload, Cd(II) overload, Hg(II) overload, or Pb(II) overload). In certain embodiments, the metal overload is tetravalent metal overload (*e.g.*, Pb(IV) overload). In certain embodiments, the metal overload is pentavalent metal overload. In certain embodiments, the metal overload is hexavalent metal overload (*e.g.*, Cr(VI) overload, Mo(VI) overload, W(VI) overload, or U(VI) overload).

[00342] The inventive compounds, pharmaceutical compositions, and methods may also be useful in treating and/or preventing metal poisoning in a subject. Metal poisoning may be caused by metal toxicity to a subject. For example, metals with little or no endogenous function may find their way into the body of a subject and cause damage. Heavy metal ions such as Hg(II) can replace ions such as Zn(II) in metalloproteins and render them inactive, resulting in serious acute or chronic toxicity that can end in a patient's death or in birth defects. Even more significantly, radioactive isotopes of the lanthanide (*e.g.*, cerium) and actinide (*e.g.*, uranium) series can cause grave illness on an individual exposed to them by mouth, air, or skin contact. Such exposure could result not only from the detonation of a nuclear bomb or a "dirty bomb" composed of nuclear waste, but also from the destruction of a nuclear power facility. In certain embodiments, the metal poisoning is iron poisoning, aluminum poisoning, thallium poisoning, chromium poisoning, magnesium poisoning, calcium poisoning, strontium poisoning, nickel poisoning, manganese poisoning, cobalt poisoning, copper poisoning, zinc poisoning, silver poisoning, sodium poisoning, potassium poisoning, cadmium poisoning, mercury poisoning, lead poisoning, antimony poisoning, molybdenum poisoning, tungsten poisoning, lanthanide poisoning (*e.g.*, cerium poisoning), or actinide poisoning (*e.g.*, uranium poisoning). In certain embodiments, the metal poisoning is iron poisoning (*e.g.*, Fe(II) poisoning or Fe(III) poisoning). In certain embodiments, the metal

poisoning is aluminum poisoning. In certain embodiments, the metal poisoning is trivalent metal poisoning (*e.g.*, Fe(III) poisoning, Al(III) poisoning, Tl(III) poisoning, Cr(III) poisoning, Co(III) poisoning, Sb(III) poisoning, Mo(III) poisoning, or Ce(III) poisoning). In certain embodiments, the metal poisoning is monovalent metal poisoning (*e.g.*, Tl(I) poisoning, Cu(I) poisoning, Ag(I) poisoning, Na(I) poisoning, K(I) poisoning, or Hg(I) poisoning). In certain embodiments, the metal poisoning is divalent metal poisoning (*e.g.*, Fe(II) poisoning, Mg(II) poisoning, Ca(II) poisoning, Sr(II) poisoning, Ni(II) poisoning, Mn(II) poisoning, Co(II) poisoning, Cu(II) poisoning, Zn(II) poisoning, Cd(II) poisoning, Hg(II) poisoning, or Pb(II) poisoning). In certain embodiments, the metal poisoning is tetravalent metal poisoning (*e.g.*, Pb(IV) or Ce(IV) poisoning). In certain embodiments, the metal poisoning is pentavalent metal poisoning (*e.g.*, Sb(V) poisoning). In certain embodiments, the metal poisoning is hexavalent metal poisoning (*e.g.*, Cr(VI) poisoning, Mo(VI) poisoning, W(VI) poisoning, or U(VI) poisoning).

[00343] The compounds, pharmaceutical compositions, and methods of the invention are also useful in treating and/or preventing oxidative stress in a subject. In a subject who suffers from oxidative stress and thus needs oxidative stress reduction, the iron released from red blood cells of the subject may react with oxygen species produced by inflammatory cells such as neutrophils to produce hydroxyl radicals that cause cell and tissue injury. Chelation and removal of the unmanaged iron may prevent or impede these harmful reactions and, therefore, reduce oxidative stress. A subject in need of oxidative stress reduction can have one or more of the following conditions: decreased levels of reducing agents, increased levels of reactive oxygen species, mutations in or decreased levels of antioxidant enzymes (*e.g.*, Cu/Zn superoxide dismutase, Mn superoxide dismutase, glutathione reductase, glutathione peroxidase, thioredoxin, thioredoxin peroxidase, DT-diaphorase), mutations in or decreased levels of metal-binding proteins (*e.g.*, transferrin, ferritin, ceruloplasmin, albumin, metallothionein), mutated or overactive enzymes capable of producing superoxide (*e.g.*, nitric oxide synthase, NADPH oxidases, xanthine oxidase, NADH oxidase, aldehyde oxidase, dihydroorotate dehydrogenase, cytochrome c oxidase), and radiation injury. Increased or decreased levels of reducing agents, reactive oxygen species, and proteins are determined relative to the amount of such substances typically found in healthy persons. A subject in need of oxidative stress reduction can be suffering from an ischemic episode. Ischemic episodes can occur when there is mechanical obstruction of the blood supply, such as from arterial narrowing or disruption. Myocardial ischemia, which can give rise to angina pectoris and myocardial infarctions, results from inadequate circulation of blood to the myocardium,

usually due to coronary artery disease. Ischemic episodes in the brain that resolve within 24 hours are referred to as transient ischemic attacks. A longer-lasting ischemic episode, a stroke, involves irreversible brain damage, where the type and severity of symptoms depend on the location and extent of brain tissue whose access to blood circulation has been compromised. A subject at risk of suffering from an ischemic episode typically suffers from atherosclerosis, other disorders of the blood vessels, increased tendency of blood to clot, or heart disease.

[00344] A subject in need of oxidative stress reduction can be suffering from inflammation. Inflammation is a fundamental pathologic process consisting of a complex of cytologic and chemical reactions that occur in blood vessels and adjacent tissues in response to an injury or abnormal stimulation caused by a physical, chemical, or biologic agent. Inflammatory disorders are characterized inflammation that lasts for an extended period (*i.e.*, chronic inflammation) or that damages tissue. Such inflammatory disorders can affect a wide variety of tissues, such as respiratory tract, joints, bowels, and soft tissue. The compounds or pharmaceutical compositions of the invention can be used to treat these pathological conditions. Not wishing to be bound by any theory, it is believed that the compounds of the invention derive their ability to reduce oxidative stress through various mechanisms. In one mechanism, the compound binds to a metal, particularly a redox-active metal (*e.g.*, iron), and fills all of the coordination sites of the metal. When all of the metal coordination sites are filled, it is believed that oxidation and/or reducing agents have a diminished ability to interact with the metal and cause redox cycling. In another mechanism, the compound stabilizes the metal in a particular oxidation state, such that it is less likely to undergo redox cycling. In yet another mechanism, the compound itself has antioxidant activity (*e.g.*, free radical scavenging, scavenging of reactive oxygen or nitrogen species). Desferrithiocin and desazadesferrithiocin, and their derivatives and analogs, are known to have intrinsic antioxidant activity, as described in U.S. Application Publication No. 2004/0044220, published March 4, 2004 and now abandoned; U.S. Application Publication No. 2004/0132789 and now abandoned, published July 8, 2004; International PCT Application Publication No. WO2004/017959, published March 4, 2004; U.S. Application Publication No. 2005/0234113, published October 20, 2005 and now abandoned; U.S. Application Publication No. 2008/0255081, published October 16, 2008 and now abandoned; U.S. Application Publication No. 2003/0236417, published December 25, 2003 and now abandoned; U.S. Patent Application, U.S.S.N. 61/576,920, filed December 16, 2011; U.S. Patent Application, U.S.S.N. 61/576,913, filed December 16, 2011; and U.S. Patent Nos.:

6,083,966, 6,559,315, 6,525,080, 6,521,652, 7,126,004, 7,531,563, and 8,008,502; each of which are incorporated herein by reference. The compounds of the invention can be used to treat these pathological conditions. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is oxidative stress. In certain embodiments, the compounds, pharmaceutical compositions, and methods of the present invention are useful in the reduction of oxidative stress. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is radiation injury. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is inflammation.

[00345] The invention also provides compounds, pharmaceutical compositions, and methods for the treatment of macular degeneration. Without wishing to be bound by a particular theory, the compounds of the invention are able to get into the eye. See, *e.g.*, U.S. Patent Application, U.S.S.N. 61/576,920, filed December 16, 2011; U.S. Patent Application, U.S.S.N. 61/576,913, filed December 16, 2011, International PCT Application Publication No. WO 2013/090750, published June 20, 2013; and International PCT Application Publication No. WO 2013/090766, published June 20, 2013. The compounds of the invention are then able to chelate and remove iron from the eye thereby preventing Fe^{+2} from generating reactive oxygen species. The local accumulation of iron is thought to contribute to macular degeneration. Therefore, the removal of iron from the eye (including the retina) can prevent and treat macular degeneration. In the treatment of macular degeneration, the compound of the invention or a pharmaceutical composition thereof may be administered systemically or ocularly. In certain embodiments, the compound or composition is administered orally. In other embodiments, the compound or composition is administered to the eye using eyedrops or an ointment suitable for ocular administration.

[00346] The compounds of the invention and pharmaceutical compositions thereof are expected to be useful in the treatment of head injury, particularly those involving bleeding into the brain or other parts of the central nervous system. Without wishing to be bound by any particular theory, the compounds of the invention are thought to chelate the iron from red blood cells the blood resulting from the head injury, thereby preventing iron ions from generating reactive oxygen species. In the case of head injury resulting in bleeding into the central nervous system where the vasculature has been compromised a compound being used may or may not have the ability to cross the blood brain barrier. In certain embodiments, the

compound being used to treat a head injury in a subject is able to cross the blood brain barrier. In other embodiments, the compounds are not able to cross the blood brain barrier. Certain compounds of the invention have been found in the CSF after systemic administration (po and sc).

[00347] Head injuries come in various forms and results from various causes. In certain embodiments, the injury is an injury to the head that penetrates the skull. In other embodiments, the head injury being treated is a closed head injury, which does not penetrate the skull. Closed head injuries result from a variety of causes including accidents including vehicular accidents, falls, and assaults. Types of closed head injuries include concussions, brain contusions, diffuse axonal injury, and hematoma. In certain embodiments, the closed head injury being treated in the present invention includes closed head injuries that result in blood outside the blood vessels of the brain.

[00348] The local accumulation of iron from the bleeding is thought to contribute to after effects associated with closed head injury. By assisting the clearance of iron from the brain the effects of the bleeding are minimized.

[00349] In the treatment of closed head injury, the compound of the invention or a pharmaceutical composition thereof may be administered systemically, for example, parenterally or orally. In certain embodiments, the compound or composition is administered orally. In other embodiments, the compound or composition is administered parenterally (e.g., intravenously).

[00350] Reactive oxygen species have been implicated in the pathogenesis of inflammatory bowel disease (IBD). Grisham *et al.*, "Neutrophil-mediated mucosal injury. Role of reactive oxygen metabolites." *Dig. Dis. Sci.* 33:6S-15S, 1988; Allgayer "Clinical relevance of oxygen radicals in inflammatory bowel disease—facts and fashion." *Klin. Wochenschr.* 69:1001-1003, 1991; Yamamada *et al.* "Role of neutrophil-derived oxidants in the pathogenesis of intestinal inflammation." *Klin. Wochenschr.* 69:988-944, 1991; Babbs, "Oxygen radicals in ulcerative colitis." *Free Radic. Biol. Med.* 13:169-181, 1992. The present invention provides for the treatment or prevention of IBD. DFO, an iron chelator, has been discovered to prevent acetic acid-induced colitis in rats, an animal model of IBD. See, e.g., U.S. Patent Application, U.S.S.N. 61/576,920, filed December 16, 2011; U.S. Patent Application, U.S.S.N. 61/576,913, filed December 16, 2011; Bergeron *et al.*, "Prevention of Acetic Acid-Induced Colitis by Desferrithiocin Analogs in a Rat Model." *Digestive Diseases and Sciences*, 48(2):399-407, February 2003. The compounds used in the inventive treatment are thought to prevent or eliminate the generation of reactive oxygen species or other longer-

lived, more stable radicals that may be responsible for the tissue damage and inflammation seen in subjects with IBD. Another possible mechanism of action of the compounds useful in the invention is the chelation of metal, such as iron, which may contribute to the generation of reactive oxygen species, such as hydroxyl radicals and hydrogen peroxide, that cause cell damage.

[00351] The present invention may also be useful in treating a subject diagnosed with IBD. The treatment may be used to treat the subject long term or may be used to treat a subject with a flare up of IBD. A therapeutically effective amount of a compound of the invention or pharmaceutical composition thereof is administered to a subject in need thereof to treat IBD. In certain embodiments, treatment with a compound of the invention leads to reduced levels of reactive oxygen species in the intestines, specifically the intestinal mucosa. The compound or composition thereof may be administered to a subject once or multiple times in the treatment of IBD.

[00352] In the treatment of IBD, the compound of the invention or a pharmaceutical composition thereof may be administered systemically, for example, parenterally or orally. In certain embodiments, the compound or composition is administered orally. In other embodiments, the compound or composition is administered parenterally (*e.g.*, intravenously). In certain embodiments, the compound or a composition is administered rectally.

[00353] The methods of the present invention are also useful in the treatment and/or prevention of stroke. The inventive treatment typically leads to a better and/or faster recovery from stroke. The stroke being treated may be either an ischemic stroke or a hemorrhagic stroke. In the treatment of an ischemic stroke, a compound of the invention or a pharmaceutical composition thereof is administered to a subject to prevent or minimize the damage due to reperfusion injury after the blood supply to the affected part of the brain is restored. The compound is thought to prevent the generation of reactive oxygen species by either chelating iron responsible for the generation of such species and/or quenching such radical species when they do occur. In hemorrhagic stroke, the compound is thought to work by similar mechanisms although the sequestering of iron from the blood in the brain is probably the predominate mechanism by which the inventive treatment works. The mechanism of action of the compound of the invention is similar to that in the treatment of head injury.

[00354] The compound being used in the treatment may have the ability to cross the blood brain barrier. In certain embodiments, when the subject has been diagnosed with an ischemic stroke, the compound used in the treatment can pass through the blood brain barrier.

[00355] Moreover, the present invention may be useful in treating a subject after the subject has been diagnosed with having a stroke, or a subject who is susceptible to having a stroke may be administered a compound of the invention or composition thereof to prevent or minimize the stroke's effects. In certain embodiments, the compound is administered as quickly as possible after a subject has been diagnosed with having a stroke. In certain embodiments, the compound is administered to the subject while the stroke is still occurring. In certain embodiments, the compound or a composition thereof is administered to a subject who has a history of strokes or is susceptible to having a stroke because of the subject's underlying medical condition. The compound or composition thereof may be administered once or multiple times in the treatment of stroke.

[00356] In the treatment of stroke the compound of the invention or a pharmaceutical composition thereof may be administered systemically, for example, parenterally or orally. In certain embodiments, the compound or composition is administered orally. In other embodiments, the compound or composition is administered parenterally (*e.g.*, intravenously).

[00357] The present invention also provides for the treatment of reperfusion injury. Reperfusion injury may occur in any area of the body where the blood supply has been compromised. In certain embodiments, the reperfusion injury being treated occurs in the heart. In other embodiments, the reperfusion injury occurs in the brain, for example, as discussed above in the context of a stroke. The inventive treatment minimizes reperfusion injury once the blood supply to the affected organ or tissue is restored. In the treatment and/or prevention of reperfusion injury, a compound of the present invention or pharmaceutical composition thereof is administered to a subject who is suffering from ischemia of a tissue or organ. Without wishing to be bound by any particular theory, the compound of the invention is thought to prevent the generation of reactive oxygen species by either chelating iron responsible for the generation of such species and/or quenching such radical species when they do occur.

[00358] The present invention may be useful in treating a subject after the subject has been diagnosed with ischemia of a particular organ or tissue. A therapeutically effective amount of a compound of the invention or composition thereof is administered to a subject to prevent or minimize reperfusion injury. In certain embodiments, the compound is

administered as quickly as possible after a subject has been diagnosed with ischemia. In certain embodiments, the compound is administered to the subject at risk of ischemia. In certain embodiments, the compound or a composition thereof is administered to a subject who is about to undergo a procedure that may lead to ischemia of an organ or tissue (*e.g.*, cardiac surgery). In certain embodiments, the compound or a composition thereof is used to prevent reperfusion injury in a transplanted organ. In certain embodiments, the compound or composition thereof is used to perfuse an isolated organ being prepared for donation. The compound or composition thereof may be administered to a subject once or multiple times in the treatment of reperfusion injury.

[00359] In the prevention or treatment of reperfusion injury, the compound of the invention or a pharmaceutical composition thereof may be administered systemically, for example, parenterally or orally. In certain embodiments, the compound or composition is administered orally. In other embodiments, the compound or composition is administered parenterally (*e.g.*, intravenously). In certain embodiments, the compound or a composition is administered locally to the organ or tissue suffering from ischemia.

[00360] The inventive compounds, or pharmaceutical compositions thereof, may also be useful in the treatment and/or prevention of a neoplastic disease or preneoplastic condition. A neoplastic disease (*i.e.*, neoplasm) is characterized by an abnormal tissue that grows by cellular proliferation more rapidly than normal tissue. The abnormal tissue continues to grow after the stimuli that initiated the new growth cease. Neoplasms show a partial or complete lack of structural organization and functional coordination with the normal tissue, and usually form a distinct mass of tissue that may be benign or malignant. A malignant neoplastic disease is also known as cancer. Neoplasms can occur, for example, in a wide variety of tissues including brain, skin, mouth, nose, esophagus, lungs, stomach, pancreas, liver, bladder, ovary, uterus, testicles, colon, and bone, as well as the immune system (lymph nodes) and endocrine system (thyroid gland, parathyroid glands, adrenal gland, thymus, pituitary gland, pineal gland). In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is a benign neoplastic disease. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is cancer. In certain embodiments, the pathological condition that may be treated and/or prevented by the compounds, pharmaceutical compositions, and methods of the invention is acoustic neuroma; adenocarcinoma; adrenal gland cancer; anal cancer; angiosarcoma (*e.g.*, lymphangiosarcoma,

lymphangioendotheliosarcoma, hemangiosarcoma); appendix cancer; benign monoclonal gammopathy; biliary cancer (*e.g.*, cholangiocarcinoma); bladder cancer; breast cancer (*e.g.*, adenocarcinoma of the breast, papillary carcinoma of the breast, mammary cancer, medullary carcinoma of the breast); brain cancer (*e.g.*, meningioma, glioblastomas, glioma (*e.g.*, astrocytoma, oligodendroglioma), medulloblastoma); bronchus cancer; carcinoid tumor; cervical cancer (*e.g.*, cervical adenocarcinoma); choriocarcinoma; chordoma; craniopharyngioma; colorectal cancer (*e.g.*, colon cancer, rectal cancer, colorectal adenocarcinoma); connective tissue cancer; epithelial carcinoma; ependymoma; endotheliosarcoma (*e.g.*, Kaposi's sarcoma, multiple idiopathic hemorrhagic sarcoma); endometrial cancer (*e.g.*, uterine cancer, uterine sarcoma); esophageal cancer (*e.g.*, adenocarcinoma of the esophagus, Barrett's adenocarcinoma); Ewing sarcoma; eye cancer (*e.g.*, intraocular melanoma, retinoblastoma); familial hypereosinophilia; gall bladder cancer; gastric cancer (*e.g.*, stomach adenocarcinoma); gastrointestinal stromal tumor (GIST); germ cell cancer; head and neck cancer (*e.g.*, head and neck squamous cell carcinoma, oral cancer (*e.g.*, oral squamous cell carcinoma), throat cancer (*e.g.*, laryngeal cancer, pharyngeal cancer, nasopharyngeal cancer, oropharyngeal cancer)); hematopoietic cancers (*e.g.*, leukemia such as acute lymphocytic leukemia (ALL) (*e.g.*, B-cell ALL, T-cell ALL), acute myelocytic leukemia (AML) (*e.g.*, B-cell AML, T-cell AML), chronic myelocytic leukemia (CML) (*e.g.*, B-cell CML, T-cell CML), and chronic lymphocytic leukemia (CLL) (*e.g.*, B-cell CLL, T-cell CLL)); lymphoma such as Hodgkin lymphoma (HL) (*e.g.*, B-cell HL, T-cell HL) and non-Hodgkin lymphoma (NHL) (*e.g.*, B-cell NHL such as diffuse large cell lymphoma (DLCL) (*e.g.*, diffuse large B-cell lymphoma), follicular lymphoma, chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), marginal zone B-cell lymphomas (*e.g.*, mucosa-associated lymphoid tissue (MALT) lymphomas, nodal marginal zone B-cell lymphoma, splenic marginal zone B-cell lymphoma), primary mediastinal B-cell lymphoma, Burkitt lymphoma, lymphoplasmacytic lymphoma (*i.e.*, Waldenström's macroglobulinemia), hairy cell leukemia (HCL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma and primary central nervous system (CNS) lymphoma; and T-cell NHL such as precursor T-lymphoblastic lymphoma/leukemia, peripheral T-cell lymphoma (PTCL) (*e.g.*, cutaneous T-cell lymphoma (CTCL) (*e.g.*, mycosis fungoides, Sezary syndrome), angioimmunoblastic T-cell lymphoma, extranodal natural killer T-cell lymphoma, enteropathy type T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, and anaplastic large cell lymphoma); a mixture of one or more leukemia/lymphoma as described above; and multiple myeloma (MM)), heavy chain disease

(*e.g.*, alpha chain disease, gamma chain disease, mu chain disease); hemangioblastoma; hypopharynx cancer; inflammatory myofibroblastic tumors; immunocytic amyloidosis; kidney cancer (*e.g.*, nephroblastoma *a.k.a.* Wilms' tumor, renal cell carcinoma); liver cancer (*e.g.*, hepatocellular cancer (HCC), malignant hepatoma); lung cancer (*e.g.*, bronchogenic carcinoma, small cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung); leiomyosarcoma (LMS); mastocytosis (*e.g.*, systemic mastocytosis); muscle cancer; myelodysplastic syndrome (MDS); mesothelioma; myeloproliferative disorder (MPD) (*e.g.*, polycythemia Vera (PV), essential thrombocytosis (ET), agnogenic myeloid metaplasia (AMM) *a.k.a.* myelofibrosis (MF), chronic idiopathic myelofibrosis, chronic myelocytic leukemia (CML), chronic neutrophilic leukemia (CNL), hypereosinophilic syndrome (HES)); neuroblastoma; neurofibroma (*e.g.*, neurofibromatosis (NF) type 1 or type 2, schwannomatosis); neuroendocrine cancer (*e.g.*, gastroenteropancreatic neuroendocrine tumor (GEP-NET), carcinoid tumor); osteosarcoma (*e.g.*, bone cancer); ovarian cancer (*e.g.*, cystadenocarcinoma, ovarian embryonal carcinoma, ovarian adenocarcinoma); papillary adenocarcinoma; pancreatic cancer (*e.g.*, pancreatic adenocarcinoma, intraductal papillary mucinous neoplasm (IPMN), Islet cell tumors); penile cancer (*e.g.*, Paget's disease of the penis and scrotum); pinealoma; primitive neuroectodermal tumor (PNT); plasma cell neoplasia; paraneoplastic syndromes; intraepithelial neoplasms; prostate cancer (*e.g.*, prostate adenocarcinoma); rectal cancer; rhabdomyosarcoma; salivary gland cancer; skin cancer (*e.g.*, squamous cell carcinoma (SCC), keratoacanthoma (KA), melanoma, basal cell carcinoma (BCC)); small bowel cancer (*e.g.*, appendix cancer); soft tissue sarcoma (*e.g.*, malignant fibrous histiocytoma (MFH), liposarcoma, malignant peripheral nerve sheath tumor (MPNST), chondrosarcoma, fibrosarcoma, myxosarcoma); sebaceous gland carcinoma; small intestine cancer; sweat gland carcinoma; synovioma; testicular cancer (*e.g.*, seminoma, testicular embryonal carcinoma); thyroid cancer (*e.g.*, papillary carcinoma of the thyroid, papillary thyroid carcinoma (PTC), medullary thyroid cancer); urethral cancer; vaginal cancer; or vulvar cancer (*e.g.*, Paget's disease of the vulva).

[00361] A preneoplastic condition precedes the formation of a benign or malignant neoplasm. A precancerous lesion typically forms before a malignant neoplasm. Preneoplastic conditions include, but are not limited to, photodermatitis, x-ray dermatitis, tar dermatitis, arsenic dermatitis, lupus dermatitis, senile keratosis, Paget disease, condylomata, burn scar, syphilitic scar, fistula scar, ulcer cruris scar, chronic ulcer, varicose ulcer, bone fistula, rectal fistula, Barrett esophagus, gastric ulcer, gastritis, cholelithiasis, kraurosis vulvae, nevus pigmentosus, Bowen dermatosis, xeroderma pigmentosum, erythroplasia, leukoplakia, Paget

disease of bone, exostoses, ecchondroma, osteitis fibrosa, leontiasis ossea, neurofibromatosis, polyposis, hydatidiform mole, adenomatous hyperplasia, and struma nodosa. The compounds, pharmaceutical compositions, and methods of the present invention can be used to treat and/or prevent these preneoplastic conditions. .

[00362] Imaging or examining one or more organs, tissues, tumors, or a combination thereof can be conducted after a metal salt of a compound of the invention is administered to a subject. The methods of imaging and examining are intended to encompass various instrumental techniques used for diagnosis, such as x-ray methods (including CT scans and conventional x-ray images), magnetic imaging (magnetic resonance imaging, electron paramagnetic resonance imaging) and radiochemical methods. Typically, the metal salts used in imaging or examining serve as a contrast agent. Therefore in one embodiment the metal complexes or metal salts of compounds of the present invention can be used as contrast agents for example in imaging or examining one or more organs, for example, the gastrointestinal tract. Metals that can serve as contrast agents include gadolinium, iron, manganese, chromium, dysprosium, technetium, scandium, barium, aluminum and holmium, preferably as trications. Radioactive metal salts can be made from isotopes including ^{241}Am , ^{51}Cr , ^{60}Co , ^{57}Co , ^{58}Co , ^{64}Cu , ^{153}Gd , ^{67}Ga , ^{198}Au , $^{113\text{m}}\text{In}$, ^{111}In , ^{59}Fe , ^{55}Fe , ^{197}Hg , ^{203}Hg , $^{99\text{m}}\text{Tc}$, ^{201}Tl , and ^{169}Yb , again preferably when the metal is present as a trivalent cation.

EXAMPLES


[00363] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

Example 1. Synthesis of the Compounds

[00364] Novel DFT analogs were synthesized, such as (*S*)-4,5-dihydro-2-(3,5-dihydroxy-2-pyridinyl)-4-methyl-4-thiazolecarboxylic acid, ((*S*)-5'-(HO)-DFT, **I-1**), (*S*)-4,5-dihydro-2-[3-hydroxy-5-(3,6-dioxaheptyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic acid, ((*S*)-5'-(HO)-DFT-norPE, **I-2**), and (*S*)-4,5-dihydro-2-[3-hydroxy-4-(3,6-dioxaheptyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic acid, ((*S*)-4'-(HO)-DFT-norPE, **I-3**) (the chemical structures shown in *Figure 1B*).


[00365] The preparation of 5'-hydroxydesferrithiocin (**I-1**) and its 5'-nor polyether (**I-2**) began with 2-cyano-3,5-difluoropyridine (**11**), which was converted to 2-cyano-3,5-

dihydroxypyridine (**13**) in two steps (*Scheme 1*). Heating **11** with the anion of 4-methoxybenzyl alcohol (NaH, 2.5 equivalents) in DMF (Féau *et al.*, "Preparation and Optical Properties of Novel 3-Alkoxy carbonyl Aza- and Diazacoumarins." *Synth. Commun.* **2010**, 40, 3033-3045; Ornelas *et al.*, "An Efficient Synthesis of Highly Functionalized Chiral Lactams." *Tetrahedron Lett.* **2011**, 52, 4760-4763) at 95 °C for 18 h gave protected diol **12** in 73% yield. Removal of the 4-methoxybenzyl groups of **12** using excess trifluoroacetic acid (TFA) (White *et al.*, "Total Synthesis of Geodiamolide A, a Novel Cyclodepsipeptide of Marine Origin." *J. Org. Chem.* **1989**, 54, 736-738) and pentamethylbenzene (Marriott *et al.*, "Synthesis of the Farnesyl Ether 2,3,5-Trifluoro-6-hydroxy-4-[(*E,E*)-3,7,11-trimethyldodeca-2,6,10-trien-1-yloxy]nitrobenzene, and Related Compounds Containing a Substituted Hydroxytrifluorophenyl Residue: Novel Inhibitors of Protein Farnesyltransferase, Geranylgeranyltransferase I and Squalene Synthase." *J. Chem. Soc., Perkin Trans.* **2000**, 1, 4265-4278) at room temperature for 22 h provided nitrile **13** in quantitative yield. Cyclocondensation of **13** with (*S*)-2-methyl cysteine (**14**) in aqueous CH₃OH buffered at pH 6 at 75 °C for 45 h followed by esterification of crude acid **I-1** with iodoethane and *N,N*-diisopropylethylamine (DIEA) (1.3 equivalents each) in DMF produced ethyl (*S*)-4,5-dihydro-2-(3,5-dihydroxy-2-pyridinyl)-4-methyl-4-thiazolecarboxylate (**I-1-iii**) in 70% yield. Hydrolysis of **I-1-iii** with aqueous NaOH in CH₃OH at room temperature generated (*S*)-4,5-dihydro-2-(3,5-dihydroxy-2-pyridinyl)-4-methyl-4-thiazolecarboxylic acid (**I-1**) as a solid in 96% yield. Also, ester **I-1-iii** was alkylated at the less hindered phenol (Bergeron *et al.*, "The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs." *J. Med. Chem.* **2010**, 53, 2843-2853) in the presence of the pyridine nitrogen with tosylate **16** and K₂CO₃ in refluxing acetone, affording compound precursor **I-2-iii** in 65% yield. The carboxylate was unmasked under alkaline conditions to give (*S*)-4,5-dihydro-2-[3-hydroxy-5-(3,6-dioxaheptyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic acid (**I-2**) in 97% yield as an oil.

Scheme 1. Exemplary synthesis of **I-1** and **I-2**.

[00366] Synthesis of the 4'-nor polyether desferrithiocin analog **I-3** (Figure 1B), an isomer of **I-2**, started with 2-methyl-3-(benzyloxy)-4-pyridone (**18**), available in two steps from maltol (Piyamongkol *et al.*, "Novel Synthetic Approach to 2-(1'-Hydroxyalkyl)- and 2-Amido-3-Hydroxypyridin-4-ones." *Tetrahedron* **2001**, 57, 3479-3486) (Scheme 2). *O*-Alkylation of **18** with tosylate **16** and K₂CO₃ in refluxing acetonitrile (Li *et al.*, "Synthesis of Coumarin-Appended Pyridyl Tricarbonylrhenium (I) 2,2'-Bipyridyl Complexes with Oligoether Spacer and Their Fluorescence Resonance Energy Transfer Studies." *Organometallics* **2009**, 28, 1620-1630) afforded 2-methyl-3-(benzyloxy)-4-(3,6-

dioxaheptyloxy)pyridine (**19**) in 68% yield, which was oxidized to aldehyde **21** by known methodology (Piyamongkol *et al.*, “Novel Synthetic Approach to 2-(1'-Hydroxyalkyl)- and 2-Amido-3-Hydroxypyridin-4-ones.” *Tetrahedron* **2001**, 57, 3479-3486). Specifically, **19** was treated with 3-chloroperbenzoic acid in CH₂Cl₂, and the resulting *N*-oxide was heated at reflux in acetic anhydride. Cleavage of the acetate ester with base gave the 2-pyridinemethanol **20** in 87% overall yield. Primary alcohol **20** was further oxidized to aldehyde **21** in 83% yield with sulfur trioxide-pyridine complex and NEt₃ in DMSO and CHCl₃. The oxime **22**, generated in 90% yield under standard conditions, was heated at reflux with acetic anhydride, furnishing the corresponding nitrile **23** in 94% yield. Removal of the benzyl-protecting group from **23** by hydrogenolysis (1 atm, 10% Pd-C, CH₃OH) in the presence of the cyano group and pyridyl ring produced 4-(3,6-dioxaheptyloxy)-3-hydroxy-2-pyridinecarbonitrile (**24**) in 81% yield. Heating **24** with amino acid **14** in aqueous CH₃OH buffered at pH 6 generated (*S*)-4,5-dihydro-2-[3-hydroxy-4-(3,6-dioxaheptyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic acid (**I-3**) in 95% yield.



Scheme 2. Exemplary synthesis of I-3.

[00367] Reagents were purchased from Aldrich Chemical Co. (Milwaukee, WI). Compound **11** was obtained from Matrix Scientific (Columbia, SC). Fisher Optima grade solvents were routinely used. Reactions were run under a nitrogen atmosphere, and organic extracts were dried with sodium sulfate. Silica gel 40–63 from SiliCycle, Inc. (Quebec City, Quebec, Canada) was used for column chromatography. Melting points are uncorrected. Glassware that was presoaked in 3 N HCl for 15 min, washed with distilled water and distilled EtOH, and oven-dried was used during the isolation of **I-1**, **I-2**, and **I-3**. Optical rotations were run at 589 nm (sodium D line) and 20 °C on a Perkin-Elmer 341 polarimeter, with *c* being concentration in grams of compound per 100 mL of CHCl₃. ¹H NMR spectra were run in CDCl₃ at 400 MHz (unless otherwise indicated), and chemical shifts (δ) are given in parts per million downfield from tetramethylsilane. ¹³C NMR spectra were measured at 100 MHz (unless otherwise indicated), and chemical shifts (δ) are referenced to the residual

solvent resonance of δ 77.16 for CDCl_3 (not indicated) or δ 39.52 for $\text{DMSO}-d_6$. Coupling constants (J) are in hertz. ESI-FTICR mass spectra are reported. Elemental analyses were performed by Atlantic Microlabs (Norcross, GA) and were within $\pm 0.4\%$ of the calculated values. Purity of the compounds is supported by high pressure liquid chromatography (HPLC) ($\geq 95\%$ for **I-1**, **I-2**, and **I-3**) and by elemental analyses.

[00368] 3,5-Bis(4-methoxybenzyloxy)pyridine-2-carbonitrile (12). Sodium hydride (60%, 3.66 g, 91.5 mmol) was added to 4-methoxybenzyl alcohol (11.5 mL, 92.6 mmol) in DMF (89 mL). The reaction mixture was stirred for 50 min and was cooled in an ice water bath, followed by addition of **11** (5.13 g, 36.6 mmol). After stirring at room temperature for 30 min and heating at 95-100 °C for 18 h, the reaction was quenched at 0 °C with EtOH and was concentrated by rotary evaporation under high vacuum. The residue was treated with H_2O (250 mL) and extracted with warm EtOAc (400 mL, 2×100 mL). The organic extracts were washed with saturated NaCl (150 mL). Purification by flash column chromatography using 2% acetone/ CH_2Cl_2 gave 10.11 g of **12** (73%) as a white solid, mp 122-122.5 °C: ^1H NMR δ 3.81 (s, 3 H), 3.82 (s, 3 H), 5.04 (s, 2 H), 5.11 (s, 2 H), 6.85 (d, 1 H, $J = 2.0$), 6.92 (dd, 4 H, $J = 8.6, 6.6$), 7.31 (d, 4 H, $J = 8.6$), 8.01 (d, 1 H, $J = 2.0$). ^{13}C NMR δ 55.44, 55.47, 70.98, 71.00, 106.86, 114.41, 115.63, 116.26, 126.83, 126.94, 129.04, 129.56, 131.98, 158.41, 159.10, 160.00, 160.14. HRMS m/z calculated for $\text{C}_{22}\text{H}_{21}\text{N}_2\text{O}_4$, 377.1496 ($\text{M} + \text{H}$); found: 377.1500. Anal. ($\text{C}_{22}\text{H}_{20}\text{N}_2\text{O}_4$) C, H, N.

[00369] 3,5-Dihydroxy-2-pyridinecarbonitrile (13). Trifluoroacetic acid (477 g) was added over 26 min to **12** (3.448 g, 41.05 mmol) and pentamethylbenzene (38.22 g, 0.2579 mol) with ice bath cooling. The reaction mixture was stirred at room temperature for 22 h, and volatiles were removed by rotary evaporation. The residue was partitioned between cold 2 N NaOH (180 mL) and Et_2O (350 mL) and separated. The Et_2O layer was back extracted with 0.5 N NaOH (80 mL). The combined aqueous phase was extracted with Et_2O (100 mL), cooled in an ice water bath, and combined with cold 2 M HCl (220 mL) and saturated NaCl (100 mL). The aqueous layer was extracted with EtOAc (250 mL, 2×120 mL). The latter organic extracts were washed with saturated NaCl (150 mL) and concentrated *in vacuo*, giving 3.70 g of **13** (quantitative) as a light tan solid: ^1H NMR ($\text{DMSO}-d_6$) δ 6.80 (d, 1 H, $J = 2.4$), 7.74 (d, 1 H, $J = 2.0$), 10.98 (s, 1 H), 11.39 (s, 1 H). ^{13}C NMR ($\text{DMSO}-d_6$) δ 108.69, 111.17, 116.74, 132.54, 158.10, 159.21. HRMS m/z calculated for $\text{C}_6\text{H}_3\text{N}_2\text{O}_2$, 135.0200 ($\text{M} -$

H); found: 135.0196. An analytical sample was recrystallized from aqueous EtOH. At >300 °C, the sample was dark but not melted. Anal. (C₆H₄N₂O₂) C, H, N.

[00370] Ethyl (S)-4,5-Dihydro-2-(3,5-dihydroxy-2-pyridinyl)-4-methyl-4-thiazole-carboxylate (I-1-iii). A degassed solution of 0.1 M phosphate buffer (pH 6, 310 mL) and CH₃OH (300 mL) was added to **13** (4.04 g, 29.7 mmol) and **14** (6.95 g, 40.5 mmol). The pH of the reaction solution was adjusted to 6.0 with NaHCO₃ (4.92 g, 58.6 mmol). The reaction mixture was heated at 73-76 °C for 45 h with stirring, cooled to 0 °C, and reduced in volume by rotary evaporation. The residue was acidified to pH ~ 1 with cold 2 N HCl (61 mL) followed by extraction with EtOAc (300 mL, 2 × 100 mL). The organic layer was washed with saturated NaCl (100 mL), concentrated *in vacuo* and dried with toluene, resulting in 6.30 g of **I-1**. Iodoethane (3.0 mL, 37.5 mmol) and DIEA (6.5 mL, 37.3 mmol) were successively added to **I-1** in DMF (130 mL), and the solution was stirred at room temperature for 47 h. After solvent removal under high vacuum, the residue was treated with 12:5 0.5 M HCl/saturated NaCl (170 mL) followed by extraction with EtOAc (150 mL, 4 × 70 mL). The EtOAc layers were washed with 100 mL portions of 1% NaHSO₃ and saturated NaCl, and the solvent was evaporated. Purification by column chromatography using (5% acetone/CH₂Cl₂) gave 5.88 g of **I-1-iii** (70%) as a pale yellow solid, mp 85-87.5 °C: [α] +35.6° (*c* 0.74). ¹H NMR δ 1.32 (t, 3 H, *J* = 7.2), 1.69 (s, 3 H), 3.20 (d, 1 H, *J* = 11.7), 3.79 (d, 1 H, *J* = 11.7), 4.27 (q, 2 H, *J* = 7.2), 6.77 (d, 1 H, *J* = 2.4), 7.82 (d, 1 H, *J* = 2.3). ¹³C NMR δ 14.23, 24.78, 39.60, 62.33, 83.67, 110.11, 127.71, 130.80, 156.17, 157.79, 173.21, 174.02. HRMS *m/z* calculated for C₁₂H₁₅N₂O₄S, 283.0747 (*M* + *H*), 305.0567 (*M* + *Na*); found: 283.0751, 305.0573. Anal. (C₁₂H₁₄N₂O₄S) C, H, N.

[00371] (S)-4,5-Dihydro-2-(3,5-dihydroxy-2-pyridinyl)-4-methyl-4-thiazole-carboxylic Acid (I-1). A solution of 50% (w/w) NaOH (13.7 g, 0.171 mol) in CH₃OH (135 mL) was added to **I-1-iii** (4.85 g, 17.2 mmol) in CH₃OH (125 mL) over 13 min at 0 °C. The reaction mixture was warmed to room temperature over 19 h, and the bulk of the solvent was removed by rotary evaporation. The concentrate was treated with dilute NaCl (150 mL) and was extracted with Et₂O (2 × 100 mL). The aqueous layer was cooled in ice, acidified with cold 6 N HCl (30 mL), and extracted with EtOAc (250 mL, 2 × 100 mL). The EtOAc extracts were washed with saturated NaCl (80 mL). Solvent was removed *in vacuo*, providing 4.18 g of **I-1** (96%) as an off white solid, mp 226-227 °C (decomposed): [α] +46.0° (*c* 0.82, DMF). ¹H NMR (DMSO-*d*₆) δ 1.58 (s, 3 H), 3.27 (d, 1 H, *J* = 11.7), 3.69 (d, 1 H, *J* = 11.7), 6.72 (d,

1 H, $J = 2.4$), 7.80 (d, 1 H, $J = 2.0$), 10.82 (s, 1 H), 12.32 (s, 1 H), 13.20 (s, 1 H). ^{13}C NMR (DMSO- d_6) δ 24.32, 38.48, 82.98, 108.59, 125.88, 131.13, 156.68, 157.58, 173.30, 173.74. HRMS m/z calculated for $\text{C}_{10}\text{H}_{11}\text{N}_2\text{O}_4\text{S}$, 255.0434 ($M + H$), 277.0253 ($M + \text{Na}$), 299.0073 ($M - H + 2\text{Na}$), 320.9892 ($M - 2H + 3\text{Na}$); found: 255.0439, 277.0255, 299.0077, 320.9899. Anal. ($\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}_4\text{S}$) C, H, N.

[00372] Ethyl (S)-4,5-Dihydro-2-[3-hydroxy-5-(3,6-dioxahexyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylate (I-2-iii). Flame activated K_2CO_3 (0.72 g, 5.25 mmol) was added to a mixture of **16** (0.96 g, 3.5 mmol) and **I-1-iii** (0.90 g, 3.19 mmol) in dry acetone (25 mL). The reaction mixture was heated at reflux for 24 h. After cooling to room temperature the solvent was removed by rotary evaporation. The residue was treated with 0.2 N HCl/saturated NaCl (50 mL) and was extracted with EtOAc (4×30 mL). The organic extracts were washed with saturated NaCl (50 mL) and solvent was removed *in vacuo*. Column chromatography using 1:2:7 CH_3OH /hexane/ CH_2Cl_2 furnished 0.80 g of **I-2-iii** (65%) as a viscous oil: $[\alpha] +30.9^\circ$ (c 1.12). ^1H NMR δ 1.30 (t, 3 H, $J = 7.0$), 1.67 (s, 3 H), 3.19 (d, 1 H, $J = 11.3$), 3.40 (s, 3 H), 3.56-3.62 (m, 2 H), 3.70-3.75 (m, 2 H), 3.80 (d, 1 H, $J = 11.7$), 3.86-3.93 (m, 2 H), 4.19 (t, 2 H, $J = 4.7$), 4.25 (q, 2 H, $J = 7.0$), 6.80 (d, 1 H, $J = 2.3$), 7.95 (d, 1 H, $J = 2.3$), 12.37 (s, 1 H). ^{13}C NMR δ 14.23, 24.77, 39.45, 59.25, 62.04, 68.11, 69.46, 70.99, 72.01, 83.84, 107.63, 127.72, 131.49, 157.39, 158.22, 172.87, 173.96. HRMS m/z calculated for $\text{C}_{17}\text{H}_{25}\text{N}_2\text{O}_6\text{S}$, 385.1428 ($M + H$), 407.1247 ($M + \text{Na}$); found: 385.1432, 407.1266. Anal. ($\text{C}_{17}\text{H}_{24}\text{N}_2\text{O}_6\text{S}$) C, H, N.

[00373] (S)-4,5-Dihydro-2-[3-hydroxy-5-(3,6-dioxahexyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic Acid (I-2). A solution of 50% (w/w) NaOH (1.46 mL, 47.0 mmol) in CH_3OH (40 mL) was added dropwise to a solution of **I-2-iii** (1.66 g, 4.31 mmol) in CH_3OH (20 mL) at 0 °C. The reaction mixture was stirred at room temperature for 6 h, and the bulk of the solvent was removed under reduced pressure. The residue was treated with dilute NaCl (50 mL) and was extracted with Et_2O (2×30 mL). The aqueous layer was cooled in ice, acidified with 2 N HCl to pH = 2, and extracted with EtOAc (5×40 mL). Combined EtOAc layers were washed with saturated NaCl (60 mL). Solvent removal *in vacuo* furnished 1.49 g of **I-2** (97%) as a yellow oil: $[\alpha] +25.3^\circ$ (c 0.88). ^1H NMR δ 1.73 (s, 3 H), 3.22 (d, 1 H, $J = 12.0$), 3.41 (s, 3 H), 3.59-3.61 (m, 2 H), 3.72-3.74 (m, 2 H), 3.83 (d, 1 H, $J = 11.6$), 3.88 (t, 2 H, $J = 4.8$), 4.19 (t, 2 H, $J = 4.4$), 6.84 (d, 1 H, $J = 2.4$), 7.94 (d, 1 H, $J = 2.4$). ^{13}C NMR δ 24.62, 39.13, 58.98, 67.99, 69.28, 70.63, 71.81, 82.83, 107.67, 126.98, 131.64, 158.15,

158.44, 174.59, 175.94. HRMS m/z calculated for $C_{15}H_{21}N_2O_6S$, 357.1115 ($M + H$); found: 357.1125. Anal. ($C_{15}H_{20}N_2O_6S$) C, H, N.

[00374] 2-Methyl-3-(benzyloxy)-4-(3,6-dioxaheptyloxy)pyridine (19). Flame activated K_2CO_3 (27.6 g, 0.20 mol) and **16** (27.4 g, 0.10 mol) were added to **18** (21.5 g, 0.10 mol) in dry CH_3CN (500 mL). The reaction mixture was heated at reflux for 24 h. After cooling to room temperature, the solvent was evaporated by rotary evaporation. The residue was treated with 10% NaCl (200 mL) and was extracted with CH_2Cl_2 (4×150 mL). The organic extracts were washed with saturated NaCl (300 mL). After solvent was removed *in vacuo*, column chromatography using 4:4:2 EtOAc/petroleum ether/acetone furnished 21.5 g of **19** (68%) as a colorless viscous oil: ^1H NMR δ 2.42 (s, 3 H), 3.34 (s, 3 H), 3.51-3.53 (m, 2 H), 3.69-3.71 (m, 2 H), 3.91 (t, 2 H, $J = 4.8$), 4.24 (t, 2 H, $J = 4.4$), 5.02 (s, 2 H), 6.72 (d, 1 H, $J = 5.6$), 7.31-7.40 (m, 3 H), 7.44-7.49 (m, 2 H), 8.12 (d, 1 H, $J = 5.6$). ^{13}C NMR δ 19.34, 59.16, 67.86, 69.45, 70.92, 71.99, 74.57, 106.68, 128.21, 128.45, 128.49, 137.53, 142.32, 145.41, 153.40, 157.64. HRMS m/z calculated for $C_{18}H_{24}NO_4$, 318.1700 ($M + H$); found: 318.1714. Anal. ($C_{18}H_{23}NO_4 \cdot 0.2 H_2O$) C, H, N.

[00375] 4-(3,6-Dioxaheptyloxy)-3-(benzyloxy)-2-pyridinemethanol (20). An ice cooled solution of 3-chloroperoxybenzoic acid (3.67 g, 36.0 mmol) in CH_2Cl_2 (75 mL) was added slowly to **19** (10.4 g, 32.8 mmol) in CH_2Cl_2 (50 mL) over 15 min at 0 °C. The reaction mixture was warmed to room temperature, stirred for 6 h, and diluted with CH_2Cl_2 (150 mL). The reaction mixture was washed with 5% Na_2CO_3 (3×100 mL) and saturated NaCl (100 mL) and was concentrated under reduced pressure to give a colorless oil. Acetic anhydride (80 mL, 0.85 mol) was added, and the reaction mixture was heated at 130 °C for 2 h. The solvent was removed under reduced pressure, and the residue was dissolved in H_2O (100 mL). The pH of the aqueous solution was adjusted to 8 with 2 N sodium hydroxide, and the aqueous solution was extracted with CH_2Cl_2 (3×100 mL). The organic fractions were combined, washed with saturated NaCl (100 mL), and concentrated *in vacuo*. The residue was dissolved in CH_3OH , treated with decolorizing charcoal, filtered, and concentrated to yield a brown oil, which was dissolved in EtOH (40 mL). Sodium hydroxide (1 M, 80 mL) was added and the reaction mixture was refluxed for 4 h and cooled. Extraction with CH_2Cl_2 (4×100 mL), washing with saturated NaCl (100 mL), concentration under reduced pressure, and column chromatography using 10% $CH_3OH/CHCl_3$ provided 9.52 g (87%) of **20** as a light brown oil: ^1H NMR δ 3.34 (s, 3 H), 3.52-3.54 (m, 2 H), 3.69-3.71 (m, 2 H), 3.92 (t, 2

H, $J = 5.2$), 4.28 (t, 2 H, $J = 4.4$), 4.65 (s, 2 H), 5.09 (s, 2 H), 7.32-7.39 (m, 3 H), 7.40-7.44 (m, 2 H), 8.19 (d, 1 H, $J = 5.6$). ^{13}C NMR δ 59.20, 60.23, 68.13, 69.44, 70.96, 72.04, 74.79, 107.83, 128.49, 128.54, 128.63, 137.15, 140.53, 144.66, 152.98, 157.52. HRMS m/z calculated for $\text{C}_{18}\text{H}_{24}\text{NO}_5$, 334.1649 ($M + \text{H}$), 356.1468 ($M + \text{Na}$); found: 334.1648, 356.1455. Anal. ($\text{C}_{18}\text{H}_{23}\text{NO}_5$) C, H, N.

[00376] 4-(3,6-Dioxaheptyloxy)-3-(benzyloxy)pyridine-2-carboxaldehyde (21). Triethylamine (70 mL, 0.29 mol) followed by DMSO (70 mL) was added to **20** (16.5 g, 49.0 mmol) in CHCl_3 (100 mL). Sulfur trioxide-pyridine complex (35 g, 0.22 mol) was slowly added over 35 min to the reaction mixture with ice bath cooling. After warming to room temperature, the reaction mixture was stirred overnight and was diluted with CHCl_3 (200 mL). The organic phase was washed with H_2O (3×200 mL) and saturated NaCl (100 mL). After solvent was removed *in vacuo*, column chromatography using 5:5:1 EtOAc/ CHCl_3 / CH_3OH furnished 13.61 g of **21** (83%) as a viscous colorless oil: ^1H NMR δ 3.34 (s, 3 H), 3.52-3.54 (m, 2 H), 3.70-3.72 (m, 2 H), 3.95 (t, 2 H, $J = 4.4$), 4.30 (t, 2 H, $J = 4.4$), 5.24 (s, 2 H), 7.02 (d, 1 H, $J = 5.2$), 7.32-7.39 (m, 3 H), 7.41-7.46 (m, 2 H), 8.39 (d, 1 H, $J = 5.6$), 10.25 (s, 1 H). ^{13}C NMR δ 59.17, 68.54, 69.23, 70.94, 71.97, 76.24, 111.69, 128.68, 128.75, 128.87, 136.16, 145.87, 146.90, 148.14, 159.49, 189.87. HRMS m/z calculated for $\text{C}_{18}\text{H}_{21}\text{NNaO}_5$, 354.1312 ($M + \text{Na}$); found: 354.1326. Anal. ($\text{C}_{18}\text{H}_{21}\text{NO}_5$) C, H, N.

[00377] 4-(3,6-Dioxaheptyloxy)-3-(benzyloxy)pyridine-2-carboxaldehyde oxime (22). Hydroxylamine hydrochloride (4.2 g, 60.0 mmol) and NaOAc (5.2 g, 60.0 mmol) were added to a solution of **21** (13.5 g, 40.7 mmol) in CH_3OH (50 mL), and the reaction mixture was heated at reflux for 2 h. The reaction mixture was concentrated by rotary evaporation, and the residue was treated with saturated NaCl (100 mL) and 0.1 M citric acid (100 mL) and then was extracted with EtOAc (2×100 mL). The organic layers were washed with H_2O (100 mL) and saturated NaCl (100 mL). Solvent was removed *in vacuo*, providing 12.7 g (90%) of **22** as a pale solid, mp 72-73 °C: ^1H NMR δ 3.34 (s, 3 H), 3.46-3.51 (m, 2 H), 3.64-3.71 (m, 2 H), 3.92 (t, 2 H, $J = 4.4$), 4.27 (t, 2 H, $J = 4.4$), 5.10 (s, 2 H), 6.85 (d, 1 H, $J = 5.6$), 7.29-7.46 (m, 5 H), 8.28 (d, 1 H, $J = 5.2$), 8.46 (s, 1 H). ^{13}C NMR δ 59.14, 68.12, 69.28, 70.86, 71.94, 75.67, 108.62, 128.42, 128.55, 128.59, 136.70, 143.43, 144.81, 145.23, 146.71, 158.66. HRMS m/z calculated for $\text{C}_{18}\text{H}_{21}\text{NaNO}_5$, 354.1312 ($M + \text{Na}$); found: 352.1326. Anal. ($\text{C}_{18}\text{H}_{21}\text{NO}_5$) C, H, N.

[00378] 4-(3,6-Dioxaheptyloxy)-3-(benzyloxy)pyridine-2-carbonitrile (23).

Compound **22** was dissolved in Ac₂O (40 mL) and heated at reflux for 8 h under a Drierite tube. The reaction mixture was concentrated by rotary evaporation, dissolved in 8% NaHCO₃ (100 mL), and extracted with CHCl₃ (100 mL, 2 × 50 mL). Combined organic fractions were washed with 4% NaHCO₃ (50 mL) and saturated NaCl (100 mL) followed by solvent removal *in vacuo*. Purification by flash chromatography eluting with 10% CH₃OH/CH₂Cl₂ gave 11.31 g (94%) of **23** as a pale solid, mp 34-35 °C: ¹H NMR δ 3.34 (s, 3 H), 3.53-3.55 (m, 2 H), 3.70-3.72 (m, 2 H), 3.93 (t, 2 H, *J* = 4.8), 4.27 (t, 2 H, *J* = 4.4), 5.31 (s, 2 H), 6.98 (d, 1 H, *J* = 5.6), 7.31-7.38 (m, 3 H), 7.49-7.52 (m, 2 H), 8.21 (d, 1 H, *J* = 5.2). ¹³C NMR δ 59.14, 68.59, 69.09, 70.92, 71.94, 75.84, 111.27, 115.43, 128.60, 128.66, 128.73, 128.84, 135.78, 147.21, 148.28, 158.42. HRMS *m/z* calculated for C₁₈H₂₀N₂NaO₄, 351.1315 (M + Na); found: 351.1325. Anal. (C₁₈H₂₀N₂O₄) C, H, N.

[00379] 4-(3,6-Dioxaheptyloxy)-3-hydroxy-2-pyridinecarbonitrile (24). Palladium on carbon (10%, 0.065 g) was added to a solution of **23** (1.3 g 3.95 mmol) in CH₃OH (15 mL), and the mixture was stirred under H₂ at atmospheric pressure for 2 h. The reaction mixture was filtered through Celite[®], and the residue was washed with CH₃OH (3 × 5 mL). The filtrate was concentrated under reduced pressure, and the residue was subjected to column chromatography eluting with 10% CH₃OH/EtOAc furnishing 0.80 g (85%) of **24** as a colorless oil: ¹H NMR δ 3.42 (s, 3 H), 3.61-3.63 (m, 2 H), 3.75-3.77 (m, 2 H), 3.92 (t, 2 H, *J* = 4.8), 4.24 (t, 2 H, *J* = 4.4), 6.91 (d, 1 H, *J* = 4.8), 8.10 (d, 1 H, *J* = 5.6). ¹³C NMR δ 58.92, 68.66, 69.05, 70.52, 71.78, 110.68, 115.28, 120.29, 143.49, 148.84, 153.72. HRMS *m/z* calculated for C₁₁H₁₄N₂NaO₄, 261.0846 (M + Na); found: 261.0849. Anal. (C₁₁H₁₄N₂O₄) C, H, N.

[00380] (S)-4,5-Dihydro-2-[3-hydroxy-4-(3,6-dioxaheptyloxy)-2-pyridinyl]-4-methyl-4-thiazolecarboxylic Acid (I-3). Compound **14** (0.78 g, 4.58 mmol), pH 6 phosphate buffer (30 mL), and NaHCO₃ (0.44 g, 5.23 mmol) were successively added to a solution of **24** (0.78 g, 3.27 mmol) in degassed CH₃OH (30 mL). The reaction mixture was heated at 75 °C for 48 h with stirring, cooled to room temperature, and concentrated by rotary evaporation. The residue was dissolved in distilled H₂O (25 mL) and the aqueous layer was acidified with cold 2 N HCl to pH <2 followed by extraction with EtOAc (5 × 50 mL). Concentration *in vacuo* resulted in 1.15 g of **I-3** (95%) as a light yellow oil: [α]_D²⁰ +52.8° (*c* 0.40). ¹H NMR δ 1.73 (s, 3 H), 3.24 (d, 1 H, *J* = 11.6), 3.39 (s, 3 H), 3.56-3.58 (m, 2 H),

3.73-3.75 (m, 2 H), 3.85 (d, 1 H, $J = 11.6$), 3.94 (t, 2 H, $J = 4.8$), 4.27 (t, 2 H, $J = 4.8$), 6.88 (d, 1 H, $J = 5.2$), 8.08 (d, 1 H, $J = 4.8$). ^{13}C NMR δ 24.65, 39.52, 59.08, 68.57, 69.31, 70.87, 71.94, 83.67, 109.81, 133.20, 141.56, 147.16, 154.39, 175.11, 176.26. HRMS m/z calculated for $\text{C}_{15}\text{H}_{21}\text{N}_2\text{O}_6\text{S}$, 357.1115 ($\text{M} + \text{H}$); found: 357.1115. Anal. ($\text{C}_{15}\text{H}_{20}\text{N}_2\text{O}_6\text{S}$) C, H, N.

Example 2. Stoichiometry of the Complexes of Fe(III) and the Compounds

[00381] Earlier studies with **1** by Anderegg and Räber showed the chelator to form a 2:1 complex with Fe(III) (Anderegg *et al.*, “Metal Complex Formation of a New Siderophore Desferrithiocin and of Three Related Ligands.” *J. Chem. Soc., Chem. Commun.* **1990**, 1194-1196). The cumulative formation constant for this complex was determined to be $4 \times 10^{29} \text{ M}^{-1}$. Hahn *et al.* were ultimately able to isolate both the Δ and λ 1-Cr(III) complexes, with chromium serving as a surrogate for Fe(III) (Hahn *et al.*, “Coordination Chemistry of Microbial Iron Transport. 42. Structural and Spectroscopic Characterization of Diastereomeric Cr(III) and Co(III) Complexes of Desferriferrithiocin.” *J. Am. Chem. Soc.* **1990**, *112*, 1854-1860). As expected, the crystal structures of the complexes unequivocally demonstrated a 2:1 ligand to metal ratio. In later studies, Job’s plots with **3** (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogues.” *J. Med. Chem.* **1999**, *42*, 2432-2440) and the corresponding desmethyl analog (Bergeron *et al.*, “Desazadesmethyldesferrithiocin Analogues as Orally Effective Iron Chelators.” *J. Med. Chem.* **1999**, *42*, 95-108) also showed that these ligands formed 2:1 complexes with Fe(III). This is in keeping with the fact that the donor groups of the chelators, the aromatic hydroxyl, the thiazoline nitrogen, and the carboxylate are the same as in **1** itself. Furthermore, a comparison of structure **3** (Bergeron *et al.*, “Iron Chelation Promoted by Desazadesferrithiocin Analogs: An Enantioselective Barrier.” *Chirality* **2003**, *15*, 593-599) with **1** reveals that the disposition of the coordinating sites is essentially the same.









[00382] The stoichiometries of the complexes of Fe(III) and DFT analogs **I-1**, **I-2**, and **I-3** were determined spectrophotometrically using Job’s plots (*Figure 2*). In each instance, the DFT analogs formed 2:1 complexes with Fe(III). Solutions were monitored at the visible λ_{max} of the Fe(III) complexes (498 nm for **I-1**, 484 nm for **I-2**, and 485 nm for **I-3**). A 100 mM Tris HCl buffer was used to maintain the pH at 7.4. Solutions containing different DFT analog/Fe(III) ratios were prepared by mixing appropriate volumes of 1.0 mM DFT analog solution and 1.0 mM Fe(III)-nitriloacetate (NTA) in Tris-HCl buffer. The 1.0 mM Fe(III)-

NTA solution was prepared immediately prior to use by dilution of a 41.6 mM Fe(III)-NTA stock solution with the Tris HCl buffer, whereas the DFT analog stock solution was prepared by dissolving the DFT analog as its monosodium salt in Tris HCl buffer at pH 7.4. The Fe(III)-NTA stock solution was prepared by mixing equal volumes of 90 mM of FeCl₃ and 180 mM trisodium NTA. The iron content of the Fe(III)-NTA solution was verified by atomic absorption spectrophotometry (AAS).

Example 3. Partition Properties of the Compounds

[00383] The partition values between octanol and water (at pH 7.4, Tris buffer) were determined using a “shake flask” direct method of measuring log P_{app} values (Sangster *et al.*, *Octanol-Water Partition Coefficients: Fundamentals and Physical Chemistry*; John Wiley and Sons: West Sussex, England, 1997; Vol. 2). The fraction of compound in the octanol is then expressed as log P_{app} . While the values vary widely (*Table 1*), one observation stands out: DFT and its analogs are always more hydrophilic than their DADFT counterparts, *i.e.*, **1** vs. **2**, **I-1** vs. **3**, **I-2** vs. **7**, and **I-3** vs. **9**. This is likely due to the presence of the aromatic nitrogen, a moderately good hydrogen bond acceptor, on the DFT analogs. Relative to the differences in lipophilicity between DFT and DADFT, fixing a polyether backbone to either the DFT or the DADFT pharmacophore had a much more moderate effect (*Table 1*).

Table 1. Iron-Clearing Efficiency of Desferrithiocin Analogs Administered to Rodents and Primates with the Respective LogP_{app} values.

Compound Structure	Compound No.	Rodent Iron-Clearing Efficiency ^a (%)	Primate Iron-Clearing Efficiency ^c (%)	LogP _{app}	PR ^d
	1	5.5 ± 3.2 [93/7]	16.1 ± 8.5 [78/22]	-1.77	2.9
	2	2.7 ± 0.5 [100/0]	21.5 ± 12.0 [76/24]	-0.34	8.0
	3	1.1 ± 0.8 [100/0]	16.8 ± 7.2 [88/12]	-1.05	15.3
	I-1	9.0 ± 3.8 [97/3]	10.0 ± 2.9 [58/42]	-1.68	1.1
	7	26.7 ± 4.7 ^b [97/3]	26.3 ± 9.9 [93/7] (capsule) 28.7 ± 12.4 [83/17] (sodium salt)	-0.89	1.0 1.1
	I-2	11.7 ± 1.2 [97/3]	18 ± 5.2 [63/37]	-1.59	1.5
	9	15.1 ± 2.0 ^b [99/1]	22.5 ± 6.4 [86/14]	-0.96	1.5
	I-3	14.2 ± 2.4 [98/2]	6.1 ± 1.8 (po) [40/60] 16.9 ± 7.3 (sc) [64/36]	-1.38	0.4 1.2

^a In the rodents [$n = 3$ (**7**), 4 (**2**, **I-3**), 5 (**1**, **I-1**, **I-2**, **9**), or 8 (**3**)], the compounds were given po at a dose of 150 $\mu\text{mol/kg}$ (**1–2**) or 300 $\mu\text{mol/kg}$ (**3**, **I-1**, **7**, **I-2**, **9**, and **I-3**). The compounds were administered in capsules (**7**), solubilized in 40% Cremophor RH-40/water (**1**, **2**), or were given as their monosodium salts, prepared by the addition of 1 equiv. of NaOH to a suspension of the free acid in distilled water (**3**, **I-1**, **I-2**, **9**, and **I-3**). The efficiency of each compound was calculated by subtracting the 24 or 48-h iron excretion of control animals from the iron excretion of the treated animals. The number was then divided by the

theoretical output; the result is expressed as a percent. The relative percentages of the iron excreted in the bile and urine are in brackets. The iron-clearing efficiency (ICE) data for: **1** is from Bergeron, R. J.; Wiegand, J.; Dionis, J. B.; Egli-Karmakka, M.; Frei, J.; Huxley-Tencer, A.; Peter, H. H. Evaluation of Desferrithiocin and Its Synthetic Analogs as Orally Effective Iron Chelators. *J. Med. Chem.* **1991**, *34*, 2072-2078; **2** is from Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, *81*, 2166-2173; **3** is from Bergeron *et al.*, “Iron Chelators and Therapeutic Uses.” In: Abraham, ed. *Burger's Medicinal Chemistry*. 6th. Wiley; New York: **2003**. pp. 479-561; **7** is from Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, *53*, 2843-2853; and **9** is from Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, *24*, 239-258.

^b ICE is based on a 48-h sample collection period.

^c In the primates [$n = 4$ (**1**, **2**, **I-1**, **7** in capsules, **I-2**, **9**, and **I-3**), or 6 (**3**), or 7 (**7** as the monosodium salt)], the compounds were given po at a dose of 75 $\mu\text{mol/kg}$ (**2**, **I-1**, **7**, **I-2**, **9**, and **I-3**) or 150 $\mu\text{mol/kg}$ (**1**, **3**). Compound **I-3** was also given to the primates sc at a dose of 75 $\mu\text{mol/kg}$. The compounds were administered in capsules (**7**), solubilized in 40% Cremophor RH-40/water (**1**, **2**), or were given as their monosodium salts, prepared by the addition of 1 equiv of NaOH to a suspension of the free acid in distilled water (**1**, **2**, **3**, **I-1**, **7**, **I-2**, **9**, and **I-3**). The efficiency was calculated by averaging the iron output for 4 days before the compound, subtracting these numbers from the 2-day iron clearance after the administration of the compound, and then dividing by the theoretical output; the result is expressed as a percent. The ICE data for: **1–3** are from Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, *42*, 2432-2440; **7** is from Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, *53*, 2843-2853; and **9** is from Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, *24*, 239-258. The relative percentages of the iron excreted in the feces and urine are in brackets.

^d Performance ratio (PR) is defined as the mean $\text{ICE}_{\text{primates}}/\text{ICE}_{\text{rodents}}$.

Example 4. Biological Assays of the Compounds

[00384] All animal experimental treatment protocols were reviewed and approved by the University of Florida's Institutional Animal Care and Use Committee.

[00385] Male Sprague-Dawley rats were procured from Harlan Sprague-Dawley (Indianapolis, IN). Male *Cebus apella* monkeys (3.5-4 kg) were obtained from World Wide Primates (Miami, FL). Ultrapure salts were obtained from Johnson Matthey Electronics (Royston, UK). All hematological and biochemical studies were performed by Antech Diagnostics (Tampa, FL). Atomic absorption (AA) measurements were made on a Perkin-Elmer model 5100 PC (Norwalk, CT). An R-Rena-strip Lateral-flow Kit for the detection of kidney injury molecule-1 (Kim-1) in rat urine was obtained from BioAssay Works (Ijamsville, MD). A Chromatoreader ReaScan (Otsuka Electronics Co., Japan) was utilized to read the test strips and to allow for the quantitation of Kim-1 in rat urine.

¹H NMR Hydrogen-Deuterium Exchange Study

[00386] Compounds **3**, **I-1**, and **I-2** were dissolved in 0.1 M pD 7.0 phosphate buffer at a concentration of 5.4 mM: **3** (1.1 mg, 4.34 μ mol) and **I-1** (1.1 mg, 4.33 μ mol) in buffer (0.80 mL) and **I-2** (1.8 mg, 5.05 μ mol) in buffer (0.933 mL). The ratio of the [**3**] (unexchanged) to the [**3**] (original) was measured by the diminution of δ 6.33 (d, H-3', $J = 2.3$) relative to the integration of δ 6.39 (d, H-5', $J = 9.0$) at time points to 104 min. The ratio of the [**I-1**] (unexchanged) to the [**I-1**] (original) was measured by the diminution of δ 6.41 (d, H-4', $J = 2.3$) relative to the integration of δ 7.54 (d, H-6', $J = 2.3$) at time points to 16 h. Plots of the natural log of [compound] (unexchanged) versus time are linear, with first order rate constants for **3** and **I-1** (Table 2). No change in the 1:1 ratio of δ 6.76 (d, H-4', $J = 2.0$) to δ 7.72 (d, H-6', $J = 2.3$) in **I-2** was observed up to 17 h.

Table 2. Hydrogen-Deuterium Exchange Rates of Compounds **3**, **I-1**, and **I-2**.

Compound No.	Exchange Rate ^a (s ⁻¹)
3	5.5×10^{-4}
I-1	3.3×10^{-6}
I-2	No exchange

^a The exchange study carried out at 25 °C, pH 7.0, in phosphate buffer (0.1 M).

Cannulation of Bile Duct in Non-Iron-Overloaded Rats

[00387] The cannulation has been described previously (Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, 81, 2166-2173; Bergeron *et al.*, “A Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393). Bile samples were collected from male Sprague-Dawley rats (400-450 g) at 3 h intervals for up to 48 h. The urine sample(s) was taken at 24 h intervals. Sample collection and handling are as previously described (Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, 81, 2166-2173; Bergeron *et al.*, “A Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393).

Iron Loading of C. apella Monkeys

[00388] The monkeys were iron overloaded with intravenous iron dextran as specified in earlier publications (Bergeron *et al.*, “A Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393; Bergeron *et al.*, “A Comparison of the Iron-Clearing Properties of 1,2-Dimethyl-3-Hydroxypyrid-4-one, 1,2-Diethyl-3-Hydroxypyrid-4-one, and Deferoxamine.” *Blood* **1992**, 79, 1882-1890) to provide about 500 mg of iron per kg of body weight; the serum transferrin iron saturation rose to between 70 and 80%. At least 20 half-lives, 60 days (Wood *et al.*, “The Metabolism of Iron-Dextran Given As a Total-Dose Infusion to Iron Deficient Jamaican Subjects.” *Br. J. Haematol.* **1968**, 14, 119-129), elapsed before any of the animals were used in experiments evaluating iron-chelating agents.

Primate Fecal and Urine Samples

[00389] Fecal and urine samples were collected at 24 h intervals and processed as described previously (Bergeron *et al.*, “A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model.” *Blood* **1993**, 81, 2166-2173; Bergeron *et al.*, “A Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393; Bergeron *et al.*, “HBED: A Potential Alternative to Deferoxamine for Iron-Chelating Therapy.” *Blood* **1998**, 91, 1446-1452). Briefly, the collections began 4 days prior to the administration of the test compound and continued for an additional 5 days after the compound was given. Iron concentrations were determined by flame absorption spectroscopy as presented in other publications (Bergeron *et al.*, “A

Comparative Evaluation of Iron Clearance Models.” *Ann. N.Y. Acad. Sci.* **1990**, 612, 378-393; Bergeron *et al.*, “Synthesis and Biological Evaluation of Naphthyl-desferrithiocin Iron Chelators.” *J. Med. Chem.* **1996**, 39, 1575-1581).

Compound Preparation and Administration: Iron Clearance

[00390] In the iron clearing experiments, the rats were given **I-1**, **I-2** and **I-3** po at a dose of 300 $\mu\text{mol/kg}$. The primates were given **I-1**, **I-2** and **I-3** po at a dose of 75 $\mu\text{mol/kg}$; compound **I-3** was also given sc at a dose of 75 $\mu\text{mol/kg}$. The compounds were administered to the rats and primates as their monosodium salts (prepared by the addition of 1 equiv of NaOH to a suspension of the free acid in distilled water). Compound preparation for the rodent urinary Kim-1 excretion studies involving **1**, **I-1**, **I-2**, and **I-3** are described below.

Calculation of Iron-Clearing Efficiency

[00391] In the text below, the term “iron-clearing efficiency” (ICE) is used as a measure of the amount of iron excretion induced by a chelator. The ICE, expressed as a percent, is calculated as (compound-induced iron excretion/theoretical iron excretion) $\times$ 100. To illustrate, the theoretical iron excretion after administration of one millimole of desferrioxamine B mesylate (DFO) (*Figure 3*), a hexadentate chelator that forms a 1:1 complex with Fe(III), is one milli-g-atom of iron. Two millimoles of desferrithiocin (DFT) (*Table 1*), a tridentate iron chelator that forms a 2:1 complex with Fe(III), are required for the theoretical excretion of one milli-g-atom of iron. The theoretical iron outputs of the chelators were generated on the basis of a 2:1 compound:iron complex. The efficiencies in the rats and monkeys were calculated as set forth elsewhere (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “HBED: A Potential Alternative to Deferoxamine for Iron-Chelating Therapy.” *Blood* **1998**, 91, 1446-1452). Data are presented as the mean $\pm$ the standard error of the mean; *p*-values were generated via a one-tailed Student’s *t*-test in which the inequality of variances was assumed; and a *p*-value of < 0.05 was considered significant.

Compound Preparation and Administration: Rodent Toxicity/ Urinary Kim-1 Excretion Studies

[00392] The impact of compounds **1**, **I-1**, **I-2** and **I-3** on urinary Kim-1 excretion were evaluated in rodents. The compounds were administered to the rats po as their monosodium salts, prepared as described above, twice daily at a dose of 237 $\mu\text{mol/kg/dose}$ (474 $\mu\text{mol/kg/d}$) for up to 7 d. The studies were performed on rats with normal iron stores. The rats were fasted overnight and were given the first dose of the compound first thing in the morning. The rats were fed ~ 3 h post-dose and had access to food for ~ 5 h before being fasted overnight.

Collection of Urine for Kim-1 Studies

[00393] The rats were housed in individual metabolic cages. Urine samples were collected from the metabolic cages at 24 h intervals. A baseline (day 0) urine sample was collected and assessed for its Kim-1 content; each animal served as its own control. The urine was collected chilled as previously described (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258).

Performance of Urinary Kim-1 Studies

[00394] The chilled urine was collected, vortexed, and warmed to room temperature; any sediment in the samples was allowed to settle. Kim-1 content was assessed using a Rat Kim-1 Rapid Test Kit according to the manufacturer’s instructions. The result was read using a ReaScan Test Reader. The quantity of Kim-1 excreted in the urine per day was calculated by multiplying the concentration of Kim-1 (ng/ml urine) $\times$ 24-h urine volume, divided by the weight of the animal. The result is expressed as urinary Kim-1 (ng/kg/24 h). Data are presented as the mean $\pm$ the standard error of the mean; *p*-values were generated via a one-tailed Student’s *t*-test in which the inequality of variances was assumed; and a *p*-value of < 0.05 was considered significant.

Results

[00395] A measure of the amount of iron excretion induced by a chelator is best described by its iron-clearing efficiency (ICE). The ICE, expressed as a percent, is calculated as (compound-induced iron excretion/theoretical iron excretion) $\times$ 100. To illustrate, the

theoretical iron excretion after administration of one millimole of DFO, a hexadentate chelator that forms a 1:1 complex with Fe(III), is one milli-g-atom of iron. Two millimoles of desferrithiocin (DFT, **1**) (*Table 1*), a tridentate chelator that forms a 2:1 complex with Fe(III), are required for the theoretical expression of one milli-g-atom of iron.

[00396] The ICE values for compounds **1**, **2**, **3**, **7** and **9** (*Table 1*) are historical and included for comparative purposes (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity.” *J. Med. Chem.* **2006**, 49, 2772-2783; Bergeron *et al.*, “Evaluation of Desferrithiocin and Its Synthetic Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1991**, 34, 2072-2078; Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, 53, 2843-2853; Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258). The biliary ferrokinetics profiles of compounds **3**, **I-1**, **7**, **I-2**, **9**, and **I-3** are presented in *Figure 3*. Each of the rats in these studies was given a single po dose of the compound at 300 $\mu\text{mol/kg}$. The biliary ferrokinetics data for compounds **1** and **2** are not included, simply because these animals were dosed at 150 $\mu\text{mol/kg}$, and the curves are not strictly comparable with the 300 $\mu\text{mol/kg}$ data. These results were published previously (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Evaluation of Desferrithiocin and Its Synthetic Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1991**, 34, 2072-2078).

[00397] DFT (**1**) given to the rats po at a dose of 150 $\mu\text{mol/kg}$ had an ICE of $5.5 \pm 3.2\%$ (Bergeron *et al.*, “Evaluation of Desferrithiocin and Its Synthetic Analogs as Orally Effective Iron Chelators.” *J. Med. Chem.* **1991**, 34, 2072-2078). Maximum iron clearance (MIC) occurred at 3 h, but deferration had returned to baseline levels by 12 h. The desaza analog of DFT, **2**, at 150 $\mu\text{mol/kg}$ had an ICE of $2.7 \pm 0.5\%$ (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440). The compound reached MIC at 6 h and had returned to baseline iron excretion by 12 h post-dose.

[00398] Compound **3** was the least effective compound, with an ICE of $1.1 \pm 0.8\%$ (Bergeron *et al.*, “(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity.” *J. Med. Chem.* **2006**, 49, 2772-2783). It presented with an MIC at 3 h; deferration was virtually over at 9 h (*Figure 3*). The DFT analog of **3**, compound **I-1**, had an ICE that was significantly better than **3**, $9.0 \pm 3.8\%$ ($p < 0.005$). MIC occurred at 6 h and its iron decorporation slowly dropped to near baseline levels by 24 h. The most efficient compound, **7**, had an ICE of $26.7 \pm 4.7\%$ (Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, 53, 2843-2853). The compound also had a very protracted iron clearance; even though its MIC occurred at around 12 h, it was still active at 48 h. Although the biliary ferrokinetics curve of **7** may appear to be biphasic (*Figure 3*), the reason for this unusual line shape is that several animals had temporarily obstructed bile flow. Once the obstruction was resolved, bile volume and overall iron excretion normalized.

[00399] The DFT analog **I-2** had an ICE that was significantly less than that of **7** ($11.7 \pm 1.2\%$ vs. $26.7 \pm 4.7\%$ for **I-2** and **7**, respectively, $p < 0.02$). Compound **I-2** also achieved MIC earlier than **7** (6 h vs. 12 h, *Figure 3*), and the iron clearance induced by **I-2** was basically over by 21 h. Finally, DADFT analog **9** (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258) and its corresponding DFT analog **I-3** presented with similar ICEs (~15%), but with very different biliary ferrokinetics (*Figure 3*). Both compounds achieved MIC at 6 h. However, while compound **9**-induced iron clearance had returned to baseline by 24 h, **I-3** was still quite active.

[00400] Finally, there is an excellent correlation between ICE and $\log P_{app}$ in rodents amongst the DFT analogs **1**, **I-1**, **I-2**, and **I-3** (*Figure 4*). The most lipophilic compounds are also the most active. This ICE vs. $\log P_{app}$ pattern has been observed before (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Iron Chelators and Therapeutic Uses.” In: Abraham, ed. *Burger's Medicinal Chemistry*. 6th. Wiley; New York: **2003**. pp. 479-561; Bergeron *et al.*, “Desferrithiocin Analogs and Nephrotoxicity.” *J. Med. Chem.* **2008**, 51, 5993-6004). In keeping with previous observations, it appears that the ICE/ $\log P_{app}$ curve for DFT and its analogs is beginning to turn over with an optimum $\log P_{app}$ of around -1.4 . This is very different than with the

DADFT analogs, in which the optimum $\log P_{\text{app}}$ was approximately 0.5 (Bergeron *et al.*, “Desferrithiocin Analogs and Nephrotoxicity.” *J. Med. Chem.* **2008**, 51, 5993-6004).

[00401] The primate iron clearance data are provided in *Table 1*. The ICE values for compounds **1**, **2**, **3**, **7**, and **9** have been reported and are included for comparative purposes (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, 53, 2843-2853; Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258). The compounds were given to the primates po at a dose of 75 $\mu\text{mol/kg}$ (**2**, **I-1**, **7**, **I-2**, **9**, and **I-3**) or 150 $\mu\text{mol/kg}$ (**1**, **3**); **I-3** was also given to the primates sc at a dose of 75 $\mu\text{mol/kg}$. Compound **1** was found to have an ICE of $16.1 \pm 8.5\%$ (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440). Removal of the pyridine nitrogen to yield **2** increased the ICE to $21.5 \pm 12\%$ (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440). However, the increase in ICE was not significant ($p > 0.05$). The introduction of a hydroxyl group at the 4'-position of **2** to provide analog **3** resulted in a compound with an ICE of $16.8 \pm 7.2\%$ (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440), which is within error of the ICE found for **1** and **2** ($p > 0.05$). The reintroduction of the pyridine nitrogen into DADFT analog **3** to provide DFT analog **I-1** (*Table 1*) decreased the ICE to $10.0 \pm 2.9\%$, significantly less than its DADFT counterpart, **3**, ($p < 0.05$). When a polyether fragment was attached to the 5'-position of **I-1** to yield **I-2**, the ICE increased to $18.0 \pm 5.2\%$, again, less than that achieved by the corresponding DADFT analog **7**. Likewise, the ICE of **I-3** given po was also less than that of DADFT analog **9** (*Table 1*). In fact, the DADFT analogs were consistently better deferrating agents in the primates than the corresponding DFT analogs (*Table 1*).

[00402] Several generalizations can be derived from *Table 1*. The performance ratios, Performance ratio (PR) values (*i.e.*, $\text{ICE}_{\text{primate}}/\text{ICE}_{\text{rodent}}$) (*Table 1*), show that the compounds are either as effective or better at iron clearance in primates than in rodents. The exception to

this is compound **I-3**. The ICE of this compound given po to the primates is $6.1 \pm 1.8\%$, while in the rats is $14.2 \pm 2.4\%$. Its PR value was 0.4, showing it to be far less efficient in primates than rodents. The poor iron clearance in primates relative to rodents was surprising. Two scenarios were evaluated in search of an explanation: compound-plasma binding and a potential GI absorption problem.

[00403] A compound-plasma binding experiment was performed in which rodent and primate plasma were incubated separately with compound **I-3** at 37 °C for 4 h. Each sample was then passed through a Millipore Amicon Ultra regenerated cellulose filter (3,000 MWCO). The filtrate was assayed for **I-3**. The results indicated there was little, if any, binding of the compound to either the rodent or the primate plasma. This suggests that compound-plasma binding does not explain the difference in the rodent vs. primate ICE values. However, when primates were given compound **I-3** sc, the ICE rose to $16.9 \pm 7.3\%$, which is similar to what was seen in rodents given **I-3** orally (*Table 1*). This observation is consistent with the idea that the primates do not absorb **I-3** well when the compound is administered orally.

[00404] One of the potential problems with introduction of a second hydroxyl in **1** to produce **I-1** or in **2** to generate **3** is the potential for quinone formation owing to the 1,4-relationship between the added hydroxyl and the imine bond. Quinone formation could result in redox cycling and, at a cellular level, untoward side effects. In order to evaluate this, a hydrogen-deuterium exchange study carried out at 25 °C, pH 7.0, in phosphate buffer (0.1 M) with compounds **3**, **I-1**, and **I-2** (*Table 2*). The study revealed the 3'-(H) of **3** to be very labile, as was the 3'-(H) of the desmethyl analog of **3** (Bergeron *et al.*, "Effects of C-4 Stereochemistry and C-4' Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs." *J. Med. Chem.* **1999**, 42, 2432-2440), with less 3'-(H)-deuterium exchange at 4'-(H) with **I-1**, and none with **I-2** (*Table 2*). The exchange rate order is understandable. The most exchangeable 3'-(H) in **3** still exchanges much faster than the 4'-(H) in **I-1**, in which an increase in electron density occurs next to the aromatic nitrogen during enolate/enol formation.

[00405] In a previous study, **1** was given to rats with normal iron stores po once daily at a dose of 384 $\mu\text{mol/kg/d}$ (100 mg/kg/d). All of the rats were dead by day 5 of a planned 10-d experiment. The compound was found to be severely nephrotoxic (Bergeron *et al.*, "A Comparative Study of the Iron-Clearing Properties of Desferrithiocin Analogs with Desferrioxamine B in a *Cebus* Monkey Model." *Blood* **1993**, 81, 2166-2173). The pathologist

noted vacuolar changes of the proximal tubules that were diffuse and severe, with multifocal vacuolar degeneration and necrosis. Nevertheless, the compound's remarkable oral activity initiated a series of SAR studies aimed at the development of orally active, nontoxic DFT analogs. This led to the development of **3** (*Table 1*), which made it to clinical trials (Galanello *et al.*, "A Dose Escalation Study of the Pharmacokinetics, Safety, and Efficacy of Deferitritin, an Oral Iron Chelator in Beta Thalassaemia Patients." *ASH Annu. Meet. Abstr.* **2007**, 110, 2669). The compound, when given once daily, cleared iron from the patients and was proceeding forward. Unfortunately, it was discovered that the compound induced proximal tubule nephrotoxicity when it was administered twice daily, and the trial was halted (Galanello *et al.*, "A Dose Escalation Study of the Pharmacokinetics, Safety, and Efficacy of Deferitritin, an Oral Iron Chelator in Beta Thalassaemia Patients." *ASH Annu. Meet. Abstr.* **2007**, 110, 2669). The compound was reengineered, and it was determined that, by fixing polyether fragments to the 3'- or 4'-position of **3**, *e.g.*, **4** and **5** (*Figure 1*), or **7** (*Table 1*), the renal toxicity virtually disappeared (Bergeron *et al.*, "(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity." *J. Med. Chem.* **2006**, 49, 2772-2783; Bergeron *et al.*, "Design, Synthesis, and Testing of Non-Nephrotoxic Desazadesferrithiocin Polyether Analogs." *J. Med. Chem.* **2008**, 51, 3913-3923; Bergeron *et al.*, "The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs." *J. Med. Chem.* **2010**, 53, 2843-2853; Bergeron *et al.*, "Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity." *Biometals*, **2011**, 24, 239-258). This outcome suggested that the introduction of either a hydroxyl group or a polyether fragment directly into DFT itself might reduce the compound's nephrotoxicity. Accordingly, compounds **I-1**, **I-2** and **I-3** were synthesized and assessed for their toxicity in rodents relative to **1**.

[00406] Assessment of chelator-induced impaired renal function has traditionally relied on the detection of a rise in blood urea nitrogen (BUN) and/or serum creatinine (SCr). However, because of the functional reserve of the kidney, these parameters are often unreliable indicators of acute kidney injury; the ultimate answer requires histopathology. The Critical Path Institute's Preventive Safety Testing Consortium (PSTC) has identified kidney injury molecule-1 (Kim-1, rat) or (KIM-1, human) as an early diagnostic biomarker for monitoring acute kidney tubular toxicity (Goodsaid *et al.*, "Novel Biomarkers of Acute Kidney Toxicity." *Clin. Pharmacol. Ther.* **2009**, 86, 490-496; Hoffmann *et al.*, "Evaluation

of a Urinary Kidney Biomarker Panel in Rat Models of Acute and Subchronic Nephrotoxicity.” *Toxicology* **2010**, 277, 49-58). Kim-1 is a type 1 transmembrane protein located in the epithelial cells of proximal tubules (Han *et al.*, “Kidney Injury Molecule-1 (KIM-1): A Novel Biomarker for Human Renal Proximal Tubule Injury.” *Kidney Int.* **2002**, 62, 237-244; Bonventre, “Kidney Injury Molecule-1 (KIM-1): A Urinary Biomarker and Much More.” *Nephro. Dial. Transplant* **2009**, 24, 3265-3268). After injury, *e.g.*, exposure to a nephrotoxic agent or ischemia, the ectodomain of Kim-1 is shed from the proximal tubular kidney epithelial cells into the urine (Zhou *et al.*, “Comparison of Kidney Injury Molecule-1 and other Nephrotoxicity Biomarkers in Urine and Kidney Following Acute Exposure to Gentamicin, Mercury, and Chromium.” *Toxicol. Sci.* **2008**, 101, 159-170; Vaidya *et al.*, “Urinary Kidney Injury Molecule-1: A Sensitive Quantitative Biomarker for Early Detection of Kidney Tubular Injury.” *Am. J. Physiol. Renal Physiol.* **2006**, 290, F517-F529; Bailly *et al.*, “Shedding of Kidney Injury Molecule-1, A Putative Adhesion Protein Involved in Renal Regeneration.” *J. Biol. Chem.* **2002** 277, 39739-39748). BioAssay Works has recently developed RenaStick, a direct lateral flow immunochromato-graphic assay, which allows for the rapid detection (less than 30 minutes) and quantitation of urinary Kim-1 (rat) or KIM-1 (human) excretion (Vaidya *et al.*, “A Rapid Urine Test for Early Detection of Kidney Injury.” *Kidney. Int.* **2009**, 76, 108-114). In the present study, rats were treated with **1**, **I-1**, **I-2**, and **I-3** given po twice daily at a dose of 237 $\mu\text{mol/kg/dose}$ (474 $\mu\text{mol/kg/d}$) for up to 7 d. Urinary Kim-1 levels were assessed at 24-h intervals (*Figure 5*). The studies were performed on rats with normal iron stores; each animal served as their own control. The data for compounds **3** and **7** have been reported and are included for comparative purposes (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258).

[00407] None of the rats treated with **1** ($n = 5$) survived the planned 7-d exposure to the compound. Two rats became moribund and were sacrificed after being given the compound for four days. The three remaining animals were found dead the morning of day 6; they had received the compound for 5 days. None of the rodents produced any urine on day 5. The **1**-treated rats’ baseline (day 0) urinary Kim-1 value was $< 20 \text{ ng/kg/24 h}$ (*Figure 5A*). After one day of **1**, the Kim-1 had increased nearly 10-fold, to $192 \pm 315 \text{ ng/kg/24 h}$. After three days of **1**, the Kim-1 had further increased to $1528 \pm 539 \text{ ng/kg/24h}$. Blood was taken from the two moribund animals immediately prior to sacrifice; the serum was assessed for its BUN and SCr content. The rats’ BUN was $139 \pm 8 \text{ mg/dl}$ (the normal range has been reported to be 9-30 mg/dl (Antech Diagnostics: www.antechdiagnostics.com/#), while their SCr was

5.1 ± 0.3 mg/dl (normal 0.4-1 mg/dl) (Antech Diagnostics: www.antechdiagnostics.com/#). In addition, as no blood was obtained from the three animals that were found dead, these values likely *underestimate* the actual impact of **1** on these parameters.

[00408] In contrast, in a previous study assessing the impact of **3** on urinary Kim-1 excretion (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258), all of the treated rats ($n = 3$) survived the 237 μmol/kg twice daily (474 μmol/kg/d) × 7 d dosing period. The rats’ baseline (day 0) urinary Kim-1 content was < 20 ng/kg/24 h (*Figure 5B*). After three days of exposure to **3**, the urinary Kim-1 had increased to 69 ± 47 ng/kg/24 h. At the end of the 7 d dosing period, the urinary Kim-1 had further increased to 189 ± 187 ng/kg/24 h (*Figure 5B*). The rats were euthanized on day 8; their BUN at that time was 32 ± 13 mg/dl, while their SCr was 1.3 ± 1.0 mg/dl.

[00409] In the present study, all of the animals treated with the corresponding hydroxylated DFT analog, **I-1**, at 237 μmol/kg twice daily also survived the full 7 d of treatment. The rats’ baseline (day 0) urinary Kim-1 content was < 20 ng/kg/24 h and remained < 50 ng/kg/24h until day 5 (*Figure 5C*). On day 6, the Kim-1 increased to 125 ± 48 ng/kg/24 h, and further increased to 435 ± 269 on day 7 (*Figure 5C*). Although the increase in Kim-1 is greater with the **I-1**-treated rats than with the **3**-treated animals, the increase is not statistically significant ($p > 0.05$). The animals were euthanized on day 8; their BUN at that time was 13 ± 2 mg/dl, while their SCr was 0.5 ± 0.1 mg/dl. Thus, simple hydroxylation of the aromatic ring of both DADFT and DFT, *e.g.*, **3** and **I-1**, respectively, resulted in compounds that were much less toxic than DFT itself.

[00410] It has previously been demonstrated that introducing a polyether fragment in the 3’-, or 4’-position of the DADFT pharmacophore provided remarkably efficient orally active iron chelators that were much less toxic than **3** (Bergeron *et al.*, “(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity.” *J. Med. Chem.* **2006**, 49, 2772-2783; Bergeron, R. J.; Wiegand, J.; Bharti, N.; Singh, S.; Rocca, J. R. Impact of 3,6,9-Trioxadecyloxy Group on Desazadesferrithiocin Analog Iron Chelators and Organ Distribution. *J. Med. Chem.* **2007**, 50, 3302-3313; Bergeron *et al.*, “Design, Synthesis, and Testing of Non-Nephrotoxic Desazadesferrithiocin Polyether Analogs.” *J. Med. Chem.* **2008**, 51, 3913-3923; Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, 53, 2843-2853; Bergeron *et al.*,

“Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258). For example, the impact of compound **7** on urinary Kim-1 excretion was determined when the compound was given po: 1) once daily during the course of 28 d toxicity trials; 2) once daily at a dose of 384 $\mu\text{mol/kg/d} \times 10$ d, and 3) twice daily at a dose of 237 $\mu\text{mol/kg/dose}$ (474 $\mu\text{mol/kg/d}$) $\times 7$ d (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258). All of the rats survived the dosing period. The rats’ baseline (day 0) urinary Kim-1 content was < 20 ng/kg/24 h and stayed within error of this value for the duration of the compound exposure. The data from the 237 $\mu\text{mol/kg}$ twice daily (474 $\mu\text{mol/kg/d}$) $\times 7$ d regimen (Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, 24, 239-258) are depicted in *Figure 5E*. In addition, the BUN and SCr of all of the **7**-treated rats were well within the normal range.

[00411] In the present study, the impact was evaluated that affixing a polyether fragment to DFT itself would have on nephrotoxicity. Accordingly, groups of rats ($n = 5$) were given **I-2** or **I-3** po twice daily at 237 $\mu\text{mol/kg/dose}$ (474 $\mu\text{mol/kg/d}$) $\times 7$ d. All of the animals survived the compound dosing regimen. The animals’ urinary Kim-1 excretion remained within error of that of the baseline (day 0) levels (*Figures 5D* and *5F*). In addition, the BUN and SCr of all of the **I-2**- or **I-3**-treated rats were well within the normal range. Thus, as with the DADFT pharmacophore, fixing a polyether fragment to the DFT framework was an effective tool in further reducing nephrotoxicity.

Conclusion

[00412] It was previously demonstrated that the severe nephrotoxicity associated with **1** could be ameliorated by the removal of the pyridine nitrogen of **1** to provide **2**, and simple hydroxylation of the aromatic ring of **2** to yield **3** (*Figure 1*) (Bergeron *et al.*, “Effects of C-4 Stereochemistry and C-4’ Hydroxylation on the Iron Clearing Efficiency and Toxicity of Desferrithiocin Analogs.” *J. Med. Chem.* **1999**, 42, 2432-2440; Bergeron *et al.*, “Iron Chelators and Therapeutic Uses.” In: Abraham, ed. *Burger's Medicinal Chemistry*. 6th. Wiley; New York: **2003**. pp. 479-561). Further reduction in **3**-induced nephrotoxicity, observed when the compound was given po at 237 $\mu\text{mol/kg}$ twice daily, was accomplished by the addition of polyether fragments, *e.g.*, **4**, **5**, and **7** (*Figure 1* and *Table 1*) (Bergeron *et al.*, “(S)-4,5-Dihydro-2-(2-hydroxy-4-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic Acid Polyethers: A Solution to Nephrotoxicity.” *J. Med. Chem.* **2006**, 49, 2772-2783; Bergeron *et*

al., “Design, Synthesis, and Testing of Non-Nephrotoxic Desazadesferrithiocin Polyether Analogs.” *J. Med. Chem.* **2008**, *51*, 3913-3923; Bergeron *et al.*, “The Impact of Polyether Chain Length on the Iron Clearing Efficiency and Physiochemical Properties of Desferrithiocin Analogs.” *J. Med. Chem.* **2010**, *53*, 2843-2853; Bergeron *et al.*, “Desferrithiocin Analog Iron Chelators: Iron Clearing Efficiency, Tissue Distribution, and Renal Toxicity.” *Biometals*, **2011**, *24*, 239-258). The purpose of the present study was to determine how these same structural modifications to DFT itself would impact the new compounds’ ICE and nephrotoxicity. Accordingly, three DFT analogs, **I-1**, **I-2**, and **I-3**, were synthesized and assessed for their lipophilicity, ICE properties in rats and primates, and for their toxicity in rats.

[00413] DFT (**1**) (*Table 1*) and its analogs were all significantly more water-soluble (lower $\log P_{app}$) than the corresponding DADFT analogs, *e.g.*, **1** vs. **2**, **I-1** vs. **3**, **I-2** vs. **7**, and **I-3** vs. **9**. There was an excellent correlation between ICE and $\log P_{app}$ in rodents amongst the DFT analogs **1**, **I-1**, **I-2**, and **I-3** (*Figure 4*), with the more lipophilic compounds having a greater ICE. This trend is in keeping with previous observations that more lipophilic compounds have better ICE properties. The biliary ferrokinetics of the DFT and DADFT compounds in the bile duct-cannulated rats (*Figure 3*) have similar temporal properties, except for compound **7**. This compound has the highest ICE and a very protracted iron clearance time.

[00414] In the primates, the DADFT analogs were consistently better deferration agents than the corresponding DFT analogs (*Table 1*). The most unusual finding was with compound **I-3**, a DFT analog with a 4’-(3,6-dioxaheptyloxy) ether functionality fixed to the 4’-position of **1**. When the compound was given po to the primates, its ICE was only $6.1 \pm 1.8\%$ vs. $14.2 \pm 2.4\%$ in the rats, and a PR value of 0.4 (*Table 1*). However, when the monkeys were given the compound sc, its ICE increased to $16.9 \pm 7.3\%$, with a PR value now at 1.2. This is consistent with the idea that compound **I-3** simply was not absorbed well orally in primates.

[00415] The effects of structural modification of DFT on its renal toxicity were assessed in rats using a urinary Kim-1 (kidney injury molecule) assay (Vaidya *et al.*, “A Rapid Urine Test for Early Detection of Kidney Injury.” *Kidney. Int.* **2009**, *76*, 108-114), as well as monitoring BUN and SCr. The most notable finding was that fixing a hydroxyl group or a polyether fragment to the DFT aromatic ring resulted in a nearly identical reduction in renal toxicity as seen after the same modification to DADFT (*Figure 5*). Although some

nephrotoxicity was noted with both hydroxylated DADFT and DFT analogs, **3** and **I-1**, respectively, the introduction of polyether groups into either pharmacophore resulted in compounds with little to no impact on renal function, *e.g.*, **7**, **I-2**, and **I-3** (*Figures 5E, 5D, and 5F*).

[00416] In summary, manipulation of the DFT aromatic ring, *e.g.*, hydroxylation, or the introduction of a polyether functionality, can have a marked and unexpected effect on the compound's ICE and renal toxicity (*Table 1* and *Figure 5*). Although the resulting DFT chelators were generally as effective in the rodents as their DADFT counterparts, they were less active in the primates. However, the tissue distribution of **I-1**, **I-2**, and **I-3** in rodents remains to be elucidated. Higher levels of these analogs in the critical target organs, *i.e.*, the liver, heart, and pancreas, could easily compensate for their somewhat lower ICE values. Nevertheless, at least one of the DFT polyethers (**I-2**) was sufficiently effective at iron clearance in rodents (ICE = $11.7 \pm 1.2\%$) and primates (ICE = $18.0 \pm 5.2\%$) and had an acceptable toxicity profile to merit further studies.

EQUIVALENTS AND SCOPE

[00417] In the claims articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[00418] Furthermore, the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, *e.g.*, in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements

and/or features, certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth *in haec verba* herein. It is also noted that the terms “comprising” and “containing” are intended to be open and permits the inclusion of additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.


[00419] This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the invention can be excluded from any claim, for any reason, whether or not related to the existence of prior art.

[00420] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.

CLAIMS

What is claimed is:

1. A compound of Formula (A):



or a pharmaceutically acceptable salt thereof, wherein:

R_1 is hydrogen, alkyl, acyl, or an oxygen protecting group;

R_2 is hydrogen, alkyl, acyl, an oxygen protecting group, $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-R''$,
or $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-(CH_2)_n-NR_{10}-C(=O)O-R''$;

each occurrence of R_3 is independently alkyl, arylalkyl, or $-OR_8$;

R_4 , R_5 , and R_6 are each independently hydrogen or alkyl;

R_7 is $-OR_9$ or $-SR_9$;

R_8 is hydrogen, alkyl, acyl, or an oxygen protecting group;

R_9 is hydrogen, alkyl, an oxygen protecting group when attached to an oxygen atom,
or a sulfur protecting group when attached to a sulfur atom;

R_{10} is hydrogen, alkyl, acyl, or a nitrogen protecting group;

R' is hydrogen or an oxygen protecting group;

R'' is hydrogen, alkyl, acyl, or an oxygen protecting group;


each occurrence of n is independently an integer from 1 to 8, inclusive;

k is an integer from 0 to 2, inclusive;


x is an integer from 1 to 8, inclusive; and

y is an integer from 0 to 8, inclusive.


2. The compound of claim 1, wherein the compound is not of the formula:



3. A compound of Formula (A):



or a pharmaceutically acceptable salt thereof, wherein:




R_2 is hydrogen, alkyl, acyl, an oxygen protecting group, $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-R''$, or $-[(CH_2)_n-O]_x-[(CH_2)_n-O]_y-(CH_2)_n-NR_{10}-C(=O)O-R''$;



each occurrence of R_3 is independently alkyl, arylalkyl, or $-OR_8$;

R_4 , R_5 , and R_6 are each independently hydrogen or alkyl, ;

R_7 is $-OR_9$ or $-SR_9$;




R_8 is hydrogen, alkyl, acyl, an oxygen protecting group,




R_9 is hydrogen, alkyl, , an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

R_{10} is hydrogen, alkyl, acyl, or a nitrogen protecting group;

R' is hydrogen or an oxygen protecting group;



R'' is hydrogen, alkyl, acyl, an oxygen protecting group,




each occurrence of n is independently an integer from 1 to 8, inclusive;


k is an integer from 0 to 2, inclusive;

x is an integer from 1 to 8, inclusive; and

y is an integer from 0 to 8, inclusive;




4. The compound of any one of claims 1-3, wherein the compound is of Formula (B):




or a pharmaceutically acceptable salt thereof.

5. The compound of 1 any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

6. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

7. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

8. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

9. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

10. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof, wherein R_{30} is H, substituted or unsubstituted C_{1-6} alkyl, or an oxygen protecting group; and a is 2, 3, 4, 5, 6, 7, 8, 9, or 10.

11. The compound of claim 10, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

12. The compound of claim 10, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

13. The compound of any one of claims 1-3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof, wherein b is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

14. The compound of claim 13, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

15. The compound of claim 13, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

16. The compound of claim 3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

17. The compound of claim 3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

18. The compound of claim 3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

19. The compound of claim 3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

20. The compound of claim 3, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.



21. The compound of claim 3, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.

22. The compound of claim 3, wherein the compound is of the formula:






or a pharmaceutically acceptable salt thereof.


23. The compound of any one of claims 1-17 and 20-22, wherein R_1 is hydrogen.

24. The compound of any one of claims 1-7, 18-19, and 21-23, wherein R_2 is hydrogen.


25. The compound of any one of claims 1-7, 18-19, and 21-23, wherein R_2 is





26. The compound of claim 25, wherein R_2 is




27. The compound of claim 25, wherein R_2 is





30. The compound of any one of claims 1-7, 18-19, and 21-23, wherein R_2 is



31. The compound of any one of claims 1-7, 18-19, and 21-23, wherein R_2 is alkyl.

32. The compound of claim 31, wherein R_2 is C_{1-12} alkyl optionally substituted with one or more substituents independently selected from the group consisting of hydroxyl, alkyloxy, and $-CO_2H$.

33. The compound of claim 31, wherein R_2 is $-(CH_2)_2-OH$, $-(CH_2)_3-OH$, $-(CH_2)_4-OH$, $-(CH_2)_5-OH$, $-(CH_2)_6-OH$, $-(CH_2)_7-OH$, or $-(CH_2)_8-OH$.

34. The compound of claim 31, wherein R_2 is $-(CH_2)_2-OCH_3$, $-(CH_2)_3-OCH_3$, $-(CH_2)_4-OCH_3$, $-(CH_2)_5-OCH_3$, $-(CH_2)_6-OCH_3$, $-(CH_2)_7-OCH_3$, or $-(CH_2)_8-OCH_3$.


35. The compound of claim 31, wherein R_2 is $-CH_2-CO_2H$, $-(CH_2)_2-CO_2H$, $-(CH_2)_3-CO_2H$, $-(CH_2)_4-CO_2H$, $-(CH_2)_5-CO_2H$, $-(CH_2)_6-CO_2H$, or $-(CH_2)_7-CO_2H$.

36. The compound of any one of claims 1-7 and 21-22, wherein R_1 and R_2 are each hydrogen.

37. The compound of any one of claims 1-5 and 16-36, wherein k is 0.


38. The compound of any one of claims 1-6, 10, 13, and 16-37, wherein R_4 and R_5 are each hydrogen.
39. The compound of any one of claims 1-6, 10, 13, and 16-38, wherein R_6 is C_{1-6} alkyl.
40. The compound of claim 39, wherein R_6 is $-CH_3$.
41. The compound of any one of claims 1-6, 10, 13, and 16-38, wherein R_4 and R_5 are each hydrogen; and R_6 is $-CH_3$.
42. The compound of any one of claims 1-8, 10-11, 13-14, 16-20, and 23-41, wherein R_7 is $-OR_9$.
43. The compound of claim 42, wherein R_7 is $-OMe$, $-OEt$, $-OPr$, or $-OBu$.
44. The compound of claim 42, wherein R_7 is $-OH$.
45. The compound of any one of claims 3-11, 13-14, and 16-44, wherein each instance of R' is hydrogen.
46. The compound of any one of claims 3-11, 13-14, and 16-44, wherein each instance of R' is an oxygen protecting group.
47. The compound of any one of claims 1-9, 18-19, 21-23, 25-26, 30, and 37-46, wherein R'' is hydrogen.
48. The compound of any one of claims 1-9, 18-19, 21-23, 25-26, 30, and 37-46, wherein R'' is C_{1-6} alkyl.
49. The compound of claim 48, wherein R'' is $-CH_3$.
50. The compound of claim 48, wherein R'' is $-CH_2-CO_2H$.


51. The compound of claim 1, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.


52. The compound of claim 1, wherein the compound is of the formula:





or a pharmaceutically acceptable salt thereof.

53. A compound of Formula (J):



or a pharmaceutically acceptable salt thereof, wherein:

===== represents a single bond or double bond;

W_1 - W_3 are independently CR_{22} , NR_{23} , oxygen, or sulfur, provided that:

when W_3 is nitrogen, R_{23} is null, and

when R_{21} is methyl or hydrogen, W_1 is not sulfur;

Z is $-OR_{11}$, $-NR_{12}R_{13}$, morpholine, or optionally substituted piperazinyl;

R_{11} is $-[(CH_2)_p-O]_u-[(CH_2)_q-O]_v-R_{14}$, $-[(CH_2)_p-NH]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$, or $-[(CH_2)_p-O]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$;

R_{12} is hydrogen, alkyl, $-[(CH_2)_p-O]_u-[(CH_2)_q-O]_v-R_{14}$, $-[(CH_2)_p-NH]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$, or $-[(CH_2)_p-O]_u-[(CH_2)_q-NR_{14}]_v-R_{15}$;

R_{13} is hydrogen or alkyl;

p and q are independently an integer from 1 to 8, inclusive;

u is an integer from 0 to 8, inclusive;

v is an integer from 1 to 8, inclusive;

R_{14} and R_{15} are independently hydrogen, alkyl, or acyl;

R_{16} is hydrogen, alkyl, or alkoxy;

R_{17} is $-OR_{18}$ or $-N(OH)R_{19}$;

R_{18} is hydrogen, alkyl, or arylalkyl;

R_{19} is alkyl or $-(CH_2)_s-N(OH)C(=O)R_{20}$;

s is an integer from 1 to 8;


R_{20} is alkyl;

R_{21} is hydrogen or alkyl;

each occurrence of R_{22} is independently null, hydrogen, or alkyl; and

each occurrence of R_{23} is independently null, hydrogen, or alkyl.

54. The compound of claim 53, wherein the compound is of Formula (K):



or a pharmaceutically acceptable salt thereof.

55. The compound of any one of claims 1-54, wherein the pharmaceutically acceptable salt is a pharmaceutically acceptable alkali or alkaline earth metal salt.

56. A pharmaceutical composition comprising a compound of any one of claims 1-55, or a pharmaceutically acceptable salt thereof, and optionally a pharmaceutically acceptable excipient.

57. The pharmaceutical composition of claim 56, wherein the pharmaceutical composition is for use in treating a pathological condition selected from the group consisting of iron overload, aluminum overload, lanthanide overload, actinide overload, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and an infectious disease.

58. The pharmaceutical composition of claim 57, wherein the compound is of a therapeutically effective amount.

59. A method of treating a pathological condition selected from the group consisting of iron overload, aluminum overload, lanthanide overload, actinide overload, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and an infectious disease, in a subject, the method comprising administering to the subject a therapeutically effective amount of a compound of any one of claims 1-55, or a pharmaceutically acceptable salt, or a pharmaceutical composition of any one of claims 56-58.

60. A method of treating a pathological condition selected from the group consisting of iron overload, aluminum overload, lanthanide overload, actinide overload, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and an infectious disease, in a subject, the method comprising:

mixing with blood a therapeutically effective amount of a compound of any one of claims 1-55, or a pharmaceutically acceptable salt, or a pharmaceutical composition of any one of claims 56-58; and

administering to the subject the mixture.

61. The method of claim 59 or 60, wherein the subject is a mammal.

62. The method of claim 59 or 60, wherein the subject is a human.

63. The method of any one of claims 59-62, wherein the pathological condition is iron overload.

64. The method of any one of claims 59-62, wherein the pathological condition is aluminum overload, lanthanide overload, or actinide overload.

65. The method of any one of claims 59-62, wherein the pathological condition is oxidative stress.

66. The method of any one of claims 59-62, wherein the pathological condition is transfusional iron overload.

67. The method of any one of claims 59-62, wherein the pathological condition is thalassemia, primary hemochromatosis, or secondary hemochromatosis.

68. The method of any one of claims 59-62, wherein the pathological condition is radiation injury.

69. The method of any one of claims 59-62, wherein the pathological condition is Friedreich's ataxia (FRDA).

70. The method of any one of claims 59-62, wherein the pathological condition is diabetes, liver disease, heart disease, cancer, neurological or neurodegenerative disorder, macular degeneration, closed head injury, irritable bowel disease, or reperfusion injury.

71. The method of any one of claims 59-62, wherein the pathological condition is an infectious disease.


72. The method of any one of claims 59-62, wherein the pathological condition is malaria.


73. A compound of any one of claims 1-55, or a pharmaceutically acceptable salt, or a pharmaceutical composition of any one of claims 56-58, for use in treating a pathological condition selected from the group consisting of iron overload, aluminum overload, lanthanide overload, actinide overload, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and an infectious disease in a subject.

74. A kit for treating a pathological condition selected from the group consisting of iron overload, aluminum overload, lanthanide overload, actinide overload, oxidative stress, transfusional iron overload, thalassemia, primary hemochromatosis, secondary hemochromatosis, diabetes, liver disease, heart disease, cancer, radiation injury, neurological or neurodegenerative disorder, Friedreich's ataxia (FRDA), macular degeneration, closed head injury, irritable bowel disease, reperfusion injury, and an infectious disease in a subject, the kit comprising:

a first container comprising a therapeutically effective amount of a compound of any one of claims 1-55, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of any one of claims 56-58; and

instructions for administering the compound, or the pharmaceutically acceptable salt thereof, or the pharmaceutical composition, to the subject to treat the pathological condition.

**Figure 1A**

**Figure 1B**




Figure 2




Figure 3




Figure 4




Figure 5A




Figure 5B




Figure 5C




Figure 5D




Figure 5E







Figure 5F




I-1




9



7




I-2




I-3

Figure 1B



(A)



(J)