

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Factor VII or VIIa Gla domain variants

(51)⁶ International Patent Classification(s)
C07K 14/745 2BHAU **C12N**
 (2006.01) 9/64
A61K 38/48 (2006.01) 20060101ALI2005100
C12N 9/64 (2006.01) 8BMEP **C12N**
C12N 15/57 (2006.01) 15/57
A61K 38/00 (2006.01) 20060101ALI2006052
C07K 14/745 1BMWO **A61K**
 20060101AFI2008092 38/00
2BHAU **A61K** 20060101ALN200607
 38/48 22BMEP
 20060101ALI2008092 PCT/DK2004/000428

(21) Application No: 2004247799 (22) Application Date: 2004.06.18

(87) WIPO No: WO04/111242

(30) Priority Data

(31) Number	(32) Date	(33) Country
PA 2004 00930	2004.06.15	DK
60/479,780	2003.06.19	US

(43) Publication Date: 2004.12.23

(71) Applicant(s)

Bayer HealthCare LLC

(72) Inventor(s)

Bornæs, Claus, Haaning, Jesper Mortensen, Andersen, Kim Vilbour

(74) Agent/Attorney

Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
23 December 2004 (23.12.2004)

PCT

(10) International Publication Number
WO 2004/111242 A1

(51) International Patent Classification⁷: C12N 15/57, 9/64, C07K 14/745, A61K 38/36

(21) International Application Number: PCT/DK2004/000428

(22) International Filing Date: 18 June 2004 (18.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/479,780 19 June 2003 (19.06.2003) US PA 2004 00930 15 June 2004 (15.06.2004) DK

(71) Applicants (for all designated States except US): MAXYGEN HOLDINGS LTD.; c/o Close Brothers (Cayman) Limited, 103 South Church Street, P.O. Box 1034 GT, Grand Cayman, British West Indies (KY). MAXYGEN ApS [DK/DK]; Agern Allé 1, DK-2970 Hørsholm (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HAANING, Jesper, Mortensen [DK/DK]; Biskop Svanes Vej 2E, DK-3460 Birkeryd (DK). ANDERSEN, Kim, Vilbørg [DK/DK]; Karlslads Allé 7, DK-2700 Brønshøj (DK). BORNÆS, Claus [DK/DK]; 6, Højgaards Allé, DK-2900 Hellerup (DK).


(74) Common Representative: MAXYGEN ApS; Agern Allé 1, DK-2970 Hørsholm (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SI, SG, SK, SL, SY, TJ, TM,

[Continued on next page]

(54) Title: FACTOR VII OR VIIa GLA DOMAIN VARIANTS

WO 2004/111242 A1



(57) Abstract: GlA domain variants of human Factor VII or human Factor VIIa, comprising 1-15 amino acid modifications relative to human Factor VII or human Factor VIIa, wherein a hydrophobic amino acid residue has been introduced by substitution in position 34, or having an amino acid substitution in position 36, or having amino acid substitutions in positions 10 and 32 and at least one further amino acid substitution in a position selected from 74, 77 and 116.



TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

SK, TR), OAPI (BF, BI, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NI, SN, TD, TG).

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,

Published:
— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

FACTOR VII OR VIIa GLA DOMAIN VARIANTS**FIELD OF THE INVENTION**

The present invention relates to novel Gla domain variants of Factor FVII (FVII) or 5 Factor VIIa (FVIIa) polypeptides, as well as the use of such polypeptide variants in therapy, in particular for the treatment of a variety of coagulation-related disorders.

BACKGROUND OF THE INVENTION

Blood coagulation is a process consisting of a complex interaction of various blood 10 components (or factors) that eventually results in a fibrin clot. Generally, the blood components participating in what has been referred to as the "coagulation cascade" are proenzymes or zymogens, i.e. enzymatically inactive proteins that are converted into an active form by the action of an activator. One of these coagulation factors is FVII.

FVII is a vitamin K-dependent plasma protein synthesized in the liver and secreted 15 into the blood as a single-chain glycoprotein with a molecular weight of 53 kDa (Broze & Majerus, *J. Biol. Chem.* 1980; 255:1242-1247). The FVII zymogen is converted into an activated form (FVIIa) by proteolytic cleavage at a single site, R152-I153, resulting in two chains linked by a single disulfide bridge. FVIIa in complex with tissue factor (FVIIa complex) is able to convert both factor IX (FIX) and factor X (FX) into their activated forms, 20 followed by reactions leading to rapid thrombin production and fibrin formation (Østerud & Rapaport, *Proc Natl Acad Sci USA* 1977; 74:5260-5264).

FVII undergoes post-translational modifications, including vitamin K-dependent 25 carboxylation resulting in ten γ -carboxyglutamic acid residues in the N-terminal region of the molecule. Thus, residues number 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35 shown in SEQ ID NO:1 are γ -carboxyglutamic acid residues in the Gla domain important for FVII activity. Other post-translational modifications include sugar moiety attachment at two naturally occurring N-glycosylation sites at position 145 and 322, respectively, and at two naturally occurring O-glycosylation sites at position 52 and 60, respectively.

The gene coding for human FVII (hFVII) has been mapped to chromosome 13 at 30 q34-pter 9 (de Grouchy et al., *Hum Genet* 1984; 66:230-233). It contains nine exons and spans 12.8 Kb (O'Hara et al., *Proc Natl Acad Sci USA* 1987; 84:5158-5162). The gene organisation and protein structure of FVII are similar to those of other vitamin K-dependent procoagulant proteins, with exons 1a and 1b encoding for signal sequence; exon 2 the propeptide and Gla domain; exon 3 a short hydrophobic region; exons 4 and 5 the epidermal

growth factor-like domains; and exon 6 through 8 the serine protease catalytic domain (Yoshitake et al., *Biochemistry* 1985; 24: 3736-3750).

Reports exist on experimental three-dimensional structures of hFVIIa (Pike et al., *Proc Natl Acad Sci USA*, 1999; 96:8925-30 and Kemball-Cook et al., *J. Struct. Biol.*, 1999; 127:213-223); of hFVIIa in complex with soluble tissue factor using X-ray crystallographic methods (Banner et al., *Nature*, 1996; 380:41 and Zhang et al., *J. Mol. Biol.*, 1999; 285: 2089); and of smaller fragments of hFVII (Muranyi et al., *Biochemistry*, 1998; 37:10605 and Kao et al., *Biochemistry*, 1999; 38:7097).

Relatively few protein-engineered variants of FVII have been reported (Dickinson & Ruf, *J Biol Chem*, 1997;272:19875-19879; Kemball-Cook et al., *J Biol Chem*, 1998; 273:8516-8521; Bharadwaj et al., *J Biol Chem*, 1996; 271:30685-30691; Ruf et al., *Biochemistry*, 1999; 38:1957-1966).

Reports exist on expression of FVII in BHK or other mammalian cells (WO 92/15686, WO 91/11514 and WO 88/10295) and co-expression of FVII and kex2 endoprotease in eukaryotic cells (WO 00/28065).

Commercial preparations of recombinant human FVIIa (rhFVIIa) are sold under the trademark NovoSeven®. NovoSeven® is indicated for the treatment of bleeding episodes in hemophilia A or B patients. NovoSeven® is the only rhFVIIa for effective and reliable treatment of bleeding episodes currently available on the market.

An inactive form of FVII in which arginine 152 and/or isoleucine 153 are modified has been reported in WO 91/11514. These amino acids are located at the activation site. WO 96/12800 describes inactivation of FVIIa by a serine proteinase inhibitor. Inactivation by carbamylation of FVIIa at the α -amino acid group I153 has been described by Petersen et al., *Eur J Biochem*, 1999;261:124-129. The inactivated form is capable of competing with wild-type FVII or FVIIa for binding to tissue factor and inhibiting clotting activity. The inactivated form of FVIIa is suggested to be used for treatment of patients suffering from hypercoagulable states, such as patients with sepsis or at risk of myocardial infarction or thrombotic stroke.

In connection with treatment of uncontrolled bleedings such as trauma it is believed that FVIIa is capable of activating FX to FXa without binding to tissue factor, and this activation reaction is believed to occur primarily on activated blood platelets (Hedner et al., *Blood Coagulation & Fibrinolysis*, 2000;11:107-111). However, hFVIIa or rhFVIIa has a low activity towards FX in the absence of tissue factor and, consequently, treatment of uncontrolled bleeding, for example in trauma patients, requires relatively high and multiple

doses of hFVIIa or rhFVIIa. Therefore, in order to treat uncontrolled bleedings more efficiently (to minimize blood loss) there is need for improved FVIIa molecules which possess a high activity toward FX in the absence of tissue factor. Such improved FVIIa molecules should exhibit a lowered clotting time (faster action/increased clotting activity) as compared to rhFVIIa when administered in connection with uncontrolled bleedings.

5 Gla domain variants of FVII/FVIIa have been disclosed in WO 99/20767, US 6,017,882 and WO 00/66753, where some residues located in the Gla domain were identified as being important for phospholipid membrane binding and hence FX activation. In particular, it was found that the residues 10 and 32 were critical and that increased 10 phospholipid membrane binding affinity, and hence increased FX activation, could be achieved by performing the mutations P10Q and K32E. In particular, it was found that FX activation was enhanced as compared to rhFVIIa at marginal coagulation conditions, such as under conditions where a low level of tissue factor is present.

WO 01/58935 discloses a new strategy for developing FVII or FVIIa molecules 15 having *inter alia* an increased half-life by means of directed glycosylation or PEGylation.

WO 03/093465 discloses FVII or FVIIa variants having certain modifications in the Gla domain and having one or more N-glycosylation sites introduced outside the Gla domain.

WO 2004/029091 discloses FVII or FVIIa variants having certain modifications in the tissue factor binding site.

20 The present inventors have now identified further residues in the Gla domain which further increase the phospholipid membrane binding affinity and hence further increase FX activation. The FVII or FVIIa variants of the invention may also exhibit reduced tissue factor binding affinity.

The object of the present invention is to provide improved FVII or FVIIa molecules 25 (FVII or FVIIa variants) which are capable of activating FX to FXa more efficiently than hFVIIa, rhFVIIa or [P10Q+K32E]rhFVIIa. In particular, it is an object of the present invention to provide improved FVII or FVIIa molecules (FVII or FVIIa variants) which are capable of activating FX to FXa more efficiently than hFVIIa, rhFVIIa or [P10Q+K32E]rhFVIIa in the absence of tissue factor. These objects are addressed by the 30 FVII or FVIIa variants provided herein.

BRIEF DISCLOSURE OF THE INVENTION

In a first aspect the present invention relates to a Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence comprising 1-15 amino acid

modifications relative to human Factor VII (hFVII) or human Factor VIIa (hFVIIa) with the amino acid sequence shown in SEQ ID NO: 1, wherein a hydrophobic amino acid residue has been introduced by substitution in position 34.

In a second aspect the invention relates to a Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence comprising 1-15 amino acid modifications relative to human Factor VII (hFVII) or human Factor VIIa (hFVIIa) with the amino acid sequence shown in SEQ ID NO: 1, wherein the amino acid sequence comprises an amino acid substitution in position 36.

In a third aspect the invention relates to a Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence comprising 3-15 amino acid modifications relative to human Factor VII (hFVII) or human Factor VIIa (hFVIIa) having the amino acid sequence shown in SEQ ID NO: 1, wherein amino acid sequence comprises an amino acid substitution in positions 10 and 32 and at least one further amino acid substitution in a position selected from the group consisting of positions 74, 77 and 116.

In a further aspect the invention relates to a Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence that differs in 1-15 amino acid residues relative to the amino acid sequence of human Factor VII (hFVII) or human Factor VIIa (hFVIIa) shown in SEQ ID NO: 1, wherein a negatively charged amino acid residue has been introduced by substitution in position 36.

Further aspects of the invention relate to a nucleotide sequence encoding the polypeptide variants of the invention, an expression vector comprising the nucleotide sequence, and a host cell comprising the nucleotide sequence or expression vector.

Still further aspects of the invention relate to a pharmaceutical composition comprising the polypeptide variants of the invention, use of the polypeptide variants of the invention or the pharmaceutical composition of the invention as a medicament, as well as methods of treatment using the polypeptide variants or pharmaceutical compositions of the invention.

In a further aspect the invention relates to a method for treating a mammal having a disease or a disorder wherein clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of the variant of the invention.

In a further aspect the invention relates to use of a variant of the invention in the manufacture of a medicament for treatment of a mammal having a disease or disorder wherein clot formation is desirable.

Further aspects of the present invention will be apparent from the description below as well as from the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1 shows the clotting time vs. concentration for variants of the invention when assayed in the "Whole Blood Assay".

Figure 2 shows the maximum tissue factor-dependent thrombin generation rate for variants of the invention determined in the "Thrombogram Assay".

10 Figure 3 shows the maximum phospholipid-dependent thrombin generation rate for variants of the invention determined in the "Thrombogram Assay".

DETAILED DISCLOSURE OF THE INVENTION**Definitions**

In the context of the present description and claims the following definitions apply:

5 The term "FVII" or "FVII polypeptide" refers to a FVII molecule provided in single chain form. One example of a FVII polypeptide is the wild-type human FVII (hFVII) having the amino acid sequence shown in SEQ ID NO:1. It should be understood, however, that the term "FVII polypeptide" also covers hFVII-like molecules, such as fragments or variants of SEQ ID NO:1, in particular variants where the sequence comprises at least one, such as up to 10, 15, preferably up to 10, amino acid modifications as compared to SEQ ID NO:1.

The term "FVIIa" or "FVIIa polypeptide" refers to a FVIIa molecule provided in its activated two-chain form. When the amino acid sequence of SEQ ID NO:1 is used to describe the amino acid sequence of FVIIa it will be understood that the peptide bond between R152 and I153 of the single-chain form has been cleaved, and that one of the chains 15 comprises amino acid residues 1-152, the other chain comprises amino acid residues 153-406.

The terms "rFVII" and "rFVIIa" refer to FVII and FVIIa polypeptides produced by recombinant techniques.

20 The terms "hFVII" and "hFVIIa" refer to human wild-type FVII and FVIIa, respectively, having the amino acid sequence shown in SEQ ID NO:1.

The terms "rhFVII" and "rhFVIIa" refer to human wild-type FVII and FVIIa, having the amino acid sequence shown in SEQ ID NO:1, produced by recombinant means. An example of rhFVIIa is NovoSeven®.

25 When used herein, the term "Gla domain" is intended to cover amino acid residues 1 to 45 of SEQ ID NO:1.

Accordingly, the term "position located outside the Gla domain" covers amino acid residues 46-406 of SEQ ID NO:1.

The abbreviations "FX", "TF" and "TFPI" mean Factor X, Tissue Factor and Tissue Factor Pathway Inhibitor, respectively.

30 The term "protease domain" is used about residues 153-406 counted from the N-terminus.

The term "catalytic site" is used to mean the catalytic triad consisting of S344, D242 and H193 of the polypeptide variant.

The term "parent" is intended to indicate the molecule to be modified/improved in accordance with the present invention. Although the parent polypeptide to be modified by the present invention may be any FVII or FVIIa polypeptide, and thus be derived from any origin, e.g. a non-human mammalian origin, it is preferred that the parent polypeptide is

5 hFVII or hFVIIa.

A "variant" is a polypeptide which differs in one or more amino acid residues from its parent polypeptide, normally in 1-15 amino acid residues (e.g. in 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues), such as in 1-10 amino acid residues, e.g. in 1-8, 1-6, 1-5 or 1-3 amino acid residues. Normally, the parent polypeptide is hFVII or hFVIIa.

10 The term "conjugate" (or interchangeably "conjugated polypeptide") is intended to indicate a heterogeneous (in the sense of composite or chimeric) molecule formed by the covalent attachment of one or more polypeptides to one or more non-polypeptide moieties such as polymer molecules, lipophilic compounds, sugar moieties or organic derivatizing agents. Preferably, the conjugate is soluble at relevant concentrations and conditions, i.e. 15 soluble in physiological fluids such as blood. Examples of conjugated polypeptides of the invention include glycosylated and/or PEGylated polypeptides.

The term "covalent attachment" or "covalently attached" means that the polypeptide variant and the non-polypeptide moiety are either directly covalently joined to one another, or else are indirectly covalently joined to one another through at least one intervening moiety 20 such as a bridge, spacer, or linkage moiety.

The term "non-polypeptide moiety" is intended to mean a molecule, different from a peptide polymer composed of amino acid monomers and linked together by peptide bonds, which molecule is capable of conjugating to an attachment group of the polypeptide variant of the invention. Preferred examples of such molecules include polymer molecules, sugar 25 moieties, lipophilic compounds or organic derivatizing agents. When used in the context of a conjugated variant of the invention it will be understood that the non-polypeptide moiety is linked to the polypeptide part of the conjugated variant through an attachment group of the polypeptide. As explained above, the non-polypeptide moiety can be directly or indirectly covalently joined to the attachment group.

30 A "polymer molecule" is a molecule formed by covalent linkage of two or more monomers, wherein none of the monomers is an amino acid residue, except where the polymer is human albumin or another abundant plasma protein. The term "polymer" may be used interchangeably with the term "polymer molecule". The term is also intended to cover carbohydrate molecules attached by *in vitro* glycosylation, i.e. a synthetic glycosylation

performed *in vitro* normally involving covalently linking a carbohydrate molecule to an attachment group of the polypeptide variant, optionally using a cross-linking agent.

The term "sugar moiety" is intended to indicate a carbohydrate-containing molecule comprising one or more monosaccharide residues, capable of being attached to the

5 polypeptide variant (to produce a polypeptide variant conjugate in the form of a glycosylated polypeptide variant) by way of *in vivo* glycosylation. The term "*in vivo* glycosylation" is intended to mean any attachment of a sugar moiety occurring *in vivo*, i.e. during posttranslational processing in a glycosylating cell used for expression of the polypeptide variant, e.g. by way of N-linked and O-linked glycosylation. The exact oligosaccharide

10 structure depends, to a large extent, on the glycosylating organism in question.

An "N-glycosylation site" has the sequence N-X-S/T/C, wherein X is any amino acid residue except proline, N is asparagine and S/T/C is either serine, threonine or cysteine, preferably serine or threonine, and most preferably threonine. Preferably, the amino acid residue in position +3 relative to the asparagine residue is not a proline residue.

15 An "O-glycosylation site" is the OH-group of a serine or threonine residue.

The term "attachment group" is intended to indicate a functional group of the polypeptide variant, in particular of an amino acid residue thereof or a carbohydrate moiety, capable of attaching a non-polypeptide moiety such as a polymer molecule, a lipophilic molecule, a sugar moiety or an organic derivatizing agent. Useful attachment groups and

20 their matching non-polypeptide moieties are apparent from the table below.

Attachment group	Amino acid	Examples of non-polypeptide moiety	Conjugation method/-Activated PEG	Reference
-NH ₂	N-terminal, Lys	Polymer, e.g. PEG, with amide or imine group	mPEG-SPA Tresylated mPEG	Nektar Therapeutics Delgado et al, Critical reviews in Therapeutic Drug Carrier Systems 9(3,4):249-304 (1992)
-COOH	C-terminal, Asp, Glu	Polymer, e.g. PEG, with ester or amide group Carbohydrate moiety	mPEG-Hz <i>In vitro</i> coupling	Nektar Therapeutics

-SH	Cys	Polymer, e.g. PEG, with disulfide, maleimide or vinyl sulfone group Carbohydrate moiety	PEG-vinylsulphone PEG-maleimide <i>In vitro</i> coupling	Nektar Therapeutics Delgado et al, Critical reviews in Therapeutic Drug Carrier Systems 9(3,4):249-304 (1992)
-OH	Ser, Thr, Lys, OH-	Sugar moiety PEG with ester, ether, carbamate, carbonate	<i>In vivo</i> O-linked glycosylation	
-CONH ₂	Asn as part of an N-glycosylation site	Sugar moiety Polymer, e.g. PEG	<i>In vivo</i> N-glycosylation	
Aromatic residue	Phe, Tyr, Trp	Carbohydrate moiety	<i>In vitro</i> coupling	
-CONH ₂	Gln	Carbohydrate moiety	<i>In vitro</i> coupling	Yan and Wold, Biochemistry, 1984, Jul 31; 23(16): 3759-65
Aldehyde Ktone	Oxidized oligo-saccharide	Polymer, e.g. PEG, PEG-hydrazide	PEGylation	Andresz et al., 1978, <i>Macromol. Chem.</i> 179:301, WO 92/16555, WO 00/23114
Guanidino	Arg	Carbohydrate moiety	<i>In vitro</i> coupling	Lundblad and Noyes, Chemical Reagents for Protein Modification, CRC Press Inc., Florida, USA
Imidazole ring	His	Carbohydrate moiety	<i>In vitro</i> coupling	As for guanidine

For *in vivo* N-glycosylation, the term "attachment group" is used in an unconventional way to indicate the amino acid residues constituting a N-glycosylation site (with the sequence N-X-S/T/C as indicated above). Although the asparagine residue of the N-glycosylation site is the one to which the sugar moiety is attached during glycosylation, such attachment cannot be achieved unless the other amino acid residues of the N-glycosylation site are present.

Accordingly, when the non-polypeptide moiety is a sugar moiety and the conjugation is to be achieved by *in vivo* N-glycosylation, the term "amino acid residue comprising an attachment group for a non-polypeptide moiety" as used in connection with alterations of the amino acid sequence of the polypeptide is to be understood as meaning that

one or more amino acid residues constituting an *in vivo* N-glycosylation site are to be altered in such a manner that a functional *in vivo* N-glycosylation site is introduced into the amino acid sequence.

In the present application, amino acid names and atom names (e.g. CA, CB, CD, 5 CG, SG, NZ, N, O, C, etc) are used as defined by the Protein DataBank (PDB) (www.pdb.org) based on the IUPAC nomenclature (IUPAC Nomenclature and Symbolism for Amino Acids and Peptides (residue names, atom names, etc.), *Eur. J. Biochem.*, 138, 9-37 (1984) together with their corrections in *Eur. J. Biochem.*, 152, 1 (1985)).

The term "amino acid residue" is intended to include any natural or synthetic amino acid residue, and is primarily intended to indicate an amino acid residue contained in the group consisting of the 20 naturally occurring amino acids, i.e. selected from the group consisting of alanine (Ala or A), cysteine (Cys or C), aspartic acid (Asp or D), glutamic acid (Glu or E), phenylalanine (Phe or F), glycine (Gly or G), histidine (His or H), isoleucine (Ile or I), lysine (Lys or K), leucine (Leu or L), methionine (Met or M), asparagine (Asn or N), 10 proline (Pro or P), glutamine (Gln or Q), arginine (Arg or R), serine (Ser or S), threonine (Thr or T), valine (Val or V), tryptophan (Trp or W), and tyrosine (Tyr or Y) residues.

The terminology used for identifying amino acid positions is illustrated as follows: G124 indicates that position 124 is occupied by a glycine residue in the amino acid sequence shown in SEQ ID NO:1. G124R indicates that the glycine residue of position 124 has been 20 substituted with an arginine residue. Alternative substitutions are indicated with a "/", e.g. N145S/T means an amino acid sequence in which asparagine in position 145 is substituted with either serine or threonine. Multiple substitutions are indicated with a "+", e.g. K143N+N145S/T means an amino acid sequence which comprises a substitution of the lysine residue in position 143 with an asparagine residue and a substitution of the asparagine residue 25 in position 145 with a serine or a threonine residue. Insertion of an additional amino acid residue, e.g. insertion of an alanine residue after G124, is indicated by G124GA. Insertion of two additional alanine residues after G124 is indicated by G124GAA, etc. When used herein, the term "inserted in position X" or "inserted at position X" means that the amino acid residue(s) is (are) inserted between amino acid residue X and X+1. A deletion of an amino 30 acid residue is indicated by an asterix. For example, deletion of the glycine residue in position 124 is indicated by G124*.

Unless otherwise indicated, the numbering of amino acid residues made herein is made relative to the amino acid sequence of the hFVII/hFVIIa polypeptide (SEQ ID NO:1).

The term "differs from" as used in connection with specific mutations is intended to allow for additional differences being present apart from the specified amino acid difference. For instance, in addition to the modifications performed in the Gla domain aiming at increasing the FX activation, the polypeptide may contain other modifications that are not necessarily related to this effect.

5 Thus, in addition to the amino acid modifications disclosed herein, it will be understood that the amino acid sequence of the polypeptide variant of the invention may, if desired, contain other alterations, i.e. other substitutions, insertions or deletions. These may, for example, include truncation of the N- and/or C-terminus by one or more amino acid 10 residues (e.g. by 1-10 amino acid residues), or addition of one or more extra residues at the N- and/or C-terminus, e.g. addition of a methionine residue at the N-terminus or introduction of a cysteine residue near or at the C-terminus, as well as "conservative amino acid substitutions", i.e. substitutions performed within groups of amino acids with similar characteristics, e.g. small amino acids, acidic amino acids, polar amino acids, basic amino 15 acids, hydrophobic amino acids and aromatic amino acids.

Examples of such conservative substitutions are shown in the below table.

1	Alanine (A)	Glycine (G)	Serine (S)	Threonine (T)
2	Aspartic acid (D)	Glutamic acid (E)		
3	Asparagine (N)	Glutamine (Q)		
4	Arginine (R)	Histidine (H)	Lysine (K)	
5	Isoleucine (I)	Leucine (L)	Methionine (M)	Valine (V)
6	Phenylalanine (F)	Tyrosine (Y)	Tryptophan (W)	

Still other examples of additional modifications are disclosed in the sections entitled 20 "Modifications outside the Gla domain" and "Other modifications outside the Gla domain".

The term "nucleotide sequence" is intended to indicate a consecutive stretch of two or more nucleotide molecules. The nucleotide sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

The term "vector" refers to a plasmid or other nucleotide sequences that are capable 25 of replicating within a host cell or being integrated into the host cell genome, and as such, are useful for performing different functions in conjunction with compatible host cells (a vector-host system) to facilitate the cloning of the nucleotide sequence, i.e. to produce useful quantities of the sequence, to direct the expression of the gene product encoded by the

sequence and to integrate the nucleotide sequence into the genome of the host cell. The vector will contain different components depending upon the function it is to perform.

“Cell”, “host cell”, “cell line” and “cell culture” are used interchangeably herein and all such terms should be understood to include progeny resulting from growth or culturing of a cell.

“Transformation” and “transfection” are used interchangeably to refer to the process of introducing DNA into a cell.

“Operably linked” refers to the covalent joining of two or more nucleotide sequences, by means of enzymatic ligation or otherwise, in a configuration relative to one another such that the normal function of the sequences can be performed. Generally, “operably linked” means that the nucleotide sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used, in conjunction with standard recombinant DNA methods.

In the context of the present invention the term “modification” or “amino acid modification” is intended to cover replacement of an amino acid side chain, substitution of an amino acid residue, deletion of an amino acid residue or insertion of an amino acid residue.

The term “introduce” refers to introduction of an amino acid residue, in particular by substitution of an existing amino acid residue, or alternatively by insertion of an additional amino acid residue.

The term “remove” refers to removal of an amino acid residue, in particular by substitution of the amino acid residue to be removed by another amino acid residue, or alternatively by deletion (without substitution) of the amino acid residue to be removed.

In the present context, the term “activity” should be understood as the relevant activity associated with the assay in which the activity is actually measured.

Thus, the term “amidolytic activity” is used to mean the activity measured in the “Amidolytic Assay” described herein. In order to exhibit “amidolytic activity” a variant of the invention, in its activated form, should have at least 10% of the amidolytic activity of rhFVIIa when assayed in the “Amidolytic Assay” described herein. In a preferred embodiment of the invention the variant, in its activated form, has at least 20% of the amidolytic activity of rhFVIIa, such as at least 30%, e.g. at least 40%, more preferably at least 50%, such as at least 60%, e.g. at least 70%, even more preferably at least 80%, such as at least 90% of the amidolytic activity of rhFVIIa when assayed in the “Amidolytic Assay”

described herein. In an interesting embodiment the variant, in its activated form, has substantially the same amidolytic activity as rhFVIIa, such as an amidolytic activity of 75-125% of the amidolytic activity of rhFVIIa.

The term "clotting activity" refers to the activity measured in the "Whole Blood Assay" described herein, i.e. the time needed to obtain clot formation. Thus, a lower clotting time corresponds to a higher clotting activity.

The term "increased clotting activity" is used to indicate that the clotting time of the polypeptide variant is statistically significantly decreased relative to that generated by rhFVIIa or [P10Q+K32E]rhFVIIa as determined under comparable conditions and when measured in the "Whole Blood Assay" described herein.

In the present context, the term "activity" is also used in connection with the variants' capability of activating FX to FXa. This activity is also denoted "FX activation activity" or "FXa generation activity" and may be determined in the "TF-independent Factor X Activation Assay" described herein.

The term "increased FX activation activity" or "increased FXa generation activity" is used to indicate that a variant of the invention, in its activated form, has a statistically significantly increased capability to activate FX to FXa as compared to a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa. To what extent a variant of the invention (in its activated form) has an increased FX activation activity may conveniently be determined in the "TF-independent Factor X Activation Assay" described herein.

The term "immunogenicity" as used in connection with a given substance is intended to indicate the ability of the substance to induce a response from the immune system. The immune response may be a cell or antibody mediated response (see, e.g., Roitt: Essential Immunology (10th Edition, Blackwell) for further definition of immunogenicity). Normally, reduced antibody reactivity will be an indication of reduced immunogenicity. The immunogenicity may be determined by use of any suitable method known in the art, e.g. *in vivo* or *in vitro*.

The term "functional *in vivo* half-life" is used in its normal meaning, i.e. the time at which 50% of the biological activity of the polypeptide is still present in the body/target organ, or the time at which the activity of the polypeptide is 50% of the initial value.

As an alternative to determining functional *in vivo* half-life, "serum half-life" may be determined, i.e. the time at which 50% of the polypeptide circulates in the plasma or bloodstream prior to being cleared. Determination of serum half-life is often more simple than determining the functional *in vivo* half-life, and the magnitude of serum half-life is

usually a good indication of the magnitude of functional *in vivo* half-life. Alternative terms to serum half-life include "plasma half-life", "circulating half-life", "serum clearance", "plasma clearance" and "clearance half-life". The polypeptide is cleared by the action of one or more of the reticuloendothelial systems (RES), kidney, spleen or liver, by tissue factor, SEC

5 receptor or other receptor mediated elimination, or by specific or unspecific proteolysis.

Normally, clearance depends on size (relative to the cutoff for glomerular filtration), charge, attached carbohydrate chains, and the presence of cellular receptors for the protein. The functionality to be retained is normally selected from procoagulant, proteolytic or receptor binding activity. The functional *in vivo* half-life and the serum half-life may be determined

10 by any suitable method known in the art.

The term "increased" as used about the functional *in vivo* half-life or serum half-life is used to indicate that the relevant half-life of the polypeptide variant is statistically significantly increased relative to that of a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, as determined under comparable conditions (typically determined in

15 an experimental animal, such as rats, rabbits, pigs or monkeys).

The term "AUC_{iv}" or "Area Under the Curve when administered intravenously" is used in its normal meaning, i.e. as the area under the activity in serum-time curve, where the polypeptide variant has been administered intravenously, in particular when administered intravenously in rats. Typically, the activity measured is the "clotting activity" as defined

20 above. Once the experimental activity-time points have been determined, the AUC_{iv} may conveniently be calculated by a computer program, such as GraphPad Prism 3.01.

It will be understood that in order to make a direct comparison between the AUC_{iv}-values of different molecules (e.g. between the variants of the invention and a reference molecule such as rhFVIIa or [P10Q+K32E]rhFVIIa) the same amount of activity should be

25 administered. Consequently, the AUC_{iv}-values are typically normalized (i.e. corrected for differences in the injected dose) and expressed as AUC_{iv}/dose administered.

The term "reduced sensitivity to proteolytic degradation" is primarily intended to mean that the polypeptide variant has reduced sensitivity to proteolytic degradation in comparison to hFVIIa, rhFVIIa or [P10Q+K32E]rhFVIIa as determined under comparable

30 conditions. Preferably, the proteolytic degradation is reduced by at least 10% (e.g. by 10-25% or by 10-50%), such as at least 25% (e.g. by 25-50%, by 25-75% or by 25-100%), more preferably by at least 35%, such as at least 50%, (e.g. by 50-75% or by 50-100%) even more preferably by at least 60%, such as by at least 75% (e.g. by 75-100%) or even at least 90%.

The term "renal clearance" is used in its normal meaning to indicate any clearance taking place by the kidneys, e.g. by glomerular filtration, tubular excretion or degradation in the tubular cells. Renal clearance depends on physical characteristics of the polypeptide, including size (diameter), hydrodynamic volume, symmetry, shape/rigidity, and charge.

5 Normally, a molecular weight of about 67 kDa is considered to be a cut-off-value for renal clearance. Renal clearance may be established by any suitable assay, e.g. an established *in vivo* assay. Typically, renal clearance is determined by administering a labelled (e.g. radiolabelled or fluorescencc labelled) polypeptide to a patient and measuring the label activity in urine collected from the patient. Reduced renal clearance is determined relative to
10 a corresponding reference polypeptide, e.g. rhFVIIa or [P10Q+K32E]rhFVIIa, under comparable conditions. Preferably, the renal clearance rate of the polypeptide variant is reduced by at least 50%, preferably by at least 75%, and most preferably by at least 90% compared to rhFVIIa or [P10Q+K32E]rhFVIIa.

15 The terms "tissue factor binding site", "active site region" and "ridge of the active site binding cleft" are defined with reference to Example 1.

The term "hydrophobic amino acid residue" includes the following amino acid residues: Isoleucine (I), leucine (L), methionine (M), valine (V), phenylalanine (F), tyrosine (Y) and tryptophan (W).

20 The term "negatively charged amino acid residue" includes the following amino acid residues: Aspartic acid (D) and glutamic acid (E).

The term "positively charged amino acid residue" includes the following amino acid residues: Lysine (K), arginine (R) and histidine (H).

Variants of the invention

25 The modifications performed in the Gla domain of the parent polypeptide preferably provide the resulting molecule with an increased phospholipid membrane binding affinity, an improved capability to activate FX to Fxa, and/or an increased clotting activity. The variants of the invention may also have a reduced tissue factor binding affinity and a reduced activity when bound to tissue factor.

30 Without being limited by any particular theory, it is presently believed that enhanced phospholipid membrane binding affinity results in a higher local concentration of the activated polypeptide variants in close proximity to the other coagulation factors, particularly FX. Thus, the rate of activation of FX to FXa will be higher, simply due to a higher molar

ratio of the activated FVII variant to FX. The increased activation rate of FX then results in a higher amount of active thrombin, and thus a higher rate of cross-linking of fibrin.

Consequently, it is contemplated that medical treatment with a polypeptide variant according to the invention may provide advantages over the currently available rhFVIIa compound (NovoSeven®), such as a lower dose, increased efficacy and/or faster action.

Further, it is believed that tissue factor-independent variants, i.e. variants that have a reduced activity when bound to tissue factor compared to wild-type human Factor VIIa, may offer certain safety advantages in terms of reduced risk of undesired blood clot formation (e.g. thrombosis or thromboembolism), in particular when used for treatment of acute uncontrolled bleeding events such as trauma.

Thus, in a highly preferred embodiment of the invention, the polypeptide variant, in its activated form and when compared to a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, has an increased FX activation activity, in particular when assayed in a tissue factor-independent assay, such as the "TF-independent Factor X Activation Assay" disclosed herein. More particularly, it is preferred that the ratio between the FX activation activity of the polypeptide variant, in its activated form, and the FX activation activity of a reference molecule is at least 1.25 when assayed in the "TF-independent Factor X Activation Assay" disclosed herein. More preferably, this ratio is at least 1.5, such as at least 1.75, e.g. at least 2, even more preferably at least 3, such as at least 4, most preferably at least 5.

When the reference molecule is rhFVIIa, the ratio between the FX activation activity of the polypeptide variant, in its activated form, and the FX activation activity of rhFVIIa is preferably at least about 5, typically at least about 10, when assayed in the "TF-independent Factor X Activation Assay" disclosed herein, such as at least about 15 or 20.

In another highly preferred embodiment of the invention, the variants of the invention possess an increased clotting activity (i.e. a reduced clotting time) as compared to rhFVIIa or [P10Q+K32E]rhFVIIa. In a preferred embodiment of the invention the ratio between the time to reach clot formation for the variant ($t_{variant}$) and the time to reach clot formation for rhFVIIa (t_{wt}) or [P10Q+K32E]rhFVIIa ($t_{P10Q+K32E}$) is at the most 0.9 when assayed in the "Whole Blood Assay" described herein. More preferably this ratio is at the most 0.75, such as 0.7, even more preferably the ratio is at the most 0.6, and most preferably the ratio is at the most 0.5.

One or more of the above-mentioned properties may be achieved by the modifications described herein.

Variants of the invention comprising a hydrophobic amino acid residue in position 34

As indicated above, the present invention relates in a first aspect to a FVII or FVIIa polypeptide variant having an amino acid sequence comprising 1-15 amino acid modifications relative to hFVII or hFVIIa (SEQ ID NO:1), wherein a hydrophobic amino acid residue has been introduced by substitution in position 34.

The hydrophobic amino acid residue to be introduced in position 34 may be selected from the group consisting of I, L, M, V, F, Y and W, preferably I, L and V, in particular L.

In a preferred embodiment, the variant further comprises an amino acid substitution in position 10, in particular P10Q, and/or an amino acid substitution in position 32, in particular K32E. In a particular preferred embodiment of the invention, the variant comprises substitutions in both of positions 10 and 32, such as P10Q+K32E.

Accordingly, in an interesting embodiment of the invention, the variant comprises the substitutions P10Q+K32E+A34L.

In a particular interesting embodiment of the invention, the variant further comprises an insertion of at least one (typically one) amino acid residue between position 3 and 4. It is preferred that the inserted amino acid residue is a hydrophobic amino acid residue. Most preferably the insertion is A3AY. Accordingly, in a particular interesting embodiment of the invention, the variant comprises the modifications A3AY+P10Q+K32E+A34L.

In addition to any of the above-mentioned modifications, the variant may comprise a further substitution in position 33. Preferably, a hydrophobic amino acid residue is introduced by substitution in position 33, in particular D33F.

The Gla domain may also contain modifications in other positions, in particular in positions 8, 11 and 28, such as R28F or R28E. On the other hand it should be understood that the Gla domain should not be modified to such an extent that the membrane binding properties are impaired. Accordingly, it is preferred that no modifications are made in the residues that become γ -carboxylated, i.e. it is preferred that no modifications are made in residues 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35. In a similar way, it is in general not preferred that non-polypeptide moieties, such as sugar moieties and/or PEG groups, are introduced in the Gla domain. Consequently, it is preferred that no modifications are made in the Gla domain that create an *in vivo* N-glycosylation site.

Finally, it will be understood that the modifications in the Gla domain discussed in this section may advantageously be combined with one or more modifications in positions located outside the Gla domain (see the sections entitled "Modifications outside the Gla domain" and "Other modifications outside the Gla domain" below).

Variants of the invention comprising an amino acid substitution in position 36

As indicated above, the invention relates in a second aspect to a FVII or FVIIa polypeptide variant having an amino acid sequence comprising 1-15 amino acid modifications relative to hFVII or hFVIIa (SEQ ID NO:1), wherein said amino acid sequence comprises an amino acid substitution in position 36.

Preferably, the amino acid residue to be introduced by substitution in position 36 is a negatively charged amino acid residue, e.g. R36E or R36D, in particular R36E.

In a preferred embodiment, the variant further comprises an amino acid substitution in position 10, in particular P10Q, and/or an amino acid substitution in position 32, in particular K32E. In a particular preferred embodiment of the invention, the variant comprises substitutions in both of positions 10 and 32, such as P10Q+K32E.

The variant of the invention may further contain a substitution in position 38. It is preferred that a negatively charged amino acid residue is introduced by substitution in position 38, e.g. K38E or K38D, in particular K38E.

Accordingly, interesting variants are those that comprise the following substitutions P10Q+K32E+R36E or P10Q+K32E+R36E-K38E.

In a particular interesting embodiment, the variant further comprises an amino acid substitution in position 34 (i.e. the resulting variant comprises substitutions in the following residues 10+32+34+36 or 10+32+34+36+38). Preferably, a negatively charged amino acid residue is introduced by substitution in position 34, e.g. A34E or A34D.

Specific examples of preferred variants are those that comprise the following substitutions P10Q+K32E+A34E+R36E or P10Q+K32E+A34D+R36E+K38E.

In an interesting embodiment of the invention, the variant further comprises an insertion of at least one (typically one) amino acid residue between position 3 and 4. It is preferred that the inserted amino acid residue is a hydrophobic amino acid residue. Most preferably the insertion is A3AY.

In addition to any of the above-mentioned modifications, the variant may comprise a further substitution in position 33. Preferably, a hydrophobic amino acid residue is introduced by substitution in position 33, in particular D33F.

The Gla domain may also contain modifications in other positions, in particular in positions 8, 11 and 28, such as R28F or R28E. On the other hand it should be understood that the Gla domain should not be modified to such an extent that the membrane binding properties are impaired. Accordingly, it is preferred that no modifications are made in the

residues that become γ -carboxylated, i.e. it is preferred that no modifications are made in residues 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35. In a similar way, it is in general not preferred that non-polypeptide moieties, such as sugar moieties and/or PEG groups, are introduced in the Gla domain. Consequently, it is preferred that no modifications are made in the Gla

5 domain that create an *in vivo* N-glycosylation site.

Finally, it will be understood that the modifications in the Gla domain discussed in this section may advantageously be combined with one or more modifications in positions located outside the Gla domain (see the sections entitled "*Modifications outside the Gla domain*" and "*Other modifications outside the Gla domain*" below).

10

Variants of the invention comprising amino acid substitutions in positions 74, 77 or 116

As indicated above, the present invention relates in a third aspect to a FVII or FVIIa polypeptide variant having an amino acid sequence comprising 3-15 amino acid modifications relative to hFVII or hFVIIa (SEQ ID NO:1), wherein said amino acid sequence 15 comprises an amino acid substitution in position 10, 32 and at least one further amino acid substitution in a position selected from the group consisting of position 74, 77 and 116.

In a preferred embodiment, the amino acid substitution in position 10 is P10Q and the amino acid substitution in position 32 is K32E

It is further preferred that the substitution in position 74, 77 or 116 is selected from 20 the group consisting of P74S, E77A and E116D.

In an interesting embodiment the variant further comprises an amino acid substitution in position 34. Preferably, a negatively charged amino acid residue is introduced by substitution in position 34, e.g. A34E or A34D, in particular A34E.

In another interesting embodiment of the invention the variant further comprises an 25 insertion of at least one (typically one) amino acid residue between position 3 and 4. It is preferred that the inserted amino acid residue is a hydrophobic amino acid residue. Most preferably the insertion is A3AY.

Thus, specific examples of interesting variants include variants comprising the following modifications A3AY+P10Q+K32E+E116D, A3AY+P10Q+K32E+E77A and 30 P10Q+K32E+A34E+P74S.

In addition to any of the above-mentioned modifications, the variant may comprise a further substitution in position 33. Preferably, a hydrophobic amino acid residue is introduced by substitution in position 33, in particular D33F.

The Gla domain may also contain modifications in other positions, in particular in positions 8, 11 and 28, such as R28F or R28E. As explained above, the Gla domain should not be modified to such an extent that the membrane binding properties are impaired, i.e. preferably no modifications are made in residues 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35, and it is preferred that an *in vivo* N-glycosylation site is not created in the Gla domain.

Finally, it will be understood that the modifications in the Gla domain discussed in this section may advantageously be combined with one or more modifications in positions located outside the Gla domain (see the sections entitled "*Modifications outside the Gla domain*" and "*Other modifications outside the Gla domain*" below).

10

Modifications outside the Gla domain

A circulating rhFVIIa half-life of 2.3 hours was reported in "Summary Basis for Approval for NovoSeven®", FDA reference number 96-0597. Relatively high doses and frequent administration are necessary to reach and sustain the desired therapeutic or prophylactic effect. As a consequence, adequate dose regulation is difficult to obtain and the need for frequent intravenous administration imposes restrictions on the patient's way of living.

A molecule with a longer circulation half-life and/or increased bioavailability (such as an increased Area Under the Curve as compared to rhFVIIa when administered intravenously) would decrease the number of necessary administrations. Given the current need for frequent injections and the potential for obtaining more optimal therapeutic FVIIa levels with concomitant enhanced therapeutic effect, there is a clear need for improved FVII- or FVIIa-like molecules.

Accordingly, a further object of the present invention is to provide improved FVII or FVII molecules (FVII or FVIIa variants) with an increased bioavailability (such as an increased Area Under the Curve as compared to a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, when administered intravenously) and which are capable of activating factor X to factor Xa (without binding to tissue factor) more efficiently than a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa (thereby being able to treat uncontrolled bleedings, such as a trauma, or chronic conditions such as hemophilia more efficiently).

Thus, interesting variants of the invention are those which, in their activated forms and when compared to a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, generate an increased Area Under the Curve when administered intravenously (AUC_{iv}). This

can conveniently be determined by intravenous administration in rats. More particularly, interesting variants of the present invention are those where the ratio between the AUC_{iv} of said variant, in its activated form, and the AUC_{iv} of a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, is at least 1.25, such as at least 1.5, e.g. at least 1.75, more preferably at least 2, such as at least 3, even more preferably at least 4, such as at least 5, in particular when administered (intravenously) in rats.

5 This effect will often correspond to an increased functional *in vivo* half-life and/or an increased serum half-life as compared to a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa. Accordingly, in another interesting embodiment of the invention, the 10 ratio between the functional *in vivo* half-life or the serum half-life for the variant, in its activated form, and the functional *in vivo* half-life or the serum half-life for a reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, is at least 1.25. More preferably, the ratio between the relevant half-life for the variant, in its activated form, and the relevant half-life for the reference molecule, such as rhFVIIa or [P10Q+K32E]rhFVIIa, is at least 1.5, such 15 as at least 1.75, e.g. at least 2, even more preferably at least 3, such as at least 4, e.g. at least 5.

One way to increase the circulation half-life of a protein is to ensure that renal clearance of the protein is reduced. This may be achieved by conjugating the protein to a chemical moiety which is capable of conferring reduced renal clearance to the protein, e.g. 20 polyethylene glycol (PEG).

Furthermore, attachment of a chemical moiety to the protein or substitution of amino acids exposed to proteolysis may effectively block a proteolytic enzyme from contact that otherwise leads to proteolytic degradation of the protein.

As indicated above, instability due to proteolytic degradation is a known problem in 25 current rhFVIIa treatment. Proteolytic degradation is thus a major obstacle for obtaining a preparation in solution as opposed to a lyophilized product. The advantage of obtaining a stable soluble preparation lies in easier handling for the patient, and, in the case of emergencies, quicker action, which potentially can become life saving. Attempts to prevent proteolytic degradation by site directed mutagenesis at major proteolytic sites have been 30 disclosed in WO 88/10295.

WO 01/58935 discloses a number of suitable modifications leading to an increase in AUC_{iv} , functional *in vivo* half-life and/or serum half-life. The variants disclosed in WO 01/58935 are the result of a generally new strategy for developing improved FVII or FVIIa

molecules, which may also be used for the parent FVII or FVIIa polypeptide of the present invention.

More specifically, by removing and/or introducing an amino acid residue comprising an attachment group for a non-polypeptide moiety in the parent FVII or FVIIa polypeptide it is possible to specifically adapt the polypeptide so as to make the molecule more susceptible to conjugation to a non-polypeptide moiety of choice, to optimize the conjugation pattern (e.g. to ensure an optimal distribution and number of non-polypeptide moieties on the surface of the FVII or FVIIa polypeptide variant and to ensure that only the attachment groups intended to be conjugated is present in the molecule) and thereby obtain a new conjugate molecule which has amidolytic activity and in addition one or more improved properties as compared to rhFVIIa

In interesting embodiments of the present invention more than one amino acid residue located outside the Gla domain is altered, e.g. the alteration embraces removal as well as introduction of amino acid residues comprising an attachment group for the non-polypeptide moiety of choice. In addition to the removal and/or introduction of amino acid residues the polypeptide variant may comprise other substitutions that are not related to introduction and/or removal of amino acid residues comprising an attachment group for the non-polypeptide moiety.

Also, the polypeptide variant may be attached to a serine proteinase inhibitor to inhibit the catalytic site of the polypeptide variant. Alternatively, one or more of the amino acid residues present in the catalytic site (S344, D242 and H193) may be mutated in order to render the resulting variant inactive. One example of such a mutation is S344A.

The amino acid residue comprising an attachment group for a non-polypeptide moiety, whether it be removed or introduced, is selected on the basis of the nature of the non-polypeptide moiety of choice and, in most instances, on the basis of the method in which conjugation between the polypeptide variant and the non-polypeptide moiety is to be achieved. For instance, when the non-polypeptide moiety is a polymer molecule such as a polyethylene glycol or polyalkylene oxide derived molecule, amino acid residues comprising an attachment group may be selected from the group consisting of lysine, cysteine, aspartic acid, glutamic acid, histidine, and tyrosine, preferably lysine, cysteine, aspartic acid and glutamic acid, more preferably lysine and cysteine, in particular cysteine.

Whenever an attachment group for a non-polypeptide moiety is to be introduced into or removed from the parent polypeptide, the position of the amino acid residue to be modified is preferably located at the surface of the parent FVII or FVIIa polypeptide, and

more preferably occupied by an amino acid residue which has at least 25% of its side chain exposed to the surface (as defined in Example 1 herein), preferably at least 50% of its side chain exposed to the surface (as defined in Example 1 herein). Such positions have been identified on the basis of an analysis of a 3D structure of the hFVII or hFVIIa molecule as described in WO 01/58935.

Furthermore, the position to be modified is preferably selected from a part of the FVII or FVIIa molecule that is located outside the tissue factor binding site, and/or outside the active site region, and/or outside the ridge of the active site binding cleft. These sites/regions are identified in Example 1 herein and in WO 01/58935.

10 In case of removal of an attachment group, the relevant amino acid residue comprising such group and occupying a position as defined above is preferably substituted with a different amino acid residue that does not comprise an attachment group for the non-polypeptide moiety in question. Normally, the amino acid residue to be removed is one to which conjugation is disadvantageous, e.g. an amino acid residue located at or near a 15 functional site of the polypeptide (since conjugation at such a site may result in inactivation or reduced activity of the resulting conjugate due to, e.g., impaired receptor recognition). In the present context the term "functional site" is intended to indicate one or more amino acid residues which is/are essential for or otherwise involved in the function or performance of FVII or FVIIa. Such amino acid residues are a part of the functional site. The functional site 20 may be determined by methods known in the art and is preferably identified by analysis of a structure of the FVIIa-tissue factor complex (See Banner et al., *Nature* 1996; 380:41-46).

In case of introduction of an attachment group, an amino acid residue comprising such group is introduced into the relevant position, preferably by substitution of the amino acid residue occupying such position.

25 The exact number of attachment groups present and available for conjugation in the FVII or FVIIa polypeptide is dependent on the effect desired to be achieved by the conjugation. The effect to be obtained is, e.g., dependent on the nature and degree of conjugation (e.g. the identity of the non-polypeptide moiety, the number of non-polypeptide moieties desirable or possible to conjugate to the polypeptide variant, where they should be 30 conjugated or where conjugation should be avoided, etc.).

The total number of amino acid residues to be modified outside the Gla domain in the parent FVII or FVIIa polypeptide (as compared to the amino acid sequence shown in SEQ ID NO:1) will typically not exceed 10. Preferably, the FVII or FVIIa variant comprises an amino acid sequence which differs in 1-10 amino acid residues from amino acid residues

46-406 shown in SEQ ID NO:1, typically in 1-8 or in 2-8 amino acid residues, e.g. in 1-5 or in 2-5 amino acid residues, such as in 1-4 or in 1-3 amino acid residues, e.g. in 1, 2 or 3 amino acid residues from amino acid residues 46-406 shown in SEQ ID NO:1.

Analogously, the polypeptide variant of the invention may contain 1-10 (additional) 5 non-polypeptide moieties, typically 1-8 or 2-8 (additional) non-polypeptide moieties, preferably 1-5 or 2-5 (additional) non-polypeptide moieties, such as 1-4 or 1-3 (additional) non-polypeptide moieties, e.g. 1, 2 or 3 (additional) non-polypeptide moieties. It will be understood that such additional non-polypeptide moieties are covalently attached to an attachment group located outside the Gla domain.

10

Polypeptide variants of the invention where the non-polypeptide moiety is a sugar moiety

In a preferred embodiment of the invention, an attachment group for a sugar moiety, such as a glycosylation site, in particular an *in vivo* glycosylation site, such as an *in vivo* N-glycosylation site, has been introduced and/or removed, preferably introduced, in a position 15 located outside the Gla domain.

When used in the present context, the term "naturally occurring glycosylation site" covers the glycosylation sites at positions N145, N322, S52 and S60. The term "naturally occurring *in vivo* O-glycosylation site" includes the positions S52 and S60, whereas the term "naturally occurring *in vivo* N-glycosylation site" includes positions N145 and N322.

20 Thus, in a very interesting embodiment of the invention, the non-polypeptide moiety is a sugar moiety and the introduced attachment group is a glycosylation site, preferably an *in vivo* glycosylation site, such as an *in vivo* O-glycosylation site or an *in vivo* N-glycosylation site, in particular an *in vivo* N-glycosylation site. Typically, 1-10 glycosylation sites, in particular *in vivo* N-glycosylation sites, have been introduced, preferably 1-8, 1-6, 1-4 or 1-3 25 glycosylation sites, in particular *in vivo* N-glycosylation sites, have been introduced in one or more positions located outside the Gla domain. For example 1, 2 or 3 glycosylation sites, in particular *in vivo* N-glycosylation sites, may have been introduced outside the Gla domain, preferably by substitution.

It will be understood that in order to prepare a polypeptide variant wherein the 30 polypeptide variant comprises one or more glycosylation sites, the polypeptide variant must be expressed in a host cell capable of attaching sugar (oligosaccharide) moieties at the glycosylation site(s) or alternatively subjected to *in vitro* glycosylation. Examples of glycosylating host cells are given in the section further below entitled "*Coupling to a sugar moiety*".

Examples of positions wherein the glycosylation sites, in particular *in vivo* N-glycosylation sites, may be introduced include amino acid residues having at least 25% of their side chain exposed to the surface (as defined in Example 1 herein), such as at least 50% of the side chain exposed to the surface. The position is preferably selected from a part of the molecule that is located outside the tissue factor binding site and/or the active site region and/or outside the ridge of the active site cleft, as defined in Example 1 herein. It should be understood that when the term "at least 25% (or at least 50%) of its side chain exposed to the surface" is used in connection with introduction of an *in vivo* N-glycosylation site this term refers to the surface accessibility of the amino acid side chain in the position where the sugar moiety is actually attached. In many cases it will be necessary to introduce a serine or a threonine residue in position +2 relative to the asparagine residue to which the sugar moiety is actually attached, and these positions where the serine or threonine residues are introduced are allowed to be buried, i.e. to have less than 25% of their side chains exposed to the surface.

Specific and preferred examples of such substitutions creating an *in vivo* N-glycosylation site include a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205S, I205T, V253N, T267N, T267N+S269T, S314N+K316S, S314N+K316T, R315N+V317S, R315N+V317T, K316N+G318S, K316N+G318T, G318N, D334N and combinations thereof. More preferably, the *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205T, V253N, T267N+S269T, S314N+K316T, R315N+V317T, K316N+G318T, G318N, D334N and combinations thereof. Even more preferably, the *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of T106N, A175T, I205T, V253N, T267N+S269T and combinations thereof, in particular one, two or three of T106N, I205T and V253N.

In one embodiment, only one *in vivo* N-glycosylation site has been introduced by substitution. In another embodiment, two or more (such as two) *in vivo* N-glycosylation sites have been introduced by substitution. Examples of preferred substitutions creating two *in vivo* N-glycosylation sites include substitutions selected from the group consisting of A51N+G58N, A51N+T106N, A51N+K109N, A51N+G124N, A51N+K143N+N145T, A51N+A175T, A51N+I205T, A51N+V253N, A51N+T267N+S269T, A51N+S314N+K316T, A51N+R315N+V317T, A51N+K316N+G318T, A51N+G318N, A51N+D334N, G58N+T106N, G58N+K109N, G58N+G124N, G58N+K143N+N145T,

G58N+A175T; G58N+I205T, G58N+V253N, G58N+T267N+S269T,
G58N+S314N+K316T, G58N+R315N+V317T, G58N+K316N+G318T, G58N+G318N,
G58N+D334N, T106N+K109N, T106N+G124N, T106N+K143N+N145T, T106N+A175T,
T106N+I205T, T106N+V253N, T106N+T267N+S269T, T106N+S314N+K316T,
5 T106N+R315N+V317T, T106N+K316N+G318T, T106N+G318N, T106N+D334N,
K109N+G124N, K109N+K143N+N145T, K109N+A175T, K109N+I205T, K109N+V253N,
K109N+T267N+S269T, K109N+S314N+K316T, K109N+R315N+V317T,
K109N+K316N+G318T, K109N+G318N, K109N+D334N, G124N+K143N+N145T,
G124N+A175T, G124N+I205T, G124N+V253N, G124N+T267N+S269T,
10 G124N+S314N+K316T, G124N+R315N+V317T, G124N+K316N+G318T, G124N+G318N,
G124N+D334N, K143N+N145T+A175T, K143N+N145T+I205T, K143N+N145T+V253N,
K143N+N145T+T267N+S269T, K143N+N145T+S314N+K316T,
K143N+N145T+R315N+V317T, K143N+N145T+K316N+G318T, K143N+N145T+G318N,
K143N+N145T+D334N, A175T+I205T, A175T+V253N, A175T+T267N+S269T,
15 A175T+S314N+K316T, A175T+R315N+V317T, A175T+K316N+G318T, A175T+G318N,
A175T+D334N, I205T+V253N, I205T+T267N+S269T, I205T+S314N+K316T,
I205T+R315N+V317T, I205T+K316N+G318T, I205T+G318N, I205T+D334N,
V253N+T267N+S269T, V253N+S314N+K316T, V253N+R315N+V317T,
V253N+K316N+G318T, V253N+G318N, V253N+D334N, T267N+S269T+S314N+K316T,
20 T267N+S269T+R315N+V317T, T267N+S269T+K316N+G318T, T267N+S269T+G318N,
T267N+S269T+D334N, S314N+K316T+R315N+V317T, S314N+K316T+G318N,
S314N+K316T+D334N, R315N+V317T+K316N+G318T, R315N+V317T+G318N,
R315N+V317T+D334N and G318N+D334N. More preferably, the substitutions are selected
from the group consisting of T106N+A175T, T106N+I205T, T106N+V253N,
25 T106N+T267N+S269T, A175T+I205T, A175T+V253N, A175T+T267N+S269T,
I205T+V253N, I205T+T267N+S269T and V253N+T267N+S269T, even more preferably
from the group consisting of T106N+I205T, T106N+V253N and I205T+V253N.

In a further embodiment, three or more (such as three) *in vivo* N-glycosylation sites have been introduced by substitution. Examples of preferred substitutions creating three *in vivo* N-glycosylation sites include substitutions selected from the group consisting of I205T+V253N+T267N+S269T and T106N+I205T+V253N.

As discussed above, it is preferred that the *in vivo* N-glycosylation site is introduced in a position which does not form part of the tissue factor binding site, the active site region or the ridge of the active site binding cleft as defined herein.

It will be understood that any of the modifications mentioned in the above sections may be combined with each other, in addition to being combined with the above-described substitutions in position 34 and/or 36, in particular A34E/L and/or R36E, and preferably in combination with the above-described substitutions in position 10 and/or 32, in particular 5 P10Q and/or K32E. Among the above-identified modifications for introduction of an *in vivo* N-glycosylation site, preferred modifications include one, two or three of T106N, I205T and V253N, in particular two of these modifications, i.e. T106N+I205T, T106N+V253N or I205T+V253N.

Thus, in one preferred embodiment of the invention the FVII or FVIIa variant 10 comprises the modifications P10Q+K32E+A34E+R36E+T106N+I205T.

In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34E+R36E+T106N+V253N.

In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34E+R36E+I205T+V253N.

15 In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+T106N+I205T.

In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+T106N+V253N.

20 In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+I205T+V253N.

In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+R36E+T106N+I205T.

In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+R36E+T106N+V253N.

25 In a further preferred embodiment the FVII or FVIIa variant comprises the modifications P10Q+K32E+A34L+R36E+I205T+V253N.

As is also explained above, any one or more of these modifications may in addition be combined with insertion of at least one amino acid residue, typically a single amino acid residue, between position 3 and 4, where the inserted residue is preferably a hydrophobic 30 amino acid residue. Most preferably the insertion is A3AY. Thus, in additional preferred embodiments of the invention the FVII or FVIIa variant comprises modifications selected from:

A3AY+P10Q+K32E+A34E+R36E+T106N+I205T;

A3AY+P10Q+K32E+A34E+R36E+T106N+V253N;

A3AY+P10Q+K32E+A34E+R36E+I205T+V253N;
A3AY+P10Q+K32E+A34L+T106N+I205T;
A3AY+P10Q+K32E+A34L+T106N+V253N;
A3AY+P10Q+K32E+A34L+I205T+V253N;
5 A3AY+P10Q+K32E+A34L+R36E+T106N+I205T;
A3AY+P10Q+K32E+A34L+R36E+T106N+V253N;
A3AY+P10Q+K32E+A34L+R36E+I205T+V253N.

Other modifications outside the Gla domain

10 In a further embodiment of the present invention, the FVII or FVIIa variant may, in addition to the modifications described in the sections above, also contain mutations which are already known to increase the intrinsic activity of the polypeptide, for example those described in WO 02/22776.

For example, the variant may comprise at least one modification in a position
15 selected from the group consisting of 157, 158, 296, 298, 305, 334, 336, 337 and 374. Examples of preferred substitutions include substitutions selected from the group consisting of V158D, E296D, M298Q, L305V and K337A. More preferably, said substitutions are selected from the group consisting of V158D+E296D+M298Q+L305V+K337A, V158D+E296D+M298Q+L305V, V158D+E296D+M298Q+K337A, V158D+E296D+M298Q+L305V, 20 V158D+E296D+M298Q, M298Q, L305V+K337A, L305V and K337A.

In a further embodiment of the present invention, the FVII or FVIIa variant may, in addition to the modifications described in the sections above, also contain other mutations, such as the substitution K341Q disclosed by Neuenschwander et al, *Biochemistry*, 1995; 34:8701-8707. Other possible additional substitutions include D196K, D196N, G237L, 25 G237GAA and combinations thereof.

Additional detailed information on conjugation of FVII and FVIIa variants to non-polypeptide moieties is found in WO 01/58935 and WO 03/093465, to which reference is made and which are incorporated herein by reference.

30 *Methods of preparing a conjugated variant of the invention*

In general, a conjugated variant according to the invention may be produced by culturing an appropriate host cell under conditions conducive for the expression of the variant polypeptide, and recovering the variant polypeptide, wherein a) the variant polypeptide comprises at least one N- or O-glycosylation site and the host cell is an eukaryotic host cell

capable of *in vivo* glycosylation, and/or b) the variant polypeptide is subjected to conjugation to a non-polypeptide moiety *in vitro*.

Conjugation to a polymer molecule

5 The polymer molecule to be coupled to the variant polypeptide may be any suitable polymer molecule, such as a natural or synthetic homo-polymer or hetero-polymer, typically with a molecular weight in the range of about 300-100,000 Da, such as about 500-20,000 Da, more preferably in the range of about 500-15,000 Da, even more preferably in the range of about 2-12 kDa, such as in the range of about 3-10 kDa. When the term "about" is used
10 herein in connection with a certain molecular weight, the word "about" indicates an approximate average molecular weight and reflects the fact that there will normally be a certain molecular weight distribution in a given polymer preparation.

Examples of homo-polymers include a polyol (i.e. poly-OH), a polyamine (i.e. poly-NH₂) and a polycarboxylic acid (i.e. poly-COOH). A hetero-polymer is a polymer
15 comprising different coupling groups, such as a hydroxyl group and an amine group.

Examples of suitable polymer molecules include polymer molecules selected from the group consisting of polyalkylene oxide (PAO), including polyalkylene glycol (PAG), such as polyethylene glycol (PEG) and polypropylene glycol (PPG), branched PEGs, poly-vinyl alcohol (PVA), poly-carboxylate, poly-(vinylpyrrolidone), polyethylene-co-maleic acid
20 anhydride, polystyrene-co-maleic acid anhydride, dextran, including carboxymethyl-dextran, or any other biopolymer suitable for reducing immunogenicity and/or increasing functional *in vivo* half-life and/or serum half-life. Another example of a polymer molecule is human albumin or another abundant plasma protein. Generally, polyalkylene glycol-derived polymers are biocompatible, non-toxic, non-antigenic, non-immunogenic, water soluble, and
25 are easily excreted from living organisms.

PEG is the preferred polymer molecule, since it has only few reactive groups capable of cross-linking compared to, e.g., polysaccharides such as dextran. In particular, monofunctional PEG, e.g. methoxypolyethylene glycol (mPEG), is of interest since its coupling chemistry is relatively simple (only one reactive group is available for conjugating
30 with attachment groups on the polypeptide). Consequently, as the risk of cross-linking is eliminated, the resulting conjugated variants are more homogeneous and the reaction of the polymer molecules with the variant polypeptide is easier to control.

To effect covalent attachment of the polymer molecule(s) to the variant polypeptide, the hydroxyl end groups of the polymer molecule must be provided in activated form, i.e.

with reactive functional groups (examples of which include primary amino groups, hydrazide (HZ), thiol, succinate (SUC), succinimidyl succinate (SS), succinimidyl succinamide (SSA), succinimidyl propionate (SPA), succinimidyl butyrate (SBA), succinimidyl carboxymethylate (SCM), benzotriazole carbonate (BTC), N-hydroxysuccinimide (NHS), 5 aldehyde, nitrophenylcarbonate (NPC), and tresylate (TRES)). Suitable activated polymer molecules are commercially available, e.g. from Nektar Therapeutics, Huntsville, AL, USA, or from PolyMASC Pharmaceuticals plc, UK.

Specific examples of activated linear or branched polymer molecules for use in the present invention are described in the Nektar Molecule Engineering Catalog 2003 (Nektar 10 Therapeutics), incorporated herein by reference.

Specific examples of activated PEG polymers include the following linear PEGs: NHS-PEG (e.g. SPA-PEG, SSPA-PEG, SBA-PEG, SS-PEG, SSA-PEG, SC-PEG, SG-PEG, and SCM-PEG), and NOR-PEG, BTC-PEG, EPOX-PEG, NCO-PEG, NPC-PEG, CDI-PEG, ALD-PEG, TRES-PEG, VS-PEG, IODO-PEG, and MAL-PEG, and branched PEGs such as 15 PEG2-NHS and those disclosed in US 5,932,462 and US 5,643,575, both of which are incorporated herein by reference. Additional publications disclosing useful polymer molecules, PEGylation chemistries and conjugation methods are listed in WO 01/58935 and WO 03/093465.

Specific examples of activated PEG polymers particularly preferred for coupling to 20 cysteine residues, include the following linear PEGs: vinylsulfone-PEG (VS-PEG), preferably vinylsulfone-mPEG (VS-mPEG); maleimide-PEG (MAL-PEG), preferably maleimide-mPEG (MAL-mPEG) and orthopyridyl-disulfide-PEG (OPSS-PEG), preferably orthopyridyl-disulfide-mPEG (OPSS-mPEG). Typically, such PEG or mPEG polymers will have a size of about 5 kDa, about 10 kD, about 12 kDa or about 20 kDa.

25 The skilled person will be aware that the activation method and/or conjugation chemistry to be used depends on the attachment group(s) of the variant polypeptide (examples of which are given further above), as well as the functional groups of the polymer (e.g. being amine, hydroxyl, carboxyl, aldehyde, sulphydryl, succinimidyl, maleimide, vinylsulfone or haloacetate). The PEGylation may be directed towards conjugation to all 30 available attachment groups on the variant polypeptide (i.e. such attachment groups that are exposed at the surface of the polypeptide) or may be directed towards one or more specific attachment groups, e.g. the N-terminal amino group as described in US 5,985,265 or to cysteine residues. Furthermore, the conjugation may be achieved in one step or in a stepwise manner (e.g. as described in WO 99/55377).

For PEGylation to cysteine residues (see above) the FVII or FVIIa variant is usually treated with a reducing agent, such as dithiothreitol (DDT) prior to PEGylation. The reducing agent is subsequently removed by any conventional method, such as by desalting. Conjugation of PEG to a cysteine residue typically takes place in a suitable buffer at pH 6-9 at temperatures 5 varying from 4°C to 25°C for periods up to 16 hours.

It will be understood that the PEGylation is designed so as to produce the optimal molecule with respect to the number of PEG molecules attached, the size and form of such molecules (e.g. whether they are linear or branched), and the attachment site(s) in the variant polypeptide. The molecular weight of the polymer to be used may e.g. be chosen on the basis 10 of the desired effect to be achieved.

In connection with conjugation to only a single attachment group on the protein (e.g. the N-terminal amino group), it may be advantageous that the polymer molecule, which may be linear or branched, has a high molecular weight, preferably about 10-25 kDa, such as about 15-25 kDa, e.g. about 20 kDa.

15 Normally, the polymer conjugation is performed under conditions aimed at reacting as many of the available polymer attachment groups as possible with polymer molecules. This is achieved by means of a suitable molar excess of the polymer relative to the polypeptide. Typically, the molar ratios of activated polymer molecules to polypeptide are up to about 1000-1, such as up to about 200-1, or up to about 100-1. In some cases the ratio 20 may be somewhat lower, however, such as up to about 50-1, 10-1, 5-1, 2-1 or 1-1 in order to obtain optimal reaction.

It is also contemplated according to the invention to couple the polymer molecules to the polypeptide through a linker. Suitable linkers are well known to the skilled person; see also WO 01/58935.

25 Subsequent to the conjugation, residual activated polymer molecules are blocked according to methods known in the art, e.g. by addition of primary amine to the reaction mixture, and the resulting inactivated polymer molecules are removed by a suitable method.

It will be understood that depending on the circumstances, e.g. the amino acid sequence of the variant polypeptide, the nature of the activated PEG compound being used 30 and the specific PEGylation conditions, including the molar ratio of PEG to polypeptide, varying degrees of PEGylation may be obtained, with a higher degree of PEGylation generally being obtained with a higher ratio of PEG to variant polypeptide. The PEGylated variant polypeptides resulting from any given PEGylation process will, however, normally

comprise a stochastic distribution of conjugated polypeptide variants having slightly different degrees of PEGylation.

Coupling to a sugar moiety

5 In order to achieve *in vivo* glycosylation of a FVII molecule comprising one or more glycosylation sites the nucleotide sequence encoding the variant polypeptide must be inserted in a glycosylating, eucaryotic expression host. The expression host cell may be selected from fungal (filamentous fungal or yeast), insect or animal cells or from transgenic plant cells. In one embodiment the host cell is a mammalian cell, such as a CHO cell, BHK or HEK, e.g. 10 HEK 293, cell, or an insect cell, such as an SF9 cell, or a yeast cell, e.g. *Saccharomyces cerevisiae* or *Pichia pastoris*, or any of the host cells mentioned hereinafter.

Covalent *in vitro* coupling of sugar moieties (such as dextran) to amino acid residues of the variant polypeptide may also be used, e.g. as described, for example in WO 87/05330 and in Aplin et al., *CRC Crit Rev. Biochem.*, pp. 259-306, 1981. See also WO 03/093465 for 15 further information on *in vitro* glycosylation of variants of FVII or FVIIa.

Attachment of serine protease inhibitor

Attachment of a serine protease inhibitor can be performed in accordance with the method described in WO 96/12800.

20

Methods of preparing a polypeptide variant of the invention

The polypeptide variant of the present invention, optionally in glycosylated form, may be produced by any suitable method known in the art. Such methods include constructing a nucleotide sequence encoding the polypeptide variant and expressing the 25 sequence in a suitable transformed or transfected host. Preferably, the host cell is a gamma-carboxylating host cell such as a mammalian cell. However, polypeptide variants of the invention may be produced, albeit less efficiently, by chemical synthesis or a combination of chemical synthesis or a combination of chemical synthesis and recombinant DNA technology.

30 A nucleotide sequence encoding a polypeptide of the invention may be constructed by isolating or synthesizing a nucleotide sequence encoding the parent FVII, such as hFVII with the amino acid sequence shown in SEQ ID NO:1 and then changing the nucleotide sequence so as to effect introduction (i.e. insertion or substitution) or removal (i.e. deletion or substitution) of the relevant amino acid residue(s).

The nucleotide sequence is conveniently modified by site-directed mutagenesis in accordance with conventional methods. Alternatively, the nucleotide sequence is prepared by chemical synthesis, e.g. by using an oligonucleotide synthesizer, wherein oligonucleotides are designed based on the amino acid sequence of the desired polypeptide, and preferably 5 selecting those codons that are favored in the host cell in which the recombinant polypeptide will be produced. For example, several small oligonucleotides coding for portions of the desired polypeptide may be synthesized and assembled by PCR (polymerase chain reaction), ligation or ligation chain reaction (LCR) (Barany, *Proc Natl Acad Sci USA* 88:189-193, 1991). The individual oligonucleotides typically contain 5' or 3' overhangs for 10 complementary assembly.

Once assembled (by synthesis, site-directed mutagenesis or another method), the nucleotide sequence encoding the polypeptide is inserted into a recombinant vector and operably linked to control sequences necessary for expression of the FVII in the desired transformed host cell.

15 Persons skilled in the art will be capable of selecting suitable vectors, expression control sequences and hosts for expressing the polypeptide. The recombinant vector may be an autonomously replicating vector, i.e. a vector, which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector is one which, when introduced into a host cell, is integrated into the 20 host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector, in which the nucleotide sequence encoding the polypeptide variant of the invention is operably linked to additional segments required for transcription of the nucleotide sequence. The vector is typically derived from 25 plasmid or viral DNA. A number of suitable expression vectors for expression in the host cells mentioned herein are commercially available or described in the literature. Detailed information on suitable vectors for expressing FVII may be found in WO 01/58935, incorporated by reference.

The term "control sequences" is defined herein to include all components which are 30 necessary or advantageous for the expression of the polypeptide variant of the invention. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide variant. Such control sequences include, but are not limited to, a leader sequence, polyadenylation sequence, propeptide sequence, promoter, enhancer or upstream activating

sequence, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter.

A wide variety of expression control sequences may be used in the present invention, e.g. any of the control sequences disclosed in WO 01/58935, incorporated by reference.

The nucleotide sequence of the invention encoding a polypeptide variant, whether prepared by site-directed mutagenesis, synthesis, PCR or other methods, may optionally include a nucleotide sequence that encodes a signal peptide. The signal peptide is present when the polypeptide variant is to be secreted from the cells in which it is expressed. Such signal peptide, if present, should be one recognized by the cell chosen for expression of the polypeptide variant. The signal peptide may be homologous (i.e. normally associated with hFVII) or heterologous (i.e. originating from another source than hFVII) to the polypeptide or may be homologous or heterologous to the host cell, i.e. a signal peptide normally expressed from the host cell or one which is not normally expressed from the host cell. For further information on suitable signal peptides, see WO 01/58935.

Any suitable host may be used to produce the polypeptide variant, including bacteria (although not particularly preferred), fungi (including yeasts), plant, insect, mammal, or other appropriate animal cells or cell lines, as well as transgenic animals or plants. Mammalian cells are preferred. Examples of bacterial host cells include gram-positive bacteria such as strains of *Bacillus*, e.g. *B. brevis* or *B. subtilis*, *Pseudomonas* or *Streptomyces*, or gram-negative bacteria, such as strains of *E. coli*. Examples of suitable filamentous fungal host cells include strains of *Aspergillus*, e.g. *A. oryzae*, *A. niger*, or *A. nidulans*, *Fusarium* or *Trichoderma*. Examples of suitable yeast host cells include strains of *Saccharomyces*, e.g. *S. cerevisiae*, *Schizosaccharomyces*, *Klyveromyces*, *Pichia*, such as *P. pastoris* or *P. methanolica*, *Hansenula*, such as *H. Polymorpha* or *Yarrowia*. Examples of suitable insect host cells include a *Lepidoptera* cell line, such as *Spodoptera frugiperda* (Sf9 or Sf21) or *Trichoplusia ni* cells (High Five) (US 5,077,214). Examples of suitable mammalian host cells include Chinese hamster ovary (CHO) cell lines, (e.g. CHO-K1; ATCC CCL-61), Green Monkey cell lines (COS) (e.g. COS 1 (ATCC CRL-1650), COS 7 (ATCC CRL-1651)); mouse cells (e.g. NS/O), Baby Hamster Kidney (BHK) cell lines (e.g. ATCC CRL-1632 or ATCC CCL-10), and human cells (e.g. HEK 293 (ATCC CRL-1573)). Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Rockville, Maryland. Also, mammalian cells, such as a CHO cell,

may be modified to express sialyltransferase, e.g. 1,6-sialyltransferase, e.g. as described in US 5,047,335, in order to provide improved glycosylation of the polypeptide variant.

In order to increase secretion it may be of particular interest to produce the polypeptide variant of the invention together with an endoprotease, in particular a PACE 5 (paired basic amino acid converting enzyme) (e.g. as described in US 5,986,079), such as a Kex2 endoprotease (e.g. as described in WO 00/28065).

Methods for introducing exogenous DNA into the above cell types, as well as other information regarding expression, production and purification of FVII variants, is found in WO 01/58935, incorporated herein by reference.

10

Pharmaceutical composition of the invention and its use

In a further aspect, the present invention relates to a composition, in particular to a pharmaceutical composition, comprising a polypeptide variant of the invention and a pharmaceutically acceptable carrier or excipient.

15

The polypeptide variant or the pharmaceutical composition according to the invention may be used as a medicament.

Due to the improved properties mentioned above, the polypeptide variants of the invention, or the pharmaceutical composition of the invention, are particular useful for the treatment of uncontrollable bleeding events in trauma patients, thrombocytopenic patients, 20 patients in anticoagulant treatment, and cirrhosis patients with variceal bleeding, or other upper gastrointestinal bleedings, in patients undergoing orthotopic liver transplantation or liver resection (allowing for transfusion free surgery), or in hemophilia patients.

Trauma is defined as an injury to living tissue caused by an extrinsic agent. It is the 4th leading cause of death in the US and places a large financial burden on the economy.

25

Trauma is classified as either blunt or penetrative. Blunt trauma results in internal compression, organ damage and internal haemorrhage whereas penetrative trauma (as the consequence of an agent penetrating the body and destroying tissue, vessels and organs) results in external haemorrhage.

30

Trauma may be caused by numerous events, e.g. traffic accidents, gunshot wounds, falls, machinery accidents, and stab wounds.

Cirrhosis of the liver may be caused by direct liver injury, including chronic alcoholism, chronic viral hepatitis (types B, C, and D), and autoimmune hepatitis as well as by indirect injury by way of bile duct damage, including primary biliary cirrhosis, primary sclerosing cholangitis and biliary atresia. Less common causes of cirrhosis include direct

liver injury from inherited disease such as cystic fibrosis, alpha-1-antitrypsin deficiency, hemochromatosis, Wilson's disease, galactosemia, and glycogen storage disease.

Transplantation is the key intervention for treating late stage cirrhotic patients

Thus, in a further aspect the present invention relates to a polypeptide variant of the invention for the manufacture of a medicament for the treatment of diseases or disorder wherein clot formation is desirable. A still further aspect of the present invention relates to a method for treating a mammal having a disease or disorder wherein clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of the polypeptide variant or the pharmaceutical composition of the invention.

Examples of diseases/disorders wherein increased clot formation is desirable include, but is not limited to, hemorrhages, including brain hemorrhages, as well as patient with severe uncontrolled bleedings, such as trauma. Further examples include patients undergoing living transplantations, patients undergoing resection and patients with variceal bleeding. Another widespread disease/disorder in which it is contemplated that the polypeptides of the invention will be useful for increased clot formation is hemophilia, e.g. von Willebrand disease, hemophilia A, hemophilia B or hemophilia C.

The polypeptide variants of the invention are administered to patients in a therapeutically effective dose, normally one approximately paralleling that employed in therapy with rFVII such as NovoSeven®, or at lower dosage. By "therapeutically effective dose" herein is meant a dose that is sufficient to produce the desired effects in relation to the condition for which it is administered. The exact dose will depend on the circumstances, and will be ascertainable by one skilled in the art using known techniques. Normally, the dose should be capable of preventing or lessening the severity or spread of the condition or indication being treated. It will be apparent to those of skill in the art that an effective amount of a polypeptide variant or composition of the invention depends, *inter alia*, upon the disease, the dose, the administration schedule, whether the polypeptide variant or composition is administered alone or in conjunction with other therapeutic agents, the plasma half-life of the compositions, and the general health of the patient.

The polypeptide variant of the invention is preferably administered in a composition including a pharmaceutically acceptable carrier or excipient. "Pharmaceutically acceptable" means a carrier or excipient that does not cause any untoward effects in patients to whom it is administered. Such pharmaceutically acceptable carriers and excipients as well as suitable pharmaceutical formulation methods are well known in the art (see, for example, Remington's Pharmaceutical Sciences, 18th edition, A. R. Gennaro, Ed., Mack Publishing

Company [1990]; Pharmaceutical Formulation Development of Peptides and Proteins, S. Frokjaer and L. Hovgaard, Eds., Taylor & Francis [2000]; and Handbook of Pharmaceutical Excipients, 3rd edition, A. Kibbe, Ed., Pharmaceutical Press [2000]).

The polypeptide variant of the invention can be used "as is" and/or in a salt form thereof. Suitable salts include, but are not limited to, salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium and magnesium, as well as e.g. zinc salts. These salts or complexes may be present as a crystalline and/or amorphous structure.

The pharmaceutical composition of the invention may be administered alone or in conjunction with other therapeutic agents. These agents may be incorporated as part of the same pharmaceutical composition or may be administered separately from the polypeptide variant of the invention, either concurrently or in accordance with another treatment schedule. In addition, the polypeptide variant or pharmaceutical composition of the invention may be used as an adjuvant to other therapies.

A "patient" for the purposes of the present invention includes both humans and other mammals. Thus, the methods are applicable to both human therapy and veterinary applications, in particular to human therapy.

The pharmaceutical composition comprising the polypeptide variant of the invention may be formulated in a variety of forms, e.g. as a liquid, gel, lyophilized, or as a compressed solid. The preferred form will depend upon the particular indication being treated and will be apparent to one skilled in the art.

In particular, the pharmaceutical composition comprising the polypeptide variant of the invention may be formulated in lyophilised or stable soluble form. The polypeptide variant may be lyophilised by a variety of procedures known in the art. The polypeptide variant may be in a stable soluble form by the removal or shielding of proteolytic degradation sites as described herein. The advantage of obtaining a stable soluble preparation lies in easier handling for the patient and, in the case of emergencies, quicker action, which potentially can become life saving. The preferred form will depend upon the particular indication being treated and will be apparent to one of skill in the art.

The administration of the formulations of the present invention can be performed in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intracerebrally, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, intraocularly, or in any other acceptable manner. The formulations can be administered continuously by infusion, although bolus injection is

acceptable, using techniques well known in the art, such as pumps or implantation. In some instances the formulations may be directly applied as a solution or spray.

Parentals

5 A preferred example of a pharmaceutical composition is a solution, in particular an aqueous solution, designed for parenteral administration. Although in many cases pharmaceutical solution formulations are provided in liquid form, appropriate for immediate use, such parenteral formulations may also be provided in frozen or in lyophilized form. In the former case, the composition must be thawed prior to use. The latter form is often used to
10 enhance the stability of the active compound contained in the composition under a wider variety of storage conditions, as it is recognized by those skilled in the art that lyophilized preparations are generally more stable than their liquid counterparts. Such lyophilized preparations are reconstituted prior to use by the addition of one or more suitable pharmaceutically acceptable diluents such as sterile water for injection or sterile
15 physiological saline solution.

In case of parenterals, they are prepared for storage as lyophilized formulations or aqueous solutions by mixing, as appropriate, the polypeptide variant having the desired degree of purity with one or more pharmaceutically acceptable carriers, excipients or stabilizers typically employed in the art (all of which are termed "excipients"), for example
20 buffering agents, stabilizing agents, preservatives, isotonifiers, non-ionic surfactants or detergents, antioxidants, and/or other miscellaneous additives such as bulking agents or fillers, chelating agents, antioxidants and cosolvents.

Detailed information on parental formulations suitable for administration of FVII variants, as well as sustained release preparations, is found in WO 01/58935 and WO
25 03/093465, incorporated herein by reference.

The invention is further described by the following non-limiting examples.

MATERIALS AND METHODS

30

Active Site Region

The active site region is defined as any residues having at least one atom within 10 Å of any atom in the catalytic triad (residues H193, D242, S344).

Measurement of Reduced Sensitivity to Proteolytic Degradation

Proteolytic degradation can be measured using the assay described in US 5,580,560, Example 5, where proteolysis is autoproteolysis.

Furthermore, reduced proteolysis can be tested in an *in vivo* model using

5 radiolabelled samples and comparing proteolysis of rhFVIIa and the polypeptide variant of the invention by withdrawing blood samples and subjecting these to SDS-PAGE and autoradiography.

Irrespective of the assay used for determining proteolytic degradation, "reduced proteolytic degradation" is intended to mean a measurable reduction in cleavage compared to 10 that obtained by rhFVIIa as measured by gel scanning of Coomassie stained SDS-PAGE gels, HPLC or as measured by conserved catalytic activity in comparison to wild type using the tissue factor independent activity assay described below.

Determination of the Molecular Weight of Polypeptide Variants

15 The molecular weight of polypeptide variants is determined by either SDS-PAGE, gel filtration, Western Blots, matrix assisted laser desorption mass spectrometry or equilibrium centrifugation, e.g. SDS-PAGE according to Laemmli, U.K., *Nature* Vol 227 (1970), pp. 680-85.

20 Determination of Phospholipid Membrane Binding Affinity

Phospholipid membrane binding affinity may be determined as described in Nelsestuen et al., *Biochemistry*, 1977; 30:10819-10824 or as described in Example 1 in US 6,017,882.

25 TF-independent Factor X Activation Assay

This assay has been described in detail on page 39826 in Nelsestuen et al., *J Biol Chem*, 2001; 276:39825-39831.

Briefly, the molecule to be assayed (either hFVIIa; rhFVIIa or the polypeptide variant of the invention in its activated form) is mixed with a source of phospholipid (preferably phosphatidylcholine and phosphatidylserine in a ratio of 8:2) and relipidated Factor X in Tris 30 buffer containing BSA. After a specified incubation time the reaction is stopped by addition of excess EDTA. The concentration of factor Xa is then measured from absorbance change at 405 nm after addition of a chromogenic substrate (S-2222, Chromogenix). After correction for background the tissue factor independent activity of rhFVIIa (a_{wt}) is determined as the absorbance change after 10 minutes and the tissue factor independent activity of the polypeptide

variant of the invention ($a_{variant}$) is also determined as the absorbance change after 10 minutes. The ratio between the activity of the polypeptide variant, in its activated form, and the activity of rhFVIIa is defined as $a_{variant}/a_{wt}$.

5 *Clotting Assay*

The clotting activity of the FVIIa and variants thereof were measured in one-stage assays and the clotting times were recorded on a Thrombotrack IV coagulometer (Medinor). Factor VII-depleted human plasma (American Diagnostica) was reconstituted and equilibrated at room temperature for 15-20 minutes. 50 microliters of plasma was then 10 transferred to the coagulometer cups.

FVIIa and variants thereof were diluted in Glyoxaline Buffer (5.7 mM barbiturate, 4.3 mM sodium citrate, 117 mM NaCl, 1 mg/ml BSA, pH 7.35). The samples were added to the cup in 50 μ l and incubated at 37°C for 2 minutes.

15 Thromboplastin (Medinor) was reconstituted with water and CaCl₂ was added to a final concentration of 4.5 mM. The reaction was initiated by adding 100 μ l thromboplastin.

To measure the clotting activity in the absence of TF the same assay was used without addition of thromboplastin. Data was analysed using PRISM software.

Whole Blood Assay

20 The clotting activity of FVIIa and variants thereof were measured in one-stage assays and the clotting times were recorded on a Thrombotrack IV coagulometer (Medinor). 100 μ l of FVIIa or variants thereof were diluted in a buffer containing 10 mM glycylglycine, 50 mM NaCl, 37.5 mM CaCl₂, pH 7.35 and transferred to the reaction cup. The clotting reaction was initiated by addition of 50 μ l blood containing 10% 0.13 M tri-sodium citrate as 25 anticoagulant. Data was analysed using Excel or PRISM software.

Amidolytic Assay

The ability of the variants to cleave small peptide substrates can be measured using the chromogenic substrate S-2288 (D-Ile-Pro-Arg-p-nitroanilide). FVIIa is diluted to about 30 10-90 nM in assay buffer (50 mM Na-Hepes pH 7.5, 150 mM NaCl, 5 mM CaCl₂, 0.1% BSA, 1U/ml Heparin). Furthermore, soluble TF (sTF) is diluted to 50-450 nM in assay buffer. 120 μ l of assay buffer is mixed with 20 μ l of the FVIIa sample and 20 μ l sTF. After 5 min incubation at room temperature with gentle shaking, followed by 10 min incubation at

37°C, the reaction is started by addition of the S-2288 substrate to 1 mM and the absorption at 405 nm is determined at several time points.

ELISA Assay

5 FVII/FVIIa (or variant) concentrations are determined by ELISA. Wells of a microtiter plate are coated with an antibody directed against the protease domain using a solution of 2 µg/ml in PBS (100 µl per well). After overnight coating at R.T. (room temperature), the wells are washed 4 times with THT buffer (100 mM NaCl, 50 mM Tris-HCl pH 7.2 0.05% Tween-20). Subsequently, 200 µl of 1% Casein (diluted from 2.5% stock 10 using 100 mM NaCl, 50 mM Tris-HCl pH 7.2) is added per well for blocking. After 1 hr incubation at R.T., the wells are emptied, and 100 µl of sample (optionally diluted in dilution buffer (THT + 0.1% Casein)) is added. After another incubation of 1 hr at room temperature, the wells are washed 4 times with THT buffer, and 100 µl of a biotin-labelled antibody directed against the EGF-like domain (1 µg/ml) is added. After another 1 hr incubation at 15 R.T., followed by 4 more washes with THT buffer, 100 µl of streptavidin-horse radish peroxidase (DAKO A/S, Glostrup, Denmark, 1/10000 diluted) is added. After another 1 hr incubation at R.T., followed by 4 more washes with THT buffer, 100 µl of TMB (3,3',5,5'-tetramethylbenzidine, Kem-en-Tech A/S, Denmark) is added. After 30 min incubation at R.T. in the dark, 100 µl of 1 M H₂SO₄ is added and OD_{450nm} is determined. A standard curve is 20 prepared using rhFVIIa (NovoSeven®).

Alternatively, FVII/FVIIa or variants may be quantified through the Gla domain rather than through the protease domain. In this ELISA set-up, wells are coated overnight with an antibody directed against the EGF-like domain and for detection, a calcium-dependent biotin-labelled monoclonal anti-Gla domain antibody is used (2 µg/ml, 100 µl per 25 well). In this set-up, 5 mM CaCl₂ is added to the THT and dilution buffers.

Thrombogram Assay

The effect of hFVIIa, rhFVIIa or FVIIa variants on thrombin generation in human plasma is tested in a modified version of the assay described on page 589 in Hemker et al. 30 (2000) *Thromb Haemost* 83:589-91. Briefly, the molecule to be assayed (either hFVIIa, rhFVIIa or a variant) is mixed with FVII-depleted platelet poor plasma (PPP) containing either relipidated recombinant tissue factor (such as Innovin from Dade Behring) or phospholipid (phosphatidylcholine and phosphatidylserine in a ratio of 8:2, or phosphatidylcholine, phosphatidylserine and phosphatidylethanol in a ratio of 4:2:4).

The reaction is started by addition of a fluoresgenic thrombin substrate and calcium chloride. The fluorescence is measured continuously and the thrombin amidolytic activity is determined by calculating the slope of the fluorescence curve (the increase in fluorescence over time). In this way the time until maximum thrombin amidolytic activity is obtained (T_{max}), and 5 the thrombin generation rate (maximal increase in thrombin activity) and the total thrombin work (area under the curve (AUC)) can be calculated.

Frozen citrated FVII-depleted plasma is thawed in the presence of corn trypsin inhibitor (100 μ g/ml serum) to inhibit the contact pathway of coagulation. To each well of a 96-well microtiter plate is added 80 μ l plasma and 20 μ l buffer containing rhFVII or variant to 10 be tested in final concentrations of between 0.1 and 100 nM. Recombinant human tissue factor (rTF) is added in 5 μ l assay buffer to a final concentration of 1 pM. The assay buffer consists of 20 mM Hepes, 150 mM NaCl and 60 mg/ml BSA in distilled water. The reaction is started by adding 20 μ l of the substrate solution containing 0.1 M calcium chloride. The assay plate and reagents are pre-warmed to 37°C and the reaction takes place at this temperature. The 15 fluorimeter used is a BMG Fluorimeter with an excitation filter at 390 nm and an emission filter at 460 nm. The fluorescence is measured in each well of 96-well clear bottom plates at 20-40 second intervals over 30-180 minutes. Data are analyzed using PRISM Software.

Tissue factor binding surface plasmon resonance assay (Biacore Assay)

20 Surface plasmon resonance analysis was used to determine the relative binding of wild-type Factor VIIa and variants thereof to soluble tissue factor. Recombinant soluble tissue factor that contains the extracellular domain was coupled to 270 response units on a Biacore CM5 chip using NHS/EDC coupling. Soluble tissue factor was coupled at a pH of 4.5 to enable interaction with the chip surface.

25 In this assay, tissue factor binding of factor VII protein was compared at a single concentration of FVIIa or variant to allow a relative comparison of the variants to wild-type. This concentration was determined by means of a standard curve of wild type FVIIa that was flowed over the chip in concentrations between 75 and 0 μ g/ml. FVIIa was removed by addition of 10 mM EDTA. It was determined in this manner that a concentration of 15 μ g/ml 30 gave binding in the linear range. Variants of FVIIa were then flowed over the chip at 15 μ g/ml to determine the relative binding strength of FVIIa or variants to tissue factor.

Example 1

The X-ray structure of hFVIIa in complex with soluble tissue factor by Banner et al.,
5 *J Mol Biol*, 1996; 285:2089 is used for this example. For further information on the
calculations in this example, see WO 01/58935.

Surface Exposure

Performing fractional ASA calculations resulted in the following residues being
10 determined to have more than 25% of their side chain exposed to the surface: A1, N2, A3,
F4, L5, E6, E7, L8, R9, P10, S12, L13, E14, E16, K18, E19, E20, Q21, S23, F24, E25, E26,
R28, E29, F31, K32, D33, A34, E35, R36, K38, L39, W41, I42, S43, S45, G47, D48, Q49,
A51, S52, S53, Q56, G58, S60, K62, D63, Q64, L65, Q66, S67, I69, F71, L73, P74, A75,
E77, G78, R79, E82, T83, H84, K85, D86, D87, Q88, L89, I90, V92, N93, E94, G97, E99,
15 S103, D104, H105, T106, G107, T108, K109, S111, R113, E116, G117, S119, L120, L121,
A122, D123, G124, V125, S126, T128, P129, T130, V131, E132, I140, L141, E142, K143,
R144, N145, A146, S147, K148, P149, Q150, G151, R152, G155, K157, V158, P160, K161,
E163, L171, N173, G174, A175, N184, T185, I186, H193, K197, K199, N200, R202, N203,
I205, S214, E215, H216, D217, G218, D219, S222, R224, S232, T233, V235, P236, G237,
20 T238, T239, N240, H249, Q250, P251, V253, T255, D256, E265, R266, T267, E270, R271,
F275, V276, R277, F278, L280, L287, L288, D289, R290, G291, A292, T293, L295, E296,
N301, M306, T307, Q308, D309, L311, Q312, Q313, R315, K316, V317, G318, D319,
S320, P321, N322, T324, E325, Y326, Y332, S333, D334, S336, K337, K341, G342, H351,
R353, G354, Q366, G367, T370, V371, G372, R379, E385, Q388, K389, R392, S393, E394,
25 P395, R396, P397, G398, V399, L400, L401, R402, P404 and P406 (A1-S45 are located in
the Gla domain, the remaining positions are located outside the Gla domain).

The following residues were determined to have more than 50% of their side chain
exposed to the surface: A1, A3, F4, L5, E6, E7, L8, R9, P10, E14, E16, K18, E19, E20, Q21,
S23, E25, E26, E29, K32, A34, E35, R36, K38, L39, I42, S43, G47, D48, A51, S52, S53,
30 Q56, G58, S60, K62, L65, Q66, S67, I69, F71, L73, P74, A75, E77, G78, R79, E82, H84,
K85, D86, D87, Q88, L89, I90, V92, N93, E94, G97, T106, G107, T108, K109, S111, E116,
S119, L121, A122, D123, G124, V131, E132, L141, E142, K143, R144, N145, A146, S147,
K148, P149, Q150, G151, R152, G155, K157, P160, N173, G174, A175, K197, K199, N200,
R202, S214, E215, H216, G218, R224, V235, P236, G237, T238, H249, Q250, V253, D256,

T267, F275, R277, F278, L288, D289, R290, G291, A292, T293, L295, N301, M306, Q308, D309, L311, Q312, Q313, R315, K316, G318, D319, N322, E325, D334, K341, G354, G367, V371, E385, K389, R392, E394, R396, P397, G398, R402, P404 and P406 (A1-S43 are located in the Gla domain, the remaining positions are located outside the Gla domain).

5

Tissue Factor Binding Site

It was determined using ASA calculations that the following residues in hFVII change their ASA in the complex. These residues were defined as constituting the receptor binding site: L13, K18, F31, E35, R36, L39, F40, I42, S43, S60, K62, D63, Q64, L65, I69, C70, F71, C72, L73, P74, F76, E77, G78, R79, E82, K85, Q88, I90, V92, N93, E94, R271, A274, F275, V276, R277, F278, R304, L305, M306, T307, Q308, D309, Q312, Q313, E325 and R379.

Active Site Region

15 The active site region is defined as any residue having at least one atom within a distance of 10 Å from any atom in the catalytic triad (residues H193, D242, S344): I153, Q167, V168, L169, L170, L171, Q176, L177, C178, G179, G180, T181, V188, V189, S190, A191, A192, H193, C194, F195, D196, K197, I198, W201, V228, I229, I230, P231, S232, T233, Y234, V235, P236, G237, T238, T239, N240, H241, D242, I243, A244, L245, L246, 20 V281, S282, G283, W284, G285, Q286, T293, T324, E325, Y326, M327, F328, D338, S339, C340, K341, G342, D343, S344, G345, G346, P347, H348, L358, T359, G360, I361, V362, S363, W364, G365, C368, V376, Y377, T378, R379, V380, Q382, Y383, W386, L387, L400 and F405.

25 *The Ridge of the Active Site Binding Cleft*

The ridge of the active site binding cleft region was defined by visual inspection of the FVIIa structure 1FAK.pdb as: N173, A175, K199, N200, N203, D289, R290, G291, A292, P321 and T370.

30 **Example 2**

Design of an expression cassette for expression of rhFVII in mammalian cells

The expression cassette for expression of rhFVII was designed and cloned as described in Example 2 of WO 01/58935.

Example 3*Construction of expression cassette encoding variants of the invention*

Sequence overhang extension (SOE) PCR was used for generating constructs having variant FVII open reading frames with substituted codons by using standard methods.

5

Example 4*Expression of polypeptide variants in CHO K1 cells*

The cell line CHO K1 (ATCC # CCL-61) is seeded at 50% confluence in T-25 flasks using MEM α , 10% FCS (Gibco/BRL Cat # 10091), P/S and 5 μ g/ml phylloquinone and allowed to grow until confluent. The confluent mono cell layer is transfected with 5 μ g of the relevant plasmid described above using the Lipofectamine 2000 transfection agent (Life Technologies) according to the manufacturer's instructions. Twenty four hours post transfection a sample is drawn and quantified using e.g. an ELISA recognizing the EGF1 domain of hFVII. At this time point relevant selection (e.g. Hygromycin B) may be applied to the cells with the purpose of generating a pool of stable transfecants. When using CHO K1 cells and the Hygromycin B resistance gene as selectable marker on the plasmid, this is usually achieved within one week.

Example 520 *Generation of CHO K1 cells stably expressing polypeptide variants.*

A vial of CHO-K1 transfecant pool is thawed and the cells seeded in a 175 cm^2 tissue flask containing 25 ml of MEM α , 10% FCS, phylloquinone (5 μ g/ml), 100 U/l penicillin, 100 μ g/l streptomycin and grown for 24 hours. The cells are harvested, diluted and plated in 96-well microtiter plates at a cell density of 1/2-1 cell/well. After a week of growth, colonies of 20-100 cells are present in the wells and those wells containing only one colony are labelled. After a further two weeks, the media in all wells containing only one colony is substituted with 200 μ l fresh medium. After 24 hours, a medium sample is withdrawn and analysed by e.g. ELISA. High producing clones are selected and used for large scale production of FVII or variants.

30

Example 6*Purification of polypeptide variants and subsequent activation*

FVII and FVII variants are purified as follows: The procedure is performed at 4°C. The harvested culture media from large-scale production is ultrafiltered using a Millipore

TFF system with 30 kDa cut-off Pellicon membranes. After concentration of the medium, citrate is added to 5 mM and the pH is adjusted to 8.6. If necessary, the conductivity is lowered to below 10 mS/cm. Subsequently, the sample is applied to a Q-sepharose FF column, equilibrated with 50 mM NaCl, 10 mM Tris pH 8.6. After washing the column with 5 100 mM NaCl, 10 mM Tris pH 8.6, followed by 150 mM NaCl, 10 mM Tris pH 8.6, FVII is eluted using 10 mM Tris, 25 mM NaCl, 35 mM CaCl₂, pH 8.6.

For the second chromatographic step, an affinity column is prepared by coupling of a monoclonal Calcium-dependent antiGla-domain antibody to CNBr-activated Sepharose FF. About 5.5 mg antibody is coupled per ml resin. The column is equilibrated with 10 mM Tris, 10 100 mM NaCl, 35 mM CaCl₂, pH 7.5. NaCl is added to the sample to a concentration of 100 mM NaCl and the pH is adjusted to 7.4 -7.6. After O/N application of the sample, the column is washed with 100 mM NaCl, 35 mM CaCl₂, 10 mM Tris pH 7.5, and the FVII protein is eluted with 100 mM NaCl, 50 mM citrate, 75 mM Tris pH 7.5.

For the third chromatographic, the conductivity of the sample is lowered to below 10 15 mS/cm, if necessary, and the pH is adjusted to 8.6. The sample is then applied to a Q- sepharose column (equilibrated with 50 mM NaCl, 10 mM Tris pH 8.6) at a density around 3-5 mg protein per ml gel to obtain efficient activation. After application, the column is washed with 50 mM NaCl, 10 mM Tris pH 8.6 for about 4 hours with a flow of 3-4 column 20 volumes (cv) per hour. The FVII protein is eluted using a gradient of 0-100% of 500 mM NaCl, 10 mM Tris pH 8.6 over 40 cv. FVII containing fractions are pooled.

For the final chromatographic step, the conductivity is lowered to below 10 mS/cm. Subsequently, the sample is applied to a Q-sepharose column (equilibrated with 140 mM 25 NaCl, 10 mM glycylglycine pH 8.6) at a concentration of 3-5 mg protein per ml gel. The column is then washed with 140 mM NaCl, 10 mM glycylglycine pH 8.6 and FVII is eluted with 140 mM NaCl, 15 mM CaCl₂, 10 mM glycylglycine pH 8.6. The eluate is diluted to 10 mM CaCl₂ and the pH is adjusted 6.8-7.2. Finally, Tween-80 is added to 0.01% and the pH is adjusted to 5.5 for storage at -80°C.

Example 7

30 Experimental results - FX activation activity

Subjecting the variants of the invention to the "TF-independent Factor X Activation Assay", the following results were obtained (the results being expressed as a percentage of the activity of the P10Q+K32E variant as a reference):

Variant	TF-independent FX activation ($a_{variant}/a_{P10QK32E}$) * 100
5 rhFVIIa	10
P10Q+K32E (reference)	100
A3AY+P10Q+K32E+A34L	216
P10Q+K32E+D33F+A34E	194
P10Q+K32E+A34E+P74S	190
10 P10Q+K32E+A34E+R36E+K38E	144
P10Q+K32E+A34D+R36E	140

Table 1

15 As it appears from the above results, the variants of the invention showed a substantial improvement in FX activation activity as compared to rhFVIIa and also as compared to [P10Q+K32E]rhFVIIa.

Example 8

20 Experimental results - clotting activity in the "Whole Blood Assay"

Subj ecting variants of the invention to the "Whole Blood Assay" revealed that they exhibited a significantly increased clotting activity (i.e. reduced clotting time) as compared to rhFVIIa as well as [P10Q+K32E]rhFVIIa. The experimental results are shown in Fig. 1 and Table 2 below.

Variant	Clotting time (Whole Blood Assay) $t_{variant}/t_{wt}$
rhFVIIa (reference)	1
30 A3AY+P10Q+K32E+E116D	0.4
A3AY+P10Q+K32E+A34L	0.3
P10Q+K32E+A34E+P74S	0.3
A3AY+P10Q+K32E+E77A	0.4

Table 2

Example 9*Experimental results – clotting activity in the “Clotting Assay”*

When assayed in a TF-dependent clotting assay (the “Clotting Assay” described above in the Materials and Methods section) it was evident that variants of the invention 5 having the R36E substitution have a significantly reduced clotting activity when compared to rhFVII or to other variants of the invention. See Table 3 below. Nevertheless, as illustrated in Example 7 above, variants having the R36E substitution have an increased Factor X activation activity in the “TF-independent Factor X Activation Assay”.

10	Variant	Average Clotting Activity (units/mg _{variant} /units/mg _{wt}) (n = 2-3)
	NovoSeven® (reference)	52,119 (100 %)
15	P10Q+K32E	52,714 (101 %)
	A3AY+P10Q+K32E+A34L	56,948 (107 %)
	P10Q+K32E+A34E+R36E	1,439 (2.7 %)
	P10Q+K32E-A34D+R36E+K38E	1,232 (2.4 %)

20 Table 3

Example 10*Experimental results – thrombin generation in the Thrombogram Assay*

Using both phospholipid(PL)-dependent and tissue factor(TF)-dependent 25 thrombograms (see the description of the Thrombogram Assay above), the maximum rate of thrombin generation was determined for FVIIa variants at different concentrations of variant proteins. By plotting the maximum thrombin generation rates (expressed as FU (fluorescence units) per sec²) as a function of the variant concentration in pM, the results shown in Figure 2 (maximum tissue factor-dependent thrombin generation rate) and Figure 3 (maximum 30 phospholipid-dependent thrombin generation rate) were obtained.

From these results it is evident that the FVIIa variant P10Q K32E A34E R36E has a differentiated thrombin generation ability depending on the whether the reaction is PL-dependent or TF-dependent. The maximum TF-dependent thrombin generation rate of this variant is decreased by approximately 10-fold (punctuated line in Figure 2) when compared 35 to the FVIIa variants P10Q K32E or A3AY P10Q K32E A34L. Also, lag time, time to peak, peak height and (to a lesser extent) AUC are reduced for P10Q K32E A34E R36E compared

to the other variants (results not shown). In contrast to the TF-dependent activity, the PL-dependent activity of the P10Q K32E A34E R36E variant is equivalent to that of the other variants tested in this example (see Figure 3), i.e. this variant has full PL-dependent activity even though the TF-dependent activity is substantially reduced.

5 In the same experiment, the variant P10Q K32E A34E R36E was compared directly to the variant P10Q K32E A34E P74S, which has a high TF-dependent thrombin generation rate as shown in Figure 2. The differences in TF-binding between these two variants (i.e. the reduced TF-binding of the variant P10Q K32E A34E R36E) is believed to be directly attributable to the presence of the R36E substitution, possibly in synergy with the A34E

10 substitution.

Example 11

Experimental results – FVIIa binding to tissue factor in the Biacore Assay

15 Subjecting variants of the invention to assay by surface plasmon resonance on a Biacore system using a TF chip as described in the Materials and Methods section, the following results were obtained:

20 Variant	Average response units (n = 5)
Wild-type FVIIa	888
P10Q; K32E	714
A3AY; P10Q; K32E; A34L	967*
25 P10Q; K32E; A34E; R36E	414

Table 4

* n = 2

In consistency with the TF-dependent thrombin generation rate data from the

30 Thrombogram Assay (Example 10), the results in Table 4 indicate that the R36E substitution confers less binding to tissue factor.

In the same Biacore Assay, FVIIa variants having the same modifications as the variants listed in Table 4 together with two additional modifications introducing two glycosylation sites (T106N and either V253N or I205T) were also tested for binding to tissue

35 factor. The results are shown in Table 5 below.

Variant	Average response units (n = 5)
5 T106N; V253N	717
T106N; I205T	612
P10Q; K32E; T106N; I205T	502
P10Q; K32E; T106N; V253N	498
A3AY; P10Q; K32E; A34L; T106N; V253N	522
10 P10Q; K32E; A34E; R36E; T106N; I205T	216

Table 5

These results are consistent with those of Table 4 and show that compared to the
 15 same variants (or the wild-type) in Table 4 without the additional glycosylation sites, the
 presence of two new glycosylation sites in the variants of Table 5 provides a (further)
 reduction in tissue factor binding. As was the case for the variants of Table 4, the presence of
 the R36E substitution in a glycosylation variant also results in a level of tissue factor binding
 that is substantially lower than the tissue factor binding of the other glycosylation variants
 20 that do not have this substitution.

The claims defining the invention are as follows:

1. A Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence that differs in 1-15 amino acid residues relative to the amino acid sequence of human Factor VII (hFVII) or human Factor VIIa (hFVIIa) shown in SEQ ID NO:1, wherein a negatively charged amino acid residue has been introduced by substitution in position 36.
2. The variant of claim 1, wherein said substitution is R36D or R36E.
3. The variant of claim 1 or 2, further comprising an amino acid substitution in position 34.
4. The variant of claim 3, comprising the substitution A34L or A34E.
5. The variant of claim 3 or 4 comprising the substitutions A34E+R36E.
6. The variant of any one of claims 1 to 5, further comprising an amino acid substitution in position 10 and/or 32.
7. The variant of claim 6, comprising the substitution K32E or P10Q.
8. The variant of claim 6 or 7, comprising the substitutions P10Q+K32E.
9. The variant of claim 8, comprising the substitutions P10Q+K32E+A34E+R36E or P10Q+K32E+A34L+R36E.
10. The variant of any one of claims 1 to 9, wherein at least one amino acid residue comprising an attachment group for a non-polypeptide moiety has been introduced in a position located outside the Gla domain.
11. The variant of claim 10, wherein said attachment group is an *in vivo* glycosylation site.
12. The variant of claim 11, wherein said glycosylation site is an *in vivo* N-glycosylation site introduced by substitution.
13. The variant of claim 12, wherein said *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205S, I205T, V253N, T267N, T267N+S269T, S314N+K316S, S314N+K316T, R315N+V317S, R315N+V317T, K316N+G318S, K316N+G318T, G318N, D334N and combinations thereof.
14. The variant of claim 13, comprising at least one substitution selected from the group consisting of T106N, I205T and V253N.
15. The variant of claim 14, comprising two *in vivo* N-glycosylation sites introduced by substitutions selected from the group consisting of T106N+I205T, T106N+V253N and I205T+V253N.

16. The variant of claim 15, comprising the substitutions

P10Q+K32E+A34E+R36E+T106N+I205T or
P10Q+K32E+A34E+R36E+T106N+V253N or
P10Q+K32E+A34E+R36E+I205T+V253N or
P10Q+K32E+A34L+R36E+T106N+I205T or
P10Q+K32E+A34L+R36E+T106N+V253N or
P10Q+K32E+A34L+R36E+I205T+V253N.

17. The variant of any one of claims 1 to 16, further comprising an insertion of at least one amino acid residue between position 3 and 4.

18. The variant of claim 17, comprising an insertion of one amino acid residue between position 3 and 4.

19. The variant of claim 17 or 18, wherein a hydrophobic amino acid residue is inserted between position 3 and 4.

20. The variant of claim 19, wherein said insertion is A3AY.

21. The variant of claim 20, comprising:

(i) the insertion A3AY and the substitutions P10Q+K32E+A34E+R36E, or
(ii) the insertion A3AY and the substitutions P10Q+K32E+A34L+R36E, or
(iii) the insertion A3AY and the substitutions P10Q+K32E+A34E+R36E+T106N+I205T, or
(iv) the insertion A3AY and the substitutions P10Q+K32E+A34E+R36E+T106N+V253N, or
(v) the insertion A3AY and the substitutions P10Q+K32E+A34E+R36E+I205T+V253N, or
(vi) the insertion A3AY and the substitutions P10Q+K32E+A34L+R36E+T106N+I205T, or
(vii) the insertion A3AY and the substitutions P10Q+K32E+A34L+R36E+T106N+V253N, or
(viii) the insertion A3AY and the substitutions P10Q+K32E+A34L+R36E+I205T+V253N.

22. The variant of any one of claims 1 to 21, wherein said variant is in its activated form.

23. A nucleotide sequence encoding a variant as defined in any one of claims 1 to 22.

24. An expression vector comprising the nucleotide sequence of claim 23.

25. A host cell comprising the nucleotide sequence of claim 23 or the expression vector of claim 24, wherein said host cell is a gamma-carboxylating cell capable of *in vivo* glycosylation.
26. A composition comprising a variant as defined in any one of claims 1 to 5 22 and at least one pharmaceutical acceptable carrier or excipient.
27. A method for treating a mammal having a disease or a disorder wherein clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of the variant as defined in any one of claims 1 to 22, or the composition of claim 26.
- 10 28. The method of claim 27, wherein said disease or disorder is selected from the group consisting of hemorrhages, including brain hemorrhages, severe uncontrolled bleedings, such as trauma, bleedings in patients undergoing transplantations or resection, variceal bleeding, and hemophilia.
- 15 29. The method of claim 28, wherein said disease or disorder is trauma or hemophilia.
30. Use of a variant as defined in any one of claims 1 to 22 in the manufacture of a medicament for treatment of a mammal having a disease or disorder wherein clot formation is desirable.

20

Dated 10 November, 2009

Bayer HealthCare LLC

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

2377595

1/3




Fig. 1

2/3




Fig. 2

3/3




Fig. 3

SEQUENCE LISTING

<110> Maxygen ApS
 Maxygen Holdings Ltd.
 Haaning, Jesper Mortensen
 Andersen, Kim Vilbørv
 Bornæs, Claus

<120> FVII or FVIIa Gla Domain Variants

<130> 0274wo310

<150> US 60/479,780
 <151> 2003-06-19

<150> DK PA 2004 00930
 <151> 2004-06-15

<160> 1

<170> PatentIn version 3.2

<210> 1

<211> 406

<212> PRT

<213> Homo sapiens

<400> 2

Ala Asn Ala Phe Leu Glu Glu Leu Arg Pro Gly Ser Leu Glu Arg Glu
 1 5 10 15

Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Lys
 20 25 30

Asp Ala Glu Arg Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp
 35 40 45

Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln
 50 55 60

Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn
 65 70 75 80

Cys Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly
 85 90 95

Gly Cys Glu Gln Tyr Cys Ser Asp His Thr Gly Thr Lys Arg Ser Cys
 100 105 110

Arg Cys His Glu Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr
 115 120 125

Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg
 130 135 140

Asn Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro
 145 150 155 160

Lys Gly Glu Cys Pro Trp Gln Val Leu Leu Leu Val Asn Gly Ala Gln
 165 170 175
 Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala
 180 185 190
 His Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu
 195 200 205
 Gly Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg
 210 215 220
 Val Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn
 225 230 235 240
 His Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp
 245 250 255
 His Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr
 260 265 270
 Leu Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gln Leu Leu
 275 280 285
 Asp Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg
 290 295 300
 Leu Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser
 305 310 315 320
 Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser
 325 330 335
 Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr
 340 345 350
 Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys
 355 360 365
 Ala Thr Val Gly His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile
 370 375 380
 Glu Trp Leu Gln Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu
 385 390 395 400
 Leu Arg Ala Pro Phe Pro
 405