

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11

Publication number:

0 682 257 A1

12

EUROPEAN PATENT APPLICATION

21

Application number: **95201176.5**

51

Int. Cl.⁶: **G01N 33/68, C12N 5/20,
C12P 21/08, C07K 16/18**

22

Date of filing: **30.11.90**

This application was filed on 08 - 05 - 1995 as a
divisional application to the application
mentioned under INID code 60.

**4225 Roosevelt Way
Seattle, WA 98040 (US)**

30

Priority: **01.12.89 US 444881
21.11.90 US 614719**

72

Inventor: **Eyre, David R.
8011 East Mercer Way,
Mercer Island
WA 98040 (US)**

43

Date of publication of application:
15.11.95 Bulletin 95/46

74

Representative: **Bizley, Richard Edward et al
Hepworth, Lawrence, Bryer & Bizley
Merlin House
Falconry Court
Bakers Lane
Epping
Essex CM16 5DO (GB)**

60

Publication number of the earlier application in
accordance with Art.76 EPC: **0 502 928**

84

Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

71


Applicant: **Washington Research Foundation**

54

Methods of detecting collagen degradation in vivo.

57

The invention provides assays and kits for measuring bone resorption by determining in a sample of body fluid the presence of concentration of a carboxy-terminal type I collagen telopeptide comprising:



wherein

EP 0 682 257 A1

K

|

K

|

K

is lysyl pyridinoline or hydroxylysyl pyridinoline.

The present invention relates to a method for detecting and monitoring collagen degradation *in vivo*. More specifically, it relates to a method and kit for determining certain cross-linked telopeptides produced *in vivo* upon degradation of collagen type I.

5 Background of the Invention

Three known classes of collagens have been described to date. The Class 1 collagens, subdivided into types I, II, III, V, and XI, are known to form fibrils. These collagens are all synthesized as procollagen molecules, made up of N-terminal and C-terminal propeptides, which are attached to the core collagen
10 molecule. After removal of the propeptides, which occurs naturally *in vivo* during collagen synthesis, the remaining core of the collagen molecule consists largely of a triple helical domain having terminal telopeptide sequences which are nontriple-helical. These telopeptide sequences have an important function as sites of intermolecular cross-linking of collagen fibrils extracellularly.

The present invention relates to methods of detecting collagen degradation based on assaying for
15 particular cross-linked telopeptides produced *in vivo* upon collagen degradation. In the past, assays have been developed for monitoring degradation of collagen *in vivo* by measuring various biochemical markers, some of which have been degradation products of collagen. For example, bone turnover associated with Paget's disease has been monitored by measuring small peptides containing hydroxyproline, which are excreted in the urine following degradation of bone collagen. Russell et al., *Metab, Bone Dis. and Rel. Res.*
20 4 and 5, 255-262 (1981); and Singer, F.R., et al., *Metabolic Bone Disease*. Vol. II (eds Avioli, L.V. and Kane, S.M.), 489-575 (1978), Academic Press, New York.

Other researchers have measured the cross-linking compound pyridinoline in urine as an index of collagen degradation in joint disease. See, for background and for example, Wu and Eyre, *Biochem-
istry*, 23:1850 (1984); Black et al., *Annals of the Rheumatic Diseases*, 48:641-644 (1989); Robins et al.,
25 *Annals of the Rheumatic Diseases*, 45:969-973 (1986); and Seibel et al., *The Journal of Rheumatology*, 16:964 (1989). In contrast to the present invention, some prior researchers have hydrolyzed peptides from body fluids and then looked for the presence of individual hydroxypyridinium residues. None of these researchers has reported measuring a telopeptide containing a cross-link that is naturally produced *in vivo* upon collagen degradation, as in the present invention.

UK Patent application GB 2,205,643 reports that the degradation of type III collagen in the body is
30 quantitatively determined by measuring the concentration of an N-terminal telopeptide from type III collagen in a body fluid. In this reference, it is reported that cross-linked telopeptide regions are not desirable. In fact, this reference reports that it is necessary to use a non-cross-linked source of collagen to obtain the telopeptide. The peptides of the present invention are all cross-linked.

Collagen cross-links are discussed in greater detail below, under the heading "Collagen Cross-Linking".

There are a number of reports indicating that collagen degradation can be measured by quantitating certain procollagen peptides. The present invention involves telopeptides rather than propeptides, the two being distinguished by their location in the collagen molecule and the timing of their cleavage *in vivo*. See
40 US Patent 4,504,587; US Patent 4,312,853; Pierard et al., *Analytical Biochemistry* 141:127-136 (1984); Niemela, *Clin. Chem.*, 31/8:1301-1304 (1985); and Rohde et al., *European Journal of Clinical Investiga-
tion*, 9:451-459 (1979).

US Patent 4,778,768 relates to a method of determining changes occurring in articular cartilage involving quantifying proteoglycan monomer or antigenic fragments thereof in a synovial fluid sample. This patent does not relate to detecting cross-linked telopeptides derived from degraded collagen.

Dodge, *J. Clin. Invest.*, 83:647-661 (1981) discloses methods for analyzing type II collagen degradation
45 utilizing a polyclonal antiserum that specifically reacts with unwound alpha-chains and cyanogen bromide-derived peptides of human and bovine type II collagens. The peptides involved are not cross-linked telopeptides as in the present invention.

Amino acid sequences of human type III collagen, human proal(II) collagen, and the entire preproal(III)
50 chain of human type III collagen and corresponding cDNA clones have been investigated and determined by several groups of researchers. See Loidl et al., *Nucleic Acids Research* 12:9383-9394 (1984); Sangiorgi et al., *Nucleic Acids Research*, 13:2207-2225 (1985); Baldwin et al., *Biochem. J.*, 262:521-528 (1989); and Ala-Kokko et al., *Biochem. J.*, 260:509-516 (1989). None of these references specifies the structures of particular telopeptide degradation products that could be measured to determine the amount of degraded
55 fibrillar collagen *in vivo*.

In spite of the above-described background information, there remains a need for effective and simple assays for determining collagen degradation *in vivo*. Such assays could be used to detect and monitor disease states in humans, such as osteoarthritis (type II collagen degradation), and various inflammatory

disorders, such as vasculitis syndrome (type III collagen degradation).

Assays for type I collagen degradation, described in WO-A-85/04491, can be utilized to detect and assess bone resorption *in vivo*. Detection of bone resorption may be a factor of interest in monitoring and detecting diseases such as osteoporosis. Osteoporosis is the most common bone disease in man. Primary
5 osteoporosis, with increased susceptibility to fractures, results from a progressive net loss of skeletal bone mass. It is estimated to affect 15-20 million individuals in the United States. Its basis is an age-dependent imbalance in bone remodeling, i.e., in the rates of synthesis and degradation of bone tissue.

About 1.2 million osteoporosis-related fractures occur in the elderly each year including about 538,000 compression fractures of the spine, about 227,000 hip fractures and a substantial number of early fractured
10 peripheral bones. Twelve to 20% of the hip fractures are fatal because they cause severe trauma and bleeding, and half of the surviving patients require nursing home care. Total costs from osteoporosis-related injuries now amount to at least \$7 billion annually (Barnes, O.M., *Science*, 236:914 (1987)).

Osteoporosis is most common in postmenopausal women who, on average, lose 15% of their bone mass in the 10 years after menopause. This disease also occurs in men as they get older and in young
15 amenorrheic women athletes. Despite the major, and growing, social and economic consequences of osteoporosis, no method is available for measuring bone resorption rates in patients or normal subjects. A major difficulty in monitoring the disease is the lack of a specific assay for measuring bone resorption rates.

Methods for assessing bone mass often rely on measuring whole-body calcium by neutron activation analysis or mineral mass in a given bone by photon absorption techniques. These measurements can give
20 only long-term impressions of whether bone mass is decreasing. Measuring calcium balances by comparing intake with output is tedious, unreliable and can only indirectly appraise whether bone mineral is being lost over the long term. Other methods currently available for assessing decreased bone mass and altered bone metabolism include quantitative scanning radiometry at selected bone locations (wrist, calcaneus, etc) and histomorphometry of iliac crest biopsies. The former provides a crude measure of the bone mineral content
25 at a specific site in a single bone.

Histomorphometry gives a semi-quantitative assessment of the balance between newly deposited bone seams and resorbing surfaces.

A urinary assay for the whole-body output of degraded bone in 24 hours would be much more useful. Mineral studies (e.g., calcium balance) cannot do this reliably or easily. Since bone resorption involves
30 degradation of the mineral and the organic matrix, a specific biochemical marker for newly degraded bone products in body fluids would be the ideal index. Several potential organic indices have been tested. For example, hydroxyproline, an amino acid largely restricted to collagen, and the principal structural protein in bone and all other connective tissues, is excreted in urine. Its excretion rate is known to be increased in certain conditions, notably Paget's disease, a metabolic bone disorder in which bone turnover is greatly
35 increased, as pointed out above. For this reason, urinary hydroxyproline has been used extensively as an amino acid marker for collagen degradation. Singer, F.R., et al. (1978), cited hereinabove.

US Patent No 3,600,132 discloses a process for determination of hydroxyproline in body fluids such as serum, urine, lumbar fluid and other intercellular fluids in order to monitor deviations in collagen metabolism. In particular, this inventor notes that in pathologic conditions such as Paget's disease, Marfan's
40 syndrome, osteogenesis imperfecta, neoplastic growth in collagen tissues and in various forms of dwarfism, increased collagen anabolism or catabolism as measured by hydroxyproline content in biological fluids can be determined. This inventor measures hydroxyproline by oxidizing it to a pyrrole compound with hydrogen peroxide and ^N-chloro-_p-toluenesulphonamide followed by colorimetric determination in _p-dimethyl-amino-benzaldehyde.

In the case of Paget's disease, the increased urinary hydroxyproline probably comes largely from bone degradation; hydroxyproline, however, generally cannot be used as a specific index. Much of the hydroxyproline in urine may come from new collagen synthesis (considerable amounts of the newly made protein are degraded and excreted without ever becoming incorporated into tissue fabric), and from turnover of certain blood proteins as well as other proteins that contain hydroxyproline. Furthermore, about 80% of the
50 free hydroxyproline derived from protein degradation is metabolized in the liver and never appears in the urine. Kivirikko, K.I. *Int. Rev. Connect. Tissue Res.* 5:93 (1970), and Weiss, P.H., and Klein, L., *J. Clin. Invest.* 48:1 (1969).

Hydroxylysine and its glycoside derivatives, both peculiar to collagenous proteins, have been considered to be more accurate than hydroxyproline as markers of collagen degradation. However, for the same
55 reasons described above for hydroxyproline, hydroxylysine and its glycosides are probably equally non-specific markers of bone resorption. Krane, S.M. and Simon, L.S. *Develop. Biochem.*, 22:185 (1981).

In addition to amino acids unique to collagen, various non-collagenous proteins of bone matrix such as osteocalcin, or their breakdown products, have formed the basis of immunoassays aimed at measuring

bone metabolism. Price, P.A. et al. *J. Clin. Invest.*, 66:878 (1980) and Gundberg, C.M. et al., *Meth. Enzymol.*, 107:516 (1984). However, it is now clear that bone-derived non-collagenous proteins, though potentially a useful index of bone metabolic activity are unlikely, on their own, to provide quantitative measures of bone resorption. The concentration in serum of osteocalcin, for example, fluctuates quite widely both normally and in metabolic bone disease. Its concentration is elevated in states of high skeletal turnover but it is unclear whether this results from increased synthesis or degradation of bone. Krane, S.M., et al., *Develop. Biochem.*, 22:185 (1981), Price, P.A. et al., *J. Clin. Invest.*, 66:878 (1980); and Gundberg, C.M. et al., *Meth. Enzymol.*, 107:516 (1984).

10 Collagen Cross-Linking

The polymers of most genetic types of vertebrate collagen require the formation of aldehyde-mediated cross-links for normal function. Collagen aldehydes are derived from a few specific lysine or hydroxylysine side-chains by the action of lysyl oxidase. Various di-, tri- and tetrafunctional cross-linking amino acids are formed by the spontaneous intra- and intermolecular reactions of these aldehydes within the newly formed collagen polymers; the type of cross-linking residue varies specifically with tissue type (see Eyre, D.R. et al., *Ann. Rev. Biochem.*, 53:717-748 (1984)).

Two basic pathways of cross-linking can be differentiated for the banded (67nm repeat) fibrillar collagens, one based on lysine aldehydes, the other on hydroxylysine aldehydes. The lysine aldehyde pathway dominates in adult skin, cornea, sclera, and rat tail tendon and also frequently occurs in other soft connective tissues. The hydroxylysine aldehyde pathway dominates in bone, cartilage, ligament, most tendons and most internal connective tissues of the body, Eyre, D.R. et al. (1974) *vide supra*. The operating pathway is governed by whether lysine residues are hydroxylated in the telopeptide sites where aldehyde residues will later be formed by lysyl oxidase (Barnes, M.J. et al., *Biochem. J.*, 139:461 (1974)).


The chemical structure(s) of the mature cross-linking amino acids on the lysine aldehyde pathway are unknown, but hydroxypyridinium residues have been identified as mature products on the hydroxylysine aldehyde route. On both pathways and in most tissues the intermediate, borohydride-reducible cross-linking residues disappear as the newly formed collagen matures, suggesting that they are relatively short-lived intermediates (Bailey, A.J. et al., *FEBS Lett*, 16:86 (1971). Exceptions are bone and dentin, where the reducible residues persist in appreciable concentration throughout life, in part apparently because the rapid mineralization of the newly made collagen fibrils inhibits further spontaneous cross-linking interactions (Eyre, D.R., In: *The Chemistry and Biology of Mineralized Connective Tissues*. (Veis, A. ed.) pp. 51-55 (1981), Elsevier, New York, and Walters, C. et al., *Calc. Tiss. Intl.* 35:401-405 (1983)).

Two chemical forms of 3-hydroxypyridinium cross-link have been identified (Formula I and II). Both compounds are naturally fluorescent, with the same characteristic excitation and emission spectra (Fujimoto, D. et al. *Biochem. Biophys. Res. Commun.*, 76:1124 (1977), and Eyre, D.R., *Develop. Biochem.* 22:50 (1981)). These amino acids can be resolved and assayed directly in tissue hydrolysates with good sensitivity using reverse phase HPLC and fluorescence detection. Eyre, D.R. et al., *Analyte. Biochem.*, 137:380-388 (1984). It should be noted that the present invention involves quantitating particular peptides rather than amino acids.


45

50

55



FORMULA I



FORMULA II

25

In growing animals, it has been reported that these mature cross-links may be concentrated more in an unmineralized fraction of bone collagen than in the mineralized collagen (Banes, A.J., et al., *Biochem. Biophys. Res. Commun.*, 113:1975 (1983)). However, other studies on young bovine or adult human bone do not support this concept, Eyre, D.R., In: *The Chemistry and Biology of Mineralized Tissues* (Butler, W.T. ed.)p. 105 (1985), Ebsco Media Inc., Birmingham, Alabama.

The presence of collagen hydroxypyridinium cross-links in human urine was first reported by Gunja-Smith and Boucek (Gunja-Smith, Z. and Boucek, R.J., *Biochem J.*, 197:759-762 (1981)) using lengthy isolation procedures for peptides and conventional amino acid analysis. At that time, they were aware only of the HP forms of the cross-link. Robins (Robins, S.P., *Biochem J.*, 207:617-620 (1982)) has reported an enzyme-linked immunoassay to measure HP in urine, having raised polyclonal antibodies to the free amino acid conjugated to bovine serum albumin. This assay is intended to provide an index for monitoring increased joint destruction that occurs with arthritic diseases and is based, according to Robins, on the finding that pyridinolone is much more prevalent in cartilage than in bone collagen.

In more recent work involving enzyme-linked immunoassay, Robins reports that lysyl pyridinolone is unreactive toward antiserum to pyridinolone covalently linked to bovine serum albumin (Robins et al., *Ann. Rheum. Diseases*, 45:969-973 (1986)). Robins' urinary index for cartilage destruction is based on the discovery that hydroxylysyl pyridinolone, derived primarily from cartilage, is found in urine at concentrations proportional to the rate of joint cartilage resorption (i.e. degradation). In principle, this index could be used to measure whole body cartilage loss; however, no information on bone resorption would be available.


Also of interest is the international publication No. WO 89/12824, by The Rowett Research Institute. This later publication describes a method of monitoring collagen degradation, comprising assaying a biological fluid sample which contains a fragment of collagen including lysyl pyridinolone or hydroxylysyl pyridinolone or a substituted form thereof, and a method of determining the tissue origin of degraded collagen thereby.

A need therefore exists for a method that allows the measurement of whole body bone resorption rates in humans. The most useful such method would be one that could be applied to body fluids, especially urine. The method should be sensitive, i.e., quantifiable down to 1 picomole and rapidly measure 24-hour bone resorption rates so that the progress of various therapies (e.g., estrogen) can be assessed.

Summary of the Invention

The present invention is based on the discovery of the presence of particular cross-linked telopeptides in body fluids of patients and normal human subjects. These telopeptides are produced in vivo during collagen degradation and remodeling. The "telopeptides" are cross-linked peptides having sequences that are associated with the carboxy-terminal telopeptide region of type I collagen. Having disclosed the structures of these telopeptides herein, it will be appreciated by one of ordinary skill in the art that they may also be produced other than in vivo e.g., synthetically. These peptides will generally be provided in purified form e.g., substantially free of impurities, particularly other peptides.

The invention provides as assay for measuring bone resorption, by determining in a sample of body fluid the presence or concentration of a carboxy-terminal type I collagen telopeptide comprising:




wherein



35 is lysyl pyridinoline or hydroxylysyl pyridinoline.

In another aspect, the invention provides a kit for measuring bone resorption, comprising an immunological binding partner which binds to a carboxy-terminal type I collagen telopeptide comprising:



50 wherein

55

K

|

5

K

|

K

10

is lysyl pyridinoline or hydroxylysyl pyridinoline.

The present invention thus relates to a method for determining *in vivo* degradation of type I collagen. The method can involve quantitating in a fluid the concentration of the particular telopeptides that have a 3-hydroxypyridinium cross-link and that are derived from type I collagen degradation. The methods disclosed
15 in the present invention are analogous to those disclosed in WO-A-85/04491 for determining the absolute rate of bone resorption *in vivo*. Those methods involved quantitating in a body fluid the concentration of telopeptides having a 3-hydroxypyridinium cross-link derived from bone collagen resorption.

In a representative assay, the patient's body fluid is contacted with an immunological binding partner specific to such a telopeptide. The body fluid may be used as is or purified prior to the contacting step.
20 This purification step may be accomplished using a number of standard procedures, including cartridge adsorption and elution, molecular sieve chromatography, dialysis, ion exchange, alumina chromatography, hydroxyapatite chromatography, and combinations thereof.

Other representative embodiments of quantitating the concentration of peptide fragments having a 3-hydroxypyridinium cross-link in a body fluid include electrochemical titration, natural fluorescence spectroscopy, and ultraviolet absorbance. Electrochemical titration may be conducted directly upon a body fluid
25 without further purification. However, when this is not possible due to excessive quantities of contaminating substances, the body fluid is first purified prior to the electrochemical titration step. Suitable methods for purification prior to electrochemical detection include dialysis, ion exchange chromatography, alumina chromatography, molecular sieve chromatography, hydroxyapatite chromatography and ion exchange absorption and elution.
30

Fluorometric measurement of a body fluid containing a 3-hydroxypyridinium cross-link is an alternative way of quantitating collagen degradation (and, hence, bone resorption, if type I peptides are quantitated). The fluorometric assay can be conducted directly on a body fluid without further purification. However, for certain body fluids, particularly urine, it is preferred that purification of the body fluid be conducted prior to
35 the fluorometric assay. This purification step consists of dialyzing an aliquot of a body fluid such as urine against an aqueous solution thereby producing partially purified peptide fragments retained within the nondiffusate (retentate). The nondiffusate is then lyophilized, dissolved in an ion pairing solution and adsorbed onto an affinity chromatography column. The chromatography column is washed with a volume of ion pairing solution and, thereafter, the peptide fragments are eluted from the column with an eluting
40 solution. These purified peptide fragments may then be hydrolyzed and the hydrolysate resolved chromatographically. Chromatographic resolution may be conducted by either high-performance liquid chromatography or microbore high performance liquid chromatography.

Another aspect of the present invention involves assays for the peptides described herein in which the pyridinium rings are intact and cleaved. Since it is suspected that some cleavage of pyridinium rings occurs
45 *in vivo*, assays that detect both intact and cleaved pyridinium rings may lead to more accurate assessments of collagen degradation. In connection with this aspect of the present invention, specific binding partners to the individual peptides containing intact or cleaved pyridinium rings, may be employed in the assays. Individual specific binding partners that recognize both types of peptides (both intact and cleaved pyridinium ring containing peptides) may be employed. Alternatively, specific binding partners that discriminate
50 between peptides containing the intact pyridinium ring and those in which the pyridinium ring is cleaved, could also be used.

The invention generally includes all specific binding partners to the peptides described herein. "Specific binding partners" are molecules that are capable of binding to the peptides of the present invention. Included within this term are immunological binding partners, such as antibodies (monoclonal and poly-
55 clonal), antigen-binding fragments of antibodies (e.g., Fab and F(ab')₂ fragments), single-chain antigen-binding molecules, and the like, whether made by hybridoma or rDNA technologies.

The invention includes fused cell hybrids (hybridomas) that produce monoclonal antibodies specific for the above-described collagen peptides having 3-hydroxypyridinium cross-links (both with an intact

pyridinium ring and one that has been cleaved).

The invention further includes monoclonal antibodies produced by the fused cell hybrids, and those antibodies (as well as binding fragments thereof, e.g., Fab) coupled to a detectable marker. Examples of detectable markers include enzymes, chromophores, fluorophores, coenzymes, enzyme inhibitors, chemiluminescent materials, paramagnetic metals, spin labels, and radioisotopes. Such specific binding partners may alternatively be coupled to one member of a ligand-binding partner complex (e.g., avidin-biotin), in which case the detectable marker can be supplied bound to the complementary member of the complex.

The test kits of the invention useful for quantitating the amount of peptides having 3-hydroxypyridinium cross-links derived from collagen type I degradation in a body fluid may include a specific binding partner to a peptide derived from degraded collagen as disclosed herein. The specific binding partners of the test kits may be coupled to a detectable marker or a member of a ligand-binding partner complex, as described above.

Brief Description of the Drawings

FIGURE 1 compares the concentration of HP and LP in both cortical and cancellous human bone with age.

FIGURE 2 depicts the cross-link molar ratios of HP to LP as a function of age.

FIGURE 3A shows relative fluorescence (297nm excitation, >370nm emission) as a function of elution volume during reverse phase HPLC separation of cross-linked type I collagen N-telopeptides.

FIGURE 3B shows relative fluorescence (297nm excitation, >370nm emission) versus elution volume during reverse phase HPLC separation of cross-linked type I collagen C-telopeptides.

FIGURE 4A shows relative fluorescence (297 nm excitation, >380nm emission) as a function of elution time for the cross-linked type I collagen telopeptides.

FIGURE 4B shows relative fluorescence (297nm excitation, >380nm emission) as a function of elution time for the cross-linked type I collagen telopeptides.

Detailed Description of the Preferred Embodiment

The invention is based on the discovery that both lysyl pyridinoline (LP) and hydroxylysyl pyridinoline (HP) peptide fragments (i.e., telopeptides, as used herein) derived from reabsorbed bone collagen are excreted in the urine without being metabolized. The invention is also based on the discovery that no other connective tissues contain significant levels of LP and that the ratio of HP to LP in mature bone collagen remains relatively constant over a person's lifetime.

FIGURE 1 compares the concentration of HP and LP in both cortical and cancellous human bone with age. It is observed that the concentration of HP plus LP cross-links in bone collagen reaches a maximum by age 10 to 15 years and remains reasonably constant throughout adult life. Furthermore, the ratio of HP to LP, shown in FIGURE 2 shows little change throughout life, remaining constant at about 3.5 to 1. These baseline data demonstrate that the 3-hydroxypyridinium cross-links in bone collagen remains relatively constant and therefore that body fluids derived from bone collagen degradation will contain 3-hydroxypyridinium cross-linked peptide fragments at concentrations proportional to the absolute rate of bone resorption.

Since LP is the 3-hydroxypyridinium cross-link unique to bone collagen, the method for determining the absolute rate of bone resorption, in its simplest form, is based on quantitating the concentration of peptide fragments containing 3-hydroxypyridinium cross-links and preferably lysyl pyridinoline (LP) cross-links in a body fluid.

As used in this description and in the appended claims with respect to type 1 telopeptides, by "quantitating" is meant measuring by any suitable means, including but not limited to spectrophotometric, gravimetric, volumetric, coulometric, immunometric, potentiometric, or amperometric means the concentration of peptide fragments containing 3-hydroxypyridinium cross-links in an aliquot of a body fluid. Suitable body fluids include urine, serum, and synovial fluid. The preferred body fluid is urine.

Since the concentration of urinary peptides will decrease as the volume of urine increases, it is further preferred that when urine is the body fluid selected, the aliquot assayed be from a combined pool of urine collected over a fixed period of time, for example, 24 hours. In this way, the absolute rate of bone resorption or collagen degradation is calculated for a 24 hour period. Alternatively, urinary peptides may be measured as a ratio relative to a marker substance found in urine such as creatinine. In this way the urinary index of collagen degradation and bone resorption would remain independent of urine volume.

In one embodiment of the present invention, monoclonal or polyclonal antibodies are produced which are specific to the peptide fragments containing lysyl pyridinoline cross-links found in a body fluid such as urine. Type I telopeptide fragments may be isolated from a body fluid of any patient, however, it is preferred that these peptides are isolated from patients with Paget's disease or from rapidly growing adolescents, due to their high concentration of type I peptide fragments.

Isolation of Type I Collagen Telopeptides

Urine from patients with active Paget's disease is dialyzed in reduced porosity dialysis tubing (<3,500 mol. wt. cut off Spectropore) at 4 °C for 48h to remove bulk solutes. Under these conditions the peptides of interest are largely retained. The freeze-dried non-diffusate is then eluted (200 mg aliquots) from a column (90 cm x 2.5 cm) of Bio-Gel P2 (200-400 mesh) in 10% acetic acid at room temperature. A region of effluent that combines the cross-linked peptides is defined by measuring the fluorescence of collected fractions at 297 nm excitation/395 nm emission, and this pool is freeze-dried. Further resolution of this material is obtained on a column of Bio-Gel P-4 (200-400 mesh, 90 cm x 2.5 cm) eluted in 10% acetic acid.

Two contiguous fraction pools are defined by monitoring the fluorescence of the eluant above. The earlier fraction is enriched in peptide fragments having two amino acid sequences that derive from the C-terminal telopeptide domain of the $\alpha 1(I)$ chain of bone type I collagen linked to a third sequence derived from the triple-helical body of bone type I collagen. These three peptide sequences are cross-linked with 3-hydroxypyridinium. The overlapping later fraction is enriched in peptide fragments having an amino acid sequence that is derived from the N-terminal telopeptide domain of bone type I collagen linked through a 3-hydroxypyridinium cross-links.


Individual peptides are then resolved from each of the two fractions obtained above by ion-exchange HPLC on a TSK DEAE-5-PW column (Bio Rad 7.5 cm x 7.5 mm) eluting with a gradient of NaCl (0-0.2M) in 0.02M Tris-HCl, pH 7.5 containing 10% (v/v) acetonitrile. The N-terminal telopeptide-based and C-terminal telopeptide-based cross-linked peptides elute in a series of 3-4 peaks of fluorescence between 0.08M and 0.15M NaCl. The C-terminal telopeptide-based cross-linked peptides elute first as a series of fluorescent peaks, and the major and minor N-terminal telopeptide-based cross-linked peptides elute towards the end of the gradient as characteristic peaks. Each of these is collected, freeze-dried and chromatographed on a C-18 reverse phase HPLC column (Vydac 218TP54, 25 cm x 4.6 mm) eluted with a gradient (0-10%) of acetonitrile: n-propanol (3:1 v/v) in 0.01M trifluoroacetic acid. About 100-500g of individual peptide fragments containing 3-hydroxypyridinium cross-links can be isolated by this procedure from a single 24h collection of Paget's urine.

Amino acid compositions of the major isolated peptides confirmed purity and molecular sizes by the whole number stoichiometry of recovered amino acids. N-terminal sequence analysis by Edman degradation confirmed the basic core structures corresponding to the sequences of the known cross-linking sites in type I collagen and from the matching amino acid compositions. The N-terminal telopeptide sequence of the $\alpha 2(I)$ chain was blocked from sequencing analysis due presumably to the known cyclization of the N-terminal glutamine to pyrrolidone carboxylic acid.


A typical elution profile of N-terminal telopeptides obtained by the above procedure is shown in FIGURE 3A. The major peptide fragment obtained has an amino acid composition $(Asx)_2(Glx)_2(Gly)_5Val-Tyr-Ser-Thr$, where Asx is the amino acid Asp or Asn and Glx is the amino acid Gln or Glu. The sequence of this peptide is represented by Formula III below.

The C-terminal telopeptide-based cross-linked peptides resolved by reverse phase HPLC as described above are shown in FIGURE 3B. As can be seen from this figure, these peptides are further resolved into a series of C-terminal telopeptides each containing a 3-hydroxypyridinium cross-links. The major peptide, shown in Figure 3B, was analyzed as described above and was found to have the amino acid composition: $(Asp)_5(Glu)_4(Gly)_{10}(His)_2(Arg)_2(Hyp)_2(Ala)_5$. The sequence of this peptide is represented by formula IV below. It is believed that the other C-terminal telopeptide-based cross-linked peptides appearing as minor peaks in FIGURE 3B represent additions and deletions of amino acids to the structure shown in Formula IV.


FORMULA III



15 FORMULA IV



35 FORMULA V



and

55

as both homobifunctional and heterobifunctional reagents (see for example the Pierce 1986-87 catalog, Pierce Chemical Co., Rockford, IL). Preferred binding agents include carbodiimides and heterobifunctional reagents such as m-Maleimidobenzyl-N-hydroxysuccinimide ester (MBS).

5 Methods for binding the hapten to the carrier molecule are known in the art. See for example, Chard, T., *Laboratory Techniques in Biochemistry and Molecular Biology*, Vol. 6 (1987) Partz Elsevier, N.Y., herein incorporated by reference.

10 Either monoclonal or polyclonal antibodies to the hapten-carrier molecule immunogen can be produced. However, it is preferred that monoclonal antibodies (Mab) be prepared. For this reason it is preferred that immunization be carried out in the mouse. Immunization protocols for the mouse usually include an adjuvant. Examples of suitable protocols are described by Chard, T. (1987) *vid supra*. Spleen cells from the immunized mouse are harvested and homogenized and thereafter fused with cancer cells in the presence of polyethylene glycol to produce a fused cell hybrid which produces monoclonal antibodies specific to peptide fragments derived from collagen. Examples of such peptides are represented by the formulas given above. Suitable cancer cells include myeloma, hepatoma, carcinoma and sarcoma cells.

15 Detailed descriptions of this procedure, including screening protocols, protocols for growing selected hybrid cells and harvesting monoclonal antibodies produced by the selected hybrid cells are provided in Galfre, G. and Milstein, C., *Meth. Enzymol.*, 73:1 (1981). A preferred preliminary screening protocol involves the use of peptide fragments derived from bone collagen resorption and containing 3-hydroxypyridinium cross-links in a solid phase radioimmunoassay. A specific example is provided below.

20 Methods utilizing recombinant DNA techniques may also be adapted to construct monoclonal antibodies. These methods may be accomplished by the construction of expression libraries and primers by one of ordinary skill in the art (see William D. Huse et al., "Generation of a Large Combinational Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246:1275-1281, December 1989, see also L. Sastry et al., "Cloning of the Immunological Repertoire in *Escherichia coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library," *Proc. Natl. Acad. Sci. USA* 86:5728-5732, August 1989; see also Michelle Altine-Mees et al., "Monoclonal Antibody Expression Libraries: A Rapid Alternative to Hybridomas," *Strategies in Molecular Biology* 3:1-9, January 1990), or through the purchase of commercial kits which are available for this purpose (i.e., from Stratacyte, La Jolla, California). Briefly, within this embodiment, mRNA is isolated from a B cell population, and utilized

25 to create heavy and light chain immunoglobulin cDNA expression libraries in the λ ImmunoZap(H) and λ ImmunoZap(L) vectors. These vectors may be screened individually or co-expressed to form Fab fragments or antibodies (see Huse et al., *supra*; see also Sastry et al., *supra*). Positive plaques may subsequently be converted to a non-lytic plasmid which allows high level expression of monoclonal antibody fragments from *E. coli*.

35 The monoclonal antibodies or other immunological binding partners used in connection with the present invention are preferably specific for a particular type of collagen telopeptide.

40 However, in some cases, such selectivity will not be necessary, for example, if it is known that a patient is not suffering degradation of one type of collagen but is suspected of suffering degradation from the assayed type of collagen. Because of the differences in amino acid sequences between the type I, type II and type III families of telopeptides, cross-reactivity should not occur to a significant degree. Indeed, hybridomas can be selected for during the screening of splenocyte fusion clones that produce monoclonal antibodies specific for the cross-linked telopeptide of interest (and lack affinity for those of the other two collagen types). Based on the differences in sequence of the isolated peptide structures, such specificity is entirely feasible. Peptide fragments of the parent types I, II and III collagens, suitable for such hybridoma

45 screening, can be prepared from human bone, cartilage and other tissues and used to screen clones from mice immunized appropriately with the individual cross-linked peptide antigens isolated from body fluid.

Immunological binding partners, especially monoclonal antibodies, produced by the above procedures, or equivalent procedures, are employed in various immunometric assays to quantitate the concentration of the peptides having 3-hydroxypyridinium cross-links described above. These immunometric assays preferably comprise a monoclonal antibody or antibody fragment coupled to a detectable marker. Examples of suitable detectable markers include but are not limited to: enzymes, coenzymes, enzyme inhibitors, chromophores, fluorophores, chemiluminescent materials, paramagnetic metals, spin labels, and radionuclides. Examples of standard immunometric methods suitable for quantitating the telopeptides include, but are not limited to, enzyme linked immunosorbent assay (ELISA) (Ingvall, E., *Meth. Enzymol.*, 70 (1981)), radio-immunoassay (RIA), and "sandwich" immunoradiometric assay (IRMA).

55

In its simplest form, these immunometric methods can be used to determine the absolute rate of bone resorption or collagen degradation by simply contacting a body fluid with the immunological binding partner specific to a collagen telopeptide having a 3-hydroxypyridinium cross-link.

It is preferred that the immunometric assays described above be conducted directly on untreated body fluids (e.g. urine, blood, serum, or synovial fluid). Occasionally, however, contaminating substances may interfere with the assay necessitating partial purification of the body fluid. Partial purification procedures include, but are not limited to, cartridge adsorption and elution, molecular sieve chromatography, dialysis, ion exchange, alumina chromatography, hydroxyapatite chromatography and combinations thereof.

Test kits, suitable for use in accordance with the present invention, contain specific binding partners such as monoclonal antibodies prepared as described above, that specifically bind to peptide fragments derived from collagen degradation found in a body fluid. It is preferred that the specific binding partners of this test kit be coupled to a detectable marker of the type described above. Test kits containing a panel of two or more specific binding partners, particularly immunological binding partners, are also contemplated. Each immunological binding partner in such a test kit will preferably not cross-react substantially with a telopeptide derived from another type of collagen. For example, an immunological binding partner that binds specifically with a type I collagen telopeptide should preferably not cross-react with either a type II or type III collagen telopeptide. A small degree (e.g., 5-10%) of cross-reactivity may be tolerable. Other test kits may contain a first specific binding partner to a collagen-derived telopeptide having a cross-link containing a pyridinium ring (which may be OH-substituted), and a second specific binding partner to a telopeptide having the same structure as the first telopeptide except that the pyridinium ring has been cleaved, such as photolytically.

Monoclonal Antibody Production

The following is an example of preparation of a monoclonal antibody against a peptide immunogen.

A fraction enriched in the peptide (indicative of bone collagen degradation) is prepared from adolescent human urine using reverse phase and molecular sieve chromatography. The peptide is conjugated to keyhole limpet hemocyanin (KLH) with glutaraldehyde using standard procedures. Mice (Balb/c) are immunized subcutaneously with this conjugate (50-70 g), first in complete Freund's adjuvant, then boosted (25 μ g) at 3 weekly intervals in incomplete Freund's adjuvant intraperitoneally. After test bleeds show a high titer against the peptide conjugated to bovine serum albumin (BSA) using an ELISA format, selected mice are boosted with a low dose (5 μ g) of the immunogen in sterile PBS intravenously. Three days later, cells from the spleens of individual mice are fused with mouse myeloma cells using standard hybridoma technology. The supernatants of hybridoma clones growing in individual wells of 96-well plates are screened for reactive monoclonal antibodies, initially using a crude telopeptide preparation conjugated to BSA. After formal cloning by limiting dilution, the antibodies produced by individual hybridomas are characterized against a panel of screening antigens using ELISA analysis. An inhibition assay is used in which telopeptide conjugated to BSA is plated out in the plastic wells, and antibody is pre-incubated with a solution of the potential antigen. A secondary antibody (goat anti-mouse IgG conjugated to horseradish peroxidase, HRP) is used for color development using an appropriate substrate.

B. Electrochemical Procedure For Assaying For Peptides

An alternative procedure for assaying for the above-described peptides consists of measuring a physical property of the peptides having 3-hydroxypyridinium cross-links. One such physical property relies upon electrochemical detection. This method consists of injecting an aliquot of a body fluid, such as urine, into an electrochemical detector poised at a redox potential suitable for detection of peptides containing the 3-hydroxypyridinium ring. The 3-hydroxypyridinium ring, being aphenol, is subject to reversible oxidation and therefore the electrochemical detector (e.g., Model 5100A Coulochem sold by Esa 45 Wiggins Ave., Bedford, MA) is a highly desirable instrument suitable for quantitating the concentration of the present peptides. Two basic forms of electrochemical detector are currently commercially available: amperometric (e.g., BioAnalytical Systems) and coulometric (ESA, Inc., Bedford, MA 01730). Both are suitable for use in accordance with the present invention, however, the latter system is inherently more sensitive and therefore preferred since complete oxidation or reduction of the analyzed molecule in the column effluent is achieved. In addition, screening or guard electrodes can be placed "upstream" from the analytical electrode to selectively oxidize or reduce interfering substances thereby greatly improving selectivity. Essentially, the voltage of the analytical electrode is tuned to the redox potential of the sample molecule, and one or more pretreatment cells are set to destroy interferents in the sample.

In a preferred assay method, a standard current/voltage curve is established for standard peptides containing lysyl pyridinoline or hydroxyllysyl pyridinoline in order to determine the proper voltage to set for optimal sensitivity. This voltage is then modified depending upon the body fluid, to minimize interference

from contaminants and optimize sensitivity. Electrochemical detectors, and the optimum conditions for their use are known to those skilled in the art. Complex mixtures of body fluids can often be directly analyzed with the electrochemical detector without interference. Accordingly, for most patients no pretreatment of the body fluid is necessary. In some cases however, interfering compounds may reduce the reliability of the measurements. In such cases, pretreatment of the body fluid (e.g., urine) may be necessary.

Accordingly, in an alternative embodiment of the invention, a body fluid is first purified prior to electrochemically titrating the purified peptide fragments. The purification step may be conducted in a variety of ways including but not limited to dialysis, ion exchange chromatography, alumina chromatography, hydroxyapatite chromatography, molecular sieve chromatography, or combinations thereof. In a preferred purification protocol, a measured aliquot (25 ml) of a 24 hour urine sample is dialyzed in reduced porosity dialysis tubing to remove the bulk of contaminating fluorescent solutes. The non-diffusate is then lyophilized, redissolved in 1% heptafluorobutyric acid (HFBA), an ion pairing solution, and the peptides adsorbed on a Waters Sep-Pak C-18 cartridge. This cartridge is then washed with 5 ml of 1% HFBA, and then eluted with 3 ml of 50% methanol in 1% HFBA.

Another preferred method of purification consists of adsorbing a measured aliquot of urine onto an ion-exchange adsorption filter and eluting the adsorption filter with a buffered eluting solution. The eluate fractions containing peptide fragments having 3-hydroxypyridinium cross-links are then collected to be assayed.

Still another preferred method of purification employs molecular sieve chromatography. For example, an aliquot of urine is applied to a Bio-Gel P2 or Sephadex G-20 column and the fraction eluting in the 1000-5000 Dalton range is collected. It will be obvious to those skilled in the art that a combination of the above methods may be used to purify or partially purify urine or other body fluids in order to isolate the peptide fragments having 3-hydroxypyridinium cross-links. The purified or partially purified peptide fragments obtained by the above procedures may be subjected to additional purification procedures, further processed or assayed directly in the partially purified state. Additional purification procedures include resolving partially purified peptide fragments employing high performance liquid chromatography (HPLC) or micro-bore HPLC when increased sensitivity is desired. These peptides may then be quantitated by electrochemical titration.

A preferred electrochemical titration protocol consists of tuning the redox potential of the detecting cell of the electrochemical detector (Coulchem Model 5100A) for maximum signal with pure HP. The detector is then used to monitor the effluent from a C-18 HPLC column used to resolve the partially purified peptides.

FIGURE 4A displays the elution profile resolved by reverse phase HPLC of natural fluorescence for a hydrolysate of peptide fragments from normal human urine. Measurement of the integrated area within the envelope of a given component is used to determine the concentration of that component within the sample. The ratio of HP:LP found in normal human urine and urine from patients having Paget's disease, FIGURE 4B, are both approximately 4.5:1. This is slightly higher than the 4:1 ratio found in bone itself (Eyre, et al., 1984). The higher ratio found in urine indicates that a portion of the HP fraction in urine may come from sources other than bone, such as the diet, or other sources of collagen degradation, i.e., cartilage catabolism. It is for this reason that it is preferred that LP which derives only from bone be used to provide an absolute index of bone resorption. However, in the absence of excessive cartilage degradation such as in rheumatoid arthritis or in cases where bone is rapidly being absorbed, HP or a combination of HP plus LP may be used as an index of bone resorption.

While the invention has been described in conjunction with preferred embodiments, one of ordinary skill after reading the foregoing specification will be able to effect various changes, substitutions or equivalents, and alterations to the subject matter set forth herein. Hence, the invention can be practiced in ways other than those specifically described herein. It is therefore intended that the protection granted by Letters Patent hereon be limited only by the appended claims and equivalents thereof.

Claims

1. An assay for measuring bone resorption, by determining in a sample of body fluid the presence of concentration of a carboxy-terminal type I collagen telopeptide comprising:




Figure 1




Figure 2




Figure 3A




Figure 3B




Figure 4B




Figure 4A



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.5)
A	WO-A-89 04491 (WASHINGTON RESEARCH FOUNDATION) ---		G01N33/68 C12N5/20 C12P21/08 C07K16/18
D,A	ANNALS OF THE RHEUMATIC DISEASES, vol. 48, no. 8, August 1989 LONDON, GB, pages 641-644, D. BLACK ET AL. 'Urinary excretion of the hydroxypyridinium cross links of collagen in patients with rheumatoid arthritis.' ---		
D,A	GB-A-2 205 643 (FAMOS GROUP LTD.) ---		
P,D, A	WO-A-89 12824 (THE ROWETT RESEARCH INSTITUTE) -----		
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.5)
			G01N C07K
Place of search	Date of completion of the search	Examiner	
THE HAGUE	16 June 1995	Griffith, G	
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			