



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2679310 A1 2008/08/28

(21) **2 679 310**

(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2008/02/20
(87) Date publication PCT/PCT Publication Date: 2008/08/28
(85) Entrée phase nationale/National Entry: 2009/08/20
(86) N° demande PCT/PCT Application No.: EP 2008/052031
(87) N° publication PCT/PCT Publication No.: 2008/101939
(30) Priorité/Priority: 2007/02/21 (EP07102826.0)


(51) Cl.Int./Int.Cl. *C07D 309/10* (2006.01),
A61K 31/70 (2006.01), *C07H 7/04* (2006.01)

(71) Demandeur/Applicant:
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,
DE

(72) Inventeurs/Inventors:
HIMMELSBACH, FRANK, DE;
ECKHARDT, MATTHIAS, DE;
EICKELMANN, PETER, DE;
THOMAS, LEO, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : DERIVES DE BENZENE GLUCOPYRANOSYLES TETRASUBSTITUES, MEDICAMENTS CONTENANT
LESDITS COMPOSES, UTILISATION ET PROCEDES DE FABRICATION DE CES DERNIERS
(54) Title: TETRASUBSTITUTED GLUCOPYRANOSYLATED BENZENE DERIVATIVES, MEDICAMENTS CONTAINING
SUCH COMPOUNDS, THEIR USE AND PROCESS FOR THEIR MANUFACTURE



(57) Abrégé/Abstract:

Tetrasubstituted glucopyranosylated benzene derivatives of general formula (I) as defined according to claim 1, including the tautomers, the stereoisomers thereof, the mixtures thereof and the salts thereof. The compounds according to the invention are suitable for the treatment of metabolic disorders.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 August 2008 (28.08.2008)

PCT

(10) International Publication Number
WO 2008/101939 A1

(51) International Patent Classification:

C07D 309/10 (2006.01) A61K 31/70 (2006.01)
C07H 7/04 (2006.01)

THOMAS, Leo [DE/DE]; Boehringer Ingelheim GmbH, CD Patents, Binger Strasse 173, 55216 Ingelheim Am Rhein (DE).

(21) International Application Number:

PCT/EP2008/052031

(74) Agents: HAMMANN, Heinz et al.; Boehringer Ingelheim GmbH, Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(22) International Filing Date:

20 February 2008 (20.02.2008)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

07102826.0 21 February 2007 (21.02.2007) EP


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US):
BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim Am Rhein (DE).

Published:

— with international search report

(54) Title: TETRASUBSTITUTED GLUCOPYRANOSYLATED BENZENE DERIVATIVES, MEDICAMENTS CONTAINING SUCH COMPOUNDS, THEIR USE AND PROCESS FOR THEIR MANUFACTURE




(57) Abstract: Tetrasubstituted glucopyranosylated benzene derivatives of general formula (I) as defined according to claim 1, including the tautomers, the stereoisomers thereof, the mixtures thereof and the salts thereof. The compounds according to the invention are suitable for the treatment of metabolic disorders.

WO 2008/101939 A1

Tetrasubstituted glucopyranosylated benzene derivatives, medicaments containing such compounds, their use and process for their manufacture

The present invention relates to tetrasubstituted glucopyranosylated benzene derivatives of
5 the general formula I



wherein the groups R¹ to R⁶ and R^{7a}, R^{7b}, R^{7c} are as defined hereinafter, including the
10 tautomers, the stereoisomers, the mixtures thereof and the salts thereof. The invention further relates to pharmaceutical compositions containing a compound of formula I according to the invention as well as the use of a compound according to the invention for preparing a pharmaceutical composition for the treatment of metabolic disorders. In addition, the invention relates to processes for preparing a pharmaceutical composition as well as a
15 compound according to the invention.

In the literature, compounds which have an inhibitory effect on the sodium-dependent glucose cotransporter SGLT2 are proposed for the treatment of diseases, particularly diabetes.

20

Glucopyranosyloxy- and glucopyranosyl-substituted aromatic groups and the preparation thereof and their possible activity as SGLT2 inhibitors are known from published International applications WO 98/31697, WO 01/27128, WO 02/083066, WO 03/099836, WO 2004/063209, WO 2004/080990, WO 2004/013118, WO 2004/052902, WO 2004/052903,
25 WO 2005/092877, WO 06/010557, WO 06/018150, WO 06/037537, WO 06/089872, WO 2006/064033, WO 06/108842, WO 2006/120208, US application US 2003/0114390, WO 2006/117360, WO 2007/014894, WO 2007/025943, WO 2007/028814, WO 2007/031548, WO 2006/117359, WO 2007/093610, WO 2007/128749.

Aim of the invention

The aim of the present invention is to find new pyranosyl-substituted benzene derivatives, particularly those which are active with regard to the sodium-dependent glucose cotransporter SGLT, particularly SGLT2. A further aim of the present invention is to find

5 pyranosyl-substituted benzene derivatives which have a good to very good inhibitory effect on the sodium-dependent glucose cotransporter SGLT2 *in vitro* and/or *in vivo* and/or have good to very good pharmacological and/or pharmacokinetic and/or physicochemical properties.

10 A further aim of the present invention is to provide new pharmaceutical compositions which are suitable for the prevention and/or treatment of metabolic disorders, particularly diabetes.


The invention also sets out to provide a process for preparing the compounds according to the invention.

15

Other aims of the present invention will become apparent to the skilled man directly from the foregoing and following remarks.

Object of the invention

20 In a first aspect the present invention relates to tetrasubstituted glucopyranosylated benzene derivatives of general formula I



wherein

25

R^1 denotes chlorine or methyl,

R^2 denotes C_{3-7} -cycloalkyl or C_{3-7} -cycloalkyloxy,

30

while the above-mentioned cycloalkyl-rings may be mono- or polysubstituted by fluorine and/or mono- or disubstituted by identical or different substituents L^2 , and/or

in the above-mentioned cycloalkyl rings one methylene group may be replaced by O, and

R³ hydrogen, fluorine, chlorine, bromine, iodine, C₁₋₆-alkyl, C₂₋₆-alkynyl, C₂₋₆-alkenyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl, C₅₋₇-cycloalkenyl, C₅₋₇-cycloalkenyl-C₁₋₃-alkyl, aryl, heteroaryl, C₁₋₄-alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, aminocarbonyl, C₁₋₄-alkylaminocarbonyl, di-(C₁₋₃-alkyl)aminocarbonyl, pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl, morpholin-4-ylcarbonyl, piperazin-1-ylcarbonyl, 4-(C₁₋₄-alkyl)piperazin-1-ylcarbonyl, hydroxycarbonyl, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkyl-amino, di-(C₁₋₃-alkyl)amino, pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, piperazin-1-yl, 4-(C₁₋₄-alkyl)piperazin-1-yl, C₁₋₄-alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, C₁₋₄-alkylsulfonylamino, arylsulfonylamino, C₁₋₆-alkoxy, C₃₋₇-cycloalkyloxy, C₅₋₇-cycloalkenyloxy, aryloxy, heteroaryloxy, C₁₋₄-alkylsulfanyl, C₁₋₄-alkylsulfinyl, C₁₋₄-alkylsulfonyl, C₃₋₇-cycloalkylsulfanyl, C₃₋₇-cycloalkylsulfinyl, C₃₋₇-cycloalkylsulfonyl, C₅₋₇-cycloalkenylsulfanyl, C₅₋₇-cycloalkenylsulfinyl, C₅₋₇-cycloalkenylsulfonyl, arylsulfanyl, arylsulfinyl, arylsulfonyl, heteroarylsulfanyl, heteroarylsulfinyl, heteroarylsulfonyl, amino, hydroxy, cyano and nitro,

while the above-mentioned alkyl-, alkenyl-, alkynyl-, cycloalkyl- und cycloalkenyl-residues may be mono- or polysubstituted by fluorine and/or mono- or disubstituted by identical or different substituents L₂, and

while in the above-mentioned C₅₋₆-cycloalkyl and C₅₋₆-cycloalkenyl rings one or two methylene groups may be replaced independently of one another by O, S, CO, SO or SO₂, and

while in the above-mentioned N-heterocycloalkyl rings one methylene group may be replaced by CO or SO₂, and

R⁴, R⁵ independently of one another denote hydrogen, fluorine, chlorine, bromine, iodine, cyano, nitro, C₁₋₃-alkyl, C₁₋₃-alkoxy, or a methyl- or methoxy-group substituted by 1 to 3 fluorine atoms,

L₁ independently of one another are selected from among fluorine, chlorine, bromine, iodine, hydroxy, cyano, C₁₋₃-alkyl, difluoromethyl, trifluoromethyl, C₁₋₃-alkoxy, difluoromethoxy, trifluoromethoxy, amino, C₁₋₃-alkyl-amino and di(C₁₋₃-alkyl)-amino; and

5 L2 independently of one another are selected from among fluorine, chlorine, hydroxy, hydroxyl-C₁₋₄-alkyl, C₁₋₄-alkoxy, trifluoromethoxy, C₁₋₄-alkoxy-C₁₋₄-alkyl, cyano, hydroxycarbonyl, (C₁₋₄-alkyl)oxycarbonyl, aminocarbonyl, C₁₋₄-alkyl, trifluoromethyl, amino, C₁₋₄-alkyl-carbonylamino, C₁₋₃-alkyl-amino and di(C₁₋₃-alkyl)-amino; and

R⁶, R^{7a},

10 R^{7b}, R^{7c} independently of one another have a meaning selected from among hydrogen, (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl and aryl-(C₁₋₃-alkyl)- carbonyl, while the aryl-groups may be mono- or disubstituted independently of one another by identical or different groups L1;

15 while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups which may be substituted as defined; and

20 while, unless otherwise stated, the above-mentioned alkyl groups may be straight-chain or branched,

25 the tautomers, the stereoisomers thereof, the mixtures thereof and the salts thereof.

30 The compounds of general formula I according to the invention and the physiologically acceptable salts thereof have valuable pharmacological properties, particularly an inhibitory effect on the sodium-dependent glucose cotransporter SGLT, particularly SGLT2. Moreover compounds according to the invention may have an inhibitory effect on the sodium- dependent glucose cotransporter SGLT1. Compared with a possible inhibitory effect on SGLT1 the compounds according to the invention preferably inhibit SGLT2 selectively.

35 The present invention also relates to the physiologically acceptable salts of the compounds according to the invention with inorganic or organic acids.

40 This invention also relates to pharmaceutical compositions, containing at least one compound according to the invention or a physiologically acceptable salt according to the invention, optionally together with one or more inert carriers and/or diluents.

45 This invention also relates to the use of at least one compound according to the invention or one of the physiologically acceptable salts thereof for preparing a pharmaceutical composition which is suitable for the treatment or prevention of diseases or conditions which

can be influenced by inhibiting the sodium-dependent glucose cotransporter SGLT, particularly SGLT2.

This invention also relates to the use of at least one compound according to the invention or 5 one of the physiologically acceptable salts thereof for preparing a pharmaceutical composition which is suitable for the treatment of metabolic disorders.

In a further aspect the present invention relates to the use of at least one compound according to the invention or one of the physiologically acceptable salts thereof for preparing 10 a pharmaceutical composition for preventing the degeneration of pancreatic beta cells and/or for improving and/or restoring the functionality of pancreatic beta cells.

In a further aspect the present invention relates to a use of at least one compound according to the invention or one of the physiologically acceptable salts thereof for preparing a 15 pharmaceutical composition for preventing, slowing, delaying or treating diseases or conditions attributed to an abnormal accumulation of liver fat in a patient in need thereof.

This invention also relates to the use of at least one compound according to the invention or one of the physiologically acceptable salts thereof for preparing a pharmaceutical 20 composition for inhibiting the sodium-dependent glucose cotransporter SGLT, particularly SGLT2.

The invention further relates to a process for preparing a pharmaceutical composition according to the invention, characterised in that a compound according to the invention or 25 one of the physiologically acceptable salts thereof is incorporated in one or more inert carriers and/or diluents by a non-chemical method.


The present invention also relates to a process for preparing the compounds of general formula I according to the invention, characterised in that

30

a) in order to prepare compounds of general formula I which are defined as hereinbefore and hereinafter,

a compound of general formula II

35



II

wherein

R' denotes H, C₁₋₄-alkyl, (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl and aryl-(C₁₋₃-alkyl)-carbonyl, wherein the alkyl or aryl groups may be mono- or polysubstituted by halogen;

R^{8a}, R^{8b}, R^{8c}, R^{8d} independently of one another have one of the meanings given hereinbefore and hereinafter for the groups R⁶, R^{7a}, R^{7b}, R^{7c}, or denote a benzyl or allyl group or a R^aR^bR^cSi group or a ketal or acetal group, particularly an alkylidene or arylalkylidene ketal or acetal group, while in each case two adjacent groups R^{8a}, R^{8b}, R^{8c}, R^{8d} may form a cyclic silyl ketal, ketal or acetal group or a 1,2-di(C₁₋₃-alkoxy)-1,2-di(C₁₋₃-alkyl)-ethylene bridge, while the above-mentioned ethylene bridge forms, together with two oxygen atoms and the two associated carbon atoms of the pyranose ring, a substituted dioxane ring, particularly a 2,3-dimethyl-2,3-di(C₁₋₃-alkoxy)-1,4-dioxane ring, and while alkyl, aryl and/or benzyl groups may be mono- or polysubstituted by halogen or C₁₋₃-alkoxy, and while benzyl groups may also be substituted by a di-(C₁₋₃-alkyl)amino group; and

R^a, R^b, R^c independently of one another denote C₁₋₄-alkyl, aryl or aryl-C₁₋₃-alkyl, wherein the aryl or alkyl groups may be mono- or polysubstituted by halogen;

while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups, preferably phenyl groups;


and wherein the groups R¹ to R⁶, R^{7a}, R^{7b}, R^{7c} are defined as hereinbefore and hereinafter;

is reacted with a reducing agent in the presence of a Lewis or Brønsted acid, while any protective groups present are cleaved simultaneously or subsequently; or

b) in order to prepare compounds of general formula I wherein R^6 , R^{7a} , R^{7b} and R^{7c} denote hydrogen,

a compound of general formula III

5



wherein R^{8a} , R^{8b} , R^{8c} , R^{8d} and R^1 to R^5 are defined as hereinbefore and hereinafter, but at least one of the groups R^{8a} , R^{8b} , R^{8c} , R^{8d} does not denote hydrogen, is hydrolysed, and

10

if desired a compound of general formula I thus obtained wherein R^6 denotes a hydrogen atom, is converted by acylation into a corresponding acyl compound of general formula I, and/or


15 if necessary any protective group used in the reactions described above is cleaved and/or

if desired a compound of general formula I thus obtained is resolved into its stereoisomers and/or

20 if desired a compound of general formula I thus obtained is converted into the salts thereof,
particularly for pharmaceutical use into the physiologically acceptable salts thereof.

This invention further relates to a process for preparing compounds of general formula II

25



wherein

R' denotes H, C₁₋₄-alkyl, (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl and aryl-(C₁₋₃-alkyl)-carbonyl, wherein the alkyl or aryl groups may be mono- or polysubstituted by halogen;

R^{8a}, R^{8b},


R^{8c}, R^{8d} independently of one another has one of the meanings given for the groups R⁶, R^{7a}, R^{7b}, R^{7c}, or denote a benzyl or allyl group or a R^aR^bR^cSi group or a ketal or acetal group, while in each case two adjacent groups R^{8a}, R^{8b}, R^{8c}, R^{8d} may form a cyclic silyl ketal, ketal or acetal group or may form, with two oxygen atoms of the pyranose ring, a substituted 2,3-oxydioxane ring, particularly a 2,3-dimethyl-2,3-di(C₁₋₃-alkoxy)-1,4-dioxane ring, and while alkyl, aryl and/or benzyl groups may be mono- or polysubstituted by halogen or C₁₋₃-alkoxy, and while benzyl groups may also be substituted by a di-(C₁₋₃-alkyl)amino group; and

R^a, R^b, R^c independently of one another denote C₁₋₄-alkyl, aryl or aryl-C₁₋₃-alkyl, while the alkyl or aryl groups may be mono- or polysubstituted by halogen;

while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups, preferably phenyl groups;


and R¹ to R⁶, R^{7a}, R^{7b}, R^{7c} are defined as hereinbefore and hereinafter,

wherein an organometallic compound (V) which may be obtained by halogen-metal exchange or by inserting a metal in the carbon-halogen bond of a halogen-benzylbenzene compound of general formula IV



30

wherein Hal denotes Cl, Br and I and R¹ to R⁵ are defined as hereinbefore and hereinafter, and optionally subsequent transmetallation, is added to a gluconolactone of general formula VI



VI

wherein R^{8a}, R^{8b}, R^{8c}, R^{8d} are defined as hereinbefore and hereinafter,

5 and

then the resulting adduct is reacted with water or an alcohol R'-OH, while R' denotes optionally substituted C₁₋₄-alkyl, in the presence of an acid, such as for example methanesulfonic acid, sulfuric acid, hydrochloric acid, acetic acid or ammonium chloride, and 10 optionally the product obtained in the reaction with water wherein R' denotes H is converted, in a subsequent reaction, with an alcohol in the presence of an acid to yield the alkoxy derivative or with an acylating agent, such as for example the corresponding acid chloride or anhydride, into the product of formula II wherein R' denotes (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl or aryl-(C₁₋₃-alkyl)-carbonyl, which may be substituted as 15 specified.

The intermediate products listed, particularly those of formula IV, formula II and formula III, are also a subject of this invention.

20

Detailed Description of the invention

Unless otherwise stated, the groups, residues and substituents, particularly R¹ to R⁵, L1, L2, R⁶, R^{7a}, R^{7b}, R^{7c}, R^{8a}, R^{8b}, R^{8c}, R^{8d}, are defined as above and hereinafter.

25 If residues, substituents or groups occur several times in a compound, as for example L1 and L2, they may have the same or different meanings.

Some preferred meanings of individual groups and substituents of the compounds according to the invention will be given hereinafter.

30

The group R¹ denotes chlorine or methyl.

The group R² preferably denotes C₃₋₇-cycloalkyl or C₃₋₇-cycloalkyloxy, while in the cycloalkyl rings a methylene group may be replaced by O, and wherein the cycloalkyl rings may be mono- or poly-fluorinated and/or mono- or disubstituted with identical or different substituents L2.

5

Even more preferred meanings of R² are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydrofuranyl, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, tetrahydrofuranyloxy; particularly cyclopropyl, cyclobutyl, cyclopentyl, cyclopropyloxy and cyclobutyloxy, cyclopentyloxy, (S)-tetrahydrofuran-2-yl, (R)-tetrahydrofuran-2-yl, (S)-tetrahydrofuran-3-yl, (R)-tetrahydrofuran-3-yl, ((S)-tetrahydrofuran-3-yl)oxy, ((R)-tetrahydrofuran-3-yl)oxy.

Preferred meanings of the group R³ are chlorine, bromine, iodine, C₁₋₄-alkyl, C₃₋₇-cycloalkyl, hydroxyl, C₁₋₄-alkyloxy, C₃₋₇-cycloalkyloxy, C₁₋₄-alkylsulfanyl, C₃₋₇-cycloalkylsulfanyl, while in a C₅₋₆-cycloalkyl ring a methylene group may be replaced by O, and wherein any alkyl group and cycloalkyl ring may be mono- or polyfluorinated and/or mono- or disubstituted with identical or different substituents L2.

More preferred meanings of the group R³ are chlorine, bromine, methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, sec-butyl, iso-butyl, tert-butyl, difluoromethyl, trifluoromethyl, 2-hydroxyl-ethyl, hydroxymethyl, 3-hydroxy-propyl, 2-hydroxy-2-methyl-prop-1-yl, 3-hydroxy-3-methyl-but-1-yl, 1-hydroxy-1-methyl-ethyl, 2-methoxy-ethyl, 2-ethoxy-ethyl, hydroxyl, methoxy, ethoxy, isopropoxy, difluoromethoxy, trifluoromethoxy, cyclobutoxy, cyclopentoxy, cyclohexyloxy, (S)-tetrahydrofuran-3-yloxy, (R)-tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, methylsulfanyl and ethylsulfanyl.

Even more preferably R³ denotes methyl, ethyl, n-propyl, isopropyl, cyclopropyl, methoxy, ethoxy, isopropoxy, methylsulfanyl or ethylsulfanyl, in particular ethyl or cyclopropyl.

30 Preferred meanings of the group R⁴ are hydrogen and fluorine, particularly hydrogen.

Preferred meanings of the group R⁵ are hydrogen and fluorine, particularly hydrogen.

Preferred meanings of the group L1 independently of one another are selected from among fluorine, chlorine, bromine, cyano, hydroxy, C₁₋₃-alkyl, difluoromethyl, trifluoromethyl, C₁₋₃-alkoxy, difluoromethoxy, trifluoromethoxy and di(C₁₋₃-alkyl)-amino.

Even more preferred meanings of the group L1 are selected from fluorine, chlorine, hydroxy, trifluoromethyl, ethyl, methoxy, ethoxy and dimethylamino, particularly methyl, ethyl, methoxy, ethoxy and dimethylamino.

5 Preferred meanings of the group L2 independently of one another are selected from among fluorine, hydroxy, hydroxy-C₁₋₄-alkyl, C₁₋₄-alkoxy, C₁₋₄-alkoxy-C₁₋₄-alkyl, C₁₋₄-alkyl, trifluoromethyl, C₁₋₄-alkyl-carbonylamino, hydroxycarbonyl and C₁₋₄-alkoxycarbonyl.




Even more preferred meanings of the group L2 are selected from fluorine, hydroxy, hydroxy-C₁₋₄-alkyl, C₁₋₄-alkoxy, C₁₋₄-alkoxy-C₁₋₄-alkyl, C₁₋₄-alkyl, hydroxycarbonyl and C₁₋₄-alkoxy-carbonyl; particularly hydroxy, hydroxymethyl, methoxymethyl, methoxy, methyl, hydroxycarbonyl, methoxycarbonyl and ethoxycarbonyl.

The group R⁶ preferably denotes according to the invention hydrogen, (C₁₋₈-alkyl)oxygen-carbonyl, C₁₋₈-alkylcarbonyl or benzoyl, particularly hydrogen or (C₁₋₆-alkyl)oxycarbonyl or C₁₋₆-alkylcarbonyl, particularly preferably hydrogen, methylcarbonyl, methoxycarbonyl or ethoxycarbonyl, most particularly preferably hydrogen.

The substituents R^{7a}, R^{7b}, R^{7c} preferably represent independently of one another hydrogen, (C₁₋₈-alkyl)oxycarbonyl, (C₁₋₁₈-alkyl)carbonyl or benzoyl, particularly hydrogen, (C₁₋₆-alkyl)oxygen-carbonyl or (C₁₋₈-alkyl)carbonyl, particularly preferably hydrogen, methoxycarbonyl, ethoxycarbonyl, methylcarbonyl or ethylcarbonyl. Most particularly preferably R^{7a}, R^{7b} and R^{7c} represent hydrogen.

25 The compounds of formula I wherein R⁶, R^{7a}, R^{7b} and R^{7c} according to the invention have a meaning other than hydrogen, for example C₁₋₈-alkylcarbonyl, are preferably suitable as intermediate products for the synthesis of compounds of formula I wherein R⁶, R^{7a}, R^{7b} and R^{7c} denote hydrogen.

30 Particularly preferred compounds of general formula I are selected from among formulae I.1 to I.3, particularly I.1 and I.2:



while the groups R¹ to R⁶ and R^{7a}, R^{7b}, R^{7c} have one of the meanings given previously, particularly have one of the given meanings specified as being preferred; and particularly

R¹ denotes chlorine or methyl,

5

R² denotes C₃₋₇-cycloalkyl or C₃₋₇-cycloalkyloxy, while in the cycloalkyl rings a methylene group may be replaced by O, and wherein the cycloalkyl rings may be mono- or poly-fluorinated and/or mono- or disubstituted with identical or different substituents L2; even more preferably R² denotes cyclopropyl, cyclobutyl, cyclopentyl, tetrahydrofuranyl, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, tetrahydrofuranyloxy; and

10

R³ denotes chlorine, bromine, iodine, C₁₋₄-alkyl, C₃₋₇-cycloalkyl, hydroxyl, C₁₋₄-alkyloxy, C₃₋₇-cycloalkyloxy, C₁₋₄-alkylsulfanyl, C₃₋₇-cycloalkylsulfanyl, while in a C₅₋₆-cycloalkyl ring a methylene group may be replaced by O; and wherein any alkyl group and cycloalkyl ring may be mono- or polyfluorinated and/or mono- or disubstituted with identical or different substituents L2; even more preferably R³ denotes chlorine,

15

bromine, methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, sec-butyl, iso-butyl, tert-butyl, difluoromethyl, trifluoromethyl, 2-hydroxyl-ethyl, hydroxymethyl, 3-hydroxy-propyl, 2-hydroxy-2-methyl-prop-1-yl, 3-hydroxy-3-methyl-but-1-yl, 1-hydroxy-1-methyl-ethyl, 2-methoxy-ethyl, 2-ethoxy-ethyl, hydroxyl, methoxy, ethoxy, isopropoxy, difluoromethoxy, trifluoromethoxy, cyclobutoxy, cyclopentoxy, cyclohexyloxy, (S)-tetrahydrofuran-3-yloxy, (R)-tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, methylsulfanyl and ethylsulfanyl; most preferably R^3 denotes methyl, ethyl, n-propyl, isopropyl, cyclopropyl, methoxy, ethoxy, isopropoxy, methylsulfanyl or ethylsulfanyl; and

10

R^4 denotes hydrogen or fluorine, particularly hydrogen; and

R^5 denotes hydrogen or fluorine, particularly hydrogen; and

15 L2 independently of one another are selected from among fluorine, hydroxy, hydroxy- C_{1-4} -alkyl, C_{1-4} -alkoxy, C_{1-4} -alkoxy- C_{1-4} -alkyl, C_{1-4} -alkyl, trifluoromethyl, C_{1-4} -alkyl-carbonyl, hydroxycarbonyl and C_{1-4} -alkyloxycarbonyl; particularly hydroxy, hydroxymethyl, methoxymethyl, methoxy, methyl, hydroxycarbonyl, methoxycarbonyl and ethoxycarbonyl; and

20

R^6 denotes hydrogen, (C_{1-6} -alkyl)oxycarbonyl, (C_{1-6} -alkyl)carbonyl or benzoyl, particularly hydrogen, methylcarbonyl, methoxycarbonyl or ethoxycarbonyl, most particularly preferably hydrogen; and

25 R^{7a} , R^{7b} , R^{7c} independently of one another represent hydrogen, (C_{1-6} -alkyl)oxycarbonyl, (C_{1-8} -alkyl)carbonyl or benzoyl, particularly hydrogen, methoxycarbonyl, ethoxycarbonyl, methylcarbonyl or ethylcarbonyl, particularly preferably hydrogen;


including the tautomers, the stereoisomers, the mixtures thereof and the salts thereof.

30

The compounds of general formula I specified in the experimental section that follows, and the derivatives thereof, wherein R^6 has a meaning according to the invention other than hydrogen, particularly wherein R^6 denotes acetyl, ethoxycarbonyl or methoxycarbonyl, 35 including the tautomers, the stereoisomers thereof and the mixtures thereof, are preferred according to another variant of this invention.


In the processes according to the invention the groups R¹ to R⁵ preferably have the meanings specified hereinbefore as being preferred. Moreover R' preferably denotes H, C₁₋₃-alkyl or benzyl, particularly H, ethyl or methyl. The groups R^{8a}, R^{8b}, R^{8c} and R^{8d} independently of one another preferably denote H, C₁₋₄-alkylcarbonyl or benzyl, particularly H, 5 methylcarbonyl, ethylcarbonyl or benzyl.

The invention also relates to compounds of general formula IV



wherein Hal denotes chlorine, bromine or iodine, and the groups R¹ to R⁵ are as hereinbefore defined, as intermediate products or starting materials in the synthesis of the compounds according to the invention. Particularly preferably, the groups R¹ to R⁵ have the meanings given following formulae I.1 and I.2.

The invention also relates to compounds of general formula II



wherein R', R^{8a}, R^{8b}, R^{8c}, R^{8d} and R¹ to R⁵ are defined as hereinbefore and hereinafter; particularly wherein R' denotes H, C₁₋₃-alkyl or benzyl, particularly H, ethyl or methyl; and the groups R^{8a}, R^{8b}, R^{8c} and R^{8d} independently of one another represent H, C₁₋₄-alkylcarbonyl, allyl or benzyl, particularly H, methylcarbonyl, ethylcarbonyl or benzyl and the groups R¹ to R⁵ are as hereinbefore defined. These compounds may serve as intermediate products or starting materials in the synthesis of the compounds according to the invention. Particularly preferably the groups R¹ to R⁵ have the meanings given following formulae I.1 to I.2.

Some terms used above and hereinafter to describe the compounds according to the invention will now be defined more closely.

The term halogen denotes an atom selected from the group consisting of F, Cl, Br and I.

The term C_{1-n} -alkyl, wherein n may have a value of 2 to 18, denotes a saturated, branched or unbranched hydrocarbon group with 1 to n C atoms. Examples of such groups include methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, neo-pentyl, tert-pentyl, n-hexyl, iso-hexyl, etc.

5

The term C_{2-n} -alkynyl, wherein n has a value of 3 to 6, denotes a branched or unbranched hydrocarbon group with 2 to n C atoms and a $C\equiv C$ triple bond. Examples of such groups include ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl etc.

10 Unless otherwise stated alkynyl groups are connected to the remainder of the molecule via the C atom in position 1. Therefore terms such as 1-propynyl, 2-propynyl, 1-butynyl, etc. are equivalent to the terms 1-propyn-1-yl, 2-propyn-1-yl, 1-butyn-1-yl, etc.. This also applies analogously to C_{2-n} -alkenyl groups.

15 The term C_{1-n} -alkoxy denotes a C_{1-n} -alkyl-O group, wherein C_{1-n} -alkyl is as hereinbefore defined. Examples of such groups include methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, n-pentoxyl, iso-pentoxyl, neo-pentoxyl, tert-pentoxyl, n-hexoxy, iso-hexoxy etc.

20 The term C_{1-n} -alkylcarbonyl denotes a C_{1-n} -alkyl-C(=O) group, wherein C_{1-n} -alkyl is as hereinbefore defined. Examples of such groups include methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, iso-propylcarbonyl, n-butylcarbonyl, iso-butylcarbonyl, sec-butylcarbonyl, tert-butylcarbonyl, n-pentylcarbonyl, iso-pentylcarbonyl, neo-pentylcarbonyl, tert-pentylcarbonyl, n-hexylcarbonyl, iso-hexylcarbonyl, etc.

25

The term C_{3-n} -cycloalkyl denotes a saturated mono-, bi-, tri- or spirocyclic group with 3 to n C atoms. Examples of such groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, decalinyl, bicyclo[3.2.1.]octyl, spiro[4.5]decyl, norpinyl, norbonyl, norcaryl, adamantyl, etc. Preferably the term C_{3-n} -cycloalkyl denotes saturated monocyclic groups.

30 The term C_{5-n} -cycloalkenyl denotes a C_{5-n} -cycloalkyl group which is as hereinbefore defined and additionally has at least one unsaturated C=C double bond.

35 The term C_{3-n} -cycloalkylcarbonyl denotes a C_{3-n} -cycloalkyl-C(=O) group wherein C_{3-n} -cycloalkyl is as hereinbefore defined.

The term tri-(C₁₋₄-alkyl)silyl comprises silyl groups which have identical or two or three different alkyl groups.

5 The term di-(C₁₋₃-alkyl)amino comprises amino groups which have identical or two different C₁₋₃-alkyl groups.

The term aryl preferably denotes naphthyl or phenyl, more preferably phenyl.

10 The term heteroaryl denotes a 5- or 6-membered monocyclic aromatic ring possessing one to four identical or different heteroatoms selected from the group comprising N, O and S. Heteroaryl denotes preferably a pyrrolyl, furanyl, thienyl, pyridyl or tetrazolyl group, or


a pyrrolyl, furanyl, thienyl or pyridyl group wherein one or two methine groups are replaced in each case by a nitrogen atom.

15 The nomenclature in structural formulas used above and hereinafter, in which a bond of a substituent of a cyclic group, as e.g. a phenyl ring, is shown towards the centre of the cyclic group, denotes, unless otherwise stated, that this substituent may be bound to any free position of the cyclic group bearing an H atom.

20 The compounds according to the invention may be obtained using methods of synthesis known in principle. Preferably the compounds are obtained by the following methods according to the invention which are described in more detail hereinafter.

25 The glucose derivatives of formula II according to the invention may be synthesised from D-gluconolactone or a derivative thereof by adding the desired benzylbenzene compound in the form of an organometallic compound (Scheme 1).

Scheme 1: Addition of an Organometal Compound to a Gluconolactone



5 The reaction according to Scheme 1 is preferably carried out starting from a halogenated benzylbenzene compound of general formula IV, wherein Hal denotes chlorine, bromine, or iodine. The Grignard or lithium reagent of benzylbenzene (V) may be prepared from the corresponding chlorinated, brominated or iodinated benzylbenzene IV either via a so-called halogen-metal exchange reaction or by inserting the metal into the carbon-halogen bond.

10 The halogen-metal exchange to synthesize the corresponding lithium compound V may be carried out for example with an organolithium compound such as e.g. n-, sec- or tert-butyllithium. The analogous magnesium compound may also be generated by a halogen-metal exchange with a suitable Grignard reagent such as e.g. isopropyl- or sec-butylmagnesium bromide or chloride or diisopropyl- or di-sec-butylmagnesium without or in

15 the presence of an additional salt such as e.g. lithium chloride that may accelerate the metalation process; the specific transmetalating organomagnesium compound may also be generated in situ from suitable precursors (see e.g. *Angew. Chem.* **2004**, 116, 3396-3399 and *Angew. Chem.* **2006**, 118, 165-169 and references quoted therein). In addition, ate

20 complexes of organomagnesium compounds resulting from combining e.g. butylmagnesium chloride or bromide or isopropylmagnesium chloride or bromide and butyllithium, may be employed as well (see e.g. *Angew. Chem.* **2000**, 112, 2594-2596 and *Tetrahedron Lett.* **2001**, 42, 4841-4844 and references quoted therein). The halogen-metal exchange reactions are preferably carried out between 40 °C and -100 °C, particularly preferably between 10 °C and -80 °C, in an inert solvent or mixtures thereof, such as for example diethylether, dioxane, tetrahydrofuran, toluene, hexane, dimethylsulfoxide, dichloromethane or mixtures thereof.

25 The magnesium or lithium derivatized compounds thus obtained may optionally be transmetalated with metal salts such as e.g. cerium trichloride, zinc chloride or bromide,


indium chloride or bromide, to form alternative organometal compounds (V) suitable for addition. Alternatively, the organometal compound V may also be prepared by inserting a metal into the carbon-halogen bond of the haloaromatic compound IV. Lithium or magnesium are suitable elemental metals for this transformation. The insertion can be achieved in
5 solvents such as e.g. diethylether, dioxane, tetrahydrofuran, toluene, hexane, dimethylsulfoxide and mixtures thereof at temperatures ranging from -80 to 100 °C, preferably at -70 to 40 °C. In cases in which no spontaneous reaction takes place prior activation of the metal might be necessary such as e.g. treatment with 1,2-dibromoethane, iodine, trimethylsilylchloride, acetic acid, hydrochloric acid and/or sonication. The addition of
10 the organometal compound V to gluconolactone or derivatives thereof (VI) is preferably carried out at temperatures between 40 °C and -100 °C, particularly preferably at 0 to -80 °C, in an inert solvent or mixtures thereof, to obtain the compound of formula II. All foregoing reactions may be performed in air though execution under inert gas atmosphere such as argon and nitrogen is preferred. The metalation and/or coupling reaction may also be carried
15 out in microreactors and/or micromixers which enable high exchange rates; for example analogously to the processes described in WO 2004/076470. Suitable solvents for the addition of the metalated phenyl group V to the appropriately protected gluconolactone VI are e.g. diethylether, dimethoxyethane, benzene, toluene, methylene chloride, hexane, tetrahydrofuran, dioxane, *N*-methylpyrrolidone and mixtures thereof. The addition reactions
20 may be carried out without any further adjuvants or in the case of sluggishly reacting coupling partners in the presence of a promoter such as e.g. $\text{BF}_3^*\text{OEt}_2$ or Me_3SiCl (see M. Schlosser, *Organometallics in Synthesis*, John Wiley & Sons, Chichester/New York/Brisbane/Toronto/Singapore, 1994). Preferred definitions of the substituents R^8 in Scheme 1 are benzyl, substituted benzyl, allyl, trialkylsilyl, particularly preferably
25 trimethylsilyl, triisopropylsilyl, allyl, 4-methoxybenzyl and benzyl. If two adjacent substituents R^8 are linked together, these two substituents are preferably part of a benzylideneacetal, 4-methoxybenzylideneacetal, isopropylketal or constitute a dioxane with 2,3-dimethoxybutylene which is linked via the 2 and 3 positions of the butane with the adjacent oxygen atoms of the pyranose. The group R' preferably denotes hydrogen, C_{1-4} -alkyl, C_{1-4} -alkylcarbonyl or C_{1-4} -alkyloxycarbonyl, particularly preferably hydrogen, methyl or ethyl. The group R' is introduced after the addition of the organometallic compound V or a derivative thereof to the gluconolactone VI. If R' equals hydrogen or C_{1-4} -alkyl the reaction solution is treated with an alcohol such as e.g. methanol or ethanol or water in the presence of an acid such as e.g. acetic acid, methanesulfonic acid, toluenesulfonic acid, sulfuric acid,
30 trifluoroacetic acid, or hydrochloric acid. R' may also be attached after preparation of the hydrogen compound II by reacting the compound with an alcohol under acidic conditions. During installing R' the protective groups R^8 may be cleaved if labile under the reaction

conditions employed resulting in the corresponding protonated compound, i.e. compound II in which R⁸ equals H.

The synthesis of haloaromatic compound of formula IV may be carried out using standard 5 transformations in organic chemistry or at least methods known from the specialist literature in organic synthesis (see *inter alia* J. March, *Advanced Organic Reactions, Reactions, Mechanisms, and Structure*, 4th Edition, John Wiley & Sons, Chichester/New York/Brisbane/Toronto/Singapore, 1992 and literature cited therein). More specifically, the 10 use of transition metals and organo metal compounds for the synthesis of aromatic compounds has been detailed in different monographs (see e.g. L. Brandsma, S.F. Vasilevsky, H.D. Verkruisze, *Application of Transition Metal Catalysts in Organic Synthesis*, Springer-Verlag, Berlin/Heidelberg, 1998; M. Schlosser, *Organometallics in Synthesis*, John Wiley & Sons, Chichester/New York/Brisbane/Toronto/Singapore, 1994; P.J. Stang, F. Diederich, *Metal-Catalyzed Cross-Coupling Reactions*, Wiley-VCH, Weinheim, 1997 and 15 references quoted therein). The synthesis strategies described in the following provide a demonstration of this, by way of example. In addition, the aglycon part may also be assembled with the pyranose moiety already present using the same synthetic approaches.


Scheme 2: Synthesis of the Diarylketone Fragment

20



Scheme 2 shows the preparation of a precursor compound that may serve for the synthesis 25 of the haloaromatic compound of formula IV starting from a benzoylchloride and a second aromatic group applying Friedel-Crafts acylation conditions or variations thereof. This classic reaction has a wide substrate scope and is commonly carried out in the presence of a catalyst which is used in catalytic or stoichiometric amounts, such as e.g. AlCl₃, FeCl₃, iodine, iron, ZnCl₂, sulphuric acid, or trifluoromethanesulphonic acid. Instead of the benzoyl chloride 30 the corresponding carboxylic acid, anhydride, ester or benzonitrile may be used as well. The reactions are preferentially carried out in chlorinated hydrocarbons such as e.g. dichloromethane and 1,2-dichloroethane at temperatures from -30 to 120°C, preferably at 30 to 100°C. However, solvent-free reactions or reactions in a microwave oven are also possible.


Scheme 3: Reduction of Diarylketones and Diarylmethanols to Diarylmethanes



In Scheme 3 the substituent R denotes C₁₋₃-alkyl or aryl. Starting from the diarylketone or 5 diarylmethanol the diarylmethane is accessible in one or two reaction steps. The diarylketone may be reduced to the diarylmethane in two steps via the corresponding diphenylmethanol or in one step. In the two-step variant the ketone is reduced with a reducing agent such as for example a metal hydride such as e.g. NaBH₄, LiAlH₄ or iBu₂AlH to form the alcohol. The resulting alcohol can be converted in the presence of a Lewis acid such as for example 10 BF₃*OEt₂, InCl₃ or AlCl₃ or Brønsted acid such as for example hydrochloric acid, sulfuric acid, trifluoroacetic acid, or acetic acid with a reducing agent such as e.g. Et₃SiH, NaBH₄, or Ph₂SiClH to the desired diphenylmethane. The one-step process starting from the ketone to obtain the diphenylmethane may be carried out e.g. with a silane such as e.g. Et₃SiH, a 15 borohydride such as e.g. NaBH₄ or an aluminum hydride such as LiAlH₄ in the presence of a Lewis or Brønsted acid such as for example BF₃*OEt₂, tris(pentafluorophenyl)borane, trifluoroacetic acid, hydrochloric acid, aluminum chloride or InCl₃. The reactions are preferably carried out in solvents such as e.g. halogenated hydrocarbons such as dichloromethane, toluene, acetonitrile, or mixtures thereof at temperatures of -30 to 150°C, preferably at 20 to 100°C. Reductions with hydrogen in the presence of a transition metal 20 catalyst such as e.g. Pd on charcoal are another possible method of synthesis. Reductions

according to Wolff-Kishner or variants thereof are also possible. The ketone is firstly converted with hydrazine or a derivative thereof, such as e.g. 1,2-bis(tert-butyldimethylsilyl)hydrazine, into the hydrazone which breaks down under strongly basic reaction conditions and heating to form the diarylmethane and nitrogen. The reaction may 5 be carried out in one reaction step or after isolation of the hydrazone or a derivative thereof in two separate reaction steps. Suitable bases include e.g. KOH, NaOH or KOtBu in solvents such as e.g. ethyleneglycol, toluene, DMSO, 2-(2-butoxyethoxy)ethanol or tert-butanol; solvent-free reactions are also possible. The reactions may be carried out at temperatures between 20 to 250°C, preferably between 80 to 200°C. An alternative to the basic conditions 10 of the Wolff-Kishner reduction is the Clemmensen reduction which takes place under acidic conditions, which may also be used here. The alcohol function in diarylmethanol may also first be transformed into a better leaving group such as e.g. chloride, bromide, iodide, acetate, carbonate, phosphate, or sulfate; the subsequent reduction step to form the diarylmethane is widely described in the organic chemistry literature.

Scheme 4: Synthesis of Diarylmethane Unit and Possible Precursor Compounds thereof




In Scheme 4 the term "Alk" denotes C_{1-3} -alkyl and each substituent R is independently selected from each other from the group consisting of H, C_{1-3} -alkyl and C_{1-3} -alkoxy. Scheme 5 delineates the synthesis of diarylmethanes and possible precursor compounds thereof starting from a metalated phenyl group. Lithium or magnesium substituted aromatic

compounds may be synthesized from chlorinated, brominated, or iodinated aromatics by a halogen-metal exchange reaction with e.g. butyllithium, isopropylmagnesium halogenide, or diisopropylmagnesium or by insertion of the elemental metal into the halogen-carbon bond. The corresponding boron substituted compound such as e.g. boronic acid, boronic acid ester, or dialkylarylborane, is accessible from these metalated phenyl groups by reaction with a boron electrophile such as e.g. boronic acid ester or a derivative thereof. In addition, the borylated aromatic compound may also be prepared from the corresponding halogenated or pseudohalogenated precursor and a diboron or borane compound through a transition metal, e.g. palladium, catalyzed reaction (see e.g. *Tetrahedron Lett.* **2003**, p. 4895-4898 and references quoted therein). The lithium or magnesium substituted phenyl compounds add to benzaldehydes (step 3) and benzoic acids or derivatives thereof (step 4) such as benzoic acid esters, benzamides such as e.g. of the Weinreb type, benzonitriles, or benzoyl chlorides. These reactions may principally be conducted without an additional transition metal catalyst or transmetalation to another metal such as e.g. cerium, indium or zinc; sometimes the use of one of the latter alternatives is advantageous. Aryl boronic acids can be added to benzaldehydes by means of a rhodium catalyst furnishing the respective diarylmethanol (see e.g. *Adv. Synth. Catal.* **2001**, p. 343-350 and references quoted therein). Moreover, arylboronic acids, esters thereof, dialkylarylboranes, or aryltrifluoroborates may be coupled with benzoyl chlorides mediated by a transition metal such as e.g. palladium, a complex or a salt thereof delivering diarylketones. Metalated phenyl groups can be reacted with benzyl electrophiles such as benzyl chlorides, bromides, or iodides affording diarylmethanes. Lithium or magnesium derivatized phenyl compounds are reacted favorably but not always necessarily in the presence of a transition metal such as e.g. copper, iron, or palladium (see e.g. *Org. Lett.* **2001**, 3, 2871-2874 and references quoted therein).

Transmetalation from lithium or magnesium to e.g. boron, tin, silicon, or zinc furnishes e.g. the corresponding aromatic boronic acids, stannanes, silanes or zinc compounds, respectively, that may undergo coupling with benzyl electrophiles, e.g. benzyl halogenides, carbonates, phosphates, sulfonates, or carboxylic esters. The reaction is conducted in the presence of a transition metal, e.g. palladium, nickel, rhodium, copper, or iron (see e.g. *Tetrahedron Lett.* **2004**, p. 8225-8228 and *Org. Lett.* **2005**, p. 4875-4878 and references cited therein).

In order to prepare compounds of general formula I, in process a) according to the invention, a compound of general formula II



II


wherein R', R¹ to R⁵ are as hereinbefore defined and

5 R^{8a}, R^{8b}, R^{8c}, R^{8d} are as hereinbefore defined and independently of one another represent for example acetyl, pivaloyl, benzoyl, tert-butoxycarbonyl, benzyloxycarbonyl, trialkylsilyl, allyl, benzyl or substituted benzyl or in each case two adjacent groups R^{8a}, R^{8b}, R^{8c}, R^{8d} are combined a benzylideneacetal, diisopropylsilylideneeketal or isopropylideneeketal or a 2,3-dimethoxy-butylene group which is linked via position 2 and 3 of the butylene group to the 10 oxygen atoms of the pyranose ring and forms with them a substituted dioxane,

which may be obtained as hereinbefore described, is reacted with a reducing agent in the presence of a Lewis or Brønsted acid.

15 Suitable reducing agents for the reaction include for example silanes, such as triethyl-, tripropyl-, triisopropyl- or diphenylsilane, sodium borohydride, sodium cyanoborohydride, zinc borohydride, boranes, lithium aluminium hydride, diisobutylaluminium hydride or samarium iodide. The reductions are carried out without or in the presence of a suitable Brønsted acid, such as e.g. hydrochloric acid, toluenesulphonic acid, trifluoroacetic acid or acetic acid, or 20 Lewis acid, such as e.g. boron trifluoride etherate, trimethylsilyltriflate, titanium tetrachloride, tin tetrachloride, scandium triflate or zinc iodide. Depending on the reducing agent and the acid used the reaction may be carried out in a solvent, such as for example methylene chloride, chloroform, acetonitrile, toluene, hexane, diethyl ether, tetrahydrofuran, dioxane, ethanol, water or mixtures thereof at temperatures between -60°C and 120°C. One 25 particularly suitable combination of reagents consists for example of triethylsilane and boron trifluoride etherate, which is conveniently used in acetonitrile or dichloromethane at temperatures of -60°C and 60°C. Moreover, hydrogen may be used in the presence of a transition metal catalyst, such as e.g. palladium on charcoal or Raney nickel, in solvents such as tetrahydrofuran, ethyl acetate, methanol, ethanol, water or acetic acid, for the 30 transformation described.

Alternatively, in order to prepare compounds of general formula I according to process b) according to the invention, in a compound of general formula III



5

wherein R¹ to R⁵ are as hereinbefore defined and

R^{8a} to R^{8d} denote one of the protective groups defined hereinbefore, such as e.g. an acyl, allyl, arylmethyl, acetal, ketal or silyl group, and which may be obtained for example by reduction from the compound of formula II as hereinbefore described, the protective groups 10 are cleaved.

Any acyl protecting group used is cleaved for example hydrolytically in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulfuric acid or in the 15 presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or potassium hydroxide or aprotically, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 120°C, preferably at temperatures between 10 and 100°C. A trifluoroacetyl group is preferably cleaved by treating with an acid such as hydrochloric acid, optionally in the presence of a solvent such as acetic acid at temperatures between 50 and 120°C or by 20 treating with sodium hydroxide solution optionally in the presence of a solvent such as tetrahydrofuran or methanol at temperatures between 0 and 50°C.

Any acetal or ketal protecting group used is cleaved for example hydrolytically in an aqueous solvent, e.g. in water, isopropanol/water, acetic acid/water, tetrahydrofuran/water or 25 dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulfuric acid or aprotically, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 120°C, preferably at temperatures between 10 and 100°C.

A trimethylsilyl group is cleaved for example in water, an aqueous solvent mixture or a lower 30 alcohol such as methanol or ethanol in the presence of a base such as lithium hydroxide, sodium hydroxide, potassium carbonate or sodium methoxide.

In aqueous or alcoholic solvents, acids such as e.g. hydrochloric acid, trifluoroacetic acid or acetic acid are also suitable. For cleaving in organic solvents, such as for example diethyl ether, tetrahydrofuran or dichloromethane, it is also suitable to use fluoride reagents, such as e.g. tetrabutylammonium fluoride.

5

A benzyl, methoxybenzyl or benzyloxycarbonyl group is advantageously cleaved hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a suitable solvent such as methanol, ethanol, ethyl acetate or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures 10 between 0 and 100°C, but preferably at ambient temperatures between 20 and 60°C, and at a hydrogen pressure of 1 to 7 bar, but preferably 3 to 5 bar. A 2,4-dimethoxybenzyl group, however, is preferably cleaved in trifluoroacetic acid in the presence of anisole.

A tert.butyl or tert.butyloxycarbonyl group is preferably cleaved by treating with an acid such 15 as trifluoroacetic acid or hydrochloric acid or by treating with iodotrimethylsilane optionally using a solvent such as methylene chloride, dioxane, methanol or diethylether.

In the reactions described hereinbefore, any reactive groups present such as ethynyl, hydroxy, amino, alkylamino or imino groups may be protected during the reaction by 20 conventional protecting groups which are cleaved again after the reaction.

For example, a protecting group for an ethynyl group may be trialkylsilyl such as e.g. trimethylsilyl and triisopropylsilyl or dialkyl-hydroxymethyl such as e.g. 2-hydroxyisoprop-2-yl.

25 For example, a protecting group for a hydroxy group may be a trimethylsilyl, acetyl, trityl, benzyl or tetrahydropyranyl group.

Protecting groups for an amino, alkylamino or imino group may be, for example, a formyl, acetyl, trifluoroacetyl, ethoxycarbonyl, tert.butoxycarbonyl, benzyloxycarbonyl, benzyl, 30 methoxybenzyl or 2,4-dimethoxybenzyl group.

Moreover, the compounds of general formula I obtained may be resolved into their enantiomers and/or diastereomers, as mentioned hereinbefore. Thus, for example, *cis/trans* mixtures may be resolved into their *cis* and *trans* isomers, and compounds with at least one 35 optically active carbon atom may be separated into their enantiomers.

Thus, for example, the *cis/trans* mixtures may be resolved by chromatography into the *cis* and *trans* isomers thereof, the compounds of general formula I obtained which occur as racemates may be separated by methods known *per se* (cf. Allinger N. L. and Eliel E. L. in "Topics in Stereochemistry", Vol. 6, Wiley Interscience, 1971) into their optical antipodes and 5 compounds of general formula I with at least 2 asymmetric carbon atoms may be resolved into their diastereomers on the basis of their physical-chemical differences using methods known *per se*, e.g. by chromatography and/or fractional crystallisation, and, if these compounds are obtained in racemic form, they may subsequently be resolved into the enantiomers as mentioned above.

10

The enantiomers are preferably separated by column separation on chiral phases or by recrystallisation from an optically active solvent or by reacting with an optically active substance which forms salts or derivatives such as e.g. esters or amides with the racemic compound, particularly acids and the activated derivatives or alcohols thereof, and 15 separating the diastereomeric mixture of salts or derivatives thus obtained, e.g. on the basis of their differences in solubility, whilst the free antipodes may be released from the pure diastereomeric salts or derivatives by the action of suitable agents. Optically active acids in common use are e.g. the D- and L-forms of tartaric acid or dibenzoyltartaric acid, di- 20 o-tolyltartaric acid, malic acid, mandelic acid, camphorsulphonic acid, glutamic acid, aspartic acid or quinic acid. An optically active alcohol may be for example (+) or (-)-menthol and an optically active acyl group in amides, for example, may be a (+)-or (-)-mentyloxycarbonyl.

Furthermore, the compounds of formula I may be converted into the salts thereof, particularly for pharmaceutical use into the physiologically acceptable salts with inorganic or organic 25 acids. Acids which may be used for this purpose include for example hydrochloric acid, hydrobromic acid, sulfuric acid, methanesulfonic acid, phosphoric acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid or maleic acid.

Moreover, the compounds obtained may be converted into mixtures, for example 1:1 or 1:2 30 mixtures with amino acids, particularly with alpha-amino acids such as proline or phenylalanine, which may have particularly favourable properties such as a high crystallinity.

The compounds according to the invention are advantageously also obtainable using the methods described in the examples that follow, which may also be combined for this purpose 35 with methods known to the skilled man from the literature, for example the methods described in WO 98/31697, WO 01/27128, WO 02/083066, WO 03/099836, WO 2004/063209, WO 2004/080990, WO 2004/013118, WO 2004/052902, WO 2004/052903,

WO 2005/092877, WO 06/010557, WO 06/018150, WO 06/037537, WO 06/089872, WO 2006/064033, WO 06/108842, WO 2006/120208 and US application US 2003/0114390.

As already mentioned, the compounds of general formula I according to the invention and
5 the physiologically acceptable salts thereof have valuable pharmacological properties,
particularly an inhibitory effect on the sodium-dependent glucose cotransporter SGLT,
preferably SGLT2.

The biological properties of the new compounds may be investigated as follows:

10

The ability of the substances to inhibit the SGLT-2 activity may be demonstrated in a test set-up in which a CHO-K1 cell line (ATCC No. CCL 61) or alternatively an HEK293 cell line (ATCC No. CRL-1573), which is stably transfected with an expression vector pZeoSV (Invitrogen, EMBL accession number L36849) , which contains the cDNA for the coding 15 sequence of the human sodium glucose cotransporter 2 (Genbank Acc. No.NM_003041) (CHO-hSGLT2 or HEK-hSGLT2). These cell lines transport ^{14}C -labelled alpha-methyl-glucopyranoside (^{14}C -AMG, Amersham) into the interior of the cell in sodium-dependent manner.

20 The SGLT-2 assay is carried out as follows:

CHO-hSGLT2 cells are cultivated in Ham's F12 Medium (BioWhittaker) with 10% foetal calf serum and 250 $\mu\text{g}/\text{ml}$ zeocin (Invitrogen), and HEK293-hSGLT2 cells are cultivated in DMEM medium with 10% foetal calf serum and 250 $\mu\text{g}/\text{ml}$ zeocin (Invitrogen). The cells are detached from the culture flasks by washing twice with PBS and subsequently treating with 25 trypsin/EDTA. After the addition of cell culture medium the cells are centrifuged, resuspended in culture medium and counted in a Casy cell counter. Then 40,000 cells per well are seeded into a white, 96-well plate coated with poly-D-lysine and incubated overnight at 37°C, 5% CO₂. The cells are washed twice with 250 μl of assay buffer (Hanks Balanced Salt Solution, 137 mM NaCl, 5.4 mM KCl, 2.8 mM CaCl₂, 1.2 mM MgSO₄ and 10 mM HEPES 30 (pH7.4), 50 $\mu\text{g}/\text{ml}$ of gentamycin). 250 μl of assay buffer and 5 μl of test compound are then added to each well and the plate is incubated for a further 15 minutes in the incubator. 5 μl of 10% DMSO are used as the negative control. The reaction is started by adding 5 μl of ^{14}C -AMG (0.05 μCi) to each well. After 2 hours' incubation at 37°C, 5% CO₂, the cells are washed again with 250 μl of PBS (20°C) and then lysed by the addition of 25 μl of 0.1 N 35 NaOH (5 min. at 37°C). 200 μl of MicroScint20 (Packard) are added to each well and incubation is continued for a further 20 min at 37°C. After this incubation the radioactivity of

the ^{14}C -AMG absorbed is measured in a Topcount (Packard) using a ^{14}C scintillation program.

To determine the selectivity with respect to human SGLT1 an analogous test is set up in
5 which the cDNA for hSGLT1 (Genbank Acc. No. NM000343) instead of hSGLT2 cDNA is expressed in CHO-K1 or HEK293 cells.

The compounds of general formula I according to the invention may for example have EC50 values below 1000 nM, particularly below 200 nM, most preferably below 50 nM.

10

In view of their ability to inhibit the SGLT activity, the compounds according to the invention and the corresponding pharmaceutically acceptable salts thereof are suitable for the treatment and/or preventative treatment of all those conditions or diseases which may be affected by the inhibition of the SGLT activity, particularly the SGLT-2 activity. Therefore, 15 compounds according to the invention are particularly suitable for the prevention or treatment of diseases, particularly metabolic disorders, or conditions such as type 1 and type 2 diabetes mellitus, complications of diabetes (such as e.g. retinopathy, nephropathy or neuropathies, diabetic foot, ulcers, macroangiopathies), metabolic acidosis or ketosis, reactive hypoglycaemia, hyperinsulinaemia, glucose metabolic disorder, insulin resistance, 20 metabolic syndrome, dyslipidaemias of different origins, atherosclerosis and related diseases, obesity, high blood pressure, chronic heart failure, edema and hyperuricaemia. These substances are also suitable for preventing beta-cell degeneration such as e.g. apoptosis or necrosis of pancreatic beta cells. The substances are also suitable for improving or restoring the functionality of pancreatic cells, and also of increasing the number and size 25 of pancreatic beta cells. The compounds according to the invention may also be used as diuretics or antihypertensives and are suitable for the prevention and treatment of acute renal failure.

By the administration of a compound according to the invention an abnormal accumulation of 30 fat in the liver may be reduced or inhibited. Therefore according to another aspect of the present invention there is provided a method for preventing, slowing, delaying or treating diseases or conditions attributed to an abnormal accumulation of liver fat in a patient in need thereof characterized in that a compound or a pharmaceutical composition according to the present invention is administered. Diseases or conditions which are attributed to an abnormal 35 accumulation of liver fat are particularly selected from the group consisting of general fatty liver, non-alcoholic fatty liver (NAFL), non-alcoholic steatohepatitis (NASH),

hyperalimentation-induced fatty liver, diabetic fatty liver, alcoholic-induced fatty liver or toxic fatty liver.

In particular, the compounds according to the invention, including the physiologically acceptable salts thereof, are suitable for the prevention or treatment of diabetes, particularly type 1 and type 2 diabetes mellitus, and/or diabetic complications.

In addition compounds according to the invention are particularly suitable for the prevention or treatment of overweight, obesity (including class I, class II and/or class III obesity), visceral obesity and/or abdominal obesity.

The dosage required to achieve the corresponding activity for treatment or prevention usually depends on the compound which is to be administered, the patient, the nature and gravity of the illness or condition and the method and frequency of administration and is for the patient's doctor to decide. Expediently, the dosage may be from 1 to 100 mg, preferably 1 to 30 mg, by intravenous route, and 1 to 1000 mg, preferably 1 to 100 mg, by oral route, in each case administered 1 to 4 times a day. For this purpose, the compounds according to the invention may be formulated, optionally together with other active substances, together with one or more inert conventional carriers and/or diluents, e.g. with corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, cetylstearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof, to produce conventional galenic preparations such as plain or coated tablets, capsules, powders, suspensions or suppositories.

25

The compounds according to the invention may also be used in conjunction with other active substances, particularly for the treatment and/or prevention of the diseases and conditions mentioned above. Other active substances which are suitable for such combinations include for example those which potentiate the therapeutic effect of an SGLT antagonist according to the invention with respect to one of the indications mentioned and/or which allow the dosage of an SGLT antagonist according to the invention to be reduced. Therapeutic agents which are suitable for such a combination include, for example, antidiabetic agents such as metformin, sulphonylureas (e.g. glibenclamide, tolbutamide, glimepiride), nateglinide, repaglinide, thiazolidinediones (e.g. rosiglitazone, pioglitazone), PPAR-gamma-agonists (e.g. GI 262570) and antagonists, PPAR-gamma/alpha modulators (e.g. KRP 297), alpha-glucosidase inhibitors (e.g. acarbose, voglibose), DPPIV inhibitors (e.g. LAF237, MK-431), alpha2-antagonists, insulin and insulin analogues, GLP-1 and GLP-1 analogues (e.g.

exendin-4) or amylin. The list also includes inhibitors of protein tyrosinephosphatase 1, substances that affect deregulated glucose production in the liver, such as e.g. inhibitors of glucose-6-phosphatase, or fructose-1,6-bisphosphatase, glycogen phosphorylase, glucagon receptor antagonists and inhibitors of phosphoenol pyruvate carboxykinase, glycogen

5 synthase kinase or pyruvate dehydrokinase, lipid lowering agents such as for example HMG-CoA-reductase inhibitors (e.g. simvastatin, atorvastatin), fibrates (e.g. bezafibrate, fenofibrate), nicotinic acid and the derivatives thereof, PPAR-alpha agonists, PPAR-delta agonists, ACAT inhibitors (e.g. avasimibe) or cholesterol absorption inhibitors such as, for example, ezetimibe, bile acid-binding substances such as, for example, cholestyramine, 10 inhibitors of ileac bile acid transport, HDL-raising compounds such as CETP inhibitors or ABC1 regulators or active substances for treating obesity, such as sibutramine or tetrahydrolipostatin, dextroamphetamine, axokine, antagonists of the cannabinoid1 receptor, MCH-1 receptor antagonists, MC4 receptor agonists, NPY5 or NPY2 antagonists or β 3-agonists such as SB-418790 or AD-9677 and agonists of the 5HT2c receptor.

15

Moreover, combinations with drugs for influencing high blood pressure, chronic heart failure or atherosclerosis such as e.g. A-II antagonists or ACE inhibitors, ECE inhibitors, diuretics, β -blockers, Ca-antagonists, centrally acting antihypertensives, antagonists of the alpha-2-adrenergic receptor, inhibitors of neutral endopeptidase, thrombocyte aggregation inhibitors 20 and others or combinations thereof are suitable. Examples of angiotensin II receptor antagonists are candesartan cilexetil, potassium losartan, eprosartan mesylate, valsartan, telmisartan, irbesartan, EXP-3174, L-158809, EXP-3312, olmesartan, medoxomil, tasosartan, KT-3-671, GA-0113, RU-64276, EMD-90423, BR-9701, etc. Angiotensin II receptor antagonists are preferably used for the treatment or prevention of high blood 25 pressure and complications of diabetes, often combined with a diuretic such as hydrochlorothiazide.

A combination with uric acid synthesis inhibitors or uricosurics is suitable for the treatment or prevention of gout.

30

A combination with GABA-receptor antagonists, Na-channel blockers, topiramat, protein-kinase C inhibitors, advanced glycation end product inhibitors or aldose reductase inhibitors may be used for the treatment or prevention of complications of diabetes.

35

The dosage for the combination partners mentioned above is usefully 1/5 of the lowest dose normally recommended up to 1/1 of the normally recommended dose.

Therefore, in another aspect, this invention relates to the use of a compound according to the invention or a physiologically acceptable salt of such a compound combined with at least one of the active substances described above as a combination partner, for preparing a pharmaceutical composition which is suitable for the treatment or prevention of diseases or 5 conditions which can be affected by inhibiting the sodium-dependent glucose cotransporter SGLT. These are preferably metabolic diseases, particularly one of the diseases or conditions listed above, most particularly diabetes or diabetic complications.

The use of the compound according to the invention, or a physiologically acceptable salt 10 thereof, in combination with another active substance may take place simultaneously or at staggered times, but particularly within a short space of time. If they are administered simultaneously, the two active substances are given to the patient together; while if they are used at staggered times the two active substances are given to the patient within a period of less than or equal to 12 hours, but particularly less than or equal to 6 hours.


15 Consequently, in another aspect, this invention relates to a pharmaceutical composition which comprises a compound according to the invention or a physiologically acceptable salt of such a compound and at least one of the active substances described above as combination partners, optionally together with one or more inert carriers and/or diluents.

20 Thus, for example, a pharmaceutical composition according to the invention comprises a combination of a compound of formula I according to the invention or a physiologically acceptable salt of such a compound and at least one angiotensin II receptor antagonist optionally together with one or more inert carriers and/or diluents.

25 The compound according to the invention, or a physiologically acceptable salt thereof, and the additional active substance to be combined therewith may both be present together in one formulation, for example a tablet or capsule, or separately in two identical or different formulations, for example as a so-called kit-of-parts.

30 In the foregoing and following text, H atoms of hydroxyl groups are not explicitly shown in every case in structural formulae. The term "ambient temperature" means temperatures in the range from 20 to 25°C. The Examples that follow are intended to illustrate the present invention without restricting it:

35

Preparation of the starting compounds:Example I5 3-Bromo-5-iodo-2-methyl-benzoic acid


N-Iodosuccinimide (5.8 g) is added in portions to an ice-cold solution of sulphuric acid (20 mL). The resulting mixture is stirred for 40 min before 3-bromo-2-methyl-benzoic acid (5.0 g) dissolved in sulphuric acid (20 mL) is added at such a rate that the solution temperature maintained below 5 °C. The mixture is stirred at 5-10 °C for another 3 h before warming to 10 room temperature over night. Then, the mixture is poured on crushed ice (300 g) and the resultant solution is extracted with ethyl acetate. The combined extracts are washed in succession with aqueous 10% $\text{Na}_2\text{S}_2\text{O}_3$ solution (2x), water (3x), and brine (1x). After drying (MgSO_4) the organic layer, the solvent is evaporated under reduced pressure to give a solid. The solid is triturated with 70 °C-warm water, separated from the water by filtration and dried. 15 After the solid is triturated with little ether, filtered and dried, the product is obtained as a white solid.

Yield: 4.9 g (62% of theory)


Mass spectrum (ESI⁻): m/z = 339/341 (Br) [M-H]⁻

20 The following compound may be obtained analogously to Example I:

(1) 4-Bromo-5-iodo-2-methyl-benzoic acid



Mass spectrum (ESI⁻): m/z = 339/341 (Br) [M-H]⁻

Example II(3-Bromo-5-iodo-2-methyl-phenyl)-(4-ethyl-phenyl)-methanone

3-Bromo-5-iodo-2-methyl-benzoic acid (7.3 g) and SOCl_2 (70 mL) are combined in a flask connected with a reflux condenser. A few drops of DMF are added and the mixture is heated at reflux for 1 h. Then, the reaction solution is concentrated under reduced pressure and the residue is taken up in dichloromethane (80 mL) and ethylbenzene (15 mL). The resulting solution is cooled in an ice-bath and aluminum trichloride (7.2 g) is added in portions. Then, the cooling bath is removed and the reaction mixture is stirred at room temperature for 2 h. The reaction mixture is poured onto crushed ice and the organic phase is separated off. The aqueous phase is extracted with dichloromethane and the combined organic phases are washed in succession with 1 M hydrochloric acid, aqueous NaHCO_3 solution and brine. The organic phase is dried (sodium sulphate), the solvent is removed and the residue is chromatographed on silica gel (cyclohexane/ethyl acetate 20:1->9:1). The purified product is recrystallized from disopropylether.


Yield: 2.6 g (28% of theory)

Mass spectrum (ESI^+): $m/z = 429/431$ (Br) $[\text{M}+\text{H}]^+$


The following compound may be obtained analogously to Example II:

20

(1) (4-Bromo-5-iodo-2-methyl-phenyl)-(4-ethyl-phenyl)-methanone



Mass spectrum (ESI^+): $m/z = 429/431$ (Br) $[\text{M}+\text{H}]^+$


Example III3-Bromo-5-(4-ethyl-benzyl)-1-iodo-4-methyl-benzene

A solution of (3-bromo-5-iodo-2-methyl-phenyl)-(4-ethyl-phenyl)-methanone (4.2 g) and 5 triethylsilane (4.7 mL) in dichloromethane (10 mL) and acetonitrile (28 mL) is cooled in an ice-bath. Then, boron trifluoride diethyletherate (1.4 mL) is added dropwise over 3 min. The solution is stirred for 14 h at ambient temperature, before an aqueous 25% solution of KOH and diisopropylether are added. The organic phase is separated and the aqueous phase is extracted three times with diisopropylether. The combined organic phases are washed with 2 10 M potassium hydroxide solution and brine and then dried (sodium sulphate). After the solvent is evaporated, the residue is chromatographed on silica gel (cyclohexane/ethyl acetate 1:0->1:1).


Yield: 3.5 g (86% of theory)

15 The following compound may be obtained analogously to Example III:

(1) 2-Bromo-5-(4-ethyl-benzyl)-1-iodo-4-methyl-benzene



20

Example IV1-Bromo-3-(4-ethylbenzyl)-5-(1-methoxy-D-glucopyranos-1-yl)-2-methyl-benzene

A solution of iPrMgCl*LiCl in THF (1 mol/L, 10 mL) was added dropwise to a cold (-60 °C) solution of 3-bromo-5-(4-ethyl-benzyl)-1-iodo-4-methyl-benzene (3.5 g) in THF (20 mL). The solution is warmed to -20 °C over a period of 1 h and then a solution of 2,3,4,6-tetrakis-O-(trimethylsilyl)-D-glucopyranone (4.8 g) in tetrahydrofuran (3 mL) is added to the solution.


5 The resulting solution is slowly warmed to -5 °C and stirred for 6 h. The reaction is quenched with aqueous NH₄Cl solution and the resulting mixture is extracted with ethyl acetate. The combined organic extracts are washed with brine and dried (sodium sulphate). After the removal of the solvent, the residue is dissolved in methanol (30 mL) and treated with 10 methanesulfonic acid (0.3 mL). The solution is stirred at 40 °C for 6 h and then neutralized by the addition of solid NaHCO₃. The solvent is removed under reduced pressure and the residue is taken up in ethyl acetate. The organic solution is washed with water and brine and dried (sodium sulphate). After the removal of the solvent, the crude product is submitted to reduction without further purification.

Yield: 4.2 g (crude product)

15


The following compound may be obtained analogously to Example IV:

(1) 1-Bromo-4-(4-ethylbenzyl)-2-(1-methoxy-D-glucopyranos-1-yl)-5-methyl-benzene



20

Example V



1-Bromo-3-(4-ethylbenzyl)-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

25 A solution of 1-bromo-3-(4-ethylbenzyl)-5-(1-methoxy-D-glucopyranos-1-yl)-2-methylbenzene (4.2 g) and triethylsilane (3.5 mL) in dichloromethane (30 mL) and acetonitrile (90


mL) is cooled to -15 °C. Then boron trifluoride diethyletherate (2.1 mL) is added dropwise at such a rate that the solution temperature maintained below -5 °C. The resultant solution is stirred in an ice-bath for another 0.5 h and then the reaction is quenched by the addition of aqueous sodium hydrogen carbonate solution. The resulting mixture is stirred at room 5 temperature for 0.5 h and then the organic layer is separated and the aqueous layer is extracted with ethyl acetate. The combined organic layers are washed with brine and dried (sodium sulphate). The solvent is removed and the residue is taken up in dichloromethane (50 mL). The resultant solution is cooled in an ice-bath and pyridine (4.0 mL), acetic anhydride (4.3 mL) and 4-dimethylaminopyridine (0.1 g) are added in succession. The 10 solution is stirred at ambient temperature for 1 h and then diluted with dichloromethane (100 mL). The organic solution is washed twice with hydrochloric acid (1 mol/l in water) and dried (sodium sulphate). After evaporation of the solvent under reduced pressure, the residue is recrystallized from ethanol.

Yield: 2.7 g (53% of theory)


15 Mass spectrum (ESI⁺): m/z = 636/638 (Br) [M+NH₄]⁺

The following compound may be obtained analogously to Example V:

20 (1) 1-Bromo-4-(4-ethylbenzyl)-5-methyl-2-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene



Mass spectrum (ESI⁺): m/z = 636/638 (Br) [M+NH₄]⁺

Example VI


3-(4-Ethylbenzyl)-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

5 A flask kept under Ar atmosphere and charged with a stirring bar, 1-bromo-3-(4-ethylbenzyl)-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene (0.60 g), bis(pinacolato)diboron (0.37 g), 1,1'-bis(diphenylphosphino)ferrocene (43 mg), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) 1:1 complex with dichloromethane (65 mg), potassium acetate (0.29 g) and dioxane (3 mL) is heated at 100 °C. After stirring for 14

10 h, another portion of the same amount of bis(pinacolato)diboron (0.37 g), 1,1'-bis(diphenylphosphino)ferrocene (43 mg) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) 1:1 complex with dichloromethane is added and the mixture is heated at 100 °C for another 18h. After cooling to room temperature, the mixture is diluted with water and ethyl acetate and treated with charcoal.

15 After filtration, the organic layer is separated, washed with brine and dried (magnesium sulphate). The solvent is removed under reduced pressure and the resulting crude product is progressed without further purification.

Yield: 0.65 g (100% of theory)

Example VII

3-(4-Ethylbenzyl)-1-hydroxy-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene


5 A solution of 3-(4-ethylbenzyl)-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene (0.64 g) in acetic acid (3 mL) is treated with hydrogen peroxide solution (0.2 mL, 35% in water) for 3 h. Then, ethyl acetate and water are added. The organic layer is separated and washed with aqueous NaHCO_3 solution, water and brine. After drying (magnesium sulphate) of the organic solution, the solvent is

10 removed under reduced pressure and the residue is purified by chromatography on silica gel (cyclohexane/ethyl acetate 2:1->1:2).

Yield: 0.35 g (65% of theory)

Mass spectrum (ESI⁺): $m/z = 574$ [M+NH₄]⁺

15

Example VIII

1-Cyclobutoxy-3-(4-ethylbenzyl)-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

20

Diethyl azodicarboxylate (0.38 g) dissolved in THF (0.5 mL) is added to 3-(4-ethylbenzyl)-1-hydroxy-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene (0.27 g),

cyclobutanol (0.1 mL) and triphenylphosphine (0.56 g) dissolved in THF (2 mL). The mixture is heated with stirring at 70 °C overnight. After cooling to room temperature, the solvent is evaporated and the residue is purified by chromatography (cyclohexane/ethyl acetate 4:1->1:3) on silica gel.


5 Yield: 0.1 g (74% of theory)

Mass spectrum (ESI⁺): m/z = 628 [M+NH₄]⁺

The following compounds may be obtained analogously to Example VIII:

10 (1) 3-(4-Ethylbenzyl)-2-methyl-1-[(*R*)-tetrahydrofuran-3-yl]-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

(*S*)-tetrahydrofuran-3-ol is used as starting material under the conditions described above.




Mass spectrum (ESI⁺): m/z = 644 [M+NH₄]⁺


15

(2) 3-(4-Ethylbenzyl)-2-methyl-1-[(*S*)-tetrahydrofuran-3-yl]-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

(*R*)-tetrahydrofuran-3-ol is used as starting material under the conditions described above.



20 Mass spectrum (ESI⁺): m/z = 644 [M+NH₄]⁺


Example IX


3-Cyclopropyl-5-(4-ethylbenzyl)-4-methyl-1-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene

5 A flask charged with 1-bromo-3-(4-ethylbenzyl)-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene (0.30 g), potassium phosphate (0.38 g), cyclopropylboronic acid (50 mg), toluene (2 mL) and water (1 mL) is flushed with Ar for 5 min. Then, palladium acetate (11 mg) and tricyclohexylphosphonium tetrafluoroborate (36 mg) are added. The flask is tightly sealed and the mixture is stirred at 100 °C overnight. After cooling to room 10 temperature, the mixture is diluted with water and ethyl acetate and treated with charcoal. After filtration, the organic layer is separated, washed with brine and dried (magnesium sulphate). The solvent is removed under reduced pressure and the resulting crude product is submitted to deacetylation without further purification.

15 The following compound may be obtained analogously to Example IX:

(1) 2-Cyclopropyl-5-(4-ethylbenzyl)-4-methyl-1-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene




Preparation of the end compounds:**Example 1**5 **1-Cyclobutoxy-3-(4-ethylbenzyl)-5-(D-glucopyranos-1-yl)-2-methyl-benzene**

Aqueous sodium hydroxide solution (0.15 mL, 4 mol/L) is added to 1-cyclobutoxy-3-(4-ethylbenzyl)-2-methyl-5-(2,3,4,6-tetra-O-acetyl-D-glucopyranos-1-yl)-benzene (80 mg) dissolved in methanol (1 mL) and THF (0.5 mL). The solution is stirred at room temperature for 1 h and then neutralized with hydrochloric acid (1 mol/L). After removal of the organic solvents, the residue is diluted with aqueous sodium bicarbonate solution and the resulting mixture is extracted with ethyl acetate. The combined organic extracts are dried (sodium sulphate) and the solvent is evaporated. The remainder is purified by chromatography on silica gel (dichloromethane/methanol 1:0->6:1).

Yield: 50 mg (86% of theory)

15 Mass spectrum (ESI⁺): m/z = 460 [M+NH₄]⁺


The following compounds may be obtained analogously to Example 1:

(2) 5-(D-glucopyranos-1-yl)-3-(4-ethylbenzyl)-2-methyl-1-[(*R*)-tetrahydrofuran-3-yl]-benzene

20


Mass spectrum (ESI⁺): m/z = 476 [M+NH₄]⁺

(3) 5-(D-glucopyranos-1-yl)-3-(4-ethylbenzyl)-2-methyl-1-[(*S*)-tetrahydrofuran-3-yl]-benzene



Mass spectrum (ESI⁺): m/z = 476 [M+NH₄]⁺


(4) 3-Cyclopropyl-1-(D-glucopyranos-1-yl)-5-(4-ethylbenzyl)-4-methyl-benzene



5

Mass spectrum (ESI⁺): m/z = 430 [M+NH₄]⁺


(5) 2-Cyclopropyl-1-(D-glucopyranos-1-yl)-5-(4-ethylbenzyl)-4-methyl-benzene




10 Mass spectrum (ESI⁺): m/z = 430 [M+NH₄]⁺

The following compounds are also prepared analogously to the above-mentioned Examples and other methods known from the literature:

Ex.	Structure	Ex.	Structure
6		7	
8		9	
10		11	
12		13	
14		15	





Some examples of formulations will now be described in which the term "active substance" denotes one or more compounds according to the invention, including the salts thereof. In the case of one of the combinations with one or additional active substances as described previously, the term "active substance" also includes the additional active substances.

Example ATablets containing 100 mg of active substanceComposition:

1 tablet contains:

5	active substance	100.0 mg
	lactose	80.0 mg
	corn starch	34.0 mg
	polyvinylpyrrolidone	4.0 mg
	magnesium stearate	2.0 mg
10		220.0 mg

Method of Preparation:

The active substance, lactose and starch are mixed together and uniformly moistened with an aqueous solution of the polyvinylpyrrolidone. After the moist composition has been screened (2.0 mm mesh size) and dried in a rack-type drier at 50°C it is screened again (1.5 mm mesh size) and the lubricant is added. The finished mixture is compressed to form tablets.

Weight of tablet: 220 mg

Diameter: 10 mm, biplanar, faceted on both sides and notched on one side.

20 Example BTablets containing 150 mg of active substanceComposition:

1 tablet contains:

	active substance	150.0 mg
25	powdered lactose	89.0 mg
	corn starch	40.0 mg
	colloidal silica	10.0 mg
	polyvinylpyrrolidone	10.0 mg
	magnesium stearate	1.0 mg
30		300.0 mg

Preparation:

The active substance mixed with lactose, corn starch and silica is moistened with a 20% aqueous polyvinylpyrrolidone solution and passed through a screen with a mesh size of 1.5 mm. The granules, dried at 45°C, are passed through the same screen again and mixed with the specified amount of magnesium stearate. Tablets are pressed from the mixture.

Weight of tablet: 300 mg

die: 10 mm, flat

Example C

Hard gelatine capsules containing 150 mg of active substance

5 Composition:

1 capsule contains:

active substance	150.0 mg
corn starch (dried)	approx. 180.0 mg
lactose (powdered)	approx. 87.0 mg
10 magnesium stearate	3.0 mg
	approx. 420.0 mg

Preparation:

The active substance is mixed with the excipients, passed through a screen with a mesh size of 0.75 mm and homogeneously mixed using a suitable apparatus. The finished mixture is packed into size 1 hard gelatine capsules.

Capsule filling: approx. 320 mg

Capsule shell: size 1 hard gelatine capsule.

20 Example D

Suppositories containing 150 mg of active substance

Composition:

1 suppository contains:

active substance	150.0 mg
25 polyethyleneglycol 1500	550.0 mg
polyethyleneglycol 6000	460.0 mg
polyoxyethylene sorbitan monostearate	840.0 mg
	2,000.0 mg

Preparation:

30 After the suppository mass has been melted the active substance is homogeneously distributed therein and the melt is poured into chilled moulds.

Example E

Ampoules containing 10 mg active substance

35 Composition:

active substance	10.0 mg
0.01 N hydrochloric acid q.s.	

double-distilled water ad 2.0 ml

Preparation:

5 The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 2 ml ampoules.

Example F

Ampoules containing 50 mg of active substance

Composition:

10 4 active substance 50.0 mg
0.01 N hydrochloric acid q.s.
double-distilled water ad 10.0 ml


Preparation:

15 The active substance is dissolved in the necessary amount of 0.01 N HCl, made isotonic with common salt, filtered sterile and transferred into 10 ml ampoules.

Claims

1. Tetrasubstituted glucopyranosylated benzene derivatives of general formula I

5



wherein

R<sup>1</sup> denotes chlorine or methyl,

10

R<sup>2</sup> denotes C<sub>3-7</sub>-cycloalkyl or C<sub>3-7</sub>-cycloalkyloxy,

15

while the above-mentioned cycloalkyl-rings may be mono- or polysubstituted by fluorine and/or mono- or disubstituted by identical or different substituents L₂, and/or in the above-mentioned cycloalkyl rings one methylene group may be replaced by O, and

20

R³ hydrogen, fluorine, chlorine, bromine, iodine, C₁₋₆-alkyl, C₂₋₆-alkynyl, C₂₋₆-alkenyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-C₁₋₃-alkyl, C₅₋₇-cycloalkenyl, C₅₋₇-cycloalkenyl-C₁₋₃-alkyl, aryl, heteroaryl, C₁₋₄-alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, aminocarbonyl, C₁₋₄-alkylaminocarbonyl, di-(C₁₋₃-alkyl)aminocarbonyl, pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl, morpholin-4-ylcarbonyl, piperazin-1-ylcarbonyl, 4-(C₁₋₄-alkyl)piperazin-1-ylcarbonyl, hydroxycarbonyl, C₁₋₄-alkoxycarbonyl, C₁₋₄-alkyl-amino, di-(C₁₋₃-alkyl)amino, pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, piperazin-1-yl, 4-(C₁₋₄-alkyl)piperazin-1-yl, C₁₋₄-alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, C₁₋₄-alkylsulfonylamino, arylsulfonylamino, C₁₋₆-alkoxy, C₃₋₇-cycloalkyloxy, C₅₋₇-cycloalkenyloxy, aryloxy, heteroaryloxy, C₁₋₄-alkylsulfanyl, C₁₋₄-alkylsulfinyl, C₁₋₄-alkylsulfonyl, C₃₋₇-cycloalkylsulfanyl, C₃₋₇-cycloalkylsulfinyl, C₃₋₇-cycloalkylsulfonyl, C₅₋₇-cycloalkenylsulfanyl, C₅₋₇-cycloalkenylsulfinyl, C₅₋₇-cycloalkenylsulfonyl, arylsulfanyl, arylsulfinyl, arylsulfonyl, heteroarylsulfanyl, heteroarylsulfinyl, heteroarylsulfonyl, amino, hydroxy, cyano and nitro,

while the above-mentioned alkyl-, alkenyl-, alkynyl-, cycloalkyl- und cycloalkenyl- residues may be mono- or polysubstituted by fluorine and/or mono- or disubstituted by identical or different substituents L2, and

5 while in the above-mentioned C₅₋₆-cycloalkyl and C₅₋₆-cycloalkenyl rings one or two methylene groups may be replaced independently of one another by O, S, CO, SO or SO₂, and

10 while in the above-mentioned N-heterocycloalkyl rings one methylene group may be replaced by CO or SO₂, and

15 R⁴, R⁵ independently of one another denote hydrogen, fluorine, chlorine, bromine, iodine, cyano, nitro, C₁₋₃-alkyl, C₁₋₃-alkoxy, or a methyl- or methoxy-group substituted by 1 to 3 fluorine atoms,

20 L1 independently of one another are selected from among fluorine, chlorine, bromine, iodine, hydroxy, cyano, C₁₋₃-alkyl, difluoromethyl, trifluoromethyl, C₁₋₃-alkoxy, difluoromethoxy, trifluoromethoxy, amino, C₁₋₃-alkyl-amino and di(C₁₋₃-alkyl)-amino; and

25 L2 independently of one another are selected from among fluorine, chlorine, hydroxy, hydroxyl-C₁₋₄-alkyl, C₁₋₄-alkoxy, trifluoromethoxy, C₁₋₄-alkoxy-C₁₋₄-alkyl, cyano, hydroxycarbonyl, (C₁₋₄-alkyl)oxycarbonyl, aminocarbonyl, C₁₋₄-alkyl, trifluoromethyl, amino, C₁₋₄-alkyl-carbonyl-amino, C₁₋₃-alkyl-amino and di(C₁₋₃-alkyl)-amino; and

30 R⁶, R^{7a}, R^{7b}, R^{7c} independently of one another have a meaning selected from among hydrogen, (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl and aryl-(C₁₋₃-alkyl)- carbonyl, while the aryl-groups may be mono- or disubstituted independently of one another by identical or different groups L1;

while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups which may be substituted as defined; and

35 while, unless otherwise stated, the above-mentioned alkyl groups may be straight-chain or branched,

including tautomers, stereoisomers thereof or mixtures thereof, and physiologically acceptable salts thereof.

5 2. Tetrasubstituted glucopyranosylated benzene derivatives according to claim 1 characterized in that R^2 denotes C_{3-7} -cycloalkyl or C_{3-7} -cycloalkyloxy, while in the cycloalkyl rings a methylene group may be replaced by O, and wherein the cycloalkyl rings may be mono- or poly-fluorinated and/or mono- or disubstituted with identical or different substituents L_2 , wherein L_2 is defined as in claim 1.

10

3. Tetrasubstituted glucopyranosylated benzene derivatives according to claim 1 or 2 characterized in that R^3 denotes chlorine, bromine, iodine, C_{1-4} -alkyl, C_{3-7} -cycloalkyl, hydroxyl, C_{1-4} -alkyloxy, C_{3-7} -cycloalkyloxy, C_{1-4} -alkylsulfanyl, C_{3-7} -cycloalkylsulfanyl, while in a C_{5-6} -cycloalkyl ring a methylene group may be replaced by O, and wherein any alkyl group and cycloalkyl ring may be mono- or polyfluorinated and/or mono- or disubstituted with identical or different substituents L_2 , wherein L_2 is defined as in claim 1.

15

20 4. Tetrasubstituted glucopyranosylated benzene derivatives according to one or more of the claims 1 to 3 characterized in that R^6 denotes hydrogen, $(C_{1-8}$ -alkyl)oxycarbonyl, C_{1-8} -alkylcarbonyl or benzoyl and R^{7a} , R^{7b} , R^{7c} represent independently of one another hydrogen, $(C_{1-8}$ -alkyl)oxycarbonyl, $(C_{1-8}$ -alkyl)carbonyl or benzoyl.

25

5. Tetrasubstituted glucopyranosylated benzene derivatives according to claim 4 characterized in that R^6 , R^{7a} , R^{7b} , R^{7c} represent hydrogen.

30 6. Physiologically acceptable salts of the compounds according to one or more of the claims 1 to 5 with inorganic or organic acids.

7. Pharmaceutical composition, comprising a compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6, optionally together with one or more inert carriers and/or diluents.


35

8. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a pharmaceutical

composition which is suitable for the treatment or prevention of diseases or conditions which can be influenced by inhibiting the sodium-dependent glucose cotransporter SGLT.

- 5 9. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a pharmaceutical composition which is suitable for the treatment or prevention of one or more metabolic disorders.
- 10 10. Use according to claim 9, characterised in that the metabolic disorder is selected from the group consisting of type 1 and type 2 diabetes mellitus, complications of diabetes, metabolic acidosis or ketosis, reactive hypoglycaemia, hyperinsulinaemia, glucose metabolic disorder, insulin resistance, metabolic syndrome, dyslipidaemias of different origins, atherosclerosis and related diseases, obesity, high blood pressure, chronic heart failure, oedema and hyperuricaemia.
- 15 11. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a pharmaceutical composition for inhibiting the sodium-dependent glucose cotransporter SGLT2.
- 20 12. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a pharmaceutical composition for preventing the degeneration of pancreatic beta cells and/or for improving and/or restoring the functionality of pancreatic beta cells.
- 25 13. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a pharmaceutical composition for preventing, slowing, delaying or treating diseases or conditions attributed to an abnormal accumulation of liver fat in a patient in need thereof.
- 30 14. Use of at least one compound according to one or more of the claims 1 to 5 or a physiologically acceptable salt according to claim 6 for preparing a diuretic and/or antihypertensive.
- 35 15. Process for preparing a compound according to one or more of the claims 1 to 5, characterised in that

a) a compound of general formula II



5 wherein

10 R' denotes H, C_{1-4} -alkyl, $(C_{1-18}$ -alkyl)carbonyl, $(C_{1-18}$ -alkyl)oxycarbonyl, arylcarbonyl and aryl- $(C_{1-3}$ -alkyl)-carbonyl, wherein the alkyl or aryl groups may be mono- or polysubstituted by halogen;

15 R^{8a} , R^{8b} ,

R^{8c} , R^{8d} independently of one another have one of the meanings given hereinbefore and hereinafter for the groups R^6 , R^{7a} , R^{7b} , R^{7c} , or denote a benzyl or allyl group or a $R^aR^bR^cSi$ group or a ketal or acetal group, particularly an alkylidene or arylalkylidene ketal or acetal group, while in each case two adjacent groups R^{8a} , R^{8b} , R^{8c} , R^{8d} may form a cyclic silyl ketal, ketal or acetal group or a 1,2-di(C_{1-3} -alkoxy)-1,2-di(C_{1-3} -alkyl)-ethylene bridge, while the above-mentioned ethylene bridge forms, together with two oxygen atoms and the two associated carbon atoms of the pyranose ring, a substituted dioxane ring, particularly a 2,3-dimethyl-2,3-di(C_{1-3} -alkoxy)-1,4-dioxane ring, and while alkyl, aryl and/or benzyl groups may be mono- or polysubstituted by halogen or C_{1-3} -alkoxy, and while benzyl groups may also be substituted by a di- $(C_{1-3}$ -alkyl)amino group; and


20 R^a , R^b , R^c independently of one another denote C_{1-4} -alkyl, aryl or aryl- C_{1-3} -alkyl, wherein the aryl or alkyl groups may be mono- or polysubstituted by halogen;

25 while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups, preferably phenyl groups;

30 and wherein the groups R^1 to R^6 , R^{7a} , R^{7b} , R^{7c} are defined as in claim 1;

is reacted with a reducing agent in the presence of a Lewis or Brønsted acid, while any protective groups present are cleaved simultaneously or subsequently; or

5 b) a compound of general formula III



wherein R^{8a}, R^{8b}, R^{8c}, R^{8d} and R¹ to R⁵ are defined as in claim 1, but at least one of the
 10 groups R^{8a}, R^{8b}, R^{8c}, R^{8d} does not denote hydrogen, is hydrolysed to yield a compound of the formula I defined as in claim 1 wherein R⁶, R^{7a}, R^{7b} and R^{7c} denote hydrogen, and

if desired a compound of formula I thus obtained wherein R⁶ denotes a hydrogen atom, is converted by acylation into a corresponding acyl compound of general formula I, and/or


15

if necessary any protective group used in the reactions described above is cleaved and/or

if desired a compound of formula I thus obtained is resolved into its stereoisomers and/or

20 if desired a compound of formula I thus obtained is converted into the salts thereof.

16. Process for preparing compounds of general formula II



25

wherein

R' denotes H, C₁₋₄-alkyl, (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl and aryl-(C₁₋₃-alkyl)-carbonyl, wherein the alkyl or aryl groups may be mono- or polysubstituted by halogen;

5 R^{8a}, R^{8b},
R^{8c}, R^{8d} independently of one another has one of the meanings given for the groups R⁶, R^{7a}, R^{7b}, R^{7c}, or denote a benzyl or allyl group or a R^aR^bR^cSi group or a ketal or acetal group, while in each case two adjacent groups R^{8a}, R^{8b}, R^{8c}, R^{8d} may form a cyclic silyl ketal, ketal or acetal group or may form, with two oxygen atoms of the pyranose ring, a substituted 2,3-oxydioxane ring, particularly a 2,3-dimethyl-2,3-di(C₁₋₃-alkoxy)-1,4-dioxane ring, and while alkyl, aryl and/or benzyl groups may be mono- or polysubstituted by halogen or C₁₋₃-alkoxy, and while benzyl groups may also be substituted by a di-(C₁₋₃-alkyl)amino group; and


10

15 R^a, R^b, R^c independently of one another denote C₁₋₄-alkyl, aryl or aryl-C₁₋₃-alkyl, while the alkyl or aryl groups may be mono- or polysubstituted by halogen;


20 while by the aryl groups mentioned in the definition of the above groups are meant phenyl or naphthyl groups, preferably phenyl groups;

25 and R¹ to R⁶, R^{7a}, R^{7b}, R^{7c} are defined as in claim 1,

wherein an organometallic compound (V) which may be obtained by halogen-metal exchange or by inserting a metal in the carbon-halogen bond of a halogen-benzylbenzene compound of general formula IV



30 wherein Hal denotes Cl, Br and I and R¹ to R⁵ are defined as in claim 1, and optionally subsequent transmetallation, is added to a gluconolactone of general formula VI




wherein R^{8a}, R^{8b}, R^{8c}, R^{8d} are defined as in claim 1,

5 and

then the resulting adduct is reacted with water or an alcohol R'-OH, while R' denotes optionally substituted C₁₋₄-alkyl, in the presence of an acid and optionally the product obtained in the reaction with water wherein R' denotes H is converted, in a subsequent 10 reaction, with an acylating agent into the product of formula II wherein R' denotes (C₁₋₁₈-alkyl)carbonyl, (C₁₋₁₈-alkyl)oxycarbonyl, arylcarbonyl or aryl-(C₁₋₃-alkyl)-carbonyl, which may be substituted as specified.


17. A compound of general formula IV



15

wherein Hal denotes chlorine, bromine or iodine and the groups R¹ to R⁵ are defined as in claim 1.

20 18. A compound of general formula II



wherein R', R^{8a}, R^{8b}, R^{8c}, R^{8d} and R¹ to R⁵ are defined as in claim 16.

