



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2678528 A1 2008/08/21

(21) **2 678 528**

(12) **DEMANDE DE BREVET CANADIEN**  
**CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2008/02/14  
(87) Date publication PCT/PCT Publication Date: 2008/08/21  
(85) Entrée phase nationale/National Entry: 2009/08/12  
(86) N° demande PCT/PCT Application No.: US 2008/053921  
(87) N° publication PCT/PCT Publication No.: 2008/101054  
(30) Priorité/Priority: 2007/02/16 (US11/707,498)

(51) Cl.Int./Int.Cl. *A01N 29/02* (2006.01),  
*A01N 25/00* (2006.01)

(71) **Demandeur/Applicant:**  
HONEYWELL INTERNATIONAL INC., US

(72) **Inventeurs/Inventors:**  
POSS, ANDREW J., US;  
SINGH, RAJIV R., US;  
PHAM, HANG T., US


(74) **Agent:** GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSITIONS DE FUMIGANT AZEOTROPE A L'IODURE DE METHYLE

(54) Title: AZEOTROPIC FUMIGANT COMPOSITIONS OF METHYL IODIDE

**(57) Abrégé/Abstract:**

Azeotropic and azeotrope-like compositions of methyl iodide and at least one fluorocarbon or hydrofluorocarbon such as 1,1,1,3,3-pentafluoropropane (HFC-245fa). The compositions are present as a gas, at temperatures of about 30°C or below. The inventive compositions serve as a non-ozone-depleting gaseous fumigant which is useful in a variety of applications, in place of methyl bromide. These compositions serve as a drop-in replacement for gaseous methyl bromide, providing the benefits of a methyl iodide fumigant while also utilizing existing methyl bromide equipment.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
21 August 2008 (21.08.2008)

PCT

(10) International Publication Number  
WO 2008/101054 A1(51) International Patent Classification:  
A01N 29/02 (2006.01) A01N 25/00 (2006.01)(74) Agent: BEATUS, Carrie; Honeywell International Inc.,  
Law Department AB/2B, 101 Columbia Road, Morristown,  
New Jersey 07962 (US).(21) International Application Number:  
PCT/US2008/053921(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,  
EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID,  
IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC,  
LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN,  
MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH,  
PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV,  
SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,  
ZA, ZM, ZW.(22) International Filing Date:  
14 February 2008 (14.02.2008)(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MT, NL,  
NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG,  
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).(25) Filing Language:  
English(26) Publication Language:  
English(30) Priority Data:  
11/707,498 16 February 2007 (16.02.2007) US

## Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(71) Applicant (for all designated States except US): HONEY-  
WELL INTERNATIONAL INC. [US/US]; Law Department AB/2B, 101 Columbia Road, Morristown, NJ 07962  
(US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): POSS, Andrew J.  
[US/US]; 62 Deerhurst Park Boulevard, Kenmore, New  
York 14217 (US). SINGH, Rajiv R. [US/US]; 18 Foxfire  
Drive, Getzville, New York 14068 (US). PHAM, Hang T.  
[US/US]; 136 Larkspur Lane, Amherst, New York 14228  
(US).

WO 2008/101054 A1

(54) Title: AZEOTROPIC FUMIGANT COMPOSITIONS OF METHYL IODIDE

(57) Abstract: Azeotropic and azeotrope-like compositions of methyl iodide and at least one fluorocarbon or hydrofluorocarbon such as 1,1,1,3,3-pentafluoropropane (HFC-245fa). The compositions are present as a gas, at temperatures of about 30°C or below. The inventive compositions serve as a non-ozone-depleting gaseous fumigant which is useful in a variety of applications, in place of methyl bromide. These compositions serve as a drop-in replacement for gaseous methyl bromide, providing the benefits of a methyl iodide fumigant while also utilizing existing methyl bromide equipment.

## AZEOTROPIC FUMIGANT COMPOSITIONS OF METHYL IODIDE

5

BACKGROUND OF THE INVENTIONField of the Invention

The present invention relates to fumigants, and particularly to fumigant compositions having similar properties to methyl bromide, without its ozone-depleting properties. Specifically, the invention relates to azeotropic and azeotrope-like compositions of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, such as 1,1,1,3,3-pentafluoropropane (HFC-245fa).

Description of the Related Art

15

Methyl Bromide is a gaseous fumigant that has been used commercially since the early 1900's. It is known for being extremely effective as a herbicide, nematocide, insecticide and fungicide. Methyl bromide has been widely used for soil fumigation, not only for controlling a variety of pests on numerous crops, but also as a commodity quarantine treatment for imports and exports, and as a structural fumigant applied to a building surface or the like. However, it is a Restricted Use Pesticide (RUP) because of its high acute toxicity to applicators. Methyl bromide has also been designated as an ozone-depleter, and thus its production and use have been severely restricted pursuant to the Montreal Protocol.

Efforts have been made to develop an alternative or replacement for methyl bromide as a fumigant. There currently exist only a few conventional methyl bromide alternatives, such as chloropicrin, 1,3-dichloropropene, metham sodium, and methyl iodide. Two or more of these materials are commonly

applied as a mixture, to produce a product similar to methyl bromide.

However, none of these potential alternatives are an adequate “drop-in replacement” for methyl bromide, based on their physical handling requirements, performance, or economics. The term “drop-in replacement” is

5 used when the methodology, equipment, production system, and the like, of an original material do not have to be changed significantly when using a replacement material, and that a comparable amount of the replacement material can be used for the same targets as the original material.

10 A great deal of research has been conducted in evaluating methyl iodide as a drop-in replacement for methyl bromide. Methyl iodide has been found to be equal to or better than methyl bromide in combating weeds, nematodes, and soil pathogens. Further, methyl iodide is not associated with ozone depletion, and does not result in plant toxicity when used in effective concentrations.

15 However, methyl iodide is a low boiling liquid with a boiling point of 42.5 °C (108 °F), while methyl bromide is a gas at ambient temperature and pressure. Methyl iodide has a lower vapor pressure and higher density than methyl bromide. Thus, the use of methyl iodide in existing methyl bromide equipment suffers several shortcomings such as clogged tubing, material

20 remnants in system pipes, and long line purging processes for cleaning. Furthermore, the use of methyl iodide results in problems such as missed bed applications, since the methyl bromide equipment is designed for gaseous fumigant applications. Such missed bed applications may lead to significant crop loss in soil fumigation. Thus, while methyl iodide may serve well as a 25 fumigant, it is not a suitable drop-in replacement for methyl bromide.

It would be desirable to provide a fumigant alternative to methyl bromide which can also serve as a drop-in replacement, thereby allowing for the use existing methyl bromide equipment and systems. The present invention 30 provides a solution to this problem.

It has now been unexpectedly found that by combining methyl iodide with at least one fluorocarbon or hydrofluorocarbon, the resulting azeotropic or azeotrope-like composition exists as a gas at temperatures below about 30°C. The resulting gaseous composition serves as a drop-in replacement for methyl bromide, providing benefits if a methyl iodide fumigant while also utilizing existing methyl bromide equipment.

#### SUMMARY OF THE INVENTION

10

The invention provides an azeotropic or azeotrope-like composition comprising a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, which composition is a gas at temperatures of about 30°C or below.

15

The invention further provides a fumigation process which comprises:

- a) providing a fumigant comprising an azeotropic or azeotrope-like composition which comprises a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, which composition is a gas at temperatures of about 30°C or below; and
- 20 b) applying the fumigant to a material to be fumigated.

25

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to azeotropic or azeotrope-like composition comprising a  
5 mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon,  
which composition is present in the form of a gas at temperatures of about  
30°C or below.

As used herein, the term "azeotrope-like" is intended in its broad sense to  
10 include both compositions that are strictly azeotropic and compositions that  
behave like azeotropic mixtures. From fundamental principles, the  
thermodynamic state of a fluid is defined by pressure, temperature, liquid  
composition, and vapor composition. An azeotropic mixture is a system of  
two or more components in which the liquid composition and vapor  
15 composition are equal at the stated pressure and temperature. In practice, this  
means that the components of an azeotropic mixture are constant boiling and  
cannot be separated during distillation.

Azeotrope-like compositions are constant boiling or essentially constant  
20 boiling. In other words, for azeotrope-like compositions, the composition of  
the vapor formed during boiling or evaporation (under substantially isobaric  
conditions) is identical, or substantially identical, to the original liquid  
composition. Thus, with boiling or evaporation, the liquid composition  
changes, if at all, only to a minimal or negligible extent. This is to be  
25 contrasted with non-azeotrope-like compositions in which, during boiling or  
evaporation, the liquid composition changes to a substantial degree. All  
azeotrope-like compositions of the invention within the indicated ranges as  
well as certain compositions outside these ranges are azeotrope-like.

30 It is well known that at differing pressures, the composition of a given

azeotrope will vary at least slightly, as does the boiling point of the composition. Thus, an azeotrope of A and B represents a unique type of relationship, but with a variable composition depending on temperature and/or pressure. It follows that, for azeotrope-like compositions, there is a range of 5 compositions containing these components in varying proportions that are azeotrope-like. All such compositions are intended to be covered by the term azeotrope-like as used herein.

As stated above, the present invention provides azeotropic and azeotrope-like 10 compositions comprising a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon. Methyl iodide, also known as iodomethane, which is commonly abbreviated as "MeI" and has the formula CH<sub>3</sub>I. Methyl iodide has a boiling point of about 42.5°C and a density of about 2.3 g/cc. Methyl iodide has been conventionally known as a useful 15 fumigant, and serves this purpose in the inventive composition. An added benefit of methyl iodide is that it is not associated with ozone depletion.

Fluorocarbons are defined herein as any carbon molecule having at least one attached fluorine group. Hydrofluorocarbons are particularly useful in the 20 present invention. The present claims require that the inventive compositions comprise at least one fluorocarbon or hydrofluorocarbon. The at least one fluorocarbon or hydrofluorocarbon increases the overall volume of the inventive compositions, facilitating application of the composition and increasing the time that a given volume of methyl iodide is exposed to a 25 material to be contacted. The at least one fluorocarbon or hydrofluorocarbon further enables a more uniform and easily controlled application of the inventive composition. In addition, the at least one fluorocarbon or hydrofluorocarbon serves as a non-toxic portion of the composition, reducing worker exposure to toxic materials.

Several different fluorocarbons and hydrofluorocarbons may be suitable for use in the compositions formed according to this invention. Examples of suitable fluorocarbons for use in the present invention nonexclusively include 1-chloro-3,3,3 trifluoropropene (HCFC-1233xd); 2,2-dichloro-1,1,1-5 trifluoroethane (HCFC-123); 1,1,2,2-tetrafluoroethyl methyl ether (HFE-245); and cis-1,3,3,3-tetrafluoropropene (HFC-1234ze). Examples of suitable hydrofluorocarbons for use in the present invention nonexclusively include 1,1,1,3,3-pentafluoropropane (HFC-245fa); 1,1,1,3,3-pentafluorobutane (HFC-365); 1,2-difluoroethane (HFC-152); 1,2,2,3,3-pentafluoropropane 10 (245ca); and 1,2,2-trifluoroethane (HFC-143). A preferred hydrocarbon is 1,1,1,3,3-pentafluoropropane (HFC-245fa), which is a nonflammable and non-toxic compound with an ozone depletion potential of zero. It has been unexpectedly found that methyl iodide and 1,1,1,3,3-pentafluoropropane (HFC-245fa) form an azeotropic mixture which closely resembles several 15 physical properties of methyl bromide, such as specific gravity and density. It has also been found that methyl iodide and 1,1,1,3,3-pentafluorobutane (HFC-365) form an azeotropic mixture which also resembles the properties of methyl bromide. It has further been found that methyl iodide and cis-1,3,3,3-tetrafluoropropene (HFC-1234ze) form an azeotropic mixture which also 20 resembles the properties of methyl bromide. In a preferred embodiment, the at least one fluorocarbon or hydrofluorocarbon has a boiling point of from about 0° C to about 50° C.

In preferred embodiments, the at least one fluorocarbon or hydrofluorocarbon 25 has an average Ozone Depletion Potential (ODP) of about 0.05 or less. The ozone depletion potential (ODP) of a chemical compound is the relative amount of degradation to the ozone layer it can cause, with trichlorofluoromethane (R-11) being fixed at an ODP of 1.0. Chlorodifluoromethane (R-22), for example, has an ODP of 0.05.

In further preferred embodiments, the at least one fluorocarbon or hydrofluorocarbon has a 100-year Global Warming Potential (GWP) of about 1,000 or less. Global warming potential (GWP) is a measure of how much a given mass of greenhouse gas is estimated to contribute to global warming. It

5 is a relative scale which compares the gas in question to that of the same mass of carbon dioxide, whose GWP is 1 by definition. A GWP is calculated over a specific time interval and the value of this must be stated whenever a GWP is quoted. The most common time interval used today is 100 years.

10 The compositions of the present invention comprise effective amounts of the methyl iodide and the at least one fluorocarbon or hydrofluorocarbon. The term "effective amounts" as used herein refers to the amount of each component which, upon combination with the other component or components, results in the formation of the presently claimed azeotropic or

15 azeotrope-like compositions. The methyl iodide is preferably present in the inventive composition an amount from about 5 to about 70 weight percent of the composition, more preferably from about 15 to about 60 weight percent, and most preferably from about 25 to about 50 weight percent. The at least one fluorocarbon or hydrofluorocarbon is preferably present in the

20 composition in an amount from about 30 to about 95 weight percent of the composition, more preferably from about 40 to about 85 weight percent of the composition, and most preferably from about 50 to about 75 weight percent of the composition.

25 The inventive compositions may be present in the form of binary azeotropes, which consist essentially of methyl iodide and at least one fluorocarbon or hydrofluorocarbon. The azeotrope-like compositions of the invention may optionally include additional components or additives. Suitable additives for the present compositions nonexclusively include chloropicrin, acrolein, 1,3-

30 dichloropropene, dimethyl disulfide, furfural, and propylene oxide. One

preferred additive includes chloropicrin. For example, in one preferred embodiment, an azeotropic or azeotrope-like composition of the invention comprises 33% methyl iodide, 33% HFC-245fa, and 33% chloropicrin.

- 5 The methyl iodide, the at least one fluorocarbon or hydrofluorocarbon, and any optional additives, may be combined using any conventional means which results in a substantially homogeneous mixture of all components. When combined, the inventive mixture forms an azeotropic or azeotrope-like composition which is present in the form of a gas at temperatures of about
- 10 30°C or below.

The inventive compositions are present in the form of a gas at temperatures of about 30°C or below, and in certain embodiments preferably at about 20°C or below, such that the compositions may be applied in a gaseous state at room

- 15 temperature, and utilize methyl bromide equipment systems which were designed for gaseous applications. In a preferred embodiment, the compositions of the present invention have a boiling point which ranges from about 8°C to about 14.5°C, more preferably from about 8°C to about 13.8°C, and most preferably from about 8°C to about 12.7°C.

20

The inventive compositions preferably exhibit a particular density which allows for the compositions to be used in methyl bromide equipment systems without causing blockages and the like. In a preferred embodiment, the compositions of the present invention have a density of from about 1.5 g/cc to

- 25 about 2.4 g/cc, more preferably from about 1.6g/cc to about 2.0 g/cc, and most preferably from about 1.6 g/cc to about 1.8 g/cc.

The inventive compositions may be used in a variety of applications. These compositions are particularly suited for use as a fumigant or as a component of

- 30 a fumigant. Examples of fumigation applications for the inventive

compositions nonexclusively include combating insects, termites, rodents, weeds, nematodes, and soil-borne diseases. The compositions may further be used to fumigate agricultural commodities, grain elevators, mills, ships, clothes, furniture, greenhouses, and the control of pests in buildings (structural fumigation), and the like.

The inventive compositions are preferably pumped through existing fumigation pipes and systems designed for methyl bromide use. The inventive gaseous compositions eliminate the problems of clogged tubing, pooled 10 deposits of chemicals that are typically associated with liquid methyl iodide, and can be easily purged from the system. In fact, methyl iodide's particular difficulty of missed bed application is no longer a problem because the inventive azeotropic or azeotrope-like compositions, like methyl bromide, are present as a gas at ambient temperature and move easily through application 15 tubing of an existing methyl bromide system. Furthermore, the inventive azeotropic or azeotrope-like compositions have a very similar environmental efficacy and spectrum of activity as methyl bromide, while also exhibiting a low potential for ozone degradation. Thus, the inventive compositions effectively serve as a drop-in replacement for methyl bromide.

20

The present invention further relates to a fumigation process. According to this process, a fumigant is provided which comprises an azeotropic or azeotrope-like composition which comprises a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, which composition is a gas at 25 temperatures of about 30°C or below. The fumigant preferably comprises or consists essentially of the azeotropic or azeotrope-like compositions discussed at length above. The fumigant is then applied to a material to be fumigated, and is preferably applied in the form of a gas. In a preferred embodiment, the application of the fumigant is conducted at an ambient temperature of from 30 about 0°C to about 50°C. The fumigant may be applied to a variety of

different materials to be fumigated, in a variety of fumigation applications as described above. In one preferred embodiment, the fumigant is applied to soil. In a further preferred embodiment the fumigant is applied to wood. In still another preferred embodiment the fumigant is applied to a building surface.

5

The following non-limiting examples serve to illustrate the invention. It will be appreciated that variations in proportions and alternatives in elements of the components of the invention will be apparent to those skilled in the art and are within the scope of the present invention.

10

#### EXAMPLE 1

An ebulliometer consisting of vacuum jacketed tube with a condenser on top which is further equipped with a Quartz Thermometer was used. About 22 g 15 HFC-245fa was charged to the ebulliometer and then methyl iodide is added in small, measured increments. Temperature depression is observed when methyl iodide was added to HFC-245fa, indicating a binary minimum boiling azeotrope is formed. The atmospheric pressure was measured to be 14.50 psia. From greater than about 0 to about 60 weight percent methyl iodide, the 20 boiling point of the composition changed by about 3°C or less. The binary mixtures shown below in Table 1 were studied and the boiling point of the compositions from 10 wt% CH<sub>3</sub>I to about 60 wt% CH<sub>3</sub>I changed by less than about 0.5 °C. The compositions exhibit azeotrope and/or azeotrope-like properties over this range.

25

Table 1

Wt.% CH <sub>3</sub> I	Wt.% HFC-245fa	Temp (°C)
---------------------------	-------------------	--------------

0.00	100.00	14.83
0.54	99.46	14.60
1.61	98.39	14.19
4.69	95.31	13.55
9.40	90.60	12.94
14.48	85.52	12.62
19.02	80.98	12.50
23.10	76.90	12.46
26.79	73.21	12.42
30.14	69.86	12.40
33.20	66.80	12.43
36.01	63.99	12.45
38.99	61.01	12.47
41.71	58.29	12.48
44.54	55.46	12.50
47.10	52.90	12.51
49.44	50.56	12.53
51.58	48.42	12.53
53.54	46.46	12.55
55.57	44.43	12.55
57.43	42.57	12.56

## EXAMPLE 2

5 An ebulliometer consisting of vacuum jacketed tube with a condenser on top similar to the one in Example 1 was used. About 10 g HFC-365 was charged to the ebulliometer and then methyl iodide is added in small, measured increments. Temperature depression is observed when methyl iodide was added to HFC-365, with a minimum boiling temperature below the boiling

10 point of pure methyl iodide and pure HFC-365, indicating a binary minimum boiling azeotrope is formed. The atmospheric pressure was measured to be 14.29 psia. From greater than about 80 to about 45 weight percent HFC-365, the boiling point of the composition changed by about 2°C or less. The binary mixtures shown in Table 2 were studied and the boiling point of the

15 compositions from 35 wt% CH<sub>3</sub>I to about 65 wt% CH<sub>3</sub>I changed by less than

about 0.5°C. The compositions exhibit azeotrope and/or azeotrope-like properties over this range.

Table 2

5

Wt.% CH <sub>3</sub> I	Wt.% HFC-365	Temp (° C)
0.0	100.0	39.2
0.5	99.5	39.2
1.1	98.9	39.2
1.6	98.4	39.1
2.1	97.9	39.0
2.6	97.4	38.4
3.2	96.8	37.8
3.7	96.3	37.5
4.2	95.8	36.8
4.7	95.3	36.2
5.1	94.9	35.9
5.6	94.4	35.6
6.1	93.9	35.5
6.6	93.4	35.4
7.1	92.9	35.0
7.5	92.5	35.1
8.0	92.0	35.4
8.4	91.6	35.1
8.9	91.1	34.8
9.3	90.7	34.5
9.8	90.2	34.2
10.2	89.8	34.1
10.7	89.3	32.8
11.1	88.9	32.7
11.5	88.5	32.6
11.9	88.1	32.5
12.4	87.6	32.4
12.8	87.2	32.4
13.6	86.4	32.3
14.4	85.6	32.2
15.2	84.8	32.0
16.0	84.0	32.0
16.7	83.3	31.8
17.5	82.5	31.6
18.9	81.1	31.4

20.3	79.7	31.0
21.7	78.3	30.9
23.0	77.0	30.6
24.3	75.7	30.5
25.5	74.5	30.3
26.7	73.3	30.3
27.8	72.2	30.3
28.9	71.1	30.2
30.0	70.0	30.1
31.1	68.9	29.9
32.1	67.9	29.9
33.1	66.9	29.8
34.0	66.0	29.7
35.0	65.0	29.6
35.9	64.1	29.5
36.7	63.3	29.5
37.6	62.4	29.5
38.4	61.6	29.5
39.2	60.8	29.5
40.0	60.0	29.5
40.8	59.2	29.5
41.6	58.4	29.5
42.3	57.7	29.5
43.0	57.0	29.4
43.7	56.3	29.4
44.4	55.6	29.4
45.0	55.0	29.4
45.7	54.3	29.4
46.3	53.7	29.3
46.9	53.1	29.3
47.5	52.5	29.2
48.1	51.9	29.2
48.7	51.3	29.2
49.3	50.7	29.2
49.8	50.2	29.2
50.4	49.6	29.2
50.9	49.1	29.2
51.4	48.6	29.2
51.9	48.1	29.2
52.4	47.6	29.2
52.9	47.1	29.2
53.4	46.6	29.1
53.9	46.1	29.1
54.3	45.7	29.1

54.8	45.2	29.1
55.2	44.8	29.1
55.6	44.4	29.1
56.1	43.9	29.1
56.5	43.5	29.0
56.9	43.1	29.0
57.3	42.7	29.0
57.7	42.3	29.0
58.1	41.9	29.0
58.4	41.6	29.0
58.8	41.2	29.0
59.2	40.8	29.0
59.5	40.5	29.0
59.9	40.1	29.0
60.2	39.8	29.0
60.6	39.4	29.0
60.9	39.1	29.0
61.2	38.8	29.0
61.6	38.4	29.0
61.9	38.1	29.0
62.2	37.8	29.0
62.5	37.5	29.0
62.8	37.2	29.0
63.1	36.9	29.0
63.4	36.6	29.0
63.7	36.3	29.0
64.0	36.0	29.0
64.2	35.8	29.0
64.5	35.5	29.0
64.8	35.2	29.0

### EXAMPLE 3

5 An ebulliometer consisting of vacuum jacketed tube with a condenser on top which is further equipped with a Quartz Thermometer was used. About 19.5 g cis isomer of 1,3,3,3-tetrafluoropropene (cis-HFC-1234ze) was charged to the ebulliometer. The cis isomer contained approximately 9 % HFC-245fa as an impurity. Then methyl iodide is added in small, measured increments.

Temperature depression is observed when methyl iodide was added to cis-HFC-1234ze, indicating that a binary minimum boiling azeotrope is formed. The atmospheric pressure was measured to be 14.42 psia. From greater than about 0 to about 55 weight percent methyl iodide, the boiling point of the 5 composition changed by about 1°C or less. The binary mixtures shown below in Table 3 were studied and the boiling point of the compositions from 5 wt% CH<sub>3</sub>I to about 55 wt% CH<sub>3</sub>I changed by less than about 0.5 °C. The compositions exhibit azeotrope and/or azeotrope-like properties over this range.

10

Table 3

Wt.% CH <sub>3</sub> I	Wt.% cis-1234ze	Temp (°C)
0.00	100.00	9.68
0.61	99.39	9.61
1.82	98.18	9.46
5.27	94.73	9.20
8.48	91.52	9.03
12.44	87.56	8.81
16.07	83.93	8.81
19.41	80.59	8.75
23.23	76.77	8.74
26.71	73.29	8.73
29.88	70.12	8.78
32.79	67.21	8.88
36.98	63.02	8.90
40.67	59.33	8.94
43.96	56.04	8.98
46.90	53.10	9.11
49.55	50.45	9.14
51.94	48.06	9.16
54.12	45.88	9.17

EXAMPLE 4

Methyl iodide and HFC-245fa are combined to form an azeotrope or azeotrope-like composition according to the present invention. The composition is applied as a fumigant to a field prior to planting the soil, by

5 injection through shank-mounted tubes that are pulled through the soil, followed by covering the soil with a plastic film. The fumigant is applied at a broadcast rate sufficient to rid the soil of any diseases and pests found therein. After the fumigant is applied, the soil is to remain undisturbed for a sufficient time to allow the fumigant to rid the soil of all deleterious organisms.

10 Applications to the soil include attention to soil conditions such as high moisture content, soil temperatures below 13°C/55°F at a 20cm/8-inch depth, and/or cloddy soils with high plant /weed trash content. The treated field may be planted after the elapsed planting interval time.

EXAMPLE 5

15 Methyl iodide, HFC-245fa, and chloropicrin are combined to form an azeotrope or azeotrope-like composition according to the present invention. The composition is applied as a fumigant to a field prior to planting the soil, by injection through shank-mounted tubes that are pulled through the soil, followed by covering the soil with a plastic film. The fumigant is applied at a broadcast rate sufficient to rid the soil of any diseases and pests found therein.

20 After the fumigant is applied, the soil is to remain undisturbed for a sufficient time to allow the fumigant to rid the soil of all deleterious organisms. Applications to the soil include attention to soil conditions such as high moisture content, soil temperatures below 13°C/55°F at a 20cm/8-inch depth,

25 and/or cloddy soils with high plant /weed trash content. The treated field may be planted after the elapsed planting interval time.

EXAMPLE 6

Methyl iodide and HFC-245fa are combined to form an azeotrope or azeotrope-like composition according to the present invention. The composition is applied as a fumigant to a field, prior to planting the soil, in

5 water through the drip irrigation system under a plastic tarp. The fumigant is applied at a broadcast rate sufficient to treat to rid the soil of any diseases and pests found therein. After the fumigant is applied, the soil should remain undisturbed for a sufficient time to allow the fumigant to rid the soil of all deleterious organisms. Applications to the soil include attention to soil

10 conditions such as high moisture content, soil temperatures below 13°C/55°F at a 20cm/8-inch depth, and/or cloddy soils with high plant /weed trash content. The treated field may be planted after the elapsed planting interval time.

EXAMPLE 7

15 Methyl iodide, HFC-245fa, and chloropicrin are combined to form an azeotrope or azeotrope-like composition according to the present invention. The composition is applied as a fumigant to a field, prior to planting the soil, in water through the drip irrigation system under a plastic tarp. The fumigant is applied at a broadcast rate sufficient to treat to rid the soil of any diseases

20 and pests found therein. After the fumigant is applied, the soil should remain undisturbed for a sufficient time to allow the fumigant to rid the soil of all deleterious organisms. Applications to the soil include attention to soil conditions such as high moisture content, soil temperatures below 13°C/55°F at a 20cm/8-inch depth, and/or cloddy soils with high plant /weed trash

25 content. The treated field may be planted after the elapsed planting interval time.

EXAMPLE 8

This example relates to a structural fumigation process. Prior to structural fumigation, all open flames and glowing heat filaments are turned off or disconnected. Methyl iodide and HFC-245fa are combined to form an azeotrope or azeotrope-like composition according to the present invention.

- 5 The composition is applied as a fumigant to tarped or sealed structures for an exposure period necessary to rid the structure of all deleterious organisms, followed by an aeration period long enough to flush the unused fumigant and any warning gas from the structure. Because the methyl iodide/245fa mixture is odorless and does not irritate the eyes or skin, trace amounts of a warning
- 10 agent (e.g. chloropicrin) are introduced into the structure prior to fumigation to act as a warning agent. The required dosage of the fumigant is influenced by the temperature at the site of the pest, the length of the exposure period, containment or the rate the fumigant is lost from the structure, and the susceptibility of the pest to be controlled.

15

While the present invention has been particularly shown and described with reference to preferred embodiments, it will be readily appreciated by those of ordinary skill in the art that various changes and modifications may be made without departing from the spirit and scope of the invention. It is intended that

- 20 the claims be interpreted to cover the disclosed embodiment, those alternatives which have been discussed above and all equivalents thereto.

What is claimed is:

1. An azeotropic or azeotrope-like composition comprising a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, which 5 composition is a gas at temperatures of about 30°C or below.
2. The composition of claim 1 wherein the at least one fluorocarbon or hydrofluorocarbon has a boiling point of from about 0° C to about 50° C.
- 10 3. The composition of claim 1 wherein the at least one fluorocarbon or hydrofluorocarbon has an average Ozone Depletion Potential of about 0.05 or less.
4. The composition of claim 1 wherein the at least one fluorocarbon or 15 hydrofluorocarbon has a 100-year Global Warming Potential of about 1,000 or less.
5. The composition of claim 1 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluoropropane.
- 20 6. The composition of claim 1 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluorobutane.
7. The composition of claim 1 wherein the at least one fluorocarbon or 25 hydrofluorocarbon comprises cis-1,3,3,3-tetrafluoropropene.
8. The composition of claim 1 which comprises from about 5 to about 70 weight percent methyl iodide, and from about 30 to about 95 weight percent of the at least one fluorocarbon or hydrofluorocarbon.

30

9. The composition of claim 1 which has a boiling point of from about 8°C to about 14.5°C.
10. The composition of claim 1 which has a density of from about 1.5 g/cc to about 2.4 g/cc.  
5
11. A fumigant comprising the composition of claim 1.
12. A fumigant comprising the composition of claim 5.  
10
13. A fumigant comprising the composition of claim 6.
14. A fumigant comprising the composition of claim 7.
15. A fumigant comprising the composition of claim 1, which composition comprises from about 5 to about 70 weight percent methyl iodide and about 30 to about 95 weight percent of the at least one fluorocarbon or hydrofluorocarbon; a density of from about 1.5 g/cc to about 2.4 g/cc; and wherein the at least one fluorocarbon or hydrofluorocarbon of the composition  
20 has an average Ozone Depletion Potential of about 0.05 or less, and a 100-year Global Warming Potential of about 1,000 or less.
16. The fumigant of claim 15 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluoropropane.  
25
17. The fumigant of claim 15 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluorobutane.
18. The fumigant of claim 15 wherein the at least one fluorocarbon or  
30 hydrofluorocarbon comprises cis-1,3,3,3-tetrafluoropropene (HFC-1234ze).

19. A fumigation process which comprises:
  - a) providing a fumigant comprising an azeotropic or azeotrope-like composition which comprises a mixture of methyl iodide and at least one fluorocarbon or hydrofluorocarbon, which composition is a gas at temperatures of about 30°C or below; and
  - b) applying the fumigant to a material to be fumigated.
20. The process of claim 19 wherein the fumigant is applied in step (b) as a gas.
21. The fumigation process of claim 19 wherein step (b) is conducted at an ambient temperature of from about 0°C to about 50°C.
- 15 22. The process of claim 19 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluoropropane.
23. The process of claim 19 wherein the at least one fluorocarbon or hydrofluorocarbon comprises 1,1,1,3,3-pentafluorobutane.
- 20 24. The process of claim 19 wherein the at least one fluorocarbon or hydrofluorocarbon comprises cis-1,3,3,3-tetrafluoropropene (HFC-1234ze).
- 25 25. The process of claim 19 wherein the fumigant is applied to soil in step (b).
26. The process of claim 19 wherein the fumigant is applied to wood in step (b).
- 30 27. The process of claim 19 wherein the fumigant is applied to a building surface in step (b).