

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : G01N 33/68, C07K 7/06, 14/78, 16/18, C12N 5/20	A1	(11) International Publication Number: WO 96/30765 (43) International Publication Date: 3 October 1996 (03.10.96)
(21) International Application Number: PCT/EP96/01228 (22) International Filing Date: 21 March 1996 (21.03.96) (30) Priority Data: 9506050.5 24 March 1995 (24.03.95) GB (71) Applicant (for all designated States except US): OSTEOMETER BIOTECH A/S [DK/DK]; OsteoPark, Herlev Hovedgade 207, DK-2730 Herlev (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): FLEDELIUS, Christian [DK/DK]; Nordre Frihavns-gade 101, 2. th., DK-2100 Copenhagen Ø (DK). BONDE, Martin [DK/DK]; Lystoftevej 28, DK-2800 Lyngby (DK). QVIST, Per [DK/DK]; Taarbæk Strandvej 103G, DK-2930 Klampenborg (DK). (74) Agent: SMART, Peter, J.; W.H. Beck, Greener & Co., 7 Stone Buildings, Lincoln's Inn, London WC2A 3SZ (GB).	(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: ASSAYING PROTEIN FRAGMENTS IN BODY FLUIDS		
(57) Abstract <p>Degradation of body proteins such as bone collagen is studied by detecting in body fluids isomerized peptide fragments of said proteins in which peptide bonding to the α-carboxylic acid group of aspartic acid or asparagine has become isomerized to an amide bond to the side chain carboxylic acid group of aspartic acid. Antibodies specific for the isomerized peptide sequences are used in assays.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

ASSAYING PROTEIN FRAGMENTS IN BODY FLUIDS

The present invention relates to the assaying of collagen or other protein degradation products and materials useful therefor.

Collagens and Disorders of Collagen Metabolism

Osteoporosis is the most common bone disease in humans. Primary osteoporosis, accompanied by increased susceptibility to fractures, results from a progressive reduction in skeletal bone mass. It is estimated to affect 15-20 million individuals in the USA alone. Its basis is an age-dependant imbalance in bone remodelling, i.e. in the rates of formation and resorption of bone tissue.

In the USA about 1.2 million osteoporosis-related fractures occur in the elderly each year including about 538,000 compression fractures of the spine, about 227,000 hip fractures and a substantial number of early fractured peripheral bones. Between 12 and 20% of the hip fractures are fatal because they cause severe trauma and bleeding, and half of the surviving patients require nursing home care. Total costs from osteoporosis-related injuries now amount to at least \$10 billion annually in the USA (Riggs, New England Journal of Medicine, 327:620-627 (1992)).

Osteoporosis is most common in postmenopausal women who, on average, lose 15% of their bone mass in the 10 years after menopause. This disease also occurs in men as they get older and in young amenorrhic women athletes. Despite the major, and growing, social and economic consequences of osteoporosis, the availability of reliable assays for measuring bone resorption rates in patients or in healthy subjects is very limited. Other disorders entailing (and correlated with) abnormalities in collagen metabolism include Paget's disease, Marfan's syndrome, osteogenesis imperfecta, neoplastic growth in collagenous tissue, dwarfism, rheumatoid arthritis, osteoarthritis and vasculitis syndrome.

-2-

Three known classes of human collagen have been described to date. The Class I collagens, subdivided into types I, II, III, V, and XI, are known to form fibrils. The amino-acid sequence of type I-III (to the extent it has been elucidated) is given in Appendix A of WO95/08115.

Collagen type I accounts for more than 90% of the organic matrix of bone. Therefore, in principle, it is possible to estimate the rate of bone resorption by monitoring the degradation of collagen type I. Likewise, a number of other disease states involving connective tissue can be monitored by determining the degradation of collagen. Examples are collagen type II degradation associated with rheumatoid arthritis and osteoarthritis and collagen type III degradation in vasculitis syndrome.

Amino acid sequences of human type III collagen, human pro $\alpha 1$ (II) collagen, and the entire prepro $\alpha 1$ (III) chain of human type III collagen and corresponding cDNA clones have been investigated and determined by several groups of researchers; see Loil et al., *Nucleic Acid Research* 12:9383-9394 (1984); Sangiorgi et al., *Nucleic Acids Research*, 13:2207-2225 (1985); Baldwin et al., *Biochem J.*, 262:521-528 (1989); and Ala-Kokko et al., *Biochem. J.*, 260:509-516 (1989).

Type I, II, and III collagens are all formed in the organism as procollagen molecules, comprising N-terminal and C-terminal propeptide sequences, which are attached to the core collagen molecules. After removal of the propeptides, which occurs naturally in vivo during collagen synthesis, the remaining core of the collagen molecules consists largely of a triple-helical domain having terminal telopeptide sequences which are non-triple-helical. These telopeptide sequences have an important function as sites of intermolecular cross-linking of collagen fibrils extracellularly. The alpha-helical region also includes crosslinkable sites.

Intermolecular cross-links provide collagen fibrils with biomechanical stability. The formation of these cross-links is initiated by modification of lysine and hydroxylysine

residues to the corresponding aldehydes. Several of these residues located on adjacent chains of collagen will spontaneously form different intermolecular cross-links. The exact position of the sites for cross-linking on collagen
5 telopeptides and from the helical region has been previously described. See, for example, Kühn, K., in *Immunochemistry of the extracellular matrix*, 1:1-29, CRC Press, Inc., Boca Raton, Florida (1982), Eyre, D.R., *Ann.Rev. Biochem.*, 53:717-48 (1984) or US Patent No. 5140103 and 5455179. Furthermore, the
10 amino acid sequences of some potential sites for cross-linking in type I, II, and III collagen are given in Table 1 below.

The fibrous proteins, collagen and elastin, are cross-linked by a unique mechanism based on aldehyde formation from lysine or hydroxylysine side chains. Four homologous loci of
15 cross-linking are evident in molecules of type I, II and III collagens (for review see Kühn, K., in *Immunochemistry of the extracellular matrix*, 1:1-29 (1982)). Two are aldehyde sites, one in each telopeptide region. The other two sites are hydroxylysine symmetrically placed at about 90 residues from
20 each end of the molecule. When collagen molecules pack into fibrils, these latter sites in the helical region align and react with telopeptide aldehydes in adjacent molecules. There is now strong evidence that 3-hydroxypyridinium residues are the mature cross-link coming from hydroxylysine-derived
25 aldehydes. The mature cross-linking residues of the other pathway, i.e. from aldehyde formation of lysine residues, are however, still unknown.

As illustrated by formula in EP-0394296 discussed below, the two 3-hydroxypyridinium cross-links have been found to be
30 hydroxylysyl pyridinoline (also known simply as "pyridinoline") and lysyl pyridinoline (also known as "deoxypyridinoline"). These cross-linking compounds are naturally fluorescent. Some hydroxylysyl pyridinoline cross-link are found to be glycosylated as discussed for instance
35 in EP-A-0424428.

However, as described in Last et al, Int. J. Biochem. Vol. 22, No. 6, pp 559-564, 1990 other crosslinks occur naturally in collagen.

5 Prior Art Assays for Collagen Degradation

In the past, assays have been developed for monitoring degradation of collagen in vivo by measuring various biochemical markers, some of which have been degradation products of collagen.

10 For example, hydroxyproline, an amino acid largely restricted to collagen, and the principal structural protein in bone and all other connective tissues, is excreted in urine. Its excretion rate is known to be increased in certain conditions, notably Paget's disease, a metabolic bone disorder
15 in which bone turnover is greatly increased, as discussed further below.

For this reason, urinary hydroxyproline has been used extensively as an amino acid marker for collagen degradation; Singer, F.R. et al., Metabolic Bone Disease, Vol. II (eds. Avioli, L.V., and Kane, S.M.), 489-575 (1978), Academic Press,
20 New York.

US Patent No. 3600132 discloses a process for the determination of hydroxyproline in body fluids such as serum, urine, lumbar fluid and other intercellular fluids in order
25 to monitor deviations in collagen metabolism. The Patent states that hydroxyproline correlates with increased collagen anabolism or catabolism associated with pathological conditions such as Paget's disease, Marfan's syndrome, osteogenesis imperfecta, neoplastic growth in collagen tissues and in
30 various forms of dwarfism.

Bone resorption associated with Paget's disease has also been monitored by measuring small peptides containing hydroxyproline, which are excreted in the urine following degradation of bone collagen; Russell et al., Metab. Bone Dis. and Rel.
35 Res. 4 and 5, 2250262 (1981), and Singer, F.R., et al., supra.

In the case of Paget's disease, the increased urinary hydroxyproline probably comes largely from bone degradation; hydroxyproline, however, generally cannot be used as a specific index for bone degradation. Much of the hydroxyproline in urine may come from new collagen synthesis (considerable amounts of the newly made protein are degraded and excreted without ever becoming incorporated into tissue fabric), and from turnover of certain blood proteins as well as other proteins that contain hydroxyproline.

Furthermore, about 80% of the free hydroxyproline derived from protein degradation is metabolised in the liver and never appears in the urine. Kivirikko, K.I., *Int. Rev. Connect. Tissue Res.* 5:93 (1970), and Weiss, P.H. and Klein, L., *J. Clin. Invest.* 48:1 (1969). Hydroxyproline is a good marker for osteoporosis as it is specific for collagen in bones even if it is not specific for bone resorption, but it is troublesome to handle.

Hydroxylysine and its glycoside derivatives, both peculiar to collagenous proteins, have been considered to be more accurate than hydroxyproline as markers of collagen degradation. However, for the same reasons described above for hydroxyproline, hydroxylysine and its glycosides are probably equally non-specific markers of bone resorption; Krane, S.M. and Simon, L.S., *Develop. Biochem.* 22:185 (1981).

Other researchers have measured the cross-linking compound 3-hydroxypyridinium in urine as an index of collagen degradation in joint diseases. See, for back-ground and as examples, Wu and Eyre, *Biochemistry*, 23:1850 (1984); Black et al., *Annals of the Rheumatic Diseases*, 45:969-973 (1986); and Seibel et al., *The Journal of Dermatology*, 16:964 (1989). In contrast to the present invention, these prior researchers have hydrolysed peptides from body fluids and then looked for the presence of free 3-hydroxypyridinium residues.

Assays for determination of the degradation of type I, II, and III collagen are disclosed in EP-0394296 and US Patent No. 4973666 and US Patent No. 5140103. However, these Patents

-6-

are restricted to collagen fragments containing the cross-linker 3-hydroxypyridinium. Furthermore, the above mentioned assays require tedious and complicated purifications from urine of collagen fragments containing 3-hydroxypyridinium to
5 be used for the production of antibodies and for antigens in the assays.

At present very few clinical data using the approach described in US Patent No. 4973666 and US Patent No. 5140103 are available. Particularly, no data concerning the correlation between the urinary concentration (as determined by
10 methods described in the above mentioned patents) of 3-hydroxypyridinium containing telopeptides of type I collagen and the actual bone loss (as determined by repeated measurements by bone densitometry) have been published. The presence
15 of 3-hydroxypyridinium containing telopeptides in urine requires the proper formation in bone tissue of this specific cross-linking structure at various times before the bone resorbing process. Very little information on these processes is available and it would be desirable to avoid this
20 dependence of the correct formation of the cross-linking structure.

GB Patent Application No. 2205643 reports that the degradation of type III collagen in the body can be quantitatively determined by measuring the concentration of an N-terminal telopeptide from type III collagen in a body fluid.
25 This method uses antibodies generated to N-terminal telopeptides released by bacterial collagenase degradation of type III collagen, said telopeptides being labelled and used in the assay.

Schröter-Kermani et al., Immunol. Invest. 19:475-491 (1990) describe immunological measurement systems based on CNBr fragments of collagen type I and II. Use is made of pepsin-solubilised collagen, leaving the telopeptides in the tissue (cf. the above mentioned GB Patent Application No.
35 2205643). There is therefore no conformity between the fragments and the antibodies raised therefrom. Further, the

reference only describes measurements on extracted tissue samples.

The development of a monoclonal antibody raised against pepsin-solubilised type I collagen is described in Werkmeister et al., Eur. J. Biochem. 1987:439-443 (1990). The antibody is used for immunohistochemical staining of tissue segments and for measuring the collagen content in cell cultures. The measurements are not carried out on body fluids.

EP Patent Application No. 0505210 describes the development of antibody reagents by immunisation with purified cross-linked C-terminal telopeptides from type I collagen. The immunogen is prepared by solubilising human bone collagen with bacterial collagenase. The antibodies thus prepared are able to react with both cross-linked and non-cross-linked telopeptides, and cross-linkers other than pyridinoline.

International Patent Application No. WO 91/09114 discloses certain synthetic peptides which are used to promote cellular adhesion to a solid substrate. The use of the synthetic peptides as immunological reagents is not mentioned.

There are a number of reports indicating that collagen degradation can be measured by quantitating certain procollagen peptides. Propeptides are distinguished from telopeptides and alpha-helical region of the collagen core by their location in the procollagen molecule and the timing of their cleavage in vivo; see US Patent No. 4504587; US Patent No. 4312853; Pierard et al., Analytical Biochemistry 141:127-136 (1984); Niemela, Clin. Chem. 31/8:1301-1304 (1985); and Rohde et al., European Journal of Clinical Investigation, 9:451-459 (1979).

EP Patent Application No. 0298210 and No. 0339443 both describe immunological determination of procollagen peptide type III and fragments thereof. Further, a method based on the measurement of procollagen is disclosed in EP Patent Application No. 0465104.

The use of synthetic peptides with sequences derived from type IX collagen for the development of immunological reagents

is disclosed in PCT Patent Application No. WO90/08195. Likewise the application describes the use of the antibodies thus produced for the determination of type IX collagen fragments in body fluids.

5 US Patent No. 4778768 relates to a method of determining changes occurring in articular cartilage involving quantifying proteoglycan monomers or antigenic fragments thereof in a synovial fluid sample.

10 Dodge, J: Clin Invest 83:647-661 (1981) discloses methods for analysing type II collagen degradation utilising a polyclonal antiserum that specifically reacts with unwound alpha-chains and cyanogen bromide-derived peptides of human and bovine type II collagens. The degradation products of collagen were not detected in a body fluid, but histochemically by
15 staining of cell cultures, i.e. by "in situ" detection.

WO94/03813 describes a competitive immunoassay for detecting collagen or collagen fragments in a sample wherein a binding partner containing a synthetic linear peptide corresponding to the non-helical C-terminal or N-terminal
20 domain of collagen is incubated with an antibody to the linear synthetic peptide and the sample, and wherein the binding of the antibody to the binding partner is determined.

WO95/08115 relates to assay methods in which collagen fragments in a body fluid are determined by reaction with an
25 antibody which is reactive with a synthetic peptide. The assay may be a competition assay in which the sample and such a peptide compete for an antibody, possibly a polyclonal antibody raised against fragments of collagen obtained by collagenase degradation of collagen. Alternatively, it may
30 be an assay in which an antibody, possibly a monoclonal antibody, is used which has been raised against such a synthetic peptide.

One particular peptide fragment which we have found in body fluid, particularly urine, is of the formula:

EKAHDGGR

|

EKAHDGGR

Formula 1

|

5 K

In the above formula, K-K-K represents cross-link which may for instance be a hydroxypyridinium cross-link but may be any naturally occurring cross-link and specifically any of those discussed in the above referenced paper of Last et al.

A larger peptide fragment including the above smaller fragment is reported in EP 0394296.

We have now discovered that a proportion of the "peptide" fragments in body fluid are related to peptides of equivalent amino acid sequence, e.g. peptides of formula 1, by the isomerization of aspartic acid in the formula to isoaspartic acid. We put "peptides" in quotes here as of course the isomerization means that these species are no longer properly regarded as being peptides.


The isomerization of proteins containing aspartic acid has been reported previously to be a spontaneous reaction occurring under physiological conditions.

See for instance Brennan et al Protein Science 1993, 2, 331-338, Galletti et al, Biochem. J. 1995, 306, 313-325, Lowenson et al, Blood Cells 1988, 14, 103-117 and Oliya et al, Pharmaceutical Research, Vol. 11, No. 5, 1994, p.751.

The isomerization has the effect of transferring that part of the peptide chain which runs downstream of the aspartic acid residue in the carboxy terminus direction from the alpha carboxylic acid of the aspartic acid to which it is bonded via a peptide bond in the normal protein to the side chain carboxylic acid in a non-peptide amide bond, as shown below:

-10-


5



10



15



20 The non-peptide bonded aspartic acid residue is termed "isoaspartic acid".

Similar isomerization can occur in proteins containing asparagine residues (i.e. with $-NH_2$ instead of $-OH$ in the starting protein in the above reaction scheme).

25 The above discovery indicates that this isomerization also occurs in bone tissue and the extent of isomerization is expected therefore to be marker for the age of the bone tissue concerned.

30 Furthermore, the presence amongst such bone peptide fragments of the isomerized peptides provides confirmation that the fragments indeed derive from bone degradation and not some other source such as the degradation of newly formed collagen never incorporated into bone.

35 Accordingly, the present invention now provides in a first aspect a method of measurement of the rate of degradation of a body protein such as collagen, e.g. from bone,

comprising determining the amount of one or more isoaspartic acid containing species in a body fluid.

The isomerized peptides in question may be characteristic of type I, type II or type III collagen, but preferably are characteristic of type I collagen.

More preferably, such a method determines the amount of one or more specific isoaspartic acid containing isomerized peptides present in said body fluid.

Preferably, the method determines the amount of the isomerized peptide of formula 2 (below) present in said body fluid:



wherein one or both of * is isoaspartic acid, or of one or more isomerized peptides incorporating an epitope present in the isomerized peptide of formula 2 which contains isoaspartic acid.

In the above formula, K-K-K is a cross-link such as a hydroxypyridinium cross-link which may be pyridinoline (which may be glycosylated or non-glycosylated) or deoxypyridinoline.

Preferably, said determination is carried out using an immunological binding partner specific for an isoaspartic acid containing species present in the sample during the procedure, preferably said isomerized peptide of formula 2 or a isomerized peptide incorporating an epitope present in the isomerized peptide of formula 2 which contains isoaspartic acid.

The immunological binding partner may be a monoclonal or polyclonal antibody. By the requirement that the immunological binding partner be specific for the isoaspartic acid containing species is meant that the immunological binding

-12-

partner distinguishes between said species and the analogous aspartic acid containing species (peptide) to an extent useful in the assay.

Suitable immunological binding partners also include
5 fragments of antibodies capable of binding the same antigenic determinant including Fab, Fab' and F(ab')₂ fragments.

Preferably, the immunological binding partner is an antibody raised against a linear isomerized peptide, preferably a synthetic isomerized peptide, corresponding to a
10 sequence within collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen protein sequence

The assay may take many forms including ELISA, RIA, or IRMA, procedures for which are too well known to warrant
15 description here.

In a second aspect, the invention includes the use in an assay for collagen derived isomerized peptides of a synthetic isomerized peptide having an amino acid sequence corresponding to a sequence within collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said
20 collagen protein sequence. In a competition assay, the said synthetic isomerized peptide may be used to compete for an immunological binding partner with one or more isomerized peptides in the sample.

In an ELISA of this type, the synthetic peptide may be immobilised on a solid support. A sample may be incubated with a polyclonal antibody for the synthetic isomer of a peptide in contact with the solid support and after washing, a peroxidase-conjugated (revealing) antibody may be added.
30 After further incubation, a peroxidase substrate solution is added. By competition, peptide isomer in the sample reactive with the antibody inhibits the peroxidase reaction.

Alternatively, the synthetic peptide isomer may be used to raise a monoclonal immunological binding partner. The
35 synthetic isomerized peptide need not then be a competing agent in the assay. For instance, collagenase treated

-13-

collagen may be purified and immobilised onto the solid support and an ELISA may be carried out using a monoclonal antibody.

5 Accordingly, in a third aspect, the invention includes an antibody, preferably a monoclonal antibody, specific for an amino acid sequence corresponding to a sequence within a protein, e.g. collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said protein, e.g. collagen sequence.

10 In a preferred embodiment of this aspect of the invention, the antibody is specific for an isomerized peptide sequence including the sequence EKAHiDGGR or containing an epitope specific for the presence of iD present in said sequence, wherein iD is isoaspartic acid.

15 Accordingly, this aspect of the invention includes an antibody, preferably a monoclonal antibody reactive with an epitope containing, contained in, or constituted by the peptide isomer sequence EKAHiDGGR, wherein iD is isoaspartic acid.

20 In a fourth aspect, the invention provides an antibody, preferably a monoclonal antibody, raised against a peptide isomer having an amino acid sequence corresponding to a sequence within a protein, e.g. collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen protein sequence.

25 The invention includes cell lines producing monoclonal antibodies according to the third or fourth aspects of the invention.

30 The invention also includes antibodies according to the third or fourth aspects of the invention coupled to a detectable marker. Suitable detectable markers include, but are not limited to, enzymes, chromophores, fluorophores, coenzymes, enzyme inhibitors, chemiluminescent materials, paramagnetic materials, spin labels, radio-isotopes, nucleic acid or
35 nucleic acid analogue sequences.

In a fifth aspect, the invention includes the use in an assay for collagen or other protein derived isomerized peptides of an antibody specific for an amino acid sequence corresponding to a sequence within the protein, (e.g. collagen) with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said protein (e.g. collagen) sequence to obtain information regarding the amount of isoaspartic acid containing peptide isomer or isomers in said body fluid.

10 In a sixth aspect, the invention includes a synthetic peptide isomer having an amino acid sequence corresponding to a sequence within collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen protein sequence, preferably in at least the
15 substantial absence of the corresponding peptide.

Preferably there is a glycine residue adjacent the aspartic acid residue in the native peptide form of the amino acid sequence, as an adjacent glycine facilitates the isomerization of aspartic acid.

20 Antibodies may be prepared which are respectively selective for one or more aspartic acid containing peptides and for their isoaspartic acid containing analogues. It is then possible to carry out an assay for both variants of the peptide or peptides. The relative amount of isoaspartic acid
25 will provide an indication of the age of the protein which is being broken down protein and of the bone if the assay is for a type of collagen fragment. Accordingly, in a sixth aspect the invention provides a method of obtaining information regarding collagen resorption in a patient, comprising
30 measuring in a body fluid the relative amounts of at least one aspartic acid containing peptide derived from collagen and a corresponding isoaspartic acid containing peptide analogue.

The invention also includes test kits useful in the methods described above. Such kits may comprise an antibody
35 according to the third or fourth aspect of the invention, or

-15-

similarly specific antibody fragment, preferably in combination with any one or more of:-

a synthetic peptide analogue containing isoaspartic acid reactive with the antibody,

5 an antibody-enzyme conjugate and/or a substrate therefor,

an enzyme conjugate-substrate reaction stopping composition, or
a wash solution.

10 The invention may be applied both to humans and to animals.

Suitable body fluids include, human or animal urine, blood, serum, plasma and synovial fluid. It is contemplated that the method may also be used e.g. on saliva and sweat.

15 The body fluid may be used as it is, or it may be purified prior to the contacting step. This purification step may be accomplished using a number of standard procedures, including, but not limited to, cartridge adsorption and elution, molecular sieve

20 chromatography, dialysis, ion exchange, alumina chromatography, hydroxyapatite chromatography, and combinations thereof.

The invention is described in more detail below. Reference is made to the appended drawings.

25

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the HPLC separation of a synthetic peptide and a peptide analogue of the sequence EKAH*GGR differing at * between isoaspartic acid (peak 2) and normal aspartic acid (peak 3) as described in Example 3a;

Figure 2 shows the differing reactivity of the separated peptide and peptide analogue of Figure 1 in two forms of ELISA, as described in Example 3a; and

35 Figures 3(a) and 3(b) show the fluorescence (a) and absorbance (b) of crosslinked peptides and peptide analogues from urine separated by HPLC as described in Example 3(b).

-16-

Figures 4 and 5 show the results of assays on serum using different antibodies to monitor the effect of bisphosphonate treatment on bone resorption.

In a preferred embodiment of the method according to the invention, the assaying of type I, II or III collagen fragments in urine or other body fluid is performed by an inhibition ELISA (enzyme linked immunosorbent assay) by metering off a sample of urine body fluid contacting the sample with a synthetic peptide analogue having a sequence derived from collagen and with an antibody, which is immunoreactive with the synthetic peptide analogue. The synthetic peptide analogue is immobilised on a solid support. The antibody may be raised against the synthetic peptide analogue or may be raised against collagen degradation products and screened for by use of such a synthetic peptide analogue.

Preparation of synthetic peptide analogues

The preparation of synthetic peptides and peptide analogues may be performed according to procedures well known in the art, e.g. by solid-phase peptide synthesis techniques commonly described as "Merrifield synthesis". Also classical solution phase techniques may be used. Sequences of interest include potential sites for collagen cross-linking (see for example Kühn, K., in *Immunochemistry of the extracellular matrix*, 1:1-29 (1982), Eyre, D.R., *Ann. Rev. Biochem.* 53:717-48 (1984), or US Patent No. 5140103). Examples of such peptide sequences are given in Table 1 below. The conventional peptide synthesis method may produce a mixture of peptide (with normal peptide bonded aspartic acid) and peptide analogue with isomerization of the bonding to the aspartic acid. Generally such a mixture will be satisfactory as the peptide will be insert in the assay. However, heating such a mixture will normally produce isomerization of the peptide content to the iso-form.

Regarding the synthetic peptide analogues, it is possible to omit (or add) one or more amino acid residues from (or to) the crosslinkable site sequences without substantial loss of the ability to (a) raise antibodies recognising the isoaspartic acid analogue of the corresponding native collagen fragment or (b) inhibit the binding of such antibodies to the said analogue of the native fragment. It is possible to use longer collagen fragments and/or chimeric peptide analogues to raise the antibodies and, in principle, it is not necessary to use the same peptide analogue as the immunogen and the competitor in a competition assay.

TABLE 1

Examples of Amino Acid Sequences with Potential Sites for Cross-linking in Various Types of Collagen to be Used as a Basis for Isoaspartic Acid Containing Synthetic Peptide analogues according to the Present Invention

Collagen Type I

Potential sites

in telopeptide analogues: N C

$\alpha 1$ (I) N-term. (Iso)Asp-Glu-Lys-Ser-Thr-Gly-Gly ($\alpha 1$ (I)N1)

$\alpha 1$ (I) C-term. Glu-Lys-Ala-His- (Iso)Asp-Gly-Gly-Arg ($\alpha 1$ (I)C1)

$\alpha 2$ (I) N-term. Gln-Tyr- (Iso) -Asp-Gly-Lys-Gly-Val-Gly ($\alpha 2$ (I)N1)

Collagen Type II

Potential sites

in telopeptide analogues: N C

$\alpha 1$ (II) N-term. Gly-Asp-Ile-Lys- (Iso)Asp-Ile-Val

$\alpha 1$ (II) C-term. Glu-Lys-Gly-Pro- (Iso)Asp

-18-

Collagen Type III

Potential sites

in telopeptide analogues: N

C

5 α 1 (III) N-term. (Iso)Asp-Val-Lys-Ser-Gly-ValPreparation of antibodies

The methods for preparation of both monoclonal and polyclonal antibodies are well known in the art. For example, see
10 Campbell, A.M., Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 12 (1986). It is possible to produce antibodies to synthetic isomerized peptides by immunisation. However, because of the relatively small molecular weight of these compounds it is preferred that the hapten be conjugated
15 to a carrier molecule. Suitable carrier molecules include, but are not limited to, bovine serum albumin, thyroglobulin, ovalbumin, tetanus toxoid, and keyhole limpet hemocyanin. The preferred carrier is bovine serum albumin. To present the hapten in its most immunogenic form to the antibody producing
20 cells of the immunised animal a number of alternative coupling protocols can be used. Suitable procedures include, but are not limited to, glutaraldehyde, carbodiimide, and periodate. Preferred binding agents are glutaraldehyde and carbodiimide.

The preparation of antibodies may be carried out by
25 conventional techniques including immunisation with collagen fragments containing natural isomerization or synthetic isomerized peptides conjugated to a carrier. To improve the immunogenicity it is preferred that the immunogen be mixed with an adjuvant before injection. Examples of adjuvants
30 include, but are not limited to, aluminium hydroxide, Freund's adjuvant, and immune-stimulating complexes (ISCOMs). ISCOMs can be made according to the method described by Morein, B. et al., Nature 308:457-460 (1984).

Either monoclonal or polyclonal antibodies to the hapten-carrier molecule can be produced. For the production of
35 monoclonal antibodies it is preferred that mice are immunised.

-19-

Spleen cells from the immunised mouse are harvested, homogenised, and thereafter fused with cancer cells in the presence of polyethylene glycol to produce a cell hybrid which produces monoclonal antibodies specific for isomerized peptide fragments derived from collagen. Suitable cancer cells include, but are not limited to, myeloma, hepatoma, carcinoma, and sarcoma cells. Detailed descriptions of the production of monoclonal antibodies are provided in Goding, J.W., in Monoclonal Antibodies: Principles and Practice, (1986). A preferred preliminary screening protocol comprises the use of synthetic isomerized peptides conjugated to a carrier and coated on to the solid surface of a microtitre plate.

For the preparation of polyclonal antibodies, which are reactive with isomerized peptide fragments derived from collagen, different animal species can be immunised. Suitable species include, but are not limited to, chicken, rabbit and goat. Chicken and rabbit are preferred.

Antibodies so produced may be screened for suitability for use according to the invention by testing for reactivity with an isoaspartic acid containing synthetic peptide analogue of appropriate sequence.

Antibody fragments are prepared by methods known in the art (see E. Ishikawa. Journal of Immunoassay 3:209-327 (1983)).

Conduct of Immunoassays

Accordingly, by utilisation of an immunoassay with the antibodies prepared as above it is possible to assay a biological fluid sample without prior fractionation or hydrolysis. The specificity for the desired collagen fragments in the biological fluid may be supplied by the antibody in combination with the use of a synthetic isomerized peptide (against which the antibody was raised or in any event with which the antibody is immunochemically reactive) in the assay construction.

-20-

As an alternative the immunoassay may be performed using a monoclonal antibody. The basic idea of this assay design is to shift the specificity of the assay from the antigen (synthetic peptide isomer to collagen) to the antibody (from rabbit antiserum to monoclonal antibody). Using this construction the assay does not need to make further use of a synthetic peptide isomer. This version of the immunoassay is suitably performed by incubating the patient sample or a standard solution with a peroxidase-conjugated antibody solution in a microtiter plate precoated with purified collagenase-treated collagen. After washing, the wells of the plate are incubated in the dark with a substrate solution. The colour reaction is stopped by the addition of a stopping solution, and finally the absorbance is measured.

The immunoassays themselves may be conducted using any procedure selected from the variety of standard assay protocols generally known in the art. As it is generally understood, the assay is constructed so as to rely on the interaction between the specific immunological binding partner and the desired analyte for specificity and to utilise some means to detect the complex formed by the analyte and the immunological binding partner. The immunological binding partner may be complexed to a solid support and used as a capture immunological binding partner for the analyte. This protocol may be run in a direct form, wherein the formation of analyte-immunological binding partner complex is detected, e.g. by a fluorescent, radioactive or enzymatic label, or it may be run in a competitive format wherein a labelled standard competes with the analyte for the immunological binding partner. The format may also be constructed as an agglutination assay or the complex may be precipitated by addition of a suitable precipitant to the reaction mixture. The specific design of the immunoassay protocol is open to a wide variety of choice, and the number of clinical assay devices and protocols available in the art is multitudinous. For a variety of such protocols, see US. Patent No. 5001225.

-21-

The antibodies and revealing reagents for the conduct of an immunoassay using standard detection protocols, for example radioisotope labelling, fluorescent labelling or ELISA, either in a direct or competitive format, may conveniently be supplied as kits which include the necessary components and instructions for the assay. In one embodiment of the invention such a kit includes a microtiter plate coated with a relevant synthetic isomerized peptide, standard solutions for preparation of standard curve, a body fluid (e.g. urine) control for quality testing of the analytical run, rabbit antibodies reactive with the above mentioned synthetic peptide isomer, anti-rabbit immunoglobulins conjugated to peroxidase, a substrate solution, a stopping solution, a washing buffer and an instruction manual.

Since immunoassays can be constructed using antibodies and specific synthetic isomerized peptides, the ratios of the corresponding collagen fragment sequences in an appropriate biological fluid can be determined as well as their individual levels and their total. Thus, the assay can be designed to include antibodies which will result in determination of several isoaspartic acid containing peptide analogues and optionally the native peptide sequences or determination of a single isoaspartic acid containing peptide analogue sequence, or any desired combination thereof.

In addition to the use of the herein specified isomerized peptides as indicators of bone resorption, bone metabolic balance is advantageously determined by the substantially simultaneous determination of a marker of the formation of bone in the same or other appropriate biological fluid from the same individual. "Substantially simultaneous" means the same day, preferably within 4 hours. For example such markers include osteocalcin (also known as bone GLA protein of BGP), propeptides of procollagen type I, bone alkaline phosphatase and total alkaline phosphatase. Suitable methods for the determination of these markers can be found, for example, in Delmas, P.D., et al., J. Bone Min. Res. (1986) 1:333-337.

-22-

The assay of the present invention which provides an index to determination of the metabolic status of tissues, which generate collagen-derived peptides and isomerized peptides when degradation occurs, is useful in a variety of contexts. First, when considering the degradation of type I collagen, the assays are methods to assess an abnormal condition of a subject by indicating, for example, excessive bone resorption. This may show the presence of an osteoporotic condition or the metastatic progress of a malignancy. Other conditions characterised by excessive bone resorption include Paget's disease and hyperparathyroidism. Likewise, a number of other disease states involving connective tissue may be monitored by determination of the degradation of collagen. Examples are collagen type II degradation associated with rheumatoid arthritis and osteoarthritis and collagen type III degradation in vasculitis syndrome. Since the condition of the subject can be monitored continuously, application of these assays can also be used to monitor the progress of therapy administered to treat these or other conditions. Further, the assays can be used as a measure of toxicity, since the administration of toxic substances often results in tissue degradation.

Thus the assays may be applied in any situation wherein the metabolic condition of collagen tissues can be used as an index of the condition, treatment, or effect of substances directly administered to the subject or to which the subject is exposed in the environment.

The following examples are intended to illustrate, but not to limit the invention.

30

EXAMPLE 1: IMMUNOASSAYS FOR SPECIFIC
 PEPTIDE SEQUENCES IN URINE

Three isomerized peptides ($\alpha 1(I)C1$, $\alpha 1(I)N1$, and $\alpha 2(I)N1$) (see Table 1) prepared by solid-phase techniques are used for the preparation of immunogens. For immunisation, the peptide isomers are covalently attached to bovine serum albumin using

35

-23-

carbodiimide or glutaraldehyde reagents and methods well known in the art. Both monoclonal and polyclonal antibodies are raised against the peptide isomers. For production of monoclonal antibodies, Balb/c mice are immunised with peptide isomer-BSA conjugates, and hybridoma cell lines are prepared using standard techniques after fusion of cells from the spleen or lymph nodes with Ag8 myeloma cells. Polyclonal antibodies are raised in rabbits and chicken. Screening of both antisera and hybridoma cell media were performed by ELISA using microtiter plates coated with the appropriate peptide isomer-protein carrier conjugate prepared using carbodiimide reagents and methods well known in the art.

Assays for three of the peptide isomer sequences ($\alpha 1(I)C1$, $\alpha 1(I)N1$, and $\alpha 2(I)N1$) in urine are performed by an inhibition ELISA as follows:-

Urine samples (10 to 25 μ l) possibly containing collagen fragments or solutions containing 0.015-15 μ g peptide isomer/ml as reference standards, respectively, are added to 75 μ l of immunological binding partners for the peptide isomers diluted 1:5,000 - 1:20,000 in phosphate buffered saline containing 0.1% Tween-20 detergent (PSB-T) and including 0.1% (w/v) of BSA. Each sample is prepared in duplicate in flat-bottomed, 96-well microtiter plates previously coated with peptide isomer-protein carrier conjugate containing the appropriate peptide isomer. After 60 minutes, the plates are washed with PBS-T (3 times) and the bound antibodies are detected by standard techniques with a horseradish peroxidase labelled antibody prepared against the species of the primary antibody. Peroxidase substrate is added and the colour development is measured at 450 nm in an automated microtiter plate reader after stopping the enzyme reaction using 1 M H_3PO_4 . Samples containing the analyte decrease the binding of primary antibody to the immobilised peptide isomer on the plate and thus have a reduced colour concentration. The amount of analyte in the sample is quantified with reference to

-25-

isolated from urine, was evaluated. The MAbA7 ELISA is a competitive assay wherein the MAbA7 monoclonal raised against the 8AA peptide (EKAHDGGR in which D is normal aspartic acid) is used and collagen degraded collagen immobilised in a micro-titer tray competes with collagen fragments in the sample for binding to the monoclonal antibody (MAbA7).

The ELISAs used in this example are carried out as follows:-

(a) THE POLYCLONAL ISO-ELISA

The polyclonal ELISA is based on an immobilised synthetic peptide analogue with an amino acid sequence of eight amino acids (8AA) specific for an isomerised part of the C-telo-peptide of the $\alpha 1$ -chain of type I collagen (Glu-Lys-Ala-His-(Iso)ASP-Gly-Gly-Arg). During incubation with an antibody reactive with this sequence, a competition takes place between the immobilised peptide iso-form analogue and the breakdown products of the $\alpha 1$ -chain of type I collagen in the sample.

Briefly, a 25 μ L sample or standard is added to each well of a formula 1 antigen-coated microplate, followed by 75 μ L of antiserum raised against collagenase treated type 1 collagen. The plates are incubated for 1 hour at room temperature under agitation and washed five times with a washing buffer. A goat anti-rabbit immunoglobulin G horseradish peroxidase conjugate (100 μ L) is added to each well. After incubation for 1 hour at room temperature, plates are washed five times as before. The enzyme substrate (100 μ L/well) is added, and after 15 minutes of incubation in the dark, the reaction is stopped by adding 100 μ L phosphoric acid (1 mol/L). The optical density of 450 nm is measured with a microplate reader. Duplicate measurements are performed for each sample, and the data are expressed as nanograms per mol creatinine (Cr), measured by a standard colorimetric technique. This assay is referred to herein as ISO-ELISA.

The MAbA7 ELISA

MAbA7 is a monoclonal antibody raised as described above in mouse against the peptide of the sequence EKAHDGGR, i.e. with normal aspartic acid.

5 This assay version is carried out in three steps as follows:

In the first step the wells of a microtiter plate pre-coated with purified collagenase-treated collagen (CTC) are incubated for 1 hour at room temperature with 15 μ l standard
10 solution or sample and 100 μ l of peroxidase-conjugated monoclonal antibody solution.

After washing in the second step the wells are incubated in the dark for 15 minutes at room temperature with 100 μ l substrate solution. Finally, in the third step, the colour
15 reaction is stopped by the addition of 100 μ l stopping solution. The absorbance at 450 nm is measured within 2 hours.

EXAMPLE 3(a)

20 This example demonstrates that antibodies specific for either synthetic EKAHDGGR or EKAHiDGGR can be developed.

A mixture of synthetic EKAHDGGR and EKAHiDGGR was separated by HPLC (Figure 1, peak 3 and peak 2, respectively). Sequence analysis confirmed that peak 2 contained isomerized
25 aspartate. This was indicated by the sequencing operation being halted after the histidine residue. Peak 3 contained regular aspartate and sequencing was able to proceed normally. Fractions from the HPLC profile were collected and tested for immunoreactivity in both MAbA7 ELISA and polyclonal ISO-ELISA.
30 It was demonstrated that the two assays detected separate peaks, i.e. peak 2 was detected by ISO-ELISA and peak 3 by MAbA7 ELISA.

The results are shown in Figure 2.

EXAMPLE 3 (b)

This example demonstrates that ISO-ELISA-related molecules can be purified from urine, and that these molecules can be separated into di-peptide related species containing isomerized or regular aspartate.

5 First urine was immunopurified using MAbA7. MAbA7 was coupled to CNBr activated Sepharose™ according to the manufacturer's instructions. Urine was diluted 1:3(v/v) in PBS buffer and the pH adjusted to 8.0 using 1 M NaOH. Eight-
10 hundred ml diluted urine was recirculated on to a column (14 cm³) for 24 hr at 4°C at a flow rate of 0.8 ml/min. After washing the column with 200 ml PBS buffer, pH 8.0, bound antigens were eluted using 20 ml 50% saturated (NH₄)₂SO₄ containing 1% TFA (v/v) and stored at -20°C. The eluted
15 antigens were desalted using a 1 ml C18 Sep-Pak™ column. The Sep-Pak™ column was conditioned with 20 ml 80% methanol (v/v) and equilibrated with 20 ml water containing 1% TFA (v/v) before adding the antigen. Bound antigens were washed with 20 ml of water containing 1% TFA (v/v) and eluted with 40%
20 acetonitrile (v/v) containing 0.1% TFA (v/v), freeze-dried and stored at -20°C.

Molecules from selected peaks obtained by the above procedure were further separated by HPLC using trifluoroacetic acid (TFA) as ionpair. Fractions were analysed by amino acid
25 composition, amino acid sequencing, mass spectrometry, and immunoreactivity in MAbA7 and ISO-ELISA.

The fluorescence (3-hydroxypyridinium cross-links) and absorbance of the HFBA HPLC profile is demonstrated in Figure 3. By further separation pure preparations were obtained and
30 tested by mass spectrometry and immunoreactivity (Table 2), as well as amino acid sequencing (Table 3) and amino acid composition (Table 4). Data demonstrates that three separate molecules all with the molecular weight around 2036 kDaltons could be identified (Table 2). Amino acid sequencing was
35 blocked after histidine in peak F-24-17-10, whereas the complete sequence EKAHDGGR was obtained for the two other

-28-

peaks (F-26-18-09 and F-29-19-24, Table 3). Amino acid composition analysis (which does not separate isomerised and regular aspartate) confirmed the three peaks contained the same amino acids.

5 These data suggest that three different, cross-linked dipeptide related species were identified in urine. The molecules differed in having either isomerised or regular aspartate in the sequence EKAHDGGR. Peak F-24-17-10 is suggested to contain two peptide analogues of the sequence
10 EKAHiDGGR, peak F-26-18-09 one EKAHiDGGR and one EKAHDGGR, and peak F-29-19-24 two cross-linked peptides of the sequence EKAHDGGR.

 Additionally, a similar observation was done on molecules of molecule weight around 2039. These did not contain 3-
15 hydroxypyridinium cross-links (as fluorescence was absent), and the nature of the cross-linker was not determined.

TABLE 2

20 Detection by MAbA7 and ISO-ELISA of
Peptide related species separated by HPLC from Urine

Fraction No.	Mass	Pyr	DPyr	MAbA7 ng/ml	ISO-ELISA ng/ml
25 F-18-17-14	2038.2			58	974
F-20-18-18	2039.5			406	350
F-22-18,19,20-16	2038.9			972	118
F-24-17-10	2036.1	+		114	2110
F-26-18-09	2036.2	+		2114	2203
30 F-26-18-12	1858.9		+	85	783
F-29-19-24	2036.0	+		3149	221
F-30-20-12	1858.8		+	2108	33

-29-

TABLE 3

Sequence Data for Peptide related molecules in
F-24-17-10, F-26-18-09, and F-29-19-24

5

Cycle	F-24-17-10	F-26-18-09	F-29-19-24	Expected	
10	1	Glu	Glu	Glu	Glu
	2	-	-	-	-
	3	Ala	Ala	Ala	Ala
	4	His	His	His	His
	5	-	Asp	Asp	Asp
15	6	-	Gly	Gly	Gly
	7	-	Gly	Gly	Gly
	8	-	Arg	Arg	Arg
	9	-	-	-	-
	10	-	-	-	-

20

TABLE 4

5 Amino Acid Composition of the Peptides
F-24-17-10, F-26-18-09, and F-29-19-24 (residue/peptide)

	Amino Acid	F-24-17-10	F-26-18-09	F-29-19-24	Expected
10	Asp	1.0	1.0	0.9	1
	Glu	1.3	1.2	1.3	1
	Ser	0.4	0.2	0.6	-
	Gly	2.2	2.0	2.4	2
	His	1.4	1.0	1.0	1
15	Thr	0.1	0.0	0.2	-
	Ala	1.0	1.0	1.0	1
	Pro	0.0	0.0	0.0	-
	Tyr	0.0	0.0	0.0	-
	Val	0.1	0.1	0.1	-
20	Met	0.0	0.0	0.0	-
	Ile	0.1	0.1	0.1	-
	Leu	0.1	0.1	0.2	-
	Lys	0.3	0.1	0.3	-
	Arg	0.8	0.9	0.8	1
25	Total	8.8	7.6	9.0	7

The polyclonal assay is shown to be specific for the peptide analogue containing isoaspartic acid residues and the MAbA7 assay is specific for the peptide containing normal aspartic acid residues. Di-peptide analogues containing both the isomerised as well as the non-isomerised aspartic acid will be detected by both assays.

The application of these two assays to a urine sample therefore allows a comparison to be made of the level of

isoaspartic acid containing isomerized peptide and the level of the normal form, giving an indication of the degree to which old established bone tissue is being broken down.

5 EXAMPLE 4

The clinical significance of the presence of isomerised peptide analogues in serum was investigated by subjecting serum samples from postmenopausal women taken before and after
10 treatment with a bisphosphonate, Ibandronate (Boehringer Mannheim), to analysis in both a serum version of the ISOELISA polyclonal assay described above and the MabA7 based ELISA monoclonal assay.

The assay developed for measurement of type I degradation
15 products in serum is based on an immobilised synthetic peptide/peptide isomer mixture with the amino acid sequence EKAHDGGR present in peptide and in peptide isomer form. Rabbits were immunised with the isomerized peptide conjugated to BSA using a two step carbodiimide procedure. If the
20 isomerized peptide is substantially more antigenic in the rabbits employed than the peptide form of the amino acid sequence one can immunise with the peptide/peptide isomer mixture conjugated to BSA. For coating of microtitre plates the peptide was conjugated to thyroglobulin using glutar-
25 aldehyde. During incubation with this antibody a competition takes place between the immobilised peptide isomer (the immobilised peptide being inactive in the assay) and the breakdown products of type I collagen in serum. As the content of the peptide in the solution increases, less
30 antibody will bind to the immobilised peptide isomer leading to a decreasing optical density.

In brief, 50 μ l Standard (synthetic 8AA-peptide isomer/peptide mixture) or unknown sample in a test tube is added 100
35 μ l Assay Buffer (500 mM TRIS, 0.0% Tween 20, 1.0% BSA; pH= 6.5). From these 150 μ l, 50 μ l is pipetted into the appropriate wells in the pre-coated ELISA plate. Then 50 μ l

-32-

Antibody Solution (rabbit antiserum to EKAHiDGGR diluted 1+20.000 in Assay Buffer) is added to each well, the plate is covered with sealing tape and incubated at room temperature for 60 min on a shaking device. All the following procedures were also carried out at room temperature. After incubation the plates were washed three times with diluted Washing Buffer (25 mmol/l TRIS and 50 mmol/l NaCl pH=7.2).

Peroxidase conjugated Antibody (HRP-conjugated goat antibodies to rabbit IgG (Jackson Immunochemicals, PA) diluted 1+4000 in Assay Buffer), 100 μ l/well) was added and the sealed wells were incubated 60 min on a shaking device. Following another washing procedure, 100 μ l of TMB Substrate Solution was added to all wells which were sealed and incubated for 15 min. The enzyme reaction was stopped after 15 min by addition of 100 μ l Stopping Solution. The optical density was read in an ELISA-reader at 450 nm.

A calibration curve was constructed on a log-linear graph paper by plotting the mean absorbances of the five standards (25 ng/ml-500 ng/ml). The concentration of EKAHiDGGR equivalent in each patient specimen were determined by interpolation on the calibration curve.

The results are shown in Figures 4 and 5. In Figure 4, no significant change is seen in the serum level of molecules reactive in the monoclonal assay directed to molecules competing for antibody reactive with the peptide sequence EKAHDGGR before and after fifteen months treatment with a therapeutic expected to reduce the rate of bone resorption. In Figure 5, one sees a drop of more than 60% following treatment for six months with 2.5 mg/day of Ibandronate in the level of molecules reactive in the polyclonal assay directed to molecules competing with the non-peptide EKAHiDGGR for antibody in the polyclonal serum, reflecting the expected result of the therapeutic treatment.

This example indicates that the sum of peptides in serum incorporating the sequence EKAHDGGR, which may include cross-linked peptides of the kind described in US 5455179 and

-33-

similar molecules based on other cross-links or not cross-linked at all, are not indicators of the rate of bone resorption, but that the related non-peptide molecules produced by isomerisation of the bonding of the aspartic acid
5 in the amino acid sequence are.

It is suggested that the specificity for the isomerised analogue of the octapeptide leaves the ISOELISA insensitive to molecules present in serum which are reactive in the MabA7 ELISA. The level of isomerisation is believed to be related
10 to the age of the bone, being therefore a quantifiable marker for turnover of mature bone, whilst the molecules interfering in the monoclonal assay do not relate to bone resorption and could originate from the turnover of "young" collagen molecules, e.g. from proteolytic breakdown of extracellular
15 collagen not yet incorporated into the bone matrix, or generated during turnover of short-lived type 1 collagen at non-skeletal sites.

Furthermore, although it is believed that the body has mechanisms for repairing damage to proteins by isomerisation
20 at aspartate, it may be that in bone collagen is shielded from these repair mechanisms making isomerisation an unusually good indicator of the resorption of "old" collagen in the case of bone.

Whilst the invention has been described with reference
25 to specific examples, many variations thereof are possible within the scope of the invention. For instance the invention may be applied to the aspartic acid or asparagine containing proteins generally and not only to collagens.

CLAIMS

1. A method of measurement of the rate of degradation of a body protein comprising determining the amount of one or more
 5 isoaspartic acid containing species in a body fluid.

2. A method as claimed in Claim 1, which determines the amount of at least one isoaspartic acid containing isomerized peptide in said body fluid.
 10

3. A method as claimed in Claim 2, wherein the isomerized peptide or peptides are characteristic of collagen.

4. A method as claimed in Claim 3, comprising determining
 15 the amount of the peptide analogue of formula 2 (below) present in said body fluid:



wherein K-K-K is any naturally occurring cross-link and * is
 25 isoaspartic acid, or of one or more peptide analogues incorporating an epitope present in the peptide analogue of formula 2 which contains isoaspartic acid.

5. A method as claimed in any preceding claim, wherein said
 30 determination is carried out using an immunological binding partner specific for an isoaspartic acid containing species present in the sample during the procedure.

6. A method as claimed in Claim 5, wherein the immunological
 35 binding partner is an antibody raised against a linear peptide analogue corresponding to a sequence within collagen with

-35-

isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen sequence.

5 7. The use in an assay for peptide derived isomerized peptides of a synthetic isomer of a peptide having an amino acid sequence corresponding to a sequence within said protein with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said protein sequence.

10 8. An antibody specific for an amino acid sequence corresponding to a sequence within a protein with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said protein sequence.

15 9. An antibody as claimed in Claim 8, which is specific for a peptide isomer sequence including the amino acid sequence EKAHiDGGR, or for an epitope included in said sequence and containing iD, wherein iD is isoaspartic acid.

20 10. An antibody raised against a peptide isomer having an amino acid sequence corresponding to a sequence within collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen sequence.

25 11. A cell line producing monoclonal antibody being and antibody as claimed in any one of claims 8 to 10.

12. An antibody as claimed in any one of Claims 8 to 10, coupled to a detectable marker.

30

13. The use in an assay for protein derived isomerized peptides of an antibody specific for an amino acid sequence corresponding to a sequence within a protein with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said protein sequence to obtain information regarding

35

-36-

the amount of isoaspartic acid containing peptide analogue or peptide analogues in said body fluid.

5 14. A synthetic peptide analogue having an amino acid sequence corresponding to a sequence within collagen with isoaspartic acid substituting in said amino acid sequence for aspartic acid in said collagen sequence.

10 15. A synthetic peptide analogue as claimed in Claim 14, wherein there is a glycine residue adjacent the isoaspartic acid residue.


15 16. A method of obtaining information regarding protein degradation in a patient, comprising measuring in a body fluid the relative amounts of at least one aspartic acid containing peptide derived from a protein and a corresponding isoaspartic acid containing peptide analogue..

20 17. A test kit for use in an assay method as claimed in any one of Claims 1 to 6, comprising an antibody as claimed in any one of Claims 8 to 10, or a similarly specific antibody fragment, in combination with any one or more of:-

25 a synthetic peptide analogue containing isoaspartic acid reactive with the antibody or antibody fragment,
an antibody-enzyme conjugate and/or a substrate therefor,
an enzyme conjugate-substrate reaction stopping composition, or
30 a wash solution.


1/4

FIG. 1.



2/4

FIG. 2.



3/4

FIG. 3(a).







FIG. 3(b).





INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 96/01228

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N33/68 C07K7/06 C07K14/78 C07K16/18 C12N5/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC 6 G01N C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 122, no. 11, 13 March 1995 Columbus, Ohio, US; abstract no. 131800, H. TANAKA ET AL.: "Urinary excretion of beta-aspartylpeptide in relation to collagen catabolism." page 986; XP002008160 see abstract & HISSU AMINOSA KENKYU, vol. 136, 1992, JAPAN, pages 26-29, --- -/--	1-3, 13-16

Further documents are listed in the continuation of box C. Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search 16 July 1996	Date of mailing of the international search report 22. 07. 96
---	--

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016	Authorized officer Cartagena y Abella, P
---	---

INTERNATIONAL SEARCH REPORT

International Application No

PC./EP 96/01228

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>J. ORG. CHEM., vol. 48, 1983, pages 2014-2021, XP000574968 J. T. CAPECCHI ET AL.: "Critical examination of a method for the analysis of alpha and omega linkages in peptides containing aspartic acid and glutamic acid." see page 2014; table II</p>	14, 15
A	<p>WO,A,94 14844 (BAYLINK DAVID J) 7 July 1994</p>	
A	<p>WO,A,92 21698 (WASHINGTON RES FOUND) 10 December 1992</p>	
A	<p>WO,A,89 04491 (WASHINGTON RES FOUND) 18 May 1989 cited in the application & EP,A,0 394 296</p>	
A	<p>DE,A,42 25 038 (BOEHRINGER MANNHEIM GMBH) 3 February 1994 cited in the application & WO,A,943 813</p>	
A	<p>EP,A,0 505 210 (ORION YHTYMAE OY) 23 September 1992 cited in the application</p>	
A	<p>THE INTERNATIONAL JOURNAL OF BIOCHEMISTRY, vol. 22, no. 5, 1990, pages 559-564, XP002008159 J. A. LAST ET AL.: "Biosynthesis of collagen crosslinks" cited in the application</p>	
A	<p>WO,A,95 08115 (OSTEOMETER A S ;QVIST PER (DK); BONDE MARTIN (DK)) 23 March 1995 cited in the application</p>	
A	<p>JOURNAL OF CLINICAL ENDOCRINOLOGY AND METABOLISM, vol. 80, no. 3, March 1995, pages 864-868, XP000575691 M. BOMDE ET AL.: "Applications of an enzyme immunoassay for a new marker of bone resorption (CrossLaps): follow-up on hormone replacement therapy and osteoporosis risk assessment."</p>	

1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 96/01228

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9414844	07-07-94	CA-A- 2152667	07-07-94
		EP-A- 0675903	11-10-95
WO-A-9221698	10-12-92	US-A- 5320970	14-06-94
		AU-B- 665953	25-01-96
		AU-B- 2140992	08-01-93
		CA-A- 2110263	10-12-92
		EP-A- 0587719	23-03-94
		JP-T- 6508138	14-09-94
		US-A- 5472884	05-12-95
WO-A-8904491	18-05-89	US-A- 4973666	27-11-90
		AT-T- 117437	15-02-95
		DE-D- 3852827	02-03-95
		DE-T- 3852827	29-06-95
		DE-T- 394296	28-04-94
		EP-A- 0394296	31-10-90
		HK-A- 101395	30-06-95
		JP-T- 3500818	21-02-91
		US-A- 5473052	05-12-95
		US-A- 5455179	03-10-95
		US-A- 5472884	05-12-95
		US-A- 5140103	18-08-92
		US-A- 5300434	05-04-94
		US-A- 5320970	14-06-94
US-A- 5532169	02-07-96		
DE-A-4225038	03-02-94	CA-A- 2139592	17-02-94
		WO-A- 9403813	17-02-94
		EP-A- 0659276	28-06-95
		FI-A- 950378	27-01-95
		JP-T- 7509778	26-10-95
		NO-A- 950325	27-01-95
EP-A-0505210	23-09-92	JP-A- 7020126	24-01-95
WO-A-9508115	23-03-95	AU-B- 7652194	03-04-95