

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 December 2006 (07.12.2006)

PCT

(10) International Publication Number  
**WO 2006/129204 A2**

(51) International Patent Classification:

*C12N 15/11* (2006.01) *A01P 7/04* (2006.01)  
*A61K 31/713* (2006.01) *C12N 1/21* (2006.01)  
*A01N 63/00* (2006.01) *C07K 14/435* (2006.01)

(21) International Application Number:

PCT/IB2006/002360

(22) International Filing Date: 31 May 2006 (31.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

05253339.5 31 May 2005 (31.05.2005) EP  
60/685,765 31 May 2005 (31.05.2005) US

(71) Applicant (for all designated States except US): **DE-  
VGEN NV** [BE/BE]; Technologiepark 30, Zwijnaarde,  
B-9052 Gent (BE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KALETTA, Titus Jan**  
[DE/BE]; Boomgaardstraat 1, B-9820 Merelbeke (BE).

(74) Agents: **BALDOCK, Sharon, Claire** et al.; BOULT  
WADE TENNANT, Verulam Gardens, 70 Gray's Inn  
Road, London WC1X 8BT (GB).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,  
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,  
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,  
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,  
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: RNAI FOR THE CONTROL OF INSECTS AND ARACHNIDS

(57) Abstract: The present invention describes a new non-compound based approach for insect and/or arachnid control. The present inventors have identified for the first time novel targets for RNAi, which can effectively control insect and/or arachnid pest populations. Accordingly, the invention provides both nucleotide and amino acid sequences for the novel targets. Also provided are RNA constructs including double stranded RNA regions for mediating RNAi in insects, DNA constructs, expression vectors, host cells and compositions for controlling insects and/or arachnids using RNAi. Finally, the invention also provides for the use of the constructs, vectors, host cells and compositions in control of insects and/or arachnids populations and suitable kits for use in an RNAi based method of controlling insect and/or arachnid pests.



WO 2006/129204 A2

**RNAi for the control of insects and arachnids****Field of the invention.**

The present invention relates to the field of double-stranded RNA (dsRNA) mediated gene silencing in insect species. More particularly, the present invention relates to genetic constructs designed for the expression of dsRNA corresponding to novel targets identified for the first time herein. These constructs are particularly useful in dsRNA mediated insect pest control, especially control of household insects or arachnids, for instance cockroaches.

**Background to the invention.**

Pest control and particularly insect and/or arachnid control, especially control of household insects, ecto-parasites and insects relevant for public health and hygiene (e.g. urban protection) such as cockroaches, fleas, ants, termites, earwigs, mosquitos, flies and house crickets is an important field. The presence of insects in locations such as at home, in offices, restaurants, hospitals or warehouses undoubtedly causes distress because there is a common public perception that insects such as cockroaches or flies live in places that are dirty and not well kept.

These insects do not only causes distress but also contaminate food and eating utensils, destroy fabric and paper products and impart stains and unpleasant odours to surfaces they contact. Furthermore, these insects can pose health risks as carriers for bacteria. For example, cockroaches may transmit bacteria that cause food poisoning (*Salmonella* spp. and *Shigella* spp.). German cockroaches are believed capable of transmitting disease-causing organisms such as *Staphylococcus* spp., *Streptococcus* spp., hepatitis virus and coliform bacteria. They also have been implicated in the spread of typhoid and dysentery. Some people, especially those with asthma, are sensitive to the allergens produced by these cockroaches.

There are various chemical insecticides and capturing devices developed and commercially available for fighting household pests. However, increasing efficacy of these means is usually linked with increased health risk. Insecticides may contaminate food which is nearly unavoidable in places such as kitchens, restaurants or food storages and incorporation may cause health risks to humans.

The solution to this problem of contamination has been to use less toxic insecticides. However, when applying less toxic insecticides, there is an increased probability that the insect may become resistant over time.

Insecticides act by binding to a certain insect protein, such as an acetylcholine receptor for example, and cause death of the pest species by either deactivating or over-activating the protein. Insecticides have been developed to be safe at certain concentration, but can and do impact on human health when incorporated at higher dosages or over long periods. In contrary to agrochemicals, household insecticides are applied in places where food is stored or prepared and food contamination and contact to humans cannot be avoided.

One alternative to chemical pesticides is to utilise biological agents. Over the last few years, downregulation of genes (also referred to as "gene silencing") in multicellular organisms by means of RNA interference or "RNAi" has become a well-established technique.

In general, RNAi comprises contacting the organism with a double stranded RNA fragment or "dsRNA" (generally either as two annealed complementary single strands of RNA or as a hairpin construct) that comprises a nucleotide sequence that corresponds to (at least part of) the nucleotide sequence of the gene to be downregulated (the "target gene"). Reference is *inter alia* made to the International application WO 99/32619 (Carnegie Institute of Washington), the International application WO 99/53050 (CSIRO), the International application WO 00/01846 (Devgen) and to Fire et al., Nature, Vol. 391, pp.806-811, February 1998.

In nematodes, RNAi can be performed by feeding the nematode with the dsRNA fragment as such, or alternatively with a bacterial strain that either contains the dsRNA fragment or that upon ingestion by the nematode is capable of expressing the dsRNA fragment. For this so-called "RNAi by feeding", reference is *inter alia* made to the International application WO 00/01846 by applicant, and to WO 99/32619 cited above, in which the nematode *C. elegans* is used.

Many dsRNA constructs have been described in the art. A classic dsRNA is produced from a DNA construct comprising two convergent promoters flanking the sequence complementary to the target sequence which needs to be downregulated (see for example WO00/01846 (Devgen)). As the technology of dsRNA mediated gene silencing advanced, new constructs were designed to improve the dsRNA for various purposes.

In order to produce the dsRNA more efficiently, a stem-loop-stem structure or "hairpin" was developed. As described in, for example, document WO 99/53050

- 3 -

(CSIRO), this hairpin allows the formation of dsRNA from one single RNA transcript. The RNA transcript comprises the sense and anti-sense version of the complementary sequence, separated by a non-complementary loop structure allowing the RNA transcript to fold back and to base pair into a dsRNA stem portion.

5 DsRNA gene silencing finds application in many different areas, such as for example dsRNA mediated gene silencing in clinical applications (WO2004/001013) and in plants. In plants, dsRNA constructs useful for gene silencing have also been designed to be cleaved and to be processed into Short interfering RNAs (siRNAs).

10 RNAi has also been proposed as a means of protecting plants against plant parasitic nematodes, i.e. by expressing in the plant (e.g. in the entire plant, or in a part, tissue or cell of a plant) one or more nucleotide sequences that form a dsRNA fragment that corresponds to a target gene in the plant parasitic nematode that is essential for its growth, reproduction and/or survival. Reference may be made to the International application WO 00/01846 by the present applicant, US patent 6,506,559 (based on WO 15 99/32619), and to International applications WO 01/96584, WO 01/37654 and WO 03/052110 for a description of such techniques.

Elbashir et al. (Nature, 411, 494-498, 2001) have demonstrated effective RNAi-mediated gene silencing in mammalian cells using dsRNA fragments of 21 nucleotides in length (also termed small interfering RNAs or siRNAs).

20 WO 03/004644 describes delivery of dsRNA to arthropods in general terms and is incorporated herein by reference. WO 03/004644 details down regulation of the reporter gene GUS (Clonetech) using RNAi in *Drosophila melanogaster* and down regulation of the vATPase gene in *H. armigera*.

25 WO 01/34815 relates to baculovirus expression vectors which produce dsRNA and the use of these vectors in pest control.

Although the technique of RNAi has been generally known in the art in plants, nematodes and mammalian cells for some years, to date little is known about the use of RNAi to down-regulate gene expression in insects and/or arachnids. In addition, little is known on the application of RNAi to control pest species such as household insects, 30 ecto-parasites and insects and/or arachnids relevant for public health and hygiene.

Constructs suitable and efficient for dsRNA mediated pest control, should meet at least some of the following requirements

- (1) the dsRNA must be taken up by the pest organisms
- (2) the dsRNA must have good stability in the pest organisms

- 4 -

(3) the dsRNA must be effective in the pest organism to control its viability, growth and/or development and/or

(4) the dsRNA must guarantee maximized safety and minimized environmental impact.

5 It is now the purpose of the present invention to provide dsRNA constructs, which meet the above-mentioned requirements.

### **Description of the invention**

10 The present invention describes a new non-compound based approach for insect and/or arachnid control. The active ingredient is a nucleic acid, a double-stranded RNA (dsRNA), which can be used as an insecticidal or arachnidicidal formulation (for example in baits or gel applications). The sequence of the dsRNA matches a part of an essential insect gene and causes down-regulation of the insect target via RNA interference (RNAi). As a result of the down-regulation of mRNA, the dsRNA prevents expression of  
15 the corresponding insect protein and hence causes death, growth arrest or sterility of the insect and/or arachnid.

### **Targets**

20 The present inventors have identified for the first time novel targets for RNAi, which can effectively control insect or arachnid pest populations.

For the avoidance of doubt, a target is defined herein as a gene whose protein product is required for the insect and/or arachnid to maintain its normal physiological and biochemical functions. Inhibition of the expression of the target gene limits the insect's and/or arachnids ability to feed, grow, or survive. Examples of insect and/or arachnid  
25 genes that may be employed in the practice of the invention include essential genes, genes involved in processes such as development, metabolism, or neurotransmission, and genes whose products are targets of existing insecticides and/or arachnids. In a preferred embodiment of the invention, the target is part of pathways required for cellular function such as transcription, translation, the cytoskeleton, cell-cycle, metabolism  
30 (anabolism or catabolism), endocytosis, intracellular and intercellular transport, calcium binding, nucleus import and export, nucleic acid binding, signal peptidase-protein binding, the proteasome, vesicle transport, neuro-transmission, water-balance, ion-balance, splicing, mitosis, meiosis, chromosome organisation, stability or integrity, micro

- 5 -

RNAs, siRNAs, posttranslational protein modifications, electron transport, apoptosis, membrane integrity, and cell adhesion.

The novel target genes identified in the present invention comprise:

5 A) structural proteins, for instance tropomyosin 1 (GenBank AF260897) (SEQ ID NOs 41 and 42), actin 5C (GenBank AY004248) (SEQ ID NOs 57 and 58), and homologous or heterologous proteins having the same biological function in the same or in other insect and/or arachnid species;

10 B) metabolic enzymes, for instance the HMG Coenzyme A synthase (GenBank X73679) (SEQ ID NO 49 and 50), and homologous or heterologous proteins having the same biological function in the same or in other insect and/or arachnid species;

C) enzymes involved in ion/ pH homeostasis, such as V-ATPase and homologous or heterologous proteins having the same biological function in the same or in other insect and/or arachnid species;

15 D) enzymes involved in the transcriptional/ translational machinery, such as for instance

- o Ribosomal protein S4 homolog (SEQ ID NOs 1 and 2)
- o Ribosomal protein S9 homolog (SEQ ID NOs 11 and 12)
- o Ribosomal protein L9 homolog (SEQ ID NOs 21 and 22)
- o Ribosomal protein L19 homolog (SEQ ID NOs 31 and 32)

20 Accordingly, according to a first aspect there is provided a nucleic acid molecule comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180 and 181, or an orthologous nucleotide sequence from an insect and/or arachnid species, wherein the orthologous nucleotide sequence has at  
25 least 70%, preferably at least 75%, 80%, 85%, 90%, more preferably at least about 95% and even more preferably at least about 96%, 97%, 98%, most preferably at least 99% sequence identity with the nucleotide sequence of any one of SEQ ID NOs 1, 11, 21 and 31. Preferred orthologous sequences comprise, or if being used according to the methods of the invention include, sequences from household insects, ecto-parasites and  
30 insects and/or arachnids relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. More preferred orthologous sequences are from cockroaches (Blattodea) such as but not limited to *Blattella* spp. (e.g. *Blattella germanica*

(german cockroach)), *Periplaneta* spp. (e.g. *Periplaneta americana* (American cockroach) and *Periplaneta australiasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus* spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex* spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g. *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)), *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)) and *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)).

According to another aspect there is provided a nucleic acid molecule comprising, consisting essentially of, or consisting of the nucleotide sequence as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200.

SEQ ID NO's 4, 6, 7, 16, 27, 26, 27, 36 and 37 all represent the nucleotide sequences of specific primers which were utilised to identify the novel sequences of the invention in a selective and specific manner.

The novel identified genes all represent components of the transcriptional/translational machinery of *Blatella germanica*. By inhibiting expression of these genes or by inhibiting expression of the novel identified target genes, through RNAi, an important pest may be controlled.

It is predicted, and would be understood by the skilled person, that also orthologues of these novel target genes represent further targets for down-regulation in the control of other insect and/or arachnid species. Thus, orthologues of the novel nucleic acid molecules of the present invention are also contemplated.

Protein or nucleotide sequences are likely to be homologous if they show a "significant" level of sequence similarity or identity. Truly homologous sequences are related by divergence from a common ancestor gene. Sequence homologues can be of two types: (i) where homologues exist in different species they are known as orthologues. e.g. the  $\alpha$ -globin genes in mouse and human are orthologues. (ii) paralogues are homologous genes within a single species. e.g. the  $\alpha$ - and  $\beta$ - globin genes in mouse are paralogues. By "orthologues" is meant herein both types of homologues referred to above.

In one embodiment, the orthologue will share at least about 40%, 50% or 60 % nucleotide sequence identity with the nucleotide sequence as set forth in any one of SEQ ID NO 1, 11, 21, 31, 41, 49 or 57. Preferably, the orthologue will share at least about 70%, 75%, 80%, 85%, 90%, more preferably at least about 95% and even more preferably at least about 96%, 97%, 98% or 99% sequence identity with the nucleotide sequence as set forth in any one of SEQ ID NOs 1, 11, 21, 31, 41, 49 or 57.

According to another embodiment, the invention encompasses target genes which are insect or arachnidae orthologues of a gene comprising, consisting essentially of, or consisting of a nucleotide sequence as represented in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200. By way of example, orthologous may comprise a nucleotide sequence as represented in any of SEQ ID NOs 71 to 200, or a fragment of at least 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 or 27 nucleotides thereof. A non-limiting list of insect or arachnid orthologous genes or sequences comprising at least a fragment of 17 nucleotides of one of the sequences of the invention is given in Tables 4 and 5. The sequences presented in Tables 4 and 5 are intended to form part of the present invention. Thus, orthologues comprise, consist essentially of or consist of any of the sequences set forth in Tables 4 and 5.

According to another aspect, the invention thus encompasses any of the methods described herein for controlling insect and/or arachnid infestation or infection, comprising contacting insects and/or arachnids with a double-stranded RNA, wherein the double-stranded RNA comprises annealed complementary strands, one of which has a nucleotide sequence which is complementary to at least part of the nucleotide sequence of a target gene comprising a fragment of at least 17, 18, 19, 20 or 21 nucleotides of any of the sequences as represented in SEQ ID NOs 71 to 200, whereby the double-



stranded RNA is taken up by the insect and/or arachnid and thereby controls growth, kills or prevents infestation or infection by the insect and/or arachnid. Said insect and/or arachnid may comprise, consist essentially of or consist of any target organisms/species described herein.

5        Related nucleic acid molecules encompassed by the invention may also be defined in terms of hybridisation to a nucleic acid molecule comprising the nucleotide sequence of any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180 and 181. Preferably, the hybridisation conditions are moderate stringency hybridisation conditions  
10        and even more preferably high stringency hybridisation conditions. Such conditions of moderate and high stringency would be immediately familiar to one of skill in the art. For example, a hybridization reaction incubated at 42°C in a solution comprising 50% formamide, 5xSSC, and 1% SDS or at 65°C in a solution comprising 5xSSC and 1% SDS, with a wash in 0.2xSSC and 0.1% SDS at 65°C represent suitable high stringency  
15        conditions.

The invention also provides the protein products of these novel target genes, and orthologues thereof.

Accordingly, according to a second aspect there is provided a protein comprising the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32 or an  
20        orthologous protein having a conserved amino acid sequence from a further insect and/or arachnid species.

As mentioned above, it is predicted also that orthologues of the novel target genes will represent further targets for down-regulation in the control of other insect and/or arachnid species. Thus, orthologues of the novel protein molecules of the  
25        present invention are also contemplated.

In one embodiment, the orthologue will share at least about 40% amino acid sequence identity with the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32. Preferably, the orthologue will share at least about 40%, 50%, 60%, 65%, 70%, 80%, more preferably at least about 90% and even more preferably at least  
30        about 96%, 97%, 98% or 95% amino acid sequence identity with the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32.

In another embodiment, the invention also provides for a nucleic acid encoding a protein comprising the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32. The nucleic acid molecules encompassed by this aspect of the invention

also include those which are functionally equivalent in that they encode the same protein molecule. Thus, all nucleic acid molecules which are possible due to the degeneracy of the genetic code are intended to fall within the scope of this aspect of the invention.

5     Target organisms/species

      The "target species" as used in the present invention, may be any insect or arachnid species which represents a pest. The term also relates to the insect or arachnid at any stage in its development. Because insects have a non-living exoskeleton, they cannot grow at a uniform rate and rather grow in stages by periodically shedding their  
10    exoskeleton. This process is referred to as moulting or ecdysis. The stages between moults are referred to as "instars" and this stage may be targeted according to the invention. Also, insect eggs or live young may also be targeted according to the present invention. All stages in the developmental cycle, which includes metamorphosis in the pterygotes, may be targeted by RNAi according to the present invention. Thus, individual  
15    stages such as larvae, pupae, nymph etc stages of development may all be targeted.

      The target species may be any insect or arachnid, meaning any organism or species belonging to the Kingdom Animals, more specifically to the Phylum Arthropoda, and to the Class Insecta or the Class Arachnida. The methods of the invention are applicable to all insects and arachnids that are susceptible to gene silencing by RNA  
20    interference and that are capable of internalising double-stranded RNA from their surrounding environment.

      In one embodiment of the invention, the insect or arachnid may belong to the following orders: Acari, Arachnida, Anoplura, Blattodea, Coleoptera, Collembola, Dermaptera, Dictyoptera, Diplura, Diptera, Embioptera, Ephemeroptera, Grylloblatodea,  
25    Hemiptera, Heteroptera, Homoptera, Hymenoptera, Isoptera, Lepidoptera, Mallophaga, Mecoptera, Neuroptera, Odonata, Orthoptera, Phasmida, Phthiraptera, Plecoptera, Protura, Psocoptera, Siphonaptera, Siphunculata, Thysanura, Sternorrhyncha, Strepsiptera, Thysanoptera, Trichoptera, Zoraptera and Zygentoma.

      In preferred, but non-limiting, embodiments of the invention the insect or arachnid  
30    is chosen from the group consisting of:

- (1) Acari: mites including Ixodida (ticks)
- (2) Arachnida: Araneae (spiders) and Opiliones (harvestman), examples include: *Latrodectus mactans* (black widow) and *Loxosceles reclusa* (Brown Recluse Spider)

- (3) Anoplura: lice, such as *Pediculus humanus* (human body louse)
- (4) Blattodea: cockroaches including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australasiae*), of the genus *Blatta*, including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella longipalpa*). A most preferred target is German cockroach (*Blatella germanica*).
- (5) Coleoptera: beetles, examples include: the family of Powderpost beetle (family of Bostrichoidea); *Dendroctonus* spp. (Black Turpentine Beetle, Southern Pine Beetle, IPS Engraver Beetle); Carpet Beetles (*Anthrenus* spp, *Attagenus* spp); Old House Borer (family of Cerambycidae: *Hylotrupes bajulus*); *Anobium punctatum*; *Tribolium* spp (flour beetle); *Trogoderma granarium* (Khapra Beetle); *Oryzaephilus sarinamensis* (Toothed Grain Beetle) etc. (Bookworm)
- (6) Dermaptera: family of earwigs
- (7) Diptera: mosquitoes (Culicidae) and flies (Brachycera), examples are: Anophelinae such as *Anopheles* spp. and Culicinae such as *Aedes fulvus*; Tabanidae such as *Tabanus punctifer* (Horse Fly), *Glossina morsitans morsitans* (tsetse fly), drain flies (Psychodidae) and Calypttratae such as *Musca domestica* (House fly), flesh flies (family of Sarcophagidae) etc.
- (8) Heteroptera: bugs, such as *Cimex lectularius* (bed bug)
- (9) Hymenoptera: wasps (Apocrita), including ants (Formicoidea), bees (Apoidea): *Solenopsis invicta* (Red Fire Ant), *Monomorium pharaonis* (Pharaoh Ant), *Camponotus* spp (Carpenter Ants), *Iasius niger* (Small Black Ant), *tetramorium caespitum* (Pavement Ant), *Myrmica rubra* (Red Ant), *Formica* spp (wood ants), *Crematogaster lineolata* (Acrobat Ant), *Iridomyrmex humilis* (Argentine Ant), *Pheidole* spp. (Big Headed Ants, *Dasymutilla occidentalis* (Velvet Ant) etc.
- (10) Isoptera: termites, examples include: *Amitermes floridensis* (Florida dark-winged subterranean termite), the eastern subterranean termite (*Reticulitermes flavipes*), the *R. hesperus* (Western Subterranean Termite), *Coptotermes formosanus* (Formosan Subterranean Termite), *Incisitermes minor* (Western Drywood Termite), *Neotermes connexus* (Forest Tree Termite) and Termitidae

- 11 -

(11) Lepidoptera: moths, examples include: Tineidae & Oecophoridae such as *Tineola bisselliella* (Common Clothes Moth), and Pyralidae such as *Pyralis farinalis* (Meal Moth) etc

(12) Psocoptera: booklice (Psocids)

5 (13) Siphonaptera: fleas such as *Pulex irritans*

(14) Sternorrhyncha: aphids (Aphididae)

(15) Zygentoma: silverfish, examples are: *Thermobia domestica* and *Lepisma saccharina*

Preferred target insects or arachnids include household insects, ecto-parasites  
 10 and insects and/or arachnids relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. More preferred targets are cockroaches (Blattodea) such as but not limited to *Blatella* spp. (e.g. *Blatella germanica* (german cockroach)),  
 15 *Periplaneta* spp. (e.g. *Periplaneta americana* (American cockroach) and *Periplaneta australiasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus*  
 20 spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex* spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g.  
 25 *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite)), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)),  
 30 *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)), *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)). More preferred targets are cockroaches. A most preferred target is German cockroach (*Blatella germanica*).

### RNA constructs

By "complementary" is meant that the RNA strand represents the RNA equivalent of the specified sequence if that sequence is a DNA sequence or the RNA equivalent of the complement of the DNA sequence.

5       The present invention relates to additional targets for RNAi mediated down regulation of gene expression. For all targets identified herein, there is provided in a further aspect of the invention an RNA construct comprising a double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence of any one of any one of (i) the  
10   target nucleic acid molecules defined in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180 and 181; or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200; or an  
15   orthologous nucleotide sequence thereof from an insect and/or arachnid species. As described above, the orthologue may share at least about 50% nucleotide sequence identity with the nucleotide sequence as set forth in any one of SEQ ID NO's 1, 4, 6, 7, 11, 16, 17, 21, 26, 27, 31, 36, 37, 41, 43, 44, 49, 51, 52 and 57. Preferably, the orthologue will share at least about 70%, 75%, 80%, 85%, 90%, more preferably at least  
20   about 95% and even more preferably at least about 96%, 97%, 98% or 99% sequence identity with the nucleotide sequence as set forth in any one of SEQ ID NO's 1, 4, 6, 7, 11, 16, 17, 21, 26, 27, 31, 36, 37, 41, 43, 44, 49, 51, 52 and 57.

As aforementioned, the orthologues of targets identified herein in *Blatella germanica* are considered to be viable targets in other insect and/or arachnid species,  
25   including household insects and/or arachnids, ecto-parasites and insects relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas.

Most preferred targets are derived from cockroaches, for example cockroaches of  
30   the genus *Blatella*, including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australasiae*), of the genus *Blatta*, including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella*

*longipalpa*). The most preferred target is German cockroach (*Blattella germanica*), in which the novel targets have been identified.

It has been previously reported that the formation of short interfering RNAs (siRNAs) of about 21 bp is desirable for effective gene silencing. However, in applications of applicant it has been shown that the minimum length of dsRNA preferably is at least about 80-100 bp in order to be efficiently taken up by certain pest organisms. There are indications that in invertebrates such as the free living nematode *C. elegans* or the plant parasitic nematode *Meloidogyne incognita* these longer fragments are more effective in gene silencing, possibly due to a more efficient uptake of these long dsRNA by the invertebrate.

It has also recently been suggested that synthetic RNA duplexes consisting of either 27-mer blunt or short hairpin (sh) RNAs with 29 bp stems and 2-nt 3' overhangs are more potent inducers of RNA interference than conventional 21-mer siRNAs (see Williams, Nature Biotechnology Vol 23, 2, Feb 2005, 181 and Kim et al, Nature Biotechnology Vol 23, 2, Feb 2005, 222-229 and Siolas et al, Nature Biotechnology Vol 23, 2, Feb 2005, 227-231 which references are incorporated herein in their entirety). Thus, molecules based upon the targets identified above and being either 27-mer blunt or short hairpin (sh) RNA's with 29-bp stems and 2-nt 3'overhangs are also included within the scope of the invention.

Therefore, in one embodiment, the RNA construct has a double stranded RNA region which has a length of at least about 17 bp, preferably at least about 21 bp, more preferably between about 20-1500 bp, even more preferably between about 80 - 1000 bp and most preferably between about 17-27 bp or between about 80-250 bp; such as double stranded RNA regions of about 17 bp, 18 bp, 20 bp, 21 bp, 22 bp, 23 bp, 24 bp, 25 bp, 26 bp, 27 bp, 50 bp, 80 bp, 100 bp, 150 bp, 200 bp, 250 bp, 300 bp, 350 bp, 400 bp, 450 bp, 500 bp, 550 bp, 600 bp, 650 bp, 700 bp, 900 bp, 100 bp, 1100 bp, 1200 bp, 1300 bp, 1400 bp or 1500 bp.

The upper limit on the length of the double-stranded RNA may be dependent on i) the requirement for the dsRNA to be taken up by the insect and/or arachnid and ii) the requirement for the dsRNA to be processed within the relevant cell into fragments that direct RNAi. The chosen length may also be influenced by the method of synthesis of the RNA and the mode of delivery of the RNA to the cell. Preferably the double-stranded RNA to be used in the methods of the invention will be less than 10,000 bp in length,

- 14 -

more preferably 1000 bp or less, more preferably 500 bp or less, more preferably 300 bp or less, more preferably 100 bp or less.

Efficacy in terms of pest control may be increased by targeting multiple target genes with a single RNA construct. Thus, the pest is less likely to survive and acquire  
5 resistance because there will be multiple double stranded RNA's mediating RNA interference, possibly all at the same time or possibly in a cascaded manner.

The methods of the invention encompass the simultaneous or sequential provision of two or more different double-stranded RNAs or RNA constructs to the same insect and/or arachnid, so as to achieve down-regulation or inhibition of multiple target  
10 genes or to achieve a more potent inhibition of a single target gene.

According to a further embodiment, the RNA constructs according to the invention comprise at least one double stranded RNA region, at least one of which comprises a nucleotide sequence that is complementary to a portion of any of the nucleotide sequences described herein, wherein the complementarity of said nucleotide  
15 sequence comprises at least 70%, preferably at least 75%, 80%, 85%, 90%, more preferably at least about 95% and even more preferably at least about 96%, 97%, 98% or 99% sequence identity with (i) the portion of the nucleotide sequence of the nucleic acid molecules as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180  
20 and 181 or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect and/or arachnid species, wherein the percentage sequence identity is calculated over the same length.

25 With "over the same length" is meant that when % identity is calculated between sequences, this is done over the corresponding stretch of nucleotides in both sequences.

Alternatively, multiple target genes are down regulated by the provision of one double-stranded RNA that acts against multiple target sequences. Alternatively, a single  
30 target may be more efficiently inhibited by the presence of more than one copy of the double stranded RNA fragment corresponding to the target gene. Thus, in one embodiment of the invention, the double-stranded RNA construct comprises multiple dsRNA regions, at least one strand of each dsRNA region comprising a nucleotide sequence that is complementary to at least part of a target nucleotide sequence of an

- 15 -

insect and/or arachnid target gene. According to the invention, the dsRNA regions in the RNA construct may be complementary to the same or to different target genes and/or the dsRNA regions may be complementary to target genes from the same or from different insect and/or arachnid species.

5 Accordingly, the invention provides an isolated double stranded RNA or RNA construct of the invention which comprises at least two double stranded RNA regions, at least one strand of each of which comprises, consists essentially of, or consists of a nucleotide sequence that is complementary to a portion of the nucleotide sequence of any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87,  
10 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence from an insect and/or arachnid species, wherein the orthologous nucleotide sequence has at least 70%, 80%, 85%, 87.5%, 90%, 95% or at least 99% sequence identity with at least the relevant portion of the nucleotide  
15 sequence of any one of SEQ ID NOs 1, 11, 21 and 31 or the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 49, 51, 52 and 57. Preferably said double stranded RNA or RNA construct comprises, consists essentially of or consists of one or at least two nucleotide sequences independently chosen from any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11,  
20 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, preferably any of SEQ ID NOs 65 to 70.

Thus, in one embodiment of the invention, the RNA construct comprises multiple dsRNA regions, at least one strand of each dsRNA region comprising a nucleotide  
25 sequence that is complementary to a portion of the nucleotide sequence of a target gene from the insect and/or arachnid species. According to the invention, the dsRNA regions in the RNA construct may be complementary to the same or to different target genes; the dsRNA regions may be complementary to targets from the same or from different insect species.

30 The dsRNA regions may be combined as follows:

- a) when multiple dsRNA regions targeting a single target gene are combined, they may be combined in the original order (ie the order in which the fragments appear in the target gene) in the RNA construct,



- 16 -

- b) alternatively, the original order of the fragments may be ignored so that they are scrambled and combined randomly or deliberately in any order into the RNA construct,
- c) alternatively, one single fragment may be repeated several times, for example from 1 to 10 times, e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 times in the RNA construct, or
- d) the dsRNA regions (targeting a single or different target genes) may be combined in the sense or antisense orientation.

In addition, the target gene(s) to be combined may be chosen from one or more of the following categories of genes (including all possible combinations thereof, as appropriate):

- e) "essential" genes or "pathogenicity genes" encompass genes that are vital for one or more target insects and/or arachnids and result in a lethal or severe (e.g. movement, feeding, paralysis, drinking, fertility, reproduction, growth) phenotype when silenced. The choice of a strong lethal target gene results in a potent RNAi effect. In the RNA constructs of the invention, multiple dsRNA regions targeting the same or different (very effective) lethal genes can be combined to further increase the potency, efficacy or speed of the RNAi effect in insect and/or arachnid control.
- f) "weak" genes encompass target genes with a particularly interesting function in one of the cellular pathways described herein, but which result in a weak phenotypic effect when silenced independently. In the RNA constructs of the invention, multiple dsRNA regions targeting a single or different weak gene(s) may be combined to obtain a stronger RNAi effect.
- g) "insect and/or arachnid specific" genes encompass genes and portions of genes that have no substantially homologous counterpart in non-pest organisms as can be determined by bioinformatics homology searches, for example by BLAST searches. The choice of an insect and/or arachnid specific target gene or portion thereof results in a species specific RNAi effect, with no effect or no substantial (adverse) effect in non-target organisms.
- h) "conserved genes" encompass genes that are conserved (at the amino acid level) between the target organism and non-target organism(s). To reduce possible effects on non-target species, such effective but conserved genes

- 17 -

are analysed and target sequences from the variable regions of these conserved genes are chosen to be targeted by the dsRNA regions in the RNA construct. Here, conservation is assessed at the level of the nucleic acid sequence. Such variable regions thus encompass the least conserved sections, at the level of the nucleic acid sequence, of the conserved target gene(s).

- i) "conserved pathway" genes encompass genes that are involved in the same biological pathway or cellular process, or encompass genes that have the same functionality in different insect and/or arachnid species resulting in a specific and potent RNAi effect and more efficient pest control;
- j) alternatively, the RNA constructs according to the present invention target multiple genes from different biological pathways, resulting in a broad cellular RNAi effect and more efficient insect and/or arachnid control.

Preferably, all double stranded RNA regions comprise at least one strand that is complementary to a portion of the nucleotide sequence of any one of SEQ ID NO 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200. However, provided one of the double stranded RNA regions comprises at least one strand that is complementary to a portion of the nucleotide sequence of any one of SEQ ID NO 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, the other double stranded RNA regions may comprise at least one strand that is complementary to a portion of any target gene from the insect and/or arachnid species (including known target genes).

The invention also provides any of the RNA constructs herein described, further comprising at least one additional functional sequence and optionally a linker. In one embodiment, the multiple dsRNA regions are connected by one or more linkers. In another embodiment, the linker is present at a site in the RNA construct, separating the dsRNA regions from another region of interest. Different linker types for the dsRNA constructs are provided by the present invention.

"Conditionally self-cleaving linkers" are RNA sequences capable of being processed under certain conditions. One example of suitable conditionally self-cleaving linkers are RNA sequences that are self-cleaving at low pH conditions. Suitable

examples of such RNA sequences are described by Jayasena and Gold (Proc Natl Acad Sci U S A. 1997 Sep 30; 94(20):10612-7), which document is incorporated herein by reference.

Another example of suitable conditionally self-cleaving linkers are RNA sequences that are self-cleaving at high pH conditions. Suitable examples of such RNA sequences are described by Borda et al. (Nucleic Acids Res. 2003 May 15;31(10):2595-600), which document is incorporated herein by reference. This sequence originates from the catalytic core of the hammerhead ribozyme HH16.

In one aspect of the invention, the linkers may be used to disconnect smaller dsRNA regions in the pest organism. Advantageously, in this situation the linker sequence may promote division of a long dsRNA into smaller dsRNA regions under particular circumstances, resulting in the release of separate dsRNA regions under these circumstances and leading to more efficient gene silencing by these smaller dsRNA regions.

In another aspect of the invention, a linker is located at a site in the RNA construct, separating the dsRNA regions from another sequence of interest, which preferably provides some additional function to the RNA construct. Non-limiting examples of other functional sequences (of interest) which may be incorporated in the RNA construct are for instance (i) additional sequences to facilitate large-scale production of the dsRNA construct; (ii) additional sequences to increase/decrease stability of dsRNA; (iii) additional sequences to bind to proteins or other molecules in a composition to facilitate uptake by the pest species; (iv) additional sequences that are aptamers and that bind to receptors or to molecules in the gut of the pest species to facilitate uptake, endocytosis and/or transcytosis by the pest species; (v) additional sequences to catalyze processing of dsRNA regions.

According to a particular embodiment the pest species has a gut system, such as for example insects and/or arachnids, and the linker is self-cleaving in the gut of the insect and/or arachnid. The pH in the gut is variable ranging from extremely acid to extremely basic.

Alternatively, the linkers are self-cleaving in the endosomes. This may be advantageous when the constructs of the present invention are taken up by the pest organisms via endocytosis or transcytosis, and are therefore compartmentalized in the endosomes of the pest species. The endosomes may have a low pH environment, leading to cleavage of the linker.

- 19 -

The above mentioned linkers that are self cleaving in hydrophobic conditions are particularly useful in dsRNA constructs of the present invention when used to be transferred from one cell to another via the transit in a cell wall, for example when  
5 crossing the cell wall of an insect and/or arachnid pest organism.

An intron may also be used as a linker. An "intron" as used herein may be any non-coding RNA sequence of a messenger RNA. Particular suitable intron sequences for the constructs of the present invention (1) are U-rich (35-45%); (2) have an average length of 100 bp (varying between about 50 and about 500 bp) which base pairs may be  
10 randomly chosen or may be based on known intron sequences; (3) start at the 5' end with -AG:GT- or -CG:GT- and/or (4) have at their 3' end -AG:GC- or -AG:AA.

A non-complementary RNA sequence, ranging from about 1 base pair to about 10,000 base pairs, may also be used as a linker.

As described above the dsRNA regions of the invention may correspond to only a  
15 portion of the target gene, provided that the complementarity is such that RNAi can occur to effectively control the insect and/or arachnid pest. It is not essential to the invention that the full length sequence of the pertinent target gene is known, as long as the dsRNA region containing construct used is capable of down-regulating the target gene.

For example, it is possible to use a dsRNA fragment based on a partial gene  
20 sequence (such as an EST) from the insect and/or arachnid, as long as said partial sequence is an ortholog of one of the sequences described as SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200. The degree of sequence  
25 homology or complementarity is determined over the length of the dsRNA fragment used.

Furthermore, it is also possible in the invention to use dsRNA fragments that differ from the nucleic acid molecules comprising the sequences described in SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30,  
30 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200 in one or more nucleotide positions (e.g. by a deletion, insertion or substitution), as long as the resulting dsRNA fragment is still capable of downregulating the target gene.

- 20 -

Preferably, the dsRNA fragment in the RNA construct has a complementarity, or level of homology comprising at least about 70% nucleotide sequence identity, preferably at least about 80% sequence identity, even more preferably at least about 85% or 87.5% sequence identity, still more preferably about 90% sequence identity, still more preferably at least about 95% sequence identity and most preferably at least about 96%, 97%, 98% or 99% sequence identity with the relevant portion of the nucleotide sequence of any one of the target nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence from an insect and/or arachnid species.

Methods for determining sequence identity are routine in the art and include use of the Blast software and GAP analysis (GCG program). High levels of sequence identity (complementarity) of at least one strand of the dsRNA with the target gene are required to mediate effective RNAi, and thus pest control.

However, it is equally advantageous that the dsRNA regions of the invention are selective to the pest target sequence versus the sequences of mammalian orthologues. This is especially relevant in the present invention where the pest must be controlled in an environment, such as a kitchen, where food is present and in which humans and other mammals may be exposed to compositions designed to control the pest. A selective biological agent is preferable to a chemical agent which may be equally toxic to a mammal, as it is to the pest species.

Furthermore, for a biological agent such as the RNA constructs of the present invention, there is the advantage that the molecules will biodegrade over time and thus will pose less of an environmental and health risk to human users than a chemical agent (such as the known insecticides).

Thus, according to a preferred embodiment, the at least one strand of the double stranded RNA in the RNA construct which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence of a nucleic acid molecule as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence from an insect and/or arachnid

- 21 -

species has less than about 5%, less than about 10%, less than about 12.5 %, less than about 15%, less than about 20%, less than about 30%, less than about 40% sequence identity with the corresponding (orthologous) nucleotide sequence from a mammalian species. In one embodiment, there is no sequence identity with mammalian sequences over 21 contiguous nucleotides. In another embodiment, there is less than about 10% or less than about 12.5 % sequence identity over 24 contiguous nucleotides with the corresponding nucleotide sequence from a mammalian species. Preferably, the mammalian species is a human.

In one embodiment, the at least one strand (of the double stranded RNA in the RNA construct which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence of a nucleic acid molecule as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200 or an orthologous nucleotide sequence from an insect and/or arachnid species) comprises at least 17 nucleotides, preferably at least 18 nucleotides, 19 nucleotides, 20 nucleotides, 21 nucleotides, 24 nucleotides, 27 nucleotides, 30 nucleotides, 40 nucleotides, 50 nucleotides, 60 nucleotides, 70 nucleotides, 80 nucleotides, 90 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides, 400 nucleotides, 450 nucleotides, 500 nucleotides, 550 nucleotides, 600 nucleotides, 650 nucleotides, 700 nucleotides, 900 nucleotides, 1000 nucleotides, 1100 nucleotides, 1200 nucleotides or 1300 nucleotides of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

An RNA construct is also provided comprising at least one double stranded RNA region, at least one strand of which comprises at least about 17 nucleotides, preferably at least about 18 nucleotides, 19 nucleotides, 20 nucleotides, 21 nucleotides, 23 nucleotides, 24 nucleotides, 25 nucleotides, 27 nucleotides, 30 nucleotides, 40 nucleotides, 50 nucleotides, 60 nucleotides, 70 nucleotides, 80 nucleotides, 90 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides, 400 nucleotides, 450 nucleotides, 500 nucleotides, 550 nucleotides, 600 nucleotides, 650 nucleotides, 700 nucleotides, 900 nucleotides, 1000

nucleotides, 1100 nucleotides, 1200 nucleotides or about 1300 nucleotides of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

#### DNA and expression constructs and host cells

In a further aspect, the invention also provides a DNA construct comprising the nucleotide sequence of the novel targets of the invention, as represented in SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200.

The invention further relates to a DNA construct comprising a region encoding an RNA construct of the invention.

The invention also provides, in a still further aspect, an expression construct comprising any of the DNA constructs of the invention.

The expression construct is such that it is capable, under suitable conditions, of providing (through transcription) an RNA construct comprising a dsRNA region as referred to above.

Genetic constructs for expressing dsRNA are well-known in the art: reference is for example made to the constructs described in WO 99/32619; in WO 00/01846 and WO 01/88121 (all Devgen); WO 00/44914 and WO 01/70949, as well as the prior art already mentioned above. As mentioned therein, such constructs may be DNA or RNA (and are preferably DNA) and may be in the form of a suitable expression vector (such as an expression vector suitable for the transformation of and for expression in bacteria) or other expression system. For example, the construct may be present in (for example by transformation) a suitable bacterial or viral system for the production in bacteria or for transformation of insects and/or arachnids, and these and other host cells containing the genetic constructs form a further aspect of the invention.

An expression construct according to the invention will usually contain – besides the sequence(s) encoding the dsRNA fragment itself – suitable regulatory elements (such as promoters, terminators and enhancers) and other elements of such genetic constructs known per se; and may for example express the dsRNA regions as two

separate complementary RNA strands that hybridize to form the desired dsRNA region or may express the dsRNA region in the form of a single RNA containing the two complementary strands, that self-hybridize to form a "stem-loop" or "hairpin" structure that contains the desired dsRNA region. All such constructs may be suitably used in the present invention, which is not particularly limited as to the type of construct used, as long as said construct is suitable for expression of a dsRNA which can mediate effective RNAi in an insect and/or arachnid pest.

The constructs themselves may also be constructed in a manner known *per se*, for which again reference is made to the above prior art references, as well as to the standard handbooks such as Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd.ed.), Vols. 1-3, Cold Spring Harbor Laboratory Press (1989) and F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987).

The dsRNA regions may be expressed preferably in a bacterial host under the control of a constitutive promoter or an inducible promoter (e.g. a promoter that is induced by a specific compound, by damage to the bacteria, etc.). The constitutive or inducible promoter may be non-specific or specific (for example, for a specific part of the life cycle of the bacteria).

The bacterial host cell may need to be inactivated before being utilised as a biological pesticide. This may be done by any technique known in the art, such as by heating or by treatment with phenol or formaldehyde for example. Alternatively, an inactivated virus, such as a suitably modified baculovirus may be utilised in order to deliver the dsRNA regions of the invention to the insect and/or arachnid pest.

The expression constructs may further contain all other elements known *per se* for nucleic acid sequences or genetic constructs, such as promoters or other control elements, terminators, translation or transcription enhancers, integration factors, signal sequences, selection markers, etc., that are preferably suited for use in a bacterial cell. The sequences that encode these further elements of the construct may again be either isolated from a suitable biological source, or provided synthetically.

Some specific, but non-limiting examples of suitable promoters include, but are not limited to, promoters from an RNA Poll, an RNA PolII, an RNA PolIII, T7 RNA polymerase, T3 RNA polymerase or SP6 RNA polymerase, and also the promoters and other regulatory elements referred to in the prior art cited above, such as in WO 00/01846 for example. The invention further provides bacterial promoters that can direct



expression of single stranded RNA, which can upon expression form a hairpin secondary structure containing a loop and a double stranded RNA region are utilised.

Specific, but non-limiting examples of transformation techniques for introducing the constructs into bacterial or viral hosts include transformation, electroporation, transfection etc.

The invention thus provides an expression construct comprising: (a) a nucleic acid encoding an RNA construct as described herein; (b) one or more control sequences capable of driving expression of the nucleic acid of (a); and optionally (c) a transcription termination sequence.

The expression constructs may be inserted into a plasmid or a vector, which may be commercially available. According to one embodiment of the present invention, the expression construct is a bacterial expression vector, suitable for transformation into bacteria and suitable for maintenance and expression of an RNA construct according to the present invention in a transformed bacterial cell. Reference is hereby made to the plasmids and vectors described in WO 01/01846 by applicant, which reference is incorporated herein in its entirety. An alternative is to use a virus cell which can infect an insect species, such as the viruses described in WO 01/34815, which reference is incorporated herein in its entirety.

The term "control sequence" as used herein is to be taken in a broad context and refers to regulatory nucleic acid sequences capable of driving and/or regulating expression of the sequences to which they are ligated and/or operably linked. According to one embodiment of the present invention, the control sequence is operable in a bacterium or virus; preferably the control sequence is derived from a bacterial sequence. The term "control sequence" encompasses a promoter or a sequence capable of activating or enhancing expression of a nucleic acid molecule in a cell, tissue or organ.

Optionally, one or more transcription termination sequences may also be incorporated in the expression construct. The term "transcription termination sequence" encompasses a control sequence at the end of a transcriptional unit, which signals 3' processing and poly-adenylation of a primary transcript and termination of transcription. Additional regulatory elements, such as transcriptional or translational enhancers, may be incorporated in the expression construct.

The expression constructs of the invention may further include an origin of replication which is required for maintenance and/or replication in a specific cell type.

One example is when a expression construct is required to be maintained in a bacterial cell as an episomal genetic element (e.g. plasmid or cosmid molecule) in a cell. Preferred origins of replication include, but are not limited to, f1-ori and colE1 ori.

The expression construct may optionally comprise a selectable marker gene. As  
5 used herein, the term "selectable marker gene" includes any gene, which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells, which are transfected or transformed, with an expression construct of the invention. Examples of suitable selectable markers include resistance genes against ampicillin (Amp<sup>r</sup>), tetracycline (Tc<sup>r</sup>), kanamycin (Kan<sup>r</sup>), phosphinothricin, and  
10 chloramphenicol (CAT) gene. Other suitable marker genes provide a metabolic trait, for example manA. Visual marker genes may also be used and include for example beta-glucuronidase (GUS), luciferase and Green Fluorescent Protein (GFP).

Thus, as described above, the invention provides a host cell comprising an RNA construct and/or a DNA construct and/or an expression construct of the invention.  
15 Preferably, the host cell is a bacterial cell, but may be a virus for example. A virus such as a baculovirus may be utilised which specifically infects insects and/or arachnids. This ensures safety for mammals, especially humans, since the virus will not infect the mammal, so no unwanted RNAi effect will occur.

The bacterial cell or virus preferably should be inactivated prior to use as a  
20 delivery agent for mediating RNAi in an insect pest when the agent is to be used in an environment where contact with humans or other mammals is likely (such as a kitchen as discussed above). Inactivation may be achieved by any means, such as by heat treatment, phenol or formaldehyde treatment for example.

A method for generating the RNA constructs of the invention is also provided.  
25 This method comprises the steps of

- a. contacting a DNA construct of the invention or an expression construct of the invention with cell-free components; or
- b. administering a DNA construct of the invention or an expression construct of the invention to a cell,

30 under conditions that allow transcription of said DNA construct to produce said RNA construct.

Thus, an *in vitro* method is provided, wherein the necessary components for transcription are provided. These components would be immediately familiar to one of skill in the art and numerous *in vitro* expression kits are commercially available.

Alternatively, the expression may be driven in a host cell. Preferably, the cell is a bacterial cell, but may be a virus for example.

Furthermore, in a further aspect of the invention, the host cells of the invention may be used as source for production of the dsRNA molecules and RNA constructs of the invention. For example, bacterial host cells, containing the expression construct of the invention (as hereinbefore described) may be cultured under suitable conditions (for example at 37°C or 42°C) in order to produce the RNA constructs of the invention in effective amounts. Large scale bacterial fermentation and harvesting processes are well known in the art and are utilised commercially. Bacterial culture may be carried out in any suitable media, such as for example LB broth, optionally supplemented with suitable antibiotics such as ampicillin, carbenicillin or chloramphenicol where an antibiotic resistant host strain is being utilised.

The resultant bacterial cultures thus produce the RNA constructs of the invention in large quantities. The bacteria themselves may be formulated into a suitable pesticide composition as described herein, or may be used as a direct (food) source of the RNA constructs of the invention for uptake, for example by ingestion, by a target insect or arachnid.

Similarly, in one embodiment, the bacteria may be used as a source of dsRNA by disrupting or otherwise inactivating the bacteria, as discussed above. For example, the cells may be ruptured or lysed using any suitable means, such as by osmotic shock for example, and the lysate or other suitable cellular fraction or extract of the bacteria utilised in the compositions of the invention.

In one embodiment, the bacterial extract or lysate may be suitably purified to leave a substantially pure RNA construct containing extract. Preferably, substantially all bacterial components are removed from the final dsRNA containing extracts, which may subsequently be formulated into any one of the compositions of the invention. Suitable purification steps are well known in the art and may include, by way of example and not limitation, suitable filtration steps, for example separation on the basis of charge or molecular weight. Suitable hybridization reactions may also be employed in order to purify the dsRNA molecules of interest.

The RNA constructs may be purified to substantial purity by standard techniques, including selective precipitation; column chromatography, immunopurification methods, and others (see, e.g., Scopes, Protein Purification: Principles and Practice (1982); Ausubel et al., supra; and Sambrook et al., supra).

The RNA and DNA constructs, including the double stranded RNA molecules, will typically incorporate natural bases. However, variants are included within the scope of the invention. Thus, the scope of "RNA" and "DNA" encompasses synthetic analogues, including well known sugar-modified bases, which are capable of base pairing and mediating RNAi or of being transcribed to produce RNA respectively in an analogous manner to natural nucleic acids. For example, nucleic acid analogues incorporating non-natural, chemically modified or derivatized bases, or nucleic acid analogues having a modified backbone are envisaged. This applies equally to the linkers which may be incorporated into the constructs of the invention. In particular, the term "double-stranded RNA" or "dsRNA" is to be interpreted as encompassing dsRNA containing non-natural bases. Double stranded RNA comprising non-natural or bases or having a chemically modified backbone may provide additional advantages regarding the increase or decrease of the stability of the dsRNA construct.

#### 15 Pesticide compositions

The invention relates, in a still further aspect to a pesticide composition comprising an RNA construct of the invention and/or a DNA construct of the invention and/or expression construct of the invention and/or host cell of the invention together with a suitable carrier, excipient or diluent.

20 According to a most preferred embodiment, the composition is in a form suitable for ingestion by an insect and/or arachnid.

The composition may be in any suitable physical form for application to insects and/or arachnids. The composition may be in solid form (such as a powder, pellet or a bait), liquid form (such as a spray) or gel form for example.

25 The composition may contain further components which serve to stabilise the dsRNA and/or prevent degradation of the dsRNA during prolonged storage of the composition.

The composition may still further contain components which enhance or promote uptake of the dsRNA by the intestinal or gut cell. These may include, for example, chemical agents which generally promote the uptake of RNA into cells e.g. lipofectamine etc.

30 It is contemplated that the "composition" of the invention may be supplied as a "kit-of-parts" comprising the double-stranded RNA in one container and a suitable diluent or carrier for the RNA containing entity (such as an RNA construct, DNA construct,

expression construct or host cell) in a separate container. The invention also relates to supply of the double-stranded RNA alone without any further components. In these embodiments the dsRNA may be supplied in a concentrated form, such as a concentrated aqueous solution. It may even be supplied in frozen form or in freeze-dried or lyophilised form. The latter may be more stable for long term storage and may be de-frosted and/or reconstituted with a suitable diluent immediately prior to use.

In one specific embodiment, the composition may be a coating, paste or powder that can be applied to a substrate in order to protect said substrate from infestation by insects and/or arachnids. In this embodiment, the composition can be used to protect any substrate or material that is susceptible to infestation by or damage caused by an insect and/or arachnid, for example foodstuffs and other perishable materials, and substrates such as wood. Preferred target insect and/or arachnid species for this embodiment include, but are not limited to the pests of the invention as defined earlier (see "Target organisms/species"), i.e. household insects and/or arachnids, ecto-parasites and insects relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. Most preferred target species are cockroaches, for example cockroaches of the genus *Blatella*, including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australasiae*), of the genus *Blatta*, including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella longipalpa*). The most preferred target is German cockroach (*Blatella germanica*).

In this embodiment the composition will comprise at least one double-stranded RNA containing entity (e.g. an RNA construct as described above), wherein the double-stranded RNA region comprises annealed complementary strands, at least one of which has a nucleotide sequence which corresponds to a target nucleotide sequence of a target gene from an insect and/or arachnid to be controlled and at least one carrier, diluent or excipient suitable for the intended use.

The nature of the excipients and the physical form of the composition may vary depending upon the nature of the substrate that it is desired to treat. For example, the composition may be a liquid that is brushed or sprayed onto or imprinted into the material or substrate to be treated, or a coating or powder that is applied to the material or

substrate to be treated. Thus, in one embodiment, the composition is in the form of a coating on a suitable surface which adheres to, and is eventually ingested by an insect and/or arachnid which comes into contact with the coating.

In one embodiment, the composition is in the form of a bait. The bait is designed to lure the insect and/or arachnid to come into contact with the composition. Upon coming into contact therewith, the composition is then internalised by the insect and/or arachnid, by ingestion for example and mediates RNAi to thus kill the insect and/or arachnid. Said bait may comprise a food substance, such as a protein based food, for example fish meal. Boric acid may also be used as a bait. The bait may depend on the species being targeted. For example, *Blatella germanica* will eat almost any food substance available to them. An attractant may also be used. The attractant may be a pheromone, such as a male or female pheromone for example. As an example, the pheromones referred to in the book "Insect Pheromones and their use in Pest Management" (Howse et al, Chapman and Hall, 1998) may be used in the invention. The attractant acts to lure the insect and/or arachnid to the bait, and may be targeted for a particular insect and/or arachnid or may attract a whole range of insects. The bait may be in any suitable form, such as a solid, paste, pellet or powdered form.

The bait may also be carried away by the insect and/or arachnid back to the colony. The bait may then act as a food source for other members of the colony, thus providing an effective control of a large number of insects and/or arachnids and potentially an entire insect and/or arachnid pest colony. This is an advantage associated with use of the double stranded RNA of the invention, because the delayed action of the RNAi mediated effects on the pests allows the bait to be carried back to the colony, thus delivering maximal impact in terms of exposure to the insects and/or arachnids.

Additionally, compositions which come into contact with the insects and/or arachnids may remain on the cuticle of the insect and/or arachnid. When cleaning, either an individual insect and/or arachnid cleaning itself or insects and/or arachnids cleaning one another, the compositions may be ingested and can thus mediate their effects in the insect and/or arachnid. This requires that the composition is sufficiently stable such that the dsRNA remains intact and capable of mediating RNAi even when exposed to external environmental conditions for a length of time, which may be a period of days for example.

The baits may be provided in a suitable "housing" or "trap". Such housings and traps are commercially available and existing traps may be adapted to include the compositions of the invention. Any housing or trap which may attract an insect and/or arachnid to enter it is included within the scope of the invention. The housing or trap may be box-shaped for example, and may be provided in pre-formed condition or may be formed of foldable cardboard for example. Suitable materials for a housing or trap include plastics and cardboard, particularly corrugated cardboard. Suitable dimensions for such a housing or trap are, for example, 7-15 cm wide, 15-20 cm long and 1-5 cm high. The inside surfaces of the traps may be lined with a sticky substance in order to restrict movement of the insect and/or arachnid once inside the trap. The housing or trap may contain a suitable trough inside which can hold the bait in place. A trap is distinguished from a housing because the insect can not readily leave a trap following entry, whereas a housing acts as a "feeding station" which provides the insect and/or arachnid with a preferred environment in which they can feed and feel safe from predators.

Accordingly, in a further aspect the invention provides a housing or trap for insects and/or arachnids which contains a composition of the invention, which may incorporate any of the features of the composition described herein.

In a further alternative embodiment, the composition may be provided in the form of a spray. Thus, a human user can spray the pest directly with the composition. The composition is then internalized by the insect and/or arachnid, from where it can mediate RNA interference, thus controlling the insect and/or arachnid. The spray is preferably a pressurized/aerosolized spray or a pump spray. The particles may be of suitable size such that they adhere to the insect and/or arachnid, for example to the exoskeleton, of the insect and/or arachnid and may be absorbed therefrom. Particle size may be measured by known means, such as by use of a Mastersizer, which is a commercially available device.

In a still further embodiment, the carrier is an electrostatically charged powder or particle which adheres to the insect and/or arachnid cuticle. Suitable powders and particles which are capable of adhering to an insect and/or arachnid and thus delivering the RNA constructs of the invention are described in detail in WO 94/00980 and WO 97/33472, both of which are incorporated herein by reference.

Alternatively, the carrier may comprise magnetic particles which adhere to the insect cuticle. Suitable magnetic particles which are capable of adhering to an insect

and/or arachnid and thus delivering the RNA constructs of the invention are described in detail in WO 00/01236, which reference is incorporated herein.

In a still further embodiment, which is preferred, the carrier of the composition comprises metallic particles which are initially unmagnetised but which are capable of becoming magnetically polarised when subjected to the electrical field provided by the insect and/or arachnid body. This mode of action is described in detail in WO 2004/049807 and is incorporated by reference herein.

These compositions which come into contact with the insects and/or arachnids may remain on the cuticle of the insect and/or arachnid. When cleaning, either an individual insect and/or arachnid cleaning itself or insects and/or arachnids cleaning one another, the compositions may be ingested and can thus mediate their effects in the insect and/or arachnid. This requires that the composition is sufficiently stable such that the dsRNA remains intact and capable of mediating RNAi even when exposed to external environmental conditions for a length of time, which may be a period of days for example.

Preferably, the composition incorporates a carrier which increases the uptake of the double stranded RNA into the insect and/or arachnid pest (see "target organisms/species" above), which is preferably an insect and/or arachnid and preferably a species of cockroach. Such a carrier may be a lipid-based carrier, preferably comprising one or more of, oil-in water emulsions, micelles, cholesterol, lipopolyamines and liposomes. Other agents which promote uptake of the constructs of the invention are well known to those of skill in the art and include polycations, dextrans and (tris) cationic lipids, such as CS096, CS102 etc. Commercially available liposomes include LIPOFECTIN® and CELLFECTIN® etc. A number of suitable carriers are listed under the heading "Transfection promoting agent" in WO 03/004644 and each of the examples provided is hereby incorporated by reference.

In a further preferred embodiment, the carrier is a nucleic acid condensing agent. Preferably, the nucleic acid condensing agent comprises spermidine or protamine sulphate or a derivative thereof.

The compositions of the invention may be combined together with further active ingredients, including with a further pesticide. Thus, the composition may be provided as a "kit-of-parts" comprising the double-stranded RNA containing composition in one container and one or more suitable pesticides, which may be a chemical or biological



pesticide, in a separate container. Alternatively, the compositions may be provided as a mixture which are stable and to be used in conjunction with one another.

Suitable active ingredients which may act in a complementary manner to the double stranded RNA molecules of the present invention include, but are not limited to the following: Chlorpyrifos, Allethrin, Resmethrin, Tetrabromoethyl, Dimethol-cyclopropane carboxylic acid (which are generally included in liquid compositions); and Hydranmethylnon, Avermectin, Chlorpyrifos, Sulfuramid, Hydroprene, Fipronil (GABA receptor), Isopropylphenyl methyl carbamate, Indoxacarb (PARA), Noviflumuron (Chitinsynthesis inhibitor), Imiprothrin (PARA), Abamectin (Glutamate-gated Chloride channel), Imidacloprid (Acetylcholin receptor) (which are generally included in bait compositions).

In a preferred embodiment, the active ingredient is known to be a preferred insecticide and/or arachnicide in terms of health and environmental considerations, such as for instance Hydranmethylnon and Avermectin.

According to another embodiment, the dsRNA is expressed in a suitable host cell such as a bacterial or fungal cell and the cell is taken up or eaten by the pest species. According to a further embodiment, the dsRNA is isolated from, or purified from, the cell which is preferably bacterial or fungal cell expressing the dsRNA, and the dsRNA is provided as a pesticide or in a pesticidal formulation to the pest species. Host cells, such as bacterial and fungal host cells may be engineered to produce any of the dsRNA or RNA constructs of the invention. These host cells, which are preferably bacterial cells may be ingested or otherwise internalized by the pest species. When taken up, the dsRNA can initiate an RNAi response, leading to the degradation of the target mRNA and weakening or killing of the pest.

Therefore, in a more specific embodiment, said double-stranded RNA or RNA construct is expressed by a prokaryotic, such as a bacterial, or eukaryotic, such as a yeast, host cell or host organism. These cells or organisms may be provided in any suitable formulation to facilitate uptake by the insect and/or arachnid.

### 30 Uses and Methods of the invention

In a still further aspect, the invention relates to the use of an RNA construct of the invention and/or a DNA construct of the invention and/or an expression construct of the invention and/or a composition of the invention and/or housing or trap of the invention for controlling an insect and/or arachnid by RNA interference. The use may

apply to a number of insects and/or arachnids at all stages of development, having orthologous target genes to the novel targets identified herein, including household insects and/or arachnids, ecto-parasites and insects and/or arachnids relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. Most preferred target species are cockroaches, for example cockroaches of the genus *Blatella*, including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australiasiae*), of the genus *Blatta*, including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella longipalpa*). The most preferred target is German cockroach (*Blatella germanica*).

Preferably, the pest is combatted via RNAi, and is consequently killed, paralysed, delayed in growth, inhibited in feeding of and/or hindered in its reproduction.

In a complementary aspect, the invention also provides a method of controlling insect and/or arachnid (pests) comprising administering to an insect and/or arachnid an RNA construct comprising a dsRNA region as defined above and/or a DNA construct of the invention and/or an expression construct as defined above and/or host cells as defined above and/or a composition as defined above and/or housing or trap as defined above, wherein the double stranded RNA is capable of down regulating the expression of at least one insect gene through RNA interference.

The administration may involve, for example feeding the insect and/or arachnid or may involve contacting the insect and/or arachnid with the dsRNA (in its various forms of presentation as described and defined above). Suitable means for direct contact include baits, sticky strips, magnetic and electrically charged powders and particles, sprays, gels, ointments, surface treatments etc as defined and described above with respect to the compositions of the invention. Any means of administration is included within the scope of the present invention provided it leads to effective double stranded RNA mediated interference of target gene expression, thus controlling the insect and/or arachnid.

It may be advantageous to provide multiple double stranded RNA region containing constructs directed against multiple targets, since this increases the efficacy of the insect and/or arachnid control and also decreases the possibility of the insect and/or arachnid acquiring resistance.

Accordingly, in one embodiment of the method, multiple RNA constructs as defined above and/or DNA constructs as defined above and/or expression constructs as defined above and/or host cells as defined above and/or compositions as defined above and/or housing or trap as defined above are provided/administered to the pest in order to  
5 mediate multiple separate RNAi events.

The multiple targets may all be targeted at the same time, or may be targeted in sequential manner. Thus, in one embodiment, the multiple RNA constructs and/or DNA constructs and/or expression constructs and/or host cells and/or compositions and/or housings or traps are provided/administered sequentially in order to reduce the  
10 probability of the insect and/or arachnid acquiring resistance.

The methods of the invention may apply to a number of insects and/or arachnids at all stages of development, having orthologous target genes to the novel targets identified herein. Target insects include household insects and/or arachnids, ectoparasites and insects and/or arachnids relevant for public health and hygiene such as, by  
15 way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. Most preferred target species are cockroaches, for example cockroaches of the genus *Blatella*, including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australasiae*), of the genus *Blatta*,  
20 including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella longipalpa*). The most preferred target is German cockroach (*Blatella germanica*).

Preferably, the insect and/or arachnid pest is combatted via RNAi, and is  
25 consequently killed, paralysed, delayed in growth, inhibited in feeding of and/or hindered in its reproduction.

The host cell may be, in one embodiment, a bacterial cell which has been engineered to produce the RNA constructs of the invention.

In a still further aspect, the invention provides a method for controlling cockroach  
30 pests comprising providing/administering to the cockroach an RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence encoding a cockroach ribosomal protein. Cockroach ribosomal proteins represent a novel target for RNAi, which can mediate effective control of a cockroach infestation.

- 35 -

Preferably, at least one strand of the at least one double stranded RNA region comprises at least about 17, 18, 19, 20, 21 nucleotides, preferably at least about 23 nucleotides, 24 nucleotides, 27 nucleotides, 30 nucleotides, 40 nucleotides, 50 nucleotides, 60 nucleotides, 70 nucleotides, 80 nucleotides, 90 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides, 400 nucleotides, 450 nucleotides, 500 nucleotides, 550 nucleotides, 600 nucleotides, 650 nucleotides, 700 nucleotides, 900 nucleotides, 1000 nucleotides, 1100 nucleotides, 1200 nucleotides or about 1300 nucleotides of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180 and 181, or the complement thereof.

In an even further aspect, the invention provides a method for controlling cockroach pests comprising providing/administering to the cockroach an RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence encoding a tropomyosin, a HMG Coenzyme A synthase gene or an Actin 5C gene. Cockroach tropomyosin, HMG Coenzyme A synthase and Actin 5C proteins represent a novel target for RNAi, which can mediate effective control of a cockroach infestation.

Preferably, at least one strand of the at least one double stranded region comprises at least about 17 nucleotides, preferably at least about 18 nucleotides, 19 nucleotides, 20 nucleotides, 21 nucleotides, 23 nucleotides, 24 nucleotides, 27 nucleotides, 30 nucleotides, 40 nucleotides, 50 nucleotides, 60 nucleotides, 70 nucleotides, 80 nucleotides, 90 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, 250 nucleotides, 300 nucleotides, 350 nucleotides, 400 nucleotides, 450 nucleotides, 500 nucleotides, 550 nucleotides, 600 nucleotides, 650 nucleotides, 700 nucleotides, 900 nucleotides, 1000 nucleotides, 1100 nucleotides, 1200 nucleotides or about 1300 nucleotides of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

The methods of the invention may apply to a number of insects and/or arachnids at all stages of development, having orthologous target genes to the novel targets identified herein. Target insects include household insects, ecto-parasites and insects

and/or arachnid relevant for public health and hygiene such as, by way of example and not limitation, flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas. More preferred target species are cockroaches (Blattodea) such as but not limited to *Blatella* spp. (e.g. *Blatella germanica* (german cockroach)), *Periplaneta* spp. (e.g. *Periplaneta americana* (American cockroach) and *Periplaneta australiasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus* spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex* spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g. *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)), *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)) and *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)), cockroaches, for example cockroaches of the genus *Blatella*, including German cockroach (*Blatella germanica*), of the genus *Periplaneta*, including American cockroach (*Periplaneta americana*) and Australian cockroach (*Periplaneta australiasiae*), of the genus *Blatta*, including Oriental cockroach (*Blatta orientalis*) and of the genus *Supella*, including brown-banded cockroach (*Supella longipalpa*). More preferred targets are cockroaches. The most preferred target is German cockroach (*Blatella germanica*).

Preferably, the insect and/or arachnid pest is combatted via RNAi, and is consequently killed, paralysed, delayed in growth, inhibited in feeding of and/or hindered in its reproduction.

### Kits of the invention

The invention also provides kits for use in the methods of the invention. These kits may incorporate the RNA constructs and/or DNA constructs and/or expression constructs and/or host cells and/or compositions and/or housings or traps of the invention, all of which deliver dsRNA regions to effect RNAi against specific target genes.

Preferably, the kits will also include instructions for use of the components of the kit. The double stranded RNAs found in the kits of the invention, or produced by components of the kits of the invention are capable of down regulating the expression of at least one insect (pest) gene through RNA interference.

Preferably, in order to provide more effective pest control (as described above), the kit comprises multiple components, each of which mediates RNAi at a different target gene or insect and/or arachnid species. Thus the kit may comprise multiple RNA constructs and/or DNA constructs and/or expression constructs and/or compositions, wherein each double stranded RNA is capable of down regulating the expression of at least one insect and/or arachnid (pest) gene through RNA interference.

Preferably, the components of the kit are applied sequentially to mediate effective pest control. However, some or all of the components may be administered simultaneously if required for maximal impact.

The kit may additionally comprise known pesticides, which may be provided together or separately from the components forming part of the invention.

Suitable active ingredients which may act in a complementary manner to the double stranded RNA molecules of the present invention include, but are not limited to the following: Chlorpyrifos, Allethrin, Resmethrin, Tetrabromoethyl, Dimethol-cyclopropane carboxylic acid (which are generally included in liquid compositions); and Hydramethylnon, Avermectin, Chlorpyrifos, Sulfuramid, Hydroprene, Fipronil (GABA receptor), Isopropylphenyl methyl carbamate, Indoxacarb (PARA), Noviflumuron (Chitinsynthesis inhibitor), Imiprothrin (PARA), Abamectin (Glutamate-gated Chloride channel), Imidacloprid (Acetylcholin receptor) (which are generally included in bait compositions).

In a preferred embodiment, the active ingredient is known to be a "preferred" insecticide and/or arachnicide with respect to health and environmental considerations, such as for instance Hydramethylnon and avermectin.

The kits of the invention may thus also be directed against multiple species at the same time in order to give a broad-scale pest control option. Double stranded RNA molecules may be included in the kits (as part of the appropriate constructs etc.) to mediate RNAi of multiple targets, including inter-species orthologues of the same targets for example.

The kits may include suitable buffers and packaging etc to ensure stability and storage of the components therein.

#### Technical advantages of the invention

There are numerous major advantages associated with the present invention over the use of conventional chemical insecticides.

(1) The RNAi mediating dsRNA has to match the target with a high degree of nucleotide sequence identity in order to effectively down regulate expression and thus control the pest. Thus, specificity can be achieved by designing double stranded RNA molecules in which one strand has high homology to the target sequence but which strand has only low homology to the orthologous sequence in a mammalian species, such as a human. This specificity is greater than can be achieved with conventional chemical pesticides.

(2) A new set of targets has been identified which can be used in the control of pests. Because these targets have previously not been identified, there should be no acquired resistance in the pest species.

(3) The double stranded RNA used in RNAi against the novel targets is a biodegradable product as compared to the known chemically synthesised pesticides, such as DMSO etc. The biodegradable nature of the constructs makes them more environmentally sound.

(4) RNAi does not necessarily provide an immediate effect in terms of killing the pest, rather the effects are mediated effectively but require time for the double stranded RNA to be associated with its target. The RNAi effect may result in killing the pest at a later moment and not directly upon contact, such as Noviflumuron (which is a chitinsynthesis inhibitor, from Dow AgroSciences). Thus, the use of RNAi may allow more facile control of large infestations of pests such as insects and/or arachnids because there is less chance of a shock effect being propagated amongst the pests where they may encounter a large number of dead pests in the vicinity of the insecticide and/or arachnicide.

(5) The use of multiple targets at the same time may provide more efficacious control of pest populations and reduce the possibility of acquired resistance. The targets may be common to a number of pest species providing broad scale treatment.

(6) In contrast to conventional pesticides, no professional assistance would be required in order to treat the relevant areas, due to the more safe nature of the DNA and RNA constructs, compositions and host cells of the invention.

(7) Minimum disruption of human activity would be required since the double stranded RNA region containing constructs are designed such that they will have no adverse effects or only minor effects on gene expression outside of the target pest population.

The invention will be further understood with reference to the following experimental section:

#### **Description of Tables and Figures**

**Table 1:** Examples of novel identified insect target genes. Gene function assigned is based on the FlyBase orthologue.

**Table 2:** dsRNA fragments complementary to *Blatella germanica* target sequences

**Table 3:** Effect of dsRNA treatments on the number of cockroaches successfully moulting to the adult stage, as a percentage of live insects (means  $\pm$  standard errors, n = 4)

**Table 4:** Selected sequences\* of target genes. Fragments of at least 17 bp of the sequences\* are present in the specified orthologous sequences in insect species (represented by GI number).

**Table 5:** Selected sequences\* of target genes. Fragments of at least 17 bp of the sequences\* are present in the specified orthologous sequences in arachnid species (represented by GI number).

**Figure 1:** Mortality of *B. germanica* on artificial pellet diet. The concentration of dsRNA in the pellets was 1% w/w. The concentration of imidacloprid was 1% w/w.

**Figure 2:** Mortality of *B. germanica* on artificial pellet diet. The concentration of dsRNA (Bg001, having the sequence as represented as SEQ ID NO 9, and Bg001 concatemer 2, having the sequence as represented as SEQ ID NO 68) in the pellets was



1% w/w. In this experiment, hexaflumuron (1% w/w) was tested as a positive control and solvent as a negative control.

**Figure 3** Stability of Bg032 dsRNA in LB medium (LB) and Rnase free water (MQ) at room temperature over a period of eight months

5 **Figure 4:** Effect on cockroach mortality upon applying dsRNA (Bg001) to first instars nymphs during one week. In this experiment, miscellaneous dsRNA and solvent were tested as negative controls. The concentration of dsRNA in the pellets was 1% w/w.

**Figure 5: Sequences of the invention**

10

**Table 1**

Target ID	Dm identifier	SEQ ID NO NA	SEQ ID NO AA	Function (based on FlyBase): <a href="http://flybase.org/">http://flybase.org/</a>
Bg001	CG11276	1	2	Ribosomal protein S4 (RpS4), structural constituent of ribosome involved in protein biosynthesis which is a component of the cytosolic small ribosomal subunit
Bg003	CG3395	11	12	Ribosomal protein S9 (RpS9), structural constituent of ribosome involved in protein biosynthesis which is a component of the cytosolic small ribosomal subunit
Bg004	CG6141	21	23	Ribosomal protein L9, structural constituent of ribosome involved in protein biosynthesis which is localised to the ribosome
Bg005	CG2746	31	32	Ribosomal protein L19, structural constituent of ribosome involved in protein biosynthesis which is localised to the ribosome
Bg031	CG4898	41	42	Tropomyosin 1 (AF260897), member of the tropomyosins family which are closely related proteins with multiple functions, including the regulation of the actin-myosin interaction, transport of mRNA, and mechanical support of the cytoplasmic membrane)
Bg032	CG16796	49	50	HMG Coenzyme A synthase (X73679) catalyzes a committed step in the pathways for isoprenoid, cholesterol, and ketone body production
Bg033	CG4027	57	58	Actin 5C (AY004248) is the major gene in <i>Drosophila melanogaster</i> that encodes the cytoskeletal actin present in all cell types in all growth stages

**Table 2:**

<b>Gene</b>	<b>dsRNA fragment size (bp)</b>	<b>freefrag</b>	<b>freefrag size (position on dsRNA)</b>
Bg001	594 (SEQ ID NO 9)	best1_human_24_3	69 (19-87)
Bg001	594 (SEQ ID NO 9)	best2_human_24_3	69 (445-513)
Bg001	594 (SEQ ID NO 9)	best3_human_24_3	62 (206-267)
Bg001	594 (SEQ ID NO 9)	best1_human_21_0	573 (1-573)
Bg003	433 (SEQ ID NO 19)	best1_human_24_3	133 (141-273)
Bg003	433 (SEQ ID NO 19)	best2_human_24_3	72 (68-139)
Bg003	433 (SEQ ID NO 19)	best3_human_24_3	65 (1-65)
Bg003	433 (SEQ ID NO 19)	best1_human_21_0	412 (1-412)
Bg004	449 (SEQ ID NO 29)	best1_human_24_3	78 (276-353)
Bg004	449 (SEQ ID NO 29)	best2_human_24_3	61 (200-260)
Bg004	449 (SEQ ID NO 29)	best3_human_24_3	53 (91-143)
Bg004	449 (SEQ ID NO 29)	best1_human_21_0	428 (1-428)
Bg005	404 (SEQ ID NO 39)	best1_human_24_3	115 (40-154)
Bg005	404 (SEQ ID NO 39)	best2_human_24_3	45 (191-235)
Bg005	404 (SEQ ID NO 39)	best3_human_24_3	42 (237-278)
Bg005	404 (SEQ ID NO 39)	best1_human_21_0	383 (1-383)
Bg031	849 (SEQ ID NO 47)	best1_human_24_3	70 (756-825)
Bg031	849 (SEQ ID NO 47)	best2_human_24_3	56 (546-601)
Bg031	849 (SEQ ID NO 47)	best3_human_24_3	54 (280-333)
Bg031	849 (SEQ ID NO 47)	best1_human_21_0	821 (8-828)
Bg031	849 (SEQ ID NO 47)	best2_human_21_0	6 (1-6)
Bg032	1300 (SEQ ID NO 55)	best1_human_24_3	126 (1138-1263)
Bg032	1300 (SEQ ID NO 55)	best2_human_24_3	114 (731-844)
Bg032	1300 (SEQ ID NO 55)	best3_human_24_3	99 (259-357)
Bg032	1300 (SEQ ID NO 55)	best1_human_21_0	1279 (1-1279)
Bg033	446 (SEQ ID NO 63)	best1_human_24_3	4 (362-365)
Bg033	446 (SEQ ID NO 63)	best2_human_24_3	4 (367-370)
Bg033	446 (SEQ ID NO 63)	best3_human_24_3	3 (115-117)
Bg033	446 (SEQ ID NO 63)	best1_human_21_0	108 (88-195)
Bg033	446 (SEQ ID NO 63)	best2_human_21_0	102 (244-345)
Bg033	446 (SEQ ID NO 63)	best3_human_21_0	62 (350-411)

**Table 3**

Day	Bg001	Bg001 concatemer 2	Positive control	Negative control
38	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	2.5 ( $\pm 2.5$ )
41	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	9.6 ( $\pm 6.7$ )
45	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	64.1 ( $\pm 14.0$ )
48	33.3 ( $\pm 23.6$ )	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	77.1 ( $\pm 10.4$ )
52	41.7 ( $\pm 25.0$ )	0.0 ( $\pm 0.0$ )	0.0 ( $\pm 0.0$ )	100.0 ( $\pm 0.0$ )

Table 4

Target ID	SEQ ID NO	Example GI-number and species	Sequence*
Bg001	71	2871551 ( <i>Drosophila melanogaster</i> )	aaggcatggatgttgacaagct
Bg001	72	48927129 ( <i>Hydropsyche</i> sp.)	gcatggaigtggacaagctcgg
Bg001	73	60293875 ( <i>Homalodisca coagulata</i> ); 71547743 ( <i>Oncometopia nigricans</i> )	attaaggtgatggaaaaagtcagaac
Bg001	74	56153292 ( <i>Rhynchosciara americana</i> )	cccaactatccagctggtttatggatgtgt
Bg001	75	90820001 ( <i>Graphocephala atropunctata</i> )	gctggtttatggatgtgttacaattgaaaa
Bg001	76	25956479 ( <i>Biphyllus lunatus</i> )	atgaaaaaaactggagaatttttcog
Bg001	77	15353483 ( <i>Apis mellifera</i> )	ggtaatctctgtatgattactgg
Bg001	78	92041090 ( <i>Drosophila willistoni</i> )	cgfcatccctgggttcctttgacattgt
Bg001	79	62083410 ( <i>Lysiphlebus testaceipes</i> )	ttaaagattcacaaaggacacac
Bg003	80	76169686 ( <i>Diploptera punctata</i> )	aaaatccgtaaagctgccagagaact
Bg003	81	62083482 ( <i>Lysiphlebus testaceipes</i> )	cgtaaagctgccagagaactct
Bg003	82	2459311 ( <i>Antheraea yamamai</i> )	aggtgtttgaaggcaatgctctt
Bg003	83	22040140 ( <i>Ctenocephalides felis</i> )	cgtatggaggtgtggatgaa
Bg003	84	83664146 ( <i>Myzus persicae</i> )	ccgtaigaagcttgattacgt
Bg003	85	55909980 ( <i>Locusta migratoria</i> ); 76169686 ( <i>Diploptera punctata</i> ); 15358510 ( <i>Apis mellifera</i> ); 67890783 ( <i>Drosophila pseudoobscura</i> )	ttgggtttgaagattgaagatttcttga
Bg003	86	62240069 ( <i>Diabrotica virgifera</i> )	aagattgaagatttcttgaa
Bg003	87	57963755 ( <i>Heliconius melpomene</i> ); 83663084 ( <i>Myzus persicae</i> )	aggaacaaacgtgaagtgtggcg
Bg004	88	70909652 ( <i>Cicindela litorea</i> )	tgciccatattgagaacatg
Bg004	89	83660638 ( <i>Myzus persicae</i> )	aagggtttcctgtacaaaaatg
Bg004	90	83931139 ( <i>Lutzomyia longipalpis</i> )	gccgtgatgcccatctcccat
Bg004	91	67895088 ( <i>Drosophila pseudoobscura</i> ); 92218607 ( <i>Drosophila willistoni</i> )	tatgccattccccattaactgcgt
Bg004	92	92960248 ( <i>Drosophila ananassae</i> ); 15455304 ( <i>Drosophila melanogaster</i> ); 38047668 ( <i>Drosophila yakuba</i> )	cgtaacttctgggcgagaagt
Bg004	93	56199511 ( <i>Culicoides sonorensis</i> ); 67876239 ( <i>Drosophila</i> )	aaatggtttggaacaaagaaggag

		pseudoobscura)		
Bg005	94	92931824 ( <i>Drosophila virilis</i> )	gatccaatgaaataaacgaaat	
Bg005	95	55883492 ( <i>Locusta migratoria</i> )	aatgaaataaacgaaattgcaaatac	
Bg005	96	60296437 ( <i>Homalodisca coagulata</i> )	ggtttggcaaaaagggaagggtac	
Bg005	97	78231035 ( <i>Heliconius erato/himera</i> mixed EST library)	gcaaatgccggtatgccacagaa	
Bg005	98	76553206 ( <i>Spodoptera frugiperda</i> ); 33491424 ( <i>Trichoplusia ni</i> )	aatgcccgatgccacagaagg	
Bg005	99	55900360 ( <i>Locusta migratoria</i> )	aagaagtacaggggaagcaagaa	
Bg005	100	57963592 ( <i>Heliconius melpomene</i> )	aagaagatcgacagacatcata	
Bg005	101	92948400 ( <i>Drosophila ananassae</i> ); 2871894 ( <i>Drosophila melanogaster</i> ); 68267374 ( <i>Drosophila simulans</i> ); 33354497 ( <i>Drosophila yakuba</i> ); 83935652 ( <i>Lutzomyia longipalpis</i> ); 18866169 ( <i>Anopheles gambiae</i> ); 60307025 ( <i>Sphaerius</i> sp.); 25958948 ( <i>Curculio glandium</i> ); 90812513 ( <i>Nasonia giraulti</i> )	caagggtaacgtgttcaagaacaacagcg	
Bg005	102	18909153 ( <i>Anopheles gambiae</i> ); 60311920 ( <i>Euclidia glyphica</i> ); 25957531 ( <i>Cicindela campestris</i> ); 18948649 ( <i>Anopheles gambiae</i> ); 38048300 ( <i>Drosophila yakuba</i> ); 58385089 ( <i>Anopheles gambiae</i> str. PEST); 27556513 ( <i>Anopheles gambiae</i> ); 70909752 ( <i>Cicindela campestris</i> ); 56462221 ( <i>Lonomia obliqua</i> ); 92931824 ( <i>Drosophila virilis</i> )	aagggtaacgtgttcaagaacaacagcgtgtcct	
Bg005	103	25957246 ( <i>Carabus granulatus</i> ); 90135865 ( <i>Bicyclus anynana</i> )	gtgttcaagaacaacggtgtccigtatggagt	
Bg005	104	71538996 ( <i>Diaphorina citri</i> ); 90812513 ( <i>Nasonia giraulti</i> ); 60311920 ( <i>Euclidia glyphica</i> )	tgatggagttcatccacaagaagaaggcgtg	
Bg005	105	15511486 ( <i>Drosophila melanogaster</i> )	catccacaagaagaaggcgtgagaag	
Bg005	106	60311920 ( <i>Euclidia glyphica</i> )	acaagaagaaggcgtgagaaggc	
Bg005	107	82572137 ( <i>Acyrtosiphon pisum</i> ); 73616334 ( <i>Aphis gossypii</i> ); 37804858 ( <i>Rhopalosiphum padi</i> ); 31365253 ( <i>Toxoptera citricida</i> ); 84647391 ( <i>Myzus persicae</i> )	accaattccagacaaaataattcgttaa	
Bg005	108	55908261 ( <i>Locusta migratoria</i> ); 10764114 ( <i>Manduca sexta</i> ); 90135865 ( <i>Bicyclus anynana</i> ); 91845469 ( <i>Bombix mori</i> )	gaagaaggcgtgagaaggccaggaca	
Bg031	109	84252313 ( <i>Aedes aegypti</i> ); 78052352 ( <i>Heliconius erato</i> ); 50818693 ( <i>Heliconius melpomene</i> ); 92942003 ( <i>Drosophila ananassae</i> );	atggatgccatcaagaagaagatgcaggcgtgaagct ggagaaggacaaacgcg	

		92466045 ( <i>Drosophila erecta</i> ); 92998051 ( <i>Drosophila grimshawi</i> ); 3627588 ( <i>Drosophila melanogaster</i> ); 92985296 ( <i>Drosophila mojavensis</i> ); 92921049 ( <i>Drosophila virilis</i> ); 92230306 ( <i>Drosophila willistoni</i> ); 92983068 ( <i>Drosophila mojavensis</i> ); 60294371 ( <i>Homalodisca coagulata</i> ); 73614014 ( <i>Aphis gossypii</i> ); 90819969 ( <i>Graphocephala atropunctata</i> ); 55886387 ( <i>Locusta migratoria</i> ); 85854848 ( <i>Aedes aegypti</i> ); 19310970 ( <i>Periplaneta fuliginosa</i> ); 20387026 ( <i>Lepisma saccharina</i> ); 27621313 ( <i>Anopheles gambiae</i> ); 91838618 ( <i>Bombyx mori</i> ); 20387028 ( <i>Lepisma saccharina</i> ); 4378572 ( <i>Periplaneta americana</i> ); 71050465 ( <i>Oncometopia nigricans</i> ); 18916954 ( <i>Anopheles gambiae</i> ); 29557544 ( <i>Bombyx mori</i> ); 55911583 ( <i>Locusta migratoria</i> ); 90978993 ( <i>Aedes aegypti</i> ); 56462261 ( <i>Lonomia obliqua</i> ); 85850284 ( <i>Aedes aegypti</i> ); 78230930 ( <i>Heliconius erato/himera</i> mixed EST library); 55895968 ( <i>Locusta migratoria</i> ); 29557242 ( <i>Bombyx mori</i> ); 18926345 ( <i>Anopheles gambiae</i> ); 37663025 ( <i>Bombyx mori</i> ); 18940590 ( <i>Anopheles gambiae</i> ); 81521031 ( <i>Lutzomyia longipalpis</i> ); 55804534 ( <i>Acyrtosiphon pisum</i> ); 18898107 ( <i>Anopheles gambiae</i> ); 29557268 ( <i>Bombyx mori</i> ); 84647487 ( <i>Myzus persicae</i> ); 37664569 ( <i>Bombyx mori</i> ); 81521022 ( <i>Lutzomyia longipalpis</i> ); 70978108 ( <i>Aedes aegypti</i> )	
Bg031	110	4378572 ( <i>Periplaneta americana</i> ); 19310970 ( <i>Periplaneta fuliginosa</i> )	gggccgagaaggctgaggaggaggc
Bg031	111	4378572 ( <i>Periplaneta americana</i> ); 19310970 ( <i>Periplaneta fuliginosa</i> )	tcctgcagaagaagatccagcagattgagaatgatct
Bg031	112	50557705 ( <i>Homalodisca coagulata</i> ); 71050465 ( <i>Oncometopia nigricans</i> )	tgatgcaagtcaacgccaagct
Bg031	113	78056651 ( <i>Heliconius erato</i> ); 50818693 ( <i>Heliconius melpomene</i> )	atgcaagtcaacgccaagctgga
Bg031	114	4378572 ( <i>Periplaneta americana</i> ); 19310970 ( <i>Periplaneta fuliginosa</i> )	gtcaacgccaagctggacgagaaggacaaggccct
Bg031	115	71050465 ( <i>Oncometopia nigricans</i> )	gagaaggacaaggccctgcagaa
Bg031	116	55907164 ( <i>Locusta migratoria</i> ); 55917622 ( <i>Locusta migratoria</i> )	aaccgcgaatccaactgctggagga
Bg031	117	86462380 ( <i>Acyrtosiphon pisum</i> ); 73618346 ( <i>Aphis gossypii</i> ); 53883526 ( <i>Plutella xylostella</i> ); 25958075 ( <i>Platystomos albinus</i> ); 85854848 ( <i>Aedes aegypti</i> ); 40384866 ( <i>Nilaparvata lugens</i> ); 56085268 ( <i>Bombyx mori</i> ); 78050451 ( <i>Heliconius erato</i> ); 71535946 ( <i>Diaphorina</i> )	gogatgaagctggagaaggacaacgcgatggatcgcg

Bg031	118	citri); 20387028 (Lepisma saccharina); 4378572 (Periplaneta americana); 19310970 (Periplaneta fuliginosa); 66500379 (Apis mellifera); 45753874 (Apis mellifera); 66522385 (Apis mellifera)	
Bg031	119	34788042 (Callosobruchus maculatus)	tcctgaggaaacgtttggccacagc
Bg031	120	20387028 (Lepisma saccharina)	tggcagatgaagagcgtatgga
Bg031	121	90972767 (Aedes aegypti); 56150925 (Rhynchosciara americana); 85854848 (Aedes aegypti)	gatgaagagcgtatggatgct
Bg031	122	60296314 (Homalodisca coagulata); 71050465 (Oncometopia nigricans)	gccttgagaaccagcctaagga
Bg031	123	85850407 (Aedes aegypti)	gagaaccagctgaaggaagcc
Bg031	124	29555905 (Bombyx mori)	cagctgaaggaagccaggttc
Bg031	125	85847532 (Aedes aegypti); 77850398 (Aedes aegypti); 3627588 (Drosophila melanogaster); 56150925 (Rhynchosciara americana); 77792932 (Aedes aegypti)	ttcatgctgagggaagctgacaagaata
Bg031	126	78540242 (Glossina morsitans); 6901854 (Bombyx mori)	gctgacaagaaatatgatgaggt
Bg031	127	40384866 (Nilaparvata lugens)	gacaagaaatatgatgaggtgc
Bg031	128	84647487 (Myzus persicae)	atggttgaggccgacttggaagagcaga
Bg031	129	51979105 (Myzus persicae)	gccgacttggaagagagcagaaga
Bg031	130	55886192 (Locusta migratoria)	cgacttggaagagcagaagagcgtgc
Bg031	131	92957972 (Drosophila ananassae)	ccaagattgtggagcttgagga
Bg031	132	60312749 (Gryllus bimaculatus)	aagattgtggagcttgaggaaga
Bg031	133	70978108 (Aedes aegypti)	tggatcgcccttctctgcgaacagcagcccg
Bg031	134	67842690 (Drosophila pseudoobscura)	attgtggagcttgaggagaactgcgct
Bg031	135	92939324 (Drosophila virilis)	ctgcgcttgcggcaacaac
Bg031	136	53883608 (Plutella xylostella)	cggtgtgcggcaacaacctgaagtccttgaggt
Bg031	137	4378572 (Periplaneta americana); 19310970 (Periplaneta fuliginosa); 33354924 (Drosophila yakuba); 25957752 (Cicindela campestris); 60312749 (Gryllus bimaculatus); 55907164 (Locusta migratoria); 75726914 (Tribolium castaneum)	gtgtcggcaacaacctgaagtccttgaggtga gaaggccaacctgcgctga
Bg031	138	19310970 (Periplaneta fuliginosa)	taccaggctaaggaggctga
Bg031	139	55923520 (Locusta migratoria); 20387028 (Lepisma saccharina);	accaggctaaggaggctgaagc

Bg031	55922834 ( <i>Locusta migratoria</i> )	
Bg031	139 25958290 ( <i>Platystomos albinus</i> )	gctaaaggaggctgaagctcg
Bg031	140 45757348 ( <i>Apis mellifera</i> ); 77783094 ( <i>Aedes aegypti</i> ); 25956952 ( <i>Biphylus lunatus</i> ); 25957752 ( <i>Cicindela campestris</i> ); 90972767 ( <i>Aedes aegypti</i> ); 75722624 ( <i>Tribolium castaneum</i> ); 47519043 ( <i>Acyrtosiphon pisum</i> ); 73612504 ( <i>Aphis gossypii</i> ); 83664605 ( <i>Myzus persicae</i> ); 9055470 ( <i>Pyrocoelia rufa</i> ); 30030953 ( <i>Toxoptera citricida</i> ); 77758700 ( <i>Aedes aegypti</i> ); 33365552 ( <i>Glossina morsitans</i> ); 56154884 ( <i>Rhynchosciara americana</i> ); 78540242 ( <i>Glossina morsitans</i> )	ctaaaggaggctgaagctcgctgagatt
Bg031	141 52630932 ( <i>Toxoptera citricida</i> ); 71047185 ( <i>Oncometopia nigricans</i> )	aaggaggctgaagctcgctgagctgctga
Bg031	142 92943056 ( <i>Drosophila ananassae</i> ); 92460361 ( <i>Drosophila erecta</i> ); 49400641 ( <i>Drosophila melanogaster</i> )	gctcgctgagctgctgaa
Bg031	143 19310970 ( <i>Periplaneta fuliginosa</i> ); 60311491 ( <i>Euclidia glyphica</i> ); 60312896 ( <i>Gryllus bimaculatus</i> ); 25958290 ( <i>Platystomos albinus</i> ); 60311415 ( <i>Euclidia glyphica</i> ); 55886380 ( <i>Locusta migratoria</i> ); 60312749 ( <i>Gryllus bimaculatus</i> )	tgcaaggagggtgacaggctgaggatgaattgtaca cgagaaggagaaagtacaagt
Bg031	144 55895696 ( <i>Locusta migratoria</i> )	ttgtacacgagaaggagaagtacaagtacat
Bg031	145 60311892 ( <i>Euclidia glyphica</i> ); 55900730 ( <i>Locusta migratoria</i> ); 60311708 ( <i>Euclidia glyphica</i> )	gagaaggagaagtacaagtacattgtgacgatcttgat gacttcaccga
Bg031	146 77732463 ( <i>Aedes aegypti</i> ); 4378572 ( <i>Periplaneta americana</i> ); 19310970 ( <i>Periplaneta fuliginosa</i> )	aacagcaggcccgacgccaac
Bg031	147 19310970 ( <i>Periplaneta fuliginosa</i> ); 60311610 ( <i>Euclidia glyphica</i> ); 60313268 ( <i>Gryllus bimaculatus</i> )	cattgtgacgatcttgatgacttcaaccgaactattgg
Bg032	148 76169650 ( <i>Diptera punctata</i> )	cggacaggaggagacatcaactc
Bg032	149 18888282 ( <i>Anopheles gambiae</i> )	tggaagaagtcgaagagcgtaag
Bg032	150 91094918 ( <i>Tribolium castaneum</i> )	ctgctccaagatccagaaaca
Bg033	151 60310034 ( <i>Scarabaeus laticollis</i> ); 83933868 ( <i>Lutzomyia longipalpis</i> ); 90137292 ( <i>Spodoptera frugiperda</i> ); 82610902 ( <i>Tineola bisselliella</i> ); 5853355 ( <i>Lymantaria dispar</i> ); 50818292 ( <i>Heliconius melpomene</i> ); 22474252 ( <i>Helicoverpa armigera</i> ); 58371832 ( <i>Lononia obliqua</i> ); 3719570 ( <i>Manduca sexta</i> ); 25959205 ( <i>Meladema coriacea</i> );	gaggccagagacaagaggagtactcactcgaagtac cccat



		53883538 ( <i>Plutella xylostella</i> ); 34787974 ( <i>Callosobruchus maculatus</i> ); 16901146 ( <i>Ctenocephalides felis</i> ); 60309684 ( <i>Scarabaeus laticollis</i> ); 78050191 ( <i>Heliconius erato</i> ); 57963831 ( <i>Heliconius melpomene</i> ); 60305522 ( <i>Mycetophagus quadripustulatus</i> ); 60295481 ( <i>Homalodisca coagulata</i> ); 71539924 ( <i>Oncometopia nigricans</i> ); 29556355 ( <i>Bombyx mori</i> ); 14010638 ( <i>Heliothis virescens</i> ); 5853355 ( <i>Lymantria dispar</i> ); 293219 ( <i>Manduca sexta</i> ); 40218737 ( <i>Spodoptera exigua</i> ); 67838313 ( <i>Drosophila pseudoobscura</i> )		
Bg033	152	83662157 ( <i>Myzus persicae</i> )	gtccagagggaacacccaatcct	
Bg033	153	71543527 ( <i>Oncometopia nigricans</i> ); 60295481 ( <i>Homalodisca coagulata</i> ); 71048162 ( <i>Oncometopia nigricans</i> )	atcctgctgactgagggtccct	
Bg033	154	49206619 ( <i>Drosophila melanogaster</i> ); 22474062 ( <i>Helicoverpa armigera</i> ); 29535046 ( <i>Bombyx mori</i> ); 60295481 ( <i>Homalodisca coagulata</i> ); 60336301 ( <i>Homalodisca coagulata</i> ); 62387502 ( <i>Reticulitermes flavipes</i> ); 60311490 ( <i>Euclidia glyphica</i> ); 91082248 ( <i>Tribolium castaneum</i> ); 67838313 ( <i>Drosophila pseudoobscura</i> ); 24251124 ( <i>Culicoides sp.</i> ); 60304993 ( <i>Dascillus cervinus</i> ); 25956497 ( <i>Biphylus lunatus</i> ); 62239347 ( <i>Diabrotica virgifera</i> ); 55783599 ( <i>Apriona germari</i> ); 50818328 ( <i>Heliconius melpomene</i> ); 75723625 ( <i>Tribolium castaneum</i> ); 60305522 ( <i>Mycetophagus quadripustulatus</i> ); 49395567 ( <i>Drosophila melanogaster</i> ); 67838495 ( <i>Drosophila pseudoobscura</i> ); 50818292 ( <i>Heliconius melpomene</i> )	aaggccaacagggagaagatgacitcaaatcatgtttgag acctcaa	
Bg033	155	25959205 ( <i>Meladema coriacea</i> ); 18923947 ( <i>Anopheles gambiae</i> ); 3477239 ( <i>Drosophila melanogaster</i> ); 29556355 ( <i>Bombyx mori</i> ); 56772582 ( <i>Drosophila virilis</i> ); 34788040 ( <i>Callosobruchus maculatus</i> ); 60315015 ( <i>Tricholepisma aurea</i> ); 49395567 ( <i>Drosophila melanogaster</i> ); 60314849 ( <i>Tricholepisma aurea</i> ); 60314729 ( <i>Tricholepisma aurea</i> ); 34787974 ( <i>Callosobruchus maculatus</i> ); 60315012 ( <i>Tricholepisma aurea</i> ); 50560908 ( <i>Homalodisca coagulata</i> ); 62387502 ( <i>Reticulitermes flavipes</i> ); 60305522	caaatcatgtttgagacctcaacacccc	

		(Mycetophagus quadripustulatus) ; 62387510 (Reticulitermes flavipes) ; 60295481 (Homalodisca coagulata)	
Bg033	156	71547931 (Oncometopia nigricans); 60311490 (Euclidia glyphica); 75723625 (Tribolium castaneum); 49394847 (Drosophila melanogaster); 61949513 (Tribolium castaneum)	tcatgtttgagaccttcaacacccccgcgatgtatgt
Bg033	157	37951847 (Ips pini); 60299272 (Diaphorina citri); 73615611 (Aphis gossypii); 84648237 (Myzus persicae); 86461101 (Acyrtosiphon pisum); 52630958 (Toxoptera citricida); 34788040 (Callosobruchus maculatus); 60315015 (Tricholepisma aurea)	accttaacacccccgcgatgtatgtgccaatccaggc
Bg033	158	37804525 (Rhopalosiphum padi); 86307561 (Culex pipiens); 62238804 (Diabrotica virgifera); 40310862 (Timarcha balearica); 49005801 (Drosophila melanogaster); 83933868 (Lutzomyia longipalpis)	cccgccatgtatgttgccaatccaggccgt
Bg033	159	67782282 (Aedes aegypti); 48718502 (Anopheles funestus); 189333335 (Anopheles gambiae); 58385473 (Anopheles gambiae str. PEST); 66509773 (Apis mellifera); 45331062 (Megachile rotundata); 18923947 (Anopheles gambiae) ; 60312762 (Gryllus bimaculatus) ; 90137292 (Spodoptera frugiperda)	gccatccaggccgtgctgtccct
Bg033	160	60312762 (Gryllus bimaculatus)	tacgcttcggccgtaccactggtattgtg
Bg033	161	30031443 (Toxoptera citricida)	gcttcggccgtaccactggtat
Bg033	162	34788040 (Callosobruchus maculatus); 19613046 (Anopheles gambiae); 92043996 (Drosophila willistoni); 58375293 (Anopheles gambiae str. PEST)	cgtaccactggtattgtcctggactctgtgga
Bg033	163	18938956 (Anopheles gambiae); 92926094 (Drosophila virilis)	ggtattgtctggactctggtgacgg
Bg033	164	92473382 (Drosophila erecta); 78050191 (Heliconius erato); 78230609 (Heliconius erato/himera mixed EST library)	ggtgacggcgctctccacacgt
Bg033	165	29556355 (Bombyx mori); 55923288 (Locusta migratoria)	gtctccacacccgtaccatctatgaaggta
Bg033	166	60304032 (Eucinetus sp.); 37951847 (Ips pini); 60310034 (Scarabaeus laticollis); 55913655 (Locusta migratoria)	tcccacacccgtaccatctatgaaggttacgc
Bg033	167	60311532 (Euclidia glyphica); 60311490 (Euclidia glyphica); 62387510 (Reticulitermes flavipes); 60309684 (Scarabaeus laticollis)	tgaagtacccattgaacatggaatcatcacaactggga

Bg033	168	92948842 ( <i>Drosophila ananassae</i> ); 62387555 ( <i>Reticulitermes flavipes</i> ); 3113938 ( <i>Drosophila melanogaster</i> ); 68267390 ( <i>Drosophila simulans</i> ); 12802910 ( <i>Coptotermes acinaciformis</i> ); 55917578 ( <i>Locusta migratoria</i> ); 78050191 ( <i>Heliconius erato</i> )	tgcccatgccaatcctgctgctggactt
Bg033	169	78231052 ( <i>Heliconius erato</i> /himera mixed EST library); 29551161 ( <i>Bombyx mori</i> ); 55888553 ( <i>Locusta migratoria</i> ); 33528426 ( <i>Trichoplusia ni</i> ); 22474252 ( <i>Helicoverpa armigera</i> ); 55896579 ( <i>Locusta migratoria</i> )	gccatcctgctgctggactggccggccgt
Bg033	170	55924447 ( <i>Locusta migratoria</i> ); 57963831 ( <i>Heliconius melpomene</i> ); 293219 ( <i>Manduca sexta</i> ); 60314729 ( <i>Tricholepisma aurea</i> ); 50818292 ( <i>Heliconius melpomene</i> )	cgctggactggccggccgtgactgac
Bg033	171	55901019 ( <i>Locusta migratoria</i> ); 67877117 ( <i>Drosophila pseudoobscura</i> ); 42765807 ( <i>Armigeres subalbatus</i> ); 92044691 ( <i>Drosophila willistoni</i> ); 91718815 ( <i>Liriomyza huidobrensis</i> ); 67838313 ( <i>Drosophila pseudoobscura</i> )	cgtagtactgactaccctgatgaagatcct
Bg033	172	13761518 ( <i>Drosophila melanogaster</i> ); 18938956 ( <i>Anopheles gambiae</i> ); 18923947 ( <i>Anopheles gambiae</i> ); 18933335 ( <i>Anopheles gambiae</i> ); 18928068 ( <i>Anopheles gambiae</i> ); 77731484 ( <i>Aedes aegypti</i> ); 21260592 ( <i>Culex pipiens</i> ); 20146853 ( <i>Simulium vittatum</i> ); 51978737 ( <i>Bacillus cereus</i> )	gactacctgatgaagatcctgaccgagcgtggctac
Bg033	173	12802910 ( <i>Coptotermes acinaciformis</i> )	atgaagatcctgaccgagcgtggctacagcttcac
Bg033	174	84648237 ( <i>Myzus persicae</i> ); 86461101 ( <i>Acyrtosiphon pisum</i> ); 73618206 ( <i>Aphis gossypii</i> ); 55913634 ( <i>Locusta migratoria</i> ); 37804558 ( <i>Rhopalosiphum padi</i> ); 52630958 ( <i>Toxoptera citricida</i> ); 37593622 ( <i>Pedicularis humanus</i> ); 49395567 ( <i>Drosophila melanogaster</i> ); 37951847 ( <i>Ips pini</i> )	ggaatcatcaccacaaactgggatgacatgga
Bg033	175	55888553 ( <i>Locusta migratoria</i> ); 53883538 ( <i>Plutella xylostella</i> ); 29535046 ( <i>Bombyx mori</i> ); 2700128 ( <i>Drosophila melanogaster</i> ); 50560971 ( <i>Homalodisca coagulata</i> ); 55901019 ( <i>Locusta migratoria</i> ); 29551161 ( <i>Bombyx mori</i> ); 55901019 ( <i>Locusta migratoria</i> ); 22474062 ( <i>Helicoverpa armigera</i> ); 50560971 ( <i>Homalodisca coagulata</i> ); 55888553 ( <i>Locusta migratoria</i> ); 53883538 ( <i>Plutella xylostella</i> );	atcatcaccacactgggatgacatggagaagatctggca

Bg033	176	29556355 (Bombyx mori); 55901019 (Locusta migratoria); 55901019 (Locusta migratoria) 677900 (Aedes aegypti); 42764600 (Armigeres subalbatus); 51978737 (Bacillus cereus); 86465013 (Bombyx mori); 90811718 (Culex pipiens); 92460622 (Drosophila erecta); 67838495 (Drosophila pseudobscura); 92926494 (Drosophila virilis); 83934452 (Lutzomyia longipalpis); 90814004 (Nasonia vitripennis); 71547039 (Oncometopia nigricans); 60315012 (Tricholepisma aurea); 71048162 (Oncometopia nigricans); 82610902 (Tineola bisselliella); 60310034 (Scarabaeus laticollis); 5853355 (Lymantria dispar); 60309684 (Scarabaeus laticollis); 49005801 (Drosophila melanogaster); 60314849 (Tricholepisma aurea); 60312762 (Gryllus bimaculatus); 60314729 (Tricholepisma aurea); 60311532 (Euclidia glyphica); 3338522 (Drosophila melanogaster); 55886573 (Locusta migratoria); 34579881 (Aedes aegypti); 25959205 (Meladema coriacea); 57963831 (Heliconius melpomene); 58371832 (Lonomia obliqua); 78230609 (Heliconius erato/himera mixed EST library)	gacatggagaagatctggcatcacacaccttctacaa
Bg033	177	25957102 (Carabus granulatus); 18939947 (Anopheles gambiae); 56152104 (Rhynchosciara americana); 60315012 (Tricholepisma aurea); 60310833 (Agriotes lineatus); 60297606 (Diaprepes abbreviatus); 25958625 (Curculio glandium); 34787974 (Callosobruchus maculatus)	atggagaagatctggcatcacacaccttctacaaatgaa
Bg033	178	25956583 (Biphyllus lunatus); 37951847 (Ips pini); 40544541 (Tribolium castaneum); 56772582 (Drosophila virilis); 34788040 (Callosobruchus maculatus); 25956497 (Biphyllus lunatus)	aagatctggcatcacacaccttctacaaatgaactcog
Bg033	179	16901057 (Ctenocephalides felis); 56772662 (Drosophila virilis); 60310833 (Agriotes lineatus); 57963831 (Heliconius melpomene); 60297606 (Diaprepes abbreviatus); 60314729 (Tricholepisma aurea); 60311490 (Euclidia glyphica); 87266181 (Choristoneura fumiferana); 62239347 (Diabrotica virgifera); 60315015 (Tricholepisma aurea); 677900 (Aedes aegypti); 51978737 (Bacillus cereus); 5853355 (Lymantria dispar); 55783599 (Apriona germari); 83934452	atctggcatcacacaccttctacaaatgaactcogagt

	(Lutzomyia longipalpis) ; 19848020 (Chelonius inanitus) ; 82610902 (Tineola bisselliella) ; 42765392 (Armigeres subalbatus); 82611040 (Trox sp.)	
--	--	--

Table 5

Target ID	SEQ ID NO	Example GI-number and species	Sequence*
Bg005	180	82835847 (Boophilus microplus); 63511642 (Ixodes scapularis)	aacgtgttcaagaacaacgctgtcct
Bg005	181	82835847 (Boophilus microplus)	catccacaagaagaagctgagaagccagg
Bg031	182	21642857 (Amblyomma variegatum); 4325305 (Boophilus microplus); 49549243 (Rhipicephalus appendiculatus)	gocatcaagaagaagatgcaggcgatgaagctggagaaga ga
Bg031	183	22758956 (Haemaphysalis longicornis)	cctgcagaagaagatccagcagat
Bg031	184	83308264 (Dermanyssus gallinae); 22758956 (Haemaphysalis longicornis)	aagatgcaggcgatgaagctggagaaggacaa
Bg031	185	21642857 (Amblyomma variegatum); 29779612 (Ornithodoros porcinus);	gagaaggacaagggccctgcag
Bg031	186	10707547 (Amblyomma americanum); 21642025 (Amblyomma variegatum); 49535169 (Rhipicephalus appendiculatus)	gtgtcggcaacaacacctgaagtcct
Bg031	187	29779612 (Ornithodoros porcinus)	aaggaggctgaagctcgtctga
Bg033	188	28627064 (Mesobuthus gibbosus)	gaggccagagcaagagaggtatcctc
Bg033	189	68767268 (Acanthoscurria gomesiana)	cagagcaagagaggtatcctcac
Bg033	190	18143239 (Araneus ventricosus)	gcccagagcaagagaggtatcctcacctgaagt
Bg033	191	32423713 (Haemaphysalis longicornis)	aaggccaacaggagagaagatgac
Bg033	192	45269080 (Ornithodoros moubata)	gagaagatgactcaaatcatgtt
Bg033	193	32423713 (Haemaphysalis longicornis)	ggtatctactctgaagtacccattga
Bg033	194	68764791 (Acanthoscurria gomesiana)	atcatgtttgagacctcaac
Bg033	195	10708501 (Amblyomma americanum); 60730229 (Ixodes ricinus); 63510574 (Ixodes scapularis); 49538235 (Rhipicephalus appendiculatus); 77539276 (Ornithodoros moubata); 29779134 (Ornithodoros porcinus)	gagacctcaacaccccccgccatgta
Bg033	196	10708501 (Amblyomma americanum)	gccatccaggccgctgtctcct

Bg033	197	77539276 (Ornithodoros moubata); 29779134 (Ornithodoros porcinus); 68764791 (Acanthoscurria gomesiana)	gtctcccaacacccgtaccatctatgaaggttacgc
Bg033	198	68767268 (Acanthoscurria gomesiana); 77539276 (Ornithodoros moubata); 29779134 (Ornithodoros porcinus); 45269080 (Ornithodoros moubata); 68758323 (Acanthoscurria gomesiana)	tcaccaactgggatgacatgggagaagatctggcatcacac
Bg033	199	68764791 (Acanthoscurria gomesiana)	gacatggagaagatctggcatcacacacctctctacaa
Bg033	200	18143239 (Araneus ventricosus); 28627064 (Mesobuthus gibbosus)	atggagaagatctggcatcacacacctctctacaatgaactcgg

## EXAMPLES

### **Example 1: Cloning of a partial sequence of the *Blattella germanica* Bg001, Bg003, Bg004 and Bg005 genes via family PCR**

High quality, intact RNA was isolated from *Blattella germanica* (source: Central  
5 Science Laboratory, York) using TRIzol Reagent (Cat. N°. 15596-026/15596-018, Invitrogen, Rockville, Maryland, USA) following the manufacturers directions. Genomic DNA present in the RNA preparation was removed by Dnase treatment as prescribed by the manufacturer. cDNA was generated using a commercially available kit (SuperScript™ III Reverse Transcriptase, Cat N° 18080044, Invitrogen, Rockville,  
10 Maryland, USA) following the manufacturers directions.

To isolate cDNA sequences comprising a portion of the Bg001, Bg003, Bg004 and Bg005 genes, a series of PCR reactions with degenerate primers were performed using Amplitaq Gold (Cat. N°. N8080240; Applied Biosystems) as prescribed by the manufacturer.

15 For Bg001, the degenerate primers oGBKA002 and oGBKA020 (represented herein as SEQ ID NO 3 and SEQ ID NO 4 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 57 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick  
20 Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N°. K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 1 and is referred to as the partial sequence of the Bg001 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 2.

25 For Bg003, the degenerate primers oGBKC001 and oGBKC010 (represented herein as SEQ ID NO: 13 and SEQ ID NO: 14 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 55 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel,  
30 purified (QIAquick Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N°. K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 11 and is referred to as the partial sequence of the Bg003 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 12.

For Bg004, the degenerate primers oGBKD001 and oGBKD006 (represented herein as SEQ ID NO 23 and SEQ ID NO 24 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 50 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N° K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 21 and is referred to as the partial sequence of the Bg004 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 22.

For Bg005, the degenerate primers oGBKE002 and oGBKE009 (represented herein as SEQ ID NO 33 and SEQ ID NO 34 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 52 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N° K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 31 and is referred to as the partial sequence of the Bg005 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 32.

**Example 2: Cloning of a partial sequence of the *Blattella germanica* Bg031, Bg032 and Bg033 genes via EST sequence**

High quality, intact RNA was isolated from *Blattella germanica* (source: Central Science Laboratory, York) using TRIzol Reagent (Cat. N° 15596-026/15596-018, Invitrogen, Rockville, Maryland, USA) following the manufacturers directions. Genomic DNA present in the RNA preparation was removed by DNase treatment as prescribed by the manufacturer. cDNA was generated using a commercially available kit (SuperScript™ III Reverse Transcriptase, Cat N° 18080044, Invitrogen, Rockville, Maryland, USA) following the manufacturers directions.

To identify a partial cDNA sequence from the Bg031, Bg032 and Bg033 genes, one EST per gene was found in the public database Genbank under accession numbers AF260897, X73679 and AY004248 respectively, originating from the public database Genbank.



To isolate cDNA sequences comprising a portion of the Bg031, Bg032 and Bg033 genes, a series of PCR reactions with EST based specific primers were performed using Perfectshot™ ExTaq (Cat N° RR005A, TAKARA BIO INC.) as prescribed by the manufacturer.

5 For Bg031, the specific primers oGBLA001 and oGBLA002 (represented herein as SEQ ID NO 43 and SEQ ID NO 44 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 55 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick  
10 Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N° K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 41 and is referred to as the partial sequence of the Bg031 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 42.

15 For Bg032, the specific primers oGBLB003 and oGBLB004 (represented herein as SEQ ID NO: 51 and SEQ ID NO: 52 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 55 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick  
20 Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N° K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 49 and is referred to as the partial sequence of the Bg032 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 50.

25 For Bg033, the specific primers oGBLC001 and oGBLC004 (represented herein as SEQ ID NO 59 and SEQ ID NO 60 respectively) were used in two independent PCR reactions with the following conditions: 10 minutes at 95 °C, followed by 40 cycles of 30 seconds at 95 °C, 1 minute at 55 °C and 1 minute at 72 °C, followed by 7 minutes at 72 °C. The resulting PCR products were analyzed on agarose gel, purified (QIAquick  
30 Gel Extraction Kit; Cat. N° 28706, Qiagen), cloned into the pCR4-TOPO vector (Cat. N° K4575-40, Invitrogen) and sequenced. The consensus sequence resulting from the sequencing of both PCR products is herein represented by SEQ ID NO 57 and is referred to as the partial sequence of the Bg033 gene. The corresponding partial amino acid sequence is herein represented as SEQ ID NO 58.

**Example 3: dsRNA production of the *Blattella germanica* Bg001, Bg003, Bg004, Bg005, Bg031, Bg032 and Bg033 genes**

dsRNA was synthesized in milligram amounts using the commercially available kit T7 RiboMAX™ Express RNAi System (Cat. N°. P1700, Promega). First two separate  
5 single 5' T7 RNA polymerase promoter templates were generated in two separate PCR reactions, each reaction containing the target sequence in a different orientation relative to the T7 promotor.

For Bg001 the sense T7 template was generated using the specific T7 FW primer oGBLD001 and the specific RV primer oGBLD010 (represented herein as SEQ ID NO 5  
10 and SEQ ID NO 6 respectively) in a PCR reaction with the following conditions: 4 minutes at 95°C, followed by 35 cycles of 30 seconds at 95°C, 30 seconds at 51°C and 1 minute at 72°C, followed by 10 minutes at 72°C. The anti-sense T7 template was generated using the specific FW primer oGBLD009 and the specific T7 RV primer oGBLD002 (represented herein as SEQ ID NO 7 and SEQ ID NO 8 respectively) in a  
15 PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein  
20 represented by SEQ ID NO 9.

For Bg003 the sense T7 template was generated using the specific T7 FW primer oGBLD003 and the specific RV primer oGBLD012 (represented herein as SEQ ID NO 15 and SEQ ID NO 16 respectively) in a PCR reaction with the following conditions: 4  
25 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLD011 and the specific T7 RV primer oGBLD004 (represented herein as SEQ ID NO: 17 and SEQ ID NO: 18 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7  
30 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 19.

For Bg004 the sense T7 template was generated using the specific T7 FW primer oGBLD005 and the specific RV primer oGBLD014 (represented herein as SEQ ID NO 25 and SEQ ID NO 26 respectively) in a PCR reaction with the following conditions: 4 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLD013 and the specific T7 RV primer oGBLD006 (represented herein as SEQ ID NO: 27 and SEQ ID NO: 28 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 29.

For Bg005 the sense T7 template was generated using the specific T7 FW primer oGBLD007 and the specific RV primer oGBLD016 (represented herein as SEQ ID NO 35 and SEQ ID NO 36 respectively) in a PCR reaction with the following conditions: 4 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLD015 and the specific T7 RV primer oGBLD008 (represented herein as SEQ ID NO: 37 and SEQ ID NO: 38 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 39.

For Bg031 the sense T7 template was generated using the specific T7 FW primer oGBLA007 and the specific RV primer oGBLA002 (represented herein as SEQ ID NO 45 and SEQ ID NO 44 respectively) in a PCR reaction with the following conditions: 4 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLA001 and the specific T7 RV primer oGBLA008 (represented herein as SEQ ID NO 43 and SEQ ID NO 46 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products

were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 47.

For Bg032 the sense T7 template was generated using the specific T7 FW primer oGBLB007 and the specific RV primer oGBLB004 (represented herein as SEQ ID NO 53 and SEQ ID NO 52 respectively) in a PCR reaction with the following conditions: 4 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLB003 and the specific T7 RV primer oGBLB008 (represented herein as SEQ ID NO 51 and SEQ ID NO 54 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 55.

For Bg033 the sense T7 template was generated using the specific T7 FW primer oGBLC007 and the specific RV primer oGBLC004 (represented herein as SEQ ID NO 61 and SEQ ID NO 60 respectively) in a PCR reaction with the following conditions: 4 minutes at 95 °C, followed by 35 cycles of 30 seconds at 95 °C, 30 seconds at 51 °C and 1 minute at 72 °C, followed by 10 minutes at 72 °C. The anti-sense T7 template was generated using the specific FW primer oGBLC001 and the specific T7 RV primer oGBLC008 (represented herein as SEQ ID NO 59 and SEQ ID NO 62 respectively) in a PCR reaction with the same conditions as described above. The resulting PCR products were analyzed on agarose gel and purified by NaClO<sub>4</sub> precipitation. The generated T7 FW and RV templates were mixed to be transcribed and the resulting RNA strands were annealed, Dnase and Rnase treated, and purified by sodium acetate, following the manufacturer's directions. The sense strand of the resulting dsRNA is herein represented by SEQ ID NO 63.

**Example 4: Laboratory trials to screen dsRNA targets for activity against the German cockroach, *Blattella germanica***

Stock solutions of 1-10µg/µl dsRNA in distilled water were prepared. Each dsRNA solution was diluted to the appropriate concentration and mixed with finely ground laboratory diet (Rat and mouse standard diet, B&K Universal Ltd, Hull, UK), which was previously heat treated in order to inactivate any enzymes. The mixture or  
5 formulation was formed into small pellets of equal weight (0.3g) to achieve an end concentration of 0.1% to 2%w/w dsRNA and dried overnight at room temperature.

Newly hatched nymphs from the German cockroach, *B. germanica* were housed per 10 in plastic lidded containers (29±2°C, minimum 40% relative humidity, with a 12:12 light:dark photoperiod). Animals were starved 24 hours prior to exposure to the pellets.  
10 The cockroaches were assessed as live, moribund or dead twice a week until adulthood. The pellet was replaced with freshly prepared pellet once a week. dsRNA containing pellets, formulations, were compared with a negative control (solvent) and a positive control (1 or 2 % imidacloprid, commonly used in commercially available cockroach baits). As shown in Figure 1, at least 80 % of the cockroaches died within 24 days after  
15 first administration when treated with Bg001, Bg003 and Bg005, or within 29 days when treated with Bg004 respectively.

#### **Example 5: Testing different fragments for efficiency**

##### ***Identification of a fragment of the *Blattella germanica* Bg001, Bg003, Bg004, Bg005, Bg031, Bg032 and Bg033 genes with no substantial homology to human***

  
20

The partial sequences of the Bg001, Bg003, Bg004, Bg005, Bg031, Bg032 and Bg033 genes, herein represented respectively as SEQ ID NO 1, SEQ ID NO 11, SEQ ID NO 21, SEQ ID NO 31, SEQ ID NO 41, SEQ ID NO 49 and SEQ ID NO 57, were analyzed to find fragments with no substantial homology to non-target organisms. In particular, since the dsRNA will be diced in the organism to siRNA molecules, the  
25 sequences were scanned for siRNA sequences that would have homology to non-target species. Such siRNA could cause adverse effects in the non-target organism and should therefore preferably be avoided in the dsRNA fragment to be incorporated in the end products. The selected fragments are suitable for cockroach control by RNA interference  
30 when for instance present in the bait and taken up by a cockroach feeding from the bait. For this analysis, non-target organism was human (*Homo sapiens*). Fragments of 21 contiguous nucleotides (best1\_human\_21\_0), or 24 contiguous nucleotides allowing three mismatches (best1/2/3\_human\_24\_3), that do not occur in the non-target organism were identified and are named herein "freefrags." The longest sequence of Bg001,

Bg003, Bg004, Bg005, Bg031, Bg032 and Bg033 free of non-target organism sequences using the first selection criterium was given a SEQ ID NO and named herein "freefrag". These Bg001, Bg003, Bg004, Bg005, Bg031, Bg032 and Bg033 freefrags, are herein represented as SEQ ID NO 10, SEQ ID NO 20, SEQ ID NO 30, SEQ ID NO 40, SEQ ID NO 48, SEQ ID NO 56 and SEQ ID NO 64, respectively. The length and sequence of some examples of other freefrag sequences suitable for use in the present invention is represented in Table 2. The exact sequence can easily be deduced from the table. All freefrag sequences described in the table belong to the group of sequences of the invention.

A person skilled in the art will recognize that many more such freefrags, of various lengths, may be identified in the *Blattella germanica* sequences herein presented, as well as in sequences which are orthologues of the corresponding genes and proteins in the other pest sequences according to the invention, and accordingly, the present invention extends to these further identifiable freefrags.

#### **Example 6: Choosing the optimal fragment; testing concatemers for efficacy**

Concatemers were designed for each target gene. Concatemers are synthetic tandem repeats of 50 to 100 bp dsfragments. In the present example, concatemers were designed by selecting the best possible fragments in regions with homology in protein family as well as at nucleotide level, in regions containing the best predicted siRNAs and in regions with between 40 and 60% GC content, preferably about 50% GC content, if possible.

For Bg001 two concatemers were designed consisting of a five times repeat of a 50 bp fragment, ie represented by SEQ ID NO 65 and 66, resulting in Bg001 concatemer 1 and Bg001 concatemer 2, herein represented respectively as SEQ ID NO 67 and SEQ ID NO 68, and one concatemer was designed consisting of a three times repeat of a 100 bp repeat, ie represented by SEQ ID NO 69, resulting in Bg001 concatemer 3, herein represented as SEQ ID NO 70. XbaI and SmaI flanking sites were added for cloning in a vector to produce dsRNA. These dsRNA constructs comprising the concatemers were tested in the cockroach laboratory trials.

In a further experiment as shown in Figure 2, mortality was significant higher when treated with Bg001 and Bg001 concatemer 2 compared to the negative control (solvent)..

Besides mortality, treatment showed a significant effect on development. For example at day 48, from the surviving cockroaches treated with Bg001 only 33.3 % moulted to the adult stage whereas none of the cockroaches treated with Bg001 concatemer 2 did within this time as shown in Table 3.

5

#### **Example 7: Testing different formulations**

RNA interference (RNAi) is a potentially very powerful tool to inhibit the expression of target genes in a sequence-specific manner in many different species. However, for RNAi to be valuable and effective, specific silencing of any given target gene is essential, devoid of nonspecific knockdown and toxic side effects. Applications of dsRNA have been hindered by the inability to effectively deliver these compounds to their sites of action within cells. Progress in chemical modification of the dsRNA to enhance the strength and stability of interaction, without losing specificity, is ongoing. In this study an evaluation is made of a few concepts for delivery of dsRNA to target genes in *B. germanica*.

15

RNAi induced effects can be improved by increasing the intracellular uptake of dsRNA by facilitating endocytosis or by increasing the stability of the dsRNA in the biological environment using delivery agents such as lipids and liposomes. siRNAs have anionic phosphodiester backbones and for this reason, cationic liposome/lipid-mediated siRNA delivery (siFection) is investigated. These cationic liposome/lipid-based systems are selected from a number of commercially available products, including lipofectamine and 1,2-dioleoyl-3-trimethyl-ammonium-propane (DOTAP)-cholesterol, and test the dsRNA formulations in the cockroach laboratory trials. Parameters to be investigated include the lipid:dsRNA ratio of mixing, the extent of cationic liposome/lipid-dsRNA complex formation, the particle size, the mode of delivery and the dose-response effect.

20

25

#### **Example 8: Testing for dsRNA stability**

Application of dsRNA for gene silencing will be dependent on improvements in molecule bio-stability, specificity and delivery.

30

The stability of the generated dsRNAs was tested in TRIZMA buffer at pH 7 and pH 9 and in CAPS buffer at pH 11 to mimic the pH in the gut of some target species. dsRNA was incubated for several days and aliquots were analyzed on 20% polyacrylamide gels at different time intervals. No influence of the pH on the stability of the dsRNA could be observed based on the gel results.

The stability of the generated dsRNAs as a function of time was tested in RNase free water and in LB medium at room temperature over a period of eight months. Aliquots were taken weekly and/or monthly and stored at -20°C prior to analysis on 20% polyacrylamide gels. No significant degradation of the dsRNA could be observed on a polyacrylamide gel as shown in Figure 3;

**Example 9: Laboratory trial to test a single dose dsRNA on mortality in the early nymphal stage of the German cockroach, *Blattella germanica***

A stock solution of 10µg/µl dsRNA in distilled water was prepared and mixed with finely ground laboratory diet (Rat and mouse standard diet, B&K Universal Ltd, Hull, UK), which was previously heat treated in order to inactivate any enzymes. The mixture or formulation was formed into small pellets of equal weight (0.3g) to achieve an end concentration of 1%w/w dsRNA and dried overnight at room temperature.

Newly hatched nymphs from the German cockroach, *B. germanica* were housed per 10 in plastic lidded containers (29±2°C, minimum 40% relative humidity, with a 12:12 light:dark photoperiod). Animals were starved 24 hours prior to exposure to the pellets. After one week, this initial dose was replaced with untreated pellet. The cockroaches were assessed as live, moribund or dead twice a week until adulthood. Bg001 dsRNA containing pellet showed significant higher mortality compared to the two negative controls (solvent and miscellaneous dsRNA) as shown in Figure 4.



## Claims

1. A nucleic acid molecule comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180 and 181, or an orthologous  
5 nucleotide sequence from an insect or arachnid species, wherein the orthologous nucleotide sequence has at least 70% sequence identity with the nucleotide sequence of any one of SEQ ID NOs 1, 11, 21 and 31.
2. An RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion  
10 of the nucleotide sequence of any one of (i) the nucleic acid molecules defined in claim 1 or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect and/or arachnid species.
- 15 3. The RNA construct of claim 2 wherein said at least one double stranded RNA region has a length of at least 17 bp.
4. The RNA construct according to claim 2 or 3 wherein the complementarity of said nucleotide sequence comprises at least 70% nucleotide sequence identity with (i) the portion of the nucleotide sequence of the nucleic acid molecules defined in claim 1 or  
20 claim 2 or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect or arachnid species, wherein the percentage sequence identity is calculated over the same length.
- 25 5. The RNA construct according to any of claims 2 to 4 wherein the at least one strand which comprises a nucleotide sequence that is complementary to (i) the portion of the nucleotide sequence of the nucleic acid molecules defined in claim 1 or claim 2 or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of  
30 SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect or arachnid species has less than 12.5% sequence identity over 24 contiguous nucleotides with the corresponding nucleotide sequence from a mammalian species.

6. An RNA construct according to any of claims 2 to 5 comprising at least two nucleotide sequences independently chosen from any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, for instance any of SEQ ID NOs 65 to 70.
7. A DNA construct (i) comprising the nucleotide sequence of claim 1 or (ii) comprising a region encoding an RNA construct of any one of claims 2 to 6.
8. An expression construct comprising a DNA construct according to claim 7.
9. An expression construct according to claim 8 further comprising one or more control sequences capable of driving expression of the nucleic acid defined in claim 7; and optionally a transcription termination sequence.
10. A host cell comprising an RNA construct of any of claims 2 to 6, a DNA construct of claim 7 or 8 and/or an expression construct of claim 9.
11. A pesticide composition comprising an RNA construct as defined in any one of claims 2 to 6 and/or a DNA construct as defined in claim 7 or 8 and/or an expression construct as defined in claim 9 and/or a host cell as defined in claim 10 together with a suitable carrier.
12. The composition according to claim 10 wherein the carrier comprises electrostatically charged powder or particles and/or magnetic particles, preferably metallic particles which are initially unmagnetised but which are capable of becoming magnetically polarised when subjected to the electrical field provided by the insect or arachnid body, which powder or particles adhere to the insect or arachnid cuticle and which may be ingested by an insect or arachnid.
13. A housing or trap for insects and/or arachnids which contains a composition as defined in claim 11 or 12.
14. Use of an RNA construct as defined in any one of claims 2 to 6 and/or a DNA construct as defined in claim 7 or 8 and/or an expression construct as defined in claim 9 and/or a host cell as defined in claim 10 and/or a composition as defined in claim 11 or 12 and/or housing or trap as defined in claim 13 for controlling an insect and/or arachnid by RNA interference.

15. A method of controlling insects and/or arachnids comprising administering to an insect and/or arachnid an RNA construct as defined in any one of claims 2 to 6 and/or a DNA construct as defined in claim 7 or 8 and/or an expression construct as defined in claim 9 and/or a host cell as defined in claim 10 and/or a composition as defined in claims 11 or 12 and/or housing or trap as defined in claim 13, wherein the double stranded RNA is capable of down regulating the expression of at least one insect and/or arachnid gene through RNA interference.

16. The method of claim 15, wherein multiple RNA constructs as defined in any one of claims 2 to 6 and/or DNA constructs as defined in claim 7 or 8 and/or expression constructs as defined in claim 9 and/or host cells as defined in claim 10 and/or compositions as defined in claims 11 or 12 and/or housing or trap as defined in claim 13 are administered to said insects and/or arachnids.

17. The method of claim 16, wherein the multiple RNA constructs as defined in any one of claims 2 to 6 and/or DNA constructs as defined in claim 7 or 8 and/or expression constructs as defined in claim 9 and/or host cells as defined in claim 10 and/or compositions as defined in claims 11 or 12 and/or housing or trap as defined in claim 13 are administered sequentially in order to reduce the probability of the insect and/or arachnid acquiring resistance.

18. The use according to claim 14 or the method according to any of claims 15 to 17 wherein the insects and/or arachnids comprise household insects, ecto-parasites and insects and/or arachnids relevant for public health and hygiene, preferably flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas.

19. The use or the method according to claim 18, wherein the insect and/or arachnids comprise cockroaches (Blattodea) such as but not limited to *Blatella* spp. (e.g. *Blatella germanica* (german cockroach)), *Periplaneta* spp. (e.g. *Periplaneta americana* (American cockroach) and *Periplaneta australasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus* spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex*

spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g. *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite)), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)), *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)), *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)), and most preferably German cockroach (*Blatella germanica*).

20. A kit for use in the method of any one of claims 15 to 19, which comprises the RNA construct as defined in any one of claims 2 to 6 and/or a DNA construct as defined in claim 7 or 8 and/or an expression construct as defined in claim 9 and/or a host cell as defined in claim 10 and/or a composition as defined in claim 11 or 12 and/or housing or trap as defined in claim 13 together with instructions for use, wherein the double stranded RNA is capable of down regulating the expression of at least one pest gene through RNA interference.

21. The kit of claim 20 which comprises multiple RNA constructs as defined in any one of claims 2 to 6 and/or the DNA constructs as defined in claim 7 or 8 and/or the expression constructs as defined in claim 9 and/or the host cells as defined in claim 10 or 11 and/or the compositions as defined in claims 11 or 12 and/or housing or trap as defined in claim 13, wherein each double stranded RNA is capable of down regulating the expression of at least one pest gene through RNA interference.

22. The kit of claim 20 wherein the multiple RNA constructs as defined in any one of claims 2 to 6 and/or the DNA constructs as defined in claim 7 or 8 and/or the expression constructs as defined in claim 9 and/or the host cells as defined in claim 10 and/or the compositions as defined in claim 11 or 12 and/or housing or trap as defined in claim 13 are used sequentially in order to reduce the probability of the pest acquiring resistance.

23. A method for controlling cockroach pests comprising providing to the cockroach an RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence encoding a cockroach ribosomal protein.

24. The method according to claim 23 wherein at least one strand of the at least one double stranded RNA region comprises at least 17, 18, 19, 20, 21 or more nucleotides of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.
25. The RNA construct of any one of claims 2 to 6 which comprises at least one additional dsRNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence of a further gene from an insect and/or arachnid species.
26. The RNA construct of any one of claims 2 to 6 or 25, further comprising at least one additional functional sequence and optionally a linker.
27. The RNA construct according to claim 26 wherein said additional functional sequence is chosen from the group comprising (i) a sequence facilitating large-scale production of the RNA construct; (ii) a sequence effecting an increase or decrease in the stability of the dsRNA; (iii) a sequence allowing the binding of proteins or other molecules to facilitate uptake of the RNA construct by an insect and/or arachnid; (iv) a sequence which is an aptamer that binds to receptors or to molecules in the gut of an insect to facilitate uptake, endocytosis and/or transcytosis by the insect and/or arachnid.
28. The RNA construct according to claim 26 or 27 wherein the linker is a conditionally self-cleaving RNA sequence, preferably a pH sensitive linker or a hydrophobic sensitive linker.
29. The RNA construct according to claims 26 or 27 wherein the linker is an intron.
30. The RNA construct according to any one of claims 23 to 27 wherein the complementarity of said nucleotide sequence comprises at least 70% nucleotide sequence identity with (i) the portion of the nucleotide sequence of the nucleic acid molecules defined in claim 1 or claim 2 or (ii) the nucleic acid molecules comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect and/or arachnid species, wherein the percentage sequence identity is calculated over the same length.

31. The RNA construct according to any one of claims 25 to 30 wherein the at least one strand which comprises a nucleotide sequence that is complementary to (i) the portion of the nucleotide sequence of the nucleic acid molecules defined in claim 1 or claim 2 or (ii) the nucleic acid molecules and/or arachnid comprising the nucleotide sequence as set forth in any one of SEQ ID NOs 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or an orthologous nucleotide sequence thereof from an insect species has less than 12.5% sequence identity over 24 contiguous nucleotides with the corresponding nucleotide sequence from a mammalian species.

32. The RNA construct according to any of claims 3 to 6, or 25 to 28 wherein the at least one strand comprises at least 17, 18, 19, 20, 21 nucleotides or more of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

33. The RNA construct according to claim 31 or 32, wherein the mammalian species is a human.

34. An RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises at least 17, 18, 19, 20, 21 nucleotides or more of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

35. A DNA construct comprising a region encoding an RNA construct of any one of claims 25 to 34.

36. An expression construct comprising a DNA construct according to claim 35.

37. A host cell comprising an RNA construct of any of claims 25 to 34, a DNA construct of claim 35 or an expression construct of claim 36.

38. The host cell as defined in claim 10 or 37, which is a bacterial cell.
39. The host cell as defined in claim 38, which is an inactivated bacterial cell.
40. A method for generating the RNA construct of any of claims 2 to 6, or 25 to 34, comprising the steps of
- 5           a. contacting a DNA construct of claim 7 or 35 or an expression construct of claim 8, 9 or 36 with cell-free components; or
- b. administering a DNA construct of claim 7 or 35 or an expression construct of claim 8, 9 or 36 to a cell,
- under conditions that allow transcription of said DNA construct to produce said
- 10           RNA construct.
41. The method of claim 40 wherein the cell is a bacterial cell.
42. A pesticide composition comprising an RNA construct as defined in any one of claims 25 to 34 and/or a DNA construct as defined in claim 35 and/or an expression construct as defined in claim 36 and/or a host cell as defined in any one of claims 37 to
- 15           39 together with a suitable carrier.
43. The composition according to any of claims 10, 11, or 43 which is in a form suitable for ingestion by an insect and/or arachnid.
44. The composition according to claim 42 or 43 which is in solid form, such as a pellet or powder, liquid form or gel form.
- 20           45. The composition according to any one of claims 42 to 44 which is in the form of a bait.
46. The composition according to claim 45 wherein the bait further includes at least one food substance, such as a protein based food or boric acid and/or an attractant, such as a pheromone.
- 25           47. The composition according to any one of claims 42 to 46 wherein the composition is stored in a housing or trap which an insect and/or arachnid can enter in order to ingest the composition.
48. The composition according to any one of claims 42 to 46 which is in the form of a spray, preferably a pressurized/aerosolized spray or a pump spray.

49. The composition according to any one of claims 42 to 46 which is in the form of a coating on a suitable surface which adheres to an insect and/or arachnid when it comes into contact with the coating.
50. The composition according to any one of claims 42 to 46 wherein the carrier  
5 comprises electrostatically charged powder or particles and/or magnetic particles, preferably metallic particles which are initially unmagnetised but which are capable of becoming magnetically polarised when subjected to the electrical field provided by the insect and/or arachnid body, which adhere to the insect and/or arachnid cuticle.
51. The composition according to any one of claims 42 to 50 wherein the carrier  
10 increases the uptake of the double stranded RNA into the pest.
52. The composition of claim 51 wherein the carrier is a lipid-based carrier, preferably comprising one or more of, oil-in water emulsions, cholesterol, micelles, lipopolyamines and liposomes.
53. The composition of claim 51 wherein the carrier comprises a nucleic acid  
15 condensing agent, preferably spermidine or protamine sulphate.
54. The composition of any one of claims 11 12, or 42 to 53 in combination with a further pesticide.
55. A housing or trap for insects and/or arachnids which contains a composition as defined in any one of claims 42 to 46 or 54.
- 20 56. Use of an RNA construct as defined in any one of claims 25 to 34 and/or a DNA construct as defined in claim 35 and/or an expression construct as defined in claim 36 and/or a host cell as defined in any one of claims 37 to 39 and/or a composition as defined in any one of claims 42 to 54 and/or housing or trap as defined in claim 55 for controlling an insect and/or arachnid by RNA interference.
- 25 57. The use according to claim 56, wherein the insects and/or arachnids comprise household insects, ecto-parasites and insects and/or arachnids relevant for public health and hygiene, preferably flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas.
- 30 58. The use according to claim 57, wherein the insect and/or arachnids comprise cockroaches (Blattodea) such as but not limited to *Blatella* spp. (e.g. *Blatella germanica* (german cockroach)), *Periplaneta* spp. (e.g. *Periplaneta americana* (American



cockroach) and *Periplaneta australasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus* spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex* spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g. *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)), *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)), *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)), and most preferably German cockroach (*Blatella germanica*).

59. The use according to any one of claims 14, 18, or 56 to 58 wherein the insect and/or arachnid is growth delayed, paralysed, made infertile or killed.

60. A method of controlling insects and/or arachnids comprising administering to an insect and/or arachnid an RNA construct as defined in any one of claims 25 to 34 and/or a DNA construct as defined in claim 35 and/or an expression construct as defined in claim 36 and/or a host cell as defined in any one of claims 37 to 39 and/or a composition as defined in any one of claims 42 to 54 and/or housing or trap as defined in claim 13, wherein the double stranded RNA is capable of down regulating the expression of at least one insect and/or arachnid gene through RNA interference.

61. The method of claim 60, wherein multiple RNA constructs as defined in any one of claims 25 to 34 and/or DNA constructs as defined in claim 35 and/or expression constructs as defined in claim 36 and/or host cells as defined in any one of claims 37 to 39 and/or compositions as defined in any one of claims 42 to 54 and/or housing or trap as defined in claim 13 are administered.

62. The method of claim 61, wherein the multiple RNA constructs as defined in any one of claims 25 to 34 and/or DNA constructs as defined in claim 35 and/or expression

constructs as defined in claim 36 and/or host cells as defined in any one of claims 37 to 39 and/or compositions as defined in any one of claims 42 to 54 and/or housing or trap as defined in claim 13 are administered sequentially in order to reduce the probability of the pest acquiring resistance.

- 5 63. The method according to any one of claims 60 to 62, wherein the insects and/or arachnids comprise household insects, ecto-parasites and insects and/or arachnids relevant for public health and hygiene, preferably flies, spider mites, thrips, ticks, red poultry mite, ants, cockroaches, termites, crickets including house-crickets, silverfish, booklice, beetles, earwigs, mosquitos and fleas.
- 10 64. The method according to any one of claims 60 to 63, wherein the insects and/or arachnids comprise cockroaches (Blattodea) such as but not limited to *Blatella* spp. (e.g. *Blatella germanica* (german cockroach)), *Periplaneta* spp. (e.g. *Periplaneta americana* (American cockroach) and *Periplaneta australiasiae* (Australian cockroach)), *Blatta* spp. (e.g. *Blatta orientalis* (Oriental cockroach)) and *Supella* spp. (e.g. *Supella longipalpa* (brown-banded cockroach); ants (Formicoidea), such as but not limited to *Solenopsis* spp. (e.g. *Solenopsis invicta* (Red Fire Ant)), *Monomorium* spp. (e.g. *Monomorium pharaonis* (Pharaoh Ant)), *Camponotus* spp. (e.g. *Camponotus* spp (Carpenter Ants)), *Iasius* spp. (e.g. *Iasius niger* (Small Black Ant)), *Tetramorium* spp. (e.g. *Tetramorium caespitum* (Pavement Ant)), *Myrmica* spp. (e.g. *Myrmica rubra* (Red Ant)), *Formica* spp (wood ants), *Crematogaster* spp. (e.g. *Crematogaster lineolata* (Acrobat Ant)), *Iridomyrmex* spp. (e.g. *Iridomyrmex humilis* (Argentine Ant)), *Pheidole* spp. (Big Headed Ants), and *Dasymutilla* spp. (e.g. *Dasymutilla occidentalis* (Velvet Ant)); termites (Isoptera and/or Termitidae) such as but not limited to *Amitermes* spp. (e.g. *Amitermes floridensis* (Florida dark-winged subterranean termite)), *Reticulitermes* spp. (e.g. *Reticulitermes flavipes* (the eastern subterranean termite), *Reticulitermes hesperus* (Western Subterranean Termite)), *Coptotermes* spp. (e.g. *Coptotermes formosanus* (Formosan Subterranean Termite)), *Incisitermes* spp. (e.g. *Incisitermes minor* (Western Drywood Termite)) and *Neotermes* spp. (e.g. *Neotermes connexus* (Forest Tree Termite)).
- 20 65. The method according to any one of claims 15 to 16 or 6 to 62 wherein the insect and/or arachnid is growth delayed, paralysed, made infertile or killed.
- 25 66. A kit for use in the method of any one of claims 60 to 65, which comprises the RNA construct as defined in any one of claims 25 to 34 and/or a DNA construct as
- 30

defined in claim 35 and/or an expression construct as defined in claim 36 and/or a host cell as defined in any one of claims 37 to 39 and/or a composition as defined in any one of claims 42 to 54 and/or housing or trap as defined in claim 13 together with instructions for use, wherein the double stranded RNA is capable of down regulating the expression of at least one pest gene through RNA interference.

67. The kit of claim 64 which comprises multiple RNA constructs as defined in any one of claims 53 to 34 and/or DNA constructs as defined in claim 35 and/or expression constructs as defined in claim 36 and/or host cells as defined in any one of claims 37 to 39 and/or compositions as defined in any one of claims 42 to 54, wherein each double stranded RNA is capable of down regulating the expression of at least one pest gene through RNA interference.

68. The kit of claim 67 wherein the multiple RNA constructs as defined in any one of claims 25 to 34 and/or the DNA constructs as defined in claim 35 and/or the expression constructs as defined in claim 36 and/or the host cells as defined in any one of claims 37 to 39 and/or the compositions as defined in any one of claims 42 to 54 are used sequentially in order to reduce the probability of the pest acquiring resistance.

69. A method for controlling cockroach pests comprising providing administering to the cockroach an RNA construct comprising at least one double stranded RNA region, at least one strand of which comprises a nucleotide sequence that is complementary to a portion of the nucleotide sequence encoding a tropomyosin, a HMG Coenzyme A synthase gene or an Actin 5C gene.

70. The method according to claim 69 wherein at least one strand of the at least one double stranded region comprises at least 17, 18, 19, 20, 21 nucleotides or more of any of the nucleic acid molecules comprising the nucleotide sequence as set forth in any of SEQ ID NOs 1, 4, 6, 7, 9, 10, 65 to 70, 71 to 79, 11, 16, 17, 19, 20, 80 to 87, 21, 26, 27, 29, 30, 88 to 93, 31, 36, 37, 39, 40, 94 to 108, 180, 181, 41, 43, 44, 47, 48, 109 to 147, 182 to 187, 49, 51, 52, 55, 56, 148 to 150, 57, 63, 64, 151 to 179, and 188 to 200, or the complement thereof.

71. The method according to any of claims 23, 24, 69 or 70 wherein said cockroach pest is chosen from the species belonging to following group of genera comprising *Blatella*, *Periplaneta*, *Blatta* and *Supella*.

72. A method according to any of claims 23, 24 or 69 to 71 wherein said cockroach pest is chosen from the group comprising German cockroach (*Blatella Germanica*),

American cockroach (*Periplaneta americana*), Australian cockroach (*Periplaneta australiasiae*), Oriental cockroach (*Blatta orientalis*) and brown-banded cockroach (*Supella longipalpa*).

5 73. A protein comprising the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32 or an orthologous protein having a conserved amino acid sequence from a further insect and/or arachnid species.

74. The protein of claim 73, wherein the orthologue has at least 70% amino acid sequence identity with the amino acid sequence as set forth in any one of SEQ ID NOs 2, 12, 22 or 32.

10 75. A method for generating the RNA construct of any of claims 2 to 6 or, 25 to 34, comprising culturing a host cell as defined in any of claims 10 or 37 to 39 under conditions that permit expression of the RNA construct.

76. The method of claim 75 further comprising purification of the RNA construct.

1 / 13




FIG. 1




FIG. 2

2 / 13




FIG. 3




FIG. 4

3 / 13

SEQ ID NO 1; DNA; *Blattella germanica*

taaggcatggatgttggacaagctcgggtggagtgtatgctccaagaccaagcacaggacctcacaagttacgagagagtctgc  
 cccttgtaatatcttctgtaataggctgaaatgcatcattaaccaactgtgaggttaagaaaatgttatgcagcgcttattaagggtg  
 atggaaaagtcagaacagaccccaactatccagctggtttatggatgtgttacaattgaaaaaactggagaattttccgtctgat  
 ttatgacgtgaaaggacgtttcaccattcacagaataactgctgaagaagccaagtataaactgtgcaaggtaaagagagtgc  
 gactgggcccagggtattccattctgtgacccatgatggtagaactcttagatatcctgatcctgtcatcaaagtaataatgataca  
 gtcaacttgacatcgctactccaagattatggatagcatcaaattgataatggtaatctctgtatgattactggaggccgtaacttg  
 ggtcgtgttgaactgtagttaatcgagaacgtcatcctggttcctttgacattgtgcatgttaaagattcacaaggacacacatttgc  
 taccagattgaa

SEQ ID NO 2; PRT; *Blattella germanica*

Lys Ala Trp Met Leu Asp Lys Leu Gly Gly Val Tyr Ala Pro Arg Pro Ser Thr Gly Pro His Lys  
 Leu Arg Glu Ser Leu Pro Leu Val Ile Phe Leu Arg Asn Arg Leu Lys Tyr Ala Leu Thr Asn Cys  
 Glu Val Lys Lys Ile Val Met Gln Arg Leu Ile Lys Val Asp Gly Lys Val Arg Thr Asp Pro Asn Tyr  
 Pro Ala Gly Phe Met Asp Val Val Thr Ile Glu Lys Thr Gly Glu Phe Phe Arg Leu Ile Tyr Asp  
 Val Lys Gly Arg Phe Thr Ile His Arg Ile Thr Ala Glu Glu Ala Lys Tyr Lys Leu Cys Lys Val Lys  
 Arg Val Gln Thr Gly Pro Lys Gly Ile Pro Phe Leu Val Thr His Asp Gly Arg Thr Leu Arg Tyr  
 Pro Asp Pro Val Ile Lys Val Asn Asp Thr Val Gln Leu Asp Ile Ala Thr Ser Lys Ile Met Asp Ser  
 Ile Lys Phe Asp Asn Gly Asn Leu Cys Met Ile Thr Gly Gly Arg Asn Leu Gly Arg Val Gly Thr  
 Val Val Asn Arg Glu Arg His Pro Gly Ser Phe Asp Ile Val His Val Lys Asp Ser Gln Gly His  
 Thr Phe Ala Thr Arg Leu Asn

SEQ ID NO 3; DNA; *Blattella germanica*; degenerate primer; w is a or t; r is a or g; m is a or  
 c; y is c or t

catttgaagc gtttwrmygc ycc

SEQ ID NO 4; DNA; *Blattella germanica*; specific primer

gtgcccttgc caatgatgaa cacgttg

SEQ ID NO 5; DNA; specific T7 primer

cgctaatacg actcactata ggggagtgtg tgctccaaga ccaag

SEQ ID NO 6; DNA; *Blattella germanica*; specific primer

caatctggta gcaaatgtgt gtcc

SEQ ID NO 7; DNA; *Blattella germanica*; specific primer

ggagtgtatg ctccaagacc aag

SEQ ID NO 8; DNA; synthetic T7 primer

cgctaatacg actcactata ggcaatctgg tagcaaatgt gtgtcc

SEQ ID NO 9; RNA; *Blattella germanica*; u can be t (for DNA complement)

ggaguguaug cuccaagacc aagcacagga ccucacaagu uacgagagag ucugcccuu  
 \_

FIG. 5

---

```

guaauauuuc uucguaauag gcugaaauau gcauuuaccca acugugaggu uaagaaaauu
guuaugcagc gccuuauuaa gguugaugga aaagucagaa cagaccccaa cuauccagcu
gguuuuauug auguuguuac aaauugaaaaa acuggagaau uuuuccgucu gauuuauagc
gugaaaggac guuucaccau ucacagaaua acugcugaag aagccaagua uaaacugugc
aagguaaaga gagugcagac ugggcccag gguauuccau ucuuggugac ccaugauggu
agaacucuua gauauccuga uccugucauc aaaguuaaug auacaguuca acuugacauc
gcuacuucca agauuaugga uagcaucaa uuugauaaug guaaucucug uaugauuacu
ggaggccgua acuugggucg uguuggaacu guaguuaauc gagaacguca uccugguucc
uuugacaauug ugcauguuaa agauucacaa ggacacacau uugcuaccag auug

```

SEQ ID NO 10; RNA; *Blattella germanica*; u can be t (for DNA complement)

```

ggaguguaug cuccaagacc aagcacagga ccucacaagu uacgagagag ucugcccuu
guaauauuuc uucguaauag gcugaaauau gcauuuaccca acugugaggu uaagaaaauu
guuaugcagc gccuuauuaa gguugaugga aaagucagaa cagaccccaa cuauccagcu
gguuuuauug auguuguuac aaauugaaaaa acuggagaau uuuuccgucu gauuuauagc
gugaaaggac guuucaccau ucacagaaua acugcugaag aagccaagua uaaacugugc
aagguaaaga gagugcagac ugggcccag gguauuccau ucuuggugac ccaugauggu
agaacucuua gauauccuga uccugucauc aaaguuaaug auacaguuca acuugacauc
gcuacuucca agauuaugga uagcaucaa uuugauaaug guaaucucug uaugauuacu
ggaggccgua acuugggucg uguuggaacu guaguuaauc gagaacguca uccugguucc
uuugacaauug ugcauguuaa agauucacaa gga

```

SEQ ID NO 11; DNA; *Blattella germanica*;

```

tcccaggcgaccttatgaaaaggcacgtcttgatcaggaggtgaaaatcataggagaatatggtcttaggaacaaacgtgaagt
gtggcgagtcaagtataccttgcaaaaatccgtaaagctgccagagaacttctgactttggaagagaaagatcagcgaggtt
gttgaaggcaatgctcttctcgtcggttggtgcgtattggagtggttgatgaaacccgtatgaagcttgattacgtctgggttgaag
attgaagatttcttgaacgacgtctcacaacacaagtttcaagttggggcttgcaaatcaatccatcatgctcgtgtgctgatcc
gtcaaagacatatcagggttcgtaagcaggtcgtgaatatccaagcttcattgtgagacttgattccagaagcatattgactctc
g

```

SEQ ID NO 12; PRT; *Blattella germanica*;

```

Pro Arg Arg Pro Tyr Glu Lys Ala Arg Leu Asp Gln Glu Leu Lys Ile Ile Gly Glu Tyr Gly Leu Arg
Asn Lys Arg Glu Val Trp Arg Val Lys Tyr Thr Leu Ala Lys Ile Arg Lys Ala Ala Arg Glu Leu
Leu Thr Leu Glu Glu Lys Asp Gln Arg Arg Leu Phe Glu Gly Asn Ala Leu Leu Arg Arg Leu Val
Arg Ile Gly Val Leu Asp Glu Thr Arg Met Lys Leu Asp Tyr Val Leu Gly Leu Lys Ile Glu Asp
Phe Leu Glu Arg Arg Leu Gln Thr Gln Val Phe Lys Leu Gly Leu Ala Lys Ser Ile His His Ala
Arg Val Leu Ile Arg Gln Arg His Ile Arg Val Arg Lys Gln Val Val Asn Ile Pro Ser Phe Ile Val
Arg Leu Asp Ser Gln Lys His Ile Asp Phe
Ser

```

SEQ ID NO 13; DNA; *Blattella germanica*; degenerate primer; n is a, c, g, or t; y is c or t; k is g or t

```
tcggtcttct cgaagacnta ygtkac
```

SEQ ID NO 14; DNA; *Blattella germanica*; degenerate primer; m is a or c; y is c or t; b is c, g or t

```
ccgccgaagg gmgayttbag
```

---



5 / 13

---

 SEQ ID NO 15; DNA; specific T7 primer

cgctaatacg actcactata ggcagcgac cttatgaaa ggc

SEQ ID NO 16; DNA; *Blattella germanica*; specific primer

cgagaagtca atatgcttct ggg

SEQ ID NO 17; DNA; *Blattella germanica*; specific primer

caggcgacct tatgaaaagg c

SEQ ID NO 18; DNA; specific T7 primer

cgctaatacg actcactata ggcgagaagt caatatgctt ctggg

SEQ ID NO 19; RNA; *Blattella germanica*; u can be t (for DNA complement)

caggcgaccu uaugaaaagg cagcucuuga ucaggaguug aaaaucauag gagaauaugg  
 ucuuaggaac aaacgugaag uguggcgagu caaguauacc uuggcaaaaa uccguaaagc  
 ugccagagaa cuucugacuu uggaagagaa agaucagcgc agguuguuug aaggcaaugc  
 ucuucucgu cgguuggugc guauuggagu guuggaugaa acccguauga agcuugauua  
 cgucuuuggu uugaagauug aagauuucuu ggaacgacgu cuccaaacac aaguuuucua  
 guuggggcuu gcaaaaucua uccaucaugc ucgugugcug auccgucaaa gacauaucag  
 gguucguaag caggucguga auauuccaag cuucauugug agacuugauu cccagaagca  
 uauugacuuc ucg

SEQ ID NO 20; RNA; *Blattella germanica*; u can be t (for DNA complement)

caggcgaccu uaugaaaagg cagcucuuga ucaggaguug aaaaucauag gagaauaugg  
 ucuuaggaac aaacgugaag uguggcgagu caaguauacc uuggcaaaaa uccguaaagc  
 ugccagagaa cuucugacuu uggaagagaa agaucagcgc agguuguuug aaggcaaugc  
 ucuucucgu cgguuggugc guauuggagu guuggaugaa acccguauga agcuugauua  
 cgucuuuggu uugaagauug aagauuucuu ggaacgacgu cuccaaacac aaguuuucua  
 guuggggcuu gcaaaaucua uccaucaugc ucgugugcug auccgucaaa gacauaucag  
 gguucguaag caggucguga auauuccaag cuucauugug agacuugauu cc

SEQ ID NO 21; DNA; *Blattella germanica*;

tgtgaaaggaccacgaggcacctgaagcgcggttcaagcatcttgccttagatatccacttggtcatccaaggctcctgaagg  
 ggaaaaatggttggaacaaagaaggagttggcagccgtgcgcacgctgctctcatattgagaacatgattaaaggagtcac  
 aaagggttctgtacaaaatgcgcgcgtgtatgccatttccccattaactgcgtaaccacagaaaacaattccgttattgaagt  
 gcgtaacttctggcgagaagttcatccgcagagtgagatggctccgggagtgaccgtcaccaatttccaaagcagaaag  
 acgagctcattctggagggaacgacatcgaggatgtatcgagatcagccgcactcatccaacaatcgacgactgtgaagaa  
 caaggacatccggaattccttgac

SEQ ID NO 22; PRT; *Blattella germanica*;

Val Lys Gly Pro Arg Gly Thr Leu Lys Arg Gly Phe Lys His Leu Ala Leu Asp Ile His Leu Val  
 His Pro Arg Leu Leu Lys Val Glu Lys Trp Phe Gly Thr Lys Lys Glu Leu Ala Ala Val Arg Thr  
 Val Cys Ser His Ile Glu Asn Met Ile Lys Gly Val Thr Lys Gly Phe Leu Tyr Lys Met Arg Ala Val  
 Tyr Ala His Phe Pro Ile Asn Cys Val Thr Thr Glu Asn Asn Ser Val Ile Glu Val Arg Asn Phe  
 Leu Gly Glu Lys Phe Ile Arg Arg Val Lys Met Ala Pro Gly Val Thr Val Thr Asn Ser Pro Lys  
 Gln Lys Asp Glu Leu Ile Leu Glu Gly Asn Asp Ile Glu Asp Val Ser Arg Ser Ala Ala Leu Ile Gln  
 Gln Ser Thr Thr Val Lys Asn Lys Asp Ile

---

FIG. 5 CONT'D

6 / 13

---

Arg Lys Phe Leu Asp

SEQ ID NO 23; DNA; *Blattella germanica*; forward degenerative primer; n is a, c, g, or t  
gtgaaggccc gnntggtgac 20

SEQ ID NO 24; DNA; *Blattella germanica*; reverse degenerative primer; -d is a, g or t; h is a, c or t; r is a or g; v is a, c or g  
gtcgtcttct cdgahacrta vagacc

SEQ ID NO 25; DNA; T7 primer forward  
cgctaatacg actcactata gggtgaaagg accacgaggc acc

SEQ ID NO 26; DNA; *Blattella germanica*; specific primer reverse  
ccgtcaagga atttccggat g

SEQ ID NO 27; DNA; *Blattella germanica*; specific primer forward  
gtgaaaggac cacgaggcac c

SEQ ID NO 28; DNA; specific T7 primer reverse  
cgctaatacg actcactata ggccgtcaag gaatttccgg atg

SEQ ID NO 29; RNA; *Blattella germanica*; u can be t (for DNA complement)

gugaaaggac cacgaggcac cuugaagcgc gguuucagc aucuugcuuu agauauccac  
uugguucauc caaggcuccu gaagguggaa aaaugguuug gaacaaagaa ggaguuggca  
gccgugcgca ccgucugcuc ucauauugag aacaugauua aaggagucac aaaggguuuc  
cuguacaaaa ugcgcgccgu guauggccau uuccccauua acugcguaac cacagaaaaac  
aaauccguua uugaagugcg uaacuucuuug ggcgagaagu ucauccgcag agugaagaug  
gcuccgggag ugaccgucac caauucucca aagcagaaag acgagcucan ucuggagggc  
aacgacaucg aggauguauc gagaucagcc gcacucaucc aacaucgac gacugugaag  
aacaaggaca uccggaaaau ccuugacgg

SEQ ID NO 30; RNA; *Blattella germanica*; u can be t (for DNA complement)

gugaaaggac cacgaggcac cuugaagcgc gguuucagc aucuugcuuu agauauccac  
uugguucauc caaggcuccu gaagguggaa aaaugguuug gaacaaagaa ggaguuggca  
gccgugcgca ccgucugcuc ucauauugag aacaugauua aaggagucac aaaggguuuc  
cuguacaaaa ugcgcgccgu guauggccau uuccccauua acugcguaac cacagaaaaac  
aaauccguua uugaagugcg uaacuucuuug ggcgagaagu ucauccgcag agugaagaug  
gcuccgggag ugaccgucac caauucucca aagcagaaag acgagcucan ucuggagggc  
aacgacaucg aggauguauc gagaucagcc gcacucaucc aacaucgac gacugugaag  
aacaagga

---

FIG. 5 CONT'D

SEQ ID NO 31; DNA; *Blattella germanica*;

ggcttgatcccaatgaaataaacgaaattgcaaataccaattccagacaaaatattcgtaaactgattaaagatggctttatcatc  
aagaagcccgtagctgtactcaagggcccggtgttcgcaagaacaccgaagcaagaagaaaaggacgtcactgcggttttg  
gcaaaaggaagggtacggcaaatgccgtatgccacagaagggtctgtggattaatcgcgtgttctgagaaggcttctca  
agaagtacaggggaagcaaagaagatcgacagacatctataccaccagctgtacatgaaggccaagggtaacgtgttcaaga  
acaagcgtgtcctgatggagttcatccacaagaagaaggctgagaaggccaggacaaagatgcttaacgac

SEQ ID NO 32; PRT; *Blattella germanica*;

Leu Asp Pro Asn Glu Ile Asn Glu Ile Ala Asn Thr Asn Ser Arg Gln Asn Ile Arg Lys Leu Ile Lys  
Asp Gly Leu Ile Ile Lys Lys Pro Val Ala Val His Ser Arg Ala Arg Val Arg Lys Asn Thr Glu Ala  
Arg Arg Lys Gly Arg His Cys Gly Phe Gly Lys Arg Lys Gly Thr Ala Asn Ala Arg Met Pro Gln  
Lys Val Leu Trp Ile Asn Arg Met Arg Val Leu Arg Arg Leu Leu Lys Lys Tyr Arg Glu Ala Lys  
Lys Ile Asp Arg His Leu Tyr His Gln Leu Tyr Met Lys Ala Lys Gly Asn Val Phe Lys Asn Lys  
Arg Val Leu Met Glu Phe Ile His Lys Lys Lys Ala Glu Lys Ala Arg Thr Lys Met Leu Asn Asp

SEQ ID NO 33; DNA; *Blattella germanica*; degenerative primer forward; r is a or g; b is c or g  
or t

tgcatgctgg caaraaraag gtbttg

SEQ ID NO 34; DNA; *Blattella germanica*; degenerative primer reverse; y is c or t; r is a or g

gtcggcgagc ytcrgcytg

SEQ ID NO 35; DNA; specific T7 primer forward

cgctaatacg actcactata ggggcttgat cccaatgaaa taaacg

SEQ ID NO 36; DNA; *Blattella germanica*; specific primer reverse

gtcgttaagc atctttgtcc tggc

SEQ ID NO 37; DNA; *Blattella germanica*; specific primer forward

ggcttgatcc caatgaaata aacg

SEQ ID NO 38; DNA; specific T7 primer reverse

cgctaatacg actcactata gggtcgttaa gcatctttgt cctggc

SEQ ID NO 39; RNA; *Blattella germanica*; u can be t (for DNA complement)

ggcuugaucc caaugaaaua aacgaaaug caaaauaccaa uuccagacaa aaauauucgua  
aacugauuaa agauggucuu aucaucaaga agcccguagc uguacacuca agggcccug

---

uucgcaagaa caccgaagca agaagaaaag gacgucacug cgguuuuggc aaaaggaagg  
 guacggcaaa ugcccguaug ccacagaagg ucuuguggau uaaucgcaug cguguucuga  
 gaaggcuucu caagaaguac agggaagcaa agaagaucga cagacaucua uaccaccagc  
 uguacaugaa ggccaagggg aacguguuca agaacaagcg uguccugaug gaguuccauc  
 acaagaagaa ggcugagaag gccaggacaa agaugcuuaa cgac

SEQ ID NO 40; RNA; *Blattella germanica*; u can be t (for DNA complement)

ggcuugauc caaugaaaua aacgaaaauug caaaauaccaa uuccagacaa aaauuucgua  
 aacugauuaa agauggucuu aucaucaaga agcccguagc uguacacuca agggcccgug  
 uucgcaagaa caccgaagca agaagaaaag gacgucacug cgguuuuggc aaaaggaagg  
 guacggcaaa ugcccguaug ccacagaagg ucuuguggau uaaucgcaug cguguucuga  
 gaaggcuucu caagaaguac agggaagcaa agaagaucga cagacaucua uaccaccagc  
 uguacaugaa ggccaagggg aacguguuca agaacaagcg uguccugaug gaguuccauc  
 acaagaagaa ggcugagaag gcc

SEQ ID NO 41; DNA; *Blattella germanica*;

atggatgcatcaagaagaagatgcaggcgatgaagctggagaaggacaacgcgatggatcgcgcccttctctgcaacag  
 caggcccgacgccaacatccgggcccagagaaggctgaggaggaggcccgatccctgcagaagaagatccagcagattg  
 agaatgatcttgatcagaccatggagcagttgatgcaagtcaacgccaagctggacgagaaggacaaggccctgcagaatgc  
 tgagagtgaggtcgctgccctcaaccgccgaatccaactgctggagga  
 ggatcttgagaggtctgaggaacgttggccacagccaccgccaagttggctgaggtctccagggtccgatgagtcagagc  
 gagctcgtgaagattcttgaatccaaaggcctggcagatgaagagcgtatggatgcttggagaaccagctgaaggaagccagg  
 ttcattggctgaggaagctgacaagaaatgatgaggtcgacgtaagtggctatggtgaggccgacttgaaagagcaga  
 agagcgtgccgagactggtgaatccaagattgtggagcttgaggaagaactgcgctgttcggcaacaacctgaagtccctga  
 ggtgtctgaagagaaggccaacctgcgtgaggaagagtacaagcaacagattaagactctgaataccagggttaaaggaggc  
 tgaagctcgtgctgagttcgtgaaagatccgtgcagaaattgcagaaggaggtgacaggcttgaggatgaattggtacacga  
 gaaggagaagtaacaagtacattgtgacgatcttgatgatcttcaccgaactattggc

SEQ ID NO 42; PRT; *Blattella germanica*;

Met Asp Ala Ile Lys Lys Lys Met Gln Ala Met Lys Leu Glu Lys Asp Asn Ala Met Asp Arg Ala  
 Leu Leu Cys Glu Gln Gln Ala Arg Asp Ala Asn Ile Arg Ala Glu Lys Ala Glu Glu Glu Ala Arg  
 Ser Leu Gln Lys Lys Ile Gln Gln Ile Glu Asn Asp Leu Asp Gln Thr Met Glu Gln Leu Met Gln  
 Val Asn Ala Lys Leu Asp Glu Lys Asp Lys Ala Leu Gln Asn Ala Glu Ser Glu Val Ala Ala Leu  
 Asn Arg Arg Ile Gln Leu Leu Glu Glu Asp Leu Glu Arg Ser Glu Glu Arg Leu Ala Thr Ala Thr  
 Ala Lys Leu Ala Glu Ala Ser Gln Ala Ala Asp Glu Ser Glu Arg Ala Arg Lys Ile Leu Glu Ser  
 Lys Gly Leu Ala Asp Glu Glu Arg Met Asp Ala Leu Glu Asn Gln Leu Lys Glu Ala Arg Phe Met  
 Ala Glu Glu Ala Asp Lys Lys Tyr Asp Glu Val Ala Arg Lys Leu Ala Met Val Glu Ala Asp Leu  
 Glu Arg Ala Glu Glu Arg Ala Glu Thr Gly Glu Ser Lys Ile Val Glu Leu Glu Glu Glu Leu Arg  
 Val Val Gly Asn Asn Leu Lys Ser Leu Glu Val Ser Glu Glu Lys Ala Asn Leu Arg Glu Glu Glu  
 Tyr Lys Gln Gln Ile Lys Thr Leu Asn Thr Arg Leu Lys Glu Ala Glu Ala Arg Ala Glu Phe Ala  
 Glu Arg Ser Val Gln Lys Leu Gln Lys Glu Val Asp Arg Leu Glu Asp Glu Leu Val His Glu Lys  
 Glu Lys Tyr Lys Tyr Ile Cys Asp Asp Leu Asp Met Thr Phe Thr Glu Leu Ile Gly

SEQ ID NO 43; DNA; *Blattella germanica*; specific primer forward

atggatgcca tcaagaagaa gatgcag

---

SEQ ID NO 44; DNA; *Blattella germanica*; specific primer reverse

gccataaagt tcggtgaaag tcatatcaag

SEQ ID NO 45; DNA; specific T7 primer forward

cgctaatacg actcactata ggatggatgc catcaagaag aagatg

SEQ ID NO 46; DNA; specific T7 primer reverse

cgctaatacg actcactata gggccaataa gttcgggtgaa agtcat

SEQ ID NO 47; RNA; *Blattella germanica*; u can be t (for DNA complement)

auggaugcca ucaagaagaa gaugcaggcg augaagcugg agaaggacaa cgcgauaggau  
cgcgcccuuc ucugcgaaca gcaggcccg gacgccaaca uccgggccga gaaggcugag  
gaggaggccc gaucccugca gaagaagauc cagcagauug agaaugaucu ugaucagacc  
auggagcagu ugaugcaagu caacgccaaug cuggacgaga aggacaaggc ccugcagaau  
gcugagagug agguccguc ccucaaccgc cgaauccaac ugcuggagga ggaucuugag  
aggucugagg aacguuuggc cacagccacc gccaauguug cugaggcuuc ccaggcugcc  
gaugagucag agcagcucg uaagauucuu gaauccaaag gccuggcaga ugaagagcgu  
auggaugcuu uggagaacca gcugaaggaa gccagguuca uggcugagga agcugacaag  
aaauaugaug agguccgacg uaaguuggcu augguugagg ccgacuugga aagagcagaa  
gagcgugccg agacugguga auccaagauu guggagcuug aggaagaacu gcgcuuguc  
ggcaacaacc ugaaguccu ugaggugucu gaagagaagg ccaaccugcg ugaggaagag  
uacaagcaac agauuaagac ucugaauacc aggcuaaagg aggcugaagc ucgugcugag  
uucgcugaaa gauccgugca gaaauugcag agggagguug acaggcuuga ggaugaauug  
guacacgaga aggagaagua caaguacauu ugugacgauc ugaugaugac uuacaccgaa  
cuauuuggc

SEQ ID NO 48; RNA; *Blattella germanica*; u can be t (for DNA complement)

ccaaucaagaa gaagaugcag gcgaugaagc uggagaagga caacgcgaug gaucgcgccc  
uucucugcga acagcaggcc cgcgacgcca acauccgggc cgagaaggcu gaggaggagg  
cccgaucccu gcagaagaag auccagcaga uugagaaua ucuugaucag accauggagc  
aguugaugca agucaacgcc aagcuggacg agaaggacaa ggccugcag aaugcugaga  
gugaggucgc ugcccuaac cgccgaaucc aacugcugga ggaggauuu gagaggucug  
aggaaacguu ggccacagcc accgccaagu uggcugaggc uucccaggcu gccgaugagu  
cagagcgagc ucguaaagau cuugaaucca aaggccuggc agaugaagag cguauggau  
cuuuggagaa ccagcugaag gaagccaggu ucauggcuga ggaagcugac aagaaauaug  
augaggucgc acguaauguu gcuaugguug aggccgacu ggaagagca gaagagcgug  
ccgagacugg ugaauccaag auuguggagc uugaggaaga acugcgcuu gucggcaaca  
accugaaguc ccuugaggug ucugaagaga aggccaaacu gcugaggaa gaguacaagc  
aacagauuaa gacucugaau accaggcuua agggagcuga agcucgugcu gaguucgug  
aaagaucgu gcagaaauug cagaaggagg uugacaggcu ugaggauaa uugguacag  
agaaggagaa guacaaguac auuugugacg aucuugauau g

SEQ ID NO 49; DNA; *Blattella germanica*;

gctctggagctcatattcccttcgcagtatgtgatcaggtggacctcgaggtctacacaatgtttctgcaggaaagtacacggtg  
gggttgggacaggctcgcatgggttctgcacggacaggaggacatcaactctgtgtctaccgtcgctcagtcgactgatgg  
aacgatggagcatcccctactcgcaaatgtggcgccctggaagtaggcaccgagaccctctgg

---

acaagtgaagagcgtcaagactgtcctgatgcaactcttcaaggacaacacggacatcgaggcggtggataccgtgaacgc  
ctgttacgggggacctcggtctcttcaatgcatcttctgtgggtggagtcagctcctgggatggcaggtatgctcttgggtcgtg  
gggacattgctgtgatgctaaaggcagtgcgaggcccacccggtggagcaggggctgtggccatgctagtgggcgccaatgct  
cccctagtgttcgacagaggaggttcgttcacacatgcaacatgcttatgacttctacaaaccggaatctgctcctcgctgtaccca  
ccgtggatggcaagctgtcaattcaatgctatcttagtgcccttagatcattgttatcaactgtactgctccaagatccagaaacaact  
ggagagaagttcgatattgagcggctggatgcagttctctccacgcgcctattgtaagttgggtgcagaagctcttgcctcgtg  
cttgaacgactttgtcgggcatcagaggaggagcggacgactaaatactccagctcgtgaagcactaaaaggcgtgaagcta  
gaagatacgtacttcgaccgagaagttgagaaagcagtcacacacagcaagaacatgtttgaagagaaaacaaagccct  
cgctgttgcctcgccaaccaagtcggcaacatgtacactccttcgctttacggaggtttggtctctattggcagcaagagcgcccc  
ggagttggcaggggaagcgcgtggcctgtttcttaccggtcggactggcctcttccatgttctcttaagaatatcatcgagcgc  
agcgcgaaatcttctctgcaacgcctcgtctcgaatctctgcacatcaagccgcagctggatctgcgccacaaggtgtcaccag  
aggagtttcacaaacgatggagacgaggggaacacaaccaccacaaagctccatacaccccagagggtcgtatcgacgtct  
tgtttccaggaacttggtatctggagagcgtggacagcc

SEQ ID NO 50; PRT; *Blattella germanica*;

Ala Leu Glu Leu Ile Phe Pro Ser Gln Tyr Val Asp Gln Val Asp Leu Glu Val Tyr Asp Asn Val  
Ser Ala Gly Lys Tyr Thr Val Gly Leu Gly Gln Ala Arg Met Gly Phe Cys Thr Asp Arg Glu Asp  
Ile Asn Ser Leu Cys Leu Thr Val Val Ser Arg Leu Met Glu Arg Trp Ser Ile Pro Tyr Ser Gln Ile  
Gly Arg Leu Glu Val Gly Thr Glu Thr Leu Leu Asp Lys Ser Lys Ser Val Lys Thr Val Leu Met  
Gln Leu Phe Lys Asp Asn Thr Asp Ile Glu Gly Val Asp Thr Val Asn Ala Cys Tyr Gly Gly Thr  
Ser Ala Leu Phe Asn Ala Ile Ser Trp Val Glu Ser Ser Ser Trp Asp Gly Arg Tyr Ala Leu Val  
Val Ala Gly Asp Ile Ala Val Tyr Ala Lys Gly Ser Ala Arg Pro Thr Gly Gly Ala Gly Ala Val Ala  
Met Leu Val Gly Ala Asn Ala Pro Leu Val Phe Asp Arg Gly Val Arg Ser Ser His Met Gln His  
Ala Tyr Asp Phe Tyr Lys Pro Asp Leu Ser Ser Leu Tyr Pro Thr Val Asp Gly Lys Leu Ser Ile  
Gln Cys Tyr Leu Ser Ala Leu Asp His Cys Tyr Gln Leu Tyr Cys Ser Lys Ile Gln Lys Gln Leu  
Gly Glu Lys Phe Asp Ile Glu Arg Leu Asp Ala Val Leu Phe His Ala Pro Tyr Cys Lys Leu Val  
Gln Lys Ser Leu Ala Arg Leu Val Leu Asn Asp Phe Val Arg Ala Ser Glu Glu Glu Arg Thr Thr  
Lys Tyr Ser Ser Leu Glu Ala Leu Lys Gly Val Lys Leu Glu Asp Thr Tyr Phe Asp Arg Glu Val  
Glu Lys Ala Val Met Thr Tyr Ser Lys Asn Met Phe Glu Glu Lys Thr Lys Pro Ser Leu Leu  
Ala Asn Gln Val Gly Asn Met Tyr Thr Pro Ser Leu Tyr Gly Gly Leu Val Ser Leu Leu Val Ser  
Lys Ser Ala Gln Glu Leu Ala Gly Lys Arg Val Ala Leu Phe Ser Tyr Gly Ser Gly Leu Ala Ser  
Ser Met Phe Ser Leu Arg Ile Ser Ser Asp Ala Ser Ala Glu Ser Pro Leu Gln Arg Leu Val Ser  
Asn Leu Ser His Ile Lys Pro Gln Leu Asp Leu Arg His Lys Val Ser Pro Glu Glu Phe Ala Gln  
Thr Met Glu Thr Arg Glu His Asn His His Lys Ala Pro Tyr Thr Pro Glu Gly Ser Ile Asp Val  
Leu Phe Pro Gly Thr Trp Tyr Leu Glu Ser Val Asp Ser

SEQ ID NO 51; DNA; *Blattella germanica*; specific primer forward

gctctggagc tcatattccc ttgc

SEQ ID NO 52; DNA; *Blattella germanica*; specific primer reverse

ggctgtccac gctctccaga tacc

SEQ ID NO 53; DNA; specific T7 primer forward

cgctaatacg actcactata gggctctgga gctcatattc ccttc

SEQ ID NO 54; DNA; specific T7 primer reverse

11 / 13

cgctaatacg actcactata ggggctgtcc acgctctcca g

SEQ ID NO 55; RNA; *Blattella germanica*; u can be t (for DNA complement)

gcucuggagc ucauauuccc uucgcaguau guggaucagg uggaccucga ggucucagac  
 aauguuucug caggaaagua cacggugggg ugggacagg cucgcauggg guucugcacg  
 gacagggagg acaucaacuc ucugugucuc accgucguca gucgacugau ggaacgaugg  
 agcauccccu acucgcaau ugggcgccug gaaguaggca cggagaccu ucuggacaag  
 ucgaagagcg ucaagacugu ccugaugcaa cucuuaagg acaacacgga caucgagggc  
 guggauaccg ugaacgccug uuacgggggc accucggcuc ucuucaaugc gauuucgugg  
 guggagucca gcuccuggga uggcagguau gcucuugugg ucgcugggga cauugcugug  
 uaugcuaaag gcagugcgag gccaccggg ggagcagggg cuguggccau gcuagugggc  
 gccaaugcuc ccuaguguu cgacagagga guucguucau cacacaugca acaugcuau  
 gacuucuaca aaccggauuc guccucgcug uacccaccg uggauaggca gcugucauu  
 caaugcuauc uuagugccu agaucauugu uaucaacugu acugcucaa gaucagaaa  
 caacuuggag agaaguucga uauugagcgg cuggaugcag uucucuucca cgcgccuuau  
 uguaaguugg ugcagaaguc ucuugcucgc cucgucuuga acgacuuugu gcgggcauca  
 gaggaggagc ggacgacuaa auacuccagu cuggaagcac uaaaaggcgu gaagcuagaa  
 gauacguacu ucgaccgaga aguugagaaa gcagucauga cauacagcaa gaacauuuu  
 gaagagaaaa caaagcccuc gcuguugcuc gccaaccaag ucggaacau guacacuccu  
 ucgcuuacg gagguuuggu cucucuauug gucagcaaga gcgccagga guuggcaggg  
 aagcgcgugg ccuuguuuuc uuacggcucc ggacuggccu cuuccauguu cucucuaga  
 auaucaucgg acgccagcgc gaaucuuuc cugcaacgcc ucgucucgaa ucucucgcac  
 aucaagccgc agcuggauuc gcgccacaag gugucaccag aggaguuugc acaaacgaug  
 gagacgaggg aacacaacca ccacaaagcu ccuacaccc cagagggcuc gaucgacguc  
 uuguuuccag gaacuuggua ucuggagagc guggacagcc

SEQ ID NO 56; RNA; *Blattella germanica*; u can be t (for DNA complement)

gcucuggagc ucauauuccc uucgcaguau guggaucagg uggaccucga ggucucagac  
 aauguuucug caggaaagua cacggugggg ugggacagg cucgcauggg guucugcacg  
 gacagggagg acaucaacuc ucugugucuc accgucguca gucgacugau ggaacgaugg  
 agcauccccu acucgcaau ugggcgccug gaaguaggca cggagaccu ucuggacaag  
 ucgaagagcg ucaagacugu ccugaugcaa cucuuaagg acaacacgga caucgagggc  
 guggauaccg ugaacgccug uuacgggggc accucggcuc ucuucaaugc gauuucgugg  
 guggagucca gcuccuggga uggcagguau gcucuugugg ucgcugggga cauugcugug  
 uaugcuaaag gcagugcgag gccaccggg ggagcagggg cuguggccau gcuagugggc  
 gccaaugcuc ccuaguguu cgacagagga guucguucau cacacaugca acaugcuau  
 gacuucuaca aaccggauuc guccucgcug uacccaccg uggauaggca gcugucauu  
 caaugcuauc uuagugccu agaucauugu uaucaacugu acugcucaa gaucagaaa  
 caacuuggag agaaguucga uauugagcgg cuggaugcag uucucuucca cgcgccuuau  
 uguaaguugg ugcagaaguc ucuugcucgc cucgucuuga acgacuuugu gcgggcauca  
 gaggaggagc ggacgacuaa auacuccagu cuggaagcac uaaaaggcgu gaagcuagaa  
 gauacguacu ucgaccgaga aguugagaaa gcagucauga cauacagcaa gaacauuuu  
 gaagagaaaa caaagcccuc gcuguugcuc gccaaccaag ucggaacau guacacuccu  
 ucgcuuacg gagguuuggu cucucuauug gucagcaaga gcgccagga guuggcaggg  
 aagcgcgugg ccuuguuuuc uuacggcucc ggacuggccu cuuccauguu cucucuaga  
 auaucaucgg acgccagcgc gaaucuuuc cugcaacgcc ucgucucgaa ucucucgcac  
 aucaagccgc agcuggauuc gcgccacaag gugucaccag aggaguuugc acaaacgaug  
 gagacgaggg aacacaacca ccacaaagcu ccuacaccc cagagggcuc gaucgacguc  
 uuguuuccag gaacuuggu

SEQ ID NO 57; DNA; *Blattella germanica*;

FIG. 5 CONT'D

12 / 13

---

gaggcccagagcaagagaggtatcctcactctgaagtacccattgaacatggaatcatcaccaactgggatgacatggaga  
 agatctggcatcacaccttctacaatgaactccgagtggctccagaggaacacccaatcctgctgactgaggctcccctgaacc  
 caaaggccaacagggagaagatgactcaaatactgttgagaccttaacacccccgcatgtatgttgccatccaggccgtgc  
 tgtccctctacgcttccggccgtaccactggtattgtgctggactctggtgacggcgtctccacaccgtaccatctatgaaggta  
 cgcatgcccatccatcctgcgtctggactggccggccgtgactgactgactacctgatgaagatcctgaccgagcgtggct  
 acagcttcacaactacagcagagcgag

SEQ ID NO 58; PRT; *Blattella germanica*;

Glu Ala Gln Ser Lys Arg Gly Ile Leu Thr Leu Lys Tyr Pro Ile Glu His Gly Ile Ile Thr Asn Trp  
 Asp Asp Met Glu Lys Ile Trp His His Thr Phe Tyr Asn Glu Leu Arg Val Ala Pro Glu Glu His  
 Pro Ile Leu Leu Thr Glu Ala Pro Leu Asn Pro Lys Ala Asn Arg Glu Lys Met Thr Gln Ile Met  
 Phe Glu Thr Phe Asn Thr Pro Ala Met Tyr Val Ala Ile Gln Ala Val Leu Ser Leu Tyr Ala Ser  
 Gly Arg Thr Thr Gly Ile Val Leu Asp Ser Gly Asp Gly Val Ser His Thr Val Pro Ile Tyr Glu Gly  
 Tyr Ala Leu Pro His Ala Ile Leu Arg Leu Asp Leu Ala Gly Arg Asp Leu Thr Asp Tyr Leu Met  
 Lys Ile Leu Thr Glu Arg Gly Tyr Ser Phe Thr Thr Thr Ala Glu Arg

SEQ ID NO 59; DNA; *Blattella germanica*; degenerative primer forward

gaggcccaga gcaagagagg tatcc

SEQ ID NO 60; DNA; *Blattella germanica*; degenerative primer reverse

ctcgctctgc ttagttgtg aagctg

SEQ ID NO 61; DNA; specific T7 primer forward

cgctaatacg actcactata gggaggccca gagcaagaga gg

SEQ ID NO 62; DNA; specific T7 primer reverse

cgctaatacg actcactata ggtctgctgt agttgtgaag ctgtagcc

SEQ ID NO 63; RNA; *Blattella germanica*; u can be t (for DNA complement)

gaggcccaga gcaagagagg uauccucacu cugaaguacc ccuugaaca uggaaucauc  
 accaacuggg augacaugga gaagaucugg caucacaccu ucuacaauga acuccgagug  
 gcuccagagg aacacccaau ccugcugacu gaggcucccc ugaacccaaa ggccaacagg  
 gagaagauga cucaaaucuu guuugagacc uucaacacccc ccgccaugua uguugccauc  
 caggccgugc uguccucua cgcuuccggc cguaccacug guauugugcu ggacucuggu  
 gacggcgucu cccacaccgu acccaucuaa gaagguuacg cauugcccca ugccaucug  
 cgucuggacu uggccggccg ugacuugacu gacuaccuga ugaagaucuu gaccgagcgu  
 ggcuaacgu ucacaacuac agcaga

SEQ ID NO 64; RNA; *Blattella germanica*; u can be t (for DNA complement)

uggcaucaca ccuucucaa ugaacuccga guggcuccag aggaacaccc aaucucugcug  
 acugaggcuc ccgugaaccc aaaggccaac agggagaaga ugacuaa

SEQ ID NO 65; DNA; synthetic Bg001 fragment

tatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcc

---

FIG. 5 CONT'D



SEQ ID NO 66; DNA; synthetic Bg001 fragment

cttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcct

SEQ ID NO 67; DNA; synthetic Bg001 concatemer 1

tatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcctatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcctatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcctatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcctatgctccaagaccaagcacaggacctcacaagttacgagagagtctgcc

SEQ ID NO 68; DNA; synthetic Bg001 concatemer 2

cttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcctcttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcctcttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcctcttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcctcttgggtcgtgttggaactgtagttaatcgagaacgtcatcctggtcct

SEQ ID NO 69; DNA; synthetic Bg001 fragment

gctgaaatatgcattaaccaactgtgaggttaagaaaattgtatgcagcgccttattaagggtgatggaaaagtcagaaacagacc  
ccaactatccagct

SEQ ID NO 70; DNA; synthetic Bg001 concatemer 3

gctgaaatatgcattaaccaactgtgaggttaagaaaattgtatgcagcgccttattaagggtgatggaaaagtcagaaacagacc  
ccaactatccagctgctgaaatatgcattaaccaactgtgaggttaagaaaattgtatgcagcgccttattaagggtgatggaaa  
gtcagaaacagaccccaactatccagctgctgaaatatgcattaaccaactgtgaggttaagaaaattgtatgcagcgccttatta  
agggtgatggaaaagtcagaaacagaccccaactatccagct

FIG. 5 CONT'D

## SEQUENCE LISTING

&lt;110&gt; Devgen NV

&lt;120&gt; RNAi for the control of insects and arachnids

&lt;130&gt; P79369WO00

&lt;160&gt; 200

&lt;170&gt; PatentIn version 3.3

&lt;210&gt; 1

&lt;211&gt; 624

&lt;212&gt; DNA

&lt;213&gt; Blattella germanica

&lt;400&gt; 1

```

taaggcatgg atgttgga caagcggtgg agtgtatgct ccaagaccaa gcacaggacc      60
tcacaagtta cgagagagtc tgccccttgt aatatttctt cgtaataggc tgaaatatgc    120
attaaccaac tgtgagggtta agaaaattgt tatgcagcgc cttattaagg ttgatggaaa    180
agtcagaaca gaccccaact atccagctgg ttttatggat gttgttaca ttgaaaaaac    240
tggaagaattt ttccgtctga tttatgacgt gaaaggacgt ttcaccattc acagaataac    300
tgctgaagaa gccaagtata aactgtgcaa ggtaaagaga gtgcagactg ggccaagggt    360
tattccattc ttggtgacct atgatggtag aactcttaga tatcctgatc ctgtcatcaa    420
agttaatgat acagttcaac ttgacatcgc tacttccaag attatggata gcatcaaatt    480
tgataatggt aatctctgta tgattactgg aggccgtaac ttgggtcgtg ttggaactgt    540
agttaatcga gaacgtcatc ctggttcctt tgacattgtg catgttaaag attcacaagg    600
acacacattt gctaccagat tgaa                                           624

```

&lt;210&gt; 2

&lt;211&gt; 208

&lt;212&gt; PRT

&lt;213&gt; Blattella germanica

&lt;400&gt; 2

```

Lys Ala Trp Met Leu Asp Lys Leu Gly Gly Val Tyr Ala Pro Arg Pro
1           5           10          15

```

```

Ser Thr Gly Pro His Lys Leu Arg Glu Ser Leu Pro Leu Val Ile Phe
          20          25          30

```

```

Leu Arg Asn Arg Leu Lys Tyr Ala Leu Thr Asn Cys Glu Val Lys Lys
          35          40          45

```

```

Ile Val Met Gln Arg Leu Ile Lys Val Asp Gly Lys Val Arg Thr Asp

```

50                                      55                                      60  
 Pro Asn Tyr Pro Ala Gly Phe Met Asp Val Val Thr Ile Glu Lys Thr  
 65                                      70                                      75                                      80  
 Gly Glu Phe Phe Arg Leu Ile Tyr Asp Val Lys Gly Arg Phe Thr Ile  
                                     85                                      90                                      95  
 His Arg Ile Thr Ala Glu Glu Ala Lys Tyr Lys Leu Cys Lys Val Lys  
                                     100                                      105                                      110  
 Arg Val Gln Thr Gly Pro Lys Gly Ile Pro Phe Leu Val Thr His Asp  
                                     115                                      120                                      125  
 Gly Arg Thr Leu Arg Tyr Pro Asp Pro Val Ile Lys Val Asn Asp Thr  
                                     130                                      135                                      140  
 Val Gln Leu Asp Ile Ala Thr Ser Lys Ile Met Asp Ser Ile Lys Phe  
 145                                      150                                      155                                      160  
 Asp Asn Gly Asn Leu Cys Met Ile Thr Gly Gly Arg Asn Leu Gly Arg  
                                     165                                      170                                      175  
 Val Gly Thr Val Val Asn Arg Glu Arg His Pro Gly Ser Phe Asp Ile  
                                     180                                      185                                      190  
 Val His Val Lys Asp Ser Gln Gly His Thr Phe Ala Thr Arg Leu Asn  
                                     195                                      200                                      205

<210> 3  
 <211> 23  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Degenerate primer

<400> 3  
 catttgaagc gtttwrmygc ycc

23

<210> 4  
 <211> 27  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer

<400> 4  
 gtgcccttg caatgatgaa cacgttg

27

<210> 5  
 <211> 45  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific T7 primer

<400> 5  
 cgctaatacgc actcactata ggggagtgtg tgctccaaga ccaag

45

<210> 6  
 <211> 24  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer

<400> 6  
 caatctggta gcaaatgtgt gtcc

24

<210> 7  
 <211> 23  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer

<400> 7  
 ggagtgtatg ctccaagacc aag

23

<210> 8  
 <211> 46  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Synthetic T7 primer

<400> 8  
 cgctaatacgc actcactata ggcaatctgg tagcaaatgt gtgtcc

46

<210> 9  
 <211> 594  
 <212> RNA  
 <213> *Blattella germanica*

<400> 9  
 ggaguguaug cuccaagacc aagcacagga ccucacaagu uacgagagag ucugcccccuu

60

guaauuuuuc uucguaauag gcugaaaauau gcauuuaccca acugugaggu uaagaaaauu

120

guuaugcagc gccuuuuuaa gguugaugga aaagucagaa cagaccccaa cuauccagcu

180

gguuuuaugg auguuguuac aaugaaaaa acuggagaau uuuccgucu gauuuauagac 240  
 gugaaaggac guuucaccau ucacagaaua acugcugaag aagccaagua uaaacugugc 300  
 aagguaaaga gagugcagac ugggcccaag gguauuccau ucuuggugac ccaugauggu 360  
 agaacucuaa gauauccuga uccugucauc aaaguuaaug auacaguuca acuugacauc 420  
 gcuacuucca agauuaugga uagcaucaa uuugauaaug gaaucucug uaugauuacu 480  
 ggaggccgua acuugggucg uguuggaacu guaguuauc gagaacguca uccugguucc 540  
 uuugacauug ugcauguuaa agauucacaa ggacacacau uugcuaccag auug 594

&lt;210&gt; 10

&lt;211&gt; 573

&lt;212&gt; RNA

<213> *Blattella germanica*

&lt;400&gt; 10

ggaguguauug cuccaagacc aagcacagga ccucacaagu uacgagagag ucugcccuu 60  
 guaaauuuuc uucguauuag gcugaaauau gcauaacca acugugaggu uaagaaaauu 120  
 guuaugcagc gccuuauuaa gguugaugga aaagucagaa cagaccccaa cuauccagcu 180  
 gguuuuaugg auguuguuac aaugaaaaa acuggagaau uuuccgucu gauuuauagac 240  
 gugaaaggac guuucaccau ucacagaaua acugcugaag aagccaagua uaaacugugc 300  
 aagguaaaga gagugcagac ugggcccaag gguauuccau ucuuggugac ccaugauggu 360  
 agaacucuaa gauauccuga uccugucauc aaaguuaaug auacaguuca acuugacauc 420  
 gcuacuucca agauuaugga uagcaucaa uuugauaaug gaaucucug uaugauuacu 480  
 ggaggccgua acuugggucg uguuggaacu guaguuauc gagaacguca uccugguucc 540  
 uuugacauug ugcauguuaa agauucacaa gga 573

&lt;210&gt; 11

&lt;211&gt; 436

&lt;212&gt; DNA

<213> *Blattella germanica*

&lt;400&gt; 11

tcccaggcga cttatgaaa aggcacgtct tgatcaggag ttgaaaatca taggagaata 60  
 tggctcttagg aacaaacgtg aagtgtggcg agtcaagtat accttgga aaatccgtaa 120  
 agctgccaga gaacttctga ctttgaaga gaaagatcag cgcaggttgt ttgaaggcaa 180  
 tgctcttctt cgtcggttgg tgcgtattgg agtgttgat gaaacccgta tgaagcttga 240  
 ttacgtcttg ggtttgaaga ttgaagattt cttggaacga cgtctccaaa cacaagtttt 300  
 caagttgggg cttgcaaaat caatccatca tgctcgtgtg ctgatccgtc aaagacatat 360

cagggttcgt aagcaggtcg tgaatattcc aagcttcatt gtgagacttg attcccagaa 420

gcatattgac ttctcg 436

<210> 12

<211> 145

<212> PRT

<213> *Blattella germanica*

<400> 12

Pro Arg Arg Pro Tyr Glu Lys Ala Arg Leu Asp Gln Glu Leu Lys Ile  
1 5 10 15

Ile Gly Glu Tyr Gly Leu Arg Asn Lys Arg Glu Val Trp Arg Val Lys  
20 25 30

Tyr Thr Leu Ala Lys Ile Arg Lys Ala Ala Arg Glu Leu Leu Thr Leu  
35 40 45

Glu Glu Lys Asp Gln Arg Arg Leu Phe Glu Gly Asn Ala Leu Leu Arg  
50 55 60

Arg Leu Val Arg Ile Gly Val Leu Asp Glu Thr Arg Met Lys Leu Asp  
65 70 75 80

Tyr Val Leu Gly Leu Lys Ile Glu Asp Phe Leu Glu Arg Arg Leu Gln  
85 90 95

Thr Gln Val Phe Lys Leu Gly Leu Ala Lys Ser Ile His His Ala Arg  
100 105 110

Val Leu Ile Arg Gln Arg His Ile Arg Val Arg Lys Gln Val Val Asn  
115 120 125

Ile Pro Ser Phe Ile Val Arg Leu Asp Ser Gln Lys His Ile Asp Phe  
130 135 140

Ser

145

<210> 13

<211> 26

<212> DNA

<213> Artificial sequence

<220>

<223> degenerate primer

<220>  
<221> misc\_feature  
<222> (18)..(18)  
<223> n = a, c, g or t

<400> 13  
tcggtcttct cgaagacnta ygtkac 26

<210> 14  
<211> 20  
<212> DNA  
<213> Artificial sequence

<220>  
<223> degenerative primer

<400> 14  
ccgccgaagg gmgayttbag 20

<210> 15  
<211> 43  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer

<400> 15  
cgctaatacg actcactata ggcaggcgac cttatgaaaa ggc 43

<210> 16  
<211> 23  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer

<400> 16  
cgagaagtca atatgcttct ggg 23

<210> 17  
<211> 21  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer

<400> 17  
caggcgacct tatgaaaagg c 21

<210> 18  
<211> 45  
<212> DNA  
<213> Artificial sequence

&lt;220&gt;

&lt;223&gt; specific T7 primer

&lt;400&gt; 18

cgctaatacg actcactata ggcgagaagt caatatgctt ctggg 45

&lt;210&gt; 19

&lt;211&gt; 433

&lt;212&gt; RNA

<213> *Blattella germanica*

&lt;400&gt; 19

caggcgaccu uaugaaaagg cacgucuuga ucaggaguug aaaaucauag gagaauaugg 60

ucuuaggaac aaacgugaag uguggcgagu caaguauacc uuggcaaaaa uccguaaaagc 120

ugccagagaa cuucugacuu uggaagagaa agaucagcgc agguuguuug aaggcaaugc 180

ucuucucgu cgguuggugc guauuggagu guuggaugaa acccguauga agcuugauua 240

cgucuugggu uugaagauug aagauuucuu ggaacgacgu cuccaaacac aaguuuucuaa 300

guuggggcuu gcaaaaucuaa uccaucaugc ucgugugcug auccgucaaa gacauaucag 360

gguucguaag caggucguga auauuccaag cuucauugug agacuugauu cccagaagca 420

uauugacuuc ucg 433

&lt;210&gt; 20

&lt;211&gt; 412

&lt;212&gt; RNA

<213> *Blattella germanica*

&lt;400&gt; 20

caggcgaccu uaugaaaagg cacgucuuga ucaggaguug aaaaucauag gagaauaugg 60

ucuuaggaac aaacgugaag uguggcgagu caaguauacc uuggcaaaaa uccguaaaagc 120

ugccagagaa cuucugacuu uggaagagaa agaucagcgc agguuguuug aaggcaaugc 180

ucuucucgu cgguuggugc guauuggagu guuggaugaa acccguauga agcuugauua 240

cgucuugggu uugaagauug aagauuucuu ggaacgacgu cuccaaacac aaguuuucuaa 300

guuggggcuu gcaaaaucuaa uccaucaugc ucgugugcug auccgucaaa gacauaucag 360

gguucguaag caggucguga auauuccaag cuucauugug agacuugauu cc 412

&lt;210&gt; 21

&lt;211&gt; 448

&lt;212&gt; DNA

<213> *Blattella germanica*

&lt;400&gt; 21

tgtgaaagga ccacgaggca ccttgaagcg cggtttcaag catcttgctt tagatatcca 60

cttggttcat ccaaggctcc tgaaggtgga aaaatggttt ggaacaaaga aggagtggc 120



```

agccgtgctgc accgtctgct ctcattattga gaacatgatt aaaggagtca caaaggggttt      180
cctgtacaaa atgcgcgccg tgtatgccca tttccccatt aactgcgtaa ccacagaaaa      240
caattccgtt attgaagtgc gtaacttctt gggcgagaag ttcatccgca gagtgaagat      300
ggctccggga gtgaccgtca ccaattctcc aaagcagaaa gacgagctca ttctggaggg      360
caacgacatc gaggatgtat cgagatcagc cgcactcatc caacaatcga cgactgtgaa      420
gaacaaggac atccggaaat tccttgac                                          448

```

&lt;210&gt; 22

&lt;211&gt; 149

&lt;212&gt; PRT

<213> *Blattella germanica*

&lt;400&gt; 22

```

Val Lys Gly Pro Arg Gly Thr Leu Lys Arg Gly Phe Lys His Leu Ala
1              5              10              15

```

```

Leu Asp Ile His Leu Val His Pro Arg Leu Leu Lys Val Glu Lys Trp
              20              25              30

```

```

Phe Gly Thr Lys Lys Glu Leu Ala Ala Val Arg Thr Val Cys Ser His
              35              40              45

```

```

Ile Glu Asn Met Ile Lys Gly Val Thr Lys Gly Phe Leu Tyr Lys Met
50              55              60

```

```

Arg Ala Val Tyr Ala His Phe Pro Ile Asn Cys Val Thr Thr Glu Asn
65              70              75              80

```

```

Asn Ser Val Ile Glu Val Arg Asn Phe Leu Gly Glu Lys Phe Ile Arg
              85              90              95

```

```

Arg Val Lys Met Ala Pro Gly Val Thr Val Thr Asn Ser Pro Lys Gln
              100              105              110

```

```

Lys Asp Glu Leu Ile Leu Glu Gly Asn Asp Ile Glu Asp Val Ser Arg
              115              120              125

```

```

Ser Ala Ala Leu Ile Gln Gln Ser Thr Thr Val Lys Asn Lys Asp Ile
              130              135              140

```

```

Arg Lys Phe Leu Asp
145

```

<210> 23  
<211> 20  
<212> DNA  
<213> Artificial sequence

<220>  
<223> forward degenerative primer

<220>  
<221> misc\_feature  
<222> (12)..(13)  
<223> n = a, c, g or t

<400> 23  
gtgaaggccc gnntggtgac 20

<210> 24  
<211> 26  
<212> DNA  
<213> Artificial sequence

<220>  
<223> reverse degenerative primer

<400> 24  
gtcgtcttct cdgahacrta vagacc 26

<210> 25  
<211> 43  
<212> DNA  
<213> Artificial sequence

<220>  
<223> T7 primer forward

<400> 25  
cgctaatacg actcactata gggtgaaagg accacgaggc acc 43

<210> 26  
<211> 21  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer reverse

<400> 26  
ccgtcaagga atttccggat g 21

<210> 27  
<211> 21  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer forward

<400> 27  
gtgaaaggac cacgaggcac c 21

<210> 28  
<211> 43  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer reverse

<400> 28  
cgctaatacg actcactata ggccgtcaag gaatttccgg atg 43

<210> 29  
<211> 449  
<212> RNA  
<213> Blattella germanica

<400> 29  
gugaaaggac cacgaggcac cuugaagcgc gguuucaggc aucuugcuuu agauauccac 60  
uugguucauc caaggcuccu gaagguggaa aaaugguuug gaacaaagaa ggaguuggca 120  
gccgugcgca ccgucugcuc ucauauugag aacaugauua aaggagucac aaaggguuuc 180  
cuguacaaaa ugcgcgccgu guauggccau uuccccauua acugcguaac cacagaaaac 240  
aaauccguua uugaagugcg uaacuucuug ggcgagaagu ucauccgcag agugaagaug 300  
gcuccggggag ugaccgucac caauucucca aagcagaaaag acgagcucac ucuggagggc 360  
aacgacaucg aggauguauc gagaucagcc gcacucaucc aacaucgac gacugugaag 420  
aacaaggaca uccggaaaau ccuugacgg 449

<210> 30  
<211> 428  
<212> RNA  
<213> Blattella germanica

<400> 30  
gugaaaggac cacgaggcac cuugaagcgc gguuucaggc aucuugcuuu agauauccac 60  
uugguucauc caaggcuccu gaagguggaa aaaugguuug gaacaaagaa ggaguuggca 120  
gccgugcgca ccgucugcuc ucauauugag aacaugauua aaggagucac aaaggguuuc 180  
cuguacaaaa ugcgcgccgu guauggccau uuccccauua acugcguaac cacagaaaac 240  
aaauccguua uugaagugcg uaacuucuug ggcgagaagu ucauccgcag agugaagaug 300  
gcuccggggag ugaccgucac caauucucca aagcagaaaag acgagcucac ucuggagggc 360  
aacgacaucg aggauguauc gagaucagcc gcacucaucc aacaucgac gacugugaag 420

aacaagga

428

&lt;210&gt; 31

&lt;211&gt; 404

&lt;212&gt; DNA

&lt;213&gt; Blattella germanica

&lt;400&gt; 31

ggcttgatcc caatgaaata aacgaaattg caaataccaa ttccagacaa aatattcgta 60

aactgattaa agatgggtctt atcatcaaga agcccgtagc tgtacactca agggcccgtg 120

ttcgcaagaa caccgaagca agaagaaaag gacgtcactg cggttttggc aaaaggaagg 180

gtacggcaaa tgcccgtatg ccacagaagg tcttgtggat taatcgcacg cgtgttctga 240

gaaggcttct caagaagtac agggaagcaa agaagatcga cagacatcta taccaccagc 300

tgtacatgaa ggccaagggg aacgtgttca agaacaagcg tgtcctgatg gagttcatcc 360

acaagaagaa ggctgagaag gccaggacaa agatgcttaa cgac 404

&lt;210&gt; 32

&lt;211&gt; 134

&lt;212&gt; PRT

&lt;213&gt; Blattella germanica

&lt;400&gt; 32

Leu Asp Pro Asn Glu Ile Asn Glu Ile Ala Asn Thr Asn Ser Arg Gln  
 1 5 10 15

Asn Ile Arg Lys Leu Ile Lys Asp Gly Leu Ile Ile Lys Lys Pro Val  
 20 25 30

Ala Val His Ser Arg Ala Arg Val Arg Lys Asn Thr Glu Ala Arg Arg  
 35 40 45

Lys Gly Arg His Cys Gly Phe Gly Lys Arg Lys Gly Thr Ala Asn Ala  
 50 55 60

Arg Met Pro Gln Lys Val Leu Trp Ile Asn Arg Met Arg Val Leu Arg  
 65 70 75 80

Arg Leu Leu Lys Lys Tyr Arg Glu Ala Lys Lys Ile Asp Arg His Leu  
 85 90 95

Tyr His Gln Leu Tyr Met Lys Ala Lys Gly Asn Val Phe Lys Asn Lys  
 100 105 110

Arg Val Leu Met Glu Phe Ile His Lys Lys Lys Ala Glu Lys Ala Arg  
 115 120 125

Thr Lys Met Leu Asn Asp  
130

<210> 33  
<211> 26  
<212> DNA  
<213> Artificial sequence

<220>  
<223> degenerative primer forward

<400> 33  
tgcgatgcgg caaraaraag gtbttg 26

<210> 34  
<211> 19  
<212> DNA  
<213> Artificial sequence

<220>  
<223> degenerative primer reverse

<400> 34  
gtcggcgagc ytcrgcytg 19

<210> 35  
<211> 46  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer forward

<400> 35  
cgctaatacg actcactata ggggcttgat cccaatgaaa taaacg 46

<210> 36  
<211> 24  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer reverse

<400> 36  
gtcgttaagc atctttgtcc tggc 24

<210> 37  
<211> 24  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific primer forward

<400> 37  
ggcttgatcc caatgaaata aacg 24

<210> 38  
<211> 46  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer reverse

<400> 38  
cgctaatacg actcactata gggtcgttaa gcatctttgt cctggc 46

<210> 39  
<211> 404  
<212> RNA  
<213> Blattella germanica

<400> 39  
ggcuugaucc caaugaaaua aacgaaauug caaaauaccaa uuccagacaa aaauuucgua 60  
aacugauuaa agauggucuu aucaucaaga agcccguagc uguacacuca agggcccug 120  
uucgcaagaa caccgaagca agaagaaaag gacgucacug cgguuuuggc aaaaggaagg 180  
guacggcaaa ugcccguaug ccacagaagg ucuuguggau uaaucgcaug cguguucuga 240  
gaaggcuucu caagaaguac aggggaagcaa agaagaucga cagacaucua uaccaccagc 300  
uguacaugaa ggccaagggg aacguguuca agaacaagcg uguccugaug gaguucaucc 360  
acaagaagaa ggcugagaag gccaggacaa agaugcuuaa cgac 404

<210> 40  
<211> 383  
<212> RNA  
<213> Blattella germanica

<400> 40  
ggcuugaucc caaugaaaua aacgaaauug caaaauaccaa uuccagacaa aaauuucgua 60  
aacugauuaa agauggucuu aucaucaaga agcccguagc uguacacuca agggcccug 120  
uucgcaagaa caccgaagca agaagaaaag gacgucacug cgguuuuggc aaaaggaagg 180  
guacggcaaa ugcccguaug ccacagaagg ucuuguggau uaaucgcaug cguguucuga 240  
gaaggcuucu caagaaguac aggggaagcaa agaagaucga cagacaucua uaccaccagc 300  
uguacaugaa ggccaagggg aacguguuca agaacaagcg uguccugaug gaguucaucc 360  
acaagaagaa ggcugagaag gcc 383

<210> 41  
<211> 849

&lt;212&gt; DNA

<213> *Blattella germanica*

&lt;400&gt; 41

```

atggatgcc tcaagaagaa gatgcaggcg atgaagctgg agaaggacaa cgcgatggat      60
cgcgcccttc tctgcgaaca gcaggcccg gacgccaaca tccgggccga gaaggctgag      120
gaggaggccc gatccctgca gaagaagatc cagcagattg agaatgatct tgatcagacc      180
atggagcagt tgatgcaagt caacgccaaag ctggacgaga aggacaaggc cctgcagaat      240
gctgagagtg aggtcgctgc cctcaaccgc cgaatccaac tgctggagga ggatcttgag      300
aggtctgagg aacgtttggc cacagccacc gccaaagtgg ctgaggcttc ccaggctgcc      360
gatgagtcag agcgagctcg taagattctt gaatccaaag gcctggcaga tgaagagcgt      420
atggatgctt tggagaacca gctgaaggaa gccaggttca tggctgagga agctgacaag      480
aaatatgatg aggtcgcacg taagttggct atggttgagg ccgacttgga aagagcagaa      540
gagcgtgccg agactggtga atccaagatt gtggagcttg aggaagaact gcgcgttgct      600
ggcaacaacc tgaagtcctt tgaggtgtct gaagagaagg ccaacctgcg tgaggaagag      660
tacaagcaac agattaagac tctgaatacc aggctaaagg aggctgaagc tcgtgctgag      720
ttcgctgaaa gatccgtgca gaaattgcag aaggaggttg acaggcttga ggatgaattg      780
gtacacgaga aggagaagta caagtacatt tgtgacgata ttgatatgac tttcacccga      840
cttattggc                                     849

```

&lt;210&gt; 42

&lt;211&gt; 283

&lt;212&gt; PRT

<213> *Blattella germanica*

&lt;400&gt; 42

```

Met Asp Ala Ile Lys Lys Lys Met Gln Ala Met Lys Leu Glu Lys Asp
1           5           10           15

Asn Ala Met Asp Arg Ala Leu Leu Cys Glu Gln Gln Ala Arg Asp Ala
          20           25           30

Asn Ile Arg Ala Glu Lys Ala Glu Glu Glu Ala Arg Ser Leu Gln Lys
          35           40           45

Lys Ile Gln Gln Ile Glu Asn Asp Leu Asp Gln Thr Met Glu Gln Leu
          50           55           60

Met Gln Val Asn Ala Lys Leu Asp Glu Lys Asp Lys Ala Leu Gln Asn
65           70           75           80

```

Ala Glu Ser Glu Val Ala Ala Leu Asn Arg Arg Ile Gln Leu Leu Glu  
85 90 95

Glu Asp Leu Glu Arg Ser Glu Glu Arg Leu Ala Thr Ala Thr Ala Lys  
100 105 110

Leu Ala Glu Ala Ser Gln Ala Ala Asp Glu Ser Glu Arg Ala Arg Lys  
115 120 125

Ile Leu Glu Ser Lys Gly Leu Ala Asp Glu Glu Arg Met Asp Ala Leu  
130 135 140

Glu Asn Gln Leu Lys Glu Ala Arg Phe Met Ala Glu Glu Ala Asp Lys  
145 150 155 160

Lys Tyr Asp Glu Val Ala Arg Lys Leu Ala Met Val Glu Ala Asp Leu  
165 170 175

Glu Arg Ala Glu Glu Arg Ala Glu Thr Gly Glu Ser Lys Ile Val Glu  
180 185 190

Leu Glu Glu Glu Leu Arg Val Val Gly Asn Asn Leu Lys Ser Leu Glu  
195 200 205

Val Ser Glu Glu Lys Ala Asn Leu Arg Glu Glu Glu Tyr Lys Gln Gln  
210 215 220

Ile Lys Thr Leu Asn Thr Arg Leu Lys Glu Ala Glu Ala Arg Ala Glu  
225 230 235 240

Phe Ala Glu Arg Ser Val Gln Lys Leu Gln Lys Glu Val Asp Arg Leu  
245 250 255

Glu Asp Glu Leu Val His Glu Lys Glu Lys Tyr Lys Tyr Ile Cys Asp  
260 265 270

Asp Leu Asp Met Thr Phe Thr Glu Leu Ile Gly  
275 280

<210> 43

<211> 27

<212> DNA

<213> Artificial sequence

<220>

<223> specific primer forward



<400> 43  
 atggatgcca tcaagaagaa gatgcag 27

<210> 44  
 <211> 30  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer reverse

<400> 44  
 gccataagt tcggtgaaag tcatatcaag 30

<210> 45  
 <211> 46  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific T7 primer forward

<400> 45  
 cgctaatacg actcactata ggatggatgc catcaagaag aagatg 46

<210> 46  
 <211> 46  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific T7 primer reverse

<400> 46  
 cgctaatacg actcactata gggccaataa gtccggtgaa agtcat 46

<210> 47  
 <211> 849  
 <212> RNA  
 <213> Blattella germanica

<400> 47  
 auggaugcca ucaagaagaa gaugcaggcg augaagcugg agaaggacaa cgcgauaggau 60  
 cgcgcccuuc ucugcgaaca gcaggcccg cagcccaaca uccgggcca gaaggcugag 120  
 gaggaggccc gauccugca gaagaagauc cagcagauug agaaugaucu ugaucagacc 180  
 auggagcagu ugaugcaagu caacgccaag cuggacgaga aggacaaggc ccugcagaau 240  
 gcugagagug agguccgucg ccucaaccgc cgaauccaac ugcuggagga ggaucuuag 300  
 agguccugagg aacguuuggc cacagccacc gccaauguug cugaggcuuc ccaggcugcc 360  
 gaugagucag agcgagcucg uaagauucuu gaaucxaaag gccuggcaga ugaagagcgu 420  
 auggaugcuu uggagaacca gcugaaggaa gccagguuca uggcugagga agcugacaag 480

aaauaugaug agguccgacg uaaguuggcu augguugagg ccgacuugga aagagcagaa 540  
 gagcgugccg agacugguga auccaagauu guggagcuug aggaagaacu gcgcguuguc 600  
 ggcaacaacc ugaagucccu ugaggugucu gaagagaagg ccaaccugcg ugaggaagag 660  
 uacaagcaac agauuaagac ucugaauacc aggcuaaagg aggcugaagc ucgugcugag 720  
 uucgcugaaa gauccgugca gaaauugcag aaggagguug acaggcuuga ggaugaauug 780  
 guacacgaga aggagaagua caaguacauu ugugacgauc uugauaugac uuucaccgaa 840  
 cuuauuggc 849

&lt;210&gt; 48

&lt;211&gt; 821

&lt;212&gt; RNA

&lt;213&gt; Blattella germanica

&lt;400&gt; 48

ccaucaagaa gaagaugcag gcgaugaagc uggagaagga caacgcgaug gaucgcgccc 60  
 uucucugcga acagcaggcc cgcgacgcca acauccgggc cgagaaggcu gaggaggagg 120  
 cccgaucccu gcagaagaag auccagcaga uugagaauga ucuugaucag accauggagc 180  
 aguugaugca agucaacgcc aagcuggacg agaaggacaa ggcccugcag aaugcugaga 240  
 gugaggucgc ugcccucaac cgccgaaucc aacugcugga ggaggauuu gagaggucug 300  
 aggaacguuu ggccacagcc accgccaagu uggcugaggc uucccaggcu gccgaugagu 360  
 cagagcgagc ucguaagauu cuugaaucca aaggccuggc agaugaagag cguauggaug 420  
 cuuuggagaa ccagcugaag gaagccaggu ucauggcuga ggaagcugac aagaaauaug 480  
 augaggucgc acguaaguug gcuaugguug aggccgacuu ggaaagagca gaagagcgug 540  
 ccgagacugg ugaauccaag auuguggagc uugaggaaga acugcgcguu gucggcaaca 600  
 accugaaguc ccuugaggug ucugaagaga aggccaaaccu gcgugaggaa gaguacaagc 660  
 aacagauuaa gacucugaau accaggcuuaa aggaggcuga agcucgugcu gaguucgcug 720  
 aaagauccgu gcagaaauug cagaaggagg uugacaggcu ugaggaukaa uugguacacg 780  
 agaaggagaa guacaaguac auuugugacg aucuugauau g 821

&lt;210&gt; 49

&lt;211&gt; 1300

&lt;212&gt; DNA

&lt;213&gt; Blattella germanica

&lt;400&gt; 49

gctctggagc tcatattccc ttgcagtat gtggatcagg tggacctga ggtctacgac 60  
 aatgtttctg caggaaagta cacggtgggg ttgggacagg ctgcgatggg gttctgcacg 120

gacagggagg acatcaactc tctgtgtctc accgtcgtca gtcgactgat ggaacgatgg 180  
 agcatcccct actcgcaaat tgggcgcctg gaagtaggca ccgagaccct tctggacaag 240  
 tcgaagagcg tcaagactgt cctgatgcaa ctcttcaagg acaacacgga catcgagggc 300  
 gtggataccg tgaacgcctg ttacgggggc acctcggctc tcttcaatgc gatttcgtgg 360  
 gtggagtcca gctcctggga tggcaggtat gctcttgtgg tcgctgggga cattgctgtg 420  
 tatgctaaag gcagtgcgag gccacccggg ggagcagggg ctgtggccat gctagtgggc 480  
 gccaatgctc ccctagtgtt cgacagagga gtctgttcat cacacatgca acatgcttat 540  
 gacttctaca aaccggatct gtcctcgtg taccaccg tggatggcaa gctgtcaatt 600  
 caatgctatc ttagtgcctt agatcattgt tatcaactgt actgctccaa gatccagaaa 660  
 caacttggag agaagttcga tattgagcgg ctggatgcag ttctcttcca cgcgcttat 720  
 tgtaagttgg tgcagaagtc tcttgctcgc ctctcttga acgactttgt gcgggcatca 780  
 gaggaggagc ggacgactaa atactccagt ctggaagcac taaaaggcgt gaagctagaa 840  
 gatacgta ctgaccgaga agttgagaaa gcagtcatga catacagcaa gaacatgttt 900  
 gaagagaaaa caaagccctc gctgttgctc gccaaccaag tcggcaacat gtacactcct 960  
 tcgctttacg gaggtttggg ctctctattg gtcagcaaga gcgccagga gttggcaggg 1020  
 aagcgcgtgg ccttgttttc ttacggctcc ggactggcct cttccatgtt ctctctaaga 1080  
 atatcatcgg acgccagcgc gaaatcttct ctgcaacgcc tcgtctcgaa tctctcgac 1140  
 atcaagccgc agctggatct gcgccacaag gtgtcaccag aggagtttgc acaaacgatg 1200  
 gagacgaggg aacacaacca ccacaaagct ccatacacc cagagggctc gatcgacgtc 1260  
 ttgtttccag gaacttggtg tctggagagc gtggacagcc 1300

&lt;210&gt; 50

&lt;211&gt; 433

&lt;212&gt; PRT

<213> *Blattella germanica*

&lt;400&gt; 50

Ala Leu Glu Leu Ile Phe Pro Ser Gln Tyr Val Asp Gln Val Asp Leu  
 1 5 10 15

Glu Val Tyr Asp Asn Val Ser Ala Gly Lys Tyr Thr Val Gly Leu Gly  
 20 25 30

Gln Ala Arg Met Gly Phe Cys Thr Asp Arg Glu Asp Ile Asn Ser Leu  
 35 40 45

Cys Leu Thr Val Val Ser Arg Leu Met Glu Arg Trp Ser Ile Pro Tyr

50

55

60

Ser Gln Ile Gly Arg Leu Glu Val Gly Thr Glu Thr Leu Leu Asp Lys  
 65 70 75 80

Ser Lys Ser Val Lys Thr Val Leu Met Gln Leu Phe Lys Asp Asn Thr  
 85 90 95

Asp Ile Glu Gly Val Asp Thr Val Asn Ala Cys Tyr Gly Gly Thr Ser  
 100 105 110

Ala Leu Phe Asn Ala Ile Ser Trp Val Glu Ser Ser Ser Trp Asp Gly  
 115 120 125

Arg Tyr Ala Leu Val Val Ala Gly Asp Ile Ala Val Tyr Ala Lys Gly  
 130 135 140

Ser Ala Arg Pro Thr Gly Gly Ala Gly Ala Val Ala Met Leu Val Gly  
 145 150 155 160

Ala Asn Ala Pro Leu Val Phe Asp Arg Gly Val Arg Ser Ser His Met  
 165 170 175

Gln His Ala Tyr Asp Phe Tyr Lys Pro Asp Leu Ser Ser Leu Tyr Pro  
 180 185 190

Thr Val Asp Gly Lys Leu Ser Ile Gln Cys Tyr Leu Ser Ala Leu Asp  
 195 200 205

His Cys Tyr Gln Leu Tyr Cys Ser Lys Ile Gln Lys Gln Leu Gly Glu  
 210 215 220

Lys Phe Asp Ile Glu Arg Leu Asp Ala Val Leu Phe His Ala Pro Tyr  
 225 230 235 240

Cys Lys Leu Val Gln Lys Ser Leu Ala Arg Leu Val Leu Asn Asp Phe  
 245 250 255

Val Arg Ala Ser Glu Glu Glu Arg Thr Thr Lys Tyr Ser Ser Leu Glu  
 260 265 270

Ala Leu Lys Gly Val Lys Leu Glu Asp Thr Tyr Phe Asp Arg Glu Val  
 275 280 285

Glu Lys Ala Val Met Thr Tyr Ser Lys Asn Met Phe Glu Glu Lys Thr  
 290 295 300

Lys Pro Ser Leu Leu Leu Ala Asn Gln Val Gly Asn Met Tyr Thr Pro  
 305 310 315 320

Ser Leu Tyr Gly Gly Leu Val Ser Leu Leu Val Ser Lys Ser Ala Gln  
 325 330 335

Glu Leu Ala Gly Lys Arg Val Ala Leu Phe Ser Tyr Gly Ser Gly Leu  
 340 345 350

Ala Ser Ser Met Phe Ser Leu Arg Ile Ser Ser Asp Ala Ser Ala Glu  
 355 360 365

Ser Pro Leu Gln Arg Leu Val Ser Asn Leu Ser His Ile Lys Pro Gln  
 370 375 380

Leu Asp Leu Arg His Lys Val Ser Pro Glu Glu Phe Ala Gln Thr Met  
 385 390 395 400

Glu Thr Arg Glu His Asn His His Lys Ala Pro Tyr Thr Pro Glu Gly  
 405 410 415

Ser Ile Asp Val Leu Phe Pro Gly Thr Trp Tyr Leu Glu Ser Val Asp  
 420 425 430

Ser

<210> 51  
 <211> 25  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer forward

<400> 51  
 gctctggagc tcatattccc ttcgc

25

<210> 52  
 <211> 24  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> specific primer reverse

<400> 52  
 ggctgtccac gctctccaga tacc

24

<210> 53  
<211> 45  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer forward

<400> 53  
cgctaatacg actcactata gggctctgga gctcatattc ccttc 45

<210> 54  
<211> 41  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer reverse

<400> 54  
cgctaatacg actcactata ggggctgtcc acgctctcca g 41

<210> 55  
<211> 1300  
<212> RNA  
<213> Blattella germanica

<400> 55  
gcucuggagc ucauauuccc uucgcaguau guggaucagg uggaccucga ggucuacgac 60  
aauguuucug caggaaagua cacggugggg uggggacagg cucgcauggg guucugcacg 120  
gacagggagg acaucaacuc ucugugucuc accgucguca gucgacugau ggaacgaugg 180  
agcauccccu acucgcaaau ugggcgccug gaaguaggca ccgagaccu ucuggacaag 240  
ucgaagagcg ucaagacugu ccugaugcaa cucuucagg acaacacgga caucgagggc 300  
guggauaccg ugaacgccug uuacgggggc accucggcuc ucuucaaugc gauuucgugg 360  
guggagucca gcuccuggga uggcagguau gcucuugugg ucgcugggga cauugcugug 420  
uangcuaaag gcagugcgag gccaccggu ggagcagggg cuguggccau gcuagugggc 480  
gccaaugcuc ccuaguguu cgacagagga guucguucau cacacaugca acaugcuuau 540  
gacuucuaca aaccggaucu guccucgcug uacccaccg uggauaggca gcugucauuu 600  
caaugcuau uagugccuu agaucuuu uaucaacugu acugcuccaa gauccagaaa 660  
caacuuggag agaaguucga uauugagcgg cuggaugcag uucucuucca cgcgccuuau 720  
uguaaguugg ugcagaaguc ucuugcucgc cucgucuuga acgacuugu gcgggcauca 780  
gaggaggagc ggacgacuaa auacuccagu cuggaagcac uaaaaggcgu gaagcuagaa 840  
gauacguacu ucgaccgaga aguugagaaa gcagucauga cauacagcaa gaacauguuu 900

gaagagaaaa caaagcccuc gcuguugcuc gccaaccaag ucggcaacau guacacuccu 960  
 ucgcuuuacg gagguuuggu cucucuaauug gucagcaaga gcgcccagga guuggcaggg 1020  
 aagcgcgugg ccuuguuuuc uuacggcucc ggacuggccu cuuccauguu cucucuaaga 1080  
 auaucaucgg acgccagcgc gaaauucuuc cugcaacgcc ucgucucgaa ucucucgcac 1140  
 aucaagccgc agcuggaucu gcgccacaag gugucaccag aggaguuugc acaaacgaug 1200  
 gagacgaggg aacacaacca ccacaaagcu ccuacacccc cagagggcuc gaucgacguc 1260  
 uuguuuccag gaacuuggua ucuggagagc guggacagcc 1300

<210> 56

<211> 1279

<212> RNA

<213> *Blattella germanica*

<400> 56

gcucuggagc ucauauuccc uucgcaguau guggaucagg uggaccucga ggucuaacgac 60  
 aauguuucug caggaaagua cacggugggg uugggacagg cucgcauggg guucugcacg 120  
 gacagggagg acaucaacuc ucugugucuc accgucguca gucgacugau ggaacgaugg 180  
 agcauccccu acucgcaaau ugggcgccug gaaguaggca ccgagacccu ucuggacaag 240  
 ucgaagagcg ucaagacugu ccugaugcaa cucuuaagg acaacacgga caucgagggc 300  
 guggauaccg ugaacgccug uuacgggggc accucggcuc ucuucaaugc gauuucgugg 360  
 guggagucca gcuccuggga uggcagguau gcucuugugg ucgcugggga cauugcugug 420  
 uaugcuaaag gcagugcgag gccaccggu ggagcagggg cuguggccau gcuagugggc 480  
 gccaaugcuc cccuaguguu cgacagagga guucguucau cacacaugca acaugcuuau 540  
 gacuucuaca aaccggaucu guccucgcug uaccccaccg uggauggcaa gcugucaauu 600  
 caaugcuauc uuagugccuu agaauuugu uaucaacugu acugcuccaa gauccagaaa 660  
 caacuuggag agaaguucga uauugagcgg cuggaugcag uucucuucca cgcgccuuau 720  
 uguaaguugg ugcagaaguc ucuugcucgc cucgucuuga acgacuuugu gcgggcauca 780  
 gaggaggagc ggacgacuaa auacuccagu cuggaagcac uaaaaggcgu gaagcuagaa 840  
 gauacguacu ucgaccgaga aguugagaaa gcagucauga cauacagcaa gaacauguuu 900  
 gaagagaaaa caaagcccuc gcuguugcuc gccaaccaag ucggcaacau guacacuccu 960  
 ucgcuuuacg gagguuuggu cucucuaauug gucagcaaga gcgcccagga guuggcaggg 1020  
 aagcgcgugg ccuuguuuuc uuacggcucc ggacuggccu cuuccauguu cucucuaaga 1080  
 auaucaucgg acgccagcgc gaaauucuuc cugcaacgcc ucgucucgaa ucucucgcac 1140  
 aucaagccgc agcuggaucu gcgccacaag gugucaccag aggaguuugc acaaacgaug 1200

gagacgaggg aacacaacca ccacaaagcu ccuacaccc cagagggcuc gaucgacguc 1260  
 uuguuuccag gaacuuggu 1279

<210> 57  
 <211> 451  
 <212> DNA  
 <213> *Blattella germanica*

<400> 57  
 gaggcccaga gcäagagagg taccctcact ctgaagtacc ccattgaaca tggaatcatc 60  
 accaactggg atgacatgga gaagatctgg catcacacct tctacaatga actccgagtg 120  
 gctccagagg aacacccaat cctgctgact gaggtcccc tgaacccaaa ggccaacagg 180  
 gagaagatga ctcaaactcat gtttgagacc ttcaacaccc ccgcatgta tgttgccatc 240  
 caggccgtgc tgtccctcta cgcttcgggc cgtaccactg gtattgtgct ggactctggg 300  
 gagggcgtct cccacaccgt acccatctat gaagggttacg cattgcccc tgccatcctg 360  
 cgtctggact tggccggccg tgacttgact gactacctga tgaagatcct gaccgagcgt 420  
 ggctacagct tcacaactac agcagagcga g 451

<210> 58  
 <211> 150  
 <212> PRT  
 <213> *Blattella germanica*

<400> 58  
 Glu Ala Gln Ser Lys Arg Gly Ile Leu Thr Leu Lys Tyr Pro Ile Glu  
 1 5 10 15  
 His Gly Ile Ile Thr Asn Trp Asp Asp Met Glu Lys Ile Trp His His  
 20 25 30  
 Thr Phe Tyr Asn Glu Leu Arg Val Ala Pro Glu Glu His Pro Ile Leu  
 35 40 45  
 Leu Thr Glu Ala Pro Leu Asn Pro Lys Ala Asn Arg Glu Lys Met Thr  
 50 55 60  
 Gln Ile Met Phe Glu Thr Phe Asn Thr Pro Ala Met Tyr Val Ala Ile  
 65 70 75 80  
 Gln Ala Val Leu Ser Leu Tyr Ala Ser Gly Arg Thr Thr Gly Ile Val  
 85 90 95  
 Leu Asp Ser Gly Asp Gly Val Ser His Thr Val Pro Ile Tyr Glu Gly



100 105 110

Tyr Ala Leu Pro His Ala Ile Leu Arg Leu Asp Leu Ala Gly Arg Asp  
115 120 125

Leu Thr Asp Tyr Leu Met Lys Ile Leu Thr Glu Arg Gly Tyr Ser Phe  
130 135 140

Thr Thr Thr Ala Glu Arg  
145 150

<210> 59  
<211> 25  
<212> DNA  
<213> Artificial sequence

<220>  
<223> degenerative primer forward

<400> 59  
gaggcccaga gcaagagagg tatcc

25

<210> 60  
<211> 26  
<212> DNA  
<213> Artificial sequence

<220>  
<223> degenerative primer reverse

<400> 60  
ctcgctctgc ttagttgtg aagctg

26

<210> 61  
<211> 42  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer forward

<400> 61  
cgctaatacg actcactata gggaggccca gagcaagaga gg

42

<210> 62  
<211> 48  
<212> DNA  
<213> Artificial sequence

<220>  
<223> specific T7 primer reverse

<400> 62  
cgctaatacg actcactata ggtctgctgt agttgtgaag ctgtagcc

48

<210> 63  
 <211> 446  
 <212> RNA  
 <213> *Blattella germanica*

<400> 63  
 gaggcccaga gcaagagagg uauccucacu cugaaguacc ccuugaaca uggaaucauc 60  
 accaacuggg augacaugga gaagaucugg caucacaccu ucuacaauga acuccgagug 120  
 gcuccagagg aaacaaccaau ccugcugacu gaggcucucc ugaacccaaa ggccaacagg 180  
 gagaagauga cucaaaucan guuugagacc uucaacaccc ccgccaugua uguugccauc 240  
 caggccgugc ugucccucua cgcuuccggc cguaccacug guauugugcu ggacucuggu 300  
 gacggcgucu cccacaccgu acccaucuaa gaagguuacg cauugcccca ugccaucug 360  
 cgucuggacu ugcccgccg ugacuugacu gacuaccuga ugaagaucuu gaccgagcgu 420  
 ggcuacagcu ucacaacuac agcaga 446

<210> 64  
 <211> 108  
 <212> RNA  
 <213> *Blattella germanica*

<400> 64  
 uggcaucaca ccuucuacaa ugaacuccga guggcuccag aggaacaccc aauccugcug 60  
 acugaggcuc ccugaaccc aaaggccaac agggagaaga ugacuaa 108

<210> 65  
 <211> 50  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Synthetic fragment

<400> 65  
 tatgctcaa gaccaagcac aggacctcac aagttacgag agagtctgcc 50

<210> 66  
 <211> 50  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Synthetic fragment

<400> 66  
 cttgggtcgt gttggaactg tagttaatcg agaacgtcat cctgggtcct 50

<210> 67  
 <211> 250  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Concatemer

<400> 67  
 tatgctccaa gaccaagcac aggacctcac aagttacgag agagtctgcc tatgctccaa 60  
 gaccaagcac aggacctcac aagttacgag agagtctgcc tatgctccaa gaccaagcac 120  
 aggacctcac aagttacgag agagtctgcc tatgctccaa gaccaagcac aggacctcac 180  
 aagttacgag agagtctgcc tatgctccaa gaccaagcac aggacctcac aagttacgag 240  
 agagtctgcc 250

<210> 68  
 <211> 250  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Concatemer

<400> 68  
 cttgggtcgt gttggaactg tagttaatcg agaacgtcat cctgggttcct cttgggtcgt 60  
 gttggaactg tagttaatcg agaacgtcat cctgggttcct cttgggtcgt gttggaactg 120  
 tagttaatcg agaacgtcat cctgggttcct cttgggtcgt gttggaactg tagttaatcg 180  
 agaacgtcat cctgggttcct cttgggtcgt gttggaactg tagttaatcg agaacgtcat 240  
 cctgggttcct 250

<210> 69  
 <211> 100  
 <212> DNA  
 <213> Artificial sequenc

<220>  
 <223> Synthetic fragment

<400> 69  
 gctgaaaatat gcattaacca actgtgaggt taagaaaatt gttatgcagc gccttattaa 60  
 gggtgatgga aaagtcagaa cagaccccaa ctatccagct 100

<210> 70  
 <211> 300  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <223> Concatemer

<400> 70  
gctgaaatat gcattaacca actgtgaggt taagaaaatt gttatgcagc gccttattaa 60  
ggttgatgga aaagtcagaa cagaccccaa ctatccagct gctgaaatat gcattaacca 120  
actgtgaggt taagaaaatt gttatgcagc gccttattaa ggttgatgga aaagtcagaa 180  
cagaccccaa ctatccagct gctgaaatat gcattaacca actgtgaggt taagaaaatt 240  
gttatgcagc gccttattaa ggttgatgga aaagtcagaa cagaccccaa ctatccagct 300

<210> 71  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 71  
aaggcatgga tggtggacaa gct 23

<210> 72  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 72  
gcatggatgt tggacaagct cgg 23

<210> 73  
<211> 26  
<212> DNA  
<213> *Blattella germanica*

<400> 73  
attaaggttg atggaaaagt cagaac 26

<210> 74  
<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 74  
cccaactatc cagctggttt tatggatggt gt 32

<210> 75  
<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 75  
gctggtttta tggatgttgt tacaattgaa aa 32

<210> 76  
<211> 26  
<212> DNA

<213> Blattella germanica

<400> 76

attgaaaaaa ctggagaatt tttccg

26

<210> 77

<211> 23

<212> DNA

<213> Blattella germanica

<400> 77

ggtaatctct gtatgattac tgg

23

<210> 78

<211> 26

<212> DNA

<213> Blattella germanica

<400> 78

cgtcatcctg gttcctttga cattgt

26

<210> 79

<211> 22

<212> DNA

<213> Blattella germanica

<400> 79

ttaaagattc acaaggacac ac

22

<210> 80

<211> 26

<212> DNA

<213> Blattella germanica

<400> 80

aaaatccgta aagctgccag agaact

26

<210> 81

<211> 23

<212> DNA

<213> Blattella germanica

<400> 81

cgtaaagctg ccagagaact tct

23

<210> 82

<211> 24

<212> DNA

<213> Blattella germanica

<400> 82

aggttgtttg aaggcaatgc tctt

24

<210> 83

<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 83  
cgtattggag tgttgatga a 21

<210> 84  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 84  
ccgtatgaag cttgattacg t 21

<210> 85  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 85  
ttgggtttga agattgaaga tttcttga 29

<210> 86  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 86  
aagattgaag atttcttga a 21

<210> 87  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 87  
aggaacaaac gtgaagtgtg gcg 23

<210> 88  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 88  
tgctctcata ttgagaacat g 21

<210> 89  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 89  
aagggtttcc tgtacaaaat g 21

<210> 90  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 90  
gccgtgtatg cccatttccc cat 23

<210> 91  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 91  
tatgcccatt tccccattaa ctgcgt 26

<210> 92  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 92  
cgtaacttct tgggcgagaa gt 22

<210> 93  
<211> 24  
<212> DNA  
<213> Blattella germanica

<400> 93  
aaatggtttg gaacaaagaa ggag 24

<210> 94  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 94  
gatcccaatg aaataaacga aat 23

<210> 95  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 95  
aatgaaataa acgaaattgc aaatac 26

<210> 96  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 96

ggtttttgga aaaggaaggg tac

23

<210> 97  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 97  
gcaaattgccc gtatgccaca gaa

23

<210> 98  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 98  
aatgcccgtg tgccacagaa gg

22

<210> 99  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 99  
aagaagtaca gggaagcaaa gaa

23

<210> 100  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 100  
aagaagatcg acagacatct ata

23

<210> 101  
<211> 27  
<212> DNA  
<213> Blattella germanica

<400> 101  
caagggtaac gtgttcaaga acaagcg

27

<210> 102  
<211> 32  
<212> DNA  
<213> Blattella germanica

<400> 102  
aagggtaacg tgttcaagaa caagcgtgtc ct

32

<210> 103  
<211> 31  
<212> DNA  
<213> Blattella germanica



<400> 103  
gtgttcaaga acaagcgtgt cctgatggag t 31

<210> 104  
<211> 30  
<212> DNA  
<213> Blattella germanica

<400> 104  
tgatggagtt catccacaag aagaaggctg 30

<210> 105  
<211> 25  
<212> DNA  
<213> Blattella germanica

<400> 105  
catccacaag aagaaggctg agaag 25

<210> 106  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 106  
acaagaagaa ggctgagaag gc 22

<210> 107  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 107  
accaattcca gacaaaatat tcgtaa 26

<210> 108  
<211> 25  
<212> DNA  
<213> Blattella germanica

<400> 108  
gaagaaggct gagaaggcca ggaca 25

<210> 109  
<211> 54  
<212> DNA  
<213> Blattella germanica

<400> 109  
atggatgcca tcaagaagaa gatgcaggcg atgaagctgg agaaggacaa cgcg 54

<210> 110  
<211> 25

<212> DNA  
<213> Blattella germanica

<400> 110  
gggccgagaa ggctgaggag gaggc 25

<210> 111  
<211> 38  
<212> DNA  
<213> Blattella germanica

<400> 111  
tccctgcaga agaagatcca gcagattgag aatgatct 38

<210> 112  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 112  
tgatgcaagt caacgccaaag ct 22

<210> 113  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 113  
atgcaagtca acgccaaagt gga 23

<210> 114  
<211> 35  
<212> DNA  
<213> Blattella germanica

<400> 114  
gtcaacgccaa agctggacga gaaggacaag gccct 35

<210> 115  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 115  
gagaaggaca aggccctgca gaa 23

<210> 116  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 116  
aaccgccgaa tccaactgct ggagga 26

<210> 117  
<211> 38  
<212> DNA  
<213> Blattella germanica

<400> 117  
gcgatgaagc tggagaagga caacgcgatg gatcgcg  
38

<210> 118  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 118  
tctgaggaac gtttggccac agc  
23

<210> 119  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 119  
tggcagatga agagcgtatg ga  
22

<210> 120  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 120  
gatgaagagc gtatggatgc t  
21

<210> 121  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 121  
gctttggaga accagctgaa gga  
23

<210> 122  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 122  
gagaaccagc tgaaggaagc c  
21

<210> 123  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 123  
cagctgaagg aagccagggtt c  
21

<210> 124  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 124  
ttcatggctg aggaagctga caagaaata 29

<210> 125  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 125  
gctgacaaga aatatgatga ggt 23

<210> 126  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 126  
gacaagaaat atgatgaggt cgc 23

<210> 127  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 127  
atggttgagg ccgacttgga aagagcaga 29

<210> 128  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 128  
gccgacttgg aaagagcaga aga 23

<210> 129  
<211> 27  
<212> DNA  
<213> Blattella germanica

<400> 129  
cgacttggaag agagcagaag agcgtgc 27

<210> 130  
<211> 22  
<212> DNA  
<213> Blattella germanica

<400> 130  
ccaagattgt ggagcttgag ga 22

<210> 131  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 131  
aagattgtgg agcttgagga aga 23

<210> 132  
<211> 34  
<212> DNA  
<213> Blattella germanica

<400> 132  
tggatcgcgc ccttctctgc gaacagcagg cccg 34

<210> 133  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 133  
attgtggagc ttgaggaaga actgcgcgt 29

<210> 134  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 134  
ctgcgcgttg tcggcaacaa c 21

<210> 135  
<211> 35  
<212> DNA  
<213> Blattella germanica

<400> 135  
cgcgttgctg gcaacaacct gaagtcctt gaggt 35

<210> 136  
<211> 59  
<212> DNA  
<213> Blattella germanica

<400> 136  
gttgtcggca acaacctgaa gtcccttgag gtgtctgaag agaaggccaa cctgcgtga 59

<210> 137  
<211> 21

<212> DNA  
<213> *Blattella germanica*

<400> 137  
taccaggcta aaggaggctg a 21

<210> 138  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 138  
accaggctaa aggaggctga agc 23

<210> 139  
<211> 21  
<212> DNA  
<213> *Blattella germanica*

<400> 139  
gctaaaggag gctgaagctc g 21

<210> 140  
<211> 29  
<212> DNA  
<213> *Blattella germanica*

<400> 140  
ctaaaggagg ctgaagctcg tgctgagtt 29

<210> 141  
<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 141  
aaggaggctg aagctcgtgc tgagttcgct ga 32

<210> 142  
<211> 21  
<212> DNA  
<213> *Blattella germanica*

<400> 142  
gctcgtgctg agttcgctga a 21

<210> 143  
<211> 60  
<212> DNA  
<213> *Blattella germanica*

<400> 143  
tgcagaagga gggtgacagg cttgaggatg aattggtaca cgagaaggag aagtacaagt 60

<210> 144  
<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 144  
ttggtacacg agaaggagaa gtacaagtac at 32

<210> 145  
<211> 53  
<212> DNA  
<213> *Blattella germanica*

<400> 145  
gagaaggaga agtacaagta catttgtgac gatcttgata tgactttcac cga 53

<210> 146  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 146  
aacagcaggc ccgcgacgcc aac 23

<210> 147  
<211> 42  
<212> DNA  
<213> *Blattella germanica*

<400> 147  
catttgtgac gatcttgata tgactttcac cgaacttatt gg 42

<210> 148  
<211> 22  
<212> DNA  
<213> *Blattella germanica*

<400> 148  
cggacaggga ggacatcaac tc 22

<210> 149  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 149  
tggacaagtc gaagagcgtc aag 23

<210> 150  
<211> 21  
<212> DNA  
<213> *Blattella germanica*

<400> 150  
ctgctccaag atccagaaac a 21

<210> 151  
<211> 44  
<212> DNA  
<213> *Blattella germanica*

<400> 151  
gaggcccaga gcaagagagg taccctcact ctgaagtacc ccat 44

<210> 152  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 152  
gctccagagg aacacccaat cct 23

<210> 153  
<211> 23  
<212> DNA  
<213> *Blattella germanica*

<400> 153  
atcctgctga ctgaggctcc cct 23

<210> 154  
<211> 47  
<212> DNA  
<213> *Blattella germanica*

<400> 154  
aaggccaaca gggagaagat gactcaaacc atgtttgaga ccttcaa 47

<210> 155  
<211> 29  
<212> DNA  
<213> *Blattella germanica*

<400> 155  
caaatcatgt ttgagacctt caacacccc 29

<210> 156  
<211> 37  
<212> DNA  
<213> *Blattella germanica*

<400> 156  
tcatgtttga gaccttcaac acccccgcga tgtatgt 37

<210> 157  
<211> 38  
<212> DNA  
<213> *Blattella germanica*



<400> 157  
accttcaaca ccccgccat gtatgttgcc atccaggc 38

<210> 158  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 158  
cccgccatgt atgttgccat ccaggccgt 29

<210> 159  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 159  
gccatccagg ccgtgctgtc cct 23

<210> 160  
<211> 30  
<212> DNA  
<213> Blattella germanica

<400> 160  
tacgcttccg gccgtaccac tggattgtg 30

<210> 161  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 161  
gcttccggcc gtaccactgg tat 23

<210> 162  
<211> 32  
<212> DNA  
<213> Blattella germanica

<400> 162  
cgtaccactg gtattgtgct ggactctggt ga 32

<210> 163  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 163  
ggtattgtgc tggactctgg tgacgg 26

<210> 164  
<211> 23  
<212> DNA

<213> Blattella germanica

<400> 164

ggtgacggcg tctccacac cgt

23

<210> 165

<211> 23

<212> DNA

<213> Blattella germanica

<400> 165

ggtgacggcg tctccacac cgt

23

<210> 166

<211> 32

<212> DNA

<213> Blattella germanica

<400> 166

tccacaccg tacccatcta tgaagggtac gc

32

<210> 167

<211> 40

<212> DNA

<213> Blattella germanica

<400> 167

tgaagtaccc cattgaacat ggaatcatca ccaactggga

40

<210> 168

<211> 28

<212> DNA

<213> Blattella germanica

<400> 168

tgcccatgc catcctgcgt ctggactt

28

<210> 169

<211> 30

<212> DNA

<213> Blattella germanica

<400> 169

gccatcctgc gtctggactt ggccggccgt

30

<210> 170

<211> 29

<212> DNA

<213> Blattella germanica

<400> 170

cgtctggact tggccggccg tgacttgac

29

<210> 171

<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 171  
cgtgacttga ctgactacct gatgaagatc ct 32

<210> 172  
<211> 36  
<212> DNA  
<213> *Blattella germanica*

<400> 172  
gactacctga tgaagatcct gaccgagcgt ggctac 36

<210> 173  
<211> 35  
<212> DNA  
<213> *Blattella germanica*

<400> 173  
atgaagatcc tgaccgagcg tggctacagc ttcac 35

<210> 174  
<211> 29  
<212> DNA  
<213> *Blattella germanica*

<400> 174  
ggaatcatca ccaactggga tgacatgga 29

<210> 175  
<211> 38  
<212> DNA  
<213> *Blattella germanica*

<400> 175  
atcatcacca actgggatga catggagaag atctggca 38

<210> 176  
<211> 35  
<212> DNA  
<213> *Blattella germanica*

<400> 176  
gacatggaga agatctggca tcacaccttc tacaa 35

<210> 177  
<211> 36  
<212> DNA  
<213> *Blattella germanica*

<400> 177  
atggagaaga tctggcatca caccttctac aatgaa 36

<210> 178  
<211> 35  
<212> DNA  
<213> *Blattella germanica*

<400> 178  
aagatctggc atcacacctt ctacaatgaa ctccg 35

<210> 179  
<211> 35  
<212> DNA  
<213> *Blattella germanica*

<400> 179  
atctggcatc acaccttcta caatgaactc cgagt 35

<210> 180  
<211> 26  
<212> DNA  
<213> *Blattella germanica*

<400> 180  
aacgtgttca agaacaagcg tgtcct 26

<210> 181  
<211> 31  
<212> DNA  
<213> *Blattella germanica*

<400> 181  
catccacaag aagaaggctg agaaggccag g 31

<210> 182  
<211> 41  
<212> DNA  
<213> *Blattella germanica*

<400> 182  
gccatcaaga agaagatgca ggcgatgaag ctggagaagg a 41

<210> 183  
<211> 24  
<212> DNA  
<213> *Blattella germanica*

<400> 183  
cctgcagaag aagatccagc agat 24

<210> 184  
<211> 32  
<212> DNA  
<213> *Blattella germanica*

<400> 184

aagatgcagg cgatgaagct ggagaaggac aa

32

<210> 185

<211> 21

<212> DNA

<213> Blattella germanica

<400> 185

gagaaggaca aggccctgca g

21

<210> 186

<211> 26

<212> DNA

<213> Blattella germanica

<400> 186

gttgtcggca acaacctgaa gtcctt

26

<210> 187

<211> 23

<212> DNA

<213> Blattella germanica

<400> 187

aaggaggctg aagctcgtgc tga

23

<210> 188

<211> 27

<212> DNA

<213> Blattella germanica

<400> 188

gagggccaga gcaagagagg taccctc

27

<210> 189

<211> 23

<212> DNA

<213> Blattella germanica

<400> 189

cagagcaaga gaggtaccct cac

23

<210> 190

<211> 34

<212> DNA

<213> Blattella germanica

<400> 190

gcccagagca agagaggtat cctcactctg aagt

34

<210> 191

<211> 23

<212> DNA

<213> Blattella germanica

<400> 191  
aaggccaaca gggagaagat gac 23

<210> 192  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 192  
gagaagatga ctcaaatacat gtt 23

<210> 193  
<211> 29  
<212> DNA  
<213> Blattella germanica

<400> 193  
ggtatcctca ctctgaagta cccattga 29

<210> 194  
<211> 21  
<212> DNA  
<213> Blattella germanica

<400> 194  
atcatgtttg agaccttcaa c 21

<210> 195  
<211> 26  
<212> DNA  
<213> Blattella germanica

<400> 195  
gagaccttca acacccccgc catgta 26

<210> 196  
<211> 23  
<212> DNA  
<213> Blattella germanica

<400> 196  
gccatccagg ccgtgctgtc cct 23

<210> 197  
<211> 35  
<212> DNA  
<213> Blattella germanica

<400> 197  
gtctcccaca ccgtacccat ctatgaaggt tacgc 35

<210> 198  
<211> 40

<212> DNA

<213> *Blattella germanica*

<400> 198

tcaccaactg ggatgacatg gagaagatct ggcatcacac

40

<210> 199

<211> 35

<212> DNA

<213> *Blattella germanica*

<400> 199

gacatggaga agatctggca tcacaccttc taca

35

<210> 200

<211> 41

<212> DNA

<213> *Blattella germanica*

<400> 200

atggagaaga tctggcatca caccttctac aatgaactcc g

41