

[54] 2-HYDROCARBYLOXY-PYRIDINE
COMPOUNDS[75] Inventors: **Erwin Fleckenstein**, Hofheim,
Taunus; **Ernst Heinrich**, Frankfurt
am Main-Fechenheim; **Reinhard
Mohr**, Offenbach-Rumpenheim, all
of Germany[73] Assignee: **Cassella Farbwerke Mainkur AG**,
Germany[22] Filed: **Mar. 31, 1975**[21] Appl. No.: **563,848****Related U.S. Application Data**

[62] Division of Ser. No. 372,024, June 21, 1973.

[30] **Foreign Application Priority Data**


June 22, 1972 Germany..... 2230392

[52] **U.S. Cl.** 260/294.9; 260/294.8 C; 260/294.8 D;
260/294.8 F; 260/294.8 G; 260/247.1 M;
260/268 H; 260/293.69; 260/156; 8/41 C[51] **Int. Cl.²**..... **C07D 213/57**[58] **Field of Search**..... 260/294.9[56] **References Cited****UNITED STATES PATENTS**



3,853,895 12/1975 Lamm et al..... 260/294.9

Primary Examiner—Alan L. Rotman*Attorney, Agent, or Firm*—Connolly and Hutz[57] **ABSTRACT**




Compounds of the formula






wherein X is aliphatic, cycloaliphatic, aromatic, aralkyl, hetaryl, hetarylalkyl or hydrogen; Y is aliphatic, aromatic, aralkyl, —COOR₁, —COR₂,



—CN, —NH₂, —NO, —NO₂ or hydrogen with the proviso that when Y is hydrogen, X is other than hydrogen; Z₁ is cyano,



—OR₁₂, —SR₁₂ or —SO₂R₁₂ and Z₂ is chlorine, bromine, cyano, hydroxy, mercapto, —OR₁₂, —SR₁₂, —SO₂R₁₂,



R₁ is aliphatic; R₂ is aliphatic, cycloaliphatic, aromatic, aralkyl or heterocyclic; R₃ and R₄ when taken separately are hydrogen, aliphatic, cycloaliphatic, aromatic or aralkyl; R₃ and R₄ when taken together with the nitrogen atom to which they are attached form a heterocyclic moiety; R₅ and R₆ when taken separately are aliphatic, aromatic, aralkyl, heterocyclic or hydrogen; R₅ and R₆ when taken together with the nitrogen atom to which they are attached are heterocyclic; R₇ is hydrogen, aliphatic or aromatic; R₈ and R₉ are aliphatic or aromatic, R₁₀ is hydrogen, aliphatic or aralkyl, R₁₁ is aliphatic and R₁₂ is aliphatic, aromatic or aralkyl; R₁₀ and R₁₁ when taken together with the nitrogen atom to which they are attached are heterocyclic.


2. Claims, No Drawings

2-HYDROCARBYLOXY-PYRIDINE COMPOUNDS


CROSS-REFERENCE TO RELATED APPLICATION

This application is a division of application Ser. No. 372,024 filed June 21, 1973.


This invention relates to compounds of the formula




wherein X is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl having three to eight carbon atoms, heteroaryl having four to seven carbon atoms and containing one or two heteroatoms selected from the group of O, N and S, phenyl, benzyl, phenethyl, one of said radicals substituted as hereinafter defined or hydrogen; Y is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl, phenethyl, one of said radicals substituted as hereinafter defined, $-\text{COOR}_1$, $-\text{COR}_2$,




$-\text{SO}_2\text{R}_2$,




$-\text{CN}$, $-\text{NH}_2$, $-\text{NO}$, $-\text{NO}_2$ or hydrogen with the proviso that when Y is hydrogen, X is other than hydrogen; Z_1 is cyano,




$-\text{NH}-\text{OR}_{10}$,



$-\text{OR}_{12}$, $-\text{SR}_{12}$ or $-\text{SO}_2\text{R}_{12}$ and Z_2 is chlorine, bromine, cyano, hydroxy, mercapto, $-\text{OR}_{12}$, $-\text{SR}_{12}$, $-\text{SO}_2\text{R}_{12}$,




$-\text{NH}-\text{OR}_{10}$ or




R_1 is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms or one of said radicals substituted as hereinafter defined; R_2 is alkyl having

one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl, phenethyl, pyridyl, thiazolyl, benzothiazolyl, imidazolyl, benzimidazolyl, thienyl, furyl or pyrrolyl, one of said radicals substituted as hereinafter defined; R_3 and R_4 , when taken separately, are alkyl having one to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl, phenethyl, one of said radicals substituted as hereinafter defined or hydrogen; R_3 and R_4 , when taken together with the nitrogen atom to which they are attached, form a piperidino, piperazino, morpholino, or ethyleneimino radical; R_5 and R_6 , when taken separately, are alkyl having one to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl, phenethyl, pyridyl, thiazolyl, benzothiazolyl, imidazolyl, benzimidazolyl, pyrrolyl, one of said radicals substituted as hereinafter defined or hydrogen; R_5 and R_6 when taken together with the nitrogen atom to which they are attached, form a piperidino, piperazino, morpholino or ethyleneimino radical; R_7 is hydrogen, alkyl having one to six carbon atoms, phenyl or phenyl substituted as hereinafter defined; R_8 and R_9 are alkyl having one to six carbon atoms, phenyl or phenyl substituted as hereinafter defined; R_{10} is alkyl having one to six carbon atoms, benzyl, phenethyl, one of said groups substituted as hereinafter defined or hydrogen; R_{11} is alkyl having one to six carbon atoms or said alkyl substituted as hereinafter defined; R_{12} is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl, phenethyl or one of said radicals substituted as hereinafter defined; the substituents for substituted alkyl or alkenyl radicals specified hereinabove being selected from the group consisting of cyano, hydroxy, acetoxy, phenoxyacetoxy, alkoxy having one to two carbon atoms, phenoxy, monoalkylamino having one to four carbon atoms, dialkylamino having one to four carbon atoms in each alkyl moiety, N-ethylcarbamoyloxy, N-methylcarbamoyloxy, N-phenylcarbamoyloxy, acetyl amino, benzoylamino, heteroaryl having four to seven carbon atoms and containing one or two heteroatoms selected from the group of O, N and S, and




wherein Z constitutes the atoms completing a morpholino, piperidino, piperazino, N-methyl-piperazino or cyclopentamethyleneimino group and the substituents for the substituted cycloalkyl, phenyl, benzyl, phenethyl and heterocyclic radicals specified hereinabove being selected from the group consisting of bromine, chlorine, cyano, alkyl having one to three carbon atoms and alkoxy having one to two carbon atoms. The above-mentioned heteroaryl residues having four to seven carbon atoms may be pyridyl, thiazolyl, benzthiazolyl, imidazolyl, benzimidazolyl, thienyl, furyl or pyrrolyl groups.

The starting materials for the preparation of the novel compounds of formula I are 2,6-dihydroxypyridine derivatives of the formula:



3


wherein X and Y have the above-stated meaning. These starting compounds may be prepared in a known manner analogous to the method described by Bobbitt and Scola, Journ. of Org. Chem. 25, 560, by the condensation of correspondingly substituted acetic acid amides with correspondingly substituted β -ketocarboxylic acid esters. They may also be prepared in accordance with various other processes, as described for example in the monograph "Heterocyclic Compounds Pyridine and its Derivatives Part. 3" by Klingsberg. This monograph has appeared in the frame of the series published by Arnold Weissberger "The Chemistry of Heterocyclic Components," Interscience Publishers. The 2,6-dihydroxypyridine derivatives of formula II, according to Bobbitt and Scola, *ibid.*, are converted by means of phosphoroxychloride or analogously by means of phosphoroxybromide at 180°C. into the corresponding 2,6-dichloro or 2,6-dibromopyridine derivatives of formula III:



III


wherein X and Y have the above-stated meanings and Hal is bromine or preferably chlorine.

For the preparation of compounds of formula I, wherein Z_2 stands for chlorine or bromine and to which formula Ia then pertains, a compound of formula III is reacted with a compound of formula IV in accordance with the reaction:



III

IV




Ia

(1)

Q in this instance is hydrogen or a metal, preferably an alkali metal such as sodium or potassium. In the event that Z_1 stands for $-\text{SO}_2\text{R}_{12}$, zinc is especially preferred.


Reaction (1) is carried out in equimolar ratio at temperatures of 20°–100°C. in a suitable inert solvent. The exchange of the chlorine or bromine atom in the 6-position is ascertained by analysis of the NMR absorption of the obtained compounds of formula Ia.

To the extent that in formula IV, Z_1 stands for the radicals:




4


$-\text{NH}-\text{OR}_{10}$ or



hydrogen is appropriately selected for Q, i.e., in the reaction equation (1) formula IV then stands for the compounds



$\text{H}_2\text{NOR}_{10}$ or




In this instance, the reaction is carried out in accordance with reaction equation (1), preferably in the temperature range of 20° to 50°C. and as suitable inert solvents, alkanols with a chain length of one to four carbon atoms are preferably used. In those instances where the physical properties are appropriate, an excess of one of the reaction constituents may also serve as the solvent.

To the extent that in formula IV, Z_1 stands for cyano or for $-\text{SO}_2\text{R}_{12}$, a metal is appropriately selected for Q, preferably an alkali metal such as sodium or potassium and in the event that Z_1 is SO_2R_{12} , zinc is particularly preferred. In reaction equation (1), formula IV then stands for the compounds NaCN , KCN and $\text{Zn}(\text{SO}_2\text{R}_{12})_2$. In this instance, the reaction in accordance with reaction equation (1) is carried out preferably in the temperature range of 60°–80°C. in a suitable organic solvent such as an alcohol, ether, ester, carboxylic acid amide, N-substituted carboxylic acid amide such as dimethylformamide, dialkylsulfoxide such as dimethylsulfoxide or dialkylsulfone, all of these preferably having not more than six carbon atoms. In the reaction of sulfinates, the addition of a copper-I-salt is advantageous.

To the extent that in formula IV, Z_1 stands for the radicals $-\text{OR}_{12}$ or $-\text{SR}_{12}$, a metal atom is selected for Q, preferably an alkali metal atom such as sodium or potassium. In reaction equation (1), formula IV then stands for the compounds NaOR_{12} , NaSR_{12} , KOR_{12} and KSR_{12} . In this instance, the reaction is conducted in accordance with reaction equation (1), preferably in the temperature range of 60°–100°C., in an inert organic solvent, e.g., an aromatic hydrocarbon such as benzene, toluene or xylene, or in an excess of the reaction constituent HOR_{12} or HSR_{12} as the solvent.


The compounds of formula Ia may be further reacted in accordance with the reaction:



Ia

V

5



This reaction is carried out at temperatures of 20°–200°C. in a suitable solvent. The molar ratio of reactants is 1:1.


When Z_2 is OH in the compound of formula V, there is used a suitable alkali metal hydroxide such as sodium hydroxide or potassium hydroxide in an aqueous or aqueous-alcoholic medium. However, sodium carbonate, potassium hydroxide or similar compounds which release hydroxyl ions in an aqueous medium may also be used. The solvents may be alkanols having one to four carbon atoms. The preferred temperature range is 70°–150°C.

If Z_2 is —SH in the compound of formula V, an alkali metal hydrosulfide, preferably NaSH or KSH, is used and the reaction is carried out in an aqueous medium, preferably at temperatures of 100°–150°C.


In the event Z_2 stands for —OR₁₂ or —SR₁₂, an alkali metal, preferably sodium or potassium, is selected for Q in formula V and the reaction constituents are reacted at temperatures of 60°–150°C., preferably at 60°–100°C., in a suitable inert solvent, e.g., an aromatic hydrocarbon such as benzene, toluene or xylene, or in an excess of the reaction constituent HOR₁₂ or HSR₁₂ as solvent.

In the event Z_2 is cyano or —SO₂R₁₂ in formula V, a metal is selected for Q, preferably an alkali metal such as sodium or potassium or when Z_2 is —SO₂R₁₂, zinc is also useful. The reaction between these compounds of formulae Ia and V is carried out in a suitable organic solvent at temperatures of 20°–100°C. and preferably 60°–80°C. Suitable organic solvents, for example, include alcohols, ethers, esters, carboxylic acid amides, N-substituted carboxylic acid amides such as dimethylformamide, dialkylsulfoxides such as dimethylsulfoxide or dialkylsulfones, all having preferably not more than six carbon atoms. In the use of sulfinates, the addition of a copper-I-salt is advantageous.

In the event Z_2 is




—NH—OR₁₀ or




in formula V, hydrogen is selected for Q, and the constituents are reacted at 100°–200°C., preferably at 160°–180°C., in an inert solvent, preferably an alkanol having one to four carbon atoms or, in those instances where the physical properties are appropriate, an excess of one of the reaction constituents may be used as the solvent.

6


Novel compounds wherein Z_2 has the same meaning as Z_1 may be prepared in accordance with the reaction:



wherein a compound of formula III is reacted with a compound of formula IV in a molar ratio of at least 1:2 in a suitable solvent at temperatures of 20° to 200°C. When Z_1 is




—NH—OR₁₀ or



in formula IV, hydrogen is selected for Q, and when Z_1 is —CN, —OR₁₂, —SR₁₂ or —SO₂R₁₂, a metal is selected for Q, preferably in alkali metal such as sodium or potassium. Also when Z_1 is —SO₂R₁₂, zinc is suitable. Suitable solvents are alcohols, ethers, hydrocarbons, etc. The selection of the solvent may be the same as described for reaction (1).



Novel compounds when Z_2 is —OH may also be prepared by the reaction of a 2-hydroxy-6-bromo- or 2-hydroxy-6-chloropyridine derivative of formula VI with a compound of formula IV in a molar ratio of 1:1 in accordance with reaction:



The reaction is carried out in a suitable solvent at temperatures of 20° to 100°C. The data respecting the solvents and the meaning and selection of Q set forth above for reaction equations (1) and (3), apply to reaction (4).


Y is cyano,

Such preferred compounds are especially the pyridine
15 compounds of the formulae:



As the reaction constituents of formula III such as 2,6-dichloro or 2,6-dibromo pyridine derivatives are suitable, for example, which in the X position contain hydrogen or a methyl, ethyl, n-propyl, isopropyl, vinyl, α -methylvinyl, n-butyl, isobutyl, sec-butyl, n-amyl or isoamyl, n-hexyl, 2-dimethylaminoethyl, 2-diethylaminoethyl, 2-morpholinoethyl, 2-piperidinoethyl, 2-pyrrolidinoethyl, N-methyl-N'-piperazinoethyl, 2-cyanoethyl, 2-hydroxyethyl, 2-methoxyethyl, 2-acetoxylethyl, 2-phenoxyacetoxylethyl, N-ethylcarbamoyloxyethyl, N-phenylcarbamoyloxyethyl, 2-phenoxyethyl, 3-methoxypropyl, cyclohexyl, benzyl, 3-methylbenzyl, phenyl, 2-methylphenyl, 4-methylphenyl, 2,4-dimethylphenyl, 2-chloro-4-methylphenyl, 2-chlorophenyl, 4-chlorophenyl, 2-methoxyphenyl, 4-methoxyphenyl, 2,5-dimethyl-4-chlorophenyl or the following radicals:





acetoxyethyl, 2-benzoyloxyethyl, 2-methoxyethyl, 2-phenoxyethyl, 2-monoethylaminoethyl, 2-monomethylaminoethyl, 2-dimethylaminoethyl, 2-diethylaminoethyl, 2-morpholinoethyl, 2-piperidinoethyl, 2-pyrrolidinoethyl, n-propyl, isopropyl, vinyl, α -methylvinyl, n-butyl, isobutyl, sec.-butyl, n-amy, iso-amy, n-hexyl, cyclohexyl, benzyl, phenyl, methoxycarbonyl, ethoxycarbonyl, n-propyloxycarbonyl, isopropyloxycarbonyl, n-butyloxycarbonyl, isobutyloxycarbonyl, sec.-butyloxycarbonyl, n-amyloxycarbonyl, n-isoamyloxycarbonyl, n-hexyloxycarbonyl, acetyl, acryloyl, propionyl, capronyl, capryl, hexahydrobenzoyl, phenacetyl, benzoyl, 4-methylbenzoyl, 2,4-dimethylbenzoyl, 4-methoxybenzoyl, 4-chlorbenzoyl, aminocarbonyl, monomethylaminocarbonyl, dimethylaminocarbonyl, ethyleniminocarbonyl, monoethylaminocarbonyl, monoisopropylaminocarbonyl, diethylaminocarbonyl, monooxethylaminocarbonyl, mono- γ -methoxy-



propylaminocarbonyl, morpholinocarbonyl,
piperidinocarbonyl, cyclohexylaminocarbonyl, benzylaminocarbonyl, anilino- carbonyl, 4-

11

methylanilinocarbonyl, N-methylanilinocarbonyl, methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl, n-amylsulfonyl, isoamylsulfonyl, n-hexylsulfonyl, benzylsulfonyl, phenylsulfonyl, 4-methylphenylsulfonyl, 2,4-dimethylphenylsulfonyl, 4-methoxyphenylsulfonyl, 4-chlorophenylsulfonyl, aminosulfonyl, monomethylaminosulfonyl, dimethylaminosulfonyl, ethyleniminosulfonyl, monoethylaminosulfonyl, monoisopropylaminosulfonyl, diethylaminosulfonyl, monoethoxethylaminosulfonyl, mono-γ-methoxypropylaminosulfonyl, morpholinisulfonyl, piperidinisulfonyl, cyclohexylaminosulfonyl, benzylaminosulfonyl, anilinosulfonyl, 4-methylanilinosulfonyl, N-methylanilinosulfonyl or the radicals



Of course, suitable starting compounds of formula III are also such 2,6-dichloro or 2,6-dibromopyridine derivatives which are identically substituted in the X and Y position, the preferred substituents being those which were indicated above for the X position.

Suitable reaction constituents






of formulae IV and V include, for example, the following primary and secondary amines and diamines:

Primary Amines


Ammonia, methylamine, ethylamine, 2-hydroxyethylamine, 2-methoxyethylamine, 3-phenoxyethylamine, 3-cyanoethylamine, n-propylamine, isopropylamine, 3-hydroxypropylamine-(1), 3-methoxypropylamine-(1), 3-isopropoxypropylamine-(1), 3-cyano-propylamine-(1), allylamine, 1-methylallylamine, 2-methylallylamine, n-butylamine, isobutylamine, sec-butylamine, tert-butylamine, 2-amino-2-methyl-

12

propanol-(1), crotylamine, 3-aminopentane, n-amylamine, isoamylamine, n-hexylamine, cyclohexylamine, benzylamine, 2-phenylethylamine, aniline, 4-methylaniline, 4-methoxyaniline, 2,4-dimethylaniline, 1-aminonaphthalene, 2-aminonaphthalene,



and




Secondary Amines


Dimethylamine, diethylamine, N-(2-cyanoethyl)-N-(2-hydroxyethyl)-amine, N-di-(2-hydroxyethyl)-amine, N-di-(2-cyanoethyl)-amine, N-methyl-(2-hydroxyethyl)-amine, N-isopropyl-(2-hydroxyethyl)-amine, N-n-butyl-(2-hydroxyethyl)-amine, N-cyclohexyl-(2-hydroxyethyl)-amine, N-benzyl-(2-hydroxyethyl)-amine, di-n-propylamine, di-isopropylamine, di-n-butylamine, diisobutylamine, di-n-amylamine, di-n-hexylamine, morpholine pyrrolidine, piperidine, N-methylpiperazine, N-methylcyclohexylamine, N-ethylcyclohexylamine, N-methylbenzylamine, N-methyl-3-methylbenzylamine, N-methylaniline, N-ethylaniline, N-2-hydroxyethylaniline, N-benzylaniline, N-methyl-2-chloraniline and N-ethyl-2-chloraniline.

Diamines

N-methyl-N',N'-dimethylhydrazine, N,N-dimethylhydrazine, N,N-diethylhydrazine, N-methyl-N-phenylhydrazine, N-aminopyrrolidine, N-aminopiperidine, N-aminopiperazine, N-aminomorpholine, N,N-dimethylethylenediamine, N,N-diethylethylenediamine, N,N-dimethylpropylenediamine-(1,3), or N,N-diethylpropylenediamine-(1,3), 2-morpholinoethylamine, 2-piperidinoethylamine, 2-pyrrolidinoethylamine and N-methyl-N'-3-amino-n-propylpiperazine of the formula:




Suitable reaction constituents H_2NOR_{10} or



according to the general formulae IV and V, for example, are:

Hydroxylamines



As the hetaryl amines for Z_1 and Z_2 , the following may, for example, be used:

3-aminodiphenylene oxide
3-aminodiphenylene sulfide
3-aminodiphenylene sulfone
2-aminocarbazole
3-amino-N-methylcarbazole
3-amino-N-ethylcarbazole
3-amino-N- β -hydroxyethylcarbazole
3-amino-N- β -cyanoethylcarbazole
3-amino-N-n-propylcarbazole
3-amino-N-isopropylcarbazole
3-amino-N-(β -dimethylaminoethyl)-carbazole
3-amino-N-(β -diethylaminoethyl)-carbazole
3-amino-N-(γ -dimethylaminopropyl)-carbazole
3-amino-N-(α -methyl- β -dimethylaminoethyl)-carbazole

As the compounds $HO-R_{12}$ or $HS-R_{12}$, which after their conversion into the corresponding metal compounds, particularly the corresponding alkali metal compounds (especially the corresponding sodium and potassium compounds) are suitable as reaction constituents of formulae IV and V, the following alcohols, phenols or the corresponding mercaptans and thiophenols may be utilized: methanol, ethanol, 2-cyanomethanol-(1), ethylene glycol monomethyl ether, ethylene glycol monoethylether, ethylene glycol monoisopropyl ether, ethylene glycol mono-n-butylether, ethylene glycol monophenyl ether, ethylene glycol monoxylenyl ether, diethylene glycol monomethylether, diethylene glycol monoethylether, diethylene glycol mono-n-butylether, triethylene glycol monomethylether, triethylene glycol monoethylether, triethylene glycol mono-n-butylether, n-propanol, isopropanol, propene-(1)-ol-(3), 2-methylpropene-(1)-ol-(3), n-butanol, sec.-butanol, isobutanol, tert.-butanol, 3-methoxybutanol-(1), 4-methoxybutanol-(1), butene-(1)-ol-(2), n-pentanol, isopentanol, n-hexanol, cyclohexanol, 4-methylcyclohexanol, 4-methoxycyclohexanol, phenylmethanol, (4-chlorophenyl)methanol, phenylethanol, (4-cyanophenyl)-ethanol, phenol, 2-methylphenol, 3-methylphenol, 4-methylphenol, 2,3-dimethylphenol, 2,4-dimethylphenol, 2,5-dimethyl-

phenol, 2,6-dimethylphenol, 3,4-dimethylphenol, 3,5-dimethylphenol, 2-methoxyphenol, 3-methoxyphenol, 4-methoxyphenol, 2-methoxy-3-methylphenol, 2-methoxy-4-methylphenol, 2-methoxy-5-methylphenol,

2-methoxy-6-methylphenol, 3-methoxy-5-methylphenol, 3-methoxy-6-methylphenol, 4-methoxy-5-methylphenol, 4-methoxy-6-methylphenol, 2,3-dimethoxyphenol, 2,4-dimethoxyphenol, 3,5-dimethoxyphenol, 2-chlorophenol, 3-chlorophenol, 4-chlorophenol, 2-bromophenol, 3-bromophenol, 4-bromophenol, 2-cyanophenol, 3-cyanophenol, 4-cyanophenol, 2,3-dichlorophenol, 2,4-dichlorophenol, 2,5-dichlorophenol, 2,6-dichlorophenol, 3,4-dichlorophenol, 3,5-dichlorophenol, 2,3-dibromophenol, 2,4-dibromophenol, 2,5-dibromophenol, 2,6-dibromophenol, 3,4-dibromophenol, 3,5-dibromophenol, 2-methyl-3-chlorophenol, 2-methyl-4-chlorophenol, 2-methyl-5-chlorophenol, 2-methyl-6-chlorophenol, 2-methyl-3-bromophenol, 2-methyl-4-bromophenol, 2-methyl-5-bromophenol, 2-methyl-6-bromophenol, 3-methyl-2-chlorophenol, 3-methyl-4-chlorophenol, 3-methyl-6-chlorophenol, 3-methyl-2-bromophenol, 3-methyl-4-bromophenol, 3-methyl-6-bromophenol, 4-methyl-5-chlorophenol, 4-methyl-6-chlorophenol, 4-methyl-5-bromophenol, 4-methyl-6-bromophenol, 4-chloropyrocatechol-1-methylether, 5-chloropyrocatechol-1-methylether, 4-bromopyrocatechol-1-methylether, 5-bromopyrocatechol-1-methylether, 4-chlororesorcinol-1-methylether, 4-chlororesorcinol-3-methylether, 4-bromoresorcinol-1-methylether, 4-bromoresorcinol-3-methylether, 2-chlorohydroquinone-1-methylether, 2-chlorohydroquinone-2-methylether, 5-chlororesorcinol-1-methylether, 1-naphthol, 2-naphthol, 2-methyl-1-naphthol, 3-methyl-1-naphthol, 4-methyl-1-naphthol, 7-methyl-1-naphthol, 3,6-dimethyl-1-naphthol, 3,7-dimethyl-1-naphthol, 4,6-dimethyl-1-naphthol, 4,7-dimethyl-1-naphthol, 6,7-dimethyl-1-naphthol, 4-methoxy-1-naphthol, 5-methoxy-1-naphthol, 6-methoxy-1-naphthol, 7-methoxy-1-naphthol, 8-methoxy-1-naphthol, 2-chloro-1-naphthol, 3-chloro-1-naphthol, 4-chloro-1-naphthol, 5-chloro-1-naphthol, 7-chloro-1-naphthol, 8-chloro-1-naphthol, 2,3-dichloro-1-naphthol, 2,4-dichloro-1-naphthol, 5,7-dichloro-1-naphthol, 5,8-dichloro-1-naphthol, 2-chloro-4-bromo-1-naphthol, 2-bromo-1-naphthol, 3-bromo-1-naphthol, 4-bromo-1-naphthol, 5-bromo-1-naphthol, 6-bromo-1-naphthol, 7-bromo-1-

naphthol 8-bromo-1-naphthol, 2-ethyl-4-bromo-1-naphthol, 2,4-dibromo-1-naphthol, 1-methyl-2-naphthol, 6-methyl-2-naphthol, 1,4-dimethyl-2-naphthol, 3,6-dimethyl-2-naphthol, 3,7-dimethyl-2-naphthol, 6,7-dimethyl-2-naphthol, 1-chloro-2-naphthol, 3-chloro-2-naphthol, 4-chloro-2-naphthol, 5-chloro-2-naphthol, 6-chloro-2-naphthol, 7-chloro-2-naphthol, 8-chloro-2-naphthol, 1,3-dichloro-2-naphthol, 1,4-dichloro-2-naphthol, 1-bromo-2-naphthol, 3-bromo-2-naphthol, 4-bromo-2-naphthol, 5-bromo-2-naphthol, 6-bromo-2-naphthol, 7-bromo-2-naphthol, 8-bromo-2-naphthol, 1-methyl-6-bromo-2-naphthol, 1-ethyl-6-bromo-2-naphthol, 1,6-dibromo-2-naphthol, 3,6-dibromo-2-naphthol, 3,7-dibromo-2-naphthol and 4,6-dibromo-2-naphthol.

For the preparation of the novel compounds of formula I, wherein the substituent Z_1 and/or Z_2 signifies a $-\text{SO}_2-\text{R}_{12}$ radical, sulfinates are employed, preferably zinc sulfinates of the formula $\text{Zn}(\text{SO}_2-\text{R}_{12})_2$. They allow the introduction of, for example, the following radicals: methylsulfonyl, chloromethylsulfonyl, ethylsulfonyl, 2-chloroethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl, n-amylsulfonyl, isoamylsulfonyl, n-hexylsulfonyl, cyclohexylsulfonyl, benzylsulfonyl, phenylsulfonyl, 2,4-dimethylphenylsulfonyl, 4-methylphenylsulfonyl, 4-methoxyphenylsulfonyl, 4-chlorophenylsulfonyl, 4-chloro-3-methylphenylsulfonyl and 4-bromophenylsulfonyl.

The compounds of the present invention are valuable intermediates, in particular for the preparation of dyestuffs, preferably of azo dyestuffs. As far as these azo dyes do not contain ionogenic groups, they may be employed as disperse dyes for dyeing and printing synthetic hydrophobic fiber material, whereas those azo dyes which do contain ionogenic groups are suited for dyeing and printing cotton, wool, silk, polyamide and modified polyacrylonitrile.


Moreover, the compounds of the present invention are valuable pest control agents — in particular if they are employed as insecticides — as well as valuable pharmaceuticals, above all if they are employed as anesthetics, analgetics and diuretics.

Azo dyes containing the compounds of the present invention may be prepared, for instance, by diazotizing a diazotizable aromatic amine and coupling it with a compound of the present invention. The diazotization and the coupling are carried out in the usual manner. According to the respective initial products employed, this working method leads to the preparation, for instance, of disperse dyes, reactive dyes, cationic dyes or acid dyes. All of these azo dyes exhibit outstanding tinctorial properties, in particular a high coloring strength and a very good fastness to light, perspiration, water, sea water, alkalis and to washing. The disperse dyes distinguish themselves above all by an excellent fastness to dry-heat pleating and setting, ozone, water-gas fading and to rubbing, by a high resistance when subjected to various permanent press treatments and by a good covering power even when applied onto textured polyester materials. The reactive dyes distinguish themselves in particular by an even color depth obtained in the different setting processes. The azo dyes mentioned hereinbefore are suited for dyeing and printing textile materials according to various known processes. The preparation of the dyes as well as their application for dyeing and printing are described in Examples 15 — 19.

In the following examples, "p.b.w." means parts by weight.

EXAMPLE 1

To a composition of 320 p.b.w. ethyl alcohol and 83.7 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine, which are placed in an autoclave, there are added 51.0 p.b.w. liquid ammonia and subsequently the reaction liquid is heated for 18 hours at 200°C . After cooling, the ethyl alcohol is then distilled off, the residue agitated with 300 p.b.w. water, drawn off, washed with water and dried. The resulting 2,6-diamino-3-cyano-4-methylpyridine having the formula




may be purified by crystallization from ethyl alcohol.

Analysis: $\text{C}_7\text{H}_8\text{N}_4$ — Calculated: 37.8% N Found: 37.4% N.

EXAMPLE 2

A solution of 400 p.b.w. ethyl alcohol, 93.0 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine and 274.0 p.b.w. diethylamine is agitated 24 hours at 50°C . Upon cooling of the reaction solution, the resulting 2-chloro-3-cyano-4-methyl-6-diethylaminopyridine of the formula




is crystallized out. The pyridine derivative is then drawn off and washed on the suction apparatus first with 100 p.b.w. ethyl alcohol and subsequently with 500 p.b.w. water. The substance is analytically pure. Its constitution, i.e., the exchange of the 6-position halogen atom in the 2,6-dichloro-3-cyano-4-methylpyridine against the diethylamino group was assured by NMR absorption analysis.

Analysis: $\text{C}_{11}\text{H}_{14}\text{ClN}_3$ — Calculated: 18.8% N, 15.7% Cl. Found: 18.5%, 15.7% Cl.

EXAMPLE 3

The mixture of Example 2 is agitated in an autoclave for 12 hours at 150°C . Upon cooling, the ethyl alcohol is then distilled off and the residue agitated with 50 p.b.w. water and 245 p.b.w. soda lye, 33°Be . The resulting 2,6-bis-(diethylamino)-3-cyano-4-methylpyridine of the formula




is an oil which is isolated with ether. It may be purified by vacuum distillation.

Analysis: $\text{C}_{15}\text{H}_{24}\text{N}_4$ — Calculated: 21.5% N, Found: 21.5% N.

17

EXAMPLE 4

Introduced into 200 p.b.w. methyl alcohol while cooling were 3.7 p.b.w. sodium. Added to this sodium methyl solution then were 33.5 p.b.w. 2-chloro-3-cyano-4-methyl-6-diethylaminopyridine and subsequently the reaction solution was heated to the point of boiling for 24 hours. Then the methyl alcohol was distilled off, the residue taken up with 100 p.b.w. water, drawn off, washed with water and dried. The resulting 2-methoxy-3-cyano-4-methyl-6-diethylaminopyridine of the formula




may be purified by recrystallization from methanol.

Analysis: $C_{12}H_{17}N_3O$ — Calculated: 19.2% N, 14.2% $-OCH_3$. Found: 19.0% N, 13.8% $-OCH_3$.

EXAMPLE 5

A composition consisting of 432 p.b.w. n-propylamine and 187 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine is agitated in an autoclave for 2 hours at 100°C. Upon cooling, the reaction liquor is decom-

posed in water, the resulting 2-chloro-3-cyano-4-methyl-6-n-propylaminopyridine having the formula




is drawn off and washed with water. It may be purified by recrystallization from ethyl alcohol.

Analysis: $C_{10}H_{12}ClN_3$ — Calculated: 20.1% N, 16.7% Cl. Found: 20.3% N, 17.0% Cl.

EXAMPLE 6

A composition of 100 p.b.w. 3-dimethylamino-propylamine(1) and 41.8 p.b.w. 2-chloro-3-cyano-4-methyl-6-n-propylaminopyridine are agitated for 2 hours at 140°C. The reaction melt upon cooling is stirred with water, the residue is drawn off and washed with water. The resulting 2-(3'-dimethylamino-n-propylamino)-3-cyano-4-methyl-6-n-propylaminopyridine having the formula



may be purified by crystallization from ethyl alcohol.


Analysis: $C_{15}H_{25}N_5$ — Calculated: 25.4% N, Found: 25.2% N.

EXAMPLE 7

Introduced into a solution of 155 p.b.w. monomethylamine in 1000 p.b.w. isopropylalcohol are 187 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine. Subsequently,

18

the reaction solution is heated in an autoclave for 5 hours at 200°C. Upon cooling to room temperature, the resulting 2,6-bis-(monomethylamino)-3-cyano-4-methylpyridine having the formula




is drawn off, first washed with isopropyl alcohol, then with water and dried. It may be purified by crystallization from isopropyl alcohol.

Analysis: $C_9H_{12}N_4$ — Calculated: 31.8% N, Found: 32.0% N.

EXAMPLE 8

A composition of 200 p.b.w. ethyl alcohol, 83.7 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine and 180 p.b.w. 3-methoxypropylamine(1) is heated in an autoclave for 18 hours at 180°C. Upon cooling, the ethyl alcohol is distilled off, the residue is stirred with 100 p.b.w. water with the addition of 150 p.b.w. soda lye, 33°Be, drawn off, washed with water and dried. The resulting 2,6-bis-(3'-methoxy-n-propylamino)-3-cyano-4-methylpyridine having the formula




may be purified by vacuum distillation.

Analysis: $C_{15}H_{24}N_4O_2$ — Calculated: 19.2% N, Found: 19.6% N.

EXAMPLE 9

A composition consisting of 100 p.b.w. morpholine and 37.4 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine is heated for 30 minutes at 130°C. The reaction melt is then decomposed on ice the residue is drawn off, washed with water, and dried. The resulting 2,6-bis-(morpholino)-3-cyano-4-methylpyridine having the formula



may be purified by crystallization from ethyl alcohol.


Analysis: $C_{15}H_{20}N_4O_2$ — Calculated: 19.4% N, Found: 19.7% N.

EXAMPLE 10

There are introduced 37.2 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine into 150.0 p.b.w. aniline. This reaction solution is then heated for 24 hours at 150°C.

19

and subsequently decomposed in 1,000 p.b.w. ice with the addition of 350 p.b.w. crude hydrochloric acid ($D = 1.153$). The resulting 2,6-bis-(anilino)-3-cyano-4-methylpyridine having the formula




may be purified by crystallization from ethyl alcohol.

Analysis: $C_{19}H_{16}N_4$ — Calculated: 18.7% N, Found: 18.5% N

EXAMPLE 11

Introduced into 80 p.b.w. methyl alcohol are 7.6 p.b.w. sodium during cooling, then there are added 27.9 p.b.w. 2,6-dichloro-3-cyano-4-methylpyridine to this sodium methylate solution and subsequently the reaction solution is heated to the point of boiling for 24 hours. Then the methyl alcohol is distilled off. The residue is taken up with 100 p.b.w. water, drawn off, washed with water and dried. The resulting 2,6-dimethoxy-3-cyano-4-methylpyridine having the formula




may be purified by recrystallization from methyl alcohol.

20

Analysis: $C_9H_{10}N_2O_2$ — Calculated: 15.7% N, 34.8% —OCH₃. Found: 15.5% N, 34.2% —OCH₃

EXAMPLE 12


- 5 There are introduced 27.9 p.b.w. of 2,6-dichloro-3-cyano-4-methylpyridine into a sodium ethylate solution, which had been prepared from 80 p.b.w. ethyl alcohol and 3.8 p.b.w. sodium. Subsequently, the reaction solution is heated to the point of boiling for 24
- 10 hours. The ethyl alcohol is then distilled off, the residue is stirred with 100 p.b.w. water, and the resulting 2-chloro-3-cyano-4-methyl-4-ethoxypyridine having the formula



is taken up with ether. It may be purified by vacuum distillation.

Analysis: $C_9H_9ClN_2O$ — Calculated: 14.3% N, 17.8% Cl. Found: 14.5% N, 17.6% Cl.

In the following table there are presented further pyridine compounds prepared in accordance with the herein described process:

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
1.	—NHCH ₂ CH ₃	—NHCH ₂ CH ₃	C ₁₁ H ₁₆ N ₄	27.5	27.5
2.	—NHCH ₂ CH ₂ OH	—NHCH ₂ CH ₂ OH	C ₁₁ H ₁₆ N ₄ O ₂	23.7	24.0
3.	—NHCH ₂ CH ₂ OCH ₃	—NHCH ₂ CH ₂ OCH ₃	C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.0
4.	—NHCH ₂ CH ₂ CN	—NHCH ₂ CH ₂ CN	C ₁₃ H ₁₄ N ₆	33.1	33.5
5.	—NHCH ₂ CH ₂ CH ₃	—NHCH ₂ CH ₂ CH ₃	C ₁₃ H ₂₀ N ₄	24.1	24.6
6.	—NH—CH(CH ₃) CH ₃	—NH—CH(CH ₃) CH ₃	C ₁₃ H ₂₀ N ₄	24.1	24.4
7.	—NH—(CH ₂) ₂ CH ₂ OCH ₃	—NH—(CH ₂) ₂ CH ₂ OCH ₃	C ₁₅ H ₂₄ N ₄ O ₂	19.2	19.0
8.	—NHCH ₂ CH=CH ₂	—NHCH ₂ CH=CH ₂	C ₁₃ H ₁₆ N ₄	24.6	24.2
9.	—NH—C ₆ H ₉ (n)	—NH—C ₆ H ₉ (n)	C ₁₅ H ₂₄ N ₄	21.5	21.8
10.	—NH—C ₆ H ₉ (sec)	—NH—C ₆ H ₉ (sec)	C ₁₅ H ₂₄ N ₄	21.5	21.3
11.	—NH—CH(CH ₂ —CH ₃) CH ₂ —CH ₃	—NH—CH(CH ₂ —CH ₃) CH ₂ —CH ₃	C ₁₇ H ₂₈ N ₄	19.4	19.8
12.	—NH—C ₈ H ₁₁ (iso)	—NH—C ₈ H ₁₁ (iso)	C ₁₇ H ₂₈ N ₄	19.4	19.6
13.	—NH—C ₈ H ₁₃ (n)	—NH—C ₈ H ₁₃ (n)	C ₁₉ H ₃₄ N ₄	17.7	18.1
14.	—NH— 	—NH— 	C ₁₉ H ₂₈ N ₄	17.9	17.5
15.	—NH—CH ₂ — 	—NH—CH ₂ — 	C ₂₁ H ₂₀ N ₄	17.1	17.5
16.	—NH—CH ₂ —CH ₂ — 	—NH—CH ₂ —CH ₂ — 	C ₂₃ H ₂₄ N ₄	15.7	15.5
17.	—NH— 	—NH— 	C ₂₇ H ₂₀ N ₄	14.0	14.3

or tautomeric formulae.















-continued-

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
18.			C ₁₇ H ₁₆ N ₄ O ₂	18.2	18.5
19.			C ₁₉ H ₁₈ N ₆	25.5	25.7
20.			C ₁₅ H ₂₀ N ₄ O ₄ S ₂	14.6	15.0
21.			C ₂₅ H ₂₂ N ₆	20.7	21.0
22.			C ₁₁ H ₁₆ N ₄	27.5	27.2
23.			C ₁₉ H ₂₀ N ₈	31.1	36.7
24.			C ₁₇ H ₂₂ N ₆ O ₂	26.3	26.8
25.			C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.6
26.			C ₁₉ H ₃₂ N ₄ O ₂	16.1	16.5
27.			C ₂₃ H ₃₆ N ₄ O ₂	14.0	14.6
28.			C ₂₅ H ₂₈ N ₄ O ₂	13.5	13.2
29.			C ₁₉ H ₃₂ N ₄	17.7	18.2
30.			C ₂₃ H ₄₀ N ₄	15.1	15.5
31.			C ₃₁ H ₅₆ N ₄	11.6	11.2
32.			C ₁₅ H ₂₀ N ₄ O ₂	19.4	19.6
33.			C ₁₇ H ₂₄ N ₄	19.7	19.3
34.			C ₁₇ H ₂₆ N ₆	26.8	26.2
35.			C ₁₉ H ₂₀ N ₆	24.6	25.1
36.			C ₁₅ H ₂₀ N ₄	21.8	22.2














-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
37.			C ₂₁ H ₂₂ N ₆	23.5	23.8
38.			C ₁₁ H ₁₈ N ₆	35.9	35.6
39.			C ₁₃ H ₂₂ N ₆	32.1	32.5
40.			C ₁₃ H ₂₆ N ₆	29.0	22.4
41.			C ₂₃ H ₂₆ N ₆	21.8	21.5
42.			C ₁₅ H ₂₂ N ₆ O ₂	26.4	26.2
43.			C ₁₇ H ₃₀ N ₆	26.4	26.0
44.			C ₁₉ H ₃₄ N ₆	24.3	24.5
45.			C ₁₇ H ₃₀ N ₆	26.4	26.1
46.			C ₂₁ H ₃₄ N ₆ O ₂	20.9	20.7
47.			C ₂₃ H ₃₈ N ₆	21.1	21.5
48.			C ₇ H ₈ N ₄ O ₂	31.1	31.6
49.			C ₁₁ H ₁₆ N ₄ O ₂	23.7	24.2
50.			C ₁₃ H ₂₀ N ₄ O ₄	18.9	19.3
51.			C ₂₃ H ₂₄ N ₄ O ₂	14.4	15.0
52.			C ₁₅ H ₂₄ N ₄ O ₂	19.2	19.6
53.			C ₁₃ H ₂₄ N ₄ O ₆	15.7	16.2
54.			C ₂₅ H ₂₈ N ₄ O ₄	12.5	12.2
55.			C ₁₃ H ₁₈ N ₄ O ₂	21.5	21.2
56.			C ₁₅ H ₂₀ N ₄ O ₂	19.4	19.8
57.			C ₇ H ₈ ClN ₃	25.1	25.4
58.			C ₈ H ₈ N ₄	35.7	35.5
59.			C ₇ H ₇ N ₃ O	28.2	28.6
60.			C ₈ H ₉ N ₃ O	25.8	25.2


-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
61.	-NH ₂		C ₁₃ H ₁₁ N ₃ O	18.7	19.1
62.	-NH ₂	-SH	C ₇ H ₇ N ₃ S	25.5	25.7
63.	-NH ₂	-SC ₂ H ₅	C ₉ H ₁₁ N ₃ S	21.8	22.3
64.	-NH ₂	-SO ₂ -CH ₃	C ₈ H ₉ N ₃ O ₂ S	19.9	20.5
65.	-NH ₂	-NH-CH ₃	C ₈ H ₁₀ N ₄	34.6	34.2
66.	-NH ₂	-NH-C ₂ H ₅	C ₉ H ₁₂ N ₄	31.8	31.5
67.	-NH ₂	-NHCH ₂ CH ₂ CH ₂ OCH ₃	C ₁₁ H ₁₆ N ₄ O	25.5	25.2
68.	-NH ₂		C ₁₃ H ₁₂ N ₄	25.0	25.5
69.	-NH ₂	-NH-CH ₂ - 	C ₁₄ H ₁₄ N ₄	23.6	24.1
70.	-NH ₂	 CH ₂ -NH-	C ₁₆ H ₁₄ N ₅	25.4	25.0
71.	-NH ₂		C ₉ H ₁₂ N ₄	31.8	31.6
72.	-NH ₂		C ₁₀ H ₁₄ ON ₄	27.2	27.7
73.	-NH ₂		C ₁₁ H ₁₆ O ₂ N ₄	23.7	24.2
74.	-NH ₂		C ₁₁ H ₁₄ ON ₄	25.7	26.2
75.	-NH ₂		C ₁₂ H ₁₈ N ₄	25.9	26.3
76.	-NH ₂		C ₉ H ₁₃ N ₅	36.6	36.1
77.	-NH ₂		C ₁₃ H ₂₁ N ₅	28.3	28.1
78.	-NH ₂		C ₁₁ H ₁₅ N ₅	32.3	32.2
79.	-NH ₂		C ₉ H ₁₂ ON ₄	29.2	29.0
80.	-NH ₂		C ₁₂ H ₁₈ ON ₄	23.9	23.3
81.	-NH-CH ₃	-Cl	C ₈ H ₈ N ₂ Cl	23.2	23.5
82.	-NH-CH ₃	-CN	C ₉ H ₈ N ₄	32.6	32.2
83.	-NH-CH ₃	-OH	C ₈ H ₈ ON ₃	25.8	26.2
84.	-NH-CH ₃	-OC ₂ H ₅	C ₁₀ H ₁₃ ON ₃	22.0	22.5
85.	-NH-CH ₃	-SO ₂ -C ₂ H ₅	C ₁₀ H ₁₃ N ₃ O ₂ S	17.6	17.5
86.	-NH-CH ₃	-NH-C ₃ H ₇ (n)	C ₁₁ H ₁₆ N ₄	27.5	27.1
87.	-NH-CH ₃	-NH-C ₃ H ₇ (iso)	C ₁₃ H ₂₀ N ₄	24.1	23.5
88.	-NH-CH ₃		C ₁₄ H ₂₀ N ₄	23.0	22.5
89.	-NH-CH ₃	-NH-CH ₂ - 	C ₁₅ H ₁₆ N ₄	22.2	22.0
90.	-NH-CH ₃		C ₁₈ H ₁₆ N ₄	19.4	19.0
91.	-NH-CH ₃		C ₁₃ H ₁₅ N ₄ O	23.0	22.5

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
92.	-NH-CH ₃		C ₁₂ H ₁₈ N ₄	25.7	26.1
93.	-NH-CH ₃		C ₁₃ H ₁₇ N ₃ O	27.0	26.5
94.	-NH-CH ₃		C ₁₂ H ₁₈ N ₄	25.9	26.4
95.	-NH-CH ₃		C ₁₃ H ₁₉ N ₅	28.6	28.2
96.	-NH-CH ₃		C ₁₂ H ₁₉ N ₅	30.0	30.6
97.	-NH-CH ₃		C ₁₂ H ₁₉ N ₅	30.0	30.3
98.	-NH-CH ₃		C ₁₂ H ₁₈ N ₄ O ₂	22.4	22.0
99.	-NH-C ₂ H ₅	-Cl	C ₉ H ₁₀ ClN ₃	21.5	21.3
100.	-NH-C ₂ H ₅	-CN	C ₁₀ H ₁₀ N ₄	30.1	30.5
101.	-NH-C ₂ H ₅	-OH	C ₉ H ₁₁ N ₃ O	23.7	23.2
102.	-NH-C ₂ H ₅	-OC ₃ H ₇ (n)	C ₁₂ H ₁₇ N ₃ O	19.2	19.0
103.	-NH-C ₂ H ₅	-SCH ₃	C ₁₀ H ₁₃ N ₃ S	20.3	20.7
104.	-NH-C ₂ H ₅	-SO ₂ -C ₃ H ₇ (n)	C ₁₂ H ₁₇ N ₃ O ₂ S	15.7	15.5
105.	-NH-C ₂ H ₅	-NH-CH ₃	C ₁₀ H ₁₄ N ₄	29.5	29.0
106.	-NH-C ₂ H ₅	-NH-CH ₂ -CH ₂ -OCH ₃	C ₁₂ H ₁₈ N ₄ O	23.9	23.3
107.	-NH-C ₂ H ₅	-NH-C ₆ H ₁₃ (n)	C ₁₅ H ₂₄ N ₄	21.5	21.0
108.	-NH-C ₂ H ₅		C ₁₁ H ₁₆ N ₄	27.5	27.3
109.	-NH-C ₂ H ₅		C ₁₄ H ₂₀ N ₄	23.0	22.5
110.	-NH-C ₂ H ₅		C ₁₇ H ₁₇ N ₅	32.0	31.6
111.	-NH-C ₂ H ₅		C ₁₅ H ₂₃ N ₅	25.5	26.0
112.	-NH-C ₂ H ₅		C ₂₃ H ₂₄ N ₄ O	15.1	15.5
113.	-NH-C ₃ H ₇ (n)	-OH	C ₁₀ H ₁₃ N ₃ O	22.0	22.4
114.	-NH-C ₃ H ₇ (n)	-O-C ₃ H ₇ (iso)	C ₁₃ H ₁₉ N ₃ O	18.0	18.5
115.	-NH-C ₃ H ₇ (n)	-SO ₂ -C ₃ H ₇ (iso)	C ₁₅ H ₂₃ N ₃ O ₂ S	13.6	13.3
116.	-NH-C ₃ H ₇ (n)	-NH-CH ₃	C ₁₁ H ₁₆ N ₄	27.5	28.0
117.	-NH-C ₃ H ₇ (n)	-NH-C ₃ H ₇ (iso)	C ₁₃ H ₂₀ N ₄	24.1	24.5
118.	-NH-C ₃ H ₇ (n)		C ₁₆ H ₂₄ N ₄	20.6	21.0
119.	-NH-C ₃ H ₇ (n)		C ₁₆ H ₁₈ N ₄	21.1	21.5
120.	-NH-C ₃ H ₇ (n)		C ₁₂ H ₁₈ N ₄	25.7	26.0
121.	-NH-C ₃ H ₇ (n)		C ₁₉ H ₂₄ N ₄ O	17.3	17.5
122.	-NH-C ₃ H ₇ (n)		C ₁₄ H ₂₁ N ₅ O	25.5	26.1

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
123.	—NH—C ₃ H ₇ (n)	—NHCH ₂ CH ₂ N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₄ H ₂₃ N ₅	26.8	26.4
124.	—NH—C ₃ H ₇ (n)	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{OCH}_3 \end{smallmatrix}$	C ₁₂ H ₁₈ N ₄ O	23.9	23.2
125.	—NH—C ₃ H ₇ (iso)	—NHCH ₂ CH ₂ CH ₂ OCH ₃	C ₁₄ H ₂₂ N ₄ O	21.4	21.0
126.	—NH—C ₃ H ₇ (iso)	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₄ H ₂₂ N ₄	22.8	23.0
127.	—NH—C ₄ H ₉ (n)	—Cl	C ₁₁ H ₁₄ ClN ₃	18.8	19.0
128.	—NH—C ₄ H ₉ (n)	—OH	C ₁₁ H ₁₅ N ₃ O	20.5	21.0
129.	—NH—C ₄ H ₉ (n)	—O—C ₂ H ₅ (n)	C ₁₅ H ₂₃ N ₃ O	16.1	16.4
130.	—NH—C ₄ H ₉ (n)	—NH—CH ₃	C ₁₂ H ₁₈ N ₄	25.7	25.0
131.	—NH—C ₄ H ₉ (n)	—NHCH ₂ CH ₂ OH	C ₁₃ H ₂₀ N ₄ O	22.6	23.0
132.	—NH—C ₄ H ₉ (iso)	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₃ H ₂₀ N ₄	24.1	23.8
133.	—NH—C ₄ H ₉ (iso)		C ₁₆ H ₂₅ N ₅	24.4	24.1
134.	—NH—C ₄ H ₉ (sek)	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₅ H ₂₄ N ₄	21.5	22.0
135.	—NH—C ₄ H ₉ (sek)	—NH—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₃ H ₂₁ N ₅	28.3	28.5
136.	—NH—C ₄ H ₉ (sek)	—N $\begin{smallmatrix} \text{CH}_2\text{CH}_2\text{OH} \\ \text{O—CH}_2\text{CH}_2\text{CH}_3 \end{smallmatrix}$	C ₁₆ H ₂₆ N ₄ O ₂	18.3	18.7
137.	—NH—C ₄ H ₉ (tert)	—NH—C ₂ H ₅	C ₁₃ H ₂₀ N ₄	24.1	24.5
138.	—NH—C ₄ H ₉ (tert)	—NH—CH ₂ —CH ₂ —CN	C ₁₄ H ₁₉ N ₅	27.2	27.5
139.	—NH—C ₅ H ₁₁ (n)	—NH—CH ₃	C ₁₃ H ₂₀ N ₄	24.1	24.5
140.	—NH—C ₅ H ₁₁ (n)	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₄ H ₂₂ N ₄	22.8	23.0
141.	—NH—C ₅ H ₁₁ (n)		C ₁₈ H ₂₃ N ₅	22.7	23.2
142.	—NH—C ₅ H ₁₁ (iso)	—NH—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₄ H ₂₃ N ₅	26.8	27.2
143.	—NH—C ₆ H ₁₃ (n)	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₇ H ₂₆ N ₄	19.4	19.0
144.	—NH—C ₆ H ₁₃ (n)		C ₁₂ H ₂₆ N ₄ O	18.5	18.9
145.	—NH—C ₆ H ₁₃ (n)	—NHCH ₂ CH ₂ N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₇ H ₂₉ N ₅	23.1	23.3
146.	—NHCH ₂ CH ₂ OH	—NH—CH ₃	rCH ₃ C ₁₀ H ₁₄ N ₄ O	27.2	27.5
147.	—NHCH ₂ CH ₂ OH	—NHCH ₂ CH ₂ CH ₂ OCH ₃	C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.5
148.	—NHCH ₂ CH ₂ OH	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₁ H ₁₆ N ₄ O	25.5	26.0
149.	—NHCH ₂ CH ₂ OH		C ₁₃ H ₁₈ N ₄ O ₂	21.4	21.7
150.	—NHCH ₂ CH ₂ OH		C ₁₄ H ₂₀ N ₄ O	21.5	21.8
151.	—NHCH ₂ CH ₂ OH	—NHCH ₂ CH ₂ N $\begin{smallmatrix} \text{CH}_2 \\ \text{CH}_2 \end{smallmatrix}$	C ₁₃ H ₁₉ N ₅ O	26.8	27.2
152.	—NHCH ₂ CH ₂ OCH ₃	—NH—CH ₃	C ₁₁ H ₁₆ N ₄ O	25.5	25.8
153.	—NHCH ₂ CH ₂ OCH ₂	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₂ H ₁₈ N ₄ O	23.9	24.3
154.	—NHCH ₂ CH ₂ O 	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₈ H ₂₄ N ₄ O	17.3	17.5
155.	—NHCH ₂ CH ₂ CH ₂ CN	—NH—C ₂ H ₅	C ₁₃ H ₁₇ N ₅	28.8	28.2
156.	—NHCH ₂ CH ₂ CH ₂ CN	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_2\text{CH}_2\text{OH} \end{smallmatrix}$	C ₁₄ H ₁₉ N ₅ O	25.6	26.0

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
157.	$-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{CN}$	$-\text{N} \begin{array}{l} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_{15}\text{H}_{21}\text{N}_5$	25.8	26.3
158.	$-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{CN}$	$-\text{NH}-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{C}_6\text{H}_5 \end{array}$	$\text{C}_{18}\text{H}_{20}\text{N}_6$	26.2	27.0
159.	$-\text{NH}-\text{C}_6\text{H}_4-\text{C}_6\text{H}_5$	$-\text{NH}-\text{C}_6\text{H}_4-\text{C}_6\text{H}_5$	$\text{C}_{27}\text{H}_{20}\text{N}_4$	14.0	13.5
160.	$-\text{NH}-\text{C}_2\text{H}_5$	$-\text{NH}-\text{C}_6\text{H}_{11}$	$\text{C}_{15}\text{H}_{22}\text{N}_4$	21.7	22.2
161.	$-\text{NH}-\text{C}_2\text{H}_5$	$-\text{NH}-\text{CH}_2-\text{CH}_2-\text{C}_6\text{H}_5$	$\text{C}_{17}\text{H}_{20}\text{N}_4$	20.0	20.4
162.	$-\text{NH}-\text{C}_2\text{H}_5$	$-\text{NH}-\text{C}_6\text{H}_4-\text{CH}_3$	$\text{C}_{16}\text{H}_{18}\text{N}_4$	21.1	21.5
163.	$-\text{NH}-\text{C}_6\text{H}_9(\text{n})$	$-\text{NH}-\text{C}_6\text{H}_4-\text{OCH}_3$	$\text{C}_{18}\text{H}_{22}\text{N}_4\text{O}$	18.1	18.6
164.	$-\text{NH}-\text{C}_6\text{H}_{11}$	$-\text{NH}-\text{C}_6\text{H}_4-\text{CN}$	$\text{C}_{20}\text{H}_{21}\text{N}_5$	21.1	21.7
165.	"	$-\text{NH}_2$	$\text{C}_{13}\text{H}_{18}\text{N}_4$	24.3	24.5
166.	"	$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{20}\text{N}_4$	23.0	23.5
167.	$-\text{NH}-\text{C}_6\text{H}_{11}$	$-\text{NH}-\text{CH}_2-\text{CH}=\text{CH}_2$	$\text{C}_{16}\text{H}_{22}\text{N}_4$	20.7	21.0
168.	"	$-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$	$\text{C}_{17}\text{H}_{26}\text{N}_4\text{O}$	19.2	19.5
169.	"	$-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{15}\text{H}_{22}\text{N}_4$	21.7	22.2
170.	"	$-\text{NH}-\text{N} \begin{array}{l} \text{H} \\ \text{CH}_3 \end{array}$	$\text{C}_{18}\text{H}_{24}\text{N}_6$	25.6	26.0
171.	"	$-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{19}\text{H}_{29}\text{N}_5$	22.2	22.5
172.	$-\text{NH}-\text{C}_6\text{H}_{11}$	$\text{H}_2\text{C} \begin{array}{l} \text{C} \\ \text{CH}_2 \\ \text{CH}_2 \\ \text{O} \end{array}$	$\text{C}_{17}\text{H}_{24}\text{N}_4\text{O}$	18.7	19.1
173.	$-\text{NH}-\text{CH}_2-\text{C}_6\text{H}_5$	$-\text{NH}_2$	$\text{C}_{14}\text{H}_{14}\text{N}_4$	23.5	24.0
174.	"	$-\text{NH}-\text{C}_2\text{H}_5$	$\text{C}_{16}\text{H}_{18}\text{N}_4$	21.2	22.0
175.	"	$-\text{N} \begin{array}{l} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_{18}\text{H}_{22}\text{N}_4$	19.0	19.3
176.	"	$-\text{NHCH}_2\text{CH}_2\text{OCH}_3$	$\text{C}_{17}\text{H}_{20}\text{N}_4\text{O}$	18.9	18.4
177.	"	$-\text{NHCH}_2\text{CH}_2\text{N} \begin{array}{l} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{array}$	$\text{C}_{20}\text{H}_{27}\text{N}_5$	20.8	21.4
178.	$-\text{NH}-\text{CH}_2-\text{C}_6\text{H}_5$	$-\text{NH}-\text{CH}_2-\text{CH}_2-\text{C}_6\text{H}_4-\text{N}$	$\text{C}_{21}\text{H}_{21}\text{N}_5$	20.4	20.6
179.	$-\text{NH}-\text{CH}_2-\text{C}_6\text{H}_5$	$-\text{N} \begin{array}{l} \text{CH}_2 \\ \text{OCH}_3 \end{array}-\text{C}_6\text{H}_5$	$\text{C}_{22}\text{H}_{22}\text{N}_4\text{O}$	15.6	16.0

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
180.		$-\text{NH}-\text{CH}_3$	$\text{C}_{16}\text{H}_{18}\text{N}_4$	21.1	21.5
181.	"		$\text{C}_{17}\text{H}_{20}\text{N}_4$	20.0	20.5
182.	"		$\text{C}_{17}\text{H}_{23}\text{N}_4\text{O}$	18.7	19.0
183.		$-\text{NH}_2$	$\text{C}_{13}\text{H}_{12}\text{N}_4$	25.0	24.7
184.	"	$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{14}\text{N}_4$	23.6	24.2
185.	"	$-\text{NH}-\text{C}_3\text{H}_7(\text{n})$	$\text{C}_{16}\text{H}_{18}\text{N}_4$	21.1	21.3
186.	"	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{15}\text{H}_{16}\text{N}_4\text{O}$	20.9	21.5
187.			$\text{C}_{15}\text{H}_{16}\text{N}_4$	22.2	22.0
188.	"		$\text{C}_{19}\text{H}_{18}\text{N}_6$	25.5	25.9
189.			$\text{C}_{17}\text{H}_{18}\text{N}_4\text{O}$	19.1	19.7
190.	"		$\text{C}_{17}\text{H}_{21}\text{N}_5$	23.7	24.3
191.	"		$\text{C}_{18}\text{H}_{23}\text{N}_5$	22.7	22.3
192.	"		$\text{C}_{15}\text{H}_{16}\text{N}_4\text{O}$	20.9	21.4
193.		$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{13}\text{ClN}_4$	20.6	20.2
194.		$-\text{NH}-\text{CH}_3$	$\text{C}_{15}\text{H}_{16}\text{N}_4$	22.2	22.8
195.			$\text{C}_{17}\text{H}_{20}\text{N}_4$	20.0	20.5
196.		"	$\text{C}_{16}\text{H}_{18}\text{N}_4\text{O}$	19.9	20.3
197.		$-\text{NH}-\text{CH}_3$	$\text{C}_{13}\text{H}_{14}\text{N}_4\text{O}$	23.1	23.5
198.	"		$\text{C}_{14}\text{H}_{16}\text{N}_4\text{O}$	21.9	21.5
199.	"		$\text{C}_{19}\text{H}_{27}\text{N}_5\text{O}$	20.5	20.9
200.		$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{15}\text{N}_5$	27.7	27.2
201.		$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{15}\text{N}_5$	27.7	28.2






















-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
202.		-NH-CH ₃	C ₁₅ H ₁₇ N ₅	26.2	26.0
203.		-NH-CH ₃	C ₁₅ H ₁₇ N ₅	26.2	26.6
204.		-NH-CH ₃	C ₁₂ H ₁₆ N ₄ O ₂ S	20.0	20.5
205.	"		C ₁₃ H ₁₈ N ₄ O ₂ S	19.0	18.7
206.	"		C ₁₅ H ₂₀ N ₄ O ₃ S	16.7	17.1
207.	"		C ₁₃ H ₁₉ N ₅ O ₂ S	22.7	22.2
208.		-NH-CH ₃	C ₁₇ H ₁₇ N ₅	24.1	24.5
209.	"		C ₁₈ H ₁₉ N ₅	23.0	23.5
210.	"		C ₂₀ H ₂₁ N ₅ O	20.2	20.5
211.		-Cl	C ₁₃ H ₁₆ ClN ₃	16.9	16.5
212.	"	-OH	C ₁₃ H ₁₇ N ₃ O	18.2	18.6
213.	"	-OCH ₃	C ₁₄ H ₁₉ N ₃ O	17.1	17.6
214.	"	-CN	C ₁₄ H ₁₆ N ₄	25.0	25.3
215.	"		C ₂₀ H ₂₃ N ₃ O ₂ S	11.4	11.2
216.		-Cl	C ₁₄ H ₁₂ ClN ₃	16.3	16.0
217.	"	-OH	C ₁₄ H ₁₃ N ₃ O	17.6	18.0
218.	"	-O-CH ₃	C ₁₅ H ₁₃ N ₃ O	16.6	16.0
219.	"	-S-C ₆ H ₅ (n)	C ₁₈ H ₂₁ N ₃ S	21.5	21.8
220.		-Cl	C ₁₃ H ₁₀ ClN ₃	17.3	17.0
221.		-CN	C ₁₄ H ₁₀ N ₄	23.9	24.3
222.	"	-OH	C ₁₃ H ₁₁ N ₃ O	18.7	18.5
223.	"	-OCH ₃	C ₁₄ H ₁₃ N ₃ O	17.6	17.3
224.	"	-SCH ₃	C ₁₄ H ₁₃ N ₃ S	16.5	16.0
225.		-Cl	C ₉ H ₁₀ ClN ₃	21.5	21.0
226.	"	-CN	C ₁₀ H ₁₀ N ₄	30.1	30.5
227.	"	-OH	C ₉ H ₁₁ N ₃ O	23.7	23.4
228.	"	-OCH ₃	C ₁₀ H ₁₃ N ₃ O	22.0	22.3
229.	"	-SO ₂ -C ₂ H ₅	C ₁₁ H ₁₃ N ₃ O ₂ S	16.6	17.0
230.	"	-NH-CH ₃	C ₁₀ H ₁₄ N ₄	29.5	29.1
231.	"	-NH-C ₃ H ₇ (n)	C ₁₂ H ₁₆ N ₄	25.7	26.2
232.	"	-NH-C ₆ H ₁₃ (n)	C ₁₅ H ₂₄ N ₄	21.5	21.8

-continued-

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
233.			C ₁₅ H ₂₂ N ₄	21.7	21.5
234.	"		C ₁₆ H ₁₈ N ₄	21.1	20.7
235.	"		C ₁₅ H ₁₆ N ₄	22.2	22.6
236.	"		C ₁₉ H ₁₈ N ₄	18.5	18.9
237.	"		C ₁₄ H ₁₆ N ₄ O	21.9	21.5
238.	"		C ₁₃ H ₂₀ N ₄	24.1	23.8
239.			C ₁₄ H ₁₉ N ₅ O	25.6	26.0
240.	"		C ₁₃ H ₁₈ N ₄	24.3	24.5
241.			C ₁₄ H ₂₁ N ₅	27.0	27.4
242.	"		C ₁₃ H ₂₁ N ₅	28.3	27.9
243.	"		C ₁₄ H ₂₃ N ₅	26.8	26.6
244.	"		C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.6
245.		-Cl	C ₁₁ H ₁₄ ClN ₃	18.8	18.5
246.	"	-CN	C ₁₂ H ₁₄ N ₄	26.2	26.6
247.	"	-OH	C ₁₁ H ₁₅ N ₃ O	20.5	21.0
248.	"	-OCH ₃	C ₁₂ H ₁₇ N ₃ O	19.2	19.0
249.	"	-SC ₂ H ₅	C ₁₃ H ₁₉ N ₃ S	16.8	17.2
250.		-SO ₂ -C ₄ H ₉ (n)	C ₁₅ H ₂₃ N ₃ SO ₂	13.6	13.5
251.		-NH-CH ₃	C ₁₂ H ₁₇ N ₄	25.7	25.5
252.	"	-NHCH ₂ CH ₂ CH ₂ OCH ₃	C ₁₅ H ₂₄ N ₄ O	20.3	20.0
253.	"		C ₁₇ H ₂₅ N ₄	19.6	19.5
254.	"		C ₁₈ H ₂₂ N ₄	19.0	19.4
255.	"		C ₁₇ H ₂₀ N ₄	20.0	20.2
256.	"		C ₁₆ H ₂₄ N ₄	20.6	21.0
257.	"		C ₁₃ H ₂₁ N ₅	28.3	28.0
258.	"		C ₁₆ H ₂₇ N ₅	24.2	24.5

-continued

No.	Z ₁	Z ₂	Empirical Formula	Calcd. N%	Found N%
259.			C ₁₃ H ₂₀ N ₄ O	22.6	22.2
260.			C ₁₃ H ₁₆ N ₄ O ₂	21.5	21.3
261.	"		C ₁₆ H ₂₁ N ₅ O ₂	22.2	22.6
262.			C ₁₁ H ₁₆ N ₄ O ₂	23.7	23.4
263.			C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.5
264.	"		C ₁₇ H ₂₀ N ₄ O ₂	17.9	18.2
265.			C ₁₄ H ₁₃ N ₅ O	26.0	26.3
266.	"		C ₁₅ H ₁₈ N ₆	29.8	29.5
267.			C ₁₀ H ₁₄ N ₄ O	27.2	27.5
268.			C ₁₁ H ₁₆ N ₄ O	25.5	25.8
269.	"		C ₁₆ H ₂₄ N ₅ O	23.2	23.7
270.	"		C ₁₆ H ₁₈ N ₄ O	19.9	19.5
271.	"		C ₁₂ H ₁₉ N ₅ O	28.1	27.8
272.			C ₁₆ H ₂₆ N ₄ O ₂	13.8	14.2
273.			C ₁₄ H ₂₂ N ₄ O	21.4	21.0
274.			C ₁₇ H ₂₆ N ₄ O ₂	17.6	18.0
275.			C ₁₇ H ₂₈ N ₅ O	21.9	21.7
276.			C ₁₆ H ₂₃ N ₄ O	19.4	19.0
277.			C ₂₁ H ₃₂ N ₄ O	15.7	16.3
278.			C ₁₆ H ₁₈ N ₄ O	19.9	19.5

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
279.		$-\text{NH}-\text{CH}_3$	$\text{C}_{17}\text{H}_{20}\text{N}_4\text{O}$	18.9	19.5
280.	"	$-\text{NHCH}_2\text{CH}_2-$	$\text{C}_{23}\text{H}_{25}\text{N}_5\text{O}$	18.1	18.5
281.	$-\text{N} \begin{cases} \text{C}_3\text{H}_7(\text{n}) \\ \text{C}_3\text{H}_7(\text{n}) \end{cases}$	$-\text{Cl}$	$\text{C}_{13}\text{H}_{18}\text{ClN}_3$	16.7	16.5
282.	"	$-\text{NH}_2$	$\text{C}_{13}\text{H}_{20}\text{N}_4$	24.1	24.5
283.	"	$-\text{CN}$	$\text{C}_{14}\text{H}_{18}\text{N}_4$	23.1	23.7
284.	$-\text{N} \begin{cases} \text{C}_3\text{H}_7(\text{n}) \\ \text{C}_3\text{H}_7(\text{n}) \end{cases}$	$-\text{OH}$	$\text{C}_{13}\text{H}_{19}\text{N}_3\text{O}$	18.0	18.4
285.	"	$-\text{OCH}_3$	$\text{C}_{14}\text{H}_{21}\text{N}_3\text{O}$	17.0	17.5
286.	"	$-\text{SO}_2-\text{CH}_3$	$\text{C}_{14}\text{H}_{21}\text{N}_3\text{O}_2\text{S}$	14.2	13.8
287.	"	$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{22}\text{N}_4$	22.8	23.2
288.	"	$-\text{NH}-\text{C}_3\text{H}_7(\text{n})$	$\text{C}_{16}\text{H}_{26}\text{N}_4$	20.5	21.0
289.	"	$-\text{NH}-$	$\text{C}_{19}\text{H}_{30}\text{N}_4$	17.8	17.5
290.	"	$-\text{NH}-$	$\text{C}_{19}\text{H}_{24}\text{N}_4$	18.2	18.5
291.	"	$-\text{N} \begin{cases} \text{CH}_3 \\ \text{CH}_3 \end{cases}$	$\text{C}_{15}\text{H}_{24}\text{N}_4$	21.5	20.8
292.	"	$-\text{NH}-$	$\text{C}_{17}\text{H}_{27}\text{N}_5\text{O}$	22.1	22.5
293.	$-\text{N} \begin{cases} \text{C}_3\text{H}_7(\text{n}) \\ \text{C}_3\text{H}_7(\text{n}) \end{cases}$	$-\text{NHCH}_2\text{CH}_2\text{N} \begin{cases} \text{CH}_3 \\ \text{CH}_3 \end{cases}$	$\text{C}_{17}\text{H}_{29}\text{N}_5$	23.1	23.6
294.	"	$-\text{N} \begin{cases} \text{CH}_3 \\ \text{OCH}_3 \end{cases}$	$\text{C}_{15}\text{H}_{23}\text{N}_4\text{O}$	20.3	20.0
295.	$-\text{N} \begin{cases} \text{C}_3\text{H}_7(\text{iso}) \\ \text{C}_3\text{H}_7(\text{iso}) \end{cases}$	$-\text{NH}_2$	$\text{C}_{13}\text{H}_{20}\text{N}_4$	24.2	24.6
296.	"	$-\text{OH}$	$\text{C}_{13}\text{H}_{19}\text{N}_3\text{O}$	18.0	18.5
297.	$-\text{N} \begin{cases} \text{C}_3\text{H}_7(\text{iso}) \\ \text{C}_3\text{H}_7(\text{iso}) \end{cases}$	$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{22}\text{N}_4$	22.8	22.5
298.	"	$-\text{N}$	$\text{C}_{18}\text{H}_{28}\text{N}_4$	18.7	19.1
299.	$-\text{N} \begin{cases} \text{C}_4\text{H}_9(\text{n}) \\ \text{C}_4\text{H}_9(\text{n}) \end{cases}$	$-\text{Cl}$	$\text{C}_{15}\text{H}_{22}\text{ClN}_3$	15.0	15.4
300.	"	$-\text{CN}$	$\text{C}_{16}\text{H}_{22}\text{N}_4$	20.7	21.0
301.	$-\text{N} \begin{cases} \text{C}_4\text{H}_9(\text{n}) \\ \text{C}_4\text{H}_9(\text{n}) \end{cases}$	$-\text{OH}$	$\text{C}_{15}\text{H}_{23}\text{N}_3\text{O}$	16.1	16.5
302.	"	$-\text{OCH}_3$	$\text{C}_{16}\text{H}_{25}\text{N}_3\text{O}$	15.3	15.0
303.	"	$-\text{NH}-\text{CH}_3$	$\text{C}_{16}\text{H}_{26}\text{N}_4$	20.4	20.8
304.	"	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{17}\text{H}_{28}\text{N}_4\text{O}$	18.4	18.1
305.	"	$-\text{NH}-$	$\text{C}_{21}\text{H}_{34}\text{N}_4$	16.4	16.8
306.	"	$-\text{N}$	$\text{C}_{19}\text{H}_{30}\text{N}_4\text{O}$	17.0	17.3
307.	"	$-\text{NH}-\text{N} \begin{cases} \text{CH}_3 \\ \text{CH}_3 \end{cases}$	$\text{C}_{17}\text{H}_{29}\text{N}_5$	23.1	23.5
308.	"	$-\text{NH}-$	$\text{C}_{19}\text{H}_{31}\text{N}_5\text{O}$	20.3	21.0
309.	$-\text{N} \begin{cases} \text{C}_4\text{H}_9(\text{iso}) \\ \text{C}_4\text{H}_9(\text{iso}) \end{cases}$	$-\text{NH}_2$	$\text{C}_{15}\text{H}_{24}\text{N}_4$	21.5	21.9
310.	"	$-\text{NH}-\text{CH}_3$	$\text{C}_{16}\text{H}_{26}\text{N}_4$	20.4	20.6
311.	$-\text{N} \begin{cases} \text{C}_4\text{H}_9(\text{iso}) \\ \text{C}_4\text{H}_9(\text{iso}) \end{cases}$	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{17}\text{H}_{28}\text{N}_4\text{O}$	18.4	18.5

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
312.	$\text{—N—C}_4\text{H}_9(\text{iso})$	$\text{—NHCH}_2\text{CH}_2\text{N—CH}_3$	$\text{C}_{19}\text{H}_{33}\text{N}_5$	21.1	21.5
313.	$\text{—N—C}_4\text{H}_9(\text{iso})$	—NH_2	$\text{C}_{17}\text{H}_{28}\text{N}_4$	19.4	20.0
314.	$\text{—N—C}_5\text{H}_{11}(\text{n})$	—NH—CH_3	$\text{C}_{18}\text{H}_{30}\text{N}_4$	18.5	18.2
315.	$\text{—N—C}_5\text{H}_{11}(\text{n})$	$\text{—NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{19}\text{H}_{32}\text{N}_4\text{O}$	16.9	17.5
316.	$\text{—N—C}_6\text{H}_{13}(\text{n})$	—NH_2	$\text{C}_{19}\text{H}_{32}\text{N}_4$	17.7	18.0
317.	$\text{—N—C}_6\text{H}_{13}(\text{n})$	—NH—CH_3	$\text{C}_{20}\text{H}_{34}\text{N}_4$	17.0	16.7
318.	$\text{—N—C}_6\text{H}_{13}(\text{n})$	$\text{—NH—C}_6\text{H}_5$	$\text{C}_{25}\text{H}_{36}\text{N}_4$	14.3	14.7
319.	—N—H—O	—NH_2	$\text{C}_{11}\text{H}_{14}\text{N}_4\text{O}$	25.7	26.2
320.	—N—H—O	—OH	$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_2$	19.2	19.5
321.	—N—H—O	—NH—CH_3	$\text{C}_{12}\text{H}_{16}\text{N}_4\text{O}$	24.1	24.5
322.	—N—H—O	$\text{—N—C}_2\text{H}_5$	$\text{C}_{15}\text{H}_{22}\text{N}_4\text{O}$	20.4	21.0
323.	—N—H—O	—NH_2	$\text{C}_{11}\text{H}_{14}\text{N}_4$	27.7	28.0
324.	—N—H—O	—NH—CH_3	$\text{C}_{12}\text{H}_{16}\text{N}_4$	25.9	25.7
325.	—N—H—O	—N—CH_3	$\text{C}_{13}\text{H}_{18}\text{N}_4$	24.3	25.0
326.	—N—H—O	—OH	$\text{C}_{12}\text{H}_{15}\text{N}_3\text{O}$	19.3	19.5
327.	—N—H—O	—NH_2	$\text{C}_{12}\text{H}_{16}\text{N}_4$	25.9	25.5
328.	—N—H—O	—NH—CH_3	$\text{C}_{13}\text{H}_{18}\text{N}_4$	24.3	24.7
329.	—N—H—O	$\text{—NHCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$	$\text{C}_{16}\text{H}_{24}\text{N}_4\text{O}$	19.4	19.5
330.	—N—H—O	—N—CH_3	$\text{C}_{14}\text{H}_{20}\text{N}_4$	22.9	23.0
331.	—N—H—O	$\text{—NH—C}_6\text{H}_5$	$\text{C}_{18}\text{H}_{26}\text{N}_4$	18.8	18.5
332.	—N—H—O	$\text{—NH—C}_6\text{H}_5$	$\text{C}_{18}\text{H}_{20}\text{N}_4$	19.2	19.2
333.	—N—H—N—CH_3	—NH_2	$\text{C}_{12}\text{H}_{17}\text{N}_5$	30.4	30.6
334.	—N—H—N—CH_3	—NH—CH_3	$\text{C}_{13}\text{H}_{19}\text{N}_5$	28.5	28.7
335.	—N—H—N—CH_3	—N—CH_3	$\text{C}_{14}\text{H}_{21}\text{N}_5$	27.0	27.5
336.	$\text{—N—H—N—C}_2\text{H}_5$	—NH—CH_3	$\text{C}_{14}\text{H}_{21}\text{N}_5$	27.0	27.3
337.	—N—CH_3	—NH—CH_3	$\text{C}_{11}\text{H}_{17}\text{N}_5$	32.0	32.3
338.	—N—CH_3	$\text{—NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{12}\text{H}_{18}\text{N}_5\text{O}$	28.1	28.0
339.	—NH—N—CH_3	—NH_2	$\text{C}_9\text{H}_{13}\text{N}_5$	36.7	37.0
340.	—NH—N—CH_3	—NH—CH_3	$\text{C}_{10}\text{H}_{15}\text{N}_5$	34.1	34.5
341.	$\text{—NH—N—C}_2\text{H}_5$	—N—CH_3	$\text{C}_{13}\text{H}_{21}\text{N}_5$	28.4	29.0
342.	$\text{—NH—N—C}_2\text{H}_5$	—N—CH_3	$\text{C}_{14}\text{H}_{23}\text{N}_5\text{O}$	25.3	25.5
343.	$\text{—NH—N—C}_2\text{H}_5$	$\text{—NH—C}_6\text{H}_5$	$\text{C}_{17}\text{H}_{21}\text{N}_5$	23.8	24.0

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
344.		-NH-CH ₃	C ₁₅ H ₁₇ N ₅	26.2	26.6
345.	"		C ₁₆ H ₁₉ N ₅	24.9	25.5
346.		-NH-CH ₃	C ₁₃ H ₁₇ N ₅	28.9	29.0
347.	"		C ₁₄ H ₁₉ N ₅	27.3	27.5
348.		-NH ₂	C ₁₂ H ₁₇ N ₅	30.3	31.6
349.	"	-NH-CH ₃	C ₁₃ H ₁₉ N ₅	28.6	29.0
350.	"		C ₁₄ H ₂₁ N ₅	27.1	26.8
351.		-NH-CH ₃	C ₁₂ H ₁₇ N ₅ O	28.4	28.6
352.	"		C ₁₃ H ₁₉ N ₅ O	26.8	27.0
353.		-NH-CH ₃	C ₁₂ H ₁₉ N ₅	30.1	29.5
354.		-NHCH ₂ CH ₂ OH	C ₁₅ H ₂₅ N ₅ O	24.1	24.6
355.		-NH-CH ₃	C ₁₃ H ₂₁ N ₅	28.4	29.0
356.	"	-NHCH ₂ CH ₂ OH	C ₁₄ H ₂₃ N ₅ O	25.3	25.5
357.		-NH-CH ₃	C ₁₅ H ₂₅ N ₅	25.5	26.0
358.		-NH-CH ₃	C ₁₅ H ₂₅ N ₅ O	24.1	23.7
359.		-NH-CH ₃	C ₁₆ H ₂₅ N ₅	24.3	25.0
360.		-NH-CH ₃	C ₁₅ H ₂₃ N ₅	25.7	26.0
361.		-NH-CH ₃	C ₁₆ H ₂₆ N ₆	27.8	27.5
362.	-NH-OH	-NH-CH ₃	C ₈ H ₁₀ N ₄ O	31.5	31.9
363.		-NH-CH ₃	C ₁₀ H ₁₄ N ₄ O	27.2	27.7
364.		-NH-CH ₃	C ₁₁ H ₁₆ N ₄ O	25.5	25.1
365.		-NH-CH ₃	C ₁₆ H ₁₈ N ₄ O	19.8	19.2
366.		-NH-CH ₃	C ₁₃ H ₂₀ N ₄ O	22.6	23.0













-continued-

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
367.		-NH-CH ₃	C ₁₁ H ₁₄ N ₄ O	25.6	25.6
368.		-NH-CH ₃	C ₁₂ H ₁₆ N ₄ O	24.1	24.5
369.	-CN	-Cl	C ₈ H ₄ ClN ₃	23.7	24.2
370.	-CN	-CN	C ₈ H ₄ N ₄	33.3	33.8
371.	-CN	-OH	C ₈ H ₅ N ₃ O	26.4	27.0
372.	-CN	-OCH ₃	C ₈ H ₇ N ₃ O	24.3	24.0
373.	-CN	-NH ₂	C ₈ H ₆ N ₄	35.4	35.8
374.	-CN	-NH-CH ₃	C ₉ H ₈ N ₄	32.6	33.0
375.	-CN		C ₁₀ H ₁₀ N ₄	30.1	30.5
376.	-CN		C ₁₁ H ₁₂ N ₄ O	25.9	26.4
377.	-CN		C ₁₃ H ₁₄ N ₄	24.8	25.2
378.	-CN	-NH-N	C ₁₀ H ₁₁ N ₅	34.8	35.2
379.	-CN	-NH-OCH ₃	C ₉ H ₈ N ₄ O	29.8	29.5
380.	-OCH ₃	-Cl	C ₈ H ₇ ClN ₂ O	15.4	15.0
381.	-OC ₂ H ₅	-Cl	C ₉ H ₉ ClN ₂ O	14.3	14.7
382.	-OC ₂ H ₅ (n)	-Cl	C ₁₁ H ₁₃ ClN ₂ O	12.5	12.8
383.	-OCH ₂ CH ₂ OCH ₃	-Cl	C ₁₀ H ₁₁ ClN ₂ O	12.4	12.0
384.	-OCH ₃	-OCH ₃	C ₉ H ₁₀ N ₂ O ₂	15.7	16.0
385.	-OCH ₃	-OC ₂ H ₅	C ₁₀ H ₁₂ N ₂ O ₂	14.6	14.8
386.	-OC ₂ H ₅	-OC ₂ H ₅	C ₁₁ H ₁₄ N ₂ O ₂	13.6	13.9
387.	-OCH ₃	-OCH ₂ CH ₂ OCH ₃	C ₁₁ H ₁₄ N ₂ O ₃	12.6	12.2
388.	-OCH ₂ CH ₂ OCH ₃	-OCH ₃	C ₁₁ H ₁₄ N ₂ O ₃	12.6	13.0
389.	-OCH ₃	-O-	C ₁₄ H ₁₂ N ₂ O ₂	11.7	11.9
390.	-OCH ₃	-SC ₄ H ₉ (n)	C ₁₂ H ₁₆ N ₂ OS	11.9	12.1
391.	-OC ₂ H ₅	-SC ₂ H ₅	C ₁₁ H ₁₆ N ₂ OS	12.6	13.0
392.	-OCH ₃	-S-	C ₁₄ H ₁₂ N ₂ OS	10.9	12.1
393.	-OCH ₃	-O-CH ₂ -	C ₁₅ H ₁₄ N ₂ O ₂	11.0	11.5
394.	-OCH ₃	-NH-CH ₃	C ₉ H ₁₁ N ₃ O	23.7	23.5
395.	-OCH ₃	-NHCH ₂ CH ₂ OH	C ₁₀ H ₁₃ N ₃ O ₂	20.8	21.2
396.	-OC ₂ H ₅	-NH-(CH ₂) ₃ -OCH ₃	C ₁₃ H ₁₉ N ₃ O ₂	16.9	17.3
397.	-OCH ₂ CH ₂ OCH ₃	-NH-	C ₁₆ H ₁₇ N ₃ O ₂	14.8	15.0
398.	-OC ₄ H ₉ (n)	-NH-(CH ₂) ₃ -N	C ₁₆ H ₂₆ N ₄ O	19.3	19.5
399.	-OCH ₃		C ₁₁ H ₁₅ N ₃ O ₂	19.0	19.3
400.	-OC ₂ H ₅		C ₁₃ H ₁₆ N ₃ O ₂	17.1	17.5
401.	-SCH ₃	-Cl	C ₈ H ₇ ClN ₂ S	14.1	14.5
402.	-SC ₂ H ₅	-Cl	C ₉ H ₉ ClN ₂ S	13.2	13.0
403.	-SC ₄ H ₉ (n)	-Cl	C ₁₁ H ₁₃ ClN ₂ S	11.7	12.1
404.	-SC ₄ H ₉ (n)	-OCH ₃	C ₁₂ H ₁₆ N ₂ OS	11.9	12.3
405.	-SC ₂ H ₅	-O-	C ₁₅ H ₁₄ N ₂ OS	10.4	11.0
406.	-SC ₄ H ₉ (n)	-SC ₄ H ₉ (n)	C ₁₅ H ₂₂ N ₂ S ₂	9.5	10.0
407.	-SCH ₃	-NH-CH ₃	C ₉ H ₁₁ N ₃ S	21.7	22.0
408.	-SC ₂ H ₅	-NH-(CH ₂) ₃ -OCH ₃	C ₁₃ H ₁₉ N ₃ OS	15.8	16.0
409.	-SC ₄ H ₉ (n)	-NHCH ₂ CH ₂ OH	C ₁₃ H ₁₉ N ₃ OS	15.8	16.2
410.	-SC ₄ H ₉ (n)	-NH	C ₁₇ H ₁₉ N ₃ S	14.1	14.5


-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
411.	$-\text{SC}_4\text{H}_9(\text{n})$	$-\text{NH}-(\text{CH}_2)_3-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{16}\text{H}_{26}\text{N}_4\text{S}$	18.3	18.5
412.	$-\text{SC}_4\text{H}_9(\text{n})$	$-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_2\text{CH}_2\text{OH} \end{array}$	$\text{C}_{14}\text{H}_{21}\text{N}_3\text{OS}$	15.0	15.5
413.	$-\text{SC}_4\text{H}_9(\text{n})$	$-\text{N} \begin{array}{c} \text{H} \\ \text{O} \end{array}$	$\text{C}_{15}\text{H}_{21}\text{N}_3\text{OS}$	14.4	15.0
414.	$-\text{SO}_2-\text{CH}_3$	$-\text{Cl}$	$\text{C}_{10}\text{H}_{10}\text{ClN}_3\text{O}-\text{S}$	15.5	15.0
415.	$-\text{SO}_2-\text{C}_4\text{H}_9(\text{n})$	$-\text{Cl}$	$\text{C}_{13}\text{H}_{16}\text{ClN}_3\text{O}-\text{S}$	13.4	13.8
416.	$-\text{SO}_2-\text{C}_6\text{H}_5$	$-\text{Cl}$	$\text{C}_{15}\text{H}_{12}\text{ClN}_3\text{O}-\text{S}$	12.6	12.5
417.	$-\text{SO}_2-\text{CH}_3$	$-\text{CN}$	$\text{C}_{11}\text{H}_{10}\text{N}_4\text{O}_2\text{S}$	21.4	21.0
418.	$-\text{SO}_2-\text{CH}_3$	$-\text{OH}$	$\text{C}_{10}\text{H}_{11}\text{N}_3\text{O}_3\text{S}$	16.6	17.2
419.	$-\text{SO}_2-\text{CH}_3$	$-\text{OCH}_3$	$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_3\text{S}$	15.7	15.3
420.	$-\text{SO}_2-\text{CH}_3$	$-\text{SCH}_3$	$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_2\text{S}_2$	14.8	15.4
421.	$-\text{SO}_2-\text{CH}_3$	$-\text{SO}_2-\text{CH}_3$	$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_4\text{S}_2$	13.3	12.9
422.	$-\text{SO}_2-\text{CH}_3$	$-\text{NH}_2$	$\text{C}_{10}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$	22.2	22.6
423.	$-\text{SO}_2-\text{C}_2\text{H}_5$	$-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_2\text{S}$	19.1	18.5
424.	$-\text{SO}_2-\text{CH}_3$	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$	18.1	18.7
425.	$-\text{SO}_2-\text{CH}_3$	$-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$	18.1	18.3
426.	$-\text{SO}_2-\text{CH}_3$	$-\text{N} \begin{array}{c} \text{H} \\ \text{O} \end{array}$	$\text{C}_{14}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$	17.4	17.6
427.	$-\text{SO}_2-\text{C}_2\text{H}_5$	$-\text{NH}-(\text{CH}_2)_2-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{15}\text{H}_{23}\text{N}_5\text{O}_2\text{S}$	20.8	20.1
428.	$-\text{SO}_2-\text{C}_4\text{H}_9(\text{n})$	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{15}\text{H}_{22}\text{N}_4\text{O}_2\text{S}$	17.4	17.8
429.	$-\text{SO}_2-\text{CH}_2-\text{C}_6\text{H}_5$	$-\text{NH}-\text{CH}_3$	$\text{C}_{17}\text{H}_{18}\text{N}_4\text{O}_2\text{S}$	16.4	16.0
430.	$-\text{SO}_2-\text{C}_6\text{H}_5$	$-\text{NH}-\text{CH}_3$	$\text{C}_{16}\text{H}_{16}\text{N}_4\text{O}_2\text{S}$	17.1	17.8
431.	$-\text{NH}-\text{C}_6\text{H}_4-\text{N} \begin{array}{c} \text{H} \\ \text{C}_6\text{H}_5 \end{array}$	$-\text{NH}-\text{C}_6\text{H}_4-\text{N} \begin{array}{c} \text{H} \\ \text{C}_6\text{H}_5 \end{array}$	$\text{C}_{33}\text{H}_{25}\text{N}_7$	18.9	18.5
432.	$-\text{NH}-\text{C}_6\text{H}_4-\text{N} \begin{array}{c} \text{H} \\ \text{C}_2\text{H}_5 \end{array}$	$-\text{Cl}$	$\text{C}_{23}\text{H}_{20}\text{ClN}_5$	17.5	17.9
433.	$-\text{NH}-\text{C}_6\text{H}_4-\text{N} \begin{array}{c} \text{H} \\ \text{C}_2\text{H}_5 \end{array}$	$-\text{OCH}_3$	$\text{C}_{24}\text{H}_{23}\text{N}_5\text{O}$	17.6	17.0
434.	$-\text{NH}-\text{C}_6\text{H}_4-\text{N} \begin{array}{c} \text{H} \\ \text{C}_2\text{H}_5 \end{array}$	$-\text{N} \begin{array}{l} \text{CH}_3 \\ \text{CH}_3 \end{array}$	$\text{C}_{25}\text{H}_{26}\text{N}_6$	20.5	21.2

-continued

No.	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
435.	"		C ₃₇ H ₃₃ N ₇	17.0	17.2
436.			C ₃₉ H ₃₇ N ₇	16.3	16.6
437.		-OCH ₃	C ₂₆ H ₂₈ N ₆ O	19.1	19.8
438.	"		C ₂₆ H ₂₉ N ₇	22.3	22.5
439.	"		C ₄₁ H ₄₃ N ₉	19.1	19.8
440.			C ₃₃ H ₂₅ N ₇	18.9	18.1
441.		-Cl	C ₂₁ H ₁₅ ClN ₄ O	15.0	15.4
442.	"	-OC ₂ H ₅	C ₂₃ H ₂₀ N ₄ O ₂	14.5	15.0
443.	"		C ₃₃ H ₂₃ N ₅ O ₂	13.4	13.7
444.			C ₃₃ H ₂₃ N ₅ S ₂	12.7	12.5

In the following table are presented further pyridine compounds prepared in accordance with the process of this invention:



No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
445.	-H	-NH ₂	-Cl	C ₆ H ₇ N ₂ Cl	20.1	20.7
446.	-H	-NH ₂	-NH ₂	C ₆ H ₉ N ₃	34.1	33.6
447.	-H	-NH ₂	-OH	C ₆ H ₈ N ₂ O	22.6	22.8





-continued

No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
448.	-H	-NH ₂	-OCH ₃	C ₇ H ₁₀ N ₂ O	20.3	20.1
449.	-H	-NH ₂	-SC ₂ H ₅	C ₈ H ₁₂ N ₂ S	16.6	17.1
450.	-H	-NH ₂	-SO ₂ -CH ₃	C ₇ H ₁₀ N ₂ O ₂ S	15.0	15.2
451.	-H	-NH ₂	-NH-CH ₃	C ₇ H ₁₁ N ₃	30.6	31.0
452.	-H	-NH-CH ₃	-NH-CH ₃	C ₈ H ₁₃ N ₃	27.8	28.2
453.	-H	-NHCH ₂ CH ₂ OH	-NHCH ₂ CH ₂ OH	C ₁₀ H ₁₇ N ₃ O ₂	19.9	20.2
454.	-H			C ₁₄ H ₂₅ N ₃	17.9	17.3
455.	-H			C ₁₄ H ₂₁ N ₃ O ₂	16.0	16.3
456.	-H	-NHCH ₂ CH ₂ CH ₃	-NHCH ₂ CH ₂ N	C ₁₃ H ₂₃ N ₄	23.8	23.2
457.	-NH ₂	-NH ₂	-NH ₂	C ₆ H ₁₀ N ₄	40.6	40.8
458.	-NH ₂	-NH-CH ₃	-NH-CH ₃	C ₈ H ₁₄ N ₄	33.7	33.2
459.	-NH ₂			C ₁₀ H ₁₈ N ₄	28.9	28.7
460.	-NH ₂	-NHCH ₂ CH ₂ OH	-NHCH ₂ CH ₂ OH	C ₁₀ H ₁₈ N ₄ O ₂	24.8	25.1
461.	-NO	-NH-CH ₃	-NH-CH ₃	C ₈ H ₁₂ N ₄ O	31.1	31.0
462.	-NO			C ₁₀ H ₁₆ N ₄ O	26.9	26.7
463.	-NO			C ₁₄ H ₂₀ N ₄ O ₃	19.2	19.5
464.	-NO ₂	-NH-CH ₃	-NH-CH ₃	C ₈ H ₁₂ N ₄ O ₂	28.5	28.5
465.	-NO ₂			C ₁₀ H ₁₆ N ₄ O ₂	25.0	25.3
466.	-NO ₂			C ₁₄ H ₂₀ N ₄ O ₄	18.2	18.5
467.	-CH ₃	-NH-CH ₃	-NH-CH ₃	C ₉ H ₁₅ N ₃	25.4	26.0
468.	-CH ₂ -CH ₃	-NH-CH ₃	-NH-CH ₂ -CH ₃	C ₁₁ H ₁₉ N ₃	21.7	22.0
469.	-CH ₂ CH ₂ CN	-NH ₂	-NH ₂	C ₉ H ₁₂ N ₄	31.8	32.0
470.	-CH ₂ CH ₂ O-	-NH ₂	-NH ₂	C ₁₅ H ₁₇ N ₃ O	16.5	17.0
471.	-CH ₂ CH ₂ N-	-NH-CH ₃	-NH-CH ₃	C ₁₉ H ₂₈ N ₄ O	17.2	17.5
472.	-CH ₂ CH ₂ N-	-NH-CH ₃	-OCH ₂ CH ₂ =CH ₂	C ₁₆ H ₂₇ N ₃ O	15.2	15.3
473.	-CH ₂ -CH=CH ₂ 	-NH-CH ₃	-NH-CH ₃	C ₁₁ H ₁₇ N ₃	22.0	22.4
474.	-CH ₂ -C=CH ₂	-NH-CH ₃	-SO ₂ CH ₂ CH=CH ₂	C ₁₄ H ₂₀ N ₂ O ₂ S	10.0	9.8
475.	-C ₃ H ₇ (n)	-NHCH ₂ CH ₂ CH ₃		C ₁₅ H ₂₇ N ₃	16.8	16.5
476.	-C ₄ H ₉ (n)	-NH-CH ₃		C ₁₇ H ₂₈ N ₂ O	10.1	9.7
477.	-C ₄ H ₉ (iso)	-NH-CH ₃	-O-CH ₂ -	C ₁₈ H ₂₄ N ₂ O	9.9	10.2
478.	-C ₆ H ₁₃ (n)	-NH-CH ₃	-O-	C ₁₈ H ₂₄ N ₂ O	9.9	10.4
479.		-NH-C ₃ H ₇ (iso)	-OCH ₂ CH ₂ OCH ₃	C ₁₉ H ₃₃ N ₂ O	9.4	9.5
480.	-CH ₂ -	-NH-CH ₃	-NHCH ₂ CH ₂ CH ₂ N	C ₁₉ H ₂₇ N ₄	18.0	18.3














-continued

No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
481.		-NH ₂	-O-CH ₂ -C(CH ₃)=CH ₂	C ₁₆ H ₁₈ N ₂ O	11.0	10.5
482.		-NH ₂	-S-CH(CH ₃) ₂	C ₁₆ H ₂₀ N ₂ OS	9.7	9.5
483.		-NH-CH ₃	-S-	C ₂₀ H ₁₉ ClN ₂ S	7.9	8.3
484.		-NH-CH ₃	-S-	C ₂₁ H ₂₀ Cl ₂ N ₂ S	6.9	6.8
485.	-CH ₂ CH ₂ -N	-NH ₂	-NH ₂	C ₁₂ H ₂₀ N ₄ O	23.7	23.9
486.	"	-N(CH ₃) ₂	-N(CH ₃) ₂	C ₁₆ H ₂₈ N ₄ O	19.2	19.7
487.	-CH ₂ CH ₂ -N	-NH-CH ₃	-NH-CH ₃	C ₁₅ H ₂₆ N ₄	21.4	21.6
488.	-CH ₂ CH ₂ -N	-NH-CH ₃	-NH-CH ₃	C ₁₄ H ₂₄ N ₄	22.6	22.2
489.	-COOCH ₃	-NH-CH ₃	-NH-CH ₃	C ₁₆ H ₁₅ N ₃ O ₂	20.1	20.3
490.	-COOC ₂ H ₅	-NH-CH ₃	-NH-CH ₃	C ₁₇ H ₁₇ N ₃ O ₂	18.8	18.5
491.	-COOC ₂ H ₅	-N(CH ₂ C ₂ H ₅) ₂	-N(CH ₂ C ₂ H ₅) ₂	C ₁₇ H ₂₀ N ₃ O ₂	13.7	14.0
492.	-COOC ₂ H ₅	-N	-NH-CH ₃	C ₁₄ H ₂₁ N ₃ O ₃	15.0	14.8
493.	-COOC ₄ H ₉ (n)	-NHCH ₂ CH ₂ OH		C ₁₅ H ₂₅ N ₃ O ₄	13.5	13.8
494.	-COOC ₆ H ₁₃ (n)	-NHCH ₂ CH ₂ CH ₃	-NHCH ₂ CH ₂ N(CH ₃) ₂	C ₂₀ H ₃₆ N ₄ O ₂	15.4	15.0
495.	-COCH ₃	-NH-CH ₃	-NH-CH ₃	C ₁₀ H ₁₅ N ₃ O	21.7	21.5
496.	-CO-CH=CH ₂	-NH-CH ₃	-NH-CH ₃	C ₁₁ H ₁₅ N ₃ O	20.5	20.7
497.	-CO-C ₆ H ₁₃ (n)	-NH-CH ₃	-N(CH ₃) ₂	C ₁₈ H ₂₇ N ₃ O	15.2	15.3
498.	-CO-	-NH-CH ₃	"	C ₁₆ H ₂₅ N ₃ O	15.3	15.5
499.	-CO-CH ₂ -	-NH-CH ₃	-NH-CH ₃	C ₁₆ H ₁₉ N ₃ O	15.6	15.8
500.	-CO-	-NH ₂	-NH ₂	C ₁₃ H ₁₃ N ₃ O	18.5	18.3
501.	"	-NH-CH ₃	-NH-CH ₃	C ₁₅ H ₁₇ N ₃ O	16.5	16.7
502.	"	-N(CH ₃) ₂	-N(CH ₃) ₂	C ₁₇ H ₂₁ N ₃ O	14.8	14.6
503.	"	-NH-CH ₃	-N	C ₁₈ H ₂₁ N ₃ O ₂	13.5	13.2
504.	-CO-NH ₂	-NH ₂	-Cl	C ₇ H ₈ ClN ₃ O	22.6	22.8
505.	-CO-NH ₂	-NH ₂	-NH ₂	C ₇ H ₁₀ N ₄ O	33.8	34.2
506.	-CO-NH ₂	-NH ₂	-OH	C ₇ H ₈ N ₃ O ₂	25.1	24.7

-continued

No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
507.	—CO—NH ₂	—NH ₂	—OC ₂ H ₅	C ₉ H ₁₃ N ₃ O ₂	21.5	22.1
508.	—CO—NH ₂	—NH ₂	—SC ₂ H ₅	C ₉ H ₁₃ N ₃ OS	19.9	19.6
509.	—CO—NH ₂	—NH ₂	—SO ₂ C ₂ H ₅	C ₉ H ₁₃ N ₃ O ₃ S	17.3	17.5
510.	—CO—NH ₂	—NH—CH ₃	—NH—CH ₃	C ₉ H ₁₄ N ₄ O	28.9	28.7
511.	—CO—NH ₂	—NHCH ₂ CH ₂ OCH ₃		C ₁₃ H ₂₂ N ₄ O ₃	19.9	19.6
512.	—CO—NH ₂	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₅ H ₂₆ N ₄ O	20.1	20.5
513.	—CO—NH ₂	—NHCH ₂ CH ₃		C ₁₃ H ₂₀ N ₄ O ₂	21.2	21.4
514.	—CO—NH ₂	—NHCH ₂ CH ₃	—NHCH ₂ CH ₂ CH ₂ N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₄ H ₂₅ N ₅ O	25.1	25.3
515.	—CO—NH—CH ₃	—NH ₂	—NH ₂	C ₈ H ₁₂ N ₄ O	31.1	31.0
516.	—CONHCH ₂ CH ₂ OH	—NH ₂	—S—CH ₂ —CH=CH ₂	C ₁₂ H ₁₇ N ₃ O ₃	16.7	17.0
517.	—CONHC ₄ H ₉ (n)	—NH—CH ₃	—NH—CH ₃	C ₁₃ H ₂₂ N ₄ O	22.4	22.1
518.	—CO—N $\begin{smallmatrix} \text{CH}_2 \\ \text{CH}_2 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	—NHCH ₂ CH ₂ OCH ₃	—NHCH ₂ CH ₂ OCH ₃	C ₁₅ H ₂₄ N ₄ O ₃	18.2	18.6
519.	—CO—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	—N $\begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	C ₁₉ H ₃₄ N ₄ O	16.8	17.1
520.	—CO—N 	—NH—CH ₃	—NH—C ₃ H ₇ (iso)	C ₁₅ H ₂₄ N ₄ O ₂	19.2	19.1
521.	—CO—NH— 	—NH ₂	—NH ₂	C ₁₃ H ₁₄ N ₄ O	23.2	23.6
522.	"	—NH—CH ₃	—NH—CH ₃	C ₁₅ H ₁₈ N ₄ O	20.7	20.9
523.	—CO—NH— 	—NHCH ₂ CH ₂ OH	—NHCH ₂ CH ₂ OH	C ₁₉ H ₂₆ N ₄ O ₃	15.7	16.2
524.	—CO—N 	—NH—CH ₃	—NH—CH ₃	C ₁₆ H ₂₀ N ₄ O	19.7	19.5
525.	—SO ₂ —CH ₂ —CH ₃	—NH—CH ₃	—NH—CH ₂ —CH ₃	C ₁₁ H ₁₆ N ₃ O ₃ S	15.4	15.2
526.	—SO ₂ —C ₃ H ₇ (n)	—NH—C ₃ H ₇ (n)		C ₁₅ H ₂₇ N ₃ O ₃ S	12.8	13.1
527.	—SO ₂ —C ₄ H ₉ (n)	—NH ₂	—NH ₂	C ₁₀ H ₁₇ N ₃ O ₂ S	17.3	17.6
528.	—SO ₂ —C ₄ H ₉ (iso)	—NH ₂	—NH ₂	C ₁₀ H ₁₇ N ₃ O ₂ S	17.3	17.4
529.	—SO ₂ —C ₅ H ₁₁ (iso)	—NH—CH ₃	—N $\begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	C ₁₄ H ₂₅ N ₃ O ₂ S	14.0	14.3
530.	—SO ₂ — 	—NH—CH ₃	"	C ₁₅ H ₂₅ N ₃ O ₂ S	13.5	14.0
531.	—SO ₂ — 	—NH—CH ₃	—NH—CH ₃	C ₁₄ H ₁₇ N ₃ O ₂ S	14.4	14.2
532.	—SO ₂ — 	—NHCH ₂ CH ₂ OH	—NH—CH ₃	C ₁₅ H ₁₆ ClN ₃ O ₃ S	11.8	12.2
533.	—SO ₂ — 		—NH—CH ₃	C ₁₉ H ₂₅ N ₃ O ₃ S	11.2	11.4
534.	—SO ₂ —NH ₂	—NH ₂	—NH ₂	C ₆ H ₁₀ N ₄ O ₂ S	27.7	27.5
535.	—SO ₂ —NH ₂	—NH ₂	—SO ₂ — 	C ₁₂ H ₁₃ N ₃ O ₄ S ₂	12.8	13.2


-continued

No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
536.	$-\text{SO}_2-\text{NH}_2$	$-\text{NH}-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$\text{C}_9\text{H}_{14}\text{N}_4\text{O}_2\text{S}$	24.4	24.2
537.	$-\text{SO}_2-\text{NH}_2$	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{10}\text{H}_{18}\text{N}_4\text{O}_4\text{S}$	19.3	19.5
538.	$-\text{SO}_2-\text{N} \begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	$-\text{NH}$ 	$-\text{NH}$ 	$\text{C}_{20}\text{H}_{22}\text{N}_4\text{O}_2\text{S}$	14.7	15.2
539.	$-\text{SO}_2\text{NHC}_2\text{H}_5$	$-\text{NH}-\text{CH}_3$	$-\text{S}$ 	$\text{C}_{15}\text{H}_{25}\text{N}_3\text{O}_2\text{S}_2$	12.2	12.4
540.	$-\text{SO}_2\text{NH}(\text{CH}_2)_3\text{OCH}_3$	$-\text{NH}_2$	$-\text{NH}-\text{N} \begin{smallmatrix} \text{CH}_3 \\ \text{CH}_3 \end{smallmatrix}$	$\text{C}_{12}\text{H}_{23}\text{N}_5\text{O}_3\text{S}$	22.1	22.5
541.	$-\text{SO}_2-\text{N} \begin{smallmatrix} \text{H} \\ \text{O} \end{smallmatrix}$ 	$-\text{NH}_2$	$-\text{NH}_2$	$\text{C}_{10}\text{H}_{16}\text{N}_4\text{O}_3\text{S}$	20.6	21.1
542.	"	$-\text{NH}-\text{CH}_3$	$-\text{N} \begin{smallmatrix} \text{H} \\ \text{O} \end{smallmatrix}$ 	$\text{C}_{13}\text{H}_{24}\text{N}_4\text{O}_4\text{S}$	15.7	16.0
543.	"	$-\text{N} \begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	$-\text{N} \begin{smallmatrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{smallmatrix}$	$\text{C}_{18}\text{H}_{32}\text{N}_4\text{O}_3\text{S}$	14.6	14.8
544.	$-\text{SO}_2-\text{N} \begin{smallmatrix} \text{H} \\ \text{O} \end{smallmatrix}$ 	$-\text{NH}-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$\text{C}_{13}\text{H}_{22}\text{N}_4\text{O}_2\text{S}$	18.8	18.5
545.	"	$-\text{NH}-\text{CH}_3$	$-\text{S}-\text{CH}_2$ 	$\text{C}_{19}\text{H}_{25}\text{N}_3\text{O}_2\text{S}_2$	10.7	11.2
546.	$-\text{SO}_2-\text{N} \begin{smallmatrix} \text{H} \\ \text{N}-\text{CH}_3 \end{smallmatrix}$ 	$-\text{NH}-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$\text{C}_{12}\text{H}_{23}\text{N}_5\text{O}_2\text{S}$	23.2	23.0
547.	$-\text{SO}_2-\text{NH}$ 		$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{24}\text{N}_4\text{O}_2\text{S}$	17.9	18.2
548.	"	$-\text{NH}-\text{CH}_3$	$-\text{SO}_2$ 	$\text{C}_{18}\text{H}_{31}\text{N}_3\text{O}_4\text{S}_2$	9.8	10.2
549.	$-\text{SO}_2\text{NHCH}_2$ 	$-\text{N} \begin{smallmatrix} \text{H} \\ \text{O} \end{smallmatrix}$ 	$-\text{NHCH}_2\text{CH}_2\text{OH}$	$\text{C}_{19}\text{H}_{26}\text{N}_4\text{O}_4\text{S}$	13.8	13.5
550.	$-\text{SO}_2-\text{NH}$ 	$-\text{NH}_2$	$-\text{NH}_2$	$\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_2\text{S}$	20.1	20.3
551.	"	$-\text{NH}$ 	$-\text{NH}$ 	$\text{C}_{24}\text{H}_{34}\text{N}_4\text{O}_2\text{S}$	12.7	12.5
552.	"	$-\text{NH}-\text{CH}_2$ 	$-\text{NH}-\text{CH}_2$ 	$\text{C}_{26}\text{H}_{26}\text{N}_4\text{O}_2\text{S}$	12.2	12.5
553.	$-\text{SO}_2-\text{NH}$ 	$-\text{NH}$ 	$-\text{NH}$ 	$\text{C}_{24}\text{H}_{22}\text{N}_4\text{O}_2\text{S}$	13.0	13.2
554.	"	$-\text{NH}$ 	$-\text{NH}$ 	$\text{C}_{32}\text{H}_{26}\text{N}_4\text{O}_2\text{S}$	10.6	10.2







-continued

No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
555.		-NH-CH ₃		C ₁₆ H ₁₈ N ₂ O ₃	9.8	9.5
556.	"	-NH-CH ₃		C ₁₇ H ₂₄ N ₄ O ₂	17.8	18.2
557.		-NH-CH ₃		C ₁₉ H ₂₄ N ₂ O ₃	8.5	9.0
558.		-NH ₂	-NH ₂	C ₁₂ H ₁₂ N ₄ O	24.5	25.0
559.	"	-NH-CH ₃		C ₁₄ H ₁₆ N ₄ O	21.8	22.2
560.	"			C ₁₆ H ₂₀ N ₄ O	19.7	20.1
561.		-NH-CH ₃	-NH-CH ₃	C ₁₆ H ₁₇ N ₅ O	23.7	23.5
562.	"			C ₁₈ H ₂₁ N ₅ O	21.7	21.5
563.		-NH-CH ₃	-NH-CH ₃	C ₁₆ H ₁₆ N ₄ OS	17.9	18.2
564.				C ₁₉ H ₂₂ N ₄ O ₂ S	15.1	15.5
565.		-NH-CH ₃		C ₂₀ H ₁₉ ClN ₂ O ₂ S	7.3	7.0
566.		-NH-CH ₃	-NH-CH ₃	C ₁₄ H ₁₈ N ₄	23.1	23.5
567.		-NH-CH ₃	-NH-CH ₃	C ₁₄ H ₁₈ N ₄	23.1	23.3
568.		-NH-CH ₃	-NH-CH ₃	C ₁₅ H ₂₀ N ₄	21.9	21.7
569.	-C ₄ H ₉ (n)	-CN		C ₁₃ H ₁₉ N ₃	19.4	19.0
570.	-CO-NH ₂	-OCH ₃		C ₉ H ₁₂ N ₂ O ₃	14.3	14.7
571.	-CO-CH ₃	-OC ₂ H ₅		C ₁₂ H ₁₇ N ₃ O ₃	6.7	6.5






In the following table are presented further pyridine compounds prepared in accordance with the method of this invention:



or tautomeric forms.


No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd N%	Found N%
572.	—H	—NH—CH ₃	—NH—CH ₃	C ₈ H ₁₀ N ₄	34.6	25.0
573.	—H	—NH—CH ₂ —CH ₂ —OH	—NH—CH ₃	C ₁₀ H ₁₄ N ₄ O ₂	25.2	25.0
574.	—C ₃ H ₇ (n)	—NHCH ₂ CH ₂ CH ₃	—NH(CH ₂) ₃ N 	C ₁₇ H ₂₉ N ₅	23.1	23.3
575.	—C ₃ H ₇ (iso)	—NHCH ₂ CH ₂ CH ₃	—NH(CH ₂) ₃ N 	C ₁₇ H ₂₉ N ₅	23.1	22.8
576.	—C ₆ H ₁₃ (n)	—NHCH ₂ CH ₂ CH ₂ OCH ₃	—NH(CH ₂) ₃ N 	C ₂₀ H ₃₄ N ₄ O ₂	15.5	16.0
577.	—CH ₂ CH ₂ N 	—NH—CH ₃	—NH—CH ₃	C ₁₂ H ₁₉ N ₅	30.0	30.3
578.	—CH ₂ CH ₂ N 	—NH—CH ₃	—NH—CH ₃	C ₁₃ H ₁₉ N ₅ O	26.8	27.0
579.	—CH ₂ CH ₂ —N 	—NH ₂	—NH ₂	C ₁₂ H ₁₇ N ₅ O	28.3	28.8
580.	—CH ₂ CH ₂ —N 	—NH ₂	—NH ₂	C ₁₂ H ₁₇ N ₅	30.3	30.5
581.	—CH ₂ CH ₂ OH	—NH—CH ₃	—NH—CH ₃	C ₁₀ H ₁₄ N ₄ O	27.2	27.5
582.	—CH ₂ CH ₂ O—COCH ₃	—NHCH ₂ CH ₂ CH ₂ OH	—NH—CH ₃	C ₁₆ H ₂₄ N ₄ O ₄	16.7	17.0
583.	—(CH ₂) ₂ O—CONHC ₂ H ₅	—NH—CH ₃	—NH—CH ₃	C ₁₃ H ₁₉ N ₅ O ₂	25.3	25.1
584.	—CH ₂ CH ₂ O— 	—NH(CH ₂) ₃ O—C ₃ H ₇ (iso)		C ₂₆ H ₃₈ N ₄ O ₃	12.3	12.7
585.	— 	—NH ₂	—NH ₂	C ₁₂ H ₁₆ N ₄	25.9	25.5
586.	—CH ₂ — 	—NH ₂	—NH ₂	C ₁₃ H ₁₂ N ₄	25.0	25.3
587.	— 	—H 	—NH—CH ₃	C ₁₇ H ₁₉ N ₄	20.1	20.7
588.	— 	—NH—CH ₃	—NH—CH ₃	C ₁₄ H ₁₃ N ₄ Cl	20.6	21.0
589.	— 	—NH—CH ₃	—NH—CH ₃	C ₁₆ H ₁₈ N ₄	21.0	20.4
590.		—NH—CH ₃	—NH—CH ₃	C ₁₂ H ₁₂ N ₄ O	24.5	24.7
591.		—NH—CH ₃	—NH—CH ₃	C ₁₂ H ₁₂ N ₄ S	22.9	22.5
592.		—NH—CH ₃	—NH—CH ₃	C ₁₁ H ₁₁ N ₅ S	28.6	28.5

-continued







No.	Y	Z ₁	Z ₂	Empirical Formula	Cal'd N%	Found N%
593.		-NH-CH ₃	-NH-CH ₃	C ₁₄ H ₁₅ N ₅	27.7	28.2
594.		-NH-CH ₃	-NH-CH ₃	C ₁₃ H ₁₃ N ₅	28.3	28.7
595.		-NH-CH ₃	-NH-CH ₃	C ₁₅ H ₁₄ N ₆	30.2	29.6
596.		-NH-CH ₃	-NH-CH ₃	C ₁₂ H ₁₃ N ₅ S	27.0	27.3
597.		-NH-C ₂ H ₅	-NH-C ₂ H ₅	C ₁₇ H ₁₈ N ₆	27.5	27.7
598.	-C ₃ H ₇ (n)	-CN		C ₁₂ H ₁₄ N ₄	26.2	26.5
599.	-C ₃ H ₇ (n)	-SCH ₃	-OCH ₃	C ₁₁ H ₁₄ N ₂ OS	12.6	13.0

In the following table are presented further pyridine compounds prepared in accordance with the method of this invention:




40





or tautomeric forms.

No.	X	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
600.	-C ₃ H ₇ (n)	-C ₄ H ₉ (n)	-CN		C ₁₈ H ₂₃ N ₃	17.1	17.5
601.	-C ₃ H ₇ (n)	-CN	-OCH ₃	"	C ₁₂ H ₁₇ N ₃ O	18.7	19.2
602.	-C ₃ H ₇ (n)	-C ₃ H ₇ (n)	-SCH ₃	-SCH ₃	C ₁₃ H ₂₁ NS ₂	5.5	5.8
603.	H	-CH ₃	-NH ₂	-NH ₂	C ₆ H ₉ N ₃	34.1	34.5
604.	H		-NH-CH ₃	-NH-CH ₃	C ₁₃ H ₁₆ N ₄	24.5	24.2
605.	H	-CO-NH ₂			C ₁₀ H ₁₆ N ₄ O	26.9	27.3
606.	-C ₂ H ₅	-NH ₂	-NH-CH ₃		C ₁₂ H ₁₉ N ₃ O	19.0	19.2
607.	-C ₂ H ₅	-H	-NHCH ₂ CH ₂ OH		C ₁₃ H ₂₁ N ₃ O	17.9	18.2
608.	-C ₄ H ₉ (n)	-NO	-NH-CH ₃	-NH-CH ₃	C ₁₁ H ₁₈ N ₄ O	25.2	25.0
609.	-C ₄ H ₉ (n)	-NO ₂	-NH-CH ₃	-NH-CH ₃	C ₁₁ H ₁₈ N ₄ O ₂	23.5	23.2
610.	-C ₄ H ₉ (n)	-C ₄ H ₉ (n)	-NH ₂		C ₂₀ H ₂₈ N ₂ O ₂	8.5	8.2

-continued

No.	X	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
611.	$-\text{C}_4\text{H}_9(\text{sec})$		$-\text{NH}-\text{C}_2\text{H}_5$		$\text{C}_{19}\text{H}_{28}\text{N}_4$	17.9	17.6
612.	$-\text{C}_4\text{H}_9(\text{sec})$		$-\text{N} \begin{matrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{matrix}$	$-\text{N} \begin{matrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{matrix}$	$\text{C}_{20}\text{H}_{32}\text{N}_4\text{S}$	15.5	16.1
613.	$-\text{C}_4\text{H}_9(\text{iso})$	$-\text{OC}-$ 	$-\text{NHCH}_2\text{CH}_2\text{OCH}_3$		$\text{C}_{21}\text{H}_{30}\text{N}_4\text{O}_3$	14.5	14.2
614.	$-\text{C}_4\text{H}_9(\text{tert})$	$-\text{CO}-\text{NH}_2$	$-\text{NH}-\text{CH}_3$	$-\text{SO}_2\text{CH}_2\text{CH}_2\text{OCH}_3$	$\text{C}_{14}\text{H}_{23}\text{N}_3\text{O}_4\text{S}$	12.8	12.2
615.	$-\text{C}_5\text{H}_{11}(\text{iso})$	$-\text{CO}-\text{NH}_2$	$-\text{NH}-\text{CH}_3$	$-\text{SO}_2$ 	$\text{C}_{19}\text{H}_{22}\text{N}_4\text{O}_3\text{S}$	14.5	14.3
616.	$-\text{CH}=\text{CH}_2$	$-\text{COOC}_2\text{H}_5$		$-\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$	$\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_3$	13.8	14.2
617.	$-\text{CH}=\text{CH}-\text{CH}_3$	$-\text{SO}_2-\text{NH}_2$		$-\text{NH}-$ 	$\text{C}_{20}\text{H}_{20}\text{N}_4\text{O}_2\text{S}$	14.7	15.2
618.	$-\text{CH}_2\text{CH}_2\text{N} \begin{matrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{matrix}$	$-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$\text{C}_{14}\text{H}_{26}\text{N}_4$	22.4	22.8
619.	$-\text{CH}_2\text{CH}_2\text{CN}$	$-\text{CH}_2\text{CH}_2\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$		$-\text{NH}-\text{CH}_3$	$\text{C}_{17}\text{H}_{27}\text{N}_5\text{O}$	22.1	22.5
620.	$-\text{CH}_2\text{CH}_2\text{OCH}_3$	$-\text{CO}-\text{CH}_3$	$-\text{NHCH}_2\text{CH}_2\text{CH}_3$	$-\text{SCH}_2\text{CH}_2\text{OCH}_3$	$\text{C}_{16}\text{H}_{26}\text{N}_2\text{O}_3\text{S}$	8.6	8.9
621.	$-\text{CH}_2\text{CH}_2-\text{N} \begin{matrix} \text{H} \\ \text{H} \end{matrix}$ 	$-\text{H}$	$-\text{NH}(\text{CH}_2)_3-\text{OCH}_3$	$-\text{NH}(\text{CH}_2)_3-\text{OCH}_3$	$\text{C}_{20}\text{H}_{36}\text{N}_4\text{O}_2$	15.4	15.6
622.	$-\text{CH}_2\text{CH}_2-\text{N} \begin{matrix} \text{H} \\ \text{H} \end{matrix}$ 	$-\text{CO}-\text{NH}_2$		$-\text{N} \begin{matrix} \text{C}_2\text{H}_5 \\ \text{C}_2\text{H}_5 \end{matrix}$	$\text{C}_{21}\text{H}_{37}\text{N}_5\text{O}$	18.7	18.5
623.		$-\text{CH}_2-\text{OCH}_3$	$-\text{NH}_2$	$-\text{NH}_2$	$\text{C}_{13}\text{H}_{21}\text{N}_3\text{O}$	17.9	17.6
624.		$-\text{H}$	$-\text{NH}-$ 		$\text{C}_{23}\text{H}_{19}\text{N}_3$	12.5	12.7
625.	"	$-\text{CO}-\text{NH}_2$	$-\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$	$-\text{SO}_2$ 	$\text{C}_{22}\text{H}_{23}\text{N}_3\text{O}_3\text{S}$	10.3	10.7
626.	"	$-\text{CO}-\text{CH}_3$	"	$-\text{SO}_2\text{CH}_2-$ 	$\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_2\text{S}$	10.4	10.6
627.		$-\text{H}$	$-\text{NH}-\text{CH}_3$	$-\text{NH}-\text{CH}_3$	$\text{C}_{12}\text{H}_{14}\text{N}_4$	26.2	26.8
628.	"	$-\text{CO}-\text{NH}_2$	$-\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$	$-\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$	$\text{C}_{15}\text{H}_{19}\text{N}_5\text{O}$	24.6	25.0


-continued

No.	X	Y	Z ₁	Z ₂	Empirical Formula	Cal'd. N%	Found N%
629.		-CO-NH ₂	-NHCH ₂ CH ₂ OCH ₃		C ₁₅ H ₂₁ N ₅ O ₃ S	20.0	20.3
630.		-CO-NH ₂	-NH ₂	-OH	C ₁₃ H ₁₀ N ₄ O ₂ S	19.6	19.4

In those instances in the foregoing tables where only one value is presented for Z₁ and Z₂, this means that Z₁ and Z₂ are equal.

EXAMPLE 13

There are allowed to run 100 p.b.w. 3-methoxypropylamine into a suspension of 93.0 p.b.w. of 2,6-dichloro-3-cyano-4-methylpyridine in 200 p.b.w. ethyl alcohol for 2 hours at 20°C. This product is then agitated at room temperature for 18 hours. Then the resulting 2-chloro-3-cyano-4-methyl-6-(3'-methoxypropylamino)-pyridine of the formula




is drawn off, washed on the suction apparatus first with 120 p.b.w. ethyl alcohol and subsequently with water. The substance is analytically pure.

Analysis: C₁₁H₁₄ClN₃O — Calculated: 17.6% N, 14.6% Cl. Found: 17.9% N, 14.4% Cl.

EXAMPLE 14

Introduced into 1000 parts by weight methyl alcohol during cooling are 25.3 p.b.w. sodium. Added to this sodium methylate solution are 215 p.b.w. 2-chloro-3-cyano-4-methyl-6-(3'-methoxypropylamino)-pyridine, and subsequently the reaction solution is heated to the point of boiling for 24 hours under reflux. Then the methyl alcohol is distilled off and the residue stirred with water. The resulting 2-methoxy-3-cyano-4-methyl-6-(3'-methoxypropylamino)-pyridine of the formula




is an oil, which is isolated with ether. It may be purified by vacuum distillation.

Analysis: C₁₂H₁₇N₃O₂ — Calculated: 17.9% N, 26.4% —OCH₃. Found: 18.1% N, 26.1% —OCH₃.

EXAMPLE 15

a. 13.8 p.b.w. 4-nitraniline are diazotized at 0° to +5°C. in 300 p.b.w. of water to which 36.0 p.b.w. hydrochloric acid of 30 % have been added, with a solution consisting of 7.7 p.b.w. sodium nitrite in 50 p.b.w. of water. The filtered diazo solution is allowed to run into a solution of 500 p.b.w. of water, 25.0 p.b.w. hydrochloric acid of 30 % and 32.2 p.b.w. 2,6-bis-(3'-methoxy-n-propylamino)-3-cyano-4-methyl-pyridine, whereby the reaction temperature is maintained during the coupling at 0° to +5°C. by the addition of 500 parts of ice. The pH value of the batch is neutralized by the introduction of a solution of 65.2 p.b.w. sodium acetate in 200 p.b.w. of water, whereupon the coupling is quickly completed. The dyestuff being formed is then sucked off, washed with water until neutral and dried.

b. 1.0 g. of the dyestuff thus obtained which has the following formula




is finely dispersed and introduced while stirring into 2000 g. of water. This reaction mixture is adjusted to a pH value of 5 - 6 by the addition of acetic acid and then admixed with 4 g. ammonium sulfate and 2 g. of a commercial dispersing agent on the basis of a naphthalenesulfonic acid/formaldehyde condensation product. 100 g. of a polyester fabric on the basis of polyethylene glycolterephthalate are introduced into the dye liquor thus obtained and dyed for 1 1/2 hours at 130°C. After subsequent rinsing, reductive aftertreatment with an alkaline sodium dithionite solution of 0.2 % during 15 minutes at 60° - 70°C., rinsing and drying, one obtains a deep reddish orange dyeing which shows very good fastness properties.

EXAMPLE 16

a. 16.3 p.b.w. 2-cyano-4-nitraniline are diazotized at 15°C. in 100 p.b.w. glacial acetic acid, with 34.2 p.b.w. nitrosylsulfuric acid of 41.3 %. The limpid diazo solution is then allowed to run into a suspension of 25.8 p.b.w. 2-methoxy-3-cyano-4-methyl-6-(3'-methoxypropylamino)-pyridine and 66.0 g. sodium acetate in 160 p.b.w. ethanol. During the coupling the reaction temperature is maintained at 0 to +5°C. by cooling from the outside. When the coupling is completed, the dyestuff being formed is sucked off, washed first with ethanol, then with water and dried.

b. 30 p.b.w. of the dyestuff thus obtained (related to 1000 p.b.w. of the printing paste) which has the formula




are incorporated in a finely dispersed form into a printing paste containing 45.0 p.b.w. carob bean meal, 6.0 p.b.w. 3-nitrobenzenesulfonic acid sodium and 3.0 p.b.w. citric acid. When this printing paste is applied onto a polyester fabric one obtains after printing, drying and setting on the thermosetting stenter during 45 seconds at 215°C., rinsing and working up as described in the above Example 15 (b), a full reddish orange print which has very good fastness properties.

The dyestuff yields when printed onto a triacetate fabric and if it is employed in the form of the above printing paste and the printed fabric, after drying, is steamed for 10 minutes at a pressure of 2.5 atmospheres, rinsed, soaped, rinsed again and dried, a full reddish orange print having also very good fastness properties.

EXAMPLE 17

a. 29.7 g. 4-β-sulfatoethylsulfonyl-aniline-2-sulfonic acid are pasted up with 400 c.c. water, admixed with 25 c.c. 10 normal hydrochloric acid and diazotized by the addition of 100 c.c. 4 normal sodium dithionite solution. The reaction temperature is maintained at 0° to +3°C. by the addition of about 400 g. ice. The excess sodium, if any, is destroyed with little amidosulfonic acid and the diazo compound is added to a suspension of 14.8 g. 2,6-diamino-3-cyano-4-methyl-pyridine in 200 c.c. water to which 12.5 g. sodium bicarbonate have been added. The precipitated dyestuff is sucked off, and dried in vacuo. The reactive dye thus obtained has the following formula:



b. A mixture of mercerized or alkaliized cotton is printed with the following printing paste:

50 g.	of the above dyestuff
100 g.	urea
370 g.	hot water
450 g.	alginate thickener
20 g.	sodium bicarbonate
10 g.	sodium salt of the m-nitrobenzenesulfonic acid
1000 g.	


The print may be fixed either by steaming at 103° to 105°C. or by dry heating. In both cases the fabric is subsequently rinsed and soaped. The full greenish yellow print thus obtained has very good fastness properties.

Cationic azo dyes suited for dyeing and printing polyacrylonitrile and modified polyester fibers may be prepared according to the following Example:

EXAMPLE 18

a. 4.57 p.b.w. 4'-aminophenacyltrimethyl-ammonium chloride are dissolved in 20 p.b.w. of water and 7 p.b.w. concentrated hydrochloric acid, and diazotized at 0° - 5°C. with 4 p.b.w. 5 normal sodium nitrite solution. When the diazotization is completed, 3.52 p.b.w. 3-cyano-4-methyl-2,6-bis-methylamino-pyridine, dissolved in 10 p.b.w. glacial acetic acid, are added to the diazo solution and the reaction mixture is stirred at a pH value of 3 - 5 until the coupling is completed.

The dyestuff is precipitated by the addition of sodium chloride of 10 %, sucked off and dried. It has the following formula:




b. 1 g. of this dyestuff is pasted up with 2.5 g. acetic acid of 50 % and dissolved in 6 l. of water. To this dye liquor 1 g. sodium acetate and 10 g. sodium sulfate are added. Subsequently, 100 g. of a pre-washed polyacrylonitrile staple fiber yarn are introduced into the dye liquor which has a temperature of 60°C. This temperature is slowly raised to 100°C. and dyeing is carried out during 1 hour at boiling temperature. Subsequently, the temperature is lowered down to 70°C., the treated fabric is rinsed and dried. One obtains a clear deep orange dyeing with a very good fastness to light and to wetting, and a high absorption capacity.

c. 1 g. of the dyestuff is pasted up with 2 g. acetic acid of 50 % and dissolved in 5 l. of water which contains 1 g. sodium acetate, 100 g. of a pre-washed fabric of acid modified polyester fiber are then introduced at 60°C., the temperature is slowly raised and dyeing is carried out during 1 hour at 115°C. The temperature is then lowered down to 70°C. and the fabric is rinsed and dried. Obtained is a deep orange dyeing having a very good fastness to light and wetting.

Acid azo dyes which are suited for dyeing wool, silk and polyamide may be prepared according to the following Example:

EXAMPLE 19

a. 27.7 p.b.w. 4'-amino-azobenzene-4-sulfonic acid are diazotized in the usual manner and coupled in a slightly acid pH range with 23.6 p.b.w. 2,6-(β-hydroxyethylamino)-3-cyano-4-methyl-pyridine. The dyestuff being formed is salted out by the addition of sodium chloride of 10 %, sucked off and dried at 60°C. in vacuo. In the form of its free acid it corresponds to the following formula:




b. 1 g. of the above dyestuff is pasted up with 25 g. cold distilled water and dissolved in 975 g. boiling dis-

73

tilled water. The dyestuff solution is admixed with 2 g. acetic acid of 50 % and the reaction temperature is maintained at 60°C. At this temperature 100 g. of a pre-washed fabric of wool, silk or polyamide are introduced and the temperature is slowly raised to 100°C. and then maintained for 1 hour at boiling temperature. Subsequently, the temperature is lowered down to 70°C. and the fabric is rinsed and dried. Obtained is a deep yellowish red dyeing of high brilliancy and showing a very good fastness to light and to wetting.

We claim:

1. A compound of the formula




wherein X is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl

74

having three to eight carbon atoms, phenyl, benzyl or phenethyl; Z₁ is



-NH-OR₁₀ or



R₅ and R₆ are hydrogen, alkyl having one to six carbon atoms, hydroxyalkyl having one to six carbon atoms, cycloalkyl having three to eight carbon atoms, methoxyalkyl or ethoxyalkyl having one to six carbon atoms in the alkyl moiety, benzyl or phenethyl; R₁₀ is alkyl having one to six carbon atoms, benzyl or phenethyl; R₁₁ is alkyl having one to six carbon atoms and R₁₂ is alkyl having one to six carbon atoms, alkenyl having two to six carbon atoms, cycloalkyl having three to eight carbon atoms, phenyl, benzyl or phenethyl.

2. The compounds of claim 1 wherein X is methyl.

* * * * *