

(19)



(11) Publication number:

SG 174029 A1

(43) Publication date:

29.09.2011

(51) Int. Cl:

;

(12)

Patent Application

(21) Application number: **2011055605**

(71) Applicant:

**KYORIN PHARMACEUTICAL CO., LTD.
5, KANDA SURUGADAI 2-CHOME,
CHIYODA-KU, TOKYO, 1018311 JAPAN
JP**

(22) Date of filing: **07.08.2007**

(30) Priority: **JP 2006-215281 08.08.2006**

(72) Inventor:

**KOHNO, YASUSHI C/O DISCOVERY
RESEARCH LABORATORIES, KYORIN
PHARMACEUTICAL CO., LTD., 2399-1,
NOGI, NOGI-MACHI, SHIMOTSUGA-GUN,
TOCHIGI JAPAN JP
FUJII, KIYOSHI C/O DISCOVERY
RESEARCH LABORATORIES, KYORIN
PHARMACEUTICAL CO., LTD., 2399-1,
NOGI, NOGI-MACHI, SHIMOTSUGA-GUN,
TOCHIGI JAPAN JP
SAITO, TATSUHIRO C/O DISCOVERY
RESEARCH LABORATORIES, KYORIN
PHARMACEUTICAL CO., LTD., 2399-1,
NOGI, NOGI-MACHI, SHIMOTSUGA-GUN,
TOCHIGI JAPAN JP
KURIYAMA, KAZUHIKO C/
O DISCOVERY RESEARCH
LABORATORIES, KYORIN
PHARMACEUTICAL CO., LTD., 2399-1,
NOGI, NOGI-MACHI, SHIMOTSUGA-GUN,
TOCHIGI JAPAN JP
YASUE, TOKUTAROU C/O DISCOVERY
RESEARCH LABORATORIES, KYORIN
PHARMACEUTICAL CO., LTD., 2399-1,
NOGI, NOGI-MACHI, SHIMOTSUGA-GUN,
TOCHIGI JAPAN JP**

(54) Title:
**AMINOALCOHOL DERIVATIVE AND
IMMUNOSUPPRESSANT CONTAINING THE SAME AS
ACTIVE INGREDIENT**

(57) Abstract:
ABSTRACT AMINOALCOHOL DERIVATIVE AND
IMMUNOSUPPRESSANT CONTAINING THE SAME AS
ACTIVE INGREDIENT An amino alcohol derivative useful as
an excellent immunosuppressive agent is provided. As a result
of intensive research to create a highly safe compound which
has an excellent immunosuppressive activity, it was discovered
that an amino alcohol derivative represented by the general
formula (1), [err] has excellent immunosuppressive activity. [No
Drawing]


ABSTRACT

AMINOALCOHOL DERIVATIVE AND IMMUNOSUPPRESSANT CONTAINING THE SAME AS ACTIVE INGREDIENT

5

An amino alcohol derivative useful as an excellent immunosuppressive agent is provided.

As a result of intensive research to create a highly safe compound which has an excellent immunosuppressive activity, it was
10 discovered that an amino alcohol derivative represented by the general formula (1),



has excellent immunosuppressive activity.

15 [No Drawing]

DESCRIPTION

AMINOALCOHOL DERIVATIVE AND IMMUNOSUPPRESSANT CONTAINING THE SAME AS ACTIVE INGREDIENT

TECHNICAL FIELD

5 [0001]

The present invention relates to an amino alcohol derivative useful as an immunosuppressive agent, and a salt and a hydrate thereof.

BACKGROUND ART

[0002]

10 Immunosuppressive agents are widely utilized as a treatment agent for autoimmune diseases such as rheumatoid arthritis, nephritis, degenerative knee arthritis, and systemic lupus erythematosus, chronic inflammatory diseases such as inflammatory bowel disease, and allergic diseases such as asthma and dermatitis.

15 In particular, with the progress in medical technology, in recent years large numbers of transplant operations of tissues, organs and the like are being performed. In such medical practice, whether the post-transplant rejection reaction can be well controlled determines whether the transplant is successful or not.

20 Immunosuppressive agents are playing an important role in this area as well.

[0003]

In organ transplants, antimetabolites represented by azathioprine and mycophenolate mofetil, calcineurin inhibitors

25 represented by cyclosporine A and tacrolimus, and adrenocortical

hormones represented by prednisolone are used. However, in some cases the effectiveness of these pharmaceuticals has been insufficient, or blood concentration monitoring has been essential in order to avoid serious side effects such as kidney damage.

5 Therefore, in terms of their effectiveness and side effects, these pharmaceuticals have not always been satisfactory.

[0004]

Furthermore, to alleviate the side effects of the immunosuppressive agent and to obtain a sufficient immunosuppressive activity, multiple drug therapy using a plurality of pharmaceuticals having different action mechanisms is typical. Thus, there is also a need for development of a new type of pharmaceutical which has an action mechanism different from that of the above-described immunosuppressive agents.

15 [0005]

Recently, reports of various amino alcohol derivatives, such as 2-amino-1,3-propanediol derivatives and 2-aminoethanol derivatives, as novel immunosuppressive agents have been drawing attention. However, these amino alcohol derivatives themselves do not have an immunosuppressive activity. These drugs are metabolized after being administered into the body to produce a phosphate, and it is this phosphate which is the true physiologically active substance. The produced phosphate exhibits agonistic activity and antagonist activity against various sphingosine 1-phosphate (S1P) receptors. Among them, in 2002 it was reported for the first time

that the agonistic activity against S1P₁ receptors expresses an immunosuppressive activity by regulating white blood cell migration (Non-patent Documents 1 and 2). It has also been disclosed that, in addition to being effective for various organ transplants and GVHD, the series of derivatives introduced in these Non-patent documents are effective for rheumatoid arthritis, lupus nephritis, systemic lupus erythematosus, chronic thyroiditis, multiple sclerosis, myasthenia gravis, type I and II diabetes mellitus, autoimmune diseases such as Crohn's disease, allergic diseases such as atopic dermatitis, allergic rhinitis, allergic conjunctivitis, and allergic contact dermatitis, and inflammatory diseases such as inflammatory bowel disease or ulcerative colitis (Patent Documents 1 and 2). Since these reports, not only amino alcohol derivatives, but various compounds, such as amino phosphate derivatives and amino carboxylic acid derivatives, have been disclosed as immunosuppressive agents or S1P₁ to S1P₅ receptor regulators focused on S1P₁ receptors (Patent Documents 3 to 66). [0006]

Furthermore, since S1P₄ receptors are largely concentrated in immunocompetent cells, such as leukocytes, and in organs which greatly contribute to the immune system, it is suggested that S1P₄ receptors have a strong contribution to the immune system. In fact, compounds having an S1P₄ agonistic activity have been disclosed for autoimmune diseases such as SLE and rheumatism, asthma, allergic diseases such as atopic dermatitis, and inflammatory disease

remedies (Patent Documents 34, 39, and 50).

[0007]

Thus, while a great deal of attention is being paid to S1P
receptor agonist drugs which may have hidden potential in a wide
5 variety of medical applications, not all S1P receptor agonist drugs
provide a desirable action on the body.

[0008]

For example, an S1P receptor agonist drug which has exhibited
effectiveness in clinical trials in suppressing organ transplant
10 rejection was found to produce bradycardia as a side effect after
administration. This effect was reported to probably be caused by
agonistic activity against the S1P₃ receptor (Non-patent Documents
3 and 4). Furthermore, agonistic activity against the S1P₃ receptor
has also been reported to cause in experimental animal models effects
15 such as obstruction of myocardial blood flow (Non-patent Document
5), cerebral arterial spasms (Non-patent Document 6), and pulmonary
edema (Non-patent Document 7).

[Patent Document 1] WO 0218395 pamphlet

[Patent Document 2] WO 02076995 pamphlet

20 [Patent Document 3] WO 9408943 pamphlet

[Patent Document 4] Japanese Patent Application Laid-Open No.
Hei 9-2579602

[Patent Document 5] WO 0206268 pamphlet

[Patent Document 6] Japanese Patent Application Laid-Open No.

25 2002-53572

	[Patent Document 7]	Japanese Patent Application Laid-Open No.
	2002-167382	
	[Patent Document 9]	WO 02076995 pamphlet
	[Patent Document 10]	Japanese Patent Application Laid-Open No.
5	2003-137894	
	[Patent Document 11]	WO 03040097 pamphlet
	[Patent Document 12]	WO 02064616 pamphlet
	[Patent Document 13]	WO 02062389 pamphlet
	[Patent Document 14]	Japanese Patent Application Laid-Open No.
10	2002-316985	
	[Patent Document 15]	Japanese Patent Application Laid-Open No.
	2003-267936	
	[Patent Document 16]	WO 03051876 pamphlet
	[Patent Document 17]	WO 03061567 pamphlet
15	[Patent Document 18]	WO 03062248 pamphlet
	[Patent Document 19]	WO 03062252 pamphlet
	[Patent Document 20]	WO 03073986 pamphlet
	[Patent Document 21]	WO 03074008 pamphlet
	[Patent Document 22]	WO 03105771 pamphlet
20	[Patent Document 23]	WO 04010949 pamphlet
	[Patent Document 24]	WO 04024673 pamphlet
	[Patent Document 25]	WO 04058149 pamphlet
	[Patent Document 26]	WO 04071442 pamphlet
	[Patent Document 27]	WO 04096752 pamphlet
25	[Patent Document 28]	WO 04096757 pamphlet

	[Patent Document 29]	WO 04103279 pamphlet
	[Patent Document 30]	WO 04103306 pamphlet
	[Patent Document 31]	WO 04103309 pamphlet
	[Patent Document 32]	WO 04110979 pamphlet
5	[Patent Document 33]	WO 04113330 pamphlet
	[Patent Document 34]	WO 04074297 pamphlet
	[Patent Document 35]	WO 05014603 pamphlet
	[Patent Document 36]	WO 05020882 pamphlet
	[Patent Document 37]	WO 04002531 pamphlet
10	[Patent Document 38]	WO 05032465 pamphlet
	[Patent Document 39]	WO 05041899 pamphlet
	[Patent Document 40]	WO 05058848 pamphlet
	[Patent Document 41]	WO 05070886 pamphlet
	[Patent Document 42]	WO 05082089 pamphlet
15	[Patent Document 43]	WO 05082841 pamphlet
	[Patent Document 44]	WO 05021503 pamphlet
	[Patent Document 45]	WO 05040091 pamphlet
	[Patent Document 46]	WO 05085179 pamphlet
	[Patent Document 47]	WO 05118523 pamphlet
20	[Patent Document 48]	WO 05014525 pamphlet
	[Patent Document 49]	WO 06020951 pamphlet
	[Patent Document 50]	WO 06001463 pamphlet
	[Patent Document 51]	WO 03029184 pamphlet
	[Patent Document 52]	WO 03029205 pamphlet
25	[Patent Document 53]	WO 04026817 pamphlet

[Patent Document 54] WO 04074297 pamphlet

[Patent Document 55] WO 05021503 pamphlet

[Patent Document 56] Japanese Patent Application Laid-Open No.
2004-307439

5 [Patent Document 57] Japanese Patent Application Laid-Open No.
2004-307440

[Patent Document 58] Japanese Patent Application Laid-Open No.
2004-307441

[Patent Document 59] Japanese Patent Application Laid-Open No.
10 2004-307442

[Patent Document 60] WO 06041015 pamphlet

[Patent Document 61] Japanese Patent Application Laid-Open No.
2004-137208

[Patent Document 62] Japanese Patent Application Laid-Open No.
15 2005-41867

[Patent Document 63] Japanese Patent Application Laid-Open No.
2005-47899

[Patent Document 64] WO 05040091 pamphlet

[Patent Document 65] WO 05063671 pamphlet

20 [Patent Document 66] WO 05079788 pamphlet

[Non-patent Document 1] S. Mandala et al., Science, 296, 346 (2002).

[Non-patent Document 2] V. Brinkmann et al., J. Biol. Chem., 277,
21453 (2002).

[Non-patent Document 3] M. G. Sanna et al., J. Biol. Chem., 279,
25 13839 (2004).

[Non-patent Document 4] M. Forrest et al., J. Pharmacol. Exp. Ther., 309, 758 (2004).

[Non-patent Document 5] B. Levkau et al., Circulation, 110, 3358 (2004).

5 [Non-patent Document 6] S. Salomone et al., Eur. J. Pharmacol. 469, 125 (2003).

[Non-patent Document 7] Y. Gon et al., PNAS 102, 9270 (2005).

DISCLOSURE OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

10 [0009]

It is an object of the present invention to provide an amino alcohol derivative having excellent immunosuppressive activity and few side effects.

MEANS FOR SOLVING THE PROBLEMS

15 [0010]

As a result of intensive research concerning immunosuppressive agents which have a different action mechanism to that of antimetabolites and calcineurin inhibitors, the present inventors discovered that a novel amino alcohol derivative has high safety and excellent immunosuppressive activity, thereby completing the present invention.

[0011]


Specifically, the present invention relates to:

1) An amino alcohol derivative, represented by the general formula

25 (1),

[0012]

[Chemical formula 1]



[0013]


5 [wherein R¹ represents a chlorine atom or a straight-chain alkyl group having 1 to 3 carbon atoms or trifluoromethyl group, R² represents a fluorine atom or a chlorine atom, R³ represents a straight-chain alkyl group having 1 to 3 carbon atoms, X represents an oxygen atom or a sulfur atom, and n denotes 2 or 3], or a
10 pharmaceutically acceptable salt or hydrate thereof.

[0014]

2) The amino alcohol derivative according to 1), wherein the compound represented by the general formula (1) is a compound represented by the general formula (1a),

15 [0015]

[Chemical formula 2]



[0016]

[wherein R³, X, and n are as described above], or a pharmaceutically
20 acceptable salt or hydrate thereof.

[0017]

3) The amino alcohol derivative according to 1) or 2), wherein in the general formulae (1) or (1a), R³ is a methyl group, or a pharmaceutically acceptable salt or hydrate thereof.

[0018]

5 4) The amino alcohol derivative according to 1), wherein the compound represented by the general formula (1) is,

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylpentan-1-ol,

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]

10 -2-methylpentan-1-ol,

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylbutan-1-ol,

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylbutan-1-ol,

15 (R)-2-amino-5-[2-chloro-4-(3-ethylphenylthio)phenyl]-2-methylpentan-1-ol,

(R)-2-amino-5-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol, or

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]

20 -2-propylpentan-1-ol, or a pharmaceutically acceptable salt or hydrate thereof.

[0019]


5) The amino alcohol derivative according to 1) ,or a pharmaceutically acceptable salt or hydrate thereof, which is

25 obtainable by a step of allowing a compound represented by the

general formula (2),

[0020]

[Chemical formula 3]




5 [0021]

[wherein R¹ represents a chlorine atom or a straight-chain alkyl group having 1 to 3 carbon atoms or trifluoromethyl group, R² represents a fluorine atom or a chlorine atom, A represents a halogen atom, X represents an oxygen atom or a sulfur atom, and n denotes

10 2 or 3] and a compound represented by the general formula (10),

[0022]

[Chemical formula 4]



[0023]

15 [wherein R³ represents a straight-chain alkyl group having 1 to 3 carbon atoms and R⁴ represents an alkyl group having 1 to 6 carbon atoms] to act in the presence of a base, and a step of subjecting the resultant product to acidolysis, then further protecting a nitrogen atom with a t-butoxycarbonyl group, reducing, and

20 deprotecting the nitrogen atom, or a pharmaceutically acceptable salt or hydrate thereof.

[0024]

6) A pharmaceutical comprising the amino alcohol derivative, or a pharmaceutically acceptable salt or hydrate thereof, according to any of 1) to 5) as an active ingredient.

[0025]

5 7) The pharmaceutical according to 6), which is an immunosuppressive agent.

8) The pharmaceutical according to 6), which is a rejection reaction preventive or treatment agent in organ transplants or bone marrow transplants.

10 9) A pharmaceutical comprising the amino alcohol derivative, or a pharmaceutically acceptable salt or hydrate thereof, according to any of 1) to 5) and a calcineurin inhibitor in combination.

EFFECT OF THE INVENTION

[0028]

15 According to the present invention, an amino alcohol derivative having excellent immunosuppressive activity and safety can be provided. The compound of the present invention is useful as a preventive or treatment agent for rejection in organ transplants or bone marrow transplants, a preventive or treatment agent for
20 autoimmune diseases such as inflammatory bowel disease, systemic lupus erythematosus, Crohn's disease, nephrotic syndrome, glomerular sclerosis, glomerular nephritis, multiple sclerosis, and myasthenia gravis, a preventive or treatment agent for rheumatoid arthritis, a preventive or treatment agent for psoriasis, allergic
25 contact dermatitis, and atopic dermatitis, a preventive or treatment

agent for hepatitis, hepatic steatosis, toxic liver injury, liver cirrhosis or diabetes mellitus-derived liver disease, a preventive or treatment agent for allergic rhinitis, allergic conjunctivitis and the like, and a preventive or treatment agent for pulmonary
5 fibrosis, idiopathic interstitial pneumonia, and bronchial asthma.

BEST MODE FOR CARRYING OUT THE INVENTION

[0029]

In the present invention, the straight-chain alkyl group having 1 to 3 carbon atoms of R^1 and R^3 is a methyl group, an ethyl group,
10 or an n-propyl group.

[0030]

From the perspective of obtaining high safety, R^1 is preferably an ethyl group, a propyl group, or a trifluoromethyl group, and more preferably is a trifluoromethyl group. Furthermore, R^3 is
15 preferably a methyl group, and n is preferably 3.

[0031]

Furthermore, from the perspective of obtaining high immunosuppressive activity, X is preferably a sulfur atom, and the configuration of R^3 is preferably a configuration produced as the
20 principal product via the below-described synthesis route B (using the compound (10)).

[0032]

In the present invention, examples of pharmaceutically acceptable salts include acid addition salts such as hydrochloride
25 salts, hydrobromic acid salts, acetic acid salts, trifluoroacetic

acid salts, methanesulfonic acid salts, citric acid salts, or tartaric acid salts.

[0033]


The compound according to the present invention represented
5 by the general formula (1) can be produced, for example, via the synthesis route A shown below.

[0034]

<Synthesis Route A>

[0035]

10 [Chemical formula 5]




[0036]

In the synthesis route A, the compound represented by the general formula (3),

15 [0037]

[Chemical formula 6]




[0038]

[wherein R^1 , R^2 , R^3 , R^4 , X, and n are as described above], can be produced by allowing a compound represented by the general formula (2),

5 [0039]

[Chemical formula 7]




[0040]

[wherein R^1 , R^2 , A, X, and n are as described above], and a compound represented by the general formula (7),

[0041]

[Chemical formula 8]



[0042]

15 [wherein R^3 and R^4 are as described above] to act in the presence of a base (step A-1).

[0043]

The reaction can be carried out using methanol, ethanol, 1,4-dioxane, dimethylsulfoxide (DMSO), N,N-dimethylformamide (DMF), tetrahydrofuran (THF) and the like as a reaction solvent, in the presence of an inorganic base such as sodium hydride, potassium hydride, sodium methoxide, sodium ethoxide, sodium t-butoxide, potassium methoxide, potassium ethoxide, potassium t-butoxide, and


potassium carbonate, at 0°C to reflux temperature as the reaction temperature, and preferably at 80°C to 100°C.

[0044]

In the synthesis route A, the compound represented by the
5 general formula (4),

[0045]

[Chemical formula 9]



[0046]

10 [wherein R¹, R², R³, R⁴, X, and n are as described above], can be produced by hydrolyzing the compound represented by the general formula (3) (step A-2).

[0047]

The reaction can be carried out in the presence of a base such
15 as aqueous sodium hydroxide, aqueous potassium hydroxide, and aqueous lithium hydroxide, using methanol, ethanol, 1,4-dioxane, DMF, DMSO, THF and the like as a reaction solvent, at a reaction temperature of 0°C to reflux temperature. Preferably, the reaction is carried out using potassium hydroxide as the base, in an ethanol
20 solvent, by reacting at 50°C.

[0048]

Although the compound according to the present invention is preferably a specific optically-active substance, when the optical resolution is carried out is not especially limited. At this stage,


optical resolution may be carried out by HPLC using a chiral column, whereby the desired compound having a chiral center can be obtained.

[0049]

In the synthesis route A, the compound represented by the
5 general formula (5),

[0050]

[Chemical formula 10]



[0051]

10 [wherein R⁵ represents an alkyl group having 1 to 6 carbon atoms, and R¹, R², R³, R⁴, X, and n are as described above], can be produced by subjecting the compound represented by the general formula (4) to Curtius rearrangement (step A-3).

[0052]

15 In the reaction, typical methods for converting a carboxyl group into a carbamate may be employed. For example, a method which combines, for example, chloroethyl carbonate and NaN₃, or oxalyl chloride and NaN₃, or a method which uses only diphenylphosphoryl azide (DPPA) may be utilized. The reaction is preferably carried
20 out by, after heating diphenylphosphoryl azide to reflux in the presence of an organic base, such as triethylamine, in benzene or toluene solvent, charging the resultant product with an alcohol represented by the general formula (8),

R⁵OH (8)

[herein R⁵ is as described above], and continuing to heat the resultant solution under stirring, or, after removing the solvent used in the above reaction, such as benzene or toluene, by evaporation, by heating to reflux using the alcohol represented by the general formula (8) as a reaction solvent.

[0053]


At this stage, optical resolution may be carried out by HPLC using a chiral column, whereby the desired compound having a chiral center can be obtained.

[0054]

In the synthesis route A, the compound represented by the general formula (6),

[0055]

[Chemical formula 11]



[0056]

[wherein R¹, R², R³, R⁵, X, and n are as described above], can be produced by reducing the compound represented by the general formula

(5) (step A-4).

[0057]

The reaction can be carried out using borane, an alkyl borane derivative like 9-borabicyclo[3.3.1]nonane (9-BBN), or a metal

hydride complex compound, such as diisobutylaluminum hydride ((iBu)₂AlH), sodiumborohydride (NaBH₄), lithiumborohydride (LiBH₄), and lithium aluminum hydride (LiAlH₄), preferably LiBH₄, using THF, 1,4-dioxane, ethanol, or methanol as a reaction solvent, at a temperature of 0°C to reflux temperature, and preferably at room temperature.

[0058]

Furthermore, at this stage also, optical resolution may be carried out by HPLC using a chiral column, whereby the desired compound having a chiral center can be obtained.

[0059]

In the synthesis route A, the compound represented by the general formula (1) can be produced by subjecting the compound represented by the general formula (6) to acidolysis or hydrolysis (step A-5).

[0060]

The reaction can be carried out at room temperature to reflux temperatures in an inorganic acid or organic acid, such as hydrochloric acid, hydrobromic acid, methanesulfonic acid, acetic acid, and trifluoroacetic acid, or at room temperature to reflux temperature by adding an organic solvent such as methanol, ethanol, THF, and 1,4-dioxane to an inorganic acid or organic acid, such as hydrochloric acid, hydrobromic acid, methanesulfonic acid, acetic acid, and trifluoroacetic acid. The reaction may also be carried out in the presence of a base such as aqueous sodium hydroxide,


aqueous potassium hydroxide, and aqueous lithium hydroxide, using methanol, ethanol, THF, 1,4-dioxane, DMSO, and DMF as a reaction solvent, at a temperature of 0°C to reflux temperature, and preferably 80 to 100°C.

5 [0061]

In the synthesis route A, among the compounds represented by the general formula (5), compounds in which R⁵ represents a t-butyl group, specifically, a compound represented by the general formula (5a),

10 [0062]

[Chemical formula 12]




[0063]

[wherein Boc represents a t-butoxycarbonyl group, and R¹, R², R³, R⁴, X, and n are as described above], and among the compounds represented by the general formula (6) in the synthesis route A, compounds in which R⁵ represents a t-butyl group, specifically, a compound represented by the general formula (6a),

[0064]

20 [Chemical formula 13]



[0065]


[wherein R^1 , R^2 , R^3 , X , Boc , and n are as described above], can be produced by the synthesis route B.

[0066]

5 <Synthesis Route B>

[0067]

[Chemical formula 14]




[0068]

10 In the synthesis route B, the compound represented by the general formula (9),

[0069]

[Chemical formula 15]




15 [0070]

[wherein R^1 , R^2 , R^3 , R^4 , X , and n are as described above], can be produced by allowing a compound represented by the general formula

(2) and a compound represented by the general formula (10),

[0071]

[Chemical formula 16]



5 [0072]

[wherein R³ and R⁴ are as described above] to act in the presence of a base (step B-1).

[0073]

The reaction can be carried out using a reaction solvent such
10 as 1,4- dioxane, THF, and ether, using a base such as n-butyllithium and lithium diisopropyl amide, preferably n-butyllithium, and treating a compound represented by the general formula (10) at -78°C, then allowing a compound represented by general formula (2) to act at -78°C, and reacting while gradually increasing the temperature
15 to room temperature.

[0074]

In the synthesis route B, the compound represented by the general formula (5a) can be produced by subjecting a compound represented by the general formula (9) to acidolysis, and then
20 protecting the nitrogen atom with a t-butoxycarbonyl group (Boc group) (step B-2).

[0075]

In the reaction, an amino ester can be obtained using methanol,

ethanol, THF, 1,4-dioxane, or ethyl acetate in which hydrochloric acid is dissolved, and preferably 1,4-dioxane containing hydrochloric acid, by reacting at reflux temperature, then neutralizing with a base. Furthermore, it is preferred to allowing
5 it to act with Boc_2O at 0°C to room temperature using ethyl acetate, THF, DMF, 1,4-dioxane, methylene chloride, chloroform, methanol, ethanol, acetonitrile or the like as a solvent.

[0076]

In the synthesis route B, the compound represented by the
10 general formula (6a) can be produced by reducing a compound represented by the general formula (5a) (step B-3).

[0077]

The reaction can be carried out using borane, an alkyl borane derivative like 9-BBN, or a metal hydride complex compound, such
15 as $(\text{iBu})_2\text{AlH}$, NaBH_4 , LiBH_4 , and LiAlH_4 , preferably LiBH_4 , using THF, 1,4-dioxane, ethanol, or methanol as a reaction solvent, at a temperature of 0°C to reflux temperature, and preferably at room temperature.

[0078]

20 It is noted that concerning the synthesis method of the compound represented by the general formula (2), the compound may be produced by the methods described in the respective pamphlets of WO 03029184, WO 03029205, WO 04026817, WO 04074297, and WO 050444780.

[0079]

25 The compound of the present invention is metabolized in the

body to produce a phosphate (phosphoric acid ester) . As a result,
the compound of the present invention is useful as an excellent
preventive or treatment agent for rejection in organ transplants
or bone marrow transplants, a preventive or treatment agent for
5 autoimmune diseases such as inflammatory bowel disease, systemic
lupus erythematosus, Crohn's disease, nephrotic syndrome,
glomerular sclerosis, glomerular nephritis, multiple sclerosis,
and myasthenia gravis, a preventive or treatment agent for rheumatoid
arthritis, a preventive or treatment agent for psoriasis, allergic
10 contact dermatitis, and atopic dermatitis, a preventive or treatment
agent for hepatitis, hepatic steatosis, toxic liver injury, liver
cirrhosis or diabetes mellitus-derived liver disease, a preventive
or treatment agent for allergic rhinitis, allergic conjunctivitis
and the like, and a preventive or treatment agent for pulmonary
15 fibrosis, idiopathic interstitial pneumonia, and bronchial asthma.
[0080]

In the case of using as above, the required dose of course
depends on the administration method, the specific condition to
be treated, and the desired effects. However, generally, a daily
20 dose of about 0.03 to 2.5 mg per kg of body weight is preferred.
For mammals such as humans, the recommended daily dose is in the
range of about 0.5 mg to about 100 mg. Preferably, administration
is carried out in divided doses of four times or less per day, or
in retard form. A suitable unit dose form for oral administration
25 includes about 1 to 50 mg of active ingredient.

[0081]

The compound of the present invention may be administered by an arbitrary conventional route, especially enterally, for example orally, for example in the form of a tablet or a capsule, or
5 parenterally, for example in the form of an injectable solution or a suspension, locally, for example in the form of a lotion, a gel, an ointment, or a cream, or nasally or in the form of a suppository. A pharmaceutical composition containing the compound of the present invention in free form or a pharmaceutically acceptable salt thereof
10 together with at least one kind of pharmaceutically acceptable carrier or diluent may be produced by a conventional method of mixing with the pharmaceutically acceptable carrier or diluent.

Furthermore, the compound of the present invention is also useful when used together with an immunosuppressive agent and/or a
15 pharmaceutical which has an anti-inflammatory activity having a different mechanism. Examples of pharmaceuticals which can be used together include immunosuppressive agents used in the treatment and prevention of acute or chronic rejection of allogeneic transplants and heterologous transplantats, inflammatory diseases,
20 and autoimmune diseases, immunosuppressive agents having an immunomodulatory activity and/or anti-inflammatory agents having an anti-inflammatory or malignant cell growth inhibition activity. Specific examples include the calcineurin inhibitors cyclosporin A and FK506, the mTor inhibitors rapamycin,

25 40-O-(2-hydroxymethyl)-rapamycin, CCI779, and ABT578, the

ascomycins ABT281 and ASM981 which have an immunosuppressive activity, mycophenolic acid, mycophenolate mofetil, azathioprine, mizoribine, cyclophosphamide and the like. Further examples include the antifolate methotrexate, adrenal cortical steroids which exhibit broad anti-inflammatory activity, auranofin, actarit, mesalazine, or sulfasalazine etc. which have an immunomodulatory activity, infliximab which is an anti-TNF α antibody, MRA which is an anti-IL-6 receptor antibody, Natarizumab which is an anti-integrin antibody and the like.

10 [0082]

(Examples)

Next, the present invention will be described with the following specific examples. However, the present invention is not limited by these examples.

15 [0083]

Furthermore, as the intermediates and the like represented by the general formula (2), the compounds in the pamphlets of WO 03029184, WO 03029205, WO 04026817, WO 04074297, and WO 050444780 may be utilized. Furthermore,

20 (5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine,
(5S)-3,6-dimethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine,
and (5S)-2-allyl-3,6-diethoxy-5-isopropyl-2,5-dihydropyrazine
may be synthesized according to Ulrich Shollkopf et.al, Synthesis
969 (1981) and Chunrong Ma et.al., J. Org. Chem., 66, 4525 (2001).
25 Intermediates and the like which were newly synthesized based on

the experiment procedures described in these reference documents will now be described as the following reference examples.


[0084]

<Reference Example 1>

5 2-Fluoro-4-(3-trifluoromethylphenylthio)benzaldehyde

[0085]

[Chemical formula 17]



[0086]

10 Under an argon atmosphere, ethyldiisopropylamine (7.0 mL),
tris(dibenzylideneacetone)dipalladium(0) chloroform adduct (518
mg), xantphos (578 mg), and 3-trifluoromethylthiophenol (3.56 g)
were added at room temperature into a solution of
4-bromo-2-fluorobenzaldehyde (4.06 g) in 1,4-dioxane (42 mL), and
15 the resultant solution was heated to reflux for 5 hours. To the
reaction solution added water, extracted with ethyl acetate,
washed with water and saturated brine in that order, and then dried
over anhydrous sodium sulfate. The solvent was evaporated, and the
resultant residue was purified by silica gel column chromatography
20 (hexane : ethyl acetate = 30 : 1) to obtain the target product (4.08
g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 6.86 (1H, dd, J = 10, 1.8 Hz), 7.02 (1H,
dd, J = 7.9, 1.8 Hz), 7.58 (1H, t, J = 7.9 Hz), 7.68-7.73 (2H, m),
7.76 (1H, t, J = 7.9 Hz), 7.80 (1H, s), 10.26 (1H, s)

EIMS (+) : 300 [M]⁺.


[0087]

<Reference Example 2>

2-Chloro-4-(3-chlorophenylthio)benzaldehyde

5 [0088]

[Chemical formula 18]



[0089]

3-Chlorobenzenethiol and 2-chloro-4-fluorobenzaldehyde were
10 reacted according to the same experiment procedures as in Reference
Example 1 of the pamphlet of WO 03029205 to obtain the target product
as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 7.11 (1H, dd, J = 9.2, 1.8 Hz), 7.17 (1H,
d, J = 1.8 Hz), 7.36-7.44 (3H, m), 7.52 (1H, t, J = 1.8 Hz), 7.80
15 (1H, d, J = 7.9 Hz), 10.37 (1H, s)

EIMS (+) : 282 [M]⁺.


[0090]

<Reference Example 3>

2-Chloro-4-(3-methylphenoxy)benzaldehyde

20 [0091]

[Chemical formula 19]



[0092]

m-Cresol and 2-chloro-4-fluorobenzaldehyde were reacted according to the same experiment procedures as in Reference Example 1 of the pamphlet of WO 03029184 to obtain the target product as a colorless powder.

5 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 2.38 (3H, s), 6.87-6.96 (4H, m), 7.07 (1H, d, $J = 7.3$ Hz), 7.31 (1H, t, $J = 7.6$ Hz), 7.90 (1H, d, $J = 8.6$ Hz), 10.36 (1H, s).

EIMS (+) : 246 $[\text{M}]^+$.


[0093]

10 <Reference Example 4>

2-Chloro-4-(3-ethylphenylthio)benzaldehyde

[0094]

[Chemical formula 20]



15 [0095]

3-Ethylbenzenethiol and 2-chloro-4-fluorobenzaldehyde were reacted according to the same experiment procedures as in Reference Example 1 of the pamphlet of WO 03029205 to obtain the target product as a colorless oil.

20 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.26 (3H, t, $J = 7.3$ Hz), 2.68 (2H, q, $J = 7.3$ Hz), 7.04-7.11 (2H, m), 7.28-7.40 (4H, m), 7.76 (1H, d, $J = 8.6$ Hz), 10.35 (1H, s).

EIMS (+) : 276 $[\text{M}]^+$.


[0096]

<Reference Example 5>

2-Chloro-4-(3-propylphenoxy)benzaldehyde

[0097]

[Chemical formula 21]



5

[0098]

3-Propylphenol and 2-chloro-4-fluorobenzaldehyde were reacted according to the same experiment procedures as in Reference Example 1 of the pamphlet of WO 03029184 to obtain the target product as a pale brown oil.

10

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 0.95 (3H, t, $J = 7.3$ Hz), 1.62-1.68 (2H, m), 2.61 (2H, t, $J = 7.3$ Hz), 6.89-6.94 (3H, m), 6.96 (1H, d, $J = 2.1$ Hz), 7.08 (1H, d, $J = 7.9$ Hz), 7.31-7.35 (1H, m), 7.90 (1H, d, $J = 8.9$ Hz), 10.36 (1H, d, $J = 0.6$ Hz).

15 EIMS(+) : 274 [M] $^+$.


[0099]

<Reference Example 6>

[2-Chloro-4-(3-ethylphenylthio)phenyl]acetaldehyde

[0100]

20 [Chemical formula 22]



[0101]

The compound of Reference Example 4 was reacted according to

the same experiment procedures as in Reference Example 326 of the pamphlet of WO 04074297 to obtain the target product as a pale yellow oil.


[0102]

5 <Reference Example 7>

Ethyl 3-[2-chloro-4-(3-ethylphenylthio)phenyl]acrylate

[0103]

[Chemical formula 23]



10 [0104]

The compound of Reference Example 4 was reacted according to the same experiment procedures as in Reference Example 10 of the pamphlet of WO 03029205 to obtain the target product as a pale yellow oil.

15 EIMS(+) : 346 [M]⁺.


[0105]

<Reference Example 8>

3-[2-Chloro-4-(3-ethylphenylthio)phenyl]propan-1-ol

[0106]

20 [Chemical formula 24]



[0107]

The compound of Reference Example 7 was reacted according to

the same experiment procedures as in Reference Example 19 of the pamphlet of WO 03029205, and the resultant product was then reduced according to the same experiment procedures as in Reference Example 35 of the pamphlet of WO 03029205, to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz,): δ 1.22 (3H, t, $J = 7.3$ Hz), 1.84 -1.90 (2H, m), 2.62 (2H, q, $J = 7.6$ Hz), 2.78-2.82 (2H, m), 3.69 (2H, t, $J = 6.1$ Hz), 7.10-7.18 (4H, m), 7.23-7.29 (3H, m).


[0108]

<Reference Example 9>

3-[2-Chloro-4-(3-propylphenoxy)phenyl]propan-1-ol

[0109]

[Chemical formula 25]



[0110]

The compound of Reference Example 5 was successively reacted according to the same procedures as in Reference Example 7 and then Reference Example 8 to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz,): δ 0.94 (3H, t, $J = 7.3$ Hz), 1.37 (1H, br s), 1.58-1.68 (2H, m), 1.85-1.92 (2H, m), 2.57 (2H, t, $J = 7.6$ Hz), 2.80 (2H, t, $J = 7.6$ Hz), 3.70 (2H, dt, $J = 6.1, 4.6$ Hz), 6.80-6.85 (3H, m), 6.95 (1H, d, $J = 7.9$ Hz), 7.00 (1H, d, $J = 2.8$ Hz), 7.17 (1H, d, $J = 8.3$ Hz), 7.24 (1H, t, $J = 7.9$ Hz).

EIMS(+) : 304 [M]⁺.


[0111]

<Reference Example 10>

3-[2-Fluoro-4-(3-trifluoromethylphenylthio)phenyl]propan-1-ol

5 [0112]

[Chemical formula 26]



[0113]

The compound of Reference Example 1 was successively reacted
10 according to the same procedures as in Reference Example 7 and then
Reference Example 8 to obtain the target product as a colorless
oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.88 (2H, tt, J = 6.7, 6.1 Hz), 2.75 (2H,
t, J = 6.7 Hz), 3.69 (2H, t, J = 6.1 Hz), 7.05 (1H, dd, J = 10,
15 1.8 Hz), 7.10 (1H, dd, J = 7.9, 1.8 Hz), 7.20 (1H, t, J = 7.9 Hz),
7.38-7.51 (3H, m), 7.55 (1H, s).


[0114]

<Reference Example 11>

3-[2-Chloro-4-(3-chlorophenylthio)phenyl]propan-1-ol

20 [0115]

[Chemical formula 27]



[0116]

The compound of Reference Example 2 was successively reacted according to the same procedures as in Reference Example 7 and then Reference Example 8 to obtain the target product as a colorless oil.

5 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.33 (1H, br s), 1.83-1.95 (2H, m), 2.81-2.85 (2H, m), 3.70 (2H, br s), 7.15-7.23 (5H, m), 7.24-7.29 (1H, m), 7.38 (1H, d, $J = 1.8$ Hz).


[0117]

<Reference Example 12>

10 3-[2-Chloro-4-(3-methylphenoxy)phenyl]propan-1-ol

[0118]

[Chemical formula 28]



[0119]

15 The compound of Reference Example 3 was successively reacted according to the same procedures as in Reference Example 7 and then Reference Example 8 to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.31 (1H, brs), 1.87-1.90 (2H, m), 2.34 (3H, s), 2.80 (2H, t, $J = 7.3$ Hz), 3.70 (2H, dd, $J = 11.6, 6.1$ Hz), 6.79-6.86 (3H, m), 6.94 (1H, d, $J = 7.3$ Hz), 6.99 (1H, d, $J = 2.4$ Hz), 7.18 (1H, d, $J = 7.9$ Hz), 7.22 (1H, t, $J = 7.3$ Hz).

EIMS (+) : 276 $[\text{M}]^+$.


[0120]

<Reference Example 13>

2-Chloro-4-(3-ethylphenylthio)-1-(2-iodoethyl)benzene

[0121]

[Chemical formula 29]



5

[0122]

The compound of Reference Example 6 was reacted according to the same experiment procedures as in Reference Example 327 of the pamphlet of WO 04074297 to obtain the target product as a colorless oil.

10

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.22 (3H, t, $J = 7.3$ Hz), 2.63 (2H, q, $J = 7.3$ Hz), 3.23-3.28 (2H, m), 3.32-3.35 (2H, m), 7.09-7.29 (7H, m).

EIMS (+) : 402 $[\text{M}]^+$.

15


[0123]

<Reference Example 14>

2-Chloro-4-(3-ethylphenylthio)-1-(3-iodopropyl)benzene

[0124]

[Chemical formula 30]



20

[0125]

The compound of Reference Example 8 was reacted according to the same experiment procedures as in Reference Example 164 of the

pamphlet of WO 03029184 to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.22 (3H, t, $J = 7.3$ Hz), 2.12 (2H, quintet, $J = 7.3$ Hz), 2.63 (2H, q, $J = 7.3$ Hz), 2.81 (2H, t, $J = 7.3$ Hz),
5 3.19 (2H, t, $J = 7.3$ Hz), 7.09-7.19 (4H, m), 7.24-7.28 (3H, m).

EIMS (+) : 416 [M] $^+$.


[0126]

<Reference Example 15>

2-Chloro-1-(3-iodopropyl)-4-(3-propylphenoxy)benzene

10 [0127]

[Chemical formula 31]



[0128]

The compound of Reference Example 9 was reacted according to
15 the same experiment procedures as in Reference Example 164 of the pamphlet of WO 03029184 to obtain the target product as a pale yellow oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 0.94 (3H, t, $J = 7.3$ Hz), 1.60-1.68 (2H, m), 2.10-2.17 (2H, m), 2.57 (2H, t, $J = 7.6$ Hz), 2.81 (2H, t, $J = 7.6$ Hz), 3.21 (2H, t, $J = 7.0$ Hz), 6.80-6.85 (3H, m), 6.96 (1H, d, $J = 7.9$ Hz), 6.99 (1H, d, $J = 2.4$ Hz), 7.19 (1H, d, $J = 8.3$ Hz),
20 7.25 (1H, t, $J = 7.9$ Hz).

EIMS(+) : 414 [M] $^+$.


[0129]

<Reference Example 16>

2-Fluoro-1-(3-iodopropyl)-4-(3-trifluoromethylphenylthio)benzene

[0130]

5 [Chemical formula 32]



[0131]

The compound of Reference Example 10 was reacted according to the same experiment procedures as in Reference Example 164 of the pamphlet of WO 03029184 to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 2.13 (2H, quintet, $J = 7.3$ Hz), 2.76 (2H, t, $J = 7.3$ Hz), 3.18 (2H, t, $J = 6.7$ Hz), 7.03 (1H, dd, $J = 10$, 1.8 Hz), 7.09 (1H, dd, $J = 7.9$, 1.8 Hz), 7.20 (1H, t, $J = 7.9$ Hz), 7.39-7.52 (3H, m), 7.57 (1H, s).

EIMS(+) : 404 [M] $^+$.


[0132]

<Reference Example 17>

2-Chloro-4-(3-chlorophenylthio)-1-(3-iodopropyl)benzene

20 [0133]

[Chemical formula 33]



[0134]

The compound of Reference Example 11 was reacted according to the same experiment procedures as in Reference Example 164 of the pamphlet of WO 03029184 to obtain the target product as a colorless oil.

5 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 2.14 (2H, tt, $J = 7.3, 6.7$ Hz), 2.84 (2H, t, $J = 7.3$ Hz), 3.20 (2H, t, $J = 6.7$ Hz), 7.16–7.25 (5H, m), 7.28 (1H, t, $J = 1.8$ Hz), 7.36 (1H, d, $J = 1.8$ Hz).

EIMS (+) : 422 $[\text{M}]^+$.


[0135]

10 <Reference Example 18>

2-Chloro-1-(3-iodopropyl)-4-(3-methylphenoxy)benzene

[0136]

[Chemical formula 34]



15 [0137]

The compound of Reference Example 12 was reacted according to the same experiment procedures as in Reference Example 164 of the pamphlet of WO 03029184 to obtain the target product as a yellow oil.

20 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 2.13 (2H, quint, $J = 7.3$ Hz), 2.34 (3H, s), 2.81 (2H, t, $J = 7.3$ Hz), 3.21 (2H, t, $J = 7.3$ Hz), 6.81–6.84 (3H, m), 6.95 (1H, d, $J = 7.9$ Hz), 6.99 (1H, d, $J = 2.4$ Hz), 7.18 (1H, d, $J = 7.9$ Hz), 7.23 (1H, t, $J = 7.9$ Hz).

EIMS (+) : 386 $[\text{M}]^+$.


[0138]

<Example 1>

(2R,5S)-2-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]propyl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

5 [0139]

[Chemical formula 35]



[0140]

Under an argon atmosphere, a solution of n-butyllithium in
10 hexane (1.54 mol/L, 3.59 mL) was added at -78°C into a solution
of (5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine
(905 mg) in THF (16 mL), and the resultant solution was stirred
at -78°C for 30 minutes. Next, A solution of
2-chloro-1-(3-iodopropyl)-4-(3-trifluoromethylphenoxy)benzene
15 (2.47 g) in THF (4 mL) was added to the reaction mixture, and the
resultant solution was stirred at -78°C for 30 minutes and then
at 0°C for 1 hour. To the reaction solution was added water,
extracted with ethyl acetate, washed with water and saturated brine
in that order, and then dried over anhydrous sodium sulfate. The
20 solvent was evaporated, and the resultant residue was purified by
silica gel column chromatography (hexane : ethyl acetate = 60 :
1) to obtain the target product (1.59 g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.70 (3H, d, J = 6.7 Hz), 1.05 (3H, d,

J = 6.7 Hz), 1.18-1.50 (9H, m), 1.32 (3H, s), 1.86-1.97 (1H, m), 2.21-2.30 (1H, m), 2.65 (2H, t, J = 7.6 Hz), 3.90 (1H, d, J = 2.1 Hz), 3.97-4.21 (4H, m), 6.84 (1H, dd, J = 7.9, 2.4 Hz), 7.00 (1H, d, J = 2.4 Hz), 7.15 (2H, d, J = 7.9 Hz), 7.24 (1H, br s), 7.36 (1H, d, J = 7.9 Hz), 7.44 (1H, t, J = 7.9 Hz).


[0141]

<Example 2>

(2R,5S)-2-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]propyl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0142]

[Chemical formula 36]



[0143]

(5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

and

2-chloro-1-(3-iodopropyl)-4-(3-trifluoromethylphenylthio)benzene were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.63 (3H, d, J = 6.7 Hz), 1.07 (3H, d, J = 6.7 Hz), 1.18-1.29 (10H, m), 1.34-1.66 (2H, m), 1.79-1.91 (1H, m), 2.25-2.33 (1H, m), 2.70 (2H, t, J = 7.6 Hz), 3.85 (1H, br s), 3.99-4.23 (4H, m), 7.16 (2H, d, J = 7.9 Hz), 7.20 (1H, dd, J = 7.9, 1.8 Hz), 7.36-7.42 (3H, m), 7.44-7.50 (1H, m), 7.52 (1H, br s).


[0144]

<Example 3>

(2R,5S)-2-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]ethyl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

5 [0145]

[Chemical formula 37]



[0146]

(5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

10 e and

2-chloro-1-(2-iodoethyl)-4-(3-trifluoromethylphenoxy)benzene were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.72 (3H, d, J = 6.7 Hz), 1.08 (3H, d, J = 6.7 Hz), 1.29 (6H, t, J = 7.3 Hz), 1.36 (3H, s), 1.74-1.82 (1H, m), 2.13-2.20 (1H, m), 2.25-2.32 (1H, m), 2.39-2.56 (2H, m), 3.95 (1H, d, J = 3.1 Hz), 4.02-4.22 (4H, m), 6.83 (1H, dd, J = 8.6, 2.4 Hz), 6.99 (1H, d, J = 2.4 Hz), 7.12-7.15 (2H, m), 7.23 (1H, br s), 7.35 (1H, d, J = 7.8 Hz), 7.44 (1H, t, J = 7.8 Hz).

20 EIMS (+) : 524 [M]⁺.

[0147]


<Example 4>

(2R,5S)-2-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]ethy

1-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0148]

[Chemical formula 38]



5 [0149]

(5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and

2-chloro-1-(2-iodoethyl)-4-(3-trifluoromethylphenylthio)benzene were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.72 (3H, d, J = 6.7 Hz), 1.08 (3H, d, J = 6.7 Hz), 1.28 (6H, t, J = 7.3 Hz), 1.35 (3H, s), 1.68-1.90 (1H, m), 2.10-2.19 (1H, m), 2.38-2.57 (1H, m), 3.95 (1H, d, J = 3.1 Hz), 4.02-4.22 (4H, m), 7.13 (1H, d, J = 7.9 Hz), 7.18 (1H, dd, J = 7.9, 2.4 Hz), 7.35-7.42 (3H, m), 7.43-7.48 (1H, m), 7.54 (1H, br s).


[0150]

<Example 5>

(2R,5S)-2-[2-chloro-4-(3-ethylphenylthio)phenyl]ethyl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

20 [0151]

[Chemical formula 39]



[0152]

(5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and the compound of Reference Example 13 were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

$^1\text{H-NMR}(\text{CDCl}_3, 400 \text{ MHz}): \delta$ 0.72 (3H, d, $J = 6.7 \text{ Hz}$), 1.07 (3H, d, $J = 6.7 \text{ Hz}$), 1.21 (3H, t, $J = 7.3 \text{ Hz}$), 1.28 (3H, t, $J = 7.3 \text{ Hz}$), 1.29 (3H, t, $J = 7.3 \text{ Hz}$), 1.34 (3H, s), 1.70-1.79 (1H, m), 2.09-2.16 (1H, m), 2.24-2.32 (1H, m), 2.35-2.52 (2H, m), 2.61 (2H, q, $J = 7.3 \text{ Hz}$), 3.95 (1H, d, $J = 3.1 \text{ Hz}$), 4.03-4.20 (4H, m), 7.04-7.15 (4H, m), 7.21-7.26 (3H, m).

ESIMS (+) : 501 $[\text{M}+\text{H}]^+$.


[0153]

<Example 6>

(2R,5S)-2-[2-chloro-4-(3-methylphenoxy)phenyl]propyl-3,6-dimethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0154]

[Chemical formula 40]



[0155]

(5S)-3,6-dimethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and the compound of Reference Example 18 were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

5 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 0.68 (3H, d, $J = 6.7$ Hz), 1.07 (3H, d, $J = 6.7$ Hz), 1.33 (3H, s), 1.36-1.43 (1H, m), 1.55-1.62 (1H, m), 1.86-1.92 (1H, m), 2.24-2.26 (1H, m), 2.34 (3H, s), 2.62 (2H, t, $J = 7.9$ Hz), 3.65 (3H, s), 3.66 (3H, s), 3.94 (1H, d, $J = 3.7$ Hz), 6.79-6.82 (3H, m), 6.93 (1H, d, $J = 7.3$ Hz), 6.96 (1H, d, $J = 2.4$
10 Hz), 7.09 (1H, d, $J = 7.9$ Hz), 7.22 (1H, t, $J = 7.9$ Hz).

EIMS (+) : 456 $[\text{M}]^+$.


[0156]

<Example 7>

(2R,5S)-2-[2-chloro-4-(3-ethylphenylthio)phenyl]propyl-3,6-die
15 thoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0157]

[Chemical formula 41]



[0158]

20 (5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and the compound of Reference Example 14 were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.68 (3H, d, J = 6.7 Hz), 1.04 (3H, d, J = 6.7 Hz), 1.20-1.26 (9H, m), 1.31 (3H, s), 1.36-1.43 (1H, m), 1.50-1.57 (1H, m), 1.85-1.92 (1H, m), 2.21-2.28 (1H, m), 2.60-2.65 (4H, m), 3.88 (1H, d, J = 3.7 Hz), 4.00-4.16 (4H, m), 7.06-7.16 (4H, m), 7.22-7.27 (3H, m).

ESIMS (+) : 515 [M+H]⁺.


[0159]

<Example 8>

(2R,5S)-2-[2-chloro-4-(3-chlorophenylthio)phenyl]propyl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0160]

[Chemical formula 42]



[0161]

(5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and the compound of Reference Example 17 were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.69 (3H, d, J = 6.7 Hz), 1.08 (3H, d, J = 6.7 Hz), 1.18-1.29 (7H, m), 1.31 (3H, s), 1.34-1.47 (1H, m), 1.50-1.63 (1H, m), 1.85-1.95 (1H, m), 2.20-2.30 (1H, m), 2.65 (2H, t, J = 7.6 Hz), 3.89 (1H, d, J = 3.1 Hz), 3.99-4.23 (4H, m), 7.11-7.23 (6H, m), 7.35 (1H, d, J = 1.8 Hz).

ESIMS (+) : 521 [M+H]⁺.

[0162]


<Example 9>

(2R,5S)-2-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]prop

5 yl-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine

[0163]

[Chemical formula 43]



[0164]

10 (5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine and the compound of Reference Example 16 were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.67 (3H, d, J = 6.7 Hz), 1.06 (3H, d, J = 6.7 Hz), 1.18-1.29 (7H, m), 1.33 (3H, s), 1.36-1.66 (2H, m), 1.85-1.95 (1H, m), 2.23-2.33 (1H, m), 2.67 (2H, t, J = 7.6 Hz), 3.89 (1H, d, J = 3.1 Hz), 3.99-4.23 (4H, m), 7.02 (1H, dd, J = 9.8 Hz, 1.8 Hz), 7.08 (1H, dd, J = 7.9 Hz, 2.4 Hz), 7.13 (1H, t, J = 7.9 Hz), 7.38-7.50 (3H, m), 7.55 (1H, s).


20 [0165]

<Example 10>

(2S,5S)-2-allyl-2-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]propyl-3,6-diethoxy-5-isopropyl-2,5-dihydropyrazine

[0166]

[Chemical formula 44]



[0167]

5 (5S)-2-allyl-3,6-diethoxy-5-isopropyl-2,5-dihydropyrazine
and

2-chloro-1-(3-iodopropyl)-4-(3-trifluoromethylphenylthio)benzene were reacted in the same manner as in Example 1 to obtain the target product as a colorless oil.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 0.67 (3H, d, J = 6.7 Hz), 1.05 (3H, d, J = 6.7 Hz), 1.23 (3H, t, J = 6.4 Hz), 1.25 (3H, t, J = 6.4 Hz), 1.30-1.64 (3H, m), 1.80-1.90 (1H, m), 2.23-2.39 (2H, m), 2.53 (1H, dd, J = 12.4, 7.3 Hz), 2.65 (2H, t, J = 7.6 Hz), 3.83 (1H, d, J = 3.1 Hz), 4.03-4.18 (4H, m), 4.92-5.04 (2H, m), 5.60-5.73 (1H, m),
15 7.13 (2H, d, J = 7.9 Hz), 7.18 (1H, dd, J = 7.9 Hz, 1.8 Hz), 7.36 (1H, d, J = 1.8 Hz), 7.38-7.42 (2H, m), 7.44-7.49 (1H, m), 7.55 (1H, br s).

[0168]


<Example 11>

20 Ethyl

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylpentanoate

[0169]

[Chemical formula 45]



[0170]

To a solution of the compound of Example 1 (1.59 g) in 1,4-dioxane
5 (60 mL) was added 0.5 mol/L hydrochloric acid (30 mL). The resultant
solution was stirred at room temperature for 1 hour, and then left
to stand at room temperature overnight. The solution was
concentrated, neutralized with saturated aqueous sodium hydrogen
carbonate solution, and extracted with ethyl acetate. The extract
10 was washed with water and saturated brine, and then dried over
anhydrous sodium sulfate. The extract was concentrated, and the
resultant residue was dissolved in acetonitrile (15 mL). To this
solution was added di-tert-butoxydicarbonate (1.55 g), and the
resultant solution was stirred at room temperature for 4 hours and
15 then left to stand at room temperature overnight. To the reaction
solution added water, extracted with ethyl acetate, washed with
water and saturated brine in that order, and then dried over anhydrous
sodium sulfate. The solvent was evaporated, and the resultant
residue was purified by silica gel column chromatography (hexane :
20 ethyl acetate = 9 : 1) to obtain the target product (1.00 g) as
a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.26 (3H, t, $J = 7.3$ Hz), 1.43 (9H, s),
1.53 (3H, s), 1.45–1.68 (2H, m), 1.80–1.90 (1H, m), 2.12–2.30 (1H,
m), 2.69 (2H, t, $J = 7.6$ Hz), 4.16–4.24 (2H, m), 5.33 (1H, br s),

6.85 (1H, dd, J = 7.9 Hz, 2.4 Hz), 7.02 (1H, d, J = 2.4 Hz), 7.15 (1H, dd, J = 7.9 Hz, 2.4 Hz), 7.17 (1H, d, J = 7.9 Hz), 7.24 (1H, br s), 7.37 (1H, d, J = 7.9 Hz), 7.45 (1H, t, J = 7.9 Hz).

[0171]


5 <Example 12>

Ethyl

(R)-2-t-butoxycarbonylamino-5-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentanoate

[0172]

10 [Chemical formula 46]



[0173]

The compound of Example 9 was reacted in the same manner as in Example 11 to obtain the target product as a colorless oil.

15 ¹H-NMR (CDCl₃, 400 MHz): δ 1.26 (3H, t, J = 7.3 Hz), 1.42 (9H, s), 1.51 (3H, s), 1.45-1.68 (2H, m), 1.77-1.86 (1H, m), 2.09-2.20 (1H, m), 2.69 (2H, t, J = 7.6 Hz), 4.13-4.23 (2H, m), 5.29 (1H, br s), 7.02 (1H, dd, J = 9.8 Hz, 1.8 Hz), 7.08 (1H, dd, J = 7.9 Hz, 2.4 Hz), 7.13 (1H, t, J = 7.9 Hz), 7.38-7.50 (3H, m), 7.55 (1H, s).

20 [0174]


<Example 13>

Ethyl

(S)-2-allyl-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]pentanoate

[0175]

[Chemical formula 47]



[0176]

5 The compound of Example 10 was reacted in the same manner as
in Example 11 to obtain the target product as a colorless oil.
¹H-NMR (CDCl₃, 400 MHz) δ 1.24 (3H, t, J = 7.3 Hz), 1.29 - 1.39 (1H,
m), 1.43 (9H, s), 1.60-1.70 (1H, m), 1.78-1.86 (1H, m), 2.32-2.50
(2H, m), 2.66-2.73 (2H, m), 2.99-3.10 (1H, m), 4.19 (2H, q), 5.03
10 (1H, d, J = 3.1 Hz), 5.09 (1H, s), 5.49 (1H, br s), 5.54-5.68 (1H,
m), 7.16 (1H, d, J = 7.9 Hz), 7.19 (1H, dd, J = 7.9, 1.8 Hz), 7.35
(1H, d, J = 1.8 Hz), 7.39-7.44 (2H, m), 7.45-7.50 (1H, m), 7.54
(1H, br s).

[0177]


15 <Example 14>

Ethyl

(R)-2-*t*-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylph
enylthio)phenyl]-2-propylpentanoate

[0178]

20 [Chemical formula 48]



[0179]

To a solution of the compound of Example 13 (400 mg) in ethyl acetate (20 mL) was added palladium, on activated carbon/ethylene diamine complex (100 mg), and the resultant solution was stirred at room temperature for 24 hours under hydrogen atmosphere. The reaction solution was filtered through Celite, and the solvent was evaporated. The resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 30 : 1) to obtain the target product (293 mg) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.91 (3H, t, J = 7.3 Hz), 1.42 (9H, s), 1.15-1.77 (8H, m), 2.72 (2H, t, J = 7.3 Hz), 3.63 (1H, d, J = 12 Hz), 3.67 (1H, d, J = 12 Hz), 4.52 (1H, br s), 7.19-7.22 (2H, m), 7.39 (1H, s), 7.40-7.50 (3H, m), 7.54 (1H, br s).

FABMS (+) : 532 [M+H]⁺.


[0180]

<Example 15>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylpentan-1-ol

[0181]

[Chemical formula 49]



[0182]

To a solution of the compound of Example 11 (1.00 g) in THF (14 mL) was added under ice cooling lithium borohydride (229 mg),

and then ethanol (1.4 mL) was added dropwise. The resultant solution was then stirred for 1 hour under ice cooling. To the reaction solution was added 10% aqueous citric acid, extracted with ethyl acetate, washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1) to obtain the target product (910 mg) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.16 (3H, s), 1.43 (9H, s), 1.53-1.74 (3H, m), 1.81-1.93 (1H, m), 2.73 (2H, t, J = 7.3 Hz), 3.61 (1H, d, J = 12 Hz), 3.65 (1H, d, J = 12 Hz), 4.58 (1H, br s), 4.58 (1H, br s), 6.86 (1H, dd, J = 7.9, 2.4 Hz), 7.03 (1H, d, J = 2.4 Hz), 7.16 (1H, dd, J = 7.9 Hz, 2.4 Hz), 7.21 (1H, d, J = 7.9 Hz), 7.24 (1H, br s), 7.37 (1H, d, J = 7.9 Hz), 7.45 (1H, t, J = 7.9 Hz).


[0183]

<Example 16>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol

[0184]

[Chemical formula 50]



[0185]

The compound of Example 2 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the

same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.14 (3H, s), 1.42 (9H, s), 1.48-1.76 (4H, m), 1.81-1.90 (1H, m), 2.74 (2H, t, J = 6.7 Hz), 3.61 (1H, d, J = 12 Hz), 3.65 (1H, d, J = 12 Hz), 4.56 (1H, br s), 4.58 (1H, br s), 7.20 (2H, d, J = 1.2 Hz), 7.37-7.50 (4H, m), 7.54 (1H, br s).

Optical Rotation: [α]_D²⁷+14.31 (c 0.63, CHCl₃).


[0186]

<Example 17>

(R)-2-t-butoxycarbonylamino-4-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylbutan-1-ol

[0187]

[Chemical formula 51]



[0188]

The compound of Example 3 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.26 (3H, s), 1.45 (9H, s), 1.80-1.88 (1H, m), 2.05-2.12 (1H, m), 2.66-2.80 (2H, m), 3.68 (1H, d, J = 11.6 Hz), 3.73 (1H, d, J = 11.6 Hz), 4.70 (1H, br s), 6.86 (1H,

dd, $J = 8.5, 2.5$ Hz), 7.03 (1H, d, $J = 2.5$ Hz), 7.13-7.16 (1H, m), 7.22-7.24 (2H, m), 7.37 (1H, d, $J = 7.9$ Hz), 7.45 (1H, t, $J = 7.9$ Hz).

FABMS (+) : 474 [M+H]⁺.


5 [0189]

<Example 18>

(R)-2-*t*-butoxycarbonylamino-4-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylbutan-1-ol

[0190]

10 [Chemical formula 52]



[0191]

The compound of Example 4 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.25 (3H, s), 1.44 (9H, s), 1.79-1.89 (1H, m), 2.05-2.13 (1H, m), 2.66-2.83 (2H, m), 3.68 (1H, d, $J = 12$ Hz), 3.71 (1H, d, $J = 12$ Hz), 4.69 (1H, br s), 7.20-7.23 (2H, m), 7.37-7.42 (3H, m), 7.45-7.50 (2H, m), 7.55 (1H, br s).

[0192]


<Example 19>

(R)-2-*t*-butoxycarbonylamino-4-[2-chloro-4-(3-ethylphenylthio)p

henyl]-2-methylbutan-1-ol

[0193]

[Chemical formula 53]



5 [0194]

The compound of Example 5 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 1.22 (3H, t, J = 7.3 Hz), 1.24 (3H, s), 1.44 (9H, s), 1.77-1.85 (1H, m), 2.02-2.09 (1H, m), 2.62 (2H, q, J = 7.3 Hz), 2.63-2.78 (2H, m), 3.64-3.73 (2H, m), 4.08 (1H, br), 4.68 (1H, br s), 7.10-7.17 (4H, m), 7.22-7.28 (3H, m).

ESIMS (+) : 450 [M+H]⁺.


15 [0195]

<Example 20>

(R)-2-tert-butoxycarbonylamino-5-[2-chloro-4-(3-methylphenoxy)phenyl]-2-methylpentan-1-ol

[0196]

20 [Chemical formula 54]



[0197]

The compound of Example 6 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

5 ¹H-NMR (CDCl₃, 400 MHz): δ 1.15 (3H, s), 1.43 (9H, s), 1.61-1.67 (3H, m), 1.83-1.87 (1H, m), 2.34 (3H, s), 2.70 (2H, t, J = 7.0 Hz), 3.62-3.65 (2H, m), 4.57 (1H, s), 6.81-6.84 (3H, m), 6.94 (1H, d, J = 7.3 Hz), 6.98 (1H, d, J = 3.1 Hz), 7.15 (1H, d, J = 7.9 Hz), 7.22 (1H, t, J = 7.9 Hz).

10 ESIMS (+) : 434 [M+H]⁺.


[0198]

<Example 21>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-ethylphenylthio)phenyl]-2-methylpentan-1-ol

15 [0199]

[Chemical formula 55]



[0200]

20 The compound of Example 7 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.14 (3H, s), 1.22 (3H, t, J = 7.3 Hz), 1.43 (9H, s), 1.54-1.70 (3H, m), 1.79-1.89 (1H, m), 2.62 (2H, q,

J = 7.3 Hz), 2.70 (2H, t, J = 7.0 Hz), 3.57-3.66 (2H, m), 4.05 (1H, br), 4.55 (1H, br s), 7.10-7.17 (4H, m), 7.17-7.28 (3H, m).

ESIMS (+) : 464 [M+H]⁺.


[0201]

5 <Example 22>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-propylphenoxy)phenyl]-2-methylpentan-1-ol

[0202]

[Chemical formula 56]



[0203]

The compound of Reference Example 15 and (5S)-3,6-diethoxy-5-isopropyl-2-methyl-2,5-dihydropyrazine were reacted with in the same manner as in Example 1. The resultant compound was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.94 (3H, t, J = 7.3 Hz), 1.15 (3H, s), 1.24-1.28 (2H, m), 1.43 (9H, s), 1.60-1.69 (3H, m), 1.80-1.90 (1H, m), 2.57 (2H, t, J = 7.6 Hz), 2.70 (2H, t, J = 7.6 Hz), 3.58-3.67 (2H, m), 4.11 (1H, br s), 4.58 (1H, br s), 6.79-6.85 (3H, m), 6.95 (1H, d, J = 7.9 Hz), 6.99 (1H, d, J = 2.8 Hz), 7.15 (1H, d, J = 8.3 Hz), 7.24 (1H, t, J = 7.9 Hz).


[0204]

<Example 23>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-chlorophenylthio)phenyl]-2-methylpentan-1-ol

[0205]

5 [Chemical formula 57]



[0206]

The compound of Example 8 was reacted in the same manner as in Example 11 to obtain an ester, which was then reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.14 (3H, s), 1.43 (9H, s), 1.58-1.74 (3H, m), 1.79-1.92 (1H, m), 2.73 (2H, t, $J = 6.7$ Hz), 3.61 (1H, d, $J = 12$ Hz), 3.64 (1H, d, $J = 12$ Hz), 4.08 (1H, br s), 4.57 (1H, br s), 7.17-7.27 (6H, m), 7.37 (1H, s).

ESIMS (+) : 470 $[\text{M}+\text{H}]^+$.


[0207]

<Example 24>

(R)-2-t-butoxycarbonylamino-5-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol

[0208]

[Chemical formula 58]



[0209]

The compound of Example 12 was reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.14 (3H, s), 1.42 (9H, s), 1.55-1.74
5 (3H, m), 1.75-1.85 (1H, m), 2.65 (2H, t, J = 6.7 Hz), 3.58-3.64
(2H, m), 4.03 (1H, br s), 4.55 (1H, br s), 7.04 (1H, dd, J = 9.8
Hz, 1.8 Hz), 7.10 (1H, dd, J = 7.9 Hz, 1.8 Hz), 7.17 (1H, t, J =
7.9 Hz), 7.38-7.50 (3H, m), 7.54 (1H, br s).


[0210]

10 <Example 25>

(R)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylph
enylthio)phenyl]-2-propylpentan-1-ol

[0211]

[Chemical formula 59]



15

[0212]

The compound of Example 14 was reacted in the same manner as in Example 15 to obtain the target product as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 0.92 (3H, t, J = 7.3 Hz), 1.42 (9H, s),
20 1.14-1.80 (8H, m), 2.72 (2H, t, J = 7.3 Hz), 3.62 (1H, d, J = 12
Hz), 3.66 (1H, d, J = 12 Hz), 4.54 (1H, br s), 7.16-7.22 (2H, m),
7.39 (1H, s), 7.40-7.48 (3H, m), 7.55 (1H, br s).

FABMS (+) : 532 [M+H]⁺.


[0213]

<Example 26>

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylpentan-1-ol hydrochloride

5 [0214]

[Chemical formula 60]



[0215]

To the compound of Example 15 (6.50 g) was added a 10 w/w%
10 hydrogen chloride solution in methanol (methanol containing hydrogen
chloride, 67 mL), and the resultant mixture was stirred for 1 hour
at room temperature, and then left overnight at room temperature.
The solvent was then evaporated to obtain the target product (5.15
g) as a colorless amorphous.

15 $^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): δ 1.07 (3H, s), 1.46-1.64 (4H, m), 2.62-2.72
(2H, m), 3.31-3.36 (2H, m), 7.03 (1H, dd, $J = 7.9, 2.4$ Hz), 7.20
(1H, d, $J = 2.4$ Hz), 7.30 (1H, d, $J = 7.9$ Hz), 7.34 (1H, s), 7.39
(1H, d, $J = 7.9$ Hz), 7.52 (1H, d, $J = 7.9$ Hz), 7.63 (1H, t, $J =$
7.9 Hz).

20 HREIMS (+) : 388.1281 (Calcd. for $\text{C}_{19}\text{H}_{21}\text{NClF}_3\text{O}_2$: 388.1291).

Optical Rotation: $[\alpha]_D^{23} -2.74$ (c 0.63, CHCl_3).

[0216]


<Example 27>

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]

-2-methylpentan-1-ol hydrochloride

[0217]

[Chemical formula 61]



5 [0218]

The compound of Example 16 was reacted in the same manner as in Example 26 to obtain the target product as a white powder.

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): δ 1.09 (3H, s), 1.49–1.63 (4H, m), 2.65–2.71 (2H, br s), 3.34 (1H, d, $J = 12$ Hz), 3.38 (1H, d, $J = 12$ Hz), 7.34 (1H, dd, $J = 7.9$ Hz, 2.4 Hz), 7.41 (1H, d, $J = 7.9$ Hz), 7.49 (1H, d, $J = 2.4$ Hz), 7.55 (1H, d, $J = 7.9$ Hz), 7.61 (1H, d, $J = 2.4$ Hz), 7.67 (1H, d, $J = 7.9$ Hz), 7.53–7.74 (3H, br s).

ESIMS (+) : 404 $[\text{M}+\text{H}]^+$.

Elemental Analysis: Measured: C 51.65%, H 4.86%, N 2.86%, Calcd. for $\text{C}_{19}\text{H}_{21}\text{ClF}_3\text{NOS}\cdot\text{HCl}$: C 51.82%, H 5.04%, N 3.18%.

Optical Rotation: $[\alpha]_D^{23}$ -3.45 (c 1.00, CHCl_3).


[0219]

<Example 28>

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethoxyphenoxy)phenyl]-2-methylbutan-1-ol hydrochloride

[0220]

[Chemical formula 62]



[0221]

The compound of Example 17 was reacted in the same manner as in Example 26 to obtain the target product as a white powder.

5 $^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz): δ 1.24 (3H, s), 1.70 – 1.80 (2H, m), 2.71 (2H, t, $J = 8.6$ Hz), 3.44 (1H, dd, $J = 11$ Hz, 4.9 Hz), 3.50 (1H, dd, $J = 11$ Hz, 4.9 Hz), 5.54 (1H, t, $J = 4.9$ Hz), 7.04 (1H, dd, $J = 8.6$, 2.4 Hz), 7.21 (1H, d, $J = 2.4$ Hz), 7.31 (1H, dd, $J = 8.6$, 2.4 Hz), 7.35 (1H, br s), 7.41 (1H, d, $J = 8.6$ Hz), 7.52 (1H, d, $J = 7.9$ Hz), 7.63 (1H, t, $J = 7.9$ Hz), 7.95 (3H, br s).

FABMS (+) : 374 $[\text{M}+\text{H}]^+$.

Elemental Analysis: Measured: C 52.38%, H 4.80%, N 3.42%, Calcd. for $\text{C}_{18}\text{H}_{19}\text{ClF}_3\text{NO}_2 \cdot \text{HCl}$: C 52.70%, H 4.91%, N 3.41%.


[0222]

15 <Example 29>

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylbutan-1-ol hydrochloride

[0223]

[Chemical formula 63]



20

[0224]

The compound of Example 18 was reacted in the same manner as in Example 26 to obtain the target product as a white powder.

$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz): δ 1.22 (3H, s), 1.66-1.83 (2H, m), 2.72 (2H, t, $J = 8.6$ Hz), 3.42 (1H, dd, $J = 11.0, 7.9$ Hz), 3.49 (1H, dd, $J = 11.0, 7.9$ Hz), 5.54 (1H, t, $J = 4.9$ Hz), 7.36 (1H, dd, $J = 7.9, 1.8$ Hz), 7.42 (1H, d, $J = 7.9$ Hz), 7.50 (1H, d, $J = 1.8$ Hz), 7.53-7.64 (3H, m), 7.67 (1H, d, $J = 7.9$ Hz), 7.82 (3H, br s).
FABMS (+) : 390 $[\text{M}+\text{H}]^+$.

Elemental Analysis: Measured: C 50.47%, H 4.65%, N 3.36%, Calcd.

for $\text{C}_{18}\text{H}_{19}\text{ClF}_3\text{NOS} \cdot \text{HCl}$: C 50.71%, H 4.73%, N 3.29%.

Optical Rotation: $[\alpha]_{\text{D}}^{27} +5.78$ (c 0.33, CHCl_3).


[0225]

<Example 30>

(R)-2-amino-4-[2-chloro-4-(3-ethylphenylthio)phenyl]-2-methylbutan-1-ol hydrochloride

[0226]

[Chemical formula 64]



[0227]

The compound of Example 19 was reacted in the same manner as in Example 26 to obtain the target product as a white powder.

$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz): δ 1.14 (3H, t, $J = 7.3$ Hz), 1.22 (3H, s), 1.67-1.81 (2H, m), 2.59 (2H, q, $J = 7.3$ Hz), 2.69 (2H, t, $J = 8.6$

Hz), 3.42 (1H, dd, J = 11.6, 5.5 Hz), 3.48 (1H, dd, J = 11.6, 5.5 Hz), 5.52 (1H, t, J = 4.9 Hz), 7.16-7.22 (2H, m), 7.26-7.27 (2H, m), 7.30-7.35 (2H, m), 7.93 (3H, br s).

0 ESIMS (+) : 350 [M+H]⁺.

5 Elemental Analysis: Measured: C 58.90%, H 6.42%, N 3.59%, Calcd. for C₁₉H₂₄ClNOS. HCl: C 59.06%, H 6.52%, N 3.63%.


[0228]

<Example 31>

(R)-2-amino-5-[2-chloro-4-(3-methylphenoxy)phenyl]-2-methylpen
10 tan-1-ol hydrochloride

[0229]

[Chemical formula 65]



[0230]

15 The compound of Example 20 was reacted in the same manner as in Example 26 to obtain the target product as a colorless amorphous .

¹H-NMR (DMSO-d₆, 400 MHz): δ 1.11 (3H, s), 1.57 (4H, brs), 2.29 (3H, s), 2.64 (2H, brs), 3.35-3.39 (2H, m), 5.45 (1H, t, J = 4.9 Hz), 6.81 (1H, dd, J = 8.6, 2.4 Hz), 6.85 (1H, s), 6.92 (1H, dd, J =
20 8.6, 2.4 Hz), 6.99 (1H, d, J = 8.6 Hz), 7.03 (1H, d, J = 2.4 Hz), 7.28 (1H, t, J = 8.6 Hz), 7.34 (1H, d, J = 8.6 Hz), 7.77 (3H, brs).

HRESIMS (+) : 334.15655 (Calcd. for C₁₉H₂₅ClNO₂: 334.15738).

Optical Rotation: [α]_D^{26.7}-5.75 (c 0.60, CHCl₃).


[0231]

<Example 32>

(R)-2-amino-5-[2-chloro-4-(3-ethylphenylthio)phenyl]-2-methylpentan-1-ol hydrochloride

[0232]

5 [Chemical formula 66]



[0233]

The compound of Example 21 was reacted in the same manner as in Example 26 to obtain the target product as a colorless oil.

10 ¹H-NMR (DMSO-d₆, 400 MHz): δ 1.10 (3H, s), 1.15 (3H, t, J = 7.3 Hz), 1.52-1.58 (4H, m), 2.59 (2H, q, J = 7.3 Hz), 2.62-2.66 (2H, m), 3.32-3.39 (2H, m), 5.43 (1H, br), 7.15-7.22 (3H, m), 7.26 (2H, d, J = 1.8 Hz), 7.32 (2H, dd, J = 7.3, 1.8 Hz), 7.81 (3H, br s).
HRESIMS (+) : 364.15051 (Calcd. for C₂₀H₂₇ClNOS: 364.15019).


15 [0234]

<Example 33>

(R)-2-amino-5-[2-chloro-4-(3-propylphenoxy)phenyl]-2-methylpentan-1-ol hydrochloride

[0235]

20 [Chemical formula 67]



[0236]

The compound of Example 22 was reacted in the same manner as

in Example 26 to obtain the target product as a colorless amorphous .

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): δ 0.86 (3H, t, J = 7.3 Hz), 1.11 (3H, s), 1.51-1.61 (6H, m), 2.53 (2H, t, J = 7.3 Hz), 2.63 (2H, t, J = 6.7 Hz), 3.34-3.42 (2H, m), 5.45 (1H, t, J = 4.9 Hz), 6.81 (1H, ddd, J = 7.9, 1.8, 0.9 Hz), 6.87 (1H, t, J = 1.8 Hz), 6.91 (1H, dd, J = 8.6, 2.4 Hz), 7.00 (1H, d, J = 7.9 Hz), 7.02 (1H, d, J = 2.4 Hz), 7.30 (1H, t, J = 7.9 Hz), 7.34 (1H, d, J = 8.6 Hz), 7.85 (3H, br s).

ESIMS(+) : 362 [M+H] $^+$.

10 HRESIMS(+) : 362.19198 (Calcd. for $\text{C}_{21}\text{H}_{29}\text{ClNO}_2$: 362.18868).

Optical Rotation: $[\alpha]_D^{25.1}$ -4.46 (c 1.27, CHCl_3).

[0237]


<Example 34>

(R)-2-amino-5-[2-chloro-4-(3-chlorophenylthio)phenyl]-2-methyl

15 pentan-1-ol hydrochloride

[0238]

[Chemical formula 68]



[0239]

20 The compound of Example 23 was reacted in the same manner as in Example 26 to obtain the target product as a colorless amorphous .

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): δ 1.10 (3H, s), 1.49-1.64 (4H, m), 2.68 (2H, br s), 3.33 (1H, dd, J = 12, 4.9 Hz), 3.38 (1H, dd, J = 12, 4.9 Hz), 5.45 (1H, t, J = 4.9 Hz), 7.26 (1H, dt, J = 7.3, 1.8 Hz),

7.30-7.43 (5H, m), 7.45 (1H, d, J = 1.8 Hz), 7.77 (3H, br s).

HREIMS (+) : 370.0799 (Calcd. for C₁₈H₂₁Cl₂NOS: 370.0799).

Optical Rotation: $[\alpha]_D^{27}$ -3.81 (c 0.50, CHCl₃).


[0240]

5 <Example 35>

(R)-2-amino-5-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]
-2-methylpentan-1-ol hydrochloride

[0241]

[Chemical formula 69]



10

[0242]

The compound of Example 24 was reacted in the same manner as
in Example 26 to obtain the target product as a colorless amorphous .

¹H-NMR (DMSO-d₆, 400 MHz): δ 1.09 (3H, s), 1.48-1.61 (4H, m), 2.57-2.64
(2H, br s), 3.32 (1H, dd, J = 11, 4.9 Hz), 3.37 (1H, dd, J = 11,
4.9 Hz), 5.44 (1H, t, J = 4.9 Hz), 7.20 (1H, dd, J = 7.9, 1.8 Hz),
7.26 (1H, dd, J = 9.8, 1.8 Hz), 7.37 (1H, t, J = 7.9 Hz), 7.54-7.68
(4H, m), 7.74 (3H, br s).

HRESIMS (+) : 388.1345 (Calcd. for C₁₉H₂₂F₄NOS: 388.1358).

Optical Rotation: $[\alpha]_D^{24}$ -3.23 (c 0.69, CHCl₃).


[0243]

<Example 36>

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]
-2-propylpentan-1-ol hydrochloride

[0244]

[Chemical formula 70]



[0245]

5 The compound of Example 25 was reacted in the same manner as in Example 26 to obtain the target product as a white powder.

$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz): δ 0.84 (3H, t, $J = 7.3$ Hz), 1.20 (2H, q, $J = 7.3$ Hz), 1.36-1.63 (6H, m), 2.68 (2H, t, $J = 7.3$ Hz), 3.36 (2H, d, $J = 4.9$ Hz), 5.40 (1H, d, $J = 4.9$ Hz), 7.35 (1H, dd, $J =$
10 7.9 Hz, 1.8 Hz), 7.42 (1H, d, $J = 7.9$ Hz), 7.50 (1H, d, $J = 1.8$ Hz), 7.55 (1H, d, $J = 7.9$ Hz), 7.58-7.63 (2H, m), 7.67 (1H, d, $J = 7.9$ Hz), 7.69 (3H, br s).

FABMS (+) : 432 $[\text{M}+\text{H}]^+$.

Elemental Analysis: Measured: C 53.46%, H 5.62%, N 2.98%, Calcd.
15 for $\text{C}_{21}\text{H}_{25}\text{ClF}_3\text{NOS}\cdot\text{HCl}$: C 53.85%, H 5.59%, N 2.99%.

Optical Rotation: $[\alpha]_{\text{D}}^{23} +3.85$ (c 0.63, CHCl_3).


[0246]

<Example 37>

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]
20 -2-methylpentan-1-ol

[0247]

[Chemical formula 71]



[0248]

To a solution of the compound of Example 27 (9.3 g) in ethyl acetate (450 mL) was added saturated aqueous sodium hydrogen carbonate solution (450 mL), and the resultant solution was stirred at room temperature for 10 minutes. The organic layer was washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by NH-silica gel column chromatography (ethyl acetate : methanol = 4 : 1) to obtain the target product (8.9 g) as a white powder.

$^1\text{H-NMR}$ (DMSO- d_6 , 400 MHz): δ 0.85 (3H, s), 1.21 (2H, br s), 1.28 (2H, t, $J = 8.6$ Hz), 1.46-1.67 (2H, m), 2.65 (2H, t, $J = 8.6$ Hz), 3.06 (2H, br s), 4.49 (1H, br s), 7.32 (1H, dd, $J = 7.9, 1.8$ Hz), 7.40 (1H, d, $J = 9.8$ Hz), 7.47 (1H, d, $J = 1.8$ Hz), 7.54 (1H, dd, $J = 6.7, 1.8$ Hz), 7.56-7.62 (2H, m), 7.65 (1H, dd, $J = 6.7, 1.8$ Hz).

ESIMS (+) : 404 $[\text{M}+\text{H}]^+$.

Elemental Analysis: Measured: C 56.26%, H 5.14%, N 3.40%, Calcd. for $\text{C}_{19}\text{H}_{21}\text{ClF}_3\text{NOS}$: C 56.50%, H 5.24%, N 3.47%.

[0249]

<Example 38>


Diethyl

2-{3-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]propyl}-2

-methylmalonate

[0250]

[Chemical formula 72]



5 [0251]

2-Chloro-1-(3-iodopropyl)-4-(3-trifluoromethylphenylthio) benzene and diethyl 2-methylmalonate were reacted according to the same procedures as in Example 152 of WO 04026817 to obtain the target product as a colorless oil.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 1.25 (6H, t, J = 7.4 Hz), 1.40 (3H, s), 1.51-1.63 (2H, m), 1.90-1.97 (2H, m), 2.73 (2H, t, J = 7.9 Hz), 4.17 (4H, q, J = 7.4 Hz), 7.17-7.23 (2H, m), 7.38 (1H, d, J = 2.2 Hz), 7.39-7.44 (2H, m), 7.45-7.50 (1H, m), 7.55 (1H, s).

EIMS (+) : 502 [M]⁺.


15 [0252]

<Example 39>

(±)-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-ethoxycarbonyl-2-methylpentanoic acid

[0253]

20 [Chemical formula 73]



[0254]

To a solution of the compound of Example 38 (16.8 g) in ethanol

(167 mL) was added potassium hydroxide (2.40 g), and the resultant solution was stirred at 50°C for 24 hours. To the reaction solution was added water, neutralized with 2 mol/L aqueous hydrochloric acid, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1) to obtain the target product (11.2 g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.26 (3H, t, J = 7.4 Hz), 1.47 (3H, s), 1.55-1.66 (2H, m), 1.87-2.06 (2H, m), 2.73 (2H, t, J = 7.9 Hz), 4.22 (2H, q, J = 7.4 Hz), 7.18 (1H, d, J = 7.9 Hz), 7.20 (1H, dd, J = 7.9, 1.8 Hz), 7.38 (1H, d, J = 1.8 Hz), 7.39-7.44 (2H, m), 7.45-7.50 (1H, m), 7.54 (1H, s).

ESIMS (+) : 475 [M+H]⁺.

[0255]


<Example 40>

Ethyl

(±)-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methoxycarbonylamino-2-methylpentanoate

[0256]

[Chemical formula 74]



[0257]

To a solution of the compound of Example 39 (15.8 g) in benzene (166 mL) was added diphenylphosphoryl azide (7.86 mL) and triethylamine (6.01 mL), and the resultant solution was heated to reflux for 1.5 hours. The temperature of the reaction solution was returned to room temperature, and methanol (20 mL) was added dropwise over 20 minutes. The resultant solution was heated to reflux for 30 minutes, and then further sodium methoxide (3.58 g) was added. The resultant solution was heated to reflux for 1.5 hours. To the reaction solution was added saturated aqueous ammonium chloride, and extracted with ethyl acetate. The organic layer was washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 5 : 1) to obtain the target product (15.6 g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.25 (3H, t, J = 7.3 Hz), 1.32-1.47 (1H, m), 1.52-1.67 (1H, m), 1.57 (3H, s), 1.80-1.90 (1H, m), 2.20-2.37 (1H, m), 2.62-2.76 (2H, m), 3.64 (3H, s), 4.15-4.25 (2H, m), 5.62 (1H, br s), 7.16 (1H, d, J = 7.9 Hz), 7.20 (1H, dd, J = 7.9, 1.8 Hz), 7.38 (1H, d, J = 1.8 Hz), 7.40-7.44 (2H, m), 7.45-7.50 (1H, m), 7.55 (1H, s).

ESIMS (+) : 504 [M+H]⁺.

[0258]


<Example 41>

(±)-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-metho

xycarbonylamino-2-methylpentan-1-ol

[0259]

[Chemical formula 75]



5 [0260]

To a solution of the compound of Example 40 (15.6 g) in THF (249 mL) was added under ice cooling lithium borohydride (3.75 g), and then ethanol (16.6 mL) was added dropwise . The resultant solution was then stirred for 1 hour under ice cooling. To The reaction solution was added 10% aqueous citric acid, extracted with ethyl acetate, washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1) to obtain the target product (12.9 g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.18 (3H, s), 1.54-1.74 (3H, m), 1.78-1.89 (1H, m), 2.73 (2H, t, J = 7.9 Hz), 3.63 (3H, s), 3.56-3.70 (2H, m), 4.23 (1H, br s), 7.17-7.22 (2H, m), 7.38-7.50 (4H, m), 7.54 (1H, s).

20 ESIMS (+) : 462 [M+H]⁺.


[0261]

<Example 42>

(±)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol

[0262]

[Chemical formula 76]



[0263]

5 To a solution of the compound of Example 41 (12.9 g) in THF (60 mL) and methanol (120 mL) was added under ice cooling 5 mol/L aqueous potassium hydroxide solution (60 mL), and the resultant solution was heated to reflux for 86 hours. To the reaction solution was added water, extracted with ethyl acetate, washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The extract was concentrated, the residue was dissolved in 1,4-dioxane (279 mL), and the resultant solution was charged with di-tert-butoxydicarbonate (9.13 g). The solution was stirred at room temperature for 2 hours and then left to stand at room temperature overnight. The reaction solution was added water, extracted with ethyl acetate, washed with water and saturated brine in that order, and then dried over anhydrous sodium sulfate. The solvent was evaporated, and the resultant residue was purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1) to obtain the target product (13.0 g) as a colorless oil.

¹H-NMR (CDCl₃, 400 MHz): δ 1.14 (3H, s), 1.42 (9H, s), 1.53-1.74 (3H, m), 1.79-1.92 (1H, m), 2.74 (2H, t, J = 7.9 Hz), 3.58-3.69 (2H, m), 4.05 (1H, br s), 4.57 (1H, br s), 7.20-7.22 (2H, m), 7.38-7.50 (4H, m), 7.54 (1H, s).

ESIMS (+) : 504 [M+H]⁺.

[0264]

<Examples 43 and 44>

(+)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylph
5 enylthio)phenyl]-2-methylpentan-1-ol and

(-)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylph
enylthio)phenyl]-2-methylpentan-1-ol

The compound of Example 42 was subjected to optical resolution
using high performance liquid chromatography (CHIRALCEL OJ-H,
10 hexane : isopropanol : diethylamine = 98 : 2 : 0.1 (v/v), measurement
wavelength: UV 278 nm, flow rate: 1.0 mL/min). From the pre-elution
portion, an $[\alpha]_D^{25} +15.08$ (c 0.63, CHCl₃) colorless oil was obtained
(Example 43), and from the post-elution portion, an $[\alpha]_D^{26} -13.91$
(c 0.63, CHCl₃) colorless oil was obtained (Example 44).

15 [0265]

<Example 45>

(-)-2-t-butoxycarbonylamino-5-[2-chloro-4-(3-trifluoromethylph
enylthio)phenyl]-2-methylpentan-1-ol hydrochloride

The compound of Example 43 was reacted in the same manner as
20 in Example 26 to obtain the target product as a white powder.

ESIMS (+) : 404 [M+H]⁺.

Optical Rotation: $[\alpha]_D^{25} -4.48$ (c 1.00, CHCl₃).

Next, results which support the utility of the compound of
the present invention will be shown by experiment examples.

25 [0266]

<Experiment example 1> Activity On Total Number Of White Blood Cells
in Mouse Peripheral Blood

The test compounds were dissolved or suspended in DMSO, and then a physiological saline solution was added to make dosing solutions (DMSO final concentration 1%). If the dosing solutions were not in a dissolved or a uniform suspended state, Tween 80 was added so that the final concentration thereof was 0.01 to 0.1%. The resultant solutions were then sonicated to form a fine suspension. These test compound solutions were intraperitoneally administered in respective amounts of 0.1 mL or 0.2 mL per 10 g of body weight to BALB/cCrSlc male mice (8 to 14 weeks old, Japan SLC). For the control group, only a solvent having the same composition as that used in the preparation of the test compounds was administered in the same manner. Six hours after administration of the test compound solutions, blood was collected from the postcava under ether anesthesia (10 μ L of 10% EDTA was added as an anticoagulant). Measurement of the total number of white blood cells was carried out using the automated blood cell counting apparatus Sysmex F-820 (Sysmex).

[0267]

Here, compounds having an inhibition ratio of 60% or more were marked as “+++”, compounds having an inhibition ratio of less than 60% but more than or equal to 50% were marked as “++”, and compounds having an inhibition ratio of less than 50% but more than or equal to 40% were marked as “+”. Furthermore, the inhibition ratios are

the results from a dose of 0.3 mg/kg (“[]” are the results from 3 mg/kg). The results are shown in Table 1.

[0268]

[Table 1]

Table 1

Example No.	Inhibition Ratio
26	[+]
27	+
28	++
29	+++
36	[++]

[3 mg/kg]

5

[0269]

<Experiment example 2> Inhibition Activity of the Test Compounds
On Mouse Host-Versus-Graft Rejection Reaction>

Experiment example 2 was carried out by referring to the methods
described in Transplantation, Vol. 55, No. 3, pp. 578 to 591, 1993.
The spleens from BALB/c male mice which were 6 to 23 weeks old (Japan
SLC) were collected. The spleens were extracted in RPMI-1640 medium
(Sigma), and then turned into a spleen cell suspension by grinding
them with 2 glass slides and then passing the ground matter through
a strainer (70 microns, Falcon). The spleen cell suspensions were
centrifuged, and the supernatants were removed. Then, an ammonium
chloride-tris isotonic buffer solution was added, and the red blood
cells were hemolyzed. After centrifuging and washing 3 times with
RPMI-1640 medium, the cells were suspended in RPMI-1640 medium.
Mitomycin C (Kyowa Hakko) was added thereto so that the final

concentration was 25 µg/mL, and the cells were incubated for 30 minutes at 37°C under 5% CO₂. After centrifuging and washing 3 times with RPMI-1640 medium, the cells were suspended in RPMI-1640 medium so that the concentration was 2.5×10^8 cells/mL. These cells were used as a stimulator cell suspension. 20 µL (5×10^6 cells/mouse) of the stimulator cell suspension was injected subcutaneously into the right hind-footpad of C3H/HeN male mice which were 6 to 11 weeks old (Clea Japan) using a 27G needle and a microsyringe (Hamilton). For the normal control group, RPMI-1640 medium only was injected. After 4 days, the right popliteal lymph nodes were extracted, and their weight was measured using Mettler AT201 electronic scales (Mettler Toledo). From the day of the stimulator cell suspensions being injected to 3 days thereafter, the test compounds were intraperitoneally administered on consecutive days once per day, for a total of 4 times. For the control group, only a solvent having the same composition as that used in the preparation of the test compounds was administered in the same manner. The results are shown in Table 2. As the solvent, basically a physiological saline containing 0.5% DMSO was used (suitably adjusted according to the solubility of the compound).

The inhibition ratio was calculated using the following equation.

[0270]

[Equation 1]

Inhibition ratio =

$$\frac{(\text{Right Popliteal Lymph Node's Weight of the Positive Control Group}) - (\text{Right Popliteal Lymph Node's Weight of the Test Compound Group}) \times 100}{(\text{Right Popliteal Lymph Node's Weight of the Positive Control Group}) - (\text{Right Popliteal Lymph Node's Weight of the Normal Control Group})}$$

[0271]

Compounds having an ED₅₀ value of less than 0.1 mg/kg were marked as "+++", compounds having an ED₅₀ value of less than 1 mg/kg but more than or equal to 0.1 mg/kg were marked as "++", and compounds having an ED₅₀ value of less than 10 mg/kg but more than or equal to 1 mg/kg were marked as "+".

[0272]

[Table 2]

Table 2

Example No.	ED ₅₀
26	++
27	+++
28	+
29	++
32	++
35	++
36	++

[0273]

<Experiment example 3> Activity of the Test Compounds On Rat Skin Graft Model>

Using a rat skin graft model, activity on the rejection reaction was investigated. An allogenic skin graft for rats having a matched major histocompatibility complex (MHC) was carried out by referring to the methods described in a publication (Am. J. Med. Technol.; 36, 149-157, 1970, Transplant. Proc.; 28, 1056 to 1059, 1996) etc.

[0274]

Skin from the abdomen of a LEW/CrlCrlj rat serving as the donor (male, 5 weeks old, Japan Charles River) was peeled away, and the panniculus carnosus muscle layer under the skin was removed. Then, a 1.8 cm × 1.8 cm skin specimen (graft) was prepared. Next, a F344/DuCrlCrlj rat serving as the recipient (male, 5 weeks old, Japan Charles River) was anesthetized by intraperitoneal administration of 40 mg/kg of sodium pentobarbital (nembutal injection, Dainippon Sumitomo Pharma). The hair of the back of the rat was shorn off by electric clippers, and then the remaining hair was further shaved off using a razor. Then, a 1.8 cm × 1.8 cm mark was made on the back of the rat, and a shallow cut was made along the mark with a scalpel. Next, the skin was peeled away along this cut by tweezers to produce a graft bed. Several drops of sterile physiological saline solution (Otsuka normal saline, Otsuka Pharmaceutical) were dropped onto the graft bed, and then wiped away with gauze. Several drops of penicillin (crystalline penicillin G potassium Meiji, Meiji Seika) were then dropped thereon. Next, the graft was placed on the graft bed. After absorbing excess moisture with sterile gauze, a first-aid bandaid was stuck over the graft. Furthermore, an adhesive bandage was wrapped around the trunk of the rat. On the third day after the operation, a new adhesive bandage was wrapped thereover. On the fifth day after the operation, the bandaid and the bandages were cut open with scissors and removed. Observation of the graft was carried out on consecutive days from

when the bandaid was removed (from the fifth day). A determination of graft rejection was made when 90% or more of the graft had died and turned brown. The number of days from grafting until rejection was confirmed was taken as the survival period. The mean value of the survival period for each group was calculated as the mean survival time (MST). Investigation was carried out with 5 animals per group. All of the rats were housed in individual cages. The test compounds were dissolved in ultra-pure water, and were orally administered once a day, on consecutive days, from the graft day. For the control group, ultra-pure water was administered in the same manner.

[0275]

The results are shown in Table 3. The control group MST was 8.8 days. Those grafts of animals to which the immunosuppressive agent cyclosporin (CsA) had been administered at 30 mg/kg were considered to have survived (MST: 28 days or more) throughout the observation period. In Example 27, a survival period extension effect of the graft was found by individual administration.

[0276]

Next, the effects of Example 27 (0.3 to 10 mg/kg) in combination with CsA (10 mg/kg), in which survival throughout the observation period was not obtained by individual administration, were investigated. As a result of using these substances together, the MST was 26 days or more, thus confirming a clear combined effect.

[0277]

[Table 3]

Table 3: Survival Period Extension Effect of Allogenic Skin Graft for Rats

Compound (mg/kg, oral administration)	Individual survival period (days)	MST (days)
Control	8, 8, 8, 9, 11	8.8
CsA (10)	11, 11, 12, 13, 14	12.2
CsA (30)	28<, 28<, 28<, 28<, 28<	28<
Example 27 (3)	11, 11, 12, 12, 13	11.8
Example 27 (10)	12, 12, 14, 14, 18	14
Example 27 (30)	12, 13, 21, 28<, 28<	20.4<
Example 27 (0.3) + CsA (10)	28<, 28<, 28<, 28<, 28<	28<
Example 27 (1) + CsA (10)	28<, 28<, 28<, 28<, 28<	28<
Example 27 (3) + CsA (10)	23, 23, 28<, 28<, 28<	26<
Example 27 (10) + CsA (10)	28<, 28<, 28<, 28<, 28<	28<

[0278]

<Experiment example 4> Effect on Rat Lung Weight

Cr1j:Wistar rats (male) 5 to 8 weeks old were used in the experiment. An aqueous solution of the test compounds was respectively orally administered in an amount of 0.5 mL per 100 g of rat body weight. The lungs were extracted 6 hours later, and the lung weight was measured.

[0279]

Investigation of administering the compound of Example 26 up to a dose of 30 mg/kg and the compound of Example 27 up to a dose of 100 mg/kg showed that there were no effects whatsoever on lung weight.

[0280]

As a result, it was confirmed that the compounds of Examples

26 and 27 are highly safe compounds with mitigated side effects.

INDUSTRIAL APPLICABILITY


[0281]

According to the present invention, an amino alcohol derivative
5 having excellent immunosuppressive activity and safety can be
provided. The compound of the present invention is useful as a
preventive or treatment agent for rejection in organ transplants
or bone marrow transplants, a preventive or treatment agent for
autoimmune diseases such as inflammatory bowel disease, systemic
10 lupus erythematosus, Crohn's disease, nephrotic syndrome,
glomerular sclerosis, glomerular nephritis, multiple sclerosis,
and myasthenia gravis, a preventive or treatment agent for rheumatoid
arthritis, a preventive or treatment agent for psoriasis, allergic
contact dermatitis, and atopic dermatitis, a preventive or treatment
15 agent for hepatitis, hepatic steatosis, toxic liver injury, liver
cirrhosis-derived or diabetes mellitus-derived liver disease, a
preventive or treatment agent for allergic rhinitis, allergic
conjunctivitis and the like, and a preventive or treatment agent
for pulmonary fibrosis, idiopathic interstitial pneumonia, and
20 bronchial asthma.

CLAIMS

1. An amino alcohol derivative, represented by the general formula (1),


5 [Chemical formula 1]



[wherein R¹ represents a straight-chain alkyl group having 1 to 3 carbon atoms or trifluoromethyl group, R² represents a fluorine atom or a chlorine atom, R³ represents a straight-chain alkyl group having 1 to 3 carbon atoms, X represents an oxygen atom or a sulfur atom, and n denotes 2 or 3], or a pharmaceutically acceptable salt or hydrate thereof.

2. The amino alcohol derivative according to claim 1, wherein the compound represented by the general formula (1) is a compound represented by the general formula (1a),

[Chemical formula 2]



[wherein R³, X, and n are as described above], or a pharmaceutically acceptable salt or hydrate thereof.

3. The amino alcohol derivative according to claim 1 or 2, wherein in the general formulae (1) or (1a), R³ is a methyl group, or a pharmaceutically acceptable salt or hydrate thereof.

5 4. The amino alcohol derivative according to claim 1, wherein the compound represented by the general formula (1) is,

1)

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylpentan-1-ol,

10 2)

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol,

3)

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethylphenoxy)phenyl]-2-methylbutan-1-ol,

4)

(R)-2-amino-4-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylbutan-1-ol,

5)

20 (R)-2-amino-5-[2-chloro-4-(3-ethylphenylthio)phenyl]-2-methylpentan-1-ol,

6)


(R)-2-amino-5-[2-fluoro-4-(3-trifluoromethylphenylthio)phenyl]-2-methylpentan-1-ol, or

25 7)

(R)-2-amino-5-[2-chloro-4-(3-trifluoromethylphenylthio)phenyl]-2-propylpentan-1-ol, or a pharmaceutically acceptable salt or hydrate thereof.

5 5. The amino alcohol derivative according to claim 1, or a pharmaceutically acceptable salt or hydrate thereof, being obtainable by a step of allowing a compound represented by the general formula (2),


[Chemical formula 3]



10 [wherein R¹ represents a chlorine atom or a straight-chain alkyl group having 1 to 3 carbon atoms or trifluoromethyl group, R² represents a fluorine atom or a chlorine atom, A represents a halogen atom, X represents an oxygen atom or a sulfur atom, and n denotes

15 2 or 3] and a compound represented by the general formula (10),

[Chemical formula 4]



[wherein R³ represents a straight-chain alkyl group having 1 to 3 carbon atoms and R⁴ represents an alkyl group having 1 to 6 carbon atoms] to act in the presence of a base, and a step of subjecting the resultant product to acidolysis, then further protecting a nitrogen atom with a t-butoxycarbonyl group, reducing, and

20

deprotecting the nitrogen atom.

6. A pharmaceutical comprising the amino alcohol derivative, or
a pharmaceutically acceptable salt or hydrate thereof, according
5 to any of claims 1 to 5 as an active ingredient.

7. The pharmaceutical according to claim 6, being an
immunosuppressive agent.

10 8. The pharmaceutical according to claim 6, being a rejection
reaction preventive or treatment agent in organ transplants or bone
marrow transplants.

9. A pharmaceutical comprising the amino alcohol derivative, or
15 a pharmaceutically acceptable salt or hydrate thereof, according
to any of claims 1 to 5 and a calcineurin inhibitor in combination.