

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2021282468 B2**

(54) Title
Novel lactic acid bacteria and use thereof

(51) International Patent Classification(s)
C12N 1/20 (2006.01) **A61K 35/747** (2015.01)
A23L 33/135 (2016.01) **C12R 1/225** (2006.01)

(21) Application No: **2021282468** (22) Date of Filing: **2021.12.08**

(43) Publication Date: **2022.01.06**

(43) Publication Journal Date: **2022.01.06**

(44) Accepted Journal Date: **2023.10.05**

(62) Divisional of:
2018341753

(71) Applicant(s)
University-Industry Cooperation Group of Kyung Hee University;Navipharm Co, Ltd

(72) Inventor(s)
Kim, Dong-Hyun;Han, Myung Joo

(74) Agent / Attorney
FPA Patent Attorneys Pty Ltd, Level 19, South Tower 80 Collins Street, Melbourne, VIC, 3000, AU

(56) Related Art
CA 2998841 A1

Abstract

The present invention relates to a novel lactic acid bacterium, and more particularly, to a composition comprising *Lactobacillus reuteri* NK33 (KCCM12090P) or *Bifidobacterium adolescentis* NK98 (KCCM12297P), which is the novel lactic acid bacterium, useful in preventing and treating a neurological mental disease or an inflammatory disease.

Also, the present invention relates to a composition for diagnosing a neurological mental disease, comprising a preparation for measuring a level of an intestinal microorganism, a kit comprising the same, and a method for diagnosing the neurological mental disease, comprising a step of measuring the level of the intestinal microorganism.

Title

NOVEL LACTIC ACID BACTERIA AND USE THEREOF

Related application

This application is a divisional application of Australian application no. 2018341753, the entire contents of which is incorporated herein by reference.

Technical Field

The present invention relates to *Lactobacillus reuteri* and *Bifidobacterium adolescentis*, which are novel lactic acid bacteria, and more particularly, to a composition comprising the novel lactic acid bacteria useful in preventing and treating a neurological mental disease or an inflammatory disease.

Also, the present invention relates to a composition for diagnosing the neurological mental disease, comprising a preparation for measuring a level of an intestinal microorganism.

Background

In modern society, neurological mental diseases including anxiety, depression, schizophrenia, etc. are on the increase due to a rapid rise in stress, etc. In particular, the trend is that more and more patients are suffering from mental disorders such as depression, anxiety and the like due to various causes such as social and structural ones, etc. in our modern times overflowing with individualism.

In severe cases, patients with mental disorders may end up committing suicide accidents. In particular, it is reported that more than half of depressed patients consider suicide. Indeed, it is known that 10 to 15% of those patients actually commit suicide.

Mental disorders have no clear and objective criteria for judgment yet, such that symptoms thereof may vary depending on respective patients. If there is any

suspicion about mental disorders, treatment is required according to accurate diagnosis and examination. However, the reality is that such proper treatment is not done due to negative social attitudes toward hospital treatments for mental disorders. Also, drugs such as antidepressants, etc. used to treat mental disorders have not shown a satisfactory therapeutic effect thereon and may cause serious side effects such as cardiovascular diseases, suicide and the like, such that those drugs are limited in use.

On the other hand, as a result of study involving a use of natural products, a composition for treating mental disorders using a *Fomes fomentarius* extract and a *Lithospermum erythrorhizon* extract is disclosed in Korean patent publication No. 10-2017-0061457, but there is still a need for continuous research on effective lactic acid bacteria capable of treating mental disorders.

Against such backdrops, the present inventors identified that mental disorders may be diagnosed from a change in intestinal microbial communities, while studying a preventive and therapeutic agent for mental disorders, and further identified that novel lactic acid bacteria isolated from human and mouse feces show an effect of inhibiting a neurodegenerative disease-inducing factor, reducing anxious and depressive behaviors and the like, such that the novel lactic acid bacteria may be helpfully used in preventing or treating neurological mental diseases, particularly neurodegenerative diseases and mental disorders, thereby completed the present invention.

Prior Art References

Reference to any prior art in the specification is not an acknowledgement or suggestion that this prior art forms part of the common general knowledge in any jurisdiction or that this prior art could reasonably be expected to be combined with any other piece of prior art by a skilled person in the art.

Patent Document

(Patent Document 1) Korean patent publication No. 10-2017-0061457

Detailed Description of the Invention

In any aspect, the invention provides a novel lactic acid bacterium.

In another aspect, the invention provides a composition for preventing or treating a neurological mental disease, comprising the novel lactic acid bacterium.

In another aspect, the invention provides a composition for preventing or treating an inflammatory disease, comprising the novel lactic acid bacterium.

In another aspect, the invention provides a composition for diagnosing the neurological mental disease, comprising a preparation for measuring a level of an intestinal microorganism.

In another aspect, the invention provides a method for diagnosing the neurological mental disease, comprising: a step of measuring a level of an intestinal microorganism from feces isolated from an individual suspicious about the neurological mental disease; and a step of comparing the level of the said intestinal microorganism with that of the intestinal microorganism in feces of a control group, rather than that of the neurological mental disease.

In one aspect, the present invention provides *Lactobacillus reuteri* NK33 (depository institution: the Korean Culture Center of Microorganisms (KCCM), date of deposit: Aug. 04, 2017, accession number: KCCM12090P) or *Bifidobacterium adolescentis* NK98 (depository institution: the KCCM, date of deposit: Aug. 03, 2018, accession number: KCCM12297P).

Lactobacillus reuteri NK33 or *Bifidobacterium adolescentis* NK98 according to the present invention is characterized by being a novel lactic acid bacterium isolated and identified from human or mouse feces.

A 16S rDNA sequence for identification and classification of *Lactobacillus reuteri* NK33 according to the present invention is the same as SEQ ID NO: 1 attached to the present specification. Thus, *Lactobacillus reuteri* NK33 according to the present invention may include the 16S rDNA of SEQ ID NO: 1. As a result of analyzing the 16S rDNA sequence of the said SEQ ID NO: 1, such sequence was 99% homologous to that of known *Lactobacillus reuteri* strains, thus showing the highest molecular phylogenetic relationship with *Lactobacillus reuteri*. Thus, the said lactic acid bacterium was identified as *Lactobacillus reuteri*, named as *Lactobacillus reuteri* NK33, and deposited to the KCCM on Aug. 04, 2017 (accession number: KCCM12090P).

Lactobacillus reuteri NK33 according to the present invention is a gram-positive bacterium and a cell type thereof is bacillus. More particularly, the physiological characteristics of *Lactobacillus reuteri* NK33 may be analyzed according to a conventional method in the art, wherein the results thereof are as shown in a following table 3. Particularly, *Lactobacillus reuteri* NK33 may use as a carbon source L-arabinose, D-ribose, D-xylose, D-galactose, D-glucose, D-fructose, D-mannose, mannitol, sorbitol, N-acetyl-glucosamine, amygdaline, arbutin, esculin, salicin, cellobiose, maltose, lactose, melibiose, sucrose, trehalose, melezitose, raffinose, gentiobiose, D-turanose and gluconate.

A 16S rDNA sequence for identification and classification of *Bifidobacterium adolescentis* NK98 according to the present invention is the same as SEQ ID NO: 38 attached to the present specification. Thus, *Bifidobacterium adolescentis* NK98 according to the present invention may include the 16S rDNA of SEQ ID NO: 38. As a result of analyzing the 16S rDNA sequence of the said SEQ ID NO: 38, such sequence was 98% homologous to that of a known *Bifidobacterium adolescentis* strain, thus

showing the highest molecular phylogenetic relationship with *Bifidobacterium adolescentis*. Thus, the said lactic acid bacterium was identified as *Bifidobacterium adolescentis*, named as *Bifidobacterium adolescentis* NK98, and deposited to the KCCM on Aug. 03, 2018 (accession number: KCCM12297P).

The physiological characteristics of *Bifidobacterium adolescentis* NK98 according to the present invention may be analyzed according to a conventional method in the art, wherein the results thereof are as shown in a following table 4. Particularly, *Bifidobacterium adolescentis* NK98 may use as a carbon source D-glucose, D-mannitol, D-lactose, D-saccharose, D-maltose, salicin, D-xylose, L-arabinose, gelatin, esculin ferric citrate, D-cellobiose, D-raffinose and D-trehalose.

In other aspect, the present invention provides a pharmaceutical composition for preventing or treating a neurological mental disease, comprising a novel lactic acid bacterium.

In the present invention, the said novel lactic acid bacterium may be *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof.

“*Lactobacillus reuteri* NK33” and “*Bifidobacterium adolescentis* NK98” according to the present invention are the same as described above.

Particularly, the lactic acid bacterium contained in the pharmaceutical composition according to the present invention may be a live bacterial cell thereof, a dead bacterial cell thereof, a culture product thereof, a crush thereof or an extract thereof, but any type of lactic acid bacterium may be used without limitation, as long as it may achieve a preventive or therapeutic effect on the neurological mental disease.

In the present invention, the term “culture product” means an object obtained by culturing a lactic acid bacterium in a known liquid medium or solid medium, and is a concept encompassing a novel lactic acid bacterium in the present invention.

A neurological mental disease according to the present invention may be a neurodegenerative disease or a mental disorder.

Particularly, the neurological mental disease according to the present invention may be the mental disorder, and the said mental disorder may be one or more selected from the group including anxiety, depression, mood disorder, insomnia, delusional disorder, obsessive disorder, migraine, stress, memory disorder, cognitive disorder and disturbance attention.

In an exemplary embodiment of the present invention, it was identified that, when *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 is administered to an animal model with stress induced, anxious and depressive behaviors caused by stress are remarkably reduced; an activity of NF- κ B is inhibited in a hippocampus of the animal model with stress induced; an expression of a brain derivated neurotrophic factor is increased; and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide (LPS), which are stress parameters in blood, are decreased. From the results, it was also identified that a pharmaceutical composition comprising the said lactic acid bacteria may be helpfully used in preventing or treating a neurological mental disease, particularly a mental disorder.

Particularly, the neurological mental disease according to the present invention may be a neurodegenerative disease, wherein the said neurodegenerative disease may be one or more selected from the group consisting of Parkinson's disease, Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, spinocerebellar atrophy, Tourette's syndrome, Friedrich's ataxia, Machado-Joseph's disease, dementia, dystonia, progressive supranuclear palsy and frontotemporal dementia.

In an exemplary embodiment of the present invention, it was identified that,

when nerve cells are treated with the said lactic acid bacteria along with a stress hormone corticosterone, the activity of NF- κ B, known as a substance of inducing neurodegenerative diseases such as Alzheimer's disease, is inhibited, and an expression of a brain derivated neurotrophic factor (BDNF), known to be decreased in aging, dementia and the like, is increased at the same time (Tables 5 and 6). From the results, it was also identified that the pharmaceutical composition comprising the said lactic acid bacteria may be helpfully used in preventing and treating the neurological mental disease, particularly the neurodegenerative disease.

Also, in an exemplary embodiment of the present invention, it was identified that, when co-administered with *Lactobacillus reuteri* NK33 and *Bifidobacterium adolescentis* NK98, an effect of improving a neurological mental disease, particularly the effect of reducing stress, is remarkably increased compared to each group administered with the said lactic acid bacteria alone (Table 20).

In the present invention, the said pharmaceutical composition may further comprise *Bifidobacterium adolescentis* IM38 KCCM11807P.

The said *Bifidobacterium adolescentis* IM38 KCCM11807P is a known lactic acid bacterium disclosed in Korean patent publication No. 10-2017-0090359, and may be easily available based on Korean patent publication No. 10-2017-0090359.

In an exemplary embodiment of the present invention, it was identified that, when *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 and *Bifidobacterium adolescentis* IM38 are co-administered to an animal model with stress induced, anxious and depressive behaviors caused by stress are remarkably reduced, and an amount of corticosterone, which is a stress parameter in blood, is decreased.

In another aspect, the present invention provides a pharmaceutical

composition for preventing or treating an inflammatory disease, comprising *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof.

“*Lactobacillus reuteri* NK33” and “*Bifidobacterium adolescentis* NK98” according to the present invention are the same as described above.

Particularly, the lactic acid bacterium contained in the pharmaceutical composition according to the present invention may be a live bacterial cell thereof, a dead bacterial cell thereof, a culture product thereof, a crush thereof or an extract thereof, but any type of a lactic acid bacterium may be used without limitation, as long as it may achieve a preventive or therapeutic effect on the inflammatory disease.

In the present invention, the term “culture product” means an object obtained by culturing a lactic acid bacterium in a known liquid medium or solid medium, and is a concept encompassing a novel lactic acid bacterium in the present invention.

An inflammatory disease according to the present invention may be one or more selected from the group including arthritis, gout, hepatitis, asthma, obesity, corneitis, gastritis, enteritis, nephritis, colitis, diabetes, tuberculosis, bronchitis, pleurisy, peritonitis, spondylitis, pancreatitis, inflammatory pain, urethritis, cystitis, vaginitis, arteriosclerosis, septicemia, burn, dermatitis, periodontitis and gingivitis.

In an exemplary embodiment of the present invention, it was identified that, when macrophage isolated from a mouse is treated with the said lactic acid bacteria along with lipopolysaccharide (LPS), an inflammatory reaction inducer, an inflammatory reaction is remarkably inhibited (Tables 5 and 6). From the results, it was also identified that the pharmaceutical composition comprising the said lactic acid bacteria according to the invention may be helpfully used in preventing and treating the inflammatory disease.

Particularly, the inflammatory disease may be colitis.

In an exemplary embodiment of the present invention, it was identified that, when *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 is administered to an animal model with colitis induced by stress, a colon length thereof, which is a colitis indicator, is recovered to a normal level, and amounts of myeloperoxidase, COX-2 and iNOS, which are colitis parameters, are decreased and an activity of TNF- α is decreased, too (Fig. 15 and Table 15). From the results, it was also identified that the pharmaceutical composition comprising the said *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 may be helpfully used in preventing and treating the inflammatory disease, particularly colitis.

A pharmaceutical composition for preventing or treating a neurological mental disease, or a pharmaceutical composition for preventing or treating an inflammatory disease according to the present invention may be prepared into a pharmaceutical dosage form by means of a well-known method in the art, such that an active component of such composition may be provided via a fast, suspended or prolonged release, after being administered to a mammal. In preparing a dosage form, the pharmaceutical composition according to the present invention may further comprise a pharmaceutically acceptable carrier, to the extent that such carrier does not suppress an activity of a novel lactic acid bacterium.

The pharmaceutically acceptable carrier may include, but not limited thereto, conventionally used ones, for example, lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, maltitol, starch, acacia rubber, alginate, gelatin, calcium phosphate, calcium silicate, cellulose, methyl cellulose, microcrystalline cellulose, polyvinyl pyrrolidone, water, methyl hydroxybenzoate, propylhydroxybenzoate, talc, magnesium stearate, mineral oil and the like. Also, the pharmaceutical composition according to the present invention may comprise a diluent or an excipient such as filler,

extender, binder, humectant, disintegrant, surfactant, etc., and other pharmaceutically acceptable additives.

A dosage of the pharmaceutical composition according to the present invention needs to be a pharmaceutically effective amount. The “pharmaceutically effective amount” means an amount enough to prevent or treat neurological mental diseases or inflammatory diseases at a reasonable benefit/risk ratio applicable to medical treatment. An effective dose level may be variously selected by those skilled in the art according to such factors as a formulation method, a patient's condition and weight, the patient's gender, age and degree of disease, a drug form, an administration route and period, an excretion rate, reaction sensitivity, etc. The effective amount may vary depending on a route of treatment, a use of excipient and a possibility of being used with other drugs, as recognized by those skilled in the art. However, in case of a preparation for oral administration to achieve a preferable effect, the composition according to the present invention may be generally administered to an adult in an amount of 0.0001 to 100 mg/kg a day, preferably 0.001 to 100 mg/kg a day based on 1 kg of body weight. When such preparation is administered as shown above, *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 according to the present invention may be administered in an amount of 1×10^2 CFU/60kg to 1×10^{11} CFU/60kg a day. Such administration may be done once a day, or divided into several times a day. The said dosage does not limit the scope of the present invention in any aspect.

The pharmaceutical composition for preventing or treating a neurological mental disease, or the pharmaceutical composition for preventing or treating an inflammatory disease according to the present invention may be administered to mammals such as mice, livestock, humans, etc. through various routes. Particularly,

the pharmaceutical composition according to the present invention may be orally or parenterally administered (for example, applied or injected intravenously, subcutaneously or intraperitoneally), but may be preferably orally administered. A solid preparation for oral administration may include powder, granule, tablet, capsule, soft capsule, pill, etc. A liquid preparation for oral administration may include a suspending agent, liquid for internal use, emulsion, syrup, aerosol, etc., but may also include various excipients, for example, humectant, a sweetening agent, a flavoring agent, preservative, etc. in addition to water and liquid paraffin, which are frequently used simple diluents. A preparation for parenteral administration may be used by being formulated into a dosage form of external preparation and sterilized injectable preparation such as sterilized aqueous solution, liquid, non-aqueous solvent, a suspending agent, emulsion, eye drop, eye ointment, syrup, suppository, aerosol, etc., according to respective conventional methods, and preferably may be used by preparing a pharmaceutical composition of cream, gel, patch, spray, ointment, plaster, lotion, liniment, eye ointment, eye drop, paste or cataplasma, but not limited thereto. A preparation for local administration may be an anhydrous or aqueous form depending on a clinical prescription. As the non-aqueous solvent and the suspending agent above, propylene glycol, polyethylene glycol, vegetable oil like olive oil, injectable ester like ethyl oleate, etc. may be used. As a base of the suppository above, witepsol, macrogol, tween 61, cacao butter, laurin oil, glycerogelatin, etc. may be used.

In another aspect for achieving the said objectives, the present invention provides a method for preventing or treating a neurological mental disease, comprising a step of administering *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof to an individual.

In another aspect, the present invention provides a method for preventing or

treating a neurological mental disease, comprising a step of administering *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof to an individual.

In another aspect, the present invention provides a method for preventing or treating an inflammatory disease, comprising a step of administering *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof to an individual.

The terms “*Lactobacillus reuteri* NK33,” “*Bifidobacterium adolescentis* NK98,” “administration,” “neurological mental disease,” “inflammatory disease” and the like according to the present invention are the same as described above.

The individual refers to an animal, and may be typically a mammal, on which treatment using the inventive novel lactic acid bacterium may show a beneficial effect. A preferable example of such individual may include primates like human, as well as a rat, mouse, monkey, dog, cat, cow, horse, pig, sheep or goat. Also, such individuals may include all the individuals having a symptom of a neurological mental disease or an inflammatory disease, or having a risk of having such symptom.

In another aspect, the present invention provides a health functional food for preventing or improving a neurological mental disease, comprising *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof.

In another aspect, the present invention provides a health functional food for preventing or improving an inflammatory disease, comprising *Lactobacillus reuteri* NK33, *Bifidobacterium adolescentis* NK98 or a mixture thereof.

The terms “*Lactobacillus reuteri* NK33,” “*Bifidobacterium adolescentis* NK98,” “administration,” “neurological mental disease,” “inflammatory disease” and the like according to the present invention are the same as described above.

The health functional food, which puts an emphasis on a body modulating function of food, is a food given added value by means of a physical, biochemical or bioengineering method, such that the food may act to express the functions thereof for a particular purpose. A component of such health functional food is designed and processed to fully exert its body modulating function in vivo, which is involved in defending a living body, adjusting a body rhythm, preventing a disease and helping a recovery from the disease, and may comprise food supplementary additives, which are acceptable as food, sweeteners or functional raw materials.

In case of using *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 according to the present invention as a health functional food (or health functional beverage additives), the said lactic acid bacteria may be added thereto as they are, used along with other food or food components, or appropriately used according to a conventional method. A mixed amount of the said lactic acid bacteria may be appropriately determined depending on a purpose of use thereof (prevention, health, improvement or therapeutic treatment).

The health functional food may comprise various nutrients, vitamins, minerals (electrolytes), flavoring agents such as synthetic flavoring agents, natural flavoring agents and the like, coloring agents and enhancers (cheese, chocolate, etc.), pectic acid and salts thereof, organic acid, protective colloidal thickeners, pH adjusting agents, stabilizers, preservatives, glycerin, alcohol, carbonators used in carbonated beverages, etc. Also, the health functional food according to the present invention may comprise pulp for preparing fruit and vegetable based beverages. Such components may be used alone or in combination, and a ratio of such additives is generally selected from a range of 0.001 to 50 parts by weight with regard to a total weight of the composition.

A type of the health functional food has no particular limitation. As food, to

which the said lactic acid bacteria may be added, there are sausage, meats, bread, chocolates, snacks, candies, confectionery, ramen, pizza, other noodles, chewing gums, dairy products including ice creams, various soups, beverages, teas, health drinks, alcohol beverages, vitamin complexes, etc. In case of being formulated into beverages, a liquid component, which is added to the beverages in addition to the novel lactic acid bacteria, may include, but not limited thereto, various flavors, natural carbohydrates or the like as an additional component just as contained in conventional beverages. The aforementioned natural carbohydrates may be monosaccharide (ex. glucose, fructose, etc.), disaccharide (ex. maltose, sucrose, etc.) and polysaccharide (ex. conventional sugar such as dextrin, cyclodextrin, etc.), as well as sugar alcohol such as xylitol, sorbitol, erythritol, etc.

In another aspect for, the present invention provides a composition for diagnosing a neurological mental disease, comprising a preparation for measuring a level of intestinal microorganisms.

The terms such as “neurological mental disease,” etc. according to the present invention are the same as described above.

In the present invention, the term “intestinal microorganism” refers to a microorganism present in a digestive tract, particularly the microorganism present specifically in intestines, wherein such intestinal microorganism may be one or more selected from the group including bacteroidetes, actinobacteria, firmicutes, bifidobacteria, lactobacilli, β -proteobacteria, δ -proteobacteria, γ -proteobacteria, ϵ -proteobacteria and enterobacteriaceae.

Particularly, a composition for diagnosis according to the present invention may diagnose a neurological mental disease, if there is an increase in a level of one or more selected from the group including β -proteobacteria, δ -proteobacteria, γ -

proteobacteria, ϵ -proteobacteria and enterobacteriaceae among the said intestinal microorganisms; such composition may diagnose the neurological mental disease, if there is a decrease in a level of one selected from the group including bacteroidetes, actinobacteria, firmicutes, bifidobacteria and lactobacilli among the said intestinal microorganisms; and such composition may diagnose the neurological mental disease, if there is an increase in a level of one or more selected from the group including β -proteobacteria, δ -proteobacteria, γ -proteobacteria, ϵ -proteobacteria and enterobacteriaceae among the said intestinal microorganisms, and if there is a decrease in a level of one selected from the group including bacteroidetes, actinobacteria, firmicutes, bifidobacteria and lactobacilli among the said intestinal microorganisms.

Particularly, the said enterobacteriaceae, of which level is increased, may be one or more selected from the group including *Klebsiella oxytoca*, *Escherichia coli* and *Morganella morganii*.

Also, the said lactobacilli, of which level is decreased, may be one or more selected from the group including *Lactobacillus reuteri*, *Lactobacillus johnsonii* and *Lactobacillus rhamnosus*, and the said bifidobacteria, of which level is decreased, may be *Bifidobacterium animalis*.

The composition for diagnosing a neurological mental disease, provided in the present invention, comprises a preparation capable of measuring a level of an intestinal microorganisms, wherein an occurrence of the neurological mental disease to an individual may be identified by measuring the level of the intestinal microorganism from feces isolated from the individual, who needs to be identified about the occurrence of the neurological mental disease, by means of the preparation contained in the said composition. The said individual may be an animal, preferably a

mammal, for example, primates like human, as well as a rat, mouse, monkey, dog, cat, cow, horse, pig, sheep or goat.

The preparation may be used to identify whether the neurological mental disease has occurred to the individual or not, by measuring the level of the intestinal microorganism isolated from feces of the said individual.

The preparation capable of measuring the said level of the intestinal microorganism may be particularly a primer, probe or the like, capable of measuring a level of a specific gene of the intestinal microorganism; may be an antibody, aptamer or the like, capable of measuring a level of a protein expressed from the said specific gene; may be a primer, probe or the like, to quantitatively analyze a 16S rDNA sequence of the intestinal microorganism; and may be a preparation for pyrosequencing, etc. to quantitatively analyze a genome DNA sequence of the intestinal microorganism, but not particularly limited thereto, as long as the preparation may be used to quantitatively analyze the intestinal microorganism isolated from feces of the individual.

Particularly, a primer specific to firmicutes, the said intestinal microorganism, may be a primer pair consisting of SEQ ID NO: 2 and SEQ ID NO: 3; a primer specific to bacteroidetes may be a primer pair consisting of SEQ ID NO: 4 and SEQ ID NO: 5; a primer specific to β -proteobacteria may be a primer pair consisting of SEQ ID NO: 6 and SEQ ID NO: 7; a primer specific to δ/γ -proteobacteria may be a primer pair consisting of SEQ ID NO: 8 and SEQ ID NO: 9; a primer specific to ϵ -proteobacteria may be a primer pair consisting of SEQ ID NO: 10 and SEQ ID NO: 11; a primer specific to actinobacteria may be a primer pair consisting of SEQ ID NO: 12 and SEQ ID NO: 13; and a primer specific to enterobacteriaceae may be a primer pair consisting of SEQ ID NO: 18 and SEQ ID NO: 19.

In an exemplary embodiment of the present invention, the levels of intestinal microorganisms were measured by carrying out pyrosequencing by means of a preparation for pyrosequencing containing a barcoded primer of SEQ ID NO: 2 to SEQ ID NO: 19 in Table 1 (Experimental Example 3).

The composition according to the present invention may further comprise a preparation for measuring an amount of lipopolysaccharide (LPS). Particularly, the preparation for measuring the said amount of lipopolysaccharide (LPS) may be a preparation conventionally used to measure the amount of lipopolysaccharide (LPS), more particularly a preparation included in Limulus amoebocyte lysate (LAL) assay kit, but not limited thereto.

In an exemplary embodiment of the present invention, it was identified that the amount of lipopolysaccharide (LPS) is increased as a result of analyzing feces of an animal model with a mental disorder induced (Figs. 1, 2, 7 and 11).

In another aspect, the present invention provides a kit for diagnosing a neurological mental disease, comprising the said composition.

The kit according to the present invention may include a collection container capable of collecting feces of an individual to be identified about an occurrence of the neurological mental disease thereto by using the said composition; a buffer solution for extracting an intestinal microorganism in feces; a measurement means used to measure the intestinal microorganism; and the like.

In another aspect, the present invention provides a method for diagnosing a neurological mental disease, comprising: a step of measuring a level of an intestinal microorganism from feces isolated from an individual suspicious about the neurological mental disease; and a step of comparing the level of the said intestinal microorganism with a level of the intestinal microorganism in feces of a control group,

rather than that of the neurological mental disease.

Also, the present invention provides an information-offering method for diagnosing a neurological mental disease, comprising: a step of measuring a level of an intestinal microorganism from feces isolated from an individual suspicious about the neurological mental disease; and a step of comparing the level of the said intestinal microorganism with a level of the intestinal microorganism in feces of a control group, rather than that of the neurological mental disease.

The feces mean feces isolated and collected from an individual as a sample for measuring a level of an intestinal microorganism.

Particularly, the intestinal microorganism may be one or more selected from the group including bacteroidetes, actinobacteria, firmicutes, bifidobacteria, lactobacilli, β -proteobacteria, δ -proteobacteria, γ -proteobacteria, ϵ -proteobacteria and enterobacteriaceae.

The method for diagnosing a neurological mental disease or providing information thereon according to the present invention may diagnose whether the neurological mental disease occurs or has a risk of occurrence by identifying a microorganism, of which level is increased, among the said intestinal microorganisms.

Particularly, the method for diagnosing a neurological mental disease or providing information thereon may further comprise a step of determining an individual, whose level of an intestinal microorganism is increased, as the neurological mental disease by comparing the level of the intestinal microorganism in feces of an individual suspicious about the neurological mental disease with a level of the intestinal microorganism in feces of a control group, rather than that of the neurological mental disease.

The intestinal microorganism, of which level is increased, may be one or more

selected from the group including β -proteobacteria, δ -proteobacteria, γ -proteobacteria, ϵ -proteobacteria and enterobacteriaceae.

Particularly, the enterobacteriaceae may be one or more selected from the group including *Klebsiella oxytoca*, *Escherichia coli* and *Morganella morganii*.

Also, the method for diagnosing a neurological mental disease or providing information thereon according to the invention may diagnose whether the neurological mental disease occurs or has a risk of occurrence by identifying a microorganism, of which level is decreased, among the said intestinal microorganisms.

The terms such as “neurological mental disease,” etc. according to the present invention are the same as described above.

Particularly, the method for diagnosing a neurological mental disease or providing information thereon may further comprise a step of determining an individual, whose level of an intestinal microorganism is decreased, as the neurological mental disease by comparing a level of the intestinal microorganism in feces of an individual suspicious about the neurological mental disease with a level of the intestinal microorganism in feces of a control group, rather than that of the neurological mental disease.

The intestinal microorganism, of which level is decreased, may be one or more selected from the group including bacteroidetes, actinobacteria, firmicutes, bifidobacteria and lactobacilli.

Also, the lactobacilli may be one or more selected from the group including *Lactobacillus reuteri*, *Lactobacillus johnsonii* and *Lactobacillus rhamnosus*, and the bifidobacteria may be *Bifidobacterium animalis*.

In an exemplary embodiment of the present invention, as a result of identifying a change in an intestinal microbial community by using feces isolated from an animal

model with stress induced, such as immobilization stress, antibiotic stress or the like, it was identified that levels of β -proteobacteria, δ -proteobacteria, γ -proteobacteria, ϵ -proteobacteria and enterobacteriaceae are increased, and levels of bacteroidetes, actinobacteria, firmicutes, bifidobacteria and lactobacilli are decreased compared to an animal model without stress (Figs. 2, 6, 8, 10 and 12). In particular, it was identified that, as a particular enterobacteriaceae, of which level is increased, there are *Klebsiella oxytoca*, *Escherichia coli* and *Morganella morganii*; as a particular lactobacilli, of which level is decreased, there are *Lactobacillus reuteri*, *Lactobacillus johnsonii* and *Lactobacillus rhamnosus*; and as a particular bifidobacteria, of which level is decreased, there is *Bifidobacterium animalis*.

In another aspect, the present invention provides a use of a novel lactic acid bacterium for preventing or treating a neurological mental disease. Particularly, the present invention provides a use of *Lactobacillus reuteri* NK33 for preventing or treating the neurological mental disease. Also, the present invention provides a use of *Bifidobacterium adolescentis* NK98 for preventing or treating the neurological mental disease.

In another aspect, the present invention provides a use of a novel lactic acid bacterium for preventing or treating an inflammatory disease. Particularly, the present invention provides a use of *Lactobacillus reuteri* NK33 for preventing or treating the inflammatory disease. Also, the present invention provides a use of *Bifidobacterium adolescentis* NK98 for preventing or treating the inflammatory disease.

In another aspect, the present invention provides a use of a composition comprising a novel lactic acid bacterium in preparing a drug for preventing or treating a neurological mental disease. Particularly, the present invention provides a use of a composition comprising *Lactobacillus reuteri* NK33 in preparing a drug for

preventing or treating the neurological mental disease. Also, the present invention provides a use of a composition comprising *Bifidobacterium adolescentis* NK98 in preparing a drug for preventing or treating the neurological mental disease.

The terms “*Lactobacillus reuteri* NK33,” “*Bifidobacterium adolescentis* NK98,” “neurological mental disease,” “inflammatory disease” and the like according to the present invention are the same as described above.

The numerical values described in the present specification should be interpreted to include a range of equivalents thereof, unless otherwise stated.

Hereinafter, the present invention will be described in detail through preferred Examples for better understanding of the present invention. However, the following Examples are provided only for the purpose of illustrating the present invention, and thus the present invention is not limited thereto.

Advantageous Effects

A novel lactic acid bacterium according to the present invention, i.e. *Lactobacillus reuteri* NK33 or *Bifidobacterium adolescentis* NK98 has an effect on inhibiting a neurodegenerative disease-inducing factor and reducing anxious and depressive behaviors. Thus, the novel lactic acid bacterium according to the present invention may be used as a composition for preventing or treating a neurological mental disease, and particularly effective in preventing and treating a neurodegenerative disease and a mental disorder.

Also, the novel lactic acid bacterium according to the present invention is effective in inhibiting an inflammatory reaction, and thus may be used in a composition for preventing or treating an inflammatory disease, and particularly effective in preventing and treating colitis.

Further, a composition for measuring a level of intestinal microorganisms

according to the present invention is effective in diagnosing a neurological mental disease by measuring the level of intestinal microorganisms.

Brief Description of the Drawings

Fig. 1 shows results of measuring anxious behaviors and blood indicators with regard to a mouse (IS) subjected to immobilization stress; (a) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are decreased as a result of an elevated plus maze test; (b) is a graph of identifying that time in bright area is decreased as a result of a light/dark transition test; (c) is a graph of identifying that a marble-burying behavior is increased in a marble burying test; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased in a hippocampus and an expression level of a brain derived neurotrophic factor (BDNF) is decreased; (e) is a graph of identifying that corticosterone in blood is increased; (f) is a graph of identifying that IL-6 in blood is increased; (g) is a graph of identifying that TNF- α in blood is increased; and (h) is a graph of identifying that lipopolysaccharide (LPS) in blood is increased.

Fig. 2 (a) is a diagram of identifying that there is a change in an intestinal microbial community in feces of a mouse (IS) subjected to immobilization stress; and Fig. 2 (b) is a graph of identifying that there is an increase in lipopolysaccharide (LPS) in feces of a mouse (IS) subjected to immobilization stress.

Fig. 3 shows results of measuring colitis indicators with regard to a mouse (IS) subjected to immobilization stress; (a) is a graph of identifying that there is a decrease in a length of a colon of a mouse (IS) subjected to immobilization stress; (b) is a graph of identifying that myeloperoxidase is increased in the colon; (c) is a graph of identifying that TNF- α is increased in the colon; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased and expressions of COX-2 and iNOS are

increased in the colon; and (e) is an image of identifying that occludin and claudin-1 are decreased in the colon.

Fig. 4 shows results of identifying a state of an animal model (FIS), administered with feces of an animal model with immobilization stress induced; (a) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are decreased as a result of an elevated plus maze test; (b) is a graph of identifying that time in bright area is decreased as a result of a light/dark transition test; (c) is a graph of identifying that a marble-burying behavior is increased in a marble burying test; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased in a hippocampus and an expression level of a brain derived neurotrophic factor (BDNF) is decreased; (e) is a graph of identifying that a length of a colon is decreased; and (f) is a graph of identifying that myeloperoxidase is increased in the colon.

Fig. 5 shows results of measuring blood indicators with regard to an animal model (FIS) administered with feces of an animal model with immobilization stress induced; (a) is a graph of identifying that corticosterone in blood is increased; (b) is a graph of identifying that IL-6 in blood is increased; and (c) is a graph of identifying that TNF- α in blood is increased.

Fig. 6 shows results of identifying a change in colitis indicators and an intestinal microbial community with regard to an animal model (FIS) administered with feces of an animal model with immobilization stress induced; (a) is a graph of identifying that TNF- α is increased; (b) is a graph of identifying that IL-6 is increased in a colon; (c) is a graph of identifying that IL-10 is decreased in the colon; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased and expressions of COX-2 and iNOS are increased in the colon; and (e) is a diagram of identifying that an intestinal microbial community in feces is changed.

Fig. 7 shows results of measuring anxious behaviors and blood indicators with regard to a mouse (AP) subjected to antibiotic stress; (a) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are decreased as a result of an elevated plus maze test; (b) is a graph of identifying that time in bright area is decreased as a result of a light/dark transition test; (c) is a graph of identifying that a marble-burying behavior is increased in a marble burying test; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased in a hippocampus and an expression level of a brain derived neurotrophic factor (BDNF) is decreased; (e) is a graph of identifying that corticosterone in blood is increased; (f) is a graph of identifying that IL-6 in blood is increased; (g) is a graph identifying that TNF- α in blood is increased; and (h) is a graph of identifying that lipopolysaccharide (LPS) in blood is increased.

Fig. 8 (a) is a diagram of identifying that there is a change in an intestinal microbial community in feces of a mouse (AP) subjected to antibiotic stress; and Fig. 8 (b) is a graph of identifying that there is an increase in lipopolysaccharide (LPS) in feces of a mouse (AP) subjected to antibiotic stress.

Fig. 9 shows results of measuring colitis indicators with regard to a mouse (AP) subjected to antibiotic stress; (a) is a graph of identifying that there is a decrease in a length of a colon; (b) is a graph of identifying that myeloperoxidase is increased in the colon; (c) is a graph of identifying that TNF- α is increased in the colon; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased and expressions of COX-2 and iNOS are increased in the colon; and (e) is an image of identifying that occludin and claudin-1 are decreased in the colon.

Fig. 10 (A) is a diagram of identifying which bacterium is increased as a result of culturing feces of a mouse (IS) subjected to immobilization stress and feces of a

mouse (AP) subjected to antibiotic stress in a selective medium; and Fig. 10 (B) is a diagram of identifying which bacterium is decreased as a result of culturing feces of the mouse (IS) subjected to immobilization stress and feces of the mouse (AP) subjected to antibiotic stress in a selective medium.

Fig. 11 shows results of measuring anxious behaviors and blood indicators with regard to a mouse (AP) administered with a microorganism increased in feces of mice subjected to immobilization stress and antibiotic stress; (a) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are remarkably decreased in a group administered with *Klebsiella oxytoca*, as a result of an elevated plus maze test with mice administered with *Klebsiella oxytoca* (KO), *Escherichia coli* (EC), *Aerococcus urinaeequi* (AU) and *Morganella morganii* (MM); (b) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are decreased as a result of an elevated plus maze test with a mouse administered with *Klebsiella oxytoca* (KO); (c) is a graph of identifying that time in bright area is decreased as a result of a light/dark transition test; (d) is a graph of identifying that a marble-burying behavior is increased in a marble burying test; (e) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased in a hippocampus and an expression level of a brain derived neurotrophic factor (BDNF) is decreased; (f) is a graph of identifying that corticosterone in blood is increased; (g) is a graph of identifying that IL-6 in blood is increased; (h) is a graph of identifying that TNF- α in blood is increased; and (i) is a graph of identifying that lipopolysaccharide (LPS) in blood is increased.

Fig. 12 (a) is a diagram of identifying that there is a change in an intestinal microbial community in feces of a mouse (KO) administered with *Klebsiella oxytoca*; and Fig. 12 (b) is a graph of identifying that there is an increase in lipopolysaccharide (LPS) in feces of a mouse (KO) administered with *Klebsiella oxytoca*.

Fig. 13 shows results of measuring colitis indicators with regard to a mouse (KO) administered with *Klebsiella oxytoca*; (a) is a graph of identifying that a length of a colon is decreased; (b) is a graph of identifying that myeloperoxidase is increased in the colon; (c) is a graph of identifying that TNF- α is increased in the colon; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is increased and expressions of COX-2 and iNOS are increased in the colon; and (e) is an image of identifying that occludin and claudin-1 are decreased in the colon.

Fig. 14 shows results of measuring anxious behaviors with regard to and comparing blood indicators between a group (IS), in which a mouse subjected to immobilization stress is administered with physiological saline solution, and a group administered with *Lactobacillus reuteri* NK33 (IS+LR), a novel lactic acid bacterium; (a) is a graph of identifying that time spent in open arms (OT) and open arm entries (OE) are increased as a result of an elevated plus maze test with the group administered with *Lactobacillus reuteri* NK33 (IS+LR), the novel lactic acid bacterium; (b) is a graph of identifying that time in bright area is increased as a result of a light/dark transition test; (c) is a graph of identifying that a marble-burying behavior is decreased in a marble burying test; (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is inhibited in a hippocampus and an expression level of a brain derived neurotrophic factor (BDNF) is increased; (e) is a graph of identifying that corticosterone in blood is decreased; (f) is a graph of identifying that IL-6 in blood is decreased; and (g) is a graph of identifying that TNF- α in blood is decreased.

Fig. 15 shows results of measuring colitis indicators with regard to a group (IS), in which a mouse subjected to immobilization stress is administered with physiological saline solution, and a group administered with *Lactobacillus reuteri* NK33 (IS+LR), a novel lactic acid bacterium; (a) is a graph of identifying that a length

of a colon is recovered; (b) is a graph of identifying that myeloperoxidase is decreased in the colon; (c) is a graph of identifying that TNF- α is decreased in the colon; and (d) is an image of identifying that an NF- κ B activity (p-p65/p65) is decreased and expressions of COX-2 and iNOS are decreased in the colon.

Mode for Invention

Hereinafter, the present invention will be described in detail through preferred Examples and Experimental Examples for better understanding of the present invention. However, the following Examples and Experimental Examples are provided only for the purpose of illustrating the present invention, and thus the present invention is not limited thereto.

Experimental Example 1. Animal model with a mental disorder induced

(1) Immobilization stress

In order to induce a mental disorder such as anxiety, depression, stress or the like, a mouse was fixed to a 3 x 10 cm cylindrical-shaped immobilization stress apparatus, such that the mouse may not move therein at all.

Particularly, in order to prepare a mouse with anxiety induced by immobilization stress, the said mouse, which had been fixed to the apparatus, was repeatedly subjected to immobilization stress five times, in such a way that the mouse was held with its head up for two hours once every other day, and then a behavioral experiment was performed in two hours after a final immobilization stress.

Also, in order to prepare a mouse with depression induced by immobilization stress, the said mouse, which had been fixed to the apparatus, was repeatedly subjected to continuous immobilization stress once daily for two days, in such a way that the mouse was held with its head up for 12 hours. A lactic acid bacterium was

administered once daily for five days, and a behavioral experiment was performed in one hour after a final administration.

(2) Antibiotic stress

In order to induce a mental disorder such as anxiety, depression, stress or the like, a mouse was administered ampicillin (100 mg/kg) for two days in a row. An anxiety behavior was measured in 10 days after administration.

(3) Experimental method on an animal model with a mental disorder induced

For a mouse model with immobilization stress induced, a lactic acid bacterium or physiological saline solution was administered thereto depending on an experimental group once daily for three days from Day 7 after starting the immobilization stress. For a mouse model with antibiotic stress induced, a lactic acid bacterium or physiological saline solution was administered thereto depending on an experimental group once daily for three days from Day 7 after starting the antibiotic stress.

An anxiety behavior was measured in five hours after administering the lactic acid bacterium or physiological saline solution, and blood indicators (corticosterone, IL-6, TNF- α , etc.) were measured by collecting blood in one hour after finishing an elevated plus maze test.

Experimental Example 2. Behavior measurement method for stress diagnosis

(1) Elevated plus maze (EPM) test

An elevated plus maze is an experimental apparatus to measure a degree of a mental disorder such as stress, anxiety or the like. An elevated plus maze experimental apparatus used in the present experiment is a black plexiglass apparatus, in which two

open arms (30 x 7 cm) and two enclosed arms (30 x 7 cm) with 20 cm-high walls stand 50 cm high above a floor, each extending 7 cm away from a center platform. In the present test, a mouse was placed in the elevated plus maze in a room, where a video camera of 20 Lux was installed high up there, and then a movement of the mouse was recorded.

Particularly, a C57BL/6 mouse (male, 19-22 g) was placed in the middle of the elevated plus maze, in such a way that a head of the mouse faced toward an open arm. Time spent in open and closed arms as well as the number of arm entries were measured for five minutes. An arm entry was defined as a case in which the mouse entered an arm with all four feet.

Out of a whole test time, the time spent in open arms (OT) was calculated via $[\text{Time spent in open arms (OT)} / (\text{Time spent in open arms (OT)} + \text{Time spent in closed arms})] \times 100$. And open arm entries (OE) were calculated via $[\text{Open arm entries (OE)} / (\text{Open arm entries (OE)} + \text{Closed arm entries})] \times 100$. After the end of each behavioral experiment, remaining smell was eliminated by means of 70% ethanol.

According to a known interpretation of test results, it was interpreted that a mental disorder symptom such as anxiety, depression or the like occurs, if OT and OE decrease.

(2) Light/dark transition test

An apparatus used for a light/dark transition test consisted of a bright area (21 x 42 x 25 cm, 390 Lux white diode) and a dark area (21 x 42 x 25 cm, 2 Lux), which were of a size and partitioned off with a door (7.5 x 7.5 cm). A mouse was placed in a bright area and observed for five minutes, during which time in bright area and the number of transitions into the bright area were measured. After the end of each behavioral experiment, remaining smell was eliminated by means of 70% ethanol.

According to a known interpretation of test results, it was interpreted that a mental disorder symptom such as anxiety, depression or the like occurs, if the time in bright area and the number of transitions into the bright area decrease.

(3) Marble burying test

A mouse was placed in a transparent cage filled with sawdust 5 cm deep, and stayed there for 15 minutes. Then, 25 marbles (transparent, 2 cm in diameter) were placed on the sawdust at an interval of 5 cm. The mouse was placed again in the middle of one side and the number of marbles that the mouse buried for 30 minutes was measured.

According to a known interpretation of test results, it was interpreted that a mental disorder symptom such as anxiety, depression or the like occurs, if the mouse's behavior of burying and hiding marbles increases.

(4) Forced swimming test (FST)

According to a method of Porsolt et al. (Porsolt RD, Le Pichon, Jalfre M (1977) Depression: A new animal model sensitive to antidepressant treatments, *Nature*, Vol. 266; pp. 730-732), water at $25\pm 1^{\circ}\text{C}$ was filled into a water tank 40 cm in height and 20 cm in diameter, until a water level reached 25 cm high. A laboratory mouse was put into each tank and stayed there for total six minutes, out of which first two minutes were not measured as an adaptive time, and then for the last four minutes, an immobility time of the laboratory animal was measured. The immobility means a state that the mouse is floating in the water motionlessly except the least movement to keep its head only above the water.

(5) Tail suspension test (TST)

According to a method of Steru et al. (Steru L, Chermat R, Thierry B, Simon P (1985) The tail suspension test: A new method for screening antidepressants in mice,

Psychopharmacology, Vol. 85; pp. 367-370), a mouse was hanged up in a container 35 cm in diameter and 50 cm in height by placing a fixing device approximately 1 cm from the tip of a tail thereof of a mouse, such that the mouse was suspended 50 cm above a floor. For six minutes in total, an immobility time of the laboratory animal was measured.

Experimental Example 3. Experiment on identification of an intestinal microbial community

A real-time PCR or 454 pyrosequencing was performed to measure the shares of firmicutes, proteobacteria, actinobacteria, bacteroidetes and the like in an intestinal microbial community.

Particularly, DNA of feces obtained from an animal model was isolated by means of QIAamp DNA stool mini kit (Qiagen, Germany). Pyrosequencing was performed by means of a barcoded primer (V1 to V3 domains of a bacterial 16S rDNA gene). qPCR was analyzed by means of a primer shown in a following table 1.

【Table 1】

Type	Sequence (5'-3')	
<i>Firmicutes</i>	Forward (F)	(SEQ ID NO: 2) GGA GYA TGT GGT TTA ATT CGA AGC A
	Reverse (R)	(SEQ ID NO: 3) AGC TGA CGA CAA CCA TGC AC
<i>Bacteroidetes</i>	Forward (F)	(SEQ ID NO: 4) GTT TAA TTC GAT GAT ACG CGA G
	Reverse (R)	(SEQ ID NO: 5) TTA ASC CGA CAC CTC ACG G
<i>β-Proteobacteria</i>	Forward (F)	(SEQ ID NO: 6) AAC GCG AAA AAC CTT ACC TAC C
	Reverse (R)	(SEQ ID NO: 7) TGC CCT TTC GTA GCA ACT AGT G
<i>δ/γ-Proteobacteria</i>	Forward (F)	(SEQ ID NO: 8) GCT AAC GCA TTA AGT RYC CCG
	Reverse (R)	(SEQ ID NO: 9) GCC ATG CRG CAC CTG TCT
	Forward (F)	(SEQ ID NO: 10) TAG GCT TGA CAT TGA TAG AAT C

2021282468 08 Dec 2021

<i>ε-Proteobacteria</i>	Reverse (R)	(SEQ ID NO: 11) CTT ACG AAG GCA GTC TCC TTA
<i>Actinobacteria</i>	Forward (F)	(SEQ ID NO: 12) TGT AGC GGT GGA ATG CGC
	Reverse (R)	(SEQ ID NO: 13) AAT TAA GCC ACA TGC TCC GCT
<i>TM</i>	Forward (F)	(SEQ ID NO: 14) AYT GGG CGT AAA GAG TTG C
	Reverse (R)	(SEQ ID NO: 15) TAC GGY TAC CTT GTT ACG ACT T
<i>Verrucomicrobia</i>	Forward (F)	(SEQ ID NO: 16) TCA KGT CAG TAT GGC CCT TAT
	Reverse (R)	(SEQ ID NO: 17) CAG TTT TYA GGA TTT CCT CCG CC
<i>Enterobacteriaceae</i>	Forward (F)	(SEQ ID NO: 18) CAT TGA CGT TAC CCG CAG AAG AAG C
	Reverse (R)	(SEQ ID NO: 19) CTC TAC GAG ACT CAA GCT TGC
<i>Escherichia coli</i>	Forward (F)	(SEQ ID NO: 20) CGC GTA CTA TAC GCC ATG AAC GTA
	Reverse (R)	(SEQ ID NO: 21) ACC GTT GAT CAC TTC GGT CAG G
<i>Klebsiella spp.</i>	Forward (F)	(SEQ ID NO: 22) GAT ACG GAG TAT GCC TTT ACG GTG
	Reverse (R)	(SEQ ID NO: 23) TAG CCT TTA TCA AGC GGA TAC TGG
<i>Klebsiella oxytoca</i>	Forward (F)	(SEQ ID NO: 24) GTT AAT ACC TTT GCT CAT TGA
	Reverse (R)	(SEQ ID NO: 25) ACC AGG GTA TCT AAT CCT GTT
<i>Morganella morganii</i>	Forward (F)	(SEQ ID NO: 26) CTC GCA CCA TCA GAT GAA CCC ATA T
	Reverse (R)	(SEQ ID NO: 27) CAA AGC ATC TCT GCT AAG TTC TCT GGA TG
<i>Lactobacillus reuteri</i>	Forward (F)	(SEQ ID NO: 28) GAA CGC AYT GGC CCA A
	Reverse (R)	(SEQ ID NO: 29) TCC ATT GTG GCC GAT CAG T
<i>Lactobacillus johnsonii</i>	Forward (F)	(SEQ ID NO: 30) CAC TAG ACG CAT GTC TAG AG
	Reverse (R)	(SEQ ID NO: 31) AGT CTC TCA ACT CGG CTA TG
<i>Lactobacillus plantarum</i>	Forward (F)	(SEQ ID NO: 32) TCA TGA TTT ACA TTT GAG TG
	Reverse (R)	(SEQ ID NO: 33) GAC CAT GCG GTC CAA GTT GTT
<i>Lactobacillus rhamnosus</i>	Forward (F)	(SEQ ID NO: 34) CGC CCT TAA CAG CAG TCT TC
	Reverse (R)	(SEQ ID NO: 35) GCC CTC CGT ATG CTT AAA CC
<i>Bacterial 16S rDNA</i>	Forward (F)	(SEQ ID NO: 36) TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG GTG CCA GCM GCC GCG GTA A

	Reverse (R)	(SEQ ID NO: 37) GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GGG ACT ACH VGG GTW TCT AAT
--	-------------	--

Experimental Example 4. Method for measuring lipopolysaccharide

(1) Method for measuring lipopolysaccharide in feces

Feces (20 mg) were suspended in 30 mL of PBS, then microorganisms thereof were crushed by means of ultrasonic treatment for one hour, and then centrifuged at 500 rpm for 15 minutes. A resulting supernatant was filtered with a 0.45 μm filter, then filtered again with a 0.22 μm filter, then treated at 70°C for 10 minutes, and then used as a specimen. For the said specimen, lipopolysaccharide was measured by means of Limulus amoebocyte lysate (LAL) assay kit (Cape Cod Inc., East Falmouth, MA, U.S.A.).

(2) Method for measuring lipopolysaccharide in blood

Blood was diluted 10 times with PBS, then centrifuged, and then a resulting supernatant was treated at 0°C for 10 minutes. A resulting supernatant was filtered with a 0.45 μm filter, then filtered again with a 0.22 μm filter, and then used as a specimen. For the said specimen, lipopolysaccharide was measured by means of Limulus amoebocyte lysate (LAL) assay kit (Cape Cod Inc., East Falmouth, MA, U.S.A.).

Experimental Example 5. Method for measuring colitis indicators

(1) Measurement of myeloperoxidase activity

200 μL of 10 mM potassium phosphate buffer, pH 7.0 containing 0.5% hexadecyl trimethyl ammonium bromide was put into 100 mg of a colon tissue, and then subjected to homogenization. A resulting mixture was centrifuged for 10 minutes

on condition of 4°C and 10,000 g, so as to obtain a supernatant. 50 μl of the supernatant was put into 0.95 ml of a reaction solution (containing 1.6 mM tetramethyl benzidine and 0.1 mM H_2O_2), then subjected to reaction at 37°C, and then an observance was microscopically measured at 650 nm. An activity of the said myeloperoxidase (MPO) was calculated with a resulting reactant H_2O_2 1 $\mu\text{mol}/\text{ml} = 1$ unit.

(2) Measurement of inflammatory indicators

Inflammatory reaction biomarkers such as p-p65, p65, iNOS, COX-2 and β -actin were measured by means of a western blotting method. Particularly, 50 μg of supernatant, which had been obtained by the same method as shown in the said experiment for measuring the activity of myeloperoxidase (MPO), was taken and then immune blotting was performed thereon. Also, an expression level of cytokines was measured by means of ELISA kit, and LPS was measured by means of LAL assay kit.

Example 1: Identification of a state of an animal model with immobilization stress induced

According to Experimental Examples 2 to 5, an experiment was performed on a mouse (IS), which had been subjected to immobilization stress once every other day for five times in total by the same method as shown in the Experimental Example 1-(1) above.

As a result of performing Experimental Example 2, it was identified that time spent in open arms (OT) and open arm entries (OE) are decreased in an elevated plus maze test; time in bright area is decreased in a light/dark transition test; and a marble-burying behavior is increased in a marble burying test (Figs. 1 (a) to (c)).

Also, as a result of identifying with western blotting, it was identified that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derived neurotrophic factor (BDNF) is decreased (Fig. 1 (d)); and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide in blood are increased (Figs. 1 (e) to (h)).

As a result of performing Experimental Example 3, it was identified that bacteroidetes, actinobacteria and firmicutes in feces are decreased and δ , γ -proteobacteria and ε -proteobacteria are increased. As a result of performing Experimental Example 4, it was identified that lipopolysaccharide in feces is increased (Fig. 2).

As a result of performing Experimental Example 5, it was identified that as indicators of colitis, a length of a colon is decreased; myeloperoxidase is increased; TNF- α of the colon is increased; expressions of COX-2 and iNOS of the colon are increased; an NF- κ B activity is increased; and occludin and claudin-1, tight junction proteins, are decreased (Figs. 3 (a) to (e)).

Example 2: Identification of a state of an animal model administered with feces of the animal model with immobilization stress induced

According to Experimental Examples 2 to 4, an experiment was performed on a mouse (FIS), which was administered with feces of a mouse, which had been subjected to immobilization stress once every other day for five times in total by the same method as shown in the Experimental Example 1-(1) above.

As a result of performing Experimental Example 2, it was identified that OT and OE are decreased in an elevated plus maze test; time in bright area is decreased in a light/dark transition test; and a marble-burying behavior is increased in a marble

burying test (Figs. 4 (a) to (c)).

As a result of identifying with western blotting, it was identified that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derivated neurotrophic factor (BDNF) is decreased (Fig. 4 (d)); and amounts of corticosterone, IL-6 and TNF- α in blood are increased (Figs. 5 (a) to (c)).

As a result of performing Experimental Example 5, it was identified that as indicators of colitis, a length of a colon is decreased; myeloperoxidase is increased (Figs. 4 (e) and (f); TNF- α and IL-6 of the colon are increased; IL-10 of the colon is decreased; expressions of COX-2 and iNOS of the colon are increased; and an NF- κ B activity is increased (Figs. 6 (a) to (d)).

As a result of performing Experimental Example 3, it was identified that bacteroidetes, actinobacteria and firmicutes in feces are decreased and γ -proteobacteria and ϵ -proteobacteria are increased (Fig. 6 (e)).

Example 3: Identification of a state of an animal model with antibiotic stress induced

According to Experimental Examples 2 to 5, an experiment was performed on a mouse (AP), which had been subjected to antibiotic stress with an administration of ampicillin (100 mg/kg) for consecutive two days by the same method as shown in the Experimental Example 1-(2) above.

As a result of performing Experimental Example 2, it was identified that OT and OE are decreased in an elevated plus maze test; time in bright area is decreased in a light/dark transition test; and a marble-burying behavior is increased in a marble burying test (Figs. 7 (a) to (c)).

It was identified that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derivated neurotrophic factor (BDNF) is

decreased (Fig. 7 (d)); and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide in blood are increased (Figs. 7 (e) to (h)).

As a result of performing Experimental Example 3, it was identified that bacteroidetes, actinobacteria and firmicutes in feces are decreased and δ , γ -proteobacteria and ϵ -proteobacteria are increased. As a result of performing Experimental Example 4, it was identified that lipopolysaccharide in feces is increased (Figs. 8 (a) and (b)).

As a result of performing Experimental Example 5, it was identified that as indicators of colitis, a length of a colon is decreased; myeloperoxidase is increased; TNF- α of the colon is increased; expressions of COX-2 and iNOS of the colon are increased; an NF- κ B activity is increased; and occludin and claudin-1, tight junction proteins, are decreased (Figs. 9 (a) to (e)).

Example 4: Identification of a state of an animal model administered with microorganisms isolated from feces of the animal model with stress induced

(1) Isolation of microorganisms from feces of the animal model with stress induced

Feces of a mouse subjected to immobilization stress and feces of a mouse subjected to antibiotic stress were cultured in a selective medium, respectively. As a result, it was identified that enterobacteriaceae grown in a DHL medium is increased, and in particular *Klebsiella oxytoca*, *Escherichia coli* and *Morganella morganii* are increased (Fig. 10 (A)).

On the other hand, it was identified that bifidobacteria and lactobacilli grown in a BL medium are decreased, and in particular *Lactobacillus reuteri*, *Lactobacillus johnsonii*, *Lactobacillus rhamnosus* and *Bifidobacterium animalis* are decreased (Fig.

10 (B)).

(2) Identification of a state of an animal model administered with microorganisms

After administering to each mouse 1×10^9 cfu of *Klebsiella oxytoca*, *Escherichia coli* and *Morganella morganii* respectively, which had been increased in feces of the mouse subjected to immobilization stress and the mouse subjected to antibiotic stress, an elevated plus maze test was performed according to Experimental Example 2-(1) above. As a result, it was identified that OT and OE are remarkably decreased in a group administered with *Klebsiella oxytoca* compared to a group administered with other microorganisms (Fig. 11 (a)).

After that, according to Experimental Examples 2 to 5, an experiment was performed on a mouse (KO) administered with *Klebsiella oxytoca*.

As a result of performing Experimental Example 2, it was identified that OT and OE are decreased in an elevated plus maze test; time in bright area is decreased in a light/dark transition test; and a marble-burying behavior is increased in a marble burying test (Figs. 11 (b) to (d)).

It was identified that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derived neurotrophic factor (BDNF) is decreased (Fig. 11 (e)); and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide in blood are increased (Figs. 11 (f) to (i)).

As a result of performing Experimental Example 3, it was identified that bacteroidetes, actinobacteria and firmicutes in feces are decreased and δ , γ -proteobacteria and ϵ -proteobacteria in feces are increased. As a result of performing Experimental Example 4, it was identified that lipopolysaccharide in feces is increased (Figs. 12 (a) and (b)).

As a result of performing Experimental Example 5, it was identified that as indicators of colitis, a length of a colon is decreased; myeloperoxidase is increased; TNF- α of the colon is increased; expressions of COX-2 and iNOS of the colon are increased; an NF- κ B activity is increased; and occludin and claudin-1, tight junction proteins, are decreased (Figs. 13 (a) to (e)).

Example 5: Isolation and identification of lactic acid bacteria

(1) Isolation of lactic acid bacteria from human and mouse feces

Fresh human and mouse feces were inserted and suspended in a GAM liquid medium (GAM broth; Nissui Pharmaceutical, Japan), respectively. After that, supernatant was taken and implanted to MRS, BHI (Brain-Heart-Infusion) or BL agar medium (Nissui Pharmaceutical, Japan), then anaerobically cultured at 37°C for about 48 to 72 hours, and then strains forming a colony were isolated therefrom.

(2) Identification of isolated lactic acid bacteria

Physiological characteristics and 16S rDNA sequences of strains isolated from human or mouse feces were analyzed, then species of the strains were identified, and then names were given to the strains. Strain names given to lactic acid bacteria are as shown in a following table 2. Particularly, lactic acid bacteria isolated from human or mouse feces were 9 types of *Lactobacillus* spp. and 22 types of *Bifidobacterium* spp.

【Table 2】

Management no.	Strain name
1	<i>Lactobacillus reuteri</i> NK31
2	<i>Lactobacillus reuteri</i> NK32
3	<i>Lactobacillus reuteri</i> NK33
4	<i>Lactobacillus reuteri</i> NK34
5	<i>Lactobacillus reuteri</i> NK35
6	<i>Lactobacillus johnsonii</i> NK36
7	<i>Lactobacillus johnsonii</i> NK37

8	<i>Lactobacillus johnsonii</i> NK38
9	<i>Lactobacillus johnsonii</i> NK39
10	<i>Bifidobacterium longum</i> NK81
11	<i>Bifidobacterium longum</i> NK82
12	<i>Bifidobacterium longum</i> NK83
13	<i>Bifidobacterium longum</i> NK84
14	<i>Bifidobacterium bifidum</i> NK85
15	<i>Bifidobacterium bifidum</i> NK86
16	<i>Bifidobacterium infantis</i> NK87
17	<i>Bifidobacterium infantis</i> NK88
18	<i>Bifidobacterium infantis</i> NK89
19	<i>Bifidobacterium breve</i> NK90
20	<i>Bifidobacterium breve</i> NK91
21	<i>Bifidobacterium choerinum</i> NK92
22	<i>Bifidobacterium catenulatum</i> NK93
23	<i>Bifidobacterium catenulatum</i> NK94
24	<i>Bifidobacterium adolescentis</i> NK95
25	<i>Bifidobacterium pseudolongum</i> NK96
26	<i>Bifidobacterium adolescentis</i> NK97
27	<i>Bifidobacterium adolescentis</i> NK98
28	<i>Bifidobacterium adolescentis</i> NK99
29	<i>Bifidobacterium pseudocatenulatum</i> NK100
30	<i>Bifidobacterium pseudocatenulatum</i> NK101
31	<i>Bifidobacterium minimum</i> NK102

(3) Physiological characteristics of *Lactobacillus reuteri* NK33, a novel lactic acid bacterium

Out of strains described in Table 2 above, it was identified that *Lactobacillus reuteri* NK33 (accession number KCCM12090P) is a gram-positive bacillus. Also, it was shown that a 16S rDNA of *Lactobacillus reuteri* NK33 has a sequence of SEQ ID NO: 1. As a result of comparing the 16S rDNA sequences of *Lactobacillus reuteri* NK33 by means of BLAST search, it was identified that a *Lactobacillus reuteri* strain having the same 16S rDNA sequence is not searched at all, and 99% homologous to the 16S rDNA sequence of a known *Lactobacillus reuteri* strain.

Out of physiological characteristics of *Lactobacillus reuteri* NK33, availability

of carbon source was analyzed with a sugar fermentation test using API 50 CHL kit (BioMerieux's, USA). The results thereof are as shown in a following table 3. In Table 3 below, “+” indicates that availability of carbon source is positive and “-” indicates that availability of carbon source is negative.

【Table 3】

Carbon source	<i>L. reuteri</i> NK33	Carbon source	<i>L. reuteri</i> NK33
CONTROL	-	Esculin	+
Glycerol	-	Salicin	+
Erythritol	-	Cellobiose	+
D-arabinose	-	Maltose	+
L-arabinose	+	Lactose	+
D-ribose	+	Melibiose	+
D-xylose	+	Sucrose	+
L-xylose	-	Trehalose	+
D-adonitol	-	Inulin	-
Methyl- β -D-xylopyranoside	-	Melezitose	+
D-galactose	+	Raffinose	+
D-glucose	+	Starch	-
D-fructose	+	Glycogen	-
D-mannose	+	Xylitol	-
L-sorbose	-	Gentiobiose	+
Rhamnosus	-	D-turanose	+
Dulcitol	-	D-lixose	-
Inocitol	-	D-tagatose	-
Mannitol	+	D-fucose	-
Sorbitol	$\pm$	L-fucose	-
α -methyl-D-mannoside	-	D-arabitol	-
α -methyl-D-glucoside	-	L-arabitol	-
N-acetyl-glucosamine	+	Gluconate	$\pm$
Amygdalin	+	2-keto-gluconate	-
Arbutin	+	5-keto-gluconate	-

(4) Physiological characteristics of *Bifidobacterium adolescentis* NK98, a novel lactic acid bacterium

Out of strains described in Table 2 above, it was shown that a 16S rDNA of

Bifidobacterium adolescentis NK98 (accession number KCCM12297P) has a sequence of SEQ ID NO: 38. As a result of comparing the 16S rDNA sequences of *Bifidobacterium adolescentis* NK98 by means of BLAST search, it was identified that a *Bifidobacterium adolescentis* NK98 strain having the same 16S rDNA sequence is not searched at all, and 98% homologous to the 16S rDNA sequence of a known *Bifidobacterium adolescentis* strain.

Out of physiological characteristics of *Bifidobacterium adolescentis* NK98, availability of carbon source was analyzed with a sugar fermentation test using API kit (API 20A, BioMerieux's, USA). The results thereof are as shown in a following table 4. In Table 4 below, “+” indicates that availability of carbon source is positive and “-” indicates that availability of carbon source is negative.

【Table 4】

Carbon source	Reaction/Enzyme	<i>B. adolescentis</i> NK98
L-tryptophan	Indole formation	-
Urea	Urease	-
D-glucose	Acidification (glucose)	+
D-mannitol	Acidification (mannitol)	+
D-lactose	Acidification (lactose)	+
D-saccharose	Acidification (saccharose)	+
D-maltose	Acidification (maltose)	+
Salicin	Acidification (salicin)	+
D-xylose	Acidification (xylose)	±
L-arabinose	Acidification (arabinose)	+
Gelatin	Hydrolysis (protease)	+
Esculin ferric citrate	Hydrolysis (β -glucosidase)	+
Glycerol	Acidification (glycerol)	-

D-cellobiose	Acidification (cellobiose)	±
D-mannose	Acidification (mannose)	-
D-melezitose	Acidification (melezitose)	-
D-raffinose	Acidification (raffinose)	+
D-sorbitol	Acidification (sorbitol)	-
L-rhamnose	Acidification (rhamnose)	-
D-trehalose	Acidification (trehalose)	+
	Catalase	-
	Spores	-
	Gram reaction	+
	Morphology	-

Example 7: Comparison of activity of isolated lactic acid bacteria

(1) Antioxidant activity (in vitro)

DPPH (2,2-Diphenyl-1-picrylhydrazyl) was dissolved in ethanol to reach a 0.2 mM concentration, such that a DPPH solution was prepared. A suspension of lactic acid bacteria (1×10^8 CFU/mL) or a vitamin C solution (1 g/mL) was inserted into 0.1 mL of the said DPPH solution, and then cultured at 37°C for 20 minutes. A culture fluid was centrifuged at 3000 rpm for five minutes, such that supernatant was obtained. After that, an absorbance of the supernatant was measured at 517 nm, and then antioxidant activity of isolated lactic acid bacteria was calculated accordingly. Antioxidant activity of each lactic acid bacterium is as shown in following table 5 (*Lactobacillus* spp.) and table 6 (*Bifidobacterium* spp.).

(2) Measurement of inflammatory indicators in macrophage

2 mL of sterilized 4% thioglycolate was intraperitoneally administered into a

C57BL/6 mouse (male, 6-week old, 20-23 g). In 96 hours later, the mouse was anesthetized, and then 8 ml of RPMI 1640 medium was intraperitoneally administered to the mouse. In 5 to 10 minutes later, an RPMI medium (macrophage) was intraperitoneally extracted from the mouse, then centrifuged at 1000 g for 10 minutes, and then washed twice again with an RPMI 1640 medium. The said macrophage was spread into a 24-well plate at 0.5×10^6 cells per well, then treated with isolated *Lactobacillus* spp. lactic acid bacteria and lipopolysaccharide, an inflammatory reaction inducer for 2 or 24 hours, and then supernatant and cells were obtained. The obtained cells were inserted into an RIPA buffer (Gibco) and homogenized. An expression level of cytokines such as TNF- α was measured from a culture supernatant treated for 24 hours, and then expression levels of p65 (NF- κ B), p-p65 (phosphor-NF- κ B) and β -actin were measured from the cells obtained with treatment for two hours by means of an immunoblotting method. Expression levels of inflammatory indicators for each *Lactobacillus* spp. lactic acid bacterium are as shown in a following table 5.

(3) Measurement of inflammatory indicators in microglia

A BV-2 microglia was spread into a 24-well plate at 0.5×10^6 cells per well, then treated with isolated *Bifidobacterium* spp. lactic acid bacteria and lipopolysaccharide, an inflammatory reaction inducer, for 2 or 24 hours, and then supernatant and cells were obtained. The obtained cells were inserted into an RIPA buffer (Gibco) and homogenized. Expression levels of p65 (NF- κ B), p-p65 (phosphor-NF- κ B) and β -actin were measured from the cells obtained with treatment for two hours by means of an immunoblotting method. Expression levels of inflammatory indicators for each *Bifidobacterium* spp. lactic acid bacterium are as shown in a

following table 6.

(4) Effect of brain derivated neurotrophic factor (BDNF) expression and NF- κ B activation on SH-SY5Y cells

SH-SY5Y cells, which are a nerve cell, were purchased from Korean Cell Line Bank, then cultured in a DMEM medium containing 10% FBS and 1% antibiotics, and then divided into a 12-well plate at 2×10^6 cells per well. After that, corticosterone was added into each well at a concentration of 300 mg/ml along with lactic acid bacteria (1×10^4 CFU/ml), then cultured, and then expression levels of NF- κ B (p65, p-p65) and brain derivated neurotrophic factor (BDNF) were measured by means of an immunoblotting method. The BDNF expression levels and NF- κ B activation levels for each lactic acid bacterium are as shown in following table 5 (*Lactobacillus* spp.) and table 6 (*Bifidobacterium* spp.).

【Table 5】

Strain name	Antioxidant activity	Macrophage			SH-SY5Y cell	
		TNF- α inhibitory capacity	IL-10 expression increase	NF- κ B inhibitory capacity	BDNF Expression increase	NF- κ B inhibition
<i>L. reuteri</i> NK31	+	+	+	+	+	+
<i>L. reuteri</i> NK32	++	++	++	++	+	++
<i>L. reuteri</i> NK33	+++	++	++	++	++	++
<i>L. reuteri</i> NK34	+	+	+	+	+	+
<i>L. reuteri</i> NK35	+	+	+	+	+	+
<i>L. johnsonii</i> NK36	++	++	++	+	++	+
<i>L. johnsonii</i> NK37	++	++	++	++	+	++

<i>L. johnsonii</i> NK38	+	+	+	+	+	+
<i>L. johnsonii</i> NK39	+	+	+	+	-	+

【Table 6】

Strain name	Antioxidant activity	Microglia NF-κB inhibitory capacity	SH-SY5Y cell BDNF expression capacity
Untreated	-	-	-
<i>L. reuteri</i> NK33	+++	++	++
<i>B. longum</i> NK81	+	+	+
<i>B. longum</i> NK82	+	+	+
<i>B. longum</i> NK83	++	+	+
<i>B. longum</i> NK84	++	++	+
<i>B. bifidum</i> NK85	++	++	+
<i>B. bifidum</i> NK86	+	-	+
<i>B. infantis</i> NK87	++	+	+
<i>B. infantis</i> NK88	+	+	++
<i>B. infantis</i> NK89	+	+	+
<i>B. breve</i> NK90	+	+	++
<i>B. breve</i> NK91	++	+	+
<i>B. choerinum</i> NK92	+	+	+
<i>B. catenulatum</i> NK93	+	+	+
<i>B. catenulatum</i> NK94	+	+	+
<i>B. adolescentis</i> NK95	++	+	+
<i>B. pseudolongum</i> NK96	++	+	+
<i>B. adolescentis</i> NK97	+	+	+

<i>B. adolescentis</i> NK98	++	+++	+++
<i>B. adolescentis</i> NK99	+	++	++
<i>B. pseudocatenulatum</i> NK100	++	+	++
<i>B. pseudocatenulatum</i> NK101	++	+	+
<i>B. minimum</i> NK102	+	+	+

* -, <10%; +, 10~30%; ++, 30~50%; +++, >50%

(5) Experimental results

As a result of evaluating the activity of isolated lactic acid bacteria, it was identified that out of isolated *Lactobacillus* spp. or *Bifidobacterium* spp. lactic acid bacteria, *Lactobacillus reuteri* NK33 and *Bifidobacterium adolescentis* NK98 herein have a remarkably excellent effect on promoting antioxidant activity and inhibiting inflammatory reactions. In particular, it was identified that *Lactobacillus reuteri* NK33 and *Bifidobacterium adolescentis* NK98 inhibit the activity of NF- κ B, known as a substance of inducing aging-related diseases such as Alzheimer's disease, inhibit an inflammatory reaction of microglia present in aging, dementia and the like, and increase an expression of a brain derived neurotrophic factor (BDNF) which is produced by a brain nerve, but decreased in aging, dementia and the like (Tables 5 and 6).

Example 8: Evaluation of an immunoregulatory effect of isolated *Bifidobacterium* spp. lactic acid bacteria

To evaluate an immunoregulatory effect of *Bifidobacterium* spp. lactic acid bacteria isolated from feces, the effect of *Bifidobacterium* spp. lactic acid bacteria on an immuno reaction of macrophage and spleen cells was measured.

(1) Immuno reaction in macrophage

2 ml of sterilized 4% thioglycolate was intraperitoneally administered into a

C57BL/6 mouse (male, 6-week old, 20-23 g, Raonbio Co., Ltd.). In 96 hours after administration, the mouse was anesthetized, and then 8 mL of RPMI 1640 medium was intraperitoneally administered to the mouse. In 5 to 10 minutes later, an RPMI medium (including macrophage) was intraperitoneally extracted from the mouse, then centrifuged at 1000 rpm for 10 minutes, and then washed twice again with an RPMI 1640 medium. The said macrophage was spread over a 24-well plate at 0.5×10^6 cells per well, then cultured for 24 hours, and then unattached cells were removed and used.

The said macrophage culture fluid was treated with *Bifidobacterium adolescentis* NK98 and lipopolysaccharide, an inflammatory reaction inducer, for 2 or 24 hours, and then supernatant and cells were obtained, wherein a concentration for treating lactic acid bacteria was 1×10^4 CFU/mL. The obtained cells were inserted into an RIPA buffer (Gibco) and homogenized. An expression level of TNF- α from the obtained supernatant was measured by means of ELISA kit, and then expression levels of p65 (NF- κ B), p-p65 (phosphor-NF- κ B) and β -actin from the obtained cells were measured by means of an immunoblotting method. Particularly, 50 μ g of the supernatant was taken and subjected to electrophoresis in SDS 10%(w/v) polyacrylamide gel for one and half an hour. A sample subjected to electrophoresis was transferred onto a nitrocellulose paper on condition of 100 V and 400 mA for one hour and 10 minutes. The nitrocellulose paper, onto which the sample was transferred, was subjected to blocking by means of 5% skimmed milk for 30 minutes, then washed with PBS-Tween three times each for five minutes, and then subjected to reaction overnight with primary antibodies (Santa Cruz Biotechnology, the U.S.) at a ratio of 1:100. After

that, such paper was washed three times each for 10 minutes, and subjected to reaction with secondary antibodies (Santa Cruz Biotechnology, the U.S.) at a ratio of 1:1000 for one hour and 20 minutes. Then, such paper was washed three times each for 15 minutes, then subjected to fluorescence and luminescence, then developed, and then intensity of a chromophore band was measured, wherein the results thereof are as shown in a following table 7.

【Table 7】

Strain name	NF- κ B inhibitory capacity (p-p65/p65)	TNF- α inhibitory capacity
Untreated	-	-
<i>B. adolescentis</i> NK98	+	++

* Inhibition rate: -, <10%; +, 10~30%; ++, 30~60%; +++, >60

(2) Immuno reaction in spleen cells

A spleen of a C57BL/6 mouse (male, 6-week old, 20-22 g, OrientBio Co., Ltd.) was isolated and crushed, and then suspended in an RPMI 1640 medium containing 10% FCS. CD4 T cells were isolated therefrom by means of CD4 T cell isolation kit (MiltenyiBiotec, Bergisch Gladbach, Germany), and then the isolated CD4 T cells were divided into a 12-well plate at 5×10^5 cells per well.

The cells were cultured with an addition of anti-CD3, anti-CD28, IL-2 and IL-12 to induce differentiation of T cells into Th1 cells; anti-CD3, anti-CD28, IL-2 and IL-4 to induce differentiation of T cells into Th2 cells; anti-CD3, anti-CD28, IL-6 and TGF- β to induce differentiation of T cells into Th17 cells; and anti-CD3 and anti-CD28 to induce differentiation of T cells into Treg cells, then lactic acid bacteria were inserted thereinto at 1×10^5 CFU/ml per well, and then cultured for four days.

After that, a differentiation potency of T cells isolated from the spleen into Th1 cells, Th2 cells, Th17 cells and Treg cells was measured. Particularly, cells of culture fluid were stained with anti-FoxP3 or anti-IL-17A antibodies, and then a distribution of Th1 cells, Th2 cells, Th17 cells and Treg cells was analyzed by means of FACS (fluorescence-activated cell sorting) device (C6 Flow Cytometer System, San Jose, CA, USA), wherein the results thereof are as shown in a following table 8.

【Table 8】

Strain name	Differentiation inhibition rate			Differentiation increase rate
	Th1 cell	Th2 cell	Th17 cell	Treg cell
Untreated	-	-	-	-
<i>B. longum</i> NK81	+	++	+	+
<i>B. longum</i> NK82	+	++	+	+
<i>B. longum</i> NK83	+	++	++	+
<i>B. longum</i> NK84	+++	+	+++	++
<i>B. bifidum</i> NK85	+++	+	+++	++
<i>B. bifidum</i> NK86	+	+	+++	+
<i>B. infantis</i> NK87	+	+	+++	++
<i>B. infantis</i> NK88	+	+	+++	+
<i>B. infantis</i> NK89	+	+	+++	+
<i>B. breve</i> NK90	+	+++	++	++
<i>B. breve</i> NK91	++	++	+	+
<i>B. choerinum</i> NK92	-	+	+	-
<i>B. catenulatum</i> NK93	+	++	++	+
<i>B. catenulatum</i> NK94	++	+	+	++
<i>B. adolescentis</i> NK95	++	+	+	+

<i>B. pseudolongum</i> NK96	++	++	++	+
<i>B. adolescentis</i> NK97	+	++	+	+
<i>B. adolescentis</i> NK98	++	++	+++	+++
<i>B. adolescentis</i> NK99	+	+	++	+
<i>B. pseudocatenulatum</i> NK100	++	++	+	+
<i>B. pseudocatenulatum</i> NK101	+	++	+	+
<i>B. minimum</i> NK102	+	+	+	+

* Inhibition rate: -, <10%; +, 10-30%; ++, 30-60%; +++, >60

* Increase rate: -, <10%; +, 10-50%; ++, 50-100%; +++, >100%

Also, expression rates of transcription factors and cytokines of Th1 cells, Th2 cells, Th17 cells and Treg cells isolated from the spleen T cells were measured. Particularly, the expression levels of T-bet, IFN- γ and IL-12 from a Th1 cell differentiation-inducing culture fluid; GATA3 and IL-5 from a Th2 cell differentiation-inducing culture fluid; ROR γ t and IL-17 from a Th17 cell differentiation-inducing culture fluid; and Foxp3 and IL-10 from a Treg cell differentiation-inducing culture fluid were analyzed by means of qRT-PCR, wherein the results thereof are as shown in a following table 9.

【Table 9】

Strain name	Expression inhibition rate						Expression increase rate	
	T-bet	IFN- γ	GATA3	IL-5 cell	ROR γ t	IL-17	FOXp3	IL-10
Untreated	-	-	-	-	-	-	-	-
<i>B. longum</i> NK81	++	+	+	+	+	+	-	+
<i>B. longum</i> NK82	+	+	+	+	+	+	+	+
<i>B. longum</i> NK83	+	+	+	+	+	+	+	+

<i>B. longum</i> NK84	++	++	+	+	+++	+++	++	++
<i>B. bifidum</i> NK85	+++	+++	+	+	+	+	++	++
<i>B. bifidum</i> NK86	-	-	+	+	+	+	+	+
<i>B. infantis</i> NK87	+	+	+	+	+	+	+	+
<i>B. infantis</i> NK88	+	+	++	++	+	+	+	+
<i>B. infantis</i> NK89	+	+	+	+	++	++	+	+
<i>B. breve</i> NK90	+	+	+++	+++	++	++	++	++
<i>B. breve</i> NK91	+	+	+	+	+	+++	++	+
<i>B. choerinum</i> NK92	+	+	+	+	+	++	+	++
<i>B. catenulatum</i> NK93	+	+	+	+	+	+	+	+
<i>B. catenulatum</i> NK94	++	+	+	+	+	+	+	+
<i>B. adolescentis</i> NK95	+	+	+	+	++	++	++	++
<i>B. pseudolongum</i> NK96	+	+	+	+	+	+	+	+
<i>B. adolescentis</i> NK97	+	+	+	+	+	+	+	+
<i>B. adolescentis</i> NK98	++	++	++	++	+++	+++	+++	+++
<i>B. adolescentis</i> NK99	+	++	+	+	+	+	+	+
<i>B. pseudocatenulatum</i> NK100	++	+	+	+	-	-	+	+
<i>B. pseudocatenulatum</i> NK101	++	+	+	+	-	-	+	+
<i>B. minimum</i> NK102	+	+	+	+	+	-	-	-

* Inhibition rate: -, <10%; +, 10-30%; ++, 30-60%; +++, >60

* Increase rate: -, <10%; +, 10-50%; ++, 50-100%; +++, >100%

As identified in Tables 8 and 9 above, it was identified that *Bifidobacterium adolescentis* NK98 has a high inhibition rate on the differentiation of T cells into Th1 cells, Th2 cells and Th17 cells and in particular has a high increase rate on the differentiation of T cells into Treg cells, thus effectively inhibiting an inflammatory reaction, and effectively improving an inflammatory disease accordingly.

Example 9: Effect of *Lactobacillus reuteri* NK33 on reducing stress

From Day 7 after starting a immobilization stress process with regard to a

mouse, which had been subjected to immobilization stress as shown in Experimental Example 1 above, 1×10^9 cfu of *Lactobacillus reuteri* NK33, a novel lactic acid bacterium, or physiological saline solution was administered once daily for five days to the mouse. According to Experimental Examples 2 to 5, an experiment was performed on the mouse above.

As a result of performing Experimental Example 2, it was identified in a group (IS) administered with physiological saline solution that OT and OE are decreased in an elevated plus maze test; time in bright area is decreased in a light/dark transition test; and a marble-burying behavior is increased in a marble burying test. However, it was identified in a group (IS+LR) administered with *Lactobacillus reuteri* NK33 that OT and OE are increased; time in bright area is increased; and a marble-burying behavior is decreased in a marble burying test (Figs. 14 (a) to (c)).

It was identified in the group (IS) administered with physiological saline solution that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derived neurotrophic factor (BDNF) is decreased; and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide in blood are increased. However, it was identified in the group (IS+LR) administered with *Lactobacillus reuteri* NK33 that an NF- κ B activity (p-p65/p65) is inhibited; an expression level of a brain derived neurotrophic factor (BDNF) is increased; and amounts of corticosterone, IL-6, TNF- α and lipopolysaccharide in blood are decreased (Figs. 14 (d) to (g)).

As a result of performing Experimental Example 5, it was identified in the group (IS) administered with physiological saline solution that as indicators of colitis, a length of a colon is decreased; myeloperoxidase is increased; TNF- α of the colon is increased; expressions of COX-2 and iNOS of the colon are increased; and an NF- κ B

activity is increased. However, it was identified in the group (IR) administered with *Lactobacillus reuteri* NK33 that a length of a colon is recovered to a normal level; myeloperoxidase is decreased; TNF- α of the colon is decreased; expressions of COX-2 and iNOS of the colon are decreased; and an NF- κ B activity is inhibited (Figs. 15 (a) to (d)).

From the results, it was identified that *Lactobacillus reuteri* NK33, the novel lactic acid bacterium, has an excellent effect of improving a mental disorder, such as anxiety, depression, stress and the like.

Example 10: Effect of *Bifidobacterium adolescentis* NK98 on reducing stress

From Day 7 after starting the immobilization stress process with regard to the mouse, which had been subjected to immobilization stress as shown in Experimental Example 1 above, 1×10^9 cfu of *Bifidobacterium adolescentis* NK98 (NK98), 1×10^9 cfu of *Bifidobacterium adolescentis* IM38 (IM38), physiological saline solution (IS) or 1 mg/kg of Buspirone was administered once daily for three days to a mouse. Then, according to Experimental Examples 2 and 5 above, an experiment was performed.

(1) Elevated plus maze test

Along with an elevated plus maze test of Experimental Example 2-(1) above, an amount of corticosterone (BC) in blood was measured, wherein the results thereof are as shown in a following table 10.

【Table 10】

	Anxious mouse			Depressive mouse		
	OT (%)	OE (%)	BC (ng/mL)	OT (%)	OE (%)	BC (ng/mL)
Normal control	17.1	35.2	0.8	18.2	38.5	0.7

IS	5.2	16.1	2.8	3.4	13.5	7.4
NK98	16.2	34.1	1.2	16.2	32.3	3.5
IM38	14.3	32.6	1.5	13.3	28.6	4.1
Buspirone	16.7	33.8	1.2	15.7	29.8	3.8

As identified in Table 10 above, it was identified in the group (IS) of immobilization stress models administered with physiological saline solution that both OT and OE are decreased in an elevated plus maze test compared to a normal control group without immobilization stress. However, it was identified in a group administered with *Bifidobacterium adolescentis* NK98 that time spent in open arms (OT) and frequency of open arm entries (OE) are increased; and an amount of corticosterone in blood is remarkably decreased, thus showing a more excellent effect compared to *Bifidobacterium adolescentis* IM38.

(2) Light/dark transition test

According to Experimental Example 2-(2) above, a light/dark transition test was performed, wherein the results thereof are as shown in a following table 11.

【Table 11】

	Time in bright area (%)	Number of transitions
Normal control	46.5	23.5
IS	23.1	13.8
NK98	39.4	21.9
IM38	35.8	18.3
Buspirone	40.1	19.6

As identified in Table 11 above, it was identified in a group (IS) of immobilization stress models administered with physiological saline solution that time in bright area is decreased compared to a normal control group without immobilization stress, but in a group administered with *Bifidobacterium adolescentis* NK98 that time in bright area is increased compared to a group administered with *Bifidobacterium adolescentis* IM38.

(3) Forced swimming test

According to Experimental Example 2-(4) above, a forced swimming test was

performed, wherein the results thereof are as shown in a following table 12.

【Table 12】

	Immobility time (%)
Normal control	28.5
IS	46.7
NK98	31.2
IM38	38.2
Buspirone	32.1

As identified in Table 12 above, it was identified in a group (IS) of depressed mouse models administered with physiological saline solution that an immobility time is increased compared to a normal control group without immobilization stress, and in a group administered with *Bifidobacterium adolescentis* NK98 that the immobility time is decreased compared to a group administered with *Bifidobacterium adolescentis* IM38.

(4) Tail suspension test (TST)

According to Experimental Example 2-(5) above, a tail suspension test (TST) was performed, wherein the results thereof are as shown in a following table 13.

【Table 13】

	Immobility time (%)
Normal control	21.5
IS	37.9
NK98	24.2
IM38	26.9
Buspirone	25.2

As identified in Table 13 above, it was identified in a group (IS) of depressed mouse models administered with physiological saline solution that an immobility time is increased compared to a normal control group without immobilization stress, but in a group administered with *Bifidobacterium adolescentis* NK98 that the immobility time is decreased.

(5) Measurement of biomarkers

In two hours after performing a last behavioral experiment on a depressed

mouse, the mouse was anesthetized, and corticosterone in blood was measured by means of ELISA, and a brain derived neurotrophic factor (BDNF) and NF- κ B of a brain and NF- κ B of a colon were measured by means of immunoblotting by the same method as shown in Experimental Example 5 above, wherein the results thereof are as shown in a following table 14.

【Table 14】

	NF- κ B activation inhibitory capacity	BDNF expression level	Blood corticosterone (ng/mL)
Normal control			0.7
IS	-	-	7.4
NK98	+++	+++	3.5
IM38	++	++	4.1
Buspirone	-	-	3.8

* -, <10%; +, 10~30%; ++, 30~50%; +++, >50%

As identified in Table 14 above, it was identified in a group (IS) administered with physiological saline solution that an NF- κ B activity (p-p65/p65) is increased in a hippocampus; an expression level of a brain derived neurotrophic factor (BDNF) is decreased; and an amount of corticosterone in blood is increased. However, it was identified in a group administered with NK98 that an NF- κ B activity is inhibited; an expression level of the BDNF is increased; and an amount of corticosterone in blood is decreased, wherein such effect is excellent compared to a group administered with IM38.

(6) Effect of improving colitis

In two hours after performing a last behavioral experiment on an anxious mouse in Experimental Example 1, the mouse was anesthetized, and a colon length and activations of MPO, COX-2, TNF- α and NF- κ B were measured by the same method as shown in Experimental Example 5, wherein the results thereof are as shown in a following table 15.

【Table 15】

	Colon length (cm)	MPO inhibitory capacity	Inflammatory indicator inhibition rate		
			NF- κ B	COX-2	TNF- α
Normal control	6.2				
IS	5.1	-	-	-	-
NK98	5.7	+++	+++	+++	+++
IM38	5.5	+++	++	++	++
Buspirone	5.2	+	+	+	+

* -, <10%; +, 10~30%; ++, 30~50%; +++, >50%

As identified in Table 15 above, it was identified in a group (IS) of immobilization stress models administered with physiological saline solution that a colon length is decreased and MPO and inflammatory indicators are increased compared to a normal control group without immobilization stress, but in a group administered with *Bifidobacterium adolescentis* NK98 that a colon length is recovered, and the MPO inhibitory capacity and inflammatory indicators are improved.

(7) Effect of reducing antibiotics-induced stress

According to Experimental Examples 2 to 5, an experiment was performed on a mouse, which had been subjected to antibiotic stress with an administration of ampicillin (100 mg/kg) for consecutive two days by the same method as shown in Experimental Example 1-(2) above, wherein the results thereof are as shown in following Tables 16 and 17.

【Table 16】

	Time spent in open arms (%)	Open arm entries (%)	Blood corticosterone (ng/mL)
Normal control	16.8	33.5	0.8
IS	4.9	13.4	3.2
NK98	14.3	30.5	1.6

IM38	12.9	29.2	1.9
Buspirone	14.1	31.3	1.6

【Table 17】

	Colon length (cm)	MPO inhibitory capacity	Inflammatory indicator inhibition rate		
			NF-κB	COX-2	TNF-α
Normal control	6.3				
IS	5.4	-	-	-	-
NK98	6.1	+++	+++	+++	+++
IM38	5.8	++	++	++	++
Buspirone	5.5	+	+	+	+

* -, <10%; +, 10~30%; ++, 30~50%; +++, >50%

As identified in Table 16 above, it was identified in a group (IS) of antibiotics-induced stress models administered with physiological saline solution that OT and OE are decreased compared to a normal control group in an elevated plus maze test, and in a group administered with *Bifidobacterium adolescentis* NK98 that OT and frequency of OE are increased and an amount of corticosterone in blood is remarkably decreased, wherein such effect is excellent compared to *Bifidobacterium adolescentis* IM38.

As identified in Table 17 above, it was identified in a group (IS) of antibiotics-induced stress models administered with physiological saline solution that a colon length is decreased compared to a normal control group, and MPO and inflammatory indicators are increased, but in a group administered with *Bifidobacterium adolescentis* NK98 that a colon length is recovered and MPO inhibitory capacity and inflammatory indicators are improved.

Example 11: Effect of *Bifidobacterium adolescentis* NK98 on improving a cognitive function

On a following day after intraperitoneally administering LPS isolated from E.

coli (0.5 mg/kg/day) to a mouse for five days, lactic acid bacteria were administered thereto, and an effect of lactic acid bacteria on improving a cognitive function was identified in such a way that an object recognition test and a Y maze test were performed and BDNF in a hippocampus was measured by immunoblotting, wherein the results thereof are as shown in a following table 18.

Particularly, as an object recognition test method, two objects (A, A') of the same shape and size were fixed into a box (40 × 40 × 40 cm), of which outside is not visible from the inside, then a mouse was allowed to start moving at the center of the box, and then the number of touching the two objects was recorded for 10 minutes. In 24 hours later, one of the two objects was replaced with a new one (A, B), and then the number of touching an old one and a new one was recorded and quantified.

Also, as a Y maze test method, a test apparatus consisted of the three identical arms (8 cm in length, 30 cm in width, 14 cm in height), each positioning at a certain angle of 120°. The mouse was placed at the end of one arm and allowed to freely move in a Y maze for eight minutes, after which the number and order of entering respective arms were measured, so as to evaluate a spontaneous alteration (%). The alteration was defined as an alphabetical character order, in which the mouse sequentially entered into three arms, i.e. ABC, BCA, CAB, etc.

$$\% \text{ alteration} = [\text{Total number of alterations}] / [\text{Total number of arm entries} - 2] \times 100$$

【Table 18】

	Object recognition test (%)	Y maze test (%)	Hippocampus BDNF expression capacity
Normal control	52.1	76.5	
IS	38.5	56.7	-
NK98	48.6	68.4	+++

IM38	46.5	64.5	++
Buspirone	42.9	62.4	+

* -, <10%; +, 10~30%; ++, 30~50%; +++, >50%

As identified in Table 18 above, it was identified in a group administered with *Bifidobacterium adolescentis* NK98 that a cognitive function is improved and an expression of BDNF is increased through the object recognition test and the Y maze test.

Example 12: Effect of reducing stress according to a co-administration of two types of lactic acid bacteria

Lactobacillus reuteri NK33, a novel lactic acid bacterium, *Bifidobacterium adolescentis* IM38 (accession number: KCCM11807P), a bacterium disclosed in Korean patent publication No. 10-2017-0090359, or a mixture thereof were compared to each other in terms of an effect of reducing stress.

Particularly, physiological salt solution (IS), 1×10^9 cfu of *Lactobacillus reuteri* NK33 (NK33), 1×10^9 cfu of *Bifidobacterium adolescentis* IM38 (IM38) or a combination of the said lactic acid bacteria (NK33+IM38) each 0.5×10^9 cfu was administered into a model with immobilization stress induced as shown in Experimental Example 1-(1) above, and then an elevated plus maze test was performed. After that, blood was collected from each group, and then an amount of corticosterone in blood was measured, wherein the results thereof are as shown in a following table 19.

【Table 19】

	Time spent in open arms (%)	Open arm entries (%)	Blood corticosterone (ng/mL)
Normal control	16.1	34.0	0.9
IS	5.4	16.1	2.4

NK33	14.1	33.5	1.4
IM38	14.8	31.5	1.5
NK33+IM38	17.3	36.5	1.2

As identified in Table 19 above, it was identified in a co-administration group that time spent in open arms and open arm entries are increased and an amount of corticosterone in blood is remarkably decreased compared to a group administered with lactic acid bacteria alone.

Also, an effect of a mixture of *Lactobacillus reuteri* NK33 and *Bifidobacterium adolescentis* NK98 on reducing stress was compared by the same method as shown in an experiment on a co-administration of NK33 and IM38 above, wherein the results thereof are as shown in a following table 20.

【Table 20】

	Time spent in open arms (%)	Open arm entries (%)	Blood corticosterone (ng/mL)
Normal control	18.1	34.2	0.9
IS	5.5	17.1	2.9
NK33	14.1	33.5	1.4
NK98	16.8	33.3	1.3
NK33+NK98	18.5	38.2	1.1

As identified in Table 20 above, it was identified in the co-administration group that time spent in open arms and open arm entries are increased and an amount of corticosterone in blood is remarkably decreased compared to a group administered with lactic acid bacteria alone.

<Accession information of lactic acid bacteria>

The present inventors deposited *Lactobacillus reuteri* NK33 for the purpose of patent to the Korean Culture Center of Microorganisms, a certified depository institution (address: Yulim Building, 45, Hongjeonae 2ga-gil, Seodaemun-gu, Seoul,

South Korea) on August 04, 2017, and received an accession number of KCCM12090P.

Also, the present inventors deposited *Bifidobacterium adolescentis* NK98 for the purpose of patent to the Korean Culture Center of Microorganisms, a certified depository institution (address: Yulim Building, 45, Hongjenae 2ga-gil, Seodaemun-gu, Seoul, South Korea) on August 03, 2018, and received an accession number of KCCM12297P.

2021282468 08 Dec 2021

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE
DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT
PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT issued pursuant to Rule 7.1
by the INTERNATIONAL DEPOSITARY AUTHORITY identified at the bottom of
this page

To. Dong-Hyun Kim
26, Kyungheedaero, Dongdaemun-gu, Seoul 02447, Republic of
Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: <i>Lactobacillus reuteri</i> NK33	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM12090P
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: ☐ a scientific description ☐ a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on August. 04. 2017.	
IV. NATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : Yurim BID 45, Hongjenseo-2ga-gil, Seodaemun-gu, SEOUL 03641, Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: August. 04. 2017.

This translation proves that there is no difference with the content of
the original text.

September. 28. 2018

Patent Attorney Ahn So Young



2021282468 08 Dec 2021

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE
DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT
PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT issued pursuant to Rule 7.1
by the INTERNATIONAL DEPOSITARY AUTHORITY identified at the bottom of
this page

To. Dong-Hyun Kim
26, Kyunghedae-ro, Dongdaemun-gu, Seoul 02447, Republic of
Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: <i>Bifidobacterium adolescentis</i> NK98	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM12297P
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: ☐ a scientific description ☐ a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on August. 03. 2018.	
IV. NATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : Yurim BID 45, Hongjenseong-2ga-gil, Seodaemun-gu, SEOUL 03641, Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: August. 03. 2018.

This translation proves that there is no difference with the content of
the original text.

September. 28. 2018

Patent Attorney Ahn So Young



Claims

1. Isolated *Bifidobacterium adolescentis* NK98, as deposited at the Korean Culture Center of Microorganisms under accession number KCCM12297P.
2. The isolated *Bifidobacterium adolescentis* NK98 KCCM12297P, according to claim 1, wherein the *Bifidobacterium adolescentis* NK98 KCCM12297P comprises a 16S rDNA sequence of SEQ ID NO: 38.
3. A pharmaceutical composition comprising isolated *Bifidobacterium adolescentis* NK98 KCCM12297P.
4. Use of the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P of claim 1 or 2, or the composition of claim 3, in the preparation of a drug for preventing or treating a neurological mental disease.
5. Use of the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P of claim 1 or 2, or the composition of claim 3, in the preparation of a drug for preventing or treating an inflammatory disease.
6. The use according to claim 4 or 5, wherein the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P is a live bacterial cell thereof, a dead bacterial cell thereof, a culture product thereof, a crush thereof or an extract thereof.
7. The use according to claim 4, wherein the drug is for preventing or treating a neurological mental disease, wherein the disease is a neurodegenerative disease or a

mental disorder.

8. The use according to claim 7, wherein the mental disorder is one or more selected from the group comprising anxiety, depression, mood disorder, insomnia, delusional disorder, obsessive disorder, migraine, stress, memory disorder, cognitive disorder and disturbance attention.

9. The use according to claim 7, wherein the neurodegenerative disease is one or more selected from the group comprising Parkinson's disease, Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, spinocerebellar atrophy, Tourette's syndrome, Friedrich's ataxia, Machado-Joseph's disease, dementia, dystonia and progressive supranuclear palsy.

10. The use according to any one of claims 4 to 9, wherein the composition further comprises isolated *Bifidobacterium adolescentis* IM38 KCCM11807P.

11. A method for preventing or treating a neurological mental disease comprising administering to a patient in need thereof, a therapeutically effective amount of the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P of claims 1 or 2, or the composition according claim 3.

12. A method for preventing or treating an inflammatory disease comprising administering to a patient in need thereof, a therapeutically effective amount of the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P of claims 1 or 2, or the composition according claim 3.

13. The method of claim 11 or 12, wherein the isolated *Bifidobacterium adolescentis* NK98 KCCM12297P is a live bacterial cell thereof, a dead bacterial cell thereof, a culture product thereof, a crush thereof or an extract thereof.

14. The method of claim 11, wherein the neurological mental disease is a neurodegenerative disease or a mental disorder.

15. The method of claim 14, wherein the mental disorder is one or more selected from the group comprising anxiety, depression, mood disorder, insomnia, delusional disorder, obsessive disorder, migraine, stress, memory disorder, cognitive disorder and disturbance attention; or wherein the neurodegenerative disease is one or more selected from the group comprising Parkinson's disease, Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, spinocerebellar atrophy, Tourette's syndrome, Friedrich's ataxia, Machado-Joseph's disease, dementia, dystonia and progressive supranuclear palsy.

16. A health functional food comprising isolated *Bifidobacterium adolescentis* NK98 KCCM12297P.

17. Use of the food according to claims 16, in the preparation of a composition for preventing or improving a neurological mental disease.

18. Use of the food according to claims 16, in the preparation of a composition for preventing or improving an inflammatory disease.

Drawings

Fig. 1




Fig. 2




Fig. 3





Fig. 4


Fig. 5

Fig. 6




Fig. 7




Fig. 8





Fig. 9

Fig. 10





Fig. 11


Fig. 12

Fig. 13






Fig. 14

Fig. 15



<110> UNIVERSITY-INDUSTRY COOPERATION GROUP OF KYUNG HEE UNIVERSITY
NAVIPHARM CO, LTD

<120> Novel lactic acid bacteria and use thereof

<130> P17-120-REA-KHU

<160> 38

<170> KoPatentIn 3.0

<210> 1
<211> 1490
<212> DNA
<213> Artificial Sequence

<220>
<223> NK33 16S rDNA

<400> 1

tatggctcag gatgaacgcc ggcggtgtgc ctaatacatg caagtcgtac gcactggccc	60
aactgattga tgggtgcttgc acctgattga cgatggatca ccagtgagtg gcggacgggt	120
gagtaacacg taggtaacct gccccggagc gggggataac atttggaaac agatgctaata	180
accgcataac aacaaaagcc acatggcttt tgtttgaaag atggcttttg ctatcactct	240
gggatggacc tgcggtgcat tagctagttg gtaaggtaac ggcttaccaa ggcgatgatg	300
catagccgag ttgagagact gatcggccac aatggaactg agacacggtc catactccta	360
cgggaggcag cagtagggaa tcttccacaa tgggcgcaag cctgatggag caacaccgcg	420
tgagtgaaga agggtttcgg ctcgtaaagc tctgttggtg gagaagaacg tgcgtgagag	480
taactgttca cgcagtgacg gtatccaacc agaaagtcac ggctaactac gtgccagcag	540
ccgcggtaat acgtaggtgg caagcgttat ccggatttat tgggcgtaaa gcgagcgcag	600
gcggttgctt aggtctgatg tgaaagcctt cggcttaacc gaagaagtgc atcggaacc	660
gggcgacttg agtgcagaag aggacagtgg aactccatgt gtagcggtagg aatgcgtaga	720
tatatggaag aacaccagtg gcgaaggcgg ctgtctggtc tgcaactgac gctgaggctc	780
gaaagcatgg gtagcgaaca ggattagata ccctggtagt ccatgccgta aacgatgagt	840
gctaggtgtt ggagggtttc cgcccttcag tgccggagct aacgcattaa gcactccgcc	900
tggggagtag gaccgcaagg ttgaaactca aaggaattga cgggggcccc cacaagcgg	960
ggagcatgtg gtttaattcg aagctacgcg aagaacctta ccaggtcttg acatcttgcg	1020

2021282468 08 Dec 2021

ctaaccttag agataaggcg ttcccttcgg ggacgcaatg acaggtggtg catggtcgtc	1080
gtcagctcgt gtcgtgagat gttgggttaa gtcccgaac gagcgcaacc cttgttacta	1140
gttgccagca ttaagttggg cactctagtg agactgccgg tgacaaaccg gaggaagggtg	1200
gggacgacgt cagatcatca tgccccttat gacctgggct acacacgtgc tacaatggac	1260
ggtacaacga gtcgcaagct cgcgagagta agctaatttc ttaaagccgt tctcagttcg	1320
gactgtaggc tgcaactcgc ctacacgaag tcggaatcgc tagtaatcgc ggatcagcat	1380
gccgcggtga atacgttccc gggccttgta cacaccgccc gtcacaccat ggggagtttg	1440
taacgccccaa agtcggtggc ctaaccttta tggagggagc cgcctaaggc	1490

<210> 2
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Firmicutes F_primer

<400> 2	
ggagyatgtg gtttaattcg aagca	25

<210> 3
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Firmicutes R_primer

<400> 3	
agctgacgac aaccatgcac	20

<210> 4
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Bacteroidetes F_primer

<400> 4

2021282468 08 Dec 2021

gtttaattcg atgatacgcg ag	22
<210> 5	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Bacteroidetes R_primer	
<400> 5	
ttaascgac acctcacgg	19
<210> 6	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> beta-Proteobacteria F_primer	
<400> 6	
aacgcgaaaa accttaccta cc	22
<210> 7	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> beta-Proteobacteria R_primer	
<400> 7	
tgccctttcg tagcaactag tg	22
<210> 8	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> delta/gamma-Proteobacteria F_primer	
<400> 8	
gctaacgcat taagtryccc g	21

08 Dec 2021

2021282468

<210> 9
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> delta/gamma-Proteobacteria R_primer

<400> 9
 gccatgcrgc acctgtct 18

<210> 10
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> epsilon-Proteobacteria F_primer

<400> 10
 taggcttgac attgatagaa tc 22

<210> 11
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> epsilon-Proteobacteria R_primer

<400> 11
 cttacgaagg cagtctcctt a 21

<210> 12
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Actinobacteria F_primer

<400> 12
 tgtagcgggtg gaatgcgc 18

<210> 13

2021282468 08 Dec 2021

<211>	21	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Actinobacteria R_primer	
<400>	13	
	aattaagcca catgctccgc t	21
<210>	14	
<211>	19	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	TM F_primer	
<400>	14	
	aytgggcgta aagagttgc	19
<210>	15	
<211>	22	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	TM R_primer	
<400>	15	
	tacggytacc ttgttacgac tt	22
<210>	16	
<211>	21	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Verrucomicrobia F_primer	
<400>	16	
	tcakgtcagt atggccctta t	21
<210>	17	
<211>	23	
<212>	DNA	

08 Dec 2021

2021282468

<213>	Artificial Sequence	
<220>		
<223>	Verrucomicrobia R_primer	
<400>	17	
	cagttttyag gatttcctcc gcc	23
<210>	18	
<211>	25	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Enterobacteriaceae F_primer	
<400>	18	
	cattgacgtt acccgagaa gaagc	25
<210>	19	
<211>	21	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Enterobacteriaceae R_primer	
<400>	19	
	ctctacgaga ctcaagcttg c	21
<210>	20	
<211>	24	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Escherichia coli F_primer	
<400>	20	
	cgcgtactat acgcatgaa cgta	24
<210>	21	
<211>	22	
<212>	DNA	
<213>	Artificial Sequence	

2021282468 08 Dec 2021

<220>
 <223> Escherichia coli R_primer

<400> 21
 accgttgatc acttcgggtca gg 22

<210> 22
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Klebsiella spp. F_primer

<400> 22
 gatacggagt atgcctttac ggtg 24

<210> 23
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Klebsiella spp. R_primer

<400> 23
 tagcctttat caagcggata ctgg 24

<210> 24
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Klebsiella oxytoca F_primer

<400> 24
 gttaatacct ttgctcattg a 21

<210> 25
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Klebsiella oxytoca R_primer

2021282468 08 Dec 2021

<400>	25	
accagggtat ctaatcctgt t		21
<210>	26	
<211>	25	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Morganella morganii F_primer	
<400>	26	
ctcgcaccat cagatgaacc catat		25
<210>	27	
<211>	29	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Morganella morganii R_primer	
<400>	27	
caaagcatct ctgctaagtt ctctggatg		29
<210>	28	
<211>	16	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus reuteri F_primer	
<400>	28	
gaacgcaytg gcccaa		16
<210>	29	
<211>	19	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus reuteri R_primer	

08 Dec 2021 2021282468

<400>	29	
tccattgtgg	cgcgatcagt	19
<210>	30	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus johnsonii F_primer	
<400>	30	
cactagacgc	atgtctagag	20
<210>	31	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus johnsonii R_primer	
<400>	31	
agtcttcaa	ctcggctatg	20
<210>	32	
<211>	20	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus plantarum F_primer	
<400>	32	
tcatgattta	catttgagtg	20
<210>	33	
<211>	21	
<212>	DNA	
<213>	Artificial Sequence	
<220>		
<223>	Lactobacillus plantarum R_primer	
<400>	33	
gaccatgcgg	tcgaagttgt t	21

08 Dec 2021
2021282468

<210> 34
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Lactobacillus rhamnosus F_primer

<400> 34
cgcccttaac agcagtcttc 20

<210> 35
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Lactobacillus rhamnosus R_primer

<400> 35
gccctccgta tgcttaaacc 20

<210> 36
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Bacterial 16S rDNA F_primer

<400> 36
tcgtcggcag cgtcagatgt gtataagaga caggtgccag cmgccgcggt aa 52

<210> 37
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Bacterial 16S rDNA R_primer

<400> 37
gtctcgtggg ctcggagatg tgtataagag acagggacta chvgggtwtc taat 54

2021282468 08 Dec 2021

<210> 38
 <211> 1460
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> NK98 16S rDNA

<400> 38
 ctcaggatga acgctggcgg cgtgcttaac acatgcaagt cgaacgggat cccaggagct 60
 tgctcctggg tgagagtggc gaacgggtga gtaatgcgtg accgacctgc cccatacacc 120
 ggaatagctc ctggaaacgg gtggtaatgc cggatgctcc agttggatgc atgtccttct 180
 gggaaagatt catcggatat ggatgggggtc gcgtcctatc agcttgatgg cggggtaacg 240
 gcccaccatg gcttcgacgg gtagccggcc tgagaggggcg accggccaca ttgggactga 300
 gatacggccc agactcctac gggaggcagc agtggggaat attgcacaat gggcgcaagc 360
 ctgatgcagc gacgccgctg gcgggatgac ggccttcggg ttgtaaaccg cttttgactg 420
 ggagcaagcc cttcgggggtg agtgtacctt tcgaataagc accggctaac tacgtgccag 480
 cagccgcggt aatacgtagg gtgcaagcgt tatccggaat tattgggcgt aaagggctcg 540
 taggcggttc gtcgcgtccg gtgtgaaagt ccatcgctta acggtggatc cgcgccgggt 600
 acgggcgggc ttgagtgcgg taggggagac tggaattccc ggtgtaacgg tggaatgtgt 660
 agatatcggg aagaacacca atggcgaagg caggtctctg ggccgtcact gacgctgagg 720
 agcgaaagcg tggggagcga acaggattag ataccctggg agtccacgcc gtaaacgggtg 780
 gatgctggat gtggggacca ttccacggtc tccgtgtcgg agccaacgcg ttaagcatcc 840
 cgcctgggga gtacggccgc aaggctaaaa ctcaaagaaa ttgacggggg cccgcacaag 900
 cggcggagca tgcggattaa ttcgatgcaa cgcgaagaac cttacctggg cttgacatgt 960
 tcccacagc cccagagatg gggcctccct tcggggcggg ttcacagggtg gtgcatggtc 1020
 gtcgtcagct cgtgtcgtga gatgttgggt taagtccgc aacgagcgca accctcgccc 1080
 tgtgttgcca gcacgtcgtg gtgggaactc acgggggacc gccggggtca actcggagga 1140
 aggtggggat gacgtcagat catcatgccc cttacgtcca gggcttcacg catgtacaa 1200
 tggccggtac aacgggatgc gacactgtga ggtggagcgg atcccttaaa accggtctca 1260
 gttcggattg gagtctgcaa cccgactcca tgaaggcgga gtcgctagta atcgcggatc 1320

agcaacgccg	cggtgaatgc	gttcccgggc	cttgtaaca	ccgcccgtca	agtcataaaa	1380
gtgggtagca	cccgaagccg	gtggcccaac	ctttttgggg	ggagccgtct	aaggtgagac	1440
tcgtgattgg	gactaatcta					1460