

(12) STANDARD PATENT

(11) Application No. AU 2002331210 B2

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Compositions comprising sugar beet pectin and carotenoids

(51) International Patent Classification(s)

| | |
|------------------------------|-----------------------------|
| A23L 1/0524 (2006.01) | A61K 8/33 (2006.01) |
| A23K 1/00 (2006.01) | A61K 8/34 (2006.01) |
| A23K 1/16 (2006.01) | A61K 8/35 (2006.01) |
| A23L 1/00 (2006.01) | A61K 8/362 (2006.01) |
| A23L 1/187 (2006.01) | A61K 8/37 (2006.01) |
| A23L 1/275 (2006.01) | A61K 8/60 (2006.01) |
| A23L 1/30 (2006.01) | A61K 8/73 (2006.01) |
| A23L 2/58 (2006.01) | A61K 8/97 (2006.01) |
| A61K 8/00 (2006.01) | A61Q 19/00 (2006.01) |
| A61K 8/25 (2006.01) | C09B 61/00 (2006.01) |
| A61K 8/31 (2006.01) | C09B 67/20 (2006.01) |

(21) Application No: **2002331210** (22) Date of Filing: **2002.08.07**

(87) WIPO No: **WO03/015537**

(30) Priority Data

(31) Number **01119429.7** (32) Date **2001.08.13** (33) Country **EP**

(43) Publication Date: **2003.03.03**

(43) Publication Journal Date: **2003.05.29**

(44) Accepted Journal Date: **2007.07.19**

(71) Applicant(s)
DSM IP Assets B.V.

(72) Inventor(s)
Kuhny, Kurt;Leuenberger, Bruno;Beck, Markus Ivo

(74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(56) Related Art
WO 2000/022939 A
US 5 008 254 A
WO 2000/070967 A

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 February 2003 (27.02.2003)

PCT

(10) International Publication Number
WO 03/015537 A1

(51) International Patent Classification⁷: A23L 1/0524, 1/275

(21) International Application Number: PCT/EP02/08819

(22) International Filing Date: 7 August 2002 (07.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01119429.7 13 August 2001 (13.08.2001) EP

(71) Applicant (for all designated States except US): ROCHE VITAMINS AG [CH/CH]; 124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BECK, Markus, Ivo [DE/DE]; Am Baselweg 16, 79540 Loerrach (DE). KUHNY, Kurt [CH/CH]; Allmendstrasse 29, CH-4206 Seewen (CII). LEUENBERGER, Bruno [CII/CII]; Muehlebachweg 23, CH-4123 Allschwil (CH).

(74) Agent: MUELLER, Ingrid; Grenzacherstrasse 124, CH-4070 Basle (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report
with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/015537 A1

(54) Title: COMPOSITIONS COMPRISING SUGAR BEET PECTIN AND CAROTENOIDS

(57) Abstract: Carotenoid compositions for use as colorants or additives for food, beverages, animal feeds, cosmetics or drugs comprise finely dispersed carotenoids in a sugar beet matrix.

COMPOSITIONS COMPRISING SUGAR BEET PECTIN AND CAROTENOIDS

The present invention relates to novel compositions containing finely dispersed carotenoids and to a process for the preparation of such compositions. The novel compositions of this invention can be used as colorants or additives for food, beverages, animal feeds, cosmetics or drugs.

5 More particularly, the present invention relates to novel compositions comprising sugar beet pectin and a carotenoid, to a process for preparing these compositions, their use as a colorant for food, beverages, animal feeds, cosmetics or drugs; and to food, beverages, animal feeds, cosmetics or drugs containing such compositions.

Pectins are basically $\alpha,1\rightarrow4$ linked polygalacturonic acids which are partially esterified by 10 methyl groups and which can be obtained from plants such as citrus fruit, apples and sugar beet. The term "sugar beet pectin" as used herein denotes pectins obtainable from sugar beet which are characterized and distinguished from citrus and apple pectins in that secondary hydroxy groups are partially acetylated, and by a lack of gelling properties. While such pectins might also be produced from pears and potatoes the commercially 15 readily available pectins of this type are made from sugar beet, e.g. as GENU Beta Pectin Type BETA from CP Kelco (Copenhagen Pectin A/S, DK-4623 Lille Skensved, Denmark). Thus, the term "sugar beet pectin" is intended to denote all pectins having substantially the properties of pectin obtained from sugar beet and comprises pectins obtained from other sources, e.g., pears and potatoes inasmuch as they have substantially the properties of 20 pectins obtained from sugar beet. For the purpose of the present invention, the sugar beet pectins preferably are those of which a 10 wt.-% aqueous solution has a viscosity of 20 to 10000 mPa·s at 50 °C. The average molecular weight of such pectins is assumed to be in the range of 5 to 150 kDalton although this figure is not to be regarded as crucial in view of the well-known problematics of methodology in determinations of molecular weight.

25

The term "carotenoid" as used herein comprises a carotene or structurally related polyene compound which can be used as a colorant for food, beverages, animal feeds, cosmetics or

drugs. Examples of such carotenoids are α - or β -carotene, 8'-apo- β -carotenal, 8'-apo- β -carotenoic acid esters such as the ethyl ester, canthaxanthin, astaxanthin, lycopene, lutein, zeaxanthin or crocetin, or mixtures thereof. The preferred carotenoid is β -carotene.

In the compositions of the present invention, the amount of sugar beet pectin is suitably 5 from about 0.5 to about 60.0 wt.-% and the amount of carotenoid is suitably from about 0.1 to about 20.0 wt.-%.

Suitably, the novel compositions of this invention further contain adjuvants and/or excipients such as one or more of a mono- di-, oligo- or polysaccharide, a triglyceride, a water-soluble antioxidant, a fat-soluble antioxidant, silicic acid and water.

10 Examples of mono- and disaccharides which may be present in the compositions of the present invention are saccharose, invert sugar, glucose, fructose, lactose and maltose. Examples of oligo- or polysaccharides which may be present in the compositions of the present invention are starch and starch hydrolysates, such as dextrans and maltodextrins, especially such in the range of 5-65 dextrose equivalents (hereinafter: DE) and glucose 15 syrup, especially such in the range of 20-95 DE. The term "dextrose equivalent" (DE) denotes the degree of hydrolysis and is measure for the amount of reducing sugar calculated as D-glucose based on dry weight. Native starch has DE close to 0 while glucose has a DE = 100.

The triglyceride is suitably a vegetable oil or fat, such as corn oil, sunflower oil, soybean 20 oil, safflower oil, rape seed oil, arachis oil, palm oil, palm kernel oil, cotton seed oil or cocos oil.

The water-soluble antioxidant may be ascorbic acid and salts thereof, e.g., sodium ascorbate, and the like. The fat-soluble antioxidant may be a tocopherol, e.g., dl- α -tocopherol (i.e., synthetic tocopherol), d- α -tocopherol (i.e., natural tocopherol), β - and γ -tocopherol and mixtures thereof; ascorbic acid esters of fatty acids such as ascorbyl 25 palmitate or stearate; butyl hydroxy toluene; butyl hydroxy anisol; propyl gallate; or t-butyl hydroxy quinoline.

The compositions of the present invention may be an aqueous emulsion (i.e. an oil-in-water dispersion) or a powder.

In one aspect the present invention relates to solid compositions, i.e. stable, water-soluble or dispersible powders. In another aspect, the present invention relates to liquid compositions, i.e., aqueous colloidal solutions or oil-in-water dispersions of such powders.

Typically, a powder composition according to the present invention comprises

- 5 about 1 to about 60 wt.-%, preferably about 5 to about 30 wt.-% of sugar beet pectin;
- about 0.2 to about 20 wt.-% preferably about 0.5 to about 10 wt.-% of a carotenoid;
- 0 to about 70 wt.-% preferably about 0 to about 40 wt.-% of a mono- or 10 disaccharide;
- 0 to about 50 wt.-% preferably about 0 to about 35 wt.-% of starch;
- 0 to about 70 wt.-% preferably about 0 to about 40 wt.-% of a starch hydrolysate;
- about 0.5 to about 50 wt.-% preferably about 1.5 to about 30 wt.-% of a 15 triglyceride;
- 0 to about 5 % preferably about 0.5 to about 2 wt.-% of a water-soluble anti-oxidant;
- 0 to about 5 % preferably about 0.01 to about 2 wt.-% of a fat-soluble anti-oxidant;
- 0 to about 2 wt.-% preferably about 1 wt.-% of silicic acid; and
- 0 to about 10 wt.-% preferably about 1 to about 5 wt.-% of water;

20 the percentages of all ingredients totalling 100.

In accordance with the invention, the novel carotenoid compositions can be prepared by processing the ingredients in a manner known per se for the preparation of water-soluble or dispersible carotenoid compositions. Thus, the compositions can be prepared by a 25 process which comprises homogenizing an aqueous solution or colloidal solution of the pectin and optional water-soluble excipients and adjuvants, a solution or dispersion of the carotenoid and optional fat-soluble adjuvants in a triglyceride and, if required, converting the dispersion obtained into a powder.

30 Typically, sugar beet pectin and optional water-soluble excipients and adjuvants are dissolved in water. The carotenoid and optional fat-soluble excipients and adjuvants are dissolved or suspended in triglyceride. The carotenoid solution (or dispersion) is then added to the aqueous pectin solution with stirring and the mixture is homogenized using

- 4 -

conventional technology, e.g., by high-pressure homogenization, mixing devices as described in EP 1008380-A, high shear emulsification (rotor-stator systems), micronization or wet milling.

The so-obtained oil-in-water dispersion can be converted into a solid composition, e.g. a

- 5 dry powder using conventional technology such spray-drying, spray drying in combination with fluidized-bed granulation (the latter technique commonly known as fluidized spray drying or FSD), or by a powder-catch technique where sprayed emulsion droplets are caught in a bed of an absorbant such as starch and subsequently dried.

- 10 The novel compositions of this invention can find use as colorants or vitamin A supplement for food, beverages, animal feeds, cosmetics or drugs. By the present invention there are preferably provided compositions comprising β -carotene as a colouring agent. These compositions, when dissolved, dispersed or diluted in/with water to a final β -carotene concentration of 10 ppm are typically characterized by ultraviolet/visible-
- 15 spectroscopy using deionized water as reference. At a sample thickness of 1 cm the dispersions show an extinction of at least 0.3 (preferably above 1.0) absorbance units at the wavelength of maximum optical density in the range of 400 to 600 nm. This is equivalent to a formal extinction coefficient of β -carotene in aqueous dispersion E(1%, 1cm) of 300 (preferably >1000).

20

The following Examples illustrate the invention further.

Example 1

A dry premix of 80 g of beet pectin (GENU Pectin Type Beta of Copenhagen Pectin A/S; viscosity of a 10 % aqueous solution of the pectin at 50°C around 4000 mPa·s), 160 g of sucrose and 80 g of a maltodextrin (DE 20-23) was prepared. The dry premix was 5 dissolved in 1200 ml of deionized water at 60°C and another 335 g of maltodextrin (DE 20-23) were added. After complete dissolution of the solids 8.0 g of Na-ascorbate was added to the mixture (= solution A).

136 g of a triglyceride (Durkex 500, partly hydrogenated soybean oil of Loders Croklaan B.V.; 1520 AA Wormerveer The Netherlands) and 0.9 g of dl- α -tocopherol were mixed 10 and heated to 140°C. Subsequently, 11 g of β -carotene was suspended in the mixture of triglyceride and tocopherol. By stirring for about 10 minutes at 140°C a clear solution of β -carotene was obtained (= solution B).

Solution A was heated to 70°C and a crude emulsion was prepared by adding 135 g of 15 solution B to solution A while gently stirring. A fine emulsion was obtained by a five passage high pressure homogenizing treatment of the preemulsion at a pressure of 50/300 bar (APV Lab Homogenizer Type Gaulin Lab 40-10 RBFI of APV Switzerland AG, CH-3076 Worb). The emulsion was diluted by adding an equal volume of deionized water at 60°C and then spray dried in a laboratory spray dryer (Mobile Minor of GEA Niro A/S, DK-2860 Söborg) at an inlet temperature of 200°C-210°C and an outlet temperature of 70-20 75°C. The spray-dried powder was dried in a vacuum oven at room temperature over night.

A fine powder was obtained with a water content of 2.2 %. The β -carotene content of the 15 powder was 1.1 % as determined by spectrophotometry and HPLC-analysis. The powder was dispersed in deionized water and the extinction of the dispersion was measured in a 1 cm quartz precision cell against water. For a 10 ppm dispersion of β -carotene an 20 extinction of 2.109 at a wavelength of 464 nm was calculated ($E(1\%, 1\text{cm}) = 2109$)

Example 2

A dry premix of 160 g of beet pectin (Copenhagen Pectin A/S; viscosity of a 10 % aqueous solution of the pectin at 50°C around 500 mPa·s), 160 g of sucrose and 335 g of a maltodextrin (DE 20-23) was prepared. The dry premix was dissolved in 1400 ml of deionized water at 60°C. After complete dissolution of the solids 8.0 g of Na-ascorbate was added to the mixture (= solution A).

136 g of a triglyceride (Durkex 500) and 0.9 g of dl- α -tocopherol were mixed and heated to 140°C. Subsequently, 11 g of β -carotene was suspended in the mixture of triglyceride and 10 tocopherol. By stirring for about 10 minutes at 140°C a clear solution of β -carotene was obtained (= solution B).

Solution A was heated to 70°C and a crude emulsion was prepared by adding 135 g of solution B to solution A while gently stirring. A fine emulsion was obtained by a three passage high pressure homogenizing treatment of the preemulsion at a pressure of 50/300 15 bar (APV Lab Homogenizer Type Gaulin Lab 40-10 RBF1). The emulsion was diluted by adding an equal volume of deionized water at 60°C and then spray dried in a laboratory spray drier (Mobile Minor of GEA Niro A/S) at an inlet temperature of 200°C-210°C and an outlet temperature of 70-75°C. The spray dried powder was dried in a vacuum oven at room temperature over night.

20 A fine powder was obtained with a water content of 2.5 %. The β -carotene content of the powder was 1.2 % as determined by spectrophotometry and HPLC-analysis. The powder was dispersed in deionized water and the extinction was measured in a 1 cm quartz precision cell against water. For a 10 ppm dispersion of β -carotene an extinction of 2.051 at a wavelength of 463 nm was calculated ($E(1\%,1\text{cm})=2051$).

25 Example 3

A dry premix of 28.6 g beet pectin (GENU Pectin Type Beta of Copenhagen Pectin A/S; viscosity of a 10 % aqueous solution of the pectin at 50°C around 4000 mPa·s) and 121.4 g sucrose was prepared. The dry premix was dissolved in 180 ml of deionized water at 50°C for 30 minutes under stirring (= solution A).

A 30 % suspension of β -carotene in corn oil stabilized by dl- α -tocopherol (β -Carotene 30 % FS of Roche Vitamins) was heated under stirring for about 30 minutes at a temperature of 160°C (= solution B).

An emulsion was prepared by adding solution B to solution A. By vigorously stirring for

5 30 minutes at 50°C a fine emulsion was obtained. The emulsion was diluted by adding 200 ml of deionized water.

300 g of the diluted emulsion were taken and, again, diluted with 50 ml of water. The final emulsion was sprayed into a cooled fluidized bed of corn starch. Excess corn starch was removed by sieving and a coarse powder was obtained. The powder was dried in an air

10 stream at room temperature for about 2 hours.

A powder was obtained with a water content of 6.4 %. The β -carotene content of the powder was 2.5 % as determined by spectrophotometrical assay. The starch content of the powder was 54 %. The powder was dispersed in deionized water and the extinction was measured in a 1 cm quartz precision cell against water. For a 10 ppm dispersion of β -carotene an extinction of 0.401 at a wavelength of 530 nm was calculated ($E(1\%, 1\text{cm}) = 401$).

Example 4

Instant beverage powders were prepared according to the following compositions:

| Ingredients | # 1 [g] | # 2 [g] |
|-------------------------------|---------|---------|
| Sucrose, fine crystalline | 920.0 | 920.0 |
| Ascorbic acid, fine powder | 2.0 | 2.0 |
| Citric acid anhydrous, powder | 55.0 | 55.0 |
| Orange flavor ¹ | 7.0 | 7.0 |
| Tri-Na citrate anhydrous | 6.0 | 6.0 |
| Tri-Ca phosphate | 5.0 | 5.0 |

- 8 -

| | | |
|------------------------|-----|-----|
| powder according Ex. 1 | 5.0 | - |
| powder according Ex. 2 | - | 5.0 |

e.g. Orange Flavor 76905-71 from Givaudan Duebendorf Ltd

Procedure:

All ingredients were sieved through a 0.7 mm sieve.

5 The sieved ingredients were blended in a turbula mixer for 20 minutes

Example 5

Instant pudding powders were prepared according to the following compositions:

| Ingredients | # 1 [g] | # 2 [g] |
|-----------------------------|---------|---------|
| Sucrose, fine crystalline | 840.0 | 840.0 |
| Corn starch, cold swelling | 129.0 | 129.0 |
| Stabilizer ¹ | 23.0 | 23.0 |
| Vanilla flavor ² | 4.0 | 4.0 |
| powder according Example 1 | 4.0 | - |
| powder according Example 2 | - | 4.0 |

e.g. Flanogen ADG 56 from SKW Biosystems

10 ² e.g. Vanilla flavor 75016-32 from Givaudan Dubendorf Ltd

Procedure:

All ingredients were sieved through a 0.7 mm sieve.

The sieved ingredients were blended in a turbula mixer for 20 min.

The claims defining the invention are as follows:

1. A composition comprising pectin obtainable from sugar beet, a triglyceride, a carotenoid in the amount of from about 0.1 to about 20.0 wt % and, optionally, adjuvants and/or excipients.

5 2. A composition as claimed in claim 1 wherein the carotenoid is α - or β -carotene, 8'-apo- β -carotenal, 8'-apo- β -carotenoic acid ethyl ester, canthaxanthin, astaxanthin, lycopene, lutein, zeaxanthin or crocetin or mixtures thereof.

3. A composition as claimed in claim 2 wherein the carotenoid is β -carotene.

4. A composition as claimed in any one of claims 1 to 3 wherein the pectin is 10 one of which a 10 wt.-% aqueous solution has a viscosity of 20 to 10000 mPa·s at 50°C.

5. A composition as claimed in any one of claims 1 to 4 wherein at least one of a mono-, di-, oligo- or polysaccharide, a water-soluble anti-oxidant, a fat-soluble anti-oxidant, silicic acid and water are additionally present.

6. A composition as claimed in claim 5 wherein the mono- or disaccharide is 15 saccharose, invert sugar, glucose, fructose, lactose or maltose.

7. A composition as claimed in claim 5 wherein the polysaccharide is a starch or a starch hydrolysate.

8. A composition as claimed in claim 7 wherein the starch hydrolysate is a 20 dextrin or a maltodextrin (in the range of 5-65 dextrose equivalents) or a glucose syrup (in the range of 20-95 dextrose equivalents).

9. A composition as claimed in claim 1 wherein the triglyceride is a vegetable oil or fat.

10. A composition as claimed in any one of claims 1 to 9 wherein the amount of pectin is from about 0.5 to about 60.0 wt.-%.

25 11. A composition as claimed in any one of claims 1 to 10 which is a powder.

12. A composition as claimed in claim 11 which comprises

about 1 to about 60 wt.-% of pectin;

about 0.2 to about 20 wt.-% of a carotenoid;

0 to about 70 wt.-% of a mono- or disaccharide;

30 0 to about 50 wt.-% of starch;

0 to about 70 wt.-% of a starch or a starch hydrolysate;

about 0.5 to about 50 wt.-% of a triglyceride;

0 to about 5% of a water-soluble anti-oxidant;

0 to about 5% of a fat-soluble anti-oxidant;

35 0 to about 2 wt.-% of silicic acid; and

0 to about 10 wt.-% of water.

13. A composition as claimed in any one of claims 1 to 10 which is an oil-in-water dispersion.

14. A composition as claimed in claim 13 which comprises

5 about 0.5 to about 30 wt.-% of pectin,

about 0.1 to about 10 wt.-% of a carotenoid,

0 to about 35 wt.-% of a mono- or disaccharide.

0 to about 35 wt.-% of a starch or a starch hydro-

about 0.25 to about 25 wt.-% of a triglyceride.

0 to about 2.5% of a water-soluble anti-oxidant.

0 to about 2.5% of a fat-soluble anti-oxidant, and

54-1-1-1-05-1-04-6-1

Volume 11 Number 11 November 2011

A composition as claimed

13. A composition as claimed in any one of claims 12 to 14 which when dissolved, dispersed or diluted with/in water to a final β -carotene concentration of 10 ppm
15 has an extinction coefficient E (1%, 1 cm) of ≥ 300 at the extinction maximum.

16. A process for the preparation of a composition as claimed in any one of claims 1 to 15 which comprises homogenising, in an aqueous solution or colloidal solution of the pectin and optional water-soluble excipients and adjuvants, a solution or dispersion of the carotenoid and optional fat-soluble adjuvants in a triglyceride and, if required, converting the dispersion into a powder.

17. A composition comprising pectin obtainable from sugar beet, a triglyceride, a carotenoid in the amount of from about 0.1 to about 20.0 wt % and, optionally, adjuvants and/or excipients, substantially as hereinbefore described with reference to any one of Examples 1 to 3.

25 18. The use of a composition as claimed in any one of claims 1 to 15 or 17 as a colorant for food, beverages, animal feeds, cosmetics or drugs.

19. Food, beverages, animal feeds, cosmetics or drugs containing a composition as claimed in any one of claims 1 to 15 or 17.

Dated 23 February, 2007

DSM IP Assets B.V.

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON