

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 March 2010 (04.03.2010)

(10) International Publication Number
WO 2010/025421 A3

(51) International Patent Classification:

C07K 14/82 (2006.01) A61K 38/16 (2006.01)
C07K 19/00 (2006.01)

(21) International Application Number:

PCT/US2009/055443

(22) International Filing Date:

28 August 2009 (28.08.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/092,708 28 August 2008 (28.08.2008) US

(71) Applicant (for all designated States except US): **TAIGA BIOTECHNOLOGIES, INC.** [US/US]; 12635 East Montview Boulevard, Aurora, Colorado 80045-7336 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **REFAELI, Yosef** [IL/US]; 843 Garfield Street, Denver, Colorado 80206 (US). **TURNER, Brian Curtis** [US/US]; 655 S. Glencoe Street, Denver, Colorado 80246 (US).

(74) Agents: **WILSON SONSINI GOODRICH & ROSATI** et al.; 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

19 August 2010

(54) Title: MODULATORS OF MYC, METHODS OF USING THE SAME, AND METHODS OF IDENTIFYING AGENTS THAT MODULATE MYC

(57) Abstract: Disclosed herein are methods of modulation of the viability of a cell. Further disclosed herein are methods of modulating an immune response. Further disclosed herein are methods of identifying agents capable of modulation of the viability of a cell or an immune response. Further disclosed herein are agents and compositions capable of modulation of the viability of a cell or an immune response.



WO 2010/025421 A3

MODULATORS OF MYC, METHODS OF USING THE SAME, AND METHODS OF IDENTIFYING AGENTS THAT MODULATE MYC

CROSS-REFERENCE

5 [0001] This application claims the benefit of U.S. Provisional Application No. 61/092,708, filed 28-August-2008, which application is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

10 [0002] There are multiple types of vaccines: vaccines containing dead microorganisms (e.g. the influenza and cholera vaccines); vaccines containing attenuated microorganisms (e.g. the MMR vaccine); vaccines containing toxoids (e.g. the DPT vaccine); vaccines containing subunits of the pathogen (e.g. the HPV vaccine); and nucleic acid (e.g. RNA and DNA) vaccines (e.g. avian flu vaccine, West Nile Virus vaccine, and multiple cancer
15 vaccines). Vaccine adjuvants are agents that stimulate the immune system and increase the response of the immune system to a vaccine.

SUMMARY OF THE INVENTION

[0002a] In a first aspect, the present invention provides a pharmaceutical composition for
20 modulating an immune system, comprising: (a) an antigenic moiety derived from a pathogen; (b) a fusion peptide comprising: (i) a transporter peptide sequence; (ii) a MYC sequence; and optionally; (iii) one or more molecules that link the transporter peptide sequence and the MYC sequence; and (c) a pharmacologically-acceptable excipient.

[0002b] In a second aspect, the present invention provides use of fusion peptide for the
25 manufacture of a medicament for modulating an immune response in an individual, the fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally; (c) one or more molecules that link the transporter peptide sequence and the MYC sequence; wherein the medicament is administered before, after, or during the administration of a vaccine.

[0002c] In a third aspect, the present invention provides a method for modulating an
30 immune response in an individual, the method comprising administering to the individual a fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the

MYC sequence; wherein the fusion protein is administered before, after, or during the administration of a vaccine.

[0003] There is a need for vaccine adjuvants that enhance an immunological response to an antigen. The inventors have discovered that increasing the expression of MYC or the activity of a MYC peptide enhances immunological responses to antigens. In order to increase MYC concentrations in the nucleus of cells with minimal side-effects, the inventors have designed a fusion peptide that can penetrate the nucleus of cells. This peptide can be administered topically to decrease systemic effects. This peptide also increases the viability of leukocytes, including T-cells, and B-cells. Further, the inventors have designed several assays for identifying other agents that meet this unmet need.

[0004] There is also a need for methods of treating autoimmune disorders. The inventors have discovered that decreasing the expression of MYC or the activity of a MYC peptide treats auto-immune disorders. In order to fulfill this unmet need, the inventors have designed several assays for identifying agents that inhibit (partially or fully) the activity of an immune system. In certain instances, the agents that inhibit the immune system decrease the viability of leukocytes or increase the proportion of anergic B-cells.

[0005] Disclosed hereon, in certain embodiments, is a peptide that up-regulates the expression of a MYC gene, the activity of a Myc polypeptide, or a combination thereof, comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (I):

transporter peptide sequence -MYC sequence.

In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein in X is at least one amino acid. In some embodiments, the peptide has the following amino acid sequence (SEQ ID NO: 2):

MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEE
 NFYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGLCSPSYVAVTPFSLRGDND
 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSA
 AAKLVSEKLASYQAARKDSGSPNPARGHVCSTSSLYLQDLSAAASECIDPSVVF
 PYPLN DSSSPKSCASQDSSAFSPSSDLSSTESSPQGSPEPLVLHEETPPTTSSDSE
 EEEQEDEE EIDVVSVEKRQAPGKRSESGSPSAGGHKPPHSPLVLKRCHVSTHQ
 HNYAAPPSTR KDYPAAKRVKLDSVRVLRQISNNRKCTSPRSSDTEENVKRRTH
 NVLERQRRNELK RSFFALRDQIPELENNEKAPKVVILKKATAYILSVQAEEQKL
 ISEEDLLRKRREQLKH KLEQLRKGELNSKLEGKPIP
 NPLLGLDSTRTGHHHHHH.

[0006] Disclosed herein, in certain embodiments, is a composition that induces an immune response, comprising a fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (I):

transporter peptide sequence -MYC sequence.

In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein in X is at least one amino acid. In some embodiments, the peptide has the following amino acid sequence (SEQ ID NO: 2):

MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEE
 NFYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGLCSPSYVAVTPFSLRGDND
 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSA
 AAKLVSEKLASYQAARKDSGSPNPARGHVCSTSSLYLQDLSAAASECIDPSVVF
 PYPLN DSSSPKSCASQDSSAFSPSSDLSSTESSPQGSPEPLVLHEETPPTTSSDSE
 EEEQEDEE EIDVVSVEKRQAPGKRSESGSPSAGGHKPPHSPLVLKRCHVSTHQ
 HNYAAPPSTR KDYPAAKRVKLDSVRVLRQISNNRKCTSPRSSDTEENVKRRTH
 NVLERQRRNELK RSFFALRDQIPELENNEKAPKVVILKKATAYILSVQAEEQKL
 ISEEDLLRKRREQLKH KLEQLRKGELNSKLEGKPIP
 NPLLGLDSTRTGHHHHHH.

In some embodiments, the composition further comprises an antigen, antigenic moiety, or a combination thereof. In some embodiments, the antigen is a dead microorganism, an attenuated microorganism, a toxoid, a subunit of a pathogen, a nucleic acid, a polymer of nucleic acids, or combinations thereof. In some embodiments, the antigen is derived from: hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; meningococcal; pneumonia;

smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus; or combinations thereof. In some embodiments, the composition further comprises an antigenic moiety derived from a pathogen selected from: hepatitis A ; hepatitis B ; polio ; measles ; mumps ; rubella ; diphtheria ; pertussis ; tetanus ; influenza ; varicella zoster virus ; rotavirus ; meningococcal ; pneumonia ; smallpox ; cholera ; bubonic plague; yellow fever ; tuberculosis ; human papillomavirus; or combinations thereof. In some embodiments, the composition further comprises an antigenic moiety derived from a neoplastic cell. In some embodiments, the composition is formulated for topical administration.

[0007] Disclosed herein, in certain embodiments, is a method of inducing an immune response, comprising administering to an individual in need thereof: (a) a vaccine against at least one antigen, and (b) a fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (I):

transporter peptide sequence -MYC sequence.

In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein in X is at least one amino acid. In some embodiments, the peptide has the following amino acid sequence (SEQ ID NO: 2):

MRKKRRQRRRMDFRIVENQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEE
 NFYQQQQQSELQPPAPSEDIWKKFELLPTPLSPSRRSGLCSPSYVAVTPFSLRGDND
 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSAAAK
 LVSEKLASQYQAARKDSGSPNPARGHSVCSTSSLYLQDLSAAASECIDPSVVFYPLN
 DSSSPKSCASQDSSAFSPSSDLSLSTESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEE
 EIDVVSVEKRQAPGKRSESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPPSTR
 KDYPAAKRVKLDVRLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELK
 RSFFALRDQIPELENNEKAPKVVLKATAYILSVQAEQKLISEEDLLRKRREQLKH
 KLEQLRKGELNSKLEGKPIPNPPLLGLDSTRTGHHHHHH. In some embodiments, the

vaccine is selected from the group consisting of: hepatitis A vaccine; hepatitis B vaccine; polio vaccine; measles vaccine; mumps vaccine; rubella vaccine; diphtheria vaccine; pertussis vaccine; tetanus vaccine; influenza vaccine; varicella zoster virus vaccine; rotavirus vaccine; meningococcal vaccine; pneumonia vaccine; smallpox vaccine; cholera vaccine; bubonic plague vaccine; yellow fever vaccine; tuberculosis vaccine; human papillomavirus vaccine; a cancer vaccine; or combinations thereof. In some embodiments, the vaccine is administered before, after, or simultaneously with the agent that increases the nucleic concentration of a Myc polypeptide.

[0008] Disclosed herein, in certain embodiments, is a method of increasing the viability of a cell, comprising administering to an individual in need thereof an agent that up-regulates the

expression of MYC, increases the activity of a Myc peptide, or a combination thereof. In some embodiments, the agent is a fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has

Formula (I):

transporter peptide sequence -MYC sequence.

In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein X is at least one amino acid. In some embodiments, the peptide has the following amino acid sequence (SEQ ID NO: 2):

MRKKRRQRRRMDFRIVENQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEE
 NFYQQQQQSELQPPAPSEDIWKKFELLTPPLSPSRRSGLCSPSYVAVTPFSLRGDND
 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSA
 AAKLVSEKLSYQAARKDSGSPNPARGHSVCSSTSSLYLQDLSAAASECIDPSVVF
 PYPLNDSSSPKSCASQDSSAFSPSSDLSSTESSPQGSPEPLVLHEETPPTTSSD
 SEEEQEDEEIDVVSVEKRQAPGKRSESGSPSAGGHKPPHSPLVLKRCHVSTHQ
 HNYAAPPSTRKDYPAAKRVKLDsvrvlrqisnnrkctsprssdteenvkrrthnvl
 erqrrnelkrsffalrdqiPELENNEKAPKVVILKKATAYILSVQAEEQKLISEED
 LLRKRREQLKHKLEQLRKGELNSKLEGKPIPnPLGLDSTRTGHHHHHH. In some

embodiments, the cell is a leukocyte. In some embodiments, the cell is a T-cell. In some embodiments, the cell is a B-cell. In some embodiments, the cell is a memory T-cell.

[0009] Disclosed herein, in certain embodiments, is a method of decreasing the viability of a cell, comprising administering to an individual in need thereof an agent that down-regulates the expression of MYC, decreases the activity of a Myc peptide, or a combination thereof.

[0010] Disclosed herein, in certain embodiments, is a method of treating a disorder characterized by the under-expression of a MYC gene or a deficit in the activity of a Myc polypeptide, comprising administering to an individual in need thereof an agent that up-regulates the expression of MYC, increases the activity of a Myc peptide, or a combination thereof. In some embodiments, the agent is a fusion peptide comprising: (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide

has Formula (I):

transporter peptide sequence -MYC sequence.

In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, the peptide has Formula (II):

transporter peptide sequence-X-MYC sequence,

wherein in X is at least one amino acid. In some embodiments, the peptide has the following amino acid sequence (SEQ ID NO: 2):

MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEE
 5 NFYQQQQQSELQPPAPSEDIWKKFELLTPPLSPSRRSGLCSPSYVAVTPFSLRGDND
 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSAAAK
 LVSEKLASYQAARKDSGSPNPARGHSVCSTSSLYLQDLSAAASECIDPSVVFYPLN
 DSSSPKSCASQDSSAFSPSSDLSSTESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEE
 EIDVVSVEKRQAPGKRSESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPPSTR
 10 KDYPAAKRVKLDsvrVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELK
 RSFFALRDQIPELENNEKAPKVVILKKATAYILSVQAEEQKLISEEDLLRKRREQLKH
 KLEQLRKGELNSKLEGKPIPNPLLGLDSTRTGHHHHHH.

[0011] Disclosed herein, in certain embodiments, is a method of treating a disorder characterized by the over-expression of a MYC gene or the excess activity of a Myc polypeptide comprising, administering to an individual in need thereof an agent that down-regulates the expression of MYC, decreases the activity of a Myc peptide, or a combination thereof.

[0012] Disclosed herein, in certain embodiments, is a method of identifying an agent that up-regulates the expression of a MYC gene, and/or the activity of a Myc polypeptide, the method comprising: (a) contacting a plurality of anergic B-cells with an agent; and (b) following contact with the agent, detecting and/or measuring the level of expression of one or more cell surface markers in the cell culture, wherein the presence of the cell surface marker is indicative a non-anergic B-cell. In some embodiments, the anergic B-cells are obtained from mice with the phenotype BCR^{HEL}/sHEL. In some embodiments, the cell surface marker is: IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2). In some embodiments, the level of expression of a cell surface marker is measured by contacting the plurality of cells with a detectable antibody or detectable antigen that binds to the cell surface marker.

[0013] Disclosed herein, in certain embodiments, is a method of identifying an agent that up-regulates the expression of a MYC gene, and/or the activity of a Myc polypeptide, the method comprising: (a) contacting a plurality of factor-dependent cells with an agent; and (b) following contact with the agent, detecting and measuring the level of expansion of the plurality of cells. In some embodiments, the factor-dependent cells are lymphoid cells. In some embodiments, the factor-dependent cells are: IL-2^{-/-}, IL-3^{-/-}, IL-4^{-/-}, IL-5^{-/-}, IL-6^{-/-}, IL-7^{-/-}, IL-8^{-/-}, IL-9^{-/-}, IL-10^{-/-}, IL-11^{-/-}, IL-12^{-/-}, or any combinations thereof. In some embodiments, the factor-dependent cells are derived from: CTLL-2 cells or BAF/3 cells.

[0014] Disclosed herein, in certain embodiments, is a method of identifying an agent that up-regulates the expression of a MYC gene, and/or the activity of a Myc polypeptide, comprising: (a) transforming a plurality of cells with a reporter construct, comprising: a reporter gene operably linked to an E-box sequence encoded

in a myc-responsive promoter; (b) contacting the plurality of cells with an agent; and (c) following contact with the agent, detecting and measuring the level of expression of the reporter gene. In some embodiments, the cells are lymphoid cells. In some embodiments, the myc-responsive promoter is an ornithine decarboxylase promoter. In some embodiments, the reporter gene is a β -galactosidase gene, a β -lactamase gene, a horseradish peroxidase gene, an alkaline phosphatase gene, a thymidine kinase gene, a xanthine phosphoribotransferase gene, a tyrosinase gene, a cytosine deaminase gene, an antibiotic resistance gene, or a gene having a fluorescent expression product.

DETAILED DESCRIPTION OF THE INVENTION

10 [0015] Disclosed herein, in certain embodiments, are methods of identifying agents capable of modulating (e.g., activating, increasing, inducing, supplementing, decreasing, or inhibiting) an immune system. In some embodiments, agents are utilized to increase an immune response. In some embodiments, agents are used to decrease an immune response. Further, disclosed herein are methods of modulating an immune system using agents identified by any of the means disclosed herein.

15

Certain Definitions

[0016] Unless indicated otherwise, the following terms have the following meanings when used herein and in the appended claims.

[0017] The term “vaccine” means a composition comprising antigens and/or antigenic moieties administered to a mammal in order to elicit an immune response. The antigen or antigenic moiety is a live or attenuated microorganism, a natural product purified from a microorganism (e.g. a subunit of a protein, peptide, polysaccharide, and a nucleic acid), a synthetic product designed to mimic a component of the microorganism, a genetically engineered protein, peptide, or polysaccharide, a patient's own tumor cells, a natural product purified from a tumor cell (e.g. Prostate Specific Antigen, gp96, GM2, GD2, GD3, carcinoembryonic antigen, MART-1, and tyrosinase), a synthetic molecule designed to mimic a component of a tumor cell (e.g. sialyl Tn), or combinations thereof. Further, the term vaccine includes current vaccines, and any novel and/or modified vaccines developed.

[0018] The term “lymphoid tissue” means tissue associated with the lymphatic system. By way of non-limiting example, lymphoid tissue includes tissue obtained from the lymph nodes, tonsils, spleen, bone marrow, and thymus.

[0019] The term “lymphocyte” refers to all immature, mature, undifferentiated and differentiated white lymphocyte populations including tissue specific and specialized varieties. It encompasses, by way of non-limiting example, B-cells, T-cells, NKT cells, and NK cells. In some embodiments, lymphocytes include all B-cell lineages including pre-B-cells, Progenitor B cells, Early Pro-B cells, Late Pro-B cells, Large Pre-B

cells, Small Pre-B cells, Immature B cells, Mature B cells, plasma B-cells, memory B-cells, B-1 cells, B-2 cells and anergic AN1/T3 cell populations.

[0020] The term B-cell, refers to, by way of non-limiting example, a pre-B-cell, Progenitor B cell, Early Pro-B cell, Late Pro-B cell, Large Pre-B cell, Small Pre-B cell, Immature B cell, Mature B cell, plasma B-cell, memory B-cell, B-1 cell, B-2 cells and anergic AN1/T3 cell populations. In some embodiments, the term B-cell includes a B-cell that expresses an immunoglobulin heavy chain and/or light chain on its cells surface. In some embodiments, the term B-cell includes a B-cell that expresses and secretes an immunoglobulin heavy chain and/or light chain. In some embodiments, the term B-cell includes a cell that binds an antigen on its cell-surface. In some embodiments disclosed herein, B-cells or AN1/T3 cells are utilized in the processes described. In certain embodiments, such cells are optionally substituted with any animal cell suitable for expressing, capable of expressing (e.g., inducible expression), or capable of being differentiated into a cell suitable for expressing an antibody including, e.g., a hematopoietic stem cell, a B-cell, a pre-B-cell, a Progenitor B cell, a Early Pro-B cell, a Late Pro-B cell, a Large Pre-B cell, a Small Pre-B cell, an Immature B cell, a Mature B cell, a plasma B-cell, a memory B-cell, a B-1 cell, a B-2 cell, an anergic B-cell, or an anergic AN1/T3 cell.

[0021] The term “immune response” includes the identification and neutralization of pathogens and/or tumor cells. In some embodiments, the immune response is the adaptive immune response. By way of non-limiting example, the adaptive immune response includes the development of immunological memory.

[0022] The terms “antibody” and “antibodies” refer to any form of a natural occurring (but isolated and/or purified), engineered or synthetic antibody, including monoclonal antibodies, polyclonal antibodies, bi-specific antibodies, multispecific antibodies, grafted antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, camelized antibodies, single-chain Fvs (scFv), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies and antigen-binding fragments of any of the above. In particular, antibodies include immunoglobulin molecules and immunologically active fragments of immunoglobulin molecules, i.e., molecules that contain an antigen binding site. Immunoglobulin molecules are of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or subclass. The terms “antibody” and immunoglobulin are used interchangeably in the broadest sense. In some embodiments an antibody is part of a larger molecule, formed by covalent or non-covalent association of the antibody with one or more other proteins or peptides.

[0023] The antibodies herein include monoclonal, polyclonal, recombinant, chimeric, humanized, bi-specific, grafted, human, and fragments thereof including antibodies altered by any means to be less immunogenic in humans. Thus, for example, the monoclonal antibodies and fragments, *etc.*, herein include “chimeric” antibodies and “humanized” antibodies. In general, chimeric antibodies include a portion of the heavy and/or light chain that is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass. while the remainder

of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567); Morrison *et al. Proc. Natl Acad. Sci.* **81**:6851-6855 (1984). For example, in some embodiments a chimeric antibody contains variable regions derived from a mouse and constant regions derived from human in whom the constant region contains sequences homologous to both human IgG2 and human IgG4. "Humanized" forms of non-human (*e.g.*, murine) antibodies or fragments are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include, grafted antibodies or CDR grafted antibodies wherein part or all of the amino acid sequence of one or more complementarily determining regions (CDRs) derived from a non-human animal antibody is grafted to an appropriate position of a human antibody while maintaining the desired binding specificity and/or affinity of the original non-human antibody. In some embodiments, corresponding non-human residues replace Fv framework residues of the human immunoglobulin. In some embodiments humanized antibodies comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In some embodiments, the humanized antibody comprises substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. For further details, see, *e.g.*: Jones *et al., Nature* **321**: 522-525 (1986); Reichmann *et al., Nature* **332**: 323-329 (1988) and Presta, *Curr. Op. Struct. Biol.* **2**: 593-596 (1992).

[0024] The term "antigen" refers to a substance that elicits the production of an antibody.

[0025] The term "antigenic moiety" means the portion of a molecule, organism (*e.g.*, a bacteria, virus), or cell to which an antibody binds (or complements).

[0026] The term "self antigen" refers to an antigen that originates from within an animal, tissue, or cell. In some embodiments a self antigen comprises an endogenous antigen. In some embodiments a self antigen comprises an endogenous antigen produced by an endogenous retrovirus. In some embodiments self antigens comprise neo-self antigens, microbially or parasite encoded neo-self antigens, or other neo-self antigens expressed as a result of genetic alteration to an animal or cell following whole-body irradiation and HSC transplantation (which re-starts the immune system and allows the new antigen to be considered "self"). In some embodiments a chimeric mouse expresses a neo-self antigen.

[0027] The term "auto-antigen" refers to an antigen that comprises an epitope of a self antigen or an immunologically reactive epitope that mimics that of a self antigen. In some embodiments the term auto antigen comprises antigens to which autoantibodies are produced. In some embodiments an auto antigen comprises an endogenous antigen wherein the animal from which the endogenous antigen originated is or was once immunologically tolerant to the antigen.

[0028] The term “anergy” refers to a state wherein lymphocytes are inactive (e.g. they do not respond to the binding of an antigen). In some embodiments, the lymphocyte is a B-cell. In some embodiments, a B-cell becomes anergic following exposure to a circulating soluble self-antigen. In some embodiments, a B-cell becomes anergic following exposure to a soluble circulating antigen. In some embodiments, the antigen is soluble hen egg lysozyme (sHEL).

[0029] The term “amino acid” refers to naturally occurring and non-naturally occurring amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally encoded amino acids are the 20 common amino acids (alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine) and pyrrolysine and selenocysteine. Amino acid analogs refers to agents that have the same basic chemical structure as a naturally occurring amino acid, i.e., an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, such as, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (such as, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid.

[0030] Amino acids are referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, are referred to by their commonly accepted single-letter codes.

[0031] The terms “polypeptide”, “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to naturally occurring amino acid polymers as well as amino acid polymers in which one or more amino acid residues is a non-naturally occurring amino acid, e.g., an amino acid analog. The terms encompass amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds.

[0032] The terms “Myc protein” and “Myc polypeptide” are used synonyms and refer to the polymer of amino acid residues disclosed in NCBI Accession Number NP002458.2, and functional homologs, analogs or fragments thereof. In some embodiments, a Myc polypeptide comprises an amino acid sequence that is at least 40% to 100% identical, e.g., at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 40% to about 100% identical to the sequence of NCBI Accession Numbers NP002458.2. In some embodiments, a Myc polypeptide comprises a polypeptide sequence of 40 amino acids or more in length that is at least 50% to 100% identical, e.g., at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100% identical to the sequence of NCBI Accession Numbers NP002458.2. In some embodiments, a Myc polypeptide comprises a polypeptide sequence of 40 amino acids or more in length that is at least 50% to 100% identical, e.g., at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100% identical to the sequence of NCBI Accession Numbers

NP002458.2. In some embodiments, Myc polypeptide means a polymer of 439 amino acids, a Myc polypeptide that has not undergone any post-translational modifications, and/or a Myc polypeptide that has undergone post-translational modifications. In certain instances, the Myc polypeptide is 48,804 kDa. In certain instances, the Myc polypeptide contains a basic Helix-Loop-Helix Leucine Zipper (bHLH/LZ) domain. In certain instances, the Myc polypeptide is a transcription factor (e.g. Transcription Factor 64). In certain instances, the Myc polypeptide binds to a sequence comprising CACGTG (i.e. an E-box sequence).

[0033] The term “nucleic acid” refers to deoxyribonucleotides, deoxyribonucleosides, ribonucleosides, or ribonucleotides and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless specifically limited otherwise, the term also refers to oligonucleotide analogs including PNA (peptidonucleic acid), analogs of DNA used in antisense technology (phosphorothioates, phosphoroamidates, and the like). Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (including but not limited to, degenerate codon substitutions) and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions are achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, Nucleic Acid Res. 19:5081 (1991); Ohtsuka *et al.*, J. Biol. Chem. 260:2605-2608 (1985); and Cassol *et al.* (1992); Rossolini *et al.*, Mol. Cell. Probes 8:91-98 (1994)).

[0034] The terms “MYC” and “MYC gene” are synonyms. They refer to a nucleic acid sequence that encodes a Myc polypeptide. A MYC gene comprises a nucleotide sequence of at least 120 nucleotides that is at least 60% to 100% identical or homologous, e.g., at least 60, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 70% to about 100% identical to sequences of NCBI Accession Number NM_002467. In some embodiments, the MYC gene is a proto-oncogene. In certain instances, a MYC gene is found on chromosome 8, at 8q24.21. In certain instances, a MYC gene begins at 128,816,862 bp from *pter* and ends at 128,822,856 bp from *pter*. In certain instances, a MYC gene is about 6 kb. In certain instances, a MYC gene encodes at least eight separate mRNA sequences – 5 alternatively spliced variants and 3 unspliced variants.

[0035] To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences can be aligned for optimal comparison purposes (e.g., gaps are introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions *can* then be compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent homology between the two sequences is a function of the number of identical positions

shared by the sequences (i.e., % identity = # of identical positions/total # of positions (e.g., overlapping positions) x 100). In some embodiments the two sequences are the same length.

[0036] To determine percent homology between two sequences, the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci.

5 USA 90:5873-5877 is used. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, *et al.* (1990) J. Mol. Biol. 215:403-410. BLAST nucleotide searches are performed with the

NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to a nucleic acid molecules described or disclose herein. BLAST protein searches are performed with the XBLAST program,

score=50, wordlength=3. To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized

10 as described in Altschul *et al.* (1997) Nucleic Acids Res. 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used.

See the website of the National Center for Biotechnology Information for further details (on the World Wide Web at ncbi.nlm.nih.gov). Proteins suitable for use in the methods described herein also includes proteins

having between 1 to 15 amino acid changes, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 amino acid

15 substitutions, deletions, or additions, compared to the amino acid sequence of any protein described herein. In

other embodiments, the altered amino acid sequence is at least 75% identical, e.g., 77%, 80%, 82%, 85%,

88%, 90%, 92%, 95%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of any protein

inhibitor described herein. Such sequence-variant proteins are suitable for the methods described herein as

long as the altered amino acid sequence retains sufficient biological activity to be functional in the

20 compositions and methods described herein. Where amino acid substitutions are made, the substitutions

should be conservative amino acid substitutions. Among the common amino acids, for example, a

“conservative amino acid substitution” is illustrated by a substitution among amino acids within each of the

following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and

tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and

25 (6) lysine, arginine and histidine. The BLOSUM62 table is an amino acid substitution matrix derived from

about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff *et al* (1992), *Proc. Natl Acad. Sci. USA*, 89:10915-

10919). Accordingly, the BLOSUM62 substitution frequencies are used to define conservative amino acid

substitutions that, in some embodiments, are introduced into the amino acid sequences described or disclosed

30 herein. Although it is possible to design amino acid substitutions based solely upon chemical properties (as

discussed above), the language “conservative amino acid substitution” preferably refers to a substitution

represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is

conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. According to this

system, preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 1

35 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a

BLOSUM62 value of at least 2 (e.g., 2 or 3).

[0037] The term “Myc activity” refers to binding of a Myc polypeptide to a nucleic acid sequence. In some embodiment, MYC activity further includes MYC regulation of the transcriptional activity of Myc responsive genes. In some embodiments, Myc activity induces cell proliferation and/or antibody production.

[0038] The terms “activation of Myc” and “Myc activation” refer to the induction of Myc activity. In some 5 embodiments activation of Myc is induced by over-expression of a Myc polypeptide. In some embodiments activation of Myc is induced by transport of a Myc polypeptide into the nucleus of a cell. In some embodiments activation of Myc is induced by transport of a Myc polypeptide into a cell.

[0039] The term “expression” refers to one or more of the following events: (1) production of an RNA 10 template from a DNA sequence (e.g., by transcription) within a cell; (2) processing of an RNA transcript (e.g., by splicing, editing, 5' cap formation, and/or 3' end formation) within a cell; (3) translation of an RNA sequence into a polypeptide or protein within a cell; (4) post-translational modification of a polypeptide or protein within a cell; (5) presentation of a polypeptide or protein on the cell surface; (6) secretion or release of a polypeptide or protein from a cell.

[0040] The term “endogenous” in the context of a cellular protein refers to protein naturally occurring and/or 15 expressed by the cell in the absence of recombinant manipulation; accordingly, the terms “endogenously expressed protein” or “endogenous protein” excludes cellular proteins expressed by means of recombinant technology.

[0041] The term “factor-dependent cell” means a cell that is not able to survive and/or proliferate without 20 contact with an exogenous growth factor (e.g. a cytokine). When contacted with the required growth-factors, a factor-dependent cell is able to survive and/or proliferate. In the absence of the required growth-factors, a factor-dependent cell cannot survive (e.g. will undergo apoptosis) and/or proliferate.

[0042] The term “expansion” as used herein refers to an enlargement in the size of the culture due to the 25 proliferation (i.e. division) of cells in the cell culture. In some embodiments, the cell culture exhibits expansion (i.e. it attains a larger size and/or the concentration of cell in the culture increases). In some embodiments, the cell culture exhibits negative expansion (i.e. attains a smaller size or the concentration of cell in the culture decreases). In certain instances, the expansion of a cell culture is detectable and measurable in any suitable manner (e.g. staining, flow cytometry, fluorescent microscopy, HPLC, confocal microscopy, infrared spectroscopy, autoradiography).

[0043] The phrases “E-box sequence” and “enhancer box sequence” are used interchangeably herein and 30 mean the nucleotide sequence CANNTG, wherein N is any nucleotide. In certain instances, the E-box sequence comprises CACGTG. In certain instances, the basic helix-loop-helix domain of a transcription factor encoded by MYC binds to the E-box sequence. In certain instances the E-box sequence is located upstream of a gene (e.g. p21, Bcl-2, or ornithine decarboxylase). In certain instances, the binding of the transcription factor encoded by MYC to the E-box sequence, allows RNA polymerase to transcribe the gene 35 downstream of the E-box sequence.

Methods of Modulating the Viability of a Cell

[0044] Disclosed herein, in some embodiments, are methods of modulating the viability of a cell. As used herein, “viability” means the length of time the cell survives, the rate (or amount) of cellular proliferation, or a combination thereof. In some embodiments, the viability of a cell is increased. In some embodiments, the viability of a cell is decreased.

[0045] In some embodiments, the modulation occurs *in vivo*. In some embodiments, the modulation occurs *in vitro*. In some embodiments, a method of modulating the viability of a cell comprises modulating the viability of a leukocyte. In some embodiments, the leukocyte is a T-cell. In some embodiments, the leukocyte is a CD4+ T-cell. In some embodiments, the leukocyte is a memory T-cell.

[0046] In some embodiments, modulating the viability of a cell comprises modulating the expression of a MYC gene, and/or the activity of a Myc polypeptide. In some embodiments, modulating the expression of a MYC gene, and/or the activity of a Myc polypeptide directly or indirectly results in the modulation of a gene or polypeptide regulated by MYC. In some embodiments, the gene or polypeptide directly or indirectly regulated by a MYC transcription factor is: neurogranin, MEF-2c, gfi-1, cyclin D2, CDK4, Egr-2, Nab-2, TGIF, NF-kB p105, Carma-1, A1 or Bcl-2.

Increasing the Viability of a Cell

[0047] In some embodiments, modulating the viability of a cell means increasing the viability of a cell. In some embodiments, increasing the viability of a cell means: increasing the length of time the cell survives (e.g. as compared to an identical or similar unmodulated cell of the same type), increasing the rate or amount of cellular proliferation (e.g. as compared to an identical or similar unmodulated cell of the same type), or a combination thereof.

[0048] In some embodiments, increasing the viability of a cell comprises contacting a cell with an agent that increases the nucleic concentration of a Myc polypeptide. In some embodiments, increasing the viability of a cell comprises contacting a cell with: an exogenous Myc polypeptide, an agent that increases the expression of a MYC gene, and an agent that increases the activity of a Myc polypeptide, or a combination thereof (the “MYC Increasing Agent”). In some embodiments, the MYC Increasing Agent is a small molecule, a peptide, an antibody, or a combination thereof.

[0049] In some embodiments, a method of increasing the viability of a cell comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an exogenous Myc polypeptide. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a TAT sequence; and (b) a MYC sequence.

[0050] In some embodiments, a method of increasing the viability of a cell comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a MAD-1

gene. In certain instances, the down-regulation of a MAD-1 gene and/or polypeptide upregulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0051] In some embodiments, a method of increasing the viability of a cell comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a Mxi-1 gene. In certain instances, the down-regulation of a Mxi-1 gene and/or polypeptide upregulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0052] In some embodiments, a method of increasing the viability of a cell further comprises identifying the MYC Increasing Agent. In some embodiments, the method of identifying the MYC Increasing Agent comprises identifying an agent that reverses anergy in a B-cell. In some embodiments, the method of identifying the MYC Modulating Agent comprises identifying an agent that increases the viability of a factor-dependent cell in the absence of a required growth factor. In some embodiments, the method of identifying the MYC Increasing Agent comprises identifying an agent that induces expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence).

[0053] In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by about 1 to about 20 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by about 1 to about 15 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by about 1 to about 10 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by about 1 to about 5 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by about 1 to about 4 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, following contact with the MYC Increasing Agent the viability of a cell is increased by 1 to about 2 times (e.g. as compared to an identical or similar unmodulated cell of the same type).

[0054] In some embodiments, the cell is a T-cell. In some embodiments, the T-cell is a memory T-cell. In some embodiments, modulating the viability of a T-cell means extending the life-span of a T-cell. In some embodiments, extending the life span of a memory T-cell results in a higher concentration of memory T-cells in a body. In certain instances, the higher concentration of memory T-cells results in an accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in anergic T-cells. In certain instances, the decrease in anergic T-cells results in an accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in the time it takes a T-cell to activate in response to an antigen. In certain instances, the decrease in the time it takes a T-cell to activate in response to an antigen results in an accelerated primary immune response to antigen.

Decreasing the Viability of a Cell

[0055] In some embodiments, modulating the viability of a cell means decreasing the viability of a cell. In some embodiments, decreasing the viability of a cell means: decreasing the length of time the cell survives and decreasing the rate or amount of cellular proliferation (e.g., as compared to an identical or similar unmodulated cell of the same type). In some embodiments, decreasing the viability of a cell means
5 decreasing the length of time the cell survives and decreasing the rate or amount of cellular proliferation (e.g. as compared to an identical or similar unmodulated cell of the same type).

[0056] In some embodiments, decreasing the viability of a cell comprises contacting a cell with an agent that decreases the nucleic concentration of MYC. In some embodiments, decreasing the viability of a cell comprises contacting a cell with: an agent that decreases the expression of a MYC gene, an agent that
10 decreases the activity of a Myc polypeptide, or a combination thereof (the "MYC Decreasing Agent"). In some embodiments, the MYC Decreasing Agent is a small molecule, a biologic, a peptide, an antibody, or a combination thereof.

[0057] In some embodiments, a method of decreasing the viability of a cell comprises contacting a cell with the MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is an agonist of a MAD-1
15 gene, a MAD-1 polypeptide, or a combination thereof. In certain instances, the up-regulation of a MAD-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene. In some embodiments, a method of modulating the viability of a cell comprises contacting a cell with the MYC Decreasing Agent.

[0058] In some embodiments, a method of decreasing the viability of a cell comprises contacting a cell with
20 the MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is: Mxi-1 an agonist of a Mxi-1 gene and a Mxi-1 polypeptide, or a combination thereof. In certain instances, the up-regulation of a Mxi-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0059] In some embodiments, the method of identifying the MYC Decreasing Agent comprises identifying
25 an agent that decreases the viability of a factor-dependent cell when the factor-dependent cell is contacted with the necessary growth factor. In some embodiments, the method of identifying the MYC Decreasing Agent comprises identifying an agent that inhibits expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence).

[0060] In some embodiments, the viability of a cell is decreased by about 1 to about 25 times (e.g. as
30 compared to an identical or similar unmodulated cell of the same type). In some embodiments, the viability of a cell is decreased by about 1 to about 20 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, the viability of a cell is decreased by about 1 to about 15 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, the viability of a cell is decreased by about 1 to about 10 times (e.g. as compared to an identical or similar unmodulated
35 cell of the same type). In some embodiments, the viability of a cell is decreased by about 1 to about 5 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, the

viability of a cell is decreased by about 1 to about 4 times (e.g. as compared to an identical or similar unmodulated cell of the same type). In some embodiments, the viability of a cell is decreased by about 1 to about 2 times (e.g. as compared to an identical or similar unmodulated cell of the same type).

[0061] In some embodiments, the cell is a T-cell. In some embodiments, the T-cell is a memory T-cell. In certain instances, the down-regulation of a MYC gene results in a decreased life-span for the T-cell. In certain instances, the decreased life span of a memory T-cell results in a lower concentration of memory T-cells in a body. In certain instances, the lower concentration of memory T-cells results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in anergic T-cells. In certain instances, the increase in anergic T-cells results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in the time it takes a T-cell to activate in response to an antigen. In certain instances, the increase in the time it takes a T-cell to activate in response to an antigen results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression.

Methods of Modulating an Immune Response

[0062] Disclosed herein, in certain embodiments, is a method of modulating an immune response. In some embodiments, the immune response is an adaptive immune response (e.g. memory B-cells and memory T-cells). In some embodiments, modulating an immune response means increasing (e.g., inducing, supplementing, amplifying) an immune response. In some embodiments, modulating an immune response means decreasing (e.g., inhibiting or suppressing) an immune response. In some embodiments, the modulation occurs *in vitro*. In some embodiments, the modulation occurs *in vivo*.

[0063] In some embodiments, a method of modulating an immune response comprises modulating the viability of a lymphocyte. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a CD4+ T-cell. In some embodiments, the T-cell is a memory T-cell.

[0064] In some embodiments, modulating an immune response comprises modulating the expression of a MYC gene, and/or the activity of a Myc polypeptide. In some embodiments, modulating the expression of a MYC gene, and/or the activity of a Myc polypeptide directly or indirectly results in the modulation of a gene or polypeptide regulated by MYC. In some embodiments, the gene or polypeptide directly or indirectly regulated by a MYC transcription factor is neurogranin, MEF-2c, gfi-1, cyclin D2, CDK4, Egr-2, Nab-2, TGIF, NF-kB p105, Carma-1, A1 or Bcl-2.

[0065] In some embodiments, the subject is a human.

Inducing an Immune Response

[0066] In some embodiments, modulating an immune response means increasing (e.g., inducing, supplementing, amplifying) an immune response. In some embodiments, increasing an immune response

includes increasing the lifespan of a T-cell, increasing the rate of proliferation by a T-cell, increasing the rate at which a memory T-cell responds to an antigen, decreasing the number of anergic cells, and/or increasing the rate at which a cell ends anergy (e.g. as compared to an identical or similar unmodulated cell of the same type).

5 [0067] In some embodiments, increasing an immune response comprises administering to an individual in need thereof an agent that increases the nucleic concentration of MYC. In some embodiments, increasing an immune response comprises administering to an individual in need thereof: exogenous MYC, an agent that increases the expression of a MYC gene, and an agent that increases the activity of MYC (the “MYC Increasing Agent”). In some embodiments, the MYC Increasing Agent is a small molecule, a biologic, a
10 peptide, an antibody, or a combination thereof.

[0068] In some embodiments, a method of increasing an immune response comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an exogenous Myc polypeptide. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the
15 transporter peptide sequence and the MYC sequence. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a TAT sequence; and (b) a MYC sequence.

[0069] In some embodiments, a method of increasing an immune response comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a MAD-1 gene. In certain instances, the down-regulation of a MAD-1 gene and/or polypeptide upregulates the
20 expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0070] In some embodiments, a method of increasing an immune response comprises contacting a cell with the MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a Mxi-1 gene. In certain instances, the down-regulation of a Mxi-1 gene and/or polypeptide upregulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

25 [0071] In some embodiments, following contact with the MYC Increasing Agent, an immune response is increased by more than 1 to about 20 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Increasing Agent an immune response is increased by more than 1 to about 15 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC
30 Increasing Agent an immune response is increased by more than 1 to about 10 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Increasing Agent an immune response is increased by more than 1 to about 5 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some
35 embodiments, following contact with the MYC Increasing Agent an immune response is increased by more than 1 to about 4 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Increasing Agent an immune response is

increased by more than 1 to about 2 times (e.g. as compared to an identical or similar unmodulated immune response of the same type).

[0072] In some embodiments, a method of increasing an immune response comprises administering a MYC Increasing Agent to an individual in need thereof. In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 10 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 20 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 15 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 10 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 5 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 4 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Increasing Agent to an individual in need thereof, an immune response is increased by more than 1 to about 2 times (e.g. as compared to an identical or similar unmodulated immune response of the same type).

[0073] In some embodiments, a method of increasing an immune response further comprises co-administering a vaccine. In some embodiments, the vaccine is administered before, after, or simultaneously with a MYC Increasing Agent. In some embodiments, the MYC Increasing Agent stimulates the immune system and increases the response of the immune system to a vaccine. In some embodiments, the MYC Increasing Agent augments an immune response. In some embodiments, the MYC Increasing Agent acts synergistically with the vaccine. In some embodiments, the agent is a vaccine adjuvant.

[0074] In some embodiments, a vaccine comprises dead microorganisms, attenuated microorganisms, toxoids, subunits of the pathogen, nucleic acids, or combinations thereof. In some embodiments, the vaccine is a vaccine for hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; influenza; meningococcal disorder; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus (HPV); malaria; leishmania; *Candida albicans*; an allergen; or combinations thereof. In some embodiments, the vaccine is a vaccine for a cancer (e.g. Follicular B-cell Non-Hodgkin's Lymphoma, prostate cancer, multiple myeloma, kidney cancer, cutaneous melanoma, and ocular melanoma). In some embodiments, a cancer vaccine is a patient-specific vaccine (e.g. the vaccine comprises a patient's own tumor cells). In some embodiments, a cancer vaccine

comprises Prostate Specific Antigen (PSA). In some embodiments, a cancer vaccine comprises sialyl Tn (STn). In some embodiments, a cancer vaccine comprises Heat Shock Proteins (HSPs) (e.g., gp96). In some embodiments, a cancer vaccine comprises ganglioside molecules (e.g., GM2, GD2, and GD3). In some embodiments, a cancer vaccine comprises carcinoembryonic antigen (CEA). In some embodiments, a cancer vaccine comprises MART-1 (also known as Melan-A). In some embodiments, a cancer vaccine comprises tyrosinase. In some embodiments, the vaccine is a DNA vaccine.

[0075] In some embodiments, the vaccine comprises an antigenic moiety. In some embodiments, the antigenic moiety is a toxoid, a peptide, a nucleic acid sequence, a polysaccharide, or a combination thereof. In some embodiments, the antigenic moiety is derived from a pathogen selected from: hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; meningococcal; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus; or combinations thereof. In some embodiments, the antigenic moiety is derived a neoplastic cell. In some embodiments, the antigenic moiety is a nucleic acid or a polymer of nucleic acids.

[0076] In some embodiments, increasing an immune response in an individual receiving vaccination against an antigen results in an increase in the viability (and thus concentration) of memory T-cells, B-cells, or a combination thereof against that antigen. In certain instances, increasing an immune response in an individual receiving vaccination against an antigen results in accelerated activation of the T-cell by the antigen. In certain instances, increasing an immune response in an individual receiving vaccination against an antigen results in a decrease in anergic T-cells.

[0077] In some embodiments, a method of increasing an immune response further comprises identifying the MYC Increasing Agent. In some embodiments, the method of identifying the MYC Increasing Agent comprises identifying an agent that reverses anergy in a B-cell. In some embodiments, the method of identifying the MYC Increasing Agent comprises identifying an agent that increases the viability of a factor-dependent cell in the absence of a required growth factor. In some embodiments, the method of identifying the MYC Increasing Agent comprises identifying an agent that induces expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence).

Decreasing an Immune Response

[0078] In some embodiments, modulating an immune response means decreasing (e.g., inhibiting or suppressing) an immune response. In some embodiments, decreasing an immune response includes decreasing the lifespan of a T-cell, decreasing proliferation by a T-cell, decreasing the rate at which a memory T-cell responds to an antigen, increasing the number of anergic cells, and/or decreasing the rate at which a cell ends anergy (e.g. as compared to an identical or similar unmodulated cell of the same type).

[0079] In some embodiments, decreasing an immune response comprises administering to an individual in need thereof an agent that decreases the nucleic concentration of MYC. In some embodiments, decreasing an immune response comprises administering to an individual in need thereof: an agent that decreases the expression of a MYC gene and an agent that decreases the activity of a Myc polypeptide (the "MYC

Decreasing Agent”). In some embodiments, the MYC Decreasing Agent is a small molecule, a biologic, a peptide, an antibody, or a combination thereof.

[0080] In some embodiments, a method of decreasing an immune response comprises contacting a cell with the MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is: an agonist of a MAD-1 gene, a MAD-1 polypeptide, or a combination thereof. In certain instances, the up-regulation of a MAD-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0081] In some embodiments, a method of decreasing an immune response comprises contacting a cell with the MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is: Mxi-1 an agonist of a Mxi-1 gene and a Mxi-1 polypeptide, or a combination thereof. In certain instances, the up-regulation of a Mxi-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0082] In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 25 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 20 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 15 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 10 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 5 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 4 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following contact with the MYC Decreasing Agent an immune response is decreased by more than 1 to about 2 times (e.g. as compared to an identical or similar unmodulated immune response of the same type).

[0083] In some embodiments, a method of decreasing an immune response comprises administering a MYC Decreasing Agent to an individual in need thereof. In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 25 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 20 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1

to about 15 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 10 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 5 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 4 times (e.g. as compared to an identical or similar unmodulated immune response of the same type). In some embodiments, following administration of the MYC Decreasing Agent to an individual in need thereof an immune response is increased by more than 1 to about 2 times (e.g. as compared to an identical or similar unmodulated immune response of the same type).

[0084] In some embodiments, a method of decreasing an immune response comprises administering the MYC Decreasing Agent to an individual with an autoimmune disorder. In some embodiments, the MYC Decreasing Agent decreases an immune response. In certain instances, decreasing an immune response in an individual with an autoimmune disorder ameliorates and/or prevents an immune response against self antigens by the subject's immune system. In some embodiments, the autoimmune disorder is Castleman's Disorder, lupus, multiple sclerosis, scleroderma pigmentosa, Autoimmune Lymphoproliferative Syndrome (ALPS), myasthenia gravis, diabetes, asthma, rheumatoid arthritis, vitiligo, diGeorge's syndrome, Grave's disorder, Crohn's disorder, inflammatory bowel disorder, colitis, orchitis, scleroderma pigmentosa, uveitis, Post-Transplant Lymphoproliferative Disorder (PTLD), or Autoimmune Disorder-Associated Lymphadenopathy (ADAL).

[0085] In some embodiments, a method of decreasing an immune response comprises administering the MYC Decreasing Agent to an individual that is receiving, will receive, or has received an organ or bone marrow transplant (the "Transplant"). In some embodiments, the MYC Decreasing Agent decreases an immune response. In certain instances, decreasing an immune response in an individual that is receiving, will receive, or has received an organ transplant, or a bone marrow transplant ameliorates and/or prevents an immune response against the Transplant.

[0086] In some embodiments, the method of identifying the MYC Decreasing Agent comprises identifying an agent that decreases the viability of a factor-dependent cell when the factor-dependent cell is contacted with the necessary growth factor. In some embodiments, the method of identifying the MYC Decreasing Agent comprises identifying an agent that inhibits expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence).

Methods of Modulating a Disorder Characterized by the Aberrant Expression of a MYC Gene or the Aberrant Activity of a MYC Polypeptide

[0087] Disclosed herein, in certain embodiments, are methods of treating a disorder characterized by an over-expression of MYC or an excess of Myc polypeptide activity. Further disclosed herein, in certain
5 embodiments, are methods of treating a disorder characterized by an under-expression of MYC or a dearth of Myc polypeptide activity. In some embodiments, the method comprises administering to an individual in need thereof an effective amount of an agent that modulates the expression of a MYC gene, and/or the activity of a Myc polypeptide.

[0088] Disclosed herein, in certain embodiments, are methods of treating a disorder characterized by an
10 under-expression of MYC or a dearth of Myc polypeptide activity. In some embodiments, the method comprises administering to an individual in need thereof an effective amount of a MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is a small molecule, a biologic, a peptide, an antibody, or a combination thereof.

[0089] In some embodiments, the method comprises administering to an individual in need thereof an
15 effective amount of a MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an exogenous Myc polypeptide. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a transporter peptide sequence; (b) a MYC sequence; and optionally (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the MYC Increasing Agent is a fusion peptide comprising (a) a TAT sequence; and (b) a MYC sequence.

[0090] In some embodiments, the method comprises administering to an individual in need thereof an
20 effective amount of a MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a MAD-1 gene. In certain instances, the down-regulation of a MAD-1 gene and/or polypeptide upregulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0091] In some embodiments, the method comprises administering to an individual in need thereof an
25 effective amount of a MYC Increasing Agent. In some embodiments, the MYC Increasing Agent is an antagonist of a Mxi-1 gene. In certain instances, the down-regulation of a Mxi-1 gene and/or polypeptide upregulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0092] Disclosed herein, in certain embodiments, are methods of treating a disorder characterized by an
30 over-expression of MYC or an excess of Myc polypeptide activity. In some embodiments, the method comprises administering to an individual in need thereof an effective amount of a MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is a small molecule, a biologic, a peptide, an antibody, or a combination thereof.

[0093] In some embodiments, the method comprises administering to an individual in need thereof an
35 effective amount of a MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is: an agonist of a MAD-1 gene, a MAD-1 polypeptide, or a combination thereof. In certain instances, the up-

regulation of a MAD-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0094] In some embodiments, the method comprises administering to an individual in need thereof an effective amount of a MYC Decreasing Agent. In some embodiments, a method of modulating the viability of a cell comprises contacting a cell with the MYC Decreasing Agent. In some embodiments, the MYC Decreasing Agent is: Mxi-1an agonist of a Mxi-1 gene and a Mxi-1 polypeptide, or a combination thereof. In certain instances, the up-regulation of a Mxi-1 gene and/or polypeptide down-regulates the expression of a MYC gene and/or a polypeptide encoded by a MYC gene.

[0095] In some embodiments, an indication that an individual over-expresses MYC or that a Myc polypeptide is excessively active is the development of an autoimmune disorder. In some embodiments, the autoimmune disorder is Castleman's Disease, lupus, multiple sclerosis, scleroderma pigmentosa, Autoimmune Lymphoproliferative Syndrome (ALPS), myesthenia gravis, diabetes, asthma, rheumatoid arthritis, vitiligo, diGeorge's syndrome, Grave's disease, Crohn's disease, inflammatory bowel disease, colitis, orchitis, scleroderma pigmentosa, uveitis, Post-Transplant Lymphoproliferative Disease (PTLD), Autoimmune Disease-Associated Lymphadenopathy (ADAL), rejection of an organ transplant, rejection of a tissue transplant, or combinations thereof.

[0096] In some embodiments, an indication that an individual under-expresses MYC, or that a Myc polypeptide is minimally active or inactive is an increase in the number of anergic B-cells and/or T-cells. In certain instances, the expression of IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2) is down-regulated in anergic B-cells when compared to non-anergic B-cells. In certain instances, reversing (e.g. partially or fully) anergy results in an increase in the expression of IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2).

[0097] In some embodiments, a method of treating a disorder characterized by aberrant expression of a MYC gene or aberrant activity of a Myc polypeptide comprises identifying an agent that reverses anergy in a B-cell. In some embodiments, a method of treating a disorder characterized by aberrant expression of a MYC gene or aberrant activity of a Myc polypeptide comprises identifying an agent that increases the viability of a factor-dependent cell in the absence of a required growth factor. In some embodiments, a method of treating a disorder characterized by aberrant expression of a MYC gene or aberrant activity of a Myc polypeptide comprises identifying an agent that decreases the viability of a factor-dependent cell when the factor-dependent cell is contacted with the necessary growth factor. In some embodiments, a method of treating a disorder characterized by aberrant expression of a MYC gene or aberrant activity of a Myc polypeptide comprises identifying an agent that induces expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence). In some embodiments, a method of treating a disorder characterized by aberrant expression of a MYC gene or aberrant activity of a Myc polypeptide comprises identifying an agent that inhibits expression of a reporter gene under the control of a MYC responsive promoter (e.g. an E-box sequence).

Exogenous MYC

[0098] Disclosed herein, in certain embodiments, is a fusion peptide comprising (a) a transporter peptide sequence; and (b) a MYC sequence. In some embodiments, the fusion peptide is a peptide of Formula (I):

5 transporter peptide sequence -MYC sequence.

[0099] In some embodiments, a fusion peptide disclosed herein comprises (a) a transporter peptide sequence; (b) a MYC sequence; and (c) one or more molecules that link the transporter peptide sequence and the MYC sequence. In some embodiments, the fusion peptide is a peptide of Formula (II):

transporter peptide sequence-X-MYC sequence,

10 wherein -X- is molecule that links the transporter peptide sequence and the MYC sequence. In some embodiments, -X- is an amino acid. In some embodiments, -X- is at least one amino acid.

[00100] As used herein, a “transporter peptide” means a peptide sequence that promotes peptide penetration into cells and tissues. In some embodiments, a transporter peptide is TAT. In some embodiments, a transporter peptide is TAT_[48-57]. In some embodiments, a transporter peptide is TAT_[57-48].

15 **[00101]** As used herein, a “MYC sequence” is a MYC amino acid peptide sequence. In some embodiments, the Myc polypeptide is a complete Myc polypeptide sequence. In some embodiments, the Myc polypeptide is a partial Myc polypeptide sequence. In some embodiments, the MYC is c-MYC. In some embodiments, the Myc polypeptide sequence comprises SEQ ID NO. 1:

MDFFRVVENQQPPATMPLNVSFTNRNYDL DYDSVQPYFYCDEEENF
 20 YQQQQQSELQPPASEDIWKKFELLPTPPLSPSRRSGLCSPSYVAVTP
 FSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIK
 NIIIQDCMWSGFSAAKLVSEKLASYQAARKDSGSPNPARGHSVCST
 SSLYLQDL SAAASECIDPSVVPYPLNDSSSPKSCASQDSSAFSPSSDS
 LLSTESSPQGSPEPLVLHEETPPTSSDSEEEQEDEEEIDVVSVEKRQ
 25 APGKRSESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPSTRKD
 YPAAKRVKLD SVR VLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQ
 RRNELKRSFFALRDQIPELENNEKAPKV VILKKATAYILSVQAEEQKL
 ISEEDLLRKRREQLKHKLEQLRKGELNSKLE.

[00102] In some embodiments, a fusion peptide disclosed herein comprises (a) TAT, and (b) c-MYC. In some
 30 embodiments, a fusion peptide disclosed herein comprises (a) TAT_[48-57], and (b) c-MYC. In some
 embodiments, a fusion peptide disclosed herein comprises (a) TAT_[57-48], and (b) c-MYC.

[00103] In some embodiments, a fusion peptide disclosed herein comprises (a) TAT, (b) a linker amino acid,
 and (c) c-MYC. In some embodiments, a fusion peptide disclosed herein comprises (a) TAT_[48-57], (b) a linker
 amino acid, and (c) c-MYC. In some embodiments, a fusion peptide disclosed herein comprises (a) TAT_[57-48],
 35 (b) a linker amino acid, and (c) c-MYC.

[0104] In some embodiments, a fusion peptide disclosed herein further comprises at least one amino acid sequence that facilitates purification of the fusion protein. In some embodiments, a fusion peptide disclosed herein comprises a protein tag. In some embodiments, a fusion peptide disclosed herein comprises a polyhistidine tag. In some
 5 embodiments, a fusion peptide disclosed herein comprises an epitope tag. In some embodiments, a fusion peptide disclosed herein comprises a polyhistidine tag and an epitope tag. In some embodiments, a fusion peptide disclosed herein comprises a 6-histidine tag (SEQ ID NO: 3) and a V5 epitope tag.

[0105] In some embodiments, the histidine tag is a 6-histidine tag (SEQ ID NO: 3). In some
 10 embodiments, the histidine tag comprises the sequence HHHHHH (SEQ ID NO: 3). In some embodiments, a histidine tag is added to a fusion protein disclosed herein by any suitable method. In some embodiments, a TAT-Myc polypeptide sequence is cloned into an expression vector encoding a polyHis-tag. In some embodiments, a polyhistidine tag is added by PCR (i.e., the PCR primers comprise a polyhistidine sequence).

[0106] In some embodiments, a fusion peptide disclosed herein further comprises at least
 15 one protein tag. In some embodiments, a fusion peptide disclosed herein comprises an epitope tag. In some embodiments, a fusion peptide disclosed herein further comprises a V5 epitope tag. In some embodiments, the V5 tag comprises the amino acids: GKPIPPLLGLDST (SEQ ID NO: 4). In some embodiments, the V5 tag comprises the
 20 amino acids: IPNPLLGLD (SEQ ID NO: 5). In some embodiments, a V5 tag is added to a fusion protein disclosed herein by any suitable method. In some embodiments, a TAT-Myc polypeptide sequence is cloned into an expression vector encoding a V5 tag. In some embodiments, a V5 tag is added by PCR (i.e., the PCR primers comprise a V5 sequence).

[0107] In some embodiments, the amino acids are in the D formation. In some
 25 embodiments, the amino acids are in the L formation. In some embodiments, a first plurality of amino acids are in the D formation and a second plurality are in the L formation.

[0108] In some embodiments, the MYC Increasing Agent comprises SEQ ID NO. 2:

MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTRNYDLDYDSVQPYFYCDEEE
 NFYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGLCSPSYVAVTPFSLRGDND
 30 GGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSAAAK
 LVSEKLSYQAARKDSGSPNPARGHSVCSTSSLYLQDLSAAASECIDPSVVFYPLN
 DSSSPKSCASQDSSAFSPSSDLSLSTESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEE
 EIDVVSVEKRQAPGKRSESGSPSAGGHKPPHSPLVLKRCHVSTHQHNYAAPPSTR
 KDYPAAKRVKLDVSRVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELK
 35 RSFFALRDQIPELENNEKAPKVVILKKATAYILSVQAEEQKLISEEDLLRKRREQLKH
 KLEQLRKGELNSKLEGKPIPPLLGLDSTRTGHHHHHH.

Construction of a MYC-TAT Peptide

[0109] In some embodiments, a MYC-TAT fusion peptide disclosed herein is constructed
 40 by any suitable method. In some embodiments, a nucleotide sequence encoding a MYC-TAT fusion peptide is generated by PCR. In some embodiments, a forward primer for a

human MYC sequence comprises an in frame N-terminal 9-amino-acid sequence of the TAT protein transduction domain (i.e., RKKRRQRRR (SEQ ID NO: 6)). In some embodiments, a reverse primer for a human MYC sequence is designed to remove the stop codon. In some embodiments, the PCR product is cloned into any suitable expression vector (hereinafter, p-TAT-MYC). In some embodiments, the expression vector comprises a polyhistidine tag and a V5 tag.

Assays for the Identification of Agents

Reversal of Anergy

[0110] Disclosed herein, in certain embodiments, is a method of identifying an agent that reverses anergy in a B-cell. In certain instances, an agent that reverses anergy in a B-cell increases an immune response and/or increases the viability of a cell. In certain instances, an agent that reverses anergy in a B-cell up-regulates MYC (e.g. a MYC gene and/or a Myc polypeptide).

[0111] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises (a) contacting a plurality of anergic B-cells with an agent; and (b) following contacting the plurality of cells with the agent, detecting and/or measuring the level of expression of one or more cell surface markers (e.g. cell surface markers the presence of which is indicative a non-anergic cell) in the cell culture. In some embodiments, the cell surface marker is IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2).

[0112] In certain instances, the expression of IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2) is down-regulated in anergic B-cells when compared to non-anergic B-cells. In certain instances, reversing (e.g. partially or fully) anergy results in an increase in the expression of IgM, IgMa, IgMb, B220, CD21/35, CD23, CD24 (HSA), CD40, CD69, CD80 and/or CD86 (B7-2).

[0113] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises identifying an agent as an agonist of an immune response if expression of a cell surface marker is upregulated (e.g. as compared to an anergic B-cell not contacted with an agent). In some embodiments, expression of a cell surface marker is upregulated on activated B-cells (when compared to the expression pattern on anergic B-cells) by about 10% to about 100% (e.g. at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 10% to about 100%). In some embodiments, expression of a cell surface marker is upregulated on activated B-cells (when compared to the expression pattern on anergic B-cells) by about 30% to about 100% (e.g. at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 30% to about 100%). In some embodiments, expression of a cell surface marker is upregulated on activated B-cells (when compared

to the expression pattern on anergic B-cells) by about 50% to about 100% (e.g. at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100%). In some embodiments, expression of a cell surface marker is upregulated on activated B-cells (when compared to the expression pattern on anergic B-cells) by about 70% to about 100% (e.g. at least 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 70% to about 100%). In some embodiments, expression of a cell surface marker is upregulated on activated B-cells (when compared to the expression pattern on anergic B-cells) by about 80% to about 100% (e.g. at least 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 80% to about 100%).

[00114] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises identifying an agent as an antagonist of an immune response if expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent). In some embodiments, expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent) by about 10% to about 100% (e.g. at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 10% to about 100%). In some embodiments, expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent) by about 30% to about 100% (e.g. at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 30% to about 100%). In some embodiments, expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent) by about 50% to about 100% (e.g. at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100%). In some embodiments, expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent) by about 70% to about 100% (e.g. at least 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 70% to about 100%). In some embodiments, expression of a cell surface marker is downregulated (e.g. as compared to a non-anergic B-cell not contacted with an agent) by about 80% to about 100% (e.g. at least 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 80% to about 100%).

[00115] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell further comprises comparing (1) the level of expression of the cell surface marker observed in a plurality of anergic B-cells following contacting the plurality of cells with the agent to (2) the level of cell surface marker observed in a control. In some embodiments, the control is a plurality of anergic B-cells not contacted with an agent.

[00116] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises obtaining an anergic B-cell from any suitable source (e.g. a human, a mouse, a rat, or any mammal). In some

embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises obtaining an anergic B-cell from transgenic mice with the phenotype BCR^{HEL}/sHEL. In certain instances, BCR^{HEL} mice express a mature B-cell receptor (BCR) for the antigen hen egg lysozyme (HEL). In certain instances, the HEL BCR is the dominant BCR on the B-cells of the transgenic mice. In certain instances, sHEL mice ubiquitously express a transgene for the soluble form of HEL. In certain instances, the expression of the HEL transgene is operably linked to a metallothionein promoter. In certain instances, the exposure to soluble HEL induces anergy in B-cells. U.S. Patent Application Pub. No. 2008/0070256 is hereby incorporated by reference for the methods of constructing the murine models discussed above. In some embodiments, the anergic B-cells are obtained from the spleen, thymus, bone marrow, lymph nodes, or combinations thereof of the transgenic mice.

[00117] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises detecting and/or measuring the level of expression of one or more cell surface markers (e.g. CD86). In some embodiments, detecting and/or measuring the level of expression of one or more cell surface markers comprises (a) contacting a plurality of anergic B-cells (e.g. a control, or a plurality of anergic B-cells following contact with an agent) with antibodies to a cell surface marker (e.g. antibodies to CD86); (b) washing the cell culture (e.g. rinsing) with buffer (e.g. FACS buffer) after contact with the antibodies; and (c) detecting and/or measuring the amount of antibody/cell surface marker complex. In some embodiments, the antibodies are purchased from a commercial supplier. In some embodiments, the antibodies are generated *in-house*. For methods of generating antibodies, see Kohler *et al.*, Nature, 256:495 (1975); U.S. Pat. No. 4,816,567; or Goding, Monoclonal Antibodies: Principles and Practice (Academic Press, 1986); Ward *et al.*, Nature 341: 544-546 (1989); Huse *et al.*, Science 246: 1275-1281 (1989); McCafferty *et al.*, Nature 348: 552-554 (1990); Clackson *et al.*, Nature, 352:624-628 (1991) Marks *et al.*, J. Mol. Biol., 222:581-597 (1991) all of which are hereby incorporated by reference. In some embodiments, the cell culture is incubated on ice during the contact with the antibodies. In some embodiments, the antibody is isotopically-labeled, radio-labeled, fluorophore-labeled, or biotinylated. In some embodiments, the fluorophore is fluorescein. In certain instances, the cell surface marker/antibody complex is detectable and/or measurable in any suitable manner (e.g. HPLC, fluorescence microscopy, confocal microscopy, microarray scanners, Surface Plasmon Resonance, infrared spectroscopy, or autoradiography).

[00118] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises detecting and/or measuring the level of expression of IgM. In some embodiments, detecting and/or measuring the level of expression of IgM comprises (a) contacting a cell culture (e.g. a control, or a plurality of anergic B-cells following contact with an agent) with an antigen (e.g. HEL) for a period of time sufficient to allow the antigen to bind to an IgM; (b) washing the cell culture (e.g. rinsing) with buffer (e.g. FACS buffer) after contacting the culture with the antigen; and (c) detecting and/or measuring the amount of antigen and/or antigen/IgM complex. In some embodiments, the antigen is biotinylated. In some embodiments, the cell culture is further incubated with a marker having an affinity for biotin (the "Biotin Marker"). In some

embodiments, the Biotin Marker is avidin or streptavidin. In some embodiments, the Biotin Marker is isotopically-labeled, radio-labeled, fluorophore-labeled. In some embodiments, the fluorophore is phycoerythrin. In certain instances, the IgM/antigen/Biotin Marker complex is detectable and/or measurable in any suitable manner (e.g. HPLC, fluorescence microscopy, confocal microscopy, microarray scanners, Surface Plasmon Resonance, infrared spectroscopy, or autoradiography).

[00119] In some embodiments, a method of identifying an agent that reverses anergy in a B-cell comprises detecting and/or measuring the level of expression of IgM. In some embodiments, detecting and/or measuring the level of expression of IgM comprises (a) contacting a cell culture (e.g. a control, or a plurality of anergic B-cells following contact with an agent) with an antigen (e.g. HEL) for a period of time sufficient to allow the antigen to bind to the IgM; (b) washing the cell culture (e.g. rinsing) with buffer (e.g. FACS buffer) after contacting the culture with the antigen; and (c) detecting and/or measuring the amount of antigen and/or antigen/IgM complex. In some embodiments, the culture is contacted with dilutions of antibodies to the antigen (e.g. for a period of time sufficient to allow the antibodies to bind to the antigen). In some embodiments, the cell culture is incubated on ice during contact with the antibodies. In some embodiments, the cell culture is washed (e.g. rinsed) with buffer (e.g. FACS buffer) after contact with the antibodies. In some embodiments, the antibody is isotopically-labeled, radio-labeled, fluorophore-labeled, or biotinylated. In some embodiments, the fluorophore is fluorescein. In certain instances, the antigen/antibody and/or IgM/antigen/antibody complex is detectable and/or measurable in any suitable manner (e.g. HPLC, fluorescence microscopy, confocal microscopy, microarray scanners, Surface Plasmon Resonance, infrared spectroscopy, or autoradiography).

[00120] In some embodiments, the method further comprises detecting the presence of and/or measuring the amount of the labeled antibody, or of a complex formed between the antibody and the HEL antigen. In certain instances, the HEL antigen/antibody complex is detectable and measurable in any suitable manner (e.g. HPLC, fluorescence microscopy, confocal microscopy, microarray scanners, Surface Plasmon Resonance, infrared spectroscopy, or autoradiography).

Modulation of Cytokine Function

[00121] Disclosed herein, in certain embodiments, is a method of identifying an agent that induces (e.g., activates, increases, or supplements) an immune response. In some embodiments, the method comprises identifying an agent that replaces and/or augments cytokine function (the "Cytokine Supplementing Agent").

In certain instances, an agent that replaces and/or augments cytokine function in a factor-dependent cell increases an immune response and/or increases the viability of a cell. In certain instances, an agent that replaces and/or augments cytokine function in a factor-dependent cell up-regulates MYC (e.g. a MYC gene and/or a Myc polypeptide).

[00122] In some embodiments, the method comprises (a) contacting a plurality of factor-dependent cells with an agent; and (b) following contacting the cell with the agent, detecting and/or measuring the level of expansion of the cell culture.

[00123] In some embodiments, the method further comprises identifying an agent as a Cytokine Supplementing Agent if, following contact with the agent, the cell culture expands (e.g. as compared to a cell culture not contacted with an agent). In some embodiments, following contact with the agent the cell culture expands (when compared to a cell culture not contacted with an agent) by about 10% to about 100% (e.g. at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 10% to about 100%). In some embodiments, following contact with the agent the cell culture expands (when compared to a cell culture not contacted with an agent) by about 30% to about 100% (e.g. at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 30% to about 100%). In some embodiments, following contact with the agent the cell culture expands (when compared to a cell culture not contacted with an agent) by about 50% to about 100% (e.g. at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100%). In some embodiments, following contact with the agent the cell culture expands (when compared to a cell culture not contacted with an agent) by about 70% to about 100% (e.g. at least 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 70% to about 100%). In some embodiments, following contact with the agent the cell culture expands (when compared to a cell culture not contacted with an agent) by about 80% to about 100% (e.g. at least 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 80% to about 100%).

[00124] In some embodiments, a method of identifying an agent that that replaces and/or augments cytokine function further comprises comparing (1) the level of expansion observed in a plurality of factor-dependent cells following contacting the plurality of cells with an agent to (2) the level of expansion of a control. In some embodiments, the control is a plurality of factor-dependent cells not contacted with an agent.

[00125] In some embodiments, contacting a factor-dependent cell with an agent that replaces and/or augments cytokine function results in the expression of MYC being upregulated. In certain instances, the up-regulation of MYC results in factor-dependent cells that do not require a cytokine for viability. In certain instances, the up-regulation of MYC results in factor-dependent cells that proliferate in the absence of a cytokine. In certain instances, the up-regulation of MYC results in factor-dependent cells that do not undergo apoptosis in the absence of a cytokine.

[00126] In some embodiments, the factor-dependent cells are lymphoid cells (i.e. derived from cells of the lymphatic system). In some embodiments, the factor-dependent cells is, by way of non-limiting example, IL-2^{-/-}, IL-3^{-/-}, IL-4^{-/-}, IL-5^{-/-}, IL-6^{-/-}, IL-7^{-/-}, IL-8^{-/-}, IL-9^{-/-}, IL-10^{-/-}, IL-11^{-/-}, IL-12^{-/-}, or any combinations thereof. In some embodiments, the factor-dependent cells are derived from a cell line selected from, by way of non-limiting example: 2D6; 2E8; 10B10; ATH8; B13; BAF/3; BALM-4; BCL1; CT.4S; CT6; CTL44; CTLL-2; D1; D10; D36; Da; DS-1; Ea3.17; EL4; FL5.12; HT-2; IC-2; Kitt225; KT-3; L4; L138.8A; LBRM-33; LyD9; MC/9; MLA-144; Mono Mac 6; Nb2; NKC3; PB-1; Pno; PT-18; Ramos;

Sez627; T10; T88; T1165; TALL-103; TF-1; TMD2; TS1; UT-7; XG-1; Y16; or a combination thereof. In some embodiments, the cells are derived from CTLL-2; or BAF/3 cell lines.

[00127] Disclosed herein, in certain embodiments, are methods of identifying an agent that inhibits and/or interferes with cytokine function (the "Cytokine Interfering Agent"). In certain instances, a Cytokine

5 Interfering Agent decreases an immune response and/or decreases the viability of a cell. In some embodiments, the method comprises (a) contacting a plurality of factor-dependent cells with a required growth factor such that the cell culture expands; (b) contacting the plurality of cells plurality of factor-dependent cells with an agent; and (c) following contacting the plurality of cells plurality of factor-dependent cells with the agent, detecting and/or measuring the level of expansion of the cell culture.

10 **[00128]** In some embodiments, the method further comprises identifying an agent as cytokine interfering agent if, following contact with the agent, the cell culture contracts (e.g. as compared to a cell culture not contacted with an agent). In some embodiments, following contact with the agent the cell culture contracts (when compared to a cell culture not contacted with an agent) by about 10% to about 100% (e.g. at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 15 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 10% to about 100%). In some embodiments, following contact with the agent the cell culture contracts (when compared to a cell culture not contacted with an agent) by about 30% to about 100% (e.g. at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 30% to about 100%). In some embodiments, following contact with the agent the cell 20 culture contracts (when compared to a cell culture not contacted with an agent) by about 50% to about 100% (e.g. at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 50% to about 100%). In some embodiments, following contact with the agent the cell culture contracts (when compared to a cell culture not contacted with an agent) by about 70% to about 100% (e.g. at least 70%, 75%, 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 25 95%, 96%, 97%, 98%, or any other percent from about 70% to about 100%). In some embodiments, following contact with the agent the cell culture contracts (when compared to a cell culture not contacted with an agent) by about 80% to about 100% (e.g. at least 80%, 85%, 86%, 87%, 88%, 90%, 91%, 92%, 94%, 95%, 96%, 97%, 98%, or any other percent from about 80% to about 100%).

[00129] In some embodiments, a method of identifying an agent that that replaces and/or augments cytokine 30 function further comprises comparing (1) the level of expansion observed in a plurality of factor-dependent cells following contacting the plurality of cells with an agent to (2) the level of expansion of a control. In some embodiments, the control is a plurality of factor-dependent cells not contacted with an agent.

[00130] In some embodiments, contacting a factor-dependent cell with an agent that inhibits and/or interferes with cytokine function results in the expression of MYC being down-regulated. In certain instances, the down 35 -regulation of MYC results in factor-dependent cells that require a cytokine for viability. In certain instances, the down -regulation of MYC results in factor-dependent cells that do not proliferate in the absence of a

cytokine. In certain instances, the down -regulation of MYC results in factor-dependent cells that undergo apoptosis in the absence of a cytokine.

[00131] In some embodiments, the factor-dependent cells are lymphoid cells (i.e. derived from cells of the lymphatic system). In some embodiments, the factor-dependent cells is, by way of non-limiting example, IL-2^{-/-}, IL-3^{-/-}, IL-4^{-/-}, IL-5^{-/-}, IL-6^{-/-}, IL-7^{-/-}, IL-8^{-/-}, IL-9^{-/-}, IL-10^{-/-}, IL-11^{-/-}, IL-12^{-/-}, or any combinations thereof. In some embodiments, the factor-dependent cells are derived from a cell line selected from, by way of non-limiting example: 2D6; 2E8; 10B10; ATH8; B13; BAF/3; BALM-4; BCL1; CT.4S; CT6; CTL44; CTLL-2; D1; D10; D36; Da; DS-1; Ea3.17; EL4; FL5.12; HT-2; IC-2; Kitt225; KT-3; L4; L138.8A; LBRM-33; LyD9; MC/9; MLA-144; Mono Mac 6; Nb2; NKC3; PB-1; Pno; PT-18; Ramos; Sez627; T10; T88; T1165; TALL-103; TF-1; TMD2; TS1; UT-7; XG-1; Y16; or a combination thereof. In some embodiments, the cells are derived from CTLL-2; or BAF/3 cell lines.

Reporter Assay

[00132] Disclosed herein, in certain embodiments, is a method of identifying an agent that induces (e.g., activates, increases, or supplements) an immune response. In some embodiments, the method comprises identifying an agent that modulates the function of a MYC encoded transcription factor. In some embodiments, the method comprises (a) transforming a plurality of cells in a cell culture with a reporter construct, wherein the reporter construct is a plasmid that comprises a reporter gene, and wherein the expression of the reporter gene is operably linked to one or more E-box sequences; (b) contacting the plurality of cells plurality of cells transformed with the reporter construct with an agent; and (c) following contacting the culture with an agent, detecting and/or measuring the level of expression of the reporter gene. In some embodiments, the reporter construct is obtained in any suitable manner (e.g. purchased from a vendor or fabricated *in-house*). For methods suitable for fabricating a reporter construct and transforming cells with a reporter construct see Molecular Cloning: A Laboratory Manual, second edition (Sambrook *et al.*, 1989) and Molecular Cloning: A Laboratory Manual, third edition (Sambrook and Russel, 2001), jointly referred to herein as “Sambrook”); Current Protocols in Molecular Biology (F. M. Ausubel *et al.*, eds., 1987, including supplements through 2001); Current Protocols in Nucleic Acid Chemistry John Wiley & Sons, Inc., New York, 2000) which are hereby incorporated by reference for such disclosure.

[00133] In some embodiments, a method of identifying an agent that modulates the function of a MYC encoded transcription factor further comprises comparing (1) the level of expression of the reporter gene in a plurality of cells transformed with the reporter construct following contact with an agent to (2) the level of the level of expression of the reporter gene in a control.

[00134] In some embodiments, the cells are eukaryotic cells. In some embodiments, the eukaryotic cells are human cells. In some embodiments, the cells are human lymphoid cells. In some embodiments, the lymphoid cells are B-cells.

[00135] In some embodiments, the reporter construct comprises one or more E-box sequences upstream of a reporter gene. In some embodiments, the one or more E-box sequences are encoded in a mvc-responsive

promoter sequence and/or are upstream of the myc-responsive promoter. As used here, “promoter sequence” means a nucleotide sequence upstream of a gene that allows an RNA polymerase to transcribe the gene. As used herein, “myc-responsive promoter” means a promoter sequence to which a transcription factor encoded by MYC *can* bind. In some embodiments, the myc-responsive promoter is the ornithine decarboxylase promoter.

[00136] In some embodiments, the reporter gene is a β -galactosidase gene, a β -lactamase gene, a horseradish peroxidase gene, an alkaline phosphatase gene, a thymidine kinase gene, a xanthine phosphoribotransferase gene, a tyrosinase gene, a cytosine deaminase gene, an antibiotic resistance gene, or a gene having a fluorescent expression product. In some embodiments, the gene having a fluorescent expression product is a luciferase gene, or a green fluorescent polypeptide (GFP) gene.

[00137] In some embodiments, the method of identifying an agent that modulates the function of a MYC encoded transcription factor further comprises detecting the expression of and/or measuring the amount of expression of the reporter gene. In certain instances, the expression of the reporter gene is detectable and measurable in any suitable manner (e.g. fluorescence microscopy).

[00138] In some embodiments, the cell culture further comprises serum. In some embodiments, the serum is fetal bovine serum (FBS); bovine serum; horse serum; human serum; chicken serum; goat serum; porcine serum; rabbit serum; sheep serum; a serum replacement (e.g. Serum Replacement 1(Sigma-Aldrich)); or a combination thereof. In some embodiments, the cell culture further comprises media. In some embodiments, the media is, by way of non-limiting example, phosphate buffered saline; Earle’s Balanced Salts; Hanks’ Balanced Salts; Tyrode’s Salts; derivatives thereof; or combinations thereof. In some embodiments, the growth factor is, by way of non-limiting example, a cytokine, epidermal growth factor, or platelet-derived growth factor. In some embodiments, the cytokine is, by way of non-limiting example, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, or IL-13. In certain instances, the choice of growth factor is determined based on the type of cell used in the method.

[00139] In some embodiments, the method further comprises contacting the plurality of cells with a mitogenic stimulant. In some embodiments, the mitogenic stimulant is, by way of non-limiting example, an antigen, or a mitogenic antibody. In some embodiments, the antigen is, by way of non-limiting example, phytohaemagglutinin (PHA), concanavalin A (conA), lipopolysaccharide (LPS), pokeweed mitogen (PWM), or combinations thereof. In some embodiments, the mitogenic antibody is, by way of non-limiting example, an anti-IgM, an anti-CD40, or combinations thereof. In some embodiments, the anti-IgM is anti-IgM-F(ab’)₂. In some embodiments, the anti-CD40 is anti-CD40 IgM.

In vitro validation of agents

[00140] In some embodiments, any of the methods described herein further comprise validating *in vitro* an agent (e.g. an agent identified according to any method described herein) as a regulator of an immune response. In some embodiments, a method of validating *in vitro* an agent as a regulator of an immune response comprises (a) contacting the agent with a primary T-cell activation culture; (b) staining the culture

for activation markers; and (c) staining the culture 24 hours after contact with the agent. In certain instances, the cell viability is detectable and measurable in any suitable manner (e.g. fluorescence microscopy, flow cytometry, FACS). In certain instances, an agent that up-regulates an immune response will cause an increase in activation markers as compared to a control (or the culture before contact with the agent). In certain instances, an agent that down-regulates an immune response will cause a decrease in activation markers as compared to a control (or the culture before contact with the agent).

[00141] In some embodiments, any of the methods described herein further comprise validating *in vitro* an agent as a regulator of an immune response. In some embodiments, a method of validating *in vitro* an agent as a regulator of an immune response comprises (a) contacting the agent with a primary T-cell activation culture; (b) staining the culture with CFSE (carboxyfluorescein succinimidyl ester); and (c) staining the culture 72 hours after contact with the agent. In certain instances, CFSE indicates cell proliferation. In certain instances, a cell incorporates the CFSE. In certain instances, a portion (i.e. half) of the CFSE in a cell transfers to a daughter cell following division. In certain instances, the amount of CFSE in cell decreases with each division. In certain instances, the cell viability is detectable and measurable in any suitable manner (e.g. fluorescence microscopy, flow cytometry, FACS). In certain instances, an agent that up-regulates an immune response will cause an increase in cell proliferation as compared to a control (or the culture before contact with the agent). In certain instances, an agent that down-regulates an immune response will cause a decrease in cell proliferation as compared to a control (or the culture before contact with the agent).

[00142] In some embodiments, any of the methods described herein further comprise validating *in vitro* an agent as a regulator of an immune response. In some embodiments, a method of validating *in vitro* an agent as a regulator of an immune response comprises (a) contacting the agent with a primary T-cell activation culture; (b) staining the culture with 7AAD (7-amino-actinomycin D); and (c) staining the culture 72 hours after contact with the agent. In certain instances, 7-AAD indicates cell viability. In certain instances, 7-AAD will stain cells with compromised membranes (i.e. unviable cells). In certain instances, 7-AAD will not stain cells with uncompromised membranes (i.e. viable cells). In certain instances, the cell viability is detectable and measurable in any suitable manner (e.g. fluorescence microscopy, flow cytometry, FACS). In certain instances, an agent that up-regulates an immune response will cause an increase in cell viability as compared to a control (or the culture before contact with the agent). In certain instances, an agent that down-regulates an immune response will cause a decrease in cell viability as compared to a control (or the culture before contact with the agent).

In vivo validation of agents

[00143] In some embodiments, any of the methods described herein further comprise validating *in vivo* an agent (e.g. an agent identified according to any method described herein) as a regulator of an immune response. In some embodiments, a method of validating *in vivo* an agent as a regulator of an immune response comprises (a) immunizing a mammal (e.g. a mouse, rat, rabbit) against an antigen (e.g. OVA peptides, NP-KHL) by administering the antigen to a mouse; (b) administering the agent to the mouse before.

during, or after administration of the antigen; (c) obtaining a biological sample (e.g. blood, lymph, or serum) from the mouse after administration of the antigen and agent; and (d) detecting or measuring an immune response (e.g. the rate of proliferation of memory B-cells and/or T-cells). In some embodiments, an immune response is detected or measured by detecting and measuring antibody production following the immunization. In some embodiments, antibody production is assayed once-a-day every day for a week following immunization and then once-a-week thereafter. In some embodiments, an immune response is detected or measured by detecting and measuring T-cell proliferation upon re-stimulation with the antigen. In some embodiments, T-cell proliferation is assayed one week after immunization. In certain instances, an agent that up-regulates a primary immune response will cause an increase in antibody production and T-cell proliferation as compared to a control (i.e. a mouse not administered the agent). In certain instances, an agent that down-regulates a primary immune response will cause a decrease in antibody production and T-cell proliferation as compared to a control (i.e. a mouse not administered the agent).

[00144] In some embodiments, any of the methods described herein further comprise validating *in vivo* an agent (e.g. an agent identified according to any method described herein) as a regulator of an immune response. In some embodiments, a method of validating *in vivo* an agent as a regulator of an immune response comprises (a) immunizing a mouse against an antigen (e.g. OVA peptides, NP-KHL) by administering the antigen to the mouse; (b) administering the agent to the mouse before, during, or after administration of the antigen; (c) challenging the mouse with the antigen 3 to 6 months after inoculation; (d) obtaining a biological sample (e.g. blood, lymph, or serum) from the mouse following the challenge; and (d) detecting or measuring an immune response. In some embodiments, an immune response is detected or measured by detecting and measuring antibody production following immunization. In some embodiments, antibody production is assayed once-a-day every day for a week following challenge and then once-a-week thereafter. In some embodiments, an immune response is detected or measured by detecting and measuring T-cell proliferation upon re-stimulation with the antigen. In some embodiments, T-cell proliferation is assayed one week after challenge. In certain instances, an agent that up-regulates a secondary immune response will cause an accelerated immune response (i.e. an increase rate of antibody production and T-cell proliferation) following challenge with the antigen as compared to a control (i.e. a mouse not administered the agent). In certain instances, an agent that down-regulates a secondary immune response will cause a decelerated immune response (i.e. a decreased rate of antibody production and T-cell proliferation) following challenge with the antigen as compared to a control (i.e. a mouse not administered the agent).

Agents

[00145] Disclosed herein, in certain embodiments, are agents that modulate an immune response (e.g. agents that reverse energy, agents that modulate cytokine function, and/or agents that modulate the function of a MYC encoded transcription factor). In certain embodiments, the agent is useful in administration to an individual in need thereof. In specific embodiments, the agent is any agent identified by any of the methods

disclosed herein. In some embodiments, the agent is any agent that modulates an immune response. In some embodiments, the agent is a small molecule. In some embodiments, the agent is RNAi. In some embodiments, the agent is an agonist of MYC (e.g. a MYC gene and/or a Myc polypeptide). In some embodiments, the agent is an antagonist of MYC (e.g. a MYC gene and/or a Myc polypeptide).

5 [00146] In some embodiments, the agent is identified as an agent that modulates the viability of a cell based on the agent's ability to reverse anergy in a B-cell. In some embodiments, the agent is identified as an agent that that modulates the viability of a cell based on the agent's ability to modulate the viability of factor-dependent cells. In some embodiments, the agent is identified as an agent that modulates the viability of a cell based on the agent's ability to induce expression of a reporter gene.

10 [00147] In some embodiments, the modulation of the immune response comprises the modulation of the viability of a lymphocyte. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a CD4+ T-cell. In some embodiments, the T-cell is a memory T-cell.

[00148] In some embodiments, the viability of a lymphocyte following contact with the agent is more than 1 to about 10 times greater than the viability of a lymphocyte that has not been contacted the agent. In some
15 embodiments, the agent causes the up-regulation of a MYC gene in a lymphocyte. In some embodiments, the modulation occurs *in vivo*. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a memory T-cell. In certain instances, the up-regulation of a MYC gene results in an extended life-span for the T-cell. In certain instances, the extended life span of a memory T-cell results in a higher concentration of memory T-cells in a body. In certain instances, the higher concentration of memory T-cells results in an
20 accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in anergic T-cells. In certain instances, the decrease in anergic T-cells results in an accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in the time it takes a T-cell to activate in response to an antigen. In certain instances, the decrease in the time it takes a T-cell to activate in response to an antigen results in an accelerated primary
25 immune response to antigen.

[00149] In some embodiments, the agent is administered before, during, or after administration of a vaccine to an individual. In some embodiments, the agent stimulates the immune system and increases the response of the immune system to a vaccine. In some embodiments, agent augments an immune response. In some
30 embodiments, the agent acts synergistically with the vaccine. In some embodiments, the agent is a vaccine adjuvant.

[00150] In some embodiments, a vaccine comprises dead microorganisms, attenuated microorganisms, toxoids, subunits of the pathogen, nucleic acids, or combinations thereof. In some embodiments, the vaccine is a vaccine for hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; influenza; meningococcal disorder; pneumonia; smallpox; cholera;
35 bubonic plague; yellow fever; tuberculosis; human papillomavirus; or combinations thereof. In some embodiments, the vaccine is a vaccine for a cancer (e.g. follicular B-cell Non-Hodgkin's lymphoma, prostate

cancer, multiple myeloma, kidney cancer, cutaneous melanoma, ocular melanoma, and other solid tumors, carcinomas and sarcomas). In some embodiments, a cancer vaccine is a patient-specific vaccine (e.g. the vaccine comprises a patient's own tumor cells). In some embodiments, a cancer vaccine comprises Prostate Specific Antigen (PSA). In some embodiments, a cancer vaccine comprises sialyl Tn (STn). In some
5 embodiments, a cancer vaccine comprises Heat Shock Proteins (HSPs) (e.g., gp96). In some embodiments, a cancer vaccine comprises ganglioside molecules (e.g., GM2, GD2, and GD3). In some embodiments, a cancer vaccine comprises carcinoembryonic antigen (CEA). In some embodiments, a cancer vaccine comprises MART-1 (also known as Melan-A). In some embodiments, a cancer vaccine comprises tyrosinase. In some embodiments, the vaccine is a DNA vaccine.

10 **[00151]** In some embodiments, the comprises an antigenic moiety. In some embodiments, the antigenic moiety is a toxoid, a peptide, a nucleic acid sequence, a polysaccharide, or a combination thereof. In some embodiments, the antigenic moiety is derived from a pathogen selected from: hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; meningococcal; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human
15 papillomavirus; or combinations thereof. In some embodiments, the antigenic moiety is derived a neoplastic cell. In some embodiments, the antigenic moiety is a nucleic acid or a polymer of nucleic acids.

[00152] In certain instances, up-regulating an immune response in an individual receiving vaccination against an antigen results in an increase in the viability and thus concentration of memory T-cells against that antigen. In certain instances, up-regulating an immune response in an individual receiving vaccination against
20 an antigen results in accelerated activation of the T-cell by the antigen. In certain instances, up-regulating an immune response in an individual receiving vaccination against an antigen results in a decrease in T-cells tolerant of the antigen.

[00153] In some embodiments, the viability of a lymphocyte administered the agent is more than 1 to about 25 times less than the viability of a lymphocyte that has not been administered the agent. In some
25 embodiments, the agent causes the down-regulation of a MYC gene in a lymphocyte. In some embodiments, the modulation occurs *in vivo*. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a memory T-cell. In certain instances, the down -regulation of a MYC gene results in a decreased life-span for the T-cell. In certain instances, the decreased life span of a memory T-cell results in a lower concentration of memory T-cells in a body. In certain instances, the lower concentration of memory T-cells
30 results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in anergic T-cells. In certain instances, the increase in anergic T-cells results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in the time it takes a T-cell to activate in response
35 to an antigen. In certain instances, the increase in the time it takes a T-cell to activate in response to an

antigen results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression.

[00154] In some embodiments, the agent is administered to an individual with an autoimmune disorder. In some embodiments, the autoimmune disorder is Castleman's Disease, lupus, multiple sclerosis, scleroderma pigmentosa, Autoimmune Lymphoproliferative Syndrome (ALPS), myasthenia gravis, diabetes, asthma, rheumatoid arthritis, vitiligo, diGeorge's syndrome, Grave's disease, Crohn's disease, inflammatory bowel disease, colitis, orchitis, scleroderma pigmentosa, uveitis, Post-Transplant Lymphoproliferative Disease (PTLD), or Autoimmune Disease-Associated Lymphadenopathy (ADAL). In certain instances, down-regulating an immune response in an individual with an autoimmune disorder ameliorates and/or prevents an immune response against self antigens by the subject's immune system.

[00155] In some embodiments, the agent is administered to an individual that has received an organ transplant, or a bone marrow transplant. In certain instances, down-regulating an immune response in an individual that has received an organ or bone marrow transplant ameliorates and/or prevents an immune response against the transplanted organ or bone marrow by the subject's immune system.

Pharmaceutical Compositions

[00156] Disclosed herein, in certain embodiments, are compositions that modulate the immune system in an individual in need thereof, wherein the composition comprises a therapeutically-effective amount of an agent identified by any of the methods disclosed herein. In some embodiments, the agent is a small molecule. In some embodiments, the agent is RNAi. In some embodiments, the agent is a biological molecule (e.g., a peptide, a fusion peptide). In some embodiments, the agent is a fusion peptide disclosed herein. In some embodiments, the agent is a fusion peptide comprising (a) a transporter peptide sequence; and (b) a MYC sequence. In some embodiments, the agent is a fusion peptide of Formula (I):

transporter peptide sequence -MYC sequence.

[00157] In some embodiments, the agent is identified as an agent that modulates the viability of a cell based on the agent's ability to reverse energy in a B-cell. In some embodiments, the agent is identified as an agent that modulates the viability of a cell based on the agent's ability to modulate the viability of factor-dependent cells. In some embodiments, the agent is identified as an agent that modulates the viability of a cell based on the agent's ability to induce expression of a reporter gene.

[00158] In some embodiments, the modulation of the immune response comprises the modulation of the viability of a lymphocyte. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a CD4+ T-cell. In some embodiments, the T-cell is a memory T-cell.

[00159] In some embodiments, the viability of a lymphocyte following contact with the agent is more than 1 to about 10 times greater than the viability of a lymphocyte that has not been contacted the agent. In some embodiments, the agent causes the up-regulation of a MYC gene in a lymphocyte. In some embodiments, the modulation occurs *in vivo*. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-

cell is a memory T-cell. In certain instances, the up-regulation of a MYC gene results in an extended life-span for the T-cell. In certain instances, the extended life span of a memory T-cell results in a higher concentration of memory T-cells in a body. In certain instances, the higher concentration of memory T-cells results in an accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in anergic T-cells. In certain instances, the decrease in anergic T-cells results in an accelerated primary immune response to antigen. In certain instances, the up-regulation of a MYC gene results in a decrease in the time it takes a T-cell to activate in response to an antigen. In certain instances, the decrease in the time it takes a T-cell to activate in response to an antigen results in an accelerated primary immune response to antigen.

10 **[00160]** In some embodiments, the agent is administered before, during, or after administration of a vaccine to an individual. In some embodiments, the agent stimulates the immune system and increases the response of the immune system to a vaccine. In some embodiments, the agent augments an immune response. In some embodiments, the agent acts synergistically with the vaccine. In some embodiments, the agent is a vaccine adjuvant.

15 **[00161]** In some embodiments, a vaccine comprises dead microorganisms, attenuated microorganisms, toxoids, subunits of the pathogen, nucleic acids, or combinations thereof. In some embodiments, the vaccine is a vaccine for hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; influenza; meningococcal disease; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus; or combinations thereof. In some
20 embodiments, the vaccine is a vaccine for a cancer (e.g. Follicular B-cell Non-Hodgkin's Lymphoma, prostate cancer, multiple myeloma, kidney cancer, cutaneous melanoma, and ocular melanoma). In some embodiments, a cancer vaccine is a patient-specific vaccine (e.g. the vaccine comprises a patient's own tumor cells). In some embodiments, a cancer vaccine comprises Prostate Specific Antigen (PSA). In some
25 embodiments, a cancer vaccine comprises sialyl Tn (STn). In some embodiments, a cancer vaccine comprises Heat Shock Proteins (HSPs) (e.g., gp96). In some embodiments, a cancer vaccine comprises ganglioside molecules (e.g., GM2, GD2, and GD3). In some embodiments, a cancer vaccine comprises carcinoembryonic antigen (CEA). In some embodiments, a cancer vaccine comprises MART-1 (also known as Melan-A). In some embodiments, a cancer vaccine comprises tyrosinase. In some embodiments, the vaccine is a DNA vaccine.

30 **[00162]** In some embodiments, the agent comprises an antigenic moiety. In some embodiments, the antigenic moiety is a toxoid, a peptide, a nucleic acid sequence, a polysaccharide, or a combination thereof. In some embodiments, the antigenic moiety is derived from a pathogen selected from: hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; meningococcal; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human
35 papillomavirus; or combinations thereof. In some embodiments, the antigenic moiety is derived from a neoplastic cell. In some embodiments, the antigenic moiety is a nucleic acid or a polymer of nucleic acids.

[00163] In certain instances, up-regulating an immune response in an individual receiving vaccination against an antigen results in an increase in the viability and thus concentration of memory T-cells against that antigen. In certain instances, up-regulating an immune response in an individual receiving vaccination against an antigen results in accelerated activation of the T-cell by the antigen. In certain instances, up-regulating an immune response in an individual receiving vaccination against an antigen results in a decrease in T-cells tolerant of the antigen.

[00164] In some embodiments, the viability of a lymphocyte administered the agent is more than 1 to about 25 times less than the viability of a lymphocyte that has not been administered the agent. In some embodiments, the agent causes the down-regulation of a MYC gene in a lymphocyte. In some embodiments, the modulation occurs *in vivo*. In some embodiments, the lymphocyte is a T-cell. In some embodiments, the T-cell is a memory T-cell. In certain instances, the down-regulation of a MYC gene results in a decreased life-span for the T-cell. In certain instances, the decreased life span of a memory T-cell results in a lower concentration of memory T-cells in a body. In certain instances, the lower concentration of memory T-cells results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in anergic T-cells. In certain instances, the increase in anergic T-cells results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression. In certain instances, the down-regulation of a MYC gene results in an increase in the time it takes a T-cell to activate in response to an antigen. In certain instances, the increase in the time it takes a T-cell to activate in response to an antigen results in a decelerated primary immune response to antigen, treatment of an autoimmune disorder, and/or immunosuppression.

[00165] In some embodiments, the agent is administered to an individual with an autoimmune disorder. In some embodiments, the autoimmune disorder is Castleman's Disease, lupus, multiple sclerosis, scleroderma pigmentosa, Autoimmune Lymphoproliferative Syndrome (ALPS), myasthenia gravis, diabetes, asthma, rheumatoid arthritis, vitiligo, diGeorge's syndrome, Grave's disease, Crohn's disease, inflammatory bowel disease, colitis, orchitis, scleroderma pigmentosa, uveitis, Post-Transplant Lymphoproliferative Disease (PTLD), or Autoimmune Disease-Associated Lymphadenopathy (ADAL). In certain instances, down-regulating an immune response in an individual with an autoimmune disorder ameliorates and/or prevents an immune response against self antigens by the subject's immune system.

[00166] In some embodiments, the agent is administered to an individual that has received an organ transplant, or a bone marrow transplant. In certain instances, down-regulating an immune response in an individual that has received an organ or bone marrow transplant ameliorates and/or prevents an immune response against the transplanted organ or bone marrow by the subject's immune system.

Formulations of Pharmaceutical Compositions

[00167] In some embodiments, pharmaceutical compositions are formulated in a conventional manner using one or more physiologically acceptable carriers including, e.g., excipients and auxiliaries which facilitate processing of the active compounds into preparations which are suitable for pharmaceutical use. In certain embodiments, proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999).

[00168] A pharmaceutical composition, as used herein, refers to a mixture of a compound described herein, with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. In certain instances, the pharmaceutical composition facilitates administration of the compound to an individual or cell. In certain embodiments of practicing the methods of treatment or use provided herein, therapeutically effective amounts of compounds described herein are administered in a pharmaceutical composition to an individual having a disorder, disease, or condition to be treated. In specific embodiments, the individual is a human. As discussed herein, the therapeutic compounds described herein are either utilized singly or in combination with one or more additional therapeutic agents.

[00169] In some embodiments, the pharmaceutical formulations described herein are administered to an individual in any manner, including one or more of multiple administration routes, such as, by way of non-limiting example, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

[00170] Pharmaceutical compositions including a compound described herein are optionally manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[00171] In some embodiments, a pharmaceutical composition described herein includes one or more agents described herein, as an active ingredient in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In some embodiments, the compounds described herein are utilized as an *N*-oxide or in a crystalline or amorphous form (i.e., a polymorph). In certain embodiments, an active metabolite or prodrug of a compound described herein is utilized. In some situations, a compound described herein exists as tautomers. All tautomers are included within the scope of the compounds presented herein. In certain embodiments, a compound described herein exists in an unsolvated or solvated form, wherein solvated forms comprise any

pharmaceutically acceptable solvent, e.g., water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00172] A “carrier” includes, in some embodiments, a pharmaceutically acceptable excipient and is selected on the basis of compatibility with compounds disclosed herein, such as, compounds of any of Formulas I-V, and the release profile properties of the desired dosage form. Exemplary carrier materials include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. See, e.g., *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999).

[00173] Moreover, in some embodiments, the pharmaceutical compositions described herein are formulated as a dosage form. As such, in some embodiments, provided herein is a dosage form comprising a compound described herein, e.g., a compound of any of Formulas I-V, suitable for administration to an individual. In certain embodiments, suitable dosage forms include, by way of non-limiting example, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions, solid oral dosage forms, aerosols, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[00174] The pharmaceutical solid dosage forms described herein optionally include an additional therapeutic compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof. In some aspects, using standard coating procedures, such as those described in *Remington's Pharmaceutical Sciences*, 20th Edition (2000), a film coating is provided around the formulation of the compound of any of Formula I-V. In one embodiment, a compound described herein is in the form of a particle and some or all of the particles of the compound are coated. In certain embodiments, some or all of the particles of a compound described herein are microencapsulated. In some embodiment, the particles of the compound described herein are not microencapsulated and are uncoated.

[00175] In some embodiments, the pharmaceutical composition described herein is in unit dosage forms suitable for single administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of one or more compound. In some embodiments, the unit dosage is in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. Aqueous suspension compositions are

optionally packaged in single-dose non-reclosable containers. In some embodiments, multiple-dose re-closeable containers are used. In certain instances, multiple dose containers comprise a preservative in the composition. By way of example only, formulations for parenteral injection are presented in unit dosage form, which include, but are not limited to ampoules, or in multi-dose containers, with an added preservative.

5 [00176] In some embodiments, the agents and compositions described herein are administered before, during, or after administration of a vaccine. In some embodiments, the agents and compositions described herein are incorporated into a vaccine. In some embodiments, the agents and compositions stimulate the immune system and increase the response of the immune system to a vaccine. In some embodiments, the agents and compositions augment an immune response. In some embodiments, the agents and compositions act
10 synergistically with the vaccine. In some embodiments, the agents and compositions are vaccine adjuvants. In some embodiments, the vaccine formulation comprises an antigen or antigenic moiety (e.g. a protein or polysaccharide from *B. pertussis*, *C. tetani*, *E. coli*, *C. diphtheriae*, *P. Aeruginosa*, *V. cholerae*, *H. influenzae*, *N. meningitidis*, *S. pneumoniae*, *N. gonorrhoea*) and the agent. In some embodiments, the vaccine further comprises a carrier (e.g. water, saline, PBS), a preservative (e.g. thimerosal, 2-phenoxy ethanol, phenol,
15 benzethonium chloride), a stabilizer (e.g. lactose, monosodium glutamate), an antibiotic, an antioxidant, a pH controlling agent, or combinations thereof.

Topical Formulations

[00177] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is administered
20 topically (i.e., the polypeptide is administered to the surface of the skin). In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is administered topically at the site of a vaccine injection. In certain instances, a localized antigen depot (i.e., a tissue depot formed by vaccine adjuvants (e.g., water-in-oil emulsions, or aluminum salts) that results in the slow and even release of an antigen) arises during vaccination with current approaches. In some embodiments, the transient upregulation of MYC in the
25 resident lymphoid cells in the skin, near the site of the antigen depot, drives a more robust immune response, and yields a broader response in the context of relaxed tolerance to self antigens.

[00178] Any suitable formulation is utilized. In some embodiments, the MYC fusion polypeptide is formulated as a solution, a cream, a lotion, a gel, an ointment, a foam, a microemulsion, or a combination thereof. In some embodiments, the MYC fusion polypeptide is formulated for delivery via a transdermal
30 patch.

[00179] Any suitable method of topical administration is utilized (e.g., electroporation, sonophoresis, chemical delivery, skin abrasion). In some embodiments, the MYC fusion polypeptide is formulated for chemical delivery (i.e., a chemical substance (e.g., PEG, ethanol, glycerol monolaurate, sodium dodecyl sulfate, phosphatidyl choline, or urea) is used to facilitate penetration of an active agent across the skin). In
35 some embodiments, the MYC fusion polypeptide is formulated for delivery via electroporation (i.e., the permeabilization of a barrier (e.g., the skin) via application of an electric current). In some embodiments, the

MYC fusion polypeptide is formulated for delivery via sonophoresis (i.e., the permeablization of a barrier (e.g., the skin) via application of ultrasound).

Creams and Lotions

5 **[00180]** In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as a cream. In certain instances, creams are semisolid (e.g., soft solid or thick liquid) formulations that include a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) dispersed in an oil-in-water emulsion or a water-in-oil emulsion.

[00181] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as a lotion. In certain instances, lotions are fluid emulsions (e.g., oil-in-water emulsions or a water-in-oil
10 emulsions).

[00182] In some embodiments, the hydrophobic component of a lotion and/or cream is derived from an animal (e.g., lanolin, cod liver oil, and ambergris), plant (e.g., safflower oil, castor oil, coconut oil, cottonseed oil, menhaden oil, palm kernel oil, palm oil, peanut oil, soybean oil, rapeseed oil, linseed oil, rice bran oil, pine oil, sesame oil, or sunflower seed oil), petroleum (e.g., mineral oil, or petroleum jelly), or a combination
15 thereof.

Ointments

[00183] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as an ointment. In certain instances, ointments are semisolid preparations that soften or melt at body temperature.

20 *Pastes*

[00184] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as a paste. In certain instances, pastes contain at least 20% solids. In certain instances, pastes are ointments that do not flow at body temperature.

Gels

25 **[00185]** In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as a gel. In certain instances, gels are semisolid (or semi-rigid) systems consisting of dispersions of large organic molecules dispersed in a liquid. In certain instances, gels are water-soluble and are removed using warm water or saline.

Sticks

30 **[00186]** In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated as a stick. In certain instances, sticks are solid dosage forms that melt at body temperature. In some embodiments, a stick comprises a wax, a polymer, a resin, dry solids fused into a firm mass, and/or fused crystals. In some embodiments, a topical formulation of a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is in the form of a styptic pencil (i.e., a stick prepared by (1) heating crystals until they lose their
35 water of crystallization and become molten, and (2) pouring the molten crystals into molds and allowing them to harden). In some embodiments, a topical formulation of a MYC fusion polypeptide disclosed herein (e.g.,

TAT-MYC) is in the form of stick wherein the stick comprises a wax (e.g., the wax is melted and poured into appropriate molds in which they solidify in stick form).

[00187] In some embodiments, a topical formulation of a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is in the form of stick wherein the stick comprises a melting base (i.e., a base that softens at body temperature). Examples of melting bases include, but are not limited to, waxes, oils, polymers and gels. In some embodiments, a topical formulation of a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is in the form of stick wherein the stick comprises a moisture base (i.e., a base that is activated by the addition of moisture).

Patches

[00188] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated for administration via a patch. In some embodiments, a topical formulation disclosed herein is dissolved and/or dispersed in a polymer or an adhesive. In some embodiments, a patch disclosed herein is constructed for continuous, pulsatile, or on demand delivery of a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC).

Wound Dressings

[00189] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated for administration via a wound dressing. Wound dressings include, but are not limited to gauzes, transparent film dressings, hydrogels, polyurethane foam dressings, hydrocolloids and alginates.

Dermatological Excipients

[00190] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated with a penetration enhancer. Penetration enhancers include, but are not limited to, sodium lauryl sulfate, sodium laurate, polyoxyethylene-20-cetyl ether, laureth-9, sodium dodecylsulfate, dioctyl sodium sulfosuccinate, polyoxyethylene-9-lauryl ether (PLE), Tween 80, nonylphenoxypolyethylene (NP-POE), polysorbates, sodium glycocholate, sodium deoxycholate, sodium taurocholate, sodium taurodihydrofusidate, sodium glycodihydrofusidate, oleic acid, caprylic acid, mono- and di-glycerides, lauric acids, acylcholines, caprylic acids, acylcarnitines, sodium caprates, EDTA, citric acid, salicylates, DMSO, decylmethyl sulfoxide, ethanol, isopropanol, propylene glycol, polyethylene glycol, glycerol, propanediol, and diethylene glycol monoethyl ether.

[00191] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated with a gelling (or thickening) agent. In some embodiments, a topical formulation disclosed herein further comprises from about 0.1% to about 5%, more preferably from about 0.1% to about 3%, and most preferably from about 0.25% to about 2%, of a gelling agent. In certain embodiments, the viscosity of a topical formulation disclosed herein is in the range from about 100 to about 500,000 cP, about 100 cP to about 1,000 cP, about 500 cP to about 1500 cP, about 1000 cP to about 3000 cP, about 2000 cP to about 8,000 cP, about 4,000 cP to about 10,000 cP, about 10,000 cP to about 50,000 cP.

[00192] Suitable gelling agents for use in preparation of the gel topical formulation include, but are not limited to, celluloses, cellulose derivatives, cellulose ethers (e.g., carboxymethylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxymethylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, methylcellulose), guar gum, xanthan gum, locust bean gum, alginates (e.g., alginic acid), silicates, starch, tragacanth, carboxyvinyl polymers, carrageenan, paraffin, petrolatum, acacia (gum arabic), agar, aluminum magnesium silicate, sodium alginate, sodium stearate, bladderwrack, bentonite, carbomer, carrageenan, carbopol, xanthan, cellulose, microcrystalline cellulose (MCC), ceratonia, chondrus, dextrose, furcellaran, gelatin, ghatti gum, guar gum, hectorite, lactose, sucrose, maltodextrin, mannitol, sorbitol, honey, maize starch, wheat starch, rice starch, potato starch, gelatin, sterculia gum, polyethylene glycol (e.g. PEG 200-4500), gum tragacanth, ethyl cellulose, ethylhydroxyethyl cellulose, ethylmethyl cellulose, methyl cellulose, hydroxyethyl cellulose, hydroxyethylmethyl cellulose, hydroxypropyl cellulose, poly(hydroxyethyl methacrylate), oxypolygelatin, pectin, polygeline, povidone, propylene carbonate, methyl vinyl ether/maleic anhydride copolymer (PVM/MA), poly(methoxyethyl methacrylate), poly(methoxyethoxyethyl methacrylate), hydroxypropyl cellulose, hydroxypropylmethyl-cellulose (HPMC), sodium carboxymethyl-cellulose (CMC), silicon dioxide, polyvinylpyrrolidone (PVP: povidone), or combinations thereof.

[00193] In some embodiments, a MYC fusion polypeptide disclosed herein (e.g., TAT-MYC) is formulated with an emollient. Emollients include, but are not limited to, castor oil esters, cocoa butter esters, safflower oil esters, cottonseed oil esters, corn oil esters, olive oil esters, cod liver oil esters, almond oil esters, avocado oil esters, palm oil esters, sesame oil esters, squalene esters, kikai oil esters, soybean oil esters, acetylated monoglycerides, ethoxylated glyceryl monostearate, hexyl laurate, isoheptyl laurate, isoheptyl palmitate, isopropyl palmitate, methyl palmitate, decyloleate, isodecyl oleate, hexadecyl stearate decyl stearate, isopropyl isostearate, methyl isostearate, diisopropyl adipate, diisoheptyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate, oleyl myristate, oleyl stearate, and oleyl oleate, pelargonic acid, lauric acid, myristic acid, palmitic acid, stearic acid, isostearic acid, hydroxystearic acid, oleic acid, linoleic acid, ricinoleic acid, arachidic acid, behenic acid, erucic acid, lauryl alcohol, myristyl alcohol, cetyl alcohol, hexadecyl alcohol, stearyl alcohol, isostearyl alcohol, hydroxystearyl alcohol, oleyl alcohol, ricinoleyl alcohol, behenyl alcohol, erucyl alcohol, 2-octyl dodecanyl alcohol, lanolin and lanolin derivatives, beeswax, spermaceti, myristyl myristate, stearyl stearate, carnauba wax, candelilla wax, lecithin, and cholesterol.

Combinations

[00194] In some embodiments, it is appropriate to administer at least one therapeutic agent described herein in combination with another therapeutic agent. Or, by way of example only, the benefit experienced by a patient is increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. In any case, regardless of the disorder, disease or condition being treated, the overall benefit experienced by the patient is. in some

embodiments, additive of the two therapeutic agents or in other embodiments, the patient experiences a synergistic benefit.

[00195] In some embodiments, the particular choice of compounds depends upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

5 The compounds are optionally administered concurrently (e.g., simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the disease, disorder, or condition, the condition of the patient, and the actual choice of compounds used. In certain instances, the determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is based on an evaluation of the disorder being treated and the
10 condition of the patient.

[00196] In some embodiments, therapeutically-effective dosages vary when the drugs are used in treatment combinations. Methods for experimentally determining therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens are described in the literature. For example, the use of metronomic dosing, i.e., providing more frequent, lower doses in order to minimize toxic side effects, has
15 been described extensively in the literature. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient.

[00197] In some embodiments of the combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disorder or condition being treated and so forth. In addition, when co-administered with one or more biologically
20 agents, the compound provided herein is optionally administered either simultaneously with the biologically agent(s), or sequentially. In certain instances, if administered sequentially, the attending physician will decide on the appropriate sequence of therapeutic compound described herein in combination with the additional therapeutic agent.

[00198] The multiple therapeutic agents (at least one of which is a therapeutic compound described herein)
25 are optionally administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents are optionally provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). In certain instances, one of the therapeutic agents is optionally given in multiple doses. In other instances, both are optionally given as multiple doses. If not simultaneous, the timing between the multiple doses is any suitable timing, e.g., from more than zero weeks to less than four
30 weeks. In some embodiments, the additional therapeutic agent is utilized to achieve remission (partial or complete) of a cancer, whereupon the therapeutic agent described herein (e.g., a compound of any one of Formulas I-V) is subsequently administered. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations are also envisioned (including two or more therapeutic compounds described herein).

35 **[00199]** In some embodiments, a dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is modified in accordance with a variety of factors. These factors include the disorder from

which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, in various embodiments, the dosage regimen actually employed varies and deviates from the dosage regimens set forth herein.

[00200] In some embodiments, the pharmaceutical agents which make up the combination therapy disclosed herein are provided in a combined dosage form or in separate dosage forms for substantially simultaneous administration. In certain embodiments, the pharmaceutical agents that make up the combination therapy are administered sequentially, with either therapeutic compound being administered by a regimen calling for two-step administration. In some embodiments, two-step administration regimen calls for sequential administration of the agents or spaced-apart administration of the separate agents. In certain embodiments, the time period between the multiple administration steps varies, by way of non-limiting example, from a few minutes to several hours, depending upon the properties of each pharmaceutical agent, such as potency, solubility, bioavailability, plasma half-life and kinetic profile of the pharmaceutical agent.

[00201] In addition, the compounds described herein also are optionally used in combination with procedures that provide additional or synergistic benefit to the patient. By way of example only, patients are expected to find therapeutic and/or prophylactic benefit in the methods described herein, wherein pharmaceutical composition of a compound disclosed herein and /or combinations with other therapeutics are combined with genetic testing to determine whether that individual is a carrier of a gene or gene mutation that is known to be correlated with certain disorders or conditions. In certain embodiments, prophylactic benefit is achieved by administering a therapeutic compound described herein to an individual whose proliferative disorder (e.g., cancer) is in remission (e.g., *partial* or complete).

[00202] In some embodiments, the compounds described herein and combination therapies are administered before, during or after the occurrence of a disorder or condition. Timing of administering the composition containing a compound is optionally varied to suit the needs of the individual treated. Thus, in certain embodiments, the compounds are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or disorders in order to prevent the occurrence of the disorder or condition. In some embodiments, the compounds and compositions are administered to an individual during or as soon as possible after the onset of the symptoms. The administration of the compounds is optionally initiated within the first 48 hours of the onset of the symptoms, within the first 6 hours of the onset of the symptoms, or within 3 hours of the onset of the symptoms. The initial administration is achieved by any route practical, such as, for example, an intravenous injection, a bolus injection, infusion over 5 minutes to about 5 hours, a pill, a capsule, transdermal patch, buccal delivery, and the like, or combination thereof. In some embodiments, the compound should be administered as soon as is practicable after the onset of a disorder or condition is detected or suspected, and for a length of time necessary for the treatment of the disorder, such as, for example, from more than 1 month to about 3 months. The length of treatment is optionally varied for each subject based on known criteria. In exemplary embodiments, the compound or a formulation containing

the compound is administered for at least 2 weeks, between more than 1 month to about 5 years, or from more than 1 month to about 3 years.

[00203] In some embodiments, therapeutic agents are combined with or utilized in combination with one or more of the following therapeutic agents in any combination: Akt selective inhibitors; PI3K inhibitors; polypeptide kinase C (PKC) inhibitors; farnesyltransferase inhibitors; inhibitors of sarco/endoplasmic reticulum Ca²⁺ ATPase; Ca⁺⁺/calmodulin (CaM)-dependent polypeptide kinase inhibitors; cyclin-dependent kinase inhibitors; PPD (p-phenylenediamine); D609 (tricyclodecan-9-yl xanthogenate); PP1 (4-Amino-5-(4-methylphenyl)-7-(t-butyl)pyrazolo[3,4-d]-pyrimidine); AG-879 (alpha-Cyano-(3,5-di-t-butyl-4-hydroxy)thiocinnamide); BAY 11-7082 ((E)-3-(4-Methylphenylsulfonyl)-2-propenenitrile); thapsigargin, or RK-682 (3-Hexadecanoyl-5-hydroxymethyl-tetronic acid).

[00204] Immunosuppressants include, by way of non-limiting example, Sirolimus (rapamycin), cyclosporin, FK506, cypermethrin, deltamethrin, fenvalerate, tyrphostin 8, methotrexate, piceatannol, genistein, tacrolimus, rapamicin, cyclophosphamide, azathioprine, mercaptopurine, mycophenolate, and FTY720.

[00205] Akt selective inhibitors include, by way of non-limiting example, SH4 (1L-6-hydroxymethyl-chiro-inositol 2-R-2-O-methyl-3-O-octadecylcarbonate); SH-5 (D-3-deoxy-2-O-methyl-myoinositol 1-(R)-2-methoxy-3-(octadecyloxy) propyl hydrogen phosphate); and SH-6 (D-2,3-dieoxy-myoinositol 1-(R)-2-methoxy-3-(octadecyloxy) propyl hydrogen phosphate).

[00206] PI3K inhibitors include, by way of non-limiting example, wortmannin; LY294002 (2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one); D000 (Upstate, Milton Keynes, United Kingdom); and D121 (3-phenyl-2-(9H-purin-6-ylsulfanylmethyl)-3H-quinazolin-4-one).

[00207] By way of non-limiting example, protein kinase C (PKC) inhibitors include Ro-318220 (3-[1-[3-(amidinothio)propyl]-1H-indoyl-3-yl]-3-(1-methyl-1H-indoyl-3-yl) maleimide methane sulfonate); Gö6976 (12-(2-cyanoethyl)-6,7,12,13-tetrahydro-13-methyl-5-oxo-5H-indolo(2,3-a)pyrrolo(3,4-c)-carbazole); staurosporin; rottlerin; calphastin C; GF 109203X (3-[1-(Dimethylaminopropyl)indol-3-yl]-4-(indol-3-yl)maleimide hydrochloride); hypericin; sphingosine; miltefosine; palmitoyl-DL-carnitine Cl; rottlerin; and 2,2',3,3',4,4'-hexahydroxy-1,1'-biphenyl-6,6'-dimenthol dimethylether.

[00208] By way of non-limiting example, farnesyltransferase inhibitors include L-744,832 (C₂₆H₄₅N₃O₆S₂ 2HCl); or Sarasar (SCH66336 or lonafarnib).

[00209] By way of non-limiting example, Ca⁺⁺/calmodulin (CaM)-dependent polypeptide kinase inhibitors include: 2-hydroxy-5-(2,5-dihydroxybenzylamino) benzoic acid; KN-62 (1-[N,O-bis(5-isoquinolinesulfonyl)-N-methyl-L-tyrosyl]-4-phenylpiperazine); KN-93 (2-[N-(2-hydroxyethyl)]-N-(4-methoxybenzenesulfonyl)amino-N-(4-chlorocinnamyl)-N-methylbenzylamine); or staurosporine (AM-2282).

[00210] By way of non-limiting example, cyclin-dependent kinase inhibitors include: roscovitine (Seliciclib or CYC202); purvalanol A; or indirubin.

Methods of Dosing and Treatment Regimens

[00211] In some embodiments, an agent and/or composition described herein is used in the preparation of vaccines for immunization against pathogens (e.g. hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; influenza; meningococcal disease; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus) wherein an up-regulation of an immune response would be beneficial to said treatment. In some embodiments, an agent described herein is incorporated into the formulation of a vaccine. In some embodiments, an agent described herein is a vaccine adjuvant. In some embodiments, the volume of a vaccine formulation for injection is from about 50 μ l to about 5ml. In some embodiments, the volume of a vaccine formulation for injection is from about 100 μ l to about 3.5ml. In some embodiments, the volume of a vaccine formulation for injection is from about 200 μ l to about 2ml. In some embodiments, the volume of a vaccine formulation for injection is from about 350 μ l to about 1ml. In some embodiments, the volume of a vaccine formulation for injection is about 500 μ l. The volume of the vaccine is determined by the method of administration. In certain instances, the vaccine is administered as a subcutaneous or intradermal injection. In certain instances, the vaccine is administered as intramuscular injection. In some embodiments, the vaccine is administered as 1 dose. In some embodiments, the vaccine is administered in multiple doses (e.g. in booster shots).

[00212] In some embodiments, an agent and/or composition described herein is used in the preparation of medicaments for the prophylactic and/or therapeutic treatment of disorders that would benefit, at least in part, from a down-regulation of an immune response (e.g. autoimmune disorders, transplant rejection).

[00213] In some embodiments, an agent and/or composition described herein is used in a method of treating a disorder (e.g. hepatitis A; hepatitis B; polio; measles; mumps; rubella; diphtheria; pertussis; tetanus; influenza; varicella zoster virus; rotavirus; influenza; meningococcal disease; pneumonia; smallpox; cholera; bubonic plague; yellow fever; tuberculosis; human papillomavirus, and a cancer) wherein an up-regulation of an immune response would be beneficial to said treatment. In some embodiments, an agent and/or composition described herein is used in a method of prophylactic and/or therapeutic treatment of a disorder that would benefit, at least in part, from a down-regulation of an immune response (e.g. autoimmune disorders, transplant rejection).

[00214] In certain instances wherein the patient's condition does not improve, upon the doctor's discretion the administration of an agent or composition described herein is optionally administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disorder.

[00215] In the case wherein the patient's status does improve, upon the doctor's discretion the administration of an agent or composition described herein is optionally given continuously; alternatively, the dose of drug being administered is temporarily reduced or temporarily suspended for a certain length of time (i.e., a "drug holiday"). The length of the drug holiday optionally varies between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35

days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, or 365 days. The dose reduction during a drug holiday includes from 10%-100%, including, by way of example only, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

5 **[00216]** Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In some embodiments, patients require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00217] In some embodiments, the agents or compositions described herein are in unit dosage forms suitable
10 for single administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of an agent or composition described herein. In some embodiments, the unit dosage is in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. In some embodiments, aqueous suspension compositions are packaged in single-dose non-reclosable containers. Alternatively, multiple-dose
15 reclosable containers are used, in which case it is typical to include a preservative in the composition. By way of example only, formulations for parenteral injection are presented in unit dosage form, which include, but are not limited to ampoules, or in multi dose containers, with an added preservative.

[00218] The daily dosages appropriate for an agent or composition described herein are from about 0.01 to 10.0 mg/kg per body weight. In some embodiments, the daily dosages appropriate for an agent or
20 composition described herein are from about 0.05 to 7.5 mg/kg per body weight. In some embodiments, the daily dosages appropriate for an agent or composition described herein are from about 0.1 to 5.0 mg/kg per body weight. In some embodiments, the daily dosages appropriate for an agent or composition described herein are from about 0.25 to 2.5 mg/kg per body weight. In some embodiments, the daily dosages appropriate for an agent or composition described herein are from about 0.5 to 1.0 mg/kg per body weight.
25 An indicated daily dosage in the larger mammal, including, but not limited to, humans, is in the range from about 0.5 mg to about 100 mg, conveniently administered in divided doses, including, but not limited to, up to four times a day or in extended release form. Suitable unit dosage forms for oral administration include from more than 1 to 50 mg active ingredient. The foregoing ranges are merely suggestive, as the number of variables in regard to an individual treatment regime is large, and considerable excursions from these
30 recommended values are not uncommon. Such dosages are optionally altered depending on a number of variables, not limited to the activity of the agent or composition described herein used, the disorder or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disorder or condition being treated, and the judgment of the practitioner.

[00219] Toxicity and therapeutic efficacy of such therapeutic regimens are optionally determined in cell
35 cultures or experimental animals, including, but not limited to, the determination of the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The

dose ratio between the toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD50 and ED50. An agent or compositions described herein exhibiting high therapeutic indices is preferred. The data obtained from cell culture assays and animal studies are optionally used in formulating a range of dosage for use in human. The dosage of such an agent or composition described herein lies preferably within a range of circulating concentrations that include the ED50 with minimal toxicity. The dosage optionally varies within this range depending upon the dosage form employed and the route of administration utilized.

EXAMPLES

[0220] The following examples are for illustrative purposes only and are non-limiting embodiments. Many modifications, equivalents, and variations of the present invention are possible in light of the above teachings, therefore, it is to be understood that within the scope of the appended claims, the invention may be practiced other than as specifically described.

Example 1 – Assay Using a Reporter Construct

[0221] A plurality of cells are transformed with a reporter construct. The reporter construct has FLuc under control of an ODC promoter. The cells are divided into 96 aliquots using a standard 96-well plate. Each well is administered a different small molecule. The small molecule is incubated with the cells for 6 hours. The level of expression of the luciferase is determined using fluorescent microscopy. Agents which induce the expression of the luciferase gene are validated *in vitro*.

Example 2 – Construction of a TAT-MYC Fusion Peptide

Construction of p-TAT-MYC

[0222] Plasmid pTAT-Myc-V5-6xHis ("6xHis" disclosed as SEQ ID NO: 3) was made by PCR amplification of the coding regions for human MYC using a forward primer that contains an in frame N-terminal 9-amino-acid sequence of the TAT protein transduction domain of HIV-1 (RKKRRQRRR) (SEQ ID NO: 6), and a reverse primer that removed the stop codon. The PCR product was then cloned into pET101/D-Topo (Invitrogen) vector, which includes a C-terminal V5 epitope and 6x-histidine (SEQ ID NO: 3) purification tag.

Bacterial strain used for protein expression

[0223] BL-21 RARE cells were created by transforming BL-21 StarTM E. coli strain (Invitrogen) with pRARE (CamR), isolated from BL21 Rosetta cells (Novagen), that express tRNAs for AGG, AGA, AUA, CUA, CCC, GGA codons.

Protein induction and purification

[0224] pTAT-Myc-V5-6xHis ("6xHis" disclosed as SEQ ID NO: 3) was transformed into BL21 RARE cells and grown on TB/Amp/Cam plate at 37°C overnight. An isolated colony was used to inoculate a 5ml TB/Amp/Cam starter culture and grown at

37°C overnight. 1 liter of TB/Amp/Cam broth was inoculated with the 5ml starter culture and grown to an OD600 of 0.5, and induced with 0.5mM IPTG at 37°C for 3 hrs.

[00225] Bacterial cells were pelleted by centrifugation and the cell pellet was resuspended in lysis buffer (8 M urea, 100mM NaH₂PO₄, 10mM Tris pH to 8.0) and lysed at room temperature overnight on a shaker. The lysate was cleared by centrifugation at 29,000 x g for 30 min and the supernatant was applied to a nickel nitrilotriacetate affinity column (Qiagen) using gravity flow. The column was washed with 25 volumes of lysis buffer containing 10mM imidazole followed by elution with lysis buffer containing 100mM imidazole.

[00226] Protein was concentrated using an Amicon Ultra centrifugal filter device (10,000 MWCO) and dialyzed in a stepwise fashion into dialysis buffer (50mM NaH₂PO₄, 5mM Tris pH 7.0, 450mM NaCl, 5% glycerol, 1mM DTT). The dialysis went as follows: 2 hours in dialysis buffer containing 4M urea, 2 hours in buffer with 2M urea, then 2 hrs in dialysis buffer alone. Purity and size of proteins were verified using SDS-PAGE electrophoresis and either coomassie blue staining or western blot with anti-V5 (1:5000; Invitrogen) or anti-c-Myc (N-262, 1:2000; Santa Cruz Biotechnology) antibodies.

[00227] Protein concentration was measured by Bradford protein assay (Sigma) compared to a standard curve of bovine serum albumin.

Example 3 – Effect of TAT-Myc Fusion Peptide on Immune Responses

[00228] The TAT-MYC fusion peptide is evaluated *in vitro* as a regulator of an immune response. A primary T-cell activation culture is incubated with the TAT-MYC fusion peptide for 72 hours. The culture is then stained with CFSE. Following staining with CFSE, the culture is analyzed by FACS.

Example 4 - Effect of TAT-Myc Fusion Peptide on Immune Responses

[00229] The TAT-MYC fusion peptide is evaluated *in vivo* as a regulator of an immune response. A mouse is immunized against NP-KHL by administering NP-KHL to the mouse. Immediately after immunization against the antigen, the TAT-MYC fusion peptide is administered to the mouse. Twenty-four hours after the immunization serum is taken from the mouse. The serum is analyzed for antibodies. Alternatively, the TAT-MYC fusion peptide is administered concurrently with immunization, or preceding immunization.

Example 5 – Effect of IM TAT-MYC as a Vaccine Adjuvant

[00230] This will be a double-blind, randomized, placebo-controlled study. It is designed to assess immunological memory following administration of an influenza vaccine and a TAT-MYC fusion peptide.

[00231] Participants are divided into two groups. Group I will be administered an influenza vaccine and saline. Group II will be administered an influenza vaccine and the TAT-MYC fusion peptide.

Primary Objective

[00232] To assess the immunological effects of a TAT-MYC fusion peptide administered by intramuscular (IM) injection to individuals receiving an influenza vaccine.

Secondary Objectives

[00233] •To assess the safety and tolerability of a TAT-MYC fusion peptide administered by intramuscular injection to individuals receiving an influenza vaccine.

Methodology

5 [00234] On Day = 0, both Group I and Group II are administered a single dose of the same influenza vaccine. Subjects are monitored for 2 hours for negative side-effects.

[00235] All individuals in Group I not exhibiting negative side-effects are administered 20 uL of saline by IM injection.

10 [00236] All individuals in Group II not exhibiting negative side-effects are administered 20 uL of the TAT-MYC fusion peptide by IM injection.

[00237] Blood is drawn from all individuals on Day = 1. T-cell counts are determined and recorded.

[00238] On Day = 20, all individuals are challenged with antigens from relevant influenza strains. Blood is drawn for all individuals 1 hour after challenge, 6 hours after challenge, 12 hours after challenge, and 24 hours after challenge. T-cell levels are determined.

15

Example 6 – Effect of Topical TAT-MYC as a Vaccine Adjuvant

[00239] This will be a double-blind, randomized, placebo-controlled study. It is designed to assess immunological memory following administration of an influenza vaccine and a TAT-MYC fusion peptide.

20 [00240] Participants are divided into two groups. Group I will be administered an influenza vaccine and saline. Group II will be administered an influenza vaccine and the TAT-MYC fusion peptide.

Primary Objective

[00241] To assess the immunological effects of a TAT-MYC fusion peptide administered topically to individuals receiving an influenza vaccine.

Secondary Objectives

25 [00242] •To assess the safety and tolerability of a TAT-MYC fusion peptide administered topically to individuals receiving an influenza vaccine.

Methodology

[00243] On Day = 0, both Group I and Group II are administered a single dose of the same influenza vaccine. Subjects are monitored for 2 hours for negative side-effects.

30 [00244] All individuals in Group I not exhibiting negative side-effects are given one application of the placebo lotion. Individuals are instructed to apply the lotion onto the skin at the site of the vaccination.

[00245] All individuals in Group II not exhibiting negative side-effects are given one application of the TAT-MYC lotion. Individuals are instructed to apply the lotion onto the skin at the site of the vaccination.

[00246] Blood is drawn from all individuals on Day = 1. T-cell counts are determined and recorded.

[00247] On Day = 20, all individuals are challenged with antigens from relevant influenza strains. Blood is drawn for all individuals 1 hour after challenge, 6 hours after challenge, 12 hours after challenge, and 24 hours after challenge. T-cell levels are determined.

5 **Example 7 – Effect on RA of an Agent that Down-Regulates MYC Expression**

Primary Objective

[00248] To assess the steady-state trough serum concentrations of an agent that decreases the expression of MYC as identified by a method disclosed herein (hereinafter, “Agent X”) following weekly subcutaneous dosing in subjects with active rheumatoid arthritis (RA).

10 *Secondary Objectives*

[00249] •To assess the safety and tolerability of Agent X administered subcutaneously to individuals with RA; to assess the immunogenicity of Agent X administered subcutaneously in individuals with RA; and to examine the effect of subcutaneous administration of Agent X on serum levels of rheumatoid factor (RF) in individuals with RA.

15 *Methodology*

[00250] This will be a double-blind, randomized, placebo-controlled, parallel-group, multiple-dose study. It will be designed to assess the steady-state trough serum concentrations of Agent X following subcutaneous administration in subjects with RA.

[00251] Subjects will be randomized according to a computer generated randomization scheme, in a 3:1 ratio, to receive either Agent X or placebo in 1 of 5 parallel groups based on body weight obtained at a screening visit.

[00252] On Day 1, subjects will receive a single IV infusion (loading dose) of Agent X or placebo, based on their weight range. Agent X or placebo will be administered intravenously over ~ 30 minutes using a calibrated, constant-rate infusion pump prior to starting subcutaneous treatment. Approximately 1 hour after the completion of the IV infusion, subjects will receive their assigned subcutaneous dose of Agent X or placebo drawn from a vial or using a pre-filled syringe. Agent X or placebo will be administered by clinical study site staff weekly by the subcutaneous route, at the same dose as the subcutaneous dose on Day 1 for a total of 12 subcutaneous injections.

[00253] Subjects will be monitored for adverse events (AEs) throughout the duration of the study. Physical examinations, vital sign measurements, and clinical laboratory evaluations will be performed at selected times throughout the study. Blood samples for PK analysis will be collected on Day 1 prior to and at the end of the IV infusion. In addition, blood samples will be collected prior to each weekly SC dose of agent X on Days 8, 15, 22, 29, 36, 43, 50, 57, 64, 71, and 78 for the assessment of steady-state C_{min} concentrations. A single blood sample will also be collected on Days 72, 74, 75, and 76 for evaluation of steady-state C_{max}, T_{max}, and AUC(TAU), where TAU = 7 days, and at study discharge (Day 85). Blood samples for assessment of immunogenicity will be obtained prior to the administration of agent X on Days 1, 15, 29, 43, 57, 71, and

85. Blood samples for the determination of RF will be obtained prior to dosing on Days 1, 8, 15, 29, 57, and 85.

[00254] Subjects who complete the 12 weeks of subcutaneous dosing of Agent X or placebo are eligible to enter into a long term extension (LTE). For subjects who entered the LTE, the first dose of Agent X will be administered on Day 85.

Number of Subjects

[00255] Between 48 and 72 subjects are to be randomized to study treatment.

Diagnosis and Criteria for Inclusion

[00256] Men or women with active RA who are above 18 will eligible for participation in this study. Subjects must meet the classification criteria of the American College of Rheumatology (ACR) (formerly American Rheumatism Association (ARA)), for RA and had active disorder. Subjects must have had RA for at least 1 year from the time of initial diagnosis.

[00257] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein *may* be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

The claims defining the invention are as follows:

1. A pharmaceutical composition for modulating an immune system, comprising:
- (a) an antigenic moiety derived from a pathogen;
- (b) a fusion peptide comprising:
- 5 (i) a transporter peptide sequence;
- (ii) a MYC sequence; and optionally
- (iii) one or more molecules that link the transporter peptide sequence and the MYC sequence; and
- (c) a pharmacologically-acceptable excipient.
- 10 2. The pharmaceutical composition according to claim 1, wherein the fusion peptide has Formula (I):
- transporter peptide sequence -MYC sequence.
3. The pharmaceutical composition according to claim 1, wherein the fusion peptide has Formula (II):
- 15 transporter peptide sequence-X-MYC sequence,
- wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence.
4. The pharmaceutical composition according to claim 1, wherein the fusion peptide has Formula (II):
- 20 transporter peptide sequence-X-MYC sequence,
- wherein in X is at least one amino acid.
5. The pharmaceutical composition according to any one of claims 1-4, wherein the fusion peptide has the following sequence:
- 25 MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYF
- YCDEEENFYQQQQQSELQPPAPSEDIWKKFELLPTPLSPSRRSGLCSPSY
- VAVTPFSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDET
- FIKNIIIQDCMWSGFSAAKLVSEKLASYQAARKDSGSPNPARGHSVCST
- SSLYLQDLSAAASECIDPSVFPYPLNDSSSPKSCASQDSSAFSPSSDILLS
- STESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEEEIDVVSVEKRQAPGKRS
- 30 ESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPPSTRKDYPAAKRVK
- LDSVRVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELKRSFFA
- LRDQIPELENNEKAPKVVLKATA YILSVQAEEQKLISEEDLLRKRREQ
- KHKLEQLRKGELNSKLEGKPIPPLLGLDSTRTGHHHHH.

6. The pharmaceutical composition according to any of claims 1-5, wherein the antigenic moiety is a pathogen, a toxoid, a peptide, a nucleic acid sequence, a polysaccharide, or a combination thereof.
7. The pharmaceutical composition according to any of claims 1-5, wherein the antigenic moiety is derived from a pathogen selected from: hepatitis A ; hepatitis B ; polio ; measles ; mumps ; rubella ; diphtheria ; pertussis ; tetanus ; influenza ; varicella zoster virus ; rotavirus ; meningococcal ; pneumonia ; smallpox ; cholera ; bubonic plague; yellow fever ; tuberculosis ; human papillomavirus; or combinations thereof.
8. The pharmaceutical composition according to any one of claims 1-7, formulated for topical administration.
9. Use of fusion peptide for the manufacture of a medicament for modulating an immune response in an individual, the fusion peptide comprising:
- a transporter peptide sequence;
 - a MYC sequence; and optionally
 - one or more molecules that link the transporter peptide sequence and the MYC sequence; wherein the medicament is administered before, after, or during the administration of a vaccine.
10. The use of claim 9, wherein the fusion peptide has Formula (I):
transporter peptide sequence -MYC sequence.
11. The use of claim 9, wherein the fusion peptide has Formula (II):
transporter peptide sequence-X-MYC sequence,
wherein -X- is a molecule that links the transporter peptide sequence and the MYC sequence.
12. The use of claim 9, wherein the fusion peptide has Formula (II):
transporter peptide sequence-X-MYC sequence,
wherein in X is at least one amino acid.
13. The use of any one of claims 9-12, wherein the fusion peptide has the following sequence:
- MRKKRRQRRRMDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYF
YCDDEENFYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGLCSPSY
VAVTPFSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPDET
FIKNIHQDCMWSGFSAAKLVSEKLSYQAARKDSGSPNPARGHSVCST
SSLYLQDLSAAASECIDPSVVFYPLNDSSSPKSCASQDSSAFSPSSDILLS
STESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEEIDVVSVEKRQAPGKRS
ESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPSTRKDYPAAKRVK

LDSVRVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELKRSFFA
LRDQIPELENNEKAPKVILKKATAYILSVQAEEQKLISEEDLLRKRREQL
KHKLEQLRKGELNSKLEGKPIPPLLGLDSTRTGHHHHH.

14. The use of any one of claims 9-13, wherein the vaccine is selected from the group
5 consisting of: hepatitis A vaccine; hepatitis B vaccine; polio vaccine; measles vaccine;
mumps vaccine; rubella vaccine; diphtheria vaccine; pertussis vaccine; tetanus vaccine;
influenza vaccine; varicella zoster virus vaccine; rotavirus vaccine; meningococcal
vaccine; pneumonia vaccine; smallpox vaccine; cholera vaccine; bubonic plague vaccine;
yellow fever vaccine; tuberculosis vaccine; human papillomavirus vaccine; a cancer
10 vaccine; or combinations thereof.
15. The use of any one of claims 9-14, wherein the vaccine comprises an antigenic
moiety derived from a neoplastic cell.
16. A method for modulating an immune response in an individual, the method
comprising administering to the individual a fusion peptide comprising:
- 15 (a) a transporter peptide sequence;
(b) a MYC sequence; and optionally
(c) one or more molecules that link the transporter peptide sequence and the MYC
sequence; wherein the fusion protein is administered before, after, or during the
administration of a vaccine.
- 20 17. The method of claim 16, wherein the fusion peptide has Formula (I):
transporter peptide sequence -MYC sequence.
18. The method of claim 16, wherein the fusion peptide has Formula (II):
transporter peptide sequence-X-MYC sequence,
wherein -X- is a molecule that links the transporter peptide sequence and the MYC
25 sequence.
19. The method of claim 16, wherein the fusion peptide has Formula (II):
transporter peptide sequence-X-MYC sequence,
wherein in X is at least one amino acid.
20. The method of any one of claims 16-19, wherein the fusion peptide has the
30 following sequence:
- MRKKRRQRRRMDFFRVVENQPPATMPLNVSFTNRNYDLDYDSVQPYF
YCDEEENFYQQQQSELQPPAPSEDIWKKFELLPTPLSPSRRSGLCSPSY
VAVTPFSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPDET
FIKNIHQDCMWSGFSAAKLVSEKLSYQAARKDSGSPNPARGHSVCST
35 SSLYLQDLSAAASECIDPSVVFYPLNDSSSPKSCASQDSSAFSPSSDLSLS

STESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEEEIDVVSVEKRQAPGKRS
ESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPSTRKDYPAAKRVK
LDSVRVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELKRSFFA
LRDQIPELENNEKAPKVVLKKATAYILSVQAEEQKLISEEDLLRKRREQ
5 KHKLEQLRKGELNSKLEGKPIPPLLGLDSTRTGHHHHHH.

21. The method of any one of claims 16-20, wherein the vaccine is selected from the
group consisting of: hepatitis A vaccine; hepatitis B vaccine; polio vaccine; measles
vaccine; mumps vaccine; rubella vaccine; diphtheria vaccine; pertussis vaccine; tetanus
vaccine; influenza vaccine; varicella zoster virus vaccine; rotavirus vaccine;
10 meningococcal vaccine; pneumonia vaccine; smallpox vaccine; cholera vaccine; bubonic
plague vaccine; yellow fever vaccine; tuberculosis vaccine; human papillomavirus vaccine;
a cancer vaccine; or combinations thereof.

22. The method of any one of claims 16-21, wherein the vaccine comprises an
antigenic moiety derived from a neoplastic cell.

15 23. A pharmaceutical composition for modulating an immune system according to
claim 1, substantially as hereinbefore described with reference to any one of the
examples.

20

Dated 13 December, 2010
Taiga Biotechnologies, Inc.
Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON