



US 20170044538A1

(19) **United States**

(12) **Patent Application Publication**
RIGO et al.

(10) **Pub. No.: US 2017/0044538 A1**

(43) **Pub. Date: Feb. 16, 2017**

(54) **COMPOSITIONS AND METHODS FOR
MODULATION OF SMN2 SPLICING IN A
SUBJECT**

Related U.S. Application Data

(60) Provisional application No. 61/981,139, filed on Apr. 17, 2014.

(71) Applicant: **Biogen MA Inc.**, Cambridge, MA (US)

Publication Classification

(72) Inventors: **Frank RIGO**, Carlsbad, CA (US); **C.
Frank BENNETT**, Carlsbad, CA (US)

(51) **Int. Cl.**
C12N 15/113 (2006.01)

(21) Appl. No.: **15/303,829**

(52) **U.S. Cl.**
CPC **C12N 15/113** (2013.01); **C12N 2310/315**
(2013.01); **C12N 2310/322** (2013.01); **C12N**
2310/11 (2013.01); **C12N 2320/33** (2013.01)

(22) PCT Filed: **Apr. 17, 2015**

(86) PCT No.: **PCT/US2015/026326**

(57) **ABSTRACT**

§ 371 (c)(1),

(2) Date: **Oct. 13, 2016**

Disclosed herein are compounds, compositions and methods for modulating splicing of SMN2 mRNA in a subject.

**COMPOSITIONS AND METHODS FOR
MODULATION OF SMN2 SPLICING IN A
SUBJECT**

SEQUENCE LISTING

[0001] The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled BIOL0245WOSEQ_ST25.txt, created Apr. 3, 2015, which is 44 Kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

FIELD

[0002] Provided herein are methods, compounds, and compositions for modulation of SMN2 splicing in a subject.

BACKGROUND

[0003] Newly synthesized eukaryotic mRNA molecules, known as primary transcripts or pre-mRNA are processed before translation. Processing of the pre-mRNAs includes addition of a 5' methylated cap and an approximately 200-250 base poly(A) tail to the 3' end of the transcript. Processing of mRNA from pre-mRNA also frequently involves splicing of the pre-mRNA, which occurs in the maturation of 90-95% of mammalian mRNAs. Introns (or intervening sequences) are regions of a pre-mRNA (or the DNA encoding it) that are not included in the coding sequence of the mature mRNA. Exons are regions of a primary transcript that remain in the mature mRNA. The exons are spliced together to form the mature mRNA sequence. Splice junctions are also referred to as splice sites with the 5' side of the junction often called the "5' splice site," or "splice donor site" and the 3' side the "3' splice site" or "splice acceptor site." In splicing, the 3' end of an upstream exon is joined to the 5' end of the downstream exon. Thus the unspliced pre-mRNA has an exon/intron junction at the 5' end of an intron and an intron/exon junction at the 3' end of an intron. After the intron is removed, the exons are contiguous at what is sometimes referred to as the exon/exon junction or boundary in the mature mRNA. Cryptic splice sites are those which are less often used but may be used when the usual splice site is blocked or unavailable. Alternative splicing, defined as the splicing together of different combinations of exons, often results in multiple mRNA transcripts from a single gene.

[0004] Up to 50% of human genetic diseases resulting from a point mutation result in aberrant pre-mRNA processing. Such point mutations can either disrupt a current splice site or create a new splice site, resulting in mRNA transcripts comprised of a different combination of exons or with deletions in exons. Point mutations also can result in activation of a cryptic splice site or disrupt regulatory cis elements (i.e. splicing enhancers or silencers) (Cartegni et al., *Nat. Rev. Genet.*, 2002, 3, 285-298; Drawczak et al., *Hum. Genet.*, 1992, 90, 41-54). Antisense oligonucleotides have been used to target mutations that lead to aberrant splicing in several genetic diseases in order to redirect splicing to give a desired splice product (Kole, *Acta Biochimica Polonica*, 1997, 44, 231-238).

[0005] Antisense compounds have also been used to alter the ratio of naturally occurring alternate splice variants such as the long and short forms of Bcl-x pre-mRNA (U.S. Pat. No. 6,172,216; U.S. Pat. No. 6,214,986; Taylor et al., *Nat.*

Biotechnol. 1999, 17, 1097-1100) or to force skipping of specific exons containing premature termination codons (Wilton et al., *Neuromuscul. Disord.*, 1999, 9, 330-338). U.S. Pat. No. 5,627,274 and WO 94/26887 disclose compositions and methods for combating aberrant splicing in a pre-mRNA molecule containing a mutation using antisense oligonucleotides which do not activate RNase H.

[0006] Proximal spinal muscular atrophy (SMA) is a genetic, neurodegenerative disorder characterized by the loss of spinal motor neurons. SMA is an autosomal recessive disease of early onset and is currently the leading cause of death among infants. The severity of SMA varies among patients and has thus been classified into three types. Type I SMA is the most severe form with onset at birth or within 6 months and typically results in death within 2 years. Children with type I SMA are unable to sit or walk. Type II SMA is the intermediate form and patients are able to sit, but cannot stand or walk. Patients with type III SMA, a chronic form of the disease, typically develop SMA after 18 months of age (Lefebvre et al., *Hum. Mol. Genet.*, 1998, 7, 1531-1536).

[0007] The molecular basis of SMA is caused by the loss of both copies of survival motor neuron gene 1 (SMN1), which may also be known as SMN Telomeric, a protein that is part of a multi-protein complex thought to be involved in snRNP biogenesis and recycling. A nearly identical gene, SMN2, which may also be known as SMN Centromeric, exists in a duplicated region on chromosome 5q13 and modulates disease severity. Expression of the normal SMN1 gene results solely in expression of survival motor neuron (SMN) protein. Although SMN1 and SMN2 have the potential to code for the same protein, SMN2 contains a translationally silent mutation at position +6 of exon 7, which results in inefficient inclusion of exon 7 in SMN2 transcripts. Thus, the predominant form of SMN2 is a truncated version, lacking exon 7, which is unstable and inactive (Cartegni and Krainer, *Nat. Genet.*, 2002, 30, 377-384). Expression of the SMN2 gene results in approximately 10-20% of the SMN protein and 80-90% of the unstable/non-functional SMN-delta7 protein. SMN protein plays a well-established role in assembly of the spliceosome and may also mediate mRNA trafficking in the axon and nerve terminus of neurons.

[0008] Antisense technology is an effective means for modulating the expression of one or more specific gene products, including alternative splice products, and is uniquely useful in a number of therapeutic, diagnostic, and research applications. The principle behind antisense technology is that an antisense compound, which hybridizes to a target nucleic acid, modulates gene expression activities such as transcription, splicing or translation through one of a number of antisense mechanisms. The sequence specificity of antisense compounds makes them extremely attractive as tools for target validation and gene functionalization, as well as therapeutics to selectively modulate the expression of genes involved in disease.

[0009] Certain antisense compounds complementary to SMN2 are known in the art. See for example, WO 2007/002390; U.S. 61/168,885; Hua et al., *American J. of Human Genetics* (April 2008) 82, 1-15; Singh et al., *RNA Bio.* 6:3, 1-10 (2009). Certain antisense compounds and methods disclosed herein possess desirable characteristics compared to such compounds and methods known in the art. Chimeric peptide nucleic acid molecules designed to modulate splic-

ing of SMN2 have been described (WO 02/38738; Cartegni and Krainer, *Nat. Struct. Biol.*, 2003, 10, 120-125).

SUMMARY

[0010] In certain embodiments, the present invention provides methods comprising administering to a subject an antisense compound comprising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2 pre-mRNA, wherein the antisense compound is administered into the cerebrospinal fluid. In certain embodiments, the administration is into the intrathecal space. In certain embodiments, the administration is into the cerebrospinal fluid in the brain. In certain embodiments, the administration comprises a bolus injection. In certain embodiments, the administration comprises infusion with a delivery pump.

[0011] In certain embodiments, the antisense compound is administered at a dose from 0.01 to 10 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the dose is from 0.01 to 10 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the dose is from 0.01 to 5 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the dose is from 0.05 to 1 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the dose is from 0.01 to 0.5 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the dose is from 0.05 to 0.5 milligrams of antisense compound per kilogram of body weight of the subject.

[0012] In certain embodiments, the dose is administered daily. In certain embodiments, the dose is administered weekly. In certain embodiments, the antisense compound is administered continuously and wherein the dose is the amount administered per day. In certain embodiments, the method comprises administering at least one induction dose during an induction phase and administering at least one maintenance dose during a maintenance phase. In certain embodiments, the induction dose is from 0.05 to 5.0 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the maintenance dose is from 0.01 to 1.0 milligrams of antisense compound per kilogram of body weight of the subject. In certain embodiments, the duration of the induction phase is at least 1 week. In certain embodiments, the duration of the maintenance phase is at least 1 week. In certain embodiments, each induction dose and each maintenance dose comprises a single injection. In certain embodiments, each induction dose and each maintenance dose independently comprise two or more injections. In certain embodiments, antisense compound is administered at least 2 times over a treatment period of at least 1 week. In certain embodiments, the treatment period is at least one month. In certain embodiments, the treatment period is at least 2 months. In certain embodiments, the treatment period is at least 4 months. In certain embodiments, the induction dose is administered by one or more bolus injections and the maintenance dose is administered by an infusion pump.

[0013] In certain embodiments, the method comprises assessing the tolerability and/or effectiveness of the antisense compound. In certain embodiments, dose amount or frequency of antisense compound is reduced following an indication that administration of the antisense compound is

not tolerated. In certain embodiments, the dose amount or frequency of antisense compound is maintained or reduced following an indication that administration of the antisense compound is effective. In certain embodiments, the dose of antisense compound is increased following an indication that administration of the antisense compound is not effective. In certain embodiments, frequency of administration of antisense compound is reduced following an indication that administration of the antisense compound is effective. In certain embodiments, frequency of administration of antisense compound is increased following an indication that administration of the antisense compound is not effective.

[0014] In certain embodiments, the methods comprise co-administration of the antisense compound and at least one other therapy. In certain embodiments, an antisense compound and at least one other therapy are co-administered at the same time. In certain embodiments, an antisense compound is administered prior to administration of the at least one other therapy. In certain embodiments, an antisense compound is administered after administration of the at least one other therapy. In certain embodiments, the at least one other therapy comprises administration of one or more of valproic acid, riluzole, hydroxyurea, and a butyrate. In certain embodiments, at least one other therapy comprises administration of trichostatin-A. In certain embodiments, the at least one other therapy comprises administration of stem cells. In certain embodiments, at least one other therapy is gene therapy. In certain embodiments, gene therapy is administered to the CSF and an antisense compound is administered systemically. In certain embodiments, gene therapy is administered to the CSF and an antisense compound is administered systemically and to the CSF. In certain embodiments, the invention provides treatment regimens where initially, an antisense compound is administered to the CSF and systemically, followed by gene therapy administration to the CSF and systemic administration of antisense compound. In certain such embodiments, the subject is an infant at the time of initial treatment. In certain such embodiments, the subject is less than 2 years old. In certain embodiments, antisense compound is administered to the CNS of a subject until the subject is old enough for gene therapy. In certain such embodiments, antisense compound is administered systemically throughout.

[0015] In certain embodiments, the antisense compound is administered at a concentration of about 0.01 mg/ml, about 0.05 mg/ml, about 0.1 mg/ml, about 0.5 mg/ml, about 1 mg/ml, about 5 mg/ml, about 10 mg/ml, about 50 mg/ml, or about 100 mg/ml.

[0016] In certain embodiments, inclusion of exon 7 of SMN2 mRNA in a motoneuron in the subject is increased. In certain embodiments, inclusion of exon 7 amino acids in SMN2 polypeptide in a motoneuron in the subject is increased.

[0017] In certain embodiments, the invention provides methods of increasing inclusion of exon 7 of SMN2 mRNA in a motoneuron in a subject comprising administering to the subject an antisense compound comprising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2 and thereby increasing inclusion of exon 7 of SMN2 mRNA in the motoneuron in the subject.

[0018] In certain embodiments, the invention provides methods of increasing inclusion of exon 7 amino acids in SMN2 polypeptide in a motoneuron in a subject comprising administering to the subject an antisense compound com-

prising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2 and thereby increasing inclusion of exon 7 amino acids in SMN2 polypeptide in the motoneuron in the subject.

[0019] In certain embodiments, the subject has SMA. In certain embodiments, the subject has type I SMA. In certain embodiments, the subject has type II SMA. In certain embodiments, the subject has type III SMA.

[0020] In certain embodiments, a first dose is administered in utero. In certain embodiments, the first dose is administered prior to complete formation of the blood-brain-barrier. In certain embodiments, a first dose is administered within 1 week of birth of the subject. In certain embodiments, a first dose is administered within 1 month of birth of the subject. In certain embodiments, a first dose is administered within 3 months of birth of the subject. In certain embodiments, a first dose is administered within 6 months of birth of the subject. In certain embodiments, a first dose is administered when the subject is from 1 to 2 years of age. In certain embodiments, a first dose is administered when the subject is from 1 to 15 years of age. In certain embodiments, a first dose is administered when the subject is older than 15 years of age.

[0021] In certain embodiments, the subject is a mammal. In certain embodiments, the subject is a human.

[0022] In certain embodiments, the methods comprise identifying a subject having SMA. In certain embodiments, the subject is identified by measuring electrical activity of one or more muscles of the subject. In certain embodiments, the subject is identified by a genetic test to determine whether the subject has a mutation in the subject's SMN1 gene. In certain embodiments, the subject is identified by muscle biopsy.

[0023] In certain embodiments, administering the antisense compound results in an increase in the amount of SMN2 mRNA having exon 7 of at least 10%. In certain embodiments, the increase in the amount of SMN2 mRNA having exon 7 is at least 20%. In certain embodiments, the increase in the amount of SMN2 mRNA having exon 7 is at least 50%. In certain embodiments, the amount of SMN2 mRNA having exon 7 is at least 70%.

[0024] In certain embodiments, administering of the antisense compound results in an increase in the amount of SMN2 polypeptide having exon 7 amino acids of at least 10%. In certain embodiments, wherein the increase in the amount of SMN2 polypeptide having exon 7 amino acids is at least 20%. In certain embodiments, the increase in the amount of SMN2 polypeptide having exon 7 amino acids is at least 50%. In certain embodiments, the increase in the amount of SMN2 polypeptide having exon 7 amino acids is at least 70%.

[0025] In certain embodiments, the administering of the antisense compound ameliorates at least one symptom of SMA in the subject. In certain embodiments, the administering of the antisense compound results in improved motor function in the subject. In certain embodiments, the administering of the antisense compound results in delayed or reduced loss of motor function in the subject. In certain embodiments, administering of the antisense compound results in improved respiratory function. In certain embodiments, the administering of the antisense compound results in improved survival.

[0026] In certain embodiments, at least one nucleoside of the antisense oligonucleotide comprises a modified sugar

moiety. In certain embodiments, at least one modified sugar moiety comprises a 2'-methoxyethyl sugar moiety. In certain embodiments, essentially each nucleoside of the antisense oligonucleotide comprises a modified sugar moiety. In certain embodiments, the nucleosides comprising a modified sugar moiety all comprise the same sugar modification. In certain embodiments, wherein each modified sugar moiety comprises a 2'-methoxyethyl sugar moiety. In certain embodiments, each nucleoside of the antisense oligonucleotide comprises a modified sugar moiety. In certain embodiments, the nucleosides all comprise the same sugar modification. In certain embodiments, each modified sugar moiety comprises a 2'-methoxyethyl sugar moiety. In certain embodiments, at least one internucleoside linkage is a phosphorothioate internucleoside linkage. In certain embodiments, each internucleoside linkage is a phosphorothioate internucleoside linkage.

[0027] In certain embodiments, the antisense oligonucleotide consists of 10 to 25 linked nucleosides. In certain embodiments, the antisense oligonucleotide consists of 12 to 22 linked nucleosides. In certain embodiments, the antisense oligonucleotide consists of 15 to 20 linked nucleosides. In certain embodiments, the antisense oligonucleotide consists of 18 linked nucleosides.

[0028] In certain embodiments, the antisense oligonucleotide is at least 90% complementary to the nucleic acid encoding human SMN2. In certain embodiments, the antisense oligonucleotide is fully complementary to the nucleic acid encoding human SMN2. In certain embodiments, the oligonucleotide has a nucleobase sequence comprising at least 10 contiguous nucleobases of the nucleobase sequence SEQ ID NO: 1. In certain embodiments, the oligonucleotide has a nucleobase sequence comprising at least 15 contiguous nucleobases of the nucleobase sequence SEQ ID NO: 1. In certain embodiments, the oligonucleotide has a nucleobase sequence comprising the nucleobase sequence SEQ ID NO: 1. In certain embodiments, the oligonucleotide has a nucleobase sequence consisting of the nucleobase sequence SEQ ID NO: 1.

[0029] In certain embodiments, the antisense compound comprises a conjugate group or terminal group.

[0030] In certain embodiments, the antisense compound consists of the antisense oligonucleotide.

[0031] In certain embodiments, the antisense compound is also administered systemically. In certain embodiments, the systemic administration is by intravenous or intraperitoneal injection. In certain embodiments, systemic administration and the administration into the central nervous system are performed at the same time. In certain embodiments, systemic administration and the administration into the central nervous system are performed at different times.

[0032] In certain embodiments, the invention provides systemic administration of antisense compounds, either alone or in combination with delivery into the CSF. In certain embodiments, pharmaceutical compositions are administered systemically. In certain embodiments, pharmaceutical compositions are administered subcutaneously. In certain embodiments, pharmaceutical compositions are administered intravenously. In certain embodiments, pharmaceutical compositions are administered by intramuscular injection.

[0033] In certain embodiments, pharmaceutical compositions are administered both directly to the CSF (e.g., IT and/or ICV injection and/or infusion) and systemically.

[0034] In certain embodiments, the invention provides methods of administering to a subject having at least one symptom associated with SMA, at least one dose of an antisense compound comprising an oligonucleotide consisting of 15 to 20 linked nucleosides and having a nucleobase sequence which is 100% complementary to SEQ ID NO. 7 over its entire length, and wherein each nucleoside is a 2'-MOE modified nucleoside; and wherein at least one dose is between 0.1 mg/kg and 5 mg/kg administered to the CSF. In certain such embodiments, the dose is between 0.5 mg/kg and 2 mg/kg. In certain embodiments, at least one dose is administered by bolus injection. In certain such embodiments, the dose is administered by bolus intrathecal injection. In certain embodiments, at least one second dose is administered. In certain such embodiments, the second dose is administered at least 2 weeks after the first dose. In certain embodiments, the second dose is administered at least 4 weeks after the first dose. In certain embodiments, the second dose is administered at least 8 weeks after the first dose. In certain embodiments, the second dose is administered at least 12 weeks after the first dose. In certain embodiments, the second dose is administered at least 16 weeks after the first dose. In certain embodiments, the second dose is administered at least 20 weeks after the first dose. In certain embodiments, the subject is under 2 years old at the time of the first dose. In certain embodiments, the subject is between 2 and 15 years old. In certain embodiments, the subject is between 15 and 30 years old. In certain embodiments, the subject is older than 30 years old. In certain embodiments, at least one symptom associated with SMA is reduced its progression has slowed. In certain embodiments, the oligonucleotide is ISIS396443.

[0035] In certain embodiments, the invention provides methods of administering to a subject having at least one symptom associated with SMA, at least one dose of an antisense compound comprising an oligonucleotide consisting of 15 to 20 linked nucleosides and having a nucleobase sequence comprising which is 100% complementary to SEQ ID NO. 7 over its entire length, and wherein each nucleoside is a 2'-MOE modified nucleoside; and wherein at least one dose is administered systemically. In certain such embodiments, at least one dose is administered by bolus injection. In certain such embodiments, the dose is administered by bolus subcutaneous injection. In certain embodiments, the dose administered is between 0.5 mg/kg and 50 mg/kg. In certain embodiments, the dose is between 1 mg/kg and 10 mg/kg. In certain embodiments, the dose is between 1 mg/kg and 5 mg/kg. In certain embodiments, the dose is between 0.5 mg/kg and 1 mg/kg. In certain embodiments, at least one second dose is administered. In certain such embodiments, the second dose is administered at least 2 weeks after the first dose. In certain embodiments, the second dose is administered at least 4 weeks after the first dose. In certain embodiments, the second dose is administered at least 8 weeks after the first dose. In certain embodiments, the second dose is administered at least 12 weeks after the first dose. In certain embodiments, the second dose is administered at least 16 weeks after the first dose. In certain embodiments, the second dose is administered at least 20 weeks after the first dose. In certain embodiments, the subject is under 2 years old at the time of the first dose. In certain embodiments, the subject is between 2 and 15 years old. In certain embodiments, the subject is between 15 and 30 years old. In certain embodiments, the subject is older

than 30 years old. In certain embodiments, at least one symptom associated with SMA is reduced its progression has slowed. In certain embodiments, the oligonucleotide is ISIS396443.

[0036] In certain embodiments, the invention provides methods of administering to a subject having at least one symptom associated with SMA, at least one dose to the CSF and at least one systemic dose of an antisense compound comprising an oligonucleotide consisting of 15 to 20 linked nucleosides and having a nucleobase sequence which is 100% complementary to SEQ ID NO. 7 over its entire length, and wherein each nucleoside is a 2'-MOE modified nucleoside. In certain such embodiments, the CSF dose is between 0.1 mg/kg and 5 mg/kg. In certain embodiments, the systemic dose is between 0.5 mg/kg and 50 mg/kg. In certain embodiments, at least one CSF dose is administered by bolus injection. In certain such embodiments, at least one CSF dose is administered by bolus intrathecal injection. In certain embodiments, at least one systemic dose is administered by bolus injection. In certain such embodiments, at least one systemic dose is administered by subcutaneous injection. In certain embodiments, the CSF dose and the systemic dose are administered at the same time. In certain embodiments, the CSF dose and the systemic dose are administered at different times. In certain embodiments, the subject is under 2 years old at the time of the first dose. In certain embodiments, the subject is between 2 and 15 years old. In certain embodiments, the subject is between 15 and 30 years old. In certain embodiments, the subject is older than 30 years old. In certain embodiments, at least one symptom associated with SMA is reduced its progression has slowed. In certain embodiments, the oligonucleotide is ISIS396443.

[0037] In certain embodiments, the invention provides methods of administering to a subject having at least one symptom associated with SMA, at least one systemic dose of an antisense compound comprising an oligonucleotide consisting of 15 to 20 linked nucleosides and having a nucleobase sequence which is 100% complementary to SEQ ID NO. 7 over its entire length, and wherein each nucleoside is a 2'-MOE modified nucleoside; and at least one dose of a gene therapy agent. In certain embodiments, the systemic dose is between 0.5 mg/kg and 50 mg/kg. In certain embodiments, at least one systemic dose is administered by bolus injection. In certain such embodiments, at least one systemic dose is administered by subcutaneous injection. In certain embodiments, the systemic dose and the gene therapy agent are administered at the same time. In certain embodiments, the systemic dose and the gene therapy agent are administered at different times. In certain embodiments, the gene therapy agent is administered to the CSF. In certain such embodiments, the gene therapy agent is administered by intrathecal injection and/or infusion. In certain such embodiments, the gene therapy agent is administered by intracerebroventricular injection and/or infusion. In certain embodiments, the subject is under 2 years old at the time of the first dose. In certain embodiments, the subject is between 2 and 15 years old. In certain embodiments, the subject is between 15 and 30 years old. In certain embodiments, the subject is older than 30 years old. In certain embodiments, at least one symptom associated with SMA is reduced or its progression has slowed. In certain embodiments, the oligonucleotide is ISIS396443.

[0038] In certain embodiments, the invention provides methods of selecting a subject having at least one symptom associated with SMA and administering an antisense compound according to any of the methods above. In certain such embodiments, at least one symptom of SMA is assessed after administration. In certain such embodiments, at least one symptom of SMA is improved. In certain such embodiments, at least one symptom of SMA does not progress or progresses more slowly compared to a subject who has not received administration of antisense compound.

[0039] In certain embodiments, the invention provides an antisense compound comprising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2, for use in any of the above methods. In certain embodiments, the invention provides such a compound for use in treating a disease or condition associated with survival motor neuron 1 (SMN1).

[0040] In certain embodiments, the invention provides use of an antisense compound comprising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2 in the manufacture of a medicament for use in any of the above methods. In certain embodiments, the medicament is for treating a disease or condition associated with survival motor neuron 1 (SMN1).

[0041] In certain embodiments, the present invention provides methods comprising administering to a subject an antisense compound comprising an antisense oligonucleotide complementary to intron 7 of a nucleic acid encoding human SMN2 pre-mRNA, wherein the antisense compound is administered into the cerebrospinal fluid. In certain embodiments, the administration is into the intrathecal space. In certain embodiments, the administration is into the cerebrospinal fluid in the brain. In certain embodiments, the administration comprises a bolus injection. In certain embodiments, the administration comprises infusion with a delivery pump.

[0042] The present disclosure provides the following non-limiting numbered embodiments:

Embodiment 1

[0043] A compound comprising a modified oligonucleotide consisting of 16-20 linked nucleosides, wherein the modified oligonucleotide is complementary to SMN2; and wherein each internucleoside linkage is either phosphorothioate or phosphodiester and at least one internucleoside linkage is phosphorothioate and at least one internucleoside linkage is phosphodiester.

Embodiment 2

[0044] A compound comprising a modified oligonucleotide consisting of 16-20 linked nucleosides, wherein the modified oligonucleotide has a nucleobase sequence selected from any one of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 and having 6 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 3

[0045] A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, wherein the modified

oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 4

[0046] A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 7, wherein the modified oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 5

[0047] A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 2, wherein the modified oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 6

[0048] The compound of any of embodiments 1 to 5, having 7 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 7

[0049] The compound of any of embodiments 1 to 5, having 8 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 8

[0050] The compound of any of embodiments 1 to 5, having 9 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 9

[0051] The compound of any of embodiments 1 to 5, having 10 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 10

[0052] The compound of embodiment 3, wherein the modified oligonucleotide has an NsNsNoNoNoNsNsNsNoNoNoNoNoNoNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

Embodiment 41

[0083] The compound of embodiment 5, wherein the modified oligonucleotide has an NsNsNoNsNsNoNsNsNoNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

Embodiment 42

[0084] The compound of embodiment 5, wherein the modified oligonucleotide has an NsNsNsNoNsNsNsNoNsNsNsNoNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

Embodiment 43

[0085] The compound of embodiment 5, wherein the modified oligonucleotide has an NsNsNsNsNoNsNsNsNsNoNsNsNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

Embodiment 44

[0086] The compound of any of embodiments 1 to 43, wherein each nucleoside of the modified oligonucleotide comprises a modified sugar moiety, and wherein each modified sugar moiety comprises a 2'-methoxyethyl modification.

Embodiment 45

[0087] A pharmaceutical composition comprising the compound of any of embodiments 1 to 44.

Embodiment 46

[0088] A method of promoting inclusion of exon 7 in SMN2 transcripts in a cell, tissue or organ, comprising contacting said cell, tissue or organ with the compound of any of embodiments 1 to 44 or the composition of embodiment 45.

Embodiment 47

[0089] A method of treating Spinal Muscular Atrophy Type I, comprising administering the compound of any of embodiments 1 to 44 or the composition of embodiment 45 to a patient in need thereof.

Embodiment 48

[0090] A method of treating Spinal Muscular Atrophy Type II, comprising administering the compound of any of embodiments 1 to 44 or the composition of embodiment 45 to a patient in need thereof.

Embodiment 49

[0091] A method of treating Spinal Muscular Atrophy Type III, comprising administering the compound of any of embodiments 1 to 44 or the composition of embodiment 45 to a patient in need thereof.

Embodiment 50

[0092] A method of treating Spinal Muscular Atrophy Type IV, comprising administering the compound of any of embodiments 1 to 44 or the composition of embodiment 45 to a patient in need thereof.

Embodiment 51

[0093] An antisense oligonucleotide of any of embodiments 1 to 44 or the composition of embodiment 45 for use in therapy.

Embodiment 52

[0094] Use of an antisense oligonucleotide of any of embodiments 1 to 44 or the composition of embodiment 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type I.

Embodiment 53

[0095] Use of an antisense oligonucleotide of any of embodiments 1 to 44 or the composition of embodiment 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type II

Embodiment 54

[0096] Use of an antisense oligonucleotide of any of embodiments 1 to 44 or the composition of embodiment 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type III.

Embodiment 55

[0097] Use of an antisense oligonucleotide of any of embodiments 1 to 44 or the composition of embodiment 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type IV.

Embodiment 56

[0098] Use of an antisense oligonucleotide of any of embodiments 1 to 45 for the preparation of a medicament for the treatment of spinal muscular atrophy.

DETAILED DESCRIPTION

[0099] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed. Herein, the use of the singular includes the plural unless specifically stated otherwise. As used herein, the use of “or” means “and/or” unless stated otherwise. Furthermore, the use of the term “including” as well as other forms, such as “includes” and “included”, is not limiting. Also, terms such as “element” or “component” encompass both elements and components comprising one unit and elements and components that comprise more than one subunit, unless specifically stated otherwise.

[0100] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All documents, or portions of documents, cited in this application, including, but not limited to, patents, patent applications, articles, books, and treatises, are hereby expressly incorporated by reference in their entirety for any purpose.

I. DEFINITIONS

[0101] Unless specific definitions are provided, the nomenclature utilized in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, and chemical analysis. Certain such techniques and procedures may be found for example in "Carbohydrate Modifications in Antisense Research" Edited by Sangvi and Cook, American Chemical Society, Washington D.C., 1994; "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., 18th edition, 1990; and "Antisense Drug Technology, Principles, Strategies, and Applications" Edited by Stanley T. Crooke, CRC Press, Boca Raton, Fla.; and Sambrook et al., "Molecular Cloning, A laboratory Manual," 2nd Edition, Cold Spring Harbor Laboratory Press, 1989, which are hereby incorporated by reference for any purpose. Where permitted, all patents, applications, published applications and other publications and other data referred to throughout in the disclosure herein are incorporated by reference in their entirety.

[0102] Unless otherwise indicated, the following terms have the following meanings:

[0103] "Nucleoside" means a compound comprising a heterocyclic base moiety and a sugar moiety. Nucleosides include, but are not limited to, naturally occurring nucleosides, modified nucleosides, and nucleosides having mimetic bases and/or sugar groups. Nucleosides may be modified with any of a variety of substituents.

[0104] "Sugar moiety" means a natural or modified sugar or sugar surrogate.

[0105] "Natural sugar" means a ribofuranose moiety of DNA (2'-H) or RNA (2'-OH).

[0106] "Modified sugar" means a ribofuranose moiety comprising at least one substituent other than that of a natural sugar.

[0107] "Sugar surrogate" means a structure other than a ribofuranose ring which is capable of substituting for the sugar of a nucleoside. Examples of sugar surrogates include, but are not limited to, open ring systems, 6-membered rings, sugars in which the oxygen is replaced with, for example, sulfur or nitrogen. For example, sugar surrogates include, but are not limited to morpholinos and 4'-thio-containing sugars.

[0108] "Nucleobase" means the heterocyclic base portion of a nucleoside. Nucleobases may be naturally occurring or may be modified. In certain embodiments, a nucleobase may comprise any atom or group of atoms capable of hydrogen bonding to a nucleobase of another nucleic acid.

[0109] "Nucleotide" means a nucleoside comprising a phosphate linking group. As used herein, nucleosides include nucleotides.

[0110] "Modified nucleoside" a nucleoside comprising at least one modification compared to naturally occurring RNA or DNA nucleosides. Such modification may be at the sugar moiety and/or at the nucleobase.

[0111] "Bicyclic nucleoside" or "BNA" means a nucleoside wherein the sugar moiety of the nucleoside comprises a bridge connecting two carbon atoms of the sugar ring, thereby forming a bicyclic sugar moiety.

[0112] "4'-2' bicyclic nucleoside" means a bicyclic nucleoside comprising a furanose ring comprising a bridge con-

necting two carbon atoms of the furanose ring connects the 2' carbon atom and the 4' carbon atom of the sugar ring.

[0113] "2'-modified" or "2'-substituted" means a nucleoside comprising a sugar comprising a substituent at the 2' position other than H or OH.

[0114] "2'-OMe" or "2'-OCH₃" or "2'-O-methyl" each means a nucleoside comprising a sugar comprising an —OCH₃ group at the 2' position of the sugar ring.

[0115] "MOE" or "2'-MOE" or "2'-OCH₂CH₂OCH₃" or "2'-O-methoxyethyl" each means a nucleoside comprising a sugar comprising a —OCH₂CH₂OCH₃ group at the 2' position of the sugar ring.

[0116] "Oligonucleotide" means a compound comprising a plurality of linked nucleosides. In certain embodiments, one or more of the plurality of nucleosides is modified. In certain embodiments, an oligonucleotide comprises one or more ribonucleosides (RNA) and/or deoxyribonucleosides (DNA).

[0117] "Oligonucleoside" means an oligonucleotide in which none of the internucleoside linkages contains a phosphorus atom. As used herein, oligonucleotides include oligonucleosides.

[0118] "Modified oligonucleotide" means an oligonucleotide comprising at least one modified nucleoside and/or at least one modified internucleoside linkage.

[0119] "Internucleoside linkage" means a covalent linkage between adjacent nucleosides of an oligonucleotide.

[0120] "Naturally occurring internucleoside linkage" means a 3' to 5' phosphodiester linkage.

[0121] "Modified internucleoside linkage" means any internucleoside linkage other than a naturally occurring internucleoside linkage.

[0122] "Oligomeric compound" means a compound comprising an oligonucleotide. In certain embodiments, an oligomeric compound consists of an oligonucleotide. In certain embodiments, an oligomeric compound further comprises one or more conjugate and/or terminal groups.

[0123] "Antisense compound" means an oligomeric compound, at least a portion of which is at least partially complementary to a target nucleic acid to which it hybridizes, wherein such hybridization results in at least one antisense activity.

[0124] "Antisense oligonucleotide" means an antisense compound wherein the oligomeric compound consists of an oligonucleotide.

[0125] "Antisense activity" refers to any detectable and/or measurable effect attributable to the hybridization of an antisense compound to its target nucleic acid. In certain embodiments, such antisense activity is an increase or decrease in an amount of a nucleic acid or protein. In certain embodiments, such antisense activity is a change in the ratio of splice variants of a nucleic acid or protein. In certain embodiments, such antisense activity is a phenotypic change in a cell and/or subject.

[0126] "Detecting" or "measuring" of antisense activity may be direct or indirect. For example, in certain embodiments, antisense activity is assessed by detecting and/or measuring the amount of target nucleic acid or protein or the relative amounts of splice variants of a target nucleic acid or protein. In certain embodiments, antisense activity is detected by observing a phenotypic change in a cell or animal. In connection with any activity, response, or effect, the terms "detecting" and "measuring," indicate that a test for detecting or measuring is performed. Such detection

and/or measuring may include values of zero. Thus, if a test for detection or measuring results in a finding of no activity (activity of zero), the step of detecting or measuring the activity has nevertheless been performed.

[0127] “Target nucleic acid” refers to any nucleic acid molecule the expression, amount, or activity of which is capable of being modulated by an antisense compound.

[0128] “Target mRNA” means a pre-selected RNA molecule that encodes a protein.

[0129] “Target pre-mRNA” means a pre-selected RNA transcript that has not been fully processed into mRNA. Notably, pre-mRNA includes one or more intron.

[0130] “Target protein” means a protein encoded by a target nucleic acid.

[0131] “Modulation” means to a perturbation of function or activity. In certain embodiments, modulation means an increase in gene expression. In certain embodiments, modulation means a decrease in gene expression.

[0132] “Expression” means any functions and steps by which a gene’s coded information is converted into structures present and operating in a cell.

[0133] “Nucleobase sequence” means the order of contiguous nucleobases, in a 5’ to 3’ orientation, independent of any sugar, linkage, and/or nucleobase modification.

[0134] “Contiguous nucleobases” means nucleobases immediately adjacent to each other in a nucleic acid.

[0135] “Nucleobase complementarity” means the ability of two nucleobases to pair non-covalently via hydrogen bonding.

[0136] “Complementary” means that a first nucleic acid is capable of hybridizing to a second nucleic acid under stringent hybridization conditions. For example, an antisense compound is complementary to its target nucleic acid if it is capable of hybridizing to the target nucleic acid under stringent hybridization conditions.

[0137] “Fully complementary” means each nucleobase of a first nucleic acid is capable of pairing with a nucleobase at each corresponding contiguous position in a second nucleic acid.

[0138] “Percent complementarity” of an antisense compound means the percentage of nucleobases of the antisense compound that are complementary to an equal-length portion of a target nucleic acid. Percent complementarity is calculated by dividing the number of nucleobases of the antisense oligonucleotide that are complementary to nucleobases at corresponding contiguous positions in the target nucleic acid by the total length of the antisense compound.

[0139] “Percent identity” means the number of nucleobases in first nucleic acid that are identical to nucleobases at corresponding positions in a second nucleic acid, divided by the total number of nucleobases in the first nucleic acid.

[0140] “Hybridize” means the annealing of complementary nucleic acids that occurs through nucleobase complementarity.

[0141] “Mismatch” means a nucleobase of a first nucleic acid that is not capable of pairing with a nucleobase at a corresponding position of a second nucleic acid.

[0142] “Identical nucleobase sequence” means having the same nucleobase sequence, independent of any chemical modifications to the nucleosides.

[0143] “Different modifications” or “differently modified” refer to nucleosides or internucleoside linkages that have different nucleoside modifications or internucleoside linkages than one another, including absence of modifications.

Thus, for example, a MOE nucleoside and an unmodified DNA nucleoside are “differently modified,” even though the DNA nucleoside is unmodified. Likewise, DNA and RNA are “differently modified,” even though both are naturally-occurring unmodified nucleosides. Nucleosides that are the same but for comprising different nucleobases are not differently modified, unless otherwise indicated. For example, a nucleoside comprising a 2’-OMe modified sugar and an adenine nucleobase and a nucleoside comprising a 2’-OMe modified sugar and a thymine nucleobase are not differently modified.

[0144] “The same modifications” refer to nucleosides and internucleoside linkages (including unmodified nucleosides and internucleoside linkages) that are the same as one another. Thus, for example, two unmodified DNA nucleosides have “the same modification,” even though the DNA nucleoside is unmodified.

[0145] “Type of modification” or nucleoside of a “type” means the modification of a nucleoside and includes modified and unmodified nucleosides. Accordingly, unless otherwise indicated, a “nucleoside having a modification of a first type” may be an unmodified nucleoside.

[0146] “Separate regions” of an oligonucleotide means a portion of an oligonucleotide wherein the nucleosides and internucleoside linkages within the region all comprise the same modifications; and the nucleosides and/or the internucleoside linkages of any neighboring portions include at least one different modification.

[0147] “Motif” means a pattern of modified and/or unmodified nucleobases, sugars, and/or internucleoside linkages in an oligonucleotide.

[0148] “Fully modified oligonucleotide” means each nucleobase, each sugar, and/or each internucleoside linkage is modified.

[0149] “Uniformly modified oligonucleotide” means each nucleobase, each sugar, and/or each internucleoside linkage has the same modification throughout the modified oligonucleotide.

[0150] “Alternating motif” means an oligonucleotide or a portion thereof, having at least four separate regions of modified nucleosides in a pattern $(AB)_nA_m$, where A represents a region of nucleosides having a first type of modification; B represent a region of nucleosides having a different type of modification; n is 2-15; and m is 0 or 1. Thus, in certain embodiments, alternating motifs include 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 or more alternating regions. In certain embodiments, each A region and each B region independently comprises 1-4 nucleosides.

[0151] “Subject” means a human or non-human animal selected for treatment or therapy.

[0152] “Subject in need thereof” means a subject identified as in need of a therapy or treatment. In such embodiments, a subject has one or more indications of having or developing SMA.

[0153] “Administering” means providing a pharmaceutical agent or composition to a subject, and includes, but is not limited to, administering by a medical professional and self-administering.

[0154] “Parenteral administration,” means administration through injection or infusion. Parenteral administration includes, but is not limited to, subcutaneous administration, intravenous administration, or intramuscular administration.

[0155] “Systemic administration” means administration to an area other than the intended locus of activity. Examples

or systemic administration are subcutaneous administration and intravenous administration, and intraperitoneal administration.

[0156] “Subcutaneous administration” means administration just below the skin.

[0157] “Intravenous administration” means administration into a vein.

[0158] “Cerebrospinal fluid” or “CSF” means the fluid filling the space around the brain and spinal cord.

[0159] “Administration into the cerebrospinal fluid” means any administration that delivers a substance directly into the CSF.

[0160] “Intracerebroventricular” or “ICV” mean administration into the ventricular system of the brain.

[0161] “Intrathecal” or “IT” means administration into the CSF under the arachnoid membrane which covers the brain and spinal cord. IT injection is performed through the theca of the spinal cord into the subarachnoid space, where a pharmaceutical agent is injected into the sheath surrounding the spinal cord.

[0162] “Induction phase” means a dosing phase during which administration is initiated and steady state concentrations of active pharmaceutical agent are achieved in a target tissue. For example, an induction phase is a dosing phase during which steady state concentrations of antisense oligonucleotide are achieved in liver.

[0163] “Maintenance phase” means a dosing phase after target tissue steady state concentrations of drug have been achieved.

[0164] “Duration” means the period of time during which an activity or event continues. For example, the duration of an induction phase is the period of time during which induction doses are administered.

[0165] “Maintenance dose” means a dose administered at a single administration during the maintenance phase. As used herein, “induction dose” means a dose administered at a single administration during the induction phase.

[0166] “Co-administration” means administration of two or more pharmaceutical agents to a subject. The two or more pharmaceutical agents may be in a single pharmaceutical composition, or may be in separate pharmaceutical compositions. Each of the two or more pharmaceutical agents may be administered through the same or different routes of administration. Co-administration encompasses administration in parallel or sequentially.

[0167] “Therapy” means a disease treatment method. In certain embodiments, therapy includes, but is not limited to surgical therapies, chemical therapies, and physical interventions, such as assisted respiration, feeding tubes, and physical therapy for the purpose of increasing strength.

[0168] “Treatment” means the application of one or more specific procedures used for the cure or amelioration of a disease. In certain embodiments, the specific procedure is the administration of one or more pharmaceutical agents.

[0169] “Amelioration” means a lessening of severity of at least one indicator of a condition or disease. In certain embodiments, amelioration includes a delay or slowing in the progression of one or more indicators of a condition or disease. The severity of indicators may be determined by subjective or objective measures which are known to those skilled in the art.

[0170] “Prevent the onset of” means to prevent the development a condition or disease in a subject who is at risk for developing the disease or condition. In certain embodiments,

a subject at risk for developing the disease or condition receives treatment similar to the treatment received by a subject who already has the disease or condition.

[0171] “Delay the onset of” means to delay the development of a condition or disease in a subject who is at risk for developing the disease or condition.

[0172] “Slow the progression of” means that the severity of at least one symptom associated with a disease or condition worsens less quickly.

[0173] “Exon 7 amino acids” means the portion of an SMN protein that correspond to exon 7 of the SMN RNA. Exon 7 amino acids are present in SMN protein expressed from SMN RNA where exon 7 was not excluded during splicing.

[0174] “SMN protein” means normal full length survival motor neuron protein. SMN may be expressed from either an SMN1 gene or from an SMN2 gene, provided that exon 7 is present in the mature mRNA and the exon 7 amino acids are present in the SMN protein.

[0175] “Dose” means a specified quantity of a pharmaceutical agent provided in a single administration or over a specified amount of time. In certain embodiments, a dose may be administered in two or more boluses, tablets, or injections. For example, in certain embodiments, where subcutaneous or intrathecal or ICV administration is desired, the desired dose requires a volume not easily accommodated by a single injection. In such embodiments, two or more injections may be used to achieve the desired dose. In the setting of continuous infusion, dose may be expressed as the quantity of a pharmaceutical agent delivered per unit of time.

[0176] “Dosage unit” means a form in which a pharmaceutical agent is provided. In certain embodiments, a dosage unit is a vial containing lyophilized oligonucleotide. In certain embodiments, a dosage unit is a vial containing reconstituted oligonucleotide.

[0177] “Therapeutically effective amount” means an amount of a pharmaceutical agent that provides a therapeutic benefit to an animal.

[0178] “Pharmaceutical composition” means a mixture of substances suitable for administering to an individual that includes a pharmaceutical agent. For example, a pharmaceutical composition may comprise a modified oligonucleotide and a sterile aqueous solution.

[0179] “Acceptable safety profile” means a pattern of side effects that is within clinically acceptable limits.

[0180] “Side effect” means a physiological response attributable to a treatment other than desired effects.

1. Certain Modified Oligonucleotides

[0181] In certain embodiments, the present invention provides methods and compositions involving antisense oligonucleotides comprising one or more modification compared to oligonucleotides of naturally occurring oligomers, such as DNA or RNA. Such modified antisense oligonucleotides may possess one or more desirable properties. Certain such modifications alter the antisense activity of the antisense oligonucleotide, for example by increasing affinity of the antisense oligonucleotide for its target nucleic acid, increasing its resistance to one or more nucleases, and/or altering the pharmacokinetics or tissue distribution of the oligonucleotide. In certain embodiments, such modified antisense oligonucleotides comprise one or more modified nucleo-

sides and/or one or more modified nucleoside linkages and/or one or more conjugate groups.

[0182] a. Certain Modified Nucleosides

[0183] In certain embodiments, antisense oligonucleotides comprise one or more modified nucleosides. Such modified nucleosides may include a modified sugar and/or a modified nucleobase. In certain embodiments, incorporation of such modified nucleosides in an oligonucleotide results in increased affinity for a target nucleic acid and/or increased stability, including but not limited to, increased resistance to nuclease degradation, and or improved toxicity and/or uptake properties of the modified oligonucleotide.

[0184] i. Certain Nucleobases

[0185] The naturally occurring base portion of nucleosides are heterocyclic base, typically purines and pyrimidines. In addition to “unmodified” or “natural” nucleobases such as the purine nucleobases adenine (A) and guanine (G), and the pyrimidine nucleobases thymine (T), cytosine (C) and uracil (U), many modified nucleobases or nucleobase mimetics known to those skilled in the art are amenable to incorporation into the compounds described herein. In certain embodiments, a modified nucleobase is a nucleobase that is fairly similar in structure to the parent nucleobase, such as for example a 7-deaza purine, a 5-methyl cytosine, or a G-clamp. In certain embodiments, nucleobase mimetic include more complicated structures, such as for example a tricyclic phenoxazine nucleobase mimetic. Methods for preparation of the above noted modified nucleobases are well known to those skilled in the art.

[0186] ii. Certain Modified Sugars and Sugar Surrogates

[0187] Antisense oligonucleotides of the present invention can optionally contain one or more nucleosides wherein the sugar moiety is modified, compared to a natural sugar. Oligonucleotides comprising such sugar modified nucleosides may have enhanced nuclease stability, increased binding affinity or some other beneficial biological property. Such modifications include without limitation, addition of substituent groups, bridging of non-geminal ring atoms to form a bicyclic nucleic acid (BNA), replacement of the ribosyl ring oxygen atom with S, N(R), or C(R₁)(R)₂ (R=H, C₁-C₁₂ alkyl or a protecting group) and combinations of these such as for example a 2'-F-5'-methyl substituted nucleoside (see PCT International Application WO 2008/101157 Published on Aug. 21, 2008 for other disclosed 5',2'-bis substituted nucleosides) or replacement of the ribosyl ring oxygen atom with S with further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on Jun. 16, 2005) or alternatively 5'-substitution of a BNA (see PCT International Application WO 2007/134181 Published on Nov. 22, 2007 wherein LNA is substituted with for example a 5'-methyl or a 5'-vinyl group).

[0188] Examples of nucleosides having modified sugar moieties include without limitation nucleosides comprising 5'-vinyl, 5'-methyl (R or S), 4'-S, 2'-F, 2'-OCH₃ and 2'-O(CH₂)₂OCH₃ substituent groups. The substituent at the 2' position can also be selected from allyl, amino, azido, thio, O-allyl, O—C₁-C₁₀ alkyl, OCF₃, O(CH₂)₂SCH₃, O(CH₂)₂—O—N(R_m)(R_n), and O—CH₂—C(=O)—N(R_m)(R_n), where each R_m and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl.

[0189] Examples of bicyclic nucleic acids (BNAs) include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments,

antisense compounds provided herein include one or more BNA nucleosides wherein the bridge comprises one of the formulas: 4'-β-D-(CH₂)—O-2' (β-D-LNA); 4'-(CH₂)—S-2'; 4'-α-L-(CH₂)—O-2' (α-L-LNA); 4'-(CH₂)₂—O-2' (ENA); 4'-C(CH₃)₂—O-2' (see PCT/US2008/068922); 4'-CH(CH₃)—O-2' and 4'-C—H(CH₂OCH₃)—O-2' (see U.S. Pat. No. 7,399,845, issued on Jul. 15, 2008); 4'-CH₂—N(OCH₃)-2' (see PCT/US2008/064591); 4'-CH₂—O—N(CH₃)-2' (see published U.S. Patent Application US2004-0171570, published Sep. 2, 2004); 4'-CH₂—N(R)—O-2' (see U.S. Pat. No. 7,427,672, issued on Sep. 23, 2008); 4'-CH₂—C(CH₃)-2' and 4'-CH₂—C(=CH₂)-2' (see PCT/US2008/066154); and wherein R is, independently, H, C₁-C₁₂ alkyl, or a protecting group.

[0190] In certain embodiments, the present invention provides modified nucleosides comprising modified sugar moieties that are not bicyclic sugar moieties. Certain such modified nucleosides are known. In certain embodiments, the sugar ring of a nucleoside may be modified at any position. Examples of sugar modifications useful in this invention include, but are not limited to compounds comprising a sugar substituent group selected from: OH, F, O-alkyl, S-alkyl, N-alkyl, or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. In certain such embodiments, such substituents are at the 2' position of the sugar.

[0191] In certain embodiments, modified nucleosides comprise a substituent at the 2' position of the sugar. In certain embodiments, such substituents are selected from among: a halide (including, but not limited to F), allyl, amino, azido, thio, O-allyl, O—C₁-C₁₀ alkyl, —OCF₃, O—(CH₂)₂—O—CH₃, 2'-O(CH₂)₂SCH₃, O—(CH₂)₂—O—N(R_m)(R_n), or O—CH₂—C(=O)—N(R_m)(R_n), where each R_m and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl.

[0192] In certain embodiments, modified nucleosides suitable for use in the present invention are: 2-methoxyethoxy, 2'-O-methyl (2'-O—CH₃), 2'-fluoro (2'-F).

[0193] In certain embodiments, modified nucleosides having a substituent group at the 2'-position selected from: O[(CH₂)_nO]_mCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, OCH₂C(=O)N(H)CH₃, and O(CH₂)_nON[(CH₂)CH₃]₂, where n and m are from 1 to about 10. Other 2'-sugar substituent groups include: C₁ to C₁₀ alkyl, substituted alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving pharmacokinetic properties, or a group for improving the pharmacodynamic properties of an oligomeric compound, and other substituents having similar properties.

[0194] In certain embodiments, modified nucleosides comprise a 2'-MOE side chain (Baker et al., J. Biol. Chem., 1997, 272, 11944-12000). Such 2'-MOE substitution have been described as having improved binding affinity compared to unmodified nucleosides and to other modified nucleosides, such as 2'-O-methyl, O-propyl, and O-amino-propyl. Oligonucleotides having the 2'-MOE substituent also have been shown to be antisense inhibitors of gene expression with promising features for in vivo use (Martin, P., Helv. Chim. Acta, 1995, 78, 486-504; Altmann et al.,

Chimia, 1996, 50, 168-176; Altmann et al., Biochem. Soc. Trans., 1996, 24, 630-637; and Altmann et al., Nucleosides Nucleotides, 1997, 16, 917-926).

[0195] In certain embodiments, 2'-sugar substituent groups are in either the arabino (up) position or ribo (down) position. In certain such embodiments, a 2'-arabino modification is 2'-F arabino (FANA). Similar modifications can also be made at other positions on the sugar, particularly the 3' position of the sugar on a 3' terminal nucleoside or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide.

[0196] In certain embodiments, nucleosides suitable for use in the present invention have sugar surrogates such as cyclobutyl in place of the ribofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, each of which is herein incorporated by reference in its entirety.

[0197] In certain embodiments, the present invention provides nucleosides comprising a modification at the 2'-position of the sugar. In certain embodiments, the invention provides nucleosides comprising a modification at the 5'-position of the sugar. In certain embodiments, the invention provides nucleosides comprising modifications at the 2'-position and the 5'-position of the sugar. In certain embodiments, modified nucleosides may be useful for incorporation into oligonucleotides. In certain embodiment, modified nucleosides are incorporated into oligonucleosides at the 5'-end of the oligonucleotide.

[0198] b. Certain Internucleoside Linkages

[0199] Antisense oligonucleotides of the present invention can optionally contain one or more modified internucleoside linkages. The two main classes of linking groups are defined by the presence or absence of a phosphorus atom. Representative phosphorus containing linkages include, but are not limited to, phosphodiester (P=O), phosphotriesters, methylphosphonates, phosphoramidate, and phosphorothioates (P=S). Representative non-phosphorus containing linking groups include, but are not limited to, methylenemethylimino ($-\text{CH}_2\text{-N}(\text{CH}_3)\text{-O-CH}_2-$), thiodiester ($-\text{O}-\text{C}(\text{O})-\text{S}-$), thionocarbamate ($-\text{O}-\text{C}(\text{O})(\text{NH})-\text{S}-$); siloxane ($-\text{O}-\text{Si}(\text{H})_2\text{-O}-$); and N,N'-dimethylhydrazine ($-\text{CH}_2\text{-N}(\text{CH}_3)\text{-N}(\text{CH}_3)-$). Oligonucleotides having non-phosphorus linking groups are referred to as oligonucleosides. Modified linkages, compared to natural phosphodiester linkages, can be used to alter, typically increase, nuclease resistance of the oligonucleotides. In certain embodiments, linkages having a chiral atom can be prepared as racemic mixtures, as separate enantiomers. Representative chiral linkages include, but are not limited to, alkylphosphonates and phosphorothioates. Methods of preparation of phosphorus-containing and non-phosphorus-containing linkages are well known to those skilled in the art.

[0200] The antisense oligonucleotides described herein contain one or more asymmetric centers and thus give rise to enantiomers, diastereomers, and other stereoisomeric configurations that may be defined, in terms of absolute stereochemistry, as (R) or (S), such as for sugar anomers, or as (D) or (L) such as for amino acids et al. Included in the

antisense compounds provided herein are all such possible isomers, as well as their racemic and optically pure forms.

[0201] In certain embodiments, antisense oligonucleotides have at least one modified internucleoside linkage. In certain embodiments, antisense oligonucleotides have at least 2 modified internucleoside linkages. In certain embodiments, antisense oligonucleotides have at least 3 modified internucleoside linkages. In certain embodiments, antisense oligonucleotides have at least 10 modified internucleoside linkages. In certain embodiments, each internucleoside linkage of an antisense oligonucleotide is a modified internucleoside linkage. In certain embodiments, such modified internucleoside linkages are phosphorothioate linkages.

[0202] c. Certain Internucleoside Linkage Motifs

[0203] In certain embodiments, oligonucleotides comprise modified internucleoside linkages arranged along the oligonucleotide or region thereof in a defined pattern or modified internucleoside linkage motif.

[0204] In certain embodiments, oligonucleotides comprise a region having an alternating internucleoside linkage motif. In certain embodiments, oligonucleotides of the present invention comprise a region of uniformly modified internucleoside linkages. In certain such embodiments, the oligonucleotide comprises a region that is uniformly linked by phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide is uniformly linked by phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate and at least one internucleoside linkage is phosphorothioate.

[0205] In certain embodiments, the oligonucleotide comprises at least 5 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 6 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 7 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 8 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 9 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 10 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 11 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 12 phosphorothioate internucleoside linkages.

[0206] In certain embodiments, the oligonucleotide comprises at least one block of at least 2 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 3 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 4 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 5 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 6 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 7 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least

having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 7 month intervals.

[0218] In certain embodiments, the inclusion of 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at infrequent intervals. In certain embodiments, the inclusion of 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 3 month intervals. In certain embodiments, the inclusion of 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 4 month intervals. In certain embodiments, the inclusion of 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 5 month intervals. In certain embodiments, the inclusion of 3, 4, 5, 6, 7, 8, or 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 6 month intervals. In certain embodiments, the inclusion of 9 phosphodiester internucleoside linkages into a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, allows one having skill in the art to administer the modified oligonucleotide or a pharmaceutical composition thereof at greater than 7 month intervals.

[0219] d. Lengths

[0220] In certain embodiments, the present invention provides antisense oligonucleotides of any of a variety of ranges of lengths. In certain embodiments, the invention provides antisense compounds or antisense oligonucleotides comprising or consisting of X-Y linked nucleosides, where X and Y are each independently selected from 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, and 50; provided that X<Y. For example, in certain embodiments, the invention provides antisense compounds or antisense oligonucleotides comprising or consisting of: 8-9, 8-10, 8-11, 8-12, 8-13, 8-14, 8-15, 8-16, 8-17, 8-18, 8-19, 8-20, 8-21, 8-22, 8-23, 8-24, 8-25, 8-26, 8-27, 8-28, 8-29, 8-30, 9-10, 9-11, 9-12, 9-13, 9-14, 9-15, 9-16, 9-17, 9-18, 9-19, 9-20, 9-21, 9-22, 9-23, 9-24, 9-25, 9-26, 9-27, 9-28, 9-29, 9-30, 10-11, 10-12, 10-13, 10-14, 10-15, 10-16, 10-17, 10-18, 10-19, 10-20, 10-21, 10-22, 10-23, 10-24, 10-25, 10-26, 10-27, 10-28, 10-29, 10-30, 11-12, 11-13, 11-14, 11-15, 11-16, 11-17, 11-18, 11-19, 11-20, 11-21, 11-22, 11-23, 11-24, 11-25, 11-26, 11-27, 11-28, 11-29, 11-30, 12-13, 12-14, 12-15, 12-16, 12-17, 12-18, 12-19, 12-20, 12-21, 12-22, 12-23, 12-24, 12-25, 12-26, 12-27, 12-28, 12-29, 12-30, 13-14, 13-15, 13-16, 13-17,

13-18, 13-19, 13-20, 13-21, 13-22, 13-23, 13-24, 13-25, 13-26, 13-27, 13-28, 13-29, 13-30, 14-15, 14-16, 14-17, 14-18, 14-19, 14-20, 14-21, 14-22, 14-23, 14-24, 14-25, 14-26, 14-27, 14-28, 14-29, 14-30, 15-16, 15-17, 15-18, 15-19, 15-20, 15-21, 15-22, 15-23, 15-24, 15-25, 15-26, 15-27, 15-28, 15-29, 15-30, 16-17, 16-18, 16-19, 16-20, 16-21, 16-22, 16-23, 16-24, 16-25, 16-26, 16-27, 16-28, 16-29, 16-30, 17-18, 17-19, 17-20, 17-21, 17-22, 17-23, 17-24, 17-25, 17-26, 17-27, 17-28, 17-29, 17-30, 18-19, 18-20, 18-21, 18-22, 18-23, 18-24, 18-25, 18-26, 18-27, 18-28, 18-29, 18-30, 19-20, 19-21, 19-22, 19-23, 19-24, 19-25, 19-26, 19-29, 19-28, 19-29, 19-30, 20-21, 20-22, 20-23, 20-24, 20-25, 20-26, 20-27, 20-28, 20-29, 20-30, 21-22, 21-23, 21-24, 21-25, 21-26, 21-27, 21-28, 21-29, 21-30, 22-23, 22-24, 22-25, 22-26, 22-27, 22-28, 22-29, 22-30, 23-24, 23-25, 23-26, 23-27, 23-28, 23-29, 23-30, 24-25, 24-26, 24-27, 24-28, 24-29, 24-30, 25-26, 25-27, 25-28, 25-29, 25-30, 26-27, 26-28, 26-29, 26-30, 27-28, 27-29, 27-30, 28-29, 28-30, or 29-30 linked nucleosides.

[0221] In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 15 nucleosides in length. In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 16 nucleosides in length. In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 17 nucleosides in length. In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 18 nucleosides in length. In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 19 nucleosides in length. In certain embodiments, antisense compounds or antisense oligonucleotides of the present invention are 20 nucleosides in length.

[0222] e. Certain Oligonucleotide Motifs

[0223] In certain embodiments, antisense oligonucleotides have chemically modified subunits arranged in specific orientations along their length. In certain embodiments, antisense oligonucleotides of the invention are fully modified. In certain embodiments, antisense oligonucleotides of the invention are uniformly modified. In certain embodiments, antisense oligonucleotides of the invention are uniformly modified and each nucleoside comprises a 2'-MOE sugar moiety. In certain embodiments, antisense oligonucleotides of the invention are uniformly modified and each nucleoside comprises a 2'-OMe sugar moiety. In certain embodiments, antisense oligonucleotides of the invention are uniformly modified and each nucleoside comprises a morpholino sugar moiety.

[0224] In certain embodiments, oligonucleotides of the invention comprise an alternating motif. In certain such embodiments, the alternating modification types are selected from among 2'-MOE, 2'-F, a bicyclic sugar-modified nucleoside, and DNA (unmodified 2'-deoxy). In certain such embodiments, each alternating region comprises a single nucleoside.

[0225] In certain embodiments, oligonucleotides of the invention comprise one or more block of nucleosides of a first type and one or more block of nucleosides of a second type.

[0226] In certain embodiments, one or more alternating regions in an alternating motif include more than a single nucleoside of a type. For example, oligomeric compounds of the present invention may include one or more regions of any of the following nucleoside motifs:

- [0227] Nu₁ Nu₁ Nu₂ Nu₂ Nu₁ Nu₁;
 [0228] Nu₁ Nu₂ Nu₂ Nu₁ Nu₂ Nu₂;
 [0229] Nu₁ Nu₁ Nu₂ Nu₁ Nu₁ Nu₂;
 [0230] Nu₁ Nu₂ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁ Nu₂ Nu₂;
 [0231] Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁;
 [0232] Nu₁ Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₂;
 [0233] Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁;
 [0234] Nu₁ Nu₂ Nu₂ Nu₁ Nu₁ Nu₂ Nu₂ Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁;
 [0235] Nu₂ Nu₁ Nu₂ Nu₂ Nu₁ Nu₁ Nu₂ Nu₂ Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁;
 [0236] Nu₁ Nu₂ Nu₁ Nu₂ Nu₂ Nu₁ Nu₁ Nu₂ Nu₂ Nu₁ Nu₂ Nu₁ Nu₂ Nu₁ Nu₁;

wherein Nu₁ is a nucleoside of a first type and Nu₂ is a nucleoside of a second type. In certain embodiments, one of Nu₁ and Nu₂ is a 2'-MOE nucleoside and the other of Nu₁ and Nu₂ is selected from: a 2'-OMe modified nucleoside, BNA, and an unmodified DNA or RNA nucleoside.

2. Oligomeric Compounds

[0237] In certain embodiments, the present invention provides oligomeric compounds. In certain embodiments, oligomeric compounds are comprised only of an oligonucleotide. In certain embodiments, an oligomeric compound comprises an oligonucleotide and one or more conjugate and/or terminal group. Such conjugate and/or terminal groups may be added to oligonucleotides having any of the chemical motifs discussed above. Thus, for example, an oligomeric compound comprising an oligonucleotide having region of alternating nucleosides may comprise a terminal group.

[0238] a. Certain Conjugate Groups

[0239] In certain embodiments, oligonucleotides of the present invention are modified by attachment of one or more conjugate groups. In general, conjugate groups modify one or more properties of the attached oligomeric compound including but not limited to, pharmacodynamics, pharmacokinetics, stability, binding, absorption, cellular distribution, cellular uptake, charge and clearance. Conjugate groups are routinely used in the chemical arts and are linked directly or via an optional conjugate linking moiety or conjugate linking group to a parent compound such as an oligomeric compound, such as an oligonucleotide. Conjugate groups includes without limitation, intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, thioethers, polyethers, cholesterols, thiocholesterols, cholic acid moieties, folate, lipids, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, adamantane, acridine, fluoresceins, rhodamines, coumarins and dyes. Certain conjugate groups have been described previously, for example: cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., do-decan-diol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654;

Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

[0240] In certain embodiments, a conjugate group comprises an active drug substance, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fen-bufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folic acid, a benzothiadiazide, chlorothiazide, a diazepam, indo-methicin, a barbiturate, a cephalosporin, a sulfa drug, an antiabiotic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in U.S. patent application Ser. No. 09/334,130.

[0241] Representative U.S. patents that teach the preparation of oligonucleotide conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

[0242] Conjugate groups may be attached to either or both ends of an oligonucleotide (terminal conjugate groups) and/or at any internal position.

[0243] b. Terminal Groups

[0244] In certain embodiments, oligomeric compounds comprise terminal groups at one or both ends. In certain embodiments, a terminal group may comprise any of the conjugate groups discussed above. In certain embodiments, terminal groups may comprise additional nucleosides and/or inverted abasic nucleosides. In certain embodiments, a terminal group is a stabilizing group.

[0245] In certain embodiments, oligomeric compounds comprise one or more terminal stabilizing group that enhances properties such as for example nuclease stability. Included in stabilizing groups are cap structures. The terms "cap structure" or "terminal cap moiety," as used herein, refer to chemical modifications, which can be attached to one or both of the termini of an oligomeric compound. Certain such terminal modifications protect the oligomeric compounds having terminal nucleic acid moieties from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap) or at the 3'-terminus (3'-cap) or can be present on both termini. (for more details see Wincott et al., International PCT publication No. WO 97/26270; Beaucage and Tyer, 1993, Tetrahedron 49, 1925; U.S. Patent Application Publication No. US 2005/0020525; and WO 03/004602.

[0246] In certain embodiments, one or more additional nucleosides is added to one or both terminal ends of an oligonucleotide of an oligomeric compound. Such additional terminal nucleosides are referred to herein as terminal-group

nucleosides. In a double-stranded compound, such terminal-group nucleosides are terminal (3' and/or 5') overhangs. In the setting of double-stranded antisense compounds, such terminal-group nucleosides may or may not be complementary to a target nucleic acid. In certain embodiments, the terminal group is a non-nucleoside terminal group. Such non-terminal groups may be any terminal group other than a nucleoside.

[0247] c. Oligomeric Compound Motifs

In certain embodiments, oligomeric compounds of the present invention comprise a motif:

$T_1-(Nu_1)_{n1}-(Nu_2)_{n2}-(Nu_1)_{n3}-(Nu_2)_{n4}-(Nu_1)_{n5}-T_2$, wherein:

[0248] Nu_1 , is a nucleoside of a first type;

[0249] Nu_2 , is a nucleoside of a second type;

[0250] each of $n1$ and $n5$ is, independently from 0 to 3;

[0251] the sum of $n2$ plus $n4$ is between 10 and 25;

[0252] $n3$ is from 0 and 5; and

[0253] each T_1 and T_2 is, independently, H, a hydroxyl protecting group, an optionally linked conjugate group or a capping group.

In certain such embodiments, the sum of $n2$ and $n4$ is 13 or 14; $n1$ is 2; $n3$ is 2 or 3; and $n5$ is 2. In certain such embodiments, oligomeric compounds of the present invention comprise a motif selected from Table A.

TABLE A

n1	n2	n3	n4	n5
2	16	0	0	2
2	2	3	11	2
2	5	3	8	2
2	8	3	5	2
2	11	3	2	2
2	9	3	4	2
2	10	3	3	2
2	3	3	10	2
2	4	3	9	2
2	6	3	7	2
2	7	3	6	2
2	8	6	2	2
2	2	2	12	2
2	3	2	11	2
2	4	2	10	2
2	5	2	9	2
2	6	2	8	2
2	7	2	7	2
2	8	2	6	2
2	9	2	5	2
2	10	2	4	2
2	11	2	3	2
2	12	2	2	2

[0254] Table A is intended to illustrate, but not to limit the present invention. The oligomeric compounds depicted in Table A each comprise 20 nucleosides. Oligomeric compounds comprising more or fewer nucleosides can easily be designed by selecting different numbers of nucleosides for one or more of $n1$ - $n5$. In certain embodiments, Nu_1 and Nu_2 are each selected from among: 2'-MOE, 2'-OMe, DNA, and a bicyclic nucleoside.

3. Antisense

[0255] In certain embodiments, oligomeric compounds of the present invention are antisense compounds. Accordingly, in such embodiments, oligomeric compounds hybridize with a target nucleic acid, resulting in an antisense activity.

[0256] a. Hybridization

[0257] In certain embodiments, the invention provides antisense compounds that specifically hybridize to a target nucleic acid when there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays.

[0258] Thus, "stringent hybridization conditions" or "stringent conditions" means conditions under which an antisense compounds hybridize to a target sequence, but to a minimal number of other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances, and "stringent conditions" under which antisense oligonucleotides hybridize to a target sequence are determined by the nature and composition of the antisense oligonucleotides and the assays in which they are being investigated.

[0259] It is understood in the art that incorporation of nucleotide affinity modifications may allow for a greater number of mismatches compared to an unmodified compound. Similarly, certain nucleobase sequences may be more tolerant to mismatches than other nucleobase sequences. One of ordinary skill in the art is capable of determining an appropriate number of mismatches between oligonucleotides, or between an antisense oligonucleotide and a target nucleic acid, such as by determining melting temperature (T_m). T_m or ΔT_m can be calculated by techniques that are familiar to one of ordinary skill in the art. For example, techniques described in Freier et al. (Nucleic Acids Research, 1997, 25, 22: 4429-4443) allow one of ordinary skill in the art to evaluate nucleotide modifications for their ability to increase the melting temperature of an RNA:DNA duplex.

[0260] b. pre-mRNA Processing

[0261] In certain embodiments, antisense compounds provided herein are complementary to a pre-mRNA. In certain embodiments, such antisense compounds alter splicing of the pre-mRNA. In certain such embodiments, the ratio of one variant of a mature mRNA corresponding to a target pre-mRNA to another variant of that mature mRNA is altered. In certain such embodiments, the ratio of one variant of a protein expressed from the target pre-mRNA to another variant of the protein is altered. Certain oligomeric compounds and nucleobase sequences that may be used to alter splicing of a pre-mRNA may be found for example in U.S. Pat. No. 6,210,892; U.S. Pat. No. 5,627,274; U.S. Pat. Nos. 5,665,593; 5,916,808; U.S. Pat. No. 5,976,879; US2006/0172962; US2007/002390; US2005/0074801; US2007/0105807; US2005/0054836; WO 2007/090073; WO2007/047913, Hua et al., PLoS Biol 5(4):e73; Vickers et al., J. Immunol. 2006 Mar. 15; 176(6):3652-61; and Hua et al., American J. of Human Genetics (April 2008) 82, 1-15, each of which is hereby incorporated by reference in its entirety for any purpose. In certain embodiments antisense sequences that alter splicing are modified according to motifs of the present invention.

[0262] Antisense is an effective means for modulating the expression of one or more specific gene products and is uniquely useful in a number of therapeutic, diagnostic, and research applications. Provided herein are antisense compounds useful for modulating gene expression via antisense

mechanisms of action, including antisense mechanisms based on target occupancy. In one aspect, the antisense compounds provided herein modulate splicing of a target gene. Such modulation includes promoting or inhibiting exon inclusion. Further provided herein are antisense compounds targeted to cis splicing regulatory elements present in pre-mRNA molecules, including exonic splicing enhancers, exonic splicing silencers, intronic splicing enhancers and intronic splicing silencers. Disruption of cis splicing regulatory elements is thought to alter splice site selection, which may lead to an alteration in the composition of splice products.

[0263] Processing of eukaryotic pre-mRNAs is a complex process that requires a multitude of signals and protein factors to achieve appropriate mRNA splicing. Exon definition by the spliceosome requires more than the canonical splicing signals which define intron-exon boundaries. One such additional signal is provided by cis-acting regulatory enhancer and silencer sequences. Exonic splicing enhancers (ESE), exonic splicing silencers (ESS), intronic splicing enhancers (ISE) and intron splicing silencers (ISS) have been identified which either repress or enhance usage of splice donor sites or splice acceptor sites, depending on their site and mode of action (Yeo et al. 2004, *Proc. Natl. Acad. Sci. U.S.A.* 101(44):15700-15705). Binding of specific proteins (trans factors) to these regulatory sequences directs the splicing process, either promoting or inhibiting usage of particular splice sites and thus modulating the ratio of splicing products (Scamborova et al. 2004, *Mol. Cell. Biol.* 24(5):1855-1869; Hovhannisyan and Carstens, 2005, *Mol. Cell. Biol.* 25(1):250-263; Minovitsky et al. 2005, *Nucleic Acids Res.* 33(2):714-724).

4. Pharmaceutical Compositions

[0264] In certain embodiments, the present invention provides pharmaceutical compositions comprising one or more antisense compound. In certain embodiments, such pharmaceutical composition comprises a sterile saline solution and one or more antisense compound. In certain embodiments, such pharmaceutical composition consists of a sterile saline solution and one or more antisense compound.

[0265] In certain embodiments, antisense compounds may be admixed with pharmaceutically acceptable active and/or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions depend on a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

[0266] In certain embodiments antisense compounds, can be utilized in pharmaceutical compositions by combining such oligomeric compounds with a suitable pharmaceutically acceptable diluent or carrier. A pharmaceutically acceptable diluent includes phosphate-buffered saline (PBS). PBS is a diluent suitable for use in compositions to be delivered parenterally. Accordingly, in certain embodiments, employed in the methods described herein is a pharmaceutical composition comprising an antisense compound and a pharmaceutically acceptable diluent. In certain embodiments, the pharmaceutically acceptable diluent is PBS.

[0267] Pharmaceutical compositions comprising antisense compounds encompass any pharmaceutically acceptable salts, esters, or salts of such esters. In certain embodiments, pharmaceutical compositions comprising antisense com-

pounds comprise one or more oligonucleotide which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of antisense compounds, prodrugs, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. Suitable pharmaceutically acceptable salts include, but are not limited to, sodium and potassium salts.

[0268] A prodrug can include the incorporation of additional nucleosides at one or both ends of an oligomeric compound which are cleaved by endogenous nucleases within the body, to form the active antisense oligomeric compound.

[0269] Lipid-based vectors have been used in nucleic acid therapies in a variety of methods. For example, in one method, the nucleic acid is introduced into preformed liposomes or lipoplexes made of mixtures of cationic lipids and neutral lipids. In another method, DNA complexes with mono- or poly-cationic lipids are formed without the presence of a neutral lipid.

[0270] Certain preparations are described in Akinc et al., *Nature Biotechnology* 26, 561-569 (1 May 2008), which is herein incorporated by reference in its entirety.

5. Administration to a Subject

[0271] In certain embodiments, pharmaceutical compositions comprising one or more antisense compound are administered to a subject. In certain embodiments, such pharmaceutical compositions are administered by injection. In certain embodiments, such pharmaceutical compositions are administered by infusion.

[0272] In certain embodiments, pharmaceutical compositions are administered by injection or infusion into the CSF. In certain such embodiments, pharmaceutical compositions are administered by direct injection or infusion into the spine. In certain embodiments, pharmaceutical compositions are administered by injection or infusion into the brain. In certain embodiments, pharmaceutical compositions are administered by intrathecal injection or infusion rather than into the spinal cord tissue itself. Without being limited as to theory, in certain embodiments, the antisense compound released into the surrounding CSF and may penetrate into the spinal cord parenchyma. An additional advantage of intrathecal delivery is that the intrathecal route mimics lumbar puncture administration (i.e., spinal tap) already in routine use in humans.

[0273] In certain embodiments, pharmaceutical compositions are administered by intracerebroventricular (ICV) injection or infusion. Intracerebroventricular, or intraventricular, delivery of a pharmaceutical composition comprising one or more antisense compounds may be performed in any one or more of the brain's ventricles, which are filled with cerebrospinal fluid (CSF). CSF is a clear fluid that fills the ventricles, is present in the subarachnoid space, and surrounds the brain and spinal cord. CSF is produced by the choroid plexuses and via the weeping or transmission of tissue fluid by the brain into the ventricles. The choroid plexus is a structure lining the floor of the lateral ventricle and the roof of the third and fourth ventricles. Certain studies have indicated that these structures are capable of producing 400-600 ccs of fluid per day consistent with an amount to fill the central nervous system spaces four times in a day. In adult humans, the volume of this fluid has been

calculated to be from 125 to 150 ml (4-5 oz). The CSF is in continuous formation, circulation and absorption. Certain studies have indicated that approximately 430 to 450 ml (nearly 2 cups) of CSF may be produced every day. Certain calculations estimate that production equals approximately 0.35 ml per minute in adults and 0.15 per minute in infant humans. The choroid plexuses of the lateral ventricles produce the majority of CSF. It flows through the foramina of Monro into the third ventricle where it is added to by production from the third ventricle and continues down through the aqueduct of Sylvius to the fourth ventricle. The fourth ventricle adds more CSF; the fluid then travels into the subarachnoid space through the foramina of Magendie and Luschka. It then circulates throughout the base of the brain, down around the spinal cord and upward over the cerebral hemispheres. The CSF empties into the blood via the arachnoid villi and intracranial vascular sinuses.

[0274] In certain embodiments, such pharmaceutical compositions are administered systemically. In certain embodiments, pharmaceutical compositions are administered subcutaneously. In certain embodiments, pharmaceutical compositions are administered intravenously. In certain embodiments, pharmaceutical compositions are administered by intramuscular injection.

[0275] In certain embodiments, pharmaceutical compositions are administered both directly to the CSF (e.g., IT and/or ICV injection and/or infusion) and systemically.

[0276] In certain embodiments, an antisense compound administered systemically enters neurons. In certain embodiments, systemically administered antisense compounds may penetrate the blood-brain barrier, particularly in young subjects where the blood-brain barrier is not fully formed (e.g., in subjects in utero and/or in newborn subjects). In certain embodiments, some amount of systemically administered antisense compound may be taken up by nerve cells, even in subjects in which the blood-brain barrier is fully formed. For example, antisense compounds may enter a neuron at or near the neuromuscular junction (retrograde uptake). In certain embodiments, such retrograde uptake results in antisense activity inside the neuron, including, but not limited to, a motor neuron, and provides a therapeutic benefit by antisense activity inside the neuron.

[0277] In certain embodiments, systemic administration provides therapeutic benefit by antisense activity occurring in cells and/or tissues other than neurons. While evidence suggests that functional SMN inside neurons is required for normal neuron function, the consequence of reduced functional SMN in other cells and tissues is not well characterized. In certain embodiments, antisense activity in non-neuronal cells results in restoration of SMN function in those non-neuronal cells, which in turn results in therapeutic benefit.

[0278] In certain embodiments, improved SMN function in non-neuronal cells provides improved neuronal cell function, whether or not SMN function inside neurons is improved. For example, in certain embodiments, systemic administration of pharmaceutical compositions of the present invention results in antisense activity in muscle cells. Such antisense activity in muscle cells may provide a benefit to the motor-neurons associated with that muscle cell or to neurons generally. In such embodiments, the muscle cell having restored SMN function may provide a factor that improves neuronal viability and/or function. In certain embodiments, such antisense activity is independent of

benefit from antisense activity occurring from antisense compounds inside neurons. In certain embodiments, systemic administration of pharmaceutical compositions of the present invention results in antisense activity in other non-neuronal cells, including cells not in immediate association with neurons. Such antisense activity in non-neuronal cells may improve function of neurons. For example, antisense activity in a non-neuronal cell (e.g., liver cell) may result in that cell producing a factor that improves function of neurons. Note: since the term "antisense activity" includes direct and indirect activities, a benefit to neuronal function is an "antisense activity" even if no antisense compound enters the neuron.

[0279] In certain embodiments, systemic administration of a pharmaceutical composition results in therapeutic benefit independent of direct or indirect antisense activities in neurons. Typically, in the setting of SMA, neuronal function is diminished, resulting in significant symptoms. Additional symptoms may result from diminished SMN activity in other cells. Certain such symptoms may be masked by the relative severity of symptoms from diminished neuronal function. In certain embodiments, systemic administration results in restored or improved SMN function in non-neuronal cells. In certain such embodiments, such restored or improved SMN function in non-neuronal cells has therapeutic benefit. For example, in certain instances, subjects having SMA have reduced growth. Such reduced growth may not result from diminished function in neuronal cells. Indeed, reduced growth may be related to impaired function of cells in another organ, such as the pituitary gland, and/or may be the result of SMN deficiencies throughout the cells of the body. In such embodiments, systemic administration may result in improved SMN activity in pituitary cells and/or other cells, resulting in improved growth. In certain instances, administration to the CSF restores sufficient neuronal function to allow a subject to live longer, however one or more symptoms previously unknown because subjects typically died before such symptoms appeared emerges, because the subject lives longer. Certain such emergent symptoms may be lethal. In certain embodiments, emergent symptoms are treated by systemic administration. Regardless of mechanism, in certain embodiments, a variety of symptoms of SMA, including, but not limited to symptoms previously masked by more severe symptoms associated with impaired neuronal function, may be treated by systemic administration.

[0280] In certain embodiments, systemic administration of pharmaceutical compositions of the present invention result in increased SMN activity in muscle cells. In certain embodiments, such improved SMN activity in muscle cells provides therapeutic benefit. Improved SMN activity in muscle alone has been reported to be insufficient to provide therapeutic benefit (e.g., Gravrilina, et al., *Hum Mol Genet* 2008 17(8):1063-1075). In certain embodiments, the present invention provides methods that result improve SMN function in muscle and do provide therapeutic benefit. In certain instances, therapeutic benefit may be attributable to improved SMN function in other cells (alone or in combination with muscle cells). In certain embodiments, improved SMN function in muscle alone may provide benefit.

[0281] In certain embodiments, systemic administration results in improved survival.

6. Spinal Muscular Atrophy (SMA)

[0282] SMA is a genetic disorder characterized by degeneration of spinal motor neurons. SMA is caused by the homozygous loss of both functional copies of the SMN1 gene. However, the SMN2 gene has the potential to code for the same protein as SMN1 and thus overcome the genetic defect of SMA patients. SMN2 contains a translationally silent mutation (C→T) at position+6 of exon 7, which results in inefficient inclusion of exon 7 in SMN2 transcripts. Therefore, the predominant form of SMN2, one which lacks exon 7, is unstable and inactive. Thus, therapeutic compounds capable of modulating SMN2 splicing such that the percentage of SMN2 transcripts containing exon 7 is increased, would be useful for the treatment of SMA.

[0283] In certain embodiments, the present invention provides antisense compounds complementary to a pre-mRNA encoding SMN2. In certain such embodiments, the antisense compound alters splicing of SMN2. Certain sequences and regions useful for altering splicing of SMN2 may be found in PCT/US06/024469, which is hereby incorporated by reference in its entirety for any purpose. In certain embodiments, oligomeric compounds having any motif described herein have a nucleobase sequence complementary to intron 7 of SMN2. Certain such nucleobase sequences are exemplified in the non-limiting table below.

Sequence	Length	SEQ ID
TGCTGGCAGACTTAC	15	3
CATAATGCTGGCAGA	15	4
TCATAATGCTGGCAG	15	5
TTCATAATGCTGGCA	15	6
TTTCATAATGCTGGC	15	2
ATTCACITTCATAATGCTGG	20	7
TCACITTCATAATGCTGG	18	1
CTTTCATAATGCTGG	15	8
TCATAATGCTGG	12	9
ACTTTCATAATGCTG	15	10
TTCATAATGCTG	12	11
CACTTTCATAATGCT	15	12
TTTCATAATGCT	12	13
TCACITTCATAATGC	15	14
CTTTCATAATGC	12	15
TTCACITTCATAATG	15	16
ACTTTCATAATG	12	17
ATTCACITTCATAAT	15	18
CACTTTCATAAT	12	19
GATTCACITTCATAA	15	20
TCACITTCATAA	12	21
TTCACITTCATA	12	22

-continued

Sequence	Length	SEQ ID
ATTCACITTCAT	12	23
AGTAAGATTCACITTT	15	24

[0284] Antisense compounds of the present invention can be used to modulate the expression of SMN2 in a subject, such as a human. In certain embodiments, the subject has spinal muscular atrophy. In certain such subjects, the SMN1 gene is absent or otherwise fails to produce sufficient amounts of functional SMN protein. In certain embodiments, the antisense compounds of the present invention effectively modulate splicing of SMN2, resulting in an increase in exon 7 inclusion in SMN2 mRNA and ultimately in SMN2 protein that includes the amino acids corresponding to exon 7. Such alternate SMN2 protein resembles wild-type SMN protein. Antisense compounds of the present invention that effectively modulate expression of SMN2 mRNA or protein products of expression are considered active antisense compounds.

[0285] Modulation of expression of SMN2 can be measured in a bodily fluid, which may or may not contain cells; tissue; or organ of the animal. Methods of obtaining samples for analysis, such as body fluids (e.g., sputum, serum, CSF), tissues (e.g., biopsy), or organs, and methods of preparation of the samples to allow for analysis are well known to those skilled in the art. Methods for analysis of RNA and protein levels are discussed above and are well known to those skilled in the art. The effects of treatment can be assessed by measuring biomarkers associated with the target gene expression in the aforementioned fluids, tissues or organs, collected from an animal contacted with one or more compounds of the invention, by routine clinical methods known in the art.

[0286] Methods whereby bodily fluids, organs or tissues are contacted with an effective amount of one or more of the antisense compounds or compositions of the invention are also contemplated. Bodily fluids, organs or tissues can be contacted with one or more of the compounds of the invention resulting in modulation of SMN2 expression in the cells of bodily fluids, organs or tissues. An effective amount can be determined by monitoring the modulatory effect of the antisense compound or compounds or compositions on target nucleic acids or their products by methods routine to the skilled artisan.

[0287] The invention also provides an antisense compound as described herein, for use in any of the methods as described herein. For example, the invention provides an antisense compound comprising an antisense oligonucleotide complementary to a nucleic acid encoding human SMN2, for use in treating a disease or condition associated with survival motor neuron protein (SMN), such as spinal muscular atrophy (SMA). As a further example, the invention provides an antisense compound comprising an antisense oligonucleotide complementary to a nucleic acid encoding human SMN2, for use in treating a disease or condition associated with survival motor neuron protein (SMN) by administering the antisense compound directly into the central nervous system (CNS) or CSF.

[0288] The invention also provides the use of an antisense compound as described herein in the manufacture of a

medicament for use in any of the methods as described herein. For example, the invention provides the use of an antisense compound comprising an antisense oligonucleotide complementary to a nucleic acid encoding human SMN2 in the manufacture of a medicament for treating a disease or condition associated with survival motor neuron protein (SMN), such as spinal muscular atrophy (SMA). As a further example, the invention provides the use of an antisense compound comprising an antisense oligonucleotide complementary to a nucleic acid encoding human SMN2 in the manufacture of a medicament for treating a disease or condition associated with survival motor neuron protein (SMN) by administration of the medicament directly into the central nervous system (CNS) or CSF.

[0289] In certain embodiments, oligomeric compounds having any motif described herein have a nucleobase sequence complementary to exon 7 of SMN2.

[0290] In certain embodiments, oligomeric compounds having any motif described herein have a nucleobase sequence complementary to intron 6 of SMN2.

[0291] In certain embodiments, an antisense compound comprises an antisense oligonucleotide having a nucleobase sequence comprising at least 10 nucleobases of the sequence: TCACTTTCATAATGCTGG (SEQ ID NO: 1). In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 11 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 12 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 13 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 14 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 15 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 16 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising at least 17 nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence comprising the nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide has a nucleobase sequence consisting of the nucleobases of such sequence. In certain embodiments, an antisense oligonucleotide consists of 10-18 linked nucleosides and has a nucleobase sequence 100% identical to an equal-length portion of the sequence: TCACTTTCATAATGCTGG (SEQ ID NO: 1).

[0292] In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 1. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 2. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 3. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 4. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 5. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 6. In certain embodiments, a

compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 7. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 8. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 9. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 10. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 11. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 12. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 13. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 14. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 15. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 16. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 17. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 18. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 19. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 20. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 21. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 22. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 23. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 24. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 26. In certain embodiments, a compound comprises a modified oligonucleotide having a nucleobase sequence that consists of SEQ ID NO: 27.

[0293] A compound comprising Isis 449320. A compound comprising Isis 449323. A compound comprising Isis 605918. A compound comprising Isis 605919. A compound comprising Isis 663006. A compound comprising Isis 663007. A compound comprising Isis 663008. A compound comprising Isis 663009. A compound comprising Isis 663010. A compound comprising Isis 663011. A compound comprising Isis 663012. A compound comprising Isis 663013. A compound comprising Isis 663014. A compound comprising Isis 663015. A compound comprising Isis 663016. A compound comprising Isis 663017. A compound comprising Isis 663018. A compound comprising Isis 663019. A compound comprising Isis 669541. A compound comprising Isis 710554. A compound comprising Isis 710555. A compound comprising Isis 710556. A compound

comprising Isis 710557. A compound comprising Isis 710558. A compound comprising Isis 449321. A compound comprising Isis 449322. A compound comprising Isis 710543. A compound comprising Isis 710544. A compound comprising Isis 710548. A compound comprising Isis 710549. A compound comprising Isis 710550. A compound comprising Isis 710551. A compound comprising Isis 710552. A compound comprising Isis 710553. A compound comprising Isis 758133. A compound comprising Isis 758134. A compound comprising Isis 758135. A compound comprising Isis 758136. A compound comprising Isis 758137. A compound comprising Isis 758138. A compound comprising Isis 758139. A compound comprising Isis 758140. A compound comprising Isis 494323.

[0294] A compound consisting of Isis 449320. A compound consisting of Isis 449323. A compound consisting of Isis 605918. A compound consisting of Isis 605919. A compound consisting of Isis 663006. A compound consisting of Isis 663007. A compound consisting of Isis 663008. A compound consisting of Isis 663009. A compound consisting of Isis 663010. A compound consisting of Isis 663011. A compound consisting of Isis 663012. A compound consisting of Isis 663013. A compound consisting of Isis 663014. A compound consisting of Isis 663015. A compound consisting of Isis 663016. A compound consisting of Isis 663017. A compound consisting of Isis 663018. A compound consisting of Isis 663019. A compound consisting of Isis 669541. A compound consisting of Isis 710554. A compound consisting of Isis 710555. A compound consisting of Isis 710556. A compound consisting of Isis 710557. A compound consisting of Isis 710558. A compound consisting of Isis 449321. A compound consisting of Isis 449322. A compound consisting of Isis 710543. A compound consisting of Isis 710544. A compound consisting of Isis 710548. A compound consisting of Isis 710549. A compound consisting of Isis 710550. A compound consisting of Isis 710551. A compound consisting of Isis 710552. A compound consisting of Isis 710553. A compound consisting of Isis 758133. A compound consisting of Isis 758134. A compound consisting of Isis 758135. A compound consisting of Isis 758136. A compound consisting of Isis 758137. A compound consisting of Isis 758138. A compound consisting of Isis 758139. A compound consisting of Isis 758140. A compound consisting of Isis 494323.

7. Certain Subjects

[0295] In certain embodiments, a subject has one or more indicator of SMA. In certain embodiments, the subject has reduced electrical activity of one or more muscles. In certain embodiments, the subject has a mutant SMN1 gene. In certain embodiment, the subject's SMN1 gene is absent or incapable of producing functional SMN protein. In certain embodiments, the subject is diagnosed by a genetic test. In certain embodiments, the subject is identified by muscle biopsy. In certain embodiments, a subject is unable to sit upright. In certain embodiments, a subject is unable to stand or walk. In certain embodiments, a subject requires assistance to breathe and/or eat. In certain embodiment, a subject is identified by electrophysiological measurement of muscle and/or muscle biopsy.

[0296] In certain embodiments, the subject has SMA type I. In certain embodiments, the subject has SMA type II. In certain embodiments, the subject has SMA type III. In

certain embodiments, the subject is diagnosed as having SMA in utero. In certain embodiments, the subject is diagnosed as having SMA within one week after birth. In certain embodiments, the subject is diagnosed as having SMA within one month of birth. In certain embodiments, the subject is diagnosed as having SMA by 3 months of age. In certain embodiments, the subject is diagnosed as having SMA by 6 months of age. In certain embodiments, the subject is diagnosed as having SMA by 1 year of age. In certain embodiments, the subject is diagnosed as having SMA between 1 and 2 years of age. In certain embodiments, the subject is diagnosed as having SMA between 1 and 15 years of age. In certain embodiments, the subject is diagnosed as having SMA when the subject is older than 15 years of age.

[0297] In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered in utero. In certain such embodiments, the first dose is administered before complete development of the blood-brain-barrier. In certain embodiments, the first dose is administered to the subject in utero systemically. In certain embodiments, the first dose is administered in utero after formation of the blood-brain-barrier. In certain embodiments, the first dose is administered to the CSF.

[0298] In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than one week old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than one month old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than 3 months old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than 6 months old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than one year old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than 2 years old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is less than 15 years old. In certain embodiments, the first dose of a pharmaceutical composition according to the present invention is administered when the subject is older than 15 years old.

8. Certain Doses

[0299] In certain embodiments, the present invention provides dose amounts and frequencies. In certain embodiments, pharmaceutical compositions are administered as a bolus injection. In certain such embodiments, the dose of the bolus injection is from 0.01 to 25 milligrams of antisense compound per kilogram body weight of the subject. In certain such embodiments, the dose of the bolus injection is from 0.01 to 10 milligrams of antisense compound per kilogram body weight of the subject. In certain embodiments, the dose is from 0.05 to 5 milligrams of antisense compound per kilogram body weight of the subject. In certain embodiments, the dose is from 0.1 to 2 milligrams of antisense compound per kilogram body weight of the subject. In certain embodiments, the dose is from 0.5 to 1 milligrams of antisense compound per kilogram body

weight of the subject. In certain embodiments, such doses are administered twice monthly. In certain embodiments, such doses are administered every month. In certain embodiments, such doses are administered every 2 months. In certain embodiments, such doses are administered every 6 months. In certain embodiments, such doses are administered by bolus injection into the CSF. In certain embodiments, such doses are administered by intrathecal bolus injection. In certain embodiments, such doses are administered by bolus systemic injection (e.g., subcutaneous, intramuscular, or intravenous injection). In certain embodiments, subjects receive bolus injections into the CSF and bolus systemic injections. In such embodiments, the doses of the CSF bolus and the systemic bolus may be the same or different from one another. In certain embodiments, the CSF and systemic doses are administered at different frequencies. In certain embodiments, the invention provides a dosing regimen comprising at least one bolus intrathecal injection and at least one bolus subcutaneous injection.

[0300] In certain embodiments, pharmaceutical compositions are administered by continuous infusion. Such continuous infusion may be accomplished by an infusion pump that delivers pharmaceutical compositions to the CSF. In certain embodiments, such infusion pump delivers pharmaceutical composition IT or ICV. In certain such embodiments, the dose administered is between 0.05 and 25 milligrams of antisense compound per kilogram body weight of the subject per day. In certain embodiments, the dose administered is from 0.1 to 10 milligrams of antisense compound per kilogram body weight of the subject per day. In certain embodiments, the dose administered is from 0.5 to 10 milligrams of antisense compound per kilogram body weight of the subject per day. In certain embodiments, the dose administered is from 0.5 to 5 milligrams of antisense compound per kilogram body weight of the subject per day. In certain embodiments, the invention provides a dosing regimen comprising infusion into the CNS and at least one bolus systemic injection. In certain embodiments, the invention provides a dosing regimen comprising infusion into the CNS and at least one bolus subcutaneous injection. In certain embodiments, the dose, whether by bolus or infusion, is adjusted to achieve or maintain a concentration of antisense compound from 0.1 to 100 microgram per gram of CNS tissue. In certain embodiments, the dose, whether by bolus or infusion, is adjusted to achieve or maintain a concentration of antisense compound from 1 to 10 microgram per gram of CNS tissue. In certain embodiments, the dose, whether by bolus or infusion, is adjusted to achieve or maintain a concentration of antisense compound from 0.1 to 1 microgram per gram of CNS tissue.

[0301] In certain embodiments, dosing a subject is divided into an induction phase and a maintenance phase. In certain such embodiments, the dose administered during the induction phase is greater than the dose administered during the maintenance phase. In certain embodiments, the dose administered during the induction phase is less than the dose administered during the maintenance phase. In certain embodiments, the induction phase is achieved by bolus injection and the maintenance phase is achieved by continuous infusion.

[0302] In certain embodiments, the invention provides systemic administration of antisense compounds, either

alone or in combination with delivery into the CSF. In certain embodiments, the dose for systemic administration is from 0.1 mg/kg to 200 mg/kg. In certain embodiments, the dose for systemic administration is from 0.1 mg/kg to 100 mg/kg. In certain embodiments, the dose for systemic administration is from 0.5 mg/kg to 100 mg/kg. In certain embodiments, the dose for systemic administration is from 1 mg/kg to 100 mg/kg. In certain embodiments, the dose for systemic administration is from 1 mg/kg to 50 mg/kg. In certain embodiments, the dose for systemic administration is from 1 mg/kg to 25 mg/kg. In certain embodiments, the dose for systemic administration is from 0.1 mg/kg to 25 mg/kg. In certain embodiments, the dose for systemic administration is from 0.1 mg/kg to 10 mg/kg. In certain embodiments, the dose for systemic administration is from 1 mg/kg to 10 mg/kg. In certain embodiments, the dose for systemic administration is from 1 mg/kg to 5 mg/kg. In certain embodiments comprising both systemic and CSF delivery, the doses for those two routes are independently determined.

[0303] a. Calculation of Appropriate Human Doses

[0304] In certain embodiments, the subject is a human. In certain embodiments, a human dose is calculated or estimated from data from animal experiments, such as those described herein. In certain embodiments, a human dose is calculated or estimated from data from monkey and/or mouse experiments, such as those described herein. In certain embodiments, a human dose is calculated or estimated from data from mouse experiments, such as those described herein. In certain embodiments, appropriate human doses can be calculated using pharmacokinetic data from mouse along with knowledge of brain weight and/or cerebrospinal fluid (CSF) turnover rates. For example, the mouse brain weight is approximately 0.4 g, which is approximately 2% of its body weight. In humans, the average brain weight is 1.5 kg which is approximately 2.5% of body weight. In certain embodiments, administration into the CSF results in elimination of a portion of the compound through uptake in brain tissue and subsequent metabolism. By using the ratio of human to mouse brain weight as a scaling factor an estimate of the elimination and clearance through the brain tissue can be calculated. Additionally, the CSF turnover rate can be used to estimate elimination of compound from the CSF to blood. Mouse CSF turnover rate is approximately 10-12 times per day (0.04 mL produced at 0.325 μ L/min). Human CSF turnover rate is approximately 4 times per day (100-160 mL produced at 350-400 μ L/min). Clearance, and therefore dosing requirements, can be based on brain weight elimination scaling, and/or the CSF turnover scaling. The extrapolated human CSF clearance can be used to estimate equivalent doses in humans that approximate doses in mice. In this way, human doses can be estimated that account for differences in tissue metabolism based on brain weight and CSF turnover rates. Such methods of calculation and estimate are known to those skilled in the art.

[0305] By way of non-limiting example, in certain embodiments, an equivalent human dose can be estimated from a desired mouse dose by multiplying the mg/kg mouse dose by a factor from about 0.25 to about 1.25 depending on the determined clearance and elimination of a particular compound. Thus, for example, in certain embodiments, a human dose equivalent of a 0.01 mg dose for a 20 g mouse will range from about 8.75 mg to about 43.75 mg total dose for a 70 kg human. Likewise, in certain embodiments, a human dose equivalent of a 0.01 mg dose for a 4 g newborn

mouse will range from about 1.9 mg to about 9.4 mg total dose for a 3 kg newborn human. These example doses are merely to illustrate how one of skill may determine an appropriate human dose and are not intended to limit the present invention.

[0306] In certain embodiments, a human dose for systemic delivery (whether administered alone or in combination with CSF delivery) is calculated or estimated from data from animal experiments, such as those described herein. Typically, an appropriate human dose (in mg/kg) for systemic dose is between 0.1 and 10 times an effective dose in animals. Thus, solely for example, a subcutaneous dose of 50 µg in a 2 g newborn mouse is a dose of 25 mg/kg. The corresponding dose for a human is predicted to be between 2.5 mg/kg and 250 mg/kg. For a 3 kilogram infant, the corresponding dose is between 7.5 mg and 750 mg. For a 25 kg child, the corresponding dose is from 62.5 mg to 6250 mg.

9. Treatment Regimens

[0307] In certain embodiments, the above dose amounts, dose frequencies, routes of administration, induction and maintenance phases, and timing of first dose are combined to provide dosing regimens for subjects having SMA. Such

embodiments, administration of pharmaceutical compositions, particularly by continuous infusion, presents particular challenges. Accordingly, in certain embodiments, the present invention provides for administration of pharmaceutical compositions by bolus administration while the subject is in utero or very young, followed by continuous infusion via an implanted infusion pump when the subject is older and placement of such pump is more practical. Further, in certain embodiments, as a subject grows, the absolute dose is increased to achieve the same or similar dose:body-weight ratio.

[0308] In certain embodiments, pharmaceutical compositions of the present disclosure have improved therapeutic indices. In certain embodiments, pharmaceutical compositions of the present disclosure having improved therapeutic indices allows one having skill in the art to administer the pharmaceutical compositions of the present disclosure more infrequently. In certain embodiments, pharmaceutical compositions of the present disclosure having improved therapeutic indices allows one having skill in the art to allow longer intervals between the first dose and the second dose.

[0309] The following table is intended to exemplify treatment regimens and is not intended to limit the possible combinations of treatments which will be easily accomplished by one of skill in the art.

Dosing period	First	Second	Third	Fourth	Fifth
Regimen 1					
Subject Age	In utero, prior to formation of blood-brain-barrier	In utero, after formation of blood-brain-barrier	>1 week	6 months	1.5 years
Dose Amount	50 µg	50 µg	100 µg	10 µg/day	50 µg/day
Frequency	Single admin	Single admin	Monthly	Continuous	Continuous
Route of Administration	Systemic injection	IT injection	IT injections	IT infusion	IT infusion
Duration	N/A	N/A	6 months	1 year	Ongoing
Regimen 2					
Subject Age	In utero, after formation of blood-brain-barrier	>1 week	6 months	1.5 years	N/A
Dose Amount	50 µg	100 µg	5 mg/day	10 mg/day	N/A
Frequency	Single admin	Monthly	Continuous	Continuous	N/A
Route of Administration	ICV injection	ICV injection	ICV infusion	ICV infusion	N/A
Duration	N/A	6 months	1 year	Ongoing	N/A
Regimen 3					
Subject Age	>1 week	6 months	1.5 years	2.5 years*	
Dose Amount	100 µg	500 µg/day	20 mg/day	20 mg/day	100 mg
Frequency	2xMonthly	Continuous	Continuous	Continuous	2xMonthly
Route of Administration	ICV injection	ICV infusion	ICV infusion	ICV infusion	IP
Duration	6 months	1 year	1 year	Ongoing	Ongoing

*Note:
the 4th dosing period in regimen 3 exemplifies continuous CSF infusion combined with periodic systemic administration. These treatment regimens are intended to exemplify and not to limit the present invention.

dosing regimens may be selected and adjusted to provide amelioration of one or more symptom of SMA and/or to reduce or avoid toxicity or side effects attributable to the administration of the pharmaceutical composition. In certain embodiments, subjects are in utero or newborn. In such

[0310] In certain embodiments, the dosing regimen comprises a systemic administration, either alone or in combination with administration into the CSF (for example regimen 3, above). The table, below further exemplifies such regimens.

Systemic administration			CSF administration		
Dose	Route	Frequency	Dose	Route	Frequency
1-5 mg/kg	subcutaneous	weekly	5-10 mg/kg	bolus IT	monthly
1-5 mg/kg	subcutaneous	monthly	1-5 mg/kg	bolus ICV	2 months
10-50 mg/kg	subcutaneous	monthly	0.5-1 mg/kg	bolus IT	6 months
0.5-25 mg/kg	subcutaneous	monthly	10 mg/kg/day	IT infusion	continuous for 7 days every 6 months
0.1-10 mg/kg	subcutaneous	monthly		none	
	none		0.5-1 mg/kg	bolus IT	6 months

These treatment regimens are intended to exemplify and not to limit the present invention. One of skill in the art will be able to select an appropriate combination of the doses and deliveries in view of the present disclosure and based on a variety of factors, such as the severity of the condition and the overall health and age of the subject.

10. Co-Administration

[0311] In certain embodiments, pharmaceutical compositions of the present invention are co-administered with at least one other pharmaceutical composition for treating SMA and/or for treating one or more symptom associated with SMA. In certain embodiments, such other pharmaceutical composition is selected from trichostatin-A, valproic acid, riluzole, hydroxyurea, and a butyrate or butyrate derivative. In certain embodiments, pharmaceutical compositions of the present invention are co-administered with trichostatin A. In certain embodiments, pharmaceutical compositions of the present invention are co-administered with a derivative of quinazoline, for example as described in Thurmond, et al., *J. Med Chem.* 2008, 51, 449-469. In certain embodiments, a pharmaceutical composition of the present invention and at least one other pharmaceutical composition are co-administered at the same time. In certain embodiments, a pharmaceutical composition of the present invention and at least one other pharmaceutical composition are co-administered at different times.

[0312] In certain embodiments, pharmaceutical compositions of the present invention are co-administered with a gene therapy agent. In certain such embodiments, the gene therapy agent is administered to the CSF and the pharmaceutical composition of the present invention is administered systemically. In certain such embodiments, the gene therapy agent is administered to the CSF and the pharmaceutical composition of the present invention is administered to the CSF and systemically. In certain embodiments, a pharmaceutical composition of the present invention and a gene therapy agent are co-administered at the same time. In certain embodiments, a pharmaceutical composition of the present invention and a gene therapy agent are co-administered at different times. Certain gene therapy approaches to SMA treatment have been reported (e.g., Coady et al., *PLoS ONE* 2008 3(10): e3468; Passini et al., *J Clin Invest* 2010 Apr. 1, 120(4): 1253-64).

[0313] In certain embodiments, pharmaceutical compositions of the present invention are co-administered with at least one other therapy for SMA. In certain embodiments, such other therapy for SMA is surgery. In certain embodiments, such other therapy is physical therapy, including, but not limited to exercises designed to strengthen muscles necessary for breathing, such as cough therapy. In certain

embodiments, other therapy is a physical intervention, such as a feeding tube or device for assisted breathing.

[0314] In certain embodiments, pharmaceutical compositions of the present invention are co-administered with one or more other pharmaceutical compositions that reduce an undesired side-effect of the pharmaceutical compositions of the present invention.

11. Phenotypic Effects

[0315] In certain embodiments, administration of at least one pharmaceutical composition of the present invention results in a phenotypic change in the subject. In certain embodiments, such phenotypic changes include, but are not limited to: increased absolute amount of SMN mRNA that includes exon 7; increase in the ratio SMN mRNA that includes exon 7 to SMN mRNA lacking exon 7; increased absolute amount of SMN protein that includes exon 7; increase in the ratio SMN protein that includes exon 7 to SMN protein lacking exon 7; improved muscle strength, improved electrical activity in at least one muscle; improved respiration; weight gain; and survival. In certain embodiments, at least one phenotypic change is detected in a motoneuron of the subject. In certain embodiments, administration of at least one pharmaceutical composition of the present invention results in a subject being able to sit-up, to stand, and/or to walk. In certain embodiments, administration of at least one pharmaceutical composition of the present invention results in a subject being able to eat, drink, and/or breathe without assistance. In certain embodiments, efficacy of treatment is assessed by electrophysiological assessment of muscle. In certain embodiments, administration of a pharmaceutical composition of the present invention improves at least one symptom of SMA and has little or no inflammatory effect. In certain such embodiment, absence of inflammatory effect is determined by the absence of significant increase in Aif1 levels upon treatment.

[0316] In certain embodiments, administration of at least one pharmaceutical composition of the present invention delays the onset of at least one symptom of SMA. In certain embodiments, administration of at least one pharmaceutical composition of the present invention slows the progression of at least one symptom of SMA. In certain embodiments, administration of at least one pharmaceutical composition of the present invention reduces the severity of at least one symptom of SMA.

[0317] In certain embodiments, administration of at least one pharmaceutical composition of the present invention results in an undesired side-effect. In certain embodiments, a treatment regimen is identified that results in desired amelioration of symptoms while avoiding undesired side-effects.

12. Dosage Units

[0318] In certain embodiments pharmaceutical compositions of the present invention are prepared as dosage units for administration. Certain such dosage units are at concentrations selected from 0.01 mg to 100 mg. In certain such embodiments, a pharmaceutical composition of the present invention comprises a dose of antisense compound selected from 0.01 mg, 0.1 mg, 0.5 mg, 1 mg, 5 mg, 10 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, and 200 mg. In certain embodiments, a pharmaceutical composition is comprises a dose of oligonucleotide selected from 0.1 mg, 0.5 mg, 1 mg, 5 mg, 10 mg, 25 mg, and 50 mg.

13. Kits

[0319] In certain embodiments, the present invention provides kits comprising at least one pharmaceutical composition. In certain embodiments, such kits further comprise a means of delivery, for example a syringe or infusion pump.

Nonlimiting Disclosure and Incorporation by Reference

[0320] While certain compounds, compositions and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds described herein and are not intended to limit the same. Each of the references, GenBank accession numbers, and the like recited herein is hereby incorporated by reference in its entirety.

[0321] Although the sequence listing accompanying this filing identifies each sequence as either "RNA" or "DNA" as required, in reality, those sequences may be modified with any combination of chemical modifications. One of skill in the art will readily appreciate that such designation as "RNA" or "DNA" to describe modified oligonucleotides is, in certain instances, arbitrary. For example, an oligonucleotide comprising a nucleoside comprising a 2'-OH sugar moiety and a thymine base could be described as a DNA having a modified sugar (2'-OH for the natural 2'-H of DNA) or as an RNA having a modified base (thymine (methylated uracil) for natural uracil of RNA).

[0322] Accordingly, nucleic acid sequences provided herein, including, but not limited to those in the sequence listing, are intended to encompass nucleic acids containing any combination of natural or modified RNA and/or DNA, including, but not limited to such nucleic acids having modified nucleobases. By way of further example and without limitation, an oligomeric compound having the nucleobase sequence "ATCGATCG" encompasses any oligomeric compounds having such nucleobase sequence, whether modified or unmodified, including, but not limited to, such compounds comprising RNA bases, such as those having sequence "AUCGAUCG" and those having some DNA bases and some RNA bases such as "AUCGATCG" and oligomeric compounds having other modified bases, such as "AT^{me}CGAUCG," wherein ^{me}C indicates a cytosine base comprising a methyl group at the 5-position.

EXAMPLES

[0323] The following examples illustrate certain embodiments of the present disclosure and are not limiting. Moreover, where specific embodiments are provided, the inventors have contemplated generic application of those specific embodiments. For example, disclosure of an oligonucleotide

having a particular motif provides reasonable support for additional oligonucleotides having the same or similar motif. And, for example, where a particular high-affinity modification appears at a particular position, other high-affinity modifications at the same position are considered suitable, unless otherwise indicated.

Example 1

Effect of Antisense Oligonucleotides Targeting Human SMN2 in an SMA Type III Mice Model

[0324] Taiwan strain of SMA type III mice were obtained from The Jackson Laboratory (Bar Harbor, Me.). These mice lack mouse SMN and are homozygous for human SMN2 (mSMN -/-; hSMN2+/+). These mice have been described in Hsieh-Li H M, et al., *Nature Genet.* 24, 66-70 2000. Antisense oligonucleotides targeting human SMN2 were designed and tested in these mice.

[0325] The newly designed antisense oligonucleotides in the Table below were designed as uniform 2'-MOE oligonucleotides with mixed backbone chemistry. Each antisense oligonucleotide is 18 nucleosides in length and each nucleoside has a 2'-MOE sugar modification. The internucleoside linkages throughout each gapmer are either phosphodiester or phosphorothioate linkages. The internucleoside linkages of each oligonucleotide is denoted in the Backbone Chemistry column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. Each antisense oligonucleotide listed in the Table below is targeted to the human SMN2 genomic sequence, designated herein as SEQ ID NO: 25 (GENBANK Accession No. NT_006713.14 truncated from nucleotides 19939708 to Ser. No. 19/967,777) at the target region 27062-27079.

TABLE 1

Modified oligonucleotides targeting human SMN2 with mixed backbone chemistry			
ISIS No	Sequence	Backbone Chemistry	SEQ ID NO
449320	TCACCTTCATAATGCTGG	ssooooooooooooooss	1
449323	TCACCTTCATAATGCTGG	sosososososososos	1
605918	TCACCTTCATAATGCTGG	ssssososososososs	1
605919	TCACCTTCATAATGCTGG	soooooossssssoooss	1

Treatment

[0326] Mice were administered 500 µg of ISIS oligonucleotide by intracerebroventricular (ICV) bolus injection. Control mice were administered PBS alone (dose of 0).

Tolerability Assay

[0327] At 3 hours post injection, each mouse was evaluated according to 7 different criteria. The 7 criteria are (1) the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse shows any movement without stimuli (4) the mouse demonstrates forward movement after its lifted; (5) the mouse

demonstrates any movement after its lifted; (6) the mouse responds to a tail pinch; (7) regular breathing. For each of the 7 different criteria, each mouse was given a sub-score of 0 if it met the criteria or 1 if it did not. After all of the 7 criteria were evaluated, the sub-scores were summed for each mouse and then averaged for each group. For example, if a mouse was bright, alert, and responsive 3 hours after the ICV dose, and met all other criteria, it would get a summed score of 0. If another mouse was not bright, alert, and responsive 3 hours after the ICV dose but met all other criteria, it would receive a score of 1. Saline treated mice generally receive a score of 0. As presented in the Table below, the ISIS oligonucleotides were deemed tolerable in the transgenic mice.

TABLE 2

Scoring for tolerability to ISIS oligonucleotides in SMA Type III mice	
ISIS No	3 hr Score
449320	0
449323	0
605918	3
605919	0

Exon 7 Splicing

[0328] Animals were sacrificed 2 weeks after the injection and brain and lumbar sections of the spinal cords were collected from each animal. Real time PCR was performed on each sample to determine the amount of human SMN2 message including exon 7 (exon 7⁺) and the amount of human SMN2 message lacking exon 7(exon 7⁻). Expression levels for exon 7⁺ and exon 7⁻ were normalized to total SMN2 levels. The data presented in the Table below are expressed as exon 7⁺ and exon 7⁻ levels divided by total SMN levels, and were further normalized to the levels obtained in the PBS control group (designated as 1).

[0329] Administration of ISIS oligonucleotides targeting SEQ ID NO: 25 at start site 27062, but with different backbone chemistries, all resulted in a striking increase in inclusion of exon 7.

TABLE 3

	Effect of ISIS oligonucleotides targeting SMN2 on splicing in SMA Type III Mice			
	Brain		Lumbar Cord	
	exon 7 ⁺	exon 7 ⁻	exon 7 ⁺	exon 7 ⁻
PBS	1.0	1.0	1.0	1.0
449320	2.4	0.60	2.7	0.66
449323	4.3	0.24	4.5	0.33
605918	4.4	0.23	4.5	0.28
605919	4.2	0.26	4.0	0.38

Example 2

Effect of Antisense Oligonucleotides Targeting Human SMN2 in SMA Type III Mice and WT Mice

[0330] Additional antisense oligonucleotides targeting human SMN2 were designed and tested in SMA Type III mice or a WT (C57BL/6) strain.

[0331] The newly designed antisense oligonucleotides in the Table below were designed as uniform 2'-MOE oligonucleotides with mixed backbone chemistry. Each antisense oligonucleotide is 18 nucleosides in length wherein each nucleoside has a 2'-MOE sugar modification. The internucleoside linkages throughout each gapmer are either phosphodiester or phosphorothioate linkages. The internucleoside linkages of each oligonucleotide is denoted in the Backbone Chemistry column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. Each antisense oligonucleotide listed in the Table below is targeted to the human SMN2 genomic sequence, designated herein as SEQ ID NO: 25 (GENBANK Accession No. NT_006713.14 truncated from nucleotides 19939708 to Ser. No. 19/967,777) at the target region 27062-27079.

TABLE 4

Modified oligonucleotides targeting human SMN2 with mixed backbone chemistry			
ISIS No	Sequence	Backbone chemistry	SEQ ID NO
605918	TCACTTTCATAATGCTGG	ssssososososososs	1
605919	TCACTTTCATAATGCTGG	sooooooooooooooss	1
663006	TCACTTTCATAATGCTGG	ssosososososososs	1
663007	TCACTTTCATAATGCTGG	ssosososososoooooss	1
663008	TCACTTTCATAATGCTGG	soossoossoossooss	1
663009	TCACTTTCATAATGCTGG	sooooooooooooooss	1
663010	TCACTTTCATAATGCTGG	sooooooooooooooss	1
663011	TCACTTTCATAATGCTGG	sooooooooooooooss	1
663012	TCACTTTCATAATGCTGG	sooooooooooooooss	1
663013	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663014	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663015	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663016	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663017	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663018	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
663019	TCACTTTCATAATGCTGG	sssooooooooooooooss	1
669541	TCACTTTCATAATGCTGG	sooooooooooooooss	1

Treatment

[0332] SMA Type III mice were administered 35 µg of ISIS oligonucleotide by intracerebroventricular (ICV) bolus injection. WT mice were administered 700 µg of ISIS oligonucleotide by intracerebroventricular (ICV) bolus injection. Control mice in each strain were administered PBS alone (dose of 0).

Tolerability Assay

[0333] At 3 hours post injection, each mouse was evaluated according to 7 different criteria. The 7 criteria are (1)

the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse shows any movement without stimuli (4) the mouse demonstrates forward movement after its lifted; (5) the mouse demonstrates any movement after its lifted; (6) the mouse responds to a tail pinch; (7) regular breathing. For each of the 7 different criteria, each mouse was given a sub-score of 0 if it met the criteria or 1 if it did not. After all of the 7 criteria were evaluated, the sub-scores were summed for each mouse and then averaged for each group. For example, if a mouse was bright, alert, and responsive 3 hours after the ICV dose, and met all other criteria, it would get a summed score of 0. If another mouse was not bright, alert, and responsive 3 hours after ICV dose but met all other criteria, it would receive a score of 1. Saline treated mice generally receive a score of 0. As presented in the Table below, the ISIS oligonucleotides were deemed tolerable in the mice.

TABLE 5

Scoring for tolerability to ISIS oligonucleotides in SMA Type III mice dosed with 35 μ g of ISIS Oligonucleotide	
ISIS No	3 hr Score
605918	0
605919	0
663006	0
663007	0
663008	0
663009	0
663010	0
663011	0
663012	0
663013	0
663014	0
663015	0
663016	0
663017	0
663018	0
663019	0
669541	0

TABLE 6

Scoring for tolerability to ISIS oligonucleotides in WT mice dosed with 700 μ g of ISIS Oligonucleotide	
ISIS No	3 hr Score
605918	1
605919	3.3
663006	0
663007	0
663008	0
663009	0
663010	1.5
663011	0
663012	0
663013	0
663014	0
663015	0
663016	0.5
663017	2
663018	3
663019	2.75
669541	3

Exon 7 Splicing

[0334] Animals were sacrificed 2 weeks after the injection and brain and lumbar sections of the spinal cords were

collected from each animal. Real time PCR was performed on each sample to determine the amount of human SMN2 message including exon 7 (exon 7⁺) and the amount of human SMN2 message lacking exon 7 (exon 7⁻). Expression levels for exon 7⁺ and exon 7⁻ were normalized to total SMN2 levels. The data presented in the Table below exon 7⁺ and exon 7⁻ levels divided by total SMN levels, and were further normalized to the levels obtained in the PBS control group (designated as 1).

[0335] Administration of many of the ISIS oligonucleotides targeting SEQ ID NO: 25 around target region 27062-27079, but with different backbone chemistries, resulted in an increase in inclusion of exon 7.

TABLE 7

	Effect of ISIS oligonucleotides targeting SMN2 on splicing in SMA Type III Mice			
	Brain		Lumbar Cord	
	exon 7 ⁺	exon 7 ⁻	exon 7 ⁺	exon 7 ⁻
PBS	1.0	1.0	1.0	1.0
605918	2.0	0.6	1.9	0.6
605919	1.6	0.7	1.3	0.8
663006	1.5	0.8	1.6	0.8
663007	1.2	0.8	1.2	0.9
663008	1.4	0.8	1.3	0.9
663009	1.3	0.8	1.4	0.7
663010	1.0	0.8	1.2	0.8
663011	1.2	0.7	1.1	0.8
663012	1.3	0.8	1.1	0.8
663013	1.1	0.9	1.0	0.8
663014	1.4	0.7	1.1	0.8
663015	1.3	0.7	1.1	0.8
663016	1.2	0.7	1.2	0.7
663017	1.5	0.8	1.3	0.7
663018	1.6	0.6	1.3	0.7
663019	1.4	0.7	1.4	0.7
669541	1.9	0.5	2.0	0.5

Example 3

Design of Antisense Oligonucleotides Targeting Human SMN2

[0336] Additional antisense oligonucleotides targeting human SMN2 were designed targeting the human SMN2 genomic sequence, designated herein as SEQ ID NO: 25 (GENBANK Accession No. NT_006713.14 truncated from nucleotides 19939708 to Ser. No. 19/967,777) around target region 27062-27079.

[0337] The newly designed antisense oligonucleotides in the Table below were designed as uniform 2-MOE oligonucleotides with mixed backbone chemistry. Each antisense oligonucleotide is 15, 18, or 20 nucleosides in length wherein each nucleoside has a 2'-MOE sugar modification. The internucleoside linkages throughout each gapmer are either phosphodiester or phosphorothioate linkages. The internucleoside linkages of each oligonucleotide is denoted in the Backbone Chemistry column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. "Start site" indicates the 5'-most nucleoside to which the gapmer is targeted in the human gene sequence. "Stop site" indicates the 3'-most nucleoside to which the gapmer is targeted human gene sequence.

TABLE 8

Modified oligonucleotides targeting human SMN2 with mixed backbone chemistry						
OligoID	Start Site	Stop Site	OligoSeq	Backbone Chemistry	SEQ ID NO	
710554	27061	27075	TTTCATAATGCTGGC	soososososososs	2	
710555	27061	27075	TTTCATAATGCTGGC	ssososososososs	2	
710556	27061	27075	TTTCATAATGCTGGC	ssososososososs	2	
710557	27061	27075	TTTCATAATGCTGGC	ssososososososs	2	
710558	27061	27075	TTTCATAATGCTGGC	ssososososososs	2	
449321	27062	27079	TCACTTTCATAATGCTGG	ssooooosooooos	1	
449322	27062	27079	TCACTTTCATAATGCTGG	ssooooosooooos	1	
710543	27062	27079	TCACTTTCATAATGCTGG	ssososososososs	1	
710544	27062	27079	TCACTTTCATAATGCTGG	ssososososososs	1	
710548	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	
710549	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	
710550	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	
710551	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	
710552	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	
710553	27062	27081	ATTCACCTTTCATAATGCTGG	soosososososososs	7	

Example 4

Design of Antisense Oligonucleotides Targeting Human SMN2

[0338] Additional antisense oligonucleotides targeting human SMN2 were designed targeting the human SMN2 genomic sequence, designated herein as SEQ ID NO: 25.

[0339] The newly designed antisense oligonucleotides in the Table below were designed as uniform 2-MOE oligonucleotides with mixed backbone chemistry. Each antisense oligonucleotide is 18 nucleosides in length wherein each

nucleoside has a 2'-MOE sugar modification. The internucleoside linkages throughout each gapmer are either phosphodiester or phosphorothioate linkages. The internucleoside linkages of each oligonucleotide is denoted in the Backbone Chemistry column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. "Start site" indicates the 5'-most nucleoside to which the gapmer is targeted in the human gene sequence. "Stop site" indicates the 3'-most nucleoside to which the gapmer is targeted human gene sequence.

TABLE 9

Modified oligonucleotides targeting human SMN2 with mixed backbone chemistry						
OligoID	Start Site	Stop Site	OligoSeq	Backbone Chemistry	SEQ ID NO	
758133	27060	27077	ACTTTCATAATGCTGGCA	ssososososososs	26	
758134	27060	27077	ACTTTCATAATGCTGGCA	ssososososososs	26	
758135	27060	27077	ACTTTCATAATGCTGGCA	ssososososososs	26	
758136	27060	27077	ACTTTCATAATGCTGGCA	ssososososososs	26	
758137	27061	27078	CACTTTCATAATGCTGGC	SSOSSOSSOSSOSS	27	
758138	27061	27078	CACTTTCATAATGCTGGC	SSOSSOSSOSSOSS	27	
758139	27061	27078	CACTTTCATAATGCTGGC	SSOSSOSSOSSOSS	27	
758140	27061	27078	CACTTTCATAATGCTGGC	SSOSSOSSOSSOSS	27	

Example 5

Effect of Antisense Oligonucleotides Targeting Human SMN2 in an SMA Type III Mice Model

[0340] Groups of 3 to 4 wild type and SMN transgenic mice were treated and analyzed as described in Example 2. The antisense oligonucleotides that were administered to the mice are listed in the tables below. The results are presented, as described in Example 2, in the tables below.

TABLE 10

Scoring for tolerability of ISIS oligonucleotides in WT mice dosed with 700 µg of ISIS Oligonucleotide	
ISIS No	3 hr Score
494323	0
710543	2
710544	1
710548	0
710549	0
710550	1
710551	0
710552	1
710553	0
710554	0
710555	0
710556	0
710557	0
710558	0
758133	0
758134	0
758135	0
758136	2
758137	0
758138	0
758139	0
758140	0

TABLE 11

Scoring for tolerability of ISIS oligonucleotides in SMA Type III mice dosed with 35 µg of ISIS Oligonucleotide	
ISIS No	3 hr Score
494323	0
710543	0
710544	0
710548	0
710549	0
710550	0
710551	0
710552	0
710553	0

TABLE 12

Effect of 35 µg of ISIS Oligonucleotide targeting SMN2 on splicing in SMA Type III Mice				
Isis No.	Brain		Lumbar Cord	
	exon 7 ⁺	exon 7 ⁻	exon 7 ⁺	exon 7 ⁻
PBS	1.0	1.0	1.0	1.0
494323	2.6	0.9	2.0	0.8
710543	2.7	0.7	2.1	0.7
710544	2.5	0.9	1.9	0.7
710548	2.1	1.0	1.4	0.8
710549	2.4	0.8	1.7	0.7
710550	2.4	0.9	1.9	0.7
710551	2.4	0.8	2.1	0.7
710552	2.4	0.8	1.9	0.7
710553	3.1	0.8	2.0	0.7

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 27

<210> SEQ ID NO 1
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 1

tcactttcat aatgctgg 18

<210> SEQ ID NO 2
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 2

tttcataatg ctggc 15

<210> SEQ ID NO 3
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 3

tgctggcaga cttac 15

<210> SEQ ID NO 4
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 4

cataatgctg gcaga 15

<210> SEQ ID NO 5
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 5

tcataatgct ggcag 15

<210> SEQ ID NO 6
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 6

ttcataatgc tggca 15

<210> SEQ ID NO 7
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 7

attcactttc ataatgctgg 20

<210> SEQ ID NO 8
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 8

ctttcataat gctgg 15

<210> SEQ ID NO 9
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

-continued

tcataatgct gg 12

<210> SEQ ID NO 10
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

actttcataa tgctg 15

<210> SEQ ID NO 11
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 11

ttcataatgc tg 12

<210> SEQ ID NO 12
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 12

cactttcata atgct 15

<210> SEQ ID NO 13
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 13

tttcataatg ct 12

<210> SEQ ID NO 14
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 14

tcactttcat aatgc 15

<210> SEQ ID NO 15
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 15

ctttcataat gc 12

-continued

<210> SEQ ID NO 16
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 16

ttcactttca taatg 15

<210> SEQ ID NO 17
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 17

actttcataa tg 12

<210> SEQ ID NO 18
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 18

attcactttc ataata 15

<210> SEQ ID NO 19
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 19

cactttcata at 12

<210> SEQ ID NO 20
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 20

gattcacttt cataa 15

<210> SEQ ID NO 21
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 21

tcactttcat aa 12

<210> SEQ ID NO 22
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 22

ttcactttca ta 12

<210> SEQ ID NO 23
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 23

attcactttc at 12

<210> SEQ ID NO 24
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 24

agtaagattc acttt 15

<210> SEQ ID NO 25
 <211> LENGTH: 28070
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

ccacaaatgt gggagggcga taaccactcg tagaaagcgt gagaagttac tacaagcggg 60
 cctcccggcc accgtactgt tccgctocca gaagccccgg gcgcggaag tegtactct 120
 taagaaggga cggggcccca cgtgcgcac ccgcggttt gctatggcga tgagcagcgg 180
 cggcagtggg ggcggcgtcc cggagcagga ggattccgtg ctgttccggc gcggcacagg 240
 ccaggtgagg tcgcagccag tgcagtctcc ctattagcgc tctcagcacc cttcttccgg 300
 cccaactctc cttccgcagc ctcgggacag catcaagtcg atccgctcac tggagttgtg 360
 gtcccggttt ttctacgtct tttcccactc cgttccctgc gaaccacatc cgcaagetcc 420
 ttccctcagc agtttgggct ccttgatagc gttgagtgga ggccctgccc cgacttggca 480
 gtagcttatt ttgttcaact ctctctggct ggtgtggggg aggtgggggc attaggccag 540
 ggtgaagcag gggaaacct taggagtctg ttaagatgat ctgaacttca gaacaagatg 600
 ttattaacag agtgaaagta tttggattct gggtatattt tgaaatcgga ggcaacaggt 660
 ttttcagata gattcgataa cggaggttat cctgaaatag tgaaaagata aagttgcctt 720
 ttgctgaggt gggaaagaga agattgccag tagagcaggt ttctcaggag ttcagtcttg 780
 ggcatagcat ggtaggggtg aatttggctg gactgagttg gagagtagga gaagagaaat 840
 ccaaggcaac atttgaccag cctgggcaac atagtgtgac tccgagctcg caaaaattag 900
 acgggtgttg tggtgcgcgt ctgtggtctc agctacctgg aaggttcagg ccttgggaagg 960
 ctcagggagg tggaggctgc agtgatctgt gattgcgcct ctgcactoca gcctgggcca 1020
 cagagccaga ccctgtctta aaacaaaata aacggccggg cgcgggtggct caagcctgta 1080
 atcccagcac tttgggaggg cgaggcggcc ggatcacaag gtcaggagat cgagaccatc 1140

-continued

ctggctaaca	cggtgaaacc	cgtctctac	tacaaataca	aaaaattagc	cgggcgtggt	1200
gacgggccc	tgtagtcca	gctactcggg	aggctgaggc	aggagaatgt	catgaagccg	1260
ggaggcggag	cttgacgtga	gccgagatcg	cgccactgca	ctccagcctg	ggcgatagag	1320
caagactccg	tctcaaataa	ataaataaat	aaataaataa	ataataaaaa	catcggtag	1380
catatttcaa	ggaattctat	ttaaaaaaaaa	tttttttaga	gacaagttcg	ctctctgtgg	1440
cccaggctgg	agtacagtgg	catgatccta	gcccatggca	gcgttgatct	cttgccctca	1500
agcgaccctc	ctttggagtc	gctgggacct	aaggagtggg	ccaccacgaa	atcttattat	1560
aatggagggg	tagagaaatt	gggcaataaa	tggaggggga	agtgagttaa	gaggaatctt	1620
aattatgtgt	gtgtggtttt	aaaagagggg	ggtcttgctc	tggtgccag	gctgctgggg	1680
tgccagtggc	gcaatcatga	atcactacag	ccttggaactc	ctggcctcaa	gctatcctcc	1740
cacctctgoc	tcccaaagta	ctgggattac	tagtgtgagc	cactgcacta	agataggagc	1800
aacatgtttc	agcatgtttg	tgggttgata	ggaaagatga	gaatgggaaa	gttgatgtcg	1860
gaaagaagac	aatggctaga	gcaatgtcct	agagttaggta	agaagggatg	gatttggcct	1920
ttgttgaaa	cattagcggg	tcttttggtg	acagctatat	agttaacaca	tctatgatac	1980
gtgaatgggc	agataggatg	gcaggagatt	ttgaaagttc	tcttgattct	tactgttctc	2040
ttagtgaag	aagcaaggtt	atcagctaga	agctgggatg	ggagaggaaa	gagaagatgg	2100
gaagtagata	gttctttaga	agagtgggca	agggttgac	tagggaagtt	tagtgaaat	2160
attgctaggc	aacataaaga	gcctacttga	gattcgtggt	catgagtga	aggagaccag	2220
acagcaagat	tgtgtatgag	ggcaccocaca	gagtaaatgg	agagttgaaa	ttaatgcagt	2280
tgtgatatta	ccacgtggat	atgaagaagt	gagggggaga	agtacaaagg	agttctctta	2340
atgattgacc	atggaattta	agctggctaa	gaaaggaagt	gagaggccgg	gctcggtggc	2400
tcacgcctgt	aatcccagca	ctttggggaga	ctgagggtggg	tggattacct	gaggtcaggga	2460
gtttgagacc	aacctggccg	atatggcgaa	accccatctc	taataaaaaat	acagaaaaat	2520
tagccgggaa	tgggtggcagg	tgccctgtaat	cccagctact	caagaggctg	tggcaggag	2580
atcccttgga	cccaggaggt	ggaggttgca	gtgagccgag	atcacgccac	tgtactccag	2640
cctggacgat	atagttagac	ttcacctcaa	aaaaaaaaaa	aaagaaagga	agtgaggatt	2700
ttaagacct	gagagacagt	ttaaaaagt	ggagatcgg	ccgggcgctg	tggctgacac	2760
ctgtaatccc	agcactttgg	gagggcaggt	tgggcagatc	acaaggtcag	gagttcgaga	2820
ccagcctggc	caatattggtg	aaacctgtc	tctactaaaa	atacaaaaaat	tagccgggca	2880
tgggtgcacg	tgtctataat	cccagctact	cgggaggctg	aggcagaaaa	attgcttgaa	2940
cctggggaggc	agaggttgca	gacagctgag	atcactccat	tgcactccag	cctgggcaac	3000
aagagcaaaa	ctttgtcttt	aaaaaaaaaa	aaaaaaaaag	aatacaaaaa	ttagccgggc	3060
gtggtggcgc	gtgcctataa	tcccagctac	ttgggaggct	gaggcaggag	aatcagttga	3120
acacgggagg	cgaggtttgc	agttagccga	gattgcgcca	ctgcactcca	gcctgggcga	3180
cagagcagga	ctcctcttgg	aaaaaaaaaa	ttagctgggc	atggtggcag	gtgcctgtag	3240
tctcagctac	tagggaggct	gaggcaggaa	aatcacttga	acccgggatg	tggagtttgc	3300
agtgaccoga	gatcgtgcca	ctgtactcca	tcctgggcga	caaaatgaga	ctctgcctca	3360
aaaaaaaaaa	aaaaaaaaag	tgggaggatc	aatgtactgc	cagtcctaat	gaagtggaat	3420

-continued

gattgtcccc atcaaatcac tagtaggagt aagttgcaga gcctagaagg tgatggtaa 3480
gagagtggga ttcttgaaac tgcatttatg gagaggttgt ggttattggt tataataaat 3540
aaatacagtt gaagtggatg agtagctgag atttggggat gtatcagttc attcttacac 3600
tgctacaaag acatacctga gaccaggat ttataaagat aagaggttta atcagctcac 3660
agttctgctg cctgtacagg cttctcttgt ggagocctaa ggaaacttac agtcatggtg 3720
gaaggtgaag gggaaacaag cacagtcctc acatggccag caggagagag agagaagggg 3780
gaagtgtcac atactttaaa acaaccagat cttgtgagaa cgcttatcag gaaacagcac 3840
ttggggatgg tgctaaatca ttgaaatca ccccatgat ccagtcgct cctaccatgc 3900
ccacctcaa cactggggat cacaattcag catgagattt gggtaggaac acagagctgc 3960
accacatcag aggatgtaca agattgtggt ggagaggagt ttagagacct gcaaataatag 4020
ggtaattgaa gggatcatct acatggatat ttaaatcacc aaaaattatg acaggagtag 4080
tgttggagag agaactcgga tgtaaacatt aaggaatgag gaagagtgc tcggtaggct 4140
gtaggtgact gcaataggaa acgataatag actgtgagtc tggtgacaag attttcttc 4200
tttctttttt tccccccccc cgagacaggg cctctttttg ttgcccaggt gggagtgcag 4260
tggcgcatc acggctcact acaacctcct cccaagctca agggattctc ccacttcagc 4320
ctctcaagta gctggaacta cagggtctga ccaccatgcc tggctacttt ttgtcaggat 4380
tttcaaggct gggaaatttg agaggggaat ggagagaaat aatctgaaag tgcaagtaag 4440
gagcagggaa gatttctttt ttcttttttt tttttttttt tgagtcggag tctggctcag 4500
tcgcccaggc tggagtgcag tggcgagatc tccgctcact gcaagctccg cctcccgtgt 4560
tcacgccatt ctctctcttc agcctcccca gttagctggga ctacagggc cgcaccac 4620
gcccagctaa ttgttttttt gtatttttag tagagacggg gtttcaacct gtttagccag 4680
atggtctcaa tctctgact ttgtgatccg cccaccccg cctcccaaag cgcttgggat 4740
tacaggcgtg agccaccgcg ccagccagag caggaagat ttcttcccca catctccagt 4800
aggtagcagtg atatgaagtg tgtggaggag aaaagaggaa acatctatca tttgagatgg 4860
ctgcgaaagg aaaaggcatc ctccagggagc tagattttac ttagagcaag aaatgaaggg 4920
atgattcaga ggttaaaaga gtggatttta tgaattactc aagggagcac agtggaaagt 4980
tcaggaagtg gtaggagaag gtagaagatg gcaggggtgt gggaaataatt tgagaaatct 5040
gagctactgg aatgactga gaatcagata taaaggcagt cctgggtggc cgttctggct 5100
gccgttgctg tgtaacgaat ctgccaaaac ttagtggctt gaaacaacaa agaacatttt 5160
attatctctc attgtttctg tgggtagga atttgtgaga gccgtgctgg gcagtttctg 5220
tgccgctgtc tcgtggttgc acctacatag ttgctagagc tacagtagct ggggactgag 5280
cagctagga ttggcaggct atctcttttt ttcattgagt ctcatgaaga tttctttatg 5340
tggtttcaat gtgtgggctg gtttgattt ccttatagca tgggtggcctc agttggattg 5400
ctgttttctg atccttttca tccctccttg tccgtcctcc agacaaccac tgatctactt 5460
tctgtcacca tagattagcc tgcattttta agaattttta taaacgtgga atgatagagt 5520
accttttttg tcaegtttct tttatttata atagetattt tgattttcat ccattttatt 5580
gctgagtagt atcccattgc atgtatatac tatactgtat tcattcgctt gcttgtgaac 5640
atttgggctt tttccagttt gggactgtta acaagtagag cccatgaa tattagtga 5700

-continued

taagacttca tatagccaag gctggcagat cgcttgagcc caggagtttg agaccagcct 5760
gggaaacatg gtgaaacctc tatttttatt ttaaaatcaa aaattaaaa tttctataa 5820
aaaattttaa agaagacttt gtatagacat acgctttcat tttcttgag tgaacttta 5880
ggtctcaggg tagatgtatt ttaagtcttt aaggagctgt caaaccttc ctcaaagtgg 5940
tggttgtaac atgttacttt ttaatataac agagattaat tgagcaaaga aaaattcaaa 6000
agttggacag cccccacaac taaatagggt cagaacagct cccccat ttt gcattttgac 6060
cagcaatgta tgaaggttcc atttgetcag tgteectgca aacacctggg atggtcagtc 6120
tttttaattt taggcattat aatagatata gtggcttctt gtgattttaa ttagcatttc 6180
ctaagacca gtgctgtgtg tgatcatttc atgagtgtat ttgcatccg tatatctttt 6240
ttggtgaaat gtctattcaa atcatttggg tttttttttt ttttgttttt tttttttgga 6300
gacagtgtct cactctgtca cccaggetgt tgtgcagtggt tgcaatcaca cagcctactg 6360
cagcctccac ctctctgect cagtcttctt gtctcagcct tctgagtagc tgaatttacg 6420
agcacagccc acaatgctg gctaattttt taaaattttg tagaacaag gtctcattat 6480
gttgctggg cttgtctgta actcctgggc tcaagcaatc ttctgcctc agcctcccaa 6540
agattgggat tgcaagtatg agcactgca cccggccaac ttaccatct ttaattgaa 6600
ttttttgtt gttgaggtt gagagtctt catgtttgtt gggtaacaata tctttatcag 6660
ataggttaact tgcattgtatt ttctccggg ttacactttg gtttttcatt ttgttaaaa 6720
cgtcttttta agaacagaaa atcttaattt tgctgaaatc taatttttca gttttttctt 6780
tgatggtttt gagagaggag gtaaaaaaag actaggtaag ccgatagtta gacagagtc 6840
tcggtagaac ttcccttcta acaaaaagca gcccaagaaa tcaacttctt tctaacaagg 6900
agcagcctgg aagatcgggc tgtaaacatg tataaggaag cagctctggc acagaggggg 6960
agcttctctg gtaatcagca agcttcacat acgtaagggt ggatgtgaa gtaaacacag 7020
tatgtgaagt aaacacagt gaccttagta catactcaga taaggaagct ggaagcttgc 7080
atgttgtagg ttgttggggg tgctctcagc tgcacggaga gaaaggggta cctggggcca 7140
ggcatgtcca ccatgggtggc tccacctccc cttatttagc acatgcaca taggaaagag 7200
ataagcaatg tggagtagct caggccaagg acctgcctgc ataataaaag gttgggggtg 7260
gggatgccag agattcagc tctgtgcaga tggcaacacc tggctcctaac tggttttttg 7320
ctccctatgt gtagataagc taccctctc ccattagctc atttataaaa atgcttgcat 7380
ttcactgtgg aatgggaact cttttcagga cctctctctg caggagagag ctagtctctt 7440
tcttttgect attaaacttc tgctctagcc tcacacctt ggtgtgtcag cgtccttgat 7500
ttcctcagcg tgagaccaag aacctcgggt gccaccccag gcaacaaggc catttcagtt 7560
tgttcttttg ttataggcaa tccatgatca cagatttttc tctctttttt ttttttacac 7620
agtttagagt tttagtttta cacttaggtc tgtaatccat tttgtattaa ttcttatatg 7680
tggctcagtg taggtggaaa tttggtttgt ttttgcataa ggatttccaa tagttttacc 7740
accatttctt gaaactacta tgctttctct attaaaccac atttgtaact ttagttaaaa 7800
tcagtcacat atatcacagg gctatttctg actctcaatt ctgttacatt gtctattagt 7860
gtatattgat gtcagtacta cacttttaat tactattgct tcagggtatg tcttgtaaac 7920
caaaaataaa attataggcc cccccgccc ctgcacaacc aactgaatgg acctatctc 7980

-continued

t cagccaagg gcattccaaa attaacctga aaaactagtt caagccatga tgggaagggg 8040
gagttggaca tgtctcatca caccctacta ccttttgaa ttactgatag aacagactct 8100
taaagtctga aaagaaacat ttacaaccta cctctctga agcctgctac ctgggagctt 8160
catctgcatg ataaaacctt ggtctccaca accccttatg gtaacccaaa cattcctttc 8220
tgttgataat aactctttca actagttgcc aattagaaaa tctttaaatc ttcctatgac 8280
ctagaaacct ccctaccccc actttgagtt gtcctgcctt tcctgacaga actcatgtac 8340
atcttacata tattgattga tgcctcatgt ctccetaaaa tgtataaaac aaagctgtac 8400
cccaccacct tggggacatg tcatcaggac ctccctgtgc tgtgtcatag gagcgtcttt 8460
aactttggca aaataaactt tctaaattga ttgaaacctg tcttagctac ttctggttta 8520
cagctctaaa gttagataat gtaaatgtc cagctttggt ttatttttgt ccttagtagt 8580
tccatataaa ttttagaatc agcttttcaa ttaatacac tactttcctc ttagatccac 8640
aattaaatat atttgatgct aacaattctg ttttatgttt ttcgtttttt ttttttgaga 8700
caagagtttc gctcttggtg cccaggctgg agtgcagtg cgcgatcttg gctcaccaca 8760
acctccacct cccaggttca agcaattctt ctgcctcagc ctcccgagta gctgggatta 8820
caggcatgcg ccaccacgcc cggctaattt tgtattttta gtagagacgg ggtttcacca 8880
tgttgatcag gctggctctg aactcctgac ctccaggtgat ccaccacct cggcctccca 8940
aagtgttggg attacaggcg tgaaccacca tgcctggcca gttctgttat ttttaaacc 9000
caagtttccc tggcatatc ttggttgat gaagcgtatt ttcaatagat taccctggaa 9060
aggctagtga gtacggtatt ctctacatt ttagactttt cttagtcttg ctactcaag 9120
gacagctagg ctgcatataa aattcttggc tcatactttt tccccataaa tttctatgag 9180
aaagtctaata gataactgat tttctttatt ttgtaactta gtctttttgc ttagaggctc 9240
tctgaggatg ggaggggggt ctccctocca tccctaggaa tttttctttt ttttaaatc 9300
ctaataacta gaccaccagg aagattggtt gttttgttt gtttttattc ttcagggacc 9360
ccatttatac atacgttaaa taaatactgt ttgccaatgt atcaaccatt ttgcttctta 9420
tttatttttg ttcctttggt tctttttcat ggctttgctt tgggtgctct tagattttca 9480
gtcagatgta tttgtccttg ggtacctgt aatcagatt acctttctt ctgtcgttt 9540
gttttctggt cgttttgaaa ttacttggtt cctggtctgg caataacagt tgagatatga 9600
ggagtttgag ctgccatctg tctatgtatc ttgctttaag actgcaactc tctattgata 9660
tcaactggcct tgattttgtg atttctttat ttcttcagga ccaccttca ttttctactg 9720
tttgcttctt ttttttttga gatggagtct cactctgtca ctccaggtgg agtgcagtg 9780
tcttggtcga ttgcaacctc tgcctcccggtgtccagca tctctctgct cagcctccc 9840
aagtatctgg gactacaggt gtgcaccacc atgcccggct aagttttgta tttttaatag 9900
agacgggggt ttgccacatt ggcaggtctg tctcaaactc ctgatgtcaa gtgatccacc 9960
caccaccaacc acctctgcat cccaaagtgc tgggattaca ggaatgagct gccgtgacca 10020
gcctcccccc taccctctt tttttcttc gagacagaga ttatagggtg gagccactgg 10080
accagcctg tttttattcc ttttaccaaa tctccaagga atatcttccc tccaagtgc 10140
gaatgtaacc ttaagtcagt taacctctt gtgattactt ttcttatctg caaagtgact 10200
taatgatctt aagtactttt ttttttgag acaggtctc actgtcacc tggctggagt 10260

-continued

gcagtgccac gatctctgat ctccactcac tgcaatctcc tcttccctgg tccaagcggc 10320
cctcccacct tagcctctctg ggtagctggg actacagatg tgaaccacca cgcccagcta 10380
atTTTTgtac tTTTTgtaga gatggggTtt tgccatgttg cccaggctgg gattattaag 10440
tactTTTTat catacagcaa gattgacatt ttatatggga atacatttgt ctctatataa 10500
cggagattaa caggaaaaatg acaagcctgg gtgcggtggc tcatgcctgt aatcccagca 10560
ctttgggagg ctgaggTggg aggatcactt gaggtcagga gtccgagacc agttttgcca 10620
agatgatgaa agcccctgtc tactaaaaat acaaaaatta gcccagcttg atggTgggcg 10680
cctataatcc cagctatttg agagactgag gcaggagaat cacttgaacc tgggcagcag 10740
aggTtgccagt gagccgagat catgccactg cactccagcc tgggtggcat agcgagactc 10800
ttgtctcaag agaaaaacaaa acaaaaacaaa aaaaaaacag gaaaatgaca aaaagtaata 10860
ttacaactca gtgaatttta taacaaactt ttttggatt cattgactaa tactatacca 10920
aatccaaaat actctctagt ataccaaaat caactctacc ctatagtata aattggattc 10980
tatttggaact tgtctcacta atccctcata cagtgtgttt tattTTTTat tgaagtaaaa 11040
aaatttgTca ttttaacccat ttttaagtat atagtTcagt aatattaagt atgtTcatgt 11100
tgTtgccGaa tagatctctcG gaagTtttTc gtctTgcaac ctgaaactct acccattagc 11160
aaattcccat ttctccttac acttagccct tggtaatcat cattctTTTT tttTTTTTTT 11220
tgagatggag ttttactctt gttgcccagg ctggagtgca atggTgcaat ctcgactcac 11280
cacaacctcc gcctcccagg ttcaagcaat tctacctcag cctcccagat agctgggatt 11340
acagtcatgc accaccaGcG ccggetaatt ttgtattttt agtagagaag gggTttctcc 11400
atgttgaggc tggTctcGaa ctctgacct caggTgatct gcccacctcG gcctcccaaa 11460
gtgctgggat tacaggcgtg agccactcGg cctggcccat tctttctaat tctataaatt 11520
tgactactta gttaccttac ataataaatt tcttatagtt agtGttattt ttgcttccat 11580
gcctTTTTtg ttgtTgtTca tGctcttact tggaaTcGgt tctattTgt ctacctatGc 11640
acatcctgTt gggTttTTTT tttTTTTggg ggtTTTTTTT gttTTTTTTT gttTTTTTTT 11700
cccagacaag gtctcaattt gttaccagg ctggagtgca gggGcccat ctccactcac 11760
tgcatcctca acttccTggg cccaggTgat cctctcGcct cagcccctGc aggtagctgg 11820
gactataggc atgtgcccacc atgcccagct aaattTggtt tttTgtTtg ttgtTttTg 11880
agacagagtc tcaactctGtc acccaggctg gagTgcagtg gcacaatctc agctcactGc 11940
aatctctGcc gcccgggttc aagtgattct cctGcctcag cctcccaagc agctgggatt 12000
acaggTgact gccaccagcG cagctaagtt ttgtagTttt agtagagatg gggTttccacc 12060
ttgtTggcca tGctggtctc gaactcctga cctcgtgacG tGcctGcttc tGcctcccaa 12120
agtGctggaa ttacaggcat gagccaccac gcccggccag aattTttGta tttTtagtag 12180
acacaaggTt cttacctgtT tGcctaggct ggtctGgaag tccTggactc aagcaattca 12240
cctGcctTgg cctcccaaaa tGctgggatt acaagccacc atgcccggcc taaatcctgt 12300
tgTTTTgttt tGTTTTatTT tGTTTTgtTT tGTTTTgtTT gttTTTTgag acagagTctc 12360
gctatgtctc tcaggctgta gtgcagTggc gcgatctTgg cTcactGcca cctctGcctc 12420
ccaggTtcaa gtgattctcc tGcctcagcc tcccaagtag ctgggattac aggcattgtGc 12480
tactatgtcc ggctaatttt tGtattTTta gtagagacag ggtttcacca tGttggccag 12540

-continued

gctggtctcg aactcctgac ctgctgatcc acccacctcg gccacccaaa gtgctgggat 12600
tacaggcgtg agtggttttt atttcttagg ccggtttcct ccatatgatc ttgcagtaga 12660
cattaatttc tttccttttt aattaaata ctgtttgtat ttcacatttt gatgtttgtt 12720
aagatttgtt ttatattggt ttttgttttg tcttggtga tagtcttaa tccctagtta 12780
gataataact ggagagtacc atgtttctat atatctctca gtgacttgca cagtgcctagc 12840
agatagtgct aaaaaattat ttattattat tattattttg ttattgtgtg tgttgttgtt 12900
agacagggtc ttcctctgtc acccaggcta gagggcaatg ggatgatcat agcttactgc 12960
agcctccaac aactgggctc atgtaattct cctgcctcag cttcccaagt agctgggatt 13020
acaggcatga gccaccatgt ctggacaaaa atatttccag gtgcagtgcc tcatgcctgt 13080
aattcccaca cttgggaggc cgagcgaggc tggaggatca cttgagccta ggagtccaag 13140
accagcttgg ctaagatggc gagaccccg tccctacaaaa aattttaaaa actagccagg 13200
catggtggca tgcacctata ttcaccaacta ctcagtgggc tgaggtggga gggtcatttg 13260
aacacaggaa tttgagggga gaaaaaaga agagagaaag agaagtgaag gaaggaagaa 13320
aggaaggagg gagggagaga agaaagaaac gaaagaaag aaaagaaag gaaggaagaa 13380
aaattggtac caggaaagca ggaaggga atggaagtaa aaaaaataata ataataataa 13440
aatgaaaatt ggttagtcac tattaacaat ttgtatcctt ataactgga aacattataa 13500
tttcaaaaga aaaaatattc tttgatcat aggttctgag gtcagaacag cattcccgta 13560
gtctagatga agtcaagttt tatctgatct taattgaaat aaatatagct ggcttgaac 13620
aaatctactc atggtatgtg gataggaatt aaattgtagg ggcattcact tgatggcatt 13680
cattcttaga acatttaact atgtctagct tttggagtaa agtcacataa cctctaacca 13740
ggtaagtttc ctgtggcttt atttaggatt ttaaatactc attttcagtg taattttgtt 13800
atggtgggat taagatgact cttggtacta acatacattt tctgattaaa cctatctgaa 13860
catgagttgt ttttatttct tacccttcc agagcgatga ttctgacatt tgggatgata 13920
cagcactgat aaaagcatat gataaagctg tggcttcatt taaggatga aatgcttgc 13980
tagtctgttt cttattttct cgttattcat ttggaagga attgataaca tacgataaag 14040
tgtaaagta catgttattc agttttcatt ttgaagatta gatgtagta tgagttagtt 14100
aaatcagggt atatcctcct ttagaagttg atagcctata tatgtcatcc tttgtggagg 14160
caatttaaat aaaatttaaa acattttattc ctggctgggt atggtggctc actcctgtaa 14220
tcccagcact ttgagaggct gaggcgggtg gatcacctga ggtcaggagt ttgagaccag 14280
cctggccaac atggtgaaac cccgtcttta ctaaaaatac aaaaattagc caagcatggt 14340
ggcacgtgcc tgtaatccca gctgcttggg acaactgagc aggagaattg cttgaacctg 14400
gggggcagag gttgcaatga ttgcaccact gcactccagc ctgggcgata gagtgagact 14460
ccatctcaga aaacgaacaa acaatgtatt ccttttagta tttttacatt gtatcaaact 14520
atggaagtcc tctaattgag attaataaga aaaagacaat ctgaattata attttaaaca 14580
tttaacaagc atgtagtaaa ataattgatga agataaatag cattagtaca gcaattaata 14640
ttttagcat gctgacagtg ctctgtgtgc gtttcatata ttaaattact ctaatcatcc 14700
caaatcctgt aagtgggta tcaattcaag tgttcctatt gggtaggaat atacagttct 14760
tttaggaaat gtagtatggt tctgtgtctc aaacaggaca cttacacagt tggccaacat 14820

-continued

catcaccttc tccattctct gagatgttta gtcttactga gcaactaaata tgggtcatca 14880
atagtcacaga ctaccttgag caaacaatag tccagactac cttgagcaaa cagagcatat 14940
actcatacag tgtataaaga gcaccaagca tacagatttc atgtctttct catagttact 15000
cttgtaacat gagctaaaga tcagacctct atgtcacctt tgtaactgat ttctagattt 15060
ttttttttt ttgagatggg gtcttgccct gtcacccagg ctggagtgtg gtggcgtgat 15120
catgcctcat tggagccttc aactcatgag ctcaaaaat cctcctacct cagcttctctg 15180
agtagttggg accacaggtg tgtgccacca caccagctc atttttgtat tctttgtaga 15240
gatgcagtct caccctgttg cccacgctgg cctggaactc ctgagctcaa aagatcctc 15300
cgccttgacc ttccaaagtg ctgggattac aagcatgaac cactgcaccc ggcttagatt 15360
tttaaatgtg ctttcagta taaactgaaa ctagaagtgc actaaagaat taccaagaga 15420
attctataaa atagagattg aaatggggct cgatgtggga tgggttggtg atattgcagg 15480
gagaagtaat ctgagtaaag gaggaaaaga actgatttgg gaaaacgata gttttagtag 15540
tgagtttgag tatgaattaa gttgagattg aatttgaatt aagttgaggt tgaatatgaa 15600
ttaagttgag gttgagttg aggtatgaat taagatgtga aattgatcat tggaaatgtt 15660
agattgagaa aagtcacagc tggattaata gcttcagaag tgtgtttgca gacagttgca 15720
actaaagtaa taagaataga tggccttggc cgggcgcggt ggctcacgcc tghtaatccca 15780
gtactttggg aggctgaggc gagcaaatca cgaggtcagg agttcaagac cagcctggcc 15840
cacatggtga aacccctct ttattaaaaa taaaaaatt agctgtgcac agtgggtgcac 15900
gcctgtaate ccagctactc gggaggctga gacaggagaa tcgcttgaac ctgggaggtg 15960
gaggttgacg tgagctgaga tcagtgtgac tgcactccag cccggtgaca gagtgagact 16020
ctgtgtaaaa aaataaaata aataaaataa tggccgtaag caagtaaaaga aggatggcca 16080
gctcttattg ggaatgccta aatctaagc ttgatcagaa gtaatgaaac cgttggggcc 16140
ctacattgct atgacatcca aagggccatg aatatacagga agaaagataa ttaacaggg 16200
ctaattgtac agagaggttg agagcaagga gatttgatta aaaggtcct tagagctgat 16260
gtcaggtgta tgatgccttt aagagcagtt tttatagtgc agggggtggt caaaagagaa 16320
aataggtgct ttctgaggtg acggagcctt gagactagct tatagtagta actgggttat 16380
gtcgtgactt ttattctgtg caccaccctg taacatgtac atttttattc ctattttcgt 16440
agcatgctct aaagaatggt gacatttgtg aaacttcggg taaacaaaa accacaccta 16500
aaagaaaacc tgctaagaag aataaaagcc aaaagaagaa tactgcagct tccttacaac 16560
aggttatttt aaaatgttga gatttaactt caaaggatgt ctcatagtc cttatttaat 16620
agtgtaaaat gtctttaact taagtgtgta gtacagtgtt tctattgaca tatacttata 16680
caacttcaaa aacaactatt aaattttctg ttatttagga acatgcatat tagtcatgaa 16740
agtataaaga attagatggg aatgataaat gctaaaatca ggacatgtgt tccatttgtg 16800
aatggaaggc agggagaagg tgccgttttg aaggagtacc caagagcgt aagctgaatt 16860
ggcagtgttt tacatcttaa gctgagagat agattttttt tcccccttt tctttaaaaa 16920
ctctaaaact gttaattoca aggaaccocag aagtctaggt agattatttc tgctagttaa 16980
aagcagtagt cctgaaagct gaatattttg gtgtcttttg agccaacttt agtttcatca 17040
ttaccaaggg ggaagagagc taacagttga tgagcacttg ctctaggcca gtccagagtg 17100

-continued

ctgggcacca tacgcatttt atctccctcc cgctattcac aacaaatag ggaggtagtt 17160
tatattatag ccatctaata agatggggaa actaagactc aaagagattc agaaacttgt 17220
ccatgattat aaatgtaaga gagttggaat tcagatttat gtatttagac cccaagcctt 17280
tctcattaca tcattttgcc ttccaatct ctaccctcta tctttcacct ccccactgat 17340
caaaacgaga tgatagtttg cctcttcaa aagaaatgtg tgcattgata tatctttgat 17400
ttcttttgta gtggaaagt ggggacaaat gttctgcat ttggtcagaa gacgggtgca 17460
tttaccagc taccattgct tcaattgatt ttaagagaga aacctgtgtt gtggtttaca 17520
ctggataggg aaatagagag gagcaaaatc tgtccgatct actttcccca atctgtgaag 17580
tagctaataa tatagaacaa aatgctcaag aggtaaggat acaaaaaaaaa aaaaattcaa 17640
tttctggaag cagagactag atgagaaact gttaaacagt atacacagtt gtcagtttga 17700
tccaccgagg cattaatttt ttcttaatac cacccttata acaaaaacct gcatattttt 17760
tctttttaa gaatgaaat gaaagccaag tttcaacaga tgaaagtgag aactccaggt 17820
ctcctggaaa taaatcagat aacatcaagc ccaaatctgc tccatggaac tctttctctc 17880
ctccaccacc ccccatgcca gggccaagac tgggaccagg aaaggtaaac cttctatgaa 17940
agttttccag aaaatagtta atgtcgggac atttaacctc tctgttaact aatttgtagc 18000
tctcccatga aacttttgta gcttaatac acaagaattt ttgaaaagg aaataagata 18060
atgatgcaaa atagttaatt ttttaaaaa atgttagaca ctgcagtgga tgcaacaaaa 18120
tactttatat gaaagattta tccagttaac tttgtggag tattaggtat tagactaata 18180
attagcacac ttacttaagt tagaaagtat aataatgccc cggacgcggt agctcacgcc 18240
tgtaatcca gcaacttggg aggccaaggt gggcggatca caaggtcagg agatcgagac 18300
catcctggct aacacggtga aaccccatct ctactgaaaa tacaaaaaaaa tttgcccggc 18360
gtgatggcgg gcaactgtag tccagctac tccggaggct gaggcaggag gatgggtgta 18420
accccgagg cagagcttgc agtgagtcaa gatcgtgcca ctgcaactcca acctgggcca 18480
cagaatgaga ctccatctca acaaaaaaaaa caaaacaaaa caaaaaaaag tgtaataata 18540
atztatcatt agctggatga tatgctgttg tttcccatgt cacctgtata agatatgtaa 18600
aataagaaca cattatttac atctaataa gataaaatcc tgaggcctc tcagattgtt 18660
ttgtagagtt caaatgtaaa tattgttttc atttatggtc cttttggtta taagtaacag 18720
aaatcaactc taaaagatt tttattatag gttagattat gtcattggaac cttaaggctt 18780
gtccctttct agttcttttg tgtaaagcgg tgatttctc catggaggga atggattta 18840
ggcaattttt tttttttttt cgagatggag tcttctctg tcgctcaggc tggagtgcag 18900
tggcaccatt tcagctcact gcaactcca cctcctgggt tcaagtgatt ctctgcttc 18960
agcctcccaa gtagctgaga ttacaggcac ccgccaccac acccgctta tttgtattt 19020
ttagtagaga tggggtttca ccatgttggc caggctggtc ttgaaactct gacctcaagt 19080
gatctcccca ccttggcctt ccaagtgtc aggattacag gcgcctagcc taggcagtca 19140
ttttcaaaaa acaagcatga ctaccacaaa gttttaagat tttctgtgat aatgttctta 19200
ttgaggctta cattatatta cagtttcttg aatctaaaat gatgtacctt cttagaatat 19260
atacatcatg cttcattggt ctccgggggc tgatttttat aaggagagat ttgctagttt 19320
tcacaatatg tctctaatg tggcatgat agctaaacag gctttcataa aaatatacaa 19380

-continued

tttagttaat	gaaatttggg	atatagtctt	ttatgattga	aataattttg	ctaaatagac	19440
tgtctctgat	ttattaggtg	atcaccactc	ttattttggt	ttacttcctt	aatgtctaca	19500
tagaaaggaa	atgagaaaa	tccagagggt	gtcatttgac	ttatgagtct	gtttgacttc	19560
aggatttggg	acatgaaatt	tcacttaatc	tttttgatat	gtataaaaca	aatattctgg	19620
gtaattatth	ttatcctttt	ggttttgagt	cctttttatt	cctatcatat	tgaattgggt	19680
aagttaatth	tcctttgaaa	tattccttat	agccaggctc	aaaattcaat	ggcccaccac	19740
cgccaccgcc	accaccacca	ccccacttac	tatcatgctg	gctgcctcca	tttccttctg	19800
gaccaccagt	aagtaaaaa	gagtataggt	tagattttgc	tttcacatac	aatttgataa	19860
ttagcagaat	agaggattgt	aaaatgtcat	tgtagaacat	cccttgggcc	agattctaat	19920
gggtagaaat	ttgaactaaa	cctctgggtt	ttgtttgttt	ttaatgcctt	tctggttacc	19980
agatgcagtg	ctctttagtg	cccaagtcta	agctctaggt	tgccttcttt	cctggcagaa	20040
gttggtgtct	atgcataaag	gaggtagttc	ctgtagaag	ggatttaatt	ataccttata	20100
taaggaatta	gtgtttgccc	ttctaggtat	agttggatgt	tagcttctga	tgtaaactgg	20160
atthctthth	ctthctctct	ctthththth	ththgtthth	gagggagagt	ththgcctth	20220
taccaccaggc	tggagtgcag	tgggtgtgatc	tcagctcaca	gcaacctccg	cctcctgggt	20280
tcaagcaatt	ctgcctcggc	ctcccaagta	gctgggatta	caggcgactg	ccaccacacc	20340
cggctaattt	ttgttttatt	agtagagatg	gggtttcacc	atgttgcca	gactgatctt	20400
gaactcctga	cctcagggtg	tccaccgcc	ttggcctccc	aaagcctgg	gattacaggc	20460
gtgagctgoc	gcaccagct	gtaaaactgga	tttctaattg	tagattthta	ggtattaaca	20520
atagataaaa	agatacttht	tggcactctg	tgtattggga	tggggttaga	acagggtthc	20580
tacccaagac	atthactthta	aatcgcctc	gaaatgctat	gtgagctgtg	tgtgtgtgtg	20640
tgtgtgtgtg	tgtattaagg	aaaagcatga	aagtatttat	gcttgattth	ththththth	20700
tcatagcttc	atagtggaac	agatacatag	tctaaatcaa	aatgtthtaa	ctthththth	20760
cacttctgtg	ctthctctcc	tcgttaaatt	taattthgtt	ggtctthth	tgttattggg	20820
tggththth	caaatgctag	ctatgttaag	aaatttaagg	ccagggtacag	tggctcatgc	20880
ctgtaatccc	ggcattthth	aaggctgagg	caggaggatc	acttgagctc	aggagthth	20940
gaccagctctg	ggcaacatag	caagacctcg	tctththth	ggggaaaaaa	aagaaatth	21000
agtaggggat	tatataagca	aaaatacaat	taattthcag	cattcactat	ataatataaa	21060
tctccagact	ttactththth	gththctgga	tataaacaat	atctththct	gtctccagat	21120
aattccccca	ccacctccca	tatgtccaga	ttctcttgat	gatgctgatg	ctthththth	21180
tatgttaatt	tcatggtaca	tgagtggcta	tcatactggc	tattatatgg	taagtaatca	21240
ctcagcatct	ththctgaca	atthththth	agttatgtga	ctthththth	taaatthata	21300
aaatactact	tgtctctctc	ththatactac	taaaaaataa	aaataaaaaa	atacaactgt	21360
ctgaggctta	aattactctt	gcattgtccc	taagtataat	ththththth	ththththth	21420
ththctgcta	ththththth	atthththth	tacactthth	aattgaaatt	atactthth	21480
taataatgt	ththththth	gattththth	tgattgtagg	gaatggaaaa	gatgggataa	21540
thththcataa	atgaaaaatg	aaattctthth	ththththth	ththththth	gacggagctc	21600
tgtctctgtg	cccaggctgg	agtgcaatgg	cgtgatcttg	gctcacagca	agctctgctc	21660

-continued

cctggattca	cgccattctc	ctgcctcagc	ctcagaggta	gctgggacta	caggtgctg	21720
ccaccacgoc	tgtctaattt	tttgtatttt	tttgtaaaga	cagggtttca	ctgtgttagc	21780
caggatggtc	tcaatctcct	gaccccggtga	tccaccgcgc	tggccttcc	aagagaaatg	21840
aaatTTTTT	aatgcacaaa	gatctggggt	aatgtgtacc	acattgaacc	ttggggagta	21900
tggcttcaaa	cttgtcactt	tatacgttag	tctcctacgg	acatgttcta	ttgtatttta	21960
gtcagaacat	ttaaaattat	tttattttat	tttatttttt	tttttttttt	gagacggagt	22020
ctcgtctctg	caccacggct	ggagtacagt	ggcgcagtct	cggtcactg	caagctccgc	22080
ctcccgggtt	cacgccattc	tctgcctca	gcctctccga	gtagctggga	ctacaggcgc	22140
ccgccaccac	gcccggttaa	ttttttttta	tttttagtag	agacgggggt	tcaccgtggt	22200
ctcgatctcc	tgacctctg	atccaccgc	ctcggcctcc	caaagtgtctg	ggattacaag	22260
cgtgagccac	cgccccggc	ctaaaattat	ttttaaagt	aagctctgt	gccctgctaa	22320
aattatgatg	tgatattgta	ggcacttgta	tttttagtaa	attaatatag	aagaaacaac	22380
tgacttaaag	gtgtatgtt	ttaaatgtat	catctgtgtg	tgccccatt	aatattctta	22440
tttaaaagt	aaggccagac	atggtggctt	acaactgtaa	tccaacagt	ttgtgaggcc	22500
gaggcaggca	gatcacttga	ggtcaggagt	ttgagaccag	cctggccaac	atgatgaaac	22560
cttgtctcta	ctaaaaatac	caaaaaaat	ttagccaggc	atggtggcac	atgcctgtaa	22620
tccgagctac	ttgggaggct	gtggcaggaa	aattgcttta	atctgggagg	cagaggttgc	22680
agtgagttga	gattgtgcca	ctgcactcca	cccttgggtga	cagagtgaga	ttccatctca	22740
aaaaaagaaa	aaggcctggc	acgggtggctc	acacctataa	tcccagttct	ttgggaggta	22800
gaggcaggtg	gatcacttga	ggttaggagt	tcaggaccag	cctggccaac	atggtgacta	22860
ctccatttct	actaaataca	caaaacttag	cccagttggc	ggcagttgta	atcccagcta	22920
cttgagaggt	tgaggcagga	gaatcacttg	aacctgggag	gcagaggttg	cagtgagccg	22980
agatcacacc	gctgcactct	agcctggcca	acagagtgag	aatttgcgga	gggaaaaaaa	23040
agtcacgctt	cagttgttgt	agtataacct	tggtatattg	tatgtatcat	gaattctctca	23100
ttttaatgac	caaaaagtaa	taaatcaaca	gcttgaatt	tgttttgaga	tcagttatct	23160
gactgtaaca	ctgtaggctt	ttgtgttttt	taaattatga	aatatttgaa	aaaaatacat	23220
aatgtatata	taaagtattg	gtataattta	tgttctaaat	aactttcttg	agaaataatt	23280
cacatgggtg	gcagtttacc	tttgaagta	tacaagttgg	ctgggcacaa	tggtctcacgc	23340
ctgtaatccc	agcactttgg	gaggccaggg	caggtggatc	acgaggtcag	gagatcgaga	23400
ccatcctggc	taacatgggtg	aaaccccgtc	tctactaaaa	gtacaaaaac	aaattagccg	23460
ggcatgttgg	cgggcacctt	ttgtcccagc	tgctcgggag	gctgaggcag	gagagtgccg	23520
tgaaccagg	aggtggagct	tgcagtgagc	cgagattgtg	ccagtgcact	ccagcctggg	23580
cgacagagcg	agactctgtc	tcaaaaaata	aaataaaaaa	gaaagtatac	aagtcagtg	23640
ttttggtttt	cagttatgca	accatcacta	caatttaaga	acattttcat	cacccccaaa	23700
agaaaccctg	ttaccttcat	tttcccagc	cctaggcagt	cagtacactt	tctgtctcta	23760
tgaatttgtc	tatttttagat	attatatata	aacggaatta	tacgatatgt	ggtcttttgt	23820
gtctggcttc	tttcaactag	catgctattt	tcaagattca	tccatgctgt	agaatgcacc	23880
agtactgcat	tccttcttat	tgctgaaat	tctgtgtgtt	ggttatatca	cattttatcc	23940

-continued

attcatcagt	tcatggacat	ttaggttgtt	tttatttttg	ggctataatg	aataatggtg	24000
ctatgaacat	tcgttttgtt	tctttttgtt	tttttggttt	tttgggtttt	tttgtttttg	24060
tttttgtttt	tgagacagtc	ttgctctgtc	tcctaagctg	gagtgcagtg	gcatgatcct	24120
ggcttactgc	aagctctgcc	tcccgggttc	acaccattct	cctgcctcag	cccgacaagt	24180
agctgggact	acaggcgtgt	gccaccatgc	acggctaatt	ttttgtattt	ttagtagaga	24240
tggggtttca	ccgtgttagc	caggatggtc	tcgatctcct	gacctcgtga	tctgcctgcc	24300
taggcctccc	aaagtgtctg	gattacaggc	gtgagccact	gcacctggcc	ttaagtgttt	24360
ttaatacgtc	attgccttaa	gctaacaatt	cttaaccttt	gttctactga	agccacgtgg	24420
ttgagatagg	ctctgagtct	agcttttaac	ctctatcctt	ttgtcctaga	aatctaagca	24480
gaatgcfaat	gactaagaat	aatgttgttg	aaataacata	aaataggtta	taactttgat	24540
actcattagt	aacaaatctt	tcaatacatc	ttacggctctg	ttaggtgtag	attagtaatg	24600
aagtgggaag	ccactgcaag	ctagtataca	tgtagggaaa	gatagaaagc	attgaagcca	24660
gaagagagac	agaggacatt	tgggctagat	ctgacaagaa	aaacaaatgt	tttagtatta	24720
atthttgact	ttaaattttt	ttttatttta	gtgaatactg	gtgtttaatg	gtctcatttt	24780
aataagtatg	acacaggtag	tttaagggtca	tatatthttat	ttgatgaaaa	taaggtatag	24840
gcccggcacg	gtggctcaca	cctgtaatcc	cagcactttg	ggaggccgag	gcaggcggat	24900
cacctgaggt	cgggagttag	agactagcct	caacatggag	aaaccccgtc	tctactaaaa	24960
aaaatacaaaa	attaggcggg	cgtggtgggtg	catgcctgta	atcccagcta	ctcaggaggc	25020
tgaggcagga	gaattgcttg	aacctgggag	gtggagggtg	cggtgagccg	agatcacctc	25080
attgcactcc	agcctgggca	acaagagcaa	aactccatct	caaaaaaaaa	aaaataaggt	25140
ataagcgggc	tcaggaacat	cattggacat	actgaaagaa	gaaaaatcag	ctgggcgcag	25200
tggctcacgc	cggtaatccc	aacacttttg	gaggccaagg	caggcgaatc	acctgaagtc	25260
gggagttcca	gatcagcctg	accaacatgg	agaaaccctg	tctctactaa	aaatacaaaa	25320
ctagccgggc	atggtggcgc	atgcctgtaa	tcccagctac	ttgggaggct	gaggcaggag	25380
aattgcttga	accgagaagg	cggaggttgc	ggtgagccaa	gattgcacca	ttgcactcca	25440
gcctgggcaa	caagagcgaa	actccgtctc	aaaaaaaaaa	ggaagaaaaa	tattttttta	25500
aattaattag	tttatttatt	ttttaagatg	gagttttgcc	ctgtcaccca	ggctgggggtg	25560
caatgggtgca	atctcggctc	actgcaacct	ccgcctcctg	ggttcaagtg	attctcctgc	25620
ctcagcttcc	cgagtagctg	tgattacagc	catatgccac	cacgcccagc	cagttttgtg	25680
ttttgttttg	ttttttgttt	tttttttttg	agaggggtgc	ttgctctgtc	ccccaaagctg	25740
gagtgacgcg	gcgcgatcct	ggctcactgc	aagctctgcc	tcccagggtc	acaccattct	25800
cttgcctcag	cctcccaggt	agctgggact	acaggtgccc	gccaccacac	ccggctaatt	25860
ttttgtgttt	tttagtagag	atgggggttc	actgtgttag	ccaggatggt	ctcgatctcc	25920
tgaccttttg	atccaccocg	ctcagcctcc	ccaagtctg	ggattatagg	cgtgagccac	25980
tgtgcccgcc	ctagtcttgt	atttttagta	gagtcgggat	ttctccatgt	tggtcaggct	26040
gttctccaaa	tccgacctca	ggtgatccgc	ccgccttggc	ctccaaaagt	gcaaggcaag	26100
gcattacagg	catgagccac	tgtgaccggc	aatgttttta	aattttttac	atttaaattt	26160
tattttttag	agaccaggtc	tcactctatt	gctcaggetg	gagtgcaagg	gcacattcac	26220

-continued

```

agctcactgc agccttgacc tccagggtc aagcagtcct ctcacctcag tttcccaggt 26280
agctgggaact acagtgataa tgccactgca cctgggtaaat ttttattttt atttatttat 26340
ttttttttga gacagagtct tgctctgtca cccaggctgg agtgcagtgg tgtaaatctc 26400
agctcactgc agcctccgcc tctcgggttc aagtgattct cctgcctcaa cctcccaagt 26460
agctgggatt agaggtcccc accaccatgc ctggctaatt ttttgactt tcagtagaaa 26520
cggggttttg ccatgttggc caggctgttc tcgaactcct gagctcaggt gatccaactg 26580
tctcggcctc ccaaagtgtc gggattacag gcgtgagcca ctgtgcctag cctgagccac 26640
cacgccggcc taatttttaa attttttga gagacagggt ctcatatgt tgcccagggt 26700
ggtgtcaagc tccaggctc aagtgatccc cctacctccg cctcccaaag ttgtgggatt 26760
gtaggcatga gccactgcaa gaaaacctta actgcagcct aataattgtt ttctttggga 26820
taacttttaa agtacattaa aagactatca acttaatttc tgatcatatt ttgttgaata 26880
aaataagtaa aatgtcttgt gaaaacaaat gctttttaac atccatataa agctatctat 26940
atatagetat ctatatctat atagetattt tttttaactt cctttatttt ccttacaggg 27000
ttttagacaa aatcaaaaag aaggaagggt ctcacattcc ttaaattaag gagtaagtct 27060
gccagcatta tgaagtga tcttactttt gtaaaacttt atggtttggt gaaaacaaat 27120
gtttttgaac atttaaaaag ttcagatgtt agaaagtga aaggttaatg taaaacaatc 27180
aatattaag aattttgatg ccaaaactat tagataaaa gttaatctac atccctacta 27240
gaattctcat acttaactgg ttggttgtgt ggaagaaaca tactttcaca ataaagagct 27300
ttaggatatg atgccattt atactactag taggcagacc agcagacttt tttttattgt 27360
gatatgggat aacctaggca tactgcactg tactctctga catatgaagt gctctagtca 27420
agtttaactg gtgtccacag aggacatggt ttaactggaa ttcgtcaagc ctctggttct 27480
aatttctcat ttgcaggaaa tgctggcata gacgagcact aaatgacacc actaaagaaa 27540
cgatcagaca gatctggaat gtgaagcgtt atagaagata actggcctca tttcttcaaa 27600
atatcaagtg ttgggaaaga aaaaaggaag tggaaatggg aactcttctt gattaaaagt 27660
tatgtaataa ccaaatgcaa tgtgaaatat tttactggac tctattttga aaaacctct 27720
gtaaaagact gaggtggggg tgggaggcca gcacggtggt gaggcagtgt agaaaatttg 27780
aatgtggatt agattttgaa tgatattgga taattattgg taattttatg agctgtgaga 27840
aggggtgtgt agtttataaa agactgtctt aatttgcata ctttaagcatt taggaatgaa 27900
gtgttagagt gtcttaaaat gtttcaaag gtttaacaaa atgtatgtga ggcgtatgtg 27960
gcaaaatggt acagaatcta actggtggac atggctgttc attgtactgt tttttctat 28020
ctctatatg tttaaaagta tataataaaa atatttaatt tttttttaa 28070

```

```

<210> SEQ ID NO 26
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 26

```

```

actttcataa tgctggca

```

18

-continued

```

<210> SEQ ID NO 27
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 27

```

```

cactttcata atgctggc

```

18

What is claimed is:

1. A compound comprising a modified oligonucleotide consisting of 16-20 linked nucleosides, wherein the modified oligonucleotide is complementary to SMN2; and wherein each internucleoside linkage is either phosphorothioate or phosphodiester and at least one internucleoside linkage is phosphorothioate and at least one internucleoside linkage is phosphodiester.

2. A compound comprising a modified oligonucleotide consisting of 16-20 linked nucleosides, wherein the modified oligonucleotide has a nucleobase sequence selected from any one of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 and having 6 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

3. A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 1, wherein the modified oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

4. A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 7, wherein the modified oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

5. A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide has the nucleobase sequence of SEQ ID NO.: 2, wherein the modified oligonucleotide has 6 or more phosphodiester internucleoside linkages, and wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

6. The compound of any of claims 1 to 5, having 7 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

7. The compound of any of claims 1 to 5, having 8 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

8. The compound of any of claims 1 to 5, having 9 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

9. The compound of any of claims 1 to 5, having 10 or more phosphodiester internucleoside linkages, wherein each

internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

10. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNoNoNsNsNsNoNoNoNoNoNoNoNoNoNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

11. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNoNoNoNoNoNoNsNsNsNsNoNoNoNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

12. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNsNsNsNoNsNsNsNoNsNsNsNsNsNsNsNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

13. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNsNsNsNoNsNsNsNoNsNsNsNsNsNsNsNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

14. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNsNoNsNoNsNoNsNoNsNsNsNsNsNsNsNsNsNsNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

15. The compound of claim 3, wherein the modified oligonucleotide has an NsNoNoNoNsNsNsNsNsNsNsNsNsNsNsNsNoNoNoNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

16. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNsNoNsNoNsNoNsNsNoNsNsNsNoNsNoNsNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

17. The compound of claim 3, wherein the modified oligonucleotide has an NsNsNoNsNoNsNoNsNoNsNsNoNsNsNsNoNoNoNsNsN internucleoside motif, wherein each "N" represents a nucleobase, each "s" represents a phosphorothioate internucleoside linkage, and each "o" represents a phosphodiester internucleoside linkage.

sNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

40. The compound of claim 5, wherein the modified oligonucleotide has an NsNsNoNsNsNoNsNsNoNsNoNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

41. The compound of claim 5, wherein the modified oligonucleotide has an NsNsNoNsNsNoNsNsNoNsNsNoNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

42. The compound of claim 5, wherein the modified oligonucleotide has an NsNsNsNoNsNsNsNoNsNsNsNoNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

43. The compound of claim 5, wherein the modified oligonucleotide has an NsNsNsNsNoNsNsNsNsNoNsNsNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

44. The compound of any of claims 1 to 43, wherein each nucleoside of the modified oligonucleotide comprises a modified sugar moiety, and wherein each modified sugar moiety comprises a 2'-methoxyethyl modification.

45. A pharmaceutical composition comprising the compound of any of claims 1 to 44.

46. A method of promoting inclusion of exon 7 in SMN2 transcripts in a cell, tissue or organ, comprising contacting said cell, tissue or organ with the compound of any of claims 1 to 44 or the composition of claim 45.

47. A method of treating Spinal Muscular Atrophy Type I, comprising administering the compound of any of claims 1 to 44 or the composition of claim 45 to a patient in need thereof.

48. A method of treating Spinal Muscular Atrophy Type II, comprising administering the compound of any of claims 1 to 44 or the composition of claim 45 to a patient in need thereof.

49. A method of treating Spinal Muscular Atrophy Type III, comprising administering the compound of any of claims 1 to 44 or the composition of claim 45 to a patient in need thereof.

50. A method of treating Spinal Muscular Atrophy Type IV, comprising administering the compound of any of claims 1 to 44 or the composition of claim 45 to a patient in need thereof.

51. An antisense oligonucleotide of any of claims 1 to 44 or the composition of claim 45 for use in therapy.

52. Use of an antisense oligonucleotide of any of claims 1 to 44 or the composition of claim 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type I.

53. Use of an antisense oligonucleotide of any of claims 1 to 44 or the composition of claim 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type II

54. Use of an antisense oligonucleotide of any of claims 1 to 44 or the composition of claim 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type III.

55. Use of an antisense oligonucleotide of any of claims 1 to 44 or the composition of claim 45 for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type IV.

56. Use of an antisense oligonucleotide of any of claims 1 to 45 for the preparation of a medicament for the treatment of spinal muscular atrophy.

57. A compound comprising a modified oligonucleotide consisting of 16-20 linked nucleosides, wherein the modified oligonucleotide has a nucleobase sequence selected from any one of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, or 27 and having 6 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

58. The compound of claim 57, having 7 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

59. The compound of claim 57, having 8 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

60. The compound of claim 57, having 9 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

61. The compound of claim 57, having 10 or more phosphodiester internucleoside linkages, wherein each internucleoside linkage that is not a phosphodiester internucleoside linkage is a phosphorothioate internucleoside linkage.

62. The compound of claim 57, wherein the modified oligonucleotide has an NsNsNoNoNoNsNsNsNoNoNoNoNoNoNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

63. The compound of claim 57, wherein the modified oligonucleotide has an NsNsNoNoNoNoNoNoNoNsNsNsNsNoNoNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

64. The compound of claim 57, wherein the modified oligonucleotide has an NsNsNoNsNsNsNoNsNsNsNoNsNsNsNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

65. The compound of claim 57, wherein the modified oligonucleotide has an NsNsNoNsNsNsNoNsNsNsNoNsNsNsNoNsNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

66. The compound of claim 57, wherein the modified oligonucleotide has an NsNsNsNoNsNoNsNoNsNoNsNoNsNoNsNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

88. The compound of claim **57**, wherein the modified oligonucleotide has an NsNoNsNsNoNsNsNoNsNsNoNsNsNoNsNsNoNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

89. The compound of claim **57**, wherein the modified oligonucleotide has an NsNoNsNsNoNsNsNsNoNsNsNsNoNsNsNsNoNsNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

90. The compound of claim **57**, wherein the modified oligonucleotide has an NsNoNsNsNsNsNoNsNsNsNsNoNsNsNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

91. The compound of claim **57**, wherein the modified oligonucleotide has an NsNoNsNoNsNoNsNoNsNoNsNoNsNoNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

92. The compound of claim **57**, wherein the modified oligonucleotide has an NsNsNoNsNsNoNsNsNoNsNoNsNoNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

93. The compound of claim **57**, wherein the modified oligonucleotide has an NsNsNoNsNsNoNsNsNoNsNsNoNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

94. The compound of claim **57**, wherein the modified oligonucleotide has an NsNsNsNoNsNsNsNoNsNsNsNoNsNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

95. The compound of claim **57**, wherein the modified oligonucleotide has an NsNsNsNsNoNsNsNsNsNoNsNsNsNsNoNsNsN internucleoside motif, wherein each “N” represents a nucleobase, each “s” represents a phosphorothioate internucleoside linkage, and each “o” represents a phosphodiester internucleoside linkage.

96. The compound of any of claims **57** to **95**, wherein each nucleoside of the modified oligonucleotide comprises a modified sugar moiety, and wherein each modified sugar moiety comprises a 2'-methoxyethyl modification.

97. A pharmaceutical composition comprising the compound of any of claims **57** to **96**.

98. A method of promoting inclusion of exon 7 in SMN2 transcripts in a cell, tissue or organ, comprising contacting said cell, tissue or organ with the compound of any of claims **57** to **95** or the composition of claim **96**.

99. A method of treating Spinal Muscular Atrophy Type I, comprising administering the compound of any of claims **57** to **95** or the composition of claim **96** to a patient in need thereof.

100. A method of treating Spinal Muscular Atrophy Type II, comprising administering the compound of any of claims **57** to **95** or the composition of claim **96** to a patient in need thereof.

101. A method of treating Spinal Muscular Atrophy Type III, comprising administering the compound of any of claims **57** to **95** or the composition of claim **96** to a patient in need thereof.

102. A method of treating Spinal Muscular Atrophy Type IV, comprising administering the compound of any of claims **57** to **95** or the composition of claim **96** to a patient in need thereof.

103. An antisense oligonucleotide of any of claims **57** to **95** or the composition of claim **96** for use in therapy.

104. Use of an antisense oligonucleotide of any of claims **57** to **95** or the composition of claim **96** for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type I.

105. Use of an antisense oligonucleotide of any of claims **57** to **95** or the composition of claim **96** for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type II.

106. Use of an antisense oligonucleotide of any of claims **57** to **95** or the composition of claim **96** for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type III.

107. Use of an antisense oligonucleotide of any of claims **57** to **95** or the composition of claim **96** for the preparation of a medicament for the treatment of Spinal Muscular Atrophy Type IV.

108. Use of an antisense oligonucleotide of any of claims **57** to **96** for the preparation of a medicament for the treatment of spinal muscular atrophy.

* * * * *