



(86) Date de dépôt PCT/PCT Filing Date: 2002/06/05
(87) Date publication PCT/PCT Publication Date: 2004/03/04
(45) Date de délivrance/Issue Date: 2012/10/02
(85) Entrée phase nationale/National Entry: 2004/11/29
(86) N° demande PCT/PCT Application No.: US 2002/017785
(87) N° publication PCT/PCT Publication No.: 2004/018619
(30) Priorité/Priority: 2001/06/05 (US60/296,324)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
A61K 38/17 (2006.01), *C07K 14/725* (2006.01),
C12Q 1/02 (2006.01), *G01N 33/567* (2006.01),
A61K 38/00 (2006.01)

(72) Inventeurs/Inventors:
SHERMAN, LINDA A., US;
CARD, KIMBERLYN F., US;
WEIDANZ, JON A., US;
WONG, HING C., US;
THOMSON, ELIZABETH L., US

(73) Propriétaires/Owners:
ALTOR BIOSCIENCE CORPORATION, US;
THE SCRIPPS RESEARCH INSTITUTE, US

(74) Agent: NORTON ROSE CANADA

(54) Titre : MOLECULES DU RECEPTEUR DES LYMPHOCYTES T SE LIANT A LA PROTEINE P53, ET LEURS
UTILISATIONS

(54) Title: P53 BINDING T CELL RECEPTOR MOLECULES AND USES THEREOF

(57) **Abrégé/Abstract:**

The invention provides T-cell receptor (TCR) molecules that bind peptides derived from the p53 protein, preferably, the human p53 protein. The TCR molecules include both heterodimeric molecules and single chain molecules which specifically bind a sequence preferably spanning about amino acid positions 264 to 272 of the p53 protein displayed in the context of an HLA molecule, preferably, HLA-A2.1. Also disclosed are methods for making and using such TCR molecules. The invention has a wide spectrum of useful applications including therapeutic uses and use in the detection of cells expressing p53 protein.

(74) **Agent(suite/continued):** S.E.N.C.R.L.,S.R.L./LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018619 A2

- (51) International Patent Classification⁷: **C12N**
- (74) Agents: **REES, Dianne, M.** et al.; Edwards & Angell, LLP, P.O. Box 9169, Boston, MA 02209 (US).
- (21) International Application Number: PCT/US2002/017785
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 5 June 2002 (05.06.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/296,324 5 June 2001 (05.06.2001) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **ALTOR BIOSCIENCE CORPORATION** [US/US]; 2810 North Commerce Parkway, Miramar, FL 33025 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **SHERMAN, Linda, A.** [US/US]; 7766 Hillside Drive, LaJolla, CA 92037 (US). **CARD, Kimberlyn, F.** [US/US]; 1120 SW 87 Avenue, Pembroke Pines, FL 33025 (US). **WEIDANZ, Jon, A.** [US/US]; 801 Kachina Drive, Amarillo, TX 79124 (US). **WONG, Hing, C.** [US/US]; 2966 Wentworth, Weston, FL 33332 (US). **THOMSON, Elizabeth, L.** [US/US]; 121 N.E. 45th Street, Miami, FL 33137 (US).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: P53 BINDING T CELL RECEPTOR MOLECULES AND USES THEREOF

(57) Abstract: The invention provides T-cell receptor (TCR) molecules that bind peptides derived from the p53 protein, preferably, the human p53 protein. The TCR molecules include both heterodimeric molecules and single chain molecules which specifically bind a sequence preferably spanning about amino acid positions 264 to 272 of the p53 protein displayed in the context of an HLA molecule, preferably, HLA-A2.1. Also disclosed are methods for making and using such TCR molecules. The invention has a wide spectrum of useful applications including therapeutic uses and use in the detection of cells expressing p53 protein.

WO 2004/018619 A2

**P53 BINDING T CELL RECEPTOR MOLECULES
AND USES THEREOF**

10

15

20

FIELD OF INVENTION

The present invention relates to T cell receptor (TCR) molecules that bind particular p53 protein sequences as well as methods for making and using such molecules. The TCR molecules of the invention are useful for a variety of applications including therapeutic and diagnostic purposes.

BACKGROUND OF THE INVENTION

Traditional approaches to the treatment of diseases such as cancer have included surgery, radiation, chemotherapy, antibiotics or combination therapies. However, such therapies have not proven effective against a majority of these indications. Development of alternate remedies for preventing and/or treating such human diseases is crucial. In recent years

immunotherapy and gene therapy approaches utilizing antibodies and T-lymphocytes have emerged as new and promising methods for treating human disease.

One such approach to treatment has included the use of antibodies
5 for targeting of therapeutic or diagnostic agents to particular targets. Numerous groups have made developments revolving around the use of antibodies as a targeting agent. Such developments have included construction of antibody fusion proteins and antibody conjugate molecules linking antibodies to various effector molecules, including radioactive
10 molecules, chemotherapeutics agents, toxins, and additional bioactive proteins. Therapeutics or diagnostics developed using such molecules are designed to cause a particular effect which is targeted by the linked antibody.

Just as antibodies have been developed as therapeutics, additional
15 primary effectors of the immune system, T cell receptors (TCR), have unique advantages as a platform for developing therapeutics. While antibodies are limited to recognition of pathogens in the blood and extracellular spaces or to protein targets on the cell surface, T cell receptors can recognize antigens displayed with MHC molecules on the surfaces of cells (including antigens
20 derived from intracellular proteins). Depending on the subtype of T cells that recognize displayed antigen and become activated, T cell receptors and T cells harboring T cell receptors can participate in controlling various immune responses. For instance, T cells are involved in regulation of the humoral immune response through induction of differentiation of B cells
25 into antibody producing cells. In addition, activated T cells act to initiate cell-mediated immune responses. Thus, T cell receptors can recognize additional targets not available to antibodies.

A T cell response is modulated by antigen binding to a T cell receptor molecule. One type of TCR is a membrane bound heterodimer consisting of
30 an α chain and a β chain resembling an immunoglobulin variable (V) and constant (C) region. The TCR α chain includes a covalently linked $V\alpha$ and $C\alpha$ chain, whereas the β chain includes a $V\beta$ chain covalently linked to a $C\beta$ chain. The $V\alpha$ and $V\beta$ chains form a pocket or cleft that can bind a superantigen or antigen in the context of a major histocompatibility complex

(MHC) (known in humans as an HLA complex). See generally Davis, *Ann. Rev. of Immunology* **3**: 537 (1985); *Fundamental Immunology 3rd Ed.*, W. Paul Ed. Raven Press LTD. New York (1993).

5 The TCR is believed to play an important role in the development and function of the immune system. For example, the TCR has been reported to mediate cell killing, increase B cell proliferation, and impact the development and severity of various disorders including cancer, allergies, viral infections and autoimmune disorders.

10 Human p53 has been reported to be a tumor suppressor protein and peptide epitopes from p53 are presented by particular class I MHC molecules. p53 has been further reported to be a candidate for a broad-spectrum, tumor-associated cytotoxic T-cell (CTL) target. See, *e.g.*, Theobald, M. *et al.* (1995) *PNAS (USA)* **92**: 11993 and references cited
15 therein.

 There is recognition that abnormal forms of the human p53 protein are associated with a wide variety of cancers. One belief is that the abnormal or mutated version overrides the protective features of normal (wild-type) p53 protein. See, *e.g.*, Levine, A.J. *et al.* (1991) *Nature (London)*
20 **351**: 453.

 Human class I molecules that recognize and specifically bind peptides derived from human p53 protein have been described. One such molecule is HLA-A2.1. See, Theobald, M. *et al.*, *supra*.

 It would be desirable to have TCR molecules that recognize and bind
25 peptides derived from the human p53 protein. It would be especially desirable to have heterodimeric and single chain TCR molecules that specifically bind sequence spanning about amino acid positions 264 to 272 of the human p53 protein.

SUMMARY OF THE INVENTION

30 We have now identified T-cell receptor (TCR) molecules that bind peptides derived from the human p53 protein. In one aspect, we have isolated heterodimeric TCR molecules that specifically bind sequence

preferably spanning about amino acid positions 264 to 272 of the human p53 protein displayed in the context of an HLA molecule, preferably, HLA-A2.1. In another aspect, we have made single chain TCR (sc-TCR) molecules that specifically bind the same sequence. Also disclosed are methods for
5 making and using such TCR molecules. The invention has a wide spectrum of useful applications including therapeutic uses and use in the detection of cells expressing p53 protein.

TCR molecules in accord with the invention are typically heterodimers or single chain molecules that bind sequence preferably
10 spanning between from about amino acid positions 264 to 272 of the human p53 molecule. An especially preferred sequence is the Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1) epitope that spans amino acid positions 264 to 272 of the human p53 molecule. Other suitable p53 sequences are provided below.

15 Particular TCR molecules feature a variety of useful activities. For example, the heterodimeric TCR molecules disclosed herein can be used to detect cell expression of p53 protein, especially in the context of an appropriate antigen-presenting complex. An illustration of such a complex is a primate class I major histocompatibility complex (MHC) that binds and
20 presents to CTLs immunologically relevant fragments of the p53 protein. A preferred class I MHC molecule is the human HLA-A2.1 complex disclosed below.

The invention encompasses a variety of heterodimeric TCR molecules whose context is usually pre-determined to suit an intended use. For
25 example, in one embodiment, heterodimeric TCR molecules are expressed as cell surface molecules on a transfected or genetically engineered recombinant cell. Examples of the cells are provided below. Additionally suitable heterodimeric TCR molecules are provided in a more soluble format, *e.g.*, heterodimers that include one or more immunoglobulin (Ig) sequences as
30 discussed below.

More particular heterodimeric TCR molecules of the invention feature an α chain and a β chain which chains are typically linked together via one or more covalent bonds. Preferably, such covalent bonds include one or

more disulfide linkages. More preferred heterodimers include at least one V α chain and at least one V β chain which chains preferably configure to effectively position, within or near the heterodimer binding cleft, sequence spanning about amino acid positions 264 to 272 of the human p53 molecule, preferably amino acid positions 264 to 272. By the phrase "effectively position" is meant that TCR V chains according to the invention (heterodimer or single chain format) associate to bind a specific p53 sequence as determined by the standard assays disclosed herein including preferred T cell binding and ELISA tests provided below.

10 More specific V chains of the heterodimeric TCRs include the V- α chain linked covalently to a C- α chain and the V- β chain linked to a C- β chain. In most embodiments, the C- α chain and C- β chain are each independently linked to a suitable cell transmembrane domain which domain is typically further linked independently to a suitable cytosolic domain. In instances in which soluble heterodimeric molecules are desired, 15 it may be more preferable to remove at least the transmembrane domain, preferably essentially all of the transmembrane domain using, *e.g.*, standard recombinant DNA manipulations.

The invention features other useful TCR molecules including the single chain T cell receptor (sc-TCR) molecules disclosed herein. Such 20 molecules generally include at least one V α chain bound, by at least one peptide sequence, to at least one V β chain. If desired, the sc-TCR can further include at least one C α chain fragment and optionally at least one C β chain fragment. In more particular invention embodiments, the sc-TCR will include about one V α chain bound to about one V β chain by at least one 25 peptide linker sequence. The arrangement of any V or C sequence in the sc-TCRs is not usually important so long as intended binding results are achieved. However, it is generally preferred that the V α and V β chains be sufficient to effectively bind to the human p53 sequence spanning about amino acid positions 264 to 272 as determined by standard binding tests. 30

The present invention provides important advantages.

For example, the heterodimeric and single chain TCRs provide, for the first time, TCR molecules that recognize and bind an important p53 epitope

sequence. Binding of that sequence by the molecules of the invention provides for important and reliable recognition of p53 tumor suppressor protein in cancerous or pre-cancerous cells. Thus, in one invention aspect, the molecules can be used diagnostically to detect, and quantify if desired, presence and amount of p53 in cells, tissue and organs. Such cells include cultured cells as well as primary, secondary and immortalized cell lines. The ability to detect p53 protein is highly useful as a cancer diagnostic *in vitro* and *in vivo*. Alternatively, the TCR molecules of the invention can be used to detect and optionally quantify p53 expression in cells, particularly those that can present the p53 antigen in the context of a suitable MHC class I molecule, preferably the HLA-A2.1 complex.

Accordingly, and in one aspect, the invention features an isolated T cell receptor (TCR) heterodimer that includes a V α chain and a V β chain. Preferably, the heterodimer is capable of binding, preferably specifically in the context of an HLA-A2.1 MHC molecule, the following "target" amino acid sequence: Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1) including variants of that sequence having at least one conservative amino acid replacement. Preferred binding is determined by any standard TCR binding assay where binding specificity is indicated as an increase in binding which is significantly different from binding to an irrelevant (control) TCR (where "significance" is determined using routine statistical methods known in the art, e.g., with $p \leq 0.05$). Preferably, binding is at least about 2-fold, at least about 10-fold, at least about 20-fold, at least about 50-fold, or at least about 100-fold higher than control values. A specifically preferred TCR binding assay and irrelevant TCR heterodimer are disclosed below.

In a more specific embodiment, the invention features an isolated T-cell receptor (TCR) heterodimer that includes an α -chain and a β -chain in which the α -chain comprises covalently linked in sequence: a) a V α chain and b) a C α chain; and the β -chain comprises covalently linked in sequence: c) a V β chain and a C β sequence. Preferably, the heterodimer is capable of binding, in the context of an HLA-A2.1 MHC molecule, the foregoing target amino acid of SEQ ID NO. 1 as well as variants of that target sequence that have at least one conservative amino acid replacement.

As discussed, the invention also features sc-TCR molecules that include V chains capable of specifically binding the target sequence of SEQ ID NO: 1.

In one embodiment, such sc-TCRs include at least one V α chain covalently linked to at least one V β chain by at least one peptide linker sequence. Preferably, such sc-TCRs include between from about one to five of such V chains, more preferably about one to two of such V chains. Also preferably, such V chains will be linked together by between from about one to five peptide linkers, more preferably about one to two of such linkers. A more preferred sc-TCR is capable of binding, in the context of an HLA-A2.1 MHC molecule, the following target amino acid sequence: Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1) including variants of that sequence having at least one conservative amino acid replacement.

Preferred binding is determined by any standard TCR binding assay where binding specificity is indicated as an increase in binding which is significantly different from binding to an irrelevant (control) TCR (where "significance" is determined using routine statistical methods known in the art, e.g., with $p \leq 0.05$). Preferably, binding is at least about 2-fold, at least about 10-fold, at least about 20-fold, at least about 50-fold, or at least about 100-fold higher than control values. A specifically preferred sc-TCR binding assay and irrelevant sc-TCR are disclosed below.

In another aspect, the invention features at least a pair of nucleic acid segments (typically DNA or RNA) that encode one or more of the heterodimers provided herein.

In another aspect, the invention encompasses a DNA vector that includes at least one of the DNA segments encoding the TCR heterodimers. For example, a first DNA segment can encode the α chain and a second DNA segment can encode the β chain. In some instances, it may be more desirable to provide a single DNA vector with segments that encode both the α and β chains of the heterodimer.

Also envisioned are cells that include the DNA vectors disclosed herein.

The invention also features a nucleic acid segment (DNA or RNA) that encodes at least one, preferably between from about one to five, more preferably about one to two, of the sc-TCR molecules provided herein. Also included are DNA vectors that include the nucleic acid segment.

5 In another aspect, the invention features methods for identifying a cell or tissue expressing p53 protein in the context of an HLA-A2.1 MHC molecule. In one embodiment, such methods include contacting the cell or tissue with a transduced or genetically engineered recombinant cell comprising the sc-TCR or TCR heterodimers disclosed herein. Alternatively,
10 the cell or tissue can be contacted with a soluble sc-TCR or a TCR heterodimer instead of (or in combination with) the transduced or genetically engineered cell expressing the sc-TCR or TCR heterodimer.

The invention also features methods for identifying cells or tissues expressing p53 protein in the context of an HLA-A2.1 MHC molecule. In
15 preferred examples of the invention, the methods include contacting the cell or tissue with a sc-TCR as provided herein.

Also encompassed by the present invention are methods for killing a cell expressing the following target amino acid sequence: Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1) including variants of that sequence
20 having at least one conservative amino acid replacement. More particular methods include contacting the cell with a transduced or recombinant cell expressing a sc-TCR or heterodimeric TCR molecule as provided herein. Additionally preferred methods further include contacting the cell with an amount of sc-TCR or heterodimeric TCR that is generally sufficient to injure
25 or kill the cell as determined by conventional assays (*e.g.*, trypan blue exclusion, presence of apoptotic features, *etc.*).

In another aspect, the present invention features methods for treating cancer that include administering to a mammal a therapeutically effective amount of at least one of: a) a transduced or genetically engineered
30 recombinant cell comprising a TCR heterodimer as provided herein or b) at least one of the sc-TCRs of the invention, preferably one of such sc-TCRs. Preferably, the cancer is characterized by upregulation of p53 protein by at least about two-fold, preferably at least five to 10 fold, preferably about 100

fold as determined by standard immunohistochemistry or flow cytometry as discussed below.

Other aspects and embodiments of the invention are discussed *infra*.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1 is a schematic illustration of the pNAG2 vector.

Figure 2 is a schematic drawing showing the pSUN27 vector (deposited as ATCC#209276).

Figure 3 is a schematic drawing showing regions of the vectors encoding preferred bispecific hybrid molecules pBISP/D011.10 and
10 pBISP/149 (deposited as ATCC #203186 with the designation pSUN28).

Figure 4A, 4B and 4C are drawings showing the amino acid and nucleic acid sequences of the 264 single chain TCR (264 sc-TCR) (SEQ ID NO. 2). V α 3= TCR V3 domain (nucleic acids 61-399); linker sequence (nucleic acids 400-471); V β 3= TCR V3 domain (nucleic acids 472-813);
15 C β = TCR C domain (nucleic acids 814-1191).

Figure 5 is a drawing showing the optional C α domain (SEQ ID NO. 3) of the 264 TCR.

DETAILED DESCRIPTION OF THE INVENTION

As summarized above, we have isolated a highly useful T-cell receptor
20 (TCR) heterodimer that generally includes a V α chain and a V β chain *i.e.*, a two-chain complex. More preferred heterodimers bind, typically in the context of an HLA-A2.1 MHC molecule, amino acid sequence between from about 264 to about 272 of the human p53 protein sequence, preferably spanning amino acid positions 264 to 272 of that protein *i.e.*, the following
25 "target" amino acid sequence: Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1). As also summarized previously, good binding is determined by a standard T-cell binding assay provided below.

The general structure of many naturally-occurring TCR heterodimers
30 has been reported. See, *e.g.*, Davis *Ann. Rev. of Immunology* 3: 537 (1985);

Fundamental Immunology 3rd Ed., W. Paul Ed. Raven Press LTD. New York (1993) and references disclosed therein.

In general, a T cell recognizes antigen presented on the surfaces of cells by means of the T cell receptors expressed on their cell surface. TCRs are disulfide-linked heterodimers, most consisting of α and β chain glycoproteins. T cells use mechanisms to generate diversity in their receptor molecules similar to those mechanisms for generating antibody diversity operating in B cells (Janeway and Travers; Immunobiology 1997). Similar to the immunoglobulin genes, TCR genes are composed of segments that rearrange during development of T cells. TCR polypeptides consist of amino terminal variable and carboxy terminal constant regions. While the carboxy terminal region functions as a transmembrane anchor and participates in intracellular signaling when the receptor is occupied, the variable region is responsible for recognition of antigens. The TCR α chain contains variable regions encoded by V and D segments only, while the β chain contains additional joining (J) segments. The rearrangement of these segments in a diverse repertoire of TCRs capable of recognizing an incredibly large number of different antigens displayed in the context of different MHC molecules.

There have been reports of specific TCRs which recognize particular antigens.

Additionally, particular specific TCRs have been produced by recombinant methods as soluble, single-chain TCRs (sc-TCR). Methods for production and use of sc-TCRs have been disclosed.

Preferred TCR heterodimers of the invention include an α chain and a β chain covalently linked together by virtue of at least one disulfide bond. Non-covalent binding, *e.g.*, hydrogen bonding, between the chains has been reported. Each of the chains can be between from about 150 to about 350 amino acids long, preferably between from about 200 to about 300 amino acids long, more preferably between from about 250 to about 290 amino

acids long, with about 280 amino acids being useful for most invention applications. Heterodimeric TCR molecules according to the invention are optionally glycosylated.

By the phrase "HLA-A2.1 MHC molecule" is meant a primate class I
5 MHC molecule, preferably a human molecule, that is capable of generating or of being recognized by A2.1 restricted, tumor-reactive cytotoxic T lymphocytes (CTLs) bearing TCRs specific for peptides having sequence obtained from human p53 protein. A preferred amino acid sequence is usually between from about amino acid 250 to about amino acid 290 of the
10 human p53 sequence, preferably between from about amino acid 264 to about amino acid 272 of that sequence, with sequence spanning positions amino acid 264 to amino acid 272 of the p53 protein being preferred for most applications.

More preferred HLA-A2.1 MHC molecules in accord with the invention
15 are integral membrane proteins that often include a glycoprotein heavy chain having three extracellular domains (*i.e.* $\alpha 1$, $\alpha 2$ and $\alpha 3$), a transmembrane domain and a cytoplasmic domain. The heavy chain is typically non-covalently associated with a soluble subunit $\beta 2$ -microglobulin. The $\alpha 1$ and $\alpha 2$ domains of the heavy chain fold together to form the peptide-
20 binding groove for a particular p53 sequence. The association between the heavy chain and $\beta 2$ -microglobulin may help stabilize the peptide-binding groove. The MHC molecule may consist of nearly any combination of a naturally occurring or recombinant class I heavy chain (or fragments thereof) and a naturally occurring or recombinant $\beta 2$ - microglobulin
25 molecule (or biologically active fragments thereof).

Information relating to the human p53 amino acid and nucleic acid sequence is available from the National Center for Biotechnology Information (NCBI)- Genetic Sequence Data Bank (Genbank) at the National Library of Medicine, 38A, 8N05, Rockville Pike, Bethesda, MD 20894.
30 Genbank is also available on the internet at <http://www.ncbi.nlm.nih.gov>. See, Benson, D.A. *et al.* (1997) *Nucl. Acids. Res.* **25**: 1 for a description of Genbank. See also Theobald, M *et al.* (1995), *supra* (disclosing also the p53 amino acid numbering scheme adopted in this application).

It has been reported that expression of the tumor suppressor protein p53 is upregulated on malignant cells. It has also been shown that 50% of all tumors express increased levels of p53 on their surface (Holliston, M.D., *et al.*, *Science* (1991), **253**: 49).

5 Information relating to making and using the human HLA-A2.1 MHC molecule, particularly in the context of tumor cells expressing p53 has been reported by Theobald, M *et al.* (1995), *supra*, including references disclosed therein.

10

Methods for detecting productive binding between the HLA-A2.1 MHC molecule and an amino acid sequence obtained from the p53 protein
 15 sequence have been reported, *e.g.*, by Theobald, M *et al.* (1995), *supra*. In general, the methods involve using recognized competition assays to assess binding of p53 peptide to the HLA-A2.1 molecule.

More particular heterodimeric TCR molecules in accord with the
 20 invention include a Va chain that is at least about 90% identical to the Va3 chain shown in Figures 4A-C below, preferably between from about 95% to about 100% identical. Additional heterodimers of the invention include a V β chain that is at least about 90% identical to the Vb3 chain shown in Figure 4A-C (SEQ ID NO. 2), preferably between from about 95% to about 100%
 25 identical.

Preferably, to determine the percent identity of two amino acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps are introduced in one or both of a first and a second amino acid for optimal alignment and non-homologous sequences are disregarded for
 30 comparison purposes). A "comparison window" refers to a segment of any one of the number of contiguous positions selected from the group consisting of from 25 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a

reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. To identify sequences with the appropriate % identity as disclosed herein, the comparison window may comprise any of the segment ranges described above.

5 The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap which need to be introduced for optimal alignment of the two sequences. The amino acid residues at corresponding amino acid positions are then compared. When a position in
10 the first sequence is occupied by the same amino acid residue as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid "identity" is equivalent to amino acid "homology").

Percent identity between two sequences can be determined using a
15 mathematical algorithm as is known in the art (see, *e.g.*, Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994;
20 Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). For example, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch algorithm (J. Mol. Biol. (48): 444-453, 1970) which is part of
25 the GAP program in the GCG software package

by the local homology algorithm of Smith & Waterman (Adv. Appl. Math. 2: 482, 1981), by the search for similarity methods of Pearson & Lipman (Proc. Natl. Acad. Sci. USA 85: 2444, 1988) and Altschul, et al. (Nucleic Acids Res. 25(17): 3389-3402, 1997), by
30 computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and BLAST in the Wisconsin Genetics Software Package (available from, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection (see, *e.g.*, Ausubel *et al.*, *supra*).

Gap parameters can be modified to suit a user's needs. For example, when employing the GCG software package, a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6 can be used. Exemplary gap weights using a Blossom 62 matrix or a
 5 PAM250 matrix, can be 16, 14, 12, 10, 8, 6, or 4, while exemplary length weights can be 1, 2, 3, 4, 5, or 6. The percent identity between two amino acid also can be determined using the algorithm of E. Myers and W. Miller (CABIOS 4: 11-17, 1989) which has been incorporated into the ALIGN
 10 program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

Thus, by the term "100% identical" is meant that the amino acids of a subject V chain are 100% homologous to the corresponding naturally-occurring TCR V β or V α chain or allelic variants thereof with the same binding characteristics (*e.g.*, no significant difference in binding specificity
 15 and affinity]. That is, the subject V chain has the same length and amino sequence of the corresponding and naturally-occurring chain or allelic variants thereof with the same binding characteristics.

Additionally preferred heterodimeric molecules of the invention include a C α and C β sequence having at least about 90% identity to the C α and C β chains sequences shown in Figures 5 [SEQ ID NO. 3] and 4A-C (SEQ
 20 ID NO. 2), respectively. Preferably, the C α and C β sequences are between from about 95% to about 100% identical to the C α and C β chains sequences shown in Figures 5 and 4A-C.

The general structure of sc-TCR molecules and methods of making
 25 and using same have been disclosed.

30

As provided above, the present invention features highly useful single-chain T cell receptor (sc-TCR) proteins that generally include between

from about one to about five V α chains covalently linked to between from about one to five V β chains by between from about one to five peptide linker sequences. Preferred sc-TCR include about one V α chain and about one V β chain linked together by about one peptide linker sequence as provided
5 herein.

Additionally preferred sc-TCRs are typically capable of binding, in the context of the HLA-A2.1 MHC molecule, peptides having sequence obtained from human p53 protein. A preferred amino acid sequence is usually between from about amino acid 250 to about amino acid 290 of the human
10 p53 sequence, preferably between from about amino acid 264 to about amino acid 272 of that sequence, with sequence spanning amino acid positions 264 to 272 of the p53 protein being preferred for most applications. Good binding is preferably determined by the standard T cell receptor (TCR) ELISA assay described below.

15 More specific sc-TCR molecules in accord with the invention include a V α chain that is at least about 90% identical to the Va3 chain shown in Figures 4A-C (SEQ ID NO. 2) below, preferably between from about 95% to about 100% identical. Additionally preferred sc-TCR molecules include a V β chain that is at least about 90% identical to the Vb3 chain shown in Figures
20 4A-C (SEQ ID NO. 2), preferably between from about 95% to about 100% identical.

Additionally preferred sc-TCR molecules of the invention include a C β sequence having at least about 90% identity to the C β chain sequence shown in Figures 4A-C. Preferably, the C β sequence is between from about
25 95% to about 100% identical to the C β chain sequence shown in Figures 4A-C.

It has been discovered that the Ca chain is not always required to demonstrate good sc-TCR binding in the standard T cell receptor (TCR) ELISA assay. In these embodiments, it is not necessary to include the Ca
30 chain as part of the sc-TCR molecule. For example, see Figures 4A-5C below (disclosing especially preferred 264 sc-TCR sequences). However, sc-TCR molecules may include at least one Ca chain (as shown in Figure 5, for

example) or a functional fragment thereof, preferably between from about one to five of such chains, with about one of such a Ca chain being suitable.

In invention embodiments in which a particular sc-TCR includes the Ca chain or a functional fragment thereof, that chain will preferably bear at
5 least about 90% identity to the Ca chain sequence shown in Figure 5 (SEQ ID NO. 3). Preferably, that sequence is between from about 95% to about 100% identical to the Ca chain sequence shown in Figure 5 (SEQ ID NO. 3).

More particular sc-TCR molecules in line with the invention include those having covalently linked in sequence: 1) a Va3 chain as shown in
10 Figures 4A-C (SEQ ID NO. 2); 2) a peptide linker; and 3) a Vβ3 chain as shown in Figures 4A-C (SEQ ID NO. 2). In one embodiment, the sc-TCR molecule further includes a Cβ chain as provided in Figures 4A-C (SEQ ID NO. 2) preferably linked to the C-terminus of the Vβ3 chain.

In an embodiment of the foregoing specific sc-TCR molecule, the sc-
15 TCR further includes the Ca chain as provided in Figure 5 (SEQ ID NO. 3), the chain preferably being covalently linked between the C-terminus of the Va chain and the N-terminus of a peptide linker.

Typical Vα and Vβ chains of the heterodimeric and single chain TCR molecules disclosed herein are generally about 200 to 400 amino acids in
20 length, preferably about 300 to 350 amino acids in length. Methods for determining amino acid length are known in the field and include polyacrylamide gel electrophoresis.

As discussed, preferred sc-TCR molecules of the invention include one or more peptide linker sequences preferably positioned between the Va
25 and Vβ chains. Preferably the linker sequence comprises from about 7 to 20 amino acids, more preferably from about 8 to 16 amino acids. The linker sequence is preferably flexible so as not hold sequence derived from the human p53 protein (and presented in the context of the HLA-A2.1 molecule) in a single desired conformation. Specifically, the peptide linker sequence
30 can be positioned between the TCR variable chains typically to enhance binding flexibility between those chains. The linker predominantly comprises amino acids with small side chains, such as glycine, alanine and serine, to provide for flexibility. Preferably about 80 or 90 percent or greater

of the linker sequence comprises glycine, alanine or serine residues, particularly glycine and serine residues. For heterodimeric TCRs, the linker sequence is suitably linked to the β chain of the TCR molecule, although the linker sequence also could be attached to the α chain of the TCR molecule.

5 Alternatively, the linker sequence may be linked to both α and β chains of the TCR molecule.

See the following references for supplemental disclosure relating to making and using sc-TCR molecules: Novotny, J. *et al.* *PNAS* (USA) **88**: 8646 (1991); Soo Hoo, W.F. *et al.* *PNAS* (USA) **89**: 4759 (1992); Wülfing, C. and Plückerthun, A., *J. Mol. Biol.* **242**: 655 (1994); Kurucz, I. *et al.* *PNAS* (USA) **90**: 3830 (1993); PCT WO 96/13593; Ward, E.S. *et al.*, *J. Mol. Biol.* **224**: 885, (1992); Schlueter, C.J. *et al.* *J. Mol. Biol.* **256**: 859 (1996); Mariuzza, R.A. and Winter, G., (1989) **264**:7310; Gascoigne, N.R.J., *et al.*, *PNAS* (USA) (1987), **84**: 2936.

10

15 In a particular invention embodiment, a suitable linker sequence is ASGGGGSGGG (*i.e.*, Ala Ser Gly Gly Gly Gly Ser Gly Gly Gly) (SEQ ID NO. 4] repeated as many as four or more times, preferably linked to the first amino acid of the β domain of the TCR. Different linker sequences could be used including any of a number of flexible linker designs that have been used

20 successfully to join antibody variable regions together, see Whitlow, M. *et al.*, (1991) *Methods: A Companion to Methods in Enzymology* **2**:97-105. Suitable linker sequences can be readily identified empirically. Additionally, suitable size and sequences of linker sequences also can be determined by conventional computer modeling techniques based on the predicted size and

25 shape of the TCR molecule.

Accordingly, and in one embodiment, the invention features particular sc-TCR molecules in which at least one of the peptide sequences, preferably one of same has the following sequence: Gly Gly Gly Gly Ser repeated as many as four or more times (SEQ ID NO. 5).

30 See also the co-pending application entitled *T-cell Receptor Fusions and Conjugates and Methods of Use Thereof* filed on June 5, 2001 (USSN 09/874,907 by inventors Jon A. Weidanz, Kimberlyn F. Card, and Hing C. Wong) for additional information relating to particular 264 TCR related

molecules.

In some settings it can be useful to make the sc-TCR molecules of the present invention polyvalent, *e.g.*, to increase the valency of the sc-TCR.

5 Briefly stated, the polyvalent TCR protein is made by covalently linking together between two and four proteins (the same or different) by using *e.g.*, standard biotin-streptavidin labeling techniques, or by conjugation to suitable solid supports such as latex beads. Chemically cross-linked proteins (for example cross-linked to dendrimers) are also suitable

10 polyvalent species. For example, the protein can be modified by including sequences encoding amino acid residues with chemically reactive side chains such as Cys or His. Such amino acids with chemically reactive side chains may be positioned in a variety of positions in the linked protein, preferably distal to the antigen-binding region of the TCR. For example, the

15 C-terminus of a C β chain fragment of a soluble protein can be covalently linked to a protein purification tag or other protein which includes such a reactive amino acid(s). Suitable side chains can be included to chemically link two or more proteins to a suitable dendrimer particle to give a multivalent molecule. Dendrimers are synthetic chemical polymers that can

20 have any one of a number of different functional groups of their surface (D. Tomalia, *Aldrichimica Acta*, **26**:91-101 (1993)). Exemplary dendrimers for use in accordance with the present invention include, *e.g.*, E9 starburst polyamine dendrimer and E9 combust polyamine dendrimer, which can link cysteine residues.

25 Successful presentation of a human p53 sequence as provided herein to a TCR molecule of the invention can be determined by a variety of specified assays, including the T cell binding assays and TCR ELISA discussed below. Alternatively, successful presentation can be detected and quantified if desired by monitoring the activity of a T cell by following either

30 induction of or inhibition of T cell proliferation, or initiation or inhibition of an immune response to a particular site or target. Such suitable assays include, but are not limited to, *in vitro* assays that include sequential steps of culturing T cells to proliferate same, and contacting the T cells with a MHC-peptide antigen complex and then evaluating biological response by

the cells. See the USSN 08/813,781 and PCT/US98/04274 applications for more specific examples of such assays.

In one aspect, the functionality of a TCR molecule is determined by monitoring the ability of the TCR to recognize the appropriate p53 peptide in the context of an appropriate MHC molecule (*e.g.*, HLA-2A), *e.g.*, by
5 monitoring binding of the TCR to MHC:p53 peptide complexes. Such complexes can be presented on a cell, in which case TCR functionality is measured by contacting a labeled TCR with a p53 presenting cell and measuring binding to the cell as compared to binding to a non-p53
10 presenting cell. Labeled cells can be detected microscopically or by using flow cytometry assays as are routine in the art.

In another aspect, a non-cell based assay is used, such as a TCR Enzyme Linked Immunosorbant Assay (ELISA). For example, the TCR can be bound directly to a support and its ability to bind to MHC:p53 peptide
15 complexes can be measured, or the MHC:p53 complex can be bound to the support and the ability of the complex to bind to the TCR can be measured. Suitable supports include, but are not limited to, wells of a microtiter plate, cell culture plates, smembrane, glass or polymer substrates, and the like. Instead of direct binding of the TCR to the support, the support can be
20 coated with an antibody that recognizes the TCR such that the bound antibody can capture and thereby indirectly bind the TCR to the support. Suitable controls for such assays will be obvious to those of skill in the art, and include, but are not limited to, MHC molecules bound to irrelevant antigens, non-p53 recognizing TCR's, buffer, *etc.* In a TCR ELISA, either the
25 TCR molecule or the MHC or peptide can be labeled. Preferably, the molecule bound to the support is unlabeled. As used herein, "labeled" refers to direct or indirect labeling. Thus, a "labeled TCR molecule" may comprise a label directly linked to it or may comprise a TCR bound indirectly or directly by a labeled binding partner, such as an antibody,
30 which recognizes the TCR or which recognizes an antibody bound to the TCR. As used herein, "linked" refers to a stable association between two molecules which can be covalent or non-covalent.

Assays to monitor TCR functionality may also include assays to measure TCR-mediated signal transduction. In one aspect, a nucleic acid

construct encoding a TCR heterodimer is introduced into a cell which does not express a TCR or at least does not express a TCR of the same specificity. The ability of the TCR-expressing cell to transduce the appropriate signals upon binding to a p53:MHC complex can then be monitored. For example, the ability of the TCR-expressing cell to produce IL-2 may be measured. sc-TCRs also may be transfected into cells. In such assays, the sc-TCR is preferably expressed as a fusion with a transmembrane domain polypeptide (*e.g.*, from an immunoglobulin molecule) and more preferably, also as a fusion with an appropriate cytoplasmic signaling domain. In one aspect, the cytoplasmic signaling domain is a CD3 zeta molecule. MHC:p53 complexes may be presented by natural or engineered antigen presenting cells or may be isolated complexes.

The ability of a TCR to mediate a cytolytic response also can be determined. For such an assay, a nucleic acid construct encoding a TCR molecule preferably is introduced into a cell which can express appropriate co-stimulatory molecules.

In all of the above assays, a "functional" TCR is one which demonstrates increased function (*e.g.*, increased binding, increased signal transduction, such as IL-2 production, increased cell killing, and the like] as compared to a control TCR which does not bind to an MHC:p53 complex. The amount of increased function necessary to demonstrate a "functional TCR" will necessarily depend on the type of assay used. For example, in one aspect, an assay value which indicates a functional TCR is about 10% greater, about 15% greater, about 20% greater, about 30% greater, about 40% greater, about 50% greater, about 60% greater, about 70% greater, about 80% greater, about 90% greater or about 100% greater than a value obtained for a control TCR. In other assays, an assay value which indicates a functional TCR is about 2-fold greater, about 4-fold greater, about 8 fold greater, about 10-fold greater, about 20-fold greater, about 30-fold greater, about 40-fold greater, about 50-fold greater, about 60-fold greater, about 70-fold greater, about 80-fold greater, about 90-fold greater, or about 100-fold greater than a value obtained for a control TCR. For other assays, an assay value which indicates a functional TCR is one which is statistically significantly different from a value obtained from a control assay with $p <$

0.05. One of skill in the art can routinely evaluate measures of significance for particular assays used.

In general, preparation of the TCR of the invention can be accomplished by procedures disclosed herein and by recognized
5 recombinant DNA techniques involving, *e.g.*, polymerase chain amplification reactions (PCR), preparation of plasmid DNA, cleavage of DNA with restriction enzymes, preparation of oligonucleotides, ligation of DNA, isolation of mRNA, introduction of the DNA into a suitable cell, transformation or transfection of a host, culturing of the host. Additionally,
10 the TCR molecules can be isolated and purified using chaotropic agents and well known electrophoretic, centrifugation and chromatographic methods. See generally, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd ed. (1989); and Ausubel *et al.*, *Current Protocols in Molecular Biology*, John Wiley & Sons, New York (1989) for disclosure relating to these methods.

15 See also the USSN 08/813,781 and PCT/US98/04274 and the co-pending application referenced above entitled *T-cell Receptor Fusions and Conjugates and Methods of Use Thereof* filed on June 5, 2001 (USSN 09/874,907 by inventors Jon A. Weidanz, Kimberlyn F. Card, and Hing C. Wong) for more specific background information relating to making and
20 using the molecules disclosed herein.

As discussed, preferred molecules in accord with the invention demonstrate good binding in what is referred to herein as a standard T cell binding assay or TCR ELISA. By the phrase "standard T cell binding assay" is meant a binding test that detects and preferably quantitates binding
25 between a suitable T cell and an MHC molecule complexed with an antigen. Briefly stated, a preferred test involves providing a detectably labeled MHC molecule, contacting the labeled MHC molecule (complexed with the antigen) and the T cell under conditions conducive to forming an MHC-antigen:T cell complex and monitoring formation of that complex using
30 standard detection methods. Formation of the complex can be quantified if desired. Other assays for TCR function as described above may also be used.

Preferred TCR molecules including the heterodimers and single chain molecules provided herein are generally of sufficient size to allow for specific binding of the TCR to the MHC molecule. In embodiments where the MHC is complexed with an antigen also referred to as a peptide-MHC molecule, the TCR molecules contain at least the CDR binding loops forming the MHC-peptide binding pocket. Useful α/β TCR molecules containing an MHC-peptide binding pocket preferably consists of at least the α chain variable domain (about amino acid 1 to about amino acid 110 to about amino acid 130 dependent on CDR length of the α chain) and the β chain variable domain (about amino acid 1 to about amino acid 110 to about amino acid 130 dependent on CDR length of the β chain).

More preferred TCR molecules for use in accord with this invention exhibit significant binding activity in what has been referred to herein as the standard T-cell binding test. Preferably, the TCR molecule specifically binds its cognate MHC antigen molecule complex at a level which is significantly different from binding to an irrelevant (control) TCR (where "significance" is determined using routine statistical methods known in the art, *e.g.*, with $p \leq 0.05$). Preferably, binding is at least about 2-fold, at least about 10-fold, at least-about 20-fold, at least about 50-fold, or at least about 100-fold higher than control values. Examples of suitable control molecules include a 149 TCR molecule.

Highly useful *in vitro* and *in vivo* T cell binding assays have been disclosed.

The disclosed T cell binding assays can be used or readily adapted if necessary to identify good binding between TCR molecules of this invention and the disclosed p53 amino acid sequence displayed in the context of an appropriate MHC molecule.

5

10

In particular, Example 15 of the co-pending application filed on May 16, 2001 discloses an illustration of the standard T cell binding test. Typically, the test involves producing T cells that express the subject TCR of interest, *e.g.*, a heterodimer in accord with the invention; and then staining those cells with a suitable class I MHC molecule, particularly the HLA-A2.1 molecule. Methods for staining the T cells involving conventional biotin/streptavidin technologies have been disclosed in the co-pending application filed on May 16, 2001. As disclosed, a preferred detection format is flow cytometry although other detection strategies may be more preferred for some applications.

By the phrase "standard T cell receptor (TCR) ELISA" is meant to include, but is not limited to, any one of the suitable assays disclosed, *e.g.*, in the foregoing co-pending application filed on May 16, 2001. A preferred assay involves manipulating single chain or heterodimeric TCR constructs using, *e.g.*, a plate-based ELISA. Briefly stated, the assay involves detectably labeling the single-chain or heterodimeric TCR, contacting the labeled TCR molecule with a suitable peptide-loaded MHC molecule, preferably the HLA-A2.1 molecule disclosed herein, in which the contacting is under conditions sufficient to form a TCR:MHC-peptide complex. Preferred labeling strategies are disclosed throughout the co-pending application filed on May 16, 2001 and include standard biotin/streptavidin labeling strategies. See, for instance, Example 15 of the co-pending application filed on May 16, 2001.

As discussed, preferred TCR molecules of the invention bind, typically in the context of an HLA-A2.1 MHC molecule, amino acid sequence between from about amino acid 264 to about amino acid 272 of the human p53 protein sequence, preferably spanning amino acid positions 264 to 272 of that protein, *i.e.*, the following "target" amino acid sequence: Leu Leu Gly Arg Asn Ser Phe Glu Val (SEQ ID NO. 1). Additionally contemplated are derivatives of the target amino acid, *i.e.*, amino acid sequences having at least one conservative amino acid substitution therein. In embodiments in which two or more conservative amino acids are substituted for any of the target sequence residues, those substitutions can be adjacent or non-adjacent as needed.

Preferably, conservative substitutions are amino acid substitutions that are phenotypically silent, *i.e.*, the substitutions do not significantly affect the binding of the TCR in a standard assays. An example of a conservative amino acid substituted for another amino acid is a tyrosine amino acid substituted for the phenylalanine at amino acid position 7 (amino acid position 270 with respect to human p53 protein) of the preferred target sequence. In contrast, an arginine substituted for any of the leucine residues in the target sequence would be an example of a non-conservative amino acid substitution. Preferred examples of conservative amino acid replacements have been disclosed in the U.S. Pat. No. 6,127,524 (Figures 15A-B); the disclosure of which is incorporated herein by reference.

The invention further provides nucleic acid sequences (DNA or RNA) and particularly DNA sequences that encode the present TCR molecules including preferred heterodimers and single chain constructs. Such DNA sequences are preferably carried by a vector suited for extrachromosomal replication such as a phage, virus, plasmid, phagemid, cosmid, YAC, or episome. In some embodiments, the DNA vector can encode another helper protein whose sole function is to facilitate preparative methods described herein and to obtain significant quantities of the protein. The DNA sequence can be inserted into an appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. A variety of host-vector systems may be utilized to express the protein-coding sequence. These include

mammalian cell systems infected with virus (*e.g.*, vaccinia virus, adenovirus, etc.) or transfected with an expression vector; insect cell systems infected with virus (*e.g.*, baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage DNA, plasmid
5 DNA or cosmid DNA. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used. See generally Sambrook *et al.*, *supra* and Ausubel *et al.* *supra*.

In general, a preferred DNA vector according to the invention comprises a nucleotide sequence linked by phosphodiester bonds
10 comprising, in a 5' to 3' direction a first cloning site for introduction of a first nucleotide sequence encoding a TCR chain, operatively linked to a sequence encoding an effector molecule, *i.e.* a fusion protein or conjugate.

As used herein, an "effector molecule" refers to an amino acid sequence such as a protein, polypeptide or peptide; a sugar or
15 polysaccharide; a lipid or a glycolipid, glycoprotein, lipoprotein or chemical agent that can produce the desired effects as discussed herein. Thus, suitable molecules include regulatory factors, enzymes, antibodies, or drugs as well as DNA, RNA, and oligonucleotides. The biologically active or effector molecule can be naturally-occurring or it can be synthesized from
20 known components, *e.g.*, by recombinant or chemical synthesis and can include heterologous components. A biologically active or effector molecule is generally between about 0.1 to 100 KD or greater up to about 1000 KD, preferably between about 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 30 and 50 KD as judged by standard molecule sizing techniques such as centrifugation or
25 SDS-polyacrylamide gel electrophoresis. Desired effects of the invention include, for example, either to induce cell proliferation or cell death, initiate an immune response or to act as a detection molecule for diagnostic purposes as determined by the assays disclosed below, including an assay that includes sequential steps of culturing cells to proliferate same, and
30 contacting the cells with a TCR fusion complex of the invention and then evaluating whether the TCR fusion complex inhibits further development of the cells.

In most instances, it will be preferred that each of the fusion protein components encoded by the DNA vector be provided in a "cassette" format.

By the term "cassette" is meant that each component can be readily substituted for another component by standard recombinant methods. To make the vector coding for the TCR molecules provided herein, the sequence coding for the TCR molecule is linked to a vector sequence by use of suitable
5 ligases.

If desired, other nucleotide sequences can be included in the gene construct. For example, a promoter sequence, which controls expression of the sequence coding for the TCR molecule, or a leader sequence, which directs the TCR fusion complex to the cell surface or the culture medium,
10 can be included in the construct or be present in the expression vector into which the construct is inserted. An immunoglobulin or CMV promoter is particularly preferred for mammalian cell expression.

It is emphasized that components of the TCR molecules of this invention, including, but not limited to, variable chains, transmembrane
15 domains, constant chains, *etc.*, can be organized in nearly any order provided each is capable of performing its intended function.

A number of strategies can be employed to express TCR molecules provided herein. For example, the sc-TCR molecule binding the target amino acid sequence of SEQ ID NO: 1 in the context of an HLA molecule can
20 be incorporated into a suitable vector by known means such as by use of restriction enzymes to make cuts in the vector for insertion of the construct followed by ligation. The vector containing the recombinant gene construct is then introduced into a suitable host for expression of the TCR fusion peptide. See, generally, Sambrook *et al.*, *supra*. Selection of suitable
25 vectors can be made empirically based on factors relating to the cloning protocol. For example, the vector should be compatible with, and have the proper replicon for, the host that is being employed. Further, the vector must be able to accommodate the DNA sequence coding for the TCR molecule that is to be expressed. Suitable host cells include eukaryotic and
30 prokaryotic cells, preferably those cells that can be easily transformed and exhibit rapid growth in culture medium. Specifically preferred host cells include prokaryotes such as *E. coli*, *Bacillus subtilis*, *etc.* and eukaryotes such as animal cells and yeast strains, *e.g.*, *S. cerevisiae*. Mammalian cells are generally preferred, particularly J558, NSO, SP2-O or CHO. Other

suitable hosts include, *e.g.*, insect cells such as Sf9. Conventional culturing conditions are employed. See Sambrook, *supra*. Stable transformed or transfected cell lines can then be selected. Cells expressing a TCR molecule can be determined by known procedures. For example, expression of a TCR molecule, preferably a heterodimer linked to an immunoglobulin can be determined by an ELISA specific for the linked immunoglobulin and/or by immunoblotting.

As mentioned generally above, a host cell can be used for preparative purposes to propagate nucleic acid encoding a desired fusion protein. Thus, a host cell can include a prokaryotic or eukaryotic cell in which production of the fusion protein is specifically intended. Thus, host cells specifically include bacterial, yeast, fly, worm, plant, frog, mammalian cells and organs that are capable of propagating nucleic acid encoding the fusion. Non-limiting examples of mammalian cell lines which can be used include CHO dhfr- cells (Urlaub and Chasm, *Proc. Natl. Acad. Sci. USA*, **77**:4216 (1980)), 293 cells (Graham *et al.*, *J Gen. Virol.*, **36**:59 (1977)) or myeloma cells like SP2 or NSO (Galfre and Milstein, *Meth. Enzymol.*, **73**(B):3 (1981)).

Host cells capable of propagating nucleic acid encoding a desired heterodimer or single chain TCR encompass non-mammalian eukaryotic cells as well, including insect (*e.g.*, *Sp. frugiperda*), yeast (*e.g.*, *S. cerevisiae*, *S. pombe*, *P. pastoris*, *K. lactis*, *H. polymorpha*; as generally reviewed by Fleer, R., *Current Opinion in Biotechnology*, **3**(5):486-496 (1992)), fungal and plant cells. Also contemplated are certain prokaryotes such as *E. coli* and *Bacillus*.

Nucleic acid encoding a desired fusion protein can be introduced into a host cell by standard techniques for transfecting cells. The term "transfecting" or "transfection" is intended to encompass all conventional techniques for introducing nucleic acid into host cells, including calcium phosphate co-precipitation, DEAE-dextran-mediated transfection, lipofection, electroporation, microinjection, viral transduction and/or integration. Suitable methods for transfecting host cells can be found in Sambrook *et al. supra*, and other laboratory textbooks.

The present invention further provides a production process for isolating any one of the TCR molecules disclosed herein. In the process, a host cell (*e.g.*, a yeast, fungus, insect, bacterial or animal cell), into which has been introduced a nucleic acid encoding the protein of interest
5 operatively linked to a regulatory sequence, is grown at production scale in a culture medium to stimulate transcription of the nucleotide sequence encoding the fusion protein of interest. Subsequently, the TCR molecule is isolated from harvested host cells or from the culture medium. Standard protein purification techniques can be used to isolate the protein of interest
10 from the medium or from the harvested cells. In particular, the purification techniques can be used to express and purify a desired TCR protein on a large-scale (*i.e.*, in at least milligram quantities) from a variety of implementations including roller bottles, spinner flasks, tissue culture plates, bioreactors, or fermentors.

15 An expressed TCR molecule according to the invention can be isolated and purified by known methods. Typically the culture medium is centrifuged and then the supernatant is purified by affinity or immunoaffinity chromatography, *e.g.*, Protein-A or Protein-G affinity chromatography or an immunoaffinity protocol comprising use of
20 monoclonal antibodies that bind the expressed TCR molecule. Such molecules can be separated and purified by appropriate combination of known techniques. These methods include, for example, methods utilizing solubility such as salt precipitation and solvent precipitation, methods utilizing the difference in molecular weight such as dialysis, ultra-filtration,
25 gel-filtration, and SDS-polyacrylamide gel electrophoresis, methods utilizing a difference in electrical charge such as ion-exchange column chromatography, methods utilizing specific affinity such as affinity chromatography, methods utilizing a difference in hydrophobicity such as reverse-phase high performance liquid chromatography, and methods
30 utilizing a difference in isoelectric point such as isoelectric focusing electrophoresis and metal affinity columns such as Ni-NTA. See, generally, Sambrook *et al.* and Ausubel *et al. supra* for disclosure relating to these methods.

It is preferred that particularly the single chain TCR molecules of the present invention be substantially pure. That is, the molecules have been isolated from cell constituents that naturally accompany it so that the fusion proteins are present preferably in at least 80% or 90% to 95%
5 homogeneity (w/w). Such proteins having at least 98 to 99% homogeneity (w/w) are most preferred for many pharmaceutical, clinical and research applications. Once substantially purified, the protein should be substantially free of contaminants for therapeutic applications. Once purified partially or to substantial purity, the soluble TCR molecules,
10 preferably in a single chain format, can be used therapeutically, or in performing *in vitro* or *in vivo* assays as disclosed herein. Substantial purity can be determined by a variety of standard techniques such as chromatography and gel electrophoresis.

Truncated TCR molecules of the invention contain a TCR molecule
15 that is sufficiently truncated so the TCR molecule of the invention can be secreted into culture medium after expression. Thus, a truncated TCR molecule, sc-TCR, TCR fusion or complex will typically not include regions rich in hydrophobic residues, typically the transmembrane and cytoplasmic domains of the TCR molecule. Thus, for example, for a preferred truncated
20 TCR molecule of the invention, preferably from about amino acid residues 199 to 237 of the β chain and from about amino acid residues 193 to 230 of the α chain of the TCR molecule are not included in the truncated TCR complex.

By the term "soluble" or similar term is meant that a TCR molecule of
25 the invention, usually but not exclusively a single chain construct, is not readily sedimented under low G-force centrifugation (*e.g.* less than about 30,000 revolutions per minute in a standard centrifuge) from an aqueous buffer, *e.g.*, cell media. Further, the molecule is soluble if it remains in aqueous solution at a temperature greater than about 5-37°C and at or near
30 neutral pH in the presence of low or no concentration of an anionic or non-ionic detergent. Under these conditions, a soluble protein will often have a low sedimentation value, *e.g.*, less than about 10 to 50 Svedberg units.

Aqueous solutions referenced herein typically have a buffering compound to establish pH, typically within a pH range of about 5-9, and an

ionic strength range between about 2mM and 500mM. Sometimes a protease inhibitor or mild non-ionic detergent is added. Additionally, a carrier protein may be added if desired such as bovine serum albumin (BSA) or human serum albumin (HSA) to a few mg/ml. Exemplary aqueous
5 buffers include standard phosphate buffered saline, Tris-buffered saline, or other well-known buffers and cell media formulations.

The present TCR molecules are suitable for *in vitro* or *in vivo* use with a variety of cells that are cancerous, pre-cancerous, or tumorigenic. Preferably, such cells express high levels of p53 protein when compared to
10 normal (wild-type) cells that are not known to be cancerous, pre-cancerous, or tumorigenic.

Molecules of the invention will be especially useful to a human patient who has or is suspected of having a malignant disease, disorder or condition associated with abnormal expression of p53 (*e.g.*, an at least two-
15 fold overexpression of the molecule). For example, molecules of the invention or derivatives thereof will be particularly useful in targeting tumors in human patients associated with abnormal expression of p53. Specific examples of diseases which may be treated in accordance with the invention include cancers, *e.g.*, breast, prostate, *etc.*, well as other specific
20 disorders or conditions mentioned herein.

Administration of molecules of the invention may be made by a variety of suitable routes including oral, topical (including transdermal, buccal or sublingal), nasal and parenteral (including intraperitoneal, subcutaneous, intravenous, intradermal or intramuscular injection) with
25 oral or parenteral being generally preferred. It also will be appreciated that the preferred method of administration and dosage amount may vary with, for example, the condition and age of the recipient. Effective dosages may be monitored by determining standard clinical therapeutic endpoints such as tumor regression, decrease in expression of cancer-specific markers
30 (including p53), decreased cell proliferation, improved or normal biopsy results, and the like.

Molecules of the invention may be used in therapy alone or in conjunction with other medicaments such as those with recognized

pharmacological activity to treat the desired indications. Exemplary medicaments include recognized therapeutics such as surgery, radiation, chemotherapy and other forms of immunotherapy (*e.g.*, vaccines, antibody-based therapies). The molecule of this invention can be administered
5 before, during or after such therapies as needed.

While one or more molecules of the invention may be administered alone, they also may be present as part of a pharmaceutical composition in mixture with conventional excipient, *i.e.*, pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, oral or other
10 desired administration and which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof. Pharmaceutical compositions of the invention in general comprise one or more TCR molecules of the invention or DNA constructs coding for such TCR molecules together with one or more acceptable carriers. The carriers must
15 be "acceptable" in the sense of being compatible with other ingredients of the formulation and not deleterious to the recipient thereof. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil,
20 fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, *etc.* The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or
25 aromatic substances and the like which do not deleteriously react with the active molecules of the invention.

For parenteral application, particularly suitable are solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Ampoules are convenient unit dosages.

30 For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including

those wherein the active component is protected with differentially degradable coatings, *e.g.*, by microencapsulation, multiple coatings, *etc.*

Therapeutic compounds of the invention also may be incorporated into liposomes. The incorporation can be carried out according to known
5 liposome preparation procedures, *e.g.* sonication and extrusion. Suitable conventional methods of liposome preparation are also disclosed in *e.g.* A.D. Bangham *et al.*, *J. Mol. Biol.*, **23**:238-252 (1965); F. Olson *et al.*, *Biochim. Biophys. Acta*, **557**:9-23 (1979); F. Szoka *et al.*, *Proc. Nat. Acad. Sci.*, **75**:4194-4198 (1978); S. Kim *et al.*, *Biochim. Biophys. Acta*, **728**:339-348
10 (1983); and Mayer *et al.*, *Biochim. Biophys. Acta*, **858**:161-168 (1986).

The invention also provides methods for invoking an immune response in a mammal such as a human, including vaccinating a mammal such as a human against a targeted disorder associated with the overexpression of p53 such as cancer.

15 These methods comprise administering to a mammal an effective amount of a DNA sequence that comprises a DNA vector that codes for a TCR molecules of the invention. Preparation of expression vectors of TCR molecules is described above and in the Examples which follow. Methods for administration of plasmid DNA, uptake of that DNA by cells of the
20 administered subject and expression of protein have been reported. See Ulmer, J.B., *et al.*, *Science* (1993) **259**: 1745-1749.

DNA vectors that encode TCR molecules of the invention are suitably administered to a mammal including a human preferably by intramuscular injection. Administration of cDNA to skeletal muscle of a mammal with
25 subsequent uptake of administered expression vector by the muscle cells and expression of protein encoded by the DNA has been described by Ulmer *et al.* and represents an exemplary protocol [Ulmer, J.B., *et al.*, *Science* **259**: 1745-1749]. The optimal dose for a given therapeutic application can be determined by conventional means.

30 In addition to treatment of human disorders, TCR molecules of the invention and DNA constructs of the invention that encode such TCR molecules will have significant use for veterinary applications, *e.g.*,

treatment of disorders of livestock such as cattle, sheep, *etc.* and pets such as dogs and cats using the cognate p53 antigens and MHC molecules appropriate for the animal species.

5 It will be appreciated that actual preferred amounts of a given TCR molecule of the invention or DNA construct coding for same used in a given therapy will vary according to the particular active compound or compounds being utilized, the particular compositions formulated, the mode of application, the particular site of administration, the patient's weight, general health, sex, *etc.*, the particular indication being treated, *etc.* and
10 other such factors that are recognized by those skilled in the art including the attendant physician or veterinarian. Optimal administration rates for a given protocol of administration can be readily determined by those skilled in the art using conventional dosage determination tests conducted *e.g.* with regard to the foregoing guidelines and the assays disclosed herein.

15 A "polypeptide" refers to any polymer preferably consisting essentially of any of the 20 natural amino acids regardless of its size. Although the term "protein" is often used in reference to relatively large proteins, and "peptide" is often used in reference to small polypeptides, use of these terms in the field often overlaps. The term "polypeptide" refers generally to
20 proteins, polypeptides, and peptides unless otherwise noted. Peptides useful in accordance with the present invention in general will be generally between about 0.1 to 100 KD or greater up to about 1000 KD, preferably between about 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 30 and 50 KD as judged by standard molecule sizing techniques such as centrifugation or SDS-
25 polyacrylamide gel electrophoresis.

As used herein, the term "cell" is intended to include any primary cell or immortalized cell line, any group of such cells as in, a tissue or an organ. Preferably the cells are of mammalian and particularly of human origin, and can be infected by one or more pathogens. A "host cell" in accord with the
30 invention can be an infected cell or it can be a cell such as *E. coli* that can be used to propagate a nucleic acid described herein.

All documents mentioned herein are incorporated herein by reference. The following non-limiting examples are illustrative of the invention.

Example 1: Construction of 264 single-chain (sc) TCR

5 The T cell clone, 264, recognizes a peptide fragment (aa 264-272; LLGRNSFEV) [SEQ ID NO. 1] of the human wild-type tumor suppresser protein p53 restricted by HLA-A2.1. The T cell receptor gene was cloned into a three-domain single-chain format previously shown to produce soluble TCR and functional receptor molecules.

10

 In brief, mRNA was isolated from the T cell clone and cDNA was made using the Marathon cDNA Amplification Kit (Clontech). Sequencing of cDNA clones identified two distinct V alpha chains (V alpha 3 and V alpha 13) and a single V beta chain (V beta 3). The cDNA was used as a template in
15 polymerase chain reaction (PCR) with primers KC228 and KC229 or KC226 and KC227 to produce 5' SfiI-3' SpeI V alpha 3 or V alpha 13 fragments respectively. The same DNA was then used as a PCR template with primers PRIB4 and KC176 to generate a 5' XhoI-3' XmaI V beta C beta chain
20 fragment. The C beta chain was truncated just before the cysteine residue at amino acid 127 of the full length C beta chain.

 The alpha and beta chain fragments were cloned into the pGEM-T Easy Vector System (Promega) for DNA sequence determination. Correct fragments were restriction digested and cloned into expression vector pKC60 (described previously in pending U.S. patent application no. 08/813,781) to
25 create two V alpha-(G4S)₄ V beta C beta scTCR molecules, 264-A (with V alpha 3) and 264-B (with V alpha 13).

 The DNA constructs described above (264-A and 264-B) were re-amplified by PCR with primers ET-TCRF1 and KC170 or ET-TCRF2 and KC170, respectively, to generate 5' AgeI-3' ClaI DNA fragments. The
30 fragments were cloned into the pGEM-T Easy Vector System for DNA sequence determination.

 The 5' AgeI-3' ClaI fragments were then used as the template DNA in PCR with primers KC232 and KC208 or KC231 and KC208, respectively, to

produce 5' AgeI-3' HpaI DNA fragments for cloning to produce the CD3 zeta fusion molecule or vectors comprising such molecules (described below) and eventually the 264 IL-2 fusion molecule or vectors comprising such molecules (described below).

5 Example 2: Construction of the CD3 zeta Fusion Shuttle Vector

To determine which of the two V alpha chains was functional, both the 264-A and 264-B sc-TCR were expressed as CD3 zeta fusion molecules.

Construction of a "shuttle vector" has been previously described.

10

Briefly, alpha and beta chain TCR fragments were cloned into the expression vector pKC60 to create a V alpha-(G₄ S)₄ V beta C beta scTCR molecule. The new vector was named pNAG2 (Fig. 9). pNAG2 was then re-amplified by PCR with primers KC203 and KC208 to generate a 5' AgeI-3' HpaI/BspEI/NruI/ClaI DNA fragment. The scTCR fragment was cloned into
15 the pGEM-T Easy Vector System and this new pGEM-based vector was then used as a "shuttle vector" for introduction of other DNA fragments to create a bispecific or fusion sc molecule.

Sc-Fv DNA was then restriction digested and cloned into the "shuttle
20 vector" downstream of the sc-TCR. To connect the sc-TCR and sc-Fv together as a single-chain fusion protein, the "shuttle vector" was digested with the appropriate restriction enzymes to drop out the previous linker DNA fragment and allow for ligation of linker sequences between the sc-TCR and the sc-Fv.

In the "shuttle vector" design outlined above, a stop codon and splice site were introduced between the NruI and ClaI restriction sites as part of the PCR amplification of the scTCR with "back" primer KC208. To aid in downstream purification of the bispecific sc protein, a set of annealed oligonucleotides (KC237 and KC238) was designed to introduce a 3' EE tag
25 (EEEEYMPME) (SEQ ID NO. 4) with a stop codon and splice site. The annealed oligonucleotide pair was cloned 5'NruI-3'ClaI into the "shuttle
30 vector" already encoding for the complete bispecific sc-TCR molecule.

After cloning the scTCR, sc-Fv, linker, and tag DNA fragments into the "shuttle vector" to complete the bispecific sc molecule design, the DNA was restriction digested (AgeI-ClaI) and cloned into the mammalian cell expression vector pSUN27 (Fig. 2) (previously described in the pending U.S. Application Serial No. 08/943,086 to create pBISP/149 and pBISP/D011.10 (Fig. 3). pBISP/D011.10 can be generated by one skilled in the art using pBISP/149 which is deposited as pSUN28 and any of three D011.10scTCR plasmids (pSUN18 – ATCC#97895, pSUN19 – ATCC#97896, or pSUN27 – ATCC#209276). The disclosure of the USSN 08/943, 086 is incorporated herein by reference.

Construction of the CD3 zeta Fusion Vector

In brief, murine cDNA was used as the template in polymerase chain reaction (PCR) with primers KC312 and KC304 to produce a 5'HpaI-3'ClaI murine CD3 zeta fragment.

The murine CD3 zeta fragment was cloned into the pGEM-T Easy Vector System for DNA sequence determination. The correct fragment was restriction digested and cloned into the "shuttle vector", effectively removing the existing linker, sc-FV, and EE tag.

After cloning the CD3 zeta gene into the "shuttle vector", the DNA was digested AgeI-HpaI to allow for ligation with the 264-A and 264-B sc-TCR fragments (described above), creating two new sc-TCR/CD3 zeta fusions. Lastly, the new DNA preparations were restriction digested (AgeI-ClaI) and cloned into the mammalian cell expression vector pSUN28 (pBISP/DO 11. 10 vector), Fig. 3 previously described in pending U.S. patent application no. 09/422,375.

Example 3: Expression of 264 scTCR/CD3 Zeta Fusion Molecules

Jurkat cells were prepared for transfection by washing with cold DPBS. The cells were resuspended in DPBS and mixed with 20 µg of PvuI linearized 264-A/CD3 zeta or 264-B/CD3 zeta DNA. After five minutes on ice, the cells were electroporated using a Gene Pulser (BioRad) set to deliver one pulse of 250 volts, 960 µFd. The pulsed cells were placed on ice for five minutes. The cells were diluted into 10 ml of 10% IMDM medium (IMDM,

10% FBS, 2mM glutamine) and grown in a T-25cm² TC flask overnight at 37°C with 5% CO₂. The next day, the cells were plated in 96 well plates with selective medium (10% IMDM plus 1.0 mg/ml G418). After 1 week, the concentration of G418 was increased to 2 mg/ml. The growing colonies
5 were re-fed approximately two weeks after transfection and screened about one week later.

The transfected Jurkat cells were screened for surface expression of scTCR using flow cytometry analysis. Positive transfectants were identified by staining with a fluorescent-tagged mAb (H57-597) which detects a
10 portion of the C beta domain of murine TCR.

Example 4: Identification of the Correct 264 scTCR V alpha Domain

Transfected Jurkat cells which expressed either the 264-A or 264-B version of the CD3 zeta fusion molecule were used in a cell activation assay. In the assay, the HLA-A2 presenting cell line T2 was used as the APC. The
15 T2 cells were loaded with 264 peptide (or irrelevant peptide) overnight at 37°C with 5% CO₂. The following day, the transfected Jurkat lines were added and allowed to interact with the peptide-pulsed APCs overnight.

Specific stimulation of the transfectants by 264-loaded APCs was assessed using an IL-2 ELISA. An anti-human IL-2 mAb was coated
20 passively overnight on a 96 well plate. The plate was washed and blocked with 10% FBS/DPBS for 1 hour. The blocking reagent was flicked out and supernatants from the assay were added to the plate for 1 hour at 37°C. After washing, the bound IL-2 was detected using another anti-IL-2 mAb conjugated to biotin. Following 45 minutes at 37°C, the plate was washed
25 and streptavidin-HRP was added for 15 minutes. Finally, the plate was washed and developed using ABTS substrate. Absorbance was read at 405 nm.

Based on the cell activation assay, the V alpha 3 domain is functional. Only cells expressing the 264-A molecule were stimulated to
30 produce IL-2 in the presence of 264 peptide-loaded APCs.

Table 1, shown on the following page, shows the primary sequence of various oligonucleotides used in the forgoing examples.

TABLE 1

KC228	SEQ ID NO. 7	5'- gag gtg gcc cag ccg gcc atg gcc cag tca gtg acg cag c -3'
KC229	SEQ ID NO. 8	5'- gag gtg act agt gtc tgg ctt tat aat tag -3'
KC226	SEQ ID NO. 9	5'- gag gtg gcc cag ccg gcc atg gcc gag cag gtg gag cag c-3'
KC227	SEQ ID NO. 10	5'- gag gtg act agt gtt tga ttt aac aga gag -3'
PRIB4	SEQ ID NO. 11	5'- ggg ggg ctc gag caa ttc aaa agt cat tca gac tc -3'
KC176	SEQ ID NO. 12	5'- gag gtg gag ccc ggg gtc tgc tcg gcc cca ggc -3'
ET- TCRF1	SEQ ID NO. 13	5'- ccc acc ggt cag tca gtg acg cag ccc -3'
KC170	SEQ ID NO. 14	5'- gtg gag ttc gaa aag tgt act tac gtt tgt ctg ctc ggc ccc ag -3'
ET-TCRF2	SEQ ID NO. 15	5'- ccc acc ggt gag cag gtg gag cag ctt -3'
KC232	SEQ ID NO. 16	5'- gag gtg acc ggt cag tca gtg acg cag c -3'
KC208	SEQ ID NO. 17	5'- gtg gag atc gat aag tgt act tac gtt ttc att atc gcg atc cgg agt taa cgt ctg ctc ggc ccc ag -3'
KC231	SEQ ID NO. 18	5'- gag gtg acc ggt gag cag gtg gag cag c -3'
KC312	SEQ ID NO. 19	5'- gag gtg gtt aac gat ccc aaa ctc tgc tac ttg cta gat gga atc ctc -3'
KC304	SEQ ID NO. 20	5'- gag gtg atc gat aag tgt act tac gtt ttt agc gag ggg gca ggg c -3'

Although a preferred embodiment of the invention has been described using specific terms, such description is for illustrative purposes only, and it is to be understood that changes and variations may be made without departing from the spirit or scope of the following claims.

SEQUENCE LISTING

<110> Altor Bioscience Corporation

<111> The Scripps Research Institute

<120> P53 BINDING T CELL RECEPTOR MOLECULES AND USES THEREOF

<130> 9-4820-283CA

<140> CA 2,487,779

<141> 2002-06-05

<150> 09/774,681

<151> 2001-02-01

<150> 08/812,393

<151> 1997-03-07

<150> 60/012,845

<151> 1996-03-05

<150> 60/296,324

<151> 2001-06-05

<160> 23

<170> PatentIn Ver. 2.1

<210> 1

<211> 9

<212> PRT

<213> Homo sapiens

<400> 1

Leu Leu Gly Arg Asn Ser Phe Glu Val
 1 5

<210> 2

<211> 377

<212> PRT

<213> Homo sapiens

<400> 2

Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val Ser Glu Gly Ala
 1 5 10 15

Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly Thr Pro Tyr Leu
 20 25 30

Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln Leu Leu Lys
 35 40 45

Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn Gly Phe Glu Ala
 50 55 60

Glu	Phe	Ser	Lys	Ser	Asn	Ser	Ser	Phe	His	Leu	Arg	Lys	Ala	Ser	Val	65	70	75	80
His	Trp	Ser	Asp	Ser	Ala	Val	Tyr	Phe	Cys	Val	Leu	Ser	Glu	Asp	Ser	85	90	95	
Asn	Tyr	Gln	Leu	Ile	Trp	Gly	Ser	Gly	Thr	Lys	Leu	Ile	Ile	Lys	Pro	100	105	110	
Asp	Thr	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	115	120	125	
Gly	Ser	Gly	Gly	Gly	Gly	Ser	Ser	Ser	Asn	Ser	Lys	Val	Ile	Gln	Thr	130	135	140	
Pro	Arg	Tyr	Leu	Val	Lys	Gly	Gln	Gly	Gln	Lys	Ala	Lys	Met	Arg	Cys	145	150	155	160
Ile	Pro	Glu	Lys	Gly	His	Pro	Val	Val	Phe	Trp	Tyr	Gln	Gln	Asn	Lys	165	170	175	
Asn	Asn	Glu	Phe	Lys	Phe	Leu	Ile	Asn	Phe	Gln	Asn	Gln	Glu	Val	Leu	180	185	190	
Gln	Gln	Ile	Asp	Met	Thr	Glu	Lys	Arg	Phe	Ser	Ala	Glu	Cys	Pro	Ser	195	200	205	
Asn	Ser	Pro	Cys	Ser	Leu	Glu	Ile	Gln	Ser	Ser	Glu	Ala	Gly	Asp	Ser	210	215	220	
Ala	Leu	Tyr	Leu	Cys	Ala	Ser	Ser	Leu	Ser	Gly	Gly	Gly	Thr	Glu	Val	225	230	235	240
Phe	Phe	Gly	Lys	Gly	Thr	Arg	Leu	Thr	Val	Val	Glu	Asp	Leu	Arg	Asn	245	250	255	
Val	Thr	Pro	Pro	Lys	Val	Ser	Leu	Phe	Glu	Pro	Ser	Lys	Ala	Glu	Ile	260	265	270	
Ala	Asn	Lys	Gln	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ala	Arg	Gly	Phe	Phe	275	280	285	
Pro	Asp	His	Val	Glu	Leu	Ser	Trp	Trp	Val	Asn	Gly	Lys	Glu	Val	His	290	295	300	
Ser	Gly	Val	Ser	Thr	Asp	Pro	Gln	Ala	Tyr	Lys	Glu	Ser	Asn	Tyr	Ser	305	310	315	320
Tyr	Cys	Leu	Ser	Ser	Arg	Leu	Arg	Val	Ser	Ala	Thr	Phe	Trp	His	Asn	325	330	335	
Pro	Arg	Asn	His	Phe	Arg	Cys	Gln	Val	Gln	Phe	His	Gly	Leu	Ser	Glu	340	345	350	
Glu	Asp	Lys	Trp	Pro	Glu	Gly	Ser	Pro	Lys	Pro	Val	Thr	Gln	Asn	Ile	355	360	365	

Ser Ala Glu Ala Trp Gly Arg Ala Asp
 370 375

<210> 3
 <211> 90
 <212> PRT
 <213> Homo sapiens

<400> 3
 Ile Gln Asn Pro Glu Pro Ala Val Tyr Gln Leu Lys Asp Pro Arg Ser
 1 5 10 15
 Gln Asp Ser Thr Leu Cys Leu Phe Thr Asp Phe Asp Ser Gln Ile Asn
 20 25 30
 Val Pro Lys Thr Met Glu Ser Gly Thr Phe Ile Thr Asp Lys Thr Val
 35 40 45
 Leu Asp Met Lys Ala Met Asp Ser Lys Ser Asn Gly Ala Ile Ala Trp
 50 55 60
 Ser Asn Gln Thr Ser Phe Thr Cys Gln Asp Ile Phe Lys Glu Thr Asn
 65 70 75 80
 Ala Thr Tyr Pro Ser Ser Asp Val Pro Ser
 85 90

<210> 4
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Linker peptide

<400> 4
 Ala Ser Gly Gly Gly Gly Ser Gly Gly Gly
 1 5 10

<210> 5
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Linker peptide

<400> 5
 Gly Gly Gly Gly Ser
 1 5

<210> 6
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide tag

<400> 6
 Glu Glu Glu Glu Tyr Met Pro Met Glu
 1 5

<210> 7
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 7
 gaggtggccc agccggccat ggcccagtca gtgacgcagc 40

<210> 8
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 8
 gaggtgacta gtgtctggct ttataattag 30

<210> 9
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 9
 gaggtggccc agccggccat ggccgagcag gtggagcagc 40

<210> 10
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 10

gaggtgacta gtgttttgatt taacagagag

30

<210> 11

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 11

ggggggctcg agcaattcaa aagtcattca gactc

35

<210> 12

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 12

gaggtggagc ccgggggtctg ctcggcccca ggc

33

<210> 13

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 13

cccaccggtc agtcagtgac gcagccc

27

<210> 14

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 14
gtggagttcg aaaagtgtac ttacgtttgt ctgctcggcc ccag 44

<210> 15
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 15
cccaccggtg agcaggtgga gcagctt 27

<210> 16
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 16
gaggtgaccg gtcagtcagt gacgcagc 28

<210> 17
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 17
gtggagatcg ataagtgtac ttacgttttc attatcgca tccggagtta acgtctgctc 60
ggccccag 68

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 18
gaggtgaccg gtgagcaggt ggagcagc 28

<210> 19
 <211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 19
 gaggtggtta acgatcccaa actctgctac ttgctagatg gaatcctc 48

<210> 20
 <211> 46
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 20
 gaggtgatcg ataagtgtac ttacgttttt agcgagggggg cagggc 46

<210> 21
 <211> 1131
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1131)

<400> 21
 cag tca gtg acg cag ccc gat gct cgc gtc act gtc tct gaa gga gcc 48
 Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val Ser Glu Gly Ala
 1 5 10 15
 tct ctg cag ctg aga tgc aag tat tcc tac tct ggg aca cct tat ctg 96
 Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly Thr Pro Tyr Leu
 20 25 30
 ttc tgg tat gtc cag tac ccg cgg cag ggg ctg cag ctg ctc ctc aag 144
 Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln Leu Leu Leu Lys
 35 40 45
 tac tat tca gga gac cca gtg gtt caa gga gtg aat ggc ttc gag gct 192
 Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn Gly Phe Glu Ala
 50 55 60
 gag ttc agc aag agt aac tct tcc ttc cac ctg cgg aaa gcc tct gtg 240
 Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg Lys Ala Ser Val
 65 70 75 80

cac	tgg	agc	gac	tct	gct	gtg	tac	ttc	tgt	gtt	ttg	agc	gag	gat	agc	288
His	Trp	Ser	Asp	Ser	Ala	Val	Tyr	Phe	Cys	Val	Leu	Ser	Glu	Asp	Ser	
				85					90					95		
aac	tat	cag	ttg	atc	tgg	ggc	tct	ggg	acc	aag	cta	att	ata	aag	cca	336
Asn	Tyr	Gln	Leu	Ile	Trp	Gly	Ser	Gly	Thr	Lys	Leu	Ile	Ile	Lys	Pro	
			100					105					110			
gac	act	agt	ggt	ggc	ggt	ggc	agc	ggc	ggt	ggt	ggt	tcc	ggt	ggc	ggc	384
Asp	Thr	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	
		115					120					125				
ggt	tct	ggc	ggt	ggc	ggt	tcc	tcg	agc	aat	tca	aaa	gtc	att	cag	act	432
Gly	Ser	Gly	Gly	Gly	Gly	Ser	Ser	Ser	Asn	Ser	Lys	Val	Ile	Gln	Thr	
	130					135					140					
cca	aga	tat	ctg	gtg	aaa	ggg	caa	gga	caa	aaa	gca	aag	atg	agg	tgt	480
Pro	Arg	Tyr	Leu	Val	Lys	Gly	Gln	Gly	Gln	Lys	Ala	Lys	Met	Arg	Cys	
145					150					155					160	
atc	cct	gaa	aag	gga	cat	cca	gtt	gta	ttc	tgg	tat	caa	caa	aat	aag	528
Ile	Pro	Glu	Lys	Gly	His	Pro	Val	Val	Phe	Trp	Tyr	Gln	Gln	Asn	Lys	
				165					170					175		
aac	aat	gag	ttt	aaa	ttt	ttg	att	aac	ttt	cag	aat	caa	gaa	gtt	ctt	576
Asn	Asn	Glu	Phe	Lys	Phe	Leu	Ile	Asn	Phe	Gln	Asn	Gln	Glu	Val	Leu	
			180					185					190			
cag	caa	ata	gac	atg	act	gaa	aaa	cga	ttc	tct	gct	gag	tgt	cct	tca	624
Gln	Gln	Ile	Asp	Met	Thr	Glu	Lys	Arg	Phe	Ser	Ala	Glu	Cys	Pro	Ser	
		195					200					205				
aac	tca	cct	tgc	agc	cta	gaa	att	cag	tcc	tct	gag	gca	gga	gac	tca	672
Asn	Ser	Pro	Cys	Ser	Leu	Glu	Ile	Gln	Ser	Ser	Glu	Ala	Gly	Asp	Ser	
	210					215					220					
gca	ctg	tac	ctc	tgt	gcc	agc	agt	ctg	tca	ggg	ggc	ggc	aca	gaa	gtt	720
Ala	Leu	Tyr	Leu	Cys	Ala	Ser	Ser	Leu	Ser	Gly	Gly	Gly	Thr	Glu	Val	
225					230					235					240	
ttc	ttt	ggt	aaa	gga	acc	aga	ctc	aca	gtt	gta	gag	gat	ctg	aga	aat	768
Phe	Phe	Gly	Lys	Gly	Thr	Arg	Leu	Thr	Val	Val	Glu	Asp	Leu	Arg	Asn	
				245					250					255		
gtg	act	cca	ccc	aag	gtc	tcc	ttg	ttt	gag	cca	tca	aaa	gca	gag	att	816
Val	Thr	Pro	Pro	Lys	Val	Ser	Leu	Phe	Glu	Pro	Ser	Lys	Ala	Glu	Ile	
			260					265					270			
gca	aac	aaa	caa	aag	gct	acc	ctt	gtg	tgc	ttg	gcc	agg	ggc	ttc	ttc	864
Ala	Asn	Lys	Gln	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ala	Arg	Gly	Phe	Phe	
		275					280				285					
cct	gac	cac	gtg	gag	ctg	agc	tgg	tgg	gtg	aat	ggc	aag	gaa	gtc	cac	912
Pro	Asp	His	Val	Glu	Leu	Ser	Trp	Trp	Val	Asn	Gly	Lys	Glu	Val	His	
	290					295					300					

```

agt ggg gtc agc acg gac cct cag gcc tac aag gag agc aat tat agc 960
Ser Gly Val Ser Thr Asp Pro Gln Ala Tyr Lys Glu Ser Asn Tyr Ser
305                      310                      315                      320

tac tgc ctg agc agc cgc ctg agg gtc tct gct acc ttc tgg cac aat 1008
Tyr Cys Leu Ser Ser Arg Leu Arg Val Ser Ala Thr Phe Trp His Asn
                      325                      330                      335

cct cgc aac cac ttc cgc tgc caa gtg cag ttc cat ggg ctt tca gag 1056
Pro Arg Asn His Phe Arg Cys Gln Val Gln Phe His Gly Leu Ser Glu
                      340                      345                      350

gag gac aag tgg cca gag ggc tca ccc aaa cct gtc aca cag aac atc 1104
Glu Asp Lys Trp Pro Glu Gly Ser Pro Lys Pro Val Thr Gln Asn Ile
                      355                      360                      365

agt gca gag gcc tgg ggc cga gca gac 1131
Ser Ala Glu Ala Trp Gly Arg Ala Asp
                      370                      375

<210> 22
<211> 270
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(270)

<400> 22
atc cag aac cca gaa cct gct gtg tac cag tta aaa gat cct cgg tct 48
Ile Gln Asn Pro Glu Pro Ala Val Tyr Gln Leu Lys Asp Pro Arg Ser
1                      5                      10                      15

cag gac agc acc ctc tgc ctg ttc acc gac ttt gac tcc caa atc aat 96
Gln Asp Ser Thr Leu Cys Leu Phe Thr Asp Phe Asp Ser Gln Ile Asn
                      20                      25                      30

gtg ccg aaa acc atg gaa tct gga acg ttc atc act gac aaa act gtg 144
Val Pro Lys Thr Met Glu Ser Gly Thr Phe Ile Thr Asp Lys Thr Val
                      35                      40                      45

ctg gac atg aaa gct atg gat tcc aag agc aat ggg gcc att gcc tgg 192
Leu Asp Met Lys Ala Met Asp Ser Lys Ser Asn Gly Ala Ile Ala Trp
                      50                      55                      60

agc aac cag aca agc ttc acc tgc caa gat atc ttc aaa gag acc aac 240
Ser Asn Gln Thr Ser Phe Thr Cys Gln Asp Ile Phe Lys Glu Thr Asn
65                      70                      75                      80

gcc acc tac ccc agt tca gac gtt ccc agt 270
Ala Thr Tyr Pro Ser Ser Asp Val Pro Ser
                      85                      90

```

<210> 23
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Linker peptide

<400> 23
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15

Gly Gly Gly Ser
20

What is claimed is:

1. An isolated T cell receptor (TCR) comprising a V α chain and a V β chain, being capable of specifically binding a p53 peptide in the context of an MHC molecule, wherein the p53 peptide comprises the amino acid sequence of SEQ ID NO:1, wherein the MHC molecule is HLA A2.1, and wherein the V α chain comprises amino acids 1 to 113 of SEQ ID NO:2 and the V β chain comprises amino acids 138 to 251 of SEQ ID NO:2.
2. The isolated TCR of claim 1, wherein the TCR comprises a covalently linked in sequence: a) the V α chain and b) a C α chain, wherein the C α chain comprises amino acids 1 to 90 of SEQ ID NO:3.
3. The isolated TCR of claim 1 or 2, wherein the TCR comprises a covalently linked in sequence: a) the V β chain and b) a C β sequence, wherein the C β sequence comprises amino acids 252 to 377 of SEQ ID NO:2.
4. The isolated TCR of any one of claims 1 to 3, wherein binding is determined by monitoring binding of the isolated TCR to the MHC molecule complexed with the p53 peptide.
5. The isolated TCR of claim 4, wherein the binding is monitored in a TCR ELISA or a TCR binding assay.
6. The isolated TCR of claim 4, wherein binding is monitored by measuring signal transduction by the isolated TCR.

7. The isolated TCR of any one of claims 4 to 6, wherein binding between the p53 peptide and the isolated TCR molecule is increased by at least 2 fold when compared to a control TCR heterodimer.
8. The isolated TCR of any one of claims 4 to 6, wherein binding between the p53 peptide and the isolated TCR molecule is increased by at least 10% relative to a control TCR heterodimer.
9. The isolated TCR of any one of claims 1 to 8, wherein the V α chain is at least 90% identical to amino acids 1 to 113 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 1 to 113 of SEQ ID NO:2.
10. The isolated TCR of any one of claims 1 to 9, wherein the V β chain is at least 90% identical to the amino acids 138 to 251 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 138 to 251 of SEQ ID NO:2.
11. The isolated TCR of claim 3, wherein the C β sequence has at least 90% identity to the amino acids 252 to 377 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 252 to 377 of SEQ ID NO:2.
12. The isolated TCR of claim 2, wherein the C α chain has at least 90% identity to the amino acids 1 to 90 of SEQ ID NO:3, wherein the 90% identity is determined over the entire length of amino acids 1 to 90 of SEQ ID NO:3.

13. The isolated TCR of any one of claims 1 to 12, wherein the TCR comprises a heterodimer comprising a V α chain dimerized to the V β chain.
14. The isolated TCR of any one of claims 1 to 12, wherein the TCR comprises a single chain TCR (sc TCR).
15. The isolated TCR of claim 14, wherein the V α chain is covalently linked to the V β chain by a peptide linker sequence.
16. The isolated TCR of claim 14 or 15, wherein the sc TCR comprises a transmembrane domain.
17. The isolated TCR of any one of claims 14 to 16, wherein the sc TCR comprises a cytoplasmic signaling domain.
18. The isolated TCR of claim 16, wherein the sc TCR comprises in sequence 1) the V α 3 chain as shown in SEQ ID NO:2, 2) a peptide linker, and 3) the V β 3 chain as shown in SEQ ID NO:2.
19. The isolated TCR of claim 18, further comprising a C β chain as provided in SEQ ID NO:2 linked to the C-terminus of the V β 3 chain.
20. The isolated TCR of claim 18 or 19, further comprising a fragment of the C α chain as shown in SEQ ID NO:3, the fragment being covalently linked between the C-terminus of the V α chain and the N-terminus of the peptide linker.

21. The isolated TCR of any one of claims 18 to 20, wherein the peptide linker has the amino acid sequence of SEQ ID NO:5 repeated at least four times.
22. An isolated nucleic acid encoding the isolated TCR according to any one of claims 1 to 21.
23. A vector comprising the nucleic acid according to claim 22.
24. A host cell comprising the isolated nucleic acid according to claim 22 or the vector according to claim 23.
25. A pair of DNA segments encoding the heterodimer of claim 13, wherein a first DNA segment encodes the V α chain and a second DNA segment encodes the V β chain.
26. The pair of DNA segments of claim 25, wherein the V α chain is at least 90% identical to the amino acids 1 to 113 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 1 to 113 of SEQ ID NO:2.
27. The pair of DNA segments of claim 25 or 26, wherein the encoded V β chain is at least 90% identical to the amino acids 138 to 251 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 138 to 251 of SEQ ID NO:2.

28. The pair of DNA segments of any one of claims 25 to 27, wherein the second segment further encodes a C β chain at least 90% identical to the amino acids 252 to 377 of SEQ ID NO:2, wherein the 90% identity is determined over the entire length of amino acids 252 to 377 of SEQ ID NO:2, and wherein the C-terminus of the encoded V β chain being linked to the C-terminus of the encoded C β chain.

29. An *in vitro* method of identifying a cell or tissue expressing p53, said method comprising (i) contacting the cell or tissue with the isolated TCR according to any one of claims 1 to 21 so as to form a complex between the isolated TCR and p53 and (ii) detecting the complex so as to identify the cell or tissue expressing p53.

30. An *in vitro* method of identifying a cell or tissue expressing p53 comprising (i) contacting the cell or tissue with the host cell according to claim 24 so as to form a complex between the isolated TCR and p53 and (ii) detecting the complex so as to identify the cell or tissue expressing 53.

31. Use of the cell according to claim 24 for killing a cell expressing p53 peptide in the context of an MHC molecule, wherein the MHC molecule is HLA A2.1.

32. The use of claim 31, further comprising forming an immune complex between the MHC molecule and the TCR of any one of claims 1 to 21 expressed on the cell.

33. Use of the TCR according to any one of claims 1 to 21, or the host cell according to claim 24 for the treatment of cancer, wherein the cancer is characterized by upregulation of p53.




FIGURE 1




FIGURE 2

BISP/DO11.10	DO11.10 scTCR	linker	145-2C11 scFv	EE
BISP/149	p149 scTCR	linker	145-2C11 scFv	EE

FIGURE 3

Vα3 domain

+1 Q S V T Q P D A R V T V S

51 CAGTCAGTGA CGCAGCCCGA TGCTCGCGTC ACTGTCTCTG
GTCAGTCACT GCGTCGGGCT ACGAGCGCAG TGACAGAGAC

+1 E G A S L Q L R C K Y S Y S G T P

101 AAGGAGCCTC TCTGCAGCTG AGATGCAAGT ATTCCTACTC TGGGACACCT
TTCCTCGGAG AGACGTCGAC TCTACGTTCA TAAGGATGAG ACCCTGTGGA

+1 Y L F W Y V Q Y P R Q G L Q L L L

151 TATCTGTTCT GGTATGTCCA GTACCCGCGG CAGGGGCTGC AGCTGCTCCT
ATAGACAAGA CCATACAGGT CATGGGCGCC GTCCCCGACG TCGACGAGGA

+1 K Y Y S G D P V V Q G V N G F E

201 CAAGTACTAT TCAGGAGACC CAGTGGTTCA AGGAGTGAAT GGCTTCGAGG
GTTTCATGATA AGTCCTCTGG GTCACCAAGT TCCTCACTTA CCGAAGCTCC

+1 A E F S K S N S S F H L R K A S V

251 CTGAGTTCAG CAAGAGTAAC TCTTCCTTCC ACCTGCGGAA AGCCTCTGTG
GACTCAAGTC GTTCTCATTG AGAAGGAAGG TGGACGCCTT TCGGAGACAC

+1 H W S D S A V Y F C V L S E D S N

301 CACTGGAGCG ACTCTGCTGT GTACTTCTGT GTTTTGAGCG AGGATAGCAA
GTGACCTCGC TGAGACGACA CATGAAGACA CAAAACTCGC TCCTATCGTT

+1 Y Q L I W G S G T K L I I K P D
SpeI

351 CTATCAGTTG ATCTGGGGCT CTGGGACCAA GCTAATTATA AAGCCAGACA
GATAGTCAAC TAGACCCCGA GACCCTGGTT CGATTAATAT TTCGGTCTGT

Linker Region

+1 T S G G G G S G G G G S G G G G S
SpeI

~~~~~

401 CTAGTGGTGG CGGTGGCAGC GGCGGTGGTG GTTCCGGTGG CGGCGGTTCT  
GATCACCACC GCCACCGTCG CCGCCACCAC CAAGGCCACC GCCGCCAAGA

Vβ3 Domain

+1 G G G G S S S N S K V I Q T P R Y  
XhoI

~~~~~

451 GGCGGTGGCG GTTCCTCGAG CAATTCAAAA GTCATTCAGA CTCCAAGATA
CCGCCACCGC CAAGGAGCTC GTTAAGTTTT CAGTAAGTCT GAGGTTCTAT

+1 L V K G Q G Q K A K M R C I P E

501 TCTGGTGAAA GGGCAAGGAC AAAAAGCAAA GATGAGGTGT ATCCCTGAAA
AGACCACTTT CCCGTTCTCTG TTTTTCGTTT CTACTCCACA TAGGGACTTT

FIGURE 4A

```

+1 K G H P V V F W Y Q Q N K N N E F
551 AGGGACATCC AGTTGTATTC TGGTATCAAC AAAATAAGAA CAATGAGTTT
    TCCCTGTAGG TCAACATAAG ACCATAGTTG TTTTATTCTT GTTACTCAAA

+1 K F L I N F Q N Q E V L Q Q I D M
601 AAATTTTGA TTAACTTTCA GAATCAAGAA GTTCTTCAGC AAATAGACAT
    TTTAAAACT AATTGAAAGT CTTAGTTCTT CAAGAAGTCG TTTATCTGTA

+1 T E K R F S A E C P S N S P C S
651 GACTGAAAAA CGATTCTCTG CTGAGTGTCC TTCAAACCTCA CCTTGCAGCC
    CTGACTTTTT GCTAAGAGAC GACTCACAGG AAGTTTGAGT GGAACGTCGG

+1 L E I Q S S E A G D S A L Y L C A
701 TAGAAATTCA GTCCTCTGAG GCAGGAGACT CAGCACTGTA CCTCTGTGCC
    ATCTTTAAGT CAGGAGACTC CGTCCTCTGA GTCGTGACAT GGAGACACGG

+1 S S L S G G G T E V F F G K G T R
751 AGCAGTCTGT CAGGGGGCGG CACAGAAGTT TTCTTTGGTA AAGGAACCAG
    TCGTCAGACA GTCCCCCGCC GTGTCTTCAA AAGAAACCAT TTCCTTGGTC

      cβ domain

+1 L T V V E D L R N V T P P K V S
801 ACTCACAGTT GTAGAGGATC TGAGAAATGT GACTCCACCC AAGGTCTCCT
    TGAGTGTCOA CATCTCCTAG ACTCTTTACA CTGAGGTGGG TTCCAGAGGA

+1 L F E P S K A E I A N K Q K A T L
851 TGTTTGAGCC ATCAAAAGCA GAGATTGCAA ACAAACAAAA GGCTACCCCT
    ACAAACCTCG TAGTTTTCTG CTCTAACGTT TGTTTGTTTT CCGATGGGAA

+1 V C L A R G F F P D H V E L S W W
901 GTGTGCTTGG CCAGGGGCTT CTTCCCTGAC CACGTGGAGC TGAGCTGGTG
    CACACGAACC GGTCCCCGAA GAAGGGACTG GTGCACCTCG ACTCGACCAC

+1 V N G K E V H S G V S T D P Q A
951 GGTGAATGGC AAGGAAGTCC ACAGTGGGGT CAGCACGGAC CCTCAGGCCT
    CCACTTACCG TTCCTTCAGG TGTCACCCCA GTCGTGCCTG GGAGTCCGGA

+1 Y K E S N Y S Y C L S S R L R V S
1001 ACAAGGAGAG CAATTATAGC TACTGCCTGA GCAGCCGCCT GAGGGTCTCT
    TGTTCTCTC GTTAATATCG ATGACGGACT CGTCGGCGGA CTCCCAGAGA

+1 A T F W H N P R N H F R C Q V Q F
1051 GCTACCTTCT GGCACAATCC TCGCAACCAC TTCCGCTGCC AAGTGCAGTT
    CGATGGAAGA CCGTGTTAGG AGCGTTGGTG AAGGCGACGG TTCACGTCAA

+1 H G L S E E D K W P E G S P K P
1101 CCATGGGCTT TCAGAGGAGG ACAAGTGGCC AGAGGGCTCA CCCAAACCTG
    GGTACCCGAA AGTCTCCTCC TGTTACCCGG TCTCCCGAGT GGGTTTGGAC

```

FIGURE 4B

+1 V T Q N I S A E A W G R A D
1151 TCACACAGAA CATCAGTGCA GAGGCCTGGG GCCGAGCAGA C
AGTGTGTCTT GTAGTCACGT CTCCGGACCC CGGGTCGTCT G

FIGURE 4C

+1 V T Q N I S A E A W G R A D

5 1151 TCACACAGAA CATCAGTGCA GAGGCCTGGG GCCGAGCAGA C
AGTGTGTCTT GTAGTCACGT CTCCGGACCC CGGGTCGTCT G

FIGURE 4C

10

I Q N P E P A V Y Q L
K D P R S Q D S T L
ATCCAGAACCCAGAACCTGCTGTGTACCAGTTAAAAGATCCTCGGTCTCAGGA
5 CAGCACCTC

C L F T D F D S Q I N V
P K T M E S G T F
TGCCTGTTCACCGACTTTGACTCCCAAATCAATGTGCCGAAAACCATGGAATC
10 TGGAACGTTC

I T D K T V L D M K A M
D S K S N G A I A
ATCACTGACAAAACCTGTGCTGGACATGAAAGCTATGGATTCCAAGAGCAATGG
15 GGCCATTGC

W S N Q T S F T C Q D
I F K E T N A T Y
CTGGAGCAACCAGACAAGCTTCACCTGCCAAGATATCTTCAAAGAGACCAAC
20 GCCACCTACC

P S S D V P S
CCAGTTCAGACGTTCCCAGT

25

FIGURE 5