

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2970263 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 D 471/04 (2006.01)** **C 07 C 67/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-08-13**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-05-09**
- (86) Europæisk ansøgning nr.: **14715501.4**
- (86) Europæisk indleveringsdag: **2014-03-12**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-01-20**
- (86) International ansøgning nr.: **US2014024224**
- (87) Internationalt publikationsnr.: **WO2014165044**
- (30) Prioritet: **2013-03-13 US 201361780621 P** **2014-03-04 US 201461947850 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **AbbVie Inc., 1 North Waukegan, Road, North Chicago, IL 60064, USA**
- (72) Opfinder: **BARKALOW, Jufang, 1027 Knollwood Road, Deerfield, IL 60015, USA**
CALIFANO, Jean-christophe, 6167 North Bay Ridge, Whitefish Bay, WI 53217, USA
CHAN, Vincent S., 734 Hinman Avenue 1, Evanston, IL 60202, USA
CHRISTENSEN, Alan, C., 2147 West Wicklow Lane, Round Lake, IL 60073, USA
GRIEME, Timothy A., 4942 N. Winchester Avenue, Chicago, IL 60640, USA
KU, Yi-Yin, 23 River Oaks Circle, Buffalo Grove, IL 60089, USA
MULHERN, Mathew M., 1009 Oaktree Trail, Lake Villa, IL 60046, USA
PU, Yu-ming M., 5507 Notting Hill Road, Gurnee, IL 60031, USA
- (74) Fuldmægtig i Danmark: **Patrade A/S, Ceresbyen 75, 8000 Århus C, Danmark**
- (54) Benævnelse: **PROCESSER TIL TILVIRKNING AF ET APOPTOSE-FORANLEDIGENDE VIRKEMIDDEL**
- (56) Fremdragne publikationer:
WO-A1-2011/150016
WO-A1-2012/071336
P. G. GASSMAN ET AL: "A General Procedure for the Base-Promoted Hydrolysis of Hindered Esters at Ambient Temperatures", J. ORG. CHEM, vol. 42, no. 5, 1 January 1977 (1977-01-01), pages 918-920, XP055115941,
"Preparation of t-Butyl-3-Bromo-5-Formylbenzoate Through Selective Metal-Halogen Exchange Reactions",
ORGANIC SYNTHESSES, vol. 89, 1 January 2012 (2012-01-01), page 460, XP55128994, ISSN: 0078-6209, DOI:
10.15227/orgsyn.089.0460
T. RAUSIS ET AL: "The Basicity Gradient-Driven Migration of Iodine: Conferring Regioflexibility on the
Substitution of Fluoroarenes", EUR. J. ORG. CHEM., 1 January 2002 (2002-01-01), pages 3351-3358,
XP55115669,
V. BAVETSIAS ET AL: "Synthesis of N-substituted derivatives of tert-butyl 4-aminobenzoate via a palladium-
catalysed reaction", J. CHEM. RESEARCH (S), 1 September 2000 (2000-09-01), pages 418-419, XP055324460,

Fortsættes ...

DESCRIPTION

[0001] This application claims the benefit of United States Provisional Application No. 61/780,621, filed March 13, 2013, and United States Provisional Application No. 61/947,850, filed March 4, 2014.


FIELD

[0002] Provided herein are processes for the preparation of an apoptosis-inducing agent, and chemical intermediates thereof. Also provided herein are novel chemical intermediates related to the processes provided herein.


BACKGROUND

[0003] 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide (hereafter, "Compound 1") and 4-(4-{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl)-N-({3-nitro-4-[(1R,4R)-[4-hydroxy-4-methylcyclohexyl]methyl]amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide (hereafter, "Compound 2") are each potent and selective Bcl-2 inhibitors having, *inter alia*, antitumor activity as apoptosis-inducing agents.

[0004] Compound 1 has the formula:




[0005] Compound 2 has the formula:




[0006] Compound 1 is currently the subject of ongoing clinical trials for the treatment of chronic lymphocytic leukemia. U.S. Patent Publication No. 2010/0305122 describes Compound 1, Compound 2, and other compounds which exhibit potent binding to a Bcl-2 family protein, and pharmaceutically acceptable salts thereof. WO2011/150016 discloses compounds which inhibit the activity of anti-apoptotic Bcl-2 proteins, compositions containing the compounds and methods of treating diseases during which is expressed anti-apoptotic Bcl-2 protein. WO2012/071336 discloses salts and crystalline forms of 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-ylmethyl)amino]phenyl}-sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide which are suitable active pharmaceutical ingredients for pharmaceutical compositions useful in treatment of a disease characterized by overexpression of one or more anti-apoptotic Bcl-2 family proteins, for example cancer. U.S. Patent Publication Nos. 2012/0108590 and 2012/0277210 describe pharmaceutical compositions comprising such compounds, and methods for the treatment of neoplastic, immune or autoimmune diseases comprising these compounds. U.S. Patent Publication No. 2012/0129853 describes methods for the treatment of systemic lupus erythematosus, lupus nephritis or Sjogren's Syndrome comprising these compounds. U.S. Patent Publication No. 2012/0157470 describes pharmaceutically acceptable salts and crystalline forms of Compound 1.

SUMMARY

[0007] Provided herein are processes for the preparation of compounds of formula A 1:




wherein R^2 is selected from




which comprises:

(a) combining a compound of formula (K):





wherein R is C₁ to C₁₂ alkyl,

with a tert-butoxide salt, an aprotic organic solvent, and water to provide a compound of formula (L):





and

(b'') combining the compound of formula (L) with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, 4-dimethylaminopyridine, an organic solvent, and either a compound of formula (N), to provide a compound of formula (A1) wherein R² is




or a compound of formula (P), to provide a compound of formula (A1) wherein R² is




thereby providing a compound of formula (A1);


wherein the compound of formula (K) is prepared by_:

(d) combining a compound of formula (D):



with a compound of formula (I):







a source of palladium, a tert-butoxide salt, and a phosphine ligand in an aprotic organic solvent to provide the compound of formula (K).

DETAILED DESCRIPTION

[0008] Provided herein is a process for the preparation of compounds of formula A1:




wherein R^2 is selected from




which comprises:

(a) combining a compound of formula (K):



wherein R is C_1 to C_{12} alkyl,



with a tert-butoxide salt, an aprotic organic solvent, and water to provide a compound of formula (L):





and

(b'') combining the compound of formula (L) with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDAC), 4-dimethylaminopyridine (DMAP), an organic solvent, and either a compound of formula (N), to provide a compound of formula (A1)

wherein R^2 is




or a compound of formula (P), to provide a compound of formula (A1) wherein R^2 is




thereby providing a compound of formula (A1);

wherein the compound of formula (K) is prepared by:

(d) combining a compound of formula (D):




with a compound of formula (I):




a source of palladium, a tert-butoxide salt, and a phosphine ligand in an aprotic organic solvent to provide the compound of formula (K).

[0009] In one embodiment, R^2 is




and step (b'') comprises:

(b'') combining the compound of formula (L) with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, 4-dimethylaminopyridine, an organic solvent, and the compound of formula (N):




to provide a compound of formula (A1).

[0010] In another embodiment, R^2 is




[0011] In some embodiments, R is C_1 to C_6 alkyl. In some embodiments, R is C_1 to C_4 alkyl. In some embodiments, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl. In some embodiments, R is tert-butyl.

[0012] In one embodiment, of the process provided herein, R^2 is



and the process further comprises:


(c'') combining a compound of formula (M):




with a tertiary amine base, an organic solvent, and (1R,4R)-4-(aminomethyl)-1-methylcyclohexanol or a salt thereof, to provide the compound of formula (P).

[0013] In one embodiment, the (1R,4R)-4-(aminomethyl)-1-methylcyclohexanol salt of step (c'') is the p-toluenesulfonic acid salt.

[0014] In some embodiments, the phosphine ligand is a compound of formula (J):




[0015] In other embodiments, the phosphine ligand is selected from:



[0016] In another embodiment, the process provided herein further comprises:

(e) combining a compound of formula (B) with a compound of formula (C):




wherein R is C₁ to C₁₂ alkyl,

and a tert-butoxide salt in an organic solvent to provide the compound of formula (D).

[0017] In another embodiment, the process provided herein further comprises:

(f) combining a compound of formula (A):




with R¹MgX in an aprotic organic solvent;

wherein R¹ is C₁ to C₆ alkyl; and X is Cl, Br or I; and


(g) combining a C₁ to C₁₂ alkyl chloroformate or a di-(C₁ to C₁₂ alkyl)dicarbonate with the product of step (f), to provide the compound of formula (C).

[0018] In another embodiment, the process provided herein further comprises:

(h) combining a compound of formula (E):




with DMF and POCl₃ to provide a compound of formula (F):




(i) combining the compound of formula (F) with a source of palladium and 4-chlorophenylboronic acid in an organic solvent to provide a compound of formula (G):





(j) combining the compound of formula (G) with BOC-piperazine and sodium triacetoxyborohydride in an organic solvent to provide a compound of formula (H):




(k) combining the compound of formula (H) with hydrochloric acid to provide the compound of formula (I).

[0019] In one embodiment, the process comprises step (a), step (b''), step (c'') and step (d). In one embodiment, the process comprises step (a), step (b''), step (c''), step (d) and step (e). In one embodiment, the process comprises step (a), step (b''), step (c''), step (d), step (e), step (f) and step (g). In another embodiment the process comprises step (a), step (b''), step (c''), step (d), step (e), step (f), step (g), step (h), step (i), step (j) and step (k).


[0020] In one embodiment, which is reflected in claim 1, the process comprises steps (a), (b'') and (d). In another embodiment, the process comprises steps (a), (b''), (d) and (e). In another embodiment, the process comprises steps (a), (b''), (d), (h), (i), (j) and (k). In another embodiment, the process comprises steps (a), (b''), (c''), (d), (h), (i), (j) and (k). In another embodiment, the process comprises steps (a), (b''), (d), (f), (g), (h), (i), (j) and (k). In another embodiment, the process comprises steps (a), (b''), (d), (e), (f), (g), (h), (i), (j) and (k).


[0021] Also provided herein is a process for the preparation of Compound 1, which is an embodiment of Compound (A1) as described above, -of the formula:



which comprises:


1. (a) combining a compound of formula (K):






wherein R is C₁ to C₁₂ alkyl,

with a tert-butoxide salt, an aprotic organic solvent, and water to provide a compound of formula (L):




2. (b) combining the compound of formula (L) with a compound of formula (N):




and

1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDAC), 4-dimethylaminopyridine (DMAP), and an organic solvent to provide the compound of formula (1), wherein the compound of formula (K) is prepared by:

(d) combining a compound of formula (D):



with a compound of formula (I):




a source of palladium, a tert-butoxide salt, and a phosphine ligand in an aprotic organic solvent to provide the compound of formula (K).

[0022] In some embodiments, R is C₁ to C₆ alkyl. In some embodiments, R is C₁ to C₄ alkyl. In some embodiments, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl. In some embodiments, R is tert-butyl.


[0023] In one embodiment, the process for the preparation of Compound 1 further comprises:

(c) combining a compound of formula (M):




with (tetrahydro-2H-pyran-4-yl)methanamine or a salt thereof, a tertiary amine base, and an organic solvent to provide the compound of formula (N).

[0024] In some embodiments, the phosphine ligand is a compound of formula (J):




[0025] In other embodiments, the phosphine ligand is selected from:



[0026] In another embodiment, the process for the preparation of Compound 1 further comprises:

(e) combining a compound of formula (B) with a compound of formula (C):




wherein R is C₁ to C₁₂ alkyl,

and a tert-butoxide salt in an organic solvent to provide the compound of formula (D).

[0027] In another embodiment, the process for the preparation of Compound 1 further comprises:

(f) combining a compound of formula (A):






with R^1MgX in an aprotic organic solvent;

wherein R^1 is C_1 to C_6 alkyl; and X is Cl, Br or I; and


(g) combining a C_1 to C_{12} alkyl chloroformate or a di- $(C_1$ to C_{12} alkyl)dicarbonate with the product of step (f), to provide the compound of formula (C).

[0028] In another embodiment, the process for the preparation of Compound 1 further comprises:


(h) combining a compound of formula (E):




with DMF and $POCl_3$ to provide a compound of formula (F):



(i) combining the compound of formula (F) with a source of palladium and 4-chlorophenylboronic acid in an organic solvent to provide a compound of formula (G):



(j) combining the compound of formula (G) with BOC-piperazine and sodium triacetoxyborohydride in an organic solvent to provide a compound of formula (H):




and

(k) combining the compound of formula (H) with hydrochloric acid to provide the compound of formula (I).

[0029] In one embodiment, the process for the preparation of Compound 1 comprises steps (a) through (d). In one embodiment, the process for the preparation of Compound 1 comprises steps (a) through (e). In another embodiment, the process for the preparation of Compound 1 comprises steps (a) through (g). In another embodiment, the process for the preparation of Compound 1 comprises steps (a) through (k).


[0030] In one embodiment, which is reflected in claim 1, the process for the preparation of Compound 1 comprises steps (a), (b) and (d). In another embodiment, the process for the preparation of Compound 1 comprises steps (a), (b), (d) and (e). In another embodiment, the process for the preparation of Compound 1 comprises steps (a), (b), (d), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 1 comprises steps (a), (b), (c), (d), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 1 comprises steps (a), (b), (d), (f), (g), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 1 comprises steps (a), (b), (d), (e), (f), (g), (h), (i), (j) and (k).

[0031] Also provided herein is a process for the preparation of Compound 2, which is another embodiment of Compound (A1) as described above, of the formula:




which comprises:

(a) combining a compound of formula (K):




wherein R is C₁ to C₁₂ alkyl,

with a tert-butoxide salt, an aprotic organic solvent, and water to provide a compound of formula (L):




(b') combining the compound of formula (L) with a compound of formula (P):




and

1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDAC), 4-dimethylaminopyridine (DMAP), and an organic solvent to provide the compound of formula (2), wherein the compound of formula (K) is prepared by:

(d) combining a compound of formula (D):



with a compound of formula (I):




a source of palladium, a tert-butoxide salt, and a phosphine ligand in an aprotic organic solvent to provide the compound of formula (K).

[0032] In some embodiments, R is C₁ to C₆ alkyl. In some embodiments, R is C₁ to C₄ alkyl. In some embodiments, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl. In some embodiments, R is tert-butyl.

[0033] In one embodiment, the process for the preparation of Compound 2 further comprises:

(c') combining a compound of formula (M):



with (1R,4R)-4-(aminomethyl)-1-methylcyclohexanol or a salt thereof, a tertiary amine base, and an organic solvent to provide the compound of formula (P).

[0034] In one embodiment, the (1R,4R)-4-(aminomethyl)-1-methylcyclohexanol salt of step (c') is the p-toluenesulfonic acid salt.

[0035] In some embodiments, the method for the preparation of Compound 2 further comprises step (e) as described above for the preparation of Compound 1.

[0036] In some embodiments, the method for the preparation of Compound 2 further comprises step (f) and step (g) as described above for the preparation of Compound 1.

[0037] In some embodiments, the method for the preparation of Compound 2 further comprises step (h), step (i), step (j) and step (k) as described above for the preparation of Compound 1.

[0038] In one embodiment, the process for the preparation of Compound 2 comprises step (a), step (b'), step (c') and step (d). In one embodiment, the process for the preparation of Compound 2 comprises step (a), step (b'), step (c'), step (d) and step (e). In one embodiment, the process for the preparation of Compound 2 comprises step (a), step (b'), step (c'), step (d), step (e), step (f) and step (g). In another embodiment the process for the preparation of Compound 2 comprises step (a), step (b'), step (c'), step (d), step (e), step (f), step (g), step (h), step (i), step (j) and step (k).

[0039] In one embodiment, which is reflected in claim 1, the process for the preparation of Compound 2 comprises steps (a), (b') and (d). In another embodiment, the process for the preparation of Compound 2 comprises steps (a), (b'), (d) and (e). In another embodiment, the process for the preparation of Compound 2 comprises steps (a), (b'), (d), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 2 comprises steps (a), (b'), (c'), (d), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 2 comprises steps (a), (b'), (d), (f), (g), (h), (i), (j) and (k). In another embodiment, the process for the preparation of Compound 2 comprises steps (a), (b'), (d), (e), (f), (g), (h), (i), (j) and (k).

[0040] In some embodiments, in step (a) the tert-butoxide salt is selected from the group consisting of sodium tert-butoxide and potassium tert-butoxide. In some embodiments, in step (a) the tert-butoxide salt is sodium tert-butoxide. In some embodiments, in step (a) the tert-butoxide salt is potassium tert-butoxide.

[0041] In some embodiments, in step (a) the aprotic organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (a) the aprotic organic solvent is 2-methyltetrahydrofuran.

[0042] In some embodiments, in step (b), step (b') and/or step (b'') the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (b), step (b') and/or step (b'') the organic solvent is selected from the group consisting of dichloromethane,

chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (b), step (b') and/or step (b'') the organic solvent is dichloromethane.

[0043] In some embodiments, in step (c), step (c') and/or step (c'') the tertiary amine base is N,N-diisopropylethylamine.

[0044] In some embodiments, in step (c), step (c') and/or step (c'') the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (c), step (c') and/or step (c'') the organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (c), step (c') and/or step (c'') the organic solvent is acetonitrile.

[0045] In some embodiments, in step (d) the compound of formula (I) is first combined with a base prior to the combining of step (d). In some embodiments, the base is an inorganic base. In some embodiments, the base is an organic base. In some embodiments, the base is selected from the group consisting of K_3PO_4 , Na_3PO_4 , NaOH, KOH, K_2CO_3 or Na_2CO_3 . In some embodiments, the base is K_3PO_4 . In some embodiments, in step (d) the compound of formula (I) is first combined with a base in one or more solvents prior to the combining of step (d).

[0046] In some embodiments, in step (d) the source of palladium is Pd_2dba_3 or $[(\text{cinnamyl})PdCl]_2$. In some embodiments, in step (d) the source of palladium is Pd_2dba_3 .

[0047] In some embodiments, in step (d) the tert-butoxide salt is selected from the group consisting of sodium tert-butoxide and potassium tert-butoxide.

[0048] In some embodiments, in step (d) the tert-butoxide salt is anhydrous. In some embodiments, in step (d) the tert-butoxide salt is anhydrous sodium tert-butoxide.

[0049] In some embodiments, in step (d) the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (d) the organic solvent is selected from

the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (d) the aprotic organic solvent is a mixture of THF and toluene.

[0050] In some embodiments, step (d) further comprises the following steps:

1. (1) combining the tert-butoxide salt with the compound of formula (I) in an aprotic organic solvent;
2. (2) combining the source of palladium, the compound of formula (J), and the compound of formula (D) in an aprotic organic solvent; and
3. (3) adding the mixture of step (1) to the mixture of step (2).

[0051] In some embodiments, in step (d) the mixture resulting from step (2) is filtered prior to step (3).

[0052] In some embodiments, step (d) is carried out under an atmosphere of nitrogen or argon.

[0053] In some embodiments, in step (d) a catalytic amount of the source of palladium is used relative to the amount of compound (I). In some embodiments, the source of palladium is Pd_2dba_3 and the catalytic amount of Pd_2dba_3 is from about 0.5 mole percent to about 2 mole percent. In one embodiment, the catalytic amount of Pd_2dba_3 is about 0.75 mole percent.

[0054] In some embodiments, in step (d) a catalytic amount of the compound of formula (J) is used relative to the amount of compound (I). In some embodiments, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 5 mole percent. In one embodiment, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 4 mole percent. In one embodiment, the catalytic amount of the compound of formula (J) is from about 2 mole percent to about 4 mole percent. In one embodiment, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 2 mole percent. In one embodiment, the catalytic amount of the compound of formula (J) is about 1 mole percent or about 2 mole percent.

[0055] In some embodiments, in step (e) the tert-butoxide salt is selected from the group consisting of sodium tert-butoxide and potassium tert-butoxide. In some embodiments, in step (e) the tert-butoxide salt is sodium tert-butoxide. In some embodiments, in step (e) the tert-butoxide salt is potassium tert-butoxide.

[0056] In some embodiments, in step (e) the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone,

acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (e) the organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (e) the organic solvent is DMF.

[0057] In some embodiments, in step (f), R^1 is C_1 to C_4 alkyl. In some embodiments, R^1 is isopropyl.

[0058] In some embodiments, in step (f), R is methyl and the C_1 to C_{12} alkyl chloroformate is methyl chloroformate. In some embodiments, R is ethyl and the C_1 to C_{12} alkyl chloroformate is ethyl chloroformate. In some embodiments, R is tert-butyl and the di-(C_1 to C_{12} alkyl)dicarbonate is di-tert-butyl dicarbonate.

[0059] In some embodiments, in step (f) the organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (f) the aprotic organic solvent is THF.

[0060] In some embodiments, in step (i) the source of palladium is $Pd(OAc)_2$.

[0061] In some embodiments, in step (i) the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (i) the organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (i) the organic solvent is acetonitrile.

[0062] In some embodiments, step (i) comprises combining tetrabutylammonium bromide with the compound of formula (F), a source of palladium and 4-chlorophenylboronic acid in the organic solvent.

[0063] In some embodiments, in step (j) the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran,

and mixtures thereof. In some embodiments, in step (j) the organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (j), the organic solvent is a mixture of THF and toluene. In some embodiments, the mixture of THF and toluene is about 1:1 by volume.

[0064] In some embodiments, step (j) further comprises producing the compound of formula (H) as a crystalline solid. In some embodiments, step (j) further comprises:

1. (1) adding an aqueous solution to the mixture of step (j) to produce an aqueous and an organic phase;
2. (2) separating the organic phase from the mixture of step (1);
3. (3) concentrating the organic phase; and
4. (4) adding an organic solvent to the mixture of step (3) to produce the compound of formula (H) as a crystalline solid.

[0065] In some embodiments of step (4) of step (j), the organic solvent is acetonitrile. In some embodiments of step (4) of step (j), the organic solvent is acetonitrile and the mixture is heated to about 80 °C.

[0066] In some embodiments, step (4) of step (j) further comprises cooling the mixture to about 10 °C to about -10 °C. In some embodiments, step (4) of step (j) further comprises cooling the mixture to about -10 °C, and isolating the compound of formula (H) as a crystalline solid by filtering the mixture.


[0067] In some embodiments, the combining of step (k) is in an organic solvent. In some embodiments, the organic solvent is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, the organic solvent is isopropanol.

[0068] In some embodiments, step (k) further comprises producing the compound of formula (I) as a crystalline solid. In some embodiments, the combining of step (k) is in an organic solvent, and step (k) further comprises isolating the compound of formula (I) as a crystalline solid by filtering the mixture.

[0069] In some embodiments, the combining of step (k) is in an organic solvent, and step (k) further comprises cooling the mixture to about 10 °C to about -10 °C to produce the compound of formula (I) as a crystalline solid.

[0070] In some embodiments, the combining of step (k) is in isopropanol, and step (k) further comprises cooling the mixture to about 10 °C to about -10 °C to produce the compound of formula (I) as a crystalline solid. In some embodiments, the combining of step (k) is in isopropanol, and step (k) further comprises cooling the mixture to about -5 °C to produce the compound of formula (I) as a crystalline solid, and isolating the compound of formula (I) as a crystalline solid by filtering the mixture.


[0071] Also disclosed herein is a process of preparing a compound of formula (C):



wherein R is C₁ to C₁₂ alkyl,

which comprises

1. (a) combining a compound of formula (A):



with R¹MgX in an aprotic organic solvent; wherein R¹ is C₁ to C₆ alkyl; and X is Cl, Br or I; and

2. (b) combining a C₁ to C₁₂ alkyl chloroformate or a di-(C₁ to C₁₂ alkyl)dicarbonate with the product of step (a), to provide the compound of formula (C).

[0072] In some aspects, R is C₁ to C₆ alkyl. In some aspects, R is C₁ to C₄ alkyl. In some aspects, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl. In some aspects, R is tert-butyl.

[0073] In some aspects, R¹ is C₁ to C₄ alkyl. In some aspects, R¹ is isopropyl.

[0074] In some aspects of the process of preparing a compound of formula (C), the organic solvent of step (a) is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some aspects the organic solvent of step (a) is THF.

[0075] In one aspects, R is C₁ to C₆ alkyl.

[0076] In one aspect, R is selected from the group consisting of methyl, ethyl, n-propyl,


isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl.

[0077] In one aspect, R is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, iso-butyl and neo-butyl; and R¹ is isopropyl.

[0078] In one aspect, R is tert-butyl and R¹ is isopropyl.


[0079] In some aspect of the process of preparing a compound of formula (C), in step (b), R is methyl and the C₁ to C₁₂ alkyl chloroformate is methyl chloroformate. In some aspect, R is ethyl and the C₁ to C₁₂ alkyl chloroformate is ethyl chloroformate. In some aspect, R is tert-butyl and the di-(C₁ to C₁₂ alkyl)dicarbonate is di-tert-butyl dicarbonate.

[0080] Also provided herein is a process for the preparation of a compound of formula (D):




wherein R is C₁ to C₁₂ alkyl, which comprises:

(x) combining a compound of formula (B):



with a compound of formula (C):




and a *tert*-butoxide salt in an organic solvent to provide the compound of formula (D).

[0081] In one aspect, R is tert-butyl.

[0082] In some aspects, the process of preparing the compound of formula (D) further comprises steps (x') and (x''):

(x') combining a compound of formula (A):



with R¹MgX in an aprotic organic solvent; wherein R¹ is C₁ to C₆ alkyl; and X is Cl, Br or I;

(x'') combining a C₁ to C₁₂ alkyl chloroformate or a di-(C₁ to C₁₂ alkyl)dicarbonate with the product of step (x'), to provide the compound of formula (C).

[0083] In some aspects, in step (x) the tert-butoxide salt is selected from the group consisting of sodium tert-butoxide and potassium tert-butoxide.


[0084] In some aspects, the organic solvent of step (x) is selected from the group consisting of pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, THF, DMF, HMPA, NMP, nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, MTBE, benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, the organic solvent of step (x) is DMF.

[0085] In some aspects, in step (x'), R¹ is a C₁ to C₄ alkyl. In some embodiments, R¹ is isopropyl.

[0086] In some aspects, in step (x''), the C₁ to C₁₂ alkyl chloroformate is methyl chloroformate. In some embodiments, the C₁ to C₁₂ alkyl chloroformate is ethyl chloroformate. In some embodiments, the di-(C₁ to C₁₂ alkyl)dicarbonate is di-tert-butyl dicarbonate.


[0087] In some aspects, in step (x') the aprotic organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, in step (x') the aprotic organic solvent is THF.

[0088] Also provided herein is a compound of the formula (3):




[0089] In one aspect, the compound of the formula (3) is prepared by the following steps:

(y) combining a compound of formula (B):




with a compound of formula (C):




wherein R is tert-butyl, and a *tert*-butoxide salt in an organic solvent to provide the compound

of formula (D):




wherein R is tert-butyl; and

(z) combining the compound of formula (D), wherein R is tert-butyl; with a compound of formula (I):




a source of palladium, a tert-butoxide salt, and a phosphine ligand in an aprotic organic solvent.

[0090] In one aspect, the phosphine ligand of step (z) is a compound of formula (J):



[0091] In other aspects, the phosphine ligand is selected from:



[0092] In one aspect, in step (z) the source of palladium is Pd_2dba_3 .

[0093] In some aspects, in step (z) the aprotic organic solvent is selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, the aprotic organic solvent is a mixture of THF and toluene.

[0094] In some aspects, in step (z), the tert-butoxide salt is selected from the group consisting of sodium tert-butoxide and potassium tert-butoxide.

[0095] In some aspects, in step (z) the tert-butoxide salt is anhydrous sodium tert-butoxide or anhydrous potassium tert-butoxide.

[0096] In some aspects, step (z) further comprises the following steps:

1. (1) combining the tert-butoxide salt with the compound of formula (I) in an aprotic organic solvent;
2. (2) combining the source of palladium, the compound of formula (J), and the compound of formula (D) in an aprotic organic solvent; and
3. (3) adding the mixture of step (1) to the mixture of step (2).


[0097] In some aspects, in step (z) the mixture resulting from step (2) is filtered prior to step (3).

[0098] In some aspects, step (z) is carried out under an atmosphere of nitrogen or argon.

[0099] In some aspects, in step (z) a catalytic amount of the source of palladium is used relative to the amount of compound (I). In some aspects, the source of palladium is Pd_2dba_3 and the catalytic amount of Pd_2dba_3 is from about 0.5 mole percent to about 2 mole percent. In one aspect, the catalytic amount of Pd_2dba_3 is about 0.75 mole percent.

[0100] In some aspects, when the phosphine ligand of step (z) is a compound for formula (J), a catalytic amount of the compound of formula (J) is used relative to the amount of compound (I). In some aspects, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 5 mole percent. In one aspect, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 4 mole percent. In one aspect, the catalytic amount of the compound of formula (J) is from about 2 mole percent to about 4 mole percent. In one aspect, the catalytic amount of the compound of formula (J) is from about 1 mole percent to about 2 mole percent. In one aspect, the catalytic amount of the compound of formula (J) is about 1 mole percent or about 2 mole percent.

[0101] In another aspect of the instant disclosure, are provided herein are compounds of the formulae:




[0102] The processes described herein is directed to improved methods for commercial chemical manufacturing of Compound 1 or Compound 2. Without being bound to a particular theory or mechanism of action, the processes described herein significantly improve the overall

efficiency and product yield of Compound 1 or Compound 2. Previous processes (e.g., U.S. Patent Publication Nos. 2010/0305122 and 2012/0157470, and International Patent Publication Nos. WO 2011/15096 and WO 2012/071336) were found to lack feasibility for production of Compound 1 on a commercial scale. Thus, the processes provided herein represent improved methods for the synthesis of compounds in quantities required for clinical and/or commercial development. Improvements relative to these previous processes include, but are not limited to, overall yield of Compound 1 or Compound 2, overall process efficiency and economics, mild reaction conditions, practical isolation/purification procedures, and viability for commercialization.

[0103] The improved process provided herein involves a selective nucleophilic aromatic substitution reaction ("SnAr reaction") of compounds (B) and (C), which can be carried out under milder conditions with a shorter reaction time when compared to previously described processes as found, for example, in U.S. Patent Publication Nos. 2010/0305122 and 2012/0157470, and International Patent Publication Nos. WO 2011/15096 and WO 2012/071336. Without being limited by theory, the improved SnAr reaction of compound (B) and (C) does not generate regioisomeric side products which necessitate further purification to remove the side products, as was the case in previously described processes. The SnAr reaction in the previous process also requires a longer reaction time and harsh reaction conditions which result in a low overall yield relative to the processes described herein. Furthermore, the previous processes also require tedious purification of the intermediates which is impracticable on a large, commercial scale process. The processes described herein are more convergent than prior processes, resulting in a highly efficient cross-coupling reaction of compound (D) and the free base of compound (I) in high yield. According to some aspects, the processes described herein utilize crystalline solid intermediates (H) and (I), which allow efficient purification by crystallization to remove impurities - advantages not available in previously described processes.

[0104] The following schemes illustrate one or more embodiments of the process provided herein. According to some aspects, the compound of formula (D) is prepared from compound (B) and compound (C) as shown in Scheme 1 below. The compound of formula (B) may be prepared by techniques known in the art, e.g., as shown in WO 2000/047212 and J. Am. Chem. Soc., 1959, 81: 743-747. The compound of formula (C) may be prepared by techniques known in the art, e.g., as shown in WO 2006/059801 and Tetrahedron Letters, 2008, 49(12), 2034-2037; or as shown in Scheme 2.


Scheme 1




[0105] The compound of formula (C) of Scheme 1 may be prepared from commercially available

compound (A) as shown in Scheme 2 below, wherein " R^1MgX " represents a Grignard reagent wherein R^1 is an alkyl group, and X is Cl, Br or I. The electrophilic acetylating reagent of Scheme 2 can be, but is not limited to, methyl or ethyl chloroformate or BOC_2O .

Scheme 2




[0106] An exemplary reaction according to Scheme 2 is shown below.




[0107] In another embodiment, the compound of formula (I) is prepared from compound (E) as shown in Scheme 3 below. Compound (E) is commercially available or may be prepared by techniques known in the art, e.g., as shown in U.S. 3,813,443 and Proceedings of the Chemical Society, London, 1907, 22, 302.

Scheme 3



[0108] In another embodiment, the compound of formula (N) is prepared from compound (M) as shown in Scheme 4 below. Compound (M) is commercially available or may be prepared by techniques known in the art, e.g., as shown in GB 585940 and J. Am. Chem. Soc., 1950, 72, 1215-1218.


Scheme 4



[0109] In another embodiment, the compound of formula (P) is prepared from compound (M) as shown in Scheme 4' below.


Scheme 4'






[0110] In another embodiment, the compound of formula (1) is prepared from compound (D) and compound (I) as shown in Scheme 5 below. Compound (J) may be prepared by techniques known in the art, e.g., as shown in WO 2009/117626 and Organometallics, 2008, 27(21), 5605-5611.

Scheme 5



[0111] In another embodiment, the compound of formula (2) is prepared from compound (L) and compound (P) as shown in Scheme 6 below, wherein the preparation of compound (P) is as shown in Scheme 4' and the preparation of compound (L) is as shown in Scheme 5.

Scheme 6



[0112] In some embodiments, the preparation of the compound of formula (K) from compound (D) and compound (I) is air and/or moisture sensitive, and is therefore performed under an inert atmosphere, e.g., using nitrogen or argon gas.

[0113] Without being bound to a particular theory, the use of compound (D) as an intermediate in the preparation of the compound of formula (1) and the compound of formula (2) as shown above in Schemes 1 to 6 is an improvement over previously described processes for the preparation of the compound of formula (1) and the compound of formula (2). In some embodiments, the improvements include higher product yields, shorter reaction times. In some embodiments, the improvements are provided when R is tert-butyl in compound (D).

[0114] Schemes 1 to 6 are non-limiting examples of the process provided herein. Solvents and/or reagents are known compounds and may be interchanged according to the knowledge of those skilled in the art.

[0115] Abbreviations used in Schemes 1 to 6 are as follows:

Ac	acetyl
BOC	tert-butoxycarbonyl
dba	dibenzylideneacetone
DIEA	N,N-diisopropylethylamine
DMAP	4-dimethylaminopyridine
DMF	dimethylformamide
EDAC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide HCl
IPA	isopropanol
iPr	isopropyl
Me	methyl

n-Bu	n-butyl
tBu	tert-butyl
THF	tetrahydrofuran

[0116] Unless indicated otherwise, the temperatures at which a reaction of Schemes 1 to 6 is conducted is not critical. In certain embodiments, when a temperature is indicated in a reaction, the temperature may be varied from about plus or minus 0.1 °C, 0.5 °C, 1 °C, 5 °C, or 10 °C. Depending upon which solvent is employed in a particular reaction, the optimum temperature may vary. In some embodiments, reactions are conducted in the presence of vigorous agitation sufficient to maintain an essentially uniformly dispersed mixture of the reactants.

[0117] In conducting a reaction provided herein, neither the rate, nor the order, of addition of the reactants is critical unless otherwise indicated. Unless otherwise indicated, reactions are conducted at ambient atmospheric pressure. Unless otherwise indicated, the exact amount of reactants is not critical. In some embodiments, the amount of a reactant may be varied by about 10 mole percent or about 10% by weight.

[0118] Unless otherwise indicated, the organic solvents used in the processes provided herein may be selected from those commercially available or otherwise known to those skilled in the art. Appropriate solvents for a given reaction are within the knowledge of the skilled person and include mixtures of solvents. Examples of organic solvents provided herein for use include but are not limited to: pentane, hexane, heptane, cyclohexane, methanol, ethanol, 1-propanol, isopropanol, 1-butanol, 2-butanol, tert-butanol, 2-butanone, dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, tetrahydrofuran (THF), dimethylformamide (DMF), hexamethylphosphoramide (HMPA), N-methyl-2-pyrrolidinone (NMP), nitromethane, acetone, acetic acid, acetonitrile, ethyl acetate, diethyl ether, diethylene glycol, glyme, diglyme, petroleum ether, dioxane, methyl tert-butyl ether (MTBE), benzene, toluene, xylene, pyridine, 2-methyltetrahydrofuran, and mixtures thereof.

[0119] In some embodiments, an organic solvent used in the processes provided herein is an aprotic organic solvent. As provided herein, an aprotic solvent is a solvent that does not contain an acidic hydrogen atom or a hydrogen atom that is capable of hydrogen bonding (e.g., is not bound to an oxygen or a nitrogen atom). The aprotic organic solvent may be selected from the group consisting of dichloromethane, chloroform, acetone, acetonitrile, THF, DMF, NMP, HMPA, dioxane, nitromethane, pyridine, 2-methyltetrahydrofuran, and mixtures thereof. In some embodiments, the aprotic organic solvent is THF. In some embodiments, the aprotic organic solvent is DMF. In some embodiments, the aprotic organic solvent is acetonitrile.

[0120] As provided herein, a "tertiary amine base" refers to an amine that is substituted with three alkyl groups, e.g., triethylamine or N,N-diisopropylethylamine.

[0121] As provided herein, a "catalytic amount" refers to less than one molar equivalent of a reagent or reactant in a given reaction, as determined relative to another reagent or reactant in the reaction mixture. In some embodiments, a catalytic amount is described as a mole percent relative to another reagent or reactant in the reaction mixture.

[0122] As provided herein, a "source of palladium" refers to a source of palladium in a stable oxidation state, *i.e.*, Pd(0), Pd(I), Pd(II) and/or Pd(IV). The palladium may be free metal, such as in a powder form, or may be bound to one or more ligands, *e.g.*, PdCl₂, Pd₂dba₃, PdCl₂(PPh₃)₂, Pd(PPh₃)₄, Pd(OAc)₂ or [(cinnamyl)PdCl]₂.

[0123] As provided herein, a "phosphine ligand" refers to a compound of formula PR'₃, wherein each R' is independently selected from C₁ to C₆ alkyl or phenyl, wherein the aryl group is optionally substituted by C₁ to C₆ alkyl, phenyl, trialkylamino, alkoxy or halo.

[0124] As provided herein, unless otherwise defined, the term "about" means that the value or amount to which it refers can vary by 5%, $\pm$ 2%, or $\pm$ 1%.

[0125] The products obtained by any of the processes provided herein may be recovered by conventional means, such as evaporation or extraction, and may be purified by standard procedures, such as distillation, recrystallization or chromatography

EXAMPLES

[0126] Compounds of the following examples are shown in Schemes 1 to 6 above and were named using Chemdraw® Ultra software. In addition to the abbreviations described above with respect to the schemes provided herein, the following abbreviations are used in the Examples:

[0127] "HPLC" = high pressure liquid chromatography; "IP" = in process; "ML" = mother liquor; "NLT" = no less than; "NMT" = no more than; "RB" = round bottom; "R" = room temperature; "sm" = starting material; "DCM" = dichloromethane.

[0128] Unless indicated otherwise, compounds were characterized by HPLC and ¹H NMR analysis and used in later reactions with or without purification. ¹H NMR analysis was performed at 400 MHz unless otherwise indicated. Unless specified otherwise, product yield/purity was determined by weight, qNMR, and/or HPLC analysis.

Example 1: Synthesis of tert-butyl 4-bromo-2-fluorobenzoate (Compound (C))

[0129] To a 100 ml jacketed reactor equipped with a mechanical stirrer was charged 4-bromo-2-fluorol-iodobenzene, "Compound (A)" (5 g, 1.0 eq) and THF (25 ml). The solution was

cooled to -5 °C. 2 M isopropyl magnesium chloride in THF (10.8 ml, 1.3 eq) was slowly added maintaining the internal temperature below 0 °C. The mixture was stirred at 0 °C for 1 h. Di-tert-butyl dicarbonate (5.44 g, 1.5 eq) in THF (10 ml) was added. After 1 h, the solution was quenched with 10 % citric acid (10 ml), and then diluted with 25 % NaCl (10 ml). The layers were separated and the organic layer was concentrated to near dryness and chased with THF (3 x 10 ml). The crude oil was diluted with THF (5 ml), filtered to remove inorganics, and concentrated to dryness. The crude oil (6.1 g, potency = 67%, potency adjusted yield = 88%) was taken to the next step without further purification. ¹H NMR (DMSO-d₆): δ 1.53 (s, 9H), 7.50-7.56 (m, 1H), 7.68 (dd, J=10.5, 1.9 Hz, 1H), 7.74 (t, J= 8.2 Hz, 1H).

Example 2: Synthesis of tert-butyl 2-((1H-pyrrolo[2,3-b]pyridin-5-yl)oxy)-4-bromobenzoate (Compound (D))

[0130] To a 3 L three-neck Morton flask were charged 1H-pyrrolo[2,3-b]pyridin-5-ol (80.0 g, 1.00 eq.), tert-butyl 4-bromo-2-fluorobenzoate (193 g, 1.15 eq.), and anhydrous DMF (800 mL). The mixture was stirred at 20 °C for 15 min. The resulting solution was cooled to about zero to 5 °C. A solution of sodium tert-butoxide (62.0 g) in DMF (420 mL) was added slowly over 30 min while maintaining the internal temperature at NMT 10 °C, and rinsed with DMF (30 mL). The reaction mixture was stirred at 10 °C for 1 hour (an off-white slurry) and adjusted the internal temperature to ~ 45 °C over 30 min. The reaction mixture was stirred at 45-50 °C for 7 hr and the reaction progress monitored by HPLC (IP samples: 92% conversion % by HPLC). The solution was cooled to ~ 20 °C. The solution was stirred at 20 °C overnight.

[0131] Water (1200 mL) was added slowly to the reaction mixture at <30 °C over 1 hour (slightly exothermic). The product slurry was adjusted to ~ 20 °C, and mixed for NLT 2 hours. The crude product was collected by filtration, and washed with water (400 mL). The wet-cake was washed with heptane (400 mL) and dried under vacuum at 50 °C overnight to give the crude product (236.7g).

[0132] Re-crystallization or Re-slurry: 230.7 g of the crude product, (potency adjusted: 200.7 g) was charged back to a 3L three-neck Morton flask. Ethyl acetate (700 mL) was added, and the slurry heated slowly to refluxing temperature over 1hr (small amount of solids left). Heptane (1400 mL) was added slowly, and the mixture adjusted to refluxing temperature (78 °C). The slurry was mixed at refluxing temperature for 30 min., and cooled down slowly to down to ~ -10 °C at a rate of approximate 10 °C/hour), and mixed for 2hr. The product was collected by filtration, and rinsed with heptane (200 ml).

[0133] The solid was dried under vacuum at ~ 50 °C overnight to give 194.8 g, 86% isolated yield of the product as an off-white solid. MS-ESI 389.0 (M+1); mp: 190-191°C (uncorrected). ¹H NMR (DMSO-d₆): δ 1.40 (s, 9H), 6.41 (dd, J= 3.4, 1.7 Hz, 1H), 7.06 (d, J=1.8 Hz, 1H), 7.40 (dd, J= 8.3, 1.8 Hz, 1H), 7.51 (t, J= 3.4 Hz, 1H), 7.58 (d, J=2.6 Hz, 1H), 7.66 (d, J=8.3 Hz, 1H), 8.03 (d, J=2.7 Hz, 1H), 11.72 (s, 1H, NH).

Example 3: Synthesis of 2-chloro-4,4-dimethylcyclohexanecarbaldehyde (Compound (F))

[0134] To a 500 mL RB flask were charged anhydrous DMF (33.4 g, 0.456mol) and CH₂Cl₂ (80 mL). The solution was cooled down <-5 °C, and POCl₃ (64.7 g, 0.422 mol) added slowly over 20 min @ <20 °C (exothermic), rinsed with CH₂Cl₂ (6 mL). The slightly brown solution was adjusted to 20 °C over 30 min, and mixed at 20 °C for 1 hour. The solution was cooled back to < 5 °C. 3,3-Dimethylcyclohexanone (41.0 g, 90%, ~0.292 mol) was added, and rinsed with in CH₂Cl₂ (10 mL) (slightly exothermic) at <20 °C. The solution was heated to refluxing temperature, and mixed overnight (21 hours.).

[0135] To a 1000 mL three neck RB flask provided with a mechanical stirrer were charged 130 g of 13.6 wt % sodium acetate trihydrate aqueous solution, 130 g of 12% brine, and 130 mL of CH₂Cl₂. The mixture was stirred and cooled down to <5 °C. The above reaction mixture (clear and brown) was transferred, quenched into it slowly while maintaining the internal temperature <10 °C. The reaction vessel was rinsed with CH₂Cl₂ (10 mL). The quenched reaction mixture was stirred at <10 °C for 15 min. and allowed to rise to 20 °C. The mixture was stirred 20 °C for 15 min and allowed to settle for 30 min. (some emulsion). The lower organic phase was separated. The upper aq. phase was back extracted with CH₂Cl₂ (50 mL). The combined organic was washed with a mixture of 12% brine (150 g)-20% K₃PO₄ aq. solution (40 g). The organic was dried over MgSO₄, filtered and rinsed with CH₂Cl₂ (30 mL). The filtrate was concentrated to dryness under vacuum to give a brown oil (57.0 g, potency=90.9wt% by qNMR, ~100%). ¹H NMR (CDCl₃) δ 0.98 (s, 6H), 1.43 (t, J=6.4 Hz, 2H), 2.31 (tt, J =6.4, 2.2 Hz, 2H), 2.36 (t, J =2.2 Hz, 2H), 10.19 (s, 1H).

Example 4: Synthesis of 2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-enecarbaldehyde (Compound (G))

[0136] To a 250 mL pressure bottle were charged 2-chloro-4,4-dimethylcyclohex-1-enecarbaldehyde (10.00 g), tetrabutylammonium bromide (18.67 g), and acetonitrile (10 mL). The mixture was stirred at 20 °C for 5 min. 21.0 wt% K₂CO₃ aq. solution (76.0 g) was added. The mixture was stirred at room temperature (rt) for NLT 5 min. followed by addition of 4-chlorophenylboronic acid (9.53 g) all at once. The mixture was evacuated and purged with N₂ for three times. Palladium acetate (66 mg, 0.5 mol %) was added all at once under N₂. The reaction mixture was evacuated and purged with N₂ for three times (an orange colored mixture). The bottle was back filled with N₂ and heated to ~35 °C in an oil bath (bath temp ~35 °C). The mixture was stirred at 30 °C overnight (15 hours). The reaction mixture was cooled to RT, and pulled IP sample from the upper organic phase for reaction completion, typically

starting material < 2% (orange colored mixture). Toluene (100 mL) and 5% NaHCO₃-2% L-Cysteine aq. solution (100 mL) were added. The mixture was stirred at 20 °C for 60 min. The mixture was filtered through a pad of Celite to remove black solid, rinsing the flask and pad with toluene (10 mL). The upper organic phase was washed with 5% NaHCO₃ aq. solution-2% L-Cysteine (100 mL) once more. The upper organic phase was washed with 25% brine (100 mL). The organic layer (105.0 g) was assayed (118.8 mg/g, 12.47 g product assayed, 87% assayed yield), and concentrated to ~1/3 volume (~ 35 mL). The product solution was directly used in the next step without isolation. However, an analytical sample was obtained by removal of solvent to give a brown oil. ¹HNMR (CDCl₃) δ 1.00 (s, 6H), 1.49 (t, J=6.6 Hz, 2H), 2.28 (t, J=2.1 Hz, 2H), 2.38 (m, 2H), 7.13 (m, 2H), 7.34 (m, 2H), 9.47 (s, 1H).

Example 5: Synthesis of tert-butyl 4-((4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methyl)piperazine-1-carboxylate (Compound (H))

[0137] To a 2 L three neck RB flask provided with a mechanical stirrer were charged a solution of 4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-carbaldehyde (50.0g) in toluene (250 mL), BOC-piperazine (48.2 g) and anhydrous THF (250 mL). The yellow solution was stirred at 20 °C for 5 min. Sodium triacetoxyborohydride (52.7 g) was added in portion (note: the internal temperature rose to ~29.5 °C in 15 min cooling may be needed). The yellow mixture was stirred at ~ 25 °C for NLT 4hrs. A conversion of starting material to product of 99.5% was observed by HPLC after a 3 hour reaction time.

[0138] 12.5 wt % brine (500 g) was added slowly to quench the reaction. The mixture was stirred at 20 °C for NLT 30 min and allowed to settle for NLT 15 min. The lower aq. phase (~560 mL) was separated (note: leave any emulsion in the upper organic phase). The organic phase was washed with 10% citric acid solution (500 g x2). 500 g of 5% NaHCO₃ aq. solution was charged slowly into the flask. The mixture was stirred at 20 °C for NLT 30 min., and allowed to settle for NLT 15 min. The upper organic phase was separated. 500 g of 25% brine aq. solution was charged. The mixture was stirred at 20 °C for NLT 15 min., and allowed to settle for NLT 15 min. The upper organic phase was concentrated to ~ 200 mL volume under vacuum. The solution was adjusted to ~ 30 °C, and filtered off the inorganic salt. Toluene (50 mL) was used as a rinse. The combined filtrate was concentrated to ~100 mL volume. Acetonitrile (400 mL) was added, and the mixture heated to ~ 80 °C to achieve a clear solution. The solution was cooled down slowly to 20 °C slowly at rate 10 °C/hour, and mixed at 20 °C overnight (the product is crystallized out at ~45-50 °C, if needed, seed material may be added at 50 °C). The slurry was continued to cool down slowly to ~ -10 °C at rate of 10 °C/hours. The slurry was mixed at ~ -10 °C for NLT 6 hours. The product was collected by filtration, and rinsed with pre-cooled acetonitrile (100mL). The solid was dried under vacuum at 50 °C overnight (72.0 g, 85%). MS-ESI: 419 (M+1); mp: 109-110 °C (uncorrected); ¹H NMR (CDCl₃) δ 1.00 (s, 6H), 1.46 (s, 9H), 1.48 (t, J=6.5 Hz, 2H), 2.07 (s, br, 2H), 2.18 (m, 4H), 2.24 (t, J=6.4 Hz, 2H), 2.80 (s, 2H), 3.38 (m, 4H), 6.98 (m, 2H), 7.29 (m, 2H).

Example 6: Synthesis of 1-((4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methyl)piperazine dihydrochloride (Compound (I))

[0139] To a 2.0 L three-neck RB flask equipped with a mechanical stirrer were charged the Boc reductive amination product (Compound (H), 72.0 g) and IPA (720 mL). The mixture was stirred at rt for 5 min, and 59.3g of concentrated hydrochloride aq. solution added to the slurry. The reaction mixture was adjusted to an internal temperature of ~ 65 °C (a clear and colorless solution achieved). The reaction mixture was agitated at ~ 65 °C for NLT 12 hours.

[0140] The product slurry was cooled down to -5 °C slowly (10 °C/hour). The product slurry was mixed at ~ -5 °C for NLT 2 hours, collected by filtration. The wet cake was washed with IPA (72 mL) and dried at 50 °C under vacuum overnight to give 73.8 g (95%) of the desired product as a bis-hydrochloride IPA solvate (purity >99.5% peak area at 210 nm). MS-ESI: 319 (M+1); ¹HNMR (D₂O): δ 1.00 (s, 6H), 1.19 (d, J=6.0 Hz, 6H, IPA), 1.65 (t, J=6.1 Hz, 2H), 2.14 (s, br, 2H), 2.26 (m, 2H), 3.36 (br, 4H), 3.55 (s, br, 4H), 3.82 (s, 2H), 4.02 (septet, J=6.0 Hz, 1H, IPA), 7.16 (d, J=8.1 Hz, 2H), 7.45 (d, J=8.1 Hz, 2H); ¹HNMR (CDCl₃) δ 0.86 (s, 6H), 1.05 (d, J=6.0 Hz, 6H, IPA), 1.42 (t, J=6.1 Hz, 2H), 2.02 (s, br, 2H), 2.12 (m, 2H), 3.23 (m, 4H), 3.4 (s, br, 4H), 3.68 (s, 2H), 3.89 (septet, J=6.0 Hz, 1H, IPA), 7.11 (d, J=8.1 Hz, 2H), 7.41 (d, J=8.1 Hz, 2H).

Example 7: Synthesis of 3-nitro-4-(((tetrahydro-2H-pyran-4-yl)methyl)amino)-benzenesulfonamide (Compound (N))

[0141] To a 500 mL three-neck RB flask equipped with a mechanical stirrer were charged the 4-chloro-3-nitrobenzenesulfonamide, Compound M (10.0 g), diisopropylethylamine (17.5 g), (tetrahydro-2H-pyran-4-yl)methanamine (7.0 g) and acetonitrile (150 mL). The reaction mixture was adjusted to an internal temperature of 80 °C and agitated for no less than 12 hours.

[0142] The product solution was cooled down to 40 °C and agitated for no less than 1 hour until precipitation observed. The product slurry was further cooled to 20 °C. Water (75 mL) was slowly charged over no less than 1 hour, and the mixture cooled to 10 °C and agitated for no less than 2 hours before collected by filtration. The wet cake was washed with 1:1 mix of acetonitrile:water (40 mL). The wet cake was then reslurried in water (80 mL) at 40 °C for no less than 1 hour before collected by filtration. The wet cake was rinsed with water (20 mL), and dried at 75 °C under vacuum to give 12.7 g of the desired product in 99.9% purity and in 91% weight-adjusted yield. ¹H NMR (DMSO-d₆): δ 1.25 (m, 2H), 1.60 (m, 2H), 1.89 (m, 1H), 3.25 (m, 2H), 3.33 (m, 2H), 3.83 (m, 2H), 7.27 (d, J=9.3 Hz, 1H), 7.32 (s, NH₂, 2H), 7.81 (dd, J=9.1, 2.3 Hz, 1H), 8.45 (d, J=2.2 Hz, 1H), 8.54 (t, J=5.9 Hz, 1H, NH).

Example 8: Synthesis of tert-butyl 2-((1H-pyrrolo[2,3-b]pyridin-5-yl)oxy)-4-(4-((4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methyl)piperazin-1-yl)benzoate (Compound (K))

[0143] *General Considerations:* this chemistry is considered air and moisture sensitive. While the catalyst precursors in their solid, dry form can be handled and stored in air without special precautions, contact with even small amounts of solvent may render them susceptible to decomposition. As a result, traces of oxygen or other competent oxidants (e.g., solvent peroxides) must be removed prior to combination of the catalyst precursors with solvent and care must be used to prevent ingress of oxygen during the reaction. Also, care must be taken to use dry equipment, solvents, and reagents to prevent formation of undesirable byproducts. The sodium t-butoxide used in this reaction is hygroscopic and it should be properly handled and stored prior to or during use.

[0144] To a 2.0 L three-neck RB flask equipped with a mechanical stirrer were charged the bis-hydrochloride salt (Compound (I), 42.5 g) and toluene (285 ml). 20% K₃PO₄ (285 ml) was added and the biphasic mixture was stirred for 30 min. The layers were separated and the organic layer was washed with 25 % NaCl (145 ml). The organic layer concentrated to 120 g and used in the coupling reaction without further purification.

[0145] NaOtBu (45.2 g) and Compound (I) in toluene solution (120 g solution-30 g potency adjusted) were combined in THF (180 ml) in a suitable reactor and sparged with nitrogen for NLT 45 min. Pd₂dba₃ (0.646 g), Compound (J) (0.399 g), and Compound (D) (40.3 g) were combined in a second suitable reactor and purged with nitrogen until oxygen level was NMT 40 ppm. Using nitrogen pressure, the solution containing Compound (I) and NaOtBu in toluene/THF was added through a 0.45 µm inline filter to the second reactor (catalyst, Compound (J) and Compound (D)) and rinsed with nitrogen sparged THF (30 ml.).

[0146] The resulting mixture was heated to 55 °C with stirring for NLT 16 h, then cooled to 22 °C. The mixture was diluted with 12% NaCl (300 g) followed by THF (300 ml). The layers were separated.

[0147] The organic layer was stirred with a freshly prepared solution of L-cysteine (15 g), NaHCO₃ (23 g), and water (262 ml). After 1 h the layers were separated.

[0148] The organic layer was stirred with a second freshly prepared solution of L-cysteine (15 g), NaHCO₃ (23 g), and water (262 ml). After 1 h the layers were separated. The organic layer was washed with 12 % NaCl (300 g), then filtered through a 0.45 µm inline filter. The filtered solution was concentrated *in vacuo* to ~ 300 mL, and chased three times with heptane (600 mL each) to remove THF.

[0149] The crude mixture was concentrated to 6 volumes and diluted with cyclohexane (720

ml). The mixture was heated to 75 °C, held for 15 min, and then cooled to 65 °C over NLT 15 min. Seed material was charged and the mixture was held at 65 °C for 4 hours. The suspension was cooled to 25 °C over NLT 8 h, then held at 25 °C for 4 hours. The solids were filtered and washed with cyclohexane (90 ml) and dried at 50 °C under vacuum.

[0150] Isolated 52.5 g (88.9% yield) as a white solid. Melting point (uncorrected) 154-155 °C. ¹H NMR (DMSO-d₆): δ 0.93 (s, 6H), 1.27 (s, 9H), 1.38 (t, J= 6.4 Hz, 2H), 1.94 (s, 2H), 2.08-2.28 (m, 6H), 2.74 (s, 2H), 3.02 - 3.19 (m, 4H), 6.33 (dd, J= 3.4, 1.9 Hz, 1H), 6.38 (d, J= 2.4 Hz, 1H), 6.72 (dd, J= 9.0, 2.4 Hz, 1H), 6.99 - 7.06 (m, 2H), 7.29 (d, J= 2.7 Hz, 1H), 7.30 - 7.36 (m, 2H), 7.41 - 7.44 (m, 1H), 7.64 (t, J= 6.7 Hz, 1H), 7.94 (d, J= 2.7 Hz, 1H), 11.53 (s, 1H).

Example 9: Synthesis of 2-((1H-pyrrolo[2,3-b]pyridin-5-yl)oxy)-4-(4-((4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methyl)piperazin-1-yl)benzoic acid (Compound (L))

[0151] Solution preparation: 10% KH₂PO₄ (aq): KH₂PO₄ (6 g) in water (56 g); 2:1 heptane / 2-MeTHF : heptane (16 mL) in 2-MeTHF (8 mL).

[0152] Compound (K) (5.79 g), potassium tert-butoxide (4.89 g), 2-methyltetrahydrofuran (87 mL), and water (0.45 mL) were combined in a suitable reactor under nitrogen and heated to 55 °C until reaction completion. The reaction mixture was cooled to 22 °C, washed with the 10% KH₂PO₄ solution (31 g) twice. The organic layer was then washed with water (30 g).

[0153] After removal of the aqueous layer, the organic layer was concentrated to 4 volumes (~19 mL) and heated to no less than 50 °C. Heptane (23 ml) was slowly added. Alternatively, after removal of the aqueous layer, the organic layer was concentrated to 5 volumes and heated to no less than 70 °C and 5 volumes of heptane were slowly added. The resulting suspension was cooled to 10 °C. Solids were then collected by vacuum filtration with recirculation of the liquors and the filter cake washed with 2:1 heptane / 2-MeTHF (24 ml). Drying of the solids at 80 °C under vacuum yielded 4.0 g of Compound (L) in approximately 85% weight-adjusted yield. ¹H NMR (DMSO-d₆): δ 0.91 (s, 6H), 1.37 (t, J=6.4 Hz, 2H), 1.94 (s, br, 2H), 2.15 (m, 6H), 2.71 (s, br, 2H), 3.09 (m, 4H), 6.31 (d, J=2.3 Hz, 1H), 6.34 (dd, J=3.4, 1.9 Hz, 1H), 6.7 (dd, J= 9.0, 2.4 Hz, 1H), 7.02 (m, 2H), 7.32 (m, 2H), 7.37 (d, J=2.6 Hz, 1H), 7.44 (t, J= 3.0 Hz, 1H), 7.72 (d, J=9.0 Hz, 1H), 7.96 (d, J=2.7 Hz, 1H) & 11.59 (m, 1H).

Example 10: Synthesis of 4-(4-([2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl)piperazin-1-yl)-N-({3-nitro-4-[(tetrahydro-2H-pyran-4-yl)methyl]amino}phenyl)sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide (Compound (1))

[0154] Solution preparation prior to reaction: 10% Acetic Acid: Acetic Acid (37 mL) in water (333 g); 5% NaHCO₃: NaHCO₃ (9 g) in water (176 g); 5% NaCl : NaCl (9 g) in water (176 g).

[0155] Compound (N) (13.5 g), DMAP (10.5 g), EDAC (10.7 g) and dichloromethane (300 mL) were combined in a suitable reactor and agitated at 25 °C. In a second suitable reactor was charged the Acid (Compound (L), 25 g), Et₃N (8.7 g) and dichloromethane (120 mL). The resulting Acid (Compound (L)) solution was slowly charged to the initial suspension of Compound (N) and agitated until reaction completion. N,N-dimethylethylenediamine (9.4 g) was then charged to the reaction mixture with continued agitation. The reaction mixture was warmed to 35 °C and washed with 10% Acetic acid solution (185 mL) twice. The lower organic layer was diluted with more dichloromethane (75 mL) and methanol (12.5 mL). The organic, product layer was then washed with 5% NaHCO₃ solution (185 mL) and then washed with 5% NaCl solution (185 mL) at 35 °C. The lower, organic layer was separated and then concentrated to 8 vol (~256 mL) diluted with methanol (26 mL) and warmed to 38 °C. Ethyl Acetate (230 mL) was slowly charged. The resulting suspension was slowly cooled to 10 °C and then filtered. The wet cake was washed twice with a 1:1 mix of dichloromethane and ethyl acetate (~2 vol, 64 mL). After drying the wet cake at 90°C, 32 g (84%) of Compound (1) was isolated. ¹H NMR (DMSO-d₆): δ 0.90 (s, 6H), 1.24 (m, 2H), 1.36 (t, J=6.4 Hz, 2H), 1.60 (m, 2H), 1.87 (m, 1H), 1.93 (s, br, 2H), 2.12 (m, 2H), 2.19 (m, 4H), 2.74 (s, br, 2H), 3.06 (m, 4H), 3.26 (m, 4H), 3.83 (m, 2H), 6.17 (d, J=2.1 Hz, 1H), 6.37 (dd, J= 3.4, 1.9 Hz, 1H), 6.66 (dd, J= 9.1, 2.2 Hz, 1H), 7.01 (m, 2H), 7.31 (m, 2H), 7.48 (m, 3H), 7.78 (dd, J= 9.3, 2.3 Hz, 1H), 8.02 (d, J=2. 61 Hz, 1H), 8.54 (d, J=2. 33 Hz, 1H), 8.58 (t, J=5.9 Hz, 1H, NH), 11.65 (m, 1H).

Example 11: Synthesis of ((1*R*,4*R*)-4-hydroxy-4-methylcyclohexyl)-methanaminium 4-methylbenzenesulfonate

[0156] Step A: 1.49 g of cyclohexanedione monoethylene acetal (1.0 equiv) and 15 mL of toluene were charged to a suitable reactor. The mixture was mixed for 30 minutes at 10°C. 1.4 M methylmagnesium bromide solution (2.32 eq) in Toluene-THF (75-25) was charged to another reactor and mixed at 15°C. The starting material solution was added to the Grignard solution dropwise at around 10 to 20°C in 4hrs (addition rate = 0.1mL/min). The reaction progression was monitored by TLC. Upon reaction completion, the reaction mixture was charged to a 24 % ammonium chloride solution (20 mL) slowly at a temperature of 25oC. The reaction mixture was mixed and settled, organic layer was separated and aqueous layer was extracted with ethyl acetate (3 x 20 mL). The combined organic layers were filtered over a bed of sodium sulfate and the filtrate was concentrated by distillation to dryness. 1.57 g. crude solids were isolated (95% yield) and carried to the next step. ¹H NMR (400 MHz, Chloroform-d₁) δ ppm 3.88-4.01 (m, 4H), 1.85-1.96 (m, 2H), 1.08-1.64 (m, 7H). LCMS- (MS 310 and 292). Rf = 0.074 by TLC (hexane-EtOAc = 1-1).

[0157] Step B: 18 mL of 0.005 N hydrochloric acid solution (0.02 equiv) was charged to the distillation residue from Step A. The reaction mixture was mixed at 70°C for 3 hours and

monitored by TLC. Upon reaction completion, the reaction mixture was cooled to 25°C and charged to another suitable reactor containing 22 mL of a 5% sodium chloride solution. The reaction mixture was mixed until all salt dissolved followed by extraction with Ethyl Acetate (8x200 mL). The combined organic layers were filtered over a bed of sodium sulfate and the filtrate was concentrated by distillation to dryness. The product was isolated (99.38 % yield) and was used directly in the next step. ¹H NMR (400 MHz, Chloroform-d₁) δ ppm 2.68-2.80 (m, 2H), 2.16-2.39 (m, 3H), 1.77-2.04 (m, 4H), 1.41 (s, 3H), 1.33 (s, 1H).

[0158] Step C: Step B product (0.25 g) was dissolved with toluene (5 ml) to a 25 mL three neck flask equipped with a Dean-Stark trap. Nitrogen was bubbled through the reactor to remove air. 0.585 g of nitromethane (5 equiv) was charged to the reactor followed by 0.052 g of N,N-dimethylethylenediamine (0.3 equiv). The reaction mixture was heated to reflux, the water was removed by a Dean-Stark trap. The reaction mixture was mixed at reflux for 1 h and monitored by HPLC assay. The reaction mixture was then cooled to 20°C when HPLC product assay stabilized, concentrated then chased with EtOAc and heptane to dryness. The residue was purified on a CombiFlash column (12 g column) from Hexane/EtOAc 80-20 to 60-40. Fractions were analyzed by HPLC and TLC, product containing fractions was distilled to dryness. A concentrated oil 0.23 g was obtained (68.09 % yield) and used in Step D. ¹H NMR (400 MHz, Chloroform-d₁) δ 5.88-5.90 (bs, 1H), 4.88-4.89 (bs, 2H), 2.16-2.40 (m, 4H), 1.78-1.85 (m, 1H), 1.33 (s, 3H).

[0159] Step D: Crabtree's catalyst (0.471 g; 0.585 mmol) was added under nitrogen to a 450 mL stirred SS Parr reactor. The reactor was purged with nitrogen and a solution of the (S)-1-methyl-4-(nitromethyl)cyclohex-3-enol (34.88 g; 58.5 mmol) in DCM (100 mL). Additional sparged DCM (80 mL) was added, the reactor was purged with argon, hydrogen and hydrogen pressure 100 psig. The mixture was agitated for 4 hours at 30°C. Reaction progress was monitored by NMR, concentrated to an oil, chased 2 x with THF (50 mL) then diluted with THF (50 mL). The product was carried further for subsequent RaNi reduction in Step E. ¹H NMR (Chloroform-d₁): δ 4.33 (dJ = 7.3Hz, 2H), 4.32 (J=6.5 Hz, 1H), 2.36 - 2.20 (m, 1H), 1.92 - 1.69 (m, 1H), 1.64 - 1.40 (m, 1H), 1.39 - 1.18 (m, 1H).

[0160] Step E: RaNi (* d / (d-1) or * 7/6) = 2.04 g (20 wt%) was decanted 3 times with THF. The RaNi, solution of (1R,4R)-1-methyl-4-(nitromethyl)cyclohexanol and THF (50 mL) were added under nitrogen in a 450 mL stirred SS Parr reactor. The reactor was purged with nitrogen, hydrogen and the hydrogenation was carried out at 40 psi for 4 hours at 50°C. The reaction was monitored by GC and upon completion, it was filtered through a propylene filter funnel with diatomaceous earth/polyethylene fritted disc to remove catalyst. THF was used as a rinse to extract residual product from the filter cake. The combined filtrate gave an amber solution which was carried directly to next step. ¹H NMR (400 MHz, Chloroform-d₁) δ 2.61 (d, J = 6.5 Hz, 2H), 1.25-1.50 (m, 12H), 0.80-1.17 (m, 3H).

[0161] Step F: 9.86 g of the solution of Step E was added to a 500 mL round bottom flask and

distilled to dryness, chased twice with acetonitrile and then was dissolved in acetonitrile (100 mL). To the solution was added 4-methylbenzenesulfonic acid hydrate (11.68 g) upon which a solid precipitated out and temperature rose to 40°C. The slurry was mixed at 50°C for 2 hours and cooled to 20°C for 12 hours. Solids were filtered and washed with 40 mL acetonitrile. The wetcake was dried under vacuum to give 14.24 g of product (77% yield). ¹H NMR (400 MHz, Deuterium Oxide-d₂) δ 2.79 (d, J = 7.0 Hz, 2H), 1.48-1.68 (m, 5H), 1.31-1.46 (m, 2H), 0.90-1.29 (m, 5H).

Example 12: Synthesis of 4-({[(1R,4R)-4-hydroxy-4-methylcyclohexyl]methyl}-amino)-3-nitrobenzenesulfonamide (Compound (P))

[0162] 4-chloro-3-nitrobenzenesulfonamide (6.5 g, 27.5 mmol) and ((1*R*,4*R*)-4-hydroxy-4-methylcyclohexyl)methanaminium 4-methylbenzenesulfonate (11.26 g, 35.7 mmol) were combined in 35 mL of acetonitrile and stirred. N,N-diisopropylethylamine (8.88 g, 68.7 mmol) was added to the slurry at ambient temperature to result in an endotherm (200 to 17.5°C). After 10 minutes, the reaction mixture was heated to 80°C and maintained at that temperature for 24 hours. The reaction was monitored for completion by HPLC. Upon completion of the reaction, the reaction mixture was cooled to 40°C. Water (32.5 mL) was added over 15 minutes and held for 30 minutes. An additional 74.5 mL of water was added over 30 minutes. Solid product precipitated soon after the second portion of water was added. After stirring for 1 hour at 40°C, the product mixture was allowed to cool to 20°C, stirred for 12 hours, and then cooled to 0°C with stirring for 2 additional hours. The product was filtered and dried under vacuum to afford 8.8 g of product (Yield 93%; purity >99 pa%). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 8.52 (t, J = 5.9 Hz, 1H), 8.45 (d, J = 2.2 Hz, 1H), 7.80 (dd, J = 9.1, 2.3 Hz, 1H), 7.24-7.30 (m, 3H), 4.23 (s, 1H), 1.60-1.74 (m, 3H), 1.52-1.57 (m, 2H), 1.26-1.40 (m, 2H), 1.06-1.25 (m, 5H).

Example 13: Synthesis of 4-(4-{{[2-(4-chlorophenyl)-4,4-dimethylcyclohex-1-en-1-yl]methyl}piperazin-1-yl}-N-({3-nitro-4-[(1R,4R)-[4-hydroxy-4-methylcyclohexyl]methyl}amino]phenyl}sulfonyl)-2-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)benzamide (Compound (2))

[0163] The Sulfonamide 4-((((1*R*,4*R*)-4-hydroxy-4-methylcyclohexyl)methyl)amino)-3-nitrobenzenesulfonamide (8.00 g, 23.29 mmol), EDAC-HCl (5.80 g, 30.3 mmol) and DMAP (8.54 g, 69.9 mmol) were mixed in DCM (186 mL, 14 vol) to a golden slurry. A solution of acid, 2-(((1H-pyrrolo[2,3-b]pyridin-5-yl)oxy)-4-(4'-chloro-5,5-dimethyl-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methyl)piperazin-1-yl)benzoic acid (13.3 g, 23.29 mmol) and TEA (6.49 mL, 46.6 mmol) in DCM (80mL, 6 vol) was added over 2.5 hrs by addition funnel followed by a rinse with 10 mL DCM. After mixing for 12 hours, *N*1,*N*1-dimethylethane-1,2-diamine (5.09 mL, 46.6 mmol) was added and stirring continued at 20°C for 5 hours. The reaction mixture was washed with 10% HOAc (130mL, 3x). The organic layer was washed with 5% NaHCO₃ (140mL) and

5% NaCl (140mL). The organic layer was dried over Na₂SO₄ and concentrated to 7 volume of DCM solution. Methanol (10 vol, 140mL) was added dropwise over 2 hours, and the solution cooled to 15°C upon which the product precipitated. The product mixture was cooled to 5°C and mixed for 2 hours. Upon filtration of the solid and blow drying with nitrogen for 2 hours, 17.35 g of product was obtained (Yield 83%; purity >99.5 pa %). ¹H NMR (400 MHz, DMSO-d₆) δ ppm 11.57-11.59 (bs, 1H), 8.48-8.52 (m, 2H), 7.97 (d, *J* = 2.6 Hz, 1H), 7.73 (dd, *J* = 9.2, 2.3 Hz, 1H), 7.43-7.50 (m, 3H), 7.29-7.31 (m, 2H), 6.98-7.03 (m, 3H), 6.65 (dd, *J* = 8.9, 2.3 Hz, 1H), 6.35 (dd, *J* = 3.4, 1.8 Hz, 1H), 6.16 (d, *J* = 2.2 Hz, 1H), 4.41-4.44 (m, 1H), 3.71-3.75 (m, 2H), 2.98-3.51 (m, 11H), 2.74-2.76 (m, 3H), 2.02-2.26 (m, 6H), 1.88-1.92 (m, 2H), 1.47-1.70 (m, 5H), 1.24-1.40 (m, 4H), 1.08 (s, 5H), 0.89 (s, 6H).

[0164] The embodiments described above are intended merely to be exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [US61780621B \[0001\]](#)
- [US61947850B \[0001\]](#)
- [WO2011150016A \[0006\]](#)
- [WO2012071336A \[0006\] \[0102\] \[0103\]](#)
- [US20120108590A \[0006\]](#)
- [US20120277210A \[0006\]](#)
- [US20120129853A \[0006\]](#)
- [US20120157470A \[0006\] \[0102\] \[0103\]](#)
- [US20100305122A \[0102\] \[0103\]](#)
- [WO201115096A \[0102\] \[0103\]](#)
- [WO2000047212A \[0104\]](#)
- [WO2006059801A \[0104\]](#)
- [US3813443A \[0107\]](#)
- [GB585940A \[0108\]](#)
- [WO2009117626A \[0110\]](#)

Non-patent literature cited in the description


- **J. AM.**Chem. Soc., 1959, vol. 81, 743-747 [\[0104\]](#)
- Tetrahedron Letters, 2008, vol. 49, 122034-2037 [\[0104\]](#)
- Proceedings of the Chemical Society, 1907, vol. 22, 302- [\[0107\]](#)
- J. Am. Chem. Soc., 1950, vol. 72, 1215-1218 [\[0108\]](#)
- Organometallics, 2008, vol. 27, 215605-5611 [\[0110\]](#)

Patentkrav

1. Proces til tilvirkning af en sammensætning af formel A1:




hvor R^2 er valgt fra




som omfatter:

(a) at kombinere en sammensætning af formel (K):



hvor R er C_1 til C_{12} alkyl,

10 med et tert-butoxid salt, et aprotisk organisk opløsningsmiddel og vand til at tilvejebringe en sammensætning af formel (L):




og

- 2 -


(b") at kombinere sammensætningen af formel (L) med 1-methyl-3-(3-dimethylaminopropyl)carbodiimid hydrochlorid, 4-dimethylaminopyridin, et organisk opløsningsmiddel, og

enten en sammensætning af formel (N) til at tilvejebringe en sammensætning af

5 formel (A1), hvori R^2 er




eller en sammensætning af formel (P) til at tilvejebringe en sammensætning af formel (A1), hvori R^2 er




idet der derved tilvejebringes en sammensætning af formel (A1);

hvorved sammensætningen af formel (K) bliver tilvirket ved:

(d) at kombinere en sammensætning af formel (D):




med en sammensætning af formel (I):



en kilde til palladium, et tert-butoxid salt, og en fosphin ligand i et aprotisk organisk opløsningsmiddel til at tilvejebringe sammensætningen af formel (K).


20 2. Proces ifølge krav 1, hvorved R^2 er

- 3 -




og trin (b'') omfatter:

- (b'') at kombinere sammensætningen af formel (L) med 1-ethyl-3-(3-dimethylaminopropyl)carbodiimid hydrochlorid, 4-dimethylaminopyridin, et
- 5 organisk opløsningsmiddel og sammensætningen af formel (N):




for at tilvejebringe en sammensætning af formel (A1).

3. Proces ifølge krav 1, hvorved R^2 er




- 10 og processen yderligere omfatter:

(c'') at kombinere en sammensætning af formel (M):




- med en tertiær amin base, et organisk opløsningsmiddel og (tetrahydro-2H-pyran-4-yl)methanamin eller et salt deraf for at tilvejebringe sammensætningen
- 15 af formel (N).

4. Proces ifølge krav 1, hvorved R^2 er




og processen yderligere omfatter:

(c'') at kombinere en sammensætning af formel (M):



med en tertiær amin base, et organisk opløsningsmiddel og (1R,4R)-4-(aminomethyl)-1-methylcyclohexanol eller et salt deraf til at tilvejebringe sammensætningen af formel (P).

- 5 5. Proces ifølge et hvilket som helst af kravene 1 – 4, hvorved tert-butoxid saltet i trin (a) bliver valgt fra gruppen, der består af natrium tert-butoxid and kalium tert-butoxid.
6. Proces ifølge et hvilket som helst af kravene 3 – 5, hvorved den tertiære amin base i trin (c") er N,N-diisopropylethylamin.
- 10 7. Proces ifølge et hvilket som helst af kravene 1 til 6, hvorved sammensætningen fra formel (I) bliver kombineret med en base forud for kombinationen fra trin (d).
8. Proces ifølge et hvilket som helst af kravene 1 til 6 og krav 7, hvorved kilden til palladium i trin (d) er Pd₂dba₃ eller [(cinnamyl)PdCl]₂.
- 15 9. Proces ifølge et hvilket som helst af kravene 1 til 6, 7 og 8, hvorved fosphin liganden fra trin (d) er en sammensætning af formel (J):




10. Proces ifølge et hvilket som helst af kravene 1 til 6, 7, 8 og 9, hvorved kilden til palladium er Pd₂dba₃, der bliver benyttet en katalytisk mængde af Pd₂dba₃ i
- 20 forhold til mængden af sammensætning (I), og hvorved den katalytiske mængde af Pd₂dba₃ er fra omkring 0,5 molprocent til omkring 2 molprocent.
11. Proces ifølge krav 10, hvorved der bliver benyttet en katalytisk mængde af sammensætningen af formel (J) i forhold til mængden af sammensætningen (I), og hvorved den katalytiske mængde af sammensætningen af formel (J) er fra
- 25 omkring 1 molprocent til omkring 5 molprocent.

12. Proces ifølge et hvilket som helst af kravene 1 – 11, hvorved tert-butoxid saltet i trin (d) bliver valgt fra gruppen, der består af natrium tert-butoxid and kalium tert-butoxid.

13. Proces ifølge et hvilket som helst af kravene 1 – 12, hvorved processen yderligere omfatter:

(e) at kombinere en sammensætning af formel (B) med en sammensætning af formel (C):




og et tert-butoxid salt i et organisk opløsningsmiddel til at tilvejebringe sammensætningen af formel (D).

14. Proces ifølge krav 13, hvorved tert-butoxid saltet i trin (e) bliver valgt fra gruppen, der består af natrium tert-butoxid and kalium tert-butoxid.

15. Proces ifølge krav 13 eller 14, hvorved processen yderligere omfatter:

(f) at kombinere en sammensætning af formel (A):



med R^1MgX i et aprotisk organisk opløsningsmiddel; hvorved R^1 er C_1 til C_6 alkyl og X er Cl, Br eller I;


(g) at kombinere en C_1 til C_{12} alkyl chloroformat eller en di-(C_1 til C_{12} alkyl)dicarbonat med produktet fra trin (f) for at tilvejebringe sammensætningen af formel (C).

16. Proces ifølge krav 15, hvorved R^1 er isopropyl i trin (f).


17. Proces ifølge krav 15 eller 16, hvorved R er tert-butyl i trin (f), og di-(C_1 til C_{12} alkyl)dicarbonaten er di-tert-butyl dicarbonat.

18. Proces ifølge et hvilket som helst af kravene 1 - 17, hvorved processen yderligere omfatter:


(h) at kombinere en sammensætning af formel (E):




med dimethylformamid og POCl_3 for at tilvejebringe en sammensætning af formel (F):



- 5 (i) at kombinere sammensætningen af formel (F) med en kilde til palladium og 4-chlorophenylboronsyre i et organisk opløsningsmiddel for at tilvejebringe en sammensætning af formel (G):



- (j) at kombinere sammensætningen af formel (G) med BOC-piperazin og natrium triacetoxyborohydrid i et organisk opløsningsmiddel for at tilvejebringe en sammensætning af formel (H):



og

- (k) at kombinere sammensætningen af formel (H) med saltsyre for at tilvejebringe sammensætningen af formel (I).

- 15 19. Proces ifølge krav 18, hvorved kilden til palladium i trin (i) er $\text{Pd}(\text{OAc})_2$.
20. Proces ifølge krav 18 eller 19, hvorved trin (i) omfatter at kombinere tetrabutylammonium bromid med sammensætningen af formel (F), kilden til palladium og 4-chlorophenylboronsyre i det organiske opløsningsmiddel.
- 20 21. Proces ifølge et hvilket som helst af kravene 18 - 20, hvorved trin (j) yderligere omfatter at fremstille sammensætningen af formel (H) som et krystallinsk, fast stof.

22. Proces ifølge et hvilket som helst af kravene 18 - 21, hvorved trin (k) yderligere omfatter at fremstille sammensætningen af formel (I) som et krystallinsk, fast stof.
23. Proces ifølge et hvilket som helst af kravene 1 - 22, hvorved R er valgt fra
- 5 gruppen, der består af methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, isobutyl og neo-butyl.
24. Proces ifølge et hvilket som helst af kravene 23, hvorved R er tert-butyl.