

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization

International Bureau

(43) International Publication Date

11 October 2018 (11.10.2018)



W!P O | PCT



(10) International Publication Number

**WO 2018/184816 A1**

(51) International Patent Classification:

CUD 3/386 (2006.01) CI2N 9/26 (2006.01)

CI2N 9/22 (2006.01) CI2N 9/42 (2006.01)

CI2N 9/24 (2006.01)

(21) International Application Number:

PCT/EP20 18/056730

(22) International Filing Date:

16 March 2018 (16.03.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17165343.9 06 April 2017 (06.04.2017) EP

(71) Applicant: NOVOZYMES A/S [DK/DK]; Krogshoejvej  
36, 2880 Bagsvaerd (DK).

(72) Inventor: GORI, Klaus; Krogshoejvej 36, 2880 Bagsvaerd  
(DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: CLEANING COMPOSITIONS AND USES THEREOF

(57) Abstract: The present invention relates to compositions such as cleaning compositions comprising a mix of enzymes. The invention further relates, use of compositions comprising such enzymes in cleaning processes.



WO 2018/184816 A1

**CLEANING COMPOSITIONS AND USES THEREOF****Reference to a Sequence Listing**

5 This application contains a Sequence Listing in computer readable form, which is incorporated herein by reference.

**Background of the Invention**

10 The present invention relates to compositions such as cleaning compositions comprising a mix of enzymes. The invention further relates, use of compositions comprising such enzymes in cleaning processes and/or for deep cleaning of biofilm soiling, methods for removal or reduction of biofilm related soiling.

**Description of the Related Art**

15 Enzymes have been used in detergents for decades. Usually a cocktail of various enzymes is added to detergent compositions. The enzyme cocktail often comprises various enzymes, wherein each enzyme targets its specific substrate e.g. amylases are active towards starch stains, proteases on protein stains and so forth. Textiles surface and hard surfaces, such as dishes or the inner space of a laundry machine enduring a number of wash cycles, become soiled with many different types of soiling which may compose of proteins, grease, starch etc.  
20 One type of soiling may be organic matter, such as biofilm, EPS, etc. Organic matter composes different molecules such as polysaccharides, extracellular DNA (eDNA), and proteins. Some organic matter composes an extracellular polymeric matrix, which may be sticky or glueing, which when present on textile, attracts soils and may cause redeposition or backstaining of soil resulting in a greying of the textile. Additionally, organic matters such as biofilms often cause  
25 malodor issue as various malodor molecules can be adhered by the polysaccharides, extracellular DNA (eDNA), and proteins in the complex extracellular matrix and be slowly released out to cause consumer noticeable malodor issue. There is still a need for cleaning compositions, which effectively prevent, reduce or remove components of organic stains e.g. biofilm, an effect described in the present application as "deep cleaning". The present invention  
30 provides new compositions fulfilling such need.

**Summary of the Invention**

35 A first aspect of the present invention, relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. Another aspect of the invention relates to a cleaning composition comprising at least 0.001 ppm DNase and at least 0.001 ppm carbohydrase and a cleaning component, wherein the cleaning component is selected from

- a. 0.1 to 15 wt%, e.g. from about 1% to about 40% of at least one a surfactant;
- b. 0.5 to 20 wt% e.g. from about 5% to about 50% of at least one builder; and
- c. 0.01 to 10 wt% e.g. from about 1% to about 20% of at least one bleach component

The invention further relates to the use of a composition for deep cleaning of an item, wherein  
5 the item is a textile or a surface. The invention further relates to the use of a cleaning composition  
comprising a DNase, at least one carbohydrase and a cleaning component, wherein the  
carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. The invention further  
relates to a method of formulating a cleaning composition comprising adding a DNase, a  
10 carbohydrase and at least one cleaning component. The invention further relates to a kit intended  
for deep cleaning, wherein the kit comprises a solution of an enzyme mixture comprising a DNase  
and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a  
xylanase. The invention further relates to a method of deep cleaning an item, comprising the  
steps of: a) contacting the item with a cleaning composition comprising a DNase, at least one  
15 carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase,  
a mannanase or a xylanase; and b) and optionally rinsing the item, wherein the item is preferably  
a textile. The invention further relates to a method of deep cleaning of an item, comprising the  
steps of: a) contacting the item with a solution comprising an enzyme mixture comprising a DNase  
and a carbohydrase and optionally a protease; and a cleaning component, wherein the cleaning  
20 component is selected from 0.1 to 15 wt%, e.g. from about 1% to about 40% of at least one a  
surfactant; 0.5 to 20 wt% e.g. from about 5% to about 50% of at least one builder; and 0.01 to 10  
wt% e.g. from about 1% to about 20% of at least one bleach component; and b) and optionally  
rinsing the item, wherein the item is preferably a textile.

### Detailed Description of the Invention

25 Various enzymes are applied in cleaning processes each targeting specific types of soiling such  
as protein, starch and grease soiling. Enzymes are now standard ingredients in detergents for  
laundry and dish wash. The effectiveness of these commercial enzymes provides detergents  
which removes much of the soiling. However, organic matters such as EPS (extracellular  
polymeric substance) comprised in much biofilm constitute a challenging type of staining due to  
30 the complex nature of such organic matters. None of the commercially available cleaning  
compositions effectively remove or reduce EPS and/or biofilm related stains. Biofilm may be  
produced when a group of microorganisms' cells stick to each other or stick to a surface, such as  
a textile, dishware or hard surface or another kind of surface. These adherent cells are frequently  
embedded within a self-produced matrix of extracellular polymeric substance (EPS), which  
35 constitute 50% to 90% of the biofilm's total organic matter. EPS is mostly composed of  
polysaccharides (exopolysaccharides) and proteins, but include other macro-molecules such as  
eDNA, lipids and other organic substances. Organic matter like biofilm may be sticky or glueing,

which when present on textile, may give rise to redeposition or backstaining of soil resulting in a greying of the textile. Another drawback of organic matter e.g. biofilm is the malodor as various malodor related molecules are often associated with organic matter e.g. biofilm. Further, when dirty laundry items are washed together with less dirty laundry items the dirt present in the wash liquor tend to stick to organic matter e.g. biofilm or biofilm components as a result, hereof the laundry item is more "soiled" after wash than before wash. This is effect may also be termed re-deposition.

The compositions of the invention comprise a blend of DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and effectively reduce or remove organic components, such as protein and DNA from surfaces such as textiles and hard surfaces e.g. dishes.

The compositions of the invention comprise a blend of DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and effectively reduce or limit redeposition when applied in e.g. laundry process.

The compositions of the invention comprise a blend of DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and effectively reduce or limit malodor of e.g. textiles or hard surfaces such as dishes.

The compositions of the invention comprise a blend of DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and improve whiteness of textile.

A composition of the invention is preferably a cleaning composition, the composition of the invention comprises at least one DNase and at least one a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. Examples of useful DNases and carbohydrases are mentioned below in the sections "Polypeptides having DNase activity" and "Polypeptides having mannanase, cellulase, xylanase or amylase activity" respectively.

#### Polypeptides having DNase activity

The term "DNase" means a polypeptide with DNase (deoxyribonuclease) activity that catalyzes the hydrolytic cleavage of phosphodiester linkages in a DNA backbone, thus degrading DNA. Exodeoxyribonuclease cut or cleaves residues at the end of the DNA back bone where endo-deoxyribonucleases cleaves or cut within the DNA backbone. A DNase may cleave only double-stranded DNA or may cleave double stranded and single stranded DNA. The term "DNases" and the expression "a polypeptide with DNase activity" are used interchangeably throughout the application. For purposes of the present invention, DNase activity is determined according to the procedure described in the Assay 1.

Preferably the DNase is selected from any of the enzyme classes E.C.3.1, preferably E.C.3.1.21, e.g. such as E.C.3.1.21.X, where X = 1, 2, 3, 4, 5, 6, 7, 8 or 9, or e.g. Deoxyribonuclease I, Deoxyribonuclease IV, Type I site-specific deoxyribonuclease, Type II site-specific deoxyribonuclease, Type III site-specific deoxyribonuclease, CC-preferring endo-deoxyribonuclease, Deoxyribonuclease V, T(4) deoxyribonuclease II, T(4) deoxyribonuclease IV or E.C. 3.1.22.Y where Y = 1, 2, 4 or 5, e.g. Deoxyribonuclease II, *Aspergillus* deoxyribonuclease K(1), Crossover junction endo-deoxyribonuclease, Deoxyribonuclease X.

Preferably, the polypeptide having DNase activity is obtained from a microorganism and the DNase is a microbial enzyme. The DNase is preferably of fungal or bacterial origin.

The DNase may be obtainable from *Bacillus* e.g. *Bacillus*, such as a *Bacillus licheniformis*, *Bacillus subtilis*, *Bacillus sp-62451*, *Bacillus horikoshii*, *Bacillus sp-62451*, *Bacillus sp-16840*, *Bacillus sp-62668*, *Bacillus sp-13395*, *Bacillus horneckiae*, *Bacillus sp-1 1238*, *Bacillus cibi*, *Bacillus idriensis*, *Bacillus sp-62520*, *Bacillus sp-16840*, *Bacillus sp-62668*, *Bacillus algicola*, *Bacillus vietnamensis*, *Bacillus hwajinpoensis*, *Bacillus indicus*, *Bacillus marisflavi*, *Bacillus luciferensis*, *Bacillus sp. SA2-6*.

The DNase may also be obtained from any of the following *Pyrenochaetopsis sp.*, *Vibrissea flavovirens*, *Setosphaeria rostrate*, *Endophragmiella valdina*, *Corynespora cassicola*, *Paraphoma sp. XZ1965*, *Monilinia fructicola*, *Curvularia lunata*, *Penicillium reticulisporum*, *Penicillium quercetorum*, *Setophaeosphaeria sp.*, *Alternaria*, *Alternaria sp. XZ2545*, *Trichoderma reesei*, *Chaetomium thermophilum*, *Scytalidium thermophilum*, *Metapochonia suchlasporia*, *Daldinia fissa*, *Acremonium sp. XZ2007*, *Acremonium sp. XZ2414*, *Acremonium dichromosporum*, *Sarocladium sp. XZ2014*, *Metarhizium sp. HNA15-2*, *Isaria tenuipes*, *Scytalidium circinatum*, *Metarhizium lepidiotae*, *Thermobispora bispora*, *Sporormia fimetaria*, *Pycnidophora cf. dispersa*, *Environmental sample D*, *Environmental sample O*, *Clavicipitaceae sp-70249*, *Westerdykella sp. AS85-2*, *Humicolopsis cephalosporioides*, *Neosartorya massa*, *Roussoella intermedia*, *Pleosporales*, *Phaeosphaeria* or *Didymosphaeria futilis*.

The DNases to be used in a composition of the invention preferable belong to the NUC1 group of DNases. The NUC1 group of DNases comprises polypeptides which in addition to having DNase activity, may comprise one or more of the motifs [T/D/S][G/N]PQL (SEQ ID NO 69), [F/L/Y/I]A[N/R]D[L/I/P/V] (SEQ ID NO: 70), or C[D/N]T[A/R] (SEQ ID NO: 71). One embodiment of the invention relates to a composition comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and polypeptides having DNase activity, wherein the polypeptides comprises one or more of the motifs [T/D/S][G/N]PQL (SEQ ID NO 69), [F/L/Y/I]A[N/R]D[L/I/P/V] (SEQ ID NO: 70) or C[D/N]T[A/R] (SEQ ID NO: 71).

The DNases preferably comprises a NUC1\_A domain [D/Q][IA]DH (SEQ ID NO 72). In addition to comprising any of the domain motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/P/V] or C[D/N]T[A/R] the polypeptides having DNase activity, to be used in a composition of the

invention, may comprise the NUC1\_A domain and may share the common motif [D/Q][I/A]DH (SEQ ID NO 72). One embodiment the invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and polypeptides, which comprises one or more motifs selected from the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/P/V], C[D/N]T[A/R] and [D/Q][I/A]DH, wherein the polypeptides have DNase activity.

The DNases to be added to a composition of the invention preferably belong to the group of DNases comprised in the GYS-clade, which are group of DNases on the same branch of a phylogenetic tree having both structural and functional similarities. These NUC1 and/or NUC1\_A DNases comprise the conservative motifs [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74) and share similar structural and functional properties. The DNases of the GYS-clade are preferably obtained from *Bacillus* genus.

One embodiment of the invention relates to a composition comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, a polypeptide of the GYS clade having DNase activity, optionally wherein the polypeptide comprise one or both of the motifs [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73), ASXNRSKG (SEQ ID NO: 74) and wherein the polypeptide is selected from the group of polypeptides:

- a) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 1,
- b) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 2,
- c) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 3,
- d) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 4,
- e) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 5,
- f) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 6,

- g) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 7,
- 5 h) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 8,
- i) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 9,
- 10 j) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 10,
- k) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 11,
- 15 l) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 12,
- m) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 13,
- 20 n) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 14,
- 25 o) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 15,
- p) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 16,
- 30 q) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 17,
- r) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 18,
- 35

- s) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 19,
- 5 t) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 20,
- u) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 21,
- 10 v) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 22,
- w) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 23,
- 15 x) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 24, and
- y) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 25.
- 20

Polypeptides having DNase activity and which comprise the GYS-clade motifs have shown particularly good deep cleaning properties e.g. the DNases are particularly effective in removing or reducing DNA stains e.g. associated with biofilm or dead cell debris, from an item such as a textile or a hard surface. In addition, these DNases are particularly effective in removing or reducing malodor, from an item such as a textile or a hard surface. Further, the GYS-clade DNases are particularly effective in preventing redeposition when laundering an item such as textile.

25

In one embodiment the DNases to be added in a composition of the invention preferably belong to the group of DNases comprised in the NAWK-clade, which are NUC1 and NUC1\_A DNases, which may further comprise the conservative motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76).

30

One embodiment of the invention relates to a composition comprising a carbohydrase selected from a cellulase, an amylase, a mannanase or a xylanase, and a polypeptide of the NAWK-clade having DNase activity, optionally wherein the polypeptide comprise one or both of the motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76) and wherein the polypeptide is selected from the group of polypeptides:

35



- a) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 26,
- 5 b) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 27,
- c) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 28,
- 10 d) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 29,
- e) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 30,
- 15 f) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 31,
- g) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 32,
- 20 h) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 33,
- 25 i) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 34,
- j) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 35,
- 30 k) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 36,
- 35 l) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 37, and

m) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 38.

5 Polypeptides having DNase activity and which comprise the NAWK-clade motifs have shown particularly good deep cleaning properties e.g. the DNases are particularly effective in removing or reducing DNA stains e.g. associated with biofilm or dead cell debris, from an item such as a textile or a hard surface. In addition, these DNases are particularly effective in removing or reducing malodor, from an item such as a textile or a hard surface. Further, the NAWK-clade DNases are particularly effective in preventing redeposition when laundering an item such as  
10 textile.

The DNases to be added in a composition of the invention preferably belong to the group of DNases comprised in the KNAW-clade, which are NUC1 and NUC1\_A DNases which may further comprise the conservative motifs P[Q/E]L[W/Y] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO: 78).

15 One embodiment of the invention relates to a composition comprising a carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase, and a polypeptide of the KNAW clade having DNase activity, optionally wherein the polypeptide comprise one or both of the motifs P[Q/E]L[VWY] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO: 78), and wherein the polypeptide is selected from the group of polypeptides:

20 a) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 39,

b) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity  
25 to the polypeptide shown in SEQ ID NO: 40,

c) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 41,

d) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%,  
30 at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 42,

e) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 43

35 f) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 44,

- g) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 45,
- 5 h) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 46,
- i) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 47,
- 10 j) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 48,
- k) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 49,
- 15 l) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 50, and
- m) a polypeptide having at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 100% sequence identity to the polypeptide shown in SEQ ID NO: 51.
- 20

Polypeptides having DNase activity and which comprise the KNAW-clade motifs have shown particularly good deep cleaning properties e.g. the DNases are particularly effective in removing or reducing DNA stains e.g. associated with biofilm or dead cell debris, from an item such as a textile or a hard surface. In addition, these DNases are particularly effective in removing or reducing malodor, from an item such as a textile or a hard surface. Further, the KNAW-clade DNases are particularly effective in preventing redeposition when laundering an item such as textile.

25

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-62451 and having a sequence identity to the polypeptide shown in SEQ ID NO: 1 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 1.

30

35

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus horikoshii* and having a sequence identity to the polypeptide shown in SEQ ID NO: 2 of at least 60%, e.g., at least 65%,  
5 at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%,  
at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%,  
at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%,  
at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by  
up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in  
10 SEQ ID NO: 2.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-62520 and having a sequence identity to the polypeptide shown in SEQ ID NO: 3 of at least 60%, e.g., at least 65%,  
15 at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%,  
at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%,  
at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%,  
at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by  
up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in  
20 SEQ ID NO: 3.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-62520 and having a sequence identity to the polypeptide shown in SEQ ID NO: 4 of at least 60%, e.g., at least 65%,  
25 at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%,  
at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%,  
at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%,  
at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by  
up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in  
30 SEQ ID NO: 4.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus horikoshii* and having a sequence identity to the polypeptide shown in SEQ ID NO: 5 of at least 60%, e.g., at least 65%,  
35 at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%,  
at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%,  
at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%,

at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 5.

5 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus horikoshii* and having a sequence identity to the polypeptide shown in SEQ ID NO: 6 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect,  
10 the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 6.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-16840 and having a  
15 sequence identity to the polypeptide shown in SEQ ID NO: 7 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 7.

20 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-16840 and having a sequence identity to the polypeptide shown in SEQ ID NO: 8 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%,  
25 at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 8.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a  
30 polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-62668 and having a sequence identity to the polypeptide shown in SEQ ID NO: 9 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect,  
35 the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 9.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a

polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-13395 and having a sequence identity to the polypeptide shown in SEQ ID NO: 10 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 10.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus horneckiae* and having a sequence identity to the polypeptide shown in SEQ ID NO: 11 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 11.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-1 1238 and having a sequence identity to the polypeptide shown in SEQ ID NO: 12 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 12.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus cibi* and having a sequence identity to the polypeptide shown in SEQ ID NO: 13 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 13.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp-18318 and having a sequence identity to the polypeptide shown in SEQ ID NO: 14 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect,

the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 14.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus idriensis* and having a sequence identity to the polypeptide shown in SEQ ID NO: 15 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 15.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus algicola* having a sequence identity to the polypeptide shown in SEQ ID NO: 16 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 16.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from Environmental sample J and having a sequence identity to the polypeptide shown in SEQ ID NO: 17 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 17.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus vietnamensis* and having a sequence identity to the polypeptide shown in SEQ ID NO: 18 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 18.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus hwajinpoensis* and having a

sequence identity to the polypeptide shown in SEQ ID NO: 19 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 19.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Paenibacillus mucilaginosus* and having a sequence identity to the polypeptide shown in SEQ ID NO: 20 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 20.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus indicus* and having a sequence identity to the polypeptide shown in SEQ ID NO: 21 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 21.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus marisflavi* and having a sequence identity to the polypeptide shown in SEQ ID NO: 22 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 22.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus luciferensis* and having a sequence identity to the polypeptide shown in SEQ ID NO: 23 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 23.



In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus marisflavi* and having a sequence identity to the polypeptide shown in SEQ ID NO: 24 of at least 60%, e.g., at least 65%,  
5 at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 24.

In some embodiments, the present invention relates to compositions comprising a  
10 carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus* sp. SA2-6 and having a sequence identity to the polypeptide shown in SEQ ID NO: 25 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect,  
15 the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 25.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Pyrenochaetopsis* sp. and having a sequence identity to the  
20 polypeptide shown in SEQ ID NO: 26 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 26.

In some embodiments, the present invention relates to compositions comprising a  
25 carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Vibrissia flavovirens* and having a sequence identity to the polypeptide shown in SEQ ID NO: 27 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least  
30 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 27.

In some embodiments, the present invention relates to compositions comprising a  
35 carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Setosphaeria rostrate* and having a sequence identity to the polypeptide shown in SEQ ID NO: 28 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least

98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 28.

5 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Endophragmiella valdina* and having a sequence identity to the polypeptide shown in SEQ ID NO: 29 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ  
10 by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 29.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Corynespora cassicola* and having a sequence identity to the  
15 polypeptide shown in SEQ ID NO: 30 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 30.

20 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Paraphoma* sp. XZ1965 and having a sequence identity to the polypeptide shown in SEQ ID NO: 31 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least  
25 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 31.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a  
30 polypeptide obtainable from *Monilinia fructicola* and having a sequence identity to the polypeptide shown in SEQ ID NO: 32 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID  
35 NO: 32.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a

polypeptide obtainable from *Curvularia lunata* and having a sequence identity to the polypeptide shown in SEQ ID NO: 33 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 33.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Penicillium reticulisporum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 34 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 34.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Penicillium quercetorum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 35 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 35.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Setophaeosphaeria* sp. and having a sequence identity to the polypeptide shown in SEQ ID NO: 36 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 36.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Alternaria* sp. XZ2545 and having a sequence identity to the polypeptide shown in SEQ ID NO: 37 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ

by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 37.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Alternaria* and having a sequence identity to the polypeptide shown in SEQ ID NO: 38 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 38.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Trichoderma reesei* and having a sequence identity to the polypeptide shown in SEQ ID NO: 39 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 39.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Chaetomium thermophilum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 40 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 40.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Scytalidium thermophilum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 41 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 41.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Metapochonia suchlasporia* and having a sequence identity to the polypeptide shown in SEQ ID NO: 42 of at least 60%, e.g., at least 65%, at least 70%, at least

75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 42.

5 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Daldinia fissa* and having a sequence identity to the polypeptide shown in SEQ ID NO: 43 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 10 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 43.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Acremonium* sp. XZ2007 and having a sequence identity to the polypeptide shown in SEQ ID NO: 44 of at least 60%, e.g., at least 65%, at least 70%, at least 15 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 44.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Acremonium dichromosporum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 45 of at least 60%, e.g., at least 65%, at least 70%, at least 25 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 45.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Sarocladium* sp. XZ2014 and having a sequence identity to the polypeptide shown in SEQ ID NO: 46 of at least 60%, e.g., at least 65%, at least 70%, at least 30 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 46.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Metarhizium* sp. HNA15-2 and having a sequence identity to the polypeptide shown in SEQ ID NO: 47 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 47.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Acremonium* sp. XZ2414 and having a sequence identity to the polypeptide shown in SEQ ID NO: 48 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 48.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Isaria tenuipes* and having a sequence identity to the polypeptide shown in SEQ ID NO: 49 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 49.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Scytalidium circinatum* and having a sequence identity to the polypeptide shown in SEQ ID NO: 50 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 50.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Metarhizium lepidiotae* and having a sequence identity to the polypeptide shown in SEQ ID NO: 51 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least

98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 51.

5 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Thermobispora bispora* and having a sequence identity to the polypeptide shown in SEQ ID NO: 52 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ  
10 by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 52.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Sporormia fimetaria* and having a sequence identity to the  
15 polypeptide shown in SEQ ID NO: 53 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 53.

20 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Pycnidophora* cf. *dispera* and having a sequence identity to the polypeptide shown in SEQ ID NO: 54 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least  
25 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 54.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a  
30 polypeptide obtainable from Environmental sample D and having a sequence identity to the polypeptide shown in SEQ ID NO: 55 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown  
35 in SEQ ID NO: 55.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a

polypeptide obtainable from Enviromental sample O and having a sequence identity to the polypeptide shown in SEQ ID NO: 56 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 56.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Clavicipitaceae* sp-70249 and having a sequence identity to the polypeptide shown in SEQ ID NO: 57 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 57.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Westerdykella* sp. AS85-2 and having a sequence identity to the polypeptide shown in SEQ ID NO: 58 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 58.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Humicolopsis cephalosporioides* and having a sequence identity to the polypeptide shown in SEQ ID NO: 59 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 59.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Neosartorya massa* and having a sequence identity to the polypeptide shown in SEQ ID NO: 60 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ



by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 60.

5 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Rousoella intermedia* and having a sequence identity to the polypeptide shown in SEQ ID NO: 61 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown  
10 in SEQ ID NO: 61.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Pleosporales* and having a sequence identity to the polypeptide shown in SEQ ID NO: 62 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least  
15 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 62.

In some embodiments, the present invention relates to compositions comprising a  
20 carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Phaeosphaeria* and having a sequence identity to the polypeptide shown in SEQ ID NO: 63 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10  
25 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 63.

In some embodiments, the present invention relates to compositions comprising a  
30 carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Didymosphaeria futilis* and having a sequence identity to the polypeptide shown in SEQ ID NO: 64 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown  
in SEQ ID NO: 64.

35 In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus licheniformis* having a

sequence identity to the polypeptide shown in SEQ ID NO: 65 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 65.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Bacillus* e.g. obtainable from *Bacillus subtilis* having a sequence identity to the polypeptide shown in SEQ ID NO: 66 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 66.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Aspergillus* e.g. obtainable from *Aspergillus oryzae* having a sequence identity to the polypeptide shown in SEQ ID NO: 67 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 67.

In some embodiments, the present invention relates to compositions comprising a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and a polypeptide obtainable from *Trichoderma* e.g. obtainable from *Trichoderma harzianum* having a sequence identity to the polypeptide shown in SEQ ID NO: 68 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 68.

The DNases above may be combined with any of the carbohydrases below to form a blend to be added to a composition according to the invention.

#### Polypeptides having Carbohydrase activity (Carbohydrases)

Carbohydrase is a protein/enzyme that catalyse carbohydrates to break down carbohydrates to e.g. simple sugar such as monosaccharides. Thus, carbohydrases are any of a group of enzymes that promote hydrolysis of a carbohydrate. Starch hydrolyzing carbohydrases (e.g. amylases)

work on e.g. amylose and amylopectin and non-starch carbohydrases includes enzymes which hydrolyze polymers made up of carbon sugars e.g. cellulases which will ultimately produce glucose when complete hydrolysis is achieved. Another example is lactase which hydrolyses lactose to glucose and galactose. Examples of carbohydrases includes amylases, cellulases and mannanases. The carbohydrases to be incorporated in a composition according to the invention is preferably selected from xylanases, cellulases, mannanases and amylases.

#### Polypeptides having mannanase activity

The term "mannanase" is defined here as an enzyme that hydrolyses compounds known as mannanes. The term "mannanase activity" is as an enzyme catalyzed hydrolysis of mannan, for purposes of the present invention, mannanase activity is determined according to the procedure described in the Assay II. Suitable mannanases include those of bacterial or fungal origin. Chemically or genetically modified mannanases are included. The mannanase may be an alkaline mannanase of Family 5 or 26. It may be a wild-type from *Bacillus* or *Humicola*, particularly *B. agaradhaerens*, *B. licheniformis*, *B. halodurans*, *B. clausii*, or *H. insolens*. Suitable mannanases are described in WO 1999/064619. Commercially available mannanases are Mannaway (Novozymes A/S), and EFFECTENZ™ M1000 from Dupont. In one aspect, the present invention relates to a cleaning composition comprising at least one enzyme having mannanase activity, which may be obtained from a bacterial strain of the genus *Bacillus*. Preferably, a polypeptide selected from the group of polypeptides comprising the amino acid sequence shown in SEQ ID NO: 82. In one aspect the present invention relates to a cleaning composition comprising at least one enzyme classified in the EC 3.2.1.78 and which has mannanase activity.

Useful mannanases include polypeptides that are substantially homologous to the polypeptides shown in SEQ ID NO: 82 and species homologs (paralogs or orthologs) thereof.

The term "substantially homologous" is used herein to denote polypeptides having at least 60%, at least 65%, preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90% more preferably at least 95%, more preferably at least 97%, even more preferably at least 98% sequence identity to the sequence.

In some embodiments, the present invention relates compositions comprising a DNase and a polypeptide having a sequence identity to the polypeptide shown in SEQ ID NO: 82 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have mannanase activity. In one aspect, the polypeptides differ by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the mature polypeptide shown in SEQ ID NO: 82.

The mannanase suitable for a composition of the invention may, in addition to the enzyme core comprising a catalytically domain, also comprise a cellulose binding domain (CBD), the cellulose binding domain and enzyme core (the catalytically active domain) of the enzyme being operably linked.

5 In one aspect, the present invention relates to a composition comprising a mannanase and a DNase, wherein the a mannanase is: i) polypeptide comprising an amino acid sequence as shown in SEQ ID NO:82; or ii) or a polypeptide having a sequence identity of at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% to the amino acid sequence shown in SEQ ID NO 82.

10 In one aspect, the present invention relates to a composition comprising a mannanase and a DNase, wherein the mannanase is selected from a polypeptide having at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

15

#### Polypeptides having Cellulase activity

The term "cellulase" is defined in the present context as an enzyme that hydrolyses cellulose. In a preferred embodiment of the invention, the cellulase is an endoglucanase. The term "cellulase activity" is defined herein as an enzyme catalyzed hydrolysis of 1,4-beta-D-glucosidic linkages in beta-1,4-glucan (cellulose). For purposes of the present invention, cellulase activity is determined using AZCL- HE-cellulose (from Megazyme) as the reaction substrate, as shown in Assay IV. Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

25 Especially suitable cellulases are the alkaline or neutral cellulases having color care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/1 1262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and WO99/001544.

30 Other cellulases are endo-beta-1,4-glucanase enzyme having a sequence of at least 97% identity to the amino acid sequence of position 1 to position 773 of SEQ ID NO:2 of WO 2002/099091 or a family 44 xyloglucanase, which a xyloglucanase enzyme having a sequence of at least 60% identity to positions 40-559 of SEQ ID NO: 2 of WO 2001/062903.

35 Commercially available cellulases include Celluzyme™, and Carezyme™ (Novozymes A/S) Carezyme Premium™ (Novozymes A/S), Celluclean™ (Novozymes A/S), Celluclean

Classic™ (Novozymes A/S), Cellusoft™ (Novozymes A/S), Whitezyme™ (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation), Revitalenz™ 1000, Revitalenz™ 2000, Revitalenz™ 3000 (Dupont).

5 In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprise the amino acid sequence of SEQ ID NO: 83. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more  
10 preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 83.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprise the amino acid sequence of SEQ ID NO: 84. In one aspect of the present invention, the cleaning composition comprise a DNase and a  
15 polypeptide having cellulase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 84.

20 In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprise the amino acid sequence of SEQ ID NO: 85. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%,  
25 more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 85.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having cellulase activity, which comprise the amino acid sequence of SEQ ID NO: 86. In one aspect of the present invention, the cleaning composition comprise a DNase and a  
30 polypeptide having cellulase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even  
35 most preferably at least 99% or 100% sequence identity to SEQ ID NO: 86.

In the present invention, the cleaning composition can also comprise a cellulase, which is a xyloglucanase. The term "xyloglucanase activity" is defined herein as an enzyme catalyzed

hydrolysis of xyloglucan, which is shown in Assay III. Xyloglucanase can comprise parent xyloglucanase and the variants thereof.

In one embodiment of the present invention, the xyloglucanase is a polypeptide comprising an amino acid sequence of SEQ ID NO 87. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having xyloglucanase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 87.

#### Polypeptides having Amylase activity

An amylase is an enzyme that hydrolyses starch into sugars, for purposes of the present invention, amylase activity is determined according to the procedure described in the Assay V. Suitable amylases include alpha-amylases and/or a glucoamylases and may be of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described in more detail in GB 1,296,839.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprise the amino acid sequence of SEQ ID NO: 88. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 88.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprise the amino acid sequence of SEQ ID NO: 89. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprises an amino acid sequence having at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, even most preferably at least 99% or 100% sequence identity to SEQ ID NO: 89.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprise the amino acid sequence of SEQ ID NO: 90. In one aspect of the present invention, the cleaning composition comprise a DNase and a

polypeptide having amylase activity, which comprises an amino acid sequence at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99%, identity SEQ ID NO: 90.

In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprise the amino acid sequence of SEQ ID NO: 91. In one aspect of the present invention, the cleaning composition comprise a DNase and a polypeptide having amylase activity, which comprises an amino acid sequence at least 60%, at least 65%, at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, more preferably at least 90%, more preferably at least 95%, more preferably at least 96%, even more preferably at least 97%, most preferably at least 98%, or even most preferably at least 99%, identity SEQ ID NO: 91.

Additional amylases include amylases comprising the polypeptide shown in SEQ ID NO: 2 in WO 95/10603 or variants having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred variants are described in WO 94/02597, WO 94/18314, WO 97/43424 and SEQ ID NO: 4 of WO 99/019467, such as variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305, 391, 408, and 444. Different suitable amylases include amylases having SEQ ID NO: 6 in WO 02/010355 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193. Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO 2006/066594 or variants having 90% sequence identity thereof. Preferred variants of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one or more of the following positions: 48, 49, 107, 156, 181, 190, 197, 201, 209 and 264. Most preferred variants of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;

H156Y+A181T+N190F+A209V+Q264S; or

G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S.

Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO 99/019467 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269. Particularly

preferred amylases are those having deletion in positions R181 and G182, or positions H183 and G184. Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 2 or SEQ ID NO: 7 of WO 96/023873 or variants thereof having 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred variants of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476, using SEQ ID 2 of WO 96/023873 for numbering. More preferred variants are those having a deletion in two positions selected from 181, 182, 183 and 184, such as 181 and 182, 182 and 183, or positions 183 and 184. Most preferred amylase variants of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476. Other amylases which can be used are amylases having SEQ ID NO: 2 of WO 08/153815, SEQ ID NO: 10 in WO 01/66712 or variants thereof having 90% sequence identity to SEQ ID NO: 2 of WO 08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO 01/66712. Preferred variants of SEQ ID NO: 10 in WO 01/66712 are those having a substitution, a deletion or an insertion in one or more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264. Further suitable amylases are amylases having SEQ ID NO: 2 of WO 09/061380 or variants having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred variants of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or an insertion in one or more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred variants of SEQ ID NO: 2 are those having the substitution in one or more of the following positions: Q87E,R, Q98R, S125A, N128C, T131I, T165I, K178L, T182G, M201L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase variants of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;

N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;

S125A+N128C+K178L+T182G+Y305R+G475K; or

S125A+N128C+T131I+T165I+K178L+T182G+Y305R+G475K wherein the variants are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

Further suitable amylases are amylases having SEQ ID NO: 1 of WO 2013/184577 or variants having 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one or more of the following positions: K176, R178, G179, T180, G181, E187, N192, M199, I203, S241, R458, T459, D460, G476 and G477. More preferred variants of SEQ ID NO: 1 are those having the substitution in one or more



of the following positions: K176L, E187P, N192FYH, M199L, I203YF, S241QADN, R458N, T459S, D460T, G476K and G477K and/or deletion in position R178 and/or S179 or of T180 and/or G181. Most preferred amylase variants of SEQ ID NO: 1 are those having the substitutions:

5 E187P+I203Y+G476K

E187P+I203Y+R458N+T459S+D460T+G476K,

wherein the variants optionally further comprise a substitution at position 241 and/or a deletion at position 178 and/or position 179.

Further suitable amylases are amylases having SEQ ID NO: 1 of WO1 0104675 or variants having 10 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one or more of the following positions: N21, D97, V128 K177, R179, S180, 1181, G182, M200, L204, E242, G477 and G478. More preferred variants of SEQ ID NO: 1 are those having the substitution in one or more of the following 15 positions: N21 D, D97N, V128I K177L, M200L, L204YF, E242QA, G477K and G478K and/or deletion in position R179 and/or S180 or of 1181 and/or G182. Most preferred amylase variants of SEQ ID NO: 1 are those having the substitutions:

N21 D+D97N+V128I

wherein the variants optionally further comprise a substitution at position 200 and/or a deletion at position 180 and/or position 181. Other suitable amylases are the alpha-amylase comprising 20 the polypeptide sequence shown in SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase variants are those having a substitution, a deletion or an insertion in one or more of the following positions of SEQ ID NO: 12 in WO01/66712: R28, R118, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, 25 K446, Q449, R458, N471, N484. Preferred amylases include variants having a deletion of D183 and G184 and having the substitutions R118K, N195F, R320K and R458K, and a variant additionally having substitutions in one or more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions. Other examples are amylase variants such 30 as those described in WO201 1/098531, WO20 13/00 1078 and WO201 3/001 087.

Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™, Stainzyme™, Stainzyme Plus™, Natalase™, Liquozyme X and BAN™ (from Novozymes A/S), and Rapidase™, Purastar™/Effectenz™, Powerase, Preferenz S1000, Preferenz S100 and Preferenz S110 (from Genencor International Inc./DuPont).

35

**A composition comprising:**

The invention relates to compositions, preferably cleaning compositions comprising a DNase and a carbohydrase in combination with one or more additional cleaning composition components.

5 One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. The carbohydrase may be any of the cellulases, amylases, mannanases or xylanases mentioned under the heading "Polypeptides having cellulase, amylase, mannanase or xylanase activity" respectively.

10 As shown in the examples of the present invention carbohydrases such as cellulases act synergistically with the DNase in reduction, and removal of biofilm or components hereof. Biofilm is a complex structure comprising, the target substrate e.g. the DNA may be embedded in the biofilm structure and It's believed that when the DNases and carbohydrases are acting together, the DNA components are more effectively dispersed or removed. It is thus advantageous to formulate DNases with carbohydrases such as cellulases, amylases, mannanases and xylanases  
15 in cleaning compositions e.g. for deep cleaning. One aspect of the invention relates to a method of formulating a cleaning composition comprising adding a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. The invention further relates to a kit intended for deep cleaning, wherein the kit comprises a solution of an enzyme mixture comprising a DNase and a carbohydrase, wherein  
20 the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase.

In one aspect of the invention the carbohydrase is a cellulase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the  
25 carbohydrase is a cellulase, preferably selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO:  
30 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86.

35 In one aspect of the invention the carbohydrase is an amylase.

In one aspect of the invention the carbohydrase is an amylase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component,

wherein the carbohydrase is an amylase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is an amylase, preferably selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91.

In one aspect of the invention the carbohydrase is a mannanase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is a mannanase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is a mannanase, preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

In one aspect of the invention the carbohydrase is a xylanase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is a xylanase. In one aspect, the invention relates to a cleaning composition comprising a DNase, a carbohydrase and a cleaning component, wherein the carbohydrase is a xylanase, preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87.

The DNases to be formulated together with the carbohydrases or to be used together with the carbohydrases should be compatible with cleaning components. DNases are at present not standard ingredients in cleaning compositions. However, the applicant has identified DNases suitable for use in cleaning compositions e.g. in WO2017/060475, WO2014/08701 1, WO2015/155350 and WO2015/155351. These applications also mentioned that DNases may be formulated with other enzymes e.g. carbohydrases. However, none of these applications indicate that the DNases may have synergy with e.g. cellulases. Enzymes, such as DNases should not only be compatible with the cleaning components, the DNases should also be compatible with other enzymes which may be present in a typical cleaning composition. Surprisingly, it was found that carbohydrases such as cellulases and DNases not only are compatible but even act synergistically in respect of biofilm reduction and removal e.g. in deep cleaning.

Particularly useful DNases may be those of microbial origin. One embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase and at least one cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase is microbial, preferably obtained from bacteria or fungi. In one  
5 embodiment, the cleaning composition comprise a DNase from bacteria. One embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase and at least one cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase is obtained from *Bacillus*, preferably *Bacillus cibi*, *Bacillus horikoshii*, *Bacillus licheniformis*, *Bacillus subtilis*, *Bacillus horneckiae*, *Bacillus idriensis*, *Bacillus algicola*, *Bacillus vietnamensis*, *Bacillus hwajinpoensis*, *Bacillus indicus*, *Bacillus marisflavi* or  
10 *Bacillus luciferensis*.

As mentioned above the DNases to be used in a composition of the invention preferable belong to the NUC1 group of DNases. The NUC1 group of DNases may comprise one or more of the motifs [T/D/S][G/N]PQL (SEQ ID NO 69), [F/L/Y/I]A[N/R]D[L/I/P/V] (SEQ ID NO: 70), or  
15 C[D/N]T[A/R] (SEQ ID NO: 71). One embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and at least one cleaning component, wherein the DNase comprises one or more of the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/P/V] or C[D/N]T[A/R]. The DNases preferably additionally comprises a NUC1\_A domain [D/Q][IA]DH (SEQ ID NO 72).

20 One embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and at least one cleaning component, wherein the DNase comprises one or more motifs selected from the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/P/V], C[D/N]T[A/R] and [D/Q][IA]DH.

25 One preferred embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and at least one cleaning component, wherein the DNase comprises two or more motifs selected from the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/PA], C[D/N]T[A/R] and [D/Q][IA]DH.

30 One preferred embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and at least one cleaning component, wherein the DNase comprises three or more motifs selected from the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/PA], C[D/N]T[A/R] and [D/Q][IA]DH.

35 One preferred embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and at least one cleaning component, wherein the DNase comprises four or more motifs selected from the motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/PA], C[D/N]T[A/R] and [D/Q][IA]DH.

One preferred embodiment of the invention relates to a cleaning composition comprising a DNase, a carbohydrase, selected from cellulases, amylases, mannanases and xylanases, and

at least one cleaning component, wherein the DNase comprises all five motifs [T/D/S][G/N]PQL, [F/L/Y/I]A[N/R]D[L/I/PA], C[D/N]T[A/R] and [D/Q][I/A]DH.

The DNases to be added to a composition of the invention preferably belong to the group of DNases comprised in the GYS-clade, which are NUC1 and NUC1\_A DNases further comprising the conservative motifs [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74) and which share similar structural and functional properties. The DNases of the GYS-clade are preferably obtained from *Bacillus* genus.

One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74).

In a particularly preferred embodiment the *Bacillus* DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74). In another particularly preferred embodiment the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74) and is obtained from *Bacillus cibi*. In yet another preferred embodiment the DNase comprises the amino acid sequence shown in SEQ ID NO 13 or DNases closely related hereto.

One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

Other preferred DNases include those comprising the amino acid sequence shown in SEQ ID NO 65 and 66.

One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

The DNase may also preferably be fungal. Particularly preferred are DNases obtained from *Aspergillus* in particular, *Aspergillus oryzae*.

One embodiment of the invention relates to a cleaning composition comprising a DNase,

at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

5 Other particularly preferred are DNases obtained from *Trichoderma* in particular, *Trichoderma harzianum*.

One embodiment of the invention relates to a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase and wherein the DNase has at least 60%, at least 65%,  
10 at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment relates to a cleaning composition comprising a *Bacillus* DNase, a cellulase and at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least  
15 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least  
20 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86.

One embodiment relates to a cleaning composition comprising a *Bacillus* DNase, an  
25 amylase and at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least  
30 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least  
35 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91.

One embodiment relates to a cleaning composition comprising a *Bacillus* DNase, a mannanase and at least one cleaning component, wherein the mannanase is selected from a

polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

5 One embodiment relates to a cleaning composition comprising a *Bacillus* DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87.

10 One embodiment relates to a cleaning composition comprising a DNase, a cellulase and at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

25 One embodiment relates to a cleaning composition comprising a DNase, an amylase and at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

35 One embodiment relates to a cleaning composition comprising a DNase, a mannanase

and at least one cleaning component, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment relates to a cleaning composition comprising a DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment relates to a cleaning composition comprising a DNase, a cellulase and at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment relates to a cleaning composition comprising a DNase, an amylase and at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at



least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

5 One embodiment relates to a cleaning composition comprising a DNase, a mannanase and at least one cleaning component, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at  
10 least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment relates to a cleaning composition comprising a DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least  
15 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment relates to a cleaning composition comprising a DNase, a cellulase and  
20 at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide  
25 shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein  
30 the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment relates to a cleaning composition comprising a DNase, an amylase and  
35 at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least

85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment relates to a cleaning composition comprising a DNase, a mannanase and at least one cleaning component, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment relates to a cleaning composition comprising a DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment relates to a cleaning composition comprising a DNase, a cellulase and at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment relates to a cleaning composition comprising a DNase, an amylase and at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88,  
5 a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at  
10 least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

15 One embodiment relates to a cleaning composition comprising a DNase, a mannanase and at least one cleaning component, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at  
20 least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment relates to a cleaning composition comprising a DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least  
25 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment relates to a cleaning composition comprising a DNase, a cellulase and  
30 at least one cleaning component, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least  
35 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at

least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment relates to a cleaning composition comprising a DNase, an amylase and at least one cleaning component, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment relates to a cleaning composition comprising a DNase, a mannanase and at least one cleaning component, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment relates to a cleaning composition comprising a DNase, a xylanase and at least one cleaning component, wherein the xylanase is selected from preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment of the invention relates to a composition comprising

- a) at least 0.001 ppm e.g. 0.1 ppm or 1 ppm of at least one polypeptide having DNase activity, wherein the DNase is selected for the group consisting of:
  - i) a DNase comprising one or more of the motif(s) [T/D/S][G/N]PQL (SEQ ID NO

- 69), [F/L/Y/I]A[N/R]D[L/I/PA] (SEQ ID NO: 70), C[D/N]T[A/R] (SEQ ID NO: 71);
- ii) a DNase comprising the motif [D/Q][IA]DH (SEQ ID NO 72);
- iii) a DNase comprising one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74);
- 5 iv) a DNase comprising one or both of the motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76);
- v) a DNase comprising one or both of the motifs P[Q/E]L[W/Y] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO:78);
- 10 vi) a DNase selected from: a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 1, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 2, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 3, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 4, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 5, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 6, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 7, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 8, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 9, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 10, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 11, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 12, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 13, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 14, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 15, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 16, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 17, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 18, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 19, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 20, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 21, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 22, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 23, a polypeptide having at least 80% sequence identity to the polypeptide shown



polypeptide shown in SEQ ID NO: 55, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 56, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 57, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 58, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 59, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 60, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 61, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 62, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 63, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 64, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 65, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 66, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 67, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 68; and

- b) at least 0.001 ppm e.g. 0.1 ppm or 1 ppm of one or more carbohydrase, wherein the carbohydrase is selected from the group consisting of;
- i. a cellulase selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86;
- ii. a xylanase selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87;
- iii. a mannanase selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82;
- iv. an amylase selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least

98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91;

10 and

- c) at least one additional component e.g. cleaning component, preferably selected from surfactants, builders, bleach components, polymers and dispersing agents.

The carbohydrase and DNase may be included in the cleaning composition of the present invention at a level of from 0.01 to 1000 ppm, from 1 ppm to 1000 ppm, from 10 ppm to 1000 ppm, from 50 ppm to 1000 ppm, from 100 ppm to 1000 ppm, from 150 ppm to 1000 ppm, from 200 ppm to 1000 ppm, from 250 ppm to 1000 ppm, from 250 ppm to 750 ppm, from 250 ppm to 500 ppm. The DNases above may be combined with carbohydrases to form a blend to be added to the wash liquor solution according to the invention. The concentration of the DNase in the wash liquor solution is typically in the range of wash liquor from 0.00001 ppm to 10 ppm, from 0.00002 ppm to 10 ppm, from 0.0001 ppm to 10 ppm, from 0.0002 ppm to 10 ppm, from 0.001 ppm to 10 ppm, from 0.002 ppm to 10 ppm, from 0.01 ppm to 10 ppm, from 0.02 ppm to 10 ppm, 0.1 ppm to 10 ppm, from 0.2 ppm to 10 ppm, from 0.5 ppm to 5 ppm. The concentration of the carbohydrases in the wash liquor solution is typically in the range of wash liquor from 0.00001 ppm to 10 ppm, from 0.00002 ppm to 10 ppm, from 0.0001 ppm to 10 ppm, from 0.0002 ppm to 10 ppm, from 0.001 ppm to 10 ppm, from 0.002 ppm to 10 ppm, from 0.01 ppm to 10 ppm, from 0.02 ppm to 10 ppm, 0.1 ppm to 10 ppm, from 0.2 ppm to 10 ppm, from 0.5 ppm to 5 ppm. The DNases may be combined with any of the carbohydrases below to form a blend to be added to a composition according to the invention.

30 One embodiment relates to a cleaning composition comprising a DNase, a carbohydrase and at least one cleaning component, wherein the amount of DNase in the composition is from 0.01 to 1000 ppm and the amount of carbohydrase is from 0.01 to 1000 ppm.

The invention relates to cleaning compositions comprising an enzyme combination of the present invention in combination with one or more additional cleaning composition component(s).  
35 The choice of additional components is within the skill of the artisan and includes conventional ingredients, including the exemplary non-limiting components set forth below.



The choice of cleaning components may include, for textile care, the consideration of the type of textile to be cleaned, the type and/or degree of soiling, the temperature at which cleaning is to take place, and the formulation of the detergent product. Although components mentioned below are categorized by general header according to a particular functionality, this is not to be construed as a limitation, as a component may comprise additional functionalities as will be appreciated by the skilled artisan.

### Surfactants

The detergent composition may comprise one or more surfactants, which may be anionic and/or cationic and/or non-ionic and/or semi-polar and/or zwitterionic, or a mixture thereof. In a particular embodiment, the detergent composition includes a mixture of one or more nonionic surfactants and one or more anionic surfactants. The surfactant(s) is typically present at a level of from about 0.1% to 60% by weight, such as about 0.1% to about 15%, such as about 1% to about 40%, or about 3% to about 20%, or about 3% to about 10%. The surfactant(s) is chosen based on the desired cleaning application, and may include any conventional surfactant(s) known in the art. When included therein the detergent will usually contain from about 1% to about 40% by weight of an anionic surfactant, such as from about 5% to about 30%, including from about 5% to about 15%, or from about 15% to about 20%, or from about 20% to about 25% of an anionic surfactant. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenylyl/tetradecenylyl succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or salt of fatty acids (soap), and combinations thereof.

When included therein the detergent will usually contain from about 1% to about 40% by weight of a cationic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to about 10%, such as from about 3% to about 5%, from about 8% to about 12% or from about 10% to about 12%. Non-limiting examples of cationic surfactants include alkyldimethylethanolamine quat (ADMEAQ), cetyltrimethylammonium bromide (CTAB), dimethyldistearylammonium chloride (DSDMAC), and alkylbenzyltrimethylammonium, alkyl quaternary ammonium compounds, alkoxyated quaternary ammonium (AQA) compounds, ester quats, and combinations thereof.

When included therein the detergent will usually contain from about 0.2% to about 40% by weight of a nonionic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to about 10%, such as from about 3% to about 5%, from about 8% to about 12%, or from about 10% to about 12%. Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxyated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxyated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof. When included therein the detergent will usually contain from about 0.01% to about 10 % by weight of a semipolar surfactant. Non-limiting examples of semipolar surfactants include amine oxides (AO) such as alkyldimethylamineoxide, N-(coco alkyl)-N,N-dimethylamine oxide and N-(tallow-alkyl)-N,N-bis(2-hydroxyethyl)amine oxide, and combinations thereof. When included therein the detergent will usually contain from about 0.01% to about 10 % by weight of a zwitterionic surfactant. Non-limiting examples of zwitterionic surfactants include betaines such as alkyldimethylbetaines, sulfobetaines, and combinations thereof.

#### Builders and Co-Builders

The detergent composition may contain about 0-65% by weight, such as about 5% to about 50%, such as from about 0.5 to about 20% of a detergent builder or co-builder, or a mixture thereof. In a dish wash detergent, the level of builder is typically 40-65%, particularly 50-65%. The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in cleaning detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), ethanolamines such as 2-aminoethan-1-ol (MEA), diethanolamine (DEA, also known as 2,2'-iminodiethan-1-ol), triethanolamine (TEA, also known as 2,2',2"-nitrilotriethan-1-ol), and (carboxymethyl)inulin (CMI), and combinations thereof.

The detergent composition may also contain 0-50% by weight, such as about 5% to about 30%, of a detergent co-builder. The detergent composition may include a co-builder alone, or in combination with a builder, for example a zeolite builder. Non-limiting examples of co-builders include homopolymers of polyacrylates or copolymers thereof, such as poly(acrylic acid) (PAA) or copoly(acrylic acid/maleic acid) (PAA/PMA). Further non-limiting examples include citrate, chelators

such as aminocarboxylates, aminopolycarboxylates and phosphonates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2"-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-N,N'-disuccinic acid (EDDS), methylglycinediacetic acid (MGDA), glutamic acid-N,N'-diacetic acid (GLDA), 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetra(methylenephosphonic acid) (EDTMPA), diethylenetriaminepentakis(methylenephosphonic acid) (DTMPA or DTPMPA), N-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N'-diacetic acid (ASDA), aspartic acid-N-monopropionic acid (ASMP), iminodisuccinic acid (IDA), N-(2-sulfomethyl)-aspartic acid (SMAS), N-(2-sulfoethyl)-aspartic acid (SEAS), N-(2-sulfomethyl)-glutamic acid (SMGL), N-(2-sulfoethyl)-glutamic acid (SEGL), N-methyliminodiacetic acid (MIDA), oalanine-N,N'-diacetic acid ( $\alpha$ -ALDA), serine-N,N'-diacetic acid (SEDA), isoserine-N,N'-diacetic acid (ISDA), phenylalanine-N,N'-diacetic acid (PHDA), anthranilic acid-N,N'-diacetic acid (ANDA), sulfanilic acid-N,N'-diacetic acid (SLDA), taurine-N,N'-diacetic acid (TUDA) and sulfomethyl-N,N'-diacetic acid (SMDA), N-(2-hydroxyethyl)ethylenediamine-N,N',N"-triacetic acid (HEDTA), diethanolglycine (DEG), diethylenetriamine penta(methylenephosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), and combinations and salts thereof. Further exemplary builders and/or co-builders are described in, e.g., WO 09/102854, US 5977053

## 20 Bleaching Systems

The detergent may contain 0-30% by weight, such as about 1% to about 20%, such as from about 0.01 to about 10 wt% of a bleaching system. Any bleaching system comprising components known in the art for use in cleaning detergents may be utilized. Suitable bleaching system components include sources of hydrogen peroxide; sources of peracids; and bleach catalysts or boosters.

Sources of hydrogen peroxide:

Suitable sources of hydrogen peroxide are inorganic persalts, including alkali metal salts such as sodium percarbonate and sodium perborates (usually mono- or tetrahydrate), and hydrogen peroxide—urea (1/1).

30 Sources of peracids:

Peracids may be (a) incorporated directly as preformed peracids or (b) formed in situ in the wash liquor from hydrogen peroxide and a bleach activator (perhydrolysis) or (c) formed in situ in the wash liquor from hydrogen peroxide and a perhydrolase and a suitable substrate for the latter, e.g., an ester.

35 a) Suitable preformed peracids include, but are not limited to, peroxybenzoic acid and its ring-substituted derivatives, peroxy-naphthoic acid, peroxyphthalic acid, peroxyauric acid, peroxysearic acid,  $\epsilon$ -phthalimidoperoxypropionic acid

[phthalimidoperoxyhexanoic acid (PAP)], and o-carboxybenzamidoperoxyacetic acid; aliphatic and aromatic diperoxydicarboxylic acids such as diperoxydodecanedioic acid, diperoxyazelaic acid, diperoxysebacic acid, diperoxybrassylic acid, 2-decyldiperoxybutanedioic acid, and diperoxyphthalic, -isophthalic and -terephthalic acids; perimidic acids; peroxymonosulfuric acid; peroxydisulfuric acid; peroxyphosphoric acid; peroxydicarboxylic acid; and mixtures of said compounds. It is understood that the peracids mentioned may in some cases be best added as suitable salts, such as alkali metal salts (e.g., Oxone®) or alkaline earth-metal salts.


b) Suitable bleach activators include those belonging to the class of esters, amides, imides, nitriles or anhydrides and, where applicable, salts thereof. Suitable examples are tetraacetylenediamine (TAED), sodium 4-[(3,5,5-trimethylhexanoyl)oxy]benzene-1-sulfonate (ISONOBS), sodium 4-(dodecanoyloxy)benzene-1-sulfonate (LOBS), sodium 4-(decanoyloxy)benzene-1-sulfonate, 4-(decanoyloxy)benzoic acid (DOBA), sodium 4-(nonanoyloxy)benzene-1-sulfonate (NOBS), and/or those disclosed in W098/17767. A particular family of bleach activators of interest was disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like triacetin has the advantage that they are environmentally friendly. Furthermore, acetyl triethyl citrate and triacetin have good hydrolytical stability in the product upon storage and are efficient bleach activators. Finally, ATC is multifunctional, as the citrate released in the perhydrolysis reaction may function as a builder.

#### Bleach catalysts and boosters

The bleaching system may also include a bleach catalyst or booster.

Some non-limiting examples of bleach catalysts that may be used in the compositions of the present invention include manganese oxalate, manganese acetate, manganese-collagen, cobalt-amine catalysts and manganese triazacyclononane (MnTACN) catalysts; particularly preferred are complexes of manganese with 1,4,7-trimethyl-1,4,7-triazacyclononane (Me<sub>3</sub>-TACN) or 1,2,4,7-tetramethyl-1,4,7-triazacyclononane (Me<sub>4</sub>-TACN), in particular Me<sub>3</sub>-TACN, such as the dinuclear manganese complex [(Me<sub>3</sub>-TACN)Mn(0)3Mn(Me<sub>3</sub>-TACN)](PF<sub>6</sub>)<sub>2</sub>, and [2,2',2''-nitrioltris(ethane-1,2-diylazanylylidene -KN-methanylylidene)triphenolato-K30]manganese(III). The bleach catalysts may also be other metal compounds; such as iron or cobalt complexes.

In some embodiments, where a source of a peracid is included, an organic bleach catalyst or bleach booster may be used having one of the following formulae:



(iii) and mixtures thereof; wherein each R<sub>1</sub> is independently a branched alkyl group containing from 9 to 24 carbons or linear alkyl group containing from 11 to 24 carbons, preferably each R<sub>1</sub> is independently a branched alkyl group containing from 9 to 18 carbons or linear alkyl group containing from 11 to 18 carbons, more preferably each R<sub>1</sub> is independently selected from the group consisting of 2-propylheptyl, 2-butyloctyl, 2-pentylnonyl, 2-hexyldecyl, dodecyl, tetradecyl, hexadecyl, octadecyl, isononyl, isodecyl, isotridecyl and isopentadecyl.

Other exemplary bleaching systems are described, e.g. in WO2007/087258, WO2007/087244, WO2007/087259, EP1 867708 (Vitamin K) and WO2007/087242. Suitable photobleaches may for example be sulfonated zinc or aluminium phthalocyanines.

#### Metal care agents

Metal care agents may prevent or reduce the tarnishing, corrosion or oxidation of metals, including aluminium, stainless steel and non-ferrous metals, such as silver and copper. Suitable examples include one or more of the following:

(a) benzotriazoles, including benzotriazole or bis-benzotriazole and substituted derivatives thereof. Benzotriazole derivatives are those compounds in which the available substitution sites on the aromatic ring are partially or completely substituted. Suitable substituents include linear or branch-chain C<sub>1</sub>-C<sub>20</sub>- alkyl groups (e.g., C<sub>1</sub>-C<sub>20</sub>- alkyl groups) and hydroxyl, thio, phenyl or halogen such as fluorine, chlorine, bromine and iodine.

(b) metal salts and complexes chosen from the group consisting of zinc, manganese, titanium, zirconium, hafnium, vanadium, cobalt, gallium and cerium salts and/or complexes, the metals being in one of the oxidation states II, III, IV, V or VI. In one aspect, suitable metal salts and/or metal complexes may be chosen from the group consisting of Mn(II) sulphate, Mn(II) citrate, Mn(II) stearate, Mn(II) acetylacetonate, K<sup>+</sup>TiF<sub>6</sub> (e.g., K<sub>2</sub>TiF<sub>6</sub>), K<sup>+</sup>ZrF<sub>6</sub> (e.g., K<sub>2</sub>ZrF<sub>6</sub>), CoSO<sub>4</sub>, Co(NO<sub>2</sub>)<sub>2</sub> and Ce(NO<sub>3</sub>)<sub>3</sub>, zinc salts, for example zinc sulphate, hydrozincite or zinc acetate.;

(c) silicates, including sodium or potassium silicate, sodium disilicate, sodium metasilicate, crystalline phyllosilicate and mixtures thereof.

Further suitable organic and inorganic redox-active substances that act as silver/copper corrosion inhibitors are disclosed in WO 94/26860 and WO 94/26859. Preferably the composition of the invention comprises from 0.1 to 5% by weight of the composition of a metal care agent, preferably the metal care agent is a zinc salt.

Hydrotropes

The detergent may contain 0-10% by weight, for example 0-5% by weight, such as about 0.5 to about 5%, or about 3% to about 5%, of a hydrotrope. Any hydrotrope known in the art for use in detergents may be utilized. Non-limiting examples of hydrotropes include sodium benzenesulfonate, sodium p-toluene sulfonate (STS), sodium xylene sulfonate (SXS), sodium cumene sulfonate (SCS), sodium cymene sulfonate, amine oxides, alcohols and polyglycoethers, sodium hydroxynaphthoate, sodium hydroxynaphthalene sulfonate, sodium ethylhexyl sulfate, and combinations thereof.

Polymers

The detergent may contain 0-10% by weight, such as 0.5-5%, 2-5%, 0.5-2% or 0.2-1% of a polymer. Any polymer known in the art for use in detergents may be utilized. The polymer may function as a co-builder as mentioned above, or may provide antiredeposition, fiber protection, soil release, dye transfer inhibition, grease cleaning and/or anti-foaming properties. Some polymers may have more than one of the above-mentioned properties and/or more than one of the below-mentioned motifs. Exemplary polymers include (carboxymethyl)cellulose (CMC), polyvinyl alcohol (PVA), poly(vinylpyrrolidone) (PVP), poly(ethyleneglycol) or poly(ethylene oxide) (PEG), ethoxylated poly(ethyleneimine), carboxymethyl inulin (CMI), and polycarboxylates such as PAA, PAA/PMA, poly-aspartic acid, and lauryl methacrylate/acrylic acid copolymers, hydrophobically modified CMC (HM-CMC) and silicones, copolymers of terephthalic acid and oligomeric glycols, copolymers of poly(ethylene terephthalate) and poly(oxyethene terephthalate) (PET-POET), PVP, poly(vinylimidazole) (PVI), poly(vinylpyridine-N-oxide) (PVPO or PVPNO) and polyvinylpyrrolidone-vinylimidazole (PVPVI). Suitable examples include PVP-K15, PVP-K30, ChromaBond S-400, ChromaBond S-403E and Chromabond S-100 from Ashland Aqualon, and Sokalan® HP 165, Sokalan® HP 50 (Dispersing agent), Sokalan® HP 53 (Dispersing agent), Sokalan® HP 59 (Dispersing agent), Sokalan® HP 56 (dye transfer inhibitor), Sokalan® HP 66 K (dye transfer inhibitor) from BASF. Further exemplary polymers include sulfonated polycarboxylates, polyethylene oxide and polypropylene oxide (PEO-PPO) and diquaternium ethoxy sulfate. Other exemplary polymers are disclosed in, e.g., WO 2006/130575. Salts of the above-mentioned polymers are also contemplated. Particularly preferred polymer is ethoxylated homopolymer Sokalan® HP 20 from BASF.

Fabric hueing agents

The detergent compositions of the present invention may also include fabric hueing agents such as dyes or pigments, which when formulated in detergent compositions can deposit onto a fabric when said fabric is contacted with a wash liquor comprising said detergent compositions and thus altering the tint of said fabric through absorption/reflection of visible light. Fluorescent whitening agents emit at least some visible light. In contrast, fabric hueing agents

alter the tint of a surface as they absorb at least a portion of the visible light spectrum. Suitable fabric hueing agents include dyes and dye-clay conjugates, and may also include pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and Basic Red, or mixtures thereof, for example as described in WO2005/03274, WO2005/03275, WO2005/03276 and EP1 876226 (hereby incorporated by reference). The detergent composition preferably comprises from about 0.00003 wt% to about 0.2 wt%, from about 0.00008 wt% to about 0.05 wt%, or even from about 0.0001 wt% to about 0.04 wt% fabric hueing agent. The composition may comprise from 0.0001 wt% to 0.2 wt% fabric hueing agent, this may be especially preferred when the composition is in the form of a unit dose pouch. Suitable hueing agents are also disclosed in, e.g. WO 2007/087257 and WO2007/087243.

### Enzymes

The detergent additive as well as the detergent composition may comprise one or more additional enzymes such as one or more proteases lipase, cutinase, pectinase, arabinase, galactanase, xylanase, oxidase, e.g., a laccase, and/or peroxidase.

In general, the properties of the selected enzyme(s) should be compatible with the selected detergent, (i.e., pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

### **Proteases**

The term "protease activity" means a proteolytic activity (EC 3.4). Proteases usable in cleaning compositions of the present invention are mainly endopeptidases (EC 3.4.21). There are several protease activity types: The three main activity types are: trypsin-like where there is cleavage of amide substrates following Arg or Lys at P1, chymotrypsin-like where cleavage occurs following one of the hydrophobic amino acids at P1, and elastase-like with cleavage following an Ala at P1.

The most widely used proteases in the detergent industry such as laundry and dish wash are the serine proteases. Serine proteases is a subgroup of proteases characterised by having a serine in the active site, which forms a covalent adduct with the substrate. Serine proteases are characterized by having two active site amino acid residues apart from the serine, namely a histidine residue and an aspartic acid residue. Subtilase refer to a sub-group of serine protease according to Siezen *et al.*, 1991, *Protein Engng.* 4: 719-737 and Siezen *et al.*, 1997, *Protein Science* 6: 501-523. The subtilases may be divided into 6 sub-divisions, *i.e.*, the Subtilisin family, the Thermitase family, the Proteinase K family, the Lantibiotic peptidase family, the Kexin family and the Pyrolysin family.

Suitable proteases for the compositions of the invention include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloprotease protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.

Examples of subtilases are those derived from *Bacillus* such as *Bacillus lentus*, *Bacillus alkalophilus*, *Bacillus subtilis*, *Bacillus amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US7262042 and WO09/021867, and subtilisin lentus, subtilisin Novo, subtilisin Carlsberg, *Bacillus licheniformis*, subtilisin BPN', subtilisin 309, subtilisin 147 and subtilisin 168 described in WO89/06279 and protease PD138 described in (WO93/18140). Other useful proteases may be those described in W092/175177, WO01/016285, WO02/026024 and WO02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO89/06270, W094/25583 and WO05/040372, and the chymotrypsin proteases derived from Cellulomonas described in WO05/052161 and WO05/052146.

A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in W095/23221, and variants thereof which are described in WO92/21760, W095/23221, EP1921 147 and EP1921 148.

Examples of metalloproteases are the neutral metalloproteases as described in WO07/044993 (Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*.

Examples of useful proteases are the variants described in: W092/1 9729, WO96/034946, WO98/201 15, WO98/201 16, WO99/01 1768, WO01/44452, WO03/006602, WO04/03186, WO04/041979, WO07/006305, W01 1/036263, W01 1/036264, especially protease variants comprising a substitution in one or more of the following positions: 3, 4, 9, 15, 24, 27, 42, 55, 59, 60, 66, 74, 85, 96, 97, 98, 99, 100, 101, 102, 104, 116, 118, 121, 126, 127, 128, 154, 156, 157, 158, 161, 164, 176, 179, 182, 185, 188, 189, 193, 198, 199, 200, 203, 206, 211, 212, 216, 218, 226, 229, 230, 239, 246, 255, 256, 268 and 269, wherein the positions correspond to the positions of the *Bacillus lentus* protease shown in SEQ ID NO 69. More preferred the protease variants may comprise one or more of the mutations selected from the group consisting of: S3T, V4I, S9R, S9E, A15T, S24G, S24R, K27R, N42R, S55P, G59E, G59D, N60D, N60E, V66A, N74D, S85R, A96S, S97G, S97D, S97A, S97SD, S99E, S99D, S99G, S99M, S99N, S99R, S99H, S101A, V102I, V102Y, V102N, S104A, G116V, G116R, H118D, H118N, A120S, S126L, P127Q, S128A, S154D, A156E, G157D, G157P, S158E, Y161A, R164S, Q176E, N179E, S182E, Q185N, A188P, G189E, V193M, N198D, V199I, Y203W, S206G, L21 1Q, L21 1D, N212D, N212S, M216S, A226V, K229L, Q230H, Q239R, N246K, N255W, N255D, N255E, L256E, L256D T268A and



R269H. The protease variants are preferably variants of the *Bacillus lentus* protease (Savinase®) shown in SEQ ID NO 79 or the *Bacillus amylolichenifaciens* protease (BPN') shown in SEQ ID NO 80. The protease variants preferably have at least 80 % sequence identity to SEQ ID NO 79 or SEQ ID NO 80.

5 A protease variant comprising a substitution at one or more positions corresponding to positions 171, 173, 175, 179, or 180 of SEQ ID NO: 81, wherein said protease variant has a sequence identity of at least 75% but less than 100% to SEQ ID NO: 81.

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Duralase™, Durazym™, Relase®, Relase® Ultra, Savinase®, Savinase®  
 10 Ultra, Primase®, Polarzyme®, Kannase®, Liquanase®, Liquanase® Ultra, Ovozyme®, Coronase®, Coronase® Ultra, Blaze®, Blaze Eivity® 100T, Blaze Eivity® 125T, Blaze Eivity® 150T, Neutrase®, Everlase® and Esperase® (Novozymes A/S), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Purafect Ox®, Purafect OxP®, Puramax®, FN2®, FN3®, FN4®, Excellase®, Excellenz P1000™, Excellenz P1250™, Eraser®, Preferenz P100™,  
 15 Purafect Prime®, Preferenz P110™, Effectenz P1000™, Purafect®™, Effectenz P1050™, Purafect Ox®™, Effectenz P2000™, Purafast®, Properase®, Opticlean® and Optimase® (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and variants hereof (Henkel AG) and KAP (*Bacillus alkalophilus subtilisin*) from Kao.

#### **Peroxidases/Oxidases**

20 Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from Coprinus, e.g., from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257. Commercially available peroxidases include Guardzyme™ (Novozymes A/S).

#### **Lipases and Cutinases:**

25 Suitable lipases and cutinases include those of bacterial or fungal origin. Chemically modified or protein engineered mutant enzymes are included. Examples include lipase from *Thermomyces*, e.g. from *T. lanuginosus* (previously named *Humicola lanuginosa*) as described in EP258068 and EP305216, cutinase from *Humicola*, e.g. *H. insolens* (WO96/13580), lipase  
 30 from strains of *Pseudomonas* (some of these now renamed to *Burkholderia*), e.g. *P. alcaligenes* or *P. pseudoalcaligenes* (EP218272), *P. cepacia* (EP331376), *P. sp.* strain SD705 (WO95/06720 & WO96/27002), *P. wisconsinensis* (WO96/12012), GDSL-type *Streptomyces* lipases (W0 10/065455), cutinase from *Magnaporthe grisea* (W0 10/107560), cutinase from *Pseudomonas mendocina* (US5,389,536), lipase from *Thermobifida fusca* (W0 11/084412),  
 35 *GeoBacillus stearothermophilus* lipase (W0 11/084417), lipase from *Bacillus subtilis* (W01 1/084599), and lipase from *Streptomyces griseus* (W01 1/150157) and *S. pristinaespiralis* (WQ12/137147).

Other examples are lipase variants such as those described in EP407225, WO92/05249, WO94/01541, WO94/25578, WO95/14783, WO95/30744, WO95/35381, WO95/22615, WO96/00292, WO97/04079, WO97/07202, WO00/34450, WO00/60063, WO01/92502, WO07/87508 and WO09/1 09500.

5 Preferred commercial lipase products include Lipolase™, Lipex™; Lipolex™ and Lipoclean™ (Novozymes A/S), Lumafast (originally from Genencor) and Lipomax (originally from Gist-Brocades).

10 Still other examples are lipases sometimes referred to as acyltransferases or perhydrolases, e.g. acyltransferases with homology to *Candida antarctica* lipase A (WO10/1 11143), acyltransferase from *Mycobacterium smegmatis* (WO05/56782), perhydrolases from the CE 7 family (WO09/67279), and variants of the *M. smegmatis* perhydrolase in particular the S54V variant used in the commercial product Gentle Power Bleach from Huntsman Textile Effects Pte Ltd (WO1 0/1 00028).

### Peroxidases/Oxidases

15 A peroxidase according to the invention is a peroxidase enzyme comprised by the enzyme classification EC 1.1 1.1.7, as set out by the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (IUBMB), or any fragment derived therefrom, exhibiting peroxidase activity.

20 Suitable peroxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinopsis*, e.g., from *C. cinerea* (EP 179,486), and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

25 A suitable peroxidase includes a haloperoxidase enzyme, such as chloroperoxidase, bromoperoxidase and compounds exhibiting chloroperoxidase or bromoperoxidase activity. Haloperoxidases are classified according to their specificity for halide ions. Chloroperoxidases (E.C. 1.1.1.10) catalyze formation of hypochlorite from chloride ions. Preferably, the haloperoxidase is a vanadium haloperoxidase, i.e., a vanadate-containing haloperoxidase. Haloperoxidases have been isolated from many different fungi, in particular from the fungus group dematiaceous hyphomycetes, such as *Caldariomyces*, e.g., *C. fumago*, *Alternaria*, *Curvularia*, 30 e.g., *C. verruculosa* and *C. inaequalis*, *Drechslera*, *Ulocladium* and *Botrytis*.

Haloperoxidases have also been isolated from bacteria such as *Pseudomonas*, e.g., *P. pyrrocinia* and *Streptomyces*, e.g., *S. aureofaciens*.

35 A suitable oxidase includes in particular, any laccase enzyme comprised by the enzyme classification EC 1.10.3.2, or any fragment derived therefrom exhibiting laccase activity, or a compound exhibiting a similar activity, such as a catechol oxidase (EC 1.10.3.1), an o-aminophenol oxidase (EC 1.10.3.4), or a bilirubin oxidase (EC 1.3.3.5). Preferred laccase enzymes are enzymes of microbial origin. The enzymes may be derived from plants, bacteria or

fungi (including filamentous fungi and yeasts). Suitable examples from fungi include a laccase derivable from a strain of *Aspergillus*, *Neurospora*, e.g., *N. crassa*, *Podospora*, *Botrytis*, *Collybia*, *Fomes*, *Lentinus*, *Pleurotus*, *Trametes*, e.g., *T. villosa* and *T. versicolor*, *Rhizoctonia*, e.g., *R. solani*, *Coprinopsis*, e.g., *C. cinerea*, *C. comatus*, *C. friesii*, and *C. plicatilis*, *Psathyrella*, e.g., *P. condelleana*, *Panaeolus*, e.g., *P. papilionaceus*, *Myceliophthora*, e.g., *M. thermophila*, *Schytalidium*, e.g., *S. thermophilum*, *Polyporus*, e.g., *P. pinsitus*, *Phlebia*, e.g., *P. radiata* (WO 92/01046), or *Coriolus*, e.g., *C. hirsutus* (JP 2238885). Suitable examples from bacteria include a laccase derivable from a strain of *Bacillus*. A laccase derived from *Coprinopsis* or *Myceliophthora* is preferred; in particular, a laccase derived from *Coprinopsis cinerea*, as disclosed in WO 97/08325; or from *Myceliophthora thermophila*, as disclosed in WO 95/33836.

#### Dispersants

The cleaning compositions of the present invention can also contain dispersants. In particular, powdered detergents may comprise dispersants. Suitable water-soluble organic materials include the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Suitable dispersants are for example described in *Powdered Detergents*, Surfactant science series volume 71, Marcel Dekker, Inc.

#### Dye Transfer Inhibiting Agents

The cleaning compositions of the present invention may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in a subject composition, the dye transfer inhibiting agents may be present at levels from about 0.0001 % to about 10%, from about 0.01% to about 5% or even from about 0.1% to about 3% by weight of the composition.

#### Fluorescent whitening agent

The cleaning compositions of the present invention will preferably also contain additional components that may tint articles being cleaned, such as fluorescent whitening agent or optical brighteners. Where present the brightener is preferably at a level of about 0.01% to about 0.5%. Any fluorescent whitening agent suitable for use in a laundry detergent composition may be used in the composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulfonic acid derivatives, diarylpyrazoline derivatives and bisphenyl-distyryl derivatives. Examples of the diaminostilbene-sulfonic acid derivative type of fluorescent whitening agents include the sodium salts of: 4,4'-bis-

(2-diethanolamino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(2,4-dianilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(4-phenyl-1,2,3-triazol-2-yl)stilbene-2,2'-disulfonate and sodium 5-(2H-naphtho[1,2-d][1,2,3]triazol-2-yl)-2-[(E)-2-phenylvinyl]benzenesulfonate. Preferred fluorescent whitening agents are Tinopal DMS and Tinopal CBS available from Ciba-Geigy AG, Basel, Switzerland. Tinopal DMS is the disodium salt of 4,4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate. Tinopal CBS is the disodium salt of 2,2'-bis-(phenyl-styryl)-disulfonate. Also preferred are fluorescent whitening agents is the commercially available Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India. Other fluorescers suitable for use in the invention include the 1-3-diaryl pyrazolines and the 7-alkylaminocoumarins. Suitable fluorescent brightener levels include lower levels of from about 0.01, from 0.05, from about 0.1 or even from about 0.2 wt % to upper levels of 0.5 or even 0.75 wt%.

#### Soil release polymers

The cleaning compositions of the present invention may also include one or more soil release polymers which aid the removal of soils from fabrics such as cotton and polyester based fabrics, in particular the removal of hydrophobic soils from polyester based fabrics. The soil release polymers may for example be nonionic or anionic terephthalate based polymers, polyvinyl caprolactam and related copolymers, vinyl graft copolymers, polyester polyamides see for example Chapter 7 in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc. Another type of soil release polymers is amphiphilic alkoxyated grease cleaning polymers comprising a core structure and a plurality of alkoxyate groups attached to that core structure. The core structure may comprise a polyalkylenimine structure or a polyalkanolamine structure as described in detail in WO 2009/087523 (hereby incorporated by reference). Furthermore, random graft co-polymers are suitable soil release polymers. Suitable graft co-polymers are described in more detail in WO 2007/138054, WO 2006/108856 and WO 2006/113314 (hereby incorporated by reference). Suitable polyethylene glycol polymers include random graft co-polymers comprising: (i) hydrophilic backbone comprising polyethylene glycol; and (ii) side chain(s) selected from the group consisting of: C4-C25 alkyl group, polypropylene, polybutylene, vinyl ester of a saturated C1-C6 mono-carboxylic acid, C1-C6 alkyl ester of acrylic or methacrylic acid, and mixtures thereof. Suitable polyethylene glycol polymers have a polyethylene glycol backbone with random grafted polyvinyl acetate side chains. The average molecular weight of the polyethylene glycol backbone can be in the range of from 2,000 Da to 20,000 Da, or from 4,000 Da to 8,000 Da. The molecular weight ratio of the polyethylene glycol backbone to the polyvinyl acetate side chains can be in the range of from 1:1 to 1:5, or from 1:1.2 to 1:2. The average number of graft sites per ethylene oxide units can be less than 1, or less

than 0.8, the average number of graft sites per ethylene oxide units can be in the range of from 0.5 to 0.9, or the average number of graft sites per ethylene oxide units can be in the range of from 0.1 to 0.5, or from 0.2 to 0.4. A suitable polyethylene glycol polymer is Sokalan HP22. Other soil release polymers are substituted polysaccharide structures especially substituted cellulosic structures such as modified cellulose derivatives such as those described in EP 1867808 or WO 2003/040279 (both are hereby incorporated by reference). Suitable cellulosic polymers include cellulose, cellulose ethers, cellulose esters, cellulose amides and mixtures thereof. Suitable cellulosic polymers include anionically modified cellulose, nonionically modified cellulose, cationically modified cellulose, zwitterionically modified cellulose, and mixtures thereof. Suitable cellulosic polymers include methyl cellulose, carboxy methyl cellulose, ethyl cellulose, hydroxyl ethyl cellulose, hydroxyl propyl methyl cellulose, ester carboxy methyl cellulose, and mixtures thereof.

#### Anti-redeposition agents

The cleaning compositions of the present invention may also include one or more anti-redeposition agents such as carboxymethylcellulose (CMC), polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), polyoxyethylene and/or polyethyleneglycol (PEG), homopolymers of acrylic acid, copolymers of acrylic acid and maleic acid, and ethoxylated polyethyleneimines. The cellulose based polymers described under soil release polymers above may also function as anti-redeposition agents.

#### Rheology Modifiers

The cleaning compositions of the present invention may also include one or more rheology modifiers, structurants or thickeners, as distinct from viscosity reducing agents. The rheology modifiers are selected from the group consisting of non-polymeric crystalline, hydroxy-functional materials, polymeric rheology modifiers which impart shear thinning characteristics to the aqueous liquid matrix of a liquid detergent composition. The rheology and viscosity of the detergent can be modified and adjusted by methods known in the art, for example as shown in EP 2169040.

Other suitable cleaning composition components include, but are not limited to, anti-shrink agents, anti-wrinkling agents, bactericides, binders, carriers, dyes, enzyme stabilizers, fabric softeners, fillers, foam regulators, hydrotropes, perfumes, pigments, sod suppressors, solvents, and structurants for liquid detergents and/or structure elasticizing agents.

Formulation of detergent products

The cleaning composition of the invention may be in any convenient form, e.g., a bar, a homogenous tablet, a tablet having two or more layers, a pouch having one or more compartments, a regular or compact powder, a granule, a paste, a gel, or a regular, compact or concentrated liquid.

5 Pouches can be configured as single or multicompartments. It can be of any form, shape and material which is suitable for hold the composition, e.g. without allowing the release of the composition to release of the composition from the pouch prior to water contact. The pouch is made from water soluble film which encloses an inner volume. Said inner volume can be divided into compartments of the pouch. Preferred films are polymeric materials preferably polymers which are  
10 formed into a film or sheet. Preferred polymers, copolymers or derivatives thereof are selected polyacrylates, and water soluble acrylate copolymers, methyl cellulose, carboxy methyl cellulose, sodium dextrin, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, malto dextrin, poly methacrylates, most preferably polyvinyl alcohol copolymers and, hydroxypropyl methyl cellulose (HPMC). Preferably the level of polymer in the film for example PVA is at least about 60%.  
15 Preferred average molecular weight will typically be about 20,000 to about 150,000. Films can also be of blended compositions comprising hydrolytically degradable and water soluble polymer blends such as polylactide and polyvinyl alcohol (known under the Trade reference M8630 as sold by MonoSol LLC, Indiana, USA) plus plasticisers like glycerol, ethylene glycerol, propylene glycol, sorbitol and mixtures thereof. The pouches can comprise a solid laundry cleaning composition or  
20 part components and/or a liquid cleaning composition or part components separated by the water-soluble film. The compartment for liquid components can be different in composition than compartments containing solids: US2009/001 1970 A 1.

Detergent ingredients can be separated physically from each other by compartments in water dissolvable pouches or in different layers of tablets. Thereby negative storage interaction  
25 between components can be avoided. Different dissolution profiles of each of the compartments can also give rise to delayed dissolution of selected components in the wash solution.

A liquid or gel detergent, which is not unit dosed, may be aqueous, typically containing at least 20% by weight and up to 95% water, such as up to about 70% water, up to about 65% water, up to about 55% water, up to about 45% water, up to about 35% water. Other types of liquids,  
30 including without limitation, alkanols, amines, diols, ethers and polyols may be included in an aqueous liquid or gel. An aqueous liquid or gel detergent may contain from 0-30% organic solvent. A liquid or gel detergent may be non-aqueous.

Granular detergent formulations

35 Non-dusting granulates may be produced, e.g. as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000;

ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The DNase may be formulated as a granule for example as a co-granule that combines one or more enzymes. Each enzyme will then be present in more granules securing a more uniform distribution of enzymes in the detergent. This also reduces the physical segregation of different enzymes due to different particle sizes. Methods for producing multi-enzyme co-granulate for the detergent industry is disclosed in the IP.com disclosure IPCOM000200739D.

Another example of formulation of enzymes by the use of co-granulates are disclosed in WO 2013/188331, which relates to a detergent composition comprising (a) a multi-enzyme co-granule; (b) less than 10 wt zeolite (anhydrous basis); and (c) less than 10 wt phosphate salt (anhydrous basis), wherein said enzyme co-granule comprises from 10 to 98 wt% moisture sink component and the composition additionally comprises from 20 to 80 wt% detergent moisture sink component. WO 2013/188331 also relates to a method of treating and/or cleaning a surface, preferably a fabric surface comprising the steps of (i) contacting said surface with the detergent composition as claimed and described herein in aqueous wash liquor, (ii) rinsing and/or drying the surface.

An embodiment of the invention relates to an enzyme granule/particle comprising the DNase and at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase. The granule is composed of a core, and optionally one or more coatings (outer layers) surrounding the core. Typically, the granule/particle size, measured as equivalent spherical diameter (volume based average particle size), of the granule is 20-2000  $\mu\text{m}$ , particularly 50-1500  $\mu\text{m}$ , 100-1500  $\mu\text{m}$  or 250-1200  $\mu\text{m}$ . The core may include additional materials such as fillers, fibre materials (cellulose or synthetic fibres), stabilizing agents, solubilising agents, suspension agents, viscosity regulating agents, light spheres, plasticizers, salts, lubricants and fragrances. The core may include binders, such as synthetic polymer, wax, fat, or carbohydrate. The core may comprise a salt of a multivalent cation, a reducing agent, an antioxidant, a peroxide decomposing catalyst and/or an acidic buffer component, typically as a homogenous blend. The core may consist of an inert particle with the enzyme absorbed into it, or applied onto the surface, e.g., by fluid bed coating. The core may have a diameter of 20-2000  $\mu\text{m}$ , particularly 50-1500  $\mu\text{m}$ , 100-1500  $\mu\text{m}$  or 250-1200  $\mu\text{m}$ . The core can be prepared by granulating a blend of the ingredients, e.g., by a method comprising granulation techniques such as crystallization, precipitation, pan-coating, fluid bed coating, fluid bed agglomeration, rotary

atomization, extrusion, prilling, spheronization, size reduction methods, drum granulation, and/or high shear granulation.

Methods for preparing the core can be found in Handbook of Powder Technology; Particle size enlargement by C. E. Capes; Volume 1; 1980; Elsevier.

5 The core of the enzyme granule/particle may be surrounded by at least one coating, **e.g.**, to improve the storage stability, to reduce dust formation during handling, or for coloring the granule. The optional coating(s) may include a salt coating, or other suitable coating materials, such as polyethylene glycol (PEG), methyl hydroxy-propyl cellulose (MHPC) and polyvinyl alcohol (PVA). Examples of enzyme granules with multiple coatings are shown in WO 93/07263 and WO 97/23606.

10 The coating may be applied in an amount of at least 0.1% by weight of the core, **e.g.**, at least 0.5%, 1% or 5%. The amount may be at most 100%, 70%, 50%, 40% or 30%. The coating is preferably at least 0.1  $\mu\text{m}$  thick, particularly at least 0.5  $\mu\text{m}$ , at least 1  $\mu\text{m}$  or at least 5  $\mu\text{m}$ . In a one embodiment, the thickness of the coating is below 100  $\mu\text{m}$ . In another embodiment, the thickness of the coating is below 60  $\mu\text{m}$ . In an even more particular embodiment the total thickness of the coating is below

15 40  $\mu\text{m}$ . The coating should encapsulate the core unit by forming a substantially continuous layer. A substantially continuous layer is to be understood as a coating having few or no holes, so that the core unit it is encapsulating/enclosing has few or none uncoated areas. The layer or coating should be homogeneous in thickness. The coating can further contain other materials as known in the art, **e.g.**, fillers, antisticking agents, pigments, dyes, plasticizers and/or binders, such as titanium dioxide, kaolin, calcium carbonate or talc. A salt coating may comprise at least 60% by weight w/w of a salt,

20 **e.g.**, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% by weight w/w. The salt may be added from a salt solution where the salt is completely dissolved or from a salt suspension wherein the fine particles is less than 50  $\mu\text{m}$ , such as less than 10  $\mu\text{m}$  or less than 5  $\mu\text{m}$ . The salt coating may comprise a single salt or a mixture of

25 two or more salts. The salt may be water soluble, and may have a solubility at least 0.1 grams in 100 g of water at 20°C, preferably at least 0.5 g per 100 g water, **e.g.**, at least 1 g per 100 g water, **e.g.**, at least 5 g per 100 g water. The salt may be an inorganic salt, **e.g.**, salts of sulfate, sulfite, phosphate, phosphonate, nitrate, chloride or carbonate or salts of simple organic acids (less than 10 carbon atoms, **e.g.**, 6 or less carbon atoms) such as citrate, malonate or acetate. Examples of

30 cations in these salts are alkali or earth alkali metal ions, the ammonium ion or metal ions of the first transition series, such as sodium, potassium, magnesium, calcium, zinc or aluminium. Examples of anions include chloride, bromide, iodide, sulfate, sulfite, bisulfite, thiosulfate, phosphate, monobasic phosphate, dibasic phosphate, hypophosphite, dihydrogen pyrophosphate, tetraborate, borate, carbonate, bicarbonate, metasilicate, citrate, malate, maleate, malonate, succinate, lactate, formate,

35 acetate, butyrate, propionate, benzoate, tartrate, ascorbate or gluconate. In particular alkali- or earth alkali metal salts of sulfate, sulfite, phosphate, phosphonate, nitrate, chloride or carbonate or salts of simple organic acids such as citrate, malonate or acetate may be used. The salt in the coating



may have a constant humidity at 20°C above 60%, particularly above 70%, above 80% or above 85%, or it may be another hydrate form of such a salt (e.g., anhydrate). The salt coating may be as described in WO 00/01793 or WO 2006/034710. Specific examples of suitable salts are NaCl (CH<sub>20</sub>°c=76%), Na<sub>2</sub>CO<sub>3</sub> (CH<sub>20</sub>°c=92%), NaNOS (CH<sub>20</sub>°c=73%), Na<sub>2</sub>HPO<sub>4</sub> (CH<sub>20</sub>°c=95%), Na<sub>3</sub>PO<sub>4</sub> (CH<sub>25</sub>°c=92%), NH<sub>4</sub>Cl (CH<sub>20</sub>°c = 79.5%), (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub> (CH<sub>20</sub>°c = 93,0%), NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> (CH<sub>20</sub>°c = 93.1%), (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (CH<sub>20</sub>°c=81.1%), KCl (CH<sub>20</sub>°c=85%), K<sub>2</sub>HPO<sub>4</sub> (CH<sub>20</sub>°c=92%), KH<sub>2</sub>PO<sub>4</sub> (CH<sub>20</sub>°c=96.5%), KNOs (CH<sub>20</sub>°c=93.5%), Na<sub>2</sub>SO<sub>4</sub> (CH<sub>20</sub>°c=93%), K<sub>2</sub>SO<sub>4</sub> (CH<sub>20</sub>°c=98%), KHSO<sub>4</sub> (CH<sub>20</sub>°c=86%), MgSO<sub>4</sub> (CH<sub>20</sub>°c=90%), ZnSO<sub>4</sub> (CH<sub>20</sub>°c=90%) and sodium citrate (CH<sub>25</sub>°c=86%). Other examples include NaH<sub>2</sub>PO<sub>4</sub>, (NH<sub>4</sub>)H<sub>2</sub>PO<sub>4</sub>, CuSO<sub>4</sub>, Mg(NO<sub>3</sub>)<sub>2</sub> and magnesium acetate. The salt may be in anhydrous form, or it may be a hydrated salt, i.e. a crystalline salt hydrate with bound water(s) of crystallization, such as described in WO 99/32595. Specific examples include anhydrous sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>), anhydrous magnesium sulfate (MgSO<sub>4</sub>), magnesium sulfate heptahydrate (MgSO<sub>4</sub>·7H<sub>2</sub>O), zinc sulfate heptahydrate (ZnSO<sub>4</sub>·7H<sub>2</sub>O), sodium phosphate dibasic heptahydrate (Na<sub>2</sub>HPO<sub>4</sub>·7H<sub>2</sub>O), magnesium nitrate hexahydrate (Mg(NO<sub>3</sub>)<sub>2</sub>(6H<sub>2</sub>O)), sodium citrate dihydrate and magnesium acetate tetrahydrate. Preferably the salt is applied as a solution of the salt, e.g., using a fluid bed.

One embodiment of the present invention provides a granule, which comprises:

(a) a core comprising a DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase, and

(b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

(a) a core comprising a DNase and a cellulase wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, and

(b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 5 (a) a core comprising a DNase and an amylase wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, and
- 10
- 15
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 20 (a) a core comprising a DNase and a mannanase wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, and
- 25
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 30 (a) a core comprising a DNase and an xylanase wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, and
- 35
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 5 (c) a core comprising a DNase and a cellulase wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65, and
- 10 (d) optionally a coating consisting of one or more layer(s) surrounding the core.

20 One embodiment of the invention relates to a granule, which comprises:

- (c) a core comprising a DNase and an amylase wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65, and
- 25 (d) optionally a coating consisting of one or more layer(s) surrounding the core.
- 30
- 35

One embodiment of the invention relates to a granule, which comprises:

- (c) a core comprising a DNase and a mannanase wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65, and
- (d) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (c) a core comprising a DNase and a xylanase wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65, and
- (d) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (a) a core comprising a DNase and a cellulase wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66, and
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (e) a core comprising a DNase and an amylase wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66, and
- (f) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (e) a core comprising a DNase and a mannanase wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66, and
- (f) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (e) a core comprising a DNase and a xylanase wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66, and

- (f) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 5 (c) a core comprising a DNase and a cellulase wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ  
10 ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the  
15 polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67, and

- 20 (d) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 25 (g) a core comprising a DNase and an amylase wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ  
30 ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the  
35 polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67, and

- (h) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 5 (g) a core comprising a DNase and a mannanase wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67, and
- 10 (h) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 15 (g) a core comprising a DNase and a xylanase wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67, and
- 20 (h) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- 25 (e) a core comprising a DNase and a cellulase wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68, and
- 30
- 35

- (f) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (i) a core comprising a DNase and an amylase wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68, and
- (j) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (i) a core comprising a DNase and a mannanase wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68, and
- (j) optionally a coating consisting of one or more layer(s) surrounding the core.

One embodiment of the invention relates to a granule, which comprises:

- (i) a core comprising a DNase and a xylanase wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at



least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68, and

- (j) optionally a coating consisting of one or more layer(s) surrounding the core.

## 5 Uses

The present invention is also directed to methods for using the compositions thereof. Laundry/textile/fabric (House hold laundry washing, Industrial laundry washing). Hard surface cleaning (ADW, car wash, Industrial surface). The compositions of the invention comprise a blend of DNase and carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase and effectively reduce or remove organic components, such as protein and DNA from surfaces such as textiles and hard surfaces e.g. dishes.

The compositions of the invention comprise a blend of DNase and carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase, and effectively reduce or remove organic components, such as mannan, starch, cellulose, xyloglucan and DNA from surfaces such as textiles and hard surfaces e.g. dishes. One embodiment of the invention relates to the use of a cleaning composition comprising a DNase, a carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase and at least one cleaning component for reduction or removal of components of biofilm, such as DNA and at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase, of an item, wherein the item is a textile or a hard surface.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase for deep cleaning of an item, wherein the item is a textile or a surface.

One embodiment of the invention relates to the use of a composition comprising a DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase for reduction or removal of biofilm and/or compounds such as mannan, starch, cellulose, xyloglucan and DNA of an item. One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase for reduction or removal of biofilm and/or compounds such as mannan, starch, cellulose, xyloglucan and DNA of an item such as textile. One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase for deep cleaning when the cleaning composition is applied in e.g. laundry process.

One embodiment of the invention relates to the use of a composition comprising a DNase and carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase for reduction of redeposition or reduction of malodor. One embodiment of the invention relates to

the use of a cleaning composition comprising a DNase and carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase for reduction of redeposition or reduction of malodor.

One embodiment of the invention relates to the use of a cleaning composition comprising  
5 a DNase and carbohydrase, selected from a cellulase, an amylase, a mannanase or a xylanase  
for reduction of redeposition or reduction of malodor when the cleaning composition is applied in  
e.g. laundry process. One embodiment of the invention relates to the use of a cleaning  
composition comprising a DNase and carbohydrase, selected from a cellulase, an amylase, a  
mannanase or a xylanase for reduction of redeposition or reduction of malodor on an item e.g.  
10 textile. In one embodiment, the composition is an anti-redeposition composition.

One embodiment of the invention relates to the use of a cleaning composition  
comprising a DNase and a cellulase for deep cleaning of an item or reduction of redeposition or  
malodor, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%,  
at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%  
15 or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at  
least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at  
least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84,  
a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least  
85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide  
20 shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at  
least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
identity to the polypeptide shown in SEQ ID NO: 86.

One embodiment of the invention relates to the use of a cleaning composition comprising  
a DNase and an amylase for deep cleaning of an item or reduction of redeposition or malodor,  
25 wherein the amylase selected from a polypeptide having at least 60%, at least 65%, at least 70%,  
at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100%  
sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%,  
at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%,  
at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a  
30 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least  
85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide  
shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at  
least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
identity to the polypeptide shown in SEQ ID NO: 91.

One embodiment of the invention relates to the use of a cleaning composition comprising  
a DNase and a mannanase for deep cleaning of an item or reduction of redeposition or malodor,  
wherein the mannanase selected from a polypeptide having at least 60%, at least 65%, at least

70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and a xylanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the xylanase selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and cellulase for deep cleaning of an item or reduction of redeposition or malodor, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and cellulase for deep cleaning of an item or reduction of redeposition or malodor, wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and cellulase for deep cleaning of an item or reduction of redeposition or malodor,

wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%,  
5 at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
10 identity to the polypeptide shown in SEQ ID NO: 86 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and cellulase for deep cleaning of an item or reduction of redeposition or malodor,  
15 wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a  
20 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
25 identity to the polypeptide shown in SEQ ID NO: 86 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and cellulase for deep cleaning of an item or reduction of redeposition or malodor,  
30 wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a  
35 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence

identity to the polypeptide shown in SEQ ID NO: 86 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment of the invention relates to the use of a cleaning composition comprising  
5 a DNase and amylase for deep cleaning of an item or reduction of redeposition or malodor, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%,  
10 at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
15 identity to the polypeptide shown in SEQ ID NO: 91 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and amylase for deep cleaning of an item or reduction of redeposition or malodor,  
20 wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a  
25 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence  
30 identity to the polypeptide shown in SEQ ID NO: 91 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and amylase for deep cleaning of an item or reduction of redeposition or malodor, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least  
35 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%,

at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and amylase for deep cleaning of an item or reduction of redeposition or malodor, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and amylase for deep cleaning of an item or reduction of redeposition or malodor, wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and mannanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and mannanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and mannanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and mannanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 67.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and mannanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the is DNase has at

least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

5 One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and xylanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at  
10 least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and xylanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least  
15 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 65.

20 One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and xylanase for deep cleaning of an item or reduction of redeposition or malodor, wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the is DNase has at  
25 least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 66.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and xylanase for deep cleaning of an item or reduction of redeposition or malodor,  
30 wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ  
35 ID NO 67.

One embodiment of the invention relates to the use of a cleaning composition comprising a DNase and xylanase for deep cleaning of an item or reduction of redeposition or malodor,



wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the is DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 68.

The invention further relates to a method of deep cleaning of an item, wherein the item may be textile or hard surface preferably is a textile,

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- a) contacting the item with a cleaning composition according to the invention; and
- b) and optionally rinsing the item, wherein the item is preferably a textile.

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile.

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, a cellulase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and

b) and optionally rinsing the item, wherein the item is preferably a textile and

wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at

least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86.

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- 5 a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, an amylase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile and
- 10 wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a
- 15 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91.

20 One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, mannanase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01
- 25 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile and
- wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

30 One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, xylanase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01
- 35 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile and
- wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least

70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87.

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- 5 a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, a cellulase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile and
- 10 wherein the cellulase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a
- 15 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86 and wherein the DNase has at least 60%, at
- 20 least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, SEQ ID NO 65, SEQ ID NO 66, SEQ ID NO 67 or SEQ ID NO 68.

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

- 25 a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, an amylase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and
- b) and optionally rinsing the item, wherein the item is preferably a textile and
- 30 wherein the amylase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a
- 35 polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at

least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, SEQ ID NO 65, SEQ ID NO 66, SEQ ID NO 67 or SEQ ID NO 68..

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, mannanase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and

b) and optionally rinsing the item, wherein the item is preferably a textile and wherein the mannanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, SEQ ID NO 65, SEQ ID NO 66, SEQ ID NO 67 or SEQ ID NO 68..

One embodiment of the invention relates to a method of deep cleaning on an item, comprising the steps of:

a) contacting the item with a solution comprising an enzyme mixture comprising a DNase, xylanase; and a cleaning component, wherein the cleaning component is selected from 0.1 to 50 wt% of at least one a surfactant; 0.5 to 30 wt% of at least one builder; and 0.01 to 20 wt% of at least one bleach component; and

b) and optionally rinsing the item, wherein the item is preferably a textile and wherein the xylanase is selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87 and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13, SEQ ID NO 65, SEQ ID NO 66, SEQ ID NO 67 or SEQ ID NO 68..

The invention is further described in the following paragraphs

Paragraph 1 A cleaning composition comprising at least 0.001 ppm DNase and at least 0.001 ppm carbohydrase and a cleaning component, wherein the cleaning component is selected from

a. 0.1 to 15 wt% of at least one a surfactant;

- b. 0.5 to 20 wt% of at least one builder; and
- c. 0.01 to 10 wt% of at least one bleach component.

Paragraph 2. The cleaning composition according to paragraph 1, wherein the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73), ASXNRSKG (SEQ ID NO: 74) and the carbohydrase is a cellulase.

Paragraph 3. The cleaning composition according to paragraph 1, wherein the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73), ASXNRSKG (SEQ ID NO: 74) and the carbohydrase is a mannanase.

Paragraph 4. The cleaning composition according to paragraph 1, wherein the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73), ASXNRSKG (SEQ ID NO: 74) and the carbohydrase is a amylase.

Paragraph 5. The cleaning composition according to any of paragraphs 1 to 4, wherein the DNase is selected from the group of polypeptides:

- a) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 1,
- b) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 2,
- c) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 3,
- d) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 4,
- e) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 5,
- f) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 6,
- g) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 7,
- h) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 8,
- i) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 9,
- j) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 10,

- k) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 11,
- l) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 12,
- m) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 13,
- n) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 14,
- o) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 15,
- p) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 16,
- q) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 17,
- r) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 18,
- s) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 19,
- t) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 20,
- u) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 21,
- v) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 22,
- w) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 23,
- x) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 24, and
- y) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 25,

and wherein the carbohydrase is selected from the group consisting of;

- i. a cellulase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 86;

- 5
- 10
- ii. a xylanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 87;
  - iii. a mannanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 82; and
  - iv. an amylase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 91.

Paragraph 6. The cleaning composition according to paragraph 1 wherein the DNase comprises one or both of the motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76) and the carbohydrase is a cellulase.

Paragraph 7. The cleaning composition according to paragraph 1 wherein the DNase comprises one or both of the motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76) and the carbohydrase is a mannanase.

Paragraph 8. The cleaning composition according to paragraph 1 wherein the DNase comprise one or both of the motifs [V/I]PL[S/A]NAWK (SEQ ID NO: 75) or NPQL (SEQ ID NO: 76) and the carbohydrase is a amylase.

Paragraph 9. The cleaning composition according to any of paragraphs 1 and 6 to 8, wherein the DNase is selected from the group of polypeptides:

- a) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 26,
- b) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 27,
- c) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 28,
- d) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 29,
- e) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 30,
- f) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 31,

- g) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 32,
- h) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 33,
- i) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 34,
- j) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 35,
- k) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 36,
- l) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 37,
- m) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 38,

and wherein the carbohydrase is selected from the group consisting of;

- i. a cellulase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 86;
- ii. a xylanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 87;
- iii. a mannanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 82; and
- iv. an amylase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 91.

Paragraph 10. The cleaning composition according to paragraph 1 wherein the DNase comprises one or both of the motifs P[Q/E]L[W/Y] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO: 78) and the carbohydrase is a cellulase.



Paragraph 11. The cleaning composition according to paragraph 1 wherein the DNase comprise one or both of the motifs P[Q/E]L[W/Y] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO: 78) and the carbohydrase is a mannanase.

Paragraph 12. The cleaning composition according to paragraph 1 wherein the DNase comprise one or both of the motifs P[Q/E]L[W/Y] (SEQ ID NO: 77) or [K/H/E]NAW (SEQ ID NO: 78) and the carbohydrase is an amylase.

Paragraph 13. The cleaning composition according to paragraph 1 or 10 to 12, wherein the DNase is selected from the group of polypeptides:

- a) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 39,
- b) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 40,
- c) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 41,
- d) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 42,
- e) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 43
- f) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 44,
- g) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 45,
- h) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 46,
- i) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 47,
- j) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 48,
- k) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 49,
- l) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 50,
- m) a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 51,

and wherein the carbohydrase is selected from the group consisting of;

- 5
- 10
- 15
- i. a cellulase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 86;
  - ii. a xylanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 87;
  - iii. a mannanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 82; and
  - iv. an amylase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 91.

Paragraph 14. The cleaning composition according to paragraph 1 wherein the DNase is selected from the group consisting of:

- a) polypeptide obtainable from *Bacillus licheniformis* having a sequence identity to the polypeptide shown in SEQ ID NO: 65 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity,
- b) polypeptide obtainable from *Bacillus subtilis* having a sequence identity to the polypeptide shown in SEQ ID NO: 66 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity,
- c) polypeptide obtainable from *Aspergillus oryzae* having a sequence identity to the polypeptide shown in SEQ ID NO: 67 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity,
- d) polypeptide obtainable from *Trichoderma harzianum* having a sequence identity to the polypeptide shown in SEQ ID NO: 68 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% and which have DNase activity,

and wherein the carbohydrase is selected from the group consisting of;

- i. a cellulase selected from a polypeptide having at least 80% sequence identity to

- the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 86;
- 5
- ii. a xylanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 87;
- iii. a mannanase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 82; and
- 10
- iv. an amylase selected from a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 80% sequence identity to the polypeptide shown in SEQ ID NO: 91.
- 15

Paragraph 15. The use of a composition according to any of the previous paragraphs for deep cleaning of an item, wherein the item is a textile or a surface.

20 Paragraph 16. A method of formulating a cleaning composition comprising adding a DNase, a carbohydrase and at least one cleaning component.

Paragraph 17. A kit intended for deep cleaning, wherein the kit comprises a solution of an enzyme mixture comprising a DNase, carbohydrase and optionally a protease.

25 Paragraph 18. A method of deep cleaning on an item, comprising the steps of:

a) contacting the item with a solution comprising an enzyme mixture comprising a DNase and a carbohydrase and optionally a protease; and a cleaning component, wherein the cleaning component is selected from 0.1 to 15 wt% of at least one a surfactant; 0.5 to 20 wt% of at least one builder; and 0.01 to 10 wt% of at least one bleach component; and

30 b) and optionally rinsing the item, wherein the item is preferably a textile.

## Definitions

Biofilm is produced by any group of microorganisms in which cells stick to each other or stick to a surface, such as a textile, dishware or hard surface or another kind of surface. These

35 adherent cells are frequently embedded within a self-produced matrix of extracellular polymeric substance (EPS). Biofilm EPS is a polymeric conglomeration generally composed of extracellular

DNA, proteins, and polysaccharides. Biofilms may form on living or non-living surfaces. The microbial cells growing in a biofilm are physiologically distinct from planktonic cells of the same organism, which, by contrast, are single-cells that may float or swim in a liquid medium. Bacteria living in a biofilm usually have significantly different properties from planktonic bacteria of the same species, as the dense and protected environment of the film allows them to cooperate and interact in various ways. One benefit of this environment for the microorganisms is increased resistance to detergents and antibiotics, as the dense extracellular matrix and the outer layer of cells protect the interior of the community.

On laundry biofilm producing bacteria can be found among the following species: *Acinetobacter sp.*, *Aeromicrobium sp.*, *Brevundimonas sp.*, *Microbacterium sp.*, *Micrococcus luteus*, *Pseudomonas sp.*, *Staphylococcus epidermidis*, and *Stenotrophomonas sp.* On hard surfaces biofilm producing bacteria can be found among the following species: *Acinetobacter sp.*, *Aeromicrobium sp.*, *Brevundimonas sp.*, *Microbacterium sp.*, *Micrococcus luteus*, *Pseudomonas sp.*, *Staphylococcus epidermidis*, *Staphylococcus aureus* and *Stenotrophomonas sp.* In one aspect, the biofilm producing strain is *Brevundimonas sp.* In one aspect, the biofilm producing strain is *Pseudomonas alcaliphila* or *Pseudomonas fluorescens*. In one aspect, the biofilm producing strain is *Staphylococcus aureus*.

By the term "deep cleaning" is meant reduction, disruption or removal of components of organic matter, e.g. biofilm, such as polysaccharides, proteins, DNA, soil or other components present in the organic matter.

Cleaning component: The cleaning component e.g. the detergent adjunct ingredient is different to the DNase and carbohydrase. The precise nature of these additional cleaning components e.g. adjunct components, and levels of incorporation thereof, will depend on the physical form of the composition and the nature of the operation for which it is to be used. Suitable cleaning components e.g. adjunct materials include, but are not limited to the components described below such as surfactants, builders, flocculating aid, chelating agents, dye transfer inhibitors, enzymes, enzyme stabilizers, enzyme inhibitors, catalytic materials, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, preformed peracids, polymeric agents, clay soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, perfumes, structure elasticizing agents, fabric softeners, carriers, hydrotropes, builders and co-builders, fabric huing agents, anti-foaming agents, dispersants, processing aids, and/or pigments.

Cleaning Composition: The term "cleaning composition" refers to compositions that find use in the removal of undesired compounds from items to be cleaned, such as textiles. The detergent composition may be used to e.g. clean textiles for both household cleaning and industrial cleaning. The terms encompass any materials/compounds selected for the particular type of cleaning composition desired and the form of the product (e.g., liquid, gel, powder, granulate, paste, or spray compositions) and includes, but is not limited to, detergent

compositions (e.g., liquid and/or solid laundry detergents and fine fabric detergents; fabric fresheners; fabric softeners; and textile and laundry pre-spotters/pretreatment). In addition to containing the enzyme blend of the invention, the detergent formulation may contain one or more additional enzymes (such as proteases, lipases, cutinases, endoglucanases, xyloglucanases, 5 pectinases, pectin lyases, xanthanases, peroxidases, haloperoxygenases and catalases or any mixture thereof), and/or detergent adjunct ingredients such as surfactants, builders, chelators or chelating agents, bleach system or bleach components, polymers, fabric conditioners, foam boosters, suds suppressors, dyes, perfume, tannish inhibitors, optical brighteners, bactericides, fungicides, soil suspending agents, anti-corrosion agents, enzyme inhibitors or stabilizers, 10 enzyme activators, transferase(s), hydrolytic enzymes, oxido reductases, bluing agents and fluorescent dyes, antioxidants, and solubilizers.

The term "enzyme detergency benefit" is defined herein as the advantageous effect an enzyme may add to a detergent compared to the same detergent without the enzyme. Important detergency benefits which can be provided by enzymes are stain removal with no or very little 15 visible soils after washing and/or cleaning, prevention or reduction of redeposition of soils released in the washing process (an effect that also is termed anti-redeposition), restoring fully or partly the whiteness of textiles which originally were white but after repeated use and wash have obtained a greyish or yellowish appearance (an effect that also is termed whitening). Textile care benefits, which are not directly related to catalytic stain removal or prevention of redeposition 20 of soils, are also important for enzyme detergency benefits. Examples of such textile care benefits are prevention or reduction of dye transfer from one fabric to another fabric or another part of the same fabric (an effect that is also termed dye transfer inhibition or anti-backstaining), removal of protruding or broken fibers from a fabric surface to decrease pilling tendencies or remove already existing pills or fuzz (an effect that also is termed anti-pilling), improvement of the fabric-softness, 25 colour clarification of the fabric and removal of particulate soils which are trapped in the fibers of the fabric or garment. Enzymatic bleaching is a further enzyme detergency benefit where the catalytic activity generally is used to catalyze the formation of bleaching components such as hydrogen peroxide or other peroxides. Textile care benefits, which are not directly related to catalytic stain removal or prevention of redeposition of soils, are also important for enzyme 30 detergency benefits. Examples of such textile care benefits are prevention or reduction of dye transfer from one textile to another textile or another part of the same textile (an effect that is also termed dye transfer inhibition or anti-backstaining), removal of protruding or broken fibers from a textile surface to decrease pilling tendencies or remove already existing pills or fuzz (an effect that also is termed anti-pilling), improvement of the textile-softness, colour clarification of the 35 textile and removal of particulate soils which are trapped in the fibers of the textile. Enzymatic bleaching is a further enzyme detergency benefit where the catalytic activity generally is used to

catalyze the formation of bleaching component such as hydrogen peroxide or other peroxides or other bleaching species."

The term "hard surface cleaning" is defined herein as cleaning of hard surfaces wherein hard surfaces may include floors, tables, walls, roofs etc. as well as surfaces of hard objects such as cars (car wash) and dishes (dish wash). Dish washing includes but are not limited to cleaning of plates, cups, glasses, bowls, cutlery such as spoons, knives, forks, serving utensils, ceramics, plastics, metals, china, glass and acrylics.

The term "wash performance" is used as an enzyme's ability to remove stains present on the object to be cleaned during e.g. wash or hard surface cleaning.

The term "whiteness" is defined herein as a greying, yellowing of a textile. Loss of whiteness may be due to removal of optical brighteners/hueing agents. Greying and yellowing can be due to soil redeposition, body soils, colouring from e.g. iron and copper ions or dye transfer. Whiteness might include one or several issues from the list below: colourant or dye effects; incomplete stain removal (e.g. body soils, sebum etc.); redeposition (greying, yellowing or other discolourations of the object) (removed soils reassociate with other parts of textile, soiled or unsoiled); chemical changes in textile during application; and clarification or brightening of colours.

The term "laundering" relates to both household laundering and industrial laundering and means the process of treating textiles with a solution containing a cleaning or detergent composition of the present invention. The laundering process can for example be carried out using e.g. a household or an industrial washing machine or can be carried out by hand.

By the term "malodor" is meant an odor which is not desired on clean items. The cleaned item should smell fresh and clean without malodors adhered to the item. One example of malodor is compounds with an unpleasant smell, which may be produced by microorganisms. Another example is unpleasant smells can be sweat or body odor adhered to an item which has been in contact with human or animal. Another example of malodor can be the odor from spices, which sticks to items for example curry or other exotic spices which smells strongly.

The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc.

The term "textile" means any textile material including yarns, yarn intermediates, fibers, non-woven materials, natural materials, synthetic materials, and any other textile material, fabrics made of these materials and products made from fabrics (e.g., garments and other articles). The textile or fabric may be in the form of knits, wovens, denims, non-wovens, felts, yarns, and towelling. The textile may be cellulose based such as natural cellulose, including cotton, flax/linen, jute, ramie, sisal or coir or manmade cellulose (e.g. originating from wood pulp) including viscose/rayon, cellulose acetate fibers (tricell), lyocell or blends thereof. The textile or

fabric may also be non-cellulose based such as natural polyamides including wool, camel, cashmere, mohair, rabbit and silk or synthetic polymers such as nylon, aramid, polyester, acrylic, polypropylene and spandex/elastane, or blends thereof as well as blends of cellulose based and non-cellulose based fibers. Examples of blends are blends of cotton and/or rayon/viscose with one or more companion material such as wool, synthetic fiber (e.g. polyamide fiber, acrylic fiber, polyester fiber, polyvinyl chloride fiber, polyurethane fiber, polyurea fiber, aramid fiber), and/or cellulose-containing fiber (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fiber, lyocell). Fabric may be conventional washable laundry, for example stained household laundry. When the term fabric or garment is used it is intended to include the broader term textiles as well.

The term "variant" means a polypeptide having the activity of the parent or precursor polypeptide and comprising an alteration, i.e., a substitution, insertion, and/or deletion, at one or more (e.g., several) positions compared to the precursor or parent polypeptide. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position.

Sequence identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity". For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 6.6.0 or later. The parameters used are a gap open penalty of 10, a gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

## Examples

### Assays

#### **Assay 1: testing of DNase activity**

DNase activity was determined on DNase Test Agar with Methyl Green (BD, Franklin Lakes, NJ, USA), which was prepared according to the manual from supplier. Briefly, 21 g of agar was dissolved in 500 ml water and then autoclaved for 15 min at 121 °C. Autoclaved agar was tempered to 48°C in water bath, and 20 ml of agar was poured into petridishes with and allowed to solidify by incubation o/n at room temperature. On solidified agar plates, 5 μl of enzyme solutions are added and DNase activity is observed as colorless zones around the spotted enzyme solutions.

**Assay II: testing of mannanase activity**

Mannanase activity may be tested according to standard test procedures known in the art, such as by applying a solution to be tested to 4 mm diameter holes punched out in agar plates containing 0.2% AZCL galactomannan (carob), *i.e.* substrate for the assay of endo-1,4-beta-D-mannanase available as CatNo.I-AZGMA from the company Megazyme (Megazyme's Internet address: <http://www.megazyme.com/Purchase/index.html>).

**Assay III: testing of xyloglucanase activity**

The reaction involves endo hydrolysis of 1,4-beta-D-glucosidic linkages in xyloglucan. For purposes of the present invention, xyloglucanase activity is determined using AZCL-xyloglucan (from Megazyme) as the reaction substrate. The assay can be performed in several ways, *e.g.* as described in Example 2 of the present application or as described in WO 01/62903. One unit of xyloglucanase activity (XyloU) is defined by reference to the assay method described in WO 01/62903, page 60, lines 3 - 17.

**Assay IV: testing of cellulase activity**

The term "cellulase activity" is defined herein as an enzyme catalyzed hydrolysis of 1,4-beta-D-glucosidic linkages in beta-1,4-glucan (cellulose). For purposes of the present invention, cellulase activity is determined using AZCL- HE-cellulose (from Megazyme) as the reaction substrate.

**Example 1**Isolating laundry specific bacterial strains

One strain of *Brevundimonas* sp. isolated from laundry was used in the present example. The *Brevundimonas* sp. was isolated during a study, where the bacterial diversity in laundry after washing at 15, 40 and 60°C, respectively, was investigated. The study was conducted on laundry collected from Danish households. For each wash, 20 g of laundry items (tea towel, towel, dish cloth, bib, T-shirt armpit, T-shirt collar, socks) in the range 4:3:2:2:1 :1:1 was used. Washing was performed in a Laundr-O-Meter (LOM) at 15, 40 or 60°C. For washing at 15 and 40°C, Ariel Sensitive White & Color was used, whereas WFK IEC-A\* model detergent was used for washing at 60°C. Ariel Sensitive White & Color was prepared by weighing out 5.1 g and adding tap water up to 1000 ml followed by stirring for 5 minutes. WFK IEC-A model detergent (which is available from WFK Testgewebe GmbH) was prepared by weighing out 5 g and adding tap water up to 1300 ml followed by stirring for 15 min. Washing was performed for 1 hour at 15, 40 and 60°C, respectively, followed by 2 times rinsing with tap water for 20 min at 15°C. Laundry was sampled immediately after washing at 15, 40 and 60°C, respectively. Twenty grams of laundry was added 0.9% (w/v) NaCl (1.06404; Merck, Darmstadt, Germany) with 0.5% (w/w) tween 80 to yield a 1:10 dilution in stomacher bag. The mixture was homogenized using a Stomacher for 2 minutes at medium speed. After homogenization, ten-fold dilutions were prepared in 0.9% (w/v) NaCl. Bacteria were enumerated on Tryptone Soya Agar (TSA) (CM0129, Oxoid, Basingstoke,



Hampshire, UK) incubated aerobically at 30°C for 5-7 days. To suppress growth of yeast and moulds, 0.2% sorbic acid (359769, Sigma) and 0.1% cycloheximide (18079; Sigma) were added. Bacterial colonies were selected from countable plates and purified by restreaking twice on TSA. For long time storage, purified isolates were stored at -80°C in TSB containing 20% (w/v) glycerol (49779; Sigma).

#### Preparation of swatches with biofilm

Swatches with biofilm of *Brevundimonas* sp. was included in the present study. Bacteria was pre-grown on Tryptone Soya Agar (TSA) (pH 7.3) (CM0131 ; Oxoid Ltd, Basingstoke, UK) for 2-5 days at 30°C. From a single colony, a loop-full was transferred to 10 mL of TSB and incubated for 1 day at 30°C with shaking (240 rpm). After propagation, cells were pelleted by centrifugation (Sigma Laboratory Centrifuge 6K1 5) (3000 g at 21°C in 7 min) and resuspended in 10 mL of TSB diluted twice with water. Optical density (OD) at 600 nm was measured using a spectrophotometer (POLARstar Omega (BMG Labtech, Ortenberg, Germany). Fresh TSB diluted twice with water was inoculated to an OD600nm of 0.03, and 50 mL was added into a petridish (diameter 125 mm), in which a swatch (80 mm x 120 mm) of sterile cotton (WFK10A). After incubation (48 h at 15°C with shaking (100 rpm), swatches were rinsed twice with 0.9% (w/v) NaCl and dried in LAF bench for 60 min. Swatches were stored at 4°C prior to wash.

### **Example 2**

#### Wash experiment

Wash experiment was performed using the Automatic Mechanical Stress Assay (AMSA). With AMSA, the wash performance of many small volume enzyme-detergent solutions can be examined at the same time. The AMSA plate has many slots for test solutions, and a lid that firmly squeezes the textile to be washed against the slot openings. During the wash, the plate, test solutions, textile and lid are vigorously shaken to bring the test solution in contact with the textile and apply mechanical stress in a regular, periodic, oscillating manner.

The wash experiment was conducted under the experimental conditions specified below:

Detergent dosage	3.3 g/L (liquid detergent)
Test solution volume	160 micro L
pH	pH 8
Wash time	20 minutes
Temperature	30°C
Water hardness	15°dH
Soil	Wfk09V 0.7g/L

Model detergents and test materials were as follows:

Laundry liquid model detergent	Model detergent A
--------------------------------	-------------------

Test material	Brevundimonas sp. 2-day biofilm grown on WFK10 (cotton) or WFK30A (polyester)
---------------	---

For wash experiments, Model detergent A (containing 12% LAS, 11% AEO Biosoft N25-7 (NI), 7% AEOS (SLES), 6% MPG, 3% ethanol, 3% TEA, 2.75% cocoa soap, 2.75% soya soap, 2% glycerol, 2% sodium hydroxide, 2% sodium citrate, 1% sodium formiate, 0.2% DTMPA and 0.2% PCA (all percentages are w/w)) (3.3 g/L) dissolved in water hardness 15°dH (Ca:Mg:NaHC03- = 4:1 :1.5) was used. Soil was subsequently added to reach a concentration of 0.7g soil/L (WFK09V pigment soil) to reveal biofilm. After washing, textiles were flushed in tap water and dried over night before scanning. Wash experiments were done twice.

Wash performance was measured as the brightness of the WFK09V pigment soiled, washed textile. Brightness can also be expressed as the intensity of the light reflected from the sample when illuminated with white light. When the sample is soiled, the intensity of the reflected light is lower, than that of a clean sample. Therefore, the intensity of the reflected light can be used to measure wash performance. Intensity measurements were made with a professional flatbed scanner (Kodak iQsmart, Kodak, Midtager 29, DK-2605 Br0ndby, Denmark), which was used to capture an image of the washed and dried textile. To extract a value for the light intensity from the scanned images, 24-bit pixel values from the image were converted into values for red, green and blue (RGB). The intensity value (Int) was calculated by adding the RGB values together as vectors and then taking the length of the resulting vector:

$$Int = \sqrt{r^2 + g^2 + b^2}$$

**Example 3: Wash synergy between DNase and cellulase**

To asses wash synergy between DNase (SEQ ID NO 13) and cellulase (SEQ ID NO 85) biofilm-harboring textile was AMSA washed a) in the absence of enzyme (blank), b) in the presence of DNase alone, c) in the presence of cellulase alone and d) with a mixture of DNase and cellulase. The resulting textile intensities and corresponding wash performances (WPs) are listed in Table 1 and 2. Wash performances attributable to DNase (WPDNase), cellulase (WPceiiu) and the mixture of the two (WPDNase+ceiiu) were quantified as the difference in intensity between textile washed With and WithOUt enzyme:  $WPDNase = i_{DNase} - I_{Blank}$ ,  $WPcellu = I_{cellu} - i_{Blank}$ ,  $WPDNase+Cellu = i_{DNase+Cellu} - I_{Blank}$ . The synergistic component of wash performance  $W P_{syn}$  was quantified as the extent to which wash performance of mixed DNase and cellulase (WPDNase+ceiiu) exceeded the sum of the individual wash performances of DNase alone and cellulase alone:  $W P_{syn} = WPDNase+ceiiu - (WPDNase + WPcellu)$ .

Table 1. synergistic wash effect of DNase and cellulase (wash experiment 1)

	Cellulase (SEQ ID NO 85)	I	WP	WP <sub>syn</sub>
--	--------------------------	---	----	-------------------

Blank	No enzyme	285.460	-	-
DNase	0.00002 ppm DNase	298.681	13.22	-
Cellulase	0.015% Cellulase	289.347	3.89	-
	0.15% Cellulase	290.955	5.49	-
DNase+ Cellulase	0.00002 ppm DNase + 0.015% Cellulase	304.770	19.31	2.20
	0.00002 ppm DNase + 0.15% Cellulase	310.821	25.36	6.65

Table 2. synergistic wash effect of DNase and cellulase (wash experiment 2)

	Cellulose (SEQ ID NO 91)	I	WP	WP <sub>syn</sub>
Blank	No enzyme	286.616	-	-
DNase	0.00002 ppm DNase	322.453	35.84	-
Cellulase	0.015% Cellulase	287.859	1.24	-
	0.15% Cellulase	291.329	4.71	-
DNase+ Cellulase	0.00002 ppm DNase + 0.015% Cellulase	326.890	40.27	3.19
	0.00002 ppm DNase + 0.15% Cellulase	332.886	46.27	5.72

**Claims****What is claimed is:**

1. A cleaning composition comprising a DNase, at least one carbohydrase and a cleaning component, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase.
2. A cleaning composition according to claim 1 wherein the carbohydrase is a cellulase, preferably selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 83, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 84, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 85, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 86.
3. A cleaning composition according to claim 1 wherein the carbohydrase is an amylase, preferably selected from a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 88, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 89, a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 90, and a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 91.
4. A cleaning composition according to claim 1 wherein the carbohydrase is a mannanase, preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at

least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 82.

5. A cleaning composition according to claim 1 wherein the carbohydrase is a xylanase, preferably a polypeptide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the polypeptide shown in SEQ ID NO: 87.
6. A cleaning composition according to any of the preceding claims, wherein the DNase is microbial, preferably obtained from bacteria or fungi.
7. A cleaning composition according to any of the preceding claims, wherein the DNase is obtained from *Bacillus*, preferably *Bacillus cibi*, *Bacillus horikoshii*, *Bacillus licheniformis*, *Bacillus subtilis*, *Bacillus horneckiae*, *Bacillus idriensis*, *Bacillus algicola*, *Bacillus vietnamensis*, *Bacillus hwajinpoensis*, *Bacillus indicus*, *Bacillus marisflavi* or *Bacillus luciferensis*.
8. A cleaning composition of claim 7, wherein the DNase comprises one or both of the motif(s) [D/M/L][S/T]GYSR[D/N] (SEQ ID NO: 73) or ASXNRSKG (SEQ ID NO: 74).
9. A cleaning composition according to any of claims 6 to 8, wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO 13.
10. A cleaning composition according to any of claims 6 to 8, wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO: 65.
11. A cleaning composition according to any of claims 6 to 8, wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO: 66.
12. A cleaning composition according to claim 6, wherein the DNase is fungal, preferably obtained from *Aspergillus* and even more preferably from *Aspergillus oryzae* and wherein

the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO: 67.

13. A cleaning composition according to claim 6, wherein the DNase is fungal, preferably obtained from *Trichoderma* and even more preferably from *Trichoderma harzianum* and wherein the DNase has at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% sequence identity to the amino acid sequence shown in SEQ ID NO: 68.
14. A cleaning composition according to any of the preceding claims wherein the amount of DNase in the composition is from 0.01 to 1000 ppm and the amount of carbohydrase is from 0.01 to 1000 ppm.
15. A cleaning composition according to any of the preceding claims, wherein the cleaning component is selected from surfactants, preferably anionic and/or nonionic, builders and bleach components.
16. Use of a cleaning composition according to any of claims 1 to 15 for deep cleaning of an item, wherein the item is a textile or a surface.
17. A method of formulating a cleaning composition according to any of claims 1 to 15 comprising adding a DNase, a carbohydrase and at least one cleaning component.
18. A kit intended for deep cleaning, wherein the kit comprises a solution of an enzyme mixture comprising a DNase and a carbohydrase, wherein the carbohydrase is a cellulase, an amylase, a mannanase or a xylanase.
19. A method of deep cleaning of an item, comprising the steps of:
  - a) contacting the item with a cleaning composition according to any of claims 1 to 15;
  - b) and optionally rinsing the item, wherein the item is preferably a textile.

5

10

**INTERNATIONAL SEARCH REPORT**

International application No  
PCT/EP2018/056730

A. CLASSIFICATION OF SUBJECT MATTER  
**INV. C11D3/386 C12N9/22 C12N9/24 C12N9/26 C12N9/42**  
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
**C1D C12N**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
**EPO-Internal , WPI Data**

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2017/001472 AI (NOVOZYMES AS [DK] ) 5 January 2017 (2017-01-05) abstract, page 7, line 25 - page 10, line 14; pages 15-21 and 23-26; claims 1-14 -----	1,2,6-19
X	wo 2017/001471 AI (NOVOZYMES AS [DK] ) 5 January 2017 (2017-01-05) Entire document, in particular abstract; page 7, line 27 - page 10, line 24; pages 16-22 and 25-27; claims 1-15 -----	1,2,6-19
X	wo 2016/162558 AI (NOVOZYMES AS [DK] ) 13 October 2016 (2016-10-13) abstract; page 9, line 15 - page 10, line 24; pages 13-27 and 30-32; pages 46-47; page 48, line 26 - page 51, line 15; claims 1-20 -----	1,2,6-19
	-/- .	

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search <b>17 May 2018</b>	Date of mailing of the international search report <b>16/08/2018</b>
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  <b>van Kl ompenburg, Wim</b>

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2018/056730

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2016/162556 A1 (NOVOZYMES AS [DK]) 13 October 2016 (2016-10-13) abstract; page 10, line 9 - page 12, line 8; pages 15-20 and 24-26; claims 1-17 -----	1,2,6-19
X	WO 2015/181287 A1 (NOVOZYMES AS [DK]) 3 December 2015 (2015-12-03) abstract; pages 12-13; pages 19-20; pages 36-42 and 44-46; claims 1-15 -----	1,2,6-19
X	WO 2015/181286 A1 (NOVOZYMES AS [DK]) 3 December 2015 (2015-12-03) abstract; page 2, lines 13-23; pages 12-16, 21-22, 58-63 and 66-68; claims 1-15 -----	1,2,6-19
X	WO 2015/166075 A1 (NOVOZYMES AS [DK]) 5 November 2015 (2015-11-05) abstract, pages 12-13, 17-18, 32-48, 50 and 52-55; claims 1-15 -----	1,2,6-19
X	WO 2015/155351 A1 (NOVOZYMES AS [DK]) 15 October 2015 (2015-10-15) abstract; page 2, lines 8-21; pages 11-13, 50-57 and 59-61; claims 1-20 -----	1,2,6-19
X	WO 2015/155350 A1 (NOVOZYMES AS [DK]) 15 October 2015 (2015-10-15) abstract; page 2, lines 9-22; pages 12-15, 53-57, 59 and 62-64; claims 1-20 -----	1,2,6-19
X	WO 2014/087011 A1 (NOVOZYMES AS [DK]) 12 June 2014 (2014-06-12) abstract; pages 1-2, 12-15, 17 and 19-22; claims 1-15 -----	1,2,6-19
A	WO 2014/124927 A2 (NOVOZYMES AS [DK]) 21 August 2014 (2014-08-21) the whole document -----	2



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2018/056730

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
  
2.  As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
  
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos. :
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos. :  
  
1, 2, 6-19 (al l parti al ly)

### Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1, 2, 6-19 (all partially)

Cleaning composition comprising DNase polypeptide and a carbohydrase and a cleaning component. Methods, uses and kits relating thereto. Cleaning composition comprising a cleaning component, a DNase polypeptide and a carbohydrase wherein the carbohydrase has at least 60% identity to SEQ ID NO 83. Methods, uses and kits relating thereto.

---

2-15 . claims: 1-19 (partially)

Cleaning composition etc. as in non unity subject 1, but comprising carbohydrase or DNase polypeptides exhibiting at least 80% sequence identity to SEQ.ID.NO: 84, 85, 86, 88, 89, 90, 91, 82, 87, 13, 65, 66, 67, 68 respectively

---

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/056730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2017001472 A1	05-01-2017	CN 107922895 A	17-04-2018
		EP 3313971 A1	02-05-2018
		US 2018171270 A1	21-06-2018
		WO 2017001472 A1	05-01-2017
-----			
WO 2017001471 A1	05-01-2017	CN 107969135 A	27-04-2018
		EP 3313970 A1	02-05-2018
		WO 2017001471 A1	05-01-2017
-----			
WO 2016162558 A1	13-10-2016	CN 107636134 A	26-01-2018
		EP 3280800 A1	14-02-2018
		US 2018105772 A1	19-04-2018
		WO 2016162558 A1	13-10-2016
-----			
WO 2016162556 A1	13-10-2016	CN 107567489 A	09-01-2018
		EP 3280791 A1	14-02-2018
		US 2018112156 A1	26-04-2018
		WO 2016162556 A1	13-10-2016
-----			
WO 2015181287 A1	03-12-2015	CN 106795463 A	31-05-2017
		EP 3149144 A1	05-04-2017
		US 2017081617 A1	23-03-2017
		WO 2015181287 A1	03-12-2015
-----			
WO 2015181286 A1	03-12-2015	BR 112016027289 A2	31-10-2017
		CA 2947243 A1	03-12-2015
		CN 106459839 A	22-02-2017
		EP 3149143 A1	05-04-2017
		JP 2017517602 A	29-06-2017
		US 2017081616 A1	23-03-2017
		WO 2015181286 A1	03-12-2015
-----			
WO 2015166075 A1	05-11-2015	CN 106232795 A	14-12-2016
		EP 3137587 A1	08-03-2017
		US 2017044468 A1	16-02-2017
		WO 2015166075 A1	05-11-2015
-----			
WO 2015155351 A1	15-10-2015	CN 106164237 A	23-11-2016
		EP 3129458 A1	15-02-2017
		US 2017152462 A1	01-06-2017
		WO 2015155351 A1	15-10-2015
-----			
WO 2015155350 A1	15-10-2015	BR 112016023188 A2	16-01-2018
		CA 2943029 A1	15-10-2015
		CN 106164236 A	23-11-2016
		EP 3129457 A1	15-02-2017
		JP 2017512885 A	25-05-2017
		US 2017107457 A1	20-04-2017
		WO 2015155350 A1	15-10-2015
-----			
WO 2014087011 A1	12-06-2014	BR 112015012982 A2	12-09-2017
		CA 2893454 A1	12-06-2014
		CN 104837979 A	12-08-2015
		EP 2929004 A1	14-10-2015
		JP 2016507597 A	10-03-2016
		US 2015299623 A1	22-10-2015
		WO 2014087011 A1	12-06-2014

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/056730

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014124927 A2	21-08-2014	CN 105073971 A	18-11-2015
		EP 2956535 A2	23-12-2015
		JP 2016516452 A	09-06-2016
		US 2015376554 A1	31-12-2015
		WO 2014124927 A2	21-08-2014
-----			