

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2018/023070 A1

(43) International Publication Date
01 February 2018 (01.02.2018)

(51) International Patent Classification:

A61K 45/06 (2006.01) *A61K 31/55* (2006.01)
A61P 43/00 (2006.01)

GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2017/044511

(22) International Filing Date:

28 July 2017 (28.07.2017)

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/368,575 29 July 2016 (29.07.2016) US

(71) Applicant: SUNOVION PHARMACEUTICALS, INC.

[US/US]; 84 Waterford Drive, Marlborough, MA 01752 (US).


(72) Inventors: ALEXANDROV, Vadim; 765 Old Saw Mill River Road, Tarrytown, NY 10591 (US). BOWEN, Carrie, A.; 17 Queen of Roses Lane, Uxbridge, MA 01569 (US). HANANIA, Taleen, G.; 44 Leroy Avenue, Apt. 2, Valhalla, NY 10595 (US). POWELL, Noel, Aaron; 28 Carlisle Road, Westford, MA 01886 (US).

(74) Agent: MEGERDITCHIAN, Samuel, H.; Gibbons P.C., One Gateway Center, Newark, NJ 07102 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

(54) Title: COMPOUNDS AND COMPOSITIONS AND USES THEREOF



(57) Abstract: Compounds of formula (I) are disclosed, as are pharmaceutical compositions containing such compounds. Methods of treating neurological or psychiatric diseases and disorders in a subject in need thereof are also disclosed.

COMPOUNDS AND COMPOSITIONS AND USES THEREOF

FIELD OF THE INVENTION

[0001] The invention relates to isochroman-4-amines, benzooxepinamines and benzooxacinamines as well as pharmaceutical compositions containing such compounds. The compounds and pharmaceutical compositions are useful for treating neurological or psychiatric diseases and disorders in a subject in need thereof. Such diseases and disorders include, for example, depression, bipolar disorder, pain, schizophrenia, obsessive compulsive disorder, addiction, social disorder, attention deficit hyperactivity disorder, an anxiety disorder, autism, a cognitive impairment, or a neuropsychiatric symptom such as apathy, depression, anxiety, psychosis, aggression, agitation, impulse control disorders, and sleep disorders in neurological disorders such as Alzheimer's and Parkinson's diseases.

[0002] All publications, patents, patent applications, and other references cited in this application are incorporated herein by reference in their entirety for all purposes and to the same extent as if each individual publication, patent, patent application or other reference was specifically and individually indicated to be incorporated by reference in its entirety for all purposes. Citation of a reference herein shall not be construed as an admission that such is prior art to the present invention.

BACKGROUND OF THE INVENTION


[0003] Central nervous system diseases and disorders affect a wide range of the population with differing severity. Neurological and psychiatric diseases and disorders include major depression, schizophrenia, bipolar disorder, obsessive compulsive disorder (OCD), panic disorder, and posttraumatic stress disorder (PTSD), among others. These diseases and disorders affect a person's thoughts, mood, behavior and social interactions and can significantly impair daily functioning. *See, e.g., Diagnostic and Statistical Manual of Mental Disorders*, 4th Ed., American Psychiatric Association (2000) ("DSM-IV-TR"); *Diagnostic and Statistical Manual of Mental Disorders*, 5th Ed., American Psychiatric Association (2013) ("DSM-5").

[0004] While medications exist for some aspects of these diseases and disorders, there remains a need for effective treatments for various neurological and psychiatric diseases and

disorders, including mood disorders such as bipolar and related disorders, psychosis and schizophrenia. For example, while mood stabilizers such as lithium and valproate, antidepressants and antipsychotic drugs are used to treat mood disorders, more effective medications are necessary. And current antipsychotics may be successful in treating the positive symptoms of schizophrenia but fare less well for the negative and cognitive symptoms. Additionally, current antidepressants are typically effective only for a proportion of subjects suffering from depression. Furthermore, despite the fact that the behavioral and psychiatric symptoms of neurological disease such as Parkinson's disease and Alzheimer's disease are major reasons for the institutionalization of subjects, few drugs exist to treat them.

SUMMARY OF THE INVENTION

[0005] In one aspect, the invention relates to method for treating a neurological or psychiatric disease or disorder comprising administering to a subject in need thereof an effective amount of a compound of formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

q1 is 0 or 1,

q2 is 0 or 1, and

the sum of q1 plus q2 is 0 or 1;

R¹, R² and R³ are chosen independently from H and an aliphatic (C₁-C₈)hydrocarbyl optionally substituted with one or more groups independently selected from halogen, -C≡C-, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino and di(C₁-C₆)alkylamino;

R^{4a}, R^{4b}, R^{7a}, and R^{7b} are chosen independently from H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, and (C₁-C₆)alkoxy;

R^{5a} , R^{5b} , R^{6a} and R^{6b} are chosen independently from H, halogen, (C_1 - C_6)alkyl and (C_1 - C_6)haloalkyl;

R^8 , R^9 , R^{10} and R^{11} are chosen independently from H, halogen, (C_1 - C_8)hydrocarbyl, cyano, - CF_3 , (C_1 - C_6)haloalkyl, hydroxyl, (C_1 - C_6)alkoxy, aminocarbonyl, (C_1 - C_6)alkylaminocarbonyl, di(C_1 - C_6)alkylaminocarbonyl, (C_1 - C_6)acyl, (C_1 - C_6)haloalkoxy, hydroxy(C_1 - C_6)alkyl, carboxy, (C_1 - C_6)alkoxycarbonyl, acetoxy, nitro, amino, (C_1 - C_6)alkylamino, di(C_1 - C_6)alkylamino, aminosulfonyl, benzyl, aryl, heteroaryl, phenoxy, - $OC(O)N(alkyl)_2$ or benzyloxy;

wherein said benzyl, aryl, heteroaryl, phenoxy or benzyloxy are optionally substituted with one or more substituents independently selected from halogen, (C_1 - C_6)alkyl, cyano, (C_1 - C_6)haloalkyl, hydroxyl, (C_1 - C_6)alkoxy, aminocarbonyl, (C_1 - C_6)alkylaminocarbonyl, di(C_1 - C_6)alkylaminocarbonyl, (C_1 - C_6)acyl, (C_1 - C_6)haloalkoxy, hydroxy(C_1 - C_6)alkyl, carboxy, (C_1 - C_6)alkoxycarbonyl, acetoxy, nitro, amino, (C_1 - C_6)alkylamino, and di(C_1 - C_6)alkylamino;

and further wherein:

any two of R^8 , R^9 , R^{10} and R^{11} attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C_1 - C_6)alkylamino, di(C_1 - C_6)alkylamino, (C_1 - C_6)alkyl, (C_1 - C_6)haloalkyl, (C_1 - C_6)alkoxy, and (C_1 - C_6)haloalkoxy.

[0006] In another aspect the invention relates to pharmaceutical compositions comprising compounds of formula I above, wherein at least one of R^1 , R^2 , R^3 , R^{4a} , R^{4b} , R^{5a} , R^{5b} , R^{6a} , R^{6b} , R^{7a} , R^{7b} , R^8 , R^9 , R^{10} and R^{11} is other than hydrogen.

[0007] In another aspect, the invention relates to chemical compounds of formula I, with the provisos that:

when $q1$ and $q2$ are both zero and NR^2R^3 is $NHCH_3$ or $N(CH_3)_2$, then at least one of R^1 , R^{4a} , R^{4b} , R^8 , R^9 , R^{10} and R^{11} must be other than hydrogen or methoxy, and

when $q1$ and $q2$ are both zero and NR^2R^3 is NH_2 , then one of the following four conditions must be met:


at least two of R^8 , R^9 , R^{10} and R^{11} must be other than hydrogen, or

at least one of R^8 , R^9 , R^{10} and R^{11} must be phenyl, or

at least one of R^{4a} and R^{4b} , must be other than hydrogen, or

at least one of R^{7a} and R^{7b} must be methyl and the other of R^{7a} and R^{7b} must be other than methyl, and

when q1 or q2 is one and NR^2R^3 is NH_2 , then the following species are excluded:



BRIEF DESCRIPTION OF THE DRAWINGS

[0008] Figure 1 shows the effect of representative compounds of the invention on treatment on DA release in the PFC.

[0009] Figure 2 shows the effect the effect of representative compounds of the invention on treatment on 5-HT release in the PFC.

[0010] Figure 3 shows the effect of representative compounds of the invention on treatment on NE release in the PFC.

[0011] Figure 4 shows the effect of representative compounds of the invention on treatment on ACh release in the PFC

[0012] Figure 5 shows the effect of representative compounds of the invention on treatment on DA release in the NAcc.

[0013] Figure 6 shows the effect of representative compounds of the invention on treatment on 5-HT release in the NAcc.

[0014] Figure 7 shows the effect of representative compounds of the invention on treatment on NE release in the NAcc.

[0015] Figure 8 shows the effect of representative compounds of the invention on treatment on ACh release in the NAcc.

DETAILED DESCRIPTION OF THE INVENTION

[0016] The methods of the invention relate to the use of compounds and compositions of Formula I above to treat neurological or psychiatric diseases and disorders or impairments. In some embodiments, the neurological or psychiatric disease or disorder is depression, bipolar disorder, pain, schizophrenia, obsessive compulsive disorder, addiction, social

disorder, attention deficit hyperactivity disorder, an anxiety disorder, autism or cognitive impairments. In one embodiment, the disorder is depression, particularly treatment-resistant depression (TRD), major depressive disorder (MDD), unipolar depression, bipolar depression or depression associated with another disease or disorder.

[0017] In one embodiment relating to the use of compounds of formula I in these methods, any two of R⁸, R⁹, R¹⁰ and R¹¹ attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy. In one such embodiment, any two of R⁸, R⁹, R¹⁰ and R¹¹ form a phenyl ring, which may be substituted as described.

[0018] In another embodiment relating to the use of compounds of formula I in these methods, there are four possibilities: (a) at least one of R⁸, R⁹, and R¹¹ is chosen from halogen, methyl and ethyl; or (b) R¹⁰ is methoxy; or (c) R⁸, R⁹, R¹⁰, R¹¹, R^{7a}, and R^{7b} are hydrogen and at least one of R^{4a} and R^{4b} is methyl, or (d) R⁸ and R⁹, together with the carbon atoms to which they are attached, form a 6-membered carbocycle or 7-membered bridged carbocycle. In any of these four possibilities, when q1 and q2 are both zero, R^{4a} may be hydrogen and R^{4b} may be hydrogen or methyl, and R² and R³ may be hydrogen. In any of these four possibilities, when q1 is one, R² and R³ may be hydrogen or methyl.

[0019] In another embodiment relating to the use of compounds of formula I in these methods, exemplary compounds are those in which R² and R³ are hydrogen, q1 and q2 are both zero, and (a) both of R^{7a} and R^{7b} are methyl, or (b) both of R⁸ and R⁹ are methyl, or (c) R⁹ is methoxy and at least one of R^{4a} and R^{4b} is methyl, or (d) R¹¹ is propyl.


[0020] In another embodiment relating to the use of compounds of formula I in these methods, exemplary compounds are those in which R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, and R^{7b} are all hydrogen, and (a) q1 is one and R¹, R², R³, R⁸, R¹⁰, and R¹¹ are hydrogen, or (b) q1 and q2 are both zero, R¹⁰ and R¹¹ are hydrogen, or (c) q1 and q2 are both zero, R⁸, R⁹, and R¹⁰ are hydrogen and R¹¹ is -(C₁-C₃)alkyl.

[0021] In another embodiment relating to the use of compounds of formula I in these methods, exemplary compounds are those in which R¹, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, and R^{7b} are all hydrogen, and (a) q1 is one, R² and R³ are hydrogen or methyl, and R¹⁰ and R¹¹ are

hydrogen or methyl, or (b) q1 and q2 are both zero and R⁸, R⁹, R¹⁰ and R¹¹ are chosen from hydrogen, methyl, methoxy and chloro.

[0022] In a pharmaceutical composition aspect, the invention relates to compositions comprising a pharmaceutically acceptable carrier and a compound of formula I above, wherein at least one of R¹, R², R³, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, R^{7b}, R⁸, R⁹, R¹⁰ and R¹¹ is other than hydrogen.

[0023] In a chemical compound aspect, the invention relates to compounds of formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

q1 is 0 or 1,

q2 is 0 or 1, and

the sum of q1 plus q2 is 0 or 1;

R¹, R² and R³ are chosen independently from H and an aliphatic (C₁-C₈)hydrocarbyl optionally substituted with one or more groups independently selected from halogen, -C≡C-, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino and di(C₁-C₆)alkylamino;

R^{4a}, R^{4b}, R^{7a}, and R^{7b} are chosen independently from H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, and (C₁-C₆)alkoxy;

R^{5a}, R^{5b}, R^{6a} and R^{6b} are chosen independently from H, halogen, (C₁-C₆)alkyl and (C₁-C₆)haloalkyl;

R⁸, R⁹, R¹⁰ and R¹¹ are chosen independently from H, halogen, (C₁-C₈)hydrocarbyl, cyano, -CF₃, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy,

hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, aminosulfonyl, benzyl, aryl, heteroaryl, phenoxy, -OC(O)N(alkyl)₂ or benzyloxy;

wherein said benzyl, aryl, heteroaryl, phenoxy or benzyloxy are optionally substituted with one or more substituents independently selected from halogen, (C₁-C₆)alkyl, cyano, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy, hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, and di(C₁-C₆)alkylamino;

and further wherein:

any two of R⁸, R⁹, R¹⁰ and R¹¹ attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy;

with the provisos that,

when q1 and q2 are both zero and NR²R³ is NHCH₃ or N(CH₃)₂, then at least one of R¹, R^{4a}, R^{4b}, R⁸, R⁹, R¹⁰ and R¹¹ must be other than hydrogen or methoxy; or

when q1 and q2 are both zero and NR²R³ is NH₂, then one of the following four conditions must be met:

at least two of R⁸, R⁹, R¹⁰ and R¹¹ must be other than hydrogen, or

at least one of R⁸, R⁹, R¹⁰ and R¹¹ must be phenyl, or

at least one of R⁸, R⁹, R¹⁰ and R¹¹ must be phenyl, or


at least one of R^{4a} and R^{4b} must be other than hydrogen, or

at least one of R^{7a} and R^{7b} must be methyl and the other of R^{7a} and R^{7b} must be other than methyl, and

when q1 or q2 is one, then the following species are excluded:




[0024] In one embodiment relating to the compounds, q1 and q2 are both zero, and the compounds are chroman-1-amines of formula II:




II.

[0025] In another embodiment relating to the compounds, q1 is one, and the compounds are 1,3,4,5-tetrahydrobenzo[c]oxepin-5-amines of formula III:



III.

[0026] In another embodiment relating to the compounds, q2 is one, and the compounds are 1,2,4,5-tetrahydrobenzo[d]oxepin-1-amines of formula IV:



IV.

[0027] In some embodiments, R^{4a}, R^{4b}, R^{7a}, and R^{7b} are hydrogen.

[0028] In some embodiments, R⁷ and R⁸ form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy. In some embodiments, the carbocycle

or heterocycle may be chosen from phenyl, pyridine, cyclohexene, thiazole, bicycloheptene, and dihydropyran.

[0029] In some embodiments, R⁹ and R¹⁰ form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy. In some embodiments, the carbocycle or heterocycle may be chosen from phenyl, pyridine, cyclohexene, thiazole, bicycloheptene, and dihydropyran.

[0030] In some embodiments, R⁹ is a phenyl which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy; in others, R⁹ is a 5- or 6-membered heteroaryl which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy.

[0031] In some embodiments, two of R⁸, R⁹, R¹⁰ and R¹¹ are chosen from methyl and chloro and the other two are hydrogen.

[0032] In some embodiments, R² is hydrogen and R³ is chosen from methyl, ethyl, propyl, isopropyl, and hydroxyethyl.

[0033] In some embodiments, R¹ is hydrogen or methyl; in some embodiments R¹ is hydrogen.

[0034] In some embodiments, R^{4a} and R^{4b} are chosen independently from hydrogen and (C₁-C₆)alkyl.

[0035] In some embodiments of tetrahydrobenzo[c]oxepin-5-amines of formula III or 1,2,4,5-tetrahydrobenzo[d]oxepin-1-amines of formula IV, R⁸ and R⁹ form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy. In particular embodiments, the carbocycle is phenyl, which may be substituted as described.







[0036] In some embodiments of tetrahydrobenzo[c]oxepin-5-amines of formula III or 1,2,4,5-tetrahydrobenzo[d]oxepin-1-amines of formula IV, R^{4a} and R^{4b} are chosen from hydrogen and methyl.


[0037] In some embodiments of tetrahydrobenzo[c]oxepin-5-amines of formula III or 1,2,4,5-tetrahydrobenzo[d]oxepin-1-amines of formula IV, R² is hydrogen and R³ is hydrogen or methyl.


[0038] In some embodiments of chroman-1-amines of formula II, tetrahydrobenzo[c]oxepin-5-amines of formula III or 1,2,4,5-tetrahydrobenzo[d]oxepin-1-amines of formula IV, one or two of R⁸, R⁹, R¹⁰ and R¹¹ are chosen from methyl and chloro and the remaining two or three of R⁸, R⁹, R¹⁰ and R¹¹ are hydrogen.








[0039] In one embodiment, provided are compounds of formula (I) which are greater than 90% enantiomerically pure. In another embodiment, provided are compounds of formula I which are greater than 95% enantiomerically pure.


[0040] In one embodiment, provided is a compound according to formula I, wherein said compound is:





















[0041] In one embodiment, provided are compounds according to formula I wherein said compound is:



or a pharmaceutically acceptable salt thereof.

[0042] In another embodiment, provided are compounds according to formula I wherein said compound is:



or a pharmaceutically acceptable salt thereof.

[0043] In another embodiment, provided are compounds according to formula I wherein said compound is:




and has greater than 90% enantiomeric purity. In another embodiment, said compound has greater than 95% enantiomeric purity.

Compounds and compositions and Definitions:


[0044] Compounds and compositions of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. A comprehensive list of abbreviations utilized by organic chemists (i.e. persons of ordinary skill in the art) appears in the first issue of each volume of the Journal of Organic Chemistry. The definitions therein, which are typically presented in a table entitled “Standard List of Abbreviations” are the definitions used herein.

[0045] The graphic representations of racemic, ambiscalemic and scalemic or enantiomerically pure compounds used herein are a modified version of the denotations taken from Maehr J. Chem. Ed. 62, 114-120 (1985): simple lines provide no information about stereochemistry and convey only connectivity; solid and broken wedges are used to denote the absolute configuration of a chiral element; solid and broken bold lines are geometric descriptors indicating the relative configuration shown but not necessarily denoting racemic


character; and wedge outlines and dotted or broken lines denote enantiomerically pure compounds of the indicated relative stereochemistry of indeterminate absolute configuration. For example, the graphic representation




indicates a *trans* relationship between the two chiral centers, that is, either or both of the two representations below:



in any ratio, from pure enantiomers to racemates, while the representation:



indicates a single enantiomer with the absolute configuration depicted, e.g., *((R)-1-((R)-5-fluoroisochroman-1-yl)ethan-1-amine* in the illustration above. Further, the open wedge/broken line depiction as shown in the example below:



represents a single enantiomer of the indicated relative stereochemistry of an undetermined absolute configuration. In the text describing the stereochemistry of the examples, the convention of Chemical Abstracts is used. Thus “*(R)-1-((R)-5-rel-...*” indicates that the two

chiral centers are in that relative relationship, which would be depicted in a structural diagram by solid bold and dashed lines, whereas "(R)-1-((R)-5-...)" without the "rel" indicates a single enantiomer of that absolute configuration, which would be depicted in a structural diagram by solid and broken wedges.

[0046] The "enantiomeric excess" or "% enantiomeric excess" of a composition can be calculated using the equation shown below. In the example shown below, a composition contains 90% of one enantiomer, e.g., the S enantiomer, and 10% of the other enantiomer, e.g., the R enantiomer. $ee = (90-10)/100 = 80\%$.

[0047] Thus, a composition containing 90% of one enantiomer and 10% of the other enantiomer is said to have an enantiomeric excess of 80%. Some compositions described herein contain an enantiomeric excess of at least about 50%, 75%, 90%, 95%, or 99% of the S enantiomer. In other words, the compositions contain an enantiomeric excess of the S enantiomer over the R enantiomer. In other embodiments, some compositions described herein contain an enantiomeric excess of at least about 50%, 75%, 90%, 95%, or 99% of the R enantiomer. In other words, the compositions contain an enantiomeric excess of the R enantiomer over the S enantiomer.

[0048] For instance, an isomer/enantiomer can, in some embodiments, be provided substantially free of the corresponding enantiomer, and can also be referred to as "optically enriched," "enantiomerically enriched," "enantiomerically pure" and "non-racemic," as used interchangeably herein. These terms refer to compositions in which the percent by weight of one enantiomer is greater than the amount of that one enantiomer in a control mixture of the racemic composition (e.g., greater than 1:1 by weight). For example, an enantiomerically enriched preparation of the S enantiomer, means a preparation of the compound having greater than about 50% by weight of the S enantiomer relative to the R enantiomer, such as at least about 75% by weight, further such as at least about 80% by weight. In some embodiments, the enrichment can be much greater than about 80% by weight, providing a "substantially enantiomerically enriched," "substantially enantiomerically pure" or a "substantially non-racemic" preparation, which refers to preparations of compositions which have at least about 85% by weight of one enantiomer relative to other enantiomer, such as at least about 90% by weight, and further such as at least 95% by weight. In certain embodiments, the compound provided herein is made up of at least about 90% by weight of

one enantiomer. In other embodiments, the compound is made up of at least about 95%, 98%, or 99% by weight of one enantiomer.

[0049] In some embodiments, the compound is a racemic mixture of (S)- and (R)-isomers. In other embodiments, provided herein is a mixture of compounds wherein individual compounds of the mixture exist predominately in an (S)- or (R)-isomeric configuration. For example, the compound mixture has an (S)-enantiomeric excess of greater than about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.5%, or more. In other embodiments, the compound mixture has an (S)-enantiomeric excess of greater than about 55% to about 99.5%, greater than about 60% to about 99.5%, greater than about 65% to about 99.5%, greater than about 70% to about 99.5%, greater than about 75% to about 99.5%, greater than about 80% to about 99.5%, greater than about 85% to about 99.5%, greater than about 90% to about 99.5%, greater than about 95% to about 99.5%, greater than about 96% to about 99.5%, greater than about 97% to about 99.5%, greater than about 98% to greater than about 99.5%, greater than about 99% to about 99.5%, or more.

[0050] In other embodiments, the compound mixture has an (R)-enantiomeric purity of greater than about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.5% or more. In some other embodiments, the compound mixture has an (R)-enantiomeric excess of greater than about 55% to about 99.5%, greater than about 60% to about 99.5%, greater than about 65% to about 99.5%, greater than about 70% to about 99.5%, greater than about 75% to about 99.5%, greater than about 80% to about 99.5%, greater than about 85% to about 99.5%, greater than about 90% to about 99.5%, greater than about 95% to about 99.5%, greater than about 96% to about 99.5%, greater than about 97% to about 99.5%, greater than about 98% to greater than about 99.5%, greater than about 99% to about 99.5% or more.

[0051] In other embodiments, the compound mixture contains identical chemical entities except for their stereochemical orientations, namely (S)- or (R)-isomers. For example, if a compound disclosed herein has --CH(R)-- unit, and R is not hydrogen, then the --CH(R)-- is in an (S)- or (R)-stereochemical orientation for each of the identical chemical entities. In some embodiments, the mixture of identical chemical entities is a racemic mixture of (S)- and (R)-isomers. In another embodiment, the mixture of the identical chemical entities (except for their stereochemical orientations), contain predominately (S)-isomers or predominately (R)-

isomers. For example, the (S)-isomers in the mixture of identical chemical entities are present at about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.5%, or more, relative to the (R)-isomers. In some embodiments, the (S)-isomers in the mixture of identical chemical entities are present at an (S)-enantiomeric excess of greater than about 55% to about 99.5%, greater than about 60% to about 99.5%, greater than about 65% to about 99.5%, greater than about 70% to about 99.5%, greater than about 75% to about 99.5%, greater than about 80% to about 99.5%, greater than about 85% to about 99.5%, greater than about 90% to about 99.5%, greater than about 95% to about 99.5%, greater than about 96% to about 99.5%, greater than about 97% to about 99.5%, greater than about 98% to greater than about 99.5%, greater than about 99% to about 99.5% or more.

[0052] In another embodiment, the (R)-isomers in the mixture of identical chemical entities (except for their stereochemical orientations), are present at about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.5%, or more, relative to the (S)-isomers. In some embodiments, the (R)-isomers in the mixture of identical chemical entities (except for their stereochemical orientations), are present at a (R)-enantiomeric excess greater than about 55% to about 99.5%, greater than about 60% to about 99.5%, greater than about 65% to about 99.5%, greater than about 70% to about 99.5%, greater than about 75% to about 99.5%, greater than about 80% to about 99.5%, greater than about 85% to about 99.5%, greater than about 90% to about 99.5%, greater than about 95% to about 99.5%, greater than about 96% to about 99.5%, greater than about 97% to about 99.5%, greater than about 98% to greater than about 99.5%, greater than about 99% to about 99.5% or more.

[0053] Hydrocarbyl refers to any substituent comprised of hydrogen and carbon as the only elemental constituents. C₁ to C₂₀ hydrocarbon includes, for example, alkyl, cycloalkyl, polycycloalkyl, alkenyl, alkynyl, aryl and combinations thereof. Examples include benzyl, phenethyl, cyclohexylmethyl, adamantyl, camphoryl and naphthylethyl. Aromatic hydrocarbons include benzene (phenyl), naphthalene (naphthyl), anthracene, etc. Aliphatic hydrocarbons are hydrocarbons that are not aromatic; they may be saturated or unsaturated, cyclic, linear or branched, or combinations thereof. Aliphatic hydrocarbons include, for example, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, and combinations

thereof. Non-limiting examples of aliphatic hydrocarbons include isopropyl, 2-butenyl, 2-butynyl, cyclopentyl, cyclopropylmethyl, norbornyl, and the like.

[0054] Unless otherwise specified, alkyl (or alkylene) is intended to include linear or branched saturated hydrocarbon structures and combinations thereof. Alkyl refers to alkyl groups from 1 to 20 carbon atoms, preferably 1 to 10 carbon atoms, more preferably 1 to 6 carbon atoms. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, *n*-butyl, *s*-butyl, *t*-butyl and the like.

[0055] Cycloalkyl is a subset of hydrocarbon and includes cyclic hydrocarbon groups of from 3 to 8 carbon atoms. Examples of cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, norbornyl and the like.

[0056] Unless otherwise specified, the term “carbocycle” is intended to include ring systems in which the ring atoms are all carbon but of any oxidation state. Thus (C₃-C₁₀) carbocycle refers to both non-aromatic and aromatic systems, including such systems as cyclopropane, benzene (phenyl) and cyclohexene; (C₈-C₁₂) carbopolycycle refers to such systems as norbornane, decalin, indane and naphthalene. Carbocycle, if not otherwise limited, refers to monocycles, bicycles and polycycles, including bridged structures.

[0057] Heterocycle means an aliphatic or aromatic carbocycle residue in which from one to four carbons is replaced by a heteroatom selected from the group consisting of N, O, and S. The nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. Unless otherwise specified, a heterocycle may be non-aromatic (heteroaliphatic) or aromatic (heteroaryl). Heterocycle, if not otherwise limited, refers to monocycles, bicycles and polycycles, including bridged structures. Examples of heterocycles include, but are not limited to, pyrrolidine, pyrazole, pyrrole, indole, quinoline, isoquinoline, tetrahydroisoquinoline, benzofuran, benzodioxan, benzodioxole (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazole, morpholine, thiazole, pyridine, pyridazine, pyrimidine, thiophene, furan, oxazole, oxazoline, isoxazole, atropine, dioxane, tetrahydrofuran and the like. Examples of heterocyclyl residues include piperazinyl, piperidinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyrazinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, tetrahydrofuryl,

tetrahydropyranyl, thienyl (also historically called thiophenyl), benzothienyl, thiamorpholinyl, oxadiazolyl, triazolyl and tetrahydroquinolinyl.

[0058] Hydrocarbyloxy refers to groups of from 1 to 20 carbon atoms, preferably 1 to 10 carbon atoms, more preferably 1 to 6 carbon atoms attached to the parent structure through an oxygen. Alkoxy is a subset of hydrocarbyloxy and includes groups of a straight or branched configuration. Examples include methoxy, ethoxy, propoxy, isopropoxy and the like. Lower-alkoxy refers to groups containing one to four carbons.

[0059] The term "halogen" means fluorine, chlorine, bromine or iodine. In one embodiment, halogen may be fluorine or chlorine.

[0060] Unless otherwise specified, acyl refers to formyl and to groups of 1, 2, 3, 4, 5, 6, 7 and 8 carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. Examples include acetyl, benzoyl, propionyl, isobutyryl and the like. Lower-acyl refers to groups containing one to four carbons. The double bonded oxygen, when referred to as a substituent itself is called "oxo".

[0061] As used herein, the term "optionally substituted" may be used interchangeably with "unsubstituted or substituted". The term "substituted" refers to the replacement of one or more hydrogen atoms in a specified group with a specified radical. For example, substituted alkyl, aryl, cycloalkyl, heterocyclyl etc. refer to alkyl, aryl, cycloalkyl, or heterocyclyl wherein one or more H atoms in each residue are replaced with halogen, haloalkyl, hydrocarbyl, acyl, alkoxyalkyl, hydroxy lower alkyl, carbonyl, phenyl, heteroaryl, benzenesulfonyl, hydroxy, hydrocarbyloxy, haloalkoxy, oxaalkyl, carboxy, alkoxycarbonyl [-C(=O)O-alkyl], alkoxycarbonylamino [HNC(=O)O-alkyl], aminocarbonyl (also known as carboxamido) [-C(=O)NH₂], alkylaminocarbonyl [-C(=O)NH-alkyl], dialkylaminocarbonyl [-C(=O)N(alkyl)₂], cyano, acetoxy, nitro, amino, alkylamino, dialkylamino, (alkyl)(aryl)aminoalkyl, alkylaminoalkyl (including cycloalkylaminoalkyl), dialkylaminoalkyl, dialkylaminoalkoxy, heterocyclylalkoxy, mercapto, alkylthio, sulfoxide, sulfone, sulfonylamino, alkylsulfinyl, alkylsulfonyl, acylaminoalkyl, acylaminoalkoxy, acylamino, amidino, aryl, benzyl, heterocyclyl, heterocyclylalkyl, phenoxy, benzyloxy, heteroaryloxy, hydroxyimino, alkoxyimino, oxaalkyl, aminosulfonyl, trityl, amidino, guanidino, ureido, benzyloxyphenyl, and benzyloxy. "Oxo" is also included among the

substituents referred to in “optionally substituted”; it will be appreciated by persons of skill in the art that, because oxo is a divalent radical, there are circumstances in which it will not be appropriate as a substituent (e.g. on phenyl). In one embodiment, 1, 2, or 3 hydrogen atoms are replaced with a specified radical. In the case of alkyl and cycloalkyl, more than three hydrogen atoms can be replaced by fluorine; indeed, all available hydrogen atoms could be replaced by fluorine. In particular embodiments, substituents are halogen, halo(C₁-C₄)hydrocarbyl, halo(C₁-C₄)hydrocarbyloxy, cyano, thiocyanato, (C₁-C₄)hydrocarbylsulfinyl, (C₁-C₄)hydrocarbyl-sulfonyl, aminosulfonyl, nitro, acetyl, and acetamido. Preferred substituents are halogen, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)fluoroalkyl, (C₁-C₄)fluoroalkoxy, hydroxy, amino, (C₁-C₄)alkylamino, di(C₁-C₄)alkylamino, (C₁-C₄)acylamino, (C₁-C₄)fluoroalkyl and (C₁-C₄)fluoroalkoxy.

[0062] Substituents Rⁿ are generally defined when introduced and retain that definition throughout the specification and claims.

[0063] As used herein, and as would be understood by the person of skill in the art, the recitation of “a compound” - unless expressly further limited - is intended to include salts of that compound. Thus, for example, the recitation “a compound of formula I” as depicted above, which contains a basic amine residue -NR²R³, would include salts -NHR²R³⁺X⁻ wherein X⁻ is any counterion. In a particular embodiment, the term “compound of formula I” refers to the compound or a pharmaceutically acceptable salt thereof; this term refers to a pharmaceutically acceptable salt of the compound, even if not explicitly stated. Unless otherwise stated or depicted, structures depicted herein are also meant to include all stereoisomeric (e.g., enantiomeric, diastereomeric, and cis-trans isomeric) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and cis-trans isomeric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. In addition to therapeutic

uses, such compounds are useful, for example, as analytical tools or probes in biological assays.

[0064] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Although pharmaceutically acceptable counter ions will be preferred for preparing pharmaceutical formulations, other anions are quite acceptable as synthetic intermediates. Thus X may be pharmaceutically undesirable anions, such as iodide, oxalate, trifluoromethanesulfonate and the like, when such salts are chemical intermediates.

[0065] Unless otherwise specified, the word "includes" (or any variation thereon, e.g., "include", "including", etc.) is intended to be open-ended. For example, "A includes 1, 2 and 3" means that A includes but is not limited to 1, 2 and 3.

[0066] Unless otherwise specified, the phrase "such as" is intended to be open-ended. For example, "A can be a halogen, such as chlorine or bromine" means that A can be, but is not limited to, chlorine or bromine.

[0067] According to another embodiment, the invention provides a composition comprising a compound of this invention (or its pharmaceutically acceptable salt) and a pharmaceutically acceptable carrier, adjuvant, or vehicle. In some embodiments, the amount of compound in compositions of this invention (also referred to herein as "effective amount" or "therapeutically effective amount") is such that is effective to treat, prevent, and/or manage various neurological and/or psychiatric diseases and disorders and/or symptoms in a subject. In some embodiments, a composition of this invention is formulated for administration to a subject in need of such composition. In some embodiments, a composition of this invention is formulated for oral administration to a subject.

[0068] As used herein, the term "subject" to which administration is contemplated includes, but is not limited to, humans (i.e., a male or female of any age group, e.g., a pediatric subject (e.g., infant, child, adolescent) or adult subject (e.g., young adult, middle-aged adult or senior adult)) and/or other primates (e.g., cynomolgus monkeys, rhesus monkeys); mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, goats, cats, and/or dogs; and/or birds, including commercially relevant birds such as chickens, ducks, geese, quail, and/or turkeys.

[0069] In certain embodiments, provided herein is a composition (e.g., a pharmaceutical composition) comprising a compound described herein and a pharmaceutically acceptable excipient or carrier. In some embodiments, provided herein is a method of treating neurological or psychiatric diseases and disorders in a subject in need thereof in a subject, comprising administering an effective amount of a compound or a pharmaceutical composition described herein. Examples of carriers and excipients are well known to those skilled in the art and are described in detail in, e.g., Ansel, Howard C., et al., *Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems*. Philadelphia: Lippincott, Williams & Wilkins, 2004; Gennaro, Alfonso R., et al. *Remington: The Science and Practice of Pharmacy*. Philadelphia: Lippincott, Williams & Wilkins, 2000; and Rowe, Raymond C. *Handbook of Pharmaceutical Excipients*. Chicago, Pharmaceutical Press, 2005. The formulations may also include one or more buffers, stabilizing agents, surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants,

opaquing agents, glidants, processing aids, colorants, sweeteners, perfuming agents, flavoring agents, diluents and other known additives to provide an elegant presentation of the drug (i.e., a compound of the present invention or pharmaceutical composition thereof) or aid in the manufacturing of the pharmaceutical product (i.e., medicament).

[0070] Compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. Pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including capsules, tablets, aqueous suspensions or solutions.


[0071] The amount of compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon a variety of factors, including the host treated and the particular mode of administration. It should also be understood that a specific dosage and treatment regimen for any particular subject will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

[0072] As used herein, the terms "treatment," "treat," and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment

may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.


[0073] In one embodiment, the invention provides for a method for treating a neurological or psychiatric disease or disorder, wherein the neurological or psychiatric disease or disorder is anxiety or attention deficit hyperactivity disorder.

[0074] In one embodiment, provided is a method of treating anxiety or attention deficit hyperactivity disorder in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound selected from:




or a pharmaceutically acceptable salt thereof.

[0075] In one embodiment, provided is a method of treating anxiety in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound selected from:



or a pharmaceutically acceptable salt thereof.

[0076] In one embodiment of the invention, provided is a method of treating attention deficit hyperactivity disorder in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound selected from:



or a pharmaceutically acceptable salt thereof.

[0077] In some embodiments, the invention provides a method for treating a neurological or psychiatric diseases or disorder in a subject, comprising administering to the subject an effective amount of a compound of this invention (or its pharmaceutically acceptable salt), or composition comprising a compound of this invention (or its pharmaceutically acceptable salt). Neurological and/or psychiatric diseases and disorders diseases can exhibit a variety of psychiatric and behavioral symptoms, including apathy, depression, anxiety, cognitive impairment, psychosis, aggression, agitation, poor impulse control and sleep disruptions.

[0078] In some embodiments, the neurological or psychiatric diseases or disorder is selected from a psychosis, including schizophrenia (paranoid, disorganized, catatonic or undifferentiated), schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition and substance-induced or drug-induced (e.g., phencyclidine, ketamine and other dissociative anesthetics, amphetamine and other psychostimulants and cocaine) psychosis; psychotic disorder, psychosis associated with affective disorders, brief reactive psychosis, schizoaffective psychosis, "schizophrenia-spectrum" disorders such as schizoid or schizotypal personality disorders, or illness associated with psychosis (such as major depression, manic depressive (bipolar) disorder, Alzheimer's disease and post-traumatic stress syndrome), including both positive, negative, and cognitive symptoms of schizophrenia and other psychoses; cognitive disorders including dementia (semantic dementia, frontotemporal dementia, dementia with depressive features, persisting, subcortical dementia, dementia with Lewy Bodies, Parkinsonism-ALS Dementia Complex, and dementia associated with Alzheimer's disease, ischemia, multi-infarct dementia, trauma, vascular problems, stroke, HIV disease, Parkinson's disease, Huntington's disease, Down syndrome, Pick's disease, Creutzfeldt-Jacob disease, perinatal hypoxia, or substance abuse), delirium, amnestic disorders or age related cognitive decline; anxiety disorders including acute stress disorder, agoraphobia, generalized anxiety disorder, obsessive-compulsive disorder, panic attack, panic disorder, post-traumatic stress disorder, separation anxiety disorder, social phobia, specific

phobia, substance-induced anxiety disorder and anxiety due to a general medical condition; substance-related disorders and addictive behaviors (including substance-induced delirium, persisting dementia, persisting amnestic disorder, psychotic disorder or anxiety disorder; tolerance, dependence or withdrawal from substances including alcohol, amphetamines, cannabis, cocaine, hallucinogens, inhalants, nicotine, opioids, phencyclidine, sedatives, hypnotics or anxiolytics); obesity, bulimia nervosa, pica and compulsive eating disorders; bipolar disorders, mood disorders including depressive disorders; depression including unipolar depression, seasonal depression and post-partum depression, atypical depression, catatonic depression, elderly depression, endogenous depression, melancholic depression, perinatal depression, situational depression, chronic depression, premenstrual syndrome (PMS) and premenstrual dysphoric disorder (PDD), mood disorders due to a general medical condition, and substance-induced mood disorders; learning disorders, pervasive developmental disorder including autistic disorder, attention disorders including attention-deficit hyperactivity disorder (ADHD) and conduct disorder; disorders such as autism, depression, benign forgetfulness, childhood learning disorders, specific learning disorders, intellectual development disorders, and closed head injury; movement disorders; epilepsy; urinary incontinence; neuronal damage including ocular damage, retinopathy or macular degeneration of the eye, tinnitus, hearing impairment and loss, and brain edema; emesis; and sleep disorders including insomnia and narcolepsy.

[0079] In some embodiments, the neurological or psychiatric disease or disorder is Alzheimer's disease, Parkinson's disease, depression, cognitive impairment, stroke, schizophrenia, Down syndrome, or Fetal Alcohol Syndrome. In some embodiments, the neurological or psychiatric disease or disorder is Alzheimer's disease. In some embodiments, the neurological or psychiatric disease or disorder is Parkinson's disease. In some embodiments, the neurological or psychiatric disease or disorder is depression. In some embodiments, the neurological or psychiatric disease disorder is cognitive impairment. In some embodiments, the cognitive impairment is cognitive dysfunction associated with depression, for example, major depressive disorder. In some embodiments, the neurological or psychiatric disorder is stroke. In some embodiments, the neurological or psychiatric disease or disorder is schizophrenia. In some embodiments, the neurological or psychiatric disease or disorder is Down syndrome. In some embodiments, the neurological or psychiatric disorder is Fetal Alcohol Syndrome.

[0080] In some embodiments, the neurological or psychiatric disease or disorder is bipolar disease. Bipolar disorder is a serious psychiatric disorder that has a prevalence of approximately 2% of the population, and affects both genders alike. It is a relapsing-remitting condition characterized by cycling between elevated (*i.e.*, manic) and depressed moods, which distinguishes it from other disorders such as major depressive disorder and schizophrenia. Bipolar I is defined by the occurrence of a full manic episode, although most individuals experience significant depression. Symptoms of mania include elevated or irritable mood, hyperactivity, grandiosity, decreased need for sleep, racing thoughts and in some cases, psychosis. The depressive episodes are characterized by anhedonia, sad mood, hopelessness, poor self-esteem, diminished concentration and lethargy. Bipolar II is defined as the occurrence of a major depressive episode and hypomanic (less severe mania) episode although subjects spend considerable more time in the depressive state. Other related conditions include cyclothymic disorder.

[0081] In some embodiments, the neurological or psychiatric disease or disorder is schizophrenia. Schizophrenia is a disorder of unknown origin, which usually appears for the first time in early adulthood and is marked by characteristics such as psychotic symptoms, phasic progression and development, and/or deterioration in social behavior and professional capability. Characteristic psychotic symptoms are disorders of thought content (*e.g.*, multiple, fragmentary, incoherent, implausible or simply delusional contents, or ideas of persecution) and of mentality (*e.g.*, loss of association, flight of imagination, incoherence up to incomprehensibility), as well as disorders of perceptibility (*e.g.*, hallucinations), emotions (*e.g.*, superficial or inadequate emotions), self-perceptions, intentions, impulses, and/or inter-human relationships, and psychomotoric disorders (*e.g.*, catatonia). Other symptoms are also associated with this disorder. Schizophrenia is classified into subgroups: the paranoid type, characterized by delusions and hallucinations and absence of thought disorder, disorganized behavior, and affective flattening; the disorganized type, also named "hebephrenic schizophrenia," in which thought disorder and flat affect are present together; the cataconic type, in which prominent psychomotor disturbances are evident, and symptoms may include catatonic stupor and waxy flexibility; and the undifferentiated type, in which psychotic symptoms are present but the criteria for paranoid, disorganized, or catatonic types have not been met. The symptoms of schizophrenia normally manifest themselves in three broad categories: positive, negative and cognitive symptoms. Positive symptoms are those which represent an "excess" of normal experiences, such as hallucinations and delusions. Negative

symptoms are those where the subject suffers from a lack of normal experiences, such as anhedonia and lack of social interaction. The cognitive symptoms relate to cognitive impairment in schizophrenics, such as lack of sustained attention and deficits in decision making.

[0082] In some embodiments, the neurological or psychiatric disease or disorder is anxiety disorder. Anxiety disorders are characterized by fear, worry, and uneasiness, usually generalized and unfocused as an overreaction to a situation. Anxiety disorders differ in the situations or types of objects that induce fear, anxiety, or avoidance behavior, and the associated cognitive ideation. Anxiety differs from fear in that anxiety is an emotional response to a perceived future threat while fear is associated with a perceived or real immediate threat. They also differ in the content of the associated thoughts or beliefs. Examples of anxiety disorders include separation anxiety disorder, selective mutism, specific phobia, social anxiety disorder (social phobia), panic disorder, panic attack specifier, agoraphobia, generalized anxiety disorder, substance/medication-induced anxiety disorder, anxiety disorder due to another medical condition, illness anxiety disorder, social (pragmatic) communication disorder, other specified anxiety disorder, and unspecified anxiety disorder; stressor-related disorders, including reactive attachment disorder, disinhibited social engagement disorder, posttraumatic stress disorder (PTSD), acute stress disorder, and adjustment disorders.

[0083] Cognitive impairment includes a decline in cognitive functions or cognitive domains, *e.g.*, working memory, attention and vigilance, verbal learning and memory, visual learning and memory, reasoning and problem solving (*e.g.*, executive function, speed of processing and/or social cognition). In particular, cognitive impairment may indicate deficits in attention, disorganized thinking, slow thinking, difficulty in understanding, poor concentration, impairment of problem solving, poor memory, difficulties in expressing thoughts, and/or difficulties in integrating thoughts, feelings and behavior, or difficulties in extinction of irrelevant thoughts.

[0084] In some embodiments, the neurological or psychiatric disease or disorder involves a deficit in cognition (cognitive domains as defined by the DSM-5 are: complex attention, executive function, learning and memory, language, perceptual-motor, social cognition). In some embodiments, the neurological or psychiatric disorder is associated with a deficit in dopamine signaling. In some embodiments, the neurological or psychiatric disorder is

associated with basal ganglia dysfunction. In some embodiments, the neurological or psychiatric disorder is associated with dysregulated locomotor activity. In some embodiments, the neurological or psychiatric disorder is associated with impairment of prefrontal cortex functioning.

[0085] In some embodiments, the present invention provides a method of treating one or more symptoms of a neurological and/or psychiatric disease or disorder provided herein. Such diseases and disorders include mood disorders, including bipolar I disorder, bipolar II disorder, bipolar depression, mania, cyclothymic disorder, substance/medication-induced bipolar and related disorders, bipolar and related disorder due to another medical condition, other specified bipolar and related disorder, and unspecified bipolar and related disorders; psychotic disorders, including schizophrenia, schizophrenia spectrum disorder, acute schizophrenia, chronic schizophrenia, NOS schizophrenia, schizoid personality disorder, schizotypal personality disorder, delusional disorder, psychosis, psychotic disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition, drug-induced psychosis (e.g., cocaine, alcohol, amphetamine), schizoaffective disorder, agitation, aggression, delirium, catatonia, dissociative identity disorder, paranoid personality disorder, psychotic depression, Schizotypal Personality Disorder, Childhood Disintegrative Disorder (Heller's Syndrome), Disintegrative Psychosis, Dissociative Amnesia, Somatic Symptom Disorder, Parkinson's psychosis, excitative psychosis, Tourette's syndrome, and organic or NOS psychosis; depressive disorders, including disruptive mood dysregulation disorder, major depressive disorder (MDD) (including major depressive episode), dysthymia, persistent depressive disorder (dysthymia), treatment resistant depression, premenstrual dysphoric disorder, substance/medication-induced depressive disorder, depressive disorder due to another medical condition, other specified depressive disorder, and unspecified depressive disorder; anxiety disorders; and other disorders including substance abuse or dependency (e.g., nicotine, alcohol, cocaine), addiction, internet gaming disorder, eating disorders, behavior disorder, seizure, vertigo, epilepsy, agitation, aggression, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, dyskinesias, Huntington's disease, dementia, premenstrual dysphoria, attention deficit disorder (ADD) and attention deficit hyperactivity disorder (ADHD)), hyperkinetic syndrome, autism, autism spectrum disorder, obsessive-compulsive disorder, pain, fibromyalgia, migraine, cognitive impairment, movement disorder, restless leg syndrome (RLS), multiple sclerosis, Primary Progressive Multiple Sclerosis, multiple sclerosis, sleep

disorder, sleep apnea, narcolepsy, excessive daytime sleepiness, jet lag, drowsy side effect of medications, insomnia, sexual dysfunction, hypertension, emesis, Lesche-Nyhane disease, Wilson's disease, Rett syndrome, and Huntington's chorea. In some embodiments, the neurological and/or psychiatric disorders include agitation and aggression.

[0086] In some embodiments, the agitation and aggression are associated with Alzheimer's disease, Parkinson's disease, and/or autism.

[0087] In some embodiments, the neurological and/or psychiatric diseases or disorders are obsessive-compulsive disorder and related disorders (e.g., body dysmorphic disorder, hoarding disorder, trichotillomania, excoriation disorder).

[0088] In some embodiments, the neurological and/or psychiatric diseases or disorders are disruptive, impulse-control, and conduct disorders including oppositional defiant disorder, intermittent explosive disorder, conduct disorder, antisocial personality disorder, pyromania, kleptomania, other specified disruptive, impulse-control, and conduct disorder, unspecified disruptive, impulse-control, and conduct disorder.

[0089] Depressive disorders include major depressive disorder and dysthymia, and are associated with depressed mood (sadness), poor concentration, insomnia, fatigue, appetite disturbances, excessive guilt and thoughts of suicide.

[0090] In some embodiments, the present invention provides a method of treating one or more symptoms including depression (e.g., major depressive disorder or dysthymia); bipolar disorder, seasonal affective disorder; cognitive deficit; sleep related disorder (e.g., sleep apnea, insomnia, narcolepsy, cataplexy) including those sleep disorders which are produced by psychiatric conditions; chronic fatigue syndrome; anxieties (e.g., general anxiety disorder, social anxiety disorder, panic disorder); obsessive compulsive disorder; post-menopausal vasomotor symptoms (e.g., hot flashes, night sweats); neurodegenerative disease (e.g., Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, primary lateral sclerosis, progressive muscular atrophy, progressive bulbar (atrophy) palsy, pseudobulbar palsy spinal muscular atrophy diseases (e.g., SMA type I, also called Werdnig-Hoffmann disease, SMA type II, SMA type III, also called Kugelberg-Welander disease, and Kennedy Disease, also called progressive spinobulbar muscular atrophy), Hallervorden-Spatz disease, Seitelberger disease (Infantile Neuroaxonal Dystrophy), adrenoleukodystrophy, Alexander Disease, autosomal dominant cerebellar ataxia (ADCA), pure autonomic failure (Bradbury-

Eggleston Syndrome), CADASIL Syndrome, and neuronal ceroids lipofuscinose disorders such as Batten Disease (Spielmeyer-Vogt-Sjögren)); manic disorder; dysthymic disorder; and obesity.

[0091] In some embodiments, a depressive disorder is associated with acute suicidality or suicide ideation. The United States Food and Drug Administration has adopted a "black box" label warning indicating that antidepressants may increase the risk of suicidal thinking and behavior in some children, adolescents and young adults (up to age 24) with a depressive disorder such as MDD. In some embodiments, a provided compound does not increase the risk of suicidal thinking and/or behavior in children, adolescents and/or young adults with a depressive disorder, e.g., with MDD. In some embodiments, the present invention provides a method of treating one or more symptoms of a depressive disorder (e.g., MDD) in children, adolescents and/or young adults without increasing the risk of suicidal thinking and/or behavior.

[0092] In some embodiments, the present invention provides a method of treating one or more symptoms including senile dementia, Early Onset Alzheimer's Disease, Alzheimer's type dementia, cognition, memory loss, amnesia/amnestic syndrome, disturbances of consciousness, coma, lowering of attention, speech disorder, agnosia, aphasia, apraxia , Mild Cognitive Impairment (MCI), benign forgetfulness, mild neurocognitive disorder, major neurocognitive disorder, neurocognitive disorder due to disease (e.g., Huntington's Disease, Parkinson's disease, Prion Disease, Traumatic Brain Injury, HIV or AIDS), Binswanger's Disease (subcortical leukoencephalopathy), and Capgras Syndrome.

[0093] In some embodiments, the present invention provides a method of treating one or more symptoms of pain, e.g., neuropathic pain, sensitization accompanying neuropathic pain, or inflammatory pain. In some embodiments, the pain is neuropathic pain, including post herpetic (or post-shingles) neuralgia, reflex sympathetic dystrophy/causalgia or nerve trauma, phantom limb pain, carpal tunnel syndrome, and peripheral neuropathy (such as diabetic neuropathy or neuropathy arising from chronic alcohol use). In some embodiments, the pain is acute pain, nociceptive pain, arthritis pain, rheumatoid arthritis, osteoarthritis, joint pain, muscoskeletal pain, back pain, dorsalgia, bulging disc, hip pain, visceral pain, headache, tension headache, acute tension headache, chronic tension headache, chronic cluster headache, common migraine, classic migraine, cluster headache, mixed headache, post-traumatic headache, eye strain headache, Short-lasting Unilateral Neuralgiform (SUNCT)

headache, SUNCT Syndrome, herpes zoster, acute herpes zoster, shingles, postherpetic neuralgia (shingles), causalgia, central pain, central pain syndrome, chronic back pain, neuralgia, neuropathic pain syndrome, neuropathy, diabetic neuropathy, diabetes-related neuropathy, diabetes-related nerve pain, fibrositis, peripheral neuropathy caused by chemotherapy, peripheral nerve disease, peripheral neuropathy, nerve pain, nerve trauma, sensitization accompanying neuropathic pain, complex regional pain syndrome, compression neuropathy, craniofacial pain, chronic joint pain, chronic knee pain, chronic pain syndrome, cancer pain, trigeminal neuralgia, tic doloreaux, reflex sympathetic causalgia, painful peripheral neuropathy, spinal nerve injury, arachnoiditis, spinal pain, Bernhardt-Roth Syndrome (meralgia parasthetica), carpal tunnel syndrome, cerebrospinal fluid syndrome, Charcot-Marie-tooth disease, hereditary motor and sensory neuropathy, peroneal muscular atrophy, cluster-tic syndrome, coccygeal pain syndromes, compartment syndrome, degenerative disc disease, failed back surgery syndrome, genito-pelvic pain/penetration disorder, gout, inflammatory pain, lumbar radiculopathy, neuroma (painful scar), pain associated with multiple sclerosis, pelvic floor disorders, phantom limb pain, piriformis syndrome, psychogenic pain, radicular pain syndrome, Raeder's syndrome, referred pain, reflex sympathetic dystrophy syndrome, sciatica, sciatica pain, scoliosis, slipped disc, somatic pain, spinal stenosis, stiff-person syndrome/stiff-man syndrome, stump pain, sympathetically maintained pain, tolosa-hunt syndrome, whiplash, or pain associated with Lyme disease.

[0094] In some embodiments, the present invention provides a method of treating one or more symptoms including obesity; migraine or migraine headache; and sexual dysfunction, in men or women, including without limitation sexual dysfunction caused by psychological and/or physiological factors, erectile dysfunction, premature ejaculation, vaginal dryness, lack of sexual excitement, inability to obtain orgasm, and psycho-sexual dysfunction, including without limitation, inhibited sexual desire, inhibited sexual excitement, inhibited female orgasm, inhibited male orgasm, functional dyspareunia, functional vaginismus, and atypical psychosexual dysfunction.

[0095] In some embodiments, the present invention provides a method of suppressing rapid eye movement (REM) during both sleep and daytime equivalent.

[0096] In some embodiments, the present invention provides a method of suppressing or eliminating pathological or excessive REM during the night or daytime equivalent.

[0097] In some embodiments, the present invention provides a method of treating one or more symptoms including cataplexy (sudden involuntary transient bouts of muscle weakness or paralysis while awake); nighttime sleep disturbance/sleep fragmentation associated with narcolepsy or other conditions; sleep paralysis associated with narcolepsy or other conditions; hypnagogic and hypnapompic hallucinations associated with narcolepsy or other conditions; and excessive daytime sleepiness associated with narcolepsy, sleep apnea or shift work disorder and other medical conditions such as cancer, chronic fatigue syndrome and fibromyalgia.

[0098] In some embodiments, the present invention provides a method of treating one or more symptoms of movement disorders, including akinesias, akinetic-rigid syndromes, dyskinesias and dystonias. Examples of akinesias and akinetic-rigid syndromes include Parkinson's disease, drug-induced Parkinsonism, postencephalitic Parkinsonism, secondary Parkinsonism, Parkinson plus syndromes, atypical Parkinsonism, idiopathic Parkinsonism, progressive supranuclear palsy, multiple system atrophy, corticobasal degeneration, Parkinsonism-ALS dementia complex and basal ganglia calcification, medication-induced Parkinsonism (such as neuroleptic-induced parkinsonism, neuroleptic malignant syndrome, neuroleptic-induced acute dystonia, neuroleptic-induced acute akathisia, neuroleptic-induced tardive dyskinesia and medication-induced postural tremor), Gilles de la Tourette's syndrome, epilepsy, muscular spasms and disorders associated with muscular spasticity or weakness including tremors. Examples of dyskinesias include drug (e.g. L-DOPA) induced dyskinesia tremor (such as rest tremor, postural tremor, intention tremor), chorea (such as Sydenham's chorea, Huntington's disease, benign hereditary chorea, neuroacanthocytosis, symptomatic chorea, drug-induced chorea and hemiballism), myoclonus (including generalised myoclonus and focal myoclonus), tics (including simple tics, complex tics and symptomatic tics). Examples of dystonias include generalised dystonia, idiopathic dystonia, drug-induced dystonia, symptomatic dystonia, paroxysmal dystonia, focal dystonia, blepharospasm, oromandibular dystonia, spasmoid dysphonia, spasmoid torticollis, axial dystonia, dystonic writer's cramp and hemiplegic dystonia. Other examples of movement disorders include stereotypic movement disorder, persistent (chronic) motor disorder, medication-Induced movement disorder, psychogenic movement disorders, substance/medication-Induced movement disorder, extrapyramidal movement disorders, hyperkinetic movement disorders, hypokinetic movement disorders, alternating hemiplegia, Angelman syndrome, Hallervorden-Spatz Disease, ataxia, dentate cerebellar ataxia, ataxia telangiectasia (Louis-Bar syndrome),

Friedreich's Ataxia, hereditary spinal ataxia, hereditary spinal sclerosis, Machado-Joseph Disease, spinocerebellar ataxia, progressive myoclonic ataxia, athetosis, ballismus, blepharospasm (eye twitching), cerebral palsy, tardive dystonia, tardive dyskinesia, idiopathic torsion dystonia, torsion dystonia, focal dystonia, idiopathic familial dystonia, Idiopathic nonfamilial dystonia, cervical dystonia (spasmodic torticollis), primary dystonia, orofacial dystonia, developmental coordination disorder, bulbospinal muscular atrophy (Kennedy's Disease), Shy-Drager Syndrome, and Stiff-Person (Stiff-Man) Syndrome.

[0099] In some embodiments, the present invention provides a method of treating one or more symptoms of epilepsy and/or seizures, including abdominal epilepsy, absence seizure, acquired epilepsy, acquired epileptiform aphasia, Aicardi syndrome, Alpers' disease, Alpers-Huttenlocher syndrome, Angelman syndrome, benign focal epilepsy, benign focal epilepsy of childhood, benign intracranial hypertension, benign rolandic epilepsy (BRE), CDKL5 disorder, childhood absence epilepsy, dentate cerebellar ataxia, Doose syndrome, Dravet syndrome, dyscognitive focal seizure, epilepsy with grand mal seizures, epilepsy with myoclonic-absences, epileptic hemiplegia, febrile seizures, focal seizure, frontal lobe epilepsy, generalized tonic-clonic seizures, genetic epilepsy, Glut1 deficiency syndrome, hypothalamic hamartoma, idiopathic epilepsy, idiopathic generalized epilepsy, idiopathic localization-related epilepsies, idiopathic partial epilepsy, idiopathic seizure, juvenile absence epilepsy, juvenile myoclonic epilepsy, Lafora disease, Lafora progressive myoclonus epilepsy, Landau-Kleffner syndrome, Lassueur-Graham-Little syndrome, Lennox syndrome, Lennox-Gastaut syndrome, medically refractory epilepsy, mesial-temporal lobe sclerosis, myoclonic seizure, neonatal epilepsy, occipital lobe epilepsy, Ohtahara syndrome, Panayiotopoulos syndrome, parietal lobe epilepsy, PCDH19 epilepsy, photosensitive epilepsy, progressive myoclonic epilepsies, Rasmussen's encephalitis, Rasmussen's syndrome, refractory epilepsy, seizure disorder, status epilepticus, Sturge-Weber syndrome, symptomatic generalized epilepsy, symptomatic parital epilepsy, TBCK-related ID syndrome, temporal lobe epilepsy, temporal lobe seizures, tonic-clonic seizure, West syndrome, tremor, cerebellar tremor, cerebellar outflow tremor, intention tremor, essential tremor, benign essential tremor, Parkinsonian tremor, and medication-induced postural tremor.

[0100] In some embodiments, the present invention provides a method of treating a neurological and/or psychiatric disease or disorder described herein, comprising

administering a compound of the invention in conjunction with one or more pharmaceutical agents. Suitable pharmaceutical agents that may be used in combination with the compounds of the present invention include anti-Parkinson's drugs, anti-Alzheimer's drugs, anti-depressants, anti-psychotics, anti-ischemics, CNS depressants, anti-cholinergics, nootropics, epilepsy medication, attention (e.g., ADD/ADHD) medications, sleep-promoting medications, wakefulness-promoting medications, and pain medications. In some embodiments, suitable pharmaceutical agents are anxiolytics.

[0101] Suitable anti-Parkinson's drugs include dopamine replacement therapy (e.g. L-DOPA, carbidopa, COMT inhibitors such as entacapone or tolcapone), dopamine agonists (e.g. D1 agonists, D2 agonists, mixed D1/D2 agonists, bromocriptine, pergolide, cabergoline, ropinirole, pramipexole, piribedil, or apomorphine in combination with domperidone), histamine H2 antagonists, monoamine oxidase inhibitors (such as selegiline, rasagiline, safinamide and tranylcypromine), certain atypical antipsychotics such as pimavanserin (a non-dopaminergic atypical antipsychotic and inverse agonist of the serotonin 5-HT_{2A} receptor), and amantadine.

[0102] In some embodiments, compounds of the invention can be used in combination with levodopa (with or without a selective extracerebral decarboxylase inhibitor such as carbidopa or benserazide), anticholinergics such as biperiden (optionally as its hydrochloride or lactate salt) and trihexyphenidyl(benzhexyl)hydrochloride, COMT inhibitors such as entacapone or tolcapone, MAO A/B inhibitors, antioxidants, A2a adenosine receptor antagonists, cholinergic agonists, NMDA receptor antagonists, serotonin receptor antagonists and dopamine receptor agonists such as alentemol, bromocriptine, fenoldopam, lisuride, naxagolide, pergolide and pramipexole. It will be appreciated that the dopamine agonist may be in the form of a pharmaceutically acceptable salt, for example, alentemol hydrobromide, bromocriptine mesylate, fenoldopam mesylate, naxagolide hydrochloride and pergolide mesylate. Lisuride and pramipexole are commonly used in a non-salt form.

[0103] Suitable anti-Alzheimer's drugs include beta-secretase inhibitors, gamma-secretase inhibitors, cholinesterase inhibitors such as donepezil, galantamine or rivastigmine, HMG-CoA reductase inhibitors, NSAID's including ibuprofen, vitamin E, and anti-amyloid antibodies. In some embodiments, an anti-Alzheimer's drug is memantine.

[0104] Suitable anti-depressants and anti-anxiety agents include norepinephrine reuptake inhibitors (including tertiary amine tricyclics and secondary amine tricyclics), selective serotonin reuptake inhibitors (SSRIs), monoamine oxidase inhibitors (MAOIs), reversible inhibitors of monoamine oxidase (RIMAs), serotonin and noradrenaline reuptake inhibitors (SNRIs), corticotropin releasing factor (CRF) antagonists, α -adrenoreceptor antagonists, neurokinin-1 receptor antagonists, atypical anti-depressants, benzodiazepines, 5-HT1A agonists or antagonists, especially 5-HT1A partial agonists, and corticotropin releasing factor (CRF) antagonists.

[0105] Specific suitable anti-depressant and anti-anxiety agents include amitriptyline, clomipramine, doxepin, imipramine and trimipramine; amoxapine, desipramine, citalopram, escitalopram, maprotiline, nortriptyline and protriptyline; fluoxetine, fluvoxamine, paroxetine and sertraline; isocarboxazid, phenelzine, tranylcypromine and selegiline; moclobemide; venlafaxine; desvenlafaxine, duloxetine; aprepitant; bupropion, vilazodone, mirtazapine, lithium, nefazodone, trazodone and viloxazine; alprazolam, chlordiazepoxide, clonazepam, chlorazepate, diazepam, halazepam, lorazepam, oxazepam and prazepam; buspirone, flesinoxan, gepirone and ipsapirone, reboxetine, vortioxetine, clorazepate, and pharmaceutically acceptable salts thereof. In some embodiments, suitable anti-depressant and anti-anxiety agents are tianeptine, or pharmaceutically acceptable salts thereof.

[0106] Suitable anti-psychotic and mood stabilizer agents include D2 antagonists, 5HT2A antagonists, atypical antipsychotics, lithium, and anticonvulsants.

[0107] Specific suitable anti-psychotic and mood stabilizer agents include chlorpromazine, fluphenazine, haloperidol, amisulpride, perphenazine, thioridazine, trifluoperazine, aripiprazole, asenapine, clozapine, olanzapine, paliperidone, quetiapine, risperidone, ziprasidone, lurasidone, flupentixol, levomepromazine, pericyazine, perphenazine, pimozide, prochlorperazine, zuclopentixol, olanzapine and fluoxetine, lithium, carbamazepine, lamotrigine, valproic acid, iloperidone, thiothixene and pharmaceutically acceptable salts thereof.

[0108] Suitable epilepsy medications include levetiracetam, oxcarbazepine, clobazam, retigabine, zonisamide, felbamate, esliccarbazepine acetate, lacosamide, carbamazepine, tiagabine, methsuximide, progabide, valproic acid, lamotrigine, brivaracetam, rufinamide, topiramate and perampanel.

[0109] Suitable attention medications include methyl phenidate, atomoxetine, guanfacine, D-amphetamine, lisdexamphetamine, methylamphetamine, and clonidine.

[0110] Suitable sleep-promoting medications include ramelteon, triazolam, zopiclone, eszopiclone, zolpidem, temazepam, and trazodone.

[0111] Suitable wakefulness-promoting medications include Modafinil, D-Amphetamine, caffeine, and armodafinil.

[0112] Suitable pain medications include dextromethorphan, tapentadol, buprenorphine, codeine, fentanyl, hydrocodone, hydromorphone, morphine, naloxegol, oxycodone, tramadol, gabapentil, difluprednate, pregabalin, acetyl salicyclic acid, bromfenac, diclofenac, diflunisal, indomethacin, ketorolac, meoxicam, and naproxen.

[0113] In some embodiments, compounds of the invention may be used in combination with other therapies. Suitable therapies include psychotherapy, cognitive behavioral therapy, electroconvulsive therapy, transcranial magnetic stimulation, vagus nerve stimulation, and deep-brain stimulation.

[0114] The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the condition, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the subject to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment.

[0115] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the disease or disorder being treated. In some embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[0116] In some embodiments, a combination of two or more therapeutic agents may be administered together with the compounds of the invention. In some embodiments, a combination of three or more therapeutic agents may be administered with the compounds of the invention.

[0117] Other examples of agents the compounds of this invention may also be combined with include: vitamins and nutritional supplements, antiemetics (e.g. 5-HT3 receptor antagonists, dopamine antagonists, NK1 receptor antagonists, histamine receptor antagonists, cannabinoids, benzodiazepines, or anticholinergics), agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®, dalfampridine, alemtuzumab), Copaxone®, and mitoxantrone; treatments for Huntington's disease such as tetrabenazine; treatments for asthma such as albuterol and Singulair®; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins, fibrates, cholesterol absorption inhibitors, bile acid sequestrants, and niacin; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; agents for treating immunodeficiency disorders such as gamma globulin; and anti-diabetic agents such as biguanides (metformin, phenformin, buformin), thiazolidinediones (rosiglitazone, pioglitazone, troglitazone), sulfonylureas (tolbutamide, acetohexamide, tolazamide, chlorpropamide, glipizide, glyburide, glimepiride, gliclazide), meglitinides (repaglinide, nateglinide), alpha-glucosidase inhibitors (miglitol, acarbose), incretin mimetics (exenatide, liraglutide, taspoglutide), gastric inhibitory peptide analogs, DPP-4 inhibitors (vildagliptin, sitagliptin, saxagliptin, linagliptin, alogliptin), amylin analogs (pramlintide), and insulin and insulin analogs.

[0118] In some embodiments, a compound of the present invention, or a pharmaceutically acceptable salt thereof, is administered in combination with an antisense agent, a monoclonal or polyclonal antibody, or a siRNA therapeutic.

[0119] Those additional agents may be administered separately from an inventive compound-containing composition, as part of a multiple dosage regimen. Alternatively, those agents may be part of a single dosage form, mixed together with a compound of this invention in a single composition. If administered as part of a multiple dosage regime, the two active agents may be submitted simultaneously, sequentially or within a period of time from one another, normally within five hours from one another.

[0120] As used herein, the term "combination," "combined," and related terms refers to the simultaneous or sequential administration of therapeutic agents in accordance with this invention. For example, a compound of the present invention may be administered with another therapeutic agent simultaneously or sequentially in separate unit dosage forms or together in a single unit dosage form. Accordingly, the present invention provides a single unit dosage form comprising a compound of formula I, or a pharmaceutically acceptable salt thereof, an additional therapeutic agent, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

[0121] The amount of both, an inventive compound and additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Preferably, compositions of this invention should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of an inventive can be administered.

[0122] In those compositions which comprise an additional therapeutic agent, that additional therapeutic agent and the compound of this invention may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing only that therapeutic agent. In such compositions a dosage of between 0.01 - 100 mg/kg body weight/day of the additional therapeutic agent can be administered.

[0123] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about

50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.


[0124] In some embodiments, the present invention provides a medicament comprising at least one compound of formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

[0125] In some embodiments, the present invention provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of a neurological and/or psychiatric disease or disorder.

EXAMPLES

[0126] As depicted in the Examples below, in some embodiments, compounds are prepared according to the following procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds of the present invention, the following methods, and other methods known to persons skilled in the art, can be applied to all compounds and subclasses and species of each of these, as described herein.

[0127] Example 1. 6-Chloroisochroman-4-amine hydrochloride



Step 1. Methyl 2-hydroxyacetate (1.72 g, 19.2 mmol) was dissolved in DMF (50 mL) and treated with sodium hydride (460 mg, 19.2 mmol, 95%). The reaction mixture was stirred at room temperature for 30 min, and then cooled in an ice bath. A solution of 2.53 g (9.60 mmol) of methyl 2-(bromomethyl)-5-chlorobenzoate (Thorarensen, A., *et al*

WO2004018414) in DMF (5 mL) was added, and the reaction mixture was stirred at 0°C for 10 min. Excess hydride was carefully quenched by the dropwise addition of saturated aqueous NH₄Cl (20 mL). The mixture was diluted with water (50 mL) and extracted with EtOAc (4×60 mL). The combined organics were washed with brine (2×25 mL), anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 5% EtOAc/hexanes) gave methyl 5-chloro-2-((2-methoxy-2-oxoethoxy)methyl)benzoate **1.1** (1.82 g, 70%) as a white solid. MS (ESI): *m/z* 273 [M+H].

Step 2. A solution of compound **1.1** (1.83 g, 6.71 mmol) in EtOH (10 mL) was treated with a solution of NaOH (1.2 g, 30.1 mmol) in water (7 mL), and stirred at room temperature for 1 h. The mixture was acidified to pH = 3-4 by the addition of 1M aqueous HCl. The aqueous phase was extracted with DCM (4×50 mL). The combined organics were washed with brine (2×20 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide 2-((carboxymethoxy)methyl)-5-chlorobenzoic acid **1.2** (1.59 g, 97%) as a white solid. LC-MS (ESI): *m/z* 243 [M-H].

Step 3. Potassium acetate (2.56 g, 26.1 mmol) was added to a solution of compound **1.2** (1.6 g, 6.54 mmol) in Ac₂O (30 mL). The reaction mixture was heated to 140°C and stirred for 2 h. Ice water (100 mL) and Et₂O (100 mL) was added to the reaction vessel and the resulting biphasic mixture was transferred to a separatory funnel. The layers were separated and the aqueous phase was extracted with Et₂O (5×50 mL). The combined organics were washed with brine (2×20 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide acetic 6-chloro-4-oxoisochroman-3-carboxylic anhydride **1.3** (1.02 g, 15%) as brown oil, which was used without further purification.

Step 4. A solution of compound **1.3** (1.02 g, 3.83 mmol) in EtOH (30 mL) was treated with a solution of NaOH (306 mg, 7.66 mmol) in water (10 mL). The reaction mixture was stirred at ambient temperature for 15 min and then extracted with EtOAc (4×10 mL). The combined organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 5% EtOAc/hexanes) afforded 6-chloroisochroman-4-one **1.4** (0.505 g, 72%) as a white solid. ¹H NMR (300 MHz, DMSO-*d*₆) *δ* 8.03 (d, *J* = 1.1 Hz, 1H), 7.55 (dd, *J* = 8.1, 2.1 Hz, 1H), 7.21 (d, *J* = 8.1 Hz, 1H), 4.89 (s, 2H), 4.38 (s, 2H).

Step 5. To a solution of compound **1.4** (506 mg, 2.77 mmol) in EtOH (9 mL) was added NaBH₄ (104 mg, 2.77 mmol). The reaction was stirred at ambient temperature for 1 h. EtOAc (20 mL) and saturated aqueous NH₄Cl (15 mL) was added to the reaction vessel and the resulting biphasic mixture was transferred to a separatory funnel. The layers were separated and the aqueous phase was extracted with EtOAc (3×20 mL). The combined organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide crude 6-chloroisochroman-4-ol **1.5** (478 mg, 94%) as colorless oil, which was used without further purification. ¹H NMR (300 MHz, CDCl₃) δ 7.47 (d, *J* = 2.1 Hz, 1H), 7.25 (d, *J* = 7.2 Hz, 1H), 6.98 (d, *J* = 8.4 Hz, 1H), 4.79 (d, *J* = 15.3 Hz, 1H), 4.66 (d, *J* = 15.3 Hz, 1H), 4.55 (s, 1H), 4.08 (dd, *J* = 12.0, 3.3 Hz, 1H), 3.88 (dd, *J* = 12.0, 2.7 Hz, 1H), 3.73 (dd, *J* = 14.1, 6.9 Hz, 1H), 2.36 (br s, 1H).


Step 6. A solution of compound **1.5** (480 mg, 2.59 mmol) in toluene (7 mL) was treated with diphenyl phosphoryl azide (853 mg, 3.10 mmol), and cooled in an ice bath. A solution of DBU (777 mg, 3.10 mmol) in toluene (1.5 mL) was added, and the reaction was stirred at ambient temperature for 16 h. The reaction mixture was directly concentrated under reduced pressure and purified by flash column chromatography (SiO₂, gradient elution from 5% EtOAc/hexanes to 10% EtOAc/hexanes) to afford 4-azido-6-chloroisochroman **1.6** (329 mg, 61%) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.39 (d, *J* = 2.1 Hz, 1H), 7.32 (dd, *J* = 8.1, 2.1 Hz, 1H), 7.04 (d, *J* = 8.1 Hz, 1H), 4.86 (d, *J* = 15.3 Hz, 1H), 4.70 (d, *J* = 15.3 Hz, 1H), 4.24-2.19 (m, 2H), 3.97 (dd, *J* = 11.4, 2.4 Hz, 1H).

Step 7. A solution of compound **1.6** (330 mg, 1.57 mmol) in THF (15 mL) was treated with LiAlH₄ (59.5 mg, 1.57 mmol) at room temperature. The reaction mixture was stirred at room temperature for 30 min. A solution of Na₂CO₃ (164 mg, 1.55 mmol) in water (10 mL), EtOAc (15 mL) and di-tert-butyl dicarbonate (0.676 g, 3.1 mmol) was added and the resulting mixture was stirred at ambient temperature for 4 h. The mixture was diluted with water (10 mL) was added to the reaction vessel and aqueous phase was extracted with EtOAc (3×25 mL). The combined organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, gradient elution from 100% hexanes to 5% EtOAc/hexanes) gave *tert*-butyl (6-chloroisochroman-4-yl)carbamate **1.7** (349 mg, 78%) as a white solid. LC-MS (ESI): *m/z* 284 [M+H].

Step 8. A solution of compound **1.7** (349 mg, 1.23 mmol) in Et₂O (2 mL) was treated with a saturated solution of HCl in Et₂O (5 mL). The reaction mixture was stirred at room temperature for 16 h and then concentrated under reduced pressure. The resulting solid residue was triturated with Et₂O (2×5 mL) to provide the compound of **Example 1** (165 mg, 61%) as a white solid. LC-MS (ESI): *m/z* 184 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.52 (brs, 3H), 7.64 (d, *J* = 1.8 Hz, 1H), 7.46 (dd, *J* = 8.4, 2.1 Hz, 1H), 7.24 (d, *J* = 8.4 Hz, 1H), 4.83 (d, *J* = 15.9 Hz, 1H), 4.68 (d, *J* = 15.9 Hz, 1H), 4.38 (s, 1H), 4.14 (dd, *J* = 12.3, 1.8 Hz, 1H), 3.89 (dd, *J* = 12.6, 3.0 Hz, 1H).


[0128] **Table 1.** Compounds prepared as described in **Example 1** using the appropriately substituted starting materials.

| Example | Structure | Characterization Data |
|----------|-----------|--|
| 2 | | MS (ESI): <i>m/z</i> 184 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.58 (br s, 3H), 7.58 (d, <i>J</i> = 8.4 Hz, 1H), 7.42 (dd, <i>J</i> = 8.4, 2.1 Hz, 1H), 7.32 (d, <i>J</i> = 1.8 Hz, 1H), 4.83 (d, <i>J</i> = 15.6 Hz, 1H), 4.69 (d, <i>J</i> = 15.9 Hz, 1H), 4.36 (s, 1H), 4.18 (dd, <i>J</i> = 12.6, 1.8 Hz, 1H), 3.88 (dd, <i>J</i> = 12.6, 2.4 Hz, 1H). |
| 3 | | MS (ESI): <i>m/z</i> 168 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.61 (br s, 3H), 7.65–7.61 (m, 1H), 7.23–7.12 (m, 1H), 7.01 (dd, <i>J</i> = 9.6, 2.7 Hz, 1H), 4.83 (d, <i>J</i> = 15.9 Hz, 1H), 4.69 (d, <i>J</i> = 15.6 Hz, 1H), 4.35 (s, 1H), 4.20 (dd, <i>J</i> = 12.6, 1.8 Hz, 1H), 3.88 (dd, <i>J</i> = 12.6, 2.7 Hz, 1H). |
| 4 | | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.50 (br s, 1H), 7.42 (d, <i>J</i> = 7.8 Hz, 1H), 7.14 (d, <i>J</i> = 7.8 Hz, 1H), 6.98 (s, 1H), 4.78 (d, <i>J</i> = 15.3 Hz, 1H), 4.67 (d, <i>J</i> = 15.3 Hz, 1H), 4.28 (s, 1H), 4.17 (dd, <i>J</i> = 12.3, 1.8 Hz, 1H), 3.87 (dd, <i>J</i> = 12.3, 2.7 Hz, 1H), 2.30 (s, 3H). |
| 5 | | MS (ESI): <i>m/z</i> 184 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.76 (s, 3H), 7.60 (d, <i>J</i> = 7.5 Hz, 1H), 7.51 (d, <i>J</i> = 7.2 Hz, 1H), 7.40 (t, <i>J</i> = 7.8 Hz, 1H), 4.82 (d, <i>J</i> = 15.9 Hz, 1H), 4.63 (d, <i>J</i> = 15.9 Hz, 1H), 4.39 (s, 1H), 4.24 (d, <i>J</i> = 12.3 Hz, 1H), 3.89 (dd, <i>J</i> = 12.6, 2.4 Hz, 1H). |
| 6 | | MS (ESI): <i>m/z</i> 167 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.57 (br s, 2H), 7.43 (d, <i>J</i> = 9.6 Hz, 1H), 7.24 (d, <i>J</i> = 12.0 Hz, 2H), 4.82 (d, <i>J</i> = 15.3 Hz, 1H), 4.68 (d, <i>J</i> = 15.3 Hz, 1H), 4.37 (s, 1H), 4.14 (d, <i>J</i> = 12.6 Hz, 1H) 3.90 (dd, <i>J</i> = 12.6 Hz, 3.0 Hz, 1H). |

Example 7: 6-Phenylisochroman-4-amine hydrochloride

Step 1. A solution of 170 mg (0.45 mmol) of *tert*-butyl (6-iodoisochroman-4-yl)carbamate **7.1** (prepared from methyl 2-(bromomethyl)-5-iodobenzoate (Hayakawa, I., *et al*, WO2011068171) as described in Example 1) in 1,4-dioxane (4 mL) was treated with phenyl boronic acid (110 mg, 0.90 mmol) and Pd(PPh₃)₄ (52.3 mg, 0.045 mmol) and stirred at room temperature for 15 min. Aqueous 3M K₂CO₃ (0.76 mL, 2.3 mmol) was added, and the reaction mixture was heated to 100°C with stirring for 5 h. After cooling to room temperature, the mixture was diluted with water (5 mL) and extracted with EtOAc (3×15 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, gradient elution from 1% EtOAc/hexanes to 5% EtOAc/hexanes) gave *tert*-butyl (6-phenylisochroman-4-yl)carbamate **7.2** (110 mg, 75%) as a white solid. LC-MS (ESI): *m/z* 326 [M+H].

Step 2. The compound of **Example 7** was prepared as previously described in Example 1. MS (ESI): *m/z* 226 [M+H]. ¹H NMR (300 MHz, DMSO) δ 8.50 (s, 3H), 7.87 (s, 1H), 7.11 (d, *J* = 7.5 Hz, 3H), 7.53-7.37 (m, 3H), 7.28 (d, *J* = 8.1 Hz, 1H), 4.88 (d, *J* = 15.9 Hz, 1H), 4.75 (d, *J* = 15.3 Hz, 1H), 4.42 (s, 1H), 4.17 (d, *J* = 12.6 Hz, 1H), 3.95 (dd, *J* = 12.3, 2.4 Hz, 1H).


[0129] Example 8: 6-Isopropylisochroman-4-amine hydrochloride

Step 1. 10% Pd/C (25 mg) was added to 250 mg (0.86 mmol) of *tert*-butyl (6-(prop-1-en-2-yl)isochroman-4-yl)carbamate **8.1** (prepared as described in Example 7 using isopropenylboronic acid pinacol ester) in CH₃OH (25 mL). The reaction mixture was stirred at room temperature under a H₂ atmosphere for 16 h. After filtering through a Celite pad, the

filtrate was concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 , 5% EtOAc/hexanes) afforded *tert*-butyl (6-isopropylisochroman-4-yl) carbamate **8.2** (180 mg, 72%) as a brown oil. ^1H NMR (300 MHz, CDCl_3) δ 7.29 (s, 1H), 7.15 (dd, J = 8.1, 1.5 Hz, 1H), 6.95 (d, J = 8.1 Hz, 1H), 5.10 (d, J = 9.0 Hz, 1H), 4.81-4.68 (m, 3H), 4.07 (dd, J = 9.0, 1.8 Hz, 1H), 3.87 (dd, J = 12.0, 3.0 Hz, 1H), 2.95-2.86 (m, 1H), 1.60 (s, 9H), 0.97 (d, J = 7.2 Hz, 6H).

Step 2. The compound of **Example 8** was prepared as described in Example 1. MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.38 (br s, 3H), 7.42 (s, 1H), 7.27 (d, J = 8.1 Hz, 1H), 7.09 (d, J = 8.1 Hz, 1H), 4.79 (d, J = 15.0 Hz, 1H), 4.67 (d, J = 15.6 Hz, 1H), 4.31 (s, 1H), 4.14 (d, J = 12.3 Hz, 1H), 3.87 (d, J = 12.6 Hz, 1H), 2.91-2.86 (m, 1H), 1.21 (d, J = 6.9 Hz, 6H).


[0130] **Example 9: 6-Trifluoromethylisochroman-4-amine hydrochloride**



Step 1. A solution of compound **7.1** (227 mg, 605 μmol) in anhydrous DMF (8 mL) under a N_2 atmosphere was treated with copper(I) iodide (230 mg, 1.21 mmol) and hexamethylphosphoramide (541 mg, 3.02 mmol), and stirred at ambient temperature for 10 min. Methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (580 mg, 3.02 mmol) was added, and the reaction mixture was heated to 80°C and stirred for 6 h. After cooling to room temperature, the mixture was diluted with saturated aqueous NaHCO_3 (20 mL) and partitioned with EtOAc (20 mL). The aqueous phase was extracted with EtOAc (2 \times 30 mL). The combined organics were dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 , gradient elution from 5% EtOAc/hexanes to 10% EtOAc/hexanes) afforded *tert*-butyl (6-(trifluoromethyl) isochroman-4-yl)carbamate **9.1** (85.0 mg, 44%) as a white solid. MS (ESI): m/z 340 [M+Na].

Step 2. The compound of **Example 9** was prepared as previously described in Example 1. MS (ESI): m/z 218 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.57 (s, 3H), 7.96 (s, 1H), 7.75 (d, J = 7.5 Hz, 1H), 7.44 (d, J = 8.1 Hz, 1H), 4.92 (d, J = 16.2 Hz, 1H), 4.77 (d, J = 15.6 Hz, 1H), 4.95 (s, 1H), 4.18 (d, J = 12.3 Hz, 1H), 3.93 (dd, J = 12.6, 2.4 Hz, 1H).

[0131] Example 10: 6-Methoxyisochroman-4-amine hydrochloride



Step 1. To a solution of prop-2-en-1-ol (6.38 g, 110 mmol) in DMF (400 mL) was added 95% sodium hydride (4.38 g, 110 mmol) and 2-bromo-1-(bromomethyl)-4-methoxybenzene (15.5 g, 55.3 mmol) at 0°C. The resulting suspension was stirred at room temperature for 30 min. Excess hydride was quenched by the dropwise addition of water (200 mL), and the mixture was extracted with EtOAc (3×500 mL). The combined organics were washed with brine (2×300 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, gradient elution from 100% hexanes to 5% EtOAc/hexanes) to provide 1-((allyloxy)methyl)-2-bromo-4-methoxy benzene **10.1** (12.9 g, 91%) as a yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.42 (d, *J* = 8.7 Hz, 1H), 7.10 (d, *J* = 3.3 Hz, 1H), 6.72 (dd, *J* = 8.7, 3.3 Hz, 1H), 6.05-5.94 (m, 1H), 5.40 (ddd, *J* = 17.1, 3.3, 1.5 Hz, 1H), 5.25 (dd, *J* = 10.5, 1.5 Hz, 1H), 4.57 (s, 2H), 4.14 (m, 2H), 3.82 (s, 3H).

Step 2. A solution of compound **10.1** (12.9 g, 50.1 mmol) in DMF (400 mL) was added Cs₂CO₃ (19.5 g, 60.1 mmol), triphenylphosphine (5.90 g, 22.5 mmol) and Pd(OAc)₂ (1.68 g, 7.51 mmol). The reaction mixture was heated to 90°C for 1 h. Water (300 mL) and EtOAc (300 mL) were added to the reaction vessel and the resulting biphasic mixture was transferred to a separatory funnel. The layers were separated and the aqueous phase was extracted with EtOAc (2×500 mL). The organic phase was washed with brine (3×500 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, gradient elution from 100% hexanes to 5% EtOAc/hexanes) to afford 6-methoxy-4-

methylene isochroman **10.2** (5.70 g, 64%) as a yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 7.64 (d, J = 8.7 Hz, 1H), 6.82 (dd, J = 7.2, 2.7 Hz, 1H), 6.56 (d, J = 2.7 Hz, 1H), 5.48 (s, 1H), 4.91 (s, 1H), 4.80 (s, 2H), 4.44 (s, 2H), 3.82 (s, 3H).

Step 3. NaIO_4 (20.7 g, 96.8 mmol) and OsO_4 (81.8 mg, 322 μmol) were added to a solution of **12.2** (5.7 g, 32.3 mmol) in THF (200 mL) and water (100 mL), and the reaction mixture was stirred at room temperature for 4 h. The mixture was diluted with water (200 mL) and extracted with EtOAc (3 \times 200 mL). The combined organics were dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO_2 , gradient elution from 5% EtOAc/hexanes to 10% EtOAc/hexanes) to provide 6-methoxyisochroman-4-one **10.3** (3.10 g, 54%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 7.53 (s, 1H), 7.27 (d, J = 6.6 Hz, 1H), 7.16 (d, J = 1.5 Hz, 1H), 4.87 (s, 2H), 4.37 (s, 2H), 3.88 (s, 3H).

Step 4. 6-Methoxyisochroman-4-ol **10.4** (3.00 g, 96%) was prepared as described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.03-6.98 (m, 2H), 6.88 (m, 1H), 4.79 (A of AB, J_{AB} = 14.7 1H), 4.66 (B of AB, J_{BA} = 14.7 1H), 4.54 (m, 1H), 4.12 (A of ABX, J_{AB} = 12.0 Hz, J_{AX} = 2.7 Hz, 1H), 3.88 (B of ABX, J_{BA} = 12.0 Hz, J_{BX} = 2.7 Hz, 1H), 3.80 (s, 3H), 2.23 (d, J = 9.6 Hz, 1H).


Step 5. 4-Azido-6-methoxyisochroman **10.5** (2.70 g, 79%) was prepared as described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.01 (d, J = 8.1 Hz, 1H), 6.94-6.90 (m, 2H), 4.84 (A of AB, J_{AB} = 14.7 Hz, 1H), 4.69 (B of AB, J_{BA} = 14.7 Hz, 1H), 4.33-3.97 (m, 2H), 3.92 (dd, J = 12.9, 2.4 Hz, 1H), 3.85 (s, 3H).

Step 6. *tert*-Butyl (6-methoxyisochroman-4-yl)carbamate **10.6** (2.49 g, 68%) was prepared as described in Example 1. LC-MS (ESI): m/z 316 [M+H].

Step 7. The compound of **Example 10** (90.0 mg, 94%) was prepared as described in Example 1. MS (ESI): m/z 180 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.50 (s, 3H), 7.16 (d, J = 2.1 Hz, 1H), 7.10 (d, J = 8.4 Hz, 1H), 6.97 (dd, J = 8.4, 2.4 Hz, 1H), 4.76 (d, J = 14.7 Hz, 1H), 4.64 (d, J = 15.0 Hz, 1H), 4.31 (s, 1H), 4.12 (d, J = 12.9 Hz, 1H), 3.88 (dd, J = 12.3, 2.7 Hz, 1H), 3.77 (s, 3H).

[0133] **Table 2. Compounds prepared as described in Example 10.**

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 11 | | MS (ESI): <i>m/z</i> 206 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.53 (br s, 3H), 7.60 (s, 1H), 7.41 (dd, <i>J</i> = 8.1, 1.8 Hz, 1H), 7.09 (d, <i>J</i> = 8.1 Hz, 1H), 4.79 (d, <i>J</i> = 15.0 Hz, 1H), 4.66 (d, <i>J</i> = 15.3 Hz, 1H), 4.31 (s, 1H), 4.15 (d, <i>J</i> = 12.3 Hz, 1H), 3.89 (dd, <i>J</i> = 12.6, 2.7 Hz, 1H), 1.29 (s, 9H). |
| 12 | | MS (ESI): <i>m/z</i> 150 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.48-7.34 (m, 3H), 7.20 (d, <i>J</i> = 7.5 Hz, 1H), 4.95 (d, <i>J</i> = 15.3 Hz, 1H), 4.81 (d, <i>J</i> = 15.6 Hz, 1H), 4.38 (s, 1H), 4.26 (dd, <i>J</i> = 12.9, 1.5 Hz, 1H), 3.99 (dd, <i>J</i> = 10.5, 2.1 Hz, 1H). |
| 13 | | MS (ESI): <i>m/z</i> 200 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.46 (br s, 3H), 8.17 (d, <i>J</i> = 8.4 Hz, 1H), 7.99 (d, <i>J</i> = 8.4 Hz, 2H), 7.65-7.60 (m, 2H), 7.31 (d, <i>J</i> = 8.4 Hz, 1H), 5.05-4.88 (m, 3H), 4.38 (d, <i>J</i> = 12.3 Hz, 1H), 3.96 (dd, <i>J</i> = 12.6, 1.8 Hz, 1H). |
| 14 | | MS (ESI): <i>m/z</i> 180 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.36 (d, <i>J</i> = 8.4 Hz, 1H), 6.91 (dd, <i>J</i> = 8.4, 2.7 Hz, 1H), 6.75 (d, <i>J</i> = 2.1 Hz, 1H), 4.90 (d, <i>J</i> = 15.6 Hz, 1H), 4.75 (d, <i>J</i> = 15.6 Hz, 1H), 4.30 (s, 1H), 4.22 (dd, <i>J</i> = 12.6, 1.2 Hz, 1H), 3.94 (dd, <i>J</i> = 12.6, 2.1 Hz, 1H), 3.82 (s, 3H). |
| 15 | | MS (ESI): <i>m/z</i> 201 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 9.43 (d, <i>J</i> = 8.7 Hz, 1H), 9.29 (d, <i>J</i> = 5.1 Hz, 1H), 8.32 (d, <i>J</i> = 9.0 Hz, 1H), 8.22 (dd, <i>J</i> = 8.7, 5.1 Hz, 1H), 7.99 (d, <i>J</i> = 9.0 Hz, 1H), 5.25 (d, <i>J</i> = 17.7 Hz, 2H), 5.11 (t, <i>J</i> = 17.7 Hz, 1H), 4.50 (d, <i>J</i> = 12.9 Hz, 1H), 4.12 (d, <i>J</i> = 14.7 Hz, 1H). |
| 16 | | MS (ESI): <i>m/z</i> 221 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.04 (d, <i>J</i> = 8.4 Hz, 1H), 7.22 (d, <i>J</i> = 8.4 Hz, 1H), 4.95 (d, <i>J</i> = 15.6 Hz, 1H), 4.79 (d, <i>J</i> = 16.2 Hz, 2H), 4.23 (d, <i>J</i> = 12.3 Hz, 1H), 4.00 (dd, <i>J</i> = 12.6, 2.4 Hz, 1H), 2.85 (s, 3H). |

[0134] **Example 17: 8-Ethylisochroman-4-amine hydrochloride**


Step 1. *tert*-Butyl (8-vinylisochroman-4-yl)carbamate **17.2** was prepared from

compound **17.1** (prepared from 1,3-dibromo-2-(bromomethyl)benzene as previously described in Example 10) as previously described in Example 7 using vinylboronic acid pinacol ester. ^1H NMR (300 MHz, CDCl_3) δ 7.40-7.35 (m, 2H), 7.27-7.22 (m, 1H), 6.60 (dd, J = 18.0, 12.0 Hz, 1H), 5.68 (d, J = 18.0 Hz, 1H), 5.36-5.30 (m, 1H), 5.11-5.08 (m, 1H), 4.90 (d, J = 15.0 Hz, 1H), 4.75-4.67 (m, 2H), 4.03 (dd, J = 12.0, 3.0 Hz, 1H), 3.83 (dd, J = 12.0, 3.0 Hz, 1H), 1.46 (s, 9H).

Step 2. *tert*-Butyl (8-ethylisochroman-4-yl)carbamate **17.3** was prepared as previously described in Example 8. ^1H NMR (300 MHz, CDCl_3) δ 7.29-7.19 (m, 2H), 7.13-7.10 (m, 1H), 5.10-5.07 (m, 1H), 4.86 (d, J = 15.0 Hz, 1H), 4.77-4.71 (m, 2H), 4.04 (dd, J = 12.0, 3.0 Hz, 1H), 3.86 (dd, J = 12.0, 3.0 Hz, 1H), 2.51-2.44 (m, 2H), 1.46 (s, 9H), 1.20 (t, J = 9.0 Hz, 3H).

Step 3. The compound of **Example 17** was prepared as previously described in Example 1. MS (ESI): m/z 178 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.31-7.27 (m, 3H), 5.00 (d, J = 18.0 Hz, 1H), 4.76 (d, J = 15.0 Hz, 1H), 4.33 (s, 1H), 4.20 (dd, J = 15.0, 3 Hz, 1H), 3.92 (dd, J = 12.0, 3 Hz, 1H), 2.58-2.50 (m, 2H), 1.21 (t, J = 9.0 Hz, 3H).

[0135] **Example 18: 8-Isopropylisochroman-4-amine hydrochloride**




The compound of **Example 18** was prepared as described in Example 17 using isopropenylboronic acid pinacol ester. MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.41-7.31 (m, 2H), 7.26-7.23 (m, 1H), 5.08 (d, J = 18 Hz, 1H), 4.81 (d, J = 21 Hz, 1H), 4.32 (s, 1H), 4.19 (dd, J = 12.6, 1.2 Hz, 1H), 3.92 (dd, J = 12.9, 2.1 Hz, 1H), 2.86-2.48 (m, 1H), 1.26-1.20 (m, 6H).

[0136] **Example 19: 2,4,7,8,9,10-Hexahydro-1H-benzo[f]isochromen-1-amine hydrochloride**



The compound of **Example 19** was prepared as described in Example 10 using 5-bromo-6-(bromomethyl)-1,2,3,4-tetrahydronaphthalene **19.6**. MS (ESI): *m/z* 204 [M+H]. ¹H NMR (300 MHz, DMSO-*d*6) δ 8.38 (s, 3H), 7.09 (d, *J* = 7.8 Hz, 1H), 6.89 (d, *J* = 7.8 Hz, 1H), 4.82 (d, *J* = 15.6 Hz, 1H), 4.67 (d, *J* = 15.6 Hz, 1H), 4.36 (d, *J* = 3.9 Hz, 1H), 4.23 (d, *J* = 12.3 Hz, 1H), 3.75 (d, *J* = 12.0 Hz, 1H), 2.92-2.84 (m, 1H), 2.73-2.65 (m, 3H), 1.83-1.60 (m, 4H).

[0137] **Synthesis of 5-bromo-6-(bromomethyl)-1,2,3,4-tetrahydronaphthalene (19.6)**



Step 1. Nitric acid (10.7 mL, 240 mmol) was added to a solution of 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (10.0 g, 56.7 mmol) in AcOH (70 mL) and concentrated H₂SO₄ (70 mL). The reaction was stirred at ambient temperature for 1 h and then poured onto ice. The resulting precipitate was collected by filtration to provide 1-nitro-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid **19.1** (11.2 g, 89%) as a yellow solid. MS (ESI-): *m/z* 220 [M-H]. ¹H NMR (300 MHz, DMSO-*d*6) δ 7.75 (d, *J* = 8.1 Hz, 1H), 7.41 (d, *J* = 8.1 Hz, 1H), 2.88-2.81 (m, 2H), 2.69-2.63 (m, 2H), 1.74 (m, 4H).

Step 2. A solution of compound **19.1** (5.2 g, 11.7 mmol) in CH₃CN (100 mL) was treated with K₂CO₃ (4.85 g, 35.1 mmol) and iodomethane (4.98 g, 35.1 mmol). The reaction was stirred at room temperature for 3 h. The mixture was then partitioned between water (100 mL) and EtOAc (600 mL), and the organic phase was washed with brine (2 \times 100 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, 100% hexanes) to provide methyl 1-nitro-5,6,7,8-tetrahydronaphthalene-2-carboxylate **19.2** (2.24 g, 81%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 7.77 (d, *J* = 8.1 Hz, 1H), 7.26 (d, *J* = 8.4 Hz, 1H), 3.89 (s, 3H), 2.87-2.85 (m, 2H), 2.72-2.66 (m, 2H), 1.83-1.81 (m, 4H).


Step 3. Compound **19.2** (3.2 g, 9.52 mmol) was dissolved in THF (100 mL). Pd/C (1 g, 10%) was added and the mixture was stirred at room temperature under a H₂ atmosphere for 8 h. The mixture was filtered through a Celite pad, and the filtrate was concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 100% hexanes) gave methyl 1-amino-5,6,7,8-tetrahydronaphthalene-2-carboxylate **19.3** (1.80 g, 92%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 7.65 (d, *J* = 8.4 Hz, 1H), 6.42 (d, *J* = 8.4 Hz, 1H), 5.88 (brs, 2H), 3.85 (s, 3H), 2.72 (t, *J* = 6.3 Hz, 2H), 2.42 (t, *J* = 6.0 Hz, 2H), 1.92-1.84 (m, 2H), 1.78-1.71 (m, 2H).

Step 4. A solution of **19.3** (4.3 g, 20.9 mmol) in CH₃CN (200 mL) was cooled in an ice bath, treated with copper(I) bromide (5.99 g, 41.8 mmol), and stirred for 30 min. tert-Butyl nitrite (4.31 g, 41.8 mmol) was added, and the ice bath was removed. The flask was heated to 40 °C and stirred overnight. The mixture was cooled in an ice bath and diluted with NH₄OH and EtOAc, and the resulting biphasic mixture was separated. The aqueous phase was extracted with EtOAc (3×100 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 5% EtOAc/hexanes) gave methyl 1-bromo-5,6,7,8-tetrahydronaphthalene-2-carboxylate **19.4** (3.60 g, 63%) as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 7.40 (d, *J* = 7.8 Hz, 1H), 7.06 (d, *J* = 7.8 Hz, 1H), 3.91 (s, 3H), 2.80 (t, *J* = 8.1 Hz, 4H), 1.86-1.72 (m, 4H).

Step 5. Compound **19.4** (3.6 g, 13.3 mmol) was dissolved in anhydrous THF (100 mL) under a N₂ atmosphere and cooled to -78 °C in a dry ice/acetone bath. Diisobutylaluminum hydride (17.7 mL, 26.6 mmol, 1.5M in toluene) was added in a dropwise manner. Once addition was complete, the cold bath was removed and the reaction mixture was allowed to warm to room temperature for 1 h. Excess hydride was carefully quenched by the dropwise addition of aqueous 3M HCl (50 mL) and the biphasic mixture was transferred to a separatory funnel. The aqueous phase was washed with EtOAc (3×150 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 10% EtOAc/hexanes) gave (1-bromo-5,6,7,8-tetrahydronaphthalen-2-yl)methanol **19.5** (3.0 g, 94%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 7.20 (d, *J* = 7.8 Hz, 1H), 7.06 (d, *J* = 7.8 Hz, 1H), 4.73 (d, *J* = 6.6 Hz, 2H), 2.80-2.75 (m, 4H), 2.02 (t, *J* = 6.6 Hz, 1H), 1.88-1.71 (m, 4H).


Step 6. Compound **19.5** (3.0 g, 12.4 mmol) was dissolved in 1,2-dichloroethane (120 mL) and cooled in an ice bath. Phosphorous tribromide (0.59 mL, 6.20 mmol) was added and the mixture was stirred at 0°C for 30 min. The reaction poured into ice water and extracted with CH₂Cl₂ (3×50 mL). The combined organics were washed with saturated aqueous Na₂CO₃ (100 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to afford 5-bromo-6-(bromomethyl)-1,2,3,4-tetrahydronaphthalene **19.6** (1.70 g, 45%). ¹H NMR (300 MHz, CDCl₃) δ 7.10 (d, *J* = 7.8 Hz, 1H), 7.02 (d, *J* = 7.5 Hz, 1H), 4.65 (s, 2H), 2.80-2.71 (m, 4H), 1.87-1.70 (m, 4H).

[0138] **Example 20: 6-Ethylisochroman-4-amine hydrochloride**



The compound of **Example 20** was prepared as described in Example 19 using methyl 2-amino-4-ethylbenzoate. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, DMSO-*d*6) δ 8.44 (s, 3H), 7.35 (s, 1H), 7.23 (d, *J* = 7.8 Hz, 1H), 7.09 (d, *J* = 8.1 Hz, 1H), 4.84 (d, *J* = 11.7 Hz, 1H), 4.70 (d, *J* = 15.3 Hz, 1H), 4.32 (s, 1H), 4.16 (dd, *J* = 12.6, 1.8 Hz, 1H), 3.90 (dd, *J* = 12.6, 2.7 Hz, 1H), 2.64-2.57 (m, 2H), 1.24-1.09 (m, 3H).

[0139] **Example 21: 5-Methylisochroman-4-amine hydrochloride**



Step 1. Sodium hydride (771 mg, 19.3 mmol, 95%) was added in small portions to a solution of (2-bromo-3-methylphenyl)methanol (2.6 g, 12.9 mmol) in DMF (35 mL) cooled in an ice bath. The resulting slurry was stirred for 15 min, followed by addition of 3-bromoprop-1-ene (1.70 g, 14.1 mmol). The reaction was stirred at 0°C for 30 min. Excess hydride was quenched by the dropwise addition of saturated aqueous NH₄Cl (75 mL), and the mixture was partitioned with Et₂O (3×40 mL). The combined organics were dried over

anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO_2 , 100% hexanes) to provide 1-((allyloxy)methyl)-2-bromo-3-methylbenzene **21.1** (2.30 g, 49%) as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 7.36-7.34 (m, 1H), 7.28-7.18 (m, 2H), 6.08-5.95 (m, 1H), 5.38 (d, J = 15.0 Hz, 1H), 5.25 (d, J = 10.5 Hz, 1H), 4.62 (s, 2H), 4.14 (d, J = 5.7 Hz, 2H), 2.44 (s, 3H).

Step 2. 5-Methyl-4-methyleneisochroman **21.2** was prepared as previously described in Example 10. ^1H NMR (300 MHz, CDCl_3) δ 7.19-7.15 (m, 2H), 6.96-6.92 (m, 1H), 5.47 (s, 1H), 5.37 (s, 1H), 4.81 (s, 2H), 4.41 (s, 2H), 2.53 (s, 3H).

Step 3. 5-Methylisochroman-4-one **21.3** was prepared as previously described in Example 10. ^1H NMR (300 MHz, CDCl_3) δ 7.44 (t, J = 7.5 Hz, 1H), 7.21 (d, J = 7.5 Hz, 1H), 7.08 (d, J = 7.8 Hz, 1H), 4.88 (s, 2H), 4.35 (s, 2H), 2.70 (s, 3H).



Step 4. 5-Methylisochroman-4-ol **21.4** was prepared as previously described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.24-7.19 (m, 1H), 7.13 (d, J = 7.5 Hz, 1H), 6.88 (d, J = 7.5 Hz, 1H), 4.83 (d, J = 15.0 Hz, 1H), 4.70 (d, J = 15.0 Hz, 1H), 4.60 (d, J = 10.2 Hz, 1H), 4.29 (d, J = 12.0 Hz, 1H), 3.81 (dd, J = 12.0, 3.3 Hz, 1H), 2.48 (s, 3H), 2.23 (dd, J = 13.5, 7.5 Hz, 1H).

Step 5. 4-Azido-5-methylisochroman **21.5** was prepared as previously described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.26-7.24 (m, 1H), 7.18 (d, J = 7.2 Hz, 1H), 6.94 (d, J = 7.2 Hz, 1H), 4.91 (d, J = 15.3 Hz, 1H), 4.74 (d, J = 15.3 Hz, 1H), 4.28 (d, J = 12.0 Hz, 1H), 4.08 (s, 1H), 3.92 (dd, J = 12.3, 2.1 Hz, 1H), 2.42 (s, 3H).


Step 6. The compound of **Example 21** was prepared as previously described in Example 1. MS (ESI): m/z 164 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.26 (s, 3H), 7.29 (t, J = 7.5 Hz, 1H), 7.15 (d, J = 7.2 Hz, 1H), 6.99 (d, J = 7.8 Hz, 1H), 4.87 (d, J = 15.6 Hz, 1H), 4.72 (d, J = 15.6 Hz, 1H), 4.43 (s, 1H), 4.21 (d, J = 12.3 Hz, 1H), 3.80 (dd, J = 12.3, 1.2 Hz, 1H), 2.39 (s, 3H).

[0140] **Table 3. Compounds prepared as described in Example 21 using the appropriately substituted starting material.**

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 22 | | MS (ESI): <i>m/z</i> 168 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.52-7.44 (m, 1H), 7.14 (t, <i>J</i> = 9.0 Hz, 1H), 7.06 (d, <i>J</i> = 7.8 Hz, 1H), 4.97 (d, <i>J</i> = 15.9 Hz, 1H), 4.80 (d, <i>J</i> = 15.9 Hz, 1H), 4.58 (s, 1H), 4.28 (d, <i>J</i> = 12.9 Hz, 1H), 3.95 (dd, <i>J</i> = 13.2, 2.4 Hz, 1H). |
| 23 | | MS (ESI): <i>m/z</i> 168 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.46-7.39 (m, 1H), 7.30 (d, <i>J</i> = 7.8 Hz, 1H), 7.20 (t, <i>J</i> = 9.3 Hz, 1H), 5.05 (d, <i>J</i> = 15.9 Hz, 1H), 4.77 (d, <i>J</i> = 15.9 Hz, 1H), 4.41 (s, 1H), 4.24 (d, <i>J</i> = 12.0 Hz, 1H), 3.97 (dd, <i>J</i> = 12.9, 2.1 Hz, 1H). |
| 24 | | MS (ESI): <i>m/z</i> 180 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.37 (t, <i>J</i> = 6.0 Hz, 1H), 7.05 (d, <i>J</i> = 6.3 Hz, 1H), 4.97 (d, <i>J</i> = 12.3 Hz, 1H), 4.65 (d, <i>J</i> = 12.0 Hz, 1H), 4.34 (s, 1H), 4.21 (dd, <i>J</i> = 9.6, 0.9 Hz, 1H), 3.93 (dd, <i>J</i> = 9.6, 1.5 Hz, 1H), 3.88 (s, 3H). |
| 25 | | MS (ESI): <i>m/z</i> 180 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.38 (t, <i>J</i> = 7.8 Hz, 1H), 6.96 (d, <i>J</i> = 7.5 Hz, 1H), 6.77 (d, <i>J</i> = 7.8 Hz, 1H), 4.89 (d, <i>J</i> = 15.0 Hz, 1H), 4.72 (d, <i>J</i> = 15.3 Hz, 1H), 4.44 (d, <i>J</i> = 1.2 Hz, 1H), 4.25 (d, <i>J</i> = 12.9 Hz, 1H), 3.93 (s, 3H), 3.90 (dd, <i>J</i> = 13.5, 3.0 Hz, 1H). |
| 26 | | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> 6) δ 8.69 (brs, 3H), 7.32 (s, 1H), 7.21 (d, <i>J</i> = 7.8 Hz, 1H), 7.07 (d, <i>J</i> = 7.8 Hz, 1H), 4.80 (d, <i>J</i> = 15.3 Hz, 1H), 4.67 (d, <i>J</i> = 15.3 Hz, 1H), 4.29 (s, 1H), 4.13 (dd, <i>J</i> = 12.6, 1.8 Hz, 1H), 3.88 (dd, <i>J</i> = 12.6, 2.7 Hz, 1H), 2.30 (s, 3H). |
| 27 | | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.29-7.25 (m, 3H), 4.94 (d, <i>J</i> = 17.7 Hz, 1H), 4.71 (d, <i>J</i> = 15.9 Hz, 1H), 4.35 (s, 1H), 4.23 (d, <i>J</i> = 12.6 Hz, 1H), 3.95 (dd, <i>J</i> = 12.6, 1.8 Hz, 1H), 2.21 (s, 3H). |
| 28 | | MS (ESI): <i>m/z</i> 184 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> 6) δ 8.49 (s, 3H), 7.48-7.42 (m, 2H), 7.20 (d, <i>J</i> = 6.3 Hz, 1H), 4.89 (d, <i>J</i> = 15.9 Hz, 1H), 4.71 (d, <i>J</i> = 15.6 Hz, 1H), 4.48 (s, 1H), 4.28 (d, <i>J</i> = 12.6 Hz, 1H), 3.76 (d, <i>J</i> = 12.6 Hz, 1H). |
| 29 | | MS (ESI): M/Z 178 [M+H]. ¹ H NMR (300 MHz, CDCl ₃) δ 8.78 (brs, 3 H), 7.61 (d, <i>J</i> = 7.5 Hz, 1H), 7.35 (t, <i>J</i> = 7.5 Hz, 1H), 7.26 - 7.21 (m, 1H), 7.15 (d, <i>J</i> = 7.8 Hz, 1H), 4.36 (s, 1H), 4.29 (d, <i>J</i> = 12 Hz, 1H), 4.04 (d, <i>J</i> = 12 Hz, 1H), 1.78 (s, 3H), 1.50 (s, 3H). |

| Example | Structure | Characterization Data |
|---------|---|---|
| 30 |  | MS (ESI): m/z 164 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 8.59 (brs, 3H), 7.36 - 7.23 (m, 4H), 4.41 (s, 1H), 4.12 - 4.00 (m, 2H), 3.74 (d, J = 12.6 Hz, 1H), 3.60 - 3.49 (m, 2H), 2.89 - 2.80 (m, 1H). |
| 32 |  | MS (ESI): m/z 214 [M+H]. ^1H NMR (300 MHz, CD3OD) δ 8.23 (d, J = 8.7 Hz, 1H), 7.94 (t, J = 8.4 Hz, 2H), 7.66 (t, J = 7.2 Hz, 1H), 7.56 (t, J = 7.5 Hz, 1H), 7.41 (d, J = 8.4 Hz, 1H), 5.52 (d, J = 4.2 Hz, 1H), 4.46 - 4.31 (m, 2H), 3.90 (d, J = 14.4 Hz, 1H), 3.84 - 3.70 (m, 2H), 3.10 - 3.04 (m, 1H). |

[0141] **Example 31: 8-Methyl-1,2,4,5-tetrahydrobenzo[d]oxepin-1-amine hydrochloride**




The compound of **Example 31** was prepared as previously described in Example 21 using 2-(2-bromo-4-methylphenyl)ethanol. MS (ESI): m/z 178 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 7.71 (s, 1H), 7.08 - 7.01 (m, 2H), 4.48 (d, J = 11.4 Hz, 1H), 4.37 (s, 1H), 4.16 (d, J = 12.6 Hz, 1H), 3.77 - 3.72 (m, 2H), 3.54 (t, J = 11.7 Hz, 1H), 2.77 - 2.70 (m, 1H), 1.61 (s, 3H).

[0142] Synthesis of 2-(2-bromo-4-methylphenyl)ethanol



Step 1. A solution of 2-(2-bromo-4-methylphenyl)acetic acid(7.2 g, 31.43 mmol) in THF (314 mL) was cooled in an ice bath to 0°C, and treated with $\text{BH}_3 \cdot \text{THF}$ (62.86 mL, 62.86 mmol, 1M in THF) in a dropwise manner. After addition was complete, the reaction mixture was stirred at 0°C overnight. Excess borane was quenched by the dropwise addition of 1M HCl (20 mL) at 0°C. The mixture was concentrated under reduced pressure and then partitioned between EtOAc (300 mL) and water (200 mL). The organic layer was washed with brine (200mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated to give the 2-(2-bromo-4-methylphenyl)ethanol (6.7 g, 94%) as a yellow oil. MS (ESI) m/z 214, 216 [M+H].

[0143] Example 33: 1,3,4,5-Tetrahydrobenzo[c]oxepin-5-amine hydrochloride




The compound of **Example 33** was prepared as previously described in Example 21 using 4-bromobut-1-ene in place of 3-bromoprop-1-ene. MS (ESI): m/z 164 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.43 - 7.29 (m, 4H), 4.90 - 4.76 (m, 3H), 4.22 - 4.14 (m, 1H), 4.09 - 4.03 (m, 1H), 2.28 - 2.20 (m, 2H)

[0144] Table 4. Compounds prepared as described in Example 33 using the appropriately substituted 2-bromobenzyl alcohol.

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 34 | | MS (ESI+): m/z 178 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.18 - 7.13 (m, 3H), 4.88 - 4.71 (m, 3H), 4.20 - 4.15 (m, 1H), 4.07 - 4.00 (m, 1H), 2.40 (s, 3H), 2.24 - 2.16 (m, 2H). |
| 35 | | MS (ESI+): m/z 214 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 8.26 (d, J = 8.7 Hz, 1H), 7.95 (t, J = 6.6 Hz, 2H), 7.70 (t, J = 7.2 Hz, 1H), 7.59 (t, J = 7.8 Hz, 1H), 7.35 (d, J = 8.7 Hz, 1H), 5.73 - 5.70 (m, 1H), 5.21 (d, J = 16.5 Hz, 1H), 5.05 (d, J = 16.2 Hz, 1H), 4.27 - 4.07 (m, 2H), 2.63 - 2.57 (m, 2H). |
| 36 | | MS (ESI+): m/z 178 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.21 (s, 2H), 7.12 (s, 1H), 4.85 - 4.72 (m, 3H), 4.20 - 4.12 (m, 1H), 4.07 - 3.99 (m, 1H), 2.35 (s, 3H), 2.32 - 2.20 (m, 2H). |
| 37 | | MS (ESI+): m/z 178 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.22-7.13 (m, 3H), 5.12 (d, J = 15 Hz, 1H), 4.67 - 4.62 (m, 2H), 4.16 - 4.07 (m, 1H), 3.99 - 3.90 (m, 1H), 2.30 - 2.25 (m, 5H). |
| 38 | | MS (ESI+): m/z 198 [M+H]. ^1H NMR (300 MHz, DMSO-d_6) δ 8.85 (brs, 3H), 7.48 - 7.43 (m, 2H), 7.34 - 7.30 (m, 1H), 4.78 - 4.65 (m, 3H), 4.11 - 4.06 (m, 1H), 3.97 - 3.89 (m, 1H), 2.13 - 1.91 (m, 2H). |
| 39 | | MS (ESI+): m/z 178 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 8.79 (brs, 3H), 7.28 - 7.13 (m, 2H), 6.95 - 7.00 (m, 1H), 5.12 - 5.00 (m, 2H), 4.70 (d, J = 15.3 Hz, 1H), 4.31 - 4.21 (m, 1H), 4.01 - 3.95 (m, 1H), 2.70 - 2.51 (m, 4H), 2.31 - 2.20 (m, 1H). |


| Example | Structure | Characterization Data |
|---------|-----------|---|
| 40 | | MS (ESI+): m/z 198 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.84 (brs, 3H), 7.46 (d, <i>J</i> = 6.0 Hz, 1H), 7.40 (d, <i>J</i> = 5.7 Hz, 1H), 7.33 (d, <i>J</i> = 5.7 Hz, 1H), 5.17 (d, <i>J</i> = 11.1 Hz, 1H), 4.85 (d, <i>J</i> = 5.7 Hz, 1H), 4.74 (d, <i>J</i> = 11.1 Hz, 1H), 4.11 - 4.06 (m, 1H), 3.99 - 3.93 (m, 1H), 2.25 - 2.15 (m, 1H), 2.10 - 2.00 (m, 1H). |
| 41 | | MS (ESI+): m/z 198 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.47 (d, <i>J</i> = 8.1 Hz, 1H), 7.36 (t, <i>J</i> = 7.5 Hz, 1H), 7.22 (d, <i>J</i> = 7.8 Hz, 1H), 5.26 (d, <i>J</i> = 5.4 Hz, 1H), 5.01 (d, <i>J</i> = 16.2 Hz, 1H), 4.87 (d, <i>J</i> = 16.2 Hz, 1H), 4.15 - 3.98 (m, 2H), 2.52 - 2.37 (m, 2H). |

[0145] **Example 42: 5,6-Dimethylisochroman-4-amine hydrochloride**



The compound of **Example 42** was prepared as previously described in Example 21, using (2-bromo-3,4-dimethylphenyl)methanol (prepared from methyl 2-amino-3,4-dimethylbenzoate as previously described in Example 19). MS (ESI): m/z 178 [M+H]. ^1H NMR (300 MHz, DMSO-*d*₆) δ 8.32 (s, 3H), 7.20 (d, *J* = 7.8 Hz, 1H), 6.90 (d, *J* = 7.8 Hz, 1H), 4.83 (d, *J* = 15.3 Hz, 1H), 4.68 (d, *J* = 15.3 Hz, 1H), 4.48 (s, 1H), 4.24 (d, *J* = 12.8 Hz, 1H), 3.76 (d, *J* = 11.1 Hz, 1H), 2.25 (s, 6H).

[0146] **Example 43: 6,7-Dimethyl-1,3,4,5-tetrahydrobenzo[c]oxepin-5-amine hydrochloride**




The compound of **Example 43** was prepared as previously described in Example 42 using but-3-en-1-ol. MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, DMSO-*d*₆) δ 8.44 (brs, 3H), 7.13 (d, *J* = 7.5 Hz, 1H), 6.97 (d, *J* = 7.8 Hz, 1H), 5.00 (d, *J* = 15.0 Hz, 1H), 4.94 (s, 1H), 4.60 (d, *J* = 15.0 Hz, 1H), 4.11 - 4.01 (m, 1H), 3.95 - 3.90 (m, 1H), 2.33 (s, 3H), 2.27 (s, 3H), 2.19 - 2.09 (m, 2H).

[0147] Example 44: 5,6-Dichloroisochroman-4-amine hydrochloride



The compound of **Example 44** was prepared as previously described in Example 10, using (2-bromo-3,4-dichlorophenyl)methanol **44.5** prepared as described below. MS (ESI): *m/z* 218 [M+H]. ^1H NMR (300 MHz, DMSO-*d*6) δ 8.62 (s, 3H), 7.72 (d, *J* = 7.8 Hz, 1H), 7.26 (d, *J* = 8.4 Hz, 1H), 4.90 (d, *J* = 16.2 Hz, 1H), 4.69 (d, *J* = 16.2 Hz, 1H), 4.52 (s, 1H), 4.31 (d, *J* = 12.6 Hz, 1H), 3.82 (d, *J* = 14.7 Hz, 1H).

[0148] Synthesis of (2-bromo-3,4-dichlorophenyl)methanol (44.5)



Step 1. 2-Amino-3,4-dichlorobenzoic acid **44.1** (13.6 g, 66.0 mmol, *J. Med. Chem.* 1991, 34, 218) was dissolved in DMF (150 mL) and treated with K_2CO_3 (45.6 g, 330 mmol) and iodomethane (11.2 g, 79.2 mmol). The resulting slurry was stirred at room temperature for 2 hr. The mixture was diluted with water and extracted with EtOAc (3×100 mL). The combined organics were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 , gradient elution from 100% hexanes to 10% EtOAc/hexanes) afforded methyl 2-amino-3,4-dichlorobenzoate **44.2** (12.5 g, 86%) as a yellow solid. ^1H NMR (300 MHz, CDCl_3) δ 7.74 (d, *J* = 8.7 Hz, 1H), 6.74 (d, *J* = 8.7 Hz, 1H), 6.47 (br s, 2H), 3.89 (s, 3H).

Step 2. Methyl 2-bromo-3,4-dichlorobenzoate **44.3** was prepared as previously described in Example 19. ^1H NMR (300 MHz, CDCl_3) δ 7.54 (d, *J* = 9.0 Hz, 1H), 7.49 (d, *J* = 9.0 Hz, 1H), 3.94 (s, 3H).

Step 3. Compound **44.3** (11.1 g, 39.0 mmol) was dissolved in MeOH (250 mL) and treated with aqueous 2 N NaOH (39.2 mL, 78 mmol), and stirred at room temperature for 2 hr. The mixture was filtered, diluted with water (800 mL), and made acidic to pH =2 by the addition of aqueous 3M HCl. The resulting suspension was extracted with EtOAc (4×400 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to yield 2-bromo-3,4-dichlorobenzoic acid **44.4** (9.85 g, 94%), which was used directly without further purification.

Step 4. 2-Bromo-3,4-dichlorobenzyl alcohol **44.5** was prepared as previously described in Example 31. ¹H NMR (300 MHz, CDCl₃) δ 7.47 (d, *J* = 8.4 Hz, 1H), 7.39 (d, *J* = 8.4 Hz, 1H), 4.75 (s, 2H).

[0149] **Example 45: 5-Chloro-6-methylisochroman-4-amine hydrochloride**



The compound of **Example 45** was prepared as previously described in Example 44 from 2-amino-3-chloro-4-methylbenzoic acid (Rewcastle, G. W., *et al* *J. Med. Chem.* 1991, 34, 217). MS (ESI): *m/z* 170 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.05 (d, *J* = 0.9 Hz, 1H), 5.00 (d, *J* = 15.3 Hz, 1H), 4.80 (d, *J* = 15.3 Hz, 1H), 4.33 (s, 1H), 4.25 (dd, *J* = 12.9, 1.5 Hz, 1H), 3.84 (dd, *J* = 12.6, 1.8 Hz, 1H), 2.28 (d, *J* = 1.2 Hz, 3H).

[0150] **Example 46: 6,8-Dimethylisochroman-4-amine hydrochloride**




The compound of **Example 46** was prepared as previously described in Example 44 from 2-amino-4,6-dimethylbenzoic acid. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.07 (s, 2H), 4.90 (d, *J* = 18 Hz, 1H), 4.65 (d, *J* = 18 Hz, 1H), 4.27 (s, 1H), 4.18 (dd, *J* = 12, 3 Hz, 1H), 3.90 (dd, *J* = 12, 3 Hz, 1H), 2.32 (s, 3H), 2.15 (s, 1H).

[0151] Example 47: 5,8-Dimethylisochroman-4-amine hydrochloride




The compound of **Example 47** was prepared as previously described in Example 44 from 2-amino-3,6-dimethylbenzoic acid. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.38 (s, 3H), 7.13-7.05 (m, 2H), 4.82 (d, *J* = 15 Hz, 1H), 4.62 (d, *J* = 15 Hz, 1H), 4.39 (s, 1H), 4.25 (d, *J* = 12 Hz, 1H), 3.74 (d, *J* = 12 Hz, 1H), 2.37 (s, 3H), 2.09 (s, 3H).

[0152] Example 48: 8-Chloro-5-methylisochroman-4-amine hydrochloride



The compound of **Example 48** was prepared as previously described in Example 10, using (2-bromo-6-chloro-3-methylphenyl)methanol **48.5** prepared as described below. MS (ESI): *m/z* 198 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.46 (s, 3H), 7.42 (d, *J* = 9.0 Hz, 1H), 7.23 (d, *J* = 9.0 Hz, 1H), 4.85 (d, *J* = 15 Hz, 1H), 4.65 (d, *J* = 15 Hz, 1H), 4.46 (s, 1H), 4.25 (d, *J* = 12 Hz, 1H), 3.79 (d, *J* = 12 Hz, 1H), 2.41 (s, 3H).

[0153] Synthesis of (2-bromo-6-chloro-3-methylphenyl)methanol (**48.5**)

Step 1. A solution of 5-chloro-2-methylaniline (15 g, 105 mmol) in water (750 mL) was treated with aqueous concentrated HCl (50 mL), hydroxylamine hydrochloride (36.4 g,

525 mmol), 2,2,2-trichloroethane-1,1-diol (25.9 g, 157 mmol) and Na₂SO₄ (14.9 g, 105 mmol), and heated to reflux for 2 h. After cooling to room temperature, the precipitate was collected by filtration, washed with water, and dried under vacuum to provide (E)-N-(5-chloro-2-methylphenyl)-2-(hydroxyimino)acetamide **48.1** (61.0 g) as a brown solid, which was used without further purification.

Step 2. Compound **48.1** (51 g, 239 mmol) was dissolved in methanesulfonic acid (350 mL) and heating at 50°C for 2 h. After cooling to room temperature, the mixture was poured into ice-water. The resulting suspension was collected on a fritted filter, washed with water, and dried under vacuum to provide 4-chloro-7-methylindoline-2,3-dione **48.2** (45.0 g, 96%) as a brown solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.22 (brs, 1H), 7.01 (dd, *J* = 7.8, 0.6 Hz, 1H), 6.58 (d, *J* = 8.1 Hz, 1H), 2.07 (s, 3H).


Step 3. Compound **48.2** (15 g, 76.6 mmol) was dissolved in aqueous 0.3M NaOH (1.5 L) and treated with the dropwise addition of 40% hydrogen peroxide (150 mL). The reaction was stirred at room temperature for 4 h. Residual solids were removed by filtration. The filtrate was washed with CH₂Cl₂ (3×200 mL), acidified to pH = 3-4 with aqueous concentrated HCl, and extracted with EtOAc (3×350 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to yield 2-amino-6-chloro-3-methylbenzoic acid **48.3** (12.7 g, 90%) as a yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.02 (d, *J* = 7.8 Hz, 1H), 6.58 (d, *J* = 7.8 Hz, 1H), 2.07 (s, 3H).

Step 4. Compound **48.3** (14.0 g, 75.4 mmol) was dissolved in EtOH (300 mL) and 48% hydrobromic acid (75 mL) and cooled in an ice bath. A solution of NaNO₂ (7.79 g, 113 mmol) in water (20 mL) was added in a dropwise manner, and the resulting mixture was stirred at 0°C for 20 min. Copper(I) bromide (10.8 g, 75.4 mmole) was added in a single portion, the ice bath was removed, and the suspension was heated with stirring to 95°C for 30 min. After cooling to room temperature, saturated aqueous NH₄Cl (200 mL) was added with vigorous stirring. The aqueous phase was extracted with EtOAc (3×200 mL). The combined organics were concentrated under reduced pressure. The residue was dissolved in water (30 mL) and made basic by the addition of aqueous 2N NaOH until pH = 10. The aqueous phase was washed with CH₂Cl₂ (3×200 mL), made acidic to pH = 3 by the addition of aqueous 5N HCl, and extracted with EtOAc (4×300 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to yield 2-bromo-6-

chloro-3-methylbenzoic acid **48.4**. ^1H NMR (300 MHz, DMSO-*d*₆) δ 7.40-7.49 (m, 2H), 2.36 (s, 3H).

Step 5. (2-Bromo-6-chloro-3-methylphenyl)methanol **48.5** was prepared as described previously in Example 44. ^1H NMR (300 MHz, DMSO-*d*₆) δ 7.39 (d, *J* = 8.4 Hz, 1H), 7.32 (d, *J* = 8.1 Hz, 1H), 4.75 (s, 2H), 2.36 (s, 3H).

[0154] **Example 49: 8-Chloro-6-methylisochroman-4-amine hydrochloride**




The compound of **Example 49** was prepared as previously described in Example 48 from 3-chloro-5-methylaniline. MS (ESI): *m/z* 198 [M+H]. ^1H NMR (300 MHz, CD₃OD) δ 7.32 (s, 1H), 7.24 (s, 1H), 4.95 (d, *J* = 15 Hz, 1H), 4.64 (d, *J* = 15 Hz, 1H), 4.33 (s, 1H), 4.21 (dd, *J* = 12.0, 3.0 Hz, 1H), 3.92 (dd, *J* = 12.0, 3.0 Hz, 1H), 2.37 (s, 3H).

[0155] **Example 50: 3,4-Dihydro-1H-benzo[h]isochromen-4-amine hydrochloride**




The compound of **Example 50** was prepared as previously described in Example 21 from 2-bromo-1-naphthalenemethanol (Shaik, F. H. *et al*, *Beilstein J. Org. Chem.* **2009**, 5). MS (ESI): *m/z* 200 [M+H]. ^1H NMR (300 MHz, DMSO-*d*₆) δ 8.71 (d, *J* = 6.6 Hz, 3H), 7.99-7.85 (m, 3H), 7.66-7.59 (m, 3H), 5.38 (d, *J* = 15.9 Hz, 1H), 5.10 (d, *J* = 15.9 Hz, 1H), 4.46 (s, 1H), 4.30 (d, *J* = 12.6 Hz, 1H), 4.00 (dd, *J* = 12.3, 2.1 Hz, 1H).

[0156] **Example 51: 2,4,7,8,9,10-Hexahydro-1H-7,10-methanobenzo[f]isochromen-1-amine hydrochloride**




The compound of **Example 51** was prepared as previously described in Example 48 using 1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-amine (Ehrenfreund, J.; *et al*, WO2004035589). MS (ESI): *m/z* 216 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.22 (dd, *J* = 7.5, 2.4 Hz, 1H), 6.88 (d, *J* = 7.5 Hz, 1H), 4.97-4.77 (m, 2H), 4.50 (d, *J* = 41.1 Hz, 1H), 4.36-4.24 (m, 1H), 3.91 (d, *J* = 12.9 Hz, 1H), 3.64 (d, *J* = 12.0 Hz, 1H), 3.40-3.33 (m, 1H), 2.06-1.97 (m, 2H), 1.85-1.65 (m, 1H), 1.65-1.62 (m, 1H), 1.19-1.09 (m, 2H).

[0157] **Example 52: 1,2,4,8,9,10-Hexahydropyrano[4,3-f]chromen-1-amine hydrochloride**



The compound of **Example 52** was prepared as previously described in Example 21 using (5-bromochroman-6-yl)methanol **52.5** prepared as described below. MS (ESI): *m/z* 206 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.40 (s, 3H), 6.88 (d, *J* = 8.4 Hz, 1H), 6.80 (d, *J* = 8.4 Hz, 1H), 4.79 (d, *J* = 15.3 Hz, 1H), 4.63 (d, *J* = 15.3 Hz, 1H), 4.34 (s, 1H), 4.22 (d, *J* = 12.6 Hz, 1H), 4.16-4.06 (m, 2H), 3.74 (d, *J* = 12.0 Hz, 1H), 3.00-2.90 (m, 1H), 2.78-2.68 (m, 1H), 1.98-1.89 (m, 2H).

[0158] **Synthesis of (5-bromochroman-6-yl)methanol (52.5)**



Step 1. 8-(*tert*-Butyl)-6-methylchroman **52.1** (3.5 g, 17.1 mmol, Mao, C.-H.; *et al*, *Bioorg. Med. Chem.* **2008**, *16*, 488) was dissolved in CH₃CN (50 mL) and treated with *N*-bromosuccinimide (3.04 g, 17.1 mmol). The mixture was heated to 80°C with stirring for 4 hr. After cooling to room temperature, the solvents were removed under reduced pressure,


and the residue was purified by flash column chromatography (SiO₂, 100% hexanes) to afford 5-bromo-8-(tert-butyl)-6-methylchroman **52.2** (2.5 g, 52%). ¹H NMR (300 MHz, CDCl₃) δ 7.03 (s, 1H), 4.13 (t, *J* = 5.1 Hz, 2H), 2.81 (t, *J* = 6.6 Hz, 2H), 2.37 (s, 3H), 2.08-2.00 (m, 2H), 1.41 (s, 9H).

Step 2. AlCl₃ (3.65 g, 27.4 mmol) was added to a solution of **52.2** (6.5 g, 22.9 mmol) in CH₂Cl₂ (100 mL) at room temperature and stirred for 2 h. The reaction mixture was carefully quenched by the dropwise addition of 1 M aqueous HCl (30 mL). The mixture was extracted with CH₂Cl₂ (200 mL). The organic phase was washed with brine (2×50 mL dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, 100% hexanes) to afford 5-bromo-6-methylchroman **52.3** (4.50 g, 86%) as a green oil. ¹H NMR (300 MHz, CDCl₃) δ 7.04 (d, *J* = 8.4 Hz, 1H), 6.78 (d, *J* = 8.4 Hz, 1H), 4.16 (t, *J* = 5.1 Hz, 2H), 2.83 (t, *J* = 6.6 Hz, 2H), 2.42 (s, 3H), 2.09-2.03 (m, 2H).

Step 3. Chromium(VI) oxide (9.89 g, 99.0 mmol) was added to a solution of **52.3** (4.5 g, 19.8 mmol) in AcOH (40 mL) and water (20 mL). The reaction was stirred at ambient temperature for 2 h, diluted with water (40 mL), and extracted with EtOAc (200 mL). The organic phase was washed with saturated aqueous NaHCO₃ (2×30 mL) and brine (2×30 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 5% EtOAc/hexanes) gave 5-bromochroman-6-carbaldehyde **52.4** (1 g, 21%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 10.31 (s, 1H), 7.75 (d, *J* = 8.7 Hz, 1H), 6.87 (d, *J* = 8.7 Hz, 1H), 4.24 (t, *J* = 5.1 Hz, 2H), 2.85 (t, *J* = 6.6 Hz, 2H), 2.13-2.06 (m, 2H).


Step 4. NaBH₄ (376 mg, 9.94 mmol) was added in a single portion to a solution of **52.4** (1.2 g, 4.97 mmol) in CH₃OH (30 mL). The reaction mixture was stirred at room temperature for 30 min and then partitioned between water (30 mL) and EtOAc (300 mL). The organic phase was washed with brine (2×40 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide (5-bromochroman-6-yl)methanol **52.5** (1.10 g, 91%) as a colorless oil which was used directly without further purification.

[0159] Example 53: 5-Ethylisochroman-4-amine hydrochloride



The compound of **Example 53** was prepared as previously described in Example 21 using (2-bromo-3-ethylphenyl)methanol **53.4** prepared as described below. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.36 (t, *J* = 6.0 Hz, 1H), 7.24 (d, *J* = 9.0 Hz, 1H), 7.00 (d, *J* = 6.0 Hz, 1H), 4.95 (d, *J* = 15.0 Hz, 1H), 4.84 (d, *J* = 15.0 Hz, 1H), 4.52 (s, 1H), 4.25 (dd, *J* = 12.0, 3.0 Hz, 1H), 3.92 (dd, *J* = 12.0, 3.0 Hz, 1H), 2.85-2.66 (m, 2H), 1.27 (t, *J* = 6.0 Hz, 3H).

[0160] Synthesis of (2-Bromo-3-ethylphenyl)methanol (53.4)




Step 1. Methyl 2-amino-3-vinylbenzoate **53.1** was prepared as previously described in Example 7 using methyl 2-amino-3-bromobenzoate and vinylboronic acid pinacol ester. ¹H NMR (300 MHz, CDCl₃) δ 7.84 (d, *J* = 9.0 Hz, 1H), 7.38 (d, *J* = 6.0 Hz, 1H), 6.79-6.62 (m, 2H), 5.97 (brs, 2H), 5.62 (dd, *J* = 18.0, 3.0 Hz, 1H), 5.38 (dd, *J* = 12.0, 3.0 Hz, 1H), 3.87 (s, 3H).

Step 2. Methyl 2-amino-3-ethylbenzoate **53.2** was prepared as previously described in Example 9. ¹H NMR (300 MHz, CDCl₃) δ 7.77 (d, *J* = 9.0 Hz, 1H), 7.20 (d, *J* = 6.0 Hz, 1H), 6.63 (t, *J* = 6.0 Hz, 1H), 5.89 (brs, 2H), 3.86 (s, 1H), 2.54-2.47 (m, 2H), 1.26 (t, *J* = 7.62 Hz, 3H).

Step 3. Methyl 2-bromo-3-ethylbenzoate **53.3** was prepared as previously described in Example 19. ^1H NMR (300 MHz, CDCl_3) δ 7.45–7.42 (m, 1H), 7.36–7.25 (m, 2H), 3.92 (s, 3H), 2.87–2.80 (m, 2H), 1.24 (t, J = 7.5 Hz, 3H).



Step 4. (2-Bromo-3-ethylphenyl)methanol **53.4** was prepared as previously described in Example 19. ^1H NMR (300 MHz, CDCl_3) δ 7.32–7.23 (m, 2H), 7.19–7.16 (m, 1H), 4.76 (s, 2H), 2.84–2.76 (m, 2H), 2.17 (brs, 1H), 1.23 (t, J = 15.0 Hz, 3H).

[0161] **Example 54: N-Methylisochroman-4-amine hydrochloride**




Step 1. Isochroman-4-one (200 mg, 1.35 mmol) was dissolved in EtOH (10 mL) and treated sequentially with 2M NH_3 in MeOH (3.4 mol, 6.8 mmol), $\text{Ti}(\text{O}i\text{-Pr})_4$ (0.38 g, 1.35 mmol) and NaCNBH_3 (0.34 g, 5.4 mmol). The reaction mixture was heated with stirring to reflux for 6 h and then cooled to room temperature and stirred for an additional 16 h. The reaction mixture was partitioned between EtOAc (40 mL) and saturated aqueous Na_2CO_3 (30 mL). The aqueous layer was extracted with EtOAc (2×15 mL). The combined organics were dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to afford N-methylisochroman-4-amine (100 mg, 45%) as a colorless oil, which was converted to the compound of **Example 54** as previously described in Example 1. MS (ESI): m/z 164 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.37 (brs, 1H), 9.20 (brs, 1H), 7.57 (d, J = 7.5 Hz, 1H), 7.45–7.32 (m, 2H), 7.21 (d, J = 7.5 Hz, 1H), 4.87 (d, J = 15.6 Hz, 1H), 4.73 (d, J = 15.6 Hz, 1H), 4.42 (d, J = 13.2 Hz, 1H), 4.30 (brs, 1H), 3.87 (dd, J_1 = 12.9, 2.1 Hz, 1H), 2.56–2.50 (m, 3H).

[0162] **Table 5.** Compounds prepared as described in Example 54 using the appropriate starting materials.

| Example | Structure | Characterization Data |
|---------|---|--|
| 55 |  | MS (ESI): m/z 220 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 8.52 (s, 1H), 8.34 (s, 1H), 7.23 (d, J = 9.0 Hz, 1H), 6.93 (d, J = 9.0 Hz, 1H), 4.90 (d, J = 15.0 Hz, 1H), 4.73 (d, J = 15.0 Hz, 1H), 4.61 (d, J = 9.0 Hz, 1H), 4.43 (d, J = 12.0 Hz, 1H), 3.71 (d, J = 15.0 Hz, 1H), 3.64-3.62 (m, 1H), 2.26 (d, J = 6.0 Hz, 6H), 1.38-1.09 (m, 6H). |
| 56 |  | MS (ESI): m/z 222 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 8.80 (d, J = 24.9 Hz, 2H), 7.23 (d, J = 7.5 Hz, 1H), 6.92 (d, J = 7.2 Hz, 1H), 5.32 (s, 1H), 4.88 (d, J = 15.6 Hz, 1H), 4.70 (d, J = 15.3 Hz, 1H), 4.62-4.61 (m, 2H), 4.50 (d, J = 12.6 Hz, 1H), 3.77-3.70 (m, 3H), 3.13-3.07 (m, 2H), 2.26 (d, J = 4.8 Hz, 6H). |

[0163] **Example 57. *N*,**5,6**-Trimethylisochroman-4-amine hydrochloride**










Step 1. *tert*-Butyl (5,6-dimethylisochroman-4-yl)carbamate (400 mg, 1.44 mmol, prepared as described in Example 42) was dissolved in anhydrous THF (4 mL) and treated with a 95% dispersion of sodium hydride in mineral oil (103 mg, 4.31 mmol) at room temperature. Once addition was complete, the reaction mixture was heated to 40°C and stirred for 15 min. Iodomethane (817 mg, 5.76 mmol) was added and heating at 40°C was continued overnight. After cooling to room temperature, the reaction mixture was partitioned between brine (15 mL) and EtOAc (10 mL). The aqueous phase was washed with EtOAc (2×10 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, gradient elution from 3% EtOAc/hexanes to 5% EtOAc/hexanes) gave *tert*-butyl (5,6-dimethylisochroman-4-yl) (methyl)carbamate **57.1** (400 mg, 1.37 mmol) as a yellow oil. MS (ESI): *m/z* 292 [M⁺H].


Step 2. The compound of **Example 57** was prepared as previously described in Example 1. MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.85 (brs, 2H), 7.23 (d, J = 7.8 Hz, 1H), 6.92 (d, J = 7.8 Hz, 1H), 4.87 (d, J = 15.3 Hz, 1H), 4.70 (d, J = 15.3 Hz, 1H), 4.47 (d, J = 9.3 Hz, 2H), 3.76 (dd, J = 13.2, 1.8 Hz, 1H), 2.62 (s, 3H), 2.26 (s, 6H).

[0164] **Table 6. Compounds prepared as described in Example 57 using the appropriate Boc-protected intermediates and alkyl halide.**

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 58 | | MS (ESI): m/z 206 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 8.81 (s, 1H), 8.50 (s, 1H), 7.23 (d, J = 9.0 Hz, 1H), 6.93 (d, J = 9.0 Hz, 1H), 4.88 (d, J = 15.0 Hz, 1H), 4.71 (d, J = 15.0 Hz, 1H), 4.51-4.43 (m, 2H), 3.74 (d, J = 12.0 Hz, 1H), 3.12-3.09 (m, 1H), 2.25 (d, J = 6.0 Hz, 6H), 1.26 (t, J = 9.0 Hz, 3H). |
| 59 | | MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.26 - 7.16 (m, 3H), 5.01 (d, J = 15.6 Hz, 1H), 4.83 (d, J = 15.3 Hz, 1H), 4.50 - 4.47 (m, 1H), 4.15 - 4.09 (m, 1H), 4.03 - 3.96 (m, 1H), 2.73 (s, 3H), 2.47 - 2.39 (m, 5H). |
| 258 | | MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 9.08 (s, 1H), 8.55 (s, 1H), 7.51 (d, J = 7.8 Hz, 1H), 7.45-7.42 (m, 1H), 7.33 (t, J = 7.2 Hz, 1H), 7.22 (d, J = 7.8 Hz, 1H), 4.97 (d, J = 15.9 Hz, 1H), 4.80 (d, J = 15.9 Hz, 1H), 4.32 (s, 1H), 3.75-3.70 (m, 1H), 2.42 (s, 3H), 1.80-1.70 (m, 2H), 1.02 (t, J = 7.2 Hz, 3H). |
| 259 | | MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, DMSO- d_6) δ 9.58 (s, 1H), 9.00 (s, 1H), 7.51 (d, J = 7.5 Hz, 1H), 7.43-7.40 (m, 1H), 7.35-7.30 (m, 1H), 7.21 (d, J = 7.5 Hz, 1H), 4.80 (d, J = 16.2 Hz, 1H), 4.69 (d, J = 16.5 Hz, 1H), 4.29-4.27 (m, 1H), 4.18 (s, 1H), 2.51 (s, 3H), 1.52-1.38 (m, 2H), 0.95 (t, J = 7.5 Hz, 3H). |
| 262 | | MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.48-7.43 (m, 2H), 7.39-7.34 (m, 1H), 7.21 (d, J = 7.5Hz, 1H), 4.97 (d, J = 15.6Hz, 1H), 4.81 (d, J = 15.6 Hz, 1H), 4.43 (dd, J = 13.2, 1.2Hz, 1H), 4.29 (s, 1H), 3.92 (dd, J = 13.2, 1.8Hz, 1H), 3.11-3.06 (m, 2H), 1.81-1.68 (m, 2H), 1.02 (t, J = 7.4 Hz, 3H). |
| 263 | | MS (ESI): m/z 188 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.47 (t, J = 7.5 Hz, 2H), 7.37 (t, J = 7.8 Hz, 1H), 7.22 (d, J = 7.5Hz, 1H), 4.98 (d, J = 15.6Hz, 1H), 4.83 (d, J = 15.6Hz, 1H), 4.48-4.41 (m, 2H), 4.09-3.90 (m, 3H), 3.33 (t, J = 2.4 Hz, 1H). |
| 264 | | MS (ESI): m/z 206 [M+H]. ^1H NMR (300 MHz, CDCl ₃) δ 7.42-7.39 (m, 2H), 7.32-7.27 (m, 1H), 7.15 (d, J = 7.8 Hz, 1H), 4.80-4.75 (m, 2H), 4.51-4.48 (m, 1H), 4.08 (s, 1H), 3.01-2.87 (m, 2H), 1.71-1.62 (m, 2H), 1.18 (d, J = 6.6 Hz, 3H), 0.94 (t, J = 7.5 Hz, 3H). |
| 265 | | MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.45 (t, J = 7.5 Hz, 2H), 7.37 (d, J = 7.8 Hz, 1H), 7.21 (d, J = 7.5Hz, 1H), 4.97 (d, J = 15.6 Hz, 1H), 4.81 (d, J = 15.6 Hz, 1H), 4.43 (dd, J = 13.2, 1.2Hz, 1H), 4.29 (s, 1H), 3.92 (dd, J = 13.2, 1.8Hz, 1H), 3.13-3.06 (m, 2H), 1.81-1.68 (m, 2H), 1.02 (t, J = 7.5 Hz, 3H). |




| | | |
|-----|---|--|
| 266 |  | MS (ESI): <i>m/z</i> 188 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.47 (t, <i>J</i> = 7.2 Hz, 2H), 7.37 (t, <i>J</i> = 7.5 Hz, 1H), 7.22 (d, <i>J</i> = 7.5 Hz, 1H), 4.97 (d, <i>J</i> = 15.6 Hz, 1H), 4.82 (d, <i>J</i> = 15.6 Hz, 1H), 4.48-4.41 (m, 2H), 4.08-3.90 (m, 3H), 3.32 (t, <i>J</i> = 2.4 Hz, 1H). |
| 267 |  | MS (ESI): <i>m/z</i> 202 [M+H]. ¹ H NMR (300 MHz, CDCl ₃) δ 7.62 (d, <i>J</i> = 7.5 Hz, 1H), 7.42-7.28 (m, 2H), 7.10 (d, <i>J</i> = 7.5 Hz, 1H), 5.05-5.00 (m, 2H), 4.78 (d, <i>J</i> = 15.9 Hz, 1H), 4.36 (s, 1H), 4.09 (dd, <i>J</i> = 17.1, 2.4 Hz, 1H), 3.50 (dd, <i>J</i> = 17.1, 2.4 Hz, 1H), 2.60 (t, <i>J</i> = 2.4 Hz, 1H), 1.26 (d, <i>J</i> = 6.6 Hz, 3H). |
| 268 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.42-7.39 (m, 2H), 7.32-7.27 (m, 1H), 7.15 (d, <i>J</i> = 7.8 Hz, 1H), 4.80-4.75 (m, 2H), 4.51-4.48 (m, 1H), 4.08 (s, 1H), 3.01-2.87 (m, 2H), 1.71-1.62 (m, 2H), 1.18 (d, <i>J</i> = 6.6 Hz, 3H), 0.94 (t, <i>J</i> = 7.5 Hz, 3H). |
| 269 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.27-7.21 (m, 2H), 7.16-7.11 (m, 1H), 7.01 (d, <i>J</i> = 7.8 Hz, 1H), 4.79 (d, <i>J</i> = 15.6 Hz, 1H), 4.65 (d, <i>J</i> = 15.6 Hz, 1H), 4.07 (s, 1H), 3.89-3.86 (m, 1H), 2.70 (t, <i>J</i> = 8.4 Hz, 2H), 1.61-1.41 (m, 2H), 1.26 (d, <i>J</i> = 6.9 Hz, 3H), 0.74 (t, <i>J</i> = 7.5 Hz, 3H). |
| 270 |  | MS (ESI): <i>m/z</i> 202 [M+H]. ¹ H NMR (300 MHz, CDCl ₃) δ 7.62 (d, <i>J</i> = 7.2 Hz, 1H), 7.42-7.28 (m, 2H), 7.10 (d, <i>J</i> = 7.5 Hz, 1H), 5.05-5.00 (m, 2H), 4.78 (d, <i>J</i> = 15.9 Hz, 1H), 4.36 (s, 1H), 4.09 (dd, <i>J</i> = 17.1, 2.4 Hz, 1H), 3.50 (dd, <i>J</i> = 17.1, 2.4 Hz, 1H), 2.60 (t, <i>J</i> = 2.4 Hz, 1H), 1.26 (d, <i>J</i> = 6.6 Hz, 3H). |
| 271 |  | MS (ESI): <i>m/z</i> 202 [M+H]. ¹ H NMR (300 MHz, CDCl ₃) δ 7.68 (d, <i>J</i> = 7.5 Hz, 1H), 7.41-7.26 (m, 2H), 7.09 (d, <i>J</i> = 7.5 Hz, 1H), 5.13 (d, <i>J</i> = 15.6 Hz, 1H), 4.85 (d, <i>J</i> = 15.6 Hz, 1H), 4.38 (s, 1H), 4.21 (dd, <i>J</i> = 16.8, 2.4 Hz, 1H), 4.06 (q, <i>J</i> = 6.9 Hz, 1H), 3.52 (dd, <i>J</i> = 16.8, 2.4 Hz, 1H), 2.57 (t, <i>J</i> = 2.4 Hz, 1H), 1.74 (d, <i>J</i> = 6.9 Hz, 3H). |
| 272 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CDCl ₃) δ 10.09 (brs, 1H), 9.23 (brs, 1H), 8.20-8.17 (m, 1H), 7.40-7.37 (m, 2H), 7.12-7.09 (m, 1H), 5.10 (d, <i>J</i> = 15.6 Hz, 1H), 4.82 (d, <i>J</i> = 15.6 Hz, 1H), 4.48 (s, 1H), 4.06-4.02 (m, 1H), 3.20-3.06 (m, 1H), 2.65-2.47 (m, 1H), 2.05-1.74 (m, 2H), 1.75 (d, <i>J</i> = 6.9 Hz, 3H), 0.88 (t, <i>J</i> = 7.2 Hz, 3H). |
| | | |

[0165] Example 60. N,N-dimethylisochroman-4-amine hydrochloride




Step 1. The compound of **Example 57** (200 mg, 1.04 mmol) was dissolved in CH₃OH (4 mL). Paraformaldehyde (93.6 mg, 3.12 mmol), NaBH₃CN (196 mg, 3.12 mmol), and acetic acid (187 mg, 3.12 mmol) were added, and the reaction was stirred at room temperature for 1 h. The reaction mixture was poured into saturated aqueous Na₂CO₃ (10 mL) and extracted with EtOAc (4×10 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 100% EtOAc) gave N,N-dimethylisochroman-4-amine (120 mg, 56%) as a white oil. The compound of **Example 60** was prepared as described previously in Example 1 as an off-white solid. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.65 (brs, 1H), 7.54 (d, *J* = 7.5 Hz, 1H), 7.48–7.43 (m, 1H), 7.37 (t, *J* = 7.5 Hz, 1H), 7.23 (d, *J* = 7.8 Hz, 1H), 4.89 (d, *J* = 15.6 Hz, 1H), 4.73–4.65 (m, 2H), 4.49 (s, 1H), 3.85 (dd, *J* = 13.8, 2.7 Hz, 1H), 2.82 (d, *J* = 4.8 Hz, 3H), 2.64 (d, *J* = 4.5 Hz, 3H).

[0166] **Table 7. Compounds prepared as described in Example 60 using the appropriate amine.**

| Example | Structure | Characterization Data |
|---------|---|--|
| 61 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.31 (d, <i>J</i> = 8.1 Hz, 1H), 6.98 (d, <i>J</i> = 7.8 Hz, 1H), 4.97 (d, <i>J</i> = 15.6 Hz, 1H), 4.79 (d, <i>J</i> = 15.6 Hz, 1H), 4.67 (d, <i>J</i> = 12.6 Hz, 2H), 3.88 (d, <i>J</i> = 12.6 Hz, 1H), 2.92 (s, 6H), 2.35 (d, <i>J</i> = 2.7 Hz, 6H). |
| 62 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.29 - 7.26 (m, 2H), 7.19 (d, <i>J</i> = 7.8 Hz, 1H), 5.14 (d, <i>J</i> = 15.9 Hz, 1H), 4.81 (d, <i>J</i> = 15.9 Hz, 1H), 4.45 (d, <i>J</i> = 3.6 Hz, 1H), 4.21 - 4.15 (m, 1H), 4.00 - 3.96 (m, 1H), 3.13 (s, 3H), 2.72 - 2.68 (m, 1H), 2.66 (s, 3H), 2.41 - 2.48 (m, 1H), 2.38 (s, 3H). |
| 260 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.55-7.50 (m, 2H), 7.37 (t, <i>J</i> = 7.2 Hz, 1H), 7.27 (d, <i>J</i> = 7.5 Hz, 1H), 5.02 (d, <i>J</i> = 15.9 Hz, 1H), 4.79 (d, <i>J</i> = 15.9 Hz, 1H), 4.55 (s, 1H), 3.80-3.72 (m, 1H), 2.86 (d, <i>J</i> = 5.1 Hz, 3H), 2.46 (d, <i>J</i> = 4.8 Hz, 3H), 1.97-1.87 (m, 2H), 1.03 (t, <i>J</i> = 7.2 Hz, 3H). |
| 261 |  | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.63 (d, <i>J</i> = 7.5 Hz, 1H), 7.49-7.44 (m, 1H), 7.36 (t, <i>J</i> = 7.2 Hz, 1H), 7.23 (d, <i>J</i> = 7.8 Hz, 1H), 4.83 (d, <i>J</i> = 16.5 Hz, 1H), 4.66 (d, <i>J</i> = 16.5 Hz, 1H), 4.54-4.49 (m, 1H), 4.38 (s, 1H), 2.70 (s, 6H), |


| | | |
|-----|--|---|
| | | 1.50-1.39 (m, 2H), 0.96 (t, $J = 7.2$ Hz, 3H). |
| 283 | | MS (ESI): m/z 279 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.48 (d, $J = 8.4$ Hz, 1H), 7.06 (d, $J = 8.4$ Hz, 1H), 6.94 (s, 1H), 4.87 (d, $J = 15.9$ Hz, 1H), 4.68 (d, $J = 15.9$ Hz, 1H), 4.57 (d, $J = 13.9$ Hz, 1H), 4.31 (s, 1H), 3.86-3.79 (m, 1H), 3.44-3.26 (m, 2H), 2.95 (d, $J = 33.9$ Hz, 3H), 2.85 (s, 3H), 2.81 (s, 3H), 1.17 (dt, $J = 22.5, 6.9$ Hz, 3H). |
| 284 | | MS (ESI): m/z 279 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.25 (d, $J = 1.8$ Hz, 1H), 7.20-7.15 (m, 2H), 4.88 (d, $J = 15.6$ Hz, 1H), 4.68 (d, $J = 15.6$ Hz, 1H), 4.55 (d, $J = 13.8$ Hz, 1H), 4.27 (s, 1H), 3.82 (dd, $J = 14.1, 2.4$ Hz, 1H), 3.43-3.27 (m, 2H), 2.95 (d, $J = 36.0$ Hz, 3H), 2.82 (s, 6H), 1.20-1.06 (m, 3H). |
| 285 | | MS (ESI): m/z 293 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.34-7.27 (m, 3H), 5.05 (d, $J = 15.9$ Hz, 1H), 4.89 (d, $J = 15.9$ Hz, 1H), 4.48-4.45 (m, 1H), 4.20-4.11 (m, 1H), 3.50-3.36 (m, 2H), 3.06 (d, $J = 36.3$ Hz, 3H), 3.05 (s, 3H), 2.70 (s, 3H), 1.58 (d, $J = 6.9$ Hz, 3H), 1.24 (dt, $J = 25.1, 7.2$ Hz, 3H). |
| 286 | | MS (ESI): m/z 293 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.34-7.26 (m, 3H), 4.90-4.88 (m, 2H), 4.83-4.76 (m, 1H), 4.26 (s, 1H), 3.58-3.36 (m, 2H), 3.06 (d, $J = 36.0$ Hz, 3H), 2.95 (s, 3H), 2.88 (s, 3H), 1.31-1.18 (m, 6H). |
| 287 | | MS (ESI): m/z 293 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.26-7.16 (m, 3H), 4.99 (d, $J = 15.6$ Hz, 1H), 4.85 (d, $J = 15.6$ Hz, 1H), 4.28 (s, 1H), 4.05 (q, $J = 6.6$ Hz, 1H), 3.57-3.31 (m, 2H), 3.05 (d, $J = 34.1$ Hz, 3H), 1.41 (d, $J = 6.6$ Hz, 3H), 1.24 (dt, $J = 23.7, 7.2$ Hz, 3H). |
| 288 | | MS (ESI): m/z 293 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.09-7.00 (m, 3H), 4.69 (s, 2H), 4.17-4.04 (m, 2H), 3.40-3.32 (m, 2H), 2.89 (d, $J = 34.8$ Hz, 3H), 1.17 (d, $J = 6.6$ Hz, 3H), 1.09 (dt, $J = 24.6, 7.2$ Hz, 3H). |

[0167] Example 63. 4,4,7-Trimethyl-1,3,4,5-tetrahydrobenzo[c]oxepin-5-amine hydrochloride



[0168] The compound of **Example 63** was prepared as previously described in Example 1, using 4,4,7-trimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (prepared as described below). MS (ESI): *m/z* 206 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.24 - 7.16 (m, 3H), 4.93 - 4.76 (m, 2H), 4.29 (s, 1H), 3.80 (d, *J* = 12.6 Hz, 1H), 3.61 (d, *J* = 12.6 Hz, 1H), 2.39 (s, 3H), 1.06 (s, 3H), 0.97 (s, 3H).

[0169] **Synthesis of 4,7-methyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one and 4,4,7-trimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one**




Step 1. A solution of 1.2 g (6.8 mmol) of 7-methyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (prepared as previously described in Example 34) in DMF (100 mL) was treated with iodomethane (9.63 g, 67.9 mmol) and cesium carbonate (11.0 g, 33.9 mmol). The reaction was heated with stirring at 50 °C for 2 h, and then partitioned between EtOAc (250 mL) and water (300 mL). The organic phase was washed with brine (100 mL) dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, gradient elution from 100% hexanes to 5% EtOAc/hexanes) gave 4,7-dimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (550 mg, 43%) as a colorless oil and 4,4,7-trimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (320 mg, 23%) as a colorless oil.

[0170] **4,7-dimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one:** MS (ESI): *m/z* 190 [M+H].

[0171] **4,4,7-trimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one:** MS (ESI): *m/z* 204 [M+H].

[0172] **Example 68. 6-Fluoro-4-methylisochroman-4-amine hydrochloride**



Step 1. A solution of 6-fluoroisochroman-4-one (420 mg, 2.52 mmol, prepared as described in Example 6) in diethyl ether (40 mL) was cooled in an ice bath and treated with methylmagnesium bromide (3.78 mL, 3.78 mmol, 1M in THF). The reaction was stirred at 0 °C for 1 h. Saturated aqueous NH₄Cl (60 mL) and EtOAc (40 mL) were added to the reaction


vessel and the resulting biphasic mixture was transferred to a separatory funnel. The layers were separated and the aqueous phase was washed with EtOAc (2×40 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide 6-fluoro-4-methylisochroman-4-ol **68.1** (455 mg, 99%) as yellow oil. GC-MS (ESI) *m/z* 182 [M+H].

Step 2. Compound **68.1** (250 mg, 1.37 mmol) was dissolved in toluene (6 mL) and cooled in an ice bath. Azidotrimethylsilane (394 mg, 3.42 mmol) and BF₃-Et₂O (1.24 g, 4.11 mmol) were added, the ice bath was removed, and the reaction was stirred at 30°C for 2 days. The reaction mixture was carefully quenched by the slow addition of NH₄OH and extracted with EtOAc (3×50 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 100% hexanes) afforded 4-azido-6-fluoro-4-methyl isochroman **68.2** (240 mg, 84%) as yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.19 (d, *J* = 1.8 Hz, 1H), 7.05-6.99 (m, 2H), 4.86 (d, *J* = 15.0 Hz, 1H), 4.74 (d, *J* = 14.7 Hz, 1H), 4.05 (d, *J* = 11.7 Hz, 1H), 3.78 (d, *J* = 12.0 Hz, 1H), 1.49 (s, 3H).

Step 3. Compound **68.2** (240 mg, 1.15 mmol) was dissolved in THF (10 mL), cooled in an ice bath, and treated with LiAlH₄ (87.2 mg, 2.30 mmol) in a single portion. The ice bath was removed, and the reaction mixture was stirred at room temperature for 2 h. Excess hydride was quenched by the sequential addition of water (0.5 mL) and aqueous 2M NaOH (0.5 mL). The resulting mixture was partitioned between aqueous 1M HCl (40 mL) and CH₂Cl₂ (50 mL). The aqueous phase was washed with CH₂Cl₂ (2×50 mL) and then made basic by the addition of aqueous 1M NaOH (40 mL). The aqueous phase was extracted with CH₂Cl₂ (4×50 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to give crude 6-fluoro-4-methylisochroman-4-amine, which was purified by preparative HPLC and converted to the compound of **Example 68** (67 mg, 27%) as a white solid as described previously in Example 1. MS (ESI): *m/z* 182 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.39 (dd, *J* = 9.9, 2.4 Hz, 1H), 7.24 - 7.15 (m, 2H), 4.94 (d, *J* = 15.2 Hz, 1H), 4.83 (d, *J* = 15.3 Hz, 1H), 4.06 (d, *J* = 12.6 Hz, 1H), 3.82 (d, *J* = 12.6 Hz, 1H), 1.66 (s, 3H).

[0173] **Example 72. *cis*-3-methylisochroman-4-amine hydrochloride**

[0174] **Example 73. *trans*-3-methylisochroman-4-amine hydrochloride**



Step 1. 3-Methylisochroman-4-one **72.1** was prepared as previously described in Example 69 using isochroman-4-one (prepared as previously described in Example 12) and iodomethane. MS (ESI): m/z 163 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 8.05 (d, J = 7.5 Hz, 1H), 7.59-7.53 (m, 1H), 7.49-7.39 (m, 1H), 7.21 (d, J = 7.5 Hz, 1H), 5.18 (s, 2H), 4.28 (q, J = 6.6 Hz, 1H), 1.75 (d, J = 6.6 Hz, 3H).

Step 2. 3-Methylisochroman-4-ol **72.2** was prepared as previously described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.56-7.45 (m, 1H), 7.34-7.24 (m, 2H), 7.07-7.01 (m, 1H), 4.89-4.82 (m, 2H), 4.46-4.40 (m, 0.5H), 4.40-4.25 (m, 0.5H), 4.15-4.13 (m, 0.5H), 3.69-3.64 (m, 0.5H), 1.44 (d, J = 6.3 Hz, 3H).

Step 3. 4-Azido-3-methylisochroman **72.3** was prepared as previously described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.45 (d, J = 8.7 Hz, 1H), 7.43-7.24 (m, 2H), 7.02 (d, J = 8.4 Hz, 1H), 4.80 (d, J = 2.7 Hz, 2H), 4.11 (d, J = 7.8 Hz, 1H), 3.80-3.76 (m, 1H), 1.44 (d, J = 6.3 Hz, 3H).

Step 4. LiAlH_4 (119 mg, 3.15 mmol) was added to a room temperature solution of compound **72.3** (200 mg, 1.05 mmol) in THF (10 mL) in a single portion, and the reaction was stirred for 1 h. A solution of aqueous 1M NaOH (5 mL), di-tert-butyl dicarbonate (916 mg, 4.20 mmol) and Na_2CO_3 (333 mg, 3.15 mmol) were added to the reaction vessel, and the biphasic mixture was stirred vigorously for 2 h. After transferring to a separatory funnel, the mixture was partitioned between water (20 mL) and EtOAc (100 mL). The organic phase was washed with brine (40 mL), dried over anhydrous Na_2SO_4 , filtered, and concentrated under

reduced pressure. Purification by flash column chromatography (SiO₂, gradient elution from 100% hexanes to 5% EtOAc/hexanes) provided *tert*-butyl ((3*S*,4*S*)-3-methylisochroman-4-yl) carbamate **72.4a** (90.0 mg, 32%) as a white solid and *tert*-butyl ((3*R*,4*S*)-3-methylisochroman-4-yl)carbamate **72.4b** (70.0 mg, 25%) as a white solid.

Rac-*tert*-butyl ((3*S*,4*S*)-3-methylisochroman-4-yl)carbamate **72.4a** TLC higher R_f. MS (ESI): *m/z* 286 [M+Na]. ¹H NMR (300 MHz, CDCl₃) δ 7.45-7.42 (m, 1H), 7.28-7.25 (m, 2H), 7.03-7.00 (m, 1H), 4.95-4.81 (m, 1H), 4.82 (s, 2H), 4.64 (d, *J* = 8.7 Hz, 1H), 3.91-3.84 (m, 1H), 1.46 (s, 9H), 1.34 (d, *J* = 6.3 Hz, 3H).

Rac-*tert*-butyl ((3*R*,4*S*)-3-methylisochroman-4-yl)carbamate **72.4b** TLC lower R_f. MS (ESI): *m/z* 286 [M+Na]. ¹H NMR (300 MHz, CDCl₃) δ 7.42-7.37 (m, 1H), 7.28-7.21 (m, 2H), 7.02-7.00 (m, 1H), 4.81 (s, 2H), 4.79-4.65 (m, 2H), 3.72-3.67 (m, 1H), 1.48 (s, 9H), 1.38 (d, *J* = 6.3 Hz, 3H).

Step 5. The compounds of **Example 72** and **Example 73** were prepared as previously described in Example 1.

[0175] **Example 72.** MS (ESI): *m/z* 164 [M+H]. ¹H NMR (300 MHz, CDCl₃) δ 7.42-7.33 (m, 3H), 7.18 (d, *J* = 6.9 Hz, 1H), 4.99-4.83 (m, 2H), 4.25 (s, 1H), 4.03-4.05 (m, 1H), 1.40 (d, *J* = 6.6 Hz, 3H).

[0176] **Example 73.** MS (ESI): *m/z* 164 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.46-7.37 (m, 3H), 7.21 (d, *J* = 7.2 Hz, 1H), 4.97-4.78 (m, 2H), 4.37-4.20 (m, 2H), 1.33 (d, *J* = 6.6 Hz, 3H).


[0177] **Example 74. cis-4,7-dimethyl-1,3,4,5-tetrahydrobenzo[c]oxepin-5-amine hydrochloride**



[0153] The compound of **Example 74** was prepared as previously described in Example 72, using 4,7-dimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (prepared as described in Example 63). MS (ESI): *m/z* 192 [M+H]. ¹H NMR (300 MHz, CD₃OD) δ 7.23 - 7.06 (m, 3H), 5.07

(d, $J = 15.3$ Hz, 1H), 4.79 (d, $J = 15.3$ Hz, 1H), 4.28 (s, H), 4.21 - 4.15 (m, 1H), 3.61 - 3.54 (m, 1H), 2.52 - 2.44 (m, 1H), 2.38 (s, 3H), 0.93 (d, $J = 7.2$ Hz, 3H).


[0154] **Example 75.** *trans*-4,7-dimethyl-1,3,4,5-tetrahydrobenzo[c]oxepin-5-amine hydrochloride



[0155] The compound of **Example 75** was prepared as previously described in Example 72, using 4,7-dimethyl-3,4-dihydrobenzo[c]oxepin-5(1H)-one (prepared as described in Example 63). MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.20 - 7.17 (m, 2H), 7.11 (s, 1H), 4.82 - 4.69 (m, 3H), 4.10 - 4.05 (m, 1H), 3.88 - 3.82 (m, 1H), 2.50 - 2.41 (m, 1H), 2.40 (s, 3H), 0.94 (d, $J = 6.9$ Hz, 3H).

[0156] **Example 76.** *cis*-3,6-Dimethylisochroman-4-amine hydrochloride

[0157] **Example 77.** *trans*-3,6-Dimethylisochroman-4-amine hydrochloride



Step 1. 2-Bromo-1-((but-3-en-2-yloxy)methyl)-4-methylbenzene **76.1** was prepared as previously described in Example 10 using 2-bromo-1-(bromomethyl)-4-methylbenzene and but-3-en-2-ol. MS (ESI): *m/z* 256 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 7.38 (s, 1H), 7.37 (d, *J* = 7.8 Hz, 1H), 7.12 (d, *J* = 7.8 Hz, 1H), 5.89-5.78 (m, 1H), 5.29-5.18 (m, 2H), 4.57 (d, *J* = 12.9 Hz, 1H), 4.45 (d, *J* = 12.9 Hz, 1H), 2.33 (s, 3H), 1.33 (d, *J* = 6.3 Hz, 3H).

Step 2. 3,6-Dimethyl-4-methyleneisochroman **76.2** was prepared as previously described in Example 10. MS (ESI): *m/z* 175 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 7.49 (s, 1H), 7.08 (d, *J* = 7.8 Hz, 1H), 6.95 (d, *J* = 7.8 Hz, 1H), 5.63 (d, *J* = 0.9 Hz, 1H), 5.07 (d, *J* = 1.5 Hz, 1H), 4.82 (d, *J* = 6.9 Hz, 2H), 4.45-4.36 (m, 1H), 2.37 (s, 3H), 1.51 (d, *J* = 6.6 Hz, 3H).

Step 3. 3,6-Dimethylisochroman-4-one **76.3** was prepared as previously described in Example 10. MS (ESI): *m/z* 177 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 7.86 (s, 1H), 7.37 (d, *J* = 7.5 Hz, 1H), 7.11 (d, *J* = 7.5 Hz, 1H), 4.91 (s, 2H), 4.25 (q, *J* = 6.6 Hz, 1H), 2.41 (s, 3H), 1.50 (d, *J* = 6.6 Hz, 3H).

Step 4. 3,6-Dimethylisochroman-4-ol **76.4** was prepared as previously described in Example 1. ^1H NMR (300 MHz, CDCl_3) δ 7.30-7.24 (m, 1H), 7.14-7.06 (m, 1H), 6.95-6.90 (m, 1H), 4.78-4.76 (m, 2H), 4.01-4.35 (m, 0.75H), 4.25-4.21 (m, 0.25H), 3.80-3.77 (m, 0.25H), 3.68-3.60 (m, 0.75H), 2.37 (s, 3H), 1.42 (d, *J* = 6.3 Hz, 3H).

Step 5. 4-Azido-3,3-dimethylisochroman **76.5** was prepared as previously described in Example 1.

Step 6. Compound **76.5** (500 mg, 2.46 mmol) was dissolved in THF (30 mL) and treated with PPh_3 (645 mg, 2.46 mmol). The reaction was stirred at room temperature for 30 min. Water (10 mL) was added, and the solution was heated to 60 °C for 12 h. After cooling to room temperature, the mixture was partitioned between aqueous 1M HCl (30 mL) and EtOAc (50 mL). The aqueous phase was washed with EtOAc (2×30 mL). The aqueous phase was made basic by the addition of Na_2CO_3 (521 mg, 4.92 mmol), and then treated with di-*tert*-butyl dicarbonate (1.07 g, 4.92 mmol) and THF (40 mL). The biphasic mixture was stirred vigorously at room temperature for 30 min, and then partitioned between water (30 mL) and EtOAc (50 mL). The aqueous phase was extracted with EtOAc (2×50 mL). The combined organics were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 ,

gradient elution from 100% hexanes to 97% EtOAc/hexanes) gave *tert*-butyl-(*cis*-3,6-dimethylisochroman-4-yl)carbamate **76.5a** (300 mg, 44%) as a white solid and *tert*-butyl-(*trans*-3,6-dimethylisochroman-4-yl)carbamate **76.5b** (170 mg, 25%) as a white solid.

tert-butyl-(*cis*-3,6-dimethylisochroman-4-yl)carbamate **76.5a** TLC higher Rf. MS (ESI): *m/z* 300 [M+Na].

tert-butyl-(*trans*-3,6-dimethylisochroman-4-yl)carbamate **76.5b** TLC lower Rf. MS (ESI): *m/z* 300 [M+Na].

Step 7. The compounds of **Example 76** and **Example 77** were prepared as previously described in Example 1.










[0158] **Example 76.** MS (ESI): *m/z* 178 [M+H]. ^1H NMR (300 MHz, DMSO-*d*6) δ 8.27 (brs, 3H), 7.31 (s, 1H), 7.21 (d, *J* = 7.8 Hz, 1H), 7.06 (d, *J* = 7.8 Hz, 1H), 4.86 - 4.71 (m, 2H), 4.17 (d, *J* = 4.5 Hz, 1H), 3.97 - 3.91 (m, 1H), 2.31 (s, 3H), 1.31 (d, *J* = 6.6 Hz, 3H).

[0159] **Example 77.** MS (ESI): *m/z* 178 [M+H]. ^1H NMR (300 MHz, DMSO-*d*6) δ 8.5 (brs, 3H), 7.41 (s, 1H), 7.19 (d, *J* = 7.8 Hz, 1H), 7.07 (d, *J* = 8.1 Hz, 1H), 4.71 (s, 2H), 4.11 (d, *J* = 3.6 Hz, 2H), 2.31 (s, 3H), 1.27 (d, *J* = 6.3 Hz, 3H).

[0160] **Table 8. Compounds prepared as described in Example 76 using the appropriately substituted starting materials.**

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 78 | | MS (ESI): <i>m/z</i> 192 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> 6) δ 8.17 (brs, 3H), 7.18 (d, <i>J</i> = 7.8 Hz, 1H), 6.88 (d, <i>J</i> = 7.8 Hz, 1H), 4.86-4.70 (m, 2H), 4.36 (s, 1H), 3.90-3.83 (m, 1H), 2.23 (s, 3H), 2.21 (s, 3H), 1.32 (d, <i>J</i> = 6.6 Hz, 3H). |
| 79 | | MS (ESI): <i>m/z</i> 192 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> 6) δ 8.36 (brs, 3H), 7.19 (d, <i>J</i> = 7.8 Hz, 1H), 6.90 (d, <i>J</i> = 7.8 Hz, 1H), 4.74-4.20 (d, <i>J</i> = 5.1 Hz, 2H), 4.43-4.33 (m, 2H), 2.25 (s, 6H), 1.11 (d, <i>J</i> = 6.6 Hz, 3H). |
| 80 | | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.47-7.31 (m, 2H), 7.28-7.17 (m, 1H), 5.01-4.78 (m, 2H), 4.45 (d, <i>J</i> = 2.1 Hz, 1H), 4.05-3.89 (m, 1H), 1.41 (d, <i>J</i> = 6 Hz, 3H). |

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 81 | | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.55 (brs, 3H), 7.50-7.42 (m, 2H), 7.22 (dd, <i>J</i> = 3 Hz, 6 Hz, 1H), 4.85-4.71 (m, 2H), 4.51-4.44 (m, 1H), 4.32 (d, <i>J</i> = 3 Hz, 1H), 1.15 (d, <i>J</i> = 6 Hz, 3H). |
| 82 | | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.13 (brs, 3H), 7.29 (t, <i>J</i> = 7.5 Hz, 1H), 7.16 (d, <i>J</i> = 7.5 Hz, 1H), 6.99 (d, <i>J</i> = 7.5 Hz, 1H), 4.92-4.76 (m, 2H), 4.34-4.32 (m, 1H), 3.95-3.89 (m, 1H), 2.37 (s, 3H), 1.33 (d, <i>J</i> = 6.6 Hz, 3H). |
| 83 | | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.42 (s, 1H), 7.30 (d, <i>J</i> = 9 Hz, 1H), 7.15 (d, <i>J</i> = 9 Hz, 1H), 7.00 (d, <i>J</i> = 9 Hz, 1H), 4.83 - 4.69 (m, 2H), 4.46 - 4.39 (m, 1H), 4.26 (s, 1H), 2.41 (s, 3H), 1.12 (d, <i>J</i> = 9 Hz, 3H). |
| 84 | | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.09 (d, <i>J</i> = 8.4 Hz, 1H), 7.01 - 6.97 (m, 1H), 4.92 - 4.82 (m, 2H), 4.19 (s, 1H), 4.02 - 3.99 (m, 1H), 3.82 (s, 3H), 1.38 (d, <i>J</i> = 6.9 Hz, 3H). |
| 85 | | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.12 - 7.08 (m, 1H), 7.02 - 6.97 (m, 2H), 4.78 (s, 2H), 4.25 - 4.20 (m, 1H), 4.14 (d, <i>J</i> = 2.7 Hz, 1H), 3.82 (s, 3H), 1.30 (d, <i>J</i> = 6.6 Hz, 3H). |
| 86 | | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.50 - 7.47 (m, 1H), 7.41 - 7.37 (m, 1H), 5.02 (d, <i>J</i> = 16.5 Hz, 1H), 4.77 (d, <i>J</i> = 16.2 Hz, 1H), 4.29 (s, 1H), 4.04 - 3.99 (m, 1H), 1.41 (d, <i>J</i> = 6.6 Hz, 3H). |
| 87 | | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.52 - 7.48 (m, 1H), 7.43 - 7.39 (m, 2H), 4.90 - 4.73 (m, 2H), 4.33 - 4.29 (m, 1H), 4.23 (d, <i>J</i> = 2.1 Hz, 1H), 1.29 (d, <i>J</i> = 6.9 Hz, 3H). |
| 88 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.40 (brs, 3H), 7.43 (dd, <i>J</i> = 8.4 Hz, 1.5 Hz, 1H), 7.29 - 7.25 (m, 2H), 4.87 (d, <i>J</i> = 12 Hz, 1H), 4.76 (d, <i>J</i> = 11.7 Hz, 1H), 4.27 (s, 1H), 4.00 - 3.94 (m, 1H), 1.33 (d, <i>J</i> = 4.8 Hz, 3H). |
| 89 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.64 (brs, 3H), 7.53 (d, <i>J</i> = 8.0 Hz, 1H), 7.25 (dd, <i>J</i> = 1.8 Hz, 8.0 Hz, 2H), 4.81 - 4.71 (m, 2H), 4.21 (d, <i>J</i> = 3.0 Hz, 1H), 4.11 - 4.08 (m, 1H), 1.31 (t, <i>J</i> = 4.8 Hz, 3H). |


| Example | Structure | Characterization Data |
|---------|---|---|
| 90 |  | MS (ESI): <i>m/z</i> 214 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 8.13 (d, <i>J</i> = 8.0 Hz, 1H), 8.00 (d, <i>J</i> = 8.0 Hz, 2H), 7.72 - 7.70 (m, 1H), 7.64 - 7.62 (m, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 5.04 (s, 2H), 4.97 (s, 1H), 4.59 - 4.57 (m, 1H), 1.35 (d, <i>J</i> = 4.0 Hz, 3H). |
| 91 |  | MS (ESI): <i>m/z</i> 214 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 8.10 (d, <i>J</i> = 8.0 Hz, 1H), 7.99 (d, <i>J</i> = 8.0 Hz, 2H), 7.72 - 7.70 (m, 1H), 7.64 - 7.62 (m, 1H), 7.31 (d, <i>J</i> = 8.0 Hz, 1H), 5.13 - 5.10 (m, 2H), 5.02 (s, 1H), 4.20 - 4.18 (m, 1H), 1.55 (d, <i>J</i> = 4.0 Hz, 3H). |
| 92 |  | MS (ESI): <i>m/z</i> 232 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.64 (d, <i>J</i> = 6.3 Hz, 1H), 7.20 (d, <i>J</i> = 6.3 Hz, 1H), 4.84 (s, 2H), 4.51 (s, 1H), 4.45 - 4.40 (m, 1H), 1.27 (d, <i>J</i> = 6.3 Hz, 3H). |
| 93 |  | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.43 (d, <i>J</i> = 8.4 Hz, 1H), 7.37 - 7.33 (m, 1H), 7.25 - 7.19 (m, 1H), 4.97 - 4.80 (m, 2H), 4.28 - 4.25 (m, 1H), 4.05 - 3.98 (m, 1H), 1.39 (d, <i>J</i> = 6.6 Hz, 3H). |
| 94 |  | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.44 - 7.35 (m, 2H), 7.26 (s, 1H), 4.82 (s, 2H), 4.31 - 4.23 (m, 1H), 4.24 - 4.18 (m, 1H), 1.31 (d, <i>J</i> = 6.6 Hz, 3H). |
| 95 |  | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.29 (d, <i>J</i> = 7.8 Hz, 1H), 7.16 (d, <i>J</i> = 7.8 Hz, 1H), 7.00 (s, 1H), 4.92 (d, <i>J</i> = 15.6 Hz, 1H), 4.82 (d, <i>J</i> = 15.6 Hz, 1H), 4.20 (s, 1H), 4.04 - 3.91 (m, 1H), 2.34 (s, 3H), 1.38 (d, <i>J</i> = 6.6 Hz, 3H). |
| 96 |  | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, CD ₃ OD) δ 7.31 (d, <i>J</i> = 7.8 Hz, 1H), 7.16 (d, <i>J</i> = 7.8 Hz, 1H), 7.00 (s, 1H), 4.80 (s, 2H), 4.28 - 4.21 (m, 1H), 4.13 (s, 1H), 2.34 (s, 3H), 1.29 (d, <i>J</i> = 6.9 Hz, 3H). |
| 97 |  | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.23 (brs, 3H), 7.31 - 7.20 (m, 3H), 4.82 (d, <i>J</i> = 15.9 Hz, 1H), 4.69 (d, <i>J</i> = 15.9 Hz, 2H), 4.22 - 4.21 (m, 1H), 3.97 - 3.90 (m, 1H), 2.14 (s, 3H), 1.31 (d, <i>J</i> = 6.6 Hz, 3H). |
| 98 |  | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.54 (brs, 3H), 7.42 (d, <i>J</i> = 8.4 Hz, 1H), 7.27 - 7.18 (m, 2H), 4.68 (s, 2H), 4.20 - 4.12 (m, 2H), 2.15 (s, 3H), 1.24 (d, <i>J</i> = 6.3 Hz, 3H). |

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 194 | | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.50 (d, <i>J</i> = 2.1 Hz, 1H), 7.43 (dd, <i>J</i> = 8.3, 2.1 Hz, 1H), 7.20 (d, <i>J</i> = 8.3 Hz, 1H), 4.95 (d, <i>J</i> = 15.8 Hz, 1H), 4.83 (d, <i>J</i> = 15.8 Hz, 1H), 4.28 (s, 1H), 4.03 (m, 1H), 1.40 (d, <i>J</i> = 6.6 Hz, 3H). |
| 196 | | MS (ESI): <i>m/z</i> 218 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.11 (d, <i>J</i> = 7.9 Hz, 1H), 6.89 (d, <i>J</i> = 7.8 Hz, 1H), 4.90 (d, <i>J</i> = 15.5 Hz, 1H), 4.83 (d, <i>J</i> = 15.5 Hz, 1H), 4.34 (s, 1H), 3.96 (m, 1H), 2.86-2.73 (m, 4H), 1.92-1.72 (m, 4H), 1.39 (d, <i>J</i> = 6.6 Hz, 3H). |
| 197 | | MS (ESI): <i>m/z</i> 218 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.11 (d, <i>J</i> = 7.9 Hz, 1H), 6.89 (d, <i>J</i> = 7.8 Hz, 1H), 4.90 (d, <i>J</i> = 15.5 Hz, 1H), 4.83 (d, <i>J</i> = 15.5 Hz, 1H), 4.34 (s, 1H), 3.96 (m, 1H), 2.86-2.73 (m, 4H), 1.92-1.72 (m, 4H), 1.39 (d, <i>J</i> = 6.6 Hz, 3H). |
| 212 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, D ₂ O) δ 7.42 (dd, <i>J</i> = 8.6, 5.5 Hz, 1H), 7.08 (td, <i>J</i> = 8.7, 2.6 Hz, 1H), 6.94 (dd, <i>J</i> = 9.4, 2.5 Hz, 1H), 4.99-4.86 (m, 2H), 4.35 (s, 1H), 4.16-4.08 (m, 1H), 1.37 (d, <i>J</i> = 6.7 Hz, 3H). |
| 213 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, D ₂ O) δ 7.42 (dd, <i>J</i> = 8.6, 5.5 Hz, 1H), 7.10 (td, <i>J</i> = 8.7, 2.7 Hz, 1H), 6.95 (dd, <i>J</i> = 9.3, 2.6 Hz, 1H), 4.89-4.82 (m, 2H), 4.42-4.35 (m, 1H), 4.29 (d, <i>J</i> = 1.7 Hz, 1H), 1.26 (d, <i>J</i> = 6.7 Hz, 3H). |
| 214 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.45-7.38 (m, 1H), 7.30 (d, <i>J</i> = 7.5 Hz, 1H), 7.24-7.11 (m, 1H), 4.88-4.78 (m, 2H), 4.35-4.21 (m, 2H), 1.32 (d, <i>J</i> = 6.7 Hz, 3H). |
| 215 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.45-7.35 (m, 1H), 7.27 (d, <i>J</i> = 7.5 Hz, 1H), 7.22-7.12 (m, 1H), 5.05 (d, <i>J</i> = 16.1 Hz, 1H), 4.83 (d, <i>J</i> = 16.1 Hz, 1H), 4.31 (s, 1H), 4.07-4.00 (m, 1H), 1.41 (d, <i>J</i> = 6.6 Hz, 3H). |
| 216 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.41-7.43 (m, 1H), 7.14 (t, <i>J</i> = 9.0 Hz, 1H), 7.06 (d, <i>J</i> = 7.7 Hz, 1H), 4.86 (s, 2H), 4.45-4.27 (m, 2H), 1.30 (d, <i>J</i> = 6.7 Hz, 3H). |
| 217 | | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.50-7.42 (m, 1H), 7.12 (t, <i>J</i> = 9.0 Hz, 1H), 7.04 (d, <i>J</i> = 7.8 Hz, 1H), 4.99 (d, <i>J</i> = 15.9 Hz, 1H), 4.86 (d, <i>J</i> = 16.0 Hz, 1H), 4.49 (s, 1H), 4.04-3.97 (m, 1H), 1.42 (d, <i>J</i> = 6.6 Hz, 3H). |
| 219 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.28-7.25 (m, 2H), 7.09 (d, <i>J</i> = 8.5 Hz, 1H), 4.94 (d, <i>J</i> = 12.3 Hz, 1H), 4.83 (d, <i>J</i> = 15.4 Hz, 1H), 4.21 (brs, 1H), 4.07-3.98 (m, 1H), 2.71- |

| Example | Structure | Characterization Data |
|---------|-----------|---|
| | | 2.63 (m, 2H), 1.39 (d, $J = 6.6$ Hz, 3H), 1.25 (t, $J = 7.6$ Hz, 3H). |
| 220 | | MS (ESI): m/z 192 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.28-7.25 (m, 2H), 7.10 (d, $J = 8.4$ Hz, 1H), 4.81 (s, 2H), 4.29-4.21 (m, 1H), 4.15 (d, $J = 2.5$ Hz, 1H), 2.71-2.63 (m, 2H), 1.31 (d, $J = 6.7$ Hz, 3H), 1.24 (t, $J = 7.8$ Hz, 3H). |
| 225 | | MS (ESI): m/z 212 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.35 (d, $J = 8.2$ Hz, 1H), 7.23 (d, $J = 8.2$ Hz, 1H), 4.99 (d, $J = 12.1$ Hz, 1H), 4.76 (d, $J = 16.4$ Hz, 1H), 4.38 (s, 1H), 4.00-3.94 (m, 1H), 2.39 (s, 3H), 1.41 (d, $J = 6.6$ Hz, 3H). |
| 228 | | MS (ESI): m/z 232 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.81 (s, 1H), 7.73 (d, $J = 8.1$ Hz, 1H), 7.42 (d, $J = 8.1$ Hz, 1H), 4.99 (d, $J = 16.5$ Hz, 1H), 4.90 (d, $J = 16.5$ Hz, 1H), 4.39 (brs, 1H), 4.11-4.03 (m, 1H), 1.42 (d, $J = 6.6$ Hz, 3H). |
| 229 | | MS (ESI): m/z 232 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.82 (s, 1H), 7.73 (d, $J = 8.1$ Hz, 1H), 7.42 (d, $J = 8.1$ Hz, 1H), 4.91 (s, 2H), 4.35-4.28 (m, 2H), 1.30 (d, $J = 6.6$ Hz, 3H). |
| 230 | | MS (ESI): m/z 194 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.39 (t, $J = 8.1$ Hz, 1H), 6.95 (d, $J = 8.1$ Hz, 1H), 6.77 (d, $J = 7.8$ Hz, 1H), 4.91 (d, $J = 15.3$ Hz, 1H), 4.81 (d, $J = 15.6$ Hz, 1H), 4.38 (brs, 1H), 4.03-3.93 (m, 1H), 3.93 (s, 3H), 1.40 (d, $J = 6.7$ Hz, 3H). |
| 231 | | MS (ESI): m/z 194 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.39 (t, $J = 8.0$ Hz, 1H), 6.97 (d, $J = 8.3$ Hz, 1H), 6.78 (d, $J = 7.8$ Hz, 1H), 4.86-4.70 (m, 2H), 4.38-4.19 (m, 2H), 3.92 (s, 3H), 1.30 (d, $J = 6.7$ Hz, 3H). |
| 232 | | MS (ESI): m/z 194 [M+H]. 1 H NMR (300 MHz, D ₂ O) δ 7.35 (d, $J = 8.6$ Hz, 1H), 6.95 (dd, $J = 8.6$, 2.6 Hz, 1H), 6.79 (d, $J = 2.4$ Hz, 1H), 4.83 (brs, 2H), 4.41-4.33 (m, 1H), 4.26 (s, 1H), 3.80 (s, 3H), 1.26 (d, $J = 6.7$ Hz, 3H). |
| 233 | | MS (ESI): m/z 194 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.33 (d, $J = 8.6$ Hz, 1H), 6.92 (dd, $J = 8.6$, 2.6 Hz, 1H), 6.76 (d, $J = 2.5$ Hz, 1H), 4.90-4.85 (m, 2H), 4.29 (s, 1H), 4.13-4.06 (m, 1H), 3.78 (s, 3H), 1.35 (d, $J = 6.7$ Hz, 3H). |
| 234 | | MS (ESI): m/z 194 [M+H]. 1 H NMR (300 MHz, CD ₃ OD) δ 7.34 (t, $J = 8.0$ Hz, 1H), 7.01 (dd, $J = 8.0$, 4.2 Hz, 2H), 4.95 (d, $J = 16.2$ Hz, 1H), 4.69 (d, $J = 16.3$ Hz, 1H), 4.22 (s, 1H), 4.02-3.95 (m, 1H), 3.84 (s, 3H), 1.39 (d, $J = 6.6$ Hz, 3H). |

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 235 | | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.36 (t, <i>J</i> = 8.0 Hz, 1H), 7.03 (dd, <i>J</i> = 7.9, 3.4 Hz, 2H), 4.80 (d, <i>J</i> = 16.8 Hz, 1H), 4.70 (d, <i>J</i> = 16.5 Hz, 1H), 4.26-4.21 (m, 1H), 4.15 (s, 1H), 3.85 (s, 3H), 1.29 (d, <i>J</i> = 6.7 Hz, 3H). |
| 237 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.36 (t, <i>J</i> = 8.0 Hz, 1H), 7.03 (dd, <i>J</i> = 7.9, 3.4 Hz, 2H), 4.80 (d, <i>J</i> = 16.8 Hz, 1H), 4.70 (d, <i>J</i> = 16.5 Hz, 1H), 4.26-4.21 (m, 1H), 4.15 (s, 1H), 3.85 (s, 3H), 1.29 (d, <i>J</i> = 6.7 Hz, 3H). |
| 238 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.30 (s, 3H), 7.27 (s, 1H), 6.92 (s, 1H), 4.81-4.67 (m, 2H), 4.13-4.12 (m, 1H), 3.98-3.83 (m, 1H), 2.21 (s, 6H), 1.31 (d, <i>J</i> = 6.6 Hz, 3H). |
| 200 | | MS (ESI): <i>m/z</i> 212 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.38 (d, <i>J</i> = 8.2 Hz, 1H), 7.26 (d, <i>J</i> = 8.0 Hz, 1H), 4.90 (d, <i>J</i> = 17.0 Hz, 1H), 4.72 (d, <i>J</i> = 17.0 Hz, 1H), 4.47-4.30 (m, 2H), 2.42 (s, 3H), 1.22 (d, <i>J</i> = 6.7 Hz, 3H). |
| 248 | | MS (ESI): <i>m/z</i> 220 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.15 (br.s., 3H), 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.80 (d, <i>J</i> = 8.4 Hz, 1H), 4.82 (d, <i>J</i> = 15.3 Hz, 1H), 4.71 (d, <i>J</i> = 15.0 Hz, 1H), 4.23 (s, 1H), 4.11-4.08 (m, 2H), 3.89-3.86 (m, 1H), 2.79-2.73 (m, 2H), 1.99-1.92 (m, 2H), 1.31 (d, <i>J</i> = 6.6 Hz, 3H). |
| 249 | | MS (ESI): <i>m/z</i> 220 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 6.89 (d, <i>J</i> = 8.4 Hz, 1H), 6.83 (d, <i>J</i> = 8.7 Hz, 1H), 4.78 (s, 2H), 4.38-4.2 (m, 1H), 4.27 (s, 1H), 4.16 (t, <i>J</i> = 5.4 Hz, 2H), 2.82 (t, <i>J</i> = 6.3 Hz, 2H), 2.12-2.00 (m, 2H), 1.24 (d, <i>J</i> = 6.9 Hz, 3H). |

[0161] **Example 99. *trans*-3-Ethyl-6-methylisochroman-4-amine hydrochloride**



The compound of **Example 99** was prepared as previously described in Example 76, using pent-1-en-3-ol. MS (ESI): *m/z* 192 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.39 (brs, 3H), 7.34 (s, 1H), 7.19 (d, *J* = 7.8 Hz, 1H), 7.06 (d, *J* = 7.8 Hz, 1H), 4.69 (s, 2H), 4.18 (s, 1H), 3.90 - 3.81 (m, 1H), 2.31 (s, 3H), 1.77 - 1.51 (m, 2H), 0.96 (t, *J* = 5.4 Hz, 3H).

[0162] **Example 100. *cis*-3-Ethyl-6-methylisochroman-4-amine hydrochloride**



The compound of **Example 100** was prepared as previously described in Example 76, using pent-1-en-3-ol. MS (ESI): *m/z* 192 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.18 (brs, 3H), 7.29 (s, 1H), 7.21 (d, *J* = 7.8 Hz, 1H), 7.07 (d, *J* = 7.8 Hz, 1H), 4.87 (d, *J* = 11.7 Hz, 1H), 4.73 (d, *J* = 11.7 Hz, 1H), 4.35 (s, 1H), 3.65 - 3.62 (m, 1H), 2.31 (s, 3H), 1.66 - 1.59 (m, 2H), 0.99 (t, *J* = 5.4 Hz, 3H).


[0163] **Table 9.** Compounds prepared as described in Examples 99 and 100 using the appropriately substituted starting materials.

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 198 | | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.43-7.39 (m, 2H), 7.37-7.28 (m, 1H), 7.19 (d, <i>J</i> = 7.7 Hz, 1H), 5.00 (d, <i>J</i> = 15.6 Hz, 1H), 4.87 (d, <i>J</i> = 15.6 Hz, 1H), 4.31 (s, 1H), 3.79-3.69 (m, 1H), 1.78-1.61 (m, 2H), 1.12 (t, <i>J</i> = 7.4 Hz, 3H). |
| 199 | | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.43-7.32 (m, 3H), 7.19 (d, <i>J</i> = 7.2 Hz, 1H), 4.82 (s, 2H), 4.22 (d, <i>J</i> = 2.2 Hz, 1H), 4.00-3.95 (m, 1H), 1.71-1.46 (m, 2H), 1.07 (t, <i>J</i> = 7.4 Hz, 3H). |
| 201 | | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.27-7.13 (m, 3H), 4.99 (d, <i>J</i> = 15.5 Hz, 1H), 4.82 (d, <i>J</i> = 15.4 Hz, 1H), 4.34 (s, 1H), 3.75-3.69 (m, 1H), 1.79-1.58 (m, 2H), 1.11 (t, <i>J</i> = 7.4 Hz, 3H). |
| 202 | | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.30-7.11 (m, 3H), 4.77 (s, 2H), 4.25 (d, <i>J</i> = 2.3 Hz, 1H), 4.00-3.94 (m, 1H), 1.76-1.45 (m, 2H), 1.07 (t, <i>J</i> = 7.4 Hz, 3H). |
| 207 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.30 (t, <i>J</i> = 7.6 Hz, 1H), 7.19 (d, <i>J</i> = 7.3 Hz, 1H), 7.00 (d, <i>J</i> = 7.6 Hz, 1H), 4.88 (d, <i>J</i> = 16.2 Hz, 1H), 4.78 (d, <i>J</i> = 16.2 Hz, 1H), 4.37 (d, <i>J</i> = 1.3 Hz, 1H), 4.08-4.03 (m, 1H), 2.42 (s, 3H), 1.66 - 1.38 (m, 2H), 1.05 (t, <i>J</i> = 7.4 Hz, 3H). |

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 208 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.30 (t, <i>J</i> = 7.6 Hz, 1H), 7.20 (d, <i>J</i> = 7.2 Hz, 1H), 7.00 (d, <i>J</i> = 7.6 Hz, 1H), 5.01 (d, <i>J</i> = 15.7 Hz, 1H), 4.85 (d, <i>J</i> = 15.8 Hz, 1H), 4.42 (s, 1H), 3.72-3.67 (m, 1H), 2.42 (s, 3H), 1.81-1.59 (m, 2H), 1.12 (t, <i>J</i> = 7.4 Hz, 3H). |
| 209 | | MS (ESI): <i>m/z</i> 212 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.49-7.39 (m, 2H), 7.19-7.16 (m, 1H), 4.83 (d, <i>J</i> = 6.5 Hz, 2H), 4.48 (brs, 1H), 4.13-4.07 (m, 1H), 1.67-1.39 (m, 2H), 1.06 (t, <i>J</i> = 7.4 Hz, 3H). |
| 210 | | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.25-7.22 (m, 2H), 7.06 (d, <i>J</i> = 8.1 Hz, 1), 4.79 (s, 2H), 4.37 (d, <i>J</i> = 1.9 Hz, 1H), 3.59 (dd, <i>J</i> = 9.7, 2.0 Hz, 1H), 2.36 (s, 3H), 1.82-1.72 (m, 1H), 1.09 (d, <i>J</i> = 6.5 Hz, 3H), 0.94 (d, <i>J</i> = 6.6 Hz, 3H). |
| 211 | | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.25-7.22 (m, 2H), 7.07 (d, <i>J</i> = 8.4 Hz, 1H), 4.98 (d, <i>J</i> = 15.3 Hz, 1H), 4.80 (d, <i>J</i> = 15.3 Hz, 1H), 4.43 (s, 1H), 3.36-3.31 (m, 1H), 2.36 (s, 3H), 1.86-1.76 (m, 1H), 1.15 (d, <i>J</i> = 6.4 Hz, 3H), 1.04 (d, <i>J</i> = 6.7 Hz, 3H). |
| 218 | | MS (ESI): <i>m/z</i> 212 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.49-7.38 (m, 2H), 7.22-7.15 (m, 1H), 5.02 (d, <i>J</i> = 15.9 Hz, 1H), 4.85 (d, <i>J</i> = 16.2 Hz, 1H), 4.55 (brs, 1H), 3.76-3.68 (m, 1H), 1.79-1.66 (m, 2H), 1.13 (t, <i>J</i> = 7.4 Hz, 3H). |
| 223 | | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.46-7.30 (m, 3H), 7.19 (d, <i>J</i> = 7.2 Hz, 1H), 4.98 (d, <i>J</i> = 15.6 Hz, 1H), 4.86 (d, <i>J</i> = 15.6 Hz, 1H), 4.26 (brs, 1H), 3.95-3.90 (m, 1H), 1.97-1.89 (m, 1H), 1.70-1.60 (m, 1H), 1.48-1.39 (m, 1H), 1.03 (d, <i>J</i> = 6.6 Hz, 3H), 1.00 (d, <i>J</i> = 6.6 Hz, 3H). |
| 224 | | MS (ESI): <i>m/z</i> 206 [M+H]. ¹ H NMR (300 MHz, CD ₃ OD) δ 7.49-7.29 (m, 3H), 7.19 (d, <i>J</i> = 7.6 Hz, 1H), 4.82 (s, 2H), 4.30-4.13 (m, 2H), 1.94-1.75 (m, 1H), 1.65-1.55 (m, 1H), 1.28-1.19 (m, 1H), 0.98 (t, <i>J</i> = 6.3 Hz, 6H). |
| 250 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, DMSO-d ₆) δ 8.19 (brs, 3H), 7.47 (d, <i>J</i> = 7.5 Hz, 1H), 7.40-7.32 (m, 2H), 7.18 (d, <i>J</i> = 7.5 Hz, 1H), 4.90 (d, <i>J</i> = 15.6 Hz, 1H), 4.79 (d, <i>J</i> = 15.6 Hz, 1H), 4.28 (s, 1H), 3.78 (m, 1H), 1.60-1.56 (m, 2H), 1.52-1.44 (m, 2H), 0.95 (t, <i>J</i> = 7.2 Hz, 3H). |

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 251 | | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.42 (brs, 3H), 7.52 (d, <i>J</i> = 7.5 Hz, 1H), 7.38-7.33 (m, 2H), 7.18 (d, <i>J</i> = 7.5 Hz, 1H), 4.73 (m, 2H), 4.20 (s, 1H), 4.00 (m, 1H), 1.52-1.47 (m, 4H), 0.91 (t, <i>J</i> = 6.9 Hz, 3H). |

[0164] **Example 101. 3,3-Dimethylisochroman-4-amine hydrochloride**




The compound of **Example 101** was prepared as previously described in Example 76, using 2-methylbut-3-en-2-ol. MS (ESI): *m/z* 178 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.35 (brs, 3H), 7.49 (d, *J* = 6.9 Hz, 1H), 7.41 - 7.30 (m, 2H), 7.18 (d, *J* = 7.2 Hz, 1H), 4.86 - 4.71 (m, 2H), 4.14 (s, 1H), 1.33 (s, 3H), 1.19 (s, 3H).

[0165] **Example 104. *cis*-1-Methylisochroman-4-amine hydrochloride**

[0166] **Example 105. *trans*-1-Methylisochroman-4-amine hydrochloride**



Example 104



Example 105

[0167] The compounds are **Example 104** and **Example 105** were prepared as described previously in Example 76 using 1-(1-(allyloxy)ethyl)-2-bromobenzene (prepared as previously described in Example 21 using 1-(2-bromophenyl)ethanol).

[0168] **Example 104. *cis*-1-Methylisochroman-4-amine hydrochloride** MS (ESI): *m/z* 164 [M+H]. ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.66 (brs, 3H), 7.58 (d, *J* = 5.4 Hz, 1H), 7.41 - 7.31 (m, 2H), 7.25 (d, *J* = 5.7 Hz, 1H), 4.98 (q, *J* = 5.1 Hz, 1H), 4.35 (t, *J* = 2.7, 1H), 4.17 (dd, *J* = 2.7 Hz, 8.7 Hz, 1H), 3.89 (dd, *J* = 3 Hz, 9 Hz, 1H), 1.46 (d, *J* = 5.1 Hz, 3H).


[0169] **Example 105. *trans*-(1*S*,4*S*)-1-Methylisochroman-4-amine hydrochloride.** MS (ESI): *m/z* 164 [M+H]. ^1H NMR (300 MHz, DMSO-*d*₆) δ 8.54 (brs, 3H), 7.51 (d, *J* = 7.8 Hz, 1H), 7.41 - 7.28 (m, 3H), 4.78 (q, *J* = 6.3 Hz, 1H), 4.31 - 4.29 (m, 1H), 4.21 (d, *J* = 12.6 Hz, 1H), 3.87 (dd, *J* = 2.1 Hz, 12.6 Hz, 1H), 1.53 (d, *J* = 6.6 Hz, 3H).

[0170] **Table 10. Compounds prepared as described in Examples 104 and 105 using the appropriately substituted starting materials.**

| Example | Structure | Characterization Data |
|---------|-----------|---|
| 106 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, CDCl ₃) δ 8.67 (brs, 3H), 7.43 (s, 1H), 7.12 (d, <i>J</i> = 8.1 Hz, 1H), 6.95 (d, <i>J</i> = 7.8 Hz, 1H), 5.04 - 4.98 (m, 1H), 4.30 (s, 1H), 4.10 (s, 2H), 2.30 (s, 3H), 1.45 (d, <i>J</i> = 6.6 Hz, 3H). |
| 107 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, CDCl ₃) δ 8.69 (brs, 3H), 7.46 (s, 1H), 7.14 (d, <i>J</i> = 8.1 Hz, 1H), 7.01 (d, <i>J</i> = 8.1 Hz, 1H), 4.80 - 4.74 (m, 1H), 4.50 (d, <i>J</i> = 12.3 Hz, 1H), 4.30 (s, 1H), 3.80 (dd, <i>J</i> = 1.8 Hz, 12.6 Hz, 1H), 2.30 (s, 3H), 1.59 (d, <i>J</i> = 6.6 Hz, 3H). |
| 203 | | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.60 (brs, 3H), 7.67 (d, <i>J</i> = 2.1 Hz, 1H), 7.46 (dd, <i>J</i> = 8.4, 2.2 Hz, 1H), 7.30 (d, <i>J</i> = 8.5 Hz, 1H), 4.97-4.93 (m, 1H), 4.39 (t, <i>J</i> = 3.7 Hz, 1H), 4.14 (dd, <i>J</i> = 12.4, 3.5 Hz, 1H), 3.85 (dd, <i>J</i> = 12.3, 4.2 Hz, 1H), 1.43 (d, <i>J</i> = 6.7 Hz, 3H). |
| 204 | | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.49 (brs, 3H), 7.61 (d, <i>J</i> = 2.2 Hz, 1H), 7.48 (dd, <i>J</i> = 8.4, 2.2 Hz, 1H), 7.34 (d, <i>J</i> = 8.5 Hz, 1H), 4.79-4.73 (m, 1H), 4.35 (s, 1H), 4.16 (d, <i>J</i> = 12.1 Hz, 1H), 3.86 (dd, <i>J</i> = 12.8, 2.3 Hz, 1H), 1.52 (d, <i>J</i> = 6.5 Hz, 3H). |
| 205 | | MS (ESI): <i>m/z</i> 182 [M+H]. ^1H NMR (300 MHz, D ₂ O) δ 7.26-7.13 (m, 3H), 5.16-5.09 (m, 1H), 4.43 (s, 1H), 4.26 (dd, <i>J</i> = 13.1, 2.3 Hz, 1H), 3.98 (dd, <i>J</i> = 13.2, 2.1 Hz, 1H), 1.47 (d, <i>J</i> = 6.8 Hz, 3H). |


| Example | Structure | Characterization Data |
|---------|-----------|---|
| 206 | | MS (ESI): <i>m/z</i> 182 [M+H]. ^1H NMR (300 MHz, D_2O) δ 7.32-7.27 (m, 1H), 7.22-7.12 (m, 2H), 4.92-4.84 (m, 1H), 4.42 (s, 1H), 4.22 (d, <i>J</i> = 13.2 Hz, 1H), 4.02 (dd, <i>J</i> = 13.1, 2.0 Hz, 1H), 1.53 (d, <i>J</i> = 6.5 Hz, 3H). |
| 221 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, D_2O) δ 7.30 (t, <i>J</i> = 7.7 Hz, 1H), 7.17 (d, <i>J</i> = 7.4 Hz, 1H), 7.02 (d, <i>J</i> = 7.7 Hz, 1H), 5.18-5.10 (m, 1H), 4.54 (s, 1H), 4.18 (dd, <i>J</i> = 13.2, 1.6 Hz, 1H), 4.01 (dd, <i>J</i> = 13.2, 1.5 Hz, 1H), 2.33 (s, 3H), 1.44 (d, <i>J</i> = 6.8 Hz, 3H). |
| 222 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, D_2O) δ 7.32 (t, <i>J</i> = 7.7 Hz, 1H), 7.19 (d, <i>J</i> = 7.5 Hz, 1H), 7.10 (d, <i>J</i> = 7.8 Hz, 1H), 4.94-4.89 (m, 1H), 4.55 (s, 1H), 4.19 (d, <i>J</i> = 12.9 Hz, 1H), 3.93 (dd, <i>J</i> = 12.9, 1.3 Hz, 1H), 2.33 (s, 3H), 1.52 (d, <i>J</i> = 6.5 Hz, 3H). |
| 226 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, D_2O) δ 7.30 (t, <i>J</i> = 7.7 Hz, 1H), 7.18 (d, <i>J</i> = 7.6 Hz, 1H), 7.02 (d, <i>J</i> = 7.7 Hz, 1H), 5.18-5.11 (m, 1H), 4.54 (s, 1H), 4.18 (dd, <i>J</i> = 13.2, 1.6 Hz, 1H), 4.01 (dd, <i>J</i> = 13.2, 1.6 Hz, 1H), 2.33 (s, 3H), 1.44 (d, <i>J</i> = 6.8 Hz, 3H). |
| 227 | | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (300 MHz, D_2O) δ 7.32 (t, <i>J</i> = 7.7 Hz, 1H), 7.19 (d, <i>J</i> = 7.5 Hz, 1H), 7.11 (d, <i>J</i> = 7.7 Hz, 1H), 4.95-4.88 (m, 1H), 4.55 (s, 1H), 4.21 (dd, <i>J</i> = 12.9, 1.4 Hz, 1H), 3.94 (dd, <i>J</i> = 12.9, 1.7 Hz, 1H), 2.33 (s, 3H), 1.52 (d, <i>J</i> = 6.5 Hz, 3H). |

[0171] **Example 236. *trans*-N,1-dimethylisochroman-4-amine hydrochloride**




[0172] The compound of **Example 236** was prepared as previously described in Example 57. MS (ESI): *m/z* 178 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.26 (s, 1H), 9.11 (s, 1H), 7.54 (d, *J* = 7.5 Hz, 1H), 7.45 (d, *J* = 7.5 Hz, 1H), 7.38-7.31 (m, 2H), 4.82-4.80 (m, 1H), 4.40 (d, *J* = 13.1 Hz, 1H), 4.27 (brs, 1H), 3.86 (dd, *J* = 13.1, 2.0 Hz, 1H), 2.56 (t, *J* = 5.4 Hz, 3H), 1.54 (d, *J* = 6.5 Hz, 3H).

[0173] **Example 239. *cis*-N,N,1-trimethylisochroman-4-amine hydrochloride**



[0174] The compound of **Example 239** was prepared as previously described in Example 60. MS (ESI): m/z 192 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.59-7.47 (m, 2H), 7.44-7.37 (m, 2H), 4.90-4.84 (m, 1H), 4.64 (d, J = 14.1 Hz, 1H), 4.40 (s, 1H), 3.97 (dd, J = 14.1, 2.4 Hz, 1H), 2.95 (s, 3H), 2.90 (s, 3H), 1.61 (d, J = 6.5 Hz, 3H).


[0175] **Example 240. cis-N,1-dimethylisochroman-4-amine hydrochloride**



[0176] The compound of **Example 240** was prepared as previously described in Example 57. MS (ESI): m/z 178 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.51-7.41 (m, 2H), 7.38-7.33 (m, 1H), 7.26 (d, J = 7.7 Hz, 1H), 5.18-5.11 (m, 1H), 4.23-4.11 (m, 3H), 2.74 (s, 3H), 1.52 (d, J = 6.8 Hz, 3H).

[0177] **Example 256. cis-3-Isopropylisochroman-4-amine hydrochloride**

[0178] **Example 257. trans-3-Isopropylisochroman-4-amine hydrochloride**



Step 1. A solution of 2-bromo-3-methylbutanoic acid (5 g, 27.6 mmol) in 100 mL of CH_2Cl_2 was chilled in an ice bath and treated with 1,1-carbonyldiimidazole (4.68 g, 28.9 mmol). The ice bath was removed and the reaction mixture was stirred at ambient temperature for 15 min. *N,O*-dimethylhydroxylamine (1.76 g, 28.9 mmol) was added to the reaction mixture and stirring at ambient temperature was maintained for 2 h. Water (100 mL) was added to the reaction vessel and the resulting biphasic mixture was transferred to a separatory funnel. The organic phase was washed with 10% aqueous citric acid (2x100 mL), saturated aqueous NaHCO_3 (100 mL), and saturated aqueous NaCl (2x100 mL). The organic phase was dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to provide 2-bromo-N-methoxy-N,3-dimethylbutanamide **256.1** (2.80 g, crude) as a yellow oil. GC-MS: *m/z* 223, 225 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 4.51 (d, *J* = 9.3 Hz, 1H), 3.78 (s, 3H), 3.24 (s, 3H), 2.36-2.28 (m, 1H), 1.16 (d, *J* = 6.9 Hz, 3H), 0.98 (d, *J* = 6.9 Hz, 3H).

Step 2. A solution of (2-bromophenyl)methanol (3.47 g, 18.6 mmol) in DMF (100 mL) was cooled in an ice bath and treated with sodium hydride (743 mg, 18.6 mmol, 95% in mineral oil). The resulting slurry was stirred for 30 min and then treated with Compound **256.1** (2.8 g, 12.4 mmol). The reaction was stirred at ambient temperature for 1 h and then partitioned between H_2O (100 mL) and EtOAc (100 mL). The aqueous phase was extracted

with EtOAc (2x100 mL). The combined organics were washed with saturated aqueous NaCl (2x100 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, 100% hexanes gradient to 10% EtOAc/hexanes (90%)) to provide 2-((2-bromobenzyl)oxy)-N-methoxy-N,3-dimethylbutanamide **256.2** (4.05 g, 12.2 mmol) as a colorless oil. MS (ESI): *m/z* 330 [M+H]₊,

Step 3. A solution of Compound **256.2** (4.5 g, 13.6 mmol) in THF (100 mL) was chilled in a dry ice/acetone bath and treated with *tert*-butyllithium (33.9 mL, 1.7M in pentane). The reaction was stirred in the dry ice/acetone bath for 10 min. Excess base was quenched by the dropwise addition of saturated aqueous NH₄Cl. The mixture was partitioned between saturated aqueous NH₄Cl (100 mL) and EtOAc (100 mL). The aqueous phase was extracted with EtOAc (2x100 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting oil was purified by flash column chromatography (SiO₂, 100% hexanes) to afford 3-isopropylisochroman-4-one **256.3** (600 mg, 3.15 mmol) as a yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 8.03 (d, *J* = 7.8 Hz, 1H), 7.57-7.52 (m, 1H), 7.40 (t, *J* = 7.5 Hz, 1H), 7.21 (d, *J* = 7.8 Hz, 1H), 4.95 (d, *J* = 15.0 Hz, 1H), 4.86 (d, *J* = 15.0 Hz, 1H), 3.97-3.94 (m, 1H), 2.60-2.53 (m, 1H), 1.15 (d, *J* = 7.5 Hz, 3H), 0.98 (d, *J* = 7.5 Hz, 3H).

Step 4. 3-Isopropylisochroman-4-ol **256.4** was prepared as previously described in Example 1. GC-MS: *m/z* 192 [M+H]₊. ¹H NMR (300 MHz, CDCl₃) δ 7.54-7.52 (m, 0.4 H), 7.43-7.30 (m, 0.6H), 7.31-7.23 (m, 2H), 7.04-6.99 (m, 1H), 4.90-4.49 (m, 3H), 3.33 (dd, *J* = 7.5, 4.5 Hz, 0.4H), 3.08 (dd, *J* = 9.6, 1.2 Hz, 0.6H), 2.13-2.07 (m, 1H), 1.94 (d, *J* = 10.5 Hz, 0.6H), 1.81 (d, *J* = 9.0 Hz, 0.4H), 1.22-1.02 (m, 6H).

Step 5. 4-Azido-3-isopropylisochroman **256.5** was prepared as previously described in Example 1. ¹H NMR (300 MHz, CDCl₃) δ 7.46-7.43 (m, 1H), 7.34-7.27 (m, 2H), 7.06-7.03 (m, 1H), 4.83-4.72 (m, 2H), 4.33 (d, *J* = 7.5 Hz, 1H), 3.50-3.46 (m, 1H), 2.11-2.04 (m, 1H), 1.08-1.02 (m, 6H).

Step 6. Compounds **256.5a** and **256.5b** were prepared as previously described in Example 73.

tert-butyl-(*cis*-3-isopropylisochroman-4-yl)carbamate **256.5a** TLC higher R_f. ¹H NMR (300 MHz, CDCl₃) δ 7.49-7.46 (m, 1H), 7.25-7.22 (m, 2H), 7.00-6.97 (m, 1H),

5.02-4.72 (m, 4H), 3.17 (dd, J = 9.6, 1.8 Hz, 1H), 1.91-1.83 (m, 1H), 1.42 (s, 9H), 1.08 (d, J = 6.6 Hz, 3H), 0.99 (d, J = 6.6 Hz, 3H).

tert-butyl-(*trans*-3-isopropylisochroman -4-yl)carbamate **256.5b** TLC lower Rf. 1 H NMR (300 MHz, CDCl₃) δ 7.39-7.36 (m, 1H), 7.24-7.20 (m, 2H), 6.99-6.96 (m, 1H), 4.86-4.69 (m, 4H), 3.32 (dd, J = 6.9, 5.1 Hz, 1H), 1.94-1.92 (m, 1H), 1.50 (s, 9H), 1.05 (d, J = 6.9 Hz, 6H).


Step 7. The compounds of **Example 256** and **Example 257** were prepared as previously described in Example 1.

[0179] **Example 256.** MS (ESI): *m/z* 292 [M+H]. 1 H NMR (300 MHz, CD₃OD) δ 7.49-7.36 (m, 3H), 7.23 (d, J = 7.8 Hz, 1H), 5.07 (d, J = 15.6 Hz, 1H), 4.94 (d, J = 15.6 Hz, 1H), 4.53 (s, 1H), 3.40 (dd, J = 10.2, 1.2 Hz, 1H), 1.92-1.85 (m, 1H), 1.20 (d, J = 6.3 Hz, 3H), 1.09 (d, J = 6.6 Hz, 3H).

[0180] **Example 257.** MS (ESI): *m/z* 292 [M+H]. 1 H NMR (300 MHz, CD₃OD) δ 7.49-7.38 (m, 3H), 7.23 (d, J = 7.2 Hz, 1H), 4.88 (s, 2H), 4.47 (d, J = 2.1 Hz, 1H), 3.66 (dd, J = 9.9, 2.1 Hz, 1H), 1.87-1.80 (m, 1H), 1.14 (d, J = 6.6 Hz, 3H), 0.98 (d, J = 6.6 Hz, 3H).

[0156] **Example 273.** *cis*-3-Isopropylisochroman-4-amine hydrochloride

[0157] **Example 274.** *trans*-3-Isopropylisochroman-4-amine hydrochloride



Step 1. A solution of phenylacetylene (4 g, 39.1 mmol) in 50 mL of anhydrous THF was cooled in a dry ice/acetone bath under a N₂ atmosphere. Butyllithium (18.7 mL, 46.9 mmol) was added in a dropwise fashion, and the reaction was stirred at - 78 °C for 30 min. A solution of methyl 2,2-difluoroacetate (5.16 g, 46.9 mmol) in THF (5 mL) was added, followed by boron trifluoride etherate (5.91 mL, 46.9 mmol). The cold bath was removed, and the reaction was stirred at ambient temperature for 12 h. Saturated aqueous NH₄Cl (50 mL) was added to the reaction vessel and the reaction mixture was extracted with EtOAc (3x50 mL). The combined organics were washed with saturated aqueous NaCl (2x80 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. Purification by flash column chromatography (SiO₂, 100% hexane) afford 1,1-difluoro-4-phenylbut-3-yn-2-one **273.1** (5.50 g, 30.5 mmol) as a yellow oil. GC-MS: *m/z* 180 [M+H]. ¹H NMR (300 MHz, CDCl₃) δ 7.67 (d, *J* = 7.2 Hz, 1H), 7.57-7.52 (m, 1H), 7.46-7.41 (m, 2H), 5.88 (t, *J* = 54.0 Hz, 1H).

Step 2. Compound **273.1** (5.5 g, 30.5 mmol) was dissolved in THF (100 mL) and cooled in an ice bath. LiAl₄ (1.73 g, 45.7 mmol) was added slowly in portions and the

reaction mixture was stirred 0°C for 30 min. Excess hydride was quenched by the dropwise addition of H₂O (1.8 mL), followed by 1.8 mL of 15% aqueous NaOH. After stirring at room temperature for 0.5 hour, the solid was removed by filtration. The filtrate was concentrated to provide (E)-1,1-difluoro-4-phenylbut-3-en-2-ol **273.2** (5.30 g, 28.7 mmol) as a colorless oil. GC-MS: *m/z* 184 [M+H]. ¹H NMR (300 MHz, CDCl₃) δ 7.44-7.27 (m, 5H), 6.82 (d, *J* = 15.9 Hz, 1H), 6.25-6.18 (dd, *J* = 15.9, 6.3 Hz, 1H), 5.72 (td, *J* = 56.1, 4.2 Hz, 1H), 4.53-4.43 (m, 1H), 2.25 (d, *J* = 5.1 Hz, 1H).

Step 3. Compound **273.3** was prepared as previously described in Example 10.

¹H NMR (300 MHz, CDCl₃) δ 7.58-7.50 (m, 2H), 7.49-7.43 (m, 1H), 7.39-7.25 (m, 4H), 7.21-7.15 (m, 2H), 6.81 (d, *J* = 15.9 Hz, 1H), 6.18 (dd, *J* = 16.2, 7.5 Hz, 1H), 5.84-5.64 (m, 1H), 4.79-4.63 (m, 2H), 4.23-4.19 (m, 1H).

Step 4. Compound **273.4** was prepared as previously described in Example 10. ¹H NMR (300 MHz, CDCl₃) δ 7.66-7.62 (m, 1H), 7.44-7.38 (m, 3H), 7.35-7.29 (m, 3H), 7.25-7.16 (m, 1H), 7.08-7.05 (m, 1H), 6.06-5.67 (td, *J* = 54.9, 5.4 Hz, 1H), 5.20-5.11 (m, 1H), 5.01 (d, *J* = 15.3 Hz, 1H), 4.83 (d, *J* = 15.3 Hz, 1H).

Step 5. Compound **273.5** was prepared as previously described in Example 10. ¹H NMR (300 MHz, CDCl₃) δ 8.06 (d, *J* = 6.9 Hz, 1H), 7.65-7.59 (m, 1H), 7.48-7.43 (m, 1H), 7.29-7.24 (m, 1H), 6.37 (td, *J* = 53.4, 1.8 Hz, 1H), 5.14 (d, *J* = 15.3 Hz, 1H), 5.02 (d, *J* = 15.3 Hz, 1H), 4.51-4.41 (m, 1H).

Step 6. Compound **273.6** was prepared as previously described in Example 1. ¹H NMR (300 MHz, CDCl₃) δ 7.58 (d, *J* = 7.2 Hz, 0.5H), 7.43 (d, *J* = 6.9 Hz, 0.5H), 7.35-7.30 (m, 2H), 7.09-7.02 (m, 1H), 6.24-5.86 (m, 1H), 4.98-4.65 (m, 3H), 3.81-3.74 (m, 1H), 2.34 (d, *J* = 6.6 Hz, 0.5H), 2.12 (d, *J* = 9.9 Hz, 0.5H).

Step 7. Compound **273.7** was prepared as previously described in Example 1. ¹H NMR (300 MHz, CDCl₃) δ 7.51-7.48 (m, 0.5H), 7.43-7.32 (m, 2.5H), 7.15-7.06 (m, 1H), 6.22-5.82 (m, 1H), 5.04 (d, *J* = 15.3 Hz, 0.5H), 4.91-4.86 (m, 1.5H), 4.62 (d, *J* = 7.8 Hz, 0.5H), 4.34 (s, 0.5H), 3.96-3.88 (m, 1H).

Step 8. Compounds **273.8a** and **273.8b** were prepared as previously described in Example 1.

tert-butyl-(*cis*-3-(difluoromethyl)isochroman-4-yl)carbamate **273.8a** TLC higher Rf. ^1H NMR (300 MHz, CDCl_3) δ 7.44-7.41 (m, 1H), 7.34-7.22 (m, 2H), 7.05-7.00 (m, 1H), 6.11-5.73 (m, 1H), 4.99-4.95 (m, 2H), 4.88-4.81 (m, 2H), 3.91-3.85 (m, 1H), 1.44 (s, 9H).


tert-butyl-(*trans*-3-(difluoromethyl)isochroman-4-yl)carbamate **273.8b** TLC lower Rf. ^1H NMR (300 MHz, CDCl_3) δ 7.42-7.40 (m, 1H), 7.32-7.26 (m, 2H), 7.04-7.01 (m, 1H), 6.16-5.78 (m, 1H), 5.02-4.95 (m, 1H), 4.90-4.89 (m, 2H), 4.85-4.80 (m, 1H), 3.84-3.80 (m, 1H), 1.44 (s, 9H).

Step 7. The compounds of **Example 273** and **Example 274** were prepared as previously described in Example 1.

[0158] **Example 273.** MS (ESI): m/z 200 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.51-7.45 (m, 2H), 7.42-7.38 (m, 1H), 7.25 (d, J = 7.5 Hz, 1H), 6.19 (td, J = 54.0, 3.9 Hz, 1H), 5.15 (d, J = 15.6 Hz, 1H), 5.02 (d, J = 15.6 Hz, 1H), 4.67 (s, 1H), 4.34-4.24 (m, 1H).


[0159] **Example 274.** MS (ESI): m/z 200 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.54-7.40 (m, 3H), 7.25 (d, J = 7.5 Hz, 1H), 6.13 (td, J = 54.3, 5.1 Hz, 1H), 5.00 (s, 2H), 4.64 (d, J = 2.4 Hz, 1H), 4.37-4.27 (m, 1H).

[0160] **Example 275. *cis*-3-(fluoromethyl)isochroman-4-amine hydrochloride**




[0161] The compound of **Example 275** was prepared as previously described in Example 273 using ethyl 2-fluoroacetate. MS (ESI): m/z 182 [M+H]. ^1H NMR (300 MHz, DMSO-d_6) δ 8.41 (s, 3H), 7.52-7.50 (d, J = 7.5 Hz, 1H), 7.44-7.31 (m, 2H), 7.21-7.19 (d, J = 7.2 Hz, 1H), 5.0-4.85 (m, 2.5H), 4.83-4.68 (m, 1H), 4.62-4.56 (m, 0.5H), 4.45 (s, 1H), 4.21-4.15 (d, J = 17.1 Hz, 1H).

[0162] **Example 276. *cis*-3-(fluoromethyl)isochroman-4-amine hydrochloride**



[0163] The compound of **Example 276** was prepared as previously described in Example 274 using ethyl 2-fluoroacetate. MS (ESI): *m/z* 182 [M+H]. ¹H NMR (300 MHz, DMSO-d₆) δ 8.8 (s, 3H), 7.68-7.66 (m, 1H), 7.38-7.34 (m, 2H), 7.21-7.20 (m, 1H), 4.91-4.61 (m, 4H), 4.44 (s, 1H), 4.29-4.17 (m, 1H).

[0164] **Example 277. 4-Aminoisochroman-6-yl ethyl(methyl)carbamate hydrochloride**



Example 277

Step 1. 6-Methoxyisochroman-4-one **10.3** (1 g, 5.61 mmol) was dissolved in anhydrous DMF (3 mL) under a N₂ atmosphere and treated with sodium ethanethiolate (841 mg, 8.41 mmol). The reaction mixture was heated to 140 °C and stirred at that temperature for 15 min. After cooling to room temperature, the mixture was partitioned between H₂O (15 mL) and EtOAc (10 mL). The aqueous phase was extracted with EtOAc (2x10 mL). The aqueous phase was acidified with aqueous 1N HCl until pH=2 and extracted with EtOAc (2x10 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to afford 6-hydroxyisochroman-4-one **277.1** (696 mg,

4.24 mmol) as brown oil which was used without further purification. ^1H NMR (300 MHz, CDCl_3) δ 7.45 (d, $I = 2.1$ Hz, 1H), 7.15-7.07 (m, 2H), 4.84 (s, 2H), 4.35 (s, 2H).

Step 2. A solution of Compound **277.1** (696.5 mg, 4.23 mmol) in CH_3CN (15 mL) was treated with ethyl(methyl)carbamic chloride (1.02 g, 8.46 mmol) and K_2CO_3 (876 mg, 6.34 mmol). The reaction mixture was heated to 55 °C and stirred at that temperature for 5 h. After concentration under reduced pressure, the residue was diluted with water (10 mL) and extracted with EtOAc (3x10 mL). The combined organics were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 , gradient elution from 10% $\text{EtOAc}/\text{hexanes}$ to 75% $\text{EtOAc}/\text{hexanes}$) provided 4-oxoisochroman-6-yl ethyl(methyl)carbamate **277.2** (230 mg, 0.922 mmol) as a yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 7.76 (s, 1H), 7.37 (d, $J = 8.1$ Hz, 1H), 7.23 (d, $J = 8.4$ Hz, 1H), 4.89 (s, 2H), 4.36 (s, 2H), 3.50-3.40 (m, 2H), 3.04 (d, $J = 24.3$ Hz, 3H), 1.28-1.11 (m, 3H).

Step 3. Compound **277.2** (230mg, 922 μmol) was dissolved in MeOH (3 mL) and treated with hydroxylamine hydrochloride (319 mg, 4.60 mmol) and pyridine (2 mL). The reaction mixture was heated to 65 °C and stirred at that temperature for 30 min. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure. The residue was diluted with H_2O (10 mL). The pH was adjusted to 5~6 with 1 N HCl aqueous solution, and the mixture was partitioned with EtOAc (15 mL). The aqueous phase was extracted with EtOAc (2x15 mL). The combined organics were dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo to provide crude 4-(hydroxyimino)isochroman-6-yl ethyl(methyl)carbamate **277.3** (228 mg, 862 μmol) as a colorless oil which was used without further purification. ^1H NMR (300 MHz, CDCl_3) δ 7.67 (s, 1H), 7.10 (s, 2H), 4.78 (s, 2H), 4.67 (s, 2H), 3.51-3.38 (m, 2H), 3.03 (d, $J = 22.5$ Hz, 3H), 1.29-1.22 (m, 3H).

Step 4. A flask containing solution of Compound **277.3** (228 mg, 862 μmol) in MeOH (5 mL) was flushed with H_2 . Raney Ni (0.5 mL) was added and the reaction mixture was stirred at ambient temperature for 3 h under a H_2 balloon. $(\text{Boc})_2\text{O}$ (375 mg, 1.72 mmol) was added, and the mixture was stirred at ambient temperature for 5 h. The nickel catalyst was removed by filtration through a celite pad, and the filtrate was concentrated under reduced pressure. Purification by flash column chromatography (SiO_2 , gradient elution from 5% $\text{EtOAc}/\text{hexanes}$ to 20% EtOAc) afforded 4-((tert-butoxycarbonyl)amino)isochroman-6-yl ethyl(methyl)carbamate **277.4** (132 mg, 376 μmol) as a colorless oil. ^1H NMR (300 MHz,

CDCl_3) δ 7.16 (s, 1H), 7.04-6.95 (m, 2H), 5.08 (d, J = 8.7 Hz, 1H), 4.81-4.65 (m, 3H), 4.02 (dd, J = 14.7, 2.7 Hz, 1H), 3.85 (dd, J = 14.7, 3.0 Hz, 1H), 3.45-3.39 (m, 2H), 3.02 (d, J = 21.9 Hz, 3H), 1.46 (s, 9H), 1.26-1.14 (m, 3H).

Step 5. The compound of **Example 277** was prepared as previously described in Example 1. MS (ESI): m/z 251 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.24-7.12 (m, 3H), 4.98 (d, J = 15.3 Hz, 1H), 4.75 (d, J = 15.3 Hz, 1H), 4.33 (s, 1H), 4.19 (d, J = 12.9 Hz, 1H), 3.94 (dd, J = 12.9, 2.1 Hz, 1H), 3.53-3.35 (m, 2H), 3.05 (d, J = 31.8 Hz, 3H), 1.19 (dt, J = 24.0, 7.2 Hz, 3H).

[0181] **Table 11.** Compounds were prepared as described in **Example 277** using the appropriately substituted starting materials.

| Example | Structure | Characterization Data |
|---------|-----------|--|
| 278 | | MS (ESI): m/z 178 [M+H]. ^1H NMR (300 MHz, CDCl_3) δ 8.67 (brs, 3H), 7.43 (s, 1H), 7.12 (d, J = 8.1 Hz, 1H), 6.95 (d, J = 7.8 Hz, 1H), 5.04 - 4.98 (m, 1H), 4.30 (s, 1H), 4.10 (s, 2H), 2.30 (s, 3H), 1.45 (d, J = 6.6 Hz, 3H). |
| 279 | | MS (ESI): m/z 265 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.30 (brs, 3H), 7.49 (d, J = 8.4 Hz, 1H), 7.08 (dd, J = 8.4, 2.4 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 4.85 (d, J = 15.9 Hz, 1H), 4.75 (d, J = 15.9 Hz, 1H), 4.25 (s, 1H), 3.99-3.92 (m, 1H), 3.44-3.26 (m, 2H), 2.95 (d, J = 36.0 Hz, 3H), 1.32 (d, J = 6.6 Hz, 3H), 1.12 (dt, J = 24.5, 7.2 Hz, 3H). |
| 280 | | MS (ESI): m/z 265 [M+H]. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.57 (brs, 3H), 7.59 (d, J = 8.4 Hz, 1H), 7.10 (dd, J = 8.4, 2.1 Hz, 1H), 6.97 (d, J = 2.1 Hz, 1H), 4.73 (s, 2H), 4.18-4.13 (m, 2H), 3.44-3.26 (m, 2H), 2.95 (d, J = 36.0 Hz, 3H), 1.27 (d, J = 6.3 Hz, 3H), 1.13 (dt, J = 24.4, 6.9 Hz, 3H). |
| 281 | | MS (ESI): m/z 265 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.26-7.16 (m, 3H), 4.99 (d, J = 15.6 Hz, 1H), 4.85 (d, J = 15.6 Hz, 1H), 4.28 (s, 1H), 4.05 (q, J = 6.6 Hz, 1H), 3.57-3.31 (m, 2H), 3.05 (d, J = 34.1 Hz, 3H), 1.41 (d, J = 6.6 Hz, 3H), 1.24 (dt, J = 23.7, 7.2 Hz, 3H). |
| 282 | | MS (ESI): m/z 265 [M+H]. ^1H NMR (300 MHz, CD_3OD) δ 7.09-7.00 (m, 3H), 4.69 (s, 2H), 4.17-4.04 (m, 2H), 3.40-3.32 (m, 2H), 2.89 (d, J = 34.8 Hz, 3H), 1.17 (d, J = 6.6 Hz, 3H), 1.09 (dt, J = 24.6, |



| Example | Structure | Characterization Data |
|---------|-----------|-----------------------|
| | | 7.2 Hz, 3H). |







[0165] General Procedure for Chiral Separation of Racemic Compounds.








[0166] The chiral separations were carried out on the racemic amines or N-Boc protected amine intermediates using SFC (supercritical CO_2 fluid chromatography) on a preparative Thar SFC-80 system using the specified column and co-solvent system. CO_2 total flows were between 60 to 80 g/min with 100 bar back pressure. Additional examples were separated by normal phase preparatory HPLC using the specified column and solvent system.








[0167] The Faster Moving Enantiomer (FME) was the earlier eluting enantiomer and the Slower Moving Enantiomer (SME) was the later eluting enantiomer. Following isolation of the separated enantiomers, the Boc protecting groups were removed (if needed) and HCl salts were formed as previously described in Example 1. The compounds in Table 12 are all single enantiomers prepared by chiral separation of previously described racemates. For example, Examples 111 and 112 are the individual enantiomers of Example 12. Each example is a single enantiomer, but the absolute stereochemistry has not been conclusively proven.








[0168] Table 12. Compounds Prepared by Chiral Separation








| Example | Separation Conditions | Analytical Data |
|--|--|--|
|  111 | SFC AD-H 20×250mm, 5 μm 65:35 CO_2 /MeOH (0.1% NH_4OH) FME | MS(ESI): m/z 150 [M+H]. ^1H NMR (400 MHz, CD_3OD) δ 7.36-7.49 (m, 3H), 7.20 (d, J = 7.6 Hz, 1H), 4.95 (d, J = 15.2 Hz, 1H), 4.82 (d, J = 15.6 Hz, 1H), 4.38 (s, 1H), 4.26 (dd, J = 12.8, 1.6 Hz, 1H), 3.99 (dd, J = 12.8, 2.0 Hz, 1H) |
|  112 | SFC AD-H 20×250mm, 5 μm 65:35 CO_2 /MeOH (0.1% NH_4OH) SME | MS(ESI): m/z 150 [M+H]. ^1H NMR (400 MHz, CD_3OD) δ 7.36-7.49 (m, 3H), 7.20 (d, J = 7.6 Hz, 1H), 4.95 (d, J = 15.2 Hz, 1H), 4.82 (d, J = 15.6 Hz, 1H), 4.38 (s, 1H), 4.26 (dd, J = 12.8, 1.6 Hz, 1H), 3.99 (dd, J = 12.8, 2.0 Hz, 1H) |

| Example | Separation Conditions | Analytical Data |
|---|---|---|
|  <p>113</p> | <p>SFC AD 20×250mm, 5µm 75:25 CO₂/MeOH (0.1% NH₄OH) FME</p> | <p>MS(ESI): <i>m/z</i> 168 [M+H]. ¹H NMR (500 MHz, CD₃OD) δ 7.28-7.18 (m, 3H), 4.94 (d, <i>J</i> = 15.0 Hz, 1H), 4.78 (d, <i>J</i> = 15.5 Hz, 1H), 4.40 (s, 1H), 4.25 (dd, <i>J</i> = 12.5, 1.5 Hz, 1H), 3.98 (dd, <i>J</i> = 12.5, 2.5 Hz, 1H).</p> |
|  <p>114</p> | <p>SFC AD 20×250mm, 5µm 75:25 CO₂/MeOH (0.1% NH₄OH) SME</p> | <p>MS(ESI): <i>m/z</i> 168 [M+H]. ¹H NMR (500 MHz, CD₃OD) δ 7.28-7.18 (m, 3H), 4.94 (d, <i>J</i> = 15.0 Hz, 1H), 4.78 (d, <i>J</i> = 15.5 Hz, 1H), 4.40 (s, 1H), 4.25 (dd, <i>J</i> = 12.5, 1.5 Hz, 1H), 3.98 (dd, <i>J</i> = 12.5, 2.5 Hz, 1H).</p> |
|  <p>115</p> | <p>SFC AD 20×250mm, 5µm 80:20 CO₂/MeOH (0.5% NH₃) FME</p> | <p>MS (ESI): <i>m/z</i> 177 [M+H]. ¹H NMR (500 MHz, CD₃OD) δ 7.17 (d, <i>J</i> = 11.6 Hz, 3H), 4.82 (d, <i>J</i> = 14.6 Hz, 1H), 4.78 (d, <i>J</i> = 8.6 Hz, 1H), 4.74 (d, <i>J</i> = 14.6 Hz, 1H), 4.21-4.14 (m, 1H), 4.07-4.02 (m, 1H), 2.40 (s, 3H), 2.19-2.28 (m, 2H).</p> |
|  <p>116</p> | <p>SFC AD 20×250mm, 5µm 80:20 CO₂/MeOH (0.5% NH₃) SME</p> | <p>MS (ESI): <i>m/z</i> 177 [M+H]. ¹H NMR (500 MHz, CD₃OD) δ 7.17 (d, <i>J</i> = 11.4 Hz, 3H), 4.82 (d, <i>J</i> = 14.6 Hz, 1H), 4.78 (d, <i>J</i> = 8.4 Hz, 1H), 4.74 (d, <i>J</i> = 14.6 Hz, 1H), 4.22-4.14 (m, 1H), 4.07-4.02 (m, 1H), 2.40 (s, 3H), 2.30-2.15 (m, 2H).</p> |
|  <p>117</p> | <p>SFC AD-H 20×250mm, 5µm 60:30 CO₂/MeOH (0.5% NH₄OH) FME</p> | <p>MS (ESI): <i>m/z</i> 183 [M-16]. ¹H-NMR of freebase (500 MHz, CDCl₃) δ 8.17 (d, <i>J</i> = 8.5 Hz, 1H), 7.87 (d, <i>J</i> = 8.0 Hz, 1H), 7.76 (d, <i>J</i> = 8.5 Hz, 1H), 7.63-7.59 (m, 1H), 7.53-7.50 (m, 1H), 7.12 (d, <i>J</i> = 8.5 Hz, 1H), 4.99-4.88 (m, 2H), 4.38 (s, 1H), 4.27 (d, <i>J</i> = 11.5 Hz, 1H), 4.00-3.97 (dd, <i>J</i> = 11.0, 2.5 Hz, 1H), 1.90 (s, 2H).</p> |
|  <p>118</p> | <p>SFC AD-H 20×250mm, 5µm 60:40 CO₂/MeOH (0.5% NH₄OH) SME</p> | <p>MS (ESI): <i>m/z</i> 183 [M-16]. ¹H-NMR of freebase (500 MHz, CDCl₃) δ 8.16 (d, <i>J</i> = 9.0 Hz, 1H), 7.87 (d, <i>J</i> = 8.0 Hz, 1H), 7.76 (d, <i>J</i> = 8.5 Hz, 1H), 7.63-7.59 (m, 1H), 7.53-7.50 (m, 1H), 7.12 (d, <i>J</i> = 8.5 Hz, 1H), 4.99-4.88 (m, 2H), 4.39 (s, 1H), 4.28 (d, <i>J</i> = 11.5 Hz, 1H), 4.00-3.97 (dd, <i>J</i> = 11.0, 2.5 Hz, 1H), 1.92 (s, 2H).</p> |







| Example | Separation Conditions | Analytical Data |
|---|--|---|
|  | SFC AD 20×250mm, 5µm 60:30 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 218 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.64 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (d, <i>J</i> = 8.4 Hz, 1H), 4.96 (d, <i>J</i> = 15.9 Hz, 1H), 4.78 (d, <i>J</i> = 15.9 Hz, 1H), 4.67 (s, 1H), 4.32 (d, <i>J</i> = 13.0 Hz, 1H), 3.92 (dd, <i>J</i> = 13.0, 1.8 Hz, 1H). |
|  | SFC AD 20×250mm, 5µm 60:40 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 218 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.64 (d, <i>J</i> = 8.4 Hz, 1H), 7.21 (d, <i>J</i> = 8.4 Hz, 1H), 4.96 (d, <i>J</i> = 15.9 Hz, 1H), 4.78 (d, <i>J</i> = 15.9 Hz, 1H), 4.66 (s, 1H), 4.32 (d, <i>J</i> = 13.0 Hz, 1H), 3.92 (dd, <i>J</i> = 13.0, 1.9 Hz, 1H). |
|  | SFC AD 20×250mm, 5µm 75:25 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 180 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.11 (d, <i>J</i> = 8.5 Hz, 1H), 7.05 (d, <i>J</i> = 2.5 Hz, 1H), 7.00 (dd, <i>J</i> = 8.5, 2.6 Hz, 1H), 4.85 (d, <i>J</i> = 14.9 Hz, 1H), 4.73 (d, <i>J</i> = 14.9 Hz, 1H), 4.34 (s, 1H), 4.23 (dd, <i>J</i> = 12.8, 1.4 Hz, 1H), 3.95 (dd, <i>J</i> = 12.8, 2.3 Hz, 1H), 3.83 (s, 3H). |
|  | SFC AD 20×250mm, 5µm 75:25 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 180 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.11 (d, <i>J</i> = 8.4 Hz, 1H), 7.05 – 6.96 (m, 2H), 4.85 (d, <i>J</i> = 14.9 Hz, 1H), 4.74 (d, <i>J</i> = 14.9 Hz, 1H), 4.32 (s, 1H), 4.22 (dd, <i>J</i> = 12.8, 2.0 Hz, 1H), 3.95 (dd, <i>J</i> = 12.8, 2.3 Hz, 1H), 3.83 (s, 3H). |
|  | SFC AY 20×250mm, 5µm 90:10 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.28-7.24 (m, 2H), 7.07 (d, <i>J</i> = 8.0 Hz, 1H), 4.91-4.88 (m, 1H), 4.76 (d, <i>J</i> = 15.2 Hz, 1 H), 4.32 (s, 1H), 4.23 (d, <i>J</i> = 12.8 Hz, 1H), 3.95 (dd, <i>J</i> = 12.8, 2.4Hz, 1H), 2.37 (s, 3H). |
|  | SFC AY 20×250mm, 5µm 90:10 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.28-7.23 (m, 2H), 7.07 (d, <i>J</i> = 8.0 Hz, 1H), 4.92-4.88 (m, 1H), 4.75 (d, <i>J</i> = 15.2 Hz, 1 H), 4.32 (s, 1H), 4.23 (dd, <i>J</i> = 12.8, 1.2Hz, 1H), 3.95 (dd, <i>J</i> = 12.8, 2.4 Hz, 1H), 2.37 (s, 3H). |
|  | HPLC AS-H 20×250, 10 □m 95:5 Hexanes/EtOH (0.1%DEA) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.10 (d, <i>J</i> = 7.2 Hz, 2H), 4.89 (s, 1H), 4.68 (d, <i>J</i> = 15.5 Hz, 1H), 4.30 (s, 1H), 4.22 (d, <i>J</i> = 12.7 Hz, 1H), 3.92 (d, <i>J</i> = 12.6 Hz, 1H), 2.35 (s, 3H), 2.18 (s, 3H). |

| Example | Separation Conditions | Analytical Data |
|---|--|---|
|  126 | HPLC AS-H 20×250, 10 μ m 95:5 Hexanes/EtOH (0.1%DEA) SME | MS (ESI): <i>m/z</i> 178 [M+H]. 1 H NMR (500 MHz, CD ₃ OD) δ 7.11 (d, <i>J</i> = 15.7 Hz, 2H), 4.89 (s, 1H), 4.67 (d, <i>J</i> = 15.5 Hz, 1H), 4.31 (s, 1H), 4.23 (d, <i>J</i> = 12.7 Hz, 1H), 3.92 (d, <i>J</i> = 12.6 Hz, 1H), 2.35 (s, 3H), 2.18 (s, 3H). |
|  127 | SFC AD 20×250mm, 5 μ m 70:30 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 184.0 [M+H]. 1 H NMR (400 MHz, CD ₃ OD) δ 7.47-7.42 (m, 2H), 7.19 (dd, <i>J</i> = 6.0, 3.2 Hz, 1H), 5.00-4.78 (m, 2H), 4.63 (s, 1H), 4.33 (dd, <i>J</i> = 12.8, 0.8 Hz, 1H), 3.93 (dd, <i>J</i> = 13.2, 2.0 Hz, 1H). |
|  128 | SFC AD 20×250mm, 5 μ m 70:30 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 184.0 [M+H]. 1 H NMR (400 MHz, CD ₃ OD) δ 7.46-7.42 (m, 2H), 7.18-7.20 (m, 1H), 4.99-4.78 (m, 2H), 4.62 (s, 1H), 4.34-4.31 (m, 1H), 3.93 (dd, <i>J</i> = 12.8, 1.6 Hz, 1H). |
|  129 | SFC AY 20×250mm, 5 μ m 50:50 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 164.1 [M+H]. 1 H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.73 (br s, 3H), 7.42 (d, <i>J</i> = 6.8 Hz, 1H), 7.24-7.18 (m, 2H), 4.79-4.59 (m, 2H), 4.28 (s, 1H), 4.22 (dd, <i>J</i> = 12.4, 2.0 Hz, 1H), 3.84 (dd, <i>J</i> = 12.4, 2.0 Hz, 1H), 2.13 (s, 3H). |
|  130 | SFC AY 20×250mm, 5 μ m 50:50 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 164.1 [M+H]. 1 H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.69 (br s, 3H), 7.42 (d, <i>J</i> = 6.8 Hz, 1H), 7.24-7.18 (m, 2H), 4.79-4.59 (m, 2H), 4.28 (s, 1H), 4.20 (dd, <i>J</i> = 12.4, 1.6 Hz, 1H), 3.83 (dd, <i>J</i> = 12.4, 2.4 Hz, 1H), 2.13 (s, 3H). |
|  131 | SFC OJ 20×250mm, 5 μ m 70:30 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 184 [M+H]. 1 H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.88 (br s, 3H), 7.64 (d, <i>J</i> = 7.6 Hz, 1H), 7.50-7.48 (m, 1H), 7.38 (t, <i>J</i> = 8.0 Hz, 1H), 4.82-4.61 (m, 2H), 4.36 (s, 1H), 4.28-4.25 (m, 1H), 3.88 (dd, <i>J</i> = 12.4, 2.4 Hz, 1H). |
|  132 | SFC OJ 20×250mm, 5 μ m 70:30 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 184 [M+H]. 1 H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.87 (br s, 3H), 7.64 (d, <i>J</i> = 7.2 Hz, 1H), 7.48 (dd, <i>J</i> = 8.4, 0.8 Hz, 1H), 7.38 (t, <i>J</i> = 8.0 Hz, 1H), 4.82-4.61 (m, 2H), 4.37 (s, 1H), 4.28-4.25 (m, 1H), 3.88 (dd, <i>J</i> = 12.8, 2.4 Hz, 1H). |








| Example | Separation Conditions | Analytical Data |
|---|--|---|
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.56 (br s, 3H), 7.25 (d, <i>J</i> = 7.6 Hz, 1H), 7.12 (d, <i>J</i> = 7.2 Hz, 1H), 6.97 (d, <i>J</i> = 8.0 Hz, 1H), 4.87-4.68 (m, 2H), 4.32 (s, 1H), 4.30 (d, <i>J</i> = 12.0 Hz, 1H), 3.75 (d, <i>J</i> = 11.2 Hz, 1H), 2.42 (s, 3H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, DMSO- <i>d</i> 6) δ 8.54 (br s, 3H), 7.26 (t, <i>J</i> = 7.6 Hz, 1H), 7.12 (d, <i>J</i> = 7.2 Hz, 1H), 6.97 (d, <i>J</i> = 7.6 Hz, 1H), 4.87-4.68 (m, 2H), 4.39 (s, 1H), 4.30 (d, <i>J</i> = 12.0 Hz, 1H), 3.78-3.75 (m, 1H), 2.42 (s, 3H). |
|  | SFC AD 20×250mm, 5µm 55:45 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.22 (d, <i>J</i> = 8.0 Hz, 1H), 6.92 (d, <i>J</i> = 8.0 Hz, 1H), 4.93 (d, <i>J</i> = 15.0 Hz, 1H), 4.79 (d, <i>J</i> = 15.0 Hz, 1H), 4.58 (s, 1H), 4.30 (dd, <i>J</i> = 12.5, 1.0 Hz, 1H), 3.90 (d, <i>J</i> = 12.5 Hz, 1H), 2.34 (s, 3H), 2.33 (s, 3H). |
|  | SFC AD 20×250mm, 5µm 55:45 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.23 (d, <i>J</i> = 7.5 Hz, 1H), 6.93 (d, <i>J</i> = 7.5 Hz, 1H), 4.93 (d, <i>J</i> = 15.0 Hz, 1H), 4.79 (d, <i>J</i> = 15.0 Hz, 1H), 4.61 (s, 1H), 4.30 (dd, <i>J</i> = 12.5, 1.0Hz, 1H), 3.91 (d, <i>J</i> = 12.5 Hz, 1H), 2.34 (s, 3H), 2.33 (s, 3H). |
|  | SFC AY 4.6×250mm, 5µm 85:15 CO ₂ /EtOH (0.1% NH ₃) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.35 (d, <i>J</i> = 7.6 Hz, 1H), 7.25 (d, <i>J</i> = 7.5 Hz, 1H), 7.05 (d, <i>J</i> = 7.7 Hz, 1H), 5.00 (d, <i>J</i> = 15.6 Hz, 1H), 4.84 (d, <i>J</i> = 15.6 Hz, 1H), 4.55-4.41 (m, 2H), 3.90 (dd, <i>J</i> = 13.1, 1.3 Hz, 1H), 2.85 (s, 3H), 2.47 (s, 3H). |
|  | SFC AY 4.6×250mm, 5µm 85:15 CO ₂ /EtOH (0.1% NH ₃) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.36 (t, <i>J</i> = 7.6 Hz, 1H), 7.25 (d, <i>J</i> = 7.5 Hz, 1H), 7.04 (d, <i>J</i> = 7.7 Hz, 1H), 4.99 (d, <i>J</i> = 15.6 Hz, 1H), 4.84 (d, <i>J</i> = 15.7 Hz, 1H), 4.53-4.42 (m, 2H), 3.90 (dd, <i>J</i> = 13.1, 1.2 Hz, 1H), 2.85 (s, 3H), 2.47 (s, 3H). |
|  | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) FME | MS (ESI) <i>m/z</i> 194 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.14 (d, <i>J</i> = 8.5 Hz, 1H), 7.10-7.01 (m, 2H), 4.91 (d, <i>J</i> = 12.0 Hz, 4H), 4.76 (d, <i>J</i> = 15.0 Hz, 1H), 4.43 (dd, <i>J</i> = 13.3, 0.8 Hz, 1H), 4.25 (s, 1H), 3.91 (dd, <i>J</i> = 13.3, 1.9 Hz, 1H), 3.85 (s, 3H), 2.77 (s, 3H). |









| Example | Separation Conditions | Analytical Data |
|---|--|--|
|  140 | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) SME | MS (ESI): <i>m/z</i> 194 [M+H]. ^1H NMR (400 MHz, CD_3OD) δ 7.14 (d, <i>J</i> = 8.5 Hz, 1H), 7.10-7.02 (m, 2H), 4.90 (d, <i>J</i> = 12.0 Hz, 1H), 4.76 (d, <i>J</i> = 15.0 Hz, 1H), 4.43 (dd, <i>J</i> = 13.2, 1.0 Hz, 1H), 4.25 (s, 1H), 3.91 (dd, <i>J</i> = 13.2, 1.9 Hz, 1H), 3.85 (s, 3H), 2.77 (s, 3H). |
|  141 | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (400 MHz, CD_3OD) δ 7.32 (s, 1H), 7.28 (d, <i>J</i> = 8.0 Hz, 1H), 7.10 (d, <i>J</i> = 8.0 Hz, 1H), 4.93 (d, <i>J</i> = 15.2 Hz, 1H), 4.78 (d, <i>J</i> = 15.2 Hz, 1H), 4.43 (d, <i>J</i> = 12.8 Hz, 1H), 4.23 (s, 1H), 3.92 (dd, <i>J</i> = 13.2, 2.0 Hz, 1H), 2.76 (s, 3H), 2.38 (s, 3H). |
|  142 | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ^1H NMR (400 MHz, CD_3OD) δ 7.32 (s, 1H), 7.28 (d, <i>J</i> = 8.0 Hz, 1H), 7.10 (d, <i>J</i> = 8.0 Hz, 1H), 4.93 (d, <i>J</i> = 15.2 Hz, 1H), 4.78 (d, <i>J</i> = 15.2 Hz, 1H), 4.43 (d, <i>J</i> = 12.8 Hz, 1H), 4.23 (s, 1H), 3.92 (dd, <i>J</i> = 13.2, 2.0 Hz, 1H), 2.76 (s, 3H), 2.38 (s, 3H). |
|  143 | SFC AY-H 4.6×250mm, 5µm 80:20 CO_2 /EtOH (1% NH_3) FME | MS (ESI): <i>m/z</i> 192 [M + H]. ^1H NMR (400 MHz, CD_3OD) δ 7.25 (d, <i>J</i> = 7.8 Hz, 1H), 6.93 (d, <i>J</i> = 7.8 Hz, 1H), 4.94 (d, <i>J</i> = 15.4 Hz, 1H), 4.80 (d, <i>J</i> = 15.4 Hz, 1H), 4.53~4.43 (m, 2H), 3.91~3.82 (m, 1H), 2.83 (s, 3H), 2.33 (s, 6H). |
|  144 | SFC AY-H 4.6×250mm, 5µm 80:20 CO_2 /EtOH (1% NH_3) SME | MS (ESI): <i>m/z</i> 192 [M + H]. ^1H NMR (400 MHz, CD_3OD) δ 7.27 (d, <i>J</i> = 8.0 Hz, 1H), 6.94 (d, <i>J</i> = 8.0 Hz, 1H), 4.94 (d, <i>J</i> = 15.4 Hz, 1H), 4.80 (d, <i>J</i> = 15.3 Hz, 1H), 4.48 (dd, <i>J</i> = 15.2, 1.9 Hz, 2H), 3.86 (dd, <i>J</i> = 12.9, 0.9 Hz, 1H), 2.82 (s, 3H), 2.33 (s, 6H). |
|  145 | HPLC AY-H 4.6×250mm, 5µm 80:20 Hexanes/EtOH (0.1% DEA) FME | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (500 MHz, CD_3OD) δ 7.36 (t, <i>J</i> = 7.6 Hz, 1H), 7.25 (d, <i>J</i> = 7.5 Hz, 1H), 7.04 (d, <i>J</i> = 7.7 Hz, 1H), 4.99 (d, <i>J</i> = 15.6 Hz, 1H), 4.84 (d, <i>J</i> = 15.7 Hz, 1H), 4.53 – 4.42 (m, 2H), 3.90 (dd, <i>J</i> = 13.1, 1.2 Hz, 1H), 2.85 (s, 3H), 2.47 (s, 3H). |
|  146 | HPLC AY-H 4.6×250mm, 5µm 80:20 Hexanes/EtOH (0.1% DEA) SME | MS (ESI): <i>m/z</i> 198 [M+H]. ^1H NMR (500 MHz, CD_3OD) δ 7.53 – 7.47 (m, 2H), 7.22 (dd, <i>J</i> = 8.6, 6.6 Hz, 1H), 5.01 (d, <i>J</i> = 15.9 Hz, 1H), 4.83 (d, <i>J</i> = 15.9 Hz, 1H), 4.59 – 4.46 (m, 2H), 3.92 (dd, <i>J</i> = 13.4, 1.6 Hz, 1H), 2.88 (s, 3H). |









| Example | Separation Conditions | Analytical Data |
|---------|---|---|
| | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) FME | MS (ESI): <i>m/z</i> 232 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.67 (d, <i>J</i> = 8.8 Hz, 1H), 7.23 (d, <i>J</i> = 8.4 Hz, 1H), 4.99 (d, <i>J</i> = 16.0 Hz, 1H), 4.79 (d, <i>J</i> = 16.4 Hz, 1H), 4.59 (s, 1H), 4.52 (d, <i>J</i> = 13.6 Hz, 1H), 3.89 (dd, <i>J</i> = 12.8, 1.2 Hz, 1H), 2.89 (s, 3H). |
| | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% DEA) SME | MS (ESI): <i>m/z</i> 232 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.67 (d, <i>J</i> = 8.8 Hz, 1H), 7.21 (d, <i>J</i> = 8.4 Hz, 1H), 4.99 (d, <i>J</i> = 16.0 Hz, 1H), 4.79 (d, <i>J</i> = 16.4 Hz, 1H), 4.59 (s, 1H), 4.52 (d, <i>J</i> = 13.6 Hz, 1H), 3.89 (dd, <i>J</i> = 12.8, 1.2 Hz, 1H), 2.89 (s, 3H). |
| | SFC AY 20×250mm, 5µm 80:20 CO ₂ /EtOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.52 (dd, <i>J</i> = 13.3, 8.0 Hz, 2H), 7.42 (t, <i>J</i> = 7.8 Hz, 1H), 5.03 (d, <i>J</i> = 16.3 Hz, 1H), 4.74 (d, <i>J</i> = 16.3 Hz, 1H), 4.46 (d, <i>J</i> = 13.2 Hz, 1H), 4.33 (s, 1H), 3.93 (dd, <i>J</i> = 13.2, 1.5 Hz, 1H), 2.79 (s, 3H). |
| | SFC AY-H 20×250mm, 5µm 80:20 CO ₂ /EtOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.41 (dd, <i>J</i> = 16.0, 6.6 Hz, 2H), 7.31 (t, <i>J</i> = 7.8 Hz, 1H), 4.92 (d, <i>J</i> = 16.3 Hz, 1H), 4.63 (d, <i>J</i> = 16.3 Hz, 1H), 4.35 (d, <i>J</i> = 13.3 Hz, 1H), 4.22 (s, 1H), 3.82 (dd, <i>J</i> = 13.2, 1.5 Hz, 1H), 2.68 (s, 3H). |
| | SFC OD 4.6×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.47 (m, 1H), 7.43 (ddd, <i>J</i> =7.2, 7.2, 0.8 Hz, 1H), 7.37 (dd, <i>J</i> =8.0, 8.0 Hz, 1H), 7.20 (d, <i>J</i> = 7.2 Hz, 1H), 4.86 (s, 2H), 4.30 (m, 1H), 4.21 (s, 1H), 1.33 (d, <i>J</i> = 6.8 Hz, 3H). |
| | SFC OD 4.6×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.48 (m, 1H), 7.42 (ddd, <i>J</i> =7.6, 7.6, 1.2 Hz, 1H), 7.37 (dd, <i>J</i> =7.6, 7.6, Hz, 1H), 7.20 (d, <i>J</i> = 7.2 Hz, 1H), 4.86 (s, 2H), 4.31 (m, 1H), 4.22 (s, 1H), 1.33 (d, <i>J</i> = 6.8 Hz, 3H). |
| | SFC IC 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.5% NH ₃) FME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.45-7.34 (m, 3H), 7.20 (d, <i>J</i> = 7.2 Hz, 1H), 4.98 (d, <i>J</i> = 15.6 Hz, 1H), 4.90-4.86 (m, 1H), 4.28 (s, 1H), 4.08-4.03 (m, 1H), 1.42 (d, <i>J</i> = 6.4 Hz, 3H). |

| Example | Separation Conditions | Analytical Data |
|---|---|--|
|  154 | SFC IC 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.5% NH ₃) SME | MS (ESI): <i>m/z</i> 164 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.47-7.34 (m, 3H), 7.20 (d, <i>J</i> =7.6 Hz, 1H), 4.98 (d, <i>J</i> =15.6 Hz, 1H), 4.90-4.86 (m, 1H), 4.29 (s, 1H), 4.07-4.03 (m, 1H), 1.42 (d, <i>J</i> =6.4 Hz, 3H). |
|  155 | SFC AD 4.6×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.27 (s, 1H), 7.26 (d, <i>J</i> =8.2 Hz, 1H), 7.09 (d, <i>J</i> =7.7 Hz, 1H), 4.83 (d, <i>J</i> =2.2 Hz, 2H), 4.27 (qd, <i>J</i> =6.7, 2.6 Hz, 1H), 4.15 (d, <i>J</i> =2.4 Hz, 1H), 2.39 (s, 3H), 1.33 (d, <i>J</i> =6.7 Hz, 3H). |
|  156 | SFC AD 4.6×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (500 MHz, CD ₃ OD) δ 7.27 (s, 1H), 7.26 (d, <i>J</i> =8.6 Hz, 1H), 7.09 (d, <i>J</i> =7.7 Hz, 1H), 4.83 (s, 2H), 4.28 (qd, <i>J</i> =6.6, 2.6 Hz, 1H), 4.15 (d, <i>J</i> =2.4 Hz, 1H), 2.39 (s, 3H), 1.33 (d, <i>J</i> =6.7 Hz, 3H). |
|  157 | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26 (s, 1H), 7.24 (d, <i>J</i> =8.0 Hz, 1H), 7.08 (d, <i>J</i> =8.0 Hz, 1H), 4.93 (d, <i>J</i> =12.0 Hz, 1H), 4.84 (d, <i>J</i> =15.3 Hz, 1H), 4.22 (s, 1H), 4.03 (dd, <i>J</i> =6.6, 1.6 Hz, 1H), 2.38 (s, 3H), 1.41 (d, <i>J</i> =6.6 Hz, 3H). |
|  158 | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26 (s, 1H), 7.25 (d, <i>J</i> =8.1 Hz, 1H), 7.08 (d, <i>J</i> =7.7 Hz, 1H), 4.94 (d, <i>J</i> =12.0 Hz, 1H), 4.84 (d, <i>J</i> =15.4 Hz, 1H), 4.23 (s, 1H), 4.03 (dd, <i>J</i> =6.6, 1.6 Hz, 1H), 2.38 (s, 3H), 1.41 (d, <i>J</i> =6.6 Hz, 3H). |
|  159 | SFC IC 20×250mm, 5µm 80:20 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.47-7.44 (m, 2H), 7.27-7.17 (m, 1H), 4.88 (s, 2H), 4.48-4.43 (m, 2H), 1.28 (d, <i>J</i> =6.8 Hz, 3H). |
|  160 | SFC IC 20×250mm, 5µm 80:20 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.52-7.41 (m, 2H), 7.25-7.16 (m, 1H), 4.88 (s, 2H), 4.53-4.39 (m, 2H), 1.28 (d, <i>J</i> =6.8 Hz, 3H). |

| Example | Separation Conditions | Analytical Data |
|---------|--|---|
| | SFC AY 30×250mm, 5µm 80:20 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 8.48 (bs, 3 H), 7.10 (dd, <i>J</i> ₁ = 6.4 Hz, <i>J</i> ₂ = 20 Hz, 2 H), 4.80 (d, <i>J</i> = 12.4 Hz, 1 H), 4.61 (d, <i>J</i> = 12.8 Hz, 1 H), 4.39 (s, 1 H), 4.27 (d, <i>J</i> = 9.6 Hz, 1 H), 3.73 (d, <i>J</i> = 9.6 Hz, 1 H), 2.38 (s, 3 H), 2.10 (s, 3 H). |
| | SFC AY 30×250mm, 5µm 80:20 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 8.52 (bs, 3 H), 7.09 (m, 2 H), 4.81 (d, <i>J</i> = 13.2 Hz, 1 H), 4.61 (d, <i>J</i> = 12.4 Hz, 1 H), 4.39 (d, <i>J</i> = 2.8 Hz, 1 H), 4.28 (d, <i>J</i> = 10.0 Hz, 1 H), 3.72 (d, <i>J</i> = 9.6 Hz, 1 H), 2.39 (s, 3 H), 2.09 (s, 3 H). |
| | SFC IC 4.6×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.34 (t, <i>J</i> = 7.6 Hz, 1 H), 7.23 (d, <i>J</i> = 7.5 Hz, 1 H), 7.03 (d, <i>J</i> = 7.7 Hz, 1 H), 4.88 (s, 2 H), 4.49 – 4.35 (m, 2 H), 2.45 (s, 3 H), 1.27 (d, <i>J</i> = 6.7 Hz, 3 H). |
| | SFC IC 4.6×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.34 (t, <i>J</i> = 7.6 Hz, 1 H), 7.23 (d, <i>J</i> = 7.5 Hz, 1 H), 7.03 (d, <i>J</i> = 7.7 Hz, 1 H), 4.88 (s, 2 H), 4.48 – 4.37 (m, 2 H), 2.46 (s, 3 H), 1.27 (d, <i>J</i> = 6.7 Hz, 3 H). |
| | HPLC AY-H 20×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.32 (t, <i>J</i> = 7.6 Hz, 1 H), 7.22 (d, <i>J</i> = 7.5 Hz, 1 H), 7.02 (d, <i>J</i> = 7.7 Hz, 1 H), 4.99 (d, <i>J</i> = 15.6 Hz, 1 H), 4.87 (s, 1 H), 4.44 (s, 1 H), 4.08 – 3.99 (m, 1 H), 2.45 (s, 3 H), 1.44 (d, <i>J</i> = 6.6 Hz, 3 H). |
| | HPLC AY-H 20×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.32 (t, <i>J</i> = 7.6 Hz, 1 H), 7.22 (d, <i>J</i> = 7.5 Hz, 1 H), 7.02 (d, <i>J</i> = 7.6 Hz, 1 H), 4.99 (d, <i>J</i> = 15.6 Hz, 1 H), 4.87 (s, 1 H), 4.43 (s, 1 H), 4.03 (d, <i>J</i> = 6.6 Hz, 1 H), 2.44 (s, 3 H), 1.43 (d, <i>J</i> = 6.6 Hz, 3 H). |
| | SFC OD 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.13 (d, <i>J</i> = 8.2 Hz, 1 H), 7.04-7.01 (m, 2 H), 4.86-4.75 (m, 2 H), 4.25 (tt, <i>J</i> = 6.7, 3.3 Hz, 1 H), 4.17 (d, <i>J</i> = 2.3 Hz, 1 H), 3.85 (s, 3 H), 1.34 (d, <i>J</i> = 6.7 Hz, 3 H). |
| | SFC OD 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.13 (d, <i>J</i> = 9.3 Hz, 1 H), 7.04-7.01 (m, 2 H), 4.81 (d, <i>J</i> = 1.7 Hz, 2 H), 4.25 (dd, <i>J</i> = 6.6, 2.4 Hz, 1 H), 4.17 (s, 1 H), 3.85 (s, 3 H), 1.34 (d, <i>J</i> = 6.7 Hz, 3 H). |


| Example | Separation Conditions | Analytical Data |
|---|---|---|
|  <p>173</p> | SFC AD 4.6×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.11 (s, 1H), 7.02 (d, <i>J</i> = 6.4 Hz, 2H), 4.90 (s, 1H), 4.84 (s, 1H), 4.24 (s, 1H), 4.03 (s, 1H), 1.41 (d, <i>J</i> = 6.6 Hz, 3H). |
|  <p>174</p> | SFC AD 4.6×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 194 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.12 (d, <i>J</i> = 9.2 Hz, 1H), 7.05 – 6.99 (m, 2H), 4.90 (s, 1H), 4.82 (d, <i>J</i> = 15.0 Hz, 1H), 4.24 (s, 1H), 4.03 (dd, <i>J</i> = 6.6, 1.6 Hz, 1H), 3.85 (s, 3H), 1.41 (d, <i>J</i> = 6.6 Hz, 3H). |
|  <p>175</p> | SFC IC 30×250mm, 5µm 80:20 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.39 (d, <i>J</i> = 8.0 Hz, 1H), 7.27 (d, <i>J</i> = 8.4 Hz, 1H), 5.04 (d, <i>J</i> = 16.0 Hz, 1H), 4.72 (d, <i>J</i> = 16.4 Hz, 1H), 4.59-4.58 (m, 1H), 4.35-4.30 (m, 1H), 3.90 (dd, <i>J</i> = 12.8, 1.2 Hz, 1H), 2.48-2.47 (m, 3H). |
|  <p>176</p> | SFC IC 30×250mm, 5µm 80:20 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 198 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.39 (d, <i>J</i> = 8.0 Hz, 1H), 7.26 (d, <i>J</i> = 8.4 Hz, 1H), 5.04 (d, <i>J</i> = 16.4 Hz, 1H), 4.71 (d, <i>J</i> = 16.0 Hz, 1H), 4.58-4.56 (m, 1H), 4.33-4.29 (m, 1H), 3.90 (dd, <i>J</i> = 12.4, 1.2 Hz, 1H), 2.47-2.46 (m, 3H). |
|  <p>177</p> | SFC AD-H 20×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 175 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.29 (s, 1H), 7.26-7.24 (d, <i>J</i> = 8.0 Hz, 1H), 7.09-7.07 (d, <i>J</i> = 7.9 Hz, 1H), 4.80 (s, 2H), 4.21 (s, 1H), 4.03-3.99 (m, 1H), 2.39 (s, 3H), 1.66-1.56 (m, 2H), 1.10-1.06 (t, <i>J</i> = 7.4 Hz, 3H). |
|  <p>178</p> | SFC AD-H 20×250mm, 5µm 90:10 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 175 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.30 (s, 1H), 7.26-7.24 (d, <i>J</i> = 8.0 Hz, 1H), 7.09-7.07 (d, <i>J</i> = 7.9 Hz, 1H), 4.80 (s, 2H), 4.21 (s, 1H), 4.04-4.00 (m, 1H), 2.39 (s, 3H), 1.66-1.57 (m, 2H), 1.10-1.06 (t, <i>J</i> = 7.4 Hz, 3H). |
|  <p>179</p> | HPLC AY-H 50×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) FME | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.27-7.25 (m, 2H), 7.09 (d, <i>J</i> = 7.6 Hz, 1H), 4.98 (d, <i>J</i> = 15.2 Hz, 1H), 4.83 (d, <i>J</i> = 15.6 Hz, 1H), 4.29 (s, 1H), 3.75 – 3.72 (m, 1H), 2.39 (s, 3H), 1.76-1.66 (m, 2H), 1.16 (t, <i>J</i> = 7.2 Hz, 3H). |

| Example | Separation Conditions | Analytical Data |
|---|--|--|
|  | HPLC AY-H 50×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) SME | MS (ESI): <i>m/z</i> 192 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.27– 7.25 (m, 2H), 7.09 (d, <i>J</i> = 7.6 Hz, 1H), 4.98 (d, <i>J</i> = 15.6 Hz, 1H), 4.83 (d, <i>J</i> = 15.2 Hz, 1H), 4.29 (s, 1H), 3.75 – 3.72 (m, 1H), 2.39 (s, 3H), 1.76-1.66 (m, 2H), 1.14 (t, <i>J</i> = 7.6 Hz, 3H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.30 - 7.20 (m, 3H), 4.82 (s, 2H), 4.29 (s, 1H), 4.03 - 4.00 (m, 1H), 1.70 – 1.56 (m, 2H), 1.09 (t, <i>J</i> = 7.6 Hz, 3H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.27 - 7.18 (m, 3H), 4.82 (s, 2H), 4.27 (d, <i>J</i> = 2.0 Hz, 1H), 4.02 - 3.98 (m, 1H), 1.70 – 1.54 (m, 2H), 1.09 (t, <i>J</i> = 7.6 Hz, 3H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26– 7.20 (m, 3H), 4.84 (s, 2H), 4.27-4.22 (m, 2H), 1.33 (d, <i>J</i> = 6.8 Hz, 3H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26– 7.19 (m, 3H), 4.84 (s, 2H), 4.28-4.22 (m, 2H), 1.33 (d, <i>J</i> = 6.8 Hz, 2H). |
|  | SFC IC 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) FME | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.27-7.17 (m, 3 H), 4.96 (d, <i>J</i> = 15.6 Hz, 1 H), 4.85 (d, <i>J</i> = 15.6 Hz, 1 H), 4.32 (s, 1 H), 4.07-4.02 (m, 1 H), 1.42 (d, <i>J</i> = 6.4 Hz, 3 H). |
|  | SFC IC 20×250mm, 5µm 85:15 CO ₂ /MeOH (0.2% NH ₃) SME | MS (ESI): <i>m/z</i> 182 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.27-7.17 (m, 3 H), 4.96 (d, <i>J</i> = 15.2 Hz, 1 H), 4.85 (d, <i>J</i> = 15.6 Hz, 1 H), 4.32 (s, 1 H), 4.07-4.02 (m, 1 H), 1.42 (d, <i>J</i> = 6.4 Hz, 3 H). |
|  | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) FME | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26-7.18 (m, 3H), 5.03 (d, <i>J</i> = 15.2 Hz, 1H), 4.86-4.83 (d, <i>J</i> = 15.2 Hz, 1H), 4.37 (s, 1H), 3.77-3.73 (m, 1H), 1.79-1.65 (m, 2H), 1.16-1.12 (t, <i>J</i> = 7.6 Hz, 3H). |

| Example | Separation Conditions | Analytical Data |
|---|---|--|
|  188 | SFC IC 30×250mm, 5µm 90:10 CO ₂ /MeOH (0.1% NH ₄ OH) SME | MS (ESI): <i>m/z</i> 196 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.26-7.18 (m, 3H), 5.03 (d, <i>J</i> = 15.2 Hz, 1H), 4.86-4.83 (d, <i>J</i> = 15.2 Hz, 1H), 4.37 (s, 1H), 3.77-3.73 (m, 1H), 1.79-1.65 (m, 2H), 1.16-1.12 (t, <i>J</i> = 7.6 Hz, 3H). |
|  189 | SFC AY-H 4.7×100mm, 5µm EtOH (1% NH ₃) FME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.33 (d, <i>J</i> = 8.0 Hz, 1H), 7.17 (d, <i>J</i> = 8.0 Hz, 1H), 7.01 (s, 1H), 4.95-4.82 (m, 2H), 4.23 (s, 1H), 4.05-4.04 (m, 1H), 2.35 (s, 3H), 1.41 (d, <i>J</i> = 6.8 Hz, 3H). |
|  190 | SFC AY-H 4.7×100mm, 5µm EtOH (1% NH ₃) SME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.33 (d, <i>J</i> = 8.0 Hz, 1H), 7.17 (d, <i>J</i> = 8.0 Hz, 1H), 7.01 (s, 1H), 4.95-4.82 (m, 2H), 4.23 (s, 1H), 4.05-4.04 (m, 1H), 2.35 (s, 3H), 1.41 (d, <i>J</i> = 6.8 Hz, 3H). |
|  191 | HPLC AY 20×250mm, 10µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) FME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.47-7.42 (m, 2H), 7.38 (t, <i>J</i> = 7.6 Hz, 1H), 7.22 (d, <i>J</i> = 7.6 Hz, 1H), 5.05 (d, <i>J</i> = 15.6 Hz, 1H), 4.90 (d, <i>J</i> = 15.6 Hz, 1H), 4.35 (s, 1H), 3.78-3.75 (m, 1H), 1.79-1.66 (m, 2H), 1.16 (t, <i>J</i> = 7.2 Hz, 3H). |
|  192 | HPLC AY 20×250mm, 10µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) SME | MS (ESI): <i>m/z</i> 178 [M+H]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.48-7.42 (m, 2H), 7.38 (t, <i>J</i> = 7.2 Hz, 1H), 7.22 (d, <i>J</i> = 7.6 Hz, 1H), 5.05 (d, <i>J</i> = 15.2 Hz, 1H), 4.90 (d, <i>J</i> = 15.2 Hz, 1H), 4.36 (s, 1H), 3.78-3.75 (m, 1H), 1.78-1.69 (m, 2H), 1.16 (t, <i>J</i> = 7.2 Hz, 3H). |
|  193 | SFC OD-H 20×250mm, 5µm 90:10 CO ₂ /MeOH (1% NH ₃) FME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.48-7.36 (m, 3H), 7.21-7.19 (d, <i>J</i> = 7.5 Hz, 1H), 4.85 (s, 2H), 4.27 (d, <i>J</i> = 2.1 Hz, 1H), 4.05-4.01 (m, 1H), 1.67-1.56 (m, 2H), 1.09-1.07 (t, <i>J</i> = 7.4 Hz, 3H). |
|  195 | SFC OD-H 20×250mm, 5µm 90:10 CO ₂ /MeOH (1% NH ₃) SME | MS (ESI): <i>m/z</i> 161 [M-16]. ¹ H NMR (400 MHz, CD ₃ OD) δ 7.46-7.36 (m, 3H), 7.21-7.19 (d, <i>J</i> = 7.5 Hz, 1H), 4.85 (s, 2H), 4.25 (d, <i>J</i> = 2.1 Hz, 1H), 4.04-4.00 (m, 1H), 1.66-1.55 (m, 2H), 1.11-1.07 (t, <i>J</i> = 7.4 Hz, 3H). |
|  252 | SFC OZ 20×250mm, 5µm 75:25 CO ₂ /EtOH (1% NH ₃) FME | MS (ESI): <i>m/z</i> 178 [M + H] ⁺ . ¹ H NMR (400 MHz, CD ₃ OD) δ 7.32 (d, <i>J</i> = 7.6 Hz, 1H), 7.29 (d, <i>J</i> = 8.0 Hz, 1H), 7.03 (s, 1H), 4.87 (s, 2H), 4.28-4.23 (m, 1H), 4.15 (d, <i>J</i> = 2.0 Hz, 1H), 2.37 (s, 3H), 1.32 (d, <i>J</i> = 2.8 Hz, 3H). |

| Example | Separation Conditions | Analytical Data |
|---------|--|--|
| | SFC OZ 20×250mm, 5µm 75:25 CO ₂ /EtOH (1% NH ₃) SME | MS (ESI): <i>m/z</i> 178 [M + H] ⁺ . ¹ H NMR (400 MHz, CD ₃ OD) δ d 7.35 (d, <i>J</i> = 6.0 Hz, 1H), 7.29 (d, <i>J</i> = 8.0 Hz, 1H), 7.03 (s, 1H), 4.88 (s, 2H), 4.30-4.26 (m, 1H), 4.16 (s, 1H), 2.37 (s, 3H), 1.32 (d, <i>J</i> = 2.4 Hz, 3H). |
| | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) FME | MS (ESI): <i>m/z</i> 178 [M + H] ⁺ . ¹ H NMR (400 MHz, D ₂ O) δ 7.22-7.21 (m, 2H), 7.19-7.16 (m, 1H), 5.15 (q, <i>J</i> = 6.8 Hz, 1H), 4.34-4.31 (m, 2H), 3.95 (dd, <i>J</i> = 14.0, 2.0 Hz, 1H), 2.17 (s, 3H), 1.40 (d, <i>J</i> = 7.8 Hz, 3H). |
| | HPLC AY-H 4.6×250mm, 5µm 90:10 Hexanes/EtOH (0.1% Et ₂ NH) SME | MS (ESI): <i>m/z</i> 178 [M + H] ⁺ . ¹ H NMR (400 MHz, D ₂ O) δ 7.22-7.21 (m, 2H), 7.19-7.16 (m, 1H), 5.15 (q, <i>J</i> = 6.8 Hz, 1H), 4.34-4.31 (m, 2H), 3.95 (dd, <i>J</i> = 14.0, 2.0 Hz, 1H), 2.17 (s, 3H), 1.40 (d, <i>J</i> = 7.8 Hz, 3H). |

[0169] Chiral Separation of tert-butyl (2,4,7,8,9,10-hexahydro-1H-7,10-methanobenzo-f-isochromen-1-yl)carbamate



[0170] As previously described in the General Procedure for Chiral Separation above compound **Boc-51** (prepared as previously described in Example 51) was separated into 3 peaks consisting of a mixture of enantiomers (**P1** and **P4**) and the individual enantiomers **P2** and **P3** using an AD 20×250mm, 5 um column and a mobile phase of 85:15 CO₂/MeOH/0.5% NH₃. The mixture of **P1** and **P4** was further separated by SFC into enantiomers **P1** and **P4** using a Whelk 20×250mm, 5 um column and a mobile phase of 85:15 CO₂/ MeOH(0.5% NH₃).

[0171] The individual Boc-protected enantiomers were deprotected and converted to the corresponding HCl salts as previously described in Example 1.

[0172] **Example 161.** FME on AD column, MS (ESI): *m/z* 216 [M+H]. ¹H NMR (400 MHz, CD₃OD): δ 7.22 (d, *J* = 7.6 Hz, 1H), 6.87 (d, *J* = 7.6 Hz, 1H), 4.89 (d, *J* = 15.2 Hz, 1H), 4.79 (d, *J* = 15.2 Hz, 1H), 4.56 (s, 1H), 4.32 (dd, *J* = 12.8, 1.2 Hz, 1H), 3.91 (d, *J* = 12.8, 2.0 Hz, 1H), 3.61 (s, 1H), 3.41 (d, *J* = 2.0 Hz, 1H), 2.06-2.01 (m, 2H), 1.73-1.70 (m, 1H), 1.64-1.61 (m, 1H), 1.20-1.15 (m, 2H).

[0173] **Example 162.** SME on AD column, MS (ESI): *m/z* 216 [M+H]. ¹H NMR (400 MHz, CD₃OD): δ 7.20 (d, *J* = 7.6 Hz, 1H), 6.87 (d, *J* = 7.6 Hz, 1H), 4.94 (d, *J* = 15.6 Hz, 1H), 4.77 (d, *J* = 15.2 Hz, 1H), 4.43 (s, 1H), 4.25 (dd, *J* = 12.8, 0.8 Hz, 1H), 3.91 (dd, *J* = 12.8, 2.0 Hz, 1H), 3.66 (s, 1H), 3.39 (s, 1H), 1.96-1.99 (m, 2H), 1.84-1.81 (m, 1H), 1.65-1.62 (m, 1H), 1.10-1.06 (m, 2H).

[0174] **Example 163.** FME on Whelk column, MS (ESI): *m/z* 216 [M+H]. ¹H NMR (400 MHz, CD₃OD): δ 7.20 (d, *J* = 7.6 Hz, 1H), 6.87 (d, *J* = 7.6 Hz, 1H), 4.94 (d, *J* = 15.6 Hz, 1H), 4.77 (d, *J* = 15.2 Hz, 1H), 4.43 (s, 1H), 4.25 (dd, *J* = 12.8, 0.8 Hz, 1H), 3.91 (dd, *J* = 12.8, 2.0 Hz, 1H), 3.66 (s, 1H), 3.39 (s, 1H), 1.99-1.96 (m, 2H), 1.84-1.81 (m, 1H), 1.65-1.62 (m, 1H), 1.10-1.06 (m, 2H).

[0175] **Example 164.** SME on Whelk column, MS (ESI): *m/z* 216 [M+H]. ¹H NMR (400 MHz, CD₃OD): δ 7.22 (d, *J* = 7.6 Hz, 1H), 6.87 (d, *J* = 7.6 Hz, 1H), 4.89 (d, *J* = 15.2 Hz, 1H), 4.79 (d, *J* = 15.2 Hz, 1H), 4.56 (s, 1H), 4.32 (dd, *J* = 12.8, 1.2 Hz, 1H), 3.91 (dd, *J* = 12.8, 2.0 Hz, 1H), 3.61 (s, 1H), 3.41 (d, *J* = 2.0 Hz, 1H), 2.06-2.01 (m, 2H), 1.73-1.70 (m, 1H), 1.64-1.61 (m, 1H), 1.20-1.15 (m, 2H).

Example 289
In Vivo Biological Studies

[0176] Neuropharmacological Assay (SmartCube™)

[0177] In order to demonstrate the utility of the provided compounds to treat neurological and psychiatric diseases and disorders, exemplary compounds were evaluated using the neuropharmacological screen described in S.L. Roberds *et al.*, *Front. Neurosci.* 2011 Sep 9;5:103 (doi: 10.3389/fnins.2011.00103) ("Roberds"). As reported in *Roberds*, because psychiatric diseases generally result from disorders of cell-cell communication or circuitry, intact systems are useful in detecting improvement in disease-relevant endpoints. These endpoints are typically behavioral in nature, often requiring human observation and interpretation. To facilitate testing of multiple compounds for behavioral effects relevant to psychiatric disease, PsychoGenics, Inc. (Tarrytown, NY, "PGI") developed SmartCube™, an automated system in which behaviors of compound-treated mice are captured by digital video and analyzed with computer algorithms. (D. Brunner *et al.*, *Drug Discov. Today* 2002, 7:S107-S112). PGI Analytical Systems uses data from SmartCube™ to compare the behavioral signature of a test compound to a database of behavioral signatures obtained using a large set of diverse reference compounds. (The composition of the database as well as validation of the method is further described in *Roberds*). In this way, the neuropharmacological effects of a test compound can be predicted by similarity to major classes of compounds, such as antipsychotics, anxiolytics and antidepressants.

[0178] The SmartCube™ system produces an activity signature indicating the probability that the activity of the test compound at the administered dose matches a given class of neuropharmacological agents. (See, e.g., *Roberds*, Figures 2 and 3). The test compound is simultaneously compared against multiple classes of agents; thus, a separate probability is generated for each behavioral effect measured (e.g., anxiolytic activity, analgesic activity, etc.). In the table below, these probabilities are reported for each behavioral effect measured as follows:

| | | |
|-------|------|--------|
| LOQ ≤ | + | < 5% |
| 5% ≤ | ++ | < 25 % |
| 25% ≤ | +++ | < 50 % |
| 50% ≤ | ++++ | |

where LOQ is the limit of quantification.

[0179] Provided compounds were dissolved in a mixture of Pharmasolve™ (N-methyl-2-pyrrolidone), polyethylene glycol and propylene glycol, and were injected i.p. 15 min. before the behavioral test. For each compound, injections were administered at 3 different doses. For each behavioral effect measured, results for the most efficacious dose(s) are presented. In the table below, DP: anti-depressant; AX: anxiolytic; SD: sedative hypnotic; PS: anti-psychotic; MS: mood stabilizer; AD: ADHD; CE: cognitive enhancer; AG: analgesic; UN: uncharacterized CNS activity.

[0180] The potency of many of the compounds in the table was also determined in the SmartCube™ system. Test compounds were routinely examined at dose levels of 0.3, 1, 3, 10 and 30 mg per kg (mpk), although the dose range was increased or decreased if necessary to obtain a full dose response curve. A compound's minimal effective dose (MED) is a measure of the compounds potency. The MED was defined as the dose (in mpk) having 50% or more total activity in SmartCube. The potencies of the compounds are shown in the table below, with potency values in mpk binned in the following manner:

| MED mpk range | BIN |
|----------------|-----|
| ≤3 mpk | A |
| >3 to 10 mpk | B |
| >10 to ≤30 mpk | C |
| >30 mpk | D |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|----|-----|-----|-----|----|----|-----|----|------|---------|
| 1 | + | ++ | ++ | ++ | + | + | ++ | ++ | ++ | C |
| 2 | + | ++ | + | + | + | + | + | ++ | +++ | C |
| 3 | ++ | ++ | ++ | + | + | + | + | ++ | +++ | C |
| 4 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | + | C |
| 5 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 6 | ++ | ++ | ++ | + | + | + | + | ++ | ++++ | B |
| 7 | ++ | ++ | + | ++ | ++ | + | ++ | + | +++ | C |
| 8 | ++ | ++ | +++ | ++ | + | + | ++ | ++ | ++ | C |
| 9 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 10 | ++ | ++ | ++ | +++ | + | + | +++ | ++ | ++ | B |
| 11 | ++ | + | ++ | + | + | + | ++ | + | + | C |
| 12 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 13 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 14 | ++ | ++ | + | + | + | + | + | ++ | ++++ | C |
| 15 | + | + | + | + | + | + | + | + | + | D |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|-----|-----|----|-----|----|----|-----|-----|------|---------|
| 16 | + | +++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 17 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 18 | ++ | ++ | ++ | ++ | + | + | ++ | +++ | + | C |
| 19 | + | +++ | ++ | ++ | + | + | ++ | ++ | +++ | C |
| 20 | ++ | +++ | ++ | +++ | + | + | ++ | ++ | ++ | B |
| 21 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 22 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 23 | ++ | +++ | + | ++ | + | + | ++ | ++ | ++ | C |
| 24 | ++ | ++ | ++ | ++ | + | + | + | + | + | C |
| 25 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 26 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | A |
| 27 | ++ | ++ | + | ++ | ++ | + | ++ | ++ | +++ | B |
| 28 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 29 | ++ | ++ | + | + | + | + | + | +++ | + | C |
| 30 | + | + | + | + | + | + | + | ++ | + | D |
| 31 | ++ | ++ | + | + | + | + | + | ++ | +++ | C |
| 32 | ++ | ++ | ++ | ++ | + | + | + | ++ | + | C |
| 33 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 34 | + | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 35 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 36 | ++ | ++ | ++ | ++ | + | + | +++ | ++ | ++ | B |
| 37 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 38 | ++ | ++ | + | ++ | + | + | + | ++ | +++ | C |
| 39 | ++ | +++ | + | + | + | + | ++ | ++ | ++++ | B |
| 40 | + | +++ | + | ++ | + | + | ++ | ++ | ++ | B |
| 41 | ++ | ++ | + | + | + | + | ++ | ++ | +++ | C |
| 42 | ++ | ++ | ++ | +++ | + | + | ++ | +++ | +++ | A |
| 43 | + | +++ | ++ | ++ | + | + | ++ | ++ | + | C |
| 44 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 45 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 46 | + | ++ | ++ | + | + | + | ++ | ++ | ++++ | A |
| 47 | ++ | ++ | + | ++ | ++ | + | ++ | ++ | ++++ | A |
| 48 | + | +++ | ++ | ++ | + | + | +++ | ++ | ++ | A |
| 49 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 50 | +++ | ++ | + | ++ | + | + | ++ | ++ | ++ | C |
| 51 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 52 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | + | B |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|----|------|----|-----|----|----|-----|-----|------|---------|
| 53 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 54 | ++ | ++ | + | ++ | + | + | + | ++ | +++ | C |
| 55 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 56 | ++ | ++ | + | + | + | + | ++ | +++ | + | C |
| 57 | ++ | ++ | ++ | ++ | + | + | +++ | ++ | +++ | B |
| 58 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 59 | ++ | ++ | ++ | ++ | + | + | ++ | +++ | +++ | B |
| 60 | ++ | +++ | + | ++ | + | + | ++ | ++ | + | C |
| 61 | + | ++ | + | ++ | + | + | ++ | ++ | ++++ | C |
| 62 | ++ | ++ | + | ++ | ++ | + | +++ | ++ | ++++ | B |
| 63 | ++ | ++ | + | + | + | + | ++ | ++ | ++ | C |
| 68 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++ | C |
| 72 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 73 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 74 | ++ | ++ | + | + | + | + | + | ++ | +++ | C |
| 75 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 76 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 77 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 78 | ++ | ++ | + | +++ | + | + | ++ | ++ | ++++ | B |
| 79 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++ | B |
| 80 | + | +++ | ++ | ++ | + | + | + | ++ | +++ | A |
| 81 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 82 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 83 | + | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 84 | ++ | ++++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 85 | ++ | ++ | + | + | + | + | ++ | +++ | +++ | B |
| 86 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 87 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 88 | + | +++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 89 | + | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 90 | ++ | +++ | ++ | + | + | + | ++ | ++ | + | C |
| 91 | + | + | ++ | ++ | + | + | ++ | ++ | +++ | C |
| 92 | ++ | ++ | + | + | + | + | + | + | ++++ | C |
| 93 | ++ | +++ | + | ++ | + | + | + | ++ | + | C |
| 94 | + | ++ | + | + | + | + | + | ++ | ++ | C |
| 95 | ++ | ++ | + | +++ | + | + | ++ | ++ | +++ | A |
| 96 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | A |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|-----|-----|-----|-----|----|----|-----|----|------|---------|
| 97 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | B |
| 98 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 99 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 100 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | +++ | A |
| 101 | ++ | +++ | ++ | ++ | + | + | + | ++ | + | C |
| 104 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 105 | ++ | ++ | + | ++ | ++ | + | ++ | ++ | +++ | C |
| 106 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | B |
| 107 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 111 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 112 | ++ | ++ | + | ++ | + | + | +++ | ++ | + | C |
| 113 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 114 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 115 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | +++ | B |
| 116 | ++ | ++ | +++ | ++ | + | + | ++ | ++ | +++ | B |
| 117 | ++ | ++ | + | +++ | ++ | + | ++ | ++ | + | B |
| 118 | +++ | ++ | ++ | ++ | + | + | ++ | ++ | + | B |
| 119 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 120 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++ | B |
| 121 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 122 | ++ | +++ | ++ | ++ | + | + | +++ | ++ | +++ | A |
| 123 | ++ | ++ | ++ | ++ | + | + | + | ++ | +++ | A |
| 124 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 125 | ++ | ++ | ++ | ++ | + | + | ++ | + | ++++ | A |
| 126 | ++ | ++ | + | +++ | + | + | ++ | ++ | ++++ | A |
| 127 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 128 | ++ | ++ | + | ++ | + | + | +++ | ++ | +++ | A |
| 129 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 130 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | ++++ | B |
| 131 | ++ | ++ | ++ | ++ | ++ | + | +++ | + | +++ | A |
| 132 | ++ | +++ | ++ | ++ | + | + | +++ | ++ | +++ | A |
| 133 | ++ | ++ | + | +++ | + | + | ++ | ++ | ++++ | A |
| 134 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 135 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 136 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 137 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 138 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|-----|------|----|-----|----|----|----|-----|------|---------|
| 139 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 140 | ++ | ++ | + | +++ | ++ | + | ++ | ++ | + | C |
| 141 | ++ | +++ | + | ++ | ++ | + | ++ | ++ | +++ | B |
| 142 | ++ | ++ | + | + | + | + | ++ | ++ | ++++ | B |
| 143 | ++ | ++ | ++ | + | + | + | ++ | +++ | ++ | B |
| 144 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 145 | + | ++ | + | + | + | + | ++ | ++ | +++ | C |
| 146 | + | +++ | + | + | + | + | ++ | ++ | ++ | B |
| 147 | ++ | ++ | + | + | + | + | + | ++ | ++++ | B |
| 148 | ++ | ++ | ++ | ++ | ++ | + | ++ | ++ | +++ | B |
| 149 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | +++ | A |
| 150 | + | ++ | + | ++ | + | + | + | ++ | ++++ | B |
| 151 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++ | B |
| 152 | ++ | +++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 153 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 154 | + | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 155 | ++ | ++++ | ++ | ++ | + | + | ++ | ++ | ++ | B |
| 156 | + | ++ | + | + | + | + | ++ | ++ | +++ | B |
| 157 | ++ | ++ | + | +++ | ++ | + | ++ | ++ | ++++ | A |
| 158 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | ++ | B |
| 159 | +++ | +++ | + | + | + | + | + | + | ++ | B |
| 160 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | A |
| 161 | ++ | +++ | + | ++ | + | + | ++ | ++ | ++ | B |
| 162 | + | +++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 163 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | +++ | B |
| 164 | + | +++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 165 | ++ | ++ | + | ++ | + | + | ++ | + | ++++ | B |
| 166 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | B |
| 167 | + | ++ | + | ++ | + | + | + | ++ | ++++ | A |
| 168 | ++ | ++ | + | +++ | + | + | ++ | + | ++++ | B |
| 169 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 170 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | A |
| 171 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | +++ | A |
| 172 | + | ++ | + | ++ | + | + | + | ++ | +++ | C |
| 173 | + | ++ | + | + | + | + | ++ | ++ | ++++ | B |
| 174 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 175 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | A |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|-----|------|-----|-----|----|----|----|-----|------|---------|
| 176 | ++ | ++ | + | ++ | + | + | ++ | + | ++++ | B |
| 177 | + | ++++ | + | ++ | + | + | ++ | ++ | ++++ | A |
| 178 | ++ | ++ | ++ | ++ | + | + | + | ++ | +++ | C |
| 179 | +++ | ++ | + | ++ | + | + | ++ | + | ++++ | A |
| 180 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 181 | + | ++ | +++ | ++ | + | + | ++ | ++ | ++++ | A |
| 182 | ++ | ++ | + | +++ | + | + | ++ | + | ++++ | A |
| 183 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 184 | ++ | ++ | + | ++ | + | + | ++ | + | ++++ | A |
| 185 | + | +++ | + | ++ | + | + | ++ | ++ | ++++ | A |
| 186 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 187 | + | ++ | + | +++ | + | + | ++ | ++ | ++++ | A |
| 188 | ++ | ++ | + | ++ | + | + | + | ++ | ++++ | A |
| 189 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | C |
| 190 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | B |
| 191 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 192 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | A |
| 193 | + | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 194 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 195 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 196 | + | ++ | + | ++ | + | + | ++ | + | +++ | C |
| 197 | + | ++ | + | + | + | + | + | + | ++++ | B |
| 198 | ++ | ++ | + | +++ | + | + | ++ | + | +++ | A |
| 199 | + | ++++ | ++ | ++ | + | + | ++ | ++ | ++++ | A |
| 200 | ++ | +++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 201 | + | ++ | + | ++ | + | + | ++ | + | ++++ | B |
| 202 | ++ | +++ | + | ++ | + | + | ++ | ++ | ++++ | A |
| 203 | ++ | ++ | ++ | +++ | + | + | ++ | ++ | ++++ | B |
| 204 | + | ++ | ++ | ++ | + | + | ++ | + | ++++ | B |
| 205 | ++ | +++ | ++ | ++ | + | + | ++ | + | +++ | B |
| 206 | ++ | ++++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 207 | ++ | ++ | + | ++ | + | + | ++ | ++ | ++++ | B |
| 208 | ++ | +++ | + | ++ | ++ | + | ++ | + | +++ | B |
| 209 | ++ | ++ | ++ | ++ | + | + | ++ | +++ | +++ | B |
| 210 | + | ++ | + | + | + | + | ++ | ++ | ++++ | B |
| 211 | ++ | ++ | +++ | ++ | + | + | + | + | ++++ | B |
| 212 | + | ++ | + | + | + | + | + | ++ | ++++ | C |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|----|------|-----|-----|----|----|-----|----|------|---------|
| 213 | ++ | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 214 | ++ | ++ | ++ | ++ | + | ++ | ++ | ++ | ++++ | B |
| 215 | ++ | +++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 216 | ++ | ++ | + | + | + | + | + | ++ | ++++ | B |
| 217 | ++ | +++ | + | ++ | + | + | ++ | + | ++++ | C |
| 218 | + | ++ | + | ++ | + | + | ++ | + | ++++ | A |
| 219 | + | ++ | + | + | + | + | ++ | ++ | ++++ | B |
| 220 | + | ++++ | + | + | + | + | ++ | ++ | ++++ | B |
| 221 | ++ | ++ | + | ++ | + | + | + | + | ++++ | B |
| 222 | + | ++ | + | + | + | + | + | + | ++++ | B |
| 223 | ++ | ++ | ++ | + | + | + | + | + | +++ | B |
| 224 | + | ++ | ++ | + | + | + | +++ | ++ | ++ | C |
| 225 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | ++++ | B |
| 226 | ++ | ++ | + | +++ | + | + | ++ | + | ++ | C |
| 227 | + | ++ | + | ++ | + | + | + | ++ | ++++ | B |
| 228 | ++ | ++ | + | ++ | + | + | + | ++ | +++ | C |
| 229 | ++ | ++ | + | ++ | + | + | ++ | + | +++ | B |
| 230 | + | ++ | + | ++ | + | + | ++ | + | ++++ | B |
| 231 | ++ | ++ | + | + | + | + | ++ | ++ | +++ | C |
| 232 | + | ++ | + | + | + | + | + | ++ | ++ | C |
| 233 | + | ++ | + | + | + | + | ++ | ++ | + | D |
| 234 | + | +++ | ++ | ++ | + | + | ++ | ++ | ++ | C |
| 235 | + | ++ | + | + | + | + | + | + | + | D |
| 236 | ++ | ++ | + | ++ | + | + | + | ++ | ++++ | B |
| 237 | + | ++ | +++ | ++ | + | + | + | + | ++ | C |
| 238 | + | ++ | + | ++ | + | + | ++ | + | ++++ | C |
| 239 | ++ | ++ | + | ++ | + | ++ | ++ | ++ | +++ | B |
| 240 | ++ | ++ | ++ | ++ | + | + | ++ | ++ | +++ | B |
| 248 | + | ++ | + | + | + | + | ++ | ++ | +++ | C |
| 249 | + | ++ | + | ++ | + | + | ++ | ++ | +++ | B |
| 250 | + | ++ | ++ | ++ | ++ | ++ | + | + | + | B |
| 251 | + | ++ | ++ | ++ | ++ | ++ | + | + | ++ | B |
| 252 | ++ | + | ++ | ++ | ++ | + | + | + | ++++ | A |
| 253 | ++ | + | ++ | ++ | ++ | + | + | + | ++ | B |
| 254 | + | + | ++ | + | + | + | + | + | ++ | B |
| 255 | + | + | ++ | + | + | + | + | + | ++ | C |
| 256 | + | ++ | ++ | +++ | ++ | ++ | + | + | ++ | B |

| Example | DP | AX | SD | PS | MS | AD | CE | AG | UN | Potency |
|---------|------|----|-----|------|------|----|----|----|----|---------|
| 257 | ++ | + | +++ | ++ | ++ | + | + | + | ++ | B |
| 258 | ++ | + | ++ | + | ++ | + | + | + | + | B |
| 259 | ++ | + | +++ | + | + | + | + | + | ++ | C |
| 260 | + | + | ++ | + | ++ | ++ | + | + | + | B |
| 261 | ++++ | + | ++ | + | + | ++ | + | + | ++ | C |
| 262 | ++ | + | ++ | + | + | + | + | + | ++ | C |
| 263 | + | + | ++ | + | + | + | + | + | + | A |
| 264 | ++ | ++ | ++ | +++ | ++ | ++ | + | + | + | B |
| 265 | ++ | + | ++ | + | + | + | + | + | + | B |
| 266 | + | + | ++ | + | ++ | + | + | + | + | B |
| 267 | ++ | + | ++ | + | ++ | + | + | + | ++ | A |
| 268 | + | ++ | ++ | +++ | ++ | ++ | + | + | ++ | B |
| 269 | ++ | + | ++ | + | ++ | + | + | + | ++ | B |
| 270 | ++ | ++ | ++ | + | ++++ | ++ | + | + | ++ | A |
| 271 | ++ | ++ | ++ | + | +++ | ++ | + | + | ++ | B |
| 272 | + | ++ | ++ | ++ | ++ | + | + | + | ++ | B |
| 273 | ++ | + | +++ | + | ++ | + | + | ++ | ++ | A |
| 274 | ++ | + | ++ | ++ | +++ | + | + | + | ++ | A |
| 275 | + | + | +++ | + | ++ | + | + | + | ++ | B |
| 276 | + | + | ++ | + | ++ | + | + | + | + | B |
| 277 | + | + | + | + | + | + | + | + | + | D |
| 278 | + | + | + | + | + | + | + | + | + | D |
| 279 | ++ | + | ++ | ++ | +++ | + | + | + | ++ | B |
| 280 | ++ | + | ++ | + | ++ | + | ++ | + | + | B |
| 281 | + | + | + | + | + | + | + | + | + | D |
| 282 | + | + | ++ | + | + | + | + | + | + | D |
| 283 | ++ | ++ | ++ | ++ | +++ | ++ | + | + | ++ | B |
| 284 | ++ | ++ | ++ | ++ | +++ | + | + | + | + | C |
| 285 | ++ | ++ | + | ++++ | ++++ | ++ | + | + | ++ | B |
| 286 | ++ | + | ++ | + | ++ | + | + | + | ++ | C |
| 287 | + | + | ++ | + | + | + | + | + | + | D |
| 288 | ++ | + | ++ | + | + | + | + | + | + | D |

Marble Burying Test

[0181] The marble burying assay is used to evaluate the anxiolytic activity of a test compound. After placement in a novel cage containing glass marbles, mice will bury the

marbles. Acute administration of anxiolytic agents, as well as antidepressants, decrease this activity at non-sedating doses. Male C57Bl/6J mice from Jackson Laboratories were used in this study and were acclimatized to the experimental room for at least 1 hour prior to testing. Mice (n=10 per group) were administered sterile water or test compound intraperitoneally (i.p.) or orally by gavage (p.o.), or the positive control chlordiazepoxide (15 mg/kg, i.p.), in 10 mL/kg injection volumes, and returned to their home cages for a 30-minute pretreatment period. The mice were then placed individually in clean cages, containing hard wood bedding and twenty black marbles placed in spaced rows of 4 x 5. After a 30-minute test session, the number of buried marbles was counted. A marble was considered buried if it was pushed at least two thirds into the bedding. Locomotor activity was monitored by measuring the distance traveled using an overhead camera and Video Tracker Software (ViewPoint Life Sciences Software, France). A reduction in the number of marbles buried relative to the vehicle condition indicates an anxiolytic drug-like effect.

Effects of Compounds on Marble Burying in Mice.

| Compound | Dose | Marbles Buried |
|------------|-----------------|----------------|
| 111 | 3 mg/kg (p.o.) | - |
| | 10 mg/kg (p.o.) | + |
| | 30 mg/kg (p.o.) | + |
| 123 | 3 mg/kg (p.o.) | - |
| | 10 mg/kg (p.o.) | + |
| | 30 mg/kg (p.o.) | + |
| 114 | 1 mg/kg (p.o.) | - |
| | 3 mg/kg (p.o.) | + |
| | 10 mg/kg (p.o.) | + |
| | 30 mg/kg (p.o.) | + |
| 117 | 1 mg/kg (p.o.) | - |
| | 3 mg/kg (p.o.) | + |
| | 10 mg/kg (p.o.) | + |
| | 30 mg/kg (p.o.) | + |
| 118 | 3 mg/kg (p.o.) | - |
| | 10 mg/kg (p.o.) | + |

| | | |
|------------|-----------------|---|
| | 30 mg/kg (p.o.) | + |
| 124 | 3 mg/kg (p.o.) | - |
| | 10 mg/kg (p.o.) | + |
| | 30 mg/kg (p.o.) | + |
| 152 | 1 mg/kg (i.p.) | - |
| | 3 mg/kg (i.p.) | - |
| | 10 mg/kg (i.p.) | + |
| 157 | 1 mg/kg (i.p.) | - |
| | 3 mg/kg (i.p.) | - |
| | 10 mg/kg (i.p.) | + |
| 160 | 1 mg/kg (i.p.) | - |
| | 3 mg/kg (i.p.) | - |
| | 10 mg/kg (i.p.) | + |
| 170 | 1 mg/kg (i.p.) | - |
| | 3 mg/kg (i.p.) | - |
| | 10 mg/kg (i.p.) | + |
| 195 | 1 mg/kg (i.p.) | - |
| | 3 mg/kg (i.p.) | - |
| | 10 mg/kg (i.p.) | + |

- : No reduction in number of marbles buried

+ : Significant reduction in number of marbles buried (P <0.05 vs vehicle)

Stress Induced Hyperthermia Assay

[0182] Stress-induced hyperthermia (SIH), mediated by the autonomic nervous system, is well known to occur prior to and during exposure to stress and/or anxiety-inducing situations. In many human anxiety disorders, it occurs as an integral part of the pathology and is often considered a representative symptom of the disease, e.g. in generalized anxiety disorder as defined in DSM-IV. In rodents, acute administration of anxiolytic drugs, such as buspirone or chlordiazepoxide (CDP), are known to reduce the SIH response to stressors such as handling, noise, heat, novelty, or pain. The test involves two measures of rectal temperature repeated in the same animal with a 10-minute interval. On the day prior to testing, male Sprague-Dawley rats from Envigo (n=10 per group) were brought to the experimental room approximately one hour before scheduled lights out and singly housed overnight with food

and water *ad libitum*. On the morning of the experiment, animals were first injected with either vehicle or test compound orally by gavage, or CDP (10 mg/kg, i.p.). One hour following injection, each animal was removed from the holding cage and held in a supine position while the rectal temperature was measured by slowly inserting a rectal probe into the animal's rectum at a length of approximately 0.5 cm. The rectal probe was attached to a PhysiTemp digital thermometer (Fisher Scientific) which provides temperature readings at 0.1°C accuracy. The probe remained inside the animal for approximately 5 seconds or until body temperature reaches stability. This temperature was recorded as the baseline rectal temperature (T1). The animal was immediately placed back to the holding cage and after a 10-min interval the second rectal temperature (T2) was taken using the same procedure as in measuring T1. Before each insertion, the rectal probe was cleaned with an alcohol pad and lubricated with sterile K-Y jelly. All SIH studies were conducted between 8:00 – 11:00 a.m. The SIH response was calculated as the difference in temperature between the second and first temperature reading (Delta T, ΔT). A reduction in ΔT relative to the vehicle condition indicates an anxiolytic drug-like response.

Effects of Compounds on Stress-induced Hyperthermia (SIH) in Rats.

| Compound | Dose (mg/kg, PO) | Delta T (ΔT) |
|------------|------------------|------------------------|
| 152 | 0.3 mg/kg | - |
| | 1 mg/kg | + |
| | 3 mg/kg | + |
| | 10 mg/kg | + |
| | 30 mg/kg | + |
| 157 | 1 mg/kg | - |
| | 3 mg/kg | + |
| | 10 mg/kg | + |
| | 30 mg/kg | + |
| 170 | 3 mg/kg | - |
| | 10 mg/kg | + |
| | 30 mg/kg | + |

- : No change in Delta T (ΔT)
- + : Significant decrease in Delta T (ΔT) (P <0.05 vs vehicle)

Amphetamine Induced Hyperlocomotion Assay

[0183] Psychostimulants such as amphetamine (AMPH) are frequently used to induce or mimic a manic-like or psychotic-like state. AMPH administration to rodents increases horizontal locomotion, vertical rearing and repetitive stereotyped behaviors, and typical antipsychotic drugs, such as haloperidol, can reduce these behaviors. The antimanic/antipsychotic-like effects of test compounds were evaluated in male C57Bl/6J mice from Jackson Laboratories Mice were acclimatized to the experimental room for at least 1 hour prior to testing. The mice (n=10 per group) were administered vehicle or test compound orally by gavage (10 mL/kg injection volume) and placed individually in the novel open field (OF) chambers (Med Associates) for 30 minutes of baseline activity measurement. Mice were then injected with either water or AMPH (4 mg/kg, i.p., 10 mL/kg) and placed back in the same individual OF chambers for a 60-minute testing session during which the effects of test compounds on stimulant-induced behaviors (total distance traveled, rearing and stereotypy) were measured. A reduction in total distance traveled relative to the vehicle + AMPH condition indicates an antimanic/antipsychotic drug-like response.

Effects of Compounds on AMPH-Induced Hyperlocomotion in Mice.

| Compound | Dose (PO, mg/kg) | Total Distance Traveled (cm) |
|------------|--------------------------|------------------------------|
| 111 | 3 mg/kg Compound + AMPH | # |
| | 10 mg/kg Compound + AMPH | - |
| | 30 mg/kg Compound + AMPH | + |
| 123 | 3 mg/kg Compound + AMPH | # |
| | 10 mg/kg Compound + AMPH | - |
| | 30 mg/kg Compound + AMPH | + |
| 114 | 3 mg/kg Compound + AMPH | - |
| | 10 mg/kg Compound + AMPH | - |
| | 30 mg/kg Compound + AMPH | + |
| 117 | 3 mg/kg Compound + AMPH | - |
| | 10 mg/kg Compound + AMPH | - |

| | | |
|------------|--------------------------|---|
| | 30 mg/kg Compound + AMPH | + |
| 118 | 3 mg/kg Compound + AMPH | - |
| | 10 mg/kg Compound + AMPH | - |
| | 30 mg/kg Compound + AMPH | + |
| 124 | 3 mg/kg Compound + AMPH | - |
| | 10 mg/kg Compound + AMPH | - |
| | 30 mg/kg Compound + AMPH | + |

: Significant increase in total distance traveled (P <0.05 vs vehicle + AMPH group)

- : No change in total distance traveled

+ : Significant decrease in total distance traveled (P <0.05 vs vehicle + AMPH group)

Tail Suspension Test

[0184] The tail suspension test (TST) is a rodent screening test for potential (human) antidepressant drugs. It is based on the assumption that an animal will actively try to escape an aversive (stressful) stimulus. If escape is impossible, the animal will eventually stop trying ("give up"). In the TST, a mouse is suspended by the tail so that its body dangles in the air, head downward. Mice initially struggle to face upward and climb to a solid surface. When the animal stops struggling and hangs immobile it is considered to have "given up". Shorter periods of immobility are characteristic of antidepressant-like activity. Accordingly, longer periods of immobility are considered indicative of a depressive-like state. It has been shown that treatment with an antidepressant drug will decrease the time the animal spends immobile. *See generally L. Steru et al., Psychopharmacology (Berl). 1985;85(3):367-70; B. Thierry et al., Psychopharmacology 1986;90:284-85.*

[0185] *Procedure.* Adult male AJ mice from Jackson Laboratories received vehicle (sterile water) or test compound orally by gavage, or the positive control desipramine (20 mg/kg, i.p.), in 10 mL/kg injection volumes, 30 min before being subjected to the Tail Suspension Test. In this test, mice are placed in the Tail Suspension chambers (white polyvinylchloride cubicles measuring 33 x 33 x 31.75 cm *Med Associates, Inc.* St. Albans, VT) by a piece of transparent (Scotch[®]) tape attached to the tail, from about the mid-tail, with approximately 2 cm of tape past the end of the tail for 10 min during which the time spent immobile is

measured. A reduction in total time immobile relative to the vehicle condition indicates an antidepressant drug-like response.

Effects of Compounds in a Tail-Suspension Test (TST) in Mice.

| Compound | Dose (PO, mg/kg) | Mean Total Time Immobile (sec) |
|------------|------------------|--------------------------------|
| 111 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 123 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 114 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | + |
| 117 | 3 mg/kg | - |
| | 10 mg/kg | + |
| | 30 mg/kg | - |
| 118 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 124 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |

- : No decrease in Mean Total Time Immobile

+ : Significant decrease in Mean Total Time Immobile (P <0.05 vs vehicle)

Prepulse Inhibition of Startle

[0186] The acoustic startle is an unconditioned reflex response to an external auditory stimulus. Prepulse inhibition of startle (PPI) refers to the reduction in the startle response caused by the presentation of a low-intensity auditory stimulus prior to the startle stimulus. The PPI paradigm is used for the study of schizophrenia and antipsychotic action due to the similarities between the results from human and rodent studies. PPI has been used as a tool for the assessment of deficiencies in sensory-motor gating observed in schizophrenia and to

screen for potential antipsychotic drugs. Various psychotomimetic drugs, such as phencyclidine (PCP), can disrupt PPI. In mice, antipsychotic drugs such as haloperidol can increase PPI and clozapine can reverse the disruption of PPI induced by PCP.

[0187] Male C57Bl/6J mice from Jackson Laboratories were placed in the PPI chambers (Med Associates) for a 5 min session of white noise (70 dB) habituation. After the habituation period, the test session was initiated. The session started with a habituation block of 6 presentations of the startle stimulus alone, followed by 10 PPI blocks, each of which consisted of 6 different types of trials. Trial types were: 'null' (no stimuli), 'startle' (120dB), 'startle plus prepulse' (4, 8, and 12 dB over background noise, i.e., 74, 78, or 82 dB) and 'prepulse' (82 dB). Trial types were presented in a random order within each block. Each trial started with a 50 ms stimulus-free period during which baseline movements were recorded. This was followed by a subsequent 20 ms period during which the prepulse stimulus was presented and responses to the prepulse measured. After a further 100 ms period, the startle stimulus was presented for 40 ms and responses recorded for 100 ms from startle onset. Responses were sampled every ms. The inter-trial interval was variable with an average of 15 s (range from 10 to 20 s). In 'startle' trials, the basic auditory startle response was measured. The basic startle response was calculated as the mean startle response of all 'startle' trials (i.e., excluding the first habituation block). In 'startle plus prepulse' trials, the degree of inhibition of the normal startle was calculated and expressed as a percentage of the basic startle response.

[0188] Mice were treated with vehicle or test compound orally by gavage, or haloperidol (1 mg/kg, i.p.), in 10 mL/kg injection volumes, 30 min prior to PPI test. The PPI enclosures were cleaned following each test. An increase in the percent PPI relative to the vehicle condition indicates an antipsychotic drug-like response.

Effects of Compounds on Pre-pulse Inhibition (PPI) in Mice.

| Compound | Dose (PO, mg/kg) | Effect |
|------------|------------------|--------|
| 111 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | +++ |

| | | |
|------------|----------|-----|
| 123 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 114 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 117 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | +++ |
| 118 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |
| 124 | 3 mg/kg | - |
| | 10 mg/kg | - |
| | 30 mg/kg | - |

- : No change in PPI

+ : Significant increase in PPI at one pre-pulse intensity (P <0.05 vs vehicle)

++ : Significant increase in PPI at two pre-pulse intensities (P <0.05 vs vehicle)

+++ : Significant increase in PPI at three pre-pulse intensities (P <0.05 vs vehicle)

In Vivo Microdialysis of the Prefrontal Cortex and Nucleus Accumbens

[0189] In vivo microdialysis is a biological sampling technique used to evaluate the extracellular concentrations of neurotransmitters in specific brain regions following administration of a test compound to animals. Dysfunctions of brain neurotransmitter systems, such as dopamine (DA), norepinephrine (NE), serotonin (5-HT) and/or acetylcholine (ACh), have been linked to cognitive deficits (e.g., memory impairment, inattention, etc.), which may be observed in human aging, Alzheimer's disease (AD) and Attention Deficit Hyperactivity Disorder (ADHD). Drugs that increase extracellular concentrations of DA, NE, 5-HT and/or ACh in the prefrontal cortex (PFC), such as donepezil and methylphenidate, can reverse cognitive deficits in humans and animals.

[0190] Adult male C57Bl/6 mice (n = 6 per group) from Jackson Laboratories were anesthetized using isoflurane (2%, 800 mL/min O₂). Bupivacaine/epinephrine was used for local anesthesia and carprofen was used for peri-/post-operative analgesia. The animals were placed in a stereotaxic frame (Kopf instruments, USA) and I-shaped microdialysis probes (polyacrylonitril membrane, BrainLink, the Netherlands) were inserted into the PFC (2 mm exposed membrane; coordinates for the tips of the probes: posterior (AP) = +2.0 mm to bregma, lateral (L) = -0.5 mm to midline and ventral (V) = -3.3 mm to dura, the toothbar set at 0.0 mm) and the NAcc (coordinates for the tips of the probes: posterior (AP) = +0.8 mm to bregma, lateral (L) = +0.9 mm to midline and ventral (V) = -4.2 mm to dura, the toothbar set at 0.0 mm (Paxinos and Franklin, 2001)). After surgery, animals were housed individually in cages and provided food and water *ad libitum*. In vivo microdialysis experiments were performed one day after surgery. On the day of the experiment, the microdialysis probes were connected with flexible PEEK tubing to a microperfusion pump (Harvard PHD 2000 Syringe pump, Holliston, MA or similar). Microdialysis probes were perfused with artificial cerebrospinal fluid (aCSF) containing 147 mM NaCl, 3.0 mM KCl, 1.2 mM CaCl₂ and 1.2 mM MgCl₂, at a flow rate of 1.5 µL/min. Microdialysis samples were collected for 30 minute periods by an automated fraction collector (820 Microsampler, Univentor, Malta or similar) into polystyrene mini-vials already containing 15 µL of 0.02 M formic acid (FA) and 0.04% ascorbic acid in ultrapurified H₂O. Four basal samples were collected before the administration of vehicle or test compound (i.p., 5 mL/kg injection volumes). Samples were collected for an additional 240 min following vehicle or compound administration. All the dialysis samples were stored at -80 °C awaiting analysis. After the experiment, the mice were sacrificed and brain tissue was collected for probe verification. Each dialysis sample was analyzed for concentrations of DA, NE, 5-HT and ACh as determined by HPLC with tandem mass spectrometry (MS/MS) detection using labeled isotopes of the analytes of interest as internal standards. Data was expressed as a percentage of basal output, calculated by dividing each post-dose time point with the average basal output = 100%. An increase in the post-dose percent basal output of DA, NE, 5-HT and/or ACh in the PFC relative to the vehicle condition indicates a potential cognition-enhancing drug-like response. Graphical representations of the observed changes in neurotransmitter concentrations are shown in the indicated Figures.

Effects of Compounds on Neurotransmitters in the PFC and NAcc in Mice.

| Compound | Dose (mg/kg, IP) | PFC [DA] (Figure 1) | PFC [5-HT] (Figure 2) | PFC [NE] (Figure 3) | PFC [ACh] (Figure 4) |
|------------|------------------|---------------------|-----------------------|---------------------|----------------------|
| 152 | 10 mg/kg | + | + | + | + |
| 157 | 30 mg/kg | + | + | + | - |
| 170 | 30 mg/kg | + | - | + | - |

- : No change in extracellular concentration (P <0.05 vs vehicle)

+: Significant increase in extracellular concentration (P <0.05 vs vehicle)

| Compound | Dose (mg/kg, IP) | NAcc [DA] (Figure 5) | NAcc [5-HT] (Figure 6) | NAcc [NE] (Figure 7) | NAcc [ACh] (Figure 8) |
|------------|------------------|----------------------|------------------------|----------------------|-----------------------|
| 152 | 10 mg/kg | - | + | + | - |
| 157 | 30 mg/kg | - | - | + | - |
| 170 | 30 mg/kg | - | - | + | - |

- : No change in extracellular concentration (P <0.05 vs vehicle)

+: Significant increase in extracellular concentration (P <0.05 vs vehicle)

List of Abbreviations

| | |
|-----------|--|
| 5-HT | 5-hydroxytryptophan (serotonin) |
| ACh | acetylcholine |
| ACN | acetonitrile |
| aCSF | artificial cerebrospinal fluid |
| AMPH | Amphetamine |
| AP | anterior-posterior |
| °C | degrees Celsius |
| CDP | chlordiazepoxide |
| cm | centimeter |
| DA | dopamine |
| FA | formic acid |
| g | gram |
| HPLC | high performance liquid chromatography |
| IP | intraperitoneal |
| L | lateral |
| LC | liquid chromatography |
| µL | microliter |
| MD | microdialysis |
| mg/kg | milligram per kilogram |
| mL | milliliter |
| mm | millimeter |
| mM | millimolar |
| min / Min | minute |
| MRM | multiple reaction monitoring |
| MS | mass spectrometry |


| | |
|------|-----------------------------|
| MSq | mean squared |
| NAcc | nucleus accumbens |
| NE | norepinephrine |
| nM | nanomolar |
| P | p-value |
| PCP | Phencyclidine |
| PPI | Pre-pulse Inhibition |
| % | percentage |
| PFC | prefrontal cortex |
| pmol | picomole |
| PSIG | pounds per square inch gage |
| QC | quality control |
| sec | seconds |
| SEM | standard error of the mean |
| SIH | stress-induced hyperthermia |
| TST | tail suspension test |
| V | ventral |
| UP | ultra-pure |

[0191] It may be found upon examination that additional species and genera not presently excluded from the claims to pharmaceutical compositions and chemical compounds are not patentable to the inventors in this application. In that case, the subsequent exclusion of species and genera in applicants' claims are to be considered artifacts of patent prosecution and not reflective of the inventors' concept or description of their invention. The invention, in a composition aspect, is all compounds of formula I except those that are in the public's possession.

[0192] It is to be understood that the invention is not limited to the particular embodiments of the invention described above, as variations of the particular embodiments may be made and still fall within the scope of the appended claims.

CLAIMS

1. A method for treating a neurological or psychiatric disease or disorder, or neuropsychiatric or behavior symptoms in a neurological disease or disorder, in a subject in need thereof, comprising
administering to said subject an effective amount of a compound of formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

q1 is 0 or 1,

q2 is 0 or 1, and

the sum of q1 plus q2 is 0 or 1;
 R^1, R^2 and R^3 are chosen independently from H and an aliphatic (C_1-C_8)hydrocarbyl optionally substituted with one or more groups independently selected from halogen, $-C\equiv C-$, hydroxyl, (C_1-C_8)alkoxy, amino, (C_1-C_8)alkylamino and di(C_1-C_8)alkylamino;

R^{4a} , R^{4b} , R^{7a} , and R^{7b} are chosen independently from H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, and (C₁-C₆)alkoxy;

R^{5a} , R^{5b} , R^{6a} and R^{6b} are chosen independently from H, halogen, (C_1-C_6) alkyl and (C_1-C_6) haloalkyl;

R^8 , R^9 , R^{10} and R^{11} are chosen independently from H, halogen, (C₁-C₈)hydrocarbyl, cyano, -CF₃, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy, hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, aminosulfonyl, benzyl, aryl, heteroaryl, phenoxy,

-OC(O)N(alkyl)₂ or benzyloxy;

wherein said benzyl, aryl, heteroaryl, phenoxy or benzyloxy are optionally substituted with one or more substituents independently selected from halogen, (C₁-C₆)alkyl, cyano, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy, hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, and di(C₁-C₆)alkylamino;

and further wherein:

any two of R⁸, R⁹, R¹⁰ and R¹¹ attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy.

2. The method according to claim 1, wherein the neurological or psychiatric disease or disorder is depression, bipolar disorder, pain, schizophrenia, obsessive compulsive disorder, addiction, social disorder, attention deficit hyperactivity disorder, an anxiety disorder, a movement disorder, epilepsy, autism or cognitive impairments.

3. The method according to claim 2, wherein the neurological or psychiatric disease or disease or disorder is depression.

4. The method according to claim 3, wherein the depression is treatment-resistant depression (TRD), major depressive disorder (MDD), unipolar depression, bipolar depression or depression associated with another disease or disorder.

5. The method according to claim 2, wherein the neurological or psychiatric disease or disorder is a movement disorder or epilepsy.

6. The method according to claim 1, wherein said neurological disease or disorder is selected from Alzheimer's disease and Parkinson's disease.

7. The method according to claim 1, wherein any two of R⁸, R⁹, R¹⁰ and R¹¹ attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more

substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy.

8. The method according to claim 7, wherein any two of R⁸, R⁹, R¹⁰ and R¹¹ form a phenyl ring.

9. The method according to claim 1, wherein:

(a) at least one of R⁸, R⁹, and R¹¹ is chosen from halogen, methyl and ethyl; or

(b) R¹⁰ is methoxy; or

(c) R⁸, R⁹, R¹⁰, R¹¹, R^{7a}, and R^{7b}, are hydrogen and at least one of R^{4a} and R^{4b} is methyl, or

(d) R⁸ and R⁹, together with the carbon atoms to which they are attached, form a 6-membered carbocycle or 7-membered bridged carbocycle.

10. The method of claim 9, wherein q1 and q2 are both zero, R^{4a} is hydrogen and R^{4b} is hydrogen or methyl.

11. The method of claim 10, wherein R² and R³ are hydrogen.

12. The method of claim 9, wherein q1 is one and R² and R³ are hydrogen or methyl.

13. The method according to claim 2, wherein the neurological or psychiatric disorder is pain.

14. The method of claim 13, wherein R² and R³ are hydrogen, q1 and q2 are both zero, and

(a) both of R^{7a} and R^{7b} are methyl, or

(b) both of R⁸ and R⁹ are methyl, or

(b) R⁹ is methoxy and at least one of R^{4a} and R^{4b} is methyl, or

(c) R¹¹ is propyl.

15. The method of claim 13, wherein R² is methyl, q1 is one, and R⁹ is methyl or chloro.

16. The method according to claim 2, wherein the neurological or psychiatric disorder is schizophrenia.

17. The method of claim 16, wherein R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, and R^{7b} are all hydrogen, and

- (a) q1 is one and R¹, R², R³, R⁸, R¹⁰, and R¹¹ are hydrogen, or
- (b) q1 and q2 are both zero, R¹⁰ and R¹¹ are hydrogen, or
- (c) q1 and q2 are both zero, R⁸, R⁹, and R¹⁰ are hydrogen and R¹¹ is (C₁-C₆)alkyl.

18. The method according to claim 2, wherein the neurological or psychiatric disorder is anxiety disorder.


19. The method according to claim 2, wherein the neurological or psychiatric disorder is loss of cognitive function.

20. The method of claim 19, wherein R¹, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, and R^{7b} are all hydrogen, and

- (a) q1 is one, R² and R³ are hydrogen or methyl, and R¹⁰ and R¹¹ are hydrogen or methyl, or
- (b) q1 and q2 are both zero and R⁸, R⁹, R¹⁰ and R¹¹ are chosen from hydrogen, methyl, methoxy and chloro.

21. The method according to claim 2, wherein the neurological or psychiatric disorder is autism.

22. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a compound of formula I:



I

or a pharmaceutically acceptable salt thereof,

wherein:

q1 is 0 or 1,

q2 is 0 or 1, and

the sum of q1 plus q2 is 0 or 1;

R¹, R² and R³ are chosen independently from H and an aliphatic (C₁-C₈)hydrocarbyl optionally substituted with one or more groups independently selected from halogen, -C≡C-, hydroxyl, (C₁-C₆)alkoxy, amino, (C₁-C₆)alkylamino and di(C₁-C₆)alkylamino;

R^{4a}, R^{4b}, R^{7a}, and R^{7b} are chosen independently from H, halogen, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, and (C₁-C₆)alkoxy;

R^{5a}, R^{5b}, R^{6a} and R^{6b} are chosen independently from H, halogen, (C₁-C₆)alkyl and (C₁-C₆)haloalkyl;

R⁸, R⁹, R¹⁰ and R¹¹ are chosen independently from H, halogen, (C₁-C₈)hydrocarbyl, cyano, -CF₃, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy, hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, aminosulfonyl, benzyl, aryl, heteroaryl, phenoxy, -OC(O)N(alkyl)₂ or benzyloxy;


wherein said benzyl, aryl, heteroaryl, phenoxy or benzyloxy are optionally substituted with one or more substituents independently selected from halogen, (C₁-C₆)alkyl, cyano, (C₁-C₆)haloalkyl, hydroxyl, (C₁-C₆)alkoxy, aminocarbonyl, (C₁-C₆)alkylaminocarbonyl, di(C₁-C₆)alkylaminocarbonyl, (C₁-C₆)acyl, (C₁-C₆)haloalkoxy, hydroxy(C₁-C₆)alkyl, carboxy, (C₁-C₆)alkoxycarbonyl, acetoxy, nitro, amino, (C₁-C₆)alkylamino, and di(C₁-C₆)alkylamino;

and further wherein:

any two of R⁸, R⁹, R¹⁰ and R¹¹ attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, (C₁-C₆)alkyl, (C₁-C₆)haloalkyl, (C₁-C₆)alkoxy, and (C₁-C₆)haloalkoxy;

wherein at least one of R¹, R², R³, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, R^{7b}, R⁸, R⁹, R¹⁰ and R¹¹ is other than hydrogen.

23. A compound of formula I:



or a pharmaceutically acceptable salt thereof,

wherein:

q1 is 0 or 1,

q2 is 0 or 1, and

the sum of q1 plus q2 is 0 or 1;

R^1 , R^2 and R^3 are chosen independently from H and an aliphatic (C_1 - C_8)hydrocarbyl optionally substituted with one or more groups independently selected from halogen, $-C\equiv C-$, hydroxyl, (C_1 - C_6)alkoxy, amino, (C_1 - C_6)alkylamino and di(C_1 - C_6)alkylamino;

R^{4a} , R^{4b} , R^{7a} , and R^{7b} are chosen independently from H, halogen, (C_1 - C_6)alkyl, (C_1 - C_6)haloalkyl, and (C_1 - C_6)alkoxy;

R^{5a} , R^{5b} , R^{6a} and R^{6b} are chosen independently from H, halogen, (C_1 - C_6)alkyl and (C_1 - C_6)haloalkyl;

R^8 , R^9 , R^{10} and R^{11} are chosen independently from H, halogen, (C_1 - C_8)hydrocarbyl, cyano, $-CF_3$, (C_1 - C_6)haloalkyl, hydroxyl, (C_1 - C_6)alkoxy, aminocarbonyl, (C_1 - C_6)alkylaminocarbonyl, di(C_1 - C_6)alkylaminocarbonyl, (C_1 - C_6)acyl, (C_1 - C_6)haloalkoxy, hydroxy(C_1 - C_6)alkyl, carboxy, (C_1 - C_6)alkoxycarbonyl, acetoxy, nitro, amino, (C_1 - C_6)alkylamino, di(C_1 - C_6)alkylamino, aminosulfonyl, benzyl, aryl, heteroaryl, phenoxy, $-OC(O)N(alkyl)_2$ or benzyloxy;

wherein said benzyl, aryl, heteroaryl, phenoxy or benzyloxy are optionally substituted with one or more substituents independently selected from halogen, (C_1 - C_6)alkyl, cyano, (C_1 - C_6)haloalkyl, hydroxyl, (C_1 - C_6)alkoxy, aminocarbonyl, (C_1 - C_6)alkylaminocarbonyl, di(C_1 -

C_6)alkylaminocarbonyl, (C_1-C_6) acyl, (C_1-C_6) haloalkoxy, hydroxy (C_1-C_6) alkyl, carboxy, (C_1-C_6) alkoxycarbonyl, acetoxy, nitro, amino, (C_1-C_6) alkylamino, and di (C_1-C_6) alkylamino;

and further wherein:

any two of R^8 , R^9 , R^{10} and R^{11} attached to adjacent carbon atoms, together with the carbon atoms to which they are attached, may form a 5 to 8-membered carbocycle or heterocycle which is optionally substituted with one or more substituents independently selected from halogen, hydroxyl, amino, (C_1-C_6) alkylamino, di (C_1-C_6) alkylamino, (C_1-C_6) alkyl, (C_1-C_6) haloalkyl, (C_1-C_6) alkoxy, and (C_1-C_6) haloalkoxy;

with the provisos that,


(1) when q_1 and q_2 are both zero and NR^2R^3 is $NHCH_3$ or $N(CH_3)_2$, then at least one of R^1 , R^{4a} , R^{4b} , R^8 , R^9 , R^{10} and R^{11} must be other than hydrogen or methoxy; or

(2) when q_1 and q_2 are both zero and NR^2R^3 is NH_2 , then one of the following four conditions must be met:


- (a) at least two of R^8 , R^9 , R^{10} and R^{11} must be other than hydrogen, or
- (b) at least one of R^8 , R^9 , R^{10} and R^{11} must be phenyl, or
- (c) at least one of R^{4a} and R^{4b} must be other than hydrogen, or
- (d) at least one of R^{7a} and R^{7b} must be methyl and the other of R^{7a} and R^{7b}

must be other than methyl, and

(3) when q_1 or q_2 is one, then the following species are excluded:




24. The compound according to claim 23, or a pharmaceutically acceptable salt thereof, wherein q_1 and q_2 are both zero, said compound having the formula II:




II.

25. The compound according to claim 23, or a pharmaceutically acceptable salt thereof, wherein q1 is one, said compound having the formula III:



III.

26. The compound according to claim 23, or a pharmaceutically acceptable salt thereof, wherein q2 is one, said compound having the formula IV:



IV.

27. The compound according to claim 23, wherein R^{4a}, R^{4b}, R^{7a}, and R^{7b} are hydrogen.

28. The compound according to claim 23, wherein R^{4a}, R^{4b}, R^{7a}, and R^{7b} are hydrogen.

29. The compound according to claim 23, wherein R⁸ and R⁹ form a 5 to 8-membered carbocycle or heterocycle which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy.

30. The compound according to claim 29, wherein said carbocycle or heterocycle is chosen from phenyl, pyridine, cyclohexene, thiazole, bicycloheptene, and dihydropyran.

31. The compound according to claim 23, wherein R¹⁰ and R¹¹ form a 5 to 8-membered carbocycle or heterocycle which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy.

32. The compound according to claim 31, wherein said carbocycle or heterocycle is chosen from phenyl, pyridine, cyclohexene, thiazole, bicycloheptene, and dihydropyran.

33. The compound according to claim 23, wherein R⁹ is

(a) phenyl which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy; or

(b) 5- or 6-membered heteroaryl which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy.

34. The compound according to claim 23, wherein two of R⁸, R⁹, R¹⁰ and R¹¹ are chosen from methyl and chloro and the other two are hydrogen.

35. The compound according to claim 23, wherein R² is hydrogen and R³ is chosen from methyl, ethyl, propyl, isopropyl, and hydroxyethyl.

36. The compound according to claim 23, wherein R¹ is hydrogen or methyl.

37. The compound according to claim 36, wherein R¹ is hydrogen.

38. The compound according to claim 23, wherein R^{4a} and R^{4b} are chosen independently from hydrogen and (C₁-C₆)alkyl.

39. The compound according to claim 23, wherein R⁸ and R⁹ form a 5 to 8-membered carbocycle or heterocycle which is unsubstituted or substituted with 1 to 4 substituents independently selected from halo, -OH, -NH₂, methyl, ethyl, methoxy, trifluoromethyl and trifluoromethoxy.


40. The compound according to claim 39, wherein said carbocycle is phenyl.

































41. The compound according to claim 23, wherein R^{4a} and R^{4b} are chosen from hydrogen and methyl.







42. The compound according to claim 23, wherein R² is hydrogen and R³ is hydrogen or methyl.








43. The compound according to claim 23, wherein one or two of R⁸, R⁹, R¹⁰ and R¹¹ are chosen from methyl and chloro and the remaining two or three of R⁸, R⁹, R¹⁰ and R¹¹ are hydrogen.


44. The compound according to claim 23, wherein said compound is:





















or a pharmaceutically acceptable salt thereof.

45. The compound according to claim 23, wherein said compound is:



or a pharmaceutically acceptable salt thereof.

46. The compound according to claim 23, wherein said compound is:



pharmaceutically acceptable salt thereof.

47. The method according to claim 2, wherein said neurological or psychiatric disease or disorder is anxiety.

48. The method according to claim 2, wherein said neurological or psychiatric disease or disorder is attention deficit hyperactivity disorder.

49. A method of treating anxiety or attention deficit hyperactivity disorder in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound according to claim 45, or a pharmaceutically acceptable salt thereof.

50. A method of treating anxiety in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound according to claim 45, or a pharmaceutically acceptable salt thereof.

51. A method of treating attention deficit hyperactivity disorder in a subject in need thereof, comprising the step of administering to said subject an effective amount of a compound according to claim 45, or a pharmaceutically acceptable salt thereof.

52. The compound according to claim 23, wherein said compound is greater than 90% enantiomerically pure.

53. The compound according to claim 23, wherein said compound is greater than 95% enantiomerically pure.

54. The compound according to claim 46, wherein said compound is greater than 90% enantiomerically pure.

55. The compound according to claim 46, wherein said compound is greater than 95% enantiomerically pure.

Figure 1

Effect of compound treatment on DA release in the PFC.

(Data expressed as DA concentration as % of baseline \pm SEM).




Figure 2

Effect of compound treatment on 5-HT release in the PFC.

(Data expressed as 5-HT concentration as % of baseline \pm SEM).




Figure 3

Effect of compound treatment on NE release in the PFC.

(Data expressed as NE concentration as % of baseline \pm SEM).




Figure 4

Effect of compound treatment on ACh release in the PFC
(Data expressed as Ach concentration as % of baseline \pm SEM).




Figure 5

Effect of compound treatment on DA release in the NAcc.

(Data expressed as DA concentration as % of baseline \pm SEM).




Figure 6

Effect of compound treatment on 5-HT release in the NAcc.
(Data expressed as 5-HT concentration as % of baseline \pm SEM).




Figure 7

Effect of compound treatment on NE release in the NAcc.

(Data expressed as NE concentration as % of baseline \pm SEM).




Figure 8

Effect of compound treatment on ACh release in the NAcc.

(Data expressed as Ach concentration as % of baseline \pm SEM).

