



(11) **EP 2 990 044 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
02.03.2016 Bulletin 2016/09

(51) Int Cl.:
A61K 33/40 (2006.01) A61K 33/14 (2006.01)
A61K 33/22 (2006.01) A61P 17/00 (2006.01)
A01N 39/00 (2006.01)

(21) Application number: **15186300.8**

(22) Date of filing: **04.12.2007**

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR

(72) Inventor: **KARAGEOZIAN, Hampar L.**
San Juan Capistrano, CA California 92675 (US)

(30) Priority: **04.12.2006 US 633355**

(74) Representative: **Instone, Alicia Claire**
Scott & York
Intellectual Property Limited
45 Grosvenor Road
St Albans, Hertfordshire AL1 3AW (GB)

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
07862490.5 / 2 099 296

Remarks:
This application was filed on 22-09-2015 as a divisional application to the application mentioned under INID code 62.

(71) Applicant: **S.K. Pharmaceuticals, Inc.**
San Juan Capistrano, CA 92675 (US)

(54) **SYNERGISTIC ANTIMICROBIAL PREPARATIONS CONTAINING CHLORITE AND HYDROGEN PEROXIDE**

(57) An anti-microbial preservative for use in ophthalmic and dermatologic products. The preservative includes from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound. Additionally, the preservative does not generate chlorine dioxide within the pH range of 5.0 - 8.8. Also included are an antimicrobial ophthalmic and dermatologic compositions for direct application onto an

eye or skin of a living being including from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound. Also included are methods for treating dryness of the eyes and skin disorders (e.g., wounds, burns, infections, ulcerations, psoriasis, etc.) and for disinfecting and cleansing contact lenses while in place upon an eye by applying the composition to the eye or to the contact lens.




Figure 1

EP 2 990 044 A1

Description

CROSS-REFERENCE TO RELATED APPLICATIONS

5 **[0001]** The present invention is a continuation-in-part of United States Patent Application Serial No. 10/614,646 filed on July 7, 2003, which is a continuation-in-part of United States Patent Application Serial No. 09/911,638 filed July 23, 2001, which is a continuation-in-part of United States Patent Application Serial No. 09/412,174 filed October 4, 1999, the entirety of the disclosures of which are expressly incorporated herein by reference.

10 STATEMENT RE: FEDERALLY SPONSORED RESEARCH/DEVELOPMENT

[0002] Not Applicable

BACKGROUND

15

1. Field of the Invention

[0003] The present invention relates generally to medical compositions and methods, and more particularly to certain disinfectant/antimicrobial preparations and methods for using such preparations i) to disinfect or preserve articles or surfaces, ii) as a topical antiseptic for application to body parts, iii) to prevent or deter scar formation; iv) to treat dermatological disorders such as wounds, burns, ulcers, psoriasis, acne and other scar forming lesions; and v) to treat ophthalmic disorders such as infections, inflammation, dry eye, wound healing, and allergic conjunctivitis.

20

2. Background of the Invention

25

A. Antimicrobial and Disinfectant/Antiseptic Agents Used for Disinfection/Antisepsis and Topical Treatment of Wounds, Burns, Abrasions and Infections

[0004] The prior art has included numerous antimicrobial agents which have purportedly been useable for disinfection of various articles and/or for topical application to a living being for antisepsis and/or treatment of dermal disorders (e.g., wounds, burns, abrasions, infections) wherein it is desirable to prevent or deter microbial growth to aid in healing. Such topical antimicrobial agents have contained a variety of active microbicidal ingredients such as iodine, mercurochrome, hydrogen peroxide, and chlorine dioxide.

30

i. Prior Chlorine Dioxide Preparations

[0005] Chlorite, a precursor of chlorine dioxide, is known to be useable as a disinfectant for drinking water and as a preservative for contact lens care solutions. However, chlorite exhibits only weak microbicidal activity within a concentration range that is acceptable and safe for topical application to the skin (e.g., 50-1000 parts per million). Thus, chlorite has not been routinely used as an active microbicidal ingredient in preparations for topical application to the skin.

40

[0006] In view of the limited usefulness of chlorite as an antiseptic or topical microbicide, various compositions and methods have been proposed for activation or enhancement of the microbicidal activity of chlorite. Examples of such compositions and methods for activation or enhancement of the microbicidal activity of chlorite are described in United States Patent Nos. 4,997,616 (describing general activation); 5,279,673 (describing acid activation) and 5,246,662 (describing transition metal activation).

45

[0007] Chlorine dioxide (ClO₂) and "stabilized chlorine dioxide" are known to be useable as antiseptics. Chemically, chlorine dioxide is an oxidizing agent which has strong microbicidal activity. Chlorine dioxide is generally regarded as superior even to gaseous chlorine in certain water treatment applications where it is used as to eliminate algae and other organic material and/or to remove odors or tastes. Chlorine dioxide is also effective as a microbicide, for elimination of bacteria, viruses, and microbial spores.

50

[0008] In addition to its use as a microbicide, chlorine dioxide is a highly reactive, unstable radical which is useable as an oxidizing agent in a number of other chemical and biochemical applications. For example, as described in United States Patent No. 4,855,135, chlorine dioxide can be used for (a) oxidation of double bonds between two carbon atoms; (b) oxidation of unsaturated fatty acids (lipids) via double bonds between two carbon atoms; (c) acceleration of hydrolysis of carboxylic anhydrides; (d) oxidation of aldehydes to the corresponding carboxylic acids; (e) oxidation of alcohols; (f) oxidation of amines; (g) oxidation of phenols, phenolic derivatives and thiophenolic compounds; (h) moderate oxidation of hydroquinones; (i) oxidation of amino acids, proteins and polyamides; j) oxidation of nitrates and sulfides; and (k) alteration of the CHO and CH₂OH radicals of carbohydrates to produce carboxylic functionality.

55

[0009] Concentrated chlorine dioxide in its liquid or gaseous state is highly explosive and poisonous. As a result, concentrated chlorine dioxide must be handled and transported with great caution. For this reason, it is generally not feasible to dispense pure chlorine dioxide for use as a topical antimicrobial agent or disinfectant. Instead, some antimicrobial or disinfectant preparations have been formulated to provide for "acid generation" of chlorine dioxide. Such acid generation solutions contain a metal chlorite (i.e., a precursor of chlorine dioxide available in powdered or liquid form) in combination with an acid which will react with the chlorite to liberate or release chlorine dioxide. Generally, any acid may be used for acid generation of chlorine dioxide, including strong acids such as hydrochloric acid and sulfuric acid and relatively weak acids such as citric and tartaric acid. Drawbacks or problems associated with these prior chlorine dioxide generating systems include a) the inconvenience of handling two separate containers or chemical components, b) the difficulty of delivering such two-component systems to the intended site of application, and c) the fact that these prior systems are of acid, rather than neutral, pH. Moreover, the prior chlorine dioxide generating systems which utilize acid induced generation of chlorine dioxide can, if uncontrolled, cause the generation of chlorine dioxide to occur quite rapidly and, as a result, the disinfectant or antimicrobial potency of the solution may be short lived. Increasing the concentration of chlorite and acid within the solution may prolong its disinfectant or antimicrobial shelf life, but such increased concentrations of these chemicals can result in toxicities or (in topical applications) skin irritation. Such increased concentrations may also result in the generation of more chlorine dioxide than is required.

[0010] Various methods have been described to limit or control the rate at which chlorine dioxide is produced in "acid generation" solutions. For instance, United States Patent No. Re. 31,779 (Alliger), which is a reissue of United States Patent No. 4,084,747, describes a germicidal composition which comprises a water soluble chlorite, such as sodium chlorite, in combination with lactic acid. The particular composition possesses improved disinfectant properties, properties not attained by using the same composition but replacing the lactic acid with other acids such as phosphoric acid, acetic acid, sorbic acid, fumaric acid, sulfamic acid, succinic acid, boric acid, tannic acid, and citric acid. The germ killing composition is produced by contacting an acid material containing at least 15% by weight of lactic acid with sodium chlorite in aqueous media. The methods disclosed of disinfecting and sanitizing a germ carrying substrate, such as skin, include either application of the germ killing composition, or application of the reactants to provide *in situ* production thereof. Also, United States Patent No. 5,384,134 (Kross) describes acid induced generation of chlorine dioxide from a metal chlorite wherein the chlorite concentration is limited by the amount of available chlorous acid. In particular, the Kross patent describes a method for treating dermal disorders wherein a first gel, which comprises a metal chlorite, is mixed with a second gel, which comprises a protic acid. The chlorite ions present in such solution as chlorous acid purportedly comprise no more than about 15% by weight of the total chlorite ion concentration in the composition, and the mixture of the two gels purportedly generates chlorine dioxide over an extended time of up to 24 hours.

[0011] Other prior patents have purported to describe the use of "stabilized" chlorine dioxide as a means of chlorine dioxide generation. The term stabilized chlorine dioxide refers to various compositions in which the chlorine dioxide is believed to be held in solution in the form of a labile complex. The stabilization of chlorine dioxide by the use of perborates was disclosed in United States Patent No. 2,701,781 (de Guevara). According to the de Guevara patent, an antiseptic solution of stabilized chlorine dioxide can be formed from an aqueous solution of chlorine dioxide and an inorganic boron compound with the boron compound and the chlorine dioxide being present in the solution as a labile complex. The chlorine dioxide, fixed in this stable condition, is an essential ingredient of the antiseptic solution. The de Guevara patent discloses that the chlorine dioxide may be introduced into the compositions either by *in situ* generation or it may be generated externally and introduced into the solution, as by bubbling the chlorine dioxide gas into the aqueous solution. Various methods may be employed for the external production of the chlorine dioxide, such as reaction of sulfuric acid with potassium chlorate or the reaction of the chlorate with moist oxalic acid. Alternatively, chlorine dioxide can be generated *in situ* by reaction of potassium chlorate and sulfuric acid. Note that whether the chlorine dioxide is produced *in situ* or externally, it is essentially an acid-induced liberation of the chlorine dioxide from potassium chlorate.

[0012] United States Patent No. 4,317,814 (Laso) describes stabilized chlorine dioxide preparations for treatment of burns in humans. Aqueous mixtures of perborate stabilized solutions of chlorine oxides, such as chlorine dioxide, in combination with glycerin are described for topical application to burned areas and may also be administered by oral application for treatment of burns. The aqueous solutions of perborate stabilized chlorine oxides are disclosed as being prepared by mixing with water the following: sodium chlorite, sodium hypochlorite, hydrochloric acid, sulfuric acid, an inorganic perborate, and a peroxy compound, such as sodium perborate. Thus, the solutions prepared in accordance with the Laso patent contain chlorine dioxide, hypochlorite and peroxy compounds as strong oxidizing agents and appear to utilize acid activation of the chlorine dioxide. The Laso patent states that the methods disclosed therein resulted in an immediate subsidence of burn related pain in many cases, that healing was rapid and characterized by an absence of infection or contraction, and that the burn scars were smooth and resembled normal tissue, thus eliminating the need for plastic surgery in certain cases. However, long term storage and stability are issues with the aqueous solutions described in the above-identified Laso patent, because such mixtures tend to generate chlorine dioxide very quickly, thus diminishing the long term stability of such mixtures.

[0013] United States Patent No. 3,271,242 (McNicholas et al.) describes stabilized chlorine dioxide solutions which

are formed by combining chlorine dioxide gas with an aqueous solution containing a peroxy compound, and subsequently heating the solution to a temperature which is high enough to drive off all free peroxide, but low enough not to destroy the chlorine dioxide. McNicholas et al., states that temperatures "much below" 70 degrees C are ineffective to drive off the free peroxide in the solution and that temperatures should not exceed 92 degrees C because at higher temperatures the chlorine dioxide will be driven off. McNicholas further states that, although not "entirely understood," it was believed that heating of the solution to drive off free peroxide was necessary because any free hydrogen peroxide allowed to remain in the solution would release the chlorine dioxide from the solution.

ii. Antibiotic Preparations

[0014] Antibiotic compounds have also been commonly used for the therapeutic treatment of burns, wounds, and skin and eye infections. While antibiotics may provide an effective form of treatment, several dangers are often associated with the use of antibiotics in the clinical environment. These dangers may include but are not limited to: (1) changes in the normal flora of the body, with resulting "superinfection" due to overgrowth of antibiotic resistant organisms; (2) direct antibiotic toxicity, particularly with prolonged use which can result in damage to kidneys, liver and neural tissue depending upon the type of antibiotic; (3) development of antibiotic resistant microbial populations which defy further treatment by antibiotics.

B. Difficult-To-Treat Dermal Disorders Other Than Wounds, Burns, Abrasions and Infections

[0015] While even minor wounds and abscesses can be difficult to treat in certain patients and/or under certain conditions, there are well known dermal disorders such as psoriasis and dermal ulcerations, which present particular challenges for successful treatment.

i. Psoriasis

[0016] Psoriasis is a noncontagious skin disorder that most commonly appears as inflamed swollen skin lesions covered with silvery white scale. This most common type of psoriasis is called "plaque psoriasis". Psoriasis comes in many different variations and degrees of severity. Different types of psoriasis display characteristics such as pus like blisters (pustular psoriasis), severe sloughing of the skin (erythrodermic psoriasis), drop like dots (guttate psoriasis) and smooth inflamed lesions (inverse psoriasis).

[0017] The cause of psoriasis is not presently known, though it is generally accepted that it has a genetic component, and it has recently been established that it is an autoimmune skin disorder. Approximately one in three people report a family history of psoriasis, but there is no pattern of inheritance. There are many cases in which children with no apparent family history of the disease will develop psoriasis.

[0018] The occurrence of psoriasis in any individual may depend on some precipitating event or "trigger factor". Examples of "trigger factors" believed to affect the occurrence of psoriasis include systemic infections such as strep throat, injury to the skin (the Koebner phenomenon), vaccinations, certain medications, and intramuscular injections or oral steroid medications. Once something triggers a person's genetic tendency to develop psoriasis, it is thought that in turn, the immune system triggers the excessive skin cell reproduction.

[0019] Skin cells are programmed to follow two possible programs: normal growth or wound healing. In a normal growth pattern, skin cells are created in the basal cell layer, and then move up through the epidermis to the stratum corneum, the outermost layer of the skin. Dead cells are shed from the skin at about the same rate as new cells are produced, maintaining a balance. This normal process takes about 28 days from cell birth to death. When skin is wounded, a wound healing program is triggered, also known as regenerative maturation. Cells are produced at a much faster rate, theoretically to replace and repair the wound. There is also an increased blood supply and localized inflammation. In many ways, psoriatic skin is similar to skin healing from a wound or reacting to a stimulus such as infection.

[0020] Lesional psoriasis is characterized by cell growth in the alternate growth program. Although there is no wound at a psoriatic lesion, skin cells (called "keratinocytes") behave as if there is. These keratinocytes switch from the normal growth program to regenerative maturation. Cells are created and pushed to the surface in as little as 2 4 days, and the skin cannot shed the cells fast enough. The excessive skin cells build up and form elevated, scaly lesions. The white scale (called "plaque") that usually covers the lesion is composed of dead skin cells, and the redness of the lesion is caused by increased blood supply to the area of rapidly dividing skin cells.

[0021] Although there is no known cure for psoriasis, various treatments have been demonstrated to provide temporary relief in some patients. However, the effectiveness of the currently accepted treatments for psoriasis is subject to considerable individual variation. As a result, patients and their physicians may have to experiment and/or combine therapies in order to discover the regimen that is most effective. The currently available treatments for psoriasis are often administered in step-wise fashion. Step 1 treatments include a) topical medications (e.g., topical steroids, topical retinoids),

b) systemic steroids, c) coal tar, d) anthralin, e) vitamin D3, and sunshine. Step 2 treatments include a) phototherapy (e.g., ultraviolet radiation), b) photochemotherapy (e.g., a combination of a topically applied radiation-activated agent followed by radiation to activate the agent) and c) combination therapy. Step 3 treatments include a) systemic drug therapies such as methotrexate, oral retinoids and cyclosporin and b) rotational therapy.

5

ii. Dermal Ulcerations

[0022] Dermal ulcerations are known to occur as a result of pressure, wear, or primary/secondary vascular disorders. Dermal ulcerations are generally classified according to their etiology, as follows:

10

a. Decubitus/Pressure Ulcers - A decubitus ulcer or pressure sore is a lesion caused by unrelieved pressure resulting in damage of the underlying tissue. Decubitus ulcers usually develop over a bony prominence such as the elbow or hip. The unrelieved pressure, along with numerous contributing factors, leads to the skin breakdown and persistent ulcerations.

15

b. Venous Ulcers - Venous ulcers may result from trauma or develop after chronic venous insufficiency (CVI). In CVI, venous valves do not close completely, allowing blood to flow back from the deep venous system through the perforator veins into the superficial venous system. Over time, the weight of this column of blood causes fluid and protein to exude into surrounding tissues, resulting in swollen, hyperpigmented ankles, tissue breakdown, and ulceration. Venous ulcers may be shallow or extend deep into muscle.

20

c. Arterial Ulcers - Leg ulcers also can develop in patients with arterial insufficiency caused by arterial vessel compression or obstruction, vessel wall changes, or chronic vasoconstriction. Smokers face an especially high risk of arterial disease because nicotine constricts arteries, encourages deposits of atherosclerotic plaque, and exacerbates inflammatory arterial disease (Buerger's disease) and vasoconstrictive disease (Raynaud's disease or phenomenon). Arterial ulcers, caused by trauma to an ischemic limb, can be very painful.

25

d. Diabetic Ulcers - Arterial insufficiency can be the cause of a nonhealing ulcer in a patient with diabetes. However, most diabetic ulcers result from diabetic neuropathy -- because the patient cannot feel pain in his foot, he is unaware of injuries, pressure from too tight shoes, or repetitive stress that can lead to skin breakdown.

30

[0023] There remains a need in the art for the formulation and development of new disinfectants and topically applicable preparations for the treatment of dermal disorders, such as wounds, burns, abrasions, infections, ulcerations, psoriasis and acne.

C. Contact Lens Soaking and Disinfection.

35

[0024] Whenever a contact lens is removed from an eye, it should be placed in a soaking and disinfecting solution until it is worn again. Soaking and disinfecting solutions have the following functions:

40

1. Assist in cleaning the lens of ocular secretions after the lens is removed from the eye;
2. To prevent eye infections by a bacterial contaminated lens; and
3. To maintain the state of hydrated equilibrium, which the lens achieves while it is being worn.

D. Contact Lens Cleaning.

45

[0025] During lens wear mucus material, lipids and proteins accumulate on contact lenses, making lens wear uncomfortable due to irritation, burning sensation, and redness. Accordingly, vision becomes blurry. To alleviate the discomforting problem, the soft or rigid contact lenses should be taken out of the eye, to be cleaned and disinfected regularly, using an enzymatic cleaner and a disinfecting solution. One of the serious complications associated with soft lenses can be a Giant Papillary Conjunctivitis (GPC). It is believed to be that the occurrence of the giant papillary conjunctivitis is mostly due to an inflammatory reaction associated with soft contact lens complication. This is almost always caused by protein deposits on contact lenses. GPC produces symptoms ranging from asymptomatic to itching, upper eye-lid edema, red eye, mucoid discharge, progressive contact lens intolerance. The in-the-eye cleaner of the present invention effectively cleans the protein deposits and maintains corneal epithelial cells healthy by keeping the corneal surface from microbial infection as well as by supplying molecular oxygen. Thereby, it provides convenience and benefits to both soft and rigid contact lens wearers.

55

E. Treatment of Ophthalmic Disorders.

i. Dry Eye

5 [0026] Dry eye is a syndrome in which tear production is inadequate or tear composition is inappropriate to properly wet the cornea and conjunctiva. A variety of disorders of the ocular tears causes sensations of dryness of the eyes, discomfort of presence of a foreign object to occur in the eye. In most instances, the tear film loses its normal continuity and breaks up rapidly so that it cannot maintain its structure during the interval between spontaneous blinks. All of those tear abnormalities may have multiple causes. Perhaps the most common form of dry eye is due to a decreased aqueous component in the tears. Untreated dry eye can be further deteriorated to produce more severe epithelial erosion, strands of epithelial cells, and local dry spots on the cornea, which can be further complicated by microbial infection. In its mild form, however, a feeling of dryness and irritation of the eye can be solved with artificial tears. Thus, artificial tear solution which has a broad spectrum antimicrobial activity with corneal lubricating property, can provide not only comfort but also beneficial effects on recovery of damaged corneal surface.

15

ii. Allergic Conjunctivitis

20 [0027] Airborne or hand borne allergens usually produce allergic conjunctivitis due to IgE-mediated hypersensitivity reaction. It presents itching, tearing, dry and sticky eyes, including lid-swelling, conjunctival hyperemia, papillary reaction, chemosis, and ropy mucoid discharge. The presence of hyaluronic acid in the tear, which is included in the formulation of artificial tear, would protect corneal surface from contacting the allergens. The broad spectrum antimicrobial agent of the present invention keeps the corneal surface from bacterial infection and also maintains the corneal epithelial cells healthy by supplying molecular oxygen. Thus, it provides beneficial effects on the eyes sensitive to allergens.

25

iii. Bacterial Invasion

30 [0028] Bacterial keratitis is one of the leading causes of blindness in the world. In the United States, an estimated 30,000 cases occur annually, with the popularity of contact lens wear having contributed to a rising incidence in the developed world. Statistical investigation indicates that about 30 of every 100,000 contact lens wearers develop ulcerative keratitis annually in the United States, thus making the disease a significant public health issue in view of potential blindness that can occur. While eyelids, blinking of the eyelids, and corneal and conjunctival epithelial cells provide barriers to microbial invasion, one or more of these defense mechanisms can become compromised. Such compromises can include lid abnormalities, exposure of the corneal surface, poor tear production, epithelial problems, medication toxicity, trauma, and incisional surgery. Ocular manifestations of bacterial keratitis are found in *staphylococcus* and *streptococcus* infections that tend to cause severe infiltration and necrosis which over time can lead to perforation. *Pseudomonas* keratitis tends to progress rapidly. This organism produces destructive enzymes, such as protease, lipase, and elastase, and exotoxins, which result in necrotic ulceration and perforation. *Serratia* keratitis starts as a superficial para-central ulcer, with the secretion of exotoxins and protease which can produce aggressive ulceration and perforation. In order for the bacterial keratitis to become established, microbial adhesions must bind to host cell receptors. Once this attachment has occurred, the destructive process of inflammation, necrosis, and angiogenesis can ensue.

40

45 [0029] Present treatment for bacterial keratitis relies primarily upon the use of broad spectrum antibiotic therapy. Such antibiotics include sulfonamides, trimethaprin, and quinolones. Also included are beta-lactams, penicillins, cephalosporins, aminoglycosides, tetracyclines, chloramphenicol, and erythromycin. While such antibiotics are in wide spread use, they can also become misused where antibiotic resistant pathogens emerge. Additionally, antibiotics only halt the proliferation of bacteria, but do not inhibit the activity of protease enzymes, endotoxins, or exotoxins. As is therefore apparent, a significant need is present for a bactericidal agent that addresses the proliferation of not only bacteria, but also protease enzymes, endotoxins and exotoxins.

50

BRIEF SUMMARY

55 [0030] The present invention provides antimicrobial preparations (e.g., solutions, gels, ointments, creams, etc.) for disinfection of articles or surfaces (e.g., contact lenses, counter tops, etc.), antisepsis of skin or other body parts, prevention or minimization of scarring, and/or treatment or prophylaxis of dermal (i.e., skin or mucous membrane) disorders (e.g., wounds, burns, infections, cold sores, ulcerations, psoriasis, scar forming lesions, acne), and the treatment of ophthalmic disorders (e.g., infection, inflammation, dry eye, allergic conjunctivitis, and wound healing). The antimicrobial preparations of this invention generally comprise from about 0.001% to about 0.20% by weight of a metal chlorite in combination with from 0.001% to 0.05% of a peroxy compound such as hydrogen peroxide. Additionally, the chlorite/peroxide preparations of the present invention may contain additional components such as polymeric lubricants

and surfactants, and/or may be formulated in a polymeric drug delivery system or liposomal preparation. The chlorite/peroxide preparations of the present invention have broad antimicrobial activity, including for example activity against gram negative and gram positive bacteria, yeasts and fungi. Moreover, when applied or administered to treat dermal disorders (e.g., wounds, burns, infections, ulcerations, acne and psoriasis), the chlorite/peroxide preparations of the present invention will not only prevent or lessen microbial infection, but will additionally provide oxygen to the affected tissue, assist in healing and deter scar formation.

[0031] Further, in accordance with the invention, there are provided methods for disinfection of items (e.g., contact lenses) and methods for treatment of dermal disorders (e.g., wounds, burns, infections, ulcerations and psoriasis) by application or administration of a chlorite/peroxide preparation of the present invention. With respect to contact lens disinfecting solution, as well as product formulations that will clean contact lenses in the eye without removing the lenses from the eye for cleaning, the concentration of the metal chlorite is between about 0.002% to about 0.20%. With respect to in-eye application, the present bactericidal product is a sterile, isotonic, buffered, clear, colorless solution that additionally contains polymeric lubricant and surfactant. The product has a two-year shelf life when stored in a container (e.g., a white opaque plastic bottle) at room temperature as a stabilized peroxy chloral complex of chlorite and peroxide.

[0032] In addition, the invention includes product formulations shown to have efficacy in the treatment of dry eye, wound healing, and allergic conjunctivitis.

[0033] Further in accordance with the invention, there are provided methods for deterring scar formation by application or administration of a chlorite/peroxide preparation of the present invention.

[0034] Further, in accordance with the invention, there are provided product formulations shown to have supra-additive efficacy in broad spectrum antimicrobial activity.

[0035] Furthermore, in accordance with the invention, there are provided methods for deterring eye infections, eye perforations and inflammation by application or administration of a chlorite/peroxide preparation of the present invention.

[0036] Further aspects and objects of the present invention will become apparent to those of skill in the art upon reading and understanding of the following detailed description and the examples set forth therein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0037]

Figure 1 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 7.3;

Figure 2 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 8.0;

Figure 3 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 8.8;

Figure 4 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 7.0;

Figure 5 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 6.44;

Figure 6 is a graph demonstrating the non-production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at pH level 6.0; and

Figure 7 is a graph demonstrating the production of chlorine dioxide at room temperature in the chlorite/peroxide preparation of the present invention at a pH level of 1.5.

DETAILED DESCRIPTION

[0038] The following detailed description and examples are provided for the purpose of describing certain exemplary embodiments of the invention only, and are not intended to limit the scope of the invention in any way.

[0039] The present invention provides preparations which contain chlorite (e.g., a metal chlorite such as sodium chlorite) in combination with a small amount of hydrogen peroxide in neutral aqueous (pH 5.0 - 8.8, preferably pH 7.0 - 7.8, and more preferably pH 7.0 - 7.4) solution. These preparations exhibit synergistic antimicrobial activity without generating chlorine dioxide during storage at room temperature, thereby rendering the stability of these solutions acceptable for pharmaceutical use. For example, an aqueous solution containing 400 ppm chlorite plus 100 ppm hydrogen peroxide remains stable beyond 18 months at room temperature, and is effective to reduce *Candida albicans* activity by 1.0 log within six hours of challenge, even though the individual components of such solution are ineffective when applied separately at the same concentrations to reduce *Candida albicans* activity. Additionally, the hydrogen peroxide present within the chlorite/peroxide solutions of the present invention readily decomposes into molecular oxygen and water, upon contact with the peroxidase and catalase enzymes present in tissue and/or some body fluids. Such in situ

generation of molecular oxygen contributes to cell vitality and enhances wound healing.

5 **[0040]** The chlorite/H₂O₂ solutions of the present invention are sufficiently stable to be formulated in combination with polymeric lubricants (non ionic and/or anionic; e.g., HPMC, Methocel, CMC, hyaluronic acid, etc.) and/or in combination with block polymer based surfactants (e.g., pluronics). For example, an aqueous chlorite/hydrogen peroxide system can be formulated together with methocel or hyaluronic acid as a lubricant and pluronics as a surfactant for contact lens disinfectant solution (viscosity up to 50 cps at 25 degrees C) in an ophthalmically acceptable tonicity (e.g., osmolality of at least about 200 mOsmol/kg) and a buffer to maintain the pH of the formulation within an acceptable physiological range. The formulation of the contact lens disinfection solution, artificial tear solution, and in-eye cleaner solution, contains chlorite preferably from about 0.005 to about 0.06 weight/volume percent and hydrogen peroxide preferably from about 10 0.0002 to about 0.05 weight/volume percent. Again, the presence of hydrogen peroxide provides the beneficial oxygen molecule to the cornea upon contact with catalase in the tear.

A. FORMULATIONS

15 **[0041]** The chlorite/peroxide preparations of the present invention may be formulated in various ways, including liquid solutions, gels, ointments, creams, sprays, etc. Set forth herebelow are a few examples of the types of specific formulations which may be prepared in accordance with this invention.

i. Stable Chlorite/Peroxide Liquid Solutions

20 **[0042]** The following Formula 1 is a first preferred formulation of a liquid chlorite/peroxide solution of the present invention:

FORMULA 1

25	Sodium Chlorite	0.005% - 0.10%
	Hydrogen Peroxide	0.005% - 0.05%
	Methocel A	0.05% - 0.2%
	Boric Acid	0.15%
30	Sodium Chloride	0.75%
	Pluronic F 68/F-127	0.1%
	HCl or NaOH	Adjust pH 7.4
	Purified water	Q.S. to volume

35 **[0043]** The following Formula 2 is a second preferred formulation of a liquid chlorite/peroxide solution of the present invention:

FORMULA 2

40	Sodium Chlorite	0.05%
	Hydrogen Peroxide	0.02%
	Carboxymethyl Cellulose	0.01%
	Boric Acid	0.15%
45	Sodium Chloride	0.75%
	Pluronic F 68/F-127	0.1%
	HCl or NaOH	Adjust pH 7.3
	Purified water	Q.S. to volume

50 **[0044]** The chlorite/peroxide solutions of the present invention, such as the solution of the above-shown preferred formulation, may be used for a variety of medical and non-medical applications including but not necessarily limited to a) disinfection of articles and surfaces such as contact lenses, medical/dental instruments, counter tops, treatment tables, combs and brushes, etc.; antiseptis of skin or body parts (e.g., a disinfectant hand wash, antiseptic facial scrub, etc.) and b) treatment or prophylaxis of dermal (i.e., skin or mucous membrane) disorders such as wounds, burns, infections, ulcerations, cold sores, psoriasis, acne, and c) deterrence or prevention of scar formation, and d) treatment of ophthalmic disorders (e.g., infections or inflammations caused by bacterial keratitis).

55 **[0045]** As pointed out earlier, the chlorite/hydrogen peroxide system of the present invention is sufficiently stable to

be formulated in a polymeric gel form or in a paste form. Furthermore, such polymeric gel or paste formulation can contain polymers which delay or control the release of the chlorite/hydrogen peroxide (e.g., a sustained release delivery system). Such sustained release formulations provide outstanding benefits of increasing therapeutic index by maintaining the effective concentration of chlorite/H₂O₂ for a prolonged time on the injured sites, by preventing the injured sites from external microbial contamination by forming a seal over the injured sites, and by providing oxygen molecule to the injured tissues. Unlike the conventional ointment, the polymeric gel provides a dry, clean, and comfortable coating on the injured sites upon application. Such gel formulations may contain polymeric drug delivery vehicles like hydroxypropyl methylcellulose (HPMC), methylcellulose (Methocel), hydroxyethylcellulose (HEC), hyaluronic acid, and carboxymethylcellulose (CMC), etc.

ii. A Stable Chlorite/Peroxide Gel

[0046] The following Formula 3 is a presently preferred formulation of a chlorite/peroxide gel of the present invention:
 FORMULA 3

Sodium Chlorite	0.02% - 0.10%
Hydrogen Peroxide	0.005% - 0.05%
Methocel A	2.0%
Boric Acid	0.15%
Sodium Chloride	0.75%
Pluronic F68/F-127	0.1%
HCl or NaOH	Adjust pH 7.4
Purified water	Q.S. to volume

[0047] Any of the preparations of the present invention may be formulated for sustained release of the active components by forming liposomes of the preparing in accordance with well known liposomal forming techniques and/or by adding to the formulation a pharmaceutically acceptable and effective amount (e.g., typically 1-20 percent by weight) of a sustained release component such as a polymer matrix or one or more of the following:

- a cellulose ester;
- hydroxymethylpropyl cellulose;
- methylhydroxyethyl cellulose;
- hydroxypropyl cellulose;
- hydroxyethyl cellulose;
- carboxymethyl cellulose;
- a salt of a cellulose ester;
- cellulose acetate;
- hydroxypropylmethyl cellulose phthalate;
- methacrylic acid-methyl methacrylate copolymer;
- methacrylic acid-ethyl acetate copolymer;
- polyvinylpyrrolidone;
- polyvinyl alcohol;
- hyaluronic acid;
- a phospholipid;
- cholesterol;
- a phospholipid having a neutral charge;
- a phospholipid having a negative charge;
- dipalmytoyl phoshatidyl choline;
- dipalmytoyl phoshatidyl serine; and,
- sodium salts thereof.

iii. A Stable Chlorite/Peroxide Ophthalmic Solution

[0048] The following Formula 4 is a presently preferred formulation of a chlorite/peroxide contact lens disinfecting solution for use in cleaning contact lenses residing in or out of the eye. The formulation additionally functions as a tear product for lubrication in dry-eye subjects.

EP 2 990 044 A1

FORMULA 4

5	Sodium Chlorite	0.002% - 0.20%
	Hydrogen Peroxide	0.005% - 0.05%
	Hyaluronic Acid	0.001% - 0.50%
	Boric Acid	0.15%
	Sodium Chloride	0.75%
10	Pluronic 127	0.05% - 2.0%
	HCl or NaOH	Adjust pH to 7.4
	Purified Water	Q.S. to Volume

[0049] As indicated earlier, the chlorite/peroxide preparation of the present invention, whether it be in the form of liquid solution, gel, ointment, cream, spray, etc., is specifically composed to maintain chlorite such as sodium chlorite and hydrogen peroxide as active ingredients at a pH range of 5.0 - 8.8 without generating chlorine dioxide during storage at room temperature. By way of illustration, multiple experiments were conducted on the liquid sodium chlorite/hydrogen peroxide solution in accordance with Formula 2 at different levels of pH within the specified range. However, it should be expressly stated herein that such experimentations should in no way be limited to liquid solution forms only, but are performed to illustrate the non-production of chlorine dioxide in the various forms of the present chlorite/peroxide preparation at different pH levels.

[0050] The following experimentations were designed to demonstrate the stability of chlorite such as sodium chlorite and hydrogen peroxide antibacterial formulation at neutral, basic and acidic levels of pH. More specifically, the quantitative levels of sodium chlorite and the generation of chlorine dioxide were determined at the pH levels of 7.3, 8.0, 8.8, 7.0, 6.44 and 6.0. 0.1 Normal hydrochloric acid solution and 0.1 Normal sodium hydroxide solution were applied to adjust the pH levels in the experimentations. Sterile 0.9% sodium chloride sterile solution was also applied. A placebo solution with the following formulation was further applied in a spectrophotometer (e.g., Lambda 20 Model UV - Vis. spectrophotometer) to find and measure the levels of sodium chlorite and the generation of chlorine dioxide at varying pH levels:

Placebo Solution

30	Hydrogen Peroxide	0.02%
	Carboxymethyl Cellulose	0.01%
	Boric Acid	0.15%
35	Sodium Chloride	0.75%
	Pluronic F 68/F-127	0.1%
	HCl or NaOH	Adjust pH 7.3
	Purified water	Q.S. to volume

40

Experiment 1: pH Level of 7.3

[0051] Experiment: Fill the first cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid sodium chlorite/hydrogen peroxide solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 1.

[0052] Result: The liquid solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 7.3. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH level of 7.3.

[0053] Hydrogen peroxide does not absorb in the 200nm to 400nm range. Therefore, as seen in Figure 1, absorption peaks for hydrogen peroxide were not detected.

[0054] Sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm.

[0055] Scanning the solutions that have a pH of 7.3 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

[0056] Interpretation: The liquid sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm.

EP 2 990 044 A1

[0057] This clearly indicates that at pH level of 7.3, the liquid sodium chlorite/hydrogen peroxide solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that sodium chlorite is stable at pH level of 7.3, and the sodium chlorite is not breaking up and forming the chlorine dioxide.

5 Experiment 2: pH Level of 8.0

[0058] Experiment: Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal sodium hydroxide solution to each container so as to adjust the pH of both the placebo solution as well as the liquid solution to a pH level of 8.0.

10 [0059] Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid sodium chlorite/hydrogen peroxide solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 2.

15 [0060] Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 8.0. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH level of 8.0.

[0061] As mentioned shortly above, hydrogen peroxide does not absorb in the 200nm to 400nm range. Therefore, as seen in Figure 2, absorption peaks for hydrogen peroxide were not detected. As also mentioned above, sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm.

20 [0062] Scanning the solutions that have a pH level of 8.0 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

[0063] Interpretation: The liquid sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at the pH level of 8.0, the liquid sodium chlorite/hydrogen peroxide solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that sodium chlorite is stable at the pH level of 8.0, and the chlorite is not breaking up and forming chlorine dioxide.

30 Experiment 3: pH Level of 8.8

[0064] Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal sodium hydroxide solution to each container so as to adjust the pH of both the placebo solution as well as the liquid solution to a pH level of 8.8.

35 [0065] Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid sodium chlorite/hydrogen peroxide solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 3.

[0066] Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 8.8. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH level of 8.8.

40 [0067] As already discussed, hydrogen peroxide does not absorb in the 200nm to 400nm range. Therefore, as seen in Figure 3, absorption peaks for hydrogen peroxide were not detected. As also discussed, sodium chlorite has an absorption maximum at 260nm, while chlorine Dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm.

45 [0068] Scanning the solutions that have a pH level of 8.8 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

[0069] Interpretation: The liquid sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at the pH level of 8.8, the liquid sodium chlorite/hydrogen peroxide solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that sodium chlorite is stable at the pH level of 8.8, and the chlorite is not breaking up and forming chlorine dioxide.

Experiment 4: pH Level of 7.0

55 [0070] Experiment: Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal hydrochloric acid solution to each container so as to adjust the pH of both the placebo solution as well as the liquid solution to a pH level of 7.0.

[0071] Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path

EP 2 990 044 A1

of the spectrophotometer. Fill the second cuvette with the liquid sodium chlorite/hydrogen peroxide solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 4.

5 **[0072]** Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 7.0. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH level of 7.0. Hydrogen peroxide does not absorb in the 200nm to 400nm range. Therefore, as seen in Figure 4, absorption peaks for hydrogen peroxide were not detected.

10 **[0073]** Sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm. Scanning the solutions that have a pH of 7.0 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

15 **[0074]** Interpretation: The sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at the pH level of 7.0, the liquid solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that sodium chlorite is stable at pH of 7.0, and the chlorite is not breaking up and forming chlorine dioxide.

Experiment 5: pH Level of 6.44

20 **[0075]** Experiment: Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal hydrochloric acid solution to each container so as to adjust the pH of both the placebo solution as well as the liquid solution to a pH level of 6.44.

25 **[0076]** Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 5.

[0077] Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 6.44.

30 **[0078]** The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at pH = 6.44. Hydrogen peroxide does not absorb in the 200nm to 400nm range, and thus no absorption peaks for hydrogen peroxide were detected. Sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm.

[0079] Scanning the solutions that have a pH of 6.44 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

35 **[0080]** Interpretation: The liquid sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at pH of 6.44, the liquid solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that sodium chlorite is stable at pH of 6.44, and the chlorite is not breaking up and forming chlorine dioxide.

Experiment 6: pH Level of 6.0

40 **[0081]** Experiment: Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal hydrochloric acid solution to each container so as to adjust the pH of both the placebo solution as well as the liquid solution to a pH level of 6.0.

45 **[0082]** Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid sodium chlorite/hydrogen peroxide solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 6.

50 **[0083]** Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH level of 6.0. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH level of 6.0. Hydrogen peroxide does not absorb in the 200nm to 400nm range. Therefore, as seen in Figure 6, absorption peaks for hydrogen peroxide were not detected.

55 **[0084]** Sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm. Scanning the solutions that have a pH of 6.0 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

[0085] Interpretation: The sodium chlorite/hydrogen peroxide solution does show sodium chlorite peak at 260nm, but does not show any chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at pH level of 6.0, the liquid solution has only sodium chlorite, and does not contain any quantities of chlorine dioxide. This is a clear indication that

EP 2 990 044 A1

sodium chlorite is stable at pH of 6.0, and the chlorite is not breaking up and forming chlorine dioxide.

Experiment 7: pH Level of 1.5

5 **[0086]** Experiment: Dispense 25 mL. of the placebo solution and 25 mL. of the liquid sodium chlorite/hydrogen peroxide solution into 2 clean containers. Add 0.1 Normal hydrochloric acid solution to each container so as to adjust the pH of both the placebo solution as well as the bactericidal solution to a pH of 1.5.

[0087] Fill one of the cuvette with the placebo solution, wipe it clean, and place the cuvette in the standard beam path of the spectrophotometer. Fill the second cuvette with the liquid solution, wipe it clean and place the cuvette in the sample beam path of the spectrophotometer. Scan the solutions from 200nm to 400nm and record the results. Plot and printout the results, as illustrated in the graph shown in Figure 7.

10 **[0088]** Result: The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at pH of 1.5. The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at pH of 1.5. As explained earlier, hydrogen peroxide does not absorb in the 200nm to 400nm range, and as such, no absorption peaks for hydrogen peroxide were detected.

15 **[0089]** Also explained earlier, sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide which is a degradation product of sodium chlorite has an absorption maximum at 355nm - 358nm. Scanning the solutions that have a pH of 1.5 between the 200nm and 400nm will give a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

20 **[0090]** Interpretation: The liquid sodium chlorite/hydrogen peroxide solution does not show sodium chlorite peak at 260nm, but does show a large chlorine dioxide peak at 355nm - 358nm. This clearly indicates that at the pH level of 1.5, the liquid sodium chlorite/hydrogen peroxide solution does not have any sodium chlorite. Rather, it clearly shows that the sodium chlorite has been degraded and converted to chlorine dioxide. This is a clear indication that at pH of 1.5, sodium chlorite is very unstable, and all chlorite that is present in the liquid solution is converted to chlorine dioxide.

25

Results for Experiments 1-7

[0091] The liquid sodium chlorite/hydrogen peroxide solution contained sodium chlorite and hydrogen peroxide as active ingredients, as well as buffering and tonicity agents at the pH levels of 1.5, 6.0, 6.44, 7.0, 7.3, 8.0 and 8.8.

30 **[0092]** The placebo solution contained hydrogen peroxide as active ingredient, as well as buffering and tonicity agents at the pH levels of 1.5, 6.0, 6.44, 7.0, 7.3, 8.0 and 8.8.

[0093] Hydrogen peroxide does not absorb in the 200nm to 400nm range.

[0094] Sodium chlorite has an absorption maximum at 260nm, while chlorine dioxide has an absorption maximum at 355nm - 358nm.

35 **[0095]** Scanning the solutions between the 200nm and 400nm gave a quantitative value for sodium chlorite as well as chlorine dioxide in the same scan.

Interpretation of Results for Experiments 1-7

40 **[0096]** The liquid sodium chlorite/hydrogen peroxide solutions at the pH levels of 6.0, 6.44, 7.0, 7.3, 8.0 and 8.8 does show the presence of sodium chlorite peak at 260nm, but does not show the presence of chlorine dioxide peak at 355nm - 358nm.

[0097] In contrast, the liquid sodium chlorite/hydrogen peroxide solution at pH of 1.5 does not show the presence of sodium chlorite peak at 260nm, but does show the presence of chlorine dioxide peak at 355nm - 358nm.

45

Conclusion of Results for Experiments 1-7

[0098] The results clearly show that one can quantitatively determine the level of sodium chlorite as well as chlorine dioxide which is present in the liquid sodium chlorite/hydrogen peroxide solution at the pH levels of 1.5, 6.0, 6.44, 7.0, 7.3, 8.0 and 8.8.

[0099] The results also show that the storage of the liquid sodium chlorite/hydrogen peroxide solution at about room temperature (e.g., in a white opaque bottle exposed to air at room temperature) does not produce any chlorine dioxide as determined by the absence of any absorbance at 355nm - 358nm.

50 **[0100]** In conclusion, the results of Experiments 1-7 clearly indicate that the liquid sodium chlorite/hydrogen peroxide solution retains sodium chlorite at the pH range of 6.0-8.8 without the generation of chlorine dioxide. The liquid solution, however, degrades and generates chlorine dioxide upon the acidification of the solution to pH of 1.5. Thus, these results also strongly indicate that the liquid sodium chlorite/hydrogen peroxide solution does not contain chlorine dioxide when it is manufactured, nor does the solution degrade to generate chlorine dioxide after storage at about room temperature

at the pH levels of 6.0, 6.44, 7.0, 7.3, 8.0, or 8.8.

[0101] Furthermore, these results present clear evidence that the liquid sodium chlorite/hydrogen peroxide solution of the present invention has its bactericidal properties in the pH range studied due to the sodium chlorite/ hydrogen peroxide and not due to chlorine dioxide. This is very much unlike other prior art inventions that have sodium chlorite as a starting material as, but the active bactericide is the chlorine dioxide which is generated by the acidification of the sodium chlorite.

B. EXAMPLES OF THERAPEUTIC APPLICATIONS

[0102] The following are specific examples of therapeutic applications of the chlorite/peroxide preparations of the present invention.

i. Example 1: Treatment of Psoriasis-No Crossover

[0103] A human patient having psoriasis plaques present on both arms is treated as follows:
Twice daily application to plaques on the left arm only, of a chlorite/peroxide solution having the following formulation:

Sodium Chlorite	0.06%
Hydrogen Peroxide	0.01%
HPMC	2.0%
Boric Acid	0.15%
HCl or NaOH	to adjust pH 7.4
Purified water	Q.S. to volume

Twice daily application to plaques on the right arm only of a commercially available 0.1% triamcinolone acetonide cream.

[0104] The chlorite/peroxide treated psoriatic plaques on the right arm began to become less severe within 24 hours of beginning treatment and had substantially disappeared within three days of beginning treatment. However, the triamcinolone acetonide treated psoriatic plaques present on the left arm remained unchanged and inflamed during the two week treatment period.

ii. Example 2: Treatment of Psoriasis-Crossover

[0105] A human patient having psoriasis plaques present on both arms is treated for two weeks, as follows:
Twice daily application to plaques on the left arm only, of a chlorite/peroxide solution having the following formulation:

Sodium Chlorite	0.06%
Hydrogen Peroxide	0.01%
HPMC	2.0%
Boric Acid	0.15%
HCl or NaOH	to adjust pH 7.4
Purified water	Q.S. to volume/100%

Twice daily application to plaques on the right arm only of a commercially available 0.1% triamcinolone acetonide cream.

[0106] The chlorite/peroxide treated psoriatic plaques on the right arm began to become less severe within 24 hours of beginning treatment and had substantially disappeared within one week of beginning treatment. However, the triamcinolone acetonide treated psoriatic plaques present on the left arm remained unchanged and inflamed during the two week treatment period.

[0107] Beginning the day after the end of the initial two week treatment period, and continuing for a second two week treatment period, the patient was treated as follows:

Twice daily application to plaques on the left arm only of the same commercially available 0.1% triamcinolone acetonide cream described hereabove in this example.

Twice daily application to plaques on the right arm only, of the same chlorite/peroxide sustained release gel described hereabove in this example.

5 **[0108]** Within 24 hours of commencing the second treatment period, the psoriatic lesions on the right arm began to subside. By day three and continuing through the end of the second two week treatment period, the psoriatic lesions on the right arm had substantially disappeared.

iii. Example 3: Treatment of Cold Sores

10 **[0109]** A patient with painful, fluid-containing cold sores (i.e., chancre sores) on his lips was treated twice daily by application to the lips of a chlorite/peroxide preparation prepared in accordance with Formula 1 above.

15 **[0110]** Within 6 to 12 hours of the first application of the chlorite/peroxide preparation, the patient reported that the pain had subsided. Within 24 hours of the first application of the chlorite/peroxide preparation, the fluid contained within the cold sores had substantially dissipated and the cold sores appeared dry. Within six days of the first application of the chlorite/peroxide preparation the cold sores had substantially disappeared and the lips appeared normal, whereas cold sores of such severity typically require substantially longer than six days to completely disappear and heal.

iv. Example 4: Treatment of Venous Ulcer

20 **[0111]** A patient with a venous ulcer on the right leg of 3-4 cm diameter which had been present for 9-12 months was treated by twice daily application to the ulcer of gauze soaked with a chlorite/peroxide liquid solution prepared in accordance with Formula 1 above.

25 **[0112]** Within three days after commencement of treatment the ulcer appeared clean and dry. Within 14 days of the commencement of treatment the ulcer began to decrease in size and healthy new tissue was observed about its periphery. At 35 days after commencement of treatment, the ulcer had completely healed, without scarring, and the area where the ulcer had been located was free of pain.

v. Example 5: Treatment of Diabetic Decubitus Ulcer

30 **[0113]** A non-ambulatory, diabetic patient with decubitus ulcers on both legs and some toes, of 12-18 month duration, was treated by daily application of clean, sterile gauze to the ulcers and saturation of each gauze, three times each day, with a liquid chlorite/peroxide solution prepared in accordance with Formula 1 above. Within four to seven days of commencing the chlorite/hydrogen peroxide treatments the ulcers began to appear less inflamed, clean and dry. About seven to ten days after commencement of the chlorite/hydrogen peroxide treatment, granulation tissue began to form within the ulcers. Within 12 to 14 days, re-epithelialization was observed to have begun within the ulcerated areas except for one toe ulcer which had been particularly severe and had permeated to the bone of the toe. Within 30 to 45 days of the commencement of treatment, all of the ulcers except for the severe toe ulcer had completely closed and re-epithelialized, without irregular scar formation. Also, at 30 to 45 days after the commencement of treatment, the toe ulcer had also become substantially smaller (but was not completely closed) and the patient was able to walk. The liquid and or gel formulations of the present invention, such as Formulas 1 and 2 above, may also be applied topically to prevent scar formation due to wounds, burns, acne, infections, trauma, surgical incision, or any other scar-forming lesion or disorder.

vi. Example 6:

45 a. Treatment of Dry Eye Conditions

[0114] Subjects with dry eye conditions have itchy and scratchy eyes. In extreme cases, the subjects have more serious problems that can interfere with health maintenance. Subjects were treated with a preferred tear product of the following formulation:

50

Sodium Chlorite	0.005% - 0.02%
Hydrogen Peroxide	0.01%
Methylcellulose A4M	0.075%
Hyaluronic Acid	0.10% - 0.125%
Boric Acid	0.15%
Sodium Chloride, USP	0.75%
Pluronic 127	0.10%

55

EP 2 990 044 A1

(continued)

HCl or NaOH	Adjust pH to 7.4
Purified Water	Q.S. to Volume

5

[0115] Testing of dry eye subjects with rose bengal stain or fluorescein gives a good indication regarding the condition of the corneal epithelial health, while rose bengal staining provides a good indication of the number of dead epithelial cells on the cornea as well as conjunctiva.

10

[0116] Two subjects with dry eye condition were tested with rose bengal stain, and the quantitative staining to the cornea and conjunctiva was documented by photographs. The subjects started using the above preferred tear product at a dosage of two drops three times per day. At the end of two weeks, the two subjects were tested with rose bengal stain and the level of staining was quantitatively documented by photography. The results showed a 50% to 70% reduction in rose bengal staining, which clearly indicates that the preferred tear formulation was ameliorating the corneal and conjunctival cells from dying.

15

[0117] In addition to an objective determination of the health of the epithelial cells, the two subjects were tested subjectively regarding the safety and efficacy of the preferred tear product. First of all, slit-lamp biomicroscopy of the subjects during the two-week treatment period did not show any redness, irritation, inflammation, or other signs of discomfort. Second, the subjects indicated that the application of the tear product completely removed symptoms of redness, itching, scratching, pain, and dryness due to dry eye while providing lubrication that lasted for several hours.

20

It is therefore evident that the tear product exhibits both safety and efficacy in the treatment of dry eye. As is further recognized in view of the foregoing antimicrobial activity of such compositions, the tear product will also have efficacy in enhancing wound healing within the eye such as after surgery where bacterial infections are to be avoided.

b. Treatment of Allergic Conjunctivitis

25

[0118] In addition to treating dry eye condition with the above preferred tear product, the product was also tested in the treatment of conditions from allergic conjunctivitis. In particular, two subjects suffering from allergic conjunctivitis including itchy, scratchy eyes with constant tearing applied two drops of the product three times per day. This dosage resulted in the disappearance of the symptoms.

30

C. EXAMPLES OF CONTACT LENS CLEANSING

i. Example 1: Soaking, Cleaning and Disinfecting

35

[0119] The following formulation is a preferred disinfecting solution applicable to the cleaning of contact lenses by conventional soaking.

Sodium Chlorite	0.05%
Hydrogen Peroxide	0.02%
Methylcellulose A4M	0.075%
Hyaluronic Acid	0.05% - 0.10%
Boric Acid	0.15%
Pluronic 127	0.25% - 0.50%
Sodium Chloride USP	0.75%
HCl or NaOH	Adjust pH to 7.4
Purified Water	Q.S. to Volume

40

45

50

[0120] Six subjects using soft hydrophilic contact lenses soaked the lenses in the above disinfecting solution and then placed the lenses directly into the eyes. Soaking was performed nightly or on an as-needed basis. All six subjects reported that the lenses felt very comfortable, and that no adverse effects (e.g., burning, stinging, redness, pain) were experienced. Additionally, the solution extended the comfort and clean condition of the lenses for several weeks beyond such extension experienced with other commercially available disinfecting solutions.

55

[0121] The disinfecting solution can be used with soft hydrophilic lenses of varying water content (e.g., 38% to 75%), as well as with silicone acrylate rigid gas permeable lenses. Cycling studies of soft lenses soaked daily in the solution for 30 days showed no damage or change in the physical and chemical characteristics of the lenses. Eye comfort, as earlier noted, is achieved through non-binding and non-accumulating of preservative in soft or rigid gas permeable

EP 2 990 044 A1

lenses, while such binding and accumulation can be found in certain currently commercially available formulations to cause irritation and discomfort.

ii. Example 2: Cleaning While Wearing

[0122] The following formulation is a preferred disinfecting in-eye solution applicable to the cleaning of contact lenses while they are being worn by introducing the solution into the eye:

Sodium Chlorite	0.02%
Hydrogen Peroxide	0.01% - 0.02%
Methylcellulose A4M	0.075%
Hyaluronic Acid	0.075% - 0.10%
Boric Acid	0.15%
Sodium Chloride USP	0.75%
Pluronic 127	0.75%
HCl or NaOH	Adjust pH to 7.4
Purified Water	Q.S. to Volume

[0123] Four subjects applied two drops of the above in-eye solution three times per day for 30 days to contact lenses while being worn. Examinations of all of the subjects showed no irritation, burning, stinging, or adverse effects of any kind. These subjects further reported that the solution felt soothing and lubricating.

[0124] Two subjects were involved in a comparative study where, first of all, they wore ACUVUE disposable lenses continuously for two weeks with occasional removal and cleaning with commercially available cleaning solutions followed with a saline rinse. After 14 days, the lenses became very gritty and uncomfortable, and were discarded. Second, the two subjects started with new ACUVUE lenses and practiced daily application of the present in-eye solution three times per day without removing or touching the lenses. These subjects were able to wear the lenses for three to four weeks before replacement. Additionally, the inconvenience of cleaning the lenses outside the eye was completely eliminated, as was the risk of lens loss, tearing, or contamination. It is therefore evident that the present in-eye cleaning solution provides cleansing efficacy as well as convenience.

D. IN-VITRO AND IN-VIVO ANTIMICROBIAL EFFICACY

i. Synergistic Activity

[0125] Tables I and II compare the antimicrobial effects of (a) 400 ppm sodium chlorite alone; (b) 200 ppm hydrogen peroxide alone; and (c) 400 ppm sodium chlorite and 200 ppm hydrogen peroxide in combination against antibiotic-resistant strains of *staphylococcus haemolyticus* (Table I) and *pseudomonas aeruginosa* (Table II) both isolated from human infected eyes. Tables I and II summarize the antimicrobial effects observed at time points one and two hours after introduction of the test solutions.

TABLE I

<i>(staphylococcus haemolyticus: Initial inoculum = 1.01 x 10⁷:Log 7.03)</i>			
Time (hours)	Log Reduction NaClO ₂ alone (400 ppm)	Log Reduction H ₂ O ₂ alone (200 ppm)	NaClO ₂ & H ₂ O ₂ (400 ppm & 200 ppm)
1	0.11	0.20	0.69
2	1.01	0.23	2.43

TABLE II

(pseudomonas aeruginosa: Initial inoculum = 2.22 x 10 ⁶ :Log 6.35)			
Time (hours)	Log Reduction NaClO ₂ alone (400 ppm)	Log Reduction H ₂ O ₂ alone (200 ppm)	NaClO ₂ & H ₂ O ₂ (400 ppm & 200 ppm)
1	0.351	0.01	0.04
2	1.35	0.54	6.35

5

10

15

20

25

[0126] In the experiment summarized in Table I, sodium chlorite alone caused a Log reduction in *staphylococcus haemolyticus* bacteria of 0.11 at 1 hour and 1.01 at 2 hours. Hydrogen peroxide alone caused a Log reduction in *staphylococcus haemolyticus* bacteria of 0.20 at 1 hour and 0.23 at 2 hours and the combination of sodium chlorite and hydrogen peroxide caused a Log reduction in *staphylococcus haemolyticus* bacteria of 0.69 at 1 hour and 2.43 at 2 hours. Thus, in this experiment, the antimicrobial effect of the sodium chlorite-hydrogen peroxide combination was significantly greater than the sums of the effects of the sodium chlorite and hydrogen peroxide alone, at least at the 2 hour time point. Accordingly, it is concluded that the sodium chlorite-hydrogen peroxide combination exhibited a supra-additive effect against the strain of *staphylococcus haemolyticus* used in this experiment.

[0127] In the experiment summarized in Table II, sodium chlorite alone caused a Log reduction in *pseudomonas aeruginosa* bacteria of 0.35 at 1 hour and 1.35 at 2 hours. Hydrogen peroxide alone caused a Log reduction in *pseudomonas aeruginosa* bacteria of 0.01 at 1 hour and 0.54 at 2 hours and the combination of sodium chlorite and hydrogen peroxide caused a Log reduction in *pseudomonas aeruginosa* bacteria 0.04 at 1 hour and 6.35 at 2 hours. Thus, in this experiment, the antimicrobial effect of the sodium chlorite-hydrogen peroxide combination was significantly greater than the sums of the effects of the sodium chlorite and hydrogen peroxide alone, at least in the 2 hour time point. Accordingly, it is concluded that the sodium chlorite-hydrogen peroxide combination exhibited a supra-additive effect against the strain of *pseudomonas aeruginosa* used in this experiment.

ii. Animal Testing

30

[0128] *S. haemolyticus* keratitis was induced in respective right eyes of 12 rabbits by dropping broth containing 50,000 CFU/ml of *S. haemolyticus* onto abraded corneas of these eyes. After 24 hours, all corneas were likewise infected, and the rabbits were divided randomly into three groups. The rabbits (five) of Group I then were treated with the chlorite-hydrogen peroxide formulation defined above as cleaning while wearing contact lenses (here termed "Bactericide"); the rabbits (five) of Group II were treated with commercially available 0.3% ofloxacin antibiotic ophthalmic solution; and the rabbits (two) of Group III were untreated to serve as a control.

35

[0129] At 24 and 48 hours post infection, the rabbits underwent visual eye examination, photographic documentation and biomicroscopy. After 24 hours of treatment, three animals each from Groups I and II and one animal from Group III were sacrificed. The eyes were enucleated and an 8 mm disc of cornea was homogenized and plated onto growth media for microbial isolation and quantification. After 48 hours of treatment, the same procedure was followed for the remaining animals.

40

[0130] Tables III, IV and V summarize the results of this experimentation. As is there apparent, the Bactericide of the present invention exhibited superior overall results as compared to the competing commercially available regimens. The results therefore confirm that the clinical efficacy of the Bactericide is better than the antibiotic treatment. In addition to having excellent bactericidal properties, it is demonstrated that bactericide superiority is probably attributable to inactivation of bacterial proteolytic enzymes (thus decreasing bacterial virulence) and inactivation of bacterial toxins responsible for inflammation and hyperemia.

45

TABLE III

IN-VIVO ANTIMICROBIAL EFFICACY IN INFECTIOUS <i>S. HAEMOLYTICUS</i> KERATITIS IN RABBITS			
Post Treatment Time	Group I Bactericide	Group II 0.3% Ofloxacin	Group III Untreated Control
24 hours	i) 0 CFU ii) 18,000 CFU iii) 0 CFU	i) 23,000 CFU ii) 5,000 CFU iii) 11,000 CFU	39,000 CFU

50

55

EP 2 990 044 A1

(continued)

IN-VIVO ANTIMICROBIAL EFFICACY IN INFECTIOUS <i>S. HAEMOLYTICUS</i> KERATITIS IN RABBITS			
Post Treatment Time	Group I Bactericide	Group II 0.3% Ofloxacin	Group III Untreated Control
Average	6,000 CFU	13,000 CFU	39,000 CFU
48 hours	i) 0 CFU ii) 0 CFU	i) 5,000 CFU ii) 5,200 CFU	231,000 CFU
Average	0 CFU	5,100 CFU	231,000 CFU

TABLE IV

IN-VIVO CLINICAL EFFICACY IN INFECTIOUS <i>S. HAEMOLYTICUS</i> KERATITIS IN RABBITS			
Time	Group I Bactericide	Group II 0.3% Ofloxacin	Group III Untreated Control
24 hours after infection	inflammation (+2) hyperemia (+2) corneal edema (+2)	inflammation(+2) hyperemia (+2) corneal edema (+2)	inflammation(+2) hyperemia (+2) corneal edema (+2)
24 hours after treatment	inflammation (0) hyperemia (0) corneal edema (0)	inflammation(+2) hyperemia (+2) corneal edema (+2)	inflammation(+3) hyperemia (+3) corneal edema (+3)
Time	Group I Bactericide	Group II 0.3% Ofloxacin	Group III Untreated Control
48 hours after treatment	inflammation (0) hyperemia (0) corneal edema (0)	inflammation(+1) hyperemia (+1) corneal edema (+1)	inflammation(+3) hyperemia (+3) corneal edema (+3)

TABLE V

IN-VITRO INHIBITION OF PROTEOLYTIC ENZYME ACTIVITY		
Inhibition of proteolytic enzyme activity of Trypsin and porcine pancreatic Elastase		
Enzyme	Concentration of Bactericide	% Inhibition of Enzyme activity
Elastase (porcine)	0.18 ppm	46%
Trypsin	0.12 ppm	28%

E. OCULAR TOLERABILITY AND DEGRADATION SPEED

i. Ocular Tolerability of High Levels of Hydrogen Peroxide

[0131] Previously, it was believed that the upper limit of human ocular tolerability of hydrogen peroxide is about 100ppm (0.01 wt. %). See Paugh, J.R., Brennan, N.A., and Efron, N., Ocular Response to Hydrogen Peroxide, Am. J. Optom. Physiol. Opt. 1988 Feb;65(2):91-8. The following experiments show, however, that when combined with sodium chlorite, hydrogen peroxide is well tolerated by human eyes in levels up to 500ppm (0.05 wt. %). In all of the following experiments, the hydrogen peroxide was combined with 400ppm (0.04 wt. %) of sodium chlorite in 0.2% boric acid at pH 7.4 and filtered through a 0.2µm Acrodisc syringe filter. Two drops of each formulation were then placed in the cul-de-sac of two normal human eyes. Upon instillation of the drops, the subjects were instructed to close their eyelids. The subjects' ocular symptoms of the treated eyes were observed and graded over a period of one hour post instillation, for burning and stinging sensation, pain, redness, tearing, itching, photopsia, photophobia, discharge, and foreign body sensation. The observations are presented below.

EP 2 990 044 A1

TABLE VI

Experiment 1: Human Ocular Response to 100ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Burning/Stinging	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0
Foreign body sensation	0	0	0	0	0	0	0	0
Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe								

TABLE VII

Experiment 2: Human Ocular Response to 200ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Burning/Stinging	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0
Foreign body sensation	0	0	0	0	0	0	0	0
Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe								

TABLE VIII

Experiment 3: Human Ocular Response to 300ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Burning/Stinging	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0

EP 2 990 044 A1

(continued)

Experiment 3: Human Ocular Response to 300ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Itching	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0
Foreign body sensation	0	0	0	0	0	0	0	0
Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe								

TABLE IX

Experiment 4: Human Ocular Response to 400ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 second	1 minute	2 minutes	3 minutes	5 minutes	10 minute	60 minute
Burning/Sting	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0
Foreign body sensation	0	0	0	0	0	0	0	0
Grading Scale: 0 = None; +0,5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe								

TABLE X

Experiment 5: Human Ocular Response to 500ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Burning/Stinging	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0

EP 2 990 044 A1

(continued)

Experiment 5: Human Ocular Response to 500ppm of Hydrogen Peroxide								
Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	60 minutes
Foreign body sensation	0	0	0	0	0	0	0	0
Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe								

[0132] As can be seen from the above data, hydrogen peroxide levels up to 500ppm are very well tolerated by human eyes when in the presence of sodium chlorite. There were no signs of irritation, inflammation, or any other adverse effects associated with the instillation of the formulations containing up to 500ppm hydrogen peroxide. These results show that hydrogen peroxide up to 500ppm can be very safe and free of any adverse effects to the human eye when used in conjunction with sodium chlorite. Furthermore as discussed above, an outstanding synergistic antimicrobial activity has been discovered with formulations containing sodium chlorite and hydrogen peroxide. Because the previous literature (*See Paugh*) reported that human ocular tolerability of hydrogen peroxide is about 100ppm, it is believed that the sodium chlorite must be stabilizing the hydrogen peroxide by forming a kind of transient complex molecule (e.g., peroxychlorite), which exhibits the excellent synergistic antimicrobial activity and degrades to innocuous products like water, oxygen, and salt upon contact with biological systems, as will be discussed in greater detail below.

ii. Hydrogen Peroxide/Sodium Chlorite Degradation in the Eye

[0133] The following experiments were designed to determine the speed of self degradation of the hydrogen peroxide/sodium chlorite formulation when placed in the human eye and to determine the level of ocular symptomatology associated with the formulation when used in an "in the eye" contact lens cleaner product or an artificial tear product.

Experiment 1: "In the Eye" Contact Lens Cleaner

[0134] An "in the eye" contact lens cleaner containing 0.5g carboxymethylcellulose, 0.5g pluronic, and 0.05g hydrogen peroxide/sodium chlorite mixture in 100mL sterile water was provided. The cleaner contained 400ppm sodium chlorite and 100ppm hydrogen peroxide for a total of 500ppm hydrogen peroxide/sodium chlorite mixture. Two drops of the cleaner were placed in the cul-de-sac of two normal human eyes. Upon instillation of the drops, the subjects closed their eyelids and pressed their index finger on the medial cantus, so as to block the puncta and stop the tears going into the lachrymal duct.

[0135] At 30 second, 1 minute, 2 minute, and 3 minute intervals, the subjects' tear samples were obtained by placing a fresh peroxide test strip in the cul-de-sac of the subjects' eyes. The used peroxide test strips were removed from the eye and left to dry at room temperature for 15 minutes. At the completion of the drying period, the level of hydrogen peroxide/sodium chlorite material left in the tear was estimated by comparing the color formed on the peroxide test strip to that of a standard color chart and recorded as shown below.

TABLE XI

Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes
Level of hydrogen peroxide/sodium chlorite	500ppm	>25ppm	10ppm	2ppm	0.5ppm

[0136] The data presented above shows a rapid reduction in the level of hydrogen peroxide/sodium chlorite in the tear film of the treated subjects. The placing of the index finger on the medial cantus blocks the puncta and does not allow the tears of the subjects to escape into the lachrymal duct. In addition, the closing off the eyelids stops the blinking process and thus stops the pumping action of the tear removal from the treated eyes. As such, it would appear that the rapid reduction in the level of hydrogen peroxide/sodium chlorite from the tears is not due to the loss of the tears of the subjects into the lachrymal duct. Rather, it is believed that the reduction is due to the presence of catalase and superoxide desmutase enzymes in the tears of human subjects. As the drops are placed in the eye of the patients, the catalase and other enzymes start the rapid enzymatic degradation of the hydrogen peroxide/sodium chlorite preparation, whereby in a matter of 3 minutes the level in the tears of the treated subjects is almost undetectable. The results of this experiment tend to show that upon instillation in the eye, the hydrogen peroxide/sodium chlorite mixture behaves like a self destructing

EP 2 990 044 A1

preservative with the end products being water, oxygen, and sodium chloride.

[0137] Additionally, the ocular symptoms of the treated eyes were observed and graded over a period of one hour post instillation for burning and stinging sensations, pain, redness, tearing, itching, photopsia, photophobia, discharge and for foreign body sensation as shown below.

5

10

15

20

25

30

35

40

45

50

55

5
10
15
20
25
30
35
40
45
50
55

TABLE XII

Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	15 minutes	30 minutes	60 minutes
Burning/ Stinging	0	0	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0	0	0
Foreign Body Sensation	0	+0.5	0	0	0	0	0	0	0	0

Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe

EP 2 990 044 A1

[0138] The above results show that the hydrogen peroxide/sodium chlorite mixture is very well tolerated by the human eye without presenting any signs of irritation, inflammation, or any other adverse effects.

Experiment 2: Artificial Tear Product

5

[0139] An artificial tear product containing 0.15g sodium hyaluronate, 0.50g protector, and 0.06g hydrogen peroxide/sodium chlorite mixture in 100mL sterile water was provided. The artificial tear product contained 400ppm sodium chlorite and 200ppm hydrogen peroxide for a total of 600ppm hydrogen peroxide/sodium chlorite mixture. Two drops of the cleaner were placed in the cul-de-sac of six normal human eyes. Upon instillation of the drops, the subjects closed their eyelids and pressed their index finger on the medial cantus, so as to block the puncta and stop the tears going into the lachrymal duct.

10

[0140] At zero second, 5 second, 20 second, 30 second, 60 second, 90 second, 120 second, and 180 second intervals, the subjects' tear samples were obtained by placing a fresh peroxide test strip in the cul-de-sac of the subjects' eyes. The used peroxide test strips were removed from the eye and left to dry at room temperature for 15 minutes. At the completion of the drying period, the level of hydrogen peroxide/sodium chlorite material left in the tear was estimated by comparing the color formed on the peroxide test strip to that of a standard color chart and recorded as shown below.

15

TABLE XIII

Time post instillation	Time 0	5 seconds	20 seconds	30 seconds	60 seconds	90 seconds	120 seconds	180 seconds
Subject 1			150ppm	60ppm	45ppm	38ppm		
Subject 2				75ppm	38ppm	38ppm	23ppm	
Subject 3					45ppm	30ppm	8ppm	15ppm
Subject 4			60ppm			15ppm	11ppm	15ppm
Subject 5			60ppm	23ppm				5ppm
Subject 6			150ppm	75ppm	30ppm		17ppm	
Average	600ppm	375ppm	105ppm	58ppm	40ppm	30ppm	15ppm	12ppm

20

25

30

[0141] The data presented above shows a rapid reduction in the level of hydrogen peroxide/sodium chlorite in the tear film of the treated subjects. The placing of the index finger on the medial cantus blocks the puncta and does not allow the tears of the subjects to escape into the lachrymal duct. In addition, the closing off the eyelids stops the blinking process and thus stops the pumping action of the tear removal from the treated eyes. As such, it would appear that the rapid reduction in the level of hydrogen peroxide/sodium chlorite from the tears is not due to the loss of the tears of the subjects into the lachrymal duct. Rather, it is believed that the reduction is due to the presence of catalase and superoxide desmutase enzymes in the tears of human subjects. As the drops are placed in the eye of the patients, the catalase and other enzymes start the rapid enzymatic degradation of the hydrogen peroxide/sodium chlorite preparation, whereby in a matter of 3 minutes the level in the tears of the treated subjects is almost undetectable. The results of this experiment tend to show that upon instillation in the eye, the hydrogen peroxide/sodium chlorite mixture behaves like a self destructing preservative with the end products being water, oxygen, and sodium chloride.

35

40

[0142] Additionally, the ocular symptoms of the treated eyes were observed and graded over a period of one hour post instillation for burning and stinging sensations, pain, redness, tearing, itching, photopsia, photophobia, discharge and for foreign body sensation as shown below.

45

50

55

5
10
15
20
25
30
35
40
45
50
55

TABLE XIV

Time post instillation	Zero time	30 seconds	1 minute	2 minutes	3 minutes	5 minutes	10 minutes	15 minutes	30 minutes	60 minutes
Burning/ Stinging	0	0	0	0	0	0	0	0	0	0
Pain	0	0	0	0	0	0	0	0	0	0
Redness	0	0	0	0	0	0	0	0	0	0
Tearing	0	0	0	0	0	0	0	0	0	0
Itching	0	0	0	0	0	0	0	0	0	0
Photopsia	0	0	0	0	0	0	0	0	0	0
Photophobia	0	0	0	0	0	0	0	0	0	0
Discharge	0	0	0	0	0	0	0	0	0	0
Foreign Body Sensation	0	+0.5	0	0	0	0	0	0	0	0

Grading Scale: 0 = None; +0.5 = Trace; +1 = Mild; +2 = Moderate; +3 = Moderately Severe; +4 = Severe

[0143] The above results show that the hydrogen peroxide/sodium chlorite mixture is very well tolerated by the human eye without presenting any signs of irritation, inflammation, or any other adverse effects.

[0144] It will be appreciated by those skilled in the art, that the invention has been described hereabove with reference to certain examples and specific embodiments. However, these are not the only examples and embodiments in which the invention may be practiced. Indeed, various modifications may be made to the above-described examples and embodiments without departing from the intended spirit and scope of the present invention. Accordingly, the present embodiments are to be considered on all respects as illustrative and not restrictive. It is intended that all such modifications be included within the scope of the following claims.

[0145] Preferred embodiments of the invention include:

1. An anti-microbial preservative for use in an ophthalmic product, the preservative comprising from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound, wherein the preservative does not generate chlorine dioxide, and wherein the preservative is at a pH range between about 6.0 and about 8.8.
2. The preservative of Claim 1 wherein the chlorite compound is a metal chlorite.
3. The preservative of Claim 2 wherein the metal is sodium.
4. The preservative of Claim 2 wherein the metal is selected from the group consisting of potassium, calcium, and magnesium.
5. The preservative of Claim 1 wherein the peroxy compound is hydrogen peroxide.
6. The preservative of Claim 1 wherein the ophthalmic product is an artificial tear solution.
7. The preservative of Claim 1 wherein the ophthalmic product is a contact lens disinfecting solution.
8. The preservative of Claim 1 wherein the ophthalmic product is a contact lens cleaning solution.
9. The preservative of Claim 8 wherein the cleaning solution is directly applied to a contact lens in place on an eye of a living being.
10. An antimicrobial ophthalmic composition for direct application onto an eye of a living being, the composition comprising from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH range between about 6.0 and 8.8.
11. The antimicrobial ophthalmic composition of Claim 10 wherein the chlorite compound is a metal chlorite.
12. The antimicrobial ophthalmic composition of Claim 11 wherein the metal is sodium.
13. The antimicrobial ophthalmic composition of Claim 11 wherein the metal is selected from the group consisting of potassium, calcium, and magnesium.
14. The antimicrobial ophthalmic composition of Claim 10 wherein the peroxy compound is hydrogen peroxide.
15. The antimicrobial ophthalmic composition of Claim 10 additionally comprising a lubricant chosen from the group consisting of non-ionic polymeric lubricants, anionic polymeric lubricants, and combinations thereof.
16. The antimicrobial ophthalmic composition of Claim 15 additionally comprising a block polymer based surfactant.
17. The antimicrobial ophthalmic composition of Claim 15 wherein the ophthalmic composition additionally comprises:

lubricant	0.05 wt. % to 0.3 wt. %;
boric acid	0.15 wt. % to 0.3 wt. %;
sodium chloride	0.50 wt. % to 0.8 wt. %;
surfactant	0.05 wt. % to 0.3 wt. %;
HCl or NaOH	to adjust pH; and
purified water	Q.S. to volume.

18. The antimicrobial ophthalmic composition of Claim 17 additionally comprising from about 0.001 wt. % to about 0.50 wt. % of sodium hyaluronate.

19. The antimicrobial ophthalmic composition of Claim 15 wherein the ophthalmic composition additionally comprises:

lubricant	0.05 wt. % to 0.30 wt. %;
sodium phosphate monobasic	0.10 wt. % to 0.25 wt. %;
sodium phosphate dibasic	0.10 wt. % to 0.40 wt. %;
sodium chloride	0.50 wt. % to 0.80 wt. %;
surfactant	0.05 wt. % to 0.30 wt. %;

(continued)

HCl or NaOH	to adjust pH; and
purified water	Q.S. to volume.

5

20. The antimicrobial ophthalmic composition of Claim 19 additionally comprising from about 0.001 wt. % to about 0.50 wt. % sodium hyaluronate.

21. The antimicrobial ophthalmic composition of Claim 10 wherein the composition is applied onto an eye for treating dryness of the eye.

10

22. The antimicrobial ophthalmic composition of Claim 10 wherein the composition is applied onto an eye for treating an infection of an eye.

23. The antimicrobial ophthalmic composition of Claim 22 wherein the infection is caused by bacterial keratitis.

24. The antimicrobial ophthalmic composition of Claim 22 wherein the infection is caused by a virus.

25. The antimicrobial ophthalmic composition of Claim 22 wherein the infection is caused by a fungus.

15

26. The antimicrobial ophthalmic composition of Claim 10 wherein the composition is applied onto an eye for cleansing a contact lens in place on the eye.

27. A method of treating dryness of a eye of a living being, the method comprising applying an antimicrobial ophthalmic composition onto the eye, the composition comprising from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH range between about 6.0 and 8.8.

20

28. The method of Claim 27 wherein the chlorite compound of the ophthalmic composition is a metal chlorite.

29. The method of Claim 28 wherein the metal is sodium.

30. The method of Claim 28 wherein the metal is selected from the group consisting of potassium, calcium, and magnesium.

25

31. The method of Claim 27 wherein the peroxy compound of the ophthalmic composition is hydrogen peroxide.

32. The method of Claim 27 wherein the ophthalmic composition additionally comprises a lubricant chosen from the group consisting of non-ionic polymeric lubricants, anionic polymeric lubricants, and combinations thereof.

33. The method of Claim 32 wherein the ophthalmic composition additionally comprises a block polymer based surfactant.

34. The method of Claim 33 wherein the ophthalmic composition additionally comprises:

30

lubricant	0.05 wt. % to 0.3 wt. %;
boric acid	0.15 wt. % to 0.3 wt. %;
sodium chloride	0.50 wt. % to 0.8 wt. %;
surfactant	0.05 wt. % to 0.3 wt. %;
HCl or NaOH	to adjust pH; and
purified water	Q.S. to volume.

35

40

35. The method of Claim 34 wherein the ophthalmic composition additionally comprises from about 0.001 wt. % to about 0.50 wt. % sodium hyaluronate.

36. The method of Claim 33 wherein the ophthalmic composition additionally comprises:

45

lubricant	0.05 wt. % to 0.30 wt. %;
sodium phosphate monobasic	0.10 wt. % to 0.25 wt. %;
sodium phosphate dibasic	0.10 wt. % to 0.40 wt. %;
sodium chloride	0.50 wt. % to 0.80 wt. %;
surfactant	0.05 wt. % to 0.30 wt. %;
HCl or NaOH	to adjust pH; and
purified water	Q.S. to volume.

50

55

37. The method of Claim 36 wherein the ophthalmic composition additionally comprises from about 0.001 wt. % to about 0.50 wt. % sodium hyaluronate.

38. A method of cleansing a contact lens in place upon an eye of a living being, the method comprising applying an antimicrobial ophthalmic composition onto the lens, the composition comprising from about 0.005 wt. % to about 0.20

EP 2 990 044 A1

wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH range between about 6.0 and 8.8.

39. The method of Claim 38 wherein the chlorite compound of the ophthalmic composition is a metal chlorite.

40. The method of Claim 39 wherein the metal is sodium.

5 41. The method of Claim 39 wherein the metal is selected from the group consisting of potassium, calcium, and magnesium.

42. The method of Claim 38 wherein the peroxy compound of the ophthalmic composition is hydrogen peroxide.

43. The method of Claim 38 wherein the ophthalmic composition additionally comprises a lubricant chosen from the group consisting of non-ionic polymeric lubricants, anionic polymeric lubricants, and combinations thereof.

10 44. The method of Claim 38 wherein the ophthalmic composition additionally comprises a sustained delivery component which limits the rate at which the chlorite and hydrogen peroxide become available for generation of oxygen.

45. The method of Claim 44 wherein the sustained delivery component is a polymeric matrix.

46. The method of Claim 44 wherein the sustained delivery component is a liposome.

47. The method of Claim 44 wherein the sustained delivery component is an ointment.

15 48. The method of Claim 44 wherein the sustained delivery component is selected from the group consisting of a cellulose ester, hydroxymethylpropyl cellulose, methylhydroxyethyl cellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, carboxymethyl cellulose, a salt of a cellulose ester, cellulose acetate, hydroxypropylmethyl cellulose phthalate, methacrylic acid-ethyl acetate copolymer, polyvinylpyrrolidone, polyvinyl alcohol, hyaluronic acid, a phospholipid, phospholipids having a neutral charge, phospholipids having a negative charge, dipalmytoyl phosphatidyl choline, dipalmytoyl phosphatidyl serine, and sodium salts thereof.

20 49. The method of Claim 44 wherein the sustained delivery component comprises 1 percent to 20 percent by weight of the preparation.

50. The method of Claim 47 wherein the ophthalmic composition comprises:

25	sodium chlorite	0.005 wt. % to 0.200 wt. %;
	hydrogen peroxide	0.005 wt. % to 0.050 wt. %;
	HPMC	0.05 wt. % to 0.20 wt. %;
30	sodium phosphate monobasic monohydrate	0.01 wt. % to 0.30 wt. %;
	HCl or NaOH	to adjust to pH 7.4
35	purified water	Q.S. to volume.

51. The method of Claim 38 wherein the ophthalmic composition additionally comprises a block polymer based surfactant.

52. The method of Claim 51 wherein the ophthalmic composition additionally comprises:

40	lubricant	0.05 wt. % to 0.30 wt. %;
	boric acid	0.15 wt. % to 0.30 wt. %;
	sodium chloride	0.50 wt. % to 0.80 wt. %;
45	surfactant	0.05 wt. % to 0.30 wt. %;
	HCl or NaOH	to adjust pH; and
	purified water	Q.S. to volume.

50 53. The method of Claim 52 wherein the ophthalmic composition additionally comprises from about 0.001 wt. % to about 0.50 wt. % sodium hyaluronate.

54. The method of Claim 51 wherein the ophthalmic composition additionally comprises:

55	lubricant	0.05 wt. % to 0.30 wt. %;
	sodium phosphate monobasic	0.10 wt. % to 0.25 wt. %;
	sodium phosphate dibasic	0.10 wt. % to 0.40 wt. %;

(continued)

5	sodium chloride	0.50 wt. % to 0.80 wt. %;
	surfactant	0.05 wt. % to 0.30 wt. %;
	HCl or NaOH	to adjust pH; and
	purified water	Q.S. to volume.

55. The method of Claim 54 wherein the ophthalmic composition additionally comprises from about 0.001 wt. % to about 0.50 wt. % sodium hyaluronate.

10 56. An antimicrobial preservative for use in a dermatology product, the preservative comprising from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about .005 wt. % to about 0.05 wt. % peroxy compound, wherein the preservative does not generate chlorine dioxide, and wherein the preservative is at a pH range between about 5.0 and about 8.8.

15 57. The preservative of Claim 56 wherein the chlorite compound is a metal chlorite.

58. The preservative of Claim 57 wherein the metal is sodium.

59. The preservative of Claim 57 wherein the metal is selected from the group consisting of potassium, calcium, and magnesium.

60. The preservative of Claim 56 wherein the peroxy compound is hydrogen peroxide.

61. The preservative of Claim 56 wherein the dermatology product is Decubitus gel or a cream.

20 62. A method of treating a dermatological condition, the method comprising applying an antimicrobial dermatological composition onto the affected skin area, the composition comprising from about 0.005 wt. % to about 0.20 wt. % chlorite compound and from about 0.005 wt. % to about 0.05 wt. % peroxy compound, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH range between about 5.0 and 8.8.

63. The method of Claim 62 wherein the chlorite compound of the dermatological composition is a metal chlorite.

25 64. The method of Claim 63 wherein the metal is sodium.

65. The method of Claim 63 wherein the metal is selected from a group consisting of potassium, calcium, and magnesium.

66. The method of Claim 62 wherein the peroxy compound of the dermatological composition is hydrogen peroxide.

67. The method of Claim 62 wherein the dermatological composition is Decubitus gel or a cream.

30 68. The method of Claim 67 wherein the dermatological composition additionally comprises:

35	lubricant	0.05 wt. % to 0.30 wt. %;
	boric acid	0.15 wt. % to 0.30 wt. %;
	sodium chloride	0.50 wt. % to 0.80 wt. %;
	surfactant	0.05 wt. % to 0.30 wt. %;
	HCl or NaOH	to adjust pH; and
	purified water	Q.S. to volume.

40 69. The method of Claim 67 wherein the dermatological composition additionally comprises:

45	lubricant	0.05 wt. % to 0.30 wt. %;
	sodium phosphate monobasic	0.10 wt. % to 0.25 wt. %;
	sodium phosphate dibasic	0.10 wt. % to 0.40 wt. %;
	sodium chloride	0.50 wt. % to 0.80 wt. %;
	surfactant	0.05 wt. % to 0.30 wt. %;
50	HCl or NaOH	to adjust pH; and
	purified water	Q.S. to volume.

70. The method of Claim 62 wherein the dermatological composition additionally comprises a sustained delivery component which limits the rate at which the chlorite and hydrogen peroxide become available for generation of oxygen.

55 71. The method of Claim 70 wherein the sustained delivery component comprises a polymeric matrix.

72. The method of Claim 70 wherein the sustained delivery component comprises a liposome.

73. The method of Claim 70 wherein the sustained delivery component comprises an ointment.

74. The method of Claim 70 wherein the sustained delivery component is selected from the group consisting of a cellulose ester, hydroxymethylpropyl cellulose, methylhydroxyethyl cellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, carboxymethyl cellulose, a salt of a cellulose ester, cellulose acetate, hydroxypropylmethyl cellulose phthalate, methacrylic acid-ethyl acetate copolymer, polyvinylpyrrolidone, polyvinyl alcohol, hyaluronic acid, a phospholipid, phospholipids having a neutral charge, phospholipids having a negative charge, dipalmytoyl phosphatidyl choline, dipalmytoyl phosphatidyl serine, and sodium salts thereof.

75. The method of Claim 70 wherein the sustained delivery component comprises 1 percent to 20 percent by weight of the preparation.

76. The method of Claim 73 wherein the ophthalmic composition comprises:

10	sodium chlorite	0.005 wt. % to 0.200 wt. %;
	hydrogen peroxide	0.005 wt. % to 0.050 wt. %;
15	HPMC	0.05 wt. % to 0.20 wt. %;
	sodium phosphate monobasic monohydrate	0.01 wt. % to 0.30 wt. %;
20	HCl or NaOH	to adjust to pH 7.4
	purified water	Q.S. to volume.

Claims

- 25
1. An anti-microbial preservative for use in an ophthalmic product that is well tolerated by the human eye, the preservative comprising 0.05 wt. % sodium chlorite, 0.02 wt. %, 0.03 wt. % or 0.05 wt. % hydrogen peroxide, and 0.15 wt. % boric acid, wherein the preservative does not generate chlorine dioxide and wherein the preservative is at a pH of 7.4.
 - 30
 2. The preservative of Claim 1 wherein the ophthalmic product is an artificial tear solution.
 3. The preservative of Claim 1 wherein the ophthalmic product is a contact lens disinfecting solution.
 - 35
 4. The preservative of Claim 1 wherein the ophthalmic product is a contact lens cleaning solution.
 5. The preservative of Claim 4 wherein the cleaning solution is directly applied to a contact lens in place on an eye of a living being.
 - 40
 6. An antimicrobial ophthalmic composition for direct application onto an eye of a living being that is well tolerated by the human eye, the composition comprising 0.05 wt. % sodium chlorite, 0.02 wt. %, 0.03 wt. % or 0.05 wt. % hydrogen peroxide, and 0.15 wt. % boric acid, wherein the preservative does not generate chlorine dioxide and wherein the preservative is at a pH of 7.4.
 - 45
 7. The antimicrobial ophthalmic composition of Claim 6 wherein the composition is applied onto an eye for treating dryness of the eye.
 8. The antimicrobial ophthalmic composition of Claim 6 wherein the composition is applied onto an eye for treating an infection of an eye.
 - 50
 9. The antimicrobial ophthalmic composition of Claim 8 wherein the infection is caused by bacterial keratitis.
 10. The antimicrobial ophthalmic composition of Claim 8 wherein the infection is caused by a virus.
 - 55
 11. The antimicrobial ophthalmic composition of Claim 8 wherein the infection is caused by a fungus.
 12. The antimicrobial ophthalmic composition of Claim 6 wherein the composition is applied onto an eye for cleansing a contact lens in place on the eye.

EP 2 990 044 A1

13. Use of an antimicrobial ophthalmic composition being well tolerated by the human eye and comprising 0.05 wt. % sodium chlorite, 0.02 wt. %, 0.03 wt. % or 0.05 wt. % hydrogen peroxide, and 0.15 wt. % boric acid, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH of 7.4 for the manufacture of a medicament for treating dryness of an eye on a living being.

5

14. Use of an antimicrobial ophthalmic composition being well tolerated by the human eye and comprising 0.05 wt. % sodium chlorite, 0.02 wt. %, 0.03 wt. % or 0.05 wt. % hydrogen peroxide, and 0.15 wt. % boric acid, wherein the composition does not generate chlorine dioxide, and wherein the composition is at a pH of 7.4 for the manufacture of a medicament for cleansing and disinfecting a contact lens in place upon an eye of a living being.

10

15

20

25

30

35

40

45

50

55




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7



EUROPEAN SEARCH REPORT

Application Number
EP 15 18 6300

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WO 03/009802 A (KARAGOEZIAN HAMPAR L [US]) 6 February 2003 (2003-02-06) * paragraph [0001] * * claims 1-42 * -----	1-14	INV. A61K33/40 A61K33/14 A61K33/22 A61P17/00 A01N39/00
X	WO 00/19981 A (KARAGOEZIAN HAMPAR L [US]) 13 April 2000 (2000-04-13) * paragraph [0002] * * page 13 * * claims 1-44 *	1-14	
A	CA 1 224 415 A1 (BERGER PETER) 21 July 1987 (1987-07-21) * the whole document * -----	1-14	
X	US 2006/127497 A1 (KARAGOEZIAN HAMPAR L [US]) 15 June 2006 (2006-06-15) * claim 1 * * paragraph [0086] * * paragraphs [0057], [0058] * -----	1-14	
			TECHNICAL FIELDS SEARCHED (IPC)
			A61K A01N A61P
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 21 January 2016	Examiner Albayrak, Timur
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1508 03/02 (P04CO1)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 15 18 6300

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-01-2016

10

15

20

25

30

35

40

45

50

55

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 03009802 A	06-02-2003	AT 402712 T	15-08-2008
		AU 2002322298 A1	17-02-2003
		EP 1418881 A2	19-05-2004
		ES 2310601 T3	16-01-2009
		JP 5002114 B2	15-08-2012
		JP 2004536137 A	02-12-2004
		MX PA04000764 A	17-02-2005
		US 2002064565 A1	30-05-2002
		WO 03009802 A2	06-02-2003

WO 0019981 A	13-04-2000	AT 289804 T	15-03-2005
		AU 6416999 A	26-04-2000
		DE 69923987 D1	07-04-2005
		DE 69923987 T2	02-11-2006
		EP 1119347 A1	01-08-2001
		US 6488965 B1	03-12-2002
		US 2015056272 A1	26-02-2015
		WO 0019981 A1	13-04-2000

CA 1224415 A1	21-07-1987	-----	-----
US 2006127497 A1	15-06-2006	CN 1845747 A	11-10-2006
		EP 1641472 A2	05-04-2006
		JP 2007527390 A	27-09-2007
		MX PA06000257 A	03-07-2006
		US 2004037891 A1	26-02-2004
		US 2006127497 A1	15-06-2006
		WO 2005007174 A2	27-01-2005

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 10614646 B [0001]
- US 09911638 B [0001]
- US 09412174 B [0001]
- US 4997616 A [0006]
- US 5279673 A [0006]
- US 5246662 A [0006]
- US 4855135 A [0008]
- US RE31779 E [0010]
- US 4084747 A [0010]
- US 5384134 A, Kross [0010]
- US 2701781 A, de Guevara [0011]
- US 4317814 A, Laso [0012]
- US 3271242 A, McNicholas [0013]

Non-patent literature cited in the description

- **PAUGH, J.R. ; BRENNAN, N.A. ; EFRON, N.** Ocular Response to Hydrogen Peroxide. *Am. J. Optom. Physiol. Opt.*, February 1988, vol. 65 (2), 91-8 [0131]

摘要

本发明提供了用于眼科产品和皮肤病学产品的抗微生物防腐剂。所述防腐剂包括大约 0.005 重量%-大约 0.20 重量%的亚硝酸盐化合物和大约 0.005 重量%-大约 0.05 重量%的过氧化物。另外,所述防腐剂在 5.0-8.8 的 PH 范围内不生成二氧化氯。还包括用于直接施用在生物的眼睛或皮肤上的抗微生物眼科和皮肤病学组合物,所述组合物包括大约 0.005 重量%-大约 0.20 重量%的亚硝酸盐化合物和大约 0.005 重量%-大约 0.05 重量%的过氧化物。还包括用于治疗眼睛干燥和皮肤病(例如,创伤、烧伤、感染、溃疡、银屑病等)的方法和用于通过对眼睛或对隐形眼镜施用所述组合物在隐形眼镜保持在眼睛上的情况下对隐形眼镜进行消毒和清洗的方法。