

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 2970896 T3**

-
- (51) Int.Cl.: **G 01 N 33/50 (2006.01)** **A 61 P 1/16 (2006.01)** **C 12 N 5/071 (2010.01)**
C 12 N 5/077 (2010.01)
- (45) Oversættelsen bekendtgjort den: **2024-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-04-24**
- (86) Europæisk ansøgning nr.: **14769577.9**
- (86) Europæisk indleveringsdag: **2014-03-13**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-01-20**
- (86) International ansøgning nr.: **US2014026679**
- (87) Internationalt publikationsnr.: **WO2014151921**
- (30) Prioritet: **2013-03-15 US 201313841430**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Organovo, Inc., 6275 Nancy Ridge Drive , Suite 110, San Diego, CA 92121, USA**
- (72) Opfinder: **ROBBINS, Justin, B., 4655 Ohio Street, Unit 12, San Diego, CA 92116, USA**
GORGEN, Vivian, A., 3567 Caminito Carmel Lndg, San Diego, CA 92130, USA
PRESNELL, Sharon, C., 16224 Rostrata Hill Rd., Poway, CA 92064, USA
SHEPHERD, Benjamin, R., 2/55 Dobson Street, Ferntree Gully, VIC 3156, Australien
- (74) Fuldmægtig i Danmark: **Marks & Clerk LLP, 44, rue de la Vallée , L-2661 Luxembourg, Luxembourg**
- (54) Benævnelse: **MANIPULEREDE LEVERVÆV, ARRAYS DERAFT OG FREMGANGSMÅDER TIL FREMSTILLING HERAF**
- (56) Fremdragne publikationer:
WO-A2-2012/003465
WO-A2-2013/040078
US-A1- 2003 096 411
US-A1- 2011 250 688
US-A1- 2012 045 770
US-A1- 2012 116 568
Anonymous: "Organovo describes first fully cellular 3D bioprinted liver tissue", Organovo , 22 April 2013 (2013-04-22), pages 1-2, XP002753682, Retrieved from the Internet: URL:[http://ir.organovo.com/news/press-releases/press-releases-details/2013/Organovo-Describes-First-Fully-Cellular-3D-Bioprint ed-Liver-Tissue/](http://ir.organovo.com/news/press-releases/press-releases-details/2013/Organovo-Describes-First-Fully-Cellular-3D-Bioprint-ed-Liver-Tissue/) [retrieved on 2016-02-01]
SEAN V MURPHY ET AL: "3D bioprinting of tissues and organs", NATURE BIOTECHNOLOGY, vol. 32, no. 8, 5 August 2014 (2014-08-05) , pages 773-785, XP055244641, US ISSN: 1087-0156, DOI: 10.1038/nbt.2958
KAZUO OHASHI ET AL: "Engineering functional two- and three-dimensional liver systems in vivo using hepatic tissue sheets", NATURE MEDICINE., vol. 13, no. 7, 17 June 2007 (2007-06-17), pages 880-885, XP055246201, US

Fortsættes ...

ISSN: 1078-8956, DOI: 10.1038/nm1576

NOROTTE C ET AL: "Scaffold-free vascular tissue engineering using bioprinting", BIOMATERIALS, ELSEVIER SCIENCE PUBLISHERS BV., BARKING, GB, vol. 30, no. 30, 1 October 2009 (2009-10-01), pages 5910-5917, XP026524655, ISSN: 0142-9612, DOI: 10.1016/J.BIOMATERIALS.2009.06.034 [retrieved on 2009-08-06]

SAKAI Y ET AL: "Toward engineering of vascularized three-dimensional liver tissue equivalents possessing a clinically significant mass", BIOCHEMICAL ENGINEERING JOURNAL, ELSEVIER, AMSTERDAM, NL, vol. 48, no. 3, 15 February 2010 (2010-02-15), pages 348-361, XP026886642, ISSN: 1369-703X, DOI: 10.1016/J.BEJ.2009.10.010 [retrieved on 2009-10-21]

SHIM, JIN-HYUNG ET AL.: 'Bioprinting of a mechanically enhanced three-dimensional dual cell -laden construct for osteochondral tissue engineering using a multi-head tissue/organ building system' JOURNAL OF MICROMECHANICS AND MICROENGINEERING vol. 22, no. 085014, 2012, pages 1 - 11, XP020227406

DESCRIPTION

Description

CROSS-REFERENCE TO RELATED APPLICATIONS

BACKGROUND OF THE INVENTION

[0001] A number of pressing problems confront the healthcare industry. As of June 2012 there were 114,636 patients registered by United Network for Organ Sharing (UNOS) as needing an organ transplant. According to UNOS, between January and March 2012 only 6,838 transplants were performed. Each year more patients are added to the UNOS list than transplants are performed, resulting in a net increase in the number of patients waiting for a transplant.

[0002] Additionally, the research and development cost of a new pharmaceutical compound is approximately \$1.8 billion. See Paul, et al. (2010). How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nature Reviews Drug Discovery* 9(3):203-214. Drug discovery is the process by which drugs are discovered and/or designed. The process of drug discovery generally involves at least the steps of: identification of candidates, synthesis, characterization, screening, and assays for therapeutic efficacy. Despite advances in technology and understanding of biological systems, drug discovery is still a lengthy, expensive, and inefficient process with low rate of new therapeutic discovery.

WO2013/0040078 ORGANOVO INC discloses living, three-dimensional tissue constructs for in vitro scientific and medical research, arrays thereof, and methods of making said tissues and arrays.

US2003/096411 A1 MICHALOPOULOS GEORGE et al., discloses methods and compositions for ex vivo culturing of hepatocytes and nonparenchymal cells on a matrix coated with a molecule that promotes cell adhesion, proliferation or survival, in the presence of growth factors. It is stated that the hepatic cell culture system can be used to form bio-artificial livers or may be implanted into the body of a recipient host having a hepatic disorder, including, for example, cirrhosis of the liver, induced hepatitis, chronic hepatitis, primary sclerosing cholangitis and alpha1 antitrypsin deficiency.

SUMMARY OF THE INVENTION

[0003] The invention relates to the field of regenerative medicine and tissue and/or organ

engineering. More particularly, the invention relates to engineered liver tissue constructs, arrays thereof, and methods of fabrication.

[0004] In a first aspect of the invention there is provided, an engineered, living, three-dimensional liver tissue construct, the construct comprising at least one compartment defining a planar geometry, the compartment comprising an interior defined by a border, the interior comprising parenchymal cells, the border comprising non-parenchymal cells; the cells cohered to form a living, three-dimensional liver tissue construct; provided that at least one component of the construct was bioprinted and the construct is substantially free of pre-formed scaffold at the time of use; wherein the parenchymal cells are hepatocytes or hepatocyte-like cells derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver; and wherein the non-parenchymal cells comprise one or more of: vascular cells, endothelial cells, fibroblasts, mesenchymal cells, immune cells, Kupffer cells, stellate cells, biliary epithelial cells, biliary epithelial-like cells, and sinusoidal endothelial cells.

[0005] In one embodiment there is provided an engineered, living, three-dimensional liver tissue construct according to the first aspect, comprising one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form the living, three-dimensional liver tissue construct, the construct characterized by having at least one of:

1. a. at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry, wherein the cell types comprise parenchymal and non-parenchymal cells according to claim 1; and
2. b. a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry.

[0006] In one embodiment there is provided an engineered, living, three-dimensional liver tissue construct according to the first aspect, comprising a plurality of layers; preferably wherein at least one layer is compositionally or architecturally distinct from at least one other layer to create a laminar geometry.

[0007] In certain embodiments, the engineered, living, three-dimensional liver tissue construct further comprises cancer cells.

[0008] In certain embodiments, the construct is for use in the augmentation of one or more liver functions in humans.

[0009] In a second aspect of the invention there is provided an array comprising a plurality of the engineered, living, three-dimensional liver tissue constructs of the first aspect for use in *in vitro* assays.

[0010] In certain embodiments, the *in vitro* assay is one or more of: drug discovery; drug testing; preclinical research; toxicity testing; absorption, distribution, metabolism, and excretion testing (ADME); drug metabolism and pharmacokinetics testing (DMPK); disease modeling; infectious disease modeling; host disease modeling; three-dimensional biology studies; and cell-based screening.

[0011] In a third aspect of the invention there is provided a method of fabricating a living, three-dimensional liver tissue construct comprising:

1. a. providing one or more bio-inks comprising non-parenchymal cells;
2. b. providing one or more bio-inks comprising parenchymal cells;
3. c. depositing the bio-inks onto a support; and
4. d. incubating the deposited bio-inks for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct comprising at least one compartment, the compartment comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells;

wherein the non-parenchymal cells comprise one or more of: vascular cells, endothelial cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary epithelial cells, biliary epithelial-like cells, and sinusoidal endothelial cells; and wherein the parenchymal cells are hepatocytes or hepatocyte-like cells derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver.

[0012] In certain embodiments, at least one component of the construct is bioprinted; optionally wherein the construct is substantially free of any pre-formed scaffold at the time of use; preferably wherein the construct is non-innervated.

[0013] In certain embodiments, there is provided an engineered, living, three-dimensional liver tissue construct according to the first aspect, provided that the living, three-dimensional liver tissue construct remains substantially viable *in vitro* culture for at least 7 days and has at least two of the following liver characteristics:

1. a. synthesizes cholesterol;
2. b. stores lipid;
3. c. stores glycogen;
4. d. expresses liver-specific proteins, including two or more of: albumin, fibrinogen, transferrin, CYP450, alpha-1-antitrypsin, ornithine transcarbamylase, glycogen synthase, Factor VIII, and Factor IX;
5. e. responsive to liver modulators, including one or more of: hormones, inducers, toxicants, drugs, antibodies, interfering RNAs, small RNAs, locked nucleic acids, viruses, bacteria, parasites, and inflammatory inducers;

- 6. f. responsive to liver toxicants, including one or more of: acetaminophen, alcohol, trovafloxacin, methotrexate, diclofenac, troglitazone, valproic acid, and amiodarone;
- 7. g. expresses transporter proteins, including one or more of: BSEP, OATP1B 1, NTCP, PGP, BCRP, OATP1B3, ABCB4, ATP8B1, Multi-Drug Resistance Transporter (MDR);
- 8. h. maintains at least some quiescent stellate cells after fabrication; and
- 9. i. includes microvascular structures.

[0014] In certain embodiments, the construct remains substantially viable *in vitro* culture for at least 14 days, preferably for at least 28 days; more preferably for at least 40 days.

[0015] In certain embodiments, the construct comprises cells derived from induced pluripotent stem cells (iPSC).

[0016] In certain embodiments, the construct comprises myeloid cells derived from monocytes or macrophages.

[0017] Disclosed herein but not claimed are engineered, living, three-dimensional liver tissue constructs, the construct comprising at least one compartment defining a planar geometry, the compartment comprising an interior defined by a border, the interior comprising parenchymal cells, the border comprising non-parenchymal cells; the cells cohered to form a living, three-dimensional liver tissue construct; provided that at least one component of the construct was bioprinted and the construct is substantially free of pre-formed scaffold at the time of use.

[0018] The constructs may further comprise an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting.

[0019] In some examples not claimed, the construct comprises stem/progenitor cells that were exposed to one or more differentiation signals. In some examples not claimed, the differentiation signal comprises one or more of: biomechanical signals, soluble signals, and physical signals. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals before fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals during fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals after fabrication of the construct.

[0020] In some examples not claimed, the construct is for use in *in vitro* assays. In some examples not claimed, the construct is for use in one or more of: drug discovery; drug testing; preclinical research; toxicity testing; absorption, distribution, metabolism, and excretion testing (ADME); drug metabolism and pharmacokinetics testing (DMPK); disease modeling; infectious disease modeling; host disease modeling; three-dimensional biology studies; and cell-based screening. In some examples not claimed, the construct is for use in cell-based screening, wherein the screening is for one or more infectious diseases, liver fibrosis (e.g., cirrhosis), liver

cancer, liver steatosis (e.g., fatty liver), one or more metabolic deficiencies, or one or more protein deficiencies. In some examples not claimed, the infectious diseases include viral infection or parasitic infection (e.g., plasmodium infection, etc.). In some examples not claimed, the construct is for use in long-term tissue toxicity studies, wherein analyses are conducted for periods >3 days and up to 6 months. In some examples not claimed, the construct is for use in the augmentation of one or more liver functions in humans. In some examples not claimed, the construct is for implantation in a subject at a site of injury, disease, or degeneration. In some examples not claimed, the construct is for clinical use in extracorporeal devices designed to augment or restore one or more liver functions. In some embodiments, the construct is non-innervated.

[0021] Also disclosed herein but not claimed are engineered, living, three-dimensional liver tissue constructs comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct, the construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry. In some examples not claimed, at least one component of the construct was bioprinted. In some embodiments, each layer of the engineered liver tissue comprises multiple cells in the X, Y, and Z axes. In some examples not claimed, each layer of the engineered liver tissue is at least about 50 microns in thickness in the X, Y, and Z axes. In some examples not claimed, the construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, the construct is substantially free of pre-formed scaffold at the time of use. In some examples not claimed, the liver cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, the construct further comprises one or more of the following cell types: vascular cells, endothelial cells, parenchymal cells, non-parenchymal cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the construct comprises stem/progenitor cells that were exposed to one or more differentiation signals. In some examples not claimed, the differentiation signal comprises one or more of: biomechanical signals, soluble signals, and physical signals. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals before fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals during fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals after fabrication of the construct. In some examples not claimed, the construct is for use in *in vitro* assays. In some examples not claimed the construct is for use in one or more of: drug discovery; drug testing; preclinical research; toxicity testing; absorption, distribution, metabolism, and excretion testing (ADME); drug metabolism and pharmacokinetics testing

(DMPK); disease modeling; infectious disease modeling; host disease modeling; three-dimensional biology studies; and cell-based screening. In some examples not claimed, the construct is for use in cell-based screening, wherein the screening is for one or more infectious diseases, liver fibrosis (e.g., cirrhosis), liver cancer, liver steatosis (e.g., fatty liver), one or more metabolic deficiencies, or one or more protein deficiencies.

[0022] In some examples not claimed, the infectious diseases include viral infection or parasitic infection (e.g., plasmodium infection, etc.). In some examples not claimed, the construct is for use in long-term tissue toxicity studies, wherein analyses are conducted for periods >3 days and up to 6 months. In some examples not claimed, the construct is for use in the augmentation of one or more liver functions in humans. In some examples not claimed, the construct is for implantation in a subject at a site of injury, disease, or degeneration. In some examples not claimed, the construct is for clinical use in extracorporeal devices designed to augment or restore one or more liver functions. In some examples not claimed, the construct is non-innervated.

[0023] Also disclosed herein but not claimed are engineered, living, three-dimensional liver tissue constructs comprising: a plurality of layers, each layer comprising cylindrical bio-ink, the bio-ink axially-aligned substantially in parallel, the bio-ink comprising parenchymal liver cells; and optionally, non-parenchymal cells within or among the cylindrical bio-ink; and optionally, void spaces among the cylindrical bio-ink. In some examples not claimed, at least one component of the construct was bioprinted. In some examples not claimed, the construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, the parenchymal cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, the non-parenchymal cells comprise one or more of: vascular cells, endothelial cells, fibroblasts, mesenchymal cells, immune cells, Kupffer cells, stellate cells, biliary epithelial cells, biliary epithelial-like cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the construct comprises one or more layers. In some examples not claimed, the construct comprises a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry. In some examples not claimed, the construct is for use in the augmentation of one or more liver functions in humans. In some examples not claimed, the construct is for implantation in a subject at a site of injury, disease, or degeneration. In some examples not claimed, the construct is for clinical use in extracorporeal devices designed to augment or restore one or more liver functions. In some examples not claimed, the construct is non-innervated.

[0024] Also disclosed herein but not claimed are arrays of engineered living, three-dimensional liver tissue constructs, each construct comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional

liver tissue construct, each construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry. In some examples not claimed, at least one component of each construct was bioprinted. In some examples not claimed, each construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, each construct is substantially free of pre-formed scaffold at the time of use. In some embodiments, the liver cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, each construct comprises stem/progenitor cells that were exposed to one or more differentiation signals. In some examples not claimed, the differentiation signal comprises one or more of: biomechanical signals, soluble signals, and physical signals. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals before fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals during fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals after fabrication of the construct. In some examples not claimed, each construct further comprises one or more of the following cell types: vascular cells, endothelial cells, parenchymal cells, non-parenchymal cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the array is for use in *in vitro* assays. In some examples not claimed, the array is for use in one or more of: drug discovery; drug testing; preclinical research; toxicity testing; absorption, distribution, metabolism, and excretion testing (ADME); drug metabolism and pharmacokinetics testing (DMPK); disease modeling; infectious disease modeling; host disease modeling; three-dimensional biology studies; and cell-based screening. In some examples not claimed, the array is for use in cell-based screening, wherein the screening is for one or more infectious diseases, liver fibrosis (e.g., cirrhosis), liver cancer, liver steatosis (e.g., fatty liver), one or more metabolic deficiencies, or one or more protein deficiencies. In some examples not claimed, the infectious diseases include viral infection or parasitic infection (e.g., plasmodium infection, etc.). In some examples not claimed, the array is for use in long-term tissue toxicity studies, wherein analyses are conducted for periods >3 days and up to 6 months.

[0025] Also disclosed herein but not claimed are arrays of engineered living, three-dimensional tissue constructs, wherein at least one of the constructs is a liver tissue construct, each liver tissue construct comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct, the liver tissue construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally

distinct from at least one other layer to create a laminar geometry. In some examples not claimed, at least one component of each liver tissue construct was bioprinted. In some examples not claimed, each liver tissue construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, each liver construct was substantially free of pre-formed scaffold at the time of use. In some examples not claimed, the liver cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells; induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, each liver tissue construct comprises stem/progenitor cells that were exposed to one or more differentiation signals. In some examples not claimed, the differentiation signal comprises one or more of: biomechanical signals, soluble signals, and physical signals. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals before fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals during fabrication of the construct. In some examples not claimed, the stem/progenitor cells were exposed to one or more differentiation signals after fabrication of the construct. In some examples not claimed, each liver tissue construct further comprises one or more of the following cell types: vascular cells, endothelial cells, parenchymal cells, non-parenchymal cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the array is for use in *in vitro* assays. In some examples not claimed, the array is for use in one or more of the following: tissue-tissue interactions; drug discovery; drug testing; preclinical research; toxicity testing; absorption, distribution, metabolism, and excretion testing (ADME); drug metabolism and pharmacokinetics testing (DMPK); disease modeling; infectious disease modeling; host disease modeling; three-dimensional biology studies; and cell-based screening. In some examples not claimed, the array is for use in cell-based screening, wherein the screening is for one or more infectious diseases, liver fibrosis (e.g., cirrhosis), liver cancer, liver steatosis (e.g., fatty liver), one or more metabolic deficiencies, or one or more protein deficiencies. In some examples not claimed, the infectious diseases include viral infection or parasitic infection (e.g., plasmodium infection, etc.). In some embodiments, the array is for use in long-term tissue toxicity studies, wherein analyses are conducted for periods >3 days and up to 6 months.

[0026] Also disclosed herein but not claimed are methods of fabricating a living, three-dimensional liver tissue construct comprising: preparing one or more bio-inks comprising non-parenchymal cells; preparing one or more bio-inks comprising parenchymal cells; depositing the bio-inks onto a support; and incubating the deposited bio-inks for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct comprising at least one compartment, the compartment comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells. In some examples not claimed, the non-parenchymal cells comprise one or more of: vascular cells, endothelial cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary epithelial

cells, biliary epithelial-like cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the parenchymal cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, at least one component of the construct is bioprinted. In some examples not claimed, the construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, the construct is substantially free of any pre-formed scaffold at the time of use. In some embodiments, the construct is non-innervated.

[0027] Also disclosed herein but not claimed are methods of fabricating an engineered living, three-dimensional liver tissue construct comprising: preparing one or more bio-inks comprising liver cells; depositing the one or more bio-inks onto a support; and incubating the one or more deposited bio-inks for a duration of about 1 hour to about 30 days; wherein the construct comprises one or more layers, each layer comprising one or more cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct. In some examples not claimed, the liver cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, the construct further comprising one or more of the following cell types: vascular cells, endothelial cells, parenchymal cells, non-parenchymal cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, the construct is characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry. In some examples not claimed, at least one component of the construct is bioprinted. In some examples not claimed, the construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, the construct is substantially free of any pre-formed scaffold at the time of use. In some examples not claimed, the construct is non-innervated.

[0028] Also disclosed herein but not claimed are methods of constructing a living three-dimensional liver tissue construct comprising: preparing one or more cohered multicellular aggregates comprising mammalian liver cells; placing said one or more cohered multicellular aggregates onto a support to form at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry; and incubating said one or more

multicellular aggregates for a duration of about 1 hour to about 30 days to allow them to cohere and to form a living, three-dimensional liver tissue construct. In some examples not claimed, the liver cells are derived from one or more of the following sources: adult mammalian liver tissue; fetal mammalian liver tissue; embryonic stem cells (ESC); ESC-derived hepatocyte-like cells, induced pluripotent stem cells (iPSC); iPSC-derived hepatocyte-like cells; adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples not claimed, the liver tissue construct further comprises one or more of the following cell types: vascular cells, endothelial cells, parenchymal cells, non-parenchymal cells, fibroblasts, mesenchymal cells, immune cells, cancer cells, Kupffer cells, stellate cells, biliary cells, sinusoidal endothelial cells, liver-derived stem/progenitor cells, and non-liver-derived stem/progenitor cells. In some examples not claimed, at least one component of the construct was bioprinted. In some examples not claimed, the construct further comprises an extrusion compound, the extrusion compound improving the suitability of the cells for bioprinting. In some examples not claimed, the construct is free of any pre-formed scaffold at the time of bioprinting or the time of use. In some examples not claimed, the construct is non-innervated.

[0029] Also disclosed herein but not claimed are methods of fabricating a living, three-dimensional liver tissue construct comprising incubating one or more bio-inks comprising non-parenchymal cells and one or more bio-inks comprising parenchymal cells deposited onto a support for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct, the construct comprising at least one compartment, the compartment comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells.

[0030] Also disclosed herein but not claimed are engineered, living, three-dimensional liver tissue constructs comprising one or more layers, at least one layer comprising at least two cell types, the at least two cell types comprising a parenchymal liver cell type and a non-parenchymal liver cell type, the at least two cell types spatially arranged relative to each other to create a planar geometry, the planar geometry comprising at least one parenchymal liver cell compartment and at least one non-parenchymal liver cell compartment, the one or more layers cohered to form a living, three-dimensional liver tissue construct, provided that the living, three-dimensional liver tissue construct remains substantially viable *in vitro* culture for at least 7 days and has at least two of the following liver characteristics: synthesizes cholesterol; stores lipid; stores glycogen; expresses liver-specific proteins, including two or more of: albumin, fibrinogen, transferrin, CYP450, alpha-1-antitrypsin, ornithine transcarbamylase, glycogen synthase, Factor VIII, and Factor IX; responsive to liver modulators, including one or more of: hormones, inducers, toxicants, drugs, antibodies, interfering RNAs, small RNAs, locked nucleic acids, viruses, bacteria, parasites, and inflammatory inducers; responsive to liver toxicants, including one or more of: acetaminophen, alcohol, trovafloxacin, methotrexate, diclofenac, troglitazone, valproic acid, and amiodarone; expresses transporter proteins, including one or more of: BSEP, OATP1B1, NTCP, PGP, BCRP, OATP1B3, ABCB4, ATP8B1, Multi-Drug Resistance Transporter (MDR); maintains at least some quiescent stellate cells after fabrication; and includes microvascular structures. In some examples not claimed, the

construct remains substantially viable *in vitro* culture for at least 14 days. In some examples not claimed, the construct remains substantially viable *in vitro* culture for at least 28 days. In some examples not claimed, the construct remains substantially viable *in vitro* culture for at least 40 days. In some examples not claimed, the construct remains substantially viable *in vitro* culture for at least 7, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, or more days. In some examples not claimed, the construct is produced in a fully automated procedure using a bioprinter. In some examples not claimed, the at least two cell types comprises cells derived from induced pluripotent stem cells (iPSC). In some examples not claimed, the at least two cell types comprises myeloid cells derived from monocytes or macrophages. In some examples not claimed, the construct has at least three, four, five or six, seven, eight, or all of the liver characteristics.

BRIEF DESCRIPTION OF THE DRAWINGS

[0031] Any of the following figures that do not fall within the scope of the claims do not form part of the invention.

[0032] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

Fig. 1 is an exemplary illustration of native liver tissue geometry.

Fig. 2 is an exemplary demonstration of maturation and biochemical characterization of bio-printed liver constructs containing HepaRG cells. HepaRG cells were deposited into the center of the perimeter box, followed by maturation for 96 hours (A-C). Bio-printed liver constructs are metabolically active producing albumin (D) and cholesterol (E) at 96 hours. Following dexamethasone treatment, cytochrome P450 (CYP3A4) activity is more than 5-fold higher than non-treated controls (F).

Fig. 3 is an example of bio-printed structures in defined geometric patterns (A, B) containing iPSC-derived hepatocytes (C) that synthesize significantly higher albumin levels (D), per million input cells, in the engineered, bioprinted construct than 2D culture controls.

Fig. 4 is an exemplary demonstration of maturation of a bio-printed liver construct (A) containing discrete spheroids of non-parenchymal cells (B). The bioprinted construct remained fused and stable upon incubation for an additional 24 hours (C, D).

Fig. 5 is an exemplary demonstration of dissolution of the co-printed mold which allows spatial and temporal control over the printing process of the hepatic cells and results in the generation of user-specified compartments of a defined shape and size in the x, y, and z axes. Arrows indicate regions where the co-printed mold has dissolved 24 hours post print and areas where

additional cellular inputs can be added.

Fig. 6 is a macroscopic image depicting a non-limiting example of an engineered liver tissue, in this case, a multi-layered liver tissue bioprinted using a continuous deposition mechanism using bio-ink composed of multiple liver cell types and a water-soluble extrusion compound (e.g., PF-127). (A) shows a schematic diagram of a single functional unit highlighting the planar geometry created by patterning bio-ink and negative space; (B) tessellated, functional units bioprinted with PF-127 containing 2×10^8 cells; (C) shows the construct 20 minutes after application of media; and (D) shows the construct 16 hours after application of media to the structure and dissolution of the extrusion compound. Note retention of the planar geometry over time.

Fig. 7 is a non-limiting photomicrograph of the tessellated construct of **Fig. 6**, depicting an exemplary "spoke" in the tessellated construct. This photomicrograph demonstrates Hematoxylin and Eosin staining of formalin-fixed paraffin-embedded tissue sections of stellate cells, endothelial cells, and dermal fibroblasts bioprinted by continuous deposition in a multilayer tessellated hexagonal structure and then cultured for 16 hours.

Fig. 8 is an example of dissolution of a co-printed mold allowing fusion of distinct compartmentalized hepatic cellular regions (marked by arrows). Over time (B; T=24 hours) the regions fuse into a solid bio-printed tissue.

Fig. 9 is an example of a co-printed mold (4% Gelatin and 2% Alginate) bio-printed using a Syringe Deposition Module (SDM) of a MMX Bioprinter produces constructs (A) that fuse and mature into hepatic tissues 72 hours post print (B). In this example, the bio-printed tissue is dense demonstrated by the H and E staining (C) and viable as demonstrated by relatively fewer TUNEL stained cells (D) compared to proliferating cells (E).

Fig. 10 is an example of a co-printed mold (A; 4% Gelatin and 2% Alginate) bioprinted by continuous deposition with a syringe deposition module (SDM) to produce a fused three-dimensional liver tissue construct. Addition of endothelial bio-ink in the center of the structure increases the complexity of the construct (B). E-Cadherin (C) and CD31 (D) staining demonstrate epithelial and endothelial cells present throughout the areas in which they were bioprinted at 144 hours post-print. TUNEL staining demonstrates limited cell death within the core of the engineered liver tissue following 144h incubation (E).

Fig. 11 is an example of a co-printed mold (co-mold) comprised of hydrogel (4% gelatin:2% alginate) (A) or bio-ink comprised of non-parenchymal cells and a hydrogel-based extrusion compound (B). The bio-ink co-mold lines contained 150×10^6 cells per mL of hydrogel extrusion compound. Addition of HepG2 cells to the co-mold structure and incubation for 24h, results in a fused, bioprinted liver structure (C). Dissolution of the extrusion compound occurs over time in aqueous media.

Fig. 12 is an example of a dense, cellular liver tissue, fabricated by continuous deposition using two syringe deposition modules (SDM) on the NovoGen MMX Bioprinter. The co-mold lines containing 4% gelatin and 2% alginate were deposited by the first SDM module followed

by filling of HepG2 bio-paste with a second SDM.

Fig. 13 is an example of a bioprinted neotissue with laminar geometry. A NovoGel™ hydrogel base and co-printed confining box were bioprinted, followed by deposition of a first layer comprising liver epithelial cell bio-ink (HepG2 cells), onto which a second layer was bioprinted comprised of hepatic stellate cells and endothelial cells. In this example, the stellate:EC layer was bioprinted via continuous deposition of bio-ink containing a hydrogel extrusion compound (A). Gross images of construct immediately after fabrication demonstrating the two distinct layers of bio-ink (B). Hematoxylin and Eosin staining of sections of formalin-fixed paraffin-embedded constructs (C) following 48 hours of culture reveals distinct morphology of the two layers and establishment of a laminar geometry. CD31-positive cells are restricted to the upper layer of the construct where a suspension of endothelial cells and hepatic stellate cells were bioprinted (D), while IGF-2-positive HepG2 are found only in the bottom layer (E).

Fig. 14 is an example of cell patterning and layering in bioprinted liver tissues. Hematoxylin and Eosin staining of paraffin-embedded tissue sections reveals a contiguous neotissue (A) formed by bioprinting polytypic cell populations containing vascular endothelial cells and hepatic stellate cells. Staining of the tissue sections with antibody directed at CD31 reveals the presence of centrally-located EC-lined microvessels and an external layer of CD31-positive EC (B).

Fig. 15 is an example of stimulation of bioprinted hepatic stellate cell tissues with TGF-β1. Incubation of bioprinted hepatic stellate cell sheets with increasing concentrations of TGF-β1 (0, 1, 10, 50 ng/mL), results in changes in gross observation of the bioprinted tissues as increases in cytokine concentration lead to increases in tissue outgrowth formation (A-D, 0-50 ng/mL). Trichrome staining of tissue sections from bioprinted hepatic stellate-containing tissues reveals increases in collagen deposition and construct size and dramatic decreases in cell density (E-H, 0-50 ng/mL).

Fig. 16 is an example of bioprinting in multi-well formats. Bioprinted tissue constructs are generated in multi-well plates (A) or within multi-well culture inserts (B), which are optionally placed in an appropriate multi-well plate for long-term maintenance and maturation. Here, tissue constructs were bioprinted in a 48-well polystyrene plate (A) and on the porous membrane of a 6-well cell culture insert (B).

Fig. 17 is an example of bio-printed hepatic tubular structure designed and printed for extracorporeal support. 40 mm length bio-ink cylinders comprising 70% HepG2/25% HUVEC/5% Hepatic Stellate were bio-printed and conditioned. A channel was created in the center of the structure by use of a removable, bio-inert hydrogel.

Fig. 18 is a series of non-limiting examples of planar and laminar geometries, including combinations thereof that are compatible with the methods of construction described herein, and reproduce architectural or spatial elements of native tissue architecture and biology. Exemplary geometries include an architecturally correct tissue with a vascular network (A), a zonal tissue (B), a lobulated tissue (C), a perfused/arrayed tissue (D), a tissue with a solid+liquid/liquid interface (E), a barrier tissue (F), and a layered tissue with laminar geometry

(G).

Fig. 19 is an schematic representation of an extracorporeal hepatic device containing repeating tubular structures with void spaces (depicted as occupied by temporary filler bodies; dark circles) at regular intervals allowing perfusion across and/or through the engineered liver tissue.

Fig. 20 is an example of a bioprinted human liver tissue.

Fig. 21 is a series of non-limiting photomicrographs demonstrating histological characterization of bioprinted human liver tissue; in this case, the photomicrographs show high tissue density (A), production of ECM (B), production of lipids (C), and production of glycogen (D).

Fig. 22 is a series of non-limiting photomicrographs demonstrating histological characterization of bioprinted human liver tissue; in this case, the photomicrographs show regions of hepatocytes surrounded by non-parenchymal stroma and apparent self-organization into lobule type structure that mimics native liver tissue.

Fig. 23 is an example of data demonstrating production of albumin (A) and cholesterol (B) by bioprinted human liver tissue.

Fig. 24 is an example of data demonstrating production of transferrin (A) and fibrinogen (B) by bioprinted human liver tissue.

Fig. 25 is an example of data demonstrating relative quantity of bile salt exporting protein mRNA vs. housekeeping gene RNA (18s RNA) in bioprinted human liver tissue at Day 10 post-fabrication.

Fig. 26 is an example of data demonstrating sustained induction of active CYP3A4 enzyme by rifampicin treatment (A) in bioprinted liver tissue, which is inhibitable by verapamil (B).

Fig. 27 is an example of data demonstrating induction of CYP3A4 by diclofenac treatment in bioprinted liver tissue followed by tissue damage at higher doses as measured by lactate dehydrogenase (LDH).

Fig. 28 is an example of data demonstrating decrease in albumin (A) and increase in LDH (B) in a multiple dose study of trovafloxacin vs. levofloxacin in bioprinted liver tissue.

Fig. 29 is an example of data demonstrating decrease in viability as measured by ATP (A), a decrease in albumin (B), and an increase in LDH (C) in a multiple dose study of methotrexate in bioprinted liver tissue.

Fig. 30 is an example of data demonstrating response of bioprinted liver tissue to vitamin D as measured by induction of CYP.

Fig. 31 is an example of data demonstrating response to LPS stimulation as measured by IL-6 production in Kupffer cell-containing bioprinted liver tissue.

Fig. 32 is an example of data demonstrating dose dependent response to LPS (10 µg/ml (A)

and 100 µg/ml (B)) stimulation as measured by cytokine production (IL-10, IL-1b, and TNF-α) in Kupffer cell-containing bioprinted liver tissue.

Fig. 33 is an example of a monolayer of hepatic sinusoidal endothelial cells (HSECs) at about 90% confluence.

Fig. 34 is an example of data demonstrating HSEC response to LPS as measured by IL-6 production in two-dimensional cultures.

Fig. 35 is an example of a fused three-dimensional bioprinted liver tissue incorporating HSECs in the non-parenchymal fraction.

Fig. 36 is an example of a photomicrograph (FITC; 5X) depicting small molecule penetration through a cryo-sectioned three-dimensional bioprinted liver tissue.

DETAILED DESCRIPTION OF THE INVENTION

[0033] For many patients with liver failure, restoration of even partial liver mass through the delivery of healthy cells or tissue can have a meaningful impact on survival and liver function. According to the 2011 UNOS database, there were >10,000 patients awaiting a liver transplant in the US, with <50% of those patients actually receiving a transplanted liver. Far more patients could benefit from a transplanted liver than make it onto the transplant list, which is typically reserved for the sickest patients. Furthermore, many individuals suffer from chronic degenerative diseases for which transplantation is not a current healthcare paradigm. Thus, living, functional liver tissues would be of great clinical value. Segmental, or partial liver transplants are now performed routinely in the US, wherein a single organ is divided into 2-3 sections, or lobes, with each piece transplanted into a wait-listed patient. Because healthy liver tissue has a unique ability to grow and accommodate an individual according to their size and metabolic needs, the approach of partial-mass transplant is clinically viable.

[0034] Furthermore, many compounds fail in late-stage clinical trials or post-market due to unpredicted toxicity, which accounts for 35% of compound failures in Phase III (Curr. Pharm. Des. 2005 11 :3545). Unpredicted liver toxicity accounts for the majority of those failures, followed by cardiotoxicity and renal toxicity. Thus, better systems are needed to provide a more accurate prediction of human *in vivo* performance.

[0035] There is a need for materials, tools, and techniques that substantially increase the number and quality of innovative, cost-effective new medicines, without incurring unsustainable R&D costs. Accordingly, the inventors describe herein engineered liver tissue constructs, arrays thereof, and methods of making the same that have utility in tissue and organ engineering, *in vitro* assays, drug discovery, and other areas.

[0036] Previous models of liver tissue have been focused on providing engineered tissue constructs by seeding cells onto a three-dimensional scaffold material that is pre-formed and shaped to accommodate the intended application, or by allowing liver cells to form multicellular spheroids via random assembly, wherein a spherical aggregate of cells is generated but there is no directed architecture within the aggregate. Cells seeded onto scaffold materials have been primary cells, cell lines, engineered cells, and/or stem/progenitor cells. When multipotential stem or progenitor cells are utilized, they have either undergone a differentiation program in two-dimensional monolayer culture prior to seeding on a three-dimensional scaffold material, or they have first been seeded onto a scaffold material and then been subjected to a differentiation program, *in situ* or *in vitro*, to generate the desired tissue. The traditional approach is both laborious and inefficient in terms of cell yield, the time required for terminal differentiation of the cells within the construct, and the overall cellularity and architecture of the resulting three-dimensional structure.

[0037] Importantly, previous efforts and technology have failed to generate defined tissue compartments with specific cell types existing in particular positions within the architecture and that persist in the construct for days or weeks at a time (up to 40 days). Also, previous efforts and technology have failed to demonstrate native-like structures, including microvasculature and tight junctions, which contribute to tissue longevity. The technology described herein allows fine control over architecture and allows generation of highly defined geometries that resemble native tissues. Moreover, the technology enables the construction of high cell density tissues that retain these geometries, thus extending the useful life of the tissues.

[0038] Disclosed herein, are engineered, living, three-dimensional liver tissue constructs, the construct comprising at least one compartment defining a planar geometry, the compartment comprising an interior defined by a border, the interior comprising parenchymal cells, the border comprising non-parenchymal cells; the cells cohered to form a living, three-dimensional liver tissue construct; provided that at least one component of the construct was bioprinted and the construct is substantially free of pre-formed scaffold at the time of use, wherein the parenchymal and non-parenchymal cells are as claimed.

[0039] Also disclosed herein, are engineered, living, three-dimensional liver tissue constructs comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct, the construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry, wherein the cell types are as claimed.

[0040] Also disclosed herein, but not claimed, are engineered, living, three-dimensional liver tissue constructs comprising: a plurality of layers, each layer comprising cylindrical bio-ink, the bio-ink axially-aligned substantially in parallel, the bio-ink comprising parenchymal liver cells; and optionally, non-parenchymal cells within or among the cylindrical bio-ink; and optionally, void spaces among the cylindrical bio-ink.

[0041] Also disclosed herein, are arrays of engineered living, three-dimensional liver tissue constructs, each construct comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct, each construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry, wherein the cell types are as claimed.

[0042] Also disclosed herein, but not claimed, are arrays of engineered living, three-dimensional tissue constructs, wherein at least one of the constructs is a liver tissue construct, each liver tissue construct comprising: one or more layers, each layer comprising one or more liver cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct, the liver tissue construct characterized by having at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry.

[0043] Also disclosed herein, are methods of fabricating a living, three-dimensional liver tissue construct comprising: providing one or more bio-inks comprising non-parenchymal cells; providing one or more bio-inks comprising parenchymal cells; depositing the bio-inks onto a support; and incubating the deposited bio-inks for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct comprising at least one compartment, the compartment comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells, wherein the parenchymal and non-parenchymal cells are as claimed.

[0044] Also disclosed herein, but not claimed, are methods of fabricating an engineered living, three-dimensional liver tissue construct comprising: preparing one or more bio-inks comprising liver cells; depositing the one or more bio-inks onto a support; and incubating the one or more deposited bio-inks for a duration of about 1 hour to about 30 days; wherein the construct comprises one or more layers, each layer comprising one or more cell types, the one or more layers cohered to form a living, three-dimensional liver tissue construct.

[0045] Also disclosed herein, but not claimed, are methods of constructing a living, three-dimensional liver tissue construct comprising: preparing one or more cohered multicellular aggregates comprising mammalian liver cells; placing said one or more cohered multicellular aggregates onto a support to form at least one of: at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry; and incubating said one or more multicellular aggregates for a duration of about 1 hour to about 30 days to allow them to cohere and to form a living, three-dimensional liver tissue construct.

[0046] Also disclosed herein, but not claimed, are methods of fabricating a living, three-dimensional liver tissue construct comprising incubating one or more bio-inks comprising non-parenchymal cells and one or more bio-inks comprising parenchymal cells deposited onto a support for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct, the construct comprising at least one compartment, the compartment comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells.

[0047] Also disclosed herein, but not claimed, are engineered, living, three-dimensional liver tissue constructs comprising one or more layers, at least one layer comprising at least two cell types, the at least two cell types comprising a parenchymal liver cell type and a non-parenchymal liver cell type, the at least two cell types spatially arranged relative to each other to create a planar geometry, the planar geometry comprising at least one parenchymal liver cell compartment and at least one non-parenchymal liver cell compartment, the one or more layers cohered to form a living, three-dimensional liver tissue construct, provided that the living, three-dimensional liver tissue construct remains substantially viable *in vitro* culture for at least 7 days or at least 14 days and has at least two of the following liver characteristics: synthesizes cholesterol; stores lipid; stores glycogen; expresses liver-specific proteins, including two or more of: albumin, fibrinogen, transferrin, CYP450, alpha-1-antitrypsin, ornithine transcarbamylase, glycogen synthase, Factor VIII, and Factor IX; responsive to liver modulators, including one or more of: hormones, inducers, toxicants, drugs, antibodies, interfering RNAs, small RNAs, locked nucleic acids, viruses, bacteria, parasites, and inflammatory inducers; responsive to liver toxicants, including one or more of: acetaminophen, alcohol, trovafloxacin, methotrexate, diclofenac, troglitazone, valproic acid, and amiodarone; expresses transporter proteins, including one or more of: BSEP, OATP1B1, NTCP, PGP, BCRP, OATP1B3, ABCB4, ATP8B1, Multi-Drug Resistance Transporter (MDR); maintains at least some quiescent stellate cells after fabrication; and includes microvascular structures.

Certain Definitions

[0048] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0049] As used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural references unless the context clearly dictates otherwise. Thus, for example, references to "a nucleic acid" includes one or more nucleic acids, and/or compositions of the type described herein which will become apparent to those persons skilled in the art upon reading this disclosure and so forth. Any reference to "or" herein is intended to encompass "and/or" unless otherwise stated.

[0050] As used herein, "array" means a scientific tool including an association of multiple

elements spatially arranged to allow a plurality of tests to be performed on a sample, one or more tests to be performed on a plurality of samples, or both.

[0051] As used herein, "assay" means a procedure for testing or measuring the presence or activity of a substance (e.g., a chemical, molecule, biochemical, protein, hormone, or drug, etc.) in an organic or biologic sample (e.g., cell aggregate, tissue, organ, organism, etc.).

[0052] As used herein, "biocompatible" means posing limited risk of injury or toxicity to cells. As presented in the specification and claims, "biocompatible multi-well containers" and "biocompatible membranes" pose limited risk of injury or toxicity to mammalian cells, but the definition does not extend to imply that these biocompatible elements could be implanted *in vivo* into a mammal.

[0053] As used herein, "bio-ink" means a liquid, semi-solid, or solid composition for use in bioprinting. In some examples, bio-ink comprises cell solutions, cell aggregates, cell-comprising gels, multicellular bodies, or tissues. In some examples, the bio-ink additionally comprises support material. In some examples, the bio-ink additionally comprises non-cellular materials that provide specific biomechanical properties that enable bioprinting. In some examples the bio-ink comprises an extrusion compound.

[0054] As used herein, "bioprinting" means utilizing three-dimensional, precise deposition of cells (e.g., cell solutions, cell-containing gels, cell suspensions, cell concentrations, multicellular aggregates, multicellular bodies, etc.) via methodology that is compatible with an automated or semi-automated, computer-aided, three-dimensional prototyping device (e.g., a bioprinter).

[0055] As used herein, "cohere," "cohered," and "cohesion" refer to cell-cell adhesion properties that bind cells, cell aggregates, multicellular aggregates, multicellular bodies, and/or layers thereof. The terms are used interchangeably with "fuse," "fused," and "fusion."

[0056] As used herein, "cylindrical" means having substantially the form of a cylinder. In some examples, cylindrical bio-ink has substantially the form of a cylinder and is about 99, 98, 97, 96, 95, 94, 93, 92, 91, 90, 85, 80, 75, 70, 65, or 60 percent, including increments therein, cylindrical in form. In some examples, cylindrical bio-ink has substantially the form of a cylinder along about 99, 98, 97, 96, 95, 94, 93, 92, 91, 90, 85, 80, 75, 70, 65, 60, 55, or 50 percent, including increments therein, of its length. In some examples, "cylindrical" means having substantially the form of a cylinder at the time of fabrication.

[0057] As used herein, "laminar" means a multi-layered bioprinted tissue in which two or more planar layers are combined to increase the overall thickness of the tissue in the Z-plane. In some examples, each planar layer is substantially similar in architecture and/or composition. In some examples, each planar layer is substantially distinct in architecture and/or composition.

[0058] As used herein, "layer" means an association of cells in X and Y planes that is multiple cells thick. In some examples, the engineered liver tissues describe herein include one layer. In

some examples, the engineered liver tissues describe herein include a plurality of layers. In some examples, a layer forms a contiguous, substantially contiguous, or non-contiguous sheet of cells. In some examples, each layer of an engineered liver tissue described herein comprises multiple cells in the X, Y, and Z axes. In some examples, each layer of the engineered liver tissue is at least about 50 microns in thickness in the X, Y, and Z axes.

[0059] As used herein, "multi-layered" means being comprised of two or more layers of tissue, wherein each tissue layer is one or more cell-layers in thickness. In some examples, layers of tissue are deposited one at a time. In some examples, multiple layers are deposited simultaneously. Each layer may be comprised of multiple cell types. Further, the multiple cell types within each layer are optionally arranged relative to each other in a spatially-defined architecture in the X-Y planes (i.e., horizontal planes). Furthermore, addition of layers in the Z-plane (i.e., vertical plane), in some cases, results in controlled spatial positioning of the cells within the layers relative to each other so that a spatially-defined architecture is continued in the Z-plane.

[0060] As used herein, "multi-potent cells" refers to cells that are capable of undergoing differentiation to two or more cell types. Multi-potent cells include, for example, mesenchymal stem/stromal cells, induced pluripotent stem cells, and embryonic stem cells.

[0061] As used herein, "parenchymal cells" refers to hepatocytes or hepatocyte-like cells; whereas, "non-parenchymal cells" refers to liver cells that are not hepatocytes or hepatocyte-like cells.

[0062] As used herein, "planar" means a layer of multicellular bioprinted tissue in which multiple bio-ink compositions and/or void spaces are spatially arranged into a defined pattern relative to each other substantially within the X-Y plane of the tissue layer. See, e.g., **Figs. 18A-E**. In some examples, a planar layer is substantially planar, for example, about 99, 98, 97, 96, 95, 94, 93, 92, 91, 90, 85, 80, 75, 70, 65, or 60 percent, including increments therein, spatially arranged into a defined pattern within the X-Y plane of the tissue layer.

[0063] As used herein, "scaffold" refers to synthetic scaffolds such as polymer scaffolds and porous hydrogels, non-synthetic scaffolds such as pre-formed extracellular matrix layers, dead cell layers, and decellularized tissues, and any other type of pre-formed scaffold that is integral to the physical structure of the engineered tissue and/or organ and not able to be removed from the tissue and/or organ without damage/destruction of said tissue and/or organ. In some examples, decellularized tissue scaffolds include decellularized native tissues or decellularized cellular material generated by cultured cells in any manner; for example, cell layers that are allowed to die or are decellularized, leaving behind the ECM they produced while living. The term "scaffoldless," therefore, is intended to imply that scaffold is not an integral part of the engineered tissue at the time of use, either having been removed or remaining as an inert component of the engineered tissue. "Scaffoldless" is used interchangeably with "scaffold-free" and "free of pre-formed scaffold."

[0064] As used herein, "subject" means any individual, which is a human, a non-human animal, any mammal, or any vertebrate. The term is interchangeable with "patient," "recipient" and "donor."

[0065] As used herein, "support" means any biocompatible surface capable of receiving deposited bio-ink.

[0066] As used herein, "tissue" means an aggregate of cells. Examples of tissues include, but are not limited to, connective tissue (e.g., areolar connective tissue, dense connective tissue, elastic tissue, reticular connective tissue, and adipose tissue), muscle tissue (e.g., skeletal muscle, smooth muscle and cardiac muscle), genitourinary tissue, gastrointestinal tissue, pulmonary tissue, bone tissue, nervous tissue, and epithelial tissue (e.g., simple epithelium and stratified epithelium), endoderm-derived tissue, mesoderm-derived tissue, and ectoderm-derived tissue.

Tissue Engineering

[0067] Tissue engineering is an interdisciplinary field that applies and combines the principles of engineering and life sciences toward the development of biological substitutes that restore, maintain, or improve tissue function through augmentation, repair, or replacement of an organ or tissue. The basic approach to classical tissue engineering is to seed living cells into a biocompatible and eventually biodegradable environment (e.g., a scaffold), and then culture this construct in a bioreactor so that the initial cell population expands further and matures to generate the target tissue upon implantation. With an appropriate scaffold that mimics the biological extracellular matrix (ECM), the developing tissue, in some cases, adopts both the form and function of the desired organ after *in vitro* and *in vivo* maturation. However, achieving high enough cell density with a native tissue-like architecture is challenging due to the limited ability to control the distribution and spatial arrangement of the cells throughout the scaffold. These limitations often result in tissues or organs with poor mechanical properties and/or insufficient function. Additional challenges exist with regard to biodegradation of the scaffold, entrapment of residual polymer, and industrial scale-up of manufacturing processes. Scaffoldless approaches have been attempted. Current scaffoldless approaches are subject to several limitations:

- Complex planar and/or laminar geometries, such as multi-layered structures wherein one or more layers is compositionally or architecturally distinct from other layers or wherein one or more layers comprise multiple cell types in spatially-defined positions relative to each other, often require definitive, high-resolution placement of cell types within a specific architecture to reproducibly achieve a native tissue-like outcome.
- Scale and geometry are limited by diffusion and/or the requirement for functional vascular networks for nutrient supply.
- The viability of the tissues is, in some cases, compromised by confinement material that limits diffusion and restricts the cells' access to nutrients.

[0068] Disclosed herein, are engineered mammalian tissues, engineered liver tissues/constructs, arrays thereof, and methods of fabrication. The tissue engineering methods disclosed herein have the following advantages:

- They are capable of producing cell-comprising tissues and/or organs.
- They mimic the environmental conditions found within the development, homeostasis, and/or pathogenesis of natural tissues by re-creating native tissue-like intercellular interactions.
- They optionally achieve living, three-dimensional tissues and compound tissues with a broad array of complex topologies and geometries (e.g., multilayered structures, segments, sheets, tubes, sacs, etc.).
- They are compatible with automated or semi-automated means of manufacturing and are scalable.

[0069] Bioprinting enables improved methods of generating micro-scale tissue analogues including those useful for *in vitro* assays (see below).

Bioprinting

[0070] In some examples, at least one component of the engineered liver tissues/constructs, and arrays thereof is bioprinted. In some examples, bioprinted constructs are made with a method that utilizes a rapid prototyping technology based on three-dimensional, automated, computer-aided deposition of cells, including cell solutions, cell suspensions, cell-comprising gels or pastes, cell concentrations, multicellular bodies (e.g., cylinders, spheroids, ribbons, etc.), and, optionally, confinement material onto a biocompatible support surface (e.g., composed of hydrogel and/or a porous membrane) by a three-dimensional delivery device (e.g., a bioprinter). As used herein, the term "engineered," when used to refer to tissues and/or organs means that cells, cell solutions, cell suspensions, cell-comprising gels or pastes, cell concentrates, multicellular aggregates, and layers thereof are positioned to form three-dimensional structures by a computer-aided device (e.g., a bioprinter) according to a computer script. In some examples, the computer script is, for example, one or more computer programs, computer applications, or computer modules. In some examples, three-dimensional tissue structures form through the post-printing fusion of cells or multicellular bodies which, in some cases, is similar to self-assembly phenomena in early morphogenesis.

[0071] While a number of methods are available to arrange cells, multicellular aggregates, and/or layers thereof on a biocompatible surface to produce a three-dimensional structure including manual placement, positioning by an automated, computer-aided machine such as a bioprinter is advantageous. Advantages of delivery of cells or multicellular bodies with this

technology include rapid, accurate, and reproducible placement of cells or multicellular bodies to produce constructs exhibiting planned or pre-determined orientations or patterns of cells, multicellular aggregates and/or layers thereof with various compositions. Advantages also include assured high cell density, while minimizing cell damage. Parenchymal cells of the liver are particularly susceptible to damage by shear force and other biomechanical stress; thus the combined use of bio-ink and the bioprinting process described herein provides a distinct advantage over alternative technologies as highlighted by the favorable viability of the parenchymal cells after bioprinting as highlighted in **Fig. 9**.

[0072] In some examples, the method of bioprinting is continuous and/or substantially continuous. A non-limiting example of a continuous bioprinting method is to dispense bio-ink (i.e., cells, cells combined with an excipient or extrusion compound, or aggregates of cells) from a bioprinter via a dispense tip (e.g., a syringe, needle, capillary tube, etc.) connected to a reservoir of bio-ink. In some examples, a continuous bioprinting method is to dispense bio-ink in a repeating pattern of functional units. In some examples, a repeating functional unit has any suitable geometry, including, for example, circles, squares, rectangles, triangles, polygons, and irregular geometries, thereby resulting in one or more tissue layers with planar geometry achieved via spatial patterning of distinct bio-inks and/or void spaces. In some examples, a repeating pattern of bioprinted function units comprises a layer and a plurality of layers are bioprinted adjacently (e.g., stacked) to form an engineered tissue or organ with laminar geometry. In some examples, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more layers are bioprinted adjacently (e.g., stacked) to form an engineered tissue or organ. In further embodiments, one or more layers of a tissue with laminar geometry also has planar geometry.

[0073] In some examples, a functional unit consists of a number of fused or cohered tubular structures which are arranged in both the horizontal and vertical dimensions to form a contiguous liver tissue structure that contains void spaces or channels at regular intervals which allows perfusion across and/or through the engineered liver tissue. See **Figs. 17** and **19**.

[0074] In some examples, a bioprinted functional unit repeats in a tessellated pattern. A "tessellated pattern" is a plane of figures that fills the plane with no overlaps and no gaps. **Fig. 6A** shows an example of a functional unit that is optionally repeated to produce the tessellation pattern depicted in **Figs. 6B-D** and **7**. Advantages of continuous and/or tessellated bioprinting includes, by way of example, increased productivity of bioprinted tissue. Another exemplary advantage is eliminating the need to align the bioprinter with previously deposited elements of bio-ink. In some examples, continuous bioprinting facilitates printing larger tissues from a large reservoir of bio-ink, optionally using a syringe mechanism. Continuous bioprinting is also a convenient way to co-print spatially-defined boundaries, using an extrusion compound, a hydrogel, a polymer, bio-ink, or any printable material that is capable of retaining its shape post-printing; wherein the boundaries that are created are optionally filled in via the bioprinting of a one or more bio-inks, thereby creating a mosaic tissue with spatially-defined planar geometry, see for example, **Figs. 3A, 5A, 5B, 6, 8, and 11**.

[0075] In some examples, methods in continuous bioprinting involve optimizing and/or

balancing parameters such as print height, pump speed, robot speed, or combinations thereof independently or relative to each other. In certain cases, the bioprinter head speed for deposition was 3 mm/s, with a dispense height of 0.5 mm for the first layer and dispense height was increased 0.4 mm for each subsequent layer. In some examples, the dispense height is approximately equal to the diameter of the bioprinter dispense tip. Without limitation a suitable and/or optimal dispense distance does not result in material flattening or adhering to the dispensing needle. In some examples, the bioprinter dispense tip has an inner diameter of about, 20, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 μm , or more, including increments therein. In some examples, the bio-ink reservoir of the bioprinter has a volume of about 0.05, 0.1, .5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 cubic centimeters, or more, including increments therein. The pump speed is, in some cases, suitable and/or optimal when the residual pressure build-up in the system is low. Favorable pump speeds, in some cases, depend on the ratio between the cross-sectional areas of the reservoir and dispense needle with larger ratios requiring lower pump speeds. In some embodiments, a suitable and/or optimal print speed enables the deposition of a uniform line without affecting the mechanical integrity of the material.

[0076] Also disclosed herein but not claimed are business methods. In some examples not claimed, the speed and scalability of the techniques and methods disclosed herein are utilized to design, build, and operate industrial and/or commercial facilities for production of engineered liver tissues and/or organs for implantation or use in generation of cell-based tools for research and development, such as *in vitro* assays. In some examples not claimed, the engineered liver tissues and/or organs and arrays thereof are produced, stored, distributed, marketed, advertised, and sold as, for example, cellular arrays (e.g., microarrays or chips), tissue arrays (e.g., microarrays or chips), and kits for biological assays and high-throughput drug screening. In some examples not claimed, the engineered liver tissues and/or organs and arrays thereof are produced and utilized to conduct biological assays and/or drug screening as a service.

Engineered liver tissues

[0077] Referring to **Fig. 1**, native liver tissue is comprised of multiple cell types, arranged in a distinct, spatially-patterned architecture relative to each other. Each liver lobule constitutes an operating functional unit of liver and is roughly hexagonal in shape. Parenchymal cells, or *hepatocytes*, largely constitute the lobules with various non-parenchymal cells (bile duct cells, sinusoidal endothelial cells, stellate cells, and Kupffer cells) in adjunct positions surrounding the lobule and radiating between the portal veins and bile ducts (along the periphery of the lobule) toward the central vein (in the center of the lobule). The positioning of non-parenchymal cells and the flow of blood and bile along the sinusoids and bile canaliculi, respectively, act together to form the proper micro-environment for the hepatocytes and thus influence the state of health and function of those cells. Various pathological conditions, like cirrhosis or fibrosis in the liver, lead to loss of critical metabolic functions as the hepatocyte micro environment is

disrupted and the architecture of the liver is progressively destroyed due to overproduction of extracellular matrix, inflammation, and production of pathogenic cytokines and growth factors.

[0078] Bioprinting enables compartmentalized liver tissue to be generated from multiple cellular inputs in three-dimensions, thus recapitulating key elements of native tissue architecture and function. Lobulated tissues can be generated per the examples provided herein, wherein the parenchymal and non-parenchymal cells are spatially arranged relative to each other, creating a planar geometry within each layer of tissue that is fabricated. Subsequent layers can be added that either reproduce the geometry of the first layer precisely, or introduce additional features such as void spaces or channels or additional biological elements, such as tumor cells or other biological or biochemical components associated with pathogenic or reparative / regenerative processes. In addition to lobulated patterns, one can utilize the accompanying diagrams to appreciate that bioprinted tissue could be generated that reproduces specific spatial relationships within tissues, including but not limited to: vascular / parenchymal; vascular/hepatocyte; hepatocyte/bile duct; fibrotic tissue/vasculature; fibrotic tissue/hepatocyte; hepatocyte/immune cells; parenchymal/non-parenchymal, hepatic sinusoid/blood or blood surrogate. In further embodiments, the liver tissue analogues comprise: hepatocytes or hepatocyte-like cells and optionally bile duct epithelial cells and optionally, non-parenchymal cell types including, but not limited to, stellate cells, endothelial cells, Kupffer cells, immune cells, or myofibroblasts.

[0079] Also disclosed herein, in some examples, are engineered liver tissues comprising cohered, mammalian cells, and further comprising one or more layers of mammalian cells, wherein at least one component of the tissue was bioprinted. In some examples, one or more of the tissue layers is characterized by a planar geometry, wherein multiple cell types or bio-ink types and/or void spaces exist in spatially-defined positions in the X-Y planes. In some examples, the tissues are multi-layered wherein at least one of the layers is architecturally or compositionally distinct from the other layers, giving the tissue a characteristic laminar geometry. In some examples, the layers are of similar thickness in the Z-plane. In some examples, the layers are of variable thickness in the Z-plane. In some examples, any single layer is one cell layer in thickness. In some examples, the tissues are liver analogues. In some examples, the liver tissue analogues comprise: hepatocytes or hepatocyte-like cells and optionally bile duct epithelial cells and optionally, non-parenchymal cell types including, but not limited to, stellate cells, endothelial cells, Kupffer cells, immune cells, or myofibroblasts. In some examples, the resulting liver tissue constructs are at least about 50 microns in thickness in the x, y, and z planes.

[0080] In some examples, the engineered liver tissues/constructs, are bioprinted, a methodology described herein. In some examples, at least one component of the engineered tissue is bioprinted. In some examples, additional components of the tissue are bioprinted. In some examples, the tissues are free of any pre-formed scaffold as described further herein at the time of manufacture or at the time of use. In some examples, as a result of being fabricated by tissue engineering techniques, including bioprinting, the tissues are further distinguished from tissues developed *in vivo*, as part of an organism.

[0081] In some examples, the engineered liver tissue includes any type of mammalian cell. In some examples, the engineered liver tissue includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more cell types. In some examples, the cells of the engineered liver tissue are "cohered" or "adhered" to one another. In some examples, cohesion or adhesion refers to cell-cell adhesion properties that bind cells, multicellular aggregates, multicellular bodies, and/or layers thereof.

[0082] In some examples, the engineered liver tissues/constructs, include one or more layers of cells. In some examples, one or more of the layers is characterized by having a planar geometry. In some examples, multiple layers of the engineered tissue have a planar geometry; wherein the planar geometries are variable among layers or are the same. In some examples, planar geometries (X-Y planes) in individual layers are aligned in the Z-plane during fabrication so that additional geometry is created in the Z-plane in the composite tissue. In some examples, one or more of the layers within the multi-layered architecture is characterized further by having planar geometry.

[0083] In some examples, a layer of cells comprises one or more sheets of cells. In some examples, a sheet of cells is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more cells thick, including increments therein. In some examples, a sheet of cells is about 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 or more μm thick, including increments therein. In some examples, a layer of tissue comprises fused aggregates of cells. In some examples, prior to fusion, the aggregates of cells have, a defined shape and/or architecture, being substantially spherical, elongate, substantially cylindrical and ribbon-like shape. In some examples, fused aggregates of cells form a layer about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 or more μm thick, including increments therein.

[0084] In some examples, the one or more layers include any type of mammalian cell. In some examples, each layer includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more cell types. In some examples, the engineered liver tissues/constructs include one or more layers of endothelial cells on one or more surfaces. In some examples, the engineered liver tissues/constructs include one or more layers of epithelial cells on one or more surfaces.

[0085] The engineered liver tissues, in some examples, are any suitable size. In some examples, the size of bioprinted liver tissue, change over time. In some examples, an engineered liver tissue shrinks or contracts after bioprinting due to, for example, cell migration, cell death, intercellular interactions, contraction, or other forms of shrinkage. In some examples, an engineered liver tissue grows or expands after bioprinting due to, for example, cell migration, cell growth and proliferation, production of extracellular matrix or other cell-produced components of native tissue, cell/tissue maturation or other forms of expansion.

[0086] In some examples, the physical dimensions of the engineered liver tissues are limited

by the capacity for nutrients, including oxygen, to diffuse into the interior of the construct. In some examples, the engineered liver tissues/constructs are at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, or 1000 μm in their smallest dimension at the time of bioprinting. In some examples, the engineered liver tissues are at least about 0.25, 0.5, 0.75, 1.0, 1.25, 1.5, 1.75, 2.0, 2.25, 2.5, 2.75, 3.0, 3.25, 3.5, 3.75, 4.0, 4.25, 4.5, 4.75, or 5.0 mm in their smallest dimension at the time of bioprinting. In some examples, the engineered liver tissues/constructs are between about 25 μm and about 500 μm in their smallest dimension at the time of bioprinting.

[0087] The engineered liver tissues, in some examples, are any suitable shape. In some examples, the shape is selected to mimic a particular natural tissue or organ. In some examples, the shape is selected to mimic a particular pathology, condition, or disease state. In some examples, the engineered liver tissues have a shape that is substantially planar. In some examples, planar tissues have any suitable planar geometry including, by way of non-limiting examples, square, rectangle, polygon, circle, oval, or irregular. In some examples, a planar geometry is generated in an engineered liver tissue by positioning specific cellular or bio-ink components and/or void spaces in the X-Y planes relative to each other.

[0088] In some examples, the engineered liver tissues/constructs are secured to a containment vessel by a means suitable to fix the position of the tissue in space relative to the containment vessel. In some examples, the engineered liver tissues are confined by a biocompatible material that provides physical support on one or more sides and constrains the space that the engineered liver tissue occupies. In some examples, the engineered liver tissues are affixed to a surface. In some examples, the engineered liver tissues are affixed to a biocompatible surface. In some examples, a plurality of tissues are associated by affixation to a surface and spatially arranged to form an array, as described herein. In some examples, engineered liver tissues are subjected to shear force, caused by fluid flow, on one or more sides. In some examples, application of shear force serves to facilitate the maturation and development of a tissue and/or facilitate the migration, differentiation, proliferation, deposition of extracellular matrix, or transport of proteins or molecules into or out of cells within the tissue. In some examples, the engineered liver tissues are subjected to continuous or periodic perfusion, recirculation, or agitation of liquid nutrients on one or more surfaces. In some examples, the engineered liver tissues and arrays thereof are housed in a multi-well bioreactor that provides continuous or periodic recirculation of the liquid culture media for each construct.

Tissue Geometries

[0089] Native tissues are characterized by the presence of spatial and compositional patterns driven by the cellular and extracellular (i.e., void spaces, extracellular matrices, proteinaceous matter, etc.) components of a tissue. Inherent challenges to tissue engineering strategies that deploy synthetic scaffolding to achieve three-dimensionality is the inability to reproduce both

the geometric and biologic attributes of native tissue. To date, attempts to create native tissue-like laminar or planar geometry within a scaffold structure while also enabling the incorporation of cells at a density that mimics native tissue have been hampered by technical limitations. Bioprinting overcomes both inherent challenges (planar/laminar geometry and cell density) through the spatially-defined deposition of bio-ink comprised of cells, according to the examples illustrated in **Figs. 18A-E**. In some examples, planar geometries are created from multiple bio-ink formulations, whereby two or more tissue components (i.e., stromal, epithelial, vascular, bone, cartilage, parenchymal, cortical, medullary, papillary, lobular, etc.) are fabricated in a manner that positions each tissue component / cell population / bio-ink formulation in a defined position relative to each other in the X, Y, and/or Z planes. In some examples, the planar geometry incorporates void spaces. In some examples, the void spaces within the planar geometry accommodate fluids that mimic at least one element of bodily fluids, such as blood, lymph, bile, urine, secretions, and the like. In some examples, the void spaces optionally contain non-adherent cell types or bodily-fluid-derived components (e.g., blood cells, marrow cells, lymphatic cells, immune cells, cancer cells, platelets, proteins, etc.). In some examples, non-adherent cell types of bodily-fluid-derived components optionally exist as a component of non-void spaces having been introduced into the cell-comprising components of the planar geometry before, during, or after fabrication. In some examples, non-adherent cellular components or bodily-fluid-derived components are recruited from void spaces into cell-comprising spaces within the planar geometry as a result of intercellular interactions or response to secreted factors.

[0090] In some examples, fluid flow or perfusion is optionally initiated through the void spaces within a geometry. In some examples, planar geometries enable the generation of tissue-tissue or tissue-liquid interfaces. In some examples, the tissues are fabricated into containers that are optically clear to enable real-time observation of cells at the interface(s) created by the geometry.

[0091] In some examples, tissues comprise multiple layers wherein at least one of the layers is architecturally or compositionally distinct from other layers within the construct, thereby creating a laminar architecture in the Z-plane. Examples of laminar architecture include barrier tissues that possess an endothelial or epithelial barrier to an underlying interstitial tissue as depicted by the examples shown in **Fig. 18F**. In some examples, one or more layers of a tissue incorporate vascular or microvascular components. In some examples, the incorporation of vascular or microvascular components leads to the formation of microvascular or pseudovascular networks within one or more components of the engineered tissue. In some examples, one or more components of the tissue with laminar geometry are bioprinted. In some examples, one or more tissues with laminar geometry are fabricated adjacent to each other, thereby creating a tissue interface.

[0092] In some examples, one or more layers of a multi-layered engineered tissue with laminar geometry also comprise planar geometry, according to the non-limiting examples set forth in **Figs. 18G**. In some examples, the same planar geometry is continued in each layer, resulting in a three-dimensional tissue with continuous architecture in the X, Y, and Z planes. In some

examples, the composition or planar geometry of one or more laminar layers is varied, such that the resulting three-dimensional tissue possesses a complex architecture in the X, Y and Z planes.

[0093] Referring to **Fig. 18A**, an engineered liver tissue is fabricated with a planar geometry representing an architecturally-correct tissue with a vascular network. In this example, an engineered liver tissue is fabricated with a first bio-ink including parenchymal cells, such as liver cells. Each layer of parenchymal cells optionally includes any planar pattern. In this example, an engineered liver tissue is fabricated with a second bio-ink including vascular-related cells, such as endothelial cells, smooth muscle cells, and fibroblast to form a vascular network. Optionally, this architecture may be created by first bioprinting and establishing the vascular network, and then fabricating liver tissue surrounding that network in a second step.

[0094] Referring to **Fig. 18B**, an engineered liver tissue is fabricated with a planar geometry representing a tissue zonal junction. This could be utilized, for example, to replicate the three specific zones of the hepatic acinus from the periportal region to the centrilobular region.

[0095] Referring to **Fig. 18C**, an engineered liver tissue is fabricated with a planar geometry representing a lobulated tissue. In this example, an engineered liver tissue is fabricated with a first bio-ink including liver cells to form lobules. Each geometric lobule optionally includes a spatially-directed architecture within its boundaries. In this example, an engineered liver tissue is fabricated with a second bio-ink including stromal/vascular cells to form borders around the lobules.

[0096] Referring to **Fig. 18D**, an engineered liver tissue is fabricated with a planar geometry representing a perfused/arrayed tissue. In this example, a first component forms channels, vessels, or tubes with lumens. In this example, a second component forms tissue patches that are optionally the same size/shape or different sizes/shapes, optionally include one cell type or a plurality of cell types, and optionally include one or more spatial patterns achieved by directed patterning in the X, Y, and or Z planes.

[0097] Referring to **Fig. 18E**, an engineered liver tissue is fabricated with a planar geometry representing a solid+liquid/liquid tissue interface. In this example, a first bio-ink forms the outer wall of a luminal structure, a second bio-ink forms the inner wall of a luminal structure where required, and third component is optionally a fluid containing cells or other biologically relevant components.

[0098] Referring to **Fig. 18F**, an engineered liver tissue is fabricated with a laminar geometry representing a barrier tissue. In this example, a barrier layer is optionally endothelial or epithelial and an interstitial layer forms the wall and/or surface of a luminal tissue, which rests upon a porous mesh or membrane.

[0099] Referring to **Fig. 18G**, an engineered liver tissue is fabricated with a plurality of layers, each with a planar geometry and stacked to form a tissue with a laminar geometry. In this

example, the planar geometry of the X and Y axes is continued through the Z-axis into the fabricated tissue. Features of the geometry optionally include contiguous channels or cellular compartments.

Cells

[0100] Disclosed herein, are engineered liver tissues (e.g., liver analogues) comprising one or more types of mammalian cells. In some examples, the engineered liver tissues comprise hepatocytes or hepatocyte-like cells. In some examples, suitable hepatocytes or hepatocyte-like cells include, but are not limited to primary hepatocytes, cell lines such as HepG2 cells, tissue-specific progenitors such as HepaRG cells, or combinations thereof. In some examples, the engineered liver tissues comprise bile duct epithelial or bile duct epithelial-like cells. In some examples, the engineered liver tissues comprise non-parenchymal cells. In some examples, suitable non-parenchymal cells include, vascular cells, endothelial cells, fibroblasts, myofibroblasts, mesenchymal cells, immune cells, Kupffer cells, stellate cells, biliary epithelial cells, biliary epithelial-like cells, and sinusoidal endothelial cells and combinations thereof.

[0101] In some examples not claimed, any mammalian cell is suitable for inclusion in the engineered liver tissues and arrays thereof. In further examples, at least one component of the engineered liver tissues is an adherent cell type. In some examples, the mammalian cells are, liver cells, parenchymal cells, non-parenchymal cells, fibroblasts endothelial cells, vascular cells, undifferentiated cells (e.g., embryonic cells, stem cells, and progenitor cells), and combinations thereof.

[0102] In some embodiments, the engineered liver tissues include endothelial cells, such as human endothelial cells. In some embodiments, suitable endothelial cells originate from tissue including, by way of non-limiting example, liver tissue, blood, blood vessel, lymphatic vessel, tissue of the digestive tract, tissue of the genitourinary tract, adipose tissue, tissue of the respiratory tract, tissue of the reproductive system, bone marrow, and umbilical tissue.

[0103] In some examples, the cells are adult, differentiated cells. In some examples, "differentiated cells" are cells with a tissue-specific phenotype consistent with, for example, a hepatocyte or an endothelial at the time of isolation, wherein tissue-specific phenotype (or the potential to display the phenotype) is maintained from the time of isolation to the time of use. In some examples, the cells are adult, non-differentiated cells. In some examples, "non-differentiated cells" are cells that do not have, or have lost, the definitive tissue-specific traits. In some examples, non-differentiated cells include stem cells. In some examples, "stem cells" are cells that exhibit potency and self-renewal. Stem cells include, totipotent cells, pluripotent cells, multipotent cells, oligopotent cells, unipotent cells, and progenitor cells. In some examples, stem cells are embryonic stem cells, adult stem cells, amniotic stem cells, and induced pluripotent stem cells. In some examples, the cells are a mixture of adult, differentiated cells and adult, non-differentiated cells.

[0104] In some embodiments, the engineered liver tissues include parenchymal cells are derived from multi-potent stem/progenitor cells. In various embodiments, the parenchymal cells are suitably derived from fetal mammalian liver tissue; embryonic stem cells (ESC); induced pluripotent stem cells (iPSC); adult stem/progenitor cells derived from the liver; and adult stem/progenitor cells derived from a tissue other than liver. In some examples, engineered liver tissues are derived from multi-potent stem/progenitor cells that have been either partially or fully differentiated into endodermal or hepatic phenotype prior to use in the fabrication of liver tissue constructs.

[0105] In some examples, the construct comprises stem/progenitor cells that have been exposed to one or more differentiation signals. In some examples, the differentiation signal comprises one or more of: biomechanical signals, soluble signals (e.g., a biochemical signal), and physical signals. Stem/progenitor cells are suitably exposed to one or more differentiation signals at many time points in the engineered liver tissue fabrication process. In some examples, the stem/progenitor cells were exposed to one or more differentiation signals before fabrication of the construct. In some examples, the stem/progenitor cells were exposed to one or more differentiation signals during fabrication of the construct. In some examples, the stem/progenitor cells were exposed to one or more differentiation signals after fabrication of the construct.

[0106] In some examples, the cell types and/or source of the cells are selected, configured, treated, or modulated based on a specific research goal or objective. In some examples, one or more specific cell types are selected, configured, treated, or modulated to facilitate investigation of a particular disease or condition. In some examples, one or more specific cell types are selected, configured, treated, or modulated to facilitate investigation of a disease or a condition of a particular subject. In some examples one or more specific cell types are derived from two or more distinct human donors. In some examples, one or more specific cell types are derived from a particular vertebrate subject. In some examples, one or more specific cell types are derived from a particular mammalian subject. In still further examples, one or more specific cell types are derived from a particular human subject. In some examples, one or more specific cell types are derived from a particular subject with a specific phenotype associated with disease or tissue functionality. In some examples not claimed, the subject-specific cells are isolated from the target tissue of interest by way of biopsy or tissue sampling. In some examples, the subject-specific cells are utilized to fabricate tissue immediately after isolation. In some examples, the subject-specific cells are manipulated *in vitro* prior to use in the fabrication of three-dimensional tissues; wherein the manipulation includes one or more of: expansion, differentiation, directed differentiation, proliferation, exposure to proteins or nucleic acids, incorporation of genetic vectors, incorporation of genetic or non-genetic cell-tracing moieties, de-differentiation (i.e., generation of induced pluripotent stem cells or equivalents), cryopreservation. In some examples, subject-specific cells are isolated from a tissue other than the target tissue. In some examples, the subject-specific cells require differentiation into cell types of interest within the target tissue. In some examples, subject-specific cells that require differentiation are differentiated prior to, during, or after fabrication into a three-dimensional structure.

Methods of culturing cells

[0107] The cell types used in the engineered liver tissues of the invention are suitably cultured in any manner known in the art. Methods of cell and tissue culturing are known in the art, and are described, for example, in Freshney, R., *Culture of Animal Cells: A Manual of Basic Techniques*, Wiley (1987).

[0108] General mammalian cell culture techniques, cell lines, and cell culture systems suitably used in conjunction with the present invention are also described in Doyle, A., Griffiths, J. B., Newell, D. G., (eds.) *Cell and Tissue Culture: Laboratory Procedures*, Wiley (1998).

[0109] Appropriate growth conditions for mammalian cells in culture are well known in the art. Cell culture media generally include essential nutrients and, optionally, additional elements such as growth factors, salts, minerals, vitamins, platelet-rich plasma, etc., that are optionally selected according to the cell type(s) being cultured. In some examples, particular ingredients are selected to enhance cell growth, differentiation, secretion of specific proteins, etc. In general, standard growth media include Dulbecco's Modified Eagle Medium (DMEM) with 4 g/L glucose, supplemented with 1-20% fetal bovine serum (FBS), calf serum, or human serum, insulin (4 micrograms/mL), dexamethasone (1.0 micromolar), amphotericin B (0.25 micrograms/mL) and Glutamax (1×). Williams E media supplemented with insulin (1g/L), sodium selenite (0.0067 g/L), transferrin (0.55 g/L), dexamethasone (1 micromolar), penicillin (100U/mL), streptomycin (0.1mg/mL), amphotericin B (0.25 micrograms/mL) and Glutamax (1×), or various other media known in the art. Preferably cells are cultured under sterile conditions in an atmosphere of 1-21% O₂ and preferably 3-5% CO₂, at a temperature at or near the body temperature of the animal of origin of the cell. For example, human cells are preferably cultured at approximately 37°C.

[0110] The cells are optionally cultured with cellular differentiation agents to induce differentiation of the cell along the desired line. For instance, cells are optionally cultured with growth factors, cytokines, etc. In some examples, the term "growth factor" refers to a protein, a polypeptide, or a complex of polypeptides, including cytokines that are produced by a cell and affect itself and/or a variety of other neighboring or distant cells. Typically growth factors affect the growth and/or differentiation of specific types of cells, either developmentally or in response to a multitude of physiological or environmental stimuli. Some, but not all, growth factors are hormones. Exemplary growth factors are insulin, insulin-like growth factor (IGF), nerve growth factor (NGF), vascular endothelial growth factor (VEGF), keratinocyte growth factor (KGF), fibroblast growth factors (FGFs), including basic FGF (bFGF), platelet-derived growth factors (PDGFs), including PDGF-AA and PDGF-AB, hepatocyte growth factor (HGF), transforming growth factor alpha (TGF-α), transforming growth factor beta (TGF-β), including TGFβ1 and TGFβ3, epidermal growth factor (EGF), granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte colony-stimulating factor (G-CSF), interleukin-6 (IL-6), IL-8, and the like. Growth factors are discussed in, among other places, *Molecular Cell Biology*, Scientific

American Books, Darnell et al., eds., 1986; Principles of Tissue Engineering, 2d ed., Lanza et al., eds., Academic Press, 2000. The skilled artisan will understand that any and all culture-derived growth factors in the conditioned media described herein may be suitable.

Bio-ink and multicellular aggregates

[0111] Disclosed herein, are three-dimensional living tissues, including liver tissues/constructs, arrays thereof, and methods that comprise bioprinted cells. In some examples, cells are bioprinted by depositing or extruding bio-ink from a bioprinter. In some examples, "bio-ink" includes liquid, semi-solid, or solid compositions comprising a plurality of cells. In some examples, bio-ink comprises liquid or semi-solid cell solutions, cell suspensions, or cell concentrations. In some examples, a cell solution, suspension, or concentration comprises a liquid or semi-solid (e.g., viscous) carrier and a plurality of cells. In some examples, the carrier is a suitable cell nutrient media, such as those described herein. In some examples, bio-ink comprises a plurality of cells that optionally cohere into multicellular aggregates prior to bioprinting. In some examples, bio-ink comprises a plurality of cells and is bioprinted to produce a specific planar and/or laminar geometry; wherein cohesion of the individual cells within the bio-ink takes place before, during and/or after bioprinting. In some examples, the bio-ink is produced by 1) collecting a plurality of cells in a fixed volume; wherein the cellular component(s) represent at least about 30% and at most 100% of the total volume. In some examples, bio-ink comprises semi-solid or solid multicellular aggregates or multicellular bodies. In some examples, the bio-ink is produced by 1) mixing a plurality of cells or cell aggregates and a biocompatible liquid or gel in a pre-determined ratio to result in bio-ink, and 2) compacting the bio-ink to produce the bio-ink with a desired cell density and viscosity. In some examples, the compacting of the bio-ink is achieved by centrifugation, tangential flow filtration ("TFF"), or a combination thereof. In some examples, the compacting of the bio-ink results in a composition that is extrudable, allowing formation of multicellular aggregates or multicellular bodies. In some examples, "extrudable" means able to be shaped by forcing (e.g., under pressure) through a nozzle or orifice (e.g., one or more holes or tubes). In some examples, the compacting of the bio-ink results from growing the cells to a suitable density. The cell density necessary for the bio-ink will vary with the cells being used and the tissue or organ being produced. In some examples, the cells of the bio-ink are cohered and/or adhered. In some examples, "cohere," "cohered," and "cohesion" refer to cell-cell adhesion properties that bind cells, multicellular aggregates, multicellular bodies, and/or layers thereof. In some examples, the terms are used interchangeably with "fuse," "fused," and "fusion." In some examples, the bio-ink additionally comprises support material, cell culture medium (or supplements thereof), extracellular matrix (or components thereof), cell adhesion agents, cell death inhibitors, anti-apoptotic agents, anti-oxidants, extrusion compounds, and combinations thereof.

[0112] In some examples not claimed, the cells are any suitable cell. In some examples,

the cells are vertebrate cells, mammalian cells, human cells, or combinations thereof. In some examples not claimed, the type of cell used in a method disclosed herein depends on the type

of construct or tissue being produced. In some examples, the bio-ink comprises one type of cell (also referred to as a "homogeneous" or "monotypic" bio-ink). In some examples,

the bio-ink comprises more than one type of cell (also referred to as a "heterogeneous" or "polytypic" bio-ink).

Cell culture media

[0113] In some examples, the bio-ink comprises a cell culture medium. The cell culture medium is any suitable medium. In some examples, suitable cell culture media include, Dulbecco's Phosphate Buffered Saline, Earle's Balanced Salts, Hanks' Balanced Salts, Tyrode's Salts, Alsever's Solution, Gey's Balanced Salt Solution, Kreb's-Henseleit Buffer Modified, Kreb's-Ringer Bicarbonate Buffer, Puck's Saline, Dulbecco's Modified Eagle's Medium, Dulbecco's Modified Eagle's Medium/Nutrient F-12 Ham, Nutrient Mixture F-10 Ham (Ham's F-10), Medium 199, Minimum Essential Medium Eagle, RPMI-1640 Medium, Ames' Media, BGJb Medium (Fitton-Jackson Modification), Click's Medium, CMRL-1066 Medium, Fischer's Medium, Glasgow Minimum Essential Medium (GMEM), Iscove's Modified Dulbecco's Medium (IMDM), L-15 Medium (Leibovitz), McCoy's 5A Modified Medium, NCTC Medium, Swim's S-77 Medium, Waymouth Medium, William's Medium E, or combinations thereof. In some examples, the cell culture medium is modified or supplemented. In some examples, the cell culture medium further comprises albumin, selenium, transferrins, fetuins, sugars, amino acids, vitamins, growth factors, cytokines, hormones, antibiotics, lipids, lipid carriers, cyclodextrins, platelet-rich plasma, or a combination thereof.

Extracellular matrix

[0114] In some examples, the bio-ink further comprises one or more components of an extracellular matrix or derivatives thereof. In some examples, "extracellular matrix" includes proteins that are produced by cells and transported out of the cells into the extracellular space, where they serve as a support to hold tissues together, to provide tensile strength, and/or to facilitate cell signaling. Examples, of extracellular matrix components include, collagens, fibronectin, laminins, hyaluronates, elastin, and proteoglycans. For example, the multicellular aggregates contain various ECM proteins (e.g., gelatin, fibrinogen, fibrin, collagens, fibronectin, laminins, elastin, and/or proteoglycans). The ECM components or derivatives of ECM components are optionally added to the cell paste used to form the multicellular aggregate. The ECM components or derivatives of ECM components added to the cell paste are optionally purified from a human or animal source, or produced by recombinant methods known in the art. Alternatively, the ECM components or derivatives of ECM components are naturally secreted by the cells in the elongate cellular body, or the cells used to make the elongate cellular body are optionally genetically manipulated by any suitable method known in the art to

vary the expression level of one or more ECM components or derivatives of ECM components and/or one or more cell adhesion molecules or cell-substrate adhesion molecules (e.g., selectins, integrins, immunoglobulins, and adherins). In some examples, the ECM components or derivatives of ECM components promote cohesion of the cells in the multicellular aggregates. For example, gelatin and/or fibrinogen is suitably added to the cell paste, which is used to form multicellular aggregates. The fibrinogen is converted to fibrin by the addition of thrombin.

[0115] In some examples, the bio-ink further comprises an agent that encourages cell adhesion.

[0116] In some examples, the bio-ink further comprises an agent that inhibits cell death (e.g., necrosis, apoptosis, or autophagocytosis). In some examples, the bio-ink further comprises an anti-apoptotic agent. Agents that inhibit cell death include, small molecules, antibodies, peptides, peptibodies, or combination thereof. In some examples,

the agent that inhibits cell death is selected from: anti-TNF agents, agents that inhibit the activity of an interleukin, agents that inhibit the activity of an interferon, agents that inhibit the activity of an GCSF (granulocyte colony-stimulating factor), agents that inhibit the activity of a macrophage inflammatory protein, agents that inhibit the activity of TGF-B (transforming growth factor B) (see, e.g., **Figs. 15 and 16**), agents that inhibit the activity of an MMP (matrix metalloproteinase), agents that inhibit the activity of a caspase, agents that inhibit the activity of the MAPK/JNK signaling cascade, agents that inhibit the activity of a Src kinase, agents that inhibit the activity of a JAK (Janus kinase), or a combination thereof. In some examples,

the bio-ink comprises an anti-oxidant. In some embodiments, the bio-ink comprises oxygen-carriers or other cell-specific nutrients.

Extrusion compounds

[0117] In some examples, the bio-ink further comprises an extrusion compound (i.e., a compound that modifies the extrusion properties of the bio-ink). Examples of extrusion compounds include, gels, hydrogels, peptide hydrogels, amino acid-based gels, surfactant polyols (e.g., Pluronic F-127 or PF-127), thermo-responsive polymers, hyaluronates, alginates, extracellular matrix components (and derivatives thereof), collagens, gelatin, other biocompatible natural or synthetic polymers, nano fibers, and self-assembling nanofibers. In some examples, extrusion compounds are removed after bioprinting by physical, chemical, or enzymatic means.

[0118] Gels, sometimes referred to as jellies, have been defined in various ways. For example, the United States Pharmacopoeia defines gels as semisolid systems consisting of either suspensions made up of small inorganic particles or large organic molecules interpenetrated

by a liquid. Gels include a single-phase or a two-phase system. A single-phase gel consists of organic macromolecules distributed uniformly throughout a liquid in such a manner that no apparent boundaries exist between the dispersed macromolecules and the liquid. Some single-phase gels are prepared from synthetic macromolecules (e.g., carbomer) or from natural gums (e.g., tragacanth). In some examples, single-phase gels are generally aqueous, but will also be made using alcohols and oils. Two-phase gels consist of a network of small discrete particles.

[0119] Gels, in some cases, are classified as being hydrophobic or hydrophilic. In some examples, the base of a hydrophobic gel consists of a liquid paraffin with polyethylene or fatty oils gelled with colloidal silica, or aluminum or zinc soaps. In contrast, the base of hydrophilic gels usually consists of water, glycerol, or propylene glycol gelled with a suitable gelling agent (e.g., tragacanth, starch, cellulose derivatives, carboxyvinylpolymers, and magnesium-aluminum silicates). In some examples, the rheology of the compositions or devices disclosed herein is pseudo plastic, plastic, thixotropic, or dilatant.

[0120] Suitable hydrogels include those derived from collagen, hyaluronate, hyaluronan, fibrin, alginate, agarose, chitosan, and combinations thereof. In other embodiments, suitable hydrogels are synthetic polymers. In some examples, suitable hydrogels include those derived from poly(acrylic acid) and derivatives thereof, poly(ethylene oxide) and copolymers thereof, poly(vinyl alcohol), polyphosphazene, and combinations thereof. In some examples, the confinement material is selected from: hydrogel, NovoGelTM, agarose, alginate, gelatin, MatrigelTM, hyaluronan, poloxamer, peptide hydrogel, poly(isopropyl n-polyacrylamide), polyethylene glycol diacrylate (PEG-DA), hydroxyethyl methacrylate, polydimethylsiloxane, polyacrylamide, poly(lactic acid), silicon, silk, or combinations thereof.

[0121] In some examples, hydrogel-based extrusion compounds are thermoreversible gels (also known as thermo-responsive gels or thermogels). In some examples, a suitable thermoreversible hydrogel is not a liquid at room temperature. In some examples, the gelation temperature (T_{gel}) of a suitable hydrogel is about 10°C, 11°C, 12°C, 13°C, 14°C, 15°C, 16°C, 17°C, 18°C, 19°C, 20°C, 21°C, 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 31°C, 32°C, 33°C, 34°C, 35°C, 36°C, 37°C, 38°C, 39°C, 40°C, including increments therein. In some examples, the T_{gel} of a suitable hydrogel is about 10°C to about 40°C. In some examples, the T_{gel} of a suitable hydrogel is about 20°C to about 30°C. In some examples, the bio-ink (e.g., comprising hydrogel, one or more cell types, and other additives, etc.) described herein is not a liquid at room temperature. In some examples, a suitable thermoreversible hydrogel is not a liquid at mammalian body temperature. In some examples, the gelation temperature (T_{gel}) of a suitable hydrogel is about 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 31°C, 32°C, 33°C, 34°C, 35°C, 36°C, 37°C, 38°C, 39°C, 40°C, 41°C, 42°C, 43°C, 44°C, 45°C, 46°C, 47°C, 48°C, 49°C, 50°C, 51°C, 52°C, including increments therein. In some examples, the T_{gel} of a suitable hydrogel is about 22°C to about 52°C. In some examples, the T_{gel} of a suitable hydrogel is about 32°C to about 42°C. In some examples, the bio-ink (e.g., comprising hydrogel, one or more cell types, and other additives, etc.) described herein is not a liquid at mammalian body temperature. In some examples, the gelation temperature (T_{gel}) of a bio-ink described herein is about 10°C, about 15°C, about 20°C, about 25°C, about 30°C, about 35°C,

about 40°C, about 45°C, about 50°C, about 55°C, including increments therein. In some examples, the Tgel of a bio-ink described herein is about 10°C to about 15°C. In some examples, the Tgel of a bio-ink described herein is about 15°C to about 20°C. In some examples, the Tgel of a bio-ink described herein is about 20°C to about 25°C. In some examples, the Tgel of a bio-ink described herein is about 25°C to about 30°C. In some examples, the Tgel of a bio-ink described herein is about 30°C to about 35°C. In some examples, the Tgel of a bio-ink described herein is about 35°C to about 40°C. In some examples, the Tgel of a bio-ink described herein is about 40°C to about 45°C. In some examples, the Tgel of a bio-ink described herein is about 45°C to about 50°C.

[0122] Polymers composed of polyoxypropylene and polyoxyethylene form thermoreversible gels when incorporated into aqueous solutions. These polymers have the ability to change from the liquid state to the gel state at temperatures maintainable in a bioprinter apparatus. The liquid state-to-gel state phase transition is dependent on the polymer concentration and the ingredients in the solution.

[0123] Poloxamer 407 (Pluronic F-127 or PF-127 or Lutrol) is a nonionic surfactant composed of polyoxyethylene-polyoxypropylene copolymers. Other poloxamers include 188 (F-68 grade), 237 (F-87 grade), 338 (F-108 grade). Aqueous solutions of poloxamers are stable in the presence of acids, alkalis, and metal ions. PF-127 is a commercially available polyoxyethylene-polyoxypropylene triblock copolymer of general formula E106 P70 E106, with an average molar mass of 13,000. The polymer is optionally further purified by suitable methods that will enhance gelation properties of the polymer. It contains approximately 70% ethylene oxide, which accounts for its hydrophilicity. It is one of the series of poloxamer ABA block copolymers. PF-127 has good solubilizing capacity, low toxicity and is, therefore, considered a suitable extrusion compound.

[0124] In some examples, the viscosity of the hydrogels and bio-inks presented herein is measured by any means described. For example, in some embodiments, an LVDV-II+CP Cone Plate Viscometer and a Cone Spindle CPE-40 is used to calculate the viscosity of the hydrogels and bio-inks. In some examples, a Brookfield (spindle and cup) viscometer is used to calculate the viscosity of the hydrogels and bio-inks. In some examples, the viscosity ranges referred to herein are measured at room temperature. In some examples, the viscosity ranges referred to herein are measured at body temperature (e.g., at the average body temperature of a healthy human).

[0125] In some examples, the hydrogels and/or bio-inks are characterized by having a viscosity of between about 500 and 1,000,000 centipoise, between about 750 and 1,000,000 centipoise; between about 1000 and 1,000,000 centipoise; between about 1000 and 400,000 centipoise; between about 2000 and 100,000 centipoise; between about 3000 and 50,000 centipoise; between about 4000 and 25,000 centipoise; between about 5000 and 20,000 centipoise; or between about 6000 and 15,000 centipoise.

[0126] In some examples, the bio-ink comprises cells and extrusion compounds suitable for

continuous bioprinting. In some examples, the bio-ink has a viscosity of about 1500 mPa·s. In some examples, a mixture of Pluronic F-127 and cellular material is suitable for continuous bioprinting. Such a bio-ink is suitably prepared by dissolving Pluronic F-127 powder by continuous mixing in cold (4 °C) phosphate buffered saline (PBS) over 48 hours to 30% (w/v). Pluronic F-127 is also suitably dissolved in water. In some examples, cells are cultivated and expanded using standard sterile cell culture techniques. In some examples, the cells are pelleted at 200g for example, and re-suspended in the 30% Pluronic F-127 and aspirated into a reservoir affixed to a bioprinter where it is, allowed to solidify at a gelation temperature from about 10 to about 25°C. Gelation of the bio-ink prior to bioprinting is optional. The bio-ink, including bio-ink comprising Pluronic F-127 is optionally dispensed as a liquid.

[0127] In some examples, the concentration of Pluronic F-127 is any value with suitable viscosity and/or cytotoxicity properties. In some examples, a suitable concentration of Pluronic F-127 is able to support weight while retaining its shape when bioprinted. In some examples, the concentration of Pluronic F-127 is about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, or about 50%. In some examples, the concentration of Pluronic F-127 is between about 30% and about 40%, or between about 30% and about 35%.

[0128] In some examples, the concentration of gelatin is 6% and the concentration of alginate is 0.5% in the extrusion compound or excipient. In some examples, the concentration of gelatin is 4% and the concentration of alginate is 2% in the extrusion compound or excipient.

[0129] In some examples, the non-cellular components of the bio-ink (e.g., extrusion compounds, etc.) are removed prior to use. In some examples, the non-cellular components are, for example, hydrogels, peptide hydrogels, amino acid-based gels, surfactant polyols, thermo-responsive polymers, hyaluronates, alginates, collagens, or other biocompatible natural or synthetic polymers. In some examples, the non-cellular components are removed by physical, chemical, or enzymatic means. In some examples, a proportion of the non-cellular components remain associated with the cellular components at the time of use.

[0130] In some examples, the cells are pre-treated to increase cellular interaction. For example, cells are suitably incubated inside a centrifuge tube after centrifugation in order to enhance cell-cell interactions prior to shaping the bio-ink.

Exemplary cell ratios

[0131] In some examples, the bio-ink comprises multicellular bodies, which further comprise non-parenchymal liver cells (e.g. endothelial cells, hepatic stellate cells, Kupffer cells). In further embodiments, the ratio of endothelial cells to hepatic stellate cells to Kupffer cells is any suitable ratio. In some examples, the ratio of endothelial cells to hepatic stellate cells to Kupffer cells is about 90:10:0 to about 10:90:0. In some examples, the ratio of endothelial cells to hepatic stellate cells to Kupffer cells is 45:45: 10.

[0132] In some examples, the bio-ink comprises multicellular bodies, which further comprise hepatocytes (e.g., primary hepatocytes, HepG2 or HepaRG) and additional cell types at a ratio of 100:0. In some examples, the ratio of hepatocytes to additional, non-parenchymal cell types (e.g., endothelial cells) is 95:5. In some examples, the ratio of hepatocytes to additional, non-parenchymal cell types (e.g., endothelial cells, stellate cells) is 50:50. In some examples, the ratio of parenchymal to non-parenchymal cells (e.g., endothelial cells, stellate cells, Kupffer cells) is 50:35:10:5).

Self-sorting of cells

[0133] In some examples, the multicellular aggregates used to form the construct or tissue comprises all cell types to be included in the engineered tissue (e.g., hepatocytes, endothelial cells, hepatic stellate cells and Kupffer cells); in such an example each cell type migrates to an appropriate position (e.g., during maturation) to form the engineered tissue, such as an engineered liver tissue. In some examples, the multicellular aggregates used to form the structure comprises fewer than all the cell types to be included in the engineered tissue. In some examples, the cells of each type are uniformly distributed within a multicellular aggregate, or region or layer of the tissue. In some examples, the cells of each type localize to particular regions within a multicellular aggregate or layers or regions of the tissue.

[0134] For example, in the case of an engineered liver tissue comprising hepatocytes, endothelial cells, hepatic stellate cells and Kupffer cells, in a suitable ratio (e.g., 70:15:10:5), neighboring, bioprinted cohered polytypic bio-ink units fuse. During maturation, endothelial cells localize to some extent to both the periphery and core of the construct and organize into microvascular structures. In some examples, the localization of cell types within a construct mimics the layered structure of *in vivo* or *ex vivo* mammalian tissues.

Pre-formed scaffold

[0135] Disclosed herein are engineered, implantable tissues that are free or substantially free of any pre-formed scaffold. In some examples, "scaffold" refers to synthetic scaffolds such as polymer scaffolds and porous hydrogels, non-synthetic scaffolds such as pre-formed extracellular matrix layers, dead cell layers, and decellularized tissues, and any other type of pre-formed scaffold that is integral to the physical structure of the engineered tissue and/or organ and not removed from the tissue and/or organ.

[0136] In some examples, decellularized tissue scaffolds include decellularized native tissues or decellularized cellular material generated by cultured cells in any manner; for example, cell layers that are allowed to die or are decellularized, leaving behind the ECM they produced while living.

[0137] In some examples, the engineered liver tissues/constructs and arrays thereof do not utilize any pre-formed scaffold, e.g., for the formation of the tissue, any layer of the tissue, or formation of the tissue's shape. As an example, the engineered liver tissues of the present invention do not utilize any pre-formed, synthetic scaffolds such as polymer scaffolds, pre-formed extracellular matrix layers, or any other type of pre-formed scaffold at the time of manufacture or at the time of use. In some examples, the engineered liver tissues are substantially free of any pre-formed scaffolds. In some examples, the cellular components of the tissues contain a detectable, but trace or trivial amount of scaffold, e.g., less than 2.0%, less than 1.0%, or less than 0.5% of the total composition. In some examples, trace or trivial amounts of scaffold are insufficient to affect long-term behavior of the tissue, or array thereof, or interfere with its primary biological function. In some examples, scaffold components are removed post-printing, by physical, chemical, or enzymatic methods, yielding an engineered tissue that is free or substantially-free of scaffold components.

[0138] In some examples, the engineered liver tissues free, or substantially free, of pre-formed scaffold disclosed herein are in stark contrast to those developed with certain other methods of tissue engineering in which a scaffolding material is first formed, and then cells are seeded onto the scaffold, and subsequently the cells proliferate to fill and take the shape of the scaffold for example. In some examples, the methods of bioprinting described herein allow production of viable and useful tissues that are free or substantially free of pre-formed scaffold.

[0139] In some examples, the cells held in a desired three-dimensional shape using a confinement material. The confinement material is distinct from a scaffold at least in the fact that the confinement material is temporary and/or removable from the cells and/or tissue.

Arrays

[0140] In some embodiments, disclosed herein are arrays of engineered liver tissues/constructs. In some examples, an "array" is a scientific tool including an association of multiple elements spatially arranged to allow a plurality of tests to be performed on a sample, one or more tests to be performed on a plurality of samples, or both. In some examples, the arrays are adapted for, or compatible with, screening methods and devices, including those associated with medium- or high-throughput screening. In some examples, an array allows a plurality of tests to be performed simultaneously. In some examples, an array allows a plurality of samples to be tested simultaneously. In some examples, the arrays are cellular microarrays. In some examples, a cellular microarray is a laboratory tool that allows for the multiplex interrogation of living cells on the surface of a solid support. In some examples, the arrays are tissue microarrays. In some examples, tissue microarrays include a plurality of separate tissues or tissue samples assembled in an array to allow the performance of multiple biochemical, metabolic, molecular, or histological analyses.

[0141] In some examples, the engineered liver tissues/constructs each exist in a well of a

biocompatible multi-well container (see, e.g., **Fig. 16**). In some examples, each tissue is placed into a well. In some examples, each tissue is bioprinted into a well. In some examples,

the wells are coated. In some examples, the wells are coated with one or more of: a biocompatible hydrogel, one or more proteins, one or more chemicals, one or more peptides, one or more antibodies, and one or more growth factors, including combinations thereof. In some examples, the wells are coated with NovoGel[™]. In some examples, the wells are coated with agarose. In some examples, each tissue exists on a porous, biocompatible membrane within a well of a biocompatible multi-well container. In some examples,

each well of a multi-well container contains two or more tissues.

[0142] In some examples, the engineered liver tissues/constructs are secured to a biocompatible surface on one or more sides. Many methods are suitable to secure a tissue to a biocompatible surface. In some examples, a tissue is suitably secured to a biocompatible surface, for example, along one or more entire sides, only at the edges of one or more sides, or only at the center of one or more sides. In some examples, a tissue is suitably secured to a biocompatible surface with a holder or carrier integrated into the surface or associated with the surface. In some examples, a tissue is suitably secured to a biocompatible surface with one or more pinch-clamps or plastic nubs integrated into the surface or associated with the surface. In some examples, a tissue is suitably secured to a biocompatible surface by cell-attachment to a porous membrane. In some examples, the engineered liver tissues/constructs are held in an array configuration by affixation to a biocompatible surface on one or more sides. In some examples, the tissue is affixed to a biocompatible surface on 1, 2, 3, 4, or more sides. In some examples, the biocompatible surface any surface that does not pose a significant risk of injury or toxicity to the tissue or an organism contacting the tissue. In some examples, the biocompatible surface is any surface suitable for traditional tissue culture methods. Suitable biocompatible surfaces include, treated plastics, membranes, porous membranes, coated membranes, coated plastics, metals, coated metals, glass, treated glass, and coated glass, wherein suitable coatings include hydrogels, ECM components, chemicals, proteins, etc., and coatings or treatments provide a means to stimulate or prevent cell and tissue adhesion to the biocompatible surface.

[0143] In some examples, securing of an engineered tissue to a biocompatible surface on one or more sides facilitates subjecting the tissue to shear force, caused by fluid flow.

[0144] In some examples, the engineered liver tissues/constructs, are subjected to shear force, caused by fluid flow. In some examples, the engineered liver tissues are subjected to shear force on 1, 2, 3, 4, or more sides. In some examples, the engineered liver tissues/constructs are subjected to recirculation, perfusion, or agitation of the liquid nutrients that contact the tissues on one or more exposed surfaces.

[0145] In some examples, the arrays of engineered tissues, including liver tissues/constructs,

comprise an association of two or more elements. In some examples, the arrays comprise an association of 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, or 500 elements, including increments therein. In some examples, each element comprises one or more cells, multicellular aggregates, tissues, organs, or combinations thereof.

[0146] In some examples, the arrays of engineered tissues, including liver tissues/constructs, comprise multiple elements spatially arranged in a pre-determined pattern. In some examples, the pattern is any suitable spatial arrangement of elements. In some examples, patterns of arrangement include, a two-dimensional grid, a three-dimensional grid, one or more lines, arcs, or circles, a series of rows or columns, and the like. In some examples, the pattern is chosen for compatibility with medium- or high-throughput biological assay or screening methods or devices.

[0147] In some examples, the cell types and/or source of the cells used to fabricate one or more tissues in an array are selected based on a specific research goal or objective. In some examples, the specific tissues in an array are selected based on a specific research goal or objective. In some examples, one or more specific engineered liver tissues are included in an array to facilitate investigation of a particular disease or condition. In some examples, one or more specific engineered liver tissues are included in an array to facilitate investigation of a disease or a condition of a particular subject. In some examples, one or more specific engineered liver tissues within the array are generated with one or more cell types derived from two or more distinct human donors. In some examples, each tissue within the array is substantially similar with regard to cell types, sources of cells, layers of cells, ratios of cells, methods of construction, size, shape, and the like. In some examples, one or more of the tissues within the array is unique with regard to cell types, sources of cells, layers of cells, ratios of cells, methods of construction, size, shape, and the like. In some examples, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, or more of the tissues within the array, including increments therein, is/are unique. In some examples, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% of the tissues within the array, including increments therein, is/are unique.

[0148] In some examples, each tissue within the array is maintained independently in culture. In some examples, the culture conditions of each tissue within the array are such that they are isolated from the other tissues and cannot exchange media or factors soluble in the media. In some examples, two or more individual tissues within the array exchange soluble factors. In some examples, the culture conditions of two or more individual tissues within the array are such that they exchange media and factors soluble in the media with other tissues. In some examples, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, or more of the tissues within the array, including increments therein, exchange media and/or soluble factors. In some examples, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92,

93, 94, 95, 96, 97, 98, 99, or 100% of the tissues within the array, including increments therein, exchange media and/or soluble factors.

In vitro assays

[0149] In some examples, the engineered liver tissues and arrays disclosed herein are for use in *in vitro* assays. In some examples, an "assay" is a procedure for testing or measuring the presence or activity of a substance (e.g., a chemical, molecule, biochemical, drug, etc.) in an organic or biologic sample (e.g., cell aggregate, tissue, organ, organism, etc.). In some examples, assays include qualitative assays and quantitative assays. In some examples, a quantitative assay measures the amount of a substance in a sample.

[0150] In some examples, the engineered liver tissues and arrays are for use in, image-based assays, measurement of secreted proteins, expression of markers, and production of proteins. In some examples, the engineered liver tissue and arrays are for use in assays to detect or measure one or more of: molecular binding (including radioligand binding), molecular uptake, activity (e.g., enzymatic activity and receptor activity, etc.), gene expression, protein expression, receptor agonism, receptor antagonism, cell signaling, apoptosis, chemosensitivity, transfection, cell migration, chemotaxis, cell viability, cell proliferation, safety, efficacy, metabolism, toxicity, infectivity, and abuse liability.

[0151] In some examples, the engineered liver tissues and arrays are for use in immunoassays. In some examples, immunoassays are competitive immunoassays or noncompetitive immunoassays. In a competitive immunoassay, for example, the antigen in a sample competes with labeled antigen to bind with antibodies and the amount of labeled antigen bound to the antibody site is then measured. In a noncompetitive immunoassay (also referred to as a "sandwich assay"), for example, antigen in a sample is bound to an antibody site; subsequently, labeled antibody is bound to the antigen and the amount of labeled antibody on the site is then measured.

[0152] In some examples, the engineered liver tissue and arrays are for use in enzyme-linked immunosorbent assays (ELISA). In some examples, an ELISA is a biochemical technique used to detect the presence of an antibody or an antigen in a sample. In ELISA, for example, at least one antibody with specificity for a particular antigen is utilized. By way of further example, a sample with an unknown amount of antigen is immobilized on a solid support (e.g., a polystyrene microtiter plate) either non-specifically (via adsorption to the surface) or specifically (via capture by another antibody specific to the same antigen, in a "sandwich" ELISA). By way of still further example, after the antigen is immobilized, the detection antibody is added, forming a complex with the antigen. The detection antibody is, for example, covalently linked to an enzyme, or is itself detected by a secondary antibody that is linked to an enzyme through bioconjugation.

[0153] For example, an array, microarray, or chip of cells, multicellular aggregates, or tissues

is used for drug screening or drug discovery. In some examples, an array, microarray, or chip of tissues is used as part of a kit for drug screening or drug discovery. In some examples, each engineered liver tissue/construct exists within a well of a biocompatible multi-well container, wherein the container is compatible with one or more automated drug screening procedures and/or devices. In some examples, automated drug screening procedures and/or devices include any suitable procedure or device that is computer or robot-assisted.

[0154] In some examples, arrays for drug screening assays or drug discovery assays are used to research or develop drugs potentially useful in any therapeutic area. In some examples, suitable therapeutic areas include infectious disease, hematology, oncology, pediatrics, cardiology, central nervous system disease, neurology, gastroenterology, hepatology, urology, infertility, ophthalmology, nephrology, orthopedics, pain control, psychiatry, pulmonology, vaccines, wound healing, physiology, pharmacology, dermatology, gene therapy, toxicology, and immunology.

[0155] In some examples, the engineered liver tissue and arrays are for use in cell-based screening. In some examples, the cell-based screening is for one or more infectious diseases such as viral infection or parasitic infection (e.g., plasmodium infection, etc.) . In some examples, the cell-based screening is for liver fibrosis (e.g., cirrhosis). In some examples,

the cell-based screening is for liver cancer. In some examples, the cell-based screening is for liver steatosis (e.g., fatty liver). In some examples, the cell-based screening is for one or more metabolic deficiencies. In some examples, the cell-based screening is for one or more protein deficiencies. In some examples, the engineered liver tissues and arrays are for use in the study of cancer initiation, progression, or metastasis. In some examples,

the engineered liver tissues and arrays are for use in the study of the interaction of other cell types, such as cancer cells, pathogen-bearing cells, pathogenic cells, immune cells, blood-derived cells, or stem/progenitor cells, with liver tissue and the cells comprising liver tissue.

[0156] In some examples, the constructs or arrays thereof are for use in assessing the performance of biologics, including antibodies, mammalian cells, bacteria, biologically-active proteins, hormones, etc. In some examples, the construct or arrays thereof are for use to detect, quantify, and study immunologic sampling by Kupffer and stellate cells, including the effects of gram-negative or gram-positive antigen-stimulated signaling from Kupffer or stellate cells to bordering hepatocytes (e.g., response to lipopolysaccharide (LPS)). In some examples, the constructs or arrays thereof are useful for the production of hepatotropic viruses, including HBV and HCV. In some examples, the constructs or arrays thereof are used as a vehicle for production of *Plasmodium spp.* (a parasite) in the exo-erythrocytic form. In some examples, liver construct-derived Plasmodium are utilized in comparative *in vitro* assays to identify effective therapies for exo-erythrocytic forms of the parasite. In some examples, the construct or arrays thereof are utilized as anti-host therapies for Hepatitis C Virus (HCV) or Plasmodium, including the conduct of anti-host antibody studies. In some examples, the liver

constructs or arrays thereof are useful in the study of cancer initiation, progression, or metastasis. In some examples, the liver constructs or arrays thereof are useful in the study of cell-cell and cell-tissue interactions between the mammalian liver cells/tissue comprising the construct and one or more additional cell types, including pathogen-bearing cells, living pathogenic cells, cancer cells, immune cells, blood cells, stem/progenitor cells, or genetically-manipulated cells.

[0157] In some examples, the array comprises engineered liver tissue constructs and additional tissue constructs. In some examples, the liver tissue construct is in direct contact with an additional tissue construct on one or more surfaces. In some examples, the liver tissue is connected to one or more additional tissues constructs or cells via a fluid path or common fluid reservoir. In some examples, the liquid media that contacts the engineered liver tissue construct contains living mammalian cells such as immune cells, blood-derived cells, or tumor-derived cells. In some examples, the liquid media that contacts the engineered liver tissue construct contains bacteria, viruses, parasites, or other pathogens. In some examples, the engineered liver tissue and arrays are for use as vehicle to propagate hepatotropic viruses for three-dimensional structural studies including X-ray or cryo-EM.

Extracorporeal support

[0158] In some examples, the engineered liver tissues comprise a plurality of layers, each layer comprising cylindrical bio-ink, the bio-ink axially-aligned substantially in parallel, the bio-ink comprising parenchymal liver cells; and optionally, non-parenchymal cells among the cylindrical bio-ink; and optionally, void spaces or perfusable channels among the cylindrical bio-ink. In some examples, an engineered liver tissue is used to provide extracorporeal support to a subject in need thereof. In some examples, blood is passed over or through the construct to filter the blood and supplement the subjects liver function.

[0159] Referring to **Fig. 17**, in some examples, a bioprinted liver structure is designed for providing extracorporeal support. In this example, a bioprinted liver structure is composed of a plurality of parenchymal cell cylinders. The structure also includes non-parenchymal cells and filler bodies to create compartments and/or channels through which perfusion or flow could be achieved.

[0160] Referring to **Fig. 19**, a bioprinted liver structure is used as an extracorporeal hepatic device containing repeating tubular structures with void spaces (depicted occupied by temporary filler bodies) at regular intervals allowing perfusion across and/or through the engineered liver tissue.

Methods

[0161] Disclosed herein, but not claimed are methods for constructing living, three-dimensional liver tissue constructs, the methods comprising the steps of bioprinting bio-ink comprising at least one liver cell type into or onto a form, and fusing of the bio-ink into a living, three-dimensional tissue liver construct. In some examples not claimed, the tissue construct is for *in vitro* use.

[0162] Also disclosed herein, but not claimed are methods of constructing tissues, including engineered liver tissues, comprising the steps of preparing cohered multicellular aggregates comprising either parenchymal or parenchymal and non-parenchymal cells; placing said cohered multicellular aggregates onto a support; and incubating said multicellular aggregates to allow them to cohere and form an engineered liver tissue; wherein said incubation has a duration of about 1 hours to about 30 days. In some examples not claimed, the methods utilize bioprinting. In some examples not claimed, the methods produce engineered liver tissues free or substantially free of any pre-formed scaffold at the time of use.

[0163] Also disclosed herein, but not claimed are methods of constructing living, three-dimensional liver tissues comprising the steps of preparing one or more cohered multicellular aggregates comprising mammalian liver cells; placing said one or more cohered multicellular aggregates onto a support; applying, to said one or more cohered multicellular aggregates, one or more of a layer of a first type of mammalian cells on one or more external surfaces; a layer of a second type of mammalian cells on one or more external surfaces; and incubating said one or more multicellular aggregates to allow them to cohere and to form a tissue; wherein said incubation has a duration of about 1 hour to about 30 days. In some examples not claimed the methods utilize bioprinting. In some examples not claimed the methods produce engineered liver tissues, free or substantially free of any pre-formed scaffold at the time of use.

[0164] Also disclosed herein, but not claimed are methods of constructing living, three-dimensional liver tissue constructs comprising the steps of preparing one or more cohered multicellular aggregates comprising mammalian cells; placing said one or more cohered multicellular aggregates onto a support to form at least one of at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and/or a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry; and incubating said one or more multicellular aggregates to allow them to cohere and to form a living, three-dimensional liver tissue construct.

Preparing cohered multicellular aggregates

[0165] In some examples not claimed the methods involve preparing cohered multicellular aggregates comprising one or more types of mammalian cells, wherein at least one cellular component represents liver cells. In some examples not claimed the methods involve preparing cohered multicellular aggregates comprising liver parenchymal cells. In some

examples not claimed the methods involve preparing cohered multicellular aggregates further comprising non-parenchymal cells. In some examples, the methods involve preparing cohered multicellular aggregates further comprising one or more other cell types, selected from the group: endothelial cells, stellate cells, fibroblasts, Kupffer cells, immune cells, lymphocytes, cancer cells, virus-bearing cells, pathogenic cells, adipocytes, adipogenic cells, smooth muscle cells, or stem/progenitor-derived cells.

[0166] There are various ways to make multicellular aggregates having the characteristics described herein. In some examples, a multicellular aggregate is fabricated from a cell paste containing a plurality of living cells or with a desired cell density and viscosity.

[0167] In some examples, the cell paste is shaped into a desired shape and a multicellular body formed through maturation (e.g., incubation). In some examples, the shape of the multicellular body is determined by the shape of a surrounding mold or frame. In some examples, the mold or frame is fabricated by an automated instrument in a defined geometry. In some examples not claimed the mold or frame consists of bio-ink and contains liver-derived cells. In some examples, the mold or frame comprises liver-derived non-parenchymal cells and the paste utilized to fill the frame and create the multicellular aggregate comprises parenchymal cells. In some examples, the multicellular aggregates are substantially cylindrical. In some examples, the multicellular aggregates are hexagonal, square, cuboidal, rectangular, polyhedral. In some examples, multiple multicellular aggregates are formed by fabricating the surrounding mold or frame in a tessellated pattern of repeating geometry and subsequently or simultaneously filling the molds or frames with cell paste. In some examples, the cell paste is incubated in a controlled environment to allow the cells to adhere and/or cohere to one another to form the elongate multicellular body.

[0168] In some examples, a multicellular body is produced by shaping a cell paste including a plurality of living cells in a device that holds the cell paste in a three-dimensional shape. In some examples, the cell paste is incubated in a controlled environment while it is held in the three-dimensional shape for a sufficient time to produce a body that has sufficient cohesion to support itself on a flat surface.

[0169] In some examples, a cell paste is provided by: 1) collecting cells or cell aggregates (of one or more cell types) and a biocompatible gel or liquid, such as cell culture medium (e.g., in a pre-determined ratio) to result in a cell suspension, and 2) compacting the cellular suspension to produce a cell paste with a desired cell density and viscosity. In some examples, compacting is achieved by a number of methods, such as by concentrating a particular cell suspension that resulted from cell culture to achieve the desired cell concentration (density), viscosity, and consistency required for the cell paste. In some examples, a relatively dilute cell suspension from cell culture is centrifuged for a determined time to achieve a cell concentration in the pellet that allows shaping in a mold. Tangential flow filtration ("TFF") is another suitable method of concentrating or compacting the cells. In some examples, compounds are combined with the cell suspension to lend the extrusion properties required. Suitable compounds include, surfactant polyols, collagens, hydrogels, peptide hydrogels,

amino acid-based gels, Matrigel[™], nanofibers, self-assembling nanofibers, gelatin, fibrinogen, etc.

[0170] In some examples, the cell paste is produced by mixing a plurality of living cells with a tissue culture medium, and compacting the living cells (e.g., by centrifugation). One or more ECM components (or derivative of an ECM component) is optionally included by, resuspending the cell pellet in one or more physiologically acceptable buffers containing the ECM component(s) (or derivative(s) of ECM component(s)) and the resulting cell suspension centrifuged again to form a cell paste.

[0171] In some examples, the cell density of the cell paste desired for further processing varies with cell types. In some examples, interactions between cells determine the properties of the cell paste, and different cell types will have a different relationship between cell density and cell-cell interaction. In some examples, the cells are pre-treated to increase cellular interactions before shaping the cell paste. For example, in some cases, cells are incubated inside a centrifuge tube after centrifugation in order to enhance cell-cell interactions prior to shaping the cell paste. In some examples, the cell paste is shaped concomitantly with bioprinting; wherein the cohesion of individual cells to each other to form bio-ink occurs during or after bioprinting.

[0172] In some examples, many methods are used to shape the cell paste. For example, the cell paste is manually molded or pressed (e.g., after concentration/compaction) to achieve a desired shape. By way of a further example, the cell paste is taken up (e.g., aspirated) into an instrument, such as a micropipette or syringe (e.g., a capillary pipette or a glass/plastic syringe), that shapes the cell paste to conform to an interior surface of the instrument. The cross-sectional shape of the micropipette (e.g., capillary pipette) is alternatively circular, square, rectangular, triangular, or other non-circular cross-sectional shape. In some examples, the cell paste is shaped by depositing it into a preformed mold, such as a plastic mold, metal mold, or a gel mold. In some examples, centrifugal casting or continuous casting is used to shape the cell paste. In some examples, the shaping of the bio-ink occurs concomitantly or after bioprinting. In some examples, the shaping of the bio-ink occurs as the result of a co-printed mold; wherein the mold is optionally deposited via bioprinting; wherein the mold comprises one or more of bio-ink, bio-ink that further comprises an extrusion compound, gel, hydrogel, synthetic polymer, carbohydrate, protein, or mammalian cells, or combinations thereof. In some examples, one or more components of the co-printed mold are removed after bioprinting; wherein the removal method is selected from one of physical means, solubilization with aqueous media; chemical treatment; enzymatic treatment; modulating temperature. In some examples, the co-printed mold remains associated with the tissue after fabrication. In some examples, only the cellular component(s) of the co-printed mold remain associated with the tissue after fabrication.

[0173] In some examples, multicellular aggregates of a defined shape are also suitable to build the engineered liver tissues described herein. Spherical multicellular aggregates are optionally generated by a variety of methods, including, cellular self-assembly, the use of molds, and hanging drop methods, in some examples not claimed, a method to produce substantially

spherical multicellular aggregates comprises the steps of 1) providing a cell paste containing a plurality of pre-selected cells or cell aggregates with a desired cell density and viscosity, 2) manipulating the cell paste into a cylindrical shape, 3) cutting cylinders into equal fragments, 4) optionally letting the fragments round up overnight on a gyratory shaker, and 5) optionally allowing the substantially spherical multicellular aggregates to mature over a period of 1 hour to 7 days. In some examples, multicellular aggregates are generated via acoustic focusing methodologies.

[0174] In some examples, a partially adhered and/or cohered cell paste is used for bioprinting; wherein cohesion and bio-ink formation occurs primarily post-printing. In some examples, the cellular paste is shaped in a first step prior to bioprinting. In some examples, the cell paste is transferred from the first shaping device (e.g., capillary pipette) to a second shaping device (e.g., a mold) that allows nutrients and/or oxygen to be supplied to the cells while they are retained in the second shaping device for an additional maturation period. One example of a suitable shaping device that allows the cells to be supplied with nutrients and oxygen is a mold for producing a plurality of multicellular aggregates (e.g., substantially identical multicellular aggregates). By way of further example, such a mold includes a biocompatible substrate made of a material that is resistant to migration and ingrowth of cells into the substrate and resistant to adherence of cells to the substrate. In some examples, the substrate is suitably made of Teflon[®] (PTFE), stainless steel, NovoGel[™], agarose, polyethylene glycol, glass, metal, plastic, or gel materials (e.g., agarose or other hydrogels), and similar materials. In some examples, the mold is also suitably configured to allow supplying tissue culture media to the cell paste (e.g., by dispensing tissue culture media onto the top of the mold).

[0175] Thus, in some examples, where a second shaping device is used, the partially adhered and/or cohered cell paste is transferred from the first shaping device (e.g., a capillary pipette) to the second shaping device (e.g., a mold). In some examples, the partially adhered and/or cohered cell paste is transferred by the first shaping device (e.g., the capillary pipette) into the grooves of a mold. In some examples, following a maturation period in which the mold is incubated along with the cell paste retained therein in a controlled environment to allow the cells in the cell paste to further adhere and/or cohere to one another to form the multicellular aggregate, the cohesion of the cells will be sufficiently strong to allow the resulting multicellular aggregate to be picked up with an implement (e.g., a capillary pipette). In some examples, the capillary pipette is suitably be part of a printing head of a bioprinter or similar apparatus operable to automatically place the multicellular aggregate into a three-dimensional construct.

[0176] In some examples, the cross-sectional shape and size of the multicellular aggregates will substantially correspond to the cross-sectional shapes and sizes of the first shaping device and optionally the second shaping device used to make the multicellular aggregates, and the skilled artisan will be able to select suitable shaping devices having suitable cross-sectional shapes, cross-sectional areas, diameters, and lengths suitable for creating multicellular aggregates having the cross-sectional shapes, cross-sectional areas, diameters, and lengths discussed above.

Placing cohered multicellular aggregates onto a support

[0177] A number of methods are suitable to place multicellular aggregates on a support to produce a desired three-dimensional structure. For example, the multicellular aggregates are manually placed in contact with one another, deposited in place by extrusion from a pipette, nozzle, or needle, or positioned by an automated, computer-assisted device such as a bioprinter.

[0178] As described herein, multicellular aggregates have many suitable shapes and sizes. In some examples, multicellular aggregates are elongate with any of several suitable cross-sectional shapes including, circular, oval, square, triangular, polygonal, and irregular. In some examples, multicellular aggregates are elongate and in the form of a cylinder. In some examples, elongate multicellular aggregates are of similar lengths and/or diameters. In some examples, elongate multicellular aggregates are of differing lengths and/or diameters. In some examples, multicellular aggregates are substantially spherical. In some examples, the engineered liver tissues include substantially spherical multicellular aggregates that are substantially similar in size. In some examples, the engineered liver tissues include substantially spherical multicellular aggregates that are of differing sizes. In some examples, engineered liver tissues of different shapes and sizes are formed by arranging multicellular aggregates of various shapes and sizes.

[0179] In some examples, the cohered multicellular aggregates are placed onto a support. In some examples, the support is any suitable biocompatible surface. In some examples, suitable biocompatible surfaces include, polymeric material, porous membranes, plastic, glass, metal, hydrogel, and combinations thereof. In some examples, the support is coated with a biocompatible substance including, a hydrogel, a protein, a chemical, a peptide, antibodies, growth factors, or combinations thereof. In some examples, the support is coated with NovoGel™. In some examples, the support is coated with agarose. In some examples, the cohered multicellular aggregates are placed into the wells of a biocompatible multi-well container.

[0180] Once placement of the cohered multicellular aggregates is complete,

a tissue culture medium is poured over the top of the construct. In some examples,

the tissue culture medium enters the spaces between the multicellular bodies to support the cells in the multicellular bodies.

Applying a layer of a first type of cells and/or a layer of a second type of cells

[0181] A number of methods are suitable to apply one or more layers of cells on one or more external surfaces of the cohered mammalian cell construct. For example, applying a layer of cells comprises coating one or more surfaces of said cohered multicellular aggregates with a suspension, sheet, monolayer, or fused aggregates of cells. In some examples, 1, 2, 3, 4, or more surfaces of the cohered mammalian cell construct are coated.

[0182] In some examples, applying a layer of cells comprises bioprinting an additional layer of fused multicellular aggregates. In some examples, applying a layer of cells comprises bioprinting, spraying, or ink-jetting a solution, suspension, or liquid concentrate of cells. In some examples, a suitable cell suspension comprises about 1×10^4 to about 1×10^6 cells/ μL . In some examples, a suitable cell suspension comprises about 1×10^5 to about 1.5×10^5 cells/ μL . In some examples, applying a layer of cells comprises dispensing a suspension of cells directly onto one or more surfaces of the cohered mammalian cell construct as spatially-distributed droplets. In some examples, applying a layer of cells comprises dispensing a suspension of cells directly onto one or more surfaces of the cohered mammalian cell construct as a spray. Layers of cells are, in some examples, applied at any suitable time in the construction process. In some examples, one or more layers of cells are applied on one or more external surfaces of the cohered mammalian cell construct immediately after bioprinting (e.g., up to 10 min.). In some examples, one or more layers are applied after bioprinting (e.g., after 10 min.). In some examples, one or more layers are applied during maturation of the construct.

[0183] In some examples, the methods further comprise the step of culturing a layer of cells on a support. In some examples, applying a layer of cells, in some cases, comprises placing one or more surfaces of the engineered liver tissue construct in direct contact with an established culture of cells. In some examples, the construct is bioprinted directly onto a cultured layer of cells or a monolayer of cells. Any type of cultured cell layer on a biocompatible support is suitable. In some examples, multicellular aggregates are bioprinted onto a layer of endothelial cells. In some examples, multicellular aggregates are bioprinted onto a layer of non-parenchymal cells. In some examples, the layer of cells adheres and/or coheres with the multicellular aggregates of the bioprinted construct. In some examples, each layer of a multi-layered structure are bioprinted. In some examples, the individual layers comprise variable forms of bio-ink, including but not limited to: cohered cell aggregates, cell paste, cell paste in combination with extrusion compound(s) or other additives, cell monolayers, and cell sheets.

Co-printed molds used to fabricate compartmentalized tissues

[0184] In some examples, methods comprise providing one or more bio-inks comprising non-parenchymal cells; providing one or more bio-inks comprising parenchymal cells, such as hepatocytes or hepatocyte-like cells; depositing the bio-inks onto a support; and incubating the deposited bio-inks for a duration of about 1 hour to about 30 days to form a living, three-dimensional liver tissue construct comprising at least one compartment, the compartment

comprising an interior comprising parenchymal cells confined by a border comprising non-parenchymal cells, wherein the parenchymal and non-parenchymal cells are as claimed

[0185] In some examples, the methods comprise providing one or more cohered multicellular aggregates comprising mammalian liver cells as claimed; placing the one or more cohered multicellular aggregates onto a support to form at least one of at least one layer comprising a plurality of cell types, the cell types spatially arranged relative to each other to create a planar geometry; and a plurality of layers, at least one layer compositionally or architecturally distinct from at least one other layer to create a laminar geometry; and incubating said one or more multicellular aggregates for a duration of about 1 hour to about 30 days to allow them to cohere and to form a living, three-dimensional liver tissue construct.

[0186] Referring to **Fig. 2**, liver constructs containing HepaRG cells were bio-printed, matured, and biochemically characterized. First, a perimeter box with dimensions of 4mm X 4mm X 1mm was printed with bio-ink containing endothelial cells and hepatic stellates. Second, HepRG cells were deposited into the center of the perimeter box. This was followed by maturation for 96 hours at 37°C. Third, fusion of the bio-printed construct was observed at 20 hours post print. In some examples, the bio-printed liver constructs were metabolically active producing albumin and cholesterol at 96 hours.

[0187] Referring to **Fig. 3**, liver constructs are bioprinted with borders of non-parenchymal cells and fills of parenchymal cells each made with an extrusion compound.

[0188] Referring to **Fig. 4**, liver constructs are bioprinted with a border including endothelial cells and hepatic stellates, a fill including primary hepatocytes, and a third compositional component introduced as spatially-dispersed spheroids including endothelial cells and hepatic stellates within the fill.

[0189] Referring to **Fig. 5**, liver constructs are bioprinted with a co-mold fabricated from a material that does not contain cells and dissolves in an aqueous media (e.g., PF-127) leaving only fills and establishing a negative space.

[0190] Referring to **Fig. 6**, liver constructs are bioprinted using a continuous deposition technique using bio-ink composed of multiple liver cell types encapsulated in a water-soluble extrusion compound (e.g., PF-127) to form a tessellated functional unit pattern.

[0191] Referring to **Fig. 8**, compartmentalized liver constructs are bioprinted with a co-printed mold that does not contain cells used to shape multicellular aggregates into hexagons. In some examples, the non-cellular material disappears to yield a scaffold-free tissue at the time of use.

[0192] Referring to **Fig. 10**, compartmentalized liver constructs are bioprinted with a co-mold used as a border which is filled with HepG2 cells.

[0193] Referring to **Fig. 11**, a co-printed mold containing cells is bio-printed as a border into

which parenchymal cells are used as a fill, in this case, HepG2 cells.

[0194] Referring to **Fig. 12**, compartmentalized liver constructs are bioprinted using a non-parenchymal border and a parenchymal fill to achieve a high, tissue-like density.

Incubating multicellular aggregates

[0195] In some examples, the multicellular aggregates are incubated. In some examples, the incubation allows the multicellular aggregates adhere and/or cohere to form a tissue, such as a liver tissue. In some examples, the multicellular aggregates cohere to form a tissue in a cell culture environment (e.g., a Petri dish, cell culture flask, bioreactor, etc.). In some examples, the multicellular aggregates cohere to form a tissue in an environment with conditions suitable to facilitate growth of the cell types included in the multicellular aggregates. In some examples, the multicellular aggregates are incubated at about 37°C, in a humidified atmosphere containing about 5% CO₂, in the presence of cell culture medium containing factors and/or ions to foster adherence and/or coherence. In some examples, the multicellular aggregates are maintained in an environment that contains 0.1% to 21% O₂.

[0196] The incubation, has any suitable duration. In some examples, the incubation has a duration of about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, or more minutes, including increments therein. In some examples, the incubation has a duration of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 36, 48, or more hours, including increments therein. In some examples, the incubation has a duration of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more days, including increments therein. Several factors influence the time required for multicellular aggregates to cohere to form a tissue including cell types, cell type ratios, culture conditions, and the presence of additives such as growth factors.

Additional steps for increasing viability of the engineered tissue

[0197] In some examples, the method further comprises steps for increasing the viability of the engineered tissue. In some examples, these steps involve providing direct contact between the tissue and a nutrient medium through a temporary or semi-permanent lattice of confinement material. In some examples, the tissue is constrained in a porous or gapped material. Direct access of at least some of the cells of the engineered tissue to nutrients increases the viability of the engineered tissue.

[0198] In some examples, the additional and optional steps for increasing the viability of the engineered tissue include: 1) optionally dispensing base layer of confinement material prior to placing cohered multicellular aggregates; 2) optionally dispensing a perimeter of confinement material; 3) bioprinting cells of the tissue within a defined geometry; and 4) dispensing

elongate bodies (e.g., cylinders, ribbons, etc.) of confinement material overlaying the nascent tissue in a pattern that introduces gaps in the confinement material, such as a lattice, mesh, or grid.

[0199] Many confinement materials are suitable for use in the methods described herein. In some examples, hydrogels are exemplary confinement materials possessing one or more advantageous properties including: non-adherent, biocompatible, extrudable, bioprintable, non-cellular, of suitable strength, and not soluble in aqueous conditions. In some embodiments, suitable hydrogels are natural polymers. In some examples, the confinement material is comprised of NovoGel™. In some examples, suitable hydrogels include those derived from surfactant polyols such as Pluronic F-127, collagen, hyaluronate, fibrin, alginate, agarose, chitosan, gelatin, and derivatives or combinations thereof. In some examples, suitable hydrogels are synthetic polymers. In some examples, suitable hydrogels include those derived from poly(acrylic acid) and derivatives thereof, poly(ethylene oxide) and copolymers thereof, poly(vinyl alcohol), polyphosphazene, and combinations thereof. In some examples, the confinement material is selected from: hydrogel, NovoGel™, agarose, alginate, gelatin, Matrigel™, hyaluronan, poloxamer, peptide hydrogel, poly(isopropyl n-polyacrylamide), polyethylene glycol diacrylate (PEG-DA), hydroxyethyl methacrylate, polydimethylsiloxane, polyacrylamide, poly(lactic acid), silicon, silk, and combinations thereof.

[0200] In some examples, the gaps overlaying pattern are distributed uniformly or substantially uniformly around the surface of the tissue. In some examples, the gaps are distributed non-uniformly, whereby the cells of the tissue are exposed to nutrients non-uniformly. In non-uniform examples, the differential access to nutrients is optionally exploited to influence one or more properties of the tissue. For instance, in some cases, it is desirable to have cells on one surface of a bioprinted tissue proliferate faster than cells on another surface of the bioprinted tissue. In some examples, the exposure of various parts of the tissue to nutrients is changed at various times to influence the development of the tissue toward a desired endpoint.

[0201] In some examples, the confinement material is removed at any suitable time, including, immediately after bioprinting (e.g., within 10 minutes), after bioprinting (e.g., after 10 minutes), before the cells are substantially cohered to each other, after the cells are substantially cohered to each other, before the cells produce an extracellular matrix, after the cells produce an extracellular matrix, just prior to use, and the like. In some examples, confinement material is removed by any suitable method. For example, in some embodiments, the confinement material is excised, pulled off the cells, digested, or dissolved.

[0202] In some examples, the methods further comprise the step of subjecting the engineered liver tissue to shear force, caused by fluid flow, agitation, or convective recirculation, on one or more sides.

Biological properties

[0203] The bioprinted liver tissues described herein included a multicellular structure resembling human tissue. Like native liver tissue, bioprinted liver tissues included specific cell types (e.g., hepatic stellate cells, hepatocytes, endothelial cells, etc.) enriched in specific zones, yielding a patterned, compartmentalized architecture.

[0204] In some examples, the bioprinted liver tissues described herein include important features of native liver including, high cell density, and production of ECM, lipids, and glycogen. See **Fig. 21** In some examples, the bioprinted liver tissues described herein have cell density of at least 30%. In some examples, the bioprinted liver tissues described herein have cell density of at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90%. In some examples, the bioprinted liver tissues described herein have cell density of about 30%, about 40%, about 50%, about 60%, about 70%, about 80% or about 90%.

[0205] In some examples, the three-dimensional bioprinted liver tissues include histological features such as tight junctions, maintenance of at least some quiescent hepatic stellate cells, and development of microvascular structures. In some examples, the three-dimensional bioprinted liver tissues include compartmentalization that persists well beyond initial fabrication. For example, the three-dimensional bioprinted liver tissues include compartmentalization that persists for at least 7 days, at least 10 days, at least 20 days, at least 30 days, or at least 40 days. In some examples, the three-dimensional bioprinted liver tissues include compartmentalization that persists for about 7 days, about 10 days, about 20 days, about 30 days, or about 40 days.

[0206] In some examples, post-fabrication, regions of hepatocytes surrounded by non-parenchymal "stroma" can be identified. See **Fig. 22**. Unexpectedly the bioprinted liver tissues demonstrated apparent self-organization into lobule type structure, closely mimicking native liver. The observed architecture supports function and viability for 40+ days, enabling long term testing, e.g., low-dose, repeat exposure and chronic pathological conditions.

[0207] In some examples, liver-specific functions are sustained for more than 7, 10, 20, 30, or 40 days in the three-dimensional bioprinted liver tissues described herein. For example, the bioprinted liver tissues described herein demonstrate continued production of albumin out to 6 weeks (42 days), well beyond published time course of primary hepatocytes in traditional two-dimensional technologies. See **Fig. 23**. The bioprinted liver tissues also demonstrate sustained production of other hepatocyte factors like cholesterol. See **Fig. 23**.

[0208] The three-dimensional bioprinted human liver tissues exhibit production of additional relevant metabolites not achieved in existing two-dimensional methodologies including transferring and fibrinogen. See **Fig. 24**. In some examples, the three-dimensional bioprinted human liver also exhibit markers of bile canilicular formation. See **Fig. 25**.

[0209] In contrast to two-dimensional primary hepatocytes, which lose function after a few days, the bioprinted three-dimensional liver tissue constructs described herein show sustained

inducibility of active CYP3A4 enzyme, wherein the inducibility is inhibitable by Verapamil. See **Fig. 26**. In some examples, the three-dimensional bioprinted liver tissues described herein demonstrate native-like responses to known liver toxicants including diclofenac, trovafloxacin, and methotrexate. See **Figs. 27-29**. In some examples, the three-dimensional bioprinted liver tissues described herein demonstrate native-like responses to physiologically relevant molecules such as vitamin D. See **Fig. 30**.

[0210] In some examples, the three-dimensional human liver tissues described herein are patterned with parenchymal (hepatocyte) and non-parenchymal regions (endothelial cells (EC) and hepatic stellate cells (hSC)). In some examples, as an alternative to primary hepatocytes, the patterned three-dimensional liver tissues include iPSC-derived hepatocyte-like cells (iPSC-HC) (such as iCell Hepatocytes, Cellular Dynamics Inc.). Use of iPSC allows for development of patient- and/or disease-specific *in vitro* systems, and may provide a means of generating cells that are difficult to isolate or propagate.

EXAMPLES

[0211] The following specific examples are to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

[0212] Any of the following examples that do not fall within the scope of the claims do not form part of the invention.

Example 1 - Liver tissue bioprinted using continuous deposition and tessellated functional unit structure

[0213] Engineered liver tissue was bioprinted utilizing a NovoGen MMX Bioprinter™ (Organovo, Inc., San Diego, CA) using a continuous deposition mechanism. The three-dimensional structure of the liver tissue was based on a repeating functional unit, in this case, a hexagon. The bio-ink was composed of hepatic stellate cells and endothelial cells encapsulated in an extrusion compound (surfactant polyol - PF-127).

Preparation of 30% PF-127

[0214] A 30% PF-127 solution (w/w) was made using PBS. PF-127 powder was mixed with chilled PBS using a magnetic stir plate maintained at 4°C. Complete dissolution occurred in approximately 48 hours.

Cell Preparation and bioprinting

[0215] A cell suspension comprised of 82% stellate cells (SC) and 18% human aortic endothelial cells (HAEC) and human adult dermal fibroblasts (HDFa) was separated into 15 mL tubes in order to achieve three cell concentrations: 50×10^6 cells/mL, 100×10^6 cells/mL, and 200×10^6 cells/mL following centrifugation. Each cell pellet was resuspended in 30% PF-127 and aspirated into a 3 cc reservoir using the bioprinter. With a 510 μ m dispense tip, the encapsulated cells were bioprinted onto a PDMS base plate into a single hexagon (see **Fig. 6A**) or hexagon tessellation configuration (see **Fig. 6B**). Each construct received approximately 200 μ L of media and was incubated for 20 minutes at room temperature to evaluate construct integrity.

Multi-layer bioprinting

[0216] For hexagon tessellation experiments, up to (4) sequential layers were bioprinted resulting in a taller structure with more cellular material present. Following fabrication, each construct initially received approximately 200 μ L of complete media to assess construct integrity. Constructs were incubated for 20 minutes at room temperature and were then submerged in 20 mLs of complete media.

Results

[0217] Following 18 hours of culture in growth media containing 10% fetal bovine serum (which dissolves PF127), cells contained within the bioprinted geometry were cohered to each other sufficiently to generate an intact, contiguous sheet of tissue that retained the geometrical patterning of the original design (see **Fig. 6D**). Shown in **Fig. 7** is H&E staining of a single segment of the tessellated construct, after fixation in 10% neutral buffered formalin. Cells were found to be viable, intact, and confined to their original printed geometry.

Example 2 - Forced layering

[0218] Cell populations (homogeneous or heterogeneous) were prepared for bioprinting as either cylindrical bio-ink or as a cell suspension in Pluronic F-127 (Lutrol, BASF). Briefly, for preparation of bio-ink, cells were liberated from standard tissue culture plastic using either recombinant human trypsin (75 μ g/mL, Roche) or 0.05% trypsin (Invitrogen). Following enzyme liberation, cells were washed, collected, counted and combined at desired ratios (i.e., 50:50 hepatic stellate cell (hSC):endothelial cell (EC)) and pelleted by centrifugation. Supernatant was then removed from the cell pellet and the cell mixture was aspirated into a glass microcapillary of desired diameter, typically 500 μ m or 250 μ m, internal diameter. This cylindrical cell preparation was then extruded into a mold, generated from non cell-adherent

hydrogel material with channels for bio-ink maturation. The resulting bio-ink cylinders were then cultured in complete cell culture media for an empirically determined amount of time, typically 2 to 24 hours.

[0219] Briefly, for hydrogel cell suspension preparation, cells were liberated from standard cell culture vessel using either of the enzyme-mediated protocols described herein. Liberated cells were then washed with serum containing media, collected, counted and centrifuged to form a dense cell pellet. Supernatant was removed from the resulting cell pellet and cells were then resuspended in cold PF-127 (4°C) at a concentration of 50 to 200 × 10⁶ cells/mL (ranging from 10 to 300 × 10⁶ cells/mL). This cell suspension was then aspirated into a syringe, utilizing a NovoGen MMX Bioprinter™ (Organovo, Inc., San Diego, CA).

[0220] Fabrication of tissue constructs with forced cell patterning, layering, or orientation was then accomplished using the bioprinter. Bioprinting of three-dimensional tissue constructs was performed with cylindrical bio-ink, cellular suspensions in water soluble gels, or combinations thereof. To achieve defined cell patterning or layering, combinations of relevant cell populations were included in the bio-ink or cell suspension preparation and then bioprinted in such a fashion that dissolution of the gel material supporting the cell solution, results in defined cell layering around the deposited bio-ink (see **Fig. 13**). Cell patterning, organization, or layering was also achieved through the utilization and incorporation of defined, discrete cell populations (e.g., hSC and EC), which resulted in predictable and repeatable organization of cells and cellular structures within the bioprinted tissues (see **Fig. 14**).

[0221] In some experiments, final cellular organization within the bioprinted neotissue was observed after a maturation or culture period. Constructs were maintained in a standard laboratory incubator (37°C, humidified chamber supplemented with 5% CO₂) and evaluated over time.

Results

[0222] Cell patterning, layering, or arrangement was achieved using bioprinting. By bioprinting with bio-ink containing heterogeneous (i.e., polytypic) cell populations, or by combining bio-ink (homogeneous or heterogeneous cell populations) with high density cell-gel or cell suspensions, distinct cell organization was observed. Maturation of these neotissue constructs in a humidified chamber (incubator) resulted in further establishment of distinct cell arrangement, organization and/or segregation in these bioprinted neotissues.

[0223] For example, bioprinting of EC:hSC-laden PF-127 on top of bioprinted bio-ink comprising HepG2 cells results in the establishment of distinct layers of the construct with distinct cell populations and discrete tissue morphology. In the case of bio-ink constructs containing hSC and EC, bioprinted constructs that were matured for more than 3 days in complete media were found to contain a distinct layer of EC at the periphery and organized

microvessel networks within the core of the construct. Bioprinted constructs fabricated with bio-ink comprising a homogeneous (i.e., monotypic) population of vascular smooth muscle cell onto which a highly concentrated solution of EC was bioprinted were found to contain a distinct layer of EC at the periphery of the construct.

Example 3 - Multi-well plates

[0224] Cell populations (homogeneous or heterogeneous) were prepared for bioprinting using a variety of bio-ink formats, including cylindrical bio-ink aggregates, suspensions of cellular aggregates, or cell suspensions/pastes, optionally containing extrusion compounds. Briefly, for preparation of cylindrical bio-ink, cells were liberated from standard tissue culture plastic using either recombinant human trypsin (75 µg/mL, Roche) or 0.05% trypsin (Invitrogen). Following enzyme liberation, cells were washed, collected, counted, and combined at desired ratios (i.e., 50:50 hepatic stellate cell (hSC):endothelial cell (EC)) and pelleted by centrifugation. Supernatant was then removed from the cell pellet and the cell mixture was aspirated into a glass microcapillary of desired diameter, typically 500 µm or 250 µm, internal diameter. This cylindrical cell preparation was then extruded into a mold, generated from non cell-adherent hydrogel material with channels for bio-ink maturation. The resulting bio-ink cylinders were then cultured in complete cell culture media for an empirically determined amount of time, typically 2 to 24 hours.

[0225] For preparation of a cell suspension or cell paste of cellular aggregates, the cell propagation and liberation protocols described herein were followed. At the time of cell pellet generation, supernatant was removed from the pellet and the pellet was transferred to a custom deposition syringe. This syringe was then mounted to the bioprinter deposition head for direct bioprinting of the cell aggregate solution or paste into multi-well plates.

[0226] Replicate tissue constructs were then bioprinted within the wells of either a multi-well tissue culture plate (e.g., 6-well or 24-well) or within a multi-well cell culture insert (i.e., Transwell, BD) and then placed into an appropriate multi-well plate. Following bioprinting, the three-dimensional constructs were matured/conditioned with relevant media for some period of time, typically 3 to 14 days. Following maturation, constructs were harvested, fixed and processed for routine histology and immunohistochemistry.

Results

[0227] Bioprinted tissues were successfully fabricated within multi-well culture plates or multi-well culture inserts that were then inserted into multi-well plates. This approach allows for generation of replicate bioprinted tissues that are optionally cultured and treated to generate identical or unique culture conditions. This approach results in a significant increase in bioprinting process throughput and sample generation (see **Fig. 16**).

Example 4 - Stimulation of bioprinted neotissues

[0228] Cylindrical bio-ink comprising relevant heterogeneous (i.e., polytypic) cell populations were prepared. Physiologically-relevant populations (e.g., hepatic stellate cells (hSC) and human aortic endothelial cells (EC)) of cells were combined at specific ratios to generate proper bio-ink. Cells were maintained and propagated under standard laboratory conditions and cells were cultured in media either purchased from the same vendor as the cells, or media comprising components typically found in the primary literature to be conducive to standard cell culture practices for those particular cell types. Cell processing for bio-ink preparation was as follows: briefly, cells were liberated from standard tissue culture plastic by washing with cation-free phosphate buffered saline (PBS) and then exposed to 0.1-0.05% trypsin (Invitrogen). Liberated cells were then washed in serum-containing media, collected, counted, and combined in an appropriate ratio for the stimulation assay or experiment being conducted, and pelleted by centrifugation. Supernatant was then removed and the cell pellet was aspirated into a glass microcapillary, which was then submerged in complete media for approximately 15 to 20 minutes. This cylindrical bio-ink structure was then extruded into a non cell-adherent hydrogel mold, containing linear channels and held for 2 to 18 hours. Cylindrical bio-ink containing hSC:EC was then used to fabricate bioprinted tissue constructs and maintained and/or matured in a humidified chamber. Tissue segments were fabricated with initial dimensions of 1.25 mm × 8 mm × 0.25 mm (W × L × H). During the post-bioprinting maturation period some constructs were exposed to the cytokine TGF-β1 to elicit a tissue-specific response (see Fig. 15).

[0229] For tissue constructs requiring a homogeneous (i.e., monotypic) cell layer, restricted to the upper surface, a secondary cell preparation was utilized containing the relevant cell type. Typically vascular endothelial cells or small airway epithelial cells (for blood vessel wall and human lung tissue models, respectively) were prepared in a highly concentrated cell suspension. Briefly, cells were liberated as described above, collected, enumerated and pelleted by centrifugation. Supernatant was removed and the cell pellet was resuspended in a small volume of complete media, yielding a highly concentrated cell pellet of 1×10^5 cells/μL. This cell suspension was then stored at 4°C until time of use.

[0230] The bioprinted neotissues were then submerged in serum-containing complete cell culture media and placed in a standard humidified chamber, supplemented with 5% CO₂ for maturation. The bioprinted neotissues were then cultured and stimulated with a relevant cytokine(s) for a predetermined period of time, formalin-fixed, harvested, and processed for standard histology and immunohistochemistry. The bioprinted tissues were evaluated for characteristics such as, but not limited to for tissue morphology, cell organization, extracellular matrix production, cell proliferation, cell viability, and construct integrity.

[0231] Cytokine stimulation was conducted by adding cytokine directly to the culture media and incubating the bioprinted tissues with the added protein to provide direct and prolonged cell

access to the proper stimulus. Dose-response experiments were conducted at doses typically ranging from 0.1 to 100 ng/mL, dependent on the ED50 of the experimental cytokine. For experiments in which cytokine stimulation was conducted over more than 48 hours, media was changed and fresh cytokine was added every 48 hours.

Results

[0232] Bioprinted neotissues containing physiologically-relevant populations of cells were successfully stimulated with cytokines that had been previously demonstrated to elicit cellular responses in two-dimensional *in vitro* systems. The responses observed in the bioprinted three-dimensional tissue constructs were observed to be dose-dependent and unique to the cells within the bioprinted construct (see, e.g., **Fig. 15**).

Example 5 - Bioprinting of co-molded functional liver tissue microstructure with continuous deposition

Preparation of 30% PF-127

[0233] A 30% PF-127 solution (w/w) was made using PBS. PF-127 powder was mixed with chilled PBS using a magnetic stir plate maintained at 4°C. Complete dissolution occurred in approximately 48 hours.

Cell preparation and co-printing of mold and fill

[0234] Three mL of PF-127 solution was aspirated into a 3 cc reservoir using the bioprinter and with a 510 µm dispense tip, 30% PF-127 solution was bioprinted onto a 6 well Transwell into a single hexagon shape and layered sequentially 6 times.

[0235] A cell suspension, comprised of 7.8×10^7 hepatic cells (HepG2), was centrifuged at 1000g for 6 minutes to create the cell paste. Five µL of cell paste was extruded through a 510 µm needle to fill each of the triangular molds (see **Fig. 5A**). The hexagon mold was incubated at room temperature for 15 minutes. Three mL of media (DMEM supplemented with 10% FBS and 1X penicillin, streptomycin and amphotericin B) was added to the well with the Transwell supported above followed by incubation at 37°C and 5% CO₂. Within 45 minutes the PF-127 mold dissolved into the media leaving the molded hepatic bio-ink intact to form a planar geometry of cells and void spaces (see **Fig. 5B**). To remove residual PF-127 from the media, the Transwell was transferred into a new well containing 3 mL of media and incubated for two hours. This was repeated an additional 2 times for a total media exchange of 9 mL over 6

hours.

[0236] Post 6 hours the Transwell was transferred to a new well with no media and a cell suspension of 2×10^6 cells, at a ratio of human aortic endothelial cells at 90% and 10% hepatic stellate cells, was dispensed to fill the voids created by the dissolution of PF-127 mold. The hepatic constructs were incubated for 15 minutes at room temperature. Following the 15 minute incubation, 4 mL of media containing a ratio of 85% media (DMEM supplemented with 10% FBS and 1X penicillin, streptomycin and amphotericin B, to support the hepatic and stellate cells and 15% M199 supplemented with 2% LSGS, 10% FBS, HEPES and 1X penicillin, streptomycin and amphotericin B, to support the human aortic endothelial cells). The construct was incubated at 37°C and 5% CO₂ for 48 hours to form a contiguous construct, with planar geometry comprising a lobular (triangular) arrangement of hepatic parenchyma with intervening endothelial cell-comprising tissue.

Example 6 - Bioprinting and characterization of functional liver tissue

Cell culture

[0237] Cryo-preserved primary human hepatocytes (Life Technologies) were purified per manufacturer's protocols and incubated in Hepatocyte Maintenance Media (Life Technologies) supplemented with penicillin 100 I.U./ml, streptomycin 0.25 µg/ml, and amphotericin 85 µg/ml (1XP/S/A). (Life Technologies). The following cells were utilized in the non-parenchymal cell component of the bioprinted tissues: human umbilical vein endothelial cells (HUVECs; Becton Dickenson) and hepatic stellate cells (ScienCell) and were propagated according the manufacturer's instructions. HUVECs were grown in EGM-2 media (Lonza) and hepatic stellates in DMEM (Corning) supplemented with 10% FBS (Life Technologies), penicillin 100 I.U./ml, streptomycin 0.25 µg/ml, and amphotericin 85 µg/ml (1XP/S/A; Corning).

Chemicals utilized in induction assays

[0238] The following reagents were purchased from Sigma: rifampicin, diclofenac, verapamil, 5-(and-6)-carboxy -2',7' dichlorofluorescein diacetate and vitamin D3.

Bio-ink preparation

[0239] Bio-ink was prepared based on protocols and techniques described herein. For preparation of hydrogels containing cells, NovoGel 2.0 (Organovo, Inc.) was prepared according to manufacturer's instructions and non-parenchymal cell populations were

incorporated prior to bioprinting.

Bioprinting

[0240] All tissues were fabricated directly onto the membranes of standard tissue culture well inserts (Corning Transwell) using standard Organovo bioprinting protocols and the NovoGen MMX Bioprinter or modifications thereof. The bioprinted liver tissues included borders comprising non-parenchymal (hepatic stellates, HUVEC) in Novogel 2.0. The borders defined a compartment comprising an interior. The interiors of the compartments were filled with a cell paste comprising primary hepatocytes. The bioprinted liver tissues also included a moat comprising Novogel 1.0. **Fig. 20** shows a particular example of a bioprinted human liver tissue. Bioprinted liver tissues were maintained on these membranes for the duration of the experiments. The constructs were media exchanged daily with 600µl of a blend of 90% primary hepatocyte media and 10% EGM-2 and incubated at 37°C under humidified atmospheric conditions supplemented with 5%CO₂. Spent media was collected for analysis of secreted factors and the liver tissues were harvested for endpoint analysis (e.g., gene expression, histology, tissue viability, etc.).

Secreted protein detection

[0241] Spent media was collected and stored at -80°C, then analyzed by commercially available ELISA kits for the following liver metabolites: albumin (Bethyl Laboratories), fibrinogen (GenWay Labs), and transferrin (GenWay Labs). Cholesterol biosynthesis was quantified fluorometrically (Cayman Biochemical). Lactate dehydrogenase (LDH) activity was measured colorimetrically with a commercially available reagent (Abcam).

RNA isolation and cDNA synthesis for transcriptional profiling

[0242] Bulk RNA was extracted from the three-dimensional liver constructs using Quick-RNA MiniPrep (Zymo) per manufacturer's instructions. RNA was quantified using a spectrophotometer (Beckman Coulter DU 640). 2µg input RNA was utilized in each individual cDNA synthesis SmartScribe reaction per manufacturer's instructions (CloneTech). TaqMan (Life Technologies) probe and primer pairs for CYP3A4, FAM dye-labeled, and 18S, VIC dye-labeled, were utilized in TaqMan Universal Master Mix UNG per the manufacturer's guidelines (Life Technologies). Transcript amplicons were detected using a Step One Plus system (Life Technologies).

Histological analysis: Paraffin preparation of liver tissues

[0243] Liver constructs were fixed in situ in 2% paraformaldehyde solution (2% paraformaldehyde, 10 mM calcium chloride, 50 mM sucrose in PBS) for 24 hours at room temperature. After 24 hours, the fixation solution was removed and replaced with 70% ethanol. Constructs with the attached Transwell membrane were submerged in 45°C liquid Histogel (American MasterTech, Lodi, CA, catalog # EMHGECS) in a plastic biopsy mold and allowed to solidify at room temperature. Histogel blocks with embedded liver constructs were then processed for paraffin embedding using a tissue processor. Following infiltration with paraffin, liver constructs were embedded in paraffin molds and 5 µm cross-sections were generated using a rotary microtome (Jung Biocut 2035, Leica Microsystems, Buffalo Grove, IL). For hematoxylin and eosin staining, slides were dewaxed in xylene and rehydrated through 100%, 95%, 70%, and 50% ethanol and rinsed in distilled water. Slides were immersed in Gill's hematoxylin (Fisher Scientific, Pittsburgh, PA). Following rinsing with distilled water, slides were briefly immersed in 0.2% v/v ammonium hydroxide. After rinsing with distilled water, slides were immersed in aqueous eosin solution (American MasterTech). Slides were then dehydrated through an ethanol gradient, cleared in xylene, and mounted with resinous mounting media (CytoSeal, Fisher Scientific).

[0244] Periodic acid-Sciff staining was performed according to the manufacturer's instructions (Abcam, Cambridge, MA).

[0245] For immunohistochemical analysis, slides were dewaxed in xylene and rehydrated by sequentially immersing them in 100, 95, 70, and 50% ethanol before finally washing in distilled water. Rehydrated sections were subjected to heat-mediated antigen retrieval in 10 mM sodium citrate pH 6.0 using a standard microwave oven to heat the solution and slides to a subboil followed by slow cooling for 30 minutes. Slides were then blocked with 10% goat serum in Tris-buffered saline (TBS) for 1 hour, followed by incubation with primary antibodies overnight at 4°C. The following primary antibodies were utilized: rabbit anti-E-cadherin (Abcam; 1:50); mouse anti-albumin (Sigma-Aldrich, St. Louis, MO; 1:200); rabbit anti-desmin (Abeam; 1:200); mouse anti-CD31 (Abeam; 1:25). Sections were then washed three times in TBS with 0.1% Tween 20 and incubated with AlexaFluor 488 or AlexaFluor 568-conjugated secondary antibodies (Life Technologies, Carlsbad, CA) diluted 1:200 in TBS. Sections were then washed three times in TBS-0.1% Tween 20, rinsed with distilled water, and mounted with DAPI-containing mounting media (Vector Labs, Burlingame, CA).

Preparation of constructs for cryo sectioning

[0246] Liver constructs were rinsed once with DPBS, immersed in Tissue-Tek OCT compound (Sakura Finetek Europe B.V., Netherlands), and flash frozen. Frozen blocks containing the liver construct were then sectioned at 5 µm on a cryostat (Leica Cryocut 1800, Leica Microsystems). Sectioned slides were snap fixed in -20°C liquid acetone and allowed to air dry for 20 minutes at room temperature. For Oil Red O staining, slides were rehydrated in distilled water and immersed in 60% isopropanol for 2 min. Slides were stained with 0.3% w/v Oil Red O (Sigma-Aldrich) in 60% isopropanol for 15 minutes at room temperature, followed by rinsing

with 60% isopropanol for 1 minute. Slides were immersed briefly in Gill's hematoxylin to counterstain. Slides were rinsed with distilled water and mounted with aqueous media (American MasterTech).

Microscopy

[0247] H&E, PAS, and Oil Red O-stained slides were imaged using a Zeiss Axioskop microscope (Zeiss, Jena, Germany). Images were acquired with an Insight 2 camera and Spot 5.0 software (Diagnostic Instruments, Inc., Sterling Heights, MI). Fluorescently stained slides were imaged with a Zeiss AxioImager microscope and images were acquired with a Zeiss ICM-1 camera and Zen Pro software.

Preparation of S9 fractions

[0248] Media was aspirated from the three-dimensional liver constructs and 400µl of phosphate buffered saline (Corning) supplemented with 1X protease inhibitor cocktail (Sigma) was added to the construct followed by freezing at -80°C. The constructs were thawed and homogenized utilizing a TissuRuptor (Qiagen) and disposable tips at 3000 rpm for 90 sec, followed by a 90 sec pause, followed by a homogenization at 3000 rpm for 90 sec. The homogenized supernatant was centrifuged at 20,000 X g at 4°C for 20 min. The supernatant was transferred to a fresh tube followed by protein quantification.

Protein quantification

[0249] Protein quantification was accomplished utilizing the Bradford assay (Sigma) with bovine albumin as the standard (Sigma).

CYP 3A4 assays

[0250] S9 fractions were normalized by protein content and utilized in Luciferin-IPA assays (CYP ProGlo; Promega) per manufacturer's protocol and luminescence was read utilizing a plate reader (Polarstar, BMG).

Statistics

[0251] ANOVA and students *t*-test statistical analysis were carried out with GraphPad Prism Software.

Results

[0252] The bioprinted liver tissues included a multicellular structure resembling human tissue. Like native liver tissue, bioprinted liver tissues included specific cell types (e.g., hepatic stellate cells, hepatocytes, endothelial cells, etc.) enriched in specific zones, yielding a patterned, compartmentalized architecture.

[0253] Histological characterization of the bioprinted liver tissues shows key features of native liver including, high cell density, and production of ECM, lipids, and glycogen. See Fig. 21 (Day 7 post-printing). The three-dimensional bioprinted liver tissues demonstrated attractive histological features including tight junctions (observed by visualization of E-Cadherin, Albumin, and DAPI), and quiescent stellate cells and microvasculature (observed by visualization of Desmin, CD31, and DAPI). The three-dimensional bioprinted liver tissues demonstrated compartmentalization that continued well beyond initial fabrication. At 21 Days post-fabrication, regions of hepatocytes surrounded by non-parenchymal "stroma" are identified. See **Fig. 22**. Unexpectedly the bioprinted liver tissues demonstrated apparent self-organization into lobule type structure, closely mimicking native liver.

[0254] The observed architecture supports function and viability for 40+ days, enabling long term testing, e.g., low-dose, repeat exposure and chronic pathological conditions. Liver-specific functions are sustained for more than 40 days in the three-dimensional bioprinted liver tissue (primary hepatocytes). The bioprinted liver tissues demonstrated continued production of albumin out to 6 weeks (42 days), well beyond published time course of primary hepatocytes in traditional two-dimensional technologies. See **Fig. 23**. The bioprinted liver tissues also demonstrated sustained production of other hepatocyte factors like cholesterol. See **Fig. 23**.

[0255] The three-dimensional bioprinted human liver tissues exhibit production of additional relevant metabolites not achieved in existing two-dimensional methodologies including transferring and fibrinogen. See **Fig. 24**.

[0256] The three-dimensional bioprinted human liver also exhibit markers of bile canilicular formation. Bile salt exporting protein (BSEP) is a membrane protein that is located within the canilicular membranes of hepatocytes. BSEP's main function is to eliminate unconjugated and conjugated bile acids and salts from hepatocytes. Additionally, BSEP expression is indicative of polarity within native tissue. Histological visualization of BSEP (IHC staining for BSEP protein), albumin, and DAPI demonstrates that BSEP is present in the hepatocyte albumin producing cells. **Fig. 25** shows a calculation of relative quantity of BSEP mRNA vs. housekeeping gene RNA (18s RNA) in two bioprinted liver tissues at Day 10 post-printing. Detectable levels of BSEP support the histology findings.

[0257] In contrast to two-dimensional primary hepatocytes, which lose function after a few days, the bioprinted three-dimensional liver tissue constructs show sustained inducibility of

active CYP3A4 enzyme (inhibitible by Verapamil). See **Fig. 26**.

[0258] The three-dimensional bioprinted liver tissues demonstrate native-like responses to known liver toxicants including diclofenac, trovafloxacin, and methotrexate. **Fig. 27** shows Induction of CYP3A4 at moderate doses of diclofenac, followed by damage at higher doses as measured by lactate dehydrogenase (LDH). **Fig. 28** shows a decrease in albumin and increase in LDH seen in multiple dose study of trovafloxacin vs. levofloxacin (n=3, 2 doses over 4 days). Trovafloxacin effects were seen closer to the clinical C_{max} (~4 μ M). This is in contrast to traditional two-dimensional primary hepatocyte cultures, wherein no induction of LDH seen. **Fig. 29** shows a decrease in viability (measured by ATP - Cell titer glo) and albumin and increase in LDH seen in multiple dose study of methotrexate (n=3, daily dosing for 7 days).

[0259] The three-dimensional bioprinted liver tissues demonstrate native-like responses to physiologically relevant molecules such as vitamin D. Vitamin D is important in maintaining homeostasis within the body. Vitamin D can be acquired through the dietary consumption or synthesized by cells within the skin. The liver processes Vitamin D into metabolites important for bone formation/maintenance and into forms that can be excreted by the kidney. Vitamin D also stimulates the expression of CYP3A4 in the liver and can potentially impact the bioavailability of drugs. Transcription profiling of bioprinted 3D liver tissues treated with 1, 5, 10 μ M of vitamin D was undertaken. Measurement of relative expression of CYP3A4 transcripts was performed. **Fig. 30** shows that CYP3A4 activity was stimulated ~7 and 4 fold for 5 and 10 μ M respectively of vitamin D treatments.

Example 7 - Bioprinting of functional liver tissue incorporating Kupffer cells

[0260] Incorporation of Kupffer cells allows investigation of immunological aspect of Organovo's 3D liver tissues. Kupffer cells were incorporated within the bioprinted liver constructs at a physiologically relevant percentage. The constructs respond in a dose dependent manner to LPS stimulation (LPS: *E. coli* 0127:B8; Sigma). See **Fig. 31**. Kupffer cells were bioprinted into the liver constructs at a physiologically relevant percentage (~12%). The constructs respond in a dose dependent manner to LPS stimulation. IL-10 and TNF- α are opposing cytokines. Expression of IL-10 decreases the expression of TNF- α . This is observed within the bioprinted 3D liver tissues and is most pronounced at the 100 μ g/ml dose of LPS. See **Fig. 32**.

Example 8 - Bioprinting of functional liver tissue incorporating hepatic sinusoidal endothelial cells

[0261] Incorporation of hepatic sinusoidal endothelial cells allows investigation of immunological aspects of the three-dimensional bioprinted liver tissues. Hepatic sinusoidal endothelial cells (HSECs) are a native liver endothelial cell population. **Fig. 33** shows a two-

dimensional monolayer of HSECs. These cells have unique characteristics including cell signaling for immunological responses. HSECs respond to LPS by producing IL-6 in 2D cultures, demonstrated in **Fig. 34**. In contrast, HUVECs produce no detectable amounts of IL-6 when stimulated by LPS. When incorporated into the bioprinting process, HSECs produce fused 3D liver tissue. **Fig. 35** shows a fused three-dimensional liver tissue with HSECs. To produce this tissue hepatic sinusoidal endothelial cells were incorporated into the non-parenchymal fraction produce fused structures.

Example 9 - Small molecule penetration

[0262] An experiment was conducted to determine whether small molecules can penetrate throughout the three-dimensional bioprinted liver tissues. 5-(and-6)-carboxy-2',7'dichlorofluorescein diacetate was supplemented into the media of the 3D liver tissues at 1µM for 24 hrs. The construct was rinsed with 1ml of HBSS buffer and the wash buffer was aspirated. This process was repeated 3X. OCT embedding solution was added and the construct was flash frozen and subjected to cryo-sectioning. 5 micron sections were visualized with the FITC channel at 495 nm excitation. The fluorescein component of the molecule 5-(and-6)-carboxy-2',7'dichlorofluorescein diacetate excites at this wavelength. This section was approximately 150 microns into the construct and fluorescently labeled compound is observed throughout. The formula weight for 5-(and-6)-carboxy -2',7' dichlorofluorescein diacetate is 529.28. **Fig. 36** shows FITC imaging of cryo-sectioned 3D liver tissue at 5X demonstrating the small molecule penetration.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO20130040078A](#) **[0002]**
- [US2003096411A1](#) **[0002]**

Non-patent literature cited in the description

- **PAUL et al.**How to improve R&D productivity: the pharmaceutical industry's grand challengeNature Reviews Drug Discovery, 2010, vol. 9, 3203-214 [0002]
- **FRESHNEY, R**Culture of Animal Cells: A Manual of Basic TechniquesWiley19870000 [0107]
- Cell and Tissue Culture: Laboratory ProceduresWiley19980000 [0108]
- Molecular Cell BiologyScientific American Books19860000 [0110]
- Principles of Tissue EngineeringAcademic Press20000000 [0110]

MANIPULEREDE LEVERVÆV, ARRAYS DERAFT OG FREMGANGSMÅDER TIL FREMSTILLING
HERAF

PATENTKRAV

1. Manipuleret, levende, tredimensional levervævskonstruktion, hvilken konstruktion omfatter mindst
5 ét kompartment, der definerer en planar geometri, hvor kompartmentet omfatter et indre, som defineres af en kant, hvor det indre omfatter parenkymale celler, hvor kanten omfatter ikke-parenkymale celler, hvor cellerne er sammenhængende til dannelse af en levende, tredimensional levervævskonstruktion; forudsat at mindst én komponent af konstruktionen blev fremstillet ved bioprintning, og konstruktionen i det væsentlige er fri for foruddannet skelet på anvendelsestidspunktet; hvor de parenkymale celler er hepatocytter eller
10 hepatocyt lignende celler afledt af én eller flere af følgende kilder: levervæv fra voksne pattedyr; levervæv fra føtale pattedyr; embryoniske stamceller (ESC); ESC-afledte hepatocyt lignende celler, inducerede pluripotente stamceller (iPSC); iPSC-afledte hepatocyt lignende celler; voksne stam-/progenitorceller afledt af leveren; og voksne stam-/progenitorceller afledt af et andet væv end lever; og hvor de ikke-parenkymale celler omfatter én eller flere af: vaskulære celler, endotelceller, fibroblastre, mesenkymale celler,
15 immunceller, Kupffer-celler, stjerneceller, biliære epitelceller, biliære epitellignende celler og sinusoidale endotelceller.
2. Manipuleret, levende, tredimensional levervævskonstruktion ifølge krav 1, der omfatter ét eller flere lag, hvor hvert lag omfatter én eller flere levercelletyper, hvor det ene eller flere lag er sammenhængende til dannelse af den levende, tredimensionale levervævskonstruktion, hvor konstruktionen er kendetegnet ved at
20 have mindst én af:
 - a. mindst ét lag, der omfatter en flerhed af celletyper, hvor celletyperne er rumligt anbragt i forhold til hinanden til dannelse af en planar geometri, hvor celletyperne omfatter parenkymale og ikke-parenkymale celler ifølge krav 1; og
 - b. en flerhed af lag, hvor mindst ét lag kompositorisk eller arkitektonisk adskiller sig fra
25 mindst ét andet lag til dannelse af en laminar geometri.
3. Konstruktion ifølge krav 1, der omfatter en flerhed af lag; fortrinsvis hvor mindst ét lag kompositorisk eller arkitektonisk adskiller sig fra mindst ét andet lag til dannelse af en laminar geometri.
4. Konstruktion ifølge krav 2, der endvidere omfatter cancerceller.
5. Konstruktion ifølge et hvilket som helst af kravene 1 til 4 til anvendelse ved forøgelse af én eller
30 flere leverfunktioner hos mennesker.
6. Array, der omfatter en flerhed af de manipulerede, levende, tredimensionale levervævskonstruktioner ifølge et hvilket som helst af kravene 1 til 4, til anvendelse i *in vitro*-assays.
7. Array ifølge krav 6, hvor *in vitro*-assayet er én eller flere af: lægemiddelforskning; lægemiddeltestning; præklinisk forskning; toksicitetstestning; absorptions-, fordelings-, metabolisme- og
35 udskillelsetestning (ADME); lægemiddelmetabolisme og farmakokinetiktestning (DMPK); sygdomsmodellering; modellering af infektiøse sygdomme; modellering af værtssygdomme; tredimensionale biologistudier; og cellebaseret screening.

8. Fremgangsmåde til fremstilling af en levende, tredimensional levervævskonstruktion, hvilken fremgangsmåde omfatter:
- a. tilvejebringelse af én eller flere bio-blæktyper, der omfatter ikke-parenkymale celler;
 - b. tilvejebringelse af én eller flere bio-blæktyper, der omfatter parenkymale celler;
 - 5 c. afsætning af bio-blæktyperne på en understøtning; og
 - d. inkubering af de afsatte bio-blæktyper i et tidsrum på ca. 1 time til ca. 30 dage til dannelse af en levende, tredimensional levervævskonstruktion, der omfatter mindst ét kompartment, hvilket kompartment omfatter et indre, der omfatter parenkymale celler, som er begrænset af en kant, der omfatter ikke-parenkymale celler;
- 10 hvor de ikke-parenkymale celler omfatter én eller flere af: vaskulære celler, endotelceller, fibroblastre, mesenkymale celler, immunceller, cancerceller, Kupffer-celler, stjerneceller, biliære epitelceller, biliære epitellignende celler og sinusoidale endotelceller; og hvor de parenkymale celler er hepatocytter eller hepatocytllignende celler, der er afledt af én eller flere af følgende kilder: levervæv fra voksne pattedyr; levervæv fra føtale pattedyr; embryoniske stamceller (ESC); ESC-afledte hepatocytllignende celler;
- 15 inducerede pluripotente stamceller (iPSC); iPSC-afledte hepatocytllignende celler; voksne stam-/progenitorceller afledt af leveren; og voksne stam-/progenitorceller afledt af et andet væv end lever.
9. Fremgangsmåde ifølge krav 8, hvor mindst én komponent af konstruktionen er fremstillet ved bioprintning; eventuelt hvor konstruktionen i det væsentlige er fri for et hvilket som helst foruddannede skelet på anvendelsestidspunktet; fortrinsvis hvor konstruktionen er ikke-innervet.
- 20 10. Manipuleret, levende, tredimensional levervævskonstruktion ifølge krav 1, forudsat at den levende, tredimensionale levervævskonstruktion forbliver i det væsentlige levedygtig i en *in vitro*-kultur i mindst 7 dage og har mindst to af følgende leverkarakteristika:
- a. syntetiserer kolesterol;
 - b. lagrer lipid;
 - 25 c. lagrer glycogen;
 - d. udtrykker leverspecifikke proteiner, herunder to eller flere af: albumin, fibrinogen, transferrin, CYP450, alpha-1-antitrypsin, ornitintranscarbamylase, glycogensyntase, faktor VIII og faktor IX;
 - e. responderer på levermodulatorer, herunder én eller flere af: hormoner, inducere, giftstoffer, lægemidler, antistoffer, interfererende RNA'er, små RNA'er, låste nukleinsyrer, vira, bakterier, parasitter og
 - 30 inflammatoriske udløsere;
 - f. responderer på levergiftstoffer, herunder én eller flere af: paracetamol, alkohol, trovafloxacin, methotrexat, diclofenac, troglitazon, valproinsyre og amiodaron;
 - g. udtrykker transporterproteiner, herunder én eller flere af: BSEP, OATP1B1, NTCP, PGP, BCRP, OATP1B3, ABCB4, ATP8B1, multi-lægemiddelresistens-transporter (MDR);
 - 35 h. bibeholder mindst nogle hvilende stjerneceller efter fremstilling; og
 - i. indbefatter mikrovaskulære strukturer.

- 3 -

11. Manipuleret levervævskonstruktion ifølge krav 10, hvor konstruktionen i det væsentlige forbliver levedygtig i en *in vitro*-kultur i mindst 14 dage, fortrinsvis i mindst 28 dage; mere fortrinsvis i mindst 40 dage.
12. Manipuleret levervævskonstruktion ifølge krav 10 eller krav 11, der omfatter celler afledt af
5 inducerede pluripotente stamceller (iPSC).
13. Manipuleret levervævskonstruktion ifølge krav 10 eller krav 11, der omfatter myeloide celler afledt af monocytter eller makrofager.

DRAWINGS

Drawing

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11





Fig. 12

1.3 X amplification light microscopy



1.3 X amplification light microscopy





Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17






Fig. 18

Architecturally correct tissue with
vascular network




A

Zonal tissues




B

Lobulated tissues




C

Perfused / Arrayed
tissues



D

Solid + liquid tissue / Liquid interfaces



E

Barrier tissue



F

Fig. 18 (cont.)





Fig. 19



temporary filler bodies
represented by dark circles

Fig. 20




Fig. 21





Fig. 22



A

C

B

Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30




Fig. 31




Fig. 32a

Cytokine production in 3D liver constructs
post LPS (10 μ g/ml) treatment




Fig. 32b

Cytokine production in 3D liver constructs
post LPS (100 μ g/ml) treatment




Fig. 33





Fig. 34



N=3

LPS-EB Ultrapure, *E. coli* 0111:B4

Fig. 35




Fig. 36

