



(86) Date de dépôt PCT/PCT Filing Date: 2011/09/07
(87) Date publication PCT/PCT Publication Date: 2012/03/15
(45) Date de délivrance/Issue Date: 2018/05/29
(85) Entrée phase nationale/National Entry: 2013/02/28
(86) N° demande PCT/PCT Application No.: US 2011/050728
(87) N° publication PCT/PCT Publication No.: 2012/033858
(30) Priorité/Priority: 2010/09/07 (US61/380,596)




(51) Cl.Int./Int.Cl. *A61K 31/69* (2006.01),
A61P 31/04 (2006.01), *C07F 5/02* (2006.01)

(72) Inventeurs/Inventors:
HERNANDEZ, VINCENT S., US;
DING, CHARLES, US;
PLATTNER, JACOB J., US;
ALLEY, MICHAEL RICHARD KEVIN, US;
ROCK, FERNANDO, US;
ZHANG, SUOMING, US;
...

(73) Propriétaire/Owner:
ANACOR PHARMACEUTICALS, INC., US

(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : PETITES MOLECULES CONTENANT DU BORE
(54) Title: BORON-CONTAINING SMALL MOLECULES



(57) Abrégé/Abstract:

This invention relates to, among other items, benzoxaborole compounds and their use for treating bacterial infections. The compounds are of formula (I), (II) or (III) wherein R³ is substituted or unsubstituted nitroalkyl or substituted or unsubstituted

(72) Inventeurs(suite)/Inventors(continued): EASOM, ERIC, US; LI, XIANFENG, US; ZHOU, DING, CN

(57) Abrégé(suite)/Abstract(continued):

aminoalkyl; R^4 is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl; Y is O or S; and R^5 is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
15 March 2012 (15.03.2012)

PCT

(10) International Publication Number
WO 2012/033858 A3

(51) International Patent Classification:

A61K 31/69 (2006.01) *A61P 31/04* (2006.01)
C07F 5/02 (2006.01)

(21) International Application Number:

PCT/US2011/050728

(22) International Filing Date:

7 September 2011 (07.09.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/380,596 7 September 2010 (07.09.2010) US

(71) Applicant (for all designated States except US): **ANACOR PHARMACEUTICALS, INC.** [US/US]; 1020 East Meadow Circle, Palo Alto, CA 94303 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HERNANDEZ, Vincent, S.** [US/US]; 287 Gilchrist Lane, Watsonville, CA 95076 (US). **DING, Charles** [US/US]; 23 Treetop Lane, San Mateo, CA 94402 (US). **PLATTNER, Jacob, J.** [US/US]; 119 Via Floreado, Orinda, CA 94563 (US). **ALLEY, Michael Richard, Kevin** [GB/US]; 3751 Lillick Drive, Santa Clara, CA 95051 (US). **ROCK, Fernando** [US/US]; 1183 Lisa Lane, Los Altos, CA 94024 (US). **ZHANG, Suoming** [CN/US]; 3702 Redwood Circle, Palo Alto, CA 94306 (US). **EASOM, Eric** [US/US]; 1558 Mercy Street, Mountain View, CA 94041 (US). **LI, Xianfeng** [CA/US]; 10059 Craft Drive, Cupertino, CA 95014 (US). **ZHOU, Ding** [CN/CN]; 2nd Floor, Bldg. 1B, 528 Rui Qing Road, Shanghai, Pudong 201201 (CN).(74) Agent: **ESKER, Todd**; Morgan, Lewis & Bockius, LLP, One Market, Spear Street Tower, San Francisco, CA 94105 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))


Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))


(88) Date of publication of the international search report:

10 May 2012


(54) Title: BENZOXABOROLE DERIVATIVES FOR TREATING BACTERIAL INFECTIONS



(I)



(II)



(III)


(57) Abstract: This invention relates to, among other items, benzoxaborole compounds and their use for treating bacterial infections. The compounds are of formula (I), (II) or (III) wherein R³ is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R⁴ is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl; Y is O or S; and R⁵ is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

BORON-CONTAINING SMALL MOLECULES

BACKGROUND OF THE INVENTION

[0001] The global rise of bacteria and other microorganisms resistant to antibiotics and antimicrobials in general, poses a major threat. Deployment of massive quantities
5 of antimicrobial agents into the ecosphere during the past 60 years has introduced a powerful selective pressure for the emergence and spread of antimicrobial-resistant pathogens. Thus, there is a need to discover new broad spectrum antimicrobials, such as antibiotics, useful in combating microorganisms, especially those with multidrug-resistance.


10 [0002] Boron-containing molecules, such as 1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaborole (also sometimes known as 1-hydroxy-benzo[c][1,2]oxaborole or oxaboroles or cyclic boronic esters), useful as antimicrobials have been described previously, such as in U.S. Pat. Apps. 12/142,692 (U.S. Pat. Pub. 2009/0227541); 11/505,591 (U.S. Pat. Pub. 2006/0234981) and 11/357,687 (U.S. Pat. Pub.
15 2007/0155699). Generally speaking, a 1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaborole has the following structure and substituent numbering system:



Surprisingly, it has now been discovered that certain classes of 1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaboroles are effective antibacterials. This, and other uses of
20 these 1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaboroles are described herein.

SUMMARY OF THE INVENTION

[0004] In a first aspect, the invention provides a compound having a structure according to the formula:



- 5 wherein R^3 is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R^4 is selected from the group consisting of halogen, unsubstituted alkyl and unsubstituted phenyl; Y is O or S; R^5 is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl, or a salt, hydrate or solvate thereof.
- 10 [0005] In a second aspect, the invention provides a combination comprising: a) a compound of the invention, or a pharmaceutically acceptable salt thereof; and b) a therapeutically active agent.

- [0006] In a third aspect, the invention provides a pharmaceutical formulation comprising: a) a compound of the invention, or a pharmaceutically acceptable salt thereof; and b) a pharmaceutically acceptable excipient.
- 15

- [0007] In a fourth aspect, the invention provides a method of killing or inhibiting the growth of a bacteria, said method comprising: contacting said bacteria with an effective amount of a compound of the invention or a combination of the invention, or a pharmaceutically acceptable salt thereof, thereby killing or inhibiting the growth of the bacteria.
- 20

[0008] In a fifth aspect, the invention provides a method of treating a bacterial infection comprising: administering to an animal suffering from said infection an effective amount of a compound of the invention, or a pharmaceutically-acceptable salt thereof, thereby treating the bacterial infection.

- 25 [0009] In a sixth aspect, the invention provides a method of inhibiting the editing domain of a t-RNA synthetase, comprising: contacting the synthetase with an effective amount of a compound of the invention, or a pharmaceutically-acceptable salt thereof, thereby inhibiting the synthetase.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] FIG. 1 displays biological data for exemplary compounds of the invention.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions and Abbreviations

- 5 [0011] As used herein, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. For example, reference to “an active agent” includes a single active agent as well as two or more different active agents in combination. It is to be understood that present teaching is not limited to the specific dosage forms, carriers, or the like, disclosed herein and as such may vary.
- 10 [0012] The abbreviations used herein generally have their conventional meaning within the chemical and biological arts.
- [0013] The following abbreviations have been used: Ac is acetyl; AcOH is acetic acid; ACTBr is cetyltrimethylammonium bromide; AIBN is azobisisobutyronitrile or 2,2 azobisisobutyronitrile; aq. is aqueous; Ar is aryl; B₂pin₂ is bis(pinacolato)diboron; Bn is, in general, benzyl [see Cbz for one example of an exception]; (BnS)₂ is benzyl disulfide; BnSH is benzyl thiol or benzyl mercaptan; BnBr is benzyl bromide; Boc is tert-butoxy carbonyl; Boc₂O is di-*tert*-butyl dicarbonate; Bz is, in general, benzoyl; BzOOH is benzoyl peroxide; Cbz or Z is benzyloxycarbonyl or carboxybenzyl; Cs₂CO₃ is cesium carbonate; CSA is camphor sulfonic acid; CTAB is
- 15 cetyltrimethylammonium bromide; Cy is cyclohexyl; DABCO is 1,4-diazabicyclo[2.2.2]octane; DCM is dichloromethane or methylene chloride; DHP is dihydropyran; DIAD is diisopropyl azodicarboxylate; DIEA or DIPEA is *N,N*-diisopropylethylamine; DMAP is 4-(dimethylamino)pyridine; DME is 1,2-dimethoxyethane; DMF is *N,N*-dimethylformamide; DMSO is dimethylsulfoxide;
- 20 equiv or eq. is equivalent; EtOAc is ethyl acetate; EtOH is ethanol; Et₂O is diethyl ether; EDCI is *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride; ELS is evaporative light scattering; equiv or eq is equivalent; h is hours; HATU is O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate; HOBt is *N*-hydroxybenzotriazole; HCl is hydrochloric acid; HPLC is high pressure liquid
- 25 chromatography; ISCO Companion is automated flash chromatography equipment with fraction analysis by UV absorption available from Presearch; KOAc or AcOK is potassium acetate; K₂CO₃ is potassium carbonate; LiAlH₄ or LAH is lithium
- 30

aluminum hydride; LDA is lithium diisopropylamide; LHMDs is lithium bis(trimethylsilyl) amide; KHMDs is potassium bis(trimethylsilyl) amide; LiOH is lithium hydroxide; m-CPBA is 3-chloroperoxybenzoic acid; MeCN or ACN is methyl cyanide or cyanomethane or ethanenitrile or acetonitrile which are all names for the same compound; MeOH is methanol; MgSO₄ is magnesium sulfate; mins or min is minutes; Mp or MP is melting point; NaCNBH₃ is sodium cyanoborohydride; NaOH is sodium hydroxide; Na₂SO₄ is sodium sulfate; NBS is N-bromosuccinimide; NH₄Cl is ammonium chloride; NIS is N-iodosuccinimide; N₂ is nitrogen; NMM is *N*-methylmorpholine; n-BuLi is n-butyllithium; overnight is O/N; PdCl₂(pddf) is 1,1'-Bis(diphenylphosphino) ferrocene]dichloropalladium(II); Pd/C is the catalyst known as palladium on carbon; Pd₂(dba)₃ is an organometallic catalyst known as tris(dibenzylideneacetone) dipalladium(0); Ra Ni or Raney Ni is Raney nickel; Ph is phenyl; PMB is *p*-methoxybenzyl; PrOH is 1-propanol; iPrOH is 2-propanol; POCl₃ is phosphorus chloride oxide; PTSA is *para*-toluene sulfonic acid; Pyr. or Pyr or Py as used herein means pyridine; RT or rt or r.t. is room temperature; sat. is saturated; Si-amine or Si-NH₂ is amino-functionalized silica, available from SiliCycle®; Si-pyr is pyridyl-functionalized silica, available from SiliCycle®; TEA or Et₃N is triethylamine; TFA is trifluoroacetic acid; Tf₂O is trifluoromethanesulfonic anhydride; THF is tetrahydrofuran; TFAA is trifluoroacetic anhydride; THP is tetrahydropyranyl; TMSI is trimethylsilyl iodide; H₂O is water; diNO₂PhSO₂Cl is dinitrophenyl sulfonyl chloride; 3-F-4-NO₂-PhSO₂Cl is 3-fluoro-4-nitrophenylsulfonyl chloride; 2-MeO-4-NO₂-PhSO₂Cl is 2-methoxy-4-nitrophenylsulfonyl chloride; and (EtO)₂POCH₂COOEt is a triethylester of phosphonoacetic acid known as triethyl phosphonoacetate.

[0003] "Compound of the invention," as used herein refers to the compounds discussed herein, salts (e.g. pharmaceutically acceptable salts), prodrugs, solvates and hydrates of these compounds.

[0004] The term "poly" as used herein means at least 2. For example, a polyvalent metal ion is a metal ion having a valency of at least 2.

[0005] "Moiety" refers to a radical of a molecule that is attached to the remainder of the molecule.

[0017] The symbol , whether utilized as a bond or displayed perpendicular to a bond, indicates the point at which the displayed moiety is attached to the remainder of the molecule.

[0018] The term “alkyl,” by itself or as part of another substituent, means, unless
5 otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or
combination thereof, which may be fully saturated, mono- or polyunsaturated and can
include di- and multivalent radicals, having the number of carbon atoms designated
(*i.e.* C₁-C₁₀ means one to ten carbons). In some embodiments, the term “alkyl” means
a straight or branched chain, or combinations thereof, which may be fully saturated,
10 mono- or polyunsaturated and can include di- and multivalent radicals. Examples of
saturated hydrocarbon radicals include, but are not limited to, groups such as methyl,
ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl,
(cyclohexyl)methyl, cyclopropylmethyl, homologs and isomers of, for example, n-
pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one
15 having one or more double bonds or triple bonds. Examples of unsaturated alkyl
groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-
(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-
butynyl, and the higher homologs and isomers.

[0019] The term “alkylene” by itself or as part of another substituent means a
20 divalent radical derived from an alkane, as exemplified, but not limited, by
–CH₂CH₂CH₂CH₂–. Typically, an alkyl (or alkylene) group will have from 1 to 24
carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in
the invention. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene
group, generally having eight or fewer carbon atoms.

25 [0020] The term “alkenylene” by itself or as part of another substituent means a
divalent radical derived from an alkene.

[0021] The term “cycloalkylene” by itself or as part of another substituent means a
divalent radical derived from a cycloalkyl.

[0022] The term “heteroalkylene” by itself or as part of another substituent means a
30 divalent radical derived from a heteroalkane.

[0023] The term “heterocycloalkylene” by itself or as part of another substituent means a divalent radical derived from an heterocycloalkane.

[0024] The term “arylene” by itself or as part of another substituent means a divalent radical derived from an aryl.

5 [0025] The term “heteroarylene” by itself or as part of another substituent means a divalent radical derived from heteroaryl.

[0026] The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively.

10 [0027] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and at least one heteroatom. In some embodiments, the term “heteroalkyl,” by itself or in combination with another term, means a stable straight or
15 branched chain, or combinations thereof, consisting of the stated number of carbon atoms and at least one heteroatom. In an exemplary embodiment, the heteroatoms can be selected from the group consisting of B, O, N and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) B, O, N and S may be placed at any interior
20 position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Examples include, but are not limited to, -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. Up to two heteroatoms may be consecutive, such as, for
25 example, -CH₂-NH-OCH₃. Similarly, the term “heteroalkylene” by itself or as part of another substituent means a divalent radical derived from heteroalkyl, as exemplified, but not limited by, -CH₂-CH₂-S-CH₂-CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (*e.g.*, alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the
30 like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied by the direction in which the formula of the linking group

is written. For example, the formula $-C(O)_2R'$ represents both $-C(O)_2R'$ and $R'C(O)_2-$.

[0028] The terms “cycloalkyl” and “heterocycloalkyl”, by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl”, respectively. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piprazinyl, 2-piprazinyl, and the like.

[0029] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl,” are meant to include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” is meant to include, but not be limited to, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0030] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, substituent that can be a single ring or multiple rings (preferably from 1 or 2 or 3 rings), which are fused together or linked covalently. The term “heteroaryl” refers to aryl groups (or rings) that contain from one to four heteroatoms. In an exemplary embodiment, the heteroatom is selected from B, N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 3-quinolyl, and 6-quinolyl.

Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0031] For brevity, the term “aryl” when used in combination with other terms (*e.g.*, aryloxy, arylthioxy, arylalkyl) includes those radicals in which an aryl group is attached through the next moiety to the rest of the molecule. Thus, the term “arylalkyl” is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, 1-(3-nitrophenyl)ethyl and the like). A substituent such as benzyl or 1-(3-nitrophenyl)ethyl can also be represented by ‘substituted alkyl’ wherein the ethyl radical is substituted with a 3-nitrophenyl moiety. The term “aryloxy” is meant to include those radicals in which an aryl group is attached to an oxygen atom. The term “aryloxyalkyl” is meant to include those radicals in which an aryl group is attached to an oxygen atom which is then attached to an alkyl group (*e.g.*, phenoxymethyl, 3-(1-naphthyloxy)propyl, and the like).

[0032] For brevity, the term “heteroaryl” when used in combination with other terms (*e.g.*, heteroaryloxy, heteroarylthioxy, heteroarylalkyl) includes those radicals in which a heteroaryl group is attached through the next moiety to the rest of the molecule. Thus, the term “heteroarylalkyl” is meant to include those radicals in which a heteroaryl group is attached to an alkyl group (*e.g.*, pyridylmethyl and the like). The term “heteroaryloxy” is meant to include those radicals in which a heteroaryl group is attached to an oxygen atom. The term “heteroaryloxyalkyl” is meant to include those radicals in which an aryl group is attached to an oxygen atom which is then attached to an alkyl group. (*e.g.*, 2-pyridyloxymethyl and the like).

[0033] Each of the above terms (*e.g.*, “alkyl,” “heteroalkyl,” “aryl” and “heteroaryl”) are meant to include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

[0034] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) are generically referred to as “alkyl group substituents,” and they can be one or more of a variety of groups selected from, but not limited to: -R', -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR''''-C(NR'R''R''')=NR''''',

-NR^{''''}-C(NR^{'R''})=NR^{'''}, -S(O)R', -S(O)₂R', -S(O)₂NR^{'R''}, -NR^{''}SO₂R', -CN, -NO₂, -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such radical.

R', R'', R''', R^{''''} and R^{'''''} each preferably independently refer to hydrogen,

- 5 substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl, *e.g.*, aryl substituted with 1 or 2 or 3 halogens, substituted or unsubstituted alkyl, alkoxy or thioalkoxy groups, or arylalkyl groups. When a compound of the invention includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', R^{''''} and R^{'''''} groups when more than one of these groups is
- 10 present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include, but not be limited to, 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term "alkyl" is meant to include groups including carbon atoms bound to groups
- 15 other than hydrogen groups, such as haloalkyl (*e.g.*, -CF₃ and -CH₂CF₃) and acyl (*e.g.*, -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

[0035] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups are generically referred to as "aryl group substituents."

The substituents are selected from, for example: -R', -OR', =O, =NR', =N-OR', -

- 20 NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R'', -NR''C(O)₂R', -NR^{''''}-C(NR'R''R''')=NR^{'''}, -NR^{''''}-C(NR'R''R''')=NR^{'''}, -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR^{''}SO₂R', -CN, -NO₂, -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic
- 25 ring system; and where R', R'', R''', R^{''''} and R^{'''''} are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl and substituted or unsubstituted heteroaryl. When a compound of the invention includes more than one R group, for example, each of the R groups is independently selected as are each R',
- 30 R'', R''', R^{''''} and R^{'''''} groups when more than one of these groups is present.

[0036] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CRR')_q-U-, wherein T and U are independently -NR-, -O-, -CRR'- or a single bond, and q is an

integer from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula – A-(CH₂)_r-B-, wherein A and B are independently –CRR'-, -O-, -NR-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer from 1 or 2 or 3 or 4. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula – (CRR')_s-X-(CR''R''')_d-, where s and d are independently integers from 0 or 1 or 2 or 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituents R, R', R'' and R''' are preferably independently selected from hydrogen or substituted or unsubstituted (C₁ or C₂ or C₃ or C₄ or C₅ or C₆)alkyl.

[0037] “Ring” as used herein, means a substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. A ring includes fused ring moieties. The number of atoms in a ring is typically defined by the number of members in the ring. For example, a “5- to 7-membered ring” means there are 5 or 6 or 7 atoms in the encircling arrangement. Unless otherwise specified, the ring optionally includes a heteroatom. Thus, the term “5- to 7-membered ring” includes, for example phenyl, pyridinyl and piperidinyl. The term “5- to 7-membered heterocycloalkyl ring”, on the other hand, would include pyridinyl and piperidinyl, but not phenyl. The term “ring” further includes a ring system comprising more than one “ring”, wherein each “ring” is independently defined as above.

[0038] As used herein, the term “heteroatom” includes atoms other than carbon (C) and hydrogen (H). Examples include oxygen (O), nitrogen (N) sulfur (S), silicon (Si), and boron (B).

[0039] The term “leaving group” means a functional group or atom which can be displaced by another functional group or atom in a substitution reaction, such as a nucleophilic substitution reaction. By way of example, representative leaving groups include triflate, chloro, bromo and iodo groups; sulfonic ester groups, such as mesylate, tosylate, brosylate, nosylate and the like; and acyloxy groups, such as acetoxy, trifluoroacetoxy and the like.

[0040] The symbol "R" is a general abbreviation that represents a substituent group that is selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl and substituted or unsubstituted heterocycloalkyl groups.

[0041] By "effective" amount of a drug, formulation, or permeant is meant a sufficient amount of an active agent to provide the desired local or systemic effect. A "Topically effective," "pharmaceutically effective," or "therapeutically effective" amount refers to the amount of drug needed to effect the desired result.

10 [0042] The term "pharmaceutically acceptable salt" is meant to include a salt of a compound of the invention which is prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with
15 a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino (such as choline or diethylamine or amino acids such as d-arginine, l-arginine, d-lysine or l-lysine), or magnesium salt, or a similar salt. When compounds of the invention contain relatively basic
20 functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric,
25 dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate
30 and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge et al., "Pharmaceutical Salts", Journal of Pharmaceutical Science 66: 1-19 (1977)). Certain specific compounds of the invention contain both

basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0043] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compounds in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents.

[0044] In addition to salt forms, the invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein readily undergo chemical changes under physiological conditions to provide the compounds of the invention. Additionally, prodrugs can be converted to the compounds of the invention by chemical or biochemical methods in an *ex vivo* environment.

[0045] Certain compounds of the invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of the invention. Certain compounds of the invention may exist in multiple crystalline or amorphous forms.

[0046] Certain compounds of the invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are encompassed within the scope of the invention. The graphic representations of racemic, ambiscalemic and scalemic or enantiomerically pure compounds used herein are taken from Maehr, *J. Chem. Ed.* 1985, 62: 114-120. Solid and broken wedges are used to denote the absolute configuration of a stereocenter unless otherwise noted. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms are included.

[0047] Compounds of the invention can exist in particular geometric or stereoisomeric forms. The invention contemplates all such compounds, including *cis*- and *trans*-isomers, (-)- and (+)-enantiomers, (R)- and (S)-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, such as enantiomerically or diastereomerically enriched mixtures, as falling within the scope of the invention. Additional asymmetric carbon atoms can be present in a

substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

[0048] Optically active (*R*)- and (*S*)-isomers and *d* and *l* isomers can be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques.

5 If, for instance, a particular enantiomer of a compound of the invention is desired, it can be prepared by asymmetric synthesis, or by derivatization with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group
10 cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as an amino group, or an acidic functional group, such as a carboxyl group, diastereomeric salts can be formed with an
15 appropriate optically active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means known in the art, and subsequent recovery of the pure enantiomers. In addition, separation of enantiomers and diastereomers is frequently accomplished using chromatography
employing chiral, stationary phases, optionally in combination with chemical
derivatization (e.g., formation of carbamates from amines).

[0049] The compounds of the invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for
20 example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the invention, whether radioactive or not, are intended to be encompassed within the scope of the invention.

[0050] The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable vehicle” refers to any formulation or carrier medium that provides the
25 appropriate delivery of an effective amount of an active agent as defined herein, does not interfere with the effectiveness of the biological activity of the active agent, and that is sufficiently non-toxic to the host or patient. Representative carriers include water, oils, both vegetable and mineral, cream bases, lotion bases, ointment bases and the like. These bases include suspending agents, thickeners, penetration enhancers,
30 and the like. Their formulation is well known to those in the art of cosmetics and topical pharmaceuticals. Additional information concerning carriers can be found in

Remington: The Science and Practice of Pharmacy, 21st Ed., Lippincott, Williams & Wilkins (2005).

[0006] The term “excipients” is conventionally known to mean carriers, diluents and/or vehicles used in formulating drug compositions effective for the desired use.

5 [0007] The term “microbial infection” or “infection by a microorganism” refers to any infection of a host tissue by an infectious agent including, but not limited to, bacteria or protozoa (see, e.g., Harrison's Principles of Internal Medicine, pp. 93-98 (Wilson et al., eds., 12th ed. 1991); Williams et al., J. of Medicinal Chem. 42:1481-1485 (1999)).

10 [0008] “Biological medium,” as used herein refers to both in vitro and in vivo biological milieus. Exemplary in vitro “biological media” include, but are not limited to, cell culture, tissue culture, homogenates, plasma and blood. In vivo applications are generally performed in mammals, preferably humans.

[0009] “Inhibiting” and “blocking,” are used interchangeably herein to refer to the
15 partial or full blockade of enzyme. In an exemplary embodiment, the enzyme is an editing domain of a tRNA synthetase.

[0010] Boron is able to form additional covalent or dative bonds with oxygen, sulfur or nitrogen under some circumstances in this invention.

[0011] Embodiments of the invention also encompass compounds that are poly-
20 or multi-valent species, including, for example, species such as dimers, trimers, tetramers and higher homologs of the compounds of use in the invention or reactive analogues thereof.

[0012] “Salt counterion”, as used herein, refers to positively charged ions that associate with a compound of the invention when the boron is fully negatively or
25 partially negatively charged. Examples of salt counterions include H^+ , H_3O^+ , ammonium, potassium, calcium, magnesium (such as choline or diethylamine or amino acids such as d-arginine, l-arginine, d-lysine or l-lysine) and sodium.

[0013] The compounds comprising a boron bonded to a carbon and three heteroatoms (such as three oxygens described in this section) can optionally contain a
30 fully negatively charged boron or partially negatively charged boron. Due to the

negative charge, a positively charged counterion may associate with this compound, thus forming a salt. Examples of salt counterions include H^+ , H_3O^+ , ammonium, potassium, calcium, magnesium (such as choline or diethylamine or amino acids such as d-arginine, l-arginine, d-lysine or l-lysine) and sodium. The salts of the compounds are implicitly contained in descriptions of these compounds.

II. Introduction


[0059] The invention provides novel boron compounds and methods for the preparation of these molecules. The invention further provides methods of treating bacterial infections, killing or inhibiting the growth of bacteria in part or wholly through the use of the compounds described herein. In another aspect, the invention is a combination of a compound of the invention and an antibiotic. In another aspect, the invention is a pharmaceutical formulation comprising a pharmaceutically acceptable excipient and a compound of the invention. In another aspect, the invention is a pharmaceutical formulation comprising a compound of the invention, an antibiotic, and a pharmaceutically acceptable excipient.

III. Composition of Matter

III. a.) Compounds

[0060] In one aspect the invention provides a compound of the invention. In an exemplary embodiment, the invention provides a compound described herein, or a salt thereof. In an exemplary embodiment, the salt of a compound described herein is a pharmaceutically acceptable salt. In an exemplary embodiment, the invention provides a compound described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the invention provides a compound described in a formula provided herein. In an exemplary embodiment, the invention provides a compound described herein.


[0061] In an aspect, the invention provides a compound having a structure which is:



wherein R^3 is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R^4 is selected from the group consisting of halogen, unsubstituted alkyl


unsubstituted alkoxy and unsubstituted phenyl; Y is O or S; R⁵ is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

[0062] In an aspect, the invention provides a compound having a structure which is:




wherein R³ is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R⁴ is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl; Y is O or S; R⁵ is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

[0063] In an aspect, the invention provides a compound having a structure which is:




wherein R³ is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R⁴ is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl; Y is O or S; R⁵ is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

[0064] In an exemplary embodiment, there is provided a compound having a structure according to the following formula:




wherein C* is a carbon atom stereocenter which has a configuration which is (R) or (S). In an exemplary embodiment, the C* stereocenter is in the (S) configuration.

[0065] In an exemplary embodiment, there is provided a compound having a structure according to the following formula:



wherein C* is a carbon atom stereocenter which has a configuration which is (R) or (S). In an exemplary embodiment, the C* stereocenter is in the (S) configuration.

[0066] In an exemplary embodiment, there is provided a compound having a structure according to the following formula:



wherein C* is a carbon atom stereocenter which has a configuration which is (R) or (S). In an exemplary embodiment, the C* stereocenter is in the (S) configuration.


[0067] In an exemplary embodiment, Y, R⁵ and R⁴ are as described herein, R³ is—(CR²⁰R²¹)_nNR²²R²³ in which n is an integer selected from 1 to 10; each R²⁰ and each R²¹ is independently selected from the group consisting of H, R²⁶, OR²⁶, NR²⁶R²⁷, SR²⁶, -S(O)R²⁶, -S(O)₂R²⁶, -S(O)₂NR²⁶R²⁷, -C(O)R²⁷, -C(O)OR²⁷, -C(O)NR²⁶R²⁷; R²² and R²³ are independently selected from the group consisting of H, -S(O)R²⁸, -S(O)₂R²⁸, -S(O)₂NR²⁸R²⁹, -C(O)R²⁸, -C(O)OR²⁸, -C(O)NR²⁸R²⁹, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl, wherein each R²⁶, each R²⁷, each R²⁸ and each R²⁹ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

[0068] In an exemplary embodiment, Y, R⁵ and R⁴ are as described herein, and R³ is -CH₂NH₂ or -CH₂NO₂. In an exemplary embodiment, Y, R⁵ and R⁴ are as described herein, and R³ is -CH₂NH₂. In an exemplary embodiment, Y, R⁵ and R⁴ are as described herein, R³ is -CH₂NH₂, and C* has a configuration which is (S).

[0069] In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, and *sec*-butyl. In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is selected from the group consisting of fluorine, chlorine, bromine, and iodine. In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is chlorine or bromine. In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is chlorine.

[0070] In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, and *sec*-butoxy. In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is methoxy or ethoxy. In an exemplary embodiment, Y, R⁵ and R³ are as described herein, and R⁴ is methoxy.

[0071] In an exemplary embodiment, Y, R⁴ and R³ are as described herein, and R⁵ is:



wherein a is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; each R¹⁰ and each R¹¹ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, OH and NH₂; R¹² is selected from the group consisting of H, R⁷, halogen, cyano, amidino, OR⁷, NR⁷R⁸, SR⁷, -N(R⁷)S(O)₂R⁸, -C(O)R⁷, -C(O)OR⁷, -C(O)NR⁷R⁸ wherein each R⁷ and each R⁸ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl. In an exemplary embodiment, Y, R⁴, R³, R¹⁰, R¹¹, and R¹² are as described herein, and a is 1, 2, 3, 4, or 5. In an exemplary embodiment, Y, R⁴, R³, R¹⁰, R¹¹, and R¹² are as described herein, and a is 2, 3, or 4. In an exemplary embodiment, Y, R⁴, R³, R¹⁰, R¹¹, and R¹² are as described herein, and a is 3. In an exemplary embodiment, Y, R⁴, R³, a, and R¹² are as described herein, and each R¹⁰ and each R¹¹ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, OH, and NH₂. In an exemplary embodiment, Y, R⁴, R³, a, and R¹² are as described herein, and each R¹⁰ and each R¹¹ is H. In an exemplary embodiment, Y, R⁴, R³, R¹⁰, R¹¹, and a are as

described herein, and R^{12} is selected from the group consisting of H, OH, NH_2 , methyl, ethyl, $-NHS(O)_2CH_3$, cyano, $-NHC(O)CH_3$, $-NHC(O)NHCH_2CH_3$, $-C(O)NH_2$, $-C(O)OH$, 4-(methoxy)phenyl, benzyl, benzoxy, $-NHC(O)OCH_2Ph$, $-C(O)NHCH_2CH_2OH$ and $-C(NH_2)(NH)$.

- 5 [0072] In an exemplary embodiment, R^4 , R^3 , and R^5 are as described herein, and Y is O. In an exemplary embodiment, R^4 , R^3 , and Y are as described herein, and R^5 is unsubstituted alkyl. In an exemplary embodiment, R^4 , R^3 , and Y are as described herein, and R^5 is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, and sec-butyl.
- 10 [0073] In an exemplary embodiment, R^4 is halogen, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, and sec-butyl. In an exemplary embodiment, R^4 is halogen, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is halogen, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.
- 15 [0074] In an exemplary embodiment, R^4 is chlorine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is chlorine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.
- 20 [0075] In an exemplary embodiment, R^4 is fluorine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is fluorine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.
- 25 [0076] In an exemplary embodiment, R^4 is bromine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is bromine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.
- 30 [0077] In an exemplary embodiment, R^4 is halogen, R^3 is $-CH_2NH_2$; Y is O; and R^5 is methyl. In an exemplary embodiment, R^4 is fluorine, R^3 is $-CH_2NH_2$; Y is O; and R^5 is methyl. In an exemplary embodiment, R^4 is chlorine, R^3 is $-CH_2NH_2$; Y is

O; and R⁵ is methyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is methyl.

[0078] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is ethyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—CH₂NH₂; Y is O; and
5 R⁵ is ethyl. In an exemplary embodiment, R⁴ is chlorine, R³ is—CH₂NH₂; Y is O; and R⁵ is ethyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is ethyl.

[0079] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is propyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—CH₂NH₂; Y is O;
10 and R⁵ is propyl. In an exemplary embodiment, R⁴ is chlorine, R³ is—CH₂NH₂; Y is O; and R⁵ is propyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is propyl.

[0080] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is isopropyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—CH₂NH₂; Y is O;
15 and R⁵ is isopropyl. In an exemplary embodiment, R⁴ is chlorine, R³ is—CH₂NH₂; Y is O; and R⁵ is isopropyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is isopropyl.

[0081] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₄ alkyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—
20 CH₂NH₂; Y is O; and R⁵ is unsubstituted C₄ alkyl. In an exemplary embodiment, R⁴ is chlorine, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₄ alkyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₄ alkyl.

[0082] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₅ alkyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—
25 CH₂NH₂; Y is O; and R⁵ is unsubstituted C₅ alkyl. In an exemplary embodiment, R⁴ is chlorine, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₅ alkyl. In an exemplary embodiment, R⁴ is bromine, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₅ alkyl.

[0083] In an exemplary embodiment, R⁴ is halogen, R³ is—CH₂NH₂; Y is O; and R⁵ is unsubstituted C₆ alkyl. In an exemplary embodiment, R⁴ is fluorine, R³ is—
30 CH₂NH₂; Y is O; and R⁵ is unsubstituted C₆ alkyl. In an exemplary embodiment, R⁴

is chlorine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is unsubstituted C_6 alkyl. In an exemplary embodiment, R^4 is bromine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is unsubstituted C_6 alkyl.


[0084] In an exemplary embodiment, R^4 is chlorine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is chlorine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.

[0085] In an exemplary embodiment, R^4 is fluorine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is fluorine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.


[0086] In an exemplary embodiment, R^4 is bromine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of methyl, ethyl, propyl, and isopropyl. In an exemplary embodiment, R^4 is bromine, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is selected from the group consisting of butyl, isobutyl, t-butyl, and sec-butyl.

[0087] In an exemplary embodiment, R^4 is as described herein, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is substituted or unsubstituted alkyl. In an exemplary embodiment, Y and R^5 are as described herein, R^3 is $-\text{CH}_2\text{NH}_2$; and R^4 is halogen. In an exemplary embodiment, Y is as described herein, R^4 is halogen; Y is O; and R^5 is unsubstituted alkyl. In an exemplary embodiment, R^3 is $-\text{CH}_2\text{NH}_2$; R^4 is chlorine; Y is O; and R^5 is substituted or unsubstituted alkyl. In an exemplary embodiment, R^4 is as described herein, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is ethyl.

[0088] In an exemplary embodiment, the compound has a structure which is




[0089] In an exemplary embodiment, the compound has a structure which is




, wherein R^4 , Y and R^5 are as described herein.

[0090] In an exemplary embodiment, the compound has a structure which is




, wherein R^4 , Y and R^5 are as described herein. In an exemplary embodiment, Y is O, and R^4 and R^5 are as described herein. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 are as described herein. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is unsubstituted alkyl. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is methyl or ethyl or propyl or isopropyl. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is butyl or isobutyl or neobutyl or t-butyl.

[0091] In an exemplary embodiment, the compound has a structure which is




, wherein R^4 , Y and R^5 are as described herein.

[0092] In an exemplary embodiment, the compound has a structure which is




, wherein R^4 , Y and R^5 are as described herein. In an exemplary embodiment, Y is O, and R^4 and R^5 are as described herein. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 are as described herein. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is unsubstituted alkyl. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is methyl or ethyl or propyl or isopropyl. In an exemplary embodiment, Y is O, R^4 is halogen, and R^5 is butyl or isobutyl or neobutyl or t-butyl.

[0093] In an exemplary embodiment, the compound has a structure which is



, wherein R^4 , Y and R^5 are as described herein.

[0094] In an exemplary embodiment, the compound has a structure which is



, wherein R^4 , Y and R^5 are as described herein. In an exemplary embodiment, Y is O , and R^4 and R^5 are as described herein. In an exemplary embodiment, Y is O , R^4 is halogen, and R^5 are as described herein. In an exemplary embodiment, Y is O , R^4 is halogen, and R^5 is unsubstituted alkyl. In an exemplary embodiment, Y is O , R^4 is halogen, and R^5 is methyl or ethyl or propyl or isopropyl. In an exemplary embodiment, Y is O , R^4 is halogen, and R^5 is butyl or isobutyl or neobutyl or t-butyl.

[0095] In an exemplary embodiment, said alkyl is linear alkyl or branched alkyl,

10 In an exemplary embodiment, said heteroalkyl is linear heteroalkyl or branched heteroalkyl.

[0096] In an exemplary embodiment, the invention provides poly- or multi-valent species of the compounds of the invention, including a dimer or a trimer. Another exemplary embodiment of the invention provides an anhydride of the compounds of the invention. In another exemplary embodiment, the invention provides poly- or multi-valent species of the compounds of the invention. In an exemplary embodiment, the invention provides a dimer of the compounds described herein. In an exemplary embodiment, the invention provides a dimer of the compounds described herein.

20 [0097] In an exemplary embodiment, the invention provides an anhydride of the compounds described herein. In an exemplary embodiment, the invention provides an anhydride of the compounds described herein.

[0098] In an exemplary embodiment, the invention provides a trimer of the compounds described herein. In an exemplary embodiment, the invention provides a trimer of the compounds described herein.

[0099] The compounds of the invention can form a hydrate with water, solvates with alcohols such as methanol, ethanol, propanol, and the like; adducts with amino compounds, such as ammonia, methylamine, ethylamine, and the like; adducts with

acids, such as formic acid, acetic acid and the like; complexes with ethanolamine, quinoline, amino acids, and the like.

[0100] In an exemplary embodiment, the invention provides a compound described herein, or a salt, hydrate or solvate thereof, or a combination thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt, hydrate or solvate thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt thereof. In an exemplary embodiment, the salt is a pharmaceutically acceptable salt. In an exemplary embodiment, the invention provides a compound described herein, or a hydrate thereof. In an exemplary embodiment, the invention provides a compound described herein, or a solvate thereof. In an exemplary embodiment, the invention provides a compound described herein, or a prodrug thereof. In an exemplary embodiment, the invention provides a salt of a compound described herein. In an exemplary embodiment, the invention provides a pharmaceutically acceptable salt of a compound described herein. In an exemplary embodiment, the invention provides a hydrate of a compound described herein. In an exemplary embodiment, the invention provides a solvate of a compound described herein. In an exemplary embodiment, the invention provides a prodrug of a compound described herein. In an exemplary embodiment, the invention provides a compound as described in **FIG. 1**, or a salt thereof. In an exemplary embodiment, the invention provides a compound as described in **FIG. 1**, or a pharmaceutically acceptable salt thereof.

[0101] In an exemplary embodiment, alkyl is linear alkyl. In another exemplary embodiment, alkyl is branched alkyl.

[0102] In an exemplary embodiment, heteroalkyl is linear heteroalkyl. In another exemplary embodiment, heteroalkyl is branched heteroalkyl.

III. b) Combinations comprising additional therapeutic agents

[0103] The compounds of the invention may also be used in combination with additional therapeutic agents. The invention thus provides, in a further aspect, a combination comprising a compound of the invention together with at least one additional therapeutic agent, or a salt, prodrug, hydrate or solvate thereof. In an exemplary embodiment, the compound of the invention is a compound described herein, or a salt thereof. In an exemplary embodiment, the additional therapeutic

agent is a compound of the invention. In an exemplary embodiment, the additional therapeutic agent includes a boron atom. In an exemplary embodiment, the additional therapeutic agent does not contain a boron atom. In an exemplary embodiment, the additional therapeutic agent is a compound described in sections III a) or b).

5 [0104] When a compound of the invention is used in combination with a second therapeutic agent active against the same disease state, the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily appreciated by those skilled in the art. It will be appreciated that the amount
10 of a compound of the invention required for use in treatment will vary with the nature of the condition being treated and the age and the condition of the patient and will be ultimately at the discretion of the attendant physician or veterinarian.

[0105] In an exemplary embodiment, the additional therapeutic agent is an antibacterial agent. In an exemplary embodiment, the additional therapeutic agent is an antituberculosis agent. In an exemplary embodiment, the additional therapeutic
15 agent is rifampicin. In an exemplary embodiment, the additional therapeutic agent is isoniazid. In an exemplary embodiment, the additional therapeutic agent is pyrazinamide. In an exemplary embodiment, the additional therapeutic agent is ethambutol. In an exemplary embodiment, the additional therapeutic agent is isoniazid. In an exemplary embodiment, the additional therapeutic agent is
20 streptomycin. In an exemplary embodiment, the additional therapeutic agent is an aminoglycoside. In an exemplary embodiment, the additional therapeutic agent is amikacin or kanamycin. In an exemplary embodiment, the additional therapeutic agent is a polypeptide. In an exemplary embodiment, the additional therapeutic agent is selected from the group consisting of capreomycin, viomycin, and enviomycin. In
25 an exemplary embodiment, the additional therapeutic agent is a fluoroquinolone. In an exemplary embodiment, the additional therapeutic agent is selected from the group consisting of ciprofloxacin, levofloxacin, and moxifloxacin. In an exemplary embodiment, the additional therapeutic agent is a thioamide. In an exemplary embodiment, the additional therapeutic agent is ethionamide or prothionamide. In an
30 exemplary embodiment, the additional therapeutic agent is cycloserine. In an exemplary embodiment, the additional therapeutic agent is p-aminosalicylic acid. In an exemplary embodiment, the additional therapeutic agent is selected from the group consisting of rifabutin, linezolid, thioacetazone, thioridazine, arginine, vitamin D, and

R207910. In an exemplary embodiment, the additional therapeutic agent is a macrolide.

[0106] The individual components of such combinations may be administered either simultaneously or sequentially in a unit dosage form. The unit dosage form
5 may be a single or multiple unit dosage forms. In an exemplary embodiment, the invention provides a combination in a single unit dosage form. An example of a single unit dosage form is a capsule wherein both the compound of the invention and the additional therapeutic agent are contained within the same capsule. In an exemplary embodiment, the invention provides a combination in a two unit dosage
10 form. An example of a two unit dosage form is a first capsule which contains the compound of the invention and a second capsule which contains the additional therapeutic agent. Thus the term 'single unit' or 'two unit' or 'multiple unit' refers to the object which the animal (for example, a human) ingests, not to the interior components of the object. Appropriate doses of known therapeutic agents will be
15 readily appreciated by those skilled in the art.

[0107] The combinations referred to herein may conveniently be presented for use in the form of a pharmaceutical formulation. Thus, an exemplary embodiment of the invention is a pharmaceutical formulation comprising a) a compound of the invention; b) an additional therapeutic agent and c) a pharmaceutically acceptable excipient. In
20 an exemplary embodiment, the pharmaceutical formulation is a unit dosage form. In an exemplary embodiment, the pharmaceutical formulation is a single unit dosage form. In an exemplary embodiment, the pharmaceutical formulation is a single unit dosage form which includes a compound of the invention; an antibiotic and a pharmaceutically acceptable excipient. In an exemplary embodiment, the
25 pharmaceutical formulation is a single unit dosage form which includes a compound of the invention; an antibiotic and at least one pharmaceutically acceptable excipient. In an exemplary embodiment, the pharmaceutical formulation is a two unit dosage form. In an exemplary embodiment, the pharmaceutical formulation is a two unit dosage form comprising a first unit dosage form and a second unit dosage form,
30 wherein the first unit dosage form includes a) a compound of the invention and b) a first pharmaceutically acceptable excipient; and the second unit dosage form includes c) an additional therapeutic agent and d) a second pharmaceutically acceptable excipient. In an exemplary embodiment, the pharmaceutical formulation is a two unit


dosage form comprising a first unit dosage form and a second unit dosage form, wherein the first unit dosage form includes a) a compound of the invention and b) a first pharmaceutically acceptable excipient; and the second unit dosage form includes c) an antibiotic and d) a second pharmaceutically acceptable excipient.

5 *III. c) Preparation of Boron-Containing Compounds*

[0014] Compounds of use in the invention can be prepared using commercially available starting materials, known intermediates, or by using the synthetic methods published in references described herein, such as U.S. Pat. Apps. 12/142,692 (U.S. Pat. Pub. 2009/0227541) and U.S. Pat. Pubs. US20060234981, US20070155699 and
10 US20070293457.

[0015] The following general procedures were used as indicated in generating the examples and can be applied, using the knowledge of one of skill in the art, to other appropriate compounds to obtain additional analogues.

General reaction scheme 1



15 [0016] Scheme 1 describes a synthesis for compounds of (I), wherein R^4 is fluorine or chlorine and Y and R^5 are as described herein. The fluoro or chloro compound of formula A, which may be prepared or maybe available commercially from Sigma-Aldrich, is reacted with a strong base (such as $n\text{-BuLi}$, sec-BuLi , or $t\text{-BuLi}$, 2 equiv)


followed by quenching with a formylating agent (such as DMF, dimethylformamide, formanilide, N-formylmorpholine, large excess) to give the compound of formula B. Treatment of compound B with a demethylating agent (typically BBr₃, 2 equiv) in a suitable solvent (dichloromethane, THF) gives the phenol of formula C. Compound C

5 can react with a corresponding bromide or mesylate (1-1.5 equiv) in the presence of a base (such as KOtBu, K₂CO₃, or Cs₂CO₃, 1.5-2 equiv) in an aprotic solvent such as DMF or DMSO to afford the compound of formula D. Compound D may be converted to triflate E by the reaction with 1.2 equiv of trifluoromethanesulfonic anhydride and pyridine in dichloromethane. The conversion of triflate E to boronate F

10 can be achieved by the reaction with bis(pinacolato)diborane (2 equiv), KOAc (3 equiv) and catalytic amount of PdCl₂(dppf) (4-8 mol%). The reaction of compound F with nitromethane (3 equiv) in the presence of sodium hydroxide (3 equiv) in water or THF gives the nitro compound of formula G. The compound G can be converted to the final product of formula H by the Raney-Ni reduction (Raney Ni, 2 equiv w/w, 2.0

15 M NH₃ in EtOH, absolute EtOH).

General reaction scheme 2




[0111] Scheme 2 describes a synthesis for compounds of (I), wherein R⁴ is chlorine, Y and R⁵ are as described herein. The phenol or thiophenol of formula I, which may be prepared or maybe available commercially from Sigma-Aldrich, react

20 with a solution of bromine and catalytic amount of iron powder in glacial acetic acid to give the bromo substituted compound of formula J. The alkylation of J can be achieved by reacting with a bromide in the presence of a base such as potassium

carbonate in solvents like DMF or acetonitrile. The protection of aldehyde K may be achieved by refluxing with ethylene glycol in toluene, in the presence of catalytic amount of *p*-toluenesulfonic acid. The reaction of the compound L with BuLi and triisopropyl borate, followed by treating with hydrochloric acid yields boronic acid M.

- 5 The reaction of compound M with nitromethane in the presence of sodium hydroxide gives the nitro compound of formula N. The treatment of N with 1 equivalent of sulfonyl chloride affords the chloro substituted compound O. The Raney-Ni reduction of compound O in MeOH gives the final product of formula P.


General reaction scheme 3



10 [0112] Scheme 3 describes a synthesis for compounds of (I), wherein R^4 is bromine, Y and R^5 are as described herein. The compound of formula N, which may be prepared according to Scheme 2, can be reduced to the amine of formula S, by hydrogenation in the presence of palladium hydroxide or Raney-Ni reduction as


15 described above. The amine of formula S reacts with an N-Protecting reagent such as Boc anhydride in the presence of base like triethylamine in dichloromethane to give Boc-protected compound of formula T. The treatment of T with N-bromosuccinimide and catalytic amount of AIBN in acetonitrile gives the bromo substituted compound of formula U. Deprotection of compound U in the presence of acid such as HCl in

20 dioxane will afford the final compound of formula U.

General reaction scheme 4


[0113] Scheme 4 describes a synthesis for compounds of (I), wherein R⁴ is an alkyl or aryl group, Y and R⁵ are as described herein. The compound of formula N, which may be prepared according to Scheme 2, can be brominated with N-bromosuccinimide and catalytic amount of AIBN in a solvent such as acetonitrile to give the bromide of formula W. Stille coupling of W with an organotin compound such as tetramethylstannane or tributyl-phenyl-stannane in the presence of catalytic Pd(Ph₃P)₄ in DMF affords the compound of formula X. Compound X can be reduced to the final compound of formula Y by hydrogenation in the presence of palladium on carbon or Raney-Ni reduction as described above. Alternatively, Stille reaction of W with an organotin compound such as vinyltributyltin in the presence of catalytic amount of Pd(Ph₃P)₄ in DMF affords the compound of formula Z. After Raney-Ni reduction of compound Z, further hydrogenation in the presence of palladium on carbon as described above will afford the final compound of formula AB.

General reaction scheme 5: Chiral separation



[0017] Scheme 5 describes a method to separate compounds (I) into their enantiomeric isomers, wherein R^4 , Y , and R^5 are as described herein. The compound of formula **AC**, which may be prepared according to Scheme 1 or Scheme 2 or Scheme 3 or Scheme 4, can be converted to the Boc-protected compound **AD** by the reaction with an N-Protecting reagent such as Boc anhydride in the presence of base like triethylamine in dichloromethane. Racemic compound **AD** can be resolved via chiral HPLC using a chiral column such as ChiralPak® AD-H and $SF\ CO_2$ / methanol as eluent. Two compounds collected are enantiomer **AE** and enantiomer **AF**. Analysis of the enantiomeric purity of each isomer can be achieved using a chiral column such as ChiralPak® AD column. The Boc-protected compounds **AE** and **AF** can be converted to the final chiral compounds **AG** and **AH**, by deprotection using acid such as HCl in dioxane.

General reaction scheme 6: Chiral separation



[0018] Scheme 6 describes an alternative method to separate chiral compounds AI into their enantiomeric isomers, wherein R⁴, Y, and R⁵ are as described herein. The compound of formula AI may be prepared according to Scheme 1 or Scheme 2 or Scheme 3 or Scheme 4. The separation of the two enantiomers was achieved by dissolving the racemic material AI in a suitable solvent and applying to an appropriate chiral column and eluent system. The collected separated enantiomer samples were then concentrated and used in the next step without further purification. Using this technique, it is possible to achieve a range of enantiomeric excesses of the separated enantiomers. The nitro compound AJ and AK can be converted to the final chiral compounds AL and AM, respectively, by Raney-Ni reduction (Raney Ni, 2 equiv w/w, 2.0 M NH₃ in EtOH, absolute EtOH).

IV. Assays

[0019] Art-recognized techniques of genetics and molecular biology are of use to identify compounds that bind to and/or inhibit an enzyme, such as a tRNA synthetase. Moreover, these techniques are of use to distinguish whether a compound binds to and/or inhibits a particular domain of the enzyme. For example, for leucyl tRNA synthetase (LeuRS), these techniques can distinguish whether a compound binds to and/or inhibits the synthetic domain, the editing domain, or both the editing and synthetic domains. The *Mycobacterium tuberculosis leuS* gene was synthesized by Genscript® (Piscataway, NJ) using *E. coli* optimized codons and protein was made using standard T7 RNA polymerase over-expression protocols and standard purification protocols.

IV. a) *LeuRS*

[0117] In an exemplary assay, activity of a representative compound against the editing domain was confirmed. To identify the target of a novel boron-containing antibacterial compound, mutants in *E.coli* showing resistance to the compound were isolated. Characterization of mutants showed that they have an 32-256 fold increase in resistance to the compound over wildtype. The mutants were furthermore shown to be sensitive to various antibacterial agents with known modes of action, suggesting that the cellular target of the compound is distinct from the target of the other antibacterial agents. The *leuS* gene from the mutants was cloned onto a plasmid and their resistance was confirmed by MIC. The editing domain from these mutants were sequenced and the mutations were all located in the editing domain of this enzyme.

[0118] Assays to determine whether, and how effectively, a particular compound binds to and/or inhibits the editing domain of a selected tRNA synthetase are also set forth herein, and additional assays are readily available to those of skill in the art. Briefly, in an exemplary assay, an improperly charged tRNA and a tRNA synthetase that is capable of editing the improperly charged tRNA are combined. The resulting mixture is contacted with the putative inhibitor and the degree of editing inhibition is observed.

[0119] Another assay uses genetics to show that a drug works via the editing domain. In this assay, the compound is first tested against a strain of cells over-expressing copies of the tRNA synthetase gene. The compound's effect on the over-expressing strain is compared with a control strain to determine whether the compound is active against the synthetase. If the minimum inhibitory concentration (MIC) is 2-fold higher in the strain with extra copies of the synthetase gene than the MIC of the inhibitor against a wild type cell, a further genetic screen is conducted to determine whether the increased resistance is due to mutations in the editing domain. In this second screen, the control strain is challenged against a high concentration of the inhibitor. The colonies surviving the challenge are isolated and DNA from these cells is isolated. The editing domain is amplified using a proof-reading PCR enzyme and the appropriate primers. The PCR product can be purified using standard procedures. The sequence amplified mutant DNA is compared to wild-type. If the mutant DNA bears mutations in the editing domain, such results would suggest that

the compound binds to the editing domain and affects the editing function of the molecule through this domain.

[0120] Generally, the compounds to be tested are present in the assays in ranges from about 1 pM to about 100 mM, preferably from about 1 pM to about 1 μ M.

5 Other compounds range from about 1 nM to about 100 nM, preferably from about 1 nM to about 1 μ M.

[0121] The effects of the test compounds upon the function of the enzymes can also be measured by any suitable physiological change. When the functional consequences are determined using intact cells or animals, one can also measure a
10 variety of effects such as transmitter release, hormone release, transcriptional changes to both known and uncharacterized genetic markers, changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as Ca^{2+} , or cyclic nucleotides.

[0122] Utilizing the assays set forth herein and others readily available in the art,
15 those of skill in the art will be able to readily and routinely determine other compounds and classes of compounds that operate to bind to and/or inhibit the editing domain of tRNA synthetases.

[0123] In another aspect, the invention provides a method for identifying a compound which binds to an editing domain of a tRNA synthetase comprising:
20 a) contacting said editing domain with a test compound under conditions suitable for binding; and b) detecting binding of said test compound to said editing domain. In an exemplary embodiment, detecting binding of said compound comprises use of at least one detectable element, isotope, or chemical label attached to said compound. In an exemplary embodiment, the element, isotope or chemical label is detected by a
25 fluorescent, luminescent, radioactive, or absorbance readout. In an exemplary embodiment, the contacting of said test compound with said editing domain also includes further contacting said test compound and said editing domain with a member selected from AMP and a molecule with a terminal adenosine. In an exemplary embodiment, the tRNA synthetase is derived from leucyl tRNA synthetase.
30 In an exemplary embodiment, the tRNA synthetase is derived from a mutated tRNA synthetase, wherein said mutated tRNA synthetase comprises amino acid mutations in an editing domain. In another exemplary embodiment, wherein said editing domain

of a tRNA synthetase comprises the amino acid sequence of a peptide sequence described herein.

[0124] In another aspect, the invention provides a method for identifying a compound which binds to an editing domain of a tRNA synthetase, said assay comprising: a) contacting said editing domain of a tRNA synthetase with said compound under conditions suitable for binding of said compound with said editing domain of a tRNA synthetase; b) comparing a biological activity of said editing domain of a tRNA synthetase contacting said compound to said biological activity when not contacting said compound; and c) identifying said compound as binding to said editing domain of a tRNA synthetase if said biological activity of said editing domain of a tRNA synthetase is reduced when contacting said compound. In an exemplary embodiment, the biological activity is hydrolysis of noncognate amino acid. In another exemplary embodiment, the hydrolysis of said noncognate amino acid is detected through the use of one or more labels. In another exemplary embodiment, the labels include a radiolabel, a fluorescent marker, an antibody, or a combination thereof. In another exemplary embodiment, said labels can be detected using spectroscopy. In another exemplary embodiment, said editing domain of a tRNA synthetase is derived from leucyl tRNA synthetase.

[0125] In another aspect, the invention provides a method of generating a tRNA molecule with a noncognate amino acid comprising: a) creating or isolating a mutated tRNA synthetase with altered amino acid editing domains; and b) contacting a tRNA molecule with said mutated tRNA synthetase and a noncognate amino acid. In another exemplary embodiment, the mutated tRNA synthetase contains one or more amino acid mutations in an editing domain. In another exemplary embodiment, the mutated tRNA synthetase is unable to bind with a compound of the invention. In another exemplary embodiment, the mutated tRNA synthetase is unable to bind with a compound described herein, or a pharmaceutically acceptable salt thereof. In another exemplary embodiment, the mutated tRNA synthetase is unable to bind with a compound according to a formula described herein, or a pharmaceutically acceptable salt thereof.

[0126] In another aspect, the invention provides a composition that comprises one or more tRNA molecules attached to noncognate amino acids, wherein said tRNA

molecules are synthesized using one or more mutated tRNA synthetases isolated from a microorganism or a cell line derived from a microorganism. In an exemplary embodiment, the microorganism is a bacteria. In an exemplary embodiment, wherein said mutated tRNA synthetases contain amino acid mutations in their editing domains.

5 *V. Amino acid and nucleotide sequences used in assays*

[0020] Amino acid and nucleotide sequences of use in the invention are published in references described herein, such as U.S. Pat. Apps. 12/142,692 (U.S. Pat. Pub. 2009/0227541) and U.S. Pat. Pubs. US20060234981, US20070155699 and US20070293457. The sequence for the codon optimized *M. tuberculosis leuS* gene is
10 as follows:

```
CATATGACCGAAAGCCCCGACCGCAGGTCCGGGTGGTGTGCCGCGTGCGGA
TGATGCAGATAGCGATGTGCCGCGTTATCGTTATACCGCGGAACCTGGCGG
CGCGTCTGGAACGTACCTGGCAGGAAACTGGGCGCGTCTGGGCACCTTT
AACGTGCCGAACCCGGTGGGTAGCCTGGCACCGCCGGATGGTGCAGCAGT
15 GCCGGATGATAAACTGTTTGTGCAGGATATGTTTCCGTATCCGAGCGGCG
AAGGCCTGCATGTGGGCCATCCGCTGGGCTATATTGCGACCGATGTGTAT
GCGCGTTATTTTCGTATGGTGGGCCGTAACGTGCTGCATGCGCTGGGCTTT
GATGCGTTTGGTCTGCCGGCGGAACAGTATGCGGTGCAGACCGGCACCCA
TCCGCGTACCCGTACCGAAGCGAACGTGGTGAACCTTCGTCTGTCAGCTGG
20 GCCGTCTGGGCTTTGGCCATGATAGCCGTCGTAGCTTTAGCACCAACCGATG
TGGATTTTTATCGTTGGACCCAGTGGATTTTTCTGCAGATTTATAACGCGT
GGTTTGATACCAACCGCGAACAAAGCGCGTCCGATTAGCGAACTGGTGGCG
GAATTTGAAAGCGGTGCACGTTGCCTGGATGGTGGTTCGTGATTGGGCAA
ACTGACCGCAGGTGAACGTGCGGATGTGATTGATGAATATCGTCTGGTGT
25 ATCGTGCGGATAGCCTGGTGAACCTGGTGCCCGGGTCTGGGTACCGTGCTG
GCAAACGAAGAAGTGACCGCAGATGGCCGTAGCGATCGTGGCAACTTTCC
GGTGTTCGTAAACGTCTGCGTCAGTGGATGATGCGTATTAACCGCGTATGC
GGATCGTCTGCTGGATGATCTGGATGTGCTGGATTGGCCGGAACAGGTGA
AAACCATGCAGCGTAACTGGATTGGCCGTAGCACCGGCGCGGTGGCGCTG
30 TTTAGCGCGCGTGCGGCGAGCGATGATGGCTTTGAAGTGGATATTGAAGT
GTTTACCACCCGTCGCGGATACCCTGTTTGGCGCGACCTATCTGGTGCTGGC
GCCGGAACATGATCTGGTGGATGAACTGGTGGCGGCAAGCTGGCCGGCAG
GTGTGAACCCGCTGTGGACCTATGGCGGTGGTACCCCGGGTGAAGCAATT
GCAGCATATCGTCTGTGCGATTGCGGCGAAAAGCGATCTGGAACGTCAGGA
35 AAGCCGTGAAAAAACCGGCGTGTTTCTGGGCAGCTATGCGATTAACCCGG
CGAACGGCGAACCGGTGCCGATTTTTATTGCGGATTATGTGCTGGCGGGC
TATGGCACCGGCGCGATTATGGCGGTGCCGGGCCATGATCAGCGTGATTG
GGATTTTGCGCGTGCGTTTGGCCTGCCGATTGTGGAAGTGATTGCAGGTGG
AAACATTAGCGAAAGCGCGTATACCGGCGATGGCATTCTGGTGAACAGCG
40 ATTATCTGAACGGCATGAGCGTGCCGGCAGCAAAACGTGCAATTGTGGAT
CGTCTGGAAAGCGCAGGTCGTGGTTCGTGCACGTATTGAATTTAACTGCG
TGATTGGCTGTTTGCGCGTCAGCGTTATTGGGGCGAACCGTTTCCGATTGT
GTATGATAGCGATGGCCGTCCGCATGCGCTGGATGAAGCGGCGCTGCCGG
TGGAAC TGCCGGATGTGCCGGATTATAGCCCGGTGCTGTTTGATCCGGAT
```

GATGCGGATAGCGAACCGAGCCCGCCGCTGGCGAAAGCGACCGAATGGG
 TGCATGTGGATCTGGATCTGGGCGATGGCCTGAAACCGTATAGCCGTGAT
 ACCAACGTGATGCCGCAGTGGGCGGGCAGCAGCTGGTATGAACTGCGTTA
 TACCGATCCGCATAACAGCGAACGTTTTTTCGCGGAAAGAAAACGAAGCGT
 5 ATTGGATGGGTCCGCGTCCGGCAGAACATGGTCCGGATGATCCGGGTGGT
 GTGGATCTGTATGTGGGCGGGCGCGGAACATGCGGTGCTGCATCTGCTGTA
 TAGCCGTTTTTGGCATAAAGTGCTGTATGATCTGGGGCCATGTGAGCAGCCG
 TGAACCGTATCGTCGTCTGGTGAACCAGGGCTATATTCAGGCGTATGCGT
 ATACCGATGCGCGTGGCAGCTATGTGCCGGCGGAACAAGTGATTGAACGT
 10 GGCGATCGTTTTGTGTATCCGGGGCCCGGATGGCGAAGTGGAAGTGTTTCA
 GGAATTTGGCAAAAATTGGCAAAAAGCCTGAAAAACAGCGTGAGCCCGGAT
 GAAATTTGCGATGCGTATGGCGCGGATACCCTGCGTGTGTATGAAATGAG
 CATGGGCCCCGCTGGAAGCGAGCCGTCCGTGGGCGACCAAAGATGTGGTGG
 GCGCGTATCGTTTTCTGCAGCGTGTGTGGCGTCTGGTGGTGGATGAACATA
 15 CCGGCGAAACCCGTGTGGCGGATGGCGTGGAAGTGGATATTGATACCCTG
 CGTGCGCTGCATCGTACCATTGTGGGCGTGAGCGAAGATTTTTCGGGCGCT
 GCGTAACAACACCGCGACCGCGAAACTGATTGAATATACCAACCATCTGA
 CCAAAAAACATCGTGATGCAGTGCCGCGTGCGGCAGTGGAACCGCTGGTG
 CAGATGCTGGCACCGCTGGCACCGCATATTGCGGAAGAAGTGTGGCTGCG
 20 TCTGGGCAACACCACCAGCCTGGCGCATGGCCCCGTTTCCGAAAGCGGATG
 CGGCGTATCTGGTGGATGAAACCGTGGAATATCCGGTGCAGGTGAACGGC
 AAAGTGCGTGGTCGTGTGGTGGTGGCGGCGGATACCGATGAAGAAACCCT
 GAAAGCGGCGGTGCTGACCGATGAAAAAGTGCAGGCGTTTCTGGCGGGC
 GCGACCCCGCGTAAAGTGATTGTGGTGGCGGGCCGTCTGGTGAACCTGGT
 25 GATTAACTCGAG

VI. Methods

[0128] In another aspect, the compounds of the invention can be utilized to inhibit an enzyme. In another aspect, the compounds of the invention and/or combinations of the invention exhibit potency against microorganisms, such as bacteria, and therefore have the potential to kill and/or inhibit the growth of microorganisms. In another aspect, the compounds of the invention and/or combinations of the invention exhibit potency against microorganisms, such as bacteria, and therefore have the potential to achieve therapeutic efficacy in the animals described herein.

VI. a) *LeuRS*

35 [0129] In an exemplary embodiment, the compounds of the invention exhibit the ability of inhibiting the editing domain of tRNA synthetases, such as leucyl tRNA synthetase, of microorganisms, such as bacteria, and therefore have the potential to be used as editing domain inhibitors of microorganism tRNA synthetases.

40 [0130] According to another aspect of the invention, a method for binding to and/or inhibiting the editing domain of a tRNA synthetase is provided which comprises contacting a tRNA synthetase with a compound of the invention that inhibits the

editing domain under conditions in which the tRNA synthetase interacts with its substrate to form an aminoacyl adenylate intermediate and, preferably, to form a charged tRNA. Such conditions are known to those skilled in the art. In an exemplary embodiment, the compound has a structure according to a formula

5 described herein. In an exemplary embodiment, the compound is described herein, or a salt, hydrate or solvate thereof, or a combination thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt, hydrate or solvate thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt thereof. In an exemplary embodiment, the invention

10 provides a compound described herein, or a salt thereof. The tRNA synthetase is contacted with an amount of compound of the invention sufficient to result in a detectable amount of tRNA synthetase inhibition. This method can be performed on a tRNA synthetase that is contained within an organism or which is outside an organism. In an exemplary embodiment, the method is performed on a tRNA

15 synthetase that is contained within a microorganism or a microbial cell that is in, or on the surface of, an animal. In an exemplary embodiment, the animal is a human. The method results in a decrease in the amount of charged tRNA produced by the tRNA synthetase that has an inhibited editing domain. In an exemplary embodiment, the inhibition takes place in a cell, such as a microorganism cell. In another exemplary

20 embodiment, the microorganism cell is a bacteria. In another exemplary embodiment, the tRNA synthetase is leucyl tRNA synthetase.

[0131] In an exemplary embodiment, the invention provides a method of inhibiting conversion of a tRNA molecule into a charged tRNA molecule. The method involves contacting a tRNA synthetase with a compound of the invention effective to inhibit

25 activity of an editing domain of said tRNA synthetase, under conditions sufficient to inhibit said activity, thereby inhibiting said conversion. In an exemplary embodiment, the compound of the invention is a compound described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the inhibition occurs within a cell, and the cell is a microorganism cell. In another exemplary embodiment, the

30 microorganism cell is a bacteria. In another exemplary embodiment, the microorganism cell is a bacteria which is described herein. In another exemplary embodiment, the enzyme is a leucyl tRNA synthetase of a bacteria described herein. In another exemplary embodiment, the tRNA synthetase is leucyl tRNA synthetase.

In another exemplary embodiment, the compound has a $K_{D, \text{synthesis}}$ of greater than 100 μM against a synthetic domain of said tRNA synthetase.

[0132] In certain embodiments, the mechanism of action of a compound of the invention is to inhibit the conversion of a tRNA molecule into a charged tRNA molecule by binding to and/or inhibiting at least the editing domain of the synthetase. The compounds of use in this method may also inhibit or otherwise interact with the synthetic domain (e.g., the active site of the synthetic domain). In a presently preferred embodiment, the editing domain is inhibited selectively in the presence of the synthetic domain. In a preferred embodiment, the synthetic domain is essentially uninhibited, while the editing domain is inhibited at least 50%, preferably at least 60%, more preferably at least 70%, still more preferably, at least 80% and even still more preferably at least 90% of the activity of the tRNA synthetase. In another preferred embodiment, the synthetic domain is inhibited by at most 50%, preferably at most 30%, preferably at most 20%, 10%, preferably at most 8%, more preferably at most 5%, still more preferably, at most 3% and even still more preferably at most 1%. Inhibition of the editing domain produces a decrease in the amount of the properly charged tRNA which results in retardation or cessation of cell growth and division.

[0133] In another exemplary embodiment, the ratio of a minimum concentration of said compound inhibiting said editing domain to a minimum concentration of said compound inhibiting said synthetic domain of said tRNA synthetase, represented as $K_{D, \text{edit}}/K_{D, \text{synthesis}}$, is less than one. In another exemplary embodiment, the $K_{D, \text{edit}}/K_{D, \text{synthesis}}$ of the compound is a member selected from less than 0.5, less than 0.1 and less than 0.05.

VI. b) Inhibiting Microorganism Growth or Killing Microorganisms

[0134] The compounds of the invention and/or combinations of the invention exhibit potency against microorganisms, such as bacteria, and therefore have the potential to treat, and/or prevent a microorganism infection, or kill and/or inhibit the growth of microorganisms.

[0135] In a further aspect, the invention provides a method of treating and/or preventing a microorganism infection, or a method of killing and/or inhibiting the growth of a microorganism, said method comprising: contacting said microorganism with an effective amount of a compound of the invention, thereby killing and/or

inhibiting the growth of the microorganism. In a further aspect, the invention provides a method of treating and/or preventing a microorganism infection, or a method of killing and/or inhibiting the growth of a microorganism, said method comprising: contacting said microorganism with an effective amount of a combination of the invention, thereby killing and/or inhibiting the growth of the microorganism.

[0136] In a further aspect, the invention provides a method of treating a bacterial infection comprising administering to an animal suffering from the infection an effective amount of a compound of the invention or a combination of the invention, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial infection. In an exemplary embodiment, the invention provides a method of treating a bacterial infection comprising administering to an animal suffering from the infection an effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, and an effective amount of an antibiotic, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial infection.

[0137] In a further aspect, the invention provides a method of preventing a bacterial infection comprising administering to an animal a prophylactic amount of a compound of the invention or a combination of the invention, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial infection. In an exemplary embodiment, the invention provides a method of preventing a bacterial infection comprising administering to an animal a prophylactic amount of a compound of the invention, or a pharmaceutically acceptable salt thereof.

[0138] In an exemplary embodiment, the microorganism is a bacteria. In an exemplary embodiment, the compound or combination is described herein, or a salt, prodrug, hydrate or solvate thereof, or a combination thereof. In an exemplary embodiment, the invention provides a compound or combination described herein, or a salt, hydrate or solvate thereof. In an exemplary embodiment, the invention provides a compound or combination described herein, or a salt thereof. In another exemplary embodiment, the compound or combination of the invention is a compound or combination described herein, or a pharmaceutically acceptable salt thereof. In another exemplary embodiment, the compound or combination of the combination is described by a formula listed herein, or

a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the compound is part of a combination described herein. In an exemplary embodiment, the compound is part of a pharmaceutical formulation described herein. In another exemplary embodiment, the contacting occurs under conditions which permit entry of the compound into the organism. Such conditions are known to one skilled in the art and are described herein.

[0139] In another aspect, the microorganism is inside, or on the surface of an animal. In another exemplary embodiment, the animal is described herein. In another exemplary embodiment, the animal is a human.

10 [0140] In an exemplary embodiment, the microorganism infection is treated and/or prevented, or the microorganism is killed or its growth is inhibited, through oral administration of the compound of the invention and/or the combination of the invention. In an exemplary embodiment, the microorganism infection is treated and/or prevented, or the microorganism is killed or its growth is inhibited through
15 intravenous administration of the compound of the invention and/or the combination of the invention.

[0141] In an exemplary embodiment, the microorganism is a bacterium. In an exemplary embodiment, an infection is caused by and/or associated with a microorganism, particularly a bacterium. In an exemplary embodiment, the bacterium
20 is a gram-positive bacteria. In another exemplary embodiment, the gram-positive bacterium is selected from the group consisting of *Staphylococcus* species, *Streptococcus* species, *Bacillus* species, *Mycobacterium* species, *Corynebacterium* species (*Propionibacterium* species), *Clostridium* species, *Actinomyces* species, *Enterococcus* species and *Streptomyces* species. In another exemplary embodiment,
25 the gram-positive bacterium is selected from the group consisting of *Propionibacterium acnes*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Staphylococcus haemolyticus*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Enterococcus faecalis*, *Enterococcus faecium*, *Actinomyces israelii*, *Bacillus anthracis*,
30 *Corynebacterium diphtheria*, *Clostridium perfringens*, *Clostridium botulinum*, *Clostridium tetani*, and *Clostridium difficile*. In another exemplary embodiment, the gram-positive bacterium is selected from the group consisting of *Staphylococcus*

aureus, *Staphylococcus epidermidis*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Enterococcus faecalis*, *Enterococcus faecium*, *Clostridium difficile* and *Propionibacter acnes*. In another exemplary embodiment, the bacterium is a gram-negative bacterium. In another exemplary embodiment, the gram-negative bacterium

5 is selected from the group consisting of *Acinetobacter* species, *Neisseria* species, *Pseudomonas* species, *Brucella* species, *Agrobacterium* species, *Bordetella* species, *Escherichia* species, *Shigella* species, *Yersinia* species, *Salmonella* species, *Klebsiella* species, *Enterobacter* species, *Haemophilus* species, *Pasteurella* species, *Streptobacillus* species, spirochetal species, *Campylobacter* species, *Vibrio* species,

10 *Helicobacter* species, *Bacteroides* species, *Citrobacter* species, *Proteus* species, *Providencia* species, *Serratia* species, *Stenotrophomonas* species and *Burkholderia* species. In another exemplary embodiment, the gram-negative bacterium is selected from the group consisting of *Acinetobacter* species, *Pseudomonas* species, *Escherichia* species, *Klebsiella* species, *Enterobacter* species, *Bacteroides* species,

15 *Citrobacter* species, *Proteus* species, *Providencia* species, *Serratia* species, *Stenotrophomonas* species and *Burkholderia* species. In another exemplary embodiment, the gram-negative bacterium is selected from the group consisting of *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa*, *Legionella pneumophila*, *Escherichia coli*, *Yersinia pestis*, *Haemophilus influenzae*, *Helicobacter*

20 *pylori*, *Campylobacter fetus*, *Campylobacter jejuni*, *Vibrio cholerae*, *Vibrio parahemolyticus*, *Treponema pallidum*, *Rickettsia prowazekii*, *Rickettsia rickettsii*, *Chlamydia trachomatis*, *Chlamydia psittaci*, *Brucella abortus*, *Agrobacterium tumefaciens*, *Francisella tularensis*, *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Acinetobacter baumannii*, *Bacteroides fragilis*, *Citrobacter freundii*, *Proteus*

25 *mirabilis*, *Providencia stuartii*, *Serratia marcescens*, *Stenotrophomonas maltophilia* and *Burkholderia cepacia*. In another exemplary embodiment, the gram-negative bacterium is selected from the group consisting of *Pseudomonas aeruginosa*, *Escherichia coli*, *Haemophilus influenzae*, *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Acinetobacter baumannii*, *Bacteroides fragilis*, *Citrobacter freundii*, *Proteus*

30 *mirabilis*, *Providencia stuartii*, *Serratia marcescens*, *Stenotrophomonas maltophilia* and *Burkholderia cepacia*. In another exemplary embodiment, the gram-negative bacterium is selected from the group consisting of *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *Escherichia coli*, *Klebsiella*

pneumoniae, *Proteus mirabilis*, *Serratia marcescens* and *Citrobacter freundii*. In another exemplary embodiment, the gram-negative bacterium is a *Providencia* spp..

[0142] In an exemplary embodiment, the microorganism is an acid-fast bacteria.

In another exemplary embodiment, the bacterium is a *Mycobacterium* spp.. In

5 another exemplary embodiment, the bacterium is *Mycobacterium avium*. In another exemplary embodiment, the bacterium is *Mycobacterium avium-intracellulare*. In another exemplary embodiment, the bacterium is *Mycobacterium kansasii*. In another exemplary embodiment, the bacterium is *Mycobacterium leprae*. In another exemplary embodiment, the bacterium is *Mycobacterium lepromatosis*. In another
10 exemplary embodiment, the bacterium is *Mycobacterium africanum*. In another exemplary embodiment, the bacterium is *Mycobacterium canetti*. In another exemplary embodiment, the bacterium is *Mycobacterium microti*. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis*. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is multi-
15 drug resistant. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is extensively drug resistant. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is resistant to rifampicin. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is resistant to isoniazid. In another exemplary embodiment, the bacterium is
20 *Mycobacterium tuberculosis* which is resistant to kanamycin. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is resistant to capreomycin. In another exemplary embodiment, the bacterium is *Mycobacterium tuberculosis* which is resistant to amikacin.

[0143] In another exemplary embodiment, the bacterium is a *Pseudomonas*

25 species. In another exemplary embodiment, the bacterium is *Pseudomonas aeruginosa*. In another exemplary embodiment, the bacterium is selected from the group consisting of *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Stenotrophomonas maltophilia* and *Burkholderia cepacia*. In another exemplary embodiment, the bacterium is *Acinetobacter baumannii*. In another exemplary
30 embodiment, the bacterium is *Stenotrophomonas maltophilia*. In another exemplary embodiment, the bacterium is *Burkholderia cepacia*. In another exemplary embodiment, the bacterium is *Acinetobacter* species. In another exemplary embodiment, the bacterium is *Acinetobacter anitratus*. In another exemplary

embodiment, the bacterium is selected from the group consisting of *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *E. coli*, *K. pneumoniae*, *P. mirabilis*, *Serratia marcescens*, *Citrobacter freundii* and *Providencia spp.* In another exemplary embodiment, the bacterium is selected from the group consisting of

5 *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *E. coli*, *K. pneumoniae*, *P. mirabilis*, *Serratia marcescens*, *Citrobacter freundii*, *Providencia spp.*, *S. aureus*, *S. pneumonia*, *S. pyogenes*, *E. faecalis*, and *E. faecium*. In another exemplary embodiment, the bacterium is selected from the group consisting of

10 *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*. In another exemplary embodiment, the bacterium is selected from the group consisting of *S. aureus*, *S. pneumonia*, *S. pyogenes*, *E. faecalis*, and *E. faecium*. In another exemplary embodiment, the bacterium is selected from the group consisting of Viridans group *Strep.*. In another exemplary embodiment, the bacterium is selected from the group consisting of *Strep. mitis*, *Strep. mutans*, *Strep. oralis*,

15 *Strep. sanguis*, *Strep. sobrinus* and *Strep. millari*. In another exemplary embodiment, the bacterium is *S. pneumonia*. In another exemplary embodiment, the bacterium is *H. influenzae*. In another exemplary embodiment, the bacterium is *S. aureus*. In another exemplary embodiment, the bacterium is *M. catarrhalis*. In another exemplary embodiment, the bacterium is *M. pneumoniae*. In another exemplary

20 embodiment, the bacterium is *L. pneumoniae*. In another exemplary embodiment, the bacterium is *C. pneumoniae*. In another exemplary embodiment, the bacterium is *S. pyogenes*. In another exemplary embodiment, the bacterium is an anaerobe. In another exemplary embodiment, the bacterium is an *Alcaligenes* species. In another exemplary embodiment, the bacterium is a *B. cepacia*. In another exemplary

25 embodiment, the bacterium is selected from the group consisting of *Enterobacter cloacae*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Providencia stuartii*, *Serratia marcescens*, and *Citrobacter freundii*. In another exemplary embodiment, the bacterium is resistant to methicillin. In another exemplary embodiment, the bacterium is methicillin-resistant *Staphylococcus aureus*. In another

30 exemplary embodiment, the bacterium is selected from the group consisting of *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Staphylococcus aureus*, *Mycobacterium catarrhalis*, *Mycobacterium pneumoniae*, *Legionella pneumophila* and *Chlamydia pneumoniae*. In another exemplary embodiment, the bacterium is selected from the group consisting of *Enterobacter cloacae*, *Escherichia coli*,

Klebsiella pneumoniae, *Proteus mirabilis*, *Serratia marcescens*, *Citrobacter freundii*,
Providencia stuartii, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*,
Stenotrophomonas maltophilia, *Burkholderia cepacia*, *Staphylococcus aureus*,
Streptococcus pneumoniae, *Streptococcus pyogenes*, *Enterococcus faecalis*, and
5 *Enterococcus faecium*. In another exemplary embodiment, the bacterium is selected
from the group consisting of *Staphylococcus aureus*, *Staphylococcus epidermidis*,
Staphylococcus haemolyticus, *Streptococcus pyogenes*, *Streptococcus agalactiae* and
Streptococcus pneumoniae.

[0144] In an exemplary embodiment, the microorganism is a bacterium, which is
10 selected from the group consisting of bacilli, including *Bacillus* species,
Corynebacterium species (also *Propionibacterium*) and *Clostridium* species;
filamentous bacteria, including *Actinomyces* species and *Streptomyces* species; bacilli,
such as *Pseudomonas* species, *Brucella* species, *Agrobacterium* species, *Bordetella*
species, *Escherichia* species, *Shigella* species, *Yersinia* species, *Salmonella* species,
15 *Klebsiella* species, *Enterobacter* species, *Haemophilus* species, *Pasteurella* species,
and *Streptobacillus* species; spirochetal species, *Campylobacter* species, *Vibrio*
species; and intracellular bacteria including *Rickettsiae* species and *Chlamydia*
species.

VI. b) Microorganism Infection

20 [0145] The compounds of the invention and/or combinations of the invention
exhibit potency against microorganisms, such as bacteria, and therefore have the
potential to be used to treat and/or prevent a microorganism infection, such as a
bacterial infection.

[0146] In a further aspect, the invention provides a method of treating a bacterial
25 infection comprising administering to an animal suffering from the infection an
effective amount of a compound of the invention, or a pharmaceutically acceptable salt
thereof, thereby treating the bacterial infection. In an exemplary embodiment, the
invention provides a method of treating a bacterial infection comprising administering
to an animal suffering from the infection an effective amount of a compound of the
30 invention, or a pharmaceutically acceptable salt thereof, and an effective amount of an
antibiotic, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial
infection.

[0147] In a further aspect, the invention provides a method of preventing a bacterial infection comprising administering to an animal a prophylactic amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial infection. In an exemplary embodiment, the invention provides a method of preventing a bacterial infection comprising administering to an animal a prophylactic amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, and an effective amount of an antibiotic, or a pharmaceutically acceptable salt thereof, thereby treating the bacterial infection.

VI. c) Diseases

[0148] The compounds of the invention and/or combinations of the invention exhibit potency against microorganisms, such as bacteria, and therefore have the potential to achieve therapeutic efficacy in the animals described herein.

[0149] In another aspect, the invention provides a method of treating and/or preventing a disease. In an exemplary embodiment, the method includes administering to the animal a therapeutically effective amount of a compound of the invention, thereby treating and/or preventing the disease. In an exemplary embodiment, the method includes administering to the animal a therapeutically effective amount of a combination of the invention, thereby treating and/or preventing the disease. In an exemplary embodiment, the compound of the invention or the combination of the invention can be used in human or veterinary medical therapy, particularly in the treatment or prophylaxis of bacterial-associated disease. In an exemplary embodiment, the compound is described herein, or a salt, prodrug, hydrate or solvate thereof, or a combination thereof. In an exemplary embodiment, the invention provides a compound described herein, or a prodrug thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt, hydrate or solvate thereof. In an exemplary embodiment, the invention provides a compound described herein, or a salt thereof. In another exemplary embodiment, the compound of the invention is a compound described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the compound is a compound described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the compound is according to a formula described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the compound is part of a combination described herein. In an exemplary embodiment,

the compound is part of a pharmaceutical formulation described herein. In another exemplary embodiment, the disease is a systemic disease. In another exemplary embodiment, the disease is a topical disease. In an exemplary embodiment, the animal being administered the compound is not otherwise in need of treatment with the compound.

[0150] In an exemplary embodiment, the disease is treated through oral administration of a compound of the invention and/or a combination of the invention. In an exemplary embodiment, the disease is treated through intravenous administration of a compound of the invention and/or a combination of the invention. In an exemplary embodiment, the disease is treated through subcutaneous administration of a compound of the invention and/or a combination of the invention.

Systemic Diseases

[0151] In another aspect, the invention provides a method of treating a systemic disease. The method involves contacting an animal with a compound of the invention and/or a combination of the invention.

[0152] In another exemplary embodiment, the disease is associated with a bacteria described herein. In another exemplary embodiment, the disease is associated with infection by a Gram-positive bacteria. In an exemplary embodiment, the disease is associated with a *Staphylococcus* species. In another exemplary embodiment, the disease is selected from the group consisting of pneumonia, gastroenteritis, toxic shock syndrome, community acquired pneumonia (CAP), meningitis, septic arthritis, urinary tract infection, bacteremia, endocarditis, osteomyelitis, skin and skin-structure infection. In an exemplary embodiment, the disease is associated with a *Streptococcus* species. In another exemplary embodiment, the disease is selected from the group consisting of strep throat, skin infections, necrotizing fasciitis, toxic shock syndrome, pneumonia, otitis media and sinusitis. In an exemplary embodiment, the disease is associated with an *Actinomyces* species. In another exemplary embodiment, the disease is actinomycosis. In an exemplary embodiment, the disease is associated with a *Norcardia* species. In another exemplary embodiment, the disease is pneumonia. In an exemplary embodiment, the disease is associated with a *Corynebacterium* species. In another exemplary embodiment, the disease is diphtheria. In an exemplary embodiment, the disease is associated with a

Listeria species. In another exemplary embodiment, the disease is meningitis. In an exemplary embodiment, the disease is associated with a *Bacillus* species. In another exemplary embodiment, the disease is anthrax or food poisoning. In an exemplary embodiment, the disease is associated with a *Clostridium* species. In another
 5 exemplary embodiment, the disease is selected from the group consisting of botulism, tetanus, gas gangrene and diarrhea.

[0153] In an exemplary embodiment, the disease is associated with a *Mycobacterium* species. In an exemplary embodiment, the disease is associated with *Mycobacterium tuberculosis*. In an exemplary embodiment, the disease is associated
 10 with *Mycobacterium kansasii*. In an exemplary embodiment, the disease is associated with *Mycobacterium avium-intracellulare*. In another exemplary embodiment, the disease is leprosy. In another exemplary embodiment, the disease is tuberculosis. In another exemplary embodiment, the disease is pulmonary tuberculosis. In another exemplary embodiment, the disease is extrapulmonary tuberculosis. In another
 15 exemplary embodiment, the disease is associated with multi-drug resistant tuberculosis. In another exemplary embodiment, the disease is associated with extensively drug resistant tuberculosis.

[0154] In another exemplary embodiment, the disease is associated with infection by a Gram-negative bacteria. In an exemplary embodiment, the disease is associated
 20 with a *Neisseria* species. In another exemplary embodiment, the disease is selected from the group consisting of meningitis, gonorrhea, otitis externa and folliculitis. In an exemplary embodiment, the disease is associated with an *Escherichia* species. In another exemplary embodiment, the disease is selected from the group consisting of diarrhea, urinary tract infections, meningitis, sepsis and HAP. In an exemplary
 25 embodiment, the disease is associated with a *Shigella* species. In another exemplary embodiment, the disease is selected from the group consisting of diarrhea, bacteremia, endocarditis, meningitis and gastroenteritis. In an exemplary embodiment, the disease is associated with a *Salmonella* species. In another exemplary embodiment, the disease is selected from the group consisting of Typhoid fever, sepsis, gastroenteritis,
 30 endocarditis, sinusitis and meningitis. In an exemplary embodiment, the disease is associated with a *Yersinia* species. In another exemplary embodiment, the disease is selected from the group consisting of Typhoid fever, bubonic plague, enteric fever and gastroenteritis. In an exemplary embodiment, the disease is associated with a

Klebsiella species. In another exemplary embodiment, the disease is sepsis or urinary tract infection. In an exemplary embodiment, the disease is associated with a *Proteus* species. In another exemplary embodiment, the disease is an urinary tract infection. In an exemplary embodiment, the disease is associated with an *Enterobacter* species.

5 In another exemplary embodiment, the disease is a hospital-acquired infection. In an exemplary embodiment, the disease is associated with a *Serratia* species. In another exemplary embodiment, the disease is selected from the group consisting of a urinary tract infection, skin and skin-structure infection and pneumonia. In an exemplary embodiment, the disease is associated with a *Vibrio* species. In another exemplary

10 embodiment, the disease is cholera or gastroenteritis. In an exemplary embodiment, the disease is associated with a *Campylobacter* species. In another exemplary embodiment, the disease is gastroenteritis. In an exemplary embodiment, the disease is associated with a *Helicobacter* species. In another exemplary embodiment, the disease is chronic gastritis. In an exemplary embodiment, the disease is associated

15 with a *Pseudomonas* species. In another exemplary embodiment, the disease is selected from the group consisting of pneumonia, osteomyelitis, burn-wound infections, sepsis, UTIs, endocarditis, otitis and corneal infections. In an exemplary embodiment, the disease is associated with a *Bacteroides* species. In another exemplary embodiment, the disease is periodontal disease or aspiration pneumonia.

20 In an exemplary embodiment, the disease is associated with a *Haemophilus* species. In another exemplary embodiment, the disease is selected from the group consisting of meningitis, epiglottitis, septic arthritis, sepsis, chancroid and vaginitis. In an exemplary embodiment, the disease is associated with a *Bordetella* species. In another exemplary embodiment, the disease is Whooping cough. In an exemplary

25 embodiment, the disease is associated with a *Legionella* species. In another exemplary embodiment, the disease is pneumonia or pontiac fever. In an exemplary embodiment, the disease is associated with a *Francisella* species. In another exemplary embodiment, the disease is tularemia. In an exemplary embodiment, the disease is associated with a *Brucella* species. In another exemplary embodiment, the

30 disease is brucellosis. In an exemplary embodiment, the disease is associated with a *Pasteurella* species. In another exemplary embodiment, the disease is a skin infection. In an exemplary embodiment, the disease is associated with a *Gardnerella* species. In another exemplary embodiment, the disease is vaginitis. In an exemplary embodiment, the disease is associated with a *Spirochetes* species. In another

exemplary embodiment, the disease is syphilis or Lyme disease. In an exemplary embodiment, the disease is associated with a *Chlamydia* species. In another exemplary embodiment, the disease is chlamydia. In an exemplary embodiment, the disease is associated with a *Rickettsiae* species. In another exemplary embodiment, the disease is Rocky Mountain spotted fever or typhus.

[0155] In an exemplary embodiment, the disease is associated with *Mycoplasma pneumoniae*. In another exemplary embodiment, the disease is tracheobronchitis or walking pneumonia. In an exemplary embodiment, the disease is associated with *Ureaplasma urealyticum*. In another exemplary embodiment, the disease is urethritis. In another exemplary embodiment, the disease is pyelonephritis. In another exemplary embodiment, the disease is an intra-abdominal infection. In another exemplary embodiment, the disease is febrile neutropenia. In another exemplary embodiment, the disease is a pelvic infection. In another exemplary embodiment, the disease is bacteraemia. In another exemplary embodiment, the disease is septicaemia.

[0156] In an exemplary embodiment, the disease is an acute exacerbation of chronic obstructive pulmonary disease. In an exemplary embodiment, the disease is chronic obstructive pulmonary disease. In an exemplary embodiment, the disease is pharyngitis. In an exemplary embodiment, the disease is tonsillitis. In an exemplary embodiment, the disease is Acute Exacerbation of Chronic Bronchitis (AECB). In an exemplary embodiment, the disease is cervicitis. In an exemplary embodiment, the disease is genital ulcer disease.

[0157] In an exemplary embodiment, for any of the methods described herein, the animal is selected from the group consisting of human, cattle, deer, reindeer, goat, honey bee, pig, sheep, horse, cow, bull, dog, guinea pig, gerbil, rabbit, cat, camel, yak, elephant, ostrich, otter, chicken, duck, goose, guinea fowl, pigeon, swan, and turkey. In another exemplary embodiment, for any of the methods described herein, the animal is selected from the group consisting of a human, cattle, goat, pig, sheep, horse, cow, bull, dog, guinea pig, gerbil, rabbit, cat, chicken and turkey. In another exemplary embodiment, for any of the methods described herein, the animal is a human.

[0158] In an exemplary embodiment, for any of the methods described herein, a compound of the invention, a combination of the invention, a compound described

herein or a pharmaceutically acceptable salt thereof, or combination described herein, and/or a pharmaceutical formulation described herein can be used.

VII. Pharmaceutical Formulation

5 [0159] In another aspect, the invention provides a pharmaceutical formulation comprising: a) a compound of the invention; and b) a pharmaceutically acceptable excipient. In another aspect, the invention provides a pharmaceutical formulation comprising: a) a combination of the invention; and b) a pharmaceutically acceptable excipient. In an exemplary embodiment, the compound is according to a formula described herein. In an exemplary embodiment, the compound is according to an
10 example described herein. In an exemplary embodiment, the compound of the invention or combination of the invention is a compound described herein or combination described herein, or a pharmaceutically acceptable salt thereof. In an exemplary embodiment, the compound of the invention is a compound described herein.

15 [0160] In an exemplary embodiment, the compound of the invention is present in the pharmaceutical formulation in an amount of between about 0.0001% to about 60% (w/w). In an exemplary embodiment, the amount is between about 0.01% to about 10% (w/w). In an exemplary embodiment, the amount is between about 0.1% to about 10% (w/w). In an exemplary embodiment, the amount is between about 0.25%
20 to about 6% (w/w). In an exemplary embodiment, the amount is between about 0.5% to about 5% (w/w). In an exemplary embodiment, the amount is between about 0.1% and about 1.0% (w/w). In an exemplary embodiment, the amount is between about 1.0% and about 2.0% (w/w). In an exemplary embodiment, the amount is between about 2.0% and about 3.0% (w/w). In an exemplary embodiment, the amount is
25 between about 3.0% and about 4.0% (w/w). In an exemplary embodiment, the amount is between about 4.0% and about 5.0% (w/w).

[0161] The pharmaceutical formulations of the invention can take a variety of forms adapted to the chosen route of administration. Those skilled in the art will recognize various synthetic methodologies that may be employed to prepare non-toxic
30 pharmaceutical formulations incorporating the compounds described herein. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable solvents that may be used to prepare solvates of the compounds of the

invention, such as water, ethanol, propylene glycol, mineral oil, vegetable oil and dimethylsulfoxide (DMSO).

[0162] The compositions of the invention may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing
5 conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. It is further understood that the best method of administration may be a combination of methods. Oral administration in the form of a pill, capsule, elixir, syrup, lozenge, troche, or the like is particularly preferred. The term parenteral as used herein includes subcutaneous injections, intradermal, intravascular (e.g., intravenous),
10 intramuscular, spinal, intrathecal injection or like injection or infusion techniques.

[0163] The pharmaceutical formulations containing compounds of the invention are preferably in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

15 [0164] Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical formulations, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain
20 the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or
25 acacia; lubricating agents, for example magnesium stearate, stearic acid or talc; and extenders and bulking agents, such as microcrystalline cellulose. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or
30 glyceryl distearate may be employed.

[0165] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example,

calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

[0166] Aqueous suspensions contain the active materials in admixture with
5 excipients suitable for the manufacture of aqueous suspensions. Such excipients are
suspending agents, for example sodium carboxymethylcellulose, methylcellulose,
hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum
tragacanth and gum acacia; and dispersing or wetting agents, which may be a
naturally-occurring phosphatide, for example, lecithin, or condensation products of an
10 alkylene oxide with fatty acids, for example polyoxyethylene stearate, or
condensation products of ethylene oxide with long chain aliphatic alcohols, for
example heptadecacetylencoxycetanol, or condensation products of ethylene oxide
with partial esters derived from fatty acids and a hexitol such as polyoxyethylene
sorbitol monooleate, or condensation products of ethylene oxide with partial esters
15 derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan
monooleate. The aqueous suspensions may also contain one or more preservatives,
for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one
or more flavoring agents, and one or more sweetening agents, such as sucrose or
saccharin.

20 [0167] Oily suspensions may be formulated by suspending the active ingredients
in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a
mineral oil such as liquid paraffin. The oily suspensions may contain a thickening
agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such
as those set forth above, and flavoring agents may be added to provide palatable oral
25 preparations. These compositions may be preserved by the addition of an anti-oxidant
such as ascorbic acid.

[0168] Dispersible powders and granules suitable for preparation of an aqueous
suspension by the addition of water provide the active ingredient in admixture with a
dispersing or wetting agent, suspending agent and one or more preservatives. Suitable
30 dispersing or wetting agents and suspending agents are exemplified by those already
mentioned above. Other dispersing agents include hydrophilic polymers, electrolytes,
TweenTM 60 or 80, PEG, polyvinylpyrrolidone (PVP; commercially known as

PlasdoneTM), and the carbohydrate-based dispersing agents such as, for example, hydroxypropylcellulose and hydroxypropylcellulose ethers (e.g., HPC, HPC-SL, and HPC-L), hydroxypropylmethylcellulose and hydroxypropylmethylcellulose ethers (e.g. HPMC K100, HPMC K4M, HPMC K15M, and HPMC K100M),
5 carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate stearate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), polyvinylpyrrolidone/vinyl acetate copolymer (PlasdoneTM, e.g., S-630), 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and
10 formaldehyde (also known as tyloxapol), poloxamers (e.g., Pluronic F68TM, F88TM, and F108TM, which are block copolymers of ethylene oxide and propylene oxide); and poloxamines (e.g., Tetronic 9080, also known as Poloxamine 9080, which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF® Corporation, Parsippany, N.J.)).
15 Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[0021] Pharmaceutical formulations of the invention may also be in the form of oil-in-water emulsions and water-in-oil emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or
20 mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth; naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol; anhydrides, for example sorbitan monooleate; and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan
25 monooleate. The emulsions may also contain sweetening and flavoring agents.

[0022] Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, and flavoring and coloring agents. The pharmaceutical formulations may be in the form of a sterile injectable aqueous or oleaginous
30 suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents, which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for

example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including
5 synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[0171] The composition of the invention may also be administered in the form of suppositories, e.g., for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at
10 ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

[0172] Alternatively, the compositions can be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can
15 either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

[0173] For administration to non-human animals, the composition containing the therapeutic compound may be added to the animal's feed or drinking water. Also, it will be convenient to formulate animal feed and drinking water products so that the
20 animal takes in an appropriate quantity of the compound in its diet. It will further be convenient to present the compound in a composition as a premix for addition to the feed or drinking water. The composition can also added as a food or drink supplement for humans.

[0174] Dosage levels of the order of from about 5 mg to about 250 mg per
25 kilogram of body weight per day and more preferably from about 25 mg to about 150 mg per kilogram of body weight per day, are useful in the treatment of the above-indicated conditions. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the condition being treated and the particular mode of administration. Dosage unit forms
30 will generally contain between from about 1 mg to about 500 mg of an active ingredient.

[0023] Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most disorders, a dosage regimen of 4 times daily or less is preferred. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors
5 including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

[0024] Preferred compounds of the invention will have desirable pharmacological
10 properties that include, but are not limited to, oral bioavailability, low toxicity, low serum protein binding and desirable in vitro and in vivo half-lives. Penetration of the blood brain barrier for compounds used to treat CNS disorders is necessary, while low brain levels of compounds used to treat peripheral disorders are often preferred.

[0025] The amount of the composition required for use in treatment will vary not
15 only with the particular compound selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will ultimately be at the discretion of the attendant physician or clinician.

[0026] In an exemplary embodiment, the pharmaceutical composition described herein includes an additional active ingredient. In another exemplary embodiment,
20 the additional active ingredient is a compound that has been approved for human use by the United States Food and Drug Administration. In another exemplary embodiment, the additional active ingredient is an immunosuppressive agent. In still another exemplary embodiment, the additional active ingredient is selected from the group consisting of corticosteroids, aminosalicylates, azathioprine (6-
25 mercaptopurine), methotrexate and cyclosporine, etanercept, infliximab, adalimumab, alefacept, efalizumab and anakinra.

[0027] In still another exemplary embodiment, the additional active ingredient is selected from the group consisting of betamethasone, tacrolimus and pimecrolimus. In still another exemplary embodiment, the additional active ingredient is selected
30 from the group consisting of an activated vitamin D analog and an arotinoid (an aromatic retinoic acid analog). In still another exemplary embodiment, the additional active ingredient is cariprotinol, such as Tazorac® (tazarotene).

VII. a) Topical formulations

[0180] In a preferred embodiment, the methods of the invention can be employed through the topical application of the compounds described herein. Topical administration includes for example, transmucosal, transdermal, ungual and
5 transungual routes of administration. The topical compositions useful in the subject invention can be made into a wide variety of product types. These include, but are not limited to, lotions, creams, gels, sticks, sprays, ointments, pastes, foams, mousses, masks, eye ointments, eye or ear drops, impregnated dressings, wipes, cleansers including soaps, body washes and shampoos, and make-up products, such as bases,
10 blushes, lipsticks, and eye shadows, among others. These product types can comprise several types of carrier systems including, but not limited to particles, nanoparticles, and liposomes. If desired, disintegrating agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar or alginic acid or a salt thereof such as sodium alginate. Techniques for formulation and administration can be found in Remington: The
15 Science and Practice of Pharmacy, supra. The formulation can be selected to maximize delivery to a desired target site in the body. The formulations can also include various conventional colorants, fragrances, thickeners, preservatives, humectants, emollients, demulcents, solubilizing excipients, dispersants, penetration enhancers, plasticizing agents, preservatives, stabilizers, demulsifiers, wetting agents,
20 sunscreens, emulsifiers, moisturizers, astringents, deodorants, and the like, which can be added to provide additional benefits such as, for example, improving the feel and/or appearance of the topical preparation.

[0181] Lotions, which are preparations that are to be applied to the skin, nail, hair, claw or hoof surface without friction, are typically liquid or semi-liquid preparations
25 in which finely divided solid, waxy, or liquid are dispersed. Lotions will typically contain suspending agents to produce better dispersions as well as compounds useful for localizing and holding the active agent in contact with the skin, nail, hair, claw or hoof, e.g., methylcellulose, sodium carboxymethyl-cellulose, or the like.

[0182] Creams containing the active agent for delivery according to the invention
30 are viscous liquid or semisolid emulsions, either oil-in-water or water-in-oil. Cream bases are water-washable, and contain an oil phase, an emulsifier and an aqueous phase. The oil phase is generally comprised of petrolatum or a fatty alcohol, such as cetyl- or stearyl alcohol; the aqueous phase usually, although not necessarily, exceeds

the oil phase in volume, and generally contains a humectant. The emulsifier in a cream formulation, as explained in Remington: The Science and Practice of Pharmacy, supra, is generally a nonionic, anionic, cationic or amphoteric surfactant.

[0183] Ointments, which are semisolid preparations, are typically based on petrolatum or other petroleum derivatives. As will be appreciated by the ordinarily skilled artisan, the specific ointment base to be used is one that provides for optimum delivery for the active agent chosen for a given formulation, and, preferably, provides for other desired characteristics as well, e.g., emolliency or the like. As with other carriers or vehicles, an ointment base should be inert, stable, nonirritating and non-sensitizing. As explained in Remington: The Science and Practice of Pharmacy, 19th Ed. (Easton, Pa.: Mack Publishing Co., 1995), at pages 1399-1404, ointment bases may be grouped in four classes: oleaginous bases; emulsifiable bases; emulsion bases; and water-soluble bases. Oleaginous ointment bases include, for example, vegetable oils, fats obtained from animals, and semisolid hydrocarbons obtained from petroleum. Preferred water-soluble ointment bases are prepared from polyethylene glycols of varying molecular weight; again, reference may be had to Remington: The Science and Practice of Pharmacy, supra, for further information.


[0184] Useful formulations of the invention also encompass sprays and aerosols. Sprays generally provide the active agent in an aqueous and/or alcoholic solution which can be misted onto the skin, nail, hair, claw or hoof for delivery. Such sprays include those formulated to provide for concentration of the active agent solution at the site of administration following delivery, e.g., the spray solution can be primarily composed of alcohol or other like volatile liquid in which the drug or active agent can be dissolved. Upon delivery to the skin, nail, hair, claw or hoof, the carrier evaporates, leaving concentrated active agent at the site of administration. Examples of aerosol technology are disclosed in US Patents 6,682,716; 6,716,415; 6,716,417; 6,783,753; 7,029,658; and 7,033,575.

[0185] Examples of solubilizing excipients include polyethoxylated fatty acids, PEG-fatty acid diesters, PEG-fatty acid mono-ester and di-ester mixtures, polyethylene glycol glycerol fatty acid esters, alcohol-oil transesterification products, polyglycerized fatty acids, propylene glycol fatty acid esters, mixtures of propylene glycol esters-glycerol esters, mono- and diglycerides, sterol and sterol derivatives,

polyethylene glycol sorbitan fatty acid esters, polyethylene glycol alkyl ethers, sugar esters, polyethylene glycol alkyl phenols, polyoxyethylene-polyoxypropylene block copolymers, sorbitan fatty acid esters, lower alcohol fatty acid esters, ionic surfactants, tocopherol esters, and sterol esters.


5 [0186] Exemplary embodiments are summarized herein below.

[0187] In an exemplary embodiment, the invention provides a compound having a structure according to a formula which is:



10 wherein R^3 is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl; R^4 is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl; Y is O or S; and R^5 is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl; or a salt, hydrate or solvate thereof.

15 [0188] In an exemplary embodiment, according to the above paragraph, having a structure which is



wherein C^* is a carbon atom stereocenter which has a configuration which is (R) or (S).

20 [0189] In an exemplary embodiment, according to any of the above paragraphs, wherein C^* stereocenter is in a (S) configuration.

[0190] In an exemplary embodiment, according to any of the above paragraphs, wherein R^3 is $-(CR^{20}R^{21})_nNR^{22}R^{23}$ in which n is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; each R^{20} and each R^{21} is independently selected from the group consisting of H, R^{26} , OR^{26} , $NR^{26}R^{27}$, SR^{26} , $-S(O)R^{26}$, $-S(O)_2R^{26}$, $-S(O)_2NR^{26}R^{27}$, $-C(O)R^{27}$, $-C(O)OR^{27}$, and $-C(O)NR^{26}R^{27}$

25

R^{22} and R^{23} are independently selected from the group consisting of H, $-S(O)R^{28}$, $-S(O)_2R^{28}$, $-S(O)_2NR^{28}R^{29}$, $-C(O)R^{28}$, $-C(O)OR^{28}$, $-C(O)NR^{28}R^{29}$, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl; wherein each R^{26} , each R^{27} , each R^{28} and each R^{29} is independently selected from the group consisting of H, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

10 [0191] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-CH_2NH_2$.

[0192] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-CH_2NH_2$, and C^* has a configuration which is (S).

15 [0193] In an exemplary embodiment, according to any of the above paragraphs, R^4 is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, and *sec*-butyl.


[0194] In an exemplary embodiment, according to any of the above paragraphs, R^4 is selected from the group consisting of fluorine, chlorine, bromine, and iodine.

20 [0195] In an exemplary embodiment, according to any of the above paragraphs, R^4 is fluorine.

[0196] In an exemplary embodiment, according to any of the above paragraphs, R^4 is chlorine.

[0197] In an exemplary embodiment, according to any of the above paragraphs, R^4 is bromine.

25 [0198] In an exemplary embodiment, according to any of the above paragraphs, R^5 is:



wherein a is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; each R¹⁰ and each R¹¹ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, OH and NH₂;

R¹² is selected from the group consisting of H, R⁷, halogen, cyano, amidino, OR⁷,
 5 NR⁷R⁸, SR⁷, -N(R⁷)S(O)₂R⁸, -C(O)R⁷, -C(O)OR⁷, -C(O)NR⁷R⁸ wherein each R⁷ and each R⁸ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

10 [0199] In an exemplary embodiment, according to any of the above paragraphs, a is 1, 2, 3, 4, or 5.

[0200] In an exemplary embodiment, according to any of the above paragraphs, a is 2, 3, or 4.

[0201] In an exemplary embodiment, according to any of the above paragraphs,
 15 each R¹⁰ and each R¹¹ is independently selected from the group consisting of H, substituted or unsubstituted alkyl, OH, and NH₂.

[0202] In an exemplary embodiment, according to any of the above paragraphs, each R¹⁰ and each R¹¹ is H.

[0203] In an exemplary embodiment, according to any of the above paragraphs,
 20 R¹² is selected from the group consisting of H, OH, NH₂, methyl, ethyl, -NHS(O)₂CH₃, cyano, -NHC(O)CH₃, -NHC(O)NHCH₂CH₃, -C(O)NH₂, -C(O)OH, 4-(methoxy)phenyl, benzyl, benzoxy, -NHC(O)OCH₂Ph, -C(O)NHCH₂CH₂OH and -C(NH₂)(NH).

[0204] In an exemplary embodiment, according to any of the above paragraphs, Y
 25 is O.

[0205] In an exemplary embodiment, according to any of the above paragraphs, R⁵ is unsubstituted alkyl.

[0206] In an exemplary embodiment, according to any of the above paragraphs, R⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl,
 30 isobutyl, t-butyl, and sec-butyl.

[0207] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$; and Y is O; and R^5 is substituted or unsubstituted alkyl.

[0208] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$; and R^4 is halogen.

5 [0209] In an exemplary embodiment, according to any of the above paragraphs, R^4 is halogen; Y is O; and R^5 is unsubstituted alkyl.

[0210] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$, and C^* has a configuration which is (S) and R^4 is halogen.


10 [0211] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$, and C^* has a configuration which is (S), R^5 is unsubstituted alkyl and R^4 is halogen.

[0212] In an exemplary embodiment, according to any of the above paragraphs, Y is O, R^3 is $-\text{CH}_2\text{NH}_2$, and C^* has a configuration which is (S), R^5 is unsubstituted C_1 or C_2 or C_3 or C_4 alkyl and R^4 is halogen.

15 [0213] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$; R^4 is chlorine; Y is O; and R^5 is substituted or unsubstituted alkyl.

[0214] In an exemplary embodiment, according to any of the above paragraphs, R^3 is $-\text{CH}_2\text{NH}_2$; Y is O; and R^5 is ethyl.

20 [0215] In an exemplary embodiment, according to any of the above paragraphs, the compound has a structure which is



[0216] In an exemplary embodiment, the invention provides a composition comprising: (a) a first stereoisomer of the compound according to any of the above paragraphs; b) at least one additional stereoisomer of the first stereoisomer; wherein
25 the first stereoisomer is present in an enantiomeric excess of at least 80% relative to said at least one additional stereoisomer.

[0217] In an exemplary embodiment, according to any of the above paragraphs, wherein said enantiomeric excess is at least 92%.

[0218] In an exemplary embodiment, according to any of the above paragraphs, wherein the C* stereocenter of the first stereoisomer is in a (S) configuration.

[0219] In an exemplary embodiment, according to any of the above paragraphs, wherein R³ is -CH₂NH₂.

5 [0220] In an exemplary embodiment, the invention provides a composition according to any of the above paragraphs, wherein the C* stereocenter is in a (S) configuration, and said composition is substantially free of the (R) enantiomer of the compound.

[0221] In an exemplary embodiment, the invention provides a combination
10 comprising the compound according to any of the above paragraphs, or a pharmaceutically acceptable salt thereof, together with at least one other therapeutically active agent.

[0222] In an exemplary embodiment, the invention provides a pharmaceutical formulation comprising: (a) a compound according to any of the above paragraphs, or
15 a pharmaceutically acceptable salt thereof; and (b) a pharmaceutically acceptable excipient.

[0223] In an exemplary embodiment, according to any of the above paragraphs, the formulation is in a unit dosage form.

[0224] In an exemplary embodiment, according to any of the above paragraphs,
20 the formulation is for oral or topical use.

[0225] In an exemplary embodiment, the invention provides a method of inhibiting an enzyme, comprising: contacting the enzyme with the compound according to any of the above paragraphs, thereby inhibiting the enzyme.

[0226] In an exemplary embodiment, according to any of the above paragraphs,
25 the enzyme is a t-RNA synthetase which comprises an editing domain.

[0227] In an exemplary embodiment, according to any of the above paragraphs, the enzyme is a leucyl t-RNA synthetase.

[0228] In an exemplary embodiment, the invention provides a method of killing and/or preventing the growth of a microorganism, comprising: contacting the
30 microorganism with an effective amount of a compound according to any of the

above paragraphs, or a pharmaceutically acceptable salt thereof, thereby killing and/or preventing the growth of the microorganism.

[0229] In an exemplary embodiment, according to any of the above paragraphs, the microorganism is a bacterium.

5 [0230] In an exemplary embodiment, according to any of the above paragraphs, the microorganism is *Mycobacterium tuberculosis*.

[0231] In an exemplary embodiment, the invention provides a method of treating and/or preventing a disease in an animal, comprising: administering to the animal a therapeutically effective amount of compound according to any of the above
10 paragraphs, or a pharmaceutically-acceptable salt thereof, thereby treating and/or preventing the disease.

[0232] In an exemplary embodiment, according to any of the above paragraphs, the disease is tuberculosis.

[0233] In an exemplary embodiment, according to any of the above paragraphs,
15 the animal is a human.

[0234] In an exemplary embodiment, the invention provides a method of inhibiting the editing domain of a t-RNA synthetase, comprising: contacting the synthetase with an effective amount of a compound according to any of the above paragraphs, or a pharmaceutically-acceptable salt thereof, thereby inhibiting the
20 synthetase.

[0235] In an exemplary embodiment, according to any of the above paragraphs, the synthetase is a leucyl t-RNA synthetase.

[0236] In an exemplary embodiment, according to any of the above paragraphs, the synthetase is a *Mycobacterium tuberculosis* leucyl t-RNA synthetase.

25 [0237] In an exemplary embodiment, the invention provides the use of a compound according to any of the above paragraphs or a combination according to any of the above paragraphs or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment and/or prophylaxis of bacterial infection.

[0028] It is to be understood that the present invention covers all combinations of aspects and/or embodiments, as well as suitable, convenient and preferred groups described herein.

[0029] The invention is further illustrated by the Examples that follow. The

5 Examples are not intended to define or limit the scope of the invention.

EXAMPLES


[0030] Proton NMR are recorded on Varian AS 300 spectrometer and chemical shifts are reported as δ (ppm) down field from tetramethylsilane. Mass spectra are determined on Micromass® Quattro II.

10 [0031] The *M.tuberculosis* LeuRS gene (DNA sequence listed herein) was prepared by GenScript® and cloned into the T7 expression vector pET28a(+) at the NdeI-XhoI sites. Over-expression of *M. tuberculosis* LeuRS from this construct generated a version of *M. tuberculosis* LeuRS with an N-terminal his-tag, which will use the standard procedures for purification of his-tagged proteins.

15 EXAMPLE 1

A. 3-Aminomethyl-4-fluoro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol; bis trifluoroacetic salt


6-Fluoro-2,3-dimethoxy-benzaldehyde



20 [0032] To a solution of 4-fluoro-1,2-dimethoxy-benzene (20.0 g, 128.07 mmol) in anhydrous THF (200 mL) under nitrogen at -78 °C was added dropwise a 2.5M solution in hexane of n-BuLi (102.4 mL, 256.14 mmol) for duration 30 min and the reaction mixture was further stirred at the same temperature for 3 h. The reaction mixture was quenched carefully with DMF (100 mL) at -65 °C to -40 °C and left
25 overnight. 2N HCl (300 mL) was added dropwise at -60 °C and the mixture stirred for 30 min. The two layers were separated and the aqueous layer extracted with EtOAc. The combined organic layers were washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo. Purification was completed by flash column chromatography (20% EtOAc / hexane). Yield 18.0 g (85%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.39 (s, 1 H), 7.09 (dd, *J*=9.4, 5.1 Hz, 1 H), 6.84 (t, *J*=9.6
30


Hz, 1 H), 3.98 (s, 3 H), 3.88 (s, 3 H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*): -126 ppm.

6-Fluoro-2,3-dihydroxy-benzaldehyde




- 5 [0243] To a solution of 6-fluoro-2,3-dimethoxy-benzaldehyde (18.0 g, 97.74 mmol) in anhydrous dichloromethane (100 mL) under nitrogen at -60 °C was added dropwise $\text{BBr}_3 \cdot \text{OEt}_2$ (195 mL, 195.48 mmol) over 30 min and the solution was allowed to warm up to r.t and stirred for 4h. The reaction mixture was cooled to -60 °C and 2N HCl (250 mL) was carefully added drop wise. The mixture was stirred at
- 10 r.t overnight. The two layers were separated and the aqueous layer extracted with DCM. The combined organic layers were washed with water, sat. NaHCO_3 solution, water and brine and dried over MgSO_4 . The solvent was removed in vacuo to provide the title compound which was used in the next step without further purification. Yield 11.15 g (74%). ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 11.53 (s, 1 H), 10.23
- 15 (s, 1 H), 7.11 (dd, $J=8.8, 5.3$ Hz, 1 H), 6.57 (t, $J=9.6$ Hz, 1 H), 5.48 (s, 1 H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*): -132 ppm.

3-(3-Benzoyloxy-propoxy)-6-fluoro-2-hydroxy-benzaldehyde




- [0244] To a solution of 6-fluoro-2,3-dihydroxy-benzaldehyde (11.15 g, 71.42 mmol) in anhydrous DMSO (88 mL) under nitrogen were added sequentially sodium *t*-butoxide (13.72 g, 142.84 mmol) and benzyl-3-bromopropyl ether (17.29 g, 78.56 mmol) and the reaction mixture stirred at rt for 18 h. The mixture was diluted with water (400 mL) and extracted with EtOAc (4x100 mL). The combined organic layer was washed with water and brine, dried over MgSO_4 , filtered and concentrated in
- 25 vacuo. Purification was completed by flash chromatography (20% EtOAc / hexane). Yield 11.5 g (77 %). ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 11.62 (s, 1 H), 10.23 (s, 1 H), 7.42 - 7.16 (m, 5 H), 7.05 (dd, $J=8.8, 5.3$ Hz, 1 H), 6.54 (t, $J=9.6$ Hz, 1 H), 4.52 (s, 2 H), 4.13 (t, $J=6.3$ Hz, 2 H), 3.68 (t, $J=6.1$ Hz, 2 H), 2.12 (quin, $J=6.2$ Hz, 2 H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*): -132 ppm.

Trifluoro-methanesulfonic acid 6-(3-benzyloxy-propoxy)-3-fluoro-2-formyl-phenyl ester



[0245] Trifluoromethanesulfonic anhydride (1.11 g, 3.94 mmol) was added dropwise to a solution of pyridine (389 mg, 4.92 mmol) and 3-(3-benzyloxy-propoxy)-6-fluoro-2-hydroxy-benzaldehyde (1 g, 3.28 mmol) in CH₂Cl₂ (5 mL) at 0 °C (bath temp). The reaction mixture was then allowed to warm to rt and was stirred until complete consumption of starting material (as determined by TLC). Et₂O and 2 N HCl were then added. The organic layer was separated and washed with sat. NaHCO₃ then brine. The organic layer was dried (Na₂SO₄) and filtered through a short silica gel plug, washing with Et₂O. The filtrate was concentrated in vacuo to give 1.10 g of the desired triflate (yield 76%) that was used directly without further purification. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.32 (s, 1 H), 7.44 - 7.21 (m, 7 H), 4.51 (s, 2 H), 4.18 (t, *J*=6.1 Hz, 2 H), 3.68 (t, *J*=5.7 Hz, 2 H), 2.14 (quin, *J*=5.8 Hz, 2 H).


3-(3-Benzyloxy-propoxy)-6-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde



[0246] A solution of trifluoro-methanesulfonic acid 6-(3-benzyloxy-propoxy)-3-fluoro-2-formyl-phenyl ester (1.092 g, 2.50 mmol) in anhydrous 1,4-dioxane (10 mL) was added bis(pinacolato)diborane (953 mg, 3.75 mmol) and KOAc (736 mg, 7.50 mmol) at rt, then degassed with N₂ for 20 min. PdCl₂(dppf) (46 mg, 8 mol %) was added and the resulting solution was stirred at 100 °C until the reaction was complete. The solution was cooled to rt, filtered through Celite[®] or silica gel and concentrated in vacuo. The residue was taken up in EtOAc. The organic layer was then washed with H₂O then brine, dried (Na₂SO₄), filtered, and concentrated in vacuo. The product was purified by flash chromatography (20% EtOAc / hexane) to give 0.5 g of the title compound along with detriflated by product ratio ~1:1 by H NMR spectrum. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.33 (s, 1 H), 7.41 - 7.23 (m, 6 H), 7.19 (d, *J*=9.4 Hz, 1 H), 4.57 - 4.42 (m, 2 H), 4.06 (t, *J*=6.3 Hz, 2 H), 3.69 - 3.64 (m, 2 H),


2.20 - 2.00 (m, 2 H), 1.44 (s, 12 H); ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -73.72.

7-(3-Benzoyloxy-propoxy)-4-fluoro-3-nitromethyl-3H-benzo[C][1,2]oxaborol-1-ol



- 5 [0247] NaOH (48 mg, 1.20 mmol) was added to 3-(3-benzyloxy-propoxy)-6-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (500 mg, 1.2 mmol) in H_2O (3 mL) at rt, and the reaction mixture was stirred at rt for 5 min. MeNO_2 (219 mg, 3.6 mmol) was added dropwise and the mixture was stirred at rt for 16 h. The reaction mixture was acidified with 2 N HCl and extracted with EtOAc.
- 10 The organic fraction was washed with H_2O then brine, dried (MgSO_4), and concentrated in vacuo. Purification was accomplished by flash chromatography (10-40% EtOAc / hexane) to give 120 mg of the title compound by ^1H NMR spectrum. ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.40-7.22 (m, 5 H), 7.11 (t, $J=8.8$ Hz, 1 H), 6.81 (t, $J=8.6$ Hz, 1 H), 5.93 (dd, $J=9.0, 2.3$ Hz, 1 H), 4.99 (dd, $J=13.1, 2.5$ Hz, 1 H), 4.62 - 4.53 (m, 2 H), 4.43 (dd, $J=12.9, 9.0$ Hz, 1 H), 4.19-4.01 (m, 2 H), 3.66 (dt, $J=15.9, 5.7$ Hz, 2 H), 2.18 - 1.94 (m, 2 H); ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -72.81 (s, 1 F); MS (ESI) m/z = 374 (M - 1, negative).

3-Aminomethyl-4-fluoro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol; bis-trifluoroacetic acid salt




- 20 [0248] A mixture of 7-(3-benzyloxy-propoxy)-4-fluoro-3-nitromethyl-3H-benzo[C][1,2]oxaborol-1-ol (120 mg, 0.32 mmol) and 20% $\text{Pd}(\text{OH})_2$ (120 mg, 1:1 w/w substrate to catalyst) in AcOH (10 mL) was shaken under an atmosphere of H_2 (45-50 psi) in a Parr shaker. Once the reaction was complete, the mixture was filtered
- 25 through Celite[®]. The filtrate was concentrated in vacuo to give a gummy material. Remaining AcOH was removed by co-evaporation with toluene (3 $\times$) to give the amine. Purification by preparative HPLC (0.1% aq $\text{CF}_3\text{CO}_2\text{H}$ / CH_3CN) produced 12 mg of the title compound as a white solid (yield 8.4%). ^1H NMR (400 MHz, DMSO-*d*₆) δ ppm 9.14 (br. s., 1 H), 8.02 (br. s., 3 H), 7.25 (d, $J=7.8$ Hz, 1 H), 6.93 (d, $J=7.4$

Hz, 1 H), 5.43 (br. s., 1 H), 4.55 (br. s., 1 H), 4.07 (br. s., 2 H), 3.56 (br. s., 2 H), 3.42 – 3.37 (m, 1 H), 2.95 (br. s., 1 H), 1.86 (br. s., 2 H); ^{19}F NMR (376 MHz, DMSO- d_6) δ ppm -73.90 (s, 6 F), -131.51 (s, 1 F); MS (ESI) m/z = 256 ($M + 1$, positive); HPLC purity: 95.65% (MaxPlot 200 – 400 nm), 96.63% (220 nm).


5 **B. 3-(Aminomethyl)-4-chloro-7-(3-hydroxypropoxy)benzo[c][1,2]oxaborol-1(3H)-ol hydrochloride**

3-(3-Benzoyloxy-propoxy)-2-hydroxy-benzaldehyde



[0249] NaH (2.95 g, 72.4 mmol) was added to an ice-cold solution of 2,3-
 10 dihydroxybenzaldehyde (5.0 g, 36 mmol) in anhydrous DMSO (45 mL). Benzyl-3-bromopropyl ether (6.45 mL, 36.2 mmol) was then added and the mixture was stirred at rt for 12 h. The mixture was neutralized using 1 N HCl and then extracted with EtOAc. The organic fraction was washed with H₂O and concentrated in vacuo. The residue was purified by flash chromatography (8:2 hexane/EtOAc) to give the title
 15 compound as a brown oil: yield 8.40 g (81%). ^1H NMR (400 MHz, CDCl₃) δ (ppm): 9.93 (s, 1H), 7.36-7.23 (m, 6H), 7.20-7.16 (m, 2H), 6.98-6.91 (m, 1H), 4.53 (s, 2H), 4.19 (t, J = 6.2 Hz, 2H), 3.70 (t, J = 6.1 Hz, 2H), 2.19-2.16 (m, 2H).

3-(3-Benzoyloxy-propoxy)-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde




20 [0250] Trifluoromethanesulfonic anhydride (4.60 mL, 27.9 mmol) was added dropwise to a solution of pyridine (3.42 mL, 42.5 mmol) and 3-(3-benzyloxy-propoxy)-2-hydroxy-benzaldehyde (7.6 g, 26 mmol) in CH₂Cl₂ (200 mL) at 0 °C (bath temp). The reaction mixture was then allowed to warm to rt and was stirred
 25 until complete consumption of starting material (as determined by TLC). Et₂O and 2 N HCl were then added. The organic layer was separated and washed with sat. NaHCO₃ then brine. The organic layer was dried (Na₂SO₄) and filtered through a short silica gel plug, washing with Et₂O. The filtrate was concentrated in vacuo to give 8.60 g of the desired triflate (yield 77%) that was used directly without further

purification. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 10.23 (s, 1H), 7.54-7.47 (m, 1H), 7.43 (t, $J = 8.0$ Hz, 1H), 7.36-7.22 (m, 6H), 4.52 (s, 2H), 4.23 (t, $J = 6.3$ Hz, 2H), 3.71 (t, $J = 6.1$ Hz, 2H), 2.21-2.17 (m, 2H).

[0251] A solution of trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-6-formyl-phenyl ester (8.0 g, 19 mmol) in anhydrous 1,4-dioxane (160 mL) was added bis(pinacolato)diborane (9.71 g, 38.2 mmol) and KOAc (5.71 g, 57.4 mmol) at rt, then degassed with N_2 for 20 min. $\text{PdCl}_2(\text{dppf})$ (1.39 g, 1.89 mmol) was added and the resulting solution was stirred at 100 °C until the reaction was complete. The solution was cooled to rt, filtered through Celite[®] or silica gel and concentrated in vacuo. The residue was taken up in EtOAc. The organic layer was then washed with H_2O then brine, dried (Na_2SO_4), filtered, and concentrated in vacuo. The product was purified by flash chromatography (9:1 hexane/EtOAc) to give 4.80 g of the title compound (yield 43%) along with some pinacol contamination and was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 9.93 (s, 1H), 7.46 (t, $J = 7.8$ Hz, 1H), 7.41-7.36 (m, 1H), 7.35-7.24 (m, 5H), 7.08 (d, $J = 7.8$ Hz, 1H), 4.50 (s, 2H), 4.10 (t, $J = 6.3$ Hz, 2H), 3.67 (t, $J = 6.3$ Hz, 2H), 2.11 (quin, $J = 6.2$ Hz, 2H), 1.43 (s, 12H).


7-(3-Benzyloxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol



[0252] NaOH aq. (NaOH (3.64 g, 83 mmol) was added to 3-(3-benzyloxy-propoxy)-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (36 g, 91 mmol) in H_2O (180 mL), and THF (50 mL) at rt, and the reaction mixture was stirred at rt for 5 min. MeNO_2 (16.6 g, 273 mmol) was added dropwise and the mixture was stirred at rt for 16 h. The reaction mixture was acidified with 2 N HCl and extracted with EtOAc. The organic fraction was washed with H_2O then brine, dried (MgSO_4), and concentrated in vacuo. Purification was accomplished by flash chromatography (1:1 hexane/EtOAc) to give 15.9 g of the title compound as a light yellow oil (yield 50%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm: 9.05 (s, 1H), 7.44 (t, $J = 7.8$ Hz, 1H), 7.35-7.20 (m, 5H), 7.06 (d, $J = 7.4$ Hz, 1H), 6.88 (d, $J = 8.2$ Hz, 1H), 5.70 (dd, $J = 9.4, 2.3$ Hz, 1H), 5.29 (dd, $J = 13.7, 2.7$ Hz, 1H), 4.53 (dd, $J = 13.3, 9.4$ Hz, 1H), 4.45


(s, 2H), 4.11 (t, $J = 6.1$ Hz, 2H), 3.60 (t, $J = 6.3$ Hz, 2H), 2.04-1.91 (m, 2H); MS (ESI): $m/z = 356$ (M-1, negative); HPLC purity: 99.35% (MaxPlot 200-400 nm), 97.32% (220 nm).

4-Chloro-7-(3-hydroxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol



- 5 [0253] To 7-(3-benzyloxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (1.1 g, 3.0 mmol) in glacial AcOH (10 mL) in cold water bath was added SO_2Cl_2 (0.7 mL, 9.07 mmol) dropwise over 5 minutes period. The resulting solution was stirred for 30 minutes at the same temperature then 1.5 h at room temperature. The solution was quenched with crushed ice and then diluted with EtOAc (100 mL). The organic layer was washed with water, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. To the crude residue in MeOH (20 mL) was added $\text{Pd}(\text{OH})_2$ (10% w/w on carbon, 0.7 g), conc HCl was added until pH was 1, and the reaction vessel was pressurized to 40 psi with hydrogen for 30 minutes at room temperature.
- 10 The resulting mixture was filtered through a pad of Celite[®] and washed with EtOAc. The filtrate was concentrated in vacuo, then the residue was purified by silica gel column chromatography (EtOAc:Hex, 1:1) providing the title compound (0.2 g, 24% in 2 steps). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 9.31 (br. s., 1 H), 7.49 (d, $J=8.2$ Hz, 1 H), 6.98 (d, $J=8.6$ Hz, 1 H), 5.76 (dd, $J=8.2, 2.7$ Hz, 1 H), 5.33 (dd, $J=13.2, 2.3$ Hz, 1 H), 4.70 (dd, $J=13.0, 8.4$ Hz, 1 H), 4.55 (br. s., 1 H), 4.15 - 4.05 (m, 2 H), 3.61 - 3.55 (m, 2 H), 1.95 - 1.77 (m, 2 H).
- 20

3-Aminomethyl-4-chloro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol hydrogen chloride




- 25 [0254] 4-Chloro-7-(3-hydroxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (105 mg, 0.35 mmol) in methanolic ammonia solution (2 M, 20 mL) was added Ra/Ni (0.15 g, 2800 Nickel slurry in water) and the reaction

vessel was pressurized to 40 psi with hydrogen overnight at room temperature. The resultant mixture was filtered through a pad of Celite[®] and washed with EtOAc. The filtrate was concentrated in vacuo and the residue was added water (1 mL), followed by conc HCl to pH 1. The heterogeneous mixture was lyophilized provide the title
 5 compound (130 mg, quantitative) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 9.14 (br. s., 1 H), 8.36 (br. s., 3 H), 7.49 (d, *J*=8.2 Hz, 1 H), 6.99 (d, *J*=8.2 Hz, 1 H), 5.40 (d, *J*=7.0 Hz, 1 H), 4.40 (br. s., 1 H), 4.10 (br. s., 2 H), 3.59 (br. s., 2 H), ~3.30 (hidden, 1H), 2.89 (br. s., 1 H), 1.89 (br.s, 2 H); MS (ESI) *m/z* = 272 (M+1, positive); HPLC purity: 96.92% (MaxPlot 200 – 400 nm), 97.96% (220 nm).

10 C. **3-Aminomethyl-7-ethoxy-4-fluoro-3H-benzo[*c*][1,2]-oxaborol-1-ol; hydrochloride**

6-Fluoro-2,3-dimethoxy-benzaldehyde



[0255] To a cold (-78 °C) solution of 4-fluoro-1,2-dimethoxybenzene (15.00 g, 96.05 mmol) in anhydrous THF (150 mL) was added *n*-BuLi (84.5 mL, 211.32 mmol, 2.5 M solution in hexanes) under nitrogen and stirred it for 3 h at -78 °C. Quenched the reaction with DMF (75 mL) at -65 °C, added 2N HCl (300 mL) dropwise and further stirred for 30 min. Two layers separation was observed. Aqueous layer was extracted with EtOAc. Combined organic layers were washed with water, brine and dried over MgSO₄. Filtered the ethyl acetate layer and concentrated it *in vacuo*. The title compound was purified by flash column chromatography using pure hexanes then 10 and 20% EtOAc in hexanes which provided 14.40 g (78.19 mmol, 82%) of the title compound as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 10.23 (s, 1 H), 7.36 (dd, *J*=9.0, 5.1 Hz, 1 H), 7.03 (t, *J*=9.6 Hz, 1 H), 3.87 (s, 3 H), 3.83 (s, 3 H);
 25 ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -131.68 - -131.66 (m, 1F).


6-Fluoro-2,3-dihydroxy-benzaldehyde



[0256] To a cold (-78 °C) solution of 6-fluoro-2,3-dimethoxy-benzaldehyde (4.50 g, 24.43 mmol) in anhydrous dichloromethane (30 mL) was added BBr₃ (1M in DCM,


48.8 mL, 48.87 mmol) dropwise (duration 30 min). The reaction was warmed to room temperature and stirred for 4 h. Again cooled it to -78 °C and added 2N HCl (60 mL) to it dropwise. The reaction was stirred for overnight at room temperature and extracted with DCM. Combined organic layers were washed with water, sat. NaHCO₃ solution, brine and dried over MgSO₄. Filtration and removal of solvent provided 2.42 g (15.50 mmol, 64%) of the title compound as a yellow solid. This was used in the next step without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 10.22 (s, 1 H), 7.04 (dd, *J*=8.6, 5.5 Hz, 1 H), 6.61 (dd, *J*=10.4, 8.8 Hz, 1 H); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ ppm -131.69- -131.65 (m, 1F).

10 *3-Ethoxy-6-fluoro-2-hydroxy-benzaldehyde*



[0257] To a solution of 6-fluoro-2,3-dihydroxy-benzaldehyde (1.00 g, 6.40 mmol) in dry DMSO (10 mL) was added NaOBu^t (1.23 g, 12.81 mmol) and ethyl bromide (0.77 g, 7.04 mmol) under N₂ and stirred at RT for 18 h. The resultant mixture was diluted with water, acidified to pH ~6 with 2N.HCl and extracted with EtOAc (4x25 mL). Combined organic layers were washed with water and brine and dried over MgSO₄. Filtration and removal of the solvent under reduced pressure provided 1.05 g (5.70 mmol, 89 %) of the title compound as yellow solid. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.24 (s, 1 H), 7.04 (dd, *J*=8.8, 5.3 Hz, 1 H), 6.61 - 6.50 (m, 1 H), 4.09 (q, *J*=7.0 Hz, 3 H), 1.46 (t, *J*=7.0 Hz, 3 H); ¹⁹F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -132.19- -132.15 (m, 1F).


Trifluoro-methanesulfonic acid 6-ethoxy-3-fluoro-2-formyl-phenyl ester



[0258] To a cold (0 °C) solution of 3-ethoxy-6-fluoro-2-hydroxy-benzaldehyde (1.05 g, 5.70 mmol) in dry DCM (90 ml) was added pyridine (675 mg, 8.53 mmol) under nitrogen and stirred the reaction mixture at 0 °C for 10 min. Then added triflic anhydride (1.93 g, 6.84 mmol) slowly and continued stirring for 3 h at RT. Diluted the reaction with 1N.HCl (25 mL) and extracted with DCM (2x 100 mL). The organic layer was washed with water and brine and dried over MgSO₄. Filtered and

concentrated the filtrate. Purification of the residue by flash column chromatography with 5% ethyl acetate in hexanes gave 1.12 g, (3.54 mmol, 62%) of the title compound as a white solid. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.34 (s, 1 H), 7.31 - 7.13 (m, 2 H), 4.13 (q, *J*=7.0 Hz, 2 H), 1.48 (t, *J*=7.0 Hz, 3 H); ¹⁹F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -128.15- -128.11 (m, 1F), -73.56 (s, 3F).

3-Ethoxy-6-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2] dioxaborolan-2-yl)-benzaldehyde




[0259] A solution of trifluoro-methanesulfonic acid 6-ethoxy-3-fluoro-2-formyl-phenyl ester (1.60 g, 5.05 mmol) in anhydrous THF (50 mL) was degassed for 40 min. Added bis(pinacolato)diborane (3.85 g, 15.17 mmol), KOAc (1.50 g, 15.17 mmol) and PdCl₂(dppf) (296 mg, 8 mol %) and stirred the reaction at 70 °C (bath temp) for 3 h. Another addition of bis(pinacolato)diborane (1.40 g, 5.51 mmol) and heating at 70 °C for 2h completed the reaction. The resultant mixture was cooled to room temperature and filtered through a pad of Celite®. Concentrated the filtrate.

15 Purification of the residue by flash column chromatography with hexanes and 5% EtOAc/hexanes yielded 1.65 g of title compound as white solid. ¹H NMR confirms presence of title compound but with some impurities. It was used in next step without further purification. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.29 (s, 1 H), 7.12 - 6.97 (m, 2 H), 4.00 (q, *J*=7.0 Hz, 2 H), 1.46 (s, 12 H), 1.41 (t, *J*=6.8 Hz, 3 H);

20 ¹⁹F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -133.71- -133.67 (m, 1F).


7-Ethoxy-4-fluoro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol



[0260] 3-Ethoxy-6-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2] dioxaborolan-2-yl)-benzaldehyde (1.65 g, 5.61 mmol) was added to a solution of NaOH (225 mg, 5.60 mmol) in H₂O (10 mL) and stirred for 10 min at RT. Added nitromethane (1.03 g, 16.83 mmol) dropwise and stirred for 4 h at RT. The reaction mixture was acidified with 4N HCl and extracted with ethyl acetate. Organic layer was washed with water, brine and dried over MgSO₄, filtered and concentrated *in vacuo*. Purification of the


residue by flash column chromatography with 10 to 40% EtOAc/hexanes afforded 1.7 g of a mixture of compounds by ^1H NMR spectrum. This mixture was used in next step without further purification. ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.13 (t, $J=8.8$ Hz, 1 H), 6.79 (dd, $J=8.8$, 2.5 Hz, 1 H), 5.95 (d, $J=9.0$ Hz, 1 H), 5.14 (s, 1 H), 5.01 (dd, $J=13.3$, 2.3 Hz, 1 H), 4.44 (dd, $J=13.3$, 9.0 Hz, 1 H), 4.10 (q, $J=6.8$ Hz, 2 H), 1.45 (t, $J=6.8$ Hz, 3 H); ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -131.00- -130.97 (m, 1F).

7-Ethoxy-4-fluoro-1-hydroxy-1,3-dihydro-benzo[c][1,2] oxaborol-3-ylmethyl)-carbamic acid tert-butyl ester



[0033] To a cold (0 °C) solution of 7-ethoxy-4-fluoro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (500mg, 1.96 mmol) in dry MeOH (20 mL) was added (Boc)₂O (856 mg, 3.92 mmol) followed by NiCl₂ · 6H₂O (466 mg, 1.96 mmol) under nitrogen. Stirred the reaction mixture under nitrogen for 20 min and added NaBH₄ (445 mg, 11.76 mmol) in portions and left it for overnight at RT. Evaporated the solvent and diluted the reaction with 30 ml of ethyl acetate and filtered it through Celite®. Filtrate was concentrated and residue was purified by flash column chromatography using 5% MeOH/DCM, but a mixture (950 mg) of products was obtained which was used in next step without further purification.

3-Aminomethyl-7-ethoxy-4-fluoro-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride




[0034] A solution of 7-ethoxy-4-fluoro-1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaborol-3-ylmethyl)-carbamic acid tert-butyl ester (450 mg, 1.38 mmol) in 4M HCl (in 1,4-dioxane, 15 mL) was stirred at RT for overnight. The solvent was removed under reduced pressure. Recrystallization from EtOAc/hexanes provided 245 mg (0.93 mmol, 62%) of the title compound as white solid. ^1H NMR (400 MHz, DMSO-*d*₆) δ ppm 9.15 (br. s., 1 H), 8.20 (br. s., 3 H), 7.26 (t, $J=9.0$ Hz, 1 H), 6.95 - 6.85 (m, 1 H), 5.45 (d, $J=6.3$ Hz, 1 H), 3.30 (hidden, 1H), 4.05 (q, $J=6.9$ Hz, 2 H), 2.89 (br. s., 1 H), 1.30 (t, $J=6.8$ Hz, 3 H); ^{19}F NMR (376 MHz, DMSO-*d*₆) δ ppm -131.68- -

131.66 (m, 1F); MS (ESI) m/z = 226 (M+1, positive); HPLC purity: 91.87 % (MaxPlot 200 – 400 nm), 90.33% (220 nm); Anal. Calcd for $C_{10}H_{14}BClFNO_3 \cdot 0.5 H_2O$: C 44.40%; H 5.59%; N 5.18%. Found: C 44.30%; H 5.42%; N 5.50%.


D. 3-(Aminomethyl)-4-chloro-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol

5 **2-Bromo-3-hydroxybenzaldehyde**




[0263] The suspension of 3-hydroxybenzaldehyde (5 g, 0.04 mol), iron powder (172 mg, 3 mmol) and sodium acetate (6.72 g, 0.08 mol) in acetic acid (40 mL) was warmed until a clear solution was obtained and then cooled to room temperature. To this mixture was dropwise added a solution of bromine in glacial acetic acid (10 mL) over 15 min. After the addition, the reaction mixture was stirred for 2 h and then poured into ice-water. The resulting mixture was extracted with dichloromethane (3x50 mL). The combined extracts were dried over anhydrous Na_2SO_4 and concentrated. The residue was re-crystallized from dichloromethane to afford the product (2.3 g, yield 28%). ^1H NMR (400 MHz, $DMSO-d_6$) δ 10.30 (s, 1 H), 7.54-7.51 (m, 1H), 7.39-7.35 (m, 1H), 7.31-7.27 (m, 1H), 5.90 (s, 1H).


2-Bromo-3-ethoxybenzaldehyde




[0264] The suspension of 2-bromo-3-hydroxybenzaldehyde (120 g, 0.60 mol), K_2CO_3 (247 g, 1.79 mol) and bromoethane (135 mL, 1.79 mol) in DMF (700 mL) was stirred at 70 °C for 3 h. After the reaction was quenched with water (50 mL), the resulting mixture was extracted with EtOAc (3x60 mL). The combined organic layers were washed with water (50 mL) and aqueous LiCl solution (50 mL), dried over anhydrous Na_2SO_4 and concentrated to dryness in vacuo. The residue was purified by column chromatography on silica gel to give the target compound (128 g, yield 94%). ^1H NMR (400 MHz, $DMSO-d_6$) δ 10.45 (s, 1H), 7.52-7.50 (d, 1H), 7.38-7.34 (t, 1H), 7.13-7.10 (d, 1H), 4.18-4.13 (m, 2H), 1.53-1.50 (m, 3H).

2-(2-Bromo-3-ethoxyphenyl)-1,3-dioxolane

[0265] To a solution of 2-bromo-3-ethoxybenzaldehyde (128 g, 0.56 mol) and glycol (253 mL, 4.49 mol) in toluene (600 mL) was added *p*-toluenesulfonic acid (10 g, 0.06 mol). The reaction flask had a Dean and Stark condenser attached and the reaction mixture was refluxed to remove the water for 4 h. The reaction mixture was then cooled to room temperature and concentrated in vacuo. The residue was purified by column chromatography on silica gel to give the target compound (132 g, yield 86%).

10 *Diisopropyl 2-(1,3-dioxolan-2-yl)-6-ethoxyphenylboronate*


[0266] To the solution of 2-(2-bromo-3-ethoxyphenyl)-1,3-dioxolane (132 g, 0.48 mol) in anhydrous THF (500 mL) was dropwise added *n*-BuLi (2.5 M in THF, 386 mL, 0.97 mol) at -78 °C under nitrogen protection. The mixture was stirred at -78 °C for 2 h and then triisopropyl borate (227 mL, 0.97 mol) was dropwise added. The resulting mixture was stirred at this temperature for 4 h. After the reaction was quenched by adding saturated aqueous NH₄Cl solution (200 mL), the resulting mixture was extracted with EtOAc (3x300 mL). The combined organic layers were washed with water (200 mL) and brine (200 mL), dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by column chromatography on silica gel to give the target compound (136 g, yield 87%).

2-Ethoxy-6-formylphenylboronic acid

[0267] To the mixture of diisopropyl 2-(1,3-dioxolan-2-yl)-6-ethoxyphenylboronate (136 g, 0.42 mol) in THF (500 mL) was added diluted HCl (2N, 200 mL) slowly at room temperature with stirring. After stirred for 1.5 h at room


temperature, the reaction mixture was basified with 20% aqueous solution of NaOH to pH=12 and then washed with EtOAc (2x100 mL). The aqueous layer was acidified by using the diluted HCl (2N) to pH=2 and then extracted with EtOAc (3x100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by column chromatography on silica gel to give the target compound as white solid (80 g, yield 83%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.93 (s, 1H), 7.92 (s, 2H), 7.45-7.48 (m, 2H), 7.23-7.28 (d, 1H), 4.01-4.06 (m, 2H), 1.69-1.20 (m, 3H).

7-Ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol



[0268] The mixture of 2-ethoxy-6-formylphenylboronic acid (80 g, 0.41 mol), NaOH (16.5 g, 0.41 mol) and CTAB (7.7 g, 20 mmol) in H₂O (100 mL) and THF (500 mL) was stirred for 0.5 h at room temperature. After dropwise addition of nitromethane (14 mL, 2.4 mol), the reaction mixture was stirred at room temperature for 3 h. Then the cyclization was afforded by adding the diluted HCl (2 N) to pH=2 and then extracted with EtOAc (3x300 mL). The combined organic layers were washed with brine (250 mL), dried over anhydrous Na₂SO₄ and concentrated to dryness in vacuo. The residue was purified by column chromatography on silica gel to give the target compound as white solid (92 g, yield 94%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.06 (s, 1H), 7.46-7.43 (t, 1H), 7.07-7.05 (d, 1H), 6.89-6.87 (d, 1H), 5.71-5.69 (m, 1H), 5.31-5.27 (m, 1H), 4.57-4.51 (m, 1H), 4.12-4.07 (m, 2H), 1.34-1.30 (t, 3H).


4-Chloro-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol



[0269] To a solution of 7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (42 g, 0.18 mol) in DMF (200 mL) at 80 °C was added a solution of NCS (11.8 g, 0.18 mol) in DMF (50 mL) in 30 min. The reaction was quenched with an aqueous solution of LiCl solution (500 mL) and the resulting mixture was extracted by EtOAc

(3x250 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel to give the compound as white solid. (39.7 g, contaminated with 18% C6-Cl regioisomer). The mixture was then re-crystallized from Ether/PE (1/5) to give the pure compound (28 g, yield 46.3%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.32 (s, 1H), 7.50-7.48 (d, 1H), 6.98-6.96 (d, 1H), 5.77-5.74 (d, 1H), 5.35-5.31 (d, 1H), 4.73-4.67 (m, 1H), 4.12-4.07 (m, 2H), 1.34-1.28 (t, 3H).


3-(Aminomethyl)-4-chloro-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride



[0035] A mixture of 4-chloro-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (47 g, 0.17 mol), Raney Ni (2 g) and 2 M NH₃ in EtOH (40 mL) in EtOH (200 mL) was stirred under an atmosphere of H₂ for 2 h and then filtrated. The filtrate was acidified by using 4.5 N HCl in EtOH (100 mL). After stirring for 30 min, the mixture was concentrated and the residue was washed with CH₃CN (2x50 mL) to give the product as white solid (43 g, yield 89%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.13 (s, 1 H), 8.18 (s, 3H), 7.50-7.51 (d, 1H), 6.97-7.00 (d, 1H), 5.36-5.39 (m, 1H), 4.08-4.14 (m, 2H), 3.55-3.59 (m, 1H), 2.90-2.95 (m, 1H), 1.33-1.36 (m, 3H); MS (ESI) *m/z* = 242 [M + H]⁺.

E. 3-(Aminomethyl)-4-bromo-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol 2,2,2-trifluoroacetate salt


3-(Aminomethyl)-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride salt



[0036] To the solution of 7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (2 g, 8.43 mmol), Raney Ni (200 mg) and 2 M NH₃ in EtOH (10 mL) in ethanol (35 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The crude amine was dissolved in EtOAc (10 mL) and HCl in Et₂O (30 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was


washed with acetonitrile/hexanes (2:1, 2x20 mL) to give the compound as white solid (1 g, yield 57.2%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.89 (s, 1H), 8.22 (s, 3H), 7.48-7.44 (t, 1H), 7.06-7.04 (d, 1H), 6.90-6.88 (d, 1H), 5.31-5.29 (m, 1H), 4.13-4.08 (m, 2H), 3.45-3.39 (m, 1H), 2.80-2.78 (m, 1H), 1.36-1.33 (m, 3H); MS (ESI) *m/z* = 208 [M + H]⁺.

tert-Butyl (7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methylcarbamate




[0272] To the mixture of 3-(aminomethyl)-7-ethoxybenzo[*c*][1,2]oxaborol-1(3H)-ol hydrochloride salt (300 mg, 1.23 mmol) and triethylamine (622 mg, 6.16 mmol) in dichloromethane (35 mL) at 0 °C was added di-*tert*-butyl dicarbonate (402.8 mg, 1.85 mmol) and the mixture was stirred for 2 h at room temperature. After the reaction was quenched with sat. NaHCO₃ (45 mL) and the resulting mixture was extracted with EtOAc (3x30 mL), the combined organic layers were dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by flash-column chromatography to give the product (320 mg, yield 84.6%).

tert-Butyl (4-bromo-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)-methylcarbamate



[0273] To the solution of *tert*-butyl (7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methylcarbamate (250 mg, 0.81 mmol) and 1-bromopyrrolidine-2,5-dione (173.9 mg, 0.98 mmol) in CH₃CN (50 mL) was added 2,2'-Azobis(2-methylpropionitrile) (10 mg) and the mixture was stirred for 1 h at 90 °C. The reaction mixture was then concentrated in high vacuo and the residue was purified by prep-HPLC to give the product (200 mg, yield 63.6%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.90 (s, 1H), 7.55-7.53 (d, 1H), 6.85-6.82 (d, 1H), 5.08-5.07 (d, 1H), 4.11-4.07 (m, 2H), 3.82-3.79 (d, 1H), 3.06-3.03 (m, 1H), 1.39 (s, 9H), 1.30 (t, 3H); MS (ESI) *m/z* = 387 [M + H]⁺.

3-(Aminomethyl)-4-bromo-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol 2,2,2-trifluoroacetate salt




[0274] The mixture of *tert*-butyl (4-bromo-7-ethoxy-1-hydroxy-1,3-

5 dihydrobenzo[c][1,2]oxaborol-3-yl)-methylcarbamate (200 mg, 51.8 mmol) in 2,2,2-trifluoroacetic acid and dichloromethane (1:1, 20 mL) was stirred at room temperature for 1 h and concentrated to dryness (water bath < 30 °C). The residue was washed with acetonitrile (2x5 mL) and the white solid was dried in high vacuo to give the product (190 mg, yield 91.6%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.12 (s, 1H), 8.04 (s, 3H), 7.65-7.62 (d, 1H), 6.94-6.92 (d, 1H), 5.27-5.25 (m, 1H), 4.13-4.08 (m, 2H), 3.64-3.61 (m, 1H), 2.99-2.92 (m, 1H), 1.36-1.33 (t, 3H); MS (ESI) *m/z* = 287 [M + H]⁺.

F. *3-(Aminomethyl)-7-ethoxy-4-methylbenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride salt*

15 *7-Ethoxy-4-methyl-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol*




[0275] A mixture of 4-bromo-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-

1(3H)-ol (200 mg, 0.63 mmol), tetramethylstannane (341.7 mg, 1.90 mmol) and

Pd(PPh₃)₄ (Cat. 20 mg) in DMF (35 mL) was stirred overnight at 90 °C under N₂

20 protection. The reaction was quenched by adding ice-water and the resulting mixture was extracted with ethyl acetate (3x30 mL). The combined extracts were dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by prep-TLC to give the product (72 mg, yield 45.3%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 7.23-7.21 (d, 1H), 6.83-6.81 (d, 1H), 5.77-5.75 (m, 1H), 5.27-5.24 (m, 1H), 4.50-4.44 (m, 1H), 4.08-4.03 (m, 2H), 2.25 (s, 3H), 1.33-1.29 (t, 3H).


3-(Aminomethyl)-7-ethoxy-4-methylbenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride salt



[0037] A mixture of 7-ethoxy-4-methyl-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (80 mg, 0.32 mmol), Raney Ni (50 mg) and NH₃/EtOH (2 mL) in EtOH (10 mL) was stirred under an atmosphere of H₂ for 2 h and then filtrated. The filtrate was acidified by using 4.5 N HCl in EtOH (15 mL). After stirring for 30 min, the mixture was concentrated in vacuo and the residue was washed with CH₃CN (2x3 mL) to give the product as white solid (39 mg, yield 47.5%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.15 (s, 3H), 7.24-7.22 (d, 1H), 6.83-6.81 (d, 1H), 5.37-5.35 (m, 1H), 4.08-4.03 (m, 2H), 3.36-3.28 (m, 1H), 2.73-2.70 (m, 1H), 2.23 (s, 3H), 1.34-1.30 (t, 3H); MS (ESI) *m/z* = 222 [M + H]⁺.


G. 3-(Aminomethyl)-7-ethoxy-4-ethylbenzo[c][1,2]oxaborol-1(3H)-ol

7-Ethoxy-3-(nitromethyl)-4-vinylbenzo[c][1,2]oxaborol-1(3H)-ol




[0038] A mixture of 4-bromo-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (900 mg, 2.85mmol), vinyltributyltin (5.2 g, 53 mmol) and Pd(Ph₃P)₄ (230 mg, 0.2 mmol) in DMF (45 mL) was degassed for 15 min with N₂ and then stirred at 100 °C for 30 min in microwave reactor (Biotage®). After the reaction was quenched with ice-water, the resulting mixture was extracted with EtOAc (3x30 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and then concentrated to dryness in vacuo. The residue was purified by column chromatography on silica gel to give the compound (650 mg, yield 87%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.10 (s, 1 H), 7.64-7.66 (d, 1H), 6.93-6.95 (d, 1H), 6.77-6.84 (m, 1H), 5.93-5.96 (d, 1H), 5.69-5.73 (d, 1H), 5.28-5.31 (d, 1H), 5.10-5.14 (d, 1H), 4.44-4.49 (m, 1H), 4.09-4.14 (m, 2H), 1.32-1.35 (m, 3H); MS (ESI) *m/z* = 264 [M + H]⁺.

3-(Aminomethyl)-7-ethoxy-4-vinylbenzo[c][1,2]oxaborol-1(3H)-ol



[0039] A mixture of 7-ethoxy-3-(nitromethyl)-4-vinylbenzo[c][1,2]oxaborol-1(3H)-ol (205 mg, 0.78 mmol), Raney-Ni (50 mg) and 2 M NH₃ in EtOH (5 mL) in EtOH (10 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite and the filtrate was concentrated in vacuo. The crude amine was dissolved in EtOAc (2 mL) and HCl in Et₂O (20 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was used directly for the next step without further purification.


10 *3-(Aminomethyl)-7-ethoxy-4-ethylbenzo[c][1,2]oxaborol-1(3H)-ol*



[0040] To a suspension of 3-(aminomethyl)-7-ethoxy-4-vinylbenzo[c][1,2]oxaborol-1(3H)-ol (175 mg, 0.75 mmol) with Pd/C (40 mg) in EtOH (5 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The crude amine was dissolved in EtOAc (2 mL) and HCl in Et₂O (15 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was washed with hexanes to give the target compound (23 mg, yield 13%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (s, 1 H), 8.18 (s, 3H), 7.31-7.29 (d, 1H), 6.68-6.88 (d, 1H), 5.38-5.40 (d, 1H), 4.04-4.09 (d, 2H), 3.30-3.35 (m, 1H), 2.66-2.71 (m, 1H), 1.31-1.34 (m, 3H), 1.15-1.17 (m, 3H); MS (ESI) *m/z* = 236 [M + H]⁺.


H. 3-(Aminomethyl)-7-ethoxy-4-phenylbenzo[c][1,2]oxaborol-1(3H)-ol

7-Ethoxy-3-(nitromethyl)-4-phenylbenzo[c][1,2]oxaborol-1(3H)-ol



[0041] A mixture of 4-bromo-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol (315 mg, 1 mmol), tributyl-phenyl-stannane (750 mg, 2 mmol) and Pd(Ph₃P)₄ (Cat.) in DMF (15 mL) was degassed for 15 min with N₂ and then stirred at 100 °C for 30 min in microwave reactor (Biotage®). After the reaction was quenched with ice-water, the resulting mixture was extracted with EtOAc (3x40 mL). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and then concentrated to dryness. The residue was purified by column chromatography on silica gel to give the compound as white solid (60 mg, yield 20%). ¹HNMR (400 MHz, DMSO-*d*₆) δ 9.13 (s, 1 H), 7.46-7.49 (m, 5H), 7.34-7.48 (m, 2H), 6.17-6.20 (m, 1H), 4.88-4.92 (m, 1H), 4.21-4.25 (m, 1H), 4.05-4.16 (m, 2H), 1.34-1.37 (m, 3H).


15 3-(Aminomethyl)-7-ethoxy-4-phenylbenzo[c][1,2]oxaborol-1(3H)-ol



[0042] A mixture of 7-ethoxy-3-(nitromethyl)-4-phenylbenzo[c][1,2]oxaborol-1(3H)-ol (60 mg, 0.19 mmol), Raney-Ni (~25 mg) and 2 M NH₃ in EtOH (2 mL) in EtOH (10 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The crude amine was dissolved in EtOAc (1 mL) and HCl in Et₂O (5 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was washed with hexanes to give the compound as white solid (30 mg, yield 51%). ¹HNMR (400 MHz, DMSO-*d*₆) δ 8.89 (s, 1 H), 8.04 (s, 3H), 7.43-7.46 (d, 5H), 7.36-7.38 (d, 1H), 6.98-7.00 (d, 1H), 5.78-5.81 (d, 1H), 4.09-4.14 (m, 2H), 2.56-2.59 (m, 1H), 2.24-2.30 (m, 1H), 1.33-1.36 (m, 3H); MS (ESI) *m/z* = 284 [M + H]⁺.


I. 7-(4-Aminobutoxy)-3-(aminomethyl)-4-chlorobenzo[c][1,2]oxaborol-1(3H)-ol dihydrochloride

tert-Butyl 4-hydroxybutylcarbamate




- 5 [0282] To a mixture of 4-aminobutan-1-ol (4.0 g, 45 mmol) and TEA (7.5 mL, 54 mmol) in DCM (200 mL) was added (Boc)₂O (10.2 g, 47.2 mmol). The reaction mixture was stirred for 2 h at room temperature and then washed with water (2x150 mL) and the solution of citric acid (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the product as yellow oil 7.5 g. (yield 88%).

4-(tert-Butoxycarbonyl)butyl methanesulfonate




- 15 [0283] To a mixture of *tert*-butyl 4-hydroxybutylcarbamate (7.5 g, 40 mmol) and TEA (3.6 mL, 48 mmol) in DCM (100 mL) at 0 °C was dropwise added MsCl (6.6 mL, 48 mmol). The mixture was stirred at room temperature for 1 h and then washed with water (2x100 mL) and the solution of citric acid (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the product as yellow oil (10.0 g, yield 94%).

tert-Butyl 4-(2-bromo-3-formylphenoxy)butylcarbamate




- 20 [0284] To a mixture of 2-bromo-3-hydroxybenzaldehyde (3.0 g, 15 mmol) and *tert*-butoxycarbonyl)butyl methanesulfonate (4.8 g, 18 mmol) in DMF (40 mL) was added K₂CO₃ (6.2 g, 45 mmol). The mixture was stirred at 80 °C for 45 min and quenched by addition of aqueous LiCl solution (80 mL). The mixture was extracted with EtOAc (2x80 mL) and the combined organic layers were dried over anhydrous Na₂SO₄ and then concentrated to give the crude product as brown oil (6.0 g).

tert-Butyl 4-(3-formyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)-butylcarbamate



[0285] A mixture of *tert*-butyl 4-(2-bromo-3-formylphenoxy)butylcarbamate (6.0 g, 16 mmol), KOAc (5.0 g, 48 mmol), (Pin)₂B₂ (7.7 g, 86 mmol) and Pb(dppf)Cl₂ (1.25 g, 1.6 mmol) in dioxane (100 mL) was degassed for 15 min with N₂ and refluxed for 2 h under N₂ protection. The mixture was filtered and the filtrate was concentrated in vacuo. The residue was purified by chromatography on silica gel to give the product as yellow oil (3.5 g, yield 55%).


10 *tert*-Butyl-4-(1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[*c*][1,2]oxa-borol-7-yloxy)butylcarbamate



[0286] To a solution of *tert*-butyl 4-(3-formyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenoxy)-butylcarbamate (3.5 g, 8.3 mmol) and CTAB (cat.) in THF (50 mL) was added MeNO₂ (2.8 mL, 49 mmol), followed by an aqueous solution of NaOH (0.36 g, 9.1 mmol) in H₂O (5 mL). The mixture was stirred at room temperature for 45 min. The cyclization was afforded by adding 2N HCl solution until pH=2 at 0 °C. The reaction mixture was extracted with EtOAc (3x50 mL) and the organic layers were dried over anhydrous Na₂SO₄, and then concentrated in vacuo.


20 The residue was purified by chromatography on silica gel (PE: EtOAc=3:1) to give the product as yellow oil (1.7 g, yield 53.6%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (s, 1H), 7.44-7.47 (t, 1H), 7.06-7.08 (d, 1H), 6.88-6.90 (d, 1H), 9.86 (t, 1H), 5.70-5.72 (m, 1H), 5.29-5.33 (m, 1H), 4.53-4.59 (m, 1H), 4.02-4.06 (t, 2H), 2.95-2.30 (m, 2H), 1.67-1.72 (m, 2H), 1.52-1.57 (m, 2H), 1.38 (s, 9H).

tert-Butyl-4-(4-chloro-1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[c][1,2]oxaborol-7-yloxy)butylcarbamate




[0043] A mixture of *tert*-butyl 4-(1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[c][1,2]oxaborol-7-yloxy)-butylcarbamate (640 mg, 1.7 mmol) in DMF (20 mL) was added NCS (226 mg, 1.7 mmol) in DMF (5 mL). The mixture was heated to 80 °C for 2 h. After the reaction was quenched with an aqueous LiCl solution (100 mL), the resulting mixture was extracted with EtOAc (3x50 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by prep-HPLC to give the product (280 mg, yield 67.5%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.29 (s, 1H), 7.47-7.49 (d, 1H), 6.96-6.98 (d, 1H), 6.80 (s, 1H), 5.74-5.77 (m, 1H), 5.31-5.35 (m, 1H), 4.67-4.72 (m, 1H), 4.02-4.05 (m, 2H), 2.94-2.99 (m, 2H), 1.68-1.72 (m, 2H), 1.50-1.56 (m, 2H), 1.36 (s, 9H).

tert-Butyl 4-(3-(aminomethyl)-4-chloro-1-hydroxy-1,3-dihydro-benzo[c][1,2]oxaborol-7-yloxy)butylcarbamate



[0044] A mixture of *tert*-butyl 4-(1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[c][1,2]oxaborol-7-yl-oxy)-butylcarbamate (410 mg, 1 mmol), Raney-Ni (100 mg) and 2 N NH₃ in EtOH (3 mL) in EtOH (15 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The resulting solid was used directly for the next step.

7-(4-Aminobutoxy)-3-(aminomethyl)-4-chlorobenzo[c][1,2]oxaborol-1(3H)-ol dihydrochloride




[0045] To a mixture of the crude *tert*-butyl-4-(3-(aminomethyl)-4-chloro-1-hydroxy-1,3-dihydrobenzo[c]-[1,2]-oxa-borol-7-yloxy)butylcarbamate in DCM (5

mL) was added CF₃COOH (2 mL) at room temperature. The reaction mixture was stirred for 1 h and concentrated in vacuo. The crude amine was dissolved in EtOAc (1 mL) and HCl in Et₂O (10 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was washed with hexanes to give the target

compound as white solid (180 mg, yield: 42%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.16 (s, 1H), 8.28 (s, 3H), 8.03 (d, 3H), 7.49-7.51 (d, 1H), 6.99-7.01 (d, 1H), 5.38-5.40 (m, 1H), 4.05-4.08 (m, 2H), 3.56-3.59 (d, 1H), 2.84-2.91 (m, 3H), 1.71-1.83 (m, 4H); MS (ESI) *m/z* = 285 [M + H]⁺.


J. 3-(Aminomethyl)-7-(3-aminopropoxy)-4-chlorobenzo[c][1,2]oxaborol-1(3H)-ol

tert-Butyl 3-bromopropylcarbamate




[0290] To a mixture of 3-bromopropan-1-amine (10.95 g, 50 mmol) and TEA (15.4 mL, 110 mmol) in DCM (100 mL) at 0 °C was added (Boc)₂O (11.4 g, 52.5 mmol). The reaction mixture was stirred at room temperature overnight and then washed with water (3x100 mL) and the solution of citric acid (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated to give the product as yellow oil (9.0 g, yield 76%).

tert-Butyl 3-(2-bromo-3-formylphenoxy)propylcarbamate




[0291] A mixture of 2-bromo-3-hydroxybenzaldehyde (5 g, 24.9 mmol), 3-(*tert*-butoxy carbonylamino)-propyl methanesulfonate (7.55 g, 30 mmol) and Cs₂CO₃ (24 g, 75 mmol) in DMF (60 mL) was stirred at 50 °C for 3 h and quenched with water (600 mL). The resulting mixture was extracted with EtOAc (3x60 mL) and the combined organic layers were washed with brine (60 mL), dried over anhydrous Na₂SO₄ and then concentrated in vacuo. The residue was purified by column chromatography on silica gel (PE/EtOAc=10/1) to give the product (7.2 g, yield 81%). ¹H NMR (400 MHz, CDCl₃) δ 10.43 (s, 1H), 7.53 (dd, *J*=7.8 Hz, 1.6 Hz, 1H), 7.37 (t, *J*=7.8 Hz, 1H), 7.12 (dd, *J*=8.2 Hz, 1.6 Hz, 1H), 5.16 (s, 1H), 4.15 (t, *J*=5.9 Hz, 2H), 3.42 (m, 2H), 2.10 (m, 2H), 1.44 (s, 9H); MS (ESI) *m/z* = 358 [M+H]⁺.

tert-Butyl 3-(3-formyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-phenoxy)propylcarbamate




[0292] A solution of *tert*-butyl 3-(2-bromo-3-formylphenoxy)propylcarbamate (7.1 g, 20 mmol), B₂pin₂ (10 g, 40 mmol), Pd(dppf)Cl₂ (800 mg, 2 mmol) and KOAc (5.9 g, 60 mmol) in 1,4-dioxane (30 mL) was degassed with N₂ and stirred at 80 °C for 5 h. The mixture was cooled to room temperature and diluted with EtOAc (100 mL). The organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and then concentrated. The residue was purified by chromatography on silica gel (PE/ EtOAc =5/1) to give the product (3.1 g, yield 38.3%). ¹H NMR (400 MHz, CDCl₃) δ 9.95 (s, 1H), 7.48 (t, *J*=7.8 Hz, 1H), 7.40 (m, 1H), 7.09 (d, *J*=8.2 Hz, 1H), 4.76 (s, 1H), 4.05 (t, *J*=6.3 Hz, 2H), 3.32 (m, 2H), 2.00 (m, 2H), 1.45 (s, 12H), 1.43 (s, 9H); MS (ESI) *m/z* =406 [M+H]⁺.

tert-Butyl 3-(1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[*c*][1,2]oxaborol-7-yloxy)-propylcarbamate




[0293] A mixture of *tert*-butyl 3-(3-formyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) phenoxy)propylcarbamate (3.1 g, 8.47 mmol), MeNO₂ (775 mg, 12.7 mmol), CTAB (310 mg, 0.85 mmol) and NaOH (407 mg, 10 mmol) in THF (35 mL) and H₂O (8 mL) was stirred at room temperature for 3 h. The mixture was adjusted to pH 2-3 using 2N HCl and then stirred for 30 min. The mixture was extracted with EtOAc (2x80 mL). The organic layer was washed with water (30 mL) and brine (30 mL), dried over anhydrous Na₂SO₄ and then concentrated. The residue was purified by column chromatography on silica gel (PE/ EtOAc =5/1) to give the product (2 g, yield 64.5%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (s, 1H), 7.47 (t, *J*=7.8 Hz, 1H), 7.07 (d, *J*=7.4 Hz, 1H), 6.91 (m, 2H), 5.72 (dd, *J*=9.0, 2.7 Hz, 1H), 5.31 (dd, *J*=13.3, 2.7 Hz, 1H), 4.54 (dd, *J*=13.3, 9.4 Hz, 1H), 4.05 (t, *J*=6.3 Hz, 2H), 3.09 (m, 2H), 1.83 (m, 2H), 1.37 (s, 9H); MS (ESI) *m/z* =367 [M+H]⁺.

tert-Butyl-4-(4-chloro-1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[*c*][1,2]oxaborol-7-yloxy) propylcarbamate




[0046] To a mixture of *tert*-butyl 4-(1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[*c*][1,2]oxaborol-7-yl-oxy)propylcarbamate (2.9 g, 8 mmol) in DMF (35 mL) was added NCS (1.0 g, 8 mmol) in DMF (15 mL). The reaction mixture was heated to 80 °C for 2 h and then quenched with an aqueous LiCl solution (300 mL). The resulting mixture was extracted with EtOAc (3x100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by prep-HPLC to give the product as white solid (480 mg, yield 15.2%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.28 (s, 1H), 7.49-7.51 (d, 1H), 6.96-6.98 (d, 1H), 6.85 (s, 1H), 5.74-5.77 (m, 1H), 5.31-5.35 (m, 1H), 4.67-4.72 (m, 1H), 4.03-4.06 (m, 2H), 3.07-3.11 (m, 2H), 1.82-1.86 (m, 2H), 1.37 (s, 9H); MS (ESI) *m/z* = 401 [M + H]⁺.

tert-Butyl 3-(3-(aminomethyl)-4-chloro-1-hydroxy-1,3-dihydro-benzo[*c*][1,2]oxaborol-7-yloxy)propylcarbamate



[0047] A mixture of *tert*-butyl 3-(4-chloro-1-hydroxy-3-(nitromethyl)-1,3-dihydrobenzo[*c*][1,2]oxaborol-7-yloxy)propylcarbamate (480 mg, 1.2 mmol), Raney-Ni (500 mg) and 2 M NH₃ in EtOH (3 mL) in EtOH (15 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The resulting solid was used directly for the next step.

*3-(Aminomethyl)-7-(3-aminopropoxy)-4-chlorobenzo[*c*][1,2]oxaborol-1(3H)-ol*




[0048] To a mixture of the crude *tert*-butyl 3-(3-(aminomethyl)-4-chloro-1-hydroxy-1,3-dihydrobenzo[*c*]-[1,2]oxaborol-7-yloxy)propylcarbamate in DCM (10 mL) was added CF₃COOH (2.0 mL) at 0 °C. The reaction mixture was stirred for 1 h


and concentrated in vacuo. The crude amine was dissolved in EtOH (2 mL) and HCl in Et₂O (2 mL) was added immediately. After 1 h, the mixture was concentrated in vacuo. The residue was re-crystallized by EtOH/Et₂O to give the target compound (173.4 mg, yield 42%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.28 (s, 1H), 8.36 (s, 3H), 8.17 (s, 3H), 7.50-7.52 (d, 1H), 6.98-7.00 (d, 1H), 5.39-5.42 (m, 1H), 4.13-4.16 (m, 2H), 3.56-3.59 (d, 1H), 2.98-2.99 (m, 2H), 2.87 (s, 1H), 2.04-2.10 (m, 2H); MS (ESI) *m/z* = 271 [M + H]⁺.

K. *(R)*-3-(Aminomethyl)-4-chloro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3H)-ol hydrochloride and

10 L. *(S)*-3-(Aminomethyl)-4-chloro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3H)-ol hydrochloride




tert-Butyl ((4-chloro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate



15 [0297] A solution of 3-(aminomethyl)-4-chloro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3H)-ol hydrochloride (38.4 g, 0.16 mol) and Et₃N (47.8 g, 0.47 mol) in CH₂Cl₂ (350 mL) at 0 °C was added di-*tert*-butyl dicarbonate (172 g, 0.79 mol) and the reaction was stirred for 2 h at room temperature. After the reaction was quenched by addition of sat. NaHCO₃ (100 mL) and the resulting mixture was extracted with EtOAc (3x120 mL). The combined organic layers were dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica gel to give the


compound as white solid (27 g, yield 50%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.92 (s, 1 H), 7.40-7.42 (d, 1H), 6.88-6.90 (d, 1H), 6.77-6.79 (m, 1H), 5.15-5.16 (d, 1H), 4.06-4.13 (m, 2H), 3.75-3.78 (d, 1H), 3.03-3.08 (m, 1H), 1.31-1.34 (m, 12H); MS (ESI) *m/z* = 286 [M + H]⁺.

- 5 *(S)*-tert-Butyl ((4-chloro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate and *(R)*-tert-butyl ((4-chloro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate



- [0049] 25.7 g of tert-butyl ((4-chloro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate dissolved in acetonitrile (10 mg/ml) was resolved via chiral HPLC using ChiralPak® AD-H (250×30 mm I.D.) and SF CO₂ / methanol as eluent. Flow rate is 70mL/min. UV detection was monitored at 220 nm. Two peaks were collected and evaporated to give 10.65 g of enantiomer A (faster eluting isomer) and 10.15 g of enantiomer B (slower eluting isomer). Analysis of the pooled fractions using a ChiralPak® AD-3 (150×4.6 mm I.D.) and the same mobile phase showed enantiomer A with a retention time of 3.12 min and 98.7% e.e., and enantiomer B with a retention time of 3.44 min and 98.5% e.e.


(R)-3-(Aminomethyl)-4-chloro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3*H*)-ol hydrochloride



- 20 [0050] Enantiomer A (7.0 g, 20.5 mmol) was dissolved in 30 mL of dioxane and treated with 4M HCl (26.7 mL, 106.6 mmol) in dioxane. The reaction mixture was stirred at room temperature for overnight until the reaction was completed indicated by LC/MS. After dioxane was removed in vacuo and diethyl ether was added, an off-white solid was collected and dried under high-vacuum. This material was re-
- 25 dissolved in acetonitrile and water (1:1, v/v) and lyophilized to give 5.17 g of the title compound as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.11 (s, 1 H), 8.22

(s, 3H), 7.47 (d, 1H), 6.95 (d, 1H), 5.34-5.37 (m, 1H), 4.06-4.11 (m, 2H), 3.53-3.56 (m, 1H), 2.89 (m, 1H), 1.30-1.34 (m, 3H); MS (ESI) $m/z = 242.0$ $[M + H]^+$.

(S)-3-(Aminomethyl)-4-chloro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3*H*)-ol hydrochloride




5


[0300] Enantiomer B (7.0 g, 20.5 mmol) was dissolved in 30 mL of dioxane and treated with 4M HCl (26.7 mL, 106.6 mmol) in dioxane. The reaction mixture was stirred at room temperature for overnight until the reaction was completed indicated by LC/MS. After dioxane was removed in vacuo and diethyl ether was added, an off-white solid was collected and dried under high-vacuum. This material was re-dissolved in acetonitrile and water (1:1, v/v) and lyophilized to give 5.23 g of the title compound as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 9.11 (s, 1 H), 8.25 (s, 3H), 7.47 (d, 1H), 6.95 (d, 1H), 5.35-5.38 (m, 1H), 4.06-4.11 (m, 2H), 3.53-3.56 (m, 1H), 2.88 (m, 1H), 1.30-1.33 (m, 3H); MS (ESI) $m/z = 242.0$ $[M + H]^+$.

15 **M.** *(R)*-3-(Aminomethyl)-4-fluoro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3*H*)-ol hydrochloride and

N. *(S)*-3-(Aminomethyl)-4-fluoro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3*H*)-ol hydrochloride




tert-Butyl ((4-fluoro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate




[0051] This compound was prepared from 3-aminomethyl-7-ethoxy-4-fluoro-3H-benzo[*c*][1,2]-oxaborol-1-ol, hydrochloride, using the similar procedure as described above.

(S)-*tert*-Butyl ((4-fluoro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate and *(R)*-*tert*-butyl ((4-fluoro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate



[0052] 4.5 g of *tert*-butyl ((4-fluoro-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[*c*][1,2]oxaborol-3-yl)methyl)carbamate dissolved in ethanol (100 mg/ml) was resolved via chiral HPLC using ChiralCel® OZ-H column (250×30 mm I.D.) and SF CO₂ / hexane: ethanol(1:1) as eluent. Flow rate is 70mL/min. UV detection was monitored at 220 nm. Two peaks were collected and evaporated to give 2.1 g of enantiomer A (faster eluting isomer) and 2.2 g of enantiomer B (slower eluting isomer). Analysis of the pooled fractions using a ChiralCel® OZ-H (150×4.6 mm I.D.) and SF CO₂ / ethanol (0.05% DEA) as mobile phase showed enantiomer A with a retention time of 2.66 min and 99.5% e.e, and enantiomer B with a retention time of 3.31 min and 98.1% e.e.


(R)-3-(Aminomethyl)-4-fluoro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3H)-ol hydrochloride



[0053] Enantiomer A (2.1 g) was treated with 200 mL of 1.6 N HCl in MeOH and stirred at room temperature for 5 hours until the reaction was completed indicated by LC/MS. After water (100 mL) was added, the residue was lyophilized overnight to


give 1.40 g of the title compound as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ ppm 9.16 (br. s., 1H), 8.34 (br. s., 3H), 7.26 (t, 1H), 6.70 - 6.92 (m, 1H), 5.48 (d, 1H), 4.06 (q, 2 H), 3.35 (m, 1H), 2.88 (m, 1H), 1.30 (t, 3H); MS (ESI) m/z = 226.1 (M+1, positive).

5 *(S)*-3-(Aminomethyl)-4-fluoro-7-ethoxybenzo[*c*][1,2]oxaborol-1(3*H*)-ol hydrochloride



[0304] Enantiomer B (2.1 g) was treated with 200 mL of 1.6 N HCl in MeOH and stirred at room temperature for 5 hours until the reaction was completed indicated by LC/MS. After water (100 mL) was added, the residue was lyophilized overnight to give 1.43 g of the title compound as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ ppm 9.16 (br. s., 1H), 8.31 (br. s., 3 H), 7.26 (t, 1H), 6.70 - 6.92 (m, 1 H), 5.48 (d, 1H), 4.06 (q, 2 H), 3.35 (m, 1H), 2.88 (m, 1 H), 1.31 (t, 3 H); MS (ESI) m/z = 226.1 (M+1, positive).

O. 3-Aminomethyl-5-chloro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol hydrochloride



5-Chloro-2,3-dihydroxy-benzaldehyde

- 5 [0305] To a solution of 5-chloro-2-hydroxy-3-methoxy-benzaldehyde (7 g, 37.5 mmol) in anhydrous CH₂Cl₂ (200 mL) at 0 °C was added a solution of BBr₃ in CH₂Cl₂ (1 M, 93.7 mL, 93.7 mmol) and the reaction mixture was stirred overnight at room temperature. The solution was diluted with CH₂Cl₂ (200 mL), washed with water, brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure generating the title compound (6.2 g, 36.0 mmol, 96%) as a light yellow solid. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 11.02 (s, 1 H), 9.83 (s, 1 H), 7.18 (s, 1 H), 7.14 (d, *J*=2.3 Hz, 1 H), 5.71 (s, 1 H).

3-(3-Benzyloxy-propoxy)-5-chloro-2-hydroxy-benzaldehyde

- 15 [0306] To a solution of 5-chloro-2,3-dihydroxy-benzaldchde (3.1 g, 17.7 mmol) in anhydrous DMSO (20 mL) was added NaH (60% in mineral oil, 1.50 g, 35.4 mmol) portion-wise and the mixture was stirred for 30 minutes. The solution was cooled to 0 °C and a solution of 3-benzyloxy-1-bromopropane (3.1 mL, 17.7 mmol) in DMSO (3 mL) was added dropwise over 10 minutes period. The ice bath was removed. After overnight, the solution was diluted with EtOAc (100 mL), washed

with water, brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The product was purified by silica gel column chromatography (2:1 hexanes-EtOAc mobile phase) generating the title compound (5.3 g, 16.6 mmol, 94%) as a light yellow oil. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.81 (s, 1 H), 9.87 (s, 1 H), 7.42 - 7.28 (m, 5 H), 7.17 (s, 1 H), 7.08 (s, 1 H), 4.53 (s, 2 H), 4.17 (t, *J*=6.2 Hz, 2 H), 3.69 (t, *J*=5.8 Hz, 2 H), 2.15 (t, *J*=6.2 Hz, 2 H).

Trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-4-chloro-6-formyl-phenyl ester

[0307] To a solution of 3-(3-benzyloxy-propoxy)-5-chloro-2-hydroxy-

benzaldehyde (5.3 g, 16.6 mmol) and pyridine (3.4 mL, 41.5 mmol) in CH₂Cl₂ (70 mL) at 0 °C was added Tf₂O (3.1 mL, 18.3 mmol) drop-wise over 5 minutes period and the reaction mixture was stirred for 3 h at room temperature. The solution was diluted with CH₂Cl₂ (100 mL), washed with water, brine, dried over Na₂SO₄, then concentrated under reduced pressure. The product was purified by silica gel column chromatography (2:1 hexanes-EtOAc mobile phase) generating the title compound (3.8 g, 8.5 mmol, 51%) as a light yellow oil. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 10.18 (s, 1 H), 7.48 (d, *J*=2.3 Hz, 1 H), 7.37 - 7.29 (m, 6 H), 4.52 (s, 2 H), 4.23 (t, *J*=6.2 Hz, 2 H), 3.69 (t, *J*=5.8 Hz, 2 H), 2.16 (t, *J*=6.0 Hz, 2 H); ¹⁹F NMR (376 MHz, CHLOROFORM-*d*) δ ppm -73.23 (s).

3-(3-Benzyloxy-propoxy)-5-chloro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde

[0308] To a solution of trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-4-chloro-6-formyl-phenyl ester (3.8 g, 8.4 mmol) in anhydrous 1,4-dioxane (50 mL) was added bis(pinacolato)diborane (4.3 g, 16.9 mmol) and KOAc (2.5 g, 25.4 mmol) successively and the resulting solution was degassed with N₂ for 20 minutes.

PdCl₂(dppf) (0.5 g, 0.67 mmol) was added and the resulting mixture was stirred overnight at 90 °C. The solution was diluted with EtOAc (100 mL), washed with water, brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure.


The product was purified by silica gel column chromatography (4:1 hexanes-EtOAc

mobile phase) generating the title compound (3.4 g, 7.8 mmol, 92%) as a colorless oil. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 9.87 (s, 1 H), 7.40 - 7.28 (m, 6 H), 7.03 (br s, 1 H), 4.50 (s, 2 H), 4.09 (t, *J*=6.4 Hz, 2 H), 3.70 - 3.60 (m, 2 H), 2.10 (t, *J*=6.2 Hz, 2 H), 1.42 (s, 12 H).

7-(3-Benzoyloxy-propoxy)-5-chloro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol


[0309] To a solution of 3-(3-benzyloxy-propoxy)-5-chloro-2-(4,4,5-trimethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (3.4 g, 7.8 mmol) and nitromethane (1.7 mL, 31.3 mmol) in THF (20 mL) was added a solution of NaOH (0.025 M, 40 mL). After 12 h, 2 N HCl was added until pH was 1. The solution was diluted with EtOAc (150 mL), washed with water, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The product was purified by silica gel column chromatography (4:1 hexanes-EtOAc mobile phase) to give the title compound product (1.7 g, 56%) as a colorless gel. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 7.39 - 7.28 (m, 5 H), 6.92 (s, 1 H), 6.83 (s, 1 H), 5.85 (br s, 1 H), 5.81 (dd, *J*=8.5, 3.9 Hz, 1 H), 4.70 (dd, *J*=13.2, 3.9 Hz, 1 H), 4.59 (s, 2 H), 4.47 (dd, *J*=13.0, 8.7 Hz, 1 H), 4.21 - 4.07 (m, 2 H), 3.71 - 3.60 (m, 2 H), 2.10 (quin, *J*=5.7 Hz, 2 H).

5-Chloro-7-(3-hydroxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol




[0310] 7-(3-Benzoyloxy-propoxy)-5-chloro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (0.3 g, 0.91 mmol) in MeOH (30 mL) was added conc HCl (1 mL) and Pd(OH)₂ (10% w/w on carbon, 0.2 g) and the reaction vessel was pressurized to 40 psi with hydrogen for 30 minutes at room temperature. The mixture was filtered through a pad of Celite[®] and washed with EtOAc. The filtrate was concentrated in vacuo and the product was purified by prep. HPLC (C18 column, using acetonitrile and 0.1% AcOH/water solution gradient) provided the title compound (80 mg, 33%). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 9.19 (s, 1 H), 7.22 (s, 1 H), 6.97 (s, 1 H), 5.71 (dd, *J*=8.9, 2.3 Hz, 1 H), 5.32 (dd, *J*=13.2, 2.7 Hz, 1 H), 4.64 (dd, *J*=13.6, 8.9 Hz, 1 H), 4.53 (t, *J*=4.8 Hz, 1 H), 4.12 (t, *J*=6.0 Hz, 2 H), 3.57 (q, *J*=5.5 Hz, 2 H), 1.86 (t, *J*=6.2 Hz, 2 H).

3-Aminomethyl-5-chloro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol hydrochloride



- [0311] To a 5-chloro-7-(3-hydroxy-propoxy)-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol **7** (80 mg, 0.27 mmol) in methanolic ammonia solution (2 M, 20 mL) was added Ra/Ni (~0.1 g, 2800 Nickel slurry in water) and the reaction vessel was pressurized to 40 psi with hydrogen overnight at room temperature. The mixture was filtered through a pad of Celite® and washed with EtOAc. The filtrate was concentrated in vacuo and to the resulting residue was added water (1 mL), followed by conc HCl to pH 1. The heterogeneous mixture was lyophilized providing the title compound as a hygroscopic ivory solid (79 mg, quantitative). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.20 (s, 1 H), 6.99 (s, 1 H), 5.28 (dd, *J*=8.0, 2.5 Hz, 1 H), 4.13 (t, *J*=6.0 Hz, 2 H), 3.59 (t, *J*=6.0 Hz, 2 H), 3.47 (dd, *J*=13.0, 2.5 Hz, 1 H), 2.89 (dd, *J*=13.2, 8.6 Hz, 1 H), 1.89 (t, *J*=6.0 Hz, 2 H); MS (ESI) *m/z* = 272 (M+1, positive); HPLC purity: 96.83% (MaxPlot 200 – 400 nm), 95.40% (220 nm).

P. 3-Aminomethyl-7-(3-hydroxy-propoxy)-6-methoxy-3H-benzo[c][1,2]oxaborol-1-ol hydrochloride



3-(3-Benzyloxy-propoxy)-2-bromo-4-methoxy-benzaldehyde


- [0312] Synthesized according to the methods of general procedure 4 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants

and amounts: 2-bromo-3-hydroxy-4-methoxy-benzaldehyde (1.0 g, 4.32 mmol), (3-bromo-propoxymethyl)-benzene (0.76 mL, 4.32 mmol), cesium carbonate (2.11 g, 6.5 mmol), DMF (30 mL). Purification: flash chromatography (10% EtOAc/hexanes): yield 1.54 g (95%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.26 (s, 1 H), 7.73 (d, *J*=8.6 Hz, 1 H), 7.46-7.18 (m, 5 H), 6.95 (d, *J*=8.6 Hz, 1 H), 4.56 (s, 2 H), 4.14 (t, *J*=6.1 Hz, 2 H), 3.92 (s, 3 H), 3.77 (t, *J*=6.2 Hz, 2 H), 2.15 (quin, *J*=6.5 Hz, 2 H); MS (ESI): *m/z* = 381 (M+1, positive).

3-(3-Benzyloxy-propoxy)-4-methoxy-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde

- 10 [0313] Synthesized according to the methods of general procedure 5 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-benzyloxy-propoxy)-2-bromo-4-methoxy-benzaldehyde (14.82 g, 39 mmol), bis(pinacolato)diboran (14.86 g, 58.5 mmol), KOAc (11.46 g, 117 mmol), PdCl₂(dppf) (8.5 g, 11.7 mmol), dioxane (200 mL). Purification: flash column chromatography (15% EtOAc/hexanes): yield 3.42 g (22%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.79 (s, 1 H), 7.53 (d, *J*=8.2 Hz, 1 H), 7.40-7.24 (m, 5 H), 6.98 (d, *J*=8.2 Hz, 1 H), 4.53 (s, 2 H), 4.12 (t, *J*=6.4 Hz, 2 H), 3.88 (s, 3 H), 3.69 (t, *J*=6.4 Hz, 2 H), 2.10 (quin, *J*=6.5 Hz, 2 H), 1.44 (s, 12 H).


*7-(3-Benzyloxy-propoxy)-6-methoxy-3-nitromethyl-3H-benzo[*c*][1,2]oxaborol-1-ol*



- 20 [0314] Synthesized according to the methods of general procedure 8 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-benzyloxy-propoxy)-4-methoxy-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (3.36 g, 7.88 mmol), nitromethane (1.28 mL, 23.66 mmol), NaOH (0.22 g, 5.52 mmol), THF (6 mL), water (18 mL). Purification: flash column chromatography (30% EtOAc/hexanes): yield 1.2 g (41%). ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.36 (s, 1 H), 7.40-7.22 (m, 5 H), 7.22-7.04 (m, 2 H), 5.68 (dd, *J*=9.4, 2.7 Hz, 1 H), 5.29 (dd, *J*=13.4, 2.5 Hz, 1 H), 4.52 (dd, *J*=13.3, 9.4


Hz, 1 H), 4.45 (s, 2 H), 4.25 (t, $J=6.2$ Hz, 2 H), 3.75 (s, 3 H), 3.61 (t, $J=6.2$ Hz, 2 H), 1.92 (quin, $J=6.5$ Hz, 2 H).

3-Aminomethyl-7-(3-hydroxy-propoxy)-6-methoxy-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride



5 [0054] To a solution of 7-(3-benzyloxy-propoxy)-6-methoxy-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (0.5 g, 1.29 mmol) in methanolic ammonia (10 mL) was added palladium hydroxide (0.25 g, 45 wt%) in a hydrogenation bottle and the flask was charged with hydrogen at 45 psi for 18 h. The catalyst was filtered off and the
10 solvent was evaporated under reduced pressure. In order to assure all the ammonia has been stripped off, the compound was subjected to the high vacuum for 1 h. The crude (0.4 g) obtained was further dissolved in methanol (15 mL) and transferred to a hydrogenation bottle and concentrated HCl (5-6 drops) was added to make it to pH 2. To this solution palladium hydroxide (0.11 g, 25 wt%) was added and the flask was
15 charged with hydrogen to 45 psi for 1.5 h. The catalyst was filtered off through a pad of Celite® and the solvent evaporated. Purification was accomplished by preparative HPLC generating 0.16 g (41%) of the title compound as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ (ppm): 8.06 (br. s, 1 H), 7.17 (d, $J=8.2$ Hz, 1 H), 7.12 - 7.02 (m, 1 H), 5.32 (dd, $J=7.8, 2.3$ Hz, 1 H), 4.46 - 4.27 (m, 2 H), 4.14 (t, $J=4.5$ Hz, 2 H), 3.78
20 (s, 3 H), 3.44 (dd, $J=13.3, 2.7$ Hz, 1 H), 2.88 (dd, $J=13.3, 8.2$ Hz, 1 H), 2.08 - 1.94 (m, 2 H); MS (ESI): m/z = 268 (M+1, positive); HPLC purity: 95.35% (MaxPlot 200 - 400 nm), 97.48% (220 nm).

Q. 3-Aminomethyl-7-(3-hydroxy-propoxy)-6-methyl-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride



1,2-Dimethoxy-3-methyl-benzene

- 5 [0316] To a cooled (0 °C) solution of 1,2-dimethoxy-3-methylbenzene (2.05 g, 13.45 mmol) and TMEDA (2.8 mL, 18.83 mmol) in diethyl ether (100 mL) was added t-butyllithium (1.7 M in pentane, 9.5 mL, 16.14 mmol). The color of the solution changed to light yellow and after a few minutes a white precipitate was observed. The suspension was stirred at room temperature for 18 h, cooled to 0 °C and
- 10 dimethylformamide (2.08 mL, 26.90 mmol) was added dropwise. The precipitate disappeared and the color of the solution changed to light pink. After stirring for 0.5 h, ice was added followed by 1N HCl (30 mL), the compound was extracted into ethyl acetate, dried (Na₂SO₄) and the solvent was evaporated to obtain light brown oil. Purification by flash column chromatography (5% EtOAc/hexane) generated the title
- 15 compound: yield 1.4 g (58%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.34 (s, 1 H), 7.49 (d, *J*=8.2 Hz, 1 H), 7.01 (d, *J*=7.8 Hz, 1 H), 4.00 (s, 3 H), 3.86 (s, 3 H), 2.33 (s, 3 H). MS (ESI): *m/z* = 181 (M+1, positive).

2,3-Dihydroxy-4-methyl-benzaldehyde

- [0317] To a solution of 1,2-dimethoxy-3-methyl-benzene (13.8 g, 76.66 mmol)
- 20 cooled to -30 °C (dry ice/ acetone) in dichloromethane (200 mL) was added boron

trichloride (230 mL, 230 mmol) dropwise and the mixture was left to stir overnight at room temperature. The solution was cooled to 0 °C and ice/ water was added carefully, and then extracted with excess of dichloromethane. The organic layer was washed with water, dried (Na₂SO₄) and the solvent was evaporated. Purification by silica gel column chromatography (10-20% EtOAc/hexane) gave the title compound as a crystalline solid: yield 9.2 g (80%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 11.11 (s, 1 H), 9.82 (s, 1 H), 7.05 (d, *J*=8.2 Hz, 1 H), 6.81 (d, *J*=8.6 Hz, 1 H), 5.67 (s, 1 H), 2.33 (s, 3 H).

3-(3-Benzyloxy-propoxy)-2-hydroxy-4-methyl-benzaldehyde

[0318] Synthesized according to the methods of general procedure 4 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 2,3-dihydroxy-4-methyl-benzaldehyde (9 g, 59.21 mmol), (3-bromopropoxymethyl)-benzene (11.5 mL, 65.13 mmol), sodium tert-butoxide (12.52 g, 130.26 mmol) and DMSO (100 mL). Purification: flash column chromatography (5-10% EtOAc/hexane): yield 15.1 g (85%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 11.05 (s, 1 H), 9.84 (s, 1 H), 7.38-7.26 (m, 5 H), 7.20 (d, *J*=7.8 Hz, 1 H), 6.80 (d, *J*=8.2 Hz, 1 H), 4.55 (s, 2 H), 4.15 (t, *J*=6.2 Hz, 2 H), 3.73 (t, *J*=6.2 Hz, 2 H), 2.31 (s, 3 H), 2.19-2.00 (m, 2 H).

Trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-6-formyl-3-methyl-phenyl ester

[0319] Synthesized according to the methods of general procedure 6 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-Benzyloxy-propoxy)-2-hydroxy-4-methyl-benzaldehyde (0.3 g, 1.0 mmol), trifluoromethanesulfonic acid (0.34 mL, 2.0 mmol), pyridine (0.25 mL, 3.1 mmol), dichloromethane (15 mL). Purification: flash column chromatography (10-15% EtOAc/hexane): yield 0.25 g (59%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.14 (s, 1 H), 7.61 (d, *J*=7.8 Hz, 1 H), 7.42-7.29 (m, 6 H), 4.53 (s, 2 H), 4.07 (t, *J*=6.4 Hz, 2 H), 3.71 (t, *J*=6.1 Hz, 2 H), 2.40 (s, 3 H), 2.07 - 2.22 (m, 2 H). ¹⁹F NMR (400 MHz, CDCl₃) δ (ppm): -73.63 (s).

3-(3-Benzyloxy-propoxy)-4-methyl-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde


[0320] Synthesized according to the methods of general procedure 5 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants

and amounts: trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-6-formyl-3-methyl-phenyl ester (0.26 g, 0.6 mmol), bis(pinacolato)diboran (0.31 g, 1.2 mmol), KOAc (0.18 g, 1.8 mmol), PdCl₂(dppf) (0.13 g, 0.18 mmol), THF (10 mL).

Purification: flash column chromatography (15% EtOAc/hexane): yield 0.091 g


- 5 (36%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.89 (s, 1 H), 7.47 (d, *J*=7.4 Hz, 1 H), 7.40-7.25 (m, 6 H), 4.52 (s, 2 H), 4.02 (t, *J*=6.4 Hz, 2 H), 3.70 (t, *J*=6.2 Hz, 2 H), 2.32 (s, 3 H), 2.20-2.08 (m, 2 H), 1.45 (s, 12 H).

7-(3-Benzyloxy-propoxy)-6-methyl-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol



- 10 [0321] Synthesized according to the methods of general procedure 8 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-benzyloxy-propoxy)-4-methyl-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (1.14 g, 2.78 mmol), nitromethane (0.45 mL, 8.34 mmol), NaOH (0.78 g, 1.95 mmol), THF (3 mL) and water (9 mL). Purification:
- 15 flash column chromatography (25% EtOAc/hexanes): yield 0.42 g (41%). ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.41 (s, 1 H), 7.37-7.18 (m, 6 H), 7.03 (d, *J*=7.8 Hz, 1 H), 5.69 (dd, *J*=9.2, 2.5 Hz, 1 H), 5.28 (dd, *J*=13.7, 2.7 Hz, 1 H), 4.52 (dd, *J*=13.3, 9.4 Hz, 1 H), 4.46 (s, 2 H), 4.33 (t, *J*=6.1 Hz, 2 H), 3.58 (t, *J*= 6.1 Hz, 2 H), 2.12 (s, 3 H), 2.01-1.87 (m, 2 H). MS (ESI): *m/z* = 370 (M-1, negative).


- 20 *3-Aminomethyl-7-(3-hydroxy-propoxy)-6-methyl-3H benzo[c][1,2] oxaborol-1-ol hydrochloride*



[0322] To a solution of 7-(3-benzyloxy-propoxy)-6-methyl-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (0.42 g, 1.13 mmol) in methanolic ammonia (15 mL) was

added palladium hydroxide (0.2 g, 45 wt%) in a hydrogenation bottle and the flask was charged with hydrogen at 45 psi for 18 h. The catalyst was filtered off and the solvent was evaporated under reduced pressure. In order to assure all the ammonia has been stripped off, the compound was subjected to the high vacuum for 1 h. The crude (0.38 g) obtained was further dissolved in methanol (15 mL) and transferred to a hydrogenation bottle and concentrated HCl (5-6 drops) was added to make it to pH 2. To this solution palladium hydroxide (0.1 g, 25 wt%) was added and the flask was charged with hydrogen at 45 psi for 1.5 h. The catalyst was filtered off through a pad of Celite® and the solvent evaporated. Purification by preparative HPLC provided 0.12 g (39%) of the title compound as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.40 (s, 1 H), 7.12 (d, *J*=7.4 Hz, 1 H), 6.80 (d, *J*=7.4 Hz, 1 H), 5.08-4.91 (m, 1 H), 4.49-4.21 (m, 2 H), 3.64 (t, *J*=5.7 Hz, 2 H), 3.17 (dd, *J*=12.9, 3.1 Hz, 1 H), 2.66 (dd, *J*=12.7, 8.0 Hz, 1 H), 2.14 (s, 3 H), 1.81 (quin, *J*=5.7 Hz, 2 H). MS (ESI): *m/z* = 252 (*M*+1, positive); HPLC purity: 98.25% (MaxPlot 200 – 400 nm), 98.39% (220 nm).

R. 3-Aminomethyl-6-fluoro-7-(3-hydroxy-propoxy)-3H-benzo[*c*][1,2]oxaborol-1-ol; hydrochloride salt



4-Fluoro-2,3-dihydroxy-benzaldehyde

[0055] To a solution of 3-fluoro-benzene-1,2-diol (20 g, 156 mmol) in anhydrous acetonitrile (400 mL) was added magnesium chloride (37.1 g, 312 mmol), paraformaldehyde (31.6 g) and triethylamine (134 mL, 975 mmol). The reaction mixture was heated at 80 °C for 8 h. The reaction mixture was cooled to room temperature and the solid was collected by filtration. The solid was treated with cold

2 N HCl and the aqueous layer was extracted with EtOAc. The organic layer was concentrated in vacuo yielding 20.4 g of crude. After a second run 40.8 g of crude was dissolved in DMF (1 L), cooled to 0 °C, added to Cs₂CO₃ (340 g, 1.04 mol) portion-wise. Then methyl iodide (330 mL, 5.28 mol) was added. After warming to
5 room temperature and stirring overnight the solution was filtered, ethyl acetate was added and the organic layer was washed with water (3 X). After concentration in vacuo the product was purified by Biotage® silica gel chromatography (2% to 3% to 10% to 20% EtOAc/ hexanes) resulting in 14.8 g of dimethoxy compound. This material was dissolved in DCM and cooled to -30 °C and BCl₃ (1 M in DCM, 134
10 mL, 0.1343 mol) was added to the solution at -30 °C. After overnight at room temperature, the solution was cooled to -70 °C and BBr₃ (1 M in DCM, 67.25 mL, 0.067 mol) was added. After overnight warming to room temperature, the solution was cooled in an ice bath and slowly ice water was added. The DCM layer was separated and the aqueous layer was extracted with DCM (2X). The combined
15 organic layer was extracted with brine (2X), dried over Na₂SO₄, filtered and concentrated in vacuo. After triturating the residue obtained with hexanes/DCM (6:4) the 5.60 g (11% yield) of the title compound obtained was a brownish pink solid. This material was used in the next step without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 11.36 (s, 1H), 9.83 (s, 1H), 7.16 – 7.13 (m, 1H), 6.82 –
20 6.78 (m, 1H), 5.48 (brs, 1H). ¹⁹F NMR (376 MHz, DMSO-*d*₆ with D₂O) δ (ppm): -119.03 - -119.08 (m, 1F).

3-(3-Benzyloxy-propoxy)-4-fluoro-2-hydroxy-benzaldehyde


[0056] Synthesized according to the methods of general procedure 4 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants
25 and amounts: 4-fluoro-2,3-dihydroxy-benzaldehyde (5.15 g, 32.9 mmol), NaOtBu (6.95 g, 72.3 mmol), DMSO (200 mL), (3-bromo-propoxymethyl)-benzene (8.31 g, 36.3 mmol). Purification: Biotage® silica gel chromatography (hexanes/ ethyl acetate gradient) generated 4.00 g of a mixture of the title compound and the dialkylated product. This material was used in the next step without further purification. ¹H
30 NMR (400 MHz, DMSO-*d*₆) δ (ppm): 10.13 (s, 1H), 7.66 -7.54 (m, 5H), 7.53 – 7.33 (m, 1H), 6.92 – 6.90 (m, 1H), 4.53 (s, 2H), 4.52 – 4.44 (m, 2H), 3.71 – 3.62 (m, 2H), 2.18 – 2.13 (m, 2H). ¹⁹F NMR (376 MHz, DMSO-*d*₆ with D₂O) δ (ppm): -121.03 - -121.08 (m, 1F).

3-(3-Benzyloxy-propoxy)-4-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde

[0057] Synthesized according to the methods of general procedure 6 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-benzyloxy-propoxy)-4-fluoro-2-hydroxy-benzaldehyde (4.00 g, 13.1 mmol), pyridine (2.34 mL, 28.9 mmol), DCM (100 mL), triflate anhydride (2.21 mL, 13.5 mmol). Purification: Biotage® silica gel chromatography (hexanes/ ethyl acetate gradient) generated 2.00 g of triflate that was used immediately according to general procedure 5 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692).

[0058] Synthesized according to the methods of general procedure 5 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: trifluoro-methanesulfonic acid 2-(3-benzyloxy-propoxy)-3-fluoro-6-formyl-phenyl ester (2.0 g, 4.57 mmol), THF (15 mL), B₂pin₂ (2.20 g, 8.66 mmol), KOAc (1.60 g, 16.3 mmol), PdCl₂(dppf)DCM (0.40 g, 0.55 mmol). Purification: Biotage® silica gel chromatography (hexanes/ ethyl acetate gradient) generated 0.50 g (26% yield) of the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.86 (s, 1H), 7.51 (dd, *J* = 8.2, 4.0 Hz, 1H), 7.34 – 7.31 (m, 5H), 7.21 (dd, *J* = 11.0, 8.2 Hz, 1H), 4.51 (s, 2H), 4.24 (td, *J* = 6.5, 2.0 Hz, 2H), 3.67 (t, *J* = 6.2 Hz, 2H), 2.11 – 2.07 (m, 2H); 1.33 (s, 12H); ¹⁹F NMR (376 MHz, DMSO-*d*₆ with D₂O) δ (ppm): -120.3 - -121.1 (m, 1F).


*7-(3-Benzyloxy-propoxy)-6-fluoro-3-nitromethyl-3H-benzo[*c*][1,2]oxaborol-1-ol*



[0059] Synthesized according to the methods of general procedure 8 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 3-(3-benzyloxy-propoxy)-4-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (0.40 g, 0.966 mmol), nitromethane (0.15 mL, 2.89 mmol), NaOH (0.038 g, 0.96 mmol), THF (10 mL), water (10 mL). This generated 0.34 g (94% yield) of the title compound. ¹H NMR (400 MHz, DMSO-*d*₆)


δ (ppm): 9.83 (s, 1H), 7.32 – 7.02 (m, 7H), 5.78 – 5.75 (m, 1H), 5.28 -5.23 (m, 1H), 4.60 – 4.56 (m, 1H), 4.42 (s, 2H), 4.37 (brs, 2H), 3.57 (brs, 2H), 1.92 (brs, 2H). ^{19}F NMR (376 MHz, DMSO- d_6 with D $_2$ O) δ (ppm): -132.3 (1F).

5 *3-Aminomethyl-6-fluoro-7-(3-hydroxy-propoxy)-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride salt*



[0328] To a mixture of 7-(3-benzyloxy-propoxy)-6-fluoro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (0.34 g, 0.906 mmol) and methanolic ammonia (2M, 20 mL) in a Parr apparatus was added Pd(OH) $_2$ on carbon (0.30 g). The apparatus was charged with hydrogen (~ 40 psi) and was shaken overnight at rt. The suspension was filtered through Celite[®] with methanol washing and was concentrated in vacuo. The 310 mg of cream colored solid was dissolved in methanol (20 mL), transferred to Parr apparatus and the pH was adjusted to ~ 3 with a few drops of concentrated HCl. Then Pd(OH) $_2$ on carbon (0.20 g) was added and the apparatus was charged with hydrogen (~ 40 psi). After 35 minutes, the suspension was filtered through Celite[®] with methanol washing and was concentrated in vacuo. Purification was accomplished by reverse phase preparative HPLC (acetonitrile/ water (0.1% AcOH) gradient) generating 100 mg (43% yield) of the title compound as a white solid. mp 265 - 267 °C; ^1H NMR (400 MHz, DMSO- d_6 with D $_2$ O) δ (ppm): 7.41 (dd, J = 11.1, 8.2 Hz, 1H), 7.07 (dd, J = 7.9, 2.8 Hz, 1H), 5.29 (d, J = 7.0 Hz, 1H), 4.36 (t, J = 6.0 Hz, 2H), 3.62 (br.s, 2H), 3.45 (d, J = 12.9 Hz, 1H), 2.92 – 2.86 (m, 1H), 1.93 – 1.83 (m, 2H); ^{19}F NMR (376 MHz, DMSO- d_6 with D $_2$ O) δ (ppm): -135.0 (1F); MS (ESI) m/z = 256 (M+1, positive); HPLC purity: 98.57% (MaxPlot 200-400 nm), 97.28% (220 nm).

S. **3-Aminomethyl-7-ethoxy-6-methoxy-3H-benzo[c][1,2]oxaborol-1-ol;**
hydrochloric acid salt



5 **2-Bromo-3-ethoxy-4-methoxy-benzaldehyde**


[0329] Ethyl bromide (2.88 g, 26.4 mmol) was added to a mixture of 2-bromo-3-hydroxy-4-methoxybenzaldehyde (5.08 g, 22 mmol) and potassium carbonate (4.56 g, 33 mmol) in anhydrous DMF (50 mL) at room temperature under nitrogen. The reaction mixture was stirred at 35 °C for 18 h, diluted with EtOAc (150 mL), washed with water (2x50 mL), brine, dried over Na₂SO₄ and concentrated to give crude product as a white solid. Purification by silica column chromatography (eluant : 30% EtOAc in Hexanes) to generate 5.65 g (99% yield) of the title compound as a white solid. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 10.26 (s, 1H), 7.73 (d, *J* = 8.4 Hz, 1H), 6.95 (d, *J* = 8.8 Hz, 1H), 4.08 (q, *J* = 7.1 Hz, 2H), 3.95 (s, 3H), 1.46 (t, *J* = 7.0 Hz, 3H); MS (ESI) *m/z* = 261 (M+1, positive).

[3-Ethoxy-4-methoxy-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde

[0330] Synthesized according to the methods of general procedure 5 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 2-bromo-3-ethoxy-4-methoxy-benzaldehyde (4 g, 15.43 mmol), KOAc (4.55 g, 46.29 mmol), bis(pinacolato)diboron (7.84 g, 30.86 mmol). PdCl₂(dppf) (0.91 g, 1.24 mmol) in dry dioxane (90 mL). The crude product was purified by silica gel column chromatography (eluant : EtOAc / hexanes 1:9 then 1:3) to afford the title


compound as a white solid (1.50 g, 32 % yield). ^1H NMR (400 MHz, CDCl_3) δ (ppm) 9.71 (s, 1H), 7.44 (d, $J = 8.4$ Hz, 1H), 6.89 (d, $J = 8.4$ Hz, 1H), 4.01 (q, $J = 7.1$ Hz, 2H), 3.82 (s, 3H), 1.38 (s, 12H), 1.35 (t, $J = 7.0$ Hz, 3H).

7-Ethoxy-6-methoxy-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol



5 [0331] Synthesized according to the methods of general procedure 9 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: [3-ethoxy-4-methoxy-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (1.47 g, 4.80 mmol), nitromethane (0.92 g, 14.4 mmol), CATBr (88
10 mg, 0.24 mmol) in dry THF (20 mL) and NaOH (0.025 M aqueous solution). Purification by silica gel column chromatography (eluant: 10% EtOAc/ hexane to 30% EtOAc/ hexane) to obtain the title compound as a yellow solid (0.75 g, 59 %). ^1H NMR {400 MHz, $\text{DMSO}-d_6 + \text{D}_2\text{O}$ (0.01 ml)} δ (ppm) 9.34 (s, 1H), 7.18 (d, $J = 8.1$ Hz, 1H), 7.11 (d, $J = 8.0$ Hz, 1H), 5.68 (dd, $J = 9.2, 2.0$ Hz, 1H), 5.29 (dd, $J = 13.2, 2.8$ Hz, 1H), 4.55-4.50 (m, 1H), 4.20 (q, $J = 7.0$ Hz, 2H), 3.77 (s, 3H), 1.27 (t, $J = 7.0$ Hz, 3H); MS (ESI) $m/z = 261$ (M-1, negative).

3-Aminomethyl-7-ethoxy-6-methoxy-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride salt




20 [0332] Synthesized according to the methods of general procedure 13 in U.S. Pat. Pub. No. 20090227541 (U.S. Pat. App. No. 12/142,692) using the following reactants and amounts: 7-Ethoxy-6-methoxy-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (0.97 g, 3.63 mmol), glacial acetic acid (20 mL), $\text{Pd}(\text{OH})_2$ on carbon (20% metal content, 50% weight-wet) (300 mg). Purification: preparative HPLC (C18 column, using acetonitrile and 0.1% AcOH/water solution) provided the title compound (0.28
25 g; 28% yield). m.p. 202-204 °C. ^1H NMR {400 MHz, CD_3OD } δ (ppm) 7.20 (d, $J = 8.0$ Hz, 1H), 7.08 (d, $J = 8.1$ Hz, 1H), 5.40 (dd, $J = 8.4, 2.8$ Hz, 1H), 4.23 (q, $J = 7.1$ Hz, 2H), 3.31 (s, 3H), 3.56 (dd, $J = 13.6, 7.2$ Hz, 1H), 2.92 (dd, $J = 13.2, 7.2$ Hz, 1H),

1.33 (t, $J = 7$ Hz, 3H); MS (ESI) $m/z = 238$ (M+1, positive); HPLC purity: 98.79% (MaxPlot 200 – 400 nm) and 99.13% (220 nm).

T. 3-Aminomethyl-7-ethoxy-6-fluoro-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride salt

5



4-Fluoro-2,3-dihydroxybenzaldehyde

[0333] To a solution at -78 °C of 2,3-dimethoxy-4-fluorobenzaldehyde (7.0 g, 38.0 mmol) in dry dichloromethane (150 mL) was added dropwise BBr_3 (23.8 g, 95.0 mmol) in dichloromethane (30 mL). Reaction mixture was allowed to attain room temperature and stirred for 18 h. Then reaction mixture was cooled to -78 °C, and quenched with a mixture of methanol (10 mL) and water (50 mL) and stirred at room temperature for 30 min. Precipitated solid was separated by filtration and washed with cold dichloromethane. Dichloromethane layer was concentrated to yield the title compound as a solid (5.2 g, 88%). ^1H NMR (400 MHz, $\text{CHLOROFORM-}d$) δ ppm: 11.38 (s, 1 H), 9.84 (s, 1 H), 7.15 (dd, $J=8.6, 5.5$ Hz, 1 H), 6.81 (t, $J=9.4$ Hz, 1 H), 5.47 (s, 1 H); MS (ESI) $m/z = 155$ (M-1, negative).

3-Ethoxy-4-fluoro-2-hydroxybenzaldehyde

[0334] To a solution of 2,3-dihydroxy-4-fluorobenzaldehyde (3.0 g, 19.23 mmol) in DMSO (100 mL), $\text{NaOBu-}t$ (3.692 g, 38.46 mmol) was added in portions at room temperature and stirred for 15 min. Then iodoethane was added dropwise at room


temperature and stirred for 18 h. The reaction mixture was poured onto crushed ice (200 mL) and acidified with 2.5 M HCl to pH 3.0. The product was extracted with ethyl acetate (2 x 100 mL), concentrated and the product was chromatographed on a column of silica gel (Hex:EtOAc = 95:5) to give the title compound as a crystalline solid (2.3 g, 65%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm: 11.36 (s, 1 H), 9.83 (s, 1 H), 7.39 - 7.19 (m, 1 H), 6.77 (t, *J*=9.2 Hz, 1 H), 4.22 (q, *J*=7.0 Hz, 2 H), 1.40 (t, *J*=7.0 Hz, 3 H); MS (ESI) *m/z* = 183 (M+1, positive).

Trifluoromethanesulfonic acid 2-ethoxy-3-fluoro-6-formyl-phenyl ester

[0060] To a mixture of 3-ethoxy-4-fluoro-2-hydroxybenzaldehyde (2.208 g, 12.0 mmol) and pyridine (1.986 g, 24.0 mmol) in dichloromethane (30.0 mL) at 0 °C was added dropwise trifluoromethanesulfonic anhydride (4.060, 14.4 mmol) in dichloromethane (5.0 mL). The reaction mixture was stirred at 0 °C for 2 h and room temperature for 3 h. Then diluted with dichloromethane (40 mL), washed with 2M HCl, brine and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to give the title compound as a light yellow liquid (3.3 g, 87%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm: 10.15 (s, 1 H), 7.66 (dd, *J*=8.6, 5.5 Hz, 1 H), 7.28 – 7.22 (m, 1 H), 4.36 (q, *J*=6.9 Hz, 2 H), 1.47 (t, *J*=7.0 Hz, 3 H).


3-Ethoxy-4-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde

[0061] To a solution of trifluoromethanesulfonic acid 2-ethoxy-3-fluoro-6-formyl-phenyl ester (2.2 g, 6.96 mmol) in dry THF (35.0 mL) bis(pinacolato)diboron (2.134 g, 8.4 mmol), PdCl₂(dppf) (367 mg, 0.5 mmol) and potassium acetate (1.372 g, 14.0 mmol) were added and purged with nitrogen for 15 min. The reaction mixture was heated under reflux for 24 h. Cooled to room temperature and diluted with ethyl acetate (40 mL) and filtered through Celite®. The solvent was removed under reduced pressure and the residue was chromatographed on a column of silica gel (Hex:EtOAc = 9:1) to give the title compound as an off-white solid (850.0 mg, 42 %). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm: 9.87 (s, 1 H), 7.51 (dd, *J*=8.2, 4.3 Hz, 1 H), 7.22 (dd, *J*=10.9, 8.6 Hz, 1 H), 4.20 (q, *J*=7.0 Hz, 2 H), 1.46 (s, 12 H), 1.40 (t, *J*=7.0 Hz, 3 H); MS (ESI) *m/z* = 295 (M+1, positive).

7-Ethoxy-6-fluoro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol

[0337] To a cooled solution of sodium hydroxide (80 mg, 2.0 mmol) in water (3.0 mL), nitromethane (244.0 mg, 4.0 mmol) was added at 0 °C and stirred for 10 min.


- 5 Then 3-ethoxy-4-fluoro-2-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzaldehyde (588.0 mg, 2.0 mmol) in THF (5.0 mL) was added. The reaction mixture was stirred at for 1 h at 0 °C and for 2 h at room temperature. The reaction mixture was acidified with 2.5 M HCl (1.0 mL) and the product was extracted with ethyl acetate (2 x 20 mL). The organic extracts were combined and washed with brine
- 10 and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure and product was chromatographed on a column of silica gel(CH₂Cl₂ : MeOH = 95:5) to give the title compound as a solid (350 mg, 69%). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm: 7.40 (dd, *J*=11.3, 8.2 Hz, 1 H), 7.16 (dd, *J*=8.0, 3.3 Hz, 1 H), 5.74 (d, *J*=9.0 Hz, 1 H), 5.30 (dd, *J*=13.5, 2.2 Hz, 1 H), 4.62 (dd, *J*=13.5, 9.2 Hz, 1 H),
- 15 4.35 (q, *J*=6.8 Hz, 2 H), 1.28 (t, *J*=6.8 Hz, 3 H); MS (ESI) *m/z* = 254 (M-1, negative).

3-Aminomethyl-7-ethoxy-6-fluoro-3H-benzo[c][1,2]oxaborol-1-ol; hydrochloride salt

- [0338] To a solution of 7-ethoxy-6-fluoro-3-nitromethyl-3H-benzo[c][1,2]oxaborol-1-ol (320.0 mg, 1.25 mmol) in methanol (5.0 mL), 5.0 mL of
- 20 2M ammonia in methanol and 160 mg of Pd(OH)₂ on C were added and hydrogenated at 45 PSI for 18 h. Catalyst was removed by filtration and the filtrate was concentrated to generating an off-white solid (250 mg). This solid was dissolved in methanol (3 mL) and 3.0 mL of 1.2 M HCl in methanol was added and stirred at room temperature for 3 h. Excess HCl and solvent were removed under reduced pressure
- 25 and the product was triturated with ether to give the title compound as an off-white solid (140 mg, 43%). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm: 9.43 (s, 1 H), 8.13 (br. s., 3 H), 7.40 (dd, *J*=11.5, 8.0 Hz, 1 H), 7.16 (dd, *J*=7.8, 3.1 Hz, 1 H), 5.32 (d, *J*=6.3

Hz, 1 H), 4.35 (q, $J=7.0$ Hz, 2 H), 3.43 (br. s., 1 H), 2.92 (br. s., 1 H), 1.29 (t, $J=7.0$ Hz, 3 H); ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ ppm -135 (s, 1F); MS (ESI) m/z = 226 (M+1, positive); HPLC purity: 95.81% (MaxPlot 200-400 nm), 94.73% (220 nm).

U. **3-(Aminomethyl)-5-chloro-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol**



4-Chloro-2-ethoxy-6-formylphenyl trifluoromethanesulfonate

[0339] To a solution of 3-ethoxy-2-hydroxybenzaldehyde (20 g, 120.4 mmol) in AcOH (200 mL) was added N-chlorosuccinimide (16.1 g, 120.4 mmol). The reaction mixture was heated up to 105 °C for 30 min. After cooled down to room temperature, the mixture was stirred for additional 2.5 h. Subsequently, 200 mL of water was added slowly over 10 min. The mixture was filtered and dried to give a yellow solid, which was recrystallized in ethanol to give 4 g of the target compound (4 g, 17% yield).

4-Chloro-2-ethoxy-6-formylphenyl trifluoromethanesulfonate

[0340] To a solution of 5-chloro-3-ethoxy-2-hydroxybenzaldehyde (2.0 g, 10.0 mmol) in pyridine (2 mL) and DCM (20 mL) at 0 °C was dropwise added trifluoromethanesulfonic anhydride (1 mL). The reaction was stirred for 1 h at 0 °C before quenched with ice-water. The organic layer was washed with sat. aqueous NaHCO_3 (20 mL) and brine (20 mL), dried over anhydrous Na_2SO_4 and then concentrated in vacuo. The residue was purified by column chromatography on silica gel to give target compound (2.0 g, yield: 60%).


5-Chloro-3-ethoxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde

[0062] A mixture of 4-chloro-2-ethoxy-6-formylphenyl trifluoromethanesulfonate (330 mg, 1 mmol), KOAc (350 mg, 2.0 mmol), bis(pinacolato)diborane (600 mg, 2.0 mmol) and PdCl₂(dppf)CH₂Cl₂ (65 mg, 0.08 mmol, 8 mol%) in dioxane (30 mL) was degassed for 15 min with N₂ and stirred at 100 °C for 3 h. After quenched with ice-water, the reaction mixture was extracted with EtOAc (3x30 mL). The combined organic layers were washed with sat. aqueous NaHCO₃ (20 mL) and brine (20 mL), dried over anhydrous Na₂SO₄ and then concentrated. The residue was purified by column chromatography on silica gel to give the compound (150 mg, yield: 43%).

5-Chloro-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-1(3H)-ol

[0063] The mixture of 5-chloro-3-ethoxy-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde (310 mg, 1 mmol), NaOH (40 mg, 1 mmol) and CTAB (5 mg, 0.05 mmol) in H₂O (2 mL) and THF (10 mL) was stirred for 0.5 h at room temperature. After dropwise addition of nitromethane (0.2 mL, 2 mmol), the reaction mixture was stirred at room temperature for 3 h. Then the cyclization was afforded by adding the diluted aqueous HCl solution (2 N) to pH=2 and then extracted with EtOAc (3x30 mL). The combined organic layers were washed with brine (25 mL), dried over anhydrous Na₂SO₄ and concentrated to dryness. The residue was purified by prep-HPLC to give the compound (100 mg, yield: 56%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.20 (s, 1 H), 7.22 (s, 1H), 6.96 (s, 1H), 5.69-5.72 (m, 1H), 5.30-5.34 (m, 1H), 4.61-4.67 (m, 1H), 4.10-4.15 (m, 2H), 1.31-1.34 (m, 3H);

3-(Aminomethyl)-5-chloro-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol



[0064] A mixture of 5-chloro-7-ethoxy-3-(nitromethyl)benzo[c][1,2]oxaborol-

1(3H)-ol (270 mg, 1.0 mmol), Raney-Ni (~125 mg) and 2 M NH₃ in EtOH (2 mL) in EtOH (10 mL) was shaken under an atmosphere of H₂ for 2 h at room temperature. The mixture was filtered through a bed of Celite® and the filtrate was concentrated in vacuo. The crude amine was dissolved in EtOAc (2 mL) and HCl solution in Et₂O (20 mL) was added immediately. After 1 h, the suspension was filtered and the resulting solid was washed with hexanes to give compound the target compound (100

mg, yield: 43%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.06 (s, 1H), 8.18 (s, 3H), 7.22 (s, 1H), 6.96 (s, 1H), 5.27-5.29 (m, 1H), 4.10-4.13 (m, 2H), 3.40-3.47 (m, 1H), 2.87-2.92 (m, 1H), 1.30-1.36 (m, 3H); MS (ESI) m/z = 242 $[\text{M} + \text{H}]^+$.


V. **(S)-3-(aminomethyl)-4-bromo-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride**

Step 1: *tert*-butyl (7-ethoxy-1-hydroxy-1,3-dihydrobenzo[c][1,2]oxaborol-3-yl)methyl-carbamate




[0344] To the mixture of 3-(aminomethyl)-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride salt (5.0 g, 20.5 mmol) and triethylamine (10.4 g, 103.0 mmol) in dichloromethane (250 mL) at 0 °C was added di-*tert*-butyl dicarbonate (6.7 g, 30.8 mmol). The mixture was stirred for 4 h at room temperature. After the reaction was quenched with sat. NaHCO_3 (500 mL), the resulting mixture was extracted with EtOAc (3x300 mL), and the combined organic layers were dried over anhydrous Na_2SO_4 and concentrated in vacuum. The residue was purified by flash-column chromatography (2.5 % to 5.0 % MeOH in DCM) to give the product (5.51 g, yield 87%).

Step 2: *tert*-butyl (4-bromo-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[c][1,2]oxaborol-3-yl)-methylcarbamate



[0345] To the solution of *tert*-butyl (7-ethoxy-1-hydroxy-1,3-dihydrobenzo[c][1,2]oxaborol-3-yl)methyl-carbamate (5.5 g, 17.9 mmol) and 1-bromopyrrolidine-2,5-dione (3.8 g, 21.5 mmol) in CH_3CN (1100 mL) was added 2,2'-Azobis(2-methylpropionitrile) (220 mg). The mixture was stirred for 1 h at 90 °C. The reaction mixture was then concentrated in high vacuum and the residue was purified by column chromatography (2.5 % to 5.0 % MeOH in DCM) to give the product (3.7 g, yield 54%). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.90 (s, 1H), 7.55-7.53 (d, 1H), 6.85-6.82 (d, 1H), 5.08-5.07 (d, 1H), 4.11-4.07 (m, 2H), 3.82-3.79 (bd, 1H), 3.06-3.03 (m, 1H), 1.39 (s, 9H), 1.30 (t, 3H); MS (ESI) m/z = 387 $[\text{M} + \text{H}]^+$.

Step 3: (S)-3-(aminomethyl)-4-bromo-7-ethoxybenzo[c][1,2]oxaborol-1(3H)-ol hydrochloride



[0065] The mixture of *tert*-butyl (4-bromo-7-ethoxy-1-hydroxy-1,3-dihydrobenzo[c][1,2]oxaborol-3-yl)-methylcarbamate (3.7 g, 9.6 mmol) in 4N HCl in dioxane (12 ml, 48.0 mmol) was stirred at room temperature for 2 h and then concentrated to dryness (water bath < 30 °C). The residue was triturated with DCM/ether (1/10, 2x10 mL) and the white solid was dried in high vacuum to give the product (2.96 g, yield: 92%). ¹H NMR (300 MHz, DMSO-*d*₆) 9.11 (s, 1H), 8.1 (bs, 3H), 7.63-7.60 (d, 1H), 6.92-6.89 (d, 1H), 5.27-5.24 (m, 1H), 4.12-4.05 (m, 2H), 3.62-3.57 (m, 1H), 2.99-2.92 (m, 1H), 1.34-1.30 (t, 3H); MS (ESI) *m/z* = 287 [M + H]⁺.

EXAMPLE 2

LeuRS IC50 Testing

[0066] Experiments were performed in 96-well microtiter plates, using 80 μL reaction mixtures containing 50 mM HEPES-KOH (pH 8.0), 30 mM MgCl₂ and 30 mM KCl, 13 μM [¹⁴C]leucine (306 mCi/mmol, Perkin-Elmer®), 15 uM total E.coli tRNA (Roche®, Switzerland), 0.02% (w/v) BSA, 1 mM DTT, 0.2 pM LeuRS and 4 mM ATP at 30°C. Reactions were started by the addition of 4 mM ATP. After 7 minutes, reactions were quenched and tRNA was precipitated by the addition of 50 μL of 10% (w/v) TCA and transferred to 96-well nitrocellulose membrane filter plates (Millipore® Multiscreen HTS, MSHAN4B50). Each well was then washed three times with 100 μL of 5% TCA. Filter plates were then dried under a heat lamp and the precipitated [¹⁴C]leucine tRNA^{Leu} was quantified by liquid scintillation counting using a Wallac MicroBeta Trilux model 1450 liquid scintillation counter (PerkinElmer®, Waltham MA).

[0067] To determine the inhibitor concentration which reduces enzyme activity by 50% (IC₅₀), increasing concentrations of inhibitor were incubated with LeuRS enzyme, tRNA and leucine for 20 minutes. Reactions were initiated by the addition of 4 mM ATP and stopped after 7 minutes then precipitated and counted to quantify radioactivity.

[0068] Biochemical testing results for exemplary compounds of the invention are provided in FIG. 1.

EXAMPLE 3

Antibacterial MIC Testing

5 [0069] All MIC testing of bacteria followed the Clinical and Laboratory Standards Institute (CLSI) guidelines for antimicrobial testing of aerobic bacteria (Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard - Seventh Edition)(M07-A7) and anaerobic bacteria (Methods for
10 Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard - Seventh Edition) (M11-A7).

[0070] Antibacterial MIC testing results for exemplary compounds of the invention are provided in FIG. 1.

EXAMPLE 4

Microplate Alamar Blue Assay (MABA)

15 [0071] The microplate alamar blue assay (MABA) was essentially performed as described by Collins, L., *et al.*, *Antimicrob Agents Chemother* 41: 1004-1009 (1997). For example, black, clear-bottomed 96-well microplates (black view plates; Packard Instrument Company, Meriden, Conn.) with the outer perimeter wells filled with sterile water to prevent dehydration in experimental wells. Initial drug dilutions were
20 prepared in dimethyl sulfoxide and subsequent two fold dilutions were performed in 0.1 ml of 7H9GC (no Tween® 80) in the microplates. Frozen inocula were initially diluted 1:20 in BACTEC 12B medium followed by a 1:50 dilution in 7H9GC. Addition of 100 µL to wells resulted in final bacterial titers of between 2.0×10^5 and 5×10^4 CFU/mL for H37Rv and H37Ra, respectively. Wells containing drug only were
25 used to detect autofluorescence of compounds plus additional control wells consisted of bacteria only (B) and medium only (M). Plates were incubated at 37°C. Starting at day 4 of incubation, 20 µL of 10x alamar Blue solution (Alamar Biosciences/Accumed®, Westlake, Ohio) and 12.5 µL of 20% Tween® 80 were added to one B well and one M well, and plates were reincubated at 37°C. Wells were
30 observed at 12 and 24 h for a color change from blue to pink and for a reading of greater than or equal to 50,000 fluorescence units (FU). Fluorescence was measured in a Cytofluor II microplate fluorometer (PerSeptive Biosystems®, Framingham, Mass.)

in bottom-reading mode with excitation at 530 nm and emission at 590 nm. If the B wells became pink by 24 h, reagent was added to the entire plate. If the well remained blue or 50,000 FU was measured, additional M and B wells were tested daily until a color change occurred, at which time reagents were added to all remaining wells.

5 Plates were then incubated at 37°C, and results were recorded at 24 h post-reagent addition. Visual MICs were defined as the lowest concentration of drug that prevented a color change. For fluorometric MICs, a background subtraction was performed on all wells with a mean of triplicate M wells. Percent inhibition was defined as $1 - (\text{test well FU} / \text{mean FU of triplicate B wells}) \times 100$. The lowest drug
10 concentration effecting an inhibition of 90% was considered the MIC.

[0072] Biochemical testing results for exemplary compounds of the invention are provided in FIG. 1.

EXAMPLE 5

Low Oxygen Recovery Assay (LORA)

15 [0073] The low-oxygen recovery assay (LORA) was essentially performed as described by Cho *et al. Antimicrob Agents Chemother* 51: 1380-1385 (2007). A recombinant *M. tuberculosis* H₃₇R_v bearing luxAB on a plasmid, pFCA-luxAB, was used in all the LORA experiments. Frozen aliquots from a low oxygen adapted culture were thawed, diluted in Middlebrook 7H12 broth (Middlebrook 7H9 broth
20 containing 1 mg/mL Casitone, 5.6 µg/mL palmitic acid, 5 mg/mL bovine serum albumin, and 4 µg/ml filter-sterilized catalase), and sonicated for 15s. The cultures were diluted to obtain an A₅₇₀ of 0.03 to 0.05 and 3,000 to 7,000 RLU per 100 µL. This corresponds to 5×10^5 to 2×10^6 CFU/mL. Twofold serial dilutions were prepared in a volume 100 µL in black 96-well microtiter plates, and 100 µL of the cell
25 suspension was added. For LORA, the microplate cultures were placed under anaerobic conditions (oxygen concentration, less than 0.16%) by using an Anoxomat model WS-8080 (MART Microbiology®) and three cycles of evacuation and filling with a mixture of 10% H₂, 5% CO₂, and 85% N₂. An anaerobic indicator strip was placed inside the chamber to visually confirm the removal of oxygen. The plates were
30 incubated at 37°C for 10 days and then transferred to an ambient gaseous condition (5% CO₂-enriched air) incubator for a 28-h “recovery.” On day 11 (after the 28-h aerobic recovery), 100 µL culture was transferred to white 96-well microtiter plates

for determination of luminescence. A 10% solution of n-decanal aldehyde (Sigma®) in ethanol was freshly diluted 10-fold in PBS, and 100 µl was added to each well with an autoinjector. Luminescence was measured in a Victor2 multilabel reader (Perkin-Elmer Life Sciences®) by using a reading time of 1 s. The MIC was defined as the lowest drug concentration effecting growth inhibition of 90% relative to the growth for the drug-free controls.

[0074] Biochemical testing results for exemplary compounds of the invention are provided in FIG. 1.

EXAMPLE 6

10 Tuberculosis in vivo efficacy experiments

[0075] The TB *in vivo* efficacy experiments were essentially performed as described in Lenaerts *et al. Antimicrob Agents Chemother* 47: 783-785 (2003) with a few modifications. A highly susceptible gamma interferon specific pathogen-free C57BL/6-Ifngtm1ts (GKO) mice (Jackson Laboratories®, Bar Harbor, Maine) were exposed to a low-dose aerosol infection with *M. tuberculosis* strain Erdman in a Glas-Col inhalation exposure system as previously described in Kelly *et al. Antimicrob Agents Chemother* 40: 2809-2812 (1996). Every treatment group consisted of five mice for every following time point. Treatment was started 10 days after infection. One control group of infected mice was sacrificed at the start of treatment. A second group of infected but untreated mice was sacrificed after the cessation of treatment at 24 days. C and L were formulated in saline and E was formulated in 50% water/35% PEG400/5%PG, while rifampicin was formulated in 20% cyclodextrin. All compounds were administered by oral gavage. Rifampicin was dosed at 10 mg/kg QD PO. C was dosed at 100 mg/kg BID PO. E was dosed at 100 mg/kg BID PO. L was dosed at 100 mg/kg QD PO. After completion of therapy, the mice were sacrificed by carbon dioxide inhalation. Lungs were aseptically removed and disrupted in a tissue homogenizer. The number of viable organisms was determined by serial dilution of the homogenates on nutrient Middlebrook 7H11 agar plates (GIBCO BRL®, Gaithersburg, Md.). The plates were incubated at 37°C in ambient air for 4 weeks prior to the counting of viable *M. tuberculosis* colonies (CFU).

30 [0076] On day 3, the control group had a mean log₁₀ CFU/lung of 2.83 (0.40). On day 10, the control group had a log₁₀ CFU/lung of 4.81 (0.08). On day 24, the control group had a log₁₀ CFU/lung of 8.96 (0.14). On day 24, the group treated with rifampicin had a log₁₀ CFU/lung of 6.16 (0.10). On day 24, the group treated with C

had a log₁₀ CFU/lung of 5.06 (0.26). On day 24, the group treated with **E** had a log₁₀ CFU/lung of 2.73 (0.05). On day 24, the group treated with **L** had a log₁₀ CFU/lung of 3.08 (0.06).

EXAMPLE 7

5 *Tuberculosis in vivo efficacy experiments*


- [0077] The TB *in vivo* efficacy experiments were essentially performed as described in Lenaerts *et al. Antimicrob Agents Chemother* 47: 783-785 (2003) with a few modifications. A highly susceptible gamma interferon specific pathogen-free C57BL/6-Ifngtm1ts (GKO) mice (Jackson Laboratories®, Bar Harbor, Maine) were exposed to a low-dose aerosol infection with *M. tuberculosis* strain Erdman in a Glas-Col inhalation exposure system as previously described in Kelly *et al. Antimicrob Agents Chemother* 40: 2809-2812 (1996). Every treatment group consisted of five mice for every following time point. Treatment was started 13 days after infection. One control group of infected mice was sacrificed at the start of treatment. A second group of infected but untreated mice was sacrificed after the cessation of treatment at 22 days. **N** was formulated in saline and **E** was formulated in 50% water/35% PEG400/5%PG, while Isoniazid (INH) was formulated in distilled water. All compounds were administered by oral gavage. INH was dosed at 25 mg/kg QD PO. **E** was dosed at 100 mg/kg QD PO. **N** was dosed at 100 mg/kg BID PO. After completion of therapy, the mice were sacrificed by carbon dioxide inhalation. Spleens and lungs were aseptically removed and disrupted in a tissue homogenizer. The number of viable organisms was determined by serial dilution of the homogenates on nutrient Middlebrook 7H11 agar plates (GIBCO BRL®, Gaithersburg, Md.). The plates were incubated at 37°C in ambient air for 4 weeks prior to the counting of viable *M. tuberculosis* colonies (CFU).
- 25 [0078] On day 13, the control group had a mean log₁₀ CFU of 7.02 (0.08) for lungs and mean log₁₀ CFU for spleens of 3.99 (0.21). On day 22, the control group had a log₁₀ CFU for lungs of 7.82 (0.11) and spleens of 6.69 (0.08). On day 22, the group treated with INH had a log₁₀ CFU for lungs of 5.29 (0.13) and for spleens of 4.27 (0.25). On day 22, the group treated with **E** had a log₁₀ CFU for lungs of 5.27 (0.12) and for spleens of 4.27 (0.25). On day 22, the group treated with **N** had a log₁₀ CFU for lungs of 5.51 (0.09) and spleens of 2.42 (0.48).

[0079] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light

thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound having a structure which is:



wherein R^3 is substituted or unsubstituted nitroalkyl or substituted or unsubstituted aminoalkyl;

R^4 is selected from the group consisting of halogen, unsubstituted alkyl, unsubstituted alkoxy, and unsubstituted phenyl;

Y is O or S; and


R^5 is selected from the group consisting of substituted or unsubstituted alkyl and substituted or unsubstituted heteroalkyl;

or a salt, hydrate or solvate thereof

wherein the alkyl groups have 10 or fewer carbon atoms and wherein the heteroalkyl groups include at least one heteroatom selected from the group consisting of B, O, N and S; and


wherein the substituents for the alkyl and heteroalkyl radicals are each selected from the group consisting of: $-R'$, $-OR'$, $=O$, $=NR'$, $=N-OR'$, $-NR'R''$, $-SR'$, $-halogen$, $-SiR'R''R'''$, $OC(O)R'$, $-C(O)R'$, $-CO_2R'$, $-CONR'R''$, $-OC(O)NR'R''$, $-NR''C(O)R'$, $NR' C(O)NR''R'''$, $-NR''C(O)_2R'$, $-NR''''-C(NR'R''R''')=NR''''$, $NR''''C(NR'R'')=NR''''$, $-S(O)R'$, $-S(O)_2R'$, $-S(O)_2NR'R''$, $NR''SO_2R'$, $-CN$, $-NO_2$, $-N_3$, $-CH(Ph)_2$, fluoro(C_1 - C_4)alkoxy, and fluoro(C_1 - C_4)alkyl, in a number ranging from zero to $(2m'+1)$, where m' is the total number of carbon atoms in such radical; wherein R' , R'' , R''' , R'''' and R''''' are each independently hydrogen, unsubstituted heteroalkyl, unsubstituted aryl, unsubstituted alkyl, alkoxy or thioalkoxy groups, or arylalkyl groups.

2. The compound or a salt, hydrate or solvate thereof of claim 1, having a structure which is:



wherein C* is a carbon atom stereocenter which has a configuration which is (R) or (S).

3. The compound or a salt, hydrate or solvate thereof of claim 1, having a structure which is



wherein the C* is a carbon atom stereocenter which has a configuration which is (R) or (S).

4. The compound or a salt, hydrate or solvate thereof of claim 2, wherein the C* stereocenter is in a (S) configuration.
5. The compound or a salt, hydrate or solvate thereof of claim 1, wherein R³ is -CH₂NH₂.
6. The compound or a salt, hydrate or solvate thereof of claim 1, wherein R⁴ is selected from the group consisting of fluorine, chlorine, bromine, and iodine.
7. The compound or a salt, hydrate or solvate thereof of claim 1, wherein Y is O.
8. The compound or a salt, hydrate or solvate thereof of claim 1, wherein R⁵ is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, and sec-butyl.
9. The compound or a salt, hydrate or solvate thereof of claim 1, wherein R³ is -CH₂NH₂; and R⁴ is halogen.

10. The compound or a salt, hydrate or solvate thereof of claim **1**, wherein R³ is -CH₂NH₂; R⁴ is chlorine; Y is O; and R⁵ is substituted or unsubstituted alkyl.

11. A composition comprising:

- a) a first stereoisomer of the compound of claim **2**;
- b) at least one additional stereoisomer of the first stereoisomer;

wherein the first stereoisomer is present in an enantiomeric excess of at least 80% relative to said at least one additional stereoisomer.

12. A combination comprising the compound of any one of claims **1 to 10**, or a pharmaceutically acceptable salt thereof, together with at least one other therapeutically active agent, wherein said one other therapeutically active agent is an antibacterial agent.

13. A pharmaceutical formulation comprising:

- a) the compound of any one of claims **1 to 10**, or a pharmaceutically acceptable salt thereof; and
- b) a pharmaceutically acceptable excipient.

14. Use the compound of any one of claims **1 to 10**, or a salt, hydrate or solvate thereof, for inhibiting a tRNA synthetase.

15. Use of the compound of any one of claims **1 to 10**, or a salt, hydrate or solvate thereof, in the manufacture of a medicament for inhibiting a t-RNA synthetase.

16. Use the compound of any one of claims **1 to 10**, or a salt, hydrate or solvate thereof, for killing and/or preventing the growth of a microorganism.

17. Use of the compound of any one of claims **1 to 10**, or a salt, hydrate or solvate thereof, in the manufacture of a medicament for killing and/or preventing the growth of a microorganism.

- 18.** Use of the compound of any one of claims **1 to 10**, or a pharmaceutically-acceptable salt thereof, for treating and/or preventing a bacterial-associated disease.
- 19.** Use of the compound of any one of claims **1 to 10**, or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for treating and/or preventing a bacterial-associated disease.
- 20.** The use-of claim **18** or claim **19**, wherein the disease is tuberculosis.
- 21.** The use-of claim **18** or claim **19** or claim **20**, wherein the animal is a human.
- 22.** Use of the compound of any one of claims **1 to 10**, or a pharmaceutically-acceptable salt thereof, for inhibiting the editing domain of a t-RNA synthetase.
- 23.** Use of the compound of any one of claims **1 to 10**, or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for inhibiting the editing domain of a t-RNA synthetase.
- 24.** The use of a compound of any one of claims **1 to 10** or a combination of claim **12** or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment and/or prophylaxis of bacterial infection.

Figure 1

Compound #	A. baumannii ATCC 17978 MIC	E. coli ATCC 25922 MIC	H. influenzae ATCC 49766 MIC	K. pneumoniae ATCC 13883 MIC	P. aeruginosa ATCC 27853 MIC	MABA M. tuberculosis H37Rv MIC	LORA M. tuberculosis H37Rv MIC	M. tuberculosis LeuRS IC50 (uM)
A	16	6	3	6	20	0.37	>64	0.97
B	>64	>64	2	>64	>64	<0.025	>64	0.13
C	8	4	1	4	>64	0.04	0.52	0.13
D	>64	>64	0.5	>64	>64	<0.02	0.07	0.09
E	>64	>64	1	>64	>64	0.05	0.70	0.07
F	>64	>64	NT	>64	>64	<0.2	26.20	0.2
G	8	16	NT	8	>64	<0.2	31.90	0.31
H	>64	>64	NT	>64	>64	16.80	48.46	27.6
I	>64	>64	NT	>64	>64	0.30	<0.391	0.04
J	>64	>64	NT	>64	>64	0.59	0.69	0.03
K	>64	>64	>64	>64	>64	>64	NT	NT
L	>64	64	<0.12	32	>64	<0.004	<0.039	0.02
M	>64	>64	NT	>64	>64	NT	NT	21.02
N	1.5	0.5	NT	1	3	NT	NT	0.09
O	32	0.74	8	8	64	0.28	>64	0.91
P	32	NT	4	8	32	1.91	>64	0.896
Q	32	1.1	16	8	64	3.73	>64	0.878
R	>64	1.6	16	8	64	1.01	>64	0.892
S	>64	1.1	8	8	64	0.88	>50	0.61
T	16	1.345292	16	8	16	NT	NT	0.895
U	16	1.231968	4	8	>64	0.33	5.91	NT
V	>64	>64	32	64	>64	NT	NT	0.114

