

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 2185197 T3**

-
- (51) Int.Cl.: **A 61 K 38/55 (2006.01)** **A 61 K 38/18 (2006.01)** **A 61 K 38/19 (2006.01)**
A 61 P 17/02 (2006.01) **C 12 N 5/02 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-12-10**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-10-17**
- (86) Europæisk ansøgning nr.: **08827981.5**
- (86) Europæisk indleveringsdag: **2008-08-08**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-05-19**
- (86) International ansøgning nr.: **US2008009500**
- (87) Internationalt publikationsnr.: **WO2009025730**
- (30) Prioritet: **2007-08-22 US 965707 P** **2008-04-30 US 125960**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Noveome Biotherapeutics, Inc., 100 Technology Drive, Suite 400, Pittsburgh, PA 15219, USA**
- (72) Opfinder: **SMITH, Charlotte, A., 805 Westchester Drive, Wexford, PA 15090, USA**
SING, George, L., 525 E. 80th Street, New York, NY 10021, USA
PALLADINO, Linda, O., 45 Honeysuckle Ct., Stormville, NY 12582, USA
MARSHALL, Vivienne S., 14302 Fox Fire Lane, San Antonio, Texas 78231, USA
TRUMPOWER, Catherine, J., 260 Cascade Road, Pittsburgh, PA 15221, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Hidtil ukendte cellefaktorholdige opløsningssammensætninger og anvendelse heraf til sårbehandling**
- (56) Fremdragne publikationer:
WO-A1-89/05656
WO-A1-2007/039109
WO-A2-2006/138468
US-A- 5 612 028
US-A1- 2003 191 061
US-A1- 2004 166 100
US-A1- 2004 228 853
US-A1- 2007 148 214
GOLOVNYOV ET AL: "Peculiarities of hepatic experimental wound healing in rats treated with angiogenin", MORFOLOGIA, GIPPOKRAT, SANKT PETERBURG, RU, vol. 128, no. 4, 1 January 2005 (2005-01-01), pages 98-100, XP009153008, ISSN: 1026-3543

DESCRIPTION

Field of the Invention

[0001] The field of the invention is directed to novel cellular factor-containing solution compositions (referred to herein as "CFS" compositions), including novel sustained-release cellular factor-containing solution compositions (referred to herein as "SR-CFS" compositions), methods of making such novel compositions and uses thereof.

Background of the Invention

[0002] Many individual cytokines and growth factors have been evaluated for their therapeutic utility in the treatment of many varied diseases, disorders and injuries. Unfortunately, the results have been only partially encouraging. For example, PDGF-BB has proven to be useful in the treatment of diabetic foot ulcers; GM-CSF is marketed in Europe for both venous ulcers and diabetic foot ulcers; and HGH (human growth hormone) is marketed in the US for pediatric burns. Failures include BDNF, CNTF and IGF-1 which have all been evaluated in clinical trials designed to test their efficacy in treating ALS, each with disappointing results; TGF β 2 was unsuccessful in a phase 2 study for venous ulcers; and IGF-1 and PDGF combination therapy was unsuccessful in diabetic foot ulcers.

[0003] While it is not clear why so many of these individual cytokines and growth factors have failed in the clinic, one theory is that the proteins were being administered in doses that were not physiologic, i.e. very high doses compared to the physiologic levels normally found in vivo. Also, because of the complex interaction between cytokines and growth factors in a given physiological niche, the application of just one factor, especially one at abnormally high levels, cannot recreate the physiological niche and may, in fact, grossly disturb its delicate balance.

[0004] Compounding their limited success in the clinic, cytokines and growth factors and other protein-based therapeutics are typically more difficult to administer to patients than other pharmaceuticals. Because the efficacy of a protein is related to its shape, protein-based therapeutics cannot be subjected to conditions that could cause the unfolding, or denaturing, of the protein or proteins contained therein. Consequently, special care is necessary in the preparation, storage, and administration of protein-based therapeutics.

[0005] In addition to avoiding any denaturation of the protein, it is often desirable to be able to control the amount of the protein administered to a patient over time. This helps to avoid protein concentrations within the patient that are undesirably high or low or that fluctuate too much from a desired level, and instead helps maintain a steady level of the therapeutic in the patient. To address this, sustained-release formulations for many therapeutics, including protein-based therapeutics, have been or are currently in development. Sustained-release

protein-based therapeutics can be administered by a variety of methods, including but not limited to oral delivery of tablets or capsules, inhalation of powders, implantation, incorporation into a matrix, or topical application of an encapsulated therapeutic from which the protein is gradually released over time.

[0006] The preparation of such sustained-release formulations are varied. One process includes mixing the protein with an organic solvent. For example, a powder formulation may be made by spraying a mixture of the protein and an organic solvent into liquid nitrogen. Another process involves mixing the protein with a solution of a bioerodible/biodegradable polymer in an organic solvent, resulting in the formation of microparticles which contain the protein and the polymer by coagulation of the mixture. In yet another process, proteins, powdered formulations, or microparticles can be mixed with an organic solvent to produce a liquid or gel which may be injected into a patient or applied topically. Unfortunately, drawbacks to using organic solvents are their tendency to cause protein denaturation.

[0007] Additives have been used to stabilize proteins in the presence of a denaturing organic solvent. These additives include surfactants (see U.S. Pat. No. 5,096,885), amino acids (see U.S. Pat. No. 4,297,344), polyols (see U.S. Pat. No. 5,589,167), natural polymers (see WO 8903671), synthetic polymers (see Pharm. Res. 8:285 291, 1991), and metals (see U.S. Pat. No. 6,191,107 B1).

[0008] To date, no protein-based therapeutic agent (i.e. cytokines and growth factors) is available that effectively recreates or mimics the complex combination and physiologic levels of physiologically relevant cytokines and growth factors found naturally in the body in healthy and disease or injury states. Every protein-based therapeutic currently available administers a dose many, many times higher than the levels that the cytokines or growth factors are normally found in the body. In addition, no one has yet been able to administer these physiologically relevant cytokines and growth factors at physiological levels. Further, no one has yet been able to administer these physiologically relevant cytokines and growth factors at physiological levels in a sustained-release formulation. Therefore, Applicants present herewith for the first time the instant invention whose object is to satisfy the unmet medical need of providing physiologically relevant growth factors and cytokines at physiologic levels (CFS compositions), and in some instances, delivering those physiologically relevant growth factors and cytokines at physiologic levels using a sustained-release formulation (SR-CFS compositions).

Brief Summary of the Invention

[0009] It is an object of the instant invention to provide novel cellular factor-containing solution (CFS) compositions that recreate the complex and unique combination and physiologic levels of such cytokines and growth factors found in biological niches. It is further an object of the instant invention to provide novel sustained-release cellular factor-containing solution (SR-CFS) compositions that contain the complex and unique combination and physiologic levels of the cytokines and growth factors found naturally in biological niches. Because the cellular

factors are present in levels comparable to physiological levels found in the body, they are optimal for use in therapeutic applications which require intervention to support, initiate, replace, accelerate or otherwise influence biochemical and biological processes involved in the treatment and/or healing of disease and/or injury. In the case of the SR-CFS compositions, the cellular factors are released slowly over time to provide a continual, consistent physiologic level of such factors to optimize healing and/or recovery.

[0010] In addition to the novel CFS compositions described herein, it is also an object of the invention to provide methods for making such novel CFS compositions, including pooling cell-derived compositions (i.e. pooled-ECS and pooled-ACCS compositions), and SR-CFS compositions, as well as therapeutic uses thereof.

[0011] The pooled cell-derived compositions possess several important properties and characteristics including decreased variability in the levels of physiologically relevant cellular factors necessary for therapeutic effect as compared to non-pooled compositions. The cellular factors are present in levels comparable to physiological levels found in the body and are thus optimal for use in therapeutic applications which require intervention to support, initiate, replace, accelerate or otherwise influence biochemical and biological processes involved in the treatment and/or healing of disease and/or injury. The novel methods described herein of pooling cell-derived compositions to decrease variability has the effect of optimizing levels of the secreted factors such that their full therapeutic potential can be achieved in every pool. In addition to the therapeutic value of such pooled compositions, the method of pooling samples to decrease non-pooled composition-to-composition variability has the significant commercial advantages of increasing production yields by minimizing non-pooled composition rejection for failure to meet product specifications and, consequently, decreasing production costs and increasing revenues.

[0012] Accordingly, a first aspect of the invention is an amnion-derived cellular cytokine solution composition comprising physiologic concentrations of VEGF, TGF β 2, Angiogenin, PDGF, TIMP-1 and TIMP-2, wherein the physiologic concentration is ~5.0-16ng/mL for VEGF, ~3.5-4.5ng/mL for Angiogenin, ~100-165pg/mL for PDGF, ~2.5-2.7ng/mL for TGF β 2, ~0.68 μ g/mL for TIMP-1 and ~1.04 μ g/mL for TIMP-2, wherein the amnion-derived cellular cytokine solution composition is obtainable by the method of:

1. (a) isolating amnion epithelial cells from the amnion of a placenta;
2. (b) selecting AMP cells from the amnion epithelial cells;
3. (c) culturing the AMP cells in animal-free based medium, wherein the medium is supplemented with clinical grade human albumin added up to a concentration of 10% until they reach confluence;
4. (d) changing the culture medium;
5. (e) culturing the cells in the medium for 1, 2, 3, 4, 5 or 6 days; and
6. (f) collecting the culture medium step (e) to obtain amnion-derived cellular cytokine solution;

wherein the medium of step (c) is Stemline I or II, Opti-pro, IMDM or DMEM.

[0013] Aspect two of the invention is the composition of the first aspect wherein step (f) is repeated a plurality of times and the amnion-derived cellular cytokine solution obtained in step (f) is combined to create a pooled amnion-derived cellular cytokine solution.

[0014] Aspect three of the invention is the composition of part (c) of the first aspect, wherein the medium is serum free.

[0015] Aspect four of the invention is the composition of any one of the first, second or third aspect, wherein the basal medium is further supplemented with EGF at a concentration of between 0 and 1µg/ml.

[0016] Aspect five of the invention is a sustained-release composition comprising the amnion-derived cellular cytokine solution of any one of the first or second aspect.

[0017] Aspect six of the invention is a method of making a sustained-release composition of the fifth aspect, comprising the steps of combining a cellular factor-containing solution composition with an agent capable of effecting sustained-release of the cellular factor-containing solution composition.

[0018] Aspect seven of the invention is a pharmaceutical composition comprising the composition of any one of the first to fifth aspect.

[0019] Aspect eight of the invention is a composition of any one of the first to fifth aspect or seventh aspect, for use in a method of accelerating wound healing.

Definitions

[0020] As defined herein "isolated" refers to material removed from its original environment and is thus altered "by the hand of man" from its natural state.

[0021] As used herein, the term "protein marker" means any protein molecule characteristic of the plasma membrane of a cell or in some cases of a specific cell type.

[0022] As used herein, "enriched" means to selectively concentrate or to increase the amount of one or more materials by elimination of the unwanted materials or selection and separation of desirable materials from a mixture (i.e. separate cells with specific cell markers from a heterogeneous cell population in which not all cells in the population express the marker).

[0023] As used herein, the term "substantially purified" means a population of cells substantially homogeneous for a particular marker or combination of markers. By substantially homogeneous is meant at least 90%, and preferably 95% homogeneous for a particular marker or combination of markers.

[0024] The term "placenta" as used herein means both preterm and term placenta.

[0025] As used herein, the term "totipotent cells" shall have the following meaning. In mammals, totipotent cells have the potential to become any cell type in the adult body; any cell type(s) of the extraembryonic membranes (e.g., placenta). Totipotent cells are the fertilized egg and approximately the first 4 cells produced by its cleavage.

[0026] As used herein, the term "pluripotent stem cells" shall have the following meaning. Pluripotent stem cells are true stem cells with the potential to make any differentiated cell in the body, but cannot contribute to making the components of the extraembryonic membranes which are derived from the trophoblast. The amnion develops from the epiblast, not the trophoblast. Three types of pluripotent stem cells have been confirmed to date: Embryonic Stem (ES) Cells (may also be totipotent in primates), Embryonic Germ (EG) Cells, and Embryonic Carcinoma (EC) Cells. These EC cells can be isolated from teratocarcinomas, a tumor that occasionally occurs in the gonad of a fetus. Unlike the other two, they are usually aneuploid.

[0027] As used herein, the term "multipotent stem cells" are true stem cells but can only differentiate into a limited number of types. For example, the bone marrow contains multipotent stem cells that give rise to all the cells of the blood but may not be able to differentiate into other cells types.

[0028] As used herein, the term "extraembryonic tissue" means tissue located outside the embryonic body which is involved with the embryo's protection, nutrition, waste removal, etc. Extraembryonic tissue is discarded at birth. Extraembryonic tissue includes but is not limited to the amnion, chorion (trophoblast and extraembryonic mesoderm including umbilical cord and vessels), yolk sac, allantois and amniotic fluid (including all components contained therein). Extraembryonic tissue and cells derived therefrom have the same genotype as the developing embryo.

[0029] As used herein, the term "extraembryonic cytokine secreting cells" or "ECS cells" means a population of cells derived from the extraembryonic tissue which have the characteristics of secreting a unique combination of physiologically relevant cytokines in a physiologically relevant temporal manner into the extracellular space or into surrounding culture media and which have not been cultured in the presence of any animal-derived products, making them and cell products derived from them suitable for human clinical use. In one embodiment, the ECS cells secrete at least one cytokine selected from VEGF, Angiogenin, PDGF and TGF β 2 and at least one MMP inhibitor. In a specific embodiment, the MMP inhibitor is selected from TIMP-1 and TIMP-2. In another embodiment, the ECS cells secrete more than one cytokine selected from VEGF, Angiogenin, PDGF and TGF β 2 and more than one MMP inhibitor. In a specific embodiment, the MMP inhibitor is selected from TIMP-1 and TIMP-2. In a preferred embodiment, the ECS cells secrete the cytokines VEGF, Angiogenin, PDGF and TGF β 2 and the MMP inhibitors TIMP-1 and/or TIMP-2. The physiological range of the cytokine

or cytokines in the unique combination is as follows: ~5-16ng/mL for VEGF, ~3.5-4.5 ng/mL for Angiogenin, ~100-165pg/mL for PDGF, ~2.5-2.7ng/mL for TGF β 2, ~0.68 μ g/mL for TIMP-1 and ~1.04 μ g/mL for TIMP-2. The ECS cells may optionally express Thymosin β 4. ECS cells may be selected from populations of cells and compositions described in this application and in US2003/0235563, US2004/0161419, US2005/0124003, U.S. Provisional Application Nos. 60/666,949, 60/699,257, 60/742,067, 60/813,759, U.S. Application No. 11/333,849, U.S. Application No. 11/392,892, PCTUS06/011392, US2006/0078993, PCT/US00/40052, U.S. Patent No. 7,045,148, US2004/0048372, and US2003/0032179.

[0030] As used herein, the term "amnion-derived multipotent progenitor cell" or "AMP cell" means a specific population of ECS cells that are epithelial cells derived from the amnion. In addition to the characteristics described above for ECS cells, AMP cells have the following characteristics. They have not been cultured in the presence of any animal-derived products, making them and cell products derived from them suitable for human clinical use. They grow without feeder layers, do not express the protein telomerase and are non-tumorigenic. AMP cells do not express the hematopoietic stem cell marker CD34 protein. The absence of CD34 positive cells in this population indicates the isolates are not contaminated with hematopoietic stem cells such as umbilical cord blood or embryonic fibroblasts. Virtually 100% of the cells react with antibodies to low molecular weight cytokeratins, confirming their epithelial nature. Freshly isolated AMP cells will not react with antibodies to the stem/progenitor cell markers c-kit (CD117) and Thy-1 (CD90). Several procedures used to obtain cells from full term or pre-term placenta are known in the art (see, for example, US 2004/0110287; Anker et al., 2005, Stem Cells 22:1338-1345; Ramkumar et al., 1995, Am. J. Ob. Gyn. 172:493-500). However, the methods used herein provide improved compositions and populations of cells. AMP cells have previously been described as "amnion-derived cells" (see U.S. Provisional Application Nos. 60/666,949, 60/699,257, 60/742,067, U.S. Provisional Application Nos. 60/813,759, U.S. Application No. 11/333,849, U.S. Application No. 11/392,892, and PCTUS06/011392).

[0031] By the term "animal-free" when referring to certain compositions, growth conditions, culture media, etc. described herein, is meant that no animal-derived materials, such as animal-derived serum, other than clinical grade human materials, such as recombinantly produced human proteins, are used in the preparation, growth, culturing, expansion, storage or formulation of the certain composition or process.

[0032] By the term "serum-free" when referring to certain compositions, growth conditions, culture media, etc. described herein, is meant that no animal-derived serum (i.e. no non-human) is used in the preparation, growth, culturing, expansion, storage or formulation of the certain composition or process.

[0033] By the term "expanded", in reference to cell compositions, means that the cell population constitutes a significantly higher concentration of cells than is obtained using previous methods. For example, the level of cells per gram of amniotic tissue in expanded compositions of AMP cells is at least 50 and up to 150 fold higher than the number of cells in the primary culture after 5 passages, as compared to about a 20 fold increase in such cells

using previous methods. In another example, the level of cells per gram of amniotic tissue in expanded compositions of AMP cells is at least 30 and up to 100 fold higher than the number of cells in the primary culture after 3 passages. Accordingly, an "expanded" population has at least a 2 fold, and up to a 10 fold, improvement in cell numbers per gram of amniotic tissue over previous methods. The term "expanded" is meant to cover only those situations in which a person has intervened to elevate the number of the cells.

[0034] As used herein, the term "passage" means a cell culture technique in which cells growing in culture that have attained confluence or are close to confluence in a tissue culture vessel are removed from the vessel, diluted with fresh culture media (i.e. diluted 1:5) and placed into a new tissue culture vessel to allow for their continued growth and viability. For example, cells isolated from the amnion are referred to as primary cells. Such cells are expanded in culture by being grown in the growth medium described herein. When such primary cells are subcultured, each round of subculturing is referred to as a passage. As used herein, "primary culture" means the freshly isolated cell population.

[0035] As used herein, "conditioned medium" is a medium in which a specific cell or population of cells has been cultured, and then removed. When cells are cultured in a medium, they may secrete cellular factors that can provide support to or affect the behavior of other cells. Such factors include, but are not limited to hormones, cytokines, extracellular matrix (ECM), proteins, vesicles, antibodies, chemokines, receptors, inhibitors and granules. The medium containing the cellular factors is the conditioned medium. Examples of methods of preparing conditioned media are described in U.S. Patent 6,372,494. As used herein, conditioned medium also refers to components, such as proteins, that are recovered and/or purified from conditioned medium or from ECS cells, including AMP cells.

[0036] As used herein, the term "cellular factor-containing solution" or "CFS" composition means a composition having physiologic concentrations of one or more factors selected from VEGF, Angiogenin, PDGF and TGF β 2 and at least one MMP inhibitor. Examples of suitable MMP inhibitors include but are not limited to TIMP-1 and TIMP-2. CFS compositions include conditioned media derived from ECS cells, amnion-derived cellular cytokine solution compositions (see definition below), physiologic cytokine solution compositions (see definition below), and sustained release formulations of such CFS compositions.

[0037] As used herein, the term "amnion-derived cellular cytokine solution" or "ACCS" means conditioned medium that has been derived from AMP cells or expanded AMP cells. Amnion-derived cellular cytokine solution or ACCS has previously been referred to as "amnion-derived cytokine suspension".

[0038] As used herein, the term "physiologic cytokine solution" or "PCS" composition means a composition which is not cell-derived and which has physiologic concentrations of one or more factors selected from VEGF, Angiogenin, PDGF and TGF β 2 and at least one MMP inhibitor. Examples of suitable MMP inhibitors include but are not limited to TIMP-1 and TIMP-2.

[0039] As used herein, the term "suspension" means a liquid containing dispersed components, i.e. cytokines. The dispersed components may be fully solubilized, partially solubilized, suspended or otherwise dispersed in the liquid. Suitable liquids include, but are not limited to, water, osmotic solutions such as salt and/or sugar solutions, cell culture media, and other aqueous or non-aqueous solutions.

[0040] The term "lysate" as used herein refers to the composition obtained when cells, for example, AMP cells, are lysed and optionally the cellular debris (e.g., cellular membranes) is removed. This may be achieved by mechanical means, by freezing and thawing, by sonication, by use of detergents, such as EDTA, or by enzymatic digestion using, for example, hyaluronidase, dispase, proteases, and nucleases.

[0041] The term "physiologic" or "physiological level" as used herein means the level that a substance in a living system is found and that is relevant to the proper functioning of a biochemical and/or biological process.

[0042] As used herein, the term "pooled" means a plurality of compositions that have been combined to create a new composition having more constant or consistent characteristics as compared to the non-pooled compositions. For example, pooled ACCS has more constant or consistent characteristics compared to non-pooled ACCS. Examples of pooled compositions include "SP pools" (more than one ACCS collection/one placenta), "MP1 pools" (one ACCS collection/placenta, multiple placentas), and "MP2 pools" (more than one ACCS collection/placenta, multiple placentas).

[0043] As used herein, the term "substrate" means a defined coating on a surface that cells attach to, grown on, and/or migrate on. As used herein, the term "matrix" means a substance that cells grow in or on that may or may not be defined in its components. The matrix includes both biological and non-biological substances. As used herein, the term "scaffold" means a three-dimensional (3D) structure (substrate and/or matrix) that cells grow in or on. It may be composed of biological components, synthetic components or a combination of both. Further, it may be naturally constructed by cells or artificially constructed. In addition, the scaffold may contain components that have biological activity under appropriate conditions.

[0044] The term "cell product" or "cell products" as used herein refers to any and all substances made by and secreted from a cell, including but not limited to, protein factors (i.e. growth factors, differentiation factors, engraftment factors, cytokines, morphogens, proteases (i.e. to promote endogenous cell delamination, protease inhibitors), extracellular matrix components (i.e. fibronectin, etc.).

[0045] The term "therapeutically effective amount" means that amount of a therapeutic agent necessary to achieve a desired physiological effect (i.e. accelerate wound healing).

[0046] As used herein, the term "pharmaceutically acceptable" means that the components, in addition to the therapeutic agent, comprising the formulation, are suitable for administration to

the patient being treated in accordance with the present invention.

[0047] As used herein, the term "therapeutic component" means a component of the composition which exerts a therapeutic benefit when the composition is administered to a subject.

[0048] As used herein, the term "therapeutic protein" includes a wide range of biologically active proteins including, but not limited to, growth factors, enzymes, hormones, cytokines, inhibitors of cytokines, blood clotting factors, peptide growth and differentiation factors.

[0049] As used herein, the term "tissue" refers to an aggregation of similarly specialized cells united in the performance of a particular function.

[0050] As used herein, the terms "a" or "an" means one or more; at least one.

[0051] As used herein, the term "adjunctive" means jointly, together with, in addition to, in conjunction with, and the like.

[0052] As used herein, the term "co-administer" can include simultaneous or sequential administration of two or more agents.

[0053] As used herein, the term "agent" means an active agent or an inactive agent. By the term "active agent" is meant an agent that is capable of having a physiological effect when administered to a subject. Non-limiting examples of active agents include growth factors, cytokines, antibiotics, cells, conditioned media from cells, etc. By the term "inactive agent" is meant an agent that does not have a physiological effect when administered. Such agents may alternatively be called "pharmaceutically acceptable excipients". Non-limiting examples include time release capsules and the like.

[0054] The terms "parenteral administration" and "administered parenterally" are art-recognized and refer to modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid, intraspinal, epidural, intracerebral and intrasternal injection or infusion.

[0055] The terms "sustained-release", "extended-release", "time-release", "controlled-release", or "continuous-release" as used herein means an agent, typically a therapeutic agent or drug, that is formulated to dissolve slowly and be released over time.

[0056] The terms "bioerodable" or "bioerosion" as used herein mean a combination of physical (i.e. dissolution) and chemical (i.e. chemical bond cleavage) processes that result in the breakdown of a substance.

[0057] The term "biodegradable" or "biodegradation" as used herein means a biological agent (i.e. an enzyme, microbe or cell) is responsible for the breakdown of a substance.

[0058] The terms "bioresorbable" or "bioabsorbable" as used herein mean the removal of a breakdown product by cellular activity (i.e. phagocytosis).

[0059] As used herein, the term "nanoparticle" means particles of less than 100nm in diameter that exhibit new or enhanced size-dependent properties compared with larger particles of the same material.

[0060] "Treatment," "treat," or "treating," as used herein covers any treatment of a disease or condition of a mammal, particularly a human, and includes: (a) preventing the disease or condition from occurring in a subject which may be predisposed to the disease or condition but has not yet been diagnosed as having it; (b) inhibiting the disease or condition, i.e., arresting its development; (c) relieving and or ameliorating the disease or condition, i.e., causing regression of the disease or condition; or (d) curing the disease or condition, i.e., stopping its development or progression. The population of subjects treated by the compositions of the invention includes subjects suffering from the undesirable condition or disease, as well as subjects at risk for development of the condition or disease.

[0061] As used herein, a "wound" is any disruption, from whatever cause, of normal anatomy (internal and/or external anatomy) including but not limited to traumatic injuries such as mechanical (i.e. contusion, penetrating), thermal, chemical, electrical, radiation, concussive and incisional injuries; elective injuries such as operative surgery and resultant incisional hernias, fistulas, etc.; acute wounds, chronic wounds, infected wounds, and sterile wounds, as well as wounds associated with disease states (i.e. ulcers caused by diabetic neuropathy or ulcers of the gastrointestinal or genitourinary tract). A wound is dynamic and the process of healing is a continuum requiring a series of integrated and interrelated cellular processes that begin at the time of wounding and proceed beyond initial wound closure through arrival at a stable scar. These cellular processes are mediated or modulated by humoral substances including but not limited to cytokines, lymphokines, growth factors, and hormones. In accordance with the subject invention, "wound healing" refers to improving, by some form of intervention, the natural cellular processes and humoral substances of tissue repair such that healing is faster, and/or the resulting healed area has less scarring and/or the wounded area possesses tissue strength that is closer to that of uninjured tissue and/or the wounded tissue attains some degree of functional recovery.

[0062] As used herein the term "standard animal model for wound healing" refers to any art-accepted animal model for wound healing in which the compositions of the invention exhibit efficacy as measured by accelerated wound healing. Non-limiting examples of suitable models are described in Hayward PG, Robson MC: Animal models of wound contraction. In Barbul A, et al: Clinical and Experimental Approaches to Dermal and Epidermal Repair: Normal and Chronic Wounds. John Wiley & Sons, New York, 1990; DelBecarro, et al: The use of specific thromboxane inhibitors to preserve the dermal microcirculation after burning. Surgery 87: 137-

141, 1980; Robson, et al: Increasing dermal perfusion after burning by decreasing thromboxane production. J Trauma 20: 722-725, 1980; Polo, et al: An in vivo model of human proliferative scar. J Surg Res 74: 187-195, 1998.). Skilled artisans are aware of other suitable models.

Detailed Description

[0063] In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook et al, 2001, "Molecular Cloning: A Laboratory Manual"; Ausubel, ed., 1994, "Current Protocols in Molecular Biology" Volumes I-III; Celis, ed., 1994, "Cell Biology: A Laboratory Handbook" Volumes I-III; Coligan, ed., 1994, "Current Protocols in Immunology" Volumes I-III; Gait ed., 1984, "Oligonucleotide Synthesis"; Hames & Higgins eds., 1985, "Nucleic Acid Hybridization"; Hames & Higgins, eds., 1984, "Transcription And Translation"; Freshney, ed., 1986, "Animal Cell Culture"; IRL Press, 1986, "Immobilized Cells And Enzymes"; Perbal, 1984, "A Practical Guide To Molecular Cloning."

[0064] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range is disclosed. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also disclosed, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also disclosed.

[0065] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

[0066] It must be noted that as used herein and in the appended claims, the singular forms "a," "and" and "the" include plural references unless the context clearly dictates otherwise.

Obtaining and Culturing of Cells

[0067] ECS cells- Various methods for isolating cells from the extraembryonic tissue, which may then be used to produce the ECS cells of the instant invention are described in the art (see, for example, US2003/0235563, US2004/0161419, US2005/0124003, U.S. Provisional Application Nos. 60/666,949, 60/699,257, 60/742,067, 60/813,759, U.S. Application No.

11/333,849, U.S. Application No. 11/392,892, PCTUS06/011392, US2006/0078993, PCT/US00/40052, U.S. Patent No. 7,045,148, US2004/0048372, and US2003/0032179).

[0068] Identifying ECS cells - Once extraembryonic tissue is isolated, it is necessary to identify which cells in the tissue have the characteristics associated with ECS cells (see definition above). For example, cells are assayed for their ability to secrete a unique combination of cytokines into the extracellular space or into surrounding culture medium. Suitable cells are those in which the cytokine or cytokines occurs in the physiological range of ~5.0-16ng/mL for VEGF, ~3.5-4.5 ng/mL for Angiogenin, ~100-165pg/mL for PDGF, ~2.5-2.7ng/mL for TGF β 2, ~0.68 μ g/mL for TIMP-1 and ~1.04 μ g/mL for TIMP-2. Suitable cells may optionally secrete Thymosin β 4.

[0069] AMP cells- In a particular embodiment, AMP cell compositions are prepared using the steps of a) recovery of the amnion from the placenta, b) dissociation of the amnion epithelial cells from the amniotic membrane, c) isolating AMP cells from the amnion epithelial cells, d) culturing of the AMP cells in a basal medium with the addition of a naturally derived or recombinantly produced human protein; and optionally d) further proliferation of the cells using additional additives and/or growth factors. Details are contained in US Publication No. 2006-0222634-A1.

[0070] AMP cells are cultured as follows: The AMP cells are cultured in a basal medium. Such medium includes, but is not limited to, Epilife (Cascade Biologicals), Opti-pro, VP-SFM, IMDM, Advanced DMEM, K/O DMEM, 293 SFM II (all made by Gibco; Invitrogen), HPGM, Pro 293S-CDM, Pro 293A-CDM, UltraMDCK, UltraCulture (all made by Cambrex), Stemline I and Stemline II (both made by Sigma-Aldrich), DMEM, DMEM/F-12, Ham's F12, M199, and other comparable basal media. Such media may optionally contain clinical grade human protein or be supplemented with human clinical grade protein. As used herein a "human protein" is one that is produced naturally or one that is produced using recombinant technology. In certain specific embodiments, "human protein" also is meant to include a human fluid or derivative or preparation thereof, such as human serum or amniotic fluid, which contains human protein.

[0071] In a most preferred embodiment, the cells are cultured using a system that is free of animal products to avoid xeno-contamination. In this embodiment, the culture medium is Stemline I or II, Opti-pro, IMDM or DMEM, and is optionally supplemented with clinical grade human albumin added up to concentrations of 10%. In preferred embodiments, the medium is serum-free in addition to being animal-free.

[0072] Optionally, other factors are used. In one disclosure, epidermal growth factor (EGF) at a concentration of between 0-1 μ g/mL is used. In a preferred disclosure, the EGF concentration is around 10ng/mL. Alternative growth factors which may be used include, but are not limited to, TGF α or TGF β (5ng/mL; range 0.1-100ng/mL), activin A, cholera toxin (preferably at a level of about 0.1 μ g/mL; range 0-10 μ g/mL), transferrin (5 μ g/mL; range 0.1-100 μ g/mL), fibroblast growth factors (bFGF 40ng/mL (range 0-200ng/mL), aFGF, FGF-4, FGF-8; (all in range 0-200ng/mL), bone morphogenic proteins (i.e. BMP-4) or other growth factors known to

enhance cell proliferation. All supplements are clinical grade.

Generation of Cellular Factor-Containing Solution Conditioned Medium

[0073] Generation of ECS conditioned medium- is obtained as described below for ACCS, except that ECS cells are used.

[0074] Generation of ACCS - The AMP cells of the invention can be used to generate ACCS. In one disclosure, the AMP cells are isolated as described herein and 1×10^6 cells/mL are seeded into T75 flasks containing between 5-30mL culture medium, preferably between 10-25mL culture medium, and most preferably about 10mL culture medium. The cells are cultured until confluent, the medium is changed and in one embodiment the ACCS is collected 1 day post-confluence. In another embodiment the medium is changed and ACCS is collected 2 days post-confluence. In another embodiment the medium is changed and ACCS is collected 4 days post-confluence. In another embodiment the medium is changed and ACCS is collected 5 days post-confluence. In a preferred embodiment the medium is changed and ACCS is collected 3 days post-confluence. In another preferred embodiment the medium is changed and ACCS is collected 3, 4, 5, 6 or more days post-confluence. Skilled artisans will recognize that other means for collecting ACCS from AMP cell cultures, such as using other tissue culture vessels, including but not limited to cell factories, flasks, hollow fibers, or suspension culture apparatus, or collecting ACCS from sub-confluent and/or actively proliferating cultures, are also contemplated. It is also contemplated that the ACCS be cryopreserved following collection. It is also contemplated that ACCS be lyophilized following collection. It is also contemplated by the invention that ACCS be formulated for sustained-release following collection. It is also contemplated that ACCS production be scaled up for generation of sufficient product for clinical testing and for commercialization. Skilled artisans are familiar with cryopreservation lyophilization, and sustained-release formulation methodologies.

[0075] It is also contemplated that CFS compositions such as ACCS and pooled ACCS, be diluted with appropriate diluent prior to use. Appropriate diluents include, without limitation, normal saline, PBS, lactated Ringer's solution, cell culture media, conditioned cell culture media, water, and the like. Such dilutions may be 1:2, 1:3, 1:4, 1:5, 1:10, 1:100, etc. In addition, dilutions may be less than 1:2 (i.e. 1:1, 1:0.5, etc.) The appropriate dilution required will be dependent upon the intended use and therefore will need to be empirically determined by skilled artisans.

[0076] The CFS compositions of the invention, including ACCS, and pooled ACCS, are characterized by assaying for physiologically relevant cytokines in the physiologically relevant range of ~5-16ng/mL for VEGF, ~3.5-4.5 ng/mL for Angiogenin, ~100-165pg/mL for PDGF, ~2.5-2.7ng/mL for TGF β 2, ~0.68 μ g/mL for TIMP-1 and ~1.04 μ g/mL for TIMP-2. ACCS and pooled ACCS are optionally assayed for the presence of Thymosin β 4.

Generation of Physiologic Cytokine Solution (PCS) Compositions

[0077] A non-cellular derived form of ACCS (referred to herein as Physiologic Cytokine Solution ("PCS")) composition is generated by combining physiological levels of one or more of VEGF, Angiogenin, PDGF, TGF β 2, and one or more MMP inhibitor (i.e. TIMP-1 and/or TIMP-2) in a carrier. Optionally, the PCS contains Thymosin α 4. The physiological levels for these cytokines are the same as those found in ACCS. Suitable carriers include normal saline, PBS, lactated Ringer's solution, cell culture medium, conditioned cell culture media, water, etc. Such compositions are suitable for cryopreservation, lyophilization, sustained-release formulation, scale-up, and the like. It is contemplated that PCS may be produced such that it contains more concentrated levels of the factors than those found in ACCS and that it may be subsequently diluted with appropriate diluent prior to use. Appropriate diluents include, without limitation, normal saline, PBS, lactated Ringer's solution, cell culture media, conditioned cell culture media, water, and the like. Such dilutions may be 1:2, 1:3, 1:4, 1:5, 1:10, 1:100, etc. Such dilutions may also be less than 1:2 (i.e. 1:1, 1:0.5, etc.). The appropriate dilution required will be dependent upon the intended use and therefore will need to be empirically determined by the skilled artisan.

[0078] The compositions of the invention can be prepared in a variety of ways depending on their intended use. For example, a composition may be a liquid comprising an agent of the invention, i.e. ACCS, or pooled ACCS. A liquid composition also includes a gel. The liquid composition may be aqueous or in the form of an ointment, salve, cream, or the like. The liquid composition may also be formulated in such a way as to be suitable for a spray or aerosol application.

[0079] A useful aqueous suspension may contain one or more polymers as suspending agents. Useful polymers include water-soluble polymers such as cellulosic polymers and water-insoluble polymers such as cross-linked carboxyl-containing polymers. An aqueous suspension or solution/suspension is preferably viscous or muco-adhesive, or even more preferably, both viscous and muco-adhesive.

Sustained-Release Compositions

[0080] The CFS compositions, including but not limited to ACCS, pooled ACCS and PCS, maybe formulated as sustained-release CFS compositions (referred to herein as "SR-CFS"). Skilled artisans are familiar with methodologies to create sustained-release compositions of therapeutic agents, including protein-based therapeutic agents such as ACCS, pooled ACCS or PCS.

[0081] SR-CFS, including but not limited to SR-ACCS and SR-PCS, may be made by any of the methods described herein. For example, multivesicular liposome formulation technology is useful for the sustained-release of protein and peptide therapeutics. Qui, J., et al, (ACTA

Pharmacol Sin, 2005, 26(11):1395-401) describe this methodology for the formulation of sustained-release interferon alpha-2b. Vyas, S.P., et al, (Drug Dev Ind Pharm, 2006, 32(6):699-707) describe encapsulating pegylated interferon alpha in multivesicular liposomes. ACCS (including pooled ACCS) and PCS are suitable for use in multivesicular liposome sustained-release formulation.

[0082] Nanoparticle technology is also useful for creating SR-CFS. For example, Packhaeuser, C.B., et al, (J Control Release, 2007, 123(2):131-40) describe biodegradable parenteral depot systems based on insulin loaded dialkylaminoalkyl-amine-poly(vinyl alcohol)-g-poly(lactide-co-glycolide) nanoparticules and conclude that nanoparticle-based depots are suitable candidates for the design of controlled-release devices for bioactive macromolecules (i.e. proteins). Dailey, L.A., et al, (Pharm Res 2003, 20(12):2011-20) describe surfactant-free, biodegradable nanoparticles for aerosol therapy which is based on the branched polymers DEAPA-PVAL-g-PLGA and conclude that DEAPA-PVAL-g-PLGA are versatile drug delivery systems. ACCS (including pooled ACCS) and PCS are suitable for use in nanoparticle-based sustained-release formulations.

[0083] Polymer-based sustained-release formulations are also very useful. Chan, Y.P., et al, (Expert Opin Drug Deliv, 2007, 4(4):441-51) provide a review of the Medusa system (Flamel Technologies), which is used for sustained-release of protein and peptide therapies. Thus far, the Medusa system has been applied to subcutaneous injection of IL-2 and IFN-alpha(2b), in animal models (rats, dogs, monkeys), and in clinical trials in renal cancer (IL-2) and hepatitis C (IFN-alpha(2b)) patients. Chavanpatil, M.D., et al, (Pharm Res, 2007, 24(4):803-10) describe surfactant-polymer nanoparticles as a novel platform for sustained and enhanced cellular delivery of water-soluble molecules. Takeuchi, H., et al, (Adv Drug Deliv Res, 2001, 47(1):39-54) describe mucoadhesive nanoparticulate systems for peptide drug delivery, including liposomes and polymeric nanoparticles. Wong, H.L., et al, (Pharm Res, 2006, 23(7):1574-85) describe a new polymer-lipid hybrid system which has been shown to increase cytotoxicity of doxorubicin against multidrug-resistant breast cancer cells. CFS compositions, including but not limited to ACCS (including pooled ACCS) and PCS are suitable for use in the aforementioned sustained-release formulation methodologies.

[0084] In addition, other sustained-release methodologies familiar to skilled artisans, while not specifically described herein, are also suitable for use with the CFS compositions.

Pharmaceutical Compositions of CFS Compositions Including, But not Limited to, ACCS, Pooled ACCS, PCS, SR-ACCS (including pooled ACCS), and SR-PCS

[0085] The present invention provides pharmaceutical compositions of CFS compositions and a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle

with which the composition is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin, and still others are familiar to skilled artisans.

[0086] The pharmaceutical compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

Treatment Kits Comprising CFS Compositions Including, But not Limited to, ACCS, Pooled ACCS, PCS, SR-ACCS (including pooled ACCS), and SR-PCS

[0087] Also disclosed is an article of manufacture comprising packaging material and a pharmaceutical composition of the invention contained within the packaging material, wherein the pharmaceutical composition comprises CFS compositions. The packaging material comprises a label or package insert which indicates that the CFS compositions contained therein can be used for therapeutic applications such as, for example, wound healing.

Formulation, Dosage and Administration of CFS Compositions Including, But not Limited to, ACCS, Pooled ACCS, PCS, SR-ACCS (including pooled ACCS), and SR-PCS

[0088] Compositions comprising CFS compositions may be administered to a subject to provide various cellular or tissue functions, for example, to accelerate wound healing. As used herein "subject" may mean either a human or non-human animal.

[0089] Such compositions may be formulated in any conventional manner using one or more physiologically acceptable carriers optionally comprising excipients and auxiliaries. Proper formulation is dependent upon the route of administration chosen. The compositions may be packaged with written instructions for their therapeutic use. The compositions may also be administered to the recipient in one or more physiologically acceptable carriers. Carriers for CFS compositions may include but are not limited to solutions of normal saline, phosphate

buffered saline (PBS), lactated Ringer's solution containing a mixture of salts in physiologic concentrations, or cell culture medium.

[0090] In addition, one of skill in the art may readily determine the appropriate dose of the CFS compositions for a particular purpose. A preferred dose is in the range of about 0.1-to-1000 micrograms per square centimeter of applied area. Other preferred dose ranges are 1.0-to-50.0 micrograms/ applied area. It has been found that relatively small amounts of the CFS compositions are therapeutically useful. One exemplification of such therapeutic utility is the ability for ACCS (including pooled ACCS) to accelerate wound healing (for details see U.S. Publication No. 2006/0222634 and U.S. Publication No. 2007/1231297). One of skill in the art will also recognize that the number of doses to be administered needs also to be empirically determined based on, for example, severity and type of disease, disorder or injury being treated. For example, one dose is sufficient to have a therapeutic effect (i.e. accelerate wound healing). Other disclosures contemplate, 2, 3, 4, or more doses for therapeutic effect.

[0091] The skilled artisan will recognize that a preferred dose is one which produces a therapeutic effect (a therapeutically effective amount) such as accelerating wound healing, in a patient in need thereof. Of course, proper doses of the CFS compositions will require empirical determination at time of use based on several variables including but not limited to the severity and type of injury, disorder or condition being treated; patient age, weight, sex, health; other medications and treatments being administered to the patient; and the like. One of skill in the art will also recognize that number of doses (dosing regimen) to be administered needs also to be empirically determined based on, for example, severity and type of injury, disorder or condition being treated. In addition, one of skill in the art recognizes that the frequency of dosing needs to be empirically determined based on severity and type of injury, disorder or condition being treated. In certain disclosures, one dose is administered every day for a given number of days (i.e. once a day for 7 days, etc.). In other disclosures, multiple doses may be administered in one day (every 4 hours, etc.). Multiple doses per day for multiple days is also contemplated.

[0092] In further disclosures, at least one additional agent may be combined with the CFS compositions. Such agents may act synergistically with the CFS compositions of the invention to enhance the therapeutic effect. Such agents include but are not limited to growth factors, cytokines, chemokines, antibodies, inhibitors, antibiotics, immunosuppressive agents, steroids, anti-fungals, anti-virals or other cell types (i.e. stem cells or stem-like cells, for example AMP cells). Inactive agents include carriers, diluents, stabilizers, gelling agents, delivery vehicles, ECMs (natural and synthetic), scaffolds, and the like. When the CFS compositions are administered conjointly with other pharmaceutically active agents, even less of the CFS compositions may be needed to be therapeutically effective.

[0093] CFS compositions can be administered by injection into a target site of a subject, preferably via a delivery device, such as a tube, e.g., catheter. In a preferred disclosure, the tube additionally contains a needle, e.g., a syringe, through which the CFS compositions can be introduced into the subject at a desired location. Specific, non-limiting examples of

administering the CFS compositions to subjects may also include administration by subcutaneous injection, intramuscular injection, intravenous injection, intraarterial injection, intracardiac injection, intradermal injection, intrathecal injection, epidural injection, intraperitoneal injection, or intracerebral injection,. Infusions are also contemplated (i.e. subdural, intrathecal or intracerebral infusion). If administration is intravenous, an injectable liquid suspension can be prepared and administered by a continuous drip or as a bolus. In some instances, it may be appropriate to administer the CFS compositions using an infusion pump.

[0094] The timing of administration of CFS compositions will depend upon the type and severity of the disease, disorder or injury being treated. In one disclosure, the CFS compositions are administered as soon as possible after the diagnosis or injury. In another disclosure, CFS compositions are administered more than one time following diagnosis or injury. In certain disclosures, where surgery is required, the CFS compositions are administered at surgery. In still other disclosures, the CFS compositions are administered at as well as after surgery. Such post-surgical administration may take the form of a single administration or multiple administrations.

[0095] CFS compositions may also be inserted into a delivery device, e.g., a syringe, in different forms. For example, the CFS compositions can be part of a solution contained in such a delivery device. As used herein, the term "solution" includes a pharmaceutically acceptable carrier or diluent. Pharmaceutically acceptable carriers and diluents include saline, aqueous buffer solutions, solvents and/or dispersion media. The use of such carriers and diluents is well known in the art. The solution is preferably sterile and fluid to the extent that easy syringability exists. Preferably, the solution is stable under the conditions of manufacture and storage and may optionally be preserved against the contaminating action of microorganisms such as bacteria and fungi through the use of, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. Solutions of the invention can be prepared by incorporating the CFS compositions in a pharmaceutically acceptable carrier or diluent and, as required, other ingredients enumerated above.

[0096] CFS compositions may be administered systemically (for example intravenously), locally (for example by direct application under visualization during surgery) or topically. For such administration, the compositions may be in an injectable liquid suspension preparation or in a biocompatible medium which is injectable in liquid form and becomes semi-solid at the site of damaged tissue. A controllable endoscopic delivery device can also be used.

[0097] Support matrices into which the CFS compositions can be incorporated or embedded include matrices which are recipient-compatible and which degrade into products which are not harmful to the recipient.

[0098] Natural and/or synthetic biodegradable matrices are examples of such matrices. Natural biodegradable matrices include plasma clots, e.g., derived from a mammal, collagen, fibronectin, and laminin matrices. Suitable synthetic matrix material must be biocompatible to

preclude immunological complications. It must also be resorbable. The matrix should be configurable into a variety of shapes and should have sufficient strength to prevent collapse upon implantation. Recent studies indicate that the biodegradable polyester polymers made of polyglycolic acid fulfill all of these criteria (Vacanti, et al. J. Ped. Surg. 23:3-9 (1988); Cima, et al. Biotechnol. Bioeng. 38:145 (1991); Vacanti, et al. Plast. Reconstr. Surg. 88:753-9 (1991)). Other synthetic biodegradable support matrices include synthetic polymers such as polyanhydrides, polyorthoesters, and polylactic acid. Further examples of synthetic polymers and methods of incorporating or embedding compositions into these matrices are also known in the art. See e.g., U.S. Pat. Nos. 4,298,002 and 5,308,701.

[0099] One of the advantages of a biodegradable polymeric matrix is that CFS compositions can be incorporated directly into the support matrix so that it is slowly released as the support matrix degrades in vivo. In addition to the CFS compositions, other factors, including nutrients, growth factors, inducers of differentiation or de-differentiation (i.e., causing differentiated cells to lose characteristics of differentiation and acquire characteristics such as proliferation and more general function), products of secretion, immunomodulators, inhibitors of inflammation, regression factors, biologically active compounds which enhance or allow ingrowth of the lymphatic network or nerve fibers, hyaluronic acid, and drugs, which are known to those skilled in the art and commercially available with instructions as to what constitutes an effective amount, from suppliers such as Collaborative Research, Sigma Chemical Co., growth factors such as epidermal growth factor (EGF) and heparin binding epidermal growth factor like growth factor (HB-EGF), could be incorporated into the matrix or be provided in conjunction with the matrix. Similarly, polymers containing peptides such as the attachment peptide RGD (Arg-Gly-Asp) can be synthesized for use in forming matrices (see e.g. U.S. Pat. Nos. 4,988,621, 4,792,525, 5,965,997, 4,879,237 and 4,789,734).

[0100] In another example, the CFS compositions may be incorporated in a gel matrix (such as Gelfoam from Upjohn Company). A variety of encapsulation technologies have been developed (e.g. Lacy et al., Science 254:1782-84 (1991); Sullivan et al., Science 252:718-712 (1991); WO 91/10470; WO 91/10425; U.S. Patent No. 5,837,234; U.S. Pat. No. 5,011,472; U.S. Pat. No. 4,892,538). During open surgical procedures involving direct physical access to diseased or damaged tissue, all of the described forms of the CFS composition delivery preparations are available options. These compositions can be repeatedly administered at intervals until a desired therapeutic effect, i.e. accelerated wound healing, is achieved.

[0101] The three-dimensional matrices to be used are structural matrices that provide a scaffold to guide the process of tissue healing and formation. Scaffolds can take forms ranging from fibers, gels, fabrics, sponge-like sheets, and complex 3-D structures with pores and channels fabricated using complex Solid Free Form Fabrication (SFFF) approaches. As used herein, the term "scaffold" means a three-dimensional (3D) structure (substrate and/or matrix). It may be composed of biological components, synthetic components or a combination of both. Further, it may be naturally constructed by cells or artificially constructed. In addition, the scaffold may contain components that have biological activity under appropriate conditions. The structure of the scaffold can include a mesh, a sponge or can be formed from a hydrogel.

[0102] The design and construction of the scaffolding to form a three-dimensional matrix is of primary importance. The matrix should be a pliable, non-toxic, injectable porous template for vascular ingrowth. The pores should allow vascular ingrowth. These are generally interconnected pores in the range of between approximately 100 and 300 microns, i.e., having an interstitial spacing between 100 and 300 microns, although larger openings can be used. The matrix should be shaped to maximize surface area, to allow adequate diffusion of nutrients, gases and growth factors. At the present time, a porous structure that is relatively resistant to compression is preferred, although it has been demonstrated that even if one or two of the typically six sides of the matrix are compressed, that the matrix is still effective to yield tissue growth.

[0103] The polymeric matrix may be made flexible or rigid, depending on the desired final form, structure and function. For repair of a defect, for example, a flexible fibrous mat is cut to approximate the entire defect then fitted to the surgically prepared defect as necessary during implantation. An advantage of using the fibrous matrices is the ease in reshaping and rearranging the structures at the time of implantation.

[0104] Also disclosed is the delivery of CFS compositions in conjunction with any of the above support matrices as well as amnion-derived membranes. Such membranes may be obtained as a byproduct of the process described herein for the recovery of AMP cells, or by other methods, such as are described, for example, in U.S. Patent No. 6,326,019 which describes a method for making, storing and using a surgical graft from human amniotic membrane, US 2003/0235580 which describes reconstituted and recombinant amniotic membranes for sustained delivery of therapeutic molecules, proteins or metabolites, to a site in a host, U.S. 2004/0181240, which describes an amniotic membrane covering for a tissue surface which may prevent adhesions, exclude bacteria or inhibit bacterial activity, or to promote healing or growth of tissue, and U.S. Patent No. 4,361,552, which pertains to the preparation of cross-linked amnion membranes and their use in methods for treating burns and wounds. In accordance with the present invention, CFS compositions may be incorporated into such membranes.

Exemplary Therapeutic Uses of CFS Compositions Including, But not Limited to, ACCS, Pooled ACCS, PCS, SR-ACCS (including pooled ACCS), and SR-PCS

[0105] Wound healing - The CFS compositions of the present invention are effective in accelerating wound healing of wounds caused by a number of sources, including but not limited to incisional, compression, thermal, radiation, penetrating, concussive, acute, chronic, infected, and sterile injuries. The instant invention is based upon the discovery that CFS compositions can accelerate the wound healing process for all wound types, particularly when administered topically, i.e. to the surface of the wound site. Using CFS compositions all wound types, mechanical or thermal, acute or chronic, infected or sterile, undergo healing more rapidly than similar wounds left to heal naturally or which are treated with currently available

methods. A "therapeutically effective amount" of a therapeutic agent within the meaning of the present invention will be determined by a patient's attending physician or veterinarian. Such amounts are readily ascertained by one of ordinary skill in the art and will enable accelerated wound healing when administered in accordance with the present invention. Factors which influence what a therapeutically effective amount will be include, the specific activity of the therapeutic agent being used, the wound type (mechanical or thermal, full or partial thickness, etc.), the size of the wound, the wound's depth (if full thickness), the absence or presence of infection, time elapsed since the injury's infliction, and the age, physical condition, existence of other disease states, and nutritional status of the patient. Additionally, other medication the patient may be receiving will effect the determination of the therapeutically effective amount of the therapeutic agent to administer.

Examples

[0106] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the compositions and methods of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade, and pressure is at or near atmospheric.

Example 1: Preparation of AMP cell compositions

[0107] Recovery of AMP cells - AMP cells were dissociated from starting amniotic membrane using the dissociation agents PXXIII, and trypsin. The average weight range of an amnion was 18-27 g. The number of cells recovered per g of amnion was about $10\text{-}15 \times 10^6$ for dissociation with PXXIII and $5\text{-}8 \times 10^6$ for dissociation with trypsin.

[0108] Method of obtaining selected AMP cells: Cells were plated immediately upon isolation from the amnion. After ~2 days in culture non-adherent cells were removed and the adherent cells were kept. This attachment to a plastic tissue culture vessel is the selection method used to obtain the desired population of AMP cells. Adherent and non-adherent AMP cells appear to have a similar cell surface marker expression profile but the adherent cells have greater viability and are the desired population of cells. Adherent AMP cells were cultured until they reached ~120,000-150,000 cells/cm². At this point, the cultures were confluent. Suitable cell cultures will reach this number of cells between ~5-14 days. Attaining this criterion is an indicator of the proliferative potential of the AMP cells and cells that do not achieve this criterion are not selected for further analysis and use. Once the AMP cells reached ~120,000-

150,000 cells/cm², they were collected and cryopreserved. This collection time point is called p0.

Example 2: Generation of ACCS

[0109] The AMP cells of the invention can be used to generate ACCS, including pooled ACCS.

The AMP cells were isolated as described above and $\sim 1 \times 10^6$ cells/mL were seeded into T75 flasks containing ~ 10 mL culture medium. The cells were cultured until confluent, the medium was changed and ACCS was collected 3 days post-confluence. Skilled artisans will recognize that other means for collecting ACCS from confluent cultures, such as using other tissue culture vessels, including but not limited to cell factories, flasks, hollow fibers, or suspension culture apparatus, etc. are also contemplated (see Detailed Description above). It is also contemplated that the ACCS be cryopreserved, lyophilized or formulated for sustained-release following collection. It is also contemplated that ACCS be collected at different time point (see Detailed Description for details).

Example 3: Generation of Pooled ACCS

[0110] ACCS was obtained essentially as described above. In certain embodiments, ACCS was collected multiple times from an AMP cell culture derived from one placenta and these multiple ACCS collections were pooled together. Such pools are referred to as "SP pools" (more than one ACCS collection/one placenta). In another embodiment, AMP cell cultures were derived from several placentas, i.e. from 5 or 10 placentas. The AMP cells from each placenta were cultured and one ACCS collection from each culture was collected and then they were all pooled. These pools are termed "MP1 pools" (one ACCS collection/placenta, multiple placentas). In yet another embodiment, AMP cell cultures were derived from several placentas, i.e. from 5 or 10 placentas. The AMP cells from each placenta were cultured and more than one ACCS collection was performed from each AMP cell culture and then pooled. These pools are termed "MP2 pools" (more than one ACCS collection/placenta, multiple placentas).

Example 4: Detection of Cytokines in Non-pooled and Pooled ACCS Using ELISA

[0111] Standard ELISAs familiar to skilled artisan are performed on ACCS from AMP cells obtained from 10 different placentas. In addition to assaying each ACCS sample individually, pooled ACCS samples are also tested to determine if variability of ELISA results between samples is reduced. ACCS is obtained as described above. Pools are made as follows: Pool 1 is comprised of ACCS from placentas 1-5, Pool 2 is comprised of ACCS from placentas 6-10, and Pool 3 is comprised of ACCS from placentas 1-10. In addition, ELISA of SP, MP1, MP2 pools is performed.

Example 5: Generation of PCS Compositions

[0112] The following PCS compositions are produced by combining the indicated cytokine or factor at physiologic levels in a carrier:

Composition A: VEGF and TIMP-1

Composition B: VEGF, Angiogenin and TIMP-1

Composition C: VEGF, Angiogenin, PDGF-BB and TIMP-1

Composition D: VEGF, Angiogenin, PDGF-BB, TGF β 2 and TIMP-1

Composition E: VEGF and TIMP-2

Composition F: VEGF, Angiogenin and TIMP-2

Composition G: VEGF, Angiogenin, PDGF-BB and TIMP-2

Composition H: VEGF, Angiogenin, PDGF-BB, TGF β 2 and TIMP-2

Composition I: VEGF, TIMP-1 and TIMP-2

Composition J: VEGF, Angiogenin, TIMP-1 and TIMP-2

Composition K: VEGF, Angiogenin, PDGF-BB, TIMP-1 and TIMP-2

Composition L: VEGF, Angiogenin, PDGF-BB, TGF β 2, TIMP-1 and TIMP-2

Composition M: Angiogenin and TIMP-1

Composition N: Angiogenin, PDGF-BB and TIMP-1

Composition O: Angiogenin, PDGF-BB, TGF β 2 and TIMP-1

Composition P: Angiogenin and TIMP-2

Composition Q: Angiogenin, PDGF-BB and TIMP-2

Composition R: Angiogenin, PDGF-BB, TGF β 2 and TIMP-2

Composition S: Angiogenin, PDGF-BB, TGF β 2, TIMP-1 and TIMP-2

Composition T: PDGF-BB and TIMP-1

Composition U: PDGF-BB, TGF β 2 and TIMP-1

Composition V: PDGF-BB and TIMP-2

Composition W: PDGF-BB, TGF β 2 and TIMP-2

Composition X: PDGF-BB, TIMP-1 and TIMP-2

Composition Y: PDGF-BB, TGF β 2, TIMP-1 and TIMP-2

[0113] Compositions A-Y optionally contains Thymosin β 4. Skilled artisans will recognize that in certain embodiments other MMP inhibitors (i.e. TIMP-3, TIMP-4 or synthetic MMP inhibitors) may be suitable (J. Frederick Woessner, Jr., J. Clin. Invest. 108(6): 799-800 (2001); Brew, K., et al, Biochim Biophys Acta. 2000 Mar 7;1477(1-2):267-83).

[0114] VEGF, Angiogenin, PDGF-BB, TGF β 2, TIMP-1 and TIMP-2 are added at the following physiologic levels: ~5-16ng/mL for VEGF, ~3.5-4.5 ng/mL for Angiogenin, ~100-165pg/mL for PDGF, ~2.5-2.7ng/mL for TGF β 2, ~0.68 μ g/mL for TIMP-1 and ~104 μ g/mL for TIMP-2. VEGF may be obtained from Invitrogen, catalog #PHG0144, PHG0145, PHG0146, PHG0141 or PHG0143; Angiogenin may be obtained from R&D Systems, catalog #265-AN-050 or 265-AN-250; PDGF-BB may be obtained from Invitrogen, catalog #PHG0044, #PHG0045, #PHG0046, #PHG0041, #PHG0043; TGF β 2 may be obtained from Invitrogen, catalog #PHG9114; TIMP-1 may be obtained from R&D Systems, catalog #970-TM-010; and TIMP-2 may be obtained from R&D Systems, catalog #971-TM-010. VEGF, Angiogenin, PDGF-BB, TGF β 2, TIMP-1 and TIMP-2 are added to a carrier such as normal saline, PBS, lactated Ringer's solution, cell culture media, water or other suitable aqueous solution known to skilled artisans.

[0115] The PCS compositions are tested in standard animal models for wound healing to assess activity (see Definitions above for standard animal models for wound healing).

Example 6: Generation of Sustained-Release CFS Compositions

[0116] SR-CFS compositions, such as, for example, SR-ACCS (including pooled ACCS) or SR-PCS, are produced by combining CFS compositions with any of the sustained-release formulation technologies described herein (see Detailed Description) or otherwise familiar to skilled artisans.

Example 7: Effects of ACCS in an animal model of chronic wound healing.

[0117] An art-accepted animal model for chronic granulating wound was used to study the effects of ACCS on chronic wound healing (Hayward PG, Robson MC: Animal models of wound contraction. In Barbul A, et al: Clinical and Experimental Approaches to Dermal and Epidermal Repair: Normal and Chronic Wounds. John Wiley & Sons, New York, 1990.).

[0118] Results: ACCS was effective in not allowing proliferation of tissue bacterial bioburden. ACCS allowed accelerated healing of the granulating wound significantly faster than the non-

treated infected control groups.

Example 8: Use of CFS compositions PCS, SR-ACCS and SR-PCS in an Animal Model of Chronic Wound Healing.

[0119] An art-accepted animal model for chronic granulating wound (Hayward PG, Robson MC: Animal models of wound contraction. In Barbul A, et al: Clinical and Experimental Approaches to Dermal and Epidermal Repair: Normal and Chronic Wounds. John Wiley & Sons, New York, 1990) is used to study the effects of the CFS compositions PCS, SR-ACCS or SR-PCS of the invention on chronic wound healing.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US5096885A](#) [0007]
- [US4297344A](#) [0007]
- [US5589167A](#) [0007]
- [WO8903671A](#) [0007]
- [US6191107B1](#) [0007]
- [US20030235563A](#) [0029] [0067]
- [US20040161419A](#) [0029] [0067]
- [US20050124003A](#) [0029] [0067]
- [US666949P](#) [0029] [0030] [0067]
- [US60699257B](#) [0029] [0030] [0067]
- [US60742067B](#) [0029] [0030] [0067]
- [US60813759B](#) [0029] [0067]
- [US333849A](#) [0029] [0030] [0067]
- [US392892A](#) [0029] [0030] [0067]
- [US06011392W](#) [0029] [0030] [0067]
- [US20060078993A](#) [0029] [0067]
- [US0040052W](#) [0029] [0067]
- [US7045148B](#) [0029] [0067]
- [US20040048372A](#) [0029] [0067]

- [US20030032179A](#) [0029] [0067]
- [US20040110287A](#) [0030]
- [US813759P](#) [0030]
- [US6372494B](#) [0035]
- [US20060222634A1](#) [0069]
- [US20060222634A](#) [0090]
- [US20071231297A](#) [0090]
- [US4298002A](#) [0098]
- [US5308701A](#) [0098]
- [US4988621A](#) [0099]
- [US4792525A](#) [0099]
- [US5965997A](#) [0099]
- [US4879237A](#) [0099]
- [US4789734A](#) [0099]
- [WO9110470A](#) [0100]
- [WO9110425A](#) [0100]
- [US5837234A](#) [0100]
- [US5011472A](#) [0100]
- [US4892538A](#) [0100]
- [US6326019B](#) [0104]
- [US20030235580A](#) [0104]
- [US20040181240A](#) [0104]
- [US4361552A](#) [0104]

Non-patent literature cited in the description

- Pharm. Res., 1991, vol. 8, 285-291 [0007]
- **ANKER et al.** Stem Cells, 2005, vol. 22, 1338-1345 [0030]
- **RAMKUMAR et al.** Am. J. Ob. Gyn., 1995, vol. 172, 493-500 [0030]
- Animal models of wound contraction **HAYWARD PGROBSON MC et al.** Clinical and Experimental Approaches to Dermal and Epidermal Repair: Normal and Chronic Wounds John Wiley & Sons 19900000 [0062] [0117] [0119]
- **DELBECARRO et al.** The use of specific thromboxane inhibitors to preserve the dermal microcirculation after burning Surgery, 1980, vol. 87, 137-141 [0062]
- **ROBSON et al.** Increasing dermal perfusion after burning by decreasing thromboxane production J Trauma, 1980, vol. 20, 722-725 [0062]
- **POLO et al.** An in vivo model of human proliferative scar J Surg Res, 1998, vol. 74, 187-195 [0062]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual 19940000 [0063]
- Current Protocols in Molecular Biology 19940000 vol. I-III, [0063]

- Cell Biology: A Laboratory Handbook 19940000 vol. I-III, [0063]
- Current Protocols in Immunology 19840000 vol. I-III, [0063]
- Oligonucleotide Synthesis 19850000 [0063]
- Nucleic Acid Hybridization 19840000 [0063]
- Transcription And Translation 19860000 [0063]
- Animal Cell Culture IRL Press 19860000 [0063]
- Immobilized Cells And Enzymes, [0063]
- PERBALA Practical Guide To Molecular Cloning, 1984, [0063]
- QUI, J. et al. ACTA Pharmacol Sin, 2005, vol. 26, 111395-401 [0081]
- VYAS, S.P. et al. Drug Dev Ind Pharm, 2006, vol. 32, 6699-707 [0081]
- PACKHAEUSER, C.B. et al. J Control Release, 2007, vol. 123, 2131-40 [0082]
- DAILEY, L.A. et al. Pharm Res, 2003, vol. 20, 122011-20 [0082]
- CHAN, Y.P. et al. Expert Opin Drug Deliv, 2007, vol. 4, 4441-51 [0083]
- CHAVANPATIL, M.D. et al. Pharm Res, 2007, vol. 24, 4803-10 [0083]
- TAKEUCHI, H. et al. Adv Drug Deliv Res, 2001, vol. 47, 139-54 [0083]
- WONG, H.L. et al. Pharm Res, 2006, vol. 23, 71574-85 [0083]
- VACANTI et al. J. Ped. Surg., 1988, vol. 23, 3-9 [0098]
- CIMA et al. Biotechnol. Bioeng., 1991, vol. 38, 145- [0098]
- VACANTI et al. Plast. Reconstr. Surg., 1991, vol. 88, 753-9 [0098]
- LACY et al. Science, 1991, vol. 254, 1782-84 [0100]
- SULLIVAN et al. Science, 1991, vol. 252, 718-712 [0100]
- J. FREDERICK WOESSNER, JR. J. Clin. Invest., 2001, vol. 108, 6799-800 [0113]
- BREW, K. et al. Biochim Biophys Acta, 2000, vol. 1477, 1-2267-83 [0113]

PATENTKRAV

1. Amnion-afledt cellulær cytokinopløsningssammensætning, der omfatter fysiologiske koncentrationer af VEGF, TGFβ2, angiogenin, PDGF, TIMP-1 og TIMP-2, hvor den fysiologiske koncentration er ~5,0-16 ng/mL for VEGF, ~3,5-4,5
5 ng/mL for angiogenin, ~100-165 pg/mL for PDGF, ~2,5-2,7 ng/mL for TGFβ2, ~0,68 µg/mL for TIMP-1 og ~1,04 µg/mL for TIMP-2, hvor den amnion-afledte cellulære cytokinopløsningssammensætning kan opnås ved følgende fremgangsmåde:
- (a) isolering af amnion-epitelceller fra en placentas amnion;
 - 10 (b) udvælgelse af AMP-celler fra amnion-epithelcellerne;
 - (c) dyrkning af AMP-cellerne i dyrefrit baseret medium, hvor mediet tilsættes humant albumin af klinisk kvalitet op til en koncentration på 10 % indtil sammenløb;
 - (d) ændring af dyrkningsmediet;
 - 15 (e) dyrkning af cellerne i mediet i 1, 2, 3, 4, 5 eller 6 dage; og
 - (f) opsamling af dyrkningsmediet trin (e) for at opnå amnion-afledt cellulær cytokinopløsning;
- hvor mediet fra trin (c) er Stemline I eller II, Opti-pro, IMDM eller DMEM.
2. Sammensætning ifølge krav 1, hvor trin (f) er gentaget flere gange og
20 den amnion-afledte cellulære cytokinopløsning opnået i trin (f) er kombineret for at frembringe en aggregeret amnion-afledt cellulær cytokinopløsning.
3. Sammensætning ifølge krav 1(c), hvor mediet er serumfrit.
4. Sammensætning ifølge et hvilket som helst af kravene 1 til 3, hvor det basale medium endvidere er tilsat EGF ved en koncentration på mellem 0 og 1
25 mg/µl.
5. Sammensætning med langvarig frigørelse, der omfatter den amnion-afledte cellulære cytokinopløsning ifølge et hvilket som helst af kravene 1 til 2.
6. Fremgangsmåde til fremstilling af en sammensætning med langvarig frigørelse ifølge krav 5, der omfatter trinnene med kombineret af en

cellefaktorholdig opløsningssammensætning med et middel, der er i stand til at bevirke langvarig frigørelse af den cellefaktorholdige opløsningssammensætning.

7. Farmaceutisk sammensætning, der omfatter sammensætningen ifølge et hvilket som helst af kravene 1 til 5.

- 5 **8.** Sammensætning ifølge et hvilket som helst af kravene 1 til 5 eller 7, til anvendelse i en fremgangsmåde til fremskyndelse af sårheling.