



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2410802 A1 2002/09/06

(21) **2 410 802**

**(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2001/02/21
(87) Date publication PCT/PCT Publication Date: 2002/09/06
(85) Entrée phase nationale/National Entry: 2002/11/28
(86) N° demande PCT/PCT Application No.: US 2001/005577
(87) N° publication PCT/PCT Publication No.: 2002/067935

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/44

(71) **Demandeur/Applicant:**
TEVA PHARMACEUTICAL INDUSTRIES LTD., IL
(72) **Inventeurs/Inventors:**
LEIBOVICI, MINUTZA, IL;
TENENGAUZER, RUTH, IL;
KOPEL, MIRA, IL;
ARONHIME, JUDITH, IL;
KORDOVA, MARCO, IL
(74) **Agent:** RIDOUT & MAYBEE LLP

(54) Titre : FORMULATION PHARMACEUTIQUE STABLE CONTENANT DU TORSEMIDE MODIFICATION II
(54) Title: A STABLE PHARMACEUTICAL FORMULATION COMPRISING TORSEMIDE MODIFICATION II

(57) Abrégé/Abstract:

Novel, stable pharmaceutical formulations for the oral administration of high purity torsemide modification II are disclosed. These formulations release high purity torsemide modification II in water at a constant and high purity rate, and the high purity torsemide modification II therein does not rearrange to torsemide modification I over time. Methods for their manufacture are also disclosed.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/067935 A1(51) International Patent Classification⁷:

A61K 31/44

(74) Agents: BRAINARD, Charles, R. et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).

(21) International Application Number:

PCT/US01/05577

(22) International Filing Date: 21 February 2001 (21.02.2001)

(25) Filing Language:

English

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language:

English

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except BB, US): TEVA PHARMACEUTICAL INDUSTRIES LTD. [IL/IL]; 5 Basel Street, P.O. Box 3190, 49131 Petah Tiqva (IL).

(71) Applicant (for BB only): TEVA PHARMACEUTICALS USA, INC. [US/US]; 1090 Horsham Road, P.O. Box 1090, North Wales, PA 19454-1090 (US).

(72) Inventors; and

Published:

— with international search report

(75) Inventors/Applicants (for US only): LEIBOVICI, Minutza [IL/IL]; 3/1 Gershon Shatz St., Netanya (IL). TENENGAUZER, Ruth [IL/IL]; 9/4 Ben Gamla Street, Hod HaSharon (IL). KOPEL, Mira [IL/IL]; P.O. Box 1298, Pardess Hanna (IL). ARONHIME, Judith [IL/IL]; 9/7 Hava Lutzky Street, 44244 Rehovot (IL). KORDOVA, Marco [IL/IL]; 78/14 Rothchild, Kfar Saba (IL).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/067935 A1

(54) Title: A STABLE PHARMACEUTICAL FORMULATION COMPRISING TORSEMIDE MODIFICATION II

(57) Abstract: Novel, stable pharmaceutical formulations for the oral administration of high purity torsemide modification II are disclosed. These formulations release high purity torsemide modification II in water at a constant and high purity rate, and the high purity torsemide modification II therein does not rearrange to torsemide modification I over time. Methods for their manufacture are also disclosed.

**A STABLE PHARMACEUTICAL FORMULATION COMPRISING
TORSEMIDE MODIFICATION II**

CROSS-REFERENCE TO RELATED APPLICATION

5 This application claims the benefit of provisional application serial number 60/183,288, filed February 17, 2000, the contents of which are incorporated herein by reference.


FIELD OF THE INVENTION

10 The present invention relates to the formulation of pharmaceutical compositions, and more particularly to novel pharmaceutical formulations for the oral administration of 15 tormamide.

BACKGROUND OF THE INVENTION

1-Isopropyl-3-[(4-*m*-toluidino-3-pyridyl)-sulfonyl]urea, which is represented by the structural formula

15



is approved, under the trademark DEMADEX®, by the U.S. Food and Drug Administration, for the treatment of hypertension and edema associated with congestive heart failure, renal disease, or hepatic disease. The USAN approved generic name for this 20 compound is tormamide, although this compound is also referred to as "torasemide" in the art. Tormamide is a loop diuretic that has been found to be particularly effective for the treatment of edema associated with chronic renal failure.

U.S. Patent No. Re. 30,633 describes a synthesis of tormamide. It is known that 25 tormamide can occur in at least two different crystalline forms, Acta Cryst. 1978, pp. 2659-2662 and Acta Cryst., 1978, pp. 1304-1310, in which the crystal identified by space group $P2_1/c$ is designated Dupont Form 1 herein and the crystal identified by space group $P2/n$ is

designated Dupont Form 2 herein. U.S. Patent No. 4,822,807, which reissued as U.S. Patent No. Re. 34,672, describes two crystalline forms of torsemide, designated modification I and modification II. Torsemide modification I is defined herein as the torsemide characterized by the x-ray powder diffraction pattern of Figure 1, in the 37 C.F.R. § 1.132 declaration by Dr. Fritz Topfmeier filed on December 30, 1987, which is located in the file wrapper of U.S. Patent 4,822,807 (the "Topfmeier Declaration"). Torsemide modification II is defined herein as the torsemide characterized by the x-ray powder diffraction pattern of Figure 2, in the Topfmeier Declaration.

Topfmeier and Lettenbauer have described in U.S. Patent No. 4,822,807 that when torsemide of modification II is present in very finely divided form in pharmaceutical tablets, it rearranges into torsemide modification I, with the result that the rate of dissolution of the active material upon introducing the tablets into water can be significantly changed. The dissolution rate is an important characteristic of a pharmaceutical dosage form and, in order to dose reproducibly, must not differ from one 15 tablet to the next.

Thus, there remains a need in the art for pharmaceutical formulations containing torsemide modification II, wherein the torsemide does not rearrange into torsemide modification I and that are stable with regard to dissolution rate.

20 SUMMARY OF THE INVENTION

Stable pharmaceutical formulations containing torsemide modification II, wherein, upon storage under stress conditions, the torsemide modification II does not rearrange into torsemide modification I and wherein the stable pharmaceutical formulations are also stable with regard to dissolution rate in solution, have been discovered.

25 Additionally, a high purity torsemide modification II which is substantially free of other forms of torsemide and processes for making the high purity torsemide modification II have also been discovered.

The present invention relates to high purity torsemide modification II that does not substantially rearrange into a different form of torsemide over time upon storage in bulk 30 under stress conditions, e.g., 40°C, 75% relative humidity. Preferably, the high purity

torsemide modification II does not substantially rearrange into torsemide modification I over time upon storage in bulk under stress conditions, *e.g.*, 40°C, 75% relative humidity.

The present invention also relates to a stable pharmaceutical formulation comprising an effective amount of torsemide modification II and a pharmaceutically acceptable excipients wherein the excipients have a low moisture content. Preferably, the stable pharmaceutical formulation further comprises the excipients lactose anhydrous, crospovidone, povidone, microcrystalline cellulose, and magnesium stearate all of which having a low moisture content. Preferably, the stable pharmaceutical formulation comprises torsemide modification II in an amount of about 2.5 mg to about 200 mg per tablet. More preferably, the stable pharmaceutical formulation comprises torsemide modification II in an amount of 2.5 mg, 5 mg, 10 mg, 20 mg or 100 mg.

The present invention also relates to stable pharmaceutical formulations comprising an effective amount of torsemide modification II wherein the torsemide modification II does not substantially rearrange into another form of torsemide over time upon storage under stress conditions, *e.g.*, 40°C, 75% relative humidity. Preferably, the torsemide modification II of the stable pharmaceutical formulations does not substantially rearrange into torsemide modification I over time upon storage under stress conditions. Torsemide modification II suitable for use in the present stable pharmaceutical formulations includes high purity torsemide modification II and torsemide modification II containing trace amounts of torsemide modification I. Trace amounts, as defined herein, are amounts of torsemide modification I that are about 0.5-2% by weight of the torsemide modification II (w/w % of torsemide modification I/torsemide modification II).

The present invention relates to stable pharmaceutical formulations comprising an effective amount of torsemide modification II wherein the torsemide modification II has a particle size distribution such that 100 % is below 200 μ . Preferably, the particle size distribution is such that 100% is below 100 μ . More preferably, the particle size distribution is such that 100% is below 50 μ .

The present invention also relates to stable pharmaceutical formulations comprising an effective amount of torsemide modification II wherein the dissolution rate *in vitro* of the stable pharmaceutical formulation, when measured by the U.S.P. Paddle Method at 50-90

RPM in 900 mL water, is not less than 80% (by weight) of the torsemide modification II released after 30 minutes. Preferably, the dissolution rate *in vitro* does not substantially change over time upon storage in bulk under stress conditions, *e.g.*, 40°C, 75% relative humidity. More preferably, the dissolution rate *in vitro* does not substantially change 5 during storage under stress conditions for at least 3 months.

The present invention also relates to a process for making high purity torsemide modification II wherein the high purity torsemide modification II is purified from crude modification II by the novel combination of two purification steps known in the art wherein the novel process comprises the steps of (1) reslurrying crude torsemide 10 modification II followed by (2) crystallization to yield high purity torsemide modification II by the methods of U.S. Serial No. 09/638,106; filed August 11, 2000, the contents of which are incorporated herein by reference

BRIEF DESCRIPTION OF THE DRAWINGS

15 Figure 1 is an x-ray powder diffraction pattern of a high purity torsemide modification II tablet (Batch No. K-26683).

Figure 2 is an x-ray powder diffraction pattern of bulk high purity torsemide modification II (API 851700100).

20 Figure 3 is an x-ray powder diffraction pattern of a placebo tablet corresponding to a tablet containing 100 mg of high purity torsemide modification II.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to high purity torsemide modification II wherein the 25 high purity torsemide modification II has the surprising and useful advantage of being a stable polymorphic form of torsemide, that is, it does not substantially rearrange over time, thereby making high purity torsemide modification II of the present invention useful for the manufacture of stable pharmaceutical tablets of torsemide modification II. Preferably, the high purity torsemide modification II is in the form of fine crystal. Preferably, the high 30 purity torsemide modification II does substantially rearrange over time into torsemide modification I (such as not more than 10% of torsemide modification II rearranges to torsemide

modification I).

The high purity torsemide modification II of the present invention may be in the form of fine crystals. The high purity torsemide modification II may be further characterized by having a particle size distribution such that 100 % is below 200 μ . Preferably, the particle size distribution is such that 100% is below 100 μ . More preferably, the particle size distribution is such that 100% is below 50 μ .

It was surprisingly found that when torsemide modification II is crystallized as high purity torsemide modification II, with no trace amounts of torsemide modification I, the high purity torsemide modification II is stable during storage under stress conditions for at least 3 months. In contrast, torsemide modification II that contains trace amounts of torsemide modification I is not stable during storage under stress condition for at least 3 months. The torsemide modification II containing trace amounts of torsemide modification I rearranges into torsemide modification I over time during storage under stress conditions.

Significantly, it has been found that upon storage at 40°C, 75 % relative humidity, for 3 months, the polymorphic content of high purity torsemide modification II of the tablet formulations, or the bulk active ingredient, does not undergo any significant rearrangement into different polymorphic forms of torsemide. Preferably, not more than 10% of the high purity torsemide modification II rearranges into different polymorphic forms of torsemide following storage of the tablets or bulk active ingredient. More preferably, not more than 5% of the high purity torsemide modification II rearranges into different polymorphic forms. Even more preferably, not more than 2% of the high purity torsemide modification II rearranges into different polymorphic forms and most preferably, the high purity torsemide modification II is substantially pure polymorph torsemide modification II following storage. Specifically, the high purity torsemide modification II of the present invention does not undergo a polymorphic rearrangement into torsemide modification I. The detection of torsemide modification I in bulk high purity torsemide modification II or tablets of high purity torsemide modification II may be accomplished by using x-ray powder diffraction techniques. No substantial polymorphic change of the high purity torsemide modification II of the present pharmaceutical formulations or present bulk

active ingredient can be detected
by x-ray powder diffraction techniques.

5

Without being bound by theory, it is believed that the level of purity presently achieved in the high purity torsemide modification II imparts the polymorph with unexpected stability. It is feasible that the unstable torsemide modification II described in the relevant art contains trace amounts of torsemide modification I, the presence of which facilitates the rearrangement of torsemide modification II into torsemide modification I. It has been reported in the art that trace amounts of torsemide modification I facilitates the conversion of torsemide modification II into torsemide modification I when in an aqueous suspension.

15 The present invention also relates to a novel and stable pharmaceutical formulation containing fine crystals of torsemide modification II wherein the present stable pharmaceutical formulations have the surprising and useful advantage that the active material, torsemide modification II, does not substantially rearrange into torsemide modification I (such as not more than 5% of torsemide modification II rearranges to torsemide modification I), thereby making the stable pharmaceutical formulations of the present invention useful for the administration of torsemide modification II. The pharmaceutical formulations of the present invention are solid dosage forms for the oral administration of torsemide that are presented as a tablet.

20 Surprisingly, it was also found that the pharmaceutical formulation containing use
25 of Excipient with a low water content stabilizes modification II.

30 The present invention also provides methods for making stable pharmaceutical formulations torsemide modification II which are tablets. High purity torsemide modification II tablets are prepared by mixing the active ingredient, torsemide modification II, with a combination of excipients including, lactose anhydrous NF,

crospovidone NF, povidone USP (PVP K-30), and microcrystalline cellulose NF (Avicel PH 112). Alcohol 95% USP is added to the powder mixture of torsemide modification II and excipients. The mixture is then dried

5 until only trace amounts of fluid remain in the granulate as residual moisture. Preferably, the mixture is dried to 0.5-1.5% moisture content. The granulate is then sieved, and magnesium stearate is added to the milled granulate. The final blend of torsemide modification II, excipients and magnesium stearate is compressed into tablets on a rotary tableting machine. Table 1 shows suitable ranges of active ingredients and excipients
10 (weight %) and the preferred amounts for the present pharmaceutical formulations.

While not being bound by theory, it is believed that the observed unexpected stability of torsemide modification II (which is not high purity torsemide modification II) in the present pharmaceutical formulation is achieved by the present novel formulation which

15 serves to inhibit the rearrangement of torsemide modification II into torsemide modification I.

Table 1.

Material	Range of % composition (w/w)	Preferred % composition	Function
High purity Torsemide modification II	2.5-25%	5%	active ingredient
Lactose Anhydrous NF	25.5-65%	45.5%	filler
Crospovidone NF	10-15%	12.0%	disintegrant
Povidone USP (PVP K-30)	1-3%	1.5%	binder
Microcrystalline Cellulose NF (Avicel PH 112)	25-45%	35.0%	filler and disintegrant
Alcohol 95% USP*	-	-	Granulation processing solvent
Magnesium Stearate NF	0.5-2.5%	1.0%	lubricant

* Granulation processing solvent only (dried to achieve moisture content of 0.5-1.5%).

Significantly, it has also been found that the pharmaceutical formulations of the present invention containing fine crystals of high purity torsemide modification II have a dissolution rate in water and in potassium phosphate buffer that does not substantially change over time. It has been found that the tablet formulations of the present invention, 5 during storage at 40°C, 75% relative humidity, for 6 weeks, do not undergo any substantial change in the dissolution rate. The dissolution rate was determined by the U.S.P. Paddle Method, 37°C, 90 RPM, 0.01M KH_2PO_4 , pH 4.5; and by the U.S.P. Paddle Method, 37°C, 50 RPM, purified water.

In accordance with the present invention, the pharmaceutical formulations of the 10 present invention are useful for the treatment of hypertension and edema associated with congestive heart failure, renal disease, or hepatic disease. While one of ordinary skill in the art will understand that dosages will vary according to the indication, age of the patient, and other factors, generally the formulations of the present invention will be administered at a daily dosage of the active ingredient between about 2 to about 200 mg per day, and 15 preferably about 5 mg to about 100 mg per day. As torsemide is suitable for once-daily dosing, preferably each unit dosage form will contain between about 5 mg and about 100 mg.

EXAMPLES

20 The pursuant invention will now be further explained in the following examples. However, the present invention should not be construed as limited thereby.

Determination of Polymorphic Content by X-Ray Powder Diffraction Analysis

25 **Bulk High Purity Torsemide Modification II (Bulk Nol 851700100)**

5

Polymorph Content		Length of Storage
Storage Conditions		
55°C	40°C, 75% RH	
	Polymorphic form detected (I or II)	t=0
II		1 week
II	II	2 weeks
II	II	1 month
II	II	2 months
II	II	3 months
II	II	4 months

10

15

Determination of Polymorphic Content by X-Ray Powder Diffraction Analysis

Bulk Torsemide Modification II (II) with trace amounts of Modification I (I) (Bulk No 851700200)

20

25

30

Polymorph Content		Length of Storage
Storage Conditions		
	II>>I (<0.3%)	t=0
II>>I (≈0.4%)		1 week
I+II	II>>I (≈0.5%)	2 weeks
I+II	II>>I (≈1%)	1 month
I+II	II>>I (≈6%)	2 months
I+II	I+II	3 months
I+II	I+II	4 months

Example 1**X-Ray Powder Diffraction (XRPD) Method for the Detection and Quantification of Torsemide Modification I in Torsemide Modification II**

5 1. The present procedure is used for the detection and quantitative determination of the presence of torsemide modification I in tablets wherein the active ingredient is high purity torsemide modification II. The present procedure is also used for the detection and quantitative determination of torsemide modification I in bulk high purity torsemide modification II, which is to be used as the active ingredient in tablets. The present method
10 is based on the unique x-ray powder diffraction pattern of torsemide modification I that is characterized by a strong peak at two-theta $5.7 \pm 0.2^\circ$, the presence of which indicates the presence of torsemide modification I in a sample of high purity torsemide modification II.

2. EQUIPMENT

15 2.1 Instrument: Philips x-ray powder diffractometer. Goniometer model PW 1050/70, Cu-tube, curved graphite monochromator.

2.2 Sample holder: A standard aluminum sample holder with a rectangular cavity 20*15*0.3 mm inside it.

3. RUN PARAMETERS

Scanning range: $2\theta = 4^\circ$ to at least 22°
20 Step: 0.05°
Step duration: 0.5

4. PROCEDURE FOR SAMPLE PREPARATION

4.1 Gently grind a small amount of sample powder in an agate mortar with the pestle.

25 4.2 Fill the rectangular cavity on the sample holder with the powder.

**Stability Results for Torsemide Tablets K-26058 and K-26683
Containing 100 mg of Active Ingredient**

30 Tablets containing 100 mg of high purity torsemide modification II, prepared according to the methods of Example 2, were stored under stressed conditions (40°C , 75% relative humidity). The polymorphic content of torsemide inside the tablet was monitored by x-ray powder diffraction (XRPD) techniques. Representative x-ray powder diffraction

patterns are shown in the Figures. Figure 1 is an x-ray powder diffraction pattern of a high purity torsemide tablet (Batch No. K-26683). Figure 2 is an x-ray powder diffraction pattern of bulk high purity torsemide modification II (API 851700100). Figure 3 is an x-ray powder diffraction pattern of a placebo tablet corresponding to a tablet containing 100 mg of high purity torsemide modification II and therefore contains no torsemide. The XRPD of a 100 mg tablet of Batch No. K-26683 directly following production, t=0, showed XRPD peaks typical of high purity torsemide modification II. The XRPD of the K-26683 tablet following three months of storage at 40°C and 75% relative humidity showed XRPD peaks typical of high purity torsemide modification II and did not show an XRPD peak at 5.7 degrees two-theta, which would indicate the presence of torsemide modification I. Similarly, the XRPD of a 100 mg tablet of Batch No. K-26058 directly following production, t=0, showed XRPD peaks typical of high purity torsemide modification II. The XRPD of the K-26058 tablet following three months of storage at 40°C and 75% relative humidity showed XRPD peaks typical of torsemide modification II and did not show an XRPD peak at 5.7 degrees two-theta, which would indicate the presence of torsemide modification I. The diffraction peaks at 20.4 and the broad peak at about 22.5 degrees two-theta are characteristic of the filler.

Lower dosage tablets, for example, tablets containing 10 mg of high purity torsemide modification II, were stored for 2 months at 40°C, 75% relative humidity, and were monitored by solid state NMR. The resulting solid state NMR data indicated that the high purity torsemide modification II of the lower dose tablets did not substantially rearrange.

Example 2

Manufacturing procedure

In a high speed mixer, high purity torsemide modification II was mixed with lactose anhydrous NF, crospovidone NF, povidone USP, and microcrystalline cellulose NF. Alcohol 95% USP was added to the powder mixture. The wet granulate mixture was dried in a fluid bed drier at 50°C to a loss on drying (LOD) of 0.5-2.0%. The resulting dry granulate of high purity torsemide modification II was then sifted through a 0.8 mm sieve and magnesium stearate NF was added to the milled granulate. The final blend of high purity torsemide modification II, excipients and magnesium stearate was compressed into oval shaped tablets on a rotary tableting machine.

Example 2A. High purity Torsemide Tablets (2.5 mg)

Composition (Batch No. K-26056)	grams per 15,000 tablets
High Purity Torsemide Modification II	37.5
Lactose Anhydrous NF	697.5
Crospovidone NF	150.0
Povidone USP (PVP K-30)	37.5
Microcrystalline Cellulose NF (Avicel PH 112)	52.5
Alcohol USP	500.0
Magnesium Stearate NF	12.8

5

10

Example 2B. High purity Torsemide Tablets (5 mg)

Composition (Batch No. K-26057)	grams per 15,000 tablets
High Purity Torsemide Modification II	75
Lactose Anhydrous NF	697.5
Crospovidone NF	150.0
Povidone USP (PVP K-30)	37.5
Microcrystalline Cellulose NF (Avicel PH 112)	52.5
Alcohol USP	510.0
Magnesium Stearate NF	14.6

15

20

Example 2C. High purity Torsemide Tablets (100 mg)

Composition (Batch No. K-26058)	grams per 3,750 tablets
High Purity Torsemide Modification II	375.0
Lactose Anhydrous NF	547.0
Crospovidone NF	150.0
Povidone USP (PVP K-30)	37.5
Microcrystalline Cellulose NF (Avicel PH 112)	375
Alcohol USP	616.0
Magnesium Stearate NF	15.0

25

30

Example 3
Dissolution Results

The dissolution method used was the U.S.P. Paddle Method, at 90 RPM with 0.1 M KH₂PO₄, pH 4.5 at 37°C. For the dissolution test, 6 tablets were tested in 900 mL of phosphate buffer, pH 4.5, according to the Paddle Method of the U.S.P. Examples 3A, 3B and 3C show the dissolution rates of three tablet lots directly after production and after 6 weeks of storage at 40°C at a relative humidity (RH) of 75%. The dissolution rates of high purity torsemide Form II Batch Nos. K-26056, K-26057 and K-26058 were identical under both conditions. There was no substantial change in the dissolution rates of any of the present pharmaceutical formulations containing torsemide modification II following 6 weeks of the above storage conditions.

Example 3A. Dissolution of 2.5 mg High Purity Torsemide modification II Tablets

Time (minutes)	Torsemide dissolved (%)	
	K-26056 (2.5 mg) directly after production	K-26056 (2.5 mg) after 6 weeks at 40°C/75% RH
15		
15	97	98
30	97	97
45	97	97
60	97	97

Example 3B. Dissolution of 5 mg High Purity Torsemide modification II Tablets

Time (minutes)	Torsemide dissolved (%)	
	K-26057 (5 mg) directly after production	K-26057 (5 mg) after 6 weeks at 40°C/75% RH
15		
15	98	95
30	98	96
45	97	95
60	99	95

Example 3C. Dissolution of 100 mg High Purity Torsemide modification II Tablets

Time (minutes)	Torsemide dissolved (%)

Example 3C. Dissolution of 100 mg High Purity Torsemide modification II Tablets

	K-26058 (100 mg) directly after production	K-26058 (100 mg) after 6 weeks at 40°C/75% RH
15	87	81
30	92	84
45	93	89
5	93	89

Example 4
Dissolution Results

The dissolution method used was the U.S.P. Paddle Method, at 50 RPM with purified water at 37°C. For the dissolution test, 6 tablets were tested in 900 mL of purified water according to the Paddle Method of the U.S.P. Example 4B shows the dissolution rates of one representative tablet lot directly after production and after 3 months of storage at 40°C at a relative humidity (RH) of 75%. The dissolution rates of the high purity torsemide modification II tablet Batch No. K-26683 were identical under both conditions.

10 There was no substantial change in the dissolution rates of any of the present pharmaceutical formulations containing high purity torsemide modification II following 3 months at the above storage conditions.

15

Example 4A. High purity Torsemide Tablets (100 mg)

Composition (Batch No. K-26683)	grams per 3,750 tablets
High purity Torsemide modification II	375.0
Lactose Anhydrous NF	382.5
Crospovidone NF	180.0
Povidone USP (PVP K-30)	22.5
20 Microcrystalline Cellulose NF (Avicel PH 112)	525
Alcohol USP	620.00
25 Magnesium Stearate NF	15.0

Example 4B. Dissolution of 100 mg High purity Torsemide modification II Tablets

Time (minutes)	Torsemide dissolved (%)	
	K-26683 (100 mg) directly after production	K-26683 (100 mg) after 3 months at 40°C/75% RH
15	88	88
30	97	92
45	98	97
60	99	98

10

Although certain presently preferred embodiments of the invention have been described herein, it will be apparent to those skilled in the art to which the invention pertains that variations and modifications of the described embodiment may be made 15 without departing from the spirit and scope of the invention. Accordingly, it is intended that the invention be limited only to the extent required by the appended claims and the applicable rules of law.

WE CLAIM:

1. A stable pharmaceutical formulation comprising an effective amount of high purity torsemide modification II and a pharmaceutically acceptable carrier.
2. The stable pharmaceutical formulation of claim 1 wherein said formulation further comprises lactose anhydrous, crospovidone, povidone, cellulose, and magnesium stearate.
3. The stable pharmaceutical formulation of claim 1 wherein said formulation is a tablet.
4. The stable pharmaceutical formulation of claim 7 wherein the high purity torsemide modification II is present in an amount of about 2.5 to 200 mg per tablet.
5. The stable pharmaceutical formulation of claim 8 wherein the high purity torsemide modification II is present in an amount of 100 mg per tablet.
6. The stable pharmaceutical formulation of claim 9 wherein the high purity torsemide modification II is present in an amount of 5 mg per tablet.
7. The stable pharmaceutical formulation of claim 10 wherein the high purity torsemide modification II is present in an amount of 2.5 mg per tablet.
8. The stable pharmaceutical formulation of claim 1 wherein the high purity torsemide modification II does not substantially rearrange over time.
9. The stable pharmaceutical formulation of claim 8 wherein the high purity torsemide modification II does not substantially rearrange into torsemide modification I over time.
10. The stable pharmaceutical formulation of claim 1 wherein the torsemide modification II has a particle size distribution wherein 100 % is below 200 μ .
11. The stable pharmaceutical formulation of claim 10 wherein the torsemide modification II has a particle size distribution wherein 100% is below 100 μ .
12. The stable pharmaceutical formulation of claim 11 wherein the torsemide modification II has a particle size distribution wherein 100% is below 50 μ .
13. A stable pharmaceutical formulation comprising a therapeutically effective amount of torsemide modification II wherein the dissolution rate in vitro of the formulation, when measured by the U.S.P. Paddle Method at 50-90 RPM in 900 mL water is not less than 80% (by weight) of the high purity torsemide modification II released after

30 minutes.

14. The stable pharmaceutical formulation of claim 13 wherein the dissolution rate in vitro does not substantially change over time.

15. The stable pharmaceutical formulation of claim 14 wherein the dissolution rate in vitro does not substantially change for at least 3 months.

16. A method for treating edema comprising the step of administering to a patient in need of such treatment a therapeutically effective amount of the pharmaceutical formulation of claim 1.

17. A method for treating edema comprising the step of administering to a patient in need of such treatment a therapeutically effective amount of the pharmaceutical formulation of claim 16.

1/3




Fig. 1

2/3




Fig. 2

3/3




Fig. 3