

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017206077 B2**

- (54) Title
Mesoporous silica nanoparticles with lipid bilayer coating for cargo delivery
- (51) International Patent Classification(s)
A61K 9/51 (2006.01) **A61K 47/50** (2017.01)
A61K 31/337 (2006.01)
- (21) Application No: **2017206077** (22) Date of Filing: **2017.01.06**
- (87) WIPO No: **WO17/120537**
- (30) Priority Data
- (31) Number (32) Date (33) Country
62/276,634 **2016.01.08** **US**
- (43) Publication Date: **2017.07.13**
(44) Accepted Journal Date: **2021.11.18**
- (71) Applicant(s)
The Regents of the University of California
- (72) Inventor(s)
Nel, Andre E.;Meng, Huan;Liu, Xiangsheng
- (74) Agent / Attorney
Pizzseys Patent and Trade Mark Attorneys Pty Ltd, GPO Box 1374, Brisbane, QLD, 4001, AU
- (56) Related Art
WO 2014138278 A1
D. C. DRUMMOND ET AL, "Development of a Highly Active Nanoliposomal Irinotecan Using a Novel Intraliposomal Stabilization Strategy", CANCER RESEARCH, US, (2006-03-15), vol. 66, no. 6, pages 3271 - 3277
US 20070116753 A1
HUAN MENG ET AL, "Use of a Lipid-Coated Mesoporous Silica Nanoparticle Platform for Synergistic Gemcitabine and Paclitaxel Delivery to Human Pancreatic Cancer in Mice", ACS NANO, US, (2015-04-28), vol. 9, no. 4, pages 3540 - 3557
DERRICK TARN ET AL, "Mesoporous Silica Nanoparticle Nanocarriers: Biofunctionality and Biocompatibility", ACCOUNTS OF CHEMICAL RESEARCH, (2013-03-19), vol. 46, no. 3, doi:10.1021/ar3000986, ISSN 0001-4842, pages 792 - 801



(51) International Patent Classification:

A61K 9/51 (2006.01) A61K 31/337 (2006.01)
A61K 47/50 (2017.01)

(21) International Application Number:

PCT/US2017/012625

(22) International Filing Date:

6 January 2017 (06.01.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/276,634 8 January 2016 (08.01.2016) US

(71) Applicant: THE REGENTS OF THE UNIVERSITY
OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th
Floor, Oakland, California 94607-5200 (US).(72) Inventors: NEL, Andre E.; 4212 Murietta Avenue, Sher-
man Oaks, California 91423 (US). MENG, Huan; 3340
Sawtelle Boulevard, #204, Los Angeles, California 90066
(US). LIU, Xiangsheng; 3230 Overland Avenue, #212,
Los Angeles, California 90034 (US).(74) Agents: HUNTER, Tom et al.; P.O. Box 70250, Oakland,
California 94612-0250 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: MESOPOROUS SILICA NANOPARTICLES WITH LIPID BILAYER COATING FOR CARGO DELIVERY




FIG. 5

(57) Abstract: A nanocarrier including a silica body having a surface and defining a plurality of pores that are suitable to receive molecules therein is described. The nanocarrier also includes a lipid bilayer coating the surface, and a cargo-trapping agent within the phospholipid bilayer. The phospholipid bilayer stably seals the plurality of pores. The cargo-trapping reagent can be selected to interact with a desired cargo, such as a drug.

MESOPOROUS SILICA NANOPARTICLES WITH LIPID BILAYER COATING FOR CARGO DELIVERY

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit of and priority to USSN 62/276,634, filed on
5 January 8, 2016, which is incorporated herein by reference in its entirety for all purposes.

STATEMENT OF GOVERNMENTAL SUPPORT

[0002] This invention was made with Government support under Grant Nos. R01
CA133697 and U01 CA198846, awarded by the National Institutes of Health/National
Cancer Institute. The Government has certain rights in the invention.

10

BACKGROUND

[0003] Pancreatic ductal adenocarcinoma (PDAC) is a fatal disease with a 5-year
survival rate of less than 6% (Siegel *et al.* (2014) *CA Cancer J. Clin.* 64(1): 9-29).
Currently, the major treatment regimens for chemotherapy include either a single reagent,
gemcitabine (GEM), or a four-drug regimen,. While FOLFIRINOX has a better response
15 rate than GEM (31.6% *versus* 9.4%), with improved survival (11 months *versus* 6.8
months), the former combination is significantly more toxic and restricted to a minority of
PDAC patients with good performance status (Conroy *et al.* (2011) *N. Engl. J. Med.*
364(19): 1817-1825). Irinotecan contributes significantly to this toxicity, including a severe
impact on the bone marrow (*e.g.* neutropenia), liver (*e.g.* necrosis and steatosis) and the
20 gastrointestinal (GI) tract (*e.g.* vomiting, diarrhea) (Conroy *et al.* (2011) *N. Engl. J. Med.*
364(19): 1817-1825; Ueno *et al.* (2007) *Cancer Chemother. Pharmacol.* 59(4): 447-454;
Loupakis *et al.* (2013) *Br. J. Cancer*, 108(12): 2549-2556). Thus, there is a great demand
for a treatment regimen that improves irinotecan toxicity, with a view to improving the
available drugs for first-line therapy in PDAC.

25 [0004] One approach to reducing irinotecan toxicity, with maintenance of efficacy,
is high-dose drug encapsulation in a nanocarrier with protected delivery to the cancer site
while reducing systemic drug release. Different carrier types, including polymeric particles
and liposomes, have been employed with some success for irinotecan delivery (Chou *et al.*
(2003) *J. Biosci. Bioeng.*, 95(4): 405-408; Onishi *et al.* (2003) *Biol. Pharmaceut. Bull.*
30 26(1): 116-119; Messerer *et al.* (2004) *Clin. Cancer Res*, 10(19): 6638-6649; Drummond *et*

al. (2006) *Cancer Res.*, 66(6): 3271-3277; Valencia *et al.* (2013) *Nanomed.* 8(5): 687-698; Sadzuka *et al.* (1998) *Cancer Lett.*, 127(1): 99-106; Ramsay *et al.* (2008) *Eur. J. Pharm. Biopharm.* 68(3): 607-617; Li *et al.* (2015) *Adv. Func. Mat.* 25(5): 788-798). However, while polymeric nanoparticles showed promising *in vitro* results, the limited capacity to load irinotecan (<1%, w/w) plus premature drug release (*e.g.*, 40% in 5 hours), the nanoparticles did not achieve the required toxicity reduction while improving intratumoral drug delivery (Valencia *et al.* (2013) *Nanomed.* 8(5): 687-698). While liposomes could achieve high irinotecan loading capacity through the use of ammonium sulfate or proton entrapment agents (Chou *et al.* (2003) *J. Biosci. Bioeng.*, 95(4): 405-408; Messerer *et al.* (2004) *Clin. Cancer Res.*, 10(19): 6638-6649; Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277; Sadzuka *et al.* (1998) *Cancer Lett.*, 127(1): 99-106; Ramsay *et al.* (2008) *Eur. J. Pharm. Biopharm.* 68(3): 607-617), carrier instability under shear and osmotic stress, as well as bilayer disruption by serum proteins, resulted in premature drug release and toxicity (Liu *et al.* (2000) In: *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 172(1-3): 57-67; Heurtault *et al.* (2003) *Biomaterials*, 24(23): 4283-4300; Sabin *et al.* (2006) *Eur. Phys. J. E.* 20(4): 401-408).

[0005] As noted above, many attempts to deliver cancer drugs in clinical trials or in a therapeutic setting have been based on liposomal (Messerer *et al.* (2004) *Clin. Cancer Res.*, 10(19): 6638-6649; *Cancer Res.* 2006, 66, 3271) or polymer-based systems (Onishi *et al.* (2003) *Biol. Pharm. Bull.* 26(1): 116-119). Most of these carriers are spherical particles or supramolecular assemblies in the size range of 80-200 nm, often containing PEG coating on the surface to prolong circulatory half-life, and typically exhibiting loading capacities from ~5 w/w% (*e.g.*, polymer-based nanoparticles) to ~50 w/w% (*e.g.*, liposomal carrier). At a pre-clinical level, the potential benefits of these nanocarriers in animal studies, including murine PDAC models, have been shown to include a reduction in *in vivo* toxicity, enhanced antitumor efficacy, and improved survival rate.

[0006] However, only a small number of nanocarriers have advanced to clinical trials for PDAC patients. Nanocarriers including an ionophore (A23187, also known as calimycin) enabled irinotecan delivery liposomal formulation (Irinophore C) and a protonating agent irinotecan delivering liposomal formulation (MM-398) (Baker *et al.* (2008) *Clin. Cancer Res.* 14: 7260-7271; Drummond *et al.* (2006) *Cancer Res.* 15(66): 3271-3277). The Irinophore C formulation (Champions Biotechnology) is a liposomal carrier that makes use of active irinotecan loading through the generation of transmembrane

proton gradients, using the ionophore, A23187, or ammonium sulfate (Ramsay *et al.* (2008) *Eur. J. Pharm. Biopharm.* 68: 607-617). The Irinophore C formulation was used in a clinical study that commenced in 2011, but there have been no updated information about the outcome of the study or the results.

5 **[0007]** While in a recent phase 3 clinical trial, the development of a liposomal carrier (MM-398) for irinotecan by Merrimack showed an improved survival benefit in PDAC as a 2nd-line treatment option, the relatively high rate of GI tract and bone marrow toxicity has resulted in a black box warning for severe and life-threatening diarrhea and neutropenia (Von Hoff *et al.* (2013) *Br. J. Cancer*, 109(4): 920-925;
10 www.fda.gov/newsevents/newsroom/pressannouncements/ucm468654.htm). Human subjects participating in MM-398 clinical trials also showed significant elevations of liver enzymes, including alanine aminotransferase (ALT) (*see, e.g.*,
www.accessdata.fda.gov/drugsatfda_docs/label/2015/207793LB.pdf). Nonetheless, MM-398 received FDA approval for use in PDAC for patients failing to respond to GEM
15 therapy, and is marketed as Onivyde® (*see, e.g.*,
www.fda.gov/newsevents/newsroom/pressannouncements/ucm468654.htm).

[0008] The MM-398 liposomal formulation (Merrimack Pharmaceuticals) incorporates irinotecan hydrochloride with the assistance of a polyanionic trapping agent (ESMO GI 2014, www.merrimackpharma.com). More specifically, irinotecan loading into
20 the MM-398 liposome was achieved by intra-liposomal drug encapsulation of a multivalent anionic trapping agent, triethylammonium sucrose octasulfate (TEA₈SOS). This chemical leads to irinotecan protonation and entrapment at more than 10 times the loading that can be achieved through passive drug encapsulation (Drummond *et al.* (2006) *Cancer Res.* 66(6): 3271-3277). IV administration of MM-398 liposome has been shown to induce complete
25 tumor regression in various PDAC tumor models in mice, including inhibition of metastatic tumor foci (Paz *et al.* (2102) *Cancer Res.* 72(12 Suppl): Abstract A63). MM-398 is currently in a Phase 3 clinical trial and lays claim to providing improved tumor inhibition, pharmacokinetics and efficacy compared to free irinotecan in animal and human studies (Kalra (2012) *AACR meeting*, Abstract #5622). This includes experimental data claiming
30 complete PDAC regression using a dose of 20 mg/kg MM-398 (human equivalent dose 60-120 mg/m²) in murine xenograft studies (Paz *et al.* (2102) *Cancer Res.* 72(12 Suppl): Abstract A63). MM-398 also increases the maximum tolerated dose (MTD) of free irinotecan from 80 to 324 mg/kg in mice (Drummond *et al.* (2006) *Cancer Res.* 66(6): 3271-

3277). Additionally, in the Phase 3 clinical trial by Merrimack Pharmaceuticals (Hoff *et al.* ESMO GI 2014, www.merrimackpharma.com) involving 417 PDAC patients, the combination of MM-398, 5-FU and leucovorin resulted in an overall survival (OS) of 6.1 months, which is 1.9 months longer than the control arm receiving 5-FU and leucovorin.

5 However, while the active loading of irinotecan into the MM-398 liposomal formulation enhances drug loading capacity over a passive encapsulation procedure, the synthesis technique requires multiple steps and liposomal carriers do not provide the same colloidal stability or the same amount of intracellular release compared to the LB-MSNP platform provided herein. Nonetheless, MM-398 received FDA approval for use in PDAC for
10 patients failing to respond to GEM therapy, and is marketed as ONIVYDE®. The use of polyanionic polymers to increase drug entrapment in liposomes leads to ~80 nm drug precipitation (Zhu *et al.* (1996) 39(1): 138-142; Colbern *et al.* (1998) *Clin. Cancer Res.* 4(12): 3077-3082), which constitutes one of the reasons for slow irinotecan release from the liposomal carrier compared to LB-MSNP pores.

15 **[0009]** Thus, there is still an unmet need for nanocarriers and delivery methods that enable efficient drug delivery, including chemotherapy such as irinotecan chemotherapy, with an improved margin of safety and reduced toxicity.

SUMMARY

[0010] Urgent intervention is required to improve the 5-year survival rate of
20 pancreatic ductal adenocarcinoma (PDAC). While the 4-drug regimen, FOLFIRINOX (comprised of irinotecan (IRIN), 5-fluorouracil (5-FU), oxaliplatin (OX), and leucovorin (LV)), has a better survival outcome than the more frequently used gemcitabine (GEM), the former treatment regimen is highly toxic and restricted for use in a patient with good performance status. Since irinotecan contributes significantly to FOLFIRINOX toxicity
25 (bone marrow and gastrointestinal tract), one specific aim of the present invention is to reduce the toxicity of the former drug with a custom-designed mesoporous silica nanoparticle (MSNP) platform, which uses a proton gradient for high-dose irinotecan loading across a coated lipid bilayer (LB). The improved stability of the LB-coated MSNP (LB-MSNP) carrier allows for better protected irinotecan delivery and increased tumor drug
30 concentrations when compared, for example, to a liposomal equivalent in a Kras-derived orthotopic PDAC model in immunocompetent mice. The LB-MSNP nanocarrier is also more efficient for treating tumor metastases. Equally important, the reduced leakage and slower rate of drug release by the LB-MSNP carrier dramatically reduces the rate of bone

marrow, gastrointestinal and liver toxicity compared to a liposomal carrier. The combination of high efficacy and reduced toxicity by the LB-MSNP carrier facilitates the use of irinotecan as a first-line therapeutic to improve PDAC survival.

[0011] Generally, in one aspect of the present invention, a nanocarrier is provided comprising a silica body having a surface and defining a plurality of pores that are suitable to receive molecules therein, a lipid bilayer (*e.g.*, a phospholipid bilayer) coating the surface, and a cargo-trapping agent contained in the pores by the coated lipid bilayer, where the submicron structure has a maximum dimension of less than one micron, and wherein the phospholipid bilayer stably seals the plurality of pores and can serve as a basis for additional packaging, targeting, imaging and functionalization.

[0012] One significant aspect of the present invention includes new methods for making improved cargo/agent (*e.g.* irinotecan or other molecule(s) with the chemical structures described herein) nanocarriers. These methods generally include providing an unloaded nanocarrier comprising a silica body having a surface with a plurality of pores that are suitable to receive molecules therein, and encapsulating a cargo-trapping, targeting or imaging agent within the pores by a lipid bilayer (*e.g.*, a phospholipid bilayer). As discussed in detail below, working examples include, but are not limited to new methods of making concentrated irinotecan nanocarriers. In certain embodiments these methods include selecting a nanocarrier comprising a silica body having a surface including a plurality of pores suitable to receive irinotecan therein. In these methods, an agent having an ability to specifically influence the diffusion of and/or trap the irinotecan molecule (*e.g.* triethylammonium sucrose octasulfate) is also selected and disposed within the plurality of pores. The nanocarrier and pores are completely coated with a lipid bilayer (for example using a sonication process). Typically the silica body nanocarrier is not dried and/or washed immediately prior to this coating step. Optionally, in certain embodiments, this lipid bilayer can comprise one or more bioactive molecules selected to facilitate the nanocarrier function (such as polyethylene glycol, and/or targeting ligands, and/or paclitaxel and/or activated irinotecan, SN38). Following this coating step, irinotecan is then allowed to migrate across the lipid bilayer into the pores where it is entrapped. This is also known as “remote drug loading”. In this way, irinotecan nanocarriers with surprisingly high loading capacities (*e.g.* 40 wt% or greater than 40% drug/MSNP) can be formed. In addition to these high drug loading capacities, irinotecan nanocarriers formed using these

methods have a constellation of other desirable properties, for example exhibiting <5% irinotecan leakage over 24 hours in a biological buffer with pH of 7.4 at 37 °C.

[0013] Various embodiments contemplated herein may include, but need not be limited to, one or more of the following:

5 **[0014]** Embodiment 1: A nanoparticle drug carrier comprising:

[0015] a silica nanoparticle having a surface and defining a plurality of pores that are suitable to receive molecules therein;

[0016] a lipid bilayer coating the surface;

[0017] a cargo-trapping agent within pores comprising said plurality of
10 pores; and

[0018] a cargo comprising a drug, where said cargo is associated with said cargo-trapping agent in said pores;

[0019] wherein the submicron structure has a maximum dimension of less than one micron, and wherein the lipid bilayer stably seals the plurality of pores.

15 **[0020]** Embodiment 2: The nanoparticle drug carrier of embodiment 1, wherein said lipid bilayer comprises a phospholipid, cholesterol (CHOL), and an mPEG phospholipid.

[0021] Embodiment 3: The nanoparticle drug carrier according to any one of
20 embodiments 1-2, wherein said phospholipid comprises a saturated fatty acid with a C14-C20 carbon chain, and/or an unsaturated fatty acid with a C14-C20 carbon chain, and/or a natural lipid comprising a mixture of fatty acids with C12-C20 carbon chains.

[0022] Embodiment 4: The nanoparticle drug carrier of embodiment 3, wherein
25 said phospholipid comprises a saturated fatty acid selected from the group consisting of phosphatidylcholine (DPPC), dimyristoylphosphatidylcholine (DMPC), distearoylphosphatidylcholine (DSPC), and diactylphosphatidylcholine (DAPC).

[0023] Embodiment 5: The nanoparticle drug carrier of embodiment 3, wherein said phospholipid comprises a natural lipid selected from the group consisting of egg phosphatidylcholine (egg PC), and soy phosphatidylcholine (soy PC).

[0024] Embodiment 6: The nanoparticle drug carrier of embodiment 3, wherein
30 said phospholipid comprises an unsaturated fatty acid selected from the group consisting of 1,2-dimyristoleoyl-sn-glycero-3-phosphocholine, 1,2-dipalmitoleoyl-sn-glycero-3-

phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), and 1,2-dieicosenoyl-sn-glycero-3-phosphocholine.

5 [0025] Embodiment 7: The nanoparticle drug carrier according to any one of embodiments 1-6, wherein said lipid bilayer comprises an mPEG phospholipid with a phospholipid C14-C18 carbon chain, and a PEG molecular weight ranging from about 350 Da to 5000 Da.

[0026] Embodiment 8: The nanoparticle drug carrier of embodiment 7, wherein said lipid bilayer comprises 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-PEG (DSPE-PEG).

10 [0027] Embodiment 9: The nanoparticle drug carrier of embodiment 2, wherein said lipid bilayer comprises DPPC/Chol/DSPE-PEG or DSPC/Chol/DSPE-PEG.

[0028] Embodiment 10: The nanoparticle drug carrier of embodiment 9, wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG.

15 [0029] Embodiment 11: The nanoparticle drug carrier of embodiment 10, wherein wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG2000.

[0030] Embodiment 12: The nanoparticle drug carrier according to any one of embodiments 1-11, wherein said lipid bilayer comprises a phospholipid, cholesterol, and mPEG phospholipid at a ratio of 50-90 mol% phospholipid : 10-50 mol% CHOL : 1-10 mol% mPEG phospholipid.

20 [0031] Embodiment 13: The nanoparticle drug carrier of embodiment 10, wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG in a molar ratio of about 3:2:0.15.

[0032] Embodiment 14: The nanoparticle drug carrier according to any one of embodiments 1-13, wherein said lipid bilayer forms a substantially continuous bilayer encompassing the entire nanoparticle.

25 [0033] Embodiment 15: The nanoparticle drug carrier according to any one of embodiments 1-14, wherein said lipid bilayer forms a substantially uniform and intact bilayer encompassing the entire nanoparticle.

[0034] Embodiment 16: The nanoparticle drug carrier according to any one of embodiments 1-15, wherein said silica nanoparticle is a mesoporous silica nanoparticle.

[0035] Embodiment 17: The nanoparticle drug carrier of embodiment 16, wherein said silica nanoparticle comprises a sol-gel synthesized mesoporous silica nanoparticle.

[0036] Embodiment 18: The nanoparticle drug carrier according to any one of embodiments 16-17, wherein said mesoporous silica nanoparticle is size-controlled.

5 [0037] Embodiment 19: The nanoparticle drug carrier according to any one of embodiments 16-18, wherein said mesoporous silica nanoparticle is is colloidally stable.

[0038] Embodiment 20: The nanoparticle drug carrier according to any one of embodiments 16-19, wherein said mesoporous silica has an average pore size that ranges from about 1 to about 20 nm, or from about 1 to about 10 nm, or from about 2 to about 8
10 nm.

[0039] Embodiment 21: The nanoparticle drug carrier according to any one of embodiments 1-20, wherein said mesoporous silica nanoparticles has an average size ranging from about 50 nm up to about 300 nm, or from about 50 up to about 200 nm, or from about 50 up to about 150 nm, or from about 50 up to about 100 nm, or from about 50
15 up to about 80 nm, or from about 50 up to about 70 nm, or from about 60 up to about 70 nm.

[0040] Embodiment 22: The nanoparticle drug carrier according to any one of embodiments 1-21, wherein cargo-trapping agent before reaction with said drug is selected from the group consisting of triethylammonium sucrose octasulfate (TEA_8SOS),
20 $(\text{NH}_4)_2\text{SO}_4$, an ammonium salt, a trimethylammonium salt, and a triethylammonium salt.

[0041] Embodiment 23: The nanoparticle drug carrier of embodiment 22, wherein said cargo trapping agent comprises $(\text{NH}_4)_2\text{SO}_4$.

[0042] Embodiment 24: The nanoparticle drug carrier of embodiment 22, wherein said cargo trapping agent comprises an ammonium salt selected from the group consisting
25 of ammonium sulfate; ammonium sucrose octasulfate, ammonium α -cyclodextrin sulfate, ammonium β -cyclodextrin sulfate, , ammonium γ -cyclodextrin sulfate, ammonium phosphate; ammonium α -cyclodextrin phosphate, ammonium β -cyclodextrin phosphate, ammonium γ -cyclodextrin phosphate, ammonium citrate, and ammonium acetate.

[0043] Embodiment 25: The nanoparticle drug carrier of embodiment 22, wherein
30 said cargo trapping agent comprises a trimethylammonium salt selected from the group consisting of trimethylammonium sulfate, trimethylammonium sucrose octasulfate,

trimethylammonium α -cyclodextrin sulfate, trimethylammonium β -cyclodextrin sulfate, trimethylammonium γ -cyclodextrin sulfate, trimethylammonium phosphate, trimethylammonium α -cyclodextrin phosphate, trimethylammonium β -cyclodextrin phosphate, trimethylammonium γ -cyclodextrin phosphate, trimethylammonium citrate, and
5 trimethylammonium acetate.

[0044] Embodiment 26: The nanoparticle drug carrier of embodiment 22, wherein said cargo trapping agent comprises a triethylammonium salt selected from the group consisting of triethylammonium sulfate, triethylammonium sucrose octasulfate, triethylammonium α -Cyclodextrin sulfate, triethylammonium β -Cyclodextrin sulfate,
10 triethylammonium γ -Cyclodextrin sulfate, triethylammonium phosphate, triethylammonium α -Cyclodextrin phosphate, triethylammonium β -Cyclodextrin phosphate, triethylammonium γ -Cyclodextrin phosphate, triethylammonium citrate, and triethylammonium acetate.

[0045] Embodiment 27: The nanoparticle drug carrier of embodiment 22, wherein cargo-trapping agent before reaction with said drug is triethylammonium sucrose octasulfate
15 (TEA₈SOS).

[0046] Embodiment 28: The nanoparticle drug carrier of embodiment 27, wherein said drug is protonated and trapped in said pores as a gel-like precipitate in association of SOS⁸⁻.

[0047] Embodiment 29: The nanoparticle drug carrier according to any one of
20 embodiments 1-28, wherein said drug comprises at least one weakly basic group capable of being protonated, and the cargo-trapping agent comprises at least one anionic group.

[0048] Embodiment 30: The nanoparticle drug carrier according to any one of embodiments 1-29, wherein said drug is selected to have a pKa greater than 7 and less than 11.

25 **[0049]** Embodiment 31: The nanoparticle drug carrier according to any one of embodiments 1-30, said drug comprises a primary, secondary, and tertiary amine.

[0050] Embodiment 32: The nanoparticle drug carrier according to any one of embodiments 1-31, wherein said drug is selected to have a water solubility index of about 5 to about 25 mg/mL.

[0051] Embodiment 33: The nanoparticle drug carrier according to any one of embodiments 1-32, wherein the cargo is selected to have an octanol/water partition coefficient or logP value of about -3.0 to about 3.0.

5 [0052] Embodiment 34: The nanoparticle drug carrier according to any one of embodiments 1-33, wherein the cargo is selected to be 2-8 nm and less than the average or median size of the pores of the silica nanoparticle.

[0053] Embodiment 35: The nanoparticle drug carrier according to any one of embodiments 29-34, wherein said cargo comprises an anti-cancer drug.

10 [0054] Embodiment 36: The nanoparticle drug carrier of embodiment 35, wherein said cargo comprises irinotecan.

[0055] Embodiment 37: The nanoparticle drug carrier of embodiment 35, wherein said cargo comprises one or more drugs independently selected from the group consisting of a topoisomerase inhibitor, an antitumor anthracycline antibiotic, a mitotic inhibitor, an alkaloid, an alkaline alkylating agent, a purine or pyrimidine derivative, and a protein kinase
15 inhibitor.

[0056] Embodiment 38: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises a topoisomerase inhibitor comprising topotecan.

[0057] Embodiment 39: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises an alkaloid selected from the group consisting of topotecan, 10-
20 hydroxycamptothecin, belotecan, rubitecan, vinorelbine, and LAQ824.

[0058] Embodiment 40: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises an antitumor anthracycline antibiotic selected from the group consisting of doxorubicin, and mitoxantrone.

25 [0059] Embodiment 41: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises a mitotic inhibitor selected from the group consisting of vinblastine, and vinorelbine.

[0060] Embodiment 42: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises an alkaline alkylating agent selected from the group consisting of cyclophosphamide, mechlorethamine, and temozolomide.

[0061] Embodiment 43: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises a purine or pyrimidine derivative selected from the group consisting of 5-fluorouracil, 5'-deoxy-5-fluorouridine, and gemcitabine.

5 [0062] Embodiment 44: The nanoparticle drug carrier of embodiment 37, wherein said carrier comprises a protein kinase inhibitor selected from the group consisting of imatinib, osimertinib and sunitinib pazopanib, enzastaurin, vandetanib, erlotinib, dasatinib, and nilotinib.

10 [0063] Embodiment 45: The nanoparticle drug carrier according to any one of embodiments 1-44, wherein said drug carrier is conjugated to a moiety selected from the group consisting of a targeting moiety, a fusogenic peptide, and a transport peptide.

[0064] Embodiment 46: The nanoparticle drug carrier of embodiment 45, wherein said drug carrier is conjugated to a peptide that binds a receptor on a cancer cell or tumor blood vessel.

15 [0065] Embodiment 47: The nanoparticle drug carrier of embodiment 46, wherein said drug carrier is conjugated to an iRGD peptide.

[0066] Embodiment 48: The nanoparticle drug carrier of embodiment 46, wherein said drug carrier is conjugated to a targeting peptide shown in Table 2.

20 [0067] Embodiment 49: The nanoparticle drug carrier according to any one of embodiments 45-48, wherein said drug carrier is conjugated to transferrin, and/or ApoE, and/or folate.

[0068] Embodiment 50: The nanoparticle drug carrier according to any one of embodiments 45-49, wherein said drug carrier is conjugated to a targeting moiety that comprises an antibody that binds to a cancer marker.

25 [0069] Embodiment 51: The nanoparticle drug carrier of embodiment 50, wherein said drug carrier is conjugated to a targeting moiety that comprises an antibody that binds a cancer marker shown in Table 1.

[0070] Embodiment 52: The nanoparticle drug carrier according to any one of embodiments 29-34, wherein said cargo comprises an antibiotic, an antiviral agent, or an antifungal agent.

[0071] Embodiment 53: The nanoparticle drug carrier of embodiment 52, wherein said cargo comprises an antibiotic selected from the group consisting of ciprofloxacin, and levofloxacin.

5 [0072] Embodiment 54: The nanoparticle drug carrier of embodiment 52, wherein said cargo comprises an HIV antiviral.

[0073] Embodiment 55: The nanoparticle drug carrier of embodiment 54, wherein said cargo comprises an antiviral selected from the group consisting of tenofovir, disoproxil, and fumarate.

10 [0074] Embodiment 56: The nanoparticle drug carrier of embodiment 52, wherein said cargo comprises an antifungal agent selected from the group consisting of Amphotericin B, Anidulafungin, Caspofungin, Fluconazole, Flucytosine, Isavuconazole, Itraconazole, Micafungin, Posaconazole, and Voriconazole.

15 [0075] Embodiment 57: The nanoparticle drug carrier according to any one of embodiments 1-56, wherein said drug carrier has less than about 20%, or less than about 15%, or less than about 10%, or less than about 5% leakage of the cargo over 24 hours in a biological buffer with pH of 7.4 at 37 °C.

20 [0076] Embodiment 58: The nanoparticle drug carrier according to any one of embodiments 1-57, wherein said drug carrier has a drug loading capacity of at least about 8% w/w, or at least about 10% w/w, or at least about 20% w/w, or at least about 30% w/w, or greater than about 40% w/w, or greater than about 50% w/w, or greater than about 60% w/w, or greater than about 70% w/w, or greater than about 80% w/w.

[0077] Embodiment 59: The nanoparticle drug carrier according to any one of embodiments 1-57, wherein said drug carrier has a drug loading capacity of at least 80% w/w.

25 [0078] Embodiment 60: The nanoparticle drug carrier according to any one of embodiments 1-59, wherein the lipid bilayer comprises a hydrophobic drug.

30 [0079] Embodiment 61: The nanoparticle drug carrier of embodiment 60, wherein the lipid bilayer comprises a hydrophobic drug selected from the group consisting of paclitaxel, ellipticine, camptothecin, SN-38, and a lipid prodrug (e.g., acyclovir diphosphate dimyristoylglycerol, doxorubicin conjugated phospholipid prodrug, phospholipid derivatives of nucleoside analogs, phospholipid linked chlorambucil, and the like).

[0080] Embodiment 62: The nanoparticle drug carrier of embodiment 60, wherein the lipid bilayer comprises paclitaxel.

[0081] Embodiment 63: The nanoparticle drug carrier according to any one of embodiments 1-61, wherein said drug carriers in suspension are stable for at least 1 month,
5 or at least 2 months, or at least 3 months, or at least 4 months, or at least 5 months, or at least 6 months when stored at 4°C.

[0082] Embodiment 64: The nanoparticle drug carrier according to any one of embodiments 1-63, wherein a population of said drug carriers in suspension shows a size distribution ranging in width (full width half maximum) of less than about 30nm, or less
10 than about 20 nm, or less than about 10 nm, or less than about 5 nm, or less than about 3 nm, or less than about 2 nm.

[0083] Embodiment 65: The nanoparticle drug carrier according to any one of embodiments 1-64, wherein a population of said drug carriers in suspension shows a substantially unimodal size distribution.

15 [0084] Embodiment 66: The nanoparticle drug carrier according to any one of embodiments 1-65, wherein a population of said drug carriers in suspension shows PDI less than about 0.2, or less than about 0.1.

[0085] Embodiment 67: The nanoparticle drug carrier according to any one of embodiments 1-66, wherein a population of said drug carriers in suspension shows a
20 coefficient of variation in size less than about 0.1 or less than about 0.05, or less than about 1.7/120.

[0086] Embodiment 68: The nanoparticle drug carrier according to any one of embodiments 1-67, wherein ~ 3% or more of said nanoparticle drug carriers distribute to developing tumor sites on IV injection.

25 [0087] Embodiment 69: The nanoparticle drug carrier according to any one of embodiments 1-68, wherein said nanoparticle drug carrier forms a stable suspension on rehydration after lyophilization.

[0088] Embodiment 70: The nanoparticle drug carrier according to any one of embodiments 1-69, wherein said nanoparticle drug carriers, when loaded with an anti-
30 cancer drug, provide more effective cancer cell killing than free drug, or liposomes containing said drug, in an orthotopic PDAC model.

[0089] Embodiment 71: The nanoparticle drug carrier according to any one of embodiments 1-70, wherein said nanoparticle drug carriers, when loaded with an anti-cancer drug, show reduced drug toxicity as compared to free drug and/or drug in liposomes.

5 [0090] Embodiment 72: The nanoparticle drug carrier according to any one of embodiments 1-71, wherein said nanoparticle drug carrier has colloidal stability in physiological fluids with pH 7.4 and remains monodisperse to allow systemic biodistribution and is capable of entering a disease site by vascular leakage (EPR effect) or transcytosis.

10 [0091] Embodiment 73: A pharmaceutical formulation said formulation comprising:

[0092] a plurality of nanoparticle drug carriers according to any one of embodiments 1-72; and

[0093] a pharmaceutically acceptable carrier.

15 [0094] Embodiment 74: The formulation of embodiment 73, wherein said formulation is an emulsion, dispersion, or suspension.

[0095] Embodiment 75: The formulation of embodiment 74, wherein said suspension, emulsion, or dispersion is stable for at least 1 month, or at least 2 months, or at least 3 months, or at least 4 months, or at least 5 months, or at least 6 months when stored at 4°C.

20 [0096] Embodiment 76: The formulation according to any one of embodiments 73-75, wherein the nanoscale drug carriers in said formulation show a substantially unimodal size distribution.

25 [0097] Embodiment 77: The formulation according to any one of embodiments 73-76, wherein the drug carriers in said suspension, emulsion, or dispersion shows a PDI less than about 0.2, or less than about 0.1.

30 [0098] Embodiment 78: The formulation according to any one of embodiments 73-77, wherein said formulation is formulated for administration via a route selected from the group consisting of intravenous administration, intraarterial administration, intracerebral administration, intrathecal administration, oral administration, aerosol administration, administration via inhalation (including intranasal and intratracheal delivery, intracranial administration via a cannula, and subcutaneous or intramuscular depot deposition.

[0099] Embodiment 79: The formulation according to any one of embodiments 73-77, wherein said formulation is a sterile injectable.

[0100] Embodiment 80: The formulation according to any one of embodiments 73-79, wherein said formulation is a unit dosage formulation.

5 [0101] Embodiment 81: A method of treating a cancer, said method comprising:

[0102] administering to a subject in need thereof an effective amount of a nanoparticle drug carrier according to any one of embodiments 1-51 or 57-72, or a pharmaceutical formulation according to any one of embodiments 73-80, where the drug in said nanoparticle drug carrier and/or said pharmaceutical formulation comprises an anti-cancer drug.

[0103] Embodiment 82: The method of embodiment 81, wherein said nanoparticle drug carrier and/or said pharmaceutical formulation is a primary therapy in a chemotherapeutic regimen.

15 [0104] Embodiment 83: The method of embodiment 81, wherein said nanoparticle drug carrier and/or said pharmaceutical formulation is a component in a multi-drug chemotherapeutic regimen.

[0105] Embodiment 84: The method of embodiment 83, wherein said multi-drug chemotherapeutic regimen comprises at least two drugs selected from the group consisting of irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV).

20 [0106] Embodiment 85: The method of embodiment 83, wherein said multi-drug chemotherapeutic regimen comprises at least three drugs selected from the group consisting of irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV).

[0107] Embodiment 86: The method of embodiment 83, wherein said multi-drug chemotherapeutic regimen comprises at least irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV).

25 [0108] Embodiment 87: The method according to any one of embodiments 81-86, wherein said cancer is pancreatic ductal adenocarcinoma (PDAC).

[0109] Embodiment 88: The method according to any one of embodiments 81-86, wherein said cancer is a cancer selected from the group consisting of acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), Adrenocortical carcinoma, AIDS-related cancers (*e.g.*, Kaposi sarcoma, lymphoma), anal cancer, appendix cancer, astrocytomas,

atypical teratoid/rhabdoid tumor, bile duct cancer, extrahepatic cancer, bladder cancer, bone cancer (*e.g.*, Ewing sarcoma, osteosarcoma, malignant fibrous histiocytoma), brain stem glioma, brain tumors (*e.g.*, astrocytomas, brain and spinal cord tumors, brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, central nervous system germ cell tumors, craniopharyngioma, ependymoma, breast cancer, bronchial tumors, burkitt lymphoma, carcinoid tumors (*e.g.*, childhood, gastrointestinal), cardiac tumors, cervical cancer, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative disorders, colon cancer, colorectal cancer, craniopharyngioma, cutaneous t-cell lymphoma, duct cancers *e.g.* (bile, extrahepatic), ductal carcinoma *in situ* (DCIS), embryonal tumors, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, extracranial germ cell tumor, extragonadal germ cell tumor, extrahepatic bile duct cancer, eye cancer (*e.g.*, intraocular melanoma, retinoblastoma), fibrous histiocytoma of bone, malignant, and osteosarcoma, gallbladder cancer, gastric (stomach) cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumors (*e.g.*, ovarian cancer, testicular cancer, extracranial cancers, extragonadal cancers, central nervous system), gestational trophoblastic tumor, brain stem cancer, hairy cell leukemia, head and neck cancer, heart cancer, hepatocellular (liver) cancer, histiocytosis, langerhans cell cancer, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, kaposi sarcoma, kidney cancer (*e.g.*, renal cell, Wilm's tumor, and other kidney tumors), langerhans cell histiocytosis, laryngeal cancer, leukemia, acute lymphoblastic (ALL), acute myeloid (AML), chronic lymphocytic (CLL), chronic myelogenous (CML), hairy cell, lip and oral cavity cancer, liver cancer (primary), lobular carcinoma *in situ* (LCIS), lung cancer (*e.g.*, childhood, non-small cell, small cell), lymphoma (*e.g.*, AIDS-related, Burkitt (*e.g.*, non-Hodgkin lymphoma), cutaneous T-Cell (*e.g.*, mycosis fungoides, Sézary syndrome), Hodgkin, non-Hodgkin, primary central nervous system (CNS)), macroglobulinemia, Waldenström, male breast cancer, malignant fibrous histiocytoma of bone and osteosarcoma, melanoma (*e.g.*, childhood, intraocular (eye)), merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer, midline tract carcinoma, mouth cancer, multiple endocrine neoplasia syndromes, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndromes, Myelogenous Leukemia, Chronic (CML), multiple myeloma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, oral cavity cancer, lip and oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, pancreatic

neuroendocrine tumors (islet cell tumors), papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytoma, pituitary tumor, plasma cell neoplasm, pleuropulmonary blastoma, primary central nervous system (CNS) lymphoma, prostate cancer, rectal cancer, renal cell (kidney) cancer, renal pelvis and ureter, transitional cell cancer, rhabdomyosarcoma, salivary gland cancer, sarcoma (*e.g.*, Ewing, Kaposi, osteosarcoma, rhabdomyosarcoma, soft tissue, uterine), Sézary syndrome, skin cancer (*e.g.*, melanoma, merkel cell carcinoma, basal cell carcinoma, nonmelanoma), small intestine cancer, squamous cell carcinoma, squamous neck cancer with occult primary, stomach (gastric) cancer, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, trophoblastic tumor, ureter and renal pelvis cancer, urethral cancer, uterine cancer, endometrial cancer, uterine sarcoma, vaginal cancer, vulvar cancer, Waldenström macroglobulinemia, and Wilm's tumor.

[0110] Embodiment 89: The method according to any one of embodiments 81-88, wherein said nanoparticle drug carrier is not conjugated to an iRGD peptide and the nanoparticle drug carrier is administered in conjunction with an iRGD peptide.

[0111] Embodiment 90: A method of treating an infection, said method comprising:
[0112] administering to a subject in need thereof an effective amount of a nanoparticle drug carrier according to any one of embodiments 1-34 or 52-56, or a pharmaceutical formulation according to any one of embodiments 73-80, where the drug in said nanoparticle drug carrier and/or said pharmaceutical formulation comprises an antimicrobial drug.

[0113] Embodiment 91: The method of embodiment 81, wherein said infection comprises an infection by a drug-resistant bacterium, virus, or fungus.

[0114] Embodiment 92: The method according to any one of embodiments 81-91, where the nanoparticle drug carrier and/or pharmaceutical formulation is administered via a route selected from the group consisting of intravenous administration, intraarterial administration, intracerebral administration, intrathecal administration, oral administration, aerosol administration, administration via inhalation (including intranasal and intratracheal delivery, intracranial administration via a cannula, and subcutaneous or intramuscular depot deposition.

[0115] Embodiment 93: The method according to any one of embodiments 81-91, where the nanoparticle drug carrier and/or pharmaceutical formulation is administered as an injection, from an IV drip bag, or via a drug-delivery cannula.

5 [0116] Embodiment 94: The method according to any one of embodiments 81-93, wherein said subject is a human.

[0117] Embodiment 95: The method according to any one of embodiments 81-93, wherein said subject is a non-human mammal.

10 [0118] Embodiment 96: A method of making a nanoparticle drug carrier, said method comprising: providing a nanoparticle comprising a silica having a surface and defining a plurality of pores that are suitable to receive drug molecules therein; disposing a trapping agent in pores comprising said plurality of pores where said trapping agent is selected for its ability to trap said drugs within said pores; coating the pores of the nanoparticle with a lipid bilayer; and contacting or bathing said nanoparticle coated with a lipid bilayer with a drug that can pass through said bilayer where said drug enters said
15 pores, reacts with said trapping agent and is retained within the bilayer.

[0119] Embodiment 97: The method of embodiment 96, wherein said lipid bilayer comprises a phospholipid, cholesterol (CHOL), and an mPEG phospholipid.

20 [0120] Embodiment 98: The method according to any one of embodiments 96-97, wherein said phospholipid comprises a saturated fatty acid with a C14-C20 carbon chain, and/or an unsaturated fatty acid with a C14-C20 carbon chain, and/or a natural lipid comprising a mixture of fatty acids with C12-C20 carbon chains.

25 [0121] Embodiment 99: The method of embodiment 98, wherein said phospholipid comprises a saturated fatty acid selected from the group consisting of phosphatidylcholine (DPPC), dimyristoylphosphatidylcholine (DMPC), distearoylphosphatidylcholine (DSPC), and diacylphosphatidylcholine (DAPC).

[0122] Embodiment 100: The method of embodiment 98, wherein said phospholipid comprises a natural lipid selected from the group consisting of egg phosphatidylcholine (egg PC), and soy phosphatidylcholine (soy PC).

30 [0123] Embodiment 101: The method of embodiment 98, wherein said phospholipid comprises an unsaturated fatty acid selected from the group consisting of 1,2-dimyristoleoyl-sn-glycero-3-phosphocholine, 1,2-dipalmitoleoyl-sn-glycero-3-

phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), and 1,2-dieicosenoyl-sn-glycero-3-phosphocholine.

[0124] Embodiment 102: The method according to any one of embodiments 96-101, wherein said lipid bilayer comprises an mPEG phospholipid with a phospholipid C14-18 carbon chain, and a PEG molecular weight ranging from about 350 Da to 5000 Da.

[0125] Embodiment 103: The nanoparticle drug carrier of embodiment 102, wherein said lipid bilayer comprises 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-PEG (DSPE-PEG).

[0126] Embodiment 104: The method of embodiment 97, wherein said lipid bilayer comprises DPPC/Chol/DSPE-PEG or DSPC/Chol/DSPE-PEG.

[0127] Embodiment 105: The method of embodiment 104, wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG.

[0128] Embodiment 106: The method of embodiment 105, wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG2000.

[0129] Embodiment 107: The method according to any one of embodiments 96-106, wherein said lipid bilayer comprises a phospholipid, cholesterol, and mPEG phospholipid at a ratio of 50-90 mol% phospholipid : 10-50 mol% CHOL : 1-10 mol% mPEG phospholipid.

[0130] Embodiment 108: The method of embodiment 105, wherein said lipid bilayer comprises DSPC/Chol/DSPE-PEG in a molar ratio of about 3:2:0.15.

[0131] Embodiment 109: The method according to any one of embodiments 96-108, wherein the lipids comprising said lipid bilayer are combined with said nanoparticles in a ratio sufficient to form a continuous bilayer over the entire nanoparticle.

[0132] Embodiment 110: The method according to any one of embodiments 96-109, wherein the lipids comprising said lipid bilayer are combined with said nanoparticles in particle:lipid ratio ranging from about 1.0:3.0.

[0133] Embodiment 111: The method according to any one of embodiments 96-109, wherein the lipids comprising said lipid bilayer are combined with said nanoparticles in particle:lipid ratio of about 1.0:1.1.

[0134] Embodiment 112: The method according to any one of embodiments 96-111, wherein said lipid bilayer forms a substantially continuous bilayer encompassing the entire nanoparticle.

[0135] Embodiment 113: The method according to any one of embodiments 96-112, wherein said lipid bilayer forms a substantially uniform and intact bilayer encompassing the entire nanoparticle.

[0136] Embodiment 114: The method according to any one of embodiments 96-113, wherein said silica nanoparticle is a mesoporous silica nanoparticle.

[0137] Embodiment 115: The method of embodiment 114, wherein said silica nanoparticle comprises a sol-gel synthesized mesoporous silica nanoparticle.

[0138] Embodiment 116: The method according to any one of embodiments 96-115, wherein said mesoporous silica nanoparticle is size-controlled.

[0139] Embodiment 117: The method according to any one of embodiments 96-116, wherein said mesoporous silica nanoparticle is colloidally stable.

[0140] Embodiment 118: The method according to any one of embodiments 96-117, wherein said mesoporous silica has an average pore size that ranges from about 1 to about 20 nm, or from about 1 to about 10 nm, or from about 2 to about 8 nm.

[0141] Embodiment 119: The method according to any one of embodiments 96-118, wherein said mesoporous silica nanoparticles has an average size ranging from about 50 nm up to about 300 nm, or from about 50 up to about 200 nm, or from about 50 up to about 150 nm, or from about 50 up to about 100 nm, or from about 50 up to about 80 nm, or from about 50 up to about 70 nm, or from about 60 up to about 70 nm.

[0142] Embodiment 120: The method according to any one of embodiments 96-119, wherein cargo-trapping agent before reaction with said drug is selected from the group consisting of triethylammonium sucrose octasulfate (TEA₈SOS), (NH₄)₂SO₄, an ammonium salt, a trimethylammonium salt, and a triethylammonium salt.

[0143] Embodiment 121: The method of embodiment 120, wherein said cargo trapping agent comprises (NH₄)₂SO₄.

[0144] Embodiment 122: The method of embodiment 120, wherein said cargo trapping agent comprises an ammonium salt selected from the group consisting of ammonium sulfate; ammonium sucrose octasulfate, ammonium α -cyclodextrin sulfate,

ammonium β -cyclodextrin sulfate, , ammonium γ -cyclodextrin sulfate, ammonium phosphate; ammonium α -cyclodextrin phosphate, ammonium β -cyclodextrin phosphate, ammonium γ -cyclodextrin phosphate, ammonium citrate, and ammonium acetate.

[0145] Embodiment 123: The method of embodiment 120, wherein said cargo trapping agent comprises a trimethylammonium salt selected from the group consisting of trimethylammonium sulfate, trimethylammonium sucrose octasulfate, trimethylammonium α -cyclodextrin sulfate, trimethylammonium β -cyclodextrin sulfate, trimethylammonium γ -cyclodextrin sulfate, trimethylammonium phosphate, trimethylammonium α -cyclodextrin phosphate, trimethylammonium β -cyclodextrin phosphate, trimethylammonium γ -cyclodextrin phosphate, trimethylammonium citrate, and trimethylammonium acetate.

[0146] Embodiment 124: The method of embodiment 120, wherein said cargo trapping agent comprises a triethylammonium salt selected from the group consisting of triethylammonium sulfate, triethylammonium sucrose octasulfate, triethylammonium α -Cyclodextrin sulfate, triethylammonium β -Cyclodextrin sulfate, triethylammonium γ -Cyclodextrin sulfate, triethylammonium phosphate, triethylammonium α -Cyclodextrin phosphate, triethylammonium β -Cyclodextrin phosphate, triethylammonium γ -Cyclodextrin phosphate, triethylammonium citrate, and triethylammonium acetate.

[0147] Embodiment 125: The method of embodiment 120, wherein cargo-trapping agent before reaction with said drug is triethylammonium sucrose octasulfate (TEA₈SOS).

[0148] Embodiment 126: The method of embodiment 125, wherein said drug is protonated and trapped in said pores as a gel-like precipitate in association of SOS⁸⁻.

[0149] Embodiment 127: The method according to any one of embodiments 96-126, wherein said drug comprises at least one weakly basic group capable of being protonated, and the cargo-trapping agent comprises at least one anionic group.

[0150] Embodiment 128: The method according to any one of embodiments 96-127, wherein said drug is selected to have a pKa greater than 7 and less than 11.

[0151] Embodiment 129: The method according to any one of embodiments 96-128, said drug comprises a primary, secondary, tertiary or quaternary amine.

[0152] Embodiment 130: The method according to any one of embodiments 96-129, wherein said drug is selected to have a water solubility index of about 5 to about 25 mg/mL.

[0153] Embodiment 131: The method according to any one of embodiments 96-130, wherein the cargo is selected to have an octanol/water partition coefficient or logP value of about -3.0 to about 3.0.

[0154] Embodiment 132: The method according to any one of embodiments 96-131, wherein the cargo is selected to be 2-8 nm and less than the average or median size of the pores of the silica nanoparticle.

[0155] Embodiment 133: The method according to any one of embodiments 127-132, wherein said cargo comprises an anti-cancer drug.

[0156] Embodiment 134: The method of embodiment 133, wherein said cargo comprises irinotecan.

[0157] Embodiment 135: The method of embodiment 133, wherein said cargo comprises one or more drugs independently selected from the group consisting of a topoisomerase inhibitor, an antitumor anthracycline antibiotic, a mitotic inhibitor, an alkaloid, an alkaline alkylating agent, a purine or pyrimidine derivative, a protein kinase inhibitor.

[0158] Embodiment 136: The method of embodiment 135, wherein said carrier comprises a topoisomerase inhibitor comprising topotecan.

[0159] Embodiment 137: The method of embodiment 135, wherein said carrier comprises an alkaloid selected from the group consisting of topotecan, 10-hydroxycamptothecin, belotecan, rubitecan, vinorelbine, and LAQ824.

[0160] Embodiment 138: The method of embodiment 135, wherein said carrier comprises a antitumor anthracycline antibiotic selected from the group consisting of doxorubicin, and mitoxantrone.

[0161] Embodiment 139: The method of embodiment 135, wherein said carrier comprises a mitotic inhibitor selected from the group consisting of vinblastine, and vinorelbine.

[0162] Embodiment 140: The method of embodiment 135, wherein said carrier comprises a alkaline alkylating agent selected from the group consisting of cyclophosphamide, mechlorethamine, and temozolomide.

[0163] Embodiment 141: The method of embodiment 135, wherein said carrier comprises a purine or pyrimidine derivative selected from the group consisting of 5-fluorouracil, 5'-deoxy-5-fluorouridine, and gemcitabine.

[0164] Embodiment 142: The method of embodiment 135, wherein said carrier
5 comprises a protein kinase inhibitor selected from the group consisting of imatinib, osimertinib and sunitinib pazopanib, enzastaurin, vandetanib, erlotinib, dasatinib, and nilotinib.

[0165] Embodiment 143: The method according to any one of embodiments 96-142, wherein said drug carrier is conjugated to a moiety selected from the group consisting
10 of a targeting moiety, a fusogenic peptide, and a transport peptide.

[0166] Embodiment 144: The method according to any one of embodiments 96-143, wherein said method produces a nanoparticle drug carrier according to any one of embodiments 1-72.

[0167] Embodiment 145: The method of embodiment 143, wherein said drug
15 carrier is conjugated to a peptide that binds a receptor on a cancer cell.

[0168] Embodiment 146: The method of embodiment 145, wherein said drug carrier is conjugated to an iRGD peptide.

[0169] Embodiment 147: The method of embodiment 145, wherein said drug carrier is conjugated to a targeting peptide shown in Table 2.

[0170] Embodiment 148: The method according to any one of embodiments 143-147, wherein said drug carrier is conjugated to transferrin, and/or ApoE, and/or folate.

[0171] Embodiment 149: The method according to any one of embodiments 143-148, wherein said drug carrier is conjugated to a targeting moiety that comprises an antibody that binds to a cancer marker.

[0172] Embodiment 150: The method of embodiment 149, wherein said drug
25 carrier is conjugated to a targeting moiety that comprises an antibody that binds a cancer marker shown in Table 1.

[0173] Embodiment 151: The method according to any one of embodiments 127-132, wherein said cargo comprises an antibiotic.

[0174] Embodiment 152: The method of embodiment 151, wherein said cargo comprises an antibiotic selected from the group consisting of ciprofloxacin, levofloxacin, and an HIV antiretroviral (*e.g.*, tenofovir disoproxil fumarate, *etc.*).

5 [0175] Embodiment 153: The method according to any one of embodiments 96-152, wherein said drug carrier is loaded to a capacity of at least 30% w/w, or greater than about 40% w/w, or greater than about 50% w/w, or greater than about 60% w/w, or greater than about 70% w/w, or greater than about 80% w/w.

[0176] Embodiment 154: The method according to any one of embodiments 96-152, wherein said drug carrier is loaded to a capacity of at least 80% w/w.

10 [0177] Embodiment 155: The method according to any one of embodiments 96-154, wherein the lipid bilayer comprises a hydrophobic drug.

[0178] Embodiment 156: The method of embodiment 155, wherein the lipid bilayer comprises a hydrophobic drug selected from the group consisting of paclitaxel, ellipticine, camptothecin, SN-38, and a lipid prodrug (*e.g.*, acyclovir diphosphate dimyristoylglycerol, 15 doxorubicin conjugated phospholipid prodrug, phospholipid derivatives of nucleoside analogs, phospholipid linked chlorambucil, and the like).

[0179] Embodiment 157: The method of embodiment 155, wherein the lipid bilayer comprises paclitaxel.

[0180] Embodiment 158: A method of making an irinotecan nanocarrier, 20 comprising: providing a nanocarrier comprising a silica body having a surface including a plurality of pores suitable to receive irinotecan therein, disposing an agent selected for its ability to trap irinotecan within the plurality of pores; coating the pores of the nanocarrier with a phospholipid bilayer (optionally using a sonication process); and introducing irinotecan into the phospholipid bilayer coated pores, so that a phospholipid bilayer coated 25 irinotecan nanocarrier is made.

[0181] Embodiment 159: The method of embodiment 158, wherein said silica body comprises a sol-gel synthesized, size-controlled and colloidally stable silica body.

[0182] Embodiment 160: The method of embodiment 158, wherein the irinotecan trapping agent is triethylammonium sucrose octasulfate (TEA₈SOS).

30 [0183] Embodiment 161: The method of embodiment 160, wherein the nanocarrier: (a) has an irinotecan loading capacity of at least 20% (or 30% or 40%) w/w; and/or (b) show

<5% (or <10%) irinotecan leakage over 24 hours in a biological buffer with pH of 7.4 at 37 °C.

5 **[0184]** Embodiment 162: The method of embodiment 161, wherein the nanocarrier has colloidal stability in physiological fluids with pH 7.4 and remains monodisperse to allow systemic biodistribution and is capable of entering the disease site by vascular leakage (EPR effect) or transcytosis.

[0185] Embodiment 163: The method of embodiment 161, wherein the phospholipid bilayer comprises cholesterol and/or paclitaxel.

10 **[0186]** Embodiment 164: A method of making a nanocarrier, comprising: providing an unloaded nanocarrier including a silica body having a surface and defining a plurality of pores that are suitable to receive molecules therein, and a phospholipid bilayer coating the surface; encapsulating a cargo-trapping agent within the phospholipid bilayer.

[0187] Embodiment 165: The method of embodiment 164, further comprising exposing the nanocarrier to a cargo selected to interact with the cargo-trapping agent.

15 **[0188]** Embodiment 166: The method of embodiment 165, wherein the cargo is selected to have a pKa greater than 7 and less than 11 and capable of being protonated, and the cargo-trapping agent includes at least one anionic group.

[0189] Embodiment 167: The method of embodiment 165, wherein the cargo is irinotecan and the cargo-trapping agent is triethylammonium sucrose octasulfate
20 (TEA₈SOS).

[0190] Embodiment 168: The method of embodiment 165, wherein the cargo is a topoisomerase I inhibitor, topotecan; one or more antitumor anthracycline antibiotics, doxorubicin and mitoxantrone; one or more mitotic inhibitors, vinblastine and vinorelbine; or one or more tyrosine-kinase inhibitors imatinib, osimertinib and sunitinib.

25 **[0191]** Embodiment 169: The method of embodiment 165, wherein the nanocarrier has a drug loading capacity of at least 30% w/w.

[0192] Embodiment 170: A nanocarrier, comprising: a silica body having a surface and defining a plurality of pores that are suitable to receive molecules therein; a phospholipid bilayer coating the surface; and a cargo-trapping agent within the phospholipid
30 bilayer; wherein the submicron structure has a maximum dimension of less than one micron, and wherein the phospholipid bilayer stably seals the plurality of pores.

[0193] Embodiment 171: The nanocarrier of embodiment 170, further comprising a cargo within the phospholipid bilayer.

[0194] Embodiment 172: The nanocarrier of embodiment 171, wherein the cargo is associated with the cargo-trapping agent.

5 **[0195]** Embodiment 173: The nanocarrier of embodiment 172, wherein the cargo includes at least one weakly basic group capable of being protonated, and the cargo-trapping agent includes at least one anionic group.

[0196] Embodiment 174: The nanocarrier of embodiment 171, wherein the cargo is selected to have a pKa greater than 7 and less than 11.

10 **[0197]** Embodiment 175: The nanocarrier of embodiment 171, wherein the cargo comprises a primary, secondary, tertiary or quaternary amine.

[0198] Embodiment 176: The nanocarrier of embodiment 171, wherein the cargo is selected to have a water solubility index of 5-25 mg/mL.

[0199] Embodiment 177: The nanocarrier of embodiment 171, wherein the cargo is
15 selected to have an octanol/water partition coefficient or logP value of -3.0 to 3.0.

[0200] Embodiment 178: The nanocarrier of embodiment 171, wherein the cargo is selected to be 2-8 nm and less than the size of the pores of the nanocarrier.

[0201] Embodiment 179: The nanocarrier of embodiment 171, wherein the cargo is irinotecan and the cargo-trapping agent is triethylammonium sucrose octasulfate
20 (TEA₈SOS).

[0202] Embodiment 180: The nanocarrier of embodiment 171, wherein the cargo is a topoisomerase I inhibitor, topotecan; one or more antitumor anthracycline antibiotics, doxorubicin and mitoxantrone; one or more mitotic inhibitors, vinblastine and vinorelbine; or one or more tyrosine-kinase inhibitors imatinib, osimertinib and sunitinib.

25 **[0203]** Embodiment 181: The nanocarrier of embodiment 171, wherein the nanocarrier has a less than 5% leakage of the cargo over 24 hours in a biological buffer with pH of 7.4 at 37 °C.

[0204] Embodiment 182: The nanocarrier of embodiment 171, wherein the nanocarrier has a drug loading capacity of at least 30% w/w.

[0205] Embodiment 183: The nanocarrier of embodiment 171, wherein the nanocarrier has a drug loading capacity of at least 80% w/w.

[0206] Embodiment 184: The nanocarrier of embodiment 170, wherein the phospholipid bilayer comprises paclitaxel.

5 [0207] Other objects, features and advantages of the present invention will become apparent to those skilled in the art from the following detailed description. It is to be understood, however, that the detailed description and specific examples, while indicating some embodiments of the present invention, are given by way of illustration and not
10 limitation. Many changes and modifications within the scope of the present invention may be made without departing from the spirit thereof, and the invention includes all such modifications.

DEFINITIONS

[0208] The terms "subject," "individual," and "patient" may be used interchangeably and refer to humans, as well as non-human mammals (*e.g.*, non-human primates, canines,
15 equines, felines, porcines, bovines, ungulates, lagomorphs, and the like). In various embodiments, the subject can be a human (*e.g.*, adult male, adult female, adolescent male, adolescent female, male child, female child) under the care of a physician or other health worker in a hospital, as an outpatient, or other clinical context. In certain embodiments, the subject may not be under the care or prescription of a physician or other health worker.

20 [0209] As used herein, the phrase "a subject in need thereof" refers to a subject, as described *infra*, that suffers from, or is at risk for a pathology to which the nanoparticle drug carriers described herein (silicasomes) are directed. Thus, for example, in certain embodiments the subject is a subject with a cancer (*e.g.*, pancreatic ductal adenocarcinoma (PDAC), breast cancer (*e.g.*, drug-resistant breast cancer), colon cancer, brain cancer, and
25 the like). In certain embodiments the subject is a subject with a microbial infection including, but not limited to drug-resistant microbial infections.

[0210] The term "treat" when used with reference to treating, *e.g.*, a pathology or disease refers to the mitigation and/or elimination of one or more symptoms of that pathology or disease, and/or a delay in the progression and/or a reduction in the rate of onset
30 or severity of one or more symptoms of that pathology or disease, and/or the prevention of that pathology or disease. The term treat can refer to prophylactic treatment which includes a delay in the onset or the prevention of the onset of a pathology or disease.

[0211] The terms "coadministration" or "administration in conjunction with" or "cotreatment" when used in reference to the coadministration of a first compound (*e.g.*, a silicasome containing irinotecan) and a second compound (*e.g.*, an iRGD peptide) indicates that the first compound and the second compound are administered so that there is at least
5 some chronological overlap in the biological activity of first compound and the second compound in the organism to which they are administered. Coadministration can include simultaneous administration or sequential administration. In sequential administration there may even be some substantial delay (*e.g.*, minutes or even hours) between administration of the first compound and the second compound as long as their biological activities overlap.

10 In certain embodiments the coadministration is over a time frame that permits the first compound and second compound to produce an enhanced therapeutic or prophylactic effect on the organism. In certain embodiments the enhanced effect is a synergistic effect.

[0212] The terms "nanocarrier" and "nanoparticle drug carrier" and "silicasome" are used interchangeably and refer to a nanostructure having a porous particle core, which is
15 interchangeable with the term "porous nanoparticle" as used herein, and a lipid bilayer encasing (or surrounding or enveloping) the porous particle core. In certain embodiments the silica nanoparticle is a porous silica nanoparticle (*e.g.*, mesoporous silica nanoparticle (MSNP)).

[0213] As used herein, the term "lipid" refers to conventional lipids, phospholipids,
20 cholesterol, chemically functionalized lipids for attachment of PEG and ligands, *etc.*

[0214] As used herein, the terms "lipid bilayer" or "LB" refers to any double layer of oriented amphipathic lipid molecules in which the hydrocarbon tails face inward to form a continuous non-polar phase.

[0215] As used herein, the term "liposome" refers to an aqueous compartment
25 enclosed by a lipid bilayer, as being conventionally defined (*see, e.g.*, Stryer (1981) *Biochemistry*, 2d Edition, W. H. Freeman & Co., p. 213).

[0216] Compared with the lipid bilayer defined in a silicasome, the lipid bilayer in a liposome can be referred to as an "unsupported lipid bilayer" and the liposome itself (when unloaded) can be referred to as an "empty liposome". The lipid bilayer in a silicasome can
30 be referred to as a "supported lipid bilayer" because the lipid bilayer in a silicasome is located on the surface and supported by a porous particle core. In certain embodiments, the lipid bilayer can have a thickness ranging from about 6 nm to about 7 nm which includes a

3-4 nm thickness of the hydrophobic core, plus the hydrated hydrophilic head group layers (each about 0.9 nm) plus two partially hydrated regions of about 0.3 nm each.

[0217] As used herein, the term “selective targeting” or “specific binding” refers to use of targeting ligands on the surface of silicasomes (empty or loaded), in particular, on the surface of the lipid bilayer of the silicasomes, wherein the ligands interact specifically/selectively with receptors or other biomolecular components expressed on the target, *e.g.*, a cell surface of interest. The targeting ligands can include such molecules and/or materials as peptides, antibodies, aptamers, targeting peptides, polysaccharides, and the like.

[0218] A silicasome having targeting ligands can be referred to as a “targeted silicasome”.

[0219] The term "silicasome" refers to a drug containing (drug delivery) silica nanoparticle in which the silica nanoparticle is fully covered with a lipid bilayer (*e.g.*, a phospholipid bilayer). In certain embodiments the silica nanoparticle is a porous silica nanoparticle (*e.g.*, mesoporous silica nanoparticle).

[0220] The term "about" or "approximately" as used herein refers to being within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.* the limitations of the measurement system, *i.e.* the degree of precision required for a particular purpose, such as a pharmaceutical formulation. For example, "about" can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, preferably up to 10%, more preferably up to 5% and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated, the term "about" meaning within an acceptable error range for the particular value should be assumed.

[0221] The term "drug" as used herein refers to a chemical entity of varying molecular size, small and large, naturally occurring or synthetic, that exhibits a therapeutic effect in animals and humans. A drug may include, but is not limited to, an organic molecule (*e.g.*, a small organic molecule), a therapeutic protein, peptide, antigen, or other

biomolecule, an oligonucleotide, an siRNA, a construct encoding CRISPR cas9 components and, optionally one or more guide RNAs, and the like.

[0222] A "pharmaceutically acceptable carrier" as used herein is defined as any of the standard pharmaceutically acceptable carriers. The pharmaceutical compositions of the subject invention can be formulated according to known methods for preparing pharmaceutically useful compositions. The pharmaceutically acceptable carrier can include diluents, adjuvants, and vehicles, as well as carriers, and inert, non-toxic solid or liquid fillers, diluents, or encapsulating material that does not react with the active ingredients of the invention. Examples include, but are not limited to, phosphate buffered saline, physiological saline, water, and emulsions, such as oil/water emulsions. The carrier can be a solvent or dispersing medium containing, for example, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. Formulations are described in a number of sources that are well known and readily available to those skilled in the art. For example, Remington's Pharmaceutical Sciences (Martin E W [1995] Easton Pa., Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the silicasomes described herein.

[0223] As used herein, an "antibody" refers to a protein consisting of one or more polypeptides substantially encoded by immunoglobulin genes or fragments of immunoglobulin genes or derived therefrom that is capable of binding (*e.g.*, specifically binding) to a target (*e.g.*, to a target polypeptide). The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

[0224] A typical immunoglobulin (antibody) structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (V_L) and variable heavy chain (V_H) refer to these light and heavy chains respectively.

[0225] Antibodies exist as intact immunoglobulins or as a number of well characterized fragments produced by digestion with various peptidases. Thus, for example,

pepsin digests an antibody below the disulfide linkages in the hinge region to produce $F(ab)'_2$, a dimer of Fab which itself is a light chain joined to V_H - C_H1 by a disulfide bond. The $F(ab)'_2$ may be reduced under mild conditions to break the disulfide linkage in the hinge region thereby converting the $(Fab')_2$ dimer into a Fab' monomer. The Fab' monomer is essentially a Fab with part of the hinge region (*see, Fundamental Immunology*, W.E. Paul, ed., Raven Press, N.Y. (1993), for a more detailed description of other antibody fragments). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such Fab' fragments may be synthesized *de novo* either chemically or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein also includes antibody fragments either produced by the modification of whole antibodies or synthesized *de novo* using recombinant DNA methodologies. Certain preferred antibodies include single chain antibodies (antibodies that exist as a single polypeptide chain), more preferably single chain Fv antibodies (sFv or scFv) in which a variable heavy and a variable light chain are joined together (directly or through a peptide linker) to form a continuous polypeptide. The single chain Fv antibody is a covalently linked V_H - V_L heterodimer which may be expressed from a nucleic acid including V_H - and V_L - encoding sequences either joined directly or joined by a peptide-encoding linker. Huston, *et al.* (1988) *Proc. Nat. Acad. Sci. USA*, 85: 5879-5883. While the V_H and V_L are connected to each as a single polypeptide chain, the V_H and V_L domains associate non-covalently. The first functional antibody molecules to be expressed on the surface of filamentous phage were single-chain Fv's (scFv), however, alternative expression strategies have also been successful. For example Fab molecules can be displayed on a phage if one of the chains (heavy or light) is fused to g3 capsid protein and the complementary chain exported to the periplasm as a soluble molecule. The two chains can be encoded on the same or on different replicons; the important point is that the two antibody chains in each Fab molecule assemble post-translationally and the dimer is incorporated into the phage particle via linkage of one of the chains to, *e.g.*, g3p (*see, e.g.*, U.S. Patent No: 5733743). The scFv antibodies and a number of other structures converting the naturally aggregated, but chemically separated light and heavy polypeptide chains from an antibody V region into a molecule that folds into a three dimensional structure substantially similar to the structure of an antigen-binding site are known to those of skill in the art (*see e.g.*, U.S. Patent Nos. 5,091,513, 5,132,405, and 4,956,778). In certain embodiments antibodies should include all that have been displayed on phage (*e.g.*, scFv, Fv, Fab and disulfide linked Fv (*see, e.g.*,

Reiter *et al.* (1995) *Protein Eng.* 8: 1323-1331) as well as affibodies, unibodies, and the like.

[0226] The term "specifically binds", as used herein, when referring to a biomolecule (*e.g.*, protein, nucleic acid, antibody, *etc.*), refers to a binding reaction that is determinative of the presence of a biomolecule in heterogeneous population of molecules (*e.g.*, proteins and other biologics). Thus, under designated conditions (*e.g.* immunoassay conditions in the case of an antibody or stringent hybridization conditions in the case of a nucleic acid), the specified ligand or antibody binds to its particular "target" molecule and does not bind in a significant amount to other molecules present in the sample.

BRIEF DESCRIPTION OF THE DRAWINGS

[0227] Referring now to the drawings in which like reference numbers represent corresponding parts throughout:

[0228] Figures 1A-1E illustrate the preparation of LB-MSNP and liposomal irinotecan carriers that use a protonating agent for drug loading, in accordance with one or more embodiments described herein. Fig. 1A shows schematics depicting the synthesis method and loading of irinotecan by LB-MSNPs and liposomes (FIG. 1A, panel A1). After soaking TEA₈SOS into the MSNP particles, the pores are sealed by a LB, derived from sonication of a lipid biofilm (Lu *et al.* (2007) *Small*, 3: 1341-1346). Fig. 1A, panel A2: TEA₈SOS-soaked particles are incubated in an irinotecan solution, allowing the amphipathic drug to diffuse across the lipid bilayer for protonation by TEA₈SOS (TEA₈SOS \leftrightarrow 8TEA + 20 8H⁺ + SOS⁸⁻). The lipid-soluble TEA exits the particle, while the H⁺ converts irinotecan to a hydrophilic derivative that cannot cross the LB. The protonated drug interacts with SOS⁸⁻ to form a gel-like precipitate, that is retained in the pores. Fig. 1A, panel A3: The same technique was used to produce a liposomal equivalent for irinotecan entrapment (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). Fig. 1B: Assessment of the drug loading capacity (DLC) of the Ir-MSNP and Ir-liposome carriers. DLO [the total amount of irinotecan (m₀) - non-encapsulated irinotecan (m₁)]/[the total amount of particle (m_{MSNP} or m_{lipid}) x 100%. The inclusion of TEA₈SOS had a negligible effect on hydrodynamic size and zeta potential of the particles. Hydrodynamic size and zeta potential data are shown in Table 3 (Example 2). Fig. 1C: CryoEM images of the empty, non-coated MSNP, Ir- MSNP and Ir-liposome carriers. The technique is sensitive enough to visualize irinotecan precipitation in the liposome. Fig. 1D: Carrier stability was assessed

by incubation in 100% serum at 37 °C for 24 hrs, and determining drug leakage by HPLC. Fig. 1E) Carrier stability, as determined by the change in hydrodynamic diameter and % drug leakage, following lyophilization and water resuspension.

[0229] Figures 2A-2D illustrate the biodistribution of the Ir-LB-MSNP and Ir-liposomes, in accordance with one or more embodiments described herein. Fig. 2A: Interval IVIS imaging over 48 hrs to compare the biodistribution of IV injected NIR-labeled carriers to the KPC-derived orthotopic tumor site (n=3). NIR fluorescence images in representative animals after IV injection of 100 mg/kg NIR-labeled LB-MSNPs or liposomes are shown. Fig. 2B: *Ex vivo* imaging of explanted organs in the same experiment; animals were sacrificed after 24 hrs. Confocal microscopy confirmed a higher abundance of NIR-labeled LB-MSNP compared to liposomes at the tumor site. Fig. 2C: Irinotecan tumor content was determined in an orthotopic KPC-derived xenograft model (n=3). Animals received IV injection of an irinotecan dose equivalent of 60 mg/kg for the different drug formulations. Following animal sacrifice after 24 hrs, tumor tissues were collected for the measurement of irinotecan content by HPLC. Irinotecan content was expressed as % total injected dose *per* gram of tumor tissue (%ID/g). Data represent mean \pm SD, **p*<0.05. Figure 2D: HPLC measurement of plasma irinotecan concentration in the same experiment. Data represent mean \pm SD, **p*<0.05.

[0230] Figures 3A-3E show the differential tumor inhibitory effect of the free drug and encapsulated irinotecan carriers in the KPC-derived orthotopic tumor model, in accordance with one or more embodiments described herein. Fig. 3A: Assessment of the MTD in an acute dose finding study, using a NCI protocol. Fig. 3B: Growth inhibition of KPC-derived orthotopic tumors in B6/129 mice, following IV administration of 40 mg/kg free drug or encapsulated irinotecan every 4 days for up to 8 administrations. Interval IVIS imaging was used for monitoring tumor growth, which was quantitatively expressed according to the image intensity at the operator-defined ROI. Fig. 3C: Quantitative analysis of apoptosis (using IHC staining for cleaved caspase-3) at the primary tumor site of the animals after treatment (sacrificed on day 40-47). Fig. 3D: Representative autopsy results and *ex vivo* imaging of bioluminescence intensity in the moribund animals (sacrificed on day 40-47), to show treatment impact on surrounding metastases. Visible metastatic spread could be seen in the stomach, intestines, liver, spleen, kidneys, diaphragm, and abdominal wall. There was no infiltration of the heart or lung. Fig. 3E: Heat map display to

summarize the comparative analysis of tumor spread determined by quantitative *ex vivo* imaging in (FIG. 3D). Data represent mean \pm SEM, * p <0.05.

[0231] Figures 4A-4E illustrate a comparative analysis of toxicity reduction by Ir-LB-MSNP vs. Ir-liposomes, in accordance with one or more embodiments described herein.

Fig. 4A: Liver histology obtained from representative moribund animals (sacrificed on day 40-47), using tissue from the experiment shown in Fig. 3B. The arrows in the H&E stained sections point to necrotic liver tissue, while sites marked with an asterisk denote steatosis. Bar is 200 μ m. Fig. 4B: Dual IHC staining of cleaved caspase-3 (apoptosis marker, red) and F4/80 (KC marker, green) in the livers of animal receiving different irinotecan formulations at a dose equivalent of 60 mg/kg, followed by sacrifice at 24 hrs. The nucleus was stained with Hoechst 33342 (blue). Bar = 100 μ m. Fig. 4C: IHC staining for cleaved caspase-3, with H&E counter staining to reveal the spread of apoptosis and blunting of the intestinal villi in the same treated animal groups studied in Fig. 3B. The bar represents 100 μ m. Fig. 4D: Separate experiment, in which a 40 mg/kg dose-equivalent of irinotecan, injected IV every 2nd day, 3 times, were used to study the impact on sternal bone marrow. The sternums were collected on day 7 for embedding, decalcification and H&E staining. The bar represents 200 μ m. Fig. 4E: Schematic to explain the differential hepatotoxicity of Ir-LB-MSNP and Ir-liposome formulations in the liver. Without being bound to a specific theory, it is contemplated that the injected nanocarriers are initially taken up by KCs, where carrier disintegration leads to the irinotecan release to bystander hepatocytes. The subsequent rate of carrier disintegration and drug release to the hepatocytes could determine whether the extent to which the irinotecan is metabolized and rendered inactive. It is further hypothesized that the higher instability of the liposomal carrier leads to more rapid drug release than the more stable Ir-LB-MSNP, which explains the differences in apoptosis and necrosis.

[0232] Figure 5 illustrates a scheme depicting the synthesis procedure of a lipid bilayer coated mesoporous silica nanoparticle (LB-MSNP) and mechanism of TEA₈SOS mediated irinotecan (IRIN) loading, in accordance with one or more embodiments of the invention. MSNP was synthesized using sol/gel chemistry. TEA₈SOS (synthesized in-house) was first incubated with MSNP. This was followed by depositing a LB coat using our biofilm technique (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). TEA₈SOS soaked MSNPs were added to the dry lipid biofilm (*e.g.* DSPC/cholesterol/DSPE-PEG in a molar ratio of 3 : 2 : 0.15) followed by pore sealing upon sonication. After removal of the free

trapping agent, TEA₈SOS loaded LB-MSNP was incubated with irinotecan. This allows free, lipophilic irinotecan diffusing into the particle to be protonated and trapped in the pores. This reaction proceeds as follows: Due to the presence of the trapping agent inside the LB-MSNP, the equilibrium of “TEA₈SOS \leftrightarrow 8TEA-H⁺ + SOS⁸⁻” leads to efflux of the lipid permeable triethylamine (TEA) into the particle incubation medium. The H⁺ release inside the particles creates a pH gradient that leads to the protonation of lipophilic irinotecan, which diffuses through the LB. The protonated IRIN, which is lipid impermeable, interacts with SOS⁸⁻ to form a precipitate, thus leading an efficient IRIN entrapment in LB-MSNP.

[0233] Figures 6A and 6B show morphology and drug release profiles of irinotecan loaded LB-MSNP containing the trapping reagent (Ir-LB-MSNP(+SOS)) and irinotecan loaded liposome using trapping reagent (Ir-Lipo(+SOS)), in accordance with one or more embodiments described herein. Fig. 6A: CryoEM images (TF20, FET) of Ir-LB-MSNP(+SOS) (upper panel) and Ir-Lipo(+SOS) (bottom). The coated lipid bilayer on the MSNP surface can be clearly seen. The upper zoom-in image shows a MSNP with intact lipid coating on the particle surface, while the porous interior shows the presence of the high density complex between TEO₈SOS and irinotecan. Similarly, in the liposome images (lower panel), the irinotecan-trapping agent complex can be seen as high density precipitates on the inside (red arrows marked). Fig. 6B: Drug release profiles of Ir-LB-MSNP(+SOS) and Ir-Lipo(+SOS) in PBS (pH=7.4) and phagolysosomal simulant fluid (PSF, pH 4.5). For drug release measurement, the NPs were prepared in PBS or PSF (1 mL, irinotecan 0.1 mg/mL), followed by shaking at 37°C. At the indicated time points, the released irinotecan was separated from NPs by a centrifugal filter unit with size cut off of 30 kD. The irinotecan concentration in the filtrate was determined by a microplate reader at OD of 360 nm. The experiments were repeated at least twice.

[0234] Figure 7 illustrates cellular uptake and intracellular distribution of Ir-LB-MSNP(+SOS) and Ir-Lipo(+SOS) as determined by confocal microscope, in accordance with one or more embodiments described herein. LB-MSNPs and liposomes were fluorescently labeled by 0.1% w/w Texas red-DHPE in the lipid bilayer. Both particles were incubated with PANC-1 cells for 24 hours and then washed three times in PBS. After cell fixing and washing with PBS, cell membranes were stained with WGA 488 and nuclei with Hoechst dye. The slides were visualized using a confocal microscope.

[0235] Figure 8, panels A-F, illustrates comparative cytotoxicity analysis of Ir-LB-MSNP(+SOS) *versus* Ir-Lipo(+SOS) or free drug, in accordance with one or more embodiments of the invention. Fig. 8, panels A-C: The cell viability of PANC-1 cells was determined by a MTS assay, following treatment with free IRIN, Ir-LB-MSNP(+SOS) and Ir-Lipo(+SOS) at the concentrations shown. Experiments were also carried out over different time intervals (24 hours, 48 hours and 72 hours). Fig. 8, panels D-F: The cell viability of BxPC-3 cells after treatment with free IRIN, Ir-LB-MSNP(+SOS) and Ir-LB-MSNP(+SOS) at different drug concentrations and time points as described for PANC-1 cells.

[0236] Figure 9 shows a histological analysis of the bone marrow, in accordance with one or more embodiments described herein. Bone marrow histology after treatment with free IRIN, Ir-Lipo(+SOS) and Ir-LB-MSNP(+SOS). BALB/c male mice was IV injected with IRIN, Ir-Lipo(+SOS) or Ir-LB-MSNP(+SOS) at a drug dose of 60 mg/kg. After 24 hours, the mice were sacrificed and the sternum was collected and fixed in 10% formalin. The sections were stained with hematoxylin-eosin (H&E), and examined by light microscopy. The data indicate that both LB-MSNP and liposomal formulations lowered the bone marrow depletion induced by irinotecan. Severe bone marrow damage and apoptosis can be seen in free drug group.

[0237] Figure 10 shows a histological analysis of the kidney, in accordance with one or more embodiments described herein. Histological analysis of the kidney after treatment with free IRIN, Ir-Lipo(+SOS) and Ir-LB-MSNP(+SOS). BALB/c male mice was IV injected with IRIN, Ir-Lipo(+SOS) or Ir-LB-MSNP(+SOS) at a drug dose of 60 mg/kg. After 24 hours, the mice were sacrificed and the kidneys were collected and fixed in 10% formalin, followed by paraffin embedding. Tissue sections of 4 μ m thickness were mounted on glass slides. The sections were stained with hematoxylin-eosin (H&E), and examined by light microscopy. Representative histological images showing swelling and edema of Bowman's space (arrows marked) were observed in free IRIN or Ir-Lipo(+SOS) treated animals. These lesions are indicative of acute glomerular inflammation. No significant renal abnormalities were found in the Ir-LB-MSNP(+SOS) group.

[0238] Figure 11 shows low-resolution cryoEM images of representative irinotecan-loaded LB-MSNP, in accordance with one or more embodiments of the invention. Bars represent 100 nm. Visual examination of ~500 particles confirmed the structural integrity of the LB, with successful coating of >99% particles.

[0239] Figure 12, panels A-F, shows H&E staining to show the infiltration of the KPC-derived tumor into surrounding organs, 5 weeks after implantation in B6/129 mice, in accordance with one or more embodiments of the invention. Representative images were captured to show infiltration of the pancreas (panel A), liver (panel B), spleen (panel C), kidney (panel D), intestine (panel E), and stomach (panel F). The bar size is 200 μ m.

[0240] Figure 13 shows an *ex vivo* IVIS image, captured 48 hrs post-IV injection of the NIR-labeled particles in the KPC-derived model in Fig. 2A, in accordance with one or more embodiments described herein. This demonstrates that for an equivalent particle dose, there is an increased abundance of LB-MSNPs at the tumor site compared to labeled liposomes.

[0241] Figure 14 shows the biodistribution of LB-MSNPs in the KPC-derived orthotopic tumor model in B6/129 mice (n= 3), following IV injection of LB-MSNPs (100 mg/kg) and animal sacrifice at 24 hours, in accordance with one or more embodiments of the invention. Tumor tissues and major organs were collected for the measurement of the Si content using inductively coupled plasma optical emission spectroscopy. The particle biodistribution is expressed as % of total injected Si dose (%ID) for each location. Data represent mean \pm SD.

[0242] Figure 15A shows irinotecan tumor content determined in a subcutaneous PANC-1 xenograft model in nude mice (n=3). Animals received IV injection of an irinotecan dose equivalent of 60 mg/kg for the different drug formulations. Following animal sacrifice after 24 hours, tumor tissues were collected for the measurement of irinotecan content by HPLC. Irinotecan content was expressed as % total injected dose *per* gram of tumor tissue (%ID/g). Data represent mean \pm SD, **p*<0.05 for MSNP compared to other groups. Figure 15B shows NIR fluorescence images in a representative animal 24 hours after IV injection of 100 mg/kg NIR-labeled LB-MSNP.

[0243] Figure 16 shows HPLC quantification of irinotecan content in surrounding organs in B6/129 mice. Animals received IV injection of an irinotecan dose equivalent of 60 mg/kg for the different drug formulations (n=3). Following animal sacrifice after 24 hrs, tissues were collected for the measurement of irinotecan content by HPLC. Irinotecan content was expressed as % total injected dose *per* gram of tissue (%ID/g). Data represent mean \pm SD, **p*<0.05.

[0244] Figure 17 shows representative images to show IHC staining for cleaved caspase-3 (apoptosis marker) at the primary tumor site, obtained from the primary tumor site in each animal group (sacrificed on day 40-47) in the experiment described in Fig. 3B. Apoptotic cells (brown color) are indicated by red arrows. Bar = 50 μ m.

5 [0245] Figure 18 shows death rate of the animals in each treatment group of the efficacy study described in Fig. 3B, in accordance with one or more embodiments of the invention. The arrows on the x-axis denote the injection days. The mice were monitored frequently and sacrificed based on “moribund criteria” (Zucker *et al.* (2009) *J. Control. Release*, 139(1): 73-80) instead of spontaneous death. Irinotecan loaded LB-MSNPs
10 resulted in significant ($p < 0.01$, compared by the log-rank test using SPSS 19.0, IBM SPSS Statistics, USA) survival improvement compared to saline and free irinotecan. A better survival outcome was observed for Ir-LB-MSNPs compared to Ir-liposome.

[0246] Figure 19 shows full blood count analysis was performed on samples obtained from the experiment described in Fig. 4D, in accordance with one or more
15 embodiments of the invention. The differential white cell count showed a significant decrease of the absolute neutrophil count performed for the free drug or Ir-liposome treated groups. In contrast, treatment with Ir-LB-MSNPs showed a small but non-significant drop in the neutrophil count. Data represent mean \pm SEM, $*p < 0.05$ compared to control, $^{\#}p < 0.05$ compared to LB-MSNPs group.

20 [0247] Figure 20 shows HPLC quantification of irinotecan content in liver for the different drug carriers, in accordance with one or more embodiments described herein. Animals received IV injection of an irinotecan dose equivalent of 60 mg/kg for the different drug formulations ($n=3$). Following animal sacrifice after 0.5, 4, 24 hours, respectively, liver tissues were collected for the measurement of irinotecan content by HPLC. Irinotecan
25 content was expressed as % total injected dose *per* gram of liver tissue (%ID/g). Data represent mean \pm SD, $*p < 0.05$.

[0248] Figures 21A and 21B illustrate the synthesis and characterization of silicasomes for drug loading and visualization. Fig. 21A: The top panel provides a schematic that shows the synthesis steps for constructing silicasomes and remote drug
30 loading (*see* Example 1 and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). Briefly, MSNP cores were synthesized by sol-gel chemistry and soaked in a solution containing the protonating agent, TEA₈SOS. These particles were coated with a LB, using a sonication

procedure in the presence of a lipid biofilm (*Id.*). This was followed by remote irinotecan loading across the proton gradient provided by TEA₈SOS. Box 1: Schematic of the different silicasome components. Box 2: iRGD peptide conjugation to the LB, using a thiol-maleimide reaction to link the cysteine-modified iRGD peptide to DSPE-PEG₂₀₀₀-maleimide. Box 3: Cryo-EM images to show the bare particles and the silicasomes, with and without the embedding of ~10 nm Au cores (for TEM visualization). The synthesis procedures are described in Example 3 Supplemental Materials and Methods. Bars = 50 nm. Fig. 21B: Autopsy and IVIS images of the KPC-derived orthotopic PDAC model in immunocompetent B6/129 mice. The orthotopic implantation involves minor surgery to inject of 2×10^6 KPC-luc cells in the tail of the pancreas (left panel). The autopsy and bioluminescence imaging reveal primary tumor growth after 1-2 weeks, followed by tumor metastases by 3-5 weeks. Macrometastases are marked by arrows.

[0249] Figure 22A and 22B show that co-administrated iRGD enhanced the tumor biodistribution of IV-injected silicasomes in the KPC-derived orthotopic model. Fig. 22A: The tumor-bearing mice received (i) IV 50 mg/kg of the NIR-labeled silicasome IV with co-administration of 8 μ mol/kg free iRGD (n=3, referred to as “silicasome + iRGD”), (ii) 50 mg/kg of the NIR-labeled, iRGD-conjugated silicasome (n=3, referred to as “silicasome-iRGD”), or (iii) 50 mg/kg NIR labeled silicasome without iRGD. Animals were sacrificed 24 h post injection, followed by *ex vivo* NIR imaging, using IVIS. Fig. 22B: NIR fluorescence intensity and Si content were used to quantify the nanoparticle content in the orthotopic tumors. Data represent mean \pm SD, **p*<0.05.

[0250] Figures 23A-23E show that iRGD co-administration enhances the uptake and efficacy of the Ir-silicasome in the KPC-derived orthotopic model. Fig. 23A: Schedule of the efficacy study in the luciferase expressing KPC-derived orthotopic tumor model (n=6). The chosen Ir-silicasome dose (40 mg/kg irinotecan; 80 mg/kg MSNP) is based on a previous efficacy study (Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). This dose of the Ir-silicasome was IV injected, with or without co-administration of 8 μ mol/kg iRGD. The injections were repeated every 3 days, for a total of 4 administrations. The controls involved animal groups receiving identical doses of free iRGD or the Ir-silicasome alone. Fig. 23B: Representative *ex vivo* imaging of the bioluminescence intensity in the mice prior to sacrifice to show the primary tumor burden and metastases. The images show that iRGD co-administration could enhance the silicasome efficacy. Fig. 23C: Heat map summarizing the impact on tumor and tumor metastasis inhibition in Fig. 23B. Fig. 23D: iRGD co-

administration improved the survival impact of the Ir-silicasome, as shown by the Kaplan-Meier analysis. The effect of the silicasome alone is highly significant compared to PBS and free iRGD ($p=0.001$). iRGD co-administration further enhances survival ($p=0.027$).

Fig. 23E: HPLC analysis of the irinotecan content in the tumor, 24 h after a one-time dose of the Ir-silicasome (40 mg/kg drug), was injected, with or without the co-administration of 8 $\mu\text{mol/kg}$ iRGD. Data represent mean \pm SD ($n=3$), $*p<0.05$.

[0251] Figures 24A and 24C show that iRGD-mediated silicasome uptake requires NRP-1 expression on the tumor vasculature. Fig. 24A: Schematic to demonstrate the mechanism by which iRGD initiates nanoparticle transcytosis. Fig. 24B: Multi-color IHC staining of NRP-1 (green) and CD31 (red), plus nuclear staining (blue) in a KPC-derived tumor section. The IHC staining methodology is described in the methods section. NRP-1 is expressed on both the tumor tissue as well the blood vessels. The merged image shows a high degree of co-localization (94.2%) of NRP-1 with CD31; the co-localization ratio (CR) was determined by Image J software. Bar = 100 μm . Fig. 24C: Interference of an anti-NRP-1 antibody on iRGD-mediated silicasome biodistribution. Fifty μg of the blocking antibody or a control IgG was IV injected 15 mins before IV injection of 50 mg/kg of the NIR-silicasome + 8 $\mu\text{mol/kg}$ free iRGD ($n=3$). *Ex vivo* NIR images showing nanoparticle biodistribution after 24 h. The NIR intensity as well as Si content was used to quantify the nanoparticle uptake at the orthotopic tumor site. Data represent mean \pm SD, $*p<0.05$.

[0252] Figures 25A-25C provide ultrastructural viewing of the silicasome transport system initiated by iRGD co-administration. Fig. 25A: Mice bearing orthotopic tumors were injected with 50 mg/kg Au-silicasomes, with or without co-administration of 8 $\mu\text{mol/kg}$ iRGD. Tumors were harvested at 24 h and immediately fixed for TEM analysis. At least 10 regions of interest in each group were viewed to quantitatively express the abundance of grouped, interconnected vesicles (yellow arrows) in the blood vessel endothelial cells. We calculated the number of vesicles per 1 μm^2 of the intracellular surface area (left panel). Data represent mean \pm SD, $*p<0.05$. Representative TEM pictures with high and low magnification are shown. L = lumen; R = red blood cell. Fig. 25B: TEM visualization of silicasome transcytosis in tumor-bearing mice that received 50 mg/kg Au-silicasome and then sacrificed 24 h later. The electron micrograph shows silicasomes in: (i) the lumen of a tumor blood vessel (red arrow), (ii) transport in the endothelial vesicles (pink arrow), and a deposition in the tumor interstitium (blue arrow). High magnification images of regions "1"- "3" are provided in the panels on the right. E = endothelial cell; L = lumen; P

= pericyte; R = red blood cell. Fig. 25C: TEM image showing the presence of silicasomes in a perinuclear distribution inside a cancer cell. N = nucleus; M = mitochondrion.

[0253] Figures 26A-26B illustrate iRGD-induced silicasome biodistribution in patient-derived xenografts in NSG mice. Fig. 26A: A pair of tumors (XWR#8 and XWR#187), with matched stromal abundance but differing levels of NRP-1 expression, was selected for a biodistribution study in the absence and presence of iRGD co-administration. Masson's trichrome staining shows equivalent levels of collagen expression in both tumors. Multi-color IHC staining (green fluorescent antibody for NRP-1, red fluorescent antibody for CD31) was used to determine the relative abundance of NRP-1 expression, and the extent of overlap with endothelial cells, using image J software. Data represent mean \pm SD, * p <0.05. Fig. 26B: The tumor-bearing animals received IV injection 50 mg/kg NIR-labeled silicasome with or without the co-administration of 8 μ mol/kg iRGD. Animals were sacrificed after 24 h (n=3). *Ex vivo* assessment of the uptake of silicasomes as determined by NIR fluorescence intensity and Si content. Data represent mean \pm SD, * p <0.05.

[0254] Figure 27, panels A-C, illustrate the use of a thiol-maleimide reaction for successful covalent conjugation of iRGD to the lipid bilayer on silicasomes. To validate the success of covalent conjugation, a reactive fluorescein (FAM) labeled iRGD peptide, provided by Dr. Ruoslahti (Sugahara *et al.* (2009) *Cancer Cell*, 16: 510-520), was used for binding to DSPE-PEG-maleimide. The detailed method is described in Example 3 methods section. Panel A: Fluorescence spectra of pristine and FAM-iRGD-silicasomes, suspended at 100 μ g/mL, were obtained in a microplate reader (Molecular Device M5e) at excitation wavelength of 488 nm., the significant retentions of the fluorescence signal on the washed and purified FAM-iRGD-silicasomes confirms a successful conjugation reaction (*Cancer Cell* 2009, 16:510). Panels B and C: The effect of iRGD conjugation to the silicasome was tested for its impact on cellular uptake. A batch of the silicasome-iRGD was synthesized with NIR-labeling (DyLight 680) of the MSNP framework, as described in the methods section. KPC cells were treated at a particle concentration of 100 μ g/mL at 37 °C for 2 h and uptake was determined by flow cytometry (panel B), and confocal microscopy (panel C). An iRGD-free NIR-silicasome with similar labeling efficiency was used as control. Data represent mean \pm SD, * p <0.05. Confocal microscopy confirmed the results (cell membranes were green stained with wheat germ agglutinin and nuclei blue stained with Hoechst 33342). Bars = 20 μ m.

[0255] Figure 28, panels A-B, illustrates assessment of the effect of a non-functional peptide (lacking the CendR motif) on IV silicasome biodistribution. Panel A: We used a non-CendR peptide, cyclo (RGDfK), to repeat the biodistribution experiment in orthotopic model shown in Fig. 22. Briefly, tumor-bearing animals received IV injection of 50 mg/kg NIR-labeled silicasome co-administrated with PBS or 8 μ mol/kg free cyclo (RGDfK), followed by animal sacrifice at 24 h (n=3). Representative *ex vivo* organ NIR fluorescence images were obtained to show nanoparticle biodistribution. Panel B: NIR fluorescence intensity analysis (by IVIS software) and determination of Si content (by ICP-OES) showed that the non-CendR peptide is incapable of improving silicasome uptake at the tumor site.

Data represent mean \pm SD.

[0256] Figure 29, panels A-B, illustrates assessment of silicasome transport in the tumor by confocal microscopy. Panel A: Tumor sections were obtained from experiment in Fig. 22. Representative confocal images were obtained to assess the intratumoral abundance of the NIR labeled silicasome abundance under a SP2-1P-FCS, Leica microscope, using a 633 nm laser. IHC staining of the same section to review the presence of blood vessels (CD31 in green) and the presence of nuclei (DAPI staining in blue), allowed us to determine the presence and migration of nanoparticles in the tumor tissue, using the methodology described by Sugahara *et al.* (2010) *Science*, 328: 1031-1035). Bar = 20 μ m. Panel B: Calculation of the penetration distance silicasomes from the closest tumor blood vessel was estimated for ~15 blood vessels, using Image J software. Box-and-whisker plots (Origin software) were developed to show median (horizontal line), 25th-75th percentiles (box), mean (open square) and SD (whiskers). *, $p < 0.05$ compared with silicasome alone or silicasome-iRGD.

[0257] Figure 30 shows Si contents in the liver (left) and spleen (right), using organs obtained from the animals used in Fig. 22, were determined by ICP-OES analysis. Data represent mean \pm SD, * $p < 0.05$ compared with silicasome or “silicasome + iRGD” groups.

[0258] Figure 31 shows representative TEM images of a KPC-derived tumor from an animal that was prior (24 h) injected with 50 mg/kg Au-embedded silicasomes without iRGD co-administration is shown. The image demonstrates the absence of transcytosis vesicles, with particle deposition on the adluminal side. E = endothelial cell, L = lumen.

[0259] Figure 32 illustrates the results of a comparison between protocell and silicasome technologies.

DETAILED DESCRIPTION

[0260] In order to address the high rate of toxicity while maintaining or increasing therapeutic efficacy, drug delivering nanocarriers are provided to enhance drug delivery (e.g., chemotherapy drugs, antimicrobial drugs, *etc.*) at a target site (e.g. a tumor site, a site of infection, *etc.*) while limiting the amount of free drug that can cause systemic toxicity. It is desirable that the carrier itself maintains stable drug loading in order to prevent toxicity and improve efficacy. The use of nanocarriers to deliver chemotherapeutic agents in animal tumor models established from various cancer cells has demonstrated an ability of such nanocarriers to prolong the drug circulatory half-life, deliver high drug concentrations to the tumor site with improved cytotoxic killing, as well as reduce systemic toxicity compared to free drug equivalents (*see, e.g.,* Messerer *et al.*(2004) *Clin. Cancer Res.* 10(19): 6638-6649; Drummond *et al.* (2006) *Cancer Res.* 66(6): 3271-3277; Ramsay *et al.* (2008) *Clin. Cancer Res.* 14(4): 1208-1217).

[0261] In view of the failure of many liposomal carriers to improve the safety of highly toxic drugs such as irinotecan, novel nanoparticle drug carriers are provided herein that use a lipid bilayer (LB) applied to silica nanoparticles (e.g., mesoporous silica nanoparticles (MSNPs)), thereby providing a supported lipid bilayer. These lipid bilayer coated porous silica nanoparticles (also known as silicasomes) provide a drug carrier that is significantly less leaky than liposomes (Meng *et al.* (2013) *ACS Nano*, 7(2): 994-1005; Meng *et al.* (2011) *ACS Nano*, 5(5): 4131-4144; Meng *et al.* (2013) *ACS Nano*, 7(11): 10048-10065; Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Due to a large internal surface area that can be used for drug packaging, tunable pore sizes, carrier stability, and controlled drug release abilities, MSNPs have been demonstrated to constitute a versatile and multifunctional nanocarrier platform for cancer therapy (Meng *et al.* (2013) *ACS Nano*, 7(2): 994-1005; Meng *et al.* (2011) *ACS Nano*, 5(5): 4131-4144; Meng *et al.* (2013) *ACS Nano*, 7(11): 10048-10065; Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557; Meng *et al.* (2010) *J. Am. Chem. Soc.* 132(36): 12690-12697; Li *et al.* (2012) *Chem. Soc. Rev.* 41(7): 2590-2605; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801). However methods of effectively loading lipid-bilayer coated silica particles have been limited, typically providing loading ranging from about 10 wt% up to 40 wt%.

[0262] In various embodiments new loading methods are provided herein that achieve significantly greater loading of lipid-bilayer coated nanoparticle drug carriers. In

general the new loading method utilizes a cargo trapping reagent, to retain the cargo (*e.g.*, drug or drugs of interest) within the pores of the nanoparticle inside the lipid bilayer. More specifically, in certain embodiments, the method can involve:

- 5 providing a nanocarrier comprising a porous silica body comprising a plurality of pores capable of receiving the cargo (*e.g.*, drug) of interest);
- disposing a trapping agent (a cargo trapping agent) within the plurality of pores where the agent is selected for its ability to trap the cargo within the pores;
- coating the surface pores of the nanocarrier with a lipid bilayer;
- and introducing the cargo into the pores coated by the lipid bilayer, where the cargo reacts
- 10 with the trapping agent and is retained within the pores.

[0263] In certain embodiments the trapping agent comprises a protonating agent and the methods involve providing a drug or drugs that can pass through the lipid bilayer into the bilayer-coated porous nanoparticle. The protonating agent in the porous nanoparticle converts the drug into a hydrophilic derivative that is incapable of back diffusion across the

15 lipid bilayer.

[0264] In the illustrative, but non-limiting embodiment shown in Example 1, MSNP cores were synthesized by sol-gel chemistry and soaked in a solution containing the protonating agent, TEA₈SOS. These particles were coated with a LB, using a sonication procedure in the presence of a lipid biofilm of optimal composition for bilayer stability.

20 This was followed by remote irinotecan loading by the proton gradient provided by TEA₈SOS. Irinotecan is a weak basic and amphipathic molecule that can diffuse across the LB into the MSNP interior packaging space, where proton release by prior entrapped triethylammonium sucrose octasulfate (TEA₈SOS) converts the drug into a hydrophilic derivative, incapable of back-diffusion across the LB. Using this method high loading

25 capacities (*e.g.*, greater than 40 wt%, or greater than 45 wt%, or greater than 50 wt%, or greater than 60 wt%, or greater than 70 wt%, %, or greater than 80 wt%, *etc.*) are achieved.

[0265] The lipid bilayer nanoparticle drug carriers (LB-MSNPs or "silicasomes") produced using the methods described herein offer numerous advantages. Combining an existing biofilm coating method to synthesize LB-MSNPs with a protonating agent, such as

30 TEA₈SOS, provides active and high dose loading of drugs such as irinotecan in a particulate carrier that shows numerous advantages over liposomes. Such advantages include, but are not limited to ease of synthesis, improved drug loading and release profile, improved stability and improved biodistribution. In certain embodiments illustrative, but non-limiting

embodiments, this method of remote loading leads to a 4-fold or greater increase in loading capacity of irinotecan compared to passive drug encapsulation behind the LB in MSNP.

Using the methods described herein, non-encapsulated irinotecan gains entrance to MSNP pores by diffusing through the LB, whereupon protonation renders the irinotecan

5 hydrophilic and incapable of escape.

[0266] Furthermore, the increased loading capacity of the nanocarrier leads to increased drug delivery at the cancer site, with less of the free drug becoming available at sites such as the bone marrow and gastrointestinal tract (GIT), where irinotecan causes toxicity. In one instance, the LB-MSNP has an improved loading capacity (83.5 weight%
10 irinotecan loading) versus an in-house liposomal equivalent of MM-398 (42.5 weight% loading capacity). After optimization of loading time, trapping agent concentration, and amount of irinotecan offered for loading, embodiments of the LB-MSNP formulation have demonstrated 3-5 times higher release capacity at an acidic pH (4.5) compared to the liposomal formulation. The accompanying dose adjustment allows administration of less
15 drug dose to obtain the same efficacy, leading to further reduction in systemic drug toxicity. In one instance, in the calculation of maximum tolerated dose (MTD), embodiments of the LB-MSNP showed a 5-fold improvement in the tolerated dose compared to the free drug. The increased MTD was similar to the liposomal formulation. Analysis of bone marrow histology showed that the irinotecan toxicity was greatly reduced by the LB-MSNP.

20 [0267] Additionally, the synthesis/loading methods described herein are easier to perform than either making conventional liposomes or the polymer-lipid technology used by Zhang *et al.* (2104) *Biomaterials*, 35(11): 3560-3665). This is advantageous for scale-up to GMP manufacturing, including cost savings by reducing the quantity of non-encapsulated drug.

25 [0268] While the methods and nanoparticle drug carriers (silicasomes) produced thereby described herein with respect to irinotecan, it will be understood that, as explained below, the method can be applied to a number of other cargos (*e.g.*, cargos comprising one or more drugs). Moreover, in view of the teachings provided herein, it is believed the improvements in drug delivery (on target), and/or reduced toxicity, and/or improved
30 biodistribution, and/or improved release profile will be achieved using the methods and compositions described herein with numerous other drugs.

[0269] According, in certain embodiments, a LB-MSNP-based nanocarrier delivery system is described, that allows for stable and protected loading of cargo at high loading

levels, with the assistance of a cargo-trapping agent. In some embodiments, the cargo can be a drug. Drug delivery by the modified LB-MSNP encapsulation and trapping allows for more frequent use of drug encapsulation, *e.g.*, chemotherapy drugs, because of enhanced efficacy, high drug loading capacity and reduced systemic toxicity. From a nanocarrier design perspective, various embodiments of this system, for example a polyanionic cargo-trapping agent within LB-MSNP, are employed as an effective design principle for delivering a range of additional weakly basic molecules and drugs, for the treatment of several different types of cancer and other disease processes.

[0270] In one instance, with respect to irinotecan, delivery by the modified LB-MSNP encapsulation and trapping procedure allows for more frequent use of irinotecan and FOLFIRINOX in human PDAC patients because of enhanced efficacy, high drug loading capacity and reduced systemic toxicity of irinotecan. This also allows more PDAC patients to be treated with a more potent therapeutic regimen than gemcitabine, with promise of increased survival. Ultimately, an IV injectable, efficient, biocompatible, and translationally competitive irinotecan formulation (versus MM398 liposome) for PDAC treatment is provided. Particularly in view of the methods and resulting nanoparticle drug carriers described herein, irinotecan and FOLFIRINOX therapy may also be available for the treatment of other cancers including, but not limited to colon, rectal, lung, and ovarian cancer.

[0271] The silicasome delivery system disclosed herein is a multifunctional platform. Mesoporous silica drug carriers have been demonstrated to be able to deliver a wide range of cargoes to cancer cells as well as a variety of human cancer models in animals. These include gemcitabine and paclitaxil co-delivery to xenograft and orthotopic PDAC tumors in mice. Moreover, MSNPs are biodegradable and proven to be safe in extensive animal testing (Meng *et al.* (2013) *ACS Nano*, 7(2): 994-1005; Meng *et al.* (2011) *ACS Nano*, 5(5): 4131-4144; Meng *et al.* (2013) *ACS Nano*, 7(11): 10048-10065; Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801; Zhang *et al.* (2012) *J. Am. Chem. Soc.* 134(38): 15790-15804; Lu *et al.* (2010) *Small*, 6(16): 1794-1805).

[0272] In view of the improved loading, the resulting pharmacokinetic profiles described herein, and the like, it is believed the loading methods, lipid bilayer composition and stability of the resulting nanoparticle drug carriers described herein provide significant improvements and advantages over previous nanoparticle drug carriers.

Methods of loading lipid-bilayer coated porous nanoparticles.

[0273] In various embodiments, improved methods of loading lipid bilayer-coated porous nanoparticles (*e.g.*, mesoporous silica particles) are provided along with the drug delivery nanoparticles produced by such methods. In certain embodiments the methods described herein achieve extremely high levels of drug loading *e.g.*, greater than 40 wt%, or greater than 45 wt%, or greater than 50 wt%, or greater than 60 wt%, or greater than 70 wt%, or greater than 80 wt%, *etc.*).

[0274] As noted above, in certain embodiments, the methods can involve:
providing a nanocarrier comprising a porous silica body comprising a plurality of pores capable of receiving the cargo (*e.g.*, drug) of interest);
disposing a trapping agent (a cargo trapping agent) within the plurality of pores where the agent is selected for its ability to trap the cargo within the pores;
coating the surface pores of the nanocarrier with a lipid bilayer; and
introducing the cargo into the pores coated by the lipid bilayer, where the cargo reacts with the trapping agent and is retained within the pores.

[0275] While it has never been used in a MSNP platform previously, the polyanionic compound, TEA₈SOS, was used in MM-398 to entrap irinotecan (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). Prior to the studies described herein, however, it was unknown whether or not TEA₈SOS could be used to effectively perform remote loading of porous nanoparticles at significantly greater levels than other loading or soak-in methods.

[0276] As described in the Examples, TEA₈SOS is a proton-generating agent that releases eight H⁺ ions and octavalent SOS⁸⁻ upon hydrolysis (Figure 1A, panel A2). Ion-exchange chromatography was used to generate TEA₈SOS, which was soaked into MSNP, as described below in the Methods section (*Id.*). The soaked particles were introduced to a round-bottom flask coated with a lipid biofilm, composed of DSPC/cholesterol/DSPE-PEG2000 at a molar ratio of 3:2:0.15. Sonication of the suspension yielded LB-coated particles, that contain the entrapment agent (Figure 1A, panel A1). These particles were immediately incubated in an irinotecan solution, allowing the drug to be imported, protonated, and entrapped in the pores as a gel-like precipitate in association with SOS⁸⁻ (Figure 1A, panels A1 and A2, Fig. 1B). This allowed us to achieve an irinotecan loading capacity up to 80 wt % or greater (Figure 1B), which approximates the theoretical maximum loading capacity (~100 wt %) of the porous carrier with a combined surface area

of 850 m²/g and a pore volume of ~0.7 cm³/g, as previously demonstrated (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557).

[0277] While the methods are illustrated with respect to the use of drug irinotecan and TEA8SOS as the trapping agent, it will be recognized that, having demonstrated proof of principle, the same method can be used to incorporate numerous other drugs (especially weakly basic drugs) in a silica some and numerous other trapping agents can be similarly be utilized, *e.g.*, as described herein.

[0278] Since drug loading capacity is an important manufacturing consideration that is also critical for therapeutic effectiveness and limitation of drug toxicity, much research has been devoted to optimizing the efficacy of drug encapsulation by MSNP, leading to increased loading capacity and the ability to co-deliver synergistic drug combinations. While it is possible to introduce drugs such as irinotecan into MSNPs by a traditional soaking method, this method of loading has proved to be relatively inefficient achieving loading levels of about 10 wt% (*see, e.g.*, He *et al.* (2010) *Biomaterials*, 31(12): 3335-3346), and it is difficult to retain the drug in the pores (*see, e.g.*, Meng *et al.* (2010) *J. Am. Chem. Soc.* 132(36): 12690-12697). Different from traditional drug loading approaches that depend on physical adsorption, electrostatic attachment, supramolecular assembly or use of entrapment methods (*e.g.*, stoppers, nanovalves or blocking entities), the use of an intact lipid bilayer coating disposed on the porous nanoparticle provides rapid and instantaneous pore sealing for drug entrapment. This simplifies synthesis and provides stable and high drug loading capacity. The methods described herein are effective to load porous particle with cargos at high levels and are compatible with this rapid and instantaneous pore sealing with the lipid bilayer.

[0279] A biofilm technique has been developed for the LB-coated MSNP platform that can be used to rapidly encapsulate gemcitabine (GEM) (*e.g.* a water-soluble nucleoside) by a supported LB (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Not only has this allowed the LB-MSNPs to achieve a loading capacity of up to 40 wt% GEM, but also enabled co-delivery of hydrophobic paclitaxel (PTX) that could be incorporated in the lipid bilayer (LB) (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). This has provided for a synergistic and ratiometric-designed carrier for PDAC treatment in an orthotopic human PDAC model in mice (*Id.*).

[0280] This supported LB coating method has also been demonstrated to be far superior to a comparable liposomal coating technique for MSNPs which involves several

steps, including liposomal adhesion to the MSNP surface, rupture, partial covering of the MSNP surface by an incomplete coat that is secondarily sealed by the addition of additional liposome compositions (Liu *et al.* (2009) *J. Am. Chem. Soc.* 131: 7567-7569).

[0281] However, while the presence of the lipid bilayer (LB) provided an

5 improvement on previous drug delivery nanoparticles, prior to the utilization of the remote loading methods described herein, the bilayer still provided an impediment to achieving even higher drug loading levels.

[0282] As noted above, in various embodiments drug loading by the LB-MSNP platform has been further improved by using the LB to encapsulate a protonating agent as

10 the first step, followed by irinotecan loading, which is dependent on the drug's weak basic properties ($pK_a = 8.1$). As shown in the examples herein, comparisons were made between the MSNP carrier and a liposome equivalent, in which a non-supported LB was used for irinotecan loading subsequent to encapsulation of triethylammonium sucrose octasulfate (TEA₈SOS) (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277; Von Hoff *et al.*
15 (2013) *Br. J. Cancer*, 109(4): 920-925). Not only did the MSNP carrier achieve higher loading capacity for irinotecan and tumor killing than the liposomal formulation in a robust orthotopic PDAC model, but it also prevented drug toxicity due to increased carrier stability and reduced leakage compared to liposomes. Thus, the LB-MSNP platform exhibits the properties desirable for a first-line irinotecan carrier for PDAC (or other cancer) treatment.

20 [0283] Recently, a polymer-lipid supported coating method has been described by Zhang *et al.* for irinotecan drug loading of MSNPs to treat drug-resistant breast cancer tumors in Balb/c nude mice (Zhang *et al.* (2014) *Biomaterials*, 35(11): 3650-3665).

However, the methodology described in this reference is different, and does not achieve the same drug loading capacity compared to the methods and compositions described herein.

25 First, Zhang *et al.* do not use a classical LB or a trapping agent, resulting in a carrier with only 1/5th the loading capacity (~15 %, w/w) of the carrier provided herein (~83.5%, w/w).

[0284] A second major difference is in the drug loading procedure. The authors state that "CPT-11 @MSNs" were centrifuged and washed in PBS (pH 7.4), then dried under vacuum at room temperature. In contrast, the methods described herein do not

30 require drying of the unprotected MSNPs. Drying of the MSNPs can be problematic because of the difficulty obtaining dispersion upon re-suspension in aqueous media which is a key requirement for systemic drug delivery. In addition, the present methods do not

require washing of the drug (*e.g.* irinotecan)-loaded MSNP before pore sealing which minimizes drug loss, and facilitates a high loading capacity (*e.g.*, ~83.5% (w/w)).

[0285] A third major difference is the composition of the LB formulation. In one or more embodiments, a mixture of commercially available lipids plus cholesterol (*e.g.*, DSPC/cholesterol/DSPE-PEG) is used, while Zhang *et al.* use an in-house synthesized, pH-sensitive, Pluronic P123 grafted DOPE. The use of the Pluronic is apparently based on its ability to act as an inhibitor of drug efflux (*see, e.g.*, Batrakova *et al.* (2004) *Pharm. Res.* 21(12): 2226-2233), hence the consideration to treat drug-resistant breast cancer. The absence of cholesterol in the coated bilayer decreases the fluidity and stability of the platform. In contrast, the presence of cholesterol in the silicasomes contributes to the excellent stability of the LB-MSNP disclosed herein (see Examples). Moreover, irinotecan encapsulation in the provided carrier shows <5% leakage over 24 hours in a biological buffer with pH of 7.4 at 37 °C. This is 2.5x lower than the premature leakage (~16% leakage at pH 7.4 and 37°C over 24 hours) described by Zhang *et al.* (2014) *Biomaterials*, 35: 3650-3665).

[0286] Another difference from the drug deliver platform described by Zhang *et al.* is that various embodiments of the present invention use probe sonication for biofilm rehydration and pore sealing, followed by centrifugation purification or size exclusion chromatography. On the other hand, Zhang *et al.* utilizes a membrane extrusion method, as for liposomes. Also, CTAC is used as a templating agent for MSNP synthesis, whereas Zhang *et al.* utilize CTAB. Finally, the present invention addresses the development of therapy for PDAC and other cancers and is not principally designed to overcome drug resistance, as are the drug delivery particles described by Zhang *et al.* for use in the breast cancer study. All considered, without being bound to a particular theory, it is believed the lipid bilayer coated nanoparticle drug carriers described herein (*e.g.*, the irinotecan-delivering LB-MSNP) provide a unique design that outperforms the carriers of Zhang *et al.* based on drug loading capacity, colloidal stability, ease of production, and stable drug retention in the blood and body fluids.

Nanoparticles.

[0287] In various embodiments nanoparticle drug carriers described herein comprises a porous silica nanoparticle (*e.g.*, a silica body having a surface and defining a plurality of pores that are suitable to receive molecules therein) coated with a lipid bilayer.

For example, in certain embodiments the silica nanoparticle can be a mesoporous silica nanoparticle. The fact that the nanoparticle is referred to as a silica nanoparticle does not preclude materials other than silica from also being incorporated within the silica nanoparticle. In some embodiments, the silica nanoparticle may be substantially spherical with a plurality of pore openings through the surface providing access to the pores. However, in various embodiments the silica nanoparticle can have shapes other than substantially spherical shapes. Thus, for example, in certain embodiments the silica nanoparticle can be substantially ovoid, rod-shaped, a substantially regular polygon, an irregular polygon, and the like.

[0288] Generally, the silica nanoparticle comprises a silica body that defines an outer surface between the pore openings, as well as side walls within the pores. The pores can extend through the silica body to another pore opening, or a pore can extend only partially through the silica body such that that it has a bottom surface of defined by the silica body.

[0289] In some embodiments, the silica body is mesoporous. In other embodiments, the silica body is microporous. As used herein, "mesoporous" means having pores with a diameter between about 2 nm and about 50 nm, while "microporous" means having pores with a diameter smaller than about 2 nm. In general, the pores may be of any size, but in typical embodiments are large enough to contain one or more therapeutic compounds therein. In such embodiments, the pores allow small molecules, for example, therapeutic compounds such as anticancer compounds to adhere or bind to the inside surface of the pores, and to be released from the silica body when used for therapeutic purposes. In some embodiments, the pores are substantially cylindrical.

[0290] In certain embodiments the nanoparticles comprise pores having pore diameters between about 1 nm and about 10 nm in diameter or between about 2 nm and about 8 nm. In certain embodiments the nanoparticles comprise pores having pore diameters between about 1 nm and about 6 nm, or between about 2 nm and about 5 nm. Other embodiments include particles having pore diameters less than 2.5 nm. In other embodiments, the pore diameters are between 1.5 and 2.5 nm. Silica nanoparticles having other pore sizes may be prepared, for example, by using different surfactants or swelling agents during the preparation of the silica nanoparticles.

[0291] In various embodiments the nanoparticles can include particles as large (*e.g.*, average or median diameter (or other characteristic dimension) as about 1000 nm. However

in various embodiments the nanoparticles are typically less than 500 nm or less than about 300 nm as, in general, particles larger than 300 nm may be less effective in entering living cells or blood vessel fenestrations. In certain embodiments the nanoparticles range in size from about 40 nm, or from about 50 nm, or from about 60 nm up to about 100 nm, or up to about 90 nm, or up to about 80 nm, or up to about 70 nm. In certain embodiments the nanoparticle range in size from about 60 nm to about 70 nm. Some embodiments include nanoparticles having an average maximum dimension between about 50 nm and about 1000 nm. Other embodiments include nanoparticles having an average maximum dimension between about 50 nm and about 500 nm. Other embodiments include nanoparticles having an average maximum dimension between about 50 nm and about 200 nm. In some embodiments, the average maximum dimension is greater than about 20nm, greater than about 30nm, greater than 40nm, or greater than about 50nm. Other embodiments include nanoparticles having an average maximum dimension less than about 500 nm, less than about 300nm, less than about 200nm, less than about 100nm or less than about 75 nm. As used herein, the size of the nanoparticle refers to the average or median size of the primary particles, as measured by transmission electron microscopy (TEM) or similar visualization technique.

[0292] Illustrative mesoporous silica nanoparticles include, but are not limited to MCM-41, MCM-48, and SBA-15 (*see, e.g., Katiyaret al. (2006) J. Chromatog. 1122(1-2): 13-20*).

[0293] Methods of making porous silica nanoparticles are well known to those of skill in the art. In certain embodiments mesoporous silica nanoparticle are synthesized by reacting tetraethyl orthosilicate (TEOS) with a template made of micellar rods. The result is a collection of nano-sized spheres or rods that are filled with a regular arrangement of pores. The template can then be removed by washing with a solvent adjusted to the proper pH (*see, e.g., Trewyn et al. (2007) Chem. Eng. J. 137(1): 23–29*). In certain embodiments mesoporous particles can also be synthesized using a simple sol-gel method (*see, e.g., Nandiyanto, et al. (2009) Microporous and Mesoporous Mat. 120(3): 447-453, and the like*). In certain embodiments tetraethyl orthosilicate can also be used with an additional polymer monomer (as a template). In certain embodiments 3-mercaptopropyl)trimethoxysilane (MPTMS) is used instead of TEOS.

[0294] In certain embodiments the mesoporous silica nanoparticles are cores were synthesized by a modification of the sol/gel procedure described by Meng *et al. (2015) ACS*

Nano, 9(4): 3540-3557. To synthesize a batch of ~500 mg of MSNP, 50 mL of CTAC is mixed with 150 mL of H₂O in a flask (*e.g.*, a 500 mL conical flask), followed by stirring (*e.g.*, at 350 rpm for 15 min at 85°C). This is followed by the addition of 8 mL of 10% triethanolamine for 30 min at the same temperature. Then, 7.5 mL of the silica precursor, TEOS, is added dropwise at a rate of 1 mL/min using a peristaltic pump. The solution is stirred at 350 rpm at 85°C for 20 min, leading to the formation of particles with a primary size of ~65 nm. The surfactant can be removed by washing the particles with a mixture of methanol/HCl (500:19 v/v) at room temperature for 24 h. The particles can be centrifuged at 10 000 rpm for 60 min and washed three times in methanol.

[0295] While the loading methods described herein have been demonstrated with respect to the loading of porous silica nanoparticles (*e.g.*, mesoporous silica), it will be recognized that similar loading methods can be used with other porous nanoparticles. Numerous other mesoporous materials that can be used in drug delivery nanoparticles are known to those of skill in the art. For example, in certain embodiments mesoporous carbon nanoparticles could be utilized. Mesoporous carbon nanoparticles are well known to those of skill in the art (*see, e.g.*, Huang *et al.* (2016) *Carbon*, 101: 135-142; Zhu *et al.* (2014) *Asian J. Pharm. Sci.*, 9(2): 82-91; and the like).

[0296] Similarly, in certain embodiments, mesoporous polymeric particles can be utilized. The syntheses of highly ordered mesoporous polymers and carbon frameworks from organic-organic assembly of triblock copolymers with soluble, low-molecular-weight phenolic resin precursors (resols) by an evaporation induced self-assembly strategy have been reported by Meng *et al.* (2006) *Chem. Mat.* 6(18): 4447-4464 and in the references cited therein.

[0297] The nanoparticles described herein are illustrative and non-limiting. Using the teachings provided herein numerous other lipid bilayer drug delivery nanoparticles will be available to one of skill in the art.

Lipid Bilayer.

[0298] The drug carrier nanoparticles described herein comprise a porous nanoparticle (*e.g.* a mesoporous silica nanoparticle (MSNP)) coated with a lipid bilayer. In certain embodiments the bilayer composition is optimized to provide a rapid and uniform particle coating, to provide colloidal and circulatory stability, and to provide effective cargo retention, while also permitting a desirable cargo release profile.

[0299] In certain embodiments the lipid bilayer comprises a combination of a phospholipid, cholesterol, and in certain embodiments, a pegylated lipid (*e.g.*, DSPE-PEG₂₀₀₀), or a factionalized pegylated lipid (*e.g.*, DSPE-PEG₂₀₀₀-maleimide) to facilitate conjugation with targeting or other moieties.

5 **[0300]** To attach a surface LB coating, a coated lipid film procedure was developed in which drug- or TEOS-soaked MSNP suspensions were added to a large lipid film surface, coated on, *e.g.*, a round-bottom flask. Using different lipid bilayer compositions, a series of experiments can be performed to find a composition and optimal lipid/particle ratio that provides rapid and uniform particle wrapping, coating and effective cargo retention
10 and/or release upon sonication. It is believed that this lipid composition and wrapping cannot be achieved by liposomal fusion to the particle surface under low energy vortexing conditions.

[0301] As described in Example 1, in certain embodiments, 500 mg MSNPs are soaked in a 20 mL TEOS (80 mM solution), which is added on top of the lipid biofilm,
15 comprised of a 550 mg mixture of DSPC/Chol/DSPE-PEG₂₀₀₀ (molar ratio 3:2:0.15), coated at the bottom of a round bottom flask (*see* Example 1, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). The ratio of "3:2:0.15" equals to "58.3 mol% : 38.8 mol% : 3.9 mol%" if one uses mol% to present the ratio. This provides a lipid:particle ratio of ~1.1:1. After sonication to accomplish particle wrapping and coating with a LB, free TEOS is
20 removed by size exclusion chromatography over a Sepharose CL-4B column. The TEOS loaded silicasomes are incubated in a 10 mg/mL irinotecan solution for drug loading in a water bath at 65 °C. The loading is stopped after 30 min by quenching in and ice water bath, following which the drug-loaded silicasomes were washed 3 times by centrifugation and re-suspended in PBS.

25 **[0302]** The lipid bilayer formulation described above and in Example 1 is illustrative and non-limiting. Depending on the drug(s) being loaded into the silicasome and the desired release provide, in various embodiments different lipid bilayer formulations can be used and an optimal formulation can be determined.

[0303] Accordingly, in certain embodiments the lipid bilayer can comprise: 1) one
30 or more saturated fatty acids with C14-C20 carbon chain, such as dimyristoylphosphatidylcholine (DMPC), dipalmitoylphosphatidylcholine (DPPC), distearoylphosphatidylcholine (DSPC), and dioleoylphosphatidylcholine (DOPC); and/or 2) One or more unsaturated fatty acids with a C14-C20 carbon chain, such as 1,2-

dimyristoleoyl-sn-glycero-3-phosphocholine, 1,2-dipalmitoleoyl-sn-glycero-3-phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dieicosenoyl-sn-glycero-3-phosphocholine; and/or 3) Natural lipids comprising a mixture of fatty acids with C12-C20 carbon chain, such as Egg PC, and Soy PC, sphingomyelin, and the like. These lipids are illustrative but non-limiting and numerous other lipids are known and can be incorporated into a lipid bilayer for formation of a silicasome.

[0304] In certain embodiments the silicasome contains a lipid (*e.g.*, a phospholipid), cholesterol, and a PEG functionalized lipid (*e.g.*, a mPEG phospholipid). In certain embodiments the mPEG phospholipids comprises a C14-C18 phospholipid carbon chain from, and a PEG molecular weight from 350-5000 (*e.g.*, MPEG 5000, MPEG 3000, MPEG 2000, MPEG 1000, MPEG 750, MPEG 550, MPEG 350, and the like). In certain embodiments the mPEG phospholipid comprises DSPE-PEG5000, DSPE-PEG3000, DSPE-PEG2000, DSPE-PEG1000, DSPE-PEG750, DSPE-PEG550, or DSPE-PEG350. MPEGs are commercially available (*see, e.g.*, [//avantilipids.com/product-category/products/polymers-polymerizable-lipids/mpeg-phospholipids/](http://avantilipids.com/product-category/products/polymers-polymerizable-lipids/mpeg-phospholipids/)).

[0305] In certain embodiments the ratio of phospholipid: CHOL:PEG, is about phospholipid (50-90 mol%): CHOL (10-50 mol%) : PEG (1-10 mol%).

[0306] This protocol provided above and in the Examples is illustrative. In certain embodiments the trapping agent can be altered, the lipid composition and molar ratios can be altered, and the drug or drugs can be altered to identify other silicasomes optimized for their particular cargo(s).

[0307] It is noted, for example, that an effective lipid formulation for a gemcitabine-containing silicasome comprises DPPC/cholesterol/DSPE-PEG at a molar ratio of 77.5:20:2.5, while an effective lipid formulation for an irinotecan containing silicasome comprises DSPC/Chol/DSPE-PEG₂₀₀₀ (molar ratio 3:2:0.15, which equals 58.3 mol% : 38.8 mol% : 3.9 mol%).

[0308] In certain embodiments these methods can be varied to improve drug-loading capacity (weight of drug/total weight of carrier). In certain embodiments the drug loading capacity is at least about 30%, or at least about 40%, at least about 50%, at least about 60%, at least about 70%, or at least 80% w/w. In certain embodiments drug loading is greater than 40% w/w, or greater than 45% w/w, or greater than 50% w/w, or greater than 60% w/w, or greater than 70% w/w, %, or greater than 80% w/w.

[0309] The protocols described herein provide a nanoparticle drug carrier (nanocarrier) outperforms nanocarriers made by the liposomal method of coating by fusion with the MSNP surface, as illustrated by the descriptions of the ease of synthesis and improved loading capacity and release profile provided herein. In typical embodiments, the LB coating procedure is used to rapidly encapsulate the protonation agent, *e.g.*, TEA₈SOS, which subsequently provides irinotecan loading and entrapment by protonating the incoming drug diffusing across the LB. This leads to high drug loading in the particles pores. The rapid and effective pore sealing to retain the trapping agent (*e.g.*, TEA₈SOS), without leakage, contributes to the effectiveness and stability of the carrier.

Cargo trapping reagent.

[0310] The cargo-trapping reagent can be selected to interact with a desired cargo. In some embodiments, this interaction can be an ionic or protonation reaction, although other modes of interaction are contemplated. The cargo-trapping agent can have one or more ionic sites, *i.e.*, can be mono-ionic or poly-ionic. The ionic moiety can be cationic, anionic, or in some cases, the cargo-trapping agent can include both cationic and anionic moieties. The ionic sites can be in equilibrium with corresponding uncharged forms; for example, an anionic carboxylate (-COO^-) can be in equilibrium with its corresponding carboxylic acid (-COOH); or in another example, an amine (-NH_2) can be in equilibrium with its corresponding protonated ammonium form (-NH_3^+). These equilibria are influenced by the pH of the local environment.

[0311] Likewise, in certain embodiments, the cargo can include one or more ionic sites. The cargo-trapping agent and cargo can be selected to interact inside the nanoparticle (*e.g.*, the mesoporous silica nanoparticle). This interaction can help retain the cargo within the nanoparticle until release of the cargo is desired. In some embodiments, the cargo can exist in a pH-dependent equilibrium between non-ionic and ionic forms. The non-ionic form can diffuse across the lipid bilayer and enter pores of the MSNP. There, the cargo-trapping agent (*e.g.*, a polyionic cargo-trapping agent) can interact with the ionic form of the cargo and thereby retain the cargo within the nanocarrier, *e.g.*, within pores of the MSNP (provided the ionic forms of the cargo and cargo-trapping agent have opposite charges). The interaction can be an ionic interaction, and can include formation of a precipitate. Trapping of cargo within the nanocarrier can provide higher levels of cargo loading compared to similar systems, *e.g.*, nanocarriers that omit the cargo-trapping agent, or liposomes that do include a trapping agent. Release of the cargo can be achieved by an

appropriate change in pH to disrupt the interaction between the cargo and cargo-trapping agent, for example, by returning the cargo to its non-ionic state which can more readily diffuse across the lipid bilayer. In one embodiment, the cargo is irinotecan and the cargo-trapping agent is TEA₈SOS.

5 **[0312]** The cargo trapping agent need not be limited to TEA₈SOS. In certain embodiments the cargo trapping comprises small molecules like (NH₄)₂SO₄, and the like. Other trapping agents include, but are not limited to, ammonium salts (*e.g.*, ammonium sulfate, ammonium sucrose octasulfate, ammonium α -cyclodextrin sulfate, ammonium β -cyclodextrin sulfate, , ammonium γ -cyclodextrin sulfate, ammonium phosphate, ammonium
10 α -cyclodextrin phosphate, ammonium β -cyclodextrin phosphate, ammonium γ -cyclodextrin phosphate, ammonium citrate, ammonium acetate, and the like), trimethylammonium salts (*e.g.*, trimethylammonium sulfate, trimethylammonium sucrose octasulfate, trimethylammonium α -cyclodextrin sulfate, trimethylammonium β -cyclodextrin sulfate, trimethylammonium γ -cyclodextrin sulfate, trimethylammonium phosphate,
15 trimethylammonium α -cyclodextrin phosphate, trimethylammonium β -cyclodextrin phosphate, trimethylammonium γ -cyclodextrin phosphate, trimethylammonium citrate, trimethylammonium acetate, and the like), triethylammonium salts (*e.g.*, triethylammonium sulfate, triethylammonium sucrose octasulfate, triethylammonium α -cyclodextrin sulfate, triethylammonium β -cyclodextrin sulfate, triethylammonium γ -cyclodextrin sulfate,
20 triethylammonium phosphate, triethylammonium α -cyclodextrin phosphate, triethylammonium β -cyclodextrin phosphate, triethylammonium γ -cyclodextrin phosphate, triethylammonium citrate, triethylammonium acetate, and the like).

[0313] It is also worth pointing out that, in addition to TEA₈SOS, transmembrane pH gradients can also be generated by acidic buffers (*e.g.* citrate) (Chou *et al.* (2003) *J. Biosci. Bioengineer.*, 95(4): 405-408; Nichols *et al.* (1976) *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 455(1): 269-271), proton-generating dissociable salts (*e.g.* (NH₄)₂SO₄) (Haran *et al.* (1993) *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1151(2): 201-215; Maurer-Spurej *et al.* (1999) *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1416(1): 1-10; Fritze *et al.* (2006) *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1758(10): 1633-1640), or ionophore-mediated ion gradients from metal
30 salts (*e.g.* A23187 and MnSO₄) (Messerer *et al.* (2004) *Clinical Cancer Res.* 10(19): 6638-6649; Ramsay *et al.* (2008) *Eur. J. Pharmaceut. Biopharmaceut.* 68(3): 607-617; Fenske *et al.* (1998) *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1414(1)L 188-204).

Moreover, it is possible to generate reverse pH gradients for drug loading, such as use a calcium acetate gradient to improve amphiphilic weak acid loading in LB-MSNP, a strategy that has been utilized in liposomes (Avnir *et al.* (2008) *Arthritis & Rheumatism*, 58(1): 119-129).

5 **Cargo/drug.**

[0314] In one or more embodiments, the cargo comprises an organic compound that includes at least one primary amine group, or at least one secondary amine group, or at least one tertiary amine group, or at least one quaternary amine group, or any combination thereof, capable of being protonated. We have also identified a comprehensive list of weak
10 basic drugs that can be loaded into LB-MSNPs through a proton gradient. The general characteristics of these cargo molecules include the following chemical properties:

(i) organic molecular compounds that include primary, secondary, tertiary or quaternary amine(s);

(ii) a $pK_a < 11$ to allow protonation and entrapment behind the LB (Zucker *et al.* (2009) *J. Control. Release*, 139(1): 73-80; Cern *et al.* (2012) *J. Control. Release*, 160(2):
15 147-157; Xu *et al.* (2014) *Pharmaceut. Res.* 31(10): 2583-2592);

(iii) a water solubility index of 5-25 mg/mL and amphipathic characteristics that allow diffusion across the LB;

(iv) an octanol/water partition coefficient or logP value of -3.0 to 3.0 (Zucker
20 *et al.* (2009) *J. Control. Release*, 139(1): 73-80; Cern *et al.* (2012) *J. Control. Release*, 160(2): 147-157);

(v) suitable molecular weight with a geometric size less than MSNP pore size (2-8 nm), to allow entry into the MSNP pores (Li *et al.* (2012) *Chem. Soc. Rev.* 41(7): 2590-2605; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Tarn *et al.* (2013) *Acc. Chem. Res.*
25 46(3): 792-801).

[0315] Without being all-inclusive, in various embodiments a list of potential chemotherapy agents can include irinotecan derivatives and metabolites such as SN38 together with other alkaloids (*e.g.* topotecan, 10-hydroxycamptothecin, belotecan, rubitecan, vinorelbine, LAQ824, vinblastine, vincristine, homoharringtonine, trabectedin),
30 anthracyclines (*e.g.* doxorubicin, epirubicin, pirarubicin, daunorubicin, rubidomycin, valrubicin, amrubicin), alkaline anthracenediones (*e.g.* mitoxantrone), alkaline alkylating agents (*e.g.* cyclophosphamide, mechlorethamine, temozolomide), purine or pyrimidine derivatives (*e.g.* 5-fluorouracil, 5'-deoxy-5-fluorouridine, gemcitabine, capecitabine), and

protein kinase inhibitors (*e.g.* pazopanib, enzastaurin, vandetanib, erlotinib, dasatinib, nilotinib, sunitinib).

[0316] The ability to package and deliver one or a combination of the above agents will enhance the wider utility of the multifunctional LB-MSNP platform, including
5 treatment consideration of additional cancer types such as colon, breast, lung, liver, glioma, melanoma, *etc.*

[0317] It is also possible to co-package drug combinations in the above list into a single carrier. For example, based on the success that we achieved with our GEM/ PTX co-delivery platform (*see, e.g.* Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557), it is possible to
10 consider combining drugs in the FOLFIRINOX regimen (*e.g.*, oxaliplatin with irinotecan) for synergistic and ratiometric delivery using the silicasomes described herein. Moreover, drug loading by our LB-MSNP can be used for non-cancerous applications, such as encapsulating antibiotics for infectious disease applications, *e.g.*, ciprofloxacin, levofloxacin or HIV antiretrovirals (*e.g.*, tenofovir disoproxil fumarate).

[0318] In addition to the above mentioned cancer drugs, as long as the drug
15 molecules are basic as described above, the trapping reagent facilitated LB-MSNP platform is useful in efficient drug loading and delivery. For non-basic drug molecules, while the trapping reagent will provide limited help, the provided one-step biofilm technique for MSNP pore healing is still valid for an even larger spectrum of drug molecules, such as
20 anticancer drugs, anti-viral drugs, antifungal drugs, and antibiotics.

[0319] For example, with the usage of LB biofilm for MSNP sealing, in certain embodiments efficient drug encapsulation includes, but is not limited to, everolimus, trabectedin, paclitaxel, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON
25 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457, MLN8054, PHA-739358, R-763, AT-9263, a FLT-3 inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhibitor, a c-MET inhibitor, a PARP inhibitor, a Cdk inhibitor, an EGFR TK inhibitor, an IGFR-TK inhibitor, an anti-HGF
30 antibody, a PI3 kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody, pemetrexed, erlotinib, dasatinib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab,

ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide,
 gimatecan, IL13-PE38QQR, INO 1001, IPdR1 KRX-0402, lucanthone, LY 317615,
 neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS-
 100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, 5'-deoxy-
 5 5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244,
 capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1H-pyrrolo[2,3-
 d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled
 irinotecan, tamoxifen, toremifene citrate, anastrozole, exemestane, letrozole,
 DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11,
 10 CHIR-258,); 3-[5-(methylsulfonyl)piperadinemethyl]-indolyl]-quinolone, vatalanib, AG-
 013736, AVE-0005, the acetate salt of [D-Ser(Bu t) 6,Azgly 10] (pyro-Glu-His-Trp-Ser-
 Tyr-D-Ser(Bu t)-Leu-Arg-Pro-Azgly-NH₂ acetate [C₅₉H₈₄N₁₈O_{i4}-(C₂H₄O₂)X where
 x=1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone
 acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide,
 15 flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib,
 lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016,
 Ionafernib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide
 hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KR951,
 aminoglutethimide, amsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin
 20 (BCG) vaccine, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil,
 cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin,
 daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone,
 flutamide, gemcitabine, gleevec, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide,
 levamisole, lomustine, mechlorethamine, melphalan, 6-mercaptopurine, mesna,
 25 methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin,
 pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab,
 streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin,
 vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine,
 altretamine, floxuridine, 5-deoxyuridine, cytosine arabinoside, 6-mecaptopurine,
 30 deoxycoformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan,
 razoxin, marimastat, COL-3, neovastat, BMS-275291, squalamine, endostatin, SU5416,
 SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxifene,
 spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox, gefitinib,
 bortezomib, paclitaxel, cremophor-free paclitaxel, docetaxel, epithilone B, BMS-247550,

BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA-923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR-3339, ZK186619, topotecan, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001, ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zoledronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11, dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen mustard, methylprednisolone, ibritumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89, casopitant, netupitant, an NK-1 receptor antagonists, palonosetron, aprepitant, diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, pegfilgrastim, erythropoietin, epoetin alfa, and darbepoetin alfa, boceprevir, daclatasvir, asunapavir, INX-189, FV-100, NM 283, VX-950 (telaprevir), SCH 50304, TMC435, VX-500, BX-813, SCH503034, R1626, ITMN-191 (R7227), R7128, PF-868554, TT033, CGH-759, GI 5005, MK-7009, SIRNA-034, MK-0608, A-837093, GS 9190, GS 9256, GS 9451, GS 5885, GS 6620, GS 9620, GS9669, ACH-1095, ACH-2928, GSK625433, TG4040 (MVA-HCV), A-831, F351, NS5A, NS4B, ANA598, A-689, GNI-104, IDX102, ADX184, ALS-2200, ALS-2158, BI 201335, BI 207127, BIT-225, BIT-8020, GL59728, GL60667, PSI-938, PSI-7977, PSI-7851, SCY-635, ribavirin, pegylated interferon, PHX1766, SP-30, or a mixture thereof.

[0320] In certain embodiments the cargo comprise an antifungal agent. Illustrative antifungal agents include, but are not limited to Amphotericin B (*e.g.*, for Most fungal infections except *Pseudallescheria* sp., and the like), Anidulafungin (*e.g.*, for candidiasis, including candidemia, and the like), Caspofungin (*e.g.*, for aspergillosis, candidiasis, including candidemia, and the like), Fluconazole (*e.g.*, for mucosal and systemic candidiasis, cryptococcal meningitis, coccidioidal meningitis, and the like), Flucytosine (*e.g.*, for Candidiasis (systemic), Cryptococcosis, and the like), Isavuconazole (*e.g.*, for Aspergillosis, Mucormycosis, and the like), Itraconazole (*e.g.*, for Dermatormycosis,

Histoplasmosis, blastomycosis, coccidioidomycosis, sporotrichosis, and the like), Micafungin (*e.g.*, for Candidiasis, including candidemia), Posaconazole (*e.g.*, for prophylaxis for invasive aspergillosis and candidiasis, oral candidiasis, oral candidiasis refractory to itraconazole, and the like), Voriconazole (*e.g.*, for Invasive aspergillosis, Fusariosis, Scedosporiosis, and the like), and so forth.

Dual therapeutic Silicasomes.

[0321] It will be recognized that in certain embodiments, the nanoparticle drug carriers (silicasomes) described herein can comprise two or more therapeutic agents. Thus, for example, in certain embodiments the pores in the silicasome can be loaded with two, or with three, or with four, or more different therapeutic agents. This can, in certain embodiments, permit ratiometric delivery of these therapeutic agents. By way of non-limiting illustration, numerous multi-agent therapeutic regimen are known for the treatment of cancer. These include, but are not limited to COMP (methotrexate, prednisone), LSA₂-L₂ (cyclophosphamide, vincristine, prednisone, daunomycin, methotrexate, cytarabine, thioguanine, asparaginase, and carmustine), FOLFIRINOX (irinotecan, oxaliplatin, 5-fluorouracil, leucovorin), and the like. In certain embodiments two or more agents that meet the requirements described herein for drugs to be loaded into silicasomes using the methods described herein can be provided in the silicasomes. Where the multidrug regimen include agents that are not compatible with the loading methods described herein, some agents (*e.g.*, irinotecan) can be provided in the silicasome to afford improved tolerance and other components of the treatment regimen can be administered by traditional modalities.

[0322] In certain embodiments hydrophobic (*e.g.*, lipophilic) drugs, and other agents) can be provided in the lipid bilayer component of the silicasome. Such hydrophobic drugs include, but are not limited to paclitaxel, ellipticine, camptothecin, L-asparaginase, doxorubicin, SN-38 and the like. In certain embodiments the lipid bilayer component of the silicasome can contain one or more phospholipid prodrugs (*e.g.*, drugs conjugated to a lipid). Illustrative lipid prodrugs include, but are not limited to acyclovir diphosphate dimyristoylglycerol (*see, e.g.*, Hostetler, *et al.* (1993) *Proc. Natl. Acad. Sci. USA*, 90(24): 11835-11839), doxorubicin conjugated phospholipid prodrugs (*see, e.g.*, Wang *et al.* (2015) *J. Mater. Chem. B*, 3: 3297-3305), Phospholipid Derivatives of Nucleoside Analogs (*e.g.*, 5'-diphosphate-L-1,2-dipalmitin derivatives of 1-β-D-arabinofuranosylcytosine (ara-C), 9-β-D-arabinofuranosyladenine (ara-A), tubercidin, and the like (*see, e.g.*, Matsushita *et al.*

(1981) *Cancer Res.*, 41: 2707-2713)), phospholipid linked chlorambucil (*see, e.g.,* Pederson *et al.* (2010) *J. Med. Chem.*, 53: 3782-3792), and the like.

[0323] The foregoing multi-agent silicasomes are illustrative and non-limiting. Using the teachings provided herein numerous combinations of therapeutic agents for incorporation in (or on) the silicasomes described herein will be available to one of skill in the art.

Targeting ligands and Immunoconjugates.

[0324] In certain embodiments the LB-coated nanoparticle (silicasome) can be conjugated to one or more targeting ligands, *e.g.,* to facilitate specific delivery in endothelial cells, to cancer cells, to fusogenic ligands, *e.g.,* to facilitate endosomal escape, ligands to promote transport across the blood-brain barrier, and the like.

[0325] In one illustrative, but non-limiting embodiment, the silicasome is conjugated to a fusogenic peptides such as histidine-rich H5WYG (H₂N-GLFHAI AHFIHGGWHGLIHGWYG-COOH, (SEQ ID NO:1)) (*see, e.g.,* Midoux *et al.*, (1998) *Bioconjug. Chem.* 9: 260-267).

[0326] In certain embodiments the silicasome is conjugated to targeting ligands which include antibodies as well as targeting peptides. Targeting antibodies include, but are not limited to intact immunoglobulins, immunoglobulin fragments (*e.g.,* F(ab)₂, Fab, *etc.*) single chain antibodies, diabodies, affibodies, unibodies, nanobodies, and the like. In certain embodiments antibodies will be used that specifically bind a cancer marker (*e.g.,* a tumor associated antigen). A wide variety of cancer markers are known to those of skill in the art. The markers need not be unique to cancer cells, but can also be effective where the expression of the marker is elevated in a cancer cell (as compared to normal healthy cells) or where the marker is not present at comparable levels in surrounding tissues (especially where the chimeric moiety is delivered locally).

[0327] Illustrative cancer markers include, for example, the tumor marker recognized by the ND4 monoclonal antibody. This marker is found on poorly differentiated colorectal cancer, as well as gastrointestinal neuroendocrine tumors (*see, e.g.,* Tobi *et al.* (1998) *Cancer Detection and Prevention*, 22(2): 147-152). Other important targets for cancer immunotherapy are membrane bound complement regulatory glycoproteins CD46, CD55 and CD59, which have been found to be expressed on most tumor cells *in vivo* and *in vitro*. Human mucins (*e.g.* MUC1) are known tumor markers as are gp100, tyrosinase, and

MAGE, which are found in melanoma. Wild-type Wilms' tumor gene WT1 is expressed at high levels not only in most of acute myelocytic, acute lymphocytic, and chronic myelocytic leukemia, but also in various types of solid tumors including lung cancer.

[0328] Acute lymphocytic leukemia has been characterized by the TAAs HLA-Dr, CD1, CD2, CD5, CD7, CD19, and CD20. Acute myelogenous leukemia has been characterized by the TAAs HLA-Dr, CD7, CD13, CD14, CD15, CD33, and CD34. Breast cancer has been characterized by the markers EGFR, HER2, MUC1, Tag-72. Various carcinomas have been characterized by the markers MUC1, TAG-72, and CEA. Chronic lymphocytic leukemia has been characterized by the markers CD3, CD19, CD20, CD21, CD25, and HLA-DR. Hairy cell leukemia has been characterized by the markers CD19, CD20, CD21, CD25. Hodgkin's disease has been characterized by the Leu-M1 marker. Various melanomas have been characterized by the HMB 45 marker. Non-hodgkins lymphomas have been characterized by the CD20, CD19, and Ia marker. And various prostate cancers have been characterized by the PSMA and SE10 markers.

[0329] In addition, many kinds of tumor cells display unusual antigens that are either inappropriate for the cell type and/or its environment, or are only normally present during the organisms' development (*e.g.*, fetal antigens). Examples of such antigens include the glycosphingolipid GD2, a disialoganglioside that is normally only expressed at a significant level on the outer surface membranes of neuronal cells, where its exposure to the immune system is limited by the blood-brain barrier. GD2 is expressed on the surfaces of a wide range of tumor cells including neuroblastoma, medulloblastomas, astrocytomas, melanomas, small-cell lung cancer, osteosarcomas and other soft tissue sarcomas. GD2 is thus a convenient tumor-specific target for immunotherapies.

[0330] Other kinds of tumor cells display cell surface receptors that are rare or absent on the surfaces of healthy cells, and which are responsible for activating cellular signaling pathways that cause the unregulated growth and division of the tumor cell. Examples include (ErbB2) HER2/*neu*, a constitutively active cell surface receptor that is produced at abnormally high levels on the surface of breast cancer tumor cells.

[0331] Other useful targets include, but are not limited to CD20, CD52, CD33, epidermal growth factor receptor and the like.

[0332] An illustrative, but not limiting list of suitable tumor markers is provided in Table 1. Antibodies to these and other cancer markers are known to those of skill in the art

and can be obtained commercially or readily produced, *e.g.* using phage-display technology. Such antibodies can readily be conjugated to the silicasomes described herein, *e.g.*, in the same manner that iRGD peptide is conjugated in Example 3.

5 [0333] Table 1. Illustrative cancer markers and associated references, all of which are incorporated herein by reference for the purpose of identifying the referenced tumor markers.

Marker	Reference
5 alpha reductase	Délos <i>et al.</i> (1998) <i>Int J Cancer</i> , 75:6 840-846
α -fetoprotein	Esteban <i>et al.</i> (1996) <i>Tumour Biol.</i> , 17(5): 299-305
AM-1	Harada <i>et al.</i> (1996) <i>Tohoku J Exp Med.</i> , 180(3): 273-288
APC	Dihlmann <i>et al.</i> (1997) <i>Oncol Res.</i> , 9(3) 119-127
APRIL	Sordat <i>et al.</i> (1998) <i>J Exp Med.</i> , 188(6): 1185-1190
BAGE	Böel <i>et al.</i> (1995) <i>Immunity</i> , 2: 167-175.
β -catenin	Hugh <i>et al.</i> (1999) <i>Int J Cancer</i> , 82(4): 504-11
Bcl2	Koty <i>et al.</i> (1999) <i>Lung Cancer</i> , 23(2): 115-127
bcr-abl (b3a2)	Verfaillie <i>et al.</i> (1996) <i>Blood</i> , 87(11): 4770-4779
CA-125	Bast <i>et al.</i> (1998) <i>Int J Biol Markers</i> , 13(4): 179-187
CASP-8/FLICE	Mandruzzato <i>et al.</i> (1997) <i>J Exp Med.</i> , 186(5): 785-793.
Cathepsins	Thomssen <i>et al.</i> (1995) <i>Clin Cancer Res.</i> , 1(7): 741-746
CD19	Scheuermann <i>et al.</i> (1995) <i>Leuk Lymphoma</i> , 18(5-6): 385-397
CD20	Knox <i>et al.</i> (1996) <i>Clin Cancer Res.</i> , 2(3): 457-470
CD21, CD23	Shubinsky <i>et al.</i> (1997) <i>Leuk Lymphoma</i> , 25(5-6): 521-530
CD22, CD38	French <i>et al.</i> (1995) <i>Br J Cancer</i> , 71(5): 986-994
CD33	Nakase <i>et al.</i> (1996) <i>Am J Clin Pathol.</i> , 105(6): 761-768
CD35	Yamakawa <i>et al.</i> <i>Cancer</i> , 73(11): 2808-2817
CD44	Naot <i>et al.</i> (1997) <i>Adv Cancer Res.</i> , 71: 241-319
CD45	Buzzi <i>et al.</i> (1992) <i>Cancer Res.</i> , 52(14): 4027-4035
CD46	Yamakawa <i>et al.</i> (1994) <i>Cancer</i> , 73(11): 2808-2817
CD5	Stein <i>et al.</i> (1991) <i>Clin Exp Immunol.</i> , 85(3): 418-423
CD52	Ginaldi <i>et al.</i> (1998) <i>Leuk Res.</i> , 22(2): 185-191
CD55	Spendlove <i>et al.</i> (1999) <i>Cancer Res.</i> , 59: 2282-2286.
CD59 (791Tgp72)	Jarvis <i>et al.</i> (1997) <i>Int J Cancer</i> , 71(6): 1049-1055
CDC27	Wang <i>et al.</i> (1999) <i>Science</i> , 284(5418): 1351-1354
CDK4	Wölfel <i>et al.</i> (1995) <i>Science</i> , 269(5228): 1281-1284
CEA	Kass <i>et al.</i> (1999) <i>Cancer Res.</i> , 59(3): 676-683
c-myc	Watson <i>et al.</i> (1991) <i>Cancer Res.</i> , 51(15): 3996-4000
Cox-2	Tsujii <i>et al.</i> (1998) <i>Cell</i> , 93: 705-716

DCC	Gotley <i>et al.</i> (1996) <i>Oncogene</i> , 13(4): 787-795
DcR3	Pitti <i>et al.</i> (1998) <i>Nature</i> , 396: 699-703
E6/E7	Steller <i>et al.</i> (1996) <i>Cancer Res.</i> , 56(21): 5087-5091
EGFR	Yang <i>et al.</i> (1999) <i>Cancer Res.</i> , 59(6): 1236-1243.
EMBP	Shiina <i>et al.</i> (1996) <i>Prostate</i> , 29(3): 169-176.
Ena78	Arenberg <i>et al.</i> (1998) <i>J. Clin. Invest.</i> , 102: 465-472.
FGF8b and FGF8a	Dorkin <i>et al.</i> (1999) <i>Oncogene</i> , 18(17): 2755-2761
FLK-1/KDR	Annie and Fong (1999) <i>Cancer Res.</i> , 59: 99-106
Folic Acid Receptor	Dixon <i>et al.</i> (1992) <i>J Biol Chem.</i> , 267(33): 24140-72414
G250	Divgi <i>et al.</i> (1998) <i>Clin Cancer Res.</i> , 4(11): 2729-2739
GAGE-Family	De Backer <i>et al.</i> (1999) <i>Cancer Res.</i> , 59(13): 3157-3165
gastrin 17	Watson <i>et al.</i> (1995) <i>Int J Cancer</i> , 61(2): 233-240
Gastrin-releasing hormone (bombesin)	Wang <i>et al.</i> (1996) <i>Int J Cancer</i> , 68(4): 528-534
GD2/GD3/GM2	Wiesner and Sweeley (1995) <i>Int J Cancer</i> , 60(3): 294-299
GnRH	Bahk <i>et al.</i> (1998) <i>Urol Res.</i> , 26(4): 259-264
GnTV	Hengstler <i>et al.</i> (1998) <i>Recent Results Cancer Res.</i> , 154: 47-85
gp100/Pmel17	Wagner <i>et al.</i> (1997) <i>Cancer Immunol Immunother.</i> , 44(4): 239-247
gp-100-in4	Kirkin <i>et al.</i> (1998) <i>APMIS</i> , 106(7): 665-679
gp15	Maeurer <i>et al.</i> (1996) <i>Melanoma Res.</i> , 6(1): 11-24
gp75/TRP-1	Lewis <i>et al.</i> (1995) <i>Semin Cancer Biol.</i> , 6(6): 321-327
hCG	Hoermann <i>et al.</i> (1992) <i>Cancer Res.</i> , 52(6): 1520-1524
Heparanase	Vlodavsky <i>et al.</i> (1999) <i>Nat Med.</i> , 5(7): 793-802
Her2/neu	Lewis <i>et al.</i> (1995) <i>Semin Cancer Biol.</i> , 6(6): 321-327
Her3	
HMTV	Kahl <i>et al.</i> (1991) <i>Br J Cancer</i> , 63(4): 534-540
Hsp70	Jaattela <i>et al.</i> (1998) <i>EMBO J.</i> , 17(21): 6124-6134
hTERT (telomerase)	Vonderheide <i>et al.</i> (1999) <i>Immunity</i> , 10: 673-679. 1999.
IGFR1	Ellis <i>et al.</i> (1998) <i>Breast Cancer Res. Treat.</i> , 52: 175-184
IL-13R	Murata <i>et al.</i> (1997) <i>Biochem Biophys Res Commun.</i> , 238(1): 90-94
iNOS	Klotz <i>et al.</i> (1998) <i>Cancer</i> , 82(10): 1897-1903
Ki 67	Gerdes <i>et al.</i> (1983) <i>Int J Cancer</i> , 31: 13-20
KIAA0205	Guéguen <i>et al.</i> (1998) <i>J Immunol.</i> , 160(12): 6188-6194
K-ras, H-ras, N-ras	Abrams <i>et al.</i> (1996) <i>Semin Oncol.</i> , 23(1): 118-134
KSA (CO17-1A)	Zhang <i>et al.</i> (1998) <i>Clin Cancer Res.</i> , 4(2): 295-302

LDLR-FUT	Caruso <i>et al.</i> (1998) <i>Oncol Rep.</i> , 5(4): 927-930
MAGE Family (MAGE1, MAGE3, <i>etc.</i>)	Marchand <i>et al.</i> (1999) <i>Int J Cancer</i> , 80(2): 219-230
Mammaglobin	Watson <i>et al.</i> (1999) <i>Cancer Res.</i> , 59: 13 3028-3031
MAP17	Kocher <i>et al.</i> (1996) <i>Am J Pathol.</i> , 149(2): 493-500
Melan-A/ MART-1	Lewis and Houghton (1995) <i>Semin Cancer Biol.</i> , 6(6): 321-327
mesothelin	Chang <i>et al.</i> (1996) <i>Proc. Natl. Acad. Sci., USA</i> , 93(1): 136-140
MIC A/B	Groh <i>et al.</i> (1998) <i>Science</i> , 279: 1737-1740
MT-MMP's, such as MMP2, MMP3, MMP7, MMP9	Sato and Seiki (1996) <i>J Biochem (Tokyo)</i> , 119(2): 209-215
Mox1	Candia <i>et al.</i> (1992) <i>Development</i> , 116(4): 1123-1136
Mucin, such as MUC- 1, MUC-2, MUC-3, and MUC-4	Lewis and Houghton (1995) <i>Semin Cancer Biol.</i> , 6(6): 321-327
MUM-1	Kirkin <i>et al.</i> (1998) <i>APMIS</i> , 106(7): 665-679
NY-ESO-1	Jager <i>et al.</i> (1998) <i>J. Exp. Med.</i> , 187: 265-270
Osteonectin	Graham <i>et al.</i> (1997) <i>Eur J Cancer</i> , 33(10): 1654-1660
p15	Yoshida <i>et al.</i> (1995) <i>Cancer Res.</i> , 55(13): 2756-2760
P170/MDR1	Trock <i>et al.</i> (1997) <i>J Natl Cancer Inst.</i> , 89(13): 917-931
p53	Roth <i>et al.</i> (1996) <i>Proc. Natl. Acad. Sci., USA</i> , 93(10): 4781-4786.
p97/melanotransferrin	Furukawa <i>et al.</i> (1989) <i>J Exp Med.</i> , 169(2): 585-590
PAI-1	Grøndahl-Hansen <i>et al.</i> (1993) <i>Cancer Res.</i> , 53(11): 2513-2521
PDGF	Vassbotn <i>et al.</i> (1993) <i>Mol Cell Biol.</i> , 13(7): 4066-4076
Plasminogen (uPA)	Naitoh <i>et al.</i> (1995) <i>Jpn J Cancer Res.</i> , 86(1): 48-56
PRAME	Kirkin <i>et al.</i> (1998) <i>APMIS</i> , 106(7): 665-679
Probasin	Matuo <i>et al.</i> (1985) <i>Biochem Biophys Res Commun.</i> , 130(1): 293-300
Progenipoietin	-----
PSA	Sanda <i>et al.</i> (1999) <i>Urology</i> , 53(2): 260-266.
PSM	Kawakami <i>et al.</i> (1997) <i>Cancer Res.</i> , 57(12): 2321-2324
RAGE-1	Gaugler <i>et al.</i> (1996) <i>Immunogenetics</i> , 44(5): 323-330
Rb	Dosaka-Akita <i>et al.</i> (1997) <i>Cancer</i> , 79(7): 1329-1337
RCAS1	Sonoda <i>et al.</i> (1996) <i>Cancer</i> , 77(8): 1501-1509.
SART-1	Kikuchi <i>et al.</i> (1999) <i>Int J Cancer</i> , 81(3): 459-466
SSX gene Family	Gure <i>et al.</i> (1997) <i>Int J Cancer</i> , 72(6): 965-971
STAT3	Bromberg <i>et al.</i> (1999) <i>Cell</i> , 98(3): 295-303

STn (mucin assoc.)	Sandmaier <i>et al.</i> (1999) <i>J Immunother.</i> , 22(1): 54-66
TAG-72	Kuroki <i>et al.</i> (1990) <i>Cancer Res.</i> , 50(16): 4872-4879
TGF- α	Imanishi <i>et al.</i> (1989) <i>Br J Cancer</i> , 59(5): 761-765
TGF- β	Picon <i>et al.</i> (1998) <i>Cancer Epidemiol Biomarkers Prev</i> , 7(6): 497-504
Thymosin β 15	Bao <i>et al.</i> (1996) <i>Nature Medicine</i> , 2(12), 1322-1328
IFN- α	Moradi <i>et al.</i> (1993) <i>Cancer</i> , 72(8): 2433-2440
TPA	Maulard <i>et al.</i> (1994) <i>Cancer</i> , 73(2): 394-398
TPI	Nishida <i>et al.</i> (1984) <i>Cancer Res</i> 44(8): 3324-9
TRP-2	Parkhurst <i>et al.</i> (1998) <i>Cancer Res.</i> , 58(21) 4895-4901
Tyrosinase	Kirkin <i>et al.</i> (1998) <i>APMIS</i> , 106(7): 665-679
VEGF	Hyodo <i>et al.</i> (1998) <i>Eur J Cancer</i> , 34(13): 2041-2045
ZAG	Sanchez <i>et al.</i> (1999) <i>Science</i> , 283(5409): 1914-1919
p16INK4	Quelle <i>et al.</i> (1995) <i>Oncogene</i> Aug. 17, 1995; 11(4): 635-645
Glutathione S-transferase	Hengstler (1998) <i>et al. Recent Results Cancer Res.</i> , 154: 47-85

[0334] Any of the foregoing markers can be used as targets for the targeting moieties comprising the silicasome constructs described herein. In certain embodiments the target markers include, but are not limited to members of the epidermal growth factor family (*e.g.*, HER2, HER3, EGF, HER4), CD1, CD2, CD3, CD5, CD7, CD13, CD14, CD15, CD19, CD20, CD21, CD23, CD25, CD33, CD34, CD38, 5E10, CEA, HLA-DR, HM 1.24, HMB 45, 1a, Leu-M1, MUC1, PMSA, TAG-72, phosphatidyl serine antigen, and the like.

[0335] The foregoing markers are intended to be illustrative and not limiting. Other tumor associated antigens will be known to those of skill in the art.

[0336] Where the tumor marker is a cell surface receptor, ligand to that receptor can function as targeting moieties. Similarly mimetics of such ligands can also be used as targeting moieties. Thus, in certain embodiments peptide ligands can be used in addition to or in place of various antibodies. An illustrative, but non-limiting list of suitable targeting peptides is shown in Table 2. In certain embodiments any one or more of these peptides can be conjugated to a silicasome described herein.

Table 2. Illustrative, but non-limiting peptides that target membrane receptors expressed or overexpressed by various cancer cells.

Target Membrane Receptor	Targeting Peptide	SEQ ID NO
Integrin receptor $\alpha_v\beta_3$	c(RGDfK)	2
	c(RGDfC)	3
	c(RGDyC)	4
	RGD	
GFR	GE11 (YHWYGYTPQNVI)	5
GFR	GSG-KCCYSL	6
SSTR2	Ostreotide	
GRP	QWAVGHML	7
CCK	DYMGWMDF	8
NT	RRPYIL	9
	RRPYILQLYENKPRRPYIL	10
LHRH	Gondaorelin	
GPRC family members	Antagonist G	
c() indicates cyclopeptide. Lower case indicates "D" amino acid.		

[0337] In certain embodiments the silicasomes can be conjugated to moieties that facilitate stability in circulation and/or that hide the silicasome from the reticuloendothelial system (REC) and/or that facilitate transport across a barrier (*e.g.*, a stromal barrier, the blood brain barrier, *etc.*), and/or into a tissue. In certain embodiments the silicasomes are conjugated to transferrin or ApoE to facilitate transport across the blood brain barrier. In certain embodiments the silicasomes are conjugated to folate.

[0338] Methods of coupling the silicasomes to targeting (or other) agents are well known to those of skill in the art. Examples include, but are not limited to the use of biotin and avidin or streptavidin (*see, e.g.*, U.S. Patent No: US 4,885,172 A), by traditional chemical reactions using, for example, bifunctional coupling agents such as glutaraldehyde, diimide esters, aromatic and aliphatic diisocyanates, bis-p-nitrophenyl esters of dicarboxylic acids, aromatic disulfonyl chlorides and bifunctional arylhalides such as 1,5-difluoro-2,4-dinitrobenzene; p,p'-difluoro m,m'-dinitrodiphenyl sulfone, sulfhydryl-reactive maleimides, and the like. Appropriate reactions which may be applied to such couplings are described in Williams *et al.* Methods in Immunology and Immunochemistry Vol. 1, Academic Press, New York 1967. In one illustrative but non-limiting approach described in Example 3 a peptide (in this example iRGD) is coupled to the silicasome by substituting DSPE-PEG₂₀₀₀ with DSPE-PEG₂₀₀₀-maleimide (see methods section in Example 3), allowing thiol-

maleimide coupling to the cysteine-modified peptide. It will also be recognized that in certain embodiments the targeting (and other) moieties can be conjugated to a lipid comprising the lipid bilayer.

[0339] The former conjugates and coupling methods are illustrative and non-limiting. Using the teachings provided herein, numerous other moieties can be conjugated to the silicasomes described herein by any of a variety of methods.

Pharmaceutical Formulations, Administration and Therapy

Pharmaceutical formulations.

[0340] In some embodiments, the nanoparticle drug carriers described herein are administered alone or in a mixture with a physiologically-acceptable carrier (such as physiological saline or phosphate buffer) selected in accordance with the route of administration and standard pharmaceutical practice. For example, when used as an injectable, the silicasomes can be formulated as a sterile suspension, dispersion, or emulsion with a pharmaceutically acceptable carrier. In certain embodiments normal saline can be employed as the pharmaceutically acceptable carrier. Other suitable carriers include, *e.g.*, water, buffered water, 0.4% saline, 0.3% glycine, and the like, including glycoproteins for enhanced stability, such as albumin, lipoprotein, globulin, *etc.* In compositions comprising saline or other salt-containing carriers, the carrier is preferably added following silicasome formation. Thus, after the silicasome is formed and loaded with suitable drug(s), the silicasome can be diluted into pharmaceutically acceptable carriers such as normal saline. These compositions may be sterilized by conventional, well-known sterilization techniques. The resulting aqueous solutions, suspensions, dispersions, emulsions, *etc.*, may be packaged for use or filtered under aseptic conditions. In certain embodiments the silicasomes are lyophilized, the lyophilized preparation being combined with a sterile aqueous solution prior to administration. The compositions may also contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH-adjusting and buffering agents, tonicity adjusting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, *etc.*

[0341] Additionally, in certain embodiments, the pharmaceutical formulation may include lipid-protective agents that protect lipids against free-radical and lipid-peroxidative damages on storage. Lipophilic free-radical quenchers, such as alpha-tocopherol and water-soluble iron-specific chelators, such as ferrioxamine, are suitable.

[0342] The concentration of silicasomes in the pharmaceutical formulations can vary widely, *e.g.*, from less than approximately 0.05%, usually at least approximately 2 to 5% to as much as 10 to 50%, or to 40%, or to 30% by weight and are selected primarily by fluid volumes, viscosities, *etc.*, in accordance with the particular mode of administration selected. For example, the concentration may be increased to lower the fluid load associated with treatment. This may be particularly desirable in patients having atherosclerosis-associated congestive heart failure or severe hypertension. Alternatively, silicasomes composed of irritating lipids may be diluted to low concentrations to lessen inflammation at the site of administration. The amount of silicasomes administered will depend upon the particular drug used, the disease state being treated and the judgment of the clinician but will generally be between approximately 0.01 and approximately 50 mg per kilogram of body weight, preferably between approximately 0.1 and approximately 5 mg per kg of body weight.

[0343] In some embodiments, *e.g.*, it is desirable to include polyethylene glycol (PEG)-modified phospholipids in the silicasomes. Alternatively or additionally, in certain embodiments, PEG-ceramide, or ganglioside G_{MI}-modified lipids can be incorporated in the silicasomes. Addition of such components helps prevent silicasome aggregation and provides for increasing circulation lifetime and increasing the delivery of the loaded silicasomes to the target tissues. In certain embodiments the concentration of the PEG-modified phospholipids, PEG-ceramide, or G_{MI}-modified lipids in the silicasome will be approximately 1 to 15%.

[0344] In some embodiments, overall silicasome charge is an important determinant in silicasome clearance from the blood. It is believed that charged silicasomes will be typically taken up more rapidly by the reticuloendothelial system (*see, e.g.*, Juliano (1975), *Biochem. Biophys. Res. Commun.* 63: 651-658 discussing liposome clearance by the RES) and thus have shorter half-lives in the bloodstream. Silicasomes with prolonged circulation half-lives are typically desirable for therapeutic uses. For instance, in certain embodiments, silicasomes that are maintained from 8 hrs, or 12 hrs, or 24 hrs, or greater are desirable.

[0345] In another example of their use, drug-loaded silicasomes can be incorporated into a broad range of topical dosage forms including but not limited to gels, oils, emulsions, and the like, *e.g.*, for the treatment of a topical cancer. For instance, in some embodiments the suspension containing the drug-loaded silicasomes is formulated and administered as a topical cream, paste, ointment, gel, lotion, and the like.

[0346] In some embodiments, pharmaceutical formulations comprising silicasomes described herein additionally incorporate a buffering agent. The buffering agent may be any pharmaceutically acceptable buffering agent. Buffer systems include, but are not limited to citrate buffers, acetate buffers, borate buffers, and phosphate buffers. Examples of buffers include, but are not limited to citric acid, sodium citrate, sodium acetate, acetic acid, sodium phosphate and phosphoric acid, sodium ascorbate, tartaric acid, maleic acid, glycine, sodium lactate, lactic acid, ascorbic acid, imidazole, sodium bicarbonate and carbonic acid, sodium succinate and succinic acid, histidine, and sodium benzoate, benzoic acid, and the like.

[0347] In some embodiments, pharmaceutical formulations comprising silicasomes described herein additionally incorporate a chelating agent. The chelating agent may be any pharmaceutically acceptable chelating agent. Chelating agents include, but are not limited to ethylene diaminetetraacetic acid (also synonymous with EDTA, edetic acid, versene acid, and sequestrene), and EDTA derivatives, such as dipotassium edetate, disodium edetate, edetate calcium disodium, sodium edetate, trisodium edetate, and potassium edetate. Other chelating agents include citric acid (*e.g.*, citric acid monohydrate) and derivatives thereof. Derivatives of citric acid include anhydrous citric acid, trisodiumcitrate-dihydrate, and the like. Still other chelating agents include, but are not limited to, niacinamide and derivatives thereof and sodium deoxycholate and derivatives thereof.

[0348] In some embodiments, pharmaceutical formulations comprising silicasomes described herein additionally incorporate a bioactive agent contain an antioxidant. The antioxidant may be any pharmaceutically acceptable antioxidant. Antioxidants are well known to those of ordinary skill in the art and include, but are not limited to, materials such as ascorbic acid, ascorbic acid derivatives (*e.g.*, ascorbylpalmitate, ascorbylstearate, sodium ascorbate, calcium ascorbate, *etc.*), butylated hydroxy anisole, butylated hydroxy toluene, alkylgallate, sodium meta-bisulfate, sodium bisulfate, sodium dithionite, sodium thioglycollic acid, sodium formaldehyde sulfoxylate, tocopherol and derivatives thereof, (d-alpha tocopherol, d-alpha tocopherol acetate, dl-alpha tocopherol acetate, d-alpha tocopherol succinate, beta tocopherol, delta tocopherol, gamma tocopherol, and d-alpha tocopherol polyoxyethylene glycol 1000 succinate) monothioglycerol, sodium sulfite and N-acetyl cysteine. In certain embodiments such materials, when present, are typically added in ranges from 0.01 to 2.0%.

[0349] In some embodiments, pharmaceutical formulations comprising silicasomes described herein are formulated with a cryoprotectant. The cryoprotecting agent may be any pharmaceutically acceptable cryoprotecting agent. Common cryoprotecting agents include, but are not limited to, histidine, polyethylene glycol, polyvinyl pyrrolidine, lactose, sucrose, mannitol, polyols, and the like.

[0350] In some embodiments, pharmaceutical formulations comprising silicasomes described herein are formulated with an isotonic agent. The isotonic agent can be any pharmaceutically acceptable isotonic agent. This term is used in the art interchangeably with iso-osmotic agent, and is known as a compound that is added to the pharmaceutical preparation to increase the osmotic pressure, *e.g.*, in some embodiments to that of 0.9% sodium chloride solution, which is iso-osmotic with human extracellular fluids, such as plasma. Illustrative isotonicity agents include, but are not limited to, sodium chloride, mannitol, sorbitol, lactose, dextrose and glycerol.

[0351] In certain embodiments pharmaceutical formulations of the silicasomes may optionally comprise a preservative. Common preservatives include, but are not limited to, those selected from the group consisting of chlorobutanol, parabens, thimerosal, benzyl alcohol, and phenol. Suitable preservatives include but are not limited to: chlorobutanol (*e.g.*, 0.3-0.9% w/v), parabens (*e.g.*, 0.01-5.0%), thimerosal (*e.g.*, 0.004-0.2%), benzyl alcohol (*e.g.*, 0.5-5%), phenol (*e.g.*, 0.1-1.0%), and the like.

[0352] In some embodiments, pharmaceutical formulations comprising silicasomes are formulated with a humectant, *e.g.*, to provide a pleasant mouth-feel in oral applications. Humectants known in the art include, but are not limited to, cholesterol, fatty acids, glycerin, lauric acid, magnesium stearate, pentaerythritol, and propylene glycol.

[0353] In some embodiments, an emulsifying agent is included in the formulations, for example, to ensure complete dissolution of all excipients, especially hydrophobic components such as benzyl alcohol. Many emulsifiers are known in the art, *e.g.*, polysorbate 60.

[0354] For some embodiments related to oral administration, it may be desirable to add a pharmaceutically acceptable flavoring agent and/or sweetener. Compounds such as saccharin, glycerin, simple syrup, and sorbitol are useful as sweeteners.

Administration and Therapy

[0355] The cargo (*e.g.*, drug) loaded silicasomes can be administered to a subject (*e.g.*, patient) by any of a variety of techniques.

[0356] In certain embodiments the pharmaceutical formulations are administered parenterally, *e.g.*, intraarticularly, intravenously, intraperitoneally, subcutaneously, or intramuscularly. In some embodiments, the pharmaceutical compositions are administered intravenously, intraarterially, or intraperitoneally by a bolus injection (*see, e.g.*, U.S. Pat. Nos. 3,993,754; 4,145,410; 4,235,871; 4,224,179; 4,522,803; and 4,588,578 describing administration of liposomes). Particular pharmaceutical formulations suitable for this administration are found in Remington's Pharmaceutical Sciences, Mack Publishing Company, Philadelphia, Pa., 17th ed. (1985). Typically, the formulations comprise a solution of the silicasomes suspended in an acceptable carrier, preferably an aqueous carrier. As noted above, suitable aqueous solutions include, but are not limited to physiologically compatible buffers such as Hanks solution, Ringer's solution, or physiological (*e.g.*, 0.9% isotonic) saline buffer and/or in certain emulsion formulations. The solution(s) can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. In certain embodiments the active agent(s) can be provided in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use. For transmucosal administration, and/or for blood/brain barrier passage, penetrants appropriate to the barrier to be permeated can be used in the formulation. These compositions may be sterilized by conventional, well-known sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous solution prior to administration. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, *etc.*, *e.g.*, as described above.

[0357] In other methods, the pharmaceutical formulations containing silicasomes described herein may be contacted with the target tissue by direct application of the preparation to the tissue. The application may be made by topical, "open" or "closed" procedures. By "topical" it is meant the direct application of the pharmaceutical preparation to a tissue exposed to the environment, such as the skin, oropharynx, external

auditory canal, and the like. Open procedures are those procedures that include incising the skin of a patient and directly visualizing the underlying tissue to which the pharmaceutical formulations are applied. This is generally accomplished by a surgical procedure, such as a thoracotomy to access the lungs, abdominal laparotomy to access abdominal viscera, or other direct surgical approaches to the target tissue. Closed procedures are invasive procedures in which the internal target tissues are not directly visualized, but accessed via inserting instruments through small wounds in the skin. For example, the preparations may be administered to the peritoneum by needle lavage. Likewise, the pharmaceutical preparations may be administered to the meninges or spinal cord by infusion during a lumbar puncture followed by appropriate positioning of the patient as commonly practiced for spinal anesthesia or metrizamide imaging of the spinal cord. Alternatively, the preparations may be administered through endoscopic devices. In certain embodiments the pharmaceutical formulations are introduced via a cannula.

[0358] In certain embodiments the pharmaceutical formulations comprising silicasomes described herein are administered via inhalation (*e.g.*, as an aerosol). Inhalation can be a particularly effective delivery route for administration to the lungs and/or to the brain. For administration by inhalation, the silicasomes are conveniently delivered in the form of an aerosol spray from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0359] In certain embodiments, the silicasomes described herein are formulated for oral administration. For oral administration, suitable formulations can be readily formulated by combining the silicasome(s) with pharmaceutically acceptable carriers suitable for oral delivery well known in the art. Such carriers enable the active agent(s) described herein to be formulated as tablets, pills, dragees, caplets, lozenges, gelcaps, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. For oral solid formulations such as, for example, powders, capsules and tablets, suitable excipients can include fillers such as sugars (*e.g.*, lactose, sucrose, mannitol and sorbitol), cellulose preparations (*e.g.*, maize starch, wheat starch, rice starch, potato starch, gelatin,

gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose), synthetic polymers (*e.g.*, polyvinylpyrrolidone (PVP)), granulating agents; and binding agents. If desired, disintegrating agents may be added, such as the cross-linked polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. If desired, solid dosage forms may be sugar-coated or enteric-coated using standard techniques. The preparation of enteric-coated particles is disclosed for example in U.S. Pat. Nos. 4,786,505 and 4,853,230.

[0360] In various embodiments the silicasome(s) can be formulated in rectal or vaginal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. Methods of formulating active agents for rectal or vaginal delivery are well known to those of skill in the art (*see, e.g.*, Allen (2007) *Suppositories*, Pharmaceutical Press) and typically involve combining the active agents with a suitable base (*e.g.*, hydrophilic (PEG), lipophilic materials such as cocoa butter or Witepsol W45), amphiphilic materials such as Suppocire AP and polyglycolized glyceride, and the like). The base is selected and compounded for a desired melting/delivery profile.

[0361] The route of delivery of silicasomes can also affect their distribution in the body. Passive delivery of silicasomes involves the use of various routes of administration *e.g.*, parenterally, although other effective administration forms, such as intraarticular injection, inhalant mists, orally active formulations, transdermal iontophoresis, or suppositories are also envisioned. Each route produces differences in localization of the silicasomes.

[0362] Because dosage regimens for pharmaceutical agents are well known to medical practitioners, the amount of the liposomal pharmaceutical agent formulations that is effective or therapeutic for the treatment of a disease or condition in mammals and particularly in humans will be apparent to those skilled in the art. The optimal quantity and spacing of individual dosages of the formulations herein will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and such optima can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, *e.g.*, the number of doses given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

[0363] In certain embodiments the silicasomes and/or pharmaceutical formations thereof can be used therapeutically in animals (including man) in the treatment of various cancers, or various infections, and the like including conditions that require: (1) repeated administrations, (2) the sustained delivery of the drug in its bioactive form, or (3) the decreased toxicity with suitable efficacy compared with the free drug in question. In various embodiments the silicasomes and/or pharmaceutical formations thereof are administered in a therapeutically effective dose. The term "therapeutically effective" as it pertains to the silicasomes described herein and formulations thereof means that a biologically active substance present or and/or in the silicasome provided/released in a manner sufficient to achieve a particular medical effect for which the biologically active substance (therapeutic agent) is intended. Examples, without limitation of desirable medical effects that can be attained are chemotherapy, antibiotic therapy, and regulation of metabolism. Thus, for example, a therapeutically effective dose for cancer chemotherapy may be a dose (and/or dosage regimen) effective to slow the growth and/or proliferation of cancer cells, and/or to slow, stop the growth of a solid tumor or shrink or eliminate a solid tumor, and/or slow, stop the proliferation of metastatic cells, and the like. A therapeutically effective dose for treating an infection can be a dose (and/or dosage regimen) sufficient to inhibit the growth and/or proliferation of a pathogen, and/or to kill a pathogen, and/or to mitigate one or more symptoms produced by the pathogen.

[0364] Exact dosages will vary depending upon such factors as the particular therapeutic agent and desirable medical effect, as well as patient factors such as age, sex, general condition, and the like. Those of skill in the art can readily take these factors into account and use them to establish effective therapeutic concentrations without resort to undue experimentation.

[0365] For administration to humans (or to non-human mammals) in the curative, remissive, retardive, or prophylactic treatment of diseases the prescribing physician will ultimately determine the appropriate dosage of the drug for a given human (or non-human) subject, and this can be expected to vary according to the age, weight, and response of the individual as well as the nature and severity of the patient's disease. In certain embodiments the dosage of the drug provided by the silicasome(s) can be approximately equal to that employed for the free drug. However as noted above, the silicasomes described herein can significantly reduce the toxicity of the drug(s) administered thereby and significantly

increase a therapeutic window. Accordingly, in some cases dosages in excess of those prescribed for the free drug will be utilized.

[0366] In certain embodiments, the dose of encapsulated drug administered at a particular time point will be in the range from about 1 to about 1,000 mg/m²/day, or to about 800 mg/m²/day, or to about 600 mg/m²/day, or to about 400 mg/m²/day. For example, in certain embodiments a dosage (dosage regiment) is utilized that provides a range from about 1 to about 350 mg/m²/day, 1 to about 300 mg/m²/day, 1 to about 250 mg/m²/day, 1 to about 200 mg/m²/day, 1 to about 150 mg/m²/day, 1 to about 100 mg/m²/day, from about 5 to about 80 mg/m²/day, from about 5 to about 70 mg/m²/day, from about 5 to about 60 mg/m²/day, from about 5 to about 50 mg/m²/day, from about 5 to about 40 mg/m²/day, from about 5 to about 20 mg/m²/day, from about 10 to about 80 mg/m²/day, from about 10 to about 70 mg/m²/day, from about 10 to about 60 mg/m²/day, from about 10 to about 50 mg/m²/day, from about 10 to about 40 mg/m²/day, from about 10 to about 20 mg/m²/day, from about 20 to about 40 mg/m²/day, from about 20 to about 50 mg/m²/day, from about 20 to about 90 mg/m²/day, from about 30 to about 80 mg/m²/day, from about 40 to about 90 mg/m²/day, from about 40 to about 100 mg/m²/day, from about 80 to about 150 mg/m²/day, from about 80 to about 140 mg/m²/day, from about 80 to about 135 mg/m²/day, from about 80 to about 130 mg/m²/day, from about 80 to about 120 mg/m²/day, from about 85 to about 140 mg/m²/day, from about 85 to about 135 mg/m²/day, from about 85 to about 130 mg/m²/day, from about 85 to about 120 mg/m²/day, or from about 85 to about 120 mg/m²/day. In certain embodiments the dose administered at a particular time point may also be about 130 mg/m²/day, about 120 mg/m²/day, about 100 mg/m²/day, about 90 mg/m²/day, about 85 mg/m²/day, about 80 mg/m²/day, about 70 mg/m²/day, about 60 mg/m²/day, about 50 mg/m²/day, about 40 mg/m²/day, about 30 mg/m²/day, about 20 mg/m²/day, about 15 mg/m²/day, or about 10 mg/m²/day.

[0367] Dosages may also be estimated using *in vivo* animal models, as will be appreciated by those skill in the art. In this regard, with respect to the irinotecan-loaded silicasomes described herein, it is noted that the effective therapeutic dose of the Ir-silicasome in a KPC-derived orthotopic animal model is about 40 mg/kg, which is equivalent to 120 mg/m² in a 70 Kg human subject (Liu, *et al.* (2016) *ACS Nano*, 10: 2702-2715). Fibonacci analysis indicates this dose can be achieved by starting and intermediary doses of 40 and 80 mg/m².

[0368] The dose administered may be higher or lower than the dose ranges described herein, depending upon, among other factors, the bioavailability of the composition, the tolerance of the individual to adverse side effects, the mode of administration and various factors discussed above. Dosage amount and interval may be adjusted individually to provide plasma levels of the composition that are sufficient to maintain therapeutic effect, according to the judgment of the prescribing physician. Skilled artisans will be able to optimize effective local dosages without undue experimentation in view of the teaching provided herein.

[0369] Multiple doses (*e.g.*, continuous or bolus) of the compositions as described herein may also be administered to individuals in need thereof of the course of hours, days, weeks, or months. For example, but not limited to, 1, 2, 3, 4, 5, or 6 times daily, every other day, every 10 days, weekly, monthly, twice weekly, three times a week, twice monthly, three times a month, four times a month, five times a month, every other month, every third month, every fourth month, *etc.*

Methods of treatment.

[0370] In various embodiments methods of treatment using the nanoparticle drug carrier(s) and/or pharmaceutical formulation(s) comprising nanoparticle drug carriers described herein are provided. In certain embodiments the method(s) comprise a method of treating a cancer. In certain embodiments the method can comprise administering to a subject in need thereof an effective amount of a nanoparticle drug carrier, and/or a pharmaceutical formulation comprising a nanoparticle drug carrier as described herein, where the drug in said nanoparticle drug carrier and/or said pharmaceutical formulation comprises an anti-cancer drug. In certain embodiments the nanoparticle drug carrier and/or pharmaceutical formulation is a primary therapy in a chemotherapeutic regimen. In certain embodiments the nanoparticle drug carrier and/or pharmaceutical formulation is a component in a multi-drug chemotherapeutic regimen. In certain embodiments the multi-drug chemotherapeutic regimen comprises at least two drugs selected from the group consisting of irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV). In certain embodiments the multi-drug chemotherapeutic regimen comprises at least three drugs selected from the group consisting of irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV). In certain embodiments the multi-drug chemotherapeutic regimen comprises at least irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV).

[0371] In various embodiments the nanoparticle drug carrier(s) and/or pharmaceutical formulation(s) comprising nanoparticle drug carriers described herein are effective for treating any of a variety of cancers. In certain embodiments the cancer is pancreatic ductal adenocarcinoma (PDAC). In certain embodiments the cancer is a cancer selected from the group consisting of acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical carcinoma, AIDS-related cancers (*e.g.*, Kaposi sarcoma, lymphoma), anal cancer, appendix cancer, astrocytomas, atypical teratoid/rhabdoid tumor, bile duct cancer, extrahepatic cancer, bladder cancer, bone cancer (*e.g.*, Ewing sarcoma, osteosarcoma, malignant fibrous histiocytoma), brain stem glioma, brain tumors (*e.g.*, astrocytomas, glioblastoma, brain and spinal cord tumors, brain stem glioma, central nervous system atypical teratoid/rhabdoid tumor, central nervous system embryonal tumors, central nervous system germ cell tumors, craniopharyngioma, ependymoma, breast cancer, bronchial tumors, burkitt lymphoma, carcinoid tumors (*e.g.*, childhood, gastrointestinal), cardiac tumors, cervical cancer, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative disorders, colon cancer, colorectal cancer, craniopharyngioma, cutaneous t-cell lymphoma, duct cancers *e.g.* (bile, extrahepatic), ductal carcinoma *in situ* (DCIS), embryonal tumors, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, extracranial germ cell tumor, extragonadal germ cell tumor, extrahepatic bile duct cancer, eye cancer (*e.g.*, intraocular melanoma, retinoblastoma), fibrous histiocytoma of bone, malignant, and osteosarcoma, gallbladder cancer, gastric (stomach) cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumors (*e.g.*, ovarian cancer, testicular cancer, extracranial cancers, extragonadal cancers, central nervous system), gestational trophoblastic tumor, brain stem cancer, hairy cell leukemia, head and neck cancer, heart cancer, hepatocellular (liver) cancer, histiocytosis, langerhans cell cancer, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, kaposi sarcoma, kidney cancer (*e.g.*, renal cell, Wilm's tumor, and other kidney tumors), langerhans cell histiocytosis, laryngeal cancer, leukemia, acute lymphoblastic (ALL), acute myeloid (AML), chronic lymphocytic (CLL), chronic myelogenous (CML), hairy cell, lip and oral cavity cancer, liver cancer (primary), lobular carcinoma *in situ* (LCIS), lung cancer (*e.g.*, childhood, non-small cell, small cell), lymphoma (*e.g.*, AIDS-related, Burkitt (*e.g.*, non-Hodgkin lymphoma), cutaneous T-Cell (*e.g.*, mycosis fungoides, Sézary syndrome), Hodgkin, non-Hodgkin, primary central nervous system (CNS)), macroglobulinemia, Waldenström, male breast cancer, malignant

fibrous histiocytoma of bone and osteosarcoma, melanoma (*e.g.*, childhood, intraocular (eye)), merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer, midline tract carcinoma, mouth cancer, multiple endocrine neoplasia syndromes, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndromes, chronic myeloid leukemia (CML), multiple myeloma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, oral cavity cancer, lip and oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, pancreatic neuroendocrine tumors (islet cell tumors), papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytoma, pituitary tumor, plasma cell neoplasm, pleuropulmonary blastoma, primary central nervous system (CNS) lymphoma, prostate cancer, rectal cancer, renal cell (kidney) cancer, renal pelvis and ureter, transitional cell cancer, rhabdomyosarcoma, salivary gland cancer, sarcoma (*e.g.*, Ewing, Kaposi, osteosarcoma, rhabdomyosarcoma, soft tissue, uterine), Sézary syndrome, skin cancer (*e.g.*, melanoma, merkel cell carcinoma, basal cell carcinoma, nonmelanoma), small intestine cancer, squamous cell carcinoma, squamous neck cancer with occult primary, stomach (gastric) cancer, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, trophoblastic tumor, ureter and renal pelvis cancer, urethral cancer, uterine cancer, endometrial cancer, uterine sarcoma, vaginal cancer, vulvar cancer, Waldenström macroglobulinemia, and Wilm's tumor.

[0372] In certain embodiments the nanoparticle drug carrier is not conjugated to an iRGD peptide and the nanoparticle drug carrier is administered in conjunction with an iRGD peptide (*e.g.*, the nanoparticle drug carrier and the iRGD peptide are co-administered as separate formulations).

[0373] In certain embodiments the method(s) comprise a method of treating an infection. In certain embodiments the method can comprise administering to a subject in need thereof an effective amount of a nanoparticle drug carrier, and/or a pharmaceutical formulation comprising a nanoparticle drug carrier as described herein, where the drug in said nanoparticle drug carrier and/or said pharmaceutical formulation comprises an anti-microbial or anti-viral agent. In certain embodiments the infection comprises a nosocomial infection. In certain embodiments the infection is caused by viral, bacterial, or fungal pathogens. In certain embodiments the infection comprises a bloodstream infection (BSI), pneumonia (*e.g.*, ventilator-associated pneumonia (VAP)), a gastrointestinal infection, a urinary tract infection (UTI), a surgical site infection (SSI), or a skin infection. In certain

embodiments the infection is caused by a pathogen such as *Staphylococcus aureus* (e.g., blood infection), *Escherichia coli* (e.g., UTI), Enterococci (e.g., blood, UTI, wound), *Pseudomonas aeruginosa* (e.g., kidney or respiratory infection), *Mycobacterium tuberculosis* (e.g., lung), and the like. In certain embodiments the infection is a viral
5 infection (e.g., HIV, hepatitis B, hepatitis C, etc.).

[0374] In certain embodiments the infection is caused by a drug-resistant pathogen. Illustrative drug-resistant pathogens include, but are not limited to methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant *Enterococcus* (VRE) and multi-
10 carbapenemase-producing bacteria (KPC).

[0375] In various embodiments of these treatment methods, the nanoparticle drug carrier and/or pharmaceutical formulation is administered via a route selected from the group consisting of intravenous administration, intraarterial administration, intracerebral administration, intrathecal administration, oral administration, aerosol administration,
15 administration via inhalation (including intranasal and intratracheal delivery, intracranial administration via a cannula, and subcutaneous or intramuscular depot deposition. In certain embodiments the nanoparticle drug carrier and/or pharmaceutical formulation is administered as an injection, from an IV drip bag, or via a drug-delivery cannula. In various embodiments the subject is a human and in other embodiments the subject is a non-human
20 mammal.

Kits.

[0376] In certain embodiments, kits are containing the lipid bilayer coated nanoparticle drug carriers described herein for the treatment of a pathology (e.g., a cancer, a microbial infection, a viral infection, etc.). The kits typically comprise a drug-loaded
25 silicasome as described herein and/or an immunoconjugate comprising a drug-loaded silicasome described herein. In certain embodiments the silicasome contains irinotecan. In certain embodiments the silicasome has attached thereto an iRGD peptide while in other embodiments the kit contains a separate iRGD peptide formulated for co-administration with the drug (e.g., irinotecan) loaded silicasome or silicasome immunoconjugate.

30 **[0377]** Additionally, in certain embodiments, the kits can include instructional materials disclosing means of use of the drug-loaded silicasome or silicasome

immunoconjugate (*e.g.* as a therapeutic for a pancreatic cancer, gastric cancer, cervical cancer, ovarian cancer, *etc.*).

[0378] In addition, the kits optionally include labeling and/or instructional materials providing directions (*e.g.*, protocols) for the use of the nanoparticle drug carriers described herein, *e.g.*, alone or in combination for the treatment of various cancers. Instructional materials can also include recommended dosages, description(s) of counterindications, and the like.

[0379] While the instructional materials in the various kits typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to electronic storage media (*e.g.*, magnetic discs, tapes, cartridges, chips), optical media (*e.g.*, CD ROM), and the like. Such media may include addresses to internet sites that provide such instructional materials.

EXAMPLES

[0380] The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1

Irinotecan Delivery by Lipid-Coated Mesoporous Silica Nanoparticles Shows Improved Efficacy and Safety over Liposomes for Pancreatic Cancer

[0381] Because of the concern that liposomal carriers may fail to improve the safety of highly toxic drugs such as irinotecan, we were interested in using a supported lipid bilayer (LB) that is applied to mesoporous silica nanoparticles (MSNPs) to see if this will result in a more stable carrier (than the liposome). Due to the large internal surface area that can be used for drug packaging, tunable pore sizes, carrier stability, and controlled drug release abilities, MSNPs have been demonstrated to constitute a versatile and multifunctional nanocarrier platform for cancer therapy (Lu *et al.* (2007) *Small*, 3: 1341-1346; Slowing *et al.* (2008) *Adv. Drug Deliv. Rev.* 60: 1278-1288; Meng *et al.* (2010) *J. Am. Chem. Soc.* 132(36): 12690-12697; Lee *et al.* (2010) *Angew. Chem.* 122: 8390-8395; Meng *et al.* (2011) *ACS Nano*, 5(5): 4131-4144; He and Shi (2011) *J. Mater. Chem* 21: 5845-5855; Gao *et al.* (2011) *ACS Nano*, 5: 9788-9798; Li *et al.* (2012) *Chem. Soc. Rev.* 41: 2590-2605; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Meng *et al.* (2013) *ACS Nano*, 7(2): 994-1005; Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801; Meng *et al.* (2013) *ACS*

Nano, 7(11): 10048-10065; Argyo *et al.* (2014) *Chem. Mater.* 26: 435-451; Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Moreover, MSNPs are biodegradable and proven to be safe in extensive animal testing (Meng *et al.* (2011) *ACS Nano*, 5(5): 4131-4144; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Meng *et al.* (2013) *ACS Nano*, 7(2): 994-1005; Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801; Meng *et al.* (2013) *ACS Nano*, 7(11): 10048-10065; Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557; Lu *et al.* (2010) *Small*, 6(16): 1794-18050. While it is possible to introduce irinotecan or camptothecin into MSNPs by soaking-in, this method is relatively inefficient (loading capacity <10 wt %) (Lu *et al.* (2007) *Small*, 3: 1341-1346; He *et al.* (2010) *Biomaterials* 31: 3335-3346) and may result in poor drug retention (Meng *et al.* (2010) *J. Am. Chem. Soc.* 132(36): 12690-12697). However, we have developed a biofilm technique that can be used for rapid GEM encapsulation by a supported LB (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). While a number of LB coating procedures have been published, including liposomal interaction and fusion with the particle surface (Liu, *et al.* (2009) *J. Am. Chem. Soc.*, 131(22): 7567-7569; Liu, *et al.* (2009) *J. Am. Chem. Soc.*, 131(4): 1354-1355) or solvent exchange utilizing EtOH-dispersed lipid solutions (Cauda *et al.* (2010) *Nano Lett.* 10: 2484-2492), the use of a biofilm technique provides more reproducible and complete coating, rapid encapsulation, and improved carrier stability (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Not only has this allowed LB-MSNPs to achieve a loading capacity of up to 40 wt % GEM, but it has also enabled co-delivery of paclitaxel (PTX), which could be incorporated in the LB (*Id.*). This allowed us to develop a synergistic and ratiometric-designed carrier for PDAC treatment in an orthotopic human PDAC model in mice (*Id.*). Recently, Zhang *et al.* reported a polymer-lipid-supported MSNP for delivery of irinotecan, but the drug loading was lower (*i.e.*, ~16 wt %) (Zhang *et al.* (2014) *Biomaterials*, 35: 3650-3665).

[0382] In this example, we introduce, *inter alia*, a novel design feature for the LB-MSNP platform in which the LB is used to encapsulate a protonating agent; this allows remote loading of a high irinotecan drug load that utilizes the drug's weak basic properties (pKa = 8.1). Moreover, the increased ability of the supported LB also allowed us to perform comparative analysis to a liposomal equivalent, where a nonsupported LB vesicle was used for remote loading of irinotecan by a trapping agent, triethylammonium sucrose octasulfate (TEA8SOS) (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277; Von Hoff *et al.* (2013) *Br. J. Cancer*, 109(4): 920-925). Not only did the MSNP carrier achieve higher loading capacity and tumor killing than the liposomal formulation in a robust

orthotopic PDAC model, but it also prevented irinotecan toxicity due to improved carrier stability and reduced leakage. This provides an innovative approach for reducing the toxicity of highly toxic chemotherapeutic agents by a LB-coated carrier, which allows the use of irinotecan as a first-line treatment option for PDAC rather than being reserved for GEM treatment failure (current FDA-improved indication for the use of the liposomal carrier).

RESULTS

High Irinotecan Loading in MSNP through the Use of a Coated LB and a Proton Gradient.

10 **[0383]** Our first attempt was to produce a MSNP carrier that encapsulates irinotecan by a LB coat that provides rapid and uniform pore sealing (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). This allowed passive entrapment of 22 wt % irinotecan (relative to MSNP weight). In order to further improve the loading capacity, we also determined whether entrapment of a protonating agent that converts amphipathic irinotecan, capable of
15 diffusing across the LB, into a hydrophilic derivative that is retained in the pores (Figure 1A, panel A1). We and others have used this entrapment procedure in liposomes for effective import and encapsulation of weak basic drugs, such as GEM and irinotecan (Chou *et al.* (2003) *J. Biosci. Bioeng.*, 95(4): 405-408; Meng *et al.* (2013) *ACS Nano*, 7(11): 10048-10065; Haran *et al.* (1993) *Biochim. Biophys. Acta, Biomembr.* 1151: 201-215).
20 While it has never been used in a MSNP platform previously, the polyanionic compound, TEA₈SOS, was used in MM-398 to entrap irinotecan (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). TEA₈SOS is a proton- generating agent that releases eight H⁺ ions and octavalent SOS⁸⁻ upon hydrolysis (Figure 1A, panel A2). Ion-exchange chromatography was used to generate TEA₈SOS, which was soaked into MSNP, as described below in the
25 Methods section (*Id.*). The soaked particles were introduced to a round-bottom flask coated with a lipid biofilm, composed of DSPC/cholesterol/DSPE-PEG2000 at a molar ratio of 3:2:0.15. The optimal lipid:particle ratio was determined to be 1.1:1.0. Sonication of the suspension yielded LB-coated particles, which contain the entrapment agent (Figure 1A, panel A1). These particles were immediately incubated in an irinotecan solution, allowing
30 the drug to be imported, protonated, and entrapped in the pores as a gel-like precipitate in association with SOS⁸⁻ (Figure 1A, panels A1 and A2, Fig. 1B). This allowed us to achieve an irinotecan loading capacity up to 80 wt % (Figure 1B), which approximates the

theoretical maximum loading capacity (~ 100 wt %) of the porous carrier with a combined surface area of $850 \text{ m}^2/\text{g}$ and a pore volume of $\sim 0.7 \text{ cm}^3/\text{g}$, as previously demonstrated (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Subsequent studies were performed with particles containing 50 wt % irinotecan.

5 **[0384]** In order to compare the efficacy and safety of the irinotecan- loaded LB-MSNPs (Ir-LB-MSNPs) with a liposomal carrier using TEA₈SOS, liposomes were constructed by a procedure analogous to that of MM-398 (Figure 1A, panel A3) (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). This yielded a liposome (designated “Ir-liposome”) with ~ 40 wt % irinotecan loading capacity; that is considerably
10 higher than producing a liposome without the entrapment agent (~ 5 wt %) (Figure 1B). Cryo-electron microscopy (cryoEM) revealed particle sizes of ~ 75 and ~ 80 nm for liposomes and LB-MSNPs, respectively (Figure 1C). High-magnification cryoEM images confirmed that there was uniform coating of Ir-LB-MSNPs by an intact ~ 7 nm thick bilayer. Low-magnification cryoEM images showed that the LB was intact in $>99\%$ of the particles
15 (Figure 11). This confirms the reproducibility of the one-step encapsulation protocol, which holds significant advances over other methods of MSNP bilayer coating (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557).

[0385] Since our major hypothesis is that LB-MSNPs could improve the carrier stability compared to liposomes, both carriers were incubated in serum and lyophilized to
20 determine their drug retention ability and irinotecan leakage. Both carriers were incubated for 24 h at 37°C in 100% serum, with continuous and gentle shaking. Since adsorption of serum proteins interferes with cryoEM visualization, high-performance liquid chromatography (HPLC) analysis was used, instead, to determine drug release in the carrier suspensions; this showed $<6\%$ premature irinotecan release by LB-MSNPs, compared to
25 $\sim 22\%$ from liposomes (Figure 1D). However, particle storage in a serum-free suspension for 90 days at 4°C showed that both carriers had high colloidal stability, with minor ($<5\%$) premature drug release or change in hydrodynamic particle size ($<5\%$). Conversely, particle lyophilization in a cryo-protective 5% dextrose solution⁴⁰ showed significant differences in carrier stability upon resuspension in water (Figure 1E). Thus, while Ir-LB-
30 MSNPs did not change the hydrodynamic size and only led to 2.7% drug release, liposomes were visibly aggregated and released 20.8% of their irinotecan content. This confirms that although the LB-coated carriers look morphologically similar, LB-MSNPs are significantly more stable than liposomes.

Improved Biodistribution, Pharmacokinetics (PK), and Irinotecan Delivery by LB-MSNP compared to Liposomes in a Kras-Derived Orthotopic PDAC Model

[0386] Biodistribution and PK studies were performed in immuno-competent B6/129 mice, orthotopically implanted with a luciferase-expressing cell line derived from a spontaneous PDAC tumor in a transgenic Kras^{LSL-G12D/+}; Trp53^{LSL-R172H/+}; Pdx-1-Cre (KPC) animal (Hingorani *et al.* (2005) *Cancer Cell*, 7: 469-483). The KPC-derived orthotopic model mimics human PDAC in the Kras mutation, expression of a relatively abundant stroma, local tissue invasion, and development of metastases (Tseng *et al.* (2010) *Clin. Cancer Res.* 16: 3684- 3695; Provenzano *et al.* (2013) *Br. J. Cancer*, 108: 1-8; Torres *et al.* (2013) *PLoS One*, 8: e80580). Primary tumors developed at the orthotopic implant site within 2–3 weeks, with appearance of metastases at ~5 weeks, agreeing with the literature (Figure 12) (*Id.*). In order to follow the biodistribution of the irinotecan carriers to the tumor site, animals were IV injected with near- infrared (NIR)-labeled (Dylight 680) liposomes or LB-MSNPs, 3 weeks after orthotopic implantation (Figure 2A). IVIS imaging was obtained prior to (column 2) and following IV injection of 100 mg/kg labeled LB-MSNPs or liposomes at the indicated time points (columns 3-6, Figure 2A). This was coupled with bioluminescence IVIS imaging to detect luciferase expression in the developing tumors in animals receiving intraperitoneal (IP) injection of D-Luciferin (Figure 2A, first column). Robust fluorescence intensity was observed at the tumor sites within 2 h of LB-MSNP injection, following which the signal was sustained for at least 48 h. In contrast, the NIR signal intensity was dimmer and disappeared more rapidly in mice injected with liposomes (with similar labeling efficiency). This was also confirmed by ex vivo imaging of the tumors and major organs collected from the animals, following sacrifice after 24 h postinjection (Figure 2B). Ex vivo imaging was also carried out after 48 h, which confirmed stronger NIR signal intensity in the operator-defined region of interest (ROI) in LB-MSNP compared to liposome-treated animals (Figure 13). In addition to abundant particle uptake at the tumor site, the liver and spleen were also major sites of particle distribution. Little signaling was obtained in the lung, heart, and kidney. Inductively coupled plasma optical emission spectrometry (ICP-OES) to quantify the silicon (Si) abundance in the tumor tissues and major organs demonstrated that ~3% of the total administered elemental Si dose could distribute to the developing tumor sites (Figure 14). This represents exceptionally good biodistribution compared to liposomes, polymeric micelles, gelatin, or MoS₂/Fe₃O₄ nanoparticles, where the percentage of particles distributing to the PDAC site amounted to 0.2–2.4% (Yoshida *et al.* (2012) *PLoS One*, 7:

e39545; Yu *et al.* (2015) *Theranostics*, 5: 931-945; Cabral *et al.* (2011) *Nat. Nanotechnol.* 6: 815-823; Guo *et al.* (2013) *Biomaterials* 34: 8323-8332; Xu *et al.* (2013) *Mol.*

Pharmaceutics, 10: 2031-2044). We also obtained tumor sections for visualization of the intratumoral distribution of NIR-labeled particles by confocal microscopy; this

5 demonstrated higher particle abundance at the tumor site after 24 h for LB-MSNPs compared to liposomes (Figure 2B). In order to assess irinotecan content at the tumor site, the equivalent of 60 mg/kg drug was injected as a single dose in the form of free drug, Ir-LB-MSN (60 mg/kg drug; 120 mg/kg particle dose) or Ir-liposomes (60 mg/kg drug; 150 mg/kg particle dose). Animals were sacrificed after 0.5, 4, 24, and 48 h for blood collection
10 and tumor harvesting. These tissues were used for HPLC analysis to quantify the irinotecan tissue content, which was 5- and 2-fold higher for Ir-LB-MSNPs and Ir-liposomes than for free drug at 24 h (Figure 2C). Circulatory half-life for irinotecan was calculated by using HPLC quantification of the plasma drug concentration and to 11, 8.7, and 1.0 h for LB-MSNPs, liposomes, and free drug, respectively (Figure 2D). Logistical constraint of the
15 volume of blood obtainable from mice precluded us from determining the blood residence time of the carriers. We also performed a biodistribution and PK study in a subcutaneous PANC-1 xenograft model, in which we observed 30- and 4-fold increases in tumor drug content for Ir-LB-MSNPs and Ir-liposomes, respectively, over free drug (Figure 15). We also compared the drug content of individual organs (Figure 16). This demonstrated a 0.8-,
20 0.2-, and 0.2-fold higher drug content in liver, spleen, and intestinal tissues, respectively, for the Ir-liposomes compared to the Ir-LB-MSN at 24 h postinjection. No significant difference was seen for the kidney.

Ir-LB-MSN Provides More Efficacious Killing than Free Drug or Ir-Liposomes in an Orthotopic PDAC Model.

25 [0387] In order to develop a dose-seeking schedule for efficacy assessment, we used a protocol from the U.S. National Cancer Institute to determine the maximum tolerated dose (MTD) in healthy mice (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). This assessment demonstrated a MTD of 60, 295, and 295 mg/kg for free irinotecan, Ir-liposomes, and Ir-LB-MSNs, respectively (Figure 3A). A drug dose of 40 mg/kg
30 (equivalent to 66.7 or 13.3% of the MTD for free drug or encapsulated drug quantities) was chosen for subsequent experiments. These doses are equivalent to nanoparticle doses of 80 or 100 mg/kg for Ir-LB-MSNs or Ir-liposomes, respectively. IV injection commenced on day 14 after orthotopic tumor implantation of 2×10^6 KPC cells; at this time, the mean

primary tumor size was ~5 mm, without macro-metastases. IV injection was repeated every 4 days, for up to 8 repeat repetitions (Figure 3B). The control group included animals receiving IV saline only. Orthotopic tumor growth was monitored at set time points by IVIS bioluminescence imaging (Figure 3B). Quantitative expression of tumor growth by
5 IVIS software to determine imaging intensity in the ROI, demonstrated significantly slower tumor growth by Ir-LB- MSNP treatment compared to saline, free drug, or Ir-liposomes (Figure 3B). Antitumor efficacy was also measured according to the rates of apoptosis in tumor explants, collected from animals sacrificed on days 40–47. Utilizing an
10 immunohistochemistry (IHC) staining protocol for the detection of activated (cleaved) caspase-3, we observed ~25% apoptotic cell death in Ir-LB-MSNP-treated mice, compared to ~6 and ~11% for free irinotecan or Ir-liposomes, respectively (Figure 3C).

[0388] Representative IHC images are shown in Figure 17. Although we did not set out to perform an official survival study, it was possible to use a Kaplan-Meier data display after 47 days to compare the number of surviving animals at different stages of the
15 investigation (Figure 18). This demonstrated improved survival for both carriers over free drug, including better survival rate of Ir-LB-MSNP over the liposomal preparation (Figure 18).

[0389] Animal autopsy was performed to assess local tumor spread and appearance of metastases (Figure 3D). In addition to direct tumor invasion of the stomach, intestines,
20 liver, spleen, kidneys, diaphragm, and abdominal wall, numerous macroscopic metastatic foci could be seen in saline-treated animals, which also developed hemorrhagic ascites. While free irinotecan failed to affect metastasis, Ir-liposomes provided a moderate reduction of bioluminescence intensity at the metastatic sites (Figure 3D). In striking contrast, Ir-LB-MSNP treatment exerted a potent inhibitory effect, with evidence of dim bioluminescence
25 outside the primary tumor region during IVIS imaging (Figure 3D). This was especially prominent in the region of the kidney, which showed a higher but nonsignificant content of irinotecan during HPLC analysis (Figure 16). The heat map in Figure 3E provides a quantitative display of treatment impact on tumor metastasis.

**LB-MSNPs but Not Liposomes Improve the Safety of Irinotecan Delivery in the
30 Mouse Orthotopic Model.**

[0390] Since irinotecan contributes in a major way to FOLFIRINOX toxicity, the possibility of reducing drug toxicity by encapsulated delivery is a major study objective.

MTD assessment showed that both carriers are capable of increasing the acute lethal dose ~5-fold over free drug (Figure 3A). In order to address irinotecan toxicity through encapsulated drug delivery (Figure 3B), liver, sternal bone marrow, and intestinal tissues were collected from the sacrificed animals in the efficacy study. These are the major target organs for irinotecan toxicity in PDAC patients (Conroy *et al.* (2011) *N. Engl. J. Med.* 364(19): 1817-1825; Ueno *et al.* (2007) *Cancer Chemother. Pharmacol.* 59(4): 447-454; Loupakakis *et al.* (2013) *Br. J. Cancer*, 108(12): 2549-2556). Histological examination of the liver demonstrated severe and extensive hepatocyte necrosis in animals treated with the free drug (Figure 4A). This is reflected by the presence of pyknotic and fragmented nuclei in hepatic cells (Ziegler *et al.* (2004) *News Physiol. Sci.* 19: 124-128). While the severity of liver damage was somewhat reduced by Ir-liposomes, extensive hepatic necrosis was observed (Figure 4A). In contrast, Ir-LB-MSN-treated animals only exhibited minor steatosis, a reversible drug-induced stress response, without necrosis (Figure 4A) (King and Perry (2001) *Oncologist*, 6: 162-176; Maor and Malnick (2013) *Int. J. Hepatol. Art.* 815105). The histological data were also confirmed by IHC analysis, in which liver tissue was used to stain Kupffer cells (KC) (with a FITC-labeled anti-F4/80 antibody), apoptotic cells detected with a RITC- labeled antibody (recognizing cleaved caspase-3) and cellular nuclei with Hoechst 33342 (a blue dye) (Figure 4B). Examination of the stained sections under a fluorescence microscope demonstrated extensive apoptosis throughout the liver of animals treated with the free drug. Animals receiving treatment with Ir-liposomes demonstrated less apoptosis, which frequently involved KC or adjacent hepatic tissue. In contrast, no apoptosis was seen in the livers of animals treated with Ir- LB-MSN.

[0391] While H&E staining of the GI tract did not reveal microscopic evidence of toxicity in any of the treatment groups, IHC staining for cleaved caspase-3 demonstrated the presence of apoptotic cells and blunting of the intestinal villi by the free drug (Figure 4C). While treatment with Ir-liposomes protected against blunting of the villi, IHC staining revealed the presence of apoptosis in apical enterocytes. This is in clear contrast to treatment with Ir-LB-MSNs, which did not result in any apoptosis or damage to the villi (Figure 4C).

[0392] Bone marrow toxicity was studied in a separate experiment, in which animals received three injections of a 40 mg/kg irinotecan dose equivalent, as described above. After sacrifice, sternal bone marrow was collected to evaluate myelotoxicity by H&E staining (Figure 4D) (Iusuf *et al.* (2014) *Mol. Cancer Ther.* 13: 492-503). While

animals treated with free drug or Ir-liposomes showed extensive marrow damage, evidenced by <30% of the marrow space being filled by hematopoietic cells (Travlos (2006) *Toxicol. Pathol.* 34: 566-598), there was no change in marrow cellularity in Ir-LB-MSNP-treated animals (Figure 4D). Moreover, damage to the bone marrow by the free drug or Ir-liposomes was accompanied by peripheral blood neutropenia, while Ir-LB-MSNP treatment showed a nonsignificant decline in neutrophil count (Figure 19). All considered, these data demonstrate a significant reduction in systemic and target organ toxicity during treatment with Ir-LB-MSNP, while the liposomal carrier still exhibits substantial organ toxicity.

Discussion

10 [0393] We developed a custom-designed mesoporous silica nanoparticle platform, using a supported LB and a proton-generating entrapping agent, for high dose irinotecan loading and delivery and tested it in a KPC-derived pancreatic cancer model in immunocompetent mice. The improved stability and loading capacity of the carrier improved the biodistribution, circulatory half-life, and drug tumor content of irinotecan compared to an in-house synthesized MM-398 liposome equivalent, using the same loading procedure and trapping agent (TEA₈SOS). Not only was Ir-LB-MSNP more effective at inducing tumor killing at the primary tumor site, but it was also more active for treating metastases. Equally important, the LB-MSNP carrier did not induce pronounced systemic toxicity, which is in stark contrast to liposomes that had an adverse impact on the bone marrow, GI tract, and the liver. We ascribe the toxicity reduction with maintenance of treatment efficacy to the increased carrier ability of LB-MSNPs compared to that of liposomes. Thus, the LB-MSNP carrier provides an innovative approach for the introduction of irinotecan to the treatment of PDAC, with the possibility to promote the use of this drug, as a first-line consideration rather than being reserved for patients with failure of GEM treatment, as is currently the case for the recent FDA-improved liposomal carrier.

25 [0394] We have developed a robust MSNP carrier for irinotecan, with morphological resemblance to a liposomal equivalent. However, in addition to the superior drug encapsulation provided by an intact supported LB, the LB-MSNPs also allow dense drug packaging against the porous silica sidewalls as a result of bonding and/or electrostatic interactions (Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801; Meng *et al.* (2010) *ACS Nano*, 4: 4539-4550; Ashley *et al.* (2011) *Nat. Mater.*, 10: 389-397). This allowed us to achieve improved irinotecan loading capacity, which provides superior killing efficiency at the orthotopic tumor site, compared to liposomes. The improved loading capacity is further

assisted by the supported LB stability, which improves cargo protection compared to the liposomal bilayer during blood circulation and biodistribution to the tumor site. We propose that the decreased fluctuation of the supported LB, presumably as a result of the robust electrostatic and van der Waals interactions with the particle surface, reduces the lipid loss due to serum protein interactions with a nonsupported lipid bilayer (liposome). A supported LB may also protect the carrier against complement-mediated lysis and the shear forces generated during blood flow (Liu *et al.* (2000) In: *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 172(1-3): 57-67; Heurtault *et al.* (2003) *Biomaterials*, 24(23): 4283-4300; Ashley *et al.* (2011) *Nat. Mater.*, 10: 389-397; Liang *et al.* (2004) *J. Colloid Interface Sci.* 278: 53-62; Anderson *et al.* (2009) *Langmuir*, 25: 6997-7005; Michel *et al.* (2012) *Int. J. Mol. Sci.* 13: 11610-11642; Wang *et al.* (2014) *Small* 10: 3927-3931). For instance, doxorubicin-loaded DOPC liposomes exhibit ~40% drug leakage within 24 h in a simulated serum-containing biofluid environment at 37°C (Ashley *et al.* (2011) *Nat. Mater.*, 10: 389-397). This is in accordance with our finding that, while effective as a nanocarrier, Ir-liposomes are significantly less stable than Ir-LB-MSNPs in whole serum or following particle lyophilization and resuspension (Figures 1D and 1E). Premature release of the highly toxic topoisomerase inhibitor from Ir-liposomes was also accompanied by a high rate of toxicity in the bone marrow, liver, and intestines in animal studies, which is quite different from the protective effect of Ir-LB-MSNPs. Thus, despite the morphological resemblance to liposomes, the LB-MSNPs carrier provides clear advantages over liposomal irinotecan delivery, from both efficacy and drug toxicity perspectives.

[0395] The bone marrow and the intestinal toxicity reduction by Ir- LB-MSNPs is noteworthy from the perspective that the recently FDA-approved MM-398 liposomal formulation, Onivyde, is marketed with a black box warning of possible severe neutropenia and diarrhea during treatment of PDAC patients (*see, e.g.*, www.fda.gov/newsevents/newsroom/pressannouncements/ucm468654.htm; www.accessdata.fda.gov/drugsatfda_docs/label/2015/207793LB.pdf). The warning states that fatal neutropenic sepsis was observed in 0.8% of patients, while severe or life-threatening neutropenic fever occurred in 3% (www.accessdata.fda.gov/drugsatfda_docs/label/2015/207793LB.pdf). Life-threatening neutropenia was observed in 20% of patients receiving Onivyde in combination with 5-fluorouracil and leucovorin (*Id.*). The same drug combination also caused severe diarrhea in

13% of treated subjects (*Id.*). Although we did not use Onivyde for comparison, an in-house liposomal formulation using TEA₈SOS for irinotecan loading was associated with significant bone marrow and intestinal toxicity, similar to the free drug. Similar effects were not seen during treatment with the Ir-LB- MSNP platform.

5 [0396] It is interesting that both the MSNP and liposomal carriers are sequestered by KCs in the liver, as demonstrated by confocal microscopy using NIR-labeled particles and FITC-labeled anti-F4/80 antibodies (not shown). While it is difficult to provide a direct *in vivo* demonstration, we propose that the disintegration of the LB and irinotecan release commence in acidic endo/lysosomal compartments in the KC (Hwang *et al.* (1980) *Proc. Natl. Acad. Sci. USA*, 77: 4030-4034; Derksen *et al.* (1988) *Biochim. Biophys. Acta, Bioenerg.* 971: 127-136; Kolter and Sandhoff (2010) *FEBS Lett.* 584: 1700-1712). The more rapid disintegration of the liposomes could explain why only sparse NIR-labeled particles are seen in animals injected with Ir-liposomes compared to the presence of abundant Ir-LB-MSNP (Figure 2B). The rate of carrier disintegration could, in turn, determine the rate of drug release to bystander hepatocytes. This raises the possibility that irinotecan burst release from the liposomal carrier could overwhelm the hepatic metabolism of the drug, which is carried out by CYP 3A and uridine diphosphate-glucuronosyltransferase (UGT1A1) (Mathijssen *et al.* (2001) *Clin. Cancer Res.* 7: 2182-2194). Decreased metabolism could lead to higher retention of nonmetabolized drug, which leads to hepatocyte injury and necrosis, as shown in Figure 4A. This could explain the significantly higher content of nonconjugated irinotecan in the liver tissue of animals treated with Ir-liposomes compared to Ir-LB-MSNP, that is, 26 vs 15%ID/g at 24 h post-IV injection (Figure 16). These findings are also compatible with the IHC staining of KC and activated caspase-3, which demonstrates a high cell death rate in KC and adjacent hepatocytes during treatment with Ir-liposomes, compared to the absence of apoptosis in Ir-LB-MSNP-treated animals (Figure 4B). The schematic in Figure 4E outlines our hypothesis regarding the role of carrier stability and the rate of irinotecan release from KC in determining the extent of hepatocyte apoptosis and necrosis (Zhao *et al.* (2015) *In vivo. Sci. Bull.*, 60: 3-20). It was not possible to measure hepatic enzymes, due to the limited amount of blood that could be obtained from the animals in their moribund state.

[0397] GEM frequently serves as first-line therapy for PDAC, with a survival outcome of 6.8 months (Burris *et al.* (1997) *J. Clin. Oncol.*, 15: 2403-2413; Teague *et al.* (2014) *Ther. Adv. Med. Oncol.* 7: 68-84). While FOLFIRINOX can improve the survival to

~11 months, the frequent occurrence of serious drug toxicity (mostly due to irinotecan and oxaliplatin) precludes its use as first-line therapy. Thus, FOLFIRINOX is often reserved for patients with good performance status. Unfortunately, this position has not changed with the introduction of Onivyde, which is not approved as first-line therapy and includes black box warnings of severe diarrhea and neutropenia. Since approximately 80% of PDAC patients present at an advanced stage of disease, it would be advantageous to use drug in the FOLFIRINOX regimen as first-line treatment. Although we have not used Onivyde for comparative analysis, the Ir-LB-MSN platform was more efficacious and biocompatible with lower systemic toxicity than our equivalent in-house Ir-liposomal formulation. We propose, therefore, that the Ir-LB-MSN platform could be developed for first-line therapy in PDAC. We have recently demonstrated the feasibility of a dual delivery MSNP carrier for GEM and PTX in a PDAC model (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Each treatment approach should carefully consider the design complexity, cost, and impact on good manufacturing practice before implementation (Sugahara *et al.* (2010) *Science*, 328: 1031-1035).

[0398] Although the lipid-coated MSNPs have been shown to be efficacious for cargo delivery *in vitro* and/or *in vivo* (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557; Zhang *et al.* (2014) *Biomaterials*, 35: 3650-3665; Ashley *et al.* (2011) *Nat. Mater.*, 10: 389-397; Ashley *et al.* (2012) *ACS Nano*, 6: 2174-2188; Dengler *et al.* (2013) *J. Controlled Release*, 168: 209-224), this is the first demonstration of how a protonating agent can be used to increase the loading efficiency of this carrier platform. We have also identified a comprehensive list of weak basic drugs that can be loaded into LB-MSNs through a proton gradient. The general characteristics of these cargo molecules include the following chemical properties:

(i) organic molecular compounds that include primary, secondary, tertiary, or quaternary amine(s);

(ii) a $pK_a < 11$ to allow protonation and entrapment behind the LB (Zucker *et al.* (2009) *J. Control. Release*, 139(1): 73-80; Cern Cern *et al.* (2012) *J. Control. Release*, 160(2): 147-157; Xu *et al.* (2014) *Pharmaceut. Res.* 31(10): 2583-2592);

(iii) a water solubility ranging from around 5 to 25 mg/mL and amphipathic characteristics that allow diffusion across the LB;

(iv) an octanol/water partition coefficient or log P value of -3.0 to 3.0; 71,72
(v) suitable molecular weight with a geometric size less than MSNP pore size (2-8 nm) to

allow entry into the MSNP pores (Li *et al.* (2012) *Chem. Soc. Rev.* 41: 2590-2605; Tang *et al.* (2012) *Adv. Mat.* 24(12): 1504-1534; Tarn *et al.* (2013) *Acc. Chem. Res.* 46(3): 792-801).

[0399] Without being all-inclusive, the list of potential chemotherapy agents include the topoisomerase I inhibitor, topotecan; the antitumor anthracycline antibiotics, doxorubicin and mitoxantrone; the mitotic inhibitors, vinblastine and vinorelbine; the tyrosine kinase inhibitors, imatinib, osimertinib, and sunitinib, *etc.* The ability to package and deliver one or a combination of the above agents will enhance the wider utility of our multifunctional LB-MSNP platform, including treatment consideration of additional cancer types such as colon, breast, lung, liver, glioma, melanoma, *etc.* It is also possible to co-package drug combinations in the above list into a single carrier. For example, based on the success that we have achieved with our GEM/PTX co-delivery platform (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557), it is possible to consider combining drugs in the FOLFIRINOX regimen (*e.g.*, oxaliplatin with irinotecan) for synergistic and ratiometric delivery. Moreover, drug loading by our LB-MSNP can be used for noncancerous applications, such as encapsulating antibiotics for infectious diseases, for example, ciprofloxacin, levofloxacin, or HIV anti-retrovirals (*e.g.*, tenofovir disoproxil fumarate). It is also worth pointing out that, in addition to TEA₈SOS, transmembrane pH gradients can also be generated by acidic buffers (*e.g.*, citrate) (Chou *et al.* (2003) *J. Biosci. Bioeng.*, 95(4): 405-408; Nichols *et al.* (1976) *Biochim. Biophys. Acta, Biomembr.* 455: 269-271), proton-generating dissociable salts (*e.g.*, (NH₄)₂SO₄) (Haran *et al.* (1993) *Biochim. Biophys. Acta, Biomembr.* 1151: 201-215; Maurer-Spurej *et al.* (1999) *Biochim. Biophys. Acta, Biomembr.* 1416: 1-10; Fritze *et al.* (2006) *Biochim. Biophys. Acta, Biomembr.* 1758: 1633-1640), or ionophore-mediated ion gradients from metal salts (*e.g.*, A23187 and MnSO₄) (Messerer *et al.* (2004) *Clin. Cancer Res.* 10(19): 6638-6649; Ramsay *et al.* (2008) *Eur. J. Pharm. Biopharm.* 68(3): 607-617; Fenske *et al.* (1998) *Biochim. Biophys. Acta, Biomembr.* 1414: 188-204). Moreover, it is OK possible to generate reverse pH gradients for drug loading, such as use a calcium acetate gradient, to improve amphiphilic weak acid loading in LB-MSNP, a strategy that has been utilized in liposomes (Avnir *et al.* (2008) *Arthritis Rheum.* 58: 119-129).

Conclusions

[0400] In summary, we have developed a MSNP delivery platform for irinotecan, which, in spite of its morphological resemblance to a liposome, provides biocompatibility

and therapeutic efficacy superior to an equivalent liposomal formulation or free drug. The new carrier could be used to advance irinotecan delivery and drug combinations in the FOLFIRINOX regimen to first- line treatment status for PDAC.

METHODS

5 **Materials.**

[0401] Tetraethylorthosilicate (TEOS), triethanolamine, cetyltrimethylammonium chloride solution (CTAC, 25 wt % in water), triethylamine (TEA), Dowex 50WX8 resin, Sepharose CL-4B, and Sephadex G-25 were purchased from Sigma-Aldrich, USA. 1,2-Distearoyl-*sn*-glycero-3-phosphocholine (DSPC), 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[methoxy(polyethylene glycol)-2000] (ammonium salt) (DSPE-PEG2000), and cholesterol (Chol) were purchased from Avanti Polar Lipids, USA. Sucrose octasulfate (SOS) sodium salt was purchased from Toronto Research Chemicals, Inc., Canada. Irinotecan hydrochloride trihydrate was purchased from LC Laboratories, USA. Penicillin, streptomycin, and Dulbecco's modified Eagle medium (DMEM) were obtained from Invitrogen. Fetal bovine serum (FBS) was purchased from Gemini Bio Products. A rabbit mAb antibody (catalog #9664) that detects activated (cleaved) caspase-3 was purchased from Cell Signaling. Matrigel Matrix Basement Membrane was purchased from BD Bioscience. Centrifugal filter units (cutoff size: 10K and 30K) were purchased from EMD Millipore. All chemicals were directly used without further purification.

20 **Preparation of Irinotecan-Loaded LB-MSNP Using a Trapping Reagent.**

Synthesis of MSNP.

[0402] MSNP cores were synthesized by slight modification of our previous sol/gel procedure (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). To synthesize a batch of ~500 mg of MSNP, 50 mL of CTAC was mixed with 150 mL of H₂O in a 500 mL conical flask, followed by stirring at 350 rpm for 15 min at 85°C. This was followed by the addition of 8 mL of 10% triethanolamine for 30 min at the same temperature. Then, 7.5 mL of the silica precursor, TEOS, was added dropwise at a rate of 1 mL/min using a peristaltic pump. The solution was stirred at 350 rpm at 85°C for 20 min, leading to the formation particles with a primary size of ~65 nm. The surfactant was removed by washing the particles with a mixture of methanol/HCl (500:19 v/v) at room temperature for 24 h. The particles were centrifuged at 10 000 rpm for 60 min and washed three times in methanol.

Synthesis of Trapping Agent TEA8SOS.

[0403] TEA₈SOS was synthesized from a commercially available SOS sodium salt by ion-exchange chromatography (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). Briefly, 500 mg of the salt was dissolved in 1 mL of deionized (DI) water, yielding
5 an aqueous solution of 500 mg/mL. The solution was passed through an 8 cm × 1.5 cm cationic ion- exchange resin (Dowex 50WX8) column, prepared in DI water. The SOS salt was added to the column and eluted with the water for conversion to sucrose octasulfonic (SOS) acid. The acidic SOS eluate was titrated immediately with neat triethylamine to reach a final pH of 5.8. This resulted in the formation of the TEA₈SOS. The salt
10 concentration was adjusted to ~80 mM and stored at 4°C in a refrigerator before use.

TEA8SOS Loading and Entrapment by a LB in MSNP.

[0404] We used a novel method for entrapment of TEA₈SOS in MSNP through the use of LB coating (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). The synthesis commenced by soaking 100 mg of empty MSNPs in 4 mL of an aqueous 80 mM TEA₈SOS
15 by probe sonication for 5 min, with a 15/15 s on/off working cycle and a power output of 32.5 W. The particle suspension was added immediately to a coated lipid biofilm, obtained by adding a mixture of 110 mg of DSPC/Chol/DSPE-PEG2000 (molar ratio 3:2:0.15), suspended in chloroform at 10 mg/mL, to the bottom of a 6 cm round-bottom flask. This is equivalent to a particle:lipid ratio of 1.0:1.1. Following solvent evaporation in a rotary
20 evaporator at room temperature, we obtained a biofilm with ~30 cm² surface area. Following the addition of the particle suspension to the biofilm, probe sonication was used for 30 min with a 15/15 s on/off working cycle at a power output of 32.5 W. Unentrapped TEA₈SOS was removed by size-exclusion chromatography over a Sepharose CL-4B column (1.5 × 15 cm), using a HEPES-buffered dextrose solution (5 mM HEPES, 5%
25 dextrose, pH 6.5) for elusion.

Use of TEA₈SOS-Loaded LB-MSNP for Irinotecan Loading.

[0405] Irinotecan was dissolved in HEPES-buffered dextrose (5 mM HEPES, 5% dextrose, pH 6.5) at 10 mg/mL, which was mixed with the TEA₈SOS-loaded particles to achieve an irinotecan/MSNP ratio of 1:1 (w/w). The mixture was incubated at 65°C in a
30 water bath for 30 min and then quenched in ice water for 10 min. The drug-loaded particles were washed and purified three times by centrifugation at 15 000 rpm for 10 min. The washed particles were resuspended in water, saline, or PBS as indicated.

Preparation of a Comparative Liposomal Formulation for Irinotecan Delivery.**Irinotecan Encapsulation by LB-MSNP in the Absence of a Trapping Agent.**

[0406] Empty MSNPs (40 mg) were added to 4 mL of a 10 mg/mL irinotecan solution (in water) with sonication, followed by stirring for 4 h at room temperature. The drug was then encapsulated by a LB, using sonication of lipid biofilm of similar composition as described above. After probe sonication, the particles were purified by centrifugation and washing and resuspended as described above.

Liposomal Encapsulation of Irinotecan Using TEA₈SOS.

[0407] In order to develop an in-house analogue of the MM-398 formulation (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277; Ko *et al.* (2013) *Br. J. Cancer*, 109: 920-925), 212 mg of a lipid mixture containing DSPC/Chol/DSPE-PEG2000 (at a molar ratio of 3:2:0.015) was dissolved in 0.4 mL of ethanol at 65°C. Four milliliters of a TEA₈SOS solution (80 mM) was mixed with the lipid ethanol solution at the same temperature, followed by sonication in a water bath for 2 min. The lipid suspension was extruded 15 times through a polycarbonate membrane with a 0.1 µm pore size at 65°C. Unentrapped TEA₈SOS was removed by a size-exclusion chromatography column (Sephacrose CL-4B), eluted with a HEPES-buffered dextrose solution (5 mM HEPES, 5% dextrose, pH 6.5). An irinotecan solution, prepared at 10 mg/mL in HEPES-buffered dextrose (5 mM HEPES, 5% dextrose, pH 6.5), was mixed with TEA₈SOS-loaded liposomal suspension (irinotecan/lipids = 1:2.1 w/w) and incubated in a water bath at 65°C for 30 min. The sample was immediately quenched in ice water for 10 min. The free drug was removed using a Sephadex G-25 column eluted with HEPES-buffered saline (5 mM HEPES, 145 mM NaCl, pH 6.5).

Synthesis of an Irinotecan Liposomal Carrier in the Absence of a Trapping Agent.

[0408] We also synthesized an irinotecan-delivering liposome that uses a passive encapsulation approach, instead of an entrapment agent. Briefly, a mixture of lipids (53 mg lipids, DSPC/ Chol/DSPE-PEG2000 at molar ratio of 3:2:0.015) was dissolved in 0.1 mL of ethanol at 65°C. One milliliter of an irinotecan solution (10 mg/mL in water) was mixed with the lipid ethanol solution at 65°C, followed by sonication for 2 min. The lipid suspension was extruded 15 times through a polycarbonate membrane with a 0.1 µm pore

size at 65°C. The drug-laden liposome was purified using the same method as described above.

Physicochemical Characterization.

[0409] All particles used in this example were extensively characterized for morphology, size, size distribution, shape, and surface charge. The uniformity of LB coating and its intactness were characterized by TEM (JEOL 1200-EX) and cryoEM (TF20 FEI Tecnai-G2). Particle size and ζ -potential were measured by a ZETAPALS instrument (Brookhaven Instruments Corporation). These measurements were performed with the nanoparticles suspended in water, PBS, and cell culture media plus 10% FBS at a particle concentration of 100 $\mu\text{g/mL}$.

[0410] The drug loading capacity of each carrier was determined by a subtraction method, as described previously (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Briefly, the amount of non-encapsulated irinotecan detected during purification (m_1) was determined by OD (360 nm) in a microplate reader (M5e, Molecular Device, USA) or through the use of HPLC (Raytest, Germany). Drug loading capacity was defined as $\text{DLC} = [\text{the total amount of irinotecan } (m_0) - \text{non-encapsulated irinotecan } (m_1)] / [\text{the total amount of particle } (m_{\text{MSNP or mlipid}})] \times 100\%$. The drug retention stability was tested in 100% FBS at 37°C for 24 h. Briefly, 40 μL of drug-loaded NPs (irinotecan: 10 mg/mL) was added to 1 mL of serum at 37°C, with continuous shaking. The mixture was centrifuged at set time intervals, using a centrifugal filter separation device (MW cutoff of 10K). The released irinotecan was quantified in a microplate reader or by HPLC. A serum filtrate without NPs was used as a control. The stability of both carriers was also tested through a lyophilization procedure. Briefly, Ir-MSNPs were suspended in 5% dextrose. The sample was lyophilized at -60 °C overnight and stored in a -80°C freezer. The stored samples were resuspended in water by gentle vortexing, following which the suspended solutions were characterized for size, ζ -potential, and drug release as described previously.

Cell Culture.

[0411] An immortalized cell line derived from a spontaneously developed tumor in a transgenic Kras^{LSL-G12D/+}, Trp53^{LSL-R172H/+}, Pdx1-Cre mouse was kindly provided by Dr. Andrew Lowy at UC San Diego. PANC-1 cells were purchased from American Type Culture Collection (ATCC). Both cell types were cultured in DMEM, containing 10% FBS, 100 U/mL penicillin, 100 $\mu\text{g/mL}$ streptomycin, and 2 mM L-glutamine. To allow bio-

luminescence tumor imaging in an IVIS system, both cells were permanently transfected with a luciferase-based lentiviral vector in the UCLA vector core facility, as previously described (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). Following a limiting dilution protocol to select single cell clones, the KPC-luc and PANC-1-luc cell populations were used to develop PDAC models.

Assessment of Maximum Tolerated Dose.

[0412] The MTDs of free and encapsulated irinotecan formulations were determined using a protocol from the Toxicology and Pharmacology Branch (dtp.nci.nih.gov/branches/tpb/default.htm) at the National Cancer Institute (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). Two healthy male BALB/c mice were IV injected with a 50 mg/kg (C1 dose) equivalent of free or encapsulated irinotecan. This was followed by a dose escalation factor of 1.8 to obtain the Cn dose, which results in animal death within 24 h after the last administration. The second round of dose-seeking began with the Cn-1 dose and utilized a 1.15 escalation factor (n = 2) to find the MTD, which is defined as the absence of mortality or morbidity. The MTD was validated by injecting six mice, which were followed for 15 days to determine the absence of morbidity, mortality, or >15% weight loss.

Establishment of a KPC-Derived Orthotopic Tumor Model in Immunocompetent Mice.

[0413] Female B6/129 mice (~8 weeks) were purchased from The Jackson Laboratory and maintained under pathogen-free conditions. All animal experiments were performed with protocols approved by the UCLA Animal Research Committee. To grow orthotopic xenografts, the mice were anesthetized with isoflurane, followed by IP injection of 50 mg/kg ketamine and 10 mg/kg xylazine. The surgical site was shaved to leave a margin of 1 cm around the incision site and sterilized by scrubbing with betadine and 70% ethanol. The mice were positioned for surgery on a heating pad, and the incision site in the left flank was draped with sterile gauze. A surgical incision of 0.5-0.7 cm was made to expose the pancreatic injection site, followed by an injection of 50 μ L of DMEM/Matrigel (1:1 v/v) containing 2×10^6 KPC-luc cells into the pancreatic tail by a 27 gauge needle. The fascial layers were closed with absorbable sutures (PDS II, Ethicon) and the skin with nonabsorbable sutures (PROLENE, Ethicon). The mice were kept on the warming pads until full recovery from the anesthesia and then transferred to clean cages. The efficacy study was performed in tumor-bearing mice approximately 2 weeks after implantation, at

which point the primary tumors had grown to ~0.5 cm, without evidence of macroscopic tumor metastasis. For the biodistribution experiments, the tumor-bearing mice were used 3 weeks after tumor implantation, at which point the primary tumors had grown to a size of ~0.8 cm.

5 **Systemic and Intratumoral Biodistribution of IV Injected NIR-Labeled Nanoparticles.**

[0414] An IVIS (Xenogen) *in vivo* imaging system was used to study the biodistribution of NIR-labeled MSNPs and liposomes in the KPC-derived orthotopic model (n = 3 mice). NIR labeling was performed by incorporating 0.1% w/w Dylight 680-DMPE
10 in the LB of both the liposomal and MSNP carriers without drug (Meng *et al.* (2015) *ACS Nano*, 9(4): 3540-3557). For bioluminescence imaging of the tumor site, mice were injected IP with 75 mg/kg D-Luciferin. Reference fluorescence images for the tumor-bearing mice were captured before particle injection. NIR images were obtained over 48 h in animals IV injected with 100 mg/kg NIR-labeled particles. Following animal sacrifice,
15 ex vivo images were obtained of excised tumors and major organs to quantitatively assess particle biodistribution. A small section of the tumor tissue was cryo-embedded, using the OCT reagent, and used to prepare tumor sections. The intratumoral distribution of the NIR-labeled particles was studied by confocal microscopy (SP2-1P-FCS, Leica) using a NIR laser. Nuclear staining of the same sections was performed by using Hoechst 33342 dye
20 (blue). In animals treated with LB-MSNP, the tumor tissue and major organs were also used to assess the Si content by ICP-OES. Particle biodistribution was expressed as the % total injected dose (%ID) that could be shown to distribute to individual organs. In order to measure the irinotecan content in the tumor or other normal tissues, animals were injected with 60 mg/kg free or encapsulated irinotecan (particle dose, 120 mg/kg for MSNPs and
25 150 mg/kg for liposomes), followed by sacrifice after 24 h. HPLC analysis was performed on harvested tumor tissue, liver, spleen, kidney, intestines, and blood to determine irinotecan content.

HPLC Analysis.

[0415] The harvested tumor and organ samples were weighed and homogenized on
30 ice. Following the extraction of 0.1 mL of plasma or tissue homogenate with 0.4 mL of an acidic solution (0.1 mol/L phosphoric acid/methanol, 1:4 v/v), the extracts were vortexed twice for 10 s and centrifuged at 13 000 rpm for 10 min. The irinotecan-containing

supernatants were filtrated through 0.22 µm filters for HPLC analysis in a system containing a Knauer Smartline pneumatic pump, C18 column, K-2600 spectrophotometer, and Gina data acquisition software. The mobile phase, delivered at a flow rate of 1.00 mL/min, comprised a 3% triethylammonium acetate aqueous buffer (pH 5.5) and acetonitrile (73:27 v/v) (Noble *et al.* (2006) *Cancer Res.*, 66(5): 2801-2806). Twenty microliters of irinotecan-containing sample was injected to measure the drug absorption at 254 nm, typically eluted in 4.4 min. An irinotecan standard curve was generated over the concentration range of 0.05-100 µg/mL. Irinotecan content was expressed as % of injected dose per gram of tissue (%ID/g).

10 **Efficacy Assessment Using the KPC-Derived Orthotopic Model.**

[0416] Tumor-bearing B6/129 mice were randomly assigned to four groups, including eight animals per group. The mice were IV injected with free or encapsulated irinotecan to receive a dose of 40 mg/kg (*i.e.*, a liposomal dose of 100 or 80 mg/kg MSNP) every 4 days. A maximum of eight injections were administered over a 28 day observation period. Saline injections served as a negative control. Tumor burden was monitored by IVIS imaging every 8 days and quantitatively expressed as bioluminescence imaging intensity in the operator-defined ROI. The statistical analysis of the differences between the groups was performed using a t-test (Excel software, Microsoft). All surviving mice were sacrificed at day 47, at which point all animals in the saline and drug-free group had died. The tumor tissue and major organs (GI tract, liver, spleen, heart, lung, and kidneys) were harvested for quantification of ex vivo bioluminescence image intensity.

Comparison of the Toxicity Potential of Different Irinotecan Formulations.

[0417] Major tissues collected during the efficacy testing experiment (liver, kidney, spleen, stomach, and intestines) were fixed in 10% formalin, followed by paraffin embedding. Tissue sections were stained by H&E for histological analysis and also used for IHC analysis of the expression of activated (cleaved) caspase-3. The slides were read in a blinded fashion by an experienced veterinary pathologist. The second approach for assessing bone marrow toxicity used three IV administrations, 2 days apart, of free drug and carrier formulations at 40 mg/kg in healthy mice. The animals were sacrificed on day 7, and the sternum was used for fixation in 10% formalin, decalcification, paraffin embedding, and H&E staining. We also collected blood for differential white cell counting by the UCLA Division of Laboratory Animal Medicine (DLAM) diagnostic laboratory services. To

understand the mechanism of differential liver toxicity by free drug and the carriers, mice receiving a single injection of the dose equivalent of 60 mg/kg irinotecan were sacrificed after 24 h. The liver tissues were cryo-embedded for immunofluorescence staining of KCs and the expression of activated caspase-3, using a FITC-labeled anti-F4/80 antibody or a RITC-conjugated secondary antibody to identify KC and cleaved caspase-3, respectively. Hoechst 33342 dye was used to localize the cellular nuclei in the same sections. The stained slides were examined under a confocal microscope (Observer D1, Zeiss).

Statistical Analysis.

[0418] Comparative analysis of the differences between groups was performed using the two-sided Student's t-test (Excel software, Microsoft). A statistically significant difference was determined at $p < 0.05$. Values were expressed as mean \pm SD or SEM of multiple determinations, as stated in the figure legends.

Example 2

[0419] Pancreatic ductal adenocarcinoma (PDAC) is an incurable disease with a 5-year survival rate less than 5%. Currently, first line chemotherapy for PDAC includes treatment by gemcitabine (GEM) (single agent therapy) or use of a four-drug regimen, which includes leucovorin, 5-FU, oxaliplatin, and irinotecan, a.k.a. FOLFIRINOX. While FOLFIRINOX has an increased PDAC response rate compared to GEM, *i.e.* 31.6% *versus* 9.4%, this regimen is far more toxic and therefore restricted to a minority of advanced stage PDAC patients with good functional status (Conroy *et al.* (2011) *N. Engl. J. Med.* 364: 1817-1825). Irinotecan contributes in a major way to FOLFIRINOX toxicity due to adverse effects on the bone marrow (*e.g.*, ~60% incidence of neutropenia), as well as the G.I.T. (*e.g.*, vomiting, diarrhea, nausea, anorexia in ~50% of patients) (Ueno *et al.* (2007) *Cancer Chemother. Pharmacol.* 59(4): 447-454). There is an unmet need to develop a FOLFIRINOX regimen that is less toxic, including the reducing the toxicity of irinotecan.

[0420] Use of nanoparticle carriers to deliver irinotecan provides a potential solution to reducing the toxicity of this drug, as illustrated by the delivery of irinotecan by liposomal (Messerer *et al.* (2004) *Clin. Cancer Res.* 10(19): 6638-6649; Drummond *et al.* (2006) *Cancer Res.* 66(6): 3271-3277) or polymer-based nanocarriers (Onishi *et al.* (2003) *Biol. Pharmaceut. Bull.* 26(1): 116-119). At pre-clinical level, the benefits of using these nanocarriers include toxicity reduction, increased antitumor efficacy, and enhanced survival rates of various murine PDAC models. To date, only a few irinotecan carriers have been

tested in clinical trials, including liposomal formulations that promote drug loading by an ionophore, A23187, (Irinophore C) or a protonating entrapment agent, triethylammonium sucrose octasulfate (TEA₈SOS). The therapeutic efficacy of MM-398 is dependent on the encapsulation and trapping of irinotecan in the liposome through the use of a multivalent anionic trapping agent, TEA₈SOS. TEA₈SOS leads to the protonation of irinotecan diffusing across the lipid bilayer, converting it into a hydrophilic component that cannot exit the lipid bilayer. The drug is therefore trapped at concentrations that exceed the passive encapsulation by a liposome bilayer 10-fold (Drummond *et al.* (2006) *Cancer Res.*, 66(6): 3271-3277). MM-398, is currently in Phase 3 trials by Merrimack Pharmaceuticals(www.merrimackpharma.com) (Poster presentation by Hoff *et al.*, Society for Medical Oncology, 2014). This poster can be viewed at [//merrimackpharma.com/sites/default/files/documents/ESMOGI2014MM398.pdf](http://merrimackpharma.com/sites/default/files/documents/ESMOGI2014MM398.pdf)). In the AACR 2012 meeting, the Merrimack company published an abstract, containing a statement that "MM-398 induces tumor regression in multiple mouse models of pancreatic cancer, including an orthotopic metastatic model." The subcutaneous models included BxPC3, AsPC-1, Panc-1 and MiaPaCa. The orthotopic implant in the pancreas was carried out with BxPC3. No detailed data were presented. IV administration of MM-398 has been shown to induce complete tumor regression in various PDAC tumor models in mice, including inhibition on metastatic tumor foci formation (Drummond *et al.* (2006) *Cancer Res.* 15(66): 3271-3277). Treatment of 417 patients in a Phase 3 PDAC trial with a combination of MM-398, 5-FU and leucovorin showed increased overall survival (6.1 months) compared to 1.9 months in people control receiving 5-FU and leucovorin (Hoff *et al.* ESMO Poster, www.merrimackpharma.com).

[0421] In this example, we describe a novel mesoporous silica nanoparticle (MSNP) platform, capable of achieving active loading, entrapment, and encapsulation of irinotecan by a nanocarrier that has superior capabilities compared to the liposomal equivalent. Synthesis of these particles was achieved by the rapid encapsulation of TEA₈SOS in the porous interior of the particles by a supported lipid bilayer LB. Subsequent incubation of these particles in an irinotecan solution allowed active entrapment of irinotecan, which diffuses across the bilayer and is then protonated by TEA₈SOS. Comparative analysis was performed between irinotecan-loaded LB-MSNP *versus* an in-house synthesized liposomal equivalent of MM-398. This analysis demonstrates superior loading capacity, loading efficiency, and release profile by the MSNP carrier. We also showed a similar *in vitro*

effectiveness and *in vivo* toxicity reduction compared to the in house MM-398 liposome. We are currently conducting animal efficacy studies to compare these carriers in subcutaneous and orthotopic human PDAC tumor models.

[0422] We believe the LB-MSNP mediated nano-encapsulation and targeted delivery of irinotecan will improve the clinical use of FOLFIRINOX by decreasing toxicity, including reduced toxicity of irinotecan in the bone marrow and GIT. This discovery permits an IV injectable, efficient, biocompatible, and commercially competitive irinotecan formulation capable of outperforming liposomal carriers because of loading capacity and sufficient cancer site drug release. In certain embodiments this invention could also be employed as an effective design principle for delivering other weak basic molecules ($pK_a > 7.0$) that can be used for cancer treatment as well as treatment of other diseases.

[0423] The following description illustrates certain embodiments of a carrier design for irinotecan loading as well as the comparative analysis of the LB-MSNP carrier with an in-house-synthesized liposomal formulation similar to MM-398.

Irinotecan encapsulation by a LB-MSNP that also includes a trapping agent to achieve additional active loading

[0424] In order to provide effective loading of irinotecan by LB-MSNP, we adapted our recently discovered biofilm procedure for drug loading by first encapsulating the trapping agent, TEA₈SOS, which is then responsible for irinotecan import and retention by protonating the drug diffusing across the supported LB to enter the pores. This leads to stable entrapment of irinotecan in the sealed MSNP pores, creating a gradient for further diffusion across the LB until equilibrium is reached. This actually allows the drug to accumulate at high concentration with the assistance of the large surface area of the porous interior and the trapping agent mediated drug retention. In certain embodiments the synthesis procedure is comprised of four parts, namely, MSNP core synthesis, synthesis of TEA₈SOS loaded LB-MSNP, active loading of irinotecan into the particles through the action of TEA₈SOS, and finally purification of the carrier.

Chemicals:

[0425] Tetraethyl orthosicate (TEOS, $\geq 99.0\%$, Lot #BCBK9540V), triethanolamine ($\geq 99.0\%$, Lot #BCBH9036V), cetyltrimethylammonium chloride solution (CTAC, 25 wt.% in water, Lot #STBC7888V), triethylamine (TEA, $\geq 99.0\%$, Lot #SHBD9006V) were

purchased from Sigma-Aldrich, USA. Distearoylphosphatidylcholine (DSPC, >99.0% Lot #180PC-147), methoxypoly(ethylene)glycol (PEG2000)-derivatized distearoylphosphatidylethanolamine (DSPE-PEG2000, >99.0%, Lot #180PEG2PE-122), and cholesterol (Chol, >98.0%, Lot #CH-94) were purchased from Avanti Polar Lipids, USA. Sucrose octasulfate sodium salt (Lot #1-TMH-175-4) was purchased from Toronto Research Chemicals, Inc, Canada. Irinotecan hydrochloride trihydrate (IRIN, $\geq 99.0\%$, Lot #RCN-104) was purchased LC Laboratories, USA. All chemicals were directly used without further purification.

Sol-gel synthesis of MSNP and template removal

10 [0426] MSNP cores were synthesized by a minor modification of our sol/gel procedure. The reaction was carried out in a 500 mL conical flask. 50 mL CTAC (25%) was mixed with 150 mL H₂O, while stirring at 350 rpm for 15 min at 85 °C. This was followed by the addition of 8 mL 10% triethanolamine at 85°C for 30 min. The silica precursor, TEOS (7.5 mL), was added drop-wise to the mixture at a rate of 1 mL/min
15 controlled by a peristaltic pump. To achieve the synthesis of particles with primary size of ~60 nm, the solution was stirred at 350 rpm at 85°C for 20 min. To remove the surfactant, the particles were washed by menthol and HCl (500:19, v/v) at room temperature for 24 hours. The particles were centrifuged at 10,000 rpm for 60 min and washed 3 times in menthol. In order to obtain high quality MSNP cores, frequent quality checks were
20 performed, including DLS measurements to monitor particle aggregation and contamination, TEM for visualizing particle morphology, and infrared spectroscopy & cytotoxicity assays to check whether the detergent was thoroughly removed.

Preparation of irinotecan loaded LB-MSNP using TEA₈SOS as trapping agent

[0427] In order to establish the optimal procedure for irinotecan loading by LB-
25 MSNP, we developed a novel approach in which we trapped the protonating agent in the particles, using a biofilm procedure (previously disclosed), before incubating the particles in the irinotecan solution. This allows the lipophilic irinotecan to diffuse through the LB membrane with subsequent protonation in the pores, leading to drug entrapment against a concentration gradient (see Figure 5). Considering the loading capacity and drug release
30 behavior, we optimized the irinotecan loading procedure by systemically varying loading time, trapping agent concentration, LB-MSNP concentration, irinotecan concentration, and irinotecan/LB-MSNP ratio. This allowed us to achieve an irinotecan loading capacity of

85% (irinotecan/MSNP, w/w), which exceeds the drug loading capacity in the absence of the trapping reagent ~4-fold. This amounted to twice the loading capacity of in-parallel synthesized liposomal version of MM-398. For further discussion, we have designated the irinotecan-loaded LB-MSNP as “Ir-LB-MSNP(+SOS)”.

5 **Synthesis of TEA₈SOS loaded LB-MSNP**

Preparation of trapping agent triethylammonium sucrose octasulfate (TEA₈SOS) from commercially available sucrose octasulfate sodium salt.

[0428] We used an ion-exchange reaction to synthesize TEA₈SOS from its precursor, the sodium salt of sucrose octasulfate. Briefly, 500 mg of the precursor salt was dissolved in 1 mL D.I. water, yielding an aqueous solution of 500 mg/mL. This solution was passed through a cationic Dowex 50WX8 resin ion-exchange column (15 mm in diameter and 8 cm in length, purchased from Sigma, Lot # MKBH8810V). Following the elution of the column with D.I. water, the sucrose sodium is converted to protonated octasulfate in the process. The collected sucrose octasulfonic acid was immediately titrated with trimethylamine (TEA) to a final pH of 5.5 to 6.0, resulting in the formation of TEA₈SOS. The TEA₈SOS concentration was adjusted to ~80 mM, based on titration of the amount of TEA. The multivalent anionic TEA₈SOS was stored in a 4°C refrigerator before use.

20 **Preparation of TEA₈SOS loaded LB-MSNP.**

[0429] We adapted our previously disclosed biofilm technique for LB coating (PCT/US2014/020857 (UCLA disclosure 2013-534-2), which is incorporated by reference in its entirety) to encapsulate TEA₈SOS, before proceeding to irinotecan loading. Briefly, 100 mg empty MSNPs were soaked in 4 mL TEA₈SOS aqueous solution (40 mM), using probe sonication for 30 minutes with 15/15 seconds on/off working cycle, at a power output of 32.5 W. The suspension was immediately added on top of the coated lipid biofilm, which occupied a ~30 cm² surface area at the bottom of a round-bottom flask (6 cm diameter). Following experimentation to achieve an optimal mix of lipid components (with the aim at stable coating without leaking), we decided on a mixture containing 228 mg DSPC/Chol/DSPE-PEG at molar ratio of 3:2:0.15. This mixture was dissolved in chloroform at 10 mg/mL. Biofilm formation was achieved by solvent evaporation over ~1 h, using a rotary evaporator connected to a vacuum system at room temperature. Following

the addition of the 4 mL particle suspension to the biofilm, probe sonication was used for 30 minutes with 15/15 seconds on/off working cycle, at a power output of 32.5 W. Since the sonicated suspension contains coated particles, liposomes, and free TEA₈SOS, TEA₈SOS LB-MSNP was purified using a chromatography procedure, in which a Sepharose CL-4B column (15 mm in diameter and 15 cm in length, purchased from Sigma, Lot #MKBP0885V) was eluted with a HEPES-buffer (5 mM HEPES, 5% dextrose, pH 6.5).

Use of TEA₈SOS-loaded LB-MSNP for active irinotecan loading.

[0430] Irinotecan was dissolved in a HEPES buffered solution (5 mM HEPES, 5% dextrose, pH 6.5) at concentration of 10 mg/mL. The drug solution was mixed with a TEA₈SOS loaded LB-MSNP suspension at an irinotecan/particle ratio of 1:1 (w/w), and incubated in a water bath at 60 °C for 0.5 h. The reaction was terminated by placing the container in ice water for 10 min.

Particle Purification

[0431] The Ir-LB-MSNP(+SOS) was purified from the free liposomes and free irinotecan by particle centrifugation at 15,000 rpm for 10 minutes, followed by re-suspension in saline or 5% dextrose. The washing procedure was repeated 3 times.

Preparation of control formulations for comparative studies, using the Ir-LB-MSNP (+SOS) particles

[0432] To conduct comparative analysis reflecting the unique properties of the Ir-LB-MSNP(+SOS) particles, we synthesized the LB-MSNP without the trapping agent, allowing the soaked irinotecan to be sealed off by the LB. We also prepared a liposomal equivalent of MM-398, prepared with the trapping agent, as well as a control liposome that encapsulates irinotecan, without the trapping agent.

Synthesis of Irinotecan LB-MSNP not containing the trapping agent:

[0433] Empty MSNPs (40 mg) were placed in 4 mL irinotecan solution (in water, 10 mg/mL) with sonication for 5 min, and then stirred for another 4 h at room temperature. The irinotecan-soaked MSNP suspension was added to the dry lipid membrane (91 mg of a lipid mixture containing DSPC/Cholesterol/PE-PEG2000 at molar ratio of 3:2:0.15), followed by probe sonication for 30 minutes, using a 15/15 seconds on/off working cycle at a power output of 32.5 W. The particles were purified from free liposomes and free drug using centrifugation at 15,000 rpm for 10 minutes, followed by particle washing and re-

suspension as described above. We refer to the particles without the trapping agent as “Ir-LB-MSNP (-SOS)”.

In house synthesis of a liposomal equivalent of the commercially developed liposomal preparation, MM-398.

5 **[0434]** The synthesis was carried out according to the published procedure for MM-398 by Drummond *et al.* (Cancer Res. 2006, 66, 3271-3277). Briefly, a mixture of lipids (212 mg lipid, DSPC/Chol/DSPE-PEG at a molar ratio of 3:2:0.015) was dissolved in 0.4 mL ethanol at 60-65 °C. A 4 mL TEA₈SOS solution (80 mM) was mixed with the lipid ethanol solution at the same temperature, followed by sonication in a water bath for 2
10 minutes. The lipid suspension was extruded 15 times through a polycarbonate membrane with 0.1 µm pore size at 60-65°C. Non-entrapped TEA₈SOS was removed by chromatography over a Sepharose CL-4B column, eluted with a HEPES-buffer (5 mM HEPES, 5% dextrose, pH 6.5). The irinotecan solution (10 mg/mL aqueous solution containing HEPES-buffer, 5 mM HEPES, and 5% dextrose at pH 6.5) was mixed with the
15 TEA₈SOS pre-loaded liposome suspension (irinotecan:lipid = 1:2.1 w/w), and then incubated in a water bath at 65 °C for 0.5 h. The reaction was terminated by placing the mixture in ice water for 10 min. The irinotecan-loaded liposomes were then purified by removing the free irinotecan over a Sephadex G-25 column (15 mm in diameter and 15 cm in length, Sigma, Lot #019K1077), eluted by HEPES-buffered saline (5 mmol/L HEPES,
20 145 mmol/L NaCl, pH 6.5). We will refer to the in-house synthesized liposomal formulation as “Ir-Lipo(+SOS)”.

Synthesis of irinotecan-encapsulated liposomes, without the use of a trapping agent.

25 **[0435]** We also synthesized a liposome that uses a passive encapsulation approach. Briefly, a mixture of lipids (53 mg lipid, DSPC/Chol/DSPE-PEG at molar ratio of 3:2:0.015) was dissolved in 0.1 mL ethanol at 60 °C to 65 °C. 1 mL irinotecan solution (in water, 10 mg/mL) was mixed with the lipid ethanol solution at 60 °C to 65 °C, followed by water bath sonication for 2 minutes. The lipid suspension was extruded 15 times through a polycarbonate membrane with 0.1 µm pore size at 60 °C to 65 °C. The drug-laden
30 liposomes were purified by removing the free irinotecan using a Sephadex G-25 column, eluted with HEPES-buffered saline (5 mmol/L HEPES, 145 mmol/L NaCl, pH 6.5). We refer to this liposomal preparation as “Ir-Lipo(-SOS)”.

Comparative analysis of Ir-LB-MSNP(+SOS) with control irinotecan formulations.**Physicochemical characterization of Ir-LB-MSNP(+SOS) and control formulations.**

[0436] Table 3 shows the hydrodynamic sizes, size distribution and surface charge of Ir-LB-MSNP(+SOS) compared to the control formulations. Ir-LB-MSNP(+SOS) has a hydrodynamic size of 110~130 nm with a polydispersity index (PDI) of ~0.1. This particle showed a negative zeta potential value in pure water, *i.e.* -24.7 mV. The zeta potential changed to -2.95 mV in PBS, and -3.27 mV in complete DMEM culture medium. These physicochemical parameters were similar to the control formulations, such as Ir-Lipo(-SOS), Ir-Lipo(+SOS), and Ir-LB-MSNP(-SOS), in different solutions.

Table 3. Size and zeta potential of Ir-loaded particles in different incubation media.

NP	DLS hydrodynamic size (nm) (PDI)			Zeta Potential		
	Water	PBS	DMEM+10% FBS	Water	PBS	DMEM+10% FBS
Ir-Lipo(-SOS)	101.7 (0.113)	95.5 (0.014)	76.5 (0.275)	-11.7	-4.95	-4.63
Ir-Lipo(+SOS)	102.0 (0.112)	104.7 (0.067)	98.6 (0.172)	-13.4	-2.26	-4.38
IF-LBL-MSNP(-SOS)	129.4 (0.140)	116.1 (0.087)	110.2 (0.17)	-22.4	-5.87	-3.57
Ir-LBL-MSNP(+SOS)	128.0 (0.107)	125.7 (0.112)	121.1 (0.200)	-24.7	-2.95	-3.27
Hydrodynamic size, size distribution and surface charge of different irinotecan-loaded particles. The particles were dispersed in water, PBS and DMEM media supplemented with 10% FBS at a MSNP or lipid concentrations of ~0.1 mg/mL. Size and zeta potential were measured by ZETAPALS particle sizer and zeta potential analyzer (Brookhaven instruments Corp.). The results represent the mean of three measurements						

[0437] Table 4 shows the loading capacity and loading efficiency of the LB-MSNP with and without the use of the trapping agent. The loading capacity = (Total irinotecan – irinotecan in supernatant) / (the mass of MSNP or liposome) x 100%. The loading efficiency is defined as [(Total irinotecan – irinotecan in supernatant) / total irinotecan] x 100%. For liposomes, the loading capacities were 4.12% w/w and 45.2% w/w in the absence or presence of 80 mM trapping reagent. The loading efficiencies of irinotecan in the liposome platform, increased from 8.71% to 95.5% as a result of the action of the

trapping agent. This is in agreement with the published data (Cancer Res. 2006, 66, 3271) for synthesized MM-398 liposomes. Following experimentation to obtain the most stable particle formulation, we used 40 mM of trapping agent to synthesize the LB-MSNP.

Following incubation with irinotecan, we achieved an Ir-LB-MSNP(+SOS) with loading capacity 83.5% w/w, compared to a loading capacity of 22.5% w/w for LB-MSNP without the trapping reagent. While the Ir-LB-MSNP used less trapping agent (40 mM), it actually resulted in 2x loading capacity when compared to the in house MM-398 liposome that contains 80 mM trapping agent.

Table 4. Loading capacity and efficiency of LB-MSNP vs liposomes in the absence and presence of the TEA₈SOS trapping agent

NP	IRIN in feed (g/mol phospholipid)	Loading capacity IRIN/MSNP w/w	Loading capacity IRIN/lipid w/w	Encapsulation efficiency
Ir-Lipo(-SOS)	500	N/A	4.12%	8.71%
Ir-Lipo(+SOS)	500	N/A	45.2%	95.5%
IF-LBL- MSNP(-SOS)	500	22.5%	N/A	22.5%
Ir-LBL- MSNP(+SOS)	500	83.5%	N/A	83.5%

[0438] We also performed cryoEM analysis to visualize the morphology of Ir-LB-MSNP(+SOS). This confirmed the uniform coating of the MSNP surface by a lipid bilayer (Figure 6A, upper panel). High magnification cryoEM images demonstrated primary particle sizes of ~80 nm, with complete surface coating by a bilayer ~7.1 nm thick. The zoom-in image confirmed the presence of an intact lipid coat, while also demonstrating the presence of high density material in the pores; this material represents the irinotecan-TEA₈SOS complex. We observed occasional particles that were encapsulated by a slightly larger liposome that was not tightly adhered to the particle surface, however, the LB layer remains intact. The image of Ir-Lipo(+SOS) showed a liposome structure of ~75 nm with a unilaminar bilayer ~6.5 nm thick (Figure 6A, lower panel). High magnification cryoEM provides sufficient resolution to see the high density IRIN-TEA₈SOS precipitates inside the liposome (red arrows).

[0439] In order to assess the stability of each formulation, we compared the drug stability in PBS (pH=7.4) at 37 °C (Figure 6B), with an intention to estimate the irinotecan stability in the blood stream. This demonstrated that Ir-LB-MSNP(+SOS) is very stable in

PBS, resulting in <5% release over 24 hours. A comparable stability in PBS was found in the in house MM-398 liposome.

[0440] Due to the elevated glucose uptake, glycolysis, lactic acid production (Warburg effect), abnormal blood perfusion, and presence of a dense stroma, the pH value of PDAC is acidic (Wojtkowiak, *et al.* (2012) *Cancer Res.* 72(16): 3938-3947;; Estrella *et al.* (2013) *Cancer Res.* 73(5):1524-1535). Thus, we also looked at the drug release profile in an acidic pH condition. The tumor pH classically drops to ~6.5, but can be as low as ~5.5 (Jähde *et al.* (1992) *Cancer Res.* 52: 6209-6215). The pH could further drop to <5 in lysosomal compartments when the particles were taken up by PDAC cells (Mindell *et al.* (2012) *Annu Rev Physiol.* 74: 69-86). Thus, we performed drug release experiment in the phagolysosomal simulant fluid (PSF, pH=4.5). In PSF, a rapid irinotecan release (*i.e.* 20% over 4 hours) could be seen in LB-MSNP. However, in the in house MM-398 formulation, we found a significantly slower drug release profile in PSF, *i.e.* only ~4% drug release in 24 hours.

Cellular uptake and killing efficiency of Ir-LB-MSNP(+SOS) in cultured PDAC cells

[0441] Confocal microscopy, utilizing Texas red-labeled LB-MSNP or liposomes, demonstrated abundant cellular uptake in a peri-nuclear distribution in PANC-1 cells (Figure 7). We also employed a MTS cytotoxicity assay to determine the effect of Ir-LB-MSNP(+SOS) on PANC-1 and BxPC3 cells during treatment with incremental irinotecan concentrations. We compared the effect of these particles with free drug and Ir-Lipo(+SOS) controls. While the free drug is more effective in killing over 24 hours, both Ir-LB-MSNP(+SOS) and the liposomes showed comparable killing at 72 hours in both cell types (Figure 8). Empty nanoparticles do not induce toxicity at concentrations up to 500 µg/mL over 48 hours.

Comparative analysis of the *in vivo* toxicological effects of Ir-LB-MSNP(+SOS) vs Ir-Lipo(+SOS) in mice.

Maximum tolerated dose (MTD).

[0442] We began the *in vivo* toxicity studies by determining the MTD of the various irinotecan formulations. This was accomplished using the NCI protocol from the Toxicology and Pharmacology Branch (dtp.nci.nih.gov/branches/tpb/default.htm). Two

healthy male BALB/c mice were IV injected with 50 mg/kg (C_1 dose) free or encapsulated drug. We then used a dose escalation factor of 1.8 to determine the C_n dose where both animals ($n=2$) died within 24 hours. After treatment with free or encapsulated irinotecan. The second round of dose-seeking began with C_{n-1} dose, and utilized a 1.15 escalation factor (5 $n=2$) to find the MTD, where there was no mortality or serious morbidity. The MTD was further validated by injecting 5 mice, and following them for 15 days to ascertain that there was no morbidity, mortality or $>20\%$ weight loss. Our results demonstrated that the MTD dose values of Ir-LB-MSNP(+SOS), Ir-Lipo(+SOS), and free irinotecan were 295 mg/kg, 350 mg/kg, and 60 mg/kg, respectively. Compared to free drug, nano formulations showed (10 approximately 5-fold higher MTD doses, clearly suggesting the ability of toxicity reduction in use of both LB-MSNP and liposomal formulations.

Histological analysis.

[0443] In a separate experiment, the healthy mice received single IV administration of Ir-LB-MSNP(+SOS), Ir-Lipo(+SOS), and free irinotecan at a drug dose of 60 mg/kg IV. (15 24 hours post-treatment, the mice were sacrificed for organ harvest. Appropriate size sections of the sternum, gastrointestinal (GI, stomach and intestine) tract, thymus, liver, kidney, spleen, and lung were fixed in 10% formalin and then embedded into paraffin. Tissue sections of 4 μ m thickness were mounted on glass slides. The sections were stained with hematoxylin-eosin (H&E) and examined by light microscopy. The slides were read by (20 an experienced veterinary pathologist. The results showed a much reduced bone marrow toxicity for Ir-LB-MSNP(+SOS) and Ir-Lipo(+SOS) than free IRIN (Figure 9). We are performing more detailed studies on sternal bone marrow, using immunohistochemistry to assess whether there is differential toxicity on erithroid, megacariocyte or white blood cell precursors. Free irinotecan and the liposomal formulation resulted in nephrotoxicity, which (25 manifested as glomerular swelling. No significant renal abnormalities were found in the mice treated by Ir-LB-MSNP(+SOS) particles (Figure 10).

Conclusions.

[0444] The goal of this project is to develop an IV injectable irinotecan nano delivery platform for effective PDAC killing and toxicity reduction. Through the use of (30 high drug loading capacity, nanocarrier biocompatibility, and efficient drug encapsulation, the LB-MSNP platform has shown promising results in PDAC cells and acute animal toxicity studies.

Example 3

iRGD-mediated Transcytosis Enhances Irinotecan Delivery and Efficacy of a Silicasome Nanocarrier in Murine and Human Pancreatic Cancer

[0445] Although pancreatic ductal adenocarcinoma (PDAC) is almost uniformly fatal, some improved overall survival has been achieved with the introduction of nanocarriers that deliver irinotecan or paclitaxel. We have further improved these results in animal orthotopic models, using an irinotecan silicasome carrier that is comprised of mesoporous silica nanoparticles (MSNPs) coated with a lipid bilayer. The silicasome carrier also provides major toxicity reduction compared to a liposomal carrier for irinotecan. Although it is generally assumed that nanocarriers rely principally on abnormal leaky vasculature (*a.k.a.* the enhanced permeability and retention effect) for tumor access, a transcytosis transport pathway that is regulated by the neuropilin-1 (NRP-1) receptor has recently been reported. This unique transport pathway can be triggered by the cyclical iRGD peptide, which binds to tumor-associated integrins, where the peptide is processed for subsequent binding to NRP-1. Co-administration of iRGD enhanced the uptake of irinotecan-silicasomes 3-4 fold in a robust Kras orthotopic PDAC model, leading to a survival benefit and significant decrease in metastasis. Silicasomes with imbedded gold-nanoparticle cores were used for ultrastructural viewing of the transcytosis pathway *in vivo*, demonstrating that iRGD co-administration induces a vesicular transport pathway that transports the electron-dense carriers from the blood vessel lumen to a perinuclear site in cancer cells. iRGD-mediated enhancement of silicasome uptake was also observed in patient-derived xenografts, commensurate with the level of NRP-1 expression on tumor blood vessels. These results demonstrate the utility of iRGD for a potential personalized approach to PDAC therapy, using the irinotecan-silicasome carrier.

[0446] In this example we describe the development of a lipid bilayer coated mesoporous silica nanocarrier (silicasome) for irinotecan delivery that outperforms liposomes in a robust pancreatic ductal adenocarcinoma (PDAC) animal model. We demonstrate that silicasome uptake at the orthotopic tumor site can be enhanced 3-4 fold by the co-administration of iRGD peptide, which triggers a novel transcytosis pathway through ligation of the neuropilin-1 receptor. iRGD also enhanced silicasome uptake in patient-derived xenografts that express the same vascular receptor. We provide ultrastructural evidence of the vesicle-based transcytosis pathway, which could complement the enhanced permeability and retention effect commonly used to explain nanocarrier uptake at tumor

sites. The transcytosis pathway presents the possibility of enhancing the benefit of the irinotecan-silicasome carrier in PDAC patients.

Introduction

[0447] We have developed a multifunctional mesoporous silica nanoparticle (MSNP) platform, that has been adapted to provide high dose PDAC chemotherapy using a supported lipid bilayer (LB) for drug encapsulation (*see, e.g.*, Example 1). This carrier has also been designated as a “silicasome” to distinguish it from morphologically similar liposomal carriers, which contain a non-supported LB. Compared to liposomes, including an in-house liposomal carrier for the irinotecan, silicasomes described herein have been shown to exhibit a significantly higher loading capacity for this drug, improved circulatory stability (as a result of the supported LB), and reduced drug leakage. These features allow for improved pharmacokinetics and treatment efficacy of silicasomes *vs.* liposomes in a stringent orthotopic Kras PDAC model (*see, e.g.*, Example 1). In addition, silicasomes also provide major toxicity reduction in the gastrointestinal tract, liver, and bone marrow compared to the liposomal equivalent (*Id.*). Besides the success with irinotecan, the silicasome platform has also been adapted for synergistic delivery of paclitaxel and gemcitabine, allowing it to significantly (>10-fold) outperform the combination of free gemcitabine with Abraxane[®] in an orthotopic PDAC model. Noteworthy, above results with silicasome carriers could be achieved by “passive” delivery, without the need to resort to the use of targeting ligands.

[0448] Given this background, we were interested whether the efficacy of PDAC treatment by the irinotecan-loaded silicasome (Ir-silicasome) could be improved by transcytosis, and whether this is best accomplishable by iRGD conjugation to the carrier or its co-administration. In this Example, we demonstrate the feasibility of using the co-administration of free iRGD peptide to enhance carrier uptake and treatment efficacy in a Kras orthotopic model. Moreover, we provide ultrastructural evidence of a grouped vesicle system that allows transport of Au-labeled silicasomes from the blood vessel lumen to a perinuclear site in cancer cells. We also demonstrate in patient-derived PDAC xenografts that the relative abundance of NRP-1 expression on the tumor vasculature determines the response magnitude to silicasomes through the use of iRGD co-administration.

Results

Synthesis and characterization of silicasomes for drug loading and visualization in PDAC tumors

[0449] We have demonstrated high irinotecan loading by a LB-coated MSNP (*a.k.a.* a silicasome), using a remote loading technique that relies on a protonating agent (*see, e.g.*, Fig. 21A, top panel, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). Irinotecan is a weak basic and amphipathic molecule that can diffuse across the LB into the MSNP interior packaging space, where proton release by prior entrapped triethylammonium sucrose octasulfate (TEA₈SOS) converts the drug into a hydrophilic derivative, incapable of back-diffusion across the LB (Fig. 21A, top panel). The remote loading procedure was used to synthesize an Ir-silicasome batch that achieved a drug loading capacity of 50 wt% (w/w, irinotecan/MSNP) (*Id.*). To determine whether conjugation of an iRGD peptide onto the silicasome surface can impact carrier biodistribution to the PDAC tumor site, we also synthesized a particle batch in which the DSPE-PEG₂₀₀₀ component of the LB was used for conjugation to a cysteine residue in the peptide. This was accomplished by substituting DSPE-PEG₂₀₀₀ with DSPE-PEG₂₀₀₀-maleimide (see methods section), allowing thiol-maleimide coupling to the cysteine-modified peptide, Cys-c(CRGDKGPDC) (SEQ ID NO:11) (Fig. 21A, box 2). To confirm the success of the conjugation reaction, we also used a fluorescein-labeled version of the peptide (FAM-iRGD), developed by Ruoslahti *et al.*, to perform fluorescence spectroscopy of conjugated silicasomes after extensive washing (Fig. 27, panel A) (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520). This confirmed the stable association of the fluorescent peptide with the LB. The density of iRGD conjugation was limited to ~3 mol% (of all LB components) to prevent colloidal instability and interference in carrier uptake. The uptake of the intact silicasome-iRGD carrier was confirmed by flow cytometry and confocal microscopy in KPC cells (Figs. 27, panels B and C).

[0450] In order to perform TEM visualization of the transcytosis process in PDAC tumors, we also synthesized a batch of core-shell MSNPs that include ~10 nm electron-dense Au-nanoparticles (Fig. 1A, box 3). Similar to the bare particles, the core/shell particles could be effectively coated with a LB, as demonstrated by CryoEM (Fig. 21A, box 3). The detailed synthesis and characterization procedures for all the carriers used in this communication are discussed in the methods section of the Supporting Information. The main physicochemical characteristics of silicasomes are summarized below in Table 5.

Table 5. Illustrative silicasome physiochemical properties.

Property	Value
MSNP surface area (BET)	~ 850 m ² /g
MSNP pore volume (BET)	~0.75 cm ³ /g
MSNP pore size (BET, TEM)	~ 3 nm
MSNP core size (TEM)	~ 65 nm
LB thickness (cryo-EM)	~7 nm
Silicasome size (cryo-EM)	~80 nm
Hydrodynamic size (DLS)	~130 nm
Zeta potential	~ -10 mV
Drug loading capacity (irinotecan)	~50% (w/w, drug/MSNP)

Comparison of the effect of conjugated vs. non-conjugated iRGD peptide on silicasome biodistribution in an orthotopic PDAC model

[0451] Luciferase-expressing KPC cells, derived from a spontaneous PDAC tumor from a transgenic Kras^{LSL-G12D/+}; Trp53^{LSL-R172H/+}; Pdx-1-Cre animal were orthotopically implanted in the pancreas tails in immunocompetent B6/129 mice (*see, e.g.*, Fig. 21B, left panel, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715; Tseng *et al.* (2010) *Clin. Cancer Res.* 16(14): 3684-3695). This stringent PDAC tumor model mimics human PDAC for oncogene expression, growth characteristics, metastasis, histological features and development of a dysplastic stroma. Fig. 21B summarizes the details of the tumor growth characteristics and metastasis as seen during animal autopsy and IVIS imaging (Fig. 21B) (Tseng *et al.* (2010) *Clin. Cancer Res.* 16(14): 3684-3695; Torres *et al.* (2013) *PloS One* 8: e80580). Silicasome biodistribution to the orthotopic tumor site was assessed by a one-time intravenous (IV) injection of 50 mg/kg near-infrared (NIR) labeled (DyLight 680) particles that were either non-conjugated (*i.e.*, “iRGD free” status) or peptide-conjugated (“silicasome-iRGD”) (Fig. 22A). A third group of animals received co-administration of 8 μmol/kg free peptide plus non-conjugated particles (“silicasome + iRGD”). IVIS imaging of the explanted organs, performed 24 h after initial injection and animal sacrifice, demonstrated a prominent increase in the NIR signaling intensity at the tumor site for the “silicasome + iRGD” group compared to the signaling intensity in the silicasome-iRGD or silicasome only groups (Fig. 22A). Imaging intensity was quantified by IVIS Lumina Living Image software. In contrast to their lack of an *in vivo* effect, the peptide-conjugated silicasomes could be seen to enhance carrier uptake in KPC cells (Figure 27, panels B and C). We interpret that as sufficient NRP-1 receptor density to initiate trans-membrane

uptake, while the receptor abundance at the tumor vascular site may be limited in the number of conjugated particles that are allowed to go through, as previously reported for *in vitro/in vivo* comparisons by Ruoslahti *et al.* (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Teesalu *et al.* (2009) *Proc. Natl. Acad. Sci. USA*, 106(38): 16157-16162; Hussain *et al.* (2014) *Sci Rep* 4: 5232). In order to show that the NIR intensity (Fig. 22B, left panel) reflects actual MSNP uptake, inductively coupled plasma optical emission spectrometry (ICP-OES) was used for quantification of the tumor silicon (Si) content (Fig. 22B, right panel). This demonstrated a significant (~3-fold) increase in Si content in the “silicasome + iRGD” group compared to the silicasome-iRGD or the silicasome only groups. We also demonstrated the importance of the C-terminal (CendR) motif for PDAC biodistribution by performing a separate experiment to show that silicasome co-administration with a control non-CendR peptide, cyclo (RGDfK) (Sugahara *et al.* (2010) *Science*, 328(5981): 1031-1035) is incapable of enhancing silicasome uptake (Figures 28). This result was also confirmed by ICP-OES (Figure 28).

[0452] Tumor tissue sections were used to assess the relative abundance of intratumor biodistribution of NIR-labeled silicasomes and traveling distance from the tumor blood vessels, which were elucidated by anti-CD31 Alexa Fluor[®] 488-staining (Fig. 29). This analysis showed that iRGD co-administration was the most effective strategy to enhance silicasome uptake at orthotopic tumor sites. In contrast, the iRGD-conjugated carrier failed to exert a significant effect on the number of particles as well as the traveling distance in our KPC orthotopic model (Fig. 29). The above findings resulted in the use of free iRGD to perform subsequent efficacy studies.

[0453] Because all nanocarriers are also taken up in the reticuloendothelial system (RES), it is important to comment on the observable silicasome uptake in the liver and spleen (with little appreciable effects on the lungs, hearts, and kidneys). Interestingly, IVIS imaging showed a possible increase in the biodistribution of the conjugated carrier to the RES organs (Fig. 22A). This was confirmed by an increased Si content in the liver and spleen during performance of ICP-OES (Fig. 30). While we lack an exact explanation for this observation, it is possible that peptide conjugation, directly or indirectly, leads to particle opsonization and increased uptake by scavenger receptors, independent of NRP-1 (Li and Huang (2008) *Mol. Pharm.* 5(4): 496-504).

iRGD co-administration enhances the efficacy of irinotecan delivery by the silicasome

[0454] To demonstrate the possible therapeutic benefits of iRGD co-administration in PDAC treatment with the Ir-silicasome, an efficacy experiment was performed in the same orthotopic tumor model. Animals received IV injection of the Ir-silicasomes at a drug dose of 40 mg/kg (equivalent to carrier dose of 80 mg/kg), with or without co-administration of 8 μ mol/kg iRGD. Treatment commenced 13 days after orthotopic implementation of 2×10^6 KPC-luc cells (Fig. 23A), at which time the primary tumor size was ~3-5 mm in the absence of macro-metastasis (*see above*, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). Injections were repeated every 3 days, for a total of 4 administrations (Fig. 23A). The control groups consisted of animals receiving IV PBS, the Ir-silicasome alone (same dose), or free iRGD alone (same dose). Kaplan-Meier plots were used to express animal survival (*Id.*) and animal autopsy was used to assess the local tumor spread and the presence of metastasis. Numerous metastatic foci could be seen in the spleens, intestines, stomachs, livers, and kidneys of animals treated with saline or free iRGD peptide (Fig. 23B). While the Ir-silicasome significantly reduced the tumor burden and number of metastases, iRGD co-administration further enhanced the shrinking of the primary tumor as well as inhibiting the spread to the liver, stomach and intestines (Fig. 23B). The heat map in Fig. 23C provides a quantitative display of the impact of the co-administered peptide on metastatic disease. Log-rank testing (SPSS 19.0 software, IBM SPSS Statistics, USA) also demonstrated that, compared to the PBS group, treatment with the Ir-silicasome alone improved the survival by 28.6% as compared to 57.1 % during co-administration of the iRGD peptide (Fig. 23D); this difference is statistically significant ($p = 0.027$). We also confirmed that there is a significant (~2-fold) increase in the intratumoral content of irinotecan in animals receiving a one-time administration of iRGD (8 μ mol/kg) IV together with Ir-silicasomes (40 mg/kg drug), as determined by high performance liquid chromatography (HPLC) at 24 h (Fig. 23E).

iRGD-mediated silicasome uptake requires NRP-1 expression on the tumor vasculature

[0455] The mechanism of action of iRGD in mediating drug uptake is dependent on homing to $\alpha_v\beta_3$ or $\alpha_v\beta_5$ integrins, which are preferentially expressed on cancer blood vessels (Ruoslahti and Pierschbacher (1987) *Science*, 238(4826): 491-497; Hanahan and Weinberg (2000) *Cell*, 100(1): 57-70). Binding of the cyclical peptide to the integrins is followed by

cleavage and release of the C-terminal end of the peptide, which mediates the interaction with NRP-1 (this is also known as C-end rule). NRP-1 binding leads to the triggering of a system of vesicles that can assist drug and nanoparticle transport (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Pang *et al.* (2014) *Nat. Commun.* 5: 4904). In order to
5 determine the expression of NRP-1 at the KPC tumor site, IHC staining was performed by Alexa Fluor[®] 488-labeled anti-NRP-1, while endothelial cells and nuclei were localized by staining with Alexa Fluor[®] 594-labeled anti-CD31 and DAPI, respectively. Fluorescent microscopy and Image J analysis was performed to determine % overlap of NRP-1 with CD31; this demonstrated 94.2% co-localization (Fig. 24B). To verify the role of NRP-1 in
10 silicasome uptake at the orthotopic tumor site, tumor-bearing mice were pre-injected with an antagonist antibody to the b1b2 domain of NRP-1 (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Sugahara *et al.* (2010) *Science*, 328(5981): 1031-1035). Subsequent administration of iRGD plus NIR-labeled silicasomes demonstrated a definitive reduction in the carrier uptake compared to animals not treated with a blocking antibody (Fig. 24C). The
15 same interference was not seen with the control IgG (Fig. 24C). These data confirm the role of NRP-1 in iRGD-mediated silicasome uptake.

Ultrastructural demonstration of the transport of silicasomes by transcytosis vesicles at the KPC cancer site

[0456] It has been demonstrated that binding of the C-terminal end of iRGD to
20 NRP-1 can initiate a bulk transcytosis pathway that involves a novel vesicular transport mechanism of importance for nutrient delivery to cancer (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Sugahara *et al.* (2010) *Science*, 328(5981): 1031-1035; Pang *et al.* (2014) *Nat. Commun.* 5: 4904). To the best of our knowledge, this transcytosis pathway has never been directly visualized during the transport a therapeutic nanocarrier to the site of a
25 tumor. Electron microscopy (EM), which could provide visual enhancement demonstration, was used to attempt for ultrastructural analysis of the iRGD-mediated transport pathway in the KPC model. Initially, we compared TEM images taken of the harvested tumor site at different intervals after the IV injection of silicasomes, with or without iRGD co-administration (Fig. 25A). At 24 h, iRGD co-administration could clearly be seen to induce
30 the formation of grape-like vesicles, 110~370 nm in diameter, that are spread across the endothelial cells, from the luminal to the abluminal side of the blood vessel (Fig. 25A, regions “3” and “4”). These features resemble the vesiculo-vacuolar organelle or VVO described by Dvorak *et. al* (Feng *et al.* (1996) *J. Exp. Med.* 183(5): 1981-1986). Semi-

quantitative analysis of vesicle density, determined by counting the number of vesicles in at least 10 regions of interest (ROIs) and expressing the vesicle number *per* μm^2 of interior surface area in the cell, demonstrated that iRGD can increase the vesicle density ~ 3 -fold compared to animals not receiving peptide co-administration (Fig. 25A, left panel).

5 [0457] When attempting to visualize silicasome transport by the vesicular system, the low electron-density of the MSNPs is difficult to resolve the presence of the carrier in the heterogeneous and complex PDAC microenvironment. To address this challenge, silicasomes were synthesized to include a ~ 10 nm Au-nanoparticle that can be readily visualized by TEM (Fig. 21A) (Liu *et al.* 92013) *ACS Nano.* 7(7): 6244-6257). Mice
10 expressing orthotopic KPC tumors were IV injected with 50 mg/kg Au-silicasomes, in the absence or presence of 8 $\mu\text{mol/kg}$ iRGD. Tumor tissue was harvested 24 h post-injection and fixed to perform TEM analysis. A representative electron micrograph displaying, in one image, (i) electron-dense silicasomes in the blood vessel lumen, (ii) vesicular transport in endothelial cells, and (iii) particle deposition in the tumor matrix of an animal receiving
15 iRGD co-administration is shown in Fig. 25B. Higher magnifications of regions 1-3 of the image confirm the presence of Au-containing particles. It was also possible during iRGD co-administration to demonstrate the appearance of silicasomes in a perinuclear distribution in tumor cells undergoing apoptosis (Fig. 25C). This site is several hundred μm away from the nearest tumor blood vessel (Fig. 25C). While it was possible to observe lower particle
20 density in the tumor matrix of animals not receiving iRGD treatment, we could not locate any silicasomes being carried by the vesicular transport system, nor could we identify and alternative mechanism for entry in the static images (Fig. 31).

Differential impact of iRGD co-administration on silicasome uptake in patient-derived PDAC tumors, phenotyped for NRP-1 expression

25 [0458] While it is clear that the NRP-1 pathway can be functionally engaged by iRGD in the KPC tumor model, we were interested to see if the peptide could affect silicasome uptake in patient-derived xenografts growing in NOD SCID IL2 α knockout (NSG) mice (Rückert *et al.* (2012). *J. Surg. Res.* 172(1): 29-39). One of us (TD) has established a repository of 23 human PDAC tumors in NSG mice; these tumor samples
30 were obtained from patients during Whipple's surgery. Transferred tumor tissues were phenotypically characterized for cancer features that are distinctive of the corresponding human PDAC tumors, including stromal abundance and the expression of oncogenes and signal pathway components that are characteristic of PDAC (*Id.*). We used the phenotyping

information to select a pair of tumors with equivalent stromal abundance but differing in the density and distribution of NRP-1 expression, as determined by IHC staining. The staining was performed on tumor tissue harvested from NSG mice at the 3rd or 4th tumor passage, using image J software to compare the density of NRP-1 expression, as well as its co-localization with CD31 and the cellular nucleus (DAPI) (Fig. 26A). We were able to identify two patient tumor samples, designated XWR#8 and XWR#187, that showed similar collagen density (trichrome staining) but differing in NRP-1 expression (Fig. 26A). Thus, while XWR#8 was characterized by low NRP-1 abundance, XWR#187 had high NRP-1 expression levels (Fig. 26A). XWR#187 also exhibited more NRP-1 positive tumor blood vessels (~80%) compared to XWR#8 (~35%). Subcutaneous xenografts (n = 3) were established on the flank of NSG mice before the animals were IV injected with NIR labeled silicasomes (50 mg/kg), with or without co-administration of the iRGD peptide (8 µmol/kg). Following animal sacrifice after 24 h, IVIS imaging of the explanted tissues showed a 50% increase of NIR intensity at the tumor site of XWR#187 mice receiving iRGD; similar enhancement was not seen in XWR#8 (Fig. 26B). These data were confirmed by assessing the elemental Si content of the tumor tissues using ICP-OES (Fig. 26B). All things considered, our data indicate that the density and distribution of NRP-1 expression determine the degree of silicasome biodistribution to human tumors *in vivo*.

Discussion

[0459] In this example we demonstrate that the efficacy of an irinotecan-silicasome carrier can be significantly improved by the co-administration of an unconjugated iRGD peptide that does not require to be attached to the carrier to enhance tumor uptake. Co-administration of the free iRGD peptide increased silicasome uptake at orthotopic KPC tumors sites 3-4 fold, leading to enhanced killing of the primary tumor as well as metastasis inhibition. Overall, this resulted in a significant improvement in animal survival over the Ir-silicasome alone. The iRGD effect is mediated by interacting with tumor-associated integrins initially, followed by peptide cleavage and the release of the C-terminal end that engages NRP-1. Although the physiological role of NRP-1 is to control transcytosis for nutritional purposes, the vesicular system can also be used for the transport of nanoparticles, as demonstrated by the reduction of particle transport after the injection of receptor-blocking antibodies. Moreover, EM imaging provided ultrastructural evidence that iRGD could induce the appearance of grouped vesicles in endothelial cells, with the ability to carry Au-labeled silicasomes from the blood vessel lumen to the tumor matrix. We also

obtained evidence that NRP-1 regulates a transcytosis pathway in human pancreatic tumors, which have been implanted in NSG mice. The selection of a tumor pair with differential NRP-1 expression on the tumor vasculature demonstrated differences in carrier uptake and irinotecan delivery during iRGD treatment. All things considered, these data indicate that it is possible to use a personalized approach to PDAC chemotherapy to enhance the efficacy of the irinotecan silicasome carrier by iRGD co-administration.

[0460] The utility of a transcytosis pathway to enhance irinotecan delivery in PDAC is significant for a number of reasons. The first is the display of a dysplastic stroma, which contributes to drug resistance at the tumor site, in addition to augmenting tumor growth and metastasis (Feig *et al.* (2012) *Clin. Cancer Res.* 18(16): 4266-4276; Dimou *et al.* (2012) *Ther. Adv. Med. Oncol.* 1758834012446008). While it is often understood that abnormal vascular permeability is the reason for the nanoparticle extravasation, a concept referred to as the EPR effect, we know that the pancreatic cancer stroma actively interfere in vascular permeability (Meng *et al.* (2013) *ACS Nano.* 7(11): 10048-10065; Kano *et al.* (2007) *Proc. Natl. Acad. Sci. USA*, 104(9): 3460-3465; Cabral *et al.* (2011) *Nat. Nanotechnol.* 6(12): 815-823; Liu *et al.* (2012) *Proc. Natl. Acad. Sci. USA*, 109(41): 16618-16623). This includes the presence of pericytes that tightly adhere to vascular endothelial cells (*Id.*). Thus, while the EPR effect may contribute to the nanocarrier uptake in PDAC, it is important to consider the possibility that other vascular mechanisms may contribute to the nanoparticle uptake at the tumor site, including the possible contribution of nutritional transport pathways and vascular growth factors that regulate this transport, as well as vascular leakage (Jain and Stylianopoulos (2010) *Nat. Rev. Clin. Oncol.* 7(11): 653-664; Ruoslahti *et al.* (2010) *J. Cell. Biol.* 188(6): 759-768; Maeda *et al.* (2000) *J. Control. Release* 65(1): 271-284; Li and Huang (2008) *Mol. Pharm.* 5(4): 496-504; Feng *et al.* (1996) *J. Exp. Med.* 183(5): 1981-1986; Kobayashi *et al.* (2014) *Theranostics*, 4(1): 81-89). Ruoslahti *et al.* described an endocytic pathway, which plays a role in tumor nutrition, and can also be therapeutically engaged by tumor-penetrating iRGD peptides (Pang *et al.* (2014) *Nat. Commun.* 5: 4904). In addition, vascular growth factors, such as VEGF, VEGF-A, VEGF-A165, TGF- β and semaphorin 3A, display RGD motifs that allow binding to $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins on the tumor vasculature (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Sugahara *et al.* (2010) *Science*, 328(5981): 1031-1035; Pang *et al.* (2014) *Nat. Commun.* 5: 4904). Thus, proteolytic cleavage and release of the CendR motif could trigger NRP-1 mediated transcytosis (Pang *et al.* (2014) *Nat. Commun.* 5: 4904) in addition to the

role of the signaling pathways and vascular permeability associated with growth factor receptors (Kolodkin *et al.* (1997) *Cell*. 90(4): 753-762; Ellis (2006) *Mol. Cancer Ther.* 5(5): 1099-1107; Glinka and Prud'homme (2008) *J. Leukoc. Biol.* 84(1): 302-310).

Consequently, it is possible that the NRP-1 pathway may coexist with vascular leakage, including the EPR effect, but displaying different time kinetics. While the response to the CendR motif may commence within minutes, the EPR effect typically requires 6-8 hours to peak (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520; Maeda *et al.* (2003) *Int. Immunopharmacol.* 3(3): 319-328).

[0461] Our data for silicasome transcytosis during iRGD co-administration supplement previous attempts to overcome the stromal-vascular barrier during PDAC treatment (Feig *et al.* (2012) *Clin. Cancer Res.* 18(16): 4266-4276; Dimou *et al.* (2012) *Ther. Adv. Med. Oncol.* 1758834012446008; Meng *et al.* (2013) *ACS Nano.* 7(11): 10048-10065; Meng *et al.* (2015) *ACS Nano.* 9(4): 3540-3557). Several angiogenic drugs have been introduced to improve the tumor access of nanocarriers (Dimou *et al.* (2012) *Ther. Adv. Med. Oncol.* 1758834012446008; Kobayashi *et al.* (2014) *Theranostics*, 4(1): 81-89). Among these, VEGF can provide a temporary increase in the perfusion and blood vessel leakiness at the PDAC site to augment nanoparticle uptake (Olive *et al.* (2009) *Science*, 324(5933): 1457-1461; Jacobetz *et al.* (2013) *Gut*, 62(1): 112-120). Other agents that promote vascular permeability at the tumor site include bradykinin, nitric oxide, angiotensin-converting enzyme inhibitors, tumor necrosis factor α , heme oxygenase-1, collagenase and hyaluronidase (Kobayashi *et al.* (2014) *Theranostics*, 4(1): 81-89; Eikenes *et al.* (2004) *Cancer Res.* 64(14): 4768-4773; Seynhaeve *et al.* (2007) *Cancer Res.* 67(19): 9455-9462; Fang *et al.* (2011) *Adv. Drug Deliv. Rev.* 63(3): 136-151; Fang *et al.* (2012) *Cancer Sci.* 103(3): 535-541; Maeda (2013) *Cancer Sci.* 104(7): 779-789; Eikenes *et al.* (2005) *Br. J. Cancer.* 93(1): 81-88). We have also demonstrated that the use of a nanocarrier that delivers a small molecule inhibitor of the transforming growth factor beta (TGF- β) pathway, LY364947, can rapidly (<2 h) reverse pericyte adherence to endothelial cells *in vivo* (Meng *et al.* (2013) *ACS Nano.* 7(11): 10048-10065). The accompanying multiple fold increase in vascular permeability was demonstrated to provide a dramatic increase in the egress of gemcitabine-delivering liposomes at the tumor site (19). Other approaches for enhancing blood vessel permeability by targeting of the TGF- β pathway have also been reported (Cabral *et al.* (2011) *Nat. Nanotechnol.* 6(12): 815-823; Liu *et al.* (2012) *Proc. Natl. Acad. Sci. USA*, 109(41): 16618-16623). Finally, it is worth mentioning

that the silicasome carrier can achieve stromal reduction by co-delivery of paclitaxel with gemcitabine (Meng *et al.* (2015) *ACS Nano*. 9(4): 3540-3557), to the extent that our carrier could outperform the administration of Abraxane plus free gemcitabine during treatment of orthotopic PDAC tumors in mice (Frese *et al.* (2012) *Cancer Discov.* 2(3): 260-269).

5 [0462] Based on the demonstration that co-delivery of iRGD enhances the uptake of silicasomes at PDAC tumor sites (Fig. 22), it is important to stress that the co-administration approach overcomes a major limitation of the alternative delivery mechanism where the peptide is conjugated the nanocarrier. The explanation for this difference lies in the transport capacity of the carrier system, based on the available number
10 of NRP-1 receptors (Sugahara *et al.* (2010) *Science*, 328(5981): 1031-1035; Ruoslahti (2012) *Adv. Mater.* 24(28): 3747-3756;; Ruoslahti (2016) *Adv. Drug Deliv. Rev.* pii: S0169-409X(16)30094-1. doi: 10.1016/j.addr.2016.03.008). Thus, while the transport of the conjugated silicasome is limited by the relatively small and finite number of target receptors on the vasculature, separate injection of the unconjugated peptide triggers bulk transfer of
15 bystander silicasomes (in greater number) at the tumor site. Moreover, free iRGD also harbors anti-metastatic activity through the regulation of integrin function, as demonstrated by interference in the attachment and migration of culture tumor cells on a fibronectin matrix (Sugahara *et al.* (2015) *Mol. Cancer Ther.* 14(1): 120-128). This could explain, in part, the peptide interference on tumor metastasis in our study (Figs. 23B and 23C). The
20 use of the free peptide is also more practical and affordable for clinical use, as compared to relying on a conjugation mechanism that increases the cost and the complexity of the carrier synthesis.

Materials and Methods

[0463] A more detailed description of Materials and Experimental Procedures
25 appear in the Supplemental Materials and Methods below.

Silicasome Preparation

Synthesis of irinotecan loaded silicasome:

[0464] The 65 nm MSNP core was synthesized using a sol-gel procedure as shown above (*see also*, Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). A lipid biofilm was used
30 to produce the silicasomes as previously reported (Meng *et al.* (2015) *ACS Nano*. 9(4): 3540-3557; *see Example 1* and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). Briefly,

500 mg MSNPs were soaked in a 20 mL TEA₈SOS (80 mM solution, which was added on top of the lipid biofilm, comprised of a 550 mg mixture of DSPC/Chol/DSPE-PEG₂₀₀₀ (molar ratio 3:2:0.15), coated at the bottom of a round bottom flask (*see* Example 1, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). After sonication to accomplish particle coating with a LB, free TEA₈SOS was removed by size exclusion chromatography over a Sepharose CL-4B column. The TEA₈SOS loaded silicasomes were incubated in a 10 mg/mL irinotecan solution for drug loading in a water bath at 65 °C. The loading was stopped after 30 min by quenching in and ice water bath, following which the drug-loaded silicasomes were washed 3 times by centrifugation and re-suspended in PBS.

Synthesis of iRGD conjugated silicasome:

[0465] iRGD-conjugated silicasomes were synthesized by linking the peptide to a PEG chain included in the LB. This was accomplished by using commercially available DSPE-PEG₂₀₀₀-maleimide in place of DSPE-PEG₂₀₀₀, while maintaining the molar ratio of the lipids as described above. An excess (0.15 mL, 5 mg/mL) of the cysteine-modified iRGD peptide was conjugated to the DSPE-PEG₂₀₀₀-maleimide, using a thiol-maleimide reaction, carried out at room temperature for 4 h (Sugahara *et al.* (2009) *Cancer Cell*, 16(6): 510-520). The particles were washed to remove the non-reacted iRGD. The success of the conjugation reaction was confirmed by also preparing a batch of particles conjugated to a fluorescein (FAM) labeled iRGD peptide, followed by extensive washing (*Id.*) (Fig. 27, panel A).

Synthesis of silicasomes with a gold core marker:

[0466] 10 nm Au nanoparticles were synthesized in a citrate containing solution (see Supplemental Materials and Materials below). To grow the MSNP shell on the Au nanoparticle core, 36 mL of the citrate-capped particles was rapidly injected into 12 mL CTAC solution (25 wt% in H₂O). The particles were washed and re-suspended in a CTAC solution (6.25 wt% in H₂O), with stirring at 350 rpm for 5 min at 85 °C. To this mixture, we added 0.256 mL of 10% (w/v) triethanolamine for 10 min, followed by drop wise addition of 0.32 mL of the silica precursor, TEOS. The solution was stirred at 350 rpm for 20 min, leading to the creation of Au core/MSNP shell particles with an average size of ~65 nm. The particles were purified by sequential washing in 1% NaCl in methanol (w/v) and pure methanol. The Au-labeled MSNPs were then coated with a LB, as described before.

Biodistribution study of IV injected silicasomes with or without iRGD co-administration

[0467] IVIS (Xenogen) imaging was used to study the biodistribution of NIR-labeled silicasomes in the KPC-derived orthotopic model (n=3 mice/group) (*see* Example 1, and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715). Animals were IV injected with 50 mg/kg of the conjugated and non-conjugated silicasomes, with or without the co-administration of 8 μ mol/kg iRGD. Animals were sacrificed after 24 h followed by *ex vivo* imaging of the excised tumors and major organs. Tumor biodistribution was also confirmed by assessing the Si content, using an ICP-OES protocol (*Id.*).

Assessment of Ir-silicasome efficacy by iRGD co-administration in the KPC-derived orthotopic tumor model

[0468] Tumor-bearing B6/129 mice were randomly assigned into 4 groups, with 6 animals each. The first group was IV injected with silicasomes containing an irinotecan dose of 40 mg/kg (80 mg/kg MSNP) every 3 days, for a total of 4 administrations. The second group received the same dose of the Ir-silicasome plus co-administration of 8 μ mol/kg iRGD. The third and fourth groups were treated with PBS or iRGD alone. The mice were monitored daily up to the point of spontaneous animal death or approaching moribund status (*see* Example 1 and Liu *et al.* (2016) *ACS Nano*. 10(2): 2702-2715; Olive *et al.* (2009) *Science*, 324(5933): 1457-1461). Bioluminescence imaging of the primary tumor and metastasis sites were performed by injecting the animals intraperitoneally with 75 mg/kg D-Luciferin, 10 min before sacrifice. The tumor tissue and major organs (GI tract, liver, spleen, heart, lung and kidneys) were harvested for quantitative assessment of the bioluminescence image intensity.

Ultrastructural analysis of the transcytosis pathway through TEM viewing

[0469] KPC orthotopic tumor mice were treated by IV injection of 50 mg/kg of the Au-encapsulated silicasomes, with or without the co-administration of 8 μ mol/kg iRGD. Tumor biopsies were collected after 24 h, washed in PBS, and immediately fixed at 4 °C with 2.5% glutaraldehyde. Further sample preparation and sectioning were performed by the Electron Microscopy Services Center at UCLA. After fixation in 1% OsO₄, the samples were dehydrated in propylene oxide and embedded in resin. Tissue slices of 60-80 nm thick were placed on copper grids, and viewed under a JEOL 1200-EX electron microscope.

Silicasome biodistribution in patient-derived PDAC tumors

[0470] A repository of 23 PDAC samples were collected, with institutional human subject approval, from patients undergoing Whipple's surgery, and used to establish xenografts in NSG mice in Dr. Timothy Donahue's laboratory. Utilizing phenotyping data and performance of IHC staining for NRP-1 expression, two patient samples (XWR#8 and #187) were collected for growing fresh subcutaneous xenografts in the flanks of 6 weeks old female NSG mice (Rückert *et al.* (2012). *J. Surg. Res.* 172(1): 29-39). When the tumor size grew to a diameter of ~0.8 cm, 3 animals in each group receiving silicasomes, with or without iRGD co-administration, were used to assess biodistribution to the tumor site, similar to the procedure described above. Animals were sacrificed after 24 h for the performance of *ex vivo* imaging to determine the uptake of NIR-labeled silicasomes. The imaging data were also confirmed by assessing the Si content at the tumor sites by ICP-OES.

Statistical analysis

[0471] Comparative analysis of differences between groups was performed using the two-sided Student's *t*-test (Excel software, Microsoft). A statistically significant difference was determined at $p < 0.05$. Values were expressed as mean \pm SD of multiple determinations, as stated in the figure legends. The survival data was processed by Log Rank test (Mantel-Cox) using SPSS software.

Supplemental Materials and Methods

Materials

[0472] Tetraethylorthosicate (TEOS). triethanolamine. cetyltrimethylammonium chloride solution (CTAC. 25 wt% in water). (3-aminopropyl)triethoxysilane (APTES). triethylamine (TEA). gold (III) chloride hydrate. trisodium citrate dehydrate. Dowex 50WX8 resin. and Sepharose CL-4B were purchased from Sigma-Aldrich. USA. 1, 2-Distearoyl-sn-glycero-3-phosphocholine (DSPC). 1, 2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000] (ammonium salt) (DSPE-PEG₂₀₀₀). 1, 2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[maleimide(polyethylene glycol)-2000] (ammonium salt) (DSPE-PEG₂₀₀₀-maleimide) and cholesterol (Chol) were purchased from Avanti Polar Lipids. USA. Sucrose octasulfate (SOS) sodium salt was purchased from Toronto Research Chemicals. Canada. Irinotecan hydrochloride trihydrate

was purchased from LC Laboratories. USA. Penicillin, streptomycin, and Dulbecco's modified Eagle medium (DMEM) were obtained from Invitrogen. Fetal bovine serum (FBS) was purchased from Gemini Bio Products. USA. iRGD (CRGDKGPDC, SEQ ID NO: 11) was purchased from Biomatik. USA. Reactive iRGD modified with cysteine residue and blocking anti-NPR-1 antibody (against recombinant b1b2 domain of NRP-1) were kindly provided by Dr. Ruoslahti. Cyclo (-RGDfK) (SEQ ID NO:12) was purchased from ApexBio Technology. USA. Anti-CD31 antibody (catalog #553708) was purchased from BD Pharmingen™. USA. Anti-NPR-1 antibody (ab81321) was purchased from Abcam. USA. Control normal goat IgG (sc-2028) was purchased from Santa Cruz Biotechnology. USA. Alexa Fluor® 488 conjugated goat anti-rabbit IgG (H+L) secondary antibody (A11008). Alexa Fluor® 594 conjugated goat anti-rat IgG (H+L) secondary antibody (A11007) and DyLight 680 NHS ester were purchased from Thermo Fisher Scientific Inc. USA. Matrigel™ Matrix Basement Membrane was purchased from BD Bioscience. USA. All chemicals were directly used without further purification.

15 **NIR labeling of MSNPs using DyLight 680**

[0473] The NIR fluorescent dye, DyLight 680 NHS ester was used for MSNP labeling. First, the MSNP surface was functionalized with NH₂ groups for conjugation of the NHS ester. Briefly, 10 mg MSNPs were suspended in 1 mL of ethanol and mixed with 1 µL of APTES. The reaction took place under an inert N₂ atmosphere, while stirring at 80°C overnight. Subsequently, the mixture was centrifuged and washed 3 times with ethanol. The NH₂-conjugated MSNPs were suspended in 1 mL of DMF, mixed with 0.01 mg of DyLight 680 NHS ester and stirred at room temperature for 2 h. The labeled MSNPs were washed with ethanol and deionized water. NIR-labeled MSNPs were used to make the silicasomes.

25 **Synthesis of ~10 nm gold nanoparticle**

[0474] Gold nanoparticles of ~10 nm were made by adding 5 mL HAuCl₄ (10 mM) and 45 mL Milli-Q water to a 100 mL round-bottom flask equipped with a condenser. After reaching boiling temperature, while being stirred vigorously, 5.8 mL of sodium citrate (38.8 mM) was added into the boiling solution. This was accompanied by a color change from pale yellow to burgundy. The boiling solution was stirred for 10 min at 160 °C and then stirred for an additional 15 min without heating. The gold particles were used as a core for synthesis of silicasomes, as described in the methods section of the manuscript.

Cell line

[0475] An immortalized cell line was derived from a spontaneously tumor in a transgenic Kras^{LSL-G12D/+}; Trp53^{LSL-R172H/+}; Pdx-1-Cre mouse. To allow bioluminescence tumor imaging of the growing tumors after orthotopic implantation. the cells were
5 permanently transfected with a luciferase-based lentiviral vector in the UCLA vector core facility. A representative cell clone was obtained after use of a limiting dilution protocol.

Preparation of orthotopic tumors in immune competent mice, using the KPC-derived cell line

[0476] All animal experiments were performed with protocols approved by the
10 UCLA Animal Research Committee. Female B6/129 mice (~8 weeks) were purchased from The Jackson Laboratory. To grow orthotopic xenografts. the mice were anesthetized with isoflurane. followed by IP injection of 50 mg/kg ketamine and 10 mg/kg xylazine. The surgical site was shaved to leave a margin of 1 cm around the incision site and sterilized by scrubbing with betadine and 70% ethanol. The mice were positioned for surgery on a
15 heating pad. and the incision site in the left flank was draped with sterile gauze. A surgical incision of ~0.7 cm was made to expose the injection site. followed by an injection of 50 µL of DMEM/Matrigel (1:1 v/v) containing 2×10^6 KPC-luc cells into the pancreatic tail through a 27 G needle. The fascial layers were closed with absorbable sutures (PDS II. Ethicon) and the skin with nonabsorbable sutures (PROLENE. Ethicon). The mice were
20 kept on the warming pads until full recovery from the anesthesia. and then transferred to clean cages. Artificial tear ointment was used to protect the mouse eyes during the surgery.

Use of anti-NRP-1 blocking antibody to interfere in the iRGD effect

[0477] 50 µg of a blocking anti-NRP-1 antibody or a control IgG was injected 15 min before the KPC-derived orthotopic tumor-bearing mice received IV injection of 50
25 mg/kg NIR-labeled silicasome plus 8 µmol/kg free iRGD. Animals receiving the silicasome alone were used as controls. Animals were sacrificed at 24 h post-injection. and *ex vivo* NIR imaging was used to study the biodistribution of NIR labeled silicasomes. The *ex vivo* imaging data was quantified by NIR intensity analysis using IVIS software. followed by Si content analysis using ICP-OES.

HPLC analysis

[0478] For HPLC analysis of irinotecan in tissues. the harvested tumor and organ samples were weighed and homogenized on ice. Following the extraction of 0.1 mL tissue homogenate with 0.4 mL of an acidic solution (0.1 mol/L phosphoric acid/methanol. 1:4 v/v). the extracts were vortexed twice for 10s and centrifuged at 13,000 rpm for 10 min. The irinotecan-containing supernatants were filtered through 0.22 µm filters for HPLC analysis in a system containing a Knauer Smartline pneumatic pump. C18 column. K-2600 spectrophotometer. and Gina data acquisition software. The mobile phase, delivered at a flow rate of 1.0 mL/min, was comprised of a 3% triethylammonium acetate aqueous buffer (pH=5.5) and acetonitrile (73:27 v/v). Twenty microliters of an irinotecan-containing sample was injected to measure the drug absorption at 254 nm typically eluted in ~4.4 min. An irinotecan standard curve was generated over the concentration range 0.05-100 µg/mL.

Immunofluorescence staining

[0479] Dual color immunofluorescence staining was used to determine the NPR-1 positive blood vessels in KPC tumor tissue. The tumor tissues were cryo-embedded. using the OCT reagent. and used to prepare tumor sections. The sections were first treated with anti-NRP1 monoclonal antibody (1:250) at 4 °C overnight. After removal of the primary antibody and washing in PBS 3 times. the Alexa Fluor[®] 488 secondary antibody (1:500) was added and incubated for 1 h at room temperature. The same sections were also stained with anti-CD31 antibody. followed by Alexa Fluor[®] 594-conjugated secondary antibody treatment to identify CD31 expression. DAPI was used to localize the cellular nuclei. The stained slides were examined under a fluorescence microscope (Observer D1. Zeiss). The colocalization ratio of NRP-1⁺/CD31⁺ blood vessels were determined by Imaging J software.

Example 4

Comparative Analysis of the Protocell to the Silicasome

[0480] One approach to the preparation of lipid bilayer coated nanoparticles involved the use of MSNPs synthesized by an aerosol-assisted self-assembly method. The MSNPs were coated by the use of electrostatically charged liposomes, which sequentially adhere, rupture and then fuse with the negatively charged MSNP surface (*see, e.g., Liu et al. (2009) J. Am. chem. Soc. 131: 7567-7569*). This product, developed by Dr. Brinker's group at Sandia and the University of New Mexico, is called a "Protocell", and received a US

patent (US 8,992,984 B1) with the title: “Proto cells and their use for targeted delivery of multicomponent cargos to cancer cells”. The patent teaches, *inter alia*, that the “protocell can be formed by mixing the cargo components and the porous particle with liposomes or lipids, followed by fusing the lipid bilayer on the porous particle and synergistically loading the cargo components into one or more pores of the porous particle to form the protocell compound”. An example protocell shown in Figure 1C of the 8,992,984 patent depicts “positively charged porous particles [that] can be fused with negatively charged lipid bilayers, such as a DOPS lipid bilayer, wherein the positively charged porous particles can absorb negatively charge cargo components (*e.g.*, calcein or DNA or siRNA)”.

[0481] In developing the methods and silicasomes described herein we actively avoided the procedures and ingredients used in the Protocell because we regarded this approach as unpredictable and non-enabling for a number of reasons. Not only did we fail to achieve uniform MSNP coating using the liposomal fusion method described in the 8,992,984 patent, but we could not accomplish significant drug loading in a large number of attempts. This prompted us to develop the biofilm technique, sonication procedure, and loading methods described herein to obtain a reproducible procedure for uniform particle coating with an LB of different lipid composition to the protocell.

[0482] Figure 32 depicts data generated during our attempt to produce a “Protocell” that is equivalent to the protocells described by Ashley *et al.* (2011) *Nat. Mat.*, 10:389-397. Briefly, 100 μ L 25 mg/mL MSNPs were added to 100 μ L of a liposome at 2.5 mg/mL, followed by the sequence of steps described in the Ashley *et al.* publication (*Id.*). The final product is shown in the upper panel. This product was subjected to assessment of hydrodynamic size, size distribution and colloidal stability. The picture insert on the left shows a phase separation of the “coated” product and the DLS panel in the middle shows a small peak of uncoated particles plus a large speak of an agglomerated particle mass. The TEM image at the right confirms particle aggregation for the procells and colloidal stability for the silicasomes. The lack of colloidal stability of the protocells disqualifies the product for use by IV administration. In summary, unlike the protocell, the silicasome exhibits excellent colloidal stability, low PDI, and a narrow size distribution.

[0483] Table 6 illustrates various differences between the silicasome and protocell technologies.

Table 6. Comparison of certain features of the silicasome versus the protocell technologies.

Composition, method, or outcome	Silicasome	Protocell
MSNP synthesis method	Sol-gel method , yielding uniform particle size distribution with characteristics shown in Examples 1-3	Aerosol-assisted self-assembly, yielding particles with a heterogeneous size distribution and characteristics shown above
LB composition	<p>The lipid composition of the biofilm of illustrative, but non-limiting silicasomes is:</p> <p>DPPC/Chol/DSPE-PEG (PTX/GEM co-delivery silicasome)</p> <p>DSPC/Chol/DSPE-PEG (irinotecan silicasome)</p>	<p>The actual lipid composition of the coated particles is unknown, but is made from liposomes with the following compositions:</p> <p>DOPC/Chol/DOTAP</p> <p>DOPC/Chol/DOPS</p> <p>DOPC/Chol/DSPE-PEG</p> <p>DPPC /Chol/DSPE-PEG</p> <p>DOPC/Chol/DOPE-PEG</p> <p>DSPC/Chol/DSPE-PEG</p>
MSNP: lipid ratio (wt/wt)	1:1.1	10:1 (changed in Ashley <i>et al. supra</i> to 1:2)
Coating technique and efficiency	Single step approach, using sonication with high energy input and instantaneous encapsulation of ~100% of particles, with Cry EM proof in a population of particles	Multi-step approach, using vortexing of the particle/liposome mix, with low energy input, and lack of evidence for coating most particles in a population
Particle size, size distribution	<p>In illustrative, but non-limiting embodiments, MSNP core size ranges from 60-70 nm.</p> <p>They are highly uniform with PDI <0.1.</p> <p>After LB coating, the silicasome exhibits a hydrodynamic size of 110-130 nm with a PDI of <0.1.</p> <p>CryoEM shows the primary silicasome size of ~80 nm.</p> <p>Our LB coating procedure is also effective in coating MSNPs with uniform and smaller or bigger MSNP core size, such as 50-300 nm.</p>	<p>The MSNP core is <u>not</u> made in a sol-gel procedure, therefore, the core particles are <u>not</u> uniform but a mixture of large and small particles. In the protocells, MSNP core size ranges between 50 to 300 nm. Therefore, the PDI of the sample is too heterogeneous to be calculated.</p>
Pore distribution and size	Worm like distribution, size ~3 nm	Worm like; 1.95~2.25 nm
Colloidal stability	High colloidal stability. Ongoing stability test shows	Instantaneous agglomeration, making further stability testing

	good and unchanged stability from the moment of production through storage in PBS at 4°C for more than 6 months.	impossible
Synthesis Reproducibility	Highly reproducible, even by undergraduate students performing the technique for the 1 st time	Non-reproducible in our hands as well as working side-by-side with the Brinker people
Drug loading methods	Remote loading by a proton gradient	Mostly soaking in and using electrostatic attachment to the pores before encapsulation
Active Pharmaceutical Ingredient (API) characteristics	<p>Hydrophilic drugs, hydrophobic drugs, combination of hydrophilic and hydrophobic, remote loading of drugs with weak base characteristics described according to the following paradigm:</p> <p>The general characteristics of these cargo molecules include the following chemical properties: (i) organic molecular compounds that include primary, secondary, tertiary, or quaternary amine(s); (ii) a $pK_a < 11$ to allow protonation and entrapment behind the LB; (iii) a water solubility ranging from around 5 to 25 mg/mL and amphipathic characteristics that allow diffusion across the LB; (iv) an octanol/water partition coefficient or log <i>P</i> value of -3.0 to 3.0; (v) suitable molecular weight with a geometric size less than MSNP pore size (2-8 nm) to allow entry into the MSNP pores.</p>	<p>8,992,984 patent recites a large number of payloads without clear definition on the structure of these APIs. Minimally, this includes about 360 payload structures, such as chemoagents, kinase inhibitors, antibodies and protein. The encapsulation of various DNAs and siRNAs is also claimed. While electrostatic interaction may play a role in certain cases, most of these APIs are encapsulated via a passive encapsulation process.</p>

[0484] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent

applications cited herein are hereby incorporated by reference in their entirety for all purposes.

EDITORIAL NOTE

APPLICATION NUMBER - 2017206077

Please note: Claims pages following the description should be numbered 136 to page 138.

The claims defining the invention are as follows:

1. A nanoparticle drug carrier comprising:
 - a) a silica nanoparticle having a surface and defining a plurality of pores that are suitable to receive molecules therein;
 - b) a lipid bilayer coating the surface;
 - c) a cargo-trapping agent disposed within the plurality of pores, wherein the cargo trapping agent, before reaction with the drug, is triethylammonium sucrose octasulfate (TEA8SOS); and
 - d) a cargo comprising irinotecan, where the cargo is associated with the cargo trapping agent in the plurality of pores.
2. The nanoparticle drug carrier of claim 1, wherein the nanoparticle drug carrier has a submicron structure with a maximum dimension of less than one micron.
3. The nanoparticle drug carrier of claim 1, wherein the lipid bilayer stably seals the plurality of pores.
4. The nanoparticle of drug carrier of claim 1, wherein the nanoparticle drug carrier has a drug loading capacity of at least about 20% w/w.
5. The nanoparticle drug carrier of claim 1, wherein the lipid bilayer comprises a phospholipid, cholesterol, and an mPEG phospholipid.
6. The nanoparticle drug carrier according to claim 5, wherein the lipid bilayer comprises 50-90 mol % phospholipid, 10-50 mol % cholesterol, and 1-10 mol % mPEG.
7. The nanoparticle drug carrier of claim 1, wherein the lipid bilayer comprises DSPC/Chol/DSPE-PEG in a molar ratio of about 3:2:0.15.
8. The nanoparticle drug carrier of claim 1, wherein the lipid bilayer forms a substantially continuous bilayer encompassing the nanoparticle surface.
9. The nanoparticle drug carrier of claim 1, wherein the silica nanoparticle is a mesoporous silica nanoparticle.

10. The nanoparticle drug carrier of claim 1, wherein a population of the drug carriers in suspension shows a substantially unimodal size distribution and/or a PDI less than about 0.2.
11. The nanoparticle drug carrier of claim 1, wherein the drug is protonated and trapped in the plurality of pores as a gel-like precipitate in association with SOS⁸⁻.
12. A pharmaceutical formulation comprising a plurality of nanoparticle drug carriers, each nanoparticle drug carrier comprising:
 - a) a silica nanoparticle comprising a plurality of pores that are suitable to receive molecules therein;
 - b) a lipid bilayer stably sealing the plurality of pores;
 - c) a cargo trapping agent disposed within the plurality of pores, wherein the cargo trapping agent, before reaction with the drug, is triethylammonium sucrose octasulfate (TEA8SOS); and
 - d) a drug associated with the trapping agent in the plurality of pores, where said drug comprises irinotecan.
13. The pharmaceutical formulation of claim 12, wherein the plurality of nanoparticle drug carriers have a submicron structure with a maximum dimension of less than one micron.
14. The pharmaceutical formulation of claim 12, wherein the plurality of nanoparticle drug carriers has less than about 20% leakage of the drug over 24 hours in a biological buffer with pH of 7.4 at 37° C.
15. The pharmaceutical formulation of claim 12, wherein the drug is protonated and trapped in the plurality of pores as a gel-like precipitate in association with the trapping agent.
16. A nanocarrier, comprising:
 - a) a silica body comprising a surface and a plurality of pores;
 - b) a phospholipid bilayer coating the surface; and
 - c) a cargo-trapping agent and a drug comprising irinotecan disposed within the plurality of pores; wherein the nanocarrier has a submicron structure with a maximum dimension of less than one micron, and wherein the phospholipid bilayer stably seals the plurality of pores, and wherein the cargo trapping agent, before reaction with the drug, is triethylammonium sucrose octasulfate (TEA8SOS).

17. A method of treating cancer, comprising administering to a subject in need thereof an effective amount of a nanoparticle drug carrier according to any one of claims 1-11.
18. The method of claim 17, wherein the nanoparticle drug carrier is a primary therapy in a chemotherapeutic regimen.
19. The method of claim 17, wherein the nanoparticle drug carrier is a component in a multi-drug chemotherapeutic regimen.
20. The method of claim 19, wherein the multi-drug chemotherapeutic regimen comprises at least two drugs selected from the group consisting of irinotecan (IRIN), oxaliplatin (OX), 5-fluorouracil (5-FU), and leucovorin (LV).
21. The method of claim 17, wherein the cancer is pancreatic cancer, colorectal cancer, breast cancer, lung cancer, liver cancer, glioma, or melanoma.
22. The method of claim 17, wherein the cancer is pancreatic ductal adenocarcinoma (PDAC).

1/49




Fig. 1A

2/49




Fig. 1A, cont'd.

3/49




Fig. 1B




Fig. 1C

4/49




Fig. 1D

5/49





Fig. 1E

6/49



7/49




Fig. 2B

8/49




Fig. 2C




Fig. 2D

9/49





Fig. 3A

10/49



11/49





Fig. 3C



13/49




Fig. 3E

14/49




Fig. 4A




Fig. 4B




Fig. 4C

17/49




Fig. 4D

18/49





Fig. 4E



20/49




Fig. 6A




Fig. 6B

21/49




FIG. 7




Fig. 8

23/49




FIG. 9

24/49




FIG. 10

25/49




Fig. 11




Fig. 12

27/49




Fig. 13




Fig. 14




Fig. 15A




Fig. 15B

29/49




Fig. 16




Fig. 17

30/49




Fig. 18




Fig. 19

31/49




Fig. 20

32/49





Fig. 21A

33/49




Fig. 21B

~ 30 min surgery

34/49




Fig. 22A




Fig. 22B

35/49




Fig. 23A




Fig. 23B

36/49




Fig. 23C




Fig. 23D

37/49




Fig. 23E




Fig. 24A

38/49





Fig. 24B

39/49



40/49





Fig. 25A

41/49




Fig. 25B

42/49




Fig. 25C

43/49




Fig. 26A

44/49




Fig. 26B

45/49




Fig. 27

46/49




Fig. 28

47/49





Fig. 29

48/49




Fig. 30




Fig. 31

49/49




Fig. 32