

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 November 2008 (20.11.2008)

PCT

(10) International Publication Number
WO 2008/141011 A1

(51) International Patent Classification:
C07D 209/24 (2006.01) **C07D 209/02** (2006.01)

CA 92111 (US). **SEIDERS, Thomas, Jon** [US/US]; 5257 Soledad Mountain Road, San Diego, CA 92109 (US).

(21) International Application Number:
PCT/US2008/062793

(74) Agents: **HOSTETLER, Michael, J.** et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050 (US).

(22) International Filing Date: 6 May 2008 (06.05.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
11/746,010 8 May 2007 (08.05.2007) US

(71) Applicant (for all designated States except US): **AMIRA PHARMACEUTICALS, INC.** [US/US]; 9535 Waples Street, Suite 100, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUTCHINSON, John, Howard** [US/US]; 5948 Via Zurita, La Jolla, CA 92037 (US). **WANG, Bowei** [US/US]; 5025 Manor Ridge Lane, San Diego, CA 92130 (US). **STOCK, Nicholas, Simon** [GB/US]; 7087 Camino De Grazia, #150, San Diego,

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: 5-LIPOXYGENASE-ACTIVATING PROTEIN (FLAP) INHIBITORS




Figure 14

(57) Abstract: Described herein are compounds and pharmaceutical compositions containing such compounds, which modulate the activity of 5-lipoxygenase-activating protein (FLAP). Also described herein are methods of using such FLAP modulators, alone and in combination with other compounds, for treating respiratory, cardiovascular, and other leukotriene-dependent or leukotriene mediated conditions or diseases.



European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,
NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

5-LIPOXYGENASE-ACTIVATING PROTEIN (FLAP) INHIBITORS

RELATED APPLICATIONS

[0001] This application claims the benefit of priority to U.S. Patent Application Serial No. 11/746,010 filed 8 May 2007, which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

5 [0002] The MAPEG (membrane associated proteins involved in eicosanoid and glutathione metabolism) family of proteins are involved in eicosanoid formation. Compounds described herein inhibit the activity of at least one protein in the MAPEG family of proteins. Described herein are compounds, methods of making such compounds, pharmaceutical compositions and medicaments comprising such compounds, and methods of using such
10 activity.

BACKGROUND OF THE INVENTION

[0003] The MAPEG family of proteins includes proteins that are involved in the formation of eicosanoids from arachidonic acid in the lipoxygenase and cyclooxygenase metabolic pathways. The protein 5-lipoxygenase-activating protein (FLAP) is associated with the pathway of leukotriene synthesis. In particular, 5-lipoxygenase-
15 activating protein (FLAP) is responsible for binding arachidonic acid and transferring it to 5-lipoxygenase. *See, e.g., Abramovitz, M. et al., Eur. J. Biochem.* 215:105-111 (1993). 5-lipoxygenase can then catalyze the two-step oxygenation and dehydration of arachidonic acid, converting it into the intermediate compound 5-HPETE (5-hydroperoxyeicosatetraenoic acid), and in the presence of FLAP convert the 5-HPETE to Leukotriene A₄ (LTA₄).

[0004] LTA₄ is acted on by LTC₄ synthase, which conjugates LTA₄ with reduced glutathione (GSH) to form the
20 intracellular product leukotriene C₄ (LTC₄). LTC₄ is transformed to leukotriene D₄ (LTD₄) and leukotriene E₄ (LTE₄) by the action of gamma-glutamyl-transpeptidase and dipeptidases. LTC₄ synthase plays a pivotal role as the only committed enzyme in the formation of cysteinyl leukotrienes.

[0005] Leukotrienes are biological compounds formed from arachidonic acid in the leukotriene synthesis pathway (Samuelsson *et al., Science*, 220, 568-575, 1983; Cooper, *The Cell, A Molecular Approach*, 2nd Ed.
25 Sinauer Associates, Inc., Sunderland (MA), 2000). They are synthesized primarily by eosinophils, neutrophils, mast cells, basophils, dendritic cells, macrophages and monocytes. Leukotrienes have been implicated in biological actions including, by way of example only, smooth muscle contraction, leukocyte activation, cytokine secretion, mucous secretion, and vascular function.

[0006] Arachidonic acid is transformed to prostaglandin H₂ (PGH₂) by the action of cyclooxygenase enzymes
30 (COX-1 and COX-2). Microsomal prostaglandin (PG) E synthase 1 (mPGES-1) is responsible for transforming PGH₂ to prostaglandin E₂ (PGE₂), a prostaglandin involved in pain and inflammation.

SUMMARY OF THE INVENTION


[0007] Presented herein are methods, compounds, pharmaceutical compositions, and medicaments for (a) diagnosing, preventing, or treating allergic and non-allergic inflammation, (b) controlling signs and symptoms
35 that are associated with inflammation, and/or (c) controlling proliferative or metabolic disorders. These disorders

may arise from genetic, iatrogenic, immunological, infectious, metabolic, oncologic, toxic, and/or traumatic etiology.

[0008] In one aspect, the methods, compounds, pharmaceutical compositions, and medicaments described herein comprise 5-lipoxygenase-activating protein (FLAP) inhibitors described herein.

5 [0009] In one aspect provided herein are compounds of Formula (G), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, which antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent conditions or diseases, including, but not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[0010] In one embodiment, Formula (G) is as follows:



wherein,

15 Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl, or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl, or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;

20 Y is a (substituted or unsubstituted aryl), or -(substituted or unsubstituted heteroaryl);

R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, C(O), $-CH(OH)$, $-(substituted or unsubstituted C_1-C_6alkyl)$, or $-(substituted or unsubstituted C_2-C_6alkenyl)$;

25 R_7 is $L_3-X-L_4-G_1$, wherein,

L_3 is a or substituted or unsubstituted alkyl;

X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, $-C(O)NR_9$, $-NR_9C(O)NR_9$;

L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, a substituted or unsubstituted cyclic alkyl, or a substituted or unsubstituted heterocycloalkyl;

30 G_1 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, -

$S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, -

$NR_9C(=CHR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)C(=O)R_9$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, -

$C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-C(R_9)_2(OR_9)$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, -

35 $S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is -
 $OC(O)O-$, $-NHC(O)NH-$, $-NHC(O)O$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

or G_1 is $W-G_5$, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(R₉)₂(OR₉), -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈;

each R₈ is independently selected from substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted benzyl;

each R₉ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₁-C₆fluoroalkyl, substituted or unsubstituted C₃-C₈cycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted benzyl, and substituted or unsubstituted heteroarylmethyl; or two R₉ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or R₈ and R₉ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring and

each R₁₀ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl;

R₅ is H, halogen, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted -O-C₁-C₆alkyl;

R₁₁ is L₇-L₁₀-G₆, wherein L₇ is a bond, -C(O), -C(O)NH, -NHC(O), or (substituted or unsubstituted C₁-C₆alkyl); L₁₀ is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl);

G₆ is OR₉, -C(=O)R₉, -C(=O)OR₉, -SR₈, -S(=O)R₈, -S(=O)₂R₈, N(R₉)₂, tetrazolyl, -NHS(=O)₂R₈, -S(=O)₂N(R₉)₂, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, -C(=O)N(R₉)₂, N R₉C(O)R₉, C(R₉)₂C(=O)N(R₉)₂, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -O-, C(=O), S, S(=O), S(=O)₂, -NH-, -NHC(O)O-, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O)-;

or G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl) and G₇ is H, halogen, CN, NO₂, N₃, CF₃, OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -C(R₉)₂(OR₉), -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroalkyl), -L₅-(substituted or unsubstituted heteroaryl), -L₅-(substituted or unsubstituted heterocycloalkyl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is a bond, -O-, C(=O), S, S(=O), S(=O)₂, -NH-, -NHC(O)O-, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); and

R₁₂ is H, (substituted or unsubstituted C₁-C₆ alkyl), (substituted or unsubstituted C₃-C₆ cycloalkyl);

or glucuronide metabolite, or pharmaceutically acceptable solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[0011] For any and all of the embodiments, substituents can be selected from among from a subset of the listed alternatives.

[0012] For example, in some embodiments, R_{11} comprises at least one (unsubstituted or substituted) aromatic moiety and at least one (unsubstituted or substituted) cyclic moiety, wherein the (unsubstituted or substituted) cyclic moiety is a (unsubstituted or substituted) heterocycloalkyl group or a (unsubstituted or substituted) heteroaryl group. In some embodiments, R_{11} is not a thienyl-phenyl group.

[0013] In some embodiments Z is $S(O)_m$, or $[C(R_2)_2]_n C(R_1)_2 S(O)_m$; R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[0014] In some embodiments, m is 0; and n is 0 or 1.

[0015] In some embodiments, Y is -(substituted or unsubstituted heteroaryl); and G_6 is W- G_7 .

[0016] In some embodiments, Y is a substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1 O atoms and 0-1 S atoms.

[0017] In some embodiments, Y is selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl, wherein Y is substituted or unsubstituted.

[0018] In some embodiments, L_7 is a bond; L_{10} is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl).

[0019] In some embodiments, W is a (substituted or unsubstituted heterocycloalkyl), or a (substituted or unsubstituted heteroaryl).

[0020] In some embodiments, L_{10} is a (substituted or unsubstituted aryl); and R_{12} is H.

[0021] In some embodiments, W is a (substituted or unsubstituted heterocycloalkyl containing 0-2 nitrogen atoms, 0-1 O atoms and 0-1 S atoms) or a (substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1 O atoms and 0-1 S atoms).

[0022] In some embodiments, G_7 is H, halogen, CN, NO_2 , CF_3 , OCF_3 , C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $-C_1$ - C_6 fluoroalkyl, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$.

[0023] In some embodiments, W is a substituted or unsubstituted group selected from among pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl, furopyridinyl, quinoliziny, dioxiny, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, oxazinanonyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranyl, tetrahydropyranyl, dihydrooxazolyl, oxiranyl, pyrrolidinyl, pyrazolidinyl, dihydrothiophenonyl, imidazolidinonyl,

pyrrolidinonyl, dihydrofuranonyl, dioxolanonyl, thiazolidinyl, piperidinonyl, tetrahydronaphyridinyl, tetrahydroquinolinyl, tetrahydrothiophenyl, indolinyl, tetrahydroquinolinyl, and thiazepanyl.





[0024] In some embodiments, R_6 is H, or L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, -S(O), -S(O)₂, -C(O), -CR₉(OR₉), or substituted or unsubstituted alkyl.


[0025] In some embodiments, X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(=O)-, or -C(O)NR₉.

[0026] In some embodiments, G_1 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -N(R₉)CH₂CO₂R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)C(=O)R₉, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -C(R₉)₂(OR₉), -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L_5 is -OC(O)O-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G_1 is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(R₉)₂(OR₉), -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[0027] In some embodiments, R_6 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.


[0028] In some embodiments, G_1 is selected from among H, OH, CN, CO₂H, CO₂Me, CO₂Et, CO₂NH₂,






[0029] In some embodiments, L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-$

5 $CH_2C(CH_2CH_3)H-$, $-CH_2C(isopropyl)H-$, $-CH_2C(tert-butyl)H-$, $-CH_2C(CH_3)_2-$, $-CH_2C(CH_2CH_3)_2-$,




10



[0030] In some embodiments, R_7 is selected from among







[0031] In some embodiments, R₇ is selected from among



[0032] In some embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is OR₉ or -C(O)OR₉.

[0033] In some embodiments, L₃ is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methylpropan-1,2-diyl; 2-ethylpropan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethylbutan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propylpentan-1,2-diyl; pentan-1,5-diyl; or hexan-1,6-diyl.

[0034] In some embodiments, L₃ is 2-methylpropan-1,2-diyl; or 2-ethylbutan-1,2-diyl.

[0035] In some embodiments, L₃ is methandiyl; or ethan-1,2-diyl; and L₄ is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl; and G₁ is -OR₉, or -CO₂R₉.

[0036] In some embodiments, L₃ is methandiyl; X is a bond; L₄ is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and G₁ is -CO₂R₉.

[0037] In other embodiments, Y is selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl, wherein Y is substituted or unsubstituted.

[0038] In yet other aspects, Y is a substituted or unsubstituted heteroaryl containing 1-3 nitrogen atoms.

[0039] In one embodiment, Y is a substituted or unsubstituted group selected from among pyridinyl; benzothiazolyl; thiazolyl; imidazo[1,2-*a*]pyridinyl; quinolinyl; isoquinolinyl; isoxazolyl; pyrazolyl; indolyl; pyrazinyl; pyridazinyl; pyrimidinyl; quinazolinyl; and quinoxalinyl.

[0040] In other embodiments, L₇ is a bond; L₁₀ is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl).

[0041] In one embodiment, W is a (substituted or unsubstituted heterocycloalkyl), or a (substituted or unsubstituted heteroaryl).

[0042] In some embodiments, Y is selected from among pyridin-2-yl; 3-fluoro-pyridin-2-yl; 4-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 6-fluoro-pyridin-2-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 6-methyl-pyridin-2-yl; 3,5-dimethylpyridin-2-yl; 5,6-dimethyl-pyridin-2-yl; 5-ethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-chloro-pyridin-2-yl; 5-bromo-pyridin-2-yl; 6-cyclopropyl-pyridin-2-yl; 5-methyl-1-oxy-pyridin-2-yl; N-oxido-pyridin-2-yl; benzothiazol-2-yl; 2-methylthiazol-4-yl; imidazo[1,2-*a*]pyridin-2-yl; quinolin-2-yl; 6-fluoroquinolin-2-yl; 7-fluoroquinolin-2-yl; 6-methylquinolin-2-yl; 6-bromo-quinolin-2-yl; 1-oxy-quinolin-2-yl; 5-methylisoxazol-3-yl; 1,3-dimethylpyrazol-5-yl; 1,5-dimethylpyrazol-3-yl; 1*H*-indol-2-yl; 5-methyl-pyrazin-2-yl; 6-methyl-pyridazin-3-yl; quinoxalin-2-yl; quinazolin-2-yl; pyrimidin-2-yl; and 5-methylpyrimidin-2-yl.

[0043] In other embodiments, L₁₀ is a (substituted or unsubstituted aryl).

[0044] In other embodiments, R₁₂ is H.

[0045] In some embodiments, W is a (substituted or unsubstituted heterocycloalkyl containing 0-2 nitrogen atoms, 0-1 O atoms and 0-1 S atoms) or a (substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1 O atoms and 0-1 S atoms).

[0046] In one aspect, G₇ is H, halogen, CN, NO₂, CF₃, OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[0047] In yet other embodiments, R₆ is H, or L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O), -S(O)₂, -C(O), -CR₉(OR₉), or substituted or unsubstituted alkyl.

[0048] In some embodiments, X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(=O)-, or -C(O)NR₉.


[0049] In yet other embodiments, W is a substituted or unsubstituted group selected from among pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; tetrazolyl; tetrahydropyranyl, and morpholin-4-yl.

[0050] In some embodiments, R₆ is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl;

[0051] In some embodiments, R_5 is H.

[0053] In some embodiments, X is a bond.

[illegible]



5 **[0056]** In some embodiments, G₆ is selected from among pyridin-2-yl; pyridin-3-yl; pyridin-4-yl; 3-methyl-
pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 3-methoxy-pyridin-2-yl; 4-methoxy-pyridin-2-yl; 5-
methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 6-ethoxy-pyridin-2-yl; 3-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-
yl; 3-trifluoromethyl-pyridin-2-yl; 4-trifluoromethyl-pyridin-2-yl; 5-trifluoromethyl-pyridin-2-yl; 6-
trifluoromethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-fluoromethyl-pyridin-2-yl; 5-
10 methoxymethyl-pyridin-2-yl; 5-hydroxymethyl-pyridin-2-yl; 2-methyl-pyridin-3-yl; 6-methyl-pyridin-3-yl; 6-
cyano-pyridin-3-yl; 2-methoxy-pyridin-3-yl; 5-methoxy-pyridin-3-yl; 5-fluoro-pyridin-3-yl; 6-carbamoyl-
pyridin-3-yl; 6-hydroxy-pyridin-3-yl; 6-methoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-bromo-6-methoxy-
pyridin-3-yl; 6-trifluoromethyl-pyridin-3-yl; 6-trifluoromethyl-pyridin-4-yl; 2-trifluoromethyl-pyridin-5-yl; 2-
acetyl-amino-pyridin-5-yl; pyrazin-2-yl; pyrimidin-2-yl; pyrimidin-5-yl; 5-amino-pyrazin-2-yl; 6-hydroxy-
15 pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-methyl-pyridazin-3-yl; 5-hydroxy-pyrimidin-2-yl; 2-methoxy-
pyrimidin-5-yl; 6-methyl-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-ethoxy-pyridazin-3-yl; 3-methoxy-
pyridazin-6-yl; 4-methoxy-tetrahydro-pyran-4-yl; 6-ethoxy-pyridin-3-yl; and 5-fluoro-pyridin-2-yl.

[0057] In some embodiments, Y is a substituted or unsubstituted group selected from among pyridinyl and quinolinyl.

20 **[0058]** In some embodiments, L_7 is a bond; L_{10} is a (substituted or unsubstituted aryl); and G_6 is W- G_7 , wherein W is (substituted or unsubstituted heterocycloalkyl), or a (substituted or unsubstituted heteroaryl).

[0059] In some embodiments, W is a substituted or unsubstituted group selected from among pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; and tetrazolyl.

25 **[0060]** In some embodiments, R₆ is L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O), -S(O)₂, -C(O), -CR₉(OR₉), or substituted or unsubstituted alkyl.

[0061] In some embodiments, L₁₀ is phenyl.

[0062] In some embodiments, R₆ is L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a S, -S(O)₂, -S(O)-, or -C(O).

[0063] In some embodiments, R₉ is H or C₁-C₆ alkyl; and R₁₂ is H.

[0064] In some embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is OR₉ or -C(O)OR₉.

[0065] In some embodiments, L_3 is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methylpropan-1,2-diyl; 2-ethylpropan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethylbutan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propylpentan-1,2-diyl; pentan-1,5-diyl; or hexan-1,6-diyl.

5 [0066] In some embodiments, L_3 is propan-1,2-diyl; 2-methylpropan-1,2-diyl; 2-ethylpropan-1,2-diyl; butan-1,2-diyl; 2-ethylbutan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propylpentan-1,2-diyl.

[0067] In some embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methylbutanoyl; 3,3-dimethylbutanoyl; 2-ethylbutanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; tert-butylsulfanyl; tert-butylsulfinyl; or tert-butylsulfonyl.

[0068] In some embodiments, L_3 is 2-methylpropan-1,2-diyl; or 2-ethylbutan-1,2-diyl.

[0069] In some embodiments, G_1 is $-OR_9$, $N(R_9)_2$, or $-CO_2R_9$.

[0070] In some embodiments, G_1 is $-OR_9$, or $-CO_2R_9$.

15 [0071] In some embodiments, G_1 is $-CO_2R_9$.

[0072] In some embodiments, L_3 is methandiyl; or ethan-1,2-diyl; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[0073] In some embodiments, X is a bond; and L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cyclic alkyl.

[0074] In some embodiments, L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

25 [0075] In some embodiments, L_3 is methandiyl; X is a bond; and L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[0076] In some embodiments, L_4 is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and G_1 is $-CO_2R_9$.

[0077] In another aspect, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methylbutanoyl; 3,3-dimethylbutanoyl; 2-ethylbutanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

35 [0078] In one aspect, R_9 is H.

[0079] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[0080] In one aspect, provided herein is a compound selected from among Tables 1-5.

[0081] Compounds described herein inhibit the activity of at least one protein in the MAPEG family of proteins. In one aspect, compounds described herein inhibit the activity of at least one protein in the MAPEG family of proteins selected from among FLAP, LTC₄ synthase, or mPGES-1. In another aspect, compounds
5 described herein inhibit the activity of at least one protein in the MAPEG family of proteins selected from among FLAP and LTC₄ synthase.

[0082] In another aspect, compounds described herein inhibit the activity of FLAP.

[0083] In one aspect, provided herein is a pharmaceutical composition comprising an effective amount of a compound described herein, and a pharmaceutically acceptable excipient.

10 [0084] In one aspect, described herein is the use of a compound described herein in the manufacture of a medicament for the inhibition of at least one protein member of the MAPEG family of proteins. In one aspect, the protein member of the MAPEG family of proteins is selected from among FLAP, LTC₄ synthase, and mPGES-1. In one aspect, the protein member of the MAPEG family of proteins is FLAP.

[0085] In one aspect, described herein is a method of decreasing acyl glucuronide formation of a compound
15 described herein where G₁ is CO₂H or OH, the method comprising substituting the alkyl carbon atom of L₃, X, or L₄ that is adjacent to the -CO₂H or -OH group with at least one substituent that is larger than methyl. In one aspect, the alkyl carbon atom of L₃, X, or L₄ that is adjacent to the -CO₂H or -OH group of G₁ is substituted with two ethyl groups.

[0086] In one aspect, described herein is the use of a compound described herein in the manufacture of a
20 medicament for the treatment of a leukotriene dependent or leukotriene-mediated disease or condition. In one aspect, described herein is the use of a compound described herein in the manufacture of a medicament for the treatment of inflammation in a mammal. In one aspect, described herein is the use of a compound described herein in the manufacture of a medicament for the treatment of respiratory disease in a mammal. In one aspect, described herein is the use of a compound described herein in the manufacture of a medicament for the treatment
25 of cardiovascular disease in a mammal.

[0087] Articles of manufacture, comprising packaging material, a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), which is effective for modulating the activity of 5-lipoxygenase activating protein, or for treatment, prevention or amelioration of one or more symptoms of a leukotriene dependent or leukotriene-mediated disease or condition, within the packaging
30 material, and a label that indicates that the compound or composition, or pharmaceutically acceptable salt, pharmaceutically acceptable N-oxide, pharmaceutically acceptable acyl glucuroide metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, is used for modulating the activity of 5-lipoxygenase activating protein, or for treatment, prevention or amelioration of one or more symptoms of a leukotriene dependent or leukotriene-mediated disease or condition, are provided.

35 [0088] In another aspect, provided herein is a method for treating inflammation in a mammal comprising administering a therapeutically effective amount of a compound provided herein to the mammal in need.

[0089] In yet another aspect, provided herein is a method for treating asthma in a mammal comprising administering a therapeutically effective amount of a compound provided herein to the mammal in need. In a further or alternative embodiment, provided herein is a method for treating asthma in a mammal comprising
40 administering a therapeutically effective amount of a compound provided herein, such as, for example, a

compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), wherein Z is $[C(R_2)_2]_n C(R_1)_2 O$, to the mammal in need.

[0090] In another aspect are compounds presented in any of Figures 8, 9, 10, or 11, or pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically acceptable glucuronide metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, which antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent conditions or diseases, including, but not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[0091] In another aspect are compounds presented in any of Tables 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14, pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically acceptable glucuronide metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, which antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent conditions or diseases, including, but not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[0092] In further or alternative embodiments, the compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be inhibitors of 5-lipoxygenase-activating protein (FLAP), while in still further or alternative embodiments, such inhibitors are selective for FLAP. In even further or alternative embodiments, such inhibitors have an IC_{50} below 50 μM in the FLAP binding assay.

[0093] In further or alternative embodiments, the compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be included into pharmaceutical compositions or medicaments used for treating a leukotriene-dependent or leukotriene mediated condition or disease in a patient.

[0094] In another aspect the inflammatory conditions include, but are not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, aortic aneurysm, myocardial infarction, and stroke. In other aspects the proliferative disorders include, but are not limited to, cancer and noncancerous disorders, including, but not limited to, those involving the skin or lymphatic tissues. In other aspects the metabolic disorders include, but are not limited to, bone remodeling, loss or gain. In additional aspects, such conditions are iatrogenic and increases in, or abnormal localization of, leukotrienes may be induced by other therapies or medical or surgical procedures.

[0095] In other aspects, the methods, compounds, pharmaceutical compositions, and medicaments described herein may be used to prevent the cellular activation of 5-lipoxygenase, while in other aspects the methods, compounds, pharmaceutical compositions, and medicaments described herein may be used to limit the formation of leukotrienes. In other aspects, such methods, compounds, pharmaceutical compositions, and medicaments may comprise FLAP inhibitors disclosed herein for the treatment of asthma by (a) lowering the concentrations of leukotrienes in certain tissue(s) of the body or in the entire body of a patient, (b) modulating the activity of enzymes or proteins in a patient wherein such enzymes or proteins are involved in the leukotriene pathway such

as, by way of example, 5-lipoxygenase-activating protein or 5-lipoxygenase, or (c) combining the effects of (a) and (b). In yet other aspects, the methods, compounds, pharmaceutical compositions, and medicaments described herein may be used in combination with other medical treatments or surgical modalities.

[0096] In one aspect are methods for reducing/inhibiting the leukotriene synthetic activity of 5-lipoxygenase-activating protein (FLAP) in a mammal comprising administering to the mammal at least once an effective amount of a compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[0097] In a further or alternative embodiment, the "G" group (e.g. G₁, G₅, G₆, G₇) of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is any group that is used to tailor the physical and biological properties of the molecule. Such tailoring/modifications are achieved using groups which modulate acidity, basicity, lipophilicity, solubility and other physical properties of the molecule. The physical and biological properties modulated by such modifications to "G" include, by way of example only, solubility, *in vivo* absorption, and *in vivo* metabolism. In addition, *in vivo* metabolism may include, by way of example only, controlling *in vivo* PK properties, off-target activities, potential toxicities associated with cypP450 interactions, drug-drug interactions, and the like. Further, modifications to "G" allow for the tailoring of the *in vivo* efficacy of the compound through the modulation of, by way of example, specific and non-specific protein binding to plasma proteins and lipids and tissue distribution *in vivo*. Additionally, such tailoring/modifications to "G" allow for the design of compounds selective for 5-lipoxygenase-activating protein over other proteins. In further or alternative embodiments, "G" is L₂₀-Q, wherein L₂₀ is an enzymatically cleavable linker and Q is a drug, or an affinity moiety. In further or alternative embodiments, the drug includes, by way of example only, leukotriene receptor antagonists and anti-inflammatory agents. In further or alternative embodiments, the leukotriene receptor antagonists include, but are not limited to, CysLT1/CysLT2 dual antagonists and CysLT1 antagonists. In further or alternative embodiments, the affinity moiety allows for site specific binding and include, but are not limited to, antibodies, antibody fragments, DNA, RNA, siRNA, and ligands.

[0098] In another aspect are methods for modulating, including reducing and/or inhibiting the activity of 5-lipoxygenase activating protein, directly or indirectly, in a mammal comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[0099] In another aspect are methods for modulating, including reducing and/or inhibiting, the activity of leukotrienes in a mammal, directly or indirectly, comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00100] In another aspect are methods for treating leukotriene-dependent or leukotriene mediated conditions or diseases, comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00101] In another aspect are methods for treating inflammation comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00102] In another aspect are methods for treating respiratory diseases comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). In a further embodiment of this aspect, the respiratory disease is asthma. In a further embodiment of this aspect, the respiratory disease includes, but is not limited to, adult respiratory distress syndrome and allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, seasonal asthma,

[00103] In another aspect are methods for treating chronic obstructive pulmonary disease comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). In a further embodiment of this aspect, chronic obstructive pulmonary disease includes, but is not limited to, chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation and cystic fibrosis.

[00104] In another aspect are methods for preventing increased mucosal secretion and/or edema in a disease or condition comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00105] In another aspect are methods for treating vasoconstriction, atherosclerosis and its sequelae myocardial ischemia, myocardial infarction, aortic aneurysm, vasculitis and stroke comprising administering to the mammal an effective amount of a compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00106] In another aspect are methods for treating organ reperfusion injury following organ ischemia and/or endotoxic shock comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00107] In another aspect are methods for reducing the constriction of blood vessels in a mammal comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00108] In another aspect are methods for lowering or preventing an increase in blood pressure of a mammal comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00109] In another aspect are methods for preventing eosinophil and/or basophil and/or dendritic cell and/or neutrophil and/or monocyte recruitment comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00110] A further aspect are methods for the prevention or treatment of abnormal bone remodeling, loss or gain, including diseases or conditions as, by way of example, osteopenia, osteoporosis, Paget's disease, cancer and other diseases comprising administering to the mammal at least once an effective amount of at least one

compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00111] In another aspect are methods for preventing ocular inflammation and allergic conjunctivitis, vernal keratoconjunctivitis, and papillary conjunctivitis comprising administering to the mammal at least once an effective amount of at least one having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00112] In another aspect are methods for treating CNS disorders comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). CNS disorders include, but are not limited to, multiple sclerosis, Parkinson's disease, Alzheimer's disease, stroke, cerebral ischemia, retinal ischemia, post-surgical cognitive dysfunction, migraine, peripheral neuropathy/neuropathic pain, spinal cord injury, cerebral edema and head injury.

[00113] A further aspect are methods for the treatment of cancer comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). The type of cancer may include, but is not limited to, pancreatic cancer and other solid or hematological tumors.

[00114] In another aspect are methods for treating endotoxic shock and septic shock comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00115] In another aspect are methods for treating rheumatoid arthritis and osteoarthritis comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00116] In another aspect are methods for preventing increased GI diseases comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such diseases include, by way of example only, chronic gastritis, eosinophilic gastroenteritis, and gastric motor dysfunction.

[00117] A further aspect are methods for treating kidney diseases comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such diseases include, by way of example only, glomerulonephritis, cyclosporine nephrotoxicity renal ischemia reperfusion.

[00118] In another aspect are methods for preventing or treating acute or chronic renal insufficiency comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00119] In another aspect are methods for treating type II diabetes comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00120] In another aspect are methods to diminish the inflammatory aspects of acute infections within one or more solid organs or tissues such as the kidney with acute pyelonephritis.

[00121] In another aspect are methods for preventing or treating acute or chronic disorders involving recruitment or activation of eosinophils comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

5 [00122] In another aspect are methods for preventing or treating acute or chronic erosive disease or motor dysfunction of the gastrointestinal tract caused by non-steroidal anti-inflammatory drugs (including selective or non-selective cyclooxygenase -1 or -2 inhibitors) comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

10 [00123] A further aspect are methods for the prevention or treatment of rejection or dysfunction in a transplanted organ or tissue comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

15 [00124] In another aspect are methods for treating inflammatory responses of the skin comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such inflammatory responses of the skin include, by way of example, dermatitis, contact dermatitis, eczema, urticaria, rosacea, and scarring. In another aspect are methods for reducing psoriatic lesions in the skin, joints, or other tissues or organs, comprising administering to the mammal an effective amount of a first compound having the structure of any of
20 Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00125] A further aspect are methods for the treatment of cystitis, including, by way of example only, interstitial cystitis, comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

25 [00126] A further aspect are methods for the treatment of metabolic syndromes such as Familial Mediterranean Fever comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

30 [00127] In a further aspect are methods to treat hepatorenal syndrome comprising administering to the mammal at least once an effective amount of at least one compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

35 [00128] In another aspect is the use of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in the manufacture of a medicament for treating an inflammatory disease or condition in an animal in which the activity of at least one leukotriene protein contributes to the pathology and/or symptoms of the disease or condition. In one embodiment of this aspect, the leukotriene pathway protein is 5-lipoxygenase-activating protein (FLAP). In another or further embodiment of this aspect, the inflammatory disease or conditions are respiratory, cardiovascular, or proliferative diseases.

40 [00129] In any of the aforementioned aspects are further embodiments in which administration is enteral, parenteral, or both, and wherein (a) the effective amount of the compound is systemically administered to the mammal; and/or (b) the effective amount of the compound is administered orally to the mammal; and/or (c) the

effective amount of the compound is intravenously administered to the mammal; and/or (d) the effective amount of the compound administered by inhalation; and/or (e) the effective amount of the compound is administered by nasal administration; or and/or (f) the effective amount of the compound is administered by injection to the mammal; and/or (g) the effective amount of the compound is administered topically (dermal) to the mammal; and/or (h) the effective amount of the compound is administered by ophthalmic administration; and/or (i) the effective amount of the compound is administered rectally to the mammal.

[00130] In any of the aforementioned aspects are further embodiments in which the mammal is a human, including embodiments wherein (a) the human has an asthmatic condition or one or more other condition(s) selected from the group consisting of allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, or seasonal asthma, or chronic obstructive pulmonary disease, or pulmonary hypertension or interstitial lung fibrosis. In any of the aforementioned aspects are further embodiments in which the mammal is an animal model for pulmonary inflammation, examples of which are provided herein.

[00131] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered once; (ii) the compound is administered to the mammal multiple times over the span of one day; (iii) continually; or (iv) continuously.

[00132] In any of the aforementioned aspects are further embodiments comprising multiple administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered in a single dose; (ii) the time between multiple administrations is every 6 hours; (iii) the compound is administered to the mammal every 8 hours;. In further or alternative embodiments, the method comprises a drug holiday, wherein the administration of the compound is temporarily suspended or the dose of the compound being administered is temporarily reduced; at the end of the drug holiday, dosing of the compound is resumed. The length of the drug holiday can vary from 2 days to 1 year.

[00133] In any of the aforementioned aspects involving the treatment of leukotriene dependent diseases or conditions are further embodiments comprising administering at least one additional agent, each agent may be administered in any order, including, by way of example, an anti-inflammatory agent, a different compound having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), a CysLT₁ receptor antagonist, or a CysLT₁/CysLT₂ dual receptor antagonist. In further or alternative embodiments, the CysLT₁ antagonist is selected from montelukast (Singulair™: [1-[[1-[3-[2-[(7-chloro-2-quinolyl)]vinyl]phenyl]-3-[2-(1-hydroxy-1-methyl-ethyl)phenyl]-propyl]sulfanylmethyl]cyclopropyl]acetic acid), zafirlukast (Accolate™: 3-[[2-methoxy-4-(o-tolylsulfonylcarbamoyl)phenyl]methyl]-1-methyl-1H-indol-5-yl]aminoformic acid cyclopentyl ester) or pranlukast (Onon™: 4-oxo-8-[p-(4-phenylbutoxy)benzoylamino]-2-tetrazol-5-yl)-4H-1-benzopyran)

[00134] In further or alternative embodiments, the anti-inflammatory agent includes, but is not limited to, non-steroidal anti-inflammatory drugs such as a cyclooxygenase inhibitor (COX-1 and/or COX-2), lipoxigenase inhibitors and steroids such as prednisone or dexamethasone. In further or alternative embodiments, the anti-inflammatory agent is selected from the group consisting of Arthrotec®, Asacol, Auralgan®, Azulfidine, Daypro,

etodolac, Ponstan, Salofalk, Solu-Medrol, aspirin, indomethacin (Indocin™), rofecoxib (Vioxx™), celecoxib (Celebrex™), valdecoxib (Bextra™), diclofenac, etodolac, ketoprofen, Lodine, Mobic, nabumetone, naproxen, piroxicam, Celestone, prednisone, Deltasone, or any generic equivalent thereof.

[00135] In any of the aforementioned aspects involving the treatment of proliferative disorders, including cancer,

are further embodiments comprising administering at least one additional agent selected from the group consisting of alemtuzumab, arsenic trioxide, asparaginase (pegylated or non-), bevacizumab, cetuximab, platinum-based compounds such as cisplatin, cladribine, daunorubicin/doxorubicin/idarubicin, irinotecan, fludarabine, 5-fluorouracil, gemtuzumab, methotrexate, Paclitaxel™, taxol, temozolomide, thioguanine, or classes of drugs including hormones (an antiestrogen, an antiandrogen, or gonadotropin releasing hormone analogues, interferons such as alpha interferon, nitrogen mustards such as busulfan or melphalan or mechlorethamine, retinoids such as tretinoin, topoisomerase inhibitors such as irinotecan or topotecan, tyrosine kinase inhibitors such as gefitinib or imatinib, or agents to treat signs or symptoms induced by such therapy including allopurinol, filgrastim, granisetron/ondansetron/palonosetron, dronabinol.

[00136] In any of the aforementioned aspects involving the therapy of transplanted organs or tissues or cells are further embodiments comprising administering at least one additional agent selected from the group consisting of azathioprine, a corticosteroid, cyclophosphamide, cyclosporin, dacluzimab, mycophenolate mofetil, OKT3, rapamycin, tacrolimus, or thymoglobulin.

[00137] In any of the aforementioned aspects involving the therapy of interstitial cystitis are further embodiments comprising administering at least one additional agent selected from dimethylsulfoxide, omalizumab, and pentosan polysulfate.

[00138] In any of the aforementioned aspects involving the therapy of disorders of bone are further embodiments comprising administering at least one additional agent selected from the group consisting of minerals, vitamins, bisphosphonates, anabolic steroids, parathyroid hormone or analogs, and cathepsin K inhibitors dronabinol.

[00139] In any of the aforementioned aspects involving the prevention or treatment of inflammation are further embodiments comprising: (a) monitoring inflammation in a mammal; (b) measuring bronchoconstriction in a mammal; (c) measuring eosinophil and/or basophil and/or dendritic cell and/or neutrophil and/or monocyte and/or lymphocyte recruitment in a mammal; (d) monitoring mucosal secretion in a mammal; (e) measuring mucosal edema in a mammal; (e) measuring levels of LTB₄ in the calcium ionophore-challenged blood of a mammal; (f) measuring levels of LTE₄ in the urinary excretion of a mammal; or (g) identifying a patient by measuring leukotriene-driven inflammatory biomarkers such as LTB₄, LTC₄, IL-6, CRP, SAA, MPO, EPO, MCP-1, MIP-α, sICAMs, IL-4, IL-13.

[00140] In any of the aforementioned aspects involving the prevention or treatment of leukotriene-dependent or leukotriene mediated diseases or conditions are further embodiments comprising identifying patients by screening for a leukotriene gene haplotype. In further or alternative embodiments the leukotriene gene haplotype is a leukotriene pathway gene, while in still further or alternative embodiments, the leukotriene gene haplotype is a 5-lipoxygenase-activating protein (FLAP) haplotype.

[00141] In any of the aforementioned aspects involving the prevention or treatment of leukotriene-dependent or leukotriene mediated diseases or conditions are further embodiments comprising identifying patients by monitoring the patient for either:

- i) at least one leukotriene related inflammatory biomarker; or
- ii) at least one functional marker response to a leukotriene modifying agent; or
- iii) at least one leukotriene related inflammatory biomarker and at least one functional marker response to a leukotriene modifying agent.

5 In further or alternative embodiments, the leukotriene-related inflammatory biomarkers are selected from the group consisting of LTB₄, cysteinyl leukotrienes, CRP, SAA, MPO, EPO, MCP-1, MIP- α , sICAM, IL-6, IL-4, and IL-13, while in still further or alternative embodiments, the functional marker response is significant lung volume (FEV1).

[00142] In any of the aforementioned aspects involving the prevention or treatment of leukotriene-dependent or
10 leukotriene mediated diseases or conditions are further embodiments comprising identifying patients by either:

- i) screening the patient for at least one leukotriene gene SNP and/or haplotype including SNP's in intronic or exonic locations; or
- ii) monitoring the patient for at least one leukotriene related inflammatory biomarker; or
- ii) monitoring the patient for at least one functional marker response to a leukotriene modifying agent

15 In further or alternative embodiments, the leukotriene gene SNP or haplotype is a leukotriene pathway gene. In still further or alternative embodiments, the leukotriene gene SNP or haplotype is a 5-lipoxygenase-activating protein (FLAP) SNP or haplotype. In further or alternative embodiments, the leukotriene-related inflammatory biomarkers are selected from the group consisting of LTB₄, cysteinyl leukotrienes, CRP, SAA, MPO, EPO, MCP-1, MIP- α , sICAM, IL-6, IL-4, and IL-13, while in still further or alternative embodiments, the functional
20 marker response is significant lung volume (FEV1).

[00143] In any of the aforementioned aspects involving the prevention or treatment of leukotriene-dependent or leukotriene mediated diseases or conditions are further embodiments comprising identifying patients by at least two of the following:

- i) screening the patient for at least one leukotriene gene SNP or haplotype;
- 25 ii) monitoring the patient for at least one leukotriene related inflammatory biomarker;
- ii) monitoring the patient for at least one functional marker response to a leukotriene modifying agent.

In further or alternative embodiments, the leukotriene gene SNP or haplotype is a leukotriene pathway gene. In still further or alternative embodiments, the leukotriene gene SNP or haplotype is a 5-lipoxygenase-activating protein (FLAP) SNP or haplotype. In further or alternative embodiments, the leukotriene-related inflammatory
30 biomarkers are selected from the group consisting of LTB₄, cysteinyl leukotrienes, CRP, SAA, MPO, EPO, MCP-1, MIP- α , sICAM, IL-6, IL-4, and IL-13, while in still further or alternative embodiments, the functional marker response is significant lung volume (FEV1).

[00144] In any of the aforementioned aspects involving the prevention or treatment of leukotriene-dependent or leukotriene mediated diseases or conditions are further embodiments comprising identifying patients by:

- 35 i) screening the patient for at least one leukotriene gene SNP or haplotype; and
- ii) monitoring the patient for at least one leukotriene related inflammatory biomarker; and
- ii) monitoring the patient for at least one functional marker response to a leukotriene modifying agent.

In further or alternative embodiments, the leukotriene gene SNP or haplotype is a leukotriene pathway gene. In still further or alternative embodiments, the leukotriene gene SNP or haplotype is a 5-lipoxygenase-activating
40 protein (FLAP) SNP or haplotype. In further or alternative embodiments, the leukotriene-related inflammatory

biomarkers are selected from the group consisting of LTB₄, cysteinyl leukotrienes, CRP, SAA, MPO, EPO, MCP-1, MIP- α , sICAM, IL-6, IL-4, and IL-13, while in still further or alternative embodiments, the functional marker response is significant lung volume (FEV1).

[00145] In another aspect is the prevention or treatment of leukotriene-dependent or leukotriene mediated diseases or conditions comprising administering to a patient an effective amount of a FLAP modulator, wherein the patients has been identified using information obtained by:

- i) screening the patient for at least one leukotriene gene SNP or haplotype; and
- ii) monitoring the patient for at least one leukotriene related inflammatory biomarker; and
- ii) monitoring the patient for at least one functional marker response to a leukotriene modifying agent.

In further or alternative embodiments, the FLAP modulator is a FLAP inhibitor. In further or alternative embodiments, the leukotriene gene SNP or haplotype is a leukotriene pathway gene. In still further or alternative embodiments, the leukotriene gene SNP or haplotype is a 5-lipoxygenase-activating protein (FLAP) SNP or haplotype. In further or alternative embodiments, the leukotriene-related inflammatory biomarkers are selected from the group consisting of LTB₄, cysteinyl leukotrienes, CRP, SAA, MPO, EPO, MCP-1, MIP- α , sICAM, IL-6, IL-4, and IL-13, while in still further or alternative embodiments, the functional marker response is significant lung volume (FEV1). In further or alternative embodiments, the information obtained from the three diagnostic methods may be used in an algorithm in which the information is analyzed to identify patients in need of treatment with a FLAP modulator, the treatment regimen, and the type of FLAP modulator used.

[00146] In any of the aforementioned aspects the leukotriene-dependent or leukotriene mediated diseases or conditions include, but are not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, and endotoxic shock.

[00147] Other objects, features and advantages of the methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description. All references cited herein, including patents, patent applications, and publications, are hereby incorporated by reference in their entirety.

BRIEF DESCRIPTION OF THE FIGURES

[00148] FIG. 1 presents illustrative schemes for the syntheses of compounds described herein.

[00149] FIG. 2 presents illustrative schemes for the syntheses of compounds described herein.

[00150] FIG. 3 presents illustrative schemes for the syntheses of compounds described herein.

[00151] FIG. 4 presents illustrative schemes for the syntheses of compounds described herein.

[00152] FIG. 5 presents illustrative schemes for the syntheses of compounds described herein.

[00153] FIG. 6 presents illustrative schemes for the syntheses of compounds described herein.

[00154] FIG. 7 presents illustrative schemes for the syntheses of compounds described herein.

[00155] FIG. 8 presents illustrative examples of compounds described herein.

[00156] FIG. 9 presents illustrative examples of compounds described herein.

[00157] FIG. 10 presents illustrative examples of compounds described herein.

[00158] FIG. 11 presents illustrative examples of compounds described herein.

[00159] FIG. 12 present an illustrative scheme for the treatment of patients using the compounds and methods described herein.

[00160] FIG. 13 present an illustrative scheme for the treatment of patients using the compounds and methods described herein.

5 [00161] FIG. 14 present an illustrative scheme for the treatment of patients using the compounds and methods described herein.

DETAILED DESCRIPTION OF THE INVENTION

[00162] The MAPEG (membrane associated proteins involved in eicosanoid and glutathione metabolism) family of proteins, include 5-lipoxygenase activating protein (FLAP), leukotriene C₄ synthase (LTC₄ synthase),
10 microsomal glutathione S-transferase 1 (MGST1), MGST2, and MGST3, and microsomal prostaglandin (PG) E synthase 1 (mPGES-1). Members of the MAPEG family of proteins are involved in the lipoxygenase and cyclooxygenase metabolic pathways.

[00163] There are four families of eicosanoids—the prostaglandins, prostacyclins, the thromboxanes and the leukotrienes. Leukotrienes are biological compounds formed from arachidonic acid in the leukotriene synthesis
15 pathway, which include FLAP and LTC₄ synthase. Arachidonic acid may also be transformed to prostaglandin H₂ (PGH₂) by the action of cyclooxygenase enzymes (COX-1 and COX-2) (prostaglandin endoperoxide synthase systems). Prostaglandin H₂ (PGH₂) is further metabolized to other eicosanoids, such as, PGE₂, PGF_{2α}, PGD₂, prostacyclin and thromboxane A₂. PGE₂ is formed by the action of PGES, a member of the MAPEG family.

[00164] Leukotrienes (LTs) are potent contractile and inflammatory mediators produced by release of
20 arachidonic acid from cell membranes and conversion to leukotrienes by the action of 5-lipoxygenase, 5-lipoxygenase activating protein, LTA₄ hydrolase and LTC₄ synthase. The leukotriene synthesis pathway, or 5-lipoxygenase pathway, involves a series of enzymatic reactions in which arachidonic acid is converted to leukotriene LTB₄, or the cysteinyl leukotrienes, LTC₄, LTD₄, and LTE₄. The pathway occurs mainly at the nuclear envelope and has been described. *See, e.g.,* Wood, JW *et al*, *J. Exp. Med.*, 178: 1935-1946, 1993; Peters-
25 Golden, *Am. J. Respir. Crit. Care Med.* 157:S227-S232,1998; Drazen, *et al.*, ed. *Five-Lipoxygenase Products in Asthma*, Lung Biology in Health and Disease Series, Vol. 120, Chs. 1, 2, and 7, Marcel Dekker, Inc. NY, 1998. Protein components dedicated to the leukotriene synthesis pathway include 5-lipoxygenase (5-LO), 5-lipoxygenase-activating protein, LTA₄ hydrolase, and LTC₄ synthase. The synthesis of leukotrienes has been described in the literature, e.g., by Samuelsson *et al*, *Science*, 220, 568-575, 1983; Peters-Golden, "Cell Biology
30 of the 5-Lipoxygenase Pathway" *Am J Respir Crit Care Med* 157:S227-S232 (1998). Leukotrienes are synthesized directly from arachidonic acid by different cells including eosinophils, neutrophils, basophils, lymphocytes, macrophages, monocytes and mast cells. Excess LTA₄, for example from an activated neutrophil, may enter a cell by a transcellular pathway. Most cells in the body have LTA₄ hydrolase so can produce LTB₄. Platelets and endothelial cells have LTC₄ synthase, so can make LTC₄ when presented with LTA₄ by a
35 transcellular pathway.

[00165] Arachidonic acid is a polyunsaturated fatty acid and is present mainly in the membranes of the body's cells. Upon presentation of inflammatory stimuli from the exterior of the cell, calcium is released and binds to phospholipase A₂ (PLA₂) and 5-LO. Cell activation results in the translocation of PLA₂ and 5-LO from the cytoplasm to the endoplasmic reticulum and/or nuclear membranes, where in the presence of FLAP, a 18 kDa
40 integral perinuclear membrane protein that presents the arachidonic acid released from PLA₂ to 5-LO. 5-LO

catalyzes the oxidation of arachidonic acid via a 5-HPETE intermediate to the epoxide LTA₄. Depending on the cell type, LTA₄ may be immediately converted to LTC₄ by the nuclear-bound LTC₄ synthase or to LTB₄ by the action of cytosolic LTA₄ hydrolase. LTB₄ is exported from cells by an as yet uncharacterized transporter and may activate other cells, or the cell it was made in, via high affinity binding to one of two G protein-coupled receptors (GPCRs), namely BLT₁R or BLT₂R. LTC₄ is exported to the blood via the MRP-1 anion pump and rapidly converted to LTD₄ by the action of γ -glutamyl transpeptidase and LTD₄ is then converted to LTE₄ by the action of dipeptidases. LTC₄, LTD₄ and LTE₄ are collectively referred to as the cysteinyl leukotrienes (or previously as slow reacting substance of anaphylaxis, SRS-A). The cysteinyl leukotrienes activate other cells, or the cells they are made in, via high affinity binding to one of two GPCRs, namely CysLT₁R or CysLT₂R. CysLT₁ receptors are found in the human airway eosinophils, neutrophils, macrophages, mast cells, B-lymphocytes and smooth muscle and induce bronchoconstriction. Zhu *et al*, *Am J Respir Cell Mol Biol Epub* Aug 25 (2005). CysLT₂ receptors are located in human airway eosinophils, macrophages, mast cells the human pulmonary vasculature Figueroa *et al*, *Clin Exp Allergy* 33:1380-1388 (2003). Thus, LTC₄ synthase plays a pivotal role in the formation of the cysteinyl leukotrienes.

Involvement of Leukotrienes in Diseases or Conditions

[00166] The involvement of leukotrienes in disease is described in detail in the literature. See e.g., by Busse, *Clin. Exp. Allergy* 26:868-79, 1996; O'Byrne, *Chest* 111(Supp. 2): 27S-34S, 1977; Sheftell, F.D., *et al.*, *Headache*, 40:158-163, 2000; Klickstein *et al.*, *J. Clin. Invest.*, 66:1166-1170, 1950; Davidson *et al.*, *Ann. Rheum. Dis.*, 42:677-679, 1983 Leukotrienes produce marked inflammatory responses in human skin. Evidence for the involvement of leukotrienes in a human disease is found in psoriasis, in which leukotrienes have been detected in psoriatic lesions (Kragballe *et al.*, *Arch. Dermatol.*, 119:548-552, 1983).

[00167] For example, inflammatory responses have been suggested to reflect three types of changes in the local blood vessels. The primary change is an increase in vascular diameter, which results in an increase in local blood flow and leads to an increased temperature, redness and a reduction in the velocity of blood flow, especially along the surfaces of small blood vessels. The second change is the activation of endothelial cells lining the blood vessel to express adhesion molecules that promote the binding of circulating leukocytes. The combination of slowed blood flow and induced adhesion molecules allows leukocytes to attach to the endothelium and migrate into the tissues, a process known as extravasation. These changes are initiated by cytokines and leukotrienes produced by activated macrophages. Once inflammation has begun, the first cells attracted to the site of infection are generally neutrophils. They are followed by monocytes, which differentiate into more tissue macrophages. In the latter stages of inflammation, other leukocytes, such as eosinophils and lymphocytes also enter the infected site. The third major change in the local blood vessels is an increase in vascular permeability. Instead of being tightly joined together, the endothelial cells lining the blood vessel walls become separated, leading to exit of fluid and proteins from the blood and their local accumulation in the tissue. (See Janeway, *et al.*, *Immunobiology: the immune system in health and disease*, 5th ed., Garland Publishing, New York, 2001)

[00168] LTB₄ produces relatively weak contractions of isolated trachea and lung parenchyma, and these contractions are blocked in part by inhibitors of cyclooxygenase, suggesting that the contraction are secondary to the release of prostaglandins. However, LTB₄ has been shown to be a potent chemotactic agent for eosinophils and progenitors of mast cells and the LTB₄ receptor BLT1^{-/-} knockout mouse is protected from eosinophilic

inflammation and T-cell mediated allergic airway hyperreactivity. Miyahara *et al. J Immunol* 174:4979-4784; (Weller *et al. J Exp Med* 201:1961-1971(2005).

[00169] Leukotrienes C₄ and D₄ are potent smooth muscle contractile agents, promoting bronchoconstriction in a variety of species, including humans (Dahlen *et al., Nature*, 288:484-486, 1980). These compounds have profound hemodynamic effects, constricting coronary blood vessels, and resulting in a reduction of cardiac output efficiency (Marone *et al., in Biology of Leukotrienes*, ed. By R. Levi and R.D. Krell, Ann. New York Acad. Sci. 524:321-333, 1988). Leukotrienes also act as vasoconstrictors, however, marked differences exist for different vascular beds. There are reports suggesting that leukotrienes contribute to cardiac reperfusion injury following myocardial ischemia (Barst and Mullane, *Eur. J. Pharmacol.*, 114: 383-387, 1985; Sasaki *et al., Cardiovasc. Res.*, 22: 142-148, 1988). LTC₄ and LTD₄ directly increase vascular permeability probably by promoting retraction of capillary endothelial cells via activation of the CysLT₂ receptor and possibly other as yet undefined CysLT receptors [Lotzer *et al. Arterioscler Thromb Vasc Biol* 23: e32-36.(2003)]. LTB₄ enhances atherosclerotic progression in two atherosclerotic mouse models, namely low density receptor lipoprotein receptor deficient (LDLr^{-/-}) and apolipoprotein E-deficient (ApoE^{-/-}) mice (Aiello *et al, Arterioscler Thromb Vasc Biol* 22:443-449 (2002); Subbarao *et al, Arterioscler Thromb Vasc Biol* 24:369-375 (2004); Heller *et al. Circulation* 112:578-586 (2005). LTB₄ has also been shown to increase human monocyte chemoattractant protein (MCP-1) a known enhancer of atherosclerotic progression (Huang *et al. Aterioscler Thromb Vasc Biol* 24:1783-1788 (2004).

[00170] The role of FLAP in the leukotriene synthesis pathway is significant because FLAP in concert with 5-lipoxygenase performs the first step in the pathway for the synthesis of leukotrienes. Therefore the leukotriene synthesis pathway provides a number of targets for compounds useful in the treatment of leukotriene-dependent or leukotriene mediated diseases or conditions, including, by way of example, vascular and inflammatory disorders, proliferative diseases, and non-cancerous disorders. Compounds that are inhibitors of proteins involved in leukotriene synthesis, such as FLAP, are useful in the treatment of leukotriene-dependent or leukotriene mediated diseases or conditions.

[00171] Leukotriene-dependent or leukotriene mediated conditions treated using the methods, compounds, pharmaceutical compositions and medicaments described herein, include, but are not limited to, bone diseases and disorder, cardiovascular diseases and disorders, inflammatory diseases and disorders, dermatological diseases and disorders, ocular diseases and disorders, cancer and other proliferative diseases and disorders, respiratory diseases and disorder, and non-cancerous disorders.

Treatment Options

[00172] Leukotrienes are known to contribute to the inflammation of the airways of patients with asthma. CysLT₁ receptor (CysLT₁) antagonists such as montelukast (SingulairTM) have been shown to be efficacious in asthma and allergic rhinitis [Reiss *et al. Arch Intern Med* 158:1213-1220 (1998); Phillip *et al. Clin Exp Allergy* 32:1020-1028 (2002)]. CysLT₁R antagonists pranlukast (OnonTM) and zafirlukast (AccolateTM) have also been shown to be efficacious in asthma.

[00173] A number of drugs have been designed to inhibit leukotriene formation, including the 5-lipoxygenase inhibitor zileuton (ZyfloTM) that has shown efficacy in asthma, Israel *et al. Ann Intern Med* 119:1059-1066 (1993). The 5-lipoxygenase inhibitor ZD2138 showed efficacy in inhibiting the fall of FEV1 resulting from aspirin-induced asthma, Nasser *et al, Thorax*, 49; 749-756 (1994). The following leukotriene synthesis inhibitors have shown efficacy in asthma: MK-0591, a specific inhibitor of 5-lipoxygenase-activating protein (FLAP),

Brideau, *et al.*, *Ca. J. Physiol. Pharmacol.* 70:799-807 (1992)., MK-886, a specific inhibitor of 5-lipoxygenase-activating protein (FLAP), Friedman *et al.* *Am Rev Respir Dis.*, 147: 839-844 (1993), and BAY X1005, a specific inhibitor of 5-lipoxygenase-activating protein (FLAP), Fructmann *et al.*, *Agents Actions* 38: 188-195 (1993).

[00174] FLAP inhibition will decrease LTB₄ from monocytes, neutrophils and other cells involved in vascular inflammation and thereby decrease atherosclerotic progression. The FLAP inhibitor MK-886 has been shown to decrease the postangioplasty vasoconstrictive response in a porcine carotid injury model Provost *et al.* *Brit J Pharmacol* 123: 251-258 (1998). MK-886 has also been shown to suppress femoral artery intimal hyperplasia in a rat photochemical model of endothelial injury Kondo *et al.* *Thromb Haemost* 79:635-639 (1998). The 5-lipoxygenase inhibitor zileuton has been shown to reduce renal ischemia in a mouse model, Nimesh *et al.* *Mol Pharm* 66:220-227 (2004).

[00175] FLAP modulators have been used for the treatment of a variety of diseases or conditions, including, by way of example only, (i) inflammation (see e.g. Leff AR *et al.*, "Discovery of leukotrienes and the development of antileukotriene agents", *Ann Allergy Asthma Immunol* 2001;86 (Suppl 1)4-8; Riccioni G, *et al.*, "Advances in therapy with antileukotriene drugs", *Ann Clin Lab Sci.* 2004, 34(4):379-870; (ii) respiratory diseases including asthma, adult respiratory distress syndrome and allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, seasonal asthma (see e.g. Riccioni *et al.*, *Ann. Clin. Lab. Sci.*, v34, 379-387 (2004)); (iii) chronic obstructive pulmonary disease, including chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation and cystic fibrosis (see e.g. Kostikas K *et al.*, "Leukotriene V4 in exhaled breath condensate and sputum supernatant in patients with COPD and asthma", *Chest* 2004;127:1553-9); (iv) increased mucosal secretion and/or edema in a disease or condition (see e.g. Shahab R *et al.*, "Prostaglandins, leukotrienes, and perennial rhinitis", *J Laryngol Otol.*, 2004;118;500-7); (v) vasoconstriction, atherosclerosis and its sequelae myocardial ischemia, myocardial infarction, aortic aneurysm, vasculitis and stroke (see e.g. Jala *et al.*, *Trends in Immunol.*, v25, 315-322 (2004) and Mehrabian *et al.*, *Curr. Opin. Lipidol.*, v14, 447-457 (2003)); (vi) reducing organ reperfusion injury following organ ischemia and/or endotoxic shock (see e.g. Matsui N, *et al.*, "Protective effect of the 5-lipoxygenase inhibitor ardisiaquinone A on hepatic ischemia-reperfusion injury in rats", *Planta Med.* 2005 Aug;71(8):717-20); (vii) reducing the constriction of blood vessels (see e.g. Stanke-Labesque F *et al.*, "Inhibition of leukotriene synthesis with MK-886 prevents a rise in blood pressure and reduces noradrenaline-evoked contraction in L-NAME-treated rats", *Br J Pharmacol.* 2003 Sep;140(1):186-94); (viii) lowering or preventing an increase in blood pressure (see e.g. Stanke-Labesque F *et al.*, "Inhibition of leukotriene synthesis with MK-886 prevents a rise in blood pressure and reduces noradrenaline-evoked contraction in L-NAME-treated rats", *Br J Pharmacol.* 2003 Sep;140(1):186-94, and Walch L, *et al.*, "Pharmacological evidence for a novel cysteinyl-leukotriene receptor subtype in human pulmonary artery smooth muscle", *Br J Pharmacol.* 2002 Dec; 137(8):1339-45); (ix) preventing eosinophil and/or basophil and/or dendritic cell and/or neutrophil and/or monocyte recruitment (see e.g. Miyahara N, *et al.*, "Leukotriene B4 receptor-1 is essential for allergen-mediated recruitment of CD8+ T cells and airway hyperresponsiveness", *Immunol.* 2005 Apr 15;174(8):4979-84); (x) abnormal bone remodeling, loss or gain, including osteopenia, osteoporosis, Paget's disease, cancer and other diseases (see e.g. Anderson GI, *et al.*, "Inhibition of leukotriene function can modulate particulate-induced changes in bone cell differentiation and

activity", *Biomed Mater Res.* 2001;58(4):406-140; (xi) ocular inflammation and allergic conjunctivitis, vernal keratoconjunctivitis, and papillary conjunctivitis (see e.g. Lambiase *et al*, *Arch. Ophthalmol.*, v121, 615-620 (2003)); (xii) CNS disorders, including, but are not limited to, multiple sclerosis, Parkinson's disease, Alzheimer's disease, stroke, cerebral ischemia, retinal ischemia, post-surgical cognitive dysfunction, migraine (see e.g. de Souza Carvalho D, *et al.*, "Asthma plus migraine in childhood and adolescence: prophylactic benefits with leukotriene receptor antagonist", *Headache*. 2002 Nov-Dec;42(10):1044-7; Sheftell F, *et al.*, "Montelukast in the prophylaxis of migraine: a potential role for leukotriene modifiers", *Headache*. 2000 Feb;40(2):158-63); (xiii) peripheral neuropathy/neuropathic pain, spinal cord injury (see e.g. Akpek EA, *et al.*, "A study of adenosine treatment in experimental acute spinal cord injury. Effect on arachidonic acid metabolites", *Spine*. 1999 Jan 15;24(2):128-32), cerebral edema and head injury; (xiv) cancer, including, but is not limited to, pancreatic cancer and other solid or hematological tumors, (see e.g. Poff and Balazy, *Curr. Drug Targets Inflamm. Allergy*, v3, 19-33 (2004) and Steele *et al*, *Cancer Epidemiology & Prevention*, v8, 467-483 (1999); (xv) endotoxic shock and septic shock (see e.g. Leite MS, *et al.*, "Mechanisms of increased survival after lipopolysaccharide-induced endotoxic shock in mice consuming olive oil-enriched diet", *Shock*. 2005 Feb;23(2):173-8); (xvi) rheumatoid arthritis and osteoarthritis (see e.g. Alten R, *et al.*, "Inhibition of leukotriene B₄-induced CD11B/CD18 (Mac-1) expression by BIIL 284, a new long acting LTB₄ receptor antagonist, in patients with rheumatoid arthritis", *Ann Rheum Dis*. 2004 Feb;63(2):170-6); (xvii) preventing increased GI diseases, including, by way of example only, chronic gastritis, eosinophilic gastroenteritis, and gastric motor dysfunction, (see e.g. Gyomber *et al*, *J Gastroenterol Hepatol.*, v11,922-927 (1996); Quack I *et al BMC Gastroenterol* v18,24 (2005); Cuzzocrea S, *et al.*, "5-Lipoxygenase modulates colitis through the regulation of adhesion molecule expression and neutrophil migration", *Lab Invest*. 2005 Jun;85(6):808-22); (xviii) kidney diseases, including, by way of example only, glomerulonephritis, cyclosporine nephrotoxicity renal ischemia reperfusion. (see e.g. Guasch *et al Kidney Int.*,v56, 261-267; Butterly *et al*, v 57, 2586-2593 (2000); Guasch A *et al*. "MK-591 acutely restores glomerular size selectivity and reduces proteinuria in human glomerulonephritis", *Kidney Int*. 1999;56:261-7; Butterly DW *et al*. "A role for leukotrienes in cyclosporine nephrotoxicity", *Kidney Int*. 2000;57:2586-93); (xix) preventing or treating acute or chronic renal insufficiency (see e.g. Maccarrone M, *et al.*, "Activation of 5-lipoxygenase and related cell membrane lipoperoxidation in hemodialysis patients", *J Am Soc Nephrol*. 1999;10:1991-6); (xx) type II diabetes (see e.g. Valdivielso *et al*, v16, 85-94 (2003); (xxi) diminish the inflammatory aspects of acute infections within one or more solid organs or tissues such as the kidney with acute pyelonephritis (see e.g. Tardif M, *et al.*, L-651,392, "A potent leukotriene inhibitor, controls inflammatory process in Escherichia coli pyelonephritis", *Antimicrob Agents Chemother*. 1994 Jul;38(7):1555-60); (xxii) preventing or treating acute or chronic disorders involving recruitment or activation of eosinophils (see e.g. Quack I, *et al*. "Eosinophilic gastroenteritis in a young girl - long term remission under montelukast", *BMC Gastroenterol.*, 2005;5:24; (xxiii) preventing or treating acute or chronic erosive disease or motor dysfunction of the gastrointestinal tract caused by non-steroidal anti-inflammatory drugs (including selective or non-selective cyclooxygenase -1 or -2 inhibitors) (see e.g. Marusova IB, *et al.*, "Potential gastroprotective effect of a CysLT1 receptor blocker sodium montelukast in aspirin-induced lesions of the rat stomach mucosa", *Eksp Klin Farmakol*, 2002;65:16-8 and Gyomber E, *et al.*, "Effect of lipoxygenase inhibitors and leukotriene antagonists on acute and chronic gastric haemorrhagic mucosal lesions in ulcer models in the rat", *J. Gastroenterol. Hepatol.*, 1996, 11, 922-7) and Martin St *et al.*, "Gastric motor dysfunction: is eosinophilic mural gastritis a causative factor?", *Eur J Gastroenterol. Hepatol.*, 2005,

17:983-6; (xxiv) treating type II diabetes (see e.g. Valdivielso JM, *et al.*, "Inhibition of 5-lipoxygenase activating protein decreases proteinuria in diabetic rats", *J Nephrol.* 2003 Jan-Feb;16(1):85-94; Parlapiano C, *et al.*, "The relationship between glycated hemoglobin and polymorphonuclear leukocyte leukotriene B4 release in people with diabetes mellitus", *Diabetes Res Clin Pract.* 1999 Oct;46(1):43-5; (xxv) treatment of metabolic syndromes, including, by way of example only, Familial Mediterranean Fever (see e.g. Bentancur AG, *et al.*, "Urine leukotriene B4 in familial Mediterranean fever", *Clin Exp Rheumatol.* 2004 Jul-Aug;22(4 Suppl 34):S56-8; and (xxvi) treat hepatorenal syndrome (see e.g. Capella GL, "Anti-leukotriene drugs in the prevention and treatment of hepatorenal syndrome", *Prostaglandins Leukot Essent Fatty Acids.* 2003 Apr;68(4):263-5].

[00176] Several inhibitors of FLAP have been described (Gillard *et al.*, *Can. J. Physiol. Pharmacol.*, 67, 456-464, 1989; Evans *et al.*, *Molecular Pharmacol.*, 40, 22-27, 1991; Brideau *et al.*, *Can. J. Physiol. Pharmacol.*, Musser *et al.*, *J. Med. Chem.*, 35, 2501-2524, 1992; Steinhilber, *Curr. Med. Chem.* 6(1):71-85, 1999; Riendeau, *Bioorg Med Chem Lett.*, 15(14):3352-5, 2005; Flamand, *et al.*, *Mol. Pharmacol.* 62(2):250-6, 2002; Folco, *et al.*, *Am. J. Respir. Crit. Care Med.* 161(2 Pt 2):S112-6, 2000; Hakonarson, *JAMA*, 293(18):2245-56, 2005).

Identification of Leukotriene Synthesis Pathway Inhibitors

[00177] The development and testing of novel FLAP inhibitors which are effective either alone or in combination with other drugs, and which result in minimal negative side effects would be beneficial for treating leukotriene-dependent or leukotriene mediated diseases or conditions. Inhibitors of the leukotriene synthesis pathway described herein may target any step of the pathway to prevent or reduce the formation of leukotrienes. Such leukotriene synthesis inhibitors can, by way of example, inhibit at the level of FLAP, thus minimizing the formation of various products in the leukotriene pathway, thereby decreasing the amounts of such compounds available in the cell. Leukotriene synthesis inhibitors can be identified based on their ability to bind to proteins in the leukotriene synthesis pathway. For example, FLAP inhibitors can be identified based on their binding to FLAP.

[00178] FLAP and LTC₄ synthase are two proteins of the MAPEG family that are involved in leukotriene biosynthesis.

[00179] Arachidonic acid is also metabolized to a number of different eicosanoids via cyclooxygenase enzymes (e.g. COX-1, COX-2). Arachidonic acid is metabolized to prostaglandin H₂ (PGH₂) by the action of COX enzymes. PGH₂ is a substrate for a number of different synthases that produce a spectrum of lipid mediators, including PGE₂, PGF_{2α}, PGD₂, prostacyclin and thromboxane A₂.

[00180] PGH₂ is metabolized to PGE₂ by prostaglandin E synthases (PGES). PGES isozymes have been identified: cytosolic PGES (cPGES), microsomal PGES-1 (mPGES-1) and microsomal PGES-2 (mPGES-2). cPGES is constitutively and ubiquitously expressed and selectively expressed with COX-1.

[00181] mPGES-1 catalyzes the formation of PGE₂ from PGH₂. mPGES-1 is induced by proinflammatory stimuli, downregulated by anti-inflammatory glucocorticoids, and functionally coupled with COX-2 in preference to COX-1. mPGES-1 has been shown to be inducible in various models of pain and inflammation, where it appears to be the predominant synthase involved in COX-2 mediated PGE₂ production, both in the peripheral inflamed sites and in the CNS. Mice deficient in mPGES-1 show both a reduction in the production of inflammatory responses in the collagen-induced arthritis model (Trebbino *et al.* *P.N.A.S. USA.* 2003, 100, 9044).

[00182] In another aspect, compounds that inhibit the activity of one of the proteins in MAPEG family of proteins also inhibit the activity of other proteins in the MAPEG family of proteins. In general, structure activity

relationships will be different for FLAP inhibitor compounds described herein compared to inhibitor compounds for other proteins in the MAPEG family of proteins.

[00183] Compounds described herein inhibit the activity of at least one member of the MAPEG family of proteins. In one aspect, compounds described herein inhibit the activity of at least one member of the MAPEG family of proteins selected from among FLAP, LTC₄ synthase, MGST1, MGST2, MGST3, mPGES-1, and combinations thereof. In one aspect, compounds described herein inhibit the activity of at least one member of the MAPEG family of proteins selected from among FLAP, LTC₄ synthase, mPGES-1, and combinations thereof.

[00184] In one aspect, compounds described herein are FLAP inhibitor compounds.


[00185] Compounds described herein inhibit or reduce the formation of metabolites of arachidonic acid, such as leukotrienes and prostaglandins, and thus find use in the treatment of inflammatory diseases or conditions.

Compounds

[00186] Described herein are compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H). Compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), which inhibit the activity of at least one protein from the MAPEG family of proteins. Compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), inhibit the activity of proteins in the MAPEG family of proteins, such as FLAP. In another aspect, compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), inhibit the activity of FLAP and also inhibit the activity of other proteins in the MAPEG family of proteins selected from among LTC₄ synthase and mPGES-1.

[00187] In one embodiment, provided herein is a compound of Formula (G). Compounds of Formula (G), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent or leukotriene mediated conditions or diseases, including, but not limited to, asthma, myocardial infarction, cancer, and inflammatory conditions.

[00188] In another embodiment, provided herein is a compound of Formula (G):



wherein,

Z is selected from S(O)_m, [C(R₂)₂]_nC(R₁)₂S(O)_m, S(O)_mC(R₁)₂[C(R₂)₂]_n, wherein each R₁ is independently H, CF₃, or an optionally substituted C₁-C₆alkyl, or two R₁ on the same carbon may join to form a carbonyl (=O); and each R₂ is independently H, OH, OMe, CF₃, or an optionally substituted C₁-C₆alkyl, or two R₂ on the same carbon may join to form a carbonyl (=O); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;

Y is a (substituted or unsubstituted aryl), or -(substituted or unsubstituted heteroaryl);

R₆ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycloalkyl), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl),

where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, $C(O)$, $-CH(OH)$, $-(\text{substituted or unsubstituted } C_1\text{-}C_6\text{alkyl})$, or $-(\text{substituted or unsubstituted } C_2\text{-}C_6\text{alkenyl})$;

R_7 is $L_3\text{-}X\text{-}L_4\text{-}G_1$, wherein,

L_3 is a or substituted or unsubstituted alkyl;

X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, $-C(O)NR_9$, $-NR_9C(O)NR_9-$;

L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, a substituted or unsubstituted cyclic alkyl, or a substituted or unsubstituted heterocycloalkyl;

G_1 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, -

$S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, -

$NR_9C(=CHR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)C(=O)R_9$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, -

$C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-C(R_9)_2(OR_9)$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, -

$S(=O)_2R_8$, $-L_5\text{-(substituted or unsubstituted alkyl)}$, $-L_5\text{-(substituted or unsubstituted alkenyl)}$, $-L_5\text{-(substituted or unsubstituted heteroaryl)}$, or $-L_5\text{-(substituted or unsubstituted aryl)}$, wherein L_5 is $-OC(O)O-$, $-NHC(O)NH-$, $-NHC(O)O$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

or G_1 is $W\text{-}G_5$, where W is a substituted or unsubstituted aryl, substituted or unsubstituted

heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, -

$NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(R_9)_2(OR_9)$, $-C(O)NHS(=O)_2R_8$, -

$S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, -

$NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, -

$C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$;

each R_8 is independently selected from substituted or unsubstituted $C_1\text{-}C_6\text{alkyl}$, substituted or unsubstituted $C_3\text{-}C_8\text{cycloalkyl}$, substituted or unsubstituted phenyl or substituted or unsubstituted benzyl;

each R_9 is independently selected from H, substituted or unsubstituted $C_1\text{-}C_6\text{alkyl}$, substituted or unsubstituted $C_1\text{-}C_6\text{fluoroalkyl}$, substituted or unsubstituted $C_3\text{-}C_8\text{cycloalkyl}$, substituted or unsubstituted phenyl, substituted or unsubstituted benzyl, and substituted or unsubstituted heteroarylmethyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or R_8 and R_9 can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring and

each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$,

heteroaryl, or heteroalkyl;

R_5 is H, halogen, substituted or unsubstituted $C_1\text{-}C_6\text{alkyl}$, substituted or unsubstituted $-O\text{-}C_1\text{-}C_6\text{alkyl}$;

R_{11} is $L_7\text{-}L_{10}\text{-}G_6$, wherein L_7 is a bond, $-C(O)$, $-C(O)NH$, $-NHC(O)$, or (substituted or unsubstituted $C_1\text{-}C_6\text{alkyl}$); L_{10} is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl);

G_6 is OR_9 , $-C(=O)R_9$, $-C(=O)OR_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, -

$S(=O)_2N(R_9)_2$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=O)N(R_9)_2$, $NR_9C(O)R_9$,

$C(R_9)_2C(=O)N(R_9)_2$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5\text{-(substituted or unsubstituted alkyl)}$, $-L_5\text{-(substituted or unsubstituted alkenyl)}$, $-L_5\text{-(substituted or unsubstituted heteroaryl)}$, or $-L_5\text{-(substituted or unsubstituted aryl)}$, wherein L_5 is $-O-$, $C(=O)$, S,

S(=O), S(=O)₂, -NH-, -NHC(O)O-, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O)-;

or G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl) and G₇ is H, halogen, CN, NO₂, N₃, CF₃, OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -C(R₉)₂(OR₉), -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroalkyl), -L₅-(substituted or unsubstituted heteroaryl), -L₅-(substituted or unsubstituted heterocycloalkyl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is a bond, -O-, C(=O), S, S(=O), S(=O)₂, -NH-, -NHC(O)O-, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); and

R₁₂ is H, (substituted or unsubstituted C₁-C₆ alkyl), (substituted or unsubstituted C₃-C₆ cycloalkyl);

or acyl glucuronide metabolite, or pharmaceutically acceptable solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00189] For any and all of the embodiments, substituents can be selected from among from a subset of the listed alternatives.

[00190] For example, in some embodiments, R₁₁ comprises at least one (unsubstituted or substituted) aromatic moiety and at least one (unsubstituted or substituted) cyclic moiety, wherein the (unsubstituted or substituted) cyclic moiety is a (unsubstituted or substituted) heterocycloalkyl group or a (unsubstituted or substituted) heteroaryl group. In some embodiments, R₁₁ is not a thienyl-phenyl group.

[00191] In some embodiments, Z is selected from S(O)_m, [C(R₂)₂]_nC(R₁)₂S(O)_m, S(O)_mC(R₁)₂[C(R₂)₂]_n. In other embodiments, Z is [C(R₂)₂]_nC(R₁)₂S(O)_m.

[00192] In some embodiments, Z is selected from S(O)_m, [C(R₂)₂]_nC(R₁)₂S(O)_m, and S(O)_mC(R₁)₂[C(R₂)₂]_n, wherein each R₁ is independently H, CF₃, or an optionally substituted C₁-C₆alkyl; and R₂ is H, OH, OMe, CF₃, or an optionally substituted C₁-C₆alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00193] In some embodiments, Z is selected from -S-, -[C(R₂)₂]_nC(R₁)₂S-, and -SC(R₁)₂[C(R₂)₂]_n-.

[00194] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00195] In some embodiments, each R₁ is independently H, CF₃, or an optionally substituted C₁-C₆alkyl.

[00196] In some embodiments, each R₂ is independently H, OH, OMe, CF₃, or an optionally substituted C₁-C₆alkyl.

[00197] In some embodiments, Z is -S- or [C(R₂)₂]_nC(R₁)₂S-.

[00198] In some embodiments, Z is [C(R₂)₂]_nC(R₁)₂S-.

[00199] In some embodiments, Z is -S-.

[00200] In some embodiments, Z is CH₂S-.

[00201] In some embodiments, Z is -S-, -SCH₂-, -CH₂S-, or -CH(CH₃)S-

[00202] In some embodiments, Z is -S- or -CH₂S-.

[00203] In further or alternative embodiments, Y is -(substituted or unsubstituted heteroaryl) or -(substituted or unsubstituted aryl) and G₆ is W-G₇.

[00204] In further or alternative embodiments, Y is -(substituted or unsubstituted heteroaryl).

[00205] In further or alternative embodiments, Y is selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl, wherein Y is substituted or unsubstituted.

[00206] In further or alternative embodiments, Y is selected from the group consisting of pyridinyl or quinolinyl, wherein Y is substituted or unsubstituted.

[00207] In further or alternative embodiments, R₆ is L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O)₂, -C(O), or substituted or unsubstituted alkyl.

[00208] In further or alternative embodiments, X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉.

[00209] In further or alternative embodiments, G₁ is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[00210] In further or alternative embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is -C(O)OR₉.

[00211] In further or alternative embodiments, R₉ is H or unsubstituted alkyl.

[00212] In further or alternative embodiments, L₁₀ is a substituted or unsubstituted aryl substituted or unsubstituted heteroaryl and G₆ is W-G₇ wherein W is substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl.

[00213] In further or alternative embodiments, L₁₀ is a substituted or unsubstituted aryl.

[00214] In further or alternative embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is -OR₉.

[00215] In further or alternative embodiments, G₁ is W-G₅, where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl.

[00216] In further or alternative embodiments, G₆ is W-G₇.

[00217] In some embodiments, Y is a substituted or unsubstituted aryl.

[00218] In further or alternative embodiments, Y is -(substituted or unsubstituted heteroaryl).

[00219] In further or alternative embodiments, Y is -(substituted or unsubstituted heteroaryl) and G₆ is W-G₇.

[00220] In further or alternative embodiments, Y is a substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1 O atoms and 0-1 S atoms.

[00221] In further or alternative embodiments, Y is selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl,

benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl, wherein Y is substituted or unsubstituted.

[00222] In further or alternative embodiments, Y is a substituted or unsubstituted heteroaryl containing 1-3 nitrogen atoms.

5 [00223] In further or alternative embodiments, Y is a substituted or unsubstituted group selected from among pyridinyl; benzothiazolyl; thiazolyl; imidazo[1,2-a]pyridinyl; quinolinyl; isoquinolinyl; isoxazolyl; pyrazolyl; indolyl; pyrazinyl; pyridazinyl; pyrimidinyl; quinazolinyl; and quinoxalinyl.

[00224] In further or alternative embodiments, Y is substituted with substituents selected from among H, halogen, -CN, -NO₂, -S(=O)₂NH₂, -OH, -C(O)NH₂, -C(O)OH, -C(O)OCH₃, -C(O)OCH₂CH₃, C₁-C₆ alkyl, -O-C₁-
10 C₆ alkyl, CF₃, OCF₃, heteroaryl, aryl, heterocycloalkyl, and heteroalkyl.

[00225] In further or alternative embodiments, Y is selected from among pyridin-2-yl; 3-fluoro-pyridin-2-yl; 4-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 6-fluoro-pyridin-2-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 6-methyl-pyridin-2-yl; 3,5-dimethylpyridin-2-yl; 5,6-dimethyl-pyridin-2-yl; 5-ethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-chloro-
15 pyridin-2-yl; 5-bromo-pyridin-2-yl; 6-cyclopropyl-pyridin-2-yl; 5-methyl-1-oxy-pyridin-2-yl; N-oxido-pyridin-2-yl; benzothiazol-2-yl; 2-methylthiazol-4-yl; imidazo[1,2-a]pyridin-2-yl; quinolin-2-yl; 6-fluoroquinolin-2-yl; 7-fluoroquinolin-2-yl; 6-methylquinolin-2-yl; 6-bromo-quinolin-2-yl; 1-oxy-quinolin-2-yl; 5-methylisoxazol-3-yl; 1,3-dimethylpyrazol-5-yl; 1,5-dimethylpyrazol-3-yl; 1H-indol-2-yl; 5-methyl-pyrazin-2-yl; 6-methyl-pyridazin-3-yl; quinoxalin-2-yl; quinazolin-2-yl; pyrimidin-2-yl; and 5-methylpyrimidin-2-yl..

20 [00226] In further or alternative embodiments, Y is a substituted or unsubstituted group selected from among pyridinyl and quinolinyl.

[00227] In some embodiments, L₇ is a bond; L₁₀ is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl).

25 [00228] In some other embodiments, L₇ is a bond; L₁₀ is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and G₆ is W-G₇, wherein W is a (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl).

[00229] In some embodiments, L₇ is a bond; L₁₀ is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), or a
30 (substituted or unsubstituted heteroaryl).

[00230] In some embodiments, L₁₀ is selected from among phenyl and pyridinyl.

[00231] In further or alternative embodiments, L₁₀ is a substituted or unsubstituted aryl. In yet some other embodiments, L₁₀ is a substituted or unsubstituted phenyl.

[00232] In some embodiments, L₁₀ is pyridinyl.

35 [00233] In some embodiments, G₇ is H, halogen, CN, NO₂, N₃, CF₃, OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[00234] In further or alternative embodiments, W is a (substituted or unsubstituted heterocycloalkyl containing 0-2 nitrogen atoms, 0-1 O atoms and 0-1 S atoms) or a (substituted or unsubstituted heteroaryl containing 0-4
40 nitrogen atoms, 0-1 O atoms and 0-1 S atoms).

[00235] In further or alternative embodiments, W is substituted with substituents selected from among H, halogen, -CN, -NO₂, -S(=O)₂NH₂, -OH, -C(O)NH₂, -NH₂, -NMe₂, -NHC(O)CH₃, -C(O)OH, -C(O)OCH₃, -C(O)OCH₂CH₃, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, CF₃, OCF₃, heteroaryl, aryl, heterocycloalkyl, and heteroalkyl.

[00236] In further or alternative embodiments, W is substituted with substituents selected from among H, halogen, -CN, -NO₂, -S(=O)₂NH₂, -OH, -C(O)NH₂, -NH₂, -NMe₂, -NHC(O)CH₃, -C(O)OH, -C(O)OCH₃, -C(O)OCH₂CH₃, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, CF₃, OCF₃, and heteroalkyl.

[00237] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl, furopyridinyl, quinolizine, dioxinyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, oxazinanonyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranyl, tetrahydropyranyl, dihydrooxazolyl, oxiranyl, pyrrolidinyl, pyrazolidinyl, dihydrothiophenonyl, imidazolidinonyl, pyrrolidinonyl, dihydrofuranonyl, dioxolanonyl, thiazolidinyl, piperidinonyl, tetrahydronaphthyridinyl, tetrahydroquinolinyl, tetrahydrothiophenyl, indoliny, tetrahydroquinolinyl, and thiazepanyl.

[00238] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, oxadiazolyl, thiadiazolyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, pyrazolidinyl, dioxolanonyl, and thiazolidinyl.

[00239] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; tetrazolyl; tetrahydropyranyl, and morpholin-4-yl.

[00240] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl containing 1-4 nitrogen atoms.

[00241] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl.

[00242] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, thiadiazolyl, and furazanyl.

[00243] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; and tetrazolyl.

[00244] In further or alternative embodiments, G₆ is selected from among pyridin-2-yl; pyridin-3-yl; pyridin-4-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 3-methoxy-pyridin-2-yl; 4-methoxy-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 6-ethoxy-pyridin-2-yl; 3-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 3-trifluoromethyl-pyridin-2-yl; 4-trifluoromethyl-pyridin-2-yl; 5-trifluoromethyl-pyridin-2-yl; 6-trifluoromethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-fluoromethyl-pyridin-2-yl; 5-methoxymethyl-pyridin-2-yl; 5-hydroxymethyl-pyridin-2-yl; 2-methyl-pyridin-3-yl; 6-methyl-pyridin-3-yl; 6-cyano-pyridin-3-yl; 2-methoxy-pyridin-3-yl; 5-methoxy-pyridin-3-yl; 5-fluoro-pyridin-3-yl; 6-carbamoyl-pyridin-3-yl; 6-hydroxy-pyridin-3-yl; 6-methoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-bromo-6-methoxy-pyridin-3-yl; 6-trifluoromethyl-pyridin-3-yl; 6-trifluoromethyl-pyridin-4-yl; 2-trifluoromethyl-pyridin-5-yl; 2-acetyl-amino-pyridin-5-yl; pyrazin-2-yl; pyrimidin-2-yl; pyrimidin-5-yl; 5-amino-pyrazin-2-yl; 1,3,4-oxadiazol-2-ylamine; 6-hydroxy-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-methyl-pyridazin-3-yl; 2-methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl; thiazol-2-yl; 5-methyl-thiazol-2-yl; 5-fluoro-thiazol-2-yl; 5-trifluoromethyl-thiazol-2-yl; 2,4-dimethyl-thiazol-5-yl; 5-methoxy-thiazol-2-yl; 2-methoxy-thiazol-4-yl; 2-ethoxy-thiazol-4-yl; 2-methyl-thiazol-4-yl; 2-methyl-thiazol-5-yl; 4-methyl-thiazol-2-yl; isoxazol-4-yl; 3,5-dimethyl-isoxazol-4-yl; 2-methyl-imidazol-4-yl; 1-methyl-imidazol-5-yl; 1-methyl-imidazol-4-yl; imidazol-4-yl; 4-methyl-imidazol-5-yl; pyrazol-4-yl; 1-methyl-pyrazol-4-yl; 3-methyl-pyrazol-4-yl; 5-methyl-1,2,4-oxadiazol-3-yl; 2-methyl-1,3,4-oxadiazol-5-yl; 1,3,4-oxadiazol-2-yl; 1,3,4-thiadiazol-2-yl; 3-methyl-pyrazol-5-yl; 1,2,3-thiadiazol-4-yl; tetrazol-1-yl; tetrazol-2-yl; 1-methyl-tetrazol-5-yl; 2-methyl-tetrazol-5-yl; 4-methyl-1*H*-imidazol-2-yl; 5-hydroxy-pyrimidin-2-yl; 2-methoxy-pyrimidin-5-yl; 6-methyl-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-ethoxy-pyridazin-3-yl; 3-methoxy-pyridazin-6-yl; 4-methoxy-tetrahydro-pyran-4-yl; 6-ethoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-fluoro-pyridin-2-yl, and morpholin-4-yl.

[00245] In further or alternative embodiments, R₆ is L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O), -S(O)₂, -C(O), -CR₉(OR₉), or substituted or unsubstituted alkyl.

[00246] In further or alternative embodiments, R₆ is H, L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O)₂, -S(O)-, -C(O), or substituted or unsubstituted alkyl.

[00247] In further or alternative embodiments, R₆ is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00248] In further or alternative embodiments, R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; ; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-

methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; tert-butylsulfanyl; tert-butyl-sulfinyl; or tert-butylsulfonyl.

[00249] In further or alternative embodiments, R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00250] In further or alternative embodiments, R_6 is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

[00251] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00252] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

[00253] In further or alternative embodiments, R_6 is *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00254] In further or alternative embodiments, R_6 is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00255] In further or alternative embodiments, R_6 is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00256] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00257] In further or alternative embodiments, X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, or $-C(O)NR_9$.

[00258] In further or alternative embodiments, X is a bond or $-CR_9(OR_9)$.

[00259] In further or alternative embodiments, X is a bond.

[00260] In further or alternative embodiments, R_9 is H, C_1 - C_6 alkyl, benzyl, or heteroaryl methyl.

[00261] In further or alternative embodiments, R_9 is H or C_1 - C_6 alkyl.

[00262] In further or alternative embodiments, R_9 is H.

[00263] In further or alternative embodiments, G_1 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-N(R_9)CH_2CO_2R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)C(=O)R_9$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-C(R_9)_2(OR_9)$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -

(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-OC(O)O-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_1 is $W-G_5$, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(R_9)_2(OR_9)$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$.

[00264] In further or alternative embodiments, G_1 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$.


[00265] In further or alternative embodiments, G_1 is $-OR_9$, $N(R_9)_2$, $-CO_2R_9$, $-CON(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$.


[00266] In further or alternative embodiments, G_1 is $W-G_5$, where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl. In further or alternative embodiments, G_1 is $W-G_5$, where W is a (substituted or unsubstituted heterocycloalkyl containing 0-1 O atoms and 0-2 N atoms), or (substituted or unsubstituted heteroaryl containing 0-4 N atoms).

[00267] In further or alternative embodiments, G_1 is $W-G_5$, where W is a substituted or unsubstituted group selected from among furanonyl, dihydrofuran-2-onyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, imidazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, 1,3,4-oxadiazolyl, thiazolyl, pyrazolyl, tetrazolyl, oxazolyl, or pyrrolyl.

[00268] In further or alternative embodiments, G_1 is selected from among H, OH, CN, CO_2H , CO_2Me , CO_2Et ,


CO_2NH_2 , CO_2NHMe , $CO_2N(Me)_2$, $CO_2N(Et)_2$, $-NH_2$, $-NHMe$, $-N(Me)_2$, $-N(Et)_2$, $-NMe(iPr)$,






[00269] In further or alternative embodiments, G₁ is -OR₉, N(R₉)₂, or -CO₂R₉.

[00270] In further or alternative embodiments, G_1 is selected from among H, OH, CN, CO_2H , CO_2Me , CO_2Et ,



5 CO₂NH₂, CO₂NHMe, CO₂N(Me)₂, CO₂N(Et)₂, -NH₂, -NHMe, -N(Me)₂, -N(Et)₂, -NMe(iPr),



[00271] In further or alternative embodiments, G₁ is selected from among OH, CO₂H, CO₂Me, CO₂Et, CO₂NH₂, CO₂NHMe, CO₂N(Me)₂, and CO₂N(Et)₂.

[00272] In further or alternative embodiments, G_1 is $-OR_0$, or $-CO_2R_0$.

10 [00273] In further or alternative embodiments, G_1 is $-\text{CO}_2\text{R}_9$.

[00274] In further or alternative embodiments, L₃ is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl.

15 **[00275]** In further or alternative embodiments, L₃ is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; 2-methylpropan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

[00276] In further or alternative embodiments, L₃ is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; and G₁ is OR₉, or CO₂R₉.

[00277] In further or alternative embodiments, L₃ is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L₄ is a bond; and G₁ is OR₉, or CO₂R₉.

[00278] In further or alternative embodiments, L₃ is methandiyl; or ethan-1,2-diyl.

[00279] In further or alternative embodiments, L₃ is methandiyl.

[00280] In further or alternative embodiments, L₃ is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

[00281] In further or alternative embodiments, L_3 is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00282] In further or alternative embodiments, L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cyclic alkyl.

[00283] In further or alternative embodiments, L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

[00284] In further or alternative embodiments, L_4 is a bond, methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00285] In further or alternative embodiments, L_4 is a bond, ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00286] In further or alternative embodiments, L_3 is a methandiyl; ethan-1,2-diyl; X is a bond, O , $-C(=O)$, $-CR_9(OR_9)$, S , $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, or $-C(O)NR_9$; L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

[00287] In further or alternative embodiments, L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00288] In further or alternative embodiments, L_3 is methandiyl; X is a bond; L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl;

cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00289] In further or alternative embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is -C(O)OR₉.

5 **[00290]** In further or alternative embodiments, L₃ is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L₄ is a bond; and G₁ is -C(O)OR₉.

[00291] In further or alternative embodiments, L₃ is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, X is a bond; L₄ is a bond; and G₁ is -C(O)OR₉.

[00292] In further or alternative embodiments, L₃ is 2-methyl-propan-1,2-diyl; or 2-ethyl-butan-1,2-diyl; X is a bond; L₄ is a bond; and G₁ is -C(O)OR₉.


15 **[00293]** In further or alternative embodiments, L₃ is unsubstituted alkyl; X is a bond; L₄ is a bond; and G₁ is -OR₉.

[00294] In further or alternative embodiments, L₃ is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L₄ is a bond; and G₁ is -OR₉.


20 **[00295]** In further or alternative embodiments, L₃ is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; X is a bond; L₄ is a bond; and G₁ is -OR₉.

[00296] In further or alternative embodiments, L₃ is 2-methyl-propan-1,2-diyl; 2-ethyl-butan-1,2-diyl; X is a bond; L₄ is a bond; and G₁ is -OR₉.


25 **[00297]** In some embodiments, L₃-X-L₄ is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂C(CH₃)H-, -CH₂C(CH₂CH₃)H-, -CH₂C(isopropyl)H-, -CH₂C(tert-butyl)H-, -CH₂C(CH₃)₂-, -CH₂C(CH₂CH₃)₂-,



[00298] In further or alternative embodiments, L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-CH_2C(CH_2CH_3)H-$, $-CH_2C(CH_3)_2-$, $-CH_2C(CH_2CH_3)_2-$,



[00299] In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_2CH_3)H-$, $-CH_2C(CH_2CH_3)_2-$,








5



[00300] In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_3)_2-$, or $-CH_2C(CH_2CH_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_2CH_3)_2-$.

[00301] In some embodiment, R_7 is selected from among

10





15





Chemical structures of various substituted amine derivatives, including primary, secondary, and tertiary amines, amides, sulfonamides, and heterocyclic compounds, are shown. The structures are arranged in a grid, separated by commas, and include various functional groups such as hydroxyl, amino, nitrile, and sulfonamide groups.

Chemical structures of various substituted cyclohexanecarboxylic acids and esters, showing different alkyl and aryl substituents on the cyclohexane ring.




[00303] In some embodiment, R₇ is selected from among 







5 **[00304]** In some embodiment, R_7 is selected from among




[00305] In some embodiment, R_7 is selected from among








[00306] In some embodiment, R₇ is selected from among



5






[00307] In some embodiment, R₇ is selected from among




10

[00308] In some embodiment, R₇ is selected from among





[00309] In some embodiments, compounds of Formula (G) have a structure selected from among:



Y	G ₆	R ₆
Pyridin-2-yl	1,3,4-Oxadiazol-2-ylamine	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyrimidin-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyrazin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Amino-pyrazin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Thiazol-2-yl	3,3-Dimethyl-butyryl
Pyridin-2-yl	Thiazol-2-yl	H
Pyridin-2-yl	Thiazol-2-yl	Acetyl
Pyridin-2-yl	6-Methoxy-pyridazin-3-yl	H
Pyridin-2-yl	6-Methoxy-pyridazin-3-yl	Acetyl
Pyridin-2-yl	6-Methoxy-pyridazin-3-yl	Ethyl
Pyridin-2-yl	Thiazol-2-yl	3,3-Dimethyl-butyl
Pyridin-2-yl	Thiazol-2-yl	Cyclopropane-carbonyl
Pyridin-2-yl	Thiazol-2-yl	Cyclobutane-carbonyl
Pyridin-2-yl	6-Hydroxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyridin-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Thiazol-2-yl	Cyclobutylmethyl
2-Methylthiazol-4-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-Methylthiazol-4-yl	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-Methylthiazol-4-yl	Thiazol-2-yl	H
2-Methylthiazol-4-yl	Thiazol-2-yl	3,3-Dimethyl-butyryl
2-Methylthiazol-4-yl	6-Methoxy-pyridazin-3-yl	H
2-Methylthiazol-4-yl	6-Methoxy-pyridazin-3-yl	3,3-Dimethyl-Butyryl
Pyridin-2-yl	Thiazol-2-yl	Ethyl
Benzothiazol-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-Methylthiazol-4-yl	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
Benzothiazol-2-yl	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2,4-Dimethyl-thiazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Trifluoromethyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl

Y	G ₆	R ₆
Pyridin-2-yl	2-Methyl-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methyl-thiazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	4-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Isoxazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	3,5-Dimethyl-isoxazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methyl-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1-Methyl-imidazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1-Methyl-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Imidazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	4-Methyl-imidazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyrazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1-Methyl-pyrazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	3-Methyl-pyrazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methyl-1,2,4-oxadiazol-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methyl-1,3,4-oxadiazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1,3,4-Oxadiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1,3,4-Thiadiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	3-Methyl-pyrazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1,2,3-Thiadiazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Tetrazol-1-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Tetrazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	1-Methyl-tetrazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methyl-tetrazol-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Hydroxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Cyano-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Trifluoromethyl-pyridin-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Acetyl-amino-pyridin-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methoxy-pyrimidin-5-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Methoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
3-Fluoro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
3-Fluoro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-Fluoro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Fluoro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Cyano-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methoxy-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethyl-pyridin-2-yl	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	2-Trifluoromethyl-pyridin-5-yl	<i>tert</i> -Butylsulfanyl
5-Ethyl-pyridin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Carbamoyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Cyano-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methoxy-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	4-Methyl-1 <i>H</i> -imidazol-2-yl	<i>tert</i> -Butylsulfanyl

Y	G ₆	R ₆
Pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Carbamoyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Methoxy-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Methyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Cyclopropyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	5-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Chloro-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
3-Methyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
3-Methyl-pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
3,5-Dimethylpyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
3,5-Dimethylpyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Benzothiazol-2-ylm	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Benzothiazol-2-yl	5-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Benzothiazol-2-yl	6-Methoxy-pyridin-3-yl	Cyclobutane-carbonyl
Benzothiazol-2-yl	6-Methoxy-pyridin-3-y	Cyclobutylmethyl
5-Ethylpyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	2-Methoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	Cyclobutylmethyl
5-Methylpyridin-2-yl	6-Methoxy-pyridin-3-yl	Cyclobutylmethyl
5-Methylpyridin-2-yl	6-Methoxy-pyridin-3-yl	Isobutyl
Quinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
7-Fluoroquinolin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
7-Fluoroquinolin-2-ylmethyl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
7-Fluoroquinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
7-Fluoroquinolin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Ethyl-pyridin-2-yl	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Methoxyt-hiazol-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	3-Methoxy-pyridazin-6-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl

Y	G ₆	R ₆
Quinolin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
3-Methylpyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
3-Methylpyridin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
3,5-Dimethylpyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-Methylpyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-Methylpyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-Methylpyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	Cyclobutylmethyl
6-Fluoroquinolin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Methylquinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Methylquinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Methoxy-pyridin-3-yl	Isobutyl
6-Fluoroquinolin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	2-Methyl-propane-2-sulfonyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	2-Methyl-propane-2-sulfinyl
N-Oxido-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Imidazo[1,2- <i>a</i>]pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Imidazo[1,2- <i>a</i>]pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Imidazo[1,2- <i>a</i>]pyridin-2-yl	5-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-ylmethyl	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methylisoxazol-3-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methylisoxazol-3-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methylisoxazol-3-yl	5-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
1,3-Dimethylpyrazol-5-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
1,5-Dimethylpyrazol-3-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Trifluoromethyl-pyridin-3-yl	Isobutyl
Pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	4-Methoxy-tetrahydro-pyran-4-yl	<i>tert</i> -Butylsulfanyl
6-Fluoroquinolin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl

Y	G ₆	R ₆
6-Fluoroquinolin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methylpyridin-2-yl	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Ethylpyridin-2-yl	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Bromo-pyridin-2-yl	5-Bromo-6-methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
6-Bromo-quinolin-2-yl	5-Bromo-6-methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	2-Methyl-propane-2-sulfinyl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	2-Methyl-propane-2-sulfinyl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5,6-Dimethyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Carbamoyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methoxy-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
1 <i>H</i> -Indol-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Methoxymethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-ylmethyl	6-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Hydroxymethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	4-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	2-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	3-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	H
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butyl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butyryl
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	2,2-Dimethyl-propionyl
5-Methyl-1-oxy-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
1-Oxy-quinolin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	H
5-Methyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	3,3-Dimethyl-butyryl
5-Methyl-pyridin-2-yl	6-Ethoxy-pyridin-3-yl	Phenylacetyl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	H

Y	G ₆	R ₆
5-Ethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	H
Quinolin-2-yl	5-Fluoro-pyridin-2-yl	3-Methyl-butryl
5-Ethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	3-Methyl-butryl
5-Ethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butryl
5-Ethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	2-Ethyl-butryl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	3-Methyl-butryl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butryl
5,6-Dimethyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	2-Ethyl-butryl
5-Methyl-pyrazin-2-yl	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	Isobutryl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butryl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	Propionyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	Acetyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	3-Methyl-butryl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	2,2,2-Trifluoro-acetyl
Quinoxalin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butyl
Quinoxalin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Quinoxalin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinoxalin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
3-Oxo-3,4-dihydro-quinoxalin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Trifluoromethyl-pyridin-2-yl	3,3-Dimethyl-butryl
5-Methyl-pyrazin-2-yl	5-Trifluoromethyl-pyridin-2-yl	Cyclobutanecarbonyl
6-Methyl-pyridazin-3-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
Pyridin-2-yl	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
Quinolin-2-yl	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
Quinoxalin-2-yl	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyridin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-Methyl-pyrazin-2-yl	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
5-methyl-pyrazin-2-yl	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl

wherein, G₆ is at the 3 or 4 position of the phenyl; Z is -S-, -CH(CH₃) S- or -CH₂S-; and where R₇ is as defined herein.

[00310] In further or alternative embodiments, L₃ is methandiyl; or ethan-1,2-diyl; and L₄ is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl;

cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00311] In further or alternative embodiments, X is a bond; and L₄ is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cyclic alkyl.

[00312] In further or alternative embodiments, L₃ is methandiyl; or ethan-1,2-diyl; X is a bond; and L₄ is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[00313] In further or alternative embodiments, L₃ is methandiyl; X is a bond; and L₄ is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.


[00314] In further or alternative embodiments, L₄ is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and G₁ is -CO₂R₉.

[00315] In further or alternative embodiments, L₇ is a bond; L₁₀ is a substituted or unsubstituted heteroaryl; and G₆ is W-G₇, wherein W is a (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl). In further or alternative embodiments, L₇ is a bond; L₁₀ is a substituted or unsubstituted heteroaryl; and G₆ is W-G₇, wherein W is a (substituted or unsubstituted aryl). In further or alternative embodiments, L₇ is a bond; L₁₀ is a substituted or unsubstituted pyridinyl; and G₆ is W-G₇, wherein W is a (substituted or unsubstituted phenyl). In further or alternative embodiments, L₇ is a bond; L₁₀ is a substituted or unsubstituted heteroaryl; and G₆ is W-G₇, wherein W is a (substituted or unsubstituted heteroaryl). In further or alternative embodiments, L₇ is a bond; L₁₀ is a substituted or unsubstituted pyridinyl; and G₆ is W-G₇, wherein W is a (substituted or unsubstituted pyridinyl).

[00316] In further or alternative embodiments, R₁₁ is selected from among 2-(2-methoxy-pyrid-5-yl)-pyrid-5-yl; 2-(4-methoxy phenyl)-pyrid-5-yl; 2-(4-trifluoromethoxy phenyl)-pyrid-5-yl; 5-(4-methoxy phenyl)-pyrid-2-yl; and 5-(4-trifluoromethoxyphenyl)-pyrid-2-yl.

[00317] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[00318] In some embodiments, compounds of Formula (E) are as follows:



wherein,

wherein, Z is selected from S(O)_m, [C(R₂)₂]_nC(R₁)₂S(O)_m, or S(O)_mC(R₁)₂[C(R₂)₂]_n, wherein each R₁ is independently H, CF₃, or an optionally substituted C₁-C₆alkyl, or two R₁ on the same carbon may join to form a carbonyl (=O); and each R₂ is independently H, OH, OMe, CF₃, or an optionally substituted C₁-

C₆alkyl, or two R₂ on the same carbon may join to form a carbonyl (=O); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;

Y is -L₁-(substituted or unsubstituted heterocycloalkyl), provided that when the heteroatom is directly bound to Z, the heterocycloalkyl is substituted;

where L₁ is a bond, a substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, or substituted or unsubstituted alkynyl, a substituted or unsubstituted heterocycle, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, substituted or unsubstituted heteroalkenyl, or a substituted or unsubstituted heteroalkynyl;

each R₄ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; or two R₄ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;;

R₆ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycloalkyl), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(O), -CH(OH), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₇ is L₃-X-L₄-G₁, wherein,

L₃ is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl;

X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉-, -NR₉S(=O)₂, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉-, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉-, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉-, -OC(=NR₁₀)-, or -C(=NR₁₀)O-;

L₄ is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl;

G₁ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)C(=O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -OC(O)O-, -NHC(O)NH-, -NHC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O);

or G₁ is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(R₉)₂(OR₉), -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈;

each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl;

each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl, benzyl and substituted or unsubstituted heteroarylmethyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and

each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

R_5 is H, halogen, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted O- C_1 - C_6 alkyl;

R_{11} is L_7 - L_{10} - G_6 ; wherein L_7 is a bond, $-O$, $-S$, $-S(=O)$, $-S(=O)_2$, $-NH$, $-C(O)$, $-C(O)NH$, $-NHC(O)$, (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_2 - C_6 alkenyl);

L_{10} is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl), and

G_6 is H, CN, SCN, N_3 , NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-CO_2R_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, (substituted or unsubstituted alkyl), (substituted or unsubstituted fluoroalkyl), $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-NHC(O)NH$, $-OC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

or G_6 is W - G_7 , wherein W is (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is H, halogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_6 fluoroalkyl, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroalkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NH$, $-NHC(O)O$, $-NHC(O)NH$, $-OC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

R_{12} is H, (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_2 - C_4 alkenyl);

or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00319] For any and all of the embodiments of Formula (E), substituents can be selected from among from a subset of the listed alternatives.

[00320] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$. In other embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S(O)_m$.

[00321] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, and $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00322] In some embodiments, Z is selected from $-S-$, $-[C(R_2)_2]_n C(R_1)_2 S-$, and $-SC(R_1)_2 [C(R_2)_2]_n-$.

[00323] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00324] In some embodiments, each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00325] In some embodiments, each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00326] In some embodiments, Z is $-S-$ or $[C(R_2)_2]_n C(R_1)_2 S-$.

[00327] In some embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S-$.


[00328] In some embodiments, Z is $-S-$.

[00329] In some embodiments, Z is CH_2S- .

[00330] In some embodiments, Z is $-S-$, $-SCH_2-$, $-CH_2S-$, or $-CH(CH_3)S-$.

[00331] In some embodiments, Z is $-S-$ or $-CH_2S-$.

[00332] In some embodiments, Y is $-L_1$ -(substituted or unsubstituted heterocycloalkyl). In further or alternative embodiments, the heterocycloalkyl is selected from the group consisting of quinolizinyl, dioxinyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, oxazinanonyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranyl, dihydrooxazolyl, oxiranyl, pyrrolidinyl, pyrazolidinyl, dihydrothiophenonyl, imidazolidinonyl, pyrrolidinonyl, dihydrofuranonyl, dioxolanonyl, thiazolidinyl, piperidinonyl, tetrahydronaphyridinyl, tetrahydroquinolinyl, tetrahydrothienyl, indolinyl, tetrahydroquinolinyl, and thiazepanyl. In further or alternative embodiments, the heterocycloalkyl is selected from the group consisting of:



[00333] In further or alternative embodiments, L_1 is a bond or a substituted or unsubstituted alkyl. In further or alternative embodiments, L_1 is a bond.

[00334] In further or alternative embodiments, Y is morpholin-4-yl; pyrrolidin-2-yl; 2-methyl-1,3-dioxolan-2-yl; pyrrolidin-5-yl; N-methylsulfonyl-pyrrolidin-2-yl; N-trifluoroacetyl-pyrrolidin-2-yl; N-t-butoxycarbonyl-4,5-dihydroimidazol-2-yl; 4,5-dihydroimidazol-2-yl; indolin-2-yl; N-t-butoxycarbonyl-indolin-2-yl; N-acetyl-indolin-2-yl; N-(methoxyacetyl) indolin-2-yl; N-(2-bromoethoxycarbonyl) indolin-2-yl; N-t-butoxycarbonylpyrrolidin-2-yl; N-cyclopropylcarbonyl-pyrrolidin-2-yl; N-benzoyl-pyrrolidin-2-yl; N-(2-methylpropanoyl)-pyrrolidin-2-yl; N-propanoyl-pyrrolidin-2-yl; N-acetyl-pyrrolidin-2-yl; N-(4-phenylbenzoyl)-pyrrolidin-2-yl; N-(phenylacetyl)-pyrrolidin-2-ylmethyl; N-(3-phenylpropanoyl)-pyrrolidin-2-yl; N-(3-phenoxybenzoyl)-pyrrolidin-2-yl; N-(4-phenoxybenzoyl)-pyrrolidin-2-yl; N-(nicotinoyl)-pyrrolidin-2-yl; N-(pyridin-4-ylcarbonyl)-pyrrolidin-2-yl; N-(4-phenylbenzoyl)-pyrrolidin-2-yl; N-(phenylacetyl)-pyrrolidin-2-yl; N-(phenylcyclopropylcarbonyl)-pyrrolidin-2-

yl; N-(4-chlorobenzoyl)-pyrrolidin-2-yl; N-(4-benzyloxyphenylacetyl)-pyrrolidin-2-yl; or N-(tert-butoxycarbonyl)-piperidin-2-yl.

[00335] In further or alternative embodiments, R_6 is L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or substituted or unsubstituted alkyl.

[00336] In further or alternative embodiments, R_6 is H, or L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or substituted or unsubstituted alkyl.

[00337] In further or alternative embodiments, R_6 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00338] In further or alternative embodiments, R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00339] In further or alternative embodiments, R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00340] In further or alternative embodiments, R_6 is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

[00341] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00342] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

[00343] In further or alternative embodiments, R_6 is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00344] In further or alternative embodiments, R_6 is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-

dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00345] In further or alternative embodiments, R_6 is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00346] In further or alternative embodiments, R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00347] In further or alternative embodiments, R_{12} is H and R_{11} is L_7 - L_{10} - G_6 , wherein: L_7 is a bond, (substituted or unsubstituted C_1 - C_6 alkyl); and L_{10} is a (substituted or unsubstituted aryl), (substituted or unsubstituted heteroaryl), or (substituted or unsubstituted heterocycloalkyl). In further or alternative embodiments, L_{10} is a (substituted or unsubstituted aryl).

[00348] In further or alternative embodiments, R_9 is H, C_1 - C_6 alkyl, benzyl, or heteroarylmethyl.

[00349] In further or alternative embodiments, R_9 is H or unsubstituted alkyl.

[00350] In further or alternative embodiments, L_{10} is a (substituted or unsubstituted aryl) or (substituted or unsubstituted heteroaryl). In further or alternative embodiments, L_{10} is phenyl or pyridinyl. In further or alternative embodiments, L_{10} is phenyl. In further or alternative embodiments, L_{10} is pyridinyl.

[00351] In further or alternative embodiments, G_6 is H, CN, NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-CO_2R_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, (substituted or unsubstituted alkyl), (substituted or unsubstituted fluoroalkyl), $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_6 is W - G_7 , wherein W is (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is H, halogen, C_1 - C_6 alkyl, $-C_1$ - C_6 fluoroalkyl, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$.

[00352] In further or alternative embodiments, G_6 is W - G_7 , wherein W is (substituted or unsubstituted heterocycloalkyl) or (substituted or unsubstituted heteroaryl) and G_7 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_8$, $N(R_9)_2$, $-C(=NR_{10})N(R_8)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CON(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), L_5 is $-OC(O)O$ -, $-NHC(O)NH$ -, $-NHC(O)O$ -, $-O(O)CNH$ -, $-NHC(O)$ -, $-C(O)NH$ -, $-C(O)O$ -, or $-OC(O)$.

[00353] In further or alternative embodiments, G_6 is H, CN, NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-CO_2R_9$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, (substituted or unsubstituted alkyl), (substituted or unsubstituted fluoroalkyl), $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)$ -, $-C(O)NH$ -, $-C(O)O$ -, or $-OC(O)$; or G_6 is W - G_7 , wherein W is (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7

is H, halogen, C₁-C₆ alkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, OH, -OR₈, -C(=O)CF₃, CN, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted heteroaryl), -L₅-(substituted or unsubstituted heterocycloalkyl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -NH, -NHC(O), -C(O)NH, -C(O)O, or -OC(O).

5 [00354] In some embodiments, G₇ is H, halogen, C₁-C₆ alkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, OH, -OR₈, -C(=O)CF₃, CN, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted heteroaryl), -L₅-(substituted or unsubstituted heterocycloalkyl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -NH, -NHC(O), -C(O)NH, -C(O)O, or -OC(O).

10 [00355] In further or alternative embodiments, W is a (substituted or unsubstituted heterocycloalkyl containing 0-2 nitrogen atoms, 0-1 O atoms and 0-1 S atoms) or a (substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1 O atoms and 0-1 S atoms).

[00356] In further or alternative embodiments, W is substituted with substituents selected from among H, halogen, -CN, -NO₂, -S(=O)₂NH₂, -OH, -C(O)NH₂, -NH₂, -NMe₂, -NHC(O)CH₃, -C(O)OH, -C(O)OCH₃, -C(O)OCH₂CH₃, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, CF₃, and OCF₃.

15 [00357] In further or alternative embodiments, W is substituted with substituents selected from among H, halogen, -CN, -NO₂, -S(=O)₂NH₂, -OH, -C(O)NH₂, -NH₂, -NMe₂, -NHC(O)CH₃, -C(O)OH, -C(O)OCH₃, -C(O)OCH₂CH₃, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, CF₃, and OCF₃.

[00358] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, 20 oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothienophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, imidazo[1,2-a]pyridinyl, furopyridinyl, quinolizine, dioxinyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, oxazinanonyl, dihydropyrrolyl, dihydroimidazolyl, 25 tetrahydrofuranyl, tetrahydropyranyl, dihydrooxazolyl, oxiranyl, pyrrolidinyl, pyrazolidinyl, dihydrothiophenonyl, imidazolidinonyl, pyrrolidinonyl, dihydrofuranonyl, dioxolanonyl, thiazolidinyl, piperidinonyl, tetrahydronaphthyridinyl, tetrahydroquinolinyl, tetrahydrothiophenyl, indolinyl, tetrahydroquinolinyl, and thiazepanyl.

[00359] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among 30 pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, oxadiazolyl, thiadiazolyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, pyrazolidinyl, dioxolanonyl, and thiazolidinyl.

[00360] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among 35 pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; tetrazolyl; tetrahydropyranyl, and morpholin-4-yl.

[00361] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl containing 1-4 nitrogen atoms.

[00362] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl.

[00363] In further or alternative embodiments, W is a substituted or unsubstituted heteroaryl selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, oxadiazolyl, and thiadiazolyl.

[00364] In further or alternative embodiments, W is a substituted or unsubstituted group selected from among pyridinyl; pyrazinyl; pyrimidinyl; 1,3,4-oxadiazolyl; pyridazinyl; imidazolyl; thiazolyl; isoxazolyl; pyrazolyl; 1,2,4-oxadiazolyl; 1,3,4-thiadiazolyl; and tetrazolyl.

[00365] In further or alternative embodiments, G₆ is selected from among H, CN, halogen, OR₉, -C(=O)CF₃, -C(=O)R₉, -CO₂R₉, tetrazolyl, (substituted or unsubstituted alkyl), (substituted or unsubstituted fluoroalkyl); pyridin-2-yl; pyridin-3-yl; pyridin-4-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 3-methoxy-pyridin-2-yl; 4-methoxy-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 6-ethoxy-pyridin-2-yl; 3-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 3-trifluoromethyl-pyridin-2-yl; 4-trifluoromethyl-pyridin-2-yl; 5-trifluoromethyl-pyridin-2-yl; 6-trifluoromethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-fluoromethyl-pyridin-2-yl; 5-methoxymethyl-pyridin-2-yl; 5-hydroxymethyl-pyridin-2-yl; 2-methyl-pyridin-3-yl; 6-methyl-pyridin-3-yl; 6-cyano-pyridin-3-yl; 2-methoxy-pyridin-3-yl; 5-methoxy-pyridin-3-yl; 5-fluoro-pyridin-3-yl; 6-carbamoyl-pyridin-3-yl; 6-hydroxy-pyridin-3-yl; 6-methoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-bromo-6-methoxy-pyridin-3-yl; 6-trifluoromethyl-pyridin-3-yl; 6-trifluoromethyl-pyridin-4-yl; 2-trifluoromethyl-pyridin-5-yl; 2-acetyl-amino-pyridin-5-yl; pyrazin-2-yl; pyrimidin-2-yl; pyrimidin-5-yl; 5-amino-pyrazin-2-yl; 1,3,4-oxadiazol-2-ylamine; 6-hydroxy-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-methyl-pyridazin-3-yl; 2-methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl; thiazol-2-yl; 5-methyl-thiazol-2-yl; 5-fluoro-thiazol-2-yl; 5-trifluoromethyl-thiazol-2-yl; 2,4-dimethyl-thiazol-5-yl; 5-methoxy-thiazol-2-yl; 2-methoxy-thiazol-4-yl; 2-ethoxy-thiazol-4-yl; 2-methyl-thiazol-4-yl; 2-methyl-thiazol-5-yl; 4-methyl-thiazol-2-yl; isoxazol-4-yl; 3,5-dimethyl-isoxazol-4-yl; 2-methyl-imidazol-4-yl; 1-methyl-imidazol-5-yl; 1-methyl-imidazol-4-yl; imidazol-4-yl; 4-methyl-imidazol-5-yl; pyrazol-4-yl; 1-methyl-pyrazol-4-yl; 3-methyl-pyrazol-4-yl; 5-methyl-1,2,4-oxadiazol-3-yl; 2-methyl-1,3,4-oxadiazol-5-yl; 1,3,4-oxadiazol-2-yl; 1,3,4-thiadiazol-2-yl; 3-methyl-pyrazol-5-yl; 1,2,3-thiadiazol-4-yl; tetrazol-1-yl; tetrazol-2-yl; 1-methyl-tetrazol-5-yl; 2-methyl-tetrazol-5-yl; 4-methyl-1H-imidazol-2-yl; 5-hydroxy-pyrimidin-2-yl; 2-methoxy-pyrimidin-5-yl; 6-methyl-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-ethoxy-pyridazin-3-yl; 3-methoxy-pyridazin-6-yl; 4-methoxy-tetrahydro-pyran-4-yl; 6-ethoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-fluoro-pyridin-2-yl, and morpholin-4-yl.

[00366] In some embodiments, G₆ is selected from among H; Cl; Br; pyridin-2-yl; pyridin-3-yl; pyridin-4-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 3-methoxy-pyridin-2-yl; 4-methoxy-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 6-ethoxy-pyridin-2-yl; 3-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 3-trifluoromethyl-pyridin-2-yl; 4-trifluoromethyl-pyridin-2-yl; 5-trifluoromethyl-pyridin-2-yl; 6-trifluoromethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-fluoromethyl-pyridin-2-yl; 5-methoxymethyl-pyridin-2-yl; 5-hydroxymethyl-pyridin-2-yl; 2-methyl-pyridin-3-yl; 6-methyl-pyridin-3-yl; 6-

cyano-pyridin-3-yl; 2-methoxy-pyridin-3-yl; 5-methoxy-pyridin-3-yl; 5-fluoro-pyridin-3-yl; 6-carbamoyl-pyridin-3-yl; 6-hydroxy-pyridin-3-yl; 6-methoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-bromo-6-methoxy-pyridin-3-yl; 6-trifluoromethyl-pyridin-3-yl; 6-trifluoromethyl-pyridin-4-yl; 2-trifluoromethyl-pyridin-5-yl; 2-acetylamino-pyridin-5-yl; pyrazin-2-yl; pyrimidin-2-yl; pyrimidin-5-yl; 5-amino-pyrazin-2-yl; 1,3,4-oxadiazol-2-ylamine; 6-hydroxy-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-methyl-pyridazin-3-yl; 2-methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl; thiazol-2-yl; 5-methyl-thiazol-2-yl; 5-fluoro-thiazol-2-yl; 5-trifluoromethyl-thiazol-2-yl; 2,4-dimethyl-thiazol-5-yl; 5-methoxy-thiazol-2-yl; 2-methoxy-thiazol-4-yl; 2-ethoxy-thiazol-4-yl; 2-methyl-thiazol-4-yl; 2-methyl-thiazol-5-yl; 4-methyl-thiazol-2-yl; isoxazol-4-yl; 3,5-dimethyl-isoxazol-4-yl; 2-methyl-imidazol-4-yl; 1-methyl-imidazol-5-yl; 1-methyl-imidazol-4-yl; imidazol-4-yl; 4-methyl-imidazol-5-yl; pyrazol-4-yl; 1-methyl-pyrazol-4-yl; 3-methyl-pyrazol-4-yl; 5-methyl-1,2,4-oxadiazol-3-yl; 2-methyl-1,3,4-oxadiazol-5-yl; 1,3,4-oxadiazol-2-yl; 1,3,4-thiadiazol-2-yl; 3-methyl-pyrazol-5-yl; 1,2,3-thiadiazol-4-yl; tetrazol-1-yl; tetrazol-2-yl; 1-methyl-tetrazol-5-yl; 2-methyl-tetrazol-5-yl; 4-methyl-1*H*-imidazol-2-yl; 5-hydroxy-pyrimidin-2-yl; 2-methoxy-pyrimidin-5-yl; 6-methyl-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-ethoxy-pyridazin-3-yl; 3-methoxy-pyridazin-6-yl; 4-methoxy-tetrahydro-pyran-4-yl; 6-ethoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; and 5-fluoro-pyridin-2-yl.

[00367] In some embodiments, G₆ is selected from among pyridin-2-yl; pyridin-3-yl; pyridin-4-yl; 3-methyl-pyridin-2-yl; 4-methyl-pyridin-2-yl; 5-methyl-pyridin-2-yl; 3-methoxy-pyridin-2-yl; 4-methoxy-pyridin-2-yl; 5-methoxy-pyridin-2-yl; 6-methoxy-pyridin-2-yl; 6-ethoxy-pyridin-2-yl; 3-fluoro-pyridin-2-yl; 5-fluoro-pyridin-2-yl; 3-trifluoromethyl-pyridin-2-yl; 4-trifluoromethyl-pyridin-2-yl; 5-trifluoromethyl-pyridin-2-yl; 6-trifluoromethyl-pyridin-2-yl; 5-carbamoyl-pyridin-2-yl; 5-cyano-pyridin-2-yl; 5-fluoromethyl-pyridin-2-yl; 5-methoxymethyl-pyridin-2-yl; 5-hydroxymethyl-pyridin-2-yl; 2-methyl-pyridin-3-yl; 6-methyl-pyridin-3-yl; 6-cyano-pyridin-3-yl; 2-methoxy-pyridin-3-yl; 5-methoxy-pyridin-3-yl; 5-fluoro-pyridin-3-yl; 6-carbamoyl-pyridin-3-yl; 6-hydroxy-pyridin-3-yl; 6-methoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; 5-bromo-6-methoxy-pyridin-3-yl; 6-trifluoromethyl-pyridin-3-yl; 6-trifluoromethyl-pyridin-4-yl; 2-trifluoromethyl-pyridin-5-yl; 2-acetylamino-pyridin-5-yl; pyrazin-2-yl; pyrimidin-2-yl; pyrimidin-5-yl; 5-amino-pyrazin-2-yl; 1,3,4-oxadiazol-2-ylamine; 6-hydroxy-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-methyl-pyridazin-3-yl; 2-methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl; thiazol-2-yl; 5-methyl-thiazol-2-yl; 5-fluoro-thiazol-2-yl; 5-trifluoromethyl-thiazol-2-yl; 2,4-dimethyl-thiazol-5-yl; 5-methoxy-thiazol-2-yl; 2-methoxy-thiazol-4-yl; 2-ethoxy-thiazol-4-yl; 2-methyl-thiazol-4-yl; 2-methyl-thiazol-5-yl; 4-methyl-thiazol-2-yl; isoxazol-4-yl; 3,5-dimethyl-isoxazol-4-yl; 2-methyl-imidazol-4-yl; 1-methyl-imidazol-5-yl; 1-methyl-imidazol-4-yl; imidazol-4-yl; 4-methyl-imidazol-5-yl; pyrazol-4-yl; 1-methyl-pyrazol-4-yl; 3-methyl-pyrazol-4-yl; 5-methyl-1,2,4-oxadiazol-3-yl; 2-methyl-1,3,4-oxadiazol-5-yl; 1,3,4-oxadiazol-2-yl; 1,3,4-thiadiazol-2-yl; 3-methyl-pyrazol-5-yl; 1,2,3-thiadiazol-4-yl; tetrazol-1-yl; tetrazol-2-yl; 1-methyl-tetrazol-5-yl; 2-methyl-tetrazol-5-yl; 4-methyl-1*H*-imidazol-2-yl; 5-hydroxy-pyrimidin-2-yl; 2-methoxy-pyrimidin-5-yl; 6-methyl-pyridazin-3-yl; 6-methoxy-pyridazin-3-yl; 6-ethoxy-pyridazin-3-yl; 3-methoxy-pyridazin-6-yl; 4-methoxy-tetrahydro-pyran-4-yl; 6-ethoxy-pyridin-3-yl; 6-ethoxy-pyridin-3-yl; and 5-fluoro-pyridin-2-yl.

[00368] In further or alternative embodiments, G₆ is H; Cl; Br; thiazol-2-yl; 2-methoxy-4-pyridazinyl; 2-methoxypyridin-5-yl; 2-methoxythiazol-4-yl; 5-methoxypyridin-2-yl; or 5-trifluoromethylpyridin-2-yl.

[00369] In further or alternative embodiments, R_7 is L_3 -X- L_4 - G_1 ; wherein, L_3 is a substituted or unsubstituted alkyl; X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉-, -NR₉S(=O)₂, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉-, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉-, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉-, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; and L_4 is a bond or a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or substituted or unsubstituted cycloalkyl.

[00370] In further or alternative embodiments, G_1 is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, or G_1 is W- G_5 , where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[00371] In further or alternative embodiments, X is a bond, -O-, -CR₉(OR₉), S, -S(O), -S(O)₂, -NR₈, -O-N=CH, -CH=N-O, -NHC(=O) or -C(=O)NH.

[00372] In further or alternative embodiments, R_7 is L_3 -X- L_4 - G_1 , wherein, L_3 is a bond, substituted or unsubstituted alkyl, or substituted or unsubstituted alkynyl; X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉; L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl; G_1 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)C(=O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G_1 is W- G_5 , where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(R₉)₂(OR₉), -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[00373] In further or alternative embodiments, G_1 is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈.

[00374] In further or alternative embodiments, X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(=O)-, or -C(O)NR₉.

[00375] In further or alternative embodiments, X is a bond or -CR₉(OR₉).

[00376] In further or alternative embodiments, X is a bond.


[00377] In further or alternative embodiments, R₉ is H, C₁-C₆alkyl, benzyl, or heteroarylmethyl.

[00378] In further or alternative embodiments, R₉ is H or C₁-C₆ alkyl.

[00379] In further or alternative embodiments, R₉ is H.

[00381] In further or alternative embodiments, G₁ is W-G₅, where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl. In further or alternative embodiments, G₁ is W-G₅, where W is a (substituted or unsubstituted heterocycloalkyl containing 0-1 O atoms and 0-2 N atoms), or (substituted or unsubstituted heteroaryl containing 0-4 N atoms).



[00383] In further or alternative embodiments, G₁ is selected from among H, OH, CN, CO₂H, CO₂Me, CO₂Et,

[illegible]

[00384] In further or alternative embodiments, G_1 is $-OR_9$, $N(R_9)_2$, or $-CO_2R_9$.

[00385] In further or alternative embodiments, G_1 is selected from among H, OH, CN, CO_2H , CO_2Me , CO_2Et ,

CO_2NH_2 , CO_2NHMe , $\text{CO}_2\text{N}(\text{Me})_2$, $\text{CO}_2\text{N}(\text{Et})_2$, $-\text{NH}_2$, $-\text{NHMe}$, $-\text{N}(\text{Me})_2$, $-\text{N}(\text{Et})_2$, $-\text{NMe}(\text{iPr})$,



[00386] In further or alternative embodiments, G_1 is selected from among OH, CO_2H , CO_2Me , CO_2Et , CO_2NH_2 , CO_2NHMe , $\text{CO}_2\text{N}(\text{Me})_2$, and $\text{CO}_2\text{N}(\text{Et})_2$.

[00387] In further or alternative embodiments, G_1 is $-\text{OR}_9$, or $-\text{CO}_2\text{R}_9$.

[00388] In further or alternative embodiments, G_1 is $-\text{CO}_2\text{R}_9$.

[00389] In further or alternative embodiments, L_3 is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl.

[00390] In further or alternative embodiments, L_3 is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

[00391] In further or alternative embodiments, L_3 is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00392] In further or alternative embodiments, L_3 is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00393] In further or alternative embodiments, L_3 is methandiyl; or ethan-1,2-diyl.

[00394] In further or alternative embodiments, L_3 is methandiyl.

[00395] In further or alternative embodiments, L_3 is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

[00396] In further or alternative embodiments, L_3 is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00397] In further or alternative embodiments, L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cycloalkyl.

[00398] In further or alternative embodiments, L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-

diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

[00399] In further or alternative embodiments, L_4 is a bond, methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00400] In further or alternative embodiments, L_4 is a bond, ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00401] In further or alternative embodiments, L_3 is a methandiyl; ethan-1,2-diyl; X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, or $-C(O)NR_9$; L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methylpropan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; pentan-3,3-diyl, cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

[00402] In further or alternative embodiments, L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00403] In further or alternative embodiments, L_3 is methandiyl; X is a bond; L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00404] In further or alternative embodiments, L_3 is unsubstituted alkyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00405] In further or alternative embodiments, L_3 is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00406] In further or alternative embodiments, L_3 is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00407] In further or alternative embodiments, L_3 is 2-methyl-propan-1,2-diyl; or 2-ethyl-butan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.





[00408] In further or alternative embodiments, L_3 is unsubstituted alkyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.

[00409] In further or alternative embodiments, L_3 is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.


[00410] In further or alternative embodiments, L_3 is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.

[00411] In further or alternative embodiments, L_3 is 2-methyl-propan-1,2-diyl; 2-ethyl-butan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.


[00412] In some embodiments, L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-CH_2C(CH_2CH_3)H-$, $-CH_2C(isopropyl)H-$, $-CH_2C(tert-butyl)H-$, $-CH_2C(CH_3)_2-$, $-CH_2C(CH_2CH_3)_2-$,



[00413] In further or alternative embodiments, L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-CH_2C(CH_2CH_3)H-$, $-CH_2C(CH_3)_2-$, $-CH_2C(CH_2CH_3)_2-$,




[00414] In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$,



[00415] In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_3)_2-$, or $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$.


[00416] In some embodiment, R_7 is selected from among







[00417] In some embodiment, R₇ is selected from among , , , , , , , , , , , , ,






[00418] In some embodiment, R_7 is selected from among







[00419] In some embodiment, R₇ is selected from among



5




10



- 5 **[00424]** In further or alternative embodiments, L_3 is methandiyl; or ethan-1,2-diyl; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.
- 10 **[00425]** In further or alternative embodiments, X is a bond; and L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cycloalkyl.
- [00426]** In further or alternative embodiments, L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.
- 15 **[00427]** In further or alternative embodiments, L_3 is methandiyl; X is a bond; and L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.
- 20 **[00428]** In further or alternative embodiments, L_4 is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and G_1 is $-CO_2R_9$.
- [00429]** In further or alternative embodiments, L_3 is methandiyl; X is a bond; and L_4 is propan-1,1-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[00430] In a further or alternative embodiment, compounds of Formula (E) have a structure selected from among:



Y	Z	-G ₆	R ₆
N-t-Butoxycarbonylpyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
Pyrrolidon-5-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Methylsulfonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Trifluoroacetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-t-Butoxycarbonyl-4,5-dihydroimidazol-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
4,5-Dihydroimidazol-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-t-Butoxycarbonyl indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
Morpholin-4-yl	-C(=O)CH ₂ -S-	Cl	2-methyl-2-propylthio
Indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio S,S-dioxide
N-Cyclopropylcarbonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Benzoyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(2-Methylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Propanoyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-t-Butoxycarbonyl indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
Indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Acetyl indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio-S-oxide
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	benzyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	H
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	H
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbutanoyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbutanoyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	ethyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	propyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methylpropanoyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	cyclopropylcarbonyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	benzoyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	cyclobutylcarbonyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	acetyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	propanoyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methylpropyl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbut-1-yl
N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	cyclobutylmethyl
N-(4-Phenylbenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Phenylacetyl)-pyrrolidin-2-ylmethyl	-CH ₂ S-	Cl	2-methyl-2-propylthio

Y	Z	-G ₆	R ₆
N-(3-Phenylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(3-Phenoxybenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(4-Phenoxybenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Nicotinoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Pyridin-4-ylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(4-Phenylbenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Phenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(3-Phenylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Phenylcyclopropylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Nicotinoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Pyridin-4-ylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(Phenylcyclopropylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(4-Chlorobenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(4-Benzyloxyphenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(4-Benzyloxyphenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(tert Butoxycarbonyl)piperidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(tert Butoxycarbonyl)piperidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
N-(2-Bromoethoxycarbonyl)-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
Pyrrolidin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio
2-Methyl-1,3-dioxolan-2-yl	-CH ₂ CH ₂ S-	Br	2-methyl-2-propylthio
N-t-Butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	thiazol-2-yl	2-methyl-2-propylthio
Pyrrolidin-2-yl	-CH ₂ S-	Thiazol-2-yl	2-methyl-2-propylthio
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	thiazol-2-yl	2-methyl-2-propylthio
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Thiazol-2-yl	H
N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxy-4-pyridazinyl	2-methyl-2-propylthio
N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	2-methoxy-4-pyridazinyl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxypyridin-5-yl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxythiazol-4-yl	2-methyl-2-propylthio
N-Acetyl-indolin-2-yl	-CH ₂ S-	5-methoxypyridin-2-yl	2-methyl-2-propylthio
2-Methyl-1,3-dioxolan-2-yl	-CH ₂ CH ₂ S-	2-methoxypyridin-5-yl	2-methyl-2-propylthio
N-(Methoxyacetyl) indolin-2-yl	-CH ₂ S-	5-trifluoromethylpyridin-2-yl	2-methyl-2-propylthio

where R₇ is as defined herein. In another embodiment, the "Z" group in the foregoing table is replaced with an S- group to form a new set of compounds described herein.


[00431] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[00432] Further embodiments of Formula (E), include, but are not limited to, compounds shown in Figures 8-11 and in Tables 6-8.

[00433] In another aspect, described herein are compounds of Formula (A). Compounds of Formula (A), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent or leukotriene mediated conditions or diseases, including, but not limited to, asthma, myocardial infarction, chronic obstructive pulmonary disease,

pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[00434] In an alternative or further aspect, compounds provided herein have a structure of Formula (A) as follows:



wherein, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, or $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl, or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl, or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;

Y is H, $-CO_2H$, tetrazolyl, $-NHS(=O)_2R_{3b}$, $S(=O)_2N(R_4)_2$, OH, $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-C(O)NHS(=O)_2R_{3b}$, $-S(=O)_2NHC(O)R_4$, CN, $N(R_4)_2$, $-N(R_4)C(O)R_4$, $-C(=NR_3)N(R_4)_2$, $-NR_4C(=NR_3)N(R_4)_2$, $-NR_4C(=CHR_3)N(R_4)_2$, $-C(O)NR_4C(=NR_3)N(R_4)_2$, $-C(O)NR_4C(=CHR_3)N(R_4)_2$, $-CO_2R_{3b}$, $-C(O)R_4$, $-CON(R_4)_2$, $-SR_{3b}$, $-S(=O)R_{3b}$, $-S(=O)_2R_{3b}$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted alkenyl), $-L_1$ -(substituted or unsubstituted alkynyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted aryl) or $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=CR_3)N(R_4)_2$;

where L_1 is a bond, a substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, substituted or unsubstituted heteroalkenyl, a substituted or unsubstituted heteroalkynyl, or substituted or unsubstituted aryl;

each R_3 is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl; each R_{3b} is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_4 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_4 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;

R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, C(O), $-CH(OH)$, $-(\text{substituted or unsubstituted } C_1-C_6 \text{ alkyl})$, or $-(\text{substituted or unsubstituted } C_2-C_6 \text{ alkenyl})$;

R_7 is selected from:

(i) $L_3-X-L_4-G_1$, wherein, L_3 is a substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted

heterocycloalkyl; X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉-, -NR₉S(=O)₂, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉-, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉-, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉-, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; L₄ is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G₁ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -OC(O)O-, -NHC(O)NH-, -NHC(O)O, -O(O)CNH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G₁ is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈; each R₈ is independently selected from substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; each R₉ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; or two R₉ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R₁₀ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl;

- (ii) L₃-X-L₄-G₂, wherein, L₃ is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl; X is -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉-, -NR₉S(=O)₂, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉-, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉-, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉-, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; L₄ is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G₂ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -OC(O)O-, -NHC(O)NH-, -NHC(O)O, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G₂ is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈; each R₈ is independently selected from substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl;

each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

(iii) L_3 -X- L_4 - G_3 , wherein, X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NR_9$, $-NR_9S(=O)_2$, $-OC(O)NR_9$, $-NR_9C(O)O$, $-CH=NO$, $-ON=CH$, $-NR_9C(O)NR_9$, heteroaryl, aryl, $-NR_9C(=NR_{10})NR_9$, $-NR_9C(=NR_{10})$, $-C(=NR_{10})NR_9$, $-OC(=NR_{10})$, or $-C(=NR_{10})O$; L_3 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl; L_4 is a (substituted or unsubstituted alkenyl) or (substituted or unsubstituted alkynyl); G_3 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-OC(O)O$, $-NHC(O)NH$, $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_3 is W- G_5 , where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$; each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

or (iv) L_3 -X- L_4 - G_4 , wherein, L_3 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl; X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NR_9$, $-NR_9S(=O)_2$, $-OC(O)NR_9$, $-NR_9C(O)O$, $-CH=NO$, $-ON=CH$, $-NR_9C(O)NR_9$, heteroaryl, aryl, $-NR_9C(=NR_{10})NR_9$, $-NR_9C(=NR_{10})$, $-C(=NR_{10})NR_9$, $-OC(=NR_{10})$, or $-C(=NR_{10})O$; L_4 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G_4 is $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-OC(O)NH$, $-C(O)O$, or $-OC(O)$; or G_4 is $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_4 is W- G_5 , where

W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$; each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

R_5 is H, halogen, $-N_3$, $-CN$, $-NO_2$, $-L_6$ -(substituted or unsubstituted C_1 - C_6 alkyl), $-L_6$ -(substituted or unsubstituted C_2 - C_6 alkenyl), $-L_6$ -(substituted or unsubstituted heteroaryl), or $-L_6$ -(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-NHC(O)NH-$, or $-C(O)NH$;

R_{11} is L_7 - L_{10} - G_6 ; wherein L_7 is a bond, $-O$, $-S$, $-S(=O)$, $-S(=O)_2$, $-NH$, $-C(O)$, $-C(O)NH$, $-NHC(O)$, (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_2 - C_6 alkenyl); L_{10} is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl), and G_6 is H, CN, SCN, N_3 , NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-NHC(O)NH-$, $-OC(O)O-$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_6 is W- G_7 , wherein W is (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroalkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NH$, $-NHC(O)O$, $-NHC(O)NH-$, $-OC(O)O-$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; and

R_{12} is L_8 - L_9 - R_{13} , wherein L_8 is a bond, (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_2 - C_4 alkenyl); L_9 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)NH-$, $-OC(O)O-$, $-NHC(O)-$, $-C(O)NH-$, $-C(O)O-$, or $-OC(O)-$; R_{13} is H, (substituted or unsubstituted C_1 - C_6 alkyl), (substituted or unsubstituted C_3 - C_6 cycloalkyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heteroaryl), or (substituted or unsubstituted heterocycloalkyl); or R_7 and R_{12} can together form a 4 to 8-membered heterocyclic ring;

or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00435] For any and all of the embodiments of Formula (A), substituents can be selected from among from a subset of the listed alternatives.

5 [00436] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$. In other embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S(O)_m$.

[00437] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, and $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

10 [00438] In some embodiments, Z is selected from $-S-$, $-[C(R_2)_2]_n C(R_1)_2 S-$, and $-SC(R_1)_2 [C(R_2)_2]_n-$.

[00439] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00440] In some embodiments, each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00441] In some embodiments, each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

15 [00442] In some embodiments, Z is $-S-$ or $[C(R_2)_2]_n C(R_1)_2 S-$.

[00443] In some embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S-$.

[00444] In some embodiments, Z is $-S-$.

[00445] In some embodiments, Z is CH_2S- .

[00446] In some embodiments, Z is $-S-$, $-SCH_2-$, $-CH_2S-$, or $-CH(CH_3)S-$.

20 [00447] In some embodiments, Z is $-S-$ or $-CH_2S-$.

[00448] In further or alternative embodiments of compounds of Formula (A), Y is $-L_1$ -substituted or unsubstituted aryl. In further or alternative embodiments of compounds of Formula (A), Y is $-L_1$ -substituted or unsubstituted heteroaryl. In further or alternative embodiments of compounds of Formula (A), Y is $-L_1$ -substituted or unsubstituted heterocycloalkyl. In further or alternative embodiments of compounds of Formula (A), Y is $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, or $-L_1-NR_4C(=CHR_3)N(R_4)_2$.

25 [00449] In further or alternative embodiments of compounds of Formula (A), L_1 is a bond, a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, or substituted or unsubstituted aryl.

30 [00450] In further or alternative embodiments of compounds of Formula (A), L_1 is a bond, or a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, or substituted or unsubstituted aryl.

[00451] In further or alternative embodiments of compounds of Formula (A), L_1 is a bond, or a substituted or unsubstituted alkyl, a substituted or unsubstituted heteroaryl, or substituted or unsubstituted aryl.

[00452] In further or alternative embodiments of compounds of Formula (A), L_1 is a bond, or a substituted or unsubstituted alkyl. In further or alternative embodiments of compounds of Formula (A), L_1 is a bond.

[00453] In further or alternative embodiments of compounds of Formula (A), R_6 is L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted aryl), or L_2 -(substituted or unsubstituted cycloalkyl), where

40 L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or (substituted or unsubstituted C_1 - C_6 alkyl).

[00454] In further or alternative embodiments of compounds of Formula (A), R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted aryl), or L_2 -(substituted or unsubstituted cycloalkyl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or (substituted or unsubstituted C_1 - C_6 alkyl).

[00455] In further or alternative embodiments of compounds of Formula (A), R_6 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00456] In further or alternative embodiments of compounds of Formula (A), R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00457] In further or alternative embodiments of compounds of Formula (A), R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00458] In further or alternative embodiments of compounds of Formula (A), R_6 is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; or cyclohexyloxy.

[00459] In further or alternative embodiments of compounds of Formula (A), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00460] In further or alternative embodiments of compounds of Formula (A), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

[00461] In further or alternative embodiments of compounds of Formula (A), R_6 is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00462] In further or alternative embodiments of compounds of Formula (A), R_6 H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.


[00463] In further or alternative embodiments of compounds of Formula (A), R₆ is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00464] In further or alternative embodiments of compounds of Formula (A), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.


[00465] In further or alternative embodiments of compounds of Formula (A), R₇ is L₃-X-L₄-G₁; wherein, L₃ is a substituted or unsubstituted alkyl; X is -NHC(O), -C(O)NH, -NR₈C(O), -C(O)NR₈, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₈-, -NR₈S(=O)₂-, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₈-, -NR₈C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉-, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉-, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉-, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; L₄ is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G₁ is H, -CO₂H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₈, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -NHC(O)O, -O(O)CNH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G₁ is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, -CO₂H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₈, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈; each R₈ is independently selected from substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; each R₉ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; or two R₉ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R₁₀ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl.

[00466] In further or alternative embodiments, G₁ is H, -CO₂H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₈, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈, or G₁ is W-G₅, where W is substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, -CO₂H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₈, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈. In further or alternative embodiments, X is a bond, -O-, S, -S(O), -S(O)₂, -NR₈, -O-N=CH, -CH=N-O, -NHC(=O) or -C(=O)NH.


[00467] In some embodiments, L₃-X-L₄ is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂C(CH₃)H-, -CH₂C(CH₂CH₃)H-, -CH₂C(isopropyl)H-, -CH₂C(tert-butyl)H-, -CH₂C(CH₃)₂-, -CH₂C(CH₂CH₃)₂-,



[00468] In further or alternative embodiments, L₃-X-L₄ is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂C(CH₃)H-, -CH₂C(CH₂CH₃)H-, -CH₂C(CH₃)₂-, -CH₂C(CH₂CH₃)₂-,



[00469] In further or alternative embodiments, L₃-X-L₄ is -CH₂C(CH₂CH₃)H-, -CH₂C(CH₂CH₃)₂-,



[00470] In further or alternative embodiments, L₃-X-L₄ is -CH₂C(CH₃)₂-, or -CH₂C(CH₂CH₃)₂-.

[00471] In further or alternative embodiments of compounds of Formula (A), R₁₁ is L₇-L₁₀-W-G₇. In further or alternative embodiments, W is (substituted or unsubstituted heteroaryl) or (substituted or unsubstituted heterocycloalkyl).


[00472] In further or alternative embodiments of compounds of Formula (A), R₁₂ is or L₈-L₉-R₁₃, wherein L₈ is a bond, or (substituted or unsubstituted C₁-C₆ alkyl); L₉ is a bond, -O-, -S-, -S(=O), -S(=O)₂, -NH-, -C(O)-, -(CH₂)-, -NHC(O)O-, -NHC(O)-, or -C(O)NH; R₁₃, is H, (substituted or unsubstituted C₁-C₆ alkyl) or (substituted or unsubstituted C₃-C₆ cycloalkyl).

[00473] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[00474] In another aspect, described herein are compounds of Formula (B). Compounds of Formula (B), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or inhibit

FLAP and may be used to treat patients suffering from leukotriene-dependent or leukotriene mediated conditions or diseases, including, but not limited to, asthma, myocardial infarction, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[00475] In another or alternative aspect, compounds provided herein have a structure of Formula (B) as follows:



- wherein, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; n is 0, 1, 2, or 3;
- Y is H, $-CO_2H$, tetrazolyl, $-NHS(=O)_2 R_{3b}$, $S(=O)_2 N(R_4)_2$, OH, $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-C(O)NHS(=O)_2 R_{3b}$, $-S(=O)_2 NHC(O)R_4$, CN, $N(R_4)_2$, $-N(R_4)C(O)R_4$, $-C(=NR_3)N(R_4)_2$, $-NR_4C(=NR_3)N(R_4)_2$, $-NR_4C(=CHR_3)N(R_4)_2$, $-C(O)NR_4C(=NR_3)N(R_4)_2$, $-C(O)NR_4C(=CHR_3)N(R_4)_2$, $CO_2 R_{3b}$, $-C(O)R_4$, $-CON(R_4)_2$, $-SR_{3b}$, $-S(=O)R_{3b}$, $-S(=O)_2 R_{3b}$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted alkenyl), $-L_1$ -(substituted or unsubstituted alkynyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted aryl) or $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=CHR_3)N(R_4)_2$;
- where L_1 is a bond, a substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, substituted or unsubstituted heteroalkenyl, a substituted or unsubstituted heteroalkynyl, or substituted or unsubstituted aryl;
- each R_3 is independently selected from H, $-S(=O)_2 R_8$, $-S(=O)_2 NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl; each R_{3b} is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_4 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_4 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;
- R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, C(O), $-CH(OH)$, $-(\text{substituted or unsubstituted } C_1-C_6 \text{ alkyl})$, or $-(\text{substituted or unsubstituted } C_2-C_6 \text{ alkenyl})$;
- R_7 is H or substituted or unsubstituted alkyl;
- R_5 is H, halogen, $-N_3$, $-CN$, $-NO_2$, $-L_6$ -(substituted or unsubstituted C_1 - C_6 alkyl), $-L_6$ -(substituted or unsubstituted C_2 - C_6 alkenyl), $-L_6$ -(substituted or unsubstituted heteroaryl), or $-L_6$ -(substituted or

unsubstituted aryl), wherein L_6 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-NHC(O)NH-$, or $-C(O)NH$;

R_{11} is $L_7-L_{10}-G_6$; wherein L_7 is a bond, $-O$, $-S$, $-S(=O)$, $-S(=O)_2$, $-NH$, $-C(O)$, $-C(O)NH$, $-NHC(O)$, (substituted or unsubstituted C_1-C_6 alkyl), or (substituted or unsubstituted C_2-C_6 alkenyl); L_{10} is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl), and G_6 is H, CN, SCN, N_3 , NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-NHC(O)NH-$, $-OC(O)O-$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_6 is $W-G_7$, wherein W is (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroalkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NH$, $-NHC(O)O$, $-NHC(O)NH-$, $-OC(O)O-$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

R_{12} is $L_3-X-L_4-G_1$, wherein, L_3 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl; X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NR_9-$, $-NR_9S(=O)_2$, $-OC(O)NR_9-$, $-NR_9C(O)O-$, $-CH=NO-$, $-ON=CH-$, $-NR_9C(O)NR_9-$, heteroaryl, aryl, $-NR_9C(=NR_{10})NR_9-$, $-NR_9C(=NR_{10})-$, $-C(=NR_{10})NR_9-$, $-OC(=NR_{10})-$, or $-C(=NR_{10})O-$; L_4 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G_1 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-OC(O)O-$, $-NHC(O)NH-$, $-NHC(O)O$, $-O(O)CNH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; or G_1 is $W-G_5$, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$,

or $-S(=O)_2R_8$; each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00476] For any and all of the embodiments of Formula (A), substituents can be selected from among from a subset of the listed alternatives.

[00477] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_nC(R_1)_2S(O)_m$, $S(O)_mC(R_1)_2[C(R_2)_2]_n$. In other embodiments, Z is $[C(R_2)_2]_nC(R_1)_2S(O)_m$.

[00478] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_nC(R_1)_2S(O)_m$, and $S(O)_mC(R_1)_2[C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00479] In some embodiments, Z is selected from $-S-$, $-[C(R_2)_2]_nC(R_1)_2S-$, and $-SC(R_1)_2[C(R_2)_2]_n-$.

[00480] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00481] In some embodiments, each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00482] In some embodiments, each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00483] In some embodiments, Z is $-S-$ or $[C(R_2)_2]_nC(R_1)_2S-$.

[00484] In some embodiments, Z is $[C(R_2)_2]_nC(R_1)_2S-$.

[00485] In some embodiments, Z is $-S-$.

[00486] In some embodiments, Z is CH_2S- .

[00487] In some embodiments, Z is $-S-$, $-SCH_2-$, $-CH_2S-$, or $-CH(CH_3)S-$.

[00488] In some embodiments, Z is $-S-$ or $-CH_2S-$.

[00489] In further or alternative embodiments of compounds of Formula (B), Y is $-L_1$ -substituted or unsubstituted aryl. In further or alternative embodiments, Y is $-L_1$ -substituted or unsubstituted heteroaryl. In further or alternative embodiments, Y is $-L_1$ -substituted or unsubstituted heterocycloalkyl. In further or alternative embodiments, Y is $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, or $-L_1-NR_4C(=CHR_3)N(R_4)_2$.

[00490] In further or alternative embodiments of compounds of Formula (B), L_1 is a bond, a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, or substituted or unsubstituted aryl.

[00491] In further or alternative embodiments of compounds of Formula (B), L_1 is a bond, a substituted or unsubstituted alkyl.

[00492] In further or alternative embodiments of compounds of Formula (B), L_1 is a bond.

[00493] In further or alternative embodiments of compounds of Formula (B), R_6 is L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or substituted or unsubstituted alkyl.

[00494] In further or alternative embodiments of compounds of Formula (B), R₆ is H, L₂-(substituted or unsubstituted alkyl), or L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(O)₂, -C(O), -CH(OH), or substituted or unsubstituted alkyl.

[00495] In further or alternative embodiments of compounds of Formula (B), R₆ is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00496] In further or alternative embodiments of compounds of Formula (B), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00497] In further or alternative embodiments of compounds of Formula (B), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00498] In further or alternative embodiments of compounds of Formula (B), R₆ is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

[00499] In further or alternative embodiments of compounds of Formula (B), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00500] In further or alternative embodiments of compounds of Formula (B), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

[00501] In further or alternative embodiments of compounds of Formula (B), R₆ is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00502] In further or alternative embodiments of compounds of Formula (B), R₆ is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00503] In further or alternative embodiments of compounds of Formula (B), R_6 is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00504] In further or alternative embodiments of compounds of Formula (B), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00505] In further or alternative embodiments of compounds of Formula (B), R_{11} is L_7 - L_{10} -W- G_7 . In further or alternative embodiments, W is (substituted or unsubstituted heteroaryl) or (substituted or unsubstituted heterocycloalkyl).

[00506] In further or alternative embodiments of compounds of Formula (B), R_{12} is L_3 -X- L_4 - G_1 wherein; L_3 is a substituted or unsubstituted alkyl; X is a bond, O, -C(=O), - $CR_9(OR_9)$, S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉, -NR₉S(=O)₂, -OC(O)NR₉, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; and L_4 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl. In further or alternative embodiments, G_1 is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, or G_1 is W- G_5 , where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈. In further or alternative embodiments, X is a bond, -O-, S, -S(O), -S(O)₂, -NR₈, -O-N=CH, -CH=N-O, -NHC(=O) or -C(=O)NH.

[00507] In further or alternative embodiments of compounds of Formula (B), R_7 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00508] In further or alternative embodiments of compounds of Formula (B), R_7 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; or cyclohexylmethyl.

[00509] In further or alternative embodiments of compounds of Formula (B), R_7 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; or 3,3-dimethylbutyl.


[00510] In further or alternative embodiments of compounds of Formula (B), R_7 is prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; *tert*-butyl; 3-methylbutyl; or 3,3-dimethylbutyl.

[00511] In further or alternative embodiments of compounds of Formula (B), R_7 is 2-methylpropyl.

[00512] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

- 5 [00513] In another aspect, described herein are compounds of Formula (C). Compounds of Formula (C), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent or leukotriene mediated conditions or diseases, including, but not limited to, asthma, myocardial infarction, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[00514] In another aspect, compounds provided herein have a structure of Formula (C) as follows:



- 15 wherein, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; n is 0, 1, 2, or 3;
- Y is H, $-CO_2H$, tetrazolyl, $-NHS(=O)_2R_{3b}$, $S(=O)_2N(R_4)_2$, OH, $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, -
 20 $C(O)NHS(=O)_2R_{3b}$, $-S(=O)_2NHC(O)R_4$, CN, $N(R_4)_2$, $-N(R_4)C(O)R_4$, $-C(=NR_3)N(R_4)_2$, -
 $NR_4C(=NR_3)N(R_4)_2$, $-NR_4C(=CHR_3)N(R_4)_2$, $-C(O)NR_4C(=NR_3)N(R_4)_2$, $-C(O)NR_4C(=CHR_3)N(R_4)_2$, -
 CO_2R_{3b} , $-C(O)R_4$, $-CON(R_4)_2$, $-SR_{3b}$, $-S(=O)R_{3b}$, $-S(=O)_2R_{3b}$, $-L_1$ -(substituted or unsubstituted alkyl), -
 L_1 -(substituted or unsubstituted alkenyl), $-L_1$ -(substituted or unsubstituted alkynyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted aryl) or $-L_1-C(=NR_4)N(R_4)_2$, $-L_1$ -
 25 $NR_4C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=CHR_3)N(R_4)_2$;
- where L_1 is a bond, a substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, substituted or unsubstituted heteroalkenyl, a substituted or unsubstituted heteroalkynyl, or substituted or unsubstituted aryl;
- each R_3 is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl; each R_{3b} is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_4 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_4 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;
- 35

R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, -S(=O), -S(=O)₂, C(O), -CH(OH), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R_7 is L_3 -X- L_4 - G_1 , wherein, L_3 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl; X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(O), -C(O)NR₉, -S(=O)₂NR₉, -NR₉S(=O)₂, -OC(O)NR₉, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₉C(O)NR₉, heteroaryl, aryl, -NR₉C(=NR₁₀)NR₉, -NR₉C(=NR₁₀)-, -C(=NR₁₀)NR₉, -OC(=NR₁₀)-, or -C(=NR₁₀)O-; L_4 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl; G_1 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L_5 is -OC(O)O-, -NHC(O)NH-, -NHC(O)O, -O(O)CNH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G_1 is W- G_5 , where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈; each R₈ is independently selected from substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; each R₉ is independently selected from H, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted C₃-C₈cycloalkyl, phenyl or benzyl; or two R₉ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and each R₁₀ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl;

R_5 is H, halogen, -N₃, -CN, -NO₂, -L₆-(substituted or unsubstituted C₁-C₆ alkyl), -L₆-(substituted or unsubstituted C₂-C₆ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), -NHC(O)NH-, or -C(O)NH;

R_{11} is L_7 - L_{10} - G_6 ; wherein, L_7 is a bond, -C(O), -C(O)NH, (substituted or unsubstituted C₁-C₆ alkyl), or (substituted or unsubstituted C₂-C₆ alkenyl); L_{10} is a (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl), G_6 is tetrazolyl, -NHS(=O)₂R₈, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₈, -C(=NR₁₀)N(R₈)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl),

wherein L_5 is $-\text{OC}(\text{O})\text{O}-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})\text{O}-$, $-\text{O}(\text{O})\text{CNH}-$, $-\text{NHC}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{C}(\text{O})\text{O}-$, or $-\text{OC}(\text{O})-$; or G_6 is $\text{W}-\text{G}_7$, wherein W is a (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is, tetrazolyl, $-\text{NHS}(=\text{O})_2\text{R}_8$, $\text{S}(=\text{O})_2\text{N}(\text{R}_9)$, OH , $-\text{C}(=\text{O})\text{CF}_3$, $-\text{C}(\text{O})\text{NHS}(=\text{O})_2\text{R}_8$, $-\text{S}(=\text{O})_2\text{NHC}(\text{O})\text{R}_8$, $\text{N}(\text{R}_9)_2$, $-\text{C}(=\text{NR}_{10})\text{N}(\text{R}_8)_2$, $-\text{NR}_9\text{C}(=\text{NR}_{10})\text{N}(\text{R}_9)_2$, $-\text{NR}_9\text{C}(=\text{CHR}_{10})\text{N}(\text{R}_9)_2$, $-\text{CON}(\text{R}_9)_2$, $-\text{L}_5$ -(substituted or unsubstituted alkyl), $-\text{L}_5$ -(substituted or unsubstituted alkenyl), $-\text{L}_5$ -(substituted or unsubstituted heteroaryl), $-\text{L}_5$ -(substituted or unsubstituted heterocycloalkyl), or $-\text{L}_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-\text{OC}(\text{O})\text{O}-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})\text{O}-$, $-\text{O}(\text{O})\text{CNH}-$, $-\text{NHC}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{C}(\text{O})\text{O}-$, or $-\text{OC}(\text{O})-$;

R_{12} is $\text{L}_8-\text{L}_9-\text{R}_{13}$, wherein L_8 is a bond, (substituted or unsubstituted C_1-C_6 alkyl), or (substituted or unsubstituted C_2-C_4 alkenyl); L_9 is a bond, O , S , $-\text{S}(=\text{O})$, $\text{S}(=\text{O})_2$, NH , $\text{C}(\text{O})$, $-\text{NHC}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{NH}-$, $-\text{NHC}(\text{O})\text{NH}-$, $-\text{OC}(\text{O})\text{O}-$, $-\text{NHC}(\text{O})-$, $-\text{C}(\text{O})\text{NH}-$, $-\text{C}(\text{O})\text{O}-$, or $-\text{OC}(\text{O})-$; R_{13} is H , (substituted or unsubstituted C_1-C_6 alkyl), (substituted or unsubstituted C_3-C_6 cycloalkyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heteroaryl), or (substituted or unsubstituted heterocycloalkyl); or R_7 and R_{12} can together form a 4 to 8-membered heterocyclic ring; or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00515] For any and all of the embodiments of Formula (C), substituents can be selected from among a subset of the listed alternatives.

[00516] In some embodiments, Z is selected from $\text{S}(\text{O})_m$, $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}(\text{O})_m$, $\text{S}(\text{O})_m\text{C}(\text{R}_1)_2[\text{C}(\text{R}_2)_2]_n$. In other embodiments, Z is $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}(\text{O})_m$.

[00517] In some embodiments, Z is selected from $\text{S}(\text{O})_m$, $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}(\text{O})_m$, and $\text{S}(\text{O})_m\text{C}(\text{R}_1)_2[\text{C}(\text{R}_2)_2]_n$, wherein each R_1 is independently H , CF_3 , or an optionally substituted C_1-C_6 alkyl; and R_2 is H , OH , OMe , CF_3 , or an optionally substituted C_1-C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00518] In some embodiments, Z is selected from $-\text{S}-$, $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}-$, and $-\text{SC}(\text{R}_1)_2[\text{C}(\text{R}_2)_2]_n-$.

[00519] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00520] In some embodiments, each R_1 is independently H , CF_3 , or an optionally substituted C_1-C_6 alkyl.

[00521] In some embodiments, each R_2 is independently H , OH , OMe , CF_3 , or an optionally substituted C_1-C_6 alkyl.

[00522] In some embodiments, Z is $-\text{S}-$ or $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}-$.

[00523] In some embodiments, Z is $[\text{C}(\text{R}_2)_2]_n\text{C}(\text{R}_1)_2\text{S}-$.

[00524] In some embodiments, Z is $-\text{S}-$.

[00525] In some embodiments, Z is $\text{CH}_2\text{S}-$.

[00526] In some embodiments, Z is $-\text{S}-$, $-\text{SCH}_2-$, $-\text{CH}_2\text{S}-$, or $-\text{CH}(\text{CH}_3)\text{S}-$.

[00527] In some embodiments, Z is $-\text{S}-$ or $-\text{CH}_2\text{S}-$.

[00528] In further or alternative embodiments of compounds of Formula (C), Y is $-\text{CO}_2\text{H}$, tetrazolyl, $-\text{NHS}(=\text{O})_2\text{R}_{3b}$, $\text{S}(=\text{O})_2\text{N}(\text{R}_4)_2$, OH , $-\text{OR}_{3b}$, $-\text{C}(=\text{O})(\text{C}_1-\text{C}_5 \text{ fluoroalkyl})$, $-\text{C}(\text{O})\text{NHS}(=\text{O})_2\text{R}_{3b}$, $-\text{S}(=\text{O})_2\text{NHC}(\text{O})\text{R}_4$, CN , $\text{N}(\text{R}_4)_2$, $-\text{N}(\text{R}_4)\text{C}(\text{O})\text{R}_4$, $-\text{CO}_2\text{R}_{3b}$, $-\text{C}(\text{O})\text{R}_4$, $-\text{CON}(\text{R}_4)_2$, $-\text{L}_1$ -(substituted or unsubstituted alkyl), $-\text{L}_1$ -(substituted or unsubstituted cycloalkyl), $-\text{L}_1$ -(substituted or unsubstituted heterocycloalkyl), $-\text{L}_1$ -(substituted or

unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted aryl) or $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=CHR_3)N(R_4)_2$.

[00529] In further or alternative embodiments of compounds of Formula (C), Y is $-L_1$ -substituted or unsubstituted aryl. In further or alternative embodiments, Y is $-L_1$ -substituted or unsubstituted heteroaryl. In

5 further or alternative embodiments, Y is $-L_1$ -substituted or unsubstituted heterocycloalkyl. In further or alternative embodiments, Y is $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, or $-L_1-NR_4C(=CHR_3)N(R_4)_2$.

[00530] In further or alternative embodiments of compounds of Formula (C), L_1 is a bond, a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, or substituted or unsubstituted

10 aryl. [00531] In further or alternative embodiments of compounds of Formula (C), L_1 is a bond, a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, or substituted or unsubstituted aryl.

[00532] In further or alternative embodiments of compounds of Formula (C), L_1 is a bond, or a substituted or unsubstituted alkyl. In further or alternative embodiments of compounds of Formula (C), L_1 is a bond.

[00533] In further or alternative embodiments of compounds of Formula (C), R_6 is L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or substituted or unsubstituted alkyl.

[00534] In further or alternative embodiments of compounds of Formula (C), R_7 is $L_3-X-L_4-G_1$; wherein, L_3 is a substituted or unsubstituted alkyl; X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NR_9$, $-NR_9S(=O)_2$, $-OC(O)NR_9$, $-NR_9C(O)O$, $-CH=NO$, $-ON=CH$, $-NR_9C(O)NR_9$, heteroaryl, aryl, $-NR_9C(=NR_{10})NR_9$, $-NR_9C(=NR_{10})$, $-C(=NR_{10})NR_9$, $-OC(=NR_{10})$, or $-C(=NR_{10})O$; and L_4 is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl.

25 [00535] In further or alternative embodiments, G_1 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, or G_1 is $W-G_5$, where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$. In further or alternative embodiments, X is a bond, $-O$, $-CR_9(OR_9)$, S, $-S(O)$, $-S(O)_2$, $-NR_8$, $-O-N=CH$, $-CH=N-O$, $-NHC(=O)$ or $-C(=O)NH$.

35 [00536] In further or alternative embodiments of compounds of Formula (C), R_{11} is $L_7-L_{10}-G_6$, wherein L_7 is a bond, (substituted or unsubstituted C_1-C_6 alkyl), and L_{10} is a (substituted or unsubstituted aryl), (substituted or unsubstituted heteroaryl), or (substituted or unsubstituted heterocycloalkyl). In further or alternative embodiments, G_6 is tetrazolyl, $-NHS(=O)_2R_8$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), L_5 is $-OC(O)O$, $-NHC(O)NH$, $-NHC(O)O$, $-O(O)CNH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$. In further or alternative embodiments, L_{10} is a (substituted

or unsubstituted aryl). In further or alternative embodiments, G_6 is $W-G_7$, wherein W is (substituted or unsubstituted heterocycloalkyl) or (substituted or unsubstituted heteroaryl) and G_7 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)$, OH , $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_8$, $N(R_9)_2$, $-C(=NR_{10})N(R_8)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CON(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), L_5 is $-OC(O)O-$, $-NHC(O)NH-$, $-NHC(O)O$, $-O(O)CNH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$.

[00537] In further or alternative embodiments of compounds of Formula (C), L_8 is a bond, (substituted or unsubstituted C_1 - C_6 alkyl); L_9 is a bond, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-C(O)-$, $-(CH_2)-$, $-NHC(O)O-$,

$-NHC(O)-$, or $-C(O)NH$; R_{13} is H , (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_3 - C_6 cycloalkyl).

[00538] In further or alternative embodiments of compounds of Formula (C), R_5 is H .

[00539] In further or alternative embodiments of compounds of Formula (C), R_{12} is $L_8-L_9-R_{13}$, wherein L_8 is a bond; L_9 is a bond; R_{13} is H .

[00540] In further or alternative embodiments of compounds of Formula (C), R_6 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00541] In further or alternative embodiments of compounds of Formula (C), R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00542] In further or alternative embodiments of compounds of Formula (C), R_6 is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00543] In further or alternative embodiments of compounds of Formula (C), R_6 is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

[00544] In further or alternative embodiments of compounds of Formula (C), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00545] In further or alternative embodiments of compounds of Formula (C), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

5 [00546] In further or alternative embodiments of compounds of Formula (C), R_6 is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00547] In further or alternative embodiments of compounds of Formula (C), R_6 is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-
10 butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00548] In further or alternative embodiments of compounds of Formula (C), R_6 is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-
15 butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00549] In further or alternative embodiments of compounds of Formula (C), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-
20 butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00550] In further or alternative embodiments of compounds of Formula (C), X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, or $-C(O)NR_9$.

[00551] In further or alternative embodiments of compounds of Formula (C), X is a bond or $-CR_9(OR_9)$.

[00552] In further or alternative embodiments of compounds of Formula (C), X is a bond.

25 [00553] In further or alternative embodiments of compounds of Formula (C), R_9 is H, C_1-C_6 alkyl, benzyl, or heteroaryl-methyl.

[00554] In further or alternative embodiments of compounds of Formula (C), R_9 is H or C_1-C_6 alkyl.


[00555] In further or alternative embodiments of compounds of Formula (C), R_9 is H.

[00556] In further or alternative embodiments of compounds of Formula (C), G_1 is $-OR_9$, $N(R_9)_2$, $-CO_2R_9$, $-CON(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$.

[00557] In further or alternative embodiments of compounds of Formula (C), G_1 is W- G_5 , where W is a substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl. In further or alternative
35 embodiments, G_1 is W- G_5 , where W is a (substituted or unsubstituted heterocycloalkyl containing 0-1 O atoms and 0-2 N atoms), or (substituted or unsubstituted heteroaryl containing 0-4 N atoms).


[00558] In further or alternative embodiments of compounds of Formula (C), G_1 is W- G_5 , where W is a substituted or unsubstituted group selected from among furanonyl, dihydrofuran-2-onyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, imidazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, 1,3,4-oxadiazolyl, thiazolyl, pyrazolyl, tetrazolyl, oxazolyl, or pyrrolyl.

[00559] In further or alternative embodiments of compounds of Formula (C), G₁ is selected from among H, OH, CN, CO₂H, CO₂Me, CO₂Et, CO₂NH₂, CO₂NHMe, CO₂N(Me)₂, CO₂N(Et)₂, -NH₂, -NHMe, -N(Me)₂, -N(Et)₂, -



[00560] In further or alternative embodiments of compounds of Formula (C), G_1 is $-OR_9$, $N(R_9)_2$, or $-CO_2R_9$.

10 [00561] In further or alternative embodiments of compounds of Formula (C), G₁ is selected from among H, OH, CN, CO₂H, CO₂Me, CO₂Et, CO₂NH₂, CO₂NHMe, CO₂N(Me)₂, CO₂N(Et)₂, -NH₂, -NHMe, -N(Me)₂, -N(Et)₂, -



[00562] In further or alternative embodiments of compounds of Formula (C), G₁ is selected from among OH, CO₂H, CO₂Me, CO₂Et, CO₂NH₂, CO₂NHMe, CO₂N(Me)₂, and CO₂N(Et)₂.

[00563] In further or alternative embodiments of compounds of Formula (C), G_1 is $-OR_9$, or $-CO_2R_9$.

[00564] In further or alternative embodiments of compounds of Formula (C), G_1 is $-\text{CO}_2\text{R}_9$.

[00565] In further or alternative embodiments of compounds of Formula (C), L₃ is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl.

[00566] In further or alternative embodiments of compounds of Formula (C), L_3 is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

5 [00567] In further or alternative embodiments of compounds of Formula (C), L_3 is a bond; methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; and G_1 is OR_9 , or CO_2R_9 .

10 [00568] In further or alternative embodiments of compounds of Formula (C), L_3 is a methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; pentan-1,5-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00569] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; or ethan-1,2-diyl.

[00570] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl.

15 [00571] In further or alternative embodiments of compounds of Formula (C), L_3 is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl.

20 [00572] In further or alternative embodiments of compounds of Formula (C), L_3 is 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; or 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is OR_9 , or CO_2R_9 .

[00573] In further or alternative embodiments of compounds of Formula (C), is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cyclic alkyl.

25 [00574] In further or alternative embodiments of compounds of Formula (C), L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

30 [00575] In further or alternative embodiments of compounds of Formula (C), L_4 is a bond, methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

40 [00576] In further or alternative embodiments of compounds of Formula (C), L_4 is a bond, ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl;

cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00577] In further or alternative embodiments of compounds of Formula (C), L_3 is a bond, methandiyl; ethan-1,2-diyl; X is a bond, $-C(=O)$, $-CR_9(OR_9)$, or $-C(O)NR_9$; L_4 is a bond, methandiyl; ethan-1,1-diyl; ethan-1,2-diyl; 5 propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; propan-1,3-diyl; butan-1,1-diyl; butan-1,2-diyl; butan-2,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; pentan-1,5-diyl; hexan-3,3-diyl; hexan-1,6-diyl; heptan-4,4-diyl; pentan-3,3-diyl, cyclopropan-1,1-diyl; cyclopropan-1,2-diyl; 10 cyclobutan-1,1-diyl; cyclobutan-1,3-diyl; cyclopentan-1,1-diyl; cyclopentan-1,3-diyl; cyclohexan-1,1-diyl; cyclohexan-1,4-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; tetrahydrothiopyran-4,4-diyl.

[00578] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00579] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; X is a bond; L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00580] In further or alternative embodiments of compounds of Formula (C), L_3 is unsubstituted alkyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00581] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00582] In further or alternative embodiments of compounds of Formula (C), L_3 is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.

[00583] In further or alternative embodiments of compounds of Formula (C), L_3 is 2-methyl-propan-1,2-diyl; or 2-ethyl-butan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-C(O)OR_9$.


[00584] In further or alternative embodiments of compounds of Formula (C), L_3 is unsubstituted alkyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.

[00585] In further or alternative embodiments, L_3 is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; pentan-1,5-diyl; or hexan-1,6-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.


5 [00586] In further or alternative embodiments of compounds of Formula (C), L_3 is propan-1,2-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; butan-1,2-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.

10 [00587] In further or alternative embodiments of compounds of Formula (C), L_3 is 2-methyl-propan-1,2-diyl; 2-ethyl-butan-1,2-diyl; X is a bond; L_4 is a bond; and G_1 is $-OR_9$.


[00588] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{C}(\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{isopropyl})\text{H}-$, $-\text{CH}_2\text{C}(\text{tert-butyl})\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_3)_2-$,



[00589] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{C}(\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_3)_2-$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$,





[00590] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)\text{H}-$, $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$,




[00591] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_3)_2-$, or $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2-$.


[00592] In further or alternative embodiments of compounds of Formula (C), R₇ is selected from among , and






[00593] In further or alternative embodiments of compounds of Formula (C), R₇ is selected from among ,






[00594] In further or alternative embodiments of compounds of Formula (C), R₇ is selected from among




[00595] In further or alternative embodiments of compounds of Formula (C), R_7 is selected from among




5

[00596] In further or alternative embodiments of compounds of Formula (C), R_7 is selected from among




10


[00597] In further or alternative embodiments of compounds of Formula (C), R_7 is selected from among



[00598] In further or alternative embodiments of compounds of Formula (C), R_7 is selected from among



15



and

[00599] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; or ethan-1,2-diyl; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; cycloheptan-1,1-diyl; piperidin-4,4-diyl; tetrahydropyran-4,4-diyl; or tetrahydrothiopyran-4,4-diyl.

[00600] In further or alternative embodiments of compounds of Formula (C), X is a bond; and L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, or a substituted or unsubstituted cyclic alkyl.

[00601] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; or ethan-1,2-diyl; X is a bond; and L_4 is methandiyl; ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-1,1-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[00602] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; X is a bond; and L_4 is ethan-1,1-diyl; propan-1,1-diyl; 2-methylpropan-1,1-diyl; 2,2-dimethylpropan-1,1-diyl; propan-2,2-diyl; butan-1,1-diyl; butan-2,2-diyl; pentan-2,2-diyl; pentan-3,3-diyl; hexan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[00603] In further or alternative embodiments of compounds of Formula (C), L_4 is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and G_1 is $-\text{CO}_2\text{R}_9$.


[00604] In further or alternative embodiments of compounds of Formula (C), L_3 is methandiyl; X is a bond; and L_4 is propan-1,1-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl.

[00605] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[00606] In another aspect, described herein are compounds of Formula (F). Compounds of Formula (F), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, glucuronide metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent or leukotriene mediated conditions or diseases, including, but not limited to, asthma, myocardial infarction, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel

disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[00607] In a further or alternative aspect, provided herein are compounds of Formula (F) as follows:



- 5 wherein, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;
- 10 Y is H, $-CO_2H$, tetrazolyl, $-NHS(=O)_2 R_{3b}$, $S(=O)_2 N(R_4)_2$, OH, $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-C(O)NHS(=O)_2 R_{3b}$, $-S(=O)_2 NHC(O)R_4$, CN, $N(R_4)_2$, $-N(R_4)C(O)R_4$, $-C(=NR_3)N(R_4)_2$, $-NR_4C(=NR_3)N(R_4)_2$, $-NR_4C(=CHR_3)N(R_4)_2$, $-C(O)NR_4C(=NR_3)N(R_4)_2$, $-C(O)NR_4C(=CHR_3)N(R_4)_2$, $-CO_2 R_{3b}$, $-C(O)R_4$, $-CON(R_4)_2$, $-SR_{3b}$, $-S(=O)R_{3b}$, $-S(=O)_2 R_{3b}$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted alkenyl), $-L_1$ -(substituted or unsubstituted alkynyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted aryl) or $-L_1-C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=NR_4)N(R_4)_2$, $-L_1-NR_4C(=CHR_3)N(R_4)_2$;
- 15 where L_1 is a bond, a substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, substituted or unsubstituted heteroalkenyl, a substituted or unsubstituted heteroalkynyl, or substituted or unsubstituted aryl;
- 20 each R_3 is independently selected from H, $-S(=O)_2 R_8$, $-S(=O)_2 NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl; each R_{3b} is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; each R_4 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_4 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;
- 25 R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, C(O), $-CH(OH)$, $-(\text{substituted or unsubstituted } C_1-C_6 \text{ alkyl})$, or $-(\text{substituted or unsubstituted } C_2-C_6 \text{ alkenyl})$;
- 30 R_7 is H or substituted or unsubstituted alkyl;
- R_5 is H, halogen, $-N_3$, $-CN$, $-NO_2$, $-L_6$ -(substituted or unsubstituted C_1 - C_6 alkyl), $-L_6$ -(substituted or unsubstituted C_2 - C_6 alkenyl), $-L_6$ -(substituted or unsubstituted heteroaryl), or $-L_6$ -(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-NHC(O)NH$, or $-C(O)NH$;
- 35

R_{11} is a (substituted or unsubstituted heteroaryl) or (substituted or unsubstituted heterocycloalkyl), and R_{12} is $L_8-L_9-R_{13}$, wherein L_8 is a bond, (substituted or unsubstituted C_1-C_6 alkyl), or (substituted or unsubstituted C_2-C_4 alkenyl); L_9 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)NH$, $-OC(O)O$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; R_{13} is H, (substituted or unsubstituted C_1-C_6 alkyl), (substituted or unsubstituted C_3-C_6 cycloalkyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heteroaryl), or (substituted or unsubstituted heterocycloalkyl); or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00608] For any and all of the embodiments of Formula (F), substituents can be selected from among from a subset of the listed alternatives.

[00609] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$. In other embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S(O)_m$.

[00610] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, and $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1-C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1-C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00611] In some embodiments, Z is selected from $-S-$, $-[C(R_2)_2]_n C(R_1)_2 S-$, and $-SC(R_1)_2 [C(R_2)_2]_n-$.

[00612] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00613] In some embodiments, each R_1 is independently H, CF_3 , or an optionally substituted C_1-C_6 alkyl.

[00614] In some embodiments, each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1-C_6 alkyl.

[00615] In some embodiments, Z is $-S-$ or $[C(R_2)_2]_n C(R_1)_2 S-$.

[00616] In some embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S-$.

[00617] In some embodiments, Z is $-S-$.

[00618] In some embodiments, Z is CH_2S- .

[00619] In some embodiments, Z is $-S-$, $-SCH_2-$, $-CH_2S-$, or $-CH(CH_3)S-$.

[00620] In some embodiments, Z is $-S-$ or $-CH_2S-$.

[00621] In further or alternative embodiments of compounds of Formula (F), L_1 is a bond, a substituted or unsubstituted alkyl, a substituted or unsubstituted heterocycloalkyl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, a substituted or unsubstituted heteroalkyl, or substituted or unsubstituted aryl.

[00622] In further or alternative embodiments of compounds of Formula (F), L_1 is a bond, or a substituted or unsubstituted alkyl. In further or alternative embodiments, L_1 is a bond.

[00623] In further or alternative embodiments of compounds of Formula (F), R_7 a substituted alkyl.

[00624] In further or alternative embodiments of compounds of Formula (F), R_7 a mono-substituted alkyl.

[00625] In further or alternative embodiments of compounds of Formula (F), R_7 a bi-substituted alkyl.

[00626] In further or alternative embodiments of compounds of Formula (F), the substituent on R_7 is selected from OH, C_1-C_6 alkoxy, $C(O)OH$, $C(O)O(C_1-C_6$ alkyl).

[00627] In further or alternative embodiments of compounds of Formula (F), R₆ is L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(O), -CH(OH), or -(substituted or unsubstituted C₁-C₆ alkyl).

5 [00628] In further or alternative embodiments of compounds of Formula (F), R₆ is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

10 [00629] In further or alternative embodiments of compounds of Formula (F), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; ; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

15 [00630] In further or alternative embodiments of compounds of Formula (F), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; ; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

20 [00631] In further or alternative embodiments of compounds of Formula (F), R₆ is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butoxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

25 [00632] In further or alternative embodiments of compounds of Formula (F), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

30 [00633] In further or alternative embodiments of compounds of Formula (F), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

35 [00634] In further or alternative embodiments of compounds of Formula (F), R₆ is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00635] In further or alternative embodiments of compounds of Formula (F), R₆ is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-

butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00636] In further or alternative embodiments of compounds of Formula (F), R_6 is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl;

propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00637] In further or alternative embodiments of compounds of Formula (F), R_6 is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-


butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00638] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

Compounds of Formula (H):

[00639] In another aspect are compounds of Formula (H), pharmaceutically acceptable salts, pharmaceutically acceptable N-oxides, pharmaceutically active metabolites, pharmaceutically acceptable prodrugs, and pharmaceutically acceptable solvates thereof, which antagonize or inhibit FLAP and may be used to treat patients suffering from leukotriene-dependent conditions or diseases, including, but not limited to, asthma, chronic obstructive pulmonary disease, pulmonary hypertension, interstitial lung fibrosis, rhinitis, arthritis, allergy, psoriasis, inflammatory bowel disease, adult respiratory distress syndrome, myocardial infarction, aneurysm, stroke, cancer, endotoxic shock, proliferative disorders and inflammatory conditions.

[00640] In a further or alternative aspect, provided herein are compounds of Formula (H) as follows:



wherein,

Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_1 on the same carbon may join to form a carbonyl ($=O$); and each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl or two R_2 on the same carbon may join to form a carbonyl ($=O$); m is 0, 1 or 2; each n is independently 0, 1, 2, or 3;

Y is $-CO_2H$, $-CONH_2$, $-C(=O)N(R_{4b})_2$, CO_2R_{4b} , $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-C(=NOH)R_{4b}$,

$C(=NOR_{3b})R_{4b}$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted alkenyl), $-L_1$ -(substituted or unsubstituted alkynyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), or $-L_1$ -(substituted or unsubstituted aryl);

where L_1 is $-C(=O)$, CR_8OH , CR_8OMe , $C(=NOH)$, $C(=NOR_{4b})$, $C(=O)NH$, $C(=O)NR_{4b}$, $-NHC(=O)$, $NR_{4b}C(=O)$, S , $S(=O)$, $S(=O)_2$, $-NHC(=O)NH$, or $NR_{4b}C(=O)NR_{4b}$;

each R_3 is independently selected from H , $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

each R_{3b} is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl;

each R_4 is independently selected from H , substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted benzyl; or two R_4 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;

each R_{4b} is independently selected from H , substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted aryl or substituted or unsubstituted benzyl; substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl;

R_6 is H , L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O , S , $-S(=O)$, $-S(=O)_2$, $C(O)$, $-CH(OH)$, $-(substituted or unsubstituted C_1-C_6 alkyl)$, or $-(substituted or unsubstituted C_2-C_6 alkenyl)$;

R_7 is L_3 - X - L_4 - G_1 , wherein,

L_3 is a bond, or substituted or unsubstituted alkyl;

X is a bond, O , $-C(=O)$, $-CR_9(OR_9)$, S , $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NR_9$, $-NR_9S(=O)_2$, $-OC(O)NR_9$, $-NR_9C(O)O$, $-CH=NO$, $-ON=CH$, $-NR_9C(O)NR_9$, heteroaryl, aryl, $-NR_9C(=NR_{10})NR_9$, $-NR_9C(=NR_{10})$, $-C(=NR_{10})NR_9$, $-OC(=NR_{10})$, or $-C(=NR_{10})O$;

L_4 is a bond or substituted or unsubstituted alkyl;

G_1 is H , tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN , $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-OC(O)O$, $-NHC(O)NH$, $-NHC(O)O$, $-O(O)CNH$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

or G_1 is W - G_5 , where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H , tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH , $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN , $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$;

each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl;

each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, phenyl or benzyl; or two R_9 groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; and

each R_{10} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl;

R_5 is H, halogen, $-N_3$, $-CN$, $-NO_2$, $-L_6$ -(substituted or unsubstituted C_1 - C_6 alkyl), $-L_6$ -(substituted or unsubstituted C_2 - C_6 alkenyl), $-L_6$ -(substituted or unsubstituted heteroaryl), or $-L_6$ -(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, C(O), $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, $-NHC(O)NH$ -, or $-C(O)NH$;

R_{11} is L_7 - L_{10} - G_6 ; wherein L_7 is a bond, $-O$, $-S$, $-S(=O)$, $-S(=O)_2$, $-NH$, $-C(O)$, $-C(O)NH$, $-NHC(O)$, (substituted or unsubstituted C_1 - C_6 alkyl), or (substituted or unsubstituted C_2 - C_6 alkenyl);

L_{10} is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted heterocycloalkyl);

G_6 is H, CN, SCN, N_3 , NO_2 , halogen, OR_9 , $-C(=O)CF_3$, $-C(=O)R_9$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $N(R_9)_2$, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-NHC(O)O$, $-NHC(O)NH$ -, $-OC(O)O$ -, $-OC(O)NH$ -, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;

or G_6 is W - G_7 , wherein W is (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted cycloalkenyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl) and G_7 is H, halogen, CN, NO_2 , N_3 , CF_3 , OCF_3 , C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $-C_1$ - C_6 fluoroalkyl, tetrazolyl, $-NHS(=O)_2R_8$, $-S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroalkyl), $-L_5$ -(substituted or unsubstituted heteroaryl), $-L_5$ -(substituted or unsubstituted heterocycloalkyl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is a bond, $-O$, $-C(=O)$, S, $S(=O)$, $S(=O)_2$, $-NH$, $-NHC(O)O$, $-NHC(O)NH$ -, $-OC(O)O$ -, $-OC(O)NH$ -, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$; and

R_{12} is H, (substituted or unsubstituted C_1 - C_6 alkyl), (substituted or unsubstituted C_3 - C_6 cycloalkyl); or glucuronide metabolite, or solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

[00641] For any and all of the embodiments of Formula (H), substituents can be selected from among from a subset of the listed alternatives.

[00642] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, $S(O)_m C(R_1)_2 [C(R_2)_2]_n$. In other embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S(O)_m$.

[00643] In some embodiments, Z is selected from $S(O)_m$, $[C(R_2)_2]_n C(R_1)_2 S(O)_m$, and $S(O)_m C(R_1)_2 [C(R_2)_2]_n$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; and R_2 is H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl; m is 0, 1 or 2; n is 0, 1, 2, or 3.

[00644] In some embodiments, Z is selected from $-S-$, $-[C(R_2)_2]_n C(R_1)_2 S-$, and $-SC(R_1)_2 [C(R_2)_2]_n-$.

5 [00645] In some embodiments, m is 0. In further embodiments, n is 0 or 1. In further embodiments, n is 0.

[00646] In some embodiments, each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00647] In some embodiments, each R_2 is independently H, OH, OMe, CF_3 , or an optionally substituted C_1 - C_6 alkyl.

[00648] In some embodiments, Z is $-S-$ or $[C(R_2)_2]_n C(R_1)_2 S-$.

10 [00649] In some embodiments, Z is $[C(R_2)_2]_n C(R_1)_2 S-$.

[00650] In some embodiments, Z is $-S-$.

[00651] In some embodiments, Z is CH_2S- .

[00652] In some embodiments, Z is $-S-$, $-SCH_2-$, $-CH_2S-$, or $-CH(CH_3)S-$.

[00653] In some embodiments, Z is $-S-$ or $-CH_2S-$.

15 [00654] In further or alternative embodiments of compounds of Formula (H), Y is $-CO_2H$, $-CONH_2$, -

$C(=O)N(R_{4b})_2$, CO_2R_{4b} , $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-C(=NOH)R_{4b}$, $C(=NOR_{3b})R_{4b}$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted cycloalkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), -

$-L_1$ -(substituted or unsubstituted heterocycloalkyl), or $-L_1$ -(substituted or unsubstituted aryl). In further or

20 alternative embodiments of compounds of Formula (H), G_6 is W- G_7 , wherein W is (substituted or unsubstituted cycloalkyl), (substituted or unsubstituted aryl), (substituted or unsubstituted heterocycloalkyl) or a (substituted or unsubstituted heteroaryl).

[00655] In further or alternative embodiments of compounds of Formula (H), Y is $-CO_2H$, $-CONH_2$, -

$C(=O)N(R_{4b})_2$, CO_2R_{4b} , $-OR_{3b}$, $-C(=O)(C_1-C_5 \text{ fluoroalkyl})$, $-L_1$ -(substituted or unsubstituted alkyl), $-L_1$ -(substituted or unsubstituted heteroaryl), $-L_1$ -(substituted or unsubstituted heterocycloalkyl), or $-L_1$ -(substituted or

25 unsubstituted aryl).

[00656] In further or alternative embodiments of compounds of Formula (H), L_1 is $-C(=O)$, CR_8OH , CR_8OMe , $C(=O)NH$, or $-NHC(=O)$.

[00657] In further or alternative embodiments of compounds of Formula (H), R_{11} is $L_7-L_{10}-G_6$; and L_7 is a bond.

In further or alternative embodiments of compounds of Formula (H), R_6 is L_2 -(substituted or unsubstituted alkyl),

30 or L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(O)_2$, $-C(O)$, $-CH(OH)$, or substituted or unsubstituted alkyl. In further or alternative embodiments of compounds of Formula (H), L_3 is a bond.

[00658] In further or alternative embodiments of compounds of Formula (H), R_6 is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl;

35 cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl;

40 cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00659] In further or alternative embodiments of compounds of Formula (H), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; ; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00660] In further or alternative embodiments of compounds of Formula (H), R₆ is methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; ; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; or benzyl.

[00661] In further or alternative embodiments of compounds of Formula (H), R₆ is methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; or phenoxy.

[00662] In further or alternative embodiments of compounds of Formula (H), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.

[00663] In further or alternative embodiments of compounds of Formula (H), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; or cyclohexylcarbonyl.

[00664] In further or alternative embodiments of compounds of Formula (H), R₆ is *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl.


[00665] In further or alternative embodiments of compounds of Formula (H), R₆ is H; ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00666] In further or alternative embodiments of compounds of Formula (H), R₆ is ethyl; propyl; prop-2-yl; 2-methylpropyl; *tert*-butyl; 3,3-dimethylbut-1-yl; cyclobutylmethyl; benzyl; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.


[00667] In further or alternative embodiments of compounds of Formula (H), R₆ is acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethyl-propanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; *tert*-butylsulfanyl; *tert*-butylsulfinyl; or *tert*-butylsulfonyl.

[00668] In further or alternative embodiments of compounds of Formula (H), G_1 is tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN , $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$. In further or alternative embodiments of compounds of Formula (H), X is a bond, $-O-$, $-CR_9(OR_9)$, S , $-S(O)$, $-S(O)_2$, $-NR_8$, $-NHC(=O)$, aryl or $-C(=O)NH$.


[00669] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-CH_2C(CH_2CH_3)H-$, $-CH_2C(isopropyl)H-$, $-CH_2C(tert-butyl)H-$, $-CH_2C(CH_3)_2-$,



[00670] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2C(CH_3)H-$, $-CH_2C(CH_2CH_3)H-$, $-CH_2C(CH_3)_2-$, $-CH_2C(CH_2CH_3)_2-$,




[00671] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-CH_2C(CH_2CH_3)H-$,




[00672] In further or alternative embodiments of compounds of Formula (C), L_3-X-L_4 is $-CH_2C(CH_3)_2-$, or $-CH_2C(CH_2CH_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_3)_2-$. In further or alternative embodiments, L_3-X-L_4 is $-CH_2C(CH_2CH_3)_2-$.

[00673] For any and all of the embodiments (such as, e.g. Formula (A), Formula (B), Formula (C), Formula (F), and Formula (H)), substituents are selected from among a list of alternatives. For example, in one embodiment, the heterocycloalkyl of Y is selected from quinolizines, dioxines, piperidines, morpholines, thiazines, tetrahydropyridines, piperazines, oxazinanones, dihydropyrroles, dihydroimidazoles, tetrahydrofurans, dihydrooxazoles, oxiranes, pyrrolidines, pyrazolidines, dihydrothiophenones, imidazolidinones, pyrrolidinones, dihydrofuranones, dioxolanones, thiazolidines, piperidinones, tetrahydronaphyridines, tetrahydroquinolines, tetrahydrothiophenes, and thiazepanes.

[00674] In further embodiments, the heterocycloalkyl of Y is selected from the group consisting of the following structures:



By way of example only, the heterocycloalkyl of Y is selected from



[00675] In a further or alternative embodiment, the "G" group (e.g. G₁, G₂, G₄, G₅, G₆, G₇) is any group that is used to tailor the physical and biological properties of the molecule. Such tailoring/modifications are achieved using groups which modulate acidity, basicity, lipophilicity, solubility and other physical properties of the molecule. The physical and biological properties modulated by such modifications to "G" include, by way of example only, solubility, *in vivo* absorption, and *in vivo* metabolism. In addition, *in vivo* metabolism may include, by way of example only, controlling *in vivo* PK properties, off-target activities, potential toxicities associated with cyp450 interactions, drug-drug interactions, and the like. Further, modifications to "G" allow for the tailoring of the *in vivo* efficacy of the compound through the modulation of, by way of example, specific and non-specific protein binding to plasma proteins and lipids and tissue distribution *in vivo*. Additionally, such tailoring/modifications to "G" allow for the design of compounds selective for 5-lipoxygenase-activating protein over other proteins.

[00676] In further or alternative embodiments, "G" is L₂₀-Q, wherein L₂₀ is an enzymatically cleavable linker and Q is a drug, or an affinity moiety. In further or alternative embodiments, the drug includes, by way of example only, leukotriene receptor antagonists and anti-inflammatory agents. In further or alternative embodiments, the leukotriene receptor antagonists include, but are not limited to, CysLT1/CysLT2 dual antagonists and CysLT1 antagonists. In further or alternative embodiments, the affinity moiety allow for site specific binding and include, but are not limited to, antibodies, antibody fragments, DNA, RNA, siRNA, and ligands.

[00677] Any combination of the groups described above for the various variables is contemplated herein. It is understood that substituents and substitution patterns on the compounds provided herein can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be synthesized by techniques known in the art, as well as those set forth herein.

[00678] Further embodiments of Formula (A), Formula (B), Formula (C), Formula (F), and Formula (H), include, but are not limited to, compounds shown in Figures 8-11 and in Tables 9-14.

Synthesis of Compounds

[00679] Compounds described herein (e.g. compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H)), may be synthesized using standard synthetic techniques known to those of skill in the art or using methods known in the art in combination with methods described herein. In

additions, solvents, temperatures and other reaction conditions presented herein may vary according to those of skill in the art.

[00680] The starting material used for the synthesis of the compounds described herein may be synthesized or can be obtained from commercial sources, such as, but not limited to, Aldrich Chemical Co. (Milwaukee, Wis.), or Sigma Chemical Co. (St. Louis, Mo.). The compounds described herein, and other related compounds having different substituents can be synthesized using techniques and materials known to those of skill in the art, such as described, for example, in March, ADVANCED ORGANIC CHEMISTRY 4th Ed., (Wiley 1992); Carey and Sundberg, ADVANCED ORGANIC CHEMISTRY 4th Ed., Vols. A and B (Plenum 2000, 2001), and Green and Wuts, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS 3rd Ed., (Wiley 1999) (all of which are incorporated by reference in their entirety). General methods for the preparation of compound as disclosed herein may be derived from known reactions in the field, and the reactions may be modified by the use of appropriate reagents and conditions, as would be recognized by the skilled person, for the introduction of the various moieties found in the formulae as provided herein. As a guide the following synthetic methods may be utilized.

Formation of Covalent Linkages by Reaction of an Electrophile with a Nucleophile

[00681] The compounds described herein can be modified using various electrophiles or nucleophiles to form new functional groups or substituents. Table I, entitled "Examples of Covalent Linkages and Precursors Thereof" lists selected examples of covalent linkages and precursor functional groups which yield and can be used as guidance toward the variety of electrophiles and nucleophiles combinations available. Precursor functional groups are shown as electrophilic groups and nucleophilic groups.

Table I: Examples of Covalent Linkages and Precursors Thereof

Covalent Linkage Product	Electrophile	Nucleophile
Carboxamides	Activated esters	amines/anilines
Carboxamides	acyl azides	amines/anilines
Carboxamides	acyl halides	amines/anilines
Esters	acyl halides	alcohols/phenols
Esters	acyl nitriles	alcohols/phenols
Carboxamides	acyl nitriles	amines/anilines
Imines	Aldehydes	amines/anilines
Hydrazones	aldehydes or ketones	Hydrazines
Oximes	aldehydes or ketones	Hydroxylamines
Alkyl amines	alkyl halides	amines/anilines
Esters	alkyl halides	carboxylic acids
Thioethers	alkyl halides	Thiols
Ethers	alkyl halides	alcohols/phenols
Thioethers	alkyl sulfonates	Thiols
Esters	alkyl sulfonates	carboxylic acids
Ethers	alkyl sulfonates	alcohols/phenols
Esters	Anhydrides	alcohols/phenols
Carboxamides	Anhydrides	amines/anilines
Thiophenols	aryl halides	Thiols
Aryl amines	aryl halides	Amines
Thioethers	Azindines	Thiols
Boronate esters	Boronates	Glycols

Carboxamides	carboxylic acids	amines/anilines
Esters	carboxylic acids	Alcohols
hydrazines	Hydrazides	carboxylic acids
N-acylureas or Anhydrides	carbodiimides	carboxylic acids
Esters	diazoalkanes	carboxylic acids
Thioethers	Epoxides	Thiols
Thioethers	haloacetamides	Thiols
Ammotriazines	halotriazines	amines/anilines
Triazinyl ethers	halotriazines	alcohols/phenols
Amidines	imido esters	amines/anilines
Ureas	Isocyanates	amines/anilines
Urethanes	Isocyanates	alcohols/phenols
Thioureas	isothiocyanates	amines/anilines
Thioethers	Maleimides	Thiols
Phosphite esters	phosphoramidites	Alcohols
Silyl ethers	silyl halides	Alcohols
Alkyl amines	sulfonate esters	amines/anilines
Thioethers	sulfonate esters	Thiols
Esters	sulfonate esters	carboxylic acids
Ethers	sulfonate esters	Alcohols
Sulfonamides	sulfonyl halides	amines/anilines
Sulfonate esters	sulfonyl halides	phenols/alcohols

Use of Protecting Groups


[00682] In the reactions described, it may be necessary to protect reactive functional groups, for example hydroxy, amino, imino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Protecting groups are used to block some or all reactive moieties and prevent such groups from participating in chemical reactions until the protective group is removed. It is preferred that each protective group be removable by a different means. Protective groups that are cleaved under totally disparate reaction conditions fulfill the requirement of differential removal. Protective groups can be removed by acid, base, and hydrogenolysis. Groups such as trityl, dimethoxytrityl, acetal and t-butyldimethylsilyl are acid labile and may be used to protect carboxy and hydroxy reactive moieties in the presence of amino groups protected with Cbz groups, which are removable by hydrogenolysis, and Fmoc groups, which are base labile. Carboxylic acid and hydroxy reactive moieties may be blocked with base labile groups such as, but not limited to, methyl, ethyl, and acetyl in the presence of amines blocked with acid labile groups such as t-butyl carbamate or with carbamates that are both acid and base stable but hydrolytically removable.

[00683] Carboxylic acid and hydroxy reactive moieties may also be blocked with hydrolytically removable protective groups such as the benzyl group, while amine groups capable of hydrogen bonding with acids may be blocked with base labile groups such as Fmoc. Carboxylic acid reactive moieties may be protected by conversion to simple ester compounds as exemplified herein, or they may be blocked with oxidatively-removable protective groups such as 2,4-dimethoxybenzyl, while co-existing amino groups may be blocked with fluoride labile silyl carbamates.

[00684] Allyl blocking groups are useful in the presence of acid- and base- protecting groups since the former are stable and can be subsequently removed by metal or pi-acid catalysts. For example, an allyl-blocked carboxylic acid can be deprotected with a Pd₀-catalyzed reaction in the presence of acid labile t-butyl carbamate or base-labile acetate amine protecting groups. Yet another form of protecting group is a resin to which a

compound or intermediate may be attached. As long as the residue is attached to the resin, that functional group is blocked and cannot react. Once released from the resin, the functional group is available to react.

[00685] Typically blocking/protecting groups may be selected from:



- 5 [00686] Other protecting groups, plus a detailed description of techniques applicable to the creation of protecting groups and their removal are described in Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, and Kocienski, *Protective Groups*, Thieme Verlag, New York, NY, 1994, which are incorporated herein by reference in their entirety.

- 10 [00687] Indole containing compounds can be prepared using standard literature procedures such as those found in Katritzky, "Handbook of Heterocyclic Chemistry" Pergamon Press, Oxford, 1986; Pindur *et al*, *J. Heterocyclic Chem.*, vol 25, 1, 1987, and Robinson "The Fisher Indole Synthesis", John Wiley & Sons, Chichester, New York, 1982, each of which is herein incorporated by reference in their entirety.

- 15 [00688] A non-limiting example of the synthetic approach toward indole compounds described herein (e.g. compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H)), is shown in Scheme I in Figure 1, wherein a 4-substituted anilines (I-1) can be converted to the corresponding hydrazine (I-2) using standard methodology. Reaction of hydrazine (I-2) with an appropriately substituted ketone (I-3) under standard Fisher-indolization conditions yields the indole (I-4). Indole (I-6) results from the N-alkylation of (I-4) with a benzyl halide (I-5) (or tosylate (OTs) or mesylate (OMs)) in a solvent such as tetrahydrofuran (THF) or dimethylformamide (DMF) in the presense of a base such as NaH. In the case where
- 20 the 5-substituent on the indole ring is methoxy (i.e. Z is MeO) the methyl group can be removed under standard conditions, for example using BBr₃, in a solvent such as CH₂Cl₂ to afford the phenol (I-7). This phenol can be alkylated using an electrophile (YX) to provide the alkylated product (I-8). Alternatively, in the case when the 5-substituent on the indole ring is, for example, a halide or triflate (OTf, I-7) it can be coupled with a wide variety of reagents using standard metal mediated coupling reactions well known to those skilled in the art of organic
- 25 synthesis to afford alternate compounds of structure (I-6). Such chemistry is described in Comprehensive Organometallic Chemistry II, vol 12, Pergamon, edited by Abel, Stone and Wilkinson. The Z substituent of the indole (I-6) can be further modified using standard chemical procedures. In addition, when R₇ or R₆ is a bromo or iodine, standard cross coupling reactions allow the introduction of a variety of functional groups using procedures well known to those practiced in the art of organic synthesis. Furthermore, when R₇ is H, it is possible, under

certain conditions, to regioselectively lithiate using a strong base such as nBuLi and then condense the anion with an electrophile to introduce substituents at C-2 (see Hasan *et al*, *J. Org. Chem.*, 46, 157-164, 1981).

[00689] Another non-limiting example of the synthetic approach toward indole compounds described herein is shown in Scheme II in Figure 2. Commencing with the hydrazine I-2, N-alkylation with a benzyl halide (or tosylate or mesylate; I-5) using the conditions described above, provides the hydrazine derivative (II-1). Reaction with an appropriately substituted ketone (I-3) using standard Fisher indolization conditions provides the indole (I-6).

[00690] Another non-limiting example of the synthetic approach toward indole compounds described herein is shown in reaction Scheme III in Figure 2, wherein 3-H-indoles (III-1) can be prepared directly using the procedures described above or, alternatively, they can be prepared from 3-thioindoles by treatment with moist AlCl₃ in a solvent such as CH₂Cl₂. Functionalization at the 3-position can be achieved using a variety of reactions and procedures to allow the introduction of a wide range of substituents. By way of example only, acylation using an acid chloride (or anhydride) in the presence of a Lewis acid such as AlCl₃, allows for the introduction of acyl groups (I-6; R₆ = C(O)R') see Murakami *et al. Heterocycles*, v14, 1939-1941, 1980 and references cited therein. Commencing with (III-1), and using, by way of example only, sulfenic chlorides in a suitable solvent, compounds of general structure (III-2) wherein R₆ is SR'' can be prepared (Raban, *J. Org. Chem.*, v45, 1688, 1980). Similar chemistry using indole (III-3) can be performed or, alternatively, diaryldisulfides in the presence of a base such as NaH in DMF can be used to generate (III-4) (Atkinson *et al, Synthesis*, 480-481, 1988). The reaction of electron deficient olefins with 3-H indoles (III-1) or (III-3) in the presence of a Lewis acid (such as Yb(OTf)₃·3H₂O) allows the installation of 3-alkyl substituents of general structure (III-2) or (III-4) (where R₆ is a substituted alkyl group; see Harrington and Kerr, *Synlett*, 1047-1048, 1996). Alternatively, indole (III-3) can be reacted with benzyl derivatives (I-5) in warm DMF to yield (III-4) where R₆ is a substituted benzyl group (Jacobs *et al, J. Med. Chem.*, v36, 394-409, 1993).

Further Synthesis of Indole and Indole-Type Compounds

[00691] Additional non-limiting examples of the synthetic strategy toward indole or indole-like scaffolds for compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), include modifications to various syntheses of indoles, including, but not limited to; Batcho-Leimgruber Indole Synthesis, Reissert Indole Synthesis, Hegedus Indole Synthesis, Fukuyama Indole Synthesis, Sugawara Indole Synthesis, Bischler Indole Synthesis, Gassman Indole Synthesis, Fischer Indole Synthesis, Japp-Klingemann Indole Synthesis, Buchwald Indole Synthesis, Larock Indole Synthesis, Bartoli Indole Synthesis, Castro Indole Synthesis, Hemetsberger Indole Synthesis, Mori-Ban Indole Synthesis, Madelung Indole Synthesis, Nenitzescu Indole Synthesis, and other unnamed reactions. Non-limiting examples of such synthetic methods are shown in Figures 3-7.

Further Forms of Compounds

[00692] Compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H) can be prepared as a pharmaceutically acceptable acid addition salt (which is a type of a pharmaceutically acceptable salt) by reacting the free base form of the compound with a pharmaceutically acceptable inorganic or organic acid, including, but not limited to, inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid metaphosphoric acid, and the like; and organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic

acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, p-toluenesulfonic acid, tartaric acid, trifluoroacetic acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, arylsulfonic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedithionylsulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, and muconic acid.

[00693] Alternatively, compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), can be prepared as a pharmaceutically acceptable base addition salts (which is a type of a pharmaceutically acceptable salt) by reacting the free acid form of the compound with a pharmaceutically acceptable inorganic or organic base, including, but not limited to organic bases such as ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like and inorganic bases such as aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, sodium hydroxide, and the like.

[00694] Compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), can be prepared as a pharmaceutically acceptable salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, for example an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base. In addition, the salt forms of the disclosed compounds can be prepared using salts of the starting materials or intermediates.

[00695] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be conveniently prepared or formed during the processes described herein. By way of example only, hydrates of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be conveniently prepared by recrystallization from an aqueous/organic solvent mixture, using organic solvents including, but not limited to, dioxane, tetrahydrofuran or methanol. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

[00696] Compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be in various forms, including but not limited to, amorphous forms, milled forms and nanoparticulate forms. In addition, compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), include crystalline forms, also known as polymorphs. Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound.

Polymorphs usually have different X-ray diffraction patterns, infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature may cause a single crystal form to dominate.

[00697] Compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in unoxidized form can be prepared from corresponding N-oxides of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), by treating with a reducing agent, such as, but not limited to, sulfur, sulfur dioxide, triphenyl phosphine, lithium borohydride, sodium borohydride, phosphorus trichloride, tribromide, or the like in a suitable inert organic solvent, such as, but not limited to, acetonitrile, ethanol, aqueous dioxane, or the like at 0 °C to 80 °C.

[00698] A "prodrug" refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug.

[00699] Compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be prepared as prodrugs. Prodrugs are generally drug precursors that, following administration to a subject and subsequent absorption, are converted to an active, or a more active species via some process, such as conversion by a metabolic pathway. Some prodrugs have a chemical group present on the prodrug that renders it less active and/or confers solubility or some other property to the drug. Once the chemical group has been cleaved and/or modified from the prodrug the active drug is generated.

[00700] An example, without limitation, of a prodrug would be a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), which is administered as an ester (the "prodrug") to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but which then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water-solubility is beneficial. A further example of a prodrug might be a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety.

[00701] Prodrugs may be designed as reversible drug derivatives, for use as modifiers to enhance drug transport to site-specific tissues. The design of prodrugs may increase the effective water solubility of the therapeutic compound for targeting to regions where water is the principal solvent. See, e.g., Fedorak et al., *Am. J. Physiol.*, 269:G210-218 (1995); McLoed et al., *Gastroenterol.*, 106:405-413 (1994); Hochhaus et al., *Biomed. Chrom.*, 6:283-286 (1992); J. Larsen and H. Bundgaard, *Int. J. Pharmaceutics*, 37, 87 (1987); J. Larsen et al., *Int. J. Pharmaceutics*, 47, 103 (1988); Sinkula et al., *J. Pharm. Sci.*, 64:181-210 (1975); T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series; and Edward B. Roche, *Bioreversible Carriers in Drug Design*, American Pharmaceutical Association and Pergamon Press, 1987, all incorporated herein in their entirety.

[00702] Additionally, prodrug derivatives of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be prepared by methods known to those of ordinary skill in the art (e.g., for further details see Saulnier et al., (1994), *Bioorganic and Medicinal Chemistry Letters*, Vol. 4, p. 1985). By way of example only, appropriate prodrugs can be prepared by reacting a non-derivatized compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with a suitable carbamylating agent, such as, but not limited to, 1,1-acyloxyalkylcarbanochloridate, para-nitrophenyl carbonate, or the like. Prodrug forms of the herein described compounds, wherein the prodrug is metabolized *in vivo* to produce a derivative as set forth herein are included within the scope of the claims. Indeed, some of the herein-described compounds may be a prodrug for another derivative or active compound.

[00703] Sites on the aromatic ring portion of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be susceptible to various metabolic reactions, therefore incorporation of appropriate substituents on the aromatic ring structures, such as, by way of example only, halogens can reduce, minimize or eliminate this metabolic pathway.

5 [00704] The compounds described herein may be labeled isotopically (e.g. with a radioisotope) or by another other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

[00705] Compounds described herein include isotopically-labeled compounds, which are identical to those recited in the various formulae and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, fluorine and chlorine, such as, for example, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{18}F , ^{36}Cl , respectively. Certain isotopically-labeled compounds described herein, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Further, substitution with isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements.

10

15

[00706] In additional or further embodiments, the compounds described herein are metabolized upon administration to an organism in need to produce a metabolite that is then used to produce a desired effect, including a desired therapeutic effect.

20

[00707] The compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may possess one or more stereocenters and each center may exist in the R or S configuration. The compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. Compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds, separating the diastereomers and recovering the optically pure enantiomers. While resolution of enantiomers can be carried out using covalent diastereomeric derivatives of the compounds described herein, dissociable complexes are preferred (e.g., crystalline diastereomeric salts). Diastereomers have distinct physical properties (e.g., melting points, boiling points, solubilities, reactivity, etc.) and can be readily separated by taking advantage of these dissimilarities. The diastereomers can be separated by chiral chromatography, or preferably, by separation/resolution techniques based upon differences in solubility. The optically pure enantiomer is then recovered, along with the resolving agent, by any practical means that would not result in racemization. A more detailed description of the techniques applicable to the resolution of stereoisomers of compounds from their racemic mixture can be found in Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981, herein incorporated by reference in its entirety.

25

30

35

[00708] Additionally, the compounds and methods provided herein may exist as geometric isomers. The compounds and methods provided herein include all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the appropriate mixtures thereof. In some situations, compounds may exist as tautomers. All tautomers are included within the formulas described herein are provided by compounds and methods herein. In additional

40

embodiments of the compounds and methods provided herein, mixtures of enantiomers and/or diastereoisomers, resulting from a single preparative step, combination, or interconversion may also be useful for the applications described herein.

[00709] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent

addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described herein can be conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

[00710] The screening and characterization of the pharmaceutically acceptable salts, polymorphs and/or solvates may be accomplished using a variety of techniques including, but not limited to, thermal analysis, x-ray diffraction, spectroscopy, vapor sorption, and microscopy. Thermal analysis methods address thermo chemical degradation or thermo physical processes including, but not limited to, polymorphic transitions, and such methods are used to analyze the relationships between polymorphic forms, determine weight loss, to find the glass transition temperature, or for excipient compatibility studies. Such methods include, but are not limited to, Differential scanning calorimetry (DSC), Modulated Differential Scanning Calorimetry (MDSC), Thermogravimetric analysis (TGA), and Thermogravimetric and Infrared analysis (TG/IR). X-ray diffraction methods include, but are not limited to, single crystal and powder diffractometers and synchrotron sources. The various spectroscopic techniques used include, but are not limited to, Raman, FTIR, UV-VIS, and NMR (liquid and solid state). The various microscopy techniques include, but are not limited to, polarized light microscopy, Scanning Electron Microscopy (SEM) with Energy Dispersive X-Ray Analysis (EDX), Environmental Scanning Electron Microscopy with EDX (in gas or water vapor atmosphere), IR microscopy, and Raman microscopy.

[00711] Throughout the specification, groups and substituents thereof can be chosen by one skilled in the field to provide stable moieties and compounds.

Certain Chemical Terminology

[00712] Unless otherwise stated, the following terms used in this application, including the specification and claims, have the definitions given below. It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Definition of standard chemistry terms may be found in reference works, including Carey and Sundberg "ADVANCED ORGANIC CHEMISTRY 4TH ED." Vols. A (2000) and B (2001), Plenum Press, New York. Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant DNA techniques and pharmacology, within the skill of the art are employed. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, use of the term "including" as well as other forms, such as "include", "includes," and "included," is not limiting.

[00713] An "alkyl" group refers to an aliphatic hydrocarbon group. The alkyl moiety may be a "saturated alkyl" group, which means that it does not contain any units of unsaturation (e.g. carbon-carbon double bond(s) or carbon-carbon triple bond(s)). The alkyl moiety may also be an "unsaturated alkyl" moiety, which means that it

contains at least one unit of unsaturation (e.g. carbon-carbon double bond(s) or carbon-carbon triple bond(s)).

The alkyl moiety, whether saturated or unsaturated, may be branched, straight chain, or cyclic.

[00714] The "alkyl" moiety may have 1 to 10 carbon atoms (whenever it appears herein, a numerical range such as "1 to 10" refers to each integer in the given range; e.g., "1 to 10 carbon atoms" means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the present definition also covers the occurrence of the term "alkyl" where no numerical range is designated). The alkyl group could also be a "lower alkyl" having 1 to 6 carbon atoms. The alkyl group of the compounds described herein may be designated as "C₁-C₄ alkyl" or similar designations. By way of example only, "C₁-C₄ alkyl" indicates that there are one to four carbon atoms in the alkyl chain, *i.e.*, the alkyl chain is selected from the group consisting of methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tertiary butyl, 2-methyl-butyl, 2-ethyl-butyl, 3-propyl-butyl, pentyl, neo-pentyl, 2-propyl-pentyl, hexyl, propenyl, butenyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and the like. Alkyl groups can be substituted or unsubstituted. Depending on the structure, an alkyl group can be a monoradical or a diradical (*i.e.*, an alkylene group, such as, but not limited to, methandiyl, ethan-1,2-diyl, propan-1,2-diyl, propan-2,2-diyl, butan-1,2-diyl, isobutan-1,2-diyl, 2-methyl-butan-1,2-yl, 2-ethyl-butan-1,2-diyl, 3-propyl-butan-1,2-diyl, pentan-1,2-diyl, 2-propyl-pentan-1,2-diyl, propan-2,2-diyl, pentan-3,3-diyl, and the like).

[00715] As used herein, C₁-C_x includes C₁-C₂, C₁-C₃ . . . C₁-C_x. C₁-C_x refers to the number of carbon atoms that make up the moiety to which it designates (excluding optional substituents).

[00716] An "alkoxy" group refers to a (alkyl)O- group, where alkyl is as defined herein.

[00717] The term "alkylamine" refers to the -N(alkyl)_xH_y group, where x and y are selected from the group x=1, y=1 and x=2, y=0. When x=2, the alkyl groups, taken together, can optionally form a cyclic ring system.

[00718] The term "alkenyl" refers to a type of alkyl group in which the first two atoms of the alkyl group form a double bond that is not part of an aromatic group. That is, an alkenyl group begins with the atoms -C(R)=C(R)₂, wherein R refers to the remaining portions of the alkenyl group, which may be the same or different. Non-limiting examples of an alkenyl group include -CH=CH₂, -C(CH₃)=CH₂, -CH=CHCH₃, -CH=C(CH₃)₂ and -C(CH₃)=CHCH₃. The alkenyl moiety may be branched, straight chain, or cyclic (in which case, it would also be known as a "cycloalkenyl" group). The "R" portion of the alkenyl moiety may be branched, straight chain, or cyclic. Two "R" groups on adjacent carbon atoms of the alkenyl moiety may together form a ring (in which case, it would be known as a "cycloalkenyl" group). A "lower alkenyl" refers to an alkenyl having 2 to 6 carbons. Alkenyl groups can be substituted or unsubstituted. Depending on the structure, an alkenyl group can be a monoradical or a diradical (*i.e.*, an alkenylene group).

[00719] The term "alkynyl" refers to a type of alkyl group in which the first two atoms of the alkyl group form a triple bond. That is, an alkynyl group begins with the atoms -C≡C-R, wherein R refers to the remaining portions of the alkynyl group, which may be the same or different. Non-limiting examples of an alkynyl group include -C≡CH, -C≡CCH₃ and -C≡CCH₂CH₃. The "R" portion of the alkynyl moiety may be branched, straight chain, or cyclic. Alkynyl groups can be substituted or unsubstituted. Depending on the structure, an alkynyl group can be a monoradical or a diradical (*i.e.*, an alkynylene group).

[00720] The terms “haloalkyl,” “haloalkenyl,” “haloalkynyl” and “haloalkoxy” refer to alkyl, alkenyl, alkynyl and alkoxy moieties that are substituted with one or more halo groups.


[00721] The terms “fluoroalkyl” and “fluoroalkoxy” refer to alkyl and alkoxy groups, respectively, which are substituted with one or more fluoro groups.

5 [00722] The terms “heteroalkyl” “heteroalkenyl” and “heteroalkynyl” refer to alkyl, alkenyl and alkynyl radicals that have one or more skeletal chain atoms selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorus or combinations thereof. The heteroatom(s) may be placed at any interior position of the heteroalkyl group. Examples include, but are not limited to, -CH₂-O-CH₃, -CH₂-CH₂-O-CH₃, -CH₂-NH-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-N(CH₃)-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. In addition, up to two heteroatoms may be consecutive, such as, by way of example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Excluding the number of heteroatoms, a “heteroalkyl” may have from 1 to 6 carbon atoms, a “heteroalkenyl” may have from 2 to 6 carbons atoms, and a “heteroalkynyl” may have from 2 to 6 carbon atoms.

[00723] “Halo”, halide”, or “halogen” refer to fluorine, chlorine, bromine, and iodine.

15 [00724] The term “carbocyclic” or “carbocycle” refers to a ring wherein each of the atoms forming the ring is a carbon atom. Carbocycle includes aryl and cycloalkyl. The term thus distinguishes carbocycle from heterocycle (“heterocyclic”) in which the ring backbone contains at least one atom which is different from carbon (*i.e.* a heteroatom). Heterocycle includes heteroaryl and heterocycloalkyl. Carbocycles and heterocycles can be optionally substituted.

20 [00725] The term “cycloalkyl” refers to a monocyclic or polycyclic aliphatic, non-aromatic radical, wherein each of the atoms forming the ring (*i.e.* skeletal atoms) is a carbon atom. Cycloalkyls may be saturated, or partially unsaturated. Cycloalkyls may be fused with an aromatic ring, and the point of attachment is at a carbon that is not an aromatic ring carbon atom. Cycloalkyl groups include groups having from 3 to 10 ring atoms. A “lower cycloalkyl” has 3 to 8 ring atoms. Illustrative examples of cycloalkyl groups include, but are not limited to, the following moieties:



, and the like. In some embodiments, cycloalkyl groups are


30 selected from among cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Cycloalkyl groups may be substituted or unsubstituted. Depending on the structure, a cycloalkyl group can be a monoradical or a diradical (*i.e.*, an cycloalkylene group, such as, but not limited to, cyclopropan-1,1-diyl, cyclopropan-1,2-diyl, cyclobutan-1,1-diyl, cyclobutan-1,3-diyl, cyclopentan-1,1-diyl, cyclopentan-1,3-diyl, cyclohexan-1,1-diyl, cyclohexan-1,4-diyl, cycloheptan-1,1-diyl, and the like).

[00726] The term "cycloalkenyl" refers to a type of cycloalkyl group that contains at least one carbon-carbon double bond in the ring and where the cycloalkenyl is attached at one of the carbon atoms of the carbon-carbon double bond. Non-limiting examples of a cycloalkenyl alkanyl group include cyclopenten-1-yl, cyclohexen-1-yl, cyclohepten-1-yl, and the like. Cycloalkenyl groups may be substituted or unsubstituted.

5 [00727] The term "aromatic" refers to a planar ring having a delocalized π -electron system containing $4n+2$ π electrons, where n is an integer. Aromatic rings can be formed from five, six, seven, eight, nine, ten, or more than ten atoms. Aromatics can be optionally substituted. The term "aromatic" includes both carbocyclic aryl ("aryl", e.g., phenyl) and heterocyclic aryl (or "heteroaryl" or "heteroaromatic") groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups.

10 [00728] As used herein, the term "aryl" refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. Aryl rings can be formed by five, six, seven, eight, nine, or more than nine carbon atoms. Aryl groups can be optionally substituted. Examples of aryl groups include, but are not limited to phenyl, and naphthalenyl. Depending on the structure, an aryl group can be a monoradical or a diradical (i.e., an arylene group).

15 [00729] The terms "heteroaryl" or, alternatively, "heteroaromatic" refers to an aryl group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. An *N*-containing "heteroaromatic" or "heteroaryl" moiety refers to an aromatic group in which at least one of the skeletal atoms of the ring is a nitrogen atom. An *N*-containing heteroaryl may be oxidized to the corresponding N-oxide. The polycyclic heteroaryl group may be fused or non-fused. Illustrative examples of heteroaryl groups include the following moieties:




and the like.

25 [00730] The term "heterocycle" refers to heteroaromatic and heteroalicyclic groups (heterocycloalkyl groups) containing one to four heteroatoms each selected from O, S and N, wherein each heterocyclic group has from 4 to 10 atoms in its ring system, and with the proviso that the ring of said group does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups include groups having only 4 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 4-membered heterocyclic group is azetidiny (derived from azetidine). An example of a 5-membered heterocyclic group is thiazolyl. An example of a 6-membered heterocyclic group is pyridyl, and an example of a 10-membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl,

30

tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperaziny, azetidiny, oxetanyl, thietanyl, homopiperidiny, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolany, pyrazoliny, dithianyl, dithiolany, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidiny, imidazoliny, imidazolidiny, 3-azabicyclo[3.1.0]hexany, 3-azabicyclo[4.1.0]heptany, 3H-indolyl and quinoliziny. Examples of aromatic heterocyclic groups are pyridiny, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyraziny, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinoliny, isoquinoliny, indolyl, benzimidazolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalaziny, pyridaziny, triaziny, isoindolyl, pteridinyl, puriny, oxadiazolyl, thiadiazolyl, furazany, benzofurazany, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazoliny, quinoxaliny, naphthyridiny, and furopyridiny. The foregoing groups, as derived from the groups listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl or imidazol-3-yl (both N-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems and ring systems substituted with one or two oxo (=O) moieties such as pyrrolidin-2-one.

[00731] A "heteroalicyclic" or "heterocycloalkyl" group refers to a cycloalkyl group that includes at least ring atom selected from nitrogen, oxygen and sulfur (i.e. at least one ring atom is a heteroatom). The radicals may be fused with an aryl or heteroaryl. Illustrative examples of heterocycloalkyl groups, also referred to as non-aromatic heterocycles, include:



forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. Other examples of heterocycloalkyls include, quinolizine, dioxine, piperidine, morpholine, thiazine, tetrahydropyridine, piperazine, oxazinanone, dihydropyrrole, dihydroimidazole, tetrahydrofuran, tetrahydropyran, dihydrooxazole, oxirane, pyrrolidine, pyrazolidine, imidazolidinone, pyrrolidinone,

dihydrofuranone, dioxolanone, thiazolidine, piperidinone, tetrahydroquinoline, tetrahydrothiophene, and thiazepane.

[00732] The term “membered ring” can embrace any cyclic structure. The term “membered” is meant to denote the number of skeletal atoms that constitute the ring. Thus, for example, cyclohexyl, pyridinyl, pyranyl and thiopyranyl are 6-membered rings and cyclopentyl, pyrrolyl, furanyl, and thienyl are 5-membered rings.

[00733] The term “ester” refers to a chemical moiety with formula -COOR, where R is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). Any hydroxy, or carboxyl side chain on the compounds described herein can be esterified. The procedures and specific groups to make such esters are known to those of skill in the art and can readily be found in reference sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein by reference in its entirety. The term “halo” or, alternatively, “halogen” means fluoro, chloro, bromo or iodo.

[00734] An “amide” is a chemical moiety with formula -C(O)NHR or -NHC(O)R, where R is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). An amide may be an amino acid or a peptide molecule attached to a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), thereby forming a prodrug. Any amine, or carboxyl side chain on the compounds described herein can be amidified. The procedures and specific groups to make such amides are known to those of skill in the art and can readily be found in reference sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein by reference in its entirety.

[00735] The term “bond” or “single bond” refers to a chemical bond between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure.

[00736] A “cyano” group refers to a -CN group.

[00737] An “isocyanato” group refers to a -NCO group.

[00738] An “isothiocyanato” group refers to a -NCS group.

[00739] “Sulfanyl” or “thio” group refers to a -S- moiety.

[00740] “Thiol” or “sulphydryl” refers to -SH.


[00741] The term “moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[00742] “Sulfinyl” or “sulfoxide” refers to -S(=O)-.

[00743] “Sulfonyl” refers to -S(=O)₂-.

[00744] “Thiocyanato” group refers to a -CNS group.

[00745] “Carboxy” refers to -CO₂H. In some cases, carboxy moieties may be replaced with a “carboxylic acid bioisostere”, which refers to a functional group or moiety that exhibits similar physical and/or chemical properties as a carboxylic acid moiety. A carboxylic acid bioisostere has similar biological properties to that of a carboxylic acid group. A compound with a carboxylic acid moiety can have the carboxylic acid moiety exchanged with a carboxylic acid bioisostere and have similar physical and/or biological properties when compared to the carboxylic acid-containing compound. For example, in one embodiment, a carboxylic acid bioisostere would ionize at physiological pH to roughly the same extent as a carboxylic acid group. Examples of bioisoteres of a carboxylic acid include, but are not limited to,



and the like.

[00746] The term “optionally substituted” or “substituted” means that the referenced group may be substituted with one or more additional group(s) individually and independently selected from alkyl, cycloalkyl,

5 heterocycloalkyl, aryl, heteroaryl, benzyl, heteroarylmethyl, hydroxy, alkoxy, fluoroalkoxy, aryloxy, thiol, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, arylsulfone, cyano, halo, carboxy, nitro, haloalkyl, fluoroalkyl, and amino, including mono- and di-alkyl amino groups, and the protected derivatives thereof. By way of example an optional substituents may may be L_sR_s , wherein L_sR_s is halo, amino, nitro, cyano, or each L_s is independently selected from a bond, -O-, -C(=O)-, -C(=O)O-, -OC(=O)-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -NHC(O)-, 10 -C(O)NH-, S(=O)₂NH-, -NHS(=O)₂-, -OC(O)NH-, -NHC(O)O-, and C₁-C₆alkyl; and each R_s is independently selected from H, alkyl, fluoroalkyl, cycloalkyl, heteroaryl, aryl, benzyl, heteroarylmethyl, or heteroalkyl. The protecting groups that may form the protective derivatives of the above substituents are known to those of skill in the art and may be found in references such as Greene and Wuts, above.

[00747] The compounds presented herein may possess one or more stereocenters and each center may exist in 15 the R or S configuration. The compounds presented herein include all diastereomeric, enantiomeric, and epimeric forms as well as the appropriate mixtures thereof. Stereoisomers may be obtained, if desired, by methods known in the art as, for example, the separation of stereoisomers by chiral chromatographic columns.

[00748] The methods and formulations described herein include the use of *N*-oxides, crystalline forms (also known as polymorphs), or pharmaceutically acceptable salts of compounds having the structure of any of 20 Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), as well as active metabolites of these compounds having the same type of activity. In some situations, compounds may exist as tautomers. All tautomers are included within the scope of the compounds presented herein. In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also 25 considered to be disclosed herein.

Certain Pharmaceutical Terminology

[00749] The term “acceptable” with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated.

[00750] The term “agonist,” as used herein, refers to a molecule such as a compound, a drug, an enzyme 30 activator or a hormone modulator which enhances the activity of another molecule or the activity of a receptor site.

[00751] The term “antagonist,” as used herein, refers to a molecule such as a compound, a drug, an enzyme inhibitor, or a hormone modulator, which diminishes, or prevents the action of another molecule or the activity of a receptor site.

35 [00752] The term “asthma” as used herein refers to any disorder of the lungs characterized by variations in pulmonary gas flow associated with airway constriction of whatever cause (intrinsic, extrinsic, or both; allergic or non-allergic). The term asthma may be used with one or more adjectives to indicate cause.

[00753] The term "bone disease," as used herein, refers to a disease or condition of the bone, including, but not limited to, inappropriate bone remodeling, loss or gain, osteopenia, osteomalacia, osteofibrosis, and Paget's disease [Garcia, "Leukotriene B4 stimulates osteoclastic bone resorption both in vitro and in vivo", *J Bone Miner Res.* 1996;11:1619-27].

5 [00754] The term "cardiovascular disease," as used herein refers to diseases affecting the heart or blood vessels or both, including but not limited to: arrhythmia; atherosclerosis and its sequelae; angina; myocardial ischemia; myocardial infarction; cardiac or vascular aneurysm; vasculitis, stroke; peripheral obstructive arteriopathy of a limb, an organ, or a tissue; reperfusion injury following ischemia of the brain, heart or other organ or tissue; endotoxic, surgical, or traumatic shock; hypertension, valvular heart disease, heart failure, abnormal blood pressure; shock; vasoconstriction (including that associated with migraines); vascular abnormality, inflammation, insufficiency limited to a single organ or tissue. [Lotzer K *et al.*, "The 5-lipoxygenase pathway in arterial wall biology and atherosclerosis", *Biochim Biophys Acta* 2005;1736:30-7; Helgadottir A *et al.*, "The gene encoding 5-lipoxygenase activating protein confers risk of myocardial infarction and stroke", *Nat Genet.* 2004 Mar;36(3):233-9. Epub 2004 Feb 8; [Heise CE, Evans JF *et al.*, "Characterization of the human cysteinyl leukotriene 2 receptor", *J Biol Chem.* 2000 Sep 29;275(39):30531-6].

15 [00755] The term "cancer," as used herein refers to an abnormal growth of cells which tend to proliferate in an uncontrolled way and, in some cases, to metastasize (spread). The types of cancer include, but is not limited to, solid tumors (such as those of the bladder, bowel, brain, breast, endometrium, heart, kidney, lung, lymphatic tissue (lymphoma), ovary, pancreas or other endocrine organ (thyroid), prostate, skin (melanoma) or hematological tumors (such as the leukemias) [Ding XZ *et al.*, "A novel anti-pancreatic cancer agent, LY293111", *Anticancer Drugs.* 2005 Jun;16(5):467-73. Review; Chen X *et al.*, "Overexpression of 5-lipoxygenase in rat and human esophageal adenocarcinoma and inhibitory effects of zileuton and celecoxib on carcinogenesis", *Clin Cancer Res.* 2004 Oct 1;10(19):6703-9].

25 [00756] The term "carrier," as used herein, refers to relatively nontoxic chemical compounds or agents that facilitate the incorporation of a compound into cells or tissues.

[00757] The terms "co-administration" or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

30 [00758] The term "dermatological disorder," as used herein refers to a skin disorder. Such dermatological disorders include, but are not limited to, proliferative or inflammatory disorders of the skin such as, atopic dermatitis, bullous disorders, collagenoses, contact dermatitis eczema, Kawasaki Disease, rosacea, Sjogren-Larsson Syndrome, urticaria [Wedi B *et al.*, "Pathophysiological role of leukotrienes in dermatological diseases: potential therapeutic implications", *BioDrugs.* 2001;15(11):729-43].

35 [00759] The term "diluent" refers to chemical compounds that are used to dilute the compound of interest prior to delivery. Diluents can also be used to stabilize compounds because they can provide a more stable environment. Salts dissolved in buffered solutions (which also can provide pH control or maintenance) are utilized as diluents in the art, including, but not limited to a phosphate buffered saline solution.

40 [00760] The terms "effective amount" or "therapeutically effective amount," as used herein, refer to a sufficient amount of an agent or a compound being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs,

symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic uses is the amount of the composition comprising a compound as disclosed herein required to provide a clinically significant decrease in disease symptoms. An appropriate “effective” amount in any individual case may be determined using techniques, such as a dose escalation study.

5 [00761] The terms “enhance” or “enhancing,” as used herein, means to increase or prolong either in potency or duration a desired effect. Thus, in regard to enhancing the effect of therapeutic agents, the term “enhancing” refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents on a system. An “enhancing-effective amount,” as used herein, refers to an amount adequate to enhance the effect of another therapeutic agent in a desired system.

10 [00762] The term “enzymatically cleavable linker,” as used herein refers to unstable or degradable linkages which may be degraded by one or more enzymes.

[00763] The terms “fibrosis” or “fibrosing disorder,” as used herein, refers to conditions that follow acute or chronic inflammation and are associated with the abnormal accumulation of cells and/or collagen and include but are not limited to fibrosis of individual organs or tissues such as the heart, kidney, joints, lung, or skin, and includes such disorders as idiopathic pulmonary fibrosis and cryptogenic fibrosing alveolitis [Charbeneau RP *et al.*, “Eicosanoids: mediators and therapeutic targets in fibrotic lung disease”, *Clin Sci (Lond)*. 2005 Jun;108(6):479-91].

[00764] The term “iatrogenic” means a leukotriene-dependent or leukotriene-mediated condition, disorder, or disease created or worsened by medical or surgical therapy.

20 [00765] The term “inflammatory disorders” refers to those diseases or conditions that are characterized by one or more of the signs of pain (*dolor*, from the generation of noxious substances and the stimulation of nerves), heat (*calor*, from vasodilatation), redness (*rubor*, from vasodilatation and increased blood flow), swelling (*tumor*, from excessive inflow or restricted outflow of fluid), and loss of function (*functio laesa*, which may be partial or complete, temporary or permanent). Inflammation takes many forms and includes, but is not limited to, inflammation that is one or more of the following: acute, adhesive, atrophic, catarrhal, chronic, cirrhotic, diffuse, disseminated, exudative, fibrinous, fibrosing, focal, granulomatous, hyperplastic, hypertrophic, interstitial, metastatic, necrotic, obliterative, parenchymatous, plastic, productive, proliferous, pseudomembranous, purulent, sclerosing, seroplastic, serous, simple, specific, subacute, suppurative, toxic, traumatic, and/or ulcerative. Inflammatory disorders further include, without being limited to those affecting the blood vessels (polyarteritis, 30 temporal arteritis); joints (arthritis: crystalline, osteo-, psoriatic, reactive, rheumatoid, Reiter’s); gastrointestinal tract (Disease,); skin (dermatitis); or multiple organs and tissues (systemic lupus erythematosus) [Harrison’s Principles of Internal Medicine, 16th Edition, Kasper DL, et al, Editors; McGraw-Hill, publishers].

[00766] The term “interstitial cystitis” refers to a disorder characterized by lower abdominal discomfort, frequent and sometimes painful urination that is not caused by anatomical abnormalities, infection, toxins, trauma or tumors [Bouchelouche K *et al.*, “The cysteinyl leukotriene D4 receptor antagonist montelukast for the treatment of interstitial cystitis”, *J Urol* 2001;166:1734].

[00767] The term “leukotriene-driven mediators,” as used herein, refers to molecules able to be produced in a patient that may result from excessive production of leukotriene stimulation of cells, such as, by way of example only, LTB₄, LTC₄, LTE₄, cysteinyl leukotrienes, monocyte inflammatory protein (MIP-1 α), interleukin-8 (IL-8), 40 interleukin-4 (IL-4), interleukin-13 (IL-13), monocyte chemoattractant protein (MCP-1), soluble intracellular

adhesion molecule (sICAM; soluble ICAM), myeloperoxidase (MPO), eosinophil peroxidase (EPO), and general inflammation molecules such as interleukin-6 (IL-6), C-reactive protein (CRP), and serum amyloid A protein (SAA).

[00768] The term “leukotriene-related mediators,” as used herein, refers to molecules able to be produced in a patient that may result from excessive production of leukotriene stimulation of cells, such as, by way of example only, LTB₄, LTC₄, LTE₄, cysteinyl leukotrienes, monocyte inflammatory protein (MIP-1 α), interleukin-8 (IL-8), interleukin-4 (IL-4), interleukin-13 (IL-13), monocyte chemoattractant protein (MCP-1), soluble intracellular adhesion molecule (sICAM; soluble ICAM), myeloperoxidase (MPO), eosinophil peroxidase (EPO), and general inflammation molecules such as interleukin-6 (IL-6), C-reactive protein (CRP), and serum amyloid A protein (SAA).

[00769] The term “leukotriene-dependent”, as used herein, refers to conditions or disorders that would not occur, or would not occur to the same extent, in the absence of one or more leukotrienes.

[00770] The term “leukotriene-mediated”, as used herein, refers to conditions or disorders that might occur in the absence of leukotrienes but can occur in the presence of one or more leukotrienes.

[00771] The term “leukotriene-responsive patient,” as used herein, refers to a patient who has been identified by either genotyping of FLAP haplotypes, or genotyping of one or more other genes in the leukotriene pathway and/or, by phenotyping of patients either by previous positive clinical response to another leukotriene modulator, including, by way of example only, zileuton (ZyfloTM), montelukast (SingulairTM), pranlukast (OnonTM), zafirlukast (AccolateTM), and/or by their profile of leukotriene-driven mediators that indicate excessive leukotriene stimulation of inflammatory cells, as likely to respond favorably to leukotriene modulator therapy.

[00772] “MAPEG” refers to “membrane associated proteins involved in eicosanoid and glutathione metabolism” and includes the following human proteins: 5-lipoxygenase activating protein (FLAP), leukotriene C₄ synthase (LTC₄ synthase), which are involved in leukotriene biosynthesis; microsomal glutathione S-transferase 1 (MGST1), MGST2, and MGST3, which are all glutathione transferases as well as glutathione dependent peroxidases; and prostaglandin E synthase (PGES), also referred to as MGST1-like 1 (MGST1-L1). (Bresell *et al.*, *FEBS Journal*, 272, 1688-1703, 2005; Jakobsson *et al.*, *J. Respir. Crit. Care Med.*, Volume 161, Number 2, February 2000, S20-S24; Jakobsson, *et al. Protein Sci.* 8: 689-692, 1998). PGES catalyzes the formation of PGE₂ from PGH₂, which in turn is generated from arachidonic acid by the prostaglandin endoperoxide synthase systems. PGES has also been referred to as p53 induced gene 12 (PIG12) because the gene expression was found to increase extensively following p53 expression (Polyak *et al.*, *Nature*, 389, 300-305, 1997). PGES isozymes have been identified: cytosolic PGES (cPGES), microsomal PGES-1 (mPGES-1) and microsomal PGES-2 (mPGES-2). cPGES is constitutively and ubiquitously expressed and selectively expressed with COX-1. mPGES-1 is induced by proinflammatory stimuli, downregulated by anti-inflammatory glucocorticoids, and functionally coupled with COX-2 in preference to COX-1.

[00773] The terms “kit” and “article of manufacture” are used as synonyms.

[00774] A “metabolite” of a compound disclosed herein is a derivative of that compound that is formed when the compound is metabolized. The term “active metabolite” refers to a biologically active derivative of a compound that is formed when the compound is metabolized (biotransformed). The term “metabolized,” as used herein, refers to the sum of the processes (including, but not limited to, hydrolysis reactions and reactions catalyzed by enzymes) by which a particular substance is changed by an organism. Thus, enzymes may produce specific

structural alterations to a compound. For example, cytochrome P450 catalyzes a variety of oxidative and reductive reactions while uridine diphosphate glucuronyltransferases (UGT) catalyze the transfer of an activated glucuronic-acid molecule to aromatic alcohols, aliphatic alcohols, carboxylic acids, amines and free sulphhydryl groups (e.g. conjugation reactions). Further information on metabolism may be obtained from The Pharmacological Basis of Therapeutics, 9th Edition, McGraw-Hill (1996). Metabolites of the compounds disclosed herein can be identified either by administration of compounds to a host and analysis of tissue samples from the host, or by incubation of compounds with hepatic cells in vitro and analysis of the resulting compounds. Both methods are well known in the art.

[00775] Conjugation reactions represent a common biotransformation reaction by which compounds that are absorbed in blood are eliminated from the body. After conjugation reactions have added an ionic hydrophilic moiety, such as glucuronic acid, sulfate, or glycine to the compound, water solubility is increased and lipid solubility is decreased enough to make elimination possible. In most cases, the major proportion of an administered drug dose is excreted as conjugates into the urine and bile. Conjugation may be preceded by other metabolic biotransformations or conjugation alone may be the fate of the drug dose.

[00776] Glucuronidation represents a major pathway which enhances the elimination of many lipophilic xenobiotics to more water-soluble compounds. The UDP-glucuronosyltransferase (UGT) family catalyzes the glucuronidation of the glycosyl group of a nucleotide sugar to an acceptor compound (aglycone) at a nucleophilic functional group of oxygen (eg, hydroxyl or carboxylic acid groups), nitrogen (eg, amines), sulfur (eg, thiols), and carbon, with the formation of a beta-D-glucuronide product.

[00777] As used herein, "acyl glucuronide" or "acylglucuronide" (either term used interchangeably) refers to a conjugate formed by glucuronidation at the carboxylic acid group of a xenobiotic. An acyl glucuronide is a type of glucuronide metabolite.

[00778] The liver is the principal organ for the metabolism and eventual elimination of xenobiotics and endobiotics from the human body either in the urine or in the bile. UGT isoforms have been identified in extrahepatic tissues including the kidney, gastrointestinal tract and brain.

[00779] In general, glucuronide metabolites that are released in the bile may be cleaved in the gastrointestinal tract by β -glucuronidases, to provide the glucuronide and the aglycon portion. The aglycon portion may be available for reabsorption from the duodenal-intestinal tract into the portal circulation, undergoing the process of enterohepatic cycling (Dobrinska, *J. Clin. Pharmacol.*, 1989, 29:577-580). Thus, the action of β -glucuronidases on glucuronide metabolites decreases the amount of xenobiotic that is eliminated at once and the levels of the xenobiotic in the blood stream oscillate due to this circulatory process. The result is that the pharmacokinetics of the initial drug dose may display (intermittent) spikes in the plasma drug concentration.

[00780] The detection of glucuronide metabolites, such as acylglucuronides, indicates an elimination pathway of the xenobiotic, and indicates that enterohepatic cycling may occur.

[00781] Enterohepatic cycling indicates that biliary excretion plays a major role in the elimination of a drug relative to renal clearance. In some embodiments, enterohepatic cycling is observed with compounds described herein. In some embodiments, compounds described herein that include a carboxylic acid moiety (e.g. G_1 moiety) are conjugated to glucuronic acid to provide acylglucuronides and participate in enterohepatic cycling.

[00782] Acylglucuronides are known to be metabolites of drugs bearing a carboxylic acid function.

Acylglucuronides are known to readily undergo hydrolysis to the parent drug under neutral or slightly alkaline conditions, with the rate of hydrolysis being dependent on the temperature. Acylglucuronides may accumulate in the blood of patients with renal failure.

5 [00783] In one aspect, acylglucuronides are formed by any of the compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), where G_1 is OH or CO_2H . In one aspect, acylglucuronides formed by any of the compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), participates in enterohepatic cycling. In one aspect, compounds described herein that include a carboxylic acid moiety in the R_7 moiety (i.e. G_1 is CO_2H) form acylglucuronide
10 metabolites.

[00784] Decreasing the rate of or amount of a compound dose that is conjugated to glucuronic acid provides a means to provide compounds that have a longer half in the blood after being absorbed and not provide (intermittent) spikes in blood concentration over time. Decreasing the rate of or amount of a compound dose that is conjugated to glucuronic acid decreases the amount of compound that is eliminated either in the bile or urine.

15 [00785] In one embodiment, compounds described herein that form acylglucuronide metabolites are identified and the steric bulk of substituents alpha to the carboxylic acid group in the compound are increased to decrease or slow the rate of reaction of the compound with UGT.

[00786] In one embodiment, compounds described herein that include a G_1 moiety that is CO_2H have a decreased rate of, or amount of glucuronidation when the alpha carbon relative to G_1 is substituted with at least one group that is sterically larger than hydrogen and methyl.

[00787] In one aspect, compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), where G_1 is CO_2H or OH, have a slower rate or reduced rate of glucuronidation when the carbon atom alpha to G_1 is substituted with at least one alpha group that is larger than a methyl group.

[00788] The term "modulate," as used herein, means to interact with a target either directly or indirectly so as to alter the activity of the target, including, by way of example only, to enhance the activity of the target, to inhibit the activity of the target, to limit the activity of the target, or to extend the activity of the target.

[00789] The term "modulator," as used herein, refers to a molecule that interacts with a target either directly or indirectly. The interactions include, but are not limited to, the interactions of an agonist and an antagonist.

[00790] The terms "neurogenerative disease" or "nervous system disorder," as used herein, refers to conditions that alter the structure or function of the brain, spinal cord or peripheral nervous system, including but not limited to Alzheimer's Disease, cerebral edema, cerebral ischemia, multiple sclerosis, neuropathies, Parkinson's Disease, those found after blunt or surgical trauma (including post-surgical cognitive dysfunction and spinal cord or brain stem injury), as well as the neurological aspects of disorders such as degenerative disk disease and sciatica. The acronym "CNS" refers to disorders of the central nervous system, i.e., brain and spinal cord [Sugaya K, *et al.*, "New anti-inflammatory treatment strategy in Alzheimer's disease", *Jpn J Pharmacol.* 2000 Feb;82(2):85-94; Yu GL, *et al.*, "Montelukast, a cysteinyl leukotriene receptor-1 antagonist, dose- and time-dependently protects against focal cerebral ischemia in mice", *Pharmacology.* 2005 Jan;73(1):31-40. Epub 2004 Sep 27; [Zhang WP, *et al.*, "Neuroprotective effect of ONO-1078, a leukotriene receptor antagonist, on focal cerebral ischemia in rats", *Acta Pharmacol Sin.* 2002 Oct;23(10):871-7].
30
35

[00791] The terms "ocular disease" or "ophthalmic disease," as used herein, refer to diseases which affect the eye or eyes and potentially the surrounding tissues as well. Ocular or ophthalmic diseases include, but are not limited to, conjunctivitis, retinitis, scleritis, uveitis, allergic conjunctivitis, vernal conjunctivitis, papillary conjunctivitis [Toriyama S., "Effects of leukotriene B4 receptor antagonist on experimental autoimmune

uveoretinitis in rats", *Nippon Ganka Gakkai Zasshi*. 2000 Jun;104(6):396-40; [Chen F, *et al.*, "Treatment of S antigen uveoretinitis with lipoxigenase and cyclo-oxygenase inhibitors", *Ophthalmic Res.* 1991;23(2):84-91].

[00792] The term "pharmaceutically acceptable excipient," as used herein, refers to a material, such as a carrier or diluent, which does not abrogate the desired biological activity or desired properties of the compound, and is relatively nontoxic, i.e., the material may be administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[00793] The term "pharmaceutically acceptable salt" refers to a formulation of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. Pharmaceutically acceptable salts may be obtained by reacting a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. Pharmaceutically acceptable salts may also be obtained by reacting a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a sodium or a potassium salt, an alkaline earth metal salt, such as a calcium or a magnesium salt, a salt of organic bases such as dicyclohexylamine, *N*-methyl-D-glucamine, tris(hydroxymethyl)methylamine, and salts with amino acids such as arginine, lysine, and the like, or by other methods known in the art

[00794] The term "pharmaceutical combination" as used herein, means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term "fixed combination" means that the active ingredients, e.g. a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term "non-fixed combination" means that the active ingredients, e.g. a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), and a co-agent, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[00795] The term "pharmaceutical composition" refers to a mixture of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. The pharmaceutical composition facilitates administration of the compound to an organism. Multiple techniques of administering a compound exist in the art including, but not limited to: intravenous, oral, aerosol, parenteral, ophthalmic, pulmonary and topical administration.

[00796] The term "respiratory disease," as used herein, refers to diseases affecting the organs that are involved in breathing, such as the nose, throat, larynx, trachea, bronchi, and lungs. Respiratory diseases include, but are not limited to, asthma, adult respiratory distress syndrome and allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, seasonal asthma, seasonal allergic rhinitis, perennial allergic rhinitis, chronic obstructive pulmonary disease, including chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation and cystic fibrosis, and hypoxia [Evans JF, "The Cysteinyl Leukotriene (CysLT) Pathway in Allergic Rhinitis", *Allergology International* 2005;54:187-90]; Kemp JP., "Leukotriene receptor antagonists for the treatment of asthma", *IDrugs*. 2000 Apr;3(4):430-41; Riccioni G, *et al.*, "Effect of the two different leukotriene receptor antagonists, montelukast and zafirlukast, on quality of life: a 12-week randomized study", *Allergy Asthma Proc*. 2004 Nov-Dec;25(6):445-8].

[00797] The term "subject" or "patient" encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish and the like. In one embodiment of the methods and compositions provided herein, the mammal is a human.

[00798] The terms "treat," "treating" or "treatment," as used herein, include alleviating, abating or ameliorating a disease or condition symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Pharmaceutical Composition/Formulation

[00799] Pharmaceutical compositions may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art. A summary of pharmaceutical compositions described herein may be found, for example, in *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference in their entirety.

[00800] Provided herein are pharmaceutical compositions comprising a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), and a pharmaceutically acceptable diluent(s), excipient(s), or carrier(s). In addition, the compounds described herein can be administered as pharmaceutical compositions in which compounds of any of Formula (A), Formula (B), Formula (C), Formula

(E), Formula (F), Formula (G), or Formula (H), are mixed with other active ingredients, as in combination therapy.

[00801] A pharmaceutical composition, as used herein, refers to a mixture of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. The pharmaceutical composition facilitates administration of the compound to an organism. In practicing the methods of treatment or use provided herein, therapeutically effective amounts of compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), provided herein are administered in a pharmaceutical composition to a mammal having a disease or condition to be treated. Preferably, the mammal is a human. A therapeutically effective amount can vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. The compounds can be used singly or in combination with one or more therapeutic agents as components of mixtures.

[00802] For intravenous injections, compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. For other parenteral injections, appropriate formulations may include aqueous or nonaqueous solutions, preferably with physiologically compatible buffers or excipients. Such excipients are generally known in the art.

[00803] For oral administration, compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers or excipients well known in the art. Such carriers enable the compounds described herein to be formulated as tablets, powders, pills, dragees, capsules, liquids, gels, syrups, elixirs, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

[00804] Pharmaceutical preparations for oral use can be obtained by mixing one or more solid excipient with one or more of the compounds described herein, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as: for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others such as: polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. If desired, disintegrating agents may be added, such as the cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[00805] Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

[00806] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

[00807] For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, or gels formulated in a conventional manner. Parental injections may involve bolus injection or continuous infusion.

Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The pharmaceutical composition of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be in a form suitable for parenteral injection as a sterile suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

[00808] The compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be administered topically and can be formulated into a variety of topically administrable compositions, such as solutions, suspensions, lotions, gels, pastes, medicated sticks, balms, creams or ointments. Such pharmaceutical compounds can contain solubilizers, stabilizers, tonicity enhancing agents, buffers and preservatives.

[00809] Formulations suitable for transdermal administration of compounds having the structure of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may employ transdermal delivery devices and transdermal delivery patches and can be lipophilic emulsions or buffered, aqueous solutions, dissolved and/or dispersed in a polymer or an adhesive. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents. Still further, transdermal delivery of the compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can be accomplished by means of iontophoretic patches and the like. Additionally, transdermal patches can provide controlled delivery of the compounds any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). The rate of absorption can be slowed by using rate-controlling membranes or by trapping the compound within a polymer matrix or gel. Conversely, absorption enhancers can be used to increase absorption. An absorption enhancer or carrier can include absorbable pharmaceutically acceptable solvents to assist passage through the skin. For example, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with

carriers, optionally a rate controlling barrier to deliver the compound to the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

[00810] For administration by inhalation, the compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), maybe in a form as an aerosol, a mist or a powder.

- 5 Pharmaceutical compositions of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount.

- 10 Capsules and cartridges of, such as, by way of example only, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[00811] The compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may also be formulated in rectal compositions such as enemas, rectal gels, rectal foams, rectal aerosols, suppositories, jelly suppositories, or retention enemas, containing conventional suppository bases such as cocoa butter or other glycerides, as well as synthetic polymers such as polyvinylpyrrolidone, PEG, and the like. In suppository forms of the compositions, a low-melting wax such as, but not limited to, a mixture of fatty acid glycerides, optionally in combination with cocoa butter is first melted.

- 15 [00812] Pharmaceutical compositions may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art. Pharmaceutical compositions comprising a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

25 [00813] The pharmaceutical compositions will include at least one pharmaceutically acceptable carrier, diluent or excipient and a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), described herein as an active ingredient in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In addition, the methods and pharmaceutical compositions described herein include the use of *N*-oxides, crystalline forms (also known as polymorphs), as well as active metabolites of these compounds having the same type of activity. In some situations, compounds may exist as tautomers. All tautomers are included within the scope of the compounds presented herein. Additionally, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein. In addition, the pharmaceutical compositions may include other medicinal or pharmaceutical agents, carriers, adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, and/or buffers. In addition, the pharmaceutical compositions can also contain other therapeutically valuable substances.

35

[00814] Methods for the preparation of compositions comprising the compounds described herein include formulating the compounds with one or more inert, pharmaceutically acceptable excipients or carriers to form a solid, semi-solid or liquid. Solid compositions include, but are not limited to, powders, tablets, dispersible granules, capsules, cachets, and suppositories. Liquid compositions include solutions in which a compound is dissolved, emulsions comprising a compound, or a solution containing liposomes, micelles, or nanoparticles comprising a compound as disclosed herein. Semi-solid compositions include, but are not limited to, gels, suspensions and creams. The compositions may be in liquid solutions or suspensions, solid forms suitable for solution or suspension in a liquid prior to use, or as emulsions. These compositions may also contain minor amounts of nontoxic, auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, and so forth.

[00815] A composition comprising a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), can illustratively take the form of a liquid where the agents are present in solution, in suspension or both. Typically when the composition is administered as a solution or suspension a first portion of the agent is present in solution and a second portion of the agent is present in particulate form, in suspension in a liquid matrix. In some embodiments, a liquid composition may include a gel formulation. In other embodiments, the liquid composition is aqueous.

[00816] Useful aqueous suspension can also contain one or more polymers as suspending agents. Useful polymers include water-soluble polymers such as cellulosic polymers, *e.g.*, hydroxypropyl methylcellulose, and water-insoluble polymers such as cross-linked carboxyl-containing polymers. Useful compositions can also comprise an mucoadhesive polymer, selected for example from carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate copolymer, sodium alginate and dextran.

[00817] Useful compositions may also include solubilizing agents to aid in the solubility of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). The term "solubilizing agent" generally includes agents that result in formation of a micellar solution or a true solution of the agent. Certain acceptable nonionic surfactants, for example polysorbate 80, can be useful as solubilizing agents, as can ophthalmically acceptable glycols, polyglycols, *e.g.*, polyethylene glycol 400, and glycol ethers.

[00818] Useful compositions may also include one or more pH adjusting agents or buffering agents, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

[00819] Useful compositions may also include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

[00820] Other useful compositions may also include one or more preservatives to inhibit microbial activity.

Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

- 5 [00821] Still other useful compositions may include one or more surfactants to enhance physical stability or for other purposes. Suitable nonionic surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, *e.g.*, polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, *e.g.*, octoxynol 10, octoxynol 40.

- 10 [00822] Still other useful compositions may include one or more antioxidants to enhance chemical stability where required. Suitable antioxidants include, by way of example only, ascorbic acid and sodium metabisulfite.

[00823] Aqueous suspension compositions can be packaged in single-dose non-reclosable containers.

Alternatively, multiple-dose reclosable containers can be used, in which case it is typical to include a preservative in the composition.

[00824] Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed.

- 15 Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as *N*-methylpyrrolidone also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art.
- 20 Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

- [00825] All of the formulations described herein may benefit from antioxidants, metal chelating agents, thiol containing compounds and other general stabilizing agents. Examples of such stabilizing agents, include, but are not limited to: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan polysulfate and other heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

30 **Routes of Administration**

[00826] Suitable routes of administration include, but are not limited to, intravenous, oral, rectal, aerosol, parenteral, ophthalmic, pulmonary, transmucosal, transdermal, vaginal, otic, nasal, and topical administration. In addition, by way of example only, parenteral delivery includes intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intralymphatic, and intranasal injections.

- 35 [00827] Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into an organ, often in a depot preparation or sustained release formulation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with organ-specific antibody. The liposomes will be targeted
- 40

to and taken up selectively by the organ. In addition, the drug may be provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation.

Methods of Dosing and Treatment Regimens

[00828] The compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), can be used in the preparation of medicaments for the treatment of leukotriene-dependent or leukotriene mediated diseases or conditions. In addition, a method for treating any of the diseases or conditions described herein in a subject in need of such treatment, involves administration of pharmaceutical compositions containing at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or a pharmaceutically acceptable salt, pharmaceutically acceptable N-oxide, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said subject

[00829] The compositions containing the compound(s) described herein can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest the symptoms of the disease or condition. Amounts effective for this use will depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician. It is considered well within the skill of the art for one to determine such therapeutically effective amounts by routine experimentation (including, but not limited to, a dose escalation clinical trial).

[00830] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. It is considered well within the skill of the art for one to determine such prophylactically effective amounts by routine experimentation (*e.g.*, a dose escalation clinical trial). When used in a patient, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[00831] In the case wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds may be administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00832] In the case wherein the patient's status does improve, upon the doctor's discretion the administration of the compounds may be given continuously; alternatively, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time (*i.e.*, a "drug holiday"). The length of the drug holiday can vary between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, and 365 days. The dose reduction during a drug holiday may be from 10%-100%, including by way of example only 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, and 100%.

[00833] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, can be reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. Patients can, however, require intermittent treatment on a long-term basis upon any recurrence of symptoms.

5 [00834] The amount of a given agent that will correspond to such an amount will vary depending upon factors such as the particular compound, disease condition and its severity, the identity (*e.g.*, weight) of the subject or host in need of treatment, but can nevertheless be routinely determined in a manner known in the art according to the particular circumstances surrounding the case, including, *e.g.*, the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated. In general, however, doses
10 employed for adult human treatment will typically be in the range of 0.02-5000 mg per day, preferably 1-1500 mg per day. The desired dose may conveniently be presented in a single dose or as divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00835] The pharmaceutical composition described herein may be in unit dosage forms suitable for single
15 administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of one or more compound. The unit dosage may be in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. Aqueous suspension compositions can be packaged in single-dose non-reclosable containers. Alternatively, multiple-dose reclosable containers can be used, in which case it is typical to include a preservative
20 in the composition. By way of example only, formulations for parenteral injection may be presented in unit dosage form, which include, but are not limited to ampoules, or in multi-dose containers, with an added preservative.

[00836] The daily dosages appropriate for the compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), and Formula (H), described herein are from about 0.01 to 2.5 mg/kg per body
25 weight. An indicated daily dosage in the larger mammal, including, but not limited to, humans, is in the range from about 0.5 mg to about 100 mg, conveniently administered in divided doses, including, but not limited to, up to four times a day or in extended release form. Suitable unit dosage forms for oral administration comprise from about 1 to 50 mg active ingredient. The foregoing ranges are merely suggestive, as the number of variables in regard to an individual treatment regime is large, and considerable excursions from these recommended values
30 are not uncommon. Such dosages may be altered depending on a number of variables, not limited to the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[00837] Toxicity and therapeutic efficacy of such therapeutic regimens can be determined by standard
pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the
35 determination of the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds exhibiting high therapeutic indices are preferred. The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating

concentrations that include the ED₅₀ with minimal toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

Use of FLAP Modulators to Prevent and/or Treat Leukotriene-Dependent or Leukotriene Mediated Diseases or Conditions

5 [00838] The therapy of leukotriene-dependent or leukotriene mediated diseases or conditions is designed to modulate the activity of FLAP. Such modulation may include, by way of example only, inhibiting or antagonizing FLAP activity. For example, a FLAP inhibitor can be administered in order to decrease synthesis of leukotrienes within the individual, or possibly to downregulate or decrease the expression or availability of the FLAP mRNA or specific splicing variants of the FLAP mRNA. Downregulation or decreasing expression or
10 availability of a native FLAP mRNA or of a particular splicing variant could minimize the expression or activity of a defective nucleic acid or the particular splicing variant and thereby minimize the impact of the defective nucleic acid or the particular splicing variant.

[00839] In accordance with one aspect, compositions and methods described herein include compositions and methods for treating, preventing, reversing, halting or slowing the progression of leukotriene-dependent or
15 leukotriene mediated diseases or conditions once it becomes clinically evident, or treating the symptoms associated with or related to leukotriene-dependent or leukotriene mediated diseases or conditions, by administering to the subject a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). The
20 subject may already have a leukotriene-dependent or leukotriene mediated disease or condition at the time of administration, or be at risk of developing a leukotriene-dependent or leukotriene mediated disease or condition. The symptoms of leukotriene-dependent or leukotriene mediated diseases or conditions in a subject can be determined by one skilled in the art and are described in standard textbooks.

[00840] The activity of 5-lipoxygenase activating protein in a mammal may be directly or indirectly modulated
25 by the administration of (at least once) an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), to a mammal. Such modulation includes, but is not limited to, reducing and/or inhibiting the activity of 5-lipoxygenase activating protein. In addition, the activity of
30 leukotrienes in a mammal may be directly or indirectly modulated, including reducing and/or inhibiting, by the administration of (at least once) an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), to a mammal. Such modulation includes, but is not limited to,
35 reducing and/or inhibiting the activity of 5-lipoxygenase activating protein.

[00841] Prevention and/or treatment leukotriene-dependent or leukotriene mediated diseases or conditions may comprise administering to a mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or
40 pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). By way of example, the prevention and/or

treatment of inflammation diseases or conditions may comprise administering to a mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Leukotriene-dependent or leukotriene mediated diseases or conditions that may be treated by a method comprising administering to a mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), include, but are not limited to, bone diseases and disorder, cardiovascular diseases and disorders, inflammatory diseases and disorders, dermatological diseases and disorders, ocular diseases and disorders, cancer and other proliferative diseases and disorders, respiratory diseases and disorder, and non-cancerous disorders.

[00842] By way of example only, included in the prevention/treatment methods described herein are methods for treating respiratory diseases comprising administering to the mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). By way of example the respiratory disease may be asthma; see Riccioni *et al*, *Ann. Clin. Lab. Sci.*, v34, 379-387 (2004). In addition, the respiratory disease may include, but is not limited to, adult respiratory distress syndrome and allergic (extrinsic) asthma, non-allergic (intrinsic) asthma, acute severe asthma, chronic asthma, clinical asthma, nocturnal asthma, allergen-induced asthma, aspirin-sensitive asthma, exercise-induced asthma, isocapnic hyperventilation, child-onset asthma, adult-onset asthma, cough-variant asthma, occupational asthma, steroid-resistant asthma, seasonal asthma, allergic rhinitis, vascular responses, endotoxin shock, fibrogenesis, pulmonary fibrosis, allergic diseases, chronic inflammation, and adult respiratory distress syndrome.

[00843] By way of example only, included in such treatment methods are methods for preventing chronic obstructive pulmonary disease comprising administering to the mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). In addition, chronic obstructive pulmonary disease includes, but is not limited to, chronic bronchitis or emphysema, pulmonary hypertension, interstitial lung fibrosis and/or airway inflammation and cystic fibrosis.

[00844] By way of example only, included in such treatment methods are methods for preventing increased mucosal secretion and/or edema in a disease or condition comprising administering to the mammal at least once an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00845] By way of example only, included in the prevention/treatment methods described herein are methods for preventing or treating vasoconstriction, atherosclerosis and its sequelae myocardial ischemia, myocardial infarction, aortic aneurysm, vasculitis and stroke comprising administering at least once to the mammal an

effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H); see Jala *et al*, *Trends in Immunol.*, v25, 315-322 (2004) and Mehrabian *et al*, *Curr. Opin. Lipidol.*, v14, 447-457 (2003).

[00846] By way of example only, included in the prevention/treatment methods described herein are methods for reducing cardiac reperfusion injury following myocardial ischemia and/or endotoxic shock comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00847] By way of example only, included in the prevention/treatment methods described herein are methods for reducing the constriction of blood vessels in a mammal comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00848] By way of example only, included in the prevention/treatment methods described herein are methods for lowering or preventing an increase in blood pressure of a mammal comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00849] By way of example only, included in the prevention/treatment methods described herein are methods for preventing eosinophil and/or basophil and/or dendritic cell and/or neutrophil and/or monocyte recruitment comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00850] By way of example only, included in the prevention/treatment methods described herein are methods for the prevention or treatment of abnormal bone remodeling, loss or gain, including diseases or conditions as, by way of example, osteopenia, osteoporosis, Paget's disease, cancer and other diseases comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00851] By way of example only, included in the prevention/treatment methods described herein are methods for preventing ocular inflammation and allergic conjunctivitis, vernal keratoconjunctivitis, and papillary conjunctivitis comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula

(H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H); see Lambiase *et al*, *Arch. Ophthalmol.*, v121, 615-620 (2003).

5 [00852] By way of example only, included in the prevention/treatment methods described herein are methods for preventing CNS disorders comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). CNS disorders include, but are not limited to, multiple sclerosis, Parkinson's disease, Alzheimer's disease, stroke, cerebral ischemia, retinal
10 ischemia, post-surgical cognitive dysfunction, migraine, peripheral neuropathy/neuropathic pain, spinal cord injury, cerebral edema and head injury.

[00853] By way of example only, included in the prevention/treatment methods described herein are methods for the treatment of cancer comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula
15 (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). The type of cancer may include, but is not limited to, pancreatic cancer and other solid or hematological tumors, see Poff and Balazy, *Curr. Drug Targets Inflamm. Allergy*, v3, 19-33 (2004) and Steele *et al*, *Cancer Epidemiology & Prevention*, v8, 467-483 (1999).

20 [00854] By way of example only, included in the prevention/treatment methods described herein are methods for preventing endotoxic shock and septic shock comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

25 [00855] By way of example only, included in the prevention/treatment methods described herein methods for preventing rheumatoid arthritis and osteoarthritis comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula
30 (H).

[00856] By way of example only, included in the prevention/treatment methods described herein are methods for preventing increased GI diseases comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula
35 (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such GI diseases include, by way of example only, inflammatory bowel disease (IBD), colitis and Crohn's disease.

[00857] By way of example only, included in the prevention/treatment methods described herein are methods for the reduction of inflammation while also preventing transplant rejection or preventing or treating tumors or accelerating the healing of wounds comprising administering at least once to the mammal an effective amount of
40 at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G),

or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00858] By way of example only, included in the prevention/treatment methods described herein are methods for the prevention or treatment of rejection or dysfunction in a transplanted organ or tissue comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00859] By way of example only, included in the prevention/treatment methods described herein are methods for treating type II diabetes comprising administering to at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00860] By way of example only, included in the prevention/treatment methods described herein are methods for treating inflammatory responses of the skin comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such inflammatory responses of the skin include, by way of example, psoriasis, dermatitis, contact dermatitis, eczema, urticaria, rosacea, wound healing and scarring. In another aspect are methods for reducing psoriatic lesions in the skin, joints, or other tissues or organs, comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00861] By way of example only, included in the prevention/treatment methods described herein are methods for the treatment of cystitis, including, by way of example only, interstitial cystitis, comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00862] By way of example only, included in the prevention/treatment methods described herein are methods for the treatment of metabolic syndromes such as Familial Mediterranean Fever comprising administering at least once to the mammal an effective amount of at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

Combination Treatments

[00863] In certain instances, it may be appropriate to administer at least one compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one

of the compounds herein is inflammation, then it may be appropriate to administer an anti-inflammatory agent in combination with the initial therapeutic agent. Or, by way of example only, the therapeutic effectiveness of one of the compounds described herein may be enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant may have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, by way of example only, the benefit of experienced by a patient may be increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. By way of example only, in a treatment for asthma involving administration of one of the compounds described herein, increased therapeutic benefit may result by also providing the patient with other therapeutic agents or therapies for asthma. In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

[00864] It is known to those of skill in the art that therapeutically-effective dosages can vary when the drugs are used in treatment combinations. Methods for experimentally determining therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens are described in the literature. For example, the use of metronomic dosing, *i.e.*, providing more frequent, lower doses in order to minimize toxic side effects, has been described extensively in the literature. A combination treatment regimen may encompass treatment regimens in which administration of a FLAP or 5-LO inhibitor described herein is initiated prior to, during, or after treatment with a second agent described above, and continues until any time during treatment with the second agent or after termination of treatment with the second agent. It also includes treatments in which a FLAP or 5-LO inhibitor described herein and the second agent being used in combination are administered simultaneously or at different times and/or at decreasing or increasing intervals during the treatment period. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient. For example, a FLAP or 5-LO inhibitor described herein in the combination treatment can be administered weekly at the onset of treatment, decreasing to biweekly, and decreasing further as appropriate.

[00865] Compositions and methods for combination therapy are provided herein. In accordance with one aspect, the pharmaceutical compositions disclosed herein are used to treat leukotriene-dependent or leukotriene mediated conditions. In accordance with another aspect, the pharmaceutical compositions disclosed herein are used to treat respiratory diseases, where treatment with a FLAP inhibitor is indicated, in particular asthma, and to induce bronchodilation in a subject. In one embodiment, pharmaceutical compositions disclosed herein are used to treat a subject suffering from a vascular inflammation-driven disorder. In one embodiment, the pharmaceutical compositions disclosed herein are used to treat a subject susceptible to myocardial infarction (MI).

[00866] Combination therapies described herein can be used as part of a specific treatment regimen intended to provide a beneficial effect from the co-action of a FLAP inhibitors described herein and a concurrent treatment. It is understood that the dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, can be modified in accordance with a variety of factors. These factors include the type of respiratory disorder and the type of bronchodilation from which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, the dosage regimen actually employed can vary widely and therefore can deviate from the dosage regimens set forth herein.

[00867] For combination therapies described herein, dosages of the co-administered compounds will of course vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In addition, when co-administered with one or more biologically active agents, the compound provided herein may be administered either simultaneously with the biologically active agent(s), or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein in combination with the biologically active agent(s).

[00868] In any case, the multiple therapeutic agents (one of which is one of the compounds described herein) may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may vary from more than zero weeks to less than four weeks. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations are also envisioned.

[00869] In addition, the compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may also be used in combination with procedures that may provide additional or synergistic benefit to the patient. By way of example only, patients are expected to find therapeutic and/or prophylactic benefit in the methods described herein, wherein pharmaceutical composition of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), and /or combinations with other therapeutics are combined with genetic testing to determine whether that individual is a carrier of a mutant gene that is known to be correlated with certain diseases or conditions.

[00870] The compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), and combination therapies can be administered before, during or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound can vary. Thus, for example, the compounds can be used as a prophylactic and can be administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. The compounds and compositions can be administered to a subject during or as soon as possible after the onset of the symptoms. The administration of the compounds can be initiated within the first 48 hours of the onset of the symptoms, preferably within the first 48 hours of the onset of the symptoms, more preferably within the first 6 hours of the onset of the symptoms, and most preferably within 3 hours of the onset of the symptoms. The initial administration can be via any route practical, such as, for example, an intravenous injection, a bolus injection, infusion over 5 minutes to about 5 hours, a pill, a capsule, transdermal patch, buccal delivery, and the like, or combination thereof. A compound is preferably administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease, such as, for example, from about 1 month to about 3 months. The length of treatment can vary for each subject, and the length can be determined using the known criteria. For example, the compound or a formulation containing the compound can be administered for at least 2 weeks, preferably about 1 month to about 5 years, and more preferably from about 1 month to about 3 years.

[00871] By way of example, therapies which combine compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with inhibitors of leukotriene synthesis or leukotriene receptor antagonists, either acting at the same or other points in the leukotriene synthesis pathway,

could prove to be particularly useful for treating leukotriene-dependent or leukotriene mediated diseases or conditions. In addition, by way of example, therapies which combine compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with inhibitors of inflammation could prove to be particularly useful for treating leukotriene-dependent or leukotriene mediated diseases or conditions.

5 Agents to Treat Respiratory Diseases or Conditions

[00872] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases include administering to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with other therapeutic agents that are used in the treatment of respiratory conditions or disorders, such as, but not limited to asthma. Therapeutic agents used in the treatment of respiratory conditions and disorders, such as, but not limited to asthma, include: glucocorticoids, such as, ciclesonide, beclomethasone, budesonide, flunisolide, fluticasone, mometasone, and triamcinolone; leukotriene modifiers, such as, montelukast, zafirlukast, pranlukast, and zileuton; mast cell stabilizers, such as, cromoglicate (cromolyn), and nedocromil; antimuscarinics/anticholinergics, such as, ipratropium, oxitropium, and tiotropium; methylxanthines, such as, theophylline and aminophylline; antihistamine, such as, mepyramine (pyrilamine), antazoline, diphenhydramine, carbinoxamine, doxylamine, clemastine, dimenhydrinate, pheniramine, chlorphenamine (chlorpheniramine), dexchlorphenamine, brompheniramine, triprolidine, cyclizine, chlorcyclizine, hydroxyzine, meclizine, promethazine, alimemazine (trimeprazine), cyproheptadine, azatadine, ketotifen, acrivastine, astemizole, cetirizine, loratadine, mizolastine, terfenadine, fexofenadine, levocetirizine, desloratadine, fexofenadine; omalizumab, an IgE blocker; beta2-adrenergic receptor agonists, such as: short acting beta2-adrenergic receptor agonists, such as, salbutamol (albuterol), levalbuterol, terbutaline, pirbuterol, procaterol, metaproterenol, fenoterol, bitolterol mesylate; and long-acting beta2-adrenergic receptor agonists, such as, salmeterol, formoterol, bambuterol.

Anti-Inflammatory Agents

[00873] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases include administering to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with an anti-inflammatory agent including, but not limited to, non-steroidal anti-inflammatory drugs (NSAIDs) and corticosteroids (glucocorticoids).

[00874] NSAIDs include, but are not limited to: aspirin, salicylic acid, gentisic acid, choline magnesium salicylate, choline salicylate, choline magnesium salicylate, choline salicylate, magnesium salicylate, sodium salicylate, diflunisal, carprofen, fenoprofen, fenoprofen calcium, flurobiprofen, ibuprofen, ketoprofen, nabutone, ketolorac, ketorolac tromethamine, naproxen, oxaprozin, diclofenac, etodolac, indomethacin, sulindac, tolmetin, meclofenamate, meclofenamate sodium, mefenamic acid, piroxicam, meloxicam, COX-2 specific inhibitors (such as, but not limited to, celecoxib, rofecoxib, valdecoxib, parecoxib, etoricoxib, lumiracoxib, CS-502, JTE-522, L-745,337 and NS398).

[00875] Corticosteroids, include, but are not limited to: betamethasone (Celestone), prednisone (Deltasone), alclometasone, aldosterone, amcinonide, beclomethasone, betamethasone, budesonide, ciclesonide, clobetasol, clobetasone, clocortolone, cloprednol, cortisone, cortivazol, deflazacort, deoxycorticosterone, desonide, desoximetasone, desoxycortone, dexamethasone, diflorasone, diflucortolone, difluprednate, fluclorolone, fludrocortisone, fludroxycortide, flumetasone, flunisolide, fluocinolone acetonide, fluocinonide, fluocortin, fluocortolone, fluorometholone, fluperolone, fluprednidene, fluticasone, formocortol, halcinonide, halometasone,

hydrocortisone/cortisol, hydrocortisone aceponate, hydrocortisone bupropate, hydrocortisone butyrate, loteprednol, medrysone, meprednisone, methylprednisolone, methylprednisolone aceponate, mometasone furoate, paramethasone, prednicarbate, prednisone/prednisolone, rimexolone, tixocortol, triamcinolone, and ulobetasol.

[00876] Corticosteroids do not directly inhibit leukotriene production, therefore co-dosing with steroids could provide additional anti-inflammatory benefit.

[00877] Some commercially available anti-inflammatories include, but are not limited to: Arthrotec[®] (diclofenac and misoprostol), Asacol[®], Salofalk[®] (5-aminosalicylic acid), Auralgan[®] (antipyrine and benzocaine), Azulfidine[®] (sulfasalazine), Daypro[®] (oxaprozin), Lodine[®] (etodolac), Ponstan[®] (mefenamic acid), Solumedrol[®] (methylprednisolone), Bayer[®], Bufferin[®] (aspirin), Indocin[®] (indomethacin), Vioxx[®] (rofecoxib), Celebrex[®] (celecoxib), Bextra[®] (valdecoxib), Arcoxia[®] (etoricoxib), Prexige[®] (lumiracoxib), Advil[®], Motrin[®] (ibuprofen), Voltaren[®] (diclofenac), Orudis[®] (ketoprofen), Mobic[®] (meloxicam), Relafen[®] (nabumetone), Aleve[®], Naprosyn[®] (naproxen), Feldene[®] (piroxicam).

[00878] By way of example, asthma is a chronic inflammatory disease characterized by pulmonary eosinophilia and airway hyperresponsiveness. Zhao *et al.*, *Proteomics*, July 4, 2005. In patients with asthma, leukotrienes may be released from mast cells, eosinophils, and basophils. The leukotrienes are involved in contraction of airway smooth muscle, an increase in vascular permeability and mucus secretions, and have been reported to attract and activate inflammatory cells in the airways of asthmatics (Siegel *et al.*, ed., *Basic Neurochemistry, Molecular, Cellular and Medical Aspects*, Sixth Ed., Lippincott Williams & Wilkins, 1999). Thus, in another embodiment described herein, the methods for treatment of respiratory diseases include administering to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with an anti-inflammatory agent.

Leukotriene Receptor Antagonists

[00879] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases includes administered to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with leukotriene receptor antagonists including, but are not limited to, CysLT₁/CysLT₂ dual receptor antagonists and CysLT₁ receptor antagonists. In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases includes administered to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with a CysLT₁/CysLT₂ dual receptor antagonist. CysLT₁/CysLT₂ dual receptor antagonists include, but are not limited to, BAY u9773, Cuthbert *et al* EP 00791576 (published 27 Aug 1997), DUO-LT (Galczenski *et al*, D38, Poster F4 presented at American Thoracic Society, May 2002) and Tsuji *et al*, *Org. Biomol. Chem.*, 1, 3139-3141, 2003. For a particular patient, the most appropriate formulation or method of use of such combination treatments may depend on the type of leukotriene-dependent or leukotriene mediated disorder, the time period in which the FLAP inhibitor acts to treat the disorder and the time period in which the CysLT₁/CysLT₂ dual receptor antagonist acts to inhibit CysLT receptor activity. By way of example only, such combination treatments may be used for treating a patient suffering from a respiratory disorders.

[00880] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases includes administered to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with a CysLT₁ receptor antagonist. CysLT₁ receptor antagonists include, but are not limited to, Zafirlukast ("Accolate[™]"), Montelukast ("Singulair[™]"), Pranlukast ("Onon[™]"),

and derivatives or analogs thereof. Such combinations may be used to treat leukotriene-dependent or leukotriene mediated disorder, including respiratory disorders.

[00881] The co-administration of a FLAP or 5-LO inhibitor described herein with a CysLT₁ receptor antagonist or a dual CysLT₁/CysLT₂ receptor antagonist may have therapeutic benefit over and above the benefit derived from the administration of either a FLAP or 5-LO inhibitor or a CysLT₁R antagonist alone. In the case that substantial inhibition of leukotriene production has undesired effects, partial inhibition of this pathway through the amelioration of the effects of the proinflammatory LTB₄ and cysteinyl leukotrienes combined with the block of the CysLT₁ receptor and/or dual CysLT₁/CysLT₂ receptor block may afford substantial therapeutic benefits, particularly for respiratory diseases.

Other Combination Therapies

[00882] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as proliferative disorders, including cancer, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from the group consisting of alemtuzumab, arsenic trioxide, asparaginase (pegylated or non-), bevacizumab, cetuximab, platinum-based compounds such as cisplatin, cladribine, daunorubicin/doxorubicin/idarubicin, irinotecan, fludarabine, 5-fluorouracil, gemtuzumab, methotrexate, PaclitaxelTM, taxol, temozolomide, thioguanine, or classes of drugs including hormones (an antiestrogen, an antiandrogen, or gonadotropin releasing hormone analogues, interferons such as alpha interferon, nitrogen mustards such as busulfan or melphalan or mechlorethamine, retinoids such as tretinoin, topoisomerase inhibitors such as irinotecan or topotecan, tyrosine kinase inhibitors such as gefinitinib or imatinib, or agents to treat signs or symptoms induced by such therapy including allopurinol, filgrastim, granisetron/ondansetron/palonosetron, dronabinol.

[00883] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as the therapy of transplanted organs or tissues or cells, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from the group consisting of azathioprine, a corticosteroid, cyclophosphamide, cyclosporin, dacluzimab, mycophenolate mofetil, OKT3, rapamycin, tacrolimus, thymoglobulin.

[00884] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as atherosclerosis, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from the group consisting of HMG-CoA reductase inhibitors (e.g., statins in their lactonized or dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof, including but not limited to lovastatin; simvastatin; dihydroxy open-acid simvastatin, particularly the ammonium or calcium salts thereof; pravastatin, particularly the sodium salt thereof; fluvastatin, particularly the sodium salt thereof; atorvastatin, particularly the calcium salt thereof; nisvastatin, also referred to as NK-104; rosuvastatin); agents that have both lipid-altering effects and other pharmaceutical activities; HMG-CoA synthase inhibitors; cholesterol absorption inhibitors such as ezetimibe; cholesterol ester transfer protein (CETP) inhibitors, for example JTT-705 and CP529, 414; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors); acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of

ACAT-1 or ACAT-2 as well as dual inhibitors of ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; platelet aggregation inhibitors, for example glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR γ) agonists, including the compounds commonly referred to as glitazones, for example troglitazone, pioglitazone and rosiglitazone and including those compounds included within the structural class known as thiazolidinediones as well as those PPAR γ agonists outside the thiazolidinedione structural class; PPAR α agonists such as clofibrate, fenofibrate including micronized fenofibrate, and gemfibrozil; PPAR dual α/γ agonists such as 5-[(2, 4-dioxo-5-thiazolidinyl)methyl]-2-methoxy-N-[[4-(trifluoromethyl)phenyl]methyl]-benzamide, known as KRP-297; vitamin B6 (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B12 (also known as cyanocobalamin); folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the methylglucamine salt; anti-oxidant vitamins such as vitamin C and E and beta carotene; beta-blockers; angiotensin II antagonists such as losartan; angiotensin converting enzyme inhibitors such as enalapril and captopril; calcium channel blockers such as nifedipine and diltiazem; endothelial antagonists; agents that enhance ABC1 gene expression; FXR and LXR ligands including both inhibitors and agonists; bisphosphonate compounds such as alendronate sodium; fish oils or omega-3 fatty acids (eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)) or alpha-linolenic acid (LNA) or omega-3 fatty acid esters such as OmacorTM; and cyclooxygenase-2 inhibitors such as rofecoxib and celecoxib.

[00885] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as the therapy of stroke, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from COX-2 inhibitors; nitric oxide synthase inhibitors, such as N-(3-(aminomethyl)benzyl)acetamidine; Rho kinase inhibitors, such as fasudil; angiotensin II type-1 receptor antagonists, including candesartan, losartan, irbesartan, eprosartan, telmisartan and valsartan; glycogen synthase kinase 3 inhibitors; sodium or calcium channel blockers, including clobenafite; p38 MAP kinase inhibitors, including SKB 239063; thromboxane synthetase inhibitors, including isbogrel, ozagrel, ridogrel and dazoxiben; statins (HMG CoA reductase inhibitors), including lovastatin, simvastatin, dihydroxy open-acid simvastatin, pravastatin, fluvastatin, atorvastatin, nisvastatin, and rosuvastatin; neuroprotectants, including free radical scavengers, calcium channel blockers, excitatory amino acid antagonists, growth factors, antioxidants, such as edaravone, vitamin C, TROLOXTM, citicoline and minicycline, and reactive astrocyte inhibitors, such as (2R)-2-propyloctanoic acid; beta adrenergic blockers, such as propranolol, nadolol, timolol, pindolol, labetalol, metoprolol, atenolol, esmolol and acebutolol; NMDA receptor antagonists, including memantine; NR2B antagonists, such as traxoprodil; 5-HT1A agonists; receptor platelet fibrinogen receptor antagonists, including tirofiban and lamifiban; thrombin inhibitors; antithrombotics, such as argatroban; antihypertensive agents, such as enalapril; vasodilators, such as cyclandelate; nociceptin antagonists; DP1V antagonists; GABA 5 inverse agonists; and selective androgen receptor modulators.

[00886] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as the therapy of pulmonary fibrosis, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one

additional agent selected from anti-inflammatory agents, such as corticosteroids, azathioprine or cyclophosphamide.

[00887] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as the therapy of interstitial cystitis, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from dimethylsulfoxide, omalizumab, and pentosan polysulfate.

[00888] In another embodiment described herein, methods for treatment of leukotriene-dependent or leukotriene mediated conditions or diseases, such as the therapy of disorders of bone, comprises administration to a patient compounds, pharmaceutical compositions, or medicaments described herein in combination with at least one additional agent selected from the group consisting of minerals, vitamins, bisphosphonates, anabolic steroids, parathyroid hormone or analogs, and cathepsin K inhibitors.

Treatment of Leukotriene based conditions or diseases using CysLT₁/CysLT₂ Receptor Antagonists

[00889] In accordance with another aspect, the compositions and methods described herein are designed to deliver a CysLT₁/CysLT₂ dual receptor antagonist to block the CysLT receptor activity. The term "CysLT antagonist" or "CysLT receptor antagonist" or "leukotriene receptor antagonist" refers to a therapy that decreases the signaling of CysLTs through CysLT receptors. CysLT typically refers to either LTC₄, LTD₄ or LTE₄. Cysteinyl leukotrienes are potent smooth muscle constricting agents, particularly in respiratory and circulatory systems. These are mediated via at least two cell receptors, CysLT₁ and CysLT₂. The CysLT₁ receptor and CysLT₂ receptors are G-protein-coupled receptors with seven putative transmembrane regions and an intracellular domain that interacts with G-proteins, Evans *et al*, *Prostaglandins and Other Lipid Mediators*, 68-69, p587-597, (2002). Examples of CysLT₁/CysLT₂ dual receptor antagonists are BAY u9773, Cuthbert *et al* EP 00791576 (published 27 Aug 1997), DUO-LT (Galczenski *et al*, D38, Poster F4 presented at American Thoracic Society, May 2002) and Tsuji *et al*, *Org. Biomol. Chem.*, 1, 3139-3141, 2003.

[00890] In certain embodiments, methods for treatment of leukotriene-dependent or leukotriene mediated diseases or conditions includes administering to patients compounds, pharmaceutical compositions, or medicaments comprising a CysLT₁/CysLT₂ receptor antagonist. By way of example, such compounds, pharmaceutical compositions, or medicaments may be used as treatment and/or prevention for respiratory diseases including, but not limited to, chronic stable asthma.

Diagnostic Methods for Patient Identification

[00891] The screening of "leukotriene-responsive patients" which may be selected for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical compositions or medicaments described herein which include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or other FLAP modulators, may be accomplished using techniques and methods described herein. Such techniques and methods include, by way of example, evaluation of gene haplotypes (genotype analysis), monitoring/measurement of biomarkers (phenotype analysis), monitoring/measurement of functional markers (phenotype analysis), which indicate patient response to known modulators of the leukotriene pathway, or any combination thereof.

Genotype Analysis: FLAP Polymorphisms

[00892] Human FLAP has been purified and cloned and is an 18 kilodalton membrane-bound protein which is most highly expressed in human neutrophils. The FLAP gene is located at 13q12 and the gene has been linked to increased risk for both myocardial infarction and stroke in several populations. A number of polymorphisms and haplotypes in the gene encoding FLAP have been identified in individuals (U.S. Patent Application 2005113408; Sayers, *Clin. Exp. Allergy*, 33(8):1103-10, 2003; Kedda, *et al.*, *Clin. Exp. Allergy*, 35(3):332-8, 2005). Particular FLAP haplotypes have been linked to myocardial infarction and stroke in several populations (Helgadottir A *et al.* *Nature Genet.* 36:233-239 (2004); Helgadottir A *et al.* *Am J Hum Genet* 76:505-509 (2004); Lohmussaar E *et al.* *Stroke* 36: 731-736 (2005); Kajimoto K *et al.* *Circ J* 69:1029-1034 (2005). Previously, polymorphisms in certain genes have been demonstrated to correlate with responsiveness to given therapies, for example, the responsiveness of cancers to particular chemotherapeutic agents (Erichsen, *et al.*, *Br. J. Cancer*, 90(4):747-51, 2004; Sullivan, *et al.*, *Oncogene*, 23(19):3328-37, 2004). Therefore, patients who are under consideration for treatment with the novel FLAP inhibitors described herein, or drug combinations that include such novel FLAP inhibitors, may be screened for potential responsiveness to treatment based on their FLAP polymorphisms, or haplotypes.

[00893] Additionally, polymorphisms in any of the synthetic or signaling genes dedicated to the leukotriene pathway could result in a patient who is more responsive or less responsive to leukotriene modulator therapy (either FLAP or 5-LO inhibitor or leukotriene receptor antagonists). The genes dedicated to the leukotriene pathway are 5-lipoxygenase, 5-lipoxygenase-activating protein, LTA₄ hydrolase, LTC₄ synthase, LTB₄ receptor 1 (BLT₁), LTB₄ receptor 2 (BLT₂), cysteinyl leukotriene receptor 1 (CysLT₁R), cysteinyl leukotriene receptor 2 (CysLT₂R). For example, the 5-LO gene has been linked to aspirin intolerant asthma and airway hyperresponsiveness (Choi JH *et al.* *Hum Genet* 114:337-344 (2004); Kim, SH *et al.* *Allergy* 60:760-765 (2005). Genetic variants in the promoter region of 5-LO have been shown to predict clinical responses to a 5LO inhibitor in asthmatics (Drazen *et al.*, *Nature Genetics*, 22, p168-170, (1999). The LTC₄ synthase gene has been linked to atopy and asthma (Moissidis I *et al.* *Genet Med* 7:406-410 (2005). The CysLT₂ receptor has been linked to asthma and atopy (Thompson MD *et al.* *Pharmacogenetics* 13:641-649 (2003); Pillai SG *et al.* *Pharmacogenetics* 14:627-633 (2004); Park JS *et al.* *Pharmacogenet Genomics* 15:483-492 (2005); Fukai H *et al.* *Pharmacogenetics* 14:683-690 (2004). Any polymorphisms in any leukotriene pathway gene or combination of polymorphisms or haplotypes may result in altered sensitivity of the patient to therapy aimed at reducing the pathological effects of leukotrienes. Selection of patients who might best respond to the leukotriene modulator therapies described herein may include knowledge of polymorphisms in the leukotriene pathway genes and also knowledge of the expression of leukotriene-driven mediators. Patient selection could be made on the basis of leukotriene pathway genotype alone, phenotype alone (biomarkers or functional markers) or any combination of genotype and phenotype.

[00894] A "haplotype," as described herein, refers to a combination of genetic markers ("alleles"). A haplotype can comprise one or more alleles (e.g., a haplotype containing a single SNP), two or more alleles, three or more alleles, four or more alleles, or five or more alleles. The genetic markers are particular "alleles" at "polymorphic sites" associated with FLAP. A nucleotide position at which more than one sequence is possible in a population is referred to herein as a "polymorphic site." Where a polymorphic site is a single nucleotide in length, the site is referred to as a single nucleotide polymorphism ("SNP"). For example, if at a particular chromosomal location,

one member of a population has an adenine and another member of the population has a thymine at the same position, then this position is a polymorphic site, and, more specifically, the polymorphic site is a SNP.

Polymorphic sites can allow for differences in sequences based on substitutions, insertions or deletions. Each version of the sequence with respect to the polymorphic site is referred to herein as an "allele" of the polymorphic site. Thus, in the previous example, the SNP allows for both an adenine allele and a thymine allele.

[00895] Typically, a reference sequence is referred to for a particular sequence. Alleles that differ from the reference are referred to as "variant" alleles. The term "variant FLAP" as used herein, refers to a sequence that differs from a reference FLAP sequence, but is otherwise substantially similar. The genetic markers that make up the haplotypes described herein are FLAP variants. In certain embodiments the FLAP variants are at least about 90% similar to a reference sequence. In other embodiments the FLAP variants are at least about 91% similar to a reference sequence. In other embodiments the FLAP variants are at least about 92% similar to a reference sequence. In other embodiments the FLAP variants are at least about 93% similar to a reference sequence. In other embodiments the FLAP variants are at least about 94% similar to a reference sequence. In other embodiments the FLAP variants are at least about 95% similar to a reference sequence. In other embodiments the FLAP variants are at least about 96% similar to a reference sequence. In other embodiments the FLAP variants are at least about 97% similar to a reference sequence. In other embodiments the FLAP variants are at least about 98% similar to a reference sequence. In other embodiments the FLAP variants are at least about 99% similar to a reference sequence.

[00896] Additionally, in certain embodiments the FLAP variants differ from the reference sequence by at least one base, while in other embodiments the FLAP variants differ from the reference sequence by at least two bases. In other embodiments the FLAP variants differ from the reference sequence by at least three bases, and in still other embodiments the FLAP variants differ from the reference sequence by at least four bases.

[00897] Additional variants can include changes that affect a polypeptide, e.g., the FLAP polypeptide. The polypeptide encoded by a reference nucleotide sequence is the "reference" polypeptide with a particular reference amino acid sequence, and polypeptides encoded by variant alleles are referred to as "variant" polypeptides with variant amino acid sequences. The FLAP nucleic acid sequence differences, when compared to a reference nucleotide sequence, can include the insertion or deletion of a single nucleotide, or of more than one nucleotide, resulting in a frame shift; the change of at least one nucleotide, resulting in a change in the encoded amino acid; the change of at least one nucleotide, resulting in the generation of a premature stop codon; the deletion of several nucleotides, resulting in a deletion of one or more amino acids encoded by the nucleotides; the insertion of one or several nucleotides, such as by unequal recombination or gene conversion, resulting in an interruption of the coding sequence; duplication of all or a part of a sequence; transposition; or a rearrangement of a nucleotide sequence, as described in detail above. Such sequence changes alter the polypeptide encoded by a FLAP nucleic acid. For example, if the change in the nucleic acid sequence causes a frame shift, the frame shift can result in a change in the encoded amino acids, and/or can result in the generation of a premature stop codon, causing generation of a truncated polypeptide.

[00898] By way of example, a polymorphism associated with a susceptibility to myocardial infarction (MI), acute coronary syndrome (ACS), stroke or peripheral arterial occlusive disease (PAOD) can be a synonymous change in one or more nucleotides (i.e., a change that does not result in a change in the amino acid sequence).

Such a polymorphism can, for example, alter splice sites, decrease or increase expression levels, affect the

stability or transport of mRNA, or otherwise affect the transcription or translation of the polypeptide. The haplotypes described below are found more frequently in individuals with MI, ACS, stroke or PAOD than in individuals without MI, ACS, stroke or PAOD. Therefore, these haplotypes may have predictive value for detecting a susceptibility to MI, ACS, stroke or PAOD in an individual.

- 5 [00899] Several variants of the FLAP gene have been reported to correlate with the incidence of myocardial infarction in patients (Hakonarson, *JAMA*, 293(18):2245-56, 2005), plus FLAP gene markers reportedly associated with the risk for developing asthma have been described in U.S. Patent No. 6,531,279. Methods for identifying FLAP sequence variants are described, e.g., in U.S. Publication No. 2005/0113408, and in U.S. Patent No. 6,531,279, incorporated herein by reference herein in their entirety.
- 10 [00900] By way of example only, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S99, SG13S25, SG13S377, SG13S106, SG13S32 and SG13S35 at the 13q12-13 locus. Or, the presence of the alleles T, G, G, G, A and G at SG13S99, SG13S25, SG13S377, SG13S106, SG13S32 and SG13S35, respectively (the B6 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. Or, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S99,
- 15 SG13S25, SG13S106, SG13S30 and SG13S42 at the 13q12-13 locus. Or, the presence of the alleles T, G, G, G and A at SG13S99, SG13S25, SG13S106, SG13S30 and SG13S42, respectively (the B5 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. Or, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S25, SG13S106, SG13S30 and SG13S42 at the 13q12-13 locus. Or, the presence of the alleles G, G, G and A at SG13S25, SG13S106, SG13S30 and SG13S42, respectively (the B4
- 20 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. Or, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S25, SG13S106, SG13S30 and SG13S32 at the 13q 12-13 locus. Or, the presence of the alleles G, G, G and A at SG13S25, SG13S106, SG13S30 and SG13S32, respectively (the Bs4 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. In such embodiments just described, patients who are under consideration for treatment with
- 25 compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be screened for potential responsiveness to treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), based on such haplotypes.
- 30 [00901] By way of example only, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S99, SG13S25, SG13S114, SG13S89 and SG13S32 at the 13q12-13 locus. Or, the presence of the alleles T, G, T, G and A at SG13S99, SG13S25, SG13S114, SG13S89 and SG13S32, respectively (the A5 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. Or, a haplotype associated with a susceptibility to myocardial infarction or stroke comprises markers SG13S25, SG13S114, SG13S89 and
- 35 SG13S32 at the 13q12-13 locus. Or, the presence of the alleles G, T, G and A at SG13S25, SG13S114, SG13S89 and SG13S32, respectively (the A4 haplotype), is diagnostic of susceptibility to myocardial infarction or stroke. In such embodiments just described, patients who are under consideration for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C),
- 40 Formula (E), Formula (F), Formula (G), or Formula (H), may be screened for potential responsiveness to

treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), based on such haplotypes.

[00902] Detecting haplotypes can be accomplished by methods known in the art for detecting sequences at polymorphic sites, and therefore patients may be selected using genotype selection of FLAP, 5-LO or other leukotriene pathway gene polymorphisms. The presence or absence of a leukotriene pathway gene polymorphism or haplotype can be determined by various methods, including, for example, using enzymatic amplification, restriction fragment length polymorphism analysis, nucleic acid sequencing, electrophoretic analysis of nucleic acid from the individual, or any combination thereof. In certain embodiments, determination of a SNP or haplotype may identify patients who will respond to, or gain benefit from, treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). By way of example, methods of diagnosing a susceptibility to myocardial infarction or stroke in an individual, comprises determining the presence or absence of certain single nucleotide polymorphisms (SNPs) or of certain haplotypes, wherein the presence of the SNP or the haplotype is diagnostic of susceptibility to myocardial infarction or stroke.

Phenotype Analysis: Biomarkers

[00903] Patients who are under consideration for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be screened for potential responsiveness to treatment based on leukotriene-driven inflammatory biomarker phenotypes.

[00904] Patient screening based on leukotriene-driven inflammatory biomarker phenotypes may be used as an alternative to, or it may be complimentary with, patient screening by leukotriene pathway gene haplotype detection. The term "biomarker" as used herein refers to a characteristic which can be measured and evaluated as an indicator of normal biological processes, pathological processes, or pharmacological responses to therapeutic intervention. Thus a biomarker may be any substance, structure or process which can be measured in the body, or its products, and which may influence or predict the incidence of outcome or disease. Biomarkers may be classified into markers of exposure, effect, and susceptibility. Biomarkers can be physiologic endpoints, by way of example blood pressure, or they can be analytical endpoints, by way of example, blood glucose, or cholesterol concentrations. Techniques, used to monitor and/or measure biomarkers include, but are not limited to, NMR, LC-MS, LC-MS/MS, GC-MS, GC-MS/MS, HPLC-MS, HPLC-MS/MS, FT-MS, FT-MS/MS, ICP-MS, ICP-MS/MS, peptide/protein sequencing, nucleic acid sequencing, electrophoresis techniques, immuno-assays, immuno-blotting, *in-situ* hybridization, fluorescence *in-situ* hybridization, PCR, radio-immuno assays, and enzyme-immuno assays. Single nucleotide polymorphisms (SNPs) have also been useful for the identification of biomarkers for propensity to certain diseases and also susceptibility or responsiveness to drugs such as chemotherapeutic agents and antiviral agents. These techniques, or any combination thereof, may be used to screen patients for leukotriene-dependent or leukotriene mediated diseases or conditions, wherein such patients may be beneficially treated with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H).

[00905] By way of example only, patients may be selected for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), by screening for enhanced inflammatory blood biomarkers such as, but not limited to, stimulated LTB₄, LTC₄, LTE₄, myeloperoxidase (MPO), eosinophil peroxidase (EPO), C-reactive protein (CRP), soluble intracellular adhesion molecule (sICAM), monocyte chemoattractant protein (MCP-1), monocyte inflammatory protein (MIP-1 α), interleukin-6 (IL-6), the TH2 T cell activators interleukin 4 (IL-4), and 13 (IL-13) and other inflammatory cytokines. In certain embodiments, patients with inflammatory respiratory diseases, including but not limited to, asthma and COPD, or with cardiovascular diseases, are selected as those most likely to be responsive to leukotriene synthesis inhibition using compounds of any of Formula (G), Formula (G-I), or Formula (G-II), by using a panel of leukotriene driven inflammatory biomarkers.

Phenotype Analysis: Functional Markers

[00906] Patients who are under consideration for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be screened for response to known modulators of the leukotriene pathway. Patient screening by evaluation of functional markers as indicators of a patient's response to known modulators of the leukotriene pathway may be used as an alternative to, or it may be complimentary with, patient screening by leukotriene pathway gene haplotype detection (genotype analysis) and/or monitoring/measurement of leukotriene-driven inflammatory biomarker phenotypes. Functional markers may include, but are not limited to, any physical characteristics associated with a leukotriene dependent condition or disease, or knowledge of current or past drug treatment regimens.

[00907] By way of example only, the evaluation of lung volume and/or function may be used as a functional marker for leukotriene-dependent or leukotriene mediated diseases or conditions, such as respiratory diseases.

Lung function tests may be used to screen patients, with such leukotriene-dependent or leukotriene mediated diseases or conditions, for treatment using compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical compositions or medicaments which include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). Such tests include, but are not limited to, evaluation of lung volumes and capacities, such as tidal volume, inspiratory reserve volume, expiratory reserve volume, residual volume, inspiratory capacity, functional residual capacity, vital capacity, total lung capacity, respiratory minute volume, alveolar ventilation, timed vital capacity, and ventilatory capacity. Method of measurement of lung volumes and capacities include, but are not limited to, maximum expiratory flow volume curve, forced expiratory volume in 1 sec. (FEV1), peak expiratory flow rate. In addition, other lung function tests used as functional markers for patient evaluation described herein include, but are not limited to, respiratory muscle power, maximum inspiratory pressure, maximum expiratory pressure, transdiaphragmatic pressure, distribution of ventilation, single breath nitrogen test, pulmonary nitrogen washout, and gas transfer.

[00908] Additionally, the knowledge of a patient's past or current treatment regimen may be used as a functional marker to assist in screening patients for treatment of leukotriene dependent conditions or diseases using compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or

Formula (H), or pharmaceutical compositions or medicaments which include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H). By way of example only, such treatment regimens may include past or current treatment using zileuton (ZyfloTM), montelukast (SingulairTM), pranlukast (OnonTM), zafirlukast (AccolateTM).

5 [00909] Also, patients who are under consideration for treatment with compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or drug combinations described herein that include compounds of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), may be screened for functional markers which include, but are not limited to, reduced eosinophil and/or basophil, and/or neutrophil, and/or monocyte and/or dendritic cell and/or
10 lymphocyte recruitment, decreased mucosal secretion, decreased mucosal edema, and/or increased bronchodilation.

[00910] Methods for the identification of a patient in need of treatment for leukotriene-dependent or leukotriene mediated conditions or diseases, and exemplary, non-limiting treatment methods are shown in Figure 12, Figure 13 and Figure 14, wherein a patient sample is analyzed and the information obtained is used to identify possible
15 treatment methods. It is expected that one skilled in the art will use this information in conjunction with other patient information, including, but not limited to age, weight, sex, diet, and medical condition, to choose a treatment method. It is also expected that each piece of information will be given a particular weight in the decision process. In certain embodiments, the information obtained from the diagnostic methods described above and any other patient information, including, but not limited to age, weight, sex, diet, and medical condition, are
20 incorporated into an algorithm used to elucidate a treatment method, wherein each piece of information will be given a particular weight in the decision process.

[00911] In certain embodiments a patient sample is analyzed for leukotriene gene haplotypes, by way of example only, FLAP haplotypes, and the information obtained identifies a patient in need of treatment using various treatment methods. Such treatment methods include, but are not limited to, administering a therapeutic effective
25 amount of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), administering a therapeutic effective amount of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E),
30 Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in combination with a therapeutic effective amount of a leukotriene receptor antagonist (by way of example, CysLT₁/CysLT₂ antagonist or CysLT₁ antagonist), or administering a therapeutic effective amount of a
35 compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in combination with a therapeutic
40 effective amount of another anti-inflammatory agent. In other embodiments a patient sample is analyzed for leukotriene gene haplotypes, by way of example only, FLAP haplotypes, and/or phenotype biomarkers, and/or phenotype functional marker responses to leukotriene modifying agents. The patient may then be treated using various treatment methods. Such treatment methods include, but are not limited to, administering a therapeutic effective amount of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F),

Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), administering a therapeutic effective amount of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in combination with a therapeutic effective amount of a leukotriene receptor antagonist (by way of example, CysLT₁/CysLT₂ antagonist or CysLT₁ antagonist), or administering a therapeutic effective amount of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), or pharmaceutical composition or medicament which includes a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), in combination with a therapeutic effective amount of another anti-inflammatory agent. In still other embodiments a patient sample is analyzed for leukotriene gene haplotypes, by way of example only, FLAP haplotypes, and phenotype biomarkers, and phenotype functional marker responses to leukotriene modifying agents. The patient may then be treated using various treatment methods. Such treatment methods include, but are not limited to, administering a therapeutic effective amount of a FLAP inhibitor, or pharmaceutical composition or medicament which includes a FLAP inhibitor, administering a therapeutic effective amount of a FLAP inhibitor, or pharmaceutical composition or medicament which includes a FLAP inhibitor, in combination with a therapeutic effective amount of a leukotriene receptor antagonist (by way of example, CysLT₁/CysLT₂ antagonist or CysLT₁ antagonist), or administering a therapeutic effective amount of a FLAP inhibitor, or pharmaceutical composition or medicament which includes a FLAP inhibitor, in combination with a therapeutic effective amount of another anti-inflammatory agent.

Kits/Articles of Manufacture

[00912] For use in the therapeutic applications described herein, kits and articles of manufacture are also described herein. Such kits can comprise a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers can be formed from a variety of materials such as glass or plastic.

[00913] The articles of manufacture provided herein contain packaging materials. Packaging materials for use in packaging pharmaceutical products are well known to those of skill in the art. See, e.g., U.S. Patent Nos. 5,323,907, 5,052,558 and 5,033,252. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, inhalers, pumps, bags, vials, containers, syringes, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment. A wide array of formulations of the compounds and compositions provided herein are contemplated as are a variety of treatments for any disease, disorder, or condition that would benefit by inhibition of FLAP, or in which FLAP is a mediator or contributor to the symptoms or cause.

[00914] For example, the container(s) can include one or more compounds described herein, optionally in a composition or in combination with another agent as disclosed herein. The container(s) optionally have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits optionally comprising a compound with an identifying description or label or instructions relating to its use in the methods described herein.

[00915] A kit may typically include one or more additional containers, each with one or more of various materials (such as reagents, optionally in concentrated form, and/or devices) desirable from a commercial and user standpoint for use of a compound described herein. Non-limiting examples of such materials include, but not limited to, buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use. A set of instructions will also typically be included.

[00916] A label can be on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. A label can be used to indicate that the contents are to be used for a specific therapeutic application. The label can also indicate directions for use of the contents, such as in the methods described herein.

[00917] In certain embodiments, the pharmaceutical compositions can be presented in a pack or dispenser device which can contain one or more unit dosage forms containing a compound provided herein. The pack can for example contain metal or plastic foil, such as a blister pack. The pack or dispenser device can be accompanied by instructions for administration. The pack or dispenser can also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, can be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions containing a compound provided herein formulated in a compatible pharmaceutical carrier can also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

EXAMPLES

[00918] These examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein. That is, the specific compounds disclosed herein and substitution patterns described herein (e.g. R^6 , R^7 , R^{11}) are illustrative only. That is, a particular functional group presented specifically herein can be substituted into any of the other formula or may be applied to any other set of substituents. For example only, the R^6 of compound 2-12 can be used to replace the R^6 of compound 2-23 to form a new compound. All such combinations and substitutions of substituents are herein described.













Preparation of Intermediates used in the Synthesis of compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), Formula (H).















[00919] Starting materials and intermediates used in the synthesis of compounds of compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), Formula (H) are commercially available or can be synthesized by synthetic methods known in the art or described herein. The preparation of intermediates, such as, for example, those shown in Table II, which are used herein and not commercially available is described below. Other intermediates not specifically mentioned herein and used in the synthesis of compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), Formula (H), can be prepared using the methods described herein or known in the art.

Table II. Intermediates used in the Synthesis of Compounds Described herein

Compound #	Structure	Compound Name	Method for Preparation
Int-5		C-(Di-imidazol-1-yl)-methylenamine	Route 8, Step 1
Int-10		3-Bromomethyl-azetidine-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Steps 1-3a SM: 3-Azetidinecarboxylic acid (Sigma Aldrich)
Int-19		2-Chloro- <i>N</i> -cyclopropyl-acetamide	Route 2, Step 1 SM: Cyclopropylamine (Sigma Aldrich)
Int-20		2-Chloromethyl-1,4,5,6-tetrahydropyrimidine Hydrochloride	Route 3, Steps 1-2 SM: Chloro-acetonitrile (Sigma Aldrich)
Int-21		(<i>S</i>)-2-(Toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Step 3c SM: (<i>S</i>)-(-)-1-(<i>tert</i> -Butoxycarbonyl)-2-pyrroli dinemethanol (Sigma Aldrich)
Int-22		(<i>R</i>)-2-(Toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Step 3c SM: (<i>R</i>)-(+)-1-(<i>tert</i> -Butoxycarbonyl)-2-pyrroli dinemethanol (Sigma Aldrich)
Int-23		(<i>S</i>)-2-Methanesulfonyloxymethyl-piperidine-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Step 3d SM: 1-Boc-(<i>S</i>)-2-piperidinemethanol (Chem Impex)
Int-24		Toluene-4-sulfonic acid (<i>S</i>)-5-oxo-pyrrolidin-2-ylmethyl ester	Route 1, Step 3c SM: (<i>S</i>)-(+)-5-(Hydroxymethyl)-2-pyrrolidinone (Sigma Aldrich)
Int-25		Toluene-4-sulfonic acid (<i>R</i>)-5-oxo-pyrrolidin-2-ylmethyl ester;	Route 1, Step 3c SM: (<i>R</i>)-(-)-5-(Hydroxymethyl)-2-pyrrolidinone (Acros Organics)
Int-27		3-Chloromethyl-5-methyl-isoxazole Hydrochloride	Route 4, Step 4 SM: (5-Methylisoxazol-3-yl)methanol (Acros Organics)
Int-28		3-Chloromethyl-1,5-dimethyl-1 <i>H</i> -pyrazole Hydrochloride	Route 4, Step 4 SM: (1,5-Dimethyl-1 <i>H</i> -pyrazol-3-yl)methanol (Acros Organics)
Int-29		5-Chloromethyl-1,3-dimethyl-1 <i>H</i> -pyrazole Hydrochloride	Route 4, Step 4 SM: (1,3-Dimethyl-1 <i>H</i> -pyrazol-5-yl)methanol (Acros Organics)
Int-30		2-(Toluene-4-sulfonyloxymethyl)-2,3-dihydro-indole-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Steps 1-3c SM: Indoline-2-carboxylic Acid (Sigma Aldrich)

Int-31		(S)-2-(Toluene-4-sulfonyloxymethyl)-2,3-dihydro-indole-1-carboxylic acid <i>tert</i> -Butyl ester	Route 1, Steps 1, 3c SM: (S)-(+)-2-Indolinemethanol (Sigma Aldrich)
Int-32		2-Chloromethyl-imidazo[1,2-a]pyridine	Route 4, Step 4 SM: Imidazo[1,2-a]pyridin-2-ylmethanol (Acros Organics)
Int-33		Toluene-4-sulfonic acid (S)-2- <i>tert</i> -butoxycarbonylamino-2-phenyl-ethyl ester	Route 1, Steps 1, 3c SM: (S)-(+)-2-Phenylglycinol (Sigma Aldrich)
Int-34		Toluene-4-sulfonic acid (R)-2- <i>tert</i> -butoxycarbonylamino-2-phenyl-ethyl ester	Route 1, Step 3c SM: (R)-(-)-N-(<i>tert</i> -Butoxycarbonyl)-2-phenylglycinol (Sigma Aldrich)
Int-38		2-Chloro- <i>N</i> -(4-fluoro-phenyl)-acetamide	Route 2, Step 1 SM: 4-Fluoroaniline (Sigma Aldrich)
Int-39		2-Chloro- <i>N</i> -pyridin-3-yl-acetamide	Route 2, Step 1 SM: 3-Aminopyridine (Sigma Aldrich)
Int-44		2-Chloromethyl-pyridin-1-ol	Route 4, Step 1 SM: 2-Chloromethyl-pyridine Hydrochloride (Sigma Aldrich)
Int-45		2-Chloromethyl-6-methyl-pyridine Hydrochloride	Route 4, Step 4 SM: 6-Methyl-2-pyridinemethanol (Sigma Aldrich)
Int-46		2-Chloromethyl-5-methyl-pyridine Hydrochloride	Route 4, Steps 1-4 SM: 2,5-Lutidine (Sigma Aldrich)
Int-47		2-Chloromethyl-4-methyl-pyridine Hydrochloride	Route 4, Steps 1-4 SM: 2,4-Lutidine (Sigma Aldrich)
Int-48		2-Chloromethyl-3-methyl-pyridine Hydrochloride	Route 4, Steps 1-4 SM: 2,3-Lutidine (Sigma Aldrich)
Int-49		2-Chloromethyl-3,5-dimethyl-pyridine Hydrochloride	Route 4, Steps 1-4 SM: 2,3,5-Collidine (Sigma Aldrich)
Int-50		2-Chloromethyl-6-fluoro-pyridine Hydrochloride	Route 5, Step 3c SM: 2-Fluoro-6-methylpyridine (Oakwood Product)
Int-51		2-Chloromethyl-6-bromo-pyridine Hydrochloride	Route 4, Step 4 SM: (6-Bromo-pyridin-2-yl)-methanol (Sigma Aldrich)
Int-52		2-Chloromethyl-5-ethyl-pyridine	Route 4, Steps 1-4 SM: 5-Ethyl-2-methylpyridine (Sigma Aldrich)

Int-53		2-Chloromethyl-5-chloro-pyridine	Route 1, Step 2; Route 4, Step 4 SM: 5-Chloropyridine-2-carboxylic Acid (Matrix Scientific)
Int-54		Methanesulfonic acid (S)-1-pyridin-2-yl-ethyl ester	Route 1, Step 3 SM: (R)- α -Methyl-2-pyridinemethanol (Sigma Aldrich)
Int-55		Methanesulfonic acid (R)-1-pyridin-2-yl-ethyl ester	Route 1, Step 3 SM: (S)- α -Methyl-2-pyridinemethanol (Sigma Aldrich)
Int-57		2-Bromomethyl-7-fluoro-quinoline	Route 5, Step 3a SM: 7-Fluoro-2-methylquinoline (Sigma Aldrich)
Int-58		2-Bromomethyl-6-fluoro-quinoline	Route 5, Step 3a SM: 6-Fluoro-2-methylquinoline (Sigma Aldrich)
Int-59		2-Chloromethyl-6-methyl-quinoline	Route 4, Steps 1-4 SM: 2,6-Dimethylquinoline (Sigma Aldrich)
Int-60		2-Chloro-6-bromomethyl-quinoline	Route 5, Steps 1-3a SM: Cinnamoyl chloride (Sigma Aldrich) and p-toluidine (Sigma Aldrich)
Int-71		5-Fluoro-2-(4-iodomethyl-phenyl)-thiazole	Route 6, Step 1-2a; Route 1, Step 3b
Int-72		Methanesulfonic acid 4-(5-methyl-thiazol-2-yl)-benzyl ester	Route 6, Step 1-2b; Route 1, Step 3d
Int-73		Methanesulfonic acid 4-(6-methoxy-pyridin-3-yl)-benzyl ester	Route 6, Step 1; Route 1, Step 3d
Int-74		4-(3-Bromomethyl-phenyl)-4-methoxy-tetrahydro-pyran	Route 9, Step 1; Route 5, Step 3a
Int-75		5-Bromo-2-chloromethyl-pyridine	Route 4, Step 4 SM: (5-Bromo-pyridin-2-yl)-methanol (Biofine International)

Int-76		2-Bromo-5-iodomethyl-pyridine	Route 1, Step 3b SM: (6-Bromo-pyridin-3-yl)-methanol (Biofine International)
		[4-(5-Trifluoromethyl-pyridin-2-yl)-phenyl]-methyl mesylate	Route 6, Step 1; Route 1, Step 3d
		2-(4-bromomethyl-phenyl)-5-trifluoromethylpyridine	Route 6, Step 1; Route 5, Step 3d
Int-118		5-Bromo-pyrazin-2-ylamine	Route 5, Step 3b SM: Aminopyrazine (Lancaster)
Int-135		3-Phenoxy-benzoyl chloride	Route 7, Step 1 SM: 3-Phenoxy-benzoic acid (Sigma Aldrich)
Int-136		4-Phenoxy-benzoyl chloride	Route 7, Step 1 SM: 4-Phenoxy-benzoic acid (Sigma Aldrich)
Int-140		1-tert-Butylsulfanyl-4,4-dimethyl-pentan-2-one	Route 10, Steps 1-2
Int-141		2-Bromomethyl-5-methoxy-pyridine	Route 9, Step 1b; Route 5, Step 3a SM: 5-Hydroxy-2-methylpyridine (Sigma Aldrich)
Int-142		6-Bromomethyl-nicotinonitrile	Route 5, Step 3a SM: 5-Cyano-2-methylpyridine (Alfa Aesar)
Int-143		5-Bromo-2-chloromethyl-pyridine	Route 4, Step 4 SM: 5-Bromo-2-methylpyridine (Alfa Aesar)
Int-144		6-Bromo-2-bromomethyl-quinoline	Route 5, Step 3a SM: 6-Bromoquinoline (Trans World Chemicals)
Int-145		2-Chloro-5-fluoromethyl-pyridine	Iee, K. C. <i>et al.</i> , <i>J. Org. Chem.</i> 1999 , 8576.
Int-146		6-Chloromethyl-2,3-dimethyl-pyridine	Route 4, Step 1; Route 11, Steps 1-3; Route 4, Step 4 SM: 2,3-Lutidine (Sigma Aldrich)
Int-147		2-Chloromethyl-5-methyl-pyrazine	Route 5, Step 3c SM: 2,5-Dimethylpyrazine (Sigma Aldrich)

Int-148		2-Chloromethyl-quinoxaline	Kolasa, T. <i>et al.</i> , <i>J. Med. Chem.</i> 2000 , 690.
Int-149		2-Chloromethyl-5-methyl-pyridin-1-ol	Route 4, Step 1, adding NaHCO ₃ as base to neutralize HCl salt
Int-150		2-Chloromethyl-quinolin-1-ol	Route 4, Step 1, adding NaHCO ₃ as base to neutralize HCl salt
Int-151		3-Chloromethyl-6-methyl-pyridazine	Route 12, Step 1; Route 5, Step 3c SM: Acetylacetone (Sigma Aldrich)
Int-152		2-Bromomethyl-indole-1-carboxylic acid <i>tert</i> -butyl ester	Freed, J. D. <i>et al.</i> , <i>J. Org. Chem.</i> 2001 , 839.
Int-153		5-Methylsulfanyl-4-oxo-pentanoic acid methyl ester	Prepared according to the procedures described in US patent 5,288,743 issued Feb 22, 1994
Int-154		5- <i>tert</i> -Butylsulfanyl-2-methyl-4-oxo-pentanoic acid ethyl ester	Prepared according to the procedures described in US patent 5,288,743 issued Feb 22, 1994
Int-155		5- <i>tert</i> -Butylsulfanyl-2,2-diethyl-4-oxo-pentanoic acid ethyl ester	Prepared according to the procedures described in US patent 5,288,743 issued Feb 22, 1994
Int-156		1-(3- <i>tert</i> -Butylsulfanyl-2-oxo-propyl)-cyclopentanecarboxylic acid methyl ester	Prepared according to the procedures described in US patent 5,288,743 issued Feb 22, 1994
Int-157		1- <i>tert</i> -Butylsulfanyl-propan-2-one	Bradsher <i>et al.</i> , <i>J. Am. Chem. Soc.</i> 1954 , 114.
Int-158		3- <i>tert</i> -Butylsulfanyl-2-oxo-propionic acid ethyl ester	Kolasa, T. <i>et al.</i> , <i>J. Med. Chem.</i> 2000 , 690.

Route 1:**Step 1: BOC Protection (Int-10)**

[00920] 3-Azetidinecarboxylic acid (Sigma Aldrich, 0.25 g, 2.5 mmol) was dissolved in *t*BuOH (5mL) and 1N NaOH (2.7 mL, 2.7 mmol). Di-*tert*-Butyl dicarbonate (0.59 g, 2.7 mmol) was added, and the reaction was stirred overnight at room temperature. The reaction was diluted with water, acidified slowly to pH 4 with 1N HCl, and the mixture was extracted with EtOAc until all product was removed from the aqueous layer by ninhydrin stain. The combined organic layers were dried, filtered, and concentrated to give the desired product.

Step 2: Borane Reduction (Int-10)

[00921] Acid from Step 1 (0.7 g, 3.5 mmol) was dissolved in THF and cooled to 0°C under N₂. Borane-THF complex was added to the solution, and the reaction was stirred at room temperature overnight. The reaction was

cooled to 0°C and quenched with water. The mixture was extracted 3 times with EtOAc, the combined organic layers were dried over MgSO₄, filtered, and concentrated. The crude material was filtered through a plug of silica gel and eluted with EtOAc to give the desired compound.

Step 3a: Br₂ Bromide Formation (Int-10)

5 [00922] Triphenylphosphine (1.7 g, 6.5 mmol) was dissolved in DMF and cooled to 0°C. Bromine (0.31 mL, 5.9 mmol) was added slowly, and the solution was stirred for 30 minutes. Alcohol from Step 2 (0.32 g, 2.0 mmol) was added in DMF and the reaction was stirred at room temperature overnight. The mixture was diluted with water, extracted 3 times with EtOAc, and the combined organic layers were dried over MgSO₄, filtered, and concentrated. The crude material was filtered through a plug of silica gel and eluted with EtOAc to give the
10 desired compound.

Step 3b: I₂ Iodide Formation (Int-73)

[00923] (6-Bromo-pyridin-3-yl)-methanol (0.5g, 2.7 mmol) was dissolved in toluene (20 mL). Triphenylphosphine (0.9 g, 3.5 mmol) and imidazole (0.4 g, 6.0 mmol) were added, followed by a solution of iodine (0.88 g, 3.5 mmol) in toluene dropwise. The reaction was stirred at room temperature for 15 minutes, and
15 then poured into saturated aq. Na₂CO₃. The organic layer was washed with aq. sodium thiosulfate, water, then dried over MgSO₄, filtered, and concentrated. The crude material was purified on silica gel (EtOAc:hexanes gradient) to give the desired product.

Step 3c: Tosylation (Int-21)

[00924] (S)-(-)-1-(*tert*-Butoxycarbonyl)-2-pyrrolidinemethanol (1.0 g, 5.0 mmol) was dissolved in pyridine (3
20 mL), and toluenesulfonyl chloride (1.0 g, 5.5 mmol) was added. The reaction was stirred overnight at room temperature, and diluted with water and extracted with EtOAc. The combined organic layers were washed with water, dried over MgSO₄, filtered, and concentrated. The residue was purified on silica gel (0 to 10% EtOAc in hexanes) to give the desired product.

Step 3d: Mesylation (Int-55)

25 [00925] (R)- α -Methyl-2-pyridinemethanol (1.0 g, 8.1 mmol) was dissolved in CH₂Cl₂ (20 mL) and cooled to 0°C. Triethylamine (1.7 mL, 12.2 mmol) was added, followed by methanesulfonyl chloride (0.66 mL, 8.4 mmol) dropwise. The reaction was stirred for 30 minutes, and then diluted with CH₂Cl₂, washed with water, dried over MgSO₄, filtered, and concentrated to obtain the desired product.

Route 2:

30 **Step 1: Amide Formation (Int-19)**

[00926] Cyclopropylamine (0.35 mL, 5.0 mmol) and triethylamine (0.7 mL, 5.1 mmol) were dissolved in CH₂Cl₂ (10 mL). The reaction was cooled to -10°C and chloroacetyl chloride (0.4 mL, 5.0 mmol) was added dropwise. The reaction was stirred at -10°C for 1 hour, then at room temperature for 2 hours, followed by a quench with water. The aqueous layer was extracted with CH₂Cl₂, and the organic layers were dried, filtered, and concentrated
35 to give the desired product.

Route 3:

Step 1: Imine Formation (Int-20)

[00927] Chloroacetonitrile (0.5 g, 6.6 mmol) was dissolved in Et₂O (10 mL) and cooled to 0°C. EtOH (0.43 mL, 7.3 mmol) was added, followed by 4N HCl in 1,4-dioxane (15 mL, 59.6 mmol). The reaction was stirred at 0°C
40 for 4 days, and then concentrated to give the desired product as a white solid.

Step 2: Cyclization (Int-20)

[00928] Imine from Step 1 (0.3 g, 2.0 mmol) was dissolved in EtOH (4 mL) and cooled to 0°C. 1,3-Diaminopropane (0.17 mL, 2.0 mmol) was added, followed by $i\text{Pr}_2\text{NEt}$ (0.35 mL, 2.0 mmol). The reaction was stirred at 0°C for 2 hours, and then 4N HCl in 1,4-dioxane (0.5 mL, 2 mmol) was added. The mixture was filtered, and the filtrate was concentrated to give the desired product.

Route 4:**Step 1: mCPBA Oxidation (Int-46)**

[00929] 2,5-Lutidine (5.0 g, 46.7 mmol) was dissolved in CHCl_3 (125 mL) and cooled to 0°C. m-Chloroperoxybenzoic acid (70%; 13.9 g, 55.2 mmol) was added, and the reaction was stirred overnight at room temperature. The mixture was washed with saturated aq. Na_2CO_3 , dried over Na_2SO_4 , filtered, and concentrated to give the desired product.

Step 2: Acetylation (Int-46)

[00930] The N-oxide from Step 1 (46.7 mmol) was dissolved in acetic anhydride (25 mL) and heated to reflux at 100°C for one hour. The mixture was cooled to room temperature, and ethanol (46.7 mmol) was slowly added to quench the reaction. The solution was evaporated to dryness and purified on silica gel to give the desired product.

Step 3: Hydrolysis (Int-46)

[00931] Acetate from Step 2 (46.7 mmol) was dissolved in concentrated HCl (20 mL) and refluxed for 1 hour. The reaction was cooled and evaporated to dryness to give an orange solid, which was used directly in the next reaction.

Step 4: SOCl_2 Chloride Formation (Int-46)

[00932] Alcohol from Step 3 (1.0 g, 8.1 mmol) was dissolved in thionyl chloride (3 mL) and stirred at room temperature for 30 minutes under N_2 . The mixture was evaporated to dryness to give the desired product as a hydrochloride salt, which was used directly in subsequent reactions.

Route 5:**Step 1: Condensation (Int-60)**

[00933] p-Toluidine (10 g, 60.0 mmol) and triethylamine (8.4 mL, 60.3 mmol) were dissolved in CH_2Cl_2 (200 mL) at room temperature. Cinnamoyl chloride (6.5 g, 60.7 mmol) was added, and the reaction was stirred for 1 hour. The reaction was washed with water, dried, filtered, and concentrated. To the residue was added aluminum chloride (5 g, 37.5 mmol), which was heated neat. After 45 minutes, ice was added to form a precipitate. The mixture was stirred overnight at room temperature. The precipitate was then filtered and dissolved in CH_2Cl_2 , washed with 1N HCl, brine, dried over MgSO_4 , filtered, and concentrated. The residue was recrystallized from ethanol to give the desired quinolinone product.

Step 2: POCl_3 Chloride Formation (Int-60)

[00934] Quinolinone from Step 1 (3.12 g, 19.6 mmol) was heated to 90°C in POCl_3 (10 mL). Once no starting material remained, the reaction was cooled and concentrated. The residue was diluted with EtOAc and saturated aq. NaHCO_3 , and the aqueous layer was extracted with EtOAc. The combined organics were dried, filtered, and concentrated to give the chloroquinoline product.

Step 3a: NBS Bromide Formation (Alkyl) (Int-60)

[00935] Quinoline from Step 2 (19.6 mmol) was heated to 80°C for 1 hour in benzene (200 mL) with NBS (3.6 g, 20.2 mmol) and catalytic benzoyl peroxide. The reaction mixture was concentrated and purified on silica gel to give the desired product.

5 Step 3b: NBS Bromide Formation (Aryl) (Int-118)

[00936] 2-Aminopyrazine (4 g, 42 mmol) was dissolved in water (2 mL) and DMSO (70 mL), and NBS (7.5 g, 42 mmol) was added over 1 hour at 0°C. The reaction was warmed to room temperature and stirred overnight. The mixture was poured onto ice and extracted 4 times with EtOAc. The combined organic layers were washed with 5% Na₂CO₃, water, and brine, dried over MgSO₄, filtered, and concentrated. The residue was purified on
10 silica gel to give the desired product.

Step 3c: NCS Chloride Formation (Int-50)

[00937] 2-Fluoro-6-methylpyridine (1.11 g, 10 mmol), NCS (2.0 g, 15 mmol), and catalytic benzoyl peroxide were dissolved in benzene and heated to reflux overnight. The reaction was concentrated and diluted with water and EtOAc. The organic layer was washed with saturated aq. NaHCO₃, dried, filtered, and concentrated. The
15 residue was purified on silica gel to give the desired product.

Step 3d: PBr₃ Bromide Formation

[00938] 4-(5-Trifluoromethyl-pyridin-2-yl)-phenyl-methanol (5g, 19.8mmol) was dissolved in DME (40mL) at room temperature. PBr₃ (2.8mL, 29.6mmol) was added. The reaction was stirred for 1.5 hours, and then adjusted
20 ph to 7 by adding Sat.NaHCO₃, and the aqueous phase was extracted three times with EtOAc. The combined organic layers were washed with water, dried over MgSO₄, filtered, and concentrated; the crude product was purified by column chromatography to give the desired product. (MS (ES) M+H: 317).

Route 6:**Step 1: Suzuki Coupling (Int-71)**

[00939] To (4-Hydroxymethylphenyl)boronic acid (Combi-Blocks; 1.0 g, 6.6 mmol) in DME/H₂O (16 mL, 2:1)
25 was added 2-bromothiazole (1.2 g, 7.2 mmol) and K₂CO₃ (2.7 g, 19.7 mmol). The reaction was degassed with N₂ for 20 minutes. Pd(PPh₃)₄ (0.76 g, 0.7 mmol) was added and the reaction was further degassed for 10 minutes. The reaction was then heated to 90°C overnight under N₂. LCMS confirmed the formation of the product. The reaction was partitioned between water and EtOAc and the aqueous layer was extracted twice with EtOAc. The combined organic layers were dried over MgSO₄, filtered, concentrated, and purified on silica gel
30 (EtOAc:hexanes gradient) to give the desired product.

Step 2a: Fluorination (Int-71)

[00940] Thiazole from Step 1 (0.35 g, 1.8 mmol) was dissolved in THF (15 mL) and cooled to -78°C under N₂. n-Butyllithium (1.6M; 4.6 mL, 7.3 mmol) was added dropwise, followed by NFSi (1.2 g, 3.7 mmol). The reaction
35 was quenched at -78°C with saturated aq. NH₄Cl, and diluted with EtOAc and water. The aqueous layer was extracted twice with EtOAc, and the combined organics were dried over MgSO₄, filtered, and concentrated. The residue was purified on silica gel to give the desired compound.

Step 2b: Me-Alkylation (Int-72)

[00941] Thiazole from Step 1 (0.33 g, 1.7 mmol) was dissolved in THF (15 mL) and cooled to -78°C under N₂. n-Butyllithium (1.6M; 4.3 mL, 6.7 mmol) was added dropwise, followed by iodomethane (0.16, 2.6 mmol). The
40 reaction was quenched at -78°C with saturated aq. NH₄Cl, and diluted with EtOAc and water. The aqueous layer

was extracted twice with EtOAc, and the combined organics were dried over MgSO_4 , filtered, and concentrated. The residue was purified on silica gel to give the desired compound.

Route 7:

Step 1: Acid Chloride Formation (Int-135)

- 5 [00942] 3-Phenoxy-benzoic acid (0.50 g, 0.23 mmol) was dissolved in CH_2Cl_2 . Oxalyl chloride (0.32 g, 0.25 mmol) was added, followed by 1-2 drops of DMF. The reaction was stirred at room temperature, and then concentrated to give the desired acid chloride.

Route 8:

Step 1: Alkylation (Int-5)

- 10 [00943] To imidazole (0.41 g, 6.0 mmol) in CH_2Cl_2 was added bromoacetonitrile (0.21 g, 2.0 mmol), and the reaction was refluxed for 30 minutes. The mixture was cooled to room temperature and filtered, and the filtrate was concentrated to give the desired product.

Route 9:

Step 1a: Iodomethane Methylation (Int-74)

- 15 [00944] To 4-*m*-Tolyl-tetrahydro-pyran-4-ol (2.5 g, 13.0 mmol) in THF (50 mL) was added sodium hydride (60%; 0.8 g, 20.0 mmol) at room temperature. Iodomethane (1.25 mL, 20 mmol) was added, and the reaction was stirred for 1 hour. The mixture was quenched with water, and the aqueous layer was extracted with EtOAc. The combined organic layers were washed with water, dried over MgSO_4 , filtered, and concentrated. The residue was purified on silica gel to give the desired compound.

- 20 **Step 1b: Trimethylsilyldiazomethane Methylation (Int-141)**

- To 5-Hydroxy-2-methylpyridine (1.0g, 9.16mmol) in toluene (45mL) and MeOH (45mL) at room temperature was added trimethylsilyldiazomethane (2N in ether, 9.2mL, 18.33mmol). The reaction was stirred at room temperature for 30 minutes, and then another two batches of trimethylsilyldiazomethane (2N in ether, 9.2mL, 18.33mmol) were added and the reaction was stirred overnight. Analytical tlc indicated the reaction was
25 complete, so the mixture was concentrated and purified by silica gel chromatography to give the desired methoxy product.

Route 10:

Step 1: Bromination (Int-140)

- 30 [00945] To 4,4-Dimethyl-pentan-2-one (3.7 mL, 26.3 mmol) in MeOH (2.8 mL) at 0°C was added bromine (1.34 mL, 26.3 mmol) in a single stream. The reaction was warmed slowly to 10°C for 30 minutes to initiate the reaction, and then stirred at room temperature for an additional 15 minutes. The reaction was diluted with water and diethyl ether, and the aqueous layer was extracted with diethyl ether three times. The combined organic layers were dried over MgSO_4 , filtered, and concentrated to give the desired product as a colourless liquid.

Step 2: Thiol Addition (Int-140)

- 35 [00946] Bromide from Step 1 (26.3 mmol) was dissolved in THF (50 mL), and the mixture was cooled to 0°C. 2-Methyl-2-propanethiol (2.45 mL, 21.6 mmol) was added, followed by triethylamine (7.9 mL, 56.8 mmol). The reaction was stirred at room temperature for 18 hours, then diluted with water. The aqueous layer was extracted with diethyl ether, and the combined organic layers were dried over MgSO_4 , filtered, and concentrated to give the desired product.

Route 11:**Step 1: Cyanation (Int-146)**

To 2,3-Dimethyl-pyridin-1-ol (17.6g, 0.141mol) (prepared from 2,3-Lutidine via Route 4, Step 1) in CH₂Cl₂ (250mL) was added trimethylsilyl cyanide (19.8mL, 0.148mol). After 30 minutes, N,N-diethylcarbonyl chloride (18.6mL, 0.148mol) was added, and the reaction was stirred for 3 days. The mixture was carefully quenched with 10% aq. potassium carbonate and stirred vigorously for 30 minutes. The aqueous layer was extracted three times with CH₂Cl₂, and the combined organics were dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel chromatography to give the desired nitrile product.

Step 2: Methanolysis (Int-146)

To the nitrile from Step 2 (5g, 37.8mmol) in MeOH (500mL) at -10°C was bubbled dry hydrogen chloride for 15 minutes. The vessel was sealed with a stopper and stirred at room temperature for 3 days. The mixture was diluted with water and evaporated to dryness. The residue was partitioned between EtOAc and saturated NaHCO₃ and stirred vigorously for 30 minutes, and then the aqueous layer was extracted with EtOAc, the combined organic layers were washed with water, dried over MgSO₄, filtered, and concentrated to give the desired ester product.

Step 3: DIBAL-H Reduction (Int-146)

To the ester from Step 3 (5.86g, 35.5mmol) in THF (60mL) at -78°C was added DIBAL-H (1M in THF, 100mL, 100mmol) over 5 minutes. The reaction was warmed to 0°C and quenched with saturated aq. potassium sodium tartrate. Citric acid was added to pH 8, and the mixture was extracted three times with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated, and the residue was purified by silica gel chromatography to give the desired alcohol product.

Route 12:**Step 1: Pyridazine Ring Formation (Int-151)**

Acetylacetone (58.7mL, 0.50mol) and hydrazine hydrate (24.3mL, mol) were refluxed in EtOH (500mL) for 45 minutes, and then cooled to room temperature and evaporated to dryness. The residue was dissolved in benzene (500mL), and Pd/C (3.75g) was added. The mixture was refluxed under N₂ overnight, and then cooled to room temperature, filtered through celite, and evaporated. The crude material was purified by silica gel chromatography (0-6% MeOH in CH₂Cl₂) to give the desired product.

Route 13:**Step 1: Mitsunobu Reaction**


(4-Hydroxymethyl-phenyl)-carbamic acid *tert*-butyl ester (2.6g, 11.6mmol), 2-hydroxypyridine (1.2g, 12.8mmol) and Ph₃P (3.66g, 14.0mmol) were dissolved in THF (20mL) at room temperature under N₂. The reaction mixture was cooled to 0°C, and DIAD (95%, 2.85mL, 14.4mmol) was added dropwise. The reaction mixture was then allowed to warm to room temperature slowly. After 2 hours, the reaction was quenched with saturated aqueous NaCl and diluted with EtOAc and water. The aqueous phase was extracted twice with EtOAc. The combined organic layers were dried over MgSO₄, filtered, and concentrated, and the crude product was purified by column chromatography to give the desired product.

Step 2: BOC Deprotection

[4-(Pyridin-2-yloxymethyl)-phenyl]-carbamic acid *tert*-butyl ester (1.5g, 5mmol) was treated with 4N HCl dioxane (20mL) at room temperature for 2 hours. The pH of the solution was adjusted to pH 8 with saturated aqueous NaHCO₃, and the aqueous phase was extracted three times with EtOAc. The combined organic layers were washed with water, dried over MgSO₄, filtered, and concentrated, and the crude product was purified by column chromatography to give the desired product.

Synthesis of Compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), Formula (H).

Scheme A:

**Example 1.**

[00947] Scheme A illustrates an approach to the synthesis of compounds described herein.

Preparation of Anilines of Structure A-1

[00948] Anilines of structure A-1, where Z is -S-, -CH₂S-, -CH₂CH₂S-, -C(O)CH₂S- or -SCH₂-, and Y is aryl, heteroaryl, heterocycloalkyl, alkyl, or as defined herein, are commercially available or can be readily prepared by methods described herein or known in the art. For example, in one embodiment 4-aminobenzenethiol is reacted with aryl-halides, heteroaryl-halides under transition-metal mediated reaction conditions (e.g. Pd₂dba₃, Xantphos, diisopropylethylamine, dioxane) to provide anilines of structure A-1 where Z is -S-. In other embodiments, alkyl halides, heteroarylmethylhalides, benzyl-halides, heterocycloalkylmethyl-halides, are reacted with 4-aminobenzenethiol under basic conditions to provide anilines of structure A-1, where Z is -CH₂S-, -CH₂CH₂S-, or -C(O)CH₂S-. In yet other embodiments, N-protected 4-aminobenzylhalides can be reacted with aromatic thiol compounds (e.g. benzenethiols, heteroarylthiols, such as, but not limited to, pyridine-2-thiol), alkylthiols,

heterocycloalkyl-thiols, to provide anilines of structure A-1 (after deprotection of the N-protecting group), where Z is $-\text{SCH}_2-$. Other methods of preparing anilines of structure A-1 are known in the art.

4-(5-Methyl-pyridin-2-ylsulfanyl)-phenylamine

[00949] 4-Amino-benzenethiol (2.5g, 20mmol) and 2-Bromo-5-methyl-pyridine (3.4g, 20mmol) were dissolved in 1, 4-dioxane (30mL) and degassed with N_2 for 10 minutes. iPr_2NEt (7mL, 40mmol), Pd_2dba_3 (457mg, 0.5mmol), and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (577mg, 1.0mmol) were added, and the mixture was degassed with N_2 for an additional 10 minutes. The reaction was heated to 60°C overnight, then cooled to room temperature, concentrated, and purified by silica gel chromatography to give the desired product as colorless oil.

4-(Pyridin-2-ylmethylsulfanyl)-aniline

[00950] A mixture of 4-Amino-benzenethiol (2.6g, 20mmol), 2-chloromethylpyridine hydrochloride (4.2g, 26mmol) and cesium carbonate (19.5g, 60mmol) in CH_3CN (50mL) was stirred at room temperature for overnight. The mixture was cooled, poured into water (2L) and extracted with EtOAc six times. The organic layers were washed with brine, dried over MgSO_4 , filtered and concentrated, and the crude product was purified by column chromatography to give the desired product as yellow solid. (MS (ES) $\text{M}+\text{H}$: 217).

[00951] A detailed illustrative example of the reaction conditions shown in Scheme A is described for the synthesis of 2-{3-*tert*-Butylsulfanyl-5-(5-methyl-pyridin-2-ylsulfanyl)-1-[4-(5-trifluoromethyl-pyridin-2-yl)-benzyl]-1*H*-indol-2-ylmethyl}-2-ethyl-butyric acid.

Step 1: Hydrazine formation - [4-(5-methylpyridin-2-ylsulfanyl)-phenyl]-hydrazine.

[00952] To 4-(5-Methyl-pyridin-2-ylsulfanyl)-phenylamine (2.4g, 11.1mmol) in water (32mL) and conc. HCl (2.8mL) at 0°C was added NaNO_2 (0.84g, 12.2mmol) in water (3.2mL). The diazonium salt was allowed to form over 45 minutes and then it was poured slowly over 15 minutes into a rapidly stirred mixture of $\text{Na}_2\text{S}_2\text{O}_4$ (12.2g, 6.2mmol) in water (32mL) and ether (32mL) at 0°C . Stirring continued for 40 minutes, and then the mixture was made basic using conc. KOH. After extraction using EtOAc, the organic layers were washed with water and brine, dried over MgSO_4 , and filtered. To this solution was added saturated HCl in EtOAc and a precipitate formed immediately. Collection of the solids by filtration followed by drying under vacuum provided the title compound as an off-white solid.

[00953] [4-(Pyridin-2-ylmethylsulfanyl)-phenyl]-hydrazine was prepared in the same manner (MS (ES) $\text{M}+\text{H}$: 232) starting from 4-(pyridin-2-ylmethylsulfanyl)-aniline.

Step 2: Indole Formation - 2-[3-*tert*-Butylsulfanyl-5-(5-methyl-pyridin-2-ylsulfanyl)-1*H*-indol-2-ylmethyl]-2-ethyl-butyric acid methyl ester

[00954] [4-(5-Methyl-pyridin-2-ylsulfanyl)-phenyl]-hydrazine (2.2g) and ethyl 5-(*t*-butylthio)-2, 2-dimethyl-4-oxo-pentanoate (2.3g, structure A-6 where $\text{R}_6 = \text{tBuSH}$, $\text{R}_7 = -\text{CH}_2\text{C}(\text{CH}_2\text{CH}_3)_2\text{CO}_2\text{Me}$) in *t*-BuOH (30mL) and HOAc (5mL) was stirred at 80°C for one day. The mixture was poured into water and made basic with solid Na_2CO_3 . The mixture was extracted three times with EtOAc, then washed twice with water, once with brine, dried over MgSO_4 , filtered and concentrated to give dark red-black oil. Silica gel chromatography of the mother liquor (0-20% EtOAc in hexanes) afforded the title compound as brown oil. (MS (ES) $\text{M}+\text{H}$: 471).

[00955] 2-[3-*tert*-Butylsulfanyl-5-(pyridin-2-ylmethylsulfanyl)-1*H*-indol-2-ylmethyl]-2-ethyl-butyric acid methyl ester. (MS (ES) $\text{M}+\text{H}$: 471) was prepared in an analogous manner from [4-(pyridin-2-ylmethylsulfanyl)-phenyl]-hydrazine.

Step 3 : N-alkylation of indoles of structure A-3


[00956] 2-[3-*tert*-Butylsulfanyl-5-(5-methyl-pyridin-2-ylsulfanyl)-1*H*-indol-2-ylmethyl]-2-ethyl-butyric acid methyl ester (540mg, 1.15mmol) and [4-(5-Trifluoromethyl-pyridin-2-yl)-phenyl]-methyl mesylate (380mg, 1.15mmol) were dissolved in DMF (5mL) and cooled to 0°C under N₂. Sodium hydride (60% dispersion in mineral oil; 60mg, 1.5mmol) was added portion wise. The reaction was then allowed to warm to room temperature slowly. After 16 hours, LCMS confirmed the formation of the product. The reaction was quenched with saturated NH₄Cl and diluted with methyl *tert*-Butyl ether (MTBE) and water. The aqueous phase was extracted twice with MTBE. The combined organic layers were dried over MgSO₄, filtered, and concentrated, and the crude product was purified by column chromatography to give the desired product. (MS (ES) M+H: 706).

[00957] 2-{3-*tert*-Butylsulfanyl-5-(pyridin-2-ylmethylsulfanyl)-1-[4-(5-trifluoromethyl-pyridin-2-yl)-benzyl]-1*H*-indol-2-ylmethyl}-2-ethyl-butyric acid methyl ester (MS (ES) M+H: 706) was prepared starting from 2-[3-*tert*-Butylsulfanyl-5-(pyridin-2-ylmethylsulfanyl)-1*H*-indol-2-ylmethyl]-2-ethyl-butyric acid methyl ester.

Hydrolysis of methyl ester

[00958] 2-{3-*tert*-Butylsulfanyl-5-(5-methyl-pyridin-2-ylsulfanyl)-1-[4-(5-trifluoromethyl-pyridin-2-yl)-benzyl]-1*H*-indol-2-ylmethyl}-2-ethyl-butyric acid methyl ester (25mg, 0.035mmol) was dissolved in dioxane 2ml and stirred until it became a clear solution. 1N LiOH aqueous solution (2mL, 2mmol) was added and the reaction was refluxed at 95°C for 24 hours. LCMS confirmed the formation of the product, so the reaction was cooled to room temperature and partitioned between EtOAc and water. The pH of the aqueous solution was adjusted to pH 1 with 10% HCl, and the aqueous phase was extracted three times with EtOAc. The combined organic layers were washed with water, dried over MgSO₄, filtered, and concentrated to give the desired free acid. (MS (ES) M+H: 692).

[00959] 2-{3-*tert*-Butylsulfanyl-5-(pyridin-2-ylmethylsulfanyl)-1-[4-(5-trifluoromethyl-pyridin-2-yl)-benzyl]-1*H*-indol-2-ylmethyl}-2-ethyl-butyric acid. (MS (ES) M+H: 692) was prepared by hydrolysis of 2-{3-*tert*-Butylsulfanyl-5-(pyridin-2-ylmethylsulfanyl)-1-[4-(5-trifluoromethyl-pyridin-2-yl)-benzyl]-1*H*-indol-2-ylmethyl}-2-ethyl-butyric acid methyl ester.

Scheme B:**Example 2.**

[00960] Scheme B illustrates a non-limiting example of the synthesis of indole compounds. The synthetic methodology can be adapted to produce compounds of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), Formula (H).

Step 1: 4-*tert*-Butylsulfanyl-3-oxo-butyric acid ethyl ester

- 5 [00961] Ethyl 4-chloroacetoacetate (7.5 mL, 51.9 mmol), 2-methyl-2-propanethiol (5.6 mL, 49.7 mmol), triethylamine (10.8 mL, 77.4 mmol), and catalytic tetrabutylammonium bromide were dissolved in THF (250 mL) and stirred at room temperature overnight. Silica gel was added, and the mixture was concentrated and filtered over a plug of silica gel to obtain the desired product (**B-1**), which was used without further purification.


Step 2: (3-*tert*-Butylsulfanyl-5-methoxy-1*H*-indol-2-yl)-acetic acid ethyl ester

- 10 [00962] 4-Methoxyphenylhydrazine hydrochloride (7.7 g, 44.1 mmol) and **B-1** (7.4 g, 33.9 mmol) were dissolved in 2-propanol (150 mL) and heated to reflux for 24 hours. The reaction mixture was concentrated and partitioned between EtOAc and saturated aq. NaHCO₃. The aqueous layer was extracted with EtOAc, and the combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified on silica gel (0 to 30% EtOAc in hexanes) to give the desired product (**B-2**).

15 **Step 3: (3-*tert*-Butylsulfanyl-5-hydroxy-1*H*-indol-2-yl)-acetic acid ethyl ester**

- [00963] Aluminum chloride (7.5 g 56.0 mmol) was suspended in *tert*-Butylthiol (21 mL, 186.7 mmol) at 0°C. **B-2** (6.0 g, 18.7 mmol) was added in CH₂Cl₂ (21 mL), and the reaction was allowed to warm to room temperature. After 2 hours, the reaction was complete by TLC analysis, so the solution poured into ice and acidified with 10% HCl aqueous solution. The aqueous layer was extracted three times with EtOAc, the combined organics were
20 dried over MgSO₄, filtered, and concentrated to give the desired product (**B-3**). Hydroxy indoles of structure **B-3** may be transformed to the corresponding thiol indoles, which may then be S-alkylated or used in metal mediated cross-coupling reaction with aryl halides or heteroaryl halides

Scheme C:

**Example 3.**

- 5 Indole compounds described herein may also be prepared as described in Scheme C.

Step 1:

- [00964] A solution of 4-Methoxyphenylhydrazine hydrochloride (10.0g, 57.3mmol), 4-chlorobenzylchloride (9.2g, 57.2mmol), tetrabutylammonium bromide (3.7g, 11.5mmol), and diisopropylethylamine (20mL, 115mmol) in CH₂Cl₂ (250mL) was stirred at room temperature for several days. The reaction mixture was diluted with water and the organic layer was dried over MgSO₄, filtered, and concentrated. The residue was taken up in toluene (200mL) and diethyl ether (100mL), and 1 equivalent of 4N HCl in dioxane was added at 0°C. The mixture was stirred at room temperature for 2 hours, and then evaporated to dryness to give the desired product (C-1; X=Cl) as a purple solid.

Step 2:


- 15 [00965] C-1 (~16g, 57.3mmol), ethyl 5-(tert-butylthio)-2,2-dimethyl-4-oxopentanoate (prepared according to the procedures described in US patent 5,288,743 issued Feb 22, 1994; 14.8g, 57.3mmol), NaOAc (5.2g) in toluene (120mL) and HOAc (66mL) was stirred at room temperature in the dark for 5 days. The mixture was partitioned between EtOAc and water, and the organic layer was stirred with solid NaHCO₃, filtered, and evaporated. The residue was purified on silica gel (0 to 55% CH₂Cl₂ in hexanes), and the isolated product was recrystallized from hexanes to give the desired product (C-2; X=Cl).
- 20

Step 3:

[00966] Aluminum chloride (0.820g 6.15mmol) was suspended in *tert*-Butylthiol (1.8mL, 16mmol) and cooled to 0°C. C-2 (1.0g, 2.0mmol) was added in CH₂Cl₂ (2.4mL), and the reaction was allowed to warm to room temperature. After 3 hours, the reaction was complete by TLC analysis, so the solution was diluted with CH₂Cl₂ and washed with 10% ice-cooled HCl aqueous solution. The aqueous layer was extracted three times with CH₂Cl₂, the combined organics were dried over MgSO₄, filtered, and concentrated to give the desired product (C-3; X=Cl) as a colourless foam.

Step 4:

[00967] Derivatization of hydroxy indoles of structure C-3 as an *N,N*-dimethylthiocarbamoyl indoles (structure C-4) is followed by thermal rearrangement at >200 C provides the *N,N*-dimethylcarbamoylethioindole. The hydrolysis of *N,N*-dimethylcarbamoylethioindoles may be carried out using basic conditions. Spontaneous formation of the disulfide may occur, in which case reduction of the disulfide is carried out using triphenylphosphine in aqueous dioxane, or similar reaction conditions, to provide the thiol indoles of structure C-5. Thiol indoles of structure C-5 may be S-alkylated or used in metal mediated cross-coupling reaction with aryl halides or heteroaryl halides.

Scheme D:**Example 4.**

[00968] Substituents at the C-3 position of indole skeleton of compounds described herein may be introduced, but not limited to, by the methods described in Scheme D. Treatment of indoles of structure D-1 with an acid chloride in the presence of a lewis acid, such as, but not limited to, AlCl₃, provides acylated indoles of structure D-2. Reduction of the carbonyl with a reducing agent, such as, but not limited to, sodium borohydride, provides alkyl indoles of structure D-3. If R₇ in indoles of structure D-3 have an alkyl ester group (G₁ is CO₂R₉), then treatment with a base, such as lithium hydroxide, provides hydrolyzed product (G₁ is CO₂H).

Step 1:

[00969] To D-1 (0.17mmol) in dichloroethane (5mL) is added cyclobutanecarbonyl chloride (0.50mmol) and aluminum chloride (0.66mmol). The reaction is heated under N₂ for 1.5 hours, and then cooled to room temperature and quenched with saturated aq. potassium sodium tartrate. The mixture is extracted with EtOAc, and the combined organic layers were dried over MgSO₄, filtered, concentrated, and purified on silica gel to give the desired product (D-2).

Step 2:

[00970] D-2 (0.08mmol) is suspended in CH₂Cl₂, and sodium borohydride (0.8mmol) is added dropwise in TFA (1mL) and CH₂Cl₂ (1mL). The mixture is stirred at room temperature for 4 hours, and then quenched with water and basified with solid NaOH pellets. The mixture is extracted with CH₂Cl₂, and the combined organics were


dried over MgSO_4 , filtered, and concentrated. The residue is purified on silica gel to give the desired product (D-3).

Compounds

[00971] Non-limiting examples of compounds that may be prepared by the methods described herein, as well as

5 by methods known in the art, include those in Tables 1-14 and Figures 8-11:


Table 1. R_{11} (aryl-heteroaryl) indoles with acid replacements



Compound #	Y	-G ₆	G ₁
1-1	Pyridin-2-yl	Pyridin-2-yl	C(O)NH ₂
1-2	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	CO ₂ Et
1-3	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	C(O)(CH ₂) ₆ OH
1-4	Pyridin-2-yl	Pyridin-2-yl	OH
1-5	Pyridin-2-yl	Pyridin-3-yl	OH
1-6	Pyridin-2-yl	Thiazol-2-yl	OH
1-7	Pyridin-2-yl	1,3,4-Oxadiazol-2-yl	OH
1-8	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	1,3,4-Oxadiazol-2-yl
1-9	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	1,3,4-Oxadiazol-2-ylamine
1-10	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	-C(O)NH-(Pyrazin-2-yl)
1-11	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	-C(O)NH-(Thiazol-2-yl)
1-12	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	-C(O)NH-(Pyridin-3-yl)
1-13	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	C(O)NH(CH ₂ CH ₂ NMe ₂)
1-14	Pyridin-2-yl	1,3,4-Oxadiazol-2-ylamine	CH ₃
1-15	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	C(O)NHC(=NH)NH ₂
1-16	5-Cyano-pyridin-2-yl	6-Methoxy-pyridin-3-yl	CO ₂ Et
1-17	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	C(O)N=C(NH ₂) ₂
1-18	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	1,3,4-Oxadiazol-2-ylamine
1-19	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	NHBOC
1-20	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	NH ₂
1-21	Quinolin-2-yl	5-Fluoro-pyridin-2-yl	C(O)NHSO ₂ Me
1-22	Pyridin-2-yl	6-Methoxy-pyridin-3-yl	C(O)N=C(NH ₂) ₂

Also described herein are compounds in which the -CH₂S- group between the Y and the indolyl groups is replaced with an -S- group.

10 **Table 2. R_{11} (aryl-heteroaryl/heterocycloalkyl) indoles**



Compound #	Y	position	-G ₆	R ₆
2-1	Pyridin-2-yl	4	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-2	Pyridin-2-yl	4	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
2-3	Pyridin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-4	Pyridin-2-yl	4	Pyrimidin-5-yl	<i>tert</i> -Butylsulfanyl
2-5	Pyridin-2-yl	4	Pyrazin-2-yl	<i>tert</i> -Butylsulfanyl
2-6	Pyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-7	Pyridin-2-yl	4	5-Amino-pyrazin-2-yl	<i>tert</i> -Butylsulfanyl
2-8	Pyridin-2-yl	4	Thiazol-2-yl	3,3-Dimethyl-butyl
2-9	Pyridin-2-yl	4	Thiazol-2-yl	H
2-10	Pyridin-2-yl	4	Thiazol-2-yl	Acetyl
2-11	Pyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	H
2-12	Pyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	Acetyl
2-13	Pyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	Ethyl
2-14	Pyridin-2-yl	4	Thiazol-2-yl	3,3-Dimethyl-butyl
2-15	Pyridin-2-yl	4	Thiazol-2-yl	Cyclopropane-carbonyl
2-16	Pyridin-2-yl	4	Thiazol-2-yl	Cyclobutane-carbonyl
2-17	Pyridin-2-yl	4	6-Hydroxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-18	Pyridin-2-yl	4	Pyridin-4-yl	<i>tert</i> -Butylsulfanyl
2-19	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-20	Pyridin-2-yl	4	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-21	Pyridin-2-yl	4	5-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-22	Pyridin-2-yl	4	Thiazol-2-yl	Cyclobutylmethyl
2-23	2-Methylthiazol-4-ylmethyl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-24	2-Methylthiazol-4-yl	4	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-25	2-Methylthiazol-4-yl	4	Thiazol-2-yl	H
2-26	2-Methylthiazol-4-yl	4	Thiazol-2-yl	3,3-Dimethyl-butyl
2-27	2-Methylthiazol-4-yl	4	6-Methoxy-pyridazin-3-yl	H
2-28	2-Methylthiazol-4-yl	4	6-Methoxy-pyridazin-3-yl	3,3-Dimethyl-Butyl
2-29	Pyridin-2-yl	4	Thiazol-2-yl	Ethyl
2-30	Benzothiazol-2-ylmethyl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-31	2-Methylthiazol-4-yl	4	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
2-32	Benzothiazol-2-yl	4	Pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
2-33	Pyridin-2-yl	4	2-Methyl-3-pyridin-2-ylmethyl-3H-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
2-34	Pyridin-2-yl	4	2,4-Dimethyl-thiazol-5-yl	<i>tert</i> -Butylsulfanyl
2-35	Pyridin-2-yl	4	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-36	Pyridin-2-yl	4	5-Trifluoromethyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-37	Pyridin-2-ylm	4	2-Methyl-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-38	Pyridin-2-yl	4	2-Methyl-thiazol-5-yl	<i>tert</i> -Butylsulfanyl
2-39	Pyridin-2-yl	4	4-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-40	Pyridin-2-yl	4	Isoxazol-4-yl	<i>tert</i> -Butylsulfanyl
2-41	Pyridin-2-yl	4	3,5-Dimethyl-isoxazol-4-yl	<i>tert</i> -Butylsulfanyl
2-42	Pyridin-2-yl	4	2-Methyl-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
2-43	Pyridin-2-yl	4	1-Methyl-imidazol-5-yl	<i>tert</i> -Butylsulfanyl
2-44	Pyridin-2-yl	4	1-Methyl-imidazol-4-yl	<i>tert</i> -Butylsulfanyl
2-45	Pyridin-2-yl	4	Imidazol-4-yl	<i>tert</i> -Butylsulfanyl
2-46	Pyridin-2-yl	4	4-Methyl-imidazol-5-yl	<i>tert</i> -Butylsulfanyl
2-47	Pyridin-2-yl	4	5-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-48	Pyridin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-49	Pyridin-2-yl	4	Pyrazol-4-yl	<i>tert</i> -Butylsulfanyl

Compound #	Y	position	-G ₆	R ₆
2-50	Pyridin-2-yl	4	1-Methyl-pyrazol-4-yl	<i>tert</i> -Butylsulfanyl
2-51	Pyridin-2-yl	4	3-Methyl-pyrazol-4-yl	<i>tert</i> -Butylsulfanyl
2-52	Pyridin-2-yl	4	5-Methyl-1,2,4-oxadiazol-3-yl	<i>tert</i> -Butylsulfanyl
2-53	Pyridin-2-yl	4	2-Methyl-1,3,4-oxadiazol-5-yl	<i>tert</i> -Butylsulfanyl
2-54	Pyridin-2-yl	4	1,3,4-Oxadiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-55	Pyridin-2-yl	4	1,3,4-Thiadiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-56	Pyridin-2-yl	4	3-Methyl-pyrazol-5-yl	<i>tert</i> -Butylsulfanyl
2-57	Pyridin-2-yl	4	1,2,3-Thiadiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-58	Pyridin-2-yl	4	Tetrazol-1-yl	<i>tert</i> -Butylsulfanyl
2-59	Pyridin-2-yl	4	Tetrazol-2-yl	<i>tert</i> -Butylsulfanyl
2-60	Pyridin-2-yl	4	1-Methyl-tetrazol-5-yl	<i>tert</i> -Butylsulfanyl
2-61	Pyridin-2-yl	4	2-Methyl-tetrazol-5-yl	<i>tert</i> -Butylsulfanyl
2-62	Pyridin-2-yl	4	6-Hydroxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-63	Pyridin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-64	Pyridin-2-yl	4	6-Cyano-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-65	Pyridin-2-yl	4	6-Trifluoromethyl-pyridin-4-yl	<i>tert</i> -Butylsulfanyl
2-66	Pyridin-2-yl	4	2-Acetylamino-pyridin-5-yl	<i>tert</i> -Butylsulfanyl
2-67	Pyridin-2-yl	4	2-Methoxy-pyrimidin-5-yl	<i>tert</i> -Butylsulfanyl
2-68	Pyridin-2-yl	4	2-Methoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-69	3-Fluoro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-70	3-Fluoro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-71	4-Fluoro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-72	5-Fluoro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-73	5-Methyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-74	5-Cyano-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-75	5-Methoxy-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-76	5-Ethyl-pyridin-2-yl	4	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-77	Quinolin-2-yl	4	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-78	6-Fluoroquinolin-2-yl	4	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-79	Quinolin-2-yl	3	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-80	Quinolin-2-yl	3	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-81	Quinolin-2-yl	3	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-82	5-Methyl-pyridin-2-yl	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-83	Quinolin-2-yl	3	2-Trifluoromethyl-pyridin-5-yl	<i>tert</i> -Butylsulfanyl
2-84	5-Ethyl-pyridin-2-yl	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-85	Quinolin-2-yl	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-86	Quinolin-2-yl	3	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-87	Pyridin-2-yl	4	5-Carbamoyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-88	Pyridin-2-yl	4	5-Cyano-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-89	Pyridin-2-yl	4	5-Methoxy-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-90	Pyridin-2-yl	4	6-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-91	Pyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-92	Pyridin-2-yl	4	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-93	Pyridin-2-yl	4	4-Methyl-1 <i>H</i> -imidazol-2-yl	<i>tert</i> -Butylsulfanyl
2-94	Pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-95	Pyridin-2-yl	4	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-96	Pyridin-2-yl	4	5-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-97	Pyridin-2-yl	4	6-Carbamoyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-98	Pyridin-2-yl	4	5-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-99	6-Fluoro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-100	6-Methoxy-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl

Compound #	Y	position	-G ₆	R ₆
2-101	6-Methyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-102	5-Methyl-pyridin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-103	5-Methyl-pyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-104	6-Cyclopropyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-105	5-Methyl-pyridin-2-yl	4	5-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-106	5-Methyl-pyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-107	5-Methyl-pyridin-2-ylmethoxy	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-108	5-Chloro-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-109	Pyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-110	Pyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-111	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-112	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-113	Pyridin-2-yl	4	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-114	Pyridin-2-yl	4	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-115	Pyridin-2-yl	4	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-116	Pyridin-2-yl	4	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-117	3-Methyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-118	3-Methyl-pyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-119	3,5-Dimethylpyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-120	3,5-Dimethylpyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-121	Benzothiazol-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-122	Benzothiazol-2-yl	4	5-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-123	Benzothiazol-2-yl	4	6-Methoxy-pyridin-3-yl	Cyclobutyl-carbonyl
2-124	Benzothiazol-2-yl	4	6-Methoxy-pyridin-3-y	Cyclobutylmethyl
2-125	5-Ethylpyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-126	5-Ethylpyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-127	5-Ethylpyridin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-128	5-Ethylpyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-129	5-Methylpyridin-2-yl	4	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-130	5-Methylpyridin-2-yl	4	2-Methoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-131	5-Methylpyridin-2-yl	4	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-132	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	Cyclobutylmethyl
2-133	5-Methylpyridin-2-yl	4	6-Methoxy-pyridin-3-yl	Cyclobutylmethyl
2-134	5-Methylpyridin-2-yl	4	6-Methoxy-pyridin-3-yl	Isobutyl
2-135	Quinolin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-136	Quinolin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-137	Quinolin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-138	Quinolin-2-yl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-139	Quinolin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-140	6-Fluoroquinolin-2-yl	4	6-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-141	6-Fluoroquinolin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-142	6-Fluoroquinolin-2-yl	4	2-Ethoxy-thiazol-4-yl	<i>tert</i> -Butylsulfanyl
2-143	6-Fluoroquinolin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl

Compound #	Y	position	-G ₆	R ₆
2-144	7-Fluoroquinolin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-145	7-Fluoroquinolin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-146	7-Fluoroquinolin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-147	7-Fluoroquinolin-2-y	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-148	6-Fluoroquinolin-2-yl	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-149	5-Methyl-pyridin-2-yl	4	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-150	5-Ethyl-pyridin-2-yl	4	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-151	Quinolin-2-yl	4	3-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-152	Quinolin-2-yl	3	5-Methoxy-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-153	Quinolin-2-yl	3	3-Methoxy-pyridazin-6-yl	<i>tert</i> -Butylsulfanyl
2-154	Quinolin-2-yl	3	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-155	Quinolin-2-yl	3	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-156	6-Fluoroquinolin-2-yl	4	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-157	3-Methylpyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-158	3-Methylpyridin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-159	3,5-Dimethylpyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-160	4-Methylpyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-161	4-Methylpyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-162	4-Methylpyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-163	5-Methylpyridin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	Cyclobutylmethyl
2-164	6-Fluoroquinolin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-165	6-Fluoroquinolin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-166	6-Methylquinolin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-167	6-Methylquinolin-2-yl	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-168	Quinolin-2-yl	4	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-169	Quinolin-2-yl	4	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-170	Quinolin-2-yl	4	6-Methoxy-pyridin-3-yl	Isobutyl
2-171	6-Fluoroquinolin-2-yl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-172	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	2-Methyl-propane-2-sulfonyl
2-173	Pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	2-Methyl-propane-2-sulfinyl
2-174	N-Oxido-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-175	Imidazo[1,2- <i>a</i>]pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-176	Imidazo[1,2- <i>a</i>]pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-177	Imidazo[1,2- <i>a</i>]pyridin-2-yl	4	5-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-178	Pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-179	6-Fluoroquinolin-2-yl	4	6-Methyl-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-180	5-Methylisoxazol-3-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-181	5-Methylisoxazol-3-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-182	5-Methylisoxazol-3-yl	4	5-Trifluoromethylpyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-183	1,3-Dimethylpyrazol-5-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-184	1,5-Dimethylpyrazol-3-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl


Compound #	Y	position	-G ₆	R ₆
2-185	6-Fluoroquinolin-2-yl	4	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-186	5-Ethylpyridin-2-yl	4	6-Ethoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-187	5-Ethylpyridin-2-yl	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
2-188	6-Fluoroquinolin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-189	Pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-190	6-Fluoroquinolin-2-yl	4	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-191	Pyridin-2-yl	4	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-192	5-Methylpyridin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-193	5-Methylpyridin-2-yl	4	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-194	6-Fluoroquinolin-2-yl	4	6-Trifluoromethyl-pyridin-3-yl	Isobutyl
2-195	Pyridin-2-yl	3	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-196	Pyridin-2-yl	3	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-197	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-198	Quinolin-2-yl	4	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-199	Pyridin-2-yl	4	6-Ethoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-200	6-Fluoroquinolin-2-yl	4	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-201	Pyridine-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-202	5-Methylpyridin-2-yl	4	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-203	Quinolin-2-yl	4	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-204	Pyridin-2-yl	4	6-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-205	Quinolin-2-yl	4	Thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-206	Pyridin-2-yl	3	4-Methoxy-tetrahydro-pyran-4-yl	<i>tert</i> -Butylsulfanyl
2-207	6-Fluoroquinolin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-208	5-Ethylpyridin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-209	Quinolin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-210	6-Fluoroquinolin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-211	5-Methylpyridin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-212	5-Ethylpyridin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-213	Quinolin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-214	5-Methylpyridin-2-yl	4	Pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-215	5-Methylpyridin-2-yl	4	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-216	Quinolin-2-yl	4	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-217	5-Methylpyridin-2-yl	4	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-218	5-Ethylpyridin-2-yl	4	3-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-219	5-Methylpyridin-2-yl	4	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-220	5-Ethylpyridin-2-yl	4	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-221	Quinolin-2-yl	4	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-222	5-Methylpyridin-2-yl	4	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-223	5-Ethylpyridin-2-yl	4	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-224	Quinolin-2-yl	4	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-225	5,6-Dimethyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-226	5,6-Dimethyl-pyridin-2-yl	4	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-227	5,6-Dimethyl-pyridin-2-yl	4	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-228	5,6-Dimethyl-pyridin-	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl

Compound #	Y	position	-G ₆	R ₆
	2-yl			
2-229	5,6-Dimethyl-pyridin-2-yl	4	5-Fluoro-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-230	5,6-Dimethyl-pyridin-2-yl	4	4-Methoxy-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-231	5,6-Dimethyl-pyridin-2-yl	4	Pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-232	5-Methylpyridin-2-yl	4	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-233	5-Ethylpyridin-2-yl	4	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-234	Quinolin-2-yl	4	2-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-235	5-Bromo-pyridin-2-yl	4	5-Bromo-6-methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-236	6-Bromo-quinolin-2-yl	4	5-Bromo-6-methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-237	5-Methyl-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	2-Methyl-propane-2-sulfinyl
2-238	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	2-Methyl-propane-2-sulfinyl
2-239	5,6-Dimethyl-pyridin-2-yl	4	(5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-240	5,6-Dimethyl-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-241	Quinolin-2-yl	4	5-Methyl-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-242	Quinolin-2-yl	3	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-243	Quinolin-2-yl	3	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-244	5-Carbamoyl-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-245	5-Methoxy-pyridin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-246	1 <i>H</i> -Indol-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-247	Quinolin-2-yl	4	5-Fluoro-thiazol-2-yl	<i>tert</i> -Butylsulfanyl
2-248	Quinolin-2-yl	4	5-Fluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-249	Quinolin-2-yl	4	5-Methoxymethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-250	Quinolin-2-yl	4	6-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-251	Quinolin-2-yl	4	5-Hydroxymethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-252	Quinolin-2-yl	4	4-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-253	Quinolin-2-yl	4	2-Methyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-254	Quinolin-2-yl	4	3-Methyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-255	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	H
2-256	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butyl
2-257	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butaryl
2-258	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	2,2-Dimethyl-propionyl
2-259	5-Methyl-1-oxy-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-260	1-Oxy-quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-261	5-Methyl-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	H
2-262	5-Methyl-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	3,3-Dimethyl-butaryl
2-263	5-Methyl-pyridin-2-yl	4	6-Ethoxy-pyridin-3-yl	Phenylacetyl
2-264	5,6-Dimethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	H
2-265	5-Ethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	H
2-266	Quinolin-2-yl	4	5-Fluoro-pyridin-2-yl	3-Methyl-butaryl
2-267	5-Ethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	3-Methyl-butaryl

Compound #	Y	position	-G ₆	R ₆
2-268	5-Ethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-buteryl
2-269	5-Ethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	2-Ethyl-buteryl
2-270	5,6-Dimethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	3-Methyl-buteryl
2-271	5,6-Dimethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-buteryl
2-272	5,6-Dimethyl-pyridin-2-yl	4	5-Fluoro-pyridin-2-yl	2-Ethyl-buteryl
2-273	5-Methyl-pyrazin-2-yl	4	3-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-274	5-Methyl-pyrazin-2-yl	4	4-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-275	5-Methyl-pyrazin-2-yl	4	3-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-276	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
2-277	5-Methyl-pyrazin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-278	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	Isobuteryl
2-279	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-buteryl
2-280	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	Propionyl
2-281	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	Acetyl
2-282	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	3-Methyl-buteryl
2-283	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	2,2,2-Trifluoro-acetyl
2-284	Quinoxalin-2-yl	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
2-285	5-Methyl-pyrazin-2-yl	4	5-Fluoro-pyridin-2-yl	3,3-Dimethyl-butyl
2-286	Quinoxalin-2-yl	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl

where Z is -S-, CH(CH₃)S-, or -CH₂S-.


Table 3. R₁₁ (heteroaryl-aryl) and (heteroaryl-heteroaryl) indoles



Compound #	R ₁₁
3-1	2-(2-Methoxy pyrid-5-yl)-pyrid-5-yl
3-2	2-(4-Methoxy phenyl)-pyrid-5-yl
3-3	2-(4-Trifluoromethoxy phenyl)-pyrid-5-yl
3-4	5-(4-Methoxy phenyl)-pyrid-2-yl
3-5	5-(4-Trifluoromethoxyphenyl)-pyrid-2-yl

Also described herein are compounds in which the -CH₂S- group between the pyridinyl and indolyl groups is replaced with an -S- group.

Table 4. R₁₁ (aryl-heteroaryl/heterocycloalkyl) indoles




Compound #	Y	Z	position	-G ₆	R ₆
------------	---	---	----------	-----------------	----------------

Compound #	Y	Z	position	-G ₆	R ₆
4-1	Pyridin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-2	Quinolin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-3	Quinoxalin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-4	Quinolin-2-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-5	Quinoxalin-2-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-6	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-7	Pyridin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-8	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-9	5-Methyl-pyridin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-10	Pyridin-2-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-11	5-Methyl-pyridin-2-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-12	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-13	Pyridin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
4-14	Quinolin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
4-15	5-Methyl-pyridin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
4-16	Quinoxalin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
4-17	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	6-Methoxy-pyridazin-3-yl	<i>tert</i> -Butylsulfanyl
4-18	Quinolin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-19	5-Methyl-pyridin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-20	Quinoxalin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-21	3-Oxo-3,4-dihydro-quinoxalin-2-ylmethoxy	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-22	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	3,3-Dimethyl-butyryl
4-23	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	5-Trifluoromethyl-pyridin-2-yl	Cyclobutyl-carbonyl
4-24	6-Methyl-pyridazin-3-yl	-CH ₂ S-	4	5-Fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-25	Quinolin-2-yl	-CH ₂ S-	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-26	Pyridin-2-yl	-CH ₂ S-	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-27	5-Methyl-pyridin-2-ylmethoxy	-CH ₂ S-	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-28	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	6-Trifluoromethyl-pyridin-3-yl	<i>tert</i> -Butylsulfanyl
4-29	Pyridin-2-yl	-CH ₂ S-	4	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-30	5-Methyl-pyridin-2-yl	-CH ₂ S-	4	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-31	5-Methyl-pyrazin-2-yl	-CH ₂ S-	4	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-32	Quinolin-2-yl	-CH ₂ S-	4	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-33	Quinoxalin-2-yl	-CH ₂ S-	4	5-Hydroxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-34	Pyridin-2-yl	-CH ₂ S-	4	5-trifluoromethyl-pyridin-	<i>tert</i> -Butylsulfanyl

Compound #	Y	Z	position	-G ₆	R ₆
				2-yl	
4-35	Pyrazin-2-yl	-CH ₂ S-	4	5-fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-36	pyrimidin-2-yl	-CH ₂ S-	4	5-fluoro-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-37	Pyridin-2-yl	-CH ₂ S-	4	5-methoxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-38	5-methyl-pyridin-2-yl	-CH ₂ S-	4	5-methoxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-39	5-methyl-pyrazin-2-yl	-CH ₂ S-	4	5-methoxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-40	pyrazin-2-yl	-CH ₂ S-	4	5-methoxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-41	Pyrimidin-2-yl	-CH ₂ S-	4	5-methoxy-pyrimidin-2-yl	<i>tert</i> -Butylsulfanyl
4-42	Pyridin-2-yl	-SCH ₂ -	4	5-trifluoromethyl-pyridin-2-yl	<i>tert</i> -Butylsulfanyl
4-43	5-methyl-pyridin-2-yl	-CH ₂ S-	4	Morpholin-4-yl	<i>tert</i> -Butylsulfanyl

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Table 5. R₁₁ (aryl-heteroaryl) indoles




Compound #	Y	Z	-G ₆	R ₇
5-1	5-Methyl-pyridin-2-yl	-CH ₂ S-	6-Methoxy-pyridin-3-yl	CH ₂ CH ₂ CO ₂ H
5-2	5-Methyl-pyridin-2-yl	-CH ₂ S-	6-Methoxy-pyridin-3-yl	Me
5-3	5-Methyl-pyrazin-2-yl	-CH ₂ S-	6-Methoxy-pyridin-3-yl	Me
5-4	5-Methyl-pyridin-2-yl	-CH ₂ S-	5-Fluoro-pyridin-2-yl	Me
5-5	5-Methyl-pyrazin-2-yl	-CH ₂ S-	5-Fluoro-pyridin-2-yl	Me
5-6	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-Trifluoromethyl-pyridin-2-yl	
5-7	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-Trifluoromethyl-pyridin-2-yl	
5-8	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-9	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-10	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-11	pyridin-2-yl	-CH ₂ S-	6-methoxy-pyridin-3-yl	

Compound #	Y	Z	-G ₆	R ₇
5-12	5-methyl-pyridin-2-yl	-CH ₂ S-	6-ethoxy-pyridin-3-yl	
5-13	5-methyl-pyrazin-2-yl	-CH ₂ S-	6-trifluoromethyl-pyridin-3-yl	
5-14	5-methyl-pyrazin-2-yl	-CH ₂ S-	6-trifluoromethyl-pyridin-3-yl	
5-15	5-methyl-pyridin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-16	quinolin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-17	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-18	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-19	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-20	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	
5-21	5-methyl-pyrazin-2-yl	-CH ₂ S-	5-trifluoromethyl-pyridin-2-yl	

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Table 6. Heterocycloalkyl Y substituents




Compound #	Y	Z	G ₆	R ₆	R ₉
6-1	(S)-N-t-Butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-2	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-3	(R)-N-t-Butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-4	(S)-2-Pyrrolidon-5-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-5	(R)-2-Pyrrolidon-5-ylmethyl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H

Compound #	Y	Z	G ₆	R ₆	R ₉
6-6	(R)-N-Acetyl-pyrrolidin-2-ylmethyl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-7	(R)-N-Methylsulfonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-8	(S)-N-Methylsulfonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-9	(R)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-10	N-Trifluoroacetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-11	N-t-Butoxycarbonyl-4,5-dihydroimidazol-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-12	4,5-Dihydroimidazol-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-13	(S)-N-t-Butoxycarbonyl indolin-2-ylmethyl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-14	Morpholin-4-yl	C(O)CH ₂ S	Cl	<i>tert</i> -butylsulfanyl	H
6-15	(S)-Indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-16	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-17	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio S,S-dioxide	H
6-18	(S)-N-Cyclopropylcarbonyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-19	(S)-N-Benzoyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-20	(S)-N-(2-Methylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-21	(S)-N-Propanoyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-22	N-t-Butoxycarbonyl indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-23	Indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-24	N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-25	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methyl-2-propylthio-S-oxide	H
6-26	(S)-N-Acetyl-indolin-2-ylmethyl	-CH ₂ S-	Cl	benzyl	H
6-27	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	H	H
6-28	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	H	H
6-29	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbutanoyl	H
6-30	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbutanoyl	H
6-31	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	ethyl	H
6-32	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	propyl	H
6-33	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methylpropanoyl	H
6-34	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	Cyclopropyl-carbonyl	H
6-35	(S)-N-Acetyl-indolin-2-ylmethyl	-CH ₂ S-	Cl	benzoyl	H
6-36	(S)-N-Acetyl-indolin-2-ylmethyl	-CH ₂ S-	Cl	Cyclobutyl-carbonyl	H
6-37	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	acetyl	H
6-38	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	propanoyl	H
6-39	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	2-methylpropyl	H
6-40	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	3,3-dimethylbut-1-yl	H
6-41	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	Cl	cyclobutylmethyl	H
6-42	(S)-N-(4-Phenylbenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-43	(S)-N-(Phenylacetyl)-pyrrolidin-2-ylmethyl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H

Compound #	Y	Z	G ₆	R ₆	R ₉
6-44	(S)-N-(3-Phenylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-45	(S)-N-(3-Phenoxybenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-46	(S)-N-(4-Phenoxybenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-47	(S)-N-(Nicotinoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-48	(S)-N-(Pyridin-4-ylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-49	(S)-N-(4-Phenylbenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-50	(S)-N-(Phenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-51	(S)-N-(3-Phenylpropanoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-52	(S)-N-(Phenylcyclopropylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-53	(S)-N-(Nicotinoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-54	(S)-N-(Pyridin-4-ylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-55	(S)-N-(Phenylcyclopropylcarbonyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-56	(S)-N-(4-Bhlorobenzoyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-57	(S)-N-(4-Benzoyloxyphenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-58	(S)-N-(4-Benzoyloxyphenylacetyl)-pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-59	N-(<i>tert</i> Butoxycarbonyl)piperidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-60	N-(<i>tert</i> Butoxycarbonyl)piperidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-61	(S)-N-(2-Bromoethoxycarbonyl)indolin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	Et
6-62	(S)-Pyrrolidin-2-yl	-CH ₂ S-	Cl	<i>tert</i> -butylsulfanyl	H
6-63	2-(2-Methyl-1,3-dioxolan-2-yl)	-CH ₂ S-	Br	<i>tert</i> -butylsulfanyl	H

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Table 7. Heterocycloalkyl Y substituents




Compound #	Y	Z	G ₆	R ₆	R ₇
7-1	(S)-N-t-Butoxycarbonyl-	-CH ₂ S-	2-thiazolyl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H

Compound #	Y	Z	G ₆	R ₆	R ₇
	pyrrolidin-2-yl				
7-2	(S)-Pyrrolidin-2-yl	-CH ₂ S-	2-thiazolyl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-3	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	2-thiazolyl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-4	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	2-thiazolyl	H	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-5	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxy-4-pyridazinyl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-6	(S)-N-Acetyl-pyrrolidin-2-yl	-CH ₂ S-	2-methoxy-4-pyridazinyl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-7	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxypyridin-5-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-8	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	2-methoxythiazol-4-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-9	(S)-N-Acetyl-indolin-2-yl	-CH ₂ S-	5-methoxypyridin-2-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-10	2-Methyl-1,3-dioxolan-2-yl	-CH ₂ CH ₂ S-	2-methoxypyridin-5-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-11	N-(Methoxyacetyl)indolin-2-yl	-CH ₂ S-	5-trifluoromethylpyridin-2-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
7-12	N-Acetyl-indolin-2-yl	-CH ₂ S-	5-trifluoromethylpyridin-2-yl	<i>tert</i> -butylsulfanyl	-CH ₂ C(CH ₂ CH ₃) ₂ CO ₂ H

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Table 8. Heterocycloalkyl Y substituents with tertiary alcohol R₇




Compound #	Y	Z	R ₁₁	R ₇
8-1	Morpholin-4-yl	-C(=O)CH ₂ S-	4-chloro-phenyl	2-hydroxy-2-methyprop-1-yl
8-2	N-t-butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	4-chloro-phenyl	2-hydroxy-2-methyprop-1-yl
8-3	N-t-butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	pyridin-2-yl	2-hydroxy-2-methyprop-1-yl
8-4	N-acetyl-pyrrolidin-2-yl	-CH ₂ S-	4-chlorophenyl	2-hydroxy-2-methyprop-1-yl
8-5	N-acetyl-pyrrolidin-2-yl	-CH ₂ S-	pyridin-2-yl	2-hydroxy-2-methyprop-1-yl

Compound #	Y	Z	R ₁₁	R ₇
8-6	(S)-N-t-butoxycarbonyl-pyrrolidin-2-yl	-CH ₂ S-	4-chlorophenyl	1-hydroxy-2,2-dimethylprop-3-yl

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Table 9. Non-aromatic R substituents




Compound #	R	G ₆	R ₇
9-1	2-acetamide	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-2	(S)-2-t-Butoxycarbonylamino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-3	(R)-2-t-Butoxycarbonylamino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-4	(S)-2-Amino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-5	(R)-2-Amino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-6	(S)-2-Acetylamino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-7	(R)-2-Acetylamino-2-phenylethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-8	2-[N-(3-N-t-Butoxycarbonyl aminopropyl)]acetamide	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-9	2-[N-(3-aminopropyl)]acetamide	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-10	2-(4-Fluoro)acetophenone-	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-11	2-(4-Fluorophenyl)-2-hydroxyethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-12	2-(4'-Fluoro)acetophenone oxime	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-13	2-(4'-Fluoro)acetophenone oxime methylether	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-14	2-acetamide	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ Et
9-15	cyanomethyl	Cl	-CH ₂ C(CH ₃) ₂ CO ₂ Et
9-16	2-(N-benzyl)acetamide	Br	-CH ₂ C(CH ₃) ₂ CO ₂ Et
9-17	2-acetic acid	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-18	2-hydroxyprop-1-yl	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-19	2-Acetamide	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-20	2-Methyl-2-propanamide	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-21	2-(2,2-dimethyl)acetic acid	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-22	2-methoxyprop-1-yl	2-methoxy-pyridin-5-yl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-23	2-Hydroxyprop-1-yl	2-methoxy-pyridin-5-yl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-24	2-Hydroxy-2-methylprop-1-yl	2-methoxy-pyridin-5-yl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-25	3,3-Dimethyl-2-hydroxybut-1-yl	2-methoxy-pyridin-5-yl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-26	2-(4-Fluorophenyl)-2-hydroxyethyl	2-methoxy-pyridin-5-yl	-CH ₂ C(CH ₃) ₂ CO ₂ H
9-27	2-Acetamide	2-thiazolyl	-CH ₂ C(CH ₃) ₂ CO ₂ Et
9-28	2-Acetamide	5-trifluoromethyl-	-CH ₂ C(CH ₂ CH ₃) ₂ CO ₂ H

Compound #	R	G ₆	R ₇
		pyridin-2-yl	

Also described herein are compounds in which the -S- group between the indole group and R is replaced with an -CH₂S- group.

Table 10. Substituted or unsubstituted non-cyclic Y with tertiary alcohol R₇




5

Compound #	Y	Z	R ₁₁
10-1	-C(=O)NH ₂	-CH ₂ S-	4-chlorophenyl
10-2	-C(=O)NH ₂	-CH ₂ S-	Pyridin-4-yl
10-3	-C(=O)NH ₂	-CH ₂ S-	4-cyanophenyl
10-4	-C(=O)NH ₂	-CH ₂ S-	4-iodophenyl
10-5	-C(=O)NH ₂	-CH ₂ S-	cyclopropyl
10-6	-C(=O)N(Et) ₂	-CH ₂ S-	4-chlorophenyl
10-7	-C(=O)NH(4-fluorophenyl)	-CH ₂ S-	Pyridin-4-yl
10-8	-C(=O)N(4-chlorobenzyl)(pyridin-3-yl)	-CH ₂ S-	4-chlorophenyl
10-9	-C(=O)NH(cyclopropyl)	-CH ₂ S-	Pyridin-4-yl
10-10	-C(=O)NH(4-iodobenzyl)	-CH ₂ S-	4-iodo-phenyl
10-11	-C(=O)NH ₂	-CH ₂ S-	4-(pyridin-3-yl)phenyl
10-12	-CO ₂ H	-CH ₂ S-	4-chlorophenyl
10-13	-CO ₂ Et	-CH ₂ S-	4-chlorophenyl
10-14	-CH(OH)CH ₃	-CH ₂ S-	4-chlorophenyl
10-15	-CH(OH)CH ₃	-CH ₂ S-	4-chlorophenyl
10-16	-C(=O)CH ₃	-CH ₂ S-	4-chlorophenyl

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.


Table 11. Acid replacements at G₁



Compound #	Y-	-Z-	G ₁
11-1	Quinolin-2-yl	-CH ₂ S-	2-amino-(1,3,4-oxadiazol-4-yl)
11-2	Quinolin-2-yl	-CH ₂ S-	C(O)NH-thiazol-2-yl
11-3	Quinolin-2-yl	-CH ₂ S-	C(O)NHC(O)NH ₂
11-4	Quinolin-2-yl	-CH ₂ S-	5-methyl-(1,2,4-oxadiazol-3-yl)
11-5	Quinolin-2-yl	-CH ₂ S-	C(=O)NH- pyridin-3-yl
11-6	Quinolin-2-yl	-CH ₂ S-	C(=O)NH-pyrazin-2-yl

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.


Table 12. Alkyl C-2 side chain



Compound #	Y	-Z-	G ₆	R ₆	R ₇
12-1	Pyridin-2-yl	-CH ₂ S-	CO ₂ H	<i>tert</i> -butylsulfanyl	2,2-dimethylprop-1-yl
12-2	Pyridin-2-yl	-CH ₂ S-	CO ₂ Me	<i>tert</i> -butylsulfanyl	2,2-dimethylprop-1-yl
12-3	Pyridin-2-yl	-CH ₂ S-	CO ₂ H	H	2,2-dimethylprop-1-yl
12-4	Pyridin-2-yl	-CH ₂ S-	2-dimethylamino-ethyl-aminocarbonyl	<i>tert</i> -butylsulfanyl	2,2-dimethylprop-1-yl

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.


Table 13. Heteroaryl indole tertiary alcohols











Compound #	R ₁₁
13-1	H
13-2	N-acetyl-azetidin-3-yl
13-3	cyclopropyl
13-4	Cyclobutyl
13-5	4-(N-cyclopropylNHC(=O)-)phenyl
13-6	4-(2-hydroxy-ethyl-aminocarbonyl)phenyl
13-7	-CONH ₂

Also described herein are compounds in which the -CH₂S- group between the indolyl and the pyridinyl groups is replaced with an -S- group.

Table 14. Heteroaryl Y indole compounds



Compound #	Y	Z	R ₁₁	R ₇
14-1	5-methylpyrazin-2-yl	-CH ₂ S-	Morpholin-4-ylmethyl	
14-2	5-methylpyrazin-2-yl	-CH ₂ S-	Piperidin-1-ylmethyl	
14-3	5-methylpyridin-2-yl	-CH ₂ S-	Piperidin-1-ylmethyl	
14-4	pyridin-2-yl	-CH ₂ S-	phenyl	
14-5	5-methylpyridin-2-yl	-CH ₂ S-	phenyl	
14-6	5-methylpyridin-2-yl	-CH ₂ S-	4-tertbutyl-phenyl	
14-7	5-methylpyridin-2-yl	-CH ₂ S-	2-trifluoromethyl-phenyl	
14-8	5-methylpyrazin-2-yl	-CH ₂ S-	4-(5-trifluoromethylpyridin-2-yl)-phenyl	

Also described herein are compounds in which the -CH₂S- group of Z is replaced with an -S- group.

Example 5: FLAP Binding Assays

5 [00972] A non-limiting example of such a FLAP binding assay is as follows:

Packed human polymorphonuclear cell pellets (1.8 x 10⁹ cells) (Biological Speciality Corporation) were resuspended, lysed and 100,000 g membranes prepared as described (Charleson *et al. Mol. Pharmacol.*, 41, 873-879, 1992). 100,000 xg pelleted membranes were resuspended in Tris-Tween assay buffer (100 mM Tris HCl pH 7.4, 140 mM NaCl, 2 mM EDTA, 0.5 mM DTT, 5% glycerol, 0.05% Tween 20) to yield a protein concentration of 50-100 ug/mL. 10 uL membrane suspension was added to 96 well Millipore plate, 78 uL Tris-Tween buffer, 10 uL ³H MK886 or ³H 3-[5-(pyrid-2-ylmethoxy)-3-*tert*-butylthio-1-benzyl-indol-2-yl]-2,2-dimethylpropionic acid (or ¹²⁵I MK591 derivative Egger *et al, J. Labelled Compounds and Radiopharmaceuticals*, 1994, vXXXIV, 1147)) to ~30,000 cpm, 2 uL inhibitor and incubated for 30 minutes at room temperature. 100 uL ice-cold washed buffer was added to the incubation mixture. Plates were then filtered and washed 3x with 200 uL ice cold Tris-Tween buffer, scintillation bottoms sealed, 100 uL scintillant added, shaken for 15 minutes then counted in a TopCount. Specific binding was determined as defined as total radioactive binding minus non-specific binding in the presence of 10 uM MK886. IC₅₀s were determined using Graphpad prism analysis of drug titration curves.

Example 6: Human Blood LTB₄ inhibition Assay

[00973] A non-limiting example of such a human blood LTB₄ inhibition assay is as follows:

Blood was drawn from consenting human volunteers into heparinized tubes and 125 μ L aliquots added to wells containing 2.5 μ L 50% DMSO (vehicle) or 2.5 μ L drug in 50% DMSO. Samples were incubated for 15 minutes at 37°C. 2 μ L calcium ionophore A23817 (from a 50 mM DMSO stock diluted just prior to the assay in Hanks balanced salt solution (Invitrogen)) to 1.25 mM) was added, solutions mixed and incubated for 30 minutes at 37°C. Samples were centrifuged at 1,000 rpm (~200 x g) for 10 minutes at 4°C, plasma removed and a 1:100 dilution assayed for LTB₄ concentration using ELISA (Assay Designs). Drug concentrations to achieve 50% inhibition (IC₅₀'s) of vehicle LTB₄ were determined by nonlinear regression (Graphpad Prism) of % inhibition versus log drug concentration.

Example 7: Rat peritoneal Inflammation and Edema Assay

[00974] A non-limiting example of such a rat peritoneal inflammation and edema assay is as follows:

The *in vivo* efficacy of leukotriene biosynthesis inhibitors was assessed using a rat model of peritoneal inflammation. Male Sprague-Dawley rats (weighing 200 – 300 grams) received a single intraperitoneal (i.p.) injection of 3 mL saline containing zymosan (5 mg/mL) followed immediately by an intravenous (i.v.) injection of Evans blue dye (2 mL of 1.5% solution). Compounds were administered orally (3 mL/kg in 0.5% methylcellulose vehicle) 2 to 4 hours prior to zymosan injection. One to two hours after zymosan injection, rats were euthanized, and the peritoneal cavity was flushed with 10 mL phosphate buffered saline solution (PBS). The resulting fluid was centrifuged at 1,200 rpm for 10 minutes. Vascular edema was assessed by quantifying the amount of Evans blue dye in the supernatant using a spectrophotometer (Absorbance 610 nm). LTB₄ and cysteinyl leukotriene concentrations in the supernatant were determined by ELISA. Drug concentrations to achieve 50% inhibition of plasma leakage (Evans blue dye) and inhibition of peritoneal LTB₄ and cysteinyl leukotrienes could be calculated by nonlinear regression (Graphpad Prism) of % inhibition versus log drug concentration.

Example 8: Human leukocyte inhibition assay

[00975] A non-limiting example of a human leukocyte inhibition assay is as follows:

Blood was drawn from consenting human volunteers into heparinized tubes and 3% dextran, 0.9% saline equal volume added. After sedimentation of red blood cells a hypotonic lysis of remaining red blood cells was performed and leukocytes sedimented at 1000 rpm. The pellet was resuspended at 1.25×10^5 cells/mL and aliquoted into wells containing 2.5 μ L 20% DMSO (vehicle) or 2.5 μ L drug in 20% DMSO. Samples were incubated for 5 minutes at 37 °C. and 2 μ L calcium ionophore A23817 (from a 50 mM DMSO stock diluted just prior to the assay in Hanks balanced salt solution (Invitrogen)) to 1.25 mM) was added, solutions mixed and incubated for 30 minutes at 37 °C. Samples were centrifuged at 1,000 rpm (~200 x g) for 10 minutes at 4 °C, plasma removed and a 1:4 dilution assayed for LTB₄ concentration using ELISA (Assay Designs). Drug concentrations to achieve 50% inhibition (IC₅₀'s) of vehicle LTB₄ were determined by nonlinear regression (Graphpad Prism) of % inhibition versus log drug concentration. The compounds presented in Tables 1-4 had assays of 1 nM to 5 μ M with this assay.

Example 9: Pharmaceutical Compositions**Example 9a: Parenteral Composition**

[00976] To prepare a parenteral pharmaceutical composition suitable for administration by injection, 100 mg of a water-soluble salt of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is dissolved in DMSO and then mixed with 10 mL of 0.9% sterile saline. The mixture is incorporated into a dosage unit form suitable for administration by injection.

Example 9b: Oral Composition

[00977] To prepare a pharmaceutical composition for oral delivery, 100 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is mixed with 750 mg of starch. The mixture is incorporated into an oral dosage unit for, such as a hard gelatin capsule, which is suitable for oral administration.

Example 9c: Sublingual (Hard Lozenge) Composition

[00978] To prepare a pharmaceutical composition for buccal delivery, such as a hard lozenge, mix 100 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), with 420 mg of powdered sugar mixed, with 1.6 mL of light corn syrup, 2.4 mL distilled water, and 0.42 mL mint extract. The mixture is gently blended and poured into a mold to form a lozenge suitable for buccal administration.

Example 9d: Inhalation Composition

[00979] To prepare a pharmaceutical composition for inhalation delivery, 20 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is mixed with 50 mg of anhydrous citric acid and 100 mL of 0.9% sodium chloride solution. The mixture is incorporated into an inhalation delivery unit, such as a nebulizer, which is suitable for inhalation administration.

Example 9e: Rectal Gel Composition

[00980] To prepare a pharmaceutical composition for rectal delivery, 100 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is mixed with 2.5 g of methylcellulose (1500 mPa), 100 mg of methylparaben, 5 g of glycerin and 100 mL of purified water. The resulting gel mixture is then incorporated into rectal delivery units, such as syringes, which are suitable for rectal administration.

Example 9f: Topical Gel Composition

[00981] To prepare a pharmaceutical topical gel composition, 100 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is mixed with 1.75 g of hydroxypropyl cellulose, 10 mL of propylene glycol, 10 mL of isopropyl myristate and 100 mL of purified alcohol USP. The resulting gel mixture is then incorporated into containers, such as tubes, which are suitable for topical administration.


Example 9g: Ophthalmic Solution Composition

[00982] To prepare a pharmaceutical ophthalmic solution composition, 100 mg of a compound of any of Formula (A), Formula (B), Formula (C), Formula (E), Formula (F), Formula (G), or Formula (H), is mixed with 0.9 g of NaCl in 100 mL of purified water and filtered using a 0.2 micron filter. The resulting isotonic solution is then incorporated into ophthalmic delivery units, such as eye drop containers, which are suitable for ophthalmic administration.

[00983] The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this disclosure and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

We claim:

1. A compound of Formula (G):



wherein,

- 5 Z is selected from $S(O)_m$, $C(R_1)_2S(O)_m$, $S(O)_mC(R_1)_2$, wherein each R_1 is independently H, CF_3 , or an optionally substituted C_1 - C_6 alkyl; m is 0, 1 or 2;
- Y is a (substituted or unsubstituted aryl), or -(substituted or unsubstituted heteroaryl);
- R_6 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycloalkyl), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl),
- 10 where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, $C(O)$, $-CH(OH)$, -(substituted or unsubstituted C_1 - C_6 alkyl), or -(substituted or unsubstituted C_2 - C_6 alkenyl);
- R_7 is L_3 -X- L_4 - G_1 , wherein,
- L_3 is a or substituted or unsubstituted alkyl;
- 15 X is a bond, O, $-C(=O)$, $-CR_9(OR_9)$, S, $-S(=O)$, $-S(=O)_2$, $-NR_9$, $-NR_9C(=O)-$, $-C(O)NR_9$, $-NR_9C(O)NR_9$;
- L_4 is a bond, a substituted or unsubstituted branched alkyl, a substituted or unsubstituted straight chain alkyl, a substituted or unsubstituted cyclic alkyl, or a substituted or unsubstituted heterocycloalkyl;
- G_1 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, $-OR_9$, $-C(=O)CF_3$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)C(=O)R_9$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-C(R_9)_2(OR_9)$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, $-S(=O)_2R_8$, $-L_5$ -(substituted or unsubstituted alkyl), $-L_5$ -(substituted or unsubstituted alkenyl), $-L_5$ -(substituted or unsubstituted heteroaryl), or $-L_5$ -(substituted or unsubstituted aryl), wherein L_5 is $-OC(O)O-$, $-NHC(O)NH-$, $-NHC(O)O$, $-OC(O)NH-$, $-NHC(O)$, $-C(O)NH$, $-C(O)O$, or $-OC(O)$;
- 20 or G_1 is $W-G_5$, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G_5 is H, tetrazolyl, $-NHS(=O)_2R_8$, $S(=O)_2N(R_9)_2$, OH, $-OR_8$, $-C(=O)CF_3$, $-C(R_9)_2(OR_9)$, $-C(O)NHS(=O)_2R_8$, $-S(=O)_2NHC(O)R_9$, CN, $N(R_9)_2$, $-N(R_9)C(O)R_9$, $-C(=NR_{10})N(R_9)_2$, $-NR_9C(=NR_{10})N(R_9)_2$, $-NR_9C(=CHR_{10})N(R_9)_2$, $-C(O)NR_9C(=NR_{10})N(R_9)_2$, $-C(O)NR_9C(=CHR_{10})N(R_9)_2$, $-CO_2R_9$, $-C(O)R_9$, $-CON(R_9)_2$, $-SR_8$, $-S(=O)R_8$, or $-S(=O)_2R_8$;
- 30 each R_8 is independently selected from substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted phenyl or substituted or unsubstituted benzyl;
- each R_9 is independently selected from H, substituted or unsubstituted C_1 - C_6 alkyl, substituted or unsubstituted C_1 - C_6 fluoroalkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted phenyl, substituted or unsubstituted benzyl, and substituted or unsubstituted

heteroaryl methyl; or two R₉ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring;
 or R₈ and R₉ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring and
 each R₁₀ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂,
 heteroaryl, or heteroalkyl;

- 5 R₅ is H, halogen, substituted or unsubstituted C₁-C₆alkyl, substituted or unsubstituted -O-C₁-C₆alkyl;
 R₁₁ is L₇-L₁₀-G₆, wherein L₇ is a bond, -C(O), -C(O)NH, -NHC(O), or (substituted or unsubstituted C₁-
 C₆alkyl); L₁₀ is a bond, (substituted or unsubstituted alkyl), (substituted or unsubstituted cycloalkyl),
 (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl), or (substituted or unsubstituted
 heterocycloalkyl);
- 10 G₆ is OR₉, -C(=O)R₉, -C(=O)OR₉, -SR₈, -S(=O)R₈, -S(=O)₂R₈, N(R₉)₂, tetrazolyl, -NHS(=O)₂R₈, -
 S(=O)₂N(R₉)₂, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, -C(=O)N(R₉)₂, N R₉C(O)R₉,
 C(R₉)₂C(=O)N(R₉)₂, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -L₅-(substituted
 or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted
 heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -O-, C(=O), S, S(=O), S(=O)₂, -NH,
 15 -NHC(O)O, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O)-;
 or G₆ is W-G₇, wherein W is (substituted or unsubstituted heterocycloalkyl), (substituted or
 unsubstituted aryl) or a (substituted or unsubstituted heteroaryl) and G₇ is H, halogen, CN, NO₂, N₃, CF₃,
 OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -
 OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -
 20 C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -
 C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -C(R₉)₂(OR₉), -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈,
 -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or
 unsubstituted heteroalkyl), -L₅-(substituted or unsubstituted heteroaryl), -L₅-(substituted or
 unsubstituted heterocycloalkyl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is a bond, -O-,
 25 C(=O), S, S(=O), S(=O)₂, -NH, -NHC(O)O, -NHC(O)NH-, -OC(O)O-, -OC(O)NH-, -NHC(O),
 -C(O)NH, -C(O)O, or -OC(O);

provided that R₁₁ comprises at least one (unsubstituted or substituted) aromatic moiety and at least one
 (unsubstituted or substituted) cyclic moiety, wherein the (unsubstituted or substituted) cyclic moiety is a
 (unsubstituted or substituted) heterocycloalkyl group or a (unsubstituted or substituted) heteroaryl group
 and R₁₁ is not a thienyl-phenyl group; and

30 R₁₂ is H, (substituted or unsubstituted C₁-C₆ alkyl), (substituted or unsubstituted C₃-C₆ cycloalkyl);
 or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or
 pharmaceutically acceptable prodrug thereof.

2. The compound of claim 1, wherein:

- 35 Z is S(O)_m or C(R₁)₂S(O)_m; and
 R₁ is independently H, CF₃, or an optionally substituted C₁-C₆alkyl;
 and optionally wherein m is 0; and R₁ is H or methyl;
 or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or
 pharmaceutically acceptable prodrug thereof.

3. The compound of claim 2, wherein:

Y is -(substituted or unsubstituted heteroaryl); and

G₆ is W-G₇;

and optionally wherein Y is a substituted or unsubstituted heteroaryl containing 0-4 nitrogen atoms, 0-1

O atoms and 0-1 S atoms;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

4. The compound of claim 3, wherein:

Y is selected from the group consisting of pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl,

pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl,

isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolyl, indazolyl, indolizyl,

phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl,

furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl,

quinoxalinyl, naphthyridinyl, imidazo[1,2-a]pyridinyl and furopyridinyl, wherein Y is substituted or

unsubstituted;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

5. The compound of claim 4, wherein:

L₇ is a bond;

L₁₀ is a (substituted or unsubstituted heteroaryl), (substituted or unsubstituted aryl); and

W is (substituted or unsubstituted heterocycloalkyl), (substituted or unsubstituted aryl) or a (substituted or unsubstituted heteroaryl);

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

6. The compound of claim 5, wherein:

L₁₀ is a (substituted or unsubstituted aryl); and

R₁₂ is H;

and optionally wherein W is a (substituted or unsubstituted heterocycloalkyl containing 0-2 nitrogen atoms, 0-1 O atoms and 0-1 S atoms) or a (substituted or unsubstituted heteroaryl containing 0-4

nitrogen atoms, 0-1 O atoms and 0-1 S atoms);

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

7. The compound of claim 6, wherein:

G₇ is H, halogen, CN, NO₂, CF₃, OCF₃, C₁-C₆ alkyl, C₃-C₆cycloalkyl, -C₁-C₆ fluoroalkyl, tetrazolyl, -

NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉,

N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈;

and optionally wherein W is a substituted or unsubstituted group selected from among pyridinyl,

imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl,

thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl,

benzofuranyl, cinnolyl, indazolyl, indolizyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl,

pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazoliny, quinoxaliny, naphthyridinyl, imidazo[1,2-a]pyridinyl, furopyridinyl, quinoliziny, dioxinyl, piperidinyl, morpholinyl, thiazinyl, tetrahydropyridinyl, piperazinyl, oxazinanonyl, dihydropyrrolyl, dihydroimidazolyl, tetrahydrofuranly, tetrahydropyranly, dihydrooxazolyl, oxiranyl, pyrrolidinyl, pyrazolidinyl, dihydrothiophenonyl, imidazolidinonyl, pyrrolidinonyl, dihydrofuranonyl, dioxolanonyl, thiazolidinyl, piperidinonyl, tetrahydronaphyridinyl, tetrahydroquinoliny, tetrahydrothiophenyl, indoliny, tetrahydroquinoliny, and thiazepanyl;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

8. The compound of claim 7, wherein:

R_6 is H, or L_2 -(substituted or unsubstituted alkyl), or L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, -S(O), -S(O)₂, -C(O), -CR₉(OR₉), or substituted or unsubstituted alkyl.;

and optionally wherein X is a bond, O, -C(=O), -CR₉(OR₉), S, -S(=O), -S(=O)₂, -NR₉, -NR₉C(=O)-, or -C(O)NR₉;

or glucuronide metabolite, or pharmaceutically acceptable solvate, or pharmaceutically acceptable salt, or a pharmaceutically acceptable prodrug thereof.

9. The compound of claim 8, wherein:

G_1 is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, -OR₉, -C(=O)CF₃, -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -N(R₉)CH₂CO₂R₉, -C(=NR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)₂, -NR₉C(=CHR₁₀)N(R₉)₂, -NR₉C(=NR₁₀)N(R₉)C(=O)R₉, -C(O)NR₉C(=NR₁₀)N(R₉)₂, -C(O)NR₉C(=CHR₁₀)N(R₉)₂, -CO₂R₉, -C(O)R₉, -C(R₉)₂(OR₉), -CON(R₉)₂, -SR₈, -S(=O)R₈, -S(=O)₂R₈, -L₅-(substituted or unsubstituted alkyl), -L₅-(substituted or unsubstituted alkenyl), -L₅-(substituted or unsubstituted heteroaryl), or -L₅-(substituted or unsubstituted aryl), wherein L₅ is -OC(O)O-, -NHC(O), -C(O)NH, -C(O)O, or -OC(O); or G_1 is W-G₅, where W is a substituted or unsubstituted aryl, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heteroaryl and G₅ is H, tetrazolyl, -NHS(=O)₂R₈, S(=O)₂N(R₉)₂, OH, -OR₈, -C(=O)CF₃, -C(R₉)₂(OR₉), -C(O)NHS(=O)₂R₈, -S(=O)₂NHC(O)R₉, CN, N(R₉)₂, -N(R₉)C(O)R₉, -CO₂R₉, -C(O)R₉, -CON(R₉)₂, -SR₈, -S(=O)R₈, or -S(=O)₂R₈;

and optionally wherein R₆ is hydrogen; methyl; ethyl; propyl; prop-2-yl; 2-methylpropyl; 2,2-dimethylpropyl; butyl; *tert*-butyl; 3-methylbutyl; 3,3-dimethylbutyl; cyclopropylmethyl; cyclobutylmethyl; cyclopentylmethyl; cyclohexylmethyl; benzyl; methoxy, ethoxy, propyloxy; prop-2-yloxy; *tert*-butyloxy; cyclopropylmethoxy; cyclobutylmethoxy; cyclopentylmethoxy; cyclohexylmethoxy; benzyloxy; cyclopropyloxy; cyclobutyloxy; cyclopentyloxy; cyclohexyloxy; phenoxy; acetyl; 2,2,2-trifluoro-acetyl; propanoyl; 2-methylpropanoyl; 2,2-dimethylpropanoyl; 3-methyl-butanoyl; 3,3-dimethylbutanoyl; 2-ethyl-butanoyl; benzoyl; phenylacetyl; cyclopropylcarbonyl; cyclobutylcarbonyl; cyclopentylcarbonyl; cyclohexylcarbonyl; *tert*-butylsulfanyl; *tert*-butyl-sulfinyl; or *tert*-butylsulfonyl;


or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

10. The compound of claim 9, wherein:

G₁ is selected from among H, OH, CN, CO₂H, CO₂Me, CO₂Et, CO₂NH₂, CO₂NHMe, CO₂N(Me)₂,

5

CO₂N(Et)₂, -NH₂, -NHMe, -N(Me)₂, -N(Et)₂, -NMe(iPr),





or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

11. The compound of claim 10, wherein:

15


L₃-X-L₄ is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂C(CH₃)H-, -CH₂C(CH₂CH₃)H-, -CH₂C(isopropyl)H-, -CH₂C(tert-butyl)H-, -CH₂C(CH₃)₂-, -CH₂C(CH₂CH₃)₂-, -CH₂C(CH₃)(CH₂CH₃)-,






or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.


12. The compound of claim 11, wherein:





or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

13. The compound of claim 12, wherein:



or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

14. The compound of claim 9, wherein:

L₂ is an unsubstituted linear alkyl, branched alkyl, or cycloalkyl;

X is a bond:

L_4 is a bond; and

G₁ is OR₉ or -C(O)OR₉;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

15. The compound of claim 14, wherein:

L₃ is methandiyl; ethan-1,2-diyl; propan-1,2-diyl; propan-1,3-diyl; 2-methyl-propan-1,2-diyl; 2-ethyl-propan-1,2-diyl; propan-2,2-diyl; butan-1,2-diyl; butan-1,4-diyl; 2-ethyl-butan-1,2-diyl; 2-propylbutan-1,2-diyl; 3-methylbutan-1,2-diyl; 3,3-dimethylbutan-1,2-diyl; pentan-1,2-diyl; 2-propyl-pentan-1,2-diyl, pentan-1,5-diyl; or hexan-1,6-diyl;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

16. The compound of claim 15, wherein:

L₃ is 2-methyl-propan-1,2-diyl; or 2-ethyl-butan-1,2-diyl;

or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

17. The compound of claim 9, wherein:

L_3 is methandiyl;

X is a bond;

L₄ is propan-2,2-diyl; pentan-3,3-diyl; cyclopropan-1,1-diyl; cyclobutan-1,1-diyl; cyclopentan-1,1-diyl; cyclohexan-1,1-diyl; or cycloheptan-1,1-diyl; and

G_1 is $-\text{CO}_2\text{R}_9$;


or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof.

18. A pharmaceutical composition comprising an effective amount of a compound of any of claims 1-17, or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof, and a pharmaceutically acceptable excipient.



19. A compound of claims 1-17, or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof, for use in the treatment of inflammation in a mammal.

- 5
20. The compound of claims 1-17, or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof, for use in the treatment of respiratory disease in a mammal.
 21. The compound of claims 1-17, or a glucuronide metabolite, pharmaceutically acceptable solvate, pharmaceutically acceptable salt, or pharmaceutically acceptable prodrug thereof, for use in the treatment of cardiovascular disease in a mammal.
 22. The compound of claim 20, wherein the respiratory disease is asthma or chronic obstructive pulmonary disease.

1 / 14


Scheme I**Figure 1**

2 / 14


Scheme IIScheme III**Figure 2**

3 / 14


Batcho-Leimgruber Indole Synthesis




Reissert Indole Synthesis



Hegedus Indole Synthesis



Fukuyama Indole Synthesis



Sugasawa Indole Synthesis






Figure 3

4 / 14


Bischler Indole Synthesis




Gassman Indole Synthesis



Japp-Klingemann Indole Synthesis



Buchwald Indole Synthesis



Larock Indole Synthesis






Figure 4

5 / 14


Bartoli Indole Synthesis




Castro Indole Synthesis



Heck Indole Synthesis



Hemetsberger Indole Synthesis



Mori-Ban Indole Synthesis






Figure 5

6 / 14


Madelung Indole Synthesis



McMurray Indole Synthesis



Nenitzescu Indole Synthesis



Other Syntheses







Figure 6

7 / 14

Using Palladium



Photochemical



Acid Catalyzed Cyclization




Figure 7

8 / 14





Figure 8

9 / 14

**Figure 9**

10 / 14




Figure 10

11 / 14






Figure 11


12 / 14

**Figure 12**

13 / 14

**Figure 13**

14 / 14

**Figure 14**

A. CLASSIFICATION OF SUBJECT MATTER*C07D 209/24(2006.01)i, C07D 209/02(2006.01)i*

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 8 : C07D 401/12, A61K 31/5415

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKIPASS(KIPO internal), CAS (on line), PubMed, "5-lipoxygenase", "indole", "inflammation"

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO 9316069 A1 (MERCK FROSST CANADA Inc.) 1993.08.19. see page 2-3 and claim 1-13	1-4, 18-22 5-17
X A	WO 9325546 A1 (MERCK FROSST CANADA Inc.) 1993.12.23. see page 2-3 and claim 1-18	1-4, 18-22 5-11
A	WO 2007047204 A2 (AMIRA Pharmaceuticals, Lnc.) 2007.04.26. see claim 1-134	1-22
A	Richard Frenette et al., Substituted Indoles as Potent and Orally Active 5-Lipoxygenase Activating Protein (FLAP) Inhibitors, Bioorganic & Medicinal Chemistry Letters. 1999, Vol. 9, pp. 2391-2396.	1-22



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

04 SEPTEMBER 2008 (04.09.2008)

Date of mailing of the international search report

04 SEPTEMBER 2008 (04.09.2008)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
Government Complex-Daejeon, 139 Seonsa-ro, Seo-
gu, Daejeon 302-701, Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

Na, Young Min

Telephone No. 82-42-481-8394



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2008/062793

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9316069 A1	1993.08.19.	AU 3488693 A1	1993.09.03.
WO 9325546 A1	1993.12.23.	US 5334719 A WO 9325546 A1	1994.08.02. 1993.12.23
WO 2007047204 A2	2007.04.26.	EP 01933842 A2 US 07405302 US 2007123522 A1 US 2007123522 AA WO 2007047207 A2	2008.06.25. 2008.07.29. 2007.05.31. 2007.05.31. 2007.04.26.