

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 March 2006 (23.03.2006)

PCT

(10) International Publication Number
WO 2006/031929 A2

(51) International Patent Classification:
A61K 31/4745 (2006.01) C07D 498/02 (2006.01)

(21) International Application Number:
PCT/US2005/032837

(22) International Filing Date:
15 September 2005 (15.09.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/609,992 15 September 2004 (15.09.2004) US

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA, N.V. [BE/BE];
Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CONNOLLY, Peter, J. [US/US]; 74 Delwick Lane, New Providence, NJ 07974 (US). JOHNSON, Sigmond, G. [US/US]; 97 Al-lens Corner Road, Flemington, NJ 08822 (US). PANDEY, Niranjana, B. [US/US]; 52 Barley Sheaf Road, Flemington, NJ 08822 (US). MIDDLETON, Steven, A. [US/US]; 2 Gateshead Drive, Flemington, NJ 08822 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THIAZOLOPYRIDINE KINASE INHIBITORS



(I)

(57) Abstract: The present invention is directed to novel thiazolopyridines, pharmaceutical compositions thereof, and the use thereof as inhibitors of ATP-protein kinase interactions. The thiazolopyridine compounds have the following formula (I).

WO 2006/031929 A2

THIAZOLOPYRIDINE KINASE INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

This present application claims benefit of U.S. Provisional Patent Application Serial No. 60/609992, filed September 15, 2004, which is incorporated herein by
5 reference in its entirety and for all purposes.

FIELD OF THE INVENTION

The present invention is in the area of novel thiazolopyridines and pharmaceutically acceptable forms thereof, their syntheses, and their use as kinase inhibitors.

10

BACKGROUND OF THE INVENTION

In general, protein kinases are the largest set of structurally related phosphoryl transferases, have highly conserved structures and catalytic functions and may be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, histidine and the like) and are responsible for the control of a
15 wide variety of cellular signal transduction processes.

Examples of protein-tyrosine kinases include, but are not limited to, Irk, IGFR-1, Zap-70, Bmx, Btk, CHK (Csk homologous kinase), CSK (C-terminal Src Kinase), Itk-1, Src (c-Src, Lyn, Fyn, Lck, Syk, Hck, Yes, Blk, Fgr and Frk), Tec, Txk/Rlk, Abl, EGFR (EGFR-1/ErbB-1, ErbB-2/NEU/HER-2, ErbB-3 and ErbB-4),
20 FAK, FGF1R (also FGFR1 or FGR-1), FGF2R (also FGR-2), MET (also Met-1 or c-MET), PDGFR- α , PDGFR- β , Tie-1, Tie-2 (also Tek-1 or Tek), VEGFR1 (also FLT-1), VEGFR2 (also KDR), FLT-3, FLT-4, c-KIT, JAK1, JAK2, JAK3, TYK2, LOK, RET, TRKA, PYK2, ALK (Anaplastic Lymphoma Kinase), EPHA (1-8), EPHB (1-6), RON, Fes, Fer or EPHB4 (also EPHB4-1).

25

Examples of protein-serine/threonine kinases include, but are not limited to, Ark, ATM (1-3), CamK (I-IV), CamKK, Chk1 and 2 (Checkpoint kinases), CKI, CK2, Erk, IKK- I (also IKK-ALPHA or CHUK), IKK-2 (also IKK-BETA), Ilk, Jnk (1-3), LimK (1 and 2), MLK3Raf (A, B, and C), CDK (1-10), PKC (including all PKC subtypes), Plk (1-3), NIK, Pak (1-3), PDK1, PKR, RhoK, RIP, RIP-2, GSK3 (A and
30 B), PKA, P38, Erk (1-3), PKB (including all PKB subtypes) (also AKT-1, AKT-2,

AKT-3 or AKT3-1), IRAK1, FRK, SGK, TAK1 or Tpl-2 (also COT).

Protein kinases play very important roles in the normal regulation of cell growth. However, as a result of either mutation or overexpression of the tyrosine kinases (receptor or non-receptor) or the ligands of the receptor tyrosine kinases,
5 signaling can become deregulated, resulting in uncontrolled cell proliferation leading to cancer or a related disease, disorder or syndrome.

Protein kinases catalyze and regulate the process of phosphorylation, whereby the kinases covalently attach phosphate groups to proteins or lipid targets in response to a variety of extracellular signals: hormones, neurotransmitters, growth and
10 differentiation factors, cell cycle events, environmental stresses, nutritional stresses and the like.

Phosphorylation modulates or regulates a variety of cellular processes such as proliferation, growth, differentiation, metabolism, apoptosis, motility, transcription, translation and other signaling processes. Uncontrolled signaling for cell growth due to
15 defective control of protein phosphorylation has also been implicated in a number of diseases and disease conditions, such as osteoarthritis, rheumatoid arthritis, synovial pannus invasion in arthritis, multiple sclerosis, myasthenia gravis, diabetes mellitus, diabetic angiopathies or retinopathy, inflammatory bowel disease, Crohn's disease, ulcerative colitis, transplant or bone marrow transplant rejection, lupus, chronic
20 pancreatitis, cachexia, septic shock, skin diseases or disorders (such as papilloma formation, psoriasis, dermatitis, eczema, seborrhea and the like), central nervous system diseases (such as Alzheimer's disease, Parkinson's disease, depression and the like), cancers (such as glioma cancers, epidermoid cancers, head and neck cancers, lung cancers, breast cancers, colorectal cancers, prostate cancers, gastric cancers, esophageal
25 cancers or papillocarcinomas and the like and associated pathologies such as unregulated cell proliferation, tumor growth or vascularization or metastatic cancer cell invasion and migration and the like or leukemias or lymphomas), ocular diseases (such as macular degeneration, diseases of the cornea, glaucoma and the like), viral infections (such as cytomegalovirus CMV), heart disease (such as atherosclerosis,
30 neointima formation or transplantation-induced vasculopathies (such as restenosis and the like), lung or pulmonary diseases (such as allergic-asthma, lung fibrosis or complications resulting from chronic obstructive pulmonary disorder and the like) or kidney or renal diseases (such as acute, subacute or chronic forms of

glomerulonephritis or membranoproliferative glomerulonephritis, glomerulosclerosis, congenital multicystic renal dysplasia, kidney fibrosis and the like). Therefore, kinase inhibitors have potential use as therapeutic agents.

5 The tyrosine kinases can further be categorized by whether they are receptor tyrosine kinases or non-receptor tyrosine kinases. The receptor tyrosine kinases span the cell membrane with a ligand interacting domain protruding from the cell, with a hydrophobic trans-membrane domain, and a cytoplasmic domain that contains the catalytic kinase domain and other regulatory sequences. Non-receptor tyrosine kinases are often myristylated or modified by the addition of other hydrophobic moieties that
10 allow them to be anchored to the cell membrane.

The epidermal growth factor receptor (EGFR) tyrosine-kinase family includes the receptors EGFR (also referred to as EGFR-1 or Erb-B1), HER-2 (or neu), EGFR3 and EGFR4. Epidermal Growth Factor (EGF), Transforming Growth Factor- α (TGF- α) and the HER-2 ligand heregulin are three of the ligands that bind to the EGFR
15 receptors.

For example, EGFR overexpression or mutation of one or more EGFR kinase family members has been commonly involved in cancer and other diseases characterized by uncontrolled or abnormal cell growth. Deregulation of EGFR has also been associated with epidermoid tumors, head and neck tumors, breast tumors and
20 tumors involving other major organs. Diseases associated with increased EGFR expression include proliferative glomerulonephritis, diabetes-induced renal disease and chronic pancreatitis. Overexpression of HER2 has been associated with breast and ovarian cancer. Diseases associated with the overproduction of TGF- α , rather than overexpression of EGFR, include psoriasis, a cell-proliferative skin disorder. Since
25 EGFR expression levels in uterine tissues are elevated during implantation of a fertilized egg, an EGFR inhibitor may also have potential use as a contraceptive to reduce fertility.

Human cytomegalovirus (CMV) is a widespread opportunistic human herpes virus that causes severe and fatal diseases in those who are immune compromised and
30 in transplant recipients. CMV is also a leading cause of atherosclerosis and virally mediated birth defects. The human CMV uses the EGFR receptor to enter cells during infection, EGFR is autophosphorylated and the downstream signal transduction pathway components are activated; however, the EGFR specific inhibitor tyrphostin

AG1478 has been shown to reduce the viral load in cells that were infected in the presence of the tyrphostin (Wang, et al., Nature, 24 July **2003**, Vol 424). Accordingly, potent EGFR selective inhibitors may be useful in anti-CMV therapy.

The Src family of tyrosine-kinases includes the sub-family proteins c-Src, Lyn, Fyn, Lck, Syk, Hck, Yes, Blk, Fgr and Frk. While various members of the c-Src family are important for normal cellular proliferation, their overexpression and overactivation can promote development of cancer (Yeatman TJ, Nature, June **2004**, Vol. 4). For example, the Lyn kinase has been shown to be upregulated in hormone resistant prostate cancer. Tumor xenografts of hormone resistant prostate cancer cells showed delayed growth upon treatment with peptides that specifically block Lyn kinase activity (Goldenberg-Furmanov, et al., Cancer Research, 1 Feb **2004**, 64, 1058-1064).

The Lyn and Hck Src sub-family tyrosine-kinases have both been implicated in chronic myeloid leukemia (CML). CML is caused by the BCR-Abl fusion protein that results from the t(9;22) chromosomal translocation that juxtaposes the c-Abl non-receptor tyrosine kinase gene on chromosome 9 with a breakpoint cluster region (bcr) gene on chromosome 22. The BCR-Abl fusion protein is a constitutively activated form of the Abl tyrosine kinase that drives uncontrolled growth leading to CML and many cases of adult acute lymphoblastic leukemia. Gleevec, which is an inhibitor of Abl has been successfully used to treat CML. However, Gleevec does not help patients in blast crisis because they carry mutant forms of BCR-Abl that no longer bind Gleevec. Such Gleevec resistant CML cells are sensitive to a dual src/BCR-Abl inhibitor that binds and inhibits the mutant BCR-Abl and members of the src family (Shah, et al., Science, 16 July **2004**, Vol 305, 399-401). There are also other ways that CML cells can become resistant to treatment with the tyrosine kinase Abl inhibitor Gleevec. For example, CML K562 cells that become resistant to Gleevec minimize reliance on the BCR-Abl translocation for growth and instead upregulate the Lyn and Hck kinases. This was demonstrated by expressing antisense Lyn in these cells, which reduced their rate of proliferation (Donato, et al., Blood, 15 Jan **2003**, 101(2)). c-Src and other Src family members are also involved in cellular adhesion, invasion and motility of tumor cells. Thus, small molecule inhibitors of the Src kinase family could offer new therapeutic opportunities for both leukemias and solid tumors.

U.S. Patent Nos. 6,232,320 and 6,579,882 and U.S. Patent Application Publication Nos. 20010020030 and 2003220365 describe compounds for treating

inflammation.

SUMMARY OF THE INVENTION

A first aspect of the present invention is directed to novel compounds of Formula (I).

5 A second aspect of the present invention is directed to pharmaceutical compositions comprising at least one compound of Formula (I), or a pharmaceutically acceptable form thereof and one or more pharmaceutically acceptable excipients.

A third aspect of the present invention is directed to a method of inhibiting the activity of a protein kinase, comprising contacting the protein kinase domain with one
10 or more compounds of Formula (I).

A fourth aspect of the invention is directed to a method of inhibiting unregulated kinase activity comprising administering to an animal a composition comprising a pharmaceutically effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable form thereof and one or more pharmaceutically-
15 acceptable excipients.

A fifth aspect of the present invention is directed to a method of inhibiting increased or unregulated kinase expression or signaling comprising administering to an animal a pharmaceutically effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable form thereof and one or more pharmaceutically-
20 acceptable excipients.

A sixth aspect of the present invention is directed to a method of treating diseases or conditions caused by increased kinase expression or signaling leading to unregulated cell proliferation comprising administering to an animal a pharmaceutically effective amount of at least one compound of Formula (I), or a pharmaceutically
25 acceptable form thereof and one or more pharmaceutically-acceptable excipients.

A seventh aspect of the present invention is directed to a method of treating cancer comprising administering to an animal a pharmaceutically effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable form thereof and one or more pharmaceutically-acceptable excipients.

30 An eighth aspect of the present invention is directed to describing thiazolopyridine derivatives as inhibitors of a tyrosine-kinase selected from EGFR, HER-2, c-Src, Lyn, Abl and the like.

A ninth aspect of the present invention is to describe thiazolopyridine derivatives that are useful at low dosages as inhibitors of protein kinase-induced mitogenesis. This therefore leads to a further aspect of compounds having extremely low cytotoxicity.

5 A tenth aspect of the present invention is to describe thiazolopyridine derivatives that are useful in suppressing tumors, especially tumors such as non-small-cell lung cancers, colon cancers, breast cancers and the like where mitogenesis is heavily driven by protein kinases such as EGFR, HER-2, c-Src, Lyn and the like.

10 An eleventh aspect of the present invention is to describe thiazolopyridine derivatives that have utility as chronic therapeutic agents for inhibiting protein kinase-induced responses. This therefore leads to a further aspect of such therapeutic agents being used as a long-term therapy for inducing cancer remission.

15 A twelfth aspect of the present invention is to describe thiazolopyridine derivatives that have utility as therapeutic agents against chronic or acute diseases characterized by uncontrolled cell proliferation and which may result in metastatic cancer cell invasion and migration.

 A thirteenth aspect of the present invention is to describe thiazolopyridine derivatives that have utility as therapeutic agents against EGFR protein kinase mediated cytomegalovirus (CMV) infection.


20 A fourteenth aspect of the present invention is to describe thiazolopyridine derivatives that have utility as therapeutic agents against chronic or acute kinase mediated diseases or for use as a contraceptive agent.

 A fifteenth aspect of the present invention is directed to methods of synthesizing compounds of Formula (I).

25 These and other aspects and advantages of the invention, which will become apparent in light of the detailed description below, are achieved through use of the compounds of this invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a substituted thiazolopyridine compound of Formula (I)



Formula (I)

and pharmaceutically acceptable forms thereof, wherein

- 5 L^1 is selected from the group consisting of $S(C_{1-4}\text{alkyl})$, a bond, $N(R_1)$, $N(R_1)C(O)$ and $C(O)N(R_1)$, wherein R_1 is selected from the group consisting of hydrogen, $C_{1-8}\text{alkyl}$ and $C_{1-8}\text{alkyl}(C_{1-8}\text{alkoxy})$;
- p is 0, 1, 2, 3 or 4;
- L^2 is selected from the group consisting of O, S, $N(R_1)$ and a bond;
- 10 Ar^1 is selected from the group consisting of aryl, heteroaryl, benzofused heteroaryl, heterocyclyl and benzofused heterocyclyl optionally substituted with from one to three substituents independently selected from the group consisting of
- (1) $C_{1-8}\text{alkyl}$,
 - (2) $C_{2-8}\text{alkenyl}$,
 - 15 (3) $C_{2-8}\text{alkynyl}$,
 - (4) $C_{1-8}\text{alkoxy}$,
- wherein (1), (2), (3) and (4) are optionally substituted with one to three substituents independently selected from the group consisting of
- (i) $C_{3-8}\text{cycloalkyl}$,
 - 20 (ii) aryl,
 - (iii) heteroaryl,
 - (iv) heterocyclyl,
- wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to three substituents independently selected from the group consisting of
- 25 of
- (a) $C_{1-8}\text{alkyl}$,
 - (b) $C_{1-8}\text{alkoxy}$,

- (c) C₁₋₈alkyl(C₁₋₈alkoxy),
- (d) C₁₋₈alkyl(halogen)₁₋₃,
- (e) C₁₋₈alkyl(hydroxy)₁₋₃,
- (f) CO₂(C₁₋₈alkyl),
- 5 (g) amino optionally mono or disubstituted with C₁₋₈alkyl,
- (h) cyano,
- (i) halogen,
- (j) hydroxy,
- (k) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with
- 10 C₁₋₈alkyl,
- (l) C₃₋₈cycloalkyl,
- (m) C₁₋₈alkyl(C₃₋₈cycloalkyl),
- (n) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
- 15 (o) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents, and
- (v) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (a) C₁₋₈alkyl,
- 20 (b) C₁₋₈alkyl(C₁₋₈alkoxy),
- (c) C₁₋₈alkyl(hydroxy)₁₋₃,
- (d) C₃₋₈cycloalkyl,
- (e) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
- 25 (f) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (5) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl,
- 30 (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
- (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (iv) C₁₋₈alkyl(hydroxy)₁₋₃,
- (v) C₃₋₈cycloalkyl,

- (vi) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents,
and
 - (vii) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or
two C₁₋₈alkyl substituents,
- 5 (6) cyano,
- (7) halogen,
- (8) hydroxy,
- (9) C₃₋₈cycloalkyl,
- (10) aryl,
- 10 (11) heteroaryl,
- (12) heterocyclyl,
- (13) oxy substituted with a substituent selected from the group consisting of
- (i) CF₃,
 - (ii) C₃₋₈cycloalkyl,
 - 15 (iii) aryl,
 - (iv) heteroaryl, and
 - (v) heterocyclyl,
- (14) SO₂(heterocyclyl) optionally substituted on heterocyclyl with one or two
C₁₋₈alkyl substituents,
- 20 (15) C(O) substituted with a substituent selected from the group consisting of
- (i) hydrogen,
 - (ii) hydroxy,
 - (iii) C₁₋₈alkyl,
 - (iv) C₁₋₈alkoxy, and
 - 25 (v) amino optionally mono or disubstituted with a substituent independently
selected from the group consisting of
- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (c) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with
 - 30 C₁₋₈alkyl,
 - (d) C₁₋₈alkyl(hydroxy)₁₋₃,
 - (e) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one
or two C₁₋₈alkyl substituents,

- (f) C₃₋₈cycloalkyl, and
(g) aryl; and
- (16) SO₂(amino), wherein amino is optionally mono or disubstituted with a substituent independently selected from the group consisting of
- 5 (i) C₁₋₈alkyl,
(ii) C₁₋₈alkyl(C₁₋₈alkoxy),
(iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
(iv) C₁₋₈alkyl(hydroxy)₁₋₃,
(v) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or
10 two C₁₋₈alkyl substituents,
(vi) C₃₋₈cycloalkyl, and
(vii) aryl; and
- Ar² is selected from the group consisting of aryl, heteroaryl, benzofused heteroaryl, heterocyclyl and benzofused heterocyclyl optionally substituted with from one
15 to three substituents independently selected from the group consisting of
- (1) C₁₋₈alkyl,
(2) C₂₋₈alkenyl,
(3) C₂₋₈alkynyl,
(4) C₁₋₈alkoxy,
- 20 wherein (1), (2), (3) and (4) are optionally substituted with from one to three substituents independently selected from the group consisting of
- (i) C₁₋₈alkoxy,
(ii) cyano,
(iii) halogen,
25 (iv) hydroxy,
(v) C₃₋₈cycloalkyl,
(vi) aryl,
(vii) heteroaryl,
(viii) heterocyclyl,
- 30 wherein (v), (vi), (vii) and (viii) are optionally substituted with from one to three substituents independently selected from the group consisting of
- (a) C₁₋₈alkyl,

- 5 (b) C₁₋₈alkoxy,
(c) C₁₋₈alkyl(C₁₋₈alkoxy),
(d) amino optionally mono or disubstituted with C₁₋₈alkyl,
(e) cyano,
(f) halogen,
(g) C₁₋₈alkyl(halogen)₁₋₃,
(h) hydroxy,
(i) C₁₋₈alkyl(hydroxy)₁₋₃,
(j) C₃₋₈cycloalkyl, and
10 (k) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
substituents,
(ix) amino optionally mono or disubstituted with a substituent independently
selected from the group consisting of
15 (a) C₁₋₈alkyl,
(b) C₁₋₈alkyl(C₁₋₈alkoxy),
(c) C₁₋₈alkyl(hydroxy)₁₋₃,
(d) C₃₋₈cycloalkyl,
(e) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
substituents, and
20 (f) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one
or two C₁₋₈alkyl substituents, and
(x) C(O)amino optionally mono or disubstituted with a substituent
independently selected from the group consisting of
25 (a) C₁₋₈alkyl,
(b) C₁₋₈alkyl(C₁₋₈alkoxy),
(c) C₁₋₈alkyl(hydroxy)₁₋₃,
(d) C₃₋₈cycloalkyl,
(e) aryl optionally substituted with one to three halogen substituents,
(f) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
30 substituents, and
(g) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one
or two C₁₋₈alkyl substituents,
(5) amino optionally mono or disubstituted with a substituent independently

selected from the group consisting of

- 5 (i) C₁₋₈alkyl,
(ii) C₁₋₈alkyl(C₁₋₈alkoxy),
(iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
(iv) C₁₋₈alkyl(hydroxy)₁₋₃,
(v) C₃₋₈cycloalkyl,
(vi) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents,
and
(vii) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or
10 two C₁₋₈alkyl substituents,
- (6) oxy substituted with a substituent selected from the group consisting of
(i) C₃₋₈cycloalkyl,
(ii) aryl,
(iii) heteroaryl, and
15 (iv) heterocyclyl,
wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to
three substituents independently selected from the group consisting
of
(a) C₁₋₈alkyl,
20 (b) C₁₋₈alkoxy,
(c) C₁₋₈alkyl(C₁₋₈alkoxy),
(d) amino optionally mono or disubstituted with C₁₋₈alkyl, and
(e) halogen,
- (7) C(O) substituted with a substituent independently selected from the group
25 consisting of
(i) hydrogen,
(ii) C₁₋₈alkyl optionally substituted with from one to three substituents
independently selected from the group consisting of
(a) cyano,
30 (b) halogen, and
(c) hydroxy,
(iii) C₁₋₈alkoxy,
(iv) hydroxyl, and

- (v) C₁₋₈alkoxy(C₁₋₈alkoxy),
 - (8) cyano,
 - (9) halogen,
 - (10) hydroxy,
 - 5 (11) nitro,
 - (12) C₃₋₈cycloalkyl,
 - (13) aryl,
 - (14) heteroaryl,
 - (15) benzofused heteroaryl,
 - 10 (16) heterocyclyl, and
 - (17) benzofused heterocyclyl;
- wherein (12)-(17) are optionally substituted with one or two substituents independently selected from the group consisting of
- (i) C₃₋₈cycloalkyl,
 - 15 (ii) C₁₋₈alkyl(C₃₋₈cycloalkyl),
 - (iii) aryl,
 - (iv) C₁₋₈alkyl(aryl),
 - (v) heteroaryl,
 - (vi) C₁₋₈alkyl(heteroaryl),
 - 20 (vii) heterocyclyl, and
 - (viii) C₁₋₈alkyl(heterocyclyl).

An example of a compound of Formula (I) is a compound wherein L¹ is selected from the group consisting of N(R₁), wherein R₁ is selected from the group consisting of hydrogen, C₁₋₈alkyl and C₁₋₈alkyl(C₁₋₈alkoxy).

25 An example of a compound of Formula (I) is a compound wherein L¹ is NH.

An example of a compound of Formula (I) is a compound wherein p is 0, 1, 2 or 3.

An example of a compound of Formula (I) is a compound wherein L² is selected from the group consisting of O, S and N(R₁).

30 An example of a compound of Formula (I) is a compound wherein L² is selected from the group consisting of O, S and NH.

An example of a compound of Formula (I) is a compound wherein Ar¹ is selected from the group consisting of aryl, heteroaryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of

5 (1) C₁₋₈alkyl,

(2) C₁₋₈alkoxy,

wherein (1) and (2) are optionally substituted with one to three substituents independently selected from the group consisting of

(i) C₃₋₈cycloalkyl,

10 (ii) aryl,

(iii) heteroaryl,

(iv) heterocyclyl,

wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to three substituents independently selected from the group consisting of

15

(a) C₁₋₈alkyl,

(b) C₁₋₈alkoxy,

(c) C₁₋₈alkyl(C₁₋₈alkoxy),

(d) C₁₋₈alkyl(halogen)₁₋₃,

20

(e) C₁₋₈alkyl(hydroxy)₁₋₃,

(f) amino optionally mono or disubstituted with C₁₋₈alkyl,

(g) cyano,

(h) halogen,

(i) hydroxy, and

25

(j) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl, and

(v) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of

(a) C₁₋₈alkyl,

30

(b) C₁₋₈alkyl(C₁₋₈alkoxy),

(c) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and

(d) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one

- or two C₁₋₈alkyl substituents,
- (3) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl,
 - 5 (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
 - (iii) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
 - (iv) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- 10 (4) halogen,
- (5) hydroxy,
- (6) C₃₋₈cycloalkyl,
- (7) aryl,
- (8) heteroaryl,
- 15 (9) heterocyclyl, and
- (10) SO₂(amino), wherein amino is optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
 - 20 (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
 - (iv) C₁₋₈alkyl(hydroxy)₁₋₃, and
 - (v) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents.

An example of a compound of Formula (I) is a compound wherein Ar¹ is

25 selected from the group consisting of aryl, heteroaryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of

- (1) C₁₋₈alkyl,
 - (2) C₁₋₈alkoxy,
- 30 wherein (1) and (2) are optionally substituted with one to three substituents independently selected from the group consisting of
- (i) heteroaryl, and
 - (ii) heterocyclyl,

wherein (i) and (ii) are optionally substituted with from one to three substituents independently selected from the group consisting of

- (a) C₁₋₈alkyl,
- (b) C₁₋₈alkyl(hydroxy), and
- 5 (c) amino, and
- (iii) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
 - (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 - 10 (c) heterocyclyl, and
 - (d) C₁₋₈alkyl(heterocyclyl),
- (3) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
 - (i) C₁₋₈alkyl(amino), and
 - 15 (ii) C₁₋₈alkyl(heterocyclyl),
- (4) halogen,
- (5) hydroxy,
- (6) heterocyclyl, and
- (7) SO₂(amino), wherein amino is optionally mono or disubstituted with a
- 20 substituent independently selected from the group consisting of
 - (i) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl, and
 - (iii) C₁₋₈alkyl(heterocyclyl),

25 An example of a compound of Formula (I) is a compound wherein Ar² is selected from the group consisting of aryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of

- (1) C₁₋₈alkyl,
- (2) C₂₋₈alkenyl,
- 30 (3) C₂₋₈alkynyl,
- (4) C₁₋₈alkoxy,

wherein (1), (2), (3) and (4) are optionally substituted with from one to three substituents independently selected from the group consisting of

- (i) C₁₋₈alkoxy,
 - (ii) halogen,
 - (iii) hydroxy,
 - (iv) aryl optionally substituted with from one to three substituents independently
- 5 selected from the group consisting of
- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkoxy,
 - (c) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (d) amino optionally mono or disubstituted with C₁₋₈alkyl,
 - 10 (e) halogen,
 - (f) C₁₋₈alkyl(halogen)₁₋₃,
 - (g) hydroxy, and
 - (h) C₁₋₈alkyl(hydroxy)₁₋₃, and
 - (v) amino optionally mono or disubstituted with C₁₋₈alkyl,
- 15 (5) amino optionally mono or disubstituted with a substituent independently
selected from the group consisting of
- (i) C₁₋₈alkyl, and
 - (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (6) oxy substituted with a substituent selected from the group consisting of
- 20 (i) C₃₋₈cycloalkyl,
- (ii) aryl,
 - (iii) heteroaryl, and
 - (iv) heterocyclyl,
- wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to
- 25 three substituents independently selected from the group consisting
of
- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkoxy,
 - (c) amino optionally mono or disubstituted with C₁₋₈alkyl, and
 - 30 (d) halogen,
 - (7) cyano,
 - (8) halogen, and
 - (9) hydroxy.

An example of a compound of Formula (I) is a compound wherein Ar^2 is selected from the group consisting of aryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of

- (1) C_{2-8} alkynyl,
- 5 (2) C_{1-8} alkoxy optionally substituted with from one to three substituents independently selected from the group consisting of
 - (i) halogen, and
 - (ii) aryl optionally substituted with from one to three halogen substituents,
- (3) oxyaryl optionally substituted with from one to three substituents independently
 10 selected from the group consisting of
 - (i) C_{1-8} alkyl,
 - (ii) C_{1-8} alkoxy,
 - (iii) amino optionally mono or disubstituted with C_{1-8} alkyl, and
 - (iv) halogen, and
- 15 (4) halogen.

An example of a compound of Formula (I) is a compound wherein

L^1 is NH;

p is 0, 1, 2 or 3;

L^2 is selected from the group consisting of O, S and NH;

- 20 Ar^1 is selected from the group consisting of aryl, heteroaryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of
 - (1) C_{1-8} alkyl,
 - (2) C_{1-8} alkoxy,
- 25 wherein (1) and (2) are optionally substituted with one to three substituents independently selected from the group consisting of
 - (i) heteroaryl, and
 - (ii) heterocyclyl,

wherein (i) and (ii) are optionally substituted with from one to three
- 30 substituents independently selected from the group consisting of
 - (a) C_{1-8} alkyl,
 - (b) C_{1-8} alkyl(hydroxy), and
 - (c) amino, and

- (iii) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 - 5 (c) heterocyclyl, and
 - (d) C₁₋₈alkyl(heterocyclyl),
- (3) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl(amino), and
 - 10 (ii) C₁₋₈alkyl(heterocyclyl),
 - (4) halogen,
 - (5) hydroxy,
 - (6) heterocyclyl, and
 - (7) SO₂(amino), wherein amino is optionally mono or disubstituted with a
 - 15 substituent independently selected from the group consisting of
 - (i) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl, and
 - (iii) C₁₋₈alkyl(heterocyclyl); and
- 20 Ar² is selected from the group consisting of aryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of
- (1) C₂₋₈alkynyl,
 - (2) C₁₋₈alkoxy optionally substituted with from one to three substituents
 - 25 independently selected from the group consisting of
 - (i) halogen, and
 - (ii) aryl optionally substituted with from one to three halogen substituents, and
 - (3) oxyaryl optionally substituted with from one to three substituents independently selected from the group consisting of
 - 30 (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkoxy,
 - (iii) amino optionally mono or disubstituted with C₁₋₈alkyl, and
 - (iv) halogen, and











(4) halogen.

An example of a compound of Formula (I) is a compound wherein $L^1-(CH_2)_p-Ar^1$ and L^2-Ar^2 are dependently selected from:


Cpd	$L^1-(CH_2)_p-Ar^1$	L^2-Ar^2
1	NH-[4-CH ₂ -(2-CH ₂ CH ₃ -imidazol-1-yl)]-phenyl	NH-3-Cl-4-F-phenyl
2	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-3-Cl-4-F-phenyl
3	NH-[4-CH ₂ N(CH ₃) ₂]-phenyl	NH-3-Cl-4-F-phenyl
4	NH-{4-CH ₂ N[(CH ₂) ₂ OCH ₃] ₂ }-phenyl	NH-3-Cl-4-F-phenyl
5	NH-{4-SO ₂ N[(CH ₂) ₂ OCH ₃] ₂ }-phenyl	NH-3-Cl-4-F-phenyl
6	NH-{4-CH ₂ -[(2 <i>S</i>)-2-CH ₂ OH-pyrrolidin-1-yl]}-phenyl	NH-3-Cl-4-F-phenyl
7	NH-[4-SO ₂ NH(CH ₂) ₂ -morpholin-4-yl]-phenyl	NH-3-Cl-4-F-phenyl
8	NH-(CH ₂) ₃ -morpholin-4-yl	NH-3-Cl-4-F-phenyl
9	NH-(CH ₂) ₃ -morpholin-4-yl	NH-3-C≡CH-phenyl
10	NH-[4-CH ₂ N(CH ₃) ₂]-phenyl	NH-3-C≡CH-phenyl
11	NH-{4-CH ₂ -[(2 <i>S</i>)-2-CH ₂ OH-pyrrolidin-1-yl]}-phenyl	NH-3-C≡CH-phenyl
12	NH-(CH ₂) ₂ -morpholin-4-yl	NH-3-Cl-phenyl
13	NH-(CH ₂) ₂ -morpholin-4-yl	NH-3-C≡CH-phenyl
14	NH-{4-CH ₂ N(CH ₃)CH ₂ -[(2 <i>R</i>)-tetrahydro-furan-2-yl]}-phenyl	NH-3-Cl-4-F-phenyl
15	NH-{4-CH ₂ N(CH ₃)CH ₂ -[(2 <i>S</i>)-tetrahydro-furan-2-yl]}-phenyl	NH-3-Cl-4-F-phenyl
16	NH-[4-CH ₂ N(CH ₃)-tetrahydro-pyran-4-yl]-phenyl	NH-3-Cl-4-F-phenyl
17	NH-(4-CH ₂ -pyrrolidin-1-yl)-phenyl	NH-3-Cl-4-F-phenyl
18	NH-[3-Cl-4-CH ₂ N(CH ₃) ₂]-phenyl	NH-3-Cl-4-F-phenyl
19	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-3-Cl-phenyl
20	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
21	NH-{4-CH ₂ NHCH ₂ -[(2 <i>R</i>)-tetrahydro-furan-2-yl]}-phenyl	NH-3-Cl-4-F-phenyl
22	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-3-Cl-4-F-phenyl
23	NH-[4-SO ₂ NH(CH ₂) ₃ N(CH ₃) ₂]-phenyl	NH-3-C≡CH-phenyl
24	NH-(4-CH ₂ -pyrrolidin-1-yl)-phenyl	NH-{3-Cl-4-[OCH ₂ -(3-F-phenyl)]}-phenyl
25	NH-[4-CH ₂ N(CH ₃) ₂]-phenyl	NH-2,2-F ₂ -benzo[1,3]dioxol-4-yl
26	NH-6-OCH ₃ -pyridin-3-yl	NH-3-Cl-4-F-phenyl

Cpd	L ¹ -(CH ₂) _p -Ar ¹	L ² -Ar ²
27	NH-6-OH-pyridin-3-yl	NH-3-Cl-4-F-phenyl
28	NH-6-Cl-pyridin-3-yl	NH-3-Cl-4-F-phenyl
29	NH-[6-NH(CH ₂) ₃ -morpholin-4-yl]-pyridin-3-yl	NH-3-Cl-4-F-phenyl
30	NH-6-NH(CH ₂) ₂ NH ₂ -pyridin-3-yl	NH-3-Cl-4-F-phenyl
31	NH-[4-CH ₂ -(4-NH ₂ -piperidin-1-yl)]-phenyl	NH-3-Cl-4-F-phenyl
32	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-6-Br-benzo[1,3]dioxol-4-yl
33	NH-(4-CH ₃)-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
34	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-2,4-Cl ₂ -phenyl
35	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	NH-2-Cl-5-OCH ₃ -phenyl
36	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	2,4-Cl ₂ -phenoxy
37	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	3-OCF ₃ -phenoxy
38	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	S-3-Br-phenyl
39	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	S-2-Cl-4-F-phenyl
40	NH-[4-OCH ₂ -(1-CH ₃ -piperidin-3-yl)]-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
41	NH-(4-CH ₂ -piperidin-1-yl)-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
42	NH-{3-OCH ₃ -4-[O(CH ₂) ₂ -morpholin-4-yl]}-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
43	NH-[3-OCH ₃ -4-(OCH ₂ -tetrahydro-pyran-2-yl)]-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
44	NH-{4-CH ₂ NHCH ₂ -[(2 <i>R</i>)-tetrahydro-furan-2-yl]}-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
45	NH-(4-CH ₂ -pyrrolidin-1-yl)-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
46	NH-[4-CH ₂ -(4-CH ₃ -piperazin-1-yl)]-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
47	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-2,4-Cl ₂ -5-OCH ₃ -phenyl
48	NH-(4-CH ₂ -morpholin-4-yl)-phenyl	NH-3-Br-phenyl
49	NH-[4-(CH ₂) ₂ -morpholin-4-yl]-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
50	NH-[4-(CH ₂) ₂ -piperidin-1-yl]-phenyl	NH-5-Cl-benzo[1,3]dioxol-4-yl
51	NH-[4-CH ₂ -(4-CH ₃ -piperazin-1-yl)]-phenyl	NH-(4-phenoxy)-phenyl


An example of a compound of Formula (I) includes a compound selected from the group consisting of:

Cpd. No. 1**Cpd. No. 2****Cpd. No. 3****Cpd. No. 4****Cpd. No. 5****Cpd. No. 6****Cpd. No. 7****Cpd. No. 8****Cpd. No. 9****Cpd. No. 10**


Cpd. No. 11




Cpd. No. 12




Cpd. No. 13




Cpd. No. 14




Cpd. No. 15




Cpd. No. 16




Cpd. No. 17




Cpd. No. 18




Cpd. No. 19




Cpd. No. 20




Cpd. No. 21




Cpd. No. 22




Cpd. No. 23




Cpd. No. 24




Cpd. No. 25




Cpd. No. 26




Cpd. No. 27




Cpd. No. 28























Cpd. No. 29




Cpd. No. 30



Cpd. No. 31**Cpd. No. 32****Cpd. No. 33****Cpd. No. 34****Cpd. No. 35****Cpd. No. 36****Cpd. No. 37****Cpd. No. 38****Cpd. No. 39****Cpd. No. 40**

Cpd. No. 41**Cpd. No. 42****Cpd. No. 43****Cpd. No. 44****Cpd. No. 45****Cpd. No. 46****Cpd. No. 47****Cpd. No. 48****Cpd. No. 49****Cpd. No. 50**

Cpd. No. 51

*Pharmaceutically Acceptable Forms*

Certain compounds of the Formula (I) may exist in various stereoisomeric or tautomeric forms. The present invention encompasses all such kinase inhibiting compounds. These compounds include active compounds in the form of enantiomers or diastereomers. These enantiomers or diastereomers include pure or essentially pure compounds. These compounds also encompass mixtures of enantiomers or mixtures of diastereomers. When enantiomers are present in equimolar amounts, the resulting mixtures are referred to as racemic mixtures. Some of the compounds include tautomers, or a mixture of tautomers.

The compounds of the present invention may be present in the form of pharmaceutically acceptable salts. For use in medicines, the “pharmaceutically acceptable salts” of the compounds of this invention refer to non-toxic acidic/anionic or basic/cationic salts.

Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Thus, representative pharmaceutically acceptable salts include the following: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium, camsylate (or camphosulphonate), carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, fumarate, gluconate, glutamate, hydrabamine, hydrobromine, hydrochloride, iodide, isothionate, lactate, malate, maleate, mandelate, mesylate, nitrate, oleate, pamoate, palmitate, phosphate/diphosphate, salicylate, stearate, sulfate, succinate, tartrate, tosylate.

Furthermore when the compounds of the present invention carry an acidic

moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

5 Pharmaceutically acceptable basic/cationic salts include, and are not limited to salts formed with aluminum, 2-amino-2-hydroxymethyl-propane-1,3-diol, ammonia, benzathine, t-butylamine, calcium, calcium gluconate, calcium hydroxide, chloroprocaine, choline, choline bicarbonate, choline chloride, cyclohexylamine, diethanolamine, ethylenediamine, lithium, LiOMe, L-lysine, magnesium, meglumine,
10 NH₃, NH₄OH, N-methyl-D-glucamine, piperidine, potassium, potassium-t-butoxide, potassium hydroxide (aqueous), procaine, quinine, sodium, sodium carbonate, sodium-2-ethylhexanoate, sodium hydroxide, triethanolamine or zinc.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups
15 on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, 3rd Edition, John Wiley & Sons, 1999. The protecting groups may be removed at a convenient subsequent stage using methods known in the
20 art.

The present invention contemplates compounds of various isomers and mixtures thereof. The term "isomer" refers to compounds that have the same composition and molecular weight but differ in physical and/or chemical properties. Such substances have the same number and kind of atoms but differ in structure. The structural
25 difference may be in constitution (geometric isomers) or in an ability to rotate the plane of polarized light (stereoisomers).

The term "stereoisomer" refers to isomers of identical constitution that differ in the arrangement of their atoms in space. Enantiomers and diastereomers are stereoisomers wherein an asymmetrically substituted atom acts as a chiral center. The
30 term "chiral" refers to a molecule that is not superimposable on its mirror image. The term "enantiomer" refers to one of a pair of molecular species that are mirror images of each other and are not superimposable. The term "diastereomer" refers to stereoisomers that are not related as mirror images. The symbols "R" and "S" represent

the configuration of substituents around a chiral atom(s). The symbols "R*" and "S*" denote the relative configurations of substituents around a chiral atom(s).

The term "racemate" or "racemic mixture" refers to a compound of equimolar quantities of two enantiomeric species, wherein the compound is devoid of optical activity. The term "optical activity" refers to the degree to which a chiral molecule or nonracemic mixture of chiral molecules rotates the plane of polarized light.

The term "geometric isomer" refers to isomers that differ in the orientation of substituent atoms in relationship to a carbon-carbon double bond, to a cycloalkyl ring or to a bridged bicyclic system. Substituent atoms (other than H) on each side of a carbon-carbon double bond may be in an E or Z configuration. In the "E" (opposite sided) configuration, the substituents are on opposite sides in relationship to the carbon-carbon double bond; in the "Z" (same sided) configuration, the substituents are oriented on the same side in relationship to the carbon-carbon double bond. Substituent atoms (other than H) attached to a cycloalkyl ring may be in a cis or trans configuration. In the "cis" configuration, the substituents are on the same side in relationship to the plane of the ring; in the "trans" configuration, the substituents are on opposite sides in relationship to the plane of the ring. Compounds having a mixture of "cis" and "trans" species are designated "cis/trans". Substituent atoms (other than H) attached to a bridged bicyclic system may be in an "endo" or "exo" configuration. In the "endo" configuration, the substituents attached to a bridge (not a bridgehead) point toward the larger of the two remaining bridges; in the "exo" configuration, the substituents attached to a bridge point toward the smaller of the two remaining bridges.

It is to be understood that the various substituent stereoisomers, geometric isomers and mixtures thereof used to prepare compounds of the present invention are either commercially available, can be prepared synthetically from commercially available starting materials or can be prepared as isomeric mixtures and then obtained as resolved isomers using techniques well-known to those of ordinary skill in the art.

The isomeric descriptors "R," "S," "S*," "R*," "E," "Z," "cis," "trans," "exo" and "endo" are used as described herein for indicating atom configuration(s) relative to a core molecule and are intended to be used as defined in the literature (IUPAC Recommendations for Fundamental Stereochemistry (Section E), Pure Appl. Chem., 1976, 45:13-30).

The term "tautomer" is used for isomers that are the result of tautomerism that

in turn designates a rapid and reversible interconversion of isomers associated with the actual movement of electrons as well as one or more hydrogen atoms. Each tautomer is capable of independent existence and potential isolation. An art-known example of tautomerism is keto-enol tautomerism.

5 The compounds of the present invention may be prepared as individual isomers by either isomer-specific synthesis or resolved from an isomeric mixture. Conventional resolution techniques include combining the free base (or free acid) of each isomer of an isomeric pair with an optically active acid (or base) to form an optically active salt (followed by fractional crystallization and regeneration of the free base or acid),
10 forming an ester or amide of each of the isomers of an isomeric pair by reaction with an appropriate chiral auxiliary (followed by fractional crystallization or chromatographic separation and removal of the chiral auxiliary) or resolving an isomeric mixture of either a starting material or a final product using chiral preparative TLC (thin layer chromatography) or a chiral HPLC column.

15 Furthermore, the compounds of the present invention may have one or more polymorph or amorphous crystalline forms and as such are intended to be included in the scope of the invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and as such are also intended to be encompassed within the scope of this invention.

20 *Chemical Definitions*

When any variable (e.g., aryl, heterocyclyl, R₁, R₂, etc.) occurs more than once in a substituent list, its definition on each occurrence is independent of any other occurrence.

25 Bond lines drawn into a ring system from a substituent variable (such as R₆, R₉, etc.) indicate that the substituent may be attached to any of the substitutable ring carbon atoms or heteroatoms. If the ring system is polycyclic, the substituent may be attached to any of the suitable carbon atoms or heteroatoms on the ring into which the bond line is drawn.

30 As used herein, the following terms are intended to have the following definitions. Additional definitions are provided throughout the specification as needed.

The term "C₁₋₈ alkyl," whether used alone or as part of a substituent group, refers to a saturated aliphatic branched or straight-chain monovalent hydrocarbon

radical or alkylidyl linking group having a specified number of carbon atoms, wherein the radical is derived by the removal of one hydrogen atom from a carbon atom and the alkylidyl linking group is derived by the removal of one hydrogen atom from each of two carbon atoms in the chain. The term "C₁₋₈alkyl" refers to a radical having from 1-8
5 carbon atoms in a linear or branched arrangement. For example, "C₁₋₈alkyl" specifically includes the radicals methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, *tert*-butyl, 1-butyl, 1-pentyl, 2-pentyl, 3-pentyl, 1-hexyl, 2-hexyl, 3-hexyl, 1-heptyl, 2-heptyl, 3-heptyl, 1-octyl, 2-octyl, 3-octyl and the like. Alkyl and alkylidyl radicals may be attached to a core molecule via a terminal carbon atom or via a carbon atom within
10 the chain. Similarly, any number of substituent variables may be attached to an alkyl or alkylidyl radical when allowed by available valences.

The term "C₁₋₄alkyl," whether used alone or as part of a substituent group, refers to a saturated aliphatic branched or straight-chain monovalent hydrocarbon radical or alkylidyl linking group having a specified number of carbon atoms, wherein
15 the radical is derived by the removal of one hydrogen atom from a carbon atom and the alkylidyl linking group is derived by the removal of one hydrogen atom from each of two carbon atoms in the chain. The term "C₁₋₈alkyl" refers to a radical having from 1-4 carbon atoms in a linear or branched arrangement. For example, "C₁₋₄alkyl" specifically includes the radicals methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl,
20 *tert*-butyl, 1-butyl, and the like. Alkyl and alkylidyl radicals may be attached to a core molecule via a terminal carbon atom or via a carbon atom within the chain. Similarly, any number of substituent variables may be attached to an alkyl or alkylidyl radical when allowed by available valences.

The term "C₁₋₈ alkyl(halogen)₁₋₃" refers to a C₁₋₈ alkyl radical as defined above
25 wherein one, two or three hydrogen atoms are replaced by a halogen atom. Examples of C₁₋₈ alkyl(halogen)₁₋₃ are for instance, trifluoromethyl, monochloromethane, and the like.

The term "C₁₋₈ alkyl(hydroxy)₁₋₃" refers to a C₁₋₈ alkyl radical as defined above wherein one, two or three hydrogen atoms are replaced by a hydroxyl group. Hydroxy
30 groups are preferably on different carbon atoms, although it is possible that in certain cases two hydroxyl groups will be on the same carbon atom. Examples of "C₁₋₈ alkyl(hydroxy)₁₋₃" are for instance hydroxymethyl, hydroxyethyl and the like.

The term "C₂₋₈ alkenyl," whether used alone or as part of a substituent group,

refers to an alkyl or alkylidyl radical containing at least one carbon to carbon double bond. Typically, one carbon-carbon double bond is present; and, optionally, any number of double bonds may be present whereby a stable compound results. The radical may be in either the *Z* or *E* conformation about the double bond(s).

5 The term "C₂₋₆alkenyl" refers to an alkenyl radical having from 2-6 carbon atoms. For example, "C₂₋₄alkenyl" specifically includes the radicals ethenyl, propenyl, allyl (2-propenyl), butenyl, pentenyl, hexenyl and the like. As described above, an alkenyl radical may be similarly attached to a core molecule and further substituted where indicated.

10 The term "C₂₋₈ alkynyl," whether used alone or as part of a substituent group, refers to an alkyl or alkylidyl radical containing at least one carbon to carbon triple bond. Typically, one carbon-carbon triple bond is present; and, optionally, any number of triple bonds may be present whereby a stable compound results.

 The term "C₂₋₆alkynyl" refers to an alkynyl radical having from 2-6 carbon
15 atoms. For example, "C₂₋₄alkynyl" specifically includes the radicals ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like. As described above, an alkynyl radical may be similarly attached to a core molecule and further substituted where indicated.

 The term "C₁₋₈ alkoxy," whether used alone or as part of a substituent group, refers to an alkyl or alkylidyl radical attached through an oxygen linking atom. For
20 example, "C₁₋₄alkoxy" specifically includes the radicals methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy, heptoxy, octoxy and the like. As described above, an alkoxy radical may be similarly attached to a core molecule and further substituted where indicated.

 The term "C₃₋₈ cycloalkyl," whether used alone or as part of a substituent
25 group, refers to a saturated or partially unsaturated, monocyclic or fused alkyl ring system radical. Examples of cycloalkyl ring systems include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, adamantanyl or indanyl and the like. For example, "C₃₋₈cycloalkyl" specifically includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. As
30 described above, a cycloalkyl radical may be similarly attached to a core molecule and further substituted where indicated.

 The term "aryl," whether used alone or as part of a substituent group, refers to an aromatic monocyclic or fused ring system radical. Examples of aryl ring systems

include, but are not limited to, phenyl, naphthalenyl, fluorenyl, indenyl, azulenyl, anthracenyl, naphthalenyl, fluorenyl, indenyl or anthracenyl and the like. As described above, an aryl radical may be similarly attached to a core molecule and further substituted where indicated.

5 The term "aromatic" refers to a cycloalkyl or heterocyclyl ring system having a fully unsaturated, conjugated π electron system; i.e. an aryl or heteroaryl ring system.

 The term "heterocyclyl," whether used alone or as part of a substituent group, refers to a saturated or partially saturated, monocyclic or fused heterocycloalkyl ring system radical. Partially saturated systems include rings such as those named with
10 dihydro, trihydro, tetrahydro, hexahydro and the like. Examples of heterocyclyl radicals include, and are not limited to, 2H-pyrrole (including 2-pyrrolyl or 3-pyrrolyl), pyrrolidinyl, 1,3-dioxolanyl, 2-imidazolyl (also referred to as 4,5-dihydro-1H-imidazolyl), imidazolidinyl, 2-pyrazolyl, pyrazolidinyl, tetrazolyl, tetrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl,
15 piperazinyl, azetidyl, azepanyl, hexahydro-1,4-diazepyl, hexahydro-1,4-oxazepanyl, tetrahydro-furanyl, tetrahydro-thienyl, tetrahydro-pyranyl, tetrahydro-pyridazinyl, 1,3-benzodioxolyl or 2,3-dihydro-1,4-benzodioxinyl and the like. As described above, a heterocyclyl radical may be similarly attached to a core molecule and further substituted where indicated.

20 The term "hetero," when used as a prefix for a ring system, refers to the replacement of at least one carbon atom member in a cycloalkyl or aryl ring system with a heteroatom selected from N, S, O or P. Where chemically stable and allowed by the amount of ring atoms, a ring may have 1, 2, 3 or 4 nitrogen atom members; alternatively, a ring may have 0, 1, 2 or 3 nitrogen atom members and 1 oxygen or
25 sulfur atom member. When allowed by available valences, up to four adjacent ring members may be nitrogen atoms (e.g. tetrazole); and, up to two adjacent ring members may be heteroatoms, where one heteroatom is nitrogen and the other is one selected from N, S or O.

 The term "heteroaryl," whether used alone or as part of a substituent group,
30 refers to a heteroaromatic monocyclic or fused ring system radical. Examples of heteroaryl rings include, and are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolizyl, indolyl,

azaindolyl, isoindolyl, benzo[b]furyl, benzo[b]thienyl, indazolyl, azaindazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, 4H-quinoliziny, quinoliny, isoquinoliny, cinnoliny, phthalziny, quinazoliny, quinoxaliny, naphthyridiny, pteridiny and the like.

5 The term “benzo-fused” refers to a bicyclic fused ring system radical formed by a monocyclic radical fused with a benzene ring; the benzo-fused radical may be attached to a core molecule via either ring of the bicyclic system.

 The term “benzo-fused cycloalkyl” refers to a bicyclic ring system radical wherein a monocyclic cycloalkyl radical is fused to a benzene ring. Typical benzo-
10 fused cycloalkyl radicals include indanyl, 1,2,3,4-tetrahydro-naphthalenyl, 5,6,7,8-tetrahydro-naphthalenyl, 6,7,8,9,-tetrahydro-5*H*-benzocycloheptenyl, 5,6,7,8,9,10-hexahydro-benzocyclooctenyl and the like. A benzo-fused cycloalkyl ring system is a subset of the cycloalkyl group.

 The term “benzo-fused heterocyclyl” refers to a bicyclic ring system radical
15 wherein a monocyclic heterocyclyl radical is fused to a benzene ring. Typical benzo-fused heterocyclyl radicals include 1,3-benzodioxolyl (also known as 1,3-methylenedioxyphenyl), indoliny, 2,3-dihydro-1,4-benzodioxiny (also known as 1,4-ethylenedioxyphenyl), benzo-dihydro-furyl, benzo-tetrahydro-pyrany, benzo-dihydro-thienyl and the like. A benzo-fused heterocyclyl ring system is a subset of the
20 heterocyclyl group.

 The term “benzo-fused heteroaryl” refers to a bicyclic ring system radical wherein a monocyclic heteroaryl radical is fused to a benzene ring. Typical benzo-fused heteroaryl radicals include indolyl, isoindolyl, benzo[b]furyl, benzo[b]thienyl, indazolyl, benzthiazolyl, quinoliny, isoquinoliny, cinnoliny, phthalaziny, quinazoliny, pteridiny
25 and the like. A benzo-fused heteroaryl ring is a subset of the heteroaryl group.

 The term “fused” refers to ring systems fused on adjacent ring atoms, those fused on a single ring atom and those fused on nonadjacent ring atoms. The types and amount of rings formed are limited by available ring valences, starting materials or synthetic methods. In any case, all such fused ring systems are intended to be included
30 in the scope of the present compounds and associated synthetic methods.

 Monocyclic rings forming bicyclic, tricyclic and the like ring systems are fused on adjacent ring atoms, wherein each monocyclic ring shares two ring atoms, whereby the common ring atoms are the points of attachment for the two rings, and wherein the

common atoms may be a carbon or nitrogen atom. Bicyclic, tricyclic and the like ring system radicals are derived by the removal of one hydrogen atom from a single ring carbon atom of the ring system.

Monocyclic rings forming spirocyclic ring systems are fused on a single ring atom, wherein each monocyclic ring shares one ring atom, whereby the common ring atom is the point of attachment for the two rings, and wherein the common atom is a carbon atom. Spirocyclyl radicals are derived by the removal of one hydrogen atom from a single ring carbon atom of the ring system. Typical spirocyclyl radicals include spiro[1.4]dioxolanyl, spiro[5.5]undecyl, 3-oxa-1-aza-spiro[4.4]nonenyl and the like.

Monocyclic ring systems fused on nonadjacent ring atoms are those that form a variety of bridged bicyclic ring systems. Bridged bicyclic radicals are derived by the removal of one hydrogen atom from a single ring carbon atom of the ring system. Typical bridged bicyclic radicals include bicyclo[2.2.1]heptyl, bicyclo[2.2.1]heptenyl, bicyclo[2.2.2]octyl, bicyclo[3.1.1]heptyl, bicyclo[3.2.1]octyl, bicyclo[2.2.2]octenyl, bicyclo[3.2.1]octenyl, 2-aza-bicyclo[2.2.1]heptyl, 1-aza-bicyclo[2.2.2]octyl, 8-aza-bicyclo[3.2.1]octyl, 7-oxa-bicyclo[2.2.1]heptyl and the like.

Polycyclic bridged and nonbridged ring systems are those ring systems which are further fused on nonadjacent ring atoms to form a variety of common polycyclic ring systems. Polycyclyl radicals are derived by the removal of one hydrogen atom from a single ring carbon atom of the ring system. Typical polycyclyl radicals include adamantanyl, quinuclidinyl, octahydro-4,7-methano-1*H*-indenyl, octahydro-2,5-methano-pentalenyl and the like.

The term "C(O)" refers to a carbonyl linking group moiety.

The term "substituted," refers to a core molecule in which one or more hydrogen atoms have been replaced with that amount of substituents allowed by available valences. Substitution is not limited to the core molecule, but may also occur on a substituent radical, whereby the radical becomes a linking group.

The term "independently selected" refers to two or more substituents that may be selected from a substituent variable group, wherein the selected substituents may be the same or different.

The term "dependently selected" refers to one or more substituent variables that are specified in an indicated combination for substitution in a core molecule (e.g. variables that refer to groups of substituents appearing in a tabular list of compounds).

In general, IUPAC nomenclature rules are used herein. Nomenclature for radical substituents was derived by first indicating the atom having the point of attachment, followed by the linking group atoms toward the terminal chain atom from left to right, substantially as in:

5 $-(C_{1-6})\text{alkyl}-C(O)NH-(C_{1-6})\text{alkyl}-Ph.$

Pharmaceutical Preparations & Methods Of Use

The present invention includes a first method for inhibiting unregulated protein kinase activity comprising contacting the protein kinase domain with one or more compounds of Formula (I).

10 The first method also includes inhibiting increased or unregulated protein kinase expression or signaling leading to unregulated cell proliferation comprising contacting a protein kinase receptor with one or more compounds of Formula (I).

The first method further comprises inhibiting the unregulated expression of a protein kinase such as EGFR, HER-2, c-Src, Lyn, Abl and the like.

15 The present invention includes a second method for use of a compound of Formula (I) as a therapeutic agent for treating or preventing a chronic or acute kinase mediated disease in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) or composition thereof.

The second method includes administering to the subject an effective amount of
20 a compound of Formula (I) or composition thereof in the form of a medicament.

The second method includes a chronic or acute disease mediated by a tyrosine-kinase selected from EGFR, HER-2, c-Src or Lyn.

The present invention includes a third method for use of a compound of Formula (I) as a therapeutic agent for inhibiting the effects of unregulated kinase activity in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) or composition or medicament thereof.

The third method includes inhibiting the effects of unregulated activity of a tyrosine-kinase selected from EGFR, HER-2, c-Src or Lyn.

The present invention includes a fourth method for use of a compound of
30 Formula (I) as a therapeutic agent for treating or preventing a chronic or acute kinase
mediated cancer in a subject in need thereof comprising administering to the subject an
effective amount of a compound of Formula (I) or composition thereof.

The fourth method includes suppressing a chronic or acute tumor associated with non-small-cell lung cancers, colon cancers, breast cancers and the like wherein the cancer is mediated by a tyrosine-kinase selected from EGFR, HER-2, c-Src or Lyn.

5 The fourth method also includes treating or preventing chronic unregulated cell proliferation whereby cancer remission is induced in the subject.

The present invention includes a fifth method for use of a compound of Formula (I) as a therapeutic agent for treating or preventing chronic or acute kinase mediated diseases characterized by unregulated cell proliferation or metastatic cancer cell invasion and migration.

10 The present invention includes a sixth method for use of a compound of Formula (I) as a therapeutic agent for treating or preventing viral infection by an EGFR kinase mediated cytomegalovirus.

The present invention encompasses the compounds of Formula (I) for use as a medicine.

15 The present invention further encompasses the use of the compounds of Formula (I) for the manufacture of a medicament for treating any of the conditions mentioned above.

The term "chronic or acute kinase mediated disorder" as used herein, includes, and is not limited to disorders and diseases associated with unregulated kinase activity and conditions that accompany such activity.

20 The term "unregulated kinase activity" refers to 1) increased or unregulated kinase expression or signaling, 2) increased kinase expression leading to unregulated cell proliferation, 3) increased kinase signalling leading to unregulated cell proliferation, or 4) mutations leading to constitutive kinase activation. The existence of unregulated kinase activity may be determined by procedures well known in the art.

The term "unregulated cell proliferation" refers to cell proliferation of one or more subset of cells in a multicellular organism resulting in harm (such as discomfort or decreased life expectancy) to the multicellular organism.

30 Tumor cells which result from unregulated cell proliferation use many mechanisms to enhance their survival and spread and often have high rates of proliferation because growth control signals that keep normal cells in check are defective. Many tumor cells secrete autocrine growth factors that increase proliferation rates or they induce other cells to secrete growth factors that they utilize.

Tumor cells grow and spread by dislodging from a primary tumor site, using proteases to digest the extracellular matrix, spreading in response to migration cues, allowing them to migrate to certain tissues preferentially where overexpressed adhesion molecules allow attachment and growth at the new site. The totality of these and other biological processes are responsible for the lethal effects of a tumor. A kinase inhibitor may affect one or more aspects of tumor survival mechanisms and thus be therapeutically useful. Alternatively, a kinase inhibitor may not affect one particular tumor survival mechanism but may still be therapeutically useful by affecting tumor survival by an unknown or as yet unelucidated mechanism of action.

The foregoing methods contemplate that the compounds of the present invention are therapeutically useful for treating, preventing or ameliorating diseases disorders or conditions such as, without limitation, osteoarthritis, rheumatoid arthritis, synovial pannus invasion in arthritis, multiple sclerosis, myasthenia gravis, diabetes mellitus, diabetic angiopathies or retinopathy, inflammatory bowel disease, Crohn's disease, ulcerative colitis, transplant or bone marrow transplant rejection, lupus, chronic pancreatitis, cachexia, septic shock, skin diseases or disorders (such as papilloma formation, psoriasis, dermatitis, eczema, seborrhea and the like), central nervous system diseases (such as Alzheimer's disease, Parkinson's disease, depression and the like), cancers (such as glioma cancers, epidermoid cancers, head and neck cancers, lung cancers, breast cancers, colorectal cancers, prostate cancers, gastric cancers, esophageal cancers or papilocarcinomas and the like and associated pathologies such as unregulated cell proliferation, tumor growth or vascularization or metastatic cancer cell invasion and migration and the like or leukemias or lymphomas), ocular diseases (such as macular degeneration, diseases of the cornea, glaucoma and the like), viral infections (such as cytomegalovirus), heart disease (such as atherosclerosis, neointima formation or transplantation-induced vasculopathies (such as restenosis and the like), lung or pulmonary diseases (such as allergic-asthma, lung fibrosis or complications resulting from chronic obstructive pulmonary disorder and the like) or kidney or renal diseases (such as acute, subacute or chronic forms of glomerulonephritis or membranoproliferative glomerulonephritis, glomerulosclerosis, congenital multicystic renal dysplasia, kidney fibrosis and the like).

The term "administering . . . an effective amount of a compound. . ." with respect to the methods of the present invention, refers to a means for treating,

ameliorating or preventing a disease, disorder or syndrome as described herein with a compound specifically disclosed or a compound or prodrug thereof, which would obviously be included within the scope of the invention albeit not specifically disclosed for certain of the instant compounds.

5 Such methods include therapeutically administering an effective amount of one or more compounds of Formula (I) or a composition or medicament thereof at different times during the course of a therapy or concurrently in a combination form. Such methods further include therapeutically administering an effective amount of one or more compounds of Formula (I) with one or more therapeutic agents at different times
10 during the course of a therapy or concurrently in a combination form.

 The term "prodrug" refers to a metabolic precursor of a compound of Formula (I) or pharmaceutically acceptable form thereof. In general, a prodrug is a functional derivative of a compound which may be inactive when administered to a subject but is readily convertible in vivo into an active metabolite compound.

15 The term "active metabolite" refers to a metabolic product of a compound that is pharmaceutically acceptable and effective. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

 The term "subject" as used herein, refers to an animal, preferably a mammal,
20 most preferably a human, who has been the object of treatment, observation or experiment and is at risk of (or susceptible to) developing a disease or disorder or having a disease or disorder related to unregulated kinase activity.

 The term "effective amount" refers to that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response (such as
25 inhibiting activation of unregulated kinase activity) in a tissue system, animal or human, that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated. The effective amount of a compound of Formula (I) exemplified in such a method is from about 0.001 mg/kg/day to about 300 mg/kg/day or has an IC₅₀ (50%
30 inhibition concentration) of about 25 μ M or less, or about 10 μ M or less, preferably of about 1 μ M or less, more preferably of about 0.5 μ M or less, and most preferably of about 0.1 μ M or less.

 The term "composition" refers to a product containing a compound of the

present invention (such as a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from such combinations of the specified ingredients in the specified amounts).

The term "medicament" refers to a product for use in treating or ameliorating a
5 kinase mediated disorder or condition.

The term "pharmaceutically acceptable" refers to molecular entities and compositions that are of sufficient purity and quality for use in the formulation of a composition or medicament of the present invention and that, when appropriately administered to an animal or a human, do not produce an adverse, allergic or other
10 untoward reaction. Since both human use (clinical and over-the-counter) and veterinary use are equally included within the scope of the present invention, a pharmaceutically acceptable formulation would include a composition or medicament for either human or veterinary use.

The term "combination therapy" refers to the use of one or more compounds of
15 Formula (I) or composition or medicament thereof in combination with one or more therapeutic agents for the treatment of a number of different kinase mediated disorders and advantageously may facilitate the use of a reduced effective dose of the compound of Formula (I) and/or the therapeutic agent than would be recommended for the treatment of a particular unregulated cell proliferation disorder. Therefore, it is
20 contemplated that the compounds of this invention can be used before, during or after treatment with a particular therapeutic agent.

The term "therapeutic agent" refers to chemotherapeutic agents used to treat a kinase mediated cancer or antiviral agents used to treat cytomegalovirus. Chemotherapeutic agents include and are not limited to anti-angiogenic agents, anti-
25 tumor agents, cytotoxic agents, inhibitors of cell proliferation, radiation therapy and the like or mixtures thereof.

The terms "treating" or "preventing" refer, without limitation, to facilitating the eradication of, inhibiting the progression of or promoting stasis of a malignancy.

The term "radiation therapy" refers to a therapy that comprises exposing the
30 subject in need thereof to radiation. The present invention includes a method for administering one or more compounds of Formula (I) or composition or medicament thereof in combination with radiation therapy. Procedures for administering such therapy are known to those skilled in the art. The appropriate scheme of radiation

therapy will be similar to those already employed in clinical therapies wherein the radiation therapy is used alone or in combination with other chemotherapeutic agents.

An embodiment of the present invention includes a pharmaceutical composition comprising an admixture of one or more compounds of Formula (I) and/or one or more
5 pharmaceutically acceptable forms thereof and one or more pharmaceutically acceptable excipients.

The pharmaceutically acceptable forms for a compound of Formula (I) include a pharmaceutically acceptable salt, ester, prodrug or active metabolite of a compound of Formula (I).

10 The present invention further includes the use of a process for making the composition or medicament comprising mixing one or more of the instant compounds and an optional pharmaceutically acceptable carrier; and, includes those compositions or medicaments resulting from such a process. Contemplated processes include both conventional and unconventional pharmaceutical techniques.

15 The composition or medicament may take a wide variety of forms to effectuate mode of administration, including, but not limited to, intravenous (both bolus and infusion), oral, nasal, transdermal, topical with or without occlusion, and injection intraperitoneally, subcutaneously, intramuscularly, intratumorally or parenterally. The composition or medicament may be in a dosage unit such as a tablet, pill, capsule,
20 powder, granule, sterile parenteral solution or suspension, metered aerosol or liquid spray, drop, ampoule, auto-injector device or suppository; for administration orally, parenterally, intranasally, sublingually or rectally or by inhalation or insufflation. Compositions or medicaments suitable for oral administration include solid forms such as pills, tablets, caplets, capsules (each including immediate release, timed release and
25 sustained release formulations), granules and powders; and, liquid forms such as solutions, syrups, elixirs, emulsions and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions. Alternatively, the composition or medicament may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound,
30 such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection.

The dosage form (tablet, capsule, powder, injection, suppository, teaspoonful and the like) containing the composition or medicament contains an effective amount of

the active ingredient necessary to be therapeutically or prophylactically effective as described above. The composition or medicament may contain from about 0.001 mg to about 5000 mg (preferably, from about 0.001 to about 500 mg) of the active compound or prodrug thereof and may be constituted into any form suitable for the mode of administration selected for a subject in need. A contemplated effective amount may range from about 0.001 mg to about 300 mg/kg of body weight per day. Preferably, the range is from about 0.003 to about 100 mg/kg of body weight per day. Most preferably, the range is from about 0.005 to about 15 mg/kg of body weight per day. The composition or medicament may be administered according to a dosage regimen of from about 1 to about 5 times per day.

For oral administration, the composition or medicament is preferably in the form of a tablet containing, e.g., 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. Optimal dosages will vary depending on factors associated with the particular patient being treated (e.g., age, weight, diet and time of administration), the severity of the condition being treated, the compound being employed, the mode of administration and the strength of the preparation. The use of either daily administration or post-periodic dosing may be employed.

A person skilled in the art would be able to determine the correct dose given the condition of the patient.

Synthetic Methods


Representative compounds of the present invention can be synthesized in accordance with the general synthetic schemes described below and are illustrated more particularly in the specific synthetic examples that follow. The general schemes and specific examples are offered by way of illustration; the invention should not be construed as being limited by the chemical reactions and conditions expressed. Except where indicated, starting materials and intermediates used in the schemes and examples are prepared by known methodologies well within the ordinary skill of persons versed in the art. No attempt has been made to optimize the yields obtained in any of the example reactions. One skilled in the art would also know how to increase such yields through routine variations in materials, solvents, reagents, reaction conditions and the

like. All commercially available chemicals were obtained from commercial suppliers and used without further purification. Particular equipment components used in the examples such as reaction vessels and the like are also commercially available.

The terms used in describing the invention are commonly used and known to those skilled in the art. Chemical formulas as used herein will have their ordinary meaning known to persons skilled in the art. When used herein, the following abbreviations have the indicated meanings:


AcOH	acetic acid
Cpd	compound
DCM	dichloromethane
DMF	N,N-dimethylformamide
DMSO	dimethyl sulfoxide
EtOAc	ethyl acetate
hr(s)/min(s)	hour(s)/min(s)
IPA	isopropyl alcohol
LAH	lithium aluminum hydride
MCPBA	meta-chloroperoxybenzoic acid or 3-chloroperoxybenzoic acid
MeOH	methanol
NMP	<i>n</i> -methyl pyrrolidinone or 1-methyl-2-pyrrolidinone
Ph	phenyl
PSI	pounds per square inch
RT/rt/r.t.	room temperature
sat'd	saturated
TEA	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran

SCHEME A




10 Chloroacetyl chloride Compound **A2** is reacted with a solution of t-butyl cyanoacetate Compound **A1** in a base at a temperature maintained between about -20°C and 10°C for a period of time (about 3-8 hr) to provide Compound **A3** as the product. The Compound **A1** solution is formed using a base such as sodium hydride, potassium


t-butoxide and the like or mixtures thereof and a solvent that is suitable for the given base such as THF, DMF, t-butanol and the like or mixtures thereof.



Compound **A3** is treated with an organic acid solution and is maintained at a temperature between about -20°C and 5°C for a period of time (about 18 hr) to provide Compound **A4**. The organic acid solution is formed using a preferred acid such as TFA and the like in a preferred solvent such as methylene chloride and the like).




Compound **A4** is reacted with a solution of a suitably substituted Compound **A5** at a temperature between about -20°C to 50°C for a period of time (about 5-72 hr) to provide Compound **A6**. Compound **A5** is a thioimido salt such as a potassium salt substituted with a suitable leaving group or nascent leaving group, L^{a} , such as $-\text{SCH}_3$ and the like. The Compound **A5** solution is formed using a solvent such as acetone, THF, EtOAc and the like or mixtures thereof.




15


Compound **A6** is reacted with a solvated reagent at a temperature between about 50°C to 125°C for a period of time (typically 1-2 hr) to provide Compound **A7**. The solvated reagent is formed using a reagent such as DMF-dimethyl acetal or trimethylorthoformate and the like or mixtures thereof in a solvent such as DMF and the like.




A solution of Compound **A7** is reacted with a chlorinating agent solution at a temperature maintained between about -20°C to 110°C to provide Compound **A8**. The chlorinating agent is a neat or solvated agent such as thionyl chloride, phosphorus oxychloride, phosphorus pentachloride and the like or mixtures thereof in a solvent such as DMF and the like.



Compound **A8** is reacted with a solution of Compound **A9** to provide Compound **A10**. When conducting the reaction at elevated temperatures, the Compound **A9** solution is formed using a solvent such as isopropanol; alternatively, when reacted at a temperature in the range of from about -20°C to about 50°C, the solution is formed using a strong non-nucleophilic base such as potassium t-butoxide and the like. The L^2 portion of Compound **A9** may be varied for reaction with a suitable C^7 halogen atom on Compound **A8** to provide other compounds of the present invention. The L^2 portion is a suitably reactive heteroatom such as N, NH, NH_2 , O or S.



- Compound **A10** is reacted with an oxidizing agent solution at a temperature maintained between about -20°C to 5°C to provide Compound **A11**. The oxidizing agent is an agent such as MCPBA or hydrogen peroxide and the like or mixtures thereof and the solution is formed using a solvent such as methylene chloride, chloroform or up to 10% ethanol and the like or mixtures thereof. When the Compound **A10** L^{a} group is a moiety such as SCH_3 , the reaction formed an “activated” group such as $\text{S}(\text{O})\text{CH}_3$.




- Compound **A11** is reacted with an acidic solution of Compound **A12** at a temperature maintained between about 20°C to 60°C for a period of time (typically 2-24 hr) to provide a compound of Formula (I). The Compound **A12** solution is formed using an acid such as acetic acid and the like.


- The L^1 portion moiety of Compound **A12** may be reacted with a suitable group activated on Compound **A11** to provide other compounds of Formula (I) wherein the L^1 moiety is a suitably reactive heteroatom such as NH , NH_2 , O or S .

SCHEME B


Scheme B describes a procedure whereby Compound **A7** may also be generated from **A3** by an alternative route.




Compound **A3** is reacted with a solution of a suitably substituted Compound **A5** at a temperature between about -20°C to 50°C for a period of time (about 5-48 hr) to provide Compound **B1**. Compound **A5** is a thioimido salt such as a potassium salt substituted with a suitable leaving group or nascent leaving group, L^a , such as -SCH₃ and the like. The Compound **A5** solution is formed using a solvent such as acetone, THF, EtOAc and the like or mixtures thereof.



Compound **B1** is reacted with a solvated reagent at a temperature between about 50°C to 125°C for a period of time (typically 1-2 hr) to provide Compound **B2**. The solvated reagent is formed using a reagent such as DMF-dimethyl acetal or trimethylorthoformate and the like or mixtures thereof in a solvent such as DMF and the like.



Compound **B2** is treated with an organic acid solution and is maintained at a temperature between about -20°C and 5°C for a period of time (about 3 hr) to provide Compound **B3**. The organic acid solution is formed using a preferred acid such as TFA
 5 and the like in a preferred solvent such as methylene chloride and the like.




Compound **B3** is heated in a polar protic solvent at a temperature between about 50°C and 100°C for a period of time (approximately 6 hr) to provide Compound **A7**. The reaction is carried out in a preferred solvent such as isopropanol and the like.

10


Example 1

7-(3-chloro-4-fluorophenylamino)-2-(4-pyrrolidin-1-ylmethylphenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **17**)




A solution of t-butyl cyanoacetate Compound **1a** (32.2 g, 228 mmol, 1.14 M) in
 15 THF was added dropwise to a suspension of 60% NaH (9.13 g, 228 mmol) in THF (200

mL) under nitrogen at 0 °C. The mixture was stirred until hydrogen gas evolution ceased and the solution became homogeneous. A solution of chloroacetyl chloride Compound **1b** (18.5 mL, 228 mmol) in THF (100 mL) was added at a rate sufficiently slow enough to maintain the temperature below 10°C. After stirring for 4 hr, the
 5 reaction was quenched at 0°C with water (25 mL). The resulting precipitate was filtered through Celite 545. The organic solution was then evaporated down to give a residue that was partitioned between ether and 1M NaOH. The aqueous layer was washed with ether (3X) and adjusted to pH 5 with NaH₂PO₄. The aqueous layer was extracted with EtOAc, then dried over MgSO₄ and evaporated to yield 4-chloro-2-cyano-3-hydroxy-but-2-enoic acid tert-butyl ester Compound **1c** (26.3 g) as a light
 10 brown foam. ¹H NMR (CDCl₃) δ 4.34 (br s, 2H), 1.47 (s, 9H). MS 216 (M⁻).




TFA (75 mL) in water (10 mL) was added dropwise to a solution of 4-chloro-2-cyano-3-hydroxy-but-2-enoic acid tert-butyl ester Compound **1c** (47.8 g, 220 mmol) in
 15 DCM (200 mL) at 0 °C. The mixture was then warmed to ambient temperature and stirred for 18 hr. The mixture was evaporated down under vacuum to give a residue that was partitioned between DCM and a solution of saturated sodium bicarbonate. After repeated washing, the DCM layer was dried over MgSO₄ and evaporated down to give a crude product N-tert-butyl-4-chloro-3-oxo-butamide Compound **1d** (23.9 g) as
 20 a dark brown viscous oil. ¹H NMR (CDCl₃) δ 6.32 (br s, 1H), 4.28 (s, 2H), 3.49 (s, 2H), 1.37 (s, 9H). MS 192, 194 (MH⁺).




Potassium methyl N-cyanodithioimidocarbonate Compound **1e** (21.3g, 125 mmol) was added to a solution of Compound **1d** (23.9 g, 125 mmol) at 0 °C in acetone
 25 (300 mL). The reaction was stirred for 1 hr at 0 °C and then at RT for 18 hrs. The

solvent was removed under vacuum and the product extracted into EtOAc from water. The combined organic extracts were dried over MgSO_4 , then filtered and evaporated to give 3-(4-amino-2-methylsulfanyl-thiazol-5-yl)-N-tert-butyl-3-oxo-propionamide Compound **1f** (34.6 g) as a tan solid. The material was used in the next step without further purification. ^1H NMR (DMSO) δ 7.73 (br s, 3H), 3.35 (s, 2H), 2.68 (s, 3H), 1.27 (s, 9H). MS 288 (MH^+).




A mixture of Compound **1f** (34.6 g, 120 mmol), DMF-dimethyl acetal (34.0 mL, 240 mmol, 94 % purity) and DMF (200 mL) were combined and heated to 80 °C for 90 minutes. The conversion to the cyclized intermediate, 7-hydroxy-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile was confirmed by LCMS (MS 298, MH^+). The mixture was then cooled to 0 °C and SOCl_2 was added dropwise (44 mL, 600 mmol). The reaction mixture was stirred for 2 hrs at ambient temperature. The mixture was then cooled back down to 0 °C and water (approximately 300 mL) was added dropwise. The resulting brown solid was collected by filtration and subsequently washed with water and MeOH to provide 7-chloro-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **1g** (16.1 g) as a tan solid. ^1H NMR (CDCl_3) δ 8.82 (s, 1H), 2.91 (s, 3H). MS 242, 244 (MH^+), 264, 266 (MNa^+).

Alternative Route to Compound **1g**




To a stirred solution of 4-chloro-2-cyano-3-hydroxy-but-2-enoic acid tert-butyl ester Compound **1c** (51.3 g, 236 mmol) at 0 °C in acetone (750 mL) was added potassium methyl N-cyanodithioimidocarbonate Compound **1e** (40.2g, 236 mmol). After 1 hr the reaction was allowed to warm to RT. The reaction mixture was evaporated down after stirring for 18 hrs. The resulting residue was extracted with

ethyl acetate from saturated NaHCO_3 . The combined organic layers were dried over MgSO_4 and the solvent removed under reduced pressure to yield a tan foam, 3-(4-amino-2-methylsulfanyl-thiazol-5-yl)-2-cyano-3-oxo-propionic acid tert-butyl ester Compound **1h**, 72.2 g. ^1H NMR (CDCl_3) δ 4.34 (br s, 2H), 3.14 (br s, 1H), 2.70 (br s, 3H), 1.47 (br s, 9H). MS 314 (MH^+).




A mixture of 3-(4-amino-2-methylsulfanyl-thiazol-5-yl)-2-cyano-3-oxo-propionic acid tert-butyl ester Compound **1h** (72.1 g, 230 mmol), N,N-dimethylformamide dimethyl acetal (65.3 mL, 460 mmol, 94 % purity) and DMF (400 mL) were combined and heated to 80 °C for 2 hrs. The cooled reaction was diluted with water and extracted with ethyl acetate. The combined organic extracts were then washed with a brine solution and dried over MgSO_4 . The solvent was removed under reduced pressure to give a dark brown foam Compound **1i**. MS 369 (MH^+). The foam was dissolved in DCM (200 mL) and cooled to 0 °C for the dropwise addition of TFA (200 mL). The reaction was stirred at RT for 3 hrs and then evaporated down to provide N-tert-butyl-3-[4-(dimethylamino-methyleneamino)-2-methylsulfanyl-thiazol-5-yl]-3-oxo-propionamide Compound **1j**. MS 343 (MH^+).




The crude material, Compound **1j**, was dissolved in IPA (500 mL) and refluxed for 6 hrs. The solvent was removed under reduced pressure to provide 2-

methylsulfanyl-7-oxo-4,7-dihydro-thiazolo[4,5-b]pyridine-6-carboxylic acid tert-butylamide Compound **1k**. MS 298 (MH⁺).

The crude material was dissolved up in DMF (500 mL), cooled to 0 °C, and SOCl₂ added dropwise (84 mL, 1.15 mol). The reaction mixture was stirred for 18 hrs
 5 then diluted with water and extracted with ether. The combined organic extracts were filtered through Celite 545 and then washed with water. The filtered organic solution was then dried over MgSO₄ and concentrated down to 200 mL. The tan solid, Compound **1g**, precipitated upon addition of MeOH was collected and washed consecutively with MeOH and hexane, 10.1 g. The spectral data was identical for both
 10 routes to form Compound **1g**.




Compound **1g** (2.30 g, 9.54 mmol) was combined with 3-chloro-4-fluoro-phenylamine Compound **1l** (1.39 g, 9.54 mmol) in IPA (15 mL) in a sealed vessel and heated for 4h at 145 °C. The precipitate formed upon cooling was collected by
 15 filtration and subsequently washed with MeOH and hexane to provide 7-(3-chloro-4-fluoro-phenylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **1m** (2.1 g) as a tan solid. ¹H NMR (DMSO-d₆) δ 9.77 (s, 1H), 8.66 (s, 1H), 7.55 (dd, J = 6.6, 2.8 Hz, 1H), 7.48 (t, J = 8.9 Hz, 1H), 7.30 (ddd, J = 8.9, 4.3, 2.8 Hz, 1H), 2.74 (s, 3H). MS 351, 353 (MH⁺).



20 4-nitrobenzyl chloride Compound **1n1** (4.0 g, 23.3 mmol) and pyrrolidine Compound **1n2** (4.8 mL, 58.3 mmol) were combined in THF (50 mL) at ambient temperature. After 18 hr, the reaction was diluted with 10% NH₄Cl and extracted with EtOAc. The combined organic layers were dried over MgSO₄, filtered and evaporated

in vacuo to provide 1-(4-nitrobenzyl)-pyrrolidine Compound **1n3** (4.7 g) as a pale yellow oil without further purification (prepared as described in Mitsuru Shiraishi, et al., *J. Med. Chem.*, **2000**, 43, 2049-2063). ^1H NMR (CDCl_3) δ 8.17 (d, $J=8.6$ Hz, 2H); 7.51 (d, $J=8.6$ Hz, 2H); 3.72 (s, 2H); 2.58-2.48 (m, 4H); 1.89-1.72 (m, 4H).

5 10% Pd/C (500 mg) was added to a solution of 1-(4-nitrobenzyl)-pyrrolidine Compound **1n3** (4.56 g, 22.1 mmol) in EtOAc (20 mL). The mixture was hydrogenated at 50 PSI for a period of 2 hrs and filtered through Celite. The filtrate was evaporated and the residue was dissolved in 10% NH_4Cl and washed with ethyl ether. The aqueous layer was then adjusted to pH 10 with NaOH and extracted with
10 EtOAc. The combined organic layers were dried over MgSO_4 , filtered and evaporated *in vacuo* to give 4-pyrrolidin-1-ylmethyl-phenylamine Compound **1n** (3.05 g) as a light brown oil. ^1H NMR (CDCl_3) δ 7.10 (d, $J=8.5$ Hz, 2H); 6.63 (d, $J=8.5$ Hz, 2H); 3.60 (br s, 2H); 3.50 (s, 2H); 2.52-2.42 (m, 4H); 1.83-1.69 (m, 4H).




15 MCPBA (1.36 g, 6.06 mmol, 77 % purity) was added to a solution of Compound **1m** (1.06 g, 3.03 mmol) in CHCl_3 (30 mL) at 0 °C. The reaction was diluted after 2 hr with a saturated solution of NaHCO_3 and extracted with DCM. The organic extracts were combined, dried over MgSO_4 and concentrated *in vacuo* to provide 7-(3-chloro-4-fluoro-phenylamino)-2-methanesulfinyl-thiazolo[4,5-b]pyridine-6-carbonitrile as an intermediate. MS 367 (MH^+). AcOH (30 mL) and 4-pyrrolidin-1-ylmethyl-phenylamine Compound **1n** (480 mg, 2.73 mmol) were added to the residue. The mixture was stirred for 2 hr, the solvent was evaporated and the residue was partitioned between EtOAc and saturated NaHCO_3 . The mixture was extracted with EtOAc, dried (MgSO_4), filtered and concentrated *in vacuo* to provide a yellow solid.
20 The solid was dissolved in MeOH, treated with an excess of a 2M HCl-ether solution and recrystallized from MeOH/ether to provide 7-(3-chloro-4-fluoro-phenylamino)-2-(4-pyrrolidin-1-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile

Compound **17** (708 mg) as a bis-hydrochloride salt. ^1H NMR (DMSO- d_6) δ 11.52 (br s, 1H), 10.97 (br s, 1 H), 9.99 (br s, 1H), 8.71 (s, 1H), 7.80 (d, J = 8.6 Hz, 2 H), 7.64 (d, J = 8.6 Hz, 2 H), 7.59 (dd, J = 6.8, 2.6 Hz, 1H), 7.54 (t, J = 8.8 Hz, 1 H), 7.35 (ddd, J = 8.8, 4.3, 2.6 Hz, 1 H), 4.29 (s, 2H), 3.40-3.25 (m, 2 H), 3.11-2.9 (m, 2 H), 2.09-1.80 (m, 4H). MS 479, 481 (MH^+).


Example 2

7-(3-chloro-4-fluorophenylamino)-2-[4-(2-ethyl-imidazol-1-ylmethyl)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **1**)



Using the procedure of Example 1, 2-ethyl-1*H*-imidazole Compound **2a1** was used in place of Compound **1n2** and carried forward to prepare 2-ethyl-1-(4-nitrobenzyl)-1*H*-imidazole Compound **2a2**. Using the procedure of Example 21, Compound **2a2** was used in place of Compound **21a2** and carried forward to prepare 4-(2-ethyl-imidazol-1-ylmethyl)-phenylamine Compound **2a**.

15



Using the procedure of Example 1, Compound **2a** was used in place of Compound **1n** and carried forward to prepare Compound **1** which was isolated by semi-preparative HPLC (C^{18} column, 30 to 60% $\text{CH}_3\text{CN}/\text{H}_2\text{O}/0.05\%$ TFA) as a trifluoroacetate salt upon lyophilization. ^1H NMR (DMSO- d_6) δ 10.95 (s, 1 H), 9.51 (s, 1 H), 8.55 (s, 1 H), 7.77 (d, J = 8.4 Hz, 2 H), 7.70 (d, J = 1.8 Hz, 1H), 7.66 (d, J = 1.8 Hz, 1 H), 7.54-7.45 (m, 2 H), 7.37 (d, J = 8.4 Hz, 2 H), 7.27 (ddd, J = 8.7, 4.1, 2.6 Hz, 1


20

H), 5.38 (s, 2 H), 3.00 (q, J = 7.5 Hz, 2 H), 1.21 (t, J = 7.5 Hz, 3 H). MS 504, 506 (MH⁺).

Example 3


7-(3-chloro-4-fluoro-phenylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 2)

5



Using the procedure of Example 1, morpholine Compound 3a1 was used in place of Compound 1n2 and carried forward to prepare 4-(4-nitro-benzyl)-morpholine Compound 3a2. Using the procedure of Example 46, Compound 3a2 was used in place of Compound 46a2 and carried forward to prepare 4-morpholin-4-ylmethyl-phenylamine Compound 3a.

10




Using the procedure of Example 1, Compound 3a was used in place of Compound 1n and carried forward to prepare Compound 2 which was recrystallized from methanol/EtOAc as a bis-hydrochloride salt, a yellow solid. ¹H NMR (DMSO-d₆) δ 11.30 (br s, 1 H), 10.93 (br s, 1 H), 9.78 (br s, 1 H), 8.65 (s, 1H), 7.82 (d, J = 8.4 Hz, 2 H), 7.61 (d, J = 8.4 Hz, 2 H), 7.57 (dd, J = 6.2, 2.6 Hz, 1 H), 7.54 (t, J = 9.0 Hz, 1 H), 7.36-7.31 (m, 1 H), 4.31 (s, 2 H), 3.95 (d, J = 10.9 Hz, 2 H), 3.76 (t, J = 11.8 Hz, 2 H), 3.22 (d, J = 11.8 Hz, 2 H), 3.14-3.01 (m, 2 H). MS 495, 497 (MH⁺).

20

Example 4


7-(3-chloro-4-fluoro-phenylamino)-2-(4-dimethylaminomethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 3)



Using the procedure of Example 1, dimethylamine Compound **4a1** was used in place of Compound **1n2** and carried forward to prepare dimethyl-(4-nitrobenzyl)-amine Compound **4a2**. Stannous chloride (28.1 g, 148 mmol) was added to a solution of Compound **4a2** (5.33 g, 29.6 mmol) in ethanol (200 mL) and heated to 60 °C. Sodium borohydride (0.560 g, 14.8 mmol) in ethanol (80 mL) was added dropwise. Two hrs later the reaction mixture was added to ice water. The slurry was filtered through Celite 545 and the filtrate cake rinsed with ether. The collected liquors were brought to pH 11 with 1N NaOH and extracted with ether. The combined organic layers were washed with water, dried over MgSO₄ and filtered. The dried organic solution was then reduced *in vacuo* to provide 4-dimethylaminomethyl-phenylamine Compound **4a** which was used in the next step without further purification. MS 207 (MH⁺).

10

15




Using the procedure of Example 1, Compound **4a** was used in place of Compound **1n** and carried forward to prepare Compound **3**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-d₆) δ 11.03 (br s, 1H), 9.58 (br s, 1H), 9.54 (s, 1H), 8.57 (s, 1H), 7.83 (d, J = 8.4 Hz, 2H), 7.56-7.47 (m, 4H), 7.29 (ddd, J = 8.9, 4.1, 2.8 Hz, 1H), 4.25 (s, 2H), 2.73 (s, 6H). MS 453, 455 (MH⁺).

20

Example 5


2-(4-{[bis-(2-methoxy-ethyl)-amino]-methyl}-phenylamino)-7-(3-chloro-4-fluoro-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 4)



5

Using the procedure of Example 1, bis-(2-methoxy-ethyl)-amine Compound **5a1** was used in place of Compound **1n2** and carried forward to prepare bis-(2-methoxy-ethyl)-(4-nitro-benzyl)-amine Compound **5a2**. Using the procedure of Example 46, Compound **5a2** was used in place of Compound **46a2** and carried forward to prepare 4-{[bis-(2-methoxy-ethyl)-amino]-methyl}-phenylamine Compound **5a**.

10




Using the procedure of Example 1, Compound **5a** was used in place of Compound **1n** and carried forward to prepare Compound **4**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.02 (s, 1 H), 9.53 (s, 1 H), 9.47 (br s, 1 H), 8.56 (s, 1H), 7.81 (d, J = 8.6 Hz, 2 H), 7.59-7.47 (m, 4 H), 7.29 (ddd, J = 8.7, 4.1, 2.7 Hz, 1 H), 4.36 (s, 2H), 3.75-3.53 (m, 8 H), 3.30 (s, 6H). MS 541, 534 (MH^+).


15

Example 6

4-[7-(3-chloro-4-fluoro-phenylamino)-6-cyano-thiazolo[4,5-b]pyridin-2-ylamino]-*N,N*-bis-(2-methoxy-ethyl)-benzenesulfonamide (Cpd 5)




Using the procedure of Example 1, 4-nitro-benzenesulfonyl chloride Compound **6a1**, used in place of Compound **1n1**, and bis-(2-methoxy-ethyl)-amine Compound **5a1**, used in place of Compound **1n2**, were carried forward to prepare *N,N*-bis-(2-methoxy-ethyl)-4-nitro-benzenesulfonamide Compound **6a2**. Using the procedure of Example 1, Compound **6a2** was used in place of Compound **1n3** and carried forward to prepare 4-amino-*N,N*-bis-(2-methoxy-ethyl)-benzenesulfonamide Compound **6a**.




Using the procedure of Example 1, Compound **6a** was used in place of Compound **1n** and carried forward to prepare Compound **5**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.26 (br s, 1 H), 9.60 (br s, 1 H), 8.59 (s, 1 H), 7.92 (d, J = 8.7 Hz, 2 H), 7.83 (d, J = 8.7 Hz, 2 H), 7.56-7.48 (m, 2 H), 7.31 (ddd, J = 8.7, 4.3, 2.7 Hz, 1 H), 3.41 (t, J = 5.9 Hz, 4 H), 3.27 (t, J = 5.9 Hz, 4 H), 3.20 (s, 6 H). MS 591, 593 (MH^+).

Example 7

4-[7-(3-chloro-4-fluoro-phenylamino)-6-cyano-thiazolo[4,5-b]pyridin-2-ylamino]-N-(2-morpholin-4-yl-ethyl)-benzenesulfonamide (Cpd 7)




Using the procedure of Example 1, 4-nitro-benzenesulfonyl chloride Compound **6a1**, used in place of Compound **1n1**, and 2-morpholin-4-yl-ethylamine Compound **7a1**, used in place of Compound **1n2**, were carried forward to prepare *N*-(2-morpholin-4-yl-ethyl)-4-nitro-benzenesulfonamide Compound **7a2**. Using the procedure of
 10 Example 1, Compound **7a2** was used in place of Compound **1n3** and carried forward to prepare 4-amino-*N*-(2-morpholin-4-yl-ethyl)-benzenesulfonamide Compound **7a**.




Using the procedure of Example 1, Compound **7a** was used in place of Compound **1n** and carried forward to prepare Compound **7**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 11.28 (s, 1 H), 9.84 (br s, 1 H), 9.59 (s, 1 H), 8.59 (s, 1 H), 8.01-7.82 (m, 4 H), 7.58-7.48 (m, 2 H), 7.31 (ddd, *J* = 8.9, 4.2, 2.8 Hz, 1 H), 4.10-3.91 (m, 4 H), 3.51-3.34 (m, 2 H), 3.30-3.03 (m, 6 H). MS 588, 590 (MH⁺).
 15

Example 8

7-(3-chloro-4-fluoro-phenylamino)-2-{4-[(2*S*)-2-hydroxymethyl-pyrrolidin-1-ylmethyl]-phenylamino}-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd 6)



Using the procedure of Example 1, (2*S*)-2-hydroxymethyl-pyrrolidine Compound **8a1** was used in place of Compound **1n2** and carried forward to prepare 4-[(2*S*)-2-hydroxymethyl-pyrrolidin-1-ylmethyl]-phenylamine Compound **8a**.




10 Using the procedure of Example 1, Compound **8a** was used in place of Compound **1n** and carried forward to prepare Compound **6**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 11.03 (s, 1 H), 9.54 (s, 1 H), 9.40 (br s, 1 H), 8.57 (s, 1 H), 7.81 (d, 8.5 Hz, 2 H), 7.58-7.47 (m, 4 H), 7.29 (ddd, *J* = 8.9, 4.2, 2.7 Hz, 1 H), 5.53 (br s, 1 H), 4.96 (dd, *J* = 13.0, 3.8 Hz, 1 H), 4.25 (dd, *J* = 13.0, 6.3 Hz, 1 H), 3.58 (s, 2 H), 3.35-3.10 (m, 2 H), 2.18-2.05 (m, 1 H), 2.04-1.91 (m, 1 H), 1.88-1.68 (m, 2 H). MS 509, 511 (MH⁺).

15

Example 9


7-(3-chloro-4-fluoro-phenylamino)-2-(3-morpholin-4-yl-propylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 8)



5 Using the procedure of Example 1, 3-morpholin-4-yl-propylamine Compound **9a** was used in place of Compound **1n** and carried forward to prepare Compound **8**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-d₆) δ 9.69 (br s, 1 H), 9.38 (s, 1 H), 8.76 (t, J = 5.3 Hz, 1 H), 8.48 (s, 1 H), 7.48 (t, J = 8.9 Hz, 1 H), 7.42 (dd, J = 6.5, 2.7 Hz, 1 H), 7.21 (ddd, J = 8.9, 4.1, 2.7 Hz, 1 H), 3.99 (br d, J = 12.2 Hz, 2 H),
 10 3.73-3.45 (m, 6 H), 3.22-2.95 (m, 4 H), 2.03-1.89 (m, 2 H). MS 447, 449 (MH⁺).

Example 10

7-(3-chloro-4-fluoro-phenylamino)-2-[(4-{methyl-[(2R)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 14)




15

Using the procedure of Example 1, (R)-(-)-tetrahydrofurfurylamine Compound **10a1** was used in place of Compound **1n2** and carried forward to prepare (4-nitro-benzyl)-[(2R)-tetrahydro-furan-2-ylmethyl]-amine Compound **10a2**.

Methyl iodide (0.728 mL, 11.7 mmol) was added to (4-nitro-benzyl)-[(2R)-tetrahydro-furan-2-ylmethyl]-amine Compound **10a2** (1.84 g, 7.80 mmol) and
 20 potassium carbonate (3.23 g, 23.4 mmol) in DMF (30 mL). After 4 hrs the reaction was diluted with water and extracted with ethyl ether. The combined extracts were washed with water, dried over MgSO₄, and evaporated down to give 1.43 g of methyl-

(4-nitro-benzyl)-[(2*R*)-tetrahydro-furan-2-ylmethyl]-amine Compound **10a3**. Using the procedure of Example 21, Compound **10a3** was used in place of Compound **21a2** and carried forward to prepare 4-({methyl-[(2*R*)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamine Compound **10a**.



5


Using the procedure of Example 1, Compound **10a** was used in place of Compound **1n** and carried forward to prepare Compound **14**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 9.38 (br s, 1 H), 8.41 (s, 1 H), 7.74-7.58 (m, 3 H), 7.43-7.31 (m, 3 H), 7.30-7.12 (m, 2 H), 4.34-3.99 (m, 3 H), 3.84-3.60 (m, 2 H), 3.26-2.83 (m, 2 H), 2.70 (s, 3 H), 2.05-1.91 (m, 1 H), 1.85-1.72 (m, 2 H), 1.52-1.36 (m, 1 H). MS 523, 525 (MH⁺).

10

Example 11

7-(3-chloro-4-fluoro-phenylamino)-2-[(4-{methyl-[(2*S*)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamino]-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd **15**)


15



Using the procedure of Example 1, (S)-(+)-tetrahydrofurfurylamine Compound **11a1** was used in place of Compound **1j2** and carried forward to prepare (4-nitro-benzyl)-[(2*S*)-tetrahydro-furan-2-ylmethyl]-amine Compound **11a2**. Using the procedure of Example 10, Compound **11a2**, replacing Compound **10a2**, was carried forward to give Compound **11a3**. Using the procedure of Example 21, Compound

20

11a3 was used in place of Compound **21a2** and carried forward to prepare 4-({methyl-[(2*S*)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamine Compound **11a**.




Using the procedure of Example 1, Compound **11a** was used in place of

- 5 Compound **1n** and carried forward to prepare Compound **15**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.40 (br s, 1 H), 8.80 (br s, 1 H), 8.42 (s, 1 H), 7.81-7.62 (m, 3 H), 7.42-7.32 (m, 3 H), 7.31-7.14 (m, 2 H), 4.36-3.99 (m, 3 H), 3.87-3.61 (m, 2 H), 3.26-2.83 (m, 2 H), 2.70 (s, 3 H), 2.05-1.92 (m, 1 H), 1.85-1.72 (m, 2 H), 1.54-1.37 (m, 1 H). MS 523, 525 (MH^+).

10

Example 12


7-(3-chloro-4-fluoro-phenylamino)-2-(4-{[methyl-(tetrahydro-pyran-4-yl)-amino]-methyl}-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **16**)



- 15 Using the procedure of Example 1, Compound **12a** (prepared as described in Mitsuru Shiraishi, et al., *J. Med. Chem.*, **2000**, 43, 2049-2063) was used in place of Compound **1n** and carried forward to prepare Compound **16**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.71 (br s, 1 H), 8.39 (s, 1 H), 7.93 (br s, 1 H), 7.61 (d, $J = 6.9$ Hz, 2 H), 7.41-7.15 (m, 5 H), 4.36-4.19 (m, 1 H), 4.03-3.86 (m, 3H), 3.39-3.19 (m, 3 H), 2.51 (s, 3 H), 1.98-1.63 (m, 4 H). MS 523, 525 (MH^+).
- 20

Example 13


2-(3-chloro-4-dimethylaminomethyl-phenylamino)-7-(3-chloro-4-fluoro-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 18)



5

2-chloro-4-nitrobenzoyl chloride Compound **13a1** (7.08 g, 32.2 mmol) and dimethylamine hydrochloride (2.62 g, 32.2 mmol) were combined in THF (50 mL) at 0 °C. TEA (8.97 mL, 64.4 mmol) was added and the mixture was heated at reflux for 18 hrs. When the reaction was complete, the mixture was diluted with 1N HCl and
 10 extracted with EtOAc. The product was isolated by flash chromatography (30-50% EtOAc/hexane gradient) to provide 2-chloro-N,N-dimethyl-4-nitro-benzamide Compound **13a2** (3.05 g) as a tan solid. MS 229 (MH⁺).

Using the procedure of Example 1, Compound **13a2** was used in place of Compound **1j3** and carried forward to prepare 4-amino-2-chloro-N,N-dimethyl-
 15 benzamide Compound **13a3**. 1M LAH (3.53 mL, 3.53 mmol) was added dropwise to a solution of Compound **13a3** (700 mg, 3.53 mmol) in THF (10 mL). After 10 minutes, the reaction was cooled to 0 °C and quenched with water. The mixture was extracted from ethyl ether with 1N HCl. The aqueous layer was adjusted to pH 10 with NaOH and the product was extracted with EtOAc. The combined organic layers were dried
 20 over MgSO₄ and concentrated to give 3-chloro-4-dimethylaminomethyl-phenylamine Compound **13a** (150 mg) as a viscous brown oil. MS 185 (MH⁺).




Using the procedure of Example 1, Compound **13a** was used in place of Compound **1n** and carried forward to prepare Compound **18**, which was isolated as a

trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.23 (s, 1 H), 9.69-9.56 (m, 2 H), 8.60 (s, 1 H), 8.19 (s, 1 H), 7.68 (d, J = 8.9 Hz, 1 H), 7.64 (d, J = 8.9 Hz, 1 H), 7.56 (dd, J = 6.6, 2.7 Hz, 1 H), 7.53 (t, J = 8.9 Hz, 1 H), 7.33 (ddd, J = 8.9, 4.2, 2.7 Hz, 1 H), 4.41 (s, 2H), 2.82 (s, 6 H). MS 487, 489, 491 (MH^+).


5

Example 14

7-(3-chloro-4-fluorophenylamino)-2-[4-({[(2*R*)-tetrahydrofuran-2-ylmethyl]-amino}-methyl)-phenylamino]-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd **21**)




- 10 Using the procedure of Example 21, Compound **10a2** (4-nitro-benzyl)-[(2*R*)-tetrahydro-furan-2-ylmethyl]-amine was used in place of Compound **21a2** and carried forward to prepare 4-({[(2*R*)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamine Compound **14a**.



- 15 Using the procedure of Example 1, Compound **14a** was used in place of Compound **1n** and carried forward to prepare Compound **21**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.01 (s, 1 H), 9.56 (s, 1 H), 8.91 (br s, 1 H), 8.58 (s, 1 H), 7.79 (d, J = 8.6 Hz, 2 H), 7.55-7.49 (m, 4 H), 7.30 (ddd, J = 8.8, 4.1, 2.8 Hz, 1 H), 4.21-4.06 (m, 3 H), 3.82 (dd, J = 14.8, 7.4 Hz, 1 H), 3.73 (dd, J = 14.0, 7.4 Hz, 1 H), 3.08-2.99 (m, 1 H), 2.92-2.82 (m, 1 H), 2.06-1.95 (m, 1 H), 1.91-1.80 (m, 2 H), 1.60-1.49 (m, 1 H). MS 509, 511 (MH^+).
- 20

Example 15


7-(3-chloro-4-fluoro-phenylamino)-2-(4-morpholin-4-ylmethyl-benzylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 22)



5

(4-Formylbenzyl)-carbamate tert-butyl ester Compound **15a** (555 mg, 2.37 mmol, prepared as described in Far AR, Cho YL, Rang A, Rudkewich DM, Rebek J, *Tetrahedron*, **2002**, 58(4), 741-756) was added to a solution of morpholine (207 μ L, 2.37 mmol) in MeOH (5 mL) and acetic acid (0.20 mL). The mixture was stirred for 1 hr and then 95% NaBH_3CN (198 mg, 3.0 mmol) was added. The reaction was diluted after 2 hr with saturated NaHCO_3 and extracted with EtOAc to provide (4-morpholin-4-ylmethylbenzyl)-carbamate tert-butyl ester Compound **15b** (230 mg), isolated by flash chromatography as a white solid. MS 307 (MH^+). Compound **15b** (230mg, 0.752 mmol) was deprotected with 20% TFA in DCM (10 mL) to provide 4-morpholin-4-ylmethylbenzylamine Compound **15c** (63 mg) upon work-up. MS 207 (MH^+).


15



Using the procedure of Example 1, Compound **15c** was used in place of Compound **1n** and carried forward to prepare Compound **22**, which was isolated as a trifluoroacetate salt. MS 509, 511 (MH^+).

Example 16

7-(3-chloro-4-fluoro-phenylamino)-2-(6-chloro-pyridin-3-ylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **28**)




- 5 Using the procedure of Example 1, Compound **16a** was used in place of Compound **1n** and carried forward to prepare Compound **28**, which was isolated as an acetic acid salt. ^1H NMR (DMSO- d_6) δ 11.30 (br s, 1 H), 9.67 (br s, 1 H), 8.69 (d, $J=2.9$ Hz, 1 H), 8.62 (s, 1 H), 8.32 (dd, $J = 8.8, 2.9$ Hz, 1 H), 7.61-7.49 (m, 3 H), 7.33 (ddd, $J = 8.8, 4.2, 2.8$ Hz, 1 H). MS 431, 433, 435 (MH^+).

10

Example 17


7-(3-chloro-4-fluoro-phenylamino)-2-(6-methoxy-pyridin-3-ylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **26**)



- 15 Using the procedure of Example 1, Compound **17a** was used in place of Compound **1n** and carried forward to prepare Compound **26**. ^1H NMR (DMSO- d_6) δ 10.82 (br s, 1 H), 9.45 (br s, 1 H), 8.52 (s, 1H); 8.44 (d, $J=2.5$ Hz, 1 H), 8.09 (dd, $J=9.0, 2.7$ Hz, 1 H), 7.50 (d, $J = 9.0$, 1 H), 7.50-7.45 (m, 1H), 7.26 (ddd, $J = 9.0, 4.3, 2.7$ Hz, 1 H); 6.88 (d, $J=9.0$ Hz, 1 H); 3.84 (s, 3H). MS 427, 429 (MH^+).

Example 18


7-(3-chloro-4-fluoro-phenylamino)-2-(6-hydroxy-pyridin-3-ylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **27**)



5 Compound **26** was heated with aqueous HCl in NMP (3 mL) at 80 °C to prepare Compound **27**, which was isolated as a trifluoroacetic acid salt. ¹H NMR (DMSO-d₆) δ 11.52 (br s, 1 H), 10.52 (br s, 1 H), 9.43 (s, 1H); 8.50 (s, 1H); 8.09 (d, J=2.5 Hz, 1 H), 7.52-7.42 (m, 3 H), 7.24 (ddd, J = 8.7, 4.1, 2.9 Hz, 1 H); 6.41 (d, J =9.4 Hz, 1 H). MS 413, 415 (MH⁺).


Example 19

7-(3-chloro-4-fluoro-phenylamino)-2-[6-(3-morpholin-4-yl-propylamino)-pyridin-3-ylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **29**)



2-chloro-5-nitro-pyridine Compound **16a1** (6.06 g, 38.2 mmol) and 3-morpholin-4-yl-propylamine Compound **9a** (6.27 mL, 42.1 mmol) were combined in DMSO (100 mL) and heated to 50°C. After 2.5 hours the reaction mixture was diluted with 0.5 M HCl and washed with ethyl ether. The aqueous layer was adjusted to pH 11 with NaOH and extracted with EtOAc. The combined organic layers were dried over MgSO₄ and concentrated to provide (3-morpholin-4-yl-propyl)-(5-nitro-pyridin-2-yl)-amine Compound **19a1**. Using the procedure of Example 1, Compound **19a1** was used in place of Compound **1n3** with AcOH (30 mL) replacing EtOAc as solvent. 10% Pd/C (3.1 g) was added to the solution and the reaction was carried forward to provide N²-(3-


morpholin-4-yl-propyl)-pyridine-2,5-diamine Compound **19a**. ^1H NMR (CDCl_3) δ 7.67 (d, $J=3.0$ Hz, 1H); 6.95 (dd, $J=8.6$, 3.0 Hz, 1H); 6.31(d, $J=8.6$ Hz, 1H); 3.77-3.66 (m, 4H); 2.52-2.38 (m, 8H); 2.08 (p, $J=6.8\text{Hz}$, 2H). MS 267 (MH^+).



- 5 Using the procedure of Example 1, Compound **19a** was used in place of Compound **1n** and carried forward to prepare Compound **29**, which was isolated as a trifluoroacetic acid salt. ^1H NMR (DMSO-d_6) δ 10.80 (br s, 1 H), 9.50 (s, 1 H), 8.53 (s, 1 H), 8.44 (br s, 1 H), 7.79 (dd, $J = 9.0$, 2.3 Hz, 1 H), 7.53-7.44 (m, 2 H), 7.25 (ddd, $J = 8.8$, 4.2, 2.8 Hz, 1 H); 6.79 (d, $J=9.0$ Hz, 1H); 3.89-3.74 (m, 4H); 3.34 (t, $J=6.4$ Hz, 10 2H); 3.28-3.13 (m, 6H); 2.00-1.88 (m, 2H). MS 539, 541 (MH^+).


Example 20

2-[6-(2-amino-ethylamino)-pyridin-3-ylamino]-7-(3-chloro-4-fluoro-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **30**)



15


Using the procedure of Example 1, N^1 -(5-nitro-pyridin-2-yl)-ethane-1,2-diamine Compound **20a1** was used in place of Compound **1n3** and carried forward to prepare N^2 -(2-amino-ethyl)-pyridine-2,5-diamine Compound **20a**.




Using the procedure of Example 1, Compound **20a** was used in place of Compound **1n** and carried forward to prepare Compound **30**, which was isolated as a trifluoroacetic acid salt. ^1H NMR (DMSO- d_6) δ 10.86 (br s, 1 H), 9.58 (s, 1 H), 8.57 (s, 1 H), 8.44 (s, 1 H), 7.95-7.80 (m, 3 H), 7.54-7.46 (m, 2 H), 7.28 (ddd, J = 8.7, 4.1, 2.8 Hz, 1 H); 6.80 (d, J = 8.9 Hz, 1 H); 3.52 (t, J = 5.8 Hz, 2 H); 3.04 (q, J = 5.8 Hz, 2 H). MS 455, 457 (MH^+).

Example 21

2-[4-(4-amino-piperidin-1-ylmethyl)-phenylamino]-7-(3-chloro-4-fluoro-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **31**)




Using the procedure of Example 1, piperidin-4-ylamine Compound **21a1** was used in place of Compound **1n2** and carried forward to prepare 1-(4-nitro-benzyl)-piperidin-4-ylamine Compound **21a2**. 5% Rh/C (350 mg) was added to a solution of 1-(4-nitrobenzyl)-piperidin-4-ylamine Compound **21a2** (647 mg, 2.75 mmol) in methanol (15 mL). The mixture was hydrogenated at 50 PSI for a period of 2 hrs, then filtered through Celite. The filtrate was evaporated *in vacuo* to give 1-(4-amino-benzyl)-piperidin-4-ylamine Compound **21a** (640 mg) as a light yellow oil. ^1H NMR (CDCl_3) δ 7.09 (d, J = 8.2 Hz, 2H); 6.64 (d, J = 8.2 Hz, 2H); 3.62 (br s, 1H); 3.41 (s, 2H); 2.88-2.80 (m, 2H); 2.70-2.58 (m, 1H); 2.07-1.95 (m, 2H); 1.84-1.74 (m, 2H); 1.57 (br s, 2H); 1.47-1.32 (m, 2H); 1.31-1.18 (m, 2H).




Using the procedure of Example 1, Compound **21a** was used in place of Compound **1n** and carried forward to prepare Compound **31**, which was isolated as a trifluoroacetic acid salt. ^1H NMR (DMSO- d_6) δ 11.07(br s, 1 H), 9.88 (br s, 1 H), 9.56 (s, 1 H), 8.57 (s, 1 H), 8.20-8.07 (m, 3 H), 7.82 (d, $J=8.3$ Hz, 2H); 7.55-7.45 (m, 4 H), 7.29 (ddd, $J = 8.8, 4.3, 2.7$ Hz, 1 H); 4.29 (s, 2 H); 3.44 (d, $J=11.2$ Hz, 2H); 3.34-3.16 (m, 1 H); 3.08-2.93 (m, 2H); 2.15-1.99 (m, 2H); 1.81-1.65 (m, 2H). MS 508, 510 (MH^+).

Example 22

7-(3-ethynyl-phenylamino)-2-(3-morpholin-4-yl-propylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **9**)



Using the procedure of Example 1, 3-ethynyl-phenylamine Compound **22a** was used in place of Compound **1l** and carried forward to prepare 7-(3-ethynyl-phenylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **22b**.




Using the procedure of Example 1, Compound **22b**, used in place of Compound **1m**, and Compound **9a**, used in place of Compound **1n**, were carried forward to prepare Compound **9**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.67

(br s, 1 H), 9.47 (s, 1 H), 8.82 (br s, 1 H), 8.52 (s, 1 H), 7.52-7.32 (m, 2 H), 7.26-7.17 (m, 2 H), 4.27 (s, 1 H), 3.78-3.58 (m, 4 H), 3.54-3.38 (m, 4 H), 3.22-3.00 (m, 4 H), 2.04-1.90 (m, 2 H). MS 419 (MH^+).

Example 23

5

2-(4-dimethylaminomethyl-phenylamino)-7-(3-ethynyl-phenylamino)thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 10)




Using the procedure of Example 1, Compound **22b**, used in place of Compound **1m**, and Compound **4a**, used in place of Compound **1n**, were carried forward to prepare Compound **10**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.06 (s, 1 H), 9.67 (br s, 1 H), 9.57 (s, 1 H), 8.59 (s, 1 H), 7.82 (d, $J = 8.2$ Hz, 2 H), 7.50 (d, $J = 8.2$ Hz, 2 H), 7.46 (t, $J = 7.7$ Hz, 1 H), 7.39 (d, $J = 7.7$ Hz, 1 H), 7.30 (s, 1 H), 7.27 (d, $J = 7.7$ Hz, 1 H), 4.28 (s, 1 H), 4.25 (s, 2 H), 2.73 (s, 6 H). MS 425 (MH^+).

Example 24

15

7-(3-ethynyl-phenylamino)-2-{4-[(2S)-2-hydroxymethyl-pyrrolidin-1-ylmethyl]-phenylamino}-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 11)




Using the procedure of Example 1, Compound **22b**, used in place of Compound **1m**, and Compound **8a**, used in place of Compound **1n**, were carried forward to prepare Compound **11**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.06 (s, 1 H), 9.59 (s, 1 H), 9.43 (br s, 1 H), 8.59 (s, 1 H), 7.80 (d, $J = 7.9$ Hz, 2 H), 7.54 (d, $J = 7.9$ Hz, 2 H), 7.45 (t, $J = 7.7$ Hz, 1 H), 7.38 (dd, $J = 7.7, 1.2$ Hz, 1 H), 7.29

(s, 1 H), 7.26 (d, $J = 7.7$ Hz, 1 H), 4.53-4.21 (m, 7 H), 3.33-3.23 (m, 1 H), 3.22-3.12 (m, 1 H), 2.17-2.05 (m, 1 H), 2.02-1.91 (m, 1 H), 1.88-1.69 (m, 2 H). MS 481 (MH^+).

Example 25

7-(3-ethynyl-phenylamino)-2-(2-morpholin-4-yl-ethylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 13)

5




Using the procedure of Example 1, Compound **22b**, used in place of Compound **1m**, and Compound **7a1**, used in place of Compound **1n**, were carried forward to prepare Compound **13**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.78 (br s, 1 H), 9.45 (s, 1 H), 8.86 (t, $J = 4.9$ Hz, 1 H), 8.53 (s, 1 H), 7.43 (t, $J = 8.0$ Hz, 1 H), 7.36 (d, $J = 8.0$ Hz, 1 H), 7.22 (s, 1 H), 7.21 (d, $J = 8.0$ Hz, 1 H), 4.28 (s, 1 H), 4.06-3.93 (m, 2 H), 3.83-3.74 (m, 2 H), 3.73-3.49 (m, 4 H), 3.40-3.31 (m, 2 H), 3.22-3.06 (m, 2 H). MS 405 (MH^+).


15

Example 26

4-[6-cyano-7-(3-ethynyl-phenylamino)-thiazolo[4,5-b]pyridin-2-ylamino]-N-(3-dimethylamino-propyl)-benzenesulfonamide (Cpd 23)




Using the procedure of Example 1, 4-nitro-benzenesulfonyl chloride Compound **6a1**, used in place of Compound **1n1**, and N^1,N^1 -dimethyl-propane-1,3-diamine Compound **26a1**, used in place of Compound **1n2**, were carried forward to prepare 4-amino- N -(3-dimethylamino-propyl)-benzenesulfonamide Compound **26a**.




Using the procedure of Example 1, Compound **22b**, used in place of Compound **1m**, and Compound **26a**, used in place of Compound **1n**, were carried forward to prepare Compound **23**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-
 5 d₆) δ 11.28 (s, 1 H), 9.61 (s, 1 H), 9.37 (br s, 1 H), 8.63 (s, 1 H), 7.97 (d, J = 8.9 Hz, 2 H), 7.84 (d, J = 8.9 Hz, 2 H), 7.69 (t, J = 6.1 Hz, 1 H), 7.53-7.40 (m, 2 H), 7.36-7.28 (m, 2 H), 4.30 (s, 1 H), 3.14-3.03 (m, 2 H), 2.89-2.75 (m, 8 H), 1.85-1.73 (m, 2 H). MS 532 (MH⁺).

Example 27

10 7-(3-chloro-phenylamino)-2-(2-morpholin-4-yl-ethylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **12**)



Using the procedure of Example 1, 3-chloroaniline Compound **27a** was used in place of Compound **1l** and carried forward to prepare 7-(3-chloro-phenylamino)-2-methylsulfanylstiazo[4,5-b]pyridine-6-carbonitrile Compound **27b**.
 15




Using the procedure of Example 1, Compound **27b**, used in place of Compound **1m**, and Compound **7a1**, used in place of Compound **1n**, were carried forward to

prepare Compound **12**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.91 (br s, 1 H), 9.54 (s, 1 H), 8.91 (br s, 1 H), 8.54 (s, 1 H), 7.43 (t, $J = 7.8$ Hz, 1 H), 7.29 (d, $J = 7.8$ Hz, 1 H), 7.20 (t, $J = 1.8$ Hz, 1 H), 7.11 (d, $J = 7.8$ Hz, 1 H), 4.08-3.90 (m, 2 H), 3.84-3.74 (m, 2 H), 3.71-3.33 (m, 6 H), 3.23-3.05 (m, 2 H). MS 415, 417 (MH^+).

Example 28


7-(3-chloro-phenylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **19**)



Using the procedure of Example 1, Compound **27b**, used in place of Compound **1m**, and Compound **3a**, used in place of Compound **1n**, were carried forward to prepare Compound **19**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.11 (s, 1 H), 10.03 (br s, 1 H), 9.65 (s, 1 H), 8.61 (s, 1 H), 7.83 (d, $J = 7.6$ Hz, 2 H), 7.52 (d, $J = 7.6$ Hz, 2 H), 7.45 (t, $J = 8.0$ Hz, 1 H), 7.32 (d, $J = 8.0$ Hz, 1 H), 7.28 (s, 1 H), 7.18 (d, $J = 8.0$ Hz, 1 H), 4.33 (s, 2 H), 4.04-3.92 (m, 2 H), 3.73-3.57 (m, 2 H), 3.33-3.22 (m, 2 H), 3.18-3.04 (m, 2 H). MS 477, 479 (MH^+).


Example 29

7-(2,2-difluoro-benzo[1,3]dioxol-4-ylamino)-2-(4-dimethylaminomethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **25**)



Using the procedure of Example 1, 2,2-difluoro-benzo[1,3]dioxol-4-ylamine Compound **29a** was used in place of Compound **11** and carried forward to prepare 7-

(2,2-difluoro-benzo[1,3]dioxol-4-ylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **29b**.




Using the procedure of Example 1, Compound **29b**, used in place of Compound

- 5 **1m**, and Compound **7a1**, used in place of Compound **1n**, were carried forward to prepare Compound **25**, which was isolated as a hydrochloride salt. ¹H NMR (DMSO-d₆) δ 11.42 (br s, 1 H), 10.46 (br s, 1 H), 9.97 (br s, 1 H), 8.66 (s, 1 H), 7.84 (d, J = 8.4 Hz, 2H), 7.58 (d, J = 8.4 Hz, 2 H), 7.43 (d, J = 8.2 Hz, 1 H), 7.32 (t, J = 8.2 Hz, 1 H), 7.21 (d, J=8.2 Hz, 1 H), 4.25 (s, 2 H), 2.70 (s, 6 H). MS 481 (MH⁺).

10


Example 30

7-[3-chloro-4-(3-fluorobenzyloxy)-phenylamino]-2-(4-pyrrolidin-1-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **24**)




15

1-Fluoro-3-(4-nitro-2-chlorophenoxy)methylbenzene Compound **30c** was prepared according to the procedure described in Cockerill GS, Carter MC, Guntrip SB and Smith KJ, Preparation of azolylquinazolines and related compounds as protein tyrosine kinase inhibitors, PCT Application WO98/02434.




20

Using the procedure of Example 21, Compound **30c** was used in place of Compound **21a2** and carried forward to prepare 3-chloro-4-(3-fluorobenzyloxy)-phenylamine Compound **30d**.



Using the procedure of Example 1, Compound **30d** was used in place of Compound **1l** and carried forward to prepare 7-[3-chloro-4-(3-fluorobenzoyloxy)-phenylamino]-2-methylsulfanylnicotinamide Compound **30e**.




5

Using the procedure of Example 1, Compound **30e** was used in place of Compound **1m** and carried forward to prepare Compound **24**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.0 (s, 1 H), 9.80 (br s, 1 H), 9.38 (s, 1 H), 8.54 (s, 1 H), 7.82 (d, $J = 8.6$ Hz, 2 H), 7.56-7.48 (m, 3 H), 7.46 (d, $J = 2.2$ Hz, 1 H), 7.41-7.19 (m, 5 H), 5.31 (s, 2 H), 4.33 (s, 2 H), 3.46-3.31 (m, 2 H), 3.18-3.03 (m, 2 H), 2.12-1.78 (m, 4 H). MS 585, 587 (MH^+).


10

Example 31

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile
(Cpd 20)



1M sodium hexamethyldisilazane in THF (1.42 mL, 1.42 mmol) was added to 5-chloro-benzo[1,3]dioxol-4-ylamine Compound **31a** (as described in PCT Application WO03/008409) (221 mg, 1.29 mmol) in DMF at 0°C and the mixture was stirred for 15 min. 7-chloro-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **1g** (312 mg, 1.29 mmol) in DMF (1 mL) was added and the reaction mixture stirred for an additional 2 hr at ambient temperature. The mixture was then diluted with water and filtered through Celite 545 to form a plug. The celite plug was rinsed with EtOAc and the organic layer was isolated and washed with brine. The organic layer was dried over MgSO₄, then filtered and evaporated onto silica gel. Isolation by flash chromatography (eluted with a 20 to 40 % EtOAc/hexane gradient) gave 7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **31b** (30 mg) as a yellow oil. MS 377, 379 (MH⁺).




Using the procedure of Example 1, Compound **31b**, replacing Compound **1m**, and Compound **3a**, replacing Compound **1n**, were carried forward to prepare Compound **20** as a trifluoroacetate salt. ¹H NMR (DMSO-d₆) δ 11.05 (s, 1 H), 9.92 (br s, 1 H), 9.31 (s, 1 H), 8.52 (s, 1 H), 7.82 (d, J = 8.5 Hz, 2H), 7.51 (d, J = 8.5 Hz, 2 H), 7.14 (d, J = 8.3 Hz, 1 H), 7.11 (d, J = 8.3 Hz, 1 H), 6.13 (s, 2 H), 4.32 (s, 2 H), 4.03-

3.91 (m, 2H), 3.68-3.58 (m, 2 H), 3.32-3.23 (m, 2 H), 3.17-3.05 (m, 2 H). MS 521, 523 (MH⁺).

Example 32


7-(6-bromo-benzo[1,3]dioxol-4-ylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 32)

5



Using the procedure of Example 31, 6-bromo-benzo[1,3]dioxol-4-ylamine Compound 32a (prepared as described in PCT Application WO03/008409) was used in place of Compound 31a and carried forward to prepare 7-(6-bromo-benzo[1,3]dioxol-4-ylamino)-2-methylsulfonyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound 32b.

10




Using the procedure of Example 1, Compound 32b, replacing Compound 1m, and Compound 3a, replacing Compound 1n, were carried forward to prepare Compound 32 as a trifluoroacetate salt. ¹H NMR (DMSO-d₆) δ 11.08 (s, 1 H), 9.84 (br s, 1 H), 9.49 (s, 1 H), 8.55 (s, 1 H), 7.84 (d, 2H), 7.50 (d, 2 H), 7.23 (d, 1 H), 7.04 (d, 1 H), 6.04 (s, 2 H), 4.32 (s, 2 H), 4.04-3.91 (m, 2H), 3.71-3.55 (m, 2 H), 3.36-3.01 (m, 4 H). MS 565, 567 (MH⁺).

15

Example 33

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-*p*-tolylamino-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd **33**)




5 Using the procedure of Example 1, Compound **31b**, replacing Compound **1m**, and *p*-tolylamine Compound **33a**, replacing Compound **1n**, were carried forward to prepare Compound **33** as a trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 10.75 (s, 1 H), 9.18 (s, 1 H), 8.44 (s, 1 H), 7.59 (d, 2H), 7.17 (d, 2 H), 7.11 (d, 1 H), 7.08 (d, 1 H), 6.11 (s, 2 H), 2.29 (s, 3 H). MS 436, 438 (MH⁺).


10

Example 34

7-(2,4-dichloro-phenylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd **34**)



15 Using the procedure of Example 31, 2,4-dichloroaniline Compound **34a** was used in place of Compound **31a** and carried forward to prepare 7-(2,4-dichlorophenylamino)-2-methylsulfanylmethylthiazolo[4,5-*b*]pyridine-6-carbonitrile Compound **34b**.




20 Using the procedure of Example 1, Compound **34b**, replacing Compound **1m**, and Compound **3a**, replacing Compound **1n**, were carried forward to prepare

Compound **34** as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 9.94 (br s, 1 H), 9.48 (s, 1 H), 8.53 (s, 1 H), 7.84 (s, 1H), 7.81 (d, $J = 8.0$ Hz, 2 H), 7.57-7.53 (m, 2 H), 7.50 (d, $J = 8.0$ Hz, 1 H), 4.08-3.91 (m, 2H), 3.70-3.54 (m, 2 H), 3.32-3.01 (m, 4 H). MS 511,513 (MH^+).


5

Example 35


7-(2-chloro-5-methoxy-phenylamino)-2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **35**)



10 1 M potassium t-butoxide (50.6 mL) was added dropwise to 4-nitrophenol Compound **35a** (7.03 g, 50.6 mmol) in DMF (100 mL) at 0 °C. After 20 mins, 3-chloromethyl-1-methylpiperidine hydrochloride Compound **35b** (9.31 g, 50.6 mmol, neutralized prior to addition) in DMF (10 mL) was added to the reaction mixture. The reaction was then heated at 100 °C for 5 days. The cooled reaction was diluted with 1 N NaOH and washed with ethyl ether. The aqueous layer was then adjusted to pH 5
15 with NaH_2PO_4 and extracted with ethyl acetate. After drying over MgSO_4 , the organic layer was evaporated down to give 6.0 g of 1-methyl-3-(4-nitro-phenoxy)methylpiperidine Compound **35c**. Using the procedure of Example 1, Compound **35c** was used in place of Compound **1n3** and carried forward to prepare 4-(1-methyl-piperidin-
20 3-ylmethoxy)-phenylamine Compound **35d**.



Using the procedure of Example 31, 2-chloro-5-methoxyaniline Compound **35e** was used in place of Compound **31a** and carried forward to prepare 7-(2-chloro-5-methoxy-phenylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile
25 Compound **35f**.




Using the procedure of Example 1, Compound **35f**, replacing Compound **1m**, and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **35** as a trifluoroacetate salt. MS 535, 537 (MH^+).

5


Example 36

7-(2,4-dichloro-phenoxy)-2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **36**)



10

Using the procedure of Example 31, 2,4-dichlorophenol Compound **36a** was used in place of Compound **31a** and carried forward to prepare 2-methylsulfanyl-7-(3-trifluoromethoxy-phenoxy)-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **36b**.




Using the procedure of Example 1, Compound **34b**, replacing Compound **1m**, and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **36** as a trifluoroacetate salt. MS 540, 542, 544 (MH^+).

15


Example 37

2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-7-(3-trifluoromethoxy-phenoxy)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **37**)

5



Using the procedure of Example 31, 3-trifluoromethoxyphenol Compound **37a** was used in place of Compound **31a** and carried forward to prepare 2-methylsulfanyl-7-(3-trifluoromethoxy-phenoxy)-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **37b**.




10

Using the procedure of Example 1, Compound **37b**, replacing Compound **1m**, and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **37** as a trifluoroacetate salt. MS 556 (MH^+).

Example 38


7-(3-bromo-phenylsulfanyl)-2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **38**)

15



Using the procedure of Example 31, 3-bromobenzenethiol Compound **38a** was used in place of Compound **31a** and carried forward to prepare 7-(3-bromo-


phenylsulfanyl)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **38b**.




Using the procedure of Example 1, Compound **38b**, replacing Compound **1m**, and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **38** as a trifluoroacetate salt. MS 566, 568 (MH⁺).

Example 39

7-(2-chloro-4-fluorophenylsulfanyl)-2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **39**)




Using the procedure of Example 31, 2-chloro-4-fluorobenzenethiol Compound **39a** was used in place of Compound **31a** and carried forward to prepare 7-(2-chloro-4-fluorophenylsulfanyl)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **39b**.



Using the procedure of Example 1, Compound **39b**, replacing Compound **1m**, and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **38** as a trifluoroacetate salt. MS 540, 542 (MH^+).

Example 40


5 7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[4-(1-methyl-piperidin-3-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **40**)



Using the procedure of Example 1, Compound **31b**, replacing Compound **1m**,
 10 and Compound **35d**, replacing Compound **1n**, were carried forward to prepare Compound **40** as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 10.73 (s, 1 H), 9.35 (br s, 1 H), 9.22 (s, 1 H), 8.46 (s, 1 H), 7.62 (d, $J = 8.7$ Hz, 2 H), 7.11 (d, $J = 8.1$ Hz, 1 H), 7.08 (d, $J = 8.1$ Hz, 1 H), 6.10 (s, 2H), 3.99-3.93 (m, 1 H), 3.85-3.78 (m, 1 H) 3.62-3.39 (m, 2 H), 2.90-2.77 (m, 5 H), 2.26-2.14 (m, 1 H), 1.94-1.61 (m, 3 H), 1.33-1.20 (m, 1
 15 H). MS 549, 551 (MH^+).


Example 41

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-(4-piperidin-1-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **41**)



20


Using the procedure of Example 1, piperidine Compound **41a1** was used in place of Compound **1n2** and was carried forward to prepare 4-piperidin-1-ylmethyl-phenylamine Compound **41a**.



Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **41a**, used in place of Compound **1n**, were carried forward to prepare Compound **41** which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.05(s, 1 H), 9.40-9.28 (m, 2 H), 8.50 (s, 1 H), 7.80 (d, J = 8.6 Hz, 2 H), 7.49 (d, J = 8.6 Hz, 2 H), 7.13 (d, J = 8.5 Hz, 1 H), 7.10 (d, J = 8.5 Hz, 1 H), 6.11 (s, 2H), 4.24 (s, 2 H), 3.37-3.27 (m, 2 H) 2.92-2.78 (m, 2 H), 1.87-1.76 (m, 2 H), 1.73-1.53 (m, 3 H), 1.42-1.28 (m, 1 H). MS 519, 521 (MH^+).

Example 42


7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[3-methoxy-4-(2-morpholin-4-yl-ethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **42**)



A mixture of 2-methoxy-4-nitrophenol Compound **42a1** (1.0 g, 5.4 mmol), potassium carbonate (1.5 g, 10.9 mmol) and 4-(2-chloroethyl)-morpholine hydrochloride Compound **42a2** (1.05 g, 5.64 mmol) in DMF (10 mL) was heated under microwave conditions to 140 °C for 30 minutes. The reaction mixture was filtered, diluted with EtOAc and subsequently washed with water, then brine. The organic layer was dried over sodium sulfate and concentrated to yield 4-[2-(2-methoxy-4-nitrophenoxy)-ethyl]-morpholine Compound **42a3** (1.4 g). MS 283 (MH^+).

A solution of 10% Pd/C (0.25 g) was added to a solution of 4-[2-(2-methoxy-4-nitrophenoxy)-ethyl]-morpholine Compound **42a3** (1.4g, 5.0 mmol) in EtOAc (50 mL). The solution was exposed to 40 PSI of hydrogen gas for a period of 1 hr. The


catalyst was removed by filtration through Celite 545 and the solvent was removed to give 3-methoxy-4-(2-morpholin-4-yl-ethoxy)-phenylamine Compound **42a** (1.1 g). MS 253 (MH⁺).




- 5 Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **42a**, used in place of Compound **1n**, were carried forward to prepare Compound **42**, which was isolated as a trifluoroacetate salt. MS 581, 583 (MH⁺).

Example 43

- 10 7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[3-methoxy-4-(tetrahydro-pyran-2-ylmethoxy)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **43**)




- 15 Using the procedure of Example 42, 2-chloromethyl-tetrahydropyran Compound **43a1** was used in place of Compound **42a2** and carried forward to provide 2-(2-methoxy-4-nitro-phenoxy-methyl)-tetrahydro-pyran Compound **43a2**. Using the procedure of Example 1, Compound **43a2** was used in place of Compound **1n3** and carried forward to prepare 3-methoxy-4-(tetrahydro-pyran-2-ylmethoxy)-phenylamine Compound **43a**.



Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **43a**, used in place of Compound **1n**, were carried forward to prepare Compound **43**, which was isolated as a trifluoroacetate salt. MS 566, 568 (MH⁺).

Example 44


7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[4-({[(2*R*)-tetrahydro-furan-2-ylmethyl]-amino}-methyl)-phenylamino]-thiazolo[4,5-*b*]pyridine-6-carbonitrile (Cpd **44**)



Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **14a**, used in place of Compound **1n**, were carried forward to prepare Compound **44**, which was isolated as a trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 11.00 (s, 1 H), 9.30 (s, 1 H), 8.91 (br s, 1 H), 8.50 (s, 1 H), 7.77 (d, *J* = 8.6 Hz, 2 H), 7.49 (d, *J* = 8.6 Hz, 2 H), 7.12 (d, *J* = 8.4 Hz, 1 H), 7.10 (d, *J* = 8.4 Hz, 1 H), 6.11 (s, 2H), 4.20-3.67 (m, 5 H) 3.07-2.97 (m, 1 H), 2.91-2.79 (m, 1 H), 2.05-1.94 (m, 1 H), 1.91-1.78 (m, 2 H), 1.59-1.47 (m, 1 H). MS 535, 537 (MH⁺).

Example 45

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-(4-pyrrolidin-1-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 45)




Using the procedure of Example 1, Compound **31b** was used in place of Compound **1m** and carried forward to prepare Compound **45**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.04 (s, 1 H), 9.97 (br s, 1 H), 9.32 (s, 1 H), 8.51 (s, 1 H), 7.80 (d, J = 8.4 Hz, 2 H), 7.51 (d, J = 8.4 Hz, 2 H), 7.12 (d, J = 8.2 Hz, 1 H), 7.10 (d, J = 8.2 Hz, 1 H), 6.12 (s, 2H), 4.32 (s, 2 H) 3.42-3.31 (m, 2 H), 3.14-3.02 (m, 2 H), 2.09-1.97 (m, 2 H), 1.92-1.79 (m, 2 H). MS 505, 507 (MH^+).

10

Example 46

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[4-(4-methyl-piperazin-1-ylmethyl)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 46)


15



Using the procedure of Example 1, 1-methyl-piperazine Compound **46a1** was used in place of Compound **1n2** and carried forward to prepare 1-methyl-4-(4-nitrobenzyl)-piperazine Compound **46a2**. Nickel chloride hexahydrate (45.7 g, 192 mmol) was added to a solution of Compound **46a2** (17.9 g, 87.4 mmol) in methanol (250 mL). Twenty portions of sodium borohydride (500 mg each, 264 mmol total) were then added to the reaction mixture over a period of about 2 hr. The mixture was then carefully diluted at 0°C with concentrated HCl to give a transparent green solution. The mixture was subsequently washed with ethyl ether. The cooled aqueous layer was

20


adjusted to pH 10 with NH_4OH (28%) and then extracted with EtOAc. The combined organic layers were dried over MgSO_4 , then filtered and evaporated *in vacuo* to give 4-(4-methyl-piperazin-1-ylmethyl)-phenylamine Compound **46a** (17.6 g) as a yellow foam. ^1H NMR ($\text{DMSO}-d_6$) δ 6.91 (d, $J=8.2$ Hz, 2H); 6.50 (d, $J=8.2$ Hz, 2H); 4.94 (br s, 2H); 3.28 (s, 2H); 2.44-2.26 (m, 8H); 2.19 (s, 3H).




Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **46a**, used in place of Compound **1n**, were carried forward to prepare Compound **46**, which was isolated as a trifluoroacetate salt. MS 534, 536 (MH^+).

Example 47

7-(2,4-dichloro-5-methoxy-phenylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **47**)




Using the procedure of Example 31, 2,4-dichloro-5-methoxyaniline Compound **47a** was used in place of Compound **31a** and was carried forward to prepare 7-(2,4-dichloro-5-methoxy-phenylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **47b**.




Using the procedure of Example 1, Compound **47b**, used in place of Compound **1m**, and Compound **3a**, used in place of Compound **1n**, were carried forward to prepare Compound **47**, which was isolated as a trifluoroacetic acid salt. ^1H NMR (DMSO- d_6) δ 11.02 (s, 1 H), 10.02 (br s, 1 H), 9.55 (s, 1 H), 8.54 (s, 1 H), 7.84-7.78 (m, 3 H), 7.50 (d, $J = 8.4$ Hz, 2 H), 7.35 (s, 1 H), 4.31 (s, 2H), 4.01-3.82 (m, 5 H), 3.67-3.57 (m, 2 H), 3.31-3.20 (m, 2 H), 3.16-3.03 (m, 2 H). MS 541, 543 (MH^+).

Example 48

7-(3-bromo-phenylamino)-2-(4-morpholin-4-ylmethyl-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd **48**)



Using the procedure of Example 1, 3-bromo-phenylamine Compound **48a** was used in place of Compound **1l** and was carried forward to prepare 7-(3-bromo-phenylamino)-2-methylsulfanyl-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **48b**.




Using the procedure of Example 1, Compound **48a**, used in place of Compound **1m**, and Compound **3a**, used in place of Compound **1n**, were carried forward to prepare Compound **48**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 11.07 (s, 1 H), 9.90 (br s, 1 H), 9.61 (s, 1 H), 8.59 (s, 1 H), 7.82 (d, $J = 8.6$ Hz, 2 H),

7.51 (d, $J = 8.6$ Hz, 2 H), 7.47-7.33 (m, 3 H), 7.20 (d, $J = 8.1$ Hz, 1 H), 4.32 (s, 2H), 4.02-3.91 (m, 2 H), 3.67-3.53 (m, 2 H), 3.31-3.20 (m, 2 H), 3.18-3.02 (m, 2 H). MS 521, 523 (MH^+).


Example 49

5

7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[4-(2-morpholin-4-yl-ethyl)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile (Cpd 49)



Using the procedure of Example 1, 1-(2-chloroethyl)-4-nitrobenzene Compound 10 **49a1** was used in place of Compound **1n1** and morpholine Compound **3a1** was used in place of Compound **1n2** to prepare 4-[2-(4-nitrophenyl)ethyl]-morpholine Compound **49a2**, which was carried forward to prepare 4-(2-morpholin-4-yl-ethyl)-phenylamine Compound **49a**.




15 Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **49a**, used in place of Compound **1n**, were carried forward to prepare Compound **49**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 10.87 (s, 1 H), 9.81 (br s, 1 H), 9.26 (s, 1 H), 8.49 (s, 1 H), 7.69 (d, $J = 8.1$ Hz, 2 H), 7.28 (d, $J = 8.1$ Hz, 2 H), 7.11 (s, 2 H), 6.11 (s, 2H), 4.08-3.95 (m, 2 H) 3.73-3.59 (m, 2 H), 3.56-3.44 (m, 2 H), 3.40-3.29 (m, 2 H), 3.20-3.04 (m, 2 H) 3.01-2.90 (m, 2 H). MS 535,537 (MH^+).

20

Example 50


7-(5-chloro-benzo[1,3]dioxol-4-ylamino)-2-[4-(2-piperidin-1-yl-ethyl)-phenylamino]-thiazolo[4,5-b]pyridine-6-carbonitrile
(Cpd 50)

5



Using the procedure of Example 1, Compound **49a1** was used in place of Compound **1n1** and piperidine Compound **41a1** was used in place of Compound **1n2** to prepare 1-[2-(4-nitro-phenyl)-ethyl]-piperidine Compound **50a1**, which was carried forward to prepare 4-(2-piperidin-1-yl-ethyl)-phenylamine Compound **50a**.


10




Using the procedure of Example 1, Compound **31b**, used in place of Compound **1m**, and Compound **50a**, used in place of Compound **1n**, were carried forward to prepare Compound **50**, which was isolated as a trifluoroacetate salt. ^1H NMR (DMSO- d_6) δ 10.87 (s, 1 H), 9.31-9.15 (m, 2 H), 8.48 (s, 1 H), 7.69 (d, $J = 8.5$ Hz, 2 H), 7.28 (d, $J = 8.5$ Hz, 2 H), 7.10 (s, 2 H), 6.11 (s, 2H), 3.57-3.46 (m, 2 H) 3.31-3.19 (m, 2 H), 3.00-2.84 (m, 4 H), 1.90-1.78 (m, 2 H), 1.76-1.55 (m, 3 H) 1.46-1.30 (m, 1 H),. MS 533,535 (MH^+).

Example 51

2-[4-(4-methyl-piperazin-1-ylmethyl)-phenylamino]-7-(4-phenoxy-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile
(Cpd **51**)



The procedure of Example 1 and 4-phenoxy-phenylamine Compound **51a** used in place of 3-chloro-4-fluoro-phenylamine Compound **1l** were used to prepare 2-methylsulfanyl-7-(4-phenoxy-phenylamino)-thiazolo[4,5-b]pyridine-6-carbonitrile Compound **51b**.



The procedure of Example 1, Compound **51b** used in place of Compound **1m**, and 4-(4-methyl-piperazin-1-ylmethyl)-phenylamine Compound **46a** used in place of Compound **1n** were used to prepare Compound **51**, which was isolated as a trifluoroacetate salt. MS 548 (MH⁺).

15 *Biological Examples*

The ability of the compounds to treat or ameliorate protein kinase mediated disorders was determined using the following procedures.

Example 52*EGFR Kinase Assay*

20 The EGFR kinase used was a fusion of Glutathione-S-Transferase (GST) and a PCR amplified intracellular portion of EGFR (NM_005228). The intracellular portion of EGFR started at nucleotide 2189 (corresponding to amino acid 667) and ended at the

termination codon. The portion was PCR amplified with primers that added the lambda attB sequences to each end, recombined into an entry vector, then into a GST destination vector (as described in Gateway Technologies Manual by Invitrogen Corporation, Carlsbad, California).

5 The destination vector was recombined in the DH10BAC strain of bacteria to produce a bacmid. The bacmid was transfected into Sf 9 cells and the supernatant containing the baculovirus was collected. The GSTEGFR protein was purified using large cultures of Sf 9 cells infected with stock virus. After an appropriate period of time, the cells were collected and lysed. The GSTEGFR was then purified from the
10 lysate on Glutathione-Sepharose columns (as described by Amersham Biosciences, Buckinghamshire, United Kingdom).

 The EGFR substrate was prepared by biotinylating polyGluTyr (128 mg) (Sigma, St. Louis, Missouri) in a 1X PBS buffer incubated together with a 12-fold molar excess of Sulfo-NHS-LC-Biotin on ice for at least 2 hrs. The free biotin was
15 separated from the biotinylated polyGluTyr on a gel filtration column.

 A mixture of a 10X kinase buffer (500 mM Tris at pH 8.0, 100 mM Magnesium Chloride and 1 mM Sodium Vanadate), DTT (1 mM final from 500 mM stock), ATP (5 μ M final from 10 mM stock), biotinylated polyGluTyr (10 μ g/ μ L stock), γ -³³P ATP (10 μ Ci/ μ L stock) and water was added to each well (90 μ L/well) of a Streptavidin
20 Flashplate (Perkin Elmer, Wellesley, MA).

 Test compound in 100% DMSO (2 μ L) was added to the appropriate wells. Diluted GSTEGFR (1:300 dilution in 50 mM Tris at pH 8.0 and 0.1% bovine serum albumin) (10 μ L) was added to the wells to initiate the reactions.

 The plates were incubated at 30 °C for 1 hr with shaking. The reacted contents
25 were removed and the plates were sequentially washed three times with a 1X PBS stop buffer (300 μ L without Magnesium and Calcium) and 100 mM EDTA. After the final wash, the same stop buffer (200 μ L) was added to the wells. The plates were then sealed and read on the TopCount scintillation counter.

 Test compounds were assayed in triplicate at 16 concentrations at half-log
30 dilutions starting at 200 μ M. A maximum and minimum signal for the assay was determined on each plate. The percent inhibition of a test compound was calculated according to the formula

$$\left[\frac{(\text{max signal} - \text{test compound})}{(\text{max signal} - \text{min signal})} \right] (100) = \% \text{ inhibition}$$

For a series of test concentrations, the IC₅₀ was derived by graphing percent inhibition against the log of the concentrations tested for a given compound. The IC₅₀ results are shown in Table 1. For those compounds without an IC₅₀, the percent

5 inhibition results are shown at a test concentration of 2 μM.

Table 1

EGFR IC₅₀ (μM)

Cpd	IC ₅₀	Cpd	IC ₅₀
1	0.02255	22	0.85335
2	0.078	23	0.02045
3	0.04367	24	0.0333
4	0.049	25	0.04331
5	0.06367	26	0.15845
6	0.053	27	0.20705
7	0.0285	28	0.3272
8	2.6335	29	0.0696
9	2.6305	30	0.06772
10	0.0565	31	0.06035
11	0.04173	34	1.452
12	15.85	35	0.461
13	3.28	36	8.757
14	0.0293	37	7.212
15	0.07605	38	2.199
16	0.07645	39	3.257
17	0.03994	40	46%
18	0.08575	41	44%
19	0.0795	42	18%
20	0.3776, 0.941	43	4%
21	0.04459	44	47%
		51	0%

Example 53

c-Src Kinase Assay

10 A mixture of a 10X kinase buffer (80 mM MOPS at pH 7.0, 2 mM EDTA and 100 mM Magnesium Chloride), ATP (5 μM final from a 10 mM stock), a Cdc2 peptide

KVEKIGEGTYGVVYK (100 μ M final from a 2.5 mM stock), γ - ^{33} P ATP (10 μ Ci/ μ L stock) and water (20 μ L/well) was added to each well of a Streptavidin Flashplate.

Test compound in 100% DMSO (0.5 μ L) was added to the appropriate wells. Diluted c-Src kinase (human) (Upstate Biotechnology, Lake Placid, New York) (diluted
5 in a buffer consisting of 20 mM MOPS at pH 7.0, 1 mM EDTA, β -mercaptoethanol (0.1%), Brij-35 (0.01%), glycerol (5 %), and 1 mg/mL bovine serum albumin) (2.5 μ L) was added to the wells to initiate the reactions. The reaction plates were incubated at 30 °C for 40 min. The reaction was terminated by the addition of a 3% phosphoric acid solution (5 μ L). The reaction product (10 μ L) was spotted onto a P30 filtermat and
10 washed for 5 minutes in phosphoric acid (75 mM). The wash sequence was repeated two more times, followed with one final wash in methanol. The plates were then dried, sealed and read on the TopCount scintillation counter after adding 30 μ L scintillation fluid.

The percent inhibitions and IC₅₀ values were derived according to the procedure
15 described in Example 52. The IC₅₀ results are shown in Table 2. For those compounds without an IC₅₀, the percent inhibition results are shown at a test concentration of ⁽¹⁾ 1 μ M or ⁽²⁾ 1.6 μ M.

Table 2

c-SRC IC₅₀ (μ M)

Cpd	IC ₅₀	Cpd	IC ₅₀
20	0.011 (h)	39	> 100
23	⁽¹⁾ -2% (h)	40	2.394
24	⁽¹⁾ 15% (h)	41	0.079
25	1.406	42	1.381
35	⁽²⁾ 11 %	43	> 100
36	⁽²⁾ 4 %	44	0.126
37	⁽²⁾ 3 %	45	0.145
38	⁽²⁾ 4 %	46	0.389
		51	0%

20

Example 54

Lyn Kinase Assay

A mixture of a 10X kinase buffer (500mM MOPS at pH 7.5, 1 mM EGTA, 1 mM Sodium Vanadate, 1% β -mercaptoethanol and 100 mM Magnesium Acetate), ATP

(5 μ M final from a 10 mM stock), polyGluTyr (0.1 mg/mL final from a 1 mg/mL stock), γ -³³P ATP (10 μ Ci/ μ L stock) and water (20 μ L/well) was added to each well of a Streptavidin Flashplate.

Test compound in 100% DMSO (0.5 μ L) was added to the appropriate wells.

- 5 Diluted Lyn kinase (human) (Upstate Biotechnology, Lake Placid, New York) (diluted in a buffer consisting of 50 mM Tris at pH 7.5, 0.1 mM EGTA, Sodium Vanadate (0.1 mM), β -mercaptoethanol (0.1%) and 1 mg/mL bovine serum albumin) (2.5 μ L) was added to the wells to initiate the reactions.

- 10 The reaction plates were incubated at 30 °C for 40 min. The reaction was terminated by the addition of a 3% phosphoric acid solution (5 μ L). The reaction product (10 μ L) was spotted onto a P30 filtermat and washed for 5 minutes in phosphoric acid (75 mM). The wash sequence was repeated two more times, followed with one final wash in methanol. The plates were then dried, sealed and read on the TopCount scintillation counter after adding 30 μ L scintillation fluid.

- 15 The percent inhibitions and IC₅₀ values were derived according to the procedure described in Example 52. The IC₅₀ results are shown in Table 3. For those compounds without an IC₅₀, the percent inhibition results are shown at a test concentration of ⁽¹⁾ 1 μ M or ⁽²⁾ 1.6 μ M.

Table 3

20

LYN IC₅₀ (μ M)

Cpd	IC ₅₀	Cpd	IC ₅₀
20	0.011 (h), 0.009 (m)	24	⁽¹⁾ -5% (h), ⁽¹⁾ 10% (m)
23	⁽¹⁾ 8% (h), ⁽¹⁾ 42% (m)	25	0.224 (h), 0.79 (m)
		51	⁽²⁾ 37% (h)

Example 55

HER-2 Kinase Assay

- The HER-2 kinase used was purified at Proqinase (Freiburg, Germany) from a construct that consisted of a fusion of GST (Glutathione-S-Transferase),
 25 HIS6-Thrombin and the nucleotides encoding amino acids 679 to 1255 of HER-2.

A mixture of a 10X kinase reaction buffer (600 mM Hepes at pH 7.5, 30 mM Magnesium Chloride, 0.03 mM Sodium Vanadate and 500 μ g/mL PEG 20,000), DTT (1.2 mM final from a 10 mM stock), ATP (1 μ M from a 10 mM stock), biotinylated

polyGluTyr (1.5 ng/ μ L final from stock of 1 μ g/ μ L prepared by Upstate Biotechnologies, Lake Placid, New York), Manganese Chloride (3 mM final from a 1 M stock), γ -³³P-ATP (10 μ Ci/ μ L stock) and water (70 μ L/well) was added to each well of a Streptavidin Flashplate (Cat. # SMP103, NEN, Boston, MA).

- 5 Test compound stock (1 μ L) was added to the appropriate wells. Diluted GSTHER2 kinase (6.7 ng/ μ L diluted into 50 mM Tris-HCl at pH 8.0 and 0.1% bovine serum albumin) (30 μ L) was added (total volume of 200 ng/well) to initiate the reactions.

- 10 The reaction plates were incubated at 30 °C for 1 hr. The reaction was terminated by aspirating the reaction mixture from the plate wells and washing the wells three times with a 1X PBS stop buffer (300 μ L) and 100 mM EDTA. After the final wash, the same stop buffer (200 μ L) was again added to the wells. The plates were then sealed and read on the TopCount scintillation counter.

- 15 The IC₅₀ was derived according to the procedure described in Example 52. The IC₅₀ results are shown in Table 4. For those compounds without an IC₅₀, the percent inhibition results are shown at a test concentration of 1 μ M.

Table 4

HER-2 IC ₅₀ (μ M)			
Cpd	IC ₅₀	Cpd	IC ₅₀
2	100	19	100
10	10	24	2.06
17	46.03	25	92.74
		51	>100

Example 56

20 *c-Abl Kinase Assay*

A mixture of a 10X kinase buffer (80 mM MOPS at pH 7.0, 2 mM EDTA and 100 mM Magnesium Acetate), ATP (5 μ M final from a 10 mM stock), a peptide EAIYAAPFAKKK (50 μ M final from a 0.5 mM stock), γ -³³P ATP (10 μ Ci/ μ L stock) and water is added to each well (20 μ L/well) of a Streptavidin Flashplate.

- 25 Test compound in 100% DMSO (0.5 μ L) is added to the appropriate wells. Diluted c-Abl kinase (human) (Upstate Biotechnology, Lake Placid, New York) (diluted in a buffer consisting of 20 mM MOPS at pH 7.0, 1 mM EDTA,

β -mercaptoethanol (0.1%), Brij-35 (0.01%), glycerol (5 %) and 1 mg/mL bovine serum albumin) (2.5 μ L) is added to the wells to initiate the reactions.

The reaction plates are incubated at 30 °C for 40 min. The reaction is terminated by the addition of a 3% phosphoric acid solution (5 μ L). The reaction product (10 μ L) is spotted onto a P30 filtermat and is washed for 5 minutes in phosphoric acid (75 mM). The wash sequence is repeated two more times and is followed with one final wash in methanol. The plates are then dried, sealed and read on the TopCount scintillation counter after 30 μ L scintillation fluid is added.

The percent inhibitions and IC₅₀ values are derived according to the procedure described in Example 52.

Example 57

Cell Proliferation Inhibition Assay

The ability of a test compound to inhibit unregulated cell proliferation may be determined by measuring incorporation of ¹⁴C-labelled thymidine into newly synthesized DNA within cell lines derived from carcinomas originating from several tissues. Accordingly, the anti-proliferative effect of a compound on cells with a variety of phenotypes may be determined.

Carcinoma cell lines include those such as HeLa cervical adenocarcinoma (American Type Culture Collection (ATCC), Virginia, Cat. #CCL-2), A375 malignant melanoma (ATCC CRL-1619), SK-OV-3 ovarian adenocarcinoma (ATCC HTB-77), HCT-116 colon carcinoma (CCL-247), PC-3 prostate adenocarcinoma (ATCC CRL-1435), and MDA-MB-231 (Xenogen Corp.)

The carcinoma cells are trypsinized and counted. The cells (3000-8000 count) are added to each well of a 96-well CytoStar tissue culture treated scintillating microplate (Amersham #RPNQ0160) in complete medium (100 μ L) and the plate is then incubated in complete medium for 24 hrs at 37 °C in an inert atmosphere containing 5% CO₂. Test compound (1 μ L) in 100% DMSO is added to the plate test-wells with DMSO only added to control-wells. The plate is incubated in complete medium for a second 24 hr period at 37 °C in an atmosphere containing 5% CO₂.

An aliquot of a solution of Methyl ¹⁴C-thymidine (56 mCi/mmol) (NEN #NEC568 or Amersham #CFA532) and complete medium (20 μ L to provide 0.2 μ Ci/well) is then added to each well and the plate is incubated for a third 24 hr period

at 37°C in an atmosphere containing 5% CO₂. The plate contents are then discarded, the plate is washed twice with PBS (200 µL) and then PBS (200 µL) is added to each well. The plate is sealed and the degree of methyl ¹⁴C-thymidine incorporation is quantified on a Packard Top Count.

5 Example 58

In Vivo Models – Inhibition of Tumor Growth

The ability of a test compound to inhibit unregulated growth of human tumor cells *in vivo* may be evaluated by implanting human tumor cells into the hindflank of athymic mice, administering a test compound and then quantifying any change in tumor
10 size.

Human epidermoid A431 carcinoma cells (10⁶ count) are implanted subcutaneously into the hindflank of female athymic mice (Charles River) and allowed to grow for 6-10 days. After a measurable tumor is established (as determined by baseline caliper measurement), the animal is administered an oral dose of the test
15 compound (in 10% solutol) daily for a period of 30 days. Tumor size is measured every five days and the degree of inhibition is determined by comparing drug-treated animals to vehicle-treated animals.


Variations of this method are intended to include intraperitoneal injection or intravenous infusion as the route of administration and administration of the test
20 compound either alone or in a combination therapy.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and modifications as come within the scope of the following claims and their
25 equivalents.

Throughout this application, various publications are cited. The disclosure of these publications is hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

What is Claimed is:

1. A compound of Formula (I)



Formula (I)

and pharmaceutically acceptable forms thereof, wherein

L¹ is selected from the group consisting of S(C₁-₄alkyl), a bond, N(R₁), N(R₁)C(O) and
 5 C(O)N(R₁), wherein R₁ is selected from the group consisting of hydrogen, C₁-₈alkyl and C₁-₈alkyl(C₁-₈alkoxy);

p is 0, 1, 2, 3 or 4;

L² is selected from the group consisting of O, S, N(R₁) and a bond;

Ar¹ is selected from the group consisting of aryl, heteroaryl, benzofused heteroaryl,

10 heterocyclyl and benzofused heterocyclyl optionally substituted with from one to three substituents independently selected from the group consisting of

- (1) C₁-₈alkyl,
- (2) C₂-₈alkenyl,
- (3) C₂-₈alkynyl,

15 (4) C₁-₈alkoxy,

wherein (1), (2), (3) and (4) are optionally substituted with one to three substituents independently selected from the group consisting of

- (i) C₃-₈cycloalkyl,
- (ii) aryl,
- 20 (iii) heteroaryl,
- (iv) heterocyclyl,

wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to three substituents independently selected from the group consisting of

- (a) C₁-₈alkyl,
- 25 (b) C₁-₈alkoxy,
- (c) C₁-₈alkyl(C₁-₈alkoxy),
- (d) C₁-₈alkyl(halogen)₁-₃,

- (e) C₁₋₈alkyl(hydroxy)₁₋₃,
- (f) CO₂(C₁₋₈alkyl),
- (g) amino optionally mono or disubstituted with C₁₋₈alkyl,
- (h) cyano,
- 5 (i) halogen,
- (j) hydroxy,
- (k) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (l) C₃₋₈cycloalkyl,
- 10 (m) C₁₋₈alkyl(C₃₋₈cycloalkyl),
- (n) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
- (o) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents, and
- 15 (v) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
 - (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (c) C₁₋₈alkyl(hydroxy)₁₋₃,
 - 20 (d) C₃₋₈cycloalkyl,
 - (e) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
 - (f) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- 25 (5) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
 - (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
 - 30 (iv) C₁₋₈alkyl(hydroxy)₁₋₃,
 - (v) C₃₋₈cycloalkyl,
 - (vi) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and

- (vii) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (6) cyano,
- (7) halogen,
- 5 (8) hydroxy,
- (9) C₃₋₈cycloalkyl,
- (10) aryl,
- (11) heteroaryl,
- (12) heterocyclyl,
- 10 (13) oxy substituted with a substituent selected from the group consisting of
- (i) CF₃,
- (ii) C₃₋₈cycloalkyl,
- (iii) aryl,
- (iv) heteroaryl, and
- 15 (v) heterocyclyl,
- (14) SO₂(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (15) C(O) substituted with a substituent selected from the group consisting of
- (i) hydrogen,
- 20 (ii) hydroxy,
- (iii) C₁₋₈alkyl,
- (iv) C₁₋₈alkoxy, and
- (v) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- 25 (a) C₁₋₈alkyl,
- (b) C₁₋₈alkyl(C₁₋₈alkoxy),
- (c) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (d) C₁₋₈alkyl(hydroxy)₁₋₃,
- 30 (e) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (f) C₃₋₈cycloalkyl, and
- (g) aryl; and

(16) SO₂(amino), wherein amino is optionally mono or disubstituted with a substituent independently selected from the group consisting of

- (i) C₁₋₈alkyl,
- (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
- 5 (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (iv) C₁₋₈alkyl(hydroxy)₁₋₃,
- (v) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (vi) C₃₋₈cycloalkyl, and
- 10 (vii) aryl; and

Ar² is selected from the group consisting of aryl, heteroaryl, benzofused heteroaryl, heterocyclyl and benzofused heterocyclyl optionally substituted with from one to three substituents independently selected from the group consisting of

- (1) C₁₋₈alkyl,
- 15 (2) C₂₋₈alkenyl,
- (3) C₂₋₈alkynyl,
- (4) C₁₋₈alkoxy,

wherein (1), (2), (3) and (4) are optionally substituted with from one to three substituents independently selected from the group consisting of

- 20 (i) C₁₋₈alkoxy,
- (ii) cyano,
- (iii) halogen,
- (iv) hydroxy,
- (v) C₃₋₈cycloalkyl,
- 25 (vi) aryl,
- (vii) heteroaryl,
- (viii) heterocyclyl,

wherein (v), (vi), (vii) and (viii) are optionally substituted with from one to three substituents independently selected from the group consisting of

- 30 (a) C₁₋₈alkyl,
- (b) C₁₋₈alkoxy,
- (c) C₁₋₈alkyl(C₁₋₈alkoxy),

- (d) amino optionally mono or disubstituted with C₁₋₈alkyl,
(e) cyano,
(f) halogen,
(g) C₁₋₈alkyl(halogen)₁₋₃,
5 (h) hydroxy,
(i) C₁₋₈alkyl(hydroxy)₁₋₃,
(j) C₃₋₈cycloalkyl, and
(k) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
substituents,
10 (ix) amino optionally mono or disubstituted with a substituent independently
selected from the group consisting of
(a) C₁₋₈alkyl,
(b) C₁₋₈alkyl(C₁₋₈alkoxy),
(c) C₁₋₈alkyl(hydroxy)₁₋₃,
15 (d) C₃₋₈cycloalkyl,
(e) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
substituents, and
(f) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one
or two C₁₋₈alkyl substituents, and
20 (x) C(O)amino optionally mono or disubstituted with a substituent
independently selected from the group consisting of
(a) C₁₋₈alkyl,
(b) C₁₋₈alkyl(C₁₋₈alkoxy),
(c) C₁₋₈alkyl(hydroxy)₁₋₃,
25 (d) C₃₋₈cycloalkyl,
(e) aryl optionally substituted with one to three halogen substituents,
(f) heterocyclyl optionally substituted with one or two C₁₋₈alkyl
substituents, and
(g) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one
30 or two C₁₋₈alkyl substituents,
(5) amino optionally mono or disubstituted with a substituent independently
selected from the group consisting of
(i) C₁₋₈alkyl,

- (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
- (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
- (iv) C₁₋₈alkyl(hydroxy)₁₋₃,
- (v) C₃₋₈cycloalkyl,
- 5 (vi) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents,
and
- (vii) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or
two C₁₋₈alkyl substituents,
- (6) oxy substituted with a substituent selected from the group consisting of
- 10 (i) C₃₋₈cycloalkyl,
- (ii) aryl,
- (iii) heteroaryl, and
- (iv) heterocyclyl,
wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to
- 15 three substituents independently selected from the group consisting
of
- (a) C₁₋₈alkyl,
- (b) C₁₋₈alkoxy,
- (c) C₁₋₈alkyl(C₁₋₈alkoxy),
- 20 (d) amino optionally mono or disubstituted with C₁₋₈alkyl, and
- (e) halogen,
- (7) C(O) substituted with a substituent independently selected from the group
consisting of
- (i) hydrogen,
- 25 (ii) C₁₋₈alkyl optionally substituted with from one to three substituents
independently selected from the group consisting of
- (a) cyano,
- (b) halogen, and
- (c) hydroxy,
- 30 (iii) C₁₋₈alkoxy,
- (iv) hydroxyl, and
- (v) C₁₋₈alkoxy(C₁₋₈alkoxy),
- (8) cyano,

- (9) halogen,
(10) hydroxy,
(11) nitro,
(12) C₃₋₈cycloalkyl,
5 (13) aryl,
(14) heteroaryl,
(15) benzofused heteroaryl,
(16) heterocyclyl, and
(17) benzofused heterocyclyl;
10 wherein (12)-(17) are optionally substituted with one or two substituents independently selected from the group consisting of
- (i) C₃₋₈cycloalkyl,
(ii) C₁₋₈alkyl(C₃₋₈cycloalkyl),
(iii) aryl,
15 (iv) C₁₋₈alkyl(aryl),
(v) heteroaryl,
(vi) C₁₋₈alkyl(heteroaryl),
(vii) heterocyclyl, and
(viii) C₁₋₈alkyl(heterocyclyl).
- 20 2. The compound of claim 1, wherein L¹ is selected from the group consisting of N(R₁), wherein R₁ is selected from the group consisting of hydrogen, C₁₋₈alkyl and C₁₋₈alkyl(C₁₋₈alkoxy).
3. The compound of claim 1, wherein L¹ is NH.
4. The compound of claim 1, wherein p is 0, 1, 2 or 3.
- 25 5. The compound of claim 1, wherein L² is selected from the group consisting of O, S and N(R₁).
6. The compound of claim 1, wherein L² is selected from the group consisting of O, S and NH.

7. The compound of claim 1, wherein Ar¹ is selected from the group consisting of aryl, heteroaryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of
- (1) C₁₋₈alkyl,
- 5 (2) C₁₋₈alkoxy,
- wherein (1) and (2) are optionally substituted with one to three substituents independently selected from the group consisting of
- (i) C₃₋₈cycloalkyl,
- (ii) aryl,
- 10 (iii) heteroaryl,
- (iv) heterocyclyl,
- wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to three substituents independently selected from the group consisting of
- 15 (a) C₁₋₈alkyl,
- (b) C₁₋₈alkoxy,
- (c) C₁₋₈alkyl(C₁₋₈alkoxy),
- (d) C₁₋₈alkyl(halogen)₁₋₃,
- (e) C₁₋₈alkyl(hydroxy)₁₋₃,
- 20 (f) amino optionally mono or disubstituted with C₁₋₈alkyl,
- (g) cyano,
- (h) halogen,
- (i) hydroxy, and
- (j) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with
- 25 C₁₋₈alkyl, and
- (v) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (a) C₁₋₈alkyl,
- (b) C₁₋₈alkyl(C₁₋₈alkoxy),
- 30 (c) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
- (d) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,

- (3) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
 - (iii) heterocyclyl optionally substituted with one or two C₁₋₈alkyl substituents, and
 - (iv) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents,
- (4) halogen,
- (5) hydroxy,
- (6) C₃₋₈cycloalkyl,
- (7) aryl,
- (8) heteroaryl,
- (9) heterocyclyl, and
- (10) SO₂(amino), wherein amino is optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (iii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,
 - (iv) C₁₋₈alkyl(hydroxy)₁₋₃, and
 - (v) C₁₋₈alkyl(heterocyclyl) optionally substituted on heterocyclyl with one or two C₁₋₈alkyl substituents.
8. The compound of claim 1, wherein Ar¹ is selected from the group consisting of aryl, heteroaryl and heterocyclyl, each optionally substituted with from one to three substituents independently selected from the group consisting of
- (1) C₁₋₈alkyl,
 - (2) C₁₋₈alkoxy,
- wherein (1) and (2) are optionally substituted with one to three substituents independently selected from the group consisting of
- (i) heteroaryl, and

- (ii) heterocyclyl,
 wherein (i) and (ii) are optionally substituted with from one to three
 substituents independently selected from the group consisting of
- (a) C₁₋₈alkyl,
 5 (b) C₁₋₈alkyl(hydroxy), and
 (c) amino, and
- (iii) amino optionally mono or disubstituted with a substituent independently
 selected from the group consisting of
- (a) C₁₋₈alkyl,
 10 (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 (c) heterocyclyl, and
 (d) C₁₋₈alkyl(heterocyclyl),
- (3) amino optionally mono or disubstituted with a substituent independently
 selected from the group consisting of
- 15 (i) C₁₋₈alkyl(amino), and
 (ii) C₁₋₈alkyl(heterocyclyl),
- (4) halogen,
 (5) hydroxy,
 (6) heterocyclyl, and
- 20 (7) SO₂(amino), wherein amino is optionally mono or disubstituted with a
 substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl(C₁₋₈alkoxy),
 (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with
 C₁₋₈alkyl, and
- 25 (iii) C₁₋₈alkyl(heterocyclyl).
9. The compound of claim 1, wherein Ar² is selected from the group consisting of
 aryl and heterocyclyl, each optionally substituted with from one to three
 substituents independently selected from the group consisting of
- (1) C₁₋₈alkyl,
 30 (2) C₂₋₈alkenyl,
 (3) C₂₋₈alkynyl,
 (4) C₁₋₈alkoxy,

wherein (1), (2), (3) and (4) are optionally substituted with from one to three substituents independently selected from the group consisting of

- (i) C₁₋₈alkoxy,
- (ii) halogen,
- (iii) hydroxy,
- (iv) aryl optionally substituted with from one to three substituents independently selected from the group consisting of

- (a) C₁₋₈alkyl,
- (b) C₁₋₈alkoxy,
- (c) C₁₋₈alkyl(C₁₋₈alkoxy),
- (d) amino optionally mono or disubstituted with C₁₋₈alkyl,
- (e) halogen,
- (f) C₁₋₈alkyl(halogen)₁₋₃,
- (g) hydroxy, and
- (h) C₁₋₈alkyl(hydroxy)₁₋₃, and

- (v) amino optionally mono or disubstituted with C₁₋₈alkyl,

- (5) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of

- (i) C₁₋₈alkyl, and
- (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl,

- (6) oxy substituted with a substituent selected from the group consisting of

- (i) C₃₋₈cycloalkyl,
- (ii) aryl,
- (iii) heteroaryl, and
- (iv) heterocyclyl,

wherein (i), (ii), (iii) and (iv) are optionally substituted with from one to three substituents independently selected from the group consisting of


- (a) C₁₋₈alkyl,
- (b) C₁₋₈alkoxy,
- (c) amino optionally mono or disubstituted with C₁₋₈alkyl, and
- (d) halogen,

- (7) cyano,
(8) halogen, and
(9) hydroxy.
10. The compound of claim 1, wherein Ar² is selected from the group consisting of
5 aryl and heterocyclyl, each optionally substituted with from one to three
substituents independently selected from the group consisting of
- (1) C₂₋₈alkynyl,
(2) C₁₋₈alkoxy optionally substituted with from one to three substituents
independently selected from the group consisting of
- 10 (i) halogen, and
(ii) aryl optionally substituted with from one to three halogen substituents,
(3) oxyaryl optionally substituted with from one to three substituents
independently selected from the group consisting of
- (i) C₁₋₈alkyl,
15 (ii) C₁₋₈alkoxy,
(iii) amino optionally mono or disubstituted with C₁₋₈alkyl, and
(iv) halogen, and
(4) halogen.
11. The compound of claim 1, wherein
- 20 L¹ is NH;
p is 0, 1, 2 or 3;
L² is selected from the group consisting of O, S and NH;
Ar¹ is selected from the group consisting of aryl, heteroaryl and heterocyclyl, each
optionally substituted with from one to three substituents independently selected
25 from the group consisting of
- (1) C₁₋₈alkyl,
(2) C₁₋₈alkoxy,
wherein (1) and (2) are optionally substituted with one to three substituents
independently selected from the group consisting of
- 30 (i) heteroaryl, and
(ii) heterocyclyl,
wherein (i) and (ii) are optionally substituted with from one to three


substituents independently selected from the group consisting of

- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(hydroxy), and
 - (c) amino, and
- 5 (iii) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (a) C₁₋₈alkyl,
 - (b) C₁₋₈alkyl(C₁₋₈alkoxy),
 - (c) heterocyclyl, and
 - 10 (d) C₁₋₈alkyl(heterocyclyl),
- (3) amino optionally mono or disubstituted with a substituent independently selected from the group consisting of
- (i) C₁₋₈alkyl(amino), and
 - (ii) C₁₋₈alkyl(heterocyclyl),
- 15 (4) halogen,
- (5) hydroxy,
- (6) heterocyclyl, and
- (7) SO₂(amino), wherein amino is optionally mono or disubstituted with a substituent independently selected from the group consisting of
- 20 (i) C₁₋₈alkyl(C₁₋₈alkoxy),
- (ii) C₁₋₈alkyl(amino) optionally mono or disubstituted on amino with C₁₋₈alkyl, and
- (iii) C₁₋₈alkyl(heterocyclyl); and
- Ar² is selected from the group consisting of aryl and heterocyclyl, each optionally
- 25 substituted with from one to three substituents independently selected from the group consisting of
- (1) C₂₋₈alkynyl,
 - (2) C₁₋₈alkoxy optionally substituted with from one to three substituents independently selected from the group consisting of
- 30 (i) halogen, and
- (ii) aryl optionally substituted with from one to three halogen substituents, and
- (3) oxyaryl optionally substituted with from one to three substituents independently selected from the group consisting of


- (i) C₁₋₈alkyl,
 - (ii) C₁₋₈alkoxy,
 - (iii) amino optionally mono or disubstituted with C₁₋₈alkyl, and
 - (iv) halogen, and
- 5 (4) halogen.
12. The compound of claim 1, wherein the compound is an isolated form thereof.
13. A pharmaceutical composition comprising an effective amount of a compound of any of claim 1 to 11.
14. The pharmaceutical composition of claim 13, wherein the effective amount of
10 the compound is in a range of from about 0.001 mg to about 5000 mg.
15. A process for preparing a pharmaceutical composition comprising the step of admixing a compound of any of claim 1 to 11 and a pharmaceutically acceptable carrier.
16. A process for preparing a compound of any of claim 1 to 11 comprising the
15 steps of
- a. reacting a compound of Formula A1 with a compound of Formula A2 to prepare a compound of Formula A3;



- b. transforming a compound of Formula A3 into a compound of Formula A4;




- c. reacting a compound of Formula A4 with a compound of Formula A5 to prepare a compound of Formula A6,




5

wherein L^a is a suitable leaving group or $-SCH_3$;

- d. transforming a compound of Formula A6 into a compound of Formula A7;




- e. transforming a compound of Formula A7 into a compound of Formula A8;




10

- f. reacting a compound of Formula A8 with a compound of Formula A9 to prepare a compound of Formula A10,




- wherein Ar^2 is as previously defined and L^2 is a suitably reactive heteroatom selected from N, NH, NH_2 , O or S;

- g. transforming a compound of Formula A10 into a compound of Formula A11,




- h. reacting a compound of Formula A11 with a compound of Formula A12 to prepare a compound of Formula (I),



- wherein, for a compound of Formula A12, L^1 is a suitably reactive heteroatom selected from NH, NH_2 , O or S.


17. A process for preparing a compound of any of claim 1 to 11 comprising the steps of

- a. reacting a compound of Formula A3 with a compound of Formula A5 to prepare a compound of Formula B1,




wherein L^a is a suitable leaving group or $-SCH_3$;


- b. transforming a compound of Formula B1 into a compound of Formula B2,



- 5 c. transforming a compound of Formula B2 into a compound of Formula B3;



- d. transforming a compound of Formula B3 into a compound of Formula A7;



18. A method for inhibiting unregulated protein kinase activity comprising contacting the protein kinase domain with at least one compound of any of claim 1 to 11.
- 5 19. A method for treating or preventing a chronic or acute kinase mediated disease in a subject in need thereof comprising administering to the subject an effective amount of a compound of any of claim 1 to 11.
20. The method of claim 19, wherein the effective amount of the compound is in a range of from about 0.001 mg to about 5000 mg.
- 10 21. The method of claim 19 wherein the chronic or acute kinase mediated disease is a chronic or acute tumor associated with non-small-cell lung cancers, colon cancers, breast cancers and the like wherein the cancer is mediated by a tyrosine-kinase selected from EGFR, HER-2, c-Src or Lyn.