



(86) Date de dépôt PCT/PCT Filing Date: 2000/04/19

(87) Date publication PCT/PCT Publication Date: 2000/10/26

(45) Date de délivrance/Issue Date: 2008/06/17

(85) Entrée phase nationale/National Entry: 2001/10/10

(86) N° demande PCT/PCT Application No.: US 2000/010383

(87) N° publication PCT/PCT Publication No.: 2000/063197

(30) Priorité/Priority: 1999/04/19 (US60/129,933)


(51) Cl.Int./Int.Cl. *C07D 279/16* (2006.01),
A61K 31/426 (2006.01), *A61K 31/5415* (2006.01),
A61K 31/542 (2006.01), *C07D 277/14* (2006.01),
C07D 513/04 (2006.01)

(72) Inventeurs/Inventors:
SCARLATO, GERARD ROBERT, US;
HADIDA RUAH, SARA SABINA, US;
NISHIMURA, TAMIKI, JP;
NAKATSUKA, MASASHI, JP;
SAMIZO, FUMIO, JP;
KAMIKAWA, YUMIKO, JP;
HOUTIGAI, HITOSHI, JP

(73) Propriétaire/Owner:

(54) Titre : DERIVE D'ACIDE HYDROXAMIQUE

(54) Title: HYDROXAMIC ACID DERIVATIVE



(57) Abrégé/Abstract:

A hydroxamic acid derivative represented by formula (1) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, which has a matrix metallo-proteinase inhibitor.

(73) Propriétaires(suite)/Owners(continued):DAINIPPON SUMITOMO PHARMA CO., LTD., JP


(74) Agent: KIRBY EADES GALE BAKER



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 277/14, 279/16, 417/04, 417/14, 417/06, 417/12, 513/04, A61K 31/5415, 31/426, A61P 29/00 // (C07D 513/04, 279:00, 231:00)	A1	(11) International Publication Number: WO 00/63197 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/US00/10383 (22) International Filing Date: 19 April 2000 (19.04.00) (30) Priority Data: 60/129,933 19 April 1999 (19.04.99) US (71) Applicant (for all designated States except US): SUMITOMO PHARMACEUTICALS COMPANY, LIMITED [JP/JP]; 2-8, Doshomachi 2-chome, Chuo-ku, Osaka-shi, Osaka (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): SCARLATO, Gerard, Robert [US/US]; 1121 Virginia Way, La Jolla, CA 92037 (US). HADIDA RUAH, Sara Sabina [ES/US]; 3717 Nobel Drive #1411, San Diego, CA 92122 (US). NISHIMURA, Tamiki [JP/JP]; 2-11-7-201, Sonehigashimachi, Toyonaka-shi, Osaka (JP). NAKATSUKA, Masashi [JP/JP]; 3-2-4-807, Yamazaki, Shimamoto-cho, Mishima-gun, Osaka (JP). SAMIZO, Fumio [JP/JP]; 3-11-30-313, Mukocho, Amagasaki-shi, Hyogo (JP). KAMIKAWA, Yumiko [JP/JP]; 379-12, Kitanagai-cho, Nara-shi, Nara		(JP). HOUTIGAI, Hitoshi [JP/JP]; 1-9-1-503, Tamagawa, Takatsuki-shi, Osaka (JP). (74) Agents: MEIKLE, Andrew, D. et al.; Birch, Stewart, Kolasch & Birch, LLP, P.O. Box 747, Falls Church, VA 22040-0747 (US). (81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: HYDROXAMIC ACID DERIVATIVE  (57) Abstract A hydroxamic acid derivative represented by formula (1) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, which has a matrix metallo-proteinase inhibitor.		

DESCRIPTION

HYDROXAMIC ACID DERIVATIVE

5

Technical Field

The present invention relates to hydroxamic acid derivatives useful as matrix metallo-proteinase inhibitors.

10

Background Art

15

Extra cellular matrix, composition of connective tissue, is metabolized by a family of proteinases termed matrix metallo-proteinases. It is known that there exist 16 kinds of matrix metallo-proteinases such as collagenase (matrix metallo-proteinase-1: MMP-1), gelatinase A (matrix metallo-proteinase-2: MMP-2), stromelysin (matrix metallo-proteinase-3: MMP-3), gelatinase B (matrix metallo-proteinase-9: MMP-9) and collagenase-3 (matrix metallo-proteinase-13: MMP-13).

20

Extra cellular matrix is under tight control by the expression and secretion of these matrix metallo-proteinases or endogenous inhibitors such as tissue inhibitor of matrix metallo-proteinases in normal. There are many reports about relationships between diseases characterized by excessive tissue disruption and elevated activities of matrix metallo-proteinases derived from the breakdown of these control.

25

Elevated levels of matrix metallo-proteinases,

particularly collagenase and stromelysin, have been detected in joints of osteoarthritic humans or that of rheumatoid arthritis (Arthr. Rheum., 33, 388-97 (1990); S.M. Krane et. al., "Modulation of matrix synthesis and in The Control of degradation in joint inflammation, The Control of Tissue Damage", A.B. Glauert (ed.), Elsevier Sci. Publ., Amsterdam, 1988, Ch. 14, pp 179-95; Clin.Chim.Acta, 185, 73-80(1989); Arthr.Rheum., 27, 305-312(1984); J.Clin.Invest., 84, 678-685(1989)).

Secreted proteinases such as stromelysin, collagenase and gelatinase play an important role in processes involved in the movement of cells during metastatic tumor invasion. Indeed, there is also evidence that the matrix metallo-proteinases are overexpressed in certain metastatic tumor cell line. In this context, the enzyme functions to penetrate underlying basement membranes and allow the tumor cell to escape from the site of primary tumor formation and enter circulation (FEBS J., 5, 2145-2154(1991); Trends Genet., 6, 121-125(1990); Cancer Res., 46, 1-7(1986); Cell, 64, 327-336(1990); Cancer and Metastasis Rev., 9, 305-319(1990)).

Both collagenase and stromelysin activities are observed in fibroblasts isolated from inflamed gingiva (J. Periodontal Res., 16, 417-424(1981)). Enzyme levels have been correlated to the severity of gum disease (J. Periodontal Res., 22, 81-88 (1987)).

Collagenase-3 (matrix metalloproteinase-13: MMP-13) is

expressed in synovia of rheumatoid arthritis and
chondrocyte of human osteoarthritis (J.Clin.Invest., 97,
2011-2019(1996); J.Rheumatol., 23, 509-595(1996);
J.Biol.Chem., 271, 23577-23581(1996); J.Clin.Invest., 97,
5 761-768(1996)). MMP-13 has a strong enzyme activity against
type II collagen and aggrecan. Thus, it is suggested that
MMP-13 plays an important role in osteoarthritis and
rheumatoid arthritis (J.Biol.Chem., 271, 1544-1550(1996);
FEBS Lett., 380, 17-20(1996)). Therefore, inhibitors to
10 matrix metallo-proteinase are useful as therapeutic agents
or prophylactic drugs for joint diseases such as
osteoarthritis, rheumatoid arthritis, metastasis of tumor
cell and gingivitis.

Matrix metallo-proteinases are also concerned with
15 conversion from the latent form of tumor necrosis factor- α
to the mature form (Nature, 370, 555-557(1994)),
degradation of α 1-antitrypsin (FEBS Lett., 279, 191-
194(1991)) and processing of other matrix metallo-
proteinases (Biochemistry, 29, 10261-10670(1990);
20 J.Biol.Chem., 267, 21712-21719(1992)). Therefore,
inhibitors to matrix metalloproteinase are useful as anti-
inflammatory agents.

WO 96/20936 disclose novel thiazolidin-4-one
25 derivatives which inhibit platelet-activating factor and/or
5-lipoxygenase. But, it does not disclose the compounds can
inhibit matrix metallo-proteinases.


Description of the invention

The present invention is intended to provide novel compounds useful as matrix metallo-proteinase inhibitors.

The present inventors have earnestly examined and found that hydroxamic acid derivatives have excellent inhibition activity against matrix metallo-proteinases such as MMP-3, MMP-13 and the like. Thus, the present invention has been accomplished.

That is, the present invention is as follows:

(1) A hydroxamic acid derivative represented by the formula [1] or a prodrug thereof, or a pharmaceutically acceptable salt thereof:



wherein

X is C₁-C₂ alkylene which is substituted by optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl or optionally substituted heteroaryl; optionally substituted *ortho*-arylene; or optionally substituted *ortho*-heteroarylene;

Y¹ is -O-, -S-, -S(O)- or -S(O)₂-;

Y² is O, or S;

One of R¹ and R³ is -(CHR⁴)_n-(CR⁵R⁶)-CO-NHOH;

The other of R^1 and R^3 is hydrogen, optionally substituted alkyl, or optionally substituted cycloalkyl;

R^2 is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, or optionally substituted hetero-cycloalkyl; or R^2 and R^3 may be taken together to be optionally substituted C_1 - C_{10} alkylidene;

R^4 , R^5 and R^6 are independently hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, or optionally substituted heteraryl; or R^5 may be joined with R^4 or R^6 to form, with the carbon atom which they attach, optionally substituted cycloalkane or optionally substituted heterocycloalkane;

n is an integer of 0 to 4;

provided that when R^2 and R^3 are taken together to be optionally substituted C_1 - C_{10} alkylidene, X is not methylene substituted by a phenyl or a pyridyl wherein the phenyl and the pyridyl are optionally substituted by methyl or methoxy.

(2) A hydroxamic acid derivative according to (1) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, wherein

X is C_1 - C_2 alkylene substituted by $-Z$ -Ar; optionally


substituted *ortho*-arylene; or optionally substituted
ortho-heteroarylene;

Z is single bond or alkylene;

Ar is optionally substituted aryl or optionally
 5 substituted heteroaryl.

(3) A hydroxamic acid derivative according to (1) or a
 prodrug thereof, or a pharmaceutically acceptable salt
 thereof, wherein

10 X is one of the groups represented by the formulae:



Ar¹ is optionally substituted aryl or optionally
 substituted heteroaryl;

A¹ is *ortho*-phenylene or monocyclic *ortho*-heteroarylene;

15 R⁸ and R⁹ are independently hydrogen or substituent.


(4) A hydroxamic acid derivative according to any one of
 (1)-(3) or a prodrug thereof, or a pharmaceutically
 acceptable salt thereof, wherein Y¹ is -S-, -S(O)- or -
 20 S(O)₂-.

(5) A hydroxamic acid derivative according to any one of
 (1)-(4) or a prodrug thereof, or a pharmaceutically
 acceptable salt thereof, wherein n is 0, 1 or 2.

(6) A hydroxamic acid derivative according to any one of (1)-(5) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, wherein n is 0.

5

(7) A hydroxamic acid derivative according to (1) represented by the formula [2] or a prodrug thereof, or a pharmaceutically acceptable salt thereof:



10 wherein

Y^2 , R^4 , R^5 and R^6 are as defined in (1);

Ar^2 is optionally substituted phenyl; optionally substituted naphthyl; or optionally substituted mono or bicyclic heteroaryl;


15 Y^3 is -S-, -S(O)- or -S(O)₂-;

R^{10} and R^{11} are independently hydrogen or optionally substituted alkyl;

n_1 is an integer of 0, 1 or 2.

20 (8) A hydroxamic acid derivative according to (7) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, wherein n_1 is 0.

(9) A hydroxamic acid derivative according to (1) represented by the formula [3] or a prodrug thereof, or a pharmaceutically acceptable salt thereof:



wherein

Y^2 , R^4 , R^5 and R^6 are as defined in (1);


A^1 , R^8 , and R^9 are defined in (3);

R¹² and R¹³ are independently hydrogen or optionally substituted alkyl;

Y^3 is -S-, -S(O)- or -S(O)₂-;

n2 is an integer of 0, 1 or 2.

(10) A hydroxamic acid derivative according to (1) represented by the formula [3'] or a prodrug thereof, or a pharmaceutically acceptable salt thereof:



wherein

Y^2 , R^4 , R^5 and R^6 are as defined in (1);

A^1 is defined in (3) ;

5 Y^3 , n_2 , R^{12} and R^{13} are defined in (9);

R^{14} is hydrogen or substituent;

R^{15} is substituent

10 (11) A hydroxamic acid derivative according to (9)-(10)
or a prodrug thereof, or a pharmaceutically acceptable salt
thereof;

wherein n_2 is 0, and R^{12} and R^{13} are independently hydrogen
or substituted alkyl group.

15 (12) A hydroxamic acid derivative according to any one of
claim 1-11 or a prodrug thereof, or a pharmaceutically
acceptable salt thereof, wherein Y^2 is -O-.

20 (13) A pharmaceutical composition containing a hydroxamic
acid derivative according to any one of (1)-(12) or a
prodrug thereof, or a pharmaceutically acceptable salt
thereof, and a pharmaceutically acceptable carrier or

diluent.

(14) A method of inhibiting matrix metallo-proteinases which comprises treating the matrix metallo-proteinases with a hydroxamic acid derivative according to any one of (1)-(12) or a prodrug thereof, or a pharmaceutically acceptable salt thereof.

(15) A method of inhibiting matrix metallo-proteinases according to (14), wherein the matrix metallo-proteinases are matrix metallo-proteinase 3 and/or 13.

(16) A method of treating a disease associated with excess or undesired matrix metallo-proteinases which comprises administering an effective amount of a hydroxamic acid derivative according to any one of (1)-(12) or a prodrug thereof, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

(17) A method of treating a disease associated with excess or undesired matrix metallo-proteinases according to (16), wherein the matrix metallo-proteinases are matrix metallo-proteinase 3 and/or 13.

(18) Use of a hydroxamic acid derivative according to any one of (1)-(12) or a prodrug thereof, or a

pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a disease associated with excess or undesired matrix metallo-proteinases.

"*Ortho*-arylene" includes C₆-C₁₀ *ortho*-arylene. Typical examples are *ortho*-phenylene, 1,2-naphthalenediyl, 2,3-naphthalenediyl and the like. Preferable is *ortho*-phenylene.

"*Ortho*-heteroarylene" includes mono or bicyclic 5- or 6-membered *ortho*-heteroarylene containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms. Mono or bicyclic 5-membered *ortho*-heteroarylene includes, for example, monocyclic 5-membered *ortho*-heteroarylene containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms such as pyrrol-2,3-diyl, pyrrol-3,4-diyl, thiophen-2,3-diyl, thiophen-3,4-diyl, furan-2,3-diyl, imidazol-4,5-diyl, pyrazol-3,4-diyl, thiazol-2,3-diyl, oxazole-2,3-diyl, isothiazol-3,4-diyl, isoxazol-3,4-diyl and the like; bicyclic 5-membered *ortho*-heteroarylene containing 1 or 2 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms such as indol-2,3-diyl, benzofuran-diyl, benzothiophendiyl and the like; and the like. Mono or bicyclic 6-membered *ortho*-heteroarylene includes, for example, monocyclic 6-membered *ortho*-heteroarylene containing 1 to 3 nitrogen atoms such as pyridin-2,3-diyl,

pyridin-3,4-diyl, pyrazin-2,3-diyl, pyrimidin-4,5-diyl, pyridazin-3,4-diyl and the like; bicyclic 6-membered *ortho*-heteroarylene containing 1 to 3 nitrogen atoms such as quinolin-2,3-diyl, isoquinolin-3,4-diyl, naphthyridin-3,4-diyl, quinoxalin-2,3-diyl and the like; and the like. Preferable *ortho*-heteroarylene is monocyclic 5, or 6-membered *ortho*-heteroarylene.

"Aryl" includes C₆-C₁₀ aryl. Typical examples are phenyl, 1-naphthyl, 2-naphthyl and the like.

"Heteroaryl" includes, for example, mono or bicyclic 5- or 6-membered heteroaryl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, and the like. Mono or bicyclic 5-membered heteroaryl includes, for example, monocyclic 5-membered heteroaryl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms such as pyrrolyl, thienyl, furyl, imidazolyl, pyrazolyl, isothiazolyl, isoxazolyl, furazanyl, oxazolyl, thiazolyl, triazolyl, tetrazolyl, and the like; bicyclic 5-membered heteroaryl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms such as indolyl, isoindolyl, benzofuryl, benzothienyl, thieno[2,3-*b*]thienyl and the like; and the like. Mono or bicyclic 6-membered heteroaryl includes, for example, monocyclic 6-membered heteroaryl containing 1 to 5 nitrogen atoms such as pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl and the like; bicyclic 6-membered heteroaryl containing 1

to 5 nitrogen atoms such as quinolyl, phthalazinyl, isoquinolyl, naphthyridinyl, quinoxalinyl and the like; and the like.

5 "Substituent" of the substituted aryl, the substituted heteroaryl, the substituted ortho-arylene and substituted ortho-heteroarylene, and "substituent" of R^8 , R^9 , and R^{14} includes, for example, the following groups:

halogen; alkyl; alkenyl; alkynyl; halogenated alkyl; $-Z^1-$
 10 Ar^3 ; $-Z^1-Cy^1$; $-Z^1-N(Q^2)Q^1$; $-Z^1-N(Q^2)Ar^3$; $-Z^1-N(Q^2)Cy^1$; $-Z^1-NQ^4-C(NQ^3)N(Q^2)Q^1$; $-Z^1-NQ^4-C(NQ^3)N(Q^2)Ar^3$; $-Z^1-NQ^4-C(NQ^3)N(Q^2)Cy^1$; $-Z^1-NQ^3-CON(Q^2)Q^1$; $-Z^1-NQ^3-CON(Q^2)Ar^3$; $-Z^1-NQ^3-CON(Q^2)Cy^1$; $-Z^1-NQ^2-COOQ^1$; $-Z^1-NQ^2-COOAr^3$; $-Z^1-NQ^2-COOCy^1$; $-Z^1-OCN(Q^1)Q^2$; $-Z^1-OCN(Q^2)Ar^3$; $-Z^1-OCN(Q^2)Cy^1$; $-Z^1-OCOOQ^1$; $-Z^1-OCOOAr^3$; $-Z^1-OCOOCy^1$; $-Z^1-NQ^2-COQ^1$; $-Z^1-NQ^2-COAr^3$; $-Z^1-NQ^2-COCy^1$; $-Z^1-NQ^2-SOQ^1$; $-Z^1-NQ^2-SOAr^3$; $-Z^1-NQ^2-SOCy^1$; $-Z^1-NQ^2-SO_2Q^1$; $-Z^1-NQ^2-SO_2Ar^3$; $-Z^1-NQ^2-SO_2Cy^1$; $-Z^1-OQ^1$; $-Z^1-OAr^3$; $-Z^1-O-Cy^1$; $-Z^1-OCOQ^1$; $-Z^1-OCOAr^3$; $-Z^1-OCOCy^1$; $-Z^1-COOQ^1$; $-Z^1-COOAr^3$; $-Z^1-COOCy^1$; $-Z^1-CON(Q^1)Q^2$; $-Z^1-CON(Q^2)Ar^3$; $-Z^1-CON(Q^2)Cy^1$; $-Z^1-CON(Q^2)OQ^1$; $-Z^1-CON(Q^2)OAr^3$; $-Z^1-CON(Q^2)OCy^1$; $-Z^1-COQ^1$; $-Z^1-COAr^3$; $-Z^1-COCy^1$; $-Z^1-C(NQ^3)N(Q^1)Q^2$; $-C(NQ^3)N(Q^2)Ar^3$; $-Z^1-C(NQ^3)N(Q^2)Cy^1$; $-Z^1-SQ^1$; $-Z^1-SAr^3$; $-Z^1-S-Cy^1$; $-Z^1-SOQ^1$; $-Z^1-SOAr^3$; $-Z^1-SOCy^1$; $-Z^1-SO_2Q^1$; $-Z^1-SO_2Ar^3$; $-Z^1-SO_2Cy^1$; $-Z^1-SON(Q^1)Q^2$; $-Z^1-SON(Q^2)Ar^3$; $-Z^1-SON(Q^2)Cy^1$; $-Z^1-SO_2N(Q^1)Q^2$; $-Z^1-SO_2N(Q^2)Ar^3$; $-Z^1-SO_2N(Q^2)Cy^1$; $-Z^1-SO_3H$; $-Z^1-OSO_3H$; $-Z^1-NO_2$; $-Z^1-CN$; $-CHO$;

wherein

Z^1 is single bond, alkylene, alkenylene or alkynylene;

Q^1 , Q^2 , Q^3 , Q^4 are independently hydrogen, alkyl,

alkenyl, alkynyl, wherein the alkyl, the alkenyl and the alkynyl are optionally substituted by one or more groups selected from the following groups:

halogen; $-Ar^4$; $-Cy^2$; $-N(R^{21})R^{22}$; $-N(R^{22})Ar^4$; $-N(R^{22})Cy^2$; $-NR^{24}-C(NR^{23})N(R^{22})R^{21}$; $-NR^{24}-C(NR^{23})N(R^{22})Ar^4$; $-NR^{24}-C(NR^{23})N(R^{22})Cy^2$; $-NR^{23}-CON(R^{22})R^{21}$; $-NR^{23}-CON(R^{22})Ar^4$; $-NR^{23}-CON(R^{22})Cy^2$; $-NR^{22}-COOR^{21}$; $-NR^{22}-COOAr^4$; $-NR^{22}-COOCy^2$; $-OCON(R^{22})R^{21}$; $-OCON(R^{22})Ar^4$; $-OCON(R^{22})Cy^2$; $-OCOOR^{21}$; $-OCOOAr^4$; $-OCOOCy^2$; $-NR^{22}-COR^{21}$; $-NR^{22}-COAr^4$; $-NR^{22}-COCy^2$; $-NR^{22}-SOR^{21}$; $-NR^{22}-SOAr^4$; $-NR^{22}-SOCy^2$; $-NR^{22}-SO_2R^{21}$; $-NR^{22}-SO_2Ar^4$; $-NR^{22}-SO_2Cy^2$; $-OR^{21}$; $-OAr^4$; $-OCy^2$; $-OCOR^{21}$; $-OCOAr^4$; $-OCOCy^2$; $-COOR^{21}$; $-COOAr^4$; $-COOCy^2$; $-CON(R^{21})R^{22}$; $-CON(R^{22})Ar^4$; $-CON(R^{22})Cy^2$; $-CON(R^{22})OR^{21}$; $-CON(R^{22})OAr^4$; $-CON(R^{22})OCy^2$; $-COR^{21}$; $-COAr^4$; $-COCy^2$; $-SR^{21}$; $-SAr^4$; $-SCy^2$; $-SOR^{21}$; $-SOAr^4$; $-SOCy^2$; $-SO_2R^{21}$; $-SO_2Ar^4$; $-SO_2Cy^2$; $-SON(R^{22})R^{21}$; $-SON(R^{22})Ar^4$; $-SON(R^{22})Cy^2$; $-SO_2N(R^{22})R^{21}$; $-SO_2N(R^{22})Ar^4$; $-SO_2N(R^{22})Cy^2$; $-SO_3H$; $-OSO_3H$; $-NO_2$; $-CN$; $-CHO$;

or Q^1 may be jointed with Q^2 or Q^3 to form, with the carbon atom which they attach, optionally substituted heterocycloalkane.

R^{21} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , and R^{28} are independently hydrogen, alkyl, alkenyl, alkynyl, or R^{21} may be jointed with R^{22} , R^{23} to form, with the carbon atom which they attach, heterocycloalkane, or R^{25} may be jointed with R^{26} , R^{27} to form, with the carbon atom which they attach, heterocycloalkane.

Ar^3 and Ar^4 are independently phenyl or heteroaryl, wherein the phenyl and the heteroaryl are optionally substituted by one or two groups selected from the following groups:

halogen; alkyl; alkenyl; alkynyl; halogenated alkyl; $-Z^2-N(R^{25})R^{26}$; $-Z^2-NR^{28}-C(NR^{27})N(R^{26})R^{25}$; $-Z^2-NR^{27}-CON(R^{26})R^{25}$; $-Z^2-NR^{26}-COOR^{25}$; $-Z^2-OCON(R^{26})R^{25}$; $-Z^2-NR^{26}-COR^{25}$; $-Z^2-NR^{26}-SOR^{25}$; $-Z^2-NR^{26}-SO_2R^{25}$; $-Z^2-OR^{25}$; $-Z^2-COOR^{25}$; $-Z^2-OCOR^{25}$; $-Z^2-OCOOR^{25}$; $-Z^2-CON(R^{26})R^{25}$; $-Z^2-CON(R^{26})OR^{25}$; $-Z^2-COR^{25}$; $-Z^2-C(NR^{27})N(R^{26})R^{25}$; $-Z^2-SR^{25}$; $-Z^2-SOR^{25}$; $-Z^2-SO_2R^{25}$; $-Z^2-SON(R^{25})R^{26}$; $-Z^2-SO_2N(R^{26})R^{25}$; $-Z^2-SO_3H$; $-Z^2-OSO_3H$; $-Z^2-NO_2$; $-Z^2-CN$; $-CHO-O-CH_2-O-$; $-O-CH_2-CH_2-O-$; $-O-CH_2-CH_2-$, $-CH_2-CH_2-O-CO-$, $-CH_2-O-CO-$:

Z^2 is single bond, alkylene, alkenylene or alkynylene;

Cy^1 and Cy^2 are independently cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, wherein the cycloalkyl, the cycloalkenyl, the heterocycloalkyl,

and the heterocycloalkenyl are optionally substituted by one or two groups which are the same as the substituent of the substituted phenyl as used in Ar^3 or Ar^4 .

5

"Substituents" of R^8 and R^9 may also selected from the following groups:

$-\text{O}-\text{CH}_2-\text{O}-$; $-\text{O}-\text{CH}_2-\text{CH}_2-\text{O}-$; $-\text{O}-\text{CH}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2-\text{O}-\text{CO}-$,
 $-\text{CH}_2-\text{O}-\text{CO}-$

10

"Substituent" of R^{15} includes, for example, the following groups:

$-\text{Z}^1-\text{Ar}^{3'}$; $-\text{Z}^1-\text{Cy}^{1'}$; $-\text{Z}^1-\text{N}(\text{Q}^1)\text{Q}^2$; $-\text{Z}^1-\text{N}(\text{Q}^2)\text{Ar}^3$; $-\text{Z}^1-\text{N}(\text{Q}^2)\text{Cy}^1$; $-\text{Z}^1-\text{NQ}^4-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^2)\text{Q}^1$; $-\text{Z}^1-\text{NQ}^4-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^2)\text{Ar}^3$;
 $-\text{Z}^1-\text{NQ}^4-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^2)\text{Cy}^1$; $-\text{Z}^1-\text{NQ}^3-\text{CON}(\text{Q}^2)\text{Q}^1$; $-\text{Z}^1-\text{NQ}^3-\text{CON}(\text{Q}^2)\text{Ar}^3$; $-\text{Z}^1-\text{NQ}^3-\text{CON}(\text{Q}^2)\text{Cy}^1$;
 $-\text{Z}^1-\text{NQ}^2-\text{COOQ}^1$; $-\text{Z}^1-\text{NQ}^2-\text{COOAr}^3$; $-\text{Z}^1-\text{NQ}^2-\text{COOCy}^1$; $-\text{Z}^1-\text{OCON}(\text{Q}^1)\text{Q}^2$; $-\text{Z}^1-\text{OCON}(\text{Q}^2)\text{Ar}^3$;
 $-\text{Z}^1-\text{OCON}(\text{Q}^2)\text{Cy}^1$; $-\text{Z}^1-\text{NQ}^2-\text{COQ}^1$; $-\text{Z}^1-\text{NQ}^2-\text{COAr}^3$; $-\text{Z}^1-\text{NQ}^2-\text{COCy}^1$;
 $-\text{Z}^1-\text{NQ}^2-\text{SOQ}^1$; $-\text{Z}^1-\text{NQ}^2-\text{SOAr}^3$; $-\text{Z}^1-\text{NQ}^2-\text{SOCy}^1$; $-\text{Z}^1-\text{NQ}^2-\text{SO}_2\text{Q}^1$;
 $-\text{Z}^1-\text{NQ}^2-\text{SO}_2\text{Ar}^3$; $-\text{Z}^1-\text{NQ}^2-\text{SO}_2\text{Cy}^1$; $-\text{Z}^1-\text{OQ}^{1'}$; $-\text{Z}^1-\text{OCOQ}^1$;
 $-\text{Z}^1-\text{OCOAr}^3$; $-\text{Z}^1-\text{OCOCy}^1$; $-\text{Z}^1-\text{OCOOQ}^1$; $-\text{Z}^1-\text{OCOOAr}^3$;
 $-\text{Z}^1-\text{OCOOCy}^1$; $-\text{Z}^1-\text{COOQ}^1$; $-\text{Z}^1-\text{COOAr}^3$; $-\text{Z}^1-\text{COOCy}^1$; $-\text{Z}^1-\text{CON}(\text{Q}^1)\text{Q}^2$;
 $-\text{Z}^1-\text{CON}(\text{Q}^2)\text{Ar}^3$; $-\text{Z}^1-\text{CON}(\text{Q}^2)\text{Cy}^1$; $-\text{Z}^1-\text{CON}(\text{Q}^1)\text{OQ}^2$;
 $-\text{Z}^1-\text{CON}(\text{Q}^2)\text{OAr}^3$; $-\text{Z}^1-\text{CON}(\text{Q}^2)\text{OCy}^1$; $-\text{Z}^1-\text{COQ}^1$;
 $-\text{Z}^1-\text{COAr}^3$; $-\text{Z}^1-\text{CO}-\text{Cy}^1$; $-\text{Z}^1-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^1)\text{Q}^2$; $-\text{Z}^1-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^2)\text{Ar}^3$;
 $-\text{Z}^1-\text{C}(\text{NQ}^3)\text{N}(\text{Q}^2)\text{Cy}^1$; $-\text{Z}^1-\text{SQ}^{1'}$; $-\text{Z}^1-\text{SOQ}^1$;
 $-\text{Z}^1-\text{SOAr}^3$; $-\text{Z}^1-\text{SOCy}^1$; $-\text{Z}^1-\text{SO}_2\text{Q}^1$; $-\text{Z}^1-\text{SO}_2\text{Ar}^3$; $-\text{Z}^1-\text{SO}_2\text{Cy}^1$;

15

20

25

$-Z^1-SO_2N(Q^1)Q^2$; $-Z^1-SO_2N(Q^2)Ar^3$; $-Z^1-SO_2N(Q^2)Cy^1$; $-Z^1-SO_2N(Q^1)Q^2$; $-Z^1-SO_2N(Q^2)Ar^3$; $-Z^1-SO_2N(Q^2)-Cy^1$; $-CHO$;
wherein

Q^1 , Q^2 , Q^3 , Q^4 , R^{21} , R^{22} , R^{23} , R^{24} , Z^1 , Ar^3 , Ar^4 , Cy^1 and Cy^2 are defined above.

$Q^{1'}$ is hydrogen, alkyl, alkenyl, or alkynyl, wherein the alkyl, alkenyl, and alkynyl are substituted by one or more groups selected from the following groups:

$-Ar^{4'}$; $-Cy^{2'}$; $-N(R^{21})R^{22}$; $-N(R^{22})Ar^4$; $-N(R^{22})Cy^2$; $-NR^{24}-C(NR^{23})N(R^{22})R^{21}$; $-NR^{24}-C(NR^{23})N(R^{22})Ar^4$; $-NR^{24}-C(NR^{23})N(R^{22})Cy^2$; $-NR^{23}-CON(R^{22})R^{21}$; $-NR^{23}-CON(R^{22})Ar^4$; $-NR^{23}-CON(R^{22})Cy^2$; $-NR^{22}-COOR^{21}$; $-NR^{22}-COOAr^4$; $-NR^{22}-COOCy^2$; $-OCONR^{21}$; $-OCONAr^4$; $-OCONCy^2$; $-NR^{22}-COR^{21}$; $-NR^{22}-COAr^4$; $-NR^{22}-COCy^2$; $-NR^{22}-SOR^{21}$; $-NR^{22}-SOAr^4$; $-NR^{22}-SOCy^2$; $-NR^{22}-SO_2R^{21}$; $-NR^{22}-SO_2Ar^4$; $-NR^{22}-SO_2Cy^2$; $-OCOR^{21}$; $-OCOAr^4$; $-OCOCy^2$; $-COOR^{21}$; $-COOAr^4$; $-COOCy^2$; $-OCOOR^{21}$; $-OCOOAr^4$; $-OCOOCy^2$; $-CON(R^{21})R^{22}$; $-CON(R^{22})Ar^4$; $-CON(R^{22})Cy^2$; $-CON(R^{22})OR^{21}$; $-CON(R^{22})OAr^4$; $-CON(R^{22})OCy^2$; $-COR^{21}$; $-COAr^4$; $-COCy^2$; $-SOR^{21}$; $-SOAr^4$; $-SOCy^2$; $-SO_2R^{21}$; $-SO_2Ar^4$; $-SO_2Cy^2$; $-SON(R^{22})R^{21}$; $-SON(R^{22})Ar^4$; $-SON(R^{22})Cy^2$; $-SO_2N(R^{22})R^{21}$; $-SO_2N(R^{22})Ar^4$; $-SO_2N(R^{22})Cy^2$; $-CN$; $-CHO$; $-OH$; $-SH$

$Ar^{3'}$ and $Ar^{4'}$ are heteroaryl, wherein the heteroaryl are optionally substituted by one or two groups which are the same as the substituent of the substituted phenyl

as used in Ar³ or Ar⁴.

Cy^{1'} and Cy^{2'} are heterocycloalkyl or heterocycloalkenyl, wherein the heterocycloalkyl and the heterocycloalkenyl is optionally substituted by one or two groups which are the same as the substituent of the substituted phenyl as used in Ar³ or Ar⁴.

The number of "substituent" of the substituted aryl, the substituted heteroaryl, the substituted ortho-arylene and substituted ortho-heteroarylene, and the number of "substituent" of R⁸, R⁹, R¹⁴, and R¹⁵ are one, two or three, preferably one or two.

"Alkyl" includes straight or branched C₁-C₆ alkyl. Typical examples are methyl, ethyl, propyl, 2-methylethyl, butyl, 2-methylpropyl, 1-methylpropyl, 1,1-dimethylethyl, 2,2-dimethylpropyl, pentyl, hexyl and the like.

"Halogenated alkyl" includes straight or branched C₁-C₆ alkyl substituted by one or more of halogens. Typical examples are trifluoromethyl, pentafluoroethyl, 2-chloroethyl, 3-bromopropyl, 5-fluoropentyl, 4-iodohexyl and the like.

"Alkoxy" includes straight or branched C₁-C₆ alkoxy. Typical examples are methoxy, ethoxy, propoxy, 2-methylethoxy, butoxy, 2-methylpropoxy, 1-methylpropoxy, 1,1-dimethylethoxy, 2,2-dimethylpropoxy, pentyloxy,

hexyloxy, and the like.

"Alkylthio" includes straight or branched C₁-C₆ alkylthio. Typical examples are methylthio, ethylthio, propylthio, 2-methylethylthio, butylthio, 2-methylpropylthio, 1-methylpropylthio, 1,1-dimethylethylthio, 2,2-dimethylpropylthio, pentylthio, hexylthio, and the like.

"Alkenyl" includes straight or branched C₂-C₆ alkenyl. Typical examples are vinyl, allyl, 1-propenyl, 2-butenyl and the like.

"Alkynyl" includes straight or branched C₂-C₆ alkynyl. Typical examples are ethynyl, propargyl, 1-propynyl, 2-butyne, pentynyl and the like.

"Alkylene" includes straight or branched C₁-C₆ alkylene. Typical examples are methylene, ethylene, trimethylene, 2-methyltrimethylene, tetramethylene, pentamethylene, hexamethylene and the like.

"Alkenylene" includes straight or branched C₂-C₆ alkenylene. Typical examples are vinylene, propenylene, 2-butenylene, 2-methyl-2-butenylene, 3-pentenylene, 3-hexenylene and the like.

"Alkynylene" includes straight or branched C₂-C₆ alkynylene. Typical examples are ethynylene, propynylene, 2-butyne, 3-pentynylene, 2-methyl-3-pentynylene, 3-hexynylene and the like.

"C₁-C₂ alkylene" is methylene or ethylene.

"C₁-C₁₀ alkylidene" includes straight or branched C₁-C₁₀ alkylidene, preferably straight or branched C₁-C₆ alkylidene.

Typical examples are methylenidene, ethylenidene, propylenidene, butylenidene, 2-methylbutylenidene, hexylenidene, octylenidene, nonylenidene, decylenidene and the like. Preferable is methylenidene.

5 "Cycloalkane" includes C₃-C₈ cycloalkane, which can be containing 0-2 carbonyl group. Typical examples are cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, cyclohexanone and the like.

10 "Heterocycloalkane" includes, for example, 5- or 6-membered heterocycloalkane containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, and the like. The sulfur atoms of "Heterocycloalkane" may be oxidized to form sulfoxide, or sulfone.

15 "Heterocycloalkane" may also be containing 0-2 carbonyl groups. Typical examples of 5-membered heterocycloalkane containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are pyrrolidine, imidazolidine, pyrazolidine, tetrahydrofuran, tetrahydrothiophene, dioxolane, pyrrolidinone, and the like.
20 Typical examples are 6-membered heterocycloalkane containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are piperidine, piperazine, morpholine, tetrahydropyran, dioxane, thiomorpholine, 4-piperidone and the like.
25

"Cycloalkyl" includes C₃-C₈ cycloalkyl, which may

contain 0-2 carbonyl group. Typical examples are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclohexanone-4-yl and the like.

5 "Cycloalkenyl" includes C₃-C₈ cycloalkenyl, which may contain 0-2 carbonyl group. Typical examples are cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclopenten-3-one-yl and the like.

10 "Heterocycloalkyl" includes, for example, 5- or 6-membered heterocycloalkyl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, and the like. The sulfur atoms of "Heterocycloalkyl" may be oxidized to form sulfoxide or sulfone.

15 "Heterocycloalkyl" may also be containing 0-2 carbonyl groups. Typical examples are 5-membered heterocycloalkyl containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are pyrrolidinyl, 2-pyrrolidinone-1-yl, imidazolidinyl, pyrazolidinyl, tetrahydrofuryl, tetrahydrothienyl, dioxolanyl and the like. Typical examples are 6-membered heterocycloalkyl containing 1 or 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are piperidyl, piperazinyl, morpholinyl, 20 tetrahydropyranyl, dioxanyl, thiomorpholinyl, 4-piperidone-1-yl and the like.

25 "Heterocycloalkenyl" includes, for example, 5- or 6-

membered heterocycloalkenyl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, and the like. The sulfur atoms of "Heterocycloalkenyl" may be oxidized to form sulfoxide or sulfone. "Heterocycloalkenyl" may also be containing 0-2 carbonyl groups. Typical examples are 5-membered heterocycloalkenyl containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are pyrrolinyl, imidazolinyl, pyrazolinyl, dihydrofuryl, 5-pyrazolone-4-yl and the like. Typical examples are 6-membered heterocycloalkyl containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms and oxygen atoms, are 2,3-dihydropyridyl-, 4-pyridone-1-yl and the like.

"Substituent" of the substituted alkyl, the substituted alkenyl, the substituted alkynyl, the substituted alkoxy, the substituted alkylthio, the substituted cycloalkyl, the substituted heterocycloalkyl, the cycloalkenyl, the substituted heterocycloalkenyl, the substituted cycloalkane, the substituted heterocycloalkane and the substituted C₁-C₁₀ alkylidene includes, for example, the following groups:

halogen; -Ar⁵; -Cy³; -N(Q⁵)Q⁶; -N(Q⁶)Ar⁵; -N(Q⁶)Cy³; -NQ⁸-C(NQ⁷)N(Q⁶)Q⁵; -NQ⁸-C(NQ⁷)N(Q⁶)Ar⁵; -NQ⁸-C(NQ⁷)N(Q⁶)Cy³; -NQ⁷-CON(Q⁶)Q⁵; -NQ⁶-CON(Q⁵)Ar⁵; -NQ⁷-CON(Q⁶)Cy³; -NQ⁷-COOQ⁶; -NQ⁷-COOAr⁵; -NQ⁷-COOCy³; -OCON(Q⁶)Q⁵; -OCON(Q⁶)Ar⁵; -OCON(Q⁶)Cy³; -NQ⁶-COQ⁵; -NQ⁶-COAr⁵; -NQ⁶-COCy³; -NQ⁶-SOQ⁵;

$-\text{NQ}^6-\text{SOAr}^5$; $-\text{NQ}^6-\text{SOCy}^3$; $-\text{NQ}^6-\text{SO}_2\text{Q}^5$; $-\text{NQ}^6-\text{SO}_2\text{Ar}^5$; $-\text{NQ}^6-\text{SO}_2\text{Cy}^3$; $-\text{OQ}^5$; $-\text{OAr}^5$; $-\text{OCy}^3$; $-\text{OCOQ}^5$; $-\text{OCOAr}^5$; $-\text{OCOCy}^3$; $-\text{COOQ}^5$; $-\text{COOAr}^5$; $-\text{COOCy}^3$; $-\text{OCOOQ}^5$; $-\text{OCOOAr}^5$; $-\text{OCOOCy}^3$; $-\text{CON}(\text{Q}^6)\text{Q}^5$; $-\text{CON}(\text{Q}^6)\text{Ar}^5$; $-\text{CON}(\text{Q}^6)\text{Cy}^3$; $-\text{CON}(\text{Q}^6)\text{OQ}^5$; $-\text{CON}(\text{Q}^6)\text{OAr}^5$; $-\text{CON}(\text{Q}^6)\text{OCy}^3$; $-\text{COQ}^5$; $-\text{COAr}^5$; $-\text{COCy}^3$; $-\text{SQ}^5$; $-\text{SAr}^5$; $-\text{SCy}^3$; $-\text{SOQ}^5$; $-\text{SOAr}^5$; $-\text{SOCy}^3$; $-\text{SO}_2\text{Q}^5$; $-\text{SO}_2\text{Ar}^5$; $-\text{SO}_2\text{Cy}^3$; $-\text{SON}(\text{Q}^6)\text{Q}^5$; $-\text{SON}(\text{Q}^6)\text{Ar}^5$; $-\text{SON}(\text{Q}^6)\text{Cy}^3$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Q}^5$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Ar}^5$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Cy}^3$; $-\text{SO}_3\text{H}$; $-\text{NO}_2$; $-\text{CN}$;

wherein

R^{29} , R^{30} , R^{31} , and R^{32} are independently hydrogen, alkyl, alkenyl, alkynyl;

Q^5 , Q^6 , Q^7 , and Q^8 are independently hydrogen, alkyl, alkenyl and alkynyl wherein the alkyl, the alkenyl and the alkynyl are optionally substituted by one or more groups selected from the following group:

halogen; $-\text{Ar}^6$; $-\text{Cy}^4$; $-\text{N}(\text{R}^{29})\text{R}^{30}$; $-\text{N}(\text{R}^{30})\text{Ar}^6$; $-\text{N}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{R}^{29}$; $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{Ar}^6$; $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{R}^{29}$; $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{Ar}^6$; $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{30}-\text{COOR}^{29}$; $-\text{NR}^{30}-\text{COOAr}^6$; $-\text{NR}^{30}-\text{COOCy}^4$; $-\text{OCON}(\text{R}^{30})\text{R}^{29}$; $-\text{OCON}(\text{R}^{30})\text{Ar}^6$; $-\text{OCON}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{30}-\text{COR}^{29}$; $-\text{NR}^{30}-\text{COAr}^6$; $-\text{NR}^{30}-\text{COCy}^4$; $-\text{NR}^{30}-\text{SOR}^{29}$; $-\text{NR}^{30}-\text{SOAr}^6$; $-\text{NR}^{30}-\text{SOCy}^4$; $-\text{NR}^{30}-\text{SO}_2\text{R}^{29}$; $-\text{NR}^{30}-\text{SO}_2\text{Ar}^6$; $-\text{NR}^{30}-\text{SO}_2\text{Cy}^4$; $-\text{OR}^{29}$; $-\text{OAr}^6$; $-\text{OCy}^4$; $-\text{COOR}^{29}$; $-\text{COOAr}^6$; $-\text{COOCy}^4$; $-\text{OCOR}^{29}$; $-\text{COAr}^5$; $-\text{OCOCy}^3$; $-\text{OCOOR}^{29}$; $-\text{OCOOAr}^6$; $-\text{Z}^1-\text{OCOOCy}^4$; $-\text{CON}(\text{R}^{30})\text{R}^{29}$; $-\text{CON}(\text{R}^{30})\text{Ar}^6$; $-\text{CON}(\text{R}^{30})\text{Cy}^4$; $-\text{CON}(\text{R}^{30})\text{OR}^{29}$; $-\text{CON}(\text{R}^{30})\text{OAr}^6$; $-\text{CON}(\text{R}^{30})\text{OCy}^4$; $-\text{COR}^{29}$;

COAr⁶; -COCy⁴; -SR²⁹; -SAr⁶; -SCy⁴; -SOR²⁹; -SOAr⁶; -
 SOCy⁴; -SO₂R²⁹; -SO₂Ar⁶; -SO₂Cy⁴; -SON(R³⁰)R²⁹; -
 SON(R³⁰)Ar⁶; -SON(R³⁰)Cy⁴; -SO₂N(R³⁰)R²⁹; -
 SO₂N(R³⁰)Ar⁶; -SO₂N(R³⁰)Cy⁴; -SO₃H; -NO₂; -CN;

or Q⁵ may be jointed with Q⁶ or Q⁷ to form, with the
 carbon atom which they attach, optionally substituted
 heterocycloalkane.

Ar⁵ and Ar⁶ are independently aryl or heteroaryl,
 wherein the aryl and the heteroaryl are optionally
 substituted by one or two groups which are the same as
 the "substituent" of the substituted aryl, the
 substituted heteroaryl, the substituted ortho-arylene
 and substituted ortho-heteroarylene, and "substituent"
 of R⁸, R⁹, and R¹⁴;

Cy³ and Cy⁴ are independently cycloalkyl, cycloalkenyl,
 heterocycloalkyl, or heterocycloalkenyl, wherein the
 cycloalkyl, the cycloalkenyl, the heterocycloalkyl,
 and the heterocycloalkenyl are optionally substituted
 by one or two groups which are the same as the
 substituent of the substituted phenyl as used in Ar³
 or Ar⁴.

The number of "substituent" of substituted alkyl is one
 to five, preferably one, two, or three.

When R¹ is -(CHR⁴)_n-(CR⁵R⁶)-CO-NHOH in formula 1, -
 (CHR⁴)_{n1}-(CR⁵R⁶)-CONHOH in formula 2, -(CHR⁴)_{n2}-(CR⁵R⁶)-CO-
 NHOH in formula 3 and 3', and R⁵ or R⁶ are 1-substituted

alkyl in formula 1, 2, 3, and 3', the substituents of R^5 or R^6 do not include the following groups:


$-\text{CON}(\text{Q}^6)\text{Q}^5$; $-\text{CON}(\text{Q}^6)\text{Ar}^5$; $-\text{CON}(\text{Q}^6)\text{Cy}^3$ $-\text{CON}(\text{Q}^6)\text{OQ}^5$; $-\text{CON}(\text{Q}^6)\text{OAr}^5$; $-\text{CON}(\text{Q}^6)\text{OCy}^3$; $-\text{CO-Q}^5$

wherein

Q^5 , Q^6 , Ar^5 , and Cy^3 are described above.

"Halogen" includes fluorine, chlorine, bromine and iodine. Typical examples are chlorine and fluorine.

The preferable examples of $-(\text{CHR}^4)_n-(\text{CR}^5\text{R}^6)-\text{CO-NHOH}$ as used in one of R^1 and R^3 in formula 1; $-(\text{CHR}^4)_{n1}-(\text{CR}^5\text{R}^6)-\text{CO-NHOH}$ as used in formula 2; or $-(\text{CHR}^4)_{n2}-(\text{CR}^5\text{R}^6)-\text{CO-NHOH}$ as used in formula 3 are



wherein R^5 is hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, and optionally substituted heterocycloalkyl, and more preferably hydrogen or optionally substituted alkyl;

n₃ is an integer of 0 to 3; n₄ is an integer of 0 to 2,
and more preferable n₃ and n₄ is 0;

A³ is cyclopentane or cyclohexane;

and A⁴ is cyclopentane, cyclohexane, tetrahydropyran,
piperazine and the like.

The preferable R⁶ in formula 1, 2, 3, and 3' is hydrogen
or C₁-C₃ alkyl such as methyl, ethyl.

The more preferable example of R⁵ in formula 2 is the
optionally substituted alkyl. The functional groups of the
"substituents" of substituted alkyl, such as hydroxy,
carboxyl, amino, or thiol, may be protected with typical
protective groups described in "protective groups in
organic synthesis 2nd edition by W. Greene (John Wiley &
Sons, INC.)".

The more preferable R⁵ in formula 3 and 3' is hydrogen
or alkyl such as methyl, ethyl, propyl, and isopropyl.

The preferable examples of the hydroxamic acid
derivative represented by formula 1 are

(1) the derivatives wherein R¹ is -(CHR⁴)_n-(CR⁵R⁶)-CONHOH;
and R³ is hydrogen, or optionally substituted alkyl ;
and

(2) the derivatives wherein R¹ is alkyl substituted by a)
halogen, b) optionally substituted cycloalkyl, c)
optionally substituted aryl, d) optionally substituted
hetero-cycloalkyl or e) optionally substituted
heteroaryl; and R³ is -(CHR⁴)_n-(CR⁵R⁶)-CONHOH;

wherein n , R^4 , R^5 , and R^6 are as defined above.

The preferable example of R^2 in formula 1; R^{10} in formula 2; and R^{12} in formula 3 and 3' are hydrogen or alkyl such as methyl and ethyl.

The preferable examples of the hydroxamic acid derivative represented by formula 2 are the derivatives wherein R^{11} is hydrogen or optionally substituted alkyl, such as methyl, ethyl, propyl or butyl in which the

"substituents" are selected from the following group;

$-NQ^6-COOQ^5$; $-NQ^6-COOAr^5$; $-NQ^6-COOCy^3$; $-NQ^6-COQ^5$; $-NQ^6-COAr^5$; $-NQ^6-COCy^3$; $-NQ^6-SOQ^5$; $-NQ^6-SOAr^5$; $-NQ^6-SOCy^3$; $-NQ^6-SO_2Q^5$; $-NQ^6-SO_2Ar^5$; $-NQ^6-SO_2Cy^3$; $-COOQ^5$; $-COOAr^5$; $-COOCy^3$; $-CON(Q^6)Q^5$; $-CON(Q^6)Ar^5$; $-CON(Q^6)Cy^3$; $-CON(Q^6)OQ^5$; $-CON(Q^6)OAr^5$; $-CON(Q^6)OCy^3$; $-COQ^5$; $-COAr^5$; $-COCy^3$; $-SO-Q^5$; $-SO-Ar^5$; $-SO-Cy^3$; $-SO_2-Q^5$; $-SO_2-Ar^5$; $-SO_2-Cy^3$; $-SON(Q^6)-Q^5$; $-SON(Q^6)-Ar^5$; $-SON(Q^6)-Cy^3$; $-SO_2N(Q^6)-Q^5$; $-SO_2N(Q^6)-Ar^5$; $-SO_2N(Q^6)-Cy^3$;

wherein

Ar^5 , Cy^3 , Q^5 and Q^6 are described above.

The more preferable "substituents" are selected from the following group;

$-COOQ^5$; $-COOAr^5$; $-COOCy^3$; $-CON(Q^6)Q^5$; $-CON(Q^6)Ar^5$; $-CON(Q^6)Cy^3$; $-CON(Q^6)OQ^5$; $-CON(Q^6)OAr^5$; $-CON(Q^6)OCy^3$; $-COQ^5$; $-CO-Ar^5$; $-CO-Cy^3$; $-COO-Q^5$; $-COO-Ar^5$; $-COO-Cy^3$;

wherein

Ar^5 , Cy^3 , Q^5 and Q^6 are described above.

The preferable example of the hydroxamic acid derivative represented by formula 3 and 3' are the derivatives wherein R^{13} is optionally substituted alkyl, such as methyl, ethyl, or propyl. The "substituents" are selected from the following groups;

$-Ar^5$; $-Cy^3$; $-O-Q^9$; $-OAr^5$; $-N(Q^5)Ar^5$; $-S-Q^9$; $-SAr^5$;

wherein

Ar^5 , Cy^3 , and Q^5 are described above;

Q^9 is alkyl, alkenyl or alkynyl wherein the alkyl, the alkenyl and the alkynyl are substituted by one or more groups selected from the following group:

$-Ar^6$; Cy^4 ; $-OAr^6$; $-SAr^6$;

wherein

Ar^6 , and Cy^4 are described above;

The preferable Ar^5 and Ar^6 in R^{13} in formula 3 and 3' are optionally substituted phenyl or optionally substituted monocyclic heteroaryl such as pyridyl, thienyl, furyl and pyrrolyl. The preferable "substituents" are selected from the following groups;

halogen; halogenated alkyl; $-OQ^1$; $-OAr^3$; $-SQ^1$; or $-SAr^3$ wherein Q^1 and Ar^3 are as defined above.

The preferable positions of the substituents of the substituted phenyl and the substituted monocyclic heteroaryl in R^{13} in formula 3 and 3' are *m*-position or *p*-position, especially *p*-position.

The preferable Ar^2 in formula 2 are optionally substituted phenyl, optionally substituted naphthyl, optionally substituted monocyclic heteroaryl, or optionally substituted bicyclic heteroaryl, wherein the "substituents" are selected from the following groups;

halogen, alkyl, halogenated alkyl, $-\text{OQ}^1$, $-\text{OAr}^3$, $-\text{SQ}^1$, $-\text{SAr}^3$, $-\text{SOQ}^1$, $-\text{SOAr}^3$, $-\text{SO}_2\text{Q}^1$, $-\text{SO}_2\text{Ar}^3$;


wherein

Ar^3 and Q^1 are defined above.

The preferable positions of the substituent of the substituted phenyl and the substituted monocyclic heteroaryl as used in Ar^2 in formula 2 are *m*-position or *p*-position, especially *p*-position. More preferable Ar^2 in formula 2 are phenyl or monocyclic heteroaryl.

The preferable A1 in formula 3 and 3' are phenylene or monocyclic heteroarylene, such as pyrazole-3,4-diyl, imidazole-4,5-diyl, and thiophene-2,3-diyl.

When the A1 in formula 3 and 3' is phenylene, the preferable positions of R^8 or R^9 in formula 3, and R^{14} or R^{15} in formula 3' are 6th and/or 7th positions.



The hydroxamic acid derivatives can be prepared through three processes:

[A] cyclization process;

[B] substitution process;


5 [C] process of forming hydroxamic acid group.

The order of three processes may be the same as or different from the order [A], [B] and [C] as long as the hydroxamic acid derivatives can be prepared. And protective groups may be used through these processes to protect functional groups in the hydroxamic acid derivative.

Protective groups include well known groups described in "Protective Groups in Organic Syntheses" 2nd ed. T.W.Greene and P.G.M. Wuts, John Wiley & Sons, Inc. 1991.

15 [A] Cyclization process

(1) 5-Membered ring



wherein R^A , R^B and R^L are independently hydrogen atom or substituent; and Y^2 is -O- or -S-.


20 5-Membered ring can be formed for example by cyclization reaction of imine with mercapto- or hydroxy-carboxylic acid by the same method as ones described in Chem. Rev., Vol.81, 175 (1981). One to 5 equivalents of mercapto- or hydroxy-carboxylic acid per equivalent of the

imine is preferably used. In this reaction, dehydration agent such as molecular sieves 3A (MS 3A) or triethyl orthoformate may be added. Reaction solvent includes for example ether such as THF; halogenated hydrocarbon such as chloroform, dichloromethane; amide such as DMF, N-methylpyrrolidone; sulfone or sulfoxide such as tetramethylene sulfone, DMSO; hydrocarbon such as benzene, toluene, xylene, heptane, hexane; or mixture thereof. Reaction temperature is usually 50 °C to boiling point of the solvent. The imine can be prepared by a conventional method.

(2) 6-Membered ring

6-Membered ring can be formed for example by the same methods as ones described in EP 162776 A ; Chem. Pharm. Bull., 39, 2888 (1991); *ibid.*, 42, 1264 (1994); *ibid.*, 44, 2055 (1996).

Method 1




wherein R^{C} , R^{D} and R^{M} are independently hydrogen atom or substituent, or $\text{R}^{\text{C}}-\text{C}-\text{C}-\text{R}^{\text{D}}$ is optionally substituted *ortho*-arylene or optionally substituted *ortho*-heteroarylene; and X^1 is chlorine, bromine or iodine.

6-Membered ring can be formed by amidation reaction of hydroxy-amine with halo-carboxylic acid or halo-ester,

followed by ether formation reaction. Amidation reaction can be performed by a conventional method such as dehydration method, mixed anhydride method, activated ester method and the like ("The Peptides Analysis, Synthesis, Biology", Vol.1, 2, 3, 5, ed. by E. Gross, J. Meinhofer Academic Press (1979)). Dehydration agent used in the dehydration method includes dicyclohexylcarbodiimide, N,N-dimethylaminopropyl-N'-ethylcarbodiimide, 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline, O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate. Acylation agent used in mixed anhydride method includes isobutyl chloroformate, pivaloyl chloride and the like. Ether formation reaction can be carried out by a conventional method.

Method 2




wherein R^{C} , R^{D} , R^{M} and X^1 are as defined above; and R^{E} is hydrogen or alkyl.

6-Membered ring can be formed by thioether formation reaction of mercaptoamine and halide, followed by amidation reaction. Thioether formation reaction may be performed in the presence of base such as K_2CO_3 , NaOH and triethylamine. 2-Mercapto-anilines may be commercially available or can be

prepared from 2-aminothiazoles, 2-methylthiazoles, or dithiazolium chloride by the same method as one described in Chem. Pharm. Bull., 39, 2888 (1991); *ibid*, 42, 1264 (1994); *ibid*, 44, 2055 (1996); J. Chem. Soc., 1948, 870.

The amidation reaction can be carried out by heating in the presence of acid such as HCl, HBr, H₂SO₄, acetic acid and methanesulfonic acid, or by a conventional method described above.


Method 3



wherein R^C, R^D, Y², R^E, R^M and X¹ are as defined above.

6-Membered ring can be formed by ether or thioether formation reaction of nitrohalide and thiol or alcohol, followed by reduction and amidation reaction. Ether or thioether formation reaction may be performed in the presence of base such as K₂CO₃, NaOH and triethylamine. Reduction may be carried out by a conventional method. Amidation reaction may be carried out by heating in the presence of acid or by a conventional method described above.

Method 4



wherein R^{C} , R^{D} , R^{M} , Y^2 and X^1 are as defined above; and X^2 is chlorine, bromine or iodine; and R^{F} is hydrogen or substituent.

6-Membered ring can be formed by ether or thioether formation reaction of alcohol or thiol and halide, followed by cyclization reaction. Ether or thioether formation reaction can be carried out by the method described above. Cyclization reaction can be performed in the presence of Pd catalyst by the same method as one described in J. Am. Chem. Soc., 119, 8451-8458 (1997). Base such as Cs_2CO_3 may be added in this reaction.

[B] Substitution process


(1) Substitution at the position of R^1 in the formula [1]

Introducing a substituent at the position of R^1 in the formula [1] can be performed for example by alkylation. Alkylation may be performed by reacting with corresponding alkyl halide, preferably alkyl iodide or bromide, in the presence of base such as NaH, NaOMe, potassium *t*-butoxide and Na_2CO_3 . Reaction solvent is for example ether such as THF; amide such as DMF; sulfoxide such as DMSO or mixture

thereof, preferably THF and DMF. Reaction temperature is usually 0 °C to 160 °C.


(2) Substitution at the position of R² and R³ in the formula [1]

Introducing substituent(s) at the position of R² and R³ in the formula [1] can be performed for example by aldol reaction, Wittig reaction, alkylation, Michael reaction and the like.



wherein R¹, X and Y are as defined above; and R^G is hydrogen or substituent.

Aldol reaction may be performed by reacting with the corresponding aldehyde in the presence of base or acid (J. Med. Chem., 20, 729 (1977), EP 657444 (A), JP 7-233155 (A)).



wherein R¹, X, Y and R^G are as defined above.

Wittig reaction may be performed by reacting with the corresponding aldehyde (J. Org. Chem., 40, 1731 (1975), US 3873535, US 3923709).

(2) Introduction of sulfoxide and sulfone


Introduction of sulfoxide can be carried out by oxidizing thioether group with mild oxidizing agent such as hydrogen peroxide, sodium periodate and the like.

Introduction of sulfone can be carried out by oxidizing thioether or sulfoxide with oxidizing agent such as *m*-chloroperbenzoic acid(*m*-CPBA) and potassium peroxymonosulfate.

[C] Process of forming hydroxamic acid group

Hydroxamic acid group can be formed for example by (1) amidation of ester group with hydroxylamine (J. Med. Chem., 40, 2525 (1997)) or (2) condensation of carboxylic acid group with protected hydroxylamine followed by deprotection (J. Med. Chem., 41, 1209 (1998); *Ibid.*, 41, 1745 (1998); *Ibid.*, 38, 2570 (1995)).


(1) Amidation of ester group with hydroxylamine



wherein R^{H} is optionally substituted alkyl such as methyl, ethyl and benzyl.

Preferred R^{H} is methyl or ethyl. Two to 50 equivalents of hydroxylamine per equivalent of the ester group is preferably used. Reaction solvent is for example alcohol such as methanol, ethanol; amide such as DMF; sulfoxide such as DMSO; ketone such as acetone; water or mixture thereof, especially alcohol. Reaction temperature is usually 0 °C to 80 °C.

(2) Condensation of carboxylic acid group with protected hydroxylamine followed by deprotection



wherein R^{J} is a protective group of hydroxylamine such as *t*-butyldimethylsilyl, trimethylsilyl, *t*-butyl, benzyl, 4-methoxybenzyl and tetrahydropyranyl.

Condensation of carboxylic acid group with protected hydroxylamine can be performed by a conventional method as described above. The protective group of the protected hydroxamic acid group can be removed by a conventional method ("Protective Groups in Organic Syntheses" 2nd ed. T.W.Greene and P.G.M. Wuts, John Wiley & Sons, Inc. 1991).

The prodrug of the hydroxamic acid derivative includes the prodrugs described in Chemistry and Industry, 1980, 435; Advanced Drug Discovery Reviews 3, 39(1989). The typical examples are biohydrolyzable esters such as acyloxymethyl esters, glycolates, lactates and morpholinoethyl ester of carboxyl group; hemiglutarates of phenolic hydroxyl group; N-morpholinomethyl amides; N-acyloxymethyl amines; N-ayloxyalkoxycarbonylamines.

The present invention includes every isomers such as diastereomers, enantiomers, and geometrical isomers, if the

hydroxamic acid derivative has such isomers.

Method of optical resolution of a present invention compound or an intermediate thereof which has acidic group comprises steps of

5 (1) a step of forming a salt of a compound of present invention or intermediate thereof and an optically active base;

(2) a step of recrystallization.

Examples of the solvent forming a salt are an alcohol
10 solvent (such as methanol, ethanol 2-propanol and the like), an ether solvent (such as diethyl ether and the like), an aromatic hydrocarbon solvent (such as toluene and the like), an aprotic solvent (such as acetonitrile), and a mixture of the solvent described before. Examples of the optical
15 active base are an organic amine (such as α -phenethylamine, quinine, quinidine, cinchonidine, cinchonine, strychnine, and the like), and the like.

Method of optical resolution of a present invention compound or an intermediate thereof which has basic group
20 comprises steps of

(1) a step of forming a salt of a compound of present invention or intermediate thereof and an optically active acid;

(2) a step of recrystallization.

25 Examples of the solvent forming a salt are an alcohol solvent (such as methanol, ethanol 2-propanol and the like), an ether solvent (such as diethyl ether and the like), an

aromatic hydrocarbon solvent (such as toluene and the like),
an aprotic solvent (such as acetonitrile), and a mixture of
the solvent described before. Examples of the optical
active acid are a monocarboxylic acid (such as mandelic
5 acid, N-benzyl alanine, lactic acid and the like), a
dicarboxylic acid (such as tartaric acid, O-
diisopropylidene tartaric acid, malic acid and the like), a
sulfonic acid (such as camphor sulfonic acid, bromocamphor
sulfonic acid and the like), and the like.

10 Temperature of forming the salt is selected from the
range from about room temperature to about boiling point of
the solvent. To increase the purity of optical isomer, it
is preferable to heat the solution of salts to around the
boiling point of solvent. A molar rate of optically active
15 acid and the compound or the intermediate is selected from
the range of about 0.5 to about 2.0, preferably around 1.0.
If it is necessary to improve the optical purity, it is
possible to repeat recrystallization in the solvent
described before.

20 The present invention also includes solvates such as
the hydrate and the like, of the hydroxamic acid derivative
or the prodrug thereof, or the pharmaceutically acceptable
salt thereof.

25 The pharmaceutically acceptable salts of the hydroxamic
acid derivative or the prodrug thereof include, for example,
salts with inorganic acids such as the hydrochloride,

hydrobromide, sulfate, phosphate and the like; salts with organic acids such as the acetate, oxalate, citrate, lactate, tartrate, malonate, fumarate, maleate, mathanesulfonate and the like; salts with inorganic metals such as the lithium salt, sodium salt, potassium salt, magnesium salt, aluminum salt, barium salt and the like; salts with organic bases such as the ammonium salt, triethylammonium salt, tetrabutylammonium salt, pyridinium salt, pyrrolidinium salt, piperidinium salt and the like.

The hydroxamic acid derivatives or the prodrug thereof, or the pharmaceutically acceptable salt thereof of the present invention are useful as matrix metallo-proteinase inhibitors. Therefore, they may be used to treat or prevent a disease associated with excess or undesired matrix metallo-proteinases.

The diseases associated with excess or undesired matrix metallo-proteinases include, for example, the following diseases:

Abnormal wound healing, acne, acute coronary syndrome, acute infection, AIDS, alcoholism, allergic conjunctivitis, allergic reactions, allergic rhinitis, ALS, Alzheimer's diseases, anaphylaxis, aneurysmal aortic disease, angina, angiofibromas, anorexia, aortic aneurysm, ARDS, aspirin-independent anti-thrombosis, asthma, atherosclerosis, atherosclerotic plaque rupture, atopic dermatitis, benign hyperplasia, bleeding, bone

fractures, bronchitis, burns, cachexia, cancer, cardiac
infarction, cardiac insufficiency, cardiomyopathy,
cerebral hemorrhaging, cerebral ischemia, cerebral
vascular dementia, CHF, chronic bronchitis, chronic
5 dermal wounds, chronic obstructive pulmonary disease,
cirrhosis, congestive heart failure, corneal injury,
coronary thrombosis, Crohn's disease, cystic fibrosis,
decubitis ulcer, diabetic peripheral neuropathy,
diabetic retinopathy, diabetic ulcers, Duchenne's
10 muscular dystrophy, emphysema, endometriosis,
endosclerosis, epidermolysis bullosa, eye disorders,
fibrosis, gastritis, gingivitis, glomerular diseases,
glomerulonephritis, gout, graft rejection, gum disease,
GVHD, Hashimoto's thyroiditis, head trauma, headaches,
15 heart attacks, heart failure, hemangiomas, hemorrhage,
hepatitis, hirsutism, Huntington's disease, hypertension,
insulin resistance, interstitial nephritis, ischemia,
ischemic heart disease, Kaposi's sarcoma, keratinization,
keratitis, kidney failure, leishmaniasis, leprosy,
20 leukemia, leukocyte infiltration, liver cirrhosis, loss
of appetite, macular degeneration, malaria, mandibular
joint disease, memory impairment, meningitis, migraine,
miscarriage, multi-infarct dementia, multiple sclerosis,
muscular dystrophy, myalgia, myasthenia gravis, myelinic
25 degradation, myocardial infarction, myopia, neovascular
glaucoma, neuroinflammation, ocular tumors, optic
neuritis, osteoarthritis, osteopenia, Paget's disease,

pain, pancreatitis, Parkinson's disease, periodontitis, peripheral vascular disease, polyarteritis nodositas, polychondritis, premature childbirth, premature rupture of fetal membranes, prion disease, proliferative
5 retinopathies, proteinurea, pseudogout, psoriasis, pterygium, pulmonary emphysema, radiation damage, rattlesnake bite, Reiter's syndrome, renal fibrosis, reocclusion, reperfusion injury, restenosis, scleritis, scleroderma, senile dementia, senility, sepsis, septic
10 shock, Sharp syndrome, Sjogren's syndrome, SLE, spondylosis, stenosis, sterility, stroke, system sclerosis, thrombosis, toxic effects of chemotherapy, toxic shock, tuberculosis, ulcerations (corneal, epidermal, gastric), ulcerative colitis, uremia,
15 vasculitis, ventricular dilation, vesicular epidermolysis,.

The hydroxamic acid derivative or the prodrug thereof, or the pharmaceutically acceptable salt thereof may be
20 administered orally or parenterally. Pharmaceutical forms for oral administration include for example tablets, pills, capsules, powders, granules, suspensions, emulsions, syrups and the like. Pharmaceutical forms for parenteral
administration include for example intravenous injections
25 such as drops, intramuscular injections, subcutaneous injections, intranasal preparations, eye drops, suppository, percutaneous preparations such as ointments, creams,

lotions, and the like.

The solid compositions such as tablets can be prepared by mixing the active compound with pharmaceutically acceptable conventional carriers or excipients such as lactose, sucrose, corn starch or the like; binders such as hydroxypropylcellulose, polyvinylpyrrolidone, hydroxypropylmethylcellulose or the like; disintegrating agents such as sodium carboxymethylcellulose, sodium starch glycolate or the like; lubricants such as stearic acid, magnesium stearate or the like; or preservatives or the like.

For parenteral administration, the active compound can be dissolved or suspended in a physiologically acceptable carrier such as water, saline, oil, dextrose solution or the like, which may contain auxiliary agents such as emulsifiers, stabilizers, salts for influencing osmotic pressure or buffers, if desired.

The dose for administration varies widely depending on the grade of the symptoms, the patient's age, body weight, sex, administration route, and the like. But the hydroxamic acid derivative is usually administered to an adult (ca. 60kg) in a dose of approximately 1 - 1,000 mg, preferably 5 - 300 mg once or several times per day, by the oral route. By the parenteral route, the hydroxamic acid derivative is usually administered to an adult (ca. 60kg) in a dose of approximately 0.1 - 200 mg, preferably 0.3 - 100 mg once or several times per day.

Examples


The present invention is explained below precisely with examples but is, of course, not limited by them. The mass and purity of the samples were determined by LC/MS using a column of 4.6 x 50 mm from YMC(ODS-A column). The flow rate through the column was 3.5 ml/minute. About 100 μ l/min were split into the mass-spectrometer. The mass scan ranged from 200 to 700 amu. The determination of the purity of the samples was assigned through an UV detector (wavelengths: 220 and 254 nm). The gradient employed was from 10 to 99% of mobile phase B(0.035% of TFA in acetonitrile) during a period of 5 minutes (see table below).

Duration (min)	Mobile Phase A	Mobile Phase B
0.1	90	10
0.5	90	10
3.7	1	99
0.2	1	99
1	90	10
1	90	10

Solvent system: Mobile phase A: 0.05% TFA in water

Mobile phase B: 0.035% TFA in acetonitrile

Example 1



Step1: 2-[2-(2-Fluorophenyl)-3-(2-phenylethyl)-4-oxo-thiazolidin-5-yl]-acetic acid

A solution of phenylethylamine (85 μ l), 2-fluorobenzaldehyde (143 μ l) and mercaptosuccinic acid (306 mg) and trimethyl orthoformate (0.5 mL) in THF (3 mL) was heated at 80 °C for 4 hrs. The reaction mixture was cooled, diluted with EtOAc (2 mL) and washed twice with 2 ml of 1N HCl. The organic layer was concentrated to give a residue that was used in the next step without further purification.

Step 2: 2-[2-(2-Fluorophenyl)-3-(2-phenylethyl)-4-oxo-thiazolidin-5-yl]-acetic acid N-hydroxyamide

The residue from the previous step was dissolved in dry CH₂Cl₂ (1.5 mL) and pyridine (0.5 mL). Pentafluorophenyl trifluoroacetate (175 μ l) was added and the mixture was stirred at room temperature for 1 hour. O-TBS-hydroxylamine (200 mg) was added and the agitation was continued at room temperature for 12 hours. The reaction was diluted with CH₂Cl₂ (3 mL) and quenched by adding 6N HCl (1.5 mL). The organic layer was washed with 2N HCl (2 mL). The crude reaction mixture was purified by HPLC.

The following compounds listed in the Table 1 were prepared in a similar manner.





Table 1		RR1		RR2	mol. weight		Mass (m/e)	HPLC rt (min)
Example No.								
1		-(CH ₂) ₂ -Ph		-C ₆ H ₄ -2-F	374.44		375.2, 342.2	3.05
2		-CH(benzyl)CO ₂ Me		-C ₆ H ₂ -5-Br-2,4-(OMe) ₂	553.43		555.2	3.25
3		-(CH ₂) ₂ -Ph		-4-Py	357.43		358.0, 325.2	1.77
4		-Benzyl		-2-Thiazolyl	349.43		350.2, 317.0	2.73
5		-Benzyl		-C ₆ H ₄ -4-CO ₂ Me	400.46		401.4, 368.2	3.13
6		-Benzyl		-2-Furyl-5-Br	411.28		413.0, 380.0, 265.2	3.19
7		-Benzyl			452.54		453.0, 420.4	2.83
8		-(CH ₂) ₂ -Ph		-3-Indolyl-5-OMe	425.51		426.2, 279.0	2.87
9		-CH ₂ -C ₆ H ₃ -2-F,4-Br		-C ₆ H ₄ -4-OMe	469.33		469.2, 318.0	3.15
10		-CH ₂ CHCl(CH ₂) ₃ Cl		-C ₆ H ₃ -2-NO ₂ ,3-OMe	466.34		466.2, 433.2	3.35
11		-(CH ₂) ₂ -C ₆ H ₄ -4-Me		-C ₆ H ₂ -5-Br-2,4-(OMe) ₂	509.42		511.2, 448.0	3.66
12		-(CH ₂) ₄ -Ph			474.67		474.2, 441.4	3.02
13		-(CH ₂) ₃ -Ph		-C ₆ H ₄ -4-OMe	400.50		401.2, 368.0	3.32
14		-(CH ₂) ₃ -Ph		1-Naphthyl-4-NMe ₂	463.60		464.2, 293.0, 250.0	2.23
15		-(CH ₂) ₃ N(nBu) ₂		-2-Furyl-5-(C ₆ H ₄ -4-Br)	566.56		568.2	2.99, 3.08
16		-CH ₂ CH(Me)Ph		-C ₆ H ₄ -4-O-nBu	442.58		443.0, 249.8	3.17
17		-CH ₂ C(Me) ₂ CH ₂ NMe ₂		-2-Furyl-5-(C ₆ H ₄ -4-Cl)	466.00		466.2	2.84
18		-(CH ₂) ₂ -C ₆ H ₄ -4-OH		-3-Pyrrolyl-4-CO ₂ Et,2,5-Me ₂ ,1-Ph	537.64		538.2, 505.2	3.36
19		-(CH ₂) ₂ -C ₆ H ₄ -4-OH		-C ₆ H ₃ -2,3-(OMe) ₂	432.50		433.2, 400.2	2.85
20		-(CH ₂) ₂ -C ₆ H ₃ -3,4-(OMe) ₂		-2-Furyl-5-(C ₆ H ₄ -4-Br)	561.46		563.2, 528.2	3.71

Table 1



Example 21

Step 1: 2-(4-Methoxyphenyl)-3-[(1S)-methoxycarbonyl-3-methylbutyl]-1,3-thiazolidine-4-one

A solution of amino acid methyl ester (0.68 mmol) in 1 mL of THF was poured into a sealed tube containing molecular sieves 3A and a solution of the aldehyde (1.5 mmol) in dioxane (1 mL). The mixture was heated at 80 °C for 4 hours. Mercaptoacetic acid (17 µl, 2.5 mmol, 3.7 equiv.) was added to the mixture and the heating was continued for additional 12 hours. The reaction mixture was cooled down, diluted with EtOAc (2 mL) and filtered to remove the molecular sieves. The filtrate was washed twice with 2 mL of aq Na₂CO₃ and evaporated to give an oil that was used in the next step without further purification.


Step 2: 3-[(1S)-(N-Hydroxyaminocarbonyl)-3-methylbutyl]-2-(4-methoxyphenyl)-1,3-thiazolidine-4-one




KOH (435 mg) and hydroxylamine•HCl (340.2 mg) were dissolved in MeOH (2.14 mL and 3.4 mL respectively) at room temperature. Once a solution was obtained the KOH solution was poured into the hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for





1 hour. The precipitate (KCl) was filtered off, and the filtrate was poured into the solution of the methyl ester (0.68 mmol) in MeOH (2 mL) and stirring was continued for additional 3 hours. Acetic acid (0.5 mL) was added to the mixture and the volume of the reaction was reduced to one third. The resulting viscous solution was diluted with water (2 mL) and EtOAc (5 mL). The organic layer was separated, evaporated and purified by HPLC.


The following compounds listed in the Tables 2, 2' and 2'' and Table 3 were prepared in a similar manner.

Table 2




Example No.	RR1	RR2	RR3	mol. weight	Mass (m/e)	HPLC rt (min)
21	-iBu	-C6H4-4-OMe	-H	338.43	339.0, 306.2, 278.4	2.93
22	-Benzyl	-C6H4-2-Br	-H	421.32	423.0, 390.0, 362.0	3.28
23	-Benzyl	-C6H4-4-CF3	-H	410.42	411.2, 378.2, 350.0	3.42
24	-Benzyl	-C6H3-3,4-methylenedioxy	-Me	400.46	401.4, 368.4, 340.2	3.27
25	-Benzyl	2-Furyl-5-(C6H4-4-NO2)	-H	453.48	453.8, 421.2, 393.2	3.44
26	-Benzyl		-H	496.53	497.0, 464.0, 436.2	3.63
27	-Benzyl	-C6H4-4-n-pentyl	-H	412.56	413.2, 380.2, 352.0	4.08
28	-Benzyl	-C6H3-3-NO2,4-Cl	-H	421.86	422.2, 389.2, 361.4	3.49
29	-Benzyl	-C6H4-4-CO2Me	-H	400.46	401.0, 368.0, 340.0	3.16
30	-Benzyl	-C6H4-4-CH2O-(Ph-4-NO2)	-H	493.54	494.2, 461.2, 433.2	3.47
31	-Benzyl	-4-Imidazolyl-5-Me	-H	346.41	348.2	2.20
32	-Benzyl	-C6H3-3-NO2,4-S(C6H4-4-Br)	-H	574.48	575.4	3.36
33	-iBu	-2-Pyridyl-6-Me	-H	323.42	324.2, 149.2	2.00
34	-iBu	-C6H4-4-N(CH2CH2CN)2	-H	429.55	430.0, 369.0	2.86
35	-CH2-C6H4-4-OH	-3-Furyl	-H	348.38	349.4, 288.2	2.15
36	-CH2-C6H4-4-OH	-2-Thienyl-5-Me	-H	378.47	379.2, 346.0, 282.4	2.49
37	-iPr	-CH(CN)-C6H4-4-Cl	-H	367.86	369.2, 336.2	2.47
38	-CH2-3-indolyl		-H	461.59	462.2, 429.2, 401	3.32
39	-CH2-3-indolyl		-H	487.60	488.0, 455.2	3.30
40	-CH2-3-indolyl	-2-Furyl-5-(C6H4-4-Cl)	-H	481.96	482.2, 449.0	3.19
41	-CH2-3-indolyl	-2-Furyl-5-(C6H4-4-Br)	-H	526.41	528.0, 495, 467	3.21


42	-iBu			-H	388.53	389.2, 356, 328.2	3.62, 3.68
43	-iBu			-H	414.55	415.2, 382, 354.2	3.52
44	-iBu			-H	378.52	379.2, 346, 318.0	3.25
45	-iBu		-2-Furyl-5-C6H4-3-CF3	-H	442.46	443.2, 410.2, 382	3.41
46	-iBu		-2-Furyl-5-(C6H4-4-Cl)	-H	408.91	409.2, 376, 348.2	3.40
47	-iBu		-2-Furyl-5-(C6H4-4-Br)	-H	453.36	455.2, 422, 394	3.44
48	-iBu		-3-Py	-H	309.39	310.2, 277, 249.2	1.37
49	-iBu		-4-Py	-H	309.39	310.2, 249.2	0.94
50	-iBu		-2-Naphthyl-6-OMe	-H	388.49	389.2, 356, 328.2	3.14
51	-iBu		-C6H4-4-OH	-H	324.40	325.2, 292, 264.0	2.71
52	-Benzyl			-H	384.46	385.2, 352.0, 324	3.14, 3.20
53	-Benzyl			-H	422.55	423.2, 390, 362.2	4.06
54	-Benzyl			-H	448.57	449.0, 416.2, 388	3.97
55	-Benzyl		-2-Furyl-5-(C6H4-4-Cl)	-H	442.92	443.2	3.78
56	-Benzyl		-2-Furyl-5-(C6H4-4-Br)	-H	487.38	487.0, 454, 426	3.84
57	-Benzyl		-C6H4-4-OH	-H	358.42	359.2, 326, 298.2	2.79
58	-iBu		-C6H3-3-F-4-OMe	-H	356.42	357.2, 324.2, 296.2	2.89
59	-H		-C6H2-2,4,5-(OMe)3	-H	342.37	343.2, 282.0, 251.0	2.22
60	-H			-H	406.41	407.2, 374.0	2.89
61	-(CH2)2-Ph		-C6H4-4-OMe	-H	386.47	387.2, 354.2	3.31

62	-CH ₂ -C ₆ H ₄ -4-Ph	-C ₆ H ₄ -4-OMe	-H	448.55	449.0, 416.2, 388.2	3.74
63	-CH ₂ -2-Naphthyl	-C ₆ H ₄ -4-OMe	-H	422.51	423.2, 390.2, 316.2	3.55
64	-Benzyl	-C ₆ H ₄ -4-OCF ₃	-H	408.43	409.2, 348.2, 247.2	3.14
65	-Benzyl	-C ₆ H ₄ -4-SO ₂ Me	-H	420.51	421.2, 360.2, 286.2	2.63
66	-iBu	-C ₆ H ₄ -4-Br	-H	387.30	387.2, 326.0	3.30
67	-Me	-2-Pyrrolyl-1-SO ₂ -(C ₆ H ₄ -4-Me)	-H	409.49	410.2, 377, 349.2	2.85
68	-CH ₂ -3-indolyl	-C ₆ H ₄ -4-OMe	-H	411.48	412.2, 379.2, 351	3.12
69	-CH ₂ -3-indolyl	-C ₆ H ₄ -4-O-nBu	-H	453.56	454.2, 421.2, 393	3.67
70	-(CH ₂) ₂ SCH ₃	-C ₆ H ₄ -4-OMe	-H	356.47	357.2, 324, 296.2	2.93
71	-(CH ₂) ₂ SCH ₃	-C ₆ H ₃ -3-F,4-OMe	-H	374.46	375.0, 342, 314.2	2.90
72	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-OMe	-H	378.49	379.2, 346, 318.0	3.41
73	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-O-nBu	-H	420.58	421.2, 388, 360.2	4.05
74	-CH ₂ -cyclohexyl	-C ₆ H ₃ -3-F,4-OMe	-H	396.48	397.2, 364, 336.2	3.42
75	-CH ₂ S-benzyl	-C ₆ H ₄ -4-OMe	-H	418.54	419.0, 386.2, 358	3.36
76	-CH ₂ S-benzyl	-C ₆ H ₃ -3-F,4-OMe	-H	436.53	437.2, 404.2, 376	3.39
77	-CH ₂ -C ₆ H ₄ -4-OH	-C ₆ H ₄ -4-OMe	-H	388.45	389.2, 356, 328.2	2.73, 2.83
78	-CH ₂ -C ₆ H ₄ -4-OH	-C ₆ H ₄ -4-O-nBu	-H	430.53	431.2, 398, 370.2	3.49
79	-CH ₂ O-tBu	-C ₆ H ₄ -4-OMe	-H	368.46	369.2, 313.2, 280, 252.0	3.29
80	-CH ₂ O-tBu	-C ₆ H ₃ -3-F,4-OMe	-H	386.45	387.0, 331.2, 298, 270.2	3.33
81	-nBu	-C ₆ H ₄ -4-OMe	-H	338.43	339.2, 306, 278.0	3.18
82	-nBu	-C ₆ H ₄ -4-Me	-H	322.43	323.2, 290, 262.2	3.40
83	-nBu	-C ₆ H ₃ -3-F,4-OMe	-H	356.42	357.2, 324, 296.2	3.21
84	-CH ₂ -C ₆ H ₄ -4-OH	-2-Furyl-5-(C ₆ H ₄ -3-NO ₂)	-H	469.48	470.2, 437, 409.2	3.04
85	-Ph	-C ₆ H ₄ -4-SMe	-H	374.48	375.2, 342, 314.4	3.09
86	-CH ₂ -C ₆ H ₄ -4-OH	-C ₆ H ₄ -4-Cl	-H	392.86	393.0, 360, 332.0	2.78, 2.90
87	-iBu	-C ₆ H ₄ -4-Cl	-H	342.85	343.2, 310, 282.2	3.10, 3.22
88	-iBu		-H	462.52	463.0, 430, 402.2	3.43
89	-iBu	-C ₆ H ₄ -4-O-(CH ₂) ₃ NMe ₂	-H	409.55	410.2, 349.2	2.32, 2.45
90	-iPr	-C ₆ H ₄ -4-OMe	-H	324.40	325.2, 292.2, 264	3.11
91	-iBu	-C ₆ H ₃ -3,4-methylenedioxy	-H	352.41	353.2, 320.2, 292.2	3.15
92	-iBu	-C ₆ H ₄ -4-OMe	-H	338.43	339.2, 306.2, 278.2	3.18
93	-Me	-C ₆ H ₄ -4-Br	-H	345.22	347.0, 314.0, 286.0	3.05
94	-CH(OH)CH ₃	-C ₆ H ₄ -4-Cl	-H	330.79	331.0, 298.0, 270.0	2.66
95	-nBu	-C ₆ H ₄ -4-SMe	-H	354.49	355.2, 322.2, 394.2	3.27
96	-CH ₂ O-tBu	-C ₆ H ₄ -4-O-nBu	-H	410.54	411.2, 355.0, 322.2, 294.0	3.93
97	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-OPh	-H	440.57	441.2, 408.2, 380.2	3.66

98	-CH2-cyclohexyl	-C6H4-4-Ph	-H	424.57	425.0, 392.0, 364.0	3.64
99	-CH2-cyclohexyl	-Ph	-H	348.47	349.2, 316.2, 288.2	3.21
100	-Me	-C6H4-4-OPh	-H	358.42	359.0, 326.0, 298.2	3.00
101	-nBu	-C6H4-4-OPh	-H	400.50	401.2, 368.2, 340.2	3.38
102	-nBu	-C6H4-4-Ph	-H	384.50	385.2, 352.2, 324.2	3.39
103	-CH2-3-indolyl	-C6H4-4-OMe	-H	411.48	412.2, 379.2, 351.2	2.90
104	-Ph	-C6H4-4-OMe	-H	358.42	359.0, 326.0, 298.2	2.80
105	-CH2O-tBu	-C6H4-4-OMe	-H	368.46	369.2, 313.2, 280.2, 252.0	2.96
106	-nPr	-C6H4-4-OMe	-H	324.40	325.2, 292.2, 264.0	2.71
107	-CH2-(1-benzylimidazol-4-yl)	-C6H4-4-OMe	-H	452.54	453.2, 420.2, 392.0	2.53
108	-Me	-C6H4-4-OMe	-H	296.35	297.2, 264.2, 236.2	2.41
109	-CH2-C6H4-4-O-tBu	-C6H4-4-OMe	-H	444.55	45.2, 412.2, 384.0, 328.2	3.26
110	-nBu	-C6H4-4-OMe	-H	338.43	339.2, 306.2, 278.2	2.88
111	-CH2-3-indolyl	-C6H3-3-F,4-OMe	-H	429.47	430.2, 397.2, 369.2	2.90
112	-nBu	-C6H4-4-Cl	-H	342.85	343.2, 310.2, 282.2	3.11
113	-(CH2)2SCH3	-C6H4-4-OPh	-H	418.54	419.0, 386.0, 358.0	3.06
114	-CH2S-benzyl	-C6H4-3-Me	-H	402.54	403.2, 370.0, 342.2	3.07
115	-CH2S-benzyl	-C6H4-4-SMe	-H	434.60	435.2, 402.2, 374.2	3.09
116	-CH2S-benzyl	-C6H4-3-Cl	-H	422.96	423.0, 390.0, 362.0	3.11
117	-CH2S-benzyl	-C6H4-4-Cl	-H	422.96	423.0, 390.0, 362.0	3.14
118	-CH2S-benzyl	-C6H4-4-OPh	-H	480.61	481.4, 448.2, 420.4	3.32
119	-CH2O-tBu	-C6H4-4-SMe	-H	384.52	385.2, 329.2, 296.2, 268.2	3.35
120	-CH2-3-indolyl	-C6H4-4-F	-H	399.45	400.2, 367.0, 339.2	3.74
121	-CH2-3-indolyl	-C6H4-4-Cl	-H	415.90	416.2, 383.0	3.95
122	-CH2-C6H4-4-OH	-C6H4-4-SMe	-H	404.51	405.2, 372.2, 344.0	3.59
123	-CH2-C6H4-4-OH	-C6H4-4-SMe	-H	404.51	405.2, 372.2, 344.0	2.91
124	-CH2-C6H4-4-OH	-C6H4-4-OCF3	-H	442.42	443.2, 410.2, 382.2	3.12
125	-CH2-3-indolyl	-C6H4-4-SMe	-H	427.55	428.2, 395.2, 367.2	3.30
126	-CH2-3-indolyl	-C6H4-4-F	-H	399.45	400.2, 367.2, 339.2	3.20
127	-Me	-C6H4-4-OPh	-H	358.42	359.2, 326.2, 298.2	3.23
128	-CH2-C6H4-4-OH	-C6H4-4-OPh	-H	450.52	451.0, 418.0, 390.0	3.29
129	-iBu	-C6H4-4-SMe	-H	354.49	355.2, 322.2, 294.2	2.91
130	-iBu	-C6H4-4-Cl	-H	342.85	343.2, 310.2, 282.2	2.89
131	-iBu	-C6H3-3-F,4-OMe	-H	356.42	357.0, 324.0, 296.2	2.90
132	-iBu	-C6H4-4-Cl	-H	342.85	343.2, 310.2, 282.2	3.03
133	-iBu	-C6H4-4-Br	-H	387.30	389.2, 356.2, 328.2	3.04
134	-Benzyl	-C6H4-4-SMe	-H	388.51	389.2, 356.2, 328.2	3.33, 3.37
135	-iPr	-C6H4-4-nBu	-H	350.48	351.2, 318.2, 290.0	3.42
136	-(CH2)2SCH3	-C6H4-4-SMe	-H	372.53	373.0, 340.2, 312.0	2.98
137	-(CH2)2SCH3	-C6H4-4-Cl	-H	360.88	361.0, 328.2, 300.0	3.01
138	-(CH2)2SCH3	-C6H4-4-Br	-H	405.34	407.2, 374.2, 346.0, 298.2	3.04, 3.11


139	-(CH ₂) ₂ SCH ₃	-C ₆ H ₄ -4-OPh	-H	418.54	419.0, 386, 358.0, 310.2	3.30
140	-nPr	-C ₆ H ₄ -4-SMe	-H	340.47	341.2, 308, 280.0	2.97
141	-nPr	-C ₆ H ₄ -4-Cl	-H	328.82	329.2, 296, 268.0	2.94, 3.01
142	-nPr	-C ₆ H ₄ -4-Br	-H	373.27	375.0, 342, 314	2.98, 3.07
143	-nPr	-C ₆ H ₄ -4-OPh	-H	386.47	387.0, 354, 326.2	3.30
144	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-SMe	-H	394.56	395.2, 362, 334.4	3.40
145	-CH ₂ -cyclohexyl	-C ₆ H ₃ -3-F, 4-OMe	-H	396.48	397.2, 364, 336.2	3.26
146	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-F	-H	366.46	367.0, 334, 306.2	3.29
147	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-Cl	-H	382.91	383.0, 350, 322.2	3.44
148	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-Br	-H	427.36	429.2, 396, 368.0	3.49
149	-CH ₂ -cyclohexyl	-C ₆ H ₄ -4-OPh	-H	440.57	441.2, 308, 380.2	3.69
150	-nBu	-C ₆ H ₄ -4-SMe	-H	354.49	355.2, 322, 294.0	3.14
151	-nBu	-C ₆ H ₃ -3-F, 4-OMe	-H	356.42	357.0, 324, 296.2	2.96
152	-nBu	-C ₆ H ₄ -4-F	-H	326.39	327.2, 294, 266.0	3.00
153	-nBu	-C ₆ H ₄ -4-Cl	-H	342.85	343.0, 310, 282.0	3.15
154	-nBu	-C ₆ H ₄ -4-Br	-H	387.30	389.0, 356, 328.2	3.21
155	-nBu	-C ₆ H ₄ -4-OPh	-H	400.50	401.2, 368, 340.2	3.48
156	-nBu	-C ₆ H ₄ -3-OCF ₃	-H	392.40	393.0, 360, 332.2	3.29
157	-CH ₂ -(1-benzylimidazol-4-yl)	-C ₆ H ₄ -4-F	-H	440.50	441.4, 408.2	2.63
158	-CH ₂ -(1-benzylimidazol-4-yl)	-C ₆ H ₄ -4-Br	-H	501.41	503.2	2.74, 2.80
159	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-SMe	-H	460.62	461.2, 428, 400.2	3.49
160	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₃ -3-F, 4-OMe	-H	462.54	463.2, 430, 402.2, 346.0	3.35
161	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-F	-H	432.52	433.2, 400.2, 372, 316.2	3.35
162	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	448.97	449.0, 416.2, 388, 332.0	3.51
163	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Br	-H	493.42	495.2, 462, 434.0, 378.0	3.34
164	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OPh	-H	506.63	507.2, 474, 446.2, 390.0	3.75
165	-Benzyl	-C ₆ H ₄ -4-SMe	-H	388.51	389.0, 356, 328.2	3.20
166	-Benzyl	-C ₆ H ₄ -4-Cl	-H	376.86	377.0, 344, 316.2	3.22
167	-Benzyl	-C ₆ H ₄ -4-Br	-H	421.32	423.0, 390, 362.0	3.28
168	-Benzyl	-C ₆ H ₄ -4-OPh	-H	434.52	435.2, 402, 374.0	3.49
169	-(CH ₂) ₄ NHCO ₂ -tBu	-C ₆ H ₄ -4-OMe	-H	453.56	454.2, 398.2, 354.2	3.12
170	-CH ₂ CH ₂ CO ₂ -tBu	-C ₆ H ₄ -4-OMe	-H	410.49	411.2, 355.2, 322.2	3.14
171	-CH ₂ -4-imidazolyl	-C ₆ H ₄ -4-OMe	-H	362.41	363.2, 302.2	2.02, 2.10
172	-iPr	-C ₆ H ₄ -4-SMe	-H	340.47	341.0, 308, 280.2	2.86
173	-secBu	-C ₆ H ₄ -4-Cl	-H	342.85	343.0, 310, 282.0	3.02
174	-secBu	-C ₆ H ₄ -4-F	-H	326.39	327.0, 294, 266.2	2.85
175	-secBu	-C ₆ H ₃ -3-F, 4-OMe	-H	356.42	357.2, 324, 296.2	2.89
176	-secBu	-C ₆ H ₄ -4-SMe	-H	354.49	355.2, 322.2, 294	3.32
177	-(CH ₂) ₄ NH ₂	-C ₆ H ₄ -4-OMe	-H	353.44	354.2, 321.2	2.16
178	-(CH ₂) ₂ CO ₂ H	-C ₆ H ₄ -4-OMe	-H	354.38	354.8, 337.2	2.62, 2.69
179	-iPr	-C ₆ H ₄ -4-OPh	-H	386.47	387.0, 354, 326.2	3.27

180	-iPr		-C6H4-4-O-benzyl	-H	400.50	401.2, 268, 340.0	3.31
181	-Benzyl		-C6H4-4-Cl	-H	376.86	377.2, 344, 316.2	3.22
182	-Benzyl		-C6H4-4-Br	-H	421.32	423.0, 390, 362.0	3.27
183	-Benzyl		-C6H4-4-OPh	-H	434.52	435.2, 402, 374.0	3.49
184	-CH2O-tBu		-C6H4-4-Cl	-H	372.87	373.0, 317.0, 284.0, 256	3.32
185	-CH2O-tBu		-C6H4-4-Br	-H	417.32	419.0, 363.0, 330, 302	3.38
186	-CH2-(1-benzylimidazol-4-yl)		-C6H4-4-SMe	-H	468.60	469.2	2.77
187	-CH2-(1-benzylimidazol-4-yl)		-C6H4-4-Cl	-H	456.95	457.2	2.76
188	-CH2-(1-benzylimidazol-4-yl)		-C6H4-4-OCF3	-H	506.51	507.2, 474.2	2.95
189	-CH2-C6H4-4-OH		-C6H3-3-F,4-Cl	-H	410.85	410.8, 350.2	2.79
190	-CH2-C6H4-4-OH		-2-Thienyl-5-SMe	-H	410.54	411.4, 349.8	2.83
191	-CH2-C6H4-4-OH		-C6H3-2-F,4-OMe	-H	406.44	407.2, 374, 346.0	2.90
192	-iBu		-C6H4-4-O-allyl	-H	364.47	365.2, 332, 304.2	3.40
193	-nBu		-2-Thienyl-5-Cl	-H	348.87	349.0, 316, 288.0	3.41
194	-nBu			-H	366.44	367.2, 334.2, 306	2.95
195	-nBu		2-Benzofuranyl	-H	348.42	349.2, 316, 288.2	3.31
196	-CH2-C6H4-4-OH		-C6H4-4-Br	-H	437.32	436.8, 404, 376.2	2.79
197	-Benzyl		-C6H3-2-F,5-OMe	-H	390.44	391.0, 358.2, 330	2.92
198	-Benzyl		-2-Furyl-5-(C6H4-2-Cl)	-H	442.92	443.2, 410.2, 382	3.33
199	-Benzyl		-C6H3-2-F,4-OMe	-H	390.44	391.0, 358.2, 330	2.91
200	-iBu		-C6H3-2-F,4-OMe	-H	356.42	357.2, 324.4, 296	2.80
201	-CH2-C6H4-4-OH		-2-Thienyl-4-Br	-H	443.34	445.0, 383.8	2.70
202	-CH2-C6H4-4-OH		-2-Thienyl-5-Br	-H	443.34	444.8, 384	2.72
203	-CH2-C6H4-4-OH		-2-Thienyl-5-Cl	-H	398.89	399.0, 338.0	2.67
204	-CH2-C6H4-4-OH		-C6H3-2-F,4-Br	-H	455.31	457.2, 396	2.91




Example No.	RR ¹	RR ²	mol. weight	Mass (m/e)	HPLC rt (min)
447	-Cyclohexyl	-OMe	364.5	365.2, 332.2, 304	3.48
448	-Cyclohexyl	-OMe	364.5	365.2, 332.2, 304	3.58
449	-iBu	-Cl	342.8	343.0, 310, 282.2	3.09
450	-iBu	-Cl	342.8	343.0, 310, 282.2	3.13
451	-iBu	-O-C6H4-4-F	418.5	419.2, 386, 358.2	3.45
452	-iBu	-O-C6H4-4-Cl	434.9	435.2, 374.2	3.74
453	-iBu	-Cl	342.8	343.0, 310, 282.2	3.09
454	-CH2-2-imidazolyl	-Cl	366.8	367.0, 334, 306	1.85
455	-(CH2)3NHG(NH)NH2	-Cl	385.9	386.2, 353.0	2.26
456	-(CH2)2CO-piperidino	-Cl	425.9	426.2, 393, 365.0	2.82
457	-(CH2)3NHCOOCH2Ph	-Cl	478.0	478.2	3.16
458	-(CH2)2CONMe2	-Cl	385.9	386.2, 353, 325.0	2.58
459	-(CH2)3NHCOO-tBu	-Cl	444.0	444.2, 388.2	3.29
460	-iBu	-NEt2	379.5	380.2	3.49


Table 2"



Example No.	RR ¹	RR ²	mol. weight	Mass (m/e)	HPLC rt (min)
461	-iBu	-2-Thienyl	314.4	315.0,	3.29
462	-iBu	-2-Thienyl-5-Br	393.3	393.0, 360,	3.59
463	-iBu	-2-Thienyl-5-Br	393.3	393.0, 360,	3.58
464	-iBu	-2-Thiazolyl	315.4	316.0	2.89
465	-Benzyl	-2-Thienyl	348.4	349.0, 316,	3.34
466	-Benzyl	-2-Thienyl-5-Br	427.3	429.0, 427,	3.59

Table 3

					
Example No.	RR1	RR2	mol. weight	Mass (m/e)	HPLC rt (min)
205	-(CH ₂) ₃ CONH ₂	-C ₆ H ₄ -4-Br	359.2	361.0, 328, 300	2.79
206	-(CH ₂) ₃ CONH ₂	-C ₆ H ₃ -3,4-methylenedioxy	324.4	325.2, 292, 264.2	2.46
207	-(CH ₂) ₂ CONH ₂	-C ₆ H ₄ -4-OMe	296.3	297.0, 264.0	2.30
208	-CH(CO ₂ Me)CH ₂ CONH ₂	-C ₆ H ₄ -4-Br	403.3	405.0, 371.8, 344.0	3.07

Example 209

Step 1: 2-(4-Methoxyphenyl)-3-[(1*S*)-methoxycarbonyl-2-phenylethyl]-4-oxo-thiazolidin-5-yl)-acetic acid

5 A suspension of (L)-phenylalanine methyl ester (147 mg) and triethylamine (95 μ l) in 1 mL of THF was stirred at room temperature for 5 minutes. The solution was filtrated and then poured into a sealed tube containing 3A molecular sieves and a solution of p-methoxybenzaldehyde (183 μ l) in dioxane (1 mL). The mixture was heated at 80 °C for 4 hours. Mercaptosuccinic acid (375 mg) was added to the mixture and the heating was continued for an additional 12 hours. The reaction was cooled, diluted with EtOAc (2 mL) and filtered to remove the molecular sieves. The filtrate was washed 10 twice with 2 ml of 1N HCl and evaporated to give a solid residue that was used in the next step without further purification. 15

Step 2: (2-(4-Methoxyphenyl)-3-[(1*S*)-methoxycarbonyl-2-phenylethyl]-4-oxo-thiazolidin-5-yl)-acetic acid N-methylamide

The crude carboxylic acid from step 1,
5 diisopropylethylamine (178 μ l), HOBT (138 mg) and EDC.HCl (196 mg) were dissolved in CH₂Cl₂ (2 mL) and the reaction mixture was stirred at room temperature for 1 hour.

Methylamine (254 μ l, 8.03 M in ethanol) was added to the mixture and stirring was continued for an additional 12
10 hours. The reaction was diluted with CH₂Cl₂ (5 mL) and washed with 1N HCl (2 x 1.5 mL), water (1.5 mL) and brine (1.5 mL). The organic layer was concentrated *in vacuo* and the crude product was used in the next step.


15 Step 3: 2-[3-[(1*S*)-(N-Hydroxycarbamoyl)-2-phenylethyl]-2-(4-methoxyphenyl)-4-oxo-thiazolidin-5-yl]]-N-methylacetamide

KOH (435 mg) and hydroxylamine•HCl (340 mg) were dissolved in MeOH (2.14 mL and 3.4 mL respectively) at room
20 temperature. Once a solution was obtained, the KOH solution was poured into the hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for 1 hour. The precipitate (KCl) was filtered off. The filtrate was added into the solution of the methyl ester (obtained in step 2)
25 in MeOH (2 mL) and stirring was continued for an additional 3 hours. Acetic acid (0.5 mL) was added to the mixture and the volume of the reaction was reduced to one third. The

resulting viscous solution was diluted with water (2 mL) and EtOAc (5 mL). The organic layer was separated, evaporated and purified by HPLC.

5 The following compounds listed in the Tables 4, 4' and 4'' were prepared in a similar manner.

Table 4



Example No.	RR1	RR2	RR3	RR4	mol. weight	Mass (m/e)	HPLC rt (min)
209	-Benzyl	-C6H4-4-OMe	-H	-Me	443.53	444.2, 411, 383.2	2.93, 2.99
210	-Benzyl	-C6H4-4-OMe	-H	-OH	445.50	447.2, 413.2, 386.2	3.01
211	-Benzyl	-C6H4-4-OMe	-H	-CH2-3-Py	520.61	521.4, 460.2	2.82
212	-iBu	-C6H4-4-OMe	-H	-CH2-C6H4-4-Me	499.63	500.2, 467.4, 439	3.52
213	-iBu	-C6H4-4-OMe	-H	-CH2-C6H4-4-Me	499.63	500.2, 467.4, 439	3.58
214	-iBu	-C6H4-4-OMe	-H	-CH2-3-furyl	475.57	476.2, 443.2, 415	3.24
215	-iBu	-C6H4-4-OMe	-H	-CH2-3-furyl	475.57	476.2, 443.2, 415	3.28
216	-iBu	-C6H4-4-OMe	-H	-Benzyl	485.61	486.2, 453.2, 425	3.46
217	-iBu	-C6H4-4-Cl	-H	-CH2-2-furyl	479.99	480.2, 447.0, 419	3.42
218	-iBu	-C6H4-4-Cl	-H	-CH2-2-furyl	479.99	480.2, 447.0, 419	3.50
219	-iBu	-C6H4-4-Cl	-H	-CH2-2-furyl	479.99	480.2, 447.2, 419	3.50
220	-iBu	-C6H4-4-Cl	-H	-Benzyl	490.03	490.2, 457.2, 429	3.57
221	-iBu	-C6H4-4-Cl	-H	-Benzyl	490.03	490.2, 457.2, 429	3.64
222	-iBu	-C6H4-4-Cl	-H	-Benzyl	490.03	490.2, 457.2, 429	3.67
223	-iBu	-C6H4-4-Cl	-H	-Cyclopentyl	468.02	468.4, 435.2, 407	3.64
224	-iBu	-C6H4-4-Cl	-H	-Cyclopentyl	468.02	468.4, 435.2, 407	3.60
225	-iBu	-C6H4-4-Cl	-H	-Cyclopentyl	468.02	468.4, 435.2, 407	3.62
226	-iBu	-C6H4-4-Cl	-H	-Cyclopentyl	457.98	458.2, 425.2, 397	3.17
227	-iBu	-C6H4-4-Cl	-H	-(CH2)2OMe	457.98	458.2, 425, 397	3.23
228	-iBu	-C6H4-4-Cl	-H	-(CH2)2OMe	457.98	458.2, 425, 397	3.23
229	-iBu	-C6H4-4-OMe	-H	-iPentyl	465.62	466.4, 433.2, 405	3.54
230	-iBu	-C6H4-4-OMe	-H	-3-Isooxazolyl-5-Me	476.56	477.2, 444.2	3.27
231	-iBu	-C6H4-4-OMe	-H	-CH(Me)CH2OMe	467.59	468.4, 435.2, 407	3.04
232	-iBu	-C6H4-4-OMe	-H	-Cyclopentyl	463.60	464.4, 431.2, 403	3.42
233	-iBu	-C6H4-4-OMe	-H	-Cyclopentyl	463.60	464.4, 431.2, 403	3.38
234	-iBu	-C6H4-4-OMe	-H	-Et	423.54	424.2, 391, 363.2	3.08
235	-iBu	-C6H4-4-OMe	-H	-Et	423.54	424.0, 391, 363.2	3.11
236	-iBu	-C6H4-4-OMe	-H	-Allyl	435.55	436.2, 403, 375.2	3.13
237	-iBu	-C6H4-4-OMe	-H	-Allyl	435.55	436.2, 403, 375.2	3.19


238	-iBu	-C6H4-4-OMe	-H	-iBu	451.59	452.2, 419.2, 391	3.47
239	-iBu	-C6H4-4-OMe	-H	-Cyclohexyl	477.63	478.2, 445.2, 417	3.58
240	-iBu	-C6H4-4-OMe	-H	-iPr	437.56	438.2, 405.2, 377	3.22
241	-iBu	-C6H4-4-OMe	-H	-nPr	437.56	438.2, 405, 377.2	3.22
242	-iBu	-C6H4-4-OMe	-H	-nPr	437.56	438.2, 405, 377.2	3.26
243	-iBu	-C6H4-4-OMe	-H	-Ph	471.58	472.2, 437.2, 411	3.49
244	-iBu	-C6H4-4-OMe	-H	-Ph	471.58	472.2, 739.2, 411	3.52
245	-iBu	-C6H4-4-OMe	-H	-2-Py	472.57	473.2, 412.2	2.84
246	-iBu	-C6H4-4-OMe	-H	-(CH2)3OMe	467.59	468.4, 435.2, 407	3.09
247	-iBu	-C6H4-4-OMe	-H	-(CH2)3-1-(2-Me-piperidyl)	534.72	450.2, 417.2, 389	3.21
248	-iBu	-C6H4-4-OMe	-H	-(CH2)2OMe	453.56	454.2, 421.2, 393	3.02
249	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-OMe	626.18	626.4	3.90
250	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-OMe	626.18	626.4, 593.4	4.06
251	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-OMe	626.18	626.4, 593.2, 565	4.10
252	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-OMe	610.18	610.2, 577, 549.4	4.17
253	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-Me	597.14	597.2	3.24
254	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-CH2-3-Py	587.10	587.2, 531.0, 470.2	3.79
255	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-3-Isooxazolyl-5-Me	587.10	587.2, 531, 470.2	3.88
256	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-C(Me)2OMe	578.13	578.2, 517.2	3.69
257	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-C(Me)2OMe	578.13	578.2, 545, 517.2	3.77
258	-iBu	-C6H4-4-Cl	-H	-Ph	476.00	476.0, 443.2, 415	3.77
259	-iBu	-C6H4-4-Cl	-H	-iPr	441.98	442.4, 409.2, 381	3.43
260	-iBu	-C6H4-4-Cl	-H	-nPr	441.98	442.4, 409, 381.2	3.45
261	-iBu	-C6H4-4-Cl	-H	-iBu	456.01	456.2, 423.2, 395	3.42
262	-iBu	-C6H4-4-Cl	-H	-Allyl	439.96	440.2, 407, 379.2	3.37
263	-iBu	-C6H4-4-Cl	-H	-Et	427.95	428.2, 395, 367.0	3.29
264	-iBu	-C6H4-4-Cl	-H	-CH2-cyclopropyl	453.99	454.2, 421.2, 393	3.49
265	-iBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-OMe	520.05	520.4, 487.2, 459	3.64
266	-iBu	-C6H4-4-Cl	-H	-CH2-C6H4-4-Me	504.05	504.0, 471.2, 443	3.78
267	-iBu	-C6H4-4-Cl	-H	-2-Py	476.99	477.2, 416.2	3.12
268	-iBu	-C6H4-4-Cl	-H	-(CH2)3-1-(2-Me-piperidyl)	539.14	539.4	3.04
269	-iBu	-C6H4-4-Cl	-H	-iPentyl	470.04	470.2, 437.2, 409	3.75
270	-iBu	-C6H4-4-Cl	-H	-(CH2)3OMe	472.01	472.2, 439.2, 411	3.26
271	-iBu	-C6H4-4-Cl	-H	-C(Me)2OMe	472.01	472.2, 439.2, 411	3.34
272	-iBu	-C6H4-4-Cl	-H	-(CH2)2-2-Py	505.04	505.2, 472.2	2.93
273	-iBu	-C6H4-4-Cl	-H	-CH2-3-Py	491.01	491.2, 430.2	2.92
274	-iBu	-C6H4-4-Cl	-H	-3-Isooxazolyl-5-Me	480.97	481.2, 448.0, 420	3.44, 3.52
275	-CH2-C6H4-4-O-tBu	-C6H4-4-Cl	-H	-Et	534.08	534.2, 473.2	3.66

276	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Et	534.08	534.2, 501, 473.2	3.67, 3.72
277	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Et	534.08	534.2, 501, 473.2	3.73
278	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Allyl	546.09	546.2, 485.2	3.70
279	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Allyl	546.09	546.2, 513, 485.2	3.76
280	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Cyclohexyl	588.17	588.2, 527.4	4.17
281	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-iPr	548.11	548.4, 487.2	3.78
282	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-iPr	548.11	548.4, 515, 487.2	3.88
283	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-nPr	548.11	548.4, 515, 487.2	3.88
284	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-nPr	548.11	548.4, 487.2	3.77
285	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-2-Py	583.11	583.2, 522.4	3.53
286	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-2-Py	583.11	583.2, 522.4	3.52
287	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-2-Py	583.11	583.2, 522.4	3.42
288	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Ph	582.12	582.2, 549, 521.4	4.08
289	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ -cyclopropyl	560.12	560.2, 527, 499.2	3.92
290	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ -cyclopropyl	560.12	560.2, 499.2	3.80
291	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-nPr	-nPr	590.19	590.4, 529.2	4.39
292	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-4-Py	583.11	583.2	3.30
293	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-4-Py	583.11	583.2, 527.2	3.25
294	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Benzyl	596.15	596.2, 563, 535.2	4.05
295	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Cyclopentyl	574.14	574.4, 513.4	4.06
296	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-iPentyl	576.16	576.4, 543, 515.2	4.17
297	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 3OMe	578.13	578.2, 517.2	3.64
298	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 3OMe	578.13	578.2, 517.2	3.74
299	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 3-1-(2-Me-piperidyl)	645.27	645.4	3.22
300	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 2-2-Py	611.17	611.2	3.15
301	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 2-2-Py	611.17	611.2	3.26
302	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ -2-furyl	586.11	586.2, 525.2	3.98
303	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ -2-furyl	586.11	586.2, 553, 525.2	3.85, 3.90
304	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ -2-furyl	586.11	586.2, 525.2	3.80
305	-iBu	-C ₆ H ₄ -4-Cl	-nPr	-nPr	484.06	484.2, 451, 423	4.03
306	-iBu	-C ₆ H ₄ -4-Cl	-nPr	-nPr	484.06	484.2, 451.0	4.02
307	-Benzyl	-C ₆ H ₄ -4-OMe	-H	-Me	443.53	444.2, 411, 383.2	2.85
308	-Benzyl	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	549.65	550.4, 489.2	3.46
309	-Benzyl	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	549.65	550.4, 517.4, 489	3.51
310	-iBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -cyclopropyl	449.57	450.2, 417.2, 389	3.21
311	-iBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -cyclopropyl	449.57	450.2, 417.2, 389	3.30
312	-iBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	515.63	516.2, 483.2, 455	3.45
313	-iBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	515.63	516.2, 483.2, 455	3.39, 3.44
314	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-iBu	562.13	562.4, 501.2	4.07
315	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-CH ₂ 2OMe	564.11	564.2, 503.2	3.58

316	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-(CH ₂) ₂ OMe	564.11	564.2, 503.2	3.72
317	-iBu	-C ₆ H ₄ -4-Cl	-H	-(CH ₂) ₃ SMe	488.07	488.2, 455.2, 427	3.52
318	-iBu	-C ₆ H ₄ -4-Cl	-H	-(CH ₂) ₃ SMe	488.07	488.2, 455.2, 427	3.52
319	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-(CH ₂) ₃ SMe	594.20	594.2, 533.2	3.96
320	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-nBu	562.13	562.2, 501.2	3.95
321	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-nBu	562.13	562.2, 529, 501.2	3.99
322	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-nBu	562.13	562.2, 501.2	4.04
323	-iBu	-C ₆ H ₄ -4-Cl	-H	-Cyclohexyl	482.05	482.2, 449.2, 421	3.82
324	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Me	515.63	516, 483, 455	3.43
325	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Me	520.05	520.4, 459.2	3.51
326	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-Cl	-H	-Me	520.05	520.4, 487, 459.2	3.63
327	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-iPentyl	571.74	572.4, 539, 511.4	3.91, 3.97
328	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₃ OMe	573.71	574.4, 513.4	3.49
329	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₃ OMe	573.71	574.4, 541, 513.4	3.54
330	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₃ -1-(2-Me-piperidyl)	640.85	641.4	3.21, 3.25
331	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₂ -2-Py	606.75	607.2, 574.2	3.27
332	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₂ -2-Py	606.75	607.2	3.10, 3.14
333	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -2-furyl	581.69	582.2, 549, 521.4	3.68, 3.73
334	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Benzyl	591.73	592.4, 559, 531.2	3.80, 3.87
335	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-4-Py	578.69	579.2, 518.2	3.22
336	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Cyclopentyl	569.73	570.4, 537, 509.4	3.78, 3.85
337	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Cyclopentyl	569.73	570.4, 537, 509.4	3.85
338	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₂ OMe	559.69	560.2, 527.2, 499	3.42, 3.48
339	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₃ SMe	589.78	590.2, 557.2, 529	3.68
340	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-nBu	557.71	558.2, 525, 497.4	3.81
341	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-nBu	557.71	558.2, 525, 497.4	3.84
342	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-(CH ₂) ₃ SMe	589.78	590.2, 557, 529.2	3.78
343	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-C(Me) ₂ OMe	573.71	574.4, 541, 513.4	3.61
344	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-3-Isooxazolyl-5-Me	582.68	583.2, 522	3.63
345	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-C(Me) ₂ OMe	573.71	574.4, 541, 513.4	3.52, 3.59
346	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-3-Isooxazolyl-5-Me	582.68	583.2, 550, 522.2	3.70
347	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -4-Py	592.72	593.4	3.20, 3.26
348	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Et	529.66	530.2, 497, 469.4	3.44
349	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Et	529.66	530.2, 497, 469.4	3.53
350	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Allyl	541.67	542.4, 509, 481.4	3.52
351	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Allyl	541.67	542.4, 509, 481.2	3.60
352	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-iBu	557.71	558.2, 525, 497.4	3.73
353	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-iBu	557.71	558.2, 525, 497.4	3.80

354	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Cyclohexyl	583.75	584.2, 551, 523	3.89, 3.95
355	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-iPr	543.69	544.4, 511, 483.4	3.58
356	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-iPr	543.69	544.4, 511, 483.4	3.64
357	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-nPr	543.69	544.4, 511, 483.4	3.58, 3.65
358	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-nPr	543.69	544.4, 511, 483.4	3.65
359	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-Ph	577.71	578.2, 545.2, 517	3.86
360	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-2-Py	578.69	579.2, 518.2	3.22
361	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-2-Py	578.69	579.2, 518.2	3.29
362	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -cyclopropyl	555.70	556.2, 523, 495.4	3.62, 3.69
363	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -cyclopropyl	555.70	556.2, 523, 495.4	3.70
364	-CH ₂ -C ₆ H ₄ -4-O-tBu	-C ₆ H ₄ -4-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	621.76	622.4, 589, 561	3.73, 3.81

Table 4'



Example No.	RR¹	RR²	RR³	RR⁴	mol. weight	Mass (m/e)	HPLC rt (min)
467	-Benzyl	-Cl	-H	-Benzyl	524.0	524.2 491.2, 463	4.59
468	-Benzyl	-Cl	-H	-Benzyl	524.0	524.2, 491, 463.2	4.57
469	-Benzyl	-Cl	-H	-Benzyl	524.0	524.2, 491, 463.2	4.51
470	-Benzyl	-Cl	-H	-Benzyl	524.0	524.2, 491, 463.2	4.49
471	-Benzyl	-OMe	-nPr	-nPr	513.7	514.4, 481.2, 453	4.60
472	-Benzyl	-OMe	-nPr	-nPr	513.7	514.2, 481, 453	4.75
473	-Benzyl	-OMe	-H	-iBu	485.6	486.2, 453, 425.2	4.44
474	-Benzyl	-OMe	-H	-iBu	485.6	486.2, 453.2, 425	4.30
475	-Benzyl	-OMe	-H	-(CH₂)₃OMe	501.6	502.2, 469, 441.4	4.19
476	-Benzyl	-OMe	-H	-(CH₂)₃OMe	501.6	502.2, 469, 441.4	4.14
477	-Benzyl	-OMe	-H	-(CH₂)₃OMe	501.6	502.2, 469, 441.4	4.19
478	-Benzyl	-Cl	-H	3-Isooxazoly-5-Me	515.0	515.2, 482, 454.2	4.17
479	-Benzyl	-OMe	-H	-iPr	471.6	472.2, 439, 411.2	4.32
480	-Benzyl	-OMe	-H	-iPr	471.6	472.2, 439.4, 411	4.27
481	-Benzyl	-OMe	-H	-iPr	471.6	472.2, 439.2, 411	4.28
482	-Benzyl	-Cl	-H	-iPentyl	504.1	504.2, 471, 443.2	4.32
483	-Benzyl	-Cl	-H	-iPentyl	504.1	504.2, 471.2, 443	4.46
484	-Benzyl	-OMe	-H	-C(Me)₂OMe	501.6	502.2, 469, 441.4	3.88
485	-Benzyl	-OMe	-H	-C(Me)₂OMe	501.6	502.4, 469, 441.4	3.94
486	-Benzyl	-OMe	-H	-C(Me)₂OMe	501.6	502.4, 469, 441.4	3.95
487	-Benzyl	-Cl	-H	3-Isooxazoly-5-Me	515.0	515.2, 472, 454.2	4.15
488	-Benzyl	-Cl	-H	3-Isooxazoly-5-Me	515.0	515.2, 482, 454.2	5.24
489	-Benzyl	-OMe	-H	-CH₂-cyclopropyl	483.6	484.2, 451.2, 423	4.04
490	-Benzyl	-OMe	-H	-CH₂-cyclopropyl	483.6	484.2, 451.2, 423	4.11
491	-Benzyl	-OMe	-H	-CH₂-cyclopropyl	483.6	484.2, 451.2, 423	4.17
492	-Benzyl	-OMe	-H	-Ph	505.6	506.2, 473, 445.2	4.26

493	-Benzyl	-OMe	-H	-Ph	505.6	506.2, 473, 445.2	4.28
494	-Benzyl	-OMe	-H	-Allyl	469.6	470.2, 437, 409.2	3.98
495	-CH2-cyclohexyl	-Cl	-H	-Et	468.0	468.4, 435.2, 407	4.52
496	-CH2-cyclohexyl	-Cl	-H	-iPr	482.0	482.2, 449.2, 421	4.61
497	-CH2-cyclohexyl	-Cl	-H	-iPr	482.0	482.2, 449, 421.2	4.63
498	-CH2-cyclohexyl	-Cl	-H	-nPr	482.0	482.4, 449, 421.2	4.65
499	-CH2-cyclohexyl	-Cl	-H	-nPr	482.0	482.2, 449, 421.2	4.62
500	-CH2-cyclohexyl	-Cl	-H	-Ph	516.1	516.2, 483, 455.2	3.84
501	-CH2-cyclohexyl	-Cl	-H	-CH2-C6H4-4-OMe	560.1	560.2, 527.2, 499	3.91
502	-CH2-cyclohexyl	-Cl	-H	-CH2-C6H4-4-OMe	560.1	560.2, 527.4, 499	3.76
503	-CH2-cyclohexyl	-Cl	-H	-CH2-cyclopropyl	494.1	494.4, 461, 433.2	3.61
504	-CH2-cyclohexyl	-Cl	-H	-2-Py	517.1	517.4, 456.2	3.17
505	-CH2-cyclohexyl	-Cl	-H	-2-Py	517.1	517.2, 456.2	3.13
506	-CH2-cyclohexyl	-Cl	-H	-CH2-C6H4-4-Me	544.1	544.2, 511, 483.4	3.97
507	-CH2-cyclohexyl	-Cl	-H	-(CH2)3OMe	512.1	512.2, 479, 451.2	3.41
508	-CH2-cyclohexyl	-Cl	-H	-3-Isooxazolyl-5-Me	521.0	521.2, 488, 460.2	3.65
509	-CH2-cyclohexyl	-Cl	-H	-3-Isooxazolyl-5-Me	521.0	521.4, 488, 460.2	3.67
510	-CH2-cyclohexyl	-Cl	-H	-CH2-3-Py	531.1	531.2, 470.2	2.81
511	-CH2-cyclohexyl	-Cl	-H	-CH2-3-Py	531.1	531.2, 470.2	2.60
512	-CH2-cyclohexyl	-Cl	-H	-(CH2)3OMe	512.1	512.2, 479, 451.2	3.42
513	-Benzyl	-OMe	-Me	-(CH2)2NMe2	514.6	515.2	3.92
514	-CH2-cyclohexyl	-Cl	-H	-(CH2)2-2-Py	545.1	545.2, 484.4	2.84
515	-CH2-cyclohexyl	-Cl	-H	-(CH2)2-2-Py	545.1	545.2, 484.2	2.86
516	-CH2-cyclohexyl	-Cl	-H	-CH2-2-furyl	520.1	520.4, 487, 459.2	3.66
517	-CH2-cyclohexyl	-Cl	-H	-CH2-2-furyl	520.1	520.4, 487, 459.2	3.65
518	-CH2-cyclohexyl	-Cl	-H	-Benzyl	530.1	530.2, 497, 469.4	3.83
519	-CH2-cyclohexyl	-Cl	-H	-Me	454.0	454.2, 421, 393.2	3.25
520	-CH2-cyclohexyl	-Cl	-H	-nBu	496.1	496.4, 463, 435.2	3.78
521	-CH2-cyclohexyl	-Cl	-H	-(CH2)3SMe	528.1	528.2, 495, 467.2	3.70
522	-CH2-cyclohexyl	-Cl	-H	-(CH2)2OMe	498.0	498.2, 465, 437.2	3.34
523	-CH2-cyclohexyl	-OMe	-H	-3-Isooxazolyl-5-Me	516.6	517.4, 484, 456.2	4.46
524	-CH2-cyclohexyl	-OMe	-H	-3-Isooxazolyl-5-Me	516.6	517.2, 484, 456.2	4.42
525	-CH2-cyclohexyl	-OMe	-H	-CH2-2-furyl	515.6	516.2, 483, 455.2	4.40
526	-CH2-cyclohexyl	-OMe	-H	-(CH2)2-2-Py	540.7	541.2, 508.4	4.01
527	-CH2-cyclohexyl	-OMe	-H	-(CH2)3-1-(2-Me-piperidyl)	574.8	575.6	3.98
528	-CH2-cyclohexyl	-OMe	-H	-(CH2)3OMe	507.7	508.4, 475, 447.2	4.34
529	-CH2-cyclohexyl	-OMe	-H	-(CH2)3OMe	507.7	508.4, 475.2, 447	4.29
530	-CH2-cyclohexyl	-OMe	-H	-Benzyl	525.7	526.2, 493, 465.2	4.58
531	-CH2-cyclohexyl	-OMe	-H	-Benzyl	525.7	526.2, 493.4, 465	4.56

532	-CH2-cyclohexyl	-OMe	-nPr	-nPr	519.7	520.4, 487, 459	4.83
533	-CH2-cyclohexyl	-OMe	-H	-CH2-2-furyl	515.6	516.2, 483, 455.2	4.48
534	-Benzyl	-Cl	-H	-iPentyl	504.1	504.0, 471, 443.2	4.33
535	-Benzyl	-Cl	-H	-iPentyl	504.1	504.2, 471, 443.2	4.40
536	-CH2-cyclohexyl	-OMe	-H	-Cyclopentyl	503.7	504.2, 471.2, 443	4.51
537	-CH2-cyclohexyl	-OMe	-H	-4-Py	512.6	513.4, 452.2	4.04
538	-CH2-cyclohexyl	-OMe	-H	-4-Py	512.6	513.4, 452.2	4.01
539	-CH2-cyclohexyl	-OMe	-H	-(CH2)3SMe	523.7	524.4, 491.2, 463	4.49
540	-CH2-cyclohexyl	-OMe	-H	-(CH2)2OMe	493.6	494.4, 461.2, 433	4.34
541	-CH2-cyclohexyl	-OMe	-H	-(CH2)2OMe	493.6	494.4, 461.2, 433	4.27
542	-CH2-cyclohexyl	-OMe	-H	-Cyclopentyl	503.7	504.2, 471.2, 443	4.56
543	-CH2-cyclohexyl	-OMe	-H	-nBu	491.7	492.4, 459.4, 431	4.54
544	-CH2-cyclohexyl	-OMe	-H	-Me	449.6	450.2, 417, 389.4	3.88
545	-CH2-cyclohexyl	-OMe	-H	-Et	463.6	464.2, 431.2, 403	4.32
546	-CH2-cyclohexyl	-OMe	-H	-Et	463.6	464.2, 431.2, 403	4.36
547	-CH2-cyclohexyl	-OMe	-H	-Allyl	475.6	476.2, 443.2, 415	4.41
548	-CH2-cyclohexyl	-OMe	-H	-iBu	491.7	492.4, 459.4, 431	4.47
549	-CH2-cyclohexyl	-OMe	-H	-iBu	491.7	492.4, 459.4, 431	4.51
550	-CH2-cyclohexyl	-OMe	-H	-iBu	491.7	492.4, 459.4, 431	4.56
551	-CH2-cyclohexyl	-OMe	-H	-Cyclohexyl	517.7	518.4, 485.2, 457	4.62
552	-CH2-cyclohexyl	-OMe	-H	-Cyclohexyl	517.7	518.4, 485.2, 457	4.69
553	-CH2-cyclohexyl	-OMe	-H	-iPr	477.6	478.2, 445.2, 417	4.40
554	-CH2-cyclohexyl	-OMe	-H	-iPr	477.6	478.2, 445.2, 417	4.45
555	-CH2-cyclohexyl	-OMe	-H	-nPr	477.6	478.2, 445.2, 417	4.41
556	-CH2-cyclohexyl	-OMe	-H	-nPr	477.6	478.2, 445.2, 417	4.46
557	-CH2-cyclohexyl	-OMe	-H	-Ph	511.6	512.4, 479, 451.2	4.61
558	-CH2-cyclohexyl	-OMe	-H	-2-Py	512.6	513.4, 452.2	4.13
559	-CH2-cyclohexyl	-OMe	-H	-2-Py	512.6	513.4, 452.2	4.18
560	-CH2-cyclohexyl	-OMe	-H	-CH2-cyclopropyl	489.6	490.4, 457.2, 429	4.46
561	-CH2-cyclohexyl	-OMe	-H	-CH2-cyclopropyl	489.6	490.4, 457.2, 429	4.50
562	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-OMe	555.7	556.2, 523, 495.4	4.52
563	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-OMe	555.7	556.2, 523, 495.4	4.56
564	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-OMe	555.7	556.2, 523, 495.4	4.57
565	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-Me	539.7	540.4, 507, 479.2	4.62
566	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-Me	539.7	540.4, 507, 479	4.65
567	-CH2-cyclohexyl	-OMe	-H	-CH2-C6H4-4-Me	539.7	540.4, 507, 479	4.68
568	-CH2-cyclohexyl	-OMe	-H	-CH2-3-Py	526.7	527.2, 466.4	4.00
569	-Benzyl	-Cl	-Me	-(CH2)2NMe2	519.1	519.4	3.95
570	-Benzyl	-Cl	-H	-iPr	476.0	476.2, 443, 415.4	4.34


571	-Benzyl	-Cl	-H	-iPr	476.0	476.0, 443, 415.2	4.42
572	-Benzyl	-Cl	-H	-iPr	476.0	476.2, 443.2, 415	4.48
573	-Benzyl	-Cl	-nPr	-nPr	518.1	518.2, 485.2, 457	4.65
574	-Benzyl	-Cl	-nPr	-nPr	518.1	518.2, 485, 457.2	4.81
575	-Benzyl	-OMe	-H	-CH ₂ -C ₆ H ₄ -4-Me	533.7	534.2, 501, 473.2	4.58
576	-Benzyl	-Cl	-H	-nBu	490.0	490.2, 457, 429.2	4.57
577	-Benzyl	-Cl	-H	-nBu	490.0	490.2, 457, 429.2	4.64
578	-Benzyl	-OMe	-H	-nPr	471.6	472.2, 439.2, 411	4.32
579	-Benzyl	-OMe	-H	-nPr	471.6	472.2, 439, 411.2	4.37
580	-Benzyl	-OMe	-H	-nPr	471.6	472.2, 439, 411.2	4.38
581	-Benzyl	-OMe	-H	-(CH ₂) ₃ -1-(2-Me-piperidyl)	568.7	569.4	3.98
582	-Benzyl	-OMe	-H	-Benzyl	519.6	520.4, 487, 459.2	4.37
583	-Benzyl	-Cl	-H	-Cyclohexyl	516.1	516.2, 483, 455.2	4.67
584	-Benzyl	-OMe	-H	-nBu	485.6	486.2, 453, 425.2	4.30
585	-Benzyl	-Cl	-H	-CH ₂ -2-furyl	514.0	514.2, 481, 453.2	4.51
586	-Benzyl	-OMe	-H	-Cyclohexyl	511.6	512.4, 479, 451.2	4.54
587	-Benzyl	-OMe	-H	-CH ₂ -2-furyl	509.6	510.2, 477, 449.2	4.39
588	-Benzyl	-Cl	-H	-CH ₂ -cyclopropyl	488.0	488.2, 455, 427.2	4.49
589	-Benzyl	-Cl	-H	-CH ₂ -cyclopropyl	488.0	488.2, 455.2, 427	4.56
590	-Benzyl	-Cl	-H	-Allyl	474.0	474.2, 441, 413.2	4.43
591	-Benzyl	-Cl	-H	-Allyl	474.0	474.2, 441.2, 413	4.45
592	-Benzyl	-Cl	-H	-nPr	476.0	476.2, 443.2, 415	4.50
593	-Benzyl	-OMe	-H	-Et	457.6	458.2, 425, 397.2	4.19
594	-Benzyl	-Cl	-H	-Cyclohexyl	516.1	516.2, 483, 455.2	4.60
595	-Benzyl	-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	549.7	550.4, 517.4, 489	4.40
596	-Benzyl	-OMe	-H	-CH ₂ -C ₆ H ₄ -4-OMe	549.7	550.2, 517.4, 489	4.37
598	-Benzyl	-OMe	-H	-Benzyl	519.6	520.4, 487, 459.2	4.41
599	-Benzyl	-Cl	-H	-Et	462.0	462.2, 429, 401.2	4.24
600	-Benzyl	-Cl	-H	-Et	462.0	462.2, 429, 401.2	4.33
601	-Benzyl	-OMe	-H	-(CH ₂) ₂ OMe	487.6	488.2, 455, 427.2	4.07
602	-Benzyl	-OMe	-H	-(CH ₂) ₂ OMe	487.6	488.2, 455.2, 427	4.11
603	-Benzyl	-OMe	-H	-nBu	485.6	486.2, 453, 425.2	4.37
604	-Benzyl	-OMe	-H	-nBu	485.6	486.2, 453, 425.2	4.38
605	-Benzyl	-OMe	-H	-nBu	485.6	486.2, 453, 425	4.51
606	-Benzyl	-Cl	-H	-nPr	476.0	476.2, 443, 415.2	4.38
607	-Benzyl	-Cl	-H	-Ph	510.0	510.2, 477, 449.0	4.65
608	-Benzyl	-Cl	-H	-(CH ₂) ₃ OMe	506.0	506.2, 473, 445.2	4.36
609	-Benzyl	-Cl	-H	-iBu	490.0	490.2, 457, 429.2	4.56
610	-Benzyl	-Cl	-H	-iBu	490.0	490.2, 457.2, 429	4.63

611	-Benzyl	-Cl	-H	(CH ₂) ₃ SMe	522.1	522.4, 489, 461.2	4.56
612	-Benzyl	-Cl	-H	(CH ₂) ₃ OMe	506.0	506.2, 473, 445.2	4.28
613	-Benzyl	-Cl	-H	-C(Me) ₂ OMe	506.0	506.2, 473, 445.2	4.39
614	-Benzyl	-Cl	-H	-CH ₂ -C ₆ H ₄ -4-OMe	554.1	554.2, 421, 493.4	4.60
615	-Benzyl	-Cl	-H	-CH ₂ -C ₆ H ₄ -4-OMe	554.1	554.2, 521, 493.2	4.52
616	-Benzyl	-Cl	-H	(CH ₂) ₃ -1-(2-Me-piperidyl)	573.2	573.4	4.09
617	-Benzyl	-OMe	-H	-4-Me-3-isoxazolyl	510.6	511.2, 478, 450.2	4.31
618	-Benzyl	-OMe	-H	-iPentyl	499.6	500.2, 467, 439.4	4.44
619	-Benzyl	-OMe	-H	-iPentyl	499.6	500.2, 467, 439.4	4.46
620	-Benzyl	-OMe	-H	-iPentyl	499.6	500.2, 467, 439.4	4.51
621	-Benzyl	-OMe	-H	-iPentyl	499.6	500.2, 467.4, 439	4.60
622	-Benzyl	-Cl	-H	-C(Me) ₂ OMe	506.0	506.2, 473, 445.2	4.32
623	-Benzyl	-Cl	-H	-iBu	490.0	490.2, 457, 429.2	4.51
624	-Benzyl	-OMe	-H	-Et	457.6	458.2, 425, 397.2	4.13
625	-Benzyl	-Cl	-H	-CH ₂ -2-furyl	514.0	514.2, 481, 453.2	4.45
626	-Benzyl	-Cl	-H	(CH ₂) ₂ OMe	492.0	492.2, 459, 431.2	4.24
627	-Benzyl	-Cl	-H	(CH ₂) ₂ OMe	492.0	492.2, 459, 431.2	4.29
628	-Benzyl	-OMe	-H	-4-Me-3-isoxazolyl	510.6	511.2, 478, 450.2	4.25
629	-Benzyl	-Cl	-H	-nPr	476.0	476.2, 443, 415.4	4.46
630	-Benzyl	-Cl	-H	-Allyl	474.0	474.2, 441, 413.2	4.35
631	-Benzyl	-OMe	-H	-Cyclohexyl	511.6	512.4, 479, 451.2	4.47
632	-Benzyl	-Cl	-H	-CH ₂ -cyclopropyl	488.0	488.2, 455, 427.2	4.47
633	-Benzyl	-OMe	-H	-CH ₂ -2-furyl	509.6	510.2, 477, 449.2	4.40
634	-Ph	-Cl	-H	-Et	447.9	448.0, 415, 387	4.15
635	-Ph	-Cl	-H	-2-Py	497.0	497.2	4.14
636	-Ph	-Cl	-H	-2-Py	497.0	497.2	4.18
637	-Ph	-Cl	-H	-CH ₂ -C ₆ H ₄ -4-Me	524.0	524.2, 491.2, 463	4.64
638	-Ph	-Cl	-H	(CH ₂) ₃ OMe	492.0	492.2, 459.2, 431	4.26
639	-Ph	-Cl	-H	-Allyl	460.0	460.0, 427.2, 399	4.36
640	-Benzyl	-OMe	-H	(CH ₂) ₃ SMe	517.7	518.2, 485, 457.2	4.40
641	-Benzyl	-Cl	-H	-CH ₂ -3-Py	525.0	525.2, 464.2	4.02
642	-Benzyl	-Cl	-H	(CH ₂) ₂ -2-Py	539.1	539.2, 478.2	4.02
643	-Benzyl	-Cl	-H	-2-Py	511.0	511.2, 450.0	4.20
644	-Benzyl	-OMe	-H	(CH ₂) ₂ -2-Py	534.6	535.2, 474.2	3.89
645	-Benzyl	-Cl	-H	(CH ₂) ₂ -2-Py	539.1	539.2	3.94
646	-Benzyl	-Cl	-H	-Cyclopentyl	502.0	502.2, 469, 441.2	4.50
647	-Benzyl	-Cl	-H	-Cyclopentyl	502.0	502.2, 469, 441.2	4.60
648	-Benzyl	-OMe	-H	-Cyclopentyl	497.6	498.2, 465, 437.2	4.38
649	-Benzyl	-OMe	-H	-Cyclopentyl	497.6	498.2, 465, 437.2	4.45


650	-Benzyl	-OMe	-H	-2-Py	506.6	507.2, 446.2	4.00
651	-Benzyl	-OMe	-H	-2-Py	506.6	507.2, 446.2	4.02
652	-Benzyl	-OMe	-H	-(CH ₂) ₃ SMe	517.7	518.2, 485, 457.2	4.38
653	-Ph	-OMe	-H	-Cyclohexyl	497.6	498.2, 465.2, 437	4.48
654	-Ph	-OMe	-H	-iPr	457.6	458.2, 425.2, 397	4.25
655	-Ph	-OMe	-H	-Ph	491.6	492.2, 459.2, 431	4.47
656	-Ph	-OMe	-H	-2-Py	492.6	493.4, 460.2	4.04
657	-Ph	-OMe	-H	-2-Py	492.6	493.4, 460.2	4.10
658	-Ph	-OMe	-H	-CH ₂ -cyclopropyl	469.6	470.2, 437.2, 409	4.30
659	-Ph	-OMe	-H	-CH ₂ -C ₆ H ₄ -4-Me	519.6	520.4, 487.2, 459	4.48
660	-Ph	-OMe	-H	-3-Isooxazolyl-5-Me	496.5	497.2, 464.2, 436	4.30
661	-Ph	-OMe	-H	-C(Me) ₂ OMe	487.6	488.2, 455.2, 427	4.24
662	-Ph	-OMe	-H	-(CH ₂) ₃ OMe	487.6	488.2, 455.2, 427	4.15
663	-Ph	-OMe	-H	-(CH ₂) ₂ -2-Py	520.6	521.4, 488.2	3.92
664	-Ph	-OMe	-H	-CH ₂ -2-furyl	495.6	496.2, 463.2, 435	3.11
665	-Ph	-OMe	-H	-Cyclopentyl	483.6	484.2, 451.2, 423	3.18
666	-Ph	-OMe	-H	-(CH ₂) ₂ OMe	473.6	474.2, 441.4, 413	2.81
667	-Ph	-OMe	-H	-(CH ₂) ₃ SMe	503.6	504.0, 471.2, 443	3.14
668	-Ph	-OMe	-H	-(CH ₂) ₃ -Ph	533.7	534.2, 501.2, 473	3.46
669	-Ph	-Cl	-H	-iBu	476.0	476.0, 443.2, 415	3.43
670	-Ph	-Cl	-H	-Cyclohexyl	502.0	502.2, 469.2, 441	3.60
671	-Ph	-Cl	-H	-iPr	462.0	462.2, 429.2, 401	3.26
672	-Ph	-Cl	-H	-Ph	496.0	496.2, 463.2, 435	3.59
673	-Ph	-Cl	-H	-iPentyl	490.0	490.2, 457.2, 429	3.58
674	-Ph	-Cl	-H	-(CH ₂) ₃ -1-(2-Me-piperidyl)	559.1	559.2	2.81
675	-Ph	-Cl	-H	-(CH ₂) ₃ -1-(2-Me-piperidyl)	559.1	559.2	2.89
676	-Ph	-Cl	-H	-(CH ₂) ₂ -2-Py	525.0	525.2, 492.2	2.72
677	-Ph	-Cl	-H	-4-Py	497.0	497.2, 436.2	2.79
678	-Ph	-Cl	-H	-Cyclopentyl	488.0	488.2, 455.2, 427	3.56
679	-Ph	-Cl	-H	-(CH ₂) ₂ OMe	478.0	478.2, 445.2, 417	3.10
680	-Ph	-Cl	-H	-CH ₂ -cyclopropyl	474.0	474.2, 441.2, 413	3.48
681	-Ph	-Cl	-H	-CH ₂ -C ₆ H ₄ -4-OMe	540.0	540.2, 507.2, 479	3.60
682	-Ph	-Cl	-H	-CH ₂ -2-furyl	500.0	500.2, 467.2, 439	3.51
683	-iBu	-Cl	-H	-Me	413.9	414.2, 381, 353.2	2.96
684	-iBu	-Cl	-H	-Me	413.9	414.2, 381, 353.0	3.02
685	-iBu	-Cl	-H	-Me	413.9	414.2, 381, 353.0	3.11
686	-iBu	-Cl	-H	-CH ₂ -2-furyl	480.0	480.2, 447.0, 419	3.21
687	-iBu	-Cl	-H	-CH ₂ -2-furyl	480.0	480.2, 447.0, 419	3.28
688	-iBu	-Cl	-H	-CH ₂ -2-furyl	480.0	480.2, 447.2, 419	3.30

689	-iBu	-F	-H	-CH ₂ -2-furyl	463.5	464.2, 431, 403.2	3.06
690	-iBu	-F	-H	-(CH ₂) ₂ OMe	441.5	442.4, 409, 381.2	2.89
691	-iBu	-F	-H	-Me	397.5	398.2, 365, 337.2	2.85
692	-iBu	-F	-H	-H	383.4	384.2, 351, 323.2	2.75
693	-iBu	-H	-H	-CH ₂ -2-furyl	445.5	446.2, 413, 385.2	3.21
694	-iBu	-H	-H	-(CH ₂) ₂ OMe	423.5	424.0, 391, 363.2	2.87
695	-iBu	-H	-H	-Me	379.5	380.2, 347, 319.2	2.82
696	-iBu	-O-C ₆ H ₄ -4-F	-H	-CH ₂ -2-furyl	555.6	556.2, 523, 495.4	3.25
697	-iBu	-O-C ₆ H ₄ -4-F	-H	-(CH ₂) ₂ OMe	533.6	534.2, 501.2	2.98
698	-iBu	-O-C ₆ H ₄ -4-F	-H	-Me	489.6	490, 457.2	2.97
699	-iBu	-Cl	-H	-CH ₂ -2-furyl	480.0	480.2, 447.0, 419	2.91
700	-iBu	-Cl	-H	-Me	413.9	414.2, 381, 353.0	2.49
701	-iBu	-Cl	-H	-H	399.9	400.2, 367, 339.0	2.52
702	-iBu	-Cl	-H	-(CH ₂) ₂ OMe	458.0	458.0, 425, 397.2	2.67
703	-iBu	-O-C ₆ H ₄ -4-Cl	-H	-CH ₂ -2-furyl	572.1	572.2, 539.2, 511	3.44
704	-iBu	-O-C ₆ H ₄ -4-Cl	-H	-(CH ₂) ₂ OMe	550.1	550.2, 493.4, 460.2	3.22
705	-iBu	-O-C ₆ H ₄ -4-Cl	-H	-Me	506.0	506.2, 473, 445.2	3.16

Table 4"



Example No.	RR ¹	RR ²	RR ³	mol. weight	Mass (m/e)	HPLC rt (min)
706	-Benzyl	-Cl		531.1	531.2, 470.2	3.97
707	-Benzyl	-OMe	piperidino	497.6	498.2, 465.2, 437	4.18
708	-Benzyl	-OMe	piperidino	497.6	498.2, 465.2, 437	4.19
709	-Benzyl	-OMe	piperidino	497.6	498.2, 465.2, 437	4.24
710	-CH ₂ -cyclohexyl	-Cl	morpholino	510.1	510.2, 477.2, 449	3.40
711	-CH ₂ -cyclohexyl	-Cl	piperidino	508.1	508.4, 475.2, 447	3.94
712	-CH ₂ -cyclohexyl	-Cl	piperidino	508.1	508.4, 475.2, 447	3.77
713	-CH ₂ -cyclohexyl	-Cl	1-pyrrolidinyl	494.1	494.4, 461.2, 433	3.64
714	-CH ₂ -cyclohexyl	-Cl	piperidino	508.1	508.4, 475.2, 447	3.85
715	-CH ₂ -cyclohexyl	-OMe	morpholino	505.6	506.2, 473.2, 445	4.32
716	-CH ₂ -cyclohexyl	-OMe	morpholino	505.6	506.2, 473.2, 445	4.29
717	-CH ₂ -cyclohexyl	-OMe	1-pyrrolidinyl	489.6	490.4, 457.2, 439	4.46
718	-CH ₂ -cyclohexyl	-OMe	1-pyrrolidinyl	489.6	490.4, 457.2, 429	4.40
719	-CH ₂ -cyclohexyl	-OMe	piperidino	503.7	504.4, 471.2, 443	4.59
720	-CH ₂ -cyclohexyl	-OMe	piperidino	503.7	504.4, 471.2, 443	4.62
721	-Benzyl	-OMe		526.7	527.4	3.85
722	-Benzyl	-OMe		526.7	527.4	3.90
723	-Benzyl	-Cl	1-pyrrolidinyl	488.0	488.2, 455.2, 427	4.37
724	-Benzyl	-Cl	1-pyrrolidinyl	488.0	488.2, 455.2, 427.2	4.46
725	-Benzyl	-OMe	morpholino	499.6	500.2, 467.4, 439	4.11
726	-Benzyl	-Cl	piperidino	502.0	502.2, 469.4, 441.2	4.50
727	-Benzyl	-Cl	piperidino	502.0	502.2, 469.4, 441.4	4.60
728	-Benzyl	-Cl	morpholino	504.0	504.0, 471.4, 443.2	4.29
729	-Benzyl	-OMe	1-pyrrolidinyl	483.6	484.2, 451.2, 423	4.34

Example 365**Step 1: 2H-2(Z)-(4-Methoxybenzylidene)-1,4-benzothiazin-3(4H)-one**

5 A mixture of 2H-1,4-benzothiazin-3(4H)-one (5 g), *p*-anisaldehyde (6.6 g), and sodium methoxide (11.63 g) in 260 mL of DMF was heated to reflux for 15 hrs. The reaction mixture was cooled down to r.t., and water was added. The resulting precipitate was filtered and washed with water and ethanol to give the product.

Step 2: 2H-2(Z)-(4-Methoxybenzylidene)-4-(N-ethoxycarbonylmethyl)-1,4-benzothiazin-3(4H)-one

15 To a DMF (60 mL) solution of 2H-2(Z)-(4-methoxybenzylidene)-1,4-benzothiazin-3(4H)-one (1.4 g), NaH (160 mg) was added at 4 °C. After 1 hr's stirring, ethyl bromoacetate (684 mg) was added and the reaction mixture was warmed to room temperature. After 2 hrs'

additional stirring, the reaction mixture was added ethyl bromoacetate (68 mg) and left at r.t. overnight. The reaction mixture was added 5% KHSO₄ aq. and extracted with EtOAc. The organic layer was washed with sat. NaHCO₃ aq. and brine, and dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give the product.

Step 3: 2*H*-2(*Z*)-(4-Methoxybenzylidene)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid

2*H*-2(*Z*)-(4-methoxybenzylidene)-4-(*N*-ethoxycarbonylmethyl)-1,4-benzothiazin-3(4*H*)-one (1.4 g) was dissolved in ethanol (20 mL) and THF (20 mL). 5*N* NaOH aq. was added to the solution at room temperature and stirred for 3 hrs. The reaction mixture was added 5% KHSO₄ aq. and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure to give the product.

Step 4: 2*H*-2(*Z*)-(4-Methoxybenzylidene)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid *N*-hydroxylamide


Isobutyl chloroformate (477 mg) was added dropwise to a THF (40 mL) solution of 2*H*-2(*Z*)-(4-methoxybenzylidene)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid (1.19 g) and *N*-methylmorpholine (353 mg) at -15 to -20 °C. THF (10 mL) solution of *O*-TBDMS-hydroxylamine (565 mg) was added after additional 5 minutes' stirring. The mixture was warmed to

r.t. with stirring, and stood overnight. The reaction mixture was quenched with sat. Na_2CO_3 aq. and extracted with EtOAc. The organic layer was washed with 1N HCl and brine, dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by recrystallization from mixed solvent of THF and hexane to give 1.2 g of the hydroxamic acid as colorless solid.



^1H -NMR ($\text{DMSO}-d_6$, δ) 1.82(s, 3H), 4.60(s, 1.6H), 4.88(s, 0.4H), 7.07(d, $J=8.8\text{Hz}$, 2H), 7.04-7.11(m, 1H), 7.27(dt, $J=1.4\text{Hz}$, 8.2Hz, 1H), 7.38(dd, $J=1.1\text{Hz}$, 7.7Hz, 1H), 7.68(d, $J=8.8\text{Hz}$, 2H), 7.73(s, 1H), 8.98(s, 0.8H), 9.43(s, 0.2H), 10.3(s, 0.2H), 10.8(s, 0.8H)


The following compounds listed in the Table 5 were prepared in a similar manner.

Table 5



Example No.	RR1	RR2	RR3	mol. weight	Mass (m/e)	HPLC rt (min)
365	-C6H4-4-OMe	-H	-H	356.4	357.0, 324.0, 296.2	3.05
366	-C6H4-4-OEt	-H	-H	370.4	371.0, 338.0	3.43
367	-C6H4-4-SMe	-H	-H	372.5	373.2, 339.8	3.46
368	-C6H4-4-OPh	-H	-H	418.5	419.0, 386.0	3.40
369	-C6H4-4-O-nBu	-H	-H	398.5	399.2, 366.2	3.84, 3.91
370	-C6H4-4-OMe	-H	-Cl	390.8	391.0, 358.0	3.26
371	-C6H4-4-OMe	-CF3	-H	424.4	425.0, 392.0	3.35
372	-C6H4-4-O-benzyl	-H	-H	432.5	433.2, 400.2	3.54
373	-C6H4-4-O-nPr	-H	-H	384.5	385.2, 352.2	3.42
374	-C6H4-4-OCHF2	-H	-H	392.4	393.0, 360.0	3.18, 3.26
375	-C6H4-3-OMe	-H	-H	356.4	357.2, 324.2	2.92
376	-C6H3-3,4-Cl2	-H	-H	395.3	395.0, 362.0	3.29
377	-C6H4-2-OMe	-H	-H	356.4	357.2, 324.2	2.43, 2.56
378	-2-Py	-H	-H	327.4	328.2, 267.0	1.46, 1.92
379	-2-Pyrrolyl-1-(C6H3-2,6-Cl2)	-H	-H	460.3	459.8, 427.0	3.48
380	-C6H3-3,5-(OMe)2	-H	-H	386.4	387.0, 354.2	3.00
381	-C6H3-2-O-benzyl,3-OMe	-H	-H	462.5	463.2, 430.2, 374.0	3.22
382	-C6H2-2-F,4,5-(OMe)2	-H	-H	404.4	404.5, 372.2	2.80
383	-C6H2-2-F,3-CN,4-NMe2	-H	-H	412.4	413.0, 380.2	3.00
384	-C6H2-2,3,4-(OMe)3	-H	-H	416.5	417.0, 384.0, 297.2	2.81, 2.91
385	-C6H3-3,5-(CF3)2	-H	-H	462.4	462.8, 402.2	3.45
386	-C6H4-4-F	-H	-H	344.4	345.0, 312.2	3.30
387	-C6H4-4-Me	-H	-H	340.4	341.0, 308.2	3.44
388	-Ph	-H	-H	326.4	327.0, 294.0	3.23
389	-2-Furyl-5-(C6H4-4-Br)	-H	-H	471.3	471.0, 438.0, 371.2	4.03

390		-H	-H	480.5	482.0, 420.0	3.28
391		-H	-H	416.5	417.2, 384.0	3.42
392	-C6H4-4-O(3,4-Cl2-benzyl)	-H	-H	501.4	501.0, 470.2	3.73
393	-C6H4-4-Ph	-H	-H	402.5	403.2, 370.2	3.42
394	-3-Pyrrolidyl-4-CO2Et, 2,5-Me2, 1-Ph	-H	-H	491.6	492.2, 459.2	3.72
395	-C6H4-4-CF3	-H	-H	394.4	395.2, 362.2, 315.4	3.53, 3.60
396	-C6H4-4-iBu	-H	-H	382.5	383.2, 350.2	4.02
397	C6H4-4-(1-pyrrolidinyl)	-H	-H	395.5	396.2, 363.2	3.68
398	-2-Furyl-5-(C6H4-3-NO2)	-H	-H	437.4	438.2, 405.0	3.35
399	-2-Thienyl-3-Br	-H	-H	411.3	413.0, 380.0	3.14
400	-C6H4-4-(CH=CH-Ph)	-H	-H	428.5	429.0, 396.0	3.68
401	nPentyl	-H	-H	320.4	321.2, 288.2	3.38
402	-C6H4-4-SCF3	-H	-H	426.4	427.0, 394.0	3.55
403	-C6H4-4-SO2Me	-H	-H	404.5	405.0, 372.0	2.64, 2.72
404	-1-Naphthyl-4-OMe	-H	-H	406.5	407.0, 374.0	3.33, 3.41

Example 405

Step 1: 2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one
2H-2(Z)-(4-methoxybenzylidene)-1,4-benzothiazin-3(4H)-
one (142 mg) was dissolved in MeOH (5 mL) and THF (20 mL).
10% Pd/C (350 mg) was added. The reaction mixture was
vigorously stirred at r.t. for 4 hours under H₂. The
catalyst was removed by filtration, and the filtrate was
concentrated under reduced pressure. The residue was
purified by preparative thin layer chromatography on
silica gel (EtOAc/toluene, 1/4) to give the product as a
white solid (133.5 mg) .

Step 2: 2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-
one-4-yl-acetic acid

To a solution of 2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one (428 mg) in DMF (20 mL), NaH (66 mg, 60% in oil) was added. After 1 hour's stirring, ethyl bromoacetate (263 mg, 1.57 mmol) was slowly added. The

reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc and washed with 1N HCl, 5% Na₂CO₃ aq., and brine. The organic layer was dried over Na₂SO₄ and concentrated. The crude product was used in the following step without purification.


To a solution of ethyl ester (515 mg) in MeOH (10 mL), 1N NaOH (2.58 mL) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to room temperature and left overnight. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in 5% Na₂CO₃ aq., and washed with Et₂O. The aqueous layer was acidified with 4N HCl, and the product was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated to give the acetic acid (451 mg, 88%) as a white solid.

Step 3: 2*H*-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid N-hydroxyamide

The acetic acid (343 mg), obtained as above, was dissolved in CH₂Cl₂ (4 mL) and pyridine (4 mL) and pentafluorophenyl trifluoroacetate (560 mg) was added. After 3 hours' stirring, *O*-TBDMS-hydroxylamine (736 mg) was added. The reaction mixture was stirred at r.t. overnight. The solvent was removed and the residue was dissolved in EtOAc. The organic layer was washed with 1N HCl and brine, dried over Na₂SO₄ and concentrated under

reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel ($\text{CHCl}_3/\text{MeOH}$, 10/1) to give hydroxamic acid (323 mg) as a pale orange solid. $^1\text{H-NMR}$ ($\text{DMSO-d}_6, \delta$) 2.69(1H, m), 3.16(1H, m), 3.76(3H, s), 3.83(1H, m), 4.52(2H, m), 6.87(2H, d, $J=7.0$), 7.09-7.18(4H, m), 7.33-7.40(2H, m), 9.06(1H, br-s), 10.83(1H, s).

Example 406




Step 1: 2H-2-(4-Hydroxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide

2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide (373 mg) was dissolved in CH_2Cl_2 (10 mL). BBr_3 (1M solution in CH_2Cl_2 , 5 mL) was added and the reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure, and the residue was dissolved in EtOAc. The organic layer was washed with 1N HCl and brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel ($\text{CHCl}_3/\text{MeOH}$, 10/1) to give hydroxamic acid (161 mg) as a pale yellow solid.

$^1\text{H-NMR}$ ($\text{DMSO-d}_6, \delta$) 2.58(1H, m), 3.06(1H, m), 3.75(1H, s), 4.37-4.83(2H, m), 6.66(2H, d, $J=7.86\text{Hz}$), 6.97-7.13(4H, m),

7.27-7.38(2H, m), 8.99(1H, s), 9.26(1H, s), 10.77(1H, s)

Example 407




5 **Step 1: Ethyl [2*H*-2*Z*-(4-methylbenzylidene)-1,4-benzothiazin-3(4*H*)-one-4-yl]acetate**

To a stirred solution of *p*-tolualdehyde (48 mg) and 2*H*-2-diethylphosphonyl-1,4-benzothiazine-3(4*H*)-one-4-yl-acetic acid ethyl ester (78 mg) in ethanol (3 mL), which
10 was prepared by Worley's method (J. Org. Chem., 40, 1731 (1975)), was added a solution of sodium ethoxide (28 mg, 0.42 mmol) in ethanol (1 mL). The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc and
15 washed with sat. NaHCO₃ aq., 1N HCl, and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by preparative thin layer chromatography on silica gel (CHCl₃/MeOH, 15/1) to give a yellow solid (37 mg, 52%).

20 This compound was converted to 2*H*-2-(4-Methylbenzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid N-hydroxyamide by the reactions analogous to those described in Example 405.


The following compounds listed in the Tables 6 and 6' were prepared in a similar manner.


Table 6



Example No.	RR1	NMR: ¹ H-NMR(DMSO-d ₆ ; δ)
407	-C ₆ H ₄ -4-Me	2.27(3H, s), 2.65(1H, m), 3.15(1H, m), 3.83(1H, m), 4.43(1H, d, J=16Hz), 4.51(1H, d, J=16Hz), 7.04-7.14(6H, m), 7.28-7.38(2H, m), 8.98(1H, s), 10.76(1H, s)
408	-2-Py	2.91(1H, m), 3.33(1H, m), 4.12(1H, m), 4.48(2H, s), 7.05-7.11(2H, m), 7.24-7.37(4H, m), 7.73(1H, t, J=7.5Hz), 8.49(1H, d, J=4.05Hz), 9.03(1H, s), 10.90(1H, s)
409	-C ₆ H ₄ -4-OPh	2.72(1H, m), 3.17(1H, m), 3.87(1H, m), 4.49(2H, m), 6.90-7.65(13H, m), 9.01(1H, s), 10.80(1H, s)
410	-Ph	2.70(1H, m), 3.20(1H, m), 3.88(1H, m), 4.48(2H, m), 7.05-7.38(9H, m), 8.98(1H, s), 10.76(1H, s)
411	-C ₆ H ₄ -2-Me	2.22(3H, s), 2.69(1H, m), 3.16(1H, m), 3.85(1H, m), 4.49(2H, s), 7.06-7.16(6H, m), 7.29-7.40(2H, m), 9.00(1H, s), 10.78(1H, s)
412	-C ₆ H ₄ -3-Me	2.27(3H, s), 2.65(1H, m), 3.17(1H, m), 3.85(1H, m), 4.43(1H, d, J=16Hz), 4.52(1H, d, J=16Hz), 6.98-7.19(6H, m), 7.28-7.38(2H, m), 8.99(1H, s), 10.78(1H, s)
413	-C ₆ H ₄ -2-Me	2.92(1H, m), 3.35(1H, m), 3.91(1H, m), 4.49(2H, m), 7.06-7.17(2H, m), 7.30-7.72(6H, m), 9.02(1H, s), 10.80(1H, s)
414	-C ₆ H ₄ -2-OMe	2.67(1H, m), 3.18(1H, m), 3.76(3H, s), 3.79(1H, m), 4.42(1H, d, J=16Hz), 4.53(1H, d, J=16Hz), 6.87(1H, m), 6.96(1H, d, J=8.25Hz), 7.07-7.14(3H, m), 7.21-7.37(3H, m), 8.99(1H, s), 10.79(1H, s)
415	-C ₆ H ₄ -3-OMe	2.66(1H, m), 3.17(1H, m), 3.73(3H, s), 3.87(1H, m), 4.44(1H, d, J=16Hz), 4.51(1H, d, J=16Hz), 6.76-6.79(3H, m), 7.04-7.19(3H, m), 7.28-7.38(2H, m), 8.99(1H, s), 10.77(1H, s)

416	-C6H4-4-OEt	1.31(3H, m), 2.64(1H, m), 3.09(1H, m), 3.79(1H, m), 3.98(2H, m), 4.46(2H, m), 6.82(2H, d, J=6.78Hz), 7.04-7.12(4H, m), 7.27-7.37(2H, m), 8.99(1H, s), 10.77(1H, s)
417	-C6H3-3,4-(OMe)2	2.62(1H, m), 3.12(1H, m), 3.71(3H, s), 3.72(3H, s), 3.84(1H, m), 4.47(2H, s), 6.68(1H, m), 6.83(2H, m), 7.04-7.14(2H, m), 7.28-7.39(2H, m), 8.98(1H, s), 10.77(1H, s)
418	-C6H4-4-Bu	0.89(3H, t, J=7Hz), 1.30(2H, m), 1.54(2H, m), 2.54(2H, m), 2.66(1H, m), 3.14(1H, m), 3.83(1H, m), 4.47(2H, m), 6.95-7.13(5H, m), 7.28-7.38(3H, m), 8.99(1H, s), 10.77(1H, s)
419	-C6H4-4-Ph	2.77(1H, m), 3.25(1H, m), 3.93(1H, m), 4.49(2H, m), 7.05-7.16(2H, m), 7.30-7.48(7H, m), 7.58(2H, d, J=7.14Hz), 7.65(2H, d, J=8.25Hz), 8.99(1H, s), 10.78(1H, s)
420	-(CH2)2-Ph	1.64-1.81(m, 2H), 2.56(m, 2H), 3.61(m, 3H), 4.38(d, J=16Hz, 1H), 4.50(d, J=16Hz, 1H), 7.02-7.07(m, 2H), 7.07-7.18(m, 3H), 7.18-7.30(m, 3H), 7.39(m, 1H), 8.96(s, 0.8H), 9.34(s, 0.2H), 10.3(s, 0.2H), 10.7(s, 0.8H)
421	-C6H4-4-O(CH2)2OH	2.56(1H, m), 3.05(1H, m), 3.74(1H, m), 3.83-3.92(4H, m), 4.40(2H, m), 6.76(2H, d, J=8.22Hz), 6.97-7.06(4H, m), 7.21-7.30(2H, m), 8.92(1H, s), 10.70(1H, s)
422	-C6H4-4-NEt2	1.06(6H, t, J=7Hz), 2.54(1H, m), 3.05(1H, m), 3.26-3.34(4H, m), 3.74(1H, m), 4.47(2H, m), 6.55(2H, d, J=8Hz), 6.97(2H, d, J=8Hz), 7.04-7.13(2H, m), 7.27-7.38(2H, m), 8.98(1H, s), 10.77(1H, s)
423	-3-Py	2.77(1H, m), 3.18(1H, m), 3.96(1H, m), 4.48(2H, m), 6.96-7.14(2H, m), 7.29-7.40(3H, m), 7.68(1H, d, J=7.71Hz), 8.42(2H, m), 9.00(1H, s), 10.78(1H, s)
424	-4-Py	2.75(1H, m), 3.20(1H, m), 4.01(1H, m), 4.47(2H, m), 7.06-7.15(2H, m), 7.28(2H, d, J=5.67Hz), 7.32-7.40(2H, m), 8.46(2H, d, J=5.67Hz), 9.00(1H, s), 10.78(1H, s)
425	-C6H4-4-Cl	2.73(1H, m), 3.16(1H, m), 3.90(1H, m), 4.47(2H, m), 7.05-7.14(2H, m), 7.24-7.38(6H, m), 8.99(1H, s), 10.78(1H, s)
426	-C6H4-4-Br	2.72(1H, m), 3.14(1H, m), 3.90(1H, m), 4.47(2H, m), 7.05-7.14(2H, m), 7.20(2H, d, J=8.25Hz), 7.29-7.39(2H, m), 7.47(2H, d, J=8.25Hz), 8.99(1H, s), 10.78(1H, s)

Table6'		
Example No.	RR¹	NMR: ¹H-NMR (DMSO-d6; δ)
754	-O-C6H4-4-F	2.71(1H, dd, J=9.0, 14Hz), 3.16(1H, dd, J=5.9, 14Hz), 3.85(1H, dd, J=5.9, 9.0Hz), 4.48(1.6H, br-s), 4.67(0.2H, d, J=18Hz), 4.88(0.2H, d, J=18Hz), 6.88(2H, d, J=8.4Hz), 6.95-7.39(10H, m), 8.98(0.8H, s), 9.36(0.2H, s), 10.33(0.2H, s), 10.77(0.8H, s)
755	-F	2.73(1H, m), 3.16(1H, m), 3.88(1H, m), 4.47(2H, m), 6.96-7.14(4H, m), 7.24-7.39(4H, m), 8.99(1H, s), 10.78(1H, s)
756	-CF3	2.85(1H, m), 3.27(1H, m), 3.98(1H, m), 4.48(2H, m), 7.06-7.16(2H, m), 7.30-7.40(2H, m), 7.48(2H, J=8.07Hz), 7.64(2H, J=8.07Hz), 8.99(1H, s), 10.79(1H, s)

Example 427**Step 1: 2H-2-(4-Methoxybenzyl)-7-methyl-1,4-benzothiazin-3(4H)-one**

5 A mixture of 2-amino-6-methylbenzothiazole (657 mg) in 20% NaOH aq. (10 mL) was heated at 180 °C in an autoclave for 5 hours. The mixture was cooled to 80 °C and a solution of 2-bromo-3-(4-methoxyphenyl)propionic acid (1.088 g) in 10% NaOH aq. (10 mL) was added and the
10 temperature was maintained at 80 °C for 1 hour. The mixture was cooled to r.t. and concentrated under reduced pressure. The residue was dissolved in AcOH (40 mL) and heated at 100 °C for 4 hours. The reaction mixture was cooled to r.t. and concentrated under reduced pressure.
15 The residue was dissolved in EtOAc and washed with water and 5% Na₂CO₃ aq. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by recrystallization from Et₂O and hexane to give a white solid (805 mg).

Step 2: 2H-2-(4-Methoxybenzyl)-7-methyl-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide

20 The product of Step 1 was converted to hydroxamic acid

as described in Example 405(step 2 and 3).

The following compounds listed in the Tables 7, 7' and 7'' were prepared in a similar manner.

Table 7




Table 7					
Example No.	RR1	RR2	RR3	NMR: ¹ H-NMR(DMSO-d6; δ)	
427	-H	-H	-Me	2.29(3H, s), 2.68(1H, m), 3.15(1H, m), 3.77(3H, s), 3.81(1H, m), 4.45(1H, d, J=16Hz), 4.51(1H, d, J=16Hz), 6.88(2H, d, J=8.07Hz), 7.04-7.22(5H, m), 9.02(1H, s), 10.80(1H, s)	
428	-Me	-H	-H	2.40(3H, s), 2.71(1H, m), 3.21(1H, m), 3.69(1H, m), 3.76(3H, s), 4.19(2H, br-s), 6.87(2H, d, J=7.86Hz), 7.02-7.28(5H, m), 8.90(1H, bs), 10.60(1H, s)	
429	-Cl	-H	-H	2.73(1H, m), 3.19(1H, m), 3.76(3H, s), 3.84(1H, m), 4.43(2H, s), 6.87-6.90(2H, d, J=8.61Hz), 7.13-7.19(3H, m), 7.44-7.48(2H, m), 8.89(1H, bs), 10.63(1H, s)	
430	-OMe	-H	-H	2.66(1H, m), 3.17(1H, m), 3.68(1H, m), 3.72(3H, s), 3.80(3H, s), 4.20(1H, d, J=15Hz), 4.40(1H, d, J=15Hz), 6.84(2H, d, J=8.25Hz), 6.95-7.15(5H, m), 8.76(1H, bs), 10.40(1H, s)	
431	-H	-Me	-Me	2.16(3H, s), 2.21(3H, s), 2.62(1H, m), 3.08(1H, m), 3.72(3H, s), 3.73(1H, m), 4.43(2H, s), 6.83(2H, d, J=8.61Hz), 6.93(1H, s), 7.10-7.13(3H, m), 8.98(1H, s), 10.74(1H, s)	
432	-H	-H	-F	2.70(1H, m), 3.16(1H, m), 3.77(3H, s), 3.90(1H, m), 4.47(1H, d, J=16Hz), 4.54(1H, d, J=16Hz), 6.88(2H, d, J=8.61Hz), 7.16-7.22(4H, m), 7.36(1H, d, J=7.68Hz), 9.05(1H, bs), 10.82(1H, s)	

Table 7'				
Example No.	RR ¹	RR ²	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)	
757	-OMe	-NO ₂	2.69(1H,dd,J=9,14Hz),3.08(1H,dd,J=6,14Hz),3.71(3H,s),3.99(1H,dd,J=6,9Hz),4.5-5.0(2H,m),6.82(2H,d,J=8.6Hz), 7.13(2H,d,J=8.6Hz), 7.34(1H,d,J=9Hz),8.14(1H,dd,J=2.6,9Hz),8.24(1H,d,J=2.6Hz),9.02,9.42(1H,bs),10.42,10.83(1H,bs).	
758	-OMe	-SO ₂ Me	2.78(1H,dd,J=8.8Hz,14Hz),3.19(1H,dd,J=6,14Hz),3.31(3H,s),3.68(3H,s),4.03(1H,dd,J=6,8.8Hz),4.63-5.05(2H,m),6.92(2H,d,J=8.5Hz), 7.22(2H,d,J=8.5Hz),7.44(1H,d,J=8.6Hz),7.88(1H,dd,J=2,8.6Hz),7.97(1H,d,J=2Hz),9.09,9.56(1H,bs),10.48,10.92(1H,bs).	
759	-OCF ₃	-SO ₂ Me	2.85(1H,m),3.19(1H,m),3.22(3H,s),4.06(1H,dd,J=6.5,8Hz),4.54-4.99(2H,m),7.30(5H,m),7.80(1H,d,J=8.6Hz),7.92(1H,s),9.43,9.01(1H,bs),10.83,10.41(1H,bs).	

Table 7"								
Example No.	RR¹	RR²	RR³	mol. weight	Mass (m/e)	HPLC rt (min)		
760	-H	-H	-NMe2	371.5	372.2, 339.2	2.29		
761	-H	-H	-CF3	396.4	396.8, 364.0, 336	3.00		
762	-OCF3	-H	-SO2Me	490.5	491.1, 458.0	2.71		
763	-OCF3	-H	-H	412.4	413.2, 380.2, 352	3.11		
764	-OCF3	-CF3	-H	480.4	481.2, 448.0	3.20		
765	-H	-Me	-Me	356.4	357.0, 324.2	2.97		
766	-OCF3	-H	-F	430.4	431.2, 398.0	2.99		
767	-H	-H	-OMe	358.4	359.2, 326.2, 298	2.74		
768	-H	-H	-SO2Me	406.5	407.2, 374.0, 446	2.46		
769	-OCF3	-Me	-Me	440.4	441.4, 408.2	3.17		
770	-H	-H	-Me	342.4	343.0, 310.2	2.84		
771	-H	-H	-F	346.4	347.4, 314.0	3.11		
772	-H	-H	-OEt	372.4	372.9, 340.2	3.23		
773	-H	-H	-Cl	362.8	363.2, 330.2	2.97		
774	-OCF3	-H	-Cl	446.8	446.9, 414.2	3.16		
775	-H	-H	-NO2	373.4	374.0, 341.0	2.76		
776	-H	-H	-Br	407.3	407.0, 374.0, 346	3.37		

Example 433

Step 1: 2H-2-(p-methoxybenzyl)-1,4-pyrido[3,4-e]thiazin-3(4H)-one

To a mixture of 3-Amino-4-mercaptopyridine $\cdot 2\text{HCl}$ (400 mg) and 2-bromo-3-(p-methoxyphenyl)propionic acid (520 mg) in DMF (8 mL) was added K_2CO_3 (1.38 gr) and the mixture was stirred for 3 hrs at r.t. EDC $\cdot \text{HCl}$ (800 mg) and 1-hydroxybenzotriazole (900 mg) were added to the reaction mixture and the stirring was continued overnight. The reaction mixture was diluted with aq. NaHCO_3 and extracted with EtOAc 3 times. The combined organic layers were washed with aq. NaHCO_3 , aq. NaH_2PO_4 and brine. Column chromatography on silica gel (eluent; $\text{CHCl}_3/\text{MeOH}=40/1$) gave the product as a white solid (390 mg).

Step 2: 2H-2-(p-methoxybenzyl)-1,4-pyrido[3,4-e]thiazin-3(4H)-one-4-yl-acetic acid ethyl ester

To a THF (15 mL) solution of 2H-2-(p-methoxybenzyl)-1,4-pyrido[3,4-e]thiazin-3(4H)-one (385 mg) was added NaH (60%

in oil, 39 mg) at 0 °C, and the mixture was stirred for 30 min. at the same temperature and for 30 min at r.t. After the reaction mixture was cooled to 0°C, ethyl bromoacetate (164 µL) was added and the reaction mixture was stirred for 3 hrs at r.t. The reaction was quenched by the addition of aq. NaH₂PO₄ and extracted with EtOAc several times. Column chromatography on silica gel (eluent; CHCl₃/EtOAc=5/1) gave the product almost quantitatively as a colorless oil.

Step 3: 2*H*-2-(*p*-methoxybenzyl)-1,4-pyrido[3,4-*e*]thiazin-3(4*H*)-one-4-yl-acetic acid *N*-hydroxyamide


The compound obtained above was treated with 5N NaOH (0.8 mL) in MeOH (8 mL) for 1 hr at r.t. The reaction mixture was concentrated *in vacuo*, diluted with water and the pH was adjusted to 5.2 by 1N HCl. Lyophilization gave the corresponding carboxylic acid as a white powder.

The white powder in CH₂Cl₂ (15 mL) and pyridine (0.8 mL) was treated with pentafluorophenyl trifluoroacetate (400 µL) for 3 hrs at r.t., and the resulting mixture was treated with *O*-TMS-hydroxylamine (320 µL) and stirred overnight at r.t. The reaction mixture was concentrated *in vacuo* and the residue was dispersed in water and cyclohexane. The white precipitate was filtered and purified on silica gel (eluent: CHCl₃/MeOH=20/1), affording the product as a white solid. (315 mg)

NMR(*d*₆-DMSO, δ): 2.70(1H,dd,J=14.3, 9.0), 3.11(1H,dd,J=14.3,

6.1), 3.72(3H,s), 3.99(1H,dd,J=9.0, 6.1), 4.57(2H,s),
6.83(2H,d,J=8.6), 7.13(2H,d,J=8.6), 7.42(1H,d,J=5.1),
8.18(1H,d,J=5.1), 8.35(1H,s), 9.00(1H,s), 10.81(1H,s)

5 **Example 434**



Step 1: Ethyl (2S)-2-[2'-(2-bromophenyl)thio-3'-(4-methoxyphenyl)propanoyl]amino-4-methylpentanoate

To a solution of 2-(2-bromophenyl)thio-3-(4-methoxyphenyl)propanoic acid (1.102 g) and L-leucine ethyl ester·HCl (704 mg) in DMF (10 mL), HOBt (551 mg) and EDC·HCl (690 mg) were added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc and washed with 1N HCl, 5% Na₂CO₃ aq. and brine. The organic layer was dried over Na₂SO₄ and concentrated to give the pure product as a pale yellow oil (1.561 g, quant.).

Step 2: Ethyl (2S)-2-[2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl]-4-methylpentanoate

The product obtained in Step 1 (763 mg), $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (78 mg), $\text{P}(\text{o-tolyl})_3$ (92 mg), and Cs_2CO_3 (977 mg) in toluene (12 mL) were heated to 100°C in a culture tube (PYREXPLUS[®]) under nitrogen for 40 hrs. The reaction mixture was then cooled to r.t., diluted with EtOAc, filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel (toluene/EtOAc, 20/1) to give the product as a pale yellow oil (274 mg).


Step 3: (2S)-2-[2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl]-4-methylpentanoic acid N-hydroxyamide

To a solution of the compound obtained in Step 2 (274 mg) in ethanol (5 mL) and THF (5 mL), 1N NaOH (2.56 mL) was slowly added at 0°C . After 30 minutes' stirring, the mixture was warmed to room temperature and left over 1 day. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in EtOAc and washed with 1N HCl. The organic layer was dried over Na_2SO_4 and concentrated. The crude acid was used in the following step. The acid (204 mg) was dissolved in CH_2Cl_2 (3 mL) and pyridine (3 mL) and pentafluorophenyl trifluoroacetate (716 mg) was added. After 3 hours' stirring, O-TBDMS-hydroxylamine (452 mg) was added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure, and the residue was dissolved in EtOAc. The organic layer was


washed with 1N HCl and brine, and dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel (toluene/EtOAc=1/1) to give the hydroxamic acid (208 mg) as a pale orange solid.

The following compounds listed in the Table 8 were prepared in a similar manner.

Table 8



Example No.	RR1	NMR: ¹ H-NMR(DMSO-d ₆ ; δ)
434	(S) -iBu	0.57-0.77(6H, m), 1.00(1H, m), 1.70(1.5H, m), 2.02(0.5H, m), 2.58(0.5H, m), 2.79(0.5H, m), 3.02(0.5H, m), 3.35(0.5H, m), 3.72(1.5H, s), 3.73(1.5H, s), 3.78(1H, m), 5.14(0.5H, m), 5.56(0.5H, m), 6.82-6.88(2H, m), 7.03-7.46(6H, m), 8.94(0.5H, s), 9.00(0.5H, s), 10.68(0.5H, s), 10.79(0.5H, s)
435	(R) -iBu	0.57-0.77(6H, m), 1.00(1H, m), 1.73(1.5H, m), 2.02(0.5H, m), 2.59(0.5H, m), 2.76(0.5H, m), 3.02(0.5H, m), 3.42(0.5H, m), 3.72(1.5H, s), 3.73(1.5H, s), 3.78(1H, m), 5.16(0.5H, m), 5.56(0.5H, m), 6.82-6.88(2H, m), 7.06-7.46(6H, m), 8.94(0.5H, s), 9.00(0.5H, s), 10.68(0.5H, s), 10.79(0.5H, s)
436	(S) -iPr	0.43(3H, m), 0.96(3H, m), 2.26(0.5H, m), 2.45(0.5H, m), 2.80(1H, m), 3.17(1H, m), 3.71(1.5H, s), 3.73(1.5H, s), 3.80(1H, m), 4.77(0.5H, d, J=10.9Hz), 4.94(0.5H, d, J=10.9Hz), 6.81-6.87(2H, m), 6.94-7.42(6H, m), 9.05(1H, bs), 10.81(1H, m)
437	(S) -Me	1.27(1.5H, d, J=7.32Hz), 1.39(1.5H, d, J=6.96Hz), 2.59(0.5H, m), 2.70(0.5H, m), 3.02(0.5H, m), 3.34(0.5H, m), 3.72(1.5H, s), 3.73(1.5H, s), 3.82(1H, m), 5.08(0.5H, m), 5.49(0.5H, m), 6.82-6.88(2H, m), 7.03-7.41(6H, m), 8.86(0.5H, s), 8.95(0.5H, s), 10.56(0.5H, s), 10.70(0.5H, s)
438	(R) -Me	1.27(1.5H, d, J=7.32Hz), 1.39(1.5H, d, J=6.96Hz), 2.59(0.5H, m), 2.71(0.5H, m), 3.00(0.5H, m), 3.34(0.5H, m), 3.72(1.5H, s), 3.73(1.5H, s), 3.78(1H, m), 5.08(0.5H, m), 5.49(0.5H, m), 6.82-6.88(2H, m), 7.03-7.41(6H, m), 8.86(0.5H, s), 8.95(0.5H, s), 10.56(0.5H, s), 10.70(0.5H, s)

Example 439

Step 1: 2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid t-butyl ester

To a solution of 2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one (2 g) in THF (20 mL), NaH (304 mg) and subsequently t-butyl bromoacetate (1.13 mL) were added at 0 - 5 °C. The mixture was stirred at the same temperature for 5 minutes and at r.t. for 30 minutes. The reaction was quenched by adding 150 mL of water, and extracted with EtOAc. The organic layer was washed with brine, and dried over MgSO₄ and concentrated under the reduced pressure, and residue was purified by silica gel column chromatography (Eluent: Hexane:EtOAc=10:1) to give 3.0 g of the product as oil.

Step 2: 2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid t-butyl ester 1-oxide and 1,1-

dioxide

Product in Step 1 (500 mg) was dissolved in MeOH (10 mL) and *m*-CPBA (432 mg) was added at 0 - 5 °C. The mixture was stirred at the same temperature for 5 min. and warmed to r.t. After stirred for 1 hour, 100 mL of water was added and extracted with EtOAc. The organic layer was washed with sat. NaHCO₃ aq. and brine, and dried over MgSO₄ and concentrated under the reduced pressure, and the residue was purified with silica gel column chromatography (Eluent: Hexane:EtOAc=5:1) to give 270 mg of 1,1-dioxide and 370 mg of 1-oxide as oil.

Step 3: 2*H*-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid 1,1-dioxide


To the sulfone of Step 2 (270 mg), was added TFA (2 mL) at 0 - 5 °C. The mixture was warmed to r.t. and stirred for 2 hrs. 100 mL of Et₂O was added and the resulting precipitate was filtered and washed with Et₂O, giving 245 mg of the corresponding acid.

Step 4: 2*H*-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid N-hydroxyamide 1,1-dioxide

The product of Step 3 was converted to hydroxamic acid by the method described in Example 405, giving 95.6 mg of the product as colorless solid.

¹H-NMR (DMSO-*d*₆, δ) 3.06(1H, dd, J=8.6Hz, 14.6Hz), 3.21(1H, dd, J=4.0Hz, 14.6Hz), 3.70(3H, s),

4.51(1H,d,J=17Hz), 4.57(1H,d,J=17Hz), 4.99(1H,dd,
J=4.0Hz,8.6Hz), 6.80(2H,d,J=8.4Hz), 7.20(2H,d,J=8.4Hz),
7.38(2H,m), 7.78(1H,t,J=7.3Hz), 7.88(2H,d,J=7.1Hz),
9.03(0.8H,bs), 9.42(0.2H,bs), 10.43(0.2H,bs),
10.79(0.8H,bs).

Example 440

Step 1: 2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid 1-oxide


The product (290 mg) was obtained as colorless precipitate from the sulfoxide (370 mg) of Example 439 (Step 2) by the method described in Example 439 (Step 3).

Step 2: 2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide 1-oxide

The product of Step 1 was converted to hydroxamic acid as described in Example 439 to give 95 mg of the product.

¹H-NMR (DMSO-d₆, δ) 3.09(1H,m), 3.49(1H,b), 3.76(3H,m),
4.40-5.00(3H,m), 6.30-6.50(2H,m), 7.00-7.50(4H,m), 7.60-
7.90(2H,m), 9.04-9.45(1H,m), 10.78-10.87(1H,m).

Example 441



Step1: 2H-2-(Ethoxycarbonylmethyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid t-butyl ester

To a solution of 2H-2-(ethoxycarbonylmethyl)-1,4-benzothiazin-3(4H)-one (2 g) in THF (20 mL) was added sodium hydride (335 mg), and t-butyl bromoacetate (1.23 mL) was added dropwise at 0 - 5 °C. The mixture was stirred for 5 min. at the same temperature and for 30 minutes at r.t. 150 mL of water was added and extracted with EtOAc. The organic layer was washed with brine, and dried over MgSO₄ and concentrated under the reduced pressure to give 3.1 g of the product as oil.

Step2: 2H-2-[N'-(4-methoxyphenyl)carbamoylmethyl]-1,4-benzothiazin-3(4H)-one-4-yl-acetic acid t-butyl ester

To a solution of the product of step1 in MeOH (20 ml), 1N NaOH (7.96 ml) was added at 0 °C. Stirring was continued for 3 hrs. The reaction mixture was diluted with water, and washed with Et₂O. The aqueous layer was acidified with 1N HCl and the product was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated to give the corresponding acetic acid (2.79

g) as oil.


To the solution of the acetic acid (200 mg) in DMF (20 ml), 4-anisidine (77 mg) and EDC•HCl (119 mg) were added at 0 °C. Stirring was continued for 5 hours. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with 1N HCl, sat. NaHCO₃ and brine, dried over MgSO₄ and concentrated under the reduced pressure. The residue was purified with silica gel column chromatography (Eluent: Hexane:EtOAc=3:1) to give the product (220 mg).

Step3: 2*H*-2-[N'-(4-Methoxyphenyl)-carbamoylmethyl]-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid N-hydroxylamide


The product of Step 2 was converted to the hydroxamic acid as described in Example 439 to give 141mg of the product.

The following compounds listed in the Table 9 were prepared in a similar manner.

Table 9



Example No.	RR1	NMR: ¹ H-NMR(DMSO-d ₆ ; δ)
441	-C ₆ H ₄ -4-OMe	2.55(1H,dd,J=15Hz,8.8Hz), 2.94(1H,dd,J=15Hz,5.5Hz), 3.70(3H,s), 4.01Hz(1H,dd,J=8.6Hz,5.5Hz), 4.34-4.76(2H,m), 6.86(2H,d,J=9.0Hz), 7.08(2H,m), 7.31(1H,m), 7.43(3H,m), 8.97,9.37(1H,s), 9.91,9.95(1H,s), 10.34,10.79(1H,s).
442	-CH ₂ -C ₆ H ₄ -4-OMe	2.42(1H,dd,(J=9.0Hz,15Hz), 2.81(1H,dd,J=5.0Hz,15Hz), 3.71(3H,s), 3.95(1H,m),4.17(2H,m), 4.33-4.76(2H,m), 6.87(2H,d,J=8.6Hz), 7.05-7.41(6H,m), 8.46(1H,t,J=5.6Hz), 8.97,9.39(1H,s), 10.33,10.79(1H,s).
443	-(CH ₂) ₂ -C ₆ H ₄ -4-OMe	2.30(1H,dd,J=9.5Hz,15Hz), 2.64(3H,m), 3.22(2H,m), 3.70(3H,s), 3.89(1H,m), 4.33-4.74(2H,m), 6.83(2H,d,J=8.4Hz), 7.07-4.41(6H,m), 8.08(1H,m), 8.97,9.37(1H,s), 10.34,10.79(1H,s).

Example 444**Step 1: 2H-2-(4-Methoxybenzyl)-1,4-benzoxazin-3(4H)-one**

To a solution of 2-bromo-3-(4-methoxyphenyl)propionic acid (1.6 g) in DMF (10 ml), 2-aminophenol (731 mg), EDC·HCl (1.54 g), and HOBt (500 mg) were added at 0 °C. Stirring was continued for 2 hrs. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with 1N HCl, sat. NaHCO₃ and brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (Eluent: Hexane:EtOAc=5:1).

To a solution of the obtained crystal (547 mg) in DMF (3 ml), K₂CO₃ (300 mg) was added at 0 °C. Stirring was continued at room temperature for 2 hrs. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (Eluent: Toluene:EtOAc=6:1). 266 mg of the product was obtained as oil.


Step 2: 2H-2-(4-Methoxybenzyl)-1,4-benzoxazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide

The compound obtained in Step 1 (266 mg) was converted

to the hydroxamic acid derivative by the method described above and 152 mg of the product was obtained as colorless precipitate.

¹H-NMR (DMSO-d₆, δ) 2.90(1H, dd, J=9.0Hz, 14Hz),
 3.06(1H, dd, J=4.2Hz, 14Hz), 3.71(3H, s), 4.41(1H, d, J=16Hz),
 4.47(1H, d, J=16Hz), 4.83(1H, dd, J=4.2Hz, 9.0Hz),
 6.89(2H, d, J=8.6Hz), 6.89- 7.03(4H, m), 7.15(2H, d, J=8.6 Hz),
 8.98, (0.8H, bs), 9.40(0.2H, bs), 10.35(0.2H, bs),
 10.80(0.8H, bs).

Example 445



Step 1: 3-[2H-2-(4-Methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl]propanoic acid

To a solution of 2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one (570 mg) in THF (20 mL), NaOH powder (84 mg) was added. After 10 minutes' stirring, ethyl acrylate (600 mg) was slowly added. The reaction mixture was stirred at room temperature for 5 hours. The solvent was removed under reduced pressure and the residue was acidified with 4N HCl. The product was extracted with EtOAc and the organic layer was dried over Na₂SO₄ and concentrated. The crude product was used in the following

step without purification.


To a solution of the ethyl ester in MeOH (20 mL), 1N NaOH (2.0 mL) was slowly added. The mixture was stirred at r.t. overnight and concentrated under reduced pressure. The residue was dissolved in water, and the aqueous solution was washed with Et₂O and acidified with 4N HCl. The product was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated to give the propanoic acid (618 mg) as a white solid.

Step 2:

The product of Step 1 was converted to hydroxamic acid by the method described in example 405 (step 3 and 4).

¹H-NMR (DMSO-d₆, δ) 2.30(2H, m), 2.63(1H, m), 3.03(1H, m), 3.72(3H, s), 3.77(1H, m), 4.11(2H, m), 6.84(2H, d, J=8.04Hz), 7.06-7.11(3H, m), 7.32-7.42(3H, m), 8.82(1H, s), 10.53(1H, s)

Example 446



3-[2H-2-(4-Methoxybenzylidene)-1,4-benzothiazin-3(4H)-one-4-yl]propanoic acid N-hydroxylamide


The reaction procedure of Step 1 and 2 were similar to

the example 445 (Step 1 and 2).

LC-MS: MS(m/e) 371.0, 338.0, 296.2 (M.W.=370.4);

Retention Time= 3.07 min.

5 **Example 730**



Step 1: 4-Methyl-2-(S)-[2-(RS)-(4-chlorophenyl)-4-oxo-thiazolidin-3-yl]-pentanoic acid methyl ester

A suspension of (L)-leucine methyl ester (10 g, 55 mmol) and triethylamine (7.7 ml, 55 mmol) in 120 ml of dioxane was stirred at room temperature for 5 minutes. 3 Å

molecular sieves were added followed by *p*-chlorobenzaldehyde (17 g, 121 mmol). The mixture was heated at 80 °C for 4 hrs. Mercaptoacetic acid (14.2 ml, 203.5 mmol) was added to the mixture and the heating was continued for an additional 12 hrs. The reaction mixture was cooled to room temperature, and diluted with ethyl acetate (40 ml) and water (30 ml). The reaction mixture was filtered to remove the molecular sieves. The organic layer was diluted with ethyl acetate (100 ml), washed twice with 60 ml of saturated solution of sodium carbonate, twice with aqueous 1N HCl and once with water. The organic layer was dried with MgSO₄ and evaporated to give a residue that was carried out to the next step without further purification.

Step 2: 4-Methyl-2-(*S*)-[2-(*RS*)-(4-chlorophenyl)-4-oxo-thiazolidin-3-yl]-pentanoic acid

The residue obtained in step 1 was dissolved in dioxane (50 ml). 30 ml of saturated solution of LiOH were added and the mixture was stirred at room temperature for 3h (analytical LC/MS showed no traces of starting material). The mixture was acidified with conc. HCl until acidic pH and diluted with ethyl acetate (50 ml). The 2 layers were separated and the aqueous layer was extracted 3 times with ethyl acetate (30 ml). The organic layers were combined, dried with MgSO₄, filtered and evaporated to give an oil that solidified upon standing. The reaction mixture was further purified column by

chromathography with hexane-ethyl acetate (from 95:5 to 1:1) to give 14.4 g (yield for 2 steps: 80%) of the desired product as a 60:40 mixture of diastereoisomers.

5 **Step 3: Separation of Diastereoisomers**

The mixture of pure acids obtained in Step 2 was dissolved in refluxing toluene. Upon cooling 2g of the minor isomer crystallized out of the mixture and were used for further transformations.

10 **Step 4: 4-Methyl-2-(*S*)-[2-(*S*)-(4-chlorophenyl)-1,4-dioxo-thiazolidin-3-yl]-pentanoic acid N-hydroxyamide**


The crystal obtained in Step 3 (500 mg, 1.52 mmol) was dissolved in methanol (200ml). A solution of sodium periodate (326 mg, 1.5 mmol) in 25 ml of water was added and the heterogeneous mixture was stirred at room temperature for 3 days. When no starting material was observed by LC/MS, the reaction mixture was concentrated to 80% of its volume, acidified with acetic acid (2 ml) and diluted with water (10 ml) and ethyl acetate (25 ml). The 2 layers were separated and the water layer extracted 2 additionnal times with ethyl acetate (2X 20 ml). The organic layers were combined, dried with MgSO₄, evaporated under dryness to give the product as a white solid that was taken to the next step without purification.

25 To a solution of the product in 5 ml of CH₂Cl₂ (DCM) and 2 ml of pyridine was added pentafluorophenol trifluoroacetate (914 ml, 5.32 mmol). The solution was


stirred at room temperature for 1 h. O-(*tert*-
Butyldimethylsilyl)-hydroxylamine (1.2 g, 7.6 mmol) in 2
ml of CH₂Cl₂ was added to the activated ester and the
mixture was stirred at room temperature for an additional
5 12 h. The reaction mixture was acidified with HCl (to
cleave the TBDMS protective group) and stirred at room
temperature for 1h. After dilution with DCM (4ml) and
water (2ml), the 2 layers were separated, the organic
layer dried with MgSO₄, filtered and evaporated to give an
10 oil that was dissolved in DMSO (3 ml) and purified by
reverse phase HPLC. The Product (141 mg, 26%) was
obtained with 98% purity.

The following compounds listed in the Table 13 were
15 prepared in a similar manner.

Table 13



Example No.	RR ¹	RR ²	RR ³	mol. weight	Mass (m/e)	HPLC rt (min)
730	-iBu	-Cl	-H	358.8	359.2, 326.0, 298	1.99
731	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	566.1	566.2, 508.2, 475.2	3.05
732	-iBu	-F	-H	342.4	343.0, 310.2, 282	1.98
733	-iBu	-O-C ₆ H ₄ -4-F	-H	434.5	435.2, 402.2, 374	2.76
734	-iBu	-O-C ₆ H ₄ -4-F	-H	434.5	435.2, 402.2, 374	2.75
735	-iBu	-Cl	-H	358.8	359.2, 326.0, 298	2.24
736	-iBu	-O-C ₆ H ₄ -4-Cl	-H	450.9	451.0, 418.0	3.21
737	-iBu	-H	-H	324.4	325.2, 292.2, 264	1.91
738	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONHCH ₂ -2-furyl	588.1	588.2, 509.2	3.12
739	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONHCH ₂ -2-furyl	588.1	588.2	3.14
740	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	566.1	566.2, 509.2, 476.0	2.88
741	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	566.1	566.2, 509.2, 476.0	3.15
742	-iBu	-Cl	-CH ₂ CONHCH ₂ -2-furyl	496.0	496.2	3.10
743	-iBu	-Cl	-CH ₂ CONHMe	429.9	430.2	2.81
744	-iBu	-Cl	-CH ₂ CONHMe	429.9	430.2	2.43
745	-iBu	-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	474.0	474.2, 417.2	2.86
746	-iBu	-Cl	-CH ₂ CONH ₂	415.9	416.2, 483.0	2.35
747	-iBu	-Cl	-CH ₂ CONH ₂	415.9	416.2	2.61

Example 748**Step 1: 4-Methyl-2-(*S*)-[2-(*S*)-(4-chlorophenyl)-1,1,4-trioxo-thiazolidin-3-yl]-pentanoic acid N-hydroxyamide**


5 The crystal obtained from Example 730, Step 3 (100 mg, 0.30 mmol) was dissolved in acetic acid (2ml). A solution of potassium permanganate (66.4 mg, 0.42 mmol) in 2 ml of water was added dropwise during a period of 1 hour at 0 °C. The mixture was stirred at room temperature and followed by LC/MS. After 1h, no starting material was observed; the color of the reaction was removed with NaHS₂O₃, diluted with water and extracted with EtOAc. The 2 layers were separated and the water layer extracted 2 additional times with ethyl acetate (2X 20 ml). The organic layers were combined, dried with MgSO₄, evaporated under dryness to give the product as a white solid that was taken to the next step without purification.

20 To a solution of this solid in 1 ml of DCM and 0.5 ml of pyridine was added pentafluorophenol trifluoroacetate (231 ml, 1.34 mmol). The solution was stirred at room temperature for 1 h. O-(*tert*-Butyldimethylsilyl)-


hydroxylamine (270 mg, 1.83 mmol) in 1 ml of DCM was added to the activated ester and the mixture was stirred at room temperature for an additional 12 h. The reaction was acidified with HCl (to cleave the TBDMS protective group) and stirred at room temperature for 1h. After dilution with DCM (4ml) and water (2ml), the 2 layers were separated, the organic layer dried with MgSO_4 , filtered and evaporated to give an oil that was dissolved in DMSO (2 ml) and purified by reverse phase HPLC to yield the hydroxamic acid as a white solid.

The following compounds listed in the Table 14 were prepared in a similar manner.

Table 14



Example No.	RR ¹	RR ²	RR ³	mol. weight	Mass (m/e)	HPLC rt (min)
748	-iBu	-Cl	-H	374.8	375.0,	2.37
749	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONH ₂	524.0	524.2,	2.71
750	-iBu	-O-C ₆ H ₄ -4-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	582.1	582.2,	3.36
751	-iBu	-Cl	-CH ₂ CONH(CH ₂) ₂ OMe	490.0	490.2,	2.81
752	-iBu	-O-C ₆ H ₄ -4-F	-H	450.5	451.0,	3.22
753	-iBu	-O-C ₆ H ₄ -4-Cl	-H	466.9	467.2,	3.38

Example 777

Step 1: t-Butyl 2H-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4H)-one-6-carboxylate

To 2-mercapto-3-(4-methoxyphenyl)propionic acid (14g) in ethanol(100ml), t-butyl 4-chloro-3-nitrobenzoate(17g), and potassium fluoride(3.8g) were added at 0 °C. Potassium carbonate(9.1g) was slowly added at 0°C and the mixture was heated under reflux for 7 hrs. The solvent was removed under the reduced pressure and the residue was diluted with water. The solution was washed with Et₂O and extracted with EtOAc. The organic layer was washed with brine, and dried over MgSO₄ and concentrated under the reduced pressure. The solution of the residue in MeOH(120ml) was added dropwise to the suspension of Fe powder(11.4g) in MeOH(50ml)-NH₄Cl aq.(17.6g/80ml of water) at r.t. The mixture was stirred at 80°C for 3 hrs. AcOH(50ml) was added to the reaction mixture, and the stirring was continued for 3 hrs at 80°C. The reaction mixture was cooled to room temperature, poured into ice-water, and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under

the reduced pressure. The residue was purified with silica gel column chromatography (eluent: hexane/EtOAc, 8/1,5/1,3/1) to give the product as oil. This was crystallized from EtOAc-Hexane to give a colorless crystal(12.7g).

Step 2: Ethyl 6-t-butoxycarbonyl-2*H*-2-(4-methoxy-benzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetate

To sodium hydride (3.32g) suspended with THF (30 mL) was added a solution of the product of Step 1 (30.5 g) in THF (240 mL), and ethyl bromoacetate (13.1 mL) were added dropwise at 0-5 °C. The mixture was warmed to r.t. and left at r.t. overnight. Saturated aqueous ammonium chloride was added and extracted with EtOAc. The organic layer was washed with water and brine, dried over MgSO₄, and concentrated under the reduced pressure to give 37 g of the product as oil.


Step 3: 6-Carboxyl-2*H*-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-yl-acetic acid N-hydroxyamide

To a solution of ethyl 6-t-butoxycarbonyl-2*H*-2-(4-methoxy-benzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetate (3g) was added 5*N*-aqueous sodium hydroxide (1.44mL) at r.t. and stirred for 4hr at r.t. The reaction mixture was concentrated under reduced pressure. 5% KHSO₄ aq. was added, and extracted with EtOAc. The organic layer was washed with brine, and dried over MgSO₄ and concentrated under reduced

pressure to give 6-t-butoxycarbonyl-2*H*-2-(4-methoxy-benzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl acetic acid as oil. Isobutyl chloroformate (1.04 g) was added dropwise to a THF (30 mL) solution of 6-t-butoxycarbonyl-2*H*-2-(4-Methoxy-benzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-yl acetic acid and N-methylmorpholine (769 mg) at -15 to -20 °C. *O*-TMS-hydroxylamine (1.09 g) was added after additional 15 minutes' stirring. The mixture was warmed to r.t. with stirring, and stood overnight. The reaction mixture was quenched with sat.NaHCO₃ aq. and extracted with EtOAc. The organic layer was washed with 1N HCl, sat.NaHCO₃ aq. and brine, dried over MgSO₄ and concentrated under reduced pressure to give 2.25 g of 6-t-butoxycarboxyl-2*H*-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-ylacetic acid N-hydroxyamide as pale yellow oil. 5%-ethanedithiol in TFA (10mL) was added to dissolve 6-t-butoxycarboxyl-2*H*-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-ylacetic acid N-hydroxyamide, and stood at r.t. for 2hrs. Toluene (10mL) was added and concentrated under reduced pressure. The residue was purified by recrystallization from mixed solvent of THF and hexane to give 1.2 g of the hydroxamic acid as colorless solid.

The following compounds listed in the Tables 15, and 15' were prepared in a similar manner.


Table 15




Example No.	RR ¹	RR ²	RR ³	RR ⁴	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
777	-OMe	-COOH	-H	-H	2.68(1H, dd, J=9.0, 14Hz), 3.13(1H, dd, J=6.2, 14Hz), 3.72(3H, s), 3.90(1H, dd, J=6.2, 9.0Hz), 4.50(1.6H, br-s), 4.69(0.2H, d, J=19Hz), 4.91(0.2H, d, J=19Hz), 6.83(2H, d, J=8.4Hz), 7.13(2H, d, J=8.4Hz), 7.48(1H, d, J=7.9Hz), 7.61(1H, dd, J=1.3, 7.9Hz), 7.66(1H, d, J=1.3Hz), 9.04(0.8H, s), 9.44(0.2H, s), 10.39(0.2H, s), 10.80(0.8H, s)
778	-Cl	-CH ₂ CO ₂ H	-H	-H	2.73(1H, dd, J=9.2, 14.5Hz), 3.17(1H, dd, J=6.1, 14.5Hz), 3.57(2H, s), 3.89(1H, dd, J=6.9, 2Hz), 4.38-4.81(2H, m), 6.98(1H, d, J=8Hz), 7.06(1H, s), 7.2-7.5(5H, m), 8.9-9.7(1H, bs), 10.78, 10.34(1H, bs).
779	-C ₆ H ₄ -4-Cl	-H	-H	-H	2.76(1H, m), 3.23(1H, m), 3.91(1H, m), 4.4-5.0(2H, m), 7.12(2H, m), 7.35(4H, m), 7.50(2H, d, J=8.7Hz), 7.59(2H, d, J=8.3Hz), 7.68(2H, d, J=8.7Hz), 8.98, 9.38(1H, bs), 10.35, 10.78(1H, bs).
780	-C ₆ H ₄ -4-F	-H	-H	-H	2.76(1H, dd, J=9.2, 14.3Hz), 3.23(1H, dd, J=6.0, 14.3Hz), 3.93(1H, dd, J=6.0, 9.2Hz), 4.4-5.0(2H, m), 7.0-7.4(8H, m), 7.56(2H, d, J=8.1Hz), 7.70(2H, m), 8.99, 9.38(1H, bs), 10.34, 10.78(1H, bs).
781	-O-C ₆ H ₄ -4-Cl	-H	-H	-H	2.72(1H, dd, J=9.1, 14Hz), 3.17(1H, dd, J=5.9, 14Hz), 3.87(1H, dd, J=5.9, 9.0Hz), 4.45(0.8H, d, J=16Hz), 4.50(0.8H, d, J=16Hz), 4.68(0.2H, d, J=18Hz), 4.88(0.2H, d, J=18Hz), 6.94(2H, d, J=8.5Hz), 7.01(2H, d, J=8.5Hz), 7.07(1H, td, J=0.9, 7.6Hz), 7.13(1H, d, J=8.0Hz), 7.25(2H, d, J=8.5Hz), 7.31(1H, td, J=1.3, 8.0Hz), 7.38(1H, dd, J=1.3, 7.6Hz), 7.42(2H, d, J=8.9Hz), 8.99(0.8H, s), 9.40(0.2H, s), 10.34(0.2H, s), 10.74(0.8H, s)

782	-H	-CO ₂ H	-H	-H	2.73(1H, m), 3.20(1H, m), 3.98(1H, m), 4.46(2H, m), 7.21-7.42(5H, m), 7.50(1H, d, J=8.07Hz), 7.62(1H, d, J=8.04Hz), 7.67(1H, s), 9.06(1H, s), 10.84(1H, s)
783	-O-C ₆ H ₄ -4-Cl	-COO-tBu	-H	-H	1.55(9H, s), 2.75(1H, dd, J=8.8, 14Hz), 3.15(1H, dd, J=6.1, 14Hz), 3.96(1H, dd, J=6.1, 8.8Hz), 4.48(0.8H, d, J=17Hz), 4.56(0.8H, d, J=17Hz), 4.68(0.2H, d, J=18Hz), 4.96(0.2H, d, J=18Hz), 6.93(2H, d, J=8.4Hz), 7.00(2H, d, J=9.0Hz), 7.25(2H, d, J=8.4Hz), 7.42(2H, d, J=9.0Hz), 7.50(1H, d, J=8.1Hz), 7.58(1H, dd, J=1.5, 8.1Hz), 7.64(1H, br-s), 9.04(0.8H, d), 9.44(0.2H, s), 10.42(0.2H, s), 10.85(0.8H, s)
784	-O-C ₆ H ₄ -4-Cl	-CO ₂ H	-H	-H	2.77(1H, dd, J=9.0, 14Hz), 3.19(1H, dd, J=5.9, 14Hz), 3.95(1H, m), 4.52(1.6H, br-s), 4.69(0.2H, d, J=18Hz), 4.95(0.2H, d, J=18Hz), 6.92(2H, d, J=8.4Hz), 7.00(2H, d, J=9.0Hz), 7.25(2H, d, J=8.4Hz), 7.40(2H, d, J=9.0Hz), 7.48(1H, d, J=8.1Hz), 7.62(1H, dd, J=1.7, 8.1Hz), 7.67(1H, br-s), 9.05(0.8H, s), 9.44(0.2H, s), 10.39(0.2H, s), 10.84(0.8H, s), 12.37(1H, s)
786	-H	-COONa	-H	-H	2.73(1H, m), 3.20(1H, m), 3.98(1H, m), 4.46(2H, m), 7.21-7.42(5H, m), 7.50(1H, d, J=8.07Hz), 7.62(1H, d, J=8.04Hz), 7.67(1H, s), 9.06(1H, s), 10.84(1H, s)
787	-Cl	-CO ₂ H	-H	-H	2.75 (1H, dd, J=14.1, 9.0 Hz), 3.14 (1H, dd, J=14.3Hz, 6.2Hz), 3.99 (1H, dd, J=8.6Hz, 6.2Hz), [4.49 (s), 4.67 (d, J=17.9Hz), 4.90 (d, J=17.9Hz)] 2H, 7.24 (2H, d, J=8.6Hz), 7.32 (2H, d, J=8.6Hz), 7.48 (1H, d, J=8.0Hz), 7.60 (1H, dd, J=8.1Hz, 1.5Hz), 7.65 (1H, d, J=1.5Hz), 9.44(1H, br), [10.80 (br), 10.38 (s)] 1H, 13.02 (1H, br)
788	-OCF ₃	-CO ₂ H	-H	-H	2.81(1H, dd, J=8.8, 14.5Hz), 3.20(1H, dd, J=6.4, 14.5Hz), 4.04(1H, dd, J=6.4, 8.8Hz), 4.5-4.9(2H, m), 7.26(2H, d, J=8Hz), 7.36(2H, d, J=8Hz), 7.52(1H, m), 7.61(1H, dd, J=1.5, 8Hz), 7.66(1H, d, J=1.5Hz), 9.46, 9.06(1H, bs), 10.86, 10.43(1H, bs), 13.19(1H, bs)
789	-OMe	-NO ₂	-H	-H	2.722(1H, m), 3.089(1H, dd, J=6.4, 14.4Hz), 3.716(3H, s), 4.032(1H, m), 4.544-5.02(2H, m), 6.82(2H, d, J=8.6Hz), 7.12(2H, d, J=8.6Hz), 7.68(1H, d, J=8.4Hz), 7.91(1H, d, J=8.4, 2Hz), 7.99(1H, d, J=2Hz), 9.09, 9.48(1H, bs), 10.48, 10.91(1H, bs)

790	-OMe	-CN	-H	-H	2.68(1H,dd,J=9.2,14.3Hz),3.09(1H,dd,J=6.4,14.3Hz),3.72(3H,s),3.96(1H,dd,J=6.4,9.2Hz),4.5-5.0(2H,m),6.82(2H,d,J=8.7Hz),7.12(2H,d,J=8.7Hz),7.52(1H,d,J=8.3Hz),7.59(2H,m),9.01,9.37(1H,bs),10.38,10.86(1H,bs).
791	-H	-NO2	-H	-H	2.75(1H,dd,J=9.2,14Hz),3.17(1H,dd,J=6.2,14Hz),4.08(1H,dd,J=6.2,9.2Hz),4.5-5.0(2H,m),7.10-7.30(5H,m),7.65(1H,d,J=8.6Hz),7.90(1H,dd,J=2.2,8.6Hz),8.00(1H,d,J=2.2Hz),9.05,9.47(1H,bs),10.45,10.89(1H,bs).
792	-H	-H	-H	-COONa	2.68(1H,dd,J=8.3,14Hz),3.16(1H,dd,J=10,14Hz),3.41(1H,m),4.23(0.8H,d,J=16Hz),4.55(0.8H,d,J=17Hz),4.65(0.2H,d,J=18Hz),4.76(0.2H,d,J=17Hz),6.99(1H,d,J=8.1Hz),7.00-7.30(6H,m),7.40(1H,d,J=7.5Hz),9.02(0.6H,s),9.48(0.1H,s),10.29(0.2H,s),10.77(0.8H,s)
793	-OMe	-N(SO2Me)COO-iBu	-H	-H	0.75(6H,d,J=8.4Hz),1.81(1H,m),2.61(1H,dd,J=9.5,14.5Hz),3.07(1H,dd,J=5.9,14.5Hz),3.57(3H,s),3.71(3H,s),3.78(3H,m),4.41-4.90(2H,m),6.83(2H,d,J=8.3Hz),7.10(3H,m),7.20(1H,m),7.44(1H,d,J=8.3Hz),8.98,9.36(1H,bs),10.28,10.75(1H,bs).
794	-OMe	-SO2Me	-H	-H	2.70(1H,dd,J=9.4,14Hz),3.12(1H,dd,J=6.2,14Hz),3.24(3H,s),3.73(3H,s),3.96(1H,dd,J=6.2,9.4Hz),4.58(1.6H,br-s),4.79(0.2H,d,J=18Hz),4.97(0.2H,d,J=18Hz),6.83(2H,d,J=8.6Hz),7.13(2H,d,J=8.6Hz),7.63-7.68(2H,m),7.73(1H,dd,J=1.5,8.1Hz),9.07(0.7H,s),9.46(0.3H,s),10.41(0.3H,s),10.86(0.7H,s)
795	-H	-O-CH2CONHOH	-H	-H	2.67(1H,dd,J=9.5Hz,14.5Hz),3.17(1H,dd,J=4,14.5Hz),3.81(1H,dd,J=4.9,5Hz),4.45-4.85(4H,m),6.80(2H,m),7.30(6H,m),8.95,9.35(2H,bs),10.28,10.75(2H,bs).
796	-OMe	-H	-O-CH2OMe	-H	2.64(1H,m),3.12(1H,dd,J=5.5,14Hz),3.35(3H,s),3.71(3H,s),3.78(1H,m),4.33-4.76(2H,m),5.15(2H,s),6.82(2H,d,J=8.6Hz),6.88-7.13(5H,m),8.63,8.95(1H,bs),10.30,10.73(1H,bs).
797	-OMe	-H	-O-CH2COOH	-H	2.59(1H,dd,J=9.3,14Hz),3.11(1H,dd,J=6.14Hz),3.14(3H,s),3.78(1H,dd,J=6.9,3Hz),4.30-4.80(2H,m),4.66(2H,s),6.88(4H,m),7.04(1H,d,J=9Hz),7.12(2H,d,J=8.4Hz),9.34,8.96(1H,bs),10.73,10.30(1H,bs).

Table 15'		
Example No.	Structure	NMR: ¹ H-NMR (DMSO-d6; δ)
988		2.80(1H,m),3.14(1H,dd,J=6.3,14.1Hz),4.11(1H,m),4.39-4.90(2H,m), 7.24(2H,d,J=8.4Hz),7.34(2H,d,J=8.4Hz),7.55(1H,s),9.01,9.35(1H,bs),10.36,10.78(1H,bs),1 3.44(1H,bs).

Example 798

Step 1: Ethyl 2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-6-carboxylate

5 Ethyl 4-chloro-3-nitrobenzoate (8g) was dissolved in ethanol (100mL) and potassium carbonate (9.67g) was added. Ethanol (50mL) solution of 3-(4-methoxyphenyl)-2-mercaptopropanoic acid (7.43g) was added at 4°C, and potassium fluoride(2.0g) was added. The reaction mixture was heated under reflux for 3 hrs, and concentrated under reduced pressure. 1N-hydrochloric acid was added and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was dissolved in AcOH (100mL), and added to the suspension of Fe (6.1g) in AcOH(50mL) at 60-70°C. The reaction mixture was filtered and precipitate was washed with AcOH. Filtrate was added to water, and precipitate was collected and washed with mixed solvent of THF and hexane to give 5.9g of the product as white solid.

20 **Step 2: t-Butyl 6-ethoxycarboxyl-2H-2-(4-Methoxy-benzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetate**

To sodium hydride (3.3g) suspended in THF (30 mL) was

added a solution of ethyl 2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-6-carboxylate (3g) in THF (250 mL), and t-butyl bromoacetate (1.7 mL) was added dropwise at 0 - 5 °C. The mixture was warmed to r.t. and left at r.t.

5 overnight. Saturated aqueous ammonium chloride was added and extracted with EtOAc. The organic layer was washed with water and brine, and dried over MgSO₄ and concentrated under the reduced pressure. The residue was purified by silica gel column chromatography to give 3.9 g of the
10 product as oil.

Step 3: 6-Carboxy-2*H*-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-yl-acetic acid *N*-hydroxyamide

The product of Step 2 was converted to hydroxamic acid by
15 the method described in example 777 (step 3).

The following compounds listed in the Tables 16 and 16' were prepared in a similar manner.

Table 16




					
Example No.	RR ¹	RR ²	RR ³	RR ⁴	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
798	-OMe	-CO ₂ Et	-H	-H	1.32(3H, t, J=7.0Hz), 2.68(1H, dd, J=9.2, 14Hz), 3.11(1H, dd, J=5.9, 14Hz), 3.72(3H, s), 3.91(1H, m), 4.32(2H, q, J=7.0Hz), 4.55(1.6H, br-s), 4.70(0.2H, d, J=18Hz), 4.93(0.2H, d, J=18Hz), 6.83(2H, d, J=8.4Hz), 7.13(2H, d, J=8.4Hz), 7.51(1H, d, J=7.9Hz), 7.63(1H, dd, J=1.5, 7.9Hz), 7.66(1H, br-s), 9.06(0.8H, s), 9.46(0.2H, s), 10.42(0.2H, s), 10.86(0.8H, s)
799	-H	-H	-H	-CO ₂ Et	1.30(3H, t, J=7.2Hz), 2.69(1H, dd, J=8.4, 14Hz), 3.16(1H, dd, J=6.4, 14Hz), 3.74(1H, dd, J=6.4, 8.4Hz), 4.24-4.64(2H, m), 4.44(0.8H, d, J=16Hz), 4.52(0.8H, d, J=17Hz), 4.71(0.2H, d, J=17Hz), 4.87(0.2H, d, J=18Hz), 7.16-7.28(5H, m), 7.37-7.41(2H, m), 7.64(1H, m), 9.00(0.8H, s), 9.38(0.2H, s), 10.35(0.2H, s), 10.80(0.8H, s)
800	-H	-CO ₂ Et	-H	-H	1.31(3H, t, J=7Hz), 2.71(1H, m), 3.18(1H, m), 3.97(1H, m), 4.31(2H, dd, J=7.14, 14.10Hz), 4.51(2H, m), 7.19-7.30(5H, m), 7.51(1H, d, J=8.07Hz), 7.62(1H, d, J=8.04Hz), 7.68(1H, s), 9.05(1H, br-s), 10.84(1H, s)

Table 16'



Example No.	RR¹	RR²	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
989	-OH	-CO ₂ Et	1.32(3H, t, J=7.1Hz), 2.61(1H, dd, J=9.3Hz, 14Hz), 3.05(1H, dd, J=6.0Hz, 14Hz), 3.86(1H, dd, J=6.0Hz, 9.3Hz), 4.32(2H, q, J=7.1Hz), 4.50(1.6H, br-s), 4.69(0.2H, d, J=19Hz), 4.93(0.2H, d, J=19Hz), 6.65(2H, d, J=8.4Hz), 6.99(2H, d, J=8.4Hz), 7.52(1H, d, J=8.1Hz), 7.63(1H, dd, 1.6Hz, 8.1Hz), 7.69(1H, d, J=1.6Hz), 9.05(0.8H, s), 9.26(1H, s), 9.45(0.2H, s), 10.41(0.2H, s), 10.85(0.8H, s)
990	-Cl	-C(CH ₃) ₂ -COOEt	1.12(3H, t, J=7.2Hz), 1.48(6H, s), 2.74(1H, m), 3.18(1H, dd, m), 3.90(1H, m), 4.06(2H, q, J=7.2Hz), 4.38-4.90(2H, m), 6.83-7.15(2H, m), 7.26(2H, d, J=8.4Hz), 7.32(3H, m), 9.02, 9.42(1H, bs), 10.37, 10.81(1H, bs)
991	-OCF ₃	-CO ₂ Et	1.32(3H, t, J=7.0Hz), 2.81(1H, dd, J=8.8Hz, 14Hz), 3.19(1H, dd, J=6.2Hz, 14Hz), 4.04(1H, dd, J=6.2Hz, 8.8Hz), 4.32(2H, q, J=7.0Hz), 4.49(0.8H, d, J=17Hz), 4.55(0.8H, d, J=17Hz), 4.70(0.2H, d, J=18Hz), 4.94(0.2H, d, J=18Hz), 7.25(2H, d, J=8.0Hz), 7.36(2H, d, J=8.0Hz), 7.53(1H, d, J=8.1Hz), 7.63(1H, dd, 1.3Hz, 8.1Hz), 7.69(1H, br-s), 9.06(0.8H, s), 9.46(0.2H, s), 10.44(0.2H, s), 10.87(0.8H, s)
992	-OMe	-COO-iPr	1.31(6H, t, J=6.2Hz), 2.66(1H, dd, J=9.2Hz, 14Hz), 3.08(1H, dd, J=5.9Hz, 14Hz), 3.71(3H, s), 3.90(1H, dd, J=5.9Hz, 9.2Hz), 4.48(0.8H, d, J=17Hz), 4.53(0.8H, d, J=17Hz), 4.68(0.2H, d, J=18Hz), 4.96(0.2H, d, J=18Hz), 5.11(1H, m), 6.82(2H, d, J=8.6Hz), 7.11(2H, d, J=8.6Hz), 7.50(1H, d, J=8.1Hz), 7.60(1H, dd, 1.5Hz, 8.1Hz), 7.66(1H, d, J=1.5Hz), 9.04(0.8H, s), 9.45(0.2H, s), 10.41(0.2H, s), 10.85(0.8H, s)
993	-OMe		1.20-1.45(6H, m), 1.58(3H, d, J=5.3Hz), 1.58-1.68(2H, m), 1.75-1.88(2H, m), 2.67(1H, m), 3.12(1H, m), 3.72(3H, s), 3.92(1H, m), 4.51-4.80(2.8H, m), 4.95(0.2H, d, J=18Hz), 6.75-6.89(3H, m), 7.12(2H, d, J=8.2Hz), 7.53(1H, m), 7.61(1H, m), 7.73(1H, nr-s), 9.06(0.8H, s), 9.46(0.2H, s), 10.41(0.2H, s), 10.86(0.8H, s)

Example 801**Step 1: 6-Ethoxycarbonyl-2H-1,4-benzothiazin-3(4H)-one**

To a mixture of thioglycolic acid (10.75 g) and K_2CO_3 (65.00 g) in DMF (300 mL), a solution of ethyl 4-chloro-3-nitrobenzoate (28.12 g) in DMF (100 mL) was added and heated at 80 °C. The mixture was stirred at 80 °C for 6 hrs. The solid of K_2CO_3 was filtered off, and the filtrate was concentrated under reduced pressure. To the residue, diethyl ether (50 mL) and water (100 mL) was added and the yellow solid product was collected by filtration. The product was acidified with 4N HCl and extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 and concentrated. The product (27.634 g, 83%) was used in the following step.

To a solution of the product (12.956 g) in THF (300 mL), 10% Pd/C (13.00 g) was added. The reaction mixture was vigorously stirred at r.t. for 9 hrs under H_2 . The catalyst was removed by filtration, and the filtrate was

concentrated under reduced pressure. To a solution of the crude product (9.345 g) and HOBt (5.88 g) in DMF (200 mL), EDC·HCl (7.37 g) was added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate and washed with 1N HCl, 5% Na₂CO₃ aq., and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by recrystallization from diethyl ether and hexane to give the product (8.502 g, 79% in 2 steps) as a white solid.

Step 2: 6-Ethoxycarbonyl-2*H*-2-diethylphosphoryl-1,4-benzothiazin-3(4*H*)-one

To a solution of the 6-ethoxycarbonyl-2*H*-1,4-benzothiazin-3(4*H*)-one (8.432 g) in dichloromethane (80 mL), sulfuryl chloride (4.79 g) was added dropwise. The reaction mixture was stirred at r.t. for 6 hours. The solvent was removed under reduced pressure. The residue was purified by recrystallization from chloroform and hexane to give the product (8.80 g, 91%) as a white solid.

A mixture of 6-ethoxycarbonyl-2*H*-2-chloro-1,4-benzothiazin-3(4*H*)-one (8.724 g) and triethyl phosphite (11.74 g) was heated at 120 °C and stirred for 10 hrs. The solvent was removed under reduced pressure. The residue was purified by recrystallization from THF and diethyl ether to give the product (10.534 g, 88%) as a pale yellow solid.

Step 3: *t*-Butyl 6-ethoxycarbonyl-2*H*-2-(4-phenoxybenzyl)-

1,4-benzothiazin-3(4H)-one-4-ylacetate

To a stirred solution of *p*-phenoxybenzaldehyde (796 mg) and 6-ethoxycarbonyl 2*H*-2-diethylphosphoryl-1,4-benzothiazine-3(4*H*)-one (1.00 g) in 80 % ethanol aq. (10 mL) and THF (10 mL), a solution of sodium ethoxide (400 mg) in ethanol (5 mL) was added dropwise. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure. To the residue, ethyl acetate(5 ml) and water(10 ml) was added and the solid product was collected by filtration. After drying under reduced pressure, the product(980 mg, 88%) was given as a yellow solid.

To a solution of 6-ethoxycarbonyl-2*H*-2-(4-phenoxybenzylidene)-1,4-benzothiazin-3(4*H*)-one (980 mg) was dissolved in methanol (40 mL) and THF (40 mL), 10% Pd/C (1.00 g) was added. The reaction mixture was vigorously stirred at r.t. for 4 hrs under H₂. The catalyst was removed by filtration, and the filtrate was concentrated under reduced pressure. The product (936 mg, 95%) was used in the following step without purification.

To a solution of 6-ethoxycarbonyl-2*H*-2-(4-phenoxybenzyl) -1,4-benzothiazin-3(4*H*)-one (925 mg) in DMF (15 mL), potassium *t*-butoxide (495 mg) was added at r.t.. After 20 minutes' stirring, *t*-buthyl bromoacetate (645 mg) was slowly added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure and the residue was dissolved in ethyl acetate and washed

with water. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/ethyl acetate, 30/1) to give the product (677 mg, 58%).

5

Step 4: 6-Ethoxycarbonyl-2*H*-2-(4-phenoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid *N*-*t*-butoxyamide

t-Butyl 6-ethoxycarbonyl-2*H*-2-(4-phenoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetate (677 mg) was dissolved in TFA (10 mL). The mixture was stirred r.t. for 2 hrs. The solvent was removed under reduced pressure. The residue was purified with silica gel column chromatography (eluent: chloroform/methanol, 10/1) to give the product (428 mg, 71%).

10

Isobutyl chloroformate (133 mg) was added dropwise to a THF (10 mL) solution of 6-ethoxycarbonyl-2*H*-2-(4-phethoxybenzylidene)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid (422 mg) and *N*-methylmorpholine (98 mg) at -5 °C. A solution of *O*-*t*-butyl-hydroxylamine hydrochloride (133 mg) and *N*-methylmorpholine (107 mg) in THF (2 mL) was added dropwise after additional 20 minutes' stirring. The mixture was warmed to r.t. with stirring, and stood overnight. The reaction mixture was concentrated under reduced pressure, acidified with 0.5N citric acid, and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (eluent: chloroform/methanol, 5/1) to give the product (260 mg, 54%).

15

20

25

Step 5: 6-carboxy-2*H*-2-(4-phenoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-ylacetic acid N-hydroxyamide

To a solution of 6-ethoxycarbonyl-2*H*-2-(4-phenoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-*t*-butoxyamide (256 mg) in ethanol (5 mL) and THF (5 mL), 1N NaOH (560 μ L) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to room temperature and left overnight. The reaction mixture was concentrated under reduced pressure and acidified with 1N citric acid, and the product was extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated. The crude product (241 mg) was used in the following step without purification.

6-Carboxy-2*H*-2-(4-phenoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-*t*-butoxyamide (124 mg) was dissolved in TFA (10 mL). The mixture was stirred at r.t. for 2 days. The solvent was removed under reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel (eluent: chloroform/methanol, 5/1) to give the hydroxamic acid (99 mg, 46% in 2 steps).

The following compounds listed in the Tables 17, 17', and 17'' were prepared in a similar manner.


Table 17			
Example No.	RR¹	NMR: ¹H-NMR (DMSO-d6; δ)	
801	-OPh	2.75(1H, m), 3.16(1H, m), 3.96(1H, m), 4.51(2H, m), 6.90(2H, d, J=8.25Hz), 6.99(2H, d, J=8.61Hz), 7.15(1H, m), 7.24(2H, d, J=8.43Hz), 7.36-7.42(2H, m), 7.49(1H, d, J=8.04Hz), 7.60-7.66(2H, m), 9.06(1H, br-s), 10.84(1H, s)	
802	-O-Pentyl	0.89(3H, t, J=7Hz), 1.36(4H, m), 1.68(2H, m), 2.68(1H, m), 3.13(1H, m), 3.90- 3.94(3H, m), 4.51(2H, m), 6.82(2H, d, J=8.43Hz), 7.11(2H, d, J=8.40Hz), 7.49(1H, d, J=8.04Hz), 7.61(1H, d, J=8.04Hz), 7.66(1H, s), 9.05(1H, s), 10.83(1H, s)	
803	-F	2.77(1H, m), 3.17(1H, m), 4.00(1H, m), 4.51(2H, m), 7.10(2H, t, J=8.79Hz), 7.27(2H, m), 7.50(1H, d, J=8.07Hz), 7.62(1H, d, J=8.07Hz), 7.67(1H, s), 9.05(1H, s), 10.84(1H, s)	




Table 17'			
Example No.	RR ¹	NMR: ¹ H-NMR (CDCl ₃ ; δ)	
994	-C6H3-3,4-methylenedioxy	1.41(t, J=7.2Hz, 3H), 2.81(m, 1H), 3.14(m, 1H), 3.70(m, 1H), 4.40(q, J=7.2Hz, 2H), 4.48(d, J=16Hz, 1H), 4.73(d, J=16Hz, 1H), 5.97(s, 2H), 6.49(dd, J=2.4, 8.4Hz, 1H), 6.57(d, J=2.4Hz, 1H), 6.76(d, J=8.4Hz, 1H), 6.88(m, 2H), 7.05(m, 2H), 7.44(d, J=8.0Hz, 1H), 7.78(m, 1H), 8.10(m, 1H), 9.00(br, 1H)	
995	-C6H4-3-F	1.41(t, J=6.9Hz, 3H), 2.84(m, 1H), 3.18(m, 1H), 3.73(m, 1H), 4.40(q, J=6.9Hz, 2H), 4.50(d, J=15.6Hz, 1H), 4.73(d, J=15.6Hz, 1H), 6.70(m, 1H), 6.76-6.82(m, 2H), 6.96 (m, 2H), 7.11 (m, 2H), 7.28(m, 1H), 7.45(d, J=8.0Hz, 1H), 7.78(m, 1H), 8.09(br, 1H), 9.10 (br, 1H)	
996	-C6H4-3-OMe	1.41(t, J=7.2Hz, 3H), 2.81(m, 1H), 3.18(m, 1H), 3.71(m, 1H), 3.78(s, 3H), 4.39(q, J=7.2Hz, 2H), 4.51(d, J=16Hz, 1H), 4.72(d, J=16Hz, 1H), 6.56-6.60(m, 2H), 6.65(m, 1H), 6.94(m, 2H), 7.08(m, 2H), 7.22(m, 1H), 7.43(d, J=8.0Hz, 1H), 7.49(br, 1H), 7.76(d, J=8.0Hz, 1H), 8.06(br, 1H), 9.15(br, 1H)	
997	-C6H4-4-SO ₂ Me	1.41(t, J=7.2Hz, 3H), 2.88(m, 1H), 3.06(m, 3H), 3.20(m, 1H), 3.75(m, 1H), 4.40(q, J=7.2Hz, 2H), 4.50(d, J=16Hz, 1H), 4.74(d, J=16Hz, 1H), 7.00(m, 2H), 7.08(m, 1H), 7.18(m, 2H), 7.45(d, J=8.0Hz, 1H), 7.78(d, J=8.0Hz, 1H), 7.89(m, 2H), 8.09(m, 1H), 9.03(br, 1H)	

Table 17"



Example No.	RR ¹	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
998	-O-4-pyridyl	2.82(1H,dd,J=14.2, 8.8Hz,benzyllic C-H), 3.22(1H,dd,J=14.2, 6.1Hz,another benzyllic C-H), 4.02(1H,dd,J=8.8, 6.1Hz,H-C-S), 4.52(2H,s,CH ₂ CO), 6.88 & 8.45(2H+2H,dd,J=4.7, 1.5, 3.5- & 2.6-H's of pyridine), 7.08 & 7.34(2H+2H,d,J=8.5), 7.51(1H,d,J=8.0), 7.63(1H,dd,J=8.0, 1.4), 7.68(1H,d,J=1.4), 9.06 & 10.86(1H+1H,s,NH-OH), 13.2(1H, br s,CO ₂ H)
999	-O-C ₆ H ₃ -3,4-methylenedioxy	2.72(m,1H),3.14(m,1H),3.95(m,1H),4.51+4.69+4.92(2H,NCH ₂ CO),6.03(s,2H),6.47(dd,J=2.4,8.2Hz,1H),6.71(d,J=2.4Hz,1H),6.83(m,2H),6.91(d,J=8.2Hz,1H),7.20(m,2H),7.49(d,J=8.0Hz,1H),7.61(dd,J=1.6,8.0Hz,1H),7.65(d,J=1.6Hz,1H),9.05+9.45(s,1H),10.40+10.84(s,1H) 13.14(br,1H)
1000	-O-C ₆ H ₄ -3-CF ₃	2.78(m,1H),3.19(m,1H),4.00(m,1H),4.52+4.71+4.92(2H,NCH ₂ CO),7.05(m,2H),7.23-7.33(m,4H),7.43-7.53(m,2H),7.59-7.64(m,2H),7.67(m,1H), 9.05+9.46(s,1H),10.41+10.85(s,1H),13.17(br,1H)
1001	-O-C ₆ H ₄ -3-F	2.77(m,1H),3.18(m,1H),3.99(m,1H),4.15+4.70+4.92(2H,NCH ₂ CO),6.79(m,1H),6.84(m,1H),6.94-7.01(m,3H),7.27(m,2H),7.41(m,1H), 7.51 (d,J=8.0Hz,1H), 7.62(dd,J=1.6,8.0Hz,1H),7.66(d,J=1.6Hz,1H), 9.05+ 9.46 (s,1H),10.41+10.84(s,1H),13.14(br,1H)
1002	-SO ₂ -C ₆ H ₄ -4-OMe	2.84(m,1H),3.23(m,1H),3.82(s,3H),4.06(m,1H),4.49+4.78+4.90(2H,NCH ₂ CO),7.10-7.14(m,2H),7.46-7.51(m,3H),7.59(m,1H),7.66(m,1H), 7.81-7.90(m,4H), 9.06+9.46 (s,1H),10.42+10.84(s,1H),13.10(br,1H)
1003	-O-C ₆ H ₄ -3-OMe	2.75(m,1H),3.16(m,1H),3.74(s,3H),3.98(m,1H),4.52+4.70+4.92(2H,NCH ₂ CO),6.52(m,1H),6.57(m,1H),6.72(m,1H),6.92(m,2H),7.22-7.31(m,3H), 7.51 (d,J=8Hz,1H),7.62(dd,J=4.8Hz,1H),7.66(m,1H),9.05+9.45 (s,1H),10.41 +10.85(s,1H),13.18(br,1H)

1004	-O-C6H4-4-SO2Me	2.81(m, 1H), 3.19(s, 3H), 3.40(m, 1H), 4.02(m, 1H), 4.52+4.73+4.95(2H, NCH2CO), 7.06(m, 2H), 7.13(m, 2H), 7.33(m, 2H), 7.51(d, J=8Hz, 1H), 7.62(dd, J=1.6, 8Hz, 1H), 7.67(d, J=1.6Hz, 1H), 7.91(m, 2H), 9.05+9.46 (s, 1H), 10.41+10.85(s, 1H), 13.19(br, 1H)
------	-----------------	--

Example 804

Step 1: Ethyl 6-carboxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

Ethyl 6-t-butoxycarbonyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate (32.9 g) was dissolved in 5%-ethanedithiol in TFA (100mL) and stood at r.t. for 1.5hrs. Toluene (100mL) was added and concentrated under reduced pressure. The residue was washed with mixed solvent of ether and hexane to give 30.2 g of ethyl 6-carboxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate as pale yellow solid.

Step 2: Ethyl 6-isopropylcarbamoyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

Ethyl 6-carboxyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate (2 g) was dissolved in dichloromethane (20mL), and oxalyl chloride (1.22 g) was added at 4°C. 1 drop of DMF was added and warmed to r.t., and stirred for 3hrs. Toluene (20mL) was added and concentrated under reduced pressure. This acid chloride was dissolved in


dichloromethane (15mL), and added to dichloromethane (10mL) solution of diisopropylamine (299mg) and triethylamine (511mg) at 4°C. The reaction mixture was stirred for 4hrs at r.t., diluted with 5% KHSO₄ aq., and extracted with ethyl acetate. The organic layer was washed with sat. NaHCO₃ aq. and brine, and dried over MgSO₄. The solution was concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 1.47g of ethyl 6-isopropylcarbamoyl-2*H*-2-(4-methoxy-benzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetate as pale yellow oil.

Step 3: 6-Isopropylcarbamoyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-hydroxy-amide

The product of Step 2 was converted to hydroxamic acid by the method described in example 777 (step 3).

The following compounds listed in the Table 18 were prepared in a similar manner.



Table 18






Example No.	RR ¹	RR ²	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
804	-OMe	-NHPr	1.17(6H, d, J=6.6Hz), 2.65(1H, dd, J=9.4, 14Hz), 3.10(1H, dd, J=6.1, 14Hz), 3.73(3H, s), 3.85(1H, dd, J=6.1, 9.4Hz), 4.09(1H, m), 4.53(1.6H, br-s), 4.70(0.2H, d, J=18Hz), 4.93(0.2H, d, J=18Hz), 6.83(2H, d, J=8.6Hz), 7.12(2H, d, J=8.3Hz), 7.44(1H, d, J=8.1Hz), 7.53-7.56(2H, m), 8.27(1H, d, J=7.7Hz), 9.02(0.8H, s), 9.44(0.2H, s), 10.34(0.2H, s), 10.80(0.8H, s)
805	-H		1.88(3H, m), 2.27(1H, m), 2.74(1H, m), 3.19(1H, m), 3.56(2H, m), 3.67(3H, s), 3.94(1H, m), 4.49(3H, m), 7.24-7.29(7H, m), 7.46(1H, d, J=8.04Hz), 9.04(1H, s), 10.82(1H, s)
806	-H		1.91(3H, m), 2.26(1H, m), 2.74(1H, m), 3.21(1H, m), 3.53(2H, m), 3.94(1H, m), 4.39(1H, m), 4.50(2H, m), 7.23-7.28(7H, m), 7.45(1H, d, J=7.68Hz), 9.00(1H, br-s), 10.81(1H, s)
807	-H	-Piperidino-4-COO-tBu	1.41(9H, s), 2.74(1H, m), 3.23(1H, m), 3.31-3.39(8H, m), 3.94(1H, m), 4.51(2H, m), 7.09-7.29(7H, m), 7.44(1H, d, J=8.04Hz), 9.02(1H, s), 10.79(1H, s)
808	-H	-Piperidino	2.74(1H, m), 3.15-3.26(5H, m), 3.68(4H, m), 3.94(1H, m), 4.53(2H, m), 7.14-7.29(7H, m), 7.46(1H, d, J=7.86Hz), 9.03(1H, s), 10.87(1H, s)
809	-H	-NH ₂	2.66(1H, dd, J=9.5, 14Hz), 3.14(1H, dd, J=5.7, 14Hz), 3.88(1H, dd, J=5.7, 9.5Hz), 4.48(1.6H, br-s), 4.68(0.2H, d, J=18Hz), 4.87(0.2H, d, J=18Hz), 7.14-7.25(5H, m), 7.25-7.41(2H, m), 7.50(1H, d, J=7.9Hz), 7.56(1H, br-s), 8.00(1H, br-s), 8.06(0.8H, s), 9.37(0.2H, s), 10.29(0.2H, s), 10.72(0.8H, s)
810	-OMe	-NH ₂	2.63(1H, dd, J=9.3, 14Hz), 3.12(1H, dd, J=5.9, 14Hz), 3.72(3H, s), 3.86(1H, dd, J=5.9, 9.3Hz), 4.53(1.4H, br-s), 4.72(0.3H, d, J=17Hz), 4.93(0.3H, d, J=17Hz), 6.84(2H, d, J=8.6Hz), 7.12(2H, d, J=8.6Hz), 7.42-7.45(2H, m), 10.72(0.8H, s)

811	-H	-NH-(S)-CH(Me)COOH	7.54-7.61(2H, m), 8.05(1H, br-s), 9.01(0.7H, s), 9.42(0.3H, s), 10.34(0.3H, s), 10.78(0.7H, s)
812	-OMe	-(S)-NHCH(Me)COO-t-Bu	1.40(3H, d, J=7.3Hz), 2.70(1H, m), 3.18(1H, m), 3.93(1H, m), 4.41(1H, m), 4.52(0.8H, d, J=18Hz), 4.57(0.8H, d, J=18Hz), 4.73(0.2H, dd, J=18Hz), 4.94(0.2H, dd, J=18Hz), 7.20-7.25(3H, m), 7.25-7.31(2H, m), 7.46-7.64(3H, m), 8.76(1H, dd, J=1.8Hz, 7.3Hz), 9.04(0.8H, s), 9.45(0.2H, s), 10.36(0.2H, s), 10.80(0.8H, s), 12.6(1H, s)
813	-OMe	-NH-(S)-CH(Me)COOH	1.38(3H, d, J=8.1Hz), 1.42(9H, s), 2.61(1H, m), 3.10(1H, m), 3.72(3H, s), 3.87(1H, dd, J=6.2, 9.2Hz), 4.34(1H, m), 4.53(1.6H, br-s), 4.69(0.2H, m), 4.94(0.2H, dd, J=18Hz), 6.84(2H, d, J=8.3Hz), 7.12(2H, d, J=8.3Hz), 7.43-7.59(3H, m), 8.73(1H, d, J=6.4Hz), 9.02(0.8H, s), 9.44(0.2H, s), 10.34(0.2H, s), 10.79(0.8H, s)
814	-OMe	-NH-cyclopentyl	1.40(3H, d, J=7.3Hz), 2.58-2.69(1H, m), 3.04-3.14(1H, m), 3.72(3H, s), 3.87(1H, dd, J=6.1, 9.1Hz), 4.41(1H, m), 4.53(1.6H, br-s), 4.69(0.2H, dd, J=13, 18Hz), 4.94(0.2H, dd, J=13, 18Hz), 6.84(2H, d, J=8.6Hz), 7.12(2H, dd, J=1.9, 8.3Hz), 7.45-7.60(3H, m), 8.74(1H, d, J=7.1Hz), 9.03(0.8H, s), 9.44(0.2H, s), 10.36(0.2H, s), 10.80(0.8H, s), 12.58(1H, s)
815	-OMe	-NH-cyclopropyl	1.53(4H, br-s), 1.70(2H, br-s), 1.89(2H, br-s), 2.65(1H, dd, J=9.4, 14Hz), 3.10(1H, dd, J=5.9, 14Hz), 3.73(3H, s), 3.85(1H, dd, J=5.9, 9.4Hz), 4.21(1H, m), 4.53(1.6H, br-s), 4.71(0.2H, d, J=18Hz), 4.94(0.2H, d, J=18Hz), 7.12(2H, d, J=8.4Hz), 7.40-7.55(3H, m), 8.33(1H, d, J=7.3Hz), 9.03(0.8H, s), 9.46(0.2H, s), 10.36(0.2H, s), 10.81(0.8H, s)
804	-H	-NHEt	0.53-0.59(2H, m), 0.68-0.74(2H, m), 2.65(1H, dd, J=9.4, 14Hz), 2.71(1H, m), 3.10(1H, dd, J=5.7, 14Hz), 3.72(3H, s), 3.86(1H, dd, J=5.7, 9.4Hz), 4.52(1.6H, br-s), 4.71(0.2H, d, J=17Hz), 4.92(0.2H, d, J=17Hz), 6.83(2H, d, J=8.6Hz), 7.12(2H, d, J=8.6Hz), 7.40-7.54(3H, m), 8.50(1H, d, J=3.9Hz), 9.02(0.8H, s), 9.45(0.2H, s), 10.36(0.2H, s), 10.79(0.8H, s)
817	-OMe	-NHC(Me)2CH2OH	1.12(3H, t, J=7.14Hz), 2.71(1H, m), 3.15-3.33(3H, m), 3.93(1H, m), 4.53(2H, m), 7.20-7.31(5H, m), 7.44-7.58(3H, m), 8.55(1H, m), 9.03(1H, s), 10.80(1H, s)
818	-OMe	-(S)-NHCH(iPr)CH2OH	1.31(6H, s), 2.65(1H, dd, J=9.4, 14Hz), 3.09(1H, dd, J=5.7, 14Hz), 3.73(3H, s), 3.74-3.90(3H, m), 4.53(1.4H, br-s), 4.67(0.3H, d, J=18Hz), 4.91(0.3H, d, J=18Hz), 6.84(2H, d, J=8.6Hz), 7.11(2H, d, J=8.6Hz), 7.43(1H, d, J=8.4Hz), 7.53(1H, d, J=8.4Hz), 7.61(1H, br-s), 9.03(0.7H, s), 9.45(0.3H, s), 10.36(0.3H, s), 10.81(0.7H, s)
			0.87-0.96(6H, m), 1.57(1H, m), 2.65(1H, m), 3.08(1H, m), 3.52(2H, br-s), 3.72(3H, s), 3.81(1H, m), 4.54(1.6H, br-s), 4.74(0.2H, dd, J=6.8, 18Hz), 4.95(0.2H, dd, J=12, 18Hz), 6.84(2H, d, J=7.8Hz), 6.87(1H, s),

819	-OMe	-NHMe	7.13(2H, m), 7.45(1H, d, J=7.7Hz), 7.59(1H, br-s), 7.60(1H, d, J=8.4Hz), 8.07(1H, d, J=8.7Hz), 9.03(0.8H, s), 9.49(0.2H, s), 10.37(0.2H, s), 10.81(0.8H, s)
820	-Cl	-NH-cyclopropyl	2.64(1H, dd, J=9.14Hz), 2.77(3H, d, J=4.4Hz), 3.09(1H, dd, J=5.8, 14Hz), 3.71(3H, s), 3.86(1H, dd, J=5.9Hz), 4.50-4.93(2H, m), 6.82(2H, d, J=8.6Hz), 7.11(2H, J=8.6Hz), 7.46(2H, m), 7.55(1H, s), 8.50(1H, m), 9.44, 9.01(1H, bs), 10.78, 10.34(1H, bs).
821	-Cl	-NHCH ₂ -2-furyl	0.51-0.56(2H, m), 0.66-0.73(2H, m), 2.72(1H, dd, J=14.3Hz, 9.3Hz), 2.76-2.84(1H, m), 3.12(1H, dd, J=14.4Hz, 6.4Hz), 3.93(1H, dd, J=8.9Hz, 6.4Hz), [4.50(S), 4.62(d, J=17.7Hz), 4.84(d, J=17.7Hz)] 2H, 7.23(2H, d, J=8.4Hz), 7.32(2H, d, J=8.4Hz), 7.43(1H, d, J=8.1Hz), 7.48(1H, dd, J=8.2Hz, 1.8Hz), 7.53(1H, d, J=1.8Hz), 8.49(1H, d, J=4.1Hz), [9.01(br), 9.43(s)] 1H, [10.78(br), 10.35(s)] 1H.
822	-Cl	-NHCH ₂ CF ₃	2.73(1H, dd, J=14.4, 8.8Hz), 3.14(1H, dd, J=14.3Hz, 6.6Hz), 3.94(1H, dd, J=8.9Hz, 6.4Hz), 4.46(2H, d, J=5.4Hz), [4.51(S), 4.72(d, J=17.9Hz), 4.95(d, J=3.1Hz, 0.5Hz), 6.39(1H, dd, J=3.1Hz, 1.8Hz), 7.23(2H, d, J=8.4Hz), 7.32(2H, d, J=8.4Hz), 7.45(1H, d, J=7.9Hz), 7.55(1H, dd, J=8.9Hz, 1.5Hz), 7.57(1H, d, J=1.8Hz), 7.60(1H, m), [9.01(br), 9.42(s)] 1H, 9.04(1H, t, J=5.4Hz), [10.78(br), 10.35(s)] 1H.
823	-Cl	-NH(CH ₂) ₂ OH	2.74(dd, J=14.4, 9.1Hz, 1H), 3.12(dd, J=14.3Hz, 6.6Hz, 1H), 3.96(dd, J=8.9Hz, 6.4Hz, 1H), 4.10(dd, J=10.5Hz, 6.4Hz, 2H), [4.52(S), 4.72(d, J=17.9Hz), 4.95(d, J=17.9Hz)] 2H, 7.23(d, J=8.4Hz, 2H), 7.32(d, J=8.2Hz, 2H), 7.49(d, J=8.0Hz, 1H), 7.57(dd, J=8.0Hz, 1.0Hz, 1H), 7.63(d, J=1.0Hz, 1H), [9.01(br), 9.44(s)] 1H, 9.15(t, J=6.4Hz, 1H), [10.79(br), 10.36(s)] 1H.
824	-Cl	-NH ₂	2.72(dd, J=14.1, 9.1Hz, 1H), 3.12(dd, J=13.9Hz, 6.2Hz, 1H), 3.46-3.50(m, 2H), 3.28-3.32(m, 2H), 3.94(dd, J=8.7Hz, 6.2Hz, 1H), [4.51(S), 4.72(d, J=17.9Hz), 4.95(d, J=17.9Hz)] 2H, 4.88(br, 1H), 7.28(d, J=8.4Hz, 2H), 7.32(d, J=8.4Hz, 2H), 7.44(d, J=8.0Hz, 1H), 7.53(dd, J=8.0Hz, 1.4Hz, 1H), 7.58(d, J=1.0Hz, 1H), 8.54(t, J=5.6Hz, 1H), [9.00(br), 9.42(s)] 1H., [10.78(br), 10.34(s)] 1H.
824	-Cl	-NH ₂	2.73(dd, J=13.9, 8.9Hz, 1H), 3.13(dd, J=14.1Hz, 6.1Hz, 1H), 3.94(dd, J=8.9Hz, 6.2Hz, 1H), [4.51(S), 4.72(d, J=17.9Hz), 4.90(d, J=17.9Hz)] 2H, 7.23(d, J=8.4Hz, 2H), 7.32(d, J=8.4Hz, 2H), 7.44(d, J=8.0Hz, 1H), 7.54(dd, J=8.0Hz, 1.5Hz, 1H), 7.59(d, J=1.0Hz, 1H), 8.04(br, 2H), [8.99(br), 9.41(s)] 1H, [10.76(br), 10.34(s)] 1H.

825	-Cl	-NH(CH ₂) ₂ NEt ₂	1.21 (t, J=7.1Hz, 6H), 2.74 (dd, J=14.1, 9.1 Hz, 1H), 3.13 (dd, J=14.6Hz, 6.7Hz, 1H), 3.17-3.27 (m, 6H), 3.50-3.70 (m, 2H), 3.97 (dd, J=8.7Hz, 6.4Hz, 1H), [4.51 (S), 4.69(d, J=17.3Hz), 4.93 (d, J=17.3Hz)] 2H, 7.24 (d, J=8.4Hz, 2H), 7.34 (d, J=8.3Hz, 2H), 7.51 (d, J=8.0Hz, 1H), 7.55 (d, J=8.0Hz, 1H), 7.61 (s, 1H), 8.84 (t, J=5.1Hz, 1H), 9.39 (br, 1H), [10.84 (br), 10.37 (s)] 1H.
826	-Cl		1.11 (t, J=7.1Hz, 3H), 1.50-2.20 (m, 2H), 2.73 (dd, J=14.1, 8.8 Hz, 1H), 3.06-3.17 (m, 2H), 3.45-3.52 (m, 1H), 3.54-3.64(m,2H),4.10-4.40(m,5H),3.95 (t, J=6.4Hz, 1H), [4.49 (S), 4.72(d, J=17.3Hz), 4.92 (d, J=17.3Hz)] 2H, 7.23 (d, J=8.2Hz, 2H), 7.32 (d, J=8.4Hz, 2H), 7.51 (d, J=8.0Hz, 1H), 7.55 (d, J=8.0Hz, 1H), 7.60 (s, 1H), 8.90(br, 1H), 9.20(br,1H), [10.82 (br), 10.36 (s)] 1H.
827	-Cl		1.59 (s, 6H), 2.73 (dd, J=14.1, 8.8 Hz, 1H), 3.11 (m, 2H), 3.92 (t, J=6.3Hz, 1H), [4.51 (S), 4.72(d, J=17.3Hz), 4.92 (d, J=17.3Hz)] 2H, 7.23 (d, J=8.2Hz, 2H), 7.32 (d, J=8.4Hz, 2H), 7.44 (d, J=7.8Hz,2H), 7.53 (s, 1H), 7.54(d, J=8.0Hz, 1H), 8.30(br, 1H), 10.79 (br, 1H).
828	-Cl	-NH(CH ₂) ₃ NMe ₂	1.80-1.90 (m, 2H), 2.73 (dd, J=14.1, 8.8 Hz, 1H), 2.77 (s, 6H), 3.05-3.10 (m, 2H), 3.12 (dd, J=13.9Hz, 6.4Hz, 1H), 3.27-3.36, (m, 2H), 3.94 (dd, J=8.8Hz, 6.4Hz, 1H), [4.50 (S), 4.68(d, J=17.2Hz), 4.92 (d, J=17.4Hz)] 2H, 7.23 (d, J=8.2Hz, 2H), 7.32 (d, J=8.4Hz, 2H), 7.50 (d, J=8.0Hz, 1H), 7.53 (d, J=8.0Hz, 1H), 7.58 (s, 1H), 8.70 (t, J=5.6 Hz, 1H), 9.46 (br, 1H), [10.82 (br), 10.35(s)] 1H.
829	-Cl	-NH(CH ₂) ₃ -1-imidazolyl	2.04-2.10 (m, 2H), 2.73 (dd, J=14.1, 8.8 Hz, 1H),3.12 (dd, J=13.9Hz, 6.3Hz, 1H), 3.26-3.31, (m, 2H), 3.93 (dd, J=8.8Hz, 6.4Hz, 1H), 4.21-4.26(m,2H), [4.50 (S), 4.68(d, J=17.3Hz), 4.91 (d, J=17.3Hz)] 2H, 7.23 (d, J=8.2Hz, 2H), 7.32 (d, J=8.4Hz, 2H), 7.47 (d, J=8.0Hz, 1H), 7.55 (d, J=8.0Hz, 1H), 7.57 (s, 1H), 7.69 (dd, J=1.65Hz, J=1.65Hz, 1H), 7.81 (dd, J=1.65Hz, J=1.65Hz, 1H), 8.64 (t, J=5Hz, 1H), 9.13 (s, 1H), [10.81 (br), 10.35(s)] 1H.
830	-Cl	-NH-(S)-CH(Me)COOH	1.38 (d, J=7.5Hz, 3H), 2.66-2.75 (m, 1H), 3.04-3.18 (m, 1H),, 3.95 (dd, J=8.8Hz, 6.6Hz, 1H), 4.40 (dd, J=6.9Hz, 6.4Hz, 1H), [4.52 (S), 4.66-4.84 (m), 4.88-4.96 (m)] 2H, 7.23 (d, J=8.2Hz, 2H), 7.32 (d, J=8.3Hz, 2H), 7.47 (d, J=8.4Hz, 1H), 7.58-7.63 (m 2H), 8.75 (d, J=6.4Hz, 1H), 9.45 (br,1H), [10.78 (br), 10.35(s)] 1H.
831	-Cl	-NH-(R)-CH(Me)COOH	1.38 (d, J=7.3Hz, 3H), 2.72 (dd, J=14.6Hz, 8.4Hz, 1H), 3.06-3.18 (m, 1H), 3.95 (dd, J=8.9Hz, 6.4Hz, 1H), 4.36-4.45 (m, 1H), [4.52 (S), 4.66-4.84 (m), 4.88-4.96 (m)] 2H, 7.23 (d, J=8.2Hz, 2H),

832	-Cl		7.32 (d, J=8.3Hz, 2H), 7.47 (d, J=8.2Hz, 1H), 7.58-7.63 (m 2H), 8.75 (d, J=6.4Hz, 1H), 9.44 (br, 1H), [10.79 (br), 10.35(s)] 1H. 2.30-2.46 (m, 2H), 2.70 (dd, J=13.9Hz, 9.3Hz, 1H), 3.08 (dd, J=13.7Hz, 6.2Hz, 1H), 3.95 (dd, J=8.9Hz, 6.4Hz, 1H), 4.23 (dd, J=16Hz, 9.8Hz, 1H), 4.39 (dd, J=17Hz, 9.9Hz, 1H), [4.47 (s), 4.62-4.73 (m), 4.85-4.93 (m)]2H, 4.65-4.71 (m, 1H), 7.19 (d, J=8.2Hz, 2H), 7.28 (d, J=8.2Hz, 2H), 7.48 (d, J=7.9Hz, 1H), 7.50 (d, J=8.6Hz, 1H), 7.56 (s, 1H), 9.03-9.06 (m, 2H), 9.40 (br, 1H), [10.75 (br), 10.32(s)] 1H.
833	-Cl		0.938 (d, J=3.6Hz, 2H), 1.31 (d, J=3.6Hz, 2H), 2.71 (dd, J=13.9Hz, 8.2Hz, 1H), 3.06-3.16 (m, 1H), 3.87-3.96 (m, 1H), [4.44 (d, J=16Hz), 4.53 (d, J=16Hz), 4.60-4.73 (m), 4.88-4.97 (m)] 2H, 7.23 (d, J=8.4Hz, 2H), 7.32 (d, J=8.3Hz, 2H), 7.42 (d, J=7Hz, 1H), 7.47 (d, J=7Hz, 1H), 7.60 (s, 1H), 8.88 (s, 1H), 8.89 (br, 1H), 9.44 (br, 1H), [10.77 (br), 10.34(br)] 1H, [10.62 (br), 11.28 (br)] 1H.

Example 835

Step 1: Methyl 6-hydroxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

Ethyl chloroformate (3.475 g) was added dropwise to a THF (30 mL) solution of methyl 6-carboxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate (2.795 g) and triethylamine (3.241 g) at 0 °C. A solution of sodium borohydride (3.160 g) in water (5 mL) was added dropwise after additional 1 hour's stirring. The reaction mixture was stirred at 0 °C for 1 hr. The reaction mixture was quenched with acetic acid and concentrated under reduced pressure. The residue was dissolved in ethyl acetate and washed with water. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 4/1) to give the product (2.294 g, 79%).

Step 2: Methyl 6-t-butyldimethylsiloxy-methyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

To a solution of methyl 6-hydroxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

(2.294 g) and imidazole(1.612 g) in DMF (30 mL), a solution of TBDMSCl (2.665 g) in DMF (5 ml) was slowly added. The reaction mixture was stirred at r.t. for 2 days and quenched with water. The solvent was removed under reduced pressure, and the residue was dissolved in chloroform, dried over Na₂SO₄, and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 50/1) to give the product(2.966 g, quant.).


Step 3: 6-Hydroxymethyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-hydroxyamide


To a solution of the methyl ester (2.966 g) in methanol (20 mL), 1N NaOH (8.86 mL) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to room temperature and left overnight. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in ethyl acetate and water. After the pH was adjusted to 6 with acetic acid, the product was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated to give the product (1.208 g, 42%) as a white solid.

Isobutyl chloroformate (373 mg) was added dropwise to a THF (10 mL) solution of 6-*t*-butyldimethylsiloxymethyl -2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid (1.208g) and N-methylmorpholine (276 mg) at -5 °C. THF (1 mL) solution of *O*-TMS-hydroxylamine (313 mg) was added after additional 20 minutes' stirring. The mixture was

stirred at -5 °C for 2 hrs , gradually warmed to r.t. with stirring, and stood overnight. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was washed with 1N HCl and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (eluent: chloroform/methanol, 10/1) to give the hydroxamic acid(645 mg, 70%) as a white powder.

The following compounds listed in the Table 19 were prepared in a similar manner.

Table 19					NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
Example No.	RR ¹	RR ²	RR ³		
835	-OMe	-CH ₂ OH	-H		2.63(1H, m), 3.10(1H, m), 3.72(3H, s), 3.77(1H, m), 4.45-4.49(4H, m), 5.28(1H, m), 6.84(2H, d, J=8.43Hz), 7.02(1H, d, J=8.07Hz), 7.10-7.13(3H, m), 7.31(1H, d, J=7.89Hz), 8.99(1H, s), 10.76(1H, s)
836	-OMe	-H	-CH ₂ OH		2.65(1H, m), 3.10(1H, m), 3.72(3H, s), 3.78(1H, m), 4.42-4.45(4H, m), 5.21(1H, m), 6.84(2H, d, J=8.25Hz), 7.07-7.14(3H, m), 7.23(1H, d, J=8.43Hz), 7.28(1H, s), 8.97(1H, s), 10.75(1H, s)
837	-OPh	-H	-CH ₂ OH		2.71(1H, m), 3.16(1H, m), 3.85(1H, m), 4.43-4.46(4H, m), 6.91(2H, d, J=8.61Hz), 7.00(2H, d, J=7.71Hz), 7.07-7.16(2H, m), 7.22-7.25(3H, m), 7.30(1H, s), 7.37-7.42(2H, m), 8.98(1H, s), 10.76(1H, s)
838	-Cl	-CH ₂ OH	-H		2.72(1H, dd, J=9.0, 14Hz), 3.14(1H, dd, J=6.2, 14Hz), 3.85(1H, dd, J=6.2, 9.0Hz), 4.44(1.6H, br-s), 4.49(2H, d, J=5.2Hz), 4.63(0.2H, d, J=18Hz), 4.87(0.2H, d, J=18Hz), 7.03(2H, d, J=7.8Hz), 7.11(1H, br-s), 7.24(2H, d, J=8.4Hz), 7.30-7.34(3H, m), 9.01(0.8H, s), 9.45(0.2H, s), 10.35(0.2H, s), 10.80(0.8H, s)
839	-OH	-CH ₂ OH	-H		2.58(1H, dd, J=9.5Hz, 14Hz), 3.07(1H, dd, J=5.7Hz, 14Hz), 3.79(1H, dd, J=5.7Hz, 9.5Hz), 4.69(3.8H, br-s), 4.84(0.2H, d, J=18Hz), 6.65(2H, d, J=8.4Hz), 6.98(2H, d, J=8.4Hz), 7.15(1H, dd, J=1.3Hz, 8.0Hz), 7.25(br-s, 1H), 7.35(1H, d, J=8.0Hz), 9.01(s, 1H), 9.26(0.8H, s), 9.40(0.2H, s), 10.35(0.2H, s), 10.79(0.8H, s)

Example 840

Step 1: 6-Methoxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid

To a solution of methyl 6-hydroxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-yl-acetate (335 mg) and tetra-*n*-butylammonium iodide (133 mg) in toluene (10 mL), 50% NaOH aq. (5 mL) was added. After 30 minutes' stirring, dimethyl sulfate (1.47 g) was slowly added. The reaction mixture was vigorously stirred at r.t. overnight. The solvent was removed under reduced pressure, and the residue was acidified with 4N HCl and extracted with ethyl acetate. The organic layer was dried over Na₂SO₄ and concentrated. The crude product (346 mg) was used in the following step without purification.

Step 2: 6-Methoxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-hydroxyamide


The crude product (305 mg) of step 1 was dissolved in dichloromethane (4 mL) and pyridine (3 mL) and pentafluorophenyl trifluoroacetate (664 mg) was added. After 3 hours stirring, *O*-TBDMS-hydroxylamine (465 mg) was slowly

added. The reaction mixture was stirred at r.t. overnight. The solvent was removed and the residue was dissolved in EtOAc. The organic layer was washed with 1N HCl and water, dried over Na₂SO₄ and concentrated under reduced pressure.

5 The residue was purified with preparative thin layer chromatography on silica gel (EtOAc) to give the hydroxamic acid (190 mg, 60% in 2 steps) as a pale orange solid.

¹H-NMR (DMSO-d₆, δ); 2.65(1H, m), 3.10(1H, m), 3.30(3H, s), 3.72(3H, m), 3.80(1H, m), 4.39(2H, m), 4.46(2H, m), 6.84(2H, d, J=8.61Hz), 7.02(1H, d, J=8.25Hz), 7.12(3H, m), 7.34(1H, d, J=7.89Hz), 9.00(1H, s), 10.77(1H, s)

Example 841



Step 1: 6-Methoxycarbonyl-3H-1,2,3-benzodithiazole-2-oxide

To sulphur monochloride (170 mL), a solution of methyl 4-aminobenzoate in acetic acid (200 mL) was slowly added at 0 to 10 °C. The reaction mixture was stirred at 65 °C for 4 hrs. To the cooled mixture, toluene(300 mL) was added, the solid product was collected by filtration and washed with toluene three times. After drying under reduced pressure,

this solid (30.888 g, 94 %) was dissolved in 5% sodium acetate aq. and the solution was stirred at r.t. for 2 hrs. The solid product was collected by filtration and washed with water and toluene. After drying under reduced pressure, the product (28.028 g, 98%) was obtained.

:Sawhney's procedure (Indian J. chem. 33B, 280 (1994)).

Step 2: *t*-Butyl 7-methoxycarbonyl-2*H*-2-(4-methoxy-benzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetate

To a Solution of 6-methoxycarbonyl-3*H*-1,2,3-benzodithiazole-2-oxide (11.46 g) and 2-bromo-3-(4-methoxyphenyl)propionic acid (12.96 g) in 50% MeOH aq. (100 mL), triethylamine(50 mL) was slowly added at 70 °C. The reaction mixture was stirred at 70 °C for 2 hrs. The mixture was cooled to r.t. and concentrated under reduced pressure. The residue was dissolved in EtOAc and washed with 5% Na₂CO₃ aq. and 1N HCl. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by recrystallization from Et₂O and hexane to give the product (15.131 g, 88%) as a white powder.

To a solution of 7-methoxycarbonyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one (7.00 g) in DMF (40 mL), NaH (1.47 g, 60% in oil) was added at r.t.. After 20 minutes' stirring, *t*-buthyl bromoacetate (5.56 g) was slowly added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc and washed with

water. The organic layer was dried over Na_2SO_4 and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 20/1) to give the product (6.994 g, 64%).

5

Step 3: 7-Carboxy-2*H*-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4*H*)-one-4-ylacetic acid *N*-hydroxamide

t-Butyl 7-methoxycarbonyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetate (4.81 g) was dissolved in TFA (19 mL) and ethanedithiol (1 mL). The mixture was stirred at r.t. for 2 hrs. The solvent was removed under reduced pressure, and the residue was diluted with 5% Na_2CO_3 , washed with Et_2O . The aqueous layer was acidified with 4*N* HCl and extracted with EtOAc. The organic layer was dried over Na_2SO_4 and concentrated. The residue was purified with silica gel column chromatography (eluent: EtOAc) to give the product (3.522 g, 83%).

10

15

20

25

Isobutyl chloroformate (1.304 g) was added dropwise to a THF (20 mL) solution of 7-methoxycarbonyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid (3.484 g) and *N*-methylmorpholine (0.966 g) at -5 °C. THF (1 mL) solution of *O*-TMS-hydroxylamine (1.096 g) was added after additional 20 minutes' stirring. The mixture was stirred at -5 °C for 2 hrs, gradually warmed to r.t. with stirring, and stood overnight. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was washed with 1*N* HCl and water, dried over Na_2SO_4

and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 3/1) to give the hydroxamic acid (2.943 g, 81%) as a white powder.

5 To a solution of 7-methoxycarbonyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-hydroxamide (2.827 g) in methanol (10 mL) and THF (10 mL), 1*N* NaOH (13.6 mL) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to room
10 temperature and left overnight. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in 5% Na₂CO₃ aq., and washed with Et₂O. The aqueous layer was acidified with 4*N* HCl, and the product was extracted with EtOAc. The organic layer was dried over
15 Na₂SO₄ and concentrated. The residue was purified by recrystallization from EtOAc to give the hydroxamic acid (1.234 g, 45%) as a white solid.

The following compounds listed in the Tables 20, and 20'
20 were prepared in a similar manner.

Table 20




Table 20				NMR: ¹ H-NMR (DMSO-d ₆ ; δ)	
Example No.	RR ¹	RR ²			
841	-OMe	-CO ₂ H	2.67(1H, m), 3.09(1H, m), 3.72(3H, s), 3.89(1H, m), 4.52(2H, m), 6.84(2H, d, J=8.07Hz), 7.13(2H, d, J=8.25Hz), 7.22(1H, d, J=10.26Hz), 7.85(2H, s), 9.00(1H, s), 10.80(1H, s)		
842	-OMe	-CO ₂ Me	2.67(1H, m), 3.08(1H, m), 3.73(3H, s), 3.84(3H, s), 3.92(1H, m), 4.53(2H, m), 6.84(2H, d, J=8.43Hz), 7.13(2H, d, J=8.61Hz), 7.25(1H, m), 7.84-7.90(2H, m), 9.01(1H, br-s), 10.81(1H, s)		
843	-OMe	-COONa	2.67(1H, m), 3.09(1H, m), 3.72(3H, s), 3.89(1H, m), 4.52(2H, m), 6.84(2H, d, J=8.07Hz), 7.13(2H, d, J=8.25Hz), 7.22(1H, d, J=10.26Hz), 7.85(2H, s), 9.00(1H, s), 10.80(1H, s)		
844	-Cl	-CO ₂ H	¹ H-NMR 2.75 (1H, dd, J=13.9, 8.8 Hz), 3.12 (1H, dd, J=14.3Hz, 6.2Hz), 3.98 (1H, dd, J=8.6Hz, 6.6Hz), [4.50 (s), 4.69 (d, J=17.7Hz), 4.91 (d, J=17.7Hz)] 2H, [7.22 (d, J=8.6Hz), 7.02 (d, J=8.4Hz)] 1H, 7.24 (2H, d, J=8.6Hz), 7.32 (2H, d, J=8.6Hz), 7.83 (1H, dd, J=8.4Hz, 1.8Hz), 7.86 (1H, d, J=1.8Hz), [8.99 (br), 9.39 (s)] 1H, [10.80 (br), 10.38 (s)] 1H, 13.02 (1H, br).		
845	-OPh	-CO ₂ H	2.76(1H, m), 3.13(1H, m), 4.02(1H, m), 4.53(2H, m), 6.91(2H, d, J=8.43Hz), 6.99(2H, d, J=7.50Hz), 7.14(1H, m), 7.22-7.25(3H, m), 7.39(2H, m), 7.85(1H, m), 7.88(1H, d, J=2.01Hz), 9.02(1H, s), 10.82(1H, s)		

Table 20'



Example No.	RR¹	NMR: ¹H-NMR (DMSO-d6; δ)
1005	-Cl	1.31(3H, t, J=7Hz), 2.77(1H, m), 3.14(1H, m), 4.00(1H, m), 4.29(2H, dd, J=7, 13Hz), 4.52(2H, m), 7.25-7.35(5H, m), 7.84-7.90(2H, m), 9.03(1H, br-s), 10.86(1H, br-s)
1006	-OPh	1.31(3H, t, J=7.14Hz), 2.75(1H, m), 3.13(1H, m), 3.97(1H, m), 4.29(2H, dd, J=7, 13Hz), 4.54(2H, m), 6.91(2H, d, J=8.43Hz), 6.99(2H, d, J=7.68Hz), 7.14(1H, m), 7.22-7.27(3H, m), 7.37-7.42(2H, m), 7.85(1H, dd, J=2.0, 8.6Hz), 7.91(1H, d, J=1.86Hz) 9.03(1H, br-s), 10.83(1H, br-s)

Example 846

Step 1: 7-Carboxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-allyloxyamide

To a solution of 7-methoxycarbonyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-allyloxyamide (5.067 g) in methanol (15 mL) and THF (15 mL), 1N NaOH (16.65 mL) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to room temperature and left overnight. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in 5% Na₂CO₃ aq., and washed with Et₂O. The aqueous layer was acidified with 4N HCl, and the product was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated to give the product (4.748 g, 97%) as a white solid.

Step 2: 7-Isopropylcarbamoyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-hydroxyamide

To a solution of 7-carboxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-allyl-oxyamide (221 mg) and isopropylamine (36 mg) in DMF (10 mL), HOBt

(552 mg) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide·HCl(EDC·HCl) (690 mg) were added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc and washed with 1N HCl, 5% Na₂CO₃ aq. and brine. The organic layer was dried over Na₂SO₄ and concentrated. The crude product was used in the following step without purification.

To a solution of the crude product of 7-isopropylcarbamoyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid N-allyloxyamide in 80% ethanol aq. (8 mL), Pd(OAc)₂ (12 mg), PPh₃ (53 mg), and formic acid (69 mg) were added. The reaction mixture was heated at 80 °C under nitrogen for 1 hr. The solvent was removed and the residue was dissolved in EtOAc. The organic layer was washed with water, dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by preparative thin layer chromatography on silica gel (chloroform/methanol, 8/1) to give the hydroxamic acid (207 mg, 93 % in 2 steps) as a pale orange solid.

The following compounds listed in the Tables 21 were prepared in a similar manner.






854	-OMe	-NHCH ₂ COOH	2.67(1H, m), 3.09(1H, m), 3.73(3H, s), 3.87-3.92(3H, m), 4.53(2H, m), 6.84(2H, d, J=8.43Hz), 7.14(2H, d, J=8.58Hz), 7.22(1H, d, J=8.61Hz), 7.79(1H, dd, J=1.83, 8.61Hz), 7.89(1H, d, J=2.04Hz), 8.85(1H, m), 9.00(1H, s), 10.80(1H, s), 12.63(1H, br-s)
855	-OMe	-NHC(CH ₃) ₂ COOH	1.43(6H, s), 2.67(1H, m), 3.11(1H, m), 3.73(3H, s), 3.87(1H, m), 4.51(2H, m), 6.85(2H, d, J=8.61Hz), 7.15(2H, d, J=8.61Hz), 7.20(1H, d, J=8.61Hz), 7.76(1H, d, J=8.43Hz), 7.91(1H, d, J=1.83Hz), 8.44(1H, s), 9.00(1H, s), 10.79(1H, s), 12.11(1H, br-s)
856	-OMe	-NH-(S)-CH(iBu)COOH	0.90(6H, m), 1.57-1.76(3H, m), 2.69(1H, m), 3.11(1H, m), 3.73(3H, s), 3.89(1H, m), 4.43(1H, m), 4.52(2H, m), 6.84(2H, d, J=8.43Hz), 7.15(2H, d, J=8.61Hz), 7.22(1H, d, J=8.61Hz), 7.80(1H, m), 7.94(1H, m), 8.57(1H, m), 9.00(1H, s), 10.80(1H, s)
857	-OMe		1.85-1.98(3H, m), 2.26(1H, m), 2.67(1H, m), 3.10(1H, m), 3.55(2H, m), 3.72(3H, s), 3.86(1H, m), 4.39(1H, m), 4.50(2H, d, J=8.43Hz), 6.84(2H, d, J=8.43Hz), 7.13(2H, d, J=8.43Hz), 7.18(1H, d, J=8.22Hz), 7.47-7.50(2H, m), 9.01(1H, s), 10.80(1H, s)
858	-OMe	-NH(CH ₂) ₂ CH(CH ₃) ₂	0.88(6H, d, 6.60Hz), 1.39(2H, m), 1.58(1H, m), 2.66(1H, m), 3.09(1H, m), 3.25(2H, m), 3.71(3H, s), 3.85(1H, m), 4.50(2H, m), 6.83(2H, d, J=8.61Hz), 7.12(2H, d, J=8.58Hz), 7.18(2H, d, J=8.61Hz), 7.74(1H, m), 7.83(1H, d, J=1.83Hz), 8.39(1H, m), 9.00(1H, s), 10.79(1H, s)
859	-OMe	-NH(CH ₂) ₂ N(CH ₃) ₂	2.42(6H, s), 2.63-2.72(3H, m), 3.10(1H, m), 3.37(2H, m), 3.73(3H, s), 3.87(1H, m), 4.52(2H, m), 6.84(2H, d, J=8.61Hz), 7.14(2H, d, J=8.58Hz), 7.21(2H, d, J=8.61Hz), 7.78(1H, dd, J=2.01, 8.61Hz), 7.87(1H, d, J=2.01Hz), 8.15(1H, s), 8.55(1H, m), 10.82(1H, br-s)
860	-OMe	-NH(CH ₂) ₂ NEt ₂	0.99(6H, t, J=7.14Hz), 2.55-2.70(7H, m), 3.10(1H, m), 3.35(2H, m), 3.73(3H, s), 3.88(1H, m), 4.52(2H, m), 6.84(2H, d, J=8.61Hz), 7.13(2H, d, J=8.43Hz), 7.20(2H, d, J=8.58Hz), 7.76(1H, d, J=2.01, 8.43Hz), 7.84(1H, s), 8.20(1H, s), 8.45(1H, m), 10.81(1H, br-s)
861	-OPh	-NH(CH ₂) ₂ CH(CH ₃) ₂	0.89(6H, d, 6.60Hz), 1.39(2H, m), 1.60(1H, m), 2.74(1H, m), 3.17(1H, m), 3.26(2H, m), 3.93(1H, m), 4.52(2H, m), 6.91(2H, d, J=8.43Hz), 6.99(2H, d, J=7.50Hz), 7.11-7.26(4H, m), 7.40(2H, m), 7.76(1H, d, J=8.40Hz), 7.87(1H, s), 8.41(1H, m), 9.01(1H, s), 10.80(1H, s)
862	-OPh	-NH(CH ₂) ₂ OMe	2.74(1H, m), 3.15(1H, m), 3.26(3H, s), 3.33-3.44(4H, m), 3.93(1H, m), 4.52(2H, m), 6.91(2H, d, J=8.61Hz), 7.00(2H, d, J=8.79Hz), 7.16-7.26(4H, m), 7.37-7.42(2H, m), 7.78(1H, d, J=8.61Hz), 7.89(1H, d, J=1.83Hz), 8.54(1H, m), 9.01(1H, s), 10.81(1H, s)

Table 21



Example No.	RR ¹	RR ²	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
846	-OMe	-NHIPr	1.14(6H,m), 2.67(1H, m), 3.11(1H, m), 3.73(3H, s), 3.86(1H, m), 4.09(1H, m), 4.51(2H, m), 6.85(2H, d, J=8.61Hz), 7.14(2H, d, J=8.40Hz), 7.18(1H, d, J=8.97Hz), 7.76(1H, d, J=8.61Hz), 7.87(1H, s), 8.20(1H, d, J=7.68Hz), 9.00(1H, s), 10.79(1H, s)
847	-OMe	-NH ₂	2.65(1H, m), 3.12(1H, m), 3.71(3H, s), 3.86(1H, m), 4.50(2H, d, J=8.43Hz), 7.13(2H, d, J=8.43Hz), 7.17(1H, d, J=8.79Hz), 7.36(1H, s), 7.78(1H, m), 7.86(1H, d, J=1.65Hz), 7.95(1H, s), 8.99(1H, br-s), 10.79(1H, br-s)
848	-OMe	-NH-cyclopropyl	0.56(2H, m), 0.67(2H, m), 2.65(1H, m), 2.83(1H,m), 3.11(1H, m), 3.73(3H, s), 3.86(1H, m), 4.51(2H, m), 6.84(2H, d, J=8.43Hz), 7.13(2H, d, J=8.43Hz), 7.18(1H, m), 7.74(1H, m), 7.82(1H, d, J=1.83Hz), 8.41(1H, d, J=4.05Hz), 9.00(1H, s), 10.79(1H, s)
849	-OMe	-morpholino	2.67(1H, m), 3.12(1H, m), 3.32-3.60(8H, m), 3.72(3H, s), 3.87(1H, m), 4.50(2H, m), 6.84(2H, d, J=8.61Hz), 7.13(2H, d, J=8.58Hz), 7.17(1H, m), 7.35(1H, m), 7.41(1H, d, J=1.65Hz), 9.00(1H, s), 10.80(1H, s)
850	-OMe	-Piperadino-4-CHO	2.68(1H, m), 3.15(1H, m), 3.17-3.44(8H, m), 3.72(3H, s), 3.87(1H, m), 4.51(2H, m), 6.84(2H, d, J=8.40Hz), 7.14(2H, d, J=8.04Hz), 7.18(1H, d, J=8.61Hz), 7.37(1H, d, J=8.22Hz), 7.44(1H, s), 8.06(1H, s), 9.01(1H, s), 10.81(1H, s)
851	-OMe	-NHCH(CH ₂ OH) ₂	2.66(1H, m), 3.10(1H, m), 3.48(4H, m), 3.71(3H, s), 3.83-3.94(2H, m), 4.50(2H, m), 4.63(2H, m), 6.83(2H, d, J=8.43Hz), 7.13(2H, d, J=8.61Hz), 7.18(1H, d, J=8.43Hz), 7.76(1H, d, J=6.78Hz), 7.90(1H, s), 7.98(1H, d, J=8.25Hz), 8.98(1H, s), 10.78(1H, s)
852	-OMe	-N((CH ₂) ₂ OH) ₂	2.65(1H, m), 3.09(1H, m), 3.36-3.57(8H, m), 3.71(3H, s), 3.85(1H, m), 4.48(2H, m), 4.80(2H, m), 6.83(2H, d, J=8.61Hz), 7.10-7.15(3H, m), 7.33(1H, d, J=8.25Hz), 7.41(1H, s), 9.00(1H, br-s), 10.78(1H, br-s)
853	-OMe	-NHC(Me) ₂ CH ₂ OH	1.28(6H,s), 2.66(1H, m), 3.11(1H, m), 3.48(2H, d, J=6.03Hz), 3.71(3H, s), 3.84(1H, m), 4.49(2H, m), 4.85(1H, m), 6.83(2H, d, J=8.61Hz), 7.12-7.17(3H, m), 7.54(1H, s), 7.70(1H, d, J=8.58Hz), 7.83(1H, d, J=1.83Hz), 8.98(1H, s), 10.78(1H, s)

863	-OPh	-NH(CH ₂) ₂ OEt	1.10(3H, t, J=6.96Hz), 2.74(1H, m), 3.14(1H, m), 3.31-3.42(6H, m), 3.93(1H, m), 4.52(2H, m), 6.91(2H, d, J=8.61Hz), 6.99(2H, d, J=7.50Hz), 7.11-7.25(4H, m), 7.37-7.42(2H, m), 7.78(1H, d, J=8.61Hz), 7.88(1H, s), 8.54(1H, m), 9.01(1H, s), 10.81(1H, s)
864	-OMe		0.88(6H, m), 1.30(2H, m), 1.51(1H, m), 1.97(3H, s), 2.69(1H, m), 3.12(1H, m), 3.73(3H, s), 3.88-3.92(2H, m), 4.07(1H, m), 4.27(1H, m), 4.52(2H, m), 6.85(2H, d, J=7.50Hz), 7.15(2H, d, J=8.79Hz), 7.22(1H, m), 7.76(1H, d, J=8.61Hz), 7.86(1H, s), 8.20(1H, d, J=8.61Hz), 9.00(1H, s), 10.79(1H, s)
865	-OMe		0.87(6H, m), 1.39(2H, m), 1.58(1H, m), 2.68(1H, m), 3.11(1H, m), 3.39(2H, m), 3.73(3H, s), 3.87(1H, m), 4.03(1H, m), 4.52(3H, m), 6.84(2H, d, J=8.07Hz), 7.13-7.21(3H, m), 7.78(1H, m), 7.89(1H, m), 8.00(1H, d, J=8.61Hz), 9.00(1H, br-s), 10.80(1H, s)
866	-Cl	-NHCH ₂ -2-furyl	2.73(1H, dd, J=13.9Hz, 9.3Hz), 3.13(1H, d, J=14.4Hz, 6.6Hz), 3.94(1H, dd, J=8.8Hz, 6.4Hz), 4.46(2H, d, J=5.4Hz), [4.50 (s), 4.72 (d, J=17.9Hz), 4.95 (d, J=17.9Hz)] 2H, 6.26(1H, d, J=2.7Hz), 6.38(1H, dd, J=3.0Hz, 1.6Hz), 7.20(1H, d, J=8.6Hz), 7.24 (2H, d, J=8.4Hz), 7.32(2H, d, J=8.4Hz), 7.56(1H, t, J=0.9Hz), 7.79(1H, dd, J=8.4Hz, 2.0Hz), 7.87(1H, d, J=2.2Hz), 8.95(1H, t, J=5.4Hz), [8.99 (br), 9.42 (s)] 1H, [10.78 (br), 10.35(s)] 1H.

Example 867


Step 1: 7-Amino-2H-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4H)-one-4-ylacetic acid N-hydroxyamide

5 The product in Example 757(400mg) was dissolved in MeOH (10mL) and AcOH (1 mL). 10% Pd/C (200 mg) was added and the reaction mixture was vigorously stirred at r.t. for 5hrs under H₂. The catalyst was removed by filtration, and the filtrate was concentrated under reduced pressure. The
10 residue was purified by silica gel column chromatography (chloroform/MeOH, 30/1, 10/1) to give the product as a pale yellow solid (162mg).

15 The following compounds listed in the Table 22 were prepared in a similar manner.

Table 22

Table 22				

Example 869

Step 1: Ethyl 6-azidomethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

5 Ethyl 6-hydroxymethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate (1.75 g), prepared by the procedure described in Example 835 (Step 1), was dissolved in THF (16mL), and triethylamine(0.93ml) and methanesulfonyl chloride(0.51ml) were slowly added at 0 °C. The reaction mixture was stirred at 0°C for 2 hrs. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF(30ml), and sodium azide(636mg) was added at 0 °C. Stirring was continued for 1hr at 0 °C and for 2hrs at r.t.. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (eluent: hexane:EtOAc = 5:1, 3:1). 1.83g of the product was obtained

as oil.

Step 2: Ethyl 6-N-t-butoxycarbonylaminomethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

5 The compound obtained in Step 1 (1.07 g) was dissolved in methanol(20 mL) and EtOAc(5 mL). 10% Pd/C(300 mg) was added, and the reaction mixture was vigorously stirred at r.t. for 2hrs under H₂. The catalyst was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was dissolved in MeOH (20 ml), di-t-butylidicarbonate (0.74 g) and NaHCO₃ (0.21 g) were added at 10 0 °C. The reaction mixture was stirred at r.t. overnight. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (Eluent: Hexane/EtOAc, 5/1, 3/1, 2/1). 0.67g of the product was obtained as oil.

20 Step 3: 6-N-t-Butoxycarbonylaminomethyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-hydroxyamide


25 The compound in step 2 was converted to 600mg of the hydroxamic acid by the reactions analogous to those described in Example 777.


Step 4: 6-Aminomethyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetic acid *N*-hydroxyamide hydrochloride

5 To a solution of the compound in step 3 (485 mg) in 4*N* HCl/dioxane (3 mL) was stirred for 1hr at 0 °C. The reaction mixture was diluted with Et₂O and the precipitate was filtered and washed with Et₂O. 300mg of the product was obtained as pale yellow solid.

10 The following compounds listed in the Table 23 were prepared in a similar manner.

Table 23

				NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
Example No.	RR ¹	RR ²	RR ³	
869	-OMe	-CH ₂ NH ₂	-H	2.62(1H,dd,J=9Hz,14Hz),3.06(1H,dd,J=6,14Hz),3.71(3H,s),3.80(1H,dd,J=6,9Hz),4.01(2H,m),4.48-5.00(2H,m),6.83(2H,d,J=8.6Hz),7.09(2H,d,J=8.6Hz),7.19(1H,m),7.39(2H,m),8.32(3H,b s),8.95,9.43(1H,bs),10.30,10.83(1H,bs).
870	-OMe	-CH ₂ NHBoc	-H	1.33(9H,s),2.57(1H,m),3.06(1H,m),3.66(3H,s),3.70(1H,dd,J=6,9Hz),4.05(2H,bd,J=5.7 Hz),4.3-4.9(2H,m),6.76(2H,d,J=8.5Hz),6.88(1H,d,J=8Hz),6.97(1H,s),7.05(2H,d,J=8.5Hz),7.25(1H,d,J=8Hz),7.34(1H,t,J=5.7Hz)m,8.91,9.33(1H,bs),10.24,10.69(1H,bs).
871	-H	-CH ₂ NH ₂	-H	2.62(1H,dd,J=9.5,14.2Hz),3.09(1H,dd,J=5.7,14.2Hz),3.81(1H,m),3.95,3.96(2H,s),4.44-4.88(2H,m),7.10-7.24(6H,m),7.35(2H,m),8.33(3H,bs),8.92,9.40(1H,bs),10.27,10.81(1H,bs).
872	-OMe	-H	-CH ₂ NH ₂	2.59(1H,m),3.09(1H,dd,J=5.5,14Hz),3.72(3H,s),3.83(1H,m),3.96(2H,s),6.83(2H,d,J=8.5Hz),4.48-5.00(2H,m),7.13(2H,d,J=8.5Hz),7.17(1H,d,J=8.5Hz),7.37(1H,d,J=8.5),7.49(1H,s),8.19(3H,bs),9.43,8.97(1H,bs),10.80,10.32(1H,bs).

Example 873

Step 1: Ethyl 6-tetrazolyl-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetate

5 Ethyl 6-cyano-2H-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4H)-one-4-ylacetate (700 mg) was dissolved in DMF (5 mL), and sodium azide (573 mg), ammonium chloride (641 mg) were added at r. t.. The reaction mixture was stirred at 90°C for 24 hrs, diluted with water (50 ml) and 1N HCl (5 ml),
10 and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (Eluent: chloroform/MeOH, 30/1, 20/1, chloroform/MeOH/AcOH=20/1/0.1). 800mg of the product was
15 obtained as oil.


Step 2: 6-Tetrazolyl-2H-2-benzyl-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-hydroxyamide


20 The compound in step 1 was converted to 594mg of the product by the procedure analogous to Example 777.

The following compounds listed in the Table 24 were

prepared in a similar manner.

Table 24

				NMR: ¹ H-NMR (DMSO-d6; δ)	
Example No.	RR ¹	RR ²	RR ³		
873	-OMe	-5-Tetrazolyl	-H	2.70(1H,dd,J=9.3,14.3Hz),3.11(1H,dd,J=6.0,14.3Hz),3.05(3H,s),3.94(dd,1H,J=5.9,9.1Hz),4.50-4.97(2H,m),6.84(2H,d,J=8.6Hz),7.14(2H,d,J=8.6Hz),7.60-7.82(3H,m),9.06,9.49(1H,bs),10.42,10.86(1H,bs).	
874	-H	-5-Tetrazolyl	-H	2.75(1H,dd,J=8Hz,14Hz),3.20(1H,m),3.98(1H,m),4.50-5.10(2H,m),7.30(5H,m),7.70(2H,m),7.82(1H,s),9.05,9.55(1H,bs),10.40,10.85(1H,bs).	
875	-OMe	-H	-5-Tetrazolyl	2.69(1H,dd,J=9.2,14Hz),3.11(1H,dd,J=6.14Hz),3.72(3H,s),3.94(1H,dd,J=6.9,2Hz),4.50-4.98(2H,m),6.84(2H,d,J=8.5Hz),7.14(2H,d,J=8.5Hz),7.35(1H,d,J=8.8Hz),7.94(1H,dd,J=2.8,8Hz),8.07(1H,d,J=2Hz),9.42,9.03(1H,bs),10.83,10.40(1H,bs),.	

Example 876

Step 1: Ethyl 6-(2,3-di-*t*-butoxycarbonylguanidino)-methyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-ylacetate

The compound, obtained in Example 869 (Step 2), was dissolved in THF (2 mL) and H₂O (0.04 mL). The *N,N'*-di-*t*-butoxycarbonyl-*S*-methylisothiurea (369 mg) and triethylamine (89 μ l) were added at r.t. and the stirring was continued for 2hrs at 50 °C. The reaction mixture was concentrated under reduced pressure, and the residue was purified with silica gel column chromatography (Eluent: hexane/EtOAc, 8/1, 6/1). 200 mg of the product was obtained as oil.

Step 2: 6-(2,3-Di-*t*-butoxycarbonylguanidino)methyl-2*H*-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4*H*)-one-4-yl-acetic acid *N*-hydroxyamide


The compound in step 1 was converted to 510 mg of the hydroxamic acid by the procedure analogous to Example 777.

Step 3: 6-Guanidinomethyl-2*H*-2-(4-methoxybenzyl)-1,4-

benzothiazin-3(4H)-one-4-ylacetic acid hydroxyamide

The compound in Step 2 (110 mg) was dissolved in TFA (3 ml)-ethanedithiol (100 ul) at 0°C, and stirred at r.t. for 1hr. The reaction mixture was diluted with Et₂O-Toluene (4:1) and, the product(29mg) was obtained as precipitate.

¹H-NMR (DMSO-d₆, δ); 2.62(1H, dd, J=9Hz, 14Hz), 3.08(1H, dd, J=6, 14Hz), 3.78(3H, s), 3.78(1H, dd, J=6Hz, 9Hz), 4.34(2H, d, J=6Hz), 4.45-4.90(2H, m), 6.83(2H, d, J=8.4Hz), 6.98(1H, d, J=8Hz), 7.10(2H, d, J=8.4Hz), 7.16(1H, s), 7.30(4H, bs), 7.38(1H, d, J=8Hz), 8.00(1H, t, J=6Hz), 8.97, 9.42(1H, bs), 10.31, 10.80(1H, bs).

Exemplen 877**Step 1: 2-Methyl-5H-5-(4-chlorobenzyl)-pyrazolo[4,3-b]thiazin-6(7H)-one**

A solution of 5-methylpyrazolo[3,4-d]dithiazolium chloride (930 mg), which was prepared by Chenard's procedure (J. Org. Chem. 49, 1224 (1984)), in 20% NaOH aq. (15 mL) was heated at 80 °C for 1 hr. A solution of 2-bromo-3-(4-chlorophenyl)propionic acid (1.05 g) in 20% NaOH aq. (10 mL) was added at 80 °C and the temperature was

maintained at 80 °C for 1.5 hrs. The mixture was cooled to r.t. and concentrated under reduced pressure. The residue was dissolved in acetic acid (50 mL) and heated at 80 °C for 2.5 hours. The reaction mixture was cooled to r.t. and concentrated under reduced pressure. The residue was dissolved in EtOAc and washed with water, 5% Na₂CO₃ aq. and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by recrystallization from Et₂O to give the product (512 mg, 44%).

Step 2: *t*-Butyl 2-methyl-5*H*-5-(4-chlorobenzyl)-pyrazolo[4,3-*b*]thiazin-6(7*H*)-one-7-ylacetate

To a solution of the product of Step 1 (235 mg) in DMF (10 mL), NaH (52 mg, 60% in oil) was added. After 30 minutes' stirring, *t*-buthyl bromoacetate (203 mg) was slowly added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc and washed with 1N HCl, 5% Na₂CO₃ aq., and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 20/1) to give the product(0.33 g, quant.).


Step 3: 2-Methyl-5*H*-5-(4-chlorobenzyl)-pyrazolo[4,3-*b*]thiazin-6(7*H*)-one-7-ylacetic acid N-hydroxyamide


The product of Step 2 (0.32 g) was solved in TFA (15 mL) at r.t.. The mixture was stirred for 2 hours. The

solvent was removed under reduced pressure. The crude product(0.28 g) was used in the following step without purification.

Isobutyl chloroformate (120 mg) was added dropwise to a
5 THF (10 mL) solution of 2-methyl-5*H*-5-(4-chlorobenzyl)-
pyrazolo[4,3-*b*]thiazin-6(7*H*)-one-7-ylacetic acid (0.28 g)
and N-methylmorpholine (170 mg) at -5 °C. THF (1 mL)
solution of *O*-TMS-hydroxylamine (101 mg) was added after
additional 20 minutes' stirring. The mixture was stirred at
10 -5 °C for 2 hrs, gradually warmed to r.t. with stirring, and
stood overnight. The reaction mixture was quenched with
water and extracted with EtOAc. The organic layer was
washed with saturated NaHCO₃, 1N HCl , and brine, dried
over Na₂SO₄ and concentrated under reduced pressure. The
15 residue was purified by recrystallization from Et₂O to give
the hydroxamic acid (221.3 mg, 77% in 2 steps) as a white
powder.

The following compounds listed in the Table 25 were
20 prepared in a similar manner.

Table 25			
Example No.	RR¹	NMR: ¹H-NMR (DMSO-d6; δ)	
877	-Cl	2.79(1H, m), 3.17(1H, m), 3.74(3H, s), 3.92(1H, m), 4.33(2H, m), 7.25(2H, d, J=8.40Hz), 7.34(2H, d, J=8.25Hz), 7.61(1H, s), 8.88(1H, s), 10.62(1H, s)	
878	-OPh	2.77(1H, m), 3.17(1H, m), 3.74(3H, s), 3.90(1H, m), 4.34(2H, m), 6.92(2H, d, J=8.48Hz), 6.99(2H, d, J=8.60Hz), 7.14(1H, m), 7.23(2H, d, J=8.56Hz), 7.37-7.42(2H, m), 7.62(1H, s), 8.90(1H, s), 10.64(1H, s)	

Example 879

Step 1: *t*-Butyl 2-(3-amino-1-methyl-4-pyrazolylthio)-3-(4-chlorophenyl)propanoate

A solution of 5-methylpyrazolo[3,4-*d*]dithiazolium chloride (880 mg), which was prepared by Chenard's procedure (J. Org. Chem. 49, 1224 (1984)), in 20% NaOH aq. (15 mL) was heated at 90 °C for 1.5 hrs. The mixture was cooled to r.t. and the pH was adjusted to 6.0 with AcOH, and adjusted to 10 by Na₂CO₃ sat.. To a mixture of the crude product and Na₂CO₃ (1.20 g), a solution of *t*-buthyl 2-bromo-3-(4-chlorophenyl)propionate (1.21 g) in DMF (20 mL) was added. The mixture was stirred at r.t. overnight and concentrated under reduced pressure. The residue was dissolved in EtOAc and washed with water and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 2/1) to give the product(820 mg, 59%).

Step 2: *t*-Butyl 2-(3-iodo-1-methyl-4-pyrazolylthio)-3-(4-chlorophenyl)propanoate

To a solution of isoamyl nitrite (605 mg) in diiodomethane (5 mL) at 100 °C, a solution of the product of Step 1 (380 mg) in dichloromethane (2 mL) were added dropwise. The reaction mixture was stirred at 100 °C for 1 hr. The solvent was removed under reduced pressure. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 20/1) to give the product (417 mg, 84%).

Step 3: Methyl 2-(S)-[2-methyl-5*H*-5-(4-chlorobenzyl)-pyrazolo[4,3-*b*]thiazin-6(7*H*)-one-7-yl]-propanoate

The product of Step 2 (335 mg) was dissolved in TFA (10 mL). The mixture was stirred at r.t. for 2 hrs. The solvent was removed under reduced pressure. To the solution of the residue (347 mg) and L-alanine methyl ester·HCl (215 mg) in DMF (10 mL), Et₃N (290 uL), HOBt (235 mg) and EDC·HCl (295 mg) were added. The reaction mixture was stirred at r.t. for 2 days. The solvent was removed under reduced pressure. The residue was dissolved in EtOAc and washed with 1N HCl, 5% Na₂CO₃ aq. And brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (eluent: toluene/EtOAc, 2/1) to give the alanine derivative (316 mg, 89% in 2 steps).

To a solution of the alanine derivative (205 mg) and in

DMF (10 mL), K₂CO₃ (112 mg), Cu powder (128 mg) were added. The reaction mixture was heated to 140 °C and stirred for 3 hrs. The mixture was diluted with EtOAc and filtered. The solvent was removed under reduced pressure. The residue was
5 purified with silica gel column chromatography (eluent: toluene/EtOAc, 10/1) to give the product (128 mg, 83%).


Step 4: 2-(S)-[2-Methyl-5*H*-5-(4-chlorobenzyl)-pyrazolo[4,3-*b*]thiazin-6(7*H*)-one-7-yl]-propanoic acid N-hydroxyamide

10 To a solution of the product of Step 3 (141 mg) in methanol (10 mL), 1N NaOH (2.0 mL) was slowly added at 0 °C. After 30 minutes' stirring, the mixture was warmed to r.t. and left overnight. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in EtOAc and
15 water, washed with 1N HCl and water. The organic layer was dried over Na₂SO₄ and concentrated. To a THF (10 mL) solution of the crude product(151 mg)and N-methylmorpholine (45 mg), isobutyl chloroformate (61 mg) was added dropwise at -5 °C. THF (0.5 mL) solution of *O*-TMS-hydroxylamine (51
20 mg) was added after additional 30 minutes' stirring. The mixture was stirred at -5 °C for 2 hrs , gradually warmed to r.t. with stirring, and stood overnight. The reaction mixture was quenched with water, concentrated under reduced pressure, and dissolved in EtOAc. The organic layer was
25 washed with 1N HCl and brine, dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by recrystallization from chloroform, Et₂O, and


hexane to give the hydroxamic acid(110 mg, 78% in 2 steps) as a white powder.

The following compounds listed in the Table 33 were
5 prepared in a similar manner.

Table 33



Example No.	RR ¹	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
1007	(S) -Me	1.45(3H, m), 2.76(1H, m), 3.14(0.5H, m), 3.26(0.5H, m), 3.73(1.5H, s), 3.75(1.5H, s), 3.85(0.5H, m), 3.96(0.5H, m), 5.01(0.5H, m), 5.21(0.5H, m), 7.25(2H, d, J=8.43Hz), 7.32-7.37(2H, m), 7.59(0.5H, s), 7.62(0.5H, s), 8.76(0.5H, s), 8.78(0.5H, s), 10.42(1H, s)
1008	(R) -Me	1.45(3H, m), 2.76(1H, m), 3.14(0.5H, m), 3.25(0.5H, m), 3.73(1.5H, s), 3.74(1.5H, s), 3.85(0.5H, m), 3.96(0.5H, m), 5.01(0.5H, m), 5.21(0.5H, m), 7.25(2H, d, J=8.43Hz), 7.33(2H, m), 7.59(0.5H, s), 7.61(0.5H, s), 8.76(0.5H, s), 8.78(0.5H, s), 10.42(1H, s)
1009	(R) -iPr	0.67(3H, m), 1.02(3H, m), 2.72-2.80(2H, m), 3.23(1H, m), 3.74(1.5H, s), 3.77(1.5H, s), 3.91(1H, m), 4.62(0.5H, d, J=10.62Hz), 4.75(0.5H, d, J=10.62Hz), 7.25(2H, m), 7.35(2H, m), 7.61(0.5H, s), 7.63(0.5H, s), 8.78(0.5H, s), 8.81(0.5H, s), 10.40(1H, s)

Example 880

Step 1: 7-Allyloxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one

To a solution of 7-methoxymethoxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one (1 g) in MeOH (20 ml), 4N HCl-dioxane (2 ml) was added at r.t. and stirring was continued for 24hrs. The reaction mixture was concentrated and Et₂O was added. The colorless crystal (800mg) was filtered.

The product was dissolved in DMF (5 mL), and allyl bromide (252 ul) and K₂CO₃ (366 mg) were added at 0 °C. The reaction mixture was stirred at r.t. for 24hrs, diluted with water, and extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (Eluent: hexane/EtOAc, 4/1, 3/1). 560mg of the product was obtained as oil.

Step 2: 7-Allyloxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-

3(4H)-one-4-ylacetic acid N-allyloxyamide

The product of Step 1 (560 mg) was converted to 7-allyloxy-2H-2-(4-methoxybenzyl)-1,4-benzothiazin-3(4H)-one-4-ylacetic acid N-allyloxyamide as described in Example 777.


5 The product (650 mg) was obtained as oil.

Step 3: 7-Hydroxy-2H-2-(4-methoxybenzyl)-1,4-benzo-thiazin-3(4H)-one-4-ylacetic acid N-hydroxylamide


To a solution of the product of Step 2 (650 mg) was
10 dissolved in EtOH (8 ml) and H₂O (2 ml), tetrakis-
triphenylphosphine palladium (175 mg) and formic acid (108
ul) were added at r.t. The mixture was stirred under reflux
for 2hrs. The reaction mixture was diluted with water and
extracted with EtOAc. The organic layer was washed with
15 sat.NaHCO₃, brine, 1N HCl, and brine, dried over MgSO₄ and
concentrated under reduced pressure. The residue was
purified with silica gel column chromatography (Eluent:
chloroform/MeOH, 50/1, 30/1). The oily residue was
crystalized in toluene-Et₂O, and the product (345 mg) was
20 obtained as pale yellow crystal.

The following compounds listed in the Table 26 were prepared in a similar manner.

Table 26



Example No.	RR ¹	RR ²	RR ³	NMR: ¹ H-NMR (DMSO-d ₆ ; δ)
880	-OMe	-H	-OH	2.62(1H,dd,J=9.3,14Hz),3.09(1H,dd,J=6,14Hz),3.73(3H,s),3.74(1H,m),4.30-4.74(2H,m),6.67(2H,m),6.82(2H,d,J=8.5Hz),6.94(1H,d,J=9Hz),7.10(2H,d,J=8.5Hz),9.33,8.95(1H,bs),9.55(1H,bs),10.71,10.28(1H,bs).
881	-OMe	-OH	-H	2.58(1H,dd,J=9.6,14.5Hz),3.06(1H,dd,J=5.2,14.5Hz),3.69(3H,s),3.69(1H,m),4.2-4.9(2H,m),6.47(1H,dd,J=2,8.4Hz),6.53(1H,d,J=2Hz),6.53(2H,d,J=8.8Hz),7.09(3H,m),8.97(1H,bs),9.36,9.72(1H,s),10.73,10.32(1H,bs).

Example 882

Step 1: Methyl 7-fluoro-2H-2-benzyl-2-methyl-1,4-benzothiazin-3(4H)-one-4-yl-acetate

Methyl 7-fluoro-2H-2-benzyl-1,4-benzothiazin-3(4H) - one-4-yl-acetate (100 mg, 0.29 mmol) was dissolved in THF (5 mL) at -78 ° C and lithium hexamethyl disilazane 1M solution in THF (0.64 mmol, 0.64 ml) was added. The mixture was stirred for 1 hr and methyl iodide (39.7 ml, 0.64 mmol) was added. The mixture was allowed to warm to r.t., then poured into saturated solution of sodium carbonate (5 ml), diluted with EtOAc. The 2 layers were separated and the water layer extracted 2 additional times with ethyl acetate. The combined organic layers were dried with MgSO₄, evaporated and carried out to the next step with no further purification.

Step 2: 7-Fluoro-2H-2-benzyl-2-methyl-1,4-benzo-thiazin-3(4H)-one-4-yl-acetic acid

Compound in step 1 was dissolved in dioxane (2 ml) and

saturated solution of lithium hydroxide (2 ml). The mixture was stirred at room temperature for 12 hours, then acidified with conc.HCl and extracted with ethyl acetate, dried with MgSO₄ and evaporated to give product 3 as a solid.

5

Step 3: 7-Fluoro-2H-2-benzyl-2-methyl-1,4-benzo-thiazin-3(4H)-one-4-yl-acetic acid N-hydroxyamide

Acid obtained obtained in step 2 was dissolved in 1 ml of DCM and 1 ml of pyridine. Pentafluorophenol trifluoroacetate (74.7 ml, 0.43 mmol) was added and the mixture stirred at room temperature for 1 hour. Then, O-TBDMSNH₂OH (129 mg, 0.87 mmol) was added, the mixture stirred for an additional 12 hours then acidified until pH = 1 with cHCl, diluted with DCM (3 ml) and water (2 ml). The organic layer was separated and then evaporated, the solid obtained re-dissolved in DMSO and purified using reverse phase HPLC.


10

Retention time of HPLC analysis: 2.69min

Mass: 361.2, 328.0 (M.W. =360)

15

Example 883



20

Step 1: Methyl [2H-2-benzyl-6-N-methylcarbamoyl-3(4H)-oxo-1,4-benzothiazine-4-yl]-acetate

Methyl [2H-2-benzyl-6-carboxyl-3(4H)-oxo-1,4-benzothiazine-4-yl]acetate (62mg, 0.17mmol), diisopropylethylamine (45 ml, 0.25 mmol), HOBT (35 mg, 0.25 mmol) and EDC.HCl (50 mg, 0.25 mmol) were dissolved in DCM (2 ml) and the reaction mixture was stirred at room temperature for 1 hour. Methylamine (64 ml, 0.51 mmol) (8.03 M in ethanol) was added to the mixture and stirring was continued for an additional 12 hours. The reaction mixture was diluted in DCM (5 ml) and washed with 1N HCl (2 x 2 ml), water (2 ml) and brine (2 ml). The organic layer was concentrated *in vacuo* and the crude amide was used directly in the next step.

Step 2: [2H-2-Benzyl-6-N-methylcarbamoyl-3(4H)-oxo-1,4-benzothiazine-4-yl]-acetic acid N-hydroxyamide

Potassium hydroxide (109 mg, 1.93 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) were dissolved in methanol (0.53 ml and 0.9 ml respectively) at room temperature. Once a solution was obtained the potassium hydroxide solution was poured into the hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for 1 hour. The precipitate (potassium chloride) was filtered off, the liquid added into the solution of the methyl ester, obtained in step 1,

(in methanol (1 ml) and stirring was continued for an additional 3 hours. Acetic acid (0.5 ml) was added to the mixture and the solvent was evaporated to dryness. The residue was then re-dissolved in water and extracted with ethyl acetate (2 x 3ml). The combined organic extracts were evaporated to dryness, re-dissolved in DMSO and purified *via* reverse phase chromatography to yield the major product of the reaction together with small amount of the sulfoxide.

The following compounds listed in the Table 27 and 28 were prepared in a similar manner.




Table 27					
Example No.	RR¹	RR²	mol. weight	Mass (m/e)	HPLC rt (min)
883	-OMe	-NHMe	415.5	416.4, 383.2	2.88
884	-H	-NH(CH₂)₂OMe	429.5	430.2, 397	2.62
885	-H	-morpholino	441.5	442.2, 409.2	2.71
886	-H	-NHCH₂-cyclopropyl	425.5	426.2, 393.2	2.91
887	-H	-NHCH₂-2-furyl	451.5	452.2, 419.2	2.93
888	-H	-NMe₂	399.5	400.2, 366.8	2.71
889	-H	-NHtPr	413.5	414.2, 381.0	3.85
890	-H	-NH-cyclopentyl	439.5	440.4, 407.2, 379	3.99
891	-H	-NH-3-isoxazolyl-5-Me	452.5	453.2, 420.2	3.86
892	-H	-piperidino	439.5	440.4, 407.2	3.96
893	-H	-NH-3-pyrazolyl	437.5	438.2, 405	3.77
894	-H	-NHCH₂CF₃	453.4	454.2, 421.0	3.97
895	-H	-NH(CH₂)₂OH	415.5	416.2, 383.0	3.70
896	-H	-NHMe	385.4	386.2, 353.0	3.77
897	-H	-NH-2-pyridyl	448.5	449.0	2.25
898	-H	-NH₂	371.4	372.5, 339.0	3.55
899	-H	-NH-cyclopropyl	411.5	412.2, 379.2	3.83
900	-OMe	-NH-3-pyrazolyl	467.5	468.4, 435.2	2.97
901	-OMe	-NHPH	477.5	478.2, 445.2	3.01
902	-OMe	-NHCH₂CF₃	483.5	484.3, 451.0	2.84

Table 28

Example No.	RR ¹	RR ²	RR ³	mol. weight	Mass (m/e)	HPLC rt (min)
903	-H	-NPr2	-	455.6	456.2, 423.2	2.86
904	-H	-piperidino	-	439.5	440.2, 407.2	4.02
905	-H	-piperidino	O	455.5	456.2, 437.9, 422.9, 405.2	3.95
906	-H	-NMe2	O	415.5	416.2, 398.2, 383.2, 365.2	3.74
907	-H	-NMe2	-	399.5	400.2, 367.2	3.73
908	-H	-NH-cyclopropyl	O	427.5	428.2, 410.2, 395.2	3.72
909	-H	-NH-cyclopropyl	-	411.5	412.2, 379.0	3.82
910	-H	-NHMe	-	385.4	386.2, 353.0	2.88
911	-H	-NHPr	-	413.5	414.2, 381.2	3.97
912	-H	-NH-cyclopentyl	O	455.5	455.9, 438.2, 422.9	3.90
913	-H	-NH-3-pyrazolyl	O	453.5	454.2, 436.4, 421.4	3.78
914	-H	-NH-3-pyrazolyl	-	437.5	438.2, 405.2	3.80
915	-H	-NH2	-	371.4	372.0, 339.0	3.62
916	-H	-NHPr	O	429.5	430.4, 412.2, 397.5	3.84
917	-H	-morpholino	O	457.5	458.1, 440.2, 425.2	3.59
918	-H	-morpholino	-	441.5	442.2, 409.2	3.72
919	-H	-NHPh	O	463.5	464.1, 446.2, 431.2, 413.2	4.03
920	-H	-NHPh	-	447.5	448.0, 415.2	4.10
921	-H	-NH-2-pyridyl	-	448.5	449.0, 416.2	3.80
922	-H	-NHCH2-2-furyl	O	467.5	467.9, 450.0, 434.9, 417.0	3.95
923	-H	-NHCH2-2-furyl	-	451.5	452.0, 419.2	4.04
924	-H	-NH-3-isoxazolyl-5-Me	O	468.5	469.3, 451.0, 436.4	2.43
925	-H	-NH-3-isoxazolyl-5-Me	-	452.5	453.0, 420.2	2.56



926	-H	-NH-cyclopentyl	-	439.5	440.2, 407.2	3.97
927	-H	-NHCH ₂ CF ₃	O	469.4	470.1, 452.0, 437.1	2.43
928	-H	-NHCH ₂ CF ₃	-	453.4	454.2, 421.0	2.54
929	-H	-NHCH ₂ -cyclopropyl	O	441.5	442.4, 424.0, 409.4	2.39
930	-H	-NHCH ₂ -cyclopropyl	-	425.5	426.2, 393.2	2.50
931	-H	-NH(CH ₂) ₂ OH	O	431.5	432.2, 414.2, 399.0	1.89
932	-H	-NH(CH ₂) ₂ OH	-	415.5	416.2, 383.2	2.01
933	-H	-OH	-	372.4	373.5, 340.0	2.23
934	-H	-NH-(R)-2-butyl	O	443.5	443.9, 426.2, 410.9	2.46
935	-H	-NH-(R)-2-butyl	-	427.5	428.2, 395.2	2.59
936	-H	-NH-(S)-CH(Me)COOH	-	443.5	444.2, 411.2	2.16
937	-H	-NH-(S)-CH(iBu)COOH	-	485.6	486.2, 453.2	2.56

Example 938

Step 1: Methyl 7-acetoamino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate

Methyl 7-amino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate (63 mg, 0.17 mmol), was dissolved in DCM (2 ml). Triethylamine (24 ml, 0.17 mmol) was added followed by acetyl chloride (12 ml, 0.17 mmol), and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted in DCM (5 ml) and washed with saturated Na_2CO_3 (2 x 2 ml), water (2 ml) and brine (2 ml). The organic layer was concentrated *in vacuo* and the crude amide was used directly in the next step.

Step 2: [7-Acetoamino-2H-2-benzyl-3(4H)-oxo-1,4-benzothiazine-4-yl]-acetic acid N-hydroxyamide


Potassium hydroxide (109 mg, 1.93 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) were dissolved in methanol (0.53 ml and 0.9 ml respectively) at room temperature. Once a solution was obtained the potassium hydroxide solution was poured into the


hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for 1 hour. The precipitate (potassium chloride) was filtered off, the liquid added into the solution of the methyl ester obtained in step 1 in methanol (1 ml) and stirring was continued for an additional 3 hours. Acetic acid (0.5 ml) was added to the mixture and the solvent was evaporated to dryness. The residue was then re-dissolved in water and extracted with ethyl acetate (2 x 3ml). The combined organic extracts were evaporated to dryness, dissolved in DMSO and purified by reverse phase chromatography to yield the major product of the reaction together with small amount of the sulfoxide.

The following compounds listed in the Table 29 were prepared in a similar manner.

Table 29

Example No.	RR ¹	RR ²	RR ³	RR ⁴	mol. weight	Mass (m/e)	HPLC rt (min)
938	-OMe	-H	-NHCOCH ₃	O	431.5	431.9, 414.2, 399.0	2.60
939	-OMe	-NHCOEt	-H	-	429.5	430.2, 397.0	2.88
940	-H	-NHCOtBu	-H	-	427.5	428.2, 395.2	2.98
941	-H	-NHCOEt	-H	O	415.5	416.2, 398.2, 383.2	2.31
942	-H	-NHCO-2-pyrazinyl	-H	O	465.5	466.4, 448.2	2.40
943	-H	-NHCO-2-pyrazinyl	-H	-	449.5	450.2, 417.2	2.53
944	-H	-NHCOCH ₂ -C ₆ H ₄ -4-Cl	-H	-	496.0	496.2, 463.2	2.95
945	-H	-NHAc	-H	O	401.4	402.7, 384.2	2.17
946	-H	-NHAc	-H	-	385.4	386.2, 353.0	2.33
947	-H	-NHBz	-H	O	463.5	464.1, 446.2, 431.2	2.62
948	-H	-NHBz	-H	-	447.5	448.0, 415.2	2.75
949	-H	-NHCO-CH ₂ -2-thiophenyl	-H	-	467.6	468.2, 435.2	2.72
950	-H	-NHCOCH ₂ -C ₆ H ₄ -4-OMe	-H	-	491.6	492.2, 459.2	2.76
951	-H	-NHCO-2-furyl	-H	-	437.5	438.2, 405.2	2.60
952	-H	-NHCOCH ₂ Ph	-H	O	477.5	478.3, 460.2, 445.4, 427.2	2.65
953	-H	-NHCOCH ₂ Ph	-H	-	461.5	462.2, 429.2	2.77
954	-H	-NHCOEt	-H	-	399.5	400.2, 367.2	2.46
955	-H	-NHCO-4-pyridyl	-H	O	464.5	464.9, 447.2, 431.9	2.10
956	-OMe	-H	-NHCOtBu	-	457.6	458.2, 425.2	3.48
957	-OMe	-H	-NHCOCH ₃	-	415.5	416.2, 383.2	2.33
958	-OMe	-H	-NHCO-2-pyrazinyl	-	479.5	480.2, 447.2	2.57



Example 959

Step 1: Methyl 2H-2-(4-methoxybenzyl)-6-methylsulfonylamino-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate

Methyl 6-amino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate (63 mg, 0.16 mmol), was dissolved in DCM (2 ml) and pyridine (0.5 ml). Methyl sulfonylchloride (20 ml, 0.26 mmol) was added and the reaction mixture was stirred at room temperature for 3 hour. The reaction mixture was diluted in DCM (5 ml) and washed with 1N HCl (2 x 2 ml), water (2 ml) and brine (2 ml). The organic layer was concentrated *in vacuo* and the crude sulfonamide was used directly in the next step.

Step 2: 2H-2-(4-methoxybenzyl)-6-methylsulfonylamino-3(4H)-oxo-1,4-benzothiazine-4-yl-acetic acid N-hydroxyamide


Potassium hydroxide (109 mg, 1.93 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) were dissolved in methanol (0.53 ml and 0.9 ml respectively) at room temperature. Once a solution was obtained the potassium hydroxide solution was poured into the hydroxylamine solution at 0 °C and the mixture was


maintained at this temperature for 1 hour. The precipitate (potassium chloride) was filtered off, the liquid added into the solution of the methyl ester in methanol (1 ml) and stirring was continued for an additional 3 hours.

5 Acetic acid (0.5 ml) was added to the mixture and the solvent was evaporated to dryness. The residue was then dissolved in water and extracted with ethyl acetate (2 x 3ml). The combined organic extracts were evaporated to dryness, re-dissolved in DMSO and purified by reverse phase
10 chromatography.

The following compounds listed in the Table 30 were prepared in a similar manner.

Table 30

Example No.	RR ¹	RR ²	RR ³	RR ⁴	mol. weight	Mass (m/e)	HPLC rt (min)
							
959	-OMe	-NHMs	-H	-	451.5	452.0, 419.0	2.67
960	-OMe	-NHTs	-H	-	527.6	528.2, 495.2	3.09
961	-H	-NHSO2-C6H4-4-Cl	-H	O	534.0	534.5, 516.2, 501.5	2.81
962	-H	-NHSO2-4-isoxazolyl-3,5-Me2	-H	-	502.6	503.2, 470.2	2.48
963	-H	-NHMs	-H	-	421.5	422.2, 389.2	2.40
964	-H	-NHSO2Ph	-H	O	499.6	500.1, 482.2, 467.1	2.59
965	-H	-NHSO2Ph	-H	-	483.6	484.2, 451.0	2.93
966	-H	-NHSO2-C6H4-4-F	-H	-	501.6	502.2, 469.2	2.80
967	-H	-NHSO2-C6H4-4-OMe	-H	O	529.6	530.0, 512.2, 497.1	2.64
968	-H	-NHSO2-C6H4-4-OMe	-H	-	513.6	514.3, 481.2	2.76
969	-H	-NHSO2-2-thiophenyl	-H	O	505.6	506.1, 488.2, 473.1, 455.2	2.56
970	-H	-NHSO2iPr	-H	-	449.6	450.2, 417.2	2.83
971	-OMe	-H	-NHMs	-	451.5	452.1, 419.2	2.47

Example 972

Step 1: Methyl 2H-2-(4-methoxybenzyl)-7-(N'-methylureido)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate

5 **Methyl 7-amino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate**(63 mg, 0.16 mmol) was dissolved in DCM (2 ml) and pyridine (0.5 ml). Methyl isocyanate (15 mg, 0.26 mmol) was added and the reaction mixture was stirred at room temperature for 3 hour. The reaction
10 mixture was diluted in DCM (5 ml) and washed with saturated Na₂CO₃ (2 x 2 ml), water (2 ml) and brine (2 ml). The organic layer was concentrated *in vacuo* and the crude urea was carried directly to the next step.



15 **Step 2: 2H-2-(4-methoxybenzyl)-7-(N'-methylureido)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetic acid N-hydroxyamide**


20 Potassium hydroxide (109 mg, 1.93 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) were dissolved in methanol (0.53 ml and 0.9 ml respectively) at room temperature. Once a solution was obtained the potassium hydroxide solution was poured into the

hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for 1 hour. The precipitate (potassium chloride) was filtered off, the liquid added into the solution of the methyl ester, obtained in Step 1,
5 in methanol (1 ml) and stirring was continued for an additional 3 hours. Acetic acid (0.5 ml) was added to the mixture and the solvent was evaporated to dryness. The residue was then dissolved in water and extracted with ethyl acetate (2 x 3ml). The combined organic extracts were
10 evaporated to dryness, re-dissolved in DMSO and purified by reverse phase chromatography.

The following compounds listed in the Table 31 were prepared in a similar manner.

Table 31

Example No.	RR ¹	RR ²	RR ³	RR ⁴	mol. weight	Mass (m/e)	HPLC rt (min)
							
972	-OMe	-H	-NHCONHMe	-	430.5	431.2, 398.2	2.68
973	-OMe	-NHCONHPh	-H	-	493.5	494.8, 477.0, 461.9	2.53
974	-H	-NHCONH-C6H4-4-OMe	-H	O	508.6	509.0, 491.2, 476.1	2.61
975	-H	-NHCONH-C6H4-4-OMe	-H	-	492.6	493.4, 460.2	2.73
976	-H	-NHCONHCH2-C6H4-4-F	-H	O	510.5	511.3, 493.4, 478.3, 460.2	2.66
977	-H	-NHCONHCH2-C6H4-4-F	-H	-	494.5	495.4, 462.2	2.77
978	-H	-NHCONH-(R)-CH(Me)Ph	-H	-	490.6	491.2, 458.2	2.80
979	-H	-NHCONHBzl	-H	O	492.6	493.3, 475.2, 460.4, 442.2	2.61
980	-H	-NHCONHBzl	-H	-	476.6	477.2, 444.2	2.74
981	-H	-NHCONH-(R)-CH(Me)Ph	-H	O	506.6	507.5, 489.2, 474.6	2.68
982	-H	-NHCONHMe	-H	-	400.5	401.2, 368.2	2.32
983	-H		-H	O	518.6	519.5, 501.2	2.73
984	-OMe	-H	-NHCONHBu	O	488.6	489.6, 471.2, 456.6	3.25
985	-OMe	-H	-NHCONHBu	-	472.6	473.2, 440.4	2.82

Example 986

Step 1: Methyl 7-benzylamino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate

Methyl 7-amino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetate (63 mg, 0.17 mmol) and diisopropylethylamine (45.5 ml, 0.27 mmol) were dissolved in DCM (2 ml). Benzaldehyde (27 ml, 0.26 mmol) and sodium triacetoxyborohydride (58 mg, 0.26 mmol) were added to the solution resulting in a suspension that was stirred at room temperature for 3 hour. The reaction mixture was diluted in DCM (5 ml) and washed with saturated Na_2CO_3 (2 x 2 ml), water (2 ml) and brine (2 ml). The organic layer was concentrated *in vacuo* and the crude amine was used directly in the next step.


Step 2: 7-Benzylamino-2H-2-(4-methoxybenzyl)-3(4H)-oxo-1,4-benzothiazine-4-yl-acetic acid N-hydroxyamide

Potassium hydroxide (109 mg, 1.93 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) were dissolved in methanol (0.53 ml and 0.9 ml respectively) at

room temperature. Once a solution was obtained the potassium hydroxide solution was poured into the hydroxylamine solution at 0 °C and the mixture was maintained at this temperature for 1 hour. The precipitate (potassium chloride) was filtered off, the liquid added into the solution of the methyl ester, obtained in Step 1, in methanol (1 ml) and stirring was continued for an additional 3 hours. Acetic acid (0.5 ml) was added to the mixture and the solvent was evaporated to dryness. The residue was then dissolved in water and extracted with ethyl acetate (2 x 3ml). The combined organic extracts were evaporated to dryness, re-dissolved in DMSO and purified by reverse phase chromatography.

The following compounds listed in the Table 32 were prepared in a similar manner.

Table 32

Example No.		RR ¹	RR ²	RR ³	mol. weight	Mass (m/e)	HPLC rt (min)
							
986	-OMe	-H	-H	-NHBzl	463.6	464.2, 431.2	3.03
987	-H	-NHBzl	-H	-H	433.5	434.2, 401.2	2.90

The results of biological testing of some compounds are described below.

Biological Data

5 Metallo-proteinases used and inhibition assay protocols are described below.

MMP-2 and MMP-9

Both enzymes were commercially available as active forms (Yagai-Cosmobio) and were used for inhibition assays.

10

MMP-3 (Matrix metalloproteinase-3, Stromelysin-1)

C-terminal truncated form of human prostromelysin cDNA (proMMP-3, cDNA sequence in *Nature*, 348, 699-704 (1990)) was subcloned, expressed in *E.coli* and purified as described previously (*Biochemistry* 30, 6476-6483 (1991)). Activation of proMMP-3 was achieved by treatment with 1 mM 4-aminophenylmercuric acetate at 37°C for 60 min.

15

MMP-13 (Matrix metalloproteinase-13, Collagenase-3)

20 To subclone C-terminal truncated form of procollagenase-3 (proMMP-13) cDNA (J.Biol.Chem., 269(24), 16766-16773 (1994)) ,two synthetic oligonucleotide primer (5'-GGAATTCCATATGCTGCCGCTGCCGAGTGGTGGTGATGAAGATG-3' and 5'-TTTGGATCCTTAGCCGTACAGGCTTTGAATACCTTGTACATCGTCATCAGG - 3' :the former incorporates sequences for a unique *NdeI* site (underlined) including initial methionine, and the latter has sequences for a stop codon and *BamHI* site

25

(underlined)) were used with human chondrocyte cDNA library (CLONTECH) in PCR. With these primers and *Pfu* DNA polymerase (STRATAGENE) in PCR, 767 bp fragment was generated, which encodes 84 amino acid prosequence and 164 amino acids of mature MMP-13. The fragment was digested with *Nde*I and *Bam*HI, ligated into the *Nde*I and *Bam*HI sites of pET11a (STRATAGENE), transformed in *E.coli* BL21(DE3) and cultured. The crude cell extract was prepared as described in *Biochemistry*. The extract was dialyzed against 20 mM Tris-HCl(pH7.2)/5mM CaCl₂/0.02% NaN₃ and applied to a SP-Sepharose HP column (1.6 x 10 cm, Amersham-Pharmacia Biotech), and elution was performed with 50 mL linear gradient of 0 to 0.3 M NaCl (partially purified proMMP-13 was eluted at 0.2 M approximately). The fraction was dialyzed against 20 mM Tris-HCl(pH7.9)/5mM CaCl₂/200 mM (NH₄)₂SO₄/0.02% NaN₃ and applied to a Phenyl Sepharose HP column (1.6 x 5cm, Amersham-Pharmacia Biotech), and elution was performed with 20 mL linear gradient of 0.2 to 0 M (NH₄)₂SO₄ (pure proMMP-13 was eluted at 50 mM approximately). The fraction was concentrated by YM-5 ultrafiltration membrane and activated with 4-aminophenylmercuric acetate and active MMP-13 was separated from propeptide fragments by gel filtration chromatography as described in *Biochemistry*.

Inhibition assays against MMPs

Enzymatic assays were performed in accordance with C. G.

Knight's method (*FEBS Lett.*, 296(3), 263-266 (1992)). The fluorescent MCA-labeled substrate, (7-methoxycumaline-4-yl)-Pro-Leu-Gly-Leu-L-[N-(2,4-dinitrophenyl)-L-2,3-diaminopropionyl]-Ala-Arg-NH₂ (Peptide Institute. Inc.) was used for measuring the kinetics of inhibition for each tested compound.

The substrate was incubated at 37 °C for 1.5 hours (MMP-1; for 24 hours) with active MMP (at 20 nM, except MMP-13 at 5 nM) and a test compound in assay buffer containing 100 mM Tris-HCl (pH7.5), 10 mM CaCl₂, 100 mM NaCl and 0.05% Brij-35. After the mixture was plated at 100 µL/well onto a 96-well microtiter plate and incubated at 37°C, enzymatic activity in the presence of the compound was estimated by measuring the fluorescence intensity (λ_{ex} 320nm, λ_{em} 405nm) and IC₅₀ was determined.

Inhibition against MMP-3 and MMP-13

Table 10 **IC₅₀ (μM)**

Example No.	MMP3	MMP13
10	31.6	5.3
13	>50	13.2
19	1	>50
42	>10	0.759
46	0.894	0.134
71	9.071	0.249
88	3.798	0.062
92	0.794	0.048
105	6.717	0.231
109	0.599	0.043
121	2.547	0.128
130	1.141	0.056
143	1.268	0.064
175	1.140	0.112
176	1.057	0.100
195	1.892	0.119
211	0.895	0.108
218	0.422	0.056
239	0.274	0.035
255	1.203	0.106
257	0.267	0.023
262	0.850	0.066
283	0.614	0.037
306	0.987	0.122
316	0.442	0.026
324	0.733	0.063
329	0.155	0.011
330	0.745	0.066
333	0.794	0.070
361	0.446	0.053
368	1.5	0.036
370	1.2	0.035
371	0.88	0.036
378	>50	16
388	23	1.6
389	>50	29
406	3.228	0.257
407	0.433	0.024
409	0.279	0.012
419	0.514	0.063
422	>10	4.815
423	5.134	1.216
426	0.375	0.038
429	5.616	3.601
431	0.238	0.013
436	0.938	0.671
438	1.665	0.375
441	2.959	0.118

Table 10' **IC₅₀ (μM)**

Example No.	MMP3	MMP13
740	0.459	0.012
479	0.27	0.019
739	0.301	0.017
785	0.026	0.001
865	0.053	0.0049

Table 11

IC₅₀ (μM)

Example No.	MMP-2	MMP-9
21	1.0	>1
365	0.32	0.11
368	0.30	<0.01
369	>1	0.20

Inhibition ELISA for Tumor Necrosis Factor-α (TNF-α) release

5 RAW 267.4 (derived from mouse monocytes) was seeded in
RPMI1640 containing 10% of FBS (GIBCO BRL) at 100 μL/well
(5 × 10⁵ cells/mL) onto 96-well microplates and cultured at
37°C, 5% CO₂. After culturing for 4 hours, each test
compound diluted in RPMI1640 was added at 50 μL/well. 15
10 minutes later, 4 μg/μL LPS (DIFCO) diluted in RPMI1640 was
added at 50 μL/well and cultured for 24 hours.


TNF-α produced was measured by ELISA as usual. (For
example; J. Cellular Physiol. 154, 479 (1993)) Absorbance
at 450 nm (normalized at 620 nm) was measured and
15 percentage of the amount of TNF-α release for each tested
compound (n=2) was determined (0% for untreated control
(n=4), and 100% for LPS stimulation (n=4)).

Table 12
Inhibition of TNF production

Example No.	at 5 μM (%)
365	20
367	68
368	30
387	68


CLAIMS

1. A hydroxamic acid derivative represented by the formula [1], or a pharmaceutically acceptable salt thereof:



wherein

X is one of the groups represented by the formulae [1'] and [1'']:



wherein Ar¹ is optionally substituted aryl or optionally substituted heteroaryl;

A¹ is *ortho*-phenylene or monocyclic *ortho*-heteroarylene;

R⁸ and R⁹ are independently hydrogen or substituent of the *ortho*-arylene and the *ortho*-heteroarylene;

Y¹ is -O-, -S-, -S(O)- or -S(O)₂-;

Y² is O, or S;

R¹ is -(CHR⁴)_n-(CR⁵R⁶)-CO-NHOH;

R³ is hydrogen, optionally substituted alkyl, or optionally substituted cycloalkyl;

R^2 is hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted hetero-cycloalkyl; or R^2 and R^3 may be taken together to be optionally substituted C_1 - C_{10} alkylidene;

when X is represented by formula [1'],

R^4 and R^6 are independently hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl, and R^5 is optionally substituted alkyl;

when X is represented by formula [1''],

R^4 , R^5 and R^6 are independently hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted cycloalkyl, optionally substituted heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl; or

R^5 may be joined with R^4 or R^6 to form, with the carbon atom which they attach, optionally substituted cycloalkane or optionally substituted heterocycloalkane; and

n is an integer of 0 to 4;

provided that when R^2 and R^3 are taken together to be optionally substituted C_1 - C_{10} alkylidene, X is not methylene substituted by a phenyl or a pyridyl wherein the phenyl and the pyridyl are optionally substituted by methyl or methoxy;

said cycloalkyl is C3-8 cycloalkyl containing 0-2 carbonyl group;

said cycloalkenyl is C3-8 cycloalkenyl containing 0-2 carbonyl group;

said heterocycloalkyl is 5- or 6-membered heterocycloalkyl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms, and oxygen atoms, wherein said sulfur atoms may be oxidized to form sulfoxide or sulfone;

said heterocycloalkenyl is 5- or 6-membered heterocycloalkenyl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms, and oxygen atoms, wherein said sulfur atoms may be oxidized to form sulfoxide or sulfone;

said aryl is C6-10 aryl;

said *ortho*-arylene is C6-10 *ortho*-arylene;

said heteroaryl is mono or bicyclic 5- or 6-membered heteroaryl containing 1 to 5 atoms selected independently from nitrogen atoms, sulfur atoms, and oxygen atoms;

said *ortho*-heteroarylene is mono or bicyclic 5- or 6-membered *ortho*-heteroarylene containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms, and oxygen atoms;

said cycloalkane is C3-8 cycloalkane containing 0-2 carbonyl group;

said heterocycloalkane is 5- or 6-membered heterocycloalkane containing 1 to 3 atoms selected independently from nitrogen atoms, sulfur atoms, and oxygen atoms, wherein said sulfur atoms may be oxidized to form sulfoxide or sulfone;

a substituent of said substituted aryl and said substituted heteroaryl, and a substituent represented by R⁸ and R⁹ are selected from the group consisting of

halogen; alkyl; alkenyl; alkynyl; halogenated alkyl; -Z¹-Ar³; -Z¹-Cy¹; -Z¹-N(Q²)Q¹;
-Z¹-N(Q²)Ar³; -Z¹-N(Q²)Cy¹; -Z¹-NQ⁴-C(NQ³)N(Q²)Q¹; Z¹-NQ⁴-C(NQ³)N(Q²)Ar³;
-Z¹-NQ⁴-C(NQ³)N(Q²)Cy¹; -Z¹-NQ³-CON(Q²)Q¹; -Z¹-NQ³-CON(Q²)Ar³;

$-Z^1-NQ^3-CON(Q^2)Cy^1$; $-Z^1-NQ^2-COOQ^1$; $-Z^1-NQ^2-COOAr^3$; $-Z^1-NQ^2-COOCy^1$;
 $-Z^1-OCON(Q^1)Q^2$; $-Z^1-OCON(Q^2)Ar^3$; $-Z^1-OCON(Q^2)Cy^1$; $-Z^1-OCOOQ^1$;
 $-Z^1-OCOOAr^3$; $-Z^1-OCOOCy^1$; $-Z^1-NQ^2-COQ^1$; $-Z^1-NQ^2-COAr^3$;
 $-Z^1-NQ^2-COCy^1$; $-Z^1-NQ^2-SOQ^1$; $-Z^1-NQ^2-SOAr^3$; $-Z^1-NQ^2-SOCy^1$;
 $-Z^1-NQ^2-SO_2Q^1$; $-Z^1-NQ^2-SO_2Ar^3$; $-Z^1-NQ^2-SO_2Cy^1$; $-Z^1-OQ^1$; $-Z^1-OAr^3$;
 $-Z^1-O-Cy^1$; $-Z^1-OCOQ^1$; $-Z^1-OCOAr^3$; $-Z^1-OCOCy^1$; $-Z^1-COOQ^1$; $-Z^1-COOAr^3$;
 $-Z^1-COOCy^1$; $-Z^1-CON(Q^1)Q^2$; $-Z^1-CON(Q^2)Ar^3$; $-Z^1-CON(Q^2)Cy^1$;
 $-Z^1-CON(Q^2)OQ^1$; $-Z^1-CON(Q^2)OAr^3$; $-Z^1-CON(Q^2)OCy^1$; $-Z^1-COQ^1$; $-Z^1-COAr^3$;
 $-Z^1-CO-Cy^1$; $-Z^1-C(NQ^3)N(Q^1)Q^2$; $-C(NQ^3)N(Q^2)Ar^3$; $-Z^1-C(NQ^3)N(Q^2)Cy^1$;
 $-Z^1-SQ^1$; $-Z^1-SAr^3$; $-Z^1-S-Cy^1$; $-Z^1-SOQ^1$; $-Z^1-SOAr^3$; $-Z^1-SOCy^1$; $-Z^1-SO_2Q^1$;
 $-Z^1-SO_2Ar^3$; $-Z^1-SO_2Cy^1$; $-Z^1-SON(Q^1)Q^2$; $-Z^1-SON(Q^2)Ar^3$; $-Z^1-SON(Q^2)Cy^1$;
 $-Z^1-SO_2N(Q^1)Q^2$; $-Z^1-SO_2N(Q^2)Ar^3$; $-Z^1-SO_2N(Q^2)Cy^1$; $-Z^1-SO_3H$; $-Z^1-OSO_3H$;
 $-Z^1-NO_2$; $-Z^1-CN$; and $-CHO$;

wherein

Z^1 is a single bond, alkylene, alkenylene or alkynylene;

Q^1 , Q^2 , Q^3 , Q^4 are independently hydrogen, alkyl, alkenyl, or alkynyl, wherein the alkyl, the alkenyl and the alkynyl are optionally substituted by one or more substituents selected from the group consisting of

halogen; $-Ar^4$; $-Cy^2$; $-N(R^{21})R^{22}$; $-N(R^{22})Ar^4$; $-N(R^{22})Cy^2$; $-NR^{24}-C(NR^{23})N(R^{22})R^{21}$;
 $-NR^{24}-C(NR^{23})N(R^{22})Ar^4$; $-NR^{24}-C(NR^{23})N(R^{22})Cy^2$; $-NR^{23}-CON(R^{22})R^{21}$;
 $-NR^{23}-CON(R^{22})Ar^4$; $-NR^{23}-CON(R^{22})Cy^2$; $-NR^{22}-COOR^{21}$; $-NR^{22}-COOAr^4$;
 $-NR^{22}-COOCy^2$; $-OCON(R^{22})R^{21}$; $-OCON(R^{22})Ar^4$; $-OCON(R^{22})Cy^2$; $-OCOOR^{21}$;
 $-OCOOAr^4$; $-OCOOCy^2$; $-NR^{22}-COR^{21}$; $-NR^{22}-COAr^4$; $-NR^{22}-COCy^2$;
 $-NR^{22}-SOR^{21}$; $-NR^{22}-SOAr^4$; $-NR^{22}-SOCy^2$; $-NR^{22}-SO_2R^{21}$; $-NR^{22}-SO_2Ar^4$;
 $-NR^{22}-SO_2Cy^2$; $-OR^{21}$; $-OAr^4$; $-OCy^2$; $-OCOR^{21}$; $-OCOAr^4$; $-OCOCy^2$; $-COOR^{21}$;
 $-COOAr^4$; $-COOCy^2$; $-CON(R^{21})R^{22}$; $-CON(R^{22})Ar^4$; $-CON(R^{22})Cy^2$;
 $-CON(R^{22})OR^{21}$; $-CON(R^{22})OAr^4$; $-CON(R^{22})OCy^2$; $-COR^{21}$; $-COAr^4$; $-COCy^2$;
 $-SR^{21}$; $-SAr^4$; $-SCy^2$; $-SOR^{21}$; $-SOAr^4$; $-SOCy^2$; $-SO_2R^{21}$; $-SO_2Ar^4$; $-SO_2Cy^2$;
 $-SON(R^{22})R^{21}$; $-SON(R^{22})Ar^4$; $-SON(R^{22})Cy^2$; $-SO_2N(R^{22})R^{21}$; $-SO_2N(R^{22})Ar^4$;

$-\text{SO}_2\text{N}(\text{R}^{22})\text{Cy}^2$; $-\text{SO}_3\text{H}$; $-\text{OSO}_3\text{H}$; $-\text{NO}_2$; $-\text{CN}$; and $-\text{CHO}$;

or Q^1 may be jointed with Q^2 or Q^3 to form optionally substituted heterocycloalkane, with the carbon atom which they attach;

Ar^3 and Ar^4 are independently phenyl or heteroaryl, wherein the phenyl and the heteroaryl are optionally substituted by one or two substituents selected from the group consisting of halogen; alkyl; alkenyl; alkynyl; halogenated alkyl; $-\text{Z}^2-\text{N}(\text{R}^{25})\text{R}^{26}$; $-\text{Z}^2-\text{NR}^{28}-\text{C}(\text{NR}^{27})\text{N}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{NR}^{27}-\text{CON}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{NR}^{26}-\text{COOR}^{25}$; $-\text{Z}^2-\text{OCON}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{NR}^{26}-\text{COR}^{25}$; $-\text{Z}^2-\text{NR}^{26}-\text{SOR}^{25}$; $-\text{Z}^2-\text{NR}^{26}-\text{SO}_2\text{R}^{25}$; $-\text{Z}^2-\text{OR}^{25}$; $-\text{Z}^2-\text{COOR}^{25}$; $-\text{Z}^2-\text{OCOR}^{25}$; $-\text{Z}^2-\text{OCOOR}^{25}$; $-\text{Z}^2-\text{CON}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{CON}(\text{R}^{26})\text{OR}^{25}$; $-\text{Z}^2-\text{COR}^{25}$; $-\text{Z}^2-\text{C}(\text{NR}^{27})\text{N}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{SR}^{25}$; $-\text{Z}^2-\text{SOR}^{25}$; $-\text{Z}^2-\text{SO}_2\text{R}^{25}$; $-\text{Z}^2-\text{SON}(\text{R}^{25})\text{R}^{26}$; $-\text{Z}^2-\text{SO}_2\text{N}(\text{R}^{26})\text{R}^{25}$; $-\text{Z}^2-\text{SO}_3\text{H}$; $-\text{Z}^2-\text{OSO}_3\text{H}$; $-\text{Z}^2-\text{NO}_2$; $-\text{Z}^2-\text{CN}$; $-\text{CHO}-\text{O}-\text{CH}_2-\text{O}-$; $-\text{O}-\text{CH}_2-\text{CH}_2-\text{O}-$; $-\text{O}-\text{CH}_2-\text{CH}_2-$; $-\text{CH}_2-\text{CH}_2-\text{O}-\text{CO}-$; and $-\text{CH}_2-\text{O}-\text{CO}-$;

wherein

Z^2 is a single bond, alkylene, alkenylene or alkynylene;

R^{21} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , and R^{28} are independently hydrogen, alkyl, alkenyl, alkynyl, wherein R^{21} may be jointed with R^{22} or R^{23} to form heterocycloalkane with the carbon atom which they attach, or R^{25} may be jointed with R^{26} or R^{27} to form heterocycloalkane with the carbon atom which they attach;

Cy^1 and Cy^2 are independently cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, wherein the cycloalkyl, the cycloalkenyl, the heterocycloalkyl, and the heterocycloalkenyl are optionally substituted by one or two substituents which are selected from the same group as the substituents of Ar^3 and Ar^4 ;

a substituent of said substituted alkyl, said substituted alkenyl, said substituted alkynyl, said substituted alkoxy, said substituted alkylthio, said substituted cycloalkyl, said substituted heterocycloalkyl, and said substituted $\text{C}_1\text{-C}_{10}$ alkylidene is selected from the

group consisting of

halogen; $-\text{Ar}^5$; $-\text{Cy}^3$; $-\text{N}(\text{Q}^5)\text{Q}^6$; $-\text{N}(\text{Q}^6)\text{Ar}^5$; $-\text{N}(\text{Q}^6)\text{Cy}^3$; $-\text{NQ}^8-\text{C}(\text{NQ}^7)\text{N}(\text{Q}^6)\text{Q}^5$;
 $-\text{NQ}^8-\text{C}(\text{NQ}^7)\text{N}(\text{Q}^6)\text{Ar}^5$; $-\text{NQ}^8-\text{C}(\text{NQ}^7)\text{N}(\text{Q}^6)\text{Cy}^3$; $-\text{NQ}^7-\text{CON}(\text{Q}^6)\text{Q}^5$;
 $-\text{NQ}^6-\text{CON}(\text{Q}^5)\text{Ar}^5$; $-\text{NQ}^7-\text{CON}(\text{Q}^6)\text{Cy}^3$; $-\text{NQ}^7-\text{COOQ}^6$; $-\text{NQ}^7-\text{COOAr}^5$;
 $-\text{NQ}^7-\text{COOCy}^3$; $-\text{OCON}(\text{Q}^6)\text{Q}^5$; $-\text{OCON}(\text{Q}^6)\text{Ar}^5$; $-\text{OCON}(\text{Q}^6)\text{Cy}^3$; $-\text{NQ}^6-\text{COQ}^5$;
 $-\text{NQ}^6-\text{COAr}^5$; $-\text{NQ}^6-\text{COCy}^3$; $-\text{NQ}^6-\text{SOQ}^5$; $-\text{NQ}^6-\text{SOAr}^5$; $-\text{NQ}^6-\text{SOCy}^3$;
 $-\text{NQ}^6-\text{SO}_2\text{Q}^5$; $-\text{NQ}^6-\text{SO}_2\text{Ar}^5$; $-\text{NQ}^6-\text{SO}_2\text{Cy}^3$; $-\text{OQ}^5$; $-\text{OAr}^5$; $-\text{OCy}^3$; $-\text{OCOQ}^5$;
 $-\text{OCOAr}^5$; $-\text{OCOCy}^3$; $-\text{COOQ}^5$; $-\text{COOAr}^5$; $-\text{COOCy}^3$; $-\text{OCOOQ}^5$; $-\text{OCOOAr}^5$;
 $-\text{OCOOCy}^3$; $-\text{CON}(\text{Q}^6)\text{Q}^5$; $-\text{CON}(\text{Q}^6)\text{Ar}^5$; $-\text{CON}(\text{Q}^6)\text{Cy}^3$; $-\text{CON}(\text{Q}^6)\text{OQ}^5$;
 $-\text{CON}(\text{Q}^6)\text{OAr}^5$; $-\text{CON}(\text{Q}^6)\text{OCy}^3$; $-\text{COQ}^5$; $-\text{COAr}^5$; $-\text{COCy}^3$; $-\text{SQ}^5$; $-\text{SAr}^5$; $-\text{SCy}^3$;
 $-\text{SOQ}^5$; $-\text{SOAr}^5$; $-\text{SOCy}^3$; $-\text{SO}_2\text{Q}^5$; $-\text{SO}_2\text{Ar}^5$; $-\text{SO}_2\text{Cy}^3$; $-\text{SON}(\text{Q}^6)\text{Q}^5$; $-\text{SON}(\text{Q}^6)\text{Ar}^5$;
 $-\text{SON}(\text{Q}^6)\text{Cy}^3$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Q}^5$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Ar}^5$; $-\text{SO}_2\text{N}(\text{Q}^6)\text{Cy}^3$; $-\text{SO}_3\text{H}$; $-\text{NO}_2$; and
 $-\text{CN}$;

wherein

Q^5 , Q^6 , Q^7 , and Q^8 are independently hydrogen, alkyl, alkenyl and alkynyl wherein the alkyl, the alkenyl and the alkynyl are optionally substituted by one or more substituents selected from the group consisting of

halogen; $-\text{Ar}^6$; $-\text{Cy}^4$; $-\text{N}(\text{R}^{29})\text{R}^{30}$; $-\text{N}(\text{R}^{30})\text{Ar}^6$; $-\text{N}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{R}^{29}$;
 $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{Ar}^6$; $-\text{NR}^{32}-\text{C}(\text{NR}^{31})\text{N}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{R}^{29}$;
 $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{Ar}^6$; $-\text{NR}^{31}-\text{CON}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{30}-\text{COOR}^{29}$; $-\text{NR}^{30}-\text{COOAr}^6$;
 $-\text{NR}^{30}-\text{COOCy}^4$; $-\text{OCON}(\text{R}^{30})\text{R}^{29}$; $-\text{OCON}(\text{R}^{30})\text{Ar}^6$; $-\text{OCON}(\text{R}^{30})\text{Cy}^4$; $-\text{NR}^{30}-\text{COR}^{29}$;
 $-\text{NR}^{30}-\text{COAr}^6$; $-\text{NR}^{30}-\text{COCy}^4$; $-\text{NR}^{30}-\text{SOR}^{29}$; $-\text{NR}^{30}-\text{SOAr}^6$; $-\text{NR}^{30}-\text{SOCy}^4$;
 $-\text{NR}^{30}-\text{SO}_2\text{R}^{29}$; $-\text{NR}^{30}-\text{SO}_2\text{Ar}^6$; $-\text{NR}^{30}-\text{SO}_2\text{Cy}^4$; $-\text{OR}^{29}$; $-\text{OAr}^6$; $-\text{OCy}^4$; $-\text{COOR}^{29}$;
 $-\text{COOAr}^6$; $-\text{COOCy}^4$; $-\text{OCOR}^{29}$; $-\text{COAr}^5$; $-\text{OCOCy}^3$; $-\text{OCOOAr}^{29}$; $-\text{OCOOAr}^6$;
 $-\text{Z}^1-\text{OCOOCy}^4$; $-\text{CON}(\text{R}^{30})\text{R}^{29}$; $-\text{CON}(\text{R}^{30})\text{Ar}^6$; $-\text{CON}(\text{R}^{30})\text{Cy}^4$; $-\text{CON}(\text{R}^{30})\text{OR}^{29}$;
 $-\text{CON}(\text{R}^{30})\text{OAr}^6$; $-\text{CON}(\text{R}^{30})\text{OCy}^4$; $-\text{COR}^{29}$; $-\text{COAr}^6$; $-\text{COCy}^4$; $-\text{SR}^{29}$; $-\text{SAr}^6$; $-\text{SCy}^4$;
 $-\text{SOR}^{29}$; $-\text{SOAr}^6$; $-\text{SOCy}^4$; $-\text{SO}_2\text{R}^{29}$; $-\text{SO}_2\text{Ar}^6$; $-\text{SO}_2\text{Cy}^4$; $-\text{SON}(\text{R}^{30})\text{R}^{29}$;
 $-\text{SON}(\text{R}^{30})\text{Ar}^6$; $-\text{SON}(\text{R}^{30})\text{Cy}^4$; $-\text{SO}_2\text{N}(\text{R}^{30})\text{R}^{29}$; $-\text{SO}_2\text{N}(\text{R}^{30})\text{Ar}^6$; $-\text{SO}_2\text{N}(\text{R}^{30})\text{Cy}^4$;
 $-\text{SO}_3\text{H}$; $-\text{NO}_2$; and $-\text{CN}$;

or Q^5 may be jointed with Q^6 or Q^7 to form heterocycloalkane with the carbon atom

which they attach, wherein

R^{29} , R^{30} , R^{31} , and R^{32} are independently hydrogen, alkyl, alkenyl, or alkynyl;

Ar^5 and Ar^6 are independently optionally substituted aryl or optionally substituted heteroaryl;

Cy^3 and Cy^4 are independently cycloalkyl, cycloalkenyl, heterocycloalkyl, or heterocycloalkenyl, wherein the cycloalkyl, the cycloalkenyl, the heterocycloalkyl, and the heterocycloalkenyl are optionally substituted by one or two substituents which are the same as the substituent of Ar^3 or Ar^4 ;

a substituent of said substituted cycloalkane and said substituted heterocycloalkane is selected from the group consisting of


halogen; $-Ar^5$; $-Cy^3$; $-N(Q^{5'})Q^{6'}$; $-N(Q^{6'})Ar^5$; $-N(Q^{6'})Cy^3$; $-NQ^{8'}-C(NQ^{7'})N(Q^{6'})Q^{5'}$; $-NQ^{8'}-C(NQ^{7'})N(Q^{6'})Ar^5$; $-NQ^{8'}-C(NQ^{7'})N(Q^{6'})Cy^3$; $-NQ^{7'}-CON(Q^{6'})Q^{5'}$; $-NQ^{6'}-CON(Q^{5'})Ar^5$; $-NQ^{7'}-CON(Q^{6'})Cy^3$; $-NQ^{7'}-COOQ^{6'}$; $-NQ^{7'}-COOAr^5$; $-NQ^{7'}-COOCy^3$; $-OCON(Q^{6'})Q^{5'}$; $-OCON(Q^{6'})Ar^5$; $-OCON(Q^{6'})Cy^3$; $-NQ^{6'}-COQ^{5'}$; $-NQ^{6'}-COAr^5$; $-NQ^{6'}-COCy^3$; $-NQ^{6'}-SOQ^{5'}$; $-NQ^{6'}-SOAr^5$; $-NQ^{6'}-SOCy^3$; $-NQ^{6'}-SO_2Q^{5'}$; $-NQ^{6'}-SO_2Ar^5$; $-NQ^{6'}-SO_2Cy^3$; $-OQ^{5'}$; $-OAr^5$; $-OCy^3$; $-OCOQ^{5'}$; $-OCOAr^5$; $-OCOCy^3$; $-COOQ^{5'}$; $-COOAr^5$; $-COOCy^3$; $-OCOOQ^{5'}$; $-OCOOAr^5$; $-OCOOQ^{5'}$; $-CON(Q^{6'})Q^{5'}$; $-CON(Q^{6'})Ar^5$; $-CON(Q^{6'})Cy^3$; $-CON(Q^{6'})OQ^{5'}$; $-CON(Q^{6'})OAr^5$; $-CON(Q^{6'})OCy^3$; $-COQ^{5'}$; $-COAr^5$; $-COCy^3$; $-SQ^{5'}$; $-SAr^5$; $-SCy^3$; $-SOQ^{5'}$; $-SOAr^5$; $-SOCy^3$; $-SO_2Q^{5'}$; $-SO_2Ar^5$; $-SO_2Cy^3$; $-SON(Q^{6'})Q^{5'}$; $-SON(Q^{6'})Ar^5$; $-SON(Q^{6'})Cy^3$; $-SO_2N(Q^{6'})Q^{5'}$; $-SO_2N(Q^{6'})Ar^5$; $-SO_2N(Q^{6'})Cy^3$; $-SO_3H$; $-NO_2$; and $-CN$;

wherein

$Q^{5'}$, $Q^{6'}$, $Q^{7'}$, and $Q^{8'}$ are independently hydrogen, alkyl, alkenyl or alkynyl; and

Ar^5 , Ar^6 , Cy^3 , and Cy^4 are the same as defined above.

2. A hydroxamic acid derivative according to claim 1, or a pharmaceutically acceptable salt thereof, wherein Y^1 and -S-, -S(O)- or -S(O)₂-.
3. A hydroxamic acid derivative according to claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein n is 0, 1 or 2.
4. A hydroxamic acid derivative according to any one of claims 1-3, or a pharmaceutically acceptable salt thereof, wherein n is 0.
5. A hydroxamic acid derivative according to claim 1 represented by the formula [2], or a pharmaceutically acceptable salt thereof:



wherein

Y^2 , R^4 , R^5 and R^6 are as defined in claim 1;

Ar^2 is optionally substituted phenyl; optionally substituted naphthyl; or optionally substituted mono or bicyclic heteroaryl;


Y^3 is -S-, -S(O)- or -S(O)₂-;

R^{10} and R^{11} are independently hydrogen or optionally substituted alkyl; and

$n1$ is an integer of 0, 1 or 2.

6. A hydroxamic acid derivative according to claim 5, or a pharmaceutically acceptable salt thereof, wherein $n1$ is 0.

7. A hydroxamic acid derivative according to claim 1 represented by the formula [3], or a pharmaceutically acceptable salt thereof:



wherein

Y^2 , R^4 , R^5 , R^6 , A^1 , R^8 , and R^9 are as defined in claim 1;

R^{12} and R^{13} are independently hydrogen or optionally substituted alkyl;

Y^3 is -S-, -S(O)- or -S(O)₂-; and

n_2 is an integer of 0, 1 or 2.

8. A hydroxamic acid derivative according to claim 7, or a pharmaceutically acceptable salt thereof, wherein R^8 is hydrogen or substituent; and R^9 is substituent.


9. A hydroxamic acid derivative according to claim 7 or 8, or a pharmaceutically acceptable salt thereof, wherein n_2 is 0, and R^{12} and R^{13} are independently hydrogen or substituted alkyl.

10. A hydroxamic acid derivative according to any one of claims 1-9, or a pharmaceutically acceptable salt thereof, wherein Y^2 is -O-.

11. A pharmaceutical composition containing a hydroxamic acid derivative according to any one of claims 1-10, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

12. Use of a hydroxamic acid derivative according to any one of claims 1-10, or a pharmaceutically acceptable salt thereof in the preparation of a medicament for inhibiting matrix metallo-proteinases.

13. Use according to claim 12, wherein the matrix metallo-proteinases are matrix metallo-proteinase 3 and/or 13.
14. Use of a hydroxamic acid derivative according to any one of claims 1-10, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a disease associated with excess or undesired matrix metallo-proteinases.
15. Use according to claim 14, wherein the matrix metallo-proteinases are matrix metallo-proteinase 3 and/or 13.



[1]