



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/195, C12N 15/31, 1/21, 5/10, A01H 5/00, 5/10, C12N 15/82	A2	(11) International Publication Number: WO 00/20452 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23181 (22) International Filing Date: 5 October 1999 (05.10.99) (30) Priority Data: 60/103,050 5 October 1998 (05.10.98) US (71) Applicant: EDEN BIOSCIENCE CORPORATION [US/US]; 11816 North Creek Parkway N., Bothell, WA 98011-8205 (US). (72) Inventors: WEI, Zhong-Min; 8230 125th Court, Kirkland, WA 98034 (US). FAN, Hao; 19712 6th Drive S.E., Bothell, WA 98012 (US). NIGGEMEYER, Jennifer, L.; 21315 2nd Avenue S.E., Bothell, WA 98021 (US). (74) Agents: GOLDMAN, Michael, L. et al.; Nixon Peabody LLP, Clinton Square, P.O. Box 1051, Rochester, NY 14603 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HYPERSENSITIVE RESPONSE ELICITOR FRAGMENTS WHICH ARE ACTIVE BUT DO NOT ELICIT A HYPERSENSITIVE RESPONSE (57) Abstract The present invention is directed to isolated active fragments of a hypersensitive response elicitor protein or polypeptide which fragment does not elicit a hypersensitive response in plants. Also disclosed are isolated DNA molecules which encode such fragments. Isolated fragments of hypersensitive response elicitor proteins or polypeptides in accordance with the present invention and the isolated DNA molecules that encode them have the following activities: imparting disease resistance to plants, enhancing plant growth, and/or controlling insects on plants. This can be achieved by applying the fragments of a hypersensitive response elicitor in a non-infectious form to plants or plant seeds under conditions effective to impart disease resistance, to enhance plant growth, and/or to control insects on plants or plants grown from the plant seeds. Alternatively, transgenic plants or plant seeds transformed with a DNA molecule encoding the fragment can be provided and the transgenic plants or plants resulting from the transgenic plant seeds are grown under conditions effective to impart disease resistance, to enhance plant growth, and/or to control insects on plants or plants grown from the plant seeds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HYPERSENSITIVE RESPONSE ELICITOR FRAGMENTS WHICH ARE ACTIVE BUT DO NOT ELICIT A HYPERSENSITIVE RESPONSE

This application claims benefit of U.S. Provisional Patent Application
5 Serial No. 60/103,050, filed October 5, 1998.

FIELD OF THE INVENTION

The present invention relates to active fragments of a hypersensitive
10 response elicitor which fragments do not elicit a hypersensitive response.

BACKGROUND OF THE INVENTION

Interactions between bacterial pathogens and their plant hosts generally
15 fall into two categories: (1) compatible (pathogen-host), leading to intercellular
bacterial growth, symptom development, and disease development in the host plant;
and (2) incompatible (pathogen-nonhost), resulting in the hypersensitive response, a
particular type of incompatible interaction occurring, without progressive disease
symptoms. During compatible interactions on host plants, bacterial populations
20 increase dramatically and progressive symptoms occur. During incompatible
interactions, bacterial populations do not increase, and progressive symptoms do not
occur.

The hypersensitive response is a rapid, localized necrosis that is
associated with the active defense of plants against many pathogens (Kiraly, Z.,
25 "Defenses Triggered by the Invader: Hypersensitivity," pages 201-224 in: Plant
Disease: An Advanced Treatise, Vol. 5, J.G. Horsfall and E.B. Cowling, ed.
Academic Press New York (1980); Klement, Z., "Hypersensitivity," pages 149-177
in: Phytopathogenic Prokaryotes, Vol. 2, M.S. Mount and G.H. Lacy, ed. Academic
Press, New York (1982)). The hypersensitive response elicited by bacteria is readily
30 observed as a tissue collapse if high concentrations ($\geq 10^7$ cells/ml) of a limited
host-range pathogen like *Pseudomonas syringae* or *Erwinia amylovora* are infiltrated
into the leaves of nonhost plants (necrosis occurs only in isolated plant cells at lower
levels of inoculum) (Klement, Z., "Rapid Detection of Pathogenicity of
Phytopathogenic Pseudomonads," Nature 199:299-300; Klement, et al.,

- 2 -

“Hypersensitive Reaction Induced by Phytopathogenic Bacteria in the Tobacco Leaf,” Phytopathology 54:474-477 (1963); Turner, et al., “The Quantitative Relation Between Plant and Bacterial Cells Involved in the Hypersensitive Reaction,” Phytopathology 64:885-890 (1974); Klement, Z., “Hypersensitivity,” pages 149-177 in Phytopathogenic Prokaryotes, Vol. 2., M.S. Mount and G.H. Lacy, ed. Academic Press, New York (1982)). The capacities to elicit the hypersensitive response in a nonhost and be pathogenic in a host appear linked. As noted by Klement, Z., “Hypersensitivity,” pages 149-177 in Phytopathogenic Prokaryotes, Vol. 2., M.S. Mount and G.H. Lacy, ed. Academic Press, New York, these pathogens also cause physiologically similar, albeit delayed, necroses in their interactions with compatible hosts. Furthermore, the ability to produce the hypersensitive response or pathogenesis is dependent on a common set of genes, denoted *hrp* (Lindgren, P.B., et al., “Gene Cluster of *Pseudomonas syringae* pv. ‘phaseolicola’ Controls Pathogenicity of Bean Plants and Hypersensitivity on Nonhost Plants,” J. Bacteriol. 168:512-22 (1986); Willis, D.K., et al., “*hrp* Genes of Phytopathogenic Bacteria,” Mol. Plant-Microbe Interact. 4:132-138 (1991)). Consequently, the hypersensitive response may hold clues to both the nature of plant defense and the basis for bacterial pathogenicity.

The *hrp* genes are widespread in Gram-negative plant pathogens, where they are clustered, conserved, and in some cases interchangeable (Willis, D.K., et al., “*hrp* Genes of Phytopathogenic Bacteria,” Mol. Plant-Microbe Interact. 4:132-138 (1991); Bonas, U., “*hrp* Genes of Phytopathogenic Bacteria,” pages 79-98 in: Current Topics in Microbiology and Immunology: Bacterial Pathogenesis of Plants and Animals - Molecular and Cellular Mechanisms, J.L. Dangl, ed. Springer-Verlag, Berlin (1994)). Several *hrp* genes encode components of a protein secretion pathway similar to one used by *Yersinia*, *Shigella*, and *Salmonella* spp. to secrete proteins essential in animal diseases (Van Gijsegem, et al., “Evolutionary Conservation of Pathogenicity Determinants Among Plant and Animal Pathogenic Bacteria,” Trends Microbiol. 1:175-180 (1993)). In *E. amylovora*, *P. syringae*, and *P. solanacearum*, *hrp* genes have been shown to control the production and secretion of glycine-rich, protein elicitors of the hypersensitive response (He, S.Y., et al. “*Pseudomonas Syringae* pv. *Syringae* HarpinPss: a Protein that is Secreted via the Hrp Pathway and Elicits the Hypersensitive Response in Plants,” Cell 73:1255-1266 (1993), Wei, Z.-H.,

- 3 -

et al., "HrpI of *Erwinia amylovora* Functions in Secretion of Harpin and is a Member of a New Protein Family," J. Bacteriol. 175:7958-7967 (1993); Arlat, M. et al.

"PopA1, a Protein Which Induces a Hypersensitive-like Response on Specific Petunia Genotypes, is Secreted via the Hrp Pathway of *Pseudomonas solanacearum*," EMBO J. 13:543-553 (1994)).

The first of these proteins was discovered in *E. amylovora* Ea321, a bacterium that causes fire blight of rosaceous plants, and was designated harpin (Wei, Z.-M., et al, "Harpin, Elicitor of the Hypersensitive Response Produced by the Plant Pathogen *Erwinia amylovora*," Science 257:85-88 (1992)). Mutations in the encoding *hrpN* gene revealed that harpin is required for *E. amylovora* to elicit a hypersensitive response in nonhost tobacco leaves and incite disease symptoms in highly susceptible pear fruit. The *P. solanacearum* GMI1000 PopA1 protein has similar physical properties and also elicits the hypersensitive response in leaves of tobacco, which is not a host of that strain (Arlat, et al. "PopA1, a Protein Which Induces a Hypersensitive-like Response on Specific Petunia Genotypes, is Secreted via the Hrp Pathway of *Pseudomonas solanacearum*," EMBO J. 13:543-53 (1994)). However, *P. solanacearum* *popA* mutants still elicit the hypersensitive response in tobacco and incite disease in tomato. Thus, the role of these glycine-rich hypersensitive response elicitors can vary widely among Gram-negative plant pathogens.

Other plant pathogenic hypersensitive response elicitors have been isolated, cloned, and sequenced. These include: *Erwinia chrysanthemi* (Bauer, et. al., "*Erwinia chrysanthemi* Harpin_{Ech}: Soft-Rot Pathogenesis," MPMI 8(4): 484-91 (1995)); *Erwinia carotovora* (Cui, et. al., "The RsmA⁻ Mutants of *Erwinia carotovora* subsp. *carotovora* Strain Ecc71 Overexpress *hrpN*_{Ecc} and Elicit a Hypersensitive Reaction-like Response in Tobacco Leaves," MPMI 9(7): 565-73 (1996)); *Erwinia stewartii* (Ahmad, et. al., "Harpin is not Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," 8th Int'l. Cong. Molec. Plant-Microb. Inter. July 14-19, 1996 and Ahmad, et. al., "Harpin is not Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," Ann. Mtg. Am. Phytopath. Soc. July 27-31, 1996); and *Pseudomonas syringae* pv. *syringae* (WO 94/26782 to Cornell Research Foundation, Inc.).

- 4 -

The present invention seeks to identify fragments of hypersensitive response elicitor proteins or polypeptides, which fragments do not elicit a hypersensitive response but are active when utilized in conjunction with plants.

5

SUMMARY OF THE INVENTION

The present invention is directed to isolated fragments of an *Erwinia* hypersensitive response elicitor protein or polypeptide which fragments do not elicit a hypersensitive response in plants but are otherwise active when utilized in
10 conjunction with plants. Also disclosed are isolated DNA molecules which encode such fragments.

The fragments of hypersensitive response elicitors according to the present invention have the following activity when utilized in conjunction with plants: imparting disease resistance to plants, enhancing plant growth and/or controlling
15 insects. This involves applying the fragments in a non-infectious form to plants or plant seeds under conditions effective to impart disease resistance, to enhance plant growth, and/or to control insects on plants or plants grown from the plant seeds.

As an alternative to applying the fragments to plants or plant seeds in order to impart disease resistance, to enhance plant growth, and/or to control insects
20 on plants, transgenic plants or plant seeds can be utilized. When utilizing transgenic plants, this involves providing a transgenic plant transformed with a DNA molecule encoding a fragment of a hypersensitive response elicitor protein or polypeptide in accordance with the present invention and growing the plant under conditions effective to impart disease resistance, to enhance plant growth, and/or to control
25 insects in the plants or plants grown from the plant seeds. Alternatively, a transgenic plant seed transformed with the DNA molecule encoding such a fragment can be provided and planted in soil. A plant is then propagated under conditions effective to impart disease resistance, to enhance plant growth, and/or to control insects on plants or plants grown from the plant seeds.

30

- 5 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows truncated proteins of the hypersensitive response elicitor protein or polypeptide.

5 Figure 2 shows a list of synthesized oligonucleotide primers for construction of truncated harpin proteins. N represents the N-terminus (5' region), and C represents the C-terminus (3' region). The primers correspond to the indicated sequence identification numbers for the present application: N1 (SEQ. ID. No. 1), N76 (SEQ. ID. No. 2), N99 (SEQ. ID. No. 3), N105 (SEQ. ID. No. 4), N110 (SEQ. ID. No. 5), N137 (SEQ. ID. No. 6), N150 (SEQ. ID. No. 7), N169 (SEQ. ID. No. 8),
10 N210 (SEQ. ID. No. 9), N267 (SEQ. ID. No. 10), N343 (SEQ. ID. No. 11), C75 (SEQ. ID. No. 12), C104 (SEQ. ID. No. 13), C168 (SEQ. ID. No. 14), C180 (SEQ. ID. No. 15), C204 (SEQ. ID. No. 16), C209 (SEQ. ID. No. 17), C266 (SEQ. ID. No. 18), C342 (SEQ. ID. No. 19), and C403 (SEQ. ID. No. 20).

15

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to isolated fragments of a hypersensitive response elicitor protein or polypeptide where the fragments do not
20 elicit a hypersensitive response but have other activity in plants. Also disclosed are DNA molecules encoding such fragments as well as expression systems, host cells, and plants containing such molecules. Uses of the fragments themselves and the DNA molecules encoding them are disclosed.

The fragments of hypersensitive response elicitor polypeptides or
25 proteins according to the present invention are derived from hypersensitive response elicitor polypeptides or proteins of a wide variety of fungal and bacterial pathogens. Such polypeptides or proteins are able to elicit local necrosis in plant tissue contacted by the elicitor. Examples of suitable bacterial sources of polypeptide or protein elicitors include *Erwinia*, *Pseudomonas*, and *Xanthomonas* species (e.g., the
30 following bacteria: *Erwinia amylovora*, *Erwinia chrysanthemi*, *Erwinia stewartii*, *Erwinia carotovora*, *Pseudomonas syringae*, *Pseudomonas solanacearum*, *Xanthomonas campestris*, and mixtures thereof).

- 6 -

An example of a fungal source of a hypersensitive response elicitor protein or polypeptide is *Phytophthora*. Suitable species of *Phytophthora* include *Phytophthora parasitica*, *Phytophthora cryptogea*, *Phytophthora cinnamomi*, *Phytophthora capsici*, *Phytophthora megasperma*, and *Phytophthora citrophthora*.

5 The hypersensitive response elicitor polypeptide or protein from *Erwinia chrysanthemi* has an amino acid sequence corresponding to SEQ. ID. No. 21 as follows:

10	Met	Gln	Ile	Thr	Ile	Lys	Ala	His	Ile	Gly	Gly	Asp	Leu	Gly	Val	Ser	1	5	10	15
	Gly	Leu	Gly	Ala	Gln	Gly	Leu	Lys	Gly	Leu	Asn	Ser	Ala	Ala	Ser	Ser	20	25	30	
	Leu	Gly	Ser	Ser	Val	Asp	Lys	Leu	Ser	Ser	Thr	Ile	Asp	Lys	Leu	Thr	35	40	45	
15	Ser	Ala	Leu	Thr	Ser	Met	Met	Phe	Gly	Gly	Ala	Leu	Ala	Gln	Gly	Leu	50	55	60	
	Gly	Ala	Ser	Ser	Lys	Gly	Leu	Gly	Met	Ser	Asn	Gln	Leu	Gly	Gln	Ser	65	70	75	80
20	Phe	Gly	Asn	Gly	Ala	Gln	Gly	Ala	Ser	Asn	Leu	Leu	Ser	Val	Pro	Lys	85	90	95	
	Ser	Gly	Gly	Asp	Ala	Leu	Ser	Lys	Met	Phe	Asp	Lys	Ala	Leu	Asp	Asp	100	105	110	
	Leu	Leu	Gly	His	Asp	Thr	Val	Thr	Lys	Leu	Thr	Asn	Gln	Ser	Asn	Gln	115	120	125	
25	Leu	Ala	Asn	Ser	Met	Leu	Asn	Ala	Ser	Gln	Met	Thr	Gln	Gly	Asn	Met	130	135	140	
	Asn	Ala	Phe	Gly	Ser	Gly	Val	Asn	Asn	Ala	Leu	Ser	Ser	Ile	Leu	Gly	145	150	155	160
30	Asn	Gly	Leu	Gly	Gln	Ser	Met	Ser	Gly	Phe	Ser	Gln	Pro	Ser	Leu	Gly	165	170	175	
	Ala	Gly	Gly	Leu	Gln	Gly	Leu	Ser	Gly	Ala	Gly	Ala	Phe	Asn	Gln	Leu	180	185	190	
	Gly	Asn	Ala	Ile	Gly	Met	Gly	Val	Gly	Gln	Asn	Ala	Ala	Leu	Ser	Ala	195	200	205	
35	Leu	Ser	Asn	Val	Ser	Thr	His	Val	Asp	Gly	Asn	Asn	Arg	His	Phe	Val	210	215	220	
	Asp	Lys	Glu	Asp	Arg	Gly	Met	Ala	Lys	Glu	Ile	Gly	Gln	Phe	Met	Asp	225	230	235	240

[illegible]

	CGATTTTACC	CGGGTGAACG	TGCTATGACC	GACAGCATCA	CGGTATTCTGA	CACCGTTACG	60
	GCGTTTATGG	CCGCGATGAA	CCGGCATCAG	GCGGCGCGCT	GGTCGCCGCA	ATCCGGCGTC	120
	GATCTGGTAT	TTCAGTTTGG	GGACACCGGG	CGTGAAGTCA	TGATGCAGAT	TCAGCCGGGG	180
25	CAGCAATATC	CCGGCATGTT	GCGCACGCTG	CTCGCTCGTC	GTTATCAGCA	GGCGGCAGAG	240
	TGCGATGGCT	GCCATCTGTG	CCTGAACGGC	AGCGATGTAT	TGATCCTCTG	GTGGCCGCTG	300
	CCGTCCGATC	CCGGCAGTTA	TCCGCAGGTG	ATCGAACGTT	TGTTTGAAGT	GGCGGGAATG	360
	ACGTTGCCGT	CGCTATCCAT	AGCACCGACG	GCGCGTCCGC	AGACAGGGAA	CGGACGCGCC	420
	CGATCATTAA	GATAAAGGCG	GCTTTTTTTT	TTGCAAAACG	GTAACGGTGA	GGAACCGTTT	480
30	CACCGTCGGC	GTCAGTCTAGT	AACAAGTATC	CATCATGATG	CCTACATCGG	GATCGGCGTG	540
	GGCATCCGTT	GCAGATACTT	TTGCGAACAC	CTGACATGAA	TGAGGAAACG	AAATTATGCA	600
	AATTACGATC	AAAGCGCACA	TCGGCGGTGA	TTTGGGCGTC	TCCGGTCTGG	GGCTGGGTGC	660
	TCAGGGACTG	AAAGGACTGA	ATTCCGCGGC	TTCATCGCTG	GGTTCCAGCG	TGGATAAACT	720
	GAGCAGCACC	ATCGATAAGT	TGACCTCCGC	GCTGACTTCG	ATGATGTTTG	GCGGCGCGCT	780
35	GGCGCAGGGG	CTGGGCGCCA	GCTCGAAGGG	GCTGGGGATG	AGCAATCAAC	TGGGCCAGTC	840

- 8 -

TTTCGGCAAT GGC GCGCAGG GTGCGAGCAA CCTGCTATCC GTACCGAAAT CCGGCGGCGA 900
 TCGTGTGTCA AAAATGTTTG ATAAAGCGCT GGACGATCTG CTGGGTCATG ACACCGTGAC 960
 CAAGCTGACT AACCAGAGCA ACCAACTGGC TAATTCAATG CTGAACGCCA GCCAGATGAC 1020
 CCAGGGTAAT ATGAATGCGT TCGGCAGCGG TGTGAACAAC GCACTGTCGT CCATTCTCGG 1080
 5 CAACGGTCTC GGCCAGTCTG TGAGTGGCTT CTCTCAGCCT TCTCTGGGGG CAGGCGGCTT 1140
 GCAGGGCCTG AGCGGCGCGG GTGCATTCAA CCAGTTGGGT AATGCCATCG GCATGGGCGT 1200
 GGGGCAGAAT GCTGCGCTGA GTGCGTTGAG TAACGTCAGC ACCCACGTAG ACGGTAACAA 1260
 CCGCCACTTT GTAGATAAAG AAGATCGCGG CATGGCGAAA GAGATCGGCC AGTTTATGGA 1320
 TCAGTATCCG GAAATATTCG GTAAACCGGA ATACCAGAAA GATGGCTGGA GTTCGCCGAA 1380
 10 GACGGACGAC AAATCCTGGG CTAAAGCGCT GAGTAAACCG GATGATGACG GTATGACCGG 1440
 CGCCAGCATG GACAAATTCC GTCAGGCGAT GGGTATGATC AAAAGCGCGG TGGCGGGTGA 1500
 TACCGGCAAT ACCAACCTGA ACCTGCGTGG CGCGGGCGGT GCATCGCTGG GTATCGATGC 1560
 GGCTGTCTGTC GGC GATAAAA TAGCCAACAT GTCGCTGGGT AAGCTGGCCA ACGCCTGATA 1620
 ATCTGTGCTG GCCTGATAAA GCGGAAACGA AAAAAGAGAC GGGGAAGCCT GTCTCTTTTC 1680
 15 TTATTATGCG GTTTATGCGG TTACCTGGAC CGGTTAATCA TCGTCATCGA TCTGGTACAA 1740
 ACGCACATTT TCCCGTTCAT TCGCGTCGTT ACGCGCCACA ATCGCGATGG CATCTTCCTC 1800
 GTCGCTCAGA TTGCGCGGCT GATGGGGAAC GCCGGGTGGA ATATAGAGAA ACTCGCCGGC 1860
 CAGATGGAGA CACGTCTGCG ATAAATCTGT GCCGTAACGT GTTTCTATCC GCCCCTTTAG 1920
 CAGATAGATT GCGGTTTCGT AATCAACATG GTAATGCGGT TCCGCCTGTG CGCCGGCCGG 1980
 20 GATCACCACA ATATTCATAG AAAGCTGTCT TGCACCTACC GTATCGCGGG AGATACCGAC 2040
 AAAATAGGGC AGTTTTTGCG TGGTATCCGT GGGGTGTTCC GGCCTGACAA TCTTGAGTTG 2100
 GTTCGTCATC ATCTTTCTCC ATCTGGGCGA CCTGATCGGT T 2141

25 The hypersensitive response elicitor polypeptide or protein derived
 from *Erwinia amylovora* has an amino acid sequence corresponding to SEQ. ID.
 No. 23 as follows:

30 Met Ser Leu Asn Thr Ser Gly Leu Gly Ala Ser Thr Met Gln Ile Ser
 1 5 10 15
 Ile Gly Gly Ala Gly Gly Asn Asn Gly Leu Leu Gly Thr Ser Arg Gln
 20 25 30

- 9 -

	Asn	Ala	Gly	Leu	Gly	Gly	Asn	Ser	Ala	Leu	Gly	Leu	Gly	Gly	Gly	Asn	
			35					40					45				
	Gln	Asn	Asp	Thr	Val	Asn	Gln	Leu	Ala	Gly	Leu	Leu	Thr	Gly	Met	Met	
		50					55					60					
5	Met	Met	Met	Ser	Met	Met	Gly	Gly	Gly	Gly	Leu	Met	Gly	Gly	Gly	Leu	
	65					70					75					80	
	Gly	Gly	Gly	Leu	Gly	Asn	Gly	Leu	Gly	Gly	Ser	Gly	Gly	Leu	Gly	Glu	
					85					90					95		
10	Gly	Leu	Ser	Asn	Ala	Leu	Asn	Asp	Met	Leu	Gly	Gly	Ser	Leu	Asn	Thr	
				100					105						110		
	Leu	Gly	Ser	Lys	Gly	Gly	Asn	Asn	Thr	Thr	Ser	Thr	Thr	Asn	Ser	Pro	
			115					120						125			
	Leu	Asp	Gln	Ala	Leu	Gly	Ile	Asn	Ser	Thr	Ser	Gln	Asn	Asp	Asp	Ser	
		130					135					140					
15	Thr	Ser	Gly	Thr	Asp	Ser	Thr	Ser	Asp	Ser	Ser	Asp	Pro	Met	Gln	Gln	
	145					150					155					160	
	Leu	Leu	Lys	Met	Phe	Ser	Glu	Ile	Met	Gln	Ser	Leu	Phe	Gly	Asp	Gly	
					165					170					175		
20	Gln	Asp	Gly	Thr	Gln	Gly	Ser	Ser	Ser	Gly	Gly	Lys	Gln	Pro	Thr	Glu	
				180					185					190			
	Gly	Glu	Gln	Asn	Ala	Tyr	Lys	Lys	Gly	Val	Thr	Asp	Ala	Leu	Ser	Gly	
			195					200					205				
	Leu	Met	Gly	Asn	Gly	Leu	Ser	Gln	Leu	Leu	Gly	Asn	Gly	Gly	Leu	Gly	
		210					215					220					
25	Gly	Gly	Gln	Gly	Gly	Asn	Ala	Gly	Thr	Gly	Leu	Asp	Gly	Ser	Ser	Leu	
	225					230					235					240	
	Gly	Gly	Lys	Gly	Leu	Gln	Asn	Leu	Ser	Gly	Pro	Val	Asp	Tyr	Gln	Gln	
					245					250					255		
30	Leu	Gly	Asn	Ala	Val	Gly	Thr	Gly	Ile	Gly	Met	Lys	Ala	Gly	Ile	Gln	
				260					265					270			
	Ala	Leu	Asn	Asp	Ile	Gly	Thr	His	Arg	His	Ser	Ser	Thr	Arg	Ser	Phe	
			275					280					285				
	Val	Asn	Lys	Gly	Asp	Arg	Ala	Met	Ala	Lys	Glu	Ile	Gly	Gln	Phe	Met	
		290					295					300					
35	Asp	Gln	Tyr	Pro	Glu	Val	Phe	Gly	Lys	Pro	Gln	Tyr	Gln	Lys	Gly	Pro	
	305					310					315					320	
	Gly	Gln	Glu	Val	Lys	Thr	Asp	Asp	Lys	Ser	Trp	Ala	Lys	Ala	Leu	Ser	
					325					330					335		
	Lys	Pro	Asp	Asp	Asp	Gly	Met	Thr	Pro	Ala	Ser	Met	Glu	Gln	Phe	Asn	

[illegible]

This hypersensitive response elicitor polypeptide or protein has a molecular weight of about 39 kDa, has a pI of approximately 4.3, and is heat stable at 100°C for at least 10 minutes. This hypersensitive response elicitor polypeptide or protein has substantially no cysteine. The hypersensitive response elicitor polypeptide or protein derived from *Erwinia amylovora* is more fully described in Wei, Z.-M., R. J. Laby, C. H. Zumoff, D. W. Bauer, S.-Y. He, A. Collmer, and S. V. Beer, "Harpin, Elicitor of the Hypersensitive Response Produced by the Plant Pathogen *Erwinia amylovora*," Science 257:85-88 (1992), which is hereby incorporated by reference. The DNA molecule encoding this polypeptide or protein has a nucleotide sequence corresponding to SEQ. ID. No. 24 as follows:

20	AAGCTTCGGC	ATGGCACGTT	TGACCGTTGG	GTCGGCAGGG	TACGTTTGAA	TTATTCTATA	60
	GAGGAATACG	TTATGAGTCT	GAATACAAGT	GGGCTGGGAG	CGTCAACGAT	GCAAATTTCT	120
	ATCGGCGGTG	CGGGCGGAAA	TAACGGGTTG	CTGGGTACCA	GTCGCCAGAA	TGCTGGGTTG	180
	GGTGGCAATT	CTGCACTGGG	GCTGGGCGGC	GGTAATCAAA	ATGATACCGT	CAATCAGCTG	240
25	GCTGGCTTAC	TCACCGGCAT	GATGATGATG	ATGAGCATGA	TGGGCGGTGG	TGGGCTGATG	300
	GGCGGTGGCT	TAGGCGGTGG	CTTAGGTAAT	GGCTTGGGTG	GCTCAGGTGG	CCTGGGCGAA	360
	GGACTGTCTG	ACGCGCTGAA	CGATATGTTA	GGCGGTTTCG	TGAACACGCT	GGGCTCGAAA	420
	GGCGGCAACA	ATACCACTTC	AACAACAAAT	TCCCCGCTGG	ACCAGGCGCT	GGGTATTAACT	480
30	TCAACGTCCC	AAAACGACGA	TTCCACCTCC	GGCACAGATT	CCACCTCAGA	CTCCAGCGAC	540
	CCGATGCAGC	AGCTGCTGAA	GATGTTTCAGC	GAGATAATGC	AAAGCCTGTT	TGGTGATGGG	600
	CAAGATGGCA	CCCAGGGCAG	TTCCTCTGGG	GGCAAGCAGC	CGACCGAAGG	CGAGCAGAAC	660
	GCCTATAAAA	AAGGAGTCAC	TGATGCGCTG	TCGGGCCTGA	TGGGTAATGG	TCTGAGCCAG	720
	CTCCTTGGCA	ACGGGGGACT	GGGAGGTGGT	CAGGGCGGTA	ATGCTGGCAC	GGGTCTTGAC	780
	GGTTCGTCGC	TGGGCGGCAA	AGGGCTGCAA	AACCTGAGCG	GGCCGGTGGA	CTACCAGCAG	840

- 11 -

TTAGGTAACG CCGTGGGTAC CGGTATCGGT ATGAAAGCGG GCATTTCAGGC GCTGAATGAT 900
 ATCGGTACGC ACAGGCACAG TTCAACCCGT TCTTTCGTCA ATAAAGGCGA TCGGGCGATG 960
 GCGAAGGAAA TCGGTCAGTT CATGGACCAG TATCCTGAGG TGTTTGGCAA GCCGCAGTAC 1020
 CAGAAAGGCC CGGGTCAGGA GGTGAAAACC GATGACAAAT CATGGGCAAA AGCACTGAGC 1080
 5 AAGCCAGATG ACGACGGAAT GACACCAGCC AGTATGGAGC AGTTCAACAA AGCCAAGGGC 1140
 ATGATCAAAA GGCCCATGGC GGGTGATACC GGCAACGGCA ACCTGCAGGC ACGCGGTGCC 1200
 GGTGGTTCTT CGTGGGTAT TGATGCCATG ATGGCCGGTG ATGCCATTAA CAATATGGCA 1260
 CTTGGCAAGC TGGGCGCGGC TTAAGCTT 1288

10

Another potentially suitable hypersensitive response elicitor from
Erwinia amylovora is disclosed in U.S. Patent Application Serial No. 09/120,927,
 which is hereby incorporated by reference. The protein is encoded by a DNA
 molecule having a nucleic acid sequence of SEQ. ID. No. 25 as follows:

15

ATGTCAATTC TTACGCTTAA CAACAATACC TCGTCCTCGC CGGGTCTGTT CCAGTCCGGG 60
 GGGGACAACG GGCTTGGTGG TCATAATGCA AATTCTGCGT TGGGGCAACA ACCCATCGAT 120
 20 CGGCAAACCA TTGAGCAAAT GGCTCAATTA TTGGCGGAAC TGTTAAAGTC ACTGCTATCG 180
 CCACAATCAG GTAATGCGGC AACCGGAGCC GGTGGCAATG ACCAGACTAC AGGAGTTGGT 240
 AACGCTGGCG GCCTGAACGG ACGAAAAGGC ACAGCAGGAA CCACTCCGCA GTCTGACAGT 300
 25 CAGAACATGC TGAGTGAGAT GGGCAACAAC GGGCTGGATC AGGCCATCAC GCCCATGGC 360
 CAGGGCGGCG GGCAGATCGG CGATAATCCT TTAGTAAAG CCATGCTGAA GCTTATTGCA 420
 30 CGCATGATGG ACGGCCAAAG CGATCAGTTT GGCCAACTG GTACGGGCAA CAACAGTGCC 480
 TCTTCCGGTA CTTCTTCATC TGGCGGTTCC CTTTAAACG ATCTATCAGG GGGGAAGGCC 540
 CCTTCCGGCA ACTCCCTTC CGGCAACTAC TCTCCCGTCA GTACCTTCTC ACCCCATCC 600
 35 ACGCCAACGT CCCCTACCTC ACCGCTTGAT TTCCCTTCTT CTCCCACCAA AGCAGCCGGG 660
 GGCAGCACGC CGGTAACCGA TCATCCTGAC CCTGTTGGTA GCGCGGGCAT CGGGGCCGGA 720
 40 AATTCGGTGG CCTTCACCAG CGCCGGCGCT AATCAGACGG TGCTGCATGA CACCATTACC 780
 GTGAAAGCGG GTCAGGTGTT TGATGGCAAA GGACAAACCT TCACCGCCGG TTCAGAATTA 840
 GCGGATGGCG GCCAGTCTGA AAACCAGAAA CCGCTGTTTA TACTGGAAGA CCGTGCCAGC 900
 45 CTGAAAAACG TCACCATGGG CGACGACGGG GCGGATGGTA TTCATCTTTA CCGTGATGCC 960
 AAAATAGACA ATCTGCACGT CACCAACGTG GGTGAGGACG CGATTACCGT TAAGCCAAAC 1020
 50 AGCGCGGGCA AAAAATCCCA CGTTGAAATC ACTAACAGTT CCTTCGAGCA CGCCTCTGAC 1080
 AAGATCCTGC AGCTGAATGC CGATACTAAC CTGAGCGTTG ACAACGTGAA GGCCAAAGAC 1140

- 12 -

TTTGGTACTT TTGTACGCAC TAACGGCGGT CAACAGGGTA ACTGGGATCT GAATCTGAGC 1200
 CATATCAGCG CAGAAGACGG TAAGTTCTCG TTCGTTAAAA GCGATAGCGA GGGGCTAAAC 1260
 GTCAATACCA GTGATATCTC ACTGGGTGAT GTTGAAAACC ACTACAAAGT GCCGATGTCC 1320
 GCCAACCTGA AGGTGGCTGA ATGA 1344

10

See GenBank Accession No. U94513. The isolated DNA molecule of the present invention encodes a hypersensitive response elicitor protein or polypeptide having an amino acid sequence of SEQ. ID. No. 26 as follows:

15 Met Ser Ile Leu Thr Leu Asn Asn Asn Thr Ser Ser Ser Pro Gly Leu
 1 5 10 15
 Phe Gln Ser Gly Gly Asp Asn Gly Leu Gly Gly His Asn Ala Asn Ser
 20 20 25 30
 Ala Leu Gly Gln Gln Pro Ile Asp Arg Gln Thr Ile Glu Gln Met Ala
 35 40 45
 Gln Leu Leu Ala Glu Leu Leu Lys Ser Leu Leu Ser Pro Gln Ser Gly
 25 50 55 60
 Asn Ala Ala Thr Gly Ala Gly Gly Asn Asp Gln Thr Thr Gly Val Gly
 65 70 75 80
 30 Asn Ala Gly Gly Leu Asn Gly Arg Lys Gly Thr Ala Gly Thr Thr Pro
 85 90 95
 Gln Ser Asp Ser Gln Asn Met Leu Ser Glu Met Gly Asn Asn Gly Leu
 100 105 110
 35 Asp Gln Ala Ile Thr Pro Asp Gly Gln Gly Gly Gly Gln Ile Gly Asp
 115 120 125
 Asn Pro Leu Leu Lys Ala Met Leu Lys Leu Ile Ala Arg Met Met Asp
 40 130 135 140
 Gly Gln Ser Asp Gln Phe Gly Gln Pro Gly Thr Gly Asn Asn Ser Ala
 145 150 155 160
 45 Ser Ser Gly Thr Ser Ser Ser Gly Gly Ser Pro Phe Asn Asp Leu Ser
 165 170 175
 Gly Gly Lys Ala Pro Ser Gly Asn Ser Pro Ser Gly Asn Tyr Ser Pro
 180 185 190
 50 Val Ser Thr Phe Ser Pro Pro Ser Thr Pro Thr Ser Pro Thr Ser Pro
 195 200 205
 Leu Asp Phe Pro Ser Ser Pro Thr Lys Ala Ala Gly Gly Ser Thr Pro
 55 210 215 220

- 13 -

	Val	Thr	Asp	His	Pro	Asp	Pro	Val	Gly	Ser	Ala	Gly	Ile	Gly	Ala	Gly	
	225					230					235					240	
5	Asn	Ser	Val	Ala	Phe	Thr	Ser	Ala	Gly	Ala	Asn	Gln	Thr	Val	Leu	His	
				245					250						255		
	Asp	Thr	Ile	Thr	Val	Lys	Ala	Gly	Gln	Val	Phe	Asp	Gly	Lys	Gly	Gln	
			260					265					270				
10	Thr	Phe	Thr	Ala	Gly	Ser	Glu	Leu	Gly	Asp	Gly	Gly	Gln	Ser	Glu	Asn	
		275						280					285				
	Gln	Lys	Pro	Leu	Phe	Ile	Leu	Glu	Asp	Gly	Ala	Ser	Leu	Lys	Asn	Val	
15		290					295					300					
	Thr	Met	Gly	Asp	Asp	Gly	Ala	Asp	Gly	Ile	His	Leu	Tyr	Gly	Asp	Ala	
	305					310					315					320	
	Lys	Ile	Asp	Asn	Leu	His	Val	Thr	Asn	Val	Gly	Glu	Asp	Ala	Ile	Thr	
20				325						330					335		
	Val	Lys	Pro	Asn	Ser	Ala	Gly	Lys	Lys	Ser	His	Val	Glu	Ile	Thr	Asn	
				340					345					350			
25	Ser	Ser	Phe	Glu	His	Ala	Ser	Asp	Lys	Ile	Leu	Gln	Leu	Asn	Ala	Asp	
			355					360					365				
	Thr	Asn	Leu	Ser	Val	Asp	Asn	Val	Lys	Ala	Lys	Asp	Phe	Gly	Thr	Phe	
30		370					375					380					
	Val	Arg	Thr	Asn	Gly	Gly	Gln	Gln	Gly	Asn	Trp	Asp	Leu	Asn	Leu	Ser	
	385				390						395					400	
	His	Ile	Ser	Ala	Glu	Asp	Gly	Lys	Phe	Ser	Phe	Val	Lys	Ser	Asp	Ser	
35				405						410					415		
	Glu	Gly	Leu	Asn	Val	Asn	Thr	Ser	Asp	Ile	Ser	Leu	Gly	Asp	Val	Glu	
			420						425					430			
40	Asn	His	Tyr	Lys	Val	Pro	Met	Ser	Ala	Asn	Leu	Lys	Val	Ala	Glu		
			435					440					445				

This protein or polypeptide is acidic, rich in glycine and serine, and lacks cysteine. It is also heat stable, protease sensitive, and suppressed by inhibitors of plant metabolism. The protein or polypeptide of the present invention has a predicted molecular size of ca. 4.5 kDa.

Another potentially suitable hypersensitive response elicitor from *Erwinia amylovora* is disclosed in U.S. Patent Application Serial No. 09/120,663 which is hereby incorporated by reference. The protein is encoded by a DNA molecule having a nucleic acid sequence of SEQ. ID. No. 27 as follows:

- 14 -

	ATGGAATTAA	AATCACTGGG	AACTGAACAC	AAGGCGGCAG	TACACACAGC	GGCGCACAAAC	60
	CCTGTGGGGC	ATGGTGTTC	CTTACAGCAG	GGCAGCAGCA	GCAGCAGCCC	GCAAAATGCC	120
5	GCTGCATCAT	TGGCGGCAGA	AGGCAAAAAT	CGTGGGAAAA	TGCCGAGAAT	TCACCAGCCA	180
	TCTACTGCGG	CTGATGGTAT	CAGCGCTGCT	CACCAGCAAA	AGAAATCCTT	CAGTCTCAGG	240
10	GGCTGTTTGG	GGACGAAAAA	ATTTTCCAGA	TCGGCACC GC	AGGGCCAGCC	AGGTACCACC	300
	CACAGCAAAG	GGGCAACATT	GCGCGATCTG	CTGGCGCGGG	ACGACGGCGA	AACGCAGCAT	360
	GAGGCGGCCG	CGCCAGATGC	GGCGCGTTTG	ACCCGTTTCG	GCGGCGTCAA	ACGCCGCAAT	420
15	ATGGACGACA	TGGCCGGGCG	GCCAATGGTG	AAAGGTGGCA	GCGGCGAAGA	TAAGGTACCA	480
	ACGCAGCAAA	AACGGCATCA	GCTGAACAAT	TTTGGCCAGA	TGCGCCAAAC	GATGTTGAGC	540
20	AAAATGGCTC	ACCCGGCTTC	AGCCAACGCC	GGCGATCGCC	TGCAGCATTC	ACCGCCGCAC	600
	ATCCCGGGTA	GCCACCACGA	AATCAAGGAA	GAACCGGTTG	GCTCCACCAG	CAAGGCAACA	660
	ACGGCCACG	CAGACAGAGT	GGAAATCGCT	CAGGAAGATG	ACGACAGCGA	ATTCCAGCAA	720
25	CTGCATCAAC	AGCGGCTGGC	GCGCGAACGG	GAAAATCCAC	CGCAGCCGCC	CAAACCTCGG	780
	GTTGCCACAC	CGATTAGCGC	CAGGTTTCAG	CCCAAATGTA	CTGCGGTTGC	GGAAAGCGTC	840
30	CTTGAGGGGA	CAGATACCAC	GCAGTCACCC	CTTAAGCCGC	AATCAATGCT	GAAAGGAAGT	900
	GGAGCCGGGG	TAACGCCGCT	GGCGGTAACG	CTGGATAAAG	GCAAGTTGCA	GCTGGCACCG	960
	GATAATCCAC	CCGCGCTCAA	TACGTTGTTG	AAGCAGACAT	TGGGTAAAGA	CACCCAGCAC	1020
35	TATCTGGCGC	ACCATGCCAG	CAGCGACGGT	AGCCAGCATC	TGCTGCTGGA	CAACAAAGGC	1080
	CACCTGTTTG	ATATCAAAAG	CACCGCCACC	AGCTATAGCG	TGCTGCACAA	CAGCCACCCC	1140
40	GGTGAGATAA	AGGGCAAGCT	GGCGCAGGCG	GGTACTGGCT	CCGTCAGCGT	AGACGGTAAA	1200
	AGCGGCAAGA	TCTCGCTGGG	GAGCGGTACG	CAAAGTCACA	ACAAAACAAT	GCTAAGCCAA	1260
	CCGGGGGAAG	CGCACCGTTC	CTTATTAACC	GGCATTGTC	AGCATCCTGC	TGGCGCAGCG	1320
45	CGGCCGCAGG	GCGAGTCAAT	CCGCCTGCAT	GACGACAAAA	TTCATATCCT	GCATCCGGAG	1380
	CTGGGCGTAT	GGCAATCTGC	GGATAAAGAT	ACCCACAGCC	AGCTGTCTCG	CCAGGCAGAC	1440
50	GGTAAGCTCT	ATGCGCTGAA	AGACAACCGT	ACCCTGCAAA	ACCTCTCCGA	TAATAAATCC	1500
	TCAGAAAAGC	TGGTCGATAA	AATCAAATCG	TATTCCGTTG	ATCAGCGGGG	GCAGGTGGCG	1560
	ATCCTGACGG	ATACTCCCGG	CCGCCATAAG	ATGAGTATTA	TGCCCTCGCT	GGATGCTTCC	1620
55	CCGGAGAGCC	ATATTTCCCT	CAGCCTGCAT	TTTGCCGATG	CCCACCAGGG	GTTATTGCAC	1680
	GGGAAGTCGG	AGCTTGAGGC	ACAATCTGTC	GCGATCAGCC	ATGGGCGACT	GGTTGTGGCC	1740
60	GATAGCGAAG	GCAAGCTGTT	TAGCGCCGCC	ATTCCGAAGC	AAGGGGATGG	AAACGAACTG	1800
	AAAATGAAAG	CCATGCCTCA	GCATGCGCTC	GATGAACATT	TTGGTCATGA	CCACCAGATT	1860
	TCTGGATTTT	TCCATGACGA	CCACGGCCAG	CTTAATGCGC	TGGTGAAAAA	TAACTTCAGG	1920
65	CAGCAGCATG	CCTGCCCGTT	GGGTAACGAT	CATCAGTTTC	ACCCCGGCTG	GAACCTGACT	1980

- 15 -

	GATGCGCTGG	TTATCGACAA	TCAGCTGGGG	CTGCATCATA	CCAATCCTGA	ACCGCATGAG	2040
5	ATTCTTGATA	TGGGGCATT	AGGCAGCCTG	GCGTTACAGG	AGGGCAAGCT	TCACTATTTT	2100
	GACCAGCTGA	CCAAAGGGTG	GACTGGCGCG	GAGTCAGATT	GTAAGCAGCT	GAAAAAAGGC	2160
	CTGGATGGAG	CAGCTTATCT	ACTGAAAGAC	GGTGAAGTGA	AACGCCTGAA	TATTAATCAG	2220
10	AGCACCTCCT	CTATCAAGCA	CGGAACGGAA	AACGTTTTTT	CGCTGCCGCA	TGTGCGCAAT	2280
	AAACCGGAGC	CGGGAGATGC	CCTGCAAGGG	CTGAATAAAG	ACGATAAGGC	CCAGGCCATG	2340
15	GCGGTGATTG	GGGTAAATAA	ATACCTGGCG	CTGACGGAAA	AAGGGGACAT	TCGCTCCTTC	2400
	CAGATAAAAC	CCGGCACCCA	GCAGTTGGAG	CGGCCGGCAC	AAACTCTCAG	CCGCGAAGGT	2460
	ATCAGCGGCG	AACTGAAAGA	CATTCATGTC	GACCACAAGC	AGAACCTGTA	TGCCTTGACC	2520
20	CACGAGGGAG	AGGTGTTTCA	TCAGCCGCGT	GAAGCCTGGC	AGAATGGTGC	CGAAAGCAGC	2580
	AGCTGGCACA	AACTGGCGTT	GCCACAGAGT	GAAAGTAAGC	TAAAAAGTCT	GGACATGAGC	2640
25	CATGAGCACA	AACCGATTGC	CACCTTTGAA	GACGGTAGCC	AGCATCAGCT	GAAGGCTGGC	2700
	GGCTGGCACG	CCTATGCGGC	ACCTGAACGC	GGGCCGCTGG	CGGTGGGTAC	CAGCGGTTCA	2760
	CAAACCGTCT	TTAACCGACT	AATGCAGGGG	GTGAAAGGCA	AGGTGATCCC	AGGCAGCGGG	2820
30	TTGACGGTTA	AGCTCTCGGC	TCAGACGGGG	GGAATGACCG	GCGCCGAAGG	GCGCAAGGTC	2880
	AGCAGTAAAT	TTTCCGAAAG	GATCCGCGCC	TATGCGTTCA	ACCCAACAAT	GTCCACGCCG	2940
35	CGACCATTAA	AAAATGCTGC	TTATGCCACA	CAGCACGGCT	GGCAGGGGCG	TGAGGGGTTG	3000
	AAGCCGTTGT	ACGAGATGCA	GGGAGCGCTG	ATTAAACAAC	TGGATGCGCA	TAACGTTCTG	3060
	CATAACGCGC	CACAGCCAGA	TTTGCAGAGC	AAACTGGAAA	CTCTGGATTT	AGGCGAACAT	3120
40	GGCGCAGAAT	TGCTTAACGA	CATGAAGCGC	TTCCGCGACG	AACTGGAGCA	GAGTGCAACC	3180
	CGTTCGGTGA	CCGTTTTAGG	TCAACATCAG	GGAGTGCTAA	AAAGCAACGG	TGAAATCAAT	3240
45	AGCGAATTTA	AGCCATCGCC	CGGCAAGGCG	TTGGTCCAGA	GCTTTAACGT	CAATCGCTCT	3300
	GGTCAGGATC	TAAGCAAGTC	ACTGCAACAG	GCACTACATG	CCACGCCGCC	ATCCGCAGAG	3360
	AGTAAACTGC	AATCCATGCT	GGGGCACTTT	GTCAGTGCCG	GGGTGGATAT	GAGTCATCAG	3420
50	AAGGGCGAGA	TCCCGCTGGG	CCGCCAGCGC	GATCCGAATG	ATAAAACCGC	ACTGACCAAA	3480
	TCGCGTTTAA	TTTTAGATAC	CGTGACCATC	GGTGAAGTGC	ATGAACTGGC	CGATAAGGCG	3540
55	AAACTGGTAT	CTGACCATAA	ACCCGATGCC	GATCAGATAA	AACAGCTGCG	CCAGCAGTTC	3600
	GATACGCTGC	GTGAAAAGCG	GTATGAGAGC	AATCCGGTGA	AGCATTACAC	CGATATGGGC	3660
	TTCACCCATA	ATAAGGCGCT	GGAAGCAAAC	TATGATGCGG	TCAAAGCCTT	TATCAATGCC	3720
60	TTTAAGAAAG	AGCACCACGG	CGTCAATCTG	ACCACGCGTA	CCGTACTGGA	ATCACAGGGC	3780
	AGTGCGGAGC	TGGCGAAGAA	GCTCAAGAAT	ACGCTGTTGT	CCCTGGACAG	TGGTGAAAGT	3840
65	ATGAGCTTCA	GCCGGTCATA	TGGCGGGGGC	GTCAGCACTG	TCTTTGTGCC	TACCCTTAGC	3900

- 16 -

	AAGAAGGTGC	CAGTTCCGGT	GATCCCCGGA	GCCGGCATCA	CGCTGGATCG	CGCCTATAAC	3960
	CTGAGCTTCA	GTCGTACCAG	CGGCGGATTG	AACGTCAGTT	TTGGCCGCGA	CGGCGGGGTG	4020
5	AGTGGTAACA	TCATGGTCGC	TACCGGCCAT	GATGTGATGC	CCTATATGAC	CGGTAAGAAA	4080
	ACCAGTGCAG	GTAACGCCAG	TGACTGGTTG	AGCGCAAAAC	ATAAAATCAG	CCCGGACTTG	4140
10	CGTATCGGCG	CTGCTGTGAG	TGGCACCCCTG	CAAGGAACGC	TACAAAACAG	CCTGAAGTTT	4200
	AAGCTGACAG	AGGATGAGCT	GCCTGGCTTT	ATCCATGGCT	TGACGCATGG	CACGTTGACC	4260
	CCGGCAGAAC	TGTTGCAAAA	GGGGATCGAA	CATCAGATGA	AGCAGGGCAG	CAAACGACG	4320
15	TTTAGCGTCG	ATACCTCGGC	AAATCTGGAT	CTGCGTGCCG	GTATCAATCT	GAACGAAGAC	4380
	GGCAGTAAAC	CAAATGGTGT	CACTGCCCCT	GTTTCTGCCG	GGCTAAGTGC	ATCGGCAAAC	4440
20	CTGGCCGCCG	GCTCGCGTGA	ACGCAGCACC	ACCTCTGGCC	AGTTTGGCAG	CACGACTTCG	4500
	GCCAGCAATA	ACCGCCCAAC	CTTCCTCAAC	GGGGTCGGCG	CGGGTGCTAA	CCTGACGGCT	4560
	GCTTTAGGGG	TTGCCCATTC	ATCTACGCAT	GAAGGGAAAC	CGGTCGGGAT	CTTCCCGGCA	4620
25	TTTACCTCGA	CCAATGTTTC	GGCAGCGCTG	GCGCTGGATA	ACCGTACCTC	ACAGAGTATC	4680
	AGCCTGGAAT	TGAAGCGCGC	GGAGCCGGTG	ACCAGCAACG	ATATCAGCGA	GTTGACCTCC	4740
30	ACGCTGGGAA	AACACTTTAA	GGATAGCGCC	ACAACGAAGA	TGCTTGCCGC	TCTCAAAGAG	4800
	TTAGATGACG	CTAAGCCCGC	TGAACAACTG	CATATTTTAC	AGCAGCATTT	CAGTGCAAAA	4860
	GATGTCGTCG	GTGATGAACG	CTACGAGGCG	GTGCGCAACC	TGAAAAAACT	GGTGATACGT	4920
35	CAACAGGCTG	CGGACAGCCA	CAGCATGGAA	TTAGGATCTG	CCAGTCACAG	CACGACCTAC	4980
	AATAATCTGT	CGAGAATAAA	TAATGACGGC	ATTGTCGAGC	TGCTACACAA	ACATTTTCGAT	5040
40	GCGGCATTAC	CAGCAAGCAG	TGCCAAACGT	CTTGGTGAAA	TGATGAATAA	CGATCCGGCA	5100
	CTGAAAGATA	TTATTAAGCA	GCTGCAAAGT	ACGCCGTTCA	GCAGCGCCAG	CGTGTCGATG	5160
	GAGCTGAAAG	ATGGTCTGCG	TGAGCAGACG	GAAAAAGCAA	TACTGGACGG	TAAGGTCGGT	5220
45	CGTGAAGAAG	TGGGAGTACT	TTTCCAGGAT	CGTAACAACT	TGCGTGTTAA	ATCGGTCAGC	5280
	GTCAGTCAGT	CCGTCAGCAA	AAGCGAAGGC	TTCAATACCC	CAGCGCTGTT	ACTGGGGACG	5340
50	AGCAACAGCG	CTGCTATGAG	CATGGAGCGC	AACATCGGAA	CCATTAATTT	TAAATACGGC	5400
	CAGGATCAGA	ACACCCACG	GCGATTTACC	CTGGAGGGTG	GAATAGCTCA	GGCTAATCCG	5460
	CAGGTCGCAT	CTGCGCTTAC	TGATTTGAAG	AAGGAAGGGC	TGGAAATGAA	GAGCTAA	5517

55

This DNA molecule is known as the *dspE* gene for *Erwinia amylovora*. This isolated DNA molecule of the present invention encodes a protein or polypeptide which elicits a plant pathogen's hypersensitive response having an amino acid sequence of SEQ.

ID. No. 28 as follows:

60

- 17 -

	Met	Glu	Leu	Lys	Ser	Leu	Gly	Thr	Glu	His	Lys	Ala	Ala	Val	His	Thr	
	1				5					10					15		
5	Ala	Ala	His	Asn	Pro	Val	Gly	His	Gly	Val	Ala	Leu	Gln	Gln	Gly	Ser	
				20					25					30			
	Ser	Ser	Ser	Ser	Pro	Gln	Asn	Ala	Ala	Ala	Ser	Leu	Ala	Ala	Glu	Gly	
				35				40					45				
10	Lys	Asn	Arg	Gly	Lys	Met	Pro	Arg	Ile	His	Gln	Pro	Ser	Thr	Ala	Ala	
		50					55					60					
	Asp	Gly	Ile	Ser	Ala	Ala	His	Gln	Gln	Lys	Lys	Ser	Phe	Ser	Leu	Arg	
15	65					70					75					80	
	Gly	Cys	Leu	Gly	Thr	Lys	Lys	Phe	Ser	Arg	Ser	Ala	Pro	Gln	Gly	Gln	
					85					90					95		
	Pro	Gly	Thr	Thr	His	Ser	Lys	Gly	Ala	Thr	Leu	Arg	Asp	Leu	Leu	Ala	
20					100				105								
	Arg	Asp	Asp	Gly	Glu	Thr	Gln	His	Glu	Ala	Ala	Ala	Pro	Asp	Ala	Ala	
				115				120					125				
25	Arg	Leu	Thr	Arg	Ser	Gly	Gly	Val	Lys	Arg	Arg	Asn	Met	Asp	Asp	Met	
		130					135						140				
	Ala	Gly	Arg	Pro	Met	Val	Lys	Gly	Gly	Ser	Gly	Glu	Asp	Lys	Val	Pro	
30	145					150					155					160	
	Thr	Gln	Gln	Lys	Arg	His	Gln	Leu	Asn	Asn	Phe	Gly	Gln	Met	Arg	Gln	
					165					170					175		
	Thr	Met	Leu	Ser	Lys	Met	Ala	His	Pro	Ala	Ser	Ala	Asn	Ala	Gly	Asp	
35				180					185					190			
	Arg	Leu	Gln	His	Ser	Pro	Pro	His	Ile	Pro	Gly	Ser	His	His	Glu	Ile	
			195					200					205				
40	Lys	Glu	Glu	Pro	Val	Gly	Ser	Thr	Ser	Lys	Ala	Thr	Thr	Ala	His	Ala	
		210					215					220					
	Asp	Arg	Val	Glu	Ile	Ala	Gln	Glu	Asp	Asp	Asp	Ser	Glu	Phe	Gln	Gln	
45	225					230				235						240	
	Leu	His	Gln	Gln	Arg	Leu	Ala	Arg	Glu	Arg	Glu	Asn	Pro	Pro	Gln	Pro	
					245					250					255		
	Pro	Lys	Leu	Gly	Val	Ala	Thr	Pro	Ile	Ser	Ala	Arg	Phe	Gln	Pro	Lys	
50				260					265					270			
	Leu	Thr	Ala	Val	Ala	Glu	Ser	Val	Leu	Glu	Gly	Thr	Asp	Thr	Thr	Gln	
			275					280					285				
55	Ser	Pro	Leu	Lys	Pro	Gln	Ser	Met	Leu	Lys	Gly	Ser	Gly	Ala	Gly	Val	
		290					295					300					
	Thr	Pro	Leu	Ala	Val	Thr	Leu	Asp	Lys	Gly	Lys	Leu	Gln	Leu	Ala	Pro	
60	305					310					315					320	
	Asp	Asn	Pro	Pro	Ala	Leu	Asn	Thr	Leu	Leu	Lys	Gln	Thr	Leu	Gly	Lys	
					325					330					335		
	Asp	Thr	Gln	His	Tyr	Leu	Ala	His	His	Ala	Ser	Ser	Asp	Gly	Ser	Gln	
65				340					345					350			

- 18 -

	His	Leu	Leu	Leu	Asp	Asn	Lys	Gly	His	Leu	Phe	Asp	Ile	Lys	Ser	Thr	
			355					360					365				
5	Ala	Thr	Ser	Tyr	Ser	Val	Leu	His	Asn	Ser	His	Pro	Gly	Glu	Ile	Lys	
		370					375					380					
	Gly	Lys	Leu	Ala	Gln	Ala	Gly	Thr	Gly	Ser	Val	Ser	Val	Asp	Gly	Lys	
	385					390					395					400	
10	Ser	Gly	Lys	Ile	Ser	Leu	Gly	Ser	Gly	Thr	Gln	Ser	His	Asn	Lys	Thr	
					405					410					415		
	Met	Leu	Ser	Gln	Pro	Gly	Glu	Ala	His	Arg	Ser	Leu	Leu	Thr	Gly	Ile	
15				420					425						430		
	Trp	Gln	His	Pro	Ala	Gly	Ala	Ala	Arg	Pro	Gln	Gly	Glu	Ser	Ile	Arg	
			435					440					445				
20	Leu	His	Asp	Asp	Lys	Ile	His	Ile	Leu	His	Pro	Glu	Leu	Gly	Val	Trp	
		450					455					460					
	Gln	Ser	Ala	Asp	Lys	Asp	Thr	His	Ser	Gln	Leu	Ser	Arg	Gln	Ala	Asp	
25		465				470					475					480	
	Gly	Lys	Leu	Tyr	Ala	Leu	Lys	Asp	Asn	Arg	Thr	Leu	Gln	Asn	Leu	Ser	
					485					490					495		
	Asp	Asn	Lys	Ser	Ser	Glu	Lys	Leu	Val	Asp	Lys	Ile	Lys	Ser	Tyr	Ser	
30				500					505					510			
	Val	Asp	Gln	Arg	Gly	Gln	Val	Ala	Ile	Leu	Thr	Asp	Thr	Pro	Gly	Arg	
			515					520					525				
35	His	Lys	Met	Ser	Ile	Met	Pro	Ser	Leu	Asp	Ala	Ser	Pro	Glu	Ser	His	
		530					535					540					
	Ile	Ser	Leu	Ser	Leu	His	Phe	Ala	Asp	Ala	His	Gln	Gly	Leu	Leu	His	
40		545				550					555					560	
	Gly	Lys	Ser	Glu	Leu	Glu	Ala	Gln	Ser	Val	Ala	Ile	Ser	His	Gly	Arg	
					565					570					575		
	Leu	Val	Val	Ala	Asp	Ser	Glu	Gly	Lys	Leu	Phe	Ser	Ala	Ala	Ile	Pro	
45				580					585					590			
	Lys	Gln	Gly	Asp	Gly	Asn	Glu	Leu	Lys	Met	Lys	Ala	Met	Pro	Gln	His	
			595					600					605				
50	Ala	Leu	Asp	Glu	His	Phe	Gly	His	Asp	His	Gln	Ile	Ser	Gly	Phe	Phe	
		610					615					620					
	His	Asp	Asp	His	Gly	Gln	Leu	Asn	Ala	Leu	Val	Lys	Asn	Asn	Phe	Arg	
55		625				630					635					640	
	Gln	Gln	His	Ala	Cys	Pro	Leu	Gly	Asn	Asp	His	Gln	Phe	His	Pro	Gly	
					645					650					655		
	Trp	Asn	Leu	Thr	Asp	Ala	Leu	Val	Ile	Asp	Asn	Gln	Leu	Gly	Leu	His	
60				660					665					670			
	His	Thr	Asn	Pro	Glu	Pro	His	Glu	Ile	Leu	Asp	Met	Gly	His	Leu	Gly	
			675					680					685				

- 19 -

	Ser	Leu	Ala	Leu	Gln	Glu	Gly	Lys	Leu	His	Tyr	Phe	Asp	Gln	Leu	Thr	
	690						695						700				
5	Lys	Gly	Trp	Thr	Gly	Ala	Glu	Ser	Asp	Cys	Lys	Gln	Leu	Lys	Lys	Gly	
	705					710					715					720	
	Leu	Asp	Gly	Ala	Ala	Tyr	Leu	Leu	Lys	Asp	Gly	Glu	Val	Lys	Arg	Leu	
					725					730					735		
10	Asn	Ile	Asn	Gln	Ser	Thr	Ser	Ser	Ile	Lys	His	Gly	Thr	Glu	Asn	Val	
				740					745						750		
	Phe	Ser	Leu	Pro	His	Val	Arg	Asn	Lys	Pro	Glu	Pro	Gly	Asp	Ala	Leu	
15				755				760						765			
	Gln	Gly	Leu	Asn	Lys	Asp	Asp	Lys	Ala	Gln	Ala	Met	Ala	Val	Ile	Gly	
	770						775						780				
20	Val	Asn	Lys	Tyr	Leu	Ala	Leu	Thr	Glu	Lys	Gly	Asp	Ile	Arg	Ser	Phe	
	785					790					795					800	
	Gln	Ile	Lys	Pro	Gly	Thr	Gln	Gln	Leu	Glu	Arg	Pro	Ala	Gln	Thr	Leu	
					805					810						815	
25	Ser	Arg	Glu	Gly	Ile	Ser	Gly	Glu	Leu	Lys	Asp	Ile	His	Val	Asp	His	
				820					825						830		
	Lys	Gln	Asn	Leu	Tyr	Ala	Leu	Thr	His	Glu	Gly	Glu	Val	Phe	His	Gln	
30				835					840					845			
	Pro	Arg	Glu	Ala	Trp	Gln	Asn	Gly	Ala	Glu	Ser	Ser	Ser	Trp	His	Lys	
				850				855					860				
35	Leu	Ala	Leu	Pro	Gln	Ser	Glu	Ser	Lys	Leu	Lys	Ser	Leu	Asp	Met	Ser	
	865					870					875					880	
	His	Glu	His	Lys	Pro	Ile	Ala	Thr	Phe	Glu	Asp	Gly	Ser	Gln	His	Gln	
					885					890						895	
40	Leu	Lys	Ala	Gly	Gly	Trp	His	Ala	Tyr	Ala	Ala	Pro	Glu	Arg	Gly	Pro	
				900					905						910		
	Leu	Ala	Val	Gly	Thr	Ser	Gly	Ser	Gln	Thr	Val	Phe	Asn	Arg	Leu	Met	
45				915					920					925			
	Gln	Gly	Val	Lys	Gly	Lys	Val	Ile	Pro	Gly	Ser	Gly	Leu	Thr	Val	Lys	
	930						935						940				
50	Leu	Ser	Ala	Gln	Thr	Gly	Gly	Met	Thr	Gly	Ala	Glu	Gly	Arg	Lys	Val	
	945					950					955					960	
	Ser	Ser	Lys	Phe	Ser	Glu	Arg	Ile	Arg	Ala	Tyr	Ala	Phe	Asn	Pro	Thr	
					965					970					975		
55	Met	Ser	Thr	Pro	Arg	Pro	Ile	Lys	Asn	Ala	Ala	Tyr	Ala	Thr	Gln	His	
				980					985						990		
	Gly	Trp	Gln	Gly	Arg	Glu	Gly	Leu	Lys	Pro	Leu	Tyr	Glu	Met	Gln	Gly	
				995				1000					1005				
60	Ala	Leu	Ile	Lys	Gln	Leu	Asp	Ala	His	Asn	Val	Arg	His	Asn	Ala	Pro	
				1010				1015					1020				
65	Gln	Pro	Asp	Leu	Gln	Ser	Lys	Leu	Glu	Thr	Leu	Asp	Leu	Gly	Glu	His	
	1025					1030					1035					1040	

- 20 -

Gly Ala Glu Leu Leu Asn Asp Met Lys Arg Phe Arg Asp Glu Leu Glu
 1045 1050 1055
 5 Gln Ser Ala Thr Arg Ser Val Thr Val Leu Gly Gln His Gln Gly Val
 1060 1065 1070
 Leu Lys Ser Asn Gly Glu Ile Asn Ser Glu Phe Lys Pro Ser Pro Gly
 1075 1080 1085
 10 Lys Ala Leu Val Gln Ser Phe Asn Val Asn Arg Ser Gly Gln Asp Leu
 1090 1095 1100
 Ser Lys Ser Leu Gln Gln Ala Val His Ala Thr Pro Pro Ser Ala Glu
 1105 1110 1115 1120
 15 Ser Lys Leu Gln Ser Met Leu Gly His Phe Val Ser Ala Gly Val Asp
 1125 1130 1135
 Met Ser His Gln Lys Gly Glu Ile Pro Leu Gly Arg Gln Arg Asp Pro
 1140 1145 1150
 20 Asn Asp Lys Thr Ala Leu Thr Lys Ser Arg Leu Ile Leu Asp Thr Val
 1155 1160 1165
 Thr Ile Gly Glu Leu His Glu Leu Ala Asp Lys Ala Lys Leu Val Ser
 1170 1175 1180
 25 Asp His Lys Pro Asp Ala Asp Gln Ile Lys Gln Leu Arg Gln Gln Phe
 1185 1190 1195 1200
 Asp Thr Leu Arg Glu Lys Arg Tyr Glu Ser Asn Pro Val Lys His Tyr
 1205 1210 1215
 30 Thr Asp Met Gly Phe Thr His Asn Lys Ala Leu Glu Ala Asn Tyr Asp
 1220 1225 1230
 Ala Val Lys Ala Phe Ile Asn Ala Phe Lys Lys Glu His His Gly Val
 1235 1240 1245
 40 Asn Leu Thr Thr Arg Thr Val Leu Glu Ser Gln Gly Ser Ala Glu Leu
 1250 1255 1260
 Ala Lys Lys Leu Lys Asn Thr Leu Leu Ser Leu Asp Ser Gly Glu Ser
 1265 1270 1275 1280
 45 Met Ser Phe Ser Arg Ser Tyr Gly Gly Gly Val Ser Thr Val Phe Val
 1285 1290 1295
 Pro Thr Leu Ser Lys Lys Val Pro Val Pro Val Ile Pro Gly Ala Gly
 1300 1305 1310
 50 Ile Thr Leu Asp Arg Ala Tyr Asn Leu Ser Phe Ser Arg Thr Ser Gly
 1315 1320 1325
 Gly Leu Asn Val Ser Phe Gly Arg Asp Gly Gly Val Ser Gly Asn Ile
 1330 1335 1340
 55 Met Val Ala Thr Gly His Asp Val Met Pro Tyr Met Thr Gly Lys Lys
 1345 1350 1355 1360
 Thr Ser Ala Gly Asn Ala Ser Asp Trp Leu Ser Ala Lys His Lys Ile
 1365 1370 1375

- 21 -

	Ser Pro Asp Leu Arg Ile Gly Ala Ala Val Ser Gly Thr Leu Gln Gly	1380	1385	1390
5	Thr Leu Gln Asn Ser Leu Lys Phe Lys Leu Thr Glu Asp Glu Leu Pro	1395	1400	1405
	Gly Phe Ile His Gly Leu Thr His Gly Thr Leu Thr Pro Ala Glu Leu	1410	1415	1420
10	Leu Gln Lys Gly Ile Glu His Gln Met Lys Gln Gly Ser Lys Leu Thr	1425	1430	1435
	Phe Ser Val Asp Thr Ser Ala Asn Leu Asp Leu Arg Ala Gly Ile Asn	1445	1450	1455
15	Leu Asn Glu Asp Gly Ser Lys Pro Asn Gly Val Thr Ala Arg Val Ser	1460	1465	1470
	Ala Gly Leu Ser Ala Ser Ala Asn Leu Ala Ala Gly Ser Arg Glu Arg	1475	1480	1485
20	Ser Thr Thr Ser Gly Gln Phe Gly Ser Thr Thr Ser Ala Ser Asn Asn	1490	1495	1500
	Arg Pro Thr Phe Leu Asn Gly Val Gly Ala Gly Ala Asn Leu Thr Ala	1505	1510	1515
25	Ala Leu Gly Val Ala His Ser Ser Thr His Glu Gly Lys Pro Val Gly	1525	1530	1535
30	Ile Phe Pro Ala Phe Thr Ser Thr Asn Val Ser Ala Ala Leu Ala Leu	1540	1545	1550
	Asp Asn Arg Thr Ser Gln Ser Ile Ser Leu Glu Leu Lys Arg Ala Glu	1555	1560	1565
35	Pro Val Thr Ser Asn Asp Ile Ser Glu Leu Thr Ser Thr Leu Gly Lys	1570	1575	1580
	His Phe Lys Asp Ser Ala Thr Thr Lys Met Leu Ala Ala Leu Lys Glu	1585	1590	1595
40	Leu Asp Asp Ala Lys Pro Ala Glu Gln Leu His Ile Leu Gln Gln His	1605	1610	1615
45	Phe Ser Ala Lys Asp Val Val Gly Asp Glu Arg Tyr Glu Ala Val Arg	1620	1625	1630
	Asn Leu Lys Lys Leu Val Ile Arg Gln Gln Ala Ala Asp Ser His Ser	1635	1640	1645
50	Met Glu Leu Gly Ser Ala Ser His Ser Thr Thr Tyr Asn Asn Leu Ser	1650	1655	1660
	Arg Ile Asn Asn Asp Gly Ile Val Glu Leu Leu His Lys His Phe Asp	1665	1670	1675
55	Ala Ala Leu Pro Ala Ser Ser Ala Lys Arg Leu Gly Glu Met Met Asn	1685	1690	1695
60	Asn Asp Pro Ala Leu Lys Asp Ile Ile Lys Gln Leu Gln Ser Thr Pro	1700	1705	1710
	Phe Ser Ser Ala Ser Val Ser Met Glu Leu Lys Asp Gly Leu Arg Glu	1715	1720	1725

- 22 -

5 Gln Thr Glu Lys Ala Ile Leu Asp Gly Lys Val Gly Arg Glu Glu Val
 1730 1735 1740
 Gly Val Leu Phe Gln Asp Arg Asn Asn Leu Arg Val Lys Ser Val Ser
 1745 1750 1755 1760
 Val Ser Gln Ser Val Ser Lys Ser Glu Gly Phe Asn Thr Pro Ala Leu
 1765 1770 1775
 10 Leu Leu Gly Thr Ser Asn Ser Ala Ala Met Ser Met Glu Arg Asn Ile
 1780 1785 1790
 Gly Thr Ile Asn Phe Lys Tyr Gly Gln Asp Gln Asn Thr Pro Arg Arg
 1795 1800 1805
 Phe Thr Leu Glu Gly Gly Ile Ala Gln Ala Asn Pro Gln Val Ala Ser
 1810 1815 1820
 20 Ala Leu Thr Asp Leu Lys Lys Glu Gly Leu Glu Met Lys Ser
 1825 1830 1835

This protein or polypeptide is about 198 kDa and has a pI of 8.98.

25 The present invention relates to an isolated DNA molecule having a nucleotide sequence of SEQ. ID. No. 29 as follows:

ATGACATCGT CACAGCAGCG GGTGAAAGG TTTTACAGT ATTTCTCCGC CGGGTGTAAG 60
 30 ACGCCCATAC ATCTGAAAGA CGGGGTGTGC GCCCTGTATA ACGAACAAGA TGAGGAGGCG 120
 GCGGTGCTGG AAGTACCGCA ACACAGCGAC AGCCTGTTC TACTGTCCG AATCATTGAG 180
 GCTGACCCAC AAACCTCAAT AACCTGTAT TCGATGCTAT TACAGCTGAA TTTTGAAATG 240
 35 GCGGCCATGC GCGGCTGTTG GCTGGCGCTG GATGAAGTGC ACAACGTGCG TTTATGTTTT 300
 CAGCAGTCGC TGGAGCATCT GGATGAAGCA AGTTTTAGCG ATATCGTTAG CGGCTTCATC 360
 40 GAACATGCGG CAGAAGTGCG TGAGTATATA GCGCAATTAG ACGAGAGTAG CGCGGCATAA 420

This is known as the dspF gene. This isolated DNA molecule of the present invention encodes a hypersensitive response elicitor protein or polypeptide having an amino
 45 acid sequence of SEQ. ID. No. 30 as follows:

 Met Thr Ser Ser Gln Gln Arg Val Glu Arg Phe Leu Gln Tyr Phe Ser
 1 5 10 15
 50 Ala Gly Cys Lys Thr Pro Ile His Leu Lys Asp Gly Val Cys Ala Leu
 20 25 30
 Tyr Asn Glu Gln Asp Glu Glu Ala Ala Val Leu Glu Val Pro Gln His
 35 40 45
 55 Ser Asp Ser Leu Leu Leu His Cys Arg Ile Ile Glu Ala Asp Pro Gln
 50 55 60

- 23 -

Thr Ser Ile Thr Leu Tyr Ser Met Leu Leu Gln Leu Asn Phe Glu Met
 65 70 75 80
 5 Ala Ala Met Arg Gly Cys Trp Leu Ala Leu Asp Glu Leu His Asn Val
 85 90 95
 Arg Leu Cys Phe Gln Gln Ser Leu Glu His Leu Asp Glu Ala Ser Phe
 100 105 110
 10 Ser Asp Ile Val Ser Gly Phe Ile Glu His Ala Ala Glu Val Arg Glu
 115 120 125
 Tyr Ile Ala Gln Leu Asp Glu Ser Ser Ala Ala
 130 135

This protein or polypeptide is about 16 kDa and has a pI of 4.45.

The hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas syringae* has an amino acid sequence corresponding to SEQ. ID.

20 No. 31 as follows:

Met Gln Ser Leu Ser Leu Asn Ser Ser Ser Leu Gln Thr Pro Ala Met
 1 5 10 15
 Ala Leu Val Leu Val Arg Pro Glu Ala Glu Thr Thr Gly Ser Thr Ser
 20 25 30
 25 Ser Lys Ala Leu Gln Glu Val Val Val Lys Leu Ala Glu Glu Leu Met
 35 40 45
 Arg Asn Gly Gln Leu Asp Asp Ser Ser Pro Leu Gly Lys Leu Leu Ala
 50 55 60
 30 Lys Ser Met Ala Ala Asp Gly Lys Ala Gly Gly Gly Ile Glu Asp Val
 65 70 75 80
 Ile Ala Ala Leu Asp Lys Leu Ile His Glu Lys Leu Gly Asp Asn Phe
 85 90 95
 Gly Ala Ser Ala Asp Ser Ala Ser Gly Thr Gly Gln Gln Asp Leu Met
 100 105 110
 35 Thr Gln Val Leu Asn Gly Leu Ala Lys Ser Met Leu Asp Asp Leu Leu
 115 120 125
 Thr Lys Gln Asp Gly Gly Thr Ser Phe Ser Glu Asp Asp Met Pro Met
 130 135 140
 40 Leu Asn Lys Ile Ala Gln Phe Met Asp Asp Asn Pro Ala Gln Phe Pro
 145 150 155 160
 Lys Pro Asp Ser Gly Ser Trp Val Asn Glu Leu Lys Glu Asp Asn Phe
 165 170 175
 Leu Asp Gly Asp Glu Thr Ala Ala Phe Arg Ser Ala Leu Asp Ile Ile
 180 185 190

- 24 -

	Gly	Gln	Gln	Leu	Gly	Asn	Gln	Gln	Ser	Asp	Ala	Gly	Ser	Leu	Ala	Gly	
			195					200					205				
	Thr	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Ser	Ser	Phe	Ser	Asn	Asn	Ser	Ser	
		210					215						220				
5	Val	Met	Gly	Asp	Pro	Leu	Ile	Asp	Ala	Asn	Thr	Gly	Pro	Gly	Asp	Ser	
		225				230					235					240	
	Gly	Asn	Thr	Arg	Gly	Glu	Ala	Gly	Gln	Leu	Ile	Gly	Glu	Leu	Ile	Asp	
					245					250					255		
10	Arg	Gly	Leu	Gln	Ser	Val	Leu	Ala	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Val	
				260					265					270			
	Asn	Thr	Pro	Gln	Thr	Gly	Thr	Ser	Ala	Asn	Gly	Gly	Gln	Ser	Ala	Gln	
			275					280					285				
	Asp	Leu	Asp	Gln	Leu	Leu	Gly	Gly	Leu	Leu	Leu	Lys	Gly	Leu	Glu	Ala	
		290					295					300					
15	Thr	Leu	Lys	Asp	Ala	Gly	Gln	Thr	Gly	Thr	Asp	Val	Gln	Ser	Ser	Ala	
		305				310					315					320	
	Ala	Gln	Ile	Ala	Thr	Leu	Leu	Val	Ser	Thr	Leu	Leu	Gln	Gly	Thr	Arg	
					325					330					335		
20	Asn	Gln	Ala	Ala	Ala												
					340												

This hypersensitive response elicitor polypeptide or protein has a molecular weight of 34-35 kDa. It is rich in glycine (about 13.5%) and lacks cysteine and tyrosine.

- 25 Further information about the hypersensitive response elicitor derived from *Pseudomonas syringae* is found in He, S. Y., H. C. Huang, and A. Collmer, "Pseudomonas syringae pv. syringae Harpin_{PS}: a Protein that is Secreted via the Hrp Pathway and Elicits the Hypersensitive Response in Plants," Cell 73:1255-1266 (1993), which is hereby incorporated by reference. The DNA molecule encoding the
- 30 hypersensitive response elicitor from *Pseudomonas syringae* has a nucleotide sequence corresponding to SEQ. ID. No. 32 as follows:

	ATGCAGAGTC	TCAGTCTTAA	CAGCAGCTCG	CTGCAAACCC	CGGCAATGGC	CCTTGTCTCTG	60
	GTACGTCCTG	AAGCCGAGAC	GA CTGGCAGT	ACGTCGAGCA	AGGCGCTTCA	GGAAGTTGTC	120
35	GTGAAGCTGG	CCGAGGAACT	GATGCGCAAT	GGTCAACTCG	ACGACAGCTC	GCCATTGGGA	180
	AAACTGTTGG	CCAAGTCGAT	GGCCGCAGAT	GGCAAGGCGG	GCGGCGGTAT	TGAGGATGTC	240
	ATCGCTGCGC	TGGACAAGCT	GATCCATGAA	AAGCTCGGTG	ACAACTTCGG	CGCGTCTGCG	300

- 25 -

GACAGCGCCT CGGGTACCGG ACAGCAGGAC CTGATGACTC AGGTGCTCAA TGGCCTGGCC 360
 AAGTCGATGC TCGATGATCT TCTGACCAAG CAGGATGGCG GGACAAGCTT CTCCGAAGAC 420
 GATATGCCGA TGCTGAACAA GATCGCGCAG TTCATGGATG ACAATCCCGC ACAGTTTCCC 480
 AAGCCGGACT CGGGCTCCTG GGTGAACGAA CTCAAGGAAG ACAACTTCCT TGATGGCGAC 540
 5 GAAACGGCTG CGTTCCGTTT GGCACCTCGAC ATCATTGGCC AGCAACTGGG TAATCAGCAG 600
 AGTGACGCTG GCAGTCTGGC AGGGACGGGT GGAGGTCTGG GCACTCCGAG CAGTTTTTCC 660
 AACAACTCGT CCGTGATGGG TGATCCGCTG ATCGACGCCA ATACCGGTCC CGGTGACAGC 720
 GGCAATACCC GTGGTGAAGC GGGGCAACTG ATCGGCGAGC TTATCGACCG TGGCCTGCAA 780
 TCGGTATTGG CCGGTGGTGG ACTGGGCACA CCCGTAAACA CCCCGCAGAC CGGTACGTCG 840
 10 GCGAATGGCG GACAGTCCGC TCAGGATCTT GATCAGTTGC TGGGCGGCTT GCTGCTCAAG 900
 GGCCTGGAGG CAACGCTCAA GGATGCCGGG CAAACAGGCA CCGACGTGCA GTCGAGCGCT 960
 GCGCAAATCG CCACCTTGCT GGTCAGTACG CTGCTGCAAG GCACCCGCAA TCAGGCTGCA 1020
 GCCTGA 1026

15 Another potentially suitable hypersensitive response elicitor from
Pseudomonas syringae is disclosed in U.S. Patent Application Serial No. 09/120,817,
 which is hereby incorporated by reference. The protein has a nucleotide sequence of
 SEQ. ID. No. 33 as follows:

20 TCCAATTTCG TGATTTTGAA ATTGGCAGAT TCATAGAAAC GTTCAGGTGT GGAAATCAGG 60
 CTGAGTGC GC AGATTTTCGTT GATAAGGGTG TGCTACTGGT CATTGTTGGT CATTTCAAGG 120
 CCTCTGAGTG CCGTGC GGAG CAATACCACT CTTCTGCTG GCGTGTGCAC ACTGAGTCGC 180
 25 AGGCATAGGC ATTTTCAGTTC CTTGCGTTGG TTGGGCATAT AAAAAAAGGA ACTTTTAAAA 240
 ACAGTGCAAT GAGATGCCGG CAAAACGGGA ACCGGTCGCT GCGCTTTGCC ACTCACTTCG 300
 30 AGCAAGCTCA ACCCAAACA TCCACATCCC TATCGAACGG ACAGCGATAC GGCCACTTGC 360
 TCTGGTAAAC CCTGGAGCTG GCGTCGGTCC AATTGCCCCAC TTAGCGAGGT AACGCAGCAT 420
 GAGCATCGGC ATCACACCCC GGCCGCAACA GACCACCACG CCACTCGATT TTTCGGCGCT 480
 35 AAGCGGCAAG AGTCCTCAAC CAAACACGTT CGGCGAGCAG AACACTCAGC AAGCGATCGA 540
 CCCGAGTGCA CTGTTGTTCG GCAGCGACAC ACAGAAAGAC GTCAACTTCG GCACGCCCCG 600
 40 CAGCACCGTC CAGAATCCGC AGGACGCCAG CAAGCCCAAC GACAGCCAGT CCAACATCGC 660
 TAAATTGATC AGTGCATTGA TCATGTCGTT GCTGCAGATG CTCACCAACT CCAATAAAAA 720
 GCAGGACACC AATCAGGAAC AGCCTGATAG CCAGGCTCCT TTCCAGAACA ACGGCGGGCT 780

- 26 -

CGGTACACCG TCGGCCGATA GCGGGGGCGG CGGTACACCG GATGCGACAG GTGGCGGCGG 840
 5 CGGTGATACG CCAAGCGCAA CAGGCGGTGG CGGCGGTGAT ACTCCGACCG CAACAGGCGG 900
 TGGCGGCAGC GGTGGCGGCG GCACACCCAC TGCAACAGGT GGCGGCAGCG GTGGCACACC 960
 CACTGCAACA GGCGGTGGCG AGGGTGGCGT AACACCGCAA ATCACTCCGC AGTTGGCCAA 1020
 10 CCCTAACCGT ACCTCAGGTA CTGGCTCGGT GTCGGACACC GCAGGTTCTA CCGAGCAAGC 1080
 CGGCAAGATC AATGTGGTGA AAGACACCAT CAAGGTCGGC GCTGGCGAAG TCTTTGACGG 1140
 15 CCACGGCGCA ACCTTCACTG CCGACAAATC TATGGGTAAC GGAGACCAGG GCGAAAATCA 1200
 GAAGCCCATG TTCGAGCTGG CTGAAGGCGC TACGTTGAAG AATGTGAACC TGGGTGAGAA 1260
 CGAGGTCGAT GGCATCCACG TGAAAGCCAA AAACGCTCAG GAAGTCACCA TTGACAACGT 1320
 20 GCATGCCCAG AACGTCGGTG AAGACCTGAT TACGGTCAAA GGCGAGGGAG GCGCAGCGGT 1380
 CACTAATCTG AACATCAAGA ACAGCAGTGC CAAAGGTGCA GACGACAAGG TTGTCCAGCT 1440
 CAACGCCAAC ACTCACTTGA AAATCGACAA CTTCAAGGCC GACGATTTCG GCACGATGGT 1500
 25 TCGCACCAAC GGTGGCAAGC AGTTTGATGA CATGAGCATC GAGCTGAACG GCATCGAAGC 1560
 TAACCACGGC AAGTTCGCCC TGGTGAAAAG CGACAGTGAC GATCTGAAGC TGGCAACGGG 1620
 30 CAACATCGCC ATGACCGACG TCAAACACGC CTACGATAAA ACCCAGGCAT CGACCCAACA 1680
 CACCGAGCTT TGAATCCAGA CAAGTAGCTT GAAAAAAGGG GGTGGACTC 1729

35 This DNA molecule is known as the dspE gene for *Pseudomonas syringae*. This isolated DNA molecule of the present invention encodes a protein or polypeptide which elicits a plant pathogen's hypersensitive response having an amino acid sequence of SEQ. ID. No. 34 as follows:

40 Met Ser Ile Gly Ile Thr Pro Arg Pro Gln Gln Thr Thr Thr Pro Leu
 1 5 10 15
 Asp Phe Ser Ala Leu Ser Gly Lys Ser Pro Gln Pro Asn Thr Phe Gly
 20 25 30
 45 Glu Gln Asn Thr Gln Gln Ala Ile Asp Pro Ser Ala Leu Leu Phe Gly
 35 40 45
 Ser Asp Thr Gln Lys Asp Val Asn Phe Gly Thr Pro Asp Ser Thr Val
 50 55 60
 Gln Asn Pro Gln Asp Ala Ser Lys Pro Asn Asp Ser Gln Ser Asn Ile
 65 70 75 80
 55 Ala Lys Leu Ile Ser Ala Leu Ile Met Ser Leu Leu Gln Met Leu Thr
 85 90 95

- 27 -

	Asn	Ser	Asn	Lys	Lys	Gln	Asp	Thr	Asn	Gln	Glu	Gln	Pro	Asp	Ser	Gln	
				100					105					110			
5	Ala	Pro	Phe	Gln	Asn	Asn	Gly	Gly	Leu	Gly	Thr	Pro	Ser	Ala	Asp	Ser	
			115				120						125				
	Gly	Gly	Gly	Gly	Thr	Pro	Asp	Ala	Thr	Gly	Gly	Gly	Gly	Gly	Asp	Thr	
			130				135						140				
10	Pro	Ser	Ala	Thr	Gly	Gly	Gly	Gly	Gly	Asp	Thr	Pro	Thr	Ala	Thr	Gly	
						150					155					160	
	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Thr	Pro	Thr	Ala	Thr	Gly	Gly	Gly	
					165					170					175		
15	Ser	Gly	Gly	Thr	Pro	Thr	Ala	Thr	Gly	Gly	Gly	Glu	Gly	Gly	Val	Thr	
				180					185					190			
20	Pro	Gln	Ile	Thr	Pro	Gln	Leu	Ala	Asn	Pro	Asn	Arg	Thr	Ser	Gly	Thr	
			195					200					205				
	Gly	Ser	Val	Ser	Asp	Thr	Ala	Gly	Ser	Thr	Glu	Gln	Ala	Gly	Lys	Ile	
			210				215					220					
25	Asn	Val	Val	Lys	Asp	Thr	Ile	Lys	Val	Gly	Ala	Gly	Glu	Val	Phe	Asp	
						230					235					240	
	Gly	His	Gly	Ala	Thr	Phe	Thr	Ala	Asp	Lys	Ser	Met	Gly	Asn	Gly	Asp	
					245					250					255		
30	Gln	Gly	Glu	Asn	Gln	Lys	Pro	Met	Phe	Glu	Leu	Ala	Glu	Gly	Ala	Thr	
				260					265					270			
35	Leu	Lys	Asn	Val	Asn	Leu	Gly	Glu	Asn	Glu	Val	Asp	Gly	Ile	His	Val	
			275					280					285				
	Lys	Ala	Lys	Asn	Ala	Gln	Glu	Val	Thr	Ile	Asp	Asn	Val	His	Ala	Gln	
			290				295					300					
40	Asn	Val	Gly	Glu	Asp	Leu	Ile	Thr	Val	Lys	Gly	Glu	Gly	Gly	Ala	Ala	
						310					315					320	
	Val	Thr	Asn	Leu	Asn	Ile	Lys	Asn	Ser	Ser	Ala	Lys	Gly	Ala	Asp	Asp	
					325					330					335		
45	Lys	Val	Val	Gln	Leu	Asn	Ala	Asn	Thr	His	Leu	Lys	Ile	Asp	Asn	Phe	
				340					345					350			
50	Lys	Ala	Asp	Asp	Phe	Gly	Thr	Met	Val	Arg	Thr	Asn	Gly	Gly	Lys	Gln	
			355					360					365				
	Phe	Asp	Asp	Met	Ser	Ile	Glu	Leu	Asn	Gly	Ile	Glu	Ala	Asn	His	Gly	
			370				375					380					
55	Lys	Phe	Ala	Leu	Val	Lys	Ser	Asp	Ser	Asp	Asp	Leu	Lys	Leu	Ala	Thr	
						390					395					400	
	Gly	Asn	Ile	Ala	Met	Thr	Asp	Val	Lys	His	Ala	Tyr	Asp	Lys	Thr	Gln	
					405					410						415	

- 28 -

Ala Ser Thr Gln His Thr Glu Leu
420

5

This protein or polypeptide is about 42.9 kDa.

The hypersensitive response elicitor polypeptide or protein derived
from *Pseudomonas solanacearum* has an amino acid sequence corresponding to SEQ.
10 ID. No. 35 as follows:

	Met	Ser	Val	Gly	Asn	Ile	Gln	Ser	Pro	Ser	Asn	Leu	Pro	Gly	Leu	Gln	
	1				5					10					15		
15	Asn	Leu	Asn	Leu	Asn	Thr	Asn	Thr	Asn	Ser	Gln	Gln	Ser	Gly	Gln	Ser	
				20					25					30			
	Val	Gln	Asp	Leu	Ile	Lys	Gln	Val	Glu	Lys	Asp	Ile	Leu	Asn	Ile	Ile	
			35				40						45				
	Ala	Ala	Leu	Val	Gln	Lys	Ala	Ala	Gln	Ser	Ala	Gly	Gly	Asn	Thr	Gly	
		50					55					60					
20	Asn	Thr	Gly	Asn	Ala	Pro	Ala	Lys	Asp	Gly	Asn	Ala	Asn	Ala	Gly	Ala	
	65					70					75					80	
	Asn	Asp	Pro	Ser	Lys	Asn	Asp	Pro	Ser	Lys	Ser	Gln	Ala	Pro	Gln	Ser	
					85					90					95		
25	Ala	Asn	Lys	Thr	Gly	Asn	Val	Asp	Asp	Ala	Asn	Asn	Gln	Asp	Pro	Met	
				100					105					110			
	Gln	Ala	Leu	Met	Gln	Leu	Leu	Glu	Asp	Leu	Val	Lys	Leu	Leu	Lys	Ala	
			115					120					125				
	Ala	Leu	His	Met	Gln	Gln	Pro	Gly	Gly	Asn	Asp	Lys	Gly	Asn	Gly	Val	
		130					135					140					
30	Gly	Gly	Ala	Asn	Gly	Ala	Lys	Gly	Ala	Gly	Gly	Gln	Gly	Gly	Leu	Ala	
	145					150					155					160	
	Glu	Ala	Leu	Gln	Glu	Ile	Glu	Gln	Ile	Leu	Ala	Gln	Leu	Gly	Gly	Gly	
				165					170						175		
35	Gly	Ala	Gly	Ala	Gly	Gly	Ala	Gly	Gly	Gly	Val	Gly	Gly	Ala	Gly	Gly	
				180					185					190			
	Ala	Asp	Gly	Gly	Ser	Gly	Ala	Gly	Gly	Ala	Gly	Gly	Ala	Asn	Gly	Ala	
			195					200					205				
	Asp	Gly	Gly	Asn	Gly	Val	Asn	Gly	Asn	Gln	Ala	Asn	Gly	Pro	Gln	Asn	
		210					215						220				

- 29 -

	Ala	Gly	Asp	Val	Asn	Gly	Ala	Asn	Gly	Ala	Asp	Asp	Gly	Ser	Glu	Asp	
	225					230					235					240	
	Gln	Gly	Gly	Leu	Thr	Gly	Val	Leu	Gln	Lys	Leu	Met	Lys	Ile	Leu	Asn	
					245					250					255		
5	Ala	Leu	Val	Gln	Met	Met	Gln	Gln	Gly	Gly	Leu	Gly	Gly	Gly	Asn	Gln	
				260					265					270			
	Ala	Gln	Gly	Gly	Ser	Lys	Gly	Ala	Gly	Asn	Ala	Ser	Pro	Ala	Ser	Gly	
			275					280					285				
10	Ala	Asn	Pro	Gly	Ala	Asn	Gln	Pro	Gly	Ser	Ala	Asp	Asp	Gln	Ser	Ser	
		290					295					300					
	Gly	Gln	Asn	Asn	Leu	Gln	Ser	Gln	Ile	Met	Asp	Val	Val	Lys	Glu	Val	
	305					310					315					320	
	Val	Gln	Ile	Leu	Gln	Gln	Met	Leu	Ala	Ala	Gln	Asn	Gly	Gly	Ser	Gln	
				325					330						335		
15	Gln	Ser	Thr	Ser	Thr	Gln	Pro	Met									
				340													

It is encoded by a DNA molecule having a nucleotide sequence corresponding SEQ.
ID. No. 36 as follows:

	ATGTCAGTCG	GAAACATCCA	GAGCCCGTCG	AACCTCCCGG	GTCTGCAGAA	CCTGAACCTC	60
20	AACACCAACA	CCAACAGCCA	GCAATCGGGC	CAGTCCGTGC	AAGACCTGAT	CAAGCAGGTC	120
	GAGAAGGACA	TCCTCAACAT	CATCGCAGCC	CTCGTGCAGA	AGGCCGCACA	GTCGGCGGGC	180
	GGCAACACCG	GTAACACCGG	CAACGCGCCG	GCGAAGGACG	GCAATGCCAA	CGCGGGCGCC	240
	AACGACCCGA	GCAAGAACGA	CCCAGCAAG	AGCCAGGCTC	CGCAGTCGGC	CAACAAGACC	300
	GGCAACGTCG	ACGACGCCAA	CAACCAGGAT	CCGATGCAAG	CGCTGATGCA	GCTGCTGGAA	360
25	GACCTGGTGA	AGCTGCTGAA	GGCGGCCCTG	CACATGCAGC	AGCCCGGCGG	CAATGACAAG	420
	GGCAACGGCG	TGGGCGGTGC	CAACGGCGCC	AAGGGTGCCG	GCGGCCAGGG	CGGCCTGGCC	480
	GAAGCGCTGC	AGGAGATCGA	GCAGATCCTC	GCCCAGCTCG	GCGGCGGCGG	TGCTGGCGCC	540
	GGCGGCGCGG	GTGGCGGTGT	CGGCGGTGCT	GGTGGCGCGG	ATGGCGGCTC	CGGTGCGGGT	600
	GGCGCAGGCG	GTGCGAACGG	CGCCGACGGC	GGCAATGGCG	TGAACGGCAA	CCAGGCGAAC	660
30	GGCCCGCAGA	ACGCAGGCGA	TGTCAACGGT	GCCAACGGCG	CGGATGACGG	CAGCGAAGAC	720
	CAGGGCGGCC	TCACCGGCGT	GCTGCAAAAG	CTGATGAAGA	TCCTGAACGC	GCTGGTGCAG	780
	ATGATGCAGC	AAGGCGGCCT	CGGCGGCGGC	AACCAGGCGC	AGGGCGGCTC	GAAGGGTGCC	840
	GGCAACGCCT	CGCCGGCTTC	CGGCGCGAAC	CCGGGCGCGA	ACCAGCCCGG	TTCGGCGGAT	900

- 30 -

GATCAATCGT CCGGCCAGAA CAATCTGCAA TCCCAGATCA TGGATGTGGT GAAGGAGGTC 960
 GTCCAGATCC TGCAGCAGAT GCTGGCGGCG CAGAACGGCG GCAGCCAGCA GTCCACCTCG 1020
 ACGCAGCCGA TGTA 1035

5 Further information regarding the hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas solanacearum* is set forth in Arlat, M., F. Van Gijsegem, J. C. Huet, J. C. Pemollet, and C. A. Boucher, "PopA1, a Protein which Induces a Hypersensitive-like Response in Specific Petunia Genotypes, is Secreted
 10 via the Hrp Pathway of *Pseudomonas solanacearum*," EMBO J. 13:543-533 (1994), which is hereby incorporated by reference.

The hypersensitive response elicitor polypeptide or protein from *Xanthomonas campestris* pv. glycines has an amino acid sequence corresponding to SEQ. ID. No. 37 as follows:

15 Thr Leu Ile Glu Leu Met Ile Val Val Ala Ile Ile Ala Ile Leu Ala
 1 5 10 15
 Ala Ile Ala Leu Pro Ala Tyr Gln Asp Tyr
 20 25

20 This sequence is an amino terminal sequence having only 26 residues from the hypersensitive response elicitor polypeptide or protein of *Xanthomonas campestris* pv. glycines. It matches with fimbrial subunit proteins determined in other
 25 *Xanthomonas campestris* pathovars.

The hypersensitive response elicitor polypeptide or protein from *Xanthomonas campestris* pv. *pelargonii* is heat stable, protease sensitive, and has a molecular weight of 20 kDa. It includes an amino acid sequence corresponding to SEQ. ID. No. 38 as follows:

30 Ser Ser Gln Gln Ser Pro Ser Ala Gly Ser Glu Gln Gln Leu Asp Gln
 1 5 10 15
 Leu Leu Ala Met
 20

35 Isolation of *Erwinia carotovora* hypersensitive response elicitor protein or polypeptide is described in Cui et al., "The RsmA Mutants of *Erwinia carotovora*

- 31 -

subsp. *carotovora* Strain Ecc71 Overexpress *hrp* N_{Ecc} and Elicit a Hypersensitive Reaction-like Response in Tobacco Leaves," MPMI, 9(7):565-73 (1996), which is hereby incorporated by reference. The hypersensitive response elicitor protein or polypeptide of *Erwinia stewartii* is set forth in Ahmad et al., "Harpin is Not
 5 Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," 8th Int'l. Cong. Molec. Plant-Microbe Interact., July 14-19, 1996 and Ahmad, et al., "Harpin is Not Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," Ann. Mtg. Am. Phytopath. Soc., July 27-31, 1996, which are hereby incorporated by reference.

Hypersensitive response elicitor proteins or polypeptides from
 10 *Phytophthora parasitica*, *Phytophthora cryptogea*, *Phytophthora cinnamoni*, *Phytophthora capsici*, *Phytophthora megasperma*, and *Phytophthora citrophthora* are described in Kaman, et al., "Extracellular Protein Elicitors from Phytophthora: Most Specificity and Induction of Resistance to Bacterial and Fungal Phytopathogens," Molec. Plant-Microbe Interact., 6(1):15-25 (1993), Ricci et al., "Structure and
 15 Activity of Proteins from Pathogenic Fungi Phytophthora Eliciting Necrosis and Acquired Resistance in Tobacco," Eur. J. Biochem., 183:555-63 (1989), Ricci et al., "Differential Production of Parasiticein, and Elicitor of Necrosis and Resistance in Tobacco, by Isolates of *Phytophthora parasitica*," Plant Path. 41:298-307 (1992), Baillreul et al, "A New Elicitor of the Hypersensitive Response in Tobacco: A
 20 Fungal Glycoprotein Elicits Cell Death, Expression of Defence Genes, Production of Salicylic Acid, and Induction of Systemic Acquired Resistance," Plant J., 8(4):551-60 (1995), and Bonnet et al., "Acquired Resistance Triggered by Elicitors in Tobacco and Other Plants," Eur. J. Plant Path., 102:181-92 (1996), which are hereby incorporated by reference.

25 Another hypersensitive response elicitor in accordance with the present invention is from *Clavibacter michiganensis* subsp. *sepedonicus* which is fully described in U.S. Patent Application Serial No. 09/136,625, which is hereby incorporated by reference.

The above elicitors are exemplary. Other elicitors can be identified by
 30 growing fungi or bacteria that elicit a hypersensitive response under conditions which genes encoding an elicitor are expressed. Cell-free preparations from culture

- 32 -

supernatants can be tested for elicitor activity (i.e. local necrosis) by using them to infiltrate appropriate plant tissues.

Fragments of the above hypersensitive response elicitor polypeptides or proteins as well as fragments of full length elicitors from other pathogens are encompassed by the present invention.

Suitable fragments can be produced by several means. In the first, subclones of the gene encoding a known elicitor protein are produced by conventional molecular genetic manipulation by subcloning gene fragments. The subclones then are expressed *in vitro* or *in vivo* in bacterial cells to yield a smaller protein or peptide that can be tested for elicitor activity according to the procedure described below.

As an alternative, fragments of an elicitor protein can be produced by digestion of a full-length elicitor protein with proteolytic enzymes like chymotrypsin or *Staphylococcus* proteinase A, or trypsin. Different proteolytic enzymes are likely to cleave elicitor proteins at different sites based on the amino acid sequence of the elicitor protein. Some of the fragments that result from proteolysis may be active elicitors of resistance.

In another approach, based on knowledge of the primary structure of the protein, fragments of the elicitor protein gene may be synthesized by using the PCR technique together with specific sets of primers chosen to represent particular portions of the protein. These then would be cloned into an appropriate vector for expression of a truncated peptide or protein.

Chemical synthesis can also be used to make suitable fragments. Such a synthesis is carried out using known amino acid sequences for the elicitor being produced. Alternatively, subjecting a full length elicitor to high temperatures and pressures will produce fragments. These fragments can then be separated by conventional procedures (e.g., chromatography, SDS-PAGE).

An example of suitable fragments of a hypersensitive response elicitor which do not elicit a hypersensitive response include fragments of the *Erwinia amylovora* hypersensitive response elicitor. Suitable fragments include a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 23, an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 23, or an internal fragment of the amino acid sequence of SEQ. ID. No. 23. The C-terminal fragment of the amino acid

- 33 -

sequence of SEQ. ID. No. 23 can span the following amino acids of SEQ. ID. No. 23: 169 and 403, 210 and 403, 267 and 403, or 343 and 403. The internal fragment of the amino acid sequence of SEQ. ID. No. 23 can span the following amino acids of SEQ. ID. No. 23: 105 and 179, 137 and 166, 121 and 150, or 137 and 156. Other suitable fragments can be identified in accordance with the present invention.

Another example of a useful fragment of a hypersensitive response elicitor which fragment does not itself elicit a hypersensitive response is the protein fragment containing amino acids 190 to 294 of the amino acid sequence (SEQ. ID. No. 31) for the *Pseudomonas syringae* pv. *syringae* hypersensitive response elicitor.

This fragment is useful in imparting disease resistance and enhancing plant growth.

Yet another example of a useful fragment of a hypersensitive response elicitor is the peptide having an amino acid sequence corresponding to SEQ. ID. No. 39. This peptide is derived from the hypersensitive response eliciting glycoprotein of *Phytophthora megasperma* and enhances plant growth.

Variants may be made by, for example, the deletion or addition of amino acids that have minimal influence on the properties, secondary structure, and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification, or identification of the polypeptide.

The fragment of the present invention is preferably in isolated form (i.e. separated from its host organism) and more preferably produced in purified form (preferably at least about 60%, more preferably 80%, pure) by conventional techniques. Typically, the fragment of the present invention is produced but not secreted into the growth medium of recombinant host cells. Alternatively, the protein or polypeptide of the present invention is secreted into growth medium. In the case of unsecreted protein, to isolate the protein fragment, the host cell (e.g., *E. coli*) carrying a recombinant plasmid is propagated, lysed by sonication, heat, or chemical treatment, and the homogenate is centrifuged to remove bacterial debris. The supernatant is then subjected to heat treatment and the fragment is separated by centrifugation. The supernatant fraction containing the fragment is subjected to gel filtration in an

- 34 -

appropriately sized dextran or polyacrylamide column to separate the fragment. If necessary, the protein fraction may be further purified by ion exchange or HPLC.

The DNA molecule encoding the fragment of the hypersensitive response elicitor polypeptide or protein can be incorporated in cells using conventional recombinant DNA technology. Generally, this involves inserting the DNA molecule into an expression system to which the DNA molecule is heterologous (i.e. not normally present). The heterologous DNA molecule is inserted into the expression system or vector in proper sense orientation and correct reading frame. The vector contains the necessary elements for the transcription and translation of the inserted protein-coding sequences.

U.S. Patent No. 4,237,224 to Cohen and Boyer, which is hereby incorporated by reference, describes the production of expression systems in the form of recombinant plasmids using restriction enzyme cleavage and ligation with DNA ligase. These recombinant plasmids are then introduced by means of transformation and replicated in unicellular cultures including procaryotic organisms and eucaryotic cells grown in tissue culture.

Recombinant genes may also be introduced into viruses, such as vaccinia virus. Recombinant viruses can be generated by transfection of plasmids into cells infected with virus.

Suitable vectors include, but are not limited to, the following viral vectors such as lambda vector system gt11, gt WES.tB, Charon 4, and plasmid vectors such as pBR322, pBR325, pACYC177, pACYC1084, pUC8, pUC9, pUC18, pUC19, pLG339, pR290, pKC37, pKC101, SV 40, pBluescript II SK +/- or KS +/- (see "Stratagene Cloning Systems" Catalog (1993) from Stratagene, La Jolla, Calif, which is hereby incorporated by reference), pQE, pIH821, pGEX, pET series (see F.W. Studier et. al., "Use of T7 RNA Polymerase to Direct Expression of Cloned Genes," Gene Expression Technology vol. 185 (1990), which is hereby incorporated by reference), and any derivatives thereof. Recombinant molecules can be introduced into cells via transformation, particularly transduction, conjugation, mobilization, or electroporation. The DNA sequences are cloned into the vector using standard cloning procedures in the art, as described by Sambrook et al., Molecular Cloning: A

- 35 -

Laboratory Manual, Cold Springs Laboratory, Cold Springs Harbor, New York (1989), which is hereby incorporated by reference.

A variety of host-vector systems may be utilized to express the protein-encoding sequence(s). Primarily, the vector system must be compatible with the host cell used. Host-vector systems include but are not limited to the following: bacteria transformed with bacteriophage DNA, plasmid DNA, or cosmid DNA; microorganisms such as yeast containing yeast vectors; mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); and plant cells infected by bacteria. The expression elements of these vectors vary in their strength and specificities. Depending upon the host-vector system utilized, any one of a number of suitable transcription and translation elements can be used.

Different genetic signals and processing events control many levels of gene expression (e.g., DNA transcription and messenger RNA (mRNA) translation).

Transcription of DNA is dependent upon the presence of a promotor which is a DNA sequence that directs the binding of RNA polymerase and thereby promotes mRNA synthesis. The DNA sequences of eucaryotic promoters differ from those of procaryotic promoters. Furthermore, eucaryotic promoters and accompanying genetic signals may not be recognized in or may not function in a procaryotic system, and, further, procaryotic promoters are not recognized and do not function in eucaryotic cells.

Similarly, translation of mRNA in procaryotes depends upon the presence of the proper procaryotic signals which differ from those of eucaryotes. Efficient translation of mRNA in procaryotes requires a ribosome binding site called the Shine-Dalgarno ("SD") sequence on the mRNA. This sequence is a short nucleotide sequence of mRNA that is located before the start codon, usually AUG, which encodes the amino-terminal methionine of the protein. The SD sequences are complementary to the 3'-end of the 16S rRNA (ribosomal RNA) and probably promote binding of mRNA to ribosomes by duplexing with the rRNA to allow correct positioning of the ribosome. For a review on maximizing gene expression, see Roberts and Lauer, Methods in Enzymology, 68:473 (1979), which is hereby incorporated by reference.

- 36 -

Promoters vary in their "strength" (i.e. their ability to promote transcription). For the purposes of expressing a cloned gene, it is desirable to use strong promoters in order to obtain a high level of transcription and, hence, expression of the gene. Depending upon the host cell system utilized, any one of a number of suitable promoters may be used. For instance, when cloning in *E. coli*, its bacteriophages, or plasmids, promoters such as the T7 phage promoter, *lac* promoter, *trp* promoter, *recA* promoter, ribosomal RNA promoter, the P_R and P_L promoters of coliphage lambda and others, including but not limited, to *lacUV5*, *ompF*, *bla*, *lpp*, and the like, may be used to direct high levels of transcription of adjacent DNA segments. Additionally, a hybrid *trp-lacUV5 (tac)* promoter or other *E. coli* promoters produced by recombinant DNA or other synthetic DNA techniques may be used to provide for transcription of the inserted gene.

Bacterial host cell strains and expression vectors may be chosen which inhibit the action of the promoter unless specifically induced. In certain operations, the addition of specific inducers is necessary for efficient transcription of the inserted DNA. For example, the *lac* operon is induced by the addition of lactose or IPTG (isopropylthio-beta-D-galactoside). A variety of other operons, such as *trp*, *pro*, etc., are under different controls.

Specific initiation signals are also required for efficient gene transcription and translation in procaryotic cells. These transcription and translation initiation signals may vary in "strength" as measured by the quantity of gene specific messenger RNA and protein synthesized, respectively. The DNA expression vector, which contains a promoter, may also contain any combination of various "strong" transcription and/or translation initiation signals. For instance, efficient translation in *E. coli* requires an SD sequence about 7-9 bases 5' to the initiation codon ("ATG") to provide a ribosome binding site. Thus, any SD-ATG combination that can be utilized by host cell ribosomes may be employed. Such combinations include but are not limited to the SD-ATG combination from the *cro* gene or the *N* gene of coliphage lambda, or from the *E. coli* tryptophan E, D, C, B or A genes. Additionally, any SD-ATG combination produced by recombinant DNA or other techniques involving incorporation of synthetic nucleotides may be used.

- 37 -

Once the isolated DNA molecule encoding the fragment of a hypersensitive response elicitor polypeptide or protein has been cloned into an expression system, it is ready to be incorporated into a host cell. Such incorporation can be carried out by the various forms of transformation noted above, depending upon the vector/host cell system. Suitable host cells include, but are not limited to, bacteria, virus, yeast, mammalian cells, insect, plant, and the like.

The present invention further relates to methods of imparting disease resistance to plants, enhancing plant growth, and/or effecting insect control for plants. These methods involve applying the fragment of a hypersensitive response elicitor polypeptide or protein which does not elicit a hypersensitive response in a non-infectious form to all or part of a plant or a plant seed under conditions effective for the fragment to impart disease resistance, enhance growth, and/or control insects. Alternatively, these fragments of a hypersensitive response elicitor protein or polypeptide can be applied to plants such that seeds recovered from such plants themselves are able to impart disease resistance in plants, to enhance plant growth, and/or to effect insect control.

As an alternative to applying a fragment of a hypersensitive response elicitor polypeptide or protein to plants or plant seeds in order to impart disease resistance in plants, to effect plant growth, and/or to control insects on the plants or plants grown from the seeds, transgenic plants or plant seeds can be utilized. When utilizing transgenic plants, this involves providing a transgenic plant transformed with a DNA molecule encoding a fragment of a hypersensitive response elicitor polypeptide or protein, which fragment does not elicit a hypersensitive response, and growing the plant under conditions effective to permit that DNA molecule to impart disease resistance to plants, to enhance plant growth, and/or to control insects. Alternatively, a transgenic plant seed transformed with a DNA molecule encoding a fragment of a hypersensitive response elicitor polypeptide or protein which fragment does not elicit a hypersensitive response can be provided and planted in soil. A plant is then propagated from the planted seed under conditions effective to permit that DNA molecule to impart disease resistance to plants, to enhance plant growth, and/or to control insects.

- 38 -

The embodiment of the present invention where the hypersensitive response elicitor polypeptide or protein is applied to the plant or plant seed can be carried out in a number of ways, including: 1) application of an isolated fragment or 2) application of bacteria which do not cause disease and are transformed with a gene encoding the fragment. In the latter embodiment, the fragment can be applied to plants or plant seeds by applying bacteria containing the DNA molecule encoding the fragment of the hypersensitive response elicitor polypeptide or protein which fragment does not elicit a hypersensitive response. Such bacteria must be capable of secreting or exporting the fragment so that the fragment can contact plant or plant seed cells. In these embodiments, the fragment is produced by the bacteria *in planta* or on seeds or just prior to introduction of the bacteria to the plants or plant seeds.

The methods of the present invention can be utilized to treat a wide variety of plants or their seeds to impart disease resistance, enhance growth, and/or control insects. Suitable plants include dicots and monocots. More particularly, useful crop plants can include: alfalfa, rice, wheat, barley, rye, cotton, sunflower, peanut, corn, potato, sweet potato, bean, pea, chicory, lettuce, endive, cabbage, brussel sprout, beet, parsnip, turnip, cauliflower, broccoli, radish, spinach, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane. Examples of suitable ornamental plants are: *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

With regard to the use of the fragments of the hypersensitive response elicitor protein or polypeptide of the present invention in imparting disease resistance, absolute immunity against infection may not be conferred, but the severity of the disease is reduced and symptom development is delayed. Lesion number, lesion size, and extent of sporulation of fungal pathogens are all decreased. This method of imparting disease resistance has the potential for treating previously untreatable diseases, treating diseases systemically which might not be treated separately due to cost, and avoiding the use of infectious agents or environmentally harmful materials.

The method of imparting pathogen resistance to plants in accordance with the present invention is useful in imparting resistance to a wide variety of

- 39 -

pathogens including viruses, bacteria, and fungi. Resistance, *inter alia*, to the following viruses can be achieved by the method of the present invention: *Tobacco mosaic virus* and *Tomato mosaic virus*. Resistance, *inter alia*, to the following bacteria can also be imparted to plants in accordance with present invention:

- 5 *Pseudomonas solanacearum*, *Pseudomonas syringae* pv. *tabaci*, and *Xanthomonas campestris* pv. *pelargonii*. Plants can be made resistant, *inter alia*, to the following fungi by use of the method of the present invention: *Fusarium oxysporum* and *Phytophthora infestans*.

- With regard to the use of the fragments of the hypersensitive response
- 10 elicitor protein or polypeptide of the present invention to enhance plant growth, various forms of plant growth enhancement or promotion can be achieved. This can occur as early as when plant growth begins from seeds or later in the life of a plant. For example, plant growth according to the present invention encompasses greater yield, increased quantity of seeds produced, increased percentage of seeds
- 15 germinated, increased plant size, greater biomass, more and bigger fruit, earlier fruit coloration, and earlier fruit and plant maturation. As a result, the present invention provides significant economic benefit to growers. For example, early germination and early maturation permit crops to be grown in areas where short growing seasons would otherwise preclude their growth in that locale. Increased percentage of seed
- 20 germination results in improved crop stands and more efficient seed use. Greater yield, increased size, and enhanced biomass production allow greater revenue generation from a given plot of land.

- Another aspect of the present invention is directed to effecting any form of insect control for plants. For example, insect control according to the present
- 25 invention encompasses preventing insects from contacting plants to which the hypersensitive response elicitor has been applied, preventing direct insect damage to plants by feeding injury, causing insects to depart from such plants, killing insects proximate to such plants, interfering with insect larval feeding on such plants, preventing insects from colonizing host plants, preventing colonizing insects from
- 30 releasing phytotoxins, etc. The present invention also prevents subsequent disease damage to plants resulting from insect infection.

- 40 -

The present invention is effective against a wide variety of insects. European corn borer is a major pest of corn (dent and sweet corn) but also feeds on over 200 plant species including green, wax, and lima beans and edible soybeans, peppers, potato, and tomato plus many weed species. Additional insect larval feeding pests which damage a wide variety of vegetable crops include the following: beet armyworm, cabbage looper, corn ear worm, fall armyworm, diamondback moth, cabbage root maggot, onion maggot, seed corn maggot, pickleworm (melonworm), pepper maggot, tomato pinworm, and maggots. Collectively, this group of insect pests represents the most economically important group of pests for vegetable production worldwide.

The method of the present invention involving application of the fragment of a hypersensitive response elicitor polypeptide or protein, which fragment does not elicit a hypersensitive response, can be carried out through a variety of procedures when all or part of the plant is treated, including leaves, stems, roots, propagules (e.g., cuttings), etc. This may (but need not) involve infiltration of the fragment of the hypersensitive response elicitor polypeptide or protein into the plant. Suitable application methods include high or low pressure spraying, injection, and leaf abrasion proximate to when elicitor application takes place. When treating plant seeds or propagules (e.g., cuttings), in accordance with the application embodiment of the present invention, the fragment of the hypersensitive response elicitor protein or polypeptide, in accordance with present invention, can be applied by low or high pressure spraying, coating, immersion, or injection. Other suitable application procedures can be envisioned by those skilled in the art provided they are able to effect contact of the fragment with cells of the plant or plant seed. Once treated with the fragment of the hypersensitive response elicitor of the present invention, the seeds can be planted in natural or artificial soil and cultivated using conventional procedures to produce plants. After plants have been propagated from seeds treated in accordance with the present invention, the plants may be treated with one or more applications of the fragment of the hypersensitive response elicitor protein or polypeptide or whole elicitors to impart disease resistance to plants, to enhance plant growth, and/or to control insects on the plants.

- 41 -

The fragment of the hypersensitive response elicitor polypeptide or protein, in accordance with the present invention, can be applied to plants or plant seeds alone or in a mixture with other materials. Alternatively, the fragment can be applied separately to plants with other materials being applied at different times.

5 A composition suitable for treating plants or plant seeds in accordance with the application embodiment of the present invention contains a fragment of a hypersensitive response elicitor polypeptide or protein which fragment does not elicit a hypersensitive response in a carrier. Suitable carriers include water, aqueous solutions, slurries, or dry powders. In this embodiment, the composition contains
10 greater than 500 nM of the fragment.

Although not required, this composition may contain additional additives including fertilizer, insecticide, fungicide, nematocide, and mixtures thereof. Suitable fertilizers include $(\text{NH}_4)_2\text{NO}_3$. An example of a suitable insecticide is Malathion. Useful fungicides include Captan.

15 Other suitable additives include buffering agents, wetting agents, coating agents, and abrading agents. These materials can be used to facilitate the process of the present invention. In addition, the hypersensitive response eliciting fragment can be applied to plant seeds with other conventional seed formulation and treatment materials, including clays and polysaccharides.

20 In the alternative embodiment of the present invention involving the use of transgenic plants and transgenic seeds, a fragment of a hypersensitive response elicitor need not be applied topically to the plants or seeds. Instead, transgenic plants transformed with a DNA molecule encoding such a fragment are produced according to procedures well known in the art.

25 The vector described above can be microinjected directly into plant cells by use of micropipettes to transfer mechanically the recombinant DNA. Crossway, Mol. Gen. Genetics, 202:179-85 (1985), which is hereby incorporated by reference. The genetic material may also be transferred into the plant cell using polyethylene glycol. Krens, et al., Nature, 296:72-74 (1982), which is hereby
30 incorporated by reference.

Another approach to transforming plant cells with a gene which imparts resistance to pathogens is particle bombardment (also known as biolistic

- 42 -

transformation) of the host cell. This can be accomplished in one of several ways. The first involves propelling inert or biologically active particles at cells. This technique is disclosed in U.S. Patent Nos. 4,945,050, 5,036,006, and 5,100,792, all to Sanford et al., which are hereby incorporated by reference. Generally, this procedure involves propelling inert or biologically active particles at the cells under conditions effective to penetrate the outer surface of the cell and to be incorporated within the interior thereof. When inert particles are utilized, the vector can be introduced into the cell by coating the particles with the vector containing the heterologous DNA. Alternatively, the target cell can be surrounded by the vector so that the vector is carried into the cell by the wake of the particle. Biologically active particles (e.g., dried bacterial cells containing the vector and heterologous DNA) can also be propelled into plant cells.

Yet another method of introduction is fusion of protoplasts with other entities, either minicells, cells, lysosomes, or other fusible lipid-surfaced bodies. Fraley, et al., Proc. Natl. Acad. Sci. USA, 79:1859-63 (1982), which is hereby incorporated by reference.

The DNA molecule may also be introduced into the plant cells by electroporation. Fromm et al., Proc. Natl. Acad. Sci. USA, 82:5824 (1985), which is hereby incorporated by reference. In this technique, plant protoplasts are electroporated in the presence of plasmids containing the expression cassette. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and regenerate.

Another method of introducing the DNA molecule into plant cells is to infect a plant cell with *Agrobacterium tumefaciens* or *A. rhizogenes* previously transformed with the gene. Under appropriate conditions known in the art, the transformed plant cells are grown to form shoots or roots, and develop further into plants. Generally, this procedure involves inoculating the plant tissue with a suspension of bacteria and incubating the tissue for 48 to 72 hours on regeneration medium without antibiotics at 25-28°C.

Agrobacterium is a representative genus of the Gram-negative family Rhizobiaceae. Its species are responsible for crown gall (*A. tumefaciens*) and hairy

- 43 -

root disease (*A. rhizogenes*). The plant cells in crown gall tumors and hairy roots are induced to produce amino acid derivatives known as opines, which are catabolized only by the bacteria. The bacterial genes responsible for expression of opines are a convenient source of control elements for chimeric expression cassettes. In addition, assaying for the presence of opines can be used to identify transformed tissue.

Heterologous genetic sequences can be introduced into appropriate plant cells, by means of the Ti plasmid of *A. tumefaciens* or the Ri plasmid of *A. rhizogenes*. The Ti or Ri plasmid is transmitted to plant cells on infection by *Agrobacterium* and is stably integrated into the plant genome. J. Schell, Science, 237:1176-83 (1987), which is hereby incorporated by reference.

After transformation, the transformed plant cells must be regenerated.

Plant regeneration from cultured protoplasts is described in Evans et al., Handbook of Plant Cell Cultures, Vol. 1: (MacMillan Publishing Co., New York, 1983); and Vasil I.R. (ed.), Cell Culture and Somatic Cell Genetics of Plants, Acad. Press, Orlando, Vol. I, 1984, and Vol. III (1986), which are hereby incorporated by reference.

It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to, all major species of sugarcane, sugar beets, cotton, fruit trees, and legumes.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts or a petri plate containing transformed explants is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted. Alternatively, embryo formation can be induced in the callus tissue. These embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and hormones, such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Efficient regeneration will depend on the medium, on the genotype, and on the history of the culture. If these three variables are controlled, then regeneration is usually reproducible and repeatable.

- 44 -

After the expression cassette is stably incorporated in transgenic plants, it can be transferred to other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Once transgenic plants of this type are produced, the plants themselves
5 can be cultivated in accordance with conventional procedure with the presence of the gene encoding the fragment of the hypersensitive response elicitor resulting in disease resistance, enhanced plant growth, and/or control of insects on the plant. Alternatively, transgenic seeds or propagules (e.g., cuttings) are recovered from the transgenic plants. The seeds can then be planted in the soil and cultivated using
10 conventional procedures to produce transgenic plants. The transgenic plants are propagated from the planted transgenic seeds under conditions effective to impart disease resistance to plants, to enhance plant growth, and/or to control insects. While not wishing to be bound by theory, such disease resistance, growth enhancement, and/or insect control may be RNA mediated or may result from expression of the
15 polypeptide or protein fragment.

When transgenic plants and plant seeds are used in accordance with the present invention, they additionally can be treated with the same materials as are used to treat the plants and seeds to which a fragment of a hypersensitive response elicitor in accordance with the present invention is applied. These other materials, including
20 a fragment of a hypersensitive response elicitor in accordance with the present invention, can be applied to the transgenic plants and plant seeds by the above-noted procedures, including high or low pressure spraying, injection, coating, and immersion. Similarly, after plants have been propagated from the transgenic plant seeds, the plants may be treated with one or more applications of the fragment of a
25 hypersensitive response elicitor in accordance with the present invention to impart disease resistance, enhance growth, and/or control insects. Such plants may also be treated with conventional plant treatment agents (e.g., insecticides, fertilizers, etc.).

EXAMPLES

30 Example 1 - Bacterial Strains and Plasmids

Escherichia coli strains used in the following examples include DH5 α and BL21(DE3) purchased from Gibco BRL (Grand Island, N.Y.) and Stratagene

- 45 -

(La Jolla, CA), respectively. The pET28(b) vector was purchased from Novagen (Madison, WI). Eco DH5 α /2139 contained the complete hrpN gene. The 2139 construct was produced by D. Bauer at Cornell University. The hrpN gene was cleaved from the 2139 plasmid by restriction enzyme digestion with HindIII, then purified from an agarose gel to serve as the DNA template for PCR synthesis of truncated hrpN clones. These clones were subsequently inserted into the (His)₆ vector pET28(b) which contained a Kan^r gene for selection of transformants.

Example 2 - DNA Manipulation

Restriction enzymes were obtained from Boehringer Mannheim (Indianapolis, IN) or Gibco BRL. T4 DNA ligase, Calf Intestinal Alkaline Phosphatase (CIAP), and PCR SupermixTM were obtained from Gibco BRL. The QIAprep Spin Miniprep Kit, the Qiagen Plasmid Mini Kit, and the QIAquick PCR Purification Kit were purchased from Qiagen (Hilden, Germany). The PCR primers were synthesized by Lofstrand Labs Limited (Gaithersburg, MD). The oligopeptides were synthesized by Bio-Synthesis, Inc. (Lewisville, TX). All DNA manipulations such as plasmid isolation, restriction enzyme digestion, DNA ligation, and PCR were performed according to standard techniques (Sambrook, et al., Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press (1989)) or protocols provided by the manufacturer.

Example 3 - Fragmentation of hrpN Gene

A series of N-terminal and C-terminal truncated *hrpN* genes and internal fragments were generated via PCR (Fig. 1). The full length hrpN gene was used as the DNA template and 3' and 5' primers were designed for each truncated clone (Fig. 2). The 3' primers contained an NdeI enzyme cutting site which contained the start codon ATG (methionine) and the 5' primers contained the stop codon TAA and a HindIII enzyme cutting site for ligation into the pET28(b) vector. PCR was carried out in 0.5 ml tubes in a GeneAmpTM 9700 (Perkin-Elmer, Foster City, CA). 45 μ l of SupermixTM (Life Technology, Gaithersburg, MD) were mixed with 20 pmoles of each pair of DNA primers, 10 ng of full length harpin DNA, and deionized

- 46 -

H₂O to a final volume of 50 µl. After heating the mixture at 95°C for 2 min, the PCR was performed for 30 cycles at 94°C for 1 min, 58°C for 1 min and 72°C for 1.5 min. The PCR products were verified on a 6% TBE gel (Novex, San Diego, CA). Amplified DNA was purified with the QIAquick PCR purification kit, digested with

5 Nde I and Hind III at 37°C for 5 hours, extracted once with phenol:chloroform:isoamylalcohol (25:25:1) and precipitated with ethanol. 5 µg of pET28(b) vector DNA were digested with 15 units of Nde I and 20 units of Hind III at 37°C for 3 hours followed with CIAP treatment to reduce the background resulting from incomplete single enzyme digestion. Digested vector DNA was purified with

10 the QIAquick PCR purification kit and directly used for ligation. Ligation was carried out at 14-16°C for 5-12 hours in a 15 µl mixture containing ca. 200 ng of digested pET28(b), 30 ng of targeted PCR fragment, and 1 unit T4 DNA ligase. 5 - 7.5 µl of ligation solution were added to 100 µl of DH5α competent cells in a 15 ml Falcon tube and incubated on ice for 30 min. After a heat shock at 42°C for 45 seconds, 0.9

15 ml SOC solution or 0.45 ml LB media were added to each tube and incubated at 37°C for 1 hour. 20, 100, and 200 µl of transformed cells were placed onto LB agar with 30 µg/ml of kanamycin and incubated at 37°C overnight. Single colonies were transferred to 3 ml LB-media and incubated overnight at 37°C. Plasmid DNA was prepared from 2 ml of culture with the QIAprep Miniprep kit (QIAGEN, Hilden,

20 Germany). The DNA from the transformed cells was analyzed by restriction enzyme digestion or partial sequencing to verify the success of the transformations. Plasmids with the desired DNA sequence were transferred into the BL21 strain using the standard chemical transformation method as indicated above. A clone containing the full length harpin protein in the pET28(b) vector was generated as a positive control,

25 and a clone with only the pET28(b) vector was generated as a negative control.

Example 4 - Expression of Hypersensitive Response Elicitor Truncated Proteins

Escherichia coli BL21(DE3) strains containing the hrpN clones were

30 grown in Luria broth medium (5g/L Difco Yeast extract, 10 g/L Difco Tryptone, 5 g/L NaCl, and 1 mM NaOH) containing 30 µg/ml of kanamycin at 37°C overnight. The bacteria were then inoculated into 100 volumes of the same medium and grown at

- 47 -

37°C to an OD₆₂₀ of 0.6-0.8. The bacteria were then inoculated into 250 volumes of the same medium and grown at 37°C to an OD₆₂₀ of ca. 0.3 or 0.6-0.8. One millimolar IPTG was then added and the cultures grown at 19°C overnight (ca. 18 hours). Not all of the clones were successfully expressed using this strategy. Several of the clones had to be grown in Terrific broth (12 g/L Bacto Tryptone, 24 g/L Bacto yeast, 0.4% glycerol, 0.17 M KH₂PO₄, and 0.72 K₂HPO₄), and/or grown at 37°C after IPTG induction, and/or harvested earlier than overnight (Table 1).

Table 1: Expression of hypersensitive response elicitor truncated proteins

10

Fragment	amino acids (SEQ. ID. No. 23)	Growth medium	Induction O.D.	Expression temp.	Harvest time
1 (+ control)	1-403	LB	ca. 0.3 or 0.6-0.8	19°C or 25°C	16-18 hr
2 (+ control)	-	LB and TB	ca. 0.3 or 0.6-0.8	19 C and 37 C	16-18 hr
3	105-403	LB	0.6-0.8	19°C	16-18 hr
4	169-403	TB	ca. 0.3	19°C	16-18 hr
5	210-403	LB or M9ZB	0.6-0.8	19°C	16-18 hr
6	257-403	LB or M9ZB	0.6-0.8	19°C	16-18 hr
7	343-403	LB	ca. 0.3	19°C	5 hr
8	1-75	TB	ca. 0.3	37°C	16-18 hr
9	1-104	TB	ca. 0.3	37°C	16-18 hr
10	1-168	TB	ca. 0.3	37°C	16-18 hr
11	1-266	LB	ca. 0.3	37°C	4 hr
12	1-342	LB	0.6-0.8	19°C	16-18 hr
13	76-209	LB	ca. 0.3	37°C	5 hr
14	76-168	TB or LB	ca. 0.3	37°C	3 hr or 16-18 hr
15	105-209	M9ZB	ca. 0.3	37°C	3 hr
16	169-209	no expression			
17	105-168	LB	ca. 0.3	37°C	3-5 hr
18	99-209	LB	ca. 0.3	37°C	3 hr
19	137-204	LB	ca. 0.3	37°C	3 hr
20	137-180	LB	ca. 0.3	37°C	16-18 hr.
21	105-180	LB	ca. 0.3	37°C	3 hr
22	150-209	no expression			
23	150-180	no expression			

Example 5 - Small Scale Purification of Hypersensitive Response Elicitor Truncated Proteins (Verification of Expression)

15

A 50 ml culture of a hrpN clone was grown as above to induce expression of the truncated protein. Upon harvesting of the culture, 1.5 ml of the cell

- 48 -

suspension were centrifuged at 14,000 rpm for 5 minutes, re-suspended in urea lysis buffer (8 M urea, 0.1 M Na₂HPO₄, and 0.01 M Tris -- pH 8.0), incubated at room temperature for 10 minutes, then centrifuged again at 14,000 rpm for 10 minutes, and the supernatant saved. A 50 µl aliquot of a 50% slurry of an equilibrated (His)₆-
5 binding nickel agarose resin was added to the supernatant and mixed at 4°C for one hour. The nickel agarose was then washed three times with urea washing buffer (8 M urea, 0.1 M Na₂HPO₄, and 0.01 M Tris -- pH 6.3), centrifuging at 5,000 rpm for five minutes between washings. The protein was eluted from the resin with 50 µl of urea elution buffer (8 M urea, 0.1 M Na₂HPO₄, 0.01 M Tris, and 0.1 M EDTA -- pH 6.3).
10 The eluate was run on a 4-20%, a 16%, or a 10-20% Tris-Glycine pre-cast gel depending upon the size of the truncated protein to verify the expression.

Example 6 - Induction of HR in Tobacco

A 1.5 ml aliquot from the 50 ml cultures grown for small scale
15 purification of the truncated proteins was centrifuged at 14,000 rpm for four minutes and re-suspended in an equal volume of 5 mM potassium phosphate buffer, pH 6.8. The cell suspension was sonicated for ca. 30 seconds then diluted 1:2 and 1:10 with phosphate buffer. Both dilutions plus the neat cell lysate were infiltrated into the fourth to ninth leaves of 10-15 leaf tobacco plants by making a hole in single leaf
20 panes and infiltrating the bacterial lysate into the intercellular leaf space using a syringe without a needle. The HR response was recorded 24-48 hr post infiltration. Tobacco (*Nicotiana tabacum* v. Xanthi) seedlings were grown in an environmental chamber at 20-25°C with a photoperiod of 12-h light /12-h dark and ca. 40% RH. Cell lysate was used for the initial HR assays (in order to screen the truncated proteins
25 for HR activity) as the small scale urea purification yielded very little protein which was denatured due to the purification process.

Example 7 - Large Scale Native Purification of Hypersensitive Response Elicitor Truncated Proteins for Comprehensive Biological Activity Assays

30 Six 500 ml cultures of a hrpN clone were grown as described earlier to induce expression of the truncated protein. Upon harvesting of the culture, the cells were centrifuged at 7,000 rpm for 5 minutes, re-suspended in imidazole lysis buffer (5

- 49 -

mM imidazole, 0.5 M NaCl, 20 mM Tris) plus Triton X-100 at 0.05% and lysozyme at 0.1 mg/ml, incubated at 30°C for 15 minutes, sonicated for two minutes, centrifuged again at 15,000 rpm for 20 minutes, and the supernatant was saved. A 4 ml aliquot of a 50% slurry of an equilibrated (His)₆-binding nickel agarose resin was added to the supernatant and mixed at 4°C for ca. four hours. The nickel agarose was then washed three times with imidazole washing buffer (20 mM imidazole, 0.5 M NaCl, and 20 mM Tris), centrifuging at 5,000 rpm for five minutes between washings, then placed in a disposable chromatography column. The column was centrifuged at 1100 rpm for one minute to remove any residual wash buffer and then the protein was eluted from the resin with 4 ml of imidazole elution buffer (1 M imidazole, 0.5 M NaCl, and 20 mM Tris) by incubating the column with the elution buffer for ten minutes at room temperature and then centrifuging the column at 1100 rpm for one minute. The eluate was run on a 4-20%, a 16%, or a 10-20% Tris-Glycine pre-cast gel depending upon the size of the truncated protein to verify the expression. The concentration of the proteins was determined by comparison of the protein bands with a standard protein in the Mark 12 molecular weight marker.

Example 8 - Large Scale Urea Purification of Hypersensitive Response Elicitor Truncated Proteins For Comprehensive Biological Activity Assay

The procedure was the same as the large scale native purification except that urea lysis buffer, washing buffer, and elution buffer were used, and the cells were not sonicated as in the native purification. After purification, the protein was renatured by dialyzing against lower and lower concentrations of urea over an eight hour period, then dialyzing overnight against 10 mM Tris/20 mM NaCl. The renaturing process caused the N-terminal proteins to precipitate. The precipitated 1-168 protein was solubilized by the addition of 100 mM Tris-HCl at pH 10.4 then heating the protein at 30°C for ca. one hour. The concentration of the protein was determined by comparison of the protein bands with a standard protein in the Mark 12 molecular weight marker. The 1-75 and 1-104 protein fragments were not successfully solubilized using this strategy so they were sonicated in 100 mM Tris-HCl at pH 10.4 to solubilize as much of the protein as possible and expose the active sites of the protein for the biological activity assays.

Example 9 – Induction of Growth Enhancement (GE)

Sixty tomato (*Lycopersicon spp.* cv. Marglobe) seeds were soaked
5 overnight in 10 and 20 µg/ml of the truncated protein diluted with 5mM potassium
phosphate buffer, pH 6.8. The next morning, the sixty seeds were sewn in three pots
and 12-15 days later and again 18-20 days later the heights of the 10 tallest tomato
plants per pot were measured and compared with the heights of the control plants
treated only with phosphate buffer. Analyses were done on the heights to determine if
10 there was a significant difference in the height of the plants treated with the truncated
proteins compared with the buffer control, and thereby determine whether the proteins
induced growth enhancement.

Example 10 – Induction of Systemic Acquired Resistance (SAR)

15 Three tobacco (*Nicotiana tabacum* cv. Xanthi) plants with 8-12 leaves
(ca. 75 day old plants) were used in the assay. One leaf of the tobacco plants was
covered up and the rest of the leaves were sprayed with ca. 50 ml of a 20 µg/ml
solution of the truncated proteins diluted with 5mM potassium phosphate buffer. Five
20 to seven days later two leaves (the unsprayed leaf and the sprayed leaf opposite and
just above the unsprayed leaf) were inoculated with 20 µl of a 1.8 µg/ml solution of
TMV along with a pinch of diatomaceous earth by rubbing the mixture along the top
surface of the leaves. The TMV entered the plants through tiny lesions made by the
diatomaceous earth. Ca. 3-4 days post TMV inoculation, the number of TMV lesions
25 was counted on both leaves compared with the number of lesions on the negative
control buffer treated leaves. Analyses were done to determine the efficacy of
reducing the number of TMV lesions by the protein fragments compared to the buffer
control. Percentage of efficacy was calculated as: Reduction in TMV lesions (%
efficacy) = 100 x (1 – mean # of lesions on treated leaves/mean # of lesions on buffer
30 control leaves).

Example 11 - Expression of Hypersensitive Response Elicitor Truncated Proteins

The small scale expression and purification of the fragment proteins was done to screen for expression and HR activity (Table 2).

5

Table 2

Expression and HR activity of hypersensitive response elicitor truncated proteins (small scale screening)

Fragment #	Amino Acids (SEQ. ID. No. 23)	Expression	HR activity
1(+control)	1-403	+	+
2(- control)	-	background protein only	-
3	105-403	+	+
4	169-403	+	-
5	210-403	+	-
6	267-403	+	-
7	343-403	+/-	-
8	1-75	+	-
9	1-104	+	+/-
10	1-168	+	+
11	1-266	+	+
12	1-342	+	+
13	76-209	+	+
14	76-168	+	-
15	105-209	+	+
16	169-209	-	-
17	105-168	+	-
18	99-209	+	+
19	137-204	+	+
20	137-180	+	+
21	105-180	+	+
22	150-209	-	-
23	150-180	-	-

10

All of the cloned fragment proteins were expressed at varying levels except for three small fragments (amino acids 169-209, 150-209, and 150-180). Fragments 210-403 and 267-403 were expressed very well, yielding a high concentration of protein from a small scale purification, resulting in a substantial protein band on SDS gel electrophoresis. Other fragments (such as a.a. 1-168 and 1-104) produced much less protein, resulting in faint protein bands upon electrophoresis. It was difficult to determine whether fragment 343-403, the smallest C-terminal protein, was expressed, as there were several background proteins apparent on the gel, in addition to the suspected 343-403 protein. The positive and negative control proteins, consisting of

15

- 52 -

the full length hypersensitive response elicitor protein and only background proteins, respectively, were tested for expression and HR activity as well.

The large scale expression and purification of the fragment proteins was done to determine the level of expression and titer of the HR activity (Table 3).

5

Table 3

Expression level and HR titer of hypersensitive response elicitor truncated proteins (large scale purification)

10

Fragment #	Amino acids (SEQ. ID. No. 23)	Expression	HR titer
1(+ control)	1-403	3.7 mg/ml	5-7 µg/ml
2 (- control)	-	-	1:2 dilution
4	169-403	2 mg/ml	-
5	210-403	5 mg/ml	-
6	267-403	4 mg/ml	-
7	343-402	200µg/ml	-
8	1-75	50µg/ml	-
9	1-104	50µg/ml	3 µg/ml (1:16 dilution)
10	1-168	1 mg/ml	1 µg/ml
13	76-209	2.5 mg/ml	5 µg/ml
14	76-168	2 mg/ml	-
15	105-209	5 mg/ml	5-10µg/ml
17	105-168	250µg/ml	-
19	137-204	3.6 mg/ml	3.5 µg/ml
20	137-180	250 µg/ml	16 µg/ml

15

20

The truncated proteins deemed to be the most important in characterizing the hypersensitive response elicitor were chosen for large scale expression. The positive control (full length hypersensitive response elicitor) was expressed at a relatively high level at 3.7 mg/ml. All of the C-terminal proteins were expressed at relatively high levels from 2-5 mg/ml, except for fragment 343-403 as discussed earlier. The N-terminal fragments were expressed very well also; however, during the purification process, the protein precipitated and very little was resolubilized. The concentrations in Table 3 reflect only the solubilized protein. The internal fragments were expressed in the range of 2-3.6 mg/ml. It was extremely difficult to determine the concentration of fragment 105-168 (it was suspected that the concentration was much higher than indicated), as the protein bands on the SDS gel were large, but poorly stained. The

- 53 -

negative control contained several background proteins as expected, but no obviously induced dominant protein.

Example 12 - Induction of HR in Tobacco

5

The full length positive control protein elicited HR down to only 5-7µg/ml. The negative control (pET 28) imidazole purified "protein" - which contained only background proteins - elicited an HR response down to the 1:2 dilution, which lowered the sensitivity of the assay as the 1:1 and 1:2 dilutions could not be used. This false HR was likely due to an affinity of the imidazole used in the purification process to bind to one or several of the background proteins, thereby not completely dialyzing out. Imidazole at a concentration of ca. 60 mM did elicit a false HR response.

10

One definitive domain encompassing a small internal region of the protein from a.a. 137-180 (SEQ. ID. No. 23), a mere 44 a.a, is identified as the smallest HR domain. The other potential HR domain is thought to be located in the N-terminus of the protein from a.a. 1-104 (possibly a.a. 1-75) (SEQ. ID. No. 23). It was difficult to confirm or narrow down the N-terminus HR domain due to the difficulties encountered in purifying these fragment proteins. The N-terminus fragment proteins had to be purified with urea as no protein was recovered when the native purification process was used. Consequently, these proteins precipitated during the renaturing process and were difficult or nearly impossible to get back into solution, thereby making it hard to run the proteins through the HR assay, as only soluble protein is able to elicit HR. Difficulty narrowing the N-terminus HR domain was only compounded by the fact that the negative control elicited false HR at the low dilution levels thereby reducing the sensitivity of the assay.

20

25

Surprisingly, when the internal HR domain was cleaved between a.a. 168 and 169 (fragments 76-168 and 105-168) (SEQ. ID. No. 23) the fragment lost its HR activity. This suggests that the HR activity of fragment 1-168 (SEQ. ID. No. 23) should not be attributed to the internal HR domain, but rather to some other domain, leading to the assumption that there was likely a second HR domain to be found in the N-terminal region of the protein. However, as discussed earlier it was difficult to confirm this assumption.

30

- 54 -

The hypersensitive response elicitor C-terminus (a.a. 210-403 (SEQ. ID. No. 23)) did not contain an HR domain. It did not elicit HR at a detectable level using the current HR assay. Even the large C-terminal fragment from a.a. 169-403 (SEQ. ID. No. 23) did not elicit HR even though it contained part of the internal HR domain. As stated above, cleaving the protein between amino acids 168 and 169 (SEQ. ID. No. 23) causes a loss of HR activity.

Because some of the small cloned proteins with 61 a.a. or less were not expressed, several oligopeptides were synthesized with 30 a.a. to narrow down the functional region of the internal HR domain. The oligopeptides were synthesized within the range of a.a. 121-179 (SEQ. ID. No. 23). However, these oligos did not elicit HR. It was not expected that there would be an HR from oligos 137-166, 121-150, and 137-156 (SEQ. ID. No. 23) as these fragments did not contain the imperative amino acids 168 and 169 (SEQ. ID. No. 23). It was expected that the oligo 150-179 (SEQ. ID. No. 23) would elicit an HR. It is possible that 30 a.a. is too small for the protein to elicit any activity due to a lack of folding and, therefore, a lack of binding or that during the synthesis of the peptides important amino acids were missed (either in the process, or simply by the choice of which 30 amino acids to synthesize) and, therefore, the fragments would not be able to elicit HR.

20 **Example 13 – Induction of Plant Growth Enhancement (PGE)**

The C-terminal fragments enhanced the growth of tomato by 9% to 21%. The N-terminal fragments enhanced the growth of tomato by 4% to 13%. The internal fragments enhanced growth by 9% to 20%. The 76-209 fragment enhanced growth by 18% at a concentration of 60 µg/ml, but not at the typical 20 µg/ml. This was attributed to the inaccuracy of the quantification process (Table 4).

- 55 -

Table 4

Fragment #	Amino acids	PGE ht>buffer @ 10 µg/ml	PGE ht>buffer @ 20 µg/ml
1 (+ control)	1-403	12%	11%
2 (- control)	-	-3%	-2%
4	169-403	9%	12%
5	210-403	13%	14%
			16% @ 40µg/ml
6	267-403	21%	21%
			23% @ 40µg/ml
7	343-403	7%	7%
9	1-104	4%	8%
10	1-168	13%	5%
13	76-209	7%	4%
			18% @ 60µg/ml
14	76-168	18%	20%
15	105-209	14%	19%
17	105-168	19%	16%
19	137-204	11%	13%
20	137-180	--	9%

*A height greater than 10% above the buffer control was necessary to pass the PGE assay.

The oligopeptides enhanced growth from 7.4% to 17.3% (Table 5).

Table 5

Fragment	Amino acids	Expression	HR titer	TMV efficacy	PGE ht>buffer
oligo	150-179	NA	-	72.9%	10.1%
oligo	137-166	NA	-	61.2%	12.0%
oligo	121-150	NA	-	60.0%	17.3%
oligo	137-156	NA	-	-87.7%	7.4%

The data suggests that there is more than one PGE domain, although the C-terminal and internal domains appear to be dominant over the N-terminal domain, as the N-terminal fragments enhanced growth the least amount.

Example 14 – Induction of Systemic Acquired Resistance (SAR)

All of the hypersensitive response elicitor fragments tested to date appear to have 60% efficacy or greater, except for the oligopeptide 137-156 (Tables 5 and 6).

Table 6

Fragment #	Amino acids	Efficacy of TMV control
1 (+ control)	1-403	84% & 72%
2 (- control)	-	40% & 31%
4	169-403	64% & 79%
5	210-403	77% and 78%
6	267-403	70% and 72%
9	1-104	82%
10	1-168	69%
13	76-209	44% and 84%
14	76-168	83% & 87%
15	105-209	57% and 67%
17	105-168	89%
19	137-204	89% & 77%
20	137-180	64% & 58%

These data suggest that there are multiple SAR domains within the protein.

Example 15 – Relationship Between HR, PGE, and SAR

It is clear that the hypersensitive response activity is separable from the plant growth enhancement activity. The C-terminal fragments clearly enhance the growth of tomato by ca. 20% at a concentration of only 20 µg/ml, but these same fragments were not able to elicit HR in tobacco, even at higher concentrations than 200 µg/ml. The SAR activity also appears to be separable from the HR activity. This finding is highly significant for future work on transgenic applications of the hypersensitive response elicitor technology. The fragments that induce PGE and/or SAR but do not elicit HR will be imperative for this technology, as constitutive expression of even low levels of an HR elicitor might kill a plant.

Example 16 - Non-HR Eliciting Fragments Derived from the Hypersensitive Response Elicitor from *Pseudomonas syringae* pv. *syringae* Induce Resistance in Tobacco to TMV and Promote the Growth of Tomato

To test whether non-HR eliciting fragments derived from HrpZ, the hypersensitive response elicitor from *Pseudomonas syringae* pv. *syringae*, is able to induce disease resistance, several fragment constructs were made and the expressed

- 57 -

fragment proteins were tested for HR elicitation and disease resistance induction in tobacco and growth promotion in tomato.

The following segments of *hrpZ*, the gene encoding the hypersensitive response elicitor from *Pseudomonas syringae* pv. *syringae*, were amplified by PCR using Pfu Turbo (Stratagene): Regions coding for amino acids 152-190, aa 152-294, aa 190-294, aa 301-341, and full length HrpZ (aa 1-341). The DNA fragments were cloned into pCAL-n (Stratagene) to create C-terminal fusion proteins to the calmodulin-binding peptide. pCAL-n was chosen, because the fusion protein could be easily and gently purified on calmodulin resin. The DNA was transformed into *E. coli* DH5 α , and the correct clones were identified. The clones were then transferred to *E. coli* BLR DE3 for protein expression. The bacteria were grown in Terrific Broth to an OD₆₂₀ of 0.8-1.0. Protein expression was then induced with IPTG and the bacteria were incubated for an additional 3 h. All of the HrpZ fragments were able to be expressed this way.

Amino acid fragments 152-294 and 190-294 were chosen for further analysis and characterization. It was expected that the fragment 152-294 contained a domain that elicited the HR, while fragment 190-294 contained no domain that elicited the HR. The cultures were spun down, and the bacteria resuspended in 40 ml of 10 mM Tris pH 8.0. Twenty μ l of antifoam and 40 μ l of 200 mM PMSF were added, and the bacteria was sonicated to break open the cells. The bacterial debris was removed by centrifugation, and the supernatant was placed in a boiling water bath for 10 min. The precipitate was removed by centrifugation and the supernatant, a crude protein preparation, was retained for tests.

Fifteen μ l of each supernatant was run on a gel and stained to determine if the protein was present. It was estimated that about five times as much of the 152-294 fragment was present as the 190-294 fragment. Several dilutions of each preparation were infiltrated into tobacco leaves on two plants for HR tests (Table 7). As shown in Table 7, the 152-294 fragment elicited an HR, but the 190-294 fragment did not.

- 58 -

Table 7

HR test results of HrpZ fragments

<u>HrpZ Fragment</u>	<u>Dilution of Fragment Preparation^a</u>			
	<u>1:2</u>	<u>1:5</u>	<u>1:25</u>	<u>1:125</u>
152-294	+,+ ^b	+,+	+,+	-, -
190-294	-,-	-,-	-,-	-,-

^a The preparations were diluted with MilliQ water.^b The results are indicated for each of two plants. +, HR; -, no HR.

10

The fragment preparations were then tested for inducing resistance to TMV and for growth enhancement. Due to the difference in concentration of the HrpZ fragments, the 152-294 preparation was diluted 40-fold and the 190-294 preparation was diluted 8-fold. The results showed that the 190-294 aa fragment reduced the number of TMV lesions by 85% in comparison to buffer controls (Table 8). In contrast, the 152-294 aa fragment reduced the number of TMV lesions by only 55%. As also shown in Table 8, plants treated with the 152-294 aa fragment grew 4.64% more than buffer treated plants, while plants treated with the 190-294 aa fragment grew 2.62% more than the buffer treated plants.

20

Table 8

HR test, TMV, and PGE test results

<u>HrpZ Fragment</u>	<u>HR elicitation^a</u>	<u>TMV (% efficacy)^b</u>	<u>PGE(% > buffer ht)^c</u>
152-294	+	54.64	4.64
190-294	-	85.25	2.62

^a +, elicits HR in tobacco leaves; -, no HR in tobacco leaves.^b % reduction in TMV lesions in unsprayed leaf of tobacco.^c % greater height than buffer sprayed plants.

30

The results of these tests show that amino acids 152-190 appear to be involved in HR elicitation, because their removal eliminated the ability to elicit the HR. Both fragment preparations achieved disease control and growth enhancement. Thus, the ability to elicit the HR is not the determining factor for reduction in TMV infection and growth enhancement.

35

- 59 -

Example 17 - Use of 13 Amino Acid Peptide Derived from *Phytophthora megasperma* Stimulates Tomato Seedling Growth

Parsley leaves develop a typical resistance reaction against the soybean pathogen *Phytophthora megasperma* comprising hypersensitive cell death, defense related gene activation, and phytoalexin formulation. Several years ago, a 42 kDa glycoprotein elicitor was purified from the fungal culture filtrate of *Phytophthora megasperma* (Parker et al., "An Extracellular Glycoprotein from *Phytophthora megasperma* f.sp. glycinea Elicits Phytoalexin Synthesis in Cultured Parsley Cells and Protoplasts," Mol. Plant Microbe Interact. 4:19-27 (1991), which is hereby incorporated by reference). Then, an oligopeptide of 13 amino acid was identified within the 42 kDa glycoprotein. The 13 amino acids peptide appeared to have similar biological activity as that of the full length glycoprotein (42 kDa). It is sufficient to elicit a complex defense response in parsley cells including H⁺/Ca²⁺ influxes, K⁺/Cl⁻ effluxes, active oxygen production, SAR gene induction, and phytoalexin compound accumulation (Nurnberger et al., "High Affinity Binding of a Fungal Oligopeptide Elicitor to Parsley Plasma Membranes Triggers Multiple Defense Response," Cell 78:449-460 (1994), which is hereby incorporated by reference).

To test if the 13 amino acid peptide derived from the 42 kDa protein also enhanced plant growth, 20 mg of the oligopeptide was synthesized from Biosynthesis Corp. The synthesized sequence of the peptide is NH₂-Val-Trp-Asn-Gln-Pro-Val-Arg-Gly-Phe-Lys-Val-Tyr-Glu-COOH (SEQ. ID. No. 39). The synthesized peptide was resuspended in 10 ml of 5 mM potassium phosphate buffer and, then, diluted to 1 and 100 ng/ml with the same buffer. About 100 tomato seeds (variety, Marglobe) were submerged in 20 ml of peptide solution overnight. The soaked seeds were planted in an 8 inch pot with artificial soil. Seeds soaked in the buffer without the peptide were used as a control. After seedlings emerged and the first two true leaves fully expanded, the height of the tomato seedlings was recorded. The peptide was not able to elicit the HR in tobacco and other tested plants. However, it had a profound effect on plant growth promotion. Table 9 shows that tomato seedlings treated with the peptide increased 12.6 % in height, indicating that the fungal peptide derived from the 42 kDa glycoprotein can promote tomato seedling growth. Extended studies showed that the peptide also had similar growth

- 60 -

effect in other crops including tobacco. Similar growth promotion effects were achieved by plants sprayed with the peptide solution.

Table 9

5	Treatment	Height of seedlings (cm)					Average (cm) % Change	
	Buffer	6.0	6.0	6.0	5.5	5.5	5.55	-
10		5.5	5.5	5.0	5.0	5.5		
	Peptide Solution (100ng/ml)	6.5	6.0	6.5	6.5	6.5	6.25	12.6
		6.0	6.0	6.0	6.0	6.5		

15 Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is solely for that purpose, and variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

- 61 -

WHAT IS CLAIMED:

1. An isolated fragment of a hypersensitive response elicitor protein or polypeptide, wherein said fragment does not elicit a hypersensitive response but has other activity in plants.
5
2. An isolated fragment according to claim 1, wherein the hypersensitive response elicitor protein or polypeptide is derived from an *Erwinia Pseudomonas*, *Xanthomonas*, or *Phytophthora*.
10
3. An isolated fragment according to claim 2, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Erwinia amylovora*.
15
4. An isolated fragment according to claim 3, wherein the fragment is selected from the group consisting of a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 23, an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 23, and an internal fragment of the amino acid sequence of SEQ. ID. No. 23.
20
5. An isolated fragment according to claim 4, wherein the fragment is a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 23 spanning the following amino acids of SEQ. ID. No. 23: 169 and 403, 210 and 403, 267 and 403, or 343 and 403.
25
6. An isolated fragment according to claim 4, wherein the fragment is an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 23.
7. An isolated fragment according to claim 4, wherein the
30 fragment is an internal fragment of the amino acid sequence of SEQ. ID. No. 23 spanning the following amino acids of SEQ. ID. No. 23: 105 and 179, 137 and 166, 121 and 150, or 137 and 156.

- 62 -

8. An isolated fragment according to claim 2, wherein the hypersensitive response elicitor is derived from *Pseudomonas syringae*.

9. An isolated fragment according to claim 8, wherein the
5 fragment contains amino acids 190 to 294 of SEQ. ID. No. 31.

10. An isolated DNA molecule encoding a fragment according to claim 1.

10 11. An isolated DNA molecule according to claim 10, wherein the hypersensitive response elicitor protein or polypeptide is derived from an *Erwinia Pseudomonas*, *Xanthomonas*, or *Phytophthora*.

12. An isolated DNA molecule according to claim 11, wherein the
15 hypersensitive response elicitor protein or polypeptide is derived from *Erwinia amylovora*.

13. An isolated DNA molecule according to claim 12, wherein the fragment is selected from the group consisting of a C-terminal fragment of the amino
20 acid sequence of SEQ. ID. No. 23, an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 23, and an internal fragment of the amino acid sequence of SEQ. ID. No. 23.

14. An isolated DNA molecule according to claim 12, wherein the
25 fragment is a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 23 spanning the following amino acids of SEQ. ID. No. 23: 169 and 403, 210 and 403, 267 and 403, or 343 and 403.

15. An isolated DNA molecule according to claim 12, wherein the
30 fragment is an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 23.

- 63 -

16. An isolated DNA molecule according to claim 12, wherein the fragment is an internal fragment of the amino acid sequence of SEQ. ID. No. 23 spanning the following amino acids of SEQ. ID. No. 23: 105 and 179, 137 and 166, 121 and 150, or 137 and 156.

5

17. An isolated DNA molecule according to claim 11, wherein the hypersensitive response elicitor is derived from *Pseudomonas syringae*.

18. An isolated DNA molecule according to claim 18, wherein the
10 fragment contains amino acids 190 to 294 of SEQ. ID. No. 31.

19. An expression system transformed with a DNA molecule according to claim 10.

20. An expression system according to claim 19, wherein said
15 DNA molecule is in proper sense orientation and correct reading frame.

21. A host cell transformed with a DNA molecule according to
claim 10.

20

22. A host cell according to claim 21, wherein the host cell is selected from the group consisting of a plant cell and a bacterial cell.

23. A host cell according to claim 21, wherein the DNA molecule
25 is transformed with an expression system.

24. A transgenic plant transformed with the DNA molecule of
claim 10.

25. A transgenic plant according to claim 24, wherein the plant is
30 selected from the group consisting of alfalfa, rice, wheat, barley, rye, cotton, sunflower, peanut, corn, potato, sweet potato, bean, pea, chicory, lettuce, endive,

- 64 -

cabbage, brussel sprout, beet, parsnip, turnip, cauliflower, broccoli, radish, spinach, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane.

5

26. A transgenic plant according to claim 24, wherein the plant is selected from the group consisting of *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

10

27. A transgenic plant seed transformed with the DNA molecule of claim 10.

15

28. A transgenic plant seed according to claim 27, wherein the plant seed is selected from the group consisting of alfalfa, rice, wheat, barley, rye, cotton, sunflower, peanut, corn, potato, sweet potato, bean, pea, chicory, lettuce, endive, cabbage, brussel sprout, beet, parsnip, turnip, cauliflower, broccoli, radish, spinach, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane.

20

29. A transgenic plant seed according to claim 27, wherein the plant seed is selected from the group consisting of *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

25

30. A method of imparting disease resistance to plants comprising: applying a fragment of a hypersensitive response elicitor protein or polypeptide, which fragment does not elicit a hypersensitive response, in a non-infectious form to a plant or plant seed under conditions effective to impart disease resistance.

30

31. A method according to claim 30, wherein plants are treated during said applying.

- 65 -

32. A method according to claim 30 wherein plant seeds are treated during said applying, said method further comprising:

5 planting the seeds treated with the fragment of the
hypersensitive response elicitor in natural or artificial soil and
propagating plants from the seeds planted in the soil.

33. A method of enhancing plant growth comprising:

10 applying a fragment of a hypersensitive response elicitor
protein or polypeptide, which fragment does not elicit a hypersensitive response, in a
non-infectious form to a plant or plant seed under conditions effective to enhance
plant growth.

34. A method according to claim 33, wherein plants are treated
15 during said applying.

35. A method according to claim 33, wherein plant seeds are treated during said applying, said method further comprising:

20 planting the seeds treated with the fragment of the
hypersensitive response elicitor in natural or artificial soil and
propagating plants from the seeds planted in the soil.

36. A method of insect control for plants comprising:

25 applying a fragment of a hypersensitive response elicitor protein or
polypeptide, which fragment does not elicit a hypersensitive response, in a non-
infectious form to a plant or plant seed under conditions effective to control insects.

37. A method according to claim 36, wherein plants are treated during said applying.

30

38. A method according to claim 36, wherein plant seeds are treated during said applying, said method further comprising:

- 66 -

planting the seeds treated with the fragment of the
hypersensitive response elicitor in natural or artificial soil and
propagating plants from the seeds planted in the soil.

5 39. A method of imparting disease resistance to plants comprising:
 providing a transgenic plant or plant seed transformed with a
 DNA molecule which encodes a fragment of a hypersensitive response elicitor protein
 or polypeptide, which fragment does not elicit a hypersensitive response, and
 growing the transgenic plant or transgenic plants produced
10 from the transgenic plant seeds under conditions effective to impart disease resistance.

 40. A method according to claim 39, wherein a transgenic plant is
 provided.

15 41. A method according to claim 39, wherein a transgenic plant
 seed is provided.

 42. A method of enhancing plant growth comprising:
 providing a transgenic plant or a plant seed transformed with a
20 DNA molecule which encodes a fragment of a hypersensitive response elicitor protein
 or polypeptide, which fragment does not elicit a hypersensitive response, and
 growing the transgenic plant or transgenic plants produced
 from the transgenic plant seeds under conditions effective to enhance plant growth.

25 43. A method according to claim 42, wherein a transgenic plant is
 provided.

 44. A method according to claim 42, wherein a transgenic plant
 seed is provided.


30 45. A method of insect control for plants comprising:

- 67 -

providing a transgenic plant or plant seed transformed with a
DNA molecule which encodes a fragment of a hypersensitive response elicitor protein
or polypeptide, which fragment does not elicit a hypersensitive response, and
growing the transgenic plant or transgenic plants produced
5 from the transgenic plant seeds under conditions effective to control insects.

46. A method according to claim 45, wherein a transgenic plant is
provided.

10 47. A method according to claim 45, wherein a transgenic plant
seed is provided.



Harpin fragments derived from HrpN of *Erwinia amylovora*

FIGURE 1

1; 5'-GGGAATTCATATGAGTCTGAATACAAGTGGG-3'
76; 5'-GGGAATTCATATGGGCGGTGGCTTAGGCGGT-3'
99; 5'-GGCATATGTCGAACGCGCTGAACGATATG-3'
105; 5'-GGGAATTCATATGTTAGGCGGTTCGCTGAAC-3'
110; 5'-GGCATATGCTGAACACGCTGGGCTCGAAA-3'
137; 5'-GGCATATGTCAACGTCCAAAACGACGAT-3'
150; 5'-GGCATATGTCCACCTCAGACTCCAGCG-3'
169; 5'-GGGAATTCATATGCAAAGCCTGTTTGGTGATGGG-3'
1210; 5'-GGGAATTCATATGGGTAATGGTCTGAGCAAG-3'
1267; 5'-GGGAATTCATATGAAAGCGGGCATTGAGGCG-3'
1343; 5'-GGGAATTCATATGACACCAGCCAGTATGGAGCAG-3'
175; 5'-GCAAGCTTAAACAGCCCACCACCGCCCATCAT-3'
1104; 5'-GCAAGCTTAAATCGTTCAGCGCGTTCGACAG-3'
1168; 5'-GCAAGCTTAAATATCTCGCTGAACATCTTCAGCAG-3'
1180; 5'-GCAAGCTTAAAGGTGCCATCTTGCCCATCAC-3'
1204; 5'-GCAAGCTTAAATCAGTGACTCCTTTTTTATAGGC-3'
1209; 5'-GCAAGCTTAAACAGGCCCGACAGCGCATCAGT-3'
1266; 5'-GCAAGCTTAAACCGATAACCGGTACCCACGGC-3'
1342; 5'-GCAAGCTTAAATCCGTCGTCATCTGGCTTGCTCAG-3'
1403; 5'-GCAAGCTTAAAGCCGCGCCAGCTTG-3'

Oligonucleotide primers used for the construction of the subclones of *Erwinia amylovora* hrpN.

FIGURE 2

SEQUENCE LISTING

<110> Eden Bioscience Corporation

<120> HYPERSENSITIVE RESPONSE ELICITOR FRAGMENTS WHICH ARE
ACTIVE BUT DO NOT ELICIT A HYPERSENSITIVE RESPONSE

<130> 21829/32

<140>

<141>

<150> 60/103,050

<151> 1998-10-05

<160> 39

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> DNA

<213> Erwinia amylovora

<400> 1

gggaattcat atgagtctga atacaagtgg g

31

<210> 2

<211> 31

<212> DNA

<213> Erwinia amylovora

<400> 2

gggaattcat atgggcggtg gcttaggcgg t

31

<210> 3

<211> 29

<212> DNA

<213> Erwinia amylovora

<400> 3

ggcatatgtc gaacgcgctg aacgatatg

29

<210> 4

<211> 31

<212> DNA

<213> Erwinia amylovora

<400> 4
gggaattcat atggttaggcg gttcgctgaa c 31

<210> 5
<211> 29
<212> DNA
<213> Erwinia amylovora

<400> 5
ggcatatgct gaacacgctg ggctcgaaa 29

<210> 6
<211> 29
<212> DNA
<213> Erwinia amylovora

<400> 6
ggcatatgct aacgtcccaa aacgacgat 29

<210> 7
<211> 27
<212> DNA
<213> Erwinia amylovora

<400> 7
ggcatatgct cacctcagac tccagcg 27

<210> 8
<211> 34
<212> DNA
<213> Erwinia amylovora

<400> 8
gggaattcat atgcaaagcc tgtttggtga tggg 34

<210> 9
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 9
gggaattcat atgggtaatg gtctgagcaa g 31

<210> 10
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 10
gggaattcat atgaaagcgg gcattcaggg g 31

<210> 11
<211> 34
<212> DNA
<213> Erwinia amylovora

<400> 11
gggaattcat atgacaccag ccagtatgga gcag 34

<210> 12
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 12
gcaagcttaa cagcccacca ccgcccattca t 31

<210> 13
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 13
gcaagcttaa atcggttcagc gcgttcgaca g 31

<210> 14
<211> 34
<212> DNA
<213> Erwinia amylovora

<400> 14
gcaagcttaa tatctcgtg aacatcttca gcag 34

<210> 15
<211> 30
<212> DNA
<213> Erwinia amylovora

<400> 15
gcaagcttaa ggtgccatct tgcccatcac 30

<210> 16
<211> 34
<212> DNA
<213> Erwinia amylovora

<400> 16
gcaagcttaa atcagtgact ccttttttat aggc 34

<210> 17
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 17
gcaagcttaa caggcccgac agcgcatcag t 31

<210> 18
<211> 31
<212> DNA
<213> Erwinia amylovora

<400> 18
gcaagcttaa accgataccg gtacccacgg c 31

<210> 19
<211> 34
<212> DNA
<213> Erwinia amylovora

<400> 19
gcaagcttaa tccgtcgtca tctggcttgc tcag 34

<210> 20
<211> 25
<212> DNA
<213> Erwinia amylovora

<400> 20
gcaagcttaa gccgcgccca gcttg 25

<210> 21
<211> 338
<212> PRT
<213> Erwinia chrysanthemi

<400> 21
Met Gln Ile Thr Ile Lys Ala His Ile Gly Gly Asp Leu Gly Val Ser
1 5 10 15

Gly Leu Gly Ala Gln Gly Leu Lys Gly Leu Asn Ser Ala Ala Ser Ser
20 25 30

Leu Gly Ser Ser Val Asp Lys Leu Ser Ser Thr Ile Asp Lys Leu Thr

35	40	45
Ser Ala Leu Thr Ser Met Met Phe Gly Gly Ala Leu Ala Gln Gly Leu		
50	55	60
Gly Ala Ser Ser Lys Gly Leu Gly Met Ser Asn Gln Leu Gly Gln Ser		
65	70	75
Phe Gly Asn Gly Ala Gln Gly Ala Ser Asn Leu Leu Ser Val Pro Lys		
85	90	95
Ser Gly Gly Asp Ala Leu Ser Lys Met Phe Asp Lys Ala Leu Asp Asp		
100	105	110
Leu Leu Gly His Asp Thr Val Thr Lys Leu Thr Asn Gln Ser Asn Gln		
115	120	125
Leu Ala Asn Ser Met Leu Asn Ala Ser Gln Met Thr Gln Gly Asn Met		
130	135	140
Asn Ala Phe Gly Ser Gly Val Asn Asn Ala Leu Ser Ser Ile Leu Gly		
145	150	155
Asn Gly Leu Gly Gln Ser Met Ser Gly Phe Ser Gln Pro Ser Leu Gly		
165	170	175
Ala Gly Gly Leu Gln Gly Leu Ser Gly Ala Gly Ala Phe Asn Gln Leu		
180	185	190
Gly Asn Ala Ile Gly Met Gly Val Gly Gln Asn Ala Ala Leu Ser Ala		
195	200	205
Leu Ser Asn Val Ser Thr His Val Asp Gly Asn Asn Arg His Phe Val		
210	215	220
Asp Lys Glu Asp Arg Gly Met Ala Lys Glu Ile Gly Gln Phe Met Asp		
225	230	235
Gln Tyr Pro Glu Ile Phe Gly Lys Pro Glu Tyr Gln Lys Asp Gly Trp		
245	250	255
Ser Ser Pro Lys Thr Asp Asp Lys Ser Trp Ala Lys Ala Leu Ser Lys		
260	265	270
Pro Asp Asp Asp Gly Met Thr Gly Ala Ser Met Asp Lys Phe Arg Gln		
275	280	285
Ala Met Gly Met Ile Lys Ser Ala Val Ala Gly Asp Thr Gly Asn Thr		

290 295 300
 Asn Leu Asn Leu Arg Gly Ala Gly Gly Ala Ser Leu Gly Ile Asp Ala
 305 310 315 320
 Ala Val Val Gly Asp Lys Ile Ala Asn Met Ser Leu Gly Lys Leu Ala
 325 330 335
 Asn Ala

<210> 22

<211> 2141

<212> DNA

<213> *Erwinia chrysanthemi*

<400> 22

cgattttacc cgggtgaacg tgctatgacc gacagcatca cgggtattcga caccggttacg 60
 gcgttttatgg ccgcgatgaa ccggcatcag gcggcgcgct ggctcgccgca atccggcgctc 120
 gatctggtat ttcagtttgg ggacaccggg cgtgaactca tgatgcagat tcagccggggg 180
 cagcaatata ccggcatgtt gcgcacgctg ctgcctcgtc gttatcagca ggccggcagag 240
 tgccgatggct gccatctgtg cctgaacggc agcgatgtat tgatcctctg gtggccgctg 300
 ccgtcggatc ccggcagtta tccgcagggtg atcgaacgtt tgtttgaact ggccgggaatg 360
 acgttgccgt cgctatccat agcaccgacg gcgcgtccgc agacaggga cggacgcgcc 420
 cgatcattaa gataaaggcg gcttttttta ttgcaaacg gtaacggtga ggaaccgttt 480
 caccgtcggc gtcactcagt aacaagtata catcatgatg cctacatcgg gatcggcgctg 540
 ggcacccgtt gcagataact ttgcgaacac ctgacatgaa tgaggaaacg aaattatgca 600
 aattacgatc aaagcgcaca tcggcgggtga tttgggcgtc tccgggtctgg ggctgggtgc 660
 tcagggactg aaaggactga attccgcggc ttcacgctg ggttccagcg tggataaact 720
 gagcagcacc atcgataagt tgacctccgc gctgacttcc atgatgtttg gcggcgcgct 780
 ggccgagggg ctgggcgcca gctcgaagg gctggggatg agcaatcaac tgggccagtc 840
 tttcggcaat ggccgcgagg gtgcgagcaa cctgctatcc gtaccgaaat ccggcgccga 900
 tgcggtgtca aaaatgtttg ataaagcgct ggacgatctg ctgggtcatg acaccgtgac 960
 caagctgact aaccagagca accaactggc taattcaatg ctgaacgcca gccagatgac 1020
 ccagggtaat atgaatgctg tcggcagcgg tgtgaacaac gcaactgctg ccattctcgg 1080
 caacggtctc ggccagtcga tgagtggcct ctctcagcct tctctggggg caggcggtct 1140
 gcagggcctg agcggcgcgg gtgcattcaa ccagttgggt aatgccatcg gcatgggcgt 1200
 ggggcagaat gctgcgctga gtgcgttgag taacgtcagc acccacgtag acggtaacaa 1260
 ccgccacttt gtagataaag aagatcgcgg catggcgaaa gagatcggcc agtttatgga 1320
 tcagtatccg gaaatattcg gtaaaccgga ataccagaaa gatggctgga gttcgccgaa 1380
 gacggacgac aaatcctggg ctaaagcgct gagtaaaccg gatgatgacg gtatgaccgg 1440
 cgccagcatg gacaaattcc gtcaggcgat gggatatgat aaaagcgcgg tggcgggtga 1500
 taccggcaat accaactga acctgcgtgg cgcgggcggt gcacgctgg gtatcgatgc 1560
 ggctgtcgtc ggcgataaaa tagccaacat gtcgctgggt aagctggcca acgcctgata 1620
 atctgtgctg gcctgataaa gcggaaacga aaaaagagac ggggaagcct gtctcttttc 1680
 ttattatgcg gtttatgagg ttacctggac cggttaatca tcgtcatcga tctggtacaa 1740
 acgcacattt tcccgttcat tcgcgtcgtt acgcgccaca atcgcgatgg catcttctc 1800

gtcgcctcaga ttgcgcgggct gatgggggaac gccgggtgga atatagagaa actcgccggc 1860
 cagatggaga cacgtctgcg ataaatctgt gccgtaacgt gtttctatcc gcccttttag 1920
 cagatagatt gcggtttcgt aatcaacatg gtaatgcggt tccgcctgtg cgccggccgg 1980
 gatcaccaca atattcatag aaagctgtct tgcacctacc gtatcgcggg agataccgac 2040
 aaaatagggc agtttttgcg tggatccgt ggggtgttcc ggcctgacaa tcttgagttg 2100
 gttcgtcatc atctttctcc atctgggcga cctgatcggt t 2141

<210> 23

<211> 403

<212> PRT

<213> *Erwinia amylovora*

<400> 23

Met Ser Leu Asn Thr Ser Gly Leu Gly Ala Ser Thr Met Gln Ile Ser
 1 5 10 15

Ile Gly Gly Ala Gly Gly Asn Asn Gly Leu Leu Gly Thr Ser Arg Gln
 20 25 30

Asn Ala Gly Leu Gly Gly Asn Ser Ala Leu Gly Leu Gly Gly Gly Asn
 35 40 45

Gln Asn Asp Thr Val Asn Gln Leu Ala Gly Leu Leu Thr Gly Met Met
 50 55 60

Met Met Met Ser Met Met Gly Gly Gly Gly Leu Met Gly Gly Gly Leu
 65 70 75 80

Gly Gly Gly Leu Gly Asn Gly Leu Gly Gly Ser Gly Gly Leu Gly Glu
 85 90 95

Gly Leu Ser Asn Ala Leu Asn Asp Met Leu Gly Gly Ser Leu Asn Thr
 100 105 110

Leu Gly Ser Lys Gly Gly Asn Asn Thr Thr Ser Thr Thr Asn Ser Pro
 115 120 125

Leu Asp Gln Ala Leu Gly Ile Asn Ser Thr Ser Gln Asn Asp Asp Ser
 130 135 140

Thr Ser Gly Thr Asp Ser Thr Ser Asp Ser Ser Asp Pro Met Gln Gln
 145 150 155 160

Leu Leu Lys Met Phe Ser Glu Ile Met Gln Ser Leu Phe Gly Asp Gly
 165 170 175

Gln Asp Gly Thr Gln Gly Ser Ser Ser Gly Gly Lys Gln Pro Thr Glu
 180 185 190

Gly Glu Gln Asn Ala Tyr Lys Lys Gly Val Thr Asp Ala Leu Ser Gly
 195 200 205
 Leu Met Gly Asn Gly Leu Ser Gln Leu Leu Gly Asn Gly Gly Leu Gly
 210 215 220
 Gly Gly Gln Gly Gly Asn Ala Gly Thr Gly Leu Asp Gly Ser Ser Leu
 225 230 235 240
 Gly Gly Lys Gly Leu Gln Asn Leu Ser Gly Pro Val Asp Tyr Gln Gln
 245 250 255
 Leu Gly Asn Ala Val Gly Thr Gly Ile Gly Met Lys Ala Gly Ile Gln
 260 265 270
 Ala Leu Asn Asp Ile Gly Thr His Arg His Ser Ser Thr Arg Ser Phe
 275 280 285
 Val Asn Lys Gly Asp Arg Ala Met Ala Lys Glu Ile Gly Gln Phe Met
 290 295 300
 Asp Gln Tyr Pro Glu Val Phe Gly Lys Pro Gln Tyr Gln Lys Gly Pro
 305 310 315 320
 Gly Gln Glu Val Lys Thr Asp Asp Lys Ser Trp Ala Lys Ala Leu Ser
 325 330 335
 Lys Pro Asp Asp Asp Gly Met Thr Pro Ala Ser Met Glu Gln Phe Asn
 340 345 350
 Lys Ala Lys Gly Met Ile Lys Arg Pro Met Ala Gly Asp Thr Gly Asn
 355 360 365
 Gly Asn Leu Gln Ala Arg Gly Ala Gly Gly Ser Ser Leu Gly Ile Asp
 370 375 380
 Ala Met Met Ala Gly Asp Ala Ile Asn Asn Met Ala Leu Gly Lys Leu
 385 390 395 400
 Gly Ala Ala

<210> 24

<211> 1288

<212> DNA

<213> Erwinia amylovora

<400> 24

```

aagcttcggc atggcacgtt tgaccgttgg gtcggcaggg tacgtttgaa ttattcataa 60
gaggaatacg ttatgagtct gaatacaagt gggctgggag cgtcaacgat gcaaatttct 120
atcggcggtg cgggcggaaa taacgggttg ctgggtacca gtcgccagaa tgctgggttg 180
ggtggcaatt ctgcactggg gctgggcggc ggtaatcaaa atgataccgt caatcagctg 240
gctggcttac tcaccggcat gatgatgatg atgagcatga tgggcgggtg tgggctgatg 300
ggcgggtggc taggcgggtg cttaggtaat ggcttgggtg gctcaggtgg cctgggcgaa 360
ggactgtcga acgcgctgaa cgatatgtta ggcggttcgc tgaacacgct gggctcga 420
ggcggaaca ataccacttc aacaacaaat tccccgctg accaggcgct gggattaac 480
tcaacgtccc aaaacgacga ttccacctcc ggcacagatt ccacctcaga ctccagcgac 540
ccgatgcagc agctgctgaa gatgttcagc gagataatgc aaagcctgtt tggatgatgg 600
caagatggca cccagggcag ttctctctgg ggcaagcagc cgaccgaagg cgagcagaac 660
gcctataaaa aaggagtcac tgatgcgctg tcgggcctga tgggtaatgg tctgagccag 720
ctccttggca acgggggact gggagggtgg cagggcggtg atgctggcac gggctctgac 780
ggttcgtcgc tgggcggcaa agggctgcaa aacctgagcg ggccgggtgga ctaccagcag 840
ttaggtaacg ccgtgggtac cggtatcggg atgaaagcgg gcattcaggc gctgaatgat 900
atcggtagcg acaggcacag ttcaaccctg tctttcgtca ataaaggcga tcgggcgatg 960
gcgaaggaaa tcggtcagtt catggaccag tatcctgagg tgtttggcaa gccgcagtac 1020
cagaaaggcc cgggtcagga ggtgaaaacc gatgacaaat catgggcaaa agcactgagc 1080
aagccagatg acgacggaat gacaccagcc agtatggagc agttcaaca agccaaggc 1140
atgatcaaaa ggcccatggc gggatgatac ggcaacggca acctgcaggc acgcggtgcc 1200
ggtggttctt cgctgggtat tgatgccatg atggccggtg atgccattaa caatatggca 1260
cttggcaagc tgggcgcggc ttaagctt

```

<210> 25

<211> 1344

<212> DNA

<213> *Erwinia amylovora*

<400> 25

```

atgtcaattc ttacgcttaa caacaatacc tcgtcctcgc cgggtctgtt ccagtccggg 60
ggggacaacg ggcttggtgg tcataatgca aattctcgtg tggggcaaca acccatcgat 120
cggcaaacca ttgagcaaat ggctcaatta ttggcggaac tggtaaagtc actgctatcg 180
ccacaatcag gtaatgcggc aaccggagcc ggtggcaatg accagactac aggagtgtgt 240
aacgctggcg gcctgaacgg acgaaaaggc acagcaggaa ccaactccga gtctgacagt 300
cagaacatgc tgagtgagat gggcaacaac gggctggatc aggccatcac gcccgatggc 360
cagggcgggc ggcagatcgg cgataatcct ttactgaaag ccatgctgaa gcttattgca 420
cgcagatgag acggccaaag cgatcagttt ggccaacctg gtacgggcaa caacagtgcc 480
tcttcgggta cttcttcacg tggcggttcc ctttttaacg atctatcagg ggggaaggcc 540
ccttcgggca actcccttc cggaactac tctccgtca gtaccttct accccatcc 600
acgccaacgt cccctacctc accgcttgat ttcccttctt ctcccacaa agcagccggg 660
ggcagcacgc cggtaaccga tcatcctgac cctgttggtg gcgcgggcat cggggccgga 720
aattcggtgg ccttcaccag cgccggcgct aatcagacgg tgctgcatga caccattacc 780
gtgaaagcgg gtcaggtgtt tgatggcaaa ggacaaacct tcaccgcccg ttcagaatta 840
ggcgatggcg gccagtctga aaaccagaaa ccgctgttta tactggaaga cgggtgccagc 900
ctgaaaaacg tcaccatggg cgacgacggg gcggatggta ttcatcttta cggatgatgcc 960
aaaatagaca atctgcacgt caccaacgtg ggtgaggacg cgattaccgt taagccaaac 1020

```

```

agcgcgggca aaaaatccca cgttgaaatc actaacagtt ccttcgagca cgcctctgac 1080
aagatcctgc agctgaatgc cgatactaac ctgagcggtg acaacgtgaa ggccaaagac 1140
tttggtactt ttgtacgcac taacggcggt caacagggtta actgggatct gaatctgagc 1200
catatcagcg cagaagacgg taagttctcg ttcgttaaaa gcgatagcga ggggctaaac 1260
gtcaatacca gtgatatctc actgggtgat gttgaaaacc actacaaagt gccgatgtcc 1320
gccaacctga aggtggctga atga                                     1344

```

<210> 26

<211> 447

<212> PRT

<213> *Erwinia amylovora*

<400> 26

```

Met Ser Ile Leu Thr Leu Asn Asn Asn Thr Ser Ser Ser Pro Gly Leu
  1                      5                      10                      15

```

```

Phe Gln Ser Gly Gly Asp Asn Gly Leu Gly Gly His Asn Ala Asn Ser
          20                      25                      30

```

```

Ala Leu Gly Gln Gln Pro Ile Asp Arg Gln Thr Ile Glu Gln Met Ala
      35                      40                      45

```

```

Gln Leu Leu Ala Glu Leu Leu Lys Ser Leu Leu Ser Pro Gln Ser Gly
  50                      55                      60

```

```

Asn Ala Ala Thr Gly Ala Gly Gly Asn Asp Gln Thr Thr Gly Val Gly
  65                      70                      75                      80

```

```

Asn Ala Gly Gly Leu Asn Gly Arg Lys Gly Thr Ala Gly Thr Thr Pro
          85                      90                      95

```

```

Gln Ser Asp Ser Gln Asn Met Leu Ser Glu Met Gly Asn Asn Gly Leu
      100                      105                      110

```

```

Asp Gln Ala Ile Thr Pro Asp Gly Gln Gly Gly Gly Gln Ile Gly Asp
      115                      120                      125

```

```

Asn Pro Leu Leu Lys Ala Met Leu Lys Leu Ile Ala Arg Met Met Asp
      130                      135                      140

```

```

Gly Gln Ser Asp Gln Phe Gly Gln Pro Gly Thr Gly Asn Asn Ser Ala
      145                      150                      155                      160

```

```

Ser Ser Gly Thr Ser Ser Ser Gly Gly Ser Pro Phe Asn Asp Leu Ser
          165                      170                      175

```

```

Gly Gly Lys Ala Pro Ser Gly Asn Ser Pro Ser Gly Asn Tyr Ser Pro
      180                      185                      190

```

Val Ser Thr Phe Ser Pro Pro Ser Thr Pro Thr Ser Pro Thr Ser Pro
 195 200 205

Leu Asp Phe Pro Ser Ser Pro Thr Lys Ala Ala Gly Gly Ser Thr Pro
 210 215 220

Val Thr Asp His Pro Asp Pro Val Gly Ser Ala Gly Ile Gly Ala Gly
 225 230 235 240

Asn Ser Val Ala Phe Thr Ser Ala Gly Ala Asn Gln Thr Val Leu His
 245 250 255

Asp Thr Ile Thr Val Lys Ala Gly Gln Val Phe Asp Gly Lys Gly Gln
 260 265 270

Thr Phe Thr Ala Gly Ser Glu Leu Gly Asp Gly Gly Gln Ser Glu Asn
 275 280 285

Gln Lys Pro Leu Phe Ile Leu Glu Asp Gly Ala Ser Leu Lys Asn Val
 290 295 300

Thr Met Gly Asp Asp Gly Ala Asp Gly Ile His Leu Tyr Gly Asp Ala
 305 310 315 320

Lys Ile Asp Asn Leu His Val Thr Asn Val Gly Glu Asp Ala Ile Thr
 325 330 335

Val Lys Pro Asn Ser Ala Gly Lys Lys Ser His Val Glu Ile Thr Asn
 340 345 350

Ser Ser Phe Glu His Ala Ser Asp Lys Ile Leu Gln Leu Asn Ala Asp
 355 360 365

Thr Asn Leu Ser Val Asp Asn Val Lys Ala Lys Asp Phe Gly Thr Phe
 370 375 380

Val Arg Thr Asn Gly Gly Gln Gln Gly Asn Trp Asp Leu Asn Leu Ser
 385 390 395 400

His Ile Ser Ala Glu Asp Gly Lys Phe Ser Phe Val Lys Ser Asp Ser
 405 410 415

Glu Gly Leu Asn Val Asn Thr Ser Asp Ile Ser Leu Gly Asp Val Glu
 420 425 430

Asn His Tyr Lys Val Pro Met Ser Ala Asn Leu Lys Val Ala Glu
 435 440 445

<210> 27
 <211> 5517
 <212> DNA
 <213> *Erwinia amylovora*

<400> 27

```

atggaattaa aatcactggg aactgaacac aaggcggcag tacacacagc ggcgcacaaac 60
cctgtggggc atggtgttgc cttacagcag ggcagcagca gcagcagccc gcaaaatgcc 120
gctgcatcat tggcggcaga aggcaaaaat cgtgggaaaa tgccgagaat tcaccagcca 180
tctactgcgg ctgatggtat cagcgtgct caccagcaaa agaaatcctt cagtctcagg 240
ggctgttttg ggacgaaaaa attttccaga tcggcaccgc agggccagcc aggtaccacc 300
cacagcaaa gggcaacatt gcgcatctg ctggcgcggg acgacggcga aacgcagcat 360
gaggcggccg cgccagatgc ggcgcgtttg acccgttcgg gcggcgtaa acgccgcaat 420
atggacgaca tggccgggcg gccaatggtg aaaggtggca gcggcgaaga taaggtacca 480
acgcagcaaa aacggcatca gctgaacaat tttggccaga tgccgcaaac gatgttgagc 540
aaaatggctc acccggcttc agccaacgcc ggcatcgcc tgcagcattc accgccgcac 600
atcccggtta gccaccacga aatcaaggaa gaaccggttg gctccaccag caaggcaaca 660
acggcccacg cagacagagt ggaaatcgct caggaagatg acgacagcga attccagcaa 720
ctgcatcaac agcggctggc gcggaacgg gaaaatccac cgcagccgcc caaactcggc 780
gttgccacac cgattagcgc caggtttcag cccaaactga ctgcggttgc ggaaagcgtc 840
cttgagggga cagataccac gcagtcaccc cttaaagccg aatcaatgct gaaaggaagt 900
ggagccgggg taacgccgct ggcggtaacg ctggataaag gcaagttgca gctggcaccg 960
gataatccac ccgcgctcaa tacgttgttg aagcagacat tgggtaaaga caccagcac 1020
tatctggcgc accatgccag cagcgacggg agccagcatc tgctgctgga caacaaaggc 1080
cacctgtttg atatcaaaag caccgccacc agctatagcg tgctgcacaa cagccacccc 1140
ggtgagataa agggcaagct ggcgcaggcg ggtactggct ccgtcagcgt agacggtaaa 1200
agcggcaaga tctcgtggg gagcggtacg caaagtcaca acaaaacaat gctaagccaa 1260
ccgggggaag cgcaccgttc cttattaacc ggcatttggc agcatcctgc tggcgccagc 1320
cggccgcagg gcgagtcaat ccgcctgcat gacgacaaaa ttcatactct gcatccggag 1380
ctgggcgtat ggcaatctgc ggataaagat acccacagcc agctgtctcg ccaggcagac 1440
ggtaagctct atgcgtgaa agacaaccgt accctgcaaa acctctccga taataaatcc 1500
tcagaaaagc tggtcgataa aatcaaatcg tattccgttg atcagcgggg gcaggtggcg 1560
atcctgacgg atactcccgg ccgccataag atgagtatta tgccctcgct ggatgcttcc 1620
ccggagagcc atatttccct cagcctgcat ttgcccgatg cccaccaggg gttattgcac 1680
gggaagtcgg agcttgaggc acaatctgtc gcgatcagcc atgggcgact ggttgtggcc 1740
gatagcgaag gcaagctgtt tagcgccgcc attccgaagc aaggggatgg aaacgaactg 1800
aaaatgaaag ccatgcctca gcatgcgctc gatgaacatt ttggtcatga ccaccagatt 1860
tctggatttt tccatgacga ccacggccag cttaatgcgc tggtgaaaaa taacttcagg 1920
cagcagcatg cctgcccgtt gggtaacgat catcagtttc accccggctg gaacctgact 1980
gatgcgctgg ttatcgacaa tcagctgggg ctgcatcata ccaatcctga accgcagtag 2040
attcttgata tggggcattt aggcagcctg gcgttacagg agggcaagct tcactatttt 2100
gaccagctga ccaaagggtg gactggcgcg gagtcaagatt gtaagcagct gaaaaaaggc 2160
ctggatggag cagcttatct actgaaagac ggtgaagtga aacgcctgaa tattaatcag 2220
agcacctcct ctatcaagca cggaacggaa aacgtttttt cgctgccgca tgtgcgcaat 2280
aaaccggagc cgggagatgc cctgcaaggg ctgaataaag acgataaggc ccaggccatg 2340
gcggtgattg gggtaaataa atacctggcg ctgacggaaa aaggggacat tcgctccttc 2400

```

```

cagataaaac cgggcaccca gcagttggag cggccggcac aaactctcag ccgcgaaggt 2460
atcagcggcg aactgaaaga cattcatgtc gaccacaagc agaacctgta tgccttgacc 2520
cacgagggag aggtgtttca tcagccgcgt gaagcctggc agaatgggtgc cgaaagcagc 2580
agctggcaca aactggcggt gccacagagt gaaagtaagc taaaaagtct ggacatgagc 2640
catgagcaca aaccgattgc cacctttgaa gacggtagcc agcatcagct gaaggtggc 2700
ggctggcacg cctatgcggc acctgaacgc gggccgctgg cgggtgggtac cagcggttca 2760
caaaccgtct ttaaccgact aatgcagggg gtgaaaggca aggtgatccc aggcagcggg 2820
ttgacggtta agctctcggc tcagacgggg ggaatgaccg gcgccgaagg gcgcaaggtc 2880
agcagtaaat tttccgaaag gatccgcgcc tatgcgttca acccaacaat gtccacgccg 2940
cgaccgatta aaaatgctgc ttatgccaca cagcacggct ggcagggggc tgaggggttg 3000
aagccgttgt acgagatgca gggagcgcgt attaaacaac tggatgcgca taacgttcgt 3060
cataacgcgc cacagccaga tttgcagagc aaactggaaa ctctggattt aggcgaacat 3120
ggcgcgagaat tgcttaacga catgaagcgc ttccgcgacg aactggagca gagtgaacc 3180
cgttcgggtga ccgttttagg tcaacatcag ggagtgcata aaagcaacgg tgaatcaat 3240
agcgaattta agccatcgcc cggcaaggcg ttggtccaga gctttaacgt caatcgctct 3300
ggtcaggatc taagcaagtc actgcaacag gcagtacatg ccacgcgcc atccgcagag 3360
agtaaactgc aatccatgct ggggcacttt gtcagtgcgc ggggtggatat gagtcatcag 3420
aagggcgaga tcccgcgtgg ccgccagcgc gatccgaatg ataaaaccgc actgaccaa 3480
tcgcgtttaa ttttagatac cgtgaccatc ggtgaactgc atgaactggc cgataaggcg 3540
aaactggtat ctgaccataa acccgatgcc gatcagataa aacagctgcg ccagcagttc 3600
gatacgctgc gtgaaaagcg gtatgagagc aatccggtga agcattacac cgatatggg 3660
ttcaccataa ataaggcgt ggaagcaaac tatgatgcgg tcaaagcctt tatcaatgcc 3720
tttaagaaag agcaccacgg cgtcaatctg accacgcgta ccgtactgga atcacagggc 3780
agtgcggagc tggcgaagaa gctcaagaat acgctgttgt ccctggacag tggtgaaagt 3840
atgagcttca gccggtcata tggcgggggc gtcagcactg tctttgtgcc tacccttagc 3900
aagaaggtgc cagttccggt gatccccgga gccggcatca cgctggatcg cgcctataac 3960
ctgagcttca gtcgtaccag cggcggattg aacgtcagtt ttggccgcga cggcggggtg 4020
agtggtaaca tcatggtcgc taccggccat gatgtgatgc cctatatgac cggtaagaaa 4080
accagtgcag gtaacgccag tgactgggtg agcgcaaaac ataaaatcag cccggacttg 4140
cgtatcggcg ctgctgtgag tggcaccctg caaggaacgc taaaaaacag cctgaagttt 4200
aagctgacag aggatgagct gcctggcttt atccatggct tgacgcatgg cacgttgacc 4260
ccggcagaac tgttgcaaaa ggggatcgaa catcagatga agcagggcag caaactgacg 4320
tttagcgtcg atacctcggc aaatctggat ctgcgtgcgc gtatcaatct gaacgaagac 4380
ggcagtaaac caaatgggtg cactgcccggt gtttctgccg ggctaagtgc atcggcaaac 4440
ctggccgccg gctcgcgtga acgcagcacc acctctggcc agtttggcag cacgacttcg 4500
gccagcaata accgccaac cttoctcaac ggggtcggcg cgggtgctaa cctgacggct 4560
gctttagggg ttgccattc atctacgcat gaagggaac cggtcgggat cttccggca 4620
tttacctcga ccaatgtttc ggcagcgtg gcgctggata accgtacctc acagagtatc 4680
agcctggaat tgaagcgcgc ggagccggtg accagcaacg atatcagcga gttgacctcc 4740
acgctgggaa aacactttaa ggatagcgcc acaacgaaga tgcttgccgc tctcaaagag 4800
ttagatgacg ctaagcccgc tgaacaactg catattttac agcagcattt cagtgcataa 4860
gatgtcgtcg gtgatgaacg ctacgaggcg gtgcgcaacc tgaaaaaact ggtgatacgt 4920
caacaggctg cggacagcca cagcatggaa ttaggatctg ccagtcacag cacgacctac 4980
aataatctgt cgagaataaa taatgacggc attgtcgagc tgctacacaa acatttcgat 5040
gcggcattac cagcaagcag tgccaaacgt cttggtgaaa tgatgaataa cgatccggca 5100
ctgaaagata ttattaagca gctgcaaagt acgccgttca gcagcgccag cgtgtcgatg 5160
gagctgaaag atggtctgcg tgagcagacg gaaaaagcaa tactggacgg taaggtcgg 5220
cgtgaagaag tgggagtact tttccaggat cgtaacaact tgcggtgtaa atcggtcagc 5280

```

gtcagtcagt cgcgcagcaa aagcgaaggc ttcaataccc cagcgctgtt actggggacg 5340
 agcaacagcg ctgctatgag catggagcgc aacatcggaa ccattaattt taaatacggc 5400
 caggatcaga acacccacg gcgatttacc ctggaggggtg gaatagctca ggctaataccg 5460
 caggtcgcac ctgcgcttac tgatttgaag aaggaagggc tggaaatgaa gagctaa 5517

<210> 28

<211> 1838

<212> PRT

<213> *Erwinia amylovora*

<400> 28

Met Glu Leu Lys Ser Leu Gly Thr Glu His Lys Ala Ala Val His Thr
 1 5 10 15

Ala Ala His Asn Pro Val Gly His Gly Val Ala Leu Gln Gln Gly Ser
 20 25 30

Ser Ser Ser Ser Pro Gln Asn Ala Ala Ala Ser Leu Ala Ala Glu Gly
 35 40 45

Lys Asn Arg Gly Lys Met Pro Arg Ile His Gln Pro Ser Thr Ala Ala
 50 55 60

Asp Gly Ile Ser Ala Ala His Gln Gln Lys Lys Ser Phe Ser Leu Arg
 65 70 75 80

Gly Cys Leu Gly Thr Lys Lys Phe Ser Arg Ser Ala Pro Gln Gly Gln
 85 90 95

Pro Gly Thr Thr His Ser Lys Gly Ala Thr Leu Arg Asp Leu Leu Ala
 100 105 110

Arg Asp Asp Gly Glu Thr Gln His Glu Ala Ala Ala Pro Asp Ala Ala
 115 120 125

Arg Leu Thr Arg Ser Gly Gly Val Lys Arg Arg Asn Met Asp Asp Met
 130 135 140

Ala Gly Arg Pro Met Val Lys Gly Gly Ser Gly Glu Asp Lys Val Pro
 145 150 155 160

Thr Gln Gln Lys Arg His Gln Leu Asn Asn Phe Gly Gln Met Arg Gln
 165 170 175

Thr Met Leu Ser Lys Met Ala His Pro Ala Ser Ala Asn Ala Gly Asp
 180 185 190

Arg Leu Gln His Ser Pro Pro His Ile Pro Gly Ser His His Glu Ile

195	200	205
Lys Glu Glu Pro Val Gly Ser Thr Ser Lys Ala Thr Thr Ala His Ala		
210	215	220
Asp Arg Val Glu Ile Ala Gln Glu Asp Asp Asp Ser Glu Phe Gln Gln		
225	230	235 240
Leu His Gln Gln Arg Leu Ala Arg Glu Arg Glu Asn Pro Pro Gln Pro		
	245	250 255
Pro Lys Leu Gly Val Ala Thr Pro Ile Ser Ala Arg Phe Gln Pro Lys		
	260	265 270
Leu Thr Ala Val Ala Glu Ser Val Leu Glu Gly Thr Asp Thr Thr Gln		
	275	280 285
Ser Pro Leu Lys Pro Gln Ser Met Leu Lys Gly Ser Gly Ala Gly Val		
	290	295 300
Thr Pro Leu Ala Val Thr Leu Asp Lys Gly Lys Leu Gln Leu Ala Pro		
305	310	315 320
Asp Asn Pro Pro Ala Leu Asn Thr Leu Leu Lys Gln Thr Leu Gly Lys		
	325	330 335
Asp Thr Gln His Tyr Leu Ala His His Ala Ser Ser Asp Gly Ser Gln		
	340	345 350
His Leu Leu Leu Asp Asn Lys Gly His Leu Phe Asp Ile Lys Ser Thr		
	355	360 365
Ala Thr Ser Tyr Ser Val Leu His Asn Ser His Pro Gly Glu Ile Lys		
	370	375 380
Gly Lys Leu Ala Gln Ala Gly Thr Gly Ser Val Ser Val Asp Gly Lys		
385	390	395 400
Ser Gly Lys Ile Ser Leu Gly Ser Gly Thr Gln Ser His Asn Lys Thr		
	405	410 415
Met Leu Ser Gln Pro Gly Glu Ala His Arg Ser Leu Leu Thr Gly Ile		
	420	425 430
Trp Gln His Pro Ala Gly Ala Ala Arg Pro Gln Gly Glu Ser Ile Arg		
	435	440 445
Leu His Asp Asp Lys Ile His Ile Leu His Pro Glu Leu Gly Val Trp		

450		455		460	
Gln Ser Ala Asp Lys Asp Thr His Ser Gln Leu Ser Arg Gln Ala Asp					
465		470		475	480
Gly Lys Leu Tyr Ala Leu Lys Asp Asn Arg Thr Leu Gln Asn Leu Ser					
	485		490		495
Asp Asn Lys Ser Ser Glu Lys Leu Val Asp Lys Ile Lys Ser Tyr Ser					
	500		505		510
Val Asp Gln Arg Gly Gln Val Ala Ile Leu Thr Asp Thr Pro Gly Arg					
	515		520		525
His Lys Met Ser Ile Met Pro Ser Leu Asp Ala Ser Pro Glu Ser His					
	530		535		540
Ile Ser Leu Ser Leu His Phe Ala Asp Ala His Gln Gly Leu Leu His					
	545		550		555
Gly Lys Ser Glu Leu Glu Ala Gln Ser Val Ala Ile Ser His Gly Arg					
	565		570		575
Leu Val Val Ala Asp Ser Glu Gly Lys Leu Phe Ser Ala Ala Ile Pro					
	580		585		590
Lys Gln Gly Asp Gly Asn Glu Leu Lys Met Lys Ala Met Pro Gln His					
	595		600		605
Ala Leu Asp Glu His Phe Gly His Asp His Gln Ile Ser Gly Phe Phe					
	610		615		620
His Asp Asp His Gly Gln Leu Asn Ala Leu Val Lys Asn Asn Phe Arg					
	625		630		635
Gln Gln His Ala Cys Pro Leu Gly Asn Asp His Gln Phe His Pro Gly					
	645		650		655
Trp Asn Leu Thr Asp Ala Leu Val Ile Asp Asn Gln Leu Gly Leu His					
	660		665		670
His Thr Asn Pro Glu Pro His Glu Ile Leu Asp Met Gly His Leu Gly					
	675		680		685
Ser Leu Ala Leu Gln Glu Gly Lys Leu His Tyr Phe Asp Gln Leu Thr					
	690		695		700
Lys Gly Trp Thr Gly Ala Glu Ser Asp Cys Lys Gln Leu Lys Lys Gly					

705		710		715		720
Leu Asp Gly Ala Ala Tyr Leu Leu Lys Asp Gly Glu Val Lys Arg Leu						
	725			730		735
Asn Ile Asn Gln Ser Thr Ser Ser Ile Lys His Gly Thr Glu Asn Val						
	740			745		750
Phe Ser Leu Pro His Val Arg Asn Lys Pro Glu Pro Gly Asp Ala Leu						
	755			760		765
Gln Gly Leu Asn Lys Asp Asp Lys Ala Gln Ala Met Ala Val Ile Gly						
	770			775		780
Val Asn Lys Tyr Leu Ala Leu Thr Glu Lys Gly Asp Ile Arg Ser Phe						
	785			790		795
Gln Ile Lys Pro Gly Thr Gln Gln Leu Glu Arg Pro Ala Gln Thr Leu						
		805		810		815
Ser Arg Glu Gly Ile Ser Gly Glu Leu Lys Asp Ile His Val Asp His						
	820			825		830
Lys Gln Asn Leu Tyr Ala Leu Thr His Glu Gly Glu Val Phe His Gln						
	835			840		845
Pro Arg Glu Ala Trp Gln Asn Gly Ala Glu Ser Ser Ser Trp His Lys						
	850			855		860
Leu Ala Leu Pro Gln Ser Glu Ser Lys Leu Lys Ser Leu Asp Met Ser						
	865			870		875
His Glu His Lys Pro Ile Ala Thr Phe Glu Asp Gly Ser Gln His Gln						
	885			890		895
Leu Lys Ala Gly Gly Trp His Ala Tyr Ala Ala Pro Glu Arg Gly Pro						
	900			905		910
Leu Ala Val Gly Thr Ser Gly Ser Gln Thr Val Phe Asn Arg Leu Met						
	915			920		925
Gln Gly Val Lys Gly Lys Val Ile Pro Gly Ser Gly Leu Thr Val Lys						
	930			935		940
Leu Ser Ala Gln Thr Gly Gly Met Thr Gly Ala Glu Gly Arg Lys Val						
	945			950		955
Ser Ser Lys Phe Ser Glu Arg Ile Arg Ala Tyr Ala Phe Asn Pro Thr						

965					970					975						
Met	Ser	Thr	Pro	Arg	Pro	Ile	Lys	Asn	Ala	Ala	Tyr	Ala	Thr	Gln	His	
980					985					990						
Gly	Trp	Gln	Gly	Arg	Glu	Gly	Leu	Lys	Pro	Leu	Tyr	Glu	Met	Gln	Gly	
995					1000					1005						
Ala	Leu	Ile	Lys	Gln	Leu	Asp	Ala	His	Asn	Val	Arg	His	Asn	Ala	Pro	
1010					1015					1020						
Gln	Pro	Asp	Leu	Gln	Ser	Lys	Leu	Glu	Thr	Leu	Asp	Leu	Gly	Glu	His	
1025					1030					1035					1040	
Gly	Ala	Glu	Leu	Leu	Asn	Asp	Met	Lys	Arg	Phe	Arg	Asp	Glu	Leu	Glu	
1045					1050					1055						
Gln	Ser	Ala	Thr	Arg	Ser	Val	Thr	Val	Leu	Gly	Gln	His	Gln	Gly	Val	
1060					1065					1070						
Leu	Lys	Ser	Asn	Gly	Glu	Ile	Asn	Ser	Glu	Phe	Lys	Pro	Ser	Pro	Gly	
1075					1080					1085						
Lys	Ala	Leu	Val	Gln	Ser	Phe	Asn	Val	Asn	Arg	Ser	Gly	Gln	Asp	Leu	
1090					1095					1100						
Ser	Lys	Ser	Leu	Gln	Gln	Ala	Val	His	Ala	Thr	Pro	Pro	Ser	Ala	Glu	
1105					1110					1115					1120	
Ser	Lys	Leu	Gln	Ser	Met	Leu	Gly	His	Phe	Val	Ser	Ala	Gly	Val	Asp	
1125					1130					1135						
Met	Ser	His	Gln	Lys	Gly	Glu	Ile	Pro	Leu	Gly	Arg	Gln	Arg	Asp	Pro	
1140					1145					1150						
Asn	Asp	Lys	Thr	Ala	Leu	Thr	Lys	Ser	Arg	Leu	Ile	Leu	Asp	Thr	Val	
1155					1160					1165						
Thr	Ile	Gly	Glu	Leu	His	Glu	Leu	Ala	Asp	Lys	Ala	Lys	Leu	Val	Ser	
1170					1175					1180						
Asp	His	Lys	Pro	Asp	Ala	Asp	Gln	Ile	Lys	Gln	Leu	Arg	Gln	Gln	Phe	
1185					1190					1195					1200	
Asp	Thr	Leu	Arg	Glu	Lys	Arg	Tyr	Glu	Ser	Asn	Pro	Val	Lys	His	Tyr	
1205					1210					1215						
Thr	Asp	Met	Gly	Phe	Thr	His	Asn	Lys	Ala	Leu	Glu	Ala	Asn	Tyr	Asp	

1220	1225	1230
Ala Val Lys Ala Phe Ile Asn Ala Phe Lys Lys Glu His His Gly Val		
1235	1240	1245
Asn Leu Thr Thr Arg Thr Val Leu Glu Ser Gln Gly Ser Ala Glu Leu		
1250	1255	1260
Ala Lys Lys Leu Lys Asn Thr Leu Leu Ser Leu Asp Ser Gly Glu Ser		
1265	1270	1275
Met Ser Phe Ser Arg Ser Tyr Gly Gly Gly Val Ser Thr Val Phe Val		
1285	1290	1295
Pro Thr Leu Ser Lys Lys Val Pro Val Pro Val Ile Pro Gly Ala Gly		
1300	1305	1310
Ile Thr Leu Asp Arg Ala Tyr Asn Leu Ser Phe Ser Arg Thr Ser Gly		
1315	1320	1325
Gly Leu Asn Val Ser Phe Gly Arg Asp Gly Gly Val Ser Gly Asn Ile		
1330	1335	1340
Met Val Ala Thr Gly His Asp Val Met Pro Tyr Met Thr Gly Lys Lys		
1345	1350	1355
Thr Ser Ala Gly Asn Ala Ser Asp Trp Leu Ser Ala Lys His Lys Ile		
1365	1370	1375
Ser Pro Asp Leu Arg Ile Gly Ala Ala Val Ser Gly Thr Leu Gln Gly		
1380	1385	1390
Thr Leu Gln Asn Ser Leu Lys Phe Lys Leu Thr Glu Asp Glu Leu Pro		
1395	1400	1405
Gly Phe Ile His Gly Leu Thr His Gly Thr Leu Thr Pro Ala Glu Leu		
1410	1415	1420
Leu Gln Lys Gly Ile Glu His Gln Met Lys Gln Gly Ser Lys Leu Thr		
1425	1430	1435
Phe Ser Val Asp Thr Ser Ala Asn Leu Asp Leu Arg Ala Gly Ile Asn		
1445	1450	1455
Leu Asn Glu Asp Gly Ser Lys Pro Asn Gly Val Thr Ala Arg Val Ser		
1460	1465	1470
Ala Gly Leu Ser Ala Ser Ala Asn Leu Ala Ala Gly Ser Arg Glu Arg		

1475	1480	1485
Ser Thr Thr Ser Gly Gln Phe Gly Ser Thr Thr Ser Ala Ser Asn Asn		
1490	1495	1500
Arg Pro Thr Phe Leu Asn Gly Val Gly Ala Gly Ala Asn Leu Thr Ala		
1505	1510	1515 1520
Ala Leu Gly Val Ala His Ser Ser Thr His Glu Gly Lys Pro Val Gly		
1525	1530	1535
Ile Phe Pro Ala Phe Thr Ser Thr Asn Val Ser Ala Ala Leu Ala Leu		
1540	1545	1550
Asp Asn Arg Thr Ser Gln Ser Ile Ser Leu Glu Leu Lys Arg Ala Glu		
1555	1560	1565
Pro Val Thr Ser Asn Asp Ile Ser Glu Leu Thr Ser Thr Leu Gly Lys		
1570	1575	1580
His Phe Lys Asp Ser Ala Thr Thr Lys Met Leu Ala Ala Leu Lys Glu		
1585	1590	1595 1600
Leu Asp Asp Ala Lys Pro Ala Glu Gln Leu His Ile Leu Gln Gln His		
1605	1610	1615
Phe Ser Ala Lys Asp Val Val Gly Asp Glu Arg Tyr Glu Ala Val Arg		
1620	1625	1630
Asn Leu Lys Lys Leu Val Ile Arg Gln Gln Ala Ala Asp Ser His Ser		
1635	1640	1645
Met Glu Leu Gly Ser Ala Ser His Ser Thr Thr Tyr Asn Asn Leu Ser		
1650	1655	1660
Arg Ile Asn Asn Asp Gly Ile Val Glu Leu Leu His Lys His Phe Asp		
1665	1670	1675 1680
Ala Ala Leu Pro Ala Ser Ser Ala Lys Arg Leu Gly Glu Met Met Asn		
1685	1690	1695
Asn Asp Pro Ala Leu Lys Asp Ile Ile Lys Gln Leu Gln Ser Thr Pro		
1700	1705	1710
Phe Ser Ser Ala Ser Val Ser Met Glu Leu Lys Asp Gly Leu Arg Glu		
1715	1720	1725
Gln Thr Glu Lys Ala Ile Leu Asp Gly Lys Val Gly Arg Glu Glu Val		

1730 1735 1740

Gly Val Leu Phe Gln Asp Arg Asn Asn Leu Arg Val Lys Ser Val Ser
 1745 1750 1755 1760

Val Ser Gln Ser Val Ser Lys Ser Glu Gly Phe Asn Thr Pro Ala Leu
 1765 1770 1775

Leu Leu Gly Thr Ser Asn Ser Ala Ala Met Ser Met Glu Arg Asn Ile
 1780 1785 1790

Gly Thr Ile Asn Phe Lys Tyr Gly Gln Asp Gln Asn Thr Pro Arg Arg
 1795 1800 1805

Phe Thr Leu Glu Gly Gly Ile Ala Gln Ala Asn Pro Gln Val Ala Ser
 1810 1815 1820

Ala Leu Thr Asp Leu Lys Lys Glu Gly Leu Glu Met Lys Ser
 1825 1830 1835

<210> 29

<211> 420

<212> DNA

<213> *Erwinia amylovora*

<400> 29

atgacatcgt cacagcagcg ggttgaaagg tttttacagt atttctccgc cgggtgtaaa 60
 acgcccatac atctgaaaga cgggggtgtgc gccctgtata acgaacaaga tgaggaggcg 120
 gcggtgctgg aagtaccgca acacagcgac agcctgttac tacactgccg aatcattgag 180
 gctgaccac aaacttcaat aaccctgtat tcgatgctat tacagctgaa ttttgaaatg 240
 gcggccatgc gcggtgttg gctggcgctg gatgaactgc acaacgtgcg tttatgtttt 300
 cagcagtcgc tggagcatct ggatgaagca agtttttagcg atatcgttag cggttcatc 360
 gaacatgcgg cagaagtgcg tgagtatata gcgcaattag acgagagtag cgcggcataa 420

<210> 30

<211> 139

<212> PRT

<213> *Erwinia amylovora*

<400> 30

Met Thr Ser Ser Gln Gln Arg Val Glu Arg Phe Leu Gln Tyr Phe Ser
 1 5 10 15

Ala Gly Cys Lys Thr Pro Ile His Leu Lys Asp Gly Val Cys Ala Leu
 20 25 30

Tyr Asn Glu Gln Asp Glu Glu Ala Ala Val Leu Glu Val Pro Gln His

35 40 45
 Ser Asp Ser Leu Leu Leu His Cys Arg Ile Ile Glu Ala Asp Pro Gln
 50 55 60
 Thr Ser Ile Thr Leu Tyr Ser Met Leu Leu Gln Leu Asn Phe Glu Met
 65 70 75 80
 Ala Ala Met Arg Gly Cys Trp Leu Ala Leu Asp Glu Leu His Asn Val
 85 90 95
 Arg Leu Cys Phe Gln Gln Ser Leu Glu His Leu Asp Glu Ala Ser Phe
 100 105 110
 Ser Asp Ile Val Ser Gly Phe Ile Glu His Ala Ala Glu Val Arg Glu
 115 120 125
 Tyr Ile Ala Gln Leu Asp Glu Ser Ser Ala Ala
 130 135

<210> 31

<211> 341

<212> PRT

<213> *Pseudomonas syringae*

<400> 31

Met Gln Ser Leu Ser Leu Asn Ser Ser Ser Leu Gln Thr Pro Ala Met
 1 5 10 15
 Ala Leu Val Leu Val Arg Pro Glu Ala Glu Thr Thr Gly Ser Thr Ser
 20 25 30
 Ser Lys Ala Leu Gln Glu Val Val Val Lys Leu Ala Glu Glu Leu Met
 35 40 45
 Arg Asn Gly Gln Leu Asp Asp Ser Ser Pro Leu Gly Lys Leu Leu Ala
 50 55 60
 Lys Ser Met Ala Ala Asp Gly Lys Ala Gly Gly Gly Ile Glu Asp Val
 65 70 75 80
 Ile Ala Ala Leu Asp Lys Leu Ile His Glu Lys Leu Gly Asp Asn Phe
 85 90 95
 Gly Ala Ser Ala Asp Ser Ala Ser Gly Thr Gly Gln Gln Asp Leu Met
 100 105 110

Thr Gln Val Leu Asn Gly Leu Ala Lys Ser Met Leu Asp Asp Leu Leu
 115 120 125
 Thr Lys Gln Asp Gly Gly Thr Ser Phe Ser Glu Asp Asp Met Pro Met
 130 135 140
 Leu Asn Lys Ile Ala Gln Phe Met Asp Asp Asn Pro Ala Gln Phe Pro
 145 150 155 160
 Lys Pro Asp Ser Gly Ser Trp Val Asn Glu Leu Lys Glu Asp Asn Phe
 165 170 175
 Leu Asp Gly Asp Glu Thr Ala Ala Phe Arg Ser Ala Leu Asp Ile Ile
 180 185 190
 Gly Gln Gln Leu Gly Asn Gln Gln Ser Asp Ala Gly Ser Leu Ala Gly
 195 200 205
 Thr Gly Gly Gly Leu Gly Thr Pro Ser Ser Phe Ser Asn Asn Ser Ser
 210 215 220
 Val Met Gly Asp Pro Leu Ile Asp Ala Asn Thr Gly Pro Gly Asp Ser
 225 230 235 240
 Gly Asn Thr Arg Gly Glu Ala Gly Gln Leu Ile Gly Glu Leu Ile Asp
 245 250 255
 Arg Gly Leu Gln Ser Val Leu Ala Gly Gly Gly Leu Gly Thr Pro Val
 260 265 270
 Asn Thr Pro Gln Thr Gly Thr Ser Ala Asn Gly Gly Gln Ser Ala Gln
 275 280 285
 Asp Leu Asp Gln Leu Leu Gly Gly Leu Leu Leu Lys Gly Leu Glu Ala
 290 295 300
 Thr Leu Lys Asp Ala Gly Gln Thr Gly Thr Asp Val Gln Ser Ser Ala
 305 310 315 320
 Ala Gln Ile Ala Thr Leu Leu Val Ser Thr Leu Leu Gln Gly Thr Arg
 325 330 335
 Asn Gln Ala Ala Ala
 340

<210> 32

<211> 1026

<212> DNA

<213> *Pseudomonas syringae*

<400> 32

```

atgcagagtc tcagtcttaa cagcagctcg ctgcaaacc cggcaatggc ccttgtcctg 60
gtacgtcctg aagccgagac gactggcagt acgtcgagca aggcgcttca ggaagtgtgc 120
gtgaagctgg ccgaggaact gatgcgcaat ggtcaactcg acgacagctc gccattggga 180
aaactgttgg ccaagtcgat ggccgcagat ggcaaggcgg gcggcggtat tgaggatgtc 240
atcgctgcgc tggacaagct gatccatgaa aagctcggtg acaacttcgg cgcgtctgcg 300
gacagcgctt cgggtaccgg acagcaggac ctgatgactc aggtgctcaa tggcctggcc 360
aagtcgatgc tcgatgatct tctgaccaag caggatggcg ggacaagctt ctccgaagac 420
gatatgccga tgctgaacaa gatcgcgcag ttcattggatg acaatcccgc acagtttccc 480
aagccggact cgggctcctg ggtgaacgaa ctcaagggaag acaacttcct tgatggcgac 540
gaaacggctg cgttccgttc ggactcgcac atcattggcc agcaactggg taatcagcag 600
agtacgctg gcagtctggc agggacgggt ggaggctctg gcactccgag cagtttttcc 660
aacaactcgt ccgtgatggg tgatccgctg atcgacgcca ataccggtcc cggtgacagc 720
ggcaataccc gtggtgaagc ggggcaactg atcggcgagc ttatcgaccg tggcctgcaa 780
tcggtattgg ccggtggtgg actgggcaca cccgtaaaca ccccgagac cggtagctcg 840
gcgaatggcg gacagtccgc tcaggatctt gatcagttgc tgggcggtt gctgctcaag 900
ggcctggagg caacgctcaa ggatgccggg caaacaggca ccgacgtgca gtcgagcgct 960
gcgcaaatcg ccaccttgct ggtcagtagc ctgctgcaag gcacccgcaa tcaggctgca 1020
gcctga                                     1026

```

<210> 33

<211> 1729

<212> DNA

<213> *Pseudomonas syringae*

<400> 33

```

tccacttcgc tgattttgaa attggcagat tcatagaaac gttcagggtg ggaaatcagg 60
ctgagtgcgc agatttcgtt gataagggtg tggtagctgg cattgttggc catttcaagg 120
cctctgagtg cgggtgcggag caataccagt ctctctgctg gcgtgtgcac actgagtcgc 180
aggcataggc atttcagttc cttgcgttgg ttgggcatat aaaaaaagga acttttataa 240
acagtgcgat gagatgccgg caaacgggga accggtcgct gcgctttgcc actcaacttcg 300
agcaagctca accccaaaca tccacatccc tatcgaaagg acagcgatac ggccacttgc 360
tctggtaaac cctggagctg gcgtcgggtc aattgcccac ttagcgaggt aacgcagcat 420
gagcatcggc atcacacccc ggccgcaaca gaccaccacg ccaactcgatt tttcggcgct 480
aagcggcaag agtcctcaac caaacacgtt cggcgagcag aacactcagc aagcgatcga 540
cccagtgca ctgttggtcg gcagcgacac acagaaagac gtcaacttcg gcacgcccga 600
cagcaccgtc cagaatccgc aggacgccag caagcccaac gacagccagt ccaacatcgc 660
taaattgatc agtgcattga tcatgtcgtt gctgcagatg ctcaccaact ccaataaaaa 720
gcaggacacc aatcaggaac agcctgatag ccaggctcct ttccagaaca acggcgggct 780
cggtagaccg tcggccgata gcggggggcg cggtagaccg gatgacgacg gtggcgggcg 840
cggtgatacg ccaagcgcaa caggcggtgg cggcggtgat actccgaccg caacaggcgg 900
tgccggcagc ggtggcgggc gcacaccac tgcaacaggg ggcggcagcg gtggcacacc 960
cactgcaaca ggcggtggcg aggggtggcg aacaccgcaa atcactccgc agttggccaa 1020
ccctaaccgt acctcaggta ctggctcggg gtcggacacc gcaggttcta ccgagcaagc 1080
cggcaagatc aatgtggtga aagacaccat caaggctcgg gctggcgaag tctttgacgg 1140

```

```

ccacggcgca accttcactg ccgacaaatc tatgggtaac ggagaccagg gcgaaaatca 1200
gaagcccatg ttcgagctgg ctgaaggcgc tacgttgaag aatgtgaacc tgggtgagaa 1260
cgaggtcgat ggcattccacg tgaaagccaa aaacgctcag gaagtcacca ttgacaacgt 1320
gcatgcccag aacgtcgggtg aagacctgat tacgggtcaaa ggcgagggag ggcgagcgggt 1380
cactaatctg aacatcaaga acagcagtgc caaagggtgca gacgacaagg ttgtccagct 1440
caacgccaac actcacttga aaatcgacaa cttcaaggcc gacgatttcg gcacgatgggt 1500
tcgcaccaac ggtggcaagc agtttgatga catgagcatc gagctgaacg gcacgaaagc 1560
taaccacggc aagttcgccc tggtgaaaag cgacagtgc gatctgaagc tggcaacggg 1620
caacatcgcc atgaccgacg tcaaacacgc ctacgataaa acccagggcat cgaccaaca 1680
caccgagctt tgaatccaga caagtagctt gaaaaaaggg ggtggactc 1729

```

<210> 34

<211> 424

<212> PRT

<213> *Pseudomonas syringae*

<400> 34

```

Met Ser Ile Gly Ile Thr Pro Arg Pro Gln Gln Thr Thr Thr Pro Leu
  1              5              10              15

Asp Phe Ser Ala Leu Ser Gly Lys Ser Pro Gln Pro Asn Thr Phe Gly
          20              25              30

Glu Gln Asn Thr Gln Gln Ala Ile Asp Pro Ser Ala Leu Leu Phe Gly
          35              40              45

Ser Asp Thr Gln Lys Asp Val Asn Phe Gly Thr Pro Asp Ser Thr Val
          50              55              60

Gln Asn Pro Gln Asp Ala Ser Lys Pro Asn Asp Ser Gln Ser Asn Ile
          65              70              75              80

Ala Lys Leu Ile Ser Ala Leu Ile Met Ser Leu Leu Gln Met Leu Thr
          85              90              95

Asn Ser Asn Lys Lys Gln Asp Thr Asn Gln Glu Gln Pro Asp Ser Gln
          100             105             110

Ala Pro Phe Gln Asn Asn Gly Gly Leu Gly Thr Pro Ser Ala Asp Ser
          115             120             125

Gly Gly Gly Gly Thr Pro Asp Ala Thr Gly Gly Gly Gly Gly Asp Thr
          130             135             140

Pro Ser Ala Thr Gly Gly Gly Gly Gly Asp Thr Pro Thr Ala Thr Gly
          145             150             155             160

Gly Gly Gly Ser Gly Gly Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly

```

165	170	175
Ser Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly Glu Gly Gly Val Thr		
180	185	190
Pro Gln Ile Thr Pro Gln Leu Ala Asn Pro Asn Arg Thr Ser Gly Thr		
195	200	205
Gly Ser Val Ser Asp Thr Ala Gly Ser Thr Glu Gln Ala Gly Lys Ile		
210	215	220
Asn Val Val Lys Asp Thr Ile Lys Val Gly Ala Gly Glu Val Phe Asp		
225	230	235
Gly His Gly Ala Thr Phe Thr Ala Asp Lys Ser Met Gly Asn Gly Asp		
245	250	255
Gln Gly Glu Asn Gln Lys Pro Met Phe Glu Leu Ala Glu Gly Ala Thr		
260	265	270
Leu Lys Asn Val Asn Leu Gly Glu Asn Glu Val Asp Gly Ile His Val		
275	280	285
Lys Ala Lys Asn Ala Gln Glu Val Thr Ile Asp Asn Val His Ala Gln		
290	295	300
Asn Val Gly Glu Asp Leu Ile Thr Val Lys Gly Glu Gly Gly Ala Ala		
305	310	315
Val Thr Asn Leu Asn Ile Lys Asn Ser Ser Ala Lys Gly Ala Asp Asp		
325	330	335
Lys Val Val Gln Leu Asn Ala Asn Thr His Leu Lys Ile Asp Asn Phe		
340	345	350
Lys Ala Asp Asp Phe Gly Thr Met Val Arg Thr Asn Gly Gly Lys Gln		
355	360	365
Phe Asp Asp Met Ser Ile Glu Leu Asn Gly Ile Glu Ala Asn His Gly		
370	375	380
Lys Phe Ala Leu Val Lys Ser Asp Ser Asp Asp Leu Lys Leu Ala Thr		
385	390	395
Gly Asn Ile Ala Met Thr Asp Val Lys His Ala Tyr Asp Lys Thr Gln		
405	410	415
Ala Ser Thr Gln His Thr Glu Leu		

420

<210> 35

<211> 344

<212> PRT

<213> *Pseudomonas solanacearum*

<220>

<223> Description of Unknown Organism: *Pseudomonas solanacearum*

<400> 35

```

Met Ser Val Gly Asn Ile Gln Ser Pro Ser Asn Leu Pro Gly Leu Gln
 1             5             10             15

Asn Leu Asn Leu Asn Thr Asn Thr Asn Ser Gln Gln Ser Gly Gln Ser
      20             25             30

Val Gln Asp Leu Ile Lys Gln Val Glu Lys Asp Ile Leu Asn Ile Ile
      35             40             45

Ala Ala Leu Val Gln Lys Ala Ala Gln Ser Ala Gly Gly Asn Thr Gly
      50             55             60

Asn Thr Gly Asn Ala Pro Ala Lys Asp Gly Asn Ala Asn Ala Gly Ala
      65             70             75             80

Asn Asp Pro Ser Lys Asn Asp Pro Ser Lys Ser Gln Ala Pro Gln Ser
      85             90             95

Ala Asn Lys Thr Gly Asn Val Asp Asp Ala Asn Asn Gln Asp Pro Met
      100             105             110

Gln Ala Leu Met Gln Leu Leu Glu Asp Leu Val Lys Leu Leu Lys Ala
      115             120             125

Ala Leu His Met Gln Gln Pro Gly Gly Asn Asp Lys Gly Asn Gly Val
      130             135             140

Gly Gly Ala Asn Gly Ala Lys Gly Ala Gly Gly Gln Gly Gly Leu Ala
      145             150             155             160

Glu Ala Leu Gln Glu Ile Glu Gln Ile Leu Ala Gln Leu Gly Gly Gly
      165             170             175

Gly Ala Gly Ala Gly Gly Ala Gly Gly Gly Val Gly Gly Ala Gly Gly
      180             185             190

```

Ala Asp Gly Gly Ser Gly Ala Gly Gly Ala Gly Gly Ala Asn Gly Ala
 195 200 205

Asp Gly Gly Asn Gly Val Asn Gly Asn Gln Ala Asn Gly Pro Gln Asn
 210 215 220

Ala Gly Asp Val Asn Gly Ala Asn Gly Ala Asp Asp Gly Ser Glu Asp
 225 230 235 240

Gln Gly Gly Leu Thr Gly Val Leu Gln Lys Leu Met Lys Ile Leu Asn
 245 250 255

Ala Leu Val Gln Met Met Gln Gln Gly Gly Leu Gly Gly Gly Asn Gln
 260 265 270

Ala Gln Gly Gly Ser Lys Gly Ala Gly Asn Ala Ser Pro Ala Ser Gly
 275 280 285

Ala Asn Pro Gly Ala Asn Gln Pro Gly Ser Ala Asp Asp Gln Ser Ser
 290 295 300

Gly Gln Asn Asn Leu Gln Ser Gln Ile Met Asp Val Val Lys Glu Val
 305 310 315 320

Val Gln Ile Leu Gln Gln Met Leu Ala Ala Gln Asn Gly Gly Ser Gln
 325 330 335

Gln Ser Thr Ser Thr Gln Pro Met
 340

<210> 36

<211> 1035

<212> DNA

<213> *Pseudomonas solanacearum*

<400> 36

atgtcagtcg gaaacatcca gagcccgtcg aacctcccgg gtctgcagaa cctgaacctc 60
 aacaccaaca ccaacagcca gcaatcgggc cagtccgtgc aagacctgat caagcagggtc 120
 gagaaggaca tcctcaacat catcgcagcc ctcgtgcaga aggccgcaca gtcggcgggc 180
 ggcaacaccg gtaacaccgg caacgcgcgg gcgaaggacg gcaatgcca cgcgggcggc 240
 aacgacccga gcaagaacga cccgagcaag agccaggctc cgcagtcggc caacaagacc 300
 ggcaacgtcg acgacgcca caaccaggat ccgatgcaag cgctgatgca gctgctggaa 360
 gacctggtga agctgctgaa ggccggccctg cacatgcagc agcccgggcg caatgacaag 420
 ggcaacggcg tgggcggtgc caacggcgcc aagggtgccc gcggccaggg cggcctggcc 480
 gaagcgctgc aggagatcga gcagatcctc gccagctcg gcggcgggcg tgctggcgcc 540
 ggccggcgcg gtggcggtgt cggcggtgct ggtggcgcg atggcggtc cgggtcgggg 600

```

ggcgcaggcg gtgcgaacgg cgccgacggc ggcaatggcg tgaacggcaa ccaggcgaac 660
ggccccgaga acgcaggcga tgtcaacggg gccaacggcg cggatgacgg cagcgaagac 720
cagggcgggc tcaccggcgt gctgcaaaag ctgatgaaga tcctgaacgc gctgggtgcag 780
atgatgcagc aaggcggcct cggcggcggc aaccaggcgc agggcgggctc gaagggtgcc 840
ggcaacgcct cgccgggcttc cggcgcgaac ccgggcgcga accagcccgg ttcggcggat 900
gatcaatcgt ccggccagaa caatctgcaa tcccagatca tggatgtggt gaaggaggtc 960
gtccagatcc tgcagcagat gctggcgggc cagaacggcg gcagccagca gtccacctcg 1020
acgcagccga tgtaa 1035

```

<210> 37

<211> 26

<212> PRT

<213> *Xanthomonas campestris* pv. *glycines*

<400> 37

```

Thr Leu Ile Glu Leu Met Ile Val Val Ala Ile Ile Ala Ile Leu Ala
  1             5             10             15

```

```

Ala Ile Ala Leu Pro Ala Tyr Gln Asp Tyr
          20             25

```

<210> 38

<211> 20

<212> PRT

<213> *Xanthomonas campestris* pv. *pelargonii*

<400> 38

```

Ser Ser Gln Gln Ser Pro Ser Ala Gly Ser Glu Gln Gln Leu Asp Gln
  1             5             10             15

```

```

Leu Leu Ala Met
          20

```

<210> 39

<211> 13

<212> PRT

<213> *Phytophthora megasperma*

<400> 39

```

Val Trp Asn Gln Pro Val Arg Gly Phe Lys Val Tyr Glu
  1             5             10

```