



- (51) **International Patent Classification:**
C07K 14/745 (2006.01) A61P 7/04 (2006.01)
A61K 38/36 (2006.01)
- (21) **International Application Number:**
PCT/IB2014/064564
- (22) **International Filing Date:**
16 September 2014 (16.09.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/881,834 24 September 2013 (24.09.2013) US
- (71) **Applicant:** PFIZER INC. [US/US]; 235 East 42nd Street, New York, New York 10017 (US).
- (72) **Inventors:** JANKOWSKI, Michael Anthony; 10 Paul's Way, Newbury, Massachusetts 01951 (US). JOHNSON, Keith A.; 45 Meadowlark Road, North Chelmsford, Massachusetts 01863 (US). PIACENZA, Wendy Carol; 7 Clinton Avenue, Chelmsford, Massachusetts 01824 (US). ROUSE, Jason C.; 25 Lantern Lane, Londonderry, New Hampshire 03053 (US). SHAMASHKIN, Michael; 244 Russell St., Woburn, Massachusetts 01801 (US). SHARPE, Penelope Jane; 61 Redfield Circle, Derry, New Hampshire 03038 (US). SWITZER, Mary Beth; 235 Farnum Street, North Andover, Massachusetts 01845 (US).

WESTON, Stacey B.; 114 Sesame Street, Dracut, Massachusetts 01826 (US).

- (74) **Agent:** KLEIMAN, Gabriel L.; Pfizer Inc., 235 East 42nd Street, MS 235/9/64, New York, New York 10017 (US).

- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to the identity of the inventor (Rule 4.17(i))

[Continued on next page]

- (54) **Title:** COMPOSITIONS COMPRISING HETEROGENEOUS POPULATIONS OF RECOMBINANT HUMAN CLOTTING FACTOR XA PROTEINS




FIG. 5

- (57) **Abstract:** Compositions are provided comprising recombinant variants of the human clotting Factor Xa. Such compositions include a wide variety of isoforms and post-translational modifications of FXa and are useful for treating subjects in need of hemostasis.



-
- | | |
|---|--|
| — as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) | — before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h)) |
| — as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) | — with sequence listing part of description (Rule 5.2(a)) |

Published:

- with international search report (Art. 21(3))

COMPOSITIONS COMPRISING HETEROGENEOUS POPULATIONS OF
RECOMBINANT HUMAN CLOTTING FACTOR XA PROTEINS

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of U.S. Provisional Application No. 61/881,834, filed September 24, 2013, which is incorporated by reference in its entirety.

REFERENCE TO SEQUENCE LISTING

[0002] The Sequence Listing submitted concurrently herewith under 37 CFR §1.821 in a computer readable form (CRF) via EFS-Web as file name PC72049_SEQLIST_ST25.txt is incorporated herein by reference. The electronic copy of the Sequence Listing was created on 10 Sep 2014, with a file size of 31 kilobytes.

BACKGROUND OF THE INVENTION

[0003] Effective therapies are needed to control excessive bleeding in a range of clinical conditions where bleeding cannot be adequately controlled by medical or surgical intervention. This unmet need is particularly critical among patients with hemophilia, especially those in whom factor replacement therapy is rendered less efficacious due to the production of inhibitor antibodies.

[0004] Activated clotting Factor X (FXa) occupies a central position in the coagulation cascade at the convergence of the intrinsic and extrinsic coagulation pathways. Membrane-bound FXa, in the presence of its cofactor Factor Va (FVa), converts prothrombin to thrombin, which activates platelets and converts fibrinogen to fibrin to form the thrombus. In principle, replacement therapy with direct FXa administration could correct bleeding. The therapeutic potential of FXa is limited, however, due to its very short plasma half-life and potential for inducing excessive coagulation due to activation of other coagulation factors.

[0005] Earlier work identified a FXa variant (I16L variant) in which leucine replaced isoleucine at the amino-terminus of the wild-type FXa heavy chain (position 16 in the chymotrypsin numbering scheme). The substitution yielded a FXa variant with zymogen-like characteristics. Toso, R., et al., The conformational switch from the factor X zymogen to protease state mediates exosite expression and prothrombinase assembly. J. Biol. Chem. 283, 18627–18635 (2008); Ivanciu L., et al., A zymogen-like factor Xa variant corrects the coagulation defect in hemophilia. Nat. Biotechnol. 29:1028-33 (2011).

[0006] When not incorporated into the prothrombinase complex with its cofactor Factor Va (FVa), the FXa I16L variant had no significant catalytic activity and was better protected from inactivation by serum protease inhibitors compared to wild-type FXa. As a result, the variant had longer serum half-life compared to wild-type FXa. Binding to FVa in prothrombinase, however, caused the variant to transition from the zymogen-like state to the active conformation, thereby restoring the ability of the variant to catalyze conversion of prothrombin to thrombin and thus its pro-coagulant activity. In mouse models of hemophilia A and hemophilia B, administering the FXa I16L variant before injury reduced blood loss following tail-clipping in a dose dependent manner. The results of these experiments suggest that the FXa I16L variant might be useful to treat uncontrolled bleeding in humans with hemophilia.

[0007] The FXa I16L variant used in the earlier studies, however, was made in small quantities from stably transfected HEK 293 cells followed by activation of the FX protein using Russell's viper venom protease RVVX. Toso, R., Zhu, H. & Camire, R.M. The conformational switch from the factor X zymogen to protease state mediates exosite expression and prothrombinase assembly. *J. Biol. Chem.* 283, 18627–18635 (2008). While useful for small-scale studies, this approach is not suitable for production of large quantities of purified FXa variant protein required for clinical studies and eventual supply to patients. Accordingly, there is a need in the art for preparations of the FXa I16L variant protein made in such quantity and purity that they may be tested in clinical trials, and once approved, provided to subjects in need of hemostasis.

SUMMARY OF THE INVENTION

[0008] The present disclosure addresses the unmet need in the art described above by providing compositions of FXa variant proteins produced in sufficient purity and quantity that they can be supplied to subjects in need of hemostasis. In various embodiments, these compositions comprise different isoforms and post-translational modifications of FXa variant proteins.

[0009] In one embodiment, compositions comprise the beta form of FXa variant protein in which the light and heavy chain protein sequences respectively consist of amino acids 1 to 139 and 146 to 386, amino acids 1 to 140 and 146 to 384, amino acids 1 to 140 and 146 to 386, amino acids 1 to 141 and 146 to 384, amino acids 1 to 141 and 146 to 386, amino acids 1 to 142 and 146 to 384, amino acids 1 to 142 and 146 to 386, amino acids 1 to 143 and 146

to 384, and amino acids 1 to 143 and 146 to 386, all of the amino acid sequence of SEQ ID NO:1.

[00010] In another embodiment, compositions comprise the alpha form of FXa variant protein in which the light and heavy chain protein sequences respectively consist of amino acids 1 to 139 and 146 to 398, amino acids 1 to 140 and 146 to 398, amino acids 1 to 140 and 146 to 399, amino acids 1 to 141 and 146 to 398, amino acids 1 to 141 and 146 to 399, amino acids 1 to 142 and 146 to 398, and amino acids 1 to 142 and 146 to 399, all of the amino acid sequence of SEQ ID NO:1.

[00011] In any one or more of the beta and alpha isoform embodiments of the FXa variant proteins described above, the light chains of the proteins can be modified to include β -hydroxy Asp⁶³, an O-linked hexose and 9, 10 or 11 Gla residues. In some embodiments, 9 Gla residues are present. In other embodiments, 10 Gla residues are present. And in yet other embodiments, 11 Gla residues are present.

[00012] In any one or more of the beta isoform embodiments of the FXa variant proteins described above, the heavy chains can be modified to include one or two core-1 O-linked glycans. In some embodiments, only the first, only the second, or both core-1 O-linked glycans can be non-sialylated, mono-sialylated or di-sialylated. According to some embodiments, no sialic acid groups are present. In other embodiments, one sialic acid group is present. In yet other embodiments, two sialic acid groups are present. In further embodiments, three sialic acid groups are present. And in yet further embodiments, a total of four sialic acid groups are present.

[00013] According to some embodiments, FXa variant protein compositions include at least one FXa variant protein species listed in Table 2 of the disclosure. In other embodiments, compositions of the disclosure comprise at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, or at least 60 of the FXa variant protein species listed in Table 2.

[00014] According to other embodiments, FXa variant protein compositions include at least one FXa variant protein in which the light chain is a species listed in Table 3 of the disclosure. And in yet other embodiments, FXa variant protein compositions include at least one FXa variant protein in which the heavy chain is a species listed in Table 4 of the disclosure.

[00015] In some embodiments, any one or more of the FXa variant proteins of the disclosure can be present in a composition at an average abundance that varies relative to other species that may be present. For example, a particular species may be present at major,

minor or trace abundance compared to other species that are present. Alternatively, a particular species may be present at high abundance, medium abundance, low abundance, or very low abundance compared to other species that are present.

[00016] The present disclosure also provides nucleic acids, vectors, host cells and methods of making and purifying FXa variant proteins.

[00017] Included, for example, are nucleic acid sequences encoding FX variant protein including a replacement of the native Activation Peptide (AP) sequence with a PACE processing site, thereby permitting intracellular activation of the clotting factor. In some embodiments, the AP sequence is removed entirely and replaced with the amino acid sequence Arg-Lys-Arg. According to some embodiments, a cDNA sequence encoding FX variant protein is provided by the nucleic acid sequence of SEQ ID NO:4, and the amino acid sequence encoded thereby is provided by the amino acid sequence of SEQ ID NO:1. Related methods are provided for making FXa variant protein by growing host cells co-expressing PACE (or a similar protease) and FX variant protein, and then purifying the FXa variant protein produced, processed and secreted by the host cells. Disclosed as well are FXa variant proteins produced according to these methods. In some embodiments of these methods, host cells are CHO cells.

[00018] Purification of FXa variant proteins can be performed by chromatography, including passing a solution comprising the proteins through a mixed mode chromatography (MMC) column (e.g., using Capto MMC media), followed by washing and eluting, then passing the proteins through a first ion exchange chromatography column (e.g., using Q-Sepharose Fast Flow media), followed by washing and eluting, and then passing the proteins through a second different ion exchange chromatography column (e.g., using Fractogel SO₃⁻ media), followed by washing and eluting. Other methods for purifying FXa variant proteins are also possible.

[00019] Purification methods can optionally include a step of viral inactivation, as well as the steps of ultrafiltration and diafiltration. After FXa variant proteins are purified, and in some instances concentrated, they can be diluted in a pharmaceutically acceptable diluent optionally containing other ingredients such as buffers or excipients.

[00020] The disclosure also provides methods of treating subjects in need of hemostasis by administering a hemostatically effective amount of a composition comprising one or more of the FXa variant proteins of the disclosure. In other embodiments, subjects are administered a hemostatically effective amount of a composition comprising one or more of the FXa variant proteins of the disclosure before bleeding occurs to prophylactically prevent uncontrolled

bleeding in a susceptible subject. In some embodiments, the subject is treated for Hemophilia A. In other embodiments, the subject is treated for Hemophilia B. And in yet other embodiments, the subject is treated for trauma or other types of uncontrolled bleeding.

BRIEF DESCRIPTION OF THE DRAWINGS

[00021] FIG. 1A provides the amino acid sequence (SEQ ID NO:1) of mature FX variant protein in which the wild-type isoleucine residue at position 146 is substituted with the leucine shown in bold. Position 146 corresponds to position 16 in the chymotrypsin numbering system. Potential γ -carboxyglutamic acid residues (Gla) are underlined and in italics. Predicted intrachain and interchain disulfide bonds are illustrated by lines drawn between the linked cysteines. The predicted β -hydroxylation site is enclosed by a box with solid lines. The PACE recognition and cleavage site engineered to replace the Activation Peptide is enclosed by a box with broken lines. The lysine forming the carboxy-terminal amino acid of the beta form of the heavy chain (Lys³⁸⁶) is enclosed by an oval with solid line. FIG. 1B provides the predicted amino acid sequence (SEQ ID NO:2) of mature FX variant protein light chain. FIG. 1C provides the predicted amino acid sequence (SEQ ID NO:3) of the FXa variant protein heavy chain. FIG. 1D provides the nucleotide sequence (SEQ ID NO:4) of a cDNA encoding FX variant protein, including signal sequence and propeptide.

[00022] FIG. 2 provides the mass spectrum of peptides resulting from the reduction, alkylation, and proteolytic digestion with Lys-C of a purified preparation of intact FXa variant protein. Peak assignments based on the mass spectrum after RP-HPLC/ESI-QTOF MS analysis of peptides were used to confirm the amino acid sequence of the protein and to identify certain post-translational modifications described in Table 1 and elsewhere herein. Peak labeling is as follows. “L” followed by a number refers to a particular peptide derived from the light chain. “H” followed by a number refers to a particular peptide derived from the heavy chain. “H20¹”: C-terminal alpha-form of H chain ending in Lys³⁹⁹ containing two di-sialylated core-1 O-glycans. “H20²”: C-terminal alpha-form of H chain ending in Lys³⁹⁹ containing one di-sialylated core-1 O-glycan. “H20³”: C-terminal alpha-form of H chain ending in Leu³⁹⁸ containing one di-sialylated core-1 O-glycan. “L4^{1Gla}”: L chain peptide #4 containing 1 γ -carboxyglutamic acid residue. “L10^{glc}”: L chain peptide #10 containing potential O-linked glucose (hexose). “L1^{2Gla}”: L chain peptide #1 with 2 γ -carboxyglutamic acid residues. “L3^{6-8Gla}”: L chain peptide #3 with 6, 7 or 8 γ -carboxyglutamic acid residues. “L8^{b-OH}”: L chain peptide #8 containing β -hydroxy Asp. “d”: deamidation. “R”: reagent peak, i.e., buffer related products consistent with reagent blank. “*”: overalkylation.

[00023] FIG. 3 provides an HPLC chromatogram of a purified preparation of intact FXa variant protein and demonstrates the existence in the preparation of different FXa variant protein isoforms.

[00024] FIG. 4 provides an HPLC chromatogram of a purified preparation of intact FXa variant protein and demonstrates the existence in the preparation of different FXa variant protein isoforms. The preparation was also analyzed by mass spectrometry.

[00025] FIG. 5 provides the mass spectrum of intact FXa variant protein from a purified preparation. The mass spectrum confirms the existence of two major isoforms, alpha and beta, as well as extensive mass heterogeneity attributable to variable digestion at the PACE cleavage site and the presence of different types of post-translational modification. Major peaks are identified in the figure by their observed mass in daltons. Peak assignments based on the mass spectrum after RP-HPLC/ESI-QTOF MS analysis were used to confirm the amino acid sequence of protein isoforms and to identify certain post-translational modifications as described in Table 2 and elsewhere herein. Peaks labeled with “*” indicate the presence of 11 Gla residues in the light chain Gla domain.

[00026] FIG. 6 provides an HPLC chromatogram of a purified preparation of FXa variant protein after reduction and alkylation to separate the light and heavy chains by eliminating interchain and intrachain disulfide bonds. The chromatogram demonstrates the existence of different FXa variant protein isoforms. The preparation was also analyzed by mass spectrometry.

[00027] FIG. 7A provides the mass spectrum of FXa variant protein light chain after reduction and alkylation of a purified preparation of the protein. The mass spectrum demonstrates extensive mass heterogeneity attributable to variable digestion at the PACE cleavage site and presence of different types of post-translational modifications. Major peaks are identified in the figure by their observed mass in daltons. Peak assignments based on the mass spectrum after RP-HPLC/ESI-QTOF MS analysis were used to confirm the amino acid sequence of protein isoforms and to identify certain post-translational modifications as described in Table 3 and elsewhere herein.

[00028] FIG. 7B provides the mass spectrum of FXa variant protein heavy chain after reduction and alkylation of a purified preparation of the protein. The mass spectrum demonstrates extensive mass heterogeneity attributable to presence of different types of post-translational modifications. Major peaks are identified in the figure by their observed mass in daltons. Peak assignments based on the mass spectrum after RP-HPLC/ESI-QTOF MS analysis were used to confirm the amino acid sequence of protein isoforms and to identify

certain post-translational modifications as described in Table 4 and elsewhere herein. Peaks labeled with “*” indicates overalkylation of the heavy chain.

DETAILED DESCRIPTION

[00029] Described herein are compositions comprising FXa variant protein produced in sufficient purity and quantity to be tested in clinical trials and provided to subjects in need of hemostasis. Also described are methods of making and purifying FXa variant protein.

[00030] In certain embodiments, the phrase “FX variant protein,” “FXa variant protein,” and similar terms refer (unless otherwise clear from context) to human Factor X (FX) or activated Factor X (FXa), respectively, in which the isoleucine (Ile or I) immediately following the Activation Peptide sequence in the heavy chain is substituted with leucine (Leu or L). Based on sequence alignment between the FXa heavy chain and the catalytic domain of chymotrypsin, this substitution is also referred to as the I16L substitution mutation using the chymotrypsin numbering scheme. This position corresponds to amino acid 146 in SEQ ID NO:1 and amino acid 1 in SEQ ID NO:3. As discussed above, FXa including this mutation has zymogenic properties that allow it to circulate in blood for longer periods of time compared to wild-type FXa, but is also capable of cleaving prothrombin at high rate when incorporated into the prothrombinase complex.

[00031] According to other embodiments, FX variant protein and FXa variant protein refer to human FX or FXa, respectively, in which the amino acid corresponding to position 146 in SEQ ID NO:1 (position 16 in chymotrypsin numbering scheme) is substituted with Phe (F), Asp (D), or Gly (G). According to yet other embodiments, FX variant protein and FXa variant protein refer to human FX or FXa, respectively, in which the valine (Val or V) corresponding to position 147 in SEQ ID NO:1 (position 17 in chymotrypsin numbering scheme) is substituted with Leu (L), Ala (A), or Gly (G). And, in other embodiments, FX variant protein and FXa variant protein refer to human FX or FXa, respectively, in which the amino acid corresponding to position 329 in SEQ ID NO:1 (position 194 in chymotrypsin numbering scheme) is substituted with Asn (N) or Glu (E).

[00032] Wild-type Factor X normally circulates in the blood as an inactive two-chain zymogen held together by a disulphide bond. The two-chain zymogen is formed from mature FX (i.e., lacking signal peptide and propeptide) after proteolytic removal of the peptide Arg-Lys-Arg present in the mature protein as amino acids 140-142. Activation, however, requires removal of the Activation Peptide (AP) by Factor IXa or Factor VIIa preliminary to clot formation, or by other proteases such as RVVX. The AP corresponds to amino acids 143-

194 in mature single-chain FX. In the two-chain zymogen, it is present at the amino-terminal end of the heavy chain. M. Hertzberg, Biochemistry of factor X, Blood Rev. 8(1):56-62 (1994).

[00033] In an exemplary embodiment, Factor X variant protein sequence was modified by removing the Activation Peptide and replacing it with the amino acid sequence Arg-Lys-Arg (RKR in single letter code) to create a recognition and cleavage site for paired basic amino acid cleaving enzyme (PACE; also called furin). Rather than activating FXa in a separate step, AP replacement permits intracellular activation of FX variant protein when PACE is co-expressed in host cells. This avoids the need to first purify FX variant produced by cells, separately activate the protein using a protease (such as RVVX) and then purify variant FXa. By avoiding these extra steps, FXa variant can be produced in purer form and in greater quantities.

[00034] The mature Factor X variant amino acid sequence is depicted in FIG. 1A (SEQ ID NO:1) in which the I16L leucine substitution appears at amino acid position 146 (bold), the RKR sequence replacing the Activation Peptide appears at positions 143-145 and the PACE processing site appears at positions 140-145 (box with broken line). FIG. 1B and FIG. 1C depict the predicted amino acid sequence of the FXa variant light chain (SEQ ID NO:2) and heavy chain (SEQ ID NO:3), respectively, after cleavage at the PACE site.

[00035] Analysis of multiple preparations of FXa variant protein produced according to the methods of the disclosure demonstrated an unexpected degree of heterogeneity with respect protein sequence and post-translational modification compared to the predicted structure. Nevertheless, such compositions comprising structurally heterogeneous populations of FXa variant proteins were capable of acting as potent pro-coagulants in an *in vitro* assay. As described further below, the heterogeneity was attributable to variations in light chain and heavy chain amino acid sequence and post-translational modifications present in one, the other, or both chains.

[00036] Without wishing to be bound by any particular theory of operation, it is believed that one source of structural heterogeneity is attributable to variable cleavage at the PACE/furin protease recognition and cleavage site engineered to replace the Activation Peptide. In a non-limiting embodiment, the amino acid sequence of the PACE site is RKRRKR (in single letter code) (SEQ ID NO:5) engineered into the FX variant protein by replacing the native AP amino acid sequence with the residues RKR which, in concert with the last three residues of the light chain, forms the proteolytic recognition site RKRRKR

(SEQ ID NO:5). Other PACE recognition sites can be used as well (e.g., RKR alone), as could recognition and cleavage sites for proteases other than PACE/furin.

[00037] In some embodiments, the PACE site is proteolytically cleaved to remove it entirely from the light chain leaving no residues behind (ANS...TLER¹³⁹/RKRRKR/) (SEQ ID NO:6 and SEQ ID NO:5, respectively the light chain and a peptide fragment). In the foregoing and following embodiments described in this paragraph, slash marks (“/”) indicate possible cleavage sites between the adjacent amino acids and the superscripted numeral indicates the position of the carboxy-terminal amino acid of the light chain after cleavage (based on FIG. 1A; SEQ ID NO:1). In other embodiments, the PACE site is proteolytically cleaved leaving its first residue at the C-terminus of the light chain (ANS...TLERR¹⁴⁰/KRRKR/) (SEQ ID NO:7 and SEQ ID NO:8, respectively the light chain and a peptide fragment). In other embodiments, the PACE site is proteolytically cleaved leaving its first two residues at the C-terminus of the light chain (ANS...TLERRK¹⁴¹/RRKR/) (SEQ ID NO:9 and SEQ ID NO:10, respectively the light chain and a peptide fragment). In other embodiments, the PACE site is proteolytically cleaved leaving its first three residues at the C-terminus of the light chain (ANS...TLERRKR¹⁴²/RKR/) (SEQ ID NO:11). In other embodiments, the PACE site is proteolytically cleaved leaving its first four residues at the C-terminus of the light chain (ANS...TLERRKRR¹⁴³/KR/) (SEQ ID NO:12). In other embodiments, the PACE site is proteolytically cleaved leaving its first five residues at the C-terminus of the light chain (ANS...TLERRKRRK¹⁴⁴/R/) (SEQ ID NO:13). And in yet other embodiments, the PACE site is proteolytically cleaved leaving its entire sequence at the C-terminus of the light chain (ANS...TLERRKRRKR¹⁴⁵/) (SEQ ID NO:2).

[00038] In other embodiments, structural heterogeneity in the light chain can also include presence of varying number of Gla residues in the Gla domain attributable to variations in the degree of γ -carboxylation of certain glutamic acid residues. In certain embodiments, the number of Gla residues is 9, 10, or 11. Potential sites of glutamic acid γ -carboxylation in the Gla domain are identified in FIG. 1A. According to other embodiments, additional potential sources of light chain structural heterogeneity include presence or absence of a β -hydroxylated aspartic acid residue at Asp⁶³ (FIG. 1A; SEQ ID NO:1) and presence or absence of an O-linked hexose. In some embodiments, the O-linked hexose is glucose, but in other embodiments it may be a different aldohexose, or a cyclic hemiacetal, ketohexose, or other O-linked hexose.

[00039] The heavy chain can also exhibit structural heterogeneity. According to certain embodiments variation can occur in the length of the heavy chain. For example, some heavy

chains are the longer alpha isoform terminating at K³⁹⁹ (SEQ ID NO:3) (the superscripted numeral indicating the position of the carboxy-terminal amino acid of the heavy chain based on FIG. 1A; SEQ ID NO:1), but in other instances the terminal lysine is not present so that the protein ends at L³⁹⁸ (SEQ ID NO:14). In other embodiments, the heavy chain is the shorter beta form terminating at K³⁸⁶ or K³⁸⁴ (SEQ ID NO:46 and SEQ ID NO:45, respectively).

[00040] Yet another source of heterogeneity in the heavy chains is the degree of O-linked glycosylation. Thus in some embodiments, the heavy chains are unglycosylated. In other embodiments, heavy chains include one or two core-1 O-glycans. A core-1 O-glycan is an O-glycan where N-acetylgalactosamine (GalNAc) is attached to serine or threonine and galactose (Gal) is attached to GalNAc. In some embodiments of the disclosure the GalNAc sugar is present but the Gal sugar is missing. According to yet other embodiments, each of the two core-1 O-linked glycans can separately be unsialylated, mono-sialylated or di-sialylated, such that heavy chains can possess 0-4 sialic acid groups in total. When sialic acid groups are present, they can be attached to either GalNAc or to Gal, or to both sugars.

[00041] According to other embodiments, different pairings of light chains and heavy chains are possible, thereby yielding a large combinatorial set of two-chain zymogens with different chain lengths and post-translational modifications.

[00042] In other embodiments, the different species of FXa variant proteins can vary in their relative abundance. In one embodiment, abundance of the various species can be detected by comparing peak heights in the mass spectra of intact FXa variant proteins. Alternatively, abundance of light and heavy chains can be separately analyzed by mass spectrometry after intact protein is reduced and alkylated to eliminate interchain and intrachain disulfide bonds. As described in the Examples, below, abundance of different FXa variant species can be scored as major, minor or trace by comparison to the highest peak in mass spectra.

[00043] Mass peak height comparison, however, is not the only method of measuring abundance. Other methods for measuring absolute or relative abundance of structural species is within the knowledge of those of ordinary skill in the art. In a non-limiting example of another method for categorizing relative abundance, a species is considered to be in high abundance if its mass spectrum peak height is at least about 50% that of the most abundant species (i.e., highest peak in a mass spectrum). A species is of medium abundance if its peak is between about 10-50% that of the most abundant species. A species is of low abundance if its peak is between about 2-10% that of the most abundant species, and is of very low

abundance if its peak is less than about 2% that of the most abundant species. Other conventions for comparing relative species abundance are also possible.

[00044] Specific, non-limiting embodiments of structural heterogeneity in FXa variant protein light chains, heavy chains and combinations thereof are described below.

[00045] Embodiments of the disclosure include FXa variant proteins in which the carboxy-terminus of the light chain ends at different amino acids due to variable cleavage at the PACE site. Related embodiments include presence of varying numbers of Gla residues and post-translational modifications such as β -hydroxylation of an aspartic acid residue and an O-linked hexose. Some of these embodiments are described below.

[00046] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 139 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00047] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 140 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00048] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 141 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00049] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 142 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00050] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 143 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00051] Embodiments of the disclosure include FXa variant proteins in which the carboxy-terminus of the heavy chain ends at different amino acids. Related embodiments include varying numbers of O-linked glycans including varying degrees of sialylation. Some of these embodiments are described below.

[00052] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. Compositions of the disclosure may further comprise any of the other FXa variant proteins of the disclosure.

[00053] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. Compositions of the disclosure may further comprise any of the other FXa variant proteins of the disclosure.

[00054] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. Compositions of the disclosure may comprise any of the FXa variant proteins

described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00055] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the heavy chain consists of amino acids 146 to 399 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00056] Embodiments of the disclosure include FXa variant proteins in which the carboxy-terminus of the light chain ends at different amino acids due to variable cleavage at the PACE site and in which the carboxy-terminus of the heavy chain also ends at different amino acids. Related embodiments include varying numbers of Gla residues and post-translational modifications, such as β -hydroxylation of an aspartic acid residue and an O-linked hexose in the light chains, and varying numbers of O-linked glycans including varying degrees of sialylation in the heavy chains. Some of these embodiments are described below.

[00057] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 139 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00058] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 139 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa

variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00059] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 139 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00060] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 139 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 399 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00061] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 140 of the amino acid sequence

of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00062] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 140 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00063] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 140 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00064] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 140 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 399 of the amino acid

sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00065] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 141 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00066] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 141 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00067] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 141 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9

Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00068] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 141 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 399 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00069] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 142 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa

variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00070] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 142 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00071] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 142 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00072] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 142 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 399 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated,

or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00073] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 143 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 384 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00074] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 143 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 386 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00075] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 143 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 398 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated,

or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00076] In certain embodiments, compositions of the disclosure comprise a FXa variant protein in which the light chain consists of amino acids 1 to 143 of the amino acid sequence of SEQ ID NO:1 and the heavy chain consists of amino acids 146 to 399 of the amino acid sequence of SEQ ID NO:1. In related embodiments, the light chain additionally comprises 9 Gla residues, 10 Gla residues, or 11 Gla residues. In each of these embodiments, the light chain may further comprise β -hydroxy Asp⁶³, an O-linked hexose, or both these post-translational modifications. In other related embodiments, the heavy chain additionally comprises one core-1 glycan, which may be non-sialylated, mono-sialylated, or di-sialylated, or two core-1 glycans, each of which may independently be non-sialylated, mono-sialylated, or di-sialylated. In yet other related embodiments, each possible permutation of post-translational modifications of the light chain is combined with each possible permutation of post-translational modifications of the heavy chain. Compositions of the disclosure may comprise any of the FXa variant proteins described in this paragraph alone or in combination with the others so described, as well as with any of the other FXa variant proteins of the disclosure.

[00077] The present disclosure further provides isolated nucleic acids comprising nucleic acid sequences that encode FX variant protein. According to an exemplary, non-limiting embodiment, a complementary DNA sequence (cDNA) encoding FX variant protein is disclosed herein as SEQ ID NO:4. As will be appreciated by those of ordinary skill, many other nucleic acid sequences encoding FX variant protein are possible in light of the degeneracy of the genetic code.

[00078] In some embodiments, nucleic acid encoding FX variant protein can be provided with a sequence encoding a signal peptide and/or propeptide (i.e., leader sequences) positioned at the amino-terminus of the FX protein which, among other potential functions, directs newly synthesized protein to the secretory compartment. Post-translational processing then removes the leader before mature protein is secreted from the cell. In some embodiments, these sequences are derived from native human FX. Non-native leader sequences can be used, such as that from human prothrombin. Leader sequences from other

proteins (including of non-human origin) can be used as well. In the embodiment of the nucleic acid sequence of SEQ ID NO:4, the amino-terminal leader sequence is from human prothrombin.

[00079] According to other embodiments, nucleic acid encoding FX variant protein can be modified to permit intracellular activation of the clotting factor to yield FXa variant proteins.

[00080] As noted above, FX normally circulates as a two-chain zymogen in which a 52 amino acid Activation Peptide is positioned at the amino-terminus of the FX heavy chain. Activation to form FXa capable of functioning as procoagulant in the prothrombinase complex requires proteolytic removal of the AP by proteases from the intrinsic or extrinsic clotting system, such as activated FIX or activated FVII. In prior studies, analogous methods of activation were used. Specifically, FX variant protein was expressed in cells and then activated in a separate step by treatment with Russell's Viper venom (RVVX) after which FXa variant protein was purified for subsequent analysis.

[00081] Although activated FX variant protein could in principle be produced at industrial scale using similar methods, doing so would be very inefficient and expensive. For example, rigorous purification to eliminate any residual RVVX would be required. To avoid these disadvantages, the native AP can be removed and replaced with recognition and cleavage sites of proteases capable of being expressed and functioning intracellularly. In this manner, it is possible for FX variant protein to be activated intracellularly without need for additional process steps. Activated FX variant proteins secreted by host cells into the growth medium can then be purified.

[00082] A non-limiting example of a nucleic acid sequence encoding FX variant in which the AP sequence is replaced with a PACE recognition and cleavage site is provided in SEQ ID NO:4. The protein sequence of the mature FX variant protein encoded by this exemplary cDNA sequence is provided in SEQ ID NO:1. In these embodiments, the amino acids constituting the AP (amino acids 143 to 194 of the wild-type FX protein sequence) are replaced with the sequence Arg-Lys-Arg (RKR in single letter code). This sequence, combined with the last three residues of the light chain sequence (also RKR; amino acids 140-142 of SEQ ID NO:1) creates a recognition and cleavage site (i.e., RKRRKR) (SEQ ID NO:5) for paired basic amino acid cleaving enzyme (PACE). Soluble PACE enzyme co-expressed in the same host cells expressing the modified FX variant protein can therefore activate the clotting factor intracellularly.

[00083] In some embodiments, where host cells natively express an enzyme with a similar recognition site as PACE, it may not be necessary to co-express an exogenous protease.

Rather, the native protease may be sufficient to activate FX variant protein expressed in the cells, thereby producing and secreting activated FX variant protein.

[00084] In other embodiments, recognition and cleavage sites for proteases different than PACE, but also capable of functioning intracellularly, can be used to replace the AP amino acid sequence. Such enzymes can be co-expressed in the same host cells as FX variant protein to activate the modified clotting factor. If native enzymes capable of cleaving the engineered site are produced at sufficient levels in the host cells, co-expression of exogenous proteases may not be required.

[00085] Nucleic acid sequences encoding FX variant proteins of the present disclosure may be incorporated into vectors using techniques well known to those of ordinary skill in the art.

[00086] Vectors, in certain embodiments, include plasmids generally, bacterial plasmids, eukaryotic episomes, yeast artificial chromosomes and viral genomes. Exemplary non-limiting viruses include retroviruses, adenoviruses, adeno-associated viruses (AAV), and plant viruses such as cauliflower mosaic virus, and tobacco mosaic virus. Other types of vectors are possible. In some embodiments, vectors are capable of autonomous replication in suitable hosts. In other embodiments, vectors are maintained in hosts extrachromosomally or can become integrated into the host's genome allowing the vector to replicate with the host's genome. Vectors comprising a gene and control sequences sufficient to maintain transcription and translation of the gene are called expression vectors. Vectors according to the present disclosure may be selected or designed, according to the knowledge of those ordinarily skilled in the art, to function in any cell type capable of supporting expression of FX variant protein, including bacterial cells, other prokaryotic cells, yeast cells, other fungal cells, plant cells, animal cells, insect cells, mammalian cells, CHO cells, and human cells, or others.

[00087] Vectors may optionally contain one or more control sequences. Certain control sequences permit replication, such as origins of replication. Other control sequences control or modulate transcription, such as promoters, enhancers, and transcription termination sites. Non-limiting examples of promoter or enhancers are those derived from retroviral LTRs, cytomegalovirus (CMV), Simian Virus 40 (SV40), adenovirus (e.g., the adenovirus major late promoter (AdMLP)), or polyoma virus. Additional examples include tissue specific promoters and enhancers, constitutively active promoters and enhancers, and inducible promoters and enhancers. Other promoters and enhancers are also possible. Yet other control sequences control or modulate post-transcriptional RNA processing, such as splicing

and polyadenylation signals, or signals that increase or decrease mRNA stability. Other control sequences control or modulate protein translation, such as translation initiation sequences (e.g., Kozak consensus sequence), post-translational processing, or protein stability.

[00088] Vectors can also include selectable marker genes, permitting the selection of host cells that have taken up the vectors. Non-limiting examples include selectable marker genes that confer a drug-resistant phenotype, such as the dihydrofolate reductase gene (DHFR) (for use in DHFR⁻ host cells permitting selection using methotrexate), the neo gene (permitting selection with G418 or similar drugs), the hph gene (permitting selection with hygromycin B), and the glutamate synthetase gene (permitting selection with methionine sulfoximine).

[00089] Vectors comprising nucleic acid sequences encoding the FX variant proteins of the disclosure may be introduced into one or more types of host cells capable of supporting FX variant protein expression. Methods for introducing vectors into suitable host cells are well known to those of ordinary skill in the art. Non-limiting examples include transient and stable transfection, transformation, transduction and viral infection of target host cells. Other examples include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. Methods for transforming plant cells, bacterial and yeast cells are also well known in the art.

[00090] FX or FXa variant proteins can be expressed with other proteins to optimize expression or post-translational processing of FX or FXa variant proteins. In a non-limiting example, a vector including a gene encoding γ -glutamyl carboxylase can be co-transfected with the vector encoding FX variant protein into host cells to increase carboxylation of glutamic acids in the light chain Gla domain. By so doing, the extent of carboxylation and formation of Gla residues in the Gla domain can be increased, thereby improving the yield of active clotting factor produced by the cells. In another embodiment described above, FX variant protein modified to replace the native AP amino acid sequence with a PACE cleavage site can be co-expressed with soluble PACE enzyme in the same host cells. The co-expressed PACE can then cleave FX variant protein intracellularly to form activated FX variant protein which is thereafter secreted from the cell. As noted above, this approach avoids the need to activate FX variant protein in a separate step, which would otherwise be very inefficient and expensive.

[00091] Cells capable of expressing FX variant proteins, and other proteins useful for optimizing expression of FX variant protein in an active form, include mammalian cells.

Mammalian cell lines suitable as hosts for protein expression are known in the art, including those derived from humans, rats, mice and other mammals. Exemplary non-limiting examples include certain immortalized cell lines available from the American Type Culture Collection (ATCC) or other sources, including Chinese hamster ovary (CHO) cells, NS0 cells, SP2 cells, HEK-293T cells, NIH-3T3 cells, HeLa cells, baby hamster kidney (BHK) cells, African green monkey kidney cells (e.g., COS, CV-1 or Vero cells), human hepatocellular carcinoma cells (e.g., HepG2), A549 cells, A431 cells, HeLa cells, L cells, BHK21 cells, HL-60 cells, U937 cells, HaK cells, Jurkat cells, and others. Use of other mammalian cells as host cells for protein expression is possible according to the knowledge of those ordinarily skilled.

[00092] In other embodiments, cell lines from insects, plants, bacteria or fungi may be used. Exemplary non-limiting insect cells include Sf9 or Sf21 cells, which are often used in conjunction with the baculovirus vector expression system. Exemplary non-limiting plant cells include those from nicotiana, arabidopsis, duckweed, corn, wheat, and potato species. Exemplary non-limiting bacteria include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and *Streptomyces* strains. Exemplary non-limiting fungi include *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae*, *Pichia pastoris*, *Kluyveromyces* yeast strains, and *Candida* yeast strains. Use of other insect, plant, bacterial and fungal cells as host cells for protein expression is possible according to the knowledge of those ordinarily skilled in the art.

[00093] Methods, reagents and conditions for culturing host cells to produce activated FX variant protein are within the knowledge of those ordinarily skilled in the art and are not intended to be limiting. In a non-limiting example, FXa variant proteins can be produced in 500 liter cultures of serum free medium supplemented with vitamin K. In another non-limiting example, FXa variant proteins of the disclosure can be produced at industrial scale, for example, in 2500 liter fermentation tanks, or even larger volumes, under conditions designed to optimize expression and recovery.

[00094] After growing host cells under culture conditions supporting expression and intracellular activation of FX variant proteins, the cellular growth medium can be processed to isolate and purify FXa variant proteins secreted during cell growth. In other embodiments, particularly when other than mammalian host cells are used, FXa variant protein remaining in cells can be released, for example, by breaking open cells mechanically, enzymatically, or with detergents.

[00095] Cell culture medium can be centrifuged and/or filtered to remove cells and debris, for example by depth filtration alone or followed by membrane filtration, for example through a membrane filter with average pore size of 0.45 μm , 0.2 μm , or some other pore size. After removing cellular debris by centrifugation and/or filtration, medium can be processed to further purify FXa variant proteins. Depending on the subsequent purification steps, dilution and/or buffer exchange of the clarified supernatant or filtrate may be desirable.

[00096] Methods for protein purification are well known in the art. Exemplary non-limiting methods of protein purification include salt precipitation, size exclusion chromatography, ion exchange chromatography, and affinity chromatography. For example, antibodies specifically recognizing FXa variant protein can be immobilized to purification columns to reversibly capture the protein from the surrounding medium or buffer. After washing the affinity column to remove contaminants, bound FXa variant protein can thereafter be eluted.

[00097] Partially purified FXa variant protein can be subjected to processing steps in accordance with good manufacturing practice or other regulatory requirements including, for example, removal or inactivation of viruses. Viruses can be removed by ultrafiltration or inactivated by treating a sample with alcohol and/or detergent, according to the knowledge of those ordinarily skilled in the art. For example, Virosart® brand virus filters, available from Sartorius Stedim Biotech, can be used, as can filters from other manufacturers. Other methods for removing or inactivating viruses are possible.

[00098] In some embodiments, mixed mode chromatography (MMC) can be used to purify FXa variant protein alone, or in combination with other purification steps. MMC, also called multimodal chromatography, refers to use of chromatography media that provides more than one type of interaction between ligand and sample components. Exemplary types of interactions include electrostatic (anionic or cationic exchanger), hydrophobic, π - π interaction, hydrogen bonding and thiophilic interaction. These interactions can cooperate or work independently. Specific, non-limiting examples of MMC media include Capto adhere, Capto adhere ImpRes, Capto MMC, Capto MMC ImpRes, Capto Core 700, all of which are described further in the Multimodal Chromatography Handbook published by GE Healthcare Life Sciences (2013). Capto MMC, for example, provides thiophilic, hydrophobic, hydrogen bonding, as well as cationic exchanger electrostatic properties. Use of MMC media from other sources is possible as well.

[00099] In some embodiments, ion exchange chromatography can be used to purify FXa variant protein alone, or in combination with other purification steps. Ion exchange

chromatography refers to use of chromatography media allowing separation of molecules on the basis of differences in their net surface charge, which can vary with pH, salt and other conditions. At a pH above a protein's isoelectric point, the protein will bind to a positively charged medium (anion exchanger), whereas at a pH below its isoelectric point, a protein will bind to a negatively charged medium (cation exchanger). Ion exchange media and buffer conditions (e.g., salt concentration, pH, etc.) can be chosen so that a protein to be purified (e.g., a FXa variant protein) binds preferentially to the media, permitting contaminants that bind less strongly under the same conditions to be washed away, after which the protein of interest can be eluted from the chromatography column. Additional information about ion exchange chromatography is described further in the Ion Exchange Chromatography & Chromatofocusing Principles and Methods Handbook published by GE Healthcare (2010).

[000100] Ion exchangers can be categorized as strong or weak, which indicates the degree to which they remain charged as pH changes. Strong exchangers remain charged across a wide pH range, whereas weak exchangers do not. Examples of anion exchanger functional groups include quaternary ammonium (Q), which is strong, and diethylaminoethyl (DEAE) and diethylaminopropyl (ANX), which are considered weak. Examples of cation exchanger functional groups include sulfopropyl (SP) and methyl sulfonate (S), both of which are considered strong, and carboxymethyl (CM), which is weak.

[000101] In one embodiment, FXa variant protein can be purified using an anion exchange chromatography step alone, or followed by a cation exchange chromatography step. In another embodiment, FXa variant protein can be purified using a cation exchange chromatography step alone, or followed by an anion exchange chromatography step.

[000102] Specific, non-limiting examples of ion exchange media include Capto DEAE, Capto Q ImpRes, Capto SP ImpRes, Capto S, Capto Q, SOURCE 15Q, SOURCE 15S, SOURCE 30Q, SOURCE 30S, MacroCap SP, MacroCap Q, Mini Q PC, Mini S PC, Mini Q PE, Mini S PE, HR Mono Q, HR Mono S, PC Mono Q, PC Mono S, Mono Q GL, Mono S GL, DEAE Sephadex, CM Sephadex C-25, CM Sephadex C-50, DEAE Sephadex A-25, DEAE Sephadex A-50, QAE Sephadex A-25, QAE Sephadex A-50, SP Sephadex C-25, SP Sephadex C-50, ANX Sepharose 4 Fast Flow (High Sub), ANX Sepharose 4 Fast Flow (Low Sub), CM Sepharose Fast Flow, CM Sepharose High Performance, DEAE Sepharose CL-6B, DEAE Sepharose Fast Flow, Q Sepharose Big Beads, Q Sepharose Fast Flow, Q Sepharose High Performance, Q Sepharose XL, SP Sepharose Big Beads, SP Sepharose Big Beads Food Grade, SP Sepharose High Performance, SP Sepharose Fast Flow, and SP Sepharose XL, which are available from GE Healthcare. Other ion exchange media can be obtained from

Merck Millipore including, for example, those in the Fractogel® series or the Eshmuno® series. Examples in the Fractogel® series include TMAE Resin, TMAE Hicap Resin, TMAE Medcap Resin, DEAE Resin, DMAE Resin, SO₃⁻ Resin, SE Hicap Resin, and COO⁻ Resin. Examples in the Eshmuno® series include CPX, S, Q and HCX resins. Ion exchange media from other sources can be used as well.

[000103] After FXa variant protein has been sufficiently purified, the purified protein can be concentrated using ultrafiltration and diafiltration in preparation for packaging or further processing.

[000104] In some embodiments, purified and concentrated FXa variant proteins can be diluted in aqueous solutions containing buffers and other ingredients suitable for administration to subjects, and then packaged for long term storage until delivery. Additional processing steps can be employed. For example, compositions comprising FXa variant proteins, excipients, buffers, and other pharmaceutically acceptable ingredients can be lyophilized for long term storage.

[000105] FXa variant proteins of the disclosure, whether produced according to the methods described in the Examples below, or by other methods, can be analyzed to determine the presence and degree of structural heterogeneity using methods familiar to those of ordinary skill in the art.

[000106] For example, in some embodiments, intact protein or proteolytic fragments prepared therefrom, can be analyzed by direct protein sequencing, for example, using Edman degradation. Protein or peptides can also be analyzed by liquid chromatography (LC), including reverse phase LC (RP-LC), as well as mass spectrometry, including electrospray ionization quadrupole time-of-flight mass spectrometry (ESI-QTOF MS) to detect variation in amino acid sequence and the presence of particular post-translational modifications. In addition, the presence of charged groups, for example, γ -carboxylated glutamic acid residues (Gla) can be detected using anion exchange liquid chromatography (AEX-HPLC).

[000107] Where the presence of carbohydrate moieties suspected, their existence can be confirmed by removing them with specific glycosidases. Thereafter, treated protein, which should be devoid of the carbohydrate removed by the glycosidase, can be reanalyzed, for example, by mass spectrometry, to confirm the identity of the carbohydrate hypothesized to have been present.

[000108] Other analytical techniques familiar to those of ordinary skill in the art can be used to detect the extent and identity of structural heterogeneity in FXa variant proteins.

[000109] Compositions comprising FXa variant proteins of the disclosure can be tested for pro-coagulant activity. In a non-limiting example, clotting activity is measured with a one-stage clotting assay using Factor VIII deficient human plasma as substrate. One such assay is known as the activated partial thromboplastin time (APTT) assay, but other assays may be used according to the knowledge of those ordinarily skilled in the art. APTT assay is described in more detail in Kamal, A.F., et al., How to interpret and pursue an abnormal prothrombin time, activated partial thromboplastin time, and bleeding time in adults, Mayo Clin Proc. 82(7):864-873 (2007).

[000110] In certain embodiments, compositions of the disclosure include a “therapeutically effective amount” or a “prophylactically effective amount” of FXa variant proteins. A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result, for example hemostasis of uncontrolled bleeding. Therapeutically effective amount of the FXa variant may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the FXa variant to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the FXa variant are outweighed by the therapeutically beneficial effects. A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired prophylactic result, for example, prevention of uncontrolled bleeding in a susceptible subject. For example, a dose may be given prior to a planned surgery. Examples of susceptible subjects include those with Hemophilia A or B, including those subjects producing inhibitory antibodies to factor replacement products.

[000111] Compositions of the disclosure can be administered to a subject, including a human patient, in need of treatment for or prevention of any condition characterized by insufficient coagulation or an excess of bleeding. Non-limiting examples of such conditions include Hemophilia A, Hemophilia B, Hemophilia A or B associated with inhibitory antibodies, coagulation factor deficiency, vitamin K epoxide reductase C1 deficiency, gamma-carboxylase deficiency, bleeding associated with trauma, injury, thrombosis, thrombocytopenia, stroke, coagulopathy, disseminated intravascular coagulation (DIC), over-anticoagulation treatment disorders, Bernard Soulier syndrome, Glanzman thrombasthenia, and storage pool deficiency. Treatment or prevention of other disorders characterized by insufficient coagulation or excess bleeding is also possible.

[000112] Dosage regimens can be adjusted to provide the optimum desired response (e.g., a therapeutic or prophylactic response). For example, a single bolus can be administered,

several divided doses can be administered over time or the dose can be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. According to some embodiments, parenteral compositions are formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form refers to physically discrete units to serve as unitary dosages for the subjects to be treated, with each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the appropriate pharmaceutical carrier.

[000113] In certain embodiments, a therapeutically or prophylactically effective amount of FXa variant proteins is about 0.0001 to 50 mg/kg, about 0.001 to 50 mg/kg, about 0.001 to 5 mg/kg, about 0.001 to 0.5 mg/kg, about 0.001 to 0.05 mg/kg, about 0.01 to 5 mg/kg or about 0.01 to 0.5 mg/kg. According to related embodiments, a therapeutically or prophylactically effective serum concentration of FXa variant proteins is about 0.0003 to 300 nM, about 0.003 to 300 nM, about 0.03 to 300 nM, about 0.003 to 30 nM, about 0.03 to 30 nM or about 0.3 to 3 nM. The serum concentration of the FXa variant may be measured by any method known in the art.

[000114] Unit doses of the compositions of the disclosure can be prepared to conveniently permit the administration of therapeutically or prophylactically effective amounts of FXa variant proteins to subjects or to achieve a desired serum concentration of such proteins in such subjects.

[000115] For any particular subject, specific dosage regimens or ranges may be adjusted over time according to the need of the subject being treated and the professional judgment of the individual responsible for administering or supervising the administration of the compositions. Accordingly, the dosage regimens and ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the compositions of the disclosure.

[000116] Additionally provided are kits including, for example, one or more unit doses of a composition of the disclosure. Such unit doses can be in liquid form or as a lyophilisate. If the composition is packaged in lyophilized form, kits can additionally include a container filled with diluent, such as sterile water or saline solution, for resuspending the dried pellet. Kits can also include articles to be used for administration, including for example, hypodermic syringes, or tubing and butterfly needles, and the like. Kits can additionally include directions for use, alcohol pads for sterilizing skin, or other components.

[000117] Compositions comprising FXa variant protein may be administered once or multiple times until bleeding stops or adequate coagulation is achieved according to the knowledge of one of ordinary skill. Where multiple administrations are used, they may be

hourly, daily, weekly or at other intervals. Administrations may be on a schedule such as every 10 minutes, every 15 minutes, every 20 minutes, every 30 minutes, every hour, every two hours, every three hours, every four hours, three times daily, twice daily, once daily, once every two days, once every three days, and once weekly. The FXa variant may also be administered continuously, e.g. via a minipump. The FXa variant may be administered via a parenteral route (e.g., intravenously, subcutaneously, intraperitoneally, or intramuscularly). The FXa variant may be administered once, twice, or more, or for at least the period of time required to achieve effective coagulation.

[000118] In another embodiment, compositions comprising FXa variant protein may be co-administered with another procoagulant including a different FXa variant (e.g., one having a different substitution at amino acid number 146 than leucine), Factor IX, Factor XIa, Factor XIIa, Factor VIII, Factor VIIa, FEIBA or prothrombin complex concentrate (PCC).

[000119] Compositions comprising FXa variant proteins may further comprise a pharmaceutically acceptable carrier or vehicle, which may be solvents, dispersion media, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like that are physiologically compatible. Some examples of pharmaceutically acceptable carriers merely by way of illustration, are water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. Isotonifying agents, for example, salts (e.g., sodium chloride), sugars (e.g., sucrose), or polyalcohols (e.g., mannitol or sorbitol) can also be included in the composition. Additional examples of pharmaceutically acceptable substances are wetting agents, emulsifying agents, preservatives or buffers, which enhance the stability or other attributes of the instant compositions.

[000120] Compositions for use according to the disclosure may be in any suitable form for administration to a subject, for example, as a liquid for injection or for infusion. The form depends on the intended mode of administration and therapeutic application.

[000121] Therapeutic compositions are usually sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating FXa variant proteins of the disclosure in the required amount in an appropriate solvent with one or a combination of ingredients described above, followed by filtered sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation include vacuum drying or freeze drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile filtered solution thereof.

[000122] The following examples are for illustrative purposes only and should not be construed as in any way limiting the scope of the claims or of other inventions disclosed herein.

EXAMPLES

EXAMPLE 1: Expression and Purification of Recombinant FXa Variant Protein

[000123] Using standard molecular biology techniques and reagents, cDNA encoding FX variant protein including the I16L mutation was generated in which the native Factor X Activation Peptide amino acid sequence was replaced with the amino acid sequence RKR (single letter code) which, in concert with an RKR sequence at the C-terminus of the light chain, formed a PACE proteolytic cleavage site having the amino acid sequence RKRRKR (SEQ ID NO:5). The sequence of the cDNA and mature protein encoded thereby are described in FIG. 1D (SEQ ID NO:4) and FIG. 1A (SEQ ID NO:1), respectively. In this way, the FX precursor of the FXa variant could be activated by intracellular proteolytic cleavage by PACE enzymes co-expressed with the clotting factor. Additionally, the native signal sequence and propeptide of human FX were replaced with those of human prothrombin.

[000124] The cDNA encoding the FX variant was subcloned into an expression vector under the control of a constitutive promoter. The vector also expressed the neomycin resistance gene, permitting antibiotic selection of stable transfectants. A second vector was created to express the soluble form of the PACE enzyme.

[000125] CHO K1 host cells were adapted to grow in serum-free suspension and co-transfected by lipofection with the linearized vectors for expressing FXa variant and soluble PACE. Transfected cells were selected by growth in media supplemented with G418 antibiotic and Vitamin K1. Resistant clones were screened for recombinant FXa variant protein expression and activity. Clones exhibiting relatively high recombinant FXa variant protein expression and activity were adapted to growth in serum-free suspension culture. One clone was selected and used to establish a pre-master cell bank and then a master cell bank using chemically defined medium free of animal or human-derived components.

[000126] To produce drug substance for further study, cells expressing FXa variant were grown at 500 liter (L) or 2500 L scale using chemically defined media. To start, vials of cells from a cell bank were thawed, cultured, and progressively expanded by growth in vessels of increasing volume using a chemically defined, animal component-free medium. Once expansion was complete, cultures were maintained in small stirred tank bioreactors. For

production, culturing was continued in 500 L or 2500 L scale bioreactors until harvest. To harvest, the cell culture medium was centrifuged to remove cells and debris. Medium was further clarified by depth filtration and filtration through a 0.2 µm filter.

[000127] The filtered medium was then diluted with buffer and loaded onto a Capto mixed mode chromatography (MMC) column (GE Healthcare Life Sciences). After washing, product was eluted from the column. The partially purified product was then diluted into a detergent solution for viral inactivation. After viral inactivation, product was loaded onto a Q-Sepharose Fast Flow Chromatography column (GE Healthcare Life Sciences), washed and eluted from the column. Eluted product was then diluted with buffer and loaded onto a Fractogel SO₃⁻ Chromatography column (Merck Millipore), washed and eluted from the column. Eluted product was then filtered using Virosart® CPV filters (Sartorius Stedim Biotech) to remove any residual virus that might have been present. Following viral filtration, purified product was concentrated using ultrafiltration and diafiltration. Purified drug substance was then tested to characterize the FXa variant protein produced by the CHO cells.

EXAMPLE 2: Terminal Amino Acid Sequence

[000128] Amino-terminal sequencing of FXa variant by Edman degradation was used to confirm the first ten residues of both the light chain amino-terminus and heavy chain amino-terminus. The analysis was performed by automated Edman degradation after FXa variant was resolved by SDS-PAGE under non-reducing and reducing conditions and electro-blotted onto a polyvinylidene difluoride (PVDF) membrane.

[000129] The light chain amino-terminal sequence was ANSFL(X)(X)MKK (single letter code) (SEQ ID NO:15). No signal occurred at residues 6 and 7 (shown as X), which is consistent with the presence of gamma carboxyglutamic acid (Gla) residues at those positions. The heavy chain amino-terminal sequence was LVGGQE(Z)KDG (SEQ ID NO:16). No signal occurred at residue 7 (shown as Z), which is consistent with the presence of cysteine at that position. These results confirmed the expected amino-terminal sequences of both the light chain and heavy chain compared to the theoretical sequence.

EXAMPLE 3: Peptide Mapping to Confirm Primary Structure and Identify Post-translational Modifications

[000130] To confirm the primary amino acid sequence and to detect post-translational modifications, FXa variant protein produced as described in Example 1 was analyzed by peptide mapping. Specifically, the protein was reduced, alkylated, and digested with Lys-C lysyl endopeptidase (*Achromobacter lyticus* protease 1) after which the fragments were analyzed by reverse-phase high performance liquid chromatography/electrospray ionization quadrupole time-of-flight mass spectrometry (RP-HPLC/ESI-QTOF MS).

[000131] An example of the resulting peptide map profile is shown in FIG. 2 in which peptides are labeled according to their inferred relative positions in relation to the theoretical sequence of the light chain shown in FIG. 1B and the heavy chain shown in FIG. 1C. Peaks labeled with the letter “L” indicate a light chain peptide and those labeled with the letter “H” indicate a heavy chain peptide. Minor peaks representing buffer-related products are labeled “R”. Several trace-level peaks representing deamidated (d) or overalkylated peptides are labeled “d” and “*”, respectively.

[000132] The peptides detected by peptide mapping, their theoretical and observed masses, and amino acid sequences are shown in Table 1 below. Amino acid numbering is based on FIG. 1A (SEQ ID NO:1).

TABLE 1

Peptide	Residues	Theoretical Mass (Da)	Observed Mass (Da)	Confirmed Sequence	Experimental Observations
L1 + 2 Gla	1-9	1155.475	1155.477	ANSFLEEMK (SEQ ID NO:17)	2 Gla residues observed
L2	10	146.105		K	Not detected
L3 + 8 Gla	11-36	3588.204	3588.208	GHLERECMEETCSYEEAREVFEDSDK (SEQ ID NO:18)	8 Gla residues observed; 6 or 7 Gla residues also observed at trace levels.
L4	37-43	937.429	937.434	TNEFWNK (SEQ ID NO:19)	0 Gla major species
L4 + 1 Gla	37-43	981.419	981.419	TNEFWNK (SEQ ID NO:19)	1 Gla minor species
L5	44-45	309.169		YK	Not detected
L6	46-60	1724.625	1724.625	DGDQCETSPCQNQGK (SEQ ID NO:20)	

Peptide	Residues	Theoretical Mass (Da)	Observed Mass (Da)	Confirmed Sequence	Experimental Observations
L7	61-62	249.115		CK	Not detected
L8+OH	63-79	1952.766	1952.773	<u>D</u> GLGEYTCTCLEGFEGK (SEQ ID NO:21)	
L9	80-87	1067.507	1067.512	NCELFTRK (SEQ ID NO:22)	
L10 + Hex	88-122	4245.634	4245.639	LCSLDNGDCDQFCHEEQN SVVCSCARGYTLADNGK (SEQ ID NO:23)	
L11	123-134	1321.568	1321.575	ACIPTGPYPGCK (SEQ ID NO:24)	
L12	135-141	929.541	929.542	QTLERRK (SEQ ID NO:25)	Also observed QTLERR (SEQ ID NO:26) and QTLER (SEQ ID NO:27) as minor and trace species, respectively
H1	146-153	890.417	890.417	LVGGQECK (SEQ ID NO:28)	
H2	154-193	4620.081	4620.084	DGECPWQALLINEENEGF CGGTILSEFYILTAHCLYQAK (SEQ ID NO:29)	
H3	194-196	449.275	449.276	RFK	
H4	197-221	2777.394	2777.398	VRVGDRNTEQEEGGEAV HEVEVVIK (SEQ ID NO:30)	
H5	222-227	801.424	801.425	HNRFTK (SEQ ID NO:31)	
H6	228-240	1581.840	1581.843	ETYDFDIAVLRK (SEQ ID NO:32)	
H7	241-268	3263.578	3263.581	TPITFRMNVAPACLPERD WAESTLMTQK (SEQ ID NO:33)	
H8	269-281	1387.721	1387.729	TGIVSGFGRTHEK (SEQ ID NO:34)	
H9	282-289	944.551	944.553	GRQSTRK (SEQ ID NO:35)	
H10	290-302	1610.743	1610.746	MLEVPYVDRNSCK (SEQ ID NO:36)	

Peptide	Residues	Theoretical Mass (Da)	Observed Mass (Da)	Confirmed Sequence	Experimental Observations
H11	303-321	2182.991	2182.992	LSSSFITQNMFCAGYDTK (SEQ ID NO:37)	
H12	322-339	1988.865	1988.865	QEDACQGDSGGPHVTRFK (SEQ ID NO:38)	
H13	340-357	2045.952	2045.953	DTYFVTGIVSWGEGCARK (SEQ ID NO:39)	
H14	358-359	203.127		GK	Not detected
H15	360-365	743.385	743.385	YGIYTK (SEQ ID NO:40)	
H16	366-371	677.411	677.415	VTAFLEK (SEQ ID NO:41)	
H17	372-378	934.469	934.470	WIDRSMK (SEQ ID NO:42)	
H18	379-384	670.413	670.413	TRGLPK (SEQ ID NO:43)	
H19	385-386	217.143		AK	Not detected
H20	387-399	2183.958	2183.956	SHAPEVITSSPLK (SEQ ID NO:44)	Observed with 1 and 2 di-sialylated O-glycans. Major form ends in Leu ³⁹⁸ . Minor form ends in Lys ³⁹⁹ .

[000133] Theoretical masses were calculated with PAWS (Genomic Solutions, Ann Arbor, MI). Observed masses were calculated from the most abundant multiply-charged ion in the mass spectrum. All observed masses agreed with theoretical masses to within 10 ppm. Sequence coverage among the peptides for the light chain (LC) and heavy chain (HC) was approximately 94 and 98%, respectively. Three peptides from the L chain (L2, L5, L7) and two peptides from the heavy chain (H14, H19) were not detected. Calculated differences between theoretical and observed peptide mass permitted inference of the presence of certain post-translational modifications. Those residues found to be subject to modification are denoted by underlining in column 5 of Table 1.

[000134] The N-terminal Gla domain in the light chain contained between 9 and 11 total gamma carboxylated glutamic acid (Gla) residues. The major light chain species contained 10 Gla residues. Although peptide L1 was always carboxylated, heterogeneity was detected

in peptides L3 and L4. Specifically, L3 peptide was observed with 6, 7, or 8 Gla residues and L4 peptide was observed with none or one Gla residue. See Table 1.

[000135] The heterogeneity as to gamma-carboxylation was confirmed using anion exchange high-performance liquid chromatography (AEX-HPLC) to measure charge heterogeneity. AEX-HPLC separation of FXa variant protein resolved isoforms containing 9, 10, and 11 gamma carboxyglutamic acid (Gla) residues. The relative frequency of the number of Gla residues within the FXa variant light chain was consistent among seven preparations of purified FXa variant protein tested. In particular, based on relative peak distributions, the frequency of occurrence of 9 Gla residues ranged from 10% to 17% (mean = 12.1%, SD = 2.5%), the frequency of occurrence of 10 Gla residues ranged from 41% to 46% (mean = 43.0%, SD = 1.8%) and the frequency of occurrence of 11 Gla residues ranged from 36% to 46% (mean = 42.1%, SD = 3.3%).

[000136] Purified 10 and 11 Gla isoforms were shown to be equally active in an APTT-based clotting assay, whereas the 9 Gla isoform had reduced activity (approximately 20% of the activity of the 10 and 11 Gla isoforms when normalized for mass).

[000137] Other post-translational modifications of the light chain were detected as well. In particular, the mass of peptide L8 was observed to be consistent with the presence of a beta-hydroxylated aspartic acid residue at position 63 (Asp⁶³) (SEQ ID NO:1). In addition, peptide L10 was observed with a mass difference indicating the addition of an O-linked hexose.

[000138] The heavy chain was also subject to O-linked glycosylation. In the peptide map, the major form of the heavy chain alpha isoform C-terminus was observed ending at amino acid L³⁹⁸ and the minor form of the C-terminus was observed ending at amino acid K³⁹⁹. Both of these peptides were observed containing one or two di-sialylated core-1 O-glycans, with the predominant species of both peptides containing one di-sialylated core-1 O-glycan. In the peptide map profile, the peak labeled “H20³” corresponds to the H20 peptide ending at L³⁹⁸, and the peak labeled “H20²” corresponds to the H20 peptide ending at K³⁹⁹, each containing one O-linked glycan. The peak labeled “H20¹” corresponds to the H20 peptide ending at K³⁹⁹ and containing two O-linked glycans. An H20 peptide ending at L³⁹⁸ having two O-linked glycans was detected at trace levels and is not labeled in FIG. 2. The O-glycans in H20 were localized to Thr³⁹⁴ and Ser³⁹⁵ by beta-elimination followed by MS fragmentation.

EXAMPLE 4: Structural Heterogeneity Determined by High Performance Liquid Chromatography

[000139] Reversed phase high-performance liquid chromatography (RP-HPLC) was used to measure structural heterogeneity in one preparation of intact purified FXa variant protein. Purified protein was injected onto a reversed phase column and eluted with a gradient of acidified acetonitrile. Consistent with the observations from peptide mapping analysis, the FXa variant contained structural heterogeneity at the C-terminus of the heavy chain resulting in two major structural isoforms, alpha (terminating at Lys³⁹⁹) and beta (terminating at Lys³⁸⁶). An additional minor isoform, called gamma, was observed due to clipping of the heavy chain after Lys²⁸¹ or Arg²⁸³. The HPLC chromatogram from these experiments is shown in FIG. 3.

[000140] The relative frequency of the major structural isoforms was consistent among seven preparations of purified FXa variant protein tested. In particular, based on relative peak distributions, the frequency of occurrence of the alpha isoform ranged from 54% to 74% (mean = 66.9%, SD = 7.0%), the frequency of occurrence of the beta isoform ranged from 24% to 42% (mean = 31.4%, SD = 6.5%), and the frequency of occurrence of the gamma isoform ranged from 1% to 3% (mean = 2.0%, SD = 0.8%).

[000141] Alpha and beta isoforms were purified using size-exclusion HPLC and tested for activity in a standard APTT-based clotting assay. The beta isoform demonstrated slightly lower activity (79%) compared to the alpha isoform (normalized as 100%). The activity of the minor gamma species was not tested.

EXAMPLE 5: Structural Heterogeneity of Intact Protein Determined by Mass Spectroscopy

Molecular mass (M_r) of intact FXa variant from one preparation of purified protein was analyzed by RP-HPLC followed by mass spectrometry using a high-resolution, hybrid quadrupole time-of-flight mass analyzer (ESI-QTOF MS). M_r values were determined from the zero-charge mass spectra after deconvolution of the multiply charged data with Waters MaxEnt-1 software. The relative errors between the theoretical and observed major and minor mass values were all within ± 60 ppm, which is consistent with the performance specifications of the Waters Q-ToF mass spectrometer. The HPLC chromatogram is shown in FIG. 4. The mass spectrum of intact FXa variant protein is shown in FIG. 5, and the peak assignments based on the mass spectrum are listed in Table 2 below. Amino acid numbering is based on FIG. 1A (SEQ ID NO:1). ¹ANS...LER¹³⁹ corresponds to SEQ ID NO:6;

¹ANS...ERR¹⁴⁰ corresponds to SEQ ID NO:7; ¹ANS...RRK¹⁴¹ corresponds to SEQ ID NO:9; ¹ANS...RKR¹⁴² corresponds to SEQ ID NO:11; ¹ANS...KRR¹⁴³ corresponds to SEQ ID NO:12; ¹⁴⁶LVG...LPK³⁸⁴ corresponds to SEQ ID NO:45; ¹⁴⁶LVG...KAK³⁸⁶ corresponds to SEQ ID NO:46; ¹⁴⁶LVG...SPL³⁹⁸ corresponds to SEQ ID NO:14; and ¹⁴⁶LVG...PLK³⁹⁹ corresponds to SEQ ID NO:3.

TABLE 2

Light Chain	Heavy Chain	Heavy Chain O-glycan	Glc No.	Observed M _r	Abundance
¹ ANS...ERR ¹⁴⁰	(β) ¹⁴⁶ LVG...LPK ³⁸⁴	0	9	43394.6	Trace
			10	43438.6*	Minor
			11	43482.3*	Major
¹ ANS...LER ¹³⁹	(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	9	43438.6*	Minor
			10	43482.3*	Major
¹ ANS...RRK ¹⁴¹	(β) ¹⁴⁶ LVG...LPK ³⁸⁴	0	9	43525.0	Minor
			10	43566.4	Minor
			11	43610.0	Trace
¹ ANS...ERR ¹⁴⁰	(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	9	43594.3	Trace
			10	43638.0	Major
			11	43681.1*	Major
¹ ANS...RKR ¹⁴²	(β) ¹⁴⁶ LVG...LPK ³⁸⁴	0	9	43681.1*	Major
			10	43722.3	Minor
			11	43766.0*	Major
¹ ANS...RRK ¹⁴¹	(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	9	43722.3	Minor
			10	43766.0*	Major
			11	43810.2	Major
¹ ANS...RKR ¹⁴²	(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	9	43879.8*	Trace
			10	43923.4*	Minor
			11	43966.7	Minor
¹ ANS...KRR ¹⁴³	(β) ¹⁴⁶ LVG...LPK ³⁸⁴	0	9	43879.8*	Trace
			10	43923.4*	Minor
			11	43966.7	Minor
¹ ANS...KRR ¹⁴³	(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	10	44079.7	Trace
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+1NeuAc – Gal	10	45353.0	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+1NeuAc – Gal	10	45477.6*	Minor
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	1+1NeuAc – Gal	10	45477.6*	Minor
¹ ANS...LER ¹³⁹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+1NeuAc	10	45356.0	Trace
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+1NeuAc	10	45514.4	Minor
			11	45557.5	Minor
¹ ANS...LER ¹³⁹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+2NeuAc	9	45603.5	Minor

Light Chain	Heavy Chain	Heavy Chain O-glycan	Gla No.	Observed M _r	Abundance
			10	45645.9	Minor
			11	45690.9	Minor
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+2NeuAc	10	45807.0	Minor
			11	45848.0	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+2NeuAc	9	45886.7	Minor
			10	45931.7	Minor
			11	45975.3	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	1+2NeuAc	9	46016.1	Trace
			10	46061.6	Minor
			11	46102.9	Minor
¹ ANS...RRK ¹⁴²	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1+2NeuAc	11	46132.4	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	1+2NeuAc	9	46172.3*	Trace
¹ ANS...RRK ¹⁴²	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	1+2NeuAc	9	46172.3*	Trace
			10	46218.1	Minor
			11	46261.7	Minor
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+2NeuAc	10	46297.4*	Trace
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	2+2NeuAc	10	46297.4*	Trace
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+2NeuAc	10	46427.8	Minor
			11	46469.6	Minor
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+3NeuAc	10	46591.8*	Minor
			11	46633.7*	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	2+3NeuAc	10	46591.8*	Minor
			11	46633.7*	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+3NeuAc	10	46718.1	Minor
			11	46760.5	Minor
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	2+4NeuAc	11	46796.3	Trace
¹ ANS...ERR ¹⁴⁰	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+4NeuAc	10	46879.7	Minor
			11	46924.6	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...SPL ³⁹⁸	2+4NeuAc	10	46879.7	Minor
			11	46924.6	Minor
¹ ANS...RRK ¹⁴¹	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+4NeuAc	9	46965.4	Trace
			10	47009.2	Minor
			11	47053.1	Minor
¹ ANS...RRK ¹⁴²	(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2+4NeuAc	10	47165.7	Minor

[000142] HPLC again demonstrated structural heterogeneity of the FXa variant protein (FIG. 4). The peaks labeled A correspond to gamma species. The peak labeled B correspond to the alpha form of the protein including two O-glycans. The peak labeled C correspond to the alpha form of the protein including one O-glycan. The peak labeled D corresponds to the

beta form of the protein in which the heavy chain terminates with K³⁸⁶. The peak labeled E corresponds to the beta form of the protein in which the heavy chain terminates with K³⁸⁴.

[000143] Three regions were observed in the mass spectrum (FIG. 5). The first, at the lower end of the mass spectrum (left) corresponds to intact FXa variant protein containing the heavy chain beta isoform. This region also contains so-called major species which occur at greater prevalence compared to others based on relative peak intensity. The major species include those containing the beta form of the heavy chain terminating at K³⁸⁶ in which the carboxy-terminal residue of the light chain was R¹³⁹, R¹⁴⁰, or K¹⁴¹, apparently due to variable cleavage at PACE cleavage site. Species in which the light chain terminated at R¹⁴², also apparently due to variable cleavage, were also present as a minor species. Among these major and minor species, the light chain included β -hydroxy Asp⁶³ and an O-linked hexose, as well as 10 or 11 Gla residues.

[000144] Two regions of greater mass also appear in the mass spectrum, both of which correspond to minor species of intact FXa variant protein containing the heavy chain alpha isoform (terminating at L³⁹⁸ or K³⁹⁹). One group of such species further includes a single O-glycan and a second group further includes two O-glycan post-translational modifications (center and right in FIG. 5, respectively).

[000145] Assignments expressly noted in FIG. 5 correspond to intact FXa variant protein species in which the light chain contained 10 Gla residues, β -hydroxy Asp⁶³ and O-linked glucose. Adjacent peaks in the spectrum labeled “*” represent species containing 11 Gla residues, which increases mass by 44.0 Da.

[000146] Table 2 lists the major and minor species identified from the mass spectrum described above, as well as numerous other minor and trace species identified from the mass spectrum data. In Table 2, species having the same observed M_r value that could not be differentiated from a similar isobaric isoform in the table are denoted “*”. Assignment of apparent abundance of the different species as major, minor and trace were based on relative peak intensities from the mass spectrum data. The column headed “O-glycan” lists the glycosylation status of the heavy chain of the various FXa variant protein species identified in the table. As discussed elsewhere herein, the light chain contains an O-linked hexose. In the table, “0” indicates presence of no O-linked glycans in the heavy chain; “1+1NeuAc” indicates presence of one core-1 mono-sialylated O-glycan; “1+1NeuAc – Gal” indicates the presence of one core-1 mono-sialylated O-glycan in which the terminal galactose (Gal) was not present; “1+2NeuAc” indicates presence of one core-1 di-sialylated O-glycan; “2+2NeuAc” indicates presence of two core-1 O-glycans one of which is di-sialylated or both

of which are mono-sialylated; “2+3NeuAc” indicates presence of two core-1 O-glycans one of which is mono-sialylated and the other of which is di-sialylated; and “2+4NeuAc” indicates presence of two core-1 O-glycans each of which is di-sialylated.

EXAMPLE 6: Structural Heterogeneity of Reduced and Alkylated Protein Determined by Mass Spectroscopy

[000147] To confirm the analysis described above, purified intact FXa variant protein was reduced and alkylated to eliminate interchain and intrachain disulfide bonds. The separated light and heavy chains were then analyzed by RP-HPLC/ESI-QTOF MS to determine their amino acid structure and post-translational modification.

[000148] The HPLC chromatogram is shown in FIG. 6 in which the light and heavy chains are labeled. The alpha and beta isoforms of the heavy chain elute as multiple chromatographic peaks. Trace levels of gamma species (beta heavy chain truncated after amino acid K²⁸¹ or R²⁸³) were observed eluting just before the heavy chain at approximately 48-50 minutes in the chromatogram.

[000149] The zero-charge mass spectrum of the light chain is shown in FIG. 7A and the species identified from the spectrum, including their relative abundance as major, minor or trace, are listed in Table 3 below. Amino acid numbering is based on FIG. 1A (SEQ ID NO:1). ¹ANS...LER¹³⁹ corresponds to SEQ ID NO:6; ¹ANS...ERR¹⁴⁰ corresponds to SEQ ID NO:7; ¹ANS...RRK¹⁴¹ corresponds to SEQ ID NO:9; ¹ANS...RKR¹⁴² corresponds to SEQ ID NO:11; and ¹ANS...KRR¹⁴³ corresponds to SEQ ID NO:12.

TABLE 3

Light Chain	Gla No.	Observed M _r	Abundance
¹ ANS...LER ¹³⁹	9	17169.8	Trace
	10	17213.4	Minor
	11	17257.5	Trace
¹ ANS...ERR ¹⁴⁰	9	17326.1	Minor
	10	17370.0	Major
	11	17413.7	Major
¹ ANS...RRK ¹⁴¹	9	17454.2	Minor
	10	17498.0	Major
	11	17542.0	Major
¹ ANS...RKR ¹⁴²	9	17610.0	Trace
	10	17654.2	Minor
	11	17698.0	Trace

Light Chain	Gla No.	Observed M_r	Abundance
¹ ANS...KRR ¹⁴³	10	17810.0	Trace

[000150] As seen with intact FXa variant protein, all light chain species were observed to include β -hydroxylation of Asp⁶³ and an O-linked hexose. Glutamic acid carboxylation in the Gla domain and PACE cleavage were variable, however, yielding light chain species with 9, 10 or 11 Gla residues and carboxy-terminal heterogeneity produced by ragged cleavage at the PACE site.

[000151] The zero-charge mass spectrum of the heavy chain is shown in FIG. 7B and the various species identified from the spectrum, including their relative abundance as major, minor or trace, are listed in Table 4 below. Amino acid numbering is based on FIG. 1A (SEQ ID NO:1). ¹⁴⁶LVG...LPK³⁸⁴ corresponds to SEQ ID NO:45; ¹⁴⁶LVG...KAK³⁸⁶ corresponds to SEQ ID NO:46; ¹⁴⁶LVG...SPL³⁹⁸ corresponds to SEQ ID NO:14; and ¹⁴⁶LVG...PLK³⁹⁹ corresponds to SEQ ID NO:3.

TABLE 4

Heavy chain	O-glycan	Observed M_r	Abundance
(β) ¹⁴⁶ LVG...LPK ³⁸⁴	0	27484.2	Minor
(β) ¹⁴⁶ LVG...KAK ³⁸⁶	0	27683.9	Major
(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1 core-1	29558.7	Minor
(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1 core-1 + 1 NeuAc	29687.4	Minor
(α) ¹⁴⁶ LVG...SPL ³⁹⁸	1 core-1 + 2 NeuAc	29851.0	Minor
(α) ¹⁴⁶ LVG...PLK ³⁹⁹	1 core-1 + 2 NeuAc	29978.8	Minor
(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2 core-1 + 2 NeuAc	30344.9	Minor
(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2 core-1 + 3 NeuAc	30635.2	Minor
(α) ¹⁴⁶ LVG...SPL ³⁹⁸	2 core-1 + 4 NeuAc	30799.9	Trace
(α) ¹⁴⁶ LVG...PLK ³⁹⁹	2 core-1 + 4 NeuAc	30926.7	Minor

[000152] In the mass spectrum, the heavy chain was observed as three predominant regions corresponding to the beta and alpha isoforms. The observed M_r of the most abundant heavy chain isoform (27683.9 Da) compared well with the theoretical mass of the beta heavy chain isoform (27684.3 Da; ¹⁴⁶LVG...KAK³⁸⁶) with all nine cysteine residues alkylated.

Additional mass heterogeneity was due to variation in the carboxy-terminal residue and O-glycosylation of the alpha isoforms. Specifically, certain peaks corresponded to alpha isoforms terminating at L³⁹⁸ or K³⁹⁹, whereas other peaks corresponded to heavy chains carrying one or two core-1 O-glycans that were variably non-sialylated, mono-sialylated, or

di-sialylated. In FIG. 7B, presence of the “*” indicates overalkylation of the heavy chain which is consistent with peaks observed by peptide mapping of the FXa variant protein (see Example 3).

[000153] The structural heterogeneity observed in the light and heavy chains after reduction and alkylation are consistent with the results observed using intact FXa variant protein.

[000154] Theoretical masses were calculated with PAWS software (2000.06.08, Genomic Solutions, Ann Arbor, MI). Observed masses were determined from the zero-charge mass spectra after deconvolution of the multiply charged data with Waters MaxEnt-1 software. The relative errors between the theoretical and observed major and minor mass values were all less than 60 ppm, which is consistent with the performance specifications of the Waters Q-ToF mass spectrometer for intact glycoprotein analysis. Assignments of major, minor and trace species were based on respective peak intensities, which correlate with isoform abundance.

EXAMPLE 7: In Vitro Clotting Assay

[000155] Clotting activity of seven different preparations of FXa variant protein was determined using a one-stage clotting assay using Factor VIII deficient human plasma as substrate (APTT assay). Specific activity was determined by dividing clotting activity (mg/ml) by protein concentration (mg/ml). Average clotting activity across the preparations was 1.52 mg/ml with a standard deviation of 0.09. Average specific activity was 101.3% with a standard deviation of 5.9.

[000156] The present disclosure is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the disclosure in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

[000157] Various references, including patent applications, patents, and technical publications, are cited herein. The disclosure of each such reference is hereby incorporated herein by reference in its entirety.

CLAIMS

What is claimed is:

1. A composition comprising at least one beta form of the FXa variant protein selected from among the group consisting of:

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 139 and 146 to 386 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 140 and 146 to 384 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 140 and 146 to 386 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 141 and 146 to 384 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 141 and 146 to 386 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 142 and 146 to 384 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 142 and 146 to 386 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 143 and 146 to 384 of the amino acid sequence of SEQ ID NO:1,
and

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 143 and 146 to 386 of the amino acid sequence of SEQ ID NO:1.

2. A composition comprising at least one alpha form of the FXa variant protein selected from among the group consisting of:

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 139 and 146 to 398 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 140 and 146 to 398 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 140 and 146 to 399 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 141 and 146 to 398 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 141 and 146 to 399 of the amino acid sequence of SEQ ID NO:1,

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 142 and 146 to 398 of the amino acid sequence of SEQ ID NO:1, and

A FXa variant protein wherein the light chain and heavy chain protein sequence respectively consist of amino acids 1 to 142 and 146 to 399 of the amino acid sequence of SEQ ID NO:1.

3. The composition of claim 1, wherein the light chain of said protein is modified to include β -hydroxy Asp⁶³, an O-linked hexose and 9, 10 or 11 Gla residues.
4. The composition of claim 3, wherein the light chain of said protein is modified to include 9 Gla residues.
5. The composition of claim 3, wherein the light chain of said protein is modified to include 10 Gla residues.
6. The composition of claim 3, wherein the light chain of said protein is modified to include 11 Gla residues.
7. The composition of claim 2, wherein the light chain of said protein is modified to include β hydroxy Asp⁶³, an O-linked hexose and 9, 10 or 11 Gla residues.
8. The composition of claim 7, wherein the light chain of said protein is modified to include 9 Gla residues.
9. The composition of claim 7, wherein the light chain of said protein is modified to include 10 Gla residues.
10. The composition of claim 7, wherein the light chain of said protein is modified to include 11 Gla residues.
11. The composition of claim 2, wherein the heavy chain of said protein is modified to include one core-1 O-linked glycan.
12. The composition of claim 11, wherein the core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
13. The composition of claim 12, wherein the core-1 O-linked glycan is non-sialylated.
14. The composition of claim 12, wherein the core-1 O-linked glycan is mono-sialylated.
15. The composition of claim 12, wherein the core-1 O-linked glycan is di-sialylated.
16. The composition of claim 2, wherein the heavy chain of said protein is modified to include two core-1 O-linked glycans.

17. The composition of claim 16, wherein the first of said two core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
18. The composition of claim 17, wherein the first core-1 O-linked glycan is non-sialylated.
19. The composition of claim 17, wherein the first core-1 O-linked glycan is mono-sialylated.
20. The composition of claim 17, wherein the first core-1 O-linked glycan is di-sialylated.
21. The composition of claim 16, wherein the second of said two core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
22. The composition of claim 21, wherein the second core-1 O-linked glycan is non-sialylated.
23. The composition of claim 21, wherein the second core-1 O-linked glycan is mono-sialylated.
24. The composition of claim 21, wherein the second core-1 O-linked glycan is di-sialylated.
25. The composition of claim 7, wherein the heavy chain of said protein is modified to include one core-1 O-linked glycan.
26. The composition of claim 25, wherein the core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
27. The composition of claim 26, wherein the core-1 O-linked glycan is non-sialylated.
28. The composition of claim 26, wherein the core-1 O-linked glycan is mono-sialylated.
29. The composition of claim 26, wherein the core-1 O-linked glycan is di-sialylated.
30. The composition of claim 7, wherein the heavy chain of said protein is modified to include two core-1 O-linked glycans.
31. The composition of claim 30, wherein the first of said two core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
32. The composition of claim 31, wherein the first core-1 O-linked glycan is non-sialylated.
33. The composition of claim 31, wherein the first core-1 O-linked glycan is mono-sialylated.
34. The composition of claim 31, wherein the first core-1 O-linked glycan is di-sialylated.

- 35.** The composition of claim 30, wherein the second of said two core-1 O-linked glycan is non-sialylated, mono-sialylated or di-sialylated.
- 36.** The composition of claim 35, wherein the second core-1 O-linked glycan is non-sialylated.
- 37.** The composition of claim 35, wherein the second core-1 O-linked glycan is mono-sialylated.
- 38.** The composition of claim 35, wherein the second core-1 O-linked glycan is di-sialylated.
- 39.** A composition comprising FXa variant protein, wherein said FXa variant protein is selected from among the group of FXa variant protein species listed in Table 2.
- 40.** The composition of claim 39, wherein said composition comprises at least 5 of the FXa variant protein species listed in Table 2.
- 41.** The composition of claim 39, wherein said composition comprises at least 10 of the FXa variant protein species listed in Table 2.
- 42.** The composition of claim 39, wherein said composition comprises at least 15 of the FXa variant protein species listed in Table 2.
- 43.** The composition of claim 39, wherein said composition comprises at least 20 of the FXa variant protein species listed in Table 2.
- 44.** The composition of claim 39, wherein said composition comprises at least 25 of the FXa variant protein species listed in Table 2.
- 45.** The composition of claim 39, wherein said composition comprises at least 30 of the FXa variant protein species listed in Table 2.
- 46.** The composition of claim 39, wherein said composition comprises at least 35 of the FXa variant protein species listed in Table 2.
- 47.** The composition of claim 39, wherein said composition comprises at least 40 of the FXa variant protein species listed in Table 2.
- 48.** The composition of claim 39, wherein said composition comprises at least 45 of the FXa variant protein species listed in Table 2.
- 49.** The composition of claim 39, wherein said composition comprises at least 50 of the FXa variant protein species listed in Table 2.
- 50.** The composition of claim 39, wherein said composition comprises at least 55 of the FXa variant protein species listed in Table 2.
- 51.** The composition of claim 39, wherein said composition comprises at least 60 of the FXa variant protein species listed in Table 2.

- 52.** A composition comprising FXa variant protein, wherein the light chain of said FXa variant protein is selected from among the group of light chain species listed in Table 3.
- 53.** The composition of claim 52, wherein the heavy chain of said FXa variant protein is selected from among the group of heavy chain species listed in Table 4.
- 54.** A method of treating a subject in need of hemostasis comprising the step of administering to said subject an amount of the composition of any one of claims 1, 2, 39 and 52 effective to cause hemostasis in said subject.
- 55.** The method of claim 54, wherein said subject has Hemophilia A.
- 56.** The method of claim 54, wherein said subject has Hemophilia B.
- 57.** The method of claim 54, wherein said subject has trauma.
- 58.** A method for purifying a Factor Xa (FXa) variant protein comprising the steps of:
- (i) contacting mixed mode chromatography (MMC) medium with filtered cell culture medium containing FXa variant protein,
 - (ii) eluting FXa variant protein bound to the MMC chromatography medium,
 - (iii) contacting anion exchange chromatography medium with FXa variant protein eluted at step (ii),
 - (iv) eluting FXa variant protein bound to the anion exchange chromatography medium,
 - (v) contacting cation exchange chromatography medium with FXa variant protein eluted at step (iv); and
 - (vi) eluting FXa variant protein bound to the cation exchange chromatography medium.
- 59.** The method of claim 58, wherein the mixed mode chromatography medium is capable of interacting with proteins via electrostatic interaction, hydrophobic interaction, hydrogen bonding, and thiophilic interaction.
- 60.** The method of claim 58, further comprising at least an additional step for inactivating or removing viruses.
- 61.** The method of claim 58, wherein the mixed mode chromatography medium is Capto MMC, the anion exchange chromatography medium is Q-Sepharose Fast Flow, and the cation exchange chromatography medium is Fractogel® SO₃⁻.
- 62.** The method of claim 60, further comprising steps of ultrafiltration and diafiltration to concentrate the purified FXa variant protein.

FIG. 1A

Amino acid sequence of FX Variant Protein (SEQ ID NO:1)

1 ANSFLEEMKKGHLERECMEETCSYEEAREVFEDSDKTNEFWNKYKDGDQC 50

51 ETSPCQNGKCKDGLGEYTCLEGEFGKNCELFTRKLCSLDNGDCDQFC 100

101 HEEQNSVVCARGYTLADNGKACIPTGPYPCGKQTLERRRKKRLVGGQ 150

151 ECKDGECPWQALLINEENEGFCGGTILSEFYILTAAHCLYQAKRFKVRVG 200

201 DRNTEQEEGGEAVEVEVVIKHNRFTKETYDFDIAVLRLKTPITFRMNVA 250

251 PACLPERDWAESTLMTQKTGIVSGFGRTHEKGRQSTRLKMLEVPYVDRNS 300

301 CKLSSSFIIITQNMFCAGYDTKQEDACQGDSGGPHVTRFKDTYFVTGIVSW 350

351 GEGCARKGKYGIYTKVTAFLLKWIDRSMKTRGLPKKSHAPEVITSSPLK 399

FIG. 1B

Predicted amino acid sequence of FXa variant protein light chain (SEQ ID NO:2)

001 A N S F L E E M K K G H L E R E C M E E T C S Y E E A R E V 030
031 F E D S D K T N E F W N K Y K D G D Q C E T S P C Q N Q G K 060
061 C K D G L G E Y T C T C L E G F E G K N C E L F T R K L C S 090
091 L D N G D C D Q F C H E E Q N S V V C S C A R G Y T L A D N 120
121 G K A C I P T G P Y P C G K Q T L E R R K R K R 145

FIG. 1C

Predicted amino acid sequence of FXa variant protein heavy chain (SEQ ID NO:3)

001 L V G G Q E C K D G E C P W Q A L L I N E E N E G F C G G T 030
031 I L S E F Y I L T A A H C L Y Q A K R F K V R V G D R N T E 060
061 Q E E G G E A V H E V E V I K H N R F T K E T Y D F D I A 090
091 V L R L K T P I T F R M N V A P A C L P E R D W A E S T L M 120
121 T Q K T G I V S G F G R T H E K G R Q S T R L K M L E V P Y 150
151 V D R N S C K L S S S F I I T Q N M F C A G Y D T K Q E D A 180
181 C Q G D S G G P H V T R F K D T Y F V T G I V S W G E G C A 210
211 R K G K Y G I Y T K V T A F L K W I D R S M K T R G L P K A 240
241 K S H A P E V I T S S P L K 254

3/10

FIG. 1D

cDNA encoding FX variant protein (SEQ ID NO:4)

ATGGCGCACGTCCGAGGCTTGCAGCTGCCTGGCTGCCCTGGCTGCCCTGTGTAGCCTTG
TGCACAGCCAGCATGTGTTCCTGGCTCCTCAGCAAGCACGGTCGCTGCTCCAGCGGTCCGGCG
AGCCAAATTCCTTTCTTGAAGAGATGAAGAAAGACACCTCGAAAGAGAGTGCATGGAAGAGACC
TGCTCATACGAAGAGGCCCGAGGTCTTTGAGGACAGCGACAGACGAATGAATCTGGAATA
AATACAAAGATGCGGACCAGTGTGAGACCAGTCTTGCCAGAACCCAGGCAAAATGTAAAGACGG
CCTCGGGGAATACACCTGCACCTGTTTAGAAGATTGGAAGGCAAAAACCTGTGAATTATTCACA
CGAAGCTCTGCAGCCTGGACAAACGGGACTGTGACCACTTCTGCCACGAGGAACAGAACTCTG
TGGTGTCTCCTGCGCCCGGGTACACCTGGCTGACAAACGGCAAGGCCCTGCATTCCCACAGG
GCCCTACCCCTGTGGAAACAGACCCCTGGAACGCAAGGAAGCGCTAAGCGTCTCGTGGGAGGC
CAGGAATGCAAGACGGGGAGTGTCCCTGGCAGGCCCTGCTCATCAATGAGGAAAACGAGGGTT
TCTGTGGTGGAACTATTCTGAGCGAGTTCTACATCTTAACGGCAGCCCACTGTCTTACCAAGC
CAAGAGATTCAAGGTGAGGTAGGTGACCGGAAACACGGAGCAGGAGGCGGTGAGGCGGTG
CACGAGGTGAGGTGTCATCAAGCACAAACCGGTTCACAAAGGAGACCTATGACTTCGACATCG
CCGTGCTCCGGCTCAAGACCCCATCACCTTCCGCATGAACGTGGCGCCTGCCCTCCCCGA
GGTGAATGGCCGAGTCCACGCTGATGACGCAAGACGGGATGTGAGCGGCTTCGGGCGC
ACCCACGAGAAAGGCCCGCAGTCCACCAAGGCTCAAGATGCTGAGGTGCCCTACGTGGACCGCA
ACAGCTGCAAGCTGTCCAGCAGCTTCATCATCAACCAGAACATGTTCTGTGCCGGCTACGACAC
CAAGCAGGAGGATGCCCTGCCAGGGGACAGCGGGGCCCGCACGTCACCCGCTTCAAGGACACC
TACTTCGTGACAGGCATCGTCAGCTGGGGAGAGGGCTGTGCCCGTAAGGGGAAGTACGGGATCT
ACACCAAGGTCAACCGCCTTCTCAAGTGGATCGACAGGTCCATGAAAACCCAGGGCTTGCCCCAA
GGCCAAAGAGCCATGCCCCGGAGGTCATAACGTCTCTCCATTAAAGTGA

FIG. 2




FIG. 3




FIG. 4




FIG. 5





FIG. 6



9/10

FIG. 7A
Light Chain





FIG. 7B
Heavy Chain



INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2014/064564

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07K14/745 A61K38/36 A61P7/04
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LACRAMIOARA IVANCIU ET AL: "A zymogen-like factor Xa variant corrects the coagulation defect in hemophilia", NATURE BIOTECHNOLOGY., vol. 29, no. 11, 23 October 2011 (2011-10-23), pages 1028-1033, XP055111224, US	2,39, 54-57
Y	ISSN: 1087-0156, DOI: 10.1038/nbt.1995 the whole document	1-62
X	WO 2013/049804 A1 (PHILADELPHIA CHILDREN HOSPITAL [US]) 4 April 2013 (2013-04-04)	2,39,54, 56,57
Y	page 4 - page 5 page 11 example 1	1-62
	----- -/-	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

29 January 2015

Date of mailing of the international search report

06/02/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Weisser, Dagmar

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2014/064564

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2009/175931 A1 (CAMIRE RODNEY M [US]) 9 July 2009 (2009-07-09)	2,39,54, 56,57
Y	table 1 paragraphs [0106], [0110]	1-62
X	M. W. BUNCE ET AL: "Zymogen-like factor Xa variants restore thrombin generation and effectively bypass the intrinsic pathway in vitro", BLOOD, vol. 117, no. 1, 6 January 2011 (2011-01-06), pages 290-298, XP055111227, US ISSN: 0006-4971, DOI: 10.1182/blood-2010-08-300756	2,39
Y	the whole document	1-62
X	SUN YAN-MEI ET AL: "Vitamin K epoxide reductase significantly improves carboxylation in a cell line overexpressing factor X", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 106, no. 12, 4 August 2005 (2005-08-04), pages 3811-3815, XP002377556, ISSN: 0006-4971, DOI: 10.1182/BLOOD-2005-06-2495	2,39
Y	page 3812, columns 2,6,7	1-62

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2014/064564

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013049804	A1	04-04-2013	
		AU 2012315516 A1	17-04-2014
		CA 2850603 A1	04-04-2013
		EP 2760887 A1	06-08-2014
		JP 2015501137 A	15-01-2015
		US 2014248259 A1	04-09-2014
		WO 2013049804 A1	04-04-2013

US 2009175931	A1	09-07-2009	
		AU 2006315175 A1	24-05-2007
		BR PI0618654 A2	06-09-2011
		CA 2629491 A1	24-05-2007
		CN 101356192 A	28-01-2009
		CR 9956 A	21-08-2008
		DK 1948690 T3	01-09-2014
		EC SP088452 A	30-07-2008
		EP 1948690 A2	30-07-2008
		EP 2423224 A1	29-02-2012
		EP 2431389 A1	21-03-2012
		EP 2431390 A1	21-03-2012
		ES 2496567 T3	19-09-2014
		GT 200800065 A	03-11-2008
		HK 1128699 A1	06-06-2014
		JP 2009515559 A	16-04-2009
		JP 2014073127 A	24-04-2014
		KR 20080078664 A	27-08-2008
		KR 20140005383 A	14-01-2014
		KR 20140115374 A	30-09-2014
		NZ 568108 A	30-09-2011
		PT 1948690 E	08-09-2014
		RU 2011130932 A	27-01-2013
		SI 1948690 T1	30-09-2014
		SV 2008002906 A	22-10-2008
		US 2009175931 A1	09-07-2009
		US 2014120155 A1	01-05-2014
		WO 2007059513 A2	24-05-2007
		ZA 200804026 A	28-10-2009
