




(12) PATENT ABRIDGMENT (11) Document No. AU-B-82467/91
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 636788

- (54) Title
NOVEL SULFONYLUREAS
- (51)⁵ International Patent Classification(s)
C07D 239/47 A01N 047/36 C07D 239/48 C07D 239/52
C07D 251/16 C07D 251/52
- (21) Application No. : **82467/91** (22) Application Date : **14.08.91**
- (30) Priority Data
- (31) Number (32) Date (33) Country
2654/90 15.08.90 CH SWITZERLAND
- (43) Publication Date : **11.06.92**
- (44) Publication Date of Accepted Application : **06.05.93**
- (71) Applicant(s)
CIBA-GEIGY AG
- (72) Inventor(s)
WILLY MEYER
- (74) Attorney or Agent
SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001
- (56) Prior Art Documents
EP 0044808
- (57) Claim

1. An N-phenylsulfonyl-N'-pyrimidinyl- or -triazolyl-urea of formula I



wherein

R is hydrogen or C₁-C₄alkyl;

R₁ is hydrogen, halogen, C₁-C₅alkyl, C₂-C₅alkenyl, C₁-C₄haloalkyl, COO-R₂ or A-R₃;

R₂ is C₁-C₅alkyl;

R₃ is C₁-C₅alkyl, C₂-C₄alkoxyalkyl or C₁-C₅haloalkyl;

X is C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy or C₁-C₃alkoxy mono- to tri-substituted by halogen;

A is oxygen, sulfur, SO or SO₂;

Y is halogen, C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy, C₁-C₃alkoxy mono- to tri-substituted by halogen; cyclopropyl, methylamino or dimethylamino;

E is nitrogen or the methine group; and
Z is fluorine or chlorine;
or a salt of that compound.

16. A herbicidal and plant-growth-inhibiting composition comprising one or more sulfonylureas of formula I, according to claim 1, together with a herbicidally effective diluent, carrier and/or adjuvant.

636788

S & F Ref: 188520

AUSTRALIA

PATENTS ACT 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

ORIGINAL

Name and Address
of Applicant:

Ciba-Geigy AG
Klybeckstrasse 141
CH-4002 Basle
SWITZERLAND

Actual Inventor(s): Willy Meyer

Address for Service: Spruson & Ferguson, Patent Attorneys
Level 33 St Martins Tower, 31 Market Street
Sydney, New South Wales, 2000, Australia

Invention Title: Novel Sulfonylureas

The following statement is a full description of this invention, including the best method of performing it known to me/us:-

PH/5-18197/A


Novel sulfonylureas

The present invention relates to novel herbicidally active and plant-growth-regulating N-phenylsulfonyl-N'-pyrimidinyl- and -triazinyl-ureas, to processes for the preparation thereof, to compositions comprising them as active ingredients, and to the use thereof for
5 controlling weeds, especially selectively in crops of useful plants, or for regulating and inhibiting plant growth.

Phenylsulfonylureas having herbicidal action are known from European Patent Applications No. 0 044 808 and No. 0 072 347. The compounds specifically disclosed therein are not, however, always able to satisfy requirements in respect of potency and
10 activity spectrum. There is accordingly a need for compounds having improved and more selective action.

Novel sulfonylureas having improved herbicidal and plant-growth-regulating properties have now been found.

The N-phenylsulfonyl-N'-pyrimidinyl- and -triazinyl-ureas according to the invention
15 correspond to formula I



wherein

R is hydrogen or C₁-C₄alkyl;

R₁ is hydrogen, halogen, C₁-C₅alkyl, C₂-C₅alkenyl, C₁-C₄haloalkyl, COO-R₂ or A-R₃;

R₂ is C₁-C₅alkyl;

R_3 is C_1 - C_5 alkyl, C_2 - C_4 alkoxyalkyl or C_1 - C_5 haloalkyl;

X is C_1 - C_3 alkyl, C_1 - C_3 alkyl mono- to tri-substituted by halogen; C_1 - C_3 alkoxy or C_1 - C_3 alkoxy mono- to tri-substituted by halogen;

A is oxygen, sulfur, SO or SO_2 ;

- 5 Y is halogen, C_1 - C_3 alkyl, C_1 - C_3 alkyl mono- to tri-substituted by halogen; C_1 - C_3 alkoxy, C_1 - C_3 alkoxy mono- to tri-substituted by halogen; cyclopropyl, methylamino or dimethylamino;

E is nitrogen or the methine group; and

Z is fluorine or chlorine;

- 10 and the salts of those compounds.

The alkyl groups occurring in the definitions of the substituents may be straight-chained or branched and are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl or the isomers of pentyl. Within the scope of the present invention, alkyl is preferably methyl or ethyl.

- 15 Halogen is to be understood as being fluorine, chlorine, bromine or iodine, preferably fluorine or chlorine.

Haloalkyl is, for example, trifluoromethyl, 1-fluoroethyl, 1,1-dichloroethyl, 3,3,3-trifluoropropyl, 2-fluoroisopropyl, 3-fluoropropyl, 1,1,1-trichloropentyl, 1-fluoro-3-methylpentyl, or 1-bromohexyl.

- 20 Examples of alkoxyalkyl are: methoxymethyl, methoxyethyl, methoxypropyl, ethoxymethyl, ethoxyethyl and propoxymethyl.

- The C_1 - C_3 alkyl radicals occurring as or in the substituents X and Y include, in particular, methyl, ethyl, n-propyl and isopropyl and also the haloalkyl radicals that are derived from those radicals and are mono- to tri-substituted by halogen. The alkyl radicals occurring as
25 or in the substituents X and Y preferably have one or two carbon atoms.

- Of the C_1 - C_3 alkyl groups that occur as or in the substituents X and Y and are mono- to tri-substituted by halogen, C_1 - C_2 alkyl groups mono- to tri-substituted by fluorine or chlorine are preferred. Especially preferred C_1 - C_3 alkyl radicals that occur as or in the substituents X and Y and are mono- to tri-substituted by halogen are: trifluoromethyl,
30 difluoromethyl, 2-chloroethyl, chlorodifluoromethyl, dichloromethyl, chlorofluoromethyl,

1,1-dichloroethyl, trifluoroethyl, 3,3,3-trifluoropropyl and 2,3-dichloropropyl, with fluoromethyl, chloromethyl, difluoromethyl and trifluoromethyl being particularly preferred.

5 The C₂-C₅alkenyl radicals occurring in the substituent R₁ may be in the Z form (cis) or in the E form (trans) and may be straight-chained or branched. Alkenyl radicals having a chain length of two or three carbon atoms are preferred. Examples of C₂-C₅alkenyl radicals are: vinyl, allyl, methallyl, 1-methylvinyl, 2-methylvinyl, but-2-en-1-yl and pent-3-en-1-yl. Vinyl and allyl are preferred.

10 The invention also includes the salts that the compounds of formula I can form with amines, alkali and alkaline earth metal bases or quaternary ammonium bases.

Of the alkali and alkaline earth metal hydroxides as salt-formers, prominence is to be given to the hydroxides of lithium, sodium, potassium, magnesium and calcium, but especially to the hydroxides of sodium and potassium.

15 Examples of amines that are suitable for salt formation are primary, secondary and tertiary aliphatic and aromatic amines, such as methylamine, ethylamine, propylamine, isopropylamine, the four isomers of butylamine, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, but especially ethyl-, propyl-, diethyl- or triethyl-amine, and more especially
20 isopropylamine and diethanolamine.

Examples of quaternary ammonium bases are generally the cations of haloammonium salts, for example the tetramethylammonium cation, the trimethylbenzylammonium cation, the triethylbenzylammonium cation, the tetraethylammonium cation and the trimethylethylammonium cation, but also the ammonium cation.

25 Preferred compounds of formula I are those wherein R₁ at the phenyl ring is in the 5-position and the group -S-CH₂-CHF-Z is in the 2-position.

Other preferred compounds of formula I are those wherein R₁ and/or R are (is) hydrogen.

Very especially preferred groups of compounds of formula I are those wherein

- a) X is C₁-C₃alkyl, C₁-C₃alkoxy or C₁-C₃alkoxy mono- to tri-substituted by halogen; and
Y is chlorine, C₁-C₃alkyl, C₁-C₃alkoxy, cyclopropyl, or C₁-C₃alkoxy mono- to tri-substituted by halogen; or
5 b) X is methyl, methoxy, ethoxy or difluoromethoxy; and
Y is methyl, methoxy, ethoxy, difluoromethoxy or chlorine; or
c) X is methoxy or ethoxy; and Y is methyl or methoxy; or
d) E is nitrogen.


A group of compounds of formula I that is especially prominent owing to its good
10 biological activity is that wherein R₁ is hydrogen and X is C₁-C₃alkyl, C₁-C₃alkoxy or C₁-C₃alkoxy mono- to tri-substituted by halogen; and
Y is chlorine, C₁-C₃alkyl, C₁-C₃alkoxy, cyclopropyl, or C₁-C₃alkoxy mono- to tri-substituted by halogen. Preferred among this group are also those compounds of formula
I wherein X is methyl, methoxy, ethoxy or difluoromethoxy and Y is methyl, methoxy,
15 ethoxy, difluoromethoxy or chlorine, the meanings methoxy and ethoxy being very especially preferred for X and the meanings methyl and methoxy being very especially preferred for Y. In this outstanding group of compounds, R₁ at the phenyl ring is preferably in the 5-position and the group -S-CH₂-CHF-Z is preferably in the 2-position.

The following may be mentioned as preferred individual compounds within the scope of
20 formula I:

N-(2-(2-chloro-2-fluoroethylthio)-phenyl-sulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea; and N-(2-(2,2-difluoroethylthio)-phenyl-sulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea.


The invention also provides a process for the preparation of a compound of formula I of
25 the invention, which comprises

- a) reacting a phenylsulfonamide of formula XIV




wherein R₁ and Z are as defined under formula I,
in the presence of a base, with an N-pyrimidinyl or N-triazinyl carbamate of formula XI






wherein X, Y, R and E are as defined under formula I and R₇ is C₁-C₄alkyl, or phenyl which may be substituted by C₁-C₄alkyl or by halogen; or

b) reacting a phenylsulfonamide of formula XII



5

wherein R₁ and Z are as defined in formula I, and B is $\text{R}_7\text{---O---C(=O)---NH---}$ or O=C=N- , in the presence of a base, with a 2-amino-pyrimidine or -triazine of formula XIII



wherein E, X and Y are as defined under formula I.

10 The invention further provides a herbicidal and plant-growth inhibiting composition comprising one or more sulfonylureas of formula I, of the invention together with a herbicidally effective diluent, carrier and/or adjuvant.

The invention provides a method of controlling undesirable plant growth, which comprises applying to the plants or the locus thereof an effective amount of a compound

15 of formula I, of the invention, or of a composition comprising that compound.

The invention also provides a method of inhibiting plant growth, which comprises applying to the plants or to the locus thereof an effective amount of a compound of formula I, of the invention, or of a composition comprising that active ingredient.

The reactions to form compounds of formula I are advantageously carried out in aprotic, 20 inert organic solvents. Such solvents are hydrocarbons, such as benzene, toluene, xylene




- or cyclohexane, chlorinated hydrocarbons, such as dichloromethane, trichloromethane, tetrachloromethane or chlorobenzene, ethers, such as diethyl ether, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether, tetrahydrofuran or dioxane, nitriles, such as acetonitrile or propionitrile, amides, such as dimethylformamide, diethylformamide or
- 5 N-methylpyrrolidone. The reaction temperatures are preferably from -20° to $+120^{\circ}\text{C}$.

- The reactions generally proceed in a slightly exothermic manner and can be carried out at room temperature. In order to reduce the reaction time or also to initiate the reaction, it is expedient to heat the reaction mixture briefly at boiling point. The reaction times can also be reduced by adding a few drops of base as a reaction catalyst. Suitable bases are
- 10 especially tertiary amines, such as trimethylamine, triethylamine, quinuclidine, 1,4-diazabicyclo(2.2.2)octane, 1,5-diazabicyclo(4.3.0)non-5-ene or 1,5-diazabicyclo(5.4.0)undec-7-ene. Other bases that may be used, however, are inorganic bases, such as hydrides, such as sodium or calcium hydride, hydroxides, such as sodium and potassium hydroxide, carbonates, such as sodium and potassium carbonate, or hydrogen carbonates, such as
- 15 potassium and sodium hydrogen carbonate.

The end products of formula I can be isolated by concentration and/or evaporation of the solvent and can be purified by recrystallisation or trituration of the solid residue in solvents in which they are not readily soluble, such as ethers, aromatic hydrocarbons or chlorinated hydrocarbons.

- 20 The intermediates of formulae XI, XII and XIII are known or can be prepared analogously to known processes. Processes for the preparation of N-pyrimidinyl and N-triazinyl carbamates are described, for example, in EP-A-0 101 670. Processes for the preparation of the compounds of formula XII are described in EP-A-0 044 808. Compounds of formula XIII are known from EP-A-0 070 804.

- 25 The compounds of formula XIV can be prepared by
- a) converting a 2-halophenylsulfonamide of formula II




(II)


wherein X_1 is fluorine, chlorine or bromine, in the presence of a base, with a mercaptan of formula III




wherein R_5 is C_1 - C_6 alkyl or C_1 - C_6 alkyl substituted by phenyl, into a 2-sulfenylphenylsulfonamide of formula IV




- 5 wherein R_5 is as defined under formula III,
b) oxidising that compound to the 2-sulfinylphenylsulfonamide of formula V



- wherein R_5 is as defined under formula III,
c) reacting the resulting 2-sulfinylphenylsulfonamide of formula V in the presence of an acid to form the disulfide of formula VI




- 10 d) reducing the disulfide of formula VI to the 2-mercaptophenylsulfonamide of formula VII



e) then converting that compound using a trialkylamine of formula X



wherein R_6 is C_1 - C_4 alkyl, into the 2-mercaptophenylsulfonamide trialkylamine salt of formula VIII



wherein R_6 is as defined under formula X, and

5 f) then reacting that compound with a halofluoroethane of formula IX



wherein Y_1 is chlorine or bromine and Z is fluorine or chlorine.

The compounds of formulae II, III, IV, V, VII and IX are known and some of them are commercially available.

10 The disulfides of formula VI and the 2-mercaptophenylsulfonamide trialkylamine salts of formula VIII are the subject of Swiss Patent Application No. 3 554/89-5.

The C_1 - C_6 alkyl groups occurring in the substituent R_5 may be straight-chained or branched and are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl or the isomers of pentyl, n-hexyl or the isomers of hexyl. The alkyl groups occurring in the substituent R_5 preferably have from 1 to 3 carbon atoms. If
15 the alkyl groups are substituted by phenyl, they preferably have a chain length of from 1 to 3 carbon atoms. The substituent R_5 is especially preferably benzyl.

The C_1 - C_4 alkyl groups occurring in the substituents R_6 may be straight-chained or branched and are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl or tert-butyl. R_6 is in each case especially preferably ethyl.

Process step a) is advantageously carried out in an inert solvent at a temperature of from +20°C to the boiling temperature of the solvent. The temperatures are usually from +20 to +180°C, preferably from +20 to +120°C. An especially preferred temperature range is from +50 to +70°C.

- 5 Suitable solvents are chlorinated hydrocarbons, such as dichloromethane, trichloromethane, trichloroethane or tetrachloroethane, chlorobenzene or dichlorobenzene; aromatic hydrocarbons, such as benzene, toluene or xylene; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran or dioxane; nitriles, such as acetonitrile or propionitrile; cyclohexane, pyridine, N-methylpyrrolidone or N,N-dimethylformamide, N,N-dimethyl-
10 formamide being especially preferred.

The bases used are especially hydrides, hydroxides, carbonates or alcoholates of an alkali metal or of an alkaline earth metal, a trialkylamine or a pyridine base. Especially preferred bases are pyridine, sodium hydroxide, sodium methoxide, sodium ethoxide, sodium carbonate or potassium carbonate. Potassium carbonate is very especially preferred.

- 15 Process step a) can be carried out in an especially advantageous manner when 2-fluorophenylsulfonamide is used as the 2-halophenylsulfonamide of formula II.

Permanganates, periodates, per-acids or hydrogen peroxide are especially suitable as oxidising agents for process step b).

- Preferred oxidising agents are peracetic acid, perbenzoic acid, periodic acid, potassium
20 permanganate, potassium periodate and hydrogen peroxide. A very especially preferred oxidising agent is hydrogen peroxide. The oxidation of the 2-sulphenylphenylsulfonamide of formula IV is advantageously carried out at temperatures of from 0° to +80°C, a temperature range of from 0° to +40°C being preferred.

- In a preferred variant of the process for the preparation of the compounds of formula II,
25 the oxidation of the 2-sulphenylphenylsulfonamide of formula IV is carried out with hydrogen peroxide in the presence of acetic acid at a temperature of from +5 to +15°C.

Process variant c) is preferably carried out at a temperature of from +20°C to the boiling temperature of the solvent. Solvents that have proved to be especially suitable are alcohols having a chain length of from 1 to 4 carbon atoms, for example methanol, ethanol,

propanol, isopropanol, butanol or 2-butanol. A preferred solvent is methanol.

Acid-catalysed rearrangements of sulfoxides are known in the literature under the name "Pummerer rearrangements". Examples of such reactions can be found in Adv. Org. Chem. 6, 356 (1969). Contrary to the learning known from that article, the acid-catalysed
5 rearrangement of the 2-sulfinylphenylsulfonamide of formula V does not result in 2-mercaptophenylsulfonamide but in 2,2'-bis-aminosulfonyl-diphenyl-disulfide of formula VI which can be isolated from its reaction medium in a simple manner and has a high degree of storage stability.

The nature of the acids used as catalysts in the reaction is not critical. Preferred acids are
10 mineral acids, especially hydrochloric acid or sulfuric acid.

Process steps b) and c) can also be carried out in direct succession in a reaction vessel without isolating the intermediate of formula V. The solvents suitable for that process variant correspond to those mentioned in stage c).

The reductive cleavage of the disulfide of formula VI to form the
15 2-mercaptophenylsulfonamide of formula VII (process step d)) is generally carried out at temperatures of from +20°C to +100°C.

The reduction is preferably carried out with hydrogen in the presence of noble metal catalysts, or with zinc, iron or tin in the presence of hydrochloric acid or acetic acid, or with sodium, magnesium or aluminium amalgam. Preferred reducing agents are hydrogen
20 in the presence of platinum, palladium, rhodium or nickel catalysts and also zinc, iron and tin in the presence of hydrochloric acid or acetic acid. A very especially preferred reducing agent is zinc in the presence of hydrochloric acid or acetic acid.

A trialkylamine of formula X that is especially suitable for salt formation in accordance with process step e) is triethylamine.

25 The reaction temperatures for the reaction of the 2-mercaptophenylsulfonamide trialkylamine salts with the halofluoroethane of formula IX (process step f)) are from 0° to +80°C, preferably from 0° to +40°C. The reaction proceeds in an especially advantageous manner when Y₁ in formula IX is bromine. The solvents suitable for stage f) correspond to those mentioned in stage a).

The compounds of formula I are generally used successfully at rates of application of from 0.001 to 2 kg/ha, especially from 0.005 to 1 kg/ha. The concentration required to achieve the desired effect can be determined by experiment. It is dependent on the type of action, the stage of development of the cultivated plant and of the weed, and also on the applica-
5 tion (place, time, method) and, in dependence on those parameters, can vary within wide limits.

When used at relatively low rates of application, the compounds of formula I are distinguished by growth-inhibiting and herbicidal properties, which render them excellently suitable for use in crops of useful plants, especially in cereals, cotton,
10 soybeans, rape, maize and rice, their use in cereals being very especially preferred.

The invention relates also to herbicidal and plant-growth-regulating compositions comprising a novel compound of formula I, and to methods of inhibiting plant growth.

Plant growth regulators are substances that bring about agronomically desirable biochemical and/or physiological and/or morphological changes in/to the plant.

15 The active ingredients comprised by the compositions according to the invention influence plant growth in different ways depending on the time of application, the concentration, the type of application and the environmental conditions. Plant growth regulators of formula I can, for example, inhibit the vegetative growth of plants. This type of action is valuable in the case of lawn areas, in the cultivation of ornamentals, in fruit plantations, in the case of
20 roadside embankments and in sports fields and industrial sites, but also in the specific inhibition of side-shoots, as in the case of tobacco. In agriculture, inhibition of the vegetative growth of cereals leads, owing to a strengthening of the stalk, to reduced lodging, and similar agronomic effects are achieved in rape, sunflowers, maize and other cultivated plants. Moreover, by inhibiting the vegetative growth it is possible to increase
25 the number of plants per unit area. Another field of application for growth inhibitors is the selective control of cover plants in plantations or widely spaced crops by greatly inhibiting the growth of the cover crops without killing them, so that competition with the main crop is eliminated but the agronomically positive effects, such as erosion prevention, fixing of nitrogen and loose soil structure, are preserved.

30 A method of inhibiting plant growth is to be understood as being a method of controlling a

plant's natural development without changing its life-cycle, as determined by genetic characteristics, in the sense of mutation. The method of regulating growth is applied at a time in the plant's development that has to be determined for each individual case. The compounds of formula I can be applied pre- or post-emergence, for example to the seeds
5 or seedlings, to roots, tubers, stalks, leaves, blossoms or other parts of the plant. This can be done, for example, by applying the compound as such or in the form of a composition to the plants, and/or by treating the plant's nutrient medium (soil).

Various methods and techniques are suitable for the use of the compounds of formula I or of compositions containing them for regulating plant growth, for example the following:

10 i) Seed dressing

a) Dressing the seeds with an active ingredient formulated as a wettable powder, by shaking in a container until the formulation is uniformly distributed over the surface of the seeds (dry dressing). Up to 4 g of compound of formula I (in the case of a 50 %
formulation: up to 8.0 g of wettable powder) are used per 1 kg of seed.

15 b) Dressing the seeds with an emulsifiable concentrate of the active ingredient or with an aqueous solution of the compound of formula I formulated as a wettable powder according to method a) (wet dressing).

c) Dressing by soaking the seeds for a period of from 1 to 72 hours in a liquor containing up to 1000 ppm of compound of formula I and, if desired, subsequently drying the seeds
20 (seed soaking).

Seed dressing or treatment of the germinated seedling are naturally the preferred methods of application because the treatment with the active ingredient is then directed wholly at the target crop. From 0.001 g to 4.0 g of active ingredient are normally used per 1 kg of seed, although, depending on the method employed, which also allows the addition of
25 other active ingredients or micronutrients, amounts that exceed or fall short of the specified concentration limits may be employed (repeat dressing).

ii) Controlled release of active ingredient

A solution of the active ingredient is applied to mineral granulated carriers or polymerised granules (urea/formaldehyde) and allowed to dry. If required, a coating may be applied
30 (coated granules), which allows the active ingredient to be released in metered amounts

over a specific period of time.

The compounds of formula I are used in unmodified form, as obtainable from synthesis, or, preferably, as compositions together with the adjuvants conventionally employed in formulation technology, and are therefore formulated in known manner e.g. into emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable
5 powders, soluble powders, dusts, granules, and also encapsulations in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting, wetting, scattering or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances.

The formulations, i.e. the compositions, preparations or mixtures containing the compound
10 (active ingredient) of formula I and, where appropriate, one or more solid or liquid adjuvants, are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

Suitable solvents are: aromatic hydrocarbons, preferably the fractions containing 8 to 12
15 carbon atoms, such as mixtures of alkylbenzenes, e.g. xylene mixtures or alkylated naphthalenes; aliphatic and cycloaliphatic hydrocarbons such as paraffins, cyclohexane or tetrahydronaphthalene; alcohols such as ethanol, propanol or butanol; glycols and their ethers and esters, such as propylene glycol or dipropylene glycol ether, ketones such as cyclohexanone, isophorone or diacetone alcohol, strongly polar solvents such as
20 N-methyl-2-pyrrolidone, dimethyl sulfoxide or water; vegetable oils and their esters, such as rape oil, castor oil or soybean oil; and, where appropriate, also silicone oils.

The solid carriers used, e.g. for dusts and dispersible powders, are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgit. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or
25 highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, for example pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are, for example, calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues.

30 Depending on the nature of the compound of formula I to be formulated, suitable

surface-active compounds are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

- Both so-called water-soluble soaps and water-soluble synthetic surface-active compounds
5 are suitable anionic surfactants.

Suitable soaps are the alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts of higher fatty acids (C_{10} - C_{22}), e.g. the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained e.g. from coconut oil or tallow oil. Mention may also be made of fatty acid methyltaurin salts.

- 10 More frequently, however, so-called synthetic surfactants are used, especially fatty alcohol sulfonates, fatty alcohol sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates.

- The fatty alcohol sulfonates or sulfates are usually in the form of alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts and contain a
15 C_8 - C_{22} alkyl radical, which also includes the alkyl moiety of acyl radicals, e.g. the sodium or calcium salt of lignosulfonic acid, of dodecyl sulfate or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfated and sulfonated fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain 2 sulfonic acid groups and one fatty acid radical
20 containing 8 to 22 carbon atoms. Examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic acid, dibutyl-naphthalenesulfonic acid, or of a condensate of naphthalenesulfonic acid and formaldehyde.

Also suitable are corresponding phosphates, e.g. salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 mol of ethylene oxide, or phospholipids.

- 25 Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These
5 compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Representative examples of non-ionic surfactants are nonylphenolpolyethoxyethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxy-polyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol.

Fatty acid esters of polyoxyethylene sorbitan, e.g. polyoxyethylene sorbitan trioleate, are
10 also suitable non-ionic surfactants.

Cationic surfactants are preferably quaternary ammonium salts which contain, as N-substituent, at least one C_8 - C_{22} alkyl radical and, as further substituents, unsubstituted or halogenated lower alkyl, benzyl or hydroxy-lower alkyl radicals. The salts are preferably in the form of halides, methyl sulfates or ethyl sulfates, e.g. stearyltrimethylammonium
15 chloride or benzyldi(2-chloroethyl)ethylammonium bromide.

The surfactants customarily employed in formulation technology are described inter alia in the following publications:

- "Mc Cutcheon's Detergents and Emulsifiers Annual", Mc Publishing Corp., Glen Rock, New Jersey, 1988.
- 20 - M. and J. Ash, "Encyclopedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-1981.
- Dr. Helmut Stache "Tensid-Taschenbuch", Carl Hanser Verlag, Munich/Vienna 1981.

The herbicidal compositions generally comprise 0.1 to 99 %, preferably 0.1 to 95 %, of a compound of formula I, 1 to 99 % of a solid or liquid adjuvant and 0 to 25 %, preferably
25 0.1 to 25 %, of a surfactant.

Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The compositions may also comprise further additives such as stabilisers, for example

vegetable oils and epoxidised vegetable oils (epoxidised coconut oil, rape oil or soybean oil), antifoams, for example silicone oil, preservatives, viscosity regulators, binders, tackifiers as well as fertilisers or other active ingredients for obtaining special effects.

Preferred formulations are composed in particular of the following constituents

5 (throughout, percentages are by weight):

Emulsifiable concentrates:

active ingredient:	1 to 90 %, preferably 5 to 20 %
surfactant:	1 to 30 %, preferably 10 to 20 %
liquid carrier:	5 to 94 %, preferably 70 to 85 %

10 Dusts:

active ingredient:	0.1 to 10 %, preferably 0.1 to 1 %
solid carrier:	99.9 to 90 %, preferably 99.9 to 99 %

Suspension concentrates:

active ingredient:	5 to 75 %, preferably 10 to 50 %
15 water:	94 to 24 %, preferably 88 to 30 %
surfactant:	1 to 40 %, preferably 2 to 30 %

Wettable powders:


active ingredient:	0.5 to 90 %, preferably 1 to 80 %
surfactant:	0.5 to 20 %, preferably 1 to 15 %
20 solid carrier:	5 to 95 %, preferably 15 to 90 %

Granules:

active ingredient:	0.5 to 30 %, preferably 3 to 15 %
solid carrier:	99.5 to 70 %, preferably 97 to 85 %

Preparation Examples:


Example P1: Preparation of 2-benzylthiophenylsulfonamide:



175.2 g of 2-fluorophenylsulfonamide and 160 g of potassium carbonate are added to a solution of 124.2 g of benzylmercaptan in 400 ml N,N-dimethylformamide. The reaction mixture is then heated at +60°C for 3 hours. When the mixture has cooled to +25°C it is filtered and the filtrate is then concentrated by evaporation.


1500 ml of water are added to the resulting residue, the product precipitating in the form of colourless crystals. After separating the solution off, the resulting crystals are dissolved in ethyl acetate and then treated with magnesium sulfate. After filtering and then concentrating the filtrate by evaporation, 219 g (78.5 % of the theoretical yield) of 2-benzylthiophenylsulfonamide are obtained in the form of colourless crystals having a melting point of from +104 to +106°C.

Example P2: Preparation of 2-benzylsulfinylphenylsulfonamide:




102 ml of 30 % hydrogen peroxide solution are added dropwise at a temperature of +10°C to a solution of 219 g of 2-benzylthiophenylsulfonamide obtained in accordance with Example P1 in 500 ml of concentrated acetic acid. The reaction mixture is then stirred for 5 hours at a temperature of +25°C, the product slowly crystallising out. After separating off the crystals, washing with water and drying, 219 g (94.6 %) of 2-benzylsulfinylphenylsulfonamide are obtained in the form of colourless crystals having a melting point of from +206 to +209°C.

Example P3: Preparation of 2,2'-bis-aminosulfonyl-diphenyl-disulfide:



- 5 500 ml of concentrated hydrochloric acid are added to a solution of 214 g of a 2-benzylsulfinylphenylsulfonamide prepared in accordance with Example P2 in 500 ml of methanol. After boiling the reaction mixture for 7 hours and maintaining the mixture at room temperature for 2 days, the precipitated disulfide is separated off and washed with water, isopropanol and diethyl ether to yield 133.2 g (97.7 % of the theoretical yield) of 2,2'-bis-aminosulfonyl-diphenyl-disulfide in the form of yellowish crystals having a melting point of +217°C (decomposition).

Example P4: Preparation of 2,2'-bis-aminosulfonyl-diphenyl-disulfide:




- 10 50 ml of concentrated hydrochloric acid and 3.5 g of 30 % hydrogen peroxide solution are added dropwise to a solution of 10 g of 2-benzylthiophenylsulfonamide obtained in accordance with Example P1 in 50 ml of ethanol. The reaction mixture is then heated under reflux for a period of 1 hour. After cooling to +25°C, the crystals formed are separated off and washed with isopropanol and petroleum ether to yield 4 g of
- 15 2,2'-bis-aminosulfonyl-diphenyl-disulfide in the form of yellow crystals having a melting point of from +211 to +213°C.

Example P5: Preparation of 2-mercaptophenylsulfonamide:



18 g of zinc powder are added to a suspension of 23.2 g of a 2,2'-bis-amino-sulfonyl-diphenyl-disulfide obtained in accordance with Examples P3 and P4 in 180 ml of concentrated acetic acid and the suspension is heated under reflux for 30 minutes. It is then cooled to +17°C and filtered. After washing the filter residue with ethyl acetate, the filtrate is concentrated by evaporation and the residue is then dissolved in 200 ml of ethyl acetate. After washing twice with 100 ml of water each time and drying with magnesium sulfate, the solution is concentrated by evaporation to yield 22 g (94 % of the theoretical yield) of unpurified 2-mercaptophenylsulfonamide having a melting point of from +113 to +140°C.


10 Example P6: Preparation of the triethylamine salt of 2-mercaptophenylsulfonamide:



(VIII)


5 g of triethylamine are added dropwise at a temperature of +25°C to a solution of 7.6 g of 2-mercaptosulfonamide obtained in accordance with Example P5 in 60 ml of tetrahydrofuran. The precipitated colourless crystals are then separated off and washed with diethyl ether to yield 9.2 g of the triethylamine salt of 2-mercaptophenylsulfonamide which has a melting point of from +164 to +168°C.

Example P7: Preparation of 2-(2,2-difluoroethylthio)-phenyl-sulfonamide:




A mixture of 145.2 g of the triethylamine salt of 2-mercaptophenylsulfonamide obtained in accordance with Example P6 and 79.7 g of 2-bromo-1,1-difluoroethane in 600 ml of methanol is stirred for 16 hours at a temperature of from +55°C to +60°C. The reaction mixture is then concentrated by evaporation. The residue is stirred with ice-water and the resulting suspension is filtered to yield 120 g (95 % of the theoretical yield) of 2-(2,2-difluoroethylthio)-phenyl-sulfonamide having a melting point of from +111°C to +112°C.

Example P8: Preparation of 2-(2-chloro-2-fluoroethylthio)-phenylsulfonamide:



A mixture of 217.8 g of the triethylamine salt of 2-mercaptophenylsulfonamide obtained in accordance with Example P6 and 101 g of 1,2-dichloro-1-fluoroethane in 950 ml of methanol is stirred in an autoclave for 24 hours at a temperature of from +65°C to +70°C
5 and for a further 24 hours at from +95°C to +100°C. The reaction mixture is then concentrated by evaporation and the residue is stirred with water. By extracting with ethyl acetate, washing with water, drying over sodium sulfate, concentrating by evaporation and purifying the residue by chromatography with methylene chloride, 99.5 g (49.2 % of the theoretical yield) of 2-(2-chloro-2-fluoroethylthio)-phenylsulfonamide having a melting
10 point of from +88°C to +89°C are obtained.

Example P9: Preparation of N-(2-(2-chloro-2-fluoroethylthio)-phenyl-sulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (Compound No. 1.001):




(1.001)

A solution of 1.07 g of 1,8-diazabicyclo[5.4.0]undec-7-ene in 10 ml of absolute dioxane is added dropwise at a temperature of from +20 to +25°C to a mixture of 1.9 g of
15 2-(2-chloro-2-fluoroethylthio)-phenylsulfonamide and 1.85 g of N-(4-methyl-6-methoxy-1,3,5-triazin-2-yl)-phenyl carbamate in 40 ml of absolute dioxane. The reaction mixture is then stirred for 6 hours and subsequently added to water. After acidifying with 10 % hydrochloric acid, extracting with ethyl acetate, washing with water and drying over sodium sulfate, the reaction mixture is filtered. Concentration of the
20 filtrate by evaporation and recrystallisation of the residue from ethyl acetate yield 2.4 g of

N-(2-(2-chloro-2-fluoroethylthio)-phenylsulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (Compound No. 1.001) having a melting point of from +174 to +176°C.

Example P10: Preparation of N-(2-(2,2-difluoroethylthio)-phenylsulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (Compound No. 1.008):



- 5 A solution of 1.07 g of 1,8-diazabicyclo[5.4.0]undec-7-ene in 10 ml of absolute dioxane is added dropwise at a temperature of from +20 to +25°C to a mixture of 1.77 g of 2-(2,2-difluoroethylthio)-phenylsulfonamide and 1.82 g of N-(4-methyl-6-methoxy-1,3,5-triazin-2-yl)-phenyl carbamate in 40 ml of absolute dioxane. The reaction mixture is then stirred for 6 hours and subsequently added to water. After acidifying with 10 %
- 10 hydrochloric acid, extracting with ethyl acetate, washing with water and drying over sodium sulfate, the reaction mixture is filtered. Concentration of the filtrate by evaporation and recrystallisation of the residue from ethyl acetate yield 2.3 g of N-(2-(2,2-difluoroethylthio)-phenylsulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea (Compound No. 1.008) having a melting point of from +189 to +190°C.
- 15 The compounds of formula I listed in the following Table are prepared in an analogous manner:



Table 1: Compounds of formula I:









Table 1:

	Comp. No.	Z	R	R ₁	Y	X	E	m.p. [°C]
5	1.001	Cl	H	H	CH ₃	OCH ₃	N	174-176
	1.002	Cl	CH ₃	H	CH ₃	OCH ₃	N	159-161
	1.003	Cl	H	H	CH ₃	OCH ₃	CH	
	1.004	Cl	H	H		OC ₂ H ₅	N	
	1.005	Cl	H	H		OCH ₃	N	
10	1.006	Cl	H	H	C ₂ H ₅	OCH ₃	N	189-190
	1.007	Cl	H	H	CH ₃	OC ₂ H ₅	N	
	1.008	F	H	H	CH ₃	OCH ₃	N	
	1.009	F	CH ₃	H	CH ₃	OCH ₃	N	
	1.010	F	H	H	CH ₃	OCH ₃	CH	
15	1.011	F	H	H	CH ₃	OC ₂ H ₅	N	170-174
	1.012	F	H	H		OC ₂ H ₅	N	
	1.013	F	H	H		CH ₃	CH	
	1.014	F	H	H		OCH ₃	N	
	1.015	F	H	H	C ₂ H ₅	OCH ₃	N	
20	1.016	F	H	H	NHCH ₃	OC ₂ H ₅	N	
	1.017	F	H	H	N(CH ₃) ₂	OCH ₃	N	
	1.018	F	H	H	N(CH ₃) ₂	OCH ₂ CF ₃	N	
	1.019	F	H	H	Cl	OCH ₃	N	
	1.020	F	H	H	OCHF ₂	OCHF ₂	CH	
25	1.021	F	H	H	OCH ₃	OCH ₃	N	162-167
	1.022	F	H	H	OCH ₃	OCH ₃	CH	179-181
	1.023	F	H	H	Cl	OCH ₃	CH	154-160
	1.024	F	H	H	OCHF ₂	OCH ₃	CH	170-172

Biological Examples

In order to investigate the herbicidal activity of the compounds of the present Application in comparison with compounds of the prior art (EP-A-0 044 808) the following Examples B1 and B2 were carried out:


5 Example B1: Preemergence herbicidal action

In a greenhouse, immediately after the test plants (Amarantus, Chenopodium, Sinapis, Stellaria, Chrysanthemum, Galium and Veronica) have been sown in seed trays, the surface of the soil is treated with an aqueous spray mixture in an amount corresponding to a rate of application of 30 g or 8 g of test compound/hectare. The seed trays are kept in
10 the greenhouse at 22-25°C and 45-60 % relative humidity.

After 3 weeks, the herbicidal action is evaluated according to a scale of nine ratings in comparison with an untreated control group:


- 1: plants have not germinated or are completely withered
- 2-3: very pronounced action
- 15 4-6: medium action
- 7-8: weak action
- 9: no action (as untreated control)

The compounds tested are:




Compound A: (No. 391 of EP-A-0 044 808)

20 against:




No. 1.001 of the present application,

and



against:



5 The results are indicated in Tables T1 and T2:

Table T1: Preemergence herbicidal action with 30 g a.i./ha:

Test plant	Comp. A	Comp. 1.001	Comp. B	Comp. 1.003
Amarantus	5	2	5	3
Chenopodium	9	2	5	2
10 Sinapis	8	2	8	2
Stellaria	9	2	9	2
Chrysanth.	9	2	5	2
Galium	9	3	9	2

Table T2: Preemergence herbicidal action with 8 g a.i./ha:

<u>Test plant</u>	<u>Comp. B</u>	<u>Comp. 1.003</u>
Amarantus	9	7
Sinapis	9	3
5 Stellaria	9	2
Chrysanth.	9	2
Galium	9	3
Veronica	5	2

In this test, the herbicidal activity of the compounds of the present Application proves to
10 be distinctly superior to that of the compounds of the prior art.

Example B2: Postemergence herbicidal action (contact herbicide)

A number of weeds (Amarantus, Chenopodium, Sinapis, Stellaria, Chrysanthemum,
Galium, Viola tricolor and Veronica) are sprayed postemergence (in the 4- to 6-leaf stage)
with an aqueous active ingredient dispersion at a rate of 8 or 30 g of test compound per
15 hectare and kept at 22°-25°C and 45-60 % relative humidity. The test is evaluated 15 days
after the treatment according to a scale of nine ratings in comparison with an untreated
control group:

- 1: plants have not germinated or are completely withered
- 2-3: very pronounced action
- 20 4-6: medium action
- 7-8: weak action
- 9: no action (as untreated control)

The results are indicated in Tables T3 and T4:

Table T3: Postemergence herbicidal action with 8 g a.i./ha:

Test plant	Comp. A	Comp. 1.001
Amarantus	9	3
Chenopodium	9	4
5 Sinapis	9	3
Stellaria	9	3
Chrysanth.	9	5
Galium	9	8
Viola t.	9	7
10 Veronica	9	7

Table T4: Postemergence herbicidal action with 30 g a.i./ha:

Test plant	Comp. A	Comp. 1.001	Comp. B	Comp. 1.003
Amarantus	7	2	-	-
Chenopodium	9	3	8	6
15 Sinapis	3	2	4	2
Stellaria	3	2	-	-
Chrysanth.	4	3	-	-
Galium	6	3	7	6
Viola t.	9	4	4	7
20 Veronica	9	7	-	-

In the case of postemergence application too, the compounds of formula I according to the present Application exhibit better herbicidal activity than do the comparison compounds of the prior art.

Example B3: Herbicidal action in wild rice (paddy rice)

- 25 The weeds *Echinochloa crus galli* and *Monocharia vag.*, which occur in water, are sown in plastic beakers (surface: 60 cm²; volume: 500 ml). After sowing, the beakers are filled with water up to the surface of the soil. 3 days after sowing, the water level is increased to slightly above the soil surface (3-5 mm). Application is effected 3 days after sowing by spraying the beakers with the test compounds. The rate of application corresponds to a concentration of 8-500 g of active ingredient per hectare. The beakers are then kept in the
- 30

greenhouse under optimum growth conditions for rice weeds, i.e. at 25°-30°C and at high humidity.

The evaluation of the tests takes place 3 weeks after application. The compounds of formula I damage the weeds.

5 Formulation Examples for active ingredients of formula I (throughout, percentages are by weight)

<u>1. Wettable powders</u>		a)	b)	c)
	compound of formula I	20 %	50 %	0.5 %
	sodium lignosulfonate	5 %	5 %	5 %
10	sodium laurylsulfate	3 %	-	-
	sodium diisobutylphenylsulfonate	-	6 %	6 %
	octylphenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	2 %
15	highly dispersed silicic acid	5 %	27 %	27 %
	kaolin	67 %	-	-
	sodium chloride	-	-	59.5 %

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of the desired concentration.

<u>2. Emulsifiable concentrates</u>		a)	b)
	compound of formula I	10 %	1 %
	calcium dodecylbenzenesulfonate	3 %	3 %
	octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %	3 %
25	castor oil polyethylene glycol ether (36 mol of ethylene oxide)	4 %	4 %
	cyclohexanone	30 %	10 %
	xylene mixture	50 %	79 %

30 Emulsions of any required concentration can be obtained from such concentrates by

dilution with water.

<u>3. Dusts</u>		a)	b)
	compound of formula I	0.1 %	1 %
	talcum	99.9 %	-
5	kaolin	-	99 %

Ready-for-use dusts are obtained by intimately mixing the carriers with the active ingredient.

<u>4. Extruder granules</u>		a)	b)
	compound of formula I	10 %	1 %
10	sodium lignosulfonate	2 %	2 %
	carboxymethylcellulose	1 %	1 %
	kaolin	87 %	96 %

The active ingredient is mixed and ground with the adjuvants, and the mixture is moistened with water. The mixture is extruded and then dried in a stream of air.

15	<u>5. Coated granules</u>	
	compound of formula I	3 %
	polyethylene glycol (mol. wt. 200)	3 %
	kaolin	94 %

The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

<u>6. Suspension concentrate</u>		a)	b)
	compound of formula I	5 %	40 %
	ethylene glycol	10 %	10 %
25	nonylphenol polyethylene glycol ether (15 mol of ethylene oxide)	1 %	6 %
	sodium lignosulfonate	5 %	10 %
	carboxymethylcellulose	1 %	1 %
	37 % aqueous formaldehyde		

solution	0.2 %	0.2 %
silicone oil in the form of a		
75 % aqueous emulsion	0.8 %	0.8 %
water	77 %	32 %

- 5 The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired concentration can be obtained by dilution with water.

7. Salt solution


compound of formula I	5 %
10 isopropylamine	1 %
octylphenol polyethylene glycol	
ether (78 mol of ethylene oxide)	3 %
water	91 %

- The compounds of formula I are used in unmodified form or, preferably, as compositions
15 together with the adjuvants conventionally employed in formulation technology, and are therefore formulated in known manner e.g. into emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules, and also encapsulations in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting,
20 scattering or pouring are chosen in accordance with the intended objectives and the prevailing circumstances.

What is claimed is:

The claims defining the invention are as follows:

1. An N-phenylsulfonyl-N'-pyrimidinyl- or -triazinyl-urea of formula I




wherein

- 5 R is hydrogen or C₁-C₄alkyl;
R₁ is hydrogen, halogen, C₁-C₅alkyl, C₂-C₅alkenyl, C₁-C₄haloalkyl, COO-R₂ or A-R₃;
R₂ is C₁-C₅alkyl;
R₃ is C₁-C₅alkyl, C₂-C₄alkoxyalkyl or C₁-C₅haloalkyl;
X is C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy or
10 C₁-C₃alkoxy mono- to tri-substituted by halogen;
A is oxygen, sulfur, SO or SO₂;
Y is halogen, C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy,
C₁-C₃alkoxy mono- to tri-substituted by halogen; cyclopropyl, methylamino or
dimethylamino;
15 E is nitrogen or the methine group; and
Z is fluorine or chlorine;
or a salt of that compound.
2. A compound of formula I according to claim 1, wherein
R₁ at the phenyl ring is in the 5-position and the group -S-CH₂-CHF-Z is in the 2-position.
- 20 3. A compound of formula I according to claim 1, wherein
R is hydrogen.
4. A compound of formula I according to claim 1, wherein
R₁ is hydrogen.


5. A compound of formula I according to claim 1, wherein E is nitrogen.
6. A compound of formula I according to claim 1, wherein X is C₁-C₃alkyl, C₁-C₃alkoxy or C₁-C₃alkoxy mono- to tri-substituted by halogen; and
5 Y is chlorine, C₁-C₃alkyl, C₁-C₃alkoxy, cyclopropyl, or C₁-C₃alkoxy mono- to tri-substituted by halogen.
7. A compound of formula I according to claim 1, wherein X is methyl, methoxy, ethoxy or difluoromethoxy; and Y is methyl, methoxy, ethoxy, difluoromethoxy or chlorine.
- 10 8. A compound of formula I according to claim 1, wherein X is methoxy or ethoxy; and Y is methyl or methoxy.
9. A compound of formula I according to claim 4, wherein X is C₁-C₃alkyl, C₁-C₃alkoxy or C₁-C₃alkoxy mono- to tri-substituted by halogen; and Y is chlorine, C₁-C₃alkyl, C₁-C₃alkoxy, cyclopropyl, or C₁-C₃alkoxy mono- to tri-
15 substituted by halogen.
10. A compound of formula I according to claim 9, wherein X is methyl, methoxy, ethoxy or difluoromethoxy; and Y is methyl, methoxy, ethoxy, difluoromethoxy or chlorine.
11. A compound of formula I according to claim 9, wherein
20 X is methoxy or ethoxy; and Y is methyl or methoxy.
12. A compound of formula I according to claim 9, wherein the group -S-CH₂-CHF-Z on the phenyl ring is in the 2-position.
13. N-(2-(2-chloro-2-fluoroethylthio)-phenylsulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea according to claim 1.
- 25 14. N-(2-(2,2-difluoroethylthio)-phenylsulfonyl)-N'-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)-urea according to claim 1.

15. A process for the preparation of a compound of formula I according to claim 1, which comprises

a) reacting a phenylsulfonamide of formula XIV




wherein Z is as defined under formula I, in the presence of a base, with an N-pyrimidinyl
5 or N-triazinyl carbamate of formula XI




wherein X, Y, R and E are as defined under formula I in claim 1 and R7 is C1-C4alkyl, or phenyl which may be substituted by C1-C4alkyl or by halogen; or

b) reacting a phenylsulfonamide of formula XII



wherein R1 and Z are as defined in formula I in claim 1, and B is $R_7-O-C(=O)-N(H)-$ or

10 $O=C=N-$, in the presence of a base, with a 2-amino-pyrimidine or -triazine of formula XIII



wherein E, X and Y are as defined under formula I in claim 1.

16. A herbicidal and plant-growth-inhibiting composition comprising one or more sulfonylureas of formula I, according to claim 1, together with a herbicidally effective 5 diluent, carrier and/or adjuvant.

17. A composition according to claim 16 comprising a compound of formula I according to claim 1 in an amount of from 0.1% to 95%.

18. A method of controlling undesirable plant growth, which comprises applying to the plants or to the locus thereof an effective amount of a compound of formula I, 10 according to claim 1, or of a composition comprising that compound.

19. A method according to claim 18, which comprises applying the active ingredient in an amount of from 0.001 to 2kg per hectare.

20. A method of inhibiting plant growth, which comprises applying to the plants or to the locus thereof an effective amount of a compound of formula I, according to 15 claim 1, or of a composition comprising that active ingredient.

21. A method according to claim 18 for the selective pre- or post-emergence control of weeds in crops of useful plants.

22. An N-phenylsulfonyl-N'-pyrimidinyl- or -triazinyl-urea substantially as hereinbefore described with reference to any one of the Preparation Examples.

20 23. A process for the preparation of an N-phenylsulfonyl-N'-pyrimidinyl- or -triazinyl-urea substantially as hereinbefore described with reference to any one of the Preparation Examples.

24 A herbicidal and plant-growth-inhibiting composition substantially as hereinbefore described with reference to any one of the Formulation Examples.

25 25. A method of controlling undesirable plant growth in a plant, comprising contacting the plant or locus of the plant with an effective amount of an N-phenylsulphonyl-N'-pyrimidinyl- or -triazinyl-urea compound as defined in claim 22 or a herbicidal and plant-growth-inhibiting composition as defined in claim 24.

26. A method of inhibiting plant growth, comprising applying to a plant or to the 30 locus thereof an effective amount of a N-phenylsulphonyl-N'-pyrimidinyl- or -triazinyl-urea compound as defined in claim 22 or a herbicidal and plant-growth-inhibiting composition as defined in claim 24.

Dated 24 February, 1993

Ciba-Geigy AG


Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON



Novel sulfonylureasAbstract:

N-phenylsulfonyl-N'-pyrimidinyl- and -triazinyl-ureas of formula I



wherein

- 5 R is hydrogen or C₁-C₄alkyl;
R₁ is hydrogen, halogen, C₁-C₅alkyl, C₂-C₅alkenyl, C₁-C₄haloalkyl, COO-R₂ or A-R₃;
R₂ is C₁-C₅alkyl;
R₃ is C₁-C₅alkyl, C₂-C₄alkoxyalkyl or C₁-C₅haloalkyl;
X is C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy or
- 10 C₁-C₃alkoxy mono- to tri-substituted by halogen;
A is oxygen, sulfur, SO or SO₂;
Y is halogen, C₁-C₃alkyl, C₁-C₃alkyl mono- to tri-substituted by halogen; C₁-C₃alkoxy,
C₁-C₃alkoxy mono- to tri-substituted by halogen; cyclopropyl, methylamino or
dimethylamino;
- 15 E is nitrogen or the methine group; and
Z is fluorine or chlorine; and the salts of those compounds with amines, alkali or alkaline
earth metal bases or with quaternary ammonium bases have good pre- and post-emergence
selective herbicidal and growth-regulating properties.